

**Indian Botanic Garden Library**  
**BOTANICAL SURVEY OF INDIA**

CLASS NO..... 582 .....

BOOK NO..... H00 - i : Vol - 2 . S.

ACC. NO.... A - 215 15






PLATE 2101.

CERATOSTYLIS HIMALAICA, JllooA\ /.

OHCHIDEA. Subtribe CCELOGYNEJ.


*C. himalalca*, *Hook. f. Fl. Brit. Ltd.* v. 82t>; caulibus fascicolatis brevibus simplicibus v. ramosis vaginis lanceolatis acuminatis membranaceis imbricatis opertis, pseudobulbis 0, foliis anguste lanceolatis acuminatis, floribus subsolitariis breviter pedunculatis, bractea lanceolata, c<sup>h</sup>ario sessili sepalisque araneosis, sepals lateralibus late triangulari-ovatis subacutis, petalis linear-lanceolatis acuminatis, mento brevi rotundato, labullo brevi obovato-oblongo saccato carnoRulo, dorso apicem versus gibboso et papilloso, column<sup>e</sup>D ramis brevibus. *Eria ramosissima*, *Wall. MSS.*

HAD. Eastern tropical Himalaya; E. Nepal, /, D. H.; Bhotan, *Griffith*; Khasia Hills, *Gibson (Ic. in Herb. Calcutt.)*.

*Gaules* 2-4-pollicares, dense cessitosi, simplices v. ramosi, radicibus fibrosis; vagina § - | poll. longsB, erecteo, laxe imbricate, scariosas, striato-nervosaB. *Folia* 2-2i poll, longa, subacuta v. acuminata, coriacea v. subcarnosa, sessilia v. in petiolum brevem angnstanta. *Pedunculi* vaginis immersi, uniflores, floresque sparse araneoso-pubescentes; bractea lanceolata, ovario ssquilonga. *Flores* minuti; sepala lateralia pallide rosea, 3-5-nervia; petala 3-nervia, alba, purpureo fasciata; labellum obscure 3-lobum, lobis latalibus purpureo marginatis, terminali aureo. *Colu'mna* brevissima, reniformis. *Oapsula* | poll, longa, ellipse-idea, crasse costata.

The plate is copied from a coloured drawing in the Calcutta Herbarium inscribed '*Eria ramosissima*, Wall., Mr. Gibson's Churra collection.' It represents the flowers as considerably larger than in the native specimens.—J. D. H.

Fig. 1. Flower. 2. Bract, ovary, and column. 3. Anther. 4. Pollinia. All enlarged.






PLATE 2102.

CEBATOSTYIIS LANCIFOLIA, ITtofr. /.

OBCHIDEJS. Subtribe CCELOGYNEA:

C. lancifolia, 7Io0&./..FZ. Un\$, JiaZ v. 826; caulis fasciculatis erectis simplicibua vaginis elongatis laxis ltBvibus tectis, pseudobulbis 0, foliis anguste linear-lanceolatis acatis erectis, membranaceis, floribus solitariis, pedicello ovarioque puberulis, bractea minuta tubulosa, sepalis lateralibus oblongo-lanceolatis acatis, petalis lanceolatis acatis, mento pcrbrevi 2-lobo/ labelli angasti recti hypochilo anguste oblongo epichilo angustiore pugioniforme terminate, column9 ramis erectis.

HAB. Malayan Peninsula; Perak, *Scortechini*.

*Caides* 2-4 poll, longi, radicibus fibrosis; vaginae |-|| poll, longoe, teretes, subinflatso, acutoe v. acuminatsa, striato-nervosae. *Flores* % poll, diam.; labellum carnosum, breviter ungaiculatam, epichilo tereti hypochilo planiusculo paullo longiore. *Culumnce* rami antheram longe saperantes, oblongi, obtasi.—J. D. H.

Fig. 1. Flower, with lip removed. 2. Side view of flowers. 3. Lip. *Ml* enlarged.



M.S. del et lith.

◦ *Cœlogyna occultata*, Hk f.

PLATE 2104.

CCELOGYNE OCCULTATA, *Hook. f.*

OECHIDEE Subtribe CCELOOTNEA.


*C. occultata*, *Hook.f. FL Brit. Ind. y.832*; rhizomate robusto ascende viginato, pseudobulbis turbinatim obovoideis sulcatis acute costatis, foliis ellipticis acutis, scapo e basi paeudobulbi enato ascende toto viginato, viginis coriaceis smpremis subfoliaceis, floribns paucis breviter pedicellatis viginis bracteisque obiectis albis, sepalis petalis que paullo angustioribus oblanceolatis acutis, labelli hpochilo lobis rotundatis carinis 2 crassis crenatis percurso epichilo ovato-rotundato crenato, lamina incrassata disco adnata.

HAB. Sikkim Hynalaya, *Griffith's Collectors (Ic. in Herb. Gdlcutt.)*.

*Bhizoma* crassitie penn<sup>o</sup> olorinss, internodiis brevibus. *Pseudohulbi* l~i<sup>o</sup>-pollicares, sessiles, angulis demum undulatis intervallis latis concavia lflvibus. *Folia* subaesailia v. breviter petiolata, coriacea, 2-3 poll, longa, 5-nervia. *Scapus* 1-2-pollioaris, unacum viginis crassitie perm<sup>o</sup> olorincB, viginis viridibus arete imbricatis pollicaribus acutis. *Flores* conferti; bracteaB pollicares, oblonpoe, acutae; sepala pollicaria, interdum flavo-marginata; label! nm rubro-lineolatum, lobi laterales plaga brunnea semilanai'i notatis, carinis brunneo-marginatis. *Columna* apice dilatata, troncata.

The plate is from a coloured drawing in the Calcutta Herbarium, which has supplied some characters for the description, the specimens being very incomplete.—J. D. H.

Fig. 1. Dorsal; and 2, lateral sepals. 3. Fetal. 4. Lip. 5. Column. 6. Anthor.  
*All enlarged.*



MS dele^litli.

Coelogyne Treutleri, h<sup>k.f.</sup>

PLATE 2105.

## CCELOGYNE TBBTJTLBBI, *Hook.* /.

# OBODID&B. Snbtribe CfflLOcnraia.

**C. Treutleri**, *Eooh.. Fl. Brüt. Ind. v.837*; rhfemate repente nudo,  
pseudobnlbis parvis curvis angustis teretibus basi vaginatis. » <sup>TM</sup> P <sup>TM</sup>  
elliptico-lanceolatis, scapo in pseudobnlbum terminali! <sup>b</sup> TM TM . TM £ ° °  
sepalis lateraUbus lanceolatis aoaminatis, petahs angnste \_ 1 " 1 6 " ^  
labelli hypoohUo breviusculo lobia oblongis obtnsis lamellis. 3 peroureo,  
epichilo majore orbicnlari apice obtuse tridentato disco uen.

## HAB. Sikkim Himalaya, Treutler.

*Bhizoma* crassitie pennat anatinib. *Pseudohdbi* rhizomate remoti, H PoU. longi, snerne sensim attenuati. *Folia* polh(»pia. *Scopus* fjallicaris, toii vaginatns; bractea caduco. *Flares* l poll, diam^, Sejalis 7-9-nerviis; petalis 1-nerviis; labellum basi " " »\*?• \* J ^ ohilo epiohUo breviore. *Cohmna* angnste alata, apioe rotundata apicnlata.

I have seen only the one specimen here figured.—J. D. H.

Kg. 1. Dorsal Bepal. 2. Petal. 8. Lip. *M* enlarged.



M.S. de leth.

*Coelogyne steno-iila, Hk.f.*

PLATE 2106.

CCELOGYNE STENOCHILA, *Hook. f.*

OBCHIDEJE. Subtribe GCELOOTNE<sup>^</sup>.

C. **Stenochila**, *Hook. f. Fl. Brit. Ind.* v. 837; rhizomate repente, pseudobulbis parvis teretibus basi vaginatis, foliis sessilibus ovato- v. elliptico-oblongis, scapo in pseudobulbū terminali brevi basi et apice undo, racemo elongato multifloro flexuoso, floribus inter minoribus albis, sepalis lateralibus ovato-lanceolatis, petalis adguste linearibus, labelli hypochilo elongato in lobos parvos obtusos producto, epichilo breviore latiore orbiculari, disco tricarinato.

HAB. Malayan Peninsula; Pera<sup>k</sup>. summit of Gunong Batu Pateh, alt. 6,700 feet, Wray.

*Rhizoma* crassitie pennro corvinro, nudumP. *Pseudobulbi* J-f-pol-Kcares, laves. *Folia* 1-1<sup>^</sup> poll, longa, f-§ lata, subacuta, rigida, subtus 3-5-costata. *Scapus* compressus, foliis brevior; racemus 8-pollicaris, articulatus; bracteae decidu<sup>O</sup>. *Flores* 1 poll. diam.; sepalae 3-nervia; petala uninervia; labellum basi saccatum. *Columna* apice truncata, 3-loba.

The only specimen is that here figured; the flowers were detached.—J. D. H.

Fig. 1. Flower, with one lateral sepal and a petal removed. 2. Lip. Both enlarged.



M.S.del.et. Hilt

*Coelogyne carneae*, Hk.f.

PLATE 2107.

CCELOGYNE CABNEA, *Hook. f.*

ORCHIDEJ.; Subtribe CCELOGYNEJ.


*C. carnea*, *Hook./. Fl. Brit. Tnd. v. 838*; rhizomate valido vaginato,  
P<sup>sus</sup>r<sup>ud</sup>. bulbis elongato-fusiformibus teretibus nudis, foliis petiolatis  
elhpticis obovatis v. oblanceolatis, scapo in pseudobulbum terminali  
breviusculo erecto compresso basi et apice nudo, racemo elongate multi-  
floro flexuoso, sepalis oblongo-lanceolatis acuminatis, petalis elongato-  
subulatis, labelli hypochilo brevi, lobis oblongo-rotundatis epichilo rotun-  
dato emarginato incumbentibus, disco basin versus lamellis 2 instructo,  
columna superne dilatata apice obtusa.

HAB. Malayan Peninsula; Perak, *Scortechini*.

Rhizoma crassitie pennao olorinsB; vaginis brevibus, terminali longiore  
Dasin pseudobulbi vaginante. *Pseudobulbi* 3-3<sup>1</sup>/<sub>2</sub>-pollicares, J poll, diam.,  
J<sup>2</sup>/ves. *Folia* 4-5 poll, longa, 1-1<sup>1</sup>/<sub>2</sub> poll, lata, 1-costata, multinervia.  
*Scopus* unacum racemo longiore folia superansj bractes caducss.

Fllores 1<sup>1</sup>/<sub>2</sub> poll. diam.; sepala carnea, nervis numerosis trabeculatis;  
petala 1-nervia. *Labellum* basi saccatum, carneum, venosum. *Oolumna*  
mfra medium constricta, superne 2-alata i anthera rostrata vertice  
ambonata.—J. D. H.

2. <sup>Fl?</sup> <sup>l</sup> <sup>Flowe</sup>r, with lip and portions of one lateral sepal and petal removed.  
• <sup>A</sup>P- 3. Anther. All enlarged.



itS.deletlith.

Cpelogyne Gnffithn, Hk.f.

PLATE 2108.

CCELOGYNE GRIPPITHII, *Hook. f.*

ORCHIDEJS. Subtribe CCBLOGYNEZE

C Griffithii, *Hook. f. Fl. Brit. Ind. v. 838*; rhizoma validum, pseudobulbis lineari-oblongis compressis 4-quebris vaginis magnis cymbiformibus tectis, foliis petiolatis elliptico-lanceolatis acuminatis, scapo in pseudobulbo terraiiali compresso basi nudo apice infra racemum bracteia ovatin arete distiche imbricatis tecto, racemo multifloro rachi angulatim flexuoso, floribus inter minoribus, sepalo dorsali late oblongo, lateralibus ovato-lanceolatis acutis, petalis anguste linearibus, labelli hypochilo ampio lobis rotundatis erosio disco medio lamellis 5 brevibus instructo, epichilo parvo quadrato truncate

HAB. Upper Assam; summit of the Patkoy Mts., *Griffith*; Munaiapore, alt. 4,000-5,000 feet, *Watt.*

*B*/Msowa crassitiepennceanserinaB. *Pseudobulbi* 3-4-pollicares. *Folia* 5-10 poll. loDga, 2-3 lata, 7-costata, petiolo 2-pollicari; racemi erecti i\*achis valida articulata; braotero caducBD. *Flores* 1 poll. diam.; ovario unacnm pedicello sepalis breviore; sepalala 5-7-nervia; petala 1-nervia. *Golumna* apice cacullata, obtusa.—J. D. H.

Fig. 1. Flower. 2. Lip. 3. Column. All enlarged.



M.S. del et lith.

Coelogyne ancops. Hk-f.

PLATE 2109.

CCELOGYNE AJNCEPS, *Hook. f.*

ORCHIDEJ.: Subtribe *CCBLOQYKEM*.

C. anceps, *Hodh.f. Fl. Brit. Ind.* v. 840 ; rhizomatous crasso vaginato, pseudobulbs ellipsoideis v. fnsiformibus teretibus diphyllis, foliis breviter petiolatis ellipticis acutis, scapo in pseudobulbum terminali valido compresso basi nudo apice infra racemum bracteis coriaceis arete imbricatis tecto, racemi rachi articulato, sepalis linearis-oblongis acutis, petals angusti linearibus, labelli hypochilo elongato in lobos laterales preves obtusos producto, epicnophilo Jatiore rotundato, disco lamellis 2 revibus instructo, columna subacuta.


HAB. Malayan Peninsula; Perak, Scortechini.

~~rhizoma~~ repens, crassitie penns anserines; vaginsd breves arete imbricata^ obtuse. *Pseudobulbi* nndi, substipitati. *Folia* 2^-4 poll, longa, J-1^ iata 7-9-costata, in petiolum validum angustata. *Seapus* unicum racemo erecto laxifloro 5-10-pollicaris, anceps v. anguste laiatus; bractea J poll, longa; ovarium unicum pedicello J-pollicare. •P cores 2 poll. diam. *Labelli* hypochilum epichilo duplo longior.

A handsome species, remarkable for the acute tip of the column, and very small clinandrinm.—J. D. H.

Fig. 1. Flower, with one lateral sepal removed. 2. Lip. 3. Column. All enlarged.

nwo.



M S deletlrth

Oeceogyne purascens, Hk f

PLATK 2110.

CCELOGYNE ? PURPURASCENS, *nook. f.*

ORCHIDE^. Sabtribe CCSLOGTNEJ:

. C.? purpurascens, *Jlooh f. Fl. Brit. Tnd. v.842*; ceespitosa, pseudo-bulbis angustis teretibns monophyllis, folio linearis-oblongo v.lanceolato, spapo e basi pseudobulbi erecto gracili 1-3-flore, bracteis parvis, Pepalis Hneari-oblongis obtusis, petalis linearis-subspathulatis, labello basi columnse sessili spathulato concavo basi saccato disco nudo, columnna angustis alata. *Dendrobium purpurascens*, *Thwaites, Enum. Pl. Zeyl.* 298.

HAB. Ceylon ; in the elevated part of the Central Province,  
*Thwaites.*

, Caules breves, fastigiati, radices crassas elongatas tortas epidennide piflato emittentes. JPdillbi ^§ poll longi g Folia 1-4 poll, longa, < 2 poll, lata, coriacea, rigida, soepe purpurascens, sicca marginibus jecurvis. Scapus 3-4.pollicaris, vaginis paucis parvis. Flores f poll, " ^m., fere albidi; pedicello cum ovario : } - ^ poll, longo; sepala erecto-patentia, 3-nervia; petala 1-nervia; labellum suberectam, obscure crenulatum. Colmnia erecta, apice dilatata, lateribus dentatis; Jinthera imperfecte 4-locularis, persistens, purpurea; pollinia 4, libera, znajora subpyriformia, minora ovoidea. Copsula -f pol. longa, anguste oblonga v. pyriformis, alte 6-costata.

A curions little plant, certainly not a *Dendrobium*; nearer to *Callogyne* or *Panisea* than to any other genus ; differing from the former in the entire spatulate lip not embracing the column, and from the latter in the lip having no claw.—J. D. H.

Views. 1. Front; and 2, back view of flower, with the lip depressed. 3. Lateral view of flower; 4. Lip (its concavity not represented). 6. Anther. 7. Pollinia.  
Anther enlarged.

n 2111.



WLS.del et lith.

Calarvthe diploxyphion, lik ♀

TLATK 2111.

CALANTHE DIPLOXIPHION, *Hook. f.*

OUCHIDEJ.: Subtribe CIELOGYNE\*:

*C. diploxyphion*, *Hook. f. Fl. Brit. Ind.* v. 849; caule brovi, foliis paucis amplis clongato-lanceolatis in petiolum latum angustatis, scapo clato remote vaginato, floribus in racemum brevem congestis pubescentibus, bracteis magnis late ovatis acuminatis pedicellis gracilibus multo brevioribus, sepalis lateralibus ovatis acutis v. acuminatis, petalis obovato-spathulatis v. oblanceolatis, labello subequaliter 4-lobo, lobis angustis falcato-recurvis, disco basi minute carunculato, calice gracili mcurvo pedicello breviore.

HAB. Malayan Peninsula; Perak, at Goping, *King's Collector*.

- *Caule* brevi, rhizomate lignoso. *Folia* 2, 12-20 poll, longa, 3-4 poll, lata, 5-7-costata, multinervia, petiolo valide costato. *Scapios* 1<sup>1</sup>-2-pedalis; *vaginas* basi tubulos<sup>o</sup>, inferiores pollicares, ovato-lanceolatea, acuminata<sup>b</sup>; *racemus* 2-pollicaris; *bractae* 1 poll, long<sup>o</sup>, subimbricatoe prides, pedicello cum ovario 1<sup>1</sup>-pollicari. *Flores* 1<sup>1</sup>-poll, diam., \*bi; *labellum* eepalatis longius. *Odlumna* brevis; *anthera* deppressa, rostrata.

If I have been on a <sup>a</sup>\*<sup>b</sup> one specimen were figured; it is most nearly fitted to the Himalayan *O. alismcejolia*, the lip of which has an obcordate midlobe and a large yellow callus at the base. Except in <sup>a</sup> much smaller flowers, and very short raceme and larger bracts, it resembles *G. v&ratrifolia*.—J. D. H.

*Fls* 1<sup>1</sup>-poll, <sup>a</sup>\*<sup>b</sup> wide (\* the lateral sepalis much too small and narrow). 2. Column, with Mae of hp and spur. Both enlarged.



M.S.del el lith.

*Aiundinia* Bentleyi, Hk.f.

PLATE 2112.

ARUNDINA CANTLEYI, *HooL* /.

ORCHIDEA; Subtribe CCELOGYNK<sup>A</sup>.

A. (*Dilochia*) **Cantleyi**, *Hook. f. Fl. Brit. Ind.* v. 858: caulis robustis foliosis, foliis sessilibus ovatis v. ovato-lanceolatis longe acuminatis basi rotundatis, floribns in paniculam terminalem deourvam latam foliaceo-bracteatam dispositis, bracteis cymbiformibus, sepalis oblongo-lanceolatis falcatis dorso infra apicem corniculatis, petalis nneari-oblongis, labello parvo trilobo lobis lateralibus oblongis v. subrotundatis terminali flabelliformi, disco carinis 5 crassis crenatis percurso, columna apice dentata.

J5<sup>B</sup>- Malayan Peninsula; Perak, on Gunong Batu, alt. 4,500-5,500 feet, *Oantley*<sub>g</sub>. *Wray*.

*Caulis* 2-pedalis, crassitie pennaB olorinaa, vaginis foliorara 1-1 poll. longis teotus. *Folia* 2f-3J poll, longa, subundulata, unicostata et nanltmervia. *Panicula* sessilis, ramis rigidis divaricatis basi spathaceo-oracteatis; bractea florales ^\ poll, longoe, subacutas, subcoriacetB, striataB. *Flores* 1^1-^- poll diam., straminei, sanguineo-venosi; sepalis petahsque patentibns, labello basi cuneato.

,4 <sup>re</sup>markable plant, allied to *A. (Dilochia) Wallichii* of Singapore, which has elliptic-lanceolate leaves and few erect branches of the panicle. It much resembles *Eria Scortechini* (see plate 2082) in habit and inflorescence.—J. D. H.

of <sup>1</sup>\*!?<sup>2</sup> la Ltttolal sePaL 2 and 3. Varieties of lip. 4 and 5. Side and front views <sup>\*\*</sup> column. 6. Anther. 7. Pollinia. All enlarged.



M.S. del. et lith.

Calanthe ifaimil Hk. f

PLATE 2113.

CALANTHE MANNII, *llooh f.*

ORCHIDEA.: Subtribe CCELOGYXEA;

C. Mannii, *Book. f. Fl. Brit. Ind.* v. 850; canle brevi, foliis lan-

ceolatis, scapo gracili pubescente foliis multo longiore, racemo

multifloro, bracteis minutis, floribas parvis pabescentibus, sepalis

ovato-lanceolatis, petalis oblanceolatis, labello parvo 3-lobo, lobis

lateralibus oblongo-rotundatis, terminali paullo majore orbiculari

finiarginato v. obcordato, disci carinia 3 apices versus lamellaformibus

instruct\*), calcare brevi obtuso.

—HAB. Western Himalaya; Kumaon, below Ramkhet, *Dutliie*; the

Khasia Hills, alt. 4,000 feet, *Mann, Clarke*.

• *Gaulis* 2-3-pollicaris, vaginatus. *Folia* 5-6 poll longa, sessilia,

p-costata, multinervia. *Scapus* nnacum racemo 6-10-pollicaris et

laflorescentia pabescentes; bracteeB ^ poll, long®, membranaceas.

*F*f^j^i poll, diam., nutantes; sepala 3-5-nervia; petala 1-3-nervia;

*lat~lum* basi colamns brevi adnatum, carinis disci validis, siccis

*rugulosis*.—J. D. H.

?• V. Side; and 2, front view of flower. 3. Column and lip. 4. Lip. *Ml*

*enlarged*.



M.S.del, et lith.

Calanthe Wrayi, Hi c.f.

PLATE 2114.

**CALABTTHE WBAYI,** *Uook.f.*

ORCHIDEA: Subtribe CCELOGYNEJ:

**C. Wrayi**, *HooJc. f. Fl. Brit. Ind.* v. 850; snbacaulis, foliis longe petiolatis ellipticis acuminatis, scapo interfoliaceo, racemo gracili puberulo multifloro, bracteis latis rigidis revolutis, sepalis ovato-oblongis acutis, petalis oblongo-lanceolatis, labello sub&qnaliter 4-lobo, lobis oblongis obtusis divaricatis, disco basi callis parvis pectinatis instructo, calcare gracili sepalis paullo longiore.

HAB. Malayan Peninsula; Perak, alt. 2,000 feet, *Wray.*

*Folia* 8-10-pollicaria, acuminata, 5-7-costata, multinervia. *Scapus* • unacum racemo 2^-pedalis, gracilis; racemus 3-6-pollicaris, laxiflorus; bractesB ovatce v. ovato-lanceolatse, persistentes, siccsB saope tortte; pedicello unacum ovario f poll, longo. *Flores* § poll, diam., albo-Jilacini; sepala lateralia dorso infra apicem alato-carinata; labellum lateribus column® adnatum.—J. D. H.

Fig. 1. Flower. 2. Lip and column. *Both enlarged.*



MS.del.-tith.

Eulophia Manra: Hk.f.

PLATE 2115.

**EULOPHIA MANNII, JT onhf.**

**ORCHIDÆ.** Subtribe EULOPHIEJS.

.. **E. Mannii**, *Hook. f. Fl. Brit. Ind.* vi. 4; foliis floribus coctaneis uneari-elongatis subacutis, scapo elato, bracteis majuBcnlis oblongo-lanceolatis obtusis ovario paullo brevioribus, sepalis lateralibus basi column@ insertis oblongo-lanceolatis petalisqne oblongis multinerviis, labello latiore quam longo rlicmbeo-ovato obscure 3-lobo obtuso, disco Tiurvis 3 crassis ultra medinm ramosis v. in pulvinara velutinam conuentibus percurso, calcare conico brevi. *Cyrtopera Mannii*, *Reichb.f.* in *Flora*, 1872, p. 224.

HAB. Upper Assam, *Mann*.

*Caulis* basi crassitie digitii minoris. *Folia* pedalia, \ poll. **lata**. *anajnu* nnacum racemo laxifloro 3-pedalis; bracteaa £-£~poll. longie. *iHortis* 1 poll. diam. *Lahellum* sepalis paullo brevius, lobis lateralibus rotundatis termiuali obtuso. *Columna* clavata, apoda; **anthera** de-pressa; pollinia globosa.

In the 'Flora of British India' I have followed Blume in removing *tyrppera* from *Cyrtopodium*, and placing it under *Eulophia* as a species, characterised by the column having a foot, and the insertion of the lateral sepals on the raentum or spur of the lip. *E. Mannii* was referred to *Cyrtopera* by Reichenbach, but, as I can find no trace of such insertion of the lateral sepals, I am obliged to refer it to *Eulophia* proper. Its affinity is certainly with *E. pratensis*, LindL, and *macrost'xhya*, Lindl.; but it differs from these and all others of the group to which these belong in the very narrow leaves, as well as in characters of the lip.—J. D. H.

7 ^8: if. JatoMi sepal. 2. Petal. 3. Column and lip. 5. Column. 6. Anther.  
--Pollinium AV enlarged.



M.:s.dd et lith.

*Eulophia* i holoechda. Coll. & Herasl.

PLATE 2110.

**EULOPHIA HOLOCHILA**, *Coll et JTemsl.*

ORCHIDEJE. Sabtribe EULOPHIEJS.

**E. holochila**, *Coll. et Hemsl. in Journ. Linn. Soc. xxviii. 132*; foliis hysterantheis linearibus, scapo valido elato vaginato, racemo laxifloro, bracteis lanceolatis ovario ©quilon^is, floribus magnis, sepalis erecto-patentibus oblongis obtusis lateralibus calcari adnatis, petalis obovato-oblongis obtasis, labello orbiculnri recarvo basi in calcar incnrvum obtusum producto flabellatim nervoso disco nudo ecarinato. *Hook.f.* *FL Brit. Ind. vi.* 1S|g.

HAB. Burma; on the Shan Hills, alt. 4,000 feet, *Gen. Sir B. GoJlett-*.

*Caulis* foliosns, validus, crassitie digitii minoris. *Folia* vix evoluta, 4-5 poll, longa, £ poll, lata, in apicem obtusam angustata. *Sca^ns* lateralis, unacum racemo pedalisi, validus, vaginis laxis spathaceis instructus; bracteaB § poll, longed. *Flores* 2 poll. diam.; sepala 7-nervia; petala 5-nervia; labellum petalis asquilonatum, basi tantum columnam amplectens. *Golumna* incurva; anthera apice 3-mammillata.

I have seen but one specimen, that here figured.—J. D. H.

Fig. 1. Sepal. 2. Petal. 3. Column and lip.' 4. Column. 5. Anther, seen from above. 6. Pollinia. All enlarged.



M. S. de laet & N. H.

*Cymbidium sikkimense*, Hk.f.

PLATO 2117.

CYMBIDIUM SIKKIMENSE, *Hook. f.*

OBCHIDEJE. Subtribe CYMBIDIEJG.

. C. **Bikkimense**, *Hook. f. Fl. Brit bid.* vi. 9; epiphyticum, foliis post mflorescentiam bene evolntis ensiformibus, scapo vasfinato, racemo multiicro, bracteis minutis, sepalis patentibus lateralibua lanceolatis subfalcatis, dorsali linear-i-oblongo erecto, petalis elliptico-ovatis acatis, labello rhombeo-ovato patente integro v. obscure 3-lobo, disco callis 2 parvis inatructo ceterum ecarinato et inappendicukto.

HAB. Sikkim Himalaya; Lachen Valley, alt. 6,000 feet, /. D. U.

*Rhiizomob* crassum. •*Scapus* unacura racemo pedalis, decurvns; vaginis compressis basalibus brevibus, saperioribus elongatis foliaceis; bracteae 6 poll, longas, ovato, acufaa: ovarium unaoam pedicello i-\$ poll, longo. Mores li poll, diam!

.. Tbis species I found only twice, on both occasions in full flower in the month of May, with the leaves (as represented) in a very young state. It is the only plant of the genus known to me- in which the flowers precede the foliage, except the very carious *G. macrorhizon*, final; a terrestrial parasitic sp with d r o u d rootstock, 2 <sup>WQIc</sup> b the foliage is unknown, if indeed it bears any other than sub-lonaceous sheaths at the base of the scape.—J. D. H.

\*. Anther. All enlarged.



M<sup>3</sup> del etlith.

Thecostele Maingayi, Hk.f.

PLATE 2118.

# THE COSTELE MAINGAYI, *Hooch.* f.

## ORCHIDEA: Subtribe CTMBIDIE^I.

**T. Maingayi**, *Hook. f. Fl. Brit Ind.* vi. 20; foliis ellipticis y. obovato-oblongis in petiolum angustatis, sepalis lateralibus late ovatis acunniatiforme, petalis lanceolatis acuminatis, labelli parvi lobis lateralibus flabelliforinibus terminali late ovato, auriculis basi uncinatim incurvis obtusis.

HAB. Malacca, *Maingay*.

— *Pseudobulbi* §-1-pollicares, late ovato-oblongi, compressi, altecostati.  
 • *Folia* 3-4 poll, longa, 1-1<sup>½</sup>- poll, lata, obtusa, nervosa. *Bacemus*  
 \*-o-pollicaris, decurvus, laxiflorus, breviter pedunculatus; bracte©  
 parv», ovateB. *Flares* 1 poll. diam.; sepala 5-nervia, dorsali angustiore;  
 petala 3-nervia; labellum parvum, velutinum.

Under plate 2063 of this work I have, when describing *Ollalium Wrayi*, suggested the transference of that genus to *Vanclece*, and a comparison of the structure of its flower with that of *Tliecoslele* supports this view. The column, with its long arms and tubular base, <sup>t in</sup> <sub>T^W</sub> Serti011 on the latter of the lateral sePals and the small bairv lip. with two basal auricles or lateral lobes (see note under the following piate). are identical in the two genera. A better knowledge of the pollinia <sup>wh !\* frOm livin^</sup> specimens is in both wanted, before determining ref e. et tlely stoll ^ ^ e retained as separate genera, and whether 27 & <sup>tre</sup> i\*<sup>0</sup> VandecB or EpidendrecB. The figure of the pollinia of <sup>the LinDea^ Soc iet y' (xxx- \*• 29)</sup> *Zollingeri (alata, P. et R. f.)* given in the 'Transactions of (who) requires confirmation. Maingay <sup>ficrint rega us this as a form o^ quinquejida)</sup> describes it in his manu- 8tran^ and fi nreB th C pollinia as four, terminating a long narrow Vanf <sup>lated</sup> upwards and with a glandular base: hence truly eous.—J. D. H.

**Fig.** 1. Flower. 2. Lip. 3. Column. *All enlarged.*



itS.dflLetltth

,,Thecostele quinquefida, H.<sup>kf</sup>

PLATE 2119.

THECOSTELB QITINQUEFIDA, *Hook.*/.

ORCHIDEJJ. Subtribe CYMBIDIÆ.

. T. quinquefida, *Honh.f. FL Brit. Lid* vi. 20; foliis linear-i-elongatis in petiolum angustatis, sepalis lateralibns late ovatis acutia dorsali oblongo, petalis lioeari-lanceolatis, labelli 3-lobi lobis lateralibus acmaciformibus terminali late ovato, auriculis basi linearibns erectis.

HAB. Malacca, *Maingay*.

- *Pseudohdbi* 1-1<sup>1</sup>-pollicares, ovato-oblongi, compressi, obtuse costati. *Foha* 5-7 poll, longa, 1-1<sup>1</sup>- poll, lata, apice obliquo obtuso. *Eacemvs* ^3-pollicaris, gracilis, decurvus, breviter pedunculatus, laxiflorus; bracteaB parvsB, ovat^e. *Flores* f poll. diam.; sepala lateralia patentia, o-nepvia, dorsali incurvo; petala obtusiuscula, 3-nervia; labellum pubepulum.

. In this genus the lobing of the lip has been variously described, dependent on the view taken of the small divisions at the base of the lip. T. <sup>have described nnd</sup> ~~ritish~~ <sup>er</sup> *quinquefida* in the \* Flora of India, as the normal side lobes of the lip, and regarded the broad blade as a 3-lobulate midlobe. I now think that the smaller divisions should more properly be regarded as auricles or appendages of the lip, <sup>and</sup> <sup>(\*)</sup> <sup>avenence so</sup> described them here in this and the preceding <sup>peciles</sup> <sup>For an</sup> observation on the affinities of *Thicoste*, see near the description of the preceding plate.—J. D. H.

Fig. Dorsal llZi<sup>l</sup>We<sup>i</sup>P<sup>l</sup> ^ th Po<sup>Pti</sup><><sup>n</sup> of on<sup>©</sup> lateral sopal and of one petal removed. 2. sepal, column, and lip. Both enlarged.



M. S. del et lith.

Diploprora Charapioni. Hkf.

**DIPLOPRORA CHAMPIONI, Uouk.f.****ORCHIDÆ.** Subtribe SAKCANTHHA.

*Diploprora*, *Hook. f. Ft. Brit. Ind.* vi. 26; *Sepala patentia*, late ovata, obtus-a, dorso alato-carinata. *Fetal a* ovate)- v. obovato-oblonga, obtusa. *Labellum* sepalis aequilongum, marginibus basi columns adnatis, cymbiforme, abrupte in caudam a latere corapressam apice bicaudatam angustatum, disco carina erecta lamelliforini instructo. *Oolumna brevissima*, apoda. *Anthera* 2-locularis; pollinia globoso-obovoidea 2-loba, supra apicem stipitis linearis inserta; glanduia squamaeformi.—Herba epiphytica, caule simplichwculo, basi vaginato radicante, sverne crassiuseulo foliosu. Folia disticha, ovafo- v. oblongolanceolata, subfalcata, flaccida. Pedunculus oppositifolius, crassus, erectus, pauciflorus; bractee majuscule, obtusce. Flores erecti, sessile\*, flcwidii, labello roseo-tincto ; ooario brevi.

**D. Championi**, *Hook. f. FL Brit Ind.* vi. 26. *Cottonia Championi*, *Lindl. in Hook. Journ. Bot.* vii (1855) 35: *in Journ. Linn. Soc.* iii. 39; *Benth. Fl. Hongk.* 357. *Luisia bicaudata*, *Thwaites, Enum. PL Zeijl.* 302. *Vanda bicandata*, *Thwaites*, \. c. 429.


HAB. Bhotan Himalaya, *Gammie*; Khasia Hills, alt.. 3,000-4,000 feet, *J. D. H. et T. T.*; Tenasserim, *Parish*; Ceylon, in the Ambajramowa and Saffragam districts; Hongkong, on Victoria Peak, *Champion*.

*Gaulis* 2-4 poll. *Folia* patnla, 3-4 poll, longa, 4-J poll, lata, membranacea, apice acuto obtuso v. emarginato, sicca multiuervia. *Pedunculus* foliis longior, basin versus puuei-bracteatus, strictus v. subflexuosns; racemus erectun. *Flores* \ poll, longi, sepalis 7-nerviis, petalisque 5-nerviis aureis, labello albo roseo irrorato; ovario sigmoideo.

9 f the three genera to which this curious Orchid had been referred, is perhaps nearest to *Vanda*, from which it differs in habit\* the Membranous leaves, small flowers, broad bases of the sepals, and remarkable lip. From *I/iisia* the flat membranous leaves at once distinguish it. *Cottonia* differs totally in habit, and in the large flat JP! In short, the lip of *Diploprora* is quite peculiar in its long narrow forked tip, and the broad flat erect longitudinal plate that occupies its disk.

th Th<sup>116</sup> description which is here given will supersede that of "Flora of British India," being aided by a beautiful coloured drawing by Mr. Parish, made in Moulmein in August 1869.—J. D. H.

<sup>Ej?" 1% "t,loveip-</sup> <sup>\* Sepal. 3. Column and lip. with one half of the body of the</sup>  
<sup>letter ren^vPd. 1. Anther. 6 and 6. Pollinia. All enlarged.</sup>



M.S. del e: lith.

*Sarcochihis hiriulus*, Hk.f

PLATE 2121.

SARCOCHILUS HIRTULUS, *Uvhf.*

ORCHIDÆ. Subtribe SARCANTHEJE.

*S. hirtulus*, 'Hook. f. Fl. Brit, Ind. vi. 39; caulis brevibns fasciculatis, foliis anguste lanceolatis falcatis obtuse acuminatis, scapis vaginas iuferiores perforantibns foliis mnlto brevioribus gracilibns hirtnlis remote paucifloris, bracteis minntis, sepalis ovatis v. ovato-oblongis subacutis, petal is augustioribus, labello breviter unguicnlafco basi in eaccum elongatum inflatum producto 3-lobo, lobis lateralibns falcato-recurvis, terminali 2-partito segmentis divaricatis falcatis obtnsis lobis lateralibus sequilnngis et paulJo latoribns.

HAB. Malayan Péninsula; Perak, Scortechini; Malacca, Maingay.

*Caules* ^-^<sup>o</sup>-pollicares, basi radicantes. *Folia* undiqae inserta, 3-5-pollicaria, ^-§ poll, lata, coriacea, basi et apice angustata, apicibus ssepe recnrvis, vaginis perbrevibus. *Scapi* decurvi, ^-1 poll, longi; bracteis paucis ovatis patulis. *Flares* h poll, diam., sepalis petalisque patentibns 3-nerviis pallide aurantiacis rübro-punctatis. *Labellum* sepalis brevins, quasi 4 partitum, pallidum, sacco v. calcare lobis eequilongum obtuso *Cohmmi* labello aequilonga, basi attesuata, apice utrinque calcarata et dorso stibulata incurva. *PoUinia* minuta, globosa, in apicem truncatam glandulaD majusculs triangularis sessilia.—J. D. H.

Fig. i. Portion of scape. 2. Flower. 3. Dorsal sepal. 4. Lip. 5. Follinia.  
*Gr&itly enlarged.*



U S del ethii

Sarcochilus recurvus, Hk f.

PLATE 2122.

SAKCOCHILUS RECURVUS, *Hook. f.*

OBGHIDEJB. Snbtribe SABCANTHEJE.

**S. (Fornicaria) recurvus**, *Hook. / Fl. Brit. Ind. vi.* 39; canle brovi radicante, foliis linearis-oblengis loriformibnsve planiusculis apice rotundatis dorso costatis, scapis nndis erectis foliis multo longioribus, spica (v. racemo) brevi cylindracea, bracteis dense imbricatis ovato-lanceolatis carnosulis recurvis.

HAB. Malayan Peninsula: Perak, on the Limbo Hills in the Larut Province, *King's Collector.*

*Folia* 1<sup>^</sup>-2 poll, longa, ^-\ poll, lata, basi vix angustata, vaginis £ poll, longis striatis. *Scapi* axillares v. vaginas perforantes, 3—5 poll, longi, atrictiusculi, omnino nudi; spica ^-§ poll, longa, ^ diam. l&ta; bractesB i poll, longee.

Though flowerless, I have not hesitated to name and describe this species, which is so marked in character that it cannot be confounded with any hitherto described.

In a note attached to the specimen the flowers are described as light red mottled with light brown.—J. D. H.

n.ztzs.



MS.del.flUrth.

*Sarcochilus trichoglottis*, ftk.f

PLATE 2123.

SARCOCHILUS TRICHOGLOTTIS, *nook. f.*

**ORCHIDÆ.** Subtribe SARCANTHEJ.:

8. (*Fornicaria*) *trichoglottis*, *Hook f. Fl. Brit. Ind.* vi. 39; caule elongato folioso, foliis distichis linearis oblongis emarginatis coriaceis dorso carinatis, scapis axillaribus robustis medio 1-bracteatis, racemo brevi paucifloro, bracteis panicis imbricatis, sepalis Kneari-oblongis obtusis, petalis acutioribus linearis-lanceolatis, labello sessili saccato orbiculari v. subspathulato apiculato ciliato et parce hirsute


HAB. Malayan Peninsula; Perak, *Scortechini*; Singapore, *Bidley*.

*Caules* 3-5-pollicares, inferne radicantes, vaginis foliorum J-<sup>^\wedge</sup> poll, longis striatis obtecti, *Folia* 2-2<sup>^\wedge</sup> poll, longa, £-£ *Poll*-<sup>lata</sup>» coriacea, planiuscula, enervia, basi angustata. *Scapi* axillis inferioribus erecti, norizontalen, foliis breviores; basi breviter vaginati; vaginis tubulosis; bractea ad medium trianguli-ovata; racemi bracteis £ poll, longis ovatis subacutis carnosulis undique insertis imbricatis. *Floras* pallide flavi. *Sepala* lutealia £ poll, longa, basi columnis apodæ mserta. *Labelli* saccus late conicus, obtusus, lobi laterales vix conspicui terminalis apiculiformis.

Allied to the Bunnan 8. *Hystrix*, Reichb. f., differing in the much shorter and broader leaves, and in the form of the lip, which is not glandular. I have seen no flowers, and have described them from drawings by Father Scortechini and Mr. Ridley.—J. D. H.

<sup>Fig.</sup> 1. Flower. 2. Sepal. 3. Petal. 4 Column and lip. 5. Lip. All enlarged.

K2I24.



MS.deUUith

Sarcochilus filifonnis, Hk.f.

PLATK 2124.

SARCOCHILUS FILIFORMIS, *Huuk. f.*

ORCHIDÆ. Sabtribe YANDEX.


S. (Fornicarfa) filiformis. *Hook. f. Fl. Brit Ind.* vi. 40; canle el on gat o terete vaginato, foliis elongatis filiformibus obtusiusculis, scapis foliis multo brevioribus nudis apice paucifloris, bracteis paucis minutis ovatis imbricatis, floribns albis, sepalis petalisque subsimilibus oblongo-lanceolatis obtusis, labello basi constricto lateribus involutis dein 3-lobo, lobis lateralibris alseformibus rotundatis terminali in saccum oblongum puberulum ore parvo prod a c to.

HAB. Malayan Peninsula; Perak, on trees, *King's Collector.*


*Oaulis* 3-pollicaris, vaginis inclusis ad J poll, diam., in feme radicans; vaginae ^ poll, long®, tubulosce, oblique truncataB, costatae. *Folia* 8-10 poll, longa, ^ poll, diam., teretia, laBvia, erecto-patentia. *Scapi* 2-3 poll, longi, basin versus caulis enati, vaginas perforantes, horizontales, snbflexuosi; racemus |-% poll, longus, pauciflorus; bracteoe 3-6, iV poll, long®, obtusae, carnosules; flores albi, pedicello unacum ovario 3 poll, longo. *Sepala* % poll, longa, petalaque 7-nervia. *Labellum* parv am, basi quasi in unguem brevem (columns pedem P) constricta. *Columna* brevis.

I have seen but one specimen of this singular species with a solitary flower available for analysis, and am not sure that the structure of the base of the lip is accurately described above, or represented in the Pj&te, its texture being exceedingly delicate. What appears in the plate as a claw to the lip, I regard as the prolonged foot of the column, to which the lateral sepals are adnate at their bases.—J. D. H.

**Fig.** 1. Flower. 2. Lip. *Both greatly enlarged.*



A



B.

M. Sæl, et lith.

A *Sarcochilus meruensis*, Hk.f

B. *pu^ionifolia*, Hk.f

PLATE 2125.

A. SABCOCHILUS MEBGUENSIS, *Hook. f.*

B. SAEGOCHILUS PUGIONIPOLIUS, *Hook. f.*

OBCHIDEJB. Sabtribe SARCANTHEJE.

A.—S. (*Fornicaria*) aerguensis, *Hook. f. Fl. Brit. Ind.* vi. 40; pusillus, caule brevi fblioram vaginis distiche imbricatis tecto, foliis linearis-oblongis obtusis, scapis axillaribus gracilibus, floribus in rachim brevem compressam bracteis arete imbricatis subdistichis pectinatim dispositis, capsulis anguste fusiformibus scapos ©quantibus v. iis iongioribus teretibus stylo brevi et stigmate capitellato terminatis.

HAB. Tenasserim, at Mergui, *Griffith (in Herb. Lindl.)*.

*Caules* £-f unc, suberecti, compressi; vaginre foliorum £-£ unc., truncates, striato. *Folia* 1-1 $\frac{1}{2}$  unc. longa, sabcociacea. *Scajpi* plures, filiformes, medio 1-bracteolati; bractera florales ovata, acute, suberectaB, 4s P<sup>0</sup>^\* longas. *Capsulce* 1-1<sup>1</sup> poll, long©, £ poll, diam., utrinque attenuates, breviter pedicellato, striato, stylo T<sup>1</sup>T~A P<sup>0</sup>^« longo.

A curious little species, of which the specimens are very incomplete, and in fruit only.—J. D. H.

B.—S. *pugionifolius*, *Hook.f. Fl Brit. Ind.* L c. 196; caule perbrevi compresso, foliis elongato-subulatis v, *pugionifbrmibus* acuminatis crasse coriaceis punctatis supra canaliculatis apicibus setiformibus, scapis axillaribus foliis brevioribus, floribus in rachim brevem bracteis squarroso-imbricatis dispositis, sepalis ovato-oblongis obtusis latere inferiore gibbis petalisque oblongis obtusis sti-amineis, labello albo aurantiaco maculato basi columnes inserto breviter unguiculato oblongo saccato truncato ore 3-lobo ciliato, intus bimaculato et carunculo parvo ciliato basin versus lobi medio instructo et basin versus disci sculta ciliata acuto.

HAB. Ceylon; at Varuniya, in the North Central Province, *Trimen*.


*Caulis* vaginis distiche imbricatis brnnneis tectus, radicans, apice paucifoliatus. *Folia* 1<sup>1</sup>-2 poll, longa, supra profunde canaliculata, saturate viridia. *Bcapi* foliis paullo breviores, bracteas fa poll, longte. *Flores* % poll. diam.

The plate is from a drawing made in the Peradenya Botanical Garden by W. de Alwis, kindly communicated by Dr. Trimen.

Fig. 1. Front view of flower. 2. Column and lip. 3. Lip, laid open. 4. Column.  
5. Pollinia. All enlarged.

SEE. iv. VOL. II. PAST I.

E



M.S del. et lith.

*Sarcochilus notabilis*. Hkf.

PLATE 2120.

SARCOCHILTS NOTABILIS, *Bool. f.*

ORCHIDEJE. Subtribe SARCANTHEJE.

*S. notabilis*, *TTooh. / Fl. Brit. Lid.* vi. 42; caule brevi, foliis subsessilibus oblongis v. elliptico-oblongis, scapis gracillimis, floribus in racium e bracteis arete distiche imbricatis compressam pectinatim dispositis gracile pedicellatis, sepalis petalisque consimilibus e basi anguste lanceolata 3-nervi in caudas filiformes productia, labello basi column $\odot$  sessili profunde saccato ore dilatato apice in caudam filiformem prod a cto.

HAB. Singapore, *Ridley*.

*Caulis* crassus, 1 poll, louses, foliosus. *Folia* 2-3 $\wedge$  poll, longa, §-l-j- lata, distiche patentia, ob!usa v. apice 2-dentata, coriacea, supra cupreo-brunnea, juniora viridia marginibus cupreis, subtus pallidiora, *Scapi* axillares, solitarii v. fasciculati, brunnei, medio vagina parva instructi; racemus j - 2 poll, longns, bracteae J- $\wedge$ - poll, longee. *Flares* erecti, pedicello cum cvario £ poll, longo ; sepala et petala fere pollinaria, aurantiaca, basin versus sanguineo tincta; label!urn pallide alvum, rubro-aurantiaco raarmoratum, apice caudiforri albo. *Columna* brevissima, pallide flava; rostello 0; anthera ovata, deppressa; pollinia 2, clavata, recurva, stipiti quadrato brevi sessilia, glandula brevi obovata.

A very beautiful plant, of which I am here able to give a much fuller description thaD that in 'The Flora of British India/ which was drawn up from a single leaf and three flowering scapes joined at the base and obviously so produced. Mr. Ridley hns, since the publication of that description and the preparation of this plate, sent me a beautiful coloured drawing of the plant with analyses, which differ from my description only in the solitary scapes, the raceme being young and only half an inch long, in the sac of the lip being deeper than is represented on this plate, and in its shorter filiform tip.—J. D. H.

**Fig.** 1. Flower. 2. Column and lip. 3. Lip. 4. Anther. 5. Pollinia. *Ml* enlarged.



M.S.del ©tilth.

Aerides longicorri, Hk.f.

PLATE 2127.

B RIDES LONGICORNU, *Hoolc.f.*

ORCHIDEJE. Subtribe SARCANTHE/E.

*M. longicomu*, *Hook. J. Fl. Brit. IntL* vi. 44; foliis elongatis tere-tibibus gracilibas, pedunculo gracili paucifloro, pedicellis elongatis, bracteis minutis, Bepalis petalisque liDeari-oblongis obtusis marginibns crispatis, labelli lobis lateralibus 2-partitis segmentis elongato-subulatis intermedio obcordato, calcare corniforme sepal is duplo longiore. *Mesoclastes uniflorus*, *LindL in Wall. Cat. No. 1993; Gen. et Sp. Orchid.* 45. *Lnisia uniflora*, *BLume, Bumph.* iv. 50; *Mns. Bat.* i. 64. *Lindl. Ful. Orchid.*, *Luisia*, 3.

HAB. Nepal, *Wallich*; Naga Hills, in Upper Assam, alt. 5,500 feet, *C. B. Clarke*.

*Oaules* csespitosi, pendnli, flexnosi, laxifolii, diam. pennoB anserinse. *Folia* 6-8-pollicaria, canibus tenuiora, acuta. *Pedunculi* axillaren, pollicares, pedicelli<sup>s</sup> ^-|-pollicaribus. *Flares* ad 2, albi, 1 poll. lati. *Sepnla et petala* 5-nervia, latitudine varia; petala breviora, soepissime latiora. *Labellum* sepalis aequilonga, lorum lateraliam segment a falcato-decurva, lobo intermedio membranaceo SDquilonga; calcar incurvnm, acatum. *Capsula* 2-pollicaris, angaste fasiformis, pedicello gracili.

In the absence of flowers, this interesting plant was very naturally placed in *Luisia* (from its terete leaves and long capsule) by Blume. Mr. Clarke's flowering specimen enabled me to refer it to its proper genus, where it consorts with two other terete-leaved species, viz. & *vandarum*, Eichb. f<sub>f</sub> of the Eastern Himalaya and the Khasia Hills, and *JS. cylindricum*, LindL, of the Malabar Ghats. The side lobes of the lip, divided into two long subulate decurved segments, and the long spur at once distinguish *M. longicomu* from the others. The fact of the peduncle being two-flowered obliges me to discard Lindley's specific name.—J. D. H.

Fig. 1. Flower. 2. Column. Both enlarged.



*Renanthera angustifolia*, Hk.f.

PLATE 2128.

RENANTHERA ANGUSTIFOLIA, *Hook. f.*

ORCHIDEA: Subtribe SARCANTHEJI.


*S. angustifolia*, *Hook. f. Fl. Brit. Ind.* vi. 49; caule elongate flexuoso, foliis recurvis linearibus interne angustatis apice inaequaliter bilobis, panicula elongata laxiflora, floribus 1£ poll, diam., sepalis lanceolatis lateralibus unguiculatis, petalis angustioribus, labelli minuti lobis lateralibus acatis intermedio calcare infundibulari multo breviore recurvo, columnes apice hispidulo.

HAB. Malay Peninsula; Perak, on Gunong Batu Pateh, *Wray.*

*Gardes* crassitie penna9 anserines, striati. *Folia* 4-5 poll, longa, "j-J poll. lata. *Pedunculi* laterales, elongati, graciles, parce ramosi; bracteæ parvæ; pedicel li graciles, unacum ovario pollicares. *Sepalo* 7-nervia, acuta, dorsali sessili, lateralibus basi rotundatis, abrakte breviter unguiculatis. *Petala* sepalis roquilonga, sed angustiora. *Labellum* sepalis maltoties minns, sacculum conicum ore 3-lobo lobis lateribus 2-dentatia, intermedio lingulliformi. *Golumna* brevis, crassa, apice bidentata et annulo piloram instructa; anthera ovoidea, truncata; pollinia oblonga, majascula, integra, stipite brevi lata, glandula ovata.

Near *R. elongata*, Lindl., of the Malayan Peninsula and Islands, but the flowers are very much smaller, and the spur very small.—J. D. H.

Fig. 1. Flower. 2. Front; and 3, side view of column and lip. 4. Anther. 3. Pollinia. All enlarged.



Jf.S-deletlttli.

A. *Saccolabium perpnisillum*. Hk.f.  
 B.                 . *peiiaiigiaiuim*.Hk.f

PLATE 2129.

A.--SACCOLABIUM PERPUSILLUM, *Uook.f.*

B.—SACCOLABIUM PENANGIANUM, *Hook.f.*

ORCHIDEA; Subtribe SARCANTHE<sup>^</sup>.

A.—S. perpusillum, *Hook f. Fl. Brit. Ind. vi. 56*; parvulum, caulinis curvulis, foliis recnrvis teretibus acutis, racemis erectis sessilibus multifloris, bracteis subnlati persistentibus, floribus minutis pubescentibus, sepalis ovato-oblongis, petalis obovato-oblongis, labello oblongo integro apice incrassato acuto, calcare sepalis aquilongo inflato, columna apice longe bicalcarata.

HAB. Singapore; at Sangei Buloh, *Ridley*.

*Oaules* 3-5 poll., crassitie pennre passerinse, rigidi, vaginis rigidis striatis trnncatis foliorum tecti. *Folia* disticba v. subsecunda, 3—1 .^ poll, longa. *llacemi* oppositifolii. foliis breviores, mnltiflori, r^ichi rigida; flores ?(T) poll. diam.; pedicel!is brevissimis. *Sepala* 1-nervia, lateralia acuta. *Petala* puberula, 1-uervia. *Labellum* angustum, callo erasso postice tmncato v. cordato apice incurvo acuminato instructum; calcare incurvo, obtuso, puberulo. *Columna* brevis, cornubus erectis brevior. *Capmelm* ?j poll, longce, erectce, ellipsoidero.


A very curious diminutive species, farther remarkable for the appendage terminating the lip, and the very long horns of the column, if such they are and not rostellar arms. I failed to find any calli in the spur, but, owing to the minuteness of the parts and shrinkage through drying, the analysis of the flowers was very difficult.—  
• D. H.

A. Fig. 1. Flower. 2. Dorsal sepal. 3. Fotal. 4. Column and lip. 5. Anther-case. All greatly enlarged.

B.—S. penangianum, *Hook.f. Fl. Brit. Ltd. vi. 57*; caulinis brevibus, foliis anguste linear-lanceolatis acutis carinatis infra apicem ^nidentatis, racemo brevi erecto densifloro crasse prdunculato, floribus deflexis puberulis, sepalis lateralibus late ovatis obtusis, petalis oblongis obtusis, labelli lamina brevissima 3-fida, calcare conico submflato abrupte incurvo sepalis duplo longiore.

HAH. Penang; on tlio Sonkry River, *Curtis*; Perak, *Kuustler* (ic. in *Herb. Calcutt.*).

SER. IV. VOL. II. PART II.



K.S.deletlith.

*Saccolabium Helferi Hki*".

PLATE 2130.

SACCOLABIUM HELFERI, *HooJc.f.*

OBCHIDE<sup>^</sup>. Subtribe SARCANTHEJI.

8. Helferi, *Hook.f. Fl. Brit. Ind.* vi. 57 ; caulis brevibus radicantibus, foliis loriformibus apice angustatis obtuse 2-lobis, paniculis gracilibus pendunculatis foliis brevioribus laxe ramosis, ramis curvis multifloris, floribus parvulis, sepalis orbiculari-ovatis obtusis, petalis minoribus oblongis obtusis, labelli lobis lateralibus dentiformibus terminali orbiculari-reniforme integro, calcare sacciformi sepalis paullo longiore ore ampio.

HAB. Tenasserim; at Moulmein and the Mergui Archipelago,  
*Eelfer, Griffith, Parish.*

*Caulis* 1-3-pollicares, crassitie pennae anatinæ, vaginis brevibus striatis fcliorum tecti. *Folia* 2<sup>^</sup>-3<sup>^</sup> poll, longa, |-| poll, lata, rigiria. *Paniculæ* rami ssepius curvi; bractera minutes, persistentes, patulae; flores subsessiles, 1V poll, diam., vix pubemli. *Petala* et *sepal* a 1-nervia. *Labelli* lamina calcare brevior; lobi laterales erecti, basi connati. *Golumna* brevissima, ecornuta; anthera subdidyma, truncata; pollinia minuta, subglobosa, apice stipitis magni obovati sessilia, glandula . . . ?

I am doubtful as to the real nature of what I have described as the minute lateral lobes of the lip ; which, as represented at fig. 2, sometimes appeared as if connate into a callus, representing the callus within the spur of other species.—J. D. H.

Fig. 1. Side; and 2, front view of flower. 3. Lip and column. 4. Longitudinal section of lip. 5. Anther-case. 6. Pollinia. All unlaryed.



M. S. dalet lith.

Saccolabium flavuin, Hk.f

PLATE 2131.

SACCOLABIUM FLAVUM, Hook./.

ORCHIDÆ. Subtribe SARCANTHEH.


**S. flavum**, *Hook. f. Fl. Brit. Lid.* vi. 58; caule brevi robusto, folia approximates Joriformibos apice late inasqualiter bitidis lulus acutis sinuque acutis, vaginis brovibus, racemo gracili gracile pedunculato laxifloro, bracteis minutis, pedicellis gracilibas, floribus parvulis, sepalis oblongis, labelli lobis lateralibus calcar magnum infnndibulare xnarginantibus subquadratis erosis, terminali obsolete, columna brevi crassa.

HAB. Tenasserim ; at Moulmein, *Parish*.

*Oaulis* 6-pollicaris, crassitie pennis olorinea. *Folia* dUtoha, 4-5 poll, longa, 1 poll, lata, patula, viridia, nervis saturation bus striata, lobis terminnibus £-\$ poll, longis, cuspide interdum interposito ; vaginis £-£ poll, longis, snlcatis. *Uacemus* supra folia ortus, 3^-pollicaris, decurvus; pedunculo 1^-pollicari 2-vaginato, pedicellis utiacum ovario £ poll, longis; flores \ poll, diam., pallide flavi. *bepala* 1-nervia, dorsale fornicatum ; petala subspathulata, 1-nervia *Labellum* unacum calcare sepalis duplo longius, ore lato, inter lobos laterales truncatum, intus nudam. *Columna* latior quam longa, rostello magno ut videtur carnosò; anthera acnmirvxta; pollinia 8, per paria stipirellata, stipitellis basi in unum infra apicem stipitis gracillimi elongati flexuosi insertum connatis; glandula minuta.

The plate here given is the reproduction of a coloured drawing by the Rev. G. Parish. It resembles in many respects *Sarcanthus Parishii*, Hook. (Bot. Mag. t. 5217), in habit, in the loratebifid leaves, yellow flowers, spur, and especially the curious pollen; but in that plant the leaf-lobes are rounded, the flowers are larger, the spur shorter, there is a distinct midlobe to the lip, the column is longer and narrower, and the spur is septate below the middle, and has a large dorsal callus, thus referring it to the genus *Sarcanthus*. Were it not that Mr. Parish notes that 'there is no apparent septum in the spur,' I should have regarded these two plants as congeneric. The singular structure of the pollinia is repeated with modifications in *Saccabolium obtusifolium* and *acuminatum* (Plates 2134, 2135), but it is carried (according to Parish's drawing), as it were, further in *S. flavum*, for each pollen-mass is broken up into a pair, each pair has a short stipes, and these unite into a common one that is inserted in the very long slender flat flexuous main stipes. I need scarcely add that I regard the genus *Sarcanthus* as separated from *Saccabolium* by a purely artificial character; and that a reclassification of the species of both, to get her with those of *Cleisostoma*, is to be looked for when better materials are available.—J. D. H.

Fig. 1. Flower. 2. Column and upper portion of lip. 3. Pollinia. All enlarged. (From u, drawing by the Kev. C. Purith.)



M.S. del et lith.

Saccolatmim rostellata Uira, Hk.f.

PLATE 2132.

SACCOLABIUM ROSTBLLATUM, *Iloohf.*

ORCHIDEA. Subtribe SARCANTHEJE.


8. rostellatum, *Hook. f. Fl. Brit. 2nd. vi. 59*; caule perbrevi, foliis loriformibus planis striatis apice iDaequaliter bifidis sinn acnto, r.icerao foliis multo breviore laxifloro, bracteis floribusque minutis, sepalis late oblongis obtusis, petalis paullo minoribus, labelli lobis lateralibus calcar marginantibus brevibns latia, terminali parvo ovato incurvo obtuso, calcare sepalis duplo lougiore superne infundibulari, inferne in cornu producto.

HAB. Sikkim Himalaya; at Darjeeling, alt. 7,000 feet, *Qamble* (*in Serb. CalcutL.*)

*Gaulis* vix pollicaris, radicibas confertis elongatcis planis torfcuosis. *Folia* 4-7 poll, longa, %-\$ poll. Jata, fere recta, costata; vaginis J-^ poll, lougis. *Racemus* cum pedunculo 8-10-floro 2-uncialis, gracilis; flore3 ^ poll, longi (calcare inclusio). *Sepala* et *petala* 3-5-nervia. *Columna* parva, rostello erecto prominulo; pollinia 2-loba, majnscula, apice stipitis angostioris deorsum sensim attenaati sessilia; glandula nrinuta.

The specimens of this very distinct species were hardly sufficient for a good description.—J. D. H.

Fig. 1. Flower, *nat. size.* 2. Flower. 3. Dorsal sepal, spur, and column. 4. \*«tal. 5. Pollinia. All enlarged,



M.S. del. et lith.

*Saccobodium iniuniiflorum*, Kk.f.

PLATE 2133.

**SACCOLABIUM MINIMIFLORUM**, *Booh. f.*

*ORCKIDEM.* Subtribe SABCANTHEJS.

§. *minimiflorum*, fl' oofc/. *Fl. Brit. Ind.* vi. 59; caule brevissimo, foliis elliptico-ovatis -lanceolatisve ntrinquo acutis, racemis eloDgatis ffracillimis foliis multoties longioribus multifloris, bracteis subulatis flores minutos fere sequantibus, sepalis petalisqae oblongis obtusis, jabello calcariforme sepalis longiore, basi in saccuui obtusum (calcare) Instructuni antice in cornu breve ascendentem producto.

HAB. Perak, *Father Scortechini*.

*Caulis* £-pollicaris, basi radicibus fasciculatis copiose instructus, paucifoliatus. *Folia* 1^-2 poll, longa, patula, tenniter coriacea, piana, costa tenui. *Scapi* ad basin fere bracteis peraistentibus instrncti, nnacum racemo 2-6-pollicares; bractess patnlsB; flores ad -^ poll. lati. *Sapala* et *petala* (minora) 1-nervia. *Labelli* lobi laterales 0; terminale cornu a latere compressnm simnlans; ore majusculo intus callo globoso instructo. *Columna* mediocris, basi calcare elongata armata; anthera longe rostrata.

A very curious species, quite unlike any other known to me, probably pendulous in habit. The specimens are very indifferent, but fortunately accompanied by a sketch with rude analysis and a description by its discoverer. The latter and an examination of the few available flowers enabled me to draw up the above description. Father Scortechini describes in the flower \* a something membranous, whitish, which I cannot make out whether it is a rostellum or a stipes; and this I assume to bo the spur-like body which is seen in fig. 2 projecting over the mouth of the lip, and as to the nature of which I can offer no opinion. Probably in fresh specimens it may be of a very different form.—J. D. H.

•Fig. 1. Side view; and 2, front view of flower. 3. Lip. *Ml greatly enlarged.*



ICS.WLeilith.

*Saccolabium obtusifolium*, Hk.f.

PLATE 2134.

SACCOLABIUM OBTUSIFOLIUM, *Hook. J.*

ORGHIDEJ.: Sabtribe SARCANTHE\*.


**S. obtusifolium**, *Eook. f. FL Brit. Ind. vi.* 65; cauie brevi robusto, foliis late loratis apicq profunde bifidis lobis rotundatia, raeemo elongato multilloro, bracteis ovatis acatis roflexis, sepal is pebtlisqne ohlougis obtasis concavis, labello cymbiformi v. infundibnlari in calcar incurvum apice clavatum producto, lobis lateralibus 0, terminali parvo reniforme basi callo instructo. *Uncifera obtusifolia*, *Lindl. in Journ. Linn. Soc.* iii. 40.

HAB. Sikkim Himalaya, alt. 5,000 feet, *Clarke*; Bhotan, *Col Lister*; Khasia Hills, alt. 4,000 feet, *J. D. H. and T. Thomson*.

*Caulis* 2-4-pollicares, crassitie pence anserinoe, radicibus elongatis. *Folia* 4-6-pollicaria, J-1 poll, lata, plana, crasse coriacea; vaginro i-^ poll. *Scapi* infra folia orti, unacum racemis 4-5-pollicares, bracteaB majuscneB, persistentes; pedicellis unacum ovariis J poll, longis; flores | poll, iiam., virescentes, demum straminei. *Sepala* et *petala* 3-nervia. *Labellnm* unacum calcare sepalis subduplo longius. *Columna* brevis, rostello parvulo; anthera tumida, longe rostrata; pollinia 2. parva, infra apicem late cordatam v. bialatam stipitis magni basi angustati stipitellata, gland ula elongato-lanceolata.

The singular pollinia of this and the following species induced Lindley to refer them to a genus apart, which he designated *Uncifera*. This he assuredly would not have done had he been aware of the nature of the pollinia of *S. flavum* (Plate 2131), of *Sarcanthus Parishii*, and of probably other allied plants. It is indeed a singular fact that in the tribe *Sarcanthece* the pollen and its appendages, whilst offering a maximum amount of variety in structure, seem to be of no avail for the purpose of classification.—J. D. U.

**Fig. 1.** Side view; and 2, front view of flower. 3. Apex of lip, with midlobe and callus. 4. Anther. 5 and 6. Pollinia. *M* enlarged.



*Saccolabium acurranatuia* Hkf.

PLATE 2135.

**SACCOLABIUM ACUMINATUM, Hook.f.**

ORCHIDEJE. Sabtribe SARCANTHE<sup>^</sup>.


*S. acuminatum*, Hook. f. Fl. Brit. lad. vi. 65; foliis linearibus v. oblongo-lanceolatis oblique acuminatis, racemis foliis brevioribus, bracteis minutis, sepalis lateralibus orbiculari-ovatis, petalis obovatis, labello infundibulari in calcar incurvum producto, lobis lateralibus obscuris, terminali brevi ovoido carnoso. CTncifera acuminata, Lindl. in Journ. Linn. Soc. iii. 40.

HAB. Sikkim Himalaya, alt. 6,000-7,000 feet, Gamble; Khasia Hills common, Griffith, etc.

*Caules* quam in *8. dbtusifolio* graciliores; sed folia breviora, 2-3 poll, longa, utrinque angustata, non biloba sed uno latere ultra alterum producto et acuto, pedunculi et racemi breviores, sepala latiora et petala dissimilia.

I find little difference between the pollinia and its appendages in this and in *8. ohtusifolium*. In both, the lip varies a good deal in form, and in the length and curvature of the spur.—J. D. H.

**Fig. 1. Flower. 2. Lip. 3. Front view of column and anther with the rostellar arms spread out. 4. Side view of the same. 5. Anther. 6. Pollinia. All enlarged.**



ILS dd etlith.

Sarcanthus appendiculatus<sup>Hkf</sup>

## PLATE 2136.

### SARCANTHUS APPENDICULATTJS, *Hooch.*

#### ORCHIDEJI. Subtribe SARCANTHE-E.


*S. appendiculatus*, *Hook. f. Fl. Brit. Ivd. vi. 67*; caulis elongatis, foliis elongatis cylindraceis obtusis acutis, racemis brevibus paucifloris longe pedunculatis, sepalis oblongo-lanceolatis acutis, petalis consimilibus sed minoribus, labello infundibulari a latere compresso in calcar subcylindricum obtusum lente recurvum producto, lobis lateralibus triangularibus, terminali parvo oblongo concavo, columna gracili basi antice pubescente. *S. teretifolius*, *Eichb. f. in Tram. Linn. Soc. xxx. 136* (non *Lindl.*) iErides appendicalatum, *Wall. Oat. n. 7315. Lindl Gen. et Sp. Orchid. 242.*

HAD. Tavoy, Wallich; Tenasserim, Griffith, Parish.

*Caulis* pedalis et ultra, crassitie pen nee anatina v. anserinse, maculatus, radices validos\*emittens. *Folia* 3-5 poll, longa, crassitie pennse anatince, curva. *Scapus* foliis longior, flexuosus, vaginis paucis abbreviatis instructus; racemi 1-2-pollicares, 6-8-flori; bracteaeB minutes; flores 1 poll. diam. *Petala* et *sepala* reflexa, flavo-viridia, 3-nervia et subreticulatim venulosa. *Labellum* roseum, calcare inferne septato, callo infra columnam gracilem magno lobato, callo ad basin labelli lobi terminals tumido rugoso. *Columna* incurva; anthera brevis; pollinia magna, stipite lato trianguli-ovato.

This, which Reichenbach referred to the Chinese *S. teretifolius*, differs conspicuously from that plant in the longer leaves, much longer peduncle, in the cylindric spur at the base of the funnel-shaped lip, and in the logger slender column. The plate is a reproduction of a drawing by Mr. Parish, in which the flowers are represented as very much larger than in his dried specimen, where they are hardly a quarter of an inch broad.—J. D. H.

Fig. 1. Flower. 2. Lip laid open, showing the dorsal and ventral calli of the lip, and the vertical septum of the spur. 3. The ventral callus. All enlarged.



M.SdeLetlull

*Sarcantius insectifer*, Reicftb.f.

SARCANTHUS IBTSECTIPBE, *Beichb.f.***ORCHIDEA**. Sabtribe SAKCAXTHEJS;

**8. insectifer**, *Reichb. f.* in *Bot. Zeit.* 1857, 159; caule robusto flexuoso scandente, foliis amplexicaulibus oblongis obtuse 2-lobis, racemis breviter pedunculatis paucifloris, sepalis ovato-oblongis obtusis v. subacutis, petalis oblongis, labello late infundibulari ore annulo villoso ciucto in calcar conicnm attenuate, lobis lateralibus brevibus, terminali sepalis majore triangulari-ovato v. cordato sub apice attenuate carnoso. *Hook.f. FL Brit Ltd.* vi. 68.

HAB. Behar (Western Bengal), *f. B. H.*; Cachar, *Mann* Chittagong (c. in *Herb. Calcutt.*); Tenasserim, *Parish*

*Caule* 8 pedales et ultra, crassitie pennic anatinæ, flexuosi, radices elongatas undique proferentes. *Folia* crebra, fere imbricata, distiche patentia, 1<sup>8</sup>-2 poll, longa, rigide coriacea. *Racemi* subsessiles, foliis multoties breviores, 3v-5-flores, pedunculo rachique ci\*asso decurvo; bractero minutee, deciduflB; florea ^ poll. lati. *Sepala* et *petala* 3-nervia, flavo.-viridia, rubra striat-a. *Labellum* albo-roseum. v. purpureum, ore amplio, callo basi lobi terminalis crasso, infra columnam niagu 2-fido; calcare septate. *Columna* brevissima, laba, rostellum? braciis elongatis incurvis; anthora hemispherica; poljinia sulcata, stipite brevi quadrate latiore quam longa, sessilia, glandula niagna niembranacea.

The plate here given is taken from a portrait of a fine mass of this species, attached to the bark of a billet of wood, which belongs to the Calcutta Herbarium, and is marked as having been procured at Chittagong. It represents a much larger state of the plant than that which I found in Behar, and of which the leaves were only about an inch long; the lip is larger than in any of the specimens I have examined, and in which I have not been able to determine the nature of the two hornlike processes projecting from the column represented in fig. 2, and which I assume are the two arms of the rostellum.

Fig. 1. Side view; and 2, front view of flower. 3. Lip, cut open, showing the septum. A. Front and back view\* of anther. 5. Two views of pollinia. All enlarged.



Li.S.deLetlith.

Saxcantius Scortechinn,Bk.f.

PLATE 2138.

SARCANTHUS SCORTECHINII, *Hook. f.*

**ORCHIDÆ.** Subfcribe SARCANTHEJI.


8. **Scortechinii**, *Book. f. FL Brit. Ind. vi. 68*; caule elongato robusto, foliis elliptico-lanceolatis apiculatis basi angustatis fere petiolatis crasse coriaceis, racemis erectis folia cequantibns breviter peduDcnlatis multi-floris, rachi valida, bracteis ovario eequilongis, floribus parvis, sepalis petalisque oblongis, labello infundibulari basi in calcar conico-cylindraceum obtusum producto, lobis lateralibus dentiformibus v. obsoletis, lutermedio parvo deltoid eo erecto obtuso.

HAB. Malayan Peninsula; at Perak, *Father Scortechini*, *Wray*.

*Caulis* crassitie pennce anserinee. *Folia* 3-5 poll, longa, 1-1]- poll, lata, crasse coriacea, subtns costata, costa excorrente apiculata. *Racemi* °ppositifolii; flores ^ poll. lati. *Sepala* et *petahi* 1-nervia, sordide purpurea, marginibus coRtaque purpareis. *Labellum* sepalis longius, ore ampio, callo dorsali crasso, antico puberulo, calcare infra medium \*eptato.^ *Golumna* brevis; antbera longe rostrata; pollinia biloba, apice stipitis anguste linearis sessilia, glandula minuta.

I bare seen only fruiting specimens of this species, the flowers and analysis of which are from a drawing by Scortechini.—J. D. H.

•rig. 1. Flower. 2. Lip, laid open, showing the calli and position of the septum.  
3> 4. Column. 5. Anther-case. 6. Pollinia. *M* enlarged.



*Sarcanthus lorifolius*, Parish.

PLATE 2139.

SARCANTHUS LOBIFOLIUS, *Parish.*

**ORCHIDÆ.** Subtribe SARCAHTHEJ:.


*S. lorifolius*, *Parish* mss. in *Hook. f. Fl. Brit. Ind.* vi. 60; caule brevi crasso paucifoliato, foliis elongatis lincaribus recurvis crassis canaliculatis carinatis obtusis v. breviter 2-lobis, scapo elongato flexuoso parce ramoso, ramis spicas multifloras gerentibus, bracteis reflexis, floribus parvis, sepalis lateralibns late obovatis petalis oblongo-obovatis multo latioribus, labello in calcar crassum subincurvum sepalis longius apice rotundarum producto, lobis lateralibns subulatis, intermedio parvo ovato concavo.

HAB. Tenasserim -, at Moulmein, *Parish.*

*Qaulis* crassitie fere digit. minimi, 2-3-pollicaris. *Folia* 4-5-pollicaria, -^J poll, lata ; 'Vagina brevi. *Pedunculus* 4-pollicaris; Tacemi 3-4-pollicares, decurvi, raclii valida; flores ^ poll. lati. *Sepala* et *petala* flava, rubro irrorata. *LaheUum* album, lamella longitudinali Bemiseptatum ; callo infra columnam 2-lobo. *Golumna* brevis, crassa; anthera late ovata; pollinia globosa didjma, altero cnjusvis massss multo minore, apice stipitis angusti sessilia, glandula ovata acuminata.

The accompanying plate is from an excellent drawing by the Rev. C. Parish made at Moulmein. The nearest ally of the species is *Glossostoma bipinnatum*, Lindl., in which the side-lobes of the lip are rounded, and the spur suddenly inflated and lobulate at the base, besides wanting the septum which distinguishes *Sarcanthus* from *Glossostoma*.—3. D. H.

IV 1. Flower. 2. Lip arid column. 3. Section of lip, showing the dorsal callus & longitudinal septum. 4. Pullinia. All enlarged.



*Cleisostoma andamanicum*, Hk.f.

PLATE 2140.

CLEISOSTOMA ANDAMANICUM, *Ilovh. f.*  
\*

ORCHIDEA. Subtribe SARCANTHEAS.

»


*C. andamanicum*, *Hook.* -/ *Fl. Brit. Ind.* vi. 71; caule robusto elongato, foliis loratis 2-lobis crasse coriaceis, vaginis lrevibua, pedunculis elongatis parce rarnosis, ramis apice floriferis, bracteis miimtis, Bopolis oblongis v. obovato-oblongis obtusis, labelli lobis late rali bus truncatis intermedio brevi late cordato v. reniformi apice rotundato, calcare cylindraceo inferne subinflato sepalifl dimidio longiorc.

HAB. South Andaman Islands, *Kicrz.*

*Caulis* crassitie pennos anatinie v. anserinro, radices elongatas infra folia emittentes. *Folyi* 4-G .poll, long a, i-J poll, lata, patenti-recurva, costata, lobis rotundatis. *Fedunculi* rigidj, crassitie penneo corvinu), erecto»pateiites, vaginis bast annularibus, ramis paacid divarieatis, racemis pollicaribus densiiloris; flores ^ poll, diam., pedicellis unacum ovario -|--pollicaribus. *Sepccla* et *petala* 3-nervia, fiavo-viridia, fascia intramarginali purpnrea. *Labellum* flavum, calcaro rubro maculato, callo infra columnam ligulato retuso, infra lobnm terminalem bicruri. *Columnu* brevissima rostello magno porrecto; anthera acnmiuata v. subrostrata; polluia parva, ellipsoidea, apice stipitis anguuti sesbilia, glandula parva. :

Closely allied to the Malayan *G. lalifolium* (of 'which it may be a slender state), but not nearly so stout, and with much larger flowers.—  
J. D. H.

Fig. 1. Side view; and 2» front view of flower. 3. Lip, laid open. 4. Column.  
5. Polluia. All enlarged.



M. S. del. et lith.

Cl. eisostoraabipvi TLCtatuiu, Hk.f.

PLATE 2141.

CLEISOSTOMA BIPUNCTATUM, *Book. f.*

**ORCHIDÆ.** Subtribe SARCANTHEJE.

**C. bipunctatum**, *Hook. J. Fl. Brit. bid.* vi. 73; canle gracili scandente, t'oliis angasto linearibus, apice oblique 2-dentatis, pedunculo brevissimo paacifloro, sepalis lateralibus oblongo-ianceolatis acutis, petalis sepalis aequilongis linear-i-nblongis obtusis, labelli lobis lateralibus oblongis obtusis intermedio elongato ovato-lanceolato calcare hemispherico mnltoties longiore. *Saccolabium bipunctatam*, *Var. et Ueichb.f. in Trans. Linn. Son.* xxx. 245.

HAB. TeDasserim; Moulmeio, *Pa/risk.*

*Caulis* crassitie peniisB anati\*D8B, flexnosus. *Folia* 4-6 poll, longa, 5~§ poll, lata, crassiœcula, costa valida. *Floret* cauli intra folia fere fiesailia, bracteis minntis, ad \ poll, lati, flavi v. sabparpnrei. *Sepala* 3-nervia, dorsali oblongo obtuso. *Petala* 1-nervia. *Labelli* lobum intermedium sepalis longius, pendulnm, plan am, marginibus recurvis, 3-5-nerve, album, basi callo tumido instructum ; lateralia obtusa, erecta; squama infra column am ligulfflformis erecta, erosa. *Golnmna* brevissima; antbera depresso-hemispherica; pollinia subgloboaa, bipartita, stipiti linearis affixa, glandula mediocri.

The materials for the accompanying plate are supplied by a drawing of Mr. Parish's and very imperfect specimens. The name refers to the two spots on the flower due to the lateral lobes of the lip being purple.—J. D. H.

**Fig- 1. Side Tiew; and 2, front riew of flower. 3. Lip, column, and anther. 4. Anther-case. 5. Pollinia. All enlarged.**



MIS.deLetliih.

Cleisostoma brevipes, Hk.f.

PLATE 2142.

CLEISOSTOMA BREVIPES, *ITook.f.*

ORCHIDEÆ. Subtribe SARCANTHEJB.

»


*C. brevipes*, *Jlooh.f. Fl. Brit. Tml.* vi. 73; canle crasso elongate basi rariic'inte, foliis linear-i-iancolatis acntis rigide carnosis, spica brevi, rachi crassa, bracteis in inn (is ovatis, sepalis petal is qne consimilibns ovnto-oblongis obtosis, labelli lobis lateraiibus brevi bus falcatis acutia, tcrminali brevi hastato apice membranacco disco incrassato, calcare Raccato basi rotundato.

HAD. Sikkim Himalaya, alt. 5,000 feet, *Gamble* (ie. in *Herb. Calcutt.*); Assam (*cult.*).

*Caule*\* coeapitosi, 8^12-polHcares<sub>4</sub> craRsitie pcnnie ^nserinoB; internodiis i-J-pollicaribus, e^siccatis rngalosis. *Folia* rcniformia, 3-4-j)ollicarm, erccto-pt\tentia, plana, apicibus mncroniformibns, costa inconspicna. *Spica* nnacnm pedunculo 1-1^ poll, longa, craasitie penned anatinas, vaginis infimis annnlaribufy snprioribus sparsis ovatis acutis brnnneis; ovariis ^-pollicaribns, viridibns purpureo striatis; flores ^-§ poll. diam. *Sepala* et *petdla* aurantiaca, purpureo bivittata. *Labellum* pallide flavum, callis calcaris orem claudentibus. *Columna* brevis. *Pollinia* 2, snbquadrata, apice stipitis angusti medio dilatati sessilia, snlcata; glandula masrna, membranacea, bipartita, segmentis auricula-formibas. *CapsulcB* li poll, long®, sessiles, fusiformes, teretes, striati.

The materials for the plate here given are, a drawing in the Calcutta Herbarium, a dried fruiting specimen from Mr. Gamble, and a living plant which flowered at Kew.

Fig. 1. Column, and Hction of lip 2. Pollinia. *Loth enlarged.*



M.S. del. et lith.

*Cleisostoma Maimii*, Hk.f.

CLEISOSTOMA MANNII, *Keichh.f.*OBCHIDE<sup>^</sup>. Subtribe SARGANTHEJS.

C. Mannii, *Reichb. J. in Flora*, 1872, 273; caule brevi crasso basi radicante, foliis lorutia obtase bilobis, pedunculo decnrvo, racemo elonjrato multifloro, floribus parvis, sepalis petalisque obovatis obtusis,\* labelli lobis lateralibus truncatis obtuse angulatis, intermedio orbicuiari-ovato, caicare sepalis breviore inflato v. cylindraceo. *Honk.f. FL Brit. hid.* vi. 74; in *King, Ann. Bot. Gard. Calcutt.* t. lxxxvi. (ined.).


HAD. Assam, *Mann*; Sikkim ? (fc. in *Hort. Bot. Calcutt.*).

*Caulea* 1-2-pollicares, crassitie penned olorinsa; vagina breves, aperta. *Folia* 6-pollicaria, § poll, lata, plana, striata. *Pedunculus* unacum racemo 3-pollicaris, bracteis ovatis ovaria sabcequantibus. *Flares* parvi, > poll. lati. *Sepala otpetala* pallide aurea, basi rosca. *Labdli* parvi lobas intermedium calcare brevior, lateralia obscura; squama infracolumnari linearis, erecta, apice 2-loba; callo infra lobum intermedium label li 3-partitri. *Columna* brevissima; anthera ovata, obtusa; pollinia parva, stipite lineari, glandula mediocri.

I am quite nable to distinguish satisfactorily Reichenbach's three species, *C. undulatum*, *Wendlandorum*, and *Mannii*, and doubtfully refer the plant here figured to the latter. It is copied from the type, a solitary specimen in the Kew Herbarium. According to Reichenbach's descriptions, *C. undulatum* has undulate leaves, oblong sepals, orbicular lateral lobes, a concealed obscurely 3-lobed raidlobe of the lip, and a globose spur; *G. Wendlandorum* is a much larger plant, with (according to a drawing in *Herb. Calcutt*) hairs on the lip (but which I have not found on any specimen that I have examined). I suspect that *C. Mannii* may be a small state of it.

The resemblance in the lip, its dorsal scale and ventral forked callus, to the same organ in (*T. andamanicum* (Plate 2140) is obvious.—  
J. D. H.

Fig. 1. Flower. 2. Lip, laid open. 3. Column. 4. Anther. 5. Pollinia. All



*Cleisostoma bicuspidatum*, Hk.f.

PLATK 2144.

OLEISOSTOMA BICUSPIDATUM, *Uoob.f:*

ORCHIDRS. Subtribe SABCAHTHEJI.

C. **biouspidatum**, *Hooch f. \**. *Brit. Ind.* vi. 75; ^ \* *King, Ann. BCLChJ OaLoatt.* t. tawii.; caule brevi robusto basi **radicante**, folia loriformibus profunde 2>bis lobis acntas, racemo. !**ge pedunculato simplici v.** pa<sup>TM</sup> ramoso, bracteis parr. 8 ovat<, **sepalis late obovatis obfcnsis, petalis obloDgo-obovatw,** labelh <sup>to</sup>**lateralibus parvis trnncatiB intermedio ovato acnto incurvo, calcare sepa cylindraceo obtueo.**

HAB Sikkim Himalaya, near Darjeeling, *Aiidmon* (»» *Herb. Gal-*  
*cJt.*); K h S HiS, alt. 1000 feet, Ifa»» j TenasBerim, *Pansh* (\*c. \*» *Herb. Calcutt.*).

*Caulis* 1-2-polHcaris, crassitie peniue olorin<, *Rflprias* " " ^ V ^ cibus yalidis elongatis. *Miq* 3-7 poll, longa, ^-1 poll. 1J». « « » ! \* ourva, lobis termiaalibus • erectia v. incnrvis, sinu acnto T. ?»pni8 rotundato apicolo intetpo^to, wgmis i-polhcanbus. *Bamm Mm* breviores, multi- v. panciflori; flores pam v. inajuscui!, patent<sup>es</sup> v. reflezi. *SepaZa et »i/«/« flavo-viridia, fascia media rosea.* ^<sup>TM</sup><sub>TM</sub> album, roseo maoulatum v. uroratnm; calcare recto, ore annnto nw crasBato circumdato, callo infracolnmnan 2-lpbo. *Golumm brev-*  
*Bima*, rostello elongate deeñrro; anthera tnangnlans, snbrostrato, pollbia majuscnla, globosa, Bnlcata, infra apicem stipitis angnsti Bnperne diktati inserta ; glandula minuta.

In King's » Annals of the Calcutta Botanic Gardens " I have figured and described several varieties of this plant differing much inter <>e and from that here given, -which is a copy of a drawing of <sup>a</sup> <sub>D M J</sub> <sup>X</sup> <sub>8</sub> specimen in the Calcutta Herbarium, and has much smaller flowers than the others. The acute apical teeth of the leaves are like no other species of the genus known to me.—J. D. H.

Rg. 1. Plower. 2. Aither. 3.- Bdliaia. *Mlinlargcd.*



M.S. del et lith.

*Podocarpus neriifolius* unciferus Hk.f.

PLATE 2145.

**PODOCHILUS UNCIFBEUS, Hook.f.**

ORCHIDÆA. Subtribe NOTYLIEJJ.


P. nnciferus, *Book. /. Fl. Brit Ind.* vi. 81; caulis gracilibus erectis, foliis oblongis obtusis, pedunculis latalibus numerosis gracilibus vaginatis foliis subaequilongis imo apice paucifloris, sepalis petalisque oblongis, men to elongato, labelli cuneato-ob lanceolati y. -spathulati appendice basali uncinato recirvo ungae gracili.

HAB. Malayan Peninsula; Perak, *Father Scortechini*; Borneo; Philippine Islands.

*Caules* fasciculati, 18\*24 poll, longi, foliosi. *Folia* disticha, plana, semiamplexicaulia, §-1£ poll, longa, ^-§ poll, lata, apice rotundata, s@pe denticulata et sufimentbranacea, 1-nervia, striata, nitida; *vaginae* T~t>, poll. longoD. *Pedunculi* erecfcō-paten̄tes, bracteis acicularibus vaginati, racemo brevissinio 1-paucifloro terminati, rachi capillari; bractese lat89. *Flores* ^ poll, longi. *Sepala* acuta v. acuminata, o-nervia, ineuto cylindraceo obtuso sepala aequantia. *Petala* late oblonga, 3-nervia. *Labellum* pedi brevi columnse articulatum, apice di lata turn trilobum, nngue gracili, lobis lateralibus rotundatis, termin- &lc triangulari acuto. *Columna* inermis; anthera ovato-lanceolata ; pollinia anguste pyriformia.

The only Indian species with flat (not equitant or laterally compressed) broad leaves and many lateral peduncles.—J. D. H.

**Fig. 1. Peduncle and mcm\***. 2. Flower. 3. Lip. I. Anther-case. 5. Pollinia.  
*All enlarged.*



M.S.deletlitK

*Podochilus kha.siaiuis.Hk.f.*

PLATE 2146.

PODOCHILUS KHASIANUS, Hook. f.

ORCHIDE^J. Sub tribe NOTYLIE^E.

**P. khasianus**, Hook. f. *Fl. Brit. Lid.* vi. 81; caulis gracilis foliosis, foliis linearibus acutis, racemo brevissimo terminali paucifloro, floribus minutis, sepalis ovato-oblongis acutis, mento tumido, petalis ovato-oblongis obtusis, labello oblongo-lanceolato basi profunde cordato, appendice obsoleta P. *P. microphyllus*, Wall. *Gat.* 7335 B; *Lindl. Qen. et Sp. Orchid.* 234 in part; in *Journ. Linn. Soc.* iii. 37. .

HAB. Silbet, Wallich; the Khasia Hills, at Amwee J.V.H. and T.T.

*Oaules fascicinati*, 4-6 poll, locgi, suberecti, crassitie peniiBB passerines, ad basin foliosi. *Folia* %—% poll, longa, erecto-patentia, rigid a, B8pe torta, vaginis ^-^ poll, longis. *Pedunculus* foliis brevior, bracteae rigidae nervosae. *Flo^es* .V poll, longi. *Sepala* 1-nervia, lateralia subfalcata, basi in men turn breve conico-hemisphericum iumentum. *Petala* breviora. *Labellum* breviter anguste unguiculatum, ungue basi column© inserto. *Golumna* brevissima, alata, demum in brae hi a 2 postice concava basi fissa; anthera 2-cuspidata; pollinia brevia, per paria calyprata.

Very different from *P. microphyllus*, Lindl. in Wall. Cat. 7335 A, with which it has been confounded, in the stouter habit, narrower leaves, very short spike, in apparently wanting the large basal appendage of the lip, and especially in the bicuspitate anther. *P. microphyllus* is a native of Burma and the Malay Peninsula and Islands.—J. I. H.

**Fig.** 1. Flower. 2. Lip. 3. Column after fission. 4. Anther. 5. Pollinia. All greatly enlarged.



M. S. del. et lith.

*Podochilus acicularis*, Hk.f.

PLATR 2147.

PODOCHILUS ACICULARIS, *Uoohf.*

OKCHIDEJ.: Subtribe NOTYLIEA.

*P. aciculäris*, *Hook.f. FL Brit. hid.* vi. 82; caulis filiformibus dense caespitosis ramosis, foliis aciculäribus, racemis terminalibus paucifloris brevissimis, floribus minatis, sepalis ovato-oblongis obtusis, mento tumido, petalis lineari-oblongis, labello late obovato v. oblongo, basi cordato et 2-cornuto.

HAB. Malay Peninsula; Perak, *Father Scortechini*, etc.; Penang, *Wray*.

*Caules* 6-8-pollicares, diffusi, subflexuosi, ad basin foliosi. *Flores* -JV poll. longi, ad apices ramorum subsessiles, albi, bracteis minimis. *tiepala* lateralia falcato-ovata, subacuta, 1-nervia, basi in mentum nunc obscarum rotundatum nnnc gibbosnm et 2-lobum tumescentia; dorsali nimore. *Petala* obtusa, 1-nervia. *Lahellum* sepalis brevius, 3-nerve, interdum fere rhombeo- rotundatum, ungue gracili, cornubus incurvis qbtusis. *Columna* magaa, demum bipartita; anthera ovato-cordata, a em um bifida, lobis divaricatis acuminatis; pollinia per paria calyptrata, paribus glandulad communi linearí affixa.

This species so closely reeembles *P. distichua* (*Hemmeria disticha*, *Benn. PL Jav. Bar.* t. 7, that I hesitated, and examined several flowers of each before I ventured to propose this.

The development of the column and pollinia in this genus requires a careful study. The difficulty in making out those structures in dried specimens is very great, but I think that the details which I have given on Plate 2147 are fairly accurate. The true nature of the calyprifrm cap of the pollinia baffled me. Can it be derived from the dorsal face of the column?—J. D. H.

Fig. 1. Portion of stem and leaf. 2. Flower-bract and portion of rachis of spike. 3. Mower, spread open. 4. Anthor-case in early, and 5. later, stnge. 6. Two of the pollen-masses with their calypriform shenth and gland. *Allyrdtily enlarged.*



M.S. del et lith.

Appendicula cordata, Hkf.

PLATE 2148.

APPENDICULA CORDATA, *Book. f.*

ORCHIDE-ZE. Subtribe NOTYLIEJ.:


A. cordata, *Hook. f. Fl. Brit. Ind. vi.* 83; foliis oblongis apice obtuse bidentatis, spicis lateralibus brevissime pedunculatis incurvis paucifloris vaginatis, bracteis latis, floribus parvulis punctatis, sepalis ovatis subacutis, mento rotundato, petalis obovatis, labello ovato-cordato basin versus membrana semilunari transversa instructo.

EAR. Malay Peninsula; Perak, *Father Scortechini*.

*Caulis* 7-pollicaris, crassitie penn<sup>©</sup> anatinee. *Folia* pollicaria, ad  $\frac{1}{2}$  in. lata, submembranacea, inter dentes apicales apicalata; vaginæ  $\frac{1}{2}$ -poll, longae. 8plc $<B$  1-poll, 3-4-flores; pedunculi  $\frac{1}{2}$  poll, longi, validi, vaginati, vaginis inferioribus cupuliformibus, superioribus bracteisque ovatis demura reflexis nervosis; flores  $\frac{1}{2}$ . poll, longi. *Sepala* petalaque 1-nervia. *Labellum* sepalis lateralibus minus, pedi brevi columbus sessile. *Golumna* inermis, apice dilatata, rostellum erecto ovato concavo apice 2-dentato; antera lanceolata.

I have seen but one specimen of this very distinct species.—J. I. H.

Fig. 1. Peduncle and spike. 2. Flower, spread open. 8. Column with rostellum and a pollen-mass. 4. Anther. All greatly enlarged.



NIS.d&L.ethth.

Appendicula Koenigii, Hk f.

PLATE 2149.

APPENDICULA KOENIGII, *Hook. f.*

ORCHIDEJE. Subtribe NOTTLEA.


A. Koenigii, *Hook.f. Fl. Brit. Ind.* vi. 84; foliis linear-i-oblongis oblongo-lanceolatisve apice obtusis minute 2-fidis, spicis lateralibus brevissimis 2-3-floris, floribus minutis, sepalis lateralibus ovatis acutis, mento globoso, petalis ellipticis, labello ovato v. oblongo basi appendice calceiforme instructo. *Epidendrum hexandrum*, *Koen.* in *Betz. Obs.* vi. 45.

HAB. India, *Herb. Bottler (in Herb. Kew.).*

*GavHea* dense cespitosi, 4-6-pollicares, gracileB. *Folia* crebra,  $\hat{\wedge}$ -f P<sup>o</sup>U<sup>\*</sup> longa,  $\hat{\wedge}$ -J poll, lata, membranacea, vaginis  $\hat{\wedge}$ -| poll, longis. *tipicce* brevissime pedunculatro, £-% poll, longs, patentes v. recurvae, pedunculo vaginato; bracte $\odot$  minutee, ovateB; flores iVnV P<sup>o</sup> $\wedge$ -.<sup>0D</sup>g $\circ$ -*Sepala* et *petala* 1-nervia. *Labellum* sepalis brevins, 6-nerve. *Columna* brevissima, inermis, rostello magno; antera cordato-lanceolata; pollinia 4.

<sup>a</sup> But for the solitary specimen here figured from Bottler's Herbarium, it is very improbable that the *Epidendrum hexandrum* of Koenig would have been brought to light for some time to come. Unfortunately no locality is given for the plant, and as Bottler's Herbarium contains Orchids from both Indian Peninsulas, it may have been sent to him from the Malayan, -where the genus is common. In its densely tufted habit and very minute flowers, it differs from all other species known to me.—J. D. H.

Fig. 1. Flower. 2. Lip and column. 3. Lip. 4. Same in section. 5. Anther.  
6. Pollinia. All ~~greatly~~ enlarged.



M.\$.d\*LetKth.

Appendicula lancifolia.Hk.f.

PLATE 2150.

APPENDICXTLA LANCIFOLIA, *Hook. f.*

ORC&IDE£. Subtribe NOTYLIEJ;

**A. lancifolia**, *Hook. f. FL Brit. Ind. vi.* 84; cauKbus elongatis, foliis lanceolatis apice obtuse bidentatis, raoemis terminalibus et lateralibus elongatis strictis simplicibus v. ramosis densi-multifloris, bracteis majusculis ovatis deflexis, sepalis lateralibus late ovatis, mento late cylindraceo, labello ovato-oblongo v. -obcuneato nngniculato basi znembrana concava instructo, apice trifido lobis lateralibus parvis rotundatis, intermedio late ovato.

HAB. Malay PeniDsnla: Perak, on G-unong Batn Pat eh, alt. 3,000-4,000 feet, *King's Collector.*

*Caules* 12-24 polj. longi, crassitie penure fere anserine. *Folia* 3-4 poll, longa, §-f poll, lata, basi semiamplexicaulia, sabmembranacea, dorso infra apicem mncronnlata; vaginis i-pollicaribus. *llacemi* 1-4 poll, longi, pedunculo brevi vaginis imbricatis lanceolatis erectis tecto, racbi stricta; bractere ^ poll, longee, concavse pedicellis longiores, coriacesB, nervosee; flores ^ poll, longi, aurei, disco rabro. *Sepala* lateralia sabacuta, mento breviora, petalaque mnito minora obovato-oblonga 2-nervia. *Labellum* sepalis lateralibus eequilongnm, ungue lata. *Golumna* brevissima, inermis, pede elongata; antbera late ovato-cordata; pollinia 4. *Cajpsula* ^-pollicaris, fusiformis, costata.

Very like the Javanese *A. cristata*, Blume, bat the limb of the lip is uot "rounded emarginate and crested on either side."—J. D. H.

Fig. 1. Tip of leaf, showing the dorsal mucro. 2. Flower. 3. The same laid open. 4. Lip. 5. Column. 6. Anther. 7. Pollinia. All enlarged.

Ft. 2151.



MS.diletltiL

Appendicula Mamfayi, Hk.f.

PLATE 2151.

**APPENDICULA MAINGAYI, *Hoolc. f.***

OuCHIDEiE. Subtribe NOTYLEiK.


**A. Maingayi**, *Hoolc! f. Fl. Brit. Ind.* vi. 85; canibus elongatis robustis, foliis linear-lanceolatis imo apice dilatatis et apicnlati y. bidenticulatis, racemis terminalibus erectis strictis multi-densifioris, bracteis lanceolatis patent-i incurvis, sepalis lateralibus late ovatis, men to rotundato v. suh cylindraceo, labello elliptico v. obovato-oblongo obtuso integro basi membrana lata fornicata instructo.

HAB. Malay Peninsula; Perak, *Father Scortechini*; Penang, *JUaingay*.

*Gaules* 6-16 poll, longi, crassitie fere pennse an serin 80. *Folia* 2-3 poll, longa, £-^ poll, lata, ereeto-patentia, basi angustata, vaginis  $h\sim l$  poll, longis. *liucemi* 1-2-pollicares, simplices v. basi ramosi, breviter pedunculati, rachi subdexuosa; bractes  $B_1 5^-$  poll, longae, basi deflexse dein incarvile, coriacese, 5-costattB; flores ^ poll, longi. *Sepala* lateralia obtusa, petalaque linear- v. obovato-oblonga, 3-nervia. *Labellum* sepalis paullo longias, breviter unguiculatum, 5-nerve. *Golumna* inermis, rostello brevi; anthera ovata, acuminata. *Capsvla* \$ poll, longa, fnsiformis.

The Perak specimens are more slender than those from Penang; the mentum is longer and more cylindric, and the petals narrower and 1-nerved. The species should be compared with the Javanese *- graminifolia*, Teysm. and Binnend. in *Tijdtchr. Nederl. Ind.* xxiv. (1862) 32. The specimen figured is from Penang.—**J. D. H.**

Fig. 1. Flower (of IVnang specimen). 2. Lip and column of the same. 3. Lip of the sumo. 4. Flower of Perak specimen. 5. Lip of the banie. All enlarged.



M.S. del. et lith.

Appendicula echinocarpa.Hk.f

PLATE 2152.

**APPENDICTTLA ECHINOCARPA**, *Hoohf.*


ORCHIDEJE. Subfcribe NOTYLIEE.

A. echinocarpa, *Booh.f. Fl. Brit. Ind.* vi. 85; caulis 4-6-pollicaribus gracilibus rigidis, internodiis & poll, longis, foliis ^-^-pollicaribus subimbricatis orbiculari-oblongis apice rotundatis retusis bidentatis dentibus incumbeBtibns sinu mncronulato, basi superiore reflexo, pedunculis brevibus terminalibus solitariis paacifloris, bracteis ovato-cordatis, capsula J-pollicari ellipsoidea tnrvida, valvis setis flexuosis obsitis.

**Bar.** Malayan Peninsula; Perak, on Gnnong Batu Pateh, alt 3,400 feet, T7ray.

. ThoDgh wanting the flowers, I have no hesitation in describing this interesting little species as new. It is the eighth species of the genus described from Perak, a locality that seems to abound in orchids.—  
J. D. H.

Figs. 1 and 2. Upper and lower surfaces of the upper half of a leaf. 4. Peduncle and capsule. All enlarged.



MJSdd.et.litX

Thelasis bifolia, Hk.f.

PLATE 2153.

THELASIS BIFOLIA, *Hook. J.*

ORCHIDEJS. Subtribe NOTTLIE<sup>^</sup>.

T. (Euthelasis) bifolia, *Hook. J. FL Brit. Ltd.* vi. 86; foliis pseudo-bulbo 2 nis eiliptico-lanceolatis, scapo eloDgato, spica laxiflora, sepalis lateralibus exalatis, labello ovato-oblongo v. elliptico obtuso.

HAB. Khasia Hills, *Lobb.*

*Pseudohulhus* subglobosus, § poll, diam., nudus. *Folia* 3£ poll, longa, sessilia, pa tula,\* acata, nervosa, basi angustata. *Scapus* unacum Rpica 2-pollicaris, 10 poll. Ions us, gracilis, flexuosus, vaginis' pancis brevibus remotis instructus. *B> act tee* minutce, ovatae, ovaria ©quantes. *l'erianthium* £ poll, diam., molliter cellulosum. *Sepala* obtusa, lateralia late ovata. *Pttala* oblonga v. oblongo-ovata, obtusa, 1-nervia. *Labdium* obscure 5-nerve. *Rostellum* elongatum, acuminatum, integrum.

I have seen but one specimen of this species, which was obtained in the Khasia Hills by T. Lobb when collecting orchids for Messrs. Veitch in 1850 or thereabouts.—J. D. H.

**Fig. 1. Lateral, and 2, front, view of flower. 3. Column. Ml greatly enlarged.**



M.S. del, et lith.

Thelasis khasiana, Hk.f

TLATK 2154.

THELASIS KHASIA tfA, Hook).

ORCHIDEJB. Subtribe NOTYMKJE.

T. (Euthelasis) khasiana, *Ilm*. f. *Fl*, *BHL Ltd.* vi. 87; foliis pseudobuibo solitariis 3-4-pollicaribus elongatis linearibus v. a basi pl̄tra medium sensim dilatatis obtusis, scapo uriacam spica laxiflora ioliam sequante gracili, sepalis laterali bus exalatis, labello elliptico-ovato subacuto.

HAB. Khasia HHJs, alt. 3,000-4,000 feet, *Lobb*, *L D. JET*, and *F. Thotnaon*.

*Pseudobulbs* J poll, diara., obinrbinatus v. deprefso-globosns, nudus. *folium* ^ poll. Jatum. *Scapus* flexuosus, vaginis paucis brevibus mstpuetns; spica 14—2 poll. loDga; bractee ininutie, kte ovato v. ovato-lanceolatoD, ovaria eeqaantes v. paallo breviores. *Perianthium* ♀ poll, latum. *Sepala* lateralia ovata v. ovato-lanceolata, obtuas. •*Petala* oblongo-lanceolata, obtusa. *Labellum* 3-uerve. *Rostellum* elo Dgum acuminatum, integrum.—J. D. H.

Fig. 1. Lateral, and 2. front, view of flowers. 3. Column and anther. 4. Anther.  
«greatly mlarged.



MS.del.etlith.

*Thelasis longifolia*, Hk.f.

PLATK 2155.

**THELASIS LONGIFOLIA, Houhf.**

Nat. Ord. ORCHIDEA; Subtribe NOTYLIIA.


T. (Euthelasis) longifolia, *Hook. J. Fl. Bnt. hid.* vi. p. 87; foliis pseudobulbo solitariis 6-9-pollicaribus a basi ultra medium sensim dilatatis apice rotundatis scapo valido paucivaginato longioribus, spica laxiflora, sepalis lateralibus exalatis, labello ovato acuminato.

HAB. Khasia Hills, alt. 3,000-4,000 feet, *J. D. H.* and *T. Thomson*.

*Pseudobulbus* depresso-globosns, J poll, diam., nudus. *Folium* f-1j pnll. latum, fere petiolatum. *Scapus* unicum spica laxiflora bipollucari 6-7 poll, longus. *Bracteae* minutse, late ovatse. *Perianthium*, ^re ^ poll, latum. *Sepala* Jaterialia ovato-oblonga, obtusa. *Petala* oblongo-lanceolata. *Labellum* 3-nerve. *Eostellum* elongatum, acuminatum, integrum.

*Thelasis* is a genus the flowers of which, being of a very soft, lax, succulent texture, are very difficult of examination in herbarium specimens. In this respect of texture the flowers resemble those of the very far removed genus *Phreatia* in *Epidendrees*. *T. longifolia* is very closely allied to *T. khasiana*, but is a much larger plant in foliago and flowers, and has a very robust scape.—*J. D. H.*

**Fig. 1. Flowers. 2. Column Both enlarged.**



M.:3.del,etlith.

*Thelasis elata*, Hk.f.

PLATE 2100.

OXYANTHEBA ELATA, *Iluok.* /.

OKCUIDEA. Subtribe NOTTIEIS.

**0. elata**, *TIDOJC.* f.; foliis 10-14 poll, longis loriformibus apice rotundatis, vaginis compressis coriaceis, scapo praeillimo, membro el on gato laxifloro, scpalis oblongo-lanceolatin, labello Hneari marginibns undulatis basi 2-auriculatis. Thelas<sup>a</sup>'s **elata**, *Hook. f. Fl. Brit. bid.* vi. 87.

HAB. Malayan Peninsula; Perak, on decayed wood, *Kintfs Collector.*

*Ithizoma* crassitie digit minimi. *Folia* pauca, ad 1 poll, lata; *vaginae* 3-4 poll, long<sup>o</sup>, <sup>^</sup> poll, lat», complicate. *Scapus* foliis longior, flexnosus, vspinis 2-4 <sup>^</sup> poll, longis instnictus. *Racemu*\* 6-pollicaris, praeilis; bracteo ovatas, acnniinat, ovario breviores. *Flures* breviter pedicellati; perianthio -<sup>^</sup> poll. lato. *Sepala* acnminata, nervis obscuris. *Petala* sepalis consimilia. *Labellum* subnnguiculatum, obtusum v. Bubacutum ; auriculis revolutis. *Rostellum* bipartitum.

In the ' Flora of British India,\* following Blume, Lindley, Be nth am, and Reichenbach, I referred Brongniart's genus *Oxyaiithera* to *Thelaals*. A re-examination of both genera for the \* *Icones* ' has convinced me that they shoald be kept apart, on the ground of the total difference in habit, *Oxyanthera* having no pseudobcilb, but sheathing equitant leaves on a short stem, with the blade articulate on its complicate vagina, which sheathes the base of the scape; in the lip not being flat, but concave and 2-auricled above the very short claw; and in the rostellum, which is very long in both genera, being entire in *Thela-si*\*, but bipartite in *Oxyanthera*.—J. D. H.

Fig. 1. Flower. 2 Column ami lip. 3. Lip. 4. Column with anther. 5. Anther.  
*M* greatly enlarged.



M.S. del. et lith.

*Thelasis decurva*, Hk.f.

PLATE 2157.

OX YANTHER A DECURVA, *Hook.f.*

OACHIDEJE. Subtribe NOTYLIEJ.:


0. *decurva*, *Hook. f.*; foliis 4-6 poll, longis linearibus v. anguste linear-i-lanceolatis apice angustatia, scapo gracili foliis breviore decnrvo, racemo brevi densifloro, sepalis ovatis ovato-lanceolatisve, labello carDoso pugioniformii, marginibus incarvis, basi 2-auriculato. Thelasis decarva, *Hook.f. Fl. Brit. hid.* vi. 87.

HAB. Malayan Peiinsalā: Singapore, at Krangi, *Bidley* ; Penang, on Government Hill, *Maiugay*.

*Acaulis. Folvi* £-§ poll, lata, apice snbacuta obtnsa v. oblique emarginata v. 2-loba lobis rotundatis ; vaginice 1 -1 ½-pollicares. *Scaptis* filiformis, vaginis 1-2 apicibus lanceolatis. *Raoemus* pollicaris; bractecB oblongoa ovario breviores. *Fjores* pedicellati, perianthio ^ poll. lato. *Sepal a et petala* 1-nervia. *Lahellum* basi concavum, membranacea, dein carnosam, marginibus undulatis v. crispatis interdum ad axin labelli inflexis. *Oolumna* brevis, rostello bipartito.

Closely allied to *Oxyanthera micrantha*, Brongn., which is figured as having a broadly ovate obtuse lip, narrowed into a broad short claw.—J. D. H.

Fig. 1. Apex of leaf. 2. Lateral, and 3. front, view of flower. 4. Lip and column. 5. Lip. 6. Transverse section of lip towards the tip. *All greatly enlarged.*



MS d^etlitk.

*Anaectochilus elation* Lindl.

**AN-ffICTOCHILUS ELATIOR, Lindl.**

ORCHIDEA; Subtribe **SPIRANTHEAE**.

**A. elatior**, *Lindl. in Journ. Linn. Soc.* i. 178; foliis breviter petiolatis, late ovato-rotundatis acutis, scapo elato, spica pauciflora, labelli nngue basi bastatim dilatato isthrai fimbriis utrinquend 6 lamina lonſrioribus, laminae alis oblongis divergentibua integerrimis. *Hook. f. Fl. Brit. Iml.* vi. 95.

HAR. Nilgbiri Hills; near Walla Ghat, *McIvor*.

*Caulis* basi decnmbens, dein ascendens, 1-2-pollicaris. *Folia* 1^-1 J poll, lata, 5-nervia, viridia, petiolo \^ poll. *Ion go Scapus* pedal is, vaginis pancis inflatis acuminatis auctus. *Spica* 3-4-flora; bractetB A poll. Ionge3, late ovato-lanceolatae, ovario pobescente breviores. *Fhre8* majusculi, fere 1 poll. lati. *Sepala* lateral ia ovato-lan ceo lata, acuta. *Petalu* dimidiato-obovata, acuta. *LubeUmn* § poll, longum, calcare conico inflato sepalis breviore. *Columnce* alse prominulae; rostellum elongatum, incurvnm.

I have seen only one specimen of this fine species, which resembles *A. Boxhurglei*, Lindl., a much smaller plant, with golden netted nerves on the leaf, much larger cuneately-obovate lobes of the lip, and the spur as long as the sepals.—J. D. H.

ltfg. 1. Flower. 2. Lip. 3. Column. *All enlarged.*



M.S. del, et lith.

*Anseotochilus Grifithianus Hk. f.*

PLATK 2159.

**AiTffICTOCHILUS GRIFFITH!!**, *Ilook.f.*

OBCHIDEA. Subtribe SPIRANTHE<sup>AE</sup>.

**A. Griffithii**, *Hoohf. Fh Brit. Iwt. vi. p. 96*; foliis petiolatis ovato-rotundatis acutis v. apiculatis, spica multiflora, labelli nngue basi dilatato trapeziformi, isthmo deutato y. crenato, laminae lobis cuneatis crenatis v. lobulatis.

HAB. Eastern Himalaya, *Griffith*; Naga hills in Upper Assam, *Train*.

*Caulis* basi dec urn bens, dein ascendens, 2-3-pollicaris, foliosus. *Folia* 1i-1j poll, diara., 7-nervia, viridia, petiolo ^\ poll longo. *ticapus* 4-6-pollicaris; *vaginas* 3, amplce, Jaxse, acuminatee, v. infima foliacea. *Spica* 10-20-flora, stricta; *bracteae* ovato-lanceolatre, ovaria pubernla aequanies v. super antes. *Flares* fere ^ poll. lati. *Sepala* lateralia ovata, acuta. *Petala* dimidiato-spathnlata, rostrata. *Labellimi* i poll, longum ; lobi terrinales ungui eequilongi, obcancati v. dolabri-formes, sinu acuto; calcare brevi incurvo saccato apice 2-fido. *Golumna* brevis, marginibas in appendicem plicatani dilatatis; rostellum breve.

There is no habitat for Griffith's specimens, which were probably collected in the Patkoi hills, which adjoin those of Naga where •Prain found his. In the latter habitat the claw of the lip presents rudiments of the firabrite which are so markedly developed in most of the species of the genus.—J. D. H.

Fig. 1. Flower from Griffith's specimen. 2. Petal. 3. Lip from Prain's specimen.  
\*• Column, Npur, aud lisise of lip. 5. Side view of column. All enlarged.



M.S. del et lith.

*Anaectochilus tetraphterus*, Hk.f.

PLATE 21C0.

ANJECTOCHILUS TETRAPTERUS, *Hook.*

ORCHIDEA; Sabtribe SPIRAKHKE.

**A. tetrapterus**, *Hook. f. Ft. Brit. hid.* vi. 96; foliis ovatis acufcis acumiuatisve breviter petiolatis, spica pauciflora, labelli ungrne basi anguste alato isthmo elongato gracili integerrimo, lamina) lubis bitidis bipartitisve.

HAB. Mannipore; at Eerung, alt. 2,000 feet, *Clarke*.

*Caulis* basi decumbens, vaginatus, dein ascendens, paucifoliatus. *Folia* 1^-2 poll, longa, 5-nervia, petioli vaginis iiiiflatis. *Bcups* 2-pollicaris, vaginis 2 magnis elongato-acuminatis; spica 1-2-pollicaris, tomentosa; bracteaj lanceolatae, ovaria eequantes v. superantes. *Flores* | poll. lati. *Sepala* late ovata, acuminata, *petalaque* oblique lanceolata apicibus pilis deciduis stellatim ornata. *Labellum* | poll, longum, ungue basi in auriculas breves producto, isthmo canaliculato arcuato; alis, rectangnlis divaricatis, intervallo apicalato; calcare perbrevi conico inflato iutus infra columnam appendiculis 2 conicis carnosus instructo. *Gulumna* brevis, marginibus superne in cornua ascendentia obtusa productis, basi velo bifido calcar occludente instructa.

From the shortness of its spur this species of *Anectochilus* approximates nearest of all to *Odontocilus*, and tends to invalidate the very artificial character by which these genera are kept apart—namely, that in the former genus the spur, even if reduced to a sac, is exposed, whereas in *Odmochilus* the sac is concealed by the bases of the lateral lips, which hence form a mentum.—J. D. H.

**Fig.** 1. Lateral, and 2. front, view of flower. 3. Column and spur of lip, the latter laid open. All enlarged.



M.S. del. et lith.

*Odontochilus macranthus*, Hkf.

PLATE 2161.

**ODONTOCrillus MACRANTHUS, HooLf.**

QR< HIDE£. Subtribe SPIRANTHEJE.


**O. (Myrmecis) macrantlius, Honk. f. FL Brit, Lid. vi. 98;** canle brevi robusto folioso 1-floro, foliis J-pollioaribus subsessilibns ovatis acutiR, labelli ungne angusto marginibas involatis integerrimis, limbo orbiculari emarginato marginibus basin versus undulato-crenatis.

HAB Malayan Peninsula, Perak, *Father Scortechini*; on **Maxwell's Hill, Wray.**

*Cauh's* basi decumbens, nodosus, dein 2-3-pollicaris, erectns, flexnosus, crassitie pennie corvinae. *Fulia* 1-nervia. *Fins* albus, 1^ poll, latus; ovariotblis longiore brevi ter pedunculato brae tea ovata vaginato. *Sepala* erecto-patentia, lateralia Hneari-oblonga, acuminata, basi gibba, 1-nervia. *Petala* dimidiato-ovata, acuminata. *Labellum* \$ poll. Ion gum ; nngais laminso carnusulee ima basi cordataB requilonga ; saccus rotundatus, intus carunculis 2 instructus; velum ad basin columnse 2-fidum. *Colunma* brevissima, roatello magno erecto 2-fido, lob is obtusis. *Awthnrn* linear-lanceolata; pollinia clavata; caudicula communi elongata, filiformi, gland ula elongata.

As observed in \* Fl. Brit. Ind.' this resembles the Javan *Myrmecis glabra*, Blume, but the narrower flowers, long anther, pollen, and rostellar arms distinguish it.—J. D. H.

**Fig> 1. Flower. 2. Column and lip with the sac laid open. Both enlarged.**



MS.del. etlitK  
Mo.ae.

Odoniochilus calcaratus, Hkf.

TLATE 2162.

ODONTOCHILUS CALCARATUS, *Itoohf.*

ORCHIDS. Scribe SPJRANTHEJS.

0. (*Myrmecis*) calcaratus, *Hook. f. Fl. Brit. Tnd. vi. 09*; caule robusto folioso puncifloro, foliis parvis subsessilibus ovatis acutis V. acuminatis, labelli ungue basi calcarato dein elongato dilatato crenato, alis quadratis rectangulatim divaricatis.


HAB. Malayan Peninsula; Perak, *Father Scortechini*; alt. 3,900 feet, *Wray*.

*Caulis* 3-5-pollicaris, flexuosus, crassitie penneB passerin©. *Folia* |-|-poll. longa, sessilia v. breviter petiolata, acata v. apice aristata. *Prdunculns* brevis, 2-3-florua; bracteae lanceolatce, ovaria oequantes. *Flutes* albi, 1 poll. lati. *Sepala* lateralia ovato-lanceolara, acuminata, basi in mentum calcariforme conicam obtusum connata. *Prtala* linearilanceolata. *Labellum* % poll, longum, nuge isthmo oequilongo sed anguBtiore, basi cimbriormi et in calcav conicam rectum v. apice luflexum productum. *Columna* brevis, basi processu elongato bifido in calcar descendentem instructa; rostellum elongatam, erectum, bitidurn, segnentis angustis acutis; antleria ovato-lanceolata; pollinia pyriformia, caudiculis liberis crassis striatis; glandula oblonga.

*Wray* describes the leaves as green dashed with crimson, veined with gold, and bordered with red, the flowers white. *Scortechini* says, leaves dark purple, sepals pale carnation, lip white.

This species invalidates Blume's genus *Myrmecis*, of which it has the habit, but the crenate lip-claw of *Odontochilus*.—J. D. H.

**Fig. 1.** Lateral, and 2. front, view of flower. 3. Lip with the spur removed. 4. Column showing the descending bifid appendage. 5. Pollinia. 6. Column and base of lip with spur. All enlarged.



MS. del. et lith.

Oclanlochilus purmlus, Hk.f.

PLATE 21G3.

ODONTOCHILUS PUMILUS, *Hook f.*

ORCHIDEJ.: Sabtribe SFIRANTHK<sup>^</sup>.

**0. (Myrmschis) pumilus, *Hook f. FL Brit. Lid. v\|.* 99;** caniibas brevibus subrobustis tiexuosis paucifloris, fuliis parvis breviter petiolatis ovatis, labelli ungue e basi saccato in ungnera elongatum crenulatain sensim angnstatn, laminae parvee alis oblongw v. quadratis sinn acato. Cheirostylis pasilla, partim, *LindL in Jo urn. Linn. Sac. i.* 188 (*non Gen. and Sp. Orchid.*).

HAB. Sikkim Himalaya, alt. 8,000-10,000 feet, J. P. IL and T. TJiomson, King.

*Caulis* 3-4-pollicaris, basi d ecu mben a, remote vaginatns, dein ascendens. *Folia* l<sub>2</sub>-f poll, longa, crassiuscula, nervis obscaris, acuta acuminate v. aristata. *Pedunculus* brevis, 2-3-tlorQs, vagina sol it aria ciliata; bractetB ovario breviores, oblongs, ciliatio. *Flores* \ poll, lati, ftlb. *Sepala* lateralia triangulari-ovata, obtusa, dorsuli a pice recur va. *Petala* obliqua, oblongo-lanceolata. *Labellum* \ poll, longnm ; saccas mtuscallis magnis instructus, lamina ungue brevior, alee ad angalnm 45° coiifluentes, crassinscaleB. *Columna* brevis; anthera magna.

As pointed out in the 'Flora of British India,'<sup>9</sup> this species so closely resembles *Cheirostylis pusill.n.*, that it is difficult to distinguish dried specimens of each without analysis, when the differences are clear enough. In *Cheirostylis pusilia* the sepals are connate to the middle, the lobes of the lip toothed, its claw- Huely hairy towards the tip, its sac with bristles on the nerves within, and the two frontal fleshy appendages are as long as the rostellar arms.--J. D. H.

Fig. 1. FLOWR and bract. 2. Dorsal sepal. 3. Petal. 4. Column and lip with the HHC laid open. All enlarged.



M.S. del. et lith.

*Odontochilus crispus*, Hk f.

PLATE 2164.

ODONTOCHILUS CBJSPTJS, *Hook. f.*

**ORCHIDÆ.** Subtribe SPIRANTHE.«.

0. (*Euodontoollilus*) *crispus*, *Hook. f. Fl. Brit. Intl.* vi. 99; caule ascidente paucifoliato paucifloro, foliis sessilibus late ovatis marginibus undulato-crenatis, ovario glaberrimn, bracteis sepalisque ciliolatis, labelli ungue basi saccato dein elongato angusto iutegerrimo v. obscure paucicrenato, lamina) alis divaricatis sinu acuto acinaciforuiibus margine exteriore dentatis.

HAB. Sikkim Himalaya, alt. 5,000-6,000 feot (*/c. CatJtcart*), *Clarke*; Khasia hills, at Surureem, alt. 6,000 feet, *Clarke*.

*Gaulis* basi decumbent!, nudus, dein ascendens, unacum pedunculo 6-8-pollicaris, crassitie penned corvinoe. *Folia* 2-4, f-1 poll, longa, crassiuscnla, 3-nervia, petiolo ^-1 poll. Ion go. *Peduuculus* basi vagi n is lanceolatis acuminatis erectis iustuctus, 4-8-florus; bracteaB ovario breviorcs ; ovaria ^-^ poll, longa. *Sepala* late ovata, patenti-recurva, viridia albo-margiuata, mentum saccatutn. *Petala* alba, dimidiatim ovata, in rostrum angnsta; uervo marginaii. *Lab ell am* album, ipoll. longum, s palis multo longius; lamina f poll, lata; saccus intus lamella inter calloa crassos recurves crenulatos posita instructus. *Coliimna* brevissima, rostello erecto 2-fido se^mentis acutis; pollinia globosa in apice truncato c:iudicul(s amplea triangularis coriaceee scssilia, glandula parva globosa.

The pollinia of this genus present most singular and diverse forms, which will be b»st understood by a reference to the analyses on this and Plates 2162 and 2165 to 21\*68 inclusive.—J. D. H.

Fig. I. Flower and bract. 2. Dorsal sepal. 3. Petal. 4. Lip and column. 6» Interior of sue of lip. 6. Column. 7. Pollinia. All miaryed.



M.S. del. et lith.

*Odontochilus pectmatus*, Ilk.f

PLATE 2165.

ODONTOCHILUS PECTINATUS, *Hook.f.*


ORCHIDEJE. Subtribe SPIRANTHEJS.

0. (*Euofontochilus*) *p?ctinatus*, *Hook. f. Fl. Brit. Ind.* vi. 99; canle elon<sup>^</sup>ato folioso, foliis petiolatis ovatis acutis, spica 6-8-flora pubescente, labelli ungue bast integerrimo dein elongato pectinatim dentato, lamiuse luiiatce alis oblongis subfalcatis integerrimis sinu lato.

HAB. Malayan Peninsula; Perak, alt. 3,000-4,000 feet, *King's Collector*; on Maxwell's Hill, Wray.

*Caulis* crassitie penns antierIES, basi deenmbens, dein erectus, 6-9-pollicaris. *Folia* alterna, 1J-2 poll, longa, 5-nervia, crassiuscuJa, pal tide viridia; petiolns |~<sup>^</sup> poll, longus. *Pedunculus* unacum spica 3-4-pollicarip, basi vagina foliacea instructns; bractero ^ poll, longa?, lanceolatoe, villosae, ovaris glabris aBquilonas \ *flares* § poll, lati, albi. *tiepala* lateralia ovato-lanceoiata, acominata, pnbescentia. *Pelala* oblique lanceolata, apicibus attenuatis. *Lnbedum* § poll, long am; ala3 divaricate, obtnsae, sinn lato tnberculo interposito; saccus intus callis 2 magna carnosis 2-lobis instructus. *Qolwm. ua* brevissima, auriculis 2 parvis acntis antice marginata; rostellum elongatnm, 2-partitum, lobis angastis aenminatis; anthera elongato-laoceolata; pollinia obovoidea, apice caudioala crassa duplo longiore sessilia, glandula parva oblonga.—J. D. H.

1? 1. Flower. 2. Lip with the sac laid open, showing the calli. 3. Column.  
4. Pollinia. All enlarged.



M.S. del. et lith.

*OdontochiJus brevistylis*, Hk f.

PLATE 21G6.

**ODONTOCHILUS BBBVISTYLIS**, *Huo*k.f.

ORCHIDE.FI. Subtribe SPIRANTHEJE.

**0. (Euodontochilus) brevistylis**, *Hoo*h.f. FL Brit. hid. vi. 100; oaule elongato apicem versus pauci folia to, foliis petiolata ovatis acuminatis, spica pauciflora pubescence, labello basi saccato, ungue paucispinoso, lamia $\odot$  alis seniicircularibus integris.

HAB. Malayan Peninsula; Perak, on G-anong Bata Pat eh, alt. 3,400 feet, Wruy.

*Oaulis* e basi decumbente erectua, spithameus, crassitie pennaB corvinro, remote vaginatus. *Folia* 1^2 poll, longa, 5-nervia, pal lid e viridia; petiolus £-T poll, longas. *Pedttvculus* pollicaris, nudu.s; spica 3-4-flora; bractere lanceolatoe, ovaria glabra subrostrata subsequantes ; flores ^ poll, lafcii, albi. *Sepal*\* lateralia oblique et late ovata, pilis deciduis apice barbellata; dorsali multo minore. *Petala* dimidiato-ovata, ro&trata. *Labeilum* < poll longum; unguid lamina paullo loDgior, spinis utrinque 3-4 falcatim carvis superioribus majoribus ornatus; aloe approximates, ungue paullo breviores, sinu angusto; saccus mtns basi ^emiseptatus et callis 2 cylindraceis truncatis instructus. *Columna* brevissima; rostellum breve, 2-lobum; antbera ovata; pollinia subglobosa, apice truncato caudiculee dilatatee late triangularis coriacea sessilia, glandula parva oblonga.

**The** pollinia resemble those of *O. crispus* (t. 2164).—J. D. E.

**Fig.** 1. Fl wer. 2. Lip with the sac laid open, showing the caruncles and lamella.  
3. Uilumn. 4. Anther. 5. Pollinia. *Ail* enlarged.



M.S deUlitK.

*Odontochilus Elwesii*, C.B.CI.

PLATE 2167.

ODONTOCHILUS BLWESII, *Clarke*.

OBCHIDEJJ. Subtribe SPIRANTHEJI.

0. (Euodontochilus) Elwesii, *Clarke ex Hook.f. FL Brit. Lid.* vi. 100; caule crassiusculo paucifoliato, foliis ovatis 3-costatis breviter petiolatis, pedunculo elongato et inflorescentia hirsatis, labelli ungue dilatato pectinatum paucidentato, laminae 2-alatae alis subsemicircularibus sinu lato.

HAB. Sikkim Himalaya, alt. 4,000-6,000 feet, *Clarke, King*; Khasia hills, on ShilloDg, alt. 6,100 feet, *Clarke*; Munnejpore on Kohima, *Train*.

*Caulis* basi decumbens, crassus, articulatns, ad nodos constrictus, dein erectus, 3-4-pollicaris. *Folia* 1-^2 poll, longa, crassiuscula, Jnnde viridia, costis crassis rubris; petolas b re vis, crassus. *Pedunculus* ^-4-floruR, validns, nndns v. vaginis 1-2 tomentosis instructus; bracteas 2 poll, long®, ovaria tomentosa sequantes. *Sepala* viridia apicibus purpureis, lateralia £ poll. loDga, dorsali brevi latissima acuminata; mentum latam, 2-lobum. *Petala* dimidiato orbicularia, rostrata, alba. *Labelli* iinguis sepala vix superans, irregulariter spinuloso-dentata, alba, <sup>B</sup>puinlis purpureis; lamina majuscala, alis latis integerrimis v. obscure <sup>c</sup>\*enatis albis, sinu lato; saccus scrotiformis, 2-lobus, intus semi-septatus et callis 2 crassis deformibus instructus. *Columna* brevis, crassa, basi birostrata; rostellutn breve, 2-Hdum, segmentis subulatis; ^ntheraoblonga, rostrata; pollinia elongato-pjriformia, glandula parva linear-i-obloDgl W B I W ! D. H.

<sup>Fig. \*</sup> Flower. 2. Petal. 3. Column and claw und sac of lip. 4. Sac laid open, owing the septum and caruncles. *Mt en'arged.*



M.S. del. et lith.

*Odontochilus Clarkei*. Hk.f.

PLATE 21G8.

ODONTOCHILUS CLARKEI, *Eook.f.*

ORCHIDÆ. Subtribe SPIRANTHE<sup>^</sup>.

0. (*Euodontochilus*), *Clarkei*, *Hook. f. FL Brit. Tnd.* vi. 100; caule elongato remote vaginato paucifoliato, foliis petiolatis ellipticis acutis, pedunculo elongato spicaque (sopalis exceptis) multi flora tomentosis, men to 0, labelli unguis basi facie bi lam el la to et marginibus utrinque late auriculatis dein 2-3 spinoso, laminae alis dolabriformibus integerrimis sinu angusto.

HAB. Sikkim Himalaya; at Mongpo, alt. 3,000 feet, *Clarice*.

*Gaulis* spica inclnsa<sup>^</sup>pedalis, crassitie pennse anatince, 2-3-foliatns. *Folia* 2-2f poll. longa<sub>s</sub> subobliqua, 3-5-nervia; petiolus <sup>^</sup>1-pollicaris. *Pedunculus* evaginatus. *Spica* 5-6-pollicaris, subdensiflora; bra etc<sup>©</sup> Kneari-lanceolatce, f-\$ poll. Ion gas, ovaria villosa <sup>©</sup>q nan tea. *Flores* § poll. lati. *Sepali* rosea, glabta, lateralia obliqua late ovata, basi contracta, men turn non efficientia; dorsali rotund ata, mucronata. *Petali*, sepalo dorsali cohaerentia. *Labellum* § poll, longam, lamina profnnde 2-alata, sepalis integerrimis, ungui sequilongis; saccus parvus, sepalis vix inclnsus, intus semiseptatus et carnncnlis 2 elongatis cornuformibus incurvis ad basin columnos instrnctus. *Columna* brevis, marginibus prodncntis trnnccatis; rostellum bipartitnm, segmentis sulmlatis; antleria elongata, attenuato-acumfnata ; pollinia oblongo-globosa caadicalaa validae flexuosa<sub>s</sub> striolat<sup>©</sup> affixa, glandula minuta.

This species is intermediate between *Odontochilus* and *Anectochilus*, having the small sac lamellate within and more or less covered by the sepals of the former genus, bat the sepals contracted at the base and not forming a mentnm, and the spurless lip of the latter. It appears to me that these two genera might well be combined, perhaps together with *Cheirostylis*, which only differs in the cohesion of the sepals into a tube, and the appendages of the column ; but if so, *Zevxine* roust follow. These and all the other *Spiranthea?* of India want a very careful analysis of living specimens; so long as the position of the stigma is unknown in so many, and the true natnre of appendages of the column in all, the limitation of the genera must be uncertain. I once hoped that the very remarkable differences in the structure of the pollinia would afford available generic characters, but was obliged to abandon this hope. The analyses of the orchids figured in these 'Icones,' and those described in the pages of the 'Flora of British India' have been a work of great labour, executed first by myself at various periods between 1882 and 1892; and those of the 'Icones' have been more recently, and quite independently, revised by my accomplished artist, Miss Smith, who has portrayed the results of our analyses, together with the drawings of the plants themselves.—J. D. H.

Fig. 1. Flower. 2. Column and base of lip with the sac laid open, showing the ~~sep~~turn and caruncles. 3. Pollinia. *M* enlarged.



M'.S dd, -;t hth.

Listera tenuis, Lmcll

PLATE 2169.

LISTEBA TENUIS, *LwdL*


**ORCHIDÆI**- Subtribe SPIRANTHEJE.

**L. tennis**, *Lindl. in Journ. Linn. Soc.* i. 176; gracilis, caule teretiusculo medio foliato, foliis ovatis acatis, labello cuaeato-obcordato, columna brevi. *Huok.f. FL Brit. Lid.* vi. 104.

HAB. Sikkim Himalaya; Lachen Valley, alt. 11,500 feet, /. D. H.

*Cavils* 4-5-pollicaris, glaberrimus, infra folia univaginatus. *Folia lr\ poll*, longa, sessilia, 3-5-nervia. *Pedunculus* gracillimus, nudus. *Spica* 1-2 pollicaris, puberula, 6-10-flora; flores brevissime pedicel 1 ati, a poll, lati, virides; bracteaB ovaria aequantes v. iis breviores, ovatae, aqataa, membranacese. *Sepala* lateralia ovato-lanceolata, acuminata, aorsali *petaJaque* linear-oblunga, obtusa. *Labellum* sepalis JoDgius, ^'rotundacis, sinu acuto. *Columna* vix incurva, anthera obtasa; *cusculum* breve, concavuin, obtuaura.—J. D. H.

—  
f<sup>o</sup> 1. Front, and 2. side, views of flower. 3. Front, and 4. side, views of column. All enlarged.



M. S. del. et lith.

Listera pinetorum Lmdl.

PLATE 2170.

LISTERA PINETORUM, *Lindl.*



ORGHIDEJI. Subtribe SFIRANTHEJ:.

!• pinetorum, *Lindl.* in *Journ. Linn. Soc.* i. 175; caule robusto angulato longo supra basin foliato, foliis sessilibus cordato-orbiculariis, labello e basi angusta oblanceolato bifido sinu angusto lobis subacutis. *Hook. f. Fl. Brit. Ind.* vi. 104.

**HAB.** Sikkim Himalaya; in pine woods of the Lachen Valley, alt. 10,000-11,000 feet, J. I. H.; Chumbi Valley, King.

Caulk 4-8-pollicaris, glaber, ima basi vaginatus. Folia 1-1½ poll, jata, 5-7-nervia, nervis reticulatis. Pedunculus brevis, robustus. Spica 8-flora, pubescens; flores sessiles, decarvi, ½ poll, longi; bractea <sup>Cea</sup> <sub>erba</sub> <sup>erec</sup> <sub>ab</sub> oblongae, obtusa, ovario longiores. Sepala ovato-<sup>½</sup> longa, brunnea. Petala lioearia, obtusa. Labellum sepalis bis terete, sinu acuto v. obtuso. Columna elongata, incarva, teretiuscula; anterabrevis, obtusa; rostellum rotundatum.—J. D. H.

**Fig.** 1. Flower. 2. Column. Enlarged.



LS.deLethth.

Lislera micrantha Lin <sup>dl.</sup>

PLATE 2171.

**LISTERA MICRANTHA, Lindl**

ORCHIDÆ;E. Subtribe SPIBAKTHE<sup>^</sup>.

1. micrantha, *Lindl in Journ. Linn. Soc.* i. 176; gracilliaia, foliis subradicalibus petiolata late ovatis, labello abbreviate) trim cat o breviter 3-lobo, columna brevissima. *Hook.f. Fl. Brit. Ind.* vi. 104.

HAH. Sikkim Himalaya; Lachoong Valley, alt.; 10,000 feet, J. D. ff.

*Gaulia* brevis. *Folia* \ poll, longa, 3-nervia, petiolo brevi. *Pedunculus* 2-2<sup>^</sup>-pollicaris, medio univaginatus. *Sepica* 1-1<sup>^</sup>-pollicaris, puberula, laxiflora; flores brevisaime pedicellati, minuti; bracteæ minntaD, ovatee, acuteB, ovario breviores. *Sepala* lateralia late ovata, obtusa, decurva, dorsali oblongo. *Petala* ovato-laoceolata. *Lahellum* sepalis brevias, quadrato-cuneatum, decurvum, lobis late o vat is obtusia, sinnbus rotundatis. *Columna* latior quam longa; anthera ovoidea; rostellam breve, rotundatum.—J. D. H.

*Y& 1.* Lateral, and *2.* front, view of flower. *3.* Lip and column. *All greatly enlarged.*



M. S. del. et lit.

*Zeuxme goodyeroides*, Ldl.

PLATE 2172.

ZEUXINE GOODYEROIDES, *Lindl*

ORCHIDEA. Subtribe SPIKANTHEJ.

Z. (*Monochilns*) **goodyerooides**, *Lindl Gen. and 8p. Orchid.* 486; snbro-busta, foliis breviter petiolatis ovatis, pedunculo elongatn, spia mnltitlora glabriuscula v. pilosula, floribus subsecundis, labello brevi, hypochilo magno saccato, epiochilo parvo orbicnlari-oblongo v. latissime obcordato. *Hook. f. Fl. Brit. Ind.* vi. 107; *Monochilus galeatus*, *Lindl in Journ. Linn. Soc.* i. 187.

HAB. Eastern temperate Himalaya, from Sikkim, alt. 6,000\*8,000 feet, to Mishmi, *Griffith*; and Khasia hills at Moflong, *Griffith*; Pegu, *Kurt* {in *Herb. CalcutL*}.

*Oaulis* basi decnmbens, radicibns crassis. *Folia* 1-1<sup>1</sup> poll, longa, ~~Bessilia~~ v. breviter petiolata, 3-nervia, viridia fascia media pal-pa, obtusa acuta v. acuminata, petiolo 1 poll, lon<sup>1</sup>o. *Pedunculus* 5-6-pollicaris, vaginis pancis lanceolatis. *Spica* 2-3 pollicaris, oub-uensiflora; bractea f-<sup>1</sup>\*poll. longee, ovaria glaberrima eequantea. *Flore* 8 depress. *Sepala* rosea, lateralia patentia, oblongo-lanceolata, ~~obtusa~~ <sup>orsa</sup> hypochilo incmbente ovato. *Petala* falcata, obtusa, **alba**. *Lobulum* album, sepalis eequilongum; hypochilum eubhcni-sphericum, coriaceum, medio intus 1-costatum, et callis 2 cornnformibus <sup>curvis</sup> instructum, epichilo hypocliilo minore integerrimo v. sub-bubo membranaceo. *Columna* brevis; rostellm erectum, bipartitnm; 8-nthera acuminata; pollinia pyriformia caudicnleB gracili affixa, **glandula oblonga**.

There are two drawings of the Sikkim plant in *Herb. Oalcutt.*; one <sup>Das ros</sup> 7, the other green sepals.—J. D. H.

**Fig.** 1. <sup>Ijateral</sup> » «n*i* 2. front, view of flower. 3. Column and lip. 4. Lip amoved, showing the cotte and calli. All enlarged.



MS .deletlith.

*Zeuxine reruibrrrus*, Hkf,

PLATE 2173.

ZEUXINE RENIFORMIS, *Hook.f.*

ORCHIDE<sup>A</sup>. Subtribe SPIRAKTHEJ:.


Z. (Monochilns) reniformis, *Hook. f. Fl. Brit. Ind.* vi. 107; caule elongato gracili apicent versus paucifoliato, foliis petiolatis ovato-moceolatis acuminatis, pedunculo spicae nmltiflora pnbescenbuR, labellohasi saccas\*o in unguem brevem angustato, lamina bialata v. oblata indivisa v. dentata lateribus basi iuflexis.

HAB. Malayan Peniusula; Perak, *Fatiier Scortechini*.

*Caulis* fere pedalis, erectus, crassitie penn80 anaticae, internodiis inferiob us bipolllicaribus, nndis, snperioribus brevioribus vaginatis. *Phloq* obliqua, 3-3J poll, longa, membranacea, basi rotundata, petiolo ^'Hcij ^ poll, longo. *Pedunculus* 2-3-pollicaris, vaginis linear-lanceolatis ^'4 ^ 11>structus. *Spica* 2-pollicari6, laxiflora; bracteo lanceolate, hyalinffi, ovario pnbescente longiores. *Flores* %-% poll. lati. *Sepala* lateral ia patentia, ovata, obtusa. membranacea. *Petali* dimidiato-ata cum inata, erecta. *Labellum* sepalis aequilongum, sacco intus nuse ptato et callis 2 crassis revolutis instructo, unguis mar^inibas volutis; columna brevissima, rostelli brachia erecta, fuaformia v. coiniformia; anthera lanceolata, acuminata; pollinia pyriformia, cle ca ^ ch J C lamelliformis cuneato-quadratse basi et apice truncat C segregatim affixa, g'anck magna triangnlari, cum caudicula continua.

cT1J: Inn8t refer to the figures 4 and 5 to explain the form of the most uud USf Ca,, dicle and gland of this species, which it must be difficult to fig mi\* and from tle descriiption. A modification of it is given at fig IT v and 6 of PJ 2175 » and ^ will be seen that both are extreme dev^wpmn &f that represented at fig. 5 of PI. 2174—J. D. fl.

Fig. 1. +.ow 2< a column an a IfPf wirh the sac laid open, showing the semi-septum at the 1?''' ana eal11. 3. Column and &ac of lip. 4 and 6. Pollinia. All enlarged.



US.del.ethtk

*Zeuxinerefi & Befrih.*

PLATE 2174.

ZEUXINE REGIA, *Benth.*

ORCHIDEÆ. Subfcribe SPJRANTHEIE.

**Z. (Monochilus) regia**, *Benth. in Gen. Plant* iii. 600 ; caule elongato paucifoliato, foliis petiolatis eiliptico-lanceolatis, pedunculo elongato. Risticaque pauciflora pubescenti-pilosis, labelli sepalis longioris lamina alata alis divaricatis late cuneatis v. quadrato-orbicularibus. *Hook. f.* *z<sup>b</sup>\** &\*& *fad. vi. 108.* *Monochilus regius*, *Lindl. Gen. and Sp. Orchid.* *4* » *7. Thw. Enum. PI. Zeyl* 313.

**Hab.** Ceylon; in damp woods ascending to 4,000 feet, *Walker, Sfc.*

*Gaulis gracilis*, 4-6-pollicaris, inter nodis inferioribus 1-2-pollicaribus undis, superioribus foliaceo-vaginatis. *Folia* 1^-2^ poll, longa, acuta, *viridia fascia* mediana pallida, petiolo ^ poll, longo. *Pedunculus* 3-5-pollicaris, vaginis paucis, 4-6-florus; bractero ovaria pubescentia quantes v. iis breviores. *Sejala* ^ poll, longa, ovata, obtusa, apicibus r?9urv^s^ doriali oblongo. *Petala* sepalum dorsali coherentia, linearis Dlonja, falcata, apie rotundata. *Labellum* ^ poll, longum, basi tynibiflorie, dein in ungnem v. isthmum sacco SBquilonsum angustat<sup>Dm</sup> » ^1® forma varies, plus minusve truncata, apice sinuatoe. *Perianthium* sinu lato; saccus callis 2 parvis uncinatis instructus. *Columna* levissima, rostellum brachiis elongatis; anthera ovoidea, obtusa, stictata; pollinia obovoidea in caudiculam linearem marginibus inflexis sessilia, glandula parva ovoidea.—J. D. H.

*Fig.* 1. *sejala*; 2. *petalum*; 3. *columna* and anther. 4. Anther. 5. Pollinia.



M.S.dslatlitlx

Zeuxine nervosa, Bellth..

PLATE 2175.

ZEUXINE NERVOSA, *Benth.*


ORCHIDEJS. Subtribe SPIEANTHEJI.

**Z. (Monochilns) nervosa**, *Benth.* in *Gen. Plant.* iii. 600; canle elongato folioso, foliis sparsis petiolatis late ovatis, spica pnbescents pilosa, labelli sepalis panlo longioris lamina 2-alata, alis cuneato-ovatis angulo acuto divaricatis. *Hook. f. Fl. Brit. Ind.* vi. 108. *Monochilusnervosus*, *Wall. inlndl. Qen. and Sp. Orchid.* 487; *in Journ. Linn. Soc.* i. 187. *Etaria nervosa*, *Irindl in Wall. Oat.* 7381.

HAB. Sikkim Himalaya, alt. 2,000-4,000 feet, *J. JD. H.*; Silhet and Cachar, *WaUich, Keenan*; E. Bengal, at Oomilla, &c, *Clarke*.

*Caulis* basi decnmbens, 6-10-pollicaris, longe nndus, dein ascendens, crassiusculus. *Folia* 1-2 poll, longa, acuta, 5-7-nervia. *Pedunculus* 2-5-pollicaris, gracilis v. subrobustus, vaginis 2 lanceolatis. *Spica* J-8 poll. loDga, laxi- v. subdensi-flora; bractea ovarium pilosnm. ^quantes, lanceolate; flores viridi-pnlpurei. *Sepaia* ^-J poll, longa, lateralia ovata, obtusa, patentia. *Petala* dimidiato-ovata, acuminata. *Labellum* basi breviter subsaccatum, in ungiiem isthmLmve angnstatnm; ♀ sepala panlo minores; saccus intus semiseptatus, et callis 2 nnguiwmibus recurvis instructus. *Columna* brevissima, basi antice callis ♀ prominulis instrncta; rostellum bipartitum; pollinia pyriformia, in facie caudiculae latso laminiformis (fere *Z. retiiformis* sed pollinia versns acuta) affixa, candicula ope glandnlte elongated rostello peltatim affixa.—  
**J. D. H.**

<sup>Fig. & 1\*</sup> Lateral, and 2. front, view of flower. 3. Column and sac of lip laid open.  
<sup>4a</sup> -Rostelluni, with back of caudicle. 5 and 6. FoLlinia. All enlarged.



M.S. del et lith.

Zeuxine flava, Benth.

PLATE 2176.

ZEUXINE PLAVA, *Benth.*

ORCHIDAE: Subtribe SPIRANTHEJI.

Z. (Monochilas) flava, *Benth.* in *Gen. Plant* iii. 600; caule elongato medium versus folioso, folds ovatis acutis, spica glabra v. laxe pilosa niultiflora, floribus parvulis, labelli nngne sepalis breviore ultra saccum marginato, alis late obPongo-obovatis v. dolabriformibus integerrimis v. antice dentatis. *Hook. f. Fl. Brit. Ind.* vi. 108. *Monochilus flavus*, *Wall.* in *Lindl. Gen. & Sp. orchid.* 487; *Liudl.* in *Journ. Linn. Soc.* i. 187. *Etroria flava*, *Lindl.* in *Wall. Gut.* 7380 A. B.

HAB. Nepal, *Wallich*; Assam, *Griffith*; Naga-hills, *Prain*; Tenasserim, *Heifer*, *Griffith*. Ceylon; Central Provinces, alt. 3,000 feet, *Thwaites*. Malay Isles.

*Caulk* ascendens, gracilis v. modice robustus, paucifoliatus. *Folia*, 1-1<sup>1</sup>/<sub>2</sub> poll, longa, ssepissime dissita, sessilia v. breviter petiolata, 5-nervia, vagina paullo inflata. *Scapus* 3-6-pollicaris, pauci-vaginatus, saepius laxe patentim pilosus; spica gracilis, laxiflora; bracteae ovaria cedantes v. superantes. *Flares* ad J-poll, expansi, sessiles, ovario breviores. *Sepala* -^ poll, longa, ovata, obtusa, patentia, alba. *Petala* erecta, subdimidiato-oblonga, obtusa, alba. *Labelli* uoguis basi saccatns, sacco rubro intus calcaribus 2 interdum tortis instructo, aloo divaricata. *Oolumna* brevissima; callis anticis elongatis parallels porrectis; anthera ovato-lanceolata; pollinia subgloboso-pyriformia, ope caudiculaa brevis glandules parvs lanceolatse affixa.

Judging from the specimens from many localities, extending over a large area, this speciem is a very variable one in the form of the lobes of the lip, though in this matter allowance must be made for the mutilation, by drying, of herbarium specimens. The plate is a reproduction of a drawing in the Herbarium of the Royal Gardens, Ceylon, kindly sent by the late Dr. Thwaites; it represents the flowers as white, with a red lip.—J. 1). H.

Fig. 1. Side, and 2. front, view of flower. 3. Column and lip. 4. Lip laid open. 5. Pollinium. All vn'argcd.



MIS.del.etlifh

*Zeuxine affinis*, Ben.th

PLATE 2177.

ZEUXINE AFFIWIS, *Benth.*

*ORCRIDEM*. Sab tribe SPIRANTHEJJ.

**Z. (Monochilus) affinis**, *Benth.* in *Gen. Plant ill* 600; caule elongato, foliis ovatis v. ovato-lanceolatis acutis, spica pubescente multi flora, floribus parvulis, labelli ungue ultra saccum marginato sepalis cequi-longo, alis obovato-oblongis obovatis subcaneatisve puberulis. *Hook.* /. *FL Brit. Ind.* vi. 108. *Monochilas affinis*, *Lindl. Gen. fy Sp. Orchid.* 487 (*excl Bab. Courtallwn*). *Etfuria affinis*, *Lindl. in Wall. Gat.* 7383. *E. flava*, *Wall. Cat.* 7380 0. -<*Etheria mollis*, *Lindl. in Journ. Linn. Soc.* i. 184.

HAB. Sikkim Himalaya, in the Terai, *Gamble*; Kasia Hills, alt. 5,000 feet, *Griffith*, &c.; Larma, at Taong--dong, *Wallich*; Tenasserim, »t Moulmein, *Lobb*, *Pai-ish*. \* Penang, *Maingay*.

So closely allied to *Z. flava* (PI. 2176) as to need no detailed description. It may, indeed, prove to be a form of that plant, but the characters given to it by Lindley are constant, with the exception of the lip not being so large as it is described. The flowers are a good deal larger than in *Z. flava*, more hairy, and the two broadly pyriform pollinia have connate candidles as long as themselves. The drawing is a reproduction of one in the Herbarium of the Rojal Gardens, Calcutta, lent by Dr. King.-J. D. H.

<sup>Fig-</sup> 1. Side, add 2. front<sup>\*</sup> view of flower. 3. Lip. 4. Column. 5. Pollinia. All enlarged.



VLS ddet]ith.

Zeuxine recviata.. H. k.;"

PLATE 2178.

ZEUXINE ABBBBVIATA, *Hook.f.*

OrCHIDEAE Sabtribe SPIRINTHEJJ.

**Z.** (*Monochilus*) **abbreviata**, *Book. f. Fl. Brit. Tnd. vi.* 109; **caule** elongato paucifoliato, foliis petiolatis lanceolatis acuminatis, spica jaxiflora pubescente, bracteis ovato-lanceolatis ovariis elongatis patnlia brevioribus, labelli sacco majasculo costa intus medio 2-lamellata, et basin versus callis 2 magnis instructo, alis parvis ovato-oblongis. *MaBria abbreviata*, *Lindl. in Wall. Oat.* 7385; *Oen. if Sp. Orchid.* 491 (-<*Etheria*). *B. lanceolata*, *Reichb.f. in Trans. Linn. Soc.* xxx. 142. *Dossenia lanceolata*, *Lindl. in Journ. Linn. Soc.* iii. 186.

HAB. Nepal, *Wallich*; Ebasia Hills, alt. 4,000-5,000 feet, /. D. *Uooher and T. Thomson*.

*Berba* pedalis, stricta, erecta. *Folia* 2-2<sup>1</sup>/<sub>2</sub> poll, longa, ovata v. oyato-lanceolata, mnltinervia, in petiolum  $\frac{1}{2}$  "poll, longum angustata. *bcapus* spicaque laxiflorio et bracte $\odot$  puberul $\oplus$ ; ovariu glaberrima, fera 4 poll, longa. *Flores* ad  $\frac{1}{2}$  poll, expansi. *Sepal*\* late ovata, acuminata, "08ea. *Petala* semi-orbicularia v. lunata, rosea. *Labelli* straminei  $\wedge$ nguis ultra saccum anguste alatus. *Culumna* brevis, atrinque anguste ~~slig~~  $\wedge$  a \* \*  $\wedge$ ndulatis, proceseubus 0, rostello brevi; antbera ovata; polhnia elongata, clavata, caudiculis elongatis, corpusculis lamelliB-*formib*us; glandula magna.

**R**The plate is a reproduction of a drawing which I made in the hasxa Hills in September 1850.—J. D. H.

6. **F**•  $\frac{1}{2}$ "  $\wedge$ low  $\wedge$ - 2. Sepal. 3. Petal. 4. Lip with the sac laid open. 5. Side, and •  $\wedge$ out, view of column. *All enlarged*.



M. S. del et lith.

Zeuxine moulmeir Len.sis.HICL

PLATE 2179.

ZEUXINE MOULMEINENSIS, *Hook. f.*

ORCHIDEJE. Subtribe SPIRANT HEŁ.

Z. (Monochilus) moulmeinensis, *Hook. f. Fl. Brit. Imh* vi. 100; caule gracili, foliis longe petiolatis ovatis ellipticis v. ovato-lanceolatis, vaginis brvibus laxis sacciformibnp, scapo gracili spicaqne pauciflora glanduloso-pubescentibus, brauteis ovato-lauceolatis ovario glabro aqoantibus v. eod. brevioribus, sepalis petalLsque acuminatis, labelli ungue ultra saccum rigido immargmato, sacco intus BCCUS costam late alato et basin versns utrinque callo raagno instrucro, alis falcato-CQDcatis crenatis lacerisve. Etceria moulmeinensis, *Par. et Beichb. f. in Tram. Linn. Soc. xxx. 142.*

HAB. Tenasserim; at Monlmein, *Lohb, Parish.*

*Caulis* nnacnm spica 12-18 poll, altos; inferno vnginis cnpulic-formibas instructus. *Folia* pauca, 2-3 poll, longa, inultiuervia, petiolo praciili pollicari. *Scapi* <sup>Si</sup> ~~vaginae~~ pauco<sup>3</sup>, remotas, inferiores foliacoj, Kiuperiores acuminata3. *FUrres* ad ^ poll, expans.; ovaria fere i poll, longa. *Sepala* late ovata, acuminate, sicca quasi aristata. *Petala* dimidiato-ovata v. dolabriformalia, apicibus recur vis. *Labellum* sepalis longius, sacL'i intus sub 4-loceilati lateribus extus carunculatis, unguo alisque pubescentibus. *Columna* brevissima, antice crasse bilamellosn, I'ostello brevi; anthera ovoidea; pollinia pyriformia caudiculaa parva oblongoB affixa, corpusculis lamelliBformibus.

. A remarkable species, with the interior of the sac of the lip much as in *Z.ahbreviata* (PI. 2178) and having the same thick wing^on the front of the column, and similar polliuia. The lithograph was made from a dried specimen collected by Mr. Parish in 1868, aided by a pencil sketch by that gentleman.—J. D. H.

. Fig. 1. Dorsal, and 2. latalor, sepal. 3. Petal. 4. Column and lip. ». Lip with the bac laid open. 0. Column. 7. J'olliiim. All eniarycd.



M.S. del et lith.

Zeirsine longifolia, Hk.f.

PLATE 2180.

**ZEUXINE LONGIFOLIA, Iluok.f.**

ORCHIDEJE. Subtribe SPIRANTHEJE.

2. (*H onochilus*) *longifolia*; elata, robusta, foliis lanceolatis acuminatis, *H pica* elongata laxe multiflora, rachi puberula, bracteis ovariis Kraeilibus glabris paullo brevioribns, sepal is petalisque acuminatis, labelli nngue ultra saccum gracile rigido immarginato, sacco intus secus costam anguste alato et basiti versus utrinque callo ova to instructo, alis falcato-cuneatis apice laceris. *Fl. Brit Ind.* vi. 109. *Kliombodia longifolia*, *Lindl. in Journ. Linn. Soc.* i. 181. *Hetau'ia longifolia*, *Benth. in Journ. Linn. Soc.* 88: xviii. 146.

**HAB.** Sikkim Himalaya; in hot valleys, /. *D. Hooker.*

*Caulig* G-8-pollicaris, basi procnmbcns ramosus, crassitie pcnnro olorintD. *Folia* 6-8 poll, longa, in petiolum amplum 1-1  $\frac{1}{2}$  poll, fongum anfrustata, costa valida, nervis obscuris, vaginis brevibus. *H pica* 7-pollicaris, breviter peduncnlata; flores qaaquaversi,  $\frac{1}{2}$ - $\frac{3}{4}$  poll. jam.; bracteaa ovato-lanceolatao; ovaria subrostrata. *Sepala* 1-nervia, Biecta quasi aristata, dorsali ovato; lateralia ovato-lanceolata. *Petala* dirnidiato-ovuta v. semilunata, ionge a'uminata. *Labelli* saccus antice alte bilobus quasi 2-alatus, lobis rotundatis sacco intus sub 4-Jocellato, Un'gue inter lobos projecto breviusculo rigido. *Oolumna* magna, cava, entice truncata et bicarinata; rostellnm minutum; pollinia angusre Jyriformia, glandula parva lanceclata sessilia, corpusculis lauielli-<sup>nnib</sup>us.

The ^ ^ " g 1B ^ rom a dried specimen collected by myself. It is the only one known to me.—J. D. H.

*Fixda* ^ lower\* 2- LiP ^ with tle sac laic^ oP<sup>en</sup>\* 3- Column. 4. Pollinia. All cn*wyed.*



M.S. del et lith.

Goodyera fusca, Hk.f.

PLATE 2181.

GOODYERA FUSCA, *Lindl.*

OBCHIDE£. Subtribe SPIBANTIIEX.

0. *fusca*, *Lindl. in Wall. Oat.* ft. 7395; robusta, foliis radicalibus confertis lute ovatis, spica elongata densiflora pubescente, floribus Kcundin, bracteis flores snperantibus, sepalis ovatis, petalis acinaciformibus falcatis obtusis, labelli sacco magno apice inter scpala exserto iutus alte 5-7-costato, lamina parva decarva unpuiformi v. ovata, columna brevissima. *Fl. Brit. Ind.* vi. 112 (excl. cit. Ifook./.) ^*Etheria fu<ca*, *Lindl. Gen. <& Sp. Orch.* 491% *Juurn. Linn. Soc.* i. 184. *Cyotorchis fusca*, *Benth. in Oen. Plant.* iii. 599.

Hxn. Subnlpinc Himalaya; Nopal, WalUch; Harnng, Mnwro| Garwhal, alt. 13,000 feet, Duthie; Sikkim, alt. 12,000-14,000 feet, ./!>. Hooker.

*Radices crasse fibrosse. Folia 1-1 i-pollicaria, patenti-recurva, sub-acuta, coriacea; petiolo lato brevi. ticaJw crassitie pennre anRerino?, unacum spica 5-10-polli(»ris; bracteae canlinrc spathaceoc; flores ^ poll. diam. Sepala obtusa. LaheUi costa apexque incrassattv. Culumna lata; authera parva, stipitata, didyma.*

Referred by Bentham to *Cystorchis* (which is an eminently tropical genus of very different habit) on account of the sac of the lip projecting between the lateral sepals, but i! wants the characteristic calli in the sac of the lip of that genus, und the habit is entirely that of the temperate (*Joodyenv.* In 'Fl. Brit. Ind.' when removing *G. fusca* from /*Etherw*, in which Lindley had latterly placed it, I overlooked the fact that this author had originally referred it to *Oodoxyera*.—J. D. H.

Fig 1. Bracts and flower. 2. Petal. 3. Lip ami column. 1. Limb of lip.  
A. Lateral view of column. All nilurgid.



M S del et lith

Goodyera Prairii, Hk f.

TV iiii' 2182.

GOODYERA PRAINII, *Hook. f.*

ORCHIDEJ.: Subtribe SPIRANTHEJE.

G. Prainii, *Booh. f. Fl. Brit Iml.* vi. 112; foliis petiolatis elh'pticp.  
°vut is acutis, spiea recurva glanduloso-pubescentc, floribus secundis,  
bracteis lanceolatis flores tequantibns, labello cymbiformi obtnso intus  
costato v. lainellato, rostclli biacLiis eloogatis, anthera longe rostrata  
ucuminata.

HAB. Upper Assam, on Puliabadya in the Naga Hills, nit.  
r20U feet, *Praia*.

*Caulis* nnacnm scapo et spica 6-poilicaris, robnstsns. *Folia* non  
conferta, 1-2 poll, longa, petiolata, 5-nervia, nervulis anastomosanti-  
b~~is~~ petiolas |-h poll, longus; *vagina* brevi, laxa. *Scopus* 2-3-  
~~vaginatDH~~, vavinis inferioribus foliaceis; flores erecti, ad ^ poll. lati.  
~~Spall~~<sup>Materialia</sup> oblongo-lanceolata, dorsali ovato. *Petal* a linear-  
oblonga, obtusa. *La bell urn* explanatum, ovatum, v. fere lanceolatura,  
intus 5-costatum. *Columna* brevissima; rostclli brachiis subulatis;  
anthera orbicularis, longe rostrata.

Very closely allied to Lindley's *G. recurva* from the Khasia Hills,  
^yhioh has liuear-lanceolate leaves, a sharply deflexed spike with  
linear bractn, much smaller flowers, and a beaked lip. I have seen no  
"ther specimen than that figured in the accompanying plate, where  
t he rhizome and pseudobulbs of a species of *Liparis* are seen intermixed  
with the roots of the *Goody era*.—J. D. H.

Fig. 1. Bract and flower. 2. Doraal, and 3. lateral, sepal. 4. Petal. 5. Lip and  
column. 6. Lip laid open. 7. Column. *M* enlarged.



M. S. del et lith.

*Goodyera gracilis* Hk.f.

PLATE 2183.

GOODYERA GHACILIS, *Hook. f.*

ORCHIDEJE. Sabtribe SPIBAKTHEJB.

G. gracilis, *Honk. f. FL Brit. Ind.* vi. 112; foliis pancis petiolatis (Iliptico-ovatis, scapo gracili, spica brevi puberula, bracteis floribus hrevioribus, labello recto oblongo concavo infra apicem emarginatam brcviter rostrate, disco lamella crassa percurso et basi callia 2 pyriformibus instructo.

HAB. Perak, *Father Scortechini.*

*Oatdis* nnacum scapo et spica 2-3-bracteata 8-10polHcaris, graciliH. *Folia* snbremota, 1-1<sup>^</sup> poll, longa, obtusa r. apiculata, marginibus siccis Rnbcrcnulatis, nervis obscuris. *Spica* i-1-pollicariB, pauciflora, t<sup>^></sup>rta; flores conferti, ad i poll. lati. *Sepala* linearis-oblonga v. lanceo- lata, obtusa, 1-nervia, basi gibba. *Petala* sublinearia, obtusa. *Labelum* Repalis pan Ho brevius, crassiusculam, marginibus incurvis. *Columna* brevis; rostelli brachia eloogata ; antbera lancoolata, obtusa; pollinia angaste pjriformia, breviter pcdicllata glandulce parv© linearis affixa.

The drawing is made from a specimen collected by Father Scortechini. It was accompanied with a rude figure, in which the leaves are represented as more ovate, acute, and strongly crenate.—J. D. H.

Fig. 1. Bract and flower. 2. Lip and column. 3. Interior of lip. 4. Anther.  
↳ Pollinia. All enlarged.



M.S. del et lith.

Goodyera robusta, Hkf.

PLATE 2184.

GOODYERA ROBUSTA, *Rook.*

ORCIDKE. Subtribe SPIKANTHE\*.

G. robusti. *Book.* Fl. Brit. Ind. vi. 113; data, robtista, foliis  
pi'tiolatis ovatis ovato-lauceolatisvc acatis nervosU, spica elon^ata  
~~mult~~ int l u m gUiJalrtso-töttenUwii, bract era dares secundos ~~nk~~DK  
enlos coq multibus v. brevioribua, petalift obliqno obovuto-spatiulatus,  
labello saccato lamina parva ovata terminato, sacco iiiLus scoso.

If vs. Kbasis Hills, alt. 4,000 feet, / D. Hooker and T. Thomson.

CMu/if nuacnm Bcapo et spica IS polliearis, *oruntia* [pennie olorina].  
*Folia* 3—t ptill. longa, concoloria, in petiotum ^\ poll LoDgum  
angustata, J-nervia, reticulatira venosa. *Scap*H\$ ad 3-bractealus;  
spica 7-j3ollicaris; brnetca; i poll, lungna; flores § poll. dinm. *Sopota*  
lateralia ovatn, basi jribba, d<irii<li *angnrttara*. *Petal\** BaW&'oat\*,  
obtusa. *Labethan* aepiUit bn\*viu., lamina dSOO lamullaUi. *Columna*  
brevisj rofitelli bnujliia subutata ; until era lanccolata.

Regarded by Lindley as conspecific with his *Q. seGuruiiflora*, also a native of tho Kbusia Hills, where it was first found by Griffith, and is figured in his 'oon. Pi. Asiat.' t. 347\* f. 2, and subsequently by Myself and others. But *O. uamdiflora* is a very modi smaller fewer-  
•1 I- plant, and its leaves are mottled with white. The drawing is made  
in a unique (dried) specimen collected by Dr. Thomson and nrsstf  
in 18.>0.—J. D. H.

Viz. 1. VUtw. 2. Petal. 3. Lip and column, t. Lip Uid open. 5. Frout rfaa  
of colu min. AH fad.



M.S. del et hth.

*Goodyera vittata*, B er, 1}l.




PLATE 2185.

GOODYEHA VITTATA, *Benth.*

OKCHIDEA. Subtribe SrifuxriiEJt.

G. vittata, Pftnfi. m *JJerb. Kew.*; glaberrima, robusta, folitis ovntix aculis sutunite viridilraw norvis albiiis, spica multiflnm, bracteis iiiL-ceolatis fioit'8 tcqnantibus, putalis ol:lain'ci>l;IDs fiubfulcatis obtnsis, Ia))ello Fftccato in latntnnm rcwtriforineni acabitn anf»UBtato, SACTO intns setoso, Ifltnina medio lamellis 2 *porcana*. Georchis vittata, *LindL* in *Jottrn. Linn.* 8w. i- 184.

HIB. Sikkim Hiraalaja, alt. 6,000-9,000 feet, *J. D. Hooker.*

*Cauli\** unacnm scapoepicaqno 4-G-pollic'ans, orasflitic pomiteolorinEO.. *Folia* \|-Z poll. kmg>) Wi iwwi 001 inio\*. subtun las to rubro-purpums-cetitia. *Scapu\** bi\*evis, bracteis lanceolatis dongatil TiHpnatni; spiro 2-3-poliraris, secundifiura; llores ad § poll, longt. *S-j.-tU* 3-nervia, alba, apicibtiA roseie, latcralia ovata, acaminata, dorsal! oblon^o. lanceolata, *Prtnla* niedto 3-nervia, alba. *IgabeUum* 7-ncrve, s<li 3-lobum, album. *Cviumna* brcvis, rostelli bmedia Biilmhnu. *Anthrra*. ovato-lanceolata ; polliniornm caudiculite breves, gluividulin elongate *IBx@.*

This is the only perfectly glabrous Iodian species of *Oooiiyera* known to me. The plate if taken from a coloured drawing that I made in Sikkim in Angnst 184S—J. 1). H.

Fig. 1. Brnet nnd flower. 2. Duruil sopiil. 3. FUaL 4. Lip and column, 5. Lip Inil open. 6, Side view of column. 7. Auther. Atmtargtd,



M S. del ot toll.

Goodyera hispida, L.indl.

PLATE 216C.

GOODYEHA HISPIDA, *Liudl*

ORCHIDBJ: Subtribe SPIKANTHIE\*.

**G.** *hispida*, *Liwll.in Joyrn. Linn. Soc.i.* 183; foliis ovato-lanceo-latis acuminatis glaberrimis nervosis, spiea glauiduloso-hispida torta, bracteis floribus parvis brevioribus, scpalis ovato-oblongis obtusis, petalis oblongo-ovatis obtusis, labello obovato in laciniam brevem Babrostriformem angustato, sacco intus sparse setoso. *Hook. f. Fl. Brit. bid. vi.* 114.

HAB. Kfaasia Hills; at Charra, alt. 4,000 feet, /. D. Hooker atid T. Thornton.

*Cavlis* nnacam scapo et spica 6-7-pollicaris, gracilis. *Folia* 2-2i-pollicaria, breviter petiolata, lice viridia albo striata et reticulata; *vagina* brevis. *Scaptu* 5-bracteatns, snperne cum spica pilis rigidiaculis glandulosis instructas, bracteis inferioribas foliaceis; flores in poll, diam., albi. *Sepaia* lateralia ^ poll, longa. *Petahi* obtnsa, uninervia. *Labellum* sepalis brevins. *Columna* brevis, rostellum braciis hubulatis; antbera lanceolata; pollinia anguste pyriformia caudiculis brevibus glandulaa magnaa affixa.

*O. hispida* was originally described by Lindley from the solitary specimen here figured, collected in August 1850. I have seen no other, but have a drawing of evidently the same species in the Herbarium of the Calcutta Botanical Gardens, made in 1875, probably from a cultivated specimen brought from Sikkira.—J. D. H.

Fig. 1. Bract and flower. 2. Dorsal sepal. 3. Petal. 4. Lip and column. 5. Lip laid open. 6. Column. All enlarged.



M. S. del. et lith.

Goodyera cordata, Ben.th.

PLATE 2187.

GOODYERA COEDATA, *Benth.*

ORCHIDEJ.: Sabtribe SPIRAKTHEJE.

0. cordnta, *Benth. in Herb. Kew.*; gracilis, foliis ovato-cordatis 3-flerviis, spica pauci-laxiflora pa hern la, bracteis floribus magnis sub-^rectis brevioribns, sepalis ovato-lanceolatis, petalis dimidiato-obovatis subfalcatis, labello rhotnboideo-ovato subsaccato intas setoso. *Nichols. Met. Gard. ii. 810. Hook. f. Fl. Brit. Ind. vi. 114. Georgbis cor-data, Lindl. Gen. \* \* >> Orchid. 496 (excl. Habitat.); lieichb. f. in Gard. Chron. lb79, ii. & 20.*

HAB. Khasia Hills, alt. 3,000-4,000 feet, *J. D. Hooker* and *T. Thornton*; Naga Hills, in Upper Assam, *Prain*. PPerak, *Father Scortechini*.

*Caulis* unacum scapo et spica 6-12-pollicaris, radices pansas Bimplices elongatas crassas emit tens. *Folia* H-2. ^ poll, longa, basi rotundata v. cordata, obtusa v. acnta; petiolo gracili, £-£ poll, longo. *Srcif^ut* unibracteatus, nnacum bracteis floribusqne pallide fnsco-raber; Rpica 4-8-flora; flores remoti. 1 poll. lati. *Sepala* trinervia, apicibus obtusis. *Fetala* subacinaciformia. 3-nervia, obtnsa. *Labellm* mnlti-nervp, sepalis brevius, apice acntum v. obtnsiasculam. *Oolumna* lata; rostelli bracia lanceolata, yalde elongata; antbera longa, e basi rotunda 2-loba fere snbulata ; pollinia angusta, glandale elongates linearis gracili longitndinaliter adnata.

The plate is the reproduction of a coloured drawing made at Pom-rang, in the Khasia Hills, by myself in September 1850. There is a plant very much like it found in Perak by Father Scortechini, in which the seta} nearest the tip, in the interior of the lip, are clavellato. Lindley's habitat of Ceylon (Macrae) is most probably an error; the specimen he described was probably received from the Khasia Hills, and collected by Griffith, who sent orchids to Lindley.—J. D. H.

Fig. 1. Lateral, and 2, dorsal, nepal. 3. Petal. 4. Lip laid open. 5. Column. 6. Anther. 7. Pulliuia. All tnlurged.



M.S. del et lith

*Ooodyera bifolia*, Hi. f.

PLATE 2188.

GOODYERA BIFLORA, *Hork. f.*

ORCBIDEM. Subtribe SPIRAKTHEA.

G. Kflora, *Hook. J. Fl. Brit. Ind.* vi. 114; canle breri robusto foliuso, foliiſ breviter petiolatis ovatis, spica subsessili 2-4-flora pnbescente, bracteis lanccolatis floribos magnis paacis secundis brevioribas, sepal is lanceolatis lateralibus baei connatis, labeilo basi Baccato intns setoso in laminam angastam elongatam prodncto. *Georchis biflora*, *Lindl. in Wall. Oat. n.* 7379 ; *Gen. & Sp. Orchid.* 4J6.

HAB. Western Himalaya; in the temperate region, from Nepal (Wallich) to Simla, *Gamble*, &c.

*Caulis* ascendens, crassns, radices simpliccs crassas emittens, nnacum spica 3-4-pollicaris. *Folia* 1-2 poll, longa, crassiuscula, obtusa v. auuminata; petiolo 1 poll. longo; vagina brcvis, laxa. *Scapus brevissimus*, bracteis elongatis obtectus; spica 3-4-flora; flores fere 1 poll longi. *Sepalum* doraale paullo reenrvum, lateralia recta. *Petala* linearia, obtusa, falcata, sub 3-nervia. *Labelli* lamina angusta, niedio 3-nervis, basi eaccata triplo longior. *Cnchimua* brevis, rostellli brachia setiformia valde elongata; antbera subnlata, rostello aequilonga. *Oapsula* ellipsoidea, teres, matara fere pollicaris.

The largest flowered and frnited Indian species. The drawing was Bade from a Simla specimen (dried) collected by Mr. Gamble in October 1877.—J. D. H.

Fig. 1. Dorsal, and 2. lateral, sepals. 3. Fetal. 4. Lip and column. 5. Lip with the MC laid open. 6. Column. All enlarged.



PLATE 2189.

HETJERIA HELFERI, *Rook. f.*

ORCHIDEA. Subtribe SPIRANTHEA.

E. Helferi, *Book, f Fl. Brit. Ind. vi. 115*; gracilis, foliis petiolatia ovatis acutis, spica elongata multiflora pneraia torta, bracteis floribua parvis brevioribus, sepalis ovatis, petal is linearibus obtusis subfalcatis, labello cymbiformi acuto intus trinque basin versus setulis paucis instructo.

HAB. Tenasserim (on an island of the Mergui Archipelago), *Heifer.*

*Caulis* ascendens, infeme nndns, nnacum scapo et spica 6-10'-polH-caris. *Folia* 3-4-pollicaria, 3-nervia, in petiolum subgracilem J-^pollicarem angustata, membranacea, nervis nnmerosis, primariis\* 3; *vagina* brevifl, laxa. *Scapus* gracilis, univaginatus, et bracteis plurimis Betaceis patentibus infra spicam instrncetas; spica torta; flores i poll. lati. *Lahellum* parvum. *Oolnmna* brevis, antice breviter 2-alata; rostelli brachia brevia; anthera ovata, rostrata.

The drawing is from Heifer's specimens.—J. D. H.

Fig. 1. Flower. 2. The same spread open. 3. Lip and column. 4. Column. 6. Anther. All enlarged.



M. S. del et lith.

*Hetereria elongata* Lindl.

HETJERIA ELONGATA, *Id\*dl*

ORCHIDÆ. Subtribe SPIRANTHEÆ.

H. elongate, *Limit in Watt. Cut.* ». 738i (Eia?ria) ; *Oeit. is Sp. Orchid.* 1'ji; elatu, irracilis foliis petiolatis elliptico-ovatis subsicutis, *spica* elongata **laxiflom** laxe tomentosa, **braeteia** ovario **brevioribaa** sc'latis ovato-oblongia obtusis, pctalis obovato-oblongis obtusis, **Ubello** column® adnato cymbiformi apice roatellato bay in versus intos toolh'ter setoso. ? *Trim. Cat. CeyL PL* 91. **Hook. f. F.** Brit. **huL** vi. 116. Goodyera? elongata, **LmdL** Gen. 6 8p. *Orchid.* 4'.1. ? *Rlmmphidia elongata, Thwtrilee Enum. PL Ceyl.* 313.

**HAB.** Malayan Peninsula ? *Finlayson.* ? Ceylon, Hewabetto and Ambagamowa Districts, alt. 4,000 feet, 'I hwaites.

*Cauli\** nnacnm scapo spicnqne 18-poll! car is, in feme longe foliosus, *Foh't 2-3½* poll. longa, membranae, iiiferi<>ra obtuua ; **petiolUI** ^J poll, longus ; vagiiiffi breves, lax a<sup>1</sup>, inter iurt'S hyaliue. *Scopus* pauri-bracteatus. *Spici* 4-G-pollicaris; bracieta] poll, longa?, **lanceolate**; flores fere ^ poll. hiti. *Soptda* 3-nervia. *Petala* columnt lateribus *iuv* tis niargino interiors medio incrassata, gibbti. **LttbeUum** snperins roembranacem. *Columna* brevis, antice ad basin hibelli processibus \* erectis instmcta; rostelli brachia brevia, subulata; anthera ovato-cordata; polHnia subgloboso-pynformia, caudiculis glandulat oblonguj affixis.

The ad nation of the base of the lip to the sides of the colnmn in tins species is cnrious, as is also the vertical **insertion** of the petala on th« sides of the colnmn. Fiulayson'a habitat for the plant is uncertain ; "e botanised (**daring** Sir Stamford Raffles's **miamon** to the Court of Siam) in the Malay Peninsnla and Siara, and in N. Cocliiu China; and Wallich, by whom his specimens were distributed, rarely Ka'e a locality for them in his catalogue. The plate is made from a specimen cultivated at Kew in 18^J, **trhidh** precisely agrees with 'inlayBon's. I have seen no specimen of the **Ceylon** plant (which habitat was by oversight omitted in the 'Flora of **British** India'). \*\*uless, as is most probable, the Kew **Garden** specimen was received fiom Ceylon.-J. D. H.

*Fi?*• 1. Itnct »nd flnwer 2, **Petal.** .3. Lip and column. I. Interior of Itso of lip an d portion of column, J. **Pulliaia.** All enlarged.



M.S. del et lith.

PLATE 2191.

HET-ERIA ELATA, *Hook. f.*

OKCHIDEJK. Subtribe SPIBANTHEA.

E. elata, *Hook. f.* FL Brit Ivd. vi. 116; gracilis, elata, glaberrima, foliis longe pttioiatis ovato-ianceolatis acuminatis, scapo elongate bracteis pluriniis lanceolatis vaginato, spica eloogata multi-densiflora, floribus minutis, sepal is lateral ibus late ovatis obtusis, petal is lineariorovatis, labello minimo snbhemicoperico apice rostellato, basi utrinque intas callis 2 et apicem versus lamina transversa instructo.

HAB. Perak, Father Scortechini; on Batang Padong, alt. 4,900 feet, Wray.

*Gaulis* unacnm scapn spicaqne 18-24-pollicaris. *Folia* snbdistantia, 4-5 poll, longa, in petiolum 2-3 poll, longum gracilem angustata, niultinervia, vagina brevi laxa. *Scapus* 6-pollicaria; bracte<sup>©</sup> inferiores foliaeas, spica 3-5-pollicaris; flores qaaquaversi, ^ poll, lati glaberrimi. *Labellum* a latere compressnm, profunde saccatnm, callis ad basin intnd conicis, lamina infra-apicali crassiuscula, membrana 2-fida terminata. *Columna* brevis, subquadrata, rostelli brachia brevia; anthera parva, ovata.

I have not satisfactorily ascertained the structure of the interior of the lip and of the column in this species, which can only be made out on living specimens. The tip of the transverse infra-apical plate of the labellum appeared to me to be an orbicular entire membrane ; in Miss Smith's drawing it is fleshy and bifid, and the short rostellar arms which I described in 'Fl. Brit. Ind.' are absent in her drawing or replaced (see fig. 5) by two couical projections towards the base of the column. The drawing was made from dried specimens, aided by a sketch by Father Scortechini.—J. D. H.

Fig. 1. Flower. 2 and 3. Sections of portion of lip. 4. Lip\*md column. 5. Column.  
All inlurged\*



MSdele-tlith

*Aphyllorchis; Prainii, Hk.f*

PLATE 2192.

APHYLLORCHIS PRAINII, *Hook.f.*

·ORCHIDEA: Subtribe DIURIDES.

A. Prainii, *Hook. f. Fl. Brit. Ltd.* vi. 117; elata, robusta, canle infra medium vaginato, racemo elongato laxifloro, bracteis linear-lanceolatis ovario multo brevioribus, sepalis petalisque linear-i-oblongis obtusis, labello ovato concavo, lobis lateralibus rotundatis terminali aequilongo oblongo obtuso, ungue brevissimo alis 2 parvis erectis triangularibus instructo.

HAB. Upper Assam, on the Naga Hills, *Dr. Prain.*

*Caulis* 2-pedalis, crassitie pennis olorinsa, vaginis inferioribus ovatis tubulosis, saperioribus lanceolatis. *Racemus* pedalis; bractero ^-J poll, long®, patulee v. recurves ; ovarium elongatum unicum pedicello brevi pollicare. *Sepala* % poll, longa. *Petala* linean-oblonga, apice rotundata, 3-nervia. *Labellum* sepalis brevius.

I have seen but one specimen of this plant, which belongs to the Herbarium of the Royal Garden, Calcutta; it is closely allied to *A. montana* of Sikkim and Ceylon (which are possibly different species), differing in the winged claw of the lip.—J. I). H.

Fig. 1. Flower. 2. Lip. 3. Column. 4. Anther. All enlarged.



M.S. del et lith.

AphyUorchis? va<sup>^</sup>inata,, Hk. f

**APHYLLOBCHISP VAGINATA, Eook.f.**

OBCHIPU. Subtribe DIURIDÆ.


A? vagunta, //•-/-.. f. Ft. 7Vt. Ind. ri 117; caule robusto  
vaginis iofn medium tsbalani ten ovutis uhntsis laxo vestito, ppiea  
puoiflom, bracteisovariaftabaqnantibis, sepalihoblixio-nviLtisnhtuxis,  
jn -till is luuceolatis acaminatls, labello o?ato-1<noeol;it., (<iluiiuia i  
uscu<sup>previ</sup>tn.

HAI. Klmsia Hills, in woods at Matntoo, alt 5,000 feet, J. I.  
Hooker and J. Thomson.

*Carix 6-poHioam*, erertua; vaginie ^-pollicarw, memhraniL'ere,  
^fac ea inna a obhrnscyltieaoiala, •nperiorea angntaoras. Won<sup>s</sup>  
erecti, luriilt\* rjrl>ri. *oepala* \* poll, lonpi, 3-nerviii. *Pdaia* 1-nurvia.  
*Labellum* sepalis lequilogum, stssile, pjanwiq, eeaOotttm.

I found only one specimen of this cnrions plant, wbtoh in its dried  
Mats b dificeurl of exitmirmion. Tho plate represnts it in • Ktip.  
posc• ! i-estored condit mu tlutt is, considerably en larked in; ill it spurt\*  
»ud with tho lip much nnrrmvcr thun in the flower I examined imd  
desribed in the ' Flom of British India,' in which it is brondrr thnu  
the nepau. It differs from the previously described species of the  
Renua in the Up and <\*olnmn ; but it is not safe to draw any conclusions  
fmm a solitary (proUibly distorted) KMOnMO\* the object in fi^iiriii^'  
"•(looh is to diiwt iiic ittentMB of Indian botinists to a very eurimis  
p'ant, presnnwbly a pHrophyte.—J. D. K.

"fig. I. Flower. Enlarged.



M. S. del et lith.

*Pogonia juliana*, Wall.

## Pun 21<sup>94.</sup>

### POGONIA JULIANA, Wall

ORCHIDÆ. Sllblnbo AUTHCiUt.

P. jaliaLa, 11*all*, Cat. n. 7399; f. *Uo oYao-cordato glaberi* imo,  
w»po 1-floro, aepaij\* poqiuqao unil oblanpeulit acamm rtis,  
kiwlli lobta Utermhl'iii <ib'ungi» »ub6ur< intefri\* ». ntroqoa latent  
<iint\*tj», itiUmned io elliptic x> T. rfamnbeo-Uovoklobmnpabcacri te.  
*Lindl. Gen. & Sp. Orchid.* 414. *Hook. f. Fl. Brit. Ind.* vi. 119. *Tf*  
*Enum. Pl. Zeyl.* 430. *Trim. C*  
*zb. Fl. Ind.* iii. p. 453.


HAB. AMtn (<''. fa *Herb. Calcutt.*). Silhet, C. II. Clarke, Lore.  
Lower Bengal, Itarburgh, A. C. Ceylon, ia ibo Duoubwa District,  
Theua es.

*Taber panram, f^lobcMum. Foli* <m ad apicem caiilin !-3 uncialis  
hic illio nulice\* \*. nittentis, solitarius n brrriurr prtoUt um, 2-3 p-ill.  
lot'iPim <t Inr uia, 5-7-nerve, subtus viride et p iirjuirwt'titl. *Scttpu*\*  
2-6-pollicartK, up: this ensiformibu aca minatis instru totot; bmotes\*  
parva, ovario breviter pedicellato brevior. *Flos* 1½-2½ polL (iiniii.  
*Sepala* et petab patcntta, viridia. *Lahdli* lubi k\*U'r«dw» albi, tcr-  
mina J, albo, roseo-irrorato.

Amongst a moct rniaable »ct of <1ntvingt of Moulræin ore!iiU,  
presented bj the Rev. C. Plarihk to the 1*herbari* am of Ki-w, in a figure  
«ir • *Pogtmia* (*I. cupr*M, fur. MS ) mnch immiltinfl '/. juliana, bat  
with purplo-mcu in>p~~an~~ a mi t> tain, and a mm?) shorter terminal lobe  
of tho Jip. It repreactiUi a flower on).—J. D. EL

The drawing here given is \* copy of a coloured one in Uoxburgh's  
collection.

**J6** t. Uter.il. ant 2. doml. \*\*f\*|. i. Foul. a. Bract «od flower, with ib«  
removeU A. lip. Ati mtargtd.



M. S. del. et lith.

A. *Pogonia macroglossa*. Hk.f.  
B. *maculata*. Par. & Reichb.f.

## PUT! 2195.

A POGONIA MACROOLOS8A, //.-; f.

I). POOWIA MACULATA, /<r. ami Jtrichb. /.

ORCHIDS\*. Sabtribe AurrHDIU.

A. P. maerogloua, Jl<<k. f. Fl. Brit. /<<. vi. 120; caolp mn!fl.>n>, sepalis iim-un-lrtt seolatis m-iimti atis, labello sepalis longiore line iri- ob. •mgo obtuHo ntriinjuu medium venai am! entato r. lofan Juivris lateralibus tostntr. o.

HAH. Sikkim Hi mn lava, iu hot valley\*, J. D. Hoottr; at Tamiong, fcit, ti,500 fwt, O. B. Clark:

*Foli* «m tobprqnc ignot\*. *Scoptu* 3-6-pollicar is, flexuosus, mfn\* QHsJiQni apHthis pMicci tubalontH inHructuH. *Hmetea* purrm. *Stjala* 1-1} pull, long\*, ulb\* roMtt-irmntta. *Frttilu* linear-i-nblaart>olftta. *Labe* Hum 1f poll, lonirtim, rectum, basi gibbam. *Oolumma* gracilU.

I hat© seen only two vpecimens of thin very tar^e>flowerdd species; they preaent sli^lti diftraDOH in UM lip, a\* cluiwn in tln\* drawing, which rojtresenta a ipeotmen oollecUtl hy myt\*Af in Sikkim.—J. )'. H.


H. F. macalaifc, Par. and Reichb. f. in Trans. Linn. Soc. xxx. 143; folio orbiculnri'ConUto plabro plica to, »apo floro, llorilnu\* oamoia, svpatiK petalincae ;u-<sub>2</sub>uste ob lanceolatis, labello -opaliu MqoilongO oborato.kit<x<ilht'> 8-li bo, basi breviter ob tu>c calc&rato, lobu eUiouUtt obtusis. Book./, n Brit. bd. vi. 1-20.

HAH. Tcnasserim; near Mutiluu'in, Parish.

*Tuber* globonDin radicelliH oonieil qMM tulxM-cnlaium. *Cauli*\* robus- tuH, 3 poll, lotigu\*, rjiict's vnlidas slices enrim]ns. *Folium* w& 2 M»H. dimm , sub 13-BMTT^ plicatum, viride, plicis apice late isintipu- l i b u s, Khuihiis nmriuin atroparpom notatii. *Scapus* 5-6-pollicai is, vaginis p»,icis instroelM' bxmotaa onnatm *Sepala* I pgfafa •} poll. longa, viridia. *Ltb-lluw* lmro>Tiride, niTvia tsitturatiuriljus. OPIMIM apica tubclavata ; authera pnrpurea.

The plate is from a colonred drawing by Mr. Parish.—J. D. H.

ItAT-12195 A, P. maer-gUta. Rg. I Flower. I. Pedicel, oriry, and lip. Both rged.  
- ;VATE 2195 B, P, mandate. Fig. 3. Lip. 4. Column. 6. Anthr-r. <7 enlarged.



LiS.delethtk

*Gastrodia exilis*, Hk f

PLATE 2196.

GASTRODIA EXILIS, *Hook. J.*

OBCHIDEJE. Subtribe ARETHUSK.


**0. exilis**, *Hook. f. Fl. Brit Ind.* vi. 123; gracillima, glaberrima, caule basi tuberoso, vaginis brevibus remotis, racemo paucifloro, bracteis minutifl, floribus erectis, perianthii tubo anguste subcarapanulato, sepalis orbiculatis crispatis petalis dimidio minoribus, labelli breviter nnguiculati lamina oblongo-ovata marginibus crispatus, disco apicem versus lamellis triangularibus erectis et basi callis 2 reniformibus instructo.

HAB. Khasia Hills, at Amwee, in grassy places, alt. 3,000 feet,  
*J*> *D. Hooker* and *T. Thomson*.

*Tuber* oblongnm, ^-f poll, longam. *Caulis* 6-12-pollicaris, filiformis. *Flores* 1 poll, longi, cylindracei, albidi. *Labelli* unguis brevis, laminaa apex exserta. *Columna* apice truncata, tridentata. *Capsula* immatura fusiformis.

I have had great difficulty in analysing the flowers of this singular species. The figure of the lip (3) is as that organ appeared to Miss Smith on a careful dissection. When describing it in the \* Flora of British India<sup>9</sup> I had not observed the reniform calli at its base.—J. D. H.

**Fig. 1. Flower. 2. The same with the perianth laid open. 3. Lip. 4. Column. 6. Anther. All greatly enlarged.**



M S del et lith.

A *Orchis spathulata* hb.f.B. *Stracheyi*, Hk.f.

PLATE 2197.

- A. **ORCHIS SPATHULATA**, *Reichb. f.*  
B. **ORCHIS STBACHEYI**, *Hook. f.*

OBCHLDEF. Subtribe OPHRYDE<sup>^</sup>.

A. **O. spathulata**, *Reichb. f. mss.*; folio solitario radicaii longe petiolato elliptico, scapo nudo, spica pauciflora, bracteis ttaguis ovato-oblongis obtusis, sepalis petalisque patens tenetibus, labello late rhombeo. caneato v. elliptico integro v. obscure 3-fido, calcare recto ovario breviore fusiforme. *Hook. f. FL Brit. Ind. vi. 127.* *Gymnadenia spathulata*, *Lindl. Gen. 8f Sp. Orchid. 280.*

HAD. , Alpine Himalaya, alt. 11,000-13,000 feet, from Garwhal, *Boyle, fyc.y to Sikkim, J. D. Hooker.*

*Caulis* brevis, crassinsculns, basi decumbens radices crassas emittens, Ruperne unacum folio vaginis cylindraceis truncatis involutus. *Folium* 1-3-pollicare, obtusum, crassiusculum, in petiolum validum laminse sequilongum angustatum. *Scapus* 3-5-pollicaris, valid us, apice 2-4-11 or us ;a bractea9 foliacete ilores superantes; flores \ poll, diam., purpurei. *Sepala* ovato-oblonga, obtusa v. lateralia subacuta. *Petala* elliptico-oblonga, obtusa. *Labellum* forma varium, crenulatum, calcare plus minusve elongato. *Anthem* ellipsoidea, loculis parallelis.—J. D. H.

B. **O. Stracheyi**, *Hook. f. FL Brit. Ind. vi. 128;* folio solitario radicali petiolato elliptico v. obovato, scapo nudo paucifloro, bracteis magnis ovatooblongis obtusis, sepalis petalisque patentibus, labello sepalis longiore late cuneato ad medium trinodo lobis obtusis, calcare ovario longiore incurvo obtuso.

HAB. Western Himalaya; Garwhal, near Rogile, alt. 11,000 feet, *Strachey and Winterbottom (So. 35, Gymnadenia puberula).*

Closely allied to *O. spathulata* (Fig. A), differing in the broadly 3-lobed lip and long incurved spur. In the 'Flora of British India' I have suggested the possibility of its being a hybrid between *O. spathulata* and *O. Chisna*, Don, from its having the leaf of the first-named species and the lip and spur of the second. Unfortunately tubers of both *spathulata* and *Stracheyi* are wanting, and the appearance of the dried specimens suggests their having had subterranean rhizomes. The tubers of *O. Chisna* are oblong, entire, and the leaves linear or linear-lanceolate. A good series of specimens of these three species is wanted to clear up their relations to one another.—J D. H.

PLATE 2197 A, *O. spathulata*. Fig. 1. Lateral and 2 dorsal, sepal. 3. Lip. 4. Column and spur. 5. Front view of column. All enlarged.  
PLATE 2197 13, *O. Stracheyi*. Fig. C Sepal. 7. Petal. 8. Lip. 9. Anther. All *enlarge*.



M. S. del et lith

*Herminium fallax*, Lindl.

PLATE 2198.

HERMINIUM PALLAX, *Lindl*

ORCHIDEJE. Subtribe **OPIRYDEE.**

*H. fallax*, *TAndl. in Wall. Gat n.* 7412; gracilis, foiiis 1-2 lincarihus linear-i-lanceolačisve, spica elongata multiflora, floribus subsecundis, bracteis ovarii brevioribus, sepalis ovato- v. oblongo-lanceolatis, petalis erectis angustis falcatis, labello scepius oblongo 3-lobo, lobis obtusis basi saccato, sacco polymorpho. *Hook. f. Fl. Brit. Ind.* vi. 320. *Peristylus fallax*, *Lindl. Gen. fy Sp. Orchid.* 298. *Cybele, Falc. in Lindl. Veg. Kingd.* 193, c. *G. alpina, Falc. mss.*

HAB. Alpine and Rubalpine Himalaya, alt. 7,000-12,000 ft., from Garwhal to Sikkim, *Wallich*, &c.

*Gaulis* nnacum spica 4-18-pollicaris, medio 1-2-foliatus, vaginis snperioribns aphyllis 1-2 inferioribus subfoliaeia. *Folium* pierumqne solifarium, 2-5-pollicare, i-1 poll, latum, acutum, rarius folia sub-radicalia adsant. *Spica* 2-5-pollicaris, deusi- v. laxi flora; bract ere inferiores longiores, lanceolatee; ovarium %-£ poll, longam, sab-rostratum. *iSepala* ^-poll. longa, 1-nervia. *Petala* forma varia. *Labellum*, nnnc sepalis inulto brevius crassiuscnlum breviter lobatum, nunc angustum tenuius 3-jobnm lobis brevibus v. elongatis, limbus basi trancatns v. anguslatas; calcar incurvum, subdidymum sub^lobosnra v. clavellatum. *Avthera* parva, oblonga, loculis parallelis, tubnlis 0; pollinia obovoidea, glandalis parvis nudis nitidis sessilia v. caudicnla brevi interposita.

A very common Himalayan plant of the temperate region, and exceedingly variable in stature, fuligo, lip and sac; as in the case, more or less, with the other species of that country. At p. 128 of the 'Flora of British India,'<sup>9</sup> I have stated that there is no other character by which *Senninium* can b8 distinguished from *Habenaria* but the absence of a yell-developed spur. It was, indeed, my original intention to reduce \*t to the latter genus, which accounts for my oversight in citing myself (in 'FL Brit. Ind.') as the authority for *H. fallax*, instead of Lindley. The involved synonymy which this would have entailed, however, induced me to refrain from so doing. The drawing is made from a Sikkim specimen collected by myself at about 10,000 feeb.—  
**J. D. H.**

**Fig.** 1. Flower. 2. Lip ami column. 3 and 4. Forms of lip. 5. Column.  
**6.** *P'illinum.* All enlarged.



M. S. del et lith.

A. *Herminium Duthiei*, Hkf  
 B. *orbiculare*, Hkf

**PLATE 2199.**

A. QERMINIUM DUTIUK! *Hock. f.*  
B. II OHMCULAKK. *f. I.*

**OCHIDÆ. Sc. *binl\** *Oruttntx.***

A. H Duthjifi. // *I. f. Fl. lirit. I<Li* I 30; foliis basin ver, „  
 raii dis geminis lin. • uitraa oblongii oblaaeaolaiisra, »jrtea raaltttara,  
 6or ibus sib^caDltta, UrarteM ovario brerioribo\*, •vpalu \*b*i* sis,  
 lateralibus ovato-taooaolatu qnatn duraaW late objoago btwiorib<>,  
 petalis carnosia icntia, Ubrlllo oralo •rtjtce T. orato-cordatu \*. orbieu)  
 ovato apico a

Mi<sup>n</sup>, Western Him Jajft; Oarwkal, Mar tU Kbaai Pa\*\*\*, all.  
11,000-12,000 feet, Duthie.

*OamLU tt nvam «pio» - ft. polliotri\*. Folia 2-3}-pollicis •"• \*\*\*»\* v.  
•rnntt mla, fawn rm^ina incloa\*. S»piu r»liHu«, currm\*, no dus : spica  
2-4-pollicaris ; orarik turui-lm, ourrallt, sabrr\*tr>ft; flam j ; poll.  
diam. Petala poly itwrphm, litn<n> obloti('ti v. or<U, curva, acuminata.  
Anthera ob tang\*, locali\* j<r>H<li>; pnliinm elaraU, stipitib a\* ob.  
eottieit «orneu MMO»; •t\*mioodi» elougmu, i\*u<ntt»; rosstellom  
fMi-ram, tabalatan, •rtctom.*

In the 'Flora of British Indik' (p. 128) I have indicated the singular forms which the glands of *Collomia* take in several species of this genus, from simple matt fflalmMii or ohlon<sup>^</sup> glands to extinguisher-  
»iap<sup>«</sup>d hollow ooriateow brown nhininf bodiot, wmrtnirot\* split down  
OM »id<, if to which tin- cainii.lt' «f »)- pol|i«i»1 <k>><t'U, util itn.i  
the btw if which this latter is inMrtad. Thin bod/ m a den •op-  
ment of the gland itM-ir, i. and I lik KM, axpoaed al the bM«- of the  
autherwoeU.

It. H. orbicQlan. //ook. f. Fl. Brit. Ind. vi. 130. MU1< frn. foli; foli « rrmotiM parrit obtaneeolatiK, brncU'u ovario brevioribus, Bon Iu > tiiiNutit incundis, wpalit oHMculariublongu obioib, petalis obloogtt, labelto orbiculari carnoao bast aaocaCo.

## **IIAB. Sikkim Himalaya; Rangboo, in the Chuinlit Valley, *hing's CotUttor***

*Caulis uniflorus* spica 4-6 ft; follicularis. *Folia* 1-1<sup>1/2</sup> poll, litnira. 8e<sup>ms</sup>  
 pauciflora; tlnnw, ^ poll, dia. *Sepala* et petala 1-nervia  
 membranacea. *Antera* biocellata, loculis paullo divergentibus; pollinia  
 clavata, glandula; orbiculari in M'rtta; slamiodia v. proocwoa stigma-  
 tiferi • eatilia; roateltum minntum.

A number of specimens, of which more specimens are much wanted for Mithraic description.—J. 1. 11.

Pl.v\*2!9«A.//0h'chie. Fig. 1. Flower. 2. i'tul. 3. Liriun) column. 4. An-  
etW (urn of lip. 4. Pollinium. All enlarged.

it\* **2199** IS / *orthoclare.* Fig. «. i flower. 7. Uppe, «od 8 lo«w, M rface of  
C. *All enlarged.*



M. S. del et lith.

*Hajbenaria andamamca*, Hk.f.

## PLATE 2200.

### HABENABIA ANDAMANICA, *Hoohf.*

#### ORCHIDÆ. Subtribe OPHRTDEJE

H. (Ate) **andamanica**, *Hook f. Fl. Brit. Ind.* vi. 134; subelata, foliis medium versus caulis conferfca ovato-lanceolata acuta, racemo Jaxe multifloro, bracteis lanceolatis ovariis gracilibus brevioribus, floribus amplis, sepalis louge aristatis, petalorum segmento superiore angusto inferiore in laminas 3 filiformes secto, labelli lobis lateralibus flabellatim mnltifidis, intermedio 3-secto segmentis omnibus filiformibus, calcare ovario breviore.

HAD. Andaman Islands, *Parish.*

*Tubera giobosa. Caulis* 1-1<sup>1</sup>-pedalis. *Folia* 3-4-pollicaria, sessilia. *Racemus* 6-8-pollicaris; bractesd 1-1J-pollicares, lanceolate. *Sepala* i poll, longa, linear-oblonga. *Antherarum* tubuli locnlos subaeqnantes,\* porrecti; glandules polliniornm minntas; stigmatis brachia elongata, subcjindracea, rostellum erectum.

A handsome species, most nearly allied to the Ceylon and Travancore *S. macrostachya*, Lindl., which has longer bracts and snbeqnal segments of the lip. The specimen from which the drawing is made is unique.-J. D. H.

Fig- 1- Sepal. 2. Petal. 3. Column. All enlarged.