



Indian Botanic Garden Library  
BOTANICAL SURVEY OF INDIA

CLASS NO. 580.85

BOOK NO. JA

ACC. NO. 380.4



# ICONES PLANTARUM RARIORUM.

---

EDITAE

A

NICOLAO JOSEPHO JACQUIN  
BOTANICES PROFESSORE.

VOL. EL



6/7/64



---

VINDOBONÆ

PROSTANT APUO CHRISTIANUM FRIDERICUM WAPPLBR; LONDINI APUD B, WAIN BT FILIOMJ

LUGDUNI BATAVORUM APUD S, ST J, LUCHTMANS;

BIBLIOPOLAS;

---

AB ANNO 1786 AD 1793.



# CONSPECTUS TABULARUM.

Vol 3<sup>o</sup>

## OCTANDRIA.

### MONOGYNIA.

Tah

- 455 VENOOTHERA NOCTURNA, foliis lanceolate, dentato-ferratis, plants' j  
caule laevi, fubvillofo, ramofo. *Jacq. coll. vol. 3.*  
Caulis florentis pars superior.
- 456 OENOTHERA ODORATA, foliis linear-lancebatis, fibdentatis, undulatis;  
caule villofo, fuffruticofo. *Jacq. coll. vol. 5.*  
Tola planta florens primi anni.
- 457 GAURA FRUTICOSA, caule fruticofo. *Jacq. coll. vol. 1.*  
Ramus florens & fructescens. Folium caulinum majus. Flos auctus.

## TRIGYNA.

- 458 PAULLINIA CAULIFLORA, foliis pinnatis, glabris; pari inlimo ternato.  
*Jacq. coll. vol. 4.*  
Ramulus florēns. Caulis aphylli florentis pars. Auctus Flos totus, cuius tamen  
foliola calicina duo vicina majora hie a tergo confpectura fugiunt. Auctus calyx.

## DECANDRIA.

### MONOGYNIA.

- 459 CASSIA SENSITIVA, foliis quinquejugis & quadrijugis) foliis villofe ciliatis;  
glandula pedicellata & villofa inter omnia foliolorum paria. *Jacq. coll. vol. 2.*  
Caulis florentis pars maxima. Legumen. Semen. Aucta pars petioli communis  
cum glandula.
- 460 CASSIA POLYPHYLLA, foliis a quinquejugis ad duodecimjuga, in ramis fenio-  
ribus aggregatis j foliolis oblongis, ciliatis, facie glabris, dorfo hispidulis; glan-  
dula pyramidali feffiti inter foliola infima; pedunculus fubbifloris. *Jacq. coll. vol. 4.*  
Ramus florens. Petioli pars infima aucta.
- 461 GUILANDINA MORINGA, inermis; foliis compofitis -, pinnis inferioribus  
fubbipinnatis, fuperioribus fimpliciter pinnatis. *Linn. JyJi. Jacq. coll. vol. 2.* Mo-  
ringa. *Burm. thes. Rumpk. amb. Mouringou. Rheed. mal Nux Been zeyh'  
nica. Comm. hort.*  
Ramus florens. Fructus. Ejusdempars, dempta unavalwla. Semen. Nucleus.
- 462 TRIBULUS MAXIMUS, foliis fubquadrijugis; foliolis exterioribus majoribusj  
pericarpiis decafpermis, muticis. *Linn. fyft. Jacq. coll. vol. 4. Brown, jam. Plum,  
ic. Sloan, hiji. jauL*  
Radix truncata cum ramo five eaule integro aliisque abfeiflis. Flos a tergo vifus.  
Stylus cum ftigmate. Capfula una. Eadem aucta.

- 463 LIMONIA TRIFOLIATA, foliis ternatis; Tpinis geminis. *Linn. fyfi. Jacq. coll. vol. 3. Burm. ind.*  
 Ramus florens. Ramulus fructescens. Flos. Idem auctus dempta corolla. - Auctum piftillum. Bacca transverlim fecta. Duo Semina. Semen diffectum.
- 464 LEDUM LATIFOLIUM, foliis oblongis, margine revolutis, fubtus tomentosus; floribus subpentandris. *Jacq. coll. vol. 2. Ait kew. Ledum grOnlandicum. FLor. dan. Gunn. norv. Retz.fcatid.*  
 Ramus florens & fructescens.
- 465 ANDROMEDA CORIACEA, {male MARIANA) racemis axillaribus, fimpli-cibus ; foliis ovatis , integerrimis , nitidiflavis ; ramulis triquetris. -Ait. *kew. UHerit. Jürp. Jacq. coll. vol. 2.*  
 Pars trunci cum ramo florente & fruttescente. Aucta Genitalia. Anthera aucta\*

### DIGYNIA.

- 466 SAXIFRAGA MUTATÀ, foliis radicalibus aggregatis, lingulatis, cartilagineo-ferratis, ciliatis; cauleracemofo, foliofo; calycibus villofe viscidis. *Linn.jyji\* Jacq. noil. vol. 1. Allion. pedem. HalheliK Seguier.ver.*  
 Plantae duae florentes integrae, alter a major, altera pumila. *Wulfen.*
- 467 DIANTHUS ATRORUBENS, floribus aggregatis ; fquamis calycinis ovatis, ariftatis, tubo brevioribus ; foliis tronnatis, ftriatis. *Allion. pedem. Seg. veron\**  
 Tota planta culta florens-. Flos auctus. .

### TRIGYNIA.

- 468 STELLARIA BULBOSA, foliis lanceolato-ovatis 5 caule futokchotomo 5 pedunculis elongatis , unifloris ; radicerepente, bulbifera. *Jacq. coll. vol. 3.*  
 Plantulae integrae. Calyx cum Germine atiutto. *Wulfen,*
- 469 MALPIGHIA GLANDULIFERA, foliis ovatis, subintegerrimis, utfinque pilis decumbentibus obfessis ; racemis axillaribus ; pedicellis fuperne glandula truncata infstructis. *Jacq. coll. vol. 4. et vol. 5. tab. \$fig. 3.*  
 Ramus florens. Pedicellus auctus.
- 470 MALPIGHIA COCCIGERA, foliis subovatis, dentato-spinofis, glabris. *Linn. Jyji. Jacq. coll. vol. 2. Plum. ic.*  
 Ramus florens Calyx auctus

### P J E N T A G Y N I A

- 471 OXALIS RUBELLA, (nee HIRTA) caulescens ; pedunculis unifloris ; corollis campanulatis; ftjlis intermediis 3 filamentis edentulis. *Jacq. coll. vol. 3\* Jacq. oxal. num. 29.*  
 Tota planta florens.
- 472 OXALIS MULTIFLORA, (nee HIRTAE VARIETAS) caulescens; pedunculis unifloris; corollis campanulatis; flylis brevifimis; foliolis cuneiformibus. *Jacq. coll. vol. 3. Jacq. oxal. num. 28<*  
 Tota planta florens major.
- 473 OXALIS POLYPHYLLA , (nee VERSICOLOR ) ftipitata; foliis ternatis foliolis linearibus; filamentis gibbofe denticulatis; ifylis intermediis. *Jacq. coll\* vol. 3. Jacq. oxal. num. \$6.*  
 Tota planta florens elatior. Folioli aucti pars dorfallis.

# DODECANDRIA

## TRIGYNA.

- 474 RESEDA FRUTICULOSA**, foliispinnatis, apice recurvatis; floribus tetragynis; calycibus patentibus, quinquepartitis; caule bafi fruticofo. *Linn. fyft. Jacq. coll. vol. 3.*  
 Caulis florentis & fructescens pars fuperior. Flos auctus.
- 475 RESEDA MEDITERRANEA**, foliis integris trilobisque; calycibus florem subaequantibus 3 petalis petalorumque lobis lanceolatis & linearibus. *Linn. fyft. Jacq. coll. vol. 1.*  
 Tota planta florens & fructescens. Flos auctus.
- 476 EUPHORBIA SCORDIFOLIA**, dichotoma; hirfuta; foliis crenulatis, subovatis, obtufis; pedunculis unifloris, folitariis. *Jacq. coll. vol. 5.*  
 Planta jlorens. Flos auctus.
- 477 EUPHORBIA PICTA**, dichotoma; viilofa; foliis repandis, ovatis, pictis, acutis; pedunculis unifloris, folitariis; caule erecto. *Jacq. coll. vol. 3.*  
 Caulis florentis pars fuperior. Radicis pars cum caule infimo ramofiffimo. Flos auctus. Semen auctum.
- 478 EUPHORBIA PILULIFERA**, subdichotoma; foliis ferrulatis, ovali-oblongis; pedunculis bicapitatis, axiliaribus } caule erecto. *Linn. Jyft. Jacq. coll. voL 2. Burnt, thes.*  
 Tata planta. Folium in dorfo confpectum. Flos auctus. Semen auctum.
- 479 EUPHORBIA NUDIFLORA**, subtrichotoma 3 foliis integerrimis, ovatis, subundulatis; pedunculis cymofe ramolis; caule fruticofo. *Jacq. coll. vol. 3.*  
 Caulis pars fumma. Flos auctus.
- 480 EUPHORBIA CYATHOPHORA**, (*male HETEROPHYLLA*) umbella trifida, subdichotoma; foliis panduraeformibus. *Jacq. coll. vol. 1. uhifnonyma de-leanatur,*  
 Caulis pars fuperior. Flos auctus.
- 481 EUPHORBIA ANGULATA**, umbella quinquefida; bifida; involucellis ovatis; foliis obiongis; caule ftrajp-angulato. *Jacq. coll. vol. 2.*  
 Tota planta. Flos auctus.
- 482 EUPHORBIA LITERATA**, umbella quinquefida; trifida; bifida; involucellis ovatis'; foliis lanceolatis, subpilosis, obfolete ferrulatis' petalis integris; radice \*annua. *Jacq. coll. vol. 2.*  
 Plantula tenella. Eadem primum florens. Eadem adulta. Flos auctus.
- 483 EUPHORBIA SERRATA**, umbella quinquefida; bis bifida -, involucellis diphyllis, reniformibus; foliis amplexicaulibus, ferratis. *Linn. fyft. Jacq. coll. vol. 2. Cluf.hj/i. Bauh.pin.*  
 Caulis florens. Ejusdem pars inferior. Flores aucti. Capula. Semen.
- 484 EUPHORBIA PUNICEA**, umbella quinquefida -, trifida; involucellis ovalibus, acuminatis, coloratisj capulis glabris; foliis lanceolato-cuneiformibus, subtus glaucis. *Swartz. prodr. Ait. kew. Jacq. coll. vol. 3.*  
 Trunci florentis pars fumma.

- 485 EUPHORBIA NICAEENSIS, umbella quinquefida ad dechifidam ; dichotoma; involucellis cordato-fubrotundis ; petalis bicornibus > foliis lanceolato-fpathulatis, glaucis. *Allion. pedem. Jacq. coll. vol. 2.*  
Planta fiorens. Flos auctus.

## ICOSANDRIA, MONOGYNIA.

- 486 EUGENIA BARTENSIS , foliis integerrimis, ovato-lanceolatis; pedunculis axillaribus, urifloris & aggregatis, vel multifloris. *Joey, ftirp. timer? Jacq, coll. vol. 3<*  
-Ramus florens. Flos auctus, JBacca.

## P E N T A < Y N I A.

- 4g7 MESEMBRYANTHEMUM CORDIFOLIUM, foliis oppofitis, petiolatis, cor-datis ; calycibus quadrifidis ; caule tereti. *Linn.jyji\** *Jacq. coll. vol. 2. G<2-xin. objl*  
Planta florenS.  
488 MESEMBRYANTHEMUM CUNEIFOLIUM, foliis oppofitis^ cuneiformibus, obtufis, fcabridis ; papulis oblongis ; foliolis calycinis oblongis , obtufis, medio coarctatis. *Jacqxoll. vol. 2. Mefembryanthemum limpidum. Ait. hew.*  
Planta florens & fructescens.  
48p MESEMBRYANTHEMUM POMERIDIANUM, foliis planiusculis, lato-lanceolatis, Wibus , fubciliatis, diftinctis ; caule pedunculis germinibusque hirtis. *Linh.fyft. Jadq. coll. vol. 2. Linn, decad. Ficoidea aizoides &c. Herm. lugd> Seb. mus.*  
I'ota planta florens^

## POLYGYNIA.

- 490 POTENTILLA SALISBURGENSIS, foliis hirfutis, incifo-ferratis , obtufis 5 radicalibus quinatis \$ caulinis ternatis \$ caule adfeendente, compreflb, paucifloro; calycibus fubdentatis. *Jacq. coll. vol. 2.*  
Tota planta. *Haenke,*  
491 POTENTILLA SUBACAU LIS, foliis ternatis fubquinatisve \* dentatis, utrinque tomentofis; fcapis decumbentibus. *Linn.JyJi. Jacq. coll. voL 2. Gerard\** *gatoprov. Gmel. Jibir.*  
Tota planta. *Wulfen.*

## POLYANDRIA.

### T R I G Y N I A.

- 492 ACONITUM TAURICUM, caule fubfimpliei; racemo brevi, compactoj galea breviter rofirata ; foliis nitentibus , ^ quinquepartitis \$ segmentis profunde incifis. *Jacq. coll. vol. 2. Clus. hijt.*  
Caulis florens. Radix. *Wulfen.*

# DIDYNAMIA.

## GYMttOSPERMIA.

- 493 STACHYS RUGOSA, verticillis fexfloris ; foliis lanceolatis, baft anguftatisj tomentofis\* ferratis, rugōus; caljcibus muticis. *Ait. kev>i Jacq. coll. vol. 5.*  
 Ramus florens; Aucta Calyx & Corolla:
- 494 SATUREJA RUPESTRIS, pedunculis folitariis, axillaribus; bis dichotomis, fecundis ; foliis ovato-lanceolatis, ferrulatis, fubtus punctatis; *Jacq\* coll. vol. 2\** Satureja thymifolia; *Scop. cam.*  
 Planta florens. Peduriculi frutescerites. Corolla aucta. *Wulfen.*
- 495 OCYMUM GRATISSIONUM, caule ftifFruticofo; foliis lanceolato-ovatis ; facemis teretibus. *Linn. fyft. Jacq. coll. Vol. 4. Burnt\* thes.*  
 Caulis pars fupefior. Flos auctus.

## ANGIOSPERMIA.

- 496 SELAGO FASCICULATAj corymbo multiplid; foliis dbverfe ovatis, glabris. *Linn. Jyft. Jacq. coll. vol. 3.*  
 Ramus florens. Aucta corolla cum Staminibus & Styld. Auctus Calyx Gum Bractea & Piftillo\*
- 497 CELSIA LINEARIS, foliis linedrManceblatisi *Jacq. toU. vol. 21*  
 Ramus florens.
- 498 MANULEA TOMENTOSA, foliis tomentofls \ caulibus foliofis ; pedunculis multifloris. *Linn. fyft. Jacq. coll. vol. 4.*  
 Pars caulis. Foliolum calycinum auctum.
- 499 ANTIRRHINUM PARVIFLORUM , foliis linearibus, glabris; inferibribus quinis vel quaternis \*, fuperioribus alternis ; floribus racemofis; calycibus capitato-villofis'; caulibus erēctis. *Jacq. cbl. vol. 4.*  
 Radix cum caulis abfciffis. Caulis unus florens & fructe'scens. Flos auctus.
- 500 VOLKAMERIA KAEMPFERI, foliis cordatis, fubrotundis, villofulis; flor<sup>^</sup> bus pāniculatis ; cauie erecto. *Jacq. coll. vol. 3. Keentpf. reliq.*  
 Caulis florentis pars fumma. Folium ex majoribus. Seminiis maturi Nucleus. Semen. Germen auctum, horizontaliter discitum. Fructus matus cum calyce' perfiftente.
- 501 CITHAREXYLUM ERECTUM, rāmis teretibus; racemis erēctis, terminalibusj calycibus dentatis. *Swan, prodr. Jacq. coll. vol. 3, Brown, jam. num. 5.*  
 Raiiiulus florens.
- 502 DURANTA PLUMIERI, calycibus fructescentibus contortis; *Liniufyft\* Jacq\* coll. vol. 5.* Caftorea racemofa. *Plum. ic.*  
 Ramus florens. Baccae.

## TETRADYNAMH

### SILICULOSA.

- 503 ALYSSUM GEMONENSE, caule herbaceo; ramis divaricatis ', foliis radicals bus obverfe ovatis, (ubtomentofis; filiculis inflatis. *Linn. Jyft. Jacq. coll. vol. 2. AlyfTum petraeum. Arduin. Jpec.*  
 Tola planta. *Wulfen\**

## S I L I Q U G · 8 A,

- 504 CRAMBE FILIFORMIS, fuffruticofa; foliis inferioribus pinnatis, fcabris; pinnata extima fubrotunda, magna. *Jacq. coll. vol. 5.*  
 Tota planta primi anni. Genitalia aucta. Piftillum cum pedicello suo auctum.
- 505 SISYMBRIUM LIPPIZENSE, filiis declinatis, brevibus; foliis inferioribus pinnatis; fummis inferiorumque foliolis integerrimis. *Jacq. coll. vol. 2.* Erjfimum verum. *Dalech. lugd.*  
 Tota planta. *Wulfeiu*
- 506 HELIOPHILA INTEGRIFOLIA, foliis lanceolatis, indivisis. *Limufyfi. Jacq. coll. vol. 3.*  
 Tota planta florens. Siliqua. Semina.

## MONADELPHIA. PENTANDRIA.

- 507 MftLOOHIA CÀRACCASANA > floribus capitatis; capfulis fubrotundis; foliis cordatis, crenatis; caule arborescente. *Jacq. coll. vol. 2.*  
 Ramulus florens.
- 508 ERODIUM TRILOBATUM, pedunculis multifloris; foliis cordatis, fubrotundis, tripartitis, hirfutis, ferratis. *Jacq. coll. vol. 4.*  
 Radix caulescens. Caulis pars. Genitalia aucta.
- 509 ERODIUM RIBIFOLIUM, pedunculis fubquinquefloris', calycibus ariftatis; foliis cordatis > fublobatisi ferratis. *Jacq. coll. vol. 5.*  
 Tota planta. Foliolum calycinum auctum.

## HEPTANDRIA.

- 510 PELARGONIUM RAPACEUM, acaule; radice rapacea, multicipite; foliis bipinnatis, villofis, incifisj filamentis decem, quinque fertilibus. *Jacq. coll. vol. 4.* Geranium africanum myrrhidis folio. *Comm. hort.*  
 Tota planta. Calyx cum Genitalibus\* Petalum fuperius. Idem laterale. Idem infimum.
- 511 PELARGONIUM (*nee GERANIUM*) ASTRAGALIFOLIUM, acaule; radice rapacea mullicipite \$ foliis pinnatis; foliolis ovatis; filamentis decem, quinque fertilibus. *Jacq. coll. vol. 3.* Cavanill. difj. Commel. prod.  
 Tota planta.
- 512 PELARGONIUM CARNEUM, acaule; radice rapacea, fimplice; foliis bipinnatis; pinnulis incifis, glabriusculis; filamentis decem, quinque fertilibus. *Jacq. coll. vol. 4.*  
 Tota planta.
- 513 PELARGONIUM BARBATUM, acaule; radice rapacea, fimplice; foliis pinnatis; pinnispinnatifidis, lanceolato-linearibus, acutis, apice barbatis; filamentis decem, quinque fertilibus. *Jacq\* coll. vol. 4.*  
 Tota planta.
- 514 PELARGONIUM MELANANTHON, acaule; radice rapacea, fimplice; foliis pinnatis, hispidulis; pinnis lobato-incifis, obtufis; filamentis decem, quinque fertilibus. *Jacq. coll. vol. 4.*  
 Tota planta.

- 515 PELARGONIUM TRIPHYLLUM, acaule; radice rapacea, fimplice; foliis ternatis, crenatis, glabris ; filamentis decem, quinque fertilibus. *Jacq. coll. vol. 4.*  
Tota planta.
- 516 PELARGONIUM HETEROPHYLLUM, acaule; radice rapacea, fimplice; foliis ternatis integrisquc, glabris, ciliatis; foliolis faepe lobatis; filamentis decem, quinque fertilibus. *Jacq. coll. vol. 4.*  
Tota planta. Foliola duo integra ex alia individua planta.
- 517 PELARGONIUM NERVIFOLIUM, acaule; radice rapacea, fimplice; foliis ternatis trilobis integrisque, multinerviis, glabris , ad oras aculeato-ciliatis; filamentis decem, quinque fertilibus. *Jacq. coll. vol. 4.*  
Tota planta. Marginis folii portio aucta.
- 51% PELARGONIUM LONGIFOLIUM, acaule; radice rapacea, fimplice; foliis lanceolatis, acutis, glabris; fenoribus faepe pinnatis; filamentis novem, quatuor fertilibus. *Jacq. coll. vol. 4. Cavan. diff.*  
Tota planta.
- 519 PELARGONIUM CILIATUM, acaule; radice rapacea, fubfimplice; foliis lanceolatis, utrinque acutis, ciliatis, fubtus glabris, supra pilofis, faepe appendiculatis; filamentis decem, quinque fertilibus. *Jacq. coll. vol. 4.*  
Tota planta.
- 520 PELARGONIUM DEPRESSUM, acaule 5 radice rapacea, fimplice; foliis angule lanceolatis, acutis , glabris ; umbellis tandem deprellls '3 filamentis novem, quatuor fertilibus. *Jacq. coll. vol. 4.*  
Tota planta
- 521 PELARGONIUM LONGIFLORUM, acaule; radice rapacea, fimplice; foliis lanceolatis, acutis, glabriusculis ; petalis longiflmis; filamentis decem, quatuor fertilibus. *Jacq. coll. vol. 4.*  
Tota\*planta.<sup>#</sup>
- 522 PELARGONIUM DAUCOIDES, fubcaulescens; radice tuberofa; fcapo fimplicissimo ; foliis tripinnatis, hirfutis ; f pinnulis lanceolato-linearibus ; filamentis decem, septem fertilibus. *Jacq. coll. vol. 5.*  
Tota planta.
- 523 PELARGONIUM CHAMAEDRYFOLIUM, pedunculis fubbifloris; caulis herbaceis, decumbentibus; foliis oblongis, obtufis , ferratis , villosis ; filamentis decem, quinque fertilibus. *Jacq. coll. vol. 5.*  
Tota planta. Flos auctus.
524. PELARGONIUM TRICHOSTEMON, pedunculis paucifloris ; foliis fubrotundato-ovatis, obtufe ferratis , holofericeis ; caule bienni, erectiusculo ; filamentis decem, hirfutis ; quinque fertilibus. *Jacq. coll. vol. 5.*  
Tota planta. Genitalia aucta.
- 525 PELARGONIUM O NOTHERAE (*male GERANIUM OVATUM*), pedunculis paucifloris; foliis lanceolatis, ferratis, tomentofis; caule bienni, procumbente ; filamentis decem, quinque fertilibus. *Linn. Jyft. Jacq. coll. vol. 2.*  
Tota planta.
- 526 PELARGONIUM CORONOPIFOLIUM, pedunculis paucifloris; foliis lanceolato-linearibus, dentato-ferratis, supra glabris, fubtus hirtulis ; caule fuffructifero ; filamentis decem, quinque fertilibus. *Jacq. coll. vol. 5.*

- Tota planta. Petali fumffii aucta pars inferior. Dein ex varietate Pedunculus biflorus & Folium latius, incifitm.
- 527 PELARGONIUM VIOLAREUM, pedunculis paucifloris ; foliis oblongis, obtusifolis, caule ferrato-incilis, villofis; caule bienni; filamentis decem; quinque vel quatuor fertilibus ; fterilibus fquamaeformibus. *Jacq. coll. vol. 5.*  
 Tota planta. Foliis pars aucta.
- 528 PELARGONIUM CORIANDRIFOLIUM, pedunculis bi-quinquefloris ; corollis tetra-vel pentapetalis ; caule bienni ; foliis bipinnatis ; filamentis decem, quinq-  
 tiae fertilibus. *Linn. ffft. Jacq. coll. vol. 5. Cavan. diff. Rivin. pent. Jit. kew.*  
 Tota planta florens & fructescens.
- 529 PELARGONIUM CAUCALIFOLIUM, pedunculis unifloris; corollis tetrapetalis5 caule bienni; foliis bipinnatifi, fublinearibus ; filamentis decem, septem fertilibus. *Jacq. coll. vol. 5.*  
 Tota planta.
- .530 PELARGONIUM BULLATUM, pedunculis bifloris; corollis tetrapetalis; caule bienni; foliis ternatis, J>innatifidelobatis, utrinque fubhispidulis; filamentis decem, quinque fertilibus. *Jacq. coll. vol. 5.*  
 Caulis. Duo Folia infima.
- 531 PELARGONIUM BETONICUM, pedunculis paucifloris ; corollis tetrapetalis, rarius pentapetalis; caule bienni; foliis pinnatifide lobatis , hirtulis ; filameritis decem, quinque fertilibus. *Jacq. coll. vol. 5. Cavan. diff. Riv. pent.*  
 Tota planta junior.
- 532 PELARGONIUM LACERUM, pedunculis quinquefloris; corollis pentapetalisj caule bienni, pilofo ; foliis pinnato-lobatis, incifis; filamentis decem, quinque fertilibus. *Jacq. coll. vol. 5.*  
 Caulis florens & fructescens.
- 533 PELARGONIUM LONGICAULE, pedunculis uni-quinquefloris 5 corollis te-  
 trapetalis, rarius pentapetalis; caule bienni, hirfuto ; foliis pinnato-lobatis; filar-  
 mentis decem, septem fertilibus. *Jacq. coll. vol. 5.*  
 Caulis pars.
- 534 PELARGONIUM MULTICAULE, pedunculis fubquinquefloris ; corollis te-  
 trapetalis; caule bienni> glabro ; foliis pinnato-lobatis , glatyis ; filamentis de-  
 cem, septem fertilibus. *Jacq. coll. vol. 5.*  
 Radix cum uno Caule cumque truncatis reliquis.
- 535 PELARGONIUM ANEMONAEFOLIUM, pedunculis fubquinquefloris; co-  
 rollis pentapetalis; caule bienni; foliis pinnato-lobatis, fubtus hirtulis, supra gla-  
 bris; filamentis decem, septem fertilibus. *Jacq. coll. vol. 5. Riv. pent.*  
 Tota planta.
- 52,6 PELARGONIUM HIRTUM, uiribellis fubquinquefloris; foliis tripinnatis, hirfu-  
 tisj pinnulis linearibus; caule carrofo. *Jacq. coll. vol. 5. Cavan. diff. Burm. geran\*.*  
 Caulis florens & fructescens.
- 537 PELARGONIUM TOMENTOSUM, umbellis multifloris, fimplicibus compo-  
 fitisque; foliis cordatis, fubquinquelobis, ferratis, tomentofis, mollifimis; caule carnofo. *Jacq. coll\* vol. 5\**  
 Ramus florens.

- 538 PELARGONIUM RIBIFOLIUM, umbellis multifloris, depreffis, fubdimidiatis; foliis cordatis, trilobis, ferratis, hirtutulis, lobis lobato-incisis; caule carnoſe. *Jacq. coll. vol. 5.*

Ramus florens.

- 539 PELARGONIUM CORTUSAEIFOLIUM, umbellis fubquadrifloris; foliis cordatis, inciso-lobatis, acute dentatis, glaucis, glabris; ftipulis ovatis \$ caule fruticofo, erecto. *Jacq. coll. vol. 5. Ait. kew. Geranium tabulare. Cavern, diff. Burm. ger.*

Caulis florentis pars fuperior. ' Flos parvitatem variaris.

- 540 PELARGONIUM FUSCATUM, umbeilis quadriflorisj foliis cordato-fubrotundis, fubquinquelobis, acute ferratis, glabriusculisj ftipulis ovatis; caule fruticofo, erecto, juniore fcabro; *Jacq. coll. vol. 4.*

Caulis *florentis pars.*

- 541 PELARGONIUM PATULUM, pedunculis bifloris -, foUis fubreniformibitis, lobato-incisis, acute crenatis; supra glabris^ caule fruticofo } filamentis decem, fex fertilibus. *Jacq; coll. vol. 4;*

Caulis florens.

- 542 PELARGONIUM (*nee GERANIUM*) SCABRUM, umbellis paucifloris; foliis cuneatis, femitrifidis, fcabris 3 lobis lanceolatis, laxe ferratis. *Jacq. coll. vol. 2. Ait. kevii*

Ramus florens.

- 543 PELARGONIUM BAtSAMEUM, umbellis paucifloris \ foliis quinquepartitis, hirtulis j lobis lanceolatis, acutis, incilis, rigidulii; caule fruticofo. *Jacqt coll. vol. 5.*

Caulis florentis pars fumma.

- 544 PELARGONIUM TERNATUM, pedunculis umflorisj foliis ternatis, fcabris, rigidis; foliolis cuneatis , apice dentatis ; filamentis decem, quinque fertilibus \$ daule fruticofo. *jacq. coll. vol. 5. Linn. Jjrft. Cavan. cliff.*

Ramus florens. Trunci pars.

- 545 PELARGONIUM HERMANNIAEFOLIUM , pedunculis bifloris 5 foliis cuneato-fubrotundis, incisis, hirtis, rigidis, diftichis; caule fruticofo. *Linn, fyft. Jacq. coll. vol. 5. Berg. cap.*

Ramus florens & fructescens.

## DECANDRIA.

- 546 GERANIUM ARGENTEUM, pedunculis bifloris; petalis emarginatis 5 foliis subpeltatis, septempartitis, trifidis, tomentofo-fcriceiSi *Linn, fyft. Jacq. coll. vol. 2: Bauh. hift. Ség. ver.*

Tota planta. *Wulfem*

## POLYANDRIA.

- 547 SID A PALMATA, foliis palmatis, peltatis| laciniis lanceolatis, ferratis, incis. *Cavan. diff. Jaccf. coll. vol. 2. Sida iatrophoides. Ait. kew.*

Ramus florens. Fructus. Capula.

- 548 MALACHRA FASCIATA, capitulis triphyllis, fubquinquefloris ; caule foliis-\* que hispidissimis. *Jacq. coll. vol. 2.*

Ramus florens. Caulis pars. Caljeis exterioris foliolum expanfum. Periarithium interius auctum^

- J4p MALACHRA ALCEAEFOLIA*, capitulis pntaphyllis; fabdecemfioris; caule rariter hispido; foliis glabriusculis; Jtipulis utrinque fubquaternis. *Jacq. coll. vol. 2*  
Caulis' florentispars fumma. Capitulum fructescens. Semen. Capula. Perianthii exterioris foliola duo. Periantlium interius.
- 550 HIBISCUS DOMINGENSIS**, caule arboreo, feniore inermi, juniore armato aculeis rectis', foliis fubquiaquelobis j lobis fubovatis, ferratis, acutis. *Jacq. coll. vol. 3.*  
Ramus florens.
- 551 HIBISCUS DIVERSIFOLIU&**, caule arboreo ramis petiolisque aculeatis; foliis ferratis, inferioribus quinquelobis, fubfequis trilobis; fummis oblongis integrisque. *Jacq. coll. vol. 2.*  
Caulis florentis pars fumma. Capula. Valvula una. Semen.
- 552 PAVONIA URENS**, caule fruticofo-, foliis palmatis; floribus axillaribus, glo-meratis, fubfemilibus. *Cavan. diff. Jacq. coll. vol. 3.*  
Ramus florens. Semen. Capula. Fructus totus. Filamentum auctum.
- 55\$ CAMELLIA JAPONICA**, foliis acute ferruiatis, acuminatis. *Lijm. Jyft. Jacq. coll. vol. 1. Thunb. jap\**  
Ramus florens.

## D I A D E L P H I A .

### HEXANDRIA.

- 554. FUMARIA ACAULIS'**, fcapis aphjllis, longitudine foliorum; foliis pinnatis, pinnis trilobis. *Jacq. coll. vol. 2.*  
Tota planta» Flos auctus. *JVulfeil.*

### DECANDRIA.

- 555 SPARTIUM DECUMBENS** (*nee GENISTA HUMIFUSA*), caule decumbente, ramofo; foliis folitariis, ovatis; floribus longe petiolatis. *Ait. kew. Hall, helv. num. 355. Durande. bourg. Jacq. coll. vol. 2. Tjenifta pedunculata. VHerit. Jiirp. nov.*  
Tota planta florens. *Wulfen.*
- 556 GENISTA SERICEA**, foliis lanceolatis, fubtus fericeo-villofis; ramis teretibus, ftriatis, erectis j racemis paucifloris, fecundis 5 bracteis calyce longloribus. *Jacq. coll. vol. 2.*  
Tota planta florida. *Wulfen.*
- \$51 GENISTA HISPANICA**, fpinis decompositis 5 ramis floriferis inermibus 5 foliis linearibus, pilofis. *Linn, fyft. Jacq. coll. vol. 2. Allion. pedem. Genifta fyl-vefris. Scop. earn.*  
Tota planta florens. *Wulfen.*
- 55S PHASEOLUS SEMIERECTUS**, caule femivolubii; floribus fpicatis; calycibus ebracteatis; alis expanfis, majoribus; foliolis ovatis. *Linn fyfi. Jacq. coll. vol. 1.*  
Radix cum caulis infimaparte. Caulis pars fupertor. Carina. Vexillum. Genitalia. Semen. Idem auctum. Legumen.

- 559 DOLICHOS ACINACIFORMIS**, voluhilis; leguminibus gladiatis, dorfo tricarinatis, mucrone recurvo terminatis; feminibus arillatis. *Jacq. coll. vol. 1.* an Dolichos eniformis. *Linn. fyji. pag. 6\$j. num. 8.?*  
 Caulis florentis pars. Legumen. Fruftum dorfi tricarinati. Fruftum facieL Sec-  
 tio leguminis transverfa. Valvulae pars cum feminibus contentis. Semen.
- 360 DOLICHOS GLADIATUS**, volubilis; leguminibus gladiatis, dorfo tricarinatis; in mucronem rectum terminatis 3 feminibus arillatis. *Jacq. coll. vol. 2.*  
 Ramulus florens. Legumen. Valvulae pars interior. Semen. Flos. In hujus & pre-  
 cedentis defcriptionibus error, qui irrepfit, corrigendus eft. Scilicet in Dolicho  
 acinaciformi legatur *legumen unciam latum*; in Dolicho gladiato *legumen fescun-  
 ciam latum, tricarinatum loco trimarginatum*.
- 561 ASTRAGALUS EXSCAPUS**, acaulis, exfcapus 5 leguminibus lanatis; foliis  
 villofis. *Linn. fyft. Jacq. coll. vol. 2.* Aftragalus fupinus, percnnis 5 foliis \$r  
 filiquis hispidis. *Buxb. cent.* Glaux montanum acaulon. *Bauh. pin\** Glaux la-  
 nuginofa montana acaulos. *Rupp. jén\**  
 Tota planta cum radice abfciffa. Semen. Legumen hians. Idem adhucdum claufum.
- 562 PSORALEA BIPEDUNCULATA**, foliis ternatis<sup>^</sup> foliolis oblongis, obtufis;  
 pedunculis axillaribus, racemofis, gemellis<sup>5</sup> altera multifloro, altero paucifloro.  
*Jacq. coll. vol. 4.*  
 Ramus florens & fructescens. Flos auctus.
- 563 PSORALEA PHVMATODES**, foliis pinnatis; foliolis fubtus glandulofe tuber-  
 culatis; pedunculis fpicatis, axiliaribus & terminalibus. *Jacq. coll. vol. 4. uhi di-  
 citur P. EIMPHYSODES.*  
 Ramus florens. Flos auctus. Genitalia aucta.
- 564 AESCHYNOMENE BISPINOSA(/z<sup>^</sup>SESBAN)**, caule lierbaceo, ftrigis rarif-  
 limis armato; foliolis glabris; cofta fpinofula 5 leguminibus teretibus, longiflimis<sup>^</sup>  
 glabris. *Jacq. coll. vol. 2.*  
 Caulis florentis & fructescens pars fumma. Legumen maturum. Semen.
- 565 HEDYSARUM GYRANS**, foliis ternatis, glabris; foliolis oblongis; lateralibus  
 parvis; racemo compojito, terminali. *Linn. fyjt. Jacq. coll. vol. 3.*  
 Caulis florentis pars fumma. Calyx ciim Genitalibus auctus. Vexillum. Ala. Ca-  
 rina. Legumina duo. Semen.
- 566 HEDYSARUM VESPERTILIONIS**, foliis simplicibus ternatisque; illis & ho-  
 rum foliolo medio biiobis, lobis divaricatissimis; racemo terminali, ramofulo; le-  
 guminibus retrofractis, caljci inflato inclufis. *Linn. Jyji. Jacq. coll. vol. 2.*  
 Tota planta florens & fructescens. Flos auctus. Legumen auctum. Semen auc-  
 tum. Idem in naturali magnitudine.
- 567 HEDYSARUM PICTUM**, foliis pinnatis, *in faciei disco pictis*; racemo fpicato,  
 erecto, longifimo; bracteis coloratis, caducis; leguminibus retrofractis; cau-  
 le fuffrutofo. *Jacq. coll. vol. 2.*  
 Caulis florentis pars fumma. *a.* Racemus ante florū expantionem, fpicam men-  
 tiens. *b.* Pars Pedunculi communis fructescens. *c.* Pedicellus auctus. *d.* Flos cae-  
 rulescens. *e.* Calyx auctus. *f.* Legumen. *g.* Idem auctum. *h.* Semen.
- 568 HEDYSARUM MURICATUM**, foliis pinnatis, ad oras muricatis; pedunculis  
 axiliaribus unifloris, terminalibus racemofis; leguminibus muricatis; cauie fuffru-  
 ticofo, decumbente. *Jacq. coll. vol. 5.*  
 Tota planta florens & fructescens. Folioli pars aucta. Semen. Flos auttus.

- 569 INDIGOFERA HIRSUTA\* foliis pinnatis ; hirfuta; caule erecto ; floribus fpicatis; leguminibas pendulis , lanatis. *Linn. fyft. Jacq. coll. vol. 2.* Aftragalus fpicatus, filiquis pen&uluis hirfufls. *Burnt, thes. KattuTagera. BJeed. mat*  
Tota planta. Semen. Legumen. Flos auctus.
- 570 INDIGOFERA HENDECAPHYLLA, foliis pinnatis; floribus fpicatis; legu\* minibus tetragonis, pendulis, villofis ; caulibus profratis. *Jacq. coll. vol. 2.*  
Radix cum uno caule, truncatis reliquis. Flos auctus. Semina. Spica fructescens-
- 571 INDIGOFERA DENDROIDES, foliis pinnatis, glabris; racemis axillaribus; leguminibus linearibus, glabris ; caule erecto, annuo. *Jacq. coll. vol. 2.*  
Tota planta florens. Legumen. Semen. Flos auctus, utrinque vifus.
- 572 GALEGA LONGIFOLIA, foliis ternatis; caule volubili; leguminibus rectis. *Jacq. coll. vol. 2.*  
Ramus florens & fructescens. Calyx auctus. Vexillum. Ala. Carina. Legumen\* Semen,
- 573 GALEGA EILIFORMIS, foliis ternatis 5 caule volubili; leguminibus recurvatis. *Jacq. coll. vol. 2.*  
Ramus florens. Semen. Legumem Corolla aucta\* Calyx auctus.
- 574 GALEGA CAPENSIS, leguminibus teretibus, villofulis ; racemis filifofmibus, longillimis; floribus oppofitis, diftantibus ; foliolis glabris; caule decumbente, fffruticofo. *Jacq. coll. vol. 2.*  
Tota planta florens. Calyx auctus. Semen. Legumen.
- 575 GALEGA CINEREA, leguminibus Metis, patentibus, pedunculatis; racemis oppofitifoliis ; foliolis mucronatis, fubtus villofis. *Lhtn-fjij. Jacq. coll vol. 2.*  
Galega herbacea fubcirtorea villofa. *Brown, jam.*  
Plantae florentis & fructescensis Caulis princeps. Legumen. Semem
- 576 GALEGA DUBIA, leguminibus racemolis; tota incano-viliofa; foliis fubnovenjugis, parvis. *Jacq. coll. vol. 4\**  
Tota planta florens.

## S Y N G E N E S I A POLYGAMIA AEQUALIS.

- 577 TRAGOPOGON CAPENSIS, calycibus corollae radio brevioribus, hispidis\* foliis runcinatis, hispidis, fpinofo-dentatis. *JaCq\* coll. vol. 2.*  
Caulis florentis & fructescensis pars fuperior. Semen. Folium inferius.
- 578 HIERACIUM INCARNATUM, fcapo multifloro, nudo ; foliis obovato-oblongis, repando-denticulatis, hirfutis. *Jacq. coll vol. 2. Scop, earn.*  
Tota planta. Flosculus. *Wulfeiu*
- 579 CARDUUS ACAULIS, acaulis ; calvce glabro. *Unn.jyfi. Jacq. coll vol. u* ubi Synonyma plurima leguntur.  
Tota planta. Flosculus. *Wulfen.*
- 580 CACALIA VILLOSA, caule herbaceo; tota villofa & afperula; foliis lyratis; inferioribus petiolatis ; fuperioribus amplexicaulibus ; *fummls \ ^ ^ x ^ Jacq* coll. vol. 2. Senecio purpureus. *Linn. fyji. Berg. cap.*  
Caulis pars fuperior. Flosculus auctus. Folium inferius.

- 58\* CACALIA PEUCEDANIFOLIA, caule fruticofo; foliis fubcipinnatis, filiformibus. *Jacq. coll. vol.* ^ Senecio peucedanifolius. *Linn. JyJi.*  
Ramus florens.
- 582 EUPATORIUM MYOSOTIFOLIUM, foliis lanceolatis, in petiolum angustatis, villofis, ciliatis, floribus paniculatis ; calycibus multifloris. *Jacq. coll. vol. 2.*  
Ramulus florens. Flosculus auctus.
- 583 CALEA ASPERA, floribus pedunculatis; foliis oppofitis, inaequaliter ferratis; caule tetragono. *Jacq. coll. vol. 2.*  
Ramus florens. Corolla adulta cum Stigmatibus exsertis aucta. Flos nudius aditus cum Antheris exsertis cumque Palea auctus.
- 584 SPILANTHUS EXASPERATA, foliis ovatis, obfolete ferratis, omnibus oppofitis; caule flexuoso, paniculato, verruculofo, inferne radicante. Spilanthes radicans mihi dicta fuit in *Collectamis vol. 3.*, oblitus nominis in tabula jam diu anteasculpti.  
Ramus. Trunci pars. Flosculus auctus. Flos totus auctus. Auctum fructus Receptaculum cum Calyce Paleisque exterioribus.

### P O L Y G A M I A     S U P E R F L T J A .

- 585 CONYZA CAROLINENSIS, foliis lanceolato-oblongis, acutis, integerrimis, tomentofis, viscofis ; caule arboreo. *Jacq. coll. vol. 2.*  
Ramus florens. Flos faemineus idemque hermaphroditus aucti.
- 586 ARNICA GLACIALIS, foliis alternis, repando-fudentatis; caulinis lanceolatis, fclTilibus; radicaiibus rotundato-ovatis, petiolatis. *Jacq. coll. vol. 1.*  
Plantae duae integroe. *Wulfen.*
- 587 SENECIO ROSMARINIFOLIUS, corollis radiantibus; foliis linearibus, margine revolutis; caule fruticofo. *Limufyji.* *Jacq. coll. vol. 1.*  
Caulis ramofus florens.
- 588 LEYSERA GNAPHALODES, foliis fparfis; floribus pedunculatis. *Linn. fyfi.*  
*Jacq. coll. vol. 2.* *Seb. mus.* After aethiopicus, Itaechadis folio, flore aureo.  
*Herm. lugdb.* *Breyn. prodr.*  
Caulis pars cum ramis florentibus. Calyx auctus\* Semen auctum faeminae. Idem auctum hermaphroditae.
- 589 ZINNIA ELEGANS, fpliis amplexicaulibus ; floribiis pedunculatis 3 hermaphroditarum feminibus breviffime bimucronatis. *Jacq. coll. vol. 5.*  
Ramus florens. Palea aucta. Flos hermaphroditus auctus. Flos faemineus. Squama calycina. Flos hermaphroditus, paleae fuæ inclufus.
- 590 ZINNIA TENUIFLORA/foliis effilibus; floribus pedunculatis; feminibus hermaphroditarum hinc donatis arifta longiflma, illinc gemino mucrone breviffimo. *Jacq. coll. vol. 5.*  
Caulis florentis pars fumma. Flos hermaphroditus auctus\* Palea aucta\* Flos faemineus\*

### P O L Y G A M I A     F R U S T R A N E A .

- 591 GORTERIA ASTEROIDES, foliis ellipticis; ferraturis fetaceis. *Liuuufyfi.*  
*Jacq. coll. vol. 1\** Gorteria fruticofa. *Ait. keiv.*  
Ramus florens. Flos hermaphroditus. Idem faemineus.

- 592 RUDBECKIA AMPLEXIFOLIA, foliis Mceōlato-oblongis, ferratis, amplexicaulibus, glabris. *Jacq. coll. vol. 5.*  
Caulis florentis pars fumma.
- 593 RUDBECKIA ALATA, foliis decurrentibus, inferioribus fubpinnatifidis, Tuperioribus integerrimis. *Jacq. colL vol. 5:*  
Caulis pars fuperior. Folium infimum.
- 594 COREOPSIS LIMENSIS> foliis late ovatis, fubintegerrimis, tomentofis, omnibus **alternis**. *Jacq. coll. vol. 2.*  
Cauis florentis pars. Corollulae tres ex radio. Flosculus ex disco auctus. Semen.  
Idem auctum.
- 595 COREOPSIS ARTEMISIAEFOLIA, foliis fubcipinnatis; pinnulis lanceolatis.  
*Jacq. col. vol. 5.*  
Caulis florentis pars fumma. Folium inferius. Flosculus herraphroditus auctus.  
Semen auctum. Idem in Itatu naturalL

### P O L Y 6 A M I A      N E C E S S A R I A ,

- 596 CALENDULA ARBORESCENS, foliis lanceolatis, ferrato-dentatis, fcabris;  
caule arborescente; ramis dejectis. *Jacq. coll. vol. 3. an Calendula rigida. Ait.keiv.*  
Trunci pars cum ramis truncatis. Ramulus florens. Semen\*

### MONOGAMIA.

- 597 LOBELIA SIPHILITIC A, c&ule crcto ; foliis ovato-lanceolatis, fubferratis;  
caljcum finibus refiexis. *Limufyfi. Jacq. coll. voL 1.*  
Tota planta. Flos. Genitalia. Calyx.

### GYNANDRIA.

#### D I A N D R I A.

- 598 ORCHIS MILITARIS, bulbis indivifis ; fpica oblonga; ncctarii labcllo quadridido, lobis integerrimis, disco muricato; petalis confluentibus. *Jacq. coll. vol. 2.*  
*Linn, fyji. Hall. helv.*  
Tota planta. Flos albidus ex varietate. Flos vi expanfus,
- 599 ORCHIS VARIEGATA, bulbis indivifis; fpica brevi/nectarii labello quadridido, lobis apice ferratis, disco glabro ; petaks confluentibus. *Jacq. coll. vol. 2.*  
*Ball. helv. Orchis pratenfis elatior floribus variegatis. Seg. ver. Bauh. prodr.*  
**Orchis militaris minor.** *Bupp.jen., Vaill. paris.*  
Plantse tres. Flores duo vi expanfi.
- 600 NEOTTIA SPECIOSA, foUis crenato-undulafe ; fpathis fuperioribus rubris.  
*Jacq. coll. vol. 3.*  
Tota planta. Petalum dorfale exterius. Petalum exterius laterale. Flos totus.  
Flos poll avulfa petala exteriora. GenitaUa cum Nectario. Gcnitaiia fola. Appendix ftigmatis aucta. Hac avulfa Stylus & vi reflexum Filamentum cum Anthēus aucta. Stylus, Stigma & Filamentum aucta
- 601 NEOTTIA MINOR, foliis planis ; fpathis viuentibus. *Jacq. coll vol. 3.*  
Tota planta. Flos auctus cum petalo exteriori dorfali & duobus interioribus expanfis. Genitalia aucta cum filamento vi retrorfum ducto. Capfula.

**602 LIMODORUM ALTUM** (*/tec TUBEROSUM*), floribus imberbibus; fpicis paniculatis, ex bulbi latere ortis. *Ait. hew. Linn, fyft. Jacq. coll. vol. 4.* Helleborine americana, radice tuberosa. *Martyn. cent.* Helleborine radice arundinacea. *Plum, ic*

Tota planta. Petala duo. Genitalia cum nectario. Eadem aucta. Antherae unius filamenti auctae. Capula matura. Semen auctum. Capula immatura.

**603 LIMODORUM DIURNUM**, floribus imberbibus; fpicis paniculatis, ex centro bulbi ortis. *Jacq. coll. vol. 4<*

Tota planta. Flos major. Nectarium expansum. Stylus cum ftigmate auctus.

**604 EPIDENDRUM ELONGATUM**, caule erecto; foliis oblongis; floribus fpi catis ) nectarii labio trifido, lacinia media biloba. *Jacq. coll. vol. 3..*

Tota planta florens. Flos auctus. Poll florescentiam ex fpathis caulis summi aphylli excrescunt novae plantulae, radicibus propriis donatae, quae tandem in terrain decidunt, novasque plantas producunt. Talis foboles confpicitur duplex *in figura laterali*.

**605 EPIDENDRUM COCHL ATUM**, foliis oblongis, geminis, glabris, ftriatis, bulbo innatis j fcapo multifloro; nectario cordato. *Linn. fyji. Jacq. coll. vol. 3.* Helleborine cochleato flore. *Plum. ic.* Vifcum caryophylloides, lilii albi foliis. *Catesb. carol.* Vifcum radice bulbofa minus, delphinii flore. *Sloan, jam.*

Tota planta florens. Nectarii pars fuperior. Aucta Stamen, Cucullus quadrilocularis, & compages Styli Stigmatis Nectarii & Cuculli.

## PENTANDRIA.

**606 PASSIFLORA CUPREA**, foliis indivifis, ovatis, integerrimis, trinerviis, fubtus inter nervos punctatisj petiolis eglandulofis. *Linn. fyfi. Jacq. coll. vol. 3.* *Dill, elth. Mart. cent. Catesb. car.*

Ramus florens & fructescens. Semen auctum. Idem naturale. Foliorum varietates.

**607 PASSIFLORA LUTEA**, foliis trilobis, cordatis, obtufis, integerrimis, glabris, eglandulofis ; petiolis hirfutis. *Linn. Jyji. Jacq. coll. vol. 2.* *Gron. virg. Brown, jam.*  
Ramus florens.

## HEXANDRIA.

**(308 ARISTOLOCHIA BARBATA**, foliis cordatis cum fmu profundo obtufoque'; caule volubili, fruticofo; corollae infundibiliformis appendice obverfe ovata, fupa medium antice hirfuta. *Jacq. coll. vol. 3.*

Ramus florens.

## POLYANDRU

**609 POTHOS CRASSINERVIA**, foliis lanceolatis, acuminatis, integerrimis, totis crectis, nervo medio utrinque protuberante & angulato, casterum enerviis. *Jacq. coll. vol. 4.*

Tota planta cum truncatis plurimis foliis. Aucta Flosculu\*, Germen, & unum Petalum cum inclufo Stamine. Spatha expanfa.

**610 POTHOS GRANDIFOLIA**, acaulis; foliis cordatis. *Jacq. coll. vol. 4,*  
Tota planta cum truncatis plerisque foliis fpadicibusque. Folium majus. Spatha expanfa. Spadix. Aucta Flosculus, Petalum unum cum Stamine, & Germen.

- 6n POTHOS DIGITATA (/a\* PALM ATA), foiiis digitalis. *Jacq. coll. vol.* ±  
Caulis pars. Aucta Germen, unum Petalum cum Stamine, & totus Flosculus cum  
expulfo Polline.
- 612 DRACONTIUM LANCEAEFOLIUM, foliis lanceolatis; fpatha viridi, fpa-  
dicem multoties fuperante. *Jacq. coll. vol.* 4.  
Plantae pars inferior, Scapi floreritis pars fuperior. Aucta Germen, unum Peta-  
lum Cum Stamine, & Flosculus totuS.
- \$13 ARUM HELLEBORIFOLIUM, acaule; foliis pedatis ; foliolis lanceolatis\* fu-  
perantibus fpatham fpadici fubaequalem. *Jdcq. coll. vol.* 3.  
Tota planta. Spadix fpatha exemptus. Spatha.

## MONOECIA.

### DiANDRIA.

- 6H DORSTENIA CONTRAJERVA, fcapis radicatis; foliis lobatis, ferratis,recepta-  
cuiisquadrangulis. *Linn.fyft. Jacq.coll. vol.* 3. Dorfteniafpiondyliifolio.*Plum.ic.*  
Tota planta florens & fructescens triennis. Plantula cum foliis foliis feminalibus.  
Eadem menfem agens. Receptaculi fructescens aucti portio. Receptaculi floren-  
tis & aucti pars dimidia. Auctae Capfula, fuperne feminifera, inferne vacua 5 ea-  
dem bivalvis cum femine glabro 5 & duse fteriles.

## TRIANDRU.

- 613 CAREX HERMAPHRODITA, floribus hermaphroditis; *Jdcq. coti. vol.* 4.  
Totaplanta,radice excepta. Spicula florens, eadem fructescens,Piftillum ; omnia aucta
- 616 PHYLLANTHUS SPECIOSA, foliis pinnatis ; foliolis ferrulatis, margine fl,l  
xiferis ; floribus pedunculatis ; caule arborescente\* *Jacq. coll. vol.* 2. Phyllanthus  
Epiphyllanthus. *Limufpec\* pi.* Xjlophylla latifolia. *Linnfyft. Ait.kew* Swartz\*  
prodr: anXjlophjlla arbuscula. Swartz. prodr. ? Certe hujus determinatio fpecifici-  
Ci exakte congruit.
- Trunciparsfumma. Flos fcemineus & masculus auoti. Semen. Pars folioli fruc-  
tescentis\*

## TETRANDRIA.

(f^MORUSMAURITIANA.foms lanceolata longis, utrinque ^uaUbus , intc  
gdrrimis. *Jacq. coll. vol.* 3. et *vol.* 4. pag. 224.

Ramusf^mineus florens. Spica mdscula. Mas auctus. Faemina aucta.

**82 2 11 ~ 610** *Jacq. co* **POLYANDRIA.**  
foliis dentatis, glabris ; floribus

Ramus florens. Capfula.

- 619 BEGONIA DICHOTOMA, caule fruticofo, erectivsculo ; foliis dentatis, sub-  
dichotoma, *Jacq. coll. vol.* 3.  
Ramus florens. Flores aueti. Stigma auctum. Anther\* auctae.

## MONADELPHIA.

- 620 ACALYPHA ALOPECUROIDEA, involucris faemineis femljtrilobis, *lohis Ian-*  
*ceolatis, in fetam terminate; fpicis erectis; fpicis maſculis minuti\**; foliis ovato-  
*cordatis,- ferratis, hirfutis. Jacq. coll. vol. 3.*
- Caulis pars fuperior. Aucta Capfula, Semen, Faemina, Involucrum, Mas;
- 621 CROTON PUNCTATUM, to turn ftellato-tomentofum; foliis ovatis, integerri-  
*mis. Jacq. coll. voL i;*
- Ramus florens. Aucta Faemina, Mas, & portio Folii.
- 622 CROTON PUNGENS, foliis cordato-aCuminatis, obfolete ferratis, facie aspe-  
*n̄s, dorfo ftellato-tomentofis. Jacq. coll. vol. 4.*
- Ramus florens. Mas auctus, dorfo & facie fpectandus. Folii portio dorfalis aucta.
- 6t\$ IATROPHA GOSSYPIFOLIA, foliis trilobis, rarius quinquelobis 5 lobis ova-  
*tis, ferratis; fetis ramofis, glandulofis. Linn. fyji. Jacq. coll. vol. 1. Brown, jam.*  
*Ricinus itaphisagriae folio. Sloaii. hifi. Coirim. /tort. Merian.fur.*
- Ramus florens & fructescens. Maris Calyx. Idem auctus. Ejusdem Stamina cum  
*glandulis.' Calyx hermaphroditi. Idem auctus. Ejusdem Genitalia aucta. Capfula.*  
*Semen.*
- 624 TRICHOSANTHES FOETIDISSIMA, pomis ovatis, acutis; foliis cordatis,  
*denticulatis, hirfutis, viscofis, venofis, enerviis. Jacq. coll. vol. 2.*
- Ramus florens & fructescens. Radix. Stylus auctus. Semen. Corollae lacinia auc-  
*ta. Stamen auctum.*

## DIOECIA, TETRANDRIA.

- 625 MYRICA SEGREGATA, foliis lanceolatis, integerrimis> venofis 5 floribus fe-  
*gregatis. Jacq. coll. vol. 2.*
- Ramus maſculus florens. Racemus auctus. Ramulus faemineus florens. Racemus  
*auctus. Bacca. Semen. Nucleus.*

## HE X AND R IA,

- 626 DIOSCORISA VILLOSA, foliis cordatis , alternis oppofitisque; caule laevi  
*Linn. fyft<, Jacq. coll. vol. 2. Brjoniae fimilis floridana. Plukn, aim.*
- Radix cum Caulis parte infima. Ramus masculus florens. Semina. Mas auctus.
- 627 DJOSCOREA TRIPHYLLA, foliis ternatis- *Linn. fyfi. Jacq. coll, vol. 2.*  
*Ubium fylveſtre. Hump, amb. Tsjageri-Nuren. Rheed. jnal.*
- Radix. Caulis faeminei florentis pars infima. Flos auctus. Semen.

## DECANDRIA,

- 628 KIGGELARIA INTEGRIFOLIA, foliis integerrimis. *Sacq; coll. vol. 2.*
- Ramus florens. Antherae auctae. Pifillum auctum. Poll editum Hortum kewenfem  
& denuo examinatos flores, quos inveni ob bafin corollae connatam vere elfe mo-  
nopetalos , nee pentapetalos, dubius fum, an non fit hujus Horti Royena po-  
ly andra.

## MONADELPHIA,

- 629 CISSAMPELOS SMILACINA, foliis cordatiis\* acutis, angulatis. *Linn;fyfi.*  
*Jacq. coll. vol. 4.* Smilax lenis. *Catesb. car\**  
 Ramulus mafculus florens. Calyx auctus. Flos auctus, calice avulfo. Stamen auctum.

## POLYGAMIA.

### MONOECIA.

- 630 ANDROPOGON DISTACHYOS, fpicis binis, terminalibus 5 culmo indivifo. *Linn;fyft.* *Jacq. coll. vol. 1.* *Ger;prov.* Gramen bicorne. *Bocc.Jic.*  
 Tota planta florens. Mas auctus. Flos hermaphroditus expansus cum mate convoluto.
- 631 ANDROPOGON UNDATUS, fpica pinnata j floribus fecundis, *Jacq. coll. vol. 3.*  
 Culmus florens. Flos hermaphroditus & mafculus aucti.
- 632 MIMOSA CARACASANA, inermis; foliis bipinnatis, partialibus subquinquejugis, propriis multijugis; caule glabro; calycibus ad oras glandulofis. *Jacq. noil. vol. 4.*  
 Ramulus floreris. Calyx auctus cum Corolla.
- 633 MIMOSA PORTORICENSIS, inermis; foliis bipinnatis, partialibus subquinquejugis, prdpriis multijugis) caule glabro) calycibus ad oras ciliatis. *Jacq. coll. vol. 4.*  
 Ramus florens. Calyx auctus' cum Corolla.

### DIOECIA.

- 634 PAN AX ACULEATUM, foliis ternatis; pettolis ramulisque aculeatis; caule fruticofo. *Ait. kew.* *Jacq. coll. vol. 4.* Zanthoxylum trifoliatum. *Linn;fyft.*  
 Ramuy florens hermaphroditus. Flos auctus. Calyx auctus.

## CRYPTOGAMIA,

### FILICES.

- 635 ZAMIA INTEGRIFOLIA, foliolis subintegerrimis, obtusculis, muticis rectis, nitidis ; flipite inermi. *AH. keu>*. *Jaq. coll. vol. 3.* Zamia pumila. *Linn frfi*  
 Tuta planta feminea fructescens. Spadix mafculus. Ejus duo Scuta. Scutum auctum cum Antheris. Anthera aucta bivalvis cum contento Polline cohaerente In alio latere Drupa matura. Nux. Scutum auctum cum geminis Germinibus Spadixfaemineus. Idem transverfim fectus. Idem jam maturior, longitudinaliter fectus
- 636 ZAMIA ANGUSTIFOLIA, foliolis linearibus, muticis L, aJice ^ H ^ l ginatis > ftipite tereti, inermi. *Jacq. coll. vol. 3.*
- Tota planta faeminea florens. Spadix fructiferus transverfe diftectus. Drupa Nux transverfe fecta. Eadem Integra. Spadix fructescens faemineus. In alio late're Foliorum apices aucti. Scutum antheriferum auctum. Spadix masculus. Ejusdem fectio transverfa aucta.

**637 POLYPODIUM PHYMATODES**, frondibus fimplicibus pinnatifidisque, lanceolatis, supra fructificationes in facie verrucosae. *Linn. fyft. Jacq. coll. vol. L4. Burnt, thes. Plakn. phytoogr.*

Tota planta. Folioli pars dorfalis cum fructificationibus.

**638 POLYPODIUM TRIFOLIATUM**, frondibus ternatis, finuato-lobatis, intermedia majore. *Linn. fyft. Jacq. coll. vol. 3. Hemionitis maxima trifolia. Plum, amer. etjil. Shan. hist.*

Tota planta. Fructificatio matura eademque junior auctse.

**639 POLYPODIUM FRAXINIFOLIUM**, frondibus pinnatis; foliolis lanceolatis, acuminatis, repandis, undulatis, diftantibus. *Jacq\* coll. vol. 3.*

Tota planta. Foliolum ex dorfo confpectum.

**640 POLYPODIUM MOLLE**, frondibus fubpinnatis, villofis, mollibus; foliolis fublanceolatis, acuminatis, fubpinnatifidis j lobulis integerrimis, obtufis. *Jacq. coll. vol. 3.*

Tota planta. Foliolum dorfale. Hujus portio aucta.

**641 POLYPODIUM HIPPOCREPIS**, frondibus fubpinnatis; foliolis lanceolato-oblongis, acuminatis, pinnatifide & obtufe incifis; superioribus confluentibus; floribus lunatis. *Jacq. coll. vol. 3.*

Tota planta. Fructificatio junior, dein magis provecta, tertio adulta, omnes auctse.

**642 POLYPODIUM ALPINUM**, frondibus alterne tripinnatis; pinnuiis oblongis, vage incifis; lacinulis obtufis, apice breviulme bifidis. *Jacq. coll. vol. 2. Seg. ver, Hall. help.*

Tota planta. *Wulfen.*

**643 POLYPODIUM MULTIFIDUM**, frondibus tripinnatis, glabris; pinnuiis oblongis, pinnatifide incifis in lobulos oblongos superne incifos & acuminatos. *Jacq. coll. vol. 3.*

Tota fere planta. Portio pinnae dorfalis.

**644 BLECHNUM OCCIDENTALE**, frondibus pinnatis; pinnis lanceolatis, oppositis, bafi emarginatis. *Linn. fyft. Jacq. coll. vol. 3. Brown, jam. Lonchitis juxta nervum pulverulenta. Plum. Jil. et amer.*

Tota planta. Pinnae dorfalis segmentum auctum.

**645 PTERIS CAUDATA**, frondibus supradecompositis, pinnuiis linearibus; terminalibus longissimis. *Linn. Jyft. Jacq. coll. vol. 2.*

Planta tota. Pinnae pars dorfalis. Pinnulse dorfalis pars aucta.

**646 ADIANTUM STRIATUM**, fronde bipinnata; pinnuiis rigidis, faicato-ovatis, illriatis; fructificationibus superioribus interruptis; llipite tereti, aspero. *Swartz. prodr. Jacq. coll. vol. 3.*

Tota planta. Pinnulse dorfales. Pinnula dorfalis aucta.

## ALGAE.

**647 TREMELLA JUNIPERINA**, feffilis, membranacea, auriformis, fulva. *Linn, fyft. Jacq. coll. vol. 2. Scop. earn. Leers, herb. Gunn. norv. Gmelfuc.*

Plantae plures. *Wulfen.*

**648 TREMELLA CLAVARIAEFORMIS**, gregaria, fimplex, compreffo-fubulata, subpyramidalibicornutave, aurantiaca. *Jacq. coll. vol. 2.*

Plantae plurimas. *Wulfen.*

# I N D E X

## PIANTARUM, IN TRIBUS VOLUMINIBUS CONTENTARUM.

<b>A</b>	
* <i>calyptra aiopae</i> Curoideiu	Tab. 620
<i>Aconitum tauricum.</i>	492
<i>Adiantum triatum.</i>	646
<i>Aechmonea bifinosa.</i>	564
<i>Agave foetida.</i>	379
— <i>virginica.</i>	378
<i>Agrostis tenaciflora</i> #	16
<i>Albuca abylinica.</i>	64
— <i>altillima.</i>	63
— <i>aurea.</i>	44
— <i>caudata.</i>	442
— <i>fuccida.</i>	444
— <i>major.</i>	443
— <i>petraea.</i>	440
— <i>spiralis.</i>	439
— <i>viridiflora</i> *	446
— <i>visroid.</i>	4 <sup>1</sup> 5
<i>Alliun illyricum</i>	365
— <i>obliquum.</i>	62
— <i>fuaveolens.</i>	364
— <i>triatum.</i>	3 <sup>1</sup> 6
<i>Alpinia comofa.</i>	602
<i>Althaea narbonensis</i> ^	138
<i>Alyssum gemonense</i> *	S <sup>o</sup> 3
<i>Amaryllis longifolia</i> ^	36a
<i>Andromeda coriacea.</i>	465
— — <i>lucida.</i>	79
<i>Andropogon diffachyos</i> *	1-3,0
— — <i>undatus.</i>	631
<i>Anemone fragifera.</i>	103
<i>Anthericum bipedunculatum.</i>	410
— — <i>exuviatum.</i>	415
— — <i>filibium.</i>	414
— — <i>flexibium.</i>	413
— — <i>hippidium.</i>	409
— — <i>Ian. folium.</i>	4C8
— — <i>longifolium.</i>	4 <sup>o</sup> 3
— — <i>longicupina.</i>	464
— — <i>nutans.</i>	407
— — <i>phydodes.</i>	418
— — <i>pilofum.</i>	4.6
— — <i>praemorum</i> *	406
— — <i>pugioniforme</i> *	4@5
— — <i>puliflum.</i>	4 <sup>o</sup> 7
— — <i>rothauin.</i>	403
— — <i>fubrigynuuu</i>	4 <sup>o</sup> 9
— — <i>undulatum.</i>	411
<i>Antirrhinum hirtum:</i>	117
— — <i>parviflorum</i> *	499
— — <i>verficolor.</i>	116
<i>Aquilegia viridiflora.</i>	102
<i>Arabis ovirensis.</i>	iay
<i>Atenaria biflora.</i>	8?
<i>Aristolochia barbata.</i>	603
— — <i>bilobata.</i>	188
<i>Arnica ghcialis.</i>	586
<i>Artemisia hispanica.</i>	172
<i>Arum helleborifoliumt</i>	613
<i>Afclepias citrifolia.</i>	343
<i>Afplenium anguitifolium</i> *	199
<i>Afragalus asper.</i>	.153
— — <i>excapns</i>	56i
— — <i>hians.</i>	153
— — <i>leontinus.</i>	>54
— — <i>uralensis.</i> ^	155
<i>Athamanta Matthioli,</i>	57
<b>C</b>	
<i>Avena afteriti&amp;</i>	23
<i>Axyris ceratoide&amp;</i>	19
<i>Begonia dichotoma.</i>	619
— — <i>minor.</i>	618
<i>Blechnum occidental</i> ^	644
<i>Borago zeylanica.</i>	3,4
<i>Bromelia humilis</i> *	1
<i>Buddleia capitata.</i>	237
<i>Bupleurum arborescent</i>	236
<i>Cacalia laciniata.</i>	168
— — <i>peucedanifolia</i> «	581
— — <i>villofa</i> *	58*
<i>Calea afpera.</i>	583
<i>Calendula arborecen</i> ^	596
<i>Camellia japonic</i> **	553
<i>Campanula Zoyfii</i> #	334
<i>Carduus acaulis.</i>	166
— — <i>arabisus.</i>	615
<i>Carex hermaphrodita</i> *	73
<i>Caffia chinensis</i> *	74
— — <i>Crifta.</i>	73
— — <i>multiglandulosa</i> ^	46D
— — <i>polyp&gt;hylla.</i>	70
— — <i>russifolia»</i>	459
— — <i>fennoides.</i>	5'
— — <i>fenfitiva.</i>	3 <sup>1</sup> 9
<i>Celosia procumbent</i>	57
— — <i>virgara.</i>	497
<i>Celtia linearis.</i>	»78
<i>CentaUrea Venitnm</i> *	344
<i>C^ienop odium caudatum</i> *	345
— — <i>guineenfe</i> *	629
<i>Chirrocoma dichotoma.</i>	34
<i>Ciliampelos filiacina.</i>	97
<i>Ciftus canal-lends,</i>	95
— — <i>cretenfis.</i>	98
— — <i>foetidus.</i>	99
— — <i>inutabilis</i> *	96
— — <i>friaciac.</i>	501
<i>Citharexylum erectunl</i> ^	118
— — <i>villofum</i> *	194
<i>Clematis angustifolia.</i>	294
<i>Commelinia lonicaulis</i> ^	293
— — <i>mollis.</i>	315
<i>Convolvulus crenatus,</i>	34
— — <i>floridns.</i>	S16
— — <i>pentanthus.</i>	585
<i>Conzya carolinensis.</i>	59?
<i>Coreopsis artemisiifolisu</i>	594
— — <i>limenfis.</i>	32
<i>Cortufa Mathiolu</i>	I
— — <i>Mathiolu</i>	504
<i>Cotrus arabicus</i>	128
<i>Crambe filiformis.</i>	129
— — <i>orientalis</i> #	129
— — <i>Tataria.</i>	129
<i>Craffula Umbelta.</i>	352
<i>Crepis albida.</i>	—
<i>Crinum tenellunt</i> #	—
<i>Crotalaria caerulea.</i>	—
<i>Croton glandulofum</i> *	144
— — <i>punctatum.</i>	194
— — <i>pungens.</i>	63 c
<i>Cyanella rchidiformk</i>	(2 i
<i>Cynanchum caroliqnlb</i>	4 <sup>1</sup> 7
	747
<b>Cynanchum extenfum.</b>	54
— — <i>monpeliacum.</i>	340
— — <i>obliquuin.</i>	341
<i>Cynofurus coenileus.</i>	21
— — <i>domingenfis.</i>	29
— — <i>fphaerocephalus</i> *	20
<i>Cyperns alternifluius.</i>	298
— — <i>diftans,</i>	209
— — <i>longus.</i> \	297
— — <i>tenuiflorus.</i>	296
— — <i>vifefus.</i>	295
<i>Cyrilla racemiflora.</i>	47
<i>Cytifus tener.</i>	147
<i>Dais laurifolia</i> *	77
<i>Delphinium urceolatuttl.</i>	to 1
<i>Dianthus atrorubens.</i>	467
— — <i>fylvefris.</i>	82
<i>Diodia virginica.</i>	29
<i>Dioscorea triphylla.</i>	627
— — <i>villofa»</i>	626
<i>Dolichos acinaciformis.</i>	559
— — <i>rlorius</i> ,	560
— — "Sofj	145
<i>Dorftenia Contrajerva.</i>	614
<i>Dracaena terminalis.</i>	448
<i>Dracocephalum auftriacum.</i>	112
<i>Dracontium Janceaefolium.</i>	612
<i>Drimia ciliaris.</i>	377
— — <i>elata.</i>	573
— — <i>media.</i>	375
— — <i>pufilla</i> *	374
— — <i>undulata</i> #	376
<i>Duranta Plumieri</i> *	S02
<i>Echites domingenfis</i> *	54
<i>Echium candicans.</i>	50
— — <i>glaucophyllunu</i>	318
<i>Elaeodendron orientale.</i>	48
<i>Elymus Hyfrrix.</i>	3oy
<i>Epidendrum cochleatum</i> *	ft*
— — <i>elongatum.</i>	60+
<i>Erioppermum latifolium.</i>	420
— — <i>lanceafolium</i> ^	421
— — <i>parvifolium.</i>	422
<i>Erodium ribifolium.</i>	509
— — <i>trilobatum.</i>	508
<i>Eryngium aquanticum.</i>	347
— — <i>alpinum.</i>	5?
<i>Eucomis bifolia.</i>	440
<i>Eugenia brennfis.</i>	486
<i>Eupatorium myototi folium.</i>	*R2
— — <i>fcandens.</i>	167
— — <i>friacum.</i>	170
<i>Euphorbia angulata.</i>	481
— — <i>Characias.</i>	89
— — <i>Clava.</i>	85
— — <i>cyathophora.</i>	48>
— — <i>diffusa.</i>	88
— — <i>divauicata.</i>	K
— — <i>linifolia.</i>	8*
— — <i>literata.</i>	482
— — <i>nicaenfis</i>	485
— — <i>mldiflora,</i>	479
— — <i>picta.</i>	477
— — <i>pilulifera</i> *	473
	Euphor-

Euphorbia puuicea.	484	Hypoxis obliqua.	37	Lithoppermum tenuiflorum.
— — ffordifolia,	476	— plicata.	367	Lobelia liphilitica.
— — ferrata.	483	— ferrata*	369	Lonicera tartarica.
Fothergilla Gardeni.	100	— fololifera.	\$72	Lopezia mexicana.
Fumaria acaulis.	554	— ftellata.	368	Lychais granditiora.
Galaxia ovata.	291	— villofa.	370	Manulea tomentoia.
— — plicata,	292	Hyptis capitata.	114	Mangifera indica#
Galega capensis*	574	— verticillata.	IH	Malva balfamica,
— — cinerca.	575	Jatropha goffipifolia.	623	— icoparia.
— — dubia*	576	Ilex ligustrina.	310	Malachra alceaefolia.
— — filiformis*	573	Indigofera dendroides.	57*	— fasciata.
— — longifolia,	572	— — hendecaphylla*	570	Malpighia coccigern*
— — ochroleuca.	150	— — hirtuta.	569	— glandulifera.
Gaura fruticosa.	457	Ipomoea anguitifolia.	317	Maranta lutea.
Genista hispanica*	557	— hederacea,	36	Marrubium afracanicum.
— — fericea.	556	— leucantha*	318	Medioago cat-ttienis.
Geranium argenteum.	546	— luteola.	3?	Melanthium eucomoides.
— — macrorrhizum.	134	— — peftapphylla.	319	— — junceum*
Geum aleppicum.	93	Iris flavillraia.	200	— — unifloruin.
— — hybridum	94	— tripetala.	221	Melochia caracafana.
Gladiolus alatus.	259	— tricuppis.	222	Mefembryanthemum cordifolium.
— — albibus.	256	— virginica.	—	— — — cuneifolium.
— — *anceps.	569	Juglans cinerea*	483	— — — — — polycladum.
— — angustus.	240	— nigra.	192	Mimosa caracafana.
— — bicolor.	249	Jufticia bra&eolat*.	I, I	— portoricensis.
— — brevifolius.	25?	— — caracafana*	205	— speciofa.
— — carnitis,	267	— — pulcerrima*	206	Morus mauritiana,
— — crifpus.	257	Ixia anenonaeflora.	204	Moraea collina.
— — cupidatus.	268	— angulta.	*73	— palmifolia.
— — fissifolias.	254	Bulbocodium	079	— fordescens.
— — floribundus.	258	— chloroleuca.	>71	— vegeta.
— — — — —	246	— corymbofa.	272	— virgata.
— — — — —	236	— cruciata,	288	Myagrum arborefcens,
— — — — —	250	— falcata.	29a	Mygindia Rhacomiu
— — — — —	242	— fenefrata*	276	311
— — — — —	254	— fragrans*	289	Myrica fegvegata.
— — — — —	234	— incarnata,	^74	Neottia minor,
— — — — —	235	— lancea.	282	— speciofa.
— — — — —	232	— leucantha.	281	Nerium coronanum.
— — — — —	262	— polyftachy^	57a	Ochrolia maculata
— — — — —	263	— punicea.	275	Ocymuui gratiflimuiii.
— — — — —	230	— purpurea,	Q87	Oenothera nocturna.
— — — — —	253	— — — — Varietas.	286*	— — odorata*
— — — — —	237	— radiata.	280	Onopordon acuale,
— — — — —	238	— rubrocyannea*	285	Ophrys crucigera.
— — — — —	247	— fecunda.	277	— myoies.
— — — — —	241	— uniflora.	283	Orchis maschii
— — — — —	261	— villofa.	284	— militaris*
— — — — —	270	Kiggelaria integrifolia.	628	— mbravica.
— — — — —	260	Kyllinga incompleta.	300	— palufris.
— — — — —	239	Lachenalia anguifolia^	381	— rubra.
— — — — —	^48	— — — — —	391	— variegata.
— — — — —	243	— — — — —	382	Ornithogalum caudatum*
— — — — —	244	— — — — —	401	— — — coarftacnm.
— — — — —	—*	— — — — —	402	— — — conicum.
— — — — —	—*	— — — — —	387	— — — flavefcens
— — — — —	264	— — — — —	395	— — — fuscatuin.
— — — — —	265	— — — — —	392	— — — latifolium.
— — — — —	266	— — — — —	390	— — — latifolium.
— — — — —	229	— — — — —	38?	— — — ininitiatuin.
— — — — —	251	— — — — —	384	— — — inodoratum.
— — — — —	233	— — — — —	400	— — — poly phy Hum.
Glyciie caribaea.	146	— — — — —	397	— — — pyramidale.
Gomphrena brafilienis.	546	— — — — —	393	— — — fecunduin.
Gorteria afteroides.	591	— — — — —	388	— — — fuaveolens.
Gronovia scandens.	591	— — — — —	38;	— — — tenellum.
Guilandina Moringa.	10?	— — — — —	386	Olleoppenum coeruleum.
Halaragis alata.	69	— — — — —	396	Oxalis multiflora.
Hamelia chryfantha.	— ^	— — — — —	398	— — — pblyphylla,
Hedyfarum gyrans.	565	— — — — —	399	— — — rubella.
— — muricatuin,	568	— — — — —	61	Pdnax aculeatum.
— — pictum.	567	— — — — —	389	Panicum coloratum.
— — vepertilionis.	566	— — — — —	394	— — — maximum.
Heliophila integrifolia,	506	— — — — —	162	Pafpalum racemolum.
Helonias pumila.	453	Laftuca intybacea.	58	— — — virgatum.
Hibiscus diverfrblius,	551	Laferpitium Archangelica.	349	PafTiflora cuprea.
— — doiningenis.	550	— — — peucedanoides.	106	— — — incarnata.
— — pentacarpos.	'43	Lavandula pinnata.	4^4	— — — lutea.
— — praemorfus*	141	Ledum latifolium.	588	— — — rubra,
— — virgin icns.	142	Leyfera gnaphalodes.	353	Paullinia caulinflora.
Hieracium incarnatum,	578	Linum africanuin.	602	Pavonia urens.
— — faxatile.	163	Limodorum ahum.	603	Pedicularis rofea.
Hippocrepis balearica.	149	— — — diurmum*	463	Felargoniuin aneinonaeolim.
Hyacinthus viridis.	66	— — — Limonia trifoliata.	33	— — — aftragaloliuni,
Hypecoum littorale.	309	Lidanthus Macrifo		
Hypocheiris helvetica.	16j	tu		

Pelargonium barbatum,	\$13 Poa difticha.	19 Solanum aculeatissimum.	41
— — balfatneuma	543 — perniciosa	18 — aggregatum.	323
— — betonicum.	53 k — icula.	— carolinense.	331
— — bullatum*	530 Polyantnes pygmaea*	— coccineum.	43
— — carneum.	512 Potentilla africana	— corymbosum,	40
— — caudifolium.	529 & — 'opaca.	— diphyllum,	322
— — chamaedryfoliutn.	523 Falisburgensis*	— fugax.	324
— — ciliatum.	519 **. — fubacaulis.	— fuscatum.	43
— — coriandrifolium.	528 Polypodium alpinum.	— giganteum,	328
— — coronopi folium*	526 — — fraxinifolium	— lanceaeolum.	329
— — comifaetbiura.	539 — — Hippocratea.	— lycoidea.	46
— — daucoides,	522 — — triolle*	— marginatum.	45
— — depreium.	520 — — multifidum.	— Milleri.	320
— — fuscatum.	540 — — phymatodes.	— nodiflorum*	326
— — glutinofum.	131 — — trifoliatum	— peruvianum.	327
— — hermanniaeblum.	545 Pothos craffinervii.	— petellatum.	32c
— — heterophyllum,	516 — digitata.	— framonioliuim.	44
— — hirtum.	536 — grandifolia.	— virginianum.	r12
— — lacerum.	532 Primus Ohamaeerafus*	Sonchus fruticosus.	161
— — longicaule.	533 Pteris caudata.	Spananthus paniculata,	350
— — longiflorum	521 Pforalea bipedunculata.	Spartium decumbens*	555
— — longifolium.	518 — emphydotes.	Spermacoce hirta.	308
— — melananthon*	514 Ranunculus canadenfis*	Spilanthes exasperata.	584
— — multieaule.	534 Refeda fruticulosa.	Stachys canariensis.	108
— — nervifolium.	517 — mediterranea	— lanattu	107
— — oetiotherae#	525 Rhamnus volubilis*	— mgofa^	409
— — pafulum.	541 Rhododendrum ponticum!	Stellaria bulbifera.	468
— — rapaceum.	510 Ribes petraeum.	Stellera Paflerina.	68
— — revolutum.	133 Ricinus inermis.	Strumaria angustifolia.	359
— — ribifoliunu	538 — lividus.	196 — filifolia.	t
— — fabraru	542 Robinia veicaria	148 — linguaefolia*	356
— — ternatinu.	544 Rudbeckia alata.	493 — rubella.	358
— — tetragonum.	132 — — amplexifolza.	592 — — troncata.	357
— — tomentolunu	537 Rubia fruticosa.	25 — — undulata#	360
— — trichostemon.	524 Ruellia patula.	Tabernaemontana perficariaefolia,	
— — triphyllum.	515 Rumex glaucus.	76 — —	320
— — violareunk	527 Ruta legitima*	Terminalia Catappa.	197
frhaca alpina.	151 Salvia abyfinica.	Thlaspi alliaceum	321
Phafeolus femiereftus»	558 — pseudococcinea.	Tournefortia cymofo.	
Phleum afperum.	U — ferotina*	3 —	31
— — Gerardi.	* 15 —	7 — & multiflora.	354
— — fchoenoides	110 — Viscofa.	Trajzia involucrata.	355
Phloinis caribaea.	in Sambucus racemosa.	5 Tragopogon capenfis.	190
— — zeylanica.	616 Sanicula marilandica.	59 — — mutabilis.	575
Phyllanthus speciofa.	39 Satureja rupestris.	348 — — porrifolius.	157
Phyfalys barbadensis*	38 Saxifragamata.	494 — — undulatus.	159
— — profrata.	33^ — petraea.	466 Tremella clavariaformis.	158
Phyteuina hemiphaerica*	193 — — ftoloniifera.	81 — — juniperina.	648
Pinus Mughus.	a 10 Scabiofa monspeliensis.	80 Tribulus maximus.	647
Piper aduncum.	218 Sderocarpus africanus.	4 Triglochin bulbofui.	462
— blandum.	212 Schoenus umbel lat us.	176 Trichomanes canariense.	coo
— clufiaefolium.	214 Schotia speciofa*	10 Trichofanthes foetidiflora.	^24
— cuneifolium.	213 Scilla hyacinthoides.	Turritis hirsuta.	126
— magnoliaefoliura.	51S Scorzonera taraxacifolia.	8 Verbena mutabilis.	207
— inarcinatum.	8 Selago fasciculata.	160 Verbenina gigantea%	208
— medium.	9 Senecio graminifolius.	496 Vitis vinifera.	*7)
— obtusifolium.	219 — rosmarinifolius*	*74 Volkameria Kaempferi	50
— pereskiaefoliunt	217 Sida atrofanguinea.	587 Waltheria indica.	500
— fstellatum.	211 — carpinifolia.	136 Wulfenia carinthiaca	130
— tuberculatum.	216 — mauritiana.	135 Xeranthemum fulgidum	2
— umbellatum.	23 — palmata.	137 Zamia angustifolia.	173
Plantago aegyptiaca.	27 Sinapis Millefolia.	547 — integrifolia.	636
— Cornutu	26 Sifymbrium hispanicum.	127 Zinnia elegans.	6^5
— maxima.	306 — — lippicente.	124 — tenuiflora.	589
— pangonica*	17 — molle.	505 Zoegea aleppica-	590
Poa abyffinica.	304 — — pannonicum.	x23	177
— ciliaris.			

2615



*Rudbeckia* *multiflora*  
Jacq. tunc I.H.A.



*Gaura fruticosa*  
Tug. Coll. vol. 1.



*Paullinia caudiflora.*  
Jacq. Coll. no 114



*Cassia sensitiva*  
Jacq coll vol 2

Jah. 158




*Cassia polypylla.*  
Jacq. Coll vol. 2.

*Gaudichaudia Monogynia*

Jug. Coll. vol. 1.





*Trichilia maxima L.*

Zeyl. Coll. 17



*Limonia trifoliata*  
Jug Coll vol 3



Tab. 665



Ostryomea mortona deu carinata





< r~yjia/i /ft //,(•</// v >/Y//. m.\*i

Atton. pedem. num. 1545.

3638



*Saururus*  
**Mean a** *bulbosa*.

*Taq. Co. vol. 3*





Tab. 47



*Oxalis hirta.*  
Jacq. Coll. vol. 3.

76. 172



Oxalis hirta Varietas.  
Jacq Coll. vol. 3.

36-170



*Oxalis versicolor*  
Jacq. Coll. vol. 3





*Roseda mediterranea*  
Jacq. Coll vol 1



*Euphorbia scordifolia.*  
Tayg. Coll. vol. 3.



*Euphorbia picta.*  
*Jacq Coll vols.*





*Euphorbia nudiflora*  
Jacq Coll vol 3



*Euphorbia heterophylla.*  
Jacq Coll. vol. 1.



C f' shorbia ang. fata.

Jug. Coll. vol. 2



*Cuphorbia literata.*  
Jacq. telt. vol. 2.



*(jiw/tordux serrata*  
*Troy Galvez*



*Euphorbia pumicea*

Tayl. Coll. vol. 3



C<sup>6</sup> *Cistanche nicaensis*  
Jacq. Coll. vol. 2.



*Eugenia barrenas*  
Aug Coll vol 3

*Mesembryanthemum cordifolium.*

Zap. Coll. vol. 2.






*Mesembryanthemum cuneifolium*  
Jacq. Coll. vol. 2.



*Chrysanthemum parviflorum.*  
Linn. Sp. Pl. p. 542. t. 114.



*Potentilla salisburgensis.*  
Jacq. t o\$. 33 / 1





Sal 39





*Satureja rupestris*  
Jug. Coll. vol. 2.



Oxydium gracissimum  
Jacq Coll vol 2



\*<'f< t&0      A'-<"i    v//,(/tf  
'J,,/    tl'tf. *det.*



*Celsia linearis*  
Tay Coll vol 2



*Manulea tomentosa*


Jacq. Coll. vol. 2



*Antirrhinum parviflorum*

, 250 mm. 1000 ft.

Taf. 57





*Citharexylum erodium*  
• T. & J. Coll. vol. 3



Tab. 575



*Alyssum gemonense.*  
Jaq. Coll. vol. 2.

Tab. 554.




*Crambe filiformis*  
Aug Coll vol. 5

Tab. 555



*Sisymbrium lippizense.*  
Jacq. Coll. vol. 2.

Tab. 576



*Heliophila integrifolia* Varietas  
Nag Coll vol 3



Taf. 377.



*Erodium trilobatum*  
Jacq. Coll. vol. 4.



*Erodium ribesfolium.*  
Jug. Coll. 11.5



*Paeonia rapaceum*  
Tayl. Coll. vol. 2



*G-/, Utii/// astragalifolium*  
Jacq coll vol 1

76.572




*Pelargonium carneum*  
Eng Coll vol 1





*Pelargonium melananthum*

Jacq. Coll. vol. 4.



Tab. 518



*Pelargonium heterophyllum*  
Jacq. Coll. vol. 4




*Pelargonium nervosum.*  
Jacq Coll. vol. 4

Jab. 918




*Peltaryonium longistylum.*  
Jacq. Coll. vol. 4








Pelargonium longiflorum  
Jacq. Coll. vol. 4



*Pelargonium dauricum*  
Jacq Coll vol 5



*Idangium chamaephyllum*

Zey. Coll. no. 5







*Erysimum coronarium* var. *pisiolum*.

Jay Coll vol 2



*Pelargonium violaceum*  
Jacq. Coll. vol. 5.





*Claryum cruciatum*.  
Gray t. II. pl. 5





5 Mr. S.  
*Geranium lucidum*



*Pelargonium lacrum.*

Jacq. Coll. vol. 5.



*Clypeum longistylum*


Zeyt. 127. vol. 3.



*Pelargonium multicaule*.  
Gay C. H. ad. s.



*Pelargonium anemonifolium*  
Jacq. Coll. vol. 5.



*Peltaria ludoviciana.*  
*Gouy. t. 117. vol. 3.*

Tab. 117.



*Pelargonium tomentosum*  
Lam. t. 11. v. 2.

Abb. 108



*Pedamomium ribifolium*  
Jacq. Coll. vol. 8




*Pelargonium cortusifolium.*  
Tayl. Coll. vol. 3.



*Pelargonium fuscum*  
Jacq. Coll. vol. 1

*Lobell*



*Pelargonium patulum*  
Berg Coll vol 4

*Uit Uf/P.'.' Utitt HP./?/ "}*



Tab. 523



*Pelargonium balsameum.*  
Jacq. Coll. vol. 8







( f 7, in turn arytntetan  
• Lucy Coll vol

Tab. 537.



*Sida palmata*  
Jacq Coll vol 2

Tab. 548



*Malachnia f*asciata  
Jacq. Coll.



*Malachra alveolata*  
Jacq. cat vol. 1





*Hibiscus dimorphus*

Aug 18 1831



*Pavonia urens.*  
Tay. Coll. vol. 3.

July 555



*Camellia japonica.*  
c Taq Coll vol 1





*Gentia humifusa.*

Jacq. C. M. sol. 2.



Tab. 357



*Genista hispanica.*  
Jacq. Coll. vol. 2





*Dolichos acinaciformis*  
Linn. Coll. vol. 1

Tabl. 11



*Dolichos gladiatus.*  
Jacq. Coll. vol. 2.



*A. pedatum* L. var.  
"Angulatum"



*Loralea bipedunculata.*  
Jacq. Coll vol 4.



*Psoralea phymatoidea*  
Jacq Coll. est. 9



C. hirsutum Schlecht  
Engl. 61. no. 2




*Hedysarum gynandrum*  
Jacq Coll vol 3



*Hedyotis vegetabilis*  
Tasg Coll. vol. 2.

71





*Acalypha mucicatum.*  
Gouy Coll. vol. 3.




*Indigofera hirsuta.*  
*Fag. Coll. vol. 2.*



*Indigofera heterophylla*  
*Jug. Coll. vol. 2.*

Tab. 571




*Indigofera dendroides*  
Jacq. Coll. vol. 2.



*Galega longifolia.*  
Jacq. Coll. vol. 2.



Pr/t-i/ii.jfiltrormtS,  
• n i



*Chelone obliqua*  
Heg. Coll. no. 2.



*Galega cincera.*  
Zieg. Coll. vol. 2.



*Galium dubium.*  
Zey. Cill. 1844.

1657



*Tragopogon capensis*  
Jacq Coll. vol 2.

Joh. S. G.



Hieracium incitatum  
J. & d. Coll. vol 2





*Cacalia villosa*  
Jacq. Coll. vol. 2.




*Cacalia*  
*peucedanifolia*  
Jacq. Coll. ed. 3.

1603




*Eupatorium myosotifolium.*  
Jacq. Coll. vol. 2.





*Spilanthes acutangula*  
Jacq. Coll. vol. 3.




*Conyza carolinensis*  
Jug. coll. vol. 2



Job. 57



*Senecio rosmarinifolius*  
Jacq. Coll. vol. 1. I





*Zinnia elegans*  
*Tay Coll vol 8*



Labat 91



*Gerbera astrovides.*  
Jacq. Coll. vol. 1.



*Rudbeckia amplexifolia*

Tayl. Coll. vol. 5

Tafel 97




Rudbeckia alata.  
Tasg Coll. vol. 5.



*Conoclinium* / & *encensum*

*Coreopsis artemisiifolia.*  
Jacq Coll. vol. 5





*Calendula arborea*  
Jacq. coll. vol. 3



Lobelia siphilitica

Jacq. Coll. vol. 1



{/irftfa mi/i/ati\*



*Orchis variegata*  
Jug Coll vol 2



*Neottia speciosa.*  
Jacq. Coll. vol. 3.



*Neotti* la minor:  
J. a. (1860) vol. 3.



Joh. G. C. 1753  
*Limodorum  
tuberum scutellatum*  
Jug. Coll. vol. 4



*Limodorum diurnum*  
Jacq. Coll. vol. 1.



*Epidendrum dendrogaeum*  
Tayl. Coll. vol. 5.




*Epidendrum cochleatum*  
Jacq. Coll. vol. 3.




*Passiflora cuprea.*  
Jug. coll. vol. 3








Poteriochrysa  
cerasiferae  
Tuy coll. ad.



*Pothos grandifolia*  
Jacq. Coll. vol. 1




*Lothox palmata.*  
Jacq. Coll. vol. 4



*Arum pallidofolium*  
Lam. t. 127 v. 3



*Dorstenia Contrajerva*  
Tay Coll. vol. 3



Carex //m<//>/jn',  
Taylill' coll.




*Pygmanthus speciosus*  
Tag Coll. vol. 1



*Morus mauritiana*  
Jacq Coll vol 3



*Begonia minor*  
Berg Coll vol 1





*Calypha alopecuridea*  
Jacq Coll vol 3.



*Croton punctatum*  
Jacq. Cell vol. 1



*Croton punctatus.*  
Tay Coll. wt. 3.



*Lavatera glauca*

7



*Trichosanthes fatidiflora*

< fQfJ. (off n\*)



*Myrsia segregata.*  
Tay Coll. vol. 2





*Dioclea triphylla*  
var. *latifolia*




*Kiggelaria integrifolia*  
Jacq. Coll. vol. 2.





*Andropogon distachyos*  
Jacq Coll. vol. 1.



*Andropogon undatus*  
Tug coll vols.

Tab. 605




*Mimosa pudica L.* \* v.  
Jacq. Coll. vol. 4.




*Mimosa portoricensis*  
Jacq Coll ed 1.





*Panax aculeatum*  
Jug. ill. v. 2





*Zamia udorum*  
Jacq. Coll. n. d. 3







*Polypodium plumosum*  
Troy Coll. vol. 4





*Polypodium fraxinifolium*  
Jacq Coll vol 3



*Lycopodium molle.*

'fir #if.t



*Polypodium Hypocryps*  
Gray Coll. vol. 3



*Polypodium alpinum*

Jacq. Coll. vol. 2.




*Dryopteris marginalis*



Blechnum occidentale  
Jug Coll. vol. 3



*Elephantum striatum.*  
Eng. Ed. vol. 2.



*Pteris diffusa*  
Tay Coll no -



*Tremella juniperina*.  
Jacq. Coll. vol. 2.



*Tremella clavariiformis*  
Jacq Coll. vol 2

INDIAN BOTANIC GARDEN.

LIBRARY

PRO. NO. 60



NUMBER



CARD NO. 111



CLASSIFICATION



DATE 8.7.1944

