

FLORA OF INDIA
SERIES - 4

**A Census
of The Indian
Pteridophytes**

R. D. DIXIT

BOTANICAL SURVEY OF INDIA

**A CENSUS OF
THE INDIAN
PTERIDOPHYTES**

FLORA OF INDIA

Series IV

**A CENSUS OF
THE INDIAN PTERIDOPHYTES**

R. D. DIXIT



BOTANICAL SURVEY OF INDIA

Department of Environment

© Government of India, June, 1984

Price

Published by the Director, Botanical Survey of India,
Howrah-711103 and printed at Deep Printers,
3/26, Ramesh Nagar, New Delhi-110015

C O N T E N T S

Foreword	i
Preface	ii
Acknowledgements	iii
System of classification	1
Huperziaceae	7
Lycopodiaceae	9
Selaginellaceae	11
Isoetaceae	18
Equisetaceae	19
Psilotaceae	20
Helminthostachyaceae	20
Botrychaceae	21
Ophioglossaceae	22
Angiopteridaceae	24
Marattiaceae	26
Christensiaceae	26
Osmundaceae	27
Plagiogyriaceae	27
Gleicheniaceae	29
Dicranaopteridaceae	30
Grammitaceae	31
Loxogrammaceae	33
Polypodiaceae	34
Dipteridaceae	57
Platyceriaceae	57
Drynariaceae	58

Schizaeaceae	58
Anemiacetaceae	59
Lygodiaceae	59
Sinopteridaceae	61
Cheilanthaceae	63
Cryptogrammaceae	66
Actiniopteridaceae	68
Pteriaceae	68
Acrostichaceae	73
Adiantaceae	74
Hemionitidaceae	77
Antrophyaceae	80
Vittariaceae	81
Taenitidaceae	83
Parkeriaceae	83
Marsileaceae	84
Hymenophyllaceae	85
Hymenophyllopsidaceae	90
Dicksoniaceae	92
Cyatheaceae	92
Dennstaedtiaceae	94
Monachosoraceae	97
Hypolepidaceae	97
Pteridiaceae	98
Lindsaeaceae	98
Thelypteridaceae	102
Aspleniaceae	116
Onocleaceae	123
Woodsiaceae	124

Athyriaceae	124
Hypodemataceae	137
Aspidiaceae	137
Peranemataceae	145
Dryopteridaceae	146
Lomariopsidaceae	161
Bolbitidaceae	161
Elaphoglossaceae	164
Nephrolepidaceae	166
Oleandraceae	167
Davalliaceae	168
Gymnogrammitidaceae	172
Blechnaceae	172
Stenochlaenaceae	173
Azollaceae	174
Salviniaceae	174
Index to Genera	175

F O R E W O R D

Indian Flora is rich in diversity and the Pteridophytic Flora greatly contributes to this diversity. In the early part of the Century, taxonomic studies on flowering as well as non-flowering plants did not receive the desired attention. However, during last about 30 years, particularly with the revival of the Botanical Survey of India, intensive explorations have been made in different parts of the country by the scientists of the B. S. I. and several other institutions. These expeditions have resulted in several floristic accounts and also some taxonomic revisions.

Work on Pteridophytes was also taken up in some institutions in the country which included the Botanical Survey India, Some Institutes of C S I R, like the National Botanical Research Institute, Lucknow and some Universities.

Cytological and Cyto-taxonomic studies on Pteridophytes also received attention of several scientists. With the intensification of studies on Cryptogams in the Botanical Survey of India, a few years ago, it was felt that it would be useful to bring out a census of Indian Pteridophytes.

One of our young Pteridologist, Dr. R. D. Dixit, came forward to undertake the work. The present census is the result of Dr. Dixit's bibliographic and herbarium studies of about 10 years.

Literature on floristics and taxonomy appears in such variety of journals and at so fast a speed that it is difficult to claim total completeness in any work of this nature.

I am sure Dr. Dixit will welcome constructive suggestions to this inventory and, if considered justifiable, he will attempt a supplement of addenda, after a few years.

Meantime, it is hoped that the work will be found useful by all persons interested in the natural Flora of the region.

S. K. JAIN
Director

Botanical Survey of India
10, June 1984

P R E F A C E

The Pteridophytic flora of the Indian region is very rich due to remarkable altitudinal variations ranging from sea level to the highest mountain ranges of the world, the long coastal line, hot desert in the west as well as cold desert in the north alongwith humid tropical conditions in some parts. In addition to it, adjacent countries and in many cases even far distant countries have contributed quite a large number of species to the flora of Indian subcontinent due to presence of great variety of climatic conditions, soil types and altitude, etc.

It is now more than one hundred years that Col. R.H. Beddome (1883) published "A Handbook to the ferns of British India Ceylon and the Malaya Peninsula" and its supplement in 1892 but without Azollaceae, Marsileaceae and Salviniaceae amongst the ferns and fern-allies as a whole. Inspite of these lacunae the Handbook still remains the most important reference work in Indian Fern-flora even today.

The taxonomic concepts of Pteridophytes have undergone tremendous revolutionary changes during the 20th century in comparison any other group of plants resulting changes in circumscription of species, genera and families from time to time. Many new species and records have been reported during the past one hundred years. But no serious attempt has been made so far, to consolidate all available data on Indian basis except Nayar's & Kaur. "Companion to Beddome's Handbook" listing nomenclatural changes with regard to Beddomes (l.c.) names only, a System of Classification and a chapter on Thelypteroid ferns as per names given by Prof. R.E. Holttum of the Royal Botanic Gardens, Kew.

In view of absence of even a list of Pteridophytes occurring in India, the present work has been undertaken to compile all available species of Pteridophytes found in the present political boundary of India in the form "A Census of the Indian Pteridophytes."

A broad outline of the System of Classification of Pteridophyta occurring in India mainly following Pichi Sermolli (1977, 1982) with a little modifications of Ching (1978, 1978a) have been incorporated to make the System more upto-date, comprehensive and convenient. Original reference citations for each genus alongwith reference of Monographic/ revisionary comprehensive work have been cited wherever available to avoid repetitions at species level. References to published list, enumerations have not been mentioned normally. Type species of each genus has been mentioned with full reference citations mainly following Pichi Sermolli

(1977). The genera are arranged alphabetically under each family and similarly species are also arranged alphabetically under each genus for the sake of convenience. At species level the valid name has been given in bold types, basionym and important synonyms if any, known in Indian flora have been given in italics. Cross references to Beddome (1863-1892) and Clarke (1880), have been provided. Distribution of each species in India and world has been given.

The present Census comprises of 67 families, 191 genera and more than 1,000 species of Pteridophytes from the Indian region, against 98 genera of ferns reported by Beddome (1883) from the British India, Ceylon and the Malay Peninsula (of which 6 are of non Indian distribution and 16 are being treated merely as synonyms at present) and 466 species of ferns from the present political boundary of India.

Every effort has been made to consult and incorporate recent nomenclature and publications made upto March, 1984. However, the responsibility of all judgements and errors is my own if any thing is omitted, misjudged in the text, I am sure the consultant will excuse me on the ground of human limitations.

The six dominant families of ferns as regards the number of genera given in brackets are Polypodiaceae (27), Thelypteridaceae (21), Athyriaceae (14), Hymenophyllaceae (10) and Hemionitidaceae (6).

It is hoped that this book providing uptodate data on number, nomenclature and distribution of taxa, will be useful to those who are interested in the Pteridophytic flora of this country.

Allahabad

R. D. Dixit

Dated : 29th March, 1984

A C K N O W L E D G E M E N T S

I am deeply indebted to Dr. S.K. Jain, Director, Botanical Survey of India, Howrah for the suggestion and allotment of the problem, guidance and constant encouragement throughout the period of work. I am thankful to Dr. J.N. Vohra, Regional Botanist (Cryptogams) and members of the Cryptogamic Unit, and Dr. V.S. Agarwal, Editor of Publications, Botanical Survey of India for their help rendered during the preparation and publication of the present work. Thanks are also due to Dr. N.P. Balakrishnan, Deputy Director, Botanical Survey of India, Allahabad for the facilities. I wish to express my gratitude to Prof. R.E. Holttum of England, Prof. R.C. Ching of China, Prof. R.E.G. Pichi Sermolli of Italy and Prof. K.U. Kramer of Switzerland for sending their recent literature on the subject.

SYSTEM OF CLASSIFICATION OF PTERIDOPHYTA*
(INCLUDES INDIAN PTERIDOPHYTES ONLY)

Division : Pteridophyta
Subdivision : Lycophtina
 Class : Lycopsida
 Subclass : Lycopodiidae
 Order : Lycopodiales
 Fam : Huperziaceae
 Gen : Huperzia, Phlegmariurus
 Fam : Lycopodiaceae
 Gen : Diphasiastrum, Lycopodiastrum, Lycopodium,
 Palhinhaea
 Subclass : Selaginellidae
 Order : Selaginellales
 Fam : Selaginellaceae
 Gen : Selaginella
 Subclass : Isoetidae
 Order : Isoetales
 Fam : Isoetaceae
 Gen : Isoetes
Subdivision : Sphenophytina
 Class : Equisetopsida
 Subclass : Equisetidae
 Order : Equisetales
 Fam : Equisetaceae
 Gen : Equisetum
Subdivision : Psilophytina
 Class : Psilotopsida
 Order : Psilotales
 Family : Psilotaceae
 Gen : Psilotum
Subdivision : Filicophytina
 Class : Ophioglossopsida
 Subclass : Ophioglossidae
 Fam : Helminthostachyaceae
 Gen : Helminthostachys
 Fam : Botrychiaceae
 Gen : Botrychium

*After Pic. Ser. (1977, 1982) ; Ching (1978, 1978a)

Fam : Ophioglossaceae
Gen : Ophioderma, Ophioglossum

Class : Marattiopsida

Subclass : Marattiidae

Order : Marattiales

Fam : Marattiaceae
Gen : Marattia

Fam : Angiopteridaceae
Gen : Angiopteris

Fam : Christensiaceae
Gen : Christensenia

Class : Filicopsida

Subclass : Osmundidae

Order : Osmundales

Fam : Osmundaceae
Gen : Osmunda

Subclass : Plagiogyriidae

Order : Plagiogyrlales

Fam : Plagiogyriaceae
Gen : Plagiogyria

Subclass : Gleicheniidae

Order : Gleicheniales

Fam : Gleicheniaceae
Gen : Gleichenia

Fam : Dicranopteridaceae
Gen : Dicranopteris

Order : Polypodiales

Suborder : Polypodiinae

Fam : Grammitaceae
**Gen : Ctenopteris, Prosaptia, Xiphopteris, Grammitis,
Scleroglossum**

Fam : Loxogrammaceae
Gen : Loxogramme

Fam : Polypodiaceae
**Gen : Arthromeris, Belvisia, Christopteris, Colysis,
Crypsinus, Dendroglossa, Drymotaenium,
Goniophlebium, Holcosorus, Lemmaphyllum,
Lepidogrammitis, Lepisorus, Leptochilus,
Marginaria, Metapolypodium, Microsorium,
Neocheriopteris, Paraleptochilus. Phlebodium,
Phymatopteris, Phymatosorus, Polypodiastrum,
Polypodioides, Polypodium, Pseudodrynaria,
Thylacopteris. Tricholepidium.**

Fam : Platyceriaceae
Gen : Platycerium

Fam : Drynariaceae

Gen : *Drynaria*
 Suborder : Dipteridineae
 Fam : Dipteridaceae
 Gen : *Dipteris*
 Subclass : Schizaeidae
 Order : Schizaeales
 Fam : Schizaeaceae
 Gen : *Actinostachys, Schizaea*
 Fam : Anemiaceae
 Gen : *Anemia*
 Fam : Lygodiaceae
 Gen ; *Lygodium*
 Order : Pteridales
 Suborder : Pteridnieae
 Fam : Sinopteridaceae
 Gen : *Doryopteris, Mildella, Notholaena,*
Pellaea
 Fam : Cheilanthaceae
 Gen : *Cheilanthes*
 Fam : Cryptogrammaceae
 Gen : *Cryptogramma*
 Fam : Actiniopteridaceae
 Gen : *Actiniopteris*
 Fam : Pteridaceae
 Gen : *Pteris*
 Fam : Acrostichaceae
 Gen : *Acrostichum*
 Fam : Adiantaceae
 Gen : *Adiantum*
 Fam : Hemionitidaceae
 Gen : *Annogramma, Conogramme, Hemionitis,*
Idiogramma, Pityrogramma, Syngamma
 Fam : Antrophyaceae
 Gen : *Antrophyum*
 Fam : Vittariaceae
 Gen : *Vittaria*
 Suborder : Taenitidineae
 Fam : Taenitidaceae
 Gen : *Taenitis*
 Suborder : Parkeriineae
 Fam : Parkeriaceae
 Gen : *Ceratopteris*
 Subclass : Marsileidae
 Order : Marsileales

Fam : Marsileaceae
Gen : Marsilea

Order : Hymenophyllales

Fam : Hymenophyllaceae

Gen : Gonocormous, Hymenophyllum, Mecodium, Meringium, Microgonium, Microtrichomanes, Pleuromanes, Selenodesmium, Sphaerocionum, Vandenboschia

Subclass : Hymenophyllidae

Order : Hymenophillopsidales

Fam : Hymenophyllopsidaceae

Gen : Cephalomanes, Crepidomanes

Order : Dicksoniales

Suborder : Dicksoniineae

Fam : Dicksoniaceae

Gen : Cibotium

Suborder : Cyatheineae

Fam : Cyatheaceae

Gen : Cyathea

Order : Dennstaedtiiales

Suborder : Dennstaedtiineae

Fam : Dennstaedtiaceae

Gen : Dennstaedtia, Emoditopteris, Microlepia

Fam : Monachosoraceae

Gen : Monachosorum

Fam : Hypolepidacea

Gen : Histiopteris, Hypolepis,

Fam : Pteridiaceae

Gen : Pteridium

Suborder : Lindsaeineae

Fam : Lindsaeaceae

Gen : Lindsaea, Sphenomeris, Tapeinidium

Order : Aspidiales

Suborder : Thelypteridinae

Fam : Thelypteridaceae

Gen : Ampelopteris, Amphineuron, Christella, Coryphopteris, Cyclogramma, Cyclosorus, Dictyocline, Glaphyropteridopsis, Macrothelypteris, Metathelypteris, Oreopteris, Parathelypteris, Phegopteris, Pneumatopteris, Pronephrium, Pseudocyclosorus, Pseudophegopteris,

*Sphaerostephanos, Stenogramma, Thelypteris,
Trigonospora*

Suborder : Aspleniineae

Fam : *Aspleniaceae*

Gen : *Asplenium, Ceterach, Ceterachopsis*

Suborder : Aspidiineae

Fam : *Onocleaceae*

Gen : *Onoclea*

Fam : *Woodsiaceae*

Gen : *Cheilanthopsis, Woodsia*

Fam : *Athyriaceae*

Gen : *Acystopteris, Anisocampium, Athyriopsis,*

Athyrium, Coronopteris, Cystopteris,

Dictyodroma, Diplaziopsis, Diplazium.

Dryoathyrium, Gymnocarpium, Kuiniwatsukia,

Lunathyrium, Triblemma

Fam : *Hypodemataceae*

Gen : *Hypodematum*

Fam : *Aspidiaceae*

Gen : *Acrorumohra, Ctenitis, Ctenitopsis, Cyrtomium,*

Didymochlaena, Lastreopsis,

Pleocnemia, Pteridrys, Quericiflix, Tectaria

Fam : *Peranemataceae*

Gen : *Acrophorus, Diacalpe, Lithostegia, Peranema*

Fam : *Dryopteridaceae*

Gen : *Arachniodes, Dryopteris, Nothoperanema,*

Polystichum

Fam : *Lomariopsidaceae*

Gen : *Leptogramma*

Fam : *Bolbitidaceae*

Gen : *Bolbitis, Egenolfia*

Fam : *Elaphoglossaceae*

Gen : *Elaphoglossum*

Suborder : Davalliineae

Fam : *Nephrolepidaceae*

Gen : *Arthropteris, Nephrolepis*

Fam : *Oleandraceae*

Gen : *Oleandra*

Fam : *Davalliaceae*

Gen : *Araiostegia, Davallia, Humata,*

Leucostegia, Paradavallodes

Fam : *Gymnogrammitidaceae*

Gen : *Gymnogrammitis*

Order : Blechnales

Fam : Blechnaceae

Gen : *Blechnidium*, *Blechnum*, *Brainea*
Doodia, *Woodwardia*

Fam : Stenochlaenaceae

Gen : *Stenochlaena*

Subclass : Salviniidae

Order : Salviniales

Fam : Azollaceae

Gen : *Azolla*

Fam : Salviniaceae

Gen : *Salvinia*

HUPERZIACEAE Rothm.

Huperzia Bernh. Schard. Journ. Bot. 1800 (2) : 126. 1801 ; **Mandal & Sen**, Biol. Mem. 3 (2) : 192. 1978.

Lectotype : **H. selago** (L.) Bernh. ex Schrank & Mart. Hort. Monac. 3. 1829.

H. cancellata (Spring) Trev. Atti. Soc. Ital. Sci. Nat. 17 : 247. 1875.
Lycopodium cancellatum Spring, Mém. Acad. Sci. Bélg. 24 (2) : 27. 1850.

Distrib. : India (Arunachal Pradesh), Bhutan.

H. carinata (Desv.) Trev. Atti. Soc. Ital. Sci. Nat. 17 : 247. 1875.
Lycopodium carinatum Desv. Encycl. Bot. Suppl. 3 : 559. 1813.

Distrib. : India (Tamil Nadu, Andaman & Nicobar Islands), China, Malaya Peninsula, Malesian Islands, Philippines, Formosa, Polynesia.

H. ceylanica (Spring) Trev. Atti. Soc. Ital. Sci. Nat. 17 : 248. 1875.
Lycopodium ceylanicum Spring, Mém. Acad. Sci. Bélg. 15 (1) : 37. 1843 ; 24(2) : 16. 1850.

Distrib. : India (Meghalaya, Tamil Nadu), Sri Lanka, Java.

H. cryptomerina (Maxim.) Dixit, Journ. Bomb. Nat. Hist. Soc. 77 (3) : 541. 1981. *Lycopodium cryptomerinum* Maxim. Bull. Acad. Sci. St. Petersb. 15 : 231. 1870.

Distrib. : India (Arunachal Pradesh), China, Japan.

H. fordii (Bak.) Dixit, comb. nov. *Lycopodium fordii* Bak. Handb. Fern Allies 17. 1887.

Distrib. : India (Eastern Himalayas), China, Japan.

H. hamiltonii (Spring) Trev. Atti. Soc. Ital. Sci. Nat. 17 : 248. 1875.
Lycopodium hamiltonii Spring, Syst. Veg. 5 : 429. 1828 ; Mém. Acad. Sci. Belg. 15 (1) : 35. 1843 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 590. 1880.

Distrib. : Throughout India in hilly regions, Burma, Sri Lanka, Indo-China, Malesian Islands.

H. herteriana (Kümm.) Sen & Sen, Fern Gaz. 11 (6) : 415. f. 1 a - j. 1978. *Lycopodium herterianum* Kümm. Magyar. Bot. Lap. 26 : 99. 1928. *Lycopodium lucidulum* auct. Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 589, t. 84. f. 1. 1880.

Distrib. : India (Sikkim, West Bengal, Manipur), China.

H. niligarica (Spring) Dixit, Journ. Bomb. Nat. Hist. Soc. 77 (3) : 541. 1981. *Lycopodium niligaricum* Spring, Bull. Acad. Sci. Belg. 15 (1) : 58. 1843.

Distrib. : India (Tamil Nadu, Kerala).

H. petiolata (Clarke) Dixit, Journ. Bomb. Nat. Hist. Soc. 77 (3) : 541. 1981. *Lycopodium hamiltonii* var. *petiolatum* Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 590. 1880.

Distrib. : India (Meghalaya).

H. pulcherrima (Wall. ex Hook. & Grev.) Sen & Sen, Fern Gaz. 11 (6) : 419. f. 2 i - r. 1978. *Lycopodium pulcherrimum* Wall. ex Hook. & Grev., Icon. Fil. t. 58. 1831. *L. setaceum* Buch. Ham. ex D. Don. var. *subulifolium* Wall. apud Clarke, Trans. Linn. Soc. Lond. II. Bot. 1. 590. 1880.

Distrib. : India N W Himalayas, E. (India), Nepal, Bhutan, China.

H. selago (L.) Bernh. ex Schrank et Mart. Hort. Monac. 3. 1829. *Lycopodium selago* L. Sp. Pl. 2 : 1102. 1753.

Distrib. : India (Sikkim), Arctic and north temperate zones of both hemispheres, also New Zealand, Australia, Arctic America and mountains of Central Brazil.

H. serrata (Thunb.) Rothm. Feddes Repert. 54 (1) : 59. 1944. *Lycopodium serratum* Thunb. Fl. Jap. 341. t. 38. 1784.

Distrib. : India (West Bengal, Sikkim, Arunachal Pradesh, Assam, Meghalaya, Manipur, Tamil Nadu), Nepal, Burma, Sri Lanka, China, Japan, Malaya Peninsula, Malesian Islands, Fiji, Samoa, Sandwich Islands, Tropical America.

H. squarrosa (Forst.) Trev. Atti. Soc. Ital. Sci. Nat. 17 : 247. 1875. *Lycopodium squarrosum* Forst, Prod. 86. 1786 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 591. 1880.

Distrib. : India (West Bengal, Sikkim, Meghalaya, Assam, Kerala), Nepal, Burma, Bangladesh, Sri Lanka, Malesian Islands, Philippines, Polynesia, Seychelles, Mascareen Isles.

H. subulifolia (Wall. ex Hook. & Grev.) Trev. Atti. Soc. Ital. Sci. Nat. 17 : 248. 1875. *Lycopodium subulifolium* Wall. ex Hook. & Grev. Icon. Fil. t. 49. 1831.

Distrib. : India (West Bengal, Sikkim, Meghalaya, Uttar Pradesh), Nepal, Sri Lanka.

H. vernicosa (Hook. & Grev.) Trev. Atti. Soc. Ital. Sci. Nat. 17 :

248. 1875. *Lycopodium vernicosum* Hook. & Grev. in Hook. Bot. Misc. 2 : 364. 1831.

Distrib. : India (Tamil Nadu, Kerala), Sri Lanka.

Phlegmariurus (Herter) Holub, Preslia 36 (1) : 17, 21. 1964.

Type : **P. phlegmaria** (L.) Sen & Sen, Fern Gaz. 11 (6) : 421 f. 4. a - j. 1978.

P. phlegmaria (L.) Sen & Sen, Fern Gaz. 11 (6) : 421. f. 4 a j. 1978. *Lycopodium phlegmarium* L. Sp. Pl. 2 : 1101. 1753 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 592. 1880.

Distrib. : India (West Bengal, Sikkim, Assam, Arunachal Pradesh, Kerala, Andaman & Nicobar Islands) Tropics of the old world extending up to New Zealand and Australia.

P. phyllanthum (Hook. & Arn.) Dixit, Journ Bomb. Nat. Hist. Soc. 77 (3) : 541. 1981. *Lycopodium phyllanthum* Hook. & Arn. Bot. Beech. Voy. 102. 1841.

Distrib. : India (Uttar Pradesh, Arunachal Pradesh, West Bengal, Sikkim, Assam, Manipur, Meghalaya, Kerala, Tamil Nadu), Arctic and Alpine Zones of both hemispheres ; mountains of Tropical Asia, Africa, America, Cape, Mascareen Islands, Polynesia.

LYCOPODIACEAE P. Beauv. ex Mirb.

Lycopodium L. Sp. Pl. 2 : 1101. 1753 ; Spring, Mém. Acad. Sci. Bélg. 15 (1) : 9. 1843 ; 24 (2) : 5. 1850 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 590. 1880 ; Chowdhury, Trans. Nat. Inst. Sci. India 1 : 195. 1937. Mandal & Sen, Biol. Mem. 3 (2) : 192. 1978 ; Dixit, Bull. Bot. Surv. India 24 (1-4) : 178. f. 1-3. 1982.

Lectotype : **L. clavatum** L. Sp. Pl. 2 : 1101. 1753.

L. annotinum. L. Sp. Pl. 2 : 1103. 1753.

Distrib. : India (Uttar Pradesh, Sikkim, Meghalaya), Arctic Zones and mountains of North temperate zone in both the hemispheres.

L. japonicum Thunb. Fl. Jap. 341. 1784. *L. clavatum* of auct. Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 592. 1880.

Distrib. : Throughout India in the mountainous regions, Nepal, Bhutan, Burma, Sri Lanka, China, Japan, Bangladesh.

L. veitchii Christ, Bull. Geor. Bot. Mans. 106. 1906 ; *Lycopodium*

annotinum *sensu* Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 592. 1880 (*pro parte*).

Distrib. : India (Sikkim, Arunachal Pradesh), China, Japan.

Diphasiastrum Holub, Preslia 47 : 104. 1975 ; Sen, & Sen Fern Gaz. 11 (6) : 424. f. k n. 1978.

Type : **D. complanatum** (L.) Holub, Preslia 47 : 107. 1975.

D. alpinum (L.) Holub, Preslia 47 : 107. 1975. *Lycopodium alpinum* L. Sp. Pl. 2 : 1104. 1753.

Distrib. : India (Sikkim), Arctic Zones & mountains of the North temperate zone in both the hemispheres.

D. complanatum (L.) Holub, Preslia 47. 108. 1975. *Lycopodium complanatum* L. Sp. Pl. 2 : 1102. 1753 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 593. 1880. *L. veitchii* *sensu* Mandal & Sen, Biol. Mem. 3 (2) : 248. 1978. *Diphasium complanatum* (L.) Rothm. Feddes Repert. 54 : 584. 1944 ; Sen & Sen, Fern Gaz. 11 (6) : 424. f. 5 a - j. 1978.

Distrib. : India (Eastern India, Eastern Himalayas, South India), North Temperate Zones & Alpine regions of the tropics.

D. wightianum (Wall. ex Hook. & Grev.) Holub, Preslia 47 : 108. 1975. *Lycopodium wightianum* Wall. ex Hook. & Grev. Enum. Fil. 85. 1824.

Distrib. : India (Kerala, Tamil Nadu), Sri Lanka, Malay Islands.

Lycopodiastrum Holub ex Dixit, Journ. Bomb. Nat. Hist. Soc. 77 (3) : 541. 1981.

Type : **L. casuarinoides** (Spring) Holub ex Dixit, Journ. Bomb. Nat. Hist. Soc. 77 (3) : 541. 1981.

L. casuarinoides (Spring) Holub ex Dixit, Journ. Bomb. Nat. Hist. Soc. 77 (3) : 541. 1981. *Lycopodium casuarinoides* Spring, Mém. Acad. Sci. Belg. 15 (1) : 92. 1843 ; 24 (2) : 45. 1850.

Distrb. : India (Assam, Meghalaya), Bhutan, Burma, China, Malaya Peninsula, Malaya Islands, Philippine Islands.

Palhinhaea Franco. & Vasc. in Vasc. & Franco. Bol. Soc. Broter. ser. 2. 41 : 24. 1967 ; Sen & Sen, Fern Gaz. 11 (6) : 423. 1978.

Type : **P. cernua** (L.) Franco. & Vasc. in Vasc. & Franco, Bol. Soc. Broter. ser. 2. 41 : 25. 1967.

P. cernua (L.) Franco. & Vasc. in Vasc. & Franco. Bol. Soc. Broter. ser. 2. 41 : 25. 1967. *Lycopodium cernuum* L. Sp. Pl. 2 : 1103. 1753 ;

Clarke Trans. Linn. Soc. Lond. II. Bot. 1 : 591. 1880.

Distrib. : Throughout India in mountainous regions, Tropics of both the hemispheres.

SELAGINELLACEAE Milde

Selaginella P. Beauv. Prod. Fam. Aétheog. 101. 1805 ; Spring, Mém. Acad. Sci. Bélg. 24 (2) : 55. 1850 ; Alston, Proc. Natn. Inst. Sci. India 11 : 211. 1945 ; Panigr. & Dixit, Proc. Natn. Acad. Sci. India, Sect. B. 36 : 102. 1966 ; Journ. Indian Bot. Soc. 46 : 222. 1967 ; Proc. Natn. Inst. Sci. India 34 : 191. 1968.

Type : **S. selaginoides** (L.) Link, Fil. Sp. Hort. Berol. 158. 1841.

S. adunca A. Br. ex Hieron. in Engl. & Prantl, Nat. Pfl. 1 (4) : 674. 1902. **S. yemensis** sensu Duthie, Cat. Kumaon Pl. 232. 1866.

Distrib. : India (Uttar Pradesh, Himachal Pradesh).

S. aitchisonii Hieron. in Engl. & Prantl. Nat. Pfl. 1 (4) : 674. 1902. **S. sanguinolenta** (L.) Spring f. *aitchisonii* (Hieron.) Alston, Proc. Natn. Inst. Sci. India 11 : 215. 1945.

Distrib. : India (Kashmir), Afghanistan, Turkistan.

S. biformis A. Br. ex Kuhn, Fil. Afr. 189. 1868. (*nomen. nud.*) Kuhn, Forsch Gaz. IV, Bot. 6 : 17. 1889.

Disirib. : India (Assam, Meghalaya, Nagaland, Manipur), Burma, South China, Philippines across Celebes, Java, Sumatra.

S. bisulcata Spring, Mém. Acad. Sci. Bélg. 24 (2) : 259. 1850.

Distrib. : India (Himachal Pradesh, West Bengal, Sikkim, Nagaland, Manipur, Meghalaya), Burma, China.

S. blatteri Bole & Almeida, Journ. Bomb. Nat. Hist. Soc. 74 : 320. f. 1. 1977.

Distrib. : India (Maharashtra).

S. brachystachya (Hook. & Grev.) Spring, Bull. Acad. Brux. 10 : 232. 1843. *Lycopodium brachystachyum* Hook. & Grev. in Hook. Bot. Misc 3 : 107. 1833.

Distrib. : India (Maharashtra), Sri Lanka.

S. braunii Bak. Gard. Chron. 1120. 1867.

Distrib. : India (Uttar Pradesh, Tamil Nadu), China, Georgia, Jamaica, Brazil.

S. bryopteris (L.) Bak. Journ. Bot. 22 : 376. 1884. *Lycopodium bryopteris* L. Sp. Pl. 2 : 1103. 1753. *L. imbricatum* Roxb. Hort. Beng. 75. 1814 (*nom. nud.*) ex Griff. Calc. Journ. Nat. Hist. 4 : 475. 1844 (*non* Forssk. 1775). *Selaginella imbricata* (Roxb.) J. Scott, Journ. Agr. Hort. Soc. India 1 : 260. 1868.

Distrib. : Throughout India in mountainous regions up to 1000 m.

S. cataractrum Alston, Proc. Natn. Inst. Sci. India 11 : 228. 1945.

Distrib. : India (Tamil Nadu, Kerala).

S. chrysocaulos (Hook. & Grev.) Spring, Bull. Acad. Brux. 10 : 232. 1843. *Lycopodium chrysocaulos* Hook. & Grev. in Hook. Bot. Misc. 2 : 401. 1831. *Selaginella philippina* var. *khasiensis* Bak. Journ. Bot. 22 : 298. 1884.

Distrib. : India (Himachal Pradesh, Uttar Pradesh, Sikkim, Arunachal Pradesh, Nagaland, Manipur, Meghalaya, Kerala), Nepal, Bhutan.

S. chrysorrhizos Spring, Mém. Acad. Sci. Bélg. 24 (2) : 251. 1850.

Distrib. : India (Assam, West Bengal), Nepal, Burma, Siam, Annam, Laos.

S. ciliaris (Retz.) Spring, Bull. Acad. Brux. 10 : 23. 1843. *Lycopodium ciliare* Retz. Obs. 5 : 32. 1789. *Selaginella exigua* Spring, Mém. Acad. Sci. Bélg. 24 (2) : 238. 1850 ; Haines, Bot. Bihar & Orissa 3 : 1276. 1924.

Distrib. : India (Uttar Pradesh, Bihar, West Bengal, Sikkim, Assam, Meghalaya, Tripura, Orissa, Madhya Pradesh, Kerala, Tamil Nadu, Andaman & Nicobar Islands), Burma, Sri Lanka, China, Phillipine Islands.

S. coonoriana Dixit, Bull. Bot. Surv. India 25 : t. 1. f. 1. text. fig. 1A - G. 1983.

Distrib. : India (Tamil Nadu).

S. decipiens Warb, Monsunia 1 : 127. 1900.

Distrib. : India (Arunachal Pradesh, Assam, Mizoram, Meghalaya) Malay Peninsula, Indo China, Tonkin.

S. delicatula (Desv. ex Poir.) Alston, Journ. Bot. 70 : 282. 1932.

Lycopodium delicatulum Desv. ex Poir. in Lam. Encycl. Suppl. 3 : 554. 1814. *Selaginella crassicaule* (Hook. & Grev.) Spring, Bull. Acad. Brux. 10 : 232. 1843.

Distrib. : India (Sikkim, Assam, Tripura, Maharashtra, Kerala, Tamil Nadu, Goa, Andaman & Nicobar Islands) China (eastwards) to Philippine Islands, Moluccas, Borneo, Siam, Annam, Cambodia, Tonkin, Cochin China.

S. frondosa Warb, Monsunia 1 : 105. 117. 1900. **S. superba** Alston, Journ. Bot. 70 : 63. f. A F. 1932.

Distrib. : India (Andaman & Nicobar Islands) Malaya, Malesian Islands, Indo China, Tonkin, Annam.

S. fulcrata (Buch. - Ham. ex D. Don) Spring. Bull. Acad. Sci. Brux 10 : 231. 1843. *Lycopodium fulcratum* Buch. Ham. ex D. Don, Prod. Fl. Nepal 17. 1825.

Distrib. : India (Bihar, Orissa, Maharashtra), Nepal.

S. ganguliana Dixit, Bull. Bot. Surv. India 26 : t. 1. f. 1. A I. 1984.

Distrib. : India (Kerala).

S. helferi Warb, Monsunia 1 : 107, 121. 1900. **S. plana** sensu Burkhill, Rec. Bot. Surv. India 10 : 228. 1925 (*non* Hieron. 1901). **S. willdenovii** sensu Burkhill, Rec. Bot. Surv. India 10 : 228. 1925 (*non* Bak. 1867).

Distrib. : India (Arunachal Pradesh, Assam, Nagaland, Meghalaya, Andaman & Nicobar Islands), Burma, China, Indo China.

S. hookeri Bak. Journ. Bot. 23 : 21. 1886.

Distrib. : India (Assam, Arunachal Pradesh, Nagaland, Manipur, Meghalaya), Burma, Malay Peninsula, Indo China, Annam.

S. indica (Milde) Trayon, Anal. Miss. Bot. Gard. 42 : 52. f. 27. 1955. *S. rupestris* f. *indica* Milde, Fil. Eur. Atalant. 262. 1867. *S. longipila* of authors, as Alston, Proc. Natn. Inst. Sci. India 11 : 214. 1945 ; Tagawa in Hara's Univ. Mus. Univ. Tokyo Bull. no. 2. 200. 1971.

Distrib. : India (Uttar Pradesh, Bihar, West Bengal, Arunachal Pradesh, Meghalaya, Orissa, Madhya Pradesh, Tamil Nadu), Nepal, Bhutan.

S. inequalifolia (Hook. & Grev.) Spring, Bull. Acad. Brux. 10 : 145. 1843. *Lycopodium inequalifolium* Hook. & Grev. in Hook. Bot. Misc. 2 : 391. 1831.

Distrib. : India (Assam, Kerala, Tamil Nadu), Burma.

S. integerrima (Hook. & Grev.) Spring, Bull. Acad. Brux 10 : 1843.
Lycopodium integerrimum Hook. & Grev. in Hook. Bot. Misc. 2 : 396.
 1831. *Selaginella ornithopodioides sensu* Trimen, Journ. Linn. Soc. 24 :
 152. 1887 ; Alston, Proc. Natn. Inst. Sci. India 11 : 222. 1945.

Distrib. : India (Tamil Nadu), Sri Lanka.

S. intermedia (Bl.) Spring, Bull. Acad. Brux. 10 : 144. 1843. *Lyco-*
podium intermedium Bl. Enum. Pl. Jav. 2 : 269. 1830. *L. furcatum*
 Roxb. ex Griff. Calc. Journ. Nat. Hist. 4 : 475. 1844.

Distrib. : India (Kerala, Tamil Nadu) Malay Peninsula, Java,
 Sumatra, Burma, Borneo, Celebes, Annam, Siam.

S. involvens (Sw.) Spring, Bull. Acad. Brux. 10 : 136. 1843 ; emend.
 Hieron. Hedw. 50 : 2. 1911. *Lycopodium involvens* Sw. Syn. Fil. 182. 1806.
Selaginella caulescens (Wall. ex Hook. & Grev.) Spring, Bull. Acad. Brux.
 10 : 137. 1843.

Distrib. : India (almost throughout India in mountainous regions
 except northern part), Nepal, Bhutan, Burma, Sri Lanka, Indo China,
 Malesian Islands.

S. jainii Dixit, Bull. Bot. Surv. India 25 : t. 1. f. 2. text. fig. 2 A H.
 1983.

Distrib. : India (Madhya Pradesh).

S. kashmiriana (Milde) Dixit, nom. nov. *S. borealis* f. *indica*
 Milde, Fil. Eur. 267. 1867. *S. Sanguinolenta* f. *indica* (Milde) Alston,
 Proc. Natn. Inst. Sci. India 11 : 215. 1945.

Distrib. : India (North-West) Himalayas). Afghanistan.

S. megaphylla Bak. Journ. Bot. 23 : 180. 1885. *S. atrovirides* *sensu*
 Burkitt, Rec. Bot. Surv. India 10 : 228. 1925 (*non* Spring, 1838).

Distrib. : India (Arunachal Pradesh, Assam), Bhutan, Burma.

S. miniatospora (Dalz.) Bak. Journ. Bot. 23 : 249. 1885. *Lycopodium*
miniatosporum Dalz. in Hook. Journ. Bot. 4 : 114. 1852.

Distrib. : India (Maharashtra, Goa).

S. monospora Spring, Mém. Acad. Sci. Bélg. 24 (2) : 135. 1850.

Distrib. : India (West Bengal, Sikkim, Arunachal Pradesh,
 Meghalaya, Assam, Manipur, Kerala, Tamil Nadu), Burma, China, Indo-
 China, Tonkin, Annam.

S. nairii Dixit, Bull. Bot. Surv. India 26 : t. II. f. 2 A I. 1984.

Distrib. : India (Orissa).

S. nepalensis Spring, Bull. Acad. Brux. 10 : 234. 1843. *S. pallida* (Hook. & Grev.) Spring, Bull. Acad. Brux., 10 : 234. 1843. *Lycopodium pallidum* Hook. & Grev. Hook. in Bot. Misc. II p. 389. 1831.

Distrib. : India (Arunachal Pradesh, Manipur), Nepal.

S. ornata (Hook & Grev.) Spring, Bull. Acad. Brux. 10 : 232. 1843. *Lycopodium ornatum* Hook. & Grev. in Hook. Bot. Misc. 3 : 108. 1833.

Distrib. : India (Arunachal Pradesh, Meghalaya), Malesian Islands.

S. pallidissima Spring, Bull. Acad. Brux. 10 : 231. 1843. *S. integer rima* sensu Strachey in Gaz. North West Prov. 66. 1882 (*non* Spring, 1850).

Distrib. : India (Uttar Pradesh, Himachal Pradesh, Arunachal Pradesh).

S. panchghaniana Dixit, Bull. Bot. Surv. India 25 : t. I. f. 3. text fig. 3 A H. 1983.

Distrib. : India (Maharashtra).

S. panigrahi Dixit, Bull. Bot. Surv. India 25 : t. I. f. 4. text fig. 4 A H. 1983.

Distrib. : India (Madhya Pradesh).

S. pennata (D. Don) Spring, Bull. Acad. Brux 10 : 232. 1843. *Lycopodium pennatum* D. Don, Prod. Fl. Nepal 18. 1825.

Distrib. : India (Arunachal Pradesh, Assam, Meghalaya), Burma, Siam.

S. pentagona Spring. Mém. Acad. Sci. Bélg. 24 (2) : 150. 1850.

Distrib. : India (Arunachal Pradesh, Meghalaya, Assam).

S. picta A. Br. ex Bak. Journ. Bot. 23 : 19. 1885.

Distrib. : India (Arunachal Pradesh, Assam, Manipur, Nagaland), Burma, Indo China, Laos.

S. picta A. Br. ex Bak. f. *viridis* Alston, Proc. Natn. Inst. Sci. India 11 : 223. 1945.

Distrib. : India (Assam).

S. plana (Desv. ex Poir.) Hieron. in Engl. & Prantl. Nat. pfl. 1 (4) : 703. 1902. *Lycopodium planum* Desv. ex Poir. in Lam. Encycl. Suppl. 3 : 554. 1814.

Distrib. : India (West Bengal as escape, usually cultivated in the gardens) ; cultivated as pot plant throughout the world in tropics.

S. praetermissa Alston, Journ. Bot. 70 : 65. 1932.

Distrib. : India (Arunachal Pradesh), Sri Lanka.

S. proniflora (Lam.) Bak. Journ. Bot. 22 : 156. 1885. *Lycopodium proniflorum* Lam. Encycl. 3 : 652. 1789. *Selaginella dalzellii* Bak. Journ. Bot. 23. 249 : 1885.

Distrib. : India (Maharashtra, Karnataka), Sri Lanka, Malay region to North and East Australia.

S. pubescens (Wall. ex Hook. & Grev.) Spring, Bull. Acad. Brux. 10 : 225. 1843. *Lycopodium pubescens* Wall. ex Hook. & Grev. in Hook. Bot. Misc. 2 : 383. 1831.

Distrib. : India (Assam), Siam, Indo China.

S. pulvinata (Hook. & Grev.) Maxim. Mém. Acad. Imp. Sci. Petersb. 9 : 335. 1859. *Lycopodium pulvinatum* Hook. & Grev. in Hook. Bot. Misc. 2 : 381. 1831. *Selaginella tamarscina* (P. Beauv.) Spring var. *pulvinata* (Hook. & Grev.) Alston, Bull. Fan Mem. Inst. Biol. 5 : 271. 1934.

Distrib. : India (Kumaon), Burma, China.

S. radicata (Hook. & Grov.) Spring, Mém. Acad. Sci. Bégl. 24 (2) : 114. 1850. *Lycopodium radicatum* Hook. & Grev. in Hook. Bot. Misc. 3 : 397. 1831.

Distrib. : India (Arunachal Pradesh, Tamil Nadu).

S. rajasthanensis Gena, Bharadwaja & Yadava, Amer. Fern Journ. 69 (4) : 119. 1979.

Distrib. : India (Rajasthan).

S. repanda (Desv. & Poir.) Spring, Gaud. Voy. Bonite Bot. 1 : 329. 1846. *Lycopodium repandum* Desv. & Poir. in Lamarck, Encycl. Suppl. 3 : 558. 1814. *S. implexa* J. Scott, Journ. Agr. Hort. Soc. India N. S. 1 (2) : 262. 1868.

Distrib. : India (Uttar Pradesh, Madhya Pradesh, Andhra Pradesh, Karnataka, Tamil Nadu), Nepal, Burma, China, Philippine Islands, Java, Sumatra, Tonkin, Annam, Laos, Cambodia, Cochin China, Siam.

S. reticulata (Hook. & Grev.) Spring, Bull. Acad. Brux. 10 : 233. 1843. *Lycopodium reticulatum* Hook. & Grev. in Hook. Bot. Misc. 2 : 402. 1831.

Distrib. : India (Arunachal Pradesh, Meghalaya), Burma.

S. semicordata (Wall. ex Hook. & Grev.) Spring, Mart. Fl. Bras. 1 (2) : 122. 1840. *Lycopodium semicordatum* Wall. ex Hook. & Grev. in Hook. Bot. Misc. 2 : 396. 1831.

Distrib. : India (Sikkim, Arunachal Pradesh, Assam, Tripura, Bihar), Burma, Bangladesh, Malaya and Malay Islands.

S. subdiaphana (Wall. ex Hook. & Grev.) Spring, Bull. Acad. Brux. 10 : 232. 1843. *Lycopodium subdiaphanum* Wall. ex Hook. & Grev. in Hook. Bot. Misc. 2 : 401. 1831.

Distrib. : India (Himachal Pradesh, Punjab, Uttar Pradesh, Meghalaya, Nagaland), Nepal.

S. tenera (Hook. & Grev.) Spring, Bull. Acad. Brux. 10 : 232. 1843. *Lycopodium tenerum* Hook. & Grev. in Hook. Bot. Misc. 2 : 400. 1831.

Distrib. : India (Tamil Nadu, Karnataka, Kerala).

S. tenuifolia Spring, Mém. Acad. Sci. Bélg. 24 (2) : 253. 1850.

Distrib. : India (Arunachal Pradesh, West Bengal) Burma, Laos, Siam.

S. uncinata (Desv. ex Poir.) Spring, Bull. Acad. Brux 10 : 141. 1843. *Lycopodium uncinatum* Desv. ex Poir. in Lam. Encycl. Suppl. 3 : 558. 1813.

Distrib. : India (West Bengal), China, usually cultivated as pot plant in the tropics.

S. vaginata Spring, Mém. Acad. Sci. Bélg. 24 (2) : 87. 1850. *S. imbricata* J. Scott, Journ. Agr. Hort. Soc. India (N. S.) 1 : 260. 1868 (excl. syn. Roxb. 1844).

Distrib. : Throughout India in mountainous regions up to 2000m Burma, China.

S. wallichii (Hook. & Grev.) Spring, Mart. Fl. Bras. 1 (2) : 124. 1840. *Lycopodium wallichii* Hook. & Grev. in Hook. Bot. Misc. 2 : 284. 1831.

Distrib. : India (Arunachal Pradesh, Assam, Nagaland, Manipur, Meghalaya), Burma, Malaya Peninsula, Malay Islands, Cambodia, Laos, Tonkin.

S. wattii Bak. Handb. Fern Allies 109. 1887.

Distrib. : India (Manipur), Burma.

S. wightii Hieron. Hedw. 39 : 319. 1900.

Distrib. : India (Tamil Nadu), Sri Lanka, Eastern Africa.

S. willdenovii (Desv. ex Poir) Bak. Gard. Chron. 783. 950. 1867.
Lycopodium willdenovii Desv. ex Poir. in Lam. Encycl. Suppl. 3 : 540, 552. 1814.

Distrib. : India (Arunachal Pradesh, Assam, Himalayas) Burma, Malay Peninsula, Malesian Islands.

ISOETACEAE Reichenb.

Isöetes L. Sp. Pl. 2 : 1100. 1753 ; Pant & Srivastava, Proc. Natn. Inst. Sci. India 28 B (3) : 242. 1962 ; Panigr. Biol. Mem. 6 (2) : 129. 1981.

Type : *I. locutris* L. Sp. Pl. 2 : 1100. 1753.

I. bilaspurensis Panigr. Biol. Mem. 6 (2) : 135. f. 1 - 10. 1981.

Distrib. : India (Madhya Pradesh).

I. coromandelina L. f. Suppl. Pl. Syst. veg. ed. II : 447. 1781.
 subsp. *coromandelina*.

Distrib. : India (Tamil Nadu, Kerala, Karnataka, Maharashtra, Gujarat, Madhya Pradesh, Uttar Pradesh, Bihar, Orissa and West Bengal).

I. coromandelina subsp. *brachyglossa* (A. Braun) Panigr. Biol. Mem. 6 (2) : 131. 1981. *I. brachyglossa* A. Braun, Verh. Bot. ver. Brandenb 4 : 328. 1862.

Distrib. : India (West Bengal).

I. dixitii Shende, Journ. Univ. Bombay 14 : 50. 1945.

Distrib. : India (Maharashtra).

I. indica Pant & Srivastava, Proc. Natn. Inst. Sci. India 28 B (3) : 246. t. XIV, f. 16 - 25, t. XV f. 30 - 35, t. XIV f. 30-40 text - fig. 4 A - D, 5 A - H, 7 A - M, 12. 1962. *I. mirzapurensis* Panigr. & Dixit, Proc. Indian Sci. Congr. pt. III, Abstract, Botany 243. 1965.

Distrib. : India (Uttar Pradesh, Madhya Pradesh, Tamil Nadu).

I. panchananii Pant & Srivastava, Proc. Natn. Inst. Sci. India 28 B (3) : 282 t. XIII, f. 1 - 4. f. XVI t. 41. text - fig. 1. 2 A - I. 1962.

Distrib. : India (Madhya Pradesh, Maharashtra).

I. pantii Goswami & Arya, Journ. Indian Bot. Soc. 49 : 30. 1970.

Distrib. : India (Madhya Pradesh).

I. sahyadriensis Mahabale, Curr. Sci. 7 (2) : 62. f. 1. 1938.

Distrib. : India (Maharashtra, Tamil Nadu).

I. sampathkumarnii L. N. Rao, Curr. Sci. 13 : 286. f. 1 - 3. 1944.

Distrib. : India (Karnataka, Andhra Pradesh).

I. unilocularis J. E. Smith in Rees, Cyclopaedia 19 : 150. 1819,
pro parte includo type et descr. excludo spec. labelled *Isöetes indica*
 Koenig. *I. coromandelina sensu* Roxb. ex Griff. Calc. Journ. Nat. Hist
 4 : 470. 1844.

EQUISETACEAE L. C. Richard ex De Candolle

Equisetum L. Sp. Pl. 2 : 1050. 1753 ; Clarke, Trans. Linn. Soc.
 Lond. II. Bot. 1 : 594. 1880 ; Hauke, Nova Hedw. 8 : 1. 1963 ; Page,
 Fern Gaz. 11 (i) : 29. 1974 ; Hauke, Nova Hedw. 30 : 385. 1978.

Type : **E. arvense** L. Sp. Pl. 2 : 1050. 1753.

E. arvense L. Sp. Pl. 2 : 1050. 1753.

Distrib. : India (Himalayas), throughout Canada, America except
 S. E. part, Europe, Asia South to Turkey, Iran, Nepal, China, Korea,
 Japan.

E. diffusum D. Don, Prod. Fl. Nepal. 19. 1825.

Distrib. : India (Himalayan mountains from Simla to Tibet, Sikkim,
 Assam. Meghalaya), Nepal, Bhutan, Burma, China.

E. giganteum L. Sp. Pl. ed. II. 2 : 1517. 1763.

Distrib. : India (Kashmir), Cuba, Jamaica, Hispaniola, Central &
 S. America.

E. hymale L. Sp. Pl. 2 : 1062. 1753.

Distrib. : India (Kumaon, Kashmir ?), Iceland, Europe (except
 Spain) Algeria, Asia, Russia, China.

E. palustre L. Sp. Pl. 2 : 1061. 1753.

Distrib. : India (Kashmir) N. America, South to New York,
 Michigan, Illinois, Minnesota, N. Dakota, Montesa, Idaho, California,
 Eurasia, South to Mediterranean, Greece, Turkey, Tibet, China, Japan.

E. ramossissimum Desf. Fl. Atlant. 2 : 398. 1800. subsp. **ramossissimum**.

Distrib. : North India and tropics of the world.

E. ramosissimum Desf. subsp. **debile** (Roxb. ex Vauch.) Hauke. Amer. Fern Journ. 52 : 33. 1962. *E. debile* Roxb. ex Vauch. Mon. Preles 387. 1821.

Distrib. : India (Himalayas) Nepal, Burma, S. China, Formosa, Hainan, Indo China, Philippines, Indonesia, New Guinea, New Hebrides, New Caledonia, Fiji.

PSILOTACEAE Kanitz

Psilotum Sw. Schrad. Journ. Bot. 1800 (2) : 8, 109. 1801. Thothathri, Banerjee, Hazra & Pal, Bull. Bot. Surv. India 12 (1 - 4) : 280. f. 1 - 3. 1970.

Type : **P. nudum** (L.) P. Beauv. Prod. Fam. Aéthog. 112. 1805.

P. complanatum Sw. Syn. Fil. 188, 415. t. IV. f. 5. 1806.

Distrib. : India (Nicobar Islands) Malay Peninsula, Malesian Islands, South Australia, Hawaii, Fiji Islands, Mexico and South America.

P. nudum (L.) P. Beauv. Prod. Fam. Aétheog. 112. 1805. *Lyco-podium nudum* L. Sp. Pl. 2 : 1100. 1753. *Psilotum triquetrum* Sw. Schrad. Journ. Bot. 1800. (2) : 109. 1801.

Distrib. : Throughout India in hilly regions, Burma, Malaysia.

HELMINTHOSTACHYACEAE Ching

Helminthostachys Kaulf. Enum. Fil. 103. 1822 ; Clausen, Mem. Torrey Bot. Club. 19 (2) : 108. 1938 ; Panigr. & Dixit, Proc. Natn. Inst. Sci. India 35 B (3) : 245. 1969.

Type : **H. zeylanica** (L.) Hook. Gen. Fil. t. 47. 1840.

H. zeylanica (L.) Hook. Gen. Fil. t. 47. 1840 ; Bedd. Handb. Ferns Brit. India 467. t. 292. 1883 ; Suppl. 109. 1892. *Osmunda zeylanica* L. Sp. Pl. 2 : 1063. 1753.

Distrib. : India (Bengal plains to Assam, Himalayas, Uttar Pradesh, South India) Sri Lanka, Malay Peninsula, China, Japan, Philippines, Solomon Islands, New Caledonia, New Guinea, Australia.

BOTRYCHIACEAE Nakai

Botrychium Sw. Schrad. Journ. Bot. 1800. (2) : 8, 110. 1801 ; Clausen, Mem. Torrey Bot. Club 16 (2) : 60. 1938 ; Panigr. & Dixit, Proc. Natn. Inst. Sci. India 35 B (3) : 237. 1969.

Lectotype : **B. lunaria** (L.) Sw. Schrad. Journ. Bot. 1800 (2) : 110. 1801.

B. daucifolium Wall. ex Hook. & Grev. Icon. Fil. t. 161. 1829 ; Bedd. Handb. Ferns. Brit. India 469. t. 294. 1883.

Distrib. : Throughout mountainous regions of India up to 2400 m., Burma, China, Sri Lanka, Malesian Islands, Philippines, Fiji.

B. daucifolium Wall. ex Hook. & Grev. var. **parvum** v. Ald. v. **Ros.** in Bull. Jard. Bot. Buit. 1 : 3. 1911.

Distrib. : India (Himalayas), Java.

B. lanuginosum Wall. ex Hook. & Grev. Icon. Fil. 1. t. 79. 1831. var. **lanuginosum**. **B. virginianum** (L.) Sw. var. **lanuginosum** (Wall. ex Hook. & Grev.) Bedd. Handb. Ferns Brit. India 471. t. 295. 1883 ; Suppl. 110. 1892.

Distrib. : Throughout India in mountainous regions, Nepal, Sri Lanka, Sumatra, Java, Philippines, China.

B. lanuginosum Wall. ex Hook. & Grev. var. **nepalense** (Nishida) Nair & Dixit, Journ. Bomb. Nat. Hist. Soc. 78 (3) : 447. 1981. *Japanobotrychium lanuginosum* (wall. ex Hook. & Grev.) Nishida ex Tagawa var. **nepalense** Nishida, Journ. Jap. Bot. 41 : 319. 1966.

Distrib. : India (Himalayas).

B. lunaria (L.) Sw. Schrad. Journ. Bot. 1800 (2) : 110. 1801 ; Bedd. Handb. Ferns Brit. India 469. f. 293. 1883 ; Suppl. 110. 1892. var. **lunaria**. *Osmunda lunaria* L. Sp. Pl. 2 : 1064. 1753.

Distrib. : India (N. W. Himalayas, Sikkim, E. Himalayas, South India) Europe, Afghanistan, United States, Greenland, Japan, Patagonia, New Zealand, Australia and Tasmania.

B. lunaria (L.) Sw. var. *onondaganse* (Underw.) House, Bull. N. Y. State Mus. nos. 243 244, p. 47. 1923. *B. onondaganse* Underw. Bull. Torrey Bot. Club 30 : 47. 1903.

Distrib. : India (N. W. Himalayas), Tibet, China.

B. multifidum (Gmel.) Rupr. Beitr. Zur. Pflanz. des Russ. Reich. 11 : 40. 1859. ssp. **multifidum**. *Osmunda multifida* Gmel. Nov. Comment. Acad. Petrop. 12 : 517. 1768.

Distrib. : India (Sikkim), Northern Europe, China (Yunnan), North America, British Columbia.

B. ternatum (Thunb.) Sw. Schrad. Journ. Fur di Botanik 2 : 111. 1800 ; Bedd. Handb. Ferns. Brit. India Suppl. 110. 1892. *Osmunda ternata* Thunb. Fl. Jap. 329. t. 32. 1784.

Distrib. : India (North West to Eastern Himalayas), Japan, Korea, China.

B. virginianum (L.) Sw. Schrad. Journ. Fur di Botanik 2 : 111. 1800. ssp. *virginianum*. *Osmunda virginiana* L. Sp. Pl. 2 : 1064. 1753.

Distrib. : India (North West and Eastern India) China, Japan, Europe and North America.

OPHIOGLOSSACEAE (R. Br.) Agardh

Ophioderma (Bl.) Endl. Gen. Pl. 66. 1836.

Type : *O. pendula* (L.) Presl, Suppl. Tent. Pterid. 56. 1845.

O. pendula (L.) Presl., Suppl. Tent. Pterid. 56. 1845. f. **pendulum** *Ophioglossum pendulum* L. Sp. Pl. ed. II. 2 : 1518. 1763 ; Bedd. Handb. Ferns Brit. India 465. t. 291. 1883 ; Suppl. 109. 1892.

Distrib. : India (Assam, Andaman & Nicobar Islands), Old World tropics in Madagascar, Sri Lanka, Malay Peninsula, Formosa, Philippines, Sumatra, Java, Borneo, Moluccas, New Guinea, Caroline Islands, Australia, New Hebrides, Fiji, Samoa, Tahiti and Hawaiian Islands.

Ophioglossum L. Sp. Pl. 2 : 1062. 1753 ; Clausen, Mem. Torrey Bot. Club 19 (2) : 161. 1938 ; Chakravarty, Bull. Bot. Soc. Bengal 5 : 1. 1951 ; Mahabale, Bull. Bot. Surv. India 4 : 71. 1962 ; Panigr. & Dixit, Proc. Natn. Inst. Sci. India 35 B (3) : 230. 1969.

Lectotype : *O. vulgatum* L. Sp. Pl. 2 : 1062. 1753.

O. costatum R. Br. Prod. Fl. Nov. Holl. 163. 1810. *O. pedunculosum* Desv. Mag. Ges. Nat. Fr. Berlin 5 : 306. 1811. *O. fibrosum* Schum, Beskr. Guin. Pl. 452. 1827 ; Bedd. Handb. Ferns Brit. India 465. t. 289. 1883 ; Suppl. 109. 1892. *O. brevipes* Bedd. Ferns South. India 23. t. 72. 1863. *O. bulbosum* Bedd. Ferns Brit. India Suppl. 28. 1876.

Distrib. : India (North West to Eastern Himalayas, Madhya Pradesh, West Bengal, Maharashtra, South India), Africa, Comores Archipelago, Madagascar, Sri Lanka, Sumatra to East Australia and New Zealand.

O. gramineum Willd. Nov. Act. Acad. Erf. 2 : 18 t. 1. f. 1. 1802 ; Bedd. Handb. Ferns Brit. India Suppl. 108. 1892. var. *gramineum*.

Distrib. : India (Uttar Pradesh, Madhya Pradesh, South India), pantropical.

O. gramineum var. *majus* (v. Ald. v. Ros.) Wieff. Blumea 12 (2) : 324. f. 2 b, t. 3. 1964.

Distrib. : India (Tripura), Philippines, New Guinea.

O. lusitanicum L. Sp. Pl. 2 : 1063. 1753. ssp. *lusitanicum*.

Distrib. : India (Madhya Pradesh, Tamil Nadu) Iceland, France, Portugal, Italy, Azores, Madeira, Algeria, Afghanistan, Asia.

O. nudicaule L. f. Suppl. Syst. t. 443. 1781 ; Bedd. Handb. Ferns Brit. India 464. t. 288. 1883 ; Suppl. 109. 1892. *O. parvifolium* sensu Bedd. Ferns South. India 23. t. 71. 1863.

Distrib. : Pantropic.

O. nudicaule var. *macrorrhizum* (Kunze) Clausen, Mem. Torrey Bot. Club 19 (2) : 150. 1938. *O. macrorrhizum* Kunze, Farrnkr. 1 : 57. t. 29. f. 1. 1840.

Distrib. : India (Madhya Pradesh, Gujarat, West Bengal, Uttar Pradesh, Tamil Nadu), China, Thailand, Malay Peninsula, Sumatra, Java, Lesser Sunda Island, Philippines, New Guinea, New Calendonia.

O. petiolatum Hook. Exot. Fl. 1 : t. 56. 1823.

Distrib. : India (Uttar Pradesh, Madhya Pradesh), Central Florida, Trinidad, West Indies, Mexico, America, Africa, Sri Lanka, Siam, China, Japan, Philippines, Borneo, Sumatra, Java, New Guinea, New Caledonia, New Zealand, Fiji and Samoa.

O. polypodium A. Braun apud Seubert, Fl. Azor. 17. 1844. *O. aitchisoni* (Clarke) d' Almeida, Journ. Indian Bot. Soc. 3 : 63. f. 12 13.

1922 ; *O. lusitanicum* sensu Hope, Journ. Bomb. Nat. Hist. Soc. 15 : 106. 1903.

Distrib. : India (N. W. Himalayas), Ethiopia, Kenya, Tanganyka, Rhodesia, Africa, Cape Verde Islands, Arabia.

O. reticulatum L. Sp. Pl. 2 : 1063. 1753 ; Bedd. Ferns South. India 23 t. 70. 1863 ; Handb. Ferns Brit. India 465. t. 291. 1883 ; Suppl. 109. 1892. f. ***reticulatum***. *Ophioglossum cordifolium* Roxb. ex Griff. Calc. Journ. Nat. Hist. 4 : 475. 1844.

Distrib : Throughout India from the plains to mountainous regions, pantropic.

O. reticulatum L. f. ***complicatum*** (Miq.) Wieff. Blumea 12 : 330. f. 1. a 1964. *Ophioglossum moluccanum* f. *complicatum* Miq. Ann. Mus. Bot. Lugd. Bat. 4 : 290. 1868.

Distrib. : India (Eastern India, Sikkim), Japan, Thailand, Bonin Island, Sumatra, Malay Peninsula, Java, Lesser Sunda Island, Borneo, Philippines, Moluccas, New Guinea. New Caledonia.

O. reticulatum L. f. ***dilatatum*** (Miq.) Wieff. Blumea 12 : 329. f. I. C. 1964. *O. moluccanum* f. *dilatatum* Miq. Ann. Mus. Bot. Lugd. Bat. 4 : 92. 1868.

Distrib. : India (N. W. Himalayas) Sumatra, Malay Peninsula, Java, Lesser Sunda Island, Borneo, Philippines, Fiji Islands.

O. thermale Kumarov var. ***nipponicum*** (Miyabe & Kudo) Nishida in Tagawa, Journ. Jap. Bot. 33 : 202. 1958. *O. nipponicum* Miyabe & Kudo, Trans. Sapporo Nat. Hist. Soc. 6 : 122. 1916. *O. japonicum* Prantl, Ber. deut. bot. Ges. 1 : 353. 1883 ; Mahabale, Bull. Bot. Surv. India 4 : 71. 1962.

Distrib. : India (E. Himalayas), throughout Japanese Islands, China, Korea, Japan.

O. vulgatum L. Sp. Pl. 2 : 1062. 1753 ; Bedd. Handb. Ferns Brit. India 464. 1883 ; Suppl. 109. 1892.

Distrib. : India (N. W. Himalayas), widely distributed in temperate zones on the whole northern hemisphere.

ANGIOPTERIDACEAE Féé ex Bonner

Angiopteris Hoffm. Comm. Soc. Reg. Sci. Gotting 12 : 29. 1793 94 ; de Vriese, Mon. 26. 1853 ; Dixit & Panigr. Bull. Bot. Surv. India 10 (3 & 4) : 368. 1969.

Type : **A. evecta** (Forst.) Hoffm. Comm. Soc. Reg. Sci. Gotting 12 : 29. 1793 94.

A. arnottiana Miquel, verh. Nederl. Inst. III. 4 : 53. t. 7. f. A. 1851.

Distrib. : India.

A. assamica de Vriese, Mon. 33. 1853.

Distrib. : India (Assam).

A. campsophlebia de Vriese, Mon. 31. 1853.

Distrib. : India.

A. crassipes Wall. ex Presl, Suppl. Tent. Pterid. 23. 1845.

Distrib. : India (Himalayas), Nepal, Burma, China, Australia.

A. distans Presl, Suppl. Tent. Pterid. 23. 1845.

Distrib. : India.

A. evecta (Forst.) Hoffm. Comm. Soc. Reg. Gott. 12 : 29 : t. 5. 1793-94 ; Bedd. Ferns South. India t. 78. 1863 ; Handb. Ferns Brit. India 460. t. 285. 1883. *Polypodium evectum* Forst. Prod. 81. 1786.

Distrib. : Throughout India in hilly regions, China, Japan, Tropical Australia, New Caledonia, Madagascar, Malaysia to Polynesia.

A. hookeriana d. Vriese, Mon. 29. 1853.

Distrib. : India.

A. hugeliana Presl, Suppl. Tent. Pterid. 25. 1845.

Distrib. : India.

A. laciniata de Vriese, Mon. 30. 1853.

Distrib. : India (N. W. Himalaya).

A. latifolia Presl, Suppl. Tent. Pterid. 24. 1845.

Distrib. : India (Pundoa).

A. macrocephala Presl, Tent. Pterid. 24. 1845.

Distrib. : India (Punjab).

A. repandula de Vriese, Mon. 30. t. 3. f. 4. t. 4. f. 4. 1853.

Distrib. : India (Saharnpur).

A. salicifolia (Presl) de Vriese, Mon. 34. 1853. *Psilodochea salicifolia* Presl, Suppl. Tent. Pterid. 28. 1845.

Distrib. : India (E. Himalayas), Pakistan.

A. wallichiana Presl, Suppl. Tent. Pterid. 22. 1845.

Distrib. : India (Himalayas).

A. wightiana de Vriese, Mon. 28. 1853.

Distrib. : S. India.

MARATTIACEAE Berch. et Presl

Marattia Sw. Prod. Fl. Ind. Occ. 128. 1788 ; Dixit & Panigr. Bull. Bot. Surv. India 10 (3 & 4) : 370. 1969.

Type : **M. alata** Sw. Prod. Fl. Ind. Occ. 128. 1788.

M. fraxinea Smith, Pl. Ic. in ed. 2. t. 48. 1790; Bedd. Ferns South. India 27. t. 79. 1863 ; Handb. Ferns Brit. India 460. t. 286. 1883.

Distrib. : South India, Sri Lanka, tropics of the world and a little beyond in the Southern zone.

CHRISTENSENIACEAE Ching

Christensenia Maxon, Proc. Biol. Soc. Wash. 18 : 239. 1905 ; Dixit & Panigr. Bull. Bot. Surv. India 10 (3 & 4) : 370. 1969.

Type : **C. aesculifolia** (Bl.) Maxon, Proc. Biol. Soc. Wash. 18 : 239. 1905.

C. aesculifolia (Bl.) Maxon, Proc. Biol. Soc. Wash. 18 : 239. 1905. *Aspidium aesculifolium* Bl. Enum. Pl. Jav. 2: 143. 1828. *Kaulfussia aesculifolia* (Bl.) Bl. Enum. Pl. Jav. 2 : 260. 1818 ; Bedd. Ferns Brit. India t. 185. 1865 ; Handb. Ferns. Brit. India 462. t. 267. 1883. *K. assamica* Griff. Asiat. Research. 19 : 108. t. 18. 1836.

Distrib. : India (Assam, Meghalaya, Arunachal Pradesh), Bangla Desh, Malay Islands, Philippines.

OSMUNDACEAE Berch. & Presl

Osmunda L. Sp. Pl. 2 : 1063. 1753. Panigr. & Dixit, Journ. Indian Bot. Soc. 43 (1 & 2) : 90. 1969.

Lectotype : **O. regalis** L. Sp. Pl. 2 : 1063. 1753.

O. cinnamomea L. Sp. Pl. 2 : 1066. 1753.

Distrib. : India (Arunachal Pradesh), China, Japan, Mexico, Brazil, Manchuria.

O. claytoniana L. Sp. Pl. 2 : 1066. 1753 ; Bedd. Handb. Ferns Brit. India 449. t. 275. 1883.

Distrib. : India (Himalayas from Kashmir to Eastern Himalayas), America, China.

O. claytoniana var. **vestita** Wall. ex Milde, Mon. Gen. Osmunda 102. 1868.

Distrib. : India (Eastern Himalayas).

O. japonica Thunb. Fl. Jap. 330. 1784.

Distrib. : India (Assam, Meghalaya, Uttar Pradesh), Bhutan.

O. regalis L. Sp. Pl. 2 : 1065. 1753 ; Bedd. Ferns South. India 26. t. 76. 1863 ; Handb. Ferns Brit. India 550. t. 276. 1883.

Distrib. : India (Assam, Meghalaya, Uttar Pradesh, Madhya Pradesh, Tamil Nadu, Kerala, Maharashtra, Karnataka), Pantropic.

PLAGIogyriaceae Bower

Plagiogyrea (Kunze) Mett. Abhandl. d. Senckneb. naturf. Ges. 2 : 265. 1858 ; Ching, Acta Phytotax. Sin. 7 (2) : 105. 1958 ; Dixit & Das, Proc. Indian Acad. Sci. (Plant Sci.) 90 (4) : 371. 1981.

Lectotype : **P. triquetra** Wall. ex Mett. Abhd. Senckneb. naturf. Ges. 2 : 265. 1858.

P. chinensis Ching, Acta Phytotax. Sin. 7 (2) : 140. t. 30. f. 2. 1958.

Distrib. : India (Meghalaya), China.

P. christii Copel. Philip. Journ. Sci. Suppl. 1 : 153. 1906.

Distrib. : India (Meghalaya), China.

P. coerulescens Ching, Acta Phytotax. Sin. 7(2) : 148. t. 36. f. 2. 1958.

Distrib. : India (West Bengal, Meghalaya), China.

P. decrescens Ching, Acta Phytotax. Sin. 7(2) : 146. t. 33. f. 2. 1958.

Distrib. : India (West Bengal, Sikkim), China.

P. distanta Dixit & Das, Proc. Indian Acad. Sci. (Plant Sci.) 90 (4) : 486. f. 16. 1981. *P. glaucescens* Ching var. *arguta* Ching, Acta Phytotax. Sin. 7 (2) : 151. 1958.

Distrib. : India (Nagaland), China (Yunnan).

P. elongata Dixit & Das, Proc. Indian Acad. Sci. (Plant Sci.) 90 (4) : 377. f. 5. 1981.

Distrib. : Eastern India.

P. euphlebia (Kunze) Mett. Abhandl. d. Senckneb. Natur. Ges. 2 : 274. 1858 ; Bedd. Ferns Brit. India 1866. (*pro parte* excl. t. 165) ; Handb. Ferns Brit. India 129. 1883.

Distrtb. : India (Assam, Meghalaya), China, Japan, Korea.

P. gigantea Ching, Lingnan Sc. Journ. 15 : 275. 1936.

Distrib. : Eastern India, China.

P. glaucescens Ching, Acta Phytotax. Sin. 7 (2) : 150. t. 38. f. 2. 1958. *P. glauca* sensu auct. Bedd. Ferns. Brit. India t. 90. 1865 (*descr. pro parte*) ; Handb. Ferns Brit. India 127. 1883 (*descr. pro parte*) ; Panigr. Bull. Bot. Surv. India 2 : 313. 1960. *Lomaria glauca* sensu auct. Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 472. 1880, *pro parte* (*non* Bl. 1828).

Distrib. : India (Assam, Manipur, Meghalaya), China, Burma.

P. lanuginosa Ching, Acta Phytotax. Sin. 7 (2) : 147. t. 35. f. 1. 1958.

Distrib. : India (Arunachal Pradesh), China.

P. media Ching, Acta Phytotax. Sin. 7 (2) : 149. t. 38. f. 1. 1958.

Distrib. : Eastern India, Burma, China.

P. meghalayensis Dixit & Das, Proc. Indian Acad. Sci. (PlantSci.) 90 (4) : 378. f. 7. 1981.

Distrib. : India. (Meghalaya).

P. minglingensis Dixit & Das, Proc. Indian Acad. Sci. (Plant Sci) 90 (4) : 382. f. 12. 1981.

Distrib. : Eastern India.

P. rankanensis Hayata, Icon Pl. Form. 8 : 150. f. 80. 1919. *Lomaria adnata* *sensu* auct. Hook. Sp. Fil. 3 : 19. t. 147. 1860 (*pro parte*). *P. adnata* *sensu* Bedd. Ferns Brtt. India t. 51. 1965 ; Handb. Ferns Prit. India 157. t. 65. 1883 (*non* Bl. 1828).

Distrib. : Eastern India, Burma, China, Japan.

P. scandens Mett. Abhandl. d. Senkenb. natur f. Ges. 2 : 273. t. 4. f. 21. 1858. *P. pycnophylla* *sensu* Bedd. Ferns Brit. India. t. 52. 1865 ; Handb. Ferns Brit. Indian 129. 1883 (*pro parte*). *P. communis* Ching, Acta Phytotax. Sin. 7 (2) : 147. t. 5. f. 2. 1958.

Distrib. : India (Eastern India & Eastern Himalayas), Burma, China.

P. triquetra Wall. ex Mett. Abhandl. d. Senckneb. naturf. Ges. 2. 274. 1858. *P. euphlebia* *sensu* Bedd. Ferns Brit. India t. 165. 1866 (*Descr. pro parte*).

Distrib. : Eastern India, Nepal.

GLEICHENIACEAE (R. Br.) Presl

Gleichenia J. Sm. Mém. Acad. Sc. Turin. 5 : 419. t. 9. f. 10. 1793 ; Panigr. & Dixit, Bull. Bot. Surv. India 10 : 337. 1968 ; Holtt. Fl. Males. ser. II. Pterid. 1 (5) : 1. 1959.

Type : **G. polypodioides** (L.) Sm. Mém. Acad. Sc. Turin 5 : 419. t. 9. f. 10. 1793.

G. blotiana C. Chr. Bull. Mus. Hist. Nat. Paris II. 6 : 103. 1934.

Distrib. : Eastern India, Indo China, South China, Malay Peninsula.

G. gigantea Wall. ex Hook. Sp. Fil. 2 : 5. t. 3A. 1844 ; Bedd. Ferns Brit. India t. 30. 1865. *G. glauca* auct. Hook ; Bedd. Handb. Ferns Brit. India 2. 1883. *quoad* pl. Ind. (*non* Thunb.).

Distrib. : North - East India, Burma, China, Vietnam, North Thailand, Malay Peninsula.

G. laevissima Christ, Bull. Acad. Inst. Geogr. Bot. III. 11. 268. 1902.

Distrib. : Eastern India, China, Formosa, Malaysia, Philippines.

G. longissima Bl. Enum. Pl. Jav. 2 : 250. 1828. *G. glauca sensu* Bedd.
Handb. Ferns Brit. India 2. t. 1. 1883.

Distrib. : Eastern India and Eastern Himalayas, Nepal, Bhutan,
China, Japan, Malaya, Malesian Islands, America, Polynesia, Australia.

G. volubilis Jungh. Java 1 : 592, 664. 1893.

Distrib. : India (Assam, Meghalaya, Eastern Himalayas), Indochina,
Formosa, Philippines, Malay Islands, New Guinea.

DICRANOPTERIDACEAE Ching

Dicranopteris Bernh. Schrad. Journ. Bot. 1 (2) : 38. 1805 ; Holtt.
Fl. Males ser. II. Pterid. 1 : 27. 1959 ; Panigr. & Dixit, J. Sen Mem. Vol.
469. 1969.

Type : **D. dichotoma** (Thunb. ex Murray) Bernh. Schrad. Journ.
Bot. 1 (2) : 38, 49. t. 3. f. 13. 1805.

D. curranii copel. Philip. Journ. Sci. 81 : 4. 1952.

Distrib. : India (Nicobar Islands), Malay Peninsula, Malay Islands.

D. linearis (Burm. f.) Underw. Bull. Torrey Bot. Club 34 : 250.
1907. var. *linearis*. *Polypodium linearis* Burm. f. Fl. Ind. 235. t. 67. f. 2.
1768. *Gleichenia linearis* (Burm. f.) Clarke, Trans. Linn. Soc. Lond. II.
Bot. 1 : 428. 1880.

Distrib. : Widely distributed in the tropics and subtropics of the
world.

D. linearis (Burm. f.) Underw. var. *altissima* Holtt. Reinw. 4 : 276.
1957.

Distrib. : India (Arunachal Pradesh, Assam, Meghalaya, Madhya
Pradesh, Tamil Nadu), Malay Peninsula, Philippines, Moluccas, New
Guinea, Solomon Islands.

D. linearis (Burm. f.) Underw. var. *demota* Holtt. Reinw. 4 : 272
1957.

Distrib. : India (Eastern Himalayas) Nepal, Malaysia, North Borneo,
New Guinea.

D. linearis (Burm. f.) underw. var. *latiloba* Holtt. Reinw. 4 : 277.
1957.

Distrib. : India (Arunachal Pradesh, Assam, Meghalaya) Malaysia,
Philippines, Celebes.

D. linearis (Burm. f.) underw. var. **montana** Holtt. Reinw. 4 : 276. 1957.

Distrib. : India (Himalayas, S. India), Malaysia.

D. linearis (Burm. f.) Underw. var. **sebastiana** Panigr. & Dixit. Bull. Bot. Surv. India 13 (1 & 2) : 162. 1971.

Distrib. : India (Tamil Nadu Nilgiri hills).

D. linearis (Burm. f.) Underw. var. **subferruginea** (Hieron. ex Brause) Nakai, Bull. Nat. Sci. Mus. Tokyo no. 29. 66. 1950. *Gleichenia linearis* var. *subferruginea* Hieron. ex Brause, Bot. Jahrb. 56 : 209. 1920.

Distrib. : India (Eastern Himalayas) Queensland, Polynesia, East Malaysia, New Guinea.

D. linearis (Burm. f.) Underw. var. **subpectinata** (Christ) Holtt. Reinw. 4 : 277. 1957.

Distrib. : India (Himalayas, Andaman Islands), Sumatra, Malaya Peninsula, Lingga Archipelago, Banka, Borneo.

D. linearis (Burm. f.) Underw. var. **wattii** Panigr. & Dixit, Bull. Bot. Surv. India 13 (1 & 2) : 162. 1971.

Distrib. : India (Manipur).

GRAMMITACEAE (Presl) Ching

Ctenopteris Bl. ex Kunze, Bot. Zeit 425. 1846 ; Sledge, Bull. Brit. Mus. (N. H.) 4 : 155. 1966.

Lectotype : **C. venulosa** (Bl.) Bl. ex Kunze, Bot. Zeit. 4 : 425. 1846.

C. subfalcata (Bl.) Kunze, Bot. Zeit. 120. 1848. *Polypodium sub-falcatum* Bl. Enum. Pl. Jav. 2 : 130. 1828 ; Bedd. Handb. Ferns Brit. India 314. 1883.

Distrib. : Throughout India in hilly regions, Malaya.

C. trichomanoides J. Sm. Hist. Fil. 184. 1875. *Polypodium trichomanoides* Sw. Prod. 131. 1788 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 549. 1880 ; Bedd. Handb. Ferns Brit. India 308. t. 162. 1883.

Distrib. : India (Sikkim), South America, Brazil, Ecuador, Cuba.

Grammitis Sw. Schrad. Journ. Bot. 1800 (2) : 3, 17. 1801.

Lectotype : **G. marginalis** (Sw.) Sw. Schrad. Journ. Bot. 1800 (2) : 17. 1801.

G. attenuata Kunze, Linnaea 24 : 251. 1851. *Polypodium parasiticum* Mett. Abhandl. Senekneb Naturf. Ges. 2 : 36. 1856 ; Bedd. Ferns South. India 55. t. 165. 1864 ; Handb. Ferns Brit. India 302. 1883 (*pro parte, majore*).

Distrib. : South India, Sri Lanka.

G. medialis (Bak.) Sledge, Bull. Brit. Mus. (N. H.) London 2 : 149. 1960. *Polypodium mediale* Bak. in Hook. & Bak. Syn. Fil. ed. 2. 507. 1874. *P. parasiticum sensu* Bedd., Handb. Ferns Brit. India 302. 1883. (*pro parte, minore*, non Mett.)

Distrib. : South India, Sri Lanka.

Prosaptia Presl, Tent. Pterid. 165. t. 6. f. 19. 25. 1836 ; Sledge, Bull. Brit. Mus. (N. H.) 2 : 157. 1960.

Lectotype : **P. contigua** (G. Forst) Presl, Tent. Pterid. 166. 1836.

P. alata (Bl.) Christ, Ann. Jard. Bot. Buit. II. 5 : 127. 1905. *Davallia alata* Bl. Enum. Pl. Jav. 2 : 230. 1828.

Distrib. : South India, Sri Lanka, Sumatra to Polynesia.

P. contigua (G. Forst.) Presl, Tent. Pterid. 166. 1836 ; Bedd. Ferns South. India 6. t. 19. 1863 ; Handb. Ferns Brit. India 56. 1883. *Trichomanes contiguum* G. Forst. Prod. 84. 1786.

Distrib. : South India, Sri Lanka, Sumatra to Polynesia.

P. khasyana (Hook.) C. Chr. & Tard. in Natur. Syst. 8 : 180. 1939. *Polypodium khasyanum* Hook. Ic. Pl. 10. t. 949. 1854 ; Bedd. Ferns Brit. India t. 173. 1866 ; Handb. Ferns Brit. India 308. 1883.

Distrib. : Eastern India, Burma, Hainan, Annam, Cambodia, Malaya, Borneo, Celebes, Sri Lanka.

P. obliquata (Bl.) Mett. in Reise Osterr. Novara Bot. 1 : 214. 1870. *Polypodium obliquatum* Bl. Enum. Pl. Jav. 2 : 128. 1828 ; Bedd. Handb. Ferns Brit. India 311. 1883.

Distrib. : South India, Sri Lanka, Sumatra to Polynesia.

Scleroglossum V. Ald. V. Ros. Bull. Jard. Bot. Buit. ser. II. no. 7. 37. 1912.

Lectotype : *S. pusillum* (Bl.) v. A. v. R. Bull. Jard. Bot. Buit. ser. 2. no 7. 39. 1912.

S. sulcatum (Kuhn) v. A. v. R. Bull. Jard. Bot. Buit. ser. 2. no. 7. 39. 1912. *Taeniopsis falcata* Bedd. Ferns Brit. India t. 175. 1866. (*pro parte*; *non Vittaria falcata* Fée). *Vittaria sulcata* Kuhn, Linnaea 36 : 68. 1869 ; Bedd. Handb. Ferns Brit. India 408. 1883. *Taeniopsis sulcata* (Kuhn) Bedd. Ferns Brit. India, Suppl. 25. 1876; Ferns Brit. India t. 175. 1866.

Distrib. : India (Meghalaya), Sri Lanka.

Xiphopteris Kaulf, Berlin Jahrb. Pharm. 21 : 35. 1820.

Lectotype : *X. serrulata* (Sw.) Kaulf. Enum. Fil. 85 : 1824.

X. sikkimensis (Hieron.) Copel. Gen. Fil. 215. 1947. *Polypodium sikkimense* Hieron. Hedw. 44 : 97. 1905. *Polypodium trichomanoides* sensu Bedd. Handb. Ferns Brit. India 308. 1883.

Distrib. : India (Himalayas), W. China, Tonkin.

LOXOGRAMMACEAE Ching ex Pic. Ser.

Loxogramme (Bl.) Presl, Tent. Pterid. 214. t. 9. f. 8. 1836.

Lectotype : *L. lanceolata* (Sw.) Presl, Tent. Pterid. 215. t. 9. f. 8. 1836.

L. chinensis Ching, Sinensis 1 : 13. 1929.

Distrib. : India (Eastern Himalayas, Meghalaya, Arunachal Pradesh), China, Bhutan, Burma, Thailand.

L. duclouxii Christ, Bull. Acad. Géogr. Bot. 17 : 140. 1907.

Distrib. : India (Eastern India, and Himalayas), China, Japan.

L. grammitoides (Bak.) C. Chr. Ind. Fl. Suppl. 2 : 21. 1917. *Gymnogramme grammitoides* Bak. Journ. Bot. 27 : 178. 1889.

Distrib. : Himalayas to China and Japan.

L. involuta (D. Don) Presl, Tent. Pterid. 215. 1836 ; Bedd. Ferns South. India 17. t. 50. 1863 ; Handb. Ferns Brit. India 393. t. 228. 1883. *Grammitis lanceolata* D. Don, Fl. Nepal 14. 1825. *Gymnogramme involuta* Hook. Sp. Fil. 5 : 155. 1864 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 570. 1880.

Distrib. : Throughout Indian region up to 3000 m., Nepal, Bhutan, Sri Lanka, Japan, China, Malesian Islands, Polynesia.

L. parallela Copel. in Perkins, Fragm. Fl. Philip. 3 : 182. 1905 ; Sledge, Bull. Brit. Mus. (N. H.) Bot. 2 : 141. 1960. *L. lanceolata* sensu Bedd. Handb. Ferns Brit. India 392. 1883 (*non* C. Presl)

Distrib. : India (Eastern India & Eastern Himalayas), Sri Lanka, China, Japan, Fiji, Samoa, Africa.

L. remote-frondigera (Hayata) C. Chr. Ind. Fil. Suppl. II. 27. 1917 ; Dixit, Hazra & Kar, Bull. Bot. Surv. India 18 (1 - 4) : 222. f. 1 - 3. 1976. *Polypodium remote-frondigerum* Hayata, Icon. Formos. 5 : 323. f. 135 A, B. 1915.

Distrib. : India (Eastern Himalayas), China, Japan, Formosa.

L. salicifolia (Makino) Makino in Bot. Mag. Tokyo 19 : 138. 1905. *Gymnogramme salicifolia* Makino, Phan. Pter. Jap. Icon. III. 1. t. 34. 1899.

Distrib. : India (Himalayas and Assam), China, Japan, Formosa to Ryukyu.

POLYPODIACEAE Bercht. & Presl

Arthromeris (Moore) J. Sm. Hist. Fil. 110. 1875 ; Ching, Contr. Inst. Bot. Natn. Acad. Peiping 2 : 31. 1933.

Type : **A. wallichiana** (Spr.) Ching, Contr. Inst. Bot. Natn. Acad. Peiping 2 : 92. 1933.

A. himalayensis (Hook.) Ching, Contr. Inst. Bot. Natn. Acad. Peiping 2 : 99. 1933. *Polypodium himalayanse* Hook. Sp. Fil. 5 : 91. 1863. *Pleopeltis himalayensis* (Hook.) Bedd. Ferns Brit. India t. 318. 1869 ; Handb. Ferns Brit. India 370. f. 212. 1883. *Polypodium venustum* Wall. ex Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 566. 1880. (*non* Desv. 1811.) *Polypodium venustum* var. *niphobolooides* Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 587. 1880.

Distrib. : India (Himalayas), Burma, China.

A. himalayensis (Hook.) Ching var. **furcans** Ching, Icon. Fil. Sin. 3 : 100. 1935.

Distrib. : India (Sikkim).

A. jarrettii Sastry & Chaudhury, Bull. Bot. Surv. India 11 : 442. 1969.
Distrib. : India (Arunachal Pradesh).

A. lehmanni (Mett.) Ching, Contr. Inst. Bot. Natn. Acad. Peiping 2 : 96. 1933. *Polypodium lehmanni* Mett. Abh. Senekneb. Naturf. Ges. 2 : 109. 1857. *Pleopeltis lehmanni* (Mett.) Bedd. Ferns Brit. India t. 260. 1868 ; Handb. Ferns Brit. India 370. t. 211. 1883.

Distrib. : Himalayas to China, Formosa, Burma, North Thailand.

A. lungtauensis (Ching) Ching, Contr. Inst. Bot. Natn. Acad. Peiping 2 : 98. 1933. *Polypodium lungtauensis* Ching in C. Chr. Ind. Fil. Suppl. 3 : 152. 1934.

Distrib. : India (Eastern Himalayas), Tonkin, Laos, S. China.

A. tatsiensis (Fr. & Bureau ex Christ) Ching, Contr. Inst. Bot. Natn. Acad. Peiping 2 : 93. 1933. *Polypodium tatsiense* Fr. & Bureau ex Christ, Bull. Soc. Fr. 52. Mém. 1 : 19. 1905.

Distrib. : India (Himalayas), E. Nepal, N. Thailand, China.

A. tenuicauda (Hook.) Ching, Contr. Inst. Bot. Natn. Acad. Peiping 2 : 91. 1933. *Pleopeltis juglandifolia* var. *tenuicauda* Bedd. Handb. Ferns Brit. India 370. 1883.

Distrib. : India (Himalayas from Garhwal to Bhutan, Khasia).

A. wallichiana (Spreng.) Ching, Contr. Inst. Bot. Natn. Acad. Peiping 2 : 92. 1933. *Polypodium wallichianum* Spreng. Syst. Vcg. 4 : 53. 1827. (Based on *P. juglandifolium* D. Don). *Pleopeltis juglandifolia* (D. Don) Moore, Ind. Fil. 78. 1857 ; Bedd. Handb. Ferns Brit. India 368. t. 210. 1883. *Pleopeltis capitellata* (Mett.) Bedd. Ferns Brit. India t. 12. 1865.

Distrib. : India (Himalayas), Burma, China, Thailand.

A. wardii (Clarke) Ching, Contr. Inst. Bot. Natn. Acad. Peiping 2 : 94. 1933. *Polypodium wardii* Clarke, Journ. Linn. Soc. 25 : 99. t. 43. 1889.

Distrib. : India (Naga hills), Bhutan, Burma, Tibet.

Belvisia Mirb. in Lam. & Mirb. Hist. Nat. Veg. 3 : 473. 1802 ; Sledge, Bull. Brit. Mus. (N. H.) Bot. 2 : 139. 1960.

Lectotype : **B. spicata** (L. f.) Mirb. in Lam. & Mirb. Hist. Nat. Veg. 3 : 473. 1802.

B. henryi (Hieron. ex C. Chr.) Tagawa in Hara's Fl. East. Himalaya 490. 1960. *Hymenolepis henryi* Hieron. ex C. Chr. in Dansk Bot. Ark. 6 (3) : 67. 1929. *Gymnopteris spicata* sensu Bedd. Handb. Ferns Brit. India 432. 1883. (*pro parte*).

Distrib. : India (Himalayas), Burma, Thailand, W. China.

B. mucronata (Fée) Copel. Gen. Fil. 192. 1947. *Hymenolepis mucronata* Fée, Gen. Fil. 82. t. 6 B. f. 1. 1851-52. *Gymnopteris spicata* Bedd. Ferns Brit. India 432. 1883 (*pro parte, non Presl*).

Distrib. : India (Himalayas) Sri Lanka.

B. revoluta (Bl.) Copel. Gen. Fil. 192. 1947. *Hymenolepis revoluta* Bl. Enum. Pl. Jav. 2 : 201. 1828. *Gymnopteris spicata sensu* Bedd. Handb. Ferns Brit. India 432. 1883 (*pro parte ; non Presl*.)

Distrib. : India (Himalayas), Sri Lanka.

Christopteris Copel. Perkins, Fragm. Fl. Philip. 3 : 188. 1905.

Type : **C. sagitta** (Christ) Copel. in Perkins Frag. Fl. Philip. 3 : 188. 1905.

C. tricuspidis (Hook.) Christ, Journ. de Bot. 21 : 239, 272. 1908. *Acrostichum tricuspe* Hook. Sp. Fil. 272. t. 304. 1864. *Gymnopteris tricuspidis* Bedd. Ferns Brit. India t. 53. 1866 ; Handb. Ferns Brit. India 434. t. 263. 1883.

Distrib. : India (Sikkim).

Colysis Presl, Epim. Bot. 146. 1851 ; Ching, Bull. Fan Mem. Inst. Biol. Bot. 4. 293. 1933 ; Dixit & Ghosh, Bull. Bot. Surv. India 20 : 100. 1978.

Lectotype : **C. hemionitidea** (Wall. ex Mett.) Presl, Epim. Bot. 147. 1851.

C. digitata (Bak.) Ching, Bull. Fan Mem. Inst. Biol. 4 : 328. 1933. *Gymnogramma digitata* Bak. Journ. Bot. 267. 1890.

Distrib. : India (Sikkim, Manipur), China.

C. elliptica (Thunb.) Ching, Bull. Fan Mem. Inst. Biol. 4 : 333. 1933. *Polypodium ellipticum* Thunb. Fl. Jap. 335. 1784. *Selliguea elliptica* Bedd. Ferns Brit. India t. 150. 1867 ; Handb. Ferns Brit. India 392. t. 227. 1883 (*pro parte*). *Gymnogramma elliptica* Hook. & Bak. Syn. Fil. 389. 1874 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1. 570. 1880 (*pro parte*).

Distrib. : India (Himalayas), Nepal to Bhutan, Burma, Malaya peninsula, Philippines, Queensland, Formosa.

C. hemionitidea (Wall. ex Mett.) Presl, Epim. Bot. 147. 1851. *Poly podium hemionitideum* Wall. ex Mett. Ann. Lugd. Bat. 4 (2) : 215. 1866. *Pleopeltis hemionitidea* Moore, Ind. Fil. 436. 1862 ; Bedd. Ferns Brit. India t. 182. 1867 ; Handb. Ferns Brit. India 358. t. 202. 1883.

Distrib. : Throughout India in mountainous regions, Nepal, Bhutan, Bangladesh, Malay Islands.

C. latiloba (Ching) Ching, Bull. Fan Mem. Inst. Biol. 4 : 330. 1933.
Polypodium latilobum Ching, Bull. Fan Mem. Inst. Biol. 2 : 21. t. 7. 1931.

Distrib. : India (Himalayas).

C. longisora (Bak.) Ching, Bull. Fan Mem. Inst. Biol. 4 : 331. 1933.
Gymnogramme longisora Bak. Journ. Bot. 267. 1890.

Distrib. : India (Manipur), China, Vietnam.

C. pedunculata (Hook. & Grev.) Ching, Bull. Fan Mem. Inst. Biol. 4 : 321. 1933. *Ceterach pedunculata* Hook. & Grev. Icon. Fil. t. 5. 1827. *Gymnogramma hamiltoniana* Hook. Sp. Fil. 5 : 160. 1864 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 570. 1880. *Selliguea hamiltoniana* Bedd. Ferns Brit. India t. 239. 1867 ; Handb. Ferns Brit. India 390. t. 226. 1883.

Distrib. : India (Himalayas) Bangladesh, Nepal, Burma.

C. pentaphylla (Bak.) Ching, Bull. Fan Mem. Inst. Biol. 4 : 332. 1932. *Gymnogramma pentaphylla* Bak. Kew Bull. 233. 1898.

Distrib. : India (Manipur), China (Yunnan).

C. pothifolia (D. Don) Presl, Epim. Bot. 148. 1851. *Hemionitis pothifolia* D. Don, Prod. Fl. Nepal 13. 1825. *Colysis elliptica* var. *pothifolia* (D. Don) Ching, Bull. Fan Mem. Inst. Biol. 4 : 334. 1933. *Selliguea elliptica* Bedd. Ferns Brit. India t. 150. 1867 ; Handb. Ferns Brit. India 392. 1883 (*pro parte*). *Gymnogramma elliptica* Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 570. 1880 (*pro parte*).

Distrib. : India (Himalayas), Nepal, Burma, China, Malay Peninsula, Philippines, Queensland and Formosa.

Crypsinus Presl, Epim. Bot. 123. 1851.

Type : **C. pyrolifolius** (Goldm.) Copel. Gen. Fil. 206. t. 8. f. 1. 1947.

C. yakushimensis (Mak.) Tagawa, Acta Phytotax. Geobot. 14 : 194. 1952. *Polopodium engleri* var. *yakushimense* Mak. Bot. Mag., Tokyo 23 : 248. 1909.

Distrib. : India (Himalayas), Japan.

Majority of its species has been transferred by Pic. Ser. *Webbia* 28 (2) : 445. 1973. into **Phymatopteris**, The genus requires thorough revisionary studies. Pteridologists still recognise it as good genus.

Dendroglossa Presl, Epim. Bot. 149. 1851.

Lectotype : **D. minor** (Fée) Copel. Gen. Fil. 199. 1947.

D. minutula (Fée) Copel. Gen. Fil. 199. 1947. *Leptochilus minutulus* Fée, Mèm. Fam. Foug. 8 : t. 31. f. 2. 1865. *Gymnopteris minus sensu* Bedd. Ferns Brit. India t. 116. 1865 ; Handb. Ferns Brit. India 430. f. 259. 1883.

Distrib. : Eastern India, Sri Lanka.

Drymoglossum Presl, Tent. Pterid. 227. t. 10. f. 5 6. 1836. *nom. cons.*

Type : **D. piloselloides** (L.) Presl, Tent. Pterid. 227. t. 10. f. 5 6 1836.

D. heterophyllum (L.) Trimen, Journ. Linn. Soc. Bot. 24 : 152. 1887. *Actrostichum heterophyllum* L. Sp. Pl. 2 : 1067. 1753. *Drymoglossum piloselloides* var. *beddomei* Bedd. Handb. Ferns Brit. India 413. 1883.

Distrib. : South India, Tropical Asia.

D. piloselloides Presl, Tent. Pterid. 227. 1836 ; Bedd. Ferns South. India t. 55. 1863 ; t. 186. 1865 ; Handb. Ferns Brit. India 411. 1883.

Distrb. : Throughout India in mountainous regions, Sri Lanka, Java, Philippines, China, Japan.

Drymotaenium Mak. Bot. Mag. Tokyo 15 : 102. 1901.

Type : **D. miyoshianum** (Mak.) Mak. Bot. Mag. Tokyo 15 : 102. 1901.

D. miyoshianum (Mak.) Mak. Bot. Mag. Tokyo 15 : 109. 1901 ; Dixit & Nair, Journ. Indian Bot. Soc. 54 : 256. f. 1 - 6. 1975. *Taenitis miyoshiana* Mak. Bot. Mag. Tokyo 12 : 26. 1898.

Distrib. : India (Arunachal Pradesh), China.

Goniophlebium Presl, Tent. Pterid. 185. t. 7. f. 13 14. 1836.

Lectotype : **G. articulatum** (Desv.) Presl, Tent. Pterid. 185. t. 7. f. 13 - 14. 1836.

G. amoenum (Wall. ex Mett.) J. Sm. in Hook. Gen. Fil. t. 51. 1840 ; Bedd. Ferns Brit. India t. 5. 1865 ; Handb. Ferns Brit. India 317. 1883. *Polypodium amoenum* Wall. ex Mett. Abh. Senckneb. naturf. Ges. 2 : 80. 1857.

Distrib. : India (Himalayas from Gurhwal to Bhutan, Meghalaya) China to Formosa, south to Indo China and North Thailand.

G. subauriculatum (Bl.) Presl, Tent. Pterid. 186. 1836 ; Bedd. Ferns

Brit. India t. 78. 1865 ; Handb. Ferns Brit. India 323. f. 173. 1883.
Polypodium subauriculatum Bl. Enum. Pl. Jav. 133. 1828.

Distrib. : India (Meghalaya) Malaya Peninsula, Malay Islands, Queensland.

Holcosorus Moore, Ind. Fil. XXIX 1857 ; Dixit & Nair, Proc. Indian Acad. Sci. (Plant Sci) 86 B (6) : 385. f. 1 5. 1977.

Type : *H. bisulcatus* (Hook.) Ching, Sunyat. 5 : 265. 1940.

H. bisulcatus (Hook.) Ching, Sunyat. 5 : 265. 1940. *Grammitis bisulcatus* Hook. Icon. Pl. t. 998. 1854.

Distrib. : India (Arunachal Pradesh), Borneo.

Lemaphyllum Presl, Epim. Bot. 157. 1851 ; Nayar. Bull. Natn. Bot Gard. Lucknow no. 106. 1. 1964.

Lectotype : *L. spatulatum* Presl, Epim. Bot. 158. 1851.

L. carnosum (J. Sm.) Presl, Epim. Bot. 158. 1840. *Drymoglossum carnosum* J. Sm. in Hook. Gen. Fil. t. 784 1841 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 575. 1880 ; Bedd. Handb. Ferns Brit. India 411. t. 243. 1883.

Distrib. : India (Sikkim, Meghalaya), Nepal, Tonkin, N. Thailand, China.

Lepidogrammitis Ching, Sunyat. 5 : 258. 1940.

Type : *L. drymoglossoides* (Bak.) Ching, Sunyat. 5 : 258. 1940.

L. subrostrata (C. Chr.) Ching, Sunyat. 5 : 258. 1940. *Polypodium subrostratum* C. Chr. Ind. Fil. 567. 1906. *Pleopeltis rostrata* Bedd. Ferns Brit. India t. 159. 1867. (based on *Polypodium rostratum* Hook. Icon. Pl. t. 953. 1854).

Distrib. : N. E. India to S. W. China, Indo China.

Lepiosrus (J. Sm.) Ching, Bull. Fan Mem. Inst. Biol. Bot. 4 : 47. 1933 ; Birt & Tirkha, Amer. Fern Journ. 64 (2) : 49. 1974.

Type : *L. thunbergianus* (Kaulf.) Ching, Bull. Fan Mem. Inst. Biol. Bot. 4 : 88. 1933..

L. amaurolepidus (Sledge) Bir & Tirkha, in Bir & Vasudeva, Journ. Bomb. Nat. Hist. Soc. 68 : 192. 1971. var. *amaurolepidus*. *Pleopeltis amaurolepida* Sledge, Bull. Brit. Mus. (N. H.) 2 : 136. 1960.

Distrib. : South India, Sri Lanka.

L. amaurolepidus var. **longifolius** Bir & Tirkha, Amer. Fern Journ. 64 : 63. f. 53-57. 1974.

Distrib. : India (W. Himalayas).

L. angustus Ching, Bull. Fan Mem. Inst. Biol. Bot. 4 : 86. 1933.

Distrib. : India (Arunachal Pradesh), China.

L. bicolor (Takeda) Ching, Bull. Fan Mem. Inst. Biol. 4 : 66. 1933.

Polypodium excavatum var. *bicolor* Takeda, Notes Roy. Bot. Gard. Edinb. 8 : 280. 1915.

Distrib. : India (Himalayas), China.

L. clathratus (Clarke) Ching, Bull. Fan Mem. Inst. Biol. 4 : 71. 1933. *Polypodium clathratum* Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 559. t. 82. f. 1. 1880. *Pleopeltis clathrata* (Clarke) Bedd. Handb. Ferns Brit. India 348. 1883.

Distrib. : India (Himalayas), Afghanistan.

L. contortus (Christ) Ching, Bull. Fan Mem. Inst. Biol. Bot. 4 : 90. 1933. *Polypodium lineare* var. *contortum* Christ, Nuovo. Giorn. Bot. Soc. Ital. n. s. 4 : 98. 1897.

Distrib. : India (Himalayas), China.

L. excavatus (Willd.) Ching, Bull. Fan Mem. Inst. Biol. 4 : 68. 1933, var. *excavatus*. *Polypodium excavatum* Willd. in L. Sp. Pl. 5 : 158. 1810. *Pleopeltis simplex* Bedd. Handb. Ferns Brit. India 347. 1883.

Distrib. : India (Himalayas).

L. excavatus (Willd.) Ching var. *himalayensis* Bir & Tirkha, Amer. Fern Journ. 64 : 58. f. 33-37. 1974.

Disirib. : India (N. W. Himalayas).

L. excavatus (Willd.) Ching var. *mortonianus* Bir & Tirkha, Amer. Fern Journ. 64 : 56. f. 25-32. 1974.

Distrib. : India (N. W. Himalayas).

L. excavatus var. *scolopendrius* (Buch. Ham. ex D. Don) Ching, Bull. Fan Mem. Inst. Biol. Bot. 4 : 69. 1933. *Polypodium scolopendrium* Buch. Ham. ex D. Don Prod. Fl. Nepal. 1. 1825.

Distrib. : India (Himalayas), Nepal, Tibet, China.

L. kashyapii (Mehra) Mehra in Bir, Res. Bull. (N. S.) Punjab Univ. 13 : 24. 1962. *Polypodium kashyapii* Mehra in Ferns of Mussorie, Punjab. Univ. Publ. 24. 1939.

Distrib. : India (Himalayas).

L. kuchenensis (Wu) Ching, Bull. Fan Mem. Inst. Biol. 4 : 69. 1933.
Polypodium kuchenensis Wu in Wu et al, Polypod. Yaoshan. 276. t. 129. 1932.

Distrib. : India (Eastern Himalayas), China.

L. leiopteris (Kunze) Bir & Tirkha, Amer. Fern Journ. 64 : 54. f. 21-26. 1974. *Polypodium leiopteris* Kunze, Linnaea 23 : 279, 319. 1859. *Pleopeltis scolopendrium* (D. Don) Alston & Bonner, Candollea 15 : 207. 1956.

Distrib. : India (Himalayas).

L. longifolius (Bl.) Holtt. Fl. Malaya II. 151. 1954. *Grammitis longifolia* Bl. Enum. Pl. Jav. 119. 1828. *Pleopeltis longifolia* (Bl.) Bedd. Handb. Ferns Brit. India 349. t. 196. 1883.

Distrib. : North - East India, Burma, Malay Peninsula, Singapore, Malay Islands, Philippines.

L. loriformis (Wall. ex Mett.) Ching, Bull. Fan, Mem Inst. Biol. Bot. 4 : 81. 1933. *Polypodium loriforme* Wall. ex. Mett. Poly. 92. n. 164. 1857. *Pleopeltis linearis* var. *steniste* (Clarke) Bedd. Handb. Ferns Brit. India 347. 1883.

Distrib. : India (Himalayas), China.

L. macrocarpus (Willd.) Ching, Bull. Fan Mem Inst. Biol. Bot. 4 : 80. 1933. *Polypodium macrocarpum* Willd. in Linn. Sp. Pl. 5 : 147. 1810. *Pleopeltis lepidota* Bedd. Ferns South. India 60. t. 181. 1865. (*nom. illegit*).

Distrib. : South India, Sri Lanka.

L. macrosphaerus ((Bak). Ching, Bull. Fan Mem. Inst. Biol. Bot. 4 : 73. 1933. var. *macrosphaerus*. *Polypodium macrosphaerum* Bak. in chritst Bull. Herb. Boiss. II. 3 : 509. 1903. *Pleopeltis macrosphaera* (Bak.) Panigr. & Patn. Proc. Natn. Acad. Sci. India B. 34 : 481. 1964.

Distrib. : India (Eastern India & Eastern Himalayas), China.

L. macrosphaerus var. *asterolepis* (Bak.) Dixit, comb. nov. *Polypodium asterolepis* Bak. Journ. Bot. Brit. & For. 26 : 230. 1880. *Pleopeltis macrosphaera* var. *asterolepis* (Bak.) Panigr & Patn. Proc. Natn. Acad. Sci. India B. 34 : 482. f. 1-5 1964.

Distrib. : Eastern India, Burma, China.

L. nudus (Hook.) Ching, Bull. Fan Mem. Inst. Biol. Bot. 4 : 83. 1933.

Pleopeltis nuda Hook. Exot. Fl. 1 : t. 63. 1823. *P. wightiana* (Thw.) Bedd. Ferns South. India 60. t. 180. 1865. *P. linearis* sensu Bedd. Handb. Ferns Brit. India 346. 1883 (*pro parte*) ; non *Polypodium lineare* Thunb.

Distrib. : Throughout India in mountainous regions, Sri Lanka, Malaya Peninsula, Malay Islands, China, Japan, Central & South Africa & its Islands.

L. oligolepidus (Bak.) Ching, Bull. Fan Mem. Inst. Biol. Bot. 4 : 80. 1933. *Polypodium oligolepidum* Bak. Gard. Chron. n. s. 14 : 594. 1880.

Distrib. : India (N. W. Himalayas), China.

L. oosphaerus (C. Chr.) Ching, Bull. Fan Mem. Inst. Biol. Bot. 4 : 73. 1933. *Polpyodium oosphaerium* C. Chr. Contr. Nat. Herb. 26 : 334. t. 29. 1931.

Distrib. : India (N. W. Himalayas), China.

L. pseudonudus Ching, Bull. Fan Mem. Inst. Biol. Bot. 4 : 83. 1933.

Distrib. : E. India, Burma, China.

L. pteropus (Bl.) Ching, Bull. Fan Mem. Inst. Biol. Bot. 4 : 77. 1933.

Pleopeltis pteropus Bl. Enum. Pl. Jav. 168. t. 76. 1828 ; Bedd. Handb. Ferns Brit. India 359. t. 203. 1883. *Polypodium tridactylon* Wall. ex Hook. & Grev. Icon. Fil. t. 209. 1831 ; Bedd. Ferns Brit. India t. 11. 1865. *Kaulinia pteropus* (Bl.) Nayar, Taxon 13 : 67. 1968.

Distrib. : North East India, Bhutan, Bangladesh, Malaya Peninsula, Philippines, S. China.

L. pteropus var. **minor** (Bedd.) Dixit, comb. nov. *Pleopeltis pteropus* var. *minor* Bedd. Ferns South. India t. 179. 1865 ; Handb. Ferns Brit. India 361. t. 204. 1883. *Kaulinia pteropus* var. *minor* (Bedd.) Nayar & Kaur, Comp. to Bedd. Handb. 87. 1974.

Distrib. : South India, Sri Lanka.

L. sordidus (C. Chr.) Ching, Bull. Fan Mem. Inst. Biol. Bot. 4 : 79.

1933. *Poypodium sordidum* C. Chr. Contr. U. S. Nat. Herb. 26 : 320. 1931.

Distrib. : E. India, Burma, China.

L. subconfluens Ching, Bull. Fan Mem. Inst. Biol. Bot. 4 : 85. 1933.

Distrib. : Eastern India and Eastern Himalayas, China, Thailand.

L. sublinearis (Bak. ex Takeda) Ching, Bull. Fan Mem. Inst. Biol. Bot. 4 : 78. 1933. *Polypodium sublineare* Bak. ex Takeda in Notes. Royal Bot. Gard. Edinb. 8 : 276. 1915.

Distrib. : India (Himalayas), Burma, China, Thailand, Indo-China.

L. suboligolepidus Ching, Bull. Fan Mem. Inst. Biol. Bot. 4 : 77. 1933,

Distrib. : India (N. W. Himalayas), China.

L. thunbergianus (Kaulf.) Ching, Bull. Fan Mem. Inst. Biol. Bot. 4 : 88. 1933. *Pleopeltis thunbergiana* Kaulf. Wesen. Farrnkr. 113. 1827. *Polypodium lineare* Thunb. Fl. Jap. 335. 1784 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 558. 1881 (*pro parte*).

Distrib. : India (Himalayas), China, Korea, Indo China, Formosa, Philippines.

L. ussuriensis (Regel & Mack.) Ching, Bull. Fan Mem. Inst. Biol. 4 : 91. 1933 ; Bir & Tirkha, Bull. Bot. Surv. India 11 : 268. 1969. *Pleopeltis ussuriensis* Regel & Mack. Mem. Acad. Sci. Petersb. VII. 44 : 175. 1861.

Distrib. : India (Himalayas), China.

Leptochilus Kaulf. Enum. Fil. 147. 1824 ; Ching, Bull. Fan Mem. Inst. Biol. Bot. 4 : 340. 1933.

Type : **L. axillaris** (Cav.) Kaulf. Enum. Fil. 147. t. 1. f. 10. 1824.

L. axillaris (Cav.) Kaulf. Enum. Fil. 147. t. 1. f. 10. 1824 ; *Acrostichum axillare* Cav. Anal. Nat. 1 : 101. 1799. *Gymnopteris variabilis* var. *axillaris* (Cav.) Bedd. Ferns Brit. India t. 271. 1868 ; Handb. Ferns Brit. India 430. 1883.

Distrib. : South India, plains of Bengal and Assam, Bangladesh, Burma.

L. laciniatus (Hook.) Ching, Bull. Fan Mem. Inst. Biol. Bot. 4 : 344. 1933. *Acrostichum variable* Hook. var. *laciniatum* Hook. Sp. Fil. 5 : 255. 1864. *Gymnopteris feei* var. *pinnatifida* Bedd. Ferns South. India t. 273. 1864. *Gymnopteris feei* var. *triloba* Bedd. Ferns Brit. India t. 273. 1864.

Distrib. : India (Eastern Himalayas), China, Sri Lanka.

L. lanceolatus Fée, Acrost. 87. t. 47. 1845. *Gymnopteris feei* Moore, Ind. Fil. XXX. 1857 ; Bedd. Ferns South. India t. 48. 1864. (f. typica). *Gymnopteris variabilis* var. *lanceolata* Bedd. Handb. Ferns. Brit. India 429. 1883. *G. lanceolata* Bedd. Ferns Brit. India. Suppl. 26. 1876.

Distrib. : South India, Sri Lanka, Burma

Marginaria Bory, Dict. Class d' Hist. nat. 6 : 587. 1824 ; 10 : 176. 1826.

Lectotype : **M. polypodioides** (L.) Tidstr. Torreya. 5 : 171. f. 1. 1905.

M. macrocarpa (Bory ex Willd.) Nayar & Kaur, Comp. to Bedd. Handb. Ferns Brit. India 86. 1974. *Polypodium macrocarpum* Bory ex Willd. in Linn. Sp. Pl. 5 : 147. 1810.

Distrb. : Eastern India, China.

Metapolypodium Ching, Acta Phytotax. Sin. 16 (4) : 28. 1976.

Type : **M. manmeiense** (Christ) Ching, Acta Phytotax. Sin. 16 (4) : 29. 1978.

M. manmeiense (Christ) Ching, Acta Phytotax. Sin. 16 (4) : 29. 1978.
Polypodium manmeiense Christ, Bull. Herb. Boiss. 6 : 870. 1898.

Disrib. : India (Sikkim), China.

Microsorium Link, Hort. Reg. Bot. Berol. 2 : 110. 1833 ; Ching, Bull. Fan Mem. Inst. Biol. Bot. 4 : 293. 1933 ; Nayar, Bull. Nation. Bot. Gard. Lucknow no. 58. 11. 1961.

Type : **M. punctatum** (L.) Copel. Univ. Calif. Publ. Bot. 16 : 111. 1929.

M. alternifolium (Willd.) Copel, Gen. Fil. 197. 1947. *Polypodium alternifolium* Willd. in L. Sp. Pl. 5 : 168. 1810. *Pleopeltis nigrescens* Bl. Enum. Fil. 127. 1828 ; Bedd. Handb. Ferns Brit. India 367. t. 208. 1883. *P. longissima* Bedd. Ferns. South. India t. 176. 1865.

Distrib. : India (Western ghats, plains of Assam), Sri Lanka, Cultivated with choice in the gardens.

M. cuspidatum (D. Don) Tagawa in Hara's Fl. East. Himalaya 495. 1966. *Polypodium cuspidatum* D. Don, Prod. Fl. Nepal. 2. 1825. *P. lucidum* Roxb. ex Griff. Calc. Journ. Nat. Sci. 4 : 486. 1844. *P. leiorrhizum* Wall. ex Mett. Fil. Lips. 37. t. 25. f. 7. 1856 ; Bedd. Ferns South. India t. 174. 1865. *Pleopeltis leiorrhiza* (Wall. ex Mett.) Moore, Ind. Fil. 346. 1862 ; Bedd. Handb. Ferns Brit. India 372. 1883.

Distrib. : North India to S.W. China, Tonkin, Laos.

M. dilatatum (Bedd.) Sledge, Bull. Brit. Mus. (N.H.) Bot. 2 : 143.

1960. *Pleopeltis dilatata* Bedd. Ferns Brit. India t. 122. 1866 ; Handb. Ferns Brit. India 367. 1883.

Distrib. : North India to S.W. China, Sri Lanka, Malaya Peninsula, Samoa.

M. fortunei (Moore) Ching, Bull. Fan Mem. Inst. Biol. Bot. 4 : 304
1933. *Drynaria fortunei* Moore, Gard. Chron. 1855 : 708. 1855.

Distrib. : India (Eastern Himalayas), Bhutan, Burma, Tonkin, China, Formosa.

M. heterocarpum (Bl.) Ching, Bull. Fan Mem. Inst. Biol. Bot. 4 : 295. 1933. *Grammitis heterocarpa* Bl. Enum. Pl. Jav. 2 : 118. 1828 ; 167. 1829.

Distrib. : South India, Java, Celebes.

M. hymenodes (Kunze) Ching, Bull. Fan Mem. Inst. Biol. 4 : 301. 1933. *Polypodium hymenodes* Kunze in Linnaea 23 : 279, 319. 1850. *Polypodium superficiale* Bl. Enum. Pl. Jav. 123. 1828 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 557. 1880 (excl. var.). *Polypodium superficiale* var. *semilinearis* Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 558. 1880. *Pleopeltis superficialis* Bedd. Handb. Ferns Brit. India 350. 1883.

Distrib. : India (Himalayas), Japan, Korea, China, Malaya Islands, Queensland.

M. insigne (Bl.) Ching, Bull. Fan Mem. Inst. Biol. Bot. 4 : 311. 1933. *Polypodium inisgne* Bl. Enum. Pl. Jav. 137. 1828. *Pleopeltis inisgnis* (Bl.) Bedd. Handb. Ferns Brit. India 365. t. 207. 1883.

Dsitrib. : India (Great Nicobar) Malaya, Malay Islands.

M. linguaeforme (Mett.) Copel. Univ. Calif. Publ. Bot. 16 : 116. 1929 ; Nayar & Madhusoodan, Journ. Linn. Soc. 75 : 283. 1977. *Polypodium linguaeforme* Mett. Ann. Lugd. Bat. 2 : 228. 1866.

Distrib. : South India, Malesian Islands.

M. membranaceum (D. Don) Ching, Bull. Fan Mem. Inst. Biol. 4 : 309. 1933. *Polypodium membranaceum* D. Don, Prod. Fl. Nepal 2. 1925 ; Clarke Trans. Linn. Soc. Lond. II. Bot. 1 : 580. 1880. *Pleopeltis grandifolia* Bedd. Ferns South. India t. 177. 1865. *P. membraancea* Moore, Ind. Fil. 191. 1859 ; Bedd. Handb. Ferns Brit. India 335. 1883. *Polypodium heterocarpum* (Bl.) Mett. Fil. Lips. 37. t. 25. f. 24 - 25. 1856 ; Bedd. Ferns South. India t. 177. 1865.

Distrib. : Throughout India in mountainous regions, Sri Lanka, Nepal, China.

M. phyllomanes (Christ) Koids. Acta Phytotax. Geobot. 5 : 50. 1936. *Polypodium phyllomanes* Christ, Bull. Acad. Geogr. Bot. Mans. 210. 1902. *Pleopeltis ovata* Bedd. Handb. Ferns Brit. India 354. t. 200. 1883.

Distrib. : India (Meghalaya, Sikkim), Nepal, Bhutan.

***M. pteropus** (Bl.) Copel. Univ. Calif. Publ. Bot. 16. 112. 1929. *Polypodium pteropus* Bl. Enum. Pl. Jav. ad. 3. 1828 ; Fl. Jav. Fil. 168. t. 76. 1829 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 562. 1880 ; Bedd. Handb. Ferns Brit. India 360. 1883. *Pleopeltis pteropus* Moore, Ind. Fil. LXXVIII 1867 ; Bedd. Handb. Ferns. Brit. India 359. t. 203. 1883. *Polypodium tridactylon* Wall. ex Hook. & Grev. Icon. Fil. 6. 209. 1831 ; Bedd. Ferns Brit. India t. 11. 1865.

Distrib. : India (North India, Sikkim, Meghalaya) Nepal, Bhutan, Bangladesh, Malay Peninsula. Malesia, Formosa.

***M. pteropus** var. *minor* (Bedd.) Balakr. Bull. Bot. Surv. India 22 : 137. 1980. *Pleopeltis pteropus* var. *minor* Bedd. Handb. Ferns. Brit. India 361. 1833.

Distrib. : India (South India), Andamans, Sri Lanka.

M. punctatum (L.) Copel. Orient. Gen. Polyp. in Univ. Calif. Publ. Bot. 16 : 110. 1929. *Acrostichum punctatum* L. Sp. Pl. ed. II. 2 : 1524. 1763. *Pleopeltis punctata* Bedd. Ferns Brit. India, Suppl. 22. 1876 ; Handb. Ferns Brit. India 357. 1883. *Pleopeltis irioides* Moore, Ind. Fil. LXXVII. 1857 ; Bedd. Ferns South. India t. 178. 1864.

Distrib. : South India, warmer parts of Asia, Africa to Polynesia, Formosa to S. China.

M. rubidum (Kunze) Copel. Gen. Fil. 197. 1947. *Polypodium rubidum* Kunze, Bot. Zeit. 117. 1848. *Pleopeltis longissima* Bl. Enum. Fil. 127. 1828 ; Bedd. Ferns Brit. India, Suppl. 388. 1876 ; Handb. Ferns Brit. India 366. 1883.

Distrib. : India (Plains of Assam), Bangladesh, Philippines, Formosa, Malaya Islands.

M. scolopendrium (Burm.) Copel. Gen. Fil. 195. 1947. *Polypodium scolopendria* Burm. Fl. Ind. 232. 1968. *Pleopeltis phymatodes* Moore, Ind.

*Some workers still prefer to recognise these taxa under *Microsorium*. Thorough revisionary studies of the genera *Lepisorus*, *Microsorium* and *Pyrrosia* are required urgently on Indian basis for the better understanding.

Fil. LXXVIII, 1857 ; Bedd. Ferns South. India t. 173. 1863 ; Handb. Ferns Brit. India 366. 1883.

Distrib. : Throughout the tropics of the old world, usually cultivated in the gardens.

M. superficiale (Bl.) Ching, Bull. Fan. Mem. Inst. Biol. 4 : 299. 1933. *Polypodium superficiale* Bl. Jav. Fil. 130. t. 56. f. 1. 1828. ; Bedd. Ferns Brit. India t. 75. 1865. *Pleopeltis superficiale* (Bl.) Bedd. Handb. Ferns Brit. India 350. 1883.

Distrib. : India (Eastern and Western India) Burma, Nepal, Bhutan, China, Japan, Corea, Singapore, Java, Queensland.

M. zippelii (Bl.) Ching, Bull. Fan Mem Inst. Biol. 4 : 308. 1933. *Polypodium zippelii* Bl. Enum. Pl. Jav. 172. t. 80. 1828 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 561. 1880. *Pleopeltis heterocarpa* Bedd. Ferns Brit. India t. 319. 1869. *P. zippelii* Moore, Ind. Fil. CCCVIII. 1862 ; Bedd. Handb. Ferns Brit. India 357. 1883.

Distrib. : India (Himalayas), Bhutan, Nepal, Java, Philippines, Malesia, China, Indo China.

Neocheiropteris Christ, Bull. Soc. Bot. France 52 mem 1 : 21. 1905.

Type : **N. palmatodentata** (Bak.) Christ, Bull. Soc. Bot. France 52. mem. 1 : 21. 1905.

N. normalis (D. Don) Tagawa, Journ. Jap. Bot. 27 : 217. 1952. *Polypodium normale* D. Don, Prod. Fl. Nepal. 1. 1825. *Pleopeltis normalis* (D. Don) Moore, Ind. Fil. CCCXLVII. 1862 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 558. 1880 ; Bedd. Ferns Brit. India t. 10. 1865 ; Handb. Ferns Brit. India 353. 1883.

Distrib. : India (Himalayas), Nepal, Bhutan, Burma, China.

N. ovatus (Wall. ex Hook. & Grev.) Nayar & Kaur, Comp. to Bedd. Handb. Ferns Brit. India 86. 1974. *Polypodium ovatum* Wall. ex Hook & Grev. Icon. Pl. 41. 1827 ; Bedd. Ferns Brit. India t. 157. 1865. *Pleopeltis ovata* (Wall. ex Hook. & Grev.) Bedd. Handb. Ferns Brit. India 354. 1883.

Distrib. : India (Himalayas), Nepal, Bhutan.

Paraleptochilus Copel. Gen. Fil. 198. 1947.

Type : **P. decurrens** (Bl.) Copel. Gen. Fil. 198. 1947.

P. decurrens (Bl.) Copel. Gen. Fil. 198. 1947. *Leptochilus decurrens* Bl. Enum. Pl. Jav. 2 : 206. 1828. *Gymnopteris variabilis* (Hook.) Bedd.

Ferns Brit. India t. 272. 1868 ; Handb. Ferns Brit. India 429. t. 358. 1883.
Acrostichum variable Hook. Sp. Fil. 5 : 277. 1864 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 578. 1880.

Distrib. : India (Sikkim, Eastern India, Cachar, S. India), Burma, Polynesia, Malesia, Indo China, S.W. China.

Phlebodium (R.Br.) J. Sm. in Hook. Journ. Bot. 4 : 58. 1841 ; Bhargavan & Nair, Journ. Bomb. Nat. Hist. Soc. 77 (3) : 540. t. 1. 1980.

Type : **P. aureum** (L.) J. Sm. Lond. Journ. Bot. 4 : 59. 1841.

P. aureum (L.) J. Sm. Lond. Journ. Bot. 4 : 59. 1841. *Polypodium aureum* L. Sp. Pl. 2 : 1087. 1753.

Distrib. : India (Tamil Nadu), Trop. America.

Phymatopteris Pic. Ser. Webbia 28 : 460. 1973.

Type : **P. palmata** (Bl.) Pic. Ser. Webbia 28 : 460. 1973.

P. crenato-pinnata (Clarke) Pic. Ser. Webbia 28 : 461. 1973. *Polypodium crenato - pinnatum* Clarke Journ. Linn. Soc. Lond. 25 : 99. t. 42. 1889. *Pleopeltis crenato - pinnata* (Clarke) Bedd. Handb. Ferns. Brit. India Suppl. 96. 1892.

Distrib. : India (Manipur) (China. Yunnan).

P. chrysotricha (C. Chr.) Pic. Ser. Webbia 28 : 461. 1973. *Polypodium chrysotrichum* C. Chr. Contr. U.S. Nation. Herb. 26 : 320. t. 23. 1931. *Crypsinus chrysotrichus* (C. Chr.) Tagawa, Acta Phytotax. Geobot. 14 : 193. 1952 ; Balakr. et Chowdh. Sci. & Cult. 33 : 62. 1967.

Distrib. : India (Himalayas, Andaman & Nicobar Islands), China, Japan.

P. ebenipes (Hook.) Pic. Ser. Webbia 28 : 462. 1973. *Polypodium ebenipes* Hook. Sp. Fil. 5 : 88. 1864 ; Clarke. Trans. Linn. Soc. Lond. II. Bot. 1 : 564. 1880. *Pleopeltis ebenipes* (Hook.) Bedd. Ferns Brit. India t. 138. 1867 ; Handb. Ferns. Brit. India 363. 1883. *Phymatodes ebenipes* (Hook.) Ching, Contr. Inst. Bot. Nat. Acad. Peiping 2 : 86. 1933. *Crypsinus ebenipes* (Hook.) Copel. Gen. Fil. 206. 1947.

Distrib. : India (Himalayas from Gurhwal to Bhutan), China.

P. echinospora (Tagawa) Pic. Ser. Webbia 28 (2) : 462. 1973. *Phymatodes echinospora* Tagawa, Acta Phytotax. Geobot. 3 : 95. 1934. *Crypsinus echinosporus* (Tagawa) Tagawa, Acta Phytotax. Geobot. 3 : 95. 1934.

Distrib. : India (Himalayas), Japan.

P. engleri (Luerss.) Pic. Ser. Webbia 28 : 462. 1973. *Polypodium engleri* Luerss. in Engl. Bot. Jahrb. Syst. 4 : 361. 1883. *Crypsinus engleri* (Luerss.) Copel. Gen. Fil. 206. 1947.

Distrib. : India (Himalayas), Japan.

P. erythrocarpa (Mett. ex Kuhn) Pic. Ser. Webbia 23 : 462. 1973. *Polypodium erythrocarpum* Mett. ex Kuhn, Linnaea 36 : 135. 1869 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. I : 565. 1880. *Goniophlebium erythrocarpum* (Mett. ex Kuhn) Bedd. Ferns Brit. India Suppl. 382. 1867 ; Handb. Ferns Brit. India 319. t. 171. 1883. *Phymatodes erythrocarpa* (Mett. ex Kuhn) Ching, Contr. Inst. Bot. Nat. Acad. Peiping 2 : 80. 1933. *Crypsinus erythrocarpus* (Mett. ex Kuhn) Tagawa in Hara's Fl. East. Himal. 491. 1966.

Distrib. : India (Himalayas), China.

P. griffithiana (Hook.) Pic. Ser. Webbia 28 : 462. 1973. *Polypodium griffithianum* Hook. Icon. Pl. 10 : t. 951. 1854; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 560. 1880. *Pleopeltis griffithiana* (Hook.) Bedd. Ferns Brit. India t. 158. 1867 ; Handb. Ferns. Brit. India 354. t. 199. 1883. *Crypsinus griffithianus* (Hook.) Copel. Gen. Fil. 206. 1957.

Distrib. : India (Sikkim, Meghalaya), Bhutan.

P. hastata (Thunb.) Pic. Ser. Webbia 28 : 462. 1973 *Polypodium hastatum* Thunb. Fl. Jap. 335. 1784 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 562. 1880. *Pleopeltis malacodon* sensu Bedd. Ferns Brit. India Suppl. 23. t. 387. 1876.

Distrib. : Throughout India in mountainous regions, China.

P. malacodon (Hook.) Pic. Ser. Webbia 28 : 463. 1973. *Polypodium malacodon* Hook. Sp. Fil. 5 : 87. 1864 : Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 564. 1880. *Pleopeltis malacodon* (Hook.) Bedd. Handb. Ferns Brit. India 363. 1883. *Crypsinus malacodon* (Hook.) Copel. Gen. Fil. 206. 1947.

Distrib. : India (Himalayas), Nepal, Bhutan, W. China.

P. montana (Sledge) Pic. Ser. Webbia 28 : 463. 1973. *Crypsinus montanus* Sledge. Bull. Brit. Mus. (N.H.) Bot. 2 : 145 1960. *Pleopeltis oxyloba* Bedd. Ferns South. India 59. t. 175. 1865 (*pro parte*), non *Polypodium oxylobum* Wall. ex Kunze. *Pleopeltis hastata* Bedd. Handb. Ferns Brit. India 362. 1883 (*pro parte*) ; non *Polypodium hastatum* Thunb.

Distrib. : South India, Sri Lanka.

P. nigrovenia (Christ) Pic. Ser. Webbia 28 : 463. 1973. *Polypodium shen-siense* var. *nigrovenium* Christ, Bull. Acad. Geogr. Bot. 15 : 106.

1906. *Crypsinus nigrovenius* (Christ.) K. Iwat, in Hara's Fl. East. Himal. 3rd report, Univ. Tokyo Mus. Bull. no. 8. 197. 1975.

Distrib. : India (Himalayas), Nepal, S.W. China, Japan, Vietnam.

P. oxyloba (Wall. ex Kunze) Pic. Ser. Webbia 28 : 464. 1973. *Polypodium oxylobum* Wall. ex Kunze in Linnaea 24 : 255 1851. *Pleopeltis hastata* sensu Bedd. Handb. Ferns, Brit. India 362. 1883 (*pro parte*). *Polypodium hastatum* var. *oxyloba* Clakre, Trans. Linn. Soc. Lond. II. Bot. 1 : 583. 1880 (*pro parte*). *Crypsinus oxylobus* (Wall. ex Kunze) Sledge, Bull. Brit. Mus. (N.H.) Bot. 2 : 145. 1960.

Distrib. : India (Himalayas), S.W. China.

P. quasidivaricata (Hayata) Pic. Ser. Webbia 28 : 464. 1973. *Polypodium quasidivaricatum* Hayata, Journ. Coll. Sc. Univ. Tokyo 30 : 446. 1911. *Crypsinus quasidivari catus* (Hayata) Copel. Gen. Fil. 206. 1947.

Distrib. : India (Himalayas east to China), Formosa.

P. rhynchophylla (Hook.) Pic. Ser. Webbia 28 : 464. 1973. *Polypodium rhynchophyllum* Hook. Icon Pl. 10 : t. 954. 1854. *Pleopeltis rhynco-phylla* (Hook.) Bedd. Ferns Brit. India t. 9. 1865 ; Handb. Ferns Brit. India 353. t. 198. 1883. *Crypsinus rhyncophyllus* (Hook.) Copel. Gen. Fil. 206. 1947.

Distrib. : India (Meghalaya), Burma, Bhutan, North Thailand, Laos, Vietnam, China.

P. stewartii (Bedd.) Pic. Ser. Webbia 28 : 464. 1973. *Pleopeltis stewartii* Bedd. Ferns Brit. India t. 204. 1866. *Pleopeltis malacodon* var. *majus* Bedd. Handb. Ferns Brit. India 363. 1883. *Polypodium cyrtolobum* Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 563. t. 83. 1880. *Crypsinus Stewartii* (Bedd.) Copel. Gen. Fil. 206. 1947.

Distrib. : India (Himalayas), Nepal, Bhutan, W. China.

P. stracheyi (Ching) Pic. Ser. Webbia 28 : 464. 1973. *Phymatodes stracheyi* Ching, Contr. Inst. Bot. Nat. Acad. Peiping 2 : 83. 1933.

Distrib. : India (Himalayas), China.

Phymatosorus Pic. Ser. Webbia 28 : 457. 1973.

Type : **P. scolopendria** (N.L. Burm.) Pic. Ser. Webbia 28 : 457. 1973.

P. longissimus (Bl.) Pic. Ser. Webbia 28 : 459. 1973. *Polypodium longissimum* Bl. Enum. Pl. Jav. 2 : 127. 1828. *Pleopeltis longissima* (Bl.) Bedd. Ferns Brit. India Suppl. 388. 1876 ; Handb. Ferns Brit. India 366. 1883.

Distrib. : India (Eastern India), Bangladesh, Philippines, Formosa, Malaya Islands.

P. lucidus (Roxb. ex Griff.) Pic. Ser. *Webbia* 28 : 459. 1973.
Polypodium lucidum Roxb. ex Griff. *Calc. Journ. Nat. Hist.* 4 : 486. 1844.
Pleopeltis leiorhiza Moore, *Ind. Fil.* 346. 1862 ; Bedd. *Ferns South. India* t. 174. 1863 ; *Handb. Ferns Brit. India* 372. 1883. *Polypodium leiorhizum* Wall. ex Clarke, *Trans. Linn. Soc. Lond. II. Bot.* 1 : 367. 1880.

Distrib. : India (Himalayas, S. India, Western Ghats).

P. nigrescens (Bl.) Pic. Ser. *Webbia* 28 : 459. 1973. *Polypodium nigrescens* Bl. *Enum. Pl. Jav.* 2 : 126. 1828. *Pleopeltis nigrescens* (Bl.) Bedd. *Handb. Ferns Brit. India* 367. t. 208. 1883.

Distrib. : South India, Sri Lanka.

P. scolopendria (N. L. Burm.) Pic. Ser. *Webbia* 28 : 460. 1973.
Polypodium scolopendrium N. L. Burm. *Fl. Ind.* 232. 1768. *Pleopeltis phymatodes* (L.) Bedd. *Ferns South. India* t. 173. 1865 ; *Handb. Ferns Brit. India* 366. t. 208. 1883.

Distrib. : Throughout the tropics of the old world, Sri Lanka, Malay Peninsula.

Polypodiastrum Ching, *Acta Phytotax. Sin.* 16 (4) : 28. 1978.

Type : **P. argutum** (Wall. ex Hook.) Ching, *Acta Phytotax. Sin.* 16 (4) : 28. 1978.

P. argutum (Wall. ex Hook.) Ching, *Acta Phytotax. Sin.* 16 (4) : 28. 1978. *Polypodium argutum* Wall. ex Hook. *Sp. Fil.* 5 : 32. 1883 ; Clarke, *Trans. Linn. Soc. Lond. II. Bot.* 1 : 551. 1880. *Goniophlebium argutum* (Wall. ex Hook.) J. Sm. in Hook. *Gen. ed.* t. 51. 1840 ; Bedd. *Ferns Brit. India* t. 6. 1865 ; *Handb. Ferns Brit. India* 323. t. 174. 1883.

Distrib. : India (Himalayas from Kashmir to Bhutan), China.

Polypodioides Ching, *Acta Phytotax. Sin.* 16 (4) 26 : 1978.

Type : **P. amoena** Wall. ex Ching, *Acta Phytotax. Sin.* 16 (4) : 27. 1978.

P. atkinsonii (C. Chr.) Ching, *Acta Phytotax. Sin.* 16 (4) : 27. 1978.
Polypodium atkinsonii C. Chr. *Ind. Fil.* 511. 1905. *Goniophlebium hender-*
sonii Bedd. *Ferns Brit. India Suppl.* 21. t. 383. 1876 ; *Handb. Ferns Brit. India* 320. 1883.

Distrib. : India (Himalayas), Nepal.

P. lachnopus (Wall. ex Hook.) Ching, Acta Phytotax. Sin. 16 (4) : 27. 1978. *Polypodium lachnopus* Wall. ex Hook. Icon. Pl. t. 952. 1854. *Goniophlebium lachnopus* (Wall. ex Hook.) J. Sm. in Hook. Gen. ad. t. 51. 1840 ; Bedd. Ferns Brit. India t. 163. 1867 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 551. 1880 ; Bedd. Handb. Ferns Brit. India 319. 1883.

Disrib. : India (Himalayas from Kashmir to Bhutan).

P. microrhizoma (Clarke) Ching, Acta Phytotax. Sin. 16 (4) : 27. 1978. *Polypodium microrhizomum* Clarke in Hook. et Bak. Syn. Fil. 511. 1874. *Goniophlebium microrhizoma* (Clarke) Bedd. Ferns Brit. India Suppl. t. 384. 1876 ; Handb. Ferns Brit. India 322. 1883.

Distrib. : India (Kashmir to Bhutan), China.

P. subamoenus (Clarke) Ching, Acta Phytotax. Sin. 16 (4) : 27. 1978. *Polypodium subamoena* Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 550. t. 82. f. 2. 1880. *Goniophlebium subamoenum* (Clarke) Bedd. Handb. Ferns Brit. India 317. 1883.

Disrib. : India (Himalayas), Nepal.

P. wattii (Bedd.) Ching, Acta Phytotax. Sin. 16 (4) : 27. 1978. *Polypodium nipponicum* var. *wattii* Bedd. in Journ. Bot. 235. 1888. Bedd. Handb. Ferns Brit. India Suppl. 90. 1892.

Distrib. : India (Manipur).

Polypodium L. Sp. Pl. 2 : 1082. 1753.

Lectotype : **P. vulgare** L. Sp. Pl. 2 : 1085. 1753.

P. beddomei Bak. ex Hook. & Bak. Syn. Fil. ed. II. 344. 1874. *Goniophlebium molle* sensu Bedd. Ferns Brit. India t. 206. 1867 ; Handb. Ferns Brit. India 322. 1883. (*non* Schreb 1771 ; *nec.* Jacq. 1789).

Distrib. : India (Himalayas), Malay Peninsula, Burma.

P. vulgare L. Sp. Pl. 2 : 1085. 1753.

Distrib. : India (Himalayas), China, Japan, Europe, Africa, New-Zealand.

Pseudodrynaria C. Chr. et Ching, Sunyat. 6 : 10. 1941.

Type : **P. coronans** (Wall. ex Mett.) Ching, Sunyat. 6 : 10. 1941.

P. coronans (Wall. ex Mett.) Ching, Sunyat. 6 : 10. 1941. *Polypodium coronans* Wall. ex Mett. Pol. 121. n. 242. t. 3 f. 40, 41. 1857. *Drynnaria coronans* J. Sm. in Journ. Bot. 4 : 61. 1841 ; Bedd. Ferns Brit. India t. 13. 1865 ; Handb. Ferns Brit. India 338. t. 187. 1883.

Distrib. : Throughout India, China, Nepal, Burma, Bangladesh, Malay Peninsula, Hongkong, Formosa.

Pyrrosia Mirb. in Lam. et Mirb. Hist. Nat. Veg. 5 : 91. 1802. Ching, Bull. Chin. Bot. Soc. 1 (1) : 36. 1935 ; Nayar & Chandra., Bull. Nation. Bot. Gard. Lucknow no. 117. 60. 1965.

Type : **P. lingua** (Thunb.) Farwell, Amer. Midl. Nat. 12 : 302. 1931.

P. acrostichoidea (Forst.) Ching, Bull. Chin. Bot. Soc. 1 : 69. 1935. *Polypodium acrostichoides* Forst. Prod. 81. 1786. *Niphobolus acrostichoides* Bedd. Ferns. Brit. India t. 81. 1865 ; Handb. Ferns Brit. India 327. 1883.

Distrib. : S. India, Malesia, Polynesia, Sri Lanka extending to Queensland, China.

P. adnascens (Sw.) Ching, Bull. Chin. Bot. Soc. 1 : 45. 1935. *Polypodium adnascens* Sw. Syn. Fil. 25, 222. t. 2. f. 2. 1806. *Niphobolus adnascens* (Sw.) Kaulf. Enum. Fil. 124. 1824 ; Bedd. Handb. Ferns Brit. India 325. 1824 ; (*pro parte*).

Distrib. : India to S. China, Formosa, Malaya to Polynesia.

P. assimilis (Bak.) Ching, Bull. Chin. Bot. Soc. 1 : 49. 1935. *polypodium assimilis* Bak. Journ. Bot. 1875 : 201. 1875.

Distrib. : India (Eastern Himalayas), China.

P. beddomeana (Gies.) Ching, Bull. Chin. Bot. Soc. 1 : 68. 1935. *Niphobolus beddomeanus* Gies, Farnatt. Niph. 101. 1901. *Niphobolus costatus* Bedd. Ferns Brit. India t. 120. 1865. *Polypodium stigmosum* Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 553. 1880. *Niphobolus stigmosus* Bedd. Handb. Ferns Brit. India 328. 1883 (*pro parte*).

Distrib. : India (Himalayas), China.

P. boothii (Hook.) Ching, Bull. Chin. Bot. Soc. 1 : 66. 1935. *Polypodium boothii* Hook. Sp. Fil. 5 : 53. 1863 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 555. 1880. *Niphobolus boothii* (Hook.) Bedd. Ferns Brit. India t. 258. 1868.

Distrib. : India (Himalayas), Bhutan.

P. ceylanica (Gies.) Sledge, Bull. Brit. Mus. (N. H.) 2 : 134. 1960. *Niphobolus ceylanicus* Gies. Farnatt. Niph. 216. 1901.

Distrib. : South India, Sri Lanka.

P. costata (Presl ex Bedd.) Tagawa & Iwats. in Acta Phytotax. Geobot. 22 : 100. 1967. *Niphobolus costatus* Presl ex Bedd. Ferns Brit.

India t. 120. 1865. *Pyrrosia beddomeana* sensu Tagawa in Hara's Fl. East. Himal. 498. 1966.

Distrib. : India (Himalayas), China, Burma, N. Thailand.

P. flocculosa (D. Don) Ching, Bull. Chin. Bot. Soc. 1 : 66. 1935.
Polypodium flocculosum D. Don, Prod. Fl. Nepal. 1 : 1825. Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 554. 1880. *Niphobolus flocculosus* (D. Don) Bedd. Ferns Brit. India t. 162. 1867 ; Handb. Ferns Brit. India 331. t. 180. 1883.

Distrib. : India (Himalayas), Burma, Nepal, Bhutan, Annam, Tonkin, China.

P. heteracta (Mett. ex Kuhn) Ching, Bull. Chin. Bot. Soc. 1 : 57. 1935. *Polypodium heteractis* Mett. ex Kuhn in Linnaea 36 : 140. 1869 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 553. 1880. *Niphobolus heteractus* (Mett. ex Kuhn) J. Sm. Ferns Brit. & For. ed. 2. 296. 1877 : Bedd. Handb. Ferns Brit. India 327. 1883. *N. lingua* sensu Bedd. Ferns Brit. India Suppl. 22. t. 385. 1876 (non Spr. 1827 ; Bedd. Ferns Brit. India t. 240. 1865).

Distrib. : India (Himalayas), Burma, S. W. China.

P. laevis (J. Sm.) Ching, Bull. Chin. Bot. Soc. 1 : 52. 1935. *Niphobolus laevis* J. Sm. in Bedd. Ferns Brit. India t. 161. 1866 ; Handb. Ferns Brit. India 325. 1883. *Polypodium jaintense* Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 552. t. 82. f. 4. 1880.

Distrib. : India (Himalayas).

P. lanceolata (L.) Farwell. Amer. Midl. Nat. 12 : 245. 1931. *Acrostichum laaceolatum* L. Sp. Pl. 2 : 1067. 1753. *Polypodium adnascens* auct. Hook. & Bak. Syn. Fil. 349. 1867 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 552. 1880. *Niphobolus adnascens* Bedd. Handb. Ferns Brit. India 325. t. 176. 1883. (*non* Kaulf. 1824). *N. fissus* sensu Bedd. Ferns South. India t. 184. 1865. (*non*. Bl.).

Distrib. : India (Himalayas), Sri Lanka, Polynesia, China.

P. lingua (Thunb.) Farwell, Amer. Midl. Nat. 12 : 302. 1931. *Acrostichum lingum* Thunb. Fl. Jap. 330. t. 33. 1784. *Niphobolus heteractis* sensu Bedd. Handb. Ferns Brit. India 327. 1883 (*pro parte*).

Distrib. : North East India, China, Japan, Formosa, Tonkin.

P. macrocarpa (Willd.) Kaulf. Enum. Fil. 245. 1824 ; Manickam Ninan, New Botanist 1 : 46. 1976. *Polypodium macrocarpum* Willd. in Linn. Sp. Pl. 5 : 147. 1810. *Polypodium lepidotum* Bedd. Ferns South. India 60. t. 181. 1864. *nom illejet*.

Distrib. : South India, Trop. America, Australia.

P. mannii (Gies.) Ching, Bull. Chin. Bot. Soc. 1 : 55. 1935. *Niphobolus mannii* Gies. Farngatt. Niph. 107. 1901. *N. fissus* auct. Bedd. Handb. Ferns Brit. India 330. 1883 (*pro parte*; *non* Bl. 1828), *N. floccigerus* Bedd. Ferns Brit. India Suppl. 22. t. 386. 1876.

Distrib. : North East to South India, Sri Lanka, Malaysia, Madagascar.

P. mollis (Kunze) Ching, Bull. Chin. Bot. Soc. 1 : 53. 1935. *Niphobolus mollis* Kunze, Bot. Zeit. 6: 121. 1848. *N. fissus sensu* Bedd. Ferns Brit. India Corr. II 1870; Handb. Ferns Brit. India 330. 1883 (*pro parte. non* Bl. 1828). *N. porosus* sensu Bedd. Ferns South. India t. 183. 1863. *Polypodium fissum* Hook. et Bak. Syn. Fil. 351. 1874; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 554. 1880.

Distrib. : India (Himalayas), Sri Lanka, China, Burma, Java, Tonkin.

P. nayariana Ching & Chandra, Amer. Fern Journ. 54 : 62. 1964.

Distrib. : India (Manipur).

P. nuda (Gies.) Ching, Bull. Chin. Bot. Soc. 1 : 70. 1935. *Niphobolus nudus* Gies. Farngatt. Niph. 149. 1901. *Polypodium adnascens sensu* Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 552. 1880 (*pro parte*). *Niphobolus adnascens sensu* Bedd. Handb. Ferns Brit. India 325. 1883 (*pro parte*).

Distrib. : India (Himalayas), China.

P. nummulariaefolia (Sw.) Ching, Bull. Chin. Bot. Soc. 1 : 47. 1935. *Acrostichum nummulariaefolium* Sw. Syn. Fil. 191, 418. t. 2. f. 1. 1806. *Polypodium nummulariaefolium* Mett. Farngatt. Poly. 123. t. 3. f. 9, 10. 1857; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 553. 1880; Bedd. Ferns Brit. India t. 320. 1869. *Niphobolus nummulariaefolius* J. Sm. Journ. Bot. 3 : 396. 1841; Bedd. Ferns. Brit. India. t. 320. 1869; Handb. Ferns Brit. India 334. 1883.

Distrib. : India (Himalayas), Bhutan. Bangladesh, Malaya, Philippines, China.

P. obovata (Bl.) Ching, Bull. Chin. Bot. Soc. 1 : 47. 1935. *Acrostichum obovatum* Bl. Enum. Pl. Jav. 102. 1828. *Polypodium nummulariaefolium* var. *obovatum* Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 554. 1880.

Distrib. : North-East India, extending to Java.

P. pannosa (Mett.) Ching, Bull. Chin. Bot. Soc. 1 : 58. 1933. *Polypodium pannosum* Mett. apud Kuhn in Linnaea 36 : 141. 1869. *Niph-*

bolus panosus (Mett.) Bedd. *Ferns Brit. India Suppl.* 22. 1876 ; *Handb. Ferns Brit. India* 328. t. 177. 1883. *Niphobolus lingua* Bedd. *Ferns South. India* t. 240. 1865 (*non* Spr. 1827).

Distrib. : India (Himalayas), Burma, Sri Lanka.

P. pekinensis (C. Chr.) Ching, *Bull Chin. Bot. Soc.* 1 : 51. 1935.
Cyclophorus pekinensis C. Chr. *Ind. Fil.* 200. 1905. *Polypodium davidi* Bak. *Ann. Bot.* 5 : 572. 1891.

Distrib. : India (Himalayas), China.

P. stenophylla (Bedd.) Ching, *Bull. Chin. Bot. Soc.* 1 : 55. 1935.
Niphobolus fissus var. *stenophyllus* Bedd. *Handb. Ferns Brit. India Suppl.* 92. 1892. *Polypodium fissum* var. *floccigerum* Clarke, *Trans. Linn. Soc. Lond. II. Bot.* 1 : 550. 1880.

Distrib. : (Eastern India), Bhutan.

P. stictica (Kunze) Holtt. *Novit. Bot. Inst. Univ. Carol. Prag.* 31. 1968. *Niphobolus Sticticus* Kunze in *Linnaea* 24 : 257. 1851.

Distrib. : S. India, Malay Peninsula.

P. stigmosa (Sw.) Ching, *Bull. Chin. Bot. Soc.* 1 : 67. 1935.
Polypodium stigmosum Sw. Schrad. *Journ. Bot.* 1800 (2) : 21. 1801 ; Clarke, *Trans. Linn. Soc. Lond. II. Bot.* 1 : 553. 1880 (*pro parte*). *Niphobolus stigmosus* sensu Bedd. *Handb. Ferns Brit. India* 328. t. 178. 1883 (*pro parte*).

Distrib. : India (Himalayas, S. India), Nepal, Bhutan, Burma. Malesia, northward to Cochin - China.

P. subfurfuracea (Hook.) Ching, *Bull. Chin. Bot. Soc.* 1 : 68. 1935.
Polypodium subfurfuraceum Hook. *Sp. Fil.* 5 : 52. 1863 ; Clarke, *Trans. Linn. Soc. Lond. II. Bot.* 1 : 553. 1880. *Niphobolus subfurfuraceus* Bedd. *Ferns. Brit. India* t. 259. 1868 ; *Handb. Ferns Brit. India* 329. 1883.

Distrib. : North - East India, Burma, Bhutan, China.

P. subvelutina (Christ) Ching. *Bull. Chin. Bot. Soc.* 1 : 50. 1935.
Niphobolus subvelutinus Christ, *Ann. Cons. Jard. Bot. Genéve* 7 - 8 : 331. 1904.

Distrib. : India (Himalayas)

P. varia (Kaulf.) Farwell, *Amer. Midl. Nat.* 12 : 302. 1931. *Niphobolus varius* Kaulf. *Enum. Fil.* 125. 1824. *N. adnascens* sensu Bedd. *Handb. Ferns. Brit. India* 325. 1883 (*pro parte*).

Distrb. : North - East India, Malesia to Polynesia.

Thylacopteris Kunze ex J. Sm. Hist. Fil. 87. 1875.

Type : **T. papillosa** (Bl.) J. Sm. Hist. Fil. 87. 1875.

T. papillosa (Bl.) J. Sm. Hist. Fil. 87. 1875. *Polypodium papillosum* Bl. Enum. Pl. Jav. 131. 1828.

Distrib. : India (Arunachal Pradesh), China.

Tricholepidium Ching, Acta Phototax. Geobot. 29 : 41. 1978.

Type : **T. normale** (D. Don.) Ching, Acta Phytotax. Geobot. 29 : 41. 1978.

T. normale (D. Don) Ching, Acta Phytotax. Geobot. 29 : 41. 1978.
Polypodium normale D. Don, Prod. Fl. Nepal 1. 1825. *Pleopeltis normalis* Moore, Ind. Fil. 347. 1848 ; Bedd. Handb. Ferns Brit. India 353. 1883.

Distrib. : India (Himalayas), Nepal, Bhutan, Burma, S. W. China.

DIPTERIDACEA E. (Diels) Seward & Dale

Dipteris Reinw. Syll. Nov. Ratisb. 2 : 3. 1824.

Type : **D. conjugata** Reinw. Syll. Nov. Ratisb. 2 : 3. 1824

D. wallichii (R. Br.) Moore, Ind. Fil. 80. 1857 ; Bedd. Ferns Brit. India. t. 80. 1865 ; Handb. Ferns Brit. India 334. t. 184. 1883.

Distrib. : India (Arunachal Pradesh, Assam, Meghalaya), Bangladesh, China.

PLATYCYRIACEAE (Nayar) Ching

Platycerium Desv. Mém. Linn. Soc. Paris 6 : 213. 1827.

Type : **P. alcicorne** Desv. Mém. Linn. Soc. Paris 6 : 213. 1827.

P. alcicorne Desv. Mém. Linn. Soc. Paris 6 : 213. 1827. *Acrostichum alcicorne* Wall. List no. 19. 1828 (*nom. nud.*) ex Bedd. Ferns Brit. India t. 108. 1865. *Platycerium wallichii* Hook. Gard. Chron. 1858 : 765. 1858 ; Bedd. Handb. Ferns Brit. India 445. t. 272. 1883 ; Suppl. 107. 1892.

Distrib. : India (Manipur), Burma, Malaya Peninsula.

DRYNARIACEAE Ching

Drynaria (Bory) J. Sm. in Hook. Journ. Bot. 4 : 60. 1841 ; Nayar, Bull. Nation. Bot. Gard. Lucknow no. 56.9. 1961. *nom. cons.*

Type : **D. quercifolia** (L.) J. Sm. in Hook. Journ. Bot. 3 : 398. 1841.

D. mollis Bedd. Ferns Brit. India t. 216. 1867 ; Handb. Ferns Brit. India 341. t. 190. 1883.

Distrib. : India (N.W. Himalayas), China.

D. propinqua (Wall. ex Mett.) J. Sm. apud. Bedd. Ferns Brit. India t. 160. 1866; Handb. Ferns Brit. India. 339. t. 189. 1883. *Polypodium propinquum* Wall. ex Mett. in Abh. Senckneb. Naturf. Ges. 2 : 120. t. 3. f. 50. 1857. *Pleopeltis parishii* Bedd. Ferns Brit. India t. 135. 1867. *Drynaria parishii* (Bedd.) Bedd. Ferns Brit. India Suppl. 24. 1876.

Distrib. : India (Himalayas), Burma, China, Malay Peninsula, Malesian Islands.

D. quercifolia (L.) J. Sm. in Hook. Journ. Bot. 3 : 398. 1841 ; Bedd. Handb. Ferns Brit. India 341. t. 191. 1883. *Polypodium quercifolium* L. Sp. Pl. 2 : 1087. 1753 ; Bedd. Ferns Brit. South. India t. 187. 1865.

Distrib. : Throughout India in the plains and above in hilly regions, Sri Lanka, South to Fiji and trop. Australia.

D. sparsisora (Desv.) Moore, Ind. Fil. CCCXLVIII. 1857. *Polypodium sparsisorum* Desv. Berl. Mag. 5 : 315. 1811. *Drynaria linnaei* (Bory) Bedd. Ferns Brit. India t. 315. 1869 ; Handb. Ferns Brit. India 343. 1883.

Distrib. : NE India, Sri Lanka, Malay Peninsula.

SCHIZAEACEAE Kaulf.

Actinostachys Wall. ex Hook. Gen. Fil. t. 111. 1842.

Type : **A. digitata** (L.) Wall. ex Hook. Gen. Fl. t. 111. 1842.

A. digitata (L.) Wall. ex Hook. Gen. Fil. t. 111. 1842. *Acrostichium digitatum* L. Sp. Pl. 2 ; 1068. 1753. *Schizaea digitata* (L.) Sw. Syn. Fl. 150. t. 4. 1806 ; Bedd. Ferns South. India. t. 268. 1865 ; Handb. Ferns Brit. India 452. t. 279. 1883.

Distrib. : India (North India, Meghalaya, S. India), Bangladesh, Sri Lanka, Malay Peninsula, Malesian Islands, Fiji, Philippines.

Schizaea Sm. Mém. Acad. Sci. Turin. 5 : 419. t. 19. f. 9. 1793 ; Panigr. & Dixit. Proc. Autm. Sch. Bot. 207. 1966.

Type : *S. dichotoma* (L.) Sm. Mém. Acad. Sci. Turin. 5 : 419 : 1793.

S. dichotoma (L.) Smith, Mém. Acad. Turin. 5 : 422. t. 9. 1973 ; Bedd. Ferns South. India t. 65 1863 ; Handb. Ferns Brit. India 452. t. 278. 1883. *Acrostichum dichotomum* L. Sp. Pl. 2 : 1068. 1753.

Distrib. : South India, Malay Peninsula, Tropical America, West-Indies, Australia, New Zealand, Polynesia, Philippines, Mascareen Islands.

ANEMIACEAE Link

Anemia Sw. Syn. Fil. 6, 155. 1806 ; Panigr. & Dixit, Proc. Autm. Sch. Bot. 231. 1966. *nom. cons.*

Type : *A. phyllitidis* (L.) Sw. Syn. Fil. 155. 1806.

A. tomentosa (Sav.) Sw. Syn. Fil. 156. 1806 ; Bedd. Handb. Ferns Brit. India 453. t. 280 1883. *Osmunda tomentosa* Sav. in Link Encycl. 4 : 652. 1797. *Anemia wightiana* (Gard.) Bedd. Fcrns. South. India 22. t. 66. 1863.

Distrib. : South India, Mexico, Peru, West Indies.

LYGODIACEAE Presl

Lygodium Sw. Schrad. Journ. Bot. 1800 (2) : 7, 106. 1801 : Holtt. Fl. Males. Ser. II. 1 (1) : 44. 1959, Panigr. & Dixit, Proc. Autm. Sch. Bot. 212. 1966. *nom. cons.*

Type : *L. scandens* (L.) Sw. Journ. Bot. 1806 (2) : 100. 1801.

L. altum (Clarke) v. A. v. R. Mal. Ferns 114. 1909. *L. flexuosum* var. *alta* Clarke, Trans Linn. Soc. Lond. II. Bot. 1 : 584. 1880.

Distrib. : India (Assam).

L. circinatum (Burm.) Sw. Syn. Fil. 153. 1806 ; Bedd. Handb. Ferns. Brit. India 455. f. 281. 1883. *Ophioglossum circinatum* Burm. Fl. Ind. 228. 1768. *Ophioglossum furcatum* Roxb. ex Griff. Calc. Journ. Nat. Hist. 4 : 478. 1844.

Distrib. : India (Andaman & Nicobar Islands, North East India), Sri Lanka, China, Siam, Malaya Peninsula, Malesian Islands. Philippines ; Hongkong and chusan. Its occurrence from the main land of India is doubtful.

L. flexuosum (L.) Sw. Schrad. Journ. 1800 (2) : 106. 1801 ; Bedd. Ferns South. India t. 63. 1863 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1: 584. 1880 ; Handb. Ferns Brit. India 457. f. 283. 1883. *Ophioglossum flexuosum* L. Sp. Pl. 2 : 1063. 1753.

Distrib. : Throughout India from the plains to the mountainous regions, Sri Lanka, Malay Peninsula, Malesian Islands, Philippines, China, Australia, Pacific.

L. giganteum Tagawa & Iwats. Acta Phytotax. Geobot. 22 : 97. f. 1. 1967.

Distrib. : India (Himalayas), Japan.

L. japonicum (Thunb.) Sw. Journ. Bot. 1800 (2) : 106. 1801 ; Bedd. Ferns South. India t. 64. 1863 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 584. 1880 ; Bedd. Handb. Ferns Brit. India 452. 1883. *Ophioglossum japonicum* Thunb. Fl. Jap. 328. 1784.

Distrib. : India (North-East India, South India and Western Himalayas), Japan, Sri Lanka, Korea, Japan, Siam, Indo China, China, Malesian Islands, Philippines, America.

L. longifolium (Willd.) Sw. Schrad. Journ. Bot. 1800 (2) : 305. 1801. *Hydroglossum longifolium* Willd. Abh. Kurf. Mainz. Akad. Wiss. Erfurt. 1802 : 22. t. 2. 1802.

Distrib. : South India, China, Malaya, Malaysia.

L. mearnsii Copel. Philip. Journ. Sci. Bot. 3 : 37. 1908.

Distrib. : India (Eastern Himalayas), Philippines.

L. microphyllum (Cav.) R. Br. Prod. Fl. Nov. Holl. 162. 1810 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 583. 1880 ; Bedd. Handb. Ferns. Brit. India 455. t. 282. 1883. *Ugenia microphyllum* Cav. Icon. Descr. Pl. 6 : 76. t. 595. 1801 ; Bedd. Ferns South. India t. 61. 1863. *Ophioglossum filiforme* Roxb. ex Griff. Calc. Journ. Nat. Hist. 4 : 476. t. 26. f. 3. 1844.

Distrib. : India (Assam, Sikkim, Meghalaya, Madhya Pradesh, South India, Andaman & Nicobar Islands), Malay Peninsula, Malesian Islands, Sri Lanka, Africa to Polynesia and Australia.

L. polystachyum Wall. ex Moore, Gard. Chron. 671. 1859 ; Bedd. Ferns. Brit. India t. 300. 1868 ; Handb. Ferns Brit. India 458. f. 284. 1883.

Distrib. : India (Assam), Burma, Siam, China, Indo-China, Malaysia, Malay Peninsula.

L. salicifolium Presl, Suppl. Tent. Pterid. 102 : 1845. (*pro parte* excl. Pl. Wallich, *pro parte* & *Syn. L. semibipinnatum* wall.).

Distrib. : Eastern India, Siam, Indo-China to Yunnan, Formosa, New Guinea, Malaysia, Malay Peninsula.

SINOPTERIDACEAE Koidz.

Doryopteris J. Sm. in Hook. Journ. Bot. 3 : 404. 1841.

Lectotype : **D. palmata** (Willd.) J. Sm. in Hook. Journ. Bot. 4 : 163. 1841.

D. concolor (Langsd. et Fisch.) Kuhn, v. Deck. Reis. 3 (3) : Bot. 19. 1879. *Pteris concolor* Langsd. et Fisch. Icon. Fil. 19. t. 21 1810. *Pellaea geranifolia* (Raddi) Féé. Mém. Fern. Foug. 5 : 130. 1850 52 ; Bedd. Ferns South. India t. 13. 1864.

Distrib. : S. India and tropics of the world.

D. kirkii (Hook.) Alston, Bol. Soc. Broter. ser. 2A, 30 : 14. 1956. *Cheilanthes kirkii* Hook. Cent. Ferns t. 81. 1861. *Doryopteris concolor* var. *kirkii* (Hook.) Wiss. Ergeb. Schwed. Rhod. Kongo Exp. 1 : 4. 1941.

Distrib. : S. India, Orissa, Karnataka, Andhra Pradesh, Kerala, Sri Lanka, Africa.

D. ludens (Wall. ex Hook.) J. Sm. Hist. Fil. 189. 1875 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 470. 1880 ; Handb. Ferns Brit. India 120. t. 82. 1883. *Pteris ludens* Wall. ex Hook. Sp. Fil. 2 : 210. 1858. *Litobrachia ludens* (Wall. ex Hook.) Bedd. Ferns Brit India t. 27. 1865. *L. pedata* Presl, Tent. Pterid. 149. t. 5. f. 25. 1836 ; Bedd. Ferns Brit. India t. 26. 1865. *Pteris multifida* Roxb. ex Griff. Calc. Journ. Nat. Hist. 4 : 507. 1844.

Distrib. : North - East to South India, Burma, Bangladesh, Orissa, Philippine Islands.

Notholaena R. Br. Prod. Fl. Nov. Holl. 145. 1810.

Lectotype : *N. marantae* (L.) Desv. Journ. de Bot. appl. 1 : 92. 1813.

N. lanuginosa Desv. ex Poir. in Lam. Encycl. Suppl. 4 : 110. 1816. *N. vellea* sensu Bedd. Handb. Ferns Brit. India 375. 1883 ;

Distrib. : India (North West Himalayas), Afghanistan, Pakistan, S, Europe, Africa.

N. marantae (L.) Desv. Journ. de Bot. appl. 1 : 92. 1813 ; Bedd. Handb. Ferns Brit. India 393. t. 213. 1883. *Acrostichum marantae* L. Sp. Pl. 2 : 1071. 1753.

Distrib. : India (North West Himalayas, Sikkim), S. Europe, Macronesia to the Caucasus and Abyssinia.

Mildella Trev. Rendic. Inst. Lomb. Sci. et Lett. 119 : 810. 1876.

Type : *M. intramarginalis* Trev. Rendic. Inst. t. Lomb. Sci. et lett. 119 : 810. 1876.

M. nitidula (Wall. ex Hook.) Hall & Lelling. Amer. Fern Journ. 57 (3) : 127. 1967. *Cheilanthes nitidula* Wall. ex Hook. Sp. Fil. 2 : 112. 1852 ; Bedd. Ferns Brit. India t. 222. 1867.

Distrib. : India (N. W. Himalayas), Nepal, West Pakistan.

Pellaea Link, Fil. Sp. 59. 1842. *nom cons.*

Type : *P. atropurpurea* (L.) Link, Fil. Sp. 59. 1842.

P. boivini Hook. Sp. Fil. 2 : 147. t. 118 A. 1858 ; Bedd. Handb. Ferns Brit. India t. 53. 1883. *Pteris boivini* Bedd. Ferns South. India 12. t. 36. 1863.

Distrib. : South India, Madagascar, Mauritius.

P. calomelanos (Sw.) Link, Fil. Sp. Hort. Berol. 61. 1841 ; Bedd. Handb. Ferns Brit. India 104. 1883. *Pteris calomelanos* Sw. Schrad. Journ. Bot. 1800 (2) : 70. 1801 ; Bedd. Ferns Brit. India t. 22. 1866.

Distrib. : India (North - West Himalayas), Africa.

P. seticaulis (Hook.) Ghosh, Journ. Bomb. Nat. Hist. Soc. 1984. *Pteris seticaulis* Hook. Icon. Pl. 3. t. 207. 1840. *Platyloma falcatum* var. *setosum* Bedd. Ferns South. India 7. t. 22. 1863. *Pellaea falcata* sensu Bedd. Farns South. India t. 22. 1863 ; Handb. Ferns Brit. India 102. t. 54. 1883. (*non R. Br.*)

Distrib. : South India, Sri Lanka, Malaya Peninsula, Australia, New Zealand.

CHEILANTHACEAE Nayar

Cheilanthes Sw. *Syn. Fil.* 5 : 126. 1806 ; Nayar, *Bull. Nation. Bot. Gard.* (6) : 1-35. Lucknow no. 68. 126. 1962.

Type : **C. micropteris** Sw. *Syn. Fil.* 5 : 126, 334. t. 3. f. 5. 1806.

C. albomarginata Clarke, *Trans. Linn. Soc. Lond. II. Bot.* 1 : 456. t. 52. 1880 ; Bedd. *Handb. Ferns Brit. India* 94. 1883 ; *Cheilanthes farinosa* var. *albomarginata* Bedd. *Handb. Ferns Brit. India Suppl.* 22. 1892. *Aleuritopteris albomarginata* (Clarke) Ching, *Hong Kong Nat.* 10 : 199. 1941.

Distrib. : Throughout India in hilly regions, Nepal, Bhutan, China.

C. anceps Blanf. *Journ. Simla Nat. Hist. Soc.* 1 : 21. 1886 ; *C. farinosa* var. *anceps* (Blanf.) Blanf. *Journ. Asiat. Soc. Bengal* 57. pt. II. (4) : 249. 1888 ; Bedd. *Handb. Ferns Brit. India Suppl.* 21. 1892. *Aleuritopteris farinosa* var. *anceps* (Blanf.) Ching. *Hong Kong Nat.* 194. 1941. *Aleuritopteris anceps* (Blanf.) Panigr. *Bull. Bot. Surv. India* 2 : 321. 1961.

Distrib. : Throughout India, China, Sri Lanka, Burma, Philippines, East Africa and its Islands, Trop. America, Arabia.

C. argentea (Gmel.) Kunze, *Linnaea* 23 : 242. 1850 ; Bedd. *Ferns Brit. India* t. 143. 1867 ; *Handb. Ferns Brit. India* 95. t. 48. 1883. *Pteris argentea* Gmel. *Nov. Comm. Petr.* 12 : 519. t. 12. f. 2. 1768. *Aleuritopteris argentea* (Gmel.) Fée, *Gen. Fil.* 154. 1850-52.

Distrib. : India (Meghalaya), Burma, China, Japan.

C. belangiri (Bory) C. Chr. *Ind. Fil.* 172. 1905. *Pteris belangiri* Bory, *Bél. Voy. Bot.* 2 : 44. 1833. *C. varians* Wall. ex Hook. Sp. *Fil.* 2 : 89. t. 103 A. 1852 ; Bedd. *Ferns South. India* t. 189. 1867 ; *Handb. Ferns Brit. India* 91. 1883.

Distrib. : India (Bengal plains to Assam, Meghalaya, S. India), Burma, China, Luzon.

C. bullosa Kunze, *Linnaea*. 24 : 274. 1851. *Aleuritopteris bullosa* (Kunze) Ching, *Hong Kong Nat.* 10 : 202. 1941.

Distrib. : India (Western Ghats, Nilgiri Hills). China, Australia.

C. chrysophylla Hook. Icon. Pl. 2 : 113. 1852. *Cheilanthes farinosa* var. *chrysophylla* (Hook.) Bedd. Handb. Ferns. Brit. India 93. 1883. *C. argentea* var. *chrysophylla* Hook. Syr. Fil. 142. 1865. *C. argentea* var. *sulphurea* Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 93. 1880. *Aleuritopteris chrysophylla* (Hook.) Ching, Hong Kong Nat. 10 : 201. 1941.

Distrib. : E. Himalayas.

C. dalhousiae Hook. Sp. Fil. 2 : 80. t. 78 B. 1852. *C. farinosa* var. *dalhousiae* (Hook.) Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 457. t. 51. 1880 ; Bedd. Handb. Ferns Brit. India 93. 1883. *Aleuritopteris dalhousiae* (Hook.) Ching, Hong Kong Nat. 10 : 204. 1941.

Distrib. : India (Himalayas).

C. dubia Hope, Journ. Bomb. Nat. Hist. Soc. 12 : 528. 1899. *Aleuritopteris dubia* (Hope) Ching, Hong Kong Nat. 10 : 200. 1941.

Distrib. : India (N. W. Himalayas).

C. duthei Bak. Ann. Bot. 5 : 210. 1891 ; Bedd. Handb. Ferns Brit. India Suppl. 20. 1892. *Aleuritopteris duthei* (Bak.) Ching, Bull. Fan. Mem. Inst. Biol. ser. 2. 1 : 271. 1949.

Distrib. : India (N. W. Himalayas), China.

C. farinosa (Forsk.) Kaulf. Enum. Fil. 212. 1824 ; Bedd. Ferns South. India t. 191. 1867 ; Handb. Ferns Brit. India 92. 1883. *Pteris farinosa* Forsk, Fl. Aegypt. Arab 187. 1775. *Aleuritopteris farinosa* (Forsk.) Fée, Gen. Fil. 154. t. 12 B. f. 1. 1850 52.

Distrib. : Throughout India, Sri Lanka, Burma, Trop. America, Java, Philippines, East Africa and its Islands and Arabia.

C. flaccida (Bedd.) Mehra & Bir, Res. Bull. (N. S.) Punjab Univ. 15 : 110. 1964. *C. farinosa* var. *flaccida* Bedd. Handb. Ferns Brit. India 93. 1883. *C. dalhousiae* sensu Bedd. Ferns South. India t. 192. 1867. *Aleuritopteris flaccida* (Bedd.) Nayar & Kaur, Comp. to Bedd. Handb. 25. 1974.

Distrib. : S. India, Eastern Himalayas.

C. fragrans (L. f.) Sw. Syn. Fil. 127, 325. 1806. *Adiantum fragrans* L. f. Suppl. Pl. 447. 1781.

Distrib. : India (N. W. Himalayas), Western Asia, S. Europe, N. Africa.

C. grisea (Blanf.) Blanf. Journ. Simla Nat. Hist. Soc. 1 : 21. 1886 ; *C. farinosa* var. *grisea* Blanf. Journ. Asiat. Soc. Bengal 57. t. II. (4) :

250. 1881 ; Bedd. Handb. Ferns Brit. India Suppl. 21. 1892. *Aleuritopteris grisea* (Blanf.) Panigr. Bull. Bot. Surv. 214. 1824 (*pro parte*).

Distrib. : North East to Central India, Sri Lanka, China.

C. hancockii Bak. Kew Bull. 1895: 54 : 1895.

Distrib. : India (Himalayas), China.

C. keralensis Nair & Ghosh, Journ. Indian Bot. Soc. 55 : 52. 1976.

Distrib. : S. India (Kanya Kumari distt.).

C. kuhnii Milde, Bot. Zeit. 25 : 149. 1867.

Distrib. : India (Himalayas), China, South Korea.

C. laxa Moore, Ind. Fil. 245. 1861.

Distrib. : S. India, Sri Lanka.

C. persica (Bory) Mett. ex Kuhn, Fil. Afr. 73. 1868. *Notholaena persica* Bory, Bél, Voy. Bot. 2 : 23. 1833. *C. szovitsii* Fisch. & Meyer, Bull. Soc. Misc. 6 : 260. 1883 (*nom. nud.*) ex Hook. Syn. Fil. 139. 1865 ; Bedd. Ferns Brit. India t. 145. 1867 ; Handb. Ferns Brit. India 89. 1883.

Distrib. : India (N W. Himalayas), Kabul, Asia Minor, Persia, S. Europe.

C. pteridoides (L.) C. Chr. Ind, Fil. 178. 1905. *Polypodium pteridoides* Reich. L. Syst. Pl. ed. novis 4 : 424. 1780. *Polypodium pteridoides* L. Mant. 307. 1771 ; Bedd. Ferns Brit. India t. 328. 1870. *Cheilanthes fragrans* Webb. ex Benth. Hist. nat. Canr. 3 (2) : pars. 3. 452. 1847 ; Bedd. Handb. Ferns Brit. India 88. 1883.

Distrib. : India (N W Himalayas), Cabul, all round the Mediterranean, Canaries, Maderia.

C. rufa D. Don, Prod. Fl. Nepal 16 : 1825 ; Bedd. Ferns Brit. India t. 144. 1867 ; Handb. Ferns Brit. India 94. 1883. *Aleuritopteris rufa* (D. Don) Ching, Hong Kong Nat. 10 : 200. 1941.

Distrib. : India (North West Himalayas), China.

C. subdimorpha (Clarke & Bak.) Nayar & Kaur, Comp. to Bedd. Handb. 26. 1974. *C. farinosa* var. *subdimorpha* Clarke & Bak. Journ. Linn. Soc. 24 : 411. 1888 ; Bedd Handb. Ferns Brit. India Suppl. 21. 1882.

Distrib. : India (Eastern Himalayas), China.

C. subvillosa Hook. Sp. Fil. 2 : 87. t. 98B. 1852 ; Bedd. Ferns. Brit. India t. 142. 1866 ; Handb. Ferns Brit. India 93. 1883. *Aleuritopteris subvillosa* (Hook.) Ching, Hong Kong Nat. 10 : 203. 1941.

Distrib. : India (Himalayas), China.

C. swartzii Webb. et Benth, Hist. Nat. Cenar 3 (3) : 453. 1847 ; Panigr. & Basu, Taxon 31 (1) : 103. 1982. *C. fragrans sensu* Sw. Syn. Fil. 127. 325. t. 3. f. 3. 1806. pp. maj. incl. descr. et Sp. excl. type et Syn. non *Polypodium fragrans* L. (1771) nec. L. (1783). *C. mysurensis* Wall. ex Hook. Sp. Fil. 2 : 94. 100 A 1852 ; Bedd. Ferns South, India t. 190. 1865; Handb. Ferns Brit. India 89. t. 46. 1883, (*non Asplenium mysurensis* Heyne ex Roth. 1821).

Distrib. : India (Orissa, Andhra Pradesh, Tamil Nadu, Karnataka, & Kerala), Sri Lanka, Burma.

C. tenuifolia (Burm. f.) Sw. Syn. Fil. 129, 332. 1806 ; Bedd. Ferns South. India t. 188. 1865 ; Handb. Ferns Brit. India 92. 1883. *Trichomanes tenuifolia* Burm. f. Fl. Ind. 237. 1768.

Distrib. : Throughout India, Sri Lanka, China, Formosa, Philippines, Malay Islands, Australia, New Zealand, Polynesia.

C. thwaitesii Mett. ex kuhn, Linn. Relege Mett. 36 : 82. 1869.

Distrib. : South India, Sri Lanka.

C. trichophylla Bak. Ann. Bot. 5 : 211. 1891 ; Bedd. Handb. Ferns Brit. India Suppl. 20. 1892.

Distrib. : India (Sikkim), China.

CRYPTOGRAMMACEAE Pic. Ser.

Cryptogramma R. Br. in Franklin Narr. Journ. Polar Sea 767. 1823.

Type : **C. acrostichoides** R. Br. in Franklin Narr. Journ. Polar Sea 767. 1823.

C. brunonianana Wall. ex Hook. & Grev. Icon. Fil. t. 158. 1829 ; Bedd. Ferns Brit. India t. 164. 1866. *C. crispa* sensu Hook. & Bak. Syn. Fil. 144. 1868 (*pro parte*) ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 459. 1880 ; Bedd. Handb. Ferns Brit. India 98. t. 50. 1883 ; *C. crispa* f. *indica* Hook. Sp. Fil. 2. 129 ; 1858.

Distrib. : India (Himalayas, Andaman Islands), China, Taiwan, Afghanistan.

C. stelleri (Gmel.) Prantl in Bot. Jahrb. 3 : 413. 1882. *Pteris stelleri* Gmel. in Nov. Comm. Petr. 12. 519. t. 12. f. 1. 1768. *Pellaea gracilis* Hook. Sp. Fil. 2 : 138. t. 133 B. 1858 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 460. 1880.

Distrib. : India (Eastern Himalayas), China, Nepal, Bhutan, Formosa, Japan, Siberia, Taiwan, N. America.

Onychium Kaulf. Berlin Jahrb. Pharm. 21 : 45. 1820 ; Ching, Ling-nan Sc. Journ. 13 (3) : 493 501. 1934.

Type : **O. capense** Kaulf. Enum. Pl. 145. t. 1. f. 8 1824.

O. contiguum Hope, Journ. Bomb. Nat. Hist. Soc. 13 : 444. 1901.

O. iaponicum var. *multisectum* Bedd. Handb. Ferns Brit. India Suppl. 23. 1892.

Distrib. : India (Kashmir to Sikkim), Tibet, Thai, China.

O. fragile Verma & Khullar, Nova Hedw 9 : 85. 1965.

Distrib. : India (Himalayas).

O. ipii Ching, Lingnan Sci. Journ. 15 : 282. 1936.

Distrib. : India (Sikkim), China.

O. japonicum (Thunb.) Kunze, Bot. Zeit. 6 : 507. 1848 ; Bedd. Ferns. Brit. Indin t. 2. 1969. Handb. Ferns Brit. India 96. 1883. *Trichomanes japonicum* Thunb. Fl. Jap. 340. 1784.

Distrib. : Himalayas (Punjab to Assam), Burma, China, Korea, Formosa, Japan, Philippines, Java.

O. lucidum (D. Don) Spr. Syst. 4 : 66. 1827. *Leptostegia lucida* D. Don. Prod. Fl. Nepal 14 : 1825.

Distrib. : India (North West to Eastern Himalayas), China.

O. plumosum Ching, Lingnan Sci. Journ. 13 (3) : 499. 1934 ; *Ophio-glossum japonicum* var. *parisorum* R. Bona Parte, Notes Pterid. 14. 1923.

Distrib. : India (Sikkim), China.

O. siliculosum (Desv.) C. Chr. Ind. Fil. 468. 1906. *Pteris siliculososa* Desv. Berlin Mag. 5 : 324. 1811. *O. auratum* Kaulf. Enum. Fl. 144. 1824 ; Bedd. Handb. Ferns. Brit. India 96. 1883.

Distrib. : Himalayas (Nepal to Assam), Burma, Indo China, Malaya Peninsula, China, Formosa, Philippines, Malaya Islands, New Guinea, Polynesia.

O. tenuifrons Ching, Lingnan Sci. Journ. 14 (3) : 500. 1934.

Distrib. : India (Eastern Himalayas), China.

ACTINIOPTERIDACEAE Pic. Ser.

Actiniopteris Link, Fil. Sp. 79. 1841.

Lectotype : **A. radiata** (Sw.) Link, Fil. Sp. 80. 1841.

A. radiata (Sw.) Link, Fil. Sp. 80. 1841 ; Bedd. Ferns South. India. t. 124. 1865. *Asplenium radiatum* Sw. Schrad. Journ. Bot. 1800 (2) : 50. 1801. *Actiniopteris dichotoma* Kuhn, Bot. Zeit. 504. 1871 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 505. 1880 ; Bedd. Handb. Ferns Brit. India 197. t. 98. 1883.

Distrib. : Throughout India in drier regions, Sri Lanka, Africa, Mascareen Islands, Persia, Afghanistan.

PTERIDACEAE Ching

Pteris L. Sp. Pl. 2 : 1073. 1753.

Lectotype : **P. longifolia** L. Sp. Pl. 2 : 1074. 1753.

P. almeidiana Bole & Almeida, Journ. Bomb. Nat. Hist. Soc. 74 : 320. 1977.

Distrib. : India (Maharashtra).

P. amoena Bl. Enum. Pl. Jav. 210. 1828.

Distrib. : North East India, Java.

P. argyraea Moore in Garden. Chron. 671. 1959. *P. quadriaurita* var. *argentea* Bedd. Handb. Ferns Brit. India 111. 1883.

Distrib. : South India.

P. aspercaulis Wall. ex Ag. Rec. Sp. Gen. Pterid. 22. 1839. *P. quadriaurita* var. *aspercaulis* (Wall. ex Ag.) Bedd. Handb. Brit. India 111. 1883.

Distrib. : Throughout India in mountainous regions, Nepal, Burma, Thailand, China.

P. asperula J. Sm. ex Hieron. Hedw. 55 : 361. 1914. *P. quadriaurita* var. *setigera* Bedd. Ferns Brit. India t. 202. 1867. *P. quadriaurita* var. *hamulosa* Wall. ex Bedd. Handb. Ferns Brit. India Suppl. 23. 1892. *P. quadriaurita* var. *asperula* (J. Sm. ex Hieron.) Bedd. Handb. Ferns Brit. India Suppl. 24. 1892.

Distrib. : S. E. India, Burma.

P. barbigera Ching, Lingnan Sci. Journ. 15 : 395. 1936 ; Pic. Ser. Ind. Fil. Suppl. 4 : 260. 1965.

Distrib. : North India.

P. biaurita L. Sp. Pl. 2 : 1076. 1753 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 469. 1880. *Campteris biaurita* (L.) Hook. Gen. Fil. t. 65 A. 1841 ; Bedd. Ferns South. India t. 44. 1863 ; Handb. Ferns Brit. India 115. 1883.

Distrib. : Throughout the tropics of the world.

P. blumeana Agardh, Rec. Sp. Gen. Pteridis 22. 1839. *P. quadriaurita* var. *blumeana* Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 466. 1880 ; Bedd. Handb. Ferns Brit. India 112. 1883.

Distrib. : Eastern Himalayas, Bangladesh, Burma, Singapore.

P. confusa T. G. Walker in Kew Bull. 14 : 329. f. 5. t. 5 B. 1960.

Distrib. : India (Kerala), Sri Lanka.

P. cretica L. Mant. Pl. 130. 1767 ; Bedd. Ferns South. India t. 39. 1863 ; Handb. Ferns Brit. India 106. 1883 ; Hope, Journ. Bomb. Nat. Hist. Soc. 13 : 449. 1901.

Distrib. : Widely distributed in tropics & temperate regions of the world.

P. dactylina Hook. Sp. Fil. 2 : 160. t. 130 A. 1858 ; Bedd. Ferns Brit. India t. 23. 1865 ; Handb. Ferns Brit. India 107. t. 56. 1883 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 463. 1880.

Distrib. : India (E. Himalayas), China, Formosa.

P. ensiformis Burm. f. Fl. Ind. 230. 1768 ; Clarke, Trans Linn. Soc. Lond. II. Bot. 1 : 463. 1880 ; Bedd. Handb. Ferns Brit. India 107. 1883. *P. crenata* Sw. Schrad. Journ. Bot. 1800 (2) : 65. 1801; Bedd. Ferns South. India t. 35. 1863.

Distrib. : Throughout India, Sri Lanka, Burma, Malay Peninsula, China, Formosa, Ryukyu, Philippines, Malay Islands, tropical Australia, Polynesia, Micronesia.

P. excelsa Gaud. Frey. Voy. Bot. 388. 1829 ; Bedd. Ferns Brit. India 218. 1867 ; Handb. Ferns Brit. India 114. 1883.

Distrib. : N. W. to Eastern Himalayas, Sandwich Islands.

P. furunculata Nair & Ghosh, Journ. Indian Bot. Soc. 55 : 38. 1976.

Distrib. : India (Kerala).

P. geminata Wall. ex Agardh, Rec. Sp. Gen. Pteridis 31. 1839. *Campteris kleiniana* Presl, Tent. Pterid. 147. t. 5. f. 19. 1836; Bedd. Handb. Ferns Brit. India 116. t. 60. 1883. *C. anamallyensis* Bedd. Ferns South. India t. 45. 1863.

Distrib. : S. India, America.

P. gongalensis T. G. Walker, Kew Bull. 14 : 328. f. 4. t. 5 A. 1960 ; Nair & Ghosh, Journ. Bomb. Nat. Hist. Soc. 14 : 441. 1976.

Distrib. : India (Kerala), Sri Lanka.

P. grevilleana Wall. ex Agardh, Rec. Sp. Gen. Pteridis 23. 1839 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : t. 56. 1880 ; Bedd. Handb. Ferns Brit. India 112. 1883 ; Suppl. 24. 1892.

Distrib. : E. India, Malaya Peninsula, Borneo, Bangladesh, Burma, Singapore.

P. griffithii Hook. Sp. Fil. 2 : 170. t. 123A. 1853 ; Bedd. Ferns Brit. India t. 24 : 1869 ; Handb. Ferns Brit. India 108 t. 57. 1883.

Distrib. : India (S. Andamans), Burma.

P. heteromorpha Fée, Gen. Fil. 127. 1851 52 ; Bedd. Ferns Brit. India t. 89. 1865 ; Handb. Ferns. Brit. India 106. 1883.

Distrib. : India (Orissa), Burma, Luzon, Celebes.

P. khasiana (Clarke) Hieron. Hedw. 55 : 364. 1914 ; Bedd. Handb. Ferns Brit. India 111. 1883.

Distrib. : India (Himalayas).

P. linearis Poir. in Encycl. 5 : 723. 1804. *P. normalis* D. Don, Prod. Fl. Nepal 15. 1825.

Distrib. : E. Himalayas, trop. Africa, Asia, Australia, Polynesia.

P. longipcs D. Don, Prod. Fl. Nepal 15. 1825 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 468. 1880 ; Bedd. Handb. Ferns Brit. India 115. 1883.

Distrib. : India (S. E. India, E. Himalayas), Sri Lanka, Burma, Bhutan, Nepal, Guinea coast, Guatemala.

P. longipinnula Wall. ex Agardh, Rec. Sp. Gen. Pteridis 19. 1839 ;
Bedd. Ferns South. India t. 42. 1863 ; Handb. Ferns Brit. India 115. 1883.

Distrib. : Cosmopolitan.

P. mertensiooides Willd. in Linn. Sp. Pl. 5 : 394. 1810. **P. patens**
Hook. Sp. Fil. 2 : 177. 1858 ; Bedd. Ferns South. India t. 205. 1867.

Distrib. : S. India, Sri Lanka, Borneo, Philippines, and Society Islands.

P. multiaurita Ag. Rec. Sp. Gen. Pteridis 12. 1839 ; T. G. Walker,
Kew Bull. 14 : 323. 1960.

Distrib. : S. India.

P. nemoralis Willd. Enum, 1073. 1809. et in Linn. Sp. Pl. 5 : 386.
1810.

Distrib. : India (Orissa, Kerala).

P. nepalensis Ito in Hara's Fl. East. Himalayas 466. 1966.

Distrib. : E. Himalayas.

P. pellucida Presl, Rel. Haenk. 1 : 55. 1825 ; Clarke, Trans Linn.
Soc. Lond. II. Bot. 1 : 462. 1880 ; Bedd. Ferns South. India t. 38. 1863 ;
Handb. Ferns Brit. India 106. 1883.

Distrib. : Throughout India, Burma, Malay Islands, Guinea Coast.

P. perrotteti Hieron. Hedw. 55. 374. 1914.

Distrib. : South India (Nilgiri hills).

P. perrotteti var. **brevilaciniata** Hieron. Hedw. 55 : 375. 1914.

Distrib. : South India.

P. praetermissa T. G. Walker, Kew Bull. 14 : 327. f. 3. t. 5 E. 1960.

Distrib. : India (Kerala), Sri Lanka.

P. prainii Ghosh, J. Econ. & Tox. Bot. 3 : 591. t. 1. 1982.

Distrib. : India (Nagaland).

P. pseudopellucida Ching, Lingnan Sci. Journ. 15 : 395. 1936.

Distrib. : India (Eastern India), Bangladesh.

P. quadriaurita Retz. Obs. Bot. 6 : 38. 1791 ; Clarke Trans. Linn.
Soc. Lond. II. Bot. 1 : 465. 1880 (*pro parte*) ; Bedd. Ferns South. India
t. 31. 1863 ; Handb. Ferns Brit. India 110. 1883.

Distrib. : Throughout the tropics of world.

P. rigida Sod. Sert. Fl. Ecuad. II. 15. 1908. (non Sw. 1806)

Distrib. : Eastern Himalayas, Ecuadore.

P. roseo - lilacina Hieron. Hedw. 55 : 350. 1914.

Distrib. : E. Himalayas & S. India, China.

P. semipinnata L. Sp. Pl. 2 : 1076. 1753 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 464. 1880 ; Bedd. Ferns South. India t. 34. 1863 ; Handb. Ferns Brit. India 110. t. 58. 1883.

Distrib. : Throughout India in hilly regions, Burma, Malay Islands, China, Japan,

P. setigera (Bedd.) Nair, Bull. Bot. Surv. India 11 : 187. 1969.

P. quadriaurita Retz. var. *setigera* Bedd. Ferns Brit. India t. 202. 1866; Handb. Ferns Brit. India 111. 1883.

Distrib. : South India, Burma.

P. setuloso costulata Hayata, Icon. Pl. Formos. 4 : 241. f. 168. 1914.

Distrib. : India (E. Himalayas) Burma, Indo - China, Malay Islands, Formosa.

P. stenophylla Wall. ex Hook. et. Grev. Icon. Fil. t. 130. 1829.

P. pellucida var. *stenophylla* Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 463. 1880 ; Bedd. Handb. Ferns Brit. India 107. 1883.

Distrib. : India (North - West to Eastern Himalayas), Nepal.

P. subindivisa Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 467. t. 56. f. 1. 1880. **P. quadriaurita** var. *subindivisa* (Clarke) Bedd. Handb. Ferns Brit. India 112. 1883.

Distrib. : Eastern Himalayas, Bhutan.

P. subquinta (Wall. ex Bedd.) Agardh, Rec. Sp. Gen. Pteridis 21. 1839. **P. quadriaurita** var. *subquinta* Wall. ex Bedd. Handb. Ferns Brit. India Suppl. 23. 1892.

Distrib. : India (N. W. Himalayas to Eastern Himalayas), Nepal.

P. tenuissima Ching, Lingnan Sci. Journ. 15 : 394. 1936 ; Pic. Ser. Ind. Fil. Suppl. 4 : 262. 1965.

Distrib. : India (Himalayas).

P. tremula R. Br. Prod. Fl. Nov. Holl. 154. 1810 ; Nair & Ghosh, Journ. Bomb. Nat. Hist. Soc. 73 : 240. 1977.

Distrib. : S. India, Australia, New Zealand.

P. tricolor Linden, Gardn. Chron. 123. 1860.

Distrib. : India (Sikkim), Malacca.

P. tripartita Sw. Schrad. Journ. Bot. 1800 (2) : 67. 1801. *Litobrachia tripartita* (Sw.) Presl, Tent. Pterid. 150. 1836 ; Bedd. Ferns South. India t. 220. 1865 ; Handb. Ferns. Brit. India Suppl. 25. 1892. *L. marginata* Presl, Tent. Pterid. 149. 1836 ; Bedd. Handb. Ferns Brit. India 122. 1883.

Distrib. : S. India, Sri Lanka, Malacca. Malay Peninsula, Trop. Africa, Queensland, Philippines, Polynesia.

P. venusta Kunze, Bot. Zeit. 6 : 195. 1868.

Distrib. : India (E. Himalayas), Burma, Indo China, Malay Islands, Formosa.

P. vittata L. Sp. Pl. 2 : 1074. 1753. *Pteris longifolia* auctt. quoad. Pl. asiat ; Bedd. Ferns South. India t. 33. 1863 ; Handb. Ferns Brit. India 106. t. 55. 1883 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 461. 1880.

Distrib. : Tropics and subtropics of the world.

P. wallichiana Agardh, Rec. Sp. Gen. Pteridis 69. 1839. *Campteris wallichiana* Moore, Ind. Fil. 221. 1861 ; Bedd. Ferns Brit. India t. 25 & 217. 1865 ; Handb. Ferns Brit. India 118. 1883.

Distrib. : India (Himalayas), Nepal, Bhutan, Philippines, Java, Samoa.

ACROSTICHACEAE Mett. ex Frank.

Acrostichum L. Sp. Pl. 2 : 1067. 1753.

Lectotype : **A. aureum** L. Sp. Pl. 2 : 1068. 1753.

A. aureum L. Sp. Pl. 2 : 1069. 1753 ; Bedd. Ferns South. India t. 204. 1865 ; Handb. Ferns Brit. India 440. t. 268. 1883.

Distrib. : Tidal back water throughout India and throughout the world in Warm countries near the sea.

A. speciosum Willd. in Linn. Sp. Pl. 5 : 117. 1810.

Distrib. : India (Great Nicobar Islands) ; coasts of the world.

ADIANTACEAE (Presl) Ching

Adiantum L. Sp. Pl. 2 : 1094. 1753 ; Ching, Acta Phytotax. Sin. 6 : 301. 1957 ; Nayar, Bull. Nation. Bot. Gard, Lucknow no. 52 (1) : 1. 41. 1962.

Lectotype ; **A. capillus-veneris** L. Sp. Pl. 2 : 1096. 1753.

A. aethiopicum L. Syst. Nat. ed. X. 2 : 1329. 1759 ; Bedd. Ferns South. India t. 5. 1863 ; Ferns Brit. India t. 18. 1865 ; Handb. Ferns Brit. India 84. 1883.

Distrib. : India (Tamil Nadu), Sri Lanka, Australia, New Zealand, America, Africa, East Africa Islands.

A. assamicum (Nayar) Nayar, Bull. Nation. Bot Gard. Lucknow no. 54 : 1. 1964. *A. caudatum* L. var. *assamicum* Nayar, Bull. Nation Bot. Gard. Lucknow no. 52 (1) : 10. 1962.

Distrib. : Eastern India.

A. capillus - junonis Rupr. Distr. Cr. vasc. Ross. 49. 1845 ; Panigr. Bull. Bot. Surv. India 2 : 312. 1960.

Distrib. : Eastern India, China.

A. capillus - veneris L. Sp. Pl. 2 : 1096. 1753 : Bedd. Ferns South. India t. 4. 1863 ; Handb. Ferns Brit. India 84. 1883 ; Clarke. Trans. Linn. Soc. Lond. II. Bot. 1 : 453. 1880.

Distrib. : Throughout India in mountainous regions ; Sri Lanka, Europe, America, Australia.

A. capillus veneris L. var. *dissecta* Nayar, Bull. Nation. Bot. Gard. Lucknow. no. 52 (1) : 37. 1962.

Distrib. : India (Tamil Nadu).

A. caudatum L. Mant. 308. 1771; Clarke. Trans. Linn. Soc. Lond. II. Bot. 1 : 453. 1880. (*pro parte*).

Distrib. : Throughout India from the plains to lower hills, China, Sri Lanka, Malay Peninsula, Malesian Islands, Africa, Mauritius, Cape Verde Islands.

A. caudatum L. var. *flabellatum* Nayar, Bull. Nation. Bot. Gard. Lucknow no. 52 (1) : 12. f. 3 c. 1962.

Distrib. : India (Himalayan ranges at lower elevations from Kashmir to Nepal, Western and Eastern Ghats at higher elevations, Mahadev and Aravalli ranges).

A. cuneipinnulum Nair & Ghosh, Acta Bot. Indica 2 : 78. 1974.
A. cuneatum Langsd. et Fisch. in Icon. Fil. 23. t. 26. 1810 (*non* Forst, 1786).

Distrib. : India (Darjeeling, Tamil Nadu, Kerala), Brazil, Argentina, Peru.

A. edgeworthii Hook. Sp. Fil. 2 : 14. t. 81 B. 1851 ; Hope, Journ. Bomb. Nat. Hist. Soc. 13 : 239. 1900. *A. caudatum* L. var. *edgeworthii* Bedd. Handb. Ferns Brit. India 84. 1883.

Distrib. : India (North West Himalayas).

A. flabellulatum L. Sp. Pl. 2 : 1905. 1753 ; Bedd. Ferns South. India t. 218. 1865 ; Handb. Ferns Brit. India 88. 1883. *A. amoenum* Wall. ex Hook. & Grev. Icon. Fil. 103. 1829.

Distrib. : Eastern India, Nepal, Bangladesh, Malaya Peninsula, Malesian Islands, China, Japan.

A. hispidulum Sw. Schrad. Journ. 1800 (2) : 82. 1801 ; Bedd. Ferns South. India t. 3. 1863 ; Handb. Ferns. Brit. India 87. 1883.

Distrib. : India (Tamil Nadu), Sri Lanka, Australia, New Zealand, Fiji, Africa, and Eastern Islands.

A. incisum Forsk. Fl. Aeg. 187. 1775 ; Pic. Ser. Webbia 12 : 669. f. 6. 1957. *A. caudatum* auct. Bedd. Ferns South. India t. 2. 1863 ; Handb. Ferns Brit. India 83. t. 44. 1883.

Distrib. : Throughout India ; pantropic.

A. indicum Ghatak, Bull. Bot. Surv. India 5 : 71. 1963.

Distrib. : India (Bengal, Orissa).

A. pedatum L. Sp. Pl. 2 : 1095. 1753 ; Clarke Trans. Linn. Soc. Lond. II. Bot. 1 : 453. 1880 ; Bedd. Handb. Ferns. Brit. India 86. 1883.

Distrib. : India (North West Himalayas to Sikkim), Japan, North America.

A. pectinatum Ktze. ex Ettingsh. Farrnkr. 85. t. 45. 1865.

Distrib. : Widely cultivated in the gardens as pot plant.

A. peruvianum Kl. Linn. 20 : 441. 1847.

Distrib. : Widely cultivated in the gardens as pot plant.

A. philippense L. Sp. Pl. 2 : 1094. 1753. *A. lunulatum* Burm. Fl. Ind. 235. 1768 ; Bedd. Ferns South. India t. 1. 1863 ; Handb. Ferns Brit. India 82. 1883 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 452. 1880.

Distrib. : Pantropic.

A. polyphyllum Willd. in Linn. Sp. Pl. 5 : 454. 1810.

Distrib. : Cultivated in the gardens as pot plants.

A. raddianum Presl, Tent. Pterid. 158. 1836.

Distrib. : South India, S. America.

A. refractum Christ, Bull. Ac. Géogr. Bot. Mans. 1902. 224. 1902.

Distrib. : India (Sikkim), China.

A. rubellum Moore, Gard. Chron. 866. 1868.

Distrib. : Widely cultivated throughout India as pot plant.

A. soboliferum Wall. apud Hook. Sp. Fil. 2 : 13. t. 74 A. 1851.

Adiantum lunulatum Burm var. *mittenii* (Kuhn) Bedd. Handb. Ferns Brit. India 83. 1883. *A. incisum* var. *soboliferum* Bedd. Ferns Brit. India t. 19. 1865 ; Handb. Ferns Brit. India 84. 1883.

Distrib. : South India, Tropical Africa, Burma.

A. thalictroides Willd. ex Schl. Adumbr. t. 5 : 53. 1832. Nair & Ghosh, Journ. Bom. Nat. Hist. Soc. 75 : 246. 1978.

Distrib. : Peninsular India, Africa, Mexico, America, Australia.

A. trapeziforme L. Sp. Pl. 2 : 1097. 1753.

Distrib. : Widely cultivated throughout India as pot plant.

A. venustum D. Don, Prod. Fl. Nepal 17. 1825 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 453. 1880 ; Bedd. Handb. Ferns Brit. India 86. t. 45. 1883.

Distrib. : India (North East Himalayas), Afghanistan.

A. wattii (Bedd.) Bak. Journ. Linn. Soc. 18 : 381 t. 14. f. A. 1881.

A. capillus-veneris L. var. *wattii* (Bak.) Bedd. Handb. Ferns. Brit. India 18. 1892.

Distrib. : India, (N. E. India, Eastern Himalayas), Bangladesh.

A. zollingeri Mett. ex Kuhn, Ann. Mus. Lugd. Bat. 4 : 280. 1869 ; Ghatak, Bull. Bot. Surv. India 5 : 75. 1963.

Distrib. : South India, Java, Angola.

HEMIONITIDACEAE Pic. Ser.

Anogramma Link, Fil. sp. 137. 1841.

Lectotype : **A. leptophylla** (L.) Link, Fil. sp. 137. 1841.

A. leptophylla (L.) Link, Fil. sp. 137. 1841. *Polypodium leptophyllum* L. Sp. Pl. 2 : 1092. 1753. *Gymnogramma leptophylla* Desv. Berl. Mag. 5 : 305. 1811 ; Bedd. Ferns South. India t. 270. 1865 ; Handb. Ferns Brit. India 382. t. 220. 1883.

Distrib. : India (Western Ghats, S. India) Europe, Maderia, Canarie, S. Africa, Persia, Australia, New Zealand, S. America.

Coniogramme Féé, Gen. Fil. 167. t. 4. f. 1, 2. 1850 52 ; Dixit & Das, Proc. Indian Acad. Sci. 88 (4) : 253. 1979. *nom cons.*

Type : **C. fraxinea** (D. Don) Féé ex Diels in Engl. & Prantl. Nat. Pfl. 1 (4) : 262. 1899.

C. affinis Wall. ex Hieron. Hedw. 57 : 297. 1916. *Grammitis affinis* Wall. List. no. 11. 1828 (*nom. nud.*). *Gymnogramme javanica* sensu Hook. Sp. Fil. 5 : 145. 1864 (*pro parte*) sensu Bedd. Ferns Brit. India t. 57. 1866 (*non* Bl. 1828). *Syngramme fraxinea* (D. Don) Bedd. Handb. Ferns Brit. India 386. 1883 (*pro parte*) ; non Suppl. 100. 1892.

Distrib. : India (Himalayas), Nepal.

C. caudata (Wall. ex Ettingsch) Ching in C. Chr. Ind. Fil. Suppl. 3 : 56. 1934. *Grammitis caudata* Wall. ex Ettingsch, Farrnkr. 57. t. 37. f. 7. t. 38. 1865. *Gymnogramme fraxinea* var. *pilosa* sensu Clarke, Trans. Linn. Soc. II. Bot. 1 : 569. 1880.

Distrib. : India (N W Himalayas, Eastern India), China, Nepal.

C. denticulato - serrata (Hieron.) Dixit & Das, Proc. Indian Acad. Sci. 88 (4) ; 261. f. 7. 20 : 1979. *C. fraxinea* var. *denticulo serrata* Hieron. Hedw. 57 : 290. 1916.

Distrib. : India (North - West Himalayas and Eastern India).

C. falcata (D. Don) Salom., Nomencl. 139. 1888. *Diplazium falcatum* D. Don, Prod. Fl. Nepal 13. 1825.

Distrib. : India (Eastern & Central Himalayas), Nepal.

C. faxinea (D. Don) Féé ex Diels in Engl. & Prantl, Nat. Pfl. 1 (4) : 262. 1899. *Diplazium fraxineum* D. Don, Prod. Fl. Nepal 12. 1825. *Gymnogramme fraxinea* (D. Don) Bedd. Ferns Brit. India Suppl. 24. 1876 (*pro parte*) ; *sensu* Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 569. 1880

(*pro parte*). *Syngramme fraxinea* (D. Don), Bedd. Ferns Brit. India 386. 1883 (*pro parte*) ; Suppl. 100. 1892. *Gymnogramme javanica* Bl. Enum. Pl. Jav. 112. 1828 ; *non sensu* Bedd. Ferns South. India 77 t. 232. 1865 ; Ferns Brit. India t. 57. 1866.

Distrib. : Throughout mountainous regions in India, China, Malay Peninsula, Indo China, Malesia, Formosa, Philippines.

C. indica Féé 10 mém 22. 1865.

Distrib. : Eastern India.

C. intermedia Hieron. Hedw. 57 : 310. 1916. var. **intermedia**. *Conio - gramme intermedia* Hieron. var. *villosa* Ching, Icon. Fil. Sin. 3. t. 143. 1935. *Syngramme fraxinea* (D. Don) Bedd. Handb. Ferns Brit. India 386. 1883 (*pro parte*).

Distrib. : India (Himalayas), Nepal, Bhutan, China, Japan.

C. intermedia Hieron. var. **glabra** Ching, Icon Fil. Sin. 3 : t. 143. 1935.

Distrib. : India (Himalayas), Nepal, Bhutan, China, Indo China, Formosa, Japan.

C. pitelottii Tard. Blot. Bull. Mus. Paris II. 5 : 334. 1933.

Distrib. : India (Meghalaya), China, Tonkin.

C. procera Wall. ex Féé, 10 me Mém. 22. 1865. *Grammitis procera* Wall. List. no. 3. 1828 (*nom. nud.*). *Syngramme fraxinea* (D. Don) Bedd. Handb. Ferns Brit. India 386. 1883 (*pro parte*).

Distrib. : India (Himalayas), N. Thailand, China, Formosa.

C. purpurea Dixit & Das, Proc. Indian Acad. Sci. 88 (4) : 264. f. 11, 25 27. 1979.

Distrib. : India (Eastern Himalayas).

C. serrulata (Bl.) Féé. Gen, Fil. 167. t. 14 B. f. 2. 1850 52. *Gymno - gramme serrulatum* Bl. Enum. Pl. Jav. 113. 1828. (*non* Féé 1857.)

Distrib. : Eastern India, Java.

C. subcordata Copel. Leaflets Phil. Bot. 3 : 823. 1910.

Distrib. : India (Himalayas), Nepal, Philippines.

Hemionitis L. Sp. Pl. 2 : 1077. 1753.

Lectotype : **H. palmata** L. Sp. Pl. 2 : 1077. 1753.

Hemionitis arifolia (Burm.) Moore, Ind. Fil. 114. 1859 ; Bedd. Handb. Ferns Brit. India 413. t. 245. 1883. *Asplenium arifolium* Burm. Fl. Ind. 231. 1768. *Hemionitis cornifolia* Roxb. ex Bedd. Ferns South. India t. 53. 1863.

Distrib. : South India, Bangladesh, Sri Lanka, Burma, Philippines.

Idiogramma Ghosh, J. Econ. Tax. Bot. 4 (3) : 981. 1983.

Type : **I. microphylla** (Hook.) Ghosh, J. Econ. Tax. Bot. 4 (3) : 982. 1983.

I. microphylla (Hook.) Ghosh. J. Econ. Tax. Bot. 4 (3) : 982. 1983. *Gymnogramme microphylla* Hook. Icon. Pl. 10 : t. 916. 1854 ; Bedd. Handb. Ferns Brit. India 384. t. 221. 1887. *Anogramma microphylla* Diels, Nat. Pflanz. 1 (4) : 259. 1899 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 569. 1880. *Grammitis microphylla* Bedd. Ferns Brit. India t. 148. 1866.

Distrib. : India (Eastern India, Eastern Himalayas), China.

Pityrogramma Link, Handb. Erken. Gew. 3 : 19. 1833 ; Nair & Ghosh, Jonrn. Indian Bot. Soc. 54 : 103. 1975 ; Panigr. Kew Bull. 31 : 657. 1976.

Type : **P. chrysophylla** (Sw.) Link, Handb. Erken. Gew. 3 : 19. 1833.

P. austroamericana Domin, Spisy Prirod. Fak. Karlovy Univ. 88 : 7. 1928 et Kew Bull. Misc. inf. Kew 1929 ; 221. 1929.

Distrib. : S. India, S. America, introduced and naturalised in the parts of the old world.

P. calomelanos (L.) Link, Handb. Erken. Gew. 3 : 20. 1833. *Acrostichum calomelanos* L. Sp. Pl. 2 : 1072. 1753. *Pteris calomelanos* (L.) Bedd. Ferns Brit. India t. 22. 1863. *Pellaea calomelanos* (L.) Link, Fil. Sp. 61. 1841. Bedd. Handb. Ferns Brit. India 104. 1883.

Distrib. : Throughout India, Mexico & West Indies, South to Argentina, introduced and naturalised in the parts of the old world.

P. chrysophylla (Sw.) Link, Handb. Erken. Gew. 3 : 19. 1833. *Acrostichum chrysophyllum* Sw. Schrad. Journ. Bot. 1800 (2) : 12. 1801.

Distrib. : S. India, Trop. America ; introduced to parts of Asia (Malaya, Sri Lanka, India).

Syngramme J. Sm. Lond. Journ. Bot. 4 : 168. 1845.

Lectotype : **S. alsimaeifolia** J. Sm. Lond. Journ. Bot. 4 : 168. t. 7 8 f. 5. 1845.

S. vestita Wall. ex Bedd. Ferns Brit. India t. 154. 1867 ; Handb. Ferns Brit. India 386. t. 223. 1883. *Grammitis vestita* Wall. List no. 12. 1828. (*nom. nud.*)

Distrib. : India (Himalayas), Nepal, China.

ANTROPHYACEAE (Link) Ching

Antrophyum Kaulf. Fnum. Fil. 197. 1824 ; Dixit & Nair, Journ. Indian Bot. Soc. 53 : 277. 1974.

Type : **A. reticulatum** (Forst.) Kaulf. Enum. Fil. 197. 1824.

A. callifolium Bl. Enum. Pl. Jav. 111. 1828.

Distrib. : India (South Andamans), Sri Lanka, Malaysia to Polynesia.

A. coriaceum (D. Don) Wall. ex Moore, Ind. Fil. 5 : 80. 1858 ; Bedd. Handb. Ferns Brit. India Suppl. 103. 1892. *Hemionitis coriacea* D. Don, Prod. Fl. Nepal 13. 1825. *Antrophyum reticulatum* sensu Bedd. Handb. Ferns Brit. India 401. 1883 (*pro parte*).

Distrib. : Eastern India and Eastern Himalayas, Nepal, Bhutan, China, Malay Peninsula.

A. henryi Hieron. Hedwigia 57 : 208. 1915 ; C. Chr. Asiatic Pterid. 313. 1931.

Distrib. : India (Arunachal Pradesh, Sikkim), China (Yunnan.)

A. obovatum Bak. Kew. Bull. Misc. Inf. 1898 : 238. 1898 ; *A. latifolium* sensu quoad Bedd. Handb. Ferns Brit. India 404. 1883 ; Clarke, Trans Linn. Soc. Lond. II. Bot. 1 : 573. 1880 (*non* Bl.).

Distrib. : India (Himalayas), Indo China, Japan, China

A. parvulum Bl. Pl. Jav. 110. 1828. *A. reticulatum* var. *parvula* Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 573. 1880 ; Bedd. Handb. Ferns Brit. India 403. 1883 (as *parvulum*). *A. obtusum* quoad Bl. Enum. Pl. Jav. 110. t. 34. f. 4. 1828.

Distrib. : India (Sikkim, Meghalaya), Malay Peninsula, Java.

A. plantagineum (Cav.) Kaulf. Enum. 197. 1824 ; Bedd. Handb. Ferns Brit. India 403. 1883. *Hemionitis plantaginea* Cav. Descr. 260. 1802. *Antrophyum reticulatum* sensu Bedd. Ferns South. India t. 52. 1865.

Distrib. : India (Assam, Meghalaya, Tamil Nadu, Kerala), Sri Lanka, Malay Peninsula, Malesian Islands, Philippines, Polynesian Islands.

A. reticulatum (Forst.) Kaulf. Enum. 198. 1824 ; Bedd. Handb. Ferns. Brit. India Suppl. 102. 1892. *Hemionitis reticulata* Forst. Prod. 79. 1876. *A. reticulatum* sensu Bedd. Handb. Ferns Brit. India 401. t. 235. 1883 (*pro parte*).

Distrib. : India (Assam, Meghalaya, West Bengal, Darjeeling, Sikkim, Manipur, Kerala, Tamil Nadu), Sri Lanka, Malay Peninsula, Polynesia, Queensland.

VITTARIACEAE (Presl) Ching

Vittaria J. E. Smith. Mém. Acad. Turin. 5 : 413. t. 9. 1793 ; Bir. Res. Bull. Punjab Univ. (N.S.) 13 : 15. 1962 ; Dixit. Journ. Econ. & Tax. Bot. 2 : 209. 1981.

Type : **V. lineata** (L.) Smith, Mém. Acad. Turin. 5 : 413. t. 9. f. 5. 1793.

V. amboinensis Féé, 3 Mém. 14. t. 1. f. 1. 1851 1852 ; *Taeniopsis amboinensis* (Féé), Bedd. Ferns Brit. India t. 117. 1866. *V. scolopendrina* sensu Bedd. Handb. Ferns Brit. India 408. 1883. (*pro parte*).

Distrib. : India (Eastern Himalayas), Bhutan, Vietnam, Thailand, Burma, Indonesia.

V. arunachalensis Dixit, Journ. Econ. & Tax. Bot. 3 : 959. t. 1. f. 1. 2. 1982.

Distrib. : India (Arunachal Pradesh).

V. carcina Christ, Bull. Géogr. Bot. Mans. 109. 1904.

Distrib. : India (Sikkim, Manipur, West Bengal), China.

V. doniana Hieron. Hedw. 57 : 204. 1916. var. **doniana**.

Distrib. : India (Arunachal Pradesh, Manipur, Sikkim), Philippines, Taiwan.

V. doniana Hieron. var. **intermedia** Hieron. Hedw. 57 : 204. 1916.

Distrib. : India (Eastern Himalayas).

V. elongata Sw. Syn. Fil. 109, 302. 1806. *V. elongata* sensu Bedd. Handb. Ferns Brit. India 404. 1883 (*pro parte*). *V. elogata* Sw. var. *angustifolia* sensu Panigr. Bull. Bot. Surv. India 2 : 314. 1960 (*non* Holtt. 1954). *Pteris graminifolia* Roxb. ex Griff. Calc. Journ. Sci. 4 : 502. t. 33. 1844.

Distrib. : Throughout India in mountainous regions ; Sri Lanka, Bangladesh, Burma, Vietnam, Laos, Malaysia, Taiwan, Philippines, China, Polynesia, Australia, Africa.

V. flexuosa Féé, 3 Mém. Foug. 16. 1851 1852 ; Clarke Trans. Linn. Soc. Lond. II. Bot. 1 : 575. 1880. *V. lineata* sensu Bak. in Hook. Syn. Fil. 396. 1874 (*pro parte*) ; Bedd. Handb. Ferns Brit. India 407. 1883. *Taeniopsis lineata* Bedd. Ferns South. India t. 54. 1863.

Distrib. : India (Arunachal Pradesh, Meghalaya, Assam, West Bengal, Uttar Pradesh), Sri Lanka, Nepal, Bhutan, Thailand.

V. forrestiana Ching, Sinensis 1 (12) : 191. 1931.

Distrib. : India (Assam, Manipur, Meghalaya), Burma, China, Vietnam, Laos, Khmer.

V. garhwaleensis Dixit, J. Econ. & Tax. Bot. 2 : 218. f. 18. 1981.

Distrib. : India (North West Himalayas).

V. himalayensis Ching, Sinensis 1 (12) : 190. 1931.

Distrib. : India (North West Himalayas, Meghalaya, West Bengal).

V. linearifolia Ching, Sinensis 1 (12) : 183. t. 1. f. 1 3. 1931.

Distrib. : India (Arunachal Pradesh), Burma, China.

V. microlepis Hieron. Hedw. 57 : 202. 1916 ; Manickam & Ninan, New Botanist (Suppl.) 1 : 10. 1976.

Distrib. : India (Kerala, Tamil Nadu), Sri Lanka.

V. ophiopogonoides Ching, Sinensis 1 (12) : 186. 1931.

Distrib. : India (Himalayas), Nepal, Bhutan, China.

V. sikkimensis Kuhn, Linnaea 36 : 66. 1869 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 574. t. 84. f. 3. 1880 ; Bedd. Handb. Ferns. Brit. India 406. t. 239. 1883. *V. minor* Bedd. Ferns Brit. India t. 58. 1865 (*non* Féé. 1851 52). *V. minor* var *minima* Hook. Sp. Fil. 5 : 183. 1864. *V. modesta* sensu H. Ito in Hara's Fl. East. Himal. 500. 1966.

Distrib. : India (Arunachal Pradesh, Assam, Darjeeling & Sikkim Himalayas), China.

V. taeniophylla Copel. Philip. Journ. Sci. Bot. 1. Suppl. 2 : 157. 1956. *V. mediosora* sensu Ito in Haras Fl. East. Himal. 500. 1966 (*non* Hayata, 1915).

Distrib. : India (Himalayas), Nepal, Burma, Thailand, China (Yunnan), Philippines, Taiwan.

V. wattii Dixit & Nair, New Botanist 5 : 41. f. 1 5. 1978.

Distrib. : India (Arunachal Pradesh, Nagaland, Manipur).

V. zosterifolia Willd. in Linn. Sp. Pl. 5 : 406. 1810. *V. lineata* *sensu* Bedd. Handb. Ferns Brit. India 407. 1883 (*pro parte*) : Hope, Journ. Bombay Nat. Hist. Soc. 14 : 103. 1903 (*pro parte*).

Distrib. : India (Eastern India and Eastern Himalayas, Andaman & Nicobar Islands), China.

TAENITIDACEAE (Presl) Pic. Ser.

Taenitis Willd. ex Schkuhr. krypt. Gew. 1 : 21. 1805.

Type : **T. blechnoides** (Willd.) Sw. Syn. Fil. 24, 220. 1806.

T. blechnoides (Willd.) Sw. Syn. Fil. 24, 220. 1806 ; Bedd. Ferns Brit. India t. 54. 1865 ; Handb. Ferns Brit. India 410. t. 242. 1883. *Pteris blechnoides* Willd. Phyt. 13. t. 9. f. 3. 1794.

Distrib. : South India, Sri Lanka, South East Asia.

PARKERIACEAE Hook.

Ceratopteris Ad. Brongn. Bull. Sci. Soc. Philom. Paris 1821 : 186. 1822 ; Lloyd, Britt. 26 (3) : 156. 1974.

Type : **C. thalictroides** (L.) Ad. Brongn. Bull. Sci. Soc. Philom. Paris 1821 : 186. 1822.

C. cornuta (P. Beauv.) Le Prieur, Ann. Sci. Nat. 19 : 103. t. 4A. 1830. *Pteris cornuta* P. Beauv. Fl. d'. Oware 1 : 63. t. 37 (2). 1806.

Distrib. : India (Assam), Bangladesh, Burma, Africa from Senyal to Sudan and Souto to Tanjania, Madagascar, Socotra, Saudia Arabia, Iraq, Indonesia, N. Australia.

C. pteridoides (Hook.) Hieron. in Engl. Jahrb. 84 : 561. 1905 (*pteroides*). *Parkeria pteridoides* Hook. Exot. Fl. 2 : 145. 1925.

Distrib. : India (West Bengal), South America from Amazon river basin to Peru and North to Columbia and Venezuela ; Central America, Brazil, Argentina, Paraguay, North Vietnam.

C. thalictroides (L.) Ad. Brongn. Bull. Sci. Soc. Philom. Paris 1821 : 186. 1822 ; Bedd. Ferns South. India t. 75. 1863 ; Handb. Ferns Brit. India 123. t. 83. 1883. *Acrostichum thalictroides* L. Sp. Pl. 2 : 1070. 1753.

Distrib. : Tropics and subtropics of the world.

MARSILEACEAE Mirb.

Marsilea L. Sp. Pl. 2 : 1099. 1753 ; Gupta, Marsilea Mon. 26. 1962.

Lectotype : *M. quadrifolia* L. Sp. Pl. 2 : 1099. 1753.

M. aegyptiaca Willd. in Linn. Sp. Pl. 5 : 540. 1810.

Distrib. : India (Orissa, Rajasthan), Egypt.

M. brachycarpa A. Br. Monatsber. Ak. Berl. 1863 : 420. 1864.

Distrib. : South India, Burma.

M. brachypus A. Br. Monatsber. Ak. Berl. 1863 : 421. 1864.

Distrib. : India (Nilgiris, Amritsar), Australia.

M. condensata Bak. Journ. Bot. 1886 : 281. 1886.

Distrib. : India (Rajasthan).

M. coromandelica Burm. f. Fl. Ind. t. 62. f. 3. 1768.

Distrib. : India (Coromandel Coast).

M. crenata Presl, Rel. Haenk. 1 : 84. t. 12. f. 13. 1825.

Distrib. : India (Himalayas), Malayasia.

M. gracilenta A. Br. Monatsber. Ak. Berl. 1863 : 421. 1864.

Distrib. : India (Concan, Kemnangudi).

M. maheshwarii Gupta, Amer. Fern Journ. 58 : 70. 1968.

Distrib. : India (Pondicherry).

M. major (Haines) Chowdhury, Res. Living Pterid. 49. 1971 ;
M. minuta L. var. *major* Haines, Botany Bihar & Orissa 3 : 1272. 1924.

Distrib. : India (Bihar).

M. minuta L. Mant. 308. 1771.

Distrib. : Throughout India, Java, Philippines.

M. poonensis Kolhatkar, Kew Bull. 1957 : 293. f. 1. 1957.

Distrib. : India (Poona).

M. quadrifolia L. Sp. Pl. 2 : 1099. 1753.

Distrib. : India (Kashmir), Europe, Australia, China, Japan, in temperate regions.

M. rajasthaniensis Gupta, Marsilea Bot. Mon. 29. 1962.

Distrib. : India (Rajasthan).

M. rajasthanensis Gupta var. **ballardii** (Gupta) Gupta, Marsilea Bot. Mon. 30. 1962. *M. ballardii* Gupta, Journ. Bomb. Nat. Hist. Soc. 53 : 289. 1955.

Distrib. : India (Rajasthan).

HYMENOPHYLLACEAE Link

Gonocormus v.d. Bosch, Verslag, Meded, Kon. Akad. Wetensch, Afd. Natur. 21 : 321. 1861.

Lectotype : **G. prolifer** (Bl.) Prantl, Hymen. Jav. 11 : t. 5. 1861.

G. diffusus (Bl.) v.d. Bosch, Hymen. Jav. 9 : t. 4. 1859. *Trichomanes diffusum* Bl. Enum. Pl. Jav. 225. 1828.

Distrib. : Himalayas to Java.

G. prolifer (Bl.) Prantl, Hymen. Jav. 11 : t. 6. 1861. *Trichomanes proliferum* Bl. Enum. Pl. Jav. 224. 1828 ; Bedd. Ferns South. India t. 262. 1865 ; Handb. Ferns Brit. India 39. 1883.

Distrib. : S. India, Sri Lanka, Java, Philippine Islands.

G. saxifragoides (Presl) v.d. Bosch. Hymen. Jav. 9. 1861. *Trichomanes saxifragoides* Presl, Hymen. 19. 39. 1843. *Trichomanes parvulum* Poir, Encycl. 8 : 64. 1808 ; Bedd. Ferns South. India t. 179. 1865.

Distrib. : S. India, Japan, China, Java, Polynesian Islands, Madagascar, Johanna Islands and the Moluccas.

Hymenophyllum Sm. Mém. Acad. Sci. Turin. 5 : 418. t. 9. f. 8. 1793.

Lectotype : *H. tunbrigense* (L.) Sm. in Sowerby Engl. Bot. t. 162. 1794.

H. barbatum (v. d. Bosch) Bak. Syn. Fil. 68. 1868. *Leptocionium* v. d. Bosch, Ned. kr. Arch. 5 (3) : 146. 1863.

Distrib. : India (Assam), Korea, Japan, Taiwan, China.

H. exiguum Bedd. Ferns Brit. India t. 275. 1868. *Trichomanes exiguum* (Bedd.) Bak. Syn. Fil. 464. 1874 ; Bedd. Handb. Ferns Brit. India 37. 1883.

Distrib. : South India, Sri Lanka.

H. simonsianum Hook. 2nd Cent. t. 13. 1860 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 438. 1880 ; Bedd. Ferns Brit. India t. 281. 1968. Handb. Ferns Brit. India 34. 1880.

Distrib. : Himalayas to Formosa.

Mecodium Presl ex Copel. Philip. Journ. Sci. 67 : 17. 1938.

Type : **M. sanguinolentum** (G. Forst.) Presl ex Copel. Philip. Journ. Sci. 67 : 17. 1938.

M. badium (Hook. et Grev.) Copel. Philip. Journ. Sci. 67 : 23. 1938. *Hymenophyllum badium* Hook. et Grev. Icon. Fil. t. 76. 1828. *H. javanicum* var. *badium* (Hook. et Grev.) Bedd. Ferns Brit. India t. 282. 1868 ; Handb. Ferns Brit. India 33. 1883.

Distrib. : India (Sikkim), Burma, Malay Peninsula.

M. crispatum (Wall. ex Hook. & Grev.) Copel. Philip. Journ. Sci. 67 : 23. 1938. *Hymenophyllum crispatum* Wall. ex Hook. & Grev. Icon. t. 77. 1828. *H. javanicum* *sensu* Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 437. 1880 ; Bedd. Handb. Ferns Brit. India 32. 1883. (*pro parte*).

Distrib. : Eastern Himalayas.

M. exsertum (Wall. ex Hook.) Copel. Philip. Journ. Sci. 67 : 23. 1938. *Hymenophyllum exsertum* Wall. list no. 170. 1828 (*nom. nud.*) ex Hook. Sp. Fil. 1 : 109. t. 38A. 1844 ; Bedd. Ferns South. India t. 9. 1863 ; Handb. Ferns Brit. India 30. t. 16. 1883.

Distrib. : Throughout India in mountainous regions, Burma, Sri Lanka.

M. javanicum (Spr.) Copel. Philip. Journ. Sci. 67 : 20. 1938. *Hymenophyllum javanicum* Spr. Syst. 4 : 132. 1827 ; Bedd. Handb. Ferns Brit. India 32. 1883. *H. australe* Wild. in Linn. Sp. Pl. 5 : 527. 1810.

H. crispatum Wall. List no. 169. 1828 ex Bedd. Ferns South. India t. 207. 1865.

Distrib. : Throughout India in mountainous regions ; Sri Lanka, Burma, Malay Peninsula, Australia, New Zealand, Mauritius, Bourbon, Philippines.

M. levingei (Clarke) Copel. Philip. Journ. Sci. 75 : 359. 1941.
Hymenophyllum levingei Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 439. t. 49. f. 3. 1880 ; Bedd. Handb. Ferns Brit. India 36. 1883.

Distrib. : India (Sikkim), Bhutan.

M. polyanthos (Sw.) Copel. Philip. Journ. Sci. 67 : 19. 1938.
Hymenophyllum polyanthos Sw. Schrad. Journ. Bot. 1800 (2) : 102. 1801 ;
 Bedd. Handb. Ferns Brit. India 31. 1883. *H. polyanthos* var. *minor* Bedd.
 Ferns Brit. India t. 306. 1869. *H. blumeanum* Spr. Syst. Veg. 4 : 131.
 1827 ; Bedd. Ferns South. India t. 266. 1865. *Mecodium blumeanum*
 (Spr.) Nayar et Kaur, Comp. to Bedd. Handb. 11. 1974.

Distrib. : Throughout India in mountainous regions ; Burma, Sri Lanka, Tropical and subtropical regions of the world.

M. polyanthos (Sw.) Copel. var. *blumeana* (Bedd.) Nair, Bull. Bot. Surv. India 11 : 186. 1969. *Hymenophyllum polyanthos* var. *blumeanum* Bedd. Ferns Brit. India t. 308. 1869 ; Handb. Ferns Brit. India 32. 1883.

Distrib. : S. India.

M. polyanthos (Sw.) Copel. var. *microglossum* (Bedd.) Nair, Bull. Bot. Surv. India 11 : 186. 1969. *Hymenophyllum polyanthos* var. *microglossum* Bedd. Ferns Brit. India t. 306. 1869 ; Handb. Ferns Brit. India 32. 1883.

Distrib. : S. India.

M. undulatum (Sw.) Copel. Philip. Journ. Sci. 67 : 26. 1938. *Hymenophyllum undulata* Sw. Schrad. Journ. 1800 (2) : 101. 1801. *H. dentriculatum* var. *flaccidum* Bedd. Handb. Ferns Brit. India 35. 1883.

Distrib. : India (Meghalaya), Burma.

Meringium Presl, Hymen. 19. t. 6. f. A, B. 1843.

Type : **M. meyenianum** Presl, Hymen. 24. t. 8. f. A, B. 1843.

M. acanthoides (v. d. Bosch) Copel. Philip. Journ. Sci. 67 : 42. 1938 ; Ghosh. J. Econ. Tax. Bot. 3 (2) : 578. 1982. *Didymoglossum acanthoides* v. d. Bosch. Plant Jungh. 1 : 16. 1856.

Distrib. : India (Eastern Himalayas), Malesia.

M. edentulum (v.d. Bosch) Copel. Philip. Journ. Sci. 67 : 41. 1938.
Leptocionum edentula v.d. Bosch, Ned. Kr. Arch. 5 (3) : 148. 1863.

Distrib. : India (Assam), Malesia.

M. flaccidum (v. d. Bosch) Nair, Bull. Bot. Surv. India 11 : 186. 1969. *Leptocionum flaccidum* v. d. Bosch. Ned. Kr. Arch. 5 : 149. 1863. *Hymenophyllum khasianum* Bak. in Hook. & Bak. Syn. Fil. 464. 1874 ; Bedd. Ferns Brit. India t. 276. 1868. *H. denticulatum* Sw. var. *flaccidum* Bedd. Handb. Ferns Brit. India 35. 1883.

Distrib. : India (Meghalaya).

M. denticulatum (Sw.) Copel. Philip. Journ. Sci. 67 : 26. 1938. *Hymenophyllum denticulatum* Sw. Schrad. Journ. Bot. 1800 (2) : 100. 1801 ; Bedd. Ferns Brit. India t. 278. 1868 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 438. 1880 ; Bedd. Handb. Ferns Brit. India 34. 1883. *H. neesii* Hook. Sp. Fil. 1 : 99. 1844 ; Bedd. Ferns South. India t. 279. 1865.

Distrib. : India (Meghalaya), Bhutan, Burma, Jawa.

Microgonium Presl, Hymen. 19. t. 6. f. A, B. 1843.

Type : **M. cuspidatum** (Willd.) Presl, Hymen. 19. t. 6. f. A. B. 1843.

M. bimarginatum v. d. Bosch, Hymen. Jav. 7. 1861. *Trichomanes neilgherense* Bedd. Ferns South. India t. 6. 1863 ; Handb. Ferns Brit. India 37. 1883.

Distrib. : S. India, Sri Lanka.

M. motleyi v.d. Bosch, Hymen. Jav. 5. t. 1. 1861. *Trichomanes motleyi* (v.d. Bosch) v.d. Bosch, Ned. Kr. Arch. 5 (2) : 145. 1861 ; Bedd. Handb. Fcrns. Brit. India 36. 1883. *T. henzae* Bedd. Ferns Brit. India t. 183. 1866.

Distrib. : India (Andamans) Sri Lanka, Burma, Borneo, New Caledonia.

M. sublimbatum (C. Muell.) v.d. Bosch, Hymen. Jav. 6. t. 2. 1861. *Trichomanes sublimbatum* C. Muell. Bot. Zeit. 12 : 737. 1854.

Distrib. : India (Assam), Indo China, Malaysia to Papua.

Microtrichomanes (Mett.) Copel. Philip. Journ. Sci. 67 : 35. 1938.

Type : **M. digitatum** (Sw.) Copel. Philip. Journ. Sci. 67 : 35. 1938.

M. nitidulum (v.d. Bosch) Copel. Philip. Journ. Sci. 67 : 37. 1938.

Trichomanes nitidulum v.d. Bosch, Pl. Jungh. 1 : 547. 1856. *T. corticola* Bedd. Ferns South. India. t. 264. 1865.

Distrib. : North India, Sri Lanka, Malay Peninsula, Mauritius, Bourbon, Java, Polynesian Islands.

Pleuromanes (Presl) Presl, Epim. Bot. 258. 1851.

Type *P. acutum* (Presl) Presl, Epim. Bot. 258. 1851.

P. pallidum (Bl.) Presl, Epim. Bot. 258. 1849. *Trichomanes pallidum* Bl. Enum. Pl. Jav. 225. 1828.

Distrib. : India (Eastern Himalayas), Sri Lanka, Sumatra to Polynesia.

Selenodesmium (Prantl) Copel. Philip. Journ. Sci. 67 : 80. 1938.
Selenodesmium Prantl, Hymen. 53. 1875. as section of *Trichomanes*.

Lectotype : *S. rigidum* (Sw.) Copel. Philip. Journ. Sci. 67 : 80. 1938.

S. rigidum (Sw.) Copel. Philip. Journ. Sci. 67 : 80. 1938.
Trichomanes rigidum Sw. Prod. 137. 1788 ; Bedd. Ferns South. India t. 8. 1863 ; Handb. Ferns Brit. India 46. 1883.

Distrib. : S. India, Sri Lanka, Malaya Peninsula, Tropical America, West Indies, Polynesian Islands, New Zealand, Japan, S. China, S. Africa, Angola, Fernando, Poland, the Mascareen Islands.

Sphaerocionum Presl, Hymen. 33. t. 4. f. 10 B, C. 1843.

Lectotype : *S. hirtum* (L.) Presl, Hymen. 33. t. 4. f. 10. B, C. 1843.

S. ciliatum (Sw.) Presl, Hymen. 34. 1843 ; Nayar and Kaur. Comp. to Bedd. Handb. 11. 1974. *Hymenophyllum ciliatum* Sw. Schrad. Journ. 1800 (2) : 100. 1801 ; Bedd. Ferns Brit. India t. 305. 1869 ; Handb. Ferns Brit. India 33. 1883.

Distrib. : India (Sikkim,) Tropical America, Tropical Africa, Mauritius, Madagascar, Bourbon, Seychelles, Johannan Islands, New Zealand.

Vandenboschia Copel. Philip. Journ. Sci. 67 : 51. 1938.

Type : *V. radicans* (Sw.) Copel. Philip. Journ. Sci. 67 : 51. 1938.

V. auriculata (Bl.) Copel. Philip. Journ. Sci. 67 : 55. 1938. *Trichomanes auriculatum* Bl. Jav. 225. 1828 ; Bedd. Ferns Brit. India 182. 1867 ; Handb. Ferns Brit. India 44. 1883. *Trichomanes dissectum* J. Sm. ex Hook. Sp. Fil. 1 : 140. 1844 ; Bedd. Ferns Brit. India t. 182. 1866.

Distrib. : India (Himalayas), Japan, Formosa, Philippines, Java, Guiana.

V. gigantea (Bory ex Willd.) Pic. Ser. Webbia 29 (1) : 197. 1968.
Trichomanes giganteum Bory ex Willd. in Linn. Sp. Pl. 5 : 514. 1810.

Distrib. : India (Sikkim, Arunachal Pradesh), Maritius.

V. latifrons (v.d. Bosch) Copel. Philip. Journ. Sci. 67 : 56. 104. 1938. *Trichomanes latifrons* v.d. Bosch, Ned. Kr. Arch. 5 (3) : 56. 1859. *T. pyxidiferum* L. Sp. Pl. 2 : 1098. 1753 ; Bedd. Handb. Ferns India 42. 1883. (*pro parte*). *T. pyxidiferum* var. *limbatum* Bedd. Ferns Brit. India Suppl. t. 348. 1876 ; Handb. Ferns Brit. India 43. 1883.

Distrib. : India (Himalayas), Burma, Bangladesh, Sri Lanka, Tropics of the world.

V. maxima (Bl.) Copel. Philip. Journ. Sci. 67 : 54. 1938. *Trichomanes maxima* Bl. Enum. Pl. Jav. 228. 1828.

Distrib. : India (Andaman Islands), Sumatra to Tahiti.

V. orientalis (C. Chr.) Ching, Acta Phytotax. Sinica 8 : 187. 1959. *Trichomanes radicans* var. *orientalis* (C. Chr.) Lelling. Amer. Fern Journ. 58 : 157. 1968. *T. radicans* var. *anceps* Wall. ex Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 441. 1880.

Distrib. : Eastern India and Eastern Himalayas, China.

V. radicans (Sw.) Copel. Philip. Journ. Sci. 67 : 51. 54. 1938 ; *Trichomanes radicans* Sw. Schrad. Journ. 1800 (2) : 97. 1801 ; Bedd. Ferns Brit. India t. 181. 1867 ; Handb. Ferns Brit. India 43. 1883. *T. bimarginatum* Bedd. Ferns Brit. India Suppl. t. 349. 1876 ; Handb. Ferns Brit. India 43. 1883. *T. birmanicum* Bedd. Ferns Brit. India Suppl. 3. 1876.

Distrib. : Himalayas, Nepal to Bhutan, Burma and temperate regions of both hemispheres.

V. schmidiana (Zenker ex Taschner) Copel. Philip. Journ. Sci. 67 : 53. 1938. *Trichomanes Schmidianum* Zenkerex Taschner, Dissert. 34. t. 1. f. 1. 1843.

Disirib. : S. India.

V. titibuensis Ito, Journ. Jap. Bot. 24 : 125. f. 1. 1949.

Distrib. : India, (Himalayas), Japan.

HYMENOPHYLLOPSIDACEAE Picc. - Ser.

Cephalomanes Presl, Hymen. 17. t. 5. 1843.

Type : **C. atrovirens** Presl, Hymen. 17. t. 5. 1843.

C. javanicum (Bl.) v.d. Bosch, Hymen. Jav. 30. 1861. *Trichomanes javanicum* Bl. Enum. Pl. Jav. 224. 1828 ; Bedd. Ferns Brit. India t. 180. 1867 ; Handb. Ferns Brit. India 44. t. 21. 1883.

Distrib. : Eastern India, Bangladesh, Burma, Malay Peninsula, Borneo, Java, Polynesian Islands, Madagascar.

Crepidomanes (Presl) Presl, Epim. Bot. 258. 1849.

Type : **C. intramarginale** (Hook. et Grev.) Presl, Epim. Bot. 258. 1849.

C. bilabiatum (Nees ex Bl.) Copel. Philip. Journ. Sci. 67 : 59. 1938. *Trichomanes bilabiatum* Nees ex Bl. Nova Acta 11 : 123. t. 13. 1823.

Distrib. : India (Nicobar Islands), Sri Lanka, Malesia.

C. euphlebium (v.d. Bosch) Dixit & Ghosh, *comb. nov.* *Didymoglossum euphlebium* v.d. Bosch, Ned. Kr. Arch. 5 (3) : 142. 1863. *Trichomanes euphlebium* (v.d. Bosch) Panigr. Phytologia 31 (3) : 256. 1975.

Distrib. : India (Sikkim), Bhutan.

C. griffithii (v.d. Bosch) Dixit & Ghosh, *comb. nov.* *Didymoglossum griffithii* v.d. Bosch in Ned. Kr. Arch. 5 (3) : 141. 1863. *Trichomanes griffithii* (v.d. Bosch) Panigr. Phytologia 31 (3) : 256. 1975.

Distrib. : India (Andaman Islands), Burma.

C. insigne (v.d. Bosch) Fu, III. Handb. Chin, Pl. Pterid. 39. 1957. *Didymoglossum insigne* v.d. Bosch, Ned. Kr. Arch. 5 (3) : 143. 1863. *Trichomanes insigne* (v.d. Bosch) Bedd. Ferns Brit. India t. 284. 1868. *T. bipunctatum* var. *insigne* (v.d. Bosch) Bedd. Ferns Brit. India t. 284. 1868 ; Handb. Ferns Brit. India 42. 1883.

Distrib. : Throughout the tropics of the world.

C. latealatum (v.d. Bosch) Copel. Philip. Journ. Sci. 67 : 60. 1938. *Trichomanes late-alatum* v.d. Bosch, Ned. Kr. Arch. 5 (3) : 138. 1863. *T. bipunctatum* Poir, Encycl. 8 : 69. 1808 ; Bedd. Handb. Ferns Brit. India 41. 1883.

Distrib. : India (Himalayas), Nepal, Bhutan.

C. latemarginale (D. C. Eaton) Copel. Philip. Journ. Sci. 67 : 60. 1938. *Trichomanes latemarginale* D. C. Eaton, Proc. Amer. Acad. 4 : 111. 1858. *T. kurzii* Bedd. Ferns Brit. India t. 286. 1868 ; Handb. Ferns Brit. India 40 t. 20. 1883. *T. intramarginale* Hook. & Grev. Icon. Fil. t. 211. 1831 ; Bedd. Ferns South. India t. 208. 1865 ; Handb. Ferns Brit. India 41. 1883.

Distrib. : South India. Sri Lanka.

C. racemulosum (v.d. Bosch) Ching in Chein et Chun, Fl. Republ. Pop. Sin. 2 : 170. 1959.

Distrib. : India (Assam), China.

DICKSONIACEAE Hook. ex Bower

Cibotium Kaulf. Berlin Jaharb. Pharm. 21 : 53. 1820.

Type : **C. chamissoi** Kaulf. Enum. Fil. 230. t. 1. f. 14. 1824.

C. assamicum Hook. Sp. Fil. 1 : 83. t. 29 B. 1844. *Cibotium barometz* *sensu* Bedd. Handb. Ferns Brit. India 24. 1883 (*pro parte*).

Distrib. : India (Assam), Tonkin.

C. barometz (L.) J. Sm. Lond. Journ. Bot. 1 : 437. 1842 ; Bedd. Handb. Ferns Brit. India 24. 1883 (*pro parte*). *Polypodium barometz* L. Sp. Pl. 2 : 1092. 1753.

Distrib. : Eastern India, Burma, Malay Islands, China, Australia, Malesia.

CYATHEACEAE Kaulf.

Cyathea Sm. Mém. Acad. Turin 5 : 416. 1793 ; Holtt. Kew Bull. 19 : 480. 1965.

Lectotype ; **C. arborea** (L.) Sm. Mém. Acad. Turin. 5 : 417. 1793.

C. albosetacea (Bedd.) Copel. Philip. Journ. Sci. ser. C. (Bot.) 4 : 55. 1909. *Alsophila albosetacea* Bedd. Ferns Brit. India Suppl. 2. 1876 ; Handb. Ferns Brit. India 16. 1883.

Distrib. : India (Nicobar Islands).

C. andersonii (J. Scott ex Bedd.) Copel. Journ. Sci. ser. C. (Bot.) 4 : 56. 1909. *Alsophila andersonii* J. Scott ex Bedd. Ferns Brit. India t. 310. 1869 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 433. 1880 ; Bedd. Handb. Ferns Brit. India 12. 1883.

Distrib. : India (Meghalaya, Assam, Darjeeling & Sikkim Himalayas), Bhutan, China.

C. brunoniana (Wall. ex Hook.) Clarke & Bak. in Journ. Linn. Soc. Bot. 24 : 409. 1888 ; *nomen tantum, descr. et specim. omn. excl.* *Alsophila brunoniana* Wall. ex Hook. Sp. Fil. 1 : 52. 1844 ; Bedd. Ferns Brit. India t. 86. 1865. *A. sollyana* Griff. Notulae 2 : 624. 1849. *A. contaminans* var. *brunoniana* J. Scott in Trans. Linn. Soc. 30 : 35. t. 15. 1874. *A. glauca* sensu Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 432. 1880 ; *pro parte* Bedd. Handb. Ferns Brit. India 12. 1883 ; *pro parte* (non (Bl.) J. Sm. 1841. *Hemitelia brunoniana* (Wall. ex Hook.) Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 430. 1880. *Amphicosmia brunoniana* Bedd. Handb. Ferns Brit. India 10. 1883.

Distrib. : India (Western Himalayas, Eastern India), Bangladesh, Burma, Vietnam.

C. chinensis Copel. Philip. Journ. Sci. Bot. 3 : 355. 1909. *Hemitelia brunoniana* sensu Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 430. 1880. incl. var. *scottii*, non *Alsophila brunoniana* Hook. Sp. Fil. 1 : 52. 1844. *Amphicosmia brunoniana* sensu Bedd. Handb. Ferns Brit. India 10. 1883, (non Hook.) *Cyathea brunoniana* sensu Clarke et Bak. Journ. Linn. Soc. Bot. 24 : 409. 1888.

Distrib. : India (Eastern Himalayas, Meghalaya) Bhutan, China, Burma, Vietnam.

C. contaminans (Wall. ex Hook.) Copel. Philip. Journ. Sci. ser. C. Bot. 4 : 60. 1909. *Alsophila contaminans* Wall. ex Hook. Sp. Fil. 1 : 52. t. 18. f. 2. 1844 ; Bedd. Ferns Brit. India t. 85. 1865. *A. glauca* (Bl.) J. Sm. Journ. Bot. 3 : 419. 1841 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 432. (*pro parte*) 1880 ; Bedd. Handb. Ferns Brit. India 12. 1883 (*pro parte*).

Distrib. : India (Sikkim), Malaya, Thailand.

C. crinita (Hook.) Copel. Philip. Journ. Sci. ser. C. Bot. 4 : 40. 1909. *Alsophila crinita* Hook. Icon. Pl. 7. t. 671. 1844 ; Bedd. Ferns. South. India t. 59. 1863 ; Handb. Ferns Brit. India 16. 1883.

Distrib. : S. India, Sri Lanka.

C. gigantea (Wall. ex Hook.) Holtt. Gard. Bull. Str. Settl. 8 : 318. 1935. *Alsophila gigantea* Wall. ex Hook. Sp. Fil. 1 : 53. 1844. *A. glabra* sensu Bedd. Ferns South. India t. 60. 1883 ; Handb. Ferns Brit. India 14. 1883 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 432. 1880.

Distrib. : Throughout India in hilly regions, Nepal, China, Burma, Thailand, Laos, Vietnam, Sri Lanka, Bangladesh.

C. henryi (Bak.) Copel. Philip. Journ. Sci. ser. C. Bot. 4 : 38. 1909. *Alsophila henryi* Bak. in Bull. Misc. Inf. Kew 1898. 229. 1898.

Distrib. : India (Sikkim), China.

C. khasyana (Moore ex Kuhn) Domin, Acta Bot. Bohem. 362. 1930.
Alsophila khasyana Moore ex Kuhn, Linnaea 36 : 154. 1869. *A. ornata* Scott ex Bedd. Ferns Brit. India t. 342. 1870 ; Handb. Ferns Brit. India 12. 1883 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 432. 1880. *A. oldhami* Bedd. Ferns Brit. India t. 343. 1870 ; Handb. Ferns Brit. India 13. 1883 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1. 433. 1880 ; *A. sikkimensis* Clarke & Bak. in Journ. Linn. Soc. Bot. 24 : 409. 1888 ; not *A. ornata* var. *sikkimensis* Bedd. in Journ. Bot. 51 : 225. 1893, nor *A. latebrosa* var. *ornata* Bedd. in Bull. Misc. inf. Kew 1909 : 423. 1909.

Distrib. : India (Eastern India, Eastern Himalayas), Burma.

C. nilgirensis Holtt. Kew. Bull. 19 : 468. 1965. *Alsophila latebrosa* Hook. var. *schmidiana* Kunze, Linnaea 24 : 294. 1851. *A. latebrosa* Hook. Sp. Fil. 1 : 37. 1844, *pro parte*, quoad Bedd. Handb. Ferns Brit. India 11. 1883.

Distrib. : South India.

C. spinulosa Wall. ex Hook. Sp. Fil. 1 : 25. t. 12. 1844 ; Bedd. Ferns South. India t. 57. 1863 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 430. 1880 ; Bedd, Handb. Ferns Brit. India 6. 1883 ; Suppl. 2. 1892. *Alsophila decipiens* Scott ex Bedd. Ferns Brit. India t. 311. 1869. *Hemitelia decipiens* (Scott) Scott, Trans. Linn. Soc. 30 : 33. t. 14. 1874 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 430. 1880. *Amphicosmia decipiens* (Scott) Bedd. Ferns Brit. India Suppl. 1. 1876 ; Handb. Ferns. Brit. India 10. 1883 ; Suppl. 2. 1892. *Hemitelia beddomei* Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 429. 1880. *Cyathea decipiens* (Scott) Clarke & Bak. Journ. Linn. Soc. Bot. 24 : 409. 1888.

Distrib. : Throught India in hilly regions, Nepal, Bhutan, China, Burma, Thailand, Formosa, Japan.

DENNSTAEDTIACEAE Pic. Ser.

Dennstaedtia Bernh. Schrad. Journ. Bot. 1880 (2) : 124. 1801.

Type : **D. flaccida** (Forst.) Bernh. Schrad. Journ. Bot. 1800 (2) : 124. t. 1. f. 3. 1801.

D. scabra (Wall. ex Hook.) Moore, Ind. Fil. 307. 1961 ; Bedd. Handb. Ferns Brit. India 24. t. 12. 1883. *Dicksonia scabra* Wall. ex Hook. Sp. Fil. 1 : 80. t. 28 B. 1844 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 436. 1880.

Distrib. : India (Punjab to Assam), Sri Lanka, Burma, Indo - China, China, Korea, Formosa, Japan, Philippines, Malay Peninsula, Borneo, Celebes.

Emodiopteris Ching et Wu, *Acta Phytotax. Sin.* 16 (4) : 21. 1978.

Type : **E. appendiculata** (Wall. ex Hook.) Ching et Wu, *Acta Phytotax. Sin.* 16 (4) : 21. 1978.

E. appendiculata (Wall. ex Hook.) Ching et Wu, *Acta Phytotax. sin.* 16 (4) : 21. 1978. *Dicksonia appendiculata* Wall. ex Hook. Sp. Fil. 1 : 79. t. 27 C. 1844 ; Clarke, *Trans. Linn. Soc. Lond. II. Bot.* 1 : 436. 1880. *Dennstaedtia appendiculata* (Wall. ex Hook.) J. Sm. *Hist. Fil.* 265. 1875 ; Bedd. *Handb. Ferns Brit. India* 26. 1883. *Patania appendiculata* (Wall. ex Hook.) Bedd. *Ferns Brit. India Suppl.* 2. 1876.

Distrib. : India (Himalayas), China.

E. appendiculata var. *elwesii* (Bedd.) Dixit, *comb. nov.* *Dennstaedtia elwesii* Bedd. *Ferns Brit. India Suppl.* t. 347. 1876 ; *Handb. Ferns Brit. India* 26. 1883. *D. appendiculata* var. *elwesii* (Bedd.) Bedd. *Handb. Ferns Brit. India Suppl.* 7. 1892.

Distrib. : India (Sikkim).

Histiopteris (Agardh) J. Sm. *Hist. Fil.* 294. 1875.

Type : **H. incisa** (Thunb.) J. Sm. *Hist. Fil.* 295. 1875.

H. incisa (Thunb.) J. Sm. *Hist. Fil.* 295. 1875. *Pteris incisa* Thunb. *Prod. Fl. Cap.* 171. 1800 ; Clarke, *Trans. Linn. Soc. Lond. II. Bot.* 1 : 471. 1880. *Litobrachia incisa* Presl, *Tent. Pterid.* 149. 1836 ; Bedd. *Handb. Ferns Brit. India* 120. t. 62. 1883. *L. aurita* Bedd. *Ferns South. India* t. 221. 1865.

Distrib. : Throughout India in mountainous regions, tropics and subtropics of the world.

Microlepia Presl, *Tent. Pterid.* 124. 1836 ; Nayar & Kaur, *Bull. Nation. Bot. Gard. Lucknow* no. 79. 12. 1963.

Lectotype : **M. polypodioides** (Sw.) Presl, *Tent. Pterid.* 125. 1825.

M. calvescens var. *intramarginalis* (Tagawa) Shieh in *Quart. Journ. China For.* 6 (4) : 92. 1973. **M. strigosa** (Thunb.) Presl var. *intramarginalis* Tagawa, *Acta Phytotax. Geobot.* 10 : 202. 1941.

Distrib. : India (Meghalaya), Taiwan.

M. haflangensis Nayar & Kaur, Bull. Nation. Bot. Gard. Lucknow no. 94. 8. f. 2-3. 1964.

Distrib. : Eastern India.

M. hancei Prantl, Arb. Bot. Gart. Breslau 1 : 35. 1892. *Davallia polypodioides* Benth. Fl. Hongk. 461. 1961.

Distrib. : India (Meghalaya), Hongkong.

M. khasiyana (Hook.) Presl, Epim. Bot. 95. 1849. *Davallia khasiyana* Hook. Sp. Fil. 1 : 173. t. 47 A. 1856.

Distrib. : India (Himalayas).

M. kurzii (Clarke) Bedd. Handb. Ferns Brit. India 66. 1883. *Davallia kurzii* Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 446. 1880.

Distrib. : Eastern India, Burma.

M. marginata (Houtt.) C. Chr. Ind. Fil. 212. 1905 ; 427. 1906. *polypodium marginatum* Houtt. Pfl. Syst. 12. 199. 1786. *Microlepia marginalis* Bedd. Ferns Brit. India 64. 1883.

Distrib. : India (North West Himalayas, Eastern India), Japan Formosa.

M. marginata var. **calvescens** (Hook.) Nair, Bull. Bot. Surv. India 11. 186. 1969. *Davallia calvescens* Hook. Sp. Fil. 1 : 172. t. S8B. 1846. *M. marginalis* var. **calvescens** (Hook.) Bedd. Ferns Brit. India 64. 1883.

Distrib. : India (N.W.) Himalayas.

M. pilosula Wall. ex Presl, Tent. Pterid. 125. 1836, 1964. *Davallia polypodioides* var. **pilosula** Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 448. 1880.

Distrib. : Tropics of Asia.

M. platyphylla (D. Don) J. Sm. Lond. Journ. Bot. 1 : 427. 1842 ; Bedd. Ferns South. India t. 13. 1863 ; Handb. Ferns Brit. India 66. t. 33. 1883 ; *Davallia platyphylla* D. Don, Prod. Fl. Nepal 10. 1825.

Distrib. : India (Himalayas, South India), Nepal, Bhutan, Sri Lanka.

M. speluncae (L.) Moore, Ind. Fil. 93. 1857 ; Bedd. Handb. Ferns Brit. India 67. 1883. *Polypodium speluncae* L. Sp. Pl. 2 : 1093. 1753.

Distrib. : India (Himalayas, South India), Bangladesh, Sri Lanka, Malay Peninsula, Malaya Islands, Polynesia, Tropical America.

M. strigosa (Thunb.) Presl, Epim. Bot. 95. 1849 ; Bedd. Ferns.

South. India t. 255. 1865 ; Handb. Ferns Brit. India 67. 1883. *Trichomanes strigosa* Thunb. Fl. Jap. 339. 1784. *Dicksonia strigosa* (Thunb.) Clarke, Trans. Linn. Soc. 2 : 341. 1794. *Davallia polypodioides* D. Don var. *strigosa* Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 448. 1880.

Distrib. : India (Himalayas, South India), Sri Lanka, Malay Peninsula, Malay Islands, Polynesia.

M. todayensis Christ, Philip. Journ. Sci. 3 : 272. 1908.

Distrib. : India (Himalayas), Minadano.

M. trapeziformis (Roxb. ex Griff.) Kuhn, Chaetpot. 347. 1882. *Davallia trapeziformis* Roxb. ex Griff. Calc. Journ. Sci. 4 : 516. 1844. *Microlepia speluncae* var. *rhomboidea* Bedd. Handb. Ferns Brit. India 68. 1883. *M. polypodioides* Bedd. Ferns South. India t. 15. 1863.

Distrib. : Tropics of Asia.

MONACHOSORACEAE Ching ex Ching

Monachosorum Kunze, Bot. Zeit. 6 : 119. 1848.

Type : **M. davallodes** Kunze, Bot. Zeit. 6 : 119. 1848.

M. subdigitatum (Bl.) Kuhn, Chaetopt. 345. 1882. *Aspidium subdigitatum* Bl. Enum. Pl. Jav. 171. 1828. *Phegopteris subdigitata* (Bl.) Bedd. Handb. Ferns Brit. India 295. 1883.

Distrib. : India (Himalayas), Nepal, Bhutan, Malay Peninsula, Malay Islands.

HYPOLEPIDACEAE Pic. Ser.

Hypolepis Bernh. Schrad. Neu. Journ. Bot. 1 (2) : 34. 1805.

Type : **H. tenuifolia** (G. Forst.) Benth. Schrad. Neu. Journ. Bot. 1 (2) : 34. 1805.

H. punctata (Thunb.) Mett. ex Kuhn, Fil Afr. 120. 1868 (*non* Bedd. 1892). *Polypodium punctatum* Thunb. Fl. Jap. 337. 1784. *Phegopteris punctata* Mett. Ann. Lugd. Bat. 1 : 222. 1864. *Dryopteris punctata* C. Chr. Ind. Fil. 287. 1905 (*excl. syn. Hypolepis punctata* Bedd.).

Distrib. : Throughout India in mountainous regions, Sri Lanka, Japan, Malay Peninsula.

PTERIDIACEAE Ching

Pteridium Gled. ex Scop, Fl. Carn. 169. 1760. *nom. cons.*

Type : **P. aquilinum** (L.) Kuhn, v. Deck. Reis. 3 (3) : 11. 1879.

P. aquilinum (L.) Kuhn v. Deck. Reis. 3 (3) : 11. 1879. *Pteris aquilina* L. Sp. Pl. 2 : 1075. 1753 ; Bedd. Ferns South. India t. 44. 1863 ; Handb. Ferns. Brit. India 115. 1883.

Distrib. : Throughout India in mountainous regions, Sri Lanka, Malay Peninsula, throughout the world except Arctic zones and temperate South America.

P. acqulinum var. **wightianum** (Wall.) Trayon, Rhodora 43 : 22. 1941. *Pieris Wightiana* Wall. List. no. 2178. 1829. *P. aquilina sensu* Clarke, Trans Linn. Soc. Lond. II. Bot. 1 : 468. 1880.

Distrib. : Eastern Himalayas, Sri Lanka, Indo China, Malay Peninsula, Malesian Islands, Philippines, Formosa, Nepal.

P. capense (Thunb.) Krasser var. **densum** Tagawa, Bull. Nat. Sci. Mus. no. 27 : 5. 1949.

Distrib. : Eastern Himalayas.

LINDSAEACEAE Pic. Ser.

Lindsaea Dryand. ex Smith. Mém. Acad. Turin 5 : 401. 1793 ; Dixit & Ghosh, Proc. Indian Acad. Sci. (Pl. Sci.) 92 : 233. f. 1-47. 1983.

Lectotype : **L. guianensis** (Aubl.) Dryand. Trans. Linn. Soc. 3 : 42. 1797.

L. andamanica Dixit & Ghosh, Proc. Indian Acad. Sci. (Pl. Sci.) 92 : 254. f. 40, 41. 1983.

Distrib. : India (South Andaman Islands).

L. beddomea Dixit & Ghosh, Proc. Indian Acad. Sci. (Pl. Sci.) 89 (3) : 181. f. 4-7. 1980. *Lindsaea heterophylla* Bedd. Ferns South. India 70. t. 206. 1863 (*non* Dryand. 1797) ; Handb. Ferns Brit. India 77. 1883.

Distrib. : India (South India).

L. bouillodii Christ, Not. Syst. 1 : 59. 1909. *L. tenera* auct. Bedd.

Ferns South. India 7 f. 24. 1865. *L. orbiculata* var. *tenera* Bedd. Handb. Ferns Brit. India 73. 1883.

Distrib. : India (Tamil Nadu), Sri Lanka, Thailand, Tonkin, Malay Peninsula, Malesian Islands, Indo China.

L. ensifolia Sw. Schrad. Journ. Bot. 1800 (2) : 77. 1801. *Schizoloma ensifolia* (Sw.) J. Smith in Hook. Journ. Bot. 3 : 414. 1841 ; Bedd. Ferns South. India fl 25. 1865 ; Handb. Ferns Brit. India 80. t. 41. 1883.

Distrib. : Throughout India, Bangladesh, Burma, Sri Lanka, China, Malay Peninsula, Malesian Islands, New Guinea, Australia, Africa, New Caledonia, Hawaii.

L. glandulifera V. Ald. V. Ros. Bull. Jard. Bot. Btzg. 2 (1) : 9. 1911. *L. repens* (Bory.) Thw. var. *minor* (Thw.) Bedd. Ferns South. India 72. t. 214. 1865 ; Handb. Ferns Brit. India 74. 1883.

Distrib. : India (Assam, Sikkim), Sri Lanka, Malesia.

L. gueriniana (Gaud.) Desv. Prod. 312. 1827 ; *Schizoloma gueriniana* Gaud. Ann. Sci. Nat. 3 : 508. 1824.

Distrib. : India (Nicobar Islands), Philippines, Malesian Islands, Celebes, Moluccas, New Guinea, Solomon Islands, Tahiti.

L. heterophylla Dryand. Trans. Linn. Soc. 3 : 41. t. 8. f. 1. 1797. *Adiantum heterophyllum* (Dryand.) Poir. Encycl. Suppl. 1. 139. 1810. *Schizoloma heterophyllum* (Dryand.) J. Sm. in Hook. Journ. Bot. 3 : 414. 1841 ; Bedd. Ferns South. India 9. f. 25. 1863 ; Handb. Ferns Brit. India 80. 1883. *S. savantwadiensis* Almeida, Journ. Bomb. Nat. Hist. Soc. 74 : 325. f. 4. 1977. *S. indica* Almeida, Journ. Bomb. Nat. Hist. Soc. 74 : 325. f. 3. 1977.

Distrib. : India (South India, Maharashtra, Karnataka, Nicobar Islands), Mascareen Islands, Madagascar, Sri Lanka, Ryu Kyu Islands, China, Malay Peninsula, Malesian Islands, Philippine Islands.

L. himalaica Kram. Gard. Bull. Singapore 26 (1) : 43. f. 3. 1972. *L. cultrata* (Willd.) Sw. var. *minor* Hook. Sp. Fil. 1 : 204. 1846 (*pro parte* excl. Lectotype). *L. cultrata* (Willd.) Sw. var. *assamica* Hook. loc. cit. ex. char. (no material cited).

Distrib. : India (Assam, Meghalaya, Arunachal Pradesh).

L. javanensis Bl. Enum. Pl. Jav. 219. 1828. *Schizoloma javanense* (Bl.) Holtt. Rev. Fl. Malaya 2 : 349. f. 202. 1954. *Lindsaea flabellulata* Dryand. var. *gigantea* Hook. Sp. Fil. 1. 211. f. 630. 1846. *L. flabellulata* sensu Bedd. Ferns South. India 72. f. 216. 1865 (descr. *pro parte*).

Distrib. : India (Meghalaya, Assam), Burma, China, Japan, Malay Peninsula, Malesian Islands, Philippines.

L. lucida Bl. Enum. Pl. Jav. 216. 1828. *L. lobiana* Hook. Sp. Fil. 1 : 205. 1846 ; Bedd. Ferns Brit. India t. 28. 1865. *L. cultrata sensu* Bedd. Ferns South. India 7. 1865 (descr. *pro parte*, t. 23) ; Handb. Ferns Brit. India 72. 1883 (descr. *pro parte*. excl. f. 36).

Distrib. : India (Eastern India, Andaman Islands) Burma, Bhutan, Bangladesh, China, Japan, Thailand, Malay Peninsula, Malesian Islands, Philippines, New Guinea.

L. malabarica (Bedd.) Bak. ex C. Chr. Contr. U. S. Nat. Herb. 26 : 195. 1931. *Schizoloma malabaricum* Bedd. Ferns Brit. India t. 268. 1868. with descr. *S. lobata* (Poir.) Bedd. var. *malabaricum* (Bedd.) Bedd. Handb. Ferns Brit. India 79 t. 39. 1883 (err. *malabarica*).

Distrib. : India (South India, Madhya Pradesh).

L. malayensis Holtt. Gard. Bull. S. S. 5 : 69. f. 8. 1930.

Distrib. : India (Nicobar Islands). Malay Peninsula, Malesia.

L. oblanceolata v. A. v. R. Bull, Jard. Bot. Btzg. II. 23 : 15. 1916. *Lindsaea pectinata* auct. Holtt. Rev. Fl. Malaya 2 : 324. 1954 (*non* Bl. 1828).

Distrib. : India (Nicobar Islands), Malay Peninsula, Malesian Islands, Annam, Peninsular Thailand, Philippines, Moluccas.

L. odorata Roxb. ex Griff. Calc. Journ. Nat. Hist. 4 : 511. 1844. *L. cultrata* (Willd.) Sw. var. *minor* Hook. Sp. Fil. 1 : 204. 1846 (*pro parte* excl. Lectotype). *L. cultrata* (Willd.) Sw. var. *pallens* Hook. Sp. Fil. 1 : 204. 1846. *L. cultrata* (Willd.) Sw. var. *attenuata* Hook. Sp. Fil. 1 : 204. 1846. *L. cultrata* auct. viz. Bedd. Ferns South. India 7. t. 23. 1864 ; Handb. Ferns. Brit. India 72 t. 36. 1883.

Distrib. : India (Eastern India), Burma, Bangladesh, Sri Lanka, Malaya Peninsula, Malesian Islands, New Guinea, Moluccas, Philippines.

L. odorata var. *darjeelingensis* Sen & Sen. Amer. Fern Journ. 71 : 14. f. 1 8. 1971.

Disirib. : India (Darjeeling Himalayas).

L. orbiculata (Lam.) Mett. ex Kuhn. Miq. Ann. Mus. Bot. Lugd. Bat. 4 : 279. 1869 ; Bedd. Handb. Ferns Brit. India 75. 1883. var. *orbiculata*. *Adiantum orbiculatum* Lam. Encycl. 1 : 41. 1783. *L. flabellulata* Dryand. Trans. Linn. Soc. 3 : 41. t. 8. f. 2. 1797 ; Bedd. Ferns South. India 72. 1865 (descr. *pro parte*, excl. t. 216).

Distrib. : India, Burma, China, Sri Lanka, Indo China, Japan, Thailand, Malay Peninsula, Malesian Islands, Philippines.

L. orbiculata (Lam.) Mett. ex Kuhn var. **commixta** (Tagawa) Kram. Fl. Mal. ser. II. 1 (3) : 207. 1971. *L. commixta* Tagawa, Acta phytotax. Geobot. 6 : 37. f. 3. H. J. 1937.

Distrib. : India (N. W. Himalayas), Burma, Sri Lanka, Japan, Malay Peninsula, Malesian Island, Philippines.

L. parasitica (Roxb. ex Griff.) Hieron. Hedw. 14 : 62. 1920. *Vittaria parasitica* Roxb. ex Griff. Calc. Journ. Nat. Hist. 4 : 510. 1844. *Lindsaea scandens* Hook. Sp. Fil. 1 : 205. t. 63 B. 1846 ; Bedd. Ferns Brit. India t. 298. 1868 ; Handb. Ferns Brit. India 74. t. 37. 1883.

Distrib. : India (Nicobar Islands), Malay Peninsula, Malesian Islands, Thailand.

L. repens (Bory) Thw. var. **pectinata** (Bl.) Mett. ex Kuhn, Miq. Ann. Mus. Lugd. Bat. 4 : 277. 1868. *L. pectinata* Bl. Enum. Pl. Jav. 217. 1828 (*non* Holtt. 1954). *L. repens* sensu Bedd. Ferns South. India t. 209. 1865 ; Handb. Ferns Brit. India 74. 1883.

Distrib. : India (Sikkim, West Bengal, Assam, Madhya Pradesh) Sri Lanka, Malay Peninsula, Malay Islands, Philippines, Thailand, Indo China.

L. rutlandia Dixit & Ghosh, Proc. Indian Acad. Sci. (Pl. Sci.) 92 : 244. f. 11 12. 1983.

Distrib. : India (South Andamans).

L. tenera Dryand. Trans, Linn. Soc. 3 : 42. t. 10. 1797 (*non* Kaulf. 1834, not of later authors). *Schizoloma tenerum* (Dryand.) Holtt. Rev. Fl. Malaya 2 : 348. 1954 (as to the type only).

Distrib. : India (Nicobar Islands).

L. tetragona Karam. Blumea 15 (2) : 564. 1967 ; Ghosh & Dixit, Bull. Bot. Surv. India 20 (1-4) : 170. f. 3-4. 1978.

Distrib. : India (Nicobar Islands). Minanda, Celebes, Moluccas, Louisiades, Solomon Islands, Fiji, Tahiti.

L. venusta Kaulf. ex Kuhn, Linnaea 36 : 79. 1869. *Schizoloma recurvatum* auct. (*non* Wall. ex Hook.) Bedd. Ferns South. India 9. t. 27. 1864. *Schizoloma lobata* sensu Bedd. Handb. Ferns. Brit. India 79. 1883 (*pro parte*).

Distrib. : India (South India), Sri Lanka.

Sphenomeris Maxon, Journ. Wash. Acad. Sc. 3 : 144. 1913.

Type : **S. clavata** (L.) Maxon, Journ. Wash. Acad. Sc. 3 : 144. 1913.

S. chinensis (L.) Maxon, Journ. Wash. Acad. Sc. 3 : 144. 1913.
Trichomanes chinensis L. Sp. Pl. 2 : 1099. 1753. *Davallia chinensis* Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 449. 1880. *D. tenuifolia* Sw. Schrad. Journ. Bot. 1800 (2) : 88. 1801 ; Bedd. Ferns South. India t. 16. 1863. *Stenoloma chinensis* (L.) Bedd. Handb. Ferns Brit. India 70. t. 34. 1883. *Trichomanes malayanum* Roxb. ex. Griff. Calc. Journ. Nat. Hist. 4 : 519. 1844.

Distrib. : Throughout India from the plains to hilly regions, Sri Lanka, Malay Peninsula, China, Japan, Polynesia, East African Islands.

S. chinensis var. **divaricata** (Christ) Kramer, Blumea 15 : 672. 1968.
Odontosoria chinensis var. *divaricata* Christ, Journ. de. Bot. ser. II. 2 : 23. 1909.

Distrib. : India (Assam, Sikkim), China to Malesia.

Tapeinidium (Presl) C. Chr. Ind. Fil. 631. 1906. *Tapeinidium* Presl, Epim. Bot. 96. 1849. as sec. of *Microlepia*.

Type : **T. pinnatum** (Cav.) C. Chr. Ind. Fil. 631. 1906.

T. pinnatum (Cav.) C. Chr. Ind. Fil. 631. 1906. *Davallia pinnata* Cav. Descr. 277. 1802. (*non* Mett. ex Kuhn 1869). *Microlepia pinnata* (Cav.) J. Sm. ex Hook. Journ. Bot. 3 : 416. 1841 ; Bedd. Ferns South. India t. 14. 1863 ; Handb. Ferns Brit. India 64. 1883. *Davallia serrata* Roxb. ex Griff. Calc. Journ. Nat. Hist. 4 : 514. 1844 (*non* Willd. 1810).

Distrib. : South India, Eastern Himalayas, Thailand, Ryu Kyu Island, Taiwan, Malesia, Singapore, Lingga Island, Banka, Sumatra, Java. Borneo, Celebes, Philippines.

THELYPTERIDACEAE (Presl) Pic. Ser.

Ampelopteris Kunze, Bot. Zeit. 6 : 114. 1848.

Type : **A. prolifera** (Retz.) Copel. Gen. Fil. 144. 1947.

Ampelopteris prolifera (Retz.) Copel. Gen. Fil. 144. 1947. *Hemionitis prolifera* Retz. Obs. Bot. 6 : 31. 1791. *Goniopteris prolifera* (Retz.)

Presl. Tent. Pterid. 183. 1836 ; Bedd. Ferns South. India t. 172. 1864; Handb. Ferns Brit. India 296. t. 153. 1883.

Distrib. : Throughout India, Nepal, Bhutan, Burma, North Australia, South Africa and its Islands, Philippines, New Caledonia, China.

Amphineuron Holtt. Blumea 19 : 45. 1971 ; Blumea 23 : 205. 1977.

Type : **A. opulentum** (Kaulf.) Holtt. Blumea 19 : 45. f. 19. 19a. 1971.

A. immersum (Bl.) Holtt. in Nayar & Kaur, Comp. to Bedd. Handb. 203. 1974. *Aspidium immersum* Bl. Enum Pl. Jav. 156. 1828. *Lastrea immersa* (Bl.) Moore, Ind. Fil. LXXXIX. 1857 Bedd. Ferns Brit. India t. 252. 1867 ; Handb. Ferns Brit. India 234. t. 1883.

Distrib. : India (Assam), Hainan, Thailand, throughout Malesia, Queensland, New Hebrides, New Caledonia.

A. opulentum (Kaulf.) Holtt. Blumea 19 : 45. 1971. *Aspidium opulentum* Kaulf. Enum. Fl. 238. 1824. *Nephrodium extensum* (Bl.) Moore, Ind. Fil. 91. 1858 ; Bedd. Handb. Ferns Brit. India 269. 1883.

Distrib. : S. India, Sri Lanka, East Africa, Seychelles, Burma, Thailand, Malesia, N. Queensland, Islands of the Pacific to Tahiti, Trop. America.

A. terminans (Hook.) Holtt. Amer. Fern Journ. 63 : 82. 1973. *Nephrodium terminans* Hook. Sp. Fil. 4 : 73. 1862, excl. syn. *Nephrodium pteroides* sensu Bedd. Handb. Ferns Brit. India. 269. 1883.

Distrib. : South India, Burma to Hainan and Macao, throughout Malesia, Queensland, Central Africa, Fernando Poo.

Christella Léveillé, Fl. Kouy Tscheon 472. 1915.

Lectotype : **C. parasitica** (L.) Lé v. Fl. Kouy Tscheon 472. 1915.

C. appendiculata (Presl.) Holtt. Kew Bull. 31 (2) : 311. 1977. *Nephrodium appendiculatum* Presl, Epim Bot. 47. 1851. *N. extensum* var. *microsorum* Clarke, Trans. Trans. Linn. Soc. Lond. II. Bot. 1 : 530. 1880. *N. microsorum* (Clarke) Bedd. Handb. Ferns Brit. India 270. 1883 (*non* Hook. 1862). *N. molliusculum* Bedd. Handb. Ferns Brit. India Suppl. 68. 1892, *nom nov. superfl.* *N. extensum* var. *late-repens* Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 530. 1880.

Distrib. : India (Himalayan foot hills from 78° East, eastwards to upper Assam), China.

C. arida (D. Don) Holtt. in Nayar & Kaur, Comp. to Bedd. Handb. 206. 1974. *Aspidium aridus* D. Don, Prod. Fl. Nepal. 4. 1825.

Nephrodium aridus (D. Don) Sm. Hook. Journ. Bot. 4 : 188. 1841 : Bedd. Ferns Brit. India t. 297. 1868 ; Handb. Ferns Brit. India 272. 1883 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 531. 1880.

Distrib. : North West India to S. China, Thailand, Vietnam, throughout Malesia, N. Queensland, New Hebrides, New Caledonia, Fiji, Samoa.

C. clarkei (Bedd.) Holtt. in Nayar & Kaur, Comp. to Bedd. Handb. 203. 1974, *Pleocnemia clarkei* Bedd. Ferns Brit. India Suppl. 15. t. 368. 1876 ; Handb. Ferns Brit. India 297. t. 116. 1883. *Nephrodium artinexum* Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 536. 1880.

Distrib. : India (Sikkim).

C. cylindrothrix (Rosenst.) Holtt. in Nayar & Kaur, Comp. to Bedd. Handb. 208. 1974. *Dryopteris cylindrothrix* Rosenst. in Fedde. Repert. 12 : 246. 1913. *Nephrodium parasiticum* var. *aureum* (Clarke) Bedd. Handb. Ferns Brit. India 279. 1883. *N. procurrens* sensu Bedd. Handb. Ferns Brit. India Suppl. 67. 1892 (*pro parte*).

Distrib. : India (Sikkim, Assam), Bhutan, Burma, Thailand.

C. crinipes (Hook.) Holtt. in Nayar & Kaur, Comp. to Bedd. Handb. 208. 1974. *Nephrodium crinipes* Hook. Sp. Fil. 4 : 71. 1862 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 533. 1880 ; Bedd. Ferns Brit. India t. 263. 1868 ; Handb. Ferns Brit. India 279. 1883.

Distrib. : India (Meghalaya, Darjeeling & Sikkim Himalayas), Burma, China, North Thailand, Hainan.

C. dentata (Forssk.) Brownsey & Jermy, Brit. Fern Gaz. 10 : 338. 1973. *Polypodium dentatum* Forssk. Fl. Aegypt. arab. 185. 1773.

Distrib. : Throughout the tropics and subtropics of the old world.

C. evoluta (Bedd.) Holtt. in Nayar & Kaur Comp. to Bedd. Handb. 208. 1974. *Nephrodium amboinense* var. *evolutum* Clarke & Bak. in Journ. Linn. Soc. Bot. 24 : 417. 1888. *N. evolutum* Bedd. Handb. Ferns Brit. India Suppl. 76. 1892.

Distrib. : Eastern India, North Thailand.

C. gustavii (Bedd.) Holtt. in Nayar & Kaur, Comp. to Bedd. Handb. 208. 1974. *Nephrodium gustavii* Bedd. in Journ. Bot. 31 : 227. 1893. *N. evolutum* var. *gustavii* Bedd. Handb. Ferns Brit. India Suppl. 77. 1892.

Distrib. : India (Assam), N. Thailand.

C. harveyi (Mett.) Holtt. Kew Bull. 31 : 306. 1976. *Aspidium harveyi* Mett. in Kuhn, Linnaea 36 : 115. 1869.

Distrib. : India (cultivated in Indian Botanic Gorden, Howrah) and elsewhere in the tropics, polynesia.

C. hispidula (Decne) Holtt. Kew Bull. 31 : 312. 1976. *Aspidium hispidulum* Decne, Nouv. Ann. Mus. Hist. Nat. Paris 3 : 346. 1834. *Thelypteris hilsenbergii* (Presl) Panigr. Phytologia 31 : 369. 1975. *Christella quadrangularis* (Fée) Holtt. in Nayar & Kaur, Comp. to Bedd. Handb. 207. 1974.

Distrib. : India (South India, Meghalaya) Trop. America, Africa, Sri Lanka, Malaya, Malesia.

C. hokouensis (Ching) Holtt. Kew Bull. 31 (2) : 327. 1977. *Cyclosorus hokouensis* Ching, Bull. Fan Mem. Inst. Biol. ser. 2. 1 : 289. 1949.

Distrlb. : India (Assam), China.

C. kumaunica Holtt. Kew Bull. 31 (2) : 318. 1977.

Distrib. : India (Kumaun).

C. leboefii (Bak.) Holtt. in Nayar & Kaur, Comp. to Bedd. Handb. 206. 1974. *Polypodium leboefii* Bak. in Ann. Bot. 5 : 461. 1891. *Thelypteris leboefii* (Bak.) Panigr. Notes Roy. Bot. Gard. Edinb. 33 : 498. 1975. *Nephrodium biauritum* Bedd. Handb. Ferns Brit. India Suppl. 68. 1892.

Distrib. : India (Assam), Burma, Cambodia, Thailand.

C. malabariensis (Fée) Holtt. Kew Bull. 31 (2) : 317. 1977. *Nephrodium malabariense* Fée 10 Mém 43. 1865. *Thelypteris malabariensis* (Fee) Panigr. Notes. Roy. Bot. Gard. Edinb. 33 : 497. 1975. *Christella meeboldii* (Rosenst.) Holtt. in Nayar & Kaur, Comp. to Bedd. Handb. 208. 1974.

Distrib. : North East to Southern India. China, Sri Lanka.

C. namburensis (Bedd.) Holtt. in Nayar & Kaur, Comp. to Bedd. Handb. 206. 1974. *Nephrodium namburensis* Bedd. Handb. Ferns Brit. India Suppl. 69. 1892.

Distrib. : India (Assam), Thailand.

C. papilio (Hope.) Holtt. in Nayar & Kaur, Comp. to Bedd. Handb. 208. 1974. *Nephrodium papilio* Hope, Journ. Bomb. Nat. Hist. Soc. 12 : 625. t. 12. 1899. *Nephrodium molle* var. *major* Bedd. Handb. Ferns Brit. India Suppl. 76. 1892 (*pro parte*).

Distrib. : India (North - East to South India), Sri Lanka, Thailand, Malaysia to Northern Malaya.

C. papyracea (Bedd.) Holtt. in Nayar & Kaur, Comp. to Bedd. Handb. 206. 1974. *Nephrodium papyraceum* Bedd. Handb. Ferns Brit. India Suppl. 69. 1892.

Distrib. : India (Assam), Nepal, Burma.

C. parasitica (L.) Lév. Fl. Kouy tescheon 475. 1915. *Polypodium parasiticum* L. Sp. Pl. 2 : 1090. 1753. *Nephrodium procurrens* (Mett.) Bak. Sp. Fil. 290. 1867 ; Bedd. Handb. Ferns Brit. Iddia Suppl. 67. 1892 (*pro parte*). *N. didymosorum* Parish ex Bedd. Ferns Brit. India t. 200. 1866. *N. molle* var. *didymosorum* Bedd. Handb. Ferns Brit. India 279. 1883. *Cyclosorus didymosorus* (Parish ex Bedd.) Nayar & Kaur, Comp. to Bedd Handb. 208. 1974. *Nephrodium tectum* Bedd. Handb. Ferns Brit. India Suppl. 79. 1892 excl. King 1857.

Distrib. : North East and Southern India. Wetter parts of tropics and subtropics of Asia, Malesia, Queensland, Pacific to Tahiti and Hawaii, E. Africa, St. Helena.

C. semisagittata (Roxb. ex Griff.) Holtt. Kew Bull. 31 (2) : 334. 1977. *Polypodium semisagittatum* Roxb. ex Griff. Calc. Journ. Nat. Hist. 4 : 491. 1844.

Distrib. : India (Bengal, Assam), Burma, Bangladesh.

C. siamensis (Tagawa & K. Iwats.) Holtt. Kew Bull. 31 (2) : 332. 1977. *Thelypteris siamensis* Tagawa & K. Iwats. Acta Phytotax. Geobot. 22. 101. f. 5. 1967.

Distrib. : North East India, South China, Thailand.

C. subelata (Bak.) Holtt. Kew Bull. 31 (2) : 331. 1977. *Nephrodium subelatum* Bak. Bull. Misc. Inf. Kew. 1906. 11. 1906. *N. multilineatum* var. *assamicum* Bedd. Journ. Bot. 31 : 228. 1893. *Christella assamica* (Bedd.) Holtt. in Nayar & Kaur, Comp. to Bedd. Handb. 210. 1974.

Distrib. : Eastern India, China, North Thailand.

C. subpubescens (Bl.) Holtt. Webbia 30 : 193. 1976 ; Kew Bull. 31 : 323. 1977. *Aspidium subpubescens* Bl. Fnum. Pl. Jav. 149. 1828. *Nephrodium molle* var. *major* Bedd. Handb. Ferns Brit. India Suppl. 76. 1892. *Thelypteris blumei* Panigr. Pytologia 30 : 409. 1975. *Nephrodium molle* var. *amboinense* Bedd. Handb. Ferns Brit. India 278. 1883 (*pro parte*).

Distrib. : North East India to S. W. China, Burma. Thailand, Vietnam, Malesia, N. Queensland, New Hebrides, Fiji, Samoa.

C. zeylanica (Fée) Holtt. in Nayar & Kaur, Comp. to Bedd. Handb. 201. 1974. *Nephrodium zeylanicum* Fée 10e Mém. 42. 1865. *N. srilankensis* Panigr. Notes Roy. Bot. Gard. Edinb. 33 : 429. 1975. *N. extensum* var. *minor* Bedd. Ferns Brit. India t. 201. 1866. *N. molle* sensu Bedd. Handb. Ferns Brit. India 277. 1883 (*pro parte*).

Distrib. : India (Nicobar Islands), Sri Lanka.

Coryphopteris Holtt. Blumea 19 : 33. 1971.

Type : **C. viscosa** (Bak.) Holtt. Blumea 18 : 33. 1971.

C. didymochlaenoides (Clarke) Holtt. in Nayar & Kaur, Comp. to Bedd. Handb. 203. 1974. *Nephrodium gracilescens* var. *didymochlaenoides* Clarke. Trans. Linn. Soc. Lond. II. Bot. 1 : 514. 1880. *Lastrea hirsutipes* var. *didymochlaenoides* (Clarke) Bedd. Handb. Ferns Brit. India Suppl. 52. 1892.

Distrib. : India (Meghalaya).

C. hirsutipes (Clarke) Holtt. in Nayar & Kaur, Comp. to Bedd. Handb. 203. 1974. *Nephrodium gracilescens* var. *hirsutipes* Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 514. t. 67. f. 1. 1880. *Lastrea gracilescens* sensu Bedd. Handb. Ferns Brit. India 234. 1883 (*pro parte*). *Lastrea hirsutipes* (Clarke) Bedd. Handb. Ferns Brit. India Suppl. 52. 1892. (excl. var. *didymochlaenoides*).

Distrib. : N. E. India, China, Japan. Taiwan. Ryukyu Islands, Thailand, Malaya, Sumatra.

Cyclogramma Tagawa, Acta Phytotax. Geobot. 7 : 52. 1938.

Type **C. simulans** (Ching) Tagawa, Acta Phytotax. Geobot. 7 : 52. 1938.

C. auriculata (J. Sm.) Ching, Acta Phytotax. Sinica. 317. 1963. *Phegopteris auriculata* J. Sm. Hist. Fil. 233. 1875 ; Bedd. Handb. Ferns Brit. India 290. f. 149. 1883. excl. syn. *Polypodium auriculatum* Wall. ex Hook. Sp. Fil. 4: 237. 1862, non L. nec Radd., nec Presl ; Bedd. Ferns Brit. India t. 203. 1866 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 543. 1880.

Distrib. : India (North West Himalayas, Eastern Himalayas), Bhutan, Java.

C. squamaestipes (Clarke) Tagawa, Acta Phytotax. Geobot. 7 : 52. 1938. *Polypodium appendiculatum* var. *squamaestipes* Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 543. t. 79. f. 2. 1880.

Distrib. : India (Eastern Himalayas).

Cyclosorus Link, Hort. Berol. 2 : 128. 1833 ; Holtt. Blumea 19 : (1) : 27. 1971.

Type : **C. goggilodes** (Schkuhr) Link, Hort. Berol. 2 : 128. 1833.

C. goggilodes (Schkuhr) Link, Hort. Berol. 2 : 128. 1833. *Aspidium goggilodus* Schkuhr, Kr. Gew. 1 : 193. t. 33 C. 1809.

Distrib. : Pantropic.

C. interruptus (Willd.) H. Ito, Bot. Mag. Tokyo 51 : 714. 1937. (nomen tantum). *Pteris interrupta* Willd. Phytographia 13. t. 10. f. 1. 1794. *Nephrodium propinquum* R. Br. Prod. Fl. Nov. Holl. 148. 1810 ; Bedd. Ferns South. India t. 89. 1863. *Nephrodium pteroides* Hook. & Bak. Syn, Fil. 289. 1867 ; Bedd. Handb. Ferns Brit. India 269. 1883. *Nephrodium unitum sensu* Bedd. Handb. Ferns Brit. India 268. 1883.

Distrib. : Tropics and subtropics of the world.

More thorough studies are required to understand the genus properly.

Dictyocline Moore, Gard. Chron. 854. 1855.

Lectotype ; **D. griffithii** Moore, Gard. Chron. 854. 1855.

D. griffithii Moore, Gard. Chron. 854. 1855 ; Bedd. Ferns Brit. India t. 155. 1867. *Hemionitis griffithii* Hook. et Thomson in Hook. Sp. Fil. 5 : 193. 1864 ; Bedd. Handb. Ferns Brit. India 415. t. 246. 1883.

Distrib. : India (Meghalaya, Eastern Himalayas), Tonkin, Formosa, China, Japan, Malaya.

Glaphyropteridopsis Ching, Acta phytotax. Sin. 8 : 320. 1963.

Typs ; **G. erubescens** (Wall. ex Hook.) Ching, Acta Phytotax. Sin. 8 : 320. 1963.

G. erubescens (Wall. ex Hook.) Ching. Acta Phytotax. Sin 8 : 320. 1963. *Polypodium erubescens* Wall. ex Hook. Sp. Fil. 4 : 236. 1862. *Phegopteris erubescens* (Wall. ex Hook.) J. Sm. Hist. Fil. 333. 1875 ; Bedd. Ferns Brit. India t. 213. 1867 ; Handb. Ferns Brit. India 290. 1883.

Distrib. : India (Throughout Himalayan region), China, Formosa, Malay Peninsula, Malay Islands.

Macrothelypteris (H. Ito) Ching, Acta Phytotax. Sin. 8 : 308. 1963.

Type : *M. oligophlebia* (Bak.) Ching, Acta Phytotax. Sin. 8 : 308. 1963.

M. ornata (Wall. ex Bedd.) Ching, Acta Phytotax. Sin. 8 : 309.

1963. *Polypodium ornatum* Wall. ex Bedd. Ferns Brit. India t. 171. 1864 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 545. 1880. *Phegopteris ornata* J. Sm. Hist. Fil. 233. 1875 ; Bedd. Handb. Ferns Brit. India 294. 1883.

Distrib. : N. E. India, Burma.

M. torresiana (Gaud.) Ching, Acta Phytotax. Sin. 8 : 310. 1963.

Polystichum torresianum Gaud. in Freyc. Voy. Bot. 333. 1827. *Nephrodium tenericaule* Hook. Sp. Fil. 4 : 142. 1862, (excl. t. 269) ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 528. 1880. *Lastrea tenericaulis* Moore, Ind. Fil. 99. 1858 ; Bedd. Handb. Ferns Brit. India 266. 1883. *Lastrea setigera* Bedd. Ferns Brit. India Correct. II. 1870. (non Moore 1858.)

Distrib. : South India and Himalayas, Mascareen Islands, warmer parts of mainland of Asia and Japan, Malesia, N. E. Australia, Polynesia, Hawaii, Malaya.

Metathelypteris (Bl.) Ching, Acta Phytotax. Sin. 8 : 305. 1963.

Type : *M. gracilescens* (Bl.) Ching, Acta Phytotax. Sin. 8 : 305. 1963.

M. decipiens (Clarke) Ching, Acta Phytotax. Sin. 8 : 306. 1963.

Nephrodium gracilescens var. *decipiens* Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 541. t. 65. f. 2. 1880 ; Bedd. Handb. Ferns Brit. India 234. 1883. *Lastrea gracilescens* var. *decipiens* (Clarke) Bedd. Handb. Ferns Brit. India Suppl. 51. 1892.

Distrib. : India (Eastern Himalays).

M. flaccida (Bl.) Ching, Acta Phytotax. Sin. 8 : 306. 1963. *Aspidium flaccidum* Bl. Enum. Pl. Jav. 2 : 182. 1891. *Lastrea flaccida* (Bl.) Moore, Ind. Fil. 92. 1858 ; Bedd. Ferns South. India t. 250. 1865 ; Handb. Ferns Brit. India 244. 1883.

Distrib. : India (South India and Himalays), Sri Lanka, Java, China, Philippines.

M. gracilescens (Bl.) Ching, Acta Phytotax. Sin. 8 : 305. 1963.

Aspidium gracilescens Bl. Enum. Pl. Jav. 155. 1828. *Lastrea gracilescens* (Bl.) Moore, Ind. Fil. 93. 1858 ; Bedd. Handb. Ferns Brit. India 234. 1883 (*pro parte*) ; Suppl. 51. 1892 (*pro parte*).

Distrib. : India (Himalays), Japan, Taiwan, Malesia, Malaya, Sumantra, W. Java, N. Borneo, Luzon, Minadano, New Guinea.

M. uraiensis (Ros.) Ching, Acta Phytotax. Sin. 8 : 306. 1963.

Dryopteris uraiensis Ros. Hedw. 56 : 341. 1915. *Thelypteris uraiensis* (Ros.) Ching, Bull. Fan Mem. Inst. Biol. Bot. 6 : 336. 1936.

Distrib. : Eastern India, China, Japan, Taiwan.

Oreopteris Holub, Folia Geobot. Phytotax. 4 : 49. 1969 ; Holtt. Kew Bull. 36(2) : 225. 1981.

Type : **O. limbosperma** (All.) Holub, Folia Geobot. Phytotax. 4 : 49. 1969.

O. elwesii (Bak.) Holtt. in Nayar & Kaur, Comp. to Bedd. Handb. 204. 1974. *Nephrodium elwesii* Bak. in Hook. et Bak. Syn. Fil. ed. 2 : 497. 1874. *Lastrea elwesii* (Bak.) Bedd. Ferns Brit. India Suppl. t. 376. 1876 ; Handb. Ferns Brit. India 287. 1883 ; Suppl. Handb. 54. 1892.

Distrib. : India (Sikkim).

Parathelypteris (H. Ito) Ching, Acta Phytotax. Sin. 8 : 300. 1963.

Type : **P. glanduligera** (Kze.) Anal. 44. 1857.

P. beddomei (Bak.) Ching, Acta Phytotax. Sin. 8 : 302. 1963. var. **beddomei** *Nephrodium beddomei* Bak. Syn. Fil. 267. 1867. *Lastrea gracilescens sensu* Bedd. Ferns South. India t. 110. 1863. *Lastrea beddomei* (Bak.) Bedd. Ferns Brit. India Corr. 1870 ; Handb. Ferns Brit. India 239. 1883.

Distrib. : South India, Sri Lanka, Taiwan, Formosa, Japan, Malesia, Philippines.

P. glanduligera (Kunze.) Ching, Acta Phytotax. Sin. 8 : 303. 1963. *Aspidium glanduligerum* Kunze, Anal. Pterid. 44. 1837. *Lastrea gracilescens* var. *glanduligera* Bedd. Handb. Ferns Brit. India Suppl. 51. 1892.

Distrib. : India (Eastern India and Eastern Himalays), Nepal eastwards to S. China, Korea, Formosa, Japan and South to Thailand and N. Malesia.

Phegopteris (Presl.) Fée, Gen. Fil. 242. 1850-52, emend Ching, Acta Phytotax. Sin. 8 : 312. 1963 ; Holtt. Blumea 17 : 9. 1969.

Type : **P. connectilis** (Michx.) Watt. Canad. Nat. 29. 1870.

P. connectilis (Michx.) Watt. Canad. Nat. 29. 1870. *Polypodium connectile* Michx. Fl. Bor. Amer. 2 : 271. 1803. *Phegopteris vulgaris* Mett. Fil. Hort. Lips. 83. 1856 nom. illeg. ; J. Sm. Ferns. Brit. & For. 171. 1866 ; Bedd. Handb. Ferns Brit. India 290. 1883.

Distrib. : India (Himalayas), North temperate & Boreal regions.

Pneumatopteris Nakai, Bot. Mag. Tokyo 47 : 179. 1933 ; Holtt. Blumea 21 : 293. 1973.

Type : **P. callosa** (Bl.) Nakai, Bot. Mag. Tokyo 47 : 179. 1933.

P. truncata (Poir.) Holtt. Blumea 21 (2) : 314. 1973. *Polypodium truncatum* Poir, Encycl. Meth. 5 : 534. 1804.

Distrib. : South India, N.E. India, Sri Lanka, China, Malesia, Philippines.

P. truncata var. *loyalli* Holtt. Blumea 21 (2) : 314. 1973.

Distrib. : India (Darjeeling).

Pronephrium Presl, Epim. Bot. 258. 1849 ; Holtt. Blumea 20 : 105. 1972.

Lectotype : **P. lineatum** (Bl.) presl, Epim. Bot. 259. 1849.

P. articulatum (Houlst. & Moore) Holtt. Blumea 21 (1) : 116. 1972. *Nephrodium articulatum* Houlst. & Moore, Gard. Mag. Bot. 293. 1851. *N. glandulosum* var. *late-strigosum* Clarke, Trans. Lin. Soc. II. Bot. 1 : 532. t. 74. f 2. 1880.

Distrib. : India, Bangladesh, Sri Lanka, Burma, N. Thailand, China.

P. lakhimpurensis (Rosenst.) Holtt. Blumea 20 (1) : 110. 1972. *Dryopteris lakhimpurensis* Rosenst. Meded. Rijksher. 31 : 7. 1917. *Meniscium cuspidatum* var. *longifrons* Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 572. 1880.

Distrib. : India (Himalayas), China, Tonkin, Thailand.

P. nudatum (Roxb. ex Griff.) Holtt. Blumea 21 (1) : 111. 1972. *Polypodium nudatum* Roxb. ex Griff. Calc. Journ. Nat. Hist. 4 : 491. 1844. *Nephrodium moulmeinense* Bedd. Ferns Brit. India Suppl. 18. 1876.

Distrib. : India (Himalayas), Burma, Thailand, Tonkin, China.

P. parishii (Bedd.) Holtt. Blumea 20 (1) : 123. 1972. *Meniscium triphyllum* var. *parishii* Bedd. Ferns Brit. India t. 184. 1866.

Distrib. : India (Meghalaya, Assam, Andaman & Nicobar Islands), Sri Lanka, Malaya, Tonkin.

P. penangianum (Hook.) Holtt. Blumea 20 (1) : 110. 1972. *Polypodium penangiannum* Hook. Sp. Fil. 5 : 13. 1863. *Nephrodium costatum* Bedd. Ferns Brit. India t. 220. 1867.

Distrib. : North India from Kashmir eastwards, China.

P. stenopodium Chandra, Kew Bull Bull. 26 : 81. 1971.

Distrib. : India (Assam).

P. thwaitesii (Hook.) Holtt. Blumea 20 (1) : 122. 1972. *Meniscium thwaitesii* Hook. Fil. Exot. t. 83. 1859.

Distrib. : South India, Sri Lanka.

P. triphyllum (Sw.) Holtt. Blumea 20 (1) : 122. 1972. *Meniscium triphyllum* Sw. Schrad. Journ. Bot. 1800. (2) : 16. 1801.

Distrib. : India, Sri Lanka, Burma, Thailand, China, Japan, Malesia, Queensland.

Pseudocyclosorus Ching, Acta Phytotax. Sin. 8 : 324. 1963 ; Holtt. & Grimes, Kew Bull. 34 : 499. 1979.

Type : *P. tylodes* (Kunze) Ching, Acta Phytotax. Sin. 8 : 322. 1963.

P. canus (Bak.) Holtt. & Grimes, Kew Bull. 34 (3) : 509. 1979.
Nephrodium canum Bak. Syn. Fil. 267. 1867 : Clarke, Trans Linn Soc. Lond. II Bot. 1 : 515. 1880. *Lastrea cana* Bedd. Ferns Brit. India t. 307. 1869 ; *sensu* Bedd. Handb. Ferns Brit. India 238. 1883 ; excl. Bedd. Ferns Brit. India Suppl. t. 370 1876 ; and *Aspidium eburneum* wall. List no. 389. 1828. *Nephrodium repens* Hope, Journ. Bomb. Nat. Hist. Soc. 12 : 535. t. 8. 1899.

Distrib. : India (Foot hills of Himalayas from Chamba Punjab eastwards to sikkim, Meghalaya).

P. esquirolii (C. Chr.) Ching, Acta Phytotax. Sin. 8 : 322. 1963.
Dryopteris esquirolii C. Chr. Bull. Acad. Géogr. Bot. 144. 1907. *Nephrodium canum* Bak. in Hook. Syn. Fil. 267. 1867 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 515. 1880 (*pro parte*).

Distrib. : India (Sikkim), China, Vietnam, Taiwan, Japan.

P. falciloba (Hook.) Ching, Acta Phytotax. Sin. 8 : 324. 1963.
Lastrea falciloba Hook. Journ. Bot. Kew Misc. 9 : 337. 1857 ; Bedd. Ferns South. India t. 105. 1863. *Lastrea calcarata* var. *falciloba* Bedd. Handb. Ferns Brit. India 235. 1880 (*pro parte*). *Nephrodium falcilobum* Hook. Sp. Fil. 4 : 108. 1862 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 515. 1880.

Distrib. : India (Meghalaya, Sikkim), China, Annam, Tonkin, Japan. Burma.

P. gamblei Holtt. & Grimes, Kew Bull. 34 : 511. 1979.

Distrib. : South India.

P. griseus (Bak.) Holtt. & Grimes, Kew Bull. 34 (3) : 513. 1979.
Nephrodium griseum Bak. Syn. Fil. 271. 1867. *Lastrea grisea* (Bak.) Bedd. Ferns Brit. India t. 335. 1870 ; Handb. Ferns Brit. India 241. 1883.

Distrib. : South India (Cochin).

P. ochthodes (Kze.) Holtt. in Nayar & Kaur. Comp. to Bedd. Handb. 204. 1974. *Aspidium ochthodes* Kunze, Linnaea 24 : 282. 1851 ; Bedd. Ferns South India t. 106. 1863. *Lastrea ochthodes* Moore, Ind. Fil 98. 1858 ; Bedd. Ferns South. India t. 106. 1863 ; Handb. 240. 1883 (*pro parte*). *Lastrea grisea* Bedd. Ferns Brit. India t. 335. 1870 ; Handb. Ferns Brit. India 241. 1883.

Distrib. : India Sri Lanka, China, Tonkin, Malaya, Peninsula.

P. ornatipes Holtt. et Grimes, Kew Bull. 34 : 505. 1979.

Distrib. : India (Eastern Himalayas), Nepal.

P. tylodes (Kunze) Ching, Acta Phytotax. Sin. 8 : 324. 1963. *Aspidium tylodes* Kunze, Linnaea 24 : 244. 283. 1851. *Lastrea tylodes* (Kunze) Moore, Ind. Fil. 107. 1858 ; Bedd. Ferns South India t. 107. 1863.

Nephrodium ochthodes var. *tylodes* Bedd. Handb. Ferns Brit. India 240. 1883. *N. tylodes* (Kunze) Hope, Journ. Bomb. Nat. Hist. Soc. 14 : 724. 1903.

Distrib. : India (Himalayas except North West part), Sri Lanka, Burma, China, Hong-Kong, Malesia, Philippines.

Pseudophegopteris Ching, Acta Phytotax. Sin. 8 : 313. 1963.

Type : *P. pyrrhorhachis* (Kunze) Ching, Acta Phytotax. Sin. 8 : 315. 1963.

P. aurita (Hook.) Ching, Acta Phytotax. Sin. 8 : 314. 1963. *Gymnogramme aurita* Hook. Icon. Pl. t. 974, 989. 1854 ; Bedd. Ferns Brit. India Suppl. 24. 1876 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 568. 1880. *Grammitis aurita* Moore, Gard. Chron. 389. 1856 ; Bedd. Ferns Brit. India t. 152. 1866. *Leptogramme aurita* Bedd. Handb. Ferns Brit. India 377. 1883.

Distrib. : North-East India, Burma, China, Tonkin, Malesia, New Guinea.

P. levingei (Clarke) Ching, Acta Phytotax. Sin. 8 : 314. 1963. *Gymnogramme aurita* var. *levingei* Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 568. 1880. *Leptogramme aurita* var. *levingei* Bedd. Handb. Ferns Brit. India 377. 1883. *Gymnogramme levingei* Bak. Ann. Bot. 5 : 483. 1891 ; Hope, Journ. Bomb. Nat. Hist. Soc. 15 : 99. t. 35. 1903. *Leptogramma levingei* Bedd. Handb. Ferns Brit. India Suppl. 99. 1892.

Distrib. : Kashmir to W. China.

P. pyrrhorhachis (Kunze) Ching, Acta Phytotax. Sin. 8 : 315. 1963.

Polypodium pyrrhorhachis Kunze, Linnaea 24 : 257. 1851. *Polypodium distans* D. Don. Prod. Fl. Nepal. 2. 1825. *nom. illeg. non* Kaulf. 1824 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 544. 1880 (*pro parte*). *Phegopteris distans* (D. Don.) Mett. Frangatt. 4 : 16. 1858 ; Bedd. Handb. Ferns Brit. India 292. 1883. *Lastrea microstegia* Bedd. Ferns Brit. India t. 39. 1865. *Polypodium paludosum* sensu Bedd. Ferns South. India t. 168. 1863 (*non* Bl.). *P. distans* var. *adnatum* Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 544. 1880. *P. late-repens* Trotter & Hope, Journ. Bomb. Nat. Hist. Soc. 12 : 628. 1899. *Thelypteris paludosa* (Bl.) Iwats. Acta Phytotax. Geobot. 19 : 11. 1961.

Distrib. : India, Sri Lanka, China, Tonkin.

P. pyrrhorhachis var. *glabrata* (Clarke) Holtt. Blumea 17 (1) : 24. 1969. *Polypodium distans* var. *glabratum* Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 544. 1880.

Distrib. : India (Sikkim), China.

P. rectangluaris (Zoll.) Holtt. Blumea 27 (1) : 18. 1969. *Polypodium rectangulare* Zoll. Syst. Verz. 37, 48. 1854. *P. distans* var. *minor* Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 545. t. 79. f. 1. 1880.

Distrib. : North-East India, Malesia.

Sphaerostephanos J. Sm. in Hook. Gen. Fil. t. 24. 1836.

Type : **S. asplenoides** J. Sm. in Hook. Gen. Fil. t. 14. 1839.

S. arbuscula (Willd.) Holtt. in Nayar & Kaur, Comp. to Bedd. Handb. 207. 1974. *Aspidium arbuscula* Willd. in Linn. Sp. Pl. 5 : 233. 1810.

S. kurzii Holtt. Kew Bull. 34 : 231. 1971.

Distrib. : South India, Sri Lanka, Mascareen Islands.

S. penniger (Hook.) Holtt. in Nayar & Kaur, Comp. to Bedd. Handb. 207. 1976. *Nephrodium pennigerum* Hook. Sp. Fil. 4 : 82. 1862 ; Bedd. Handb. Ferns Brit. India Suppl. 73. 1892. *N. peningerum* var. *multilineatum* Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 532. 1880. *N. multilineatum* sensu Bedd. Handb. Ferns Brit. India Suppl. 80. 1892.

Distrib. : Eastern India, Bangladesh, Sri Lanka.

S. unitus (L.) Holtt. Nayar & Kaur Comp. to Bedd. Handb. 206. 1974. var. **unitus**. *Polypodium unitum* L. Syst. Nat. ed. X 2 : 1326. 1759. *Nephrodium cucullatum* Bak. Syn. Fil. 50. 1874 ; Bedd. Handb. Ferns Brit. India 270. 1883.

Distrib. : Throughout India in mountainous regions, Burma, Sri Lanka, Thailand, Malesia, Malaya, Queensland.

Stegnogramma Bl. Enum. Pl. Jav. 172. 1828.

Type : **S. aspidioides** Bl. Enum. Pl. Jav. 172. 1828.

S. aspidioides Bl. Enum Pl. Jav. 173. 1828. *Gymnogramma stenogramma* Bl. Fl. Jav. Fil. 98. t. 44. 1829.

Distrib. : India (Meghalaya), Malesia.

S. asplenoides (C. Chr.) J. Sm. ex Ching, Sinensis 7 : 94. t. 2. 1936.

Dryopteris stenogramme var. *asplenoides* C. Chr. Ind. Fil. 294. 1905.

Distrib. : India (Assam, Meghalaya), China.

S. leptogrammoides K. Iwats. Acta Phytotax. Geobot. 19 : 119. 1963.

Distrib. : Eastern India, Japan.

Thelypteris Schmid. Icon. Pl. ed. J.C. Keller 45 t. 11, 13. 1763. *nom. cons.*

Type : **T. palustris** (Salisb.) Schott, Gen. Fil. ad. t. 10. 1834.

T. augescens (Link.) Munz et Johnston, Amer. Fern. Journ. 12 : 75. 1922 ; Chandra & Devi, Journ. Bomb. Nat. Hist. Soc. 73 : 244. 1972. *Aspidium augescens* Link, Fil. Sp. 103. 1841.

Distrib. : South India.

T. confluens (Thunb.) Morton, Contr. U.S. Nat. Herb. 38. 71. 1967 ; Holtt. Fl. Malesiana ser. II. (5) : 376. 1981. *Pteris confluens* Thunb. Prod. Fl. Cap. 171. 1800. *Lastrea thelypteris* var. *squamigera* Bedd. Handb. Ferns. Brit. India Suppl. 54. 1892. *Lastrea fairbankii* Bedd. Ferns Brit. India India t. 254. 1867 ; Handb. Ferns Brit. India 240. 1883.

Distrib : South India, Africa, Thailand, Laos, Malesia, New Zealand.

T. decursive-pinnata (van Hall) Ching, Bedd. Fan Mem. Inst. Biol. Bot. 6 : 275. 1936. *Polypodium decursive-pinnatum* van Hall, Nieuwe Verhdl. Nederl. Inst. 5 : 204. 1836. *Phegopteris decursive-pinnata* (van Hall) Fee, Gen. Fil. 242. 1852.

Distrib. : India (Kashmir), S. China, Tonkin.

T. phegopteris (L.) Slosson, Rybd. Fl. Rocky Mts. 1043. 1917. *Polypodium phegopteris* L. Sp. Pl. 2 : 1089. 1753.

Distrib. : North India, North and temperate America, Aleutian Islands, Siberia, N.W. China, Asia Minor, N. Europe.

T. palustris (Salisb.) Schott, Gen. Fil. ad t. 10. 1834. *Polypodium palustris* Salisb. Prod. 403. 1976. *Lastrea thelypteris sensu* Bedd. Ferns Brit. India t. 44. Handb. Ferns Brit. India 241. 1883.

Distrib. : South India, N. India, widely known from the temperate regions of both the hemispheres, except for South America. S. Indian plants are more scaly.

Thorough revisionary studies are required to understand the genus *Thelypteris* Schmid in India.

Trigonospora Holtt. Blumea 19 (1) : 29. 1971.

Type : **T. ciliata** (Benth.) Holtt. Blumea 19 (1) : 29 f. 4, 4a. 1971.

T. calcarata (Bl.) Holtt. in Nayar & Kaur, Comp. to Bedd. Handb. 203. 1974. *Aspidium calcaratum* Bl. Enum. Pl. Jav. 159. 1828. *Lastrea calcarata* Moore, Ind. Fil. 87. 1858 ; Bedd. Handb. Ferns Brit. India 235. 1883.

Distrib. : India, China, Australia, Malesia.

T. ciliata (Benth.) Holtt. Blumea 19 (1) : 29. f. 4, 4a. 1971. *Aspidium ciliatum* Benth. Fl. Hong Kong 455. 1861.

Distrib. : South India, and Himalayas, Burma, Sri Lanka.

T. sericea (Scott ex Bedd.) Holtt. in Nayar & Kaur, Comp. to Bedd. Handb. 24. 1974. *Lastrea sericea* Scott ex Bedd. Ferns Brit. India t. 308. 1869. *Nephrodium ciliatum* Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 514. 1880. *Lastrea calcarata* var. *sericea* (Scott ex Bedd.) Bedd. Handb. Ferns Brit. India 237. 1883.

Distrib. : South India and North India, Bangladesh, Nepal, China, Burma, Tonkin, Malay Peninsula.

ASPLENIACEAE. Mett. ex Frank

Asplenium L. Sp. Pl. 2 : 1678. 1751; Bir. Journ. Indian Bot. Soc. 43 : 556. 1964 ; Sledge. Bull. Brit. Mus. (N.H.) London Bot. 3 : 233. 1965.

Lectotype : *A. marinum* L. Sp. Pl. 2 : 1081. 1753.

A. adiantum-nigrum L. Sp. Pl. 2 : 1081. 1753 ; Bedd. Ferns Brit. India t. 62. 1865 ; Handb. Ferns Brit. India 156. 1883.

Distrib. : India (N.W. Himalayas), Europe, Asia, Africa, Sandwich Islands.

A. aethiopicum (N.L. Burm.) Becherer, Candollea 6 : 23. f. 1. 1935.
Trichomanes aethiopicum N.L. Burm. Fl. Ind. Fl. Cap. Prod. 28. 1768. *A. furcatum* Thunb. Prod. Pl. Cap. 172. 1800 ; Bedd. Handb. Ferns Brit. India 157. 1883. *A. laserpitiifolium* sensu Bedd. Ferns South. India 75. t. 225. 1865.

Distrib. : South India, Sri Lanka, Trop. America, Polynesian Islands, Cape Colony, Abyssinia, Mascareen Islands, Canaries.

A. affine Sw. Schrad. Journ. Bot. 1800 (2) : 56. 1801. f. *affine* Sledge, Bull. Brit. Mus. (N.H.) Bot. 3 (6) : 270. 1965.

Distrib. : S. India, Sri Lanka, Mascareen Islands, Java, Borneo, Islands of Jobia.

A. affine f. *majus* Sledge, Kew Bull. 15 : 408. 1962.

Distrib. : S. India.

A. auritum Sw. Schrad. Journ. Bot. 1800 (2) : 52. 1801 ; Bedd. Ferns South. India t. 137. 1865 ; Handb. Ferns Brit. India 149. 1883.

Distrib. South India, Trop. America, West Indies, Mascareen Islands.

A. bullatum Wall. ex Mett. Aspl. 51. 1859. *A. bulbiferum* auct. Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 485. 1880 ; Bedd. Handb. Ferns Brit. India 159. 1883.

Distrib. : Himalayas, Malay Peninsula, Australia, New Zealand, Mexico, New Caledonia, Natal and the east African Islands.

A. capillipes Makino, Bot. Mag. Tokyo 17 : 77. 1903.

Distrib. : India (Himalayas), Japan.

A. caudatum Forst. Fl. Insul. Austr. Prod. 80. 1786 ; Bedd. Ferns South. India t. 140. 1865 ; Handb. Ferns Brit. India 151. 1883.

Distrib. S. India, Malay Peninsula, Australia, Africa. Brazil.

A. cheilosorum Kunze. ex Mett. Aspl. 133. t. 5. f. 12-13. 1859 ; Mehra & Bir, Res. Bull. (N.S.) Punjab Univ. 15 : 157. 1964. *A. heterocarnum* Wall. ex Hook. Sp. Fil. 3 : 132. t. 175. 1860 ; Clarke, Trans. Linn.

Soc. Lond. II. Bot. 1 : 481. 1880 ; Bedd. Handb. Ferns Brit. India 153. t. 77. 1883.

Distrib. : India (Himalayas from Nepal to Bhutan, S. India, Meghalaya, Sri Lanka, Malay Peninsula, China, Borneo, Formosa, Philippines.

A. crinicaule Hance in Annal. Sci. Nat. ser. V. 5 : 254. 1866 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 479. 1880 ; Bedd. Handb. Ferns Brit. India 150. 1883. *A. falcatum* Bedd. Ferns South. India t. 141. 1865.

Distrib. : South India, Himalays, China.

A. crinicaule var. *sikkimense* Bir, Journ. Indian Bot. Soc. 43 : 564. 1964.

Distrib. : India (Sikkim).

A. decrescens Kunze in Linnaea 24 : 261. 1851. *A. contiguum* *sensu* Moore, Ind. Fil. 121. 1859 ; Bedd. Ferns South. India t. 140. 1865. *A. caudatum* *sensu* Hook. Sp. Fil. 3 : 152. 1960 ; Bedd. Handb. Ferns Brit. India 151. 1883.

Distrib. : South India, Sri Lanka, Malay Peninsula, Australia, Africa, Brazil.

A. disjunctum Sledge, Bull. Brit. Mus. (N.H.) London Bot. 3 : 273. 1965.

Distrib. : India, Sri Lanka.

A. elasticum Fée, Gen. Fil. 196. 1892.

Distrib. : India.

A. ensiforme Wall. ex Hook. & Grev. Icon. Fil. t. 71. 1829 ; Bedd. Ferns South. India t. 125. 1863 ; Handb. Ferns Brit. India 141. t. 71. 1883 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 476. 1880.

Distrib. : Throughout India in mountainous regions, Burma, Sri Lanka, China, Thailand, Indochina.

A. erectum Bory ex Willd. in Linn. Sp. Pl. ed. 4. 5 : 328. 1810. *A. lunulatum* var. *B. camptorachis* Bedd. Handb. Ferns Brit. India 148. 1883.

Distrib. : S. India, Sri Lanka, S. Africa, Java, Timor, Samoa, Hawaii, Islands.

A. falcatum Lam. Encycl. 2 : 306. 1786 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 476. 1880 ; Bedd. Handb. Eerns Brit. India 150. 1883.

A. caudatum *sensu* Bedd. Ferns South. India t. 143. 1865. *Trichomanes adiantoides* L. Sp. Pl. 2 : 1098. 1753. *Asplenium adiantoides* (L.) C. Chr. Ind. Fil. 99. 1905.

Distrib. : India (S. India. Sikkim), Sri Lanka, Bangladesh, Malay Peninsula, Australia, Africa, Polynesia.

A. falcatum var. **bipinnatum** Sledge, Bull. Brit. Mus. (N.H.) Lond. Bot. 3 : 262. 1965 ; Manickam & Ninan. New Botanist 1 : 26. 1976. *A. mysorense* Roth. Nov. Pl. 395. 1821 (*pro parte*). *A. spathulinum* sensu Bedd. Ferns South, India 75. t. 226. 1864. (*non* J. Sm. ex Hook.).

Distrib. : S. India, Madagascar, Mascareen, S. China, Malaya, Indonesia, Philippines, Polynesia, Australia, New Zealand.

A. finlaysonium Wall. ex Hook. Icon. Pl. t. 937. 1854. *Hemidictyum finnlaysonium* (Wall. ex Hook.) Moore, Ind. Fil. 50. 1857 ; Bedd. Handb. Ferns Brit. India 195. t. 96. 1883. *A. macrophyllum* sensu Ito in Hara's Fl. East. Himal. 488. 1966.

Distrib. : India (Eastern Himalayas, Eastern India), Bangladesh, Malaya Peninsula.

A. fontanum (L.) Bernh. Schrad. Journ. Bot. 1799. (1) : 314. 1799 ; Bedd. Ferns Brit. India t. 146. 1867 ; Handb. Ferns Brit. India 158. 1883. *Polyodium fontanum* L. Sp. Pl. 2 : 1089. 1753.

Distrib. : India (North-West Himalayas), Europe.

A. formosum Willd. in Linn. Sp. Pl. ed. 4. 5 : 329. 1810.

Distrib. : India (Eastern Himalayas), America, Africa, Sri Lanka, Australia, West Indies.

A. gardneri Bak. Gardn. Chron. 1873 : 712. 1873 : Bedd. Ferns Brit. India Suppl. t. 355. 1876 ; Handb. Ferns Brit. India 151. 1883.

Distrib. : India (Meghalaya), Sri Lanka.

A. germanicum Weis. Pl. Crypt. Fl. Gott. 299. 1770 ; Bedd. Handb. Ferns Brit. India Suppl. 31. 1892.

Distrib. : India (Kashmir), Scotland, Norway to Hungary and Dalmatia.

A. griffithianum Hook. Icon. Pl. 87. f. 928. 1854 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 476. 1880 ; Bedd. Ferns Brit. India t. 58. 1865 ; Handb. Ferns Brit. India 142. 1883.

Distrib. : N. E. India, Burma.

A. indicum Sledge, Bull. Brit. Mus. (N.H.) 3 : 264. 1965. *A. planicaule* Wall. List no. 189. 1829 (*nom. nud.*) ex Mett. Aspl. 157. 1859; Bedd. Ferns South. India t. 139. 1865. *A. laciniatum* var. *planicaule* Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 482. 1880. *A. laciniatum* sensu Bedd. Handb. Ferns Brit. India 154. 1883 (*pro parte*, non D. Don.)

Distrib. : India (North-West Himalayas ; Meghalaya, South India), Sri Lanka, China, Burma, Thailand, Indochina, Philippines, Japan.

A. inaequilaterale Willd. in Linn. Sp. Pl. ed. 4. 5 : 322. 1810. *A. brachyotus* Kunze in Linnaea 10 : 512. 1836. *A. trapeziforme* sensu Bedd. Ferns South. India 45. t. 134. 1865. *A. lunulatum* var. *trapeziforme* Bedd. Handb. Ferns Brit. India 148. 1883 (*pro parte*).

Distrib. : South India, Sri Lanka, S. America, S. Africa, Madagascar, Reunion.

A. khasianum Sledge, Kew Bull. 15 : 397. f. 1 A-D. 1962. *A. gardneri* Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 480. (non Bak. 1873).

Distrib. : India (Maghalaya).

A. laciniatum D. Don, Prod. Fl. Nepal 8. 1825 var. *laciniatum* ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 481. 1880. (*Pro parte*) : Bedd. Handb. Ferns Brit. India 154. 1883 (*pro parte*). Bedd. Ferns South. India t. 145. 1865.

Distrib. : Himalayas, S. India, Sri Lanka, Japan, Formosa.

A. laciniatum var. *acutipinna* Bir, Journ. Indian Bot. Soc. 43 : 558 1964.

Distrib. : India (Eastesn Himalayas).

A. laciniatum var. *subintegritolium* Mehra & Bir, Res. Bull. (N.S.) Punjab Univ. 15 : 157. 1864.

Distrib. : India (Eastern Himalayas).

A. macrophyllum Sw. Schrad. Journ. Bot. 1800 (2) : 52. 1801 ; Bedd. Handb. Ferns Brit. India 150. 1883.

Distrib. : Himalayas, Sri Lanka, Tonkin, Malay Peninsula, Malay Islands, Polynesia, Mascareen Islands.

A. neolaserpitifolium Tard. – Blot et Ching, Notul. Syst. 5 : 153. t. 6. f. 1, 2. 1936 ; Joseph & Singh, Bull. Bot. Surv. India 24 (1 – 4) : 189. t. 1. f. 1 – 4. 1982.

Distrib. : India (Arunachal Pradesh, Sikkim), S. China, Vietnam.

A. nidus L. Sp. Pl. 2 : 1079. 1753 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 457. 1880 ; Hope, Journ. Bomb. Nat. Hist. Soc. 13 : 459. 1901. *Thamnopteris nidus* Presl, Epim. Bot. 68. 1849 ; Bedd. Handb. Ferns Brit. India 137. 1883.

Distrib. : Throughout India in hilly regions, China, Japan.

A. nidus var. *acutifolium* Bir, Journ. Indian Bot. Soc. 43 : 567. 1964.

Distrib. : India (Sikkim).

A. nidus var. *phyllitidis* (D. Don) Bir, Journ. Indian Bot. Soc. 43 : 567. 1964. *A. phyllitidis* D. Don Prod. Fl. Nepal. 7. 1825. *Thamnopteris-nidus* var. *phyllitidis* (D. Don) Bedd. Handb. Ferns Brit. India 139. 1883.

Distrib. : India (North-Eastern Himalayas Andaman & Nicobar Islands), Malaya, Burma, and tropics of the old world.

A. nitidum Sw. Syn. Fil. 84. 280. 1806 ; Clarke Trans. Linn. Soc. Lond. II. Bot 1 : 483. 1880 ; Bedd. Handb. Ferns Brit. India 157. 1883. *A. nitidum* var. *obtusum* Bedd. Ferns South India t. 149. 1865.

Distrib. : India (Himalayas from Nepal to Assam, S. India), Sri Lanka, Malay Peninsula, Malay Islands, S. Africa.

A. normale D. Don, Prod. Fl. Nepal. 7. 1825 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 477. 1880 ; Bedd. Handb. Ferns Brit. India 144. 1883. *A. multijugum* Wall. ex Mett. Aspl. 135. 1859 ; Bedd. Ferns South. India f. 133. 1865.

Distrib. : South India, Himalayas, Nepal, Bhutan, Sri Lanka.

A. obscurum Bl. Enum. Pl. Jav. 181. 1828. *A. cristatum* Wall. List no. 211. 1828 (*nom. nud.*).

Distrib. : South India, North India, Sri Lanka, Africa, Burma, Thailand, Indo-china, Java.

A. paucivenosum (Ching) Bir, Bull. Bot. Surv. India 4 : 3. 1962. *Ceterach paucivenosa* Ching, Bull. Fan Mem. Inst. Biol. Bot. 2 : 210. f. 28. 1931. *Ceterachopsis paucivenosa* Ching, Bull. Fan Mem. Inst. Biol. Bot. 10 : 9. 1940.

Distrib. : India (Eastern Himalayas).

A. paucivenosum var. **majus** Bir, Amer. Fern Journ. 62 : 46. 1972.

Distrib. : India (Eastern Himalayas).

A. pellucidum Lam. Encycl. Méth. Bot. 2 : 305. 1986. *A. hirtum* Kaulf. Enum. 169. 1824 ; Bedd. Ferns Brit. India t. 194. 1867 ; Handb. Ferns Brit. India 149. 1883.

Distrib. : India (Sikkim), Burma, Mascareen Islands, Hong-Kong, Philippines, Solomon Islands.

A. perakense Mathew et Christ, Journ. Linn. Soc. 39 : 214. 1904.

Distrib. : E. India, Malay Peninsula.

A. planicaule Wall. ex Mett. Aspl. 158. 1828 ; Bedd. Ferns South. India t. 39. 1863 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 482. 1880. Bedd. Handb. Ferns Brit. India 154. 1883 (*pro parte*).

Distrib. : India (Himalayas, S. India), Sri Lanka, China, Burma, Japan, Formosa, Madagascar.

A. planicaule var. **obtusa** Bir, Journ. Indian Bot. Soc. 43 (4) : 561. f. 8. 1964.

Distrib. : India (Eastern Himalays).

A. planicaule var. **yoshinage** (Mak.) Bir, Journ. Indian Bot. Soc. 43 : 562. 1994. *A. yoshinage* Mak. Phan. Pter. Jap. IC III. t. 64. 1900. *A. planicaule* f. *viviparum* (Wu) Ching, Bull. Dept. Biol. Sun. Yatsen Univ. no. 6. 26. 1933.

Distrib. : Himalays, Japan to S.W. China, Africa.

A. prolongatum Hook. 2nd Cent. Ferns t. 42. 1860. *A. rutaefolium* Kunze in Linnaea 10 : 521. 1836. Bedd. Ferns South. India t. 138. 1856 ; Handb. Ferns Brit. India 159. 1883.

Distrib. : India (South India, Himalays), Sri Lanka, Bhutan, Nepal, Burma, Africa, Japan, Fiji.

A. pumilum var. **hymenophylloides** (Fée) Clarke, Trans. Linn. Soc. Lond. II Bot. 1 : 482. 1810.

Distrib. : India (Rajasthan) Abyssinia.

A. ruta-maria L. Sp. Pl. 2 : 1081. 1753 ; Bedd. Ferns. Brit. India t. 61. 1865 ; Handb. Ferns Brit. India 156. 1883.

Distrib. : India (Kashmir), Europe, N. Africa, Tibet, Siberia, United States.

A. rockii C. Chr. Contrib. U.S. Nat. Herb. 26 : 332. f. 27. 1931 ; Balakr, & Hazra, Bull. Bot. Surv. India 6 : 315. 1964.

Distrib. : Eastern India, Malay Islands.

A. septentrionale (L.) Hoffm. Deutschl. Fl. 2 : 12. 1795 ; Bedd. Handb. Ferns Brit. India 145. 1883. *Acrostichum septentrionale* L, Sp. Pl. 2 : 1068. 1753.

Distrib. : India (N.W. Himalayas), Arctic and alpine Europe, Asia, America.

A. sericula Fée, Mém. Fam. Foug. 5 : 196. 1852. *A. wightianum* Wall. ex Hook. Sp. Fil. 3 : 105. t. 167. 1860 ; Bedd. Ferns South. India t. 126. 1865 ; Handb. Ferns Brit. India 146. t. 73. 1887.

Distrib. : S. India, Sri Lanka, Philippines.

A. sorelii Hook. in Blakiston Five months on yangtsze 363, 364. 1862 ; Bir & Shukla, Nova Hedw. 16 : 477. 1968. *A. Saulii* Bak. in Hook. et Bak. Syn. Fil. ed. 2 : 216. 1874 ; Hope, Journ. Bomb. Nat. Hist. Soc. 13 : 661. t. 18. 1901.

Distrib. : India (N.W. Himalayas), China, Japan.

A. spathulinum J. Sm. Sp. Fil. 3 : 170. 1860.

Distrib. : India (Himalayas).

A. tenerum Forst. Prod. 80. 1786.

Distrib. : India, Australia, Sri Lanka, Malesia, Polynesia.

A. tenerum var. **terminans** Mett. in Abhandl. Senckneb. Naturf. Ges. 3 : 113. 1859.

Distrib. : South India, Sri Lanka, Malesia-Polynesia, Burma, Thailand, Indo China, Malaya, Java, Borneo, Philippines, New Guinea, polynesia.

A. tenuisolum D. Don, Prod. Fl. Nepal 8. 1825; Clarke, Trans. Linn.

Soc. Lond. II. Bot. 1 : 485. 1880 ; Bedd. Handb. Ferns. Brit. India 159 t. 79. 1883 ; Hope, Journ. Bomb. Nat. Hist. Soc. 13 : 667. 1901.

Disrib. : India (Himalayas, S. India), Nepal, Bhutan, Burma, Sri. Lanka.

A. trichomanes L. Sp. Pl. 2 : 1080. 1753 ; Bedd. Ferns South. India t. 147. 1865 ; Handb. Ferns Brit. India 144. 1883.

Distrib. : South India, North India from Nepal to Assam, W. China, Sri Lanka, Indo-china, Formosa.

A. unilaterale Lam. Encycl. Méth. 2 : 305. 1786 ; Bedd. Handb. Ferns Brit. India 152. 1883 ; Hope, Journ. Bomb. Nat. Hist Soc. 13. 658. 1901.

Distrib. : India (Himalayas), Sri Lanka, Burma, Indo - China, Malaya Peninsula, China, Formosa. Japan, Philippines, Malaya Islands, Polynesia, Hawaii, E. Africa.

A. unilaterale var. **majus** (C. Chr.) Sledge, Bull. Brit. Mus. (N.H.) Lond. Bot. 3 : 246. 1965. *A. unilaterale* f. *majus* C. Chr. in Bernice P. Bishop. Mus. Bull. 177. 67. 1943. *A. excisum* Presl, Epim Bot. 74. 1849 ; Mehra & Bir, Res. Bull. (N.S.) Punjab. Univ. 15. 158. 1964.

Distrib. : India (North India, South India), Indonesia, New Guinea, Japan, Philippines, Sri Lanka, Malaya, Indochina.

A. unilaterale var. **rivale** Bedd. Handb. Ferns Brit. India 153. 153. 1883.

Distrib. : South India.

A. unilaterale var. **udum** Atkin. ex Bedd. Handb. Ferns. Brit. India 153. 1883.

Distrib. : India (Himalayas).

A. varians Wall. ex Hook. & Grev. Icon. Fil. 2. t. 72. 1829 ; Bedd. Handb. Ferns Brit. India 158. 1883 ; Hope, Journ. Bomb. Nat. Hist. Soc. 13 : 667. 1901.

Distrib. : India (S. India & Himalayas), Sri Lanka, China, Formosa, Korea, Japan, S. Africa, Hawaii.

A. viride Hudson, Fl. Angl. 385. 1762 ; Bedd. Ferns Brit. India t. 64. 1865 ; Handb. Ferns Brit. India 143. 1883.

Distrib. : India (North-West Himalayas), Europe, Asia, N. America in Arctic and Alpine regions.

A. zenkerianum Kunze, Linnaea 24 : 259. 1851 ; Bedd. Handb. Ferns Brit. India 148. t. 75. 1883. *A. persicifolium* var. *latifolium* Hook. Sp. Fil. 3 : 109. 1860 ; Bedd. Ferns South. India t. 127. 1865.

Distrib. : South India, Sri Lanka.

Ceterach Willd. Anleit. Sclsbstud. Bot. 578. 1804. *nom. cons.*

Type : **C. officinarum** Willd. in Linn. Sp. Pl. 5 : 136. 1810.

C. officinarum Willd. in Linn. Sp. Pl. 5 : 136. 1810 ; Bedd. Ferns Brit. India t. 71. 1865. *Asplenium ceterach* L. Sp. Pl. 2 : 1080. 1753. *Hemidictyum ceterach* Bedd. Ferns Brit. India Suppl. 13. 1876 ; Handb. Ferns Brit. India 194. t. 94. 1883.

Distrib. : India (North - West Himalayas), Afghanistan, Western Asia, Europe, North - West Africa.

Ceterachopsis (J. Sm.) Ching, Bull. Fan Mem. Inst. Biol. Bot. 10 : 8. 1940.

Type : **C. dalhousiae** (Hook.) Ching, Bull. Fan Mem. Inst. Biol. Bot. 10 : 8. 1940.

C. dalhousiae (Hook.) Ching, Bull. Fan Mem. Inst. Biol. Bot. 10 : 8. 1940. *Asplenium alternans* Wall. ex Hook. Sp. Pl. 3 : 92. 1860 ; Bedd. Ferns Brit. India t. 59. 1865 ; Handb. Ferns Brit. India 142. t. 72. 1883.

Distrib. : India (N.W. Himalayas, eastward to Sikkim), Abyssinia.

Sinephropteris Mick. Britt. 28 (3) : 326. 1976.

Type : **S. delavayi** (Franch.) Mick. Britt. 28 (3) : 327. f. 1 - 2. 1976.

S. delavayi (Franch.) Mick. Britt. 28 (3) : 327. f. 1 - 2. 1976. *Scolopendrium delavayi* Franch. Bull. Soc. Bot. Fr. 32 : 28. 1885 ; Bedd. Handb. Ferns Brit. India Suppl. 41. 1892.

Distrib. : India (Sikkim, Manipur), Tibet, China.

ONOCLEACEAE Pic. Ser.

Matteuccia Tod. Giorn. Sc. Nat. Econ. Palermo. 1 : 255. 1866 ; Nayar & Kazmi, Bull. Nation. Bot. Gard. Lucknow no. 62. 11 f. 39. 1963.

Type : **M. struthiopteris** (L.) Tod. Giorn. Sc. Nat. Econ. Polermo 1 : 235. 1866.

M. orientalis (Hook.) Trev. Atti. Ist. Veneto III, 14 : 586. 1869. *Struthiopteris orientalis* Hook. 2nd Cent. Ferns t. 4 1860 ; Bedd. Ferns Brit. India t. 171. 1865 ; Handb. Ferns Brit. India 20. t. 9. 1883. *Onoclea orientalis* Hook. Sp. Pl. Fil. 4 : 161. 1862 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 434. 1880.

Distrib. : India (Sikkim, Assam, Meghalaya), China, Japan.

WOODSIACEAE (Diels) Ching ex Herter

Cheilanthes Hieron. Notizbl. Berlin 7 : 406. 1920.

Type : **C. straminea** (Brause) Hieron. Notizbl. Berlin 7 : 406. 1920.

C. straminea (Brause) Hieron. Notizbl. Berlin 7 : 406. 1920.

Cheilanthes siraminea Brause, Hedw. 54 : 205. t. 4 H. 1914.

Distrib. : India (North – East Himalayas), China.

Woodsia R. Br. Prod. Fl. Nov. Holl. 158. 1810 ; Ching, Sinensis 3 (5) : 130. 1932 ; Brown, Beech. Nova. Hedw. 16 : 102. 1956.

Lectotype : **W. ilvensis** R. Br. Prod. Fl. Nov. Holl. 158. 1810.

W. alpina R. Br. Prod. Fl. Nov. Holl. 158. 1810. **W. hyperborea** R. Br. Prod. Fl. Nov. Holl. 158. 1810 ; Clarke Trans. Linn. Soc. Lond. II. Bot. 1 : 434. 1880 ; Bedd. Handb. Ferns Brit. India 20. 1883.

Distrib. : India (Kashmir) Northern & Central Europe, N. & Central Asia, N. America, South to the Northern United States.

W. andersonii (Bedd.) Christ, Mém. Soc. Bot. France 1 : 45. 1905. **Gymnogramme andersonii** Bedd. Ferns Brit. India t. 190. 1867 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 569. 1880 ; Bedd. Handb. Ferns Brit. India Suppl. 100. 1892 (*pro parte*).

Distrib. : India (Himalayas), Tibetan plateau, China.

W. cycloloba Hand. – Mzt. Symb. Sin. 6 : 19. t. 1. f. 5. 1929.

Distrib. : India (North-West Himalayas, Sikkim), Nepal, China (Yunnan.).

W. elongata Hook. Sp. Fil. 1 : 62. t. 21 c. 1844 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 435. 1880 ; Bedd. Handb. Ferns Brit. India t. 14. 1865 ; Handb. Ferns Brit. India 22. t. 10. 1883.

Distrib. : India (Himalayas), Nepal, Bhutan.

W. hancockii Bak. Ann. Bot. 5 : 196. 1891.

Distrib. : India (North – West Himalayas), Nepal, China.

W. lanosa Hook. Syn. Fil. 47. 1866 ; Bedd. Handb. Ferns Brit. India Suppl. 22. 1892 (*non* Ferns Brit. India t. 341 1870).

Distrib. : India (N. W. Hamalayas), Sikkim, China.

W. rosthorniana Diels in Engl. Bot. Jahrb. 29 : 187. 1900. **W. delavayii** Christ, Bull. Soc. Brot. France 42. mém. 1 : 45. 1905. **W. lanosa** var. *attenuata* Christ. Bot. Gaz. 56 : 338. 1913.

Distrib. : India (Eastern Himalayas), Bhutan, China.

ATHYRIACEAE Ching

Acystopteris Nakai, Bot. Mag. Tokyo 47 : 180. 1933.

Type : *A. japonica* (Luerss.) Nakai Bot. Mag. Tokyo 47 : 180. 1933.

A. tenuisecta (Bl.) Tagawa, Acta Phytotax. Geobot. 7 : 73. 1938.
Aspidium tenuisectum Bl. Enum. Pl. Jav. 170. 1828. *Cystopteris setosa* (Bedd.) Bedd. Ferns Brit. India t. 312. 1869 ; Handb. Ferns Brit. India 71. 1883. *Lastrea setosa* Bedd. Ferns Brit. India t. 262. 1868. *non* Presl. 1848, *nec* Moore 1858. *Davallia setosa* (Bedd.) Bak. in Hook. et. Bak. Syn. Fil. 468. 1874.

Distrib. : India (Sikkim), China, Japan, Indo - China, Java.

Anisocampium Presl, Epim. Bot. 58. 1849.

Type : *A. cumingianum* Presl, Epim. Bot. 59. 1849.

A. cumingianum Presl, Epim. Bot. 59. 1849. *Nephrodium ottaria* *sensu* Bedd. Handb. Ferns. Brit. India 267 t. 137. 1883.

Distrib. : India (Assam, South India), Sri Lanka.

Athyriopsis Ching, Bull. Fan Mam. Inst. Biol. Bot. 9 : 63. 1964.

Type : *A. japonica* (Thunb.) Ching, Acta Phytotax. Sin. 9 : 63. 1964.

A. japonica (Thunb.) Ching, Acta Phytotax. Sin. 9 : 63. 1964.
Asplenium japonicum Thunb. Fl. Jap. 334. 1784. *Diplazium japonicum* (Thunb.) Bedd. Ferns Brit. India Suppl. 12. 1876 ; Handb. Ferns. Brit. India 180. 1883. *D. decussatum* Bedd. Ferns Brit. India t. 292. 1868.

Distrib. : Throughout India in mountainous regions, Sri Lanka, China, Japan, Polynesia, Malay Peninsula.

Athyrium Roth. Rom. Mag. 2(1) : 105. 1799 ; Sledge, Bull. Brit. Mus. (N. H.) Lond. Bot. 2 : 289. 1962.

Type : *A. filix-femina* (L.) Roth. Rom. Mag. 2(1) : 106. 1799.

A. anisopterum Christ, Bull. Herb. Boiss. 6 : 962. 1898. *A. macrocarpon* Bedd. Ferns South. India 51. t. 152. 1865 (*pro parte*). *Asplenium macrocarpon* var. *atkinsonii* Hook. & Bak. Syn. Fil. ed. 2 : 489. 1874. (*Atkinsoni*) ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 489. 1880.

Distrib. : India (N. W. Himalayas, eastwards to Assam, S. India), Burma, China, Indo - China, Malaya, Luzon. Borneo.

A. atkinsonii Bedd. Ferns Brit. India. Suppl. 11. t. 359. 1876 ; Handb. Ferns Brit. India 163. 1883. *Asplenium atkinsonii* (Bedd.) Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 487. 1880.

Distrib. : India (Sikkim), Japan.

A. atratum Bedd. Handb. Ferns Brit. India Suppl. 33. 1892.

Distrib. : India (Manipur).

A. attenuatum (Clarke), Tagawa, Acta Phytotax. et Geobot. 16 : 177. 1956. *Asplenium filix-femina* var. *attenuata* Clarke, Trans. Linn. Soc.

Lond. II. Bot. 1 : 462. t. 59. f. 1. 1880. *Athyrium filix-femina* var. *attenuata* (Clarke) Bedd. Handb. Ferns. Brit. India 169. 1883.

Distrib. : India (Kashmir, Sikkim).

A. brevisorum (Wall. ex Hook.) Moore, Ind. Fil. 117. 1859. *Asplenium brevisorum* Wall. ex Hook. Sp. Fil. 8 : 229. 1860.

Distrib. : India (Himalayas).

A. clarkei Bedd. Ferns Brit. India. Suppl. t. 360. 1876. *A. nigripes* var. *Clarkei* (Bedd.) Bedd. Ferns Brit. India 166. 1883. *A. nigripes* var. *tenuifrons* Wall. ex Bedd. Handb. Ferns. Brit. India Suppl. 33. 1892.

Distrib. : India (Eastern Hamalayas), Nepal.

A. crenatum (Sommerf.) Rupr. ex Nyland. Spicil. Pl. Fenn. 2 : 14. 1844. *Aspidium crenatum* Sommerf. Vet. A.K. Handl. 1834 : 104. 1835.

Distrib. : India (Kashmir), Europe, Asia temperate.

A. dentigerum (Clarke) Mehra & Bir, Amer. Fern Journ. 50. 279. 1960. *Asplenium filix-femina* var. *dentigera* Clarke, Trans Linn. Soc. Lond. II. Bot. 1 : 491. 1880. *Polypodium dentigerum* Wall. list no. 334. 1828.

Distrib. : India (Himalayas), Nepal.

A. dissitifolium (Bak.) C. Chr. Contr. U. S. Nat. Herb. 26 : 296. t. 18. 1931. *Polypodium dissitifolium* Bak. Kew Bull. Misc. 1895 : 54. 1895.

Distrib. : (Himalayas), Burma, Thailand, China.

A. drepanopterum (Kunz.) A. Br. ex Milde Fil. Fur. 49. 1867. *Polypodium drepanopterum* Kunze in Linnaea 23 : 278, 318. 1850. *Athyrium oxyphyllum* Moore Ind. Fil. XLIX. 1857 ; Bedd. Fern Brit. India t. 67. 1865 ; Handb. Ferns Brit. India 171. 1883.

Distrib. : India (Himalayas, S. India).

A. duthei (Bedd.) Bedd. Handb. Ferns. Brit. India Suppl. 34. 1892. *Asplenium duthei* Bedd. Journ. Bot. 27 : 72. 1889.

Distrib. : India (N. W. Himalayas, Sikkim).

A. falcatum Bedd. Ferns South. India t. 151 1865 ; Handb. Ferns Brit. India 164. 1883. *Athyrium drepanophyllum* Bedd. Handb. Ferns Brit. India Suppl. 32. 1892.

Distrib. : Throughout India in mountainous regions.

A. filix-femina (L.) Roth, Rom. Mag. 2 (1) : 106. 1799 ; Bedd. Handb. Ferns Brit. India 168. 1883. *Polypodium filix-femina* L. Sp. Pl. 2 : 1090. 1753.

Distrib. : India (Himalayas).

A. filix-femina var. *retusa* (Decne) Bedd. Handb. Ferns. Brit. India 170. 1883. *Cystopteris retusa* Decne, Jacq. Voy. Bot. 176. t. 177. 1844. *Asplenium filix-femina* var. *retusa* (Decne) Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 492. t. 49. f. 2. 1880.

Distrib. : India : Kashmir to Bhutan.

A. fimbriatum (Wall.) Moore, Ind. Fil. 185. 1860 ; Bedd. Ferns Brit. India t. 295. 1868 ; Handb. Ferns Brit. India t. 83. 1883 (*pro parte*). *Aspidium fimbriatum* Wall. List. no. 339. 1828 (*pro parte*). *Asplenium fimbriatum* Hook. Sp. Fil. 3 : 234. 1860. *A. andersonii* Clarke, Journ. Linn. Soc. 15 : 154. 1876. *A. atkinsoni* var. *andersonii* (Clarke) Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 494. 1880.

Distrib. : India (Himalayas), China.

A. flabellulatum (Clarke) Tard. Aspl. du Tonkin 81. t. 12. 1932. *Asplenium filix-femina* var. *flabellulata* (Clarke) Bedd. Handb. Ferns Brit. India 170. 1883.

Distrib. : India (Sikkim).

A. foliolosum (Wall.) Moore ex Sm. Cat. Hort. no. 6 : 22. 1859 ; Moore, Ind. Fil. 185. 1860 ; Bedd. Handb. Ferns Brit. India Suppl. 37. 1892. *A. fimbriatum* var. *foliosa* Wall. List no. 359. 1828 ; Bedd. Handb. Ferns Brit. India 172. 1883. *Asplenium fimbriatum* var. *foliosa* (Wall.) Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 495. t. 62. f. 2. 1880.

Distrib. : India (Sikkim), Nepal.

A. himalicum Ching ex Mehra & Bir, Res. Bull. (N. S.) Punjab. Univ. 15 : 137. f. 26 – 29. 1964.

Distrib. : India (Sikkim).

A. hohenackerianum (Kunze) Moore, Ind. Fil. XLIX. 1857 ; Bedd. Ferns South. India t. 151. 1865 ; Handb. Ferns Brit. India 163. 1883. *Altantoidea hohenackeriana* Kunze, Farrnkr. 2 : 63. t. 126. 1850. ("Hohenackeriana").

Distrib. : S. India, Sri Lanka.

A. mackinnoni (Hope) C. Chr. Ind. Fil. 143. 1905. *Asplenium mackinnoni* Hope, Journ. Bot. 34 : 124. 1896 ; Journ. Bomb. Nat. Hist. Soc. 14 : 122. 1902.

Distrib. : India (Himalayas).

A. macrocarpum (Bl.) Bedd. Ferns South. India 51. t. 153. 1865 (excl. t. 152.) ; Handb. Ferns Brit. India 165. 1883. *Aspidium macrocarpum* Bl. Enum. Pl. Jav. 162. 1828. *Asplenium macrocarpum* (Bl.) Hook. Sp. Fil. 3 : 222. 1860.

Distrib. : India (North and South India), Sri Lanka, Tonkin, Java.

A. mehrae Bir, Nova Hova Hedw. 4 : 165. 1 - 5. 1962.

Distrib. : India (Sikkim).

A. nakanoi Makino. Bot. Mag. Tokyo 23 : 247. 1909. *Asplenium macrocarpum* var. *unibinnatum* Clarke, Trans Linn. Soc. Lond. Bot. 1 : 489. 1880. *Athyrium macrocarpum* var. *unibinnatum* (Clarke) Bedd. Handb Ferns Brit. India 165. 1883.

Distrib. : India (Himalayas), Bhutan, Formosa, Japan.

A. nigripes (Bl.) Moore Ind. Fil. XLIX. 98. 1858. *Aspidium nigripes* Bl. Enum. Pl. Jav. 2 : 162. 1828. *Athyrium solenopteris* sensu Bedd. Handb. Ferns Brit. India 166. 1883 (*pro parte*), non T. Moore.

Distrib. : India (S. India, Himalayas), Sri Lanka.

A. parasnathense (Clarke) Ching ex Bir, Journ. Indian Bot. Soc. 43 : 569. 1964. *Asplenium filix-femina* var. *parsanathensis* Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 593. 1880. *Athyrium filix-femina* var. *parasnathensis* (Clarke) Bedd. Handb. Fern Brit. India 170. 1883.

Distrib. : India (Bihar : Parasnath hills).

A. pectinatum (Wall. ex Hope) Presl, Tent. Pterid. 98. 1836 ; Bedd. Handb. Ferns Brit. India Suppl. 36. 1892. *Asplenium pectinatum* Wall. List no. 231. 1828 (*nom. nud.*) ex Hope, Journ. Bombay Nat. Hist. Soc. 14 : 253. 1902. *A. filix-femina* var. *pectinata* (Wall. ex Hope) Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 492. t. 58. 1880. *Athyrium filix-femina* var. *pectinata* (Wall. ex Hope) Bedd. Handb. Ferns Brit. India 169. 1883.

Distrib. : Throughout India in mountainous regions.

A. praetarmissum Sledge in Arn & Mag. Nat. Hist. ser. 12. 9 : 457. t. 16. 1956. var. *praetermissum*. *A. costale* Moore. Ind. Fil. 180. 1860 (*pro parte*) quoad specim. ex Ceylon et Nilgiri, non *Aspidium costale* Bl. *A. nigripes* sensu Bedd. Ferns South. 52. t. 157. 1864 ; Handb. Ferns Brit. India 166. 1883 (*pro parte*), non T. Moore. *Asplenium nigripes* sensu Hook. et Bak. Syn. Fil. 227. 1867 *pro parte*, (non Hook) ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 490. 1880 (*pro parte*). *Athyrium solenopteris* sensu Bedd. Handb. Ferns Brit. India 166. 1883 (*pro parte*), quoad. Syn. *Athyrium nigripes* Bedd. non T. Moore.

Distrib. : India, Java.

A. puncticaule (Bl.) Moore, Ind. Fil. 1860. *Aspidium puncticaule* Bl. Enum. Pl. Jav. 156. 1828.

Distrib. : India (Himalayas).

A. rubricaule (Edgew.) Bir, Nova Hedw. 4 : 169. f. 12 - 13. 1862. *Asplenium filix-femina* var. *retusa* subvar. *rubricaulis* Edgw. in Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 492. t. 59. f. 2. 1880.

Distrib. : India (Eastern Himalayas).

A. rupicola (Hope) C. Chr. Ind. Fil. 145. 1905. *Asplenium rupicola* Hope, Journ. Bomb. Nat. Hist. Soc. 12 : 531. t. 5. 1899. *A. filix-femina* var. *retusa* Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 492. 1880 (*pro parte*) *Athyrium filix-femina* var. *retusa* (Clarke) Bedd. Ferns Brit. India 170. 1883 (*pro parte*).

Distrib. : Himalayas from Kashmir to Bhutan.

A. schimperi Moug. ex Féé, Mém. Gen. Fil. 187. 1850–52 ; Bedd. Handb. Ferns Brit. India Suppl. 36. 1862.

Distrib. : India (Eastern Himalayas) Abyssinia and the Cameroon mountains.

A. setiferum C. Chr. Ind. Fil. 146. 1905. *Athyrium nigripes* Mett. var. *disecta* Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 491. 1880. *Athyrium nigripes* Bedd. Handb. Ferns Brit. India 166. 1883 (*pro parte*).

Distrib. : India (Himalays), Nepal, Bhutan.

A. solenopteris (Kunze) Moore, Ind. Fil. 43. 1857 ; 187. 1860 ; Bedd. Handb. Ferns Brit. India 166. 1883. (excl. parte). var. *solenopteris*. *Allantoidea solenopteris* Kunze, Linnaea 24 : 266. 1851. *Athyrium pectinatum* Bedd. Ferns South. India 51. t. 155. 1864 ; *pro parte* ; non *Asplenium pectinatum* Wall. ex Mett.

Distrib. : S. India, Sri Lanka.

A. solenopteris var. *pusillum* (Kunze) Moore, Ind. Fil. 187. 1860. *Allantoidea solenopteris* var. *pusilla* Kunze in Linnaea 24 : 267. 1851. *Athyrium gymnogrammoides* (Klotzsch ex Mett.) Bedd. Ferns South. India 52. 1864 (excl. descr. et. fig.).

Distrib. : South India. Sri Lanka.

A. spinulosum (Maxim.) Milde, Bot. Zeit. 376. 1866 ; Bedd. Handb. Ferns Brit. India 161. 1883. *Cystopteris spinulosa* Maxim. in Mém. Acad. St. Petersb. 9 : 340. 1850.

Distrib. : India (Sikkim), Japan, China, Korea.

A. strigillosum Moore in Lowe Ferns Brit. Exot. 5 : t. 36. 1858. *Athyrium clarkei* Bedd. Ferns Brit. India Suppl. 11. t. 360. 1876. *Asplenium clarkei* Atkinson ex Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 489. 1880. *Asplenium tenuifrons* Wall. ex Hope, Journ. Bomb. Nat. Hist. Soc. 14 : 120. t. 22. 1902.

Distrib. : India (Himalayas), Burma, China.

A. subtriangulare (Hook.) Bedd. Ferns Brit. India t. 242. 1868. *Asplenium subtriangulare* Hook. Syn. Fil. 225. 1867. *A. spinulosum* var. *subtriangulare* (Hook.) Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 487. 1880.

Distrib. : India (Himalayas), China.

A. subtriangulare var. **sikkimense** Bir, Nova Hedw. 4 : 168. f. 9 – 11. 1962.

Distrib. : India (N. Sikkim).

A. wallichianum Ching, Bull. Fan. Mem. Inst. Biol. Bot. 8 : 497. 1938. *Lastrea brunoniana* (Wall. ex Mett.) Presl ex Bedd. Ferns Brit. India t. 37. 1865 ; Handb. Ferns Brit. India 246. 1883 ; Suppl. 55. 1892 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 522. 1880.

Distrib. : India (Himalayas), China.

Cornopteris Nakai, Bot. Mag. Tokyo 44 : 7. 1930.

Type **C. decurrenti-alata** (Hook.) Nakai, Bot. Mag. Tokyo 44 : 1938.

C. birii Ching ex Bir Nova Hedw. 7 : 502. f. 1 – 7. 1964.

Distrib. : India (Sikkim).

C. macdonellii (Bedd.) Tard. Amer. Fern. Journ. 48 : 32. 1958.

Asplenium macdonellii Bedd. Journ. Bot. 1889 : 73. 1889. *Athyrium macdonellii* (Bedd.) Bedd. Handb. Ferns Brit. India Suppl. 34. 1892 (Mc Donelli).

Distrib. : India (Himalayas), China.

C. opaca (D. Don) Tagawa, Acta Phytotax. Geobot. 8 : 92. 1939.

Hemionitis opaca D. Don, Prod. Fl. Nepal 13. 1825. *Leptogramma opaca* (D. Don.) Bedd. Handb. Ferns. Brit. India 379. t. 217. 1883. *Gymnogramma opaca* Spr. Syst. 4 : 39. 1827. Bed. Ferns Brit. India t. 238. 1867.

Distrib. : India (Himalayas), Nepal, Bhutan, China, Japan, Java.

C. tenuisecta (Bl.) Tard. Amer. Fern Journ. 48 : 32. 1958. *Aspidium tenuisectum* Bl. Enum. Pl. Jav. 170. 1828. *Cystopteris setosa* Bedd. Ferns Brit. India t. 312. 1869 ; Handb. Ferns Brit. India 204. 1883.

Distrib. : India (Himalayas), Tonkin, China, Formosa, Japan, Malay Islands.

Cystopteris Bernh. Schrad. Neu. Journ. 1 (2) : 26. 1805. Blasdell, Mem. Torrey Bot. Club. 21 : 1. 1963 ; Bir & Tirkha, Nova. Hedw. 47 : 1. 1964. *nom. cons.*

Type : *C. fragilis* (L.) Bernh. Neu. Journ. 1(2) : 26. t. 2. f. 9. 1805.

C. fragilis (L.) Bernh. Schrad. Neu. Journ. 1(2) : 26. t. 2. f. 9. 1805 ; Bedd. Ferns Brit. India t. 91. 1865 ; Haddb. Ferns Brit. India 70. 1883. *Polypodium fragile* L. Sp. 2 : 1091. 1753.

Distrib. : India (Himalayas) and temperate regions of the world.

C. fragilis f. **fragilis** Bir & Tirkha, Amer. Fern. Journ. 66(3) : 109. f. 1A, B. 1976.

Distrib. : India (Himachal Pradesh), Canada.

C. fragilis f. granulosa Bir. & Tirkha, Amer. Fern Journ. 66(3) : 109. f. 2A, B. 1976.

Distrib. : India (Himachal Pradesh, Kashmir.)

C. fragilis f. himalayensis Bir & Tirkha, Amer. Fern Journ. 66(3) : 110. 3A, B. 1976.

Distrib. : India (Kashmir).

C. montana (Lam.) Bernh. ex Desv. Prod. 264. 1827 ; Bedd. Handb. Ferns Brit. India Suppl. 15. 1892. *Polypodium montanum* Lam. Fl. Franc. 1 : 23. 1778.

Distrib. : India (Kashmir), mountains of Scandinavia, Scotland, Central Europe, Kamschatka, America.

C. sikkimensis Ching ex Bir, Nova Hedw. 7 : 504. f. 8 – 15. 1964.

Distrib. : India (Sikkim).

C. sudetica A. Br. & Milde, Jahrb. Schles. Ges. 1855 : 92. 1855 ; Bedd. Handb. Ferns Brit. India Suppl. 16. 1892 ; Hope, Journ. Bomb. Nat. Hist. Soc. 13 : 36. 1900. *C. moupinensis* Franch. Nouv. Arch. Mus. II. 10 : 111. 1887.

Distrib. : India (Sikkim), China, Siberia, Tibet.

Dictyodroma Ching, Acta Phytotax. Sin. 9 : 57. 1964.

Type : **D. heterophlebia** (Mett. ex Bak.) Ching, Acta Phytotax. Sin. 9 : 57. 1964.

D. heterophlebia (Mett. ex Bak.) Ching, Acta Phytotax. Sin. 9 : 57. 1964. *Aspidium heterophlebium* Mett. ex Bak. Syn. 243. 1867. *Anisogonium heterophlebium* Bedd. Ferns Brit. India t. 329. 1869 ; Handb. Ferns Brit. India 191. t. 93. 1883.

Distrib. : India (Himalayas) Nepal, Bhutan, Burma, China.

Diplaziopsis C. Chr. Ind. Fil. 227. 1905.

Type : **D. javanica** (Bl.) C. Chr. Ind. Fil. 227. 1905.

D. javanica (Bl.) C. Chr. Ind. Fil. 227. 1905. *Asplenium javanicum* Bl. Enum. Pl. Jav. 175. 1828. *Allantoidea javanica* Trev. Nu. Giorn. Bot. It. 159. (nota) 1875 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 505. 1880 ; Bedd. Handb. Ferns Brit. India 195. t. 97. 1883. *A. brunoniana* Wall. ex Bedd. Ferns South. India t. 159. 1865.

Distrib. : India (Himalayas), Nepal, Bhutan, Sri Lanka, China, Java, Sumatra.

Diplazium Sw. Schrad. Journ. Bot. 1800 (2) : 61. 1801 ; sledge, Bull. Brit. Mus. (N. H.) Lond. Bot. 2 : 275. 1962.

Lectotype : **D. plantagineum** (L.) Sw. Schrad. Journ. Bot. 1800 (2) : 62. 1802.

D. asperum Bl. Enum. Pl. Jav. 195. 1828 ; Bedd. Handb. Ferns Brit. India 184. 1883 ; *Allantoidea aspera* (Bl.) Ching, Acta Phytotax. Sin. 9 : 46. 1964. *Diplazium polypodicides* Bedd. Ferns South. India t. 163. 1865. *D. sikkimense* Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 500. t. 65. f. 1. 1880.

Distrib. : S. India, Himalayas, Sikkim, Java.

D. bellum (Clarke) Bir. Journ. Indian Bot. Soc. 43 : 57. 1964 ; *Asplenium bellum* Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 496. t. 63. f. 2. 1880.

D. cordifolium Bl. Enum. Pl. Jav. 190. 1828. *Athyrium cordifolium* (Bl.) Copel. Philip. Journ. Sci. 3 C. 300. 1908. *Anisogonium cordifolium* Bedd. Ferns Brit. India t. 331. 1870. Handb. Ferns Brit. India 191. t. 92. 1883.

Distrib. : India (Manipur), Malaya.

D. dilatatum Bl. Enum. Pl. Jav. 194. 1828 : *Asplenium latifolium* D. Don, Prod. Fl. Nepal 8. 1825 ; non *A. latifolium* Bory 1803 ; *Diplazium latifolium* Moore, Ind. Fil. 141. 1859; 331. 1861 ; Bedd. Handb. Ferns Brit. India 187. 1883. *Asplenium dilatatum* (Bl.) Hook. Sp. Fil. 3 : 258. 1860. *A. maximum sensu* Hook. & Bak. Syn. Fil. 239. 1867 *pro parte* ; non. *D. Don. Athyrium dilatatum* (Bl.) Holtt. Fl. Malaya 2 : 574. 1954.

Distrib. : India (South and East North India), China, Indo-China, Burma, Malaya, Philippines, N. Australia.

D. donianum (Mett.) Tard. Aspl. du. Tonkin 58. t. 5. 1932. *Asplenium donianum* Mett. Aspl. 177. no. 198b. 1859. *Diplazium fraxinifolium sensu* Bedd. Ferns Brit. India t. 59. 1865.

Distrib. : North-East India to China, Malay Peninsula, Malesian Islands.

D. esculentum (Retz.) Sw. Schrad. Journ. Bot. 1801 (1) : 312. 1803 ; *Hemionitis esculenta* Retz. Obs. Bot. 6 : 38. 1791. *Anisogonium esculentum* (Retz.) Presl, Tent. Pterid. 116. 1836 ; Bedd. Handb. Ferns Brit. India 192. 1883. *Callipteris esculenta* (Retz.) J. Sm. apud Houlst. & Moore in Gard. Mag. Bot. 3 : 265. 1851 ; Bedd. Ferns South. India 54. t. 164. 1865.

Distrib. : Throughout India, China, Formosa, Malaya east to Philippines, New Guinea, Samoa.

D. fieldingianum (Kunze) Panigr. Phytologia 31 (3) : 251. 1975. *Allantodia fieldingiana* Kunze, Linnaea 24 : 268. 1851. *Asplenium polymorpha* Wall. List. no. 230. 1828 (*nom. nud.*) *A. polypodioides* auct. Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 511. 1880 (*pro parte quoad*) *Asplenium (Diplazium) latifolium* D. Don var. *polymorpha* (Wall. ex Mett.) Hope, Jour. Asiatic Soc. Bengal 57 : 307. t. 17. 1888.

Distrib. : India (N.W. Himalayas), Nepal, Bhutan, China, Pakistan.

D. frondosum (Clarke) Christ. Not. Syst. Lacombe. 1 : 46. 1909 ; Panigr. Phytologia 24 : 252. 1975. *Asplenium latifolium* D. Don var. *frondosa* Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 503. 1880.

Distrib. : N.E. India, Sikkim, China, Pakistan, Nepal.

D. giganteum (Bak.) Ching in C. Chr. Ind. Fil. Suppl. 3 : 73. 1934. *Gynnogramme gigantea* Bak. in Journ. Bot. 27 : 177. 1889.

Distrib. : Himalayas to C. China.

D. himalayense (Ching) Panigr. Phytologia 31 (3) : 254. 1975. *Allantodia himalayense* Ching, Acta Phytotax. Sin. 9 : 55. 1964. *Asplenium polypodioides* Mett. var. *vestita* Clarke, Trans Linn. Soc. Lond. II. Bot. 1 : 501. 1880.

Distrib. : North-East India, Sikkim.

D. himalayense var. *effusior* (Clarke) Panigr. Phytologia 31(3) : 254. 1975. *Asplenium polypodioides* var. *effusior* Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 501. 1880. *Diplazium umbrosum* var. *multicaudatum* (Bedd.) Bedd. Handb. Ferns Brit. India 186. 1883 (*pro parte*), quoad. var. *effusior* Clarke, Syn. *tantum non Asplenium multicaudatum* (Clarke) Bak. 1880.

Distrib. : India (Sikkim), Nepal.

D. hirsutipes (Bedd.) Nayar et Kaur, Comp to Bedd. Handb. Ferns Brit. India 44. 1974. *D. stoliczkae* var. *hirsutipes* Bedd. Handb. Ferns Brit. India 182. 1883.

Distrib. : India (Himalayas), Nepal to Bhutan.

D. lasiopteris Kunze. Linnaea 27 : 568. 1843 ; Bedd. Ferns South India 53. t. 160. 1864. *D. thwaitesii* (A. Braun ex Mett.) Kletzsch ex Moore, Ind. Fil. 339. 1862 ; Bedd. Ferns Brit. India t. 291. 1868. *D. japonicum* sensu Bedd. Handb. Ferns Brit. India 180. 1883 (*pro. parte.*) non Bedd. 1876, *nec Asplenium japonicum* Thunb.

Distrib. : S. India, Sri Lanka.

D. lobulosum (Wall. ex Mett.) Presl, Tent. Pterid. 114. 1836. *Asplenium lobulosum* Wall. ex Mett. Aspl. 163 n. 170. 1859. *A. longifolium* D. Don, Prod. Fl. Nepal 7. 1825. *Diplazium longifolium* (D. Don) Moore, Ind. Fil. 141. 332. 1859 ; Bedd. Ferns Brit. India t. 247. 1865 ; Handb. Ferns Brit. India 179. t. 87. 1883.

Distrib. : India (N. W. Himalayas), Nepal.

D. muricatum (Mett.) v.A.v.R., Malayan Ferns 829. 1909; *Asplenium muricatum* Mett. Ann. Mus. Bot. Lugd.-Bat. 2 : 239. 1866. *Asplenium procerum* (Hook. & Bak.) Wall. ex Clarke, Trans. Linn. Soc. Lond. II Bot. I. 495. 1880. *non A. procerum* Bernh. 1802. *D. umbrosum* var. *australe* Bedd. Handb. Ferns Brit. India 189. 1883 (*pro parte*), *non Allantoidea australis* R. Br. *D. umbrosum* var. *procerum* (Hook. & Bak.) Bedd. Ferns. Brit. India 189. 1883.

Distrib. : S. India and E. India, Burma Nepal, Siam, Formosa, Java.

D. pin-faense Ching, Lingnan Sci. Journ. 15(2) : 279. 1936 ; Dixit & Ghosh, Bull. Bot. Surv. India 20 (1-4) : 160. f. 1-3. 1976. *Athyrium pin-faense* (Ching) Ohwi, Nat. Sci. Mus. Tokyo 3 : 100. 1956.

Distrib. : Eastern India, China, Japan.

D. pinnatifido-pinnatum (Hook.) Moore, Ind. Fil. 334. 1861.

Asplenium pinnatifido-pinnatum Hook. Sp. Fil. 3 : 238. 1860.

Distrib. : India.

D. polypodioides Bl. Enum. Pl. Jav. 194. 1828 ; Bedd. Ferns South. India 54. t. 163. 1864 ; Ferns Brit. India t. 293. 1868 ; Handb. Ferns Brit. India 184. 1883.

Distrib. : South India, Sri Lanka, East Indies, Philippines.

D. polypodioides var. **brachylobum** Sledge, Bull. Brit. Mus. (N.H.) London Bot. 2 : 307. 1962.

Distrib. : S. India, Sri Lanka.

D. simplicivenium Holtt. Gard. Bull. Str. Settl. 11 : 100. f. 6. 1940 ; Mehra and Bir, Res. Bull. (N.S.) Punjab Univ. 15 : 147. 1964. *Athyrium simplicivenium* (Holtt.) Holtt. Rev. Fl. Malaya 2 : 573. 1954.

Distrib. : India (Sikkim), Malaya Peninsula.

D. spectabile (Wall. ex Mett.) Ching, Lingnan Sci. Journ. 15 : 278. 1936. *Asplenium spectabile* Wall. ex Mett. Aspl. 196 no. 240. 1859. *Diplazium umbrosum* var. *multicaudatum* (Wall. ex Clarke) Bedd. Handb. Ferns Brit. India 190 t. 1883. *Asplenium multicaudatum* Wall. ex Clarke, Trans. Linn. Soc. Lond. II. Bot. I : 502. 1880. *Diplazium jerdoni* Bedd. Ferns Brit. India t. 327. 1869.

Distrib. : India (Himalayas from Nepal to Bhutan, Meghalaya) Bangladesh.

D. squamigerum (Mett.) Christ, Bull. Soc. Fr. 52. Mem. 1 : 51. 1905. *Asplenium squamigerum* Mett. Ann. Lugd. Bat. 2 : 239. 1866 ; Hope, Journ. Bomb. Nat. Hist. Soc. 14 : 245. 1902.

Distrib. : India (Kashmir), Chain, Japan.

D. stenochlamys C. Chr. Ind. Fil 240. 1905 ; Mehra and Bir, Res. Bull. (N.S.) Punjab Univ. 15 : 147. 1964. *Asplenium platyphyllum* Bak. Journ. Bot. 1890 : 264. 1890. (non J. Sm. 1841 ; Mett. 1859).

Distrib. : Eastern Himalayas, Tonkin.

D. stoliczkae Bedd. Ferns Brit. India Suppl. 13. t. 361. 1876 ; Handb. Ferns Brit. India 182. 1883.

Distrib. : India (N. India, Meghalaya).

D. subsinuatum (Hook. & Grev.) Tagawa, Coll. III. Jap. Pterid. 203. 1959 ; *Asplenium subsinuatum* Hook. & Grev. Icon. Fil. 1 : t. 27. 1827.

D. succulentum (Clarke). C. Chr. Ind : Fil. 240. 1905. *Asplenium succulentum* Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 502. t. 64. f. 4. 1880.

Distrib. : India (Eastern Himalayas).

D. sylvaticum (Bory) Sw. Syn. Fil. 92. 1806. Bedd. Handb. Ferns Brit. India 177. 1883. *Callipteris sylvaticum* Bory. Voy. Mars. Afr. 1 : 282. 1804. *Asplenium sylvaticum* (Bory) Presl. Rel. Haenk. 1 : 42. 1825 ; Bedd. Ferns South. India t. 161. 1865.

Distrib. : South India, Burma, Sri Lanka, Malaya Peninsula, Mauritius, Fernando Po, Java, Borneo, Philippines.

D. speciosum Bl. Enum, Pl. Jav. 193. 1828 ; Bedd. Handb. Ferns Brit. India 178. 1883. *Asplenium acuminatum* Wall. ex Bedd. Ferns Brit. India t. 290. 1868.

Distrib. : South India, Malay Peninsula.

D. torrentium (Clarke) Tard. - Blot. Aspl. du. Tonkin 69. 1932 ; Mehra and Bir, Res. Bull. N.S.) Punjab Univ. 15 : 147. 1964. *Asplenium torrentium* Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 500 t. 54. (excl. f. 2) 1880.

Distrib. : India (Sikkim).

D. travancoricum Bedd. Handb. 188. 1883 ; Sledge Bull. Brit. Mus. (N.H.) London 2 : 305. 1962.

Distrib. : S. India (Travancore Hills), Sri Lanka.

D. viridissimum Christ, Not. Syst. 1 : 45. 1909. Mehra and Bir, Res. Bull. (N.S.) Punjab Univ. 15 : 147. 1964. *Athyrium macrosorum* Copel. Leaflet Philip. Bot. 3 : 815. 1910.

Distrib. : India (Sikkim), China.

Dryoathyrium Ching, Bull. Fan Mem. Inst. Biol. Bot. 11 : 79. 1941.

Type : **D. boryanum** (Willd.) Ching, Bull. Fan Mem. Inst. Biol. Bot. 11 : 81. 1941.

D. boryanum (Willd.) Ching, Bull. Fan Mem. Inst. Biol. Bot. 11 : 81. 1941. *Aspidium boryanum* Willd. in Linn. Sp. Pl. 5 : 285. 1810. *Lastrea divisa* (Hook.) Bedd. Ferns South. India 35. t. 97. 1863. *Polypodium subtripinnatum* Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 545. 1880. *Phegopteris kingii* Bedd. Handb. Ferns Brit. India Suppl. 84. 1892. *Lastrea boryana* (Willd.) Moore, Ind. Fil. 86. 1859 ; Bedd. Handb. Ferns Brit. India 266. 1883.

Distrib. : India (Himalayas, S. India), East Africa to tropical Asia.

Gymnocarpium Newm. Phytol. 4 : 321. 1821.

Lectotype : **G. dryopteris** (L.) Newm. Phytol. 4 : 371. 1851.

G. dryopteris (L.) Newm. Phytol. 4 : 371. 1851. *Polypodium dry-*

opteris L. Sp. Pl. 2 : 1053 ; Bedd. Ferns Brit. India t. 74. 1865. *Phegopteris dryoptesis* (L.) Bedd. Handb. Ferns Brit. India 293. 1883.

Distrib. : India (N. W. Himalayas) Arctic and Alpine Europe, Asia and America.

Kuniwatsukia Pic. Ser. Webbia 28 : 455. 1975.

Type : *K. cuspidata* (Bedd.) Pic. Ser. Webbia 28 : 455. 1973.

. *K. cuspidata* (Bedd.) Pic. Ser. Webbia 28 : 454. 1973. *Lasterea cuspidata* Bedd. Ferns Brit. India 2 : III, IV et correct. 348. 1870 ; Handb. Ferns Brit. India 232. t. 119. 1883. *Polypodium elongatum* Wall. ex Hook. Sp. Fil. 4 : 234. 1862 ; Bedd. Ferns Brit. India t. 118. 1865. *Microchlaena cuspidata* (Bedd.) Ching, Acta Phytotax. Sin. 9 : 99. 1964.

Distrib. : India (Himalayas), S.W. China, Nepal, Burma.

Lunathyrium Koidz. Acta Phytotax. Geobot. 1 : 30. 1932.

Type : *L. pycnosorum* (Christ) Koidz. Acta Phytotax. Geobot. 1 : 31. 1932.

L. allantoides (Bedd.) Ching, Acta Phytotax. Sin. 9 : 72. 1964. *Athyrium allantoides* Bedd. Ferns Brit. India t. 221. 1867. *A. thelypteroides* (Michx.) Desv. Prod. 226. 1827 ; Bedd. Handb. Ferns Brit. India 164. 1883. *Asplenium thelypteroides* Michr. Fl. bor. Arn. 2 : 265. 1803 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 488. 1880.

Distrib. : Himalayas from Kashmir to Bhutan, N. America, Canada, Amurland.

L. japonicum (Thunb.) Kurata, Journ. Geobot. 9 : 99. 1961. *Asplenium japonicum* Thunb. Fl. Jap. 334. 1784. *Athyrium japonicum* (Thunb.) Copel. Philip. Journ. Sci. 3C : 290. 1908. *Diplazium japonicum* (Thunb.) Bedd. Ferns Brit. India Suppl. 12. 18/6 ; Handb. Ferns Brit. India 180. 1883. *D. decussatum* Bedd. Ferns Brit. India 292. t. 1868.

Distrib. : India (S. India, N.W. Himalayas), China, Japan, Malaya Peninsula, Polynesia.

Triblemma (J. Sm.) Ching, Acta Bot. Sin. 11 : 24. 1978.

Type : *T. lancea* (Thunb.) Ching, Acta Bot. Sin. 11. 24. 1978.

T. lancea (Thunb.) Ching, Acta Bot. Sin. 11 ; 24. 1978. *Asplenium lanceum* Thunb. Fl. Jap. 333. 1784. *Diplazium lanceum* (Thunb.) Presl, Tent. Pterid. 113. 1836 ; Bedd. Ferns South. India t. 227. 1867 ; Handb. Ferns Brit. India 174. t. 84. 1883.

Distrib. : E. India, Bangladesh, Nepal, China, Formosa, Japan, Indo-China.

HYPODEMATACEAE Ching

Hypodematuim Kunze, Fl. 1833 (2) : 690. 1833.

Type : **H. crenatum** (Forsk.) Kuhn, v. Deck. Reis. 3 (3) : 37. 1879.

H. crenatum (Forsk.) Kuhn, v. Deck. Reis. 3 (3) : 37. 1879. *Poly-podium crenatum* Forsk, Fl. Aég. arab. 185. 1775. *Lastrea crenata* Bedd. Ferns Brit. India Suppl. 18. 1876 ; Handb. Ferns Brit. India 258. 1883. *L. eriocarpa* Presl, Tent. Pterid. 77. 1836 ; Bedd. Ferns South. India t. 95. 1863.

Distrib. : India (Himalayas, S. India), Sri Lanka, Trop. Africa, Mauritius, China, Malay Peninsula.

ASPIDIACEAE Mett. ex Frank

Acrorumohra (Ito) Ito in Nakai et Honda, Nov. Fl. Jap. 4 : 101. 1939.

Type : **A. diffracta** (Bak.) Ito in Nakai et Honda, Nov. Fl. Jap. 4 : 101. 1939.

A. diffracta (Bak.) Ito in Nakai et Honda, Nov. Fl. Jap. 4 : 101. 1939. *Nephrodium diffractum* Bak. Kew Bull. 230. 1898.

Distrib. : Eastern Himalayas, Southern Asia.

Ctenitis (C. Chr.) C. Chr. ex Tard. et. C. Chr. Notul. Syst. Paris 7 : 186. 1938 ; Ching, Bull. Fan Mem. Inst. Biol. 8 : 275. 1938.

Lectotype : **C. submarginalis** (Langsd. et Fisch.) Ching, Sunyat. 5 : 250. 1940.

C. apiciflora (Wall. ex Mett.) Ching, Bull. Fan Mem. Inst. Biol. 8 : 284. 1938. *Aspidium apiciflorum* Wall ex Mett. Farnatt Pheg. et. Aspid. 54. no. 128. 1858. *Lastrea filix-mas* var. *patentissima* subvar. *apiciflora* Bedd. Handb. Ferns Brit. India 249. 1883.

Distrib. : Himalayas, Nepal, Bhutan, China.

C. clarkei (Bak.) Ching, Bull. Fan Mem. Inst. Biol. 8 : 287. 1938. *Nephrodium clarkei* Bak. Hook. et. Bak. Syn. Fil. 497. 1874. *Lastrea filix-mas* var. *clarkei* Bedd. Ferns Brit. India Suppl. 17. 1876.

Distrib. : India (Himalayas), China, Burma, Sikkim.

C. ferruginea (Bedd.) Ching, Bull. Fan Mem. Inst. Biol. 8 : 296. 1938. *Lastrea ferruginea* Bedd. Ferns South. India t. 100. 1865 ; Handb. Ferns Brit. India 284. t. 136. 1883.

Distrib. : S. India, Sri Lanka.

C. hendersonii (Bedd.) H. Ito ex Nakai et Honda, Nov. Fl. Japan 4 : 89. t. 5. (106), 10 : 212, 14 : 311. to 18 : 423. 1939. *Lastrea hendersonii* Bedd. Ferns Brit. India Suppl. t. 377. 1876. *L. spectabilis* Bedd. Handb. Ferns Brit. India 257. 1883.

Distrib. : India (Meghalaya), Nepal.

C. manii (Hope) Ching, Bull. Fan Mem. Inst. Biol. 8 : 289. 1938. *Nephrodium manii* Hope, Journ. Bot. 145. 1890. *Lastrea intermedia* Bedd. Ferns Brit. India Corr. II. 1870 ; Handb. Ferns Brit. India Suppl. 61. 1892 (*pro parte*).

Distrib. : India, Malaya Peninsula, Java, Philippines, Borneo.

C. nidus (Clark), Ching, Bull. Fan Mem. Inst. Biol. Bot. 8 : 287. 1938 ; *Nephrodium nidus* Clarke, Journ. Linn. Soc. 15 : 156. 1876. *N. apiciflorum* var. *nidus* (Clarke) Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 518. 1880. *Lastrea filix-mas* var. *nidus* Bedd. Ferns Brit. India Suppl. 171. t. 372. 1876. *L. filix-mas* var. *parallelogramma* subvar. *nidus* Bedd. Handb. Ferns Brit. India t. 249. t. 129. 1883.

Distrib. : India (Himalayas, Sikkim), China.

C. obtusiloba (Bak. ex Hook. et Bak.) Ching, Bull. Fan Mem. Inst. Biol. 8 : 296. 1938. *Nephrodium obtusilobum* Bak. ex Hook. et Bak. Syn. Fil. 284. 1867. *Lastrea obtusiloba* Bedd. Ferns Brit. India t. 296. 1868.

Distrib. : S. India, Sri Lanka.

C. rhodolepis (Clarke) Ching, Bull. Fan Mem. Inst. Biol. 8 : 300. 1938. *Nephrodium rhodolepis* Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 526. t. 72. 1880. *Lastrea intermedia* var. *Y. rhodolepis* (Clarke) Bedd. Handb. Ferns Brit. India Suppl. 62. 1892. *L. blumei* Bedd. Ferns South. India t. 249. 1864 ; Handb. Ferns Brit. India 259. 1883. *L. intermedia* var. *blumei* Bedd. Handb. Suppl. 61. 1892.

Distrib. : Himalayas, Sri Lanka, Bangladesh, China, Japan, Formosa, Fiji, Samoa, Phillipines.

C. scabrosa (Kunze) Ching, Bull. Fan Mem. Inst. Biol. 8 : 292. 1938. *Aspidium scabrosum* Kunze, Linnaea 24 : 286. 1856. *Lastrea scabrosa* Moore. Ind. Fil. 103. 1858 ; Bedd. Handb. Ferns Brit. India 263. t. 125. 1883. *Polypodium nigrocarpum* Bedd. Ferns South. India t. 169. 1865.

Distrib. : S. India.

Ctenitopsis Ching ex Tard. et C. Chr. Notul. Syst. Paris 7 : 86. 1938 ; Ching. Bull. Fan Mem. Inst. Biol. Bot. 8 : 304. 1938.

Lectotype ; **C. sagenoides** (Mett.) Ching, Bull. Fan Mem. Inst. Biol. 8 : 312. 1938.

C. dissecta (G. Forst.) Ching, Bull. Fan Mem. Inst. Biol. 8 : 321. 1938.
Polypodium dissectum G. Forst. Prod. 81. 1786. *Lastrea dissecta* (G. Forst.) Bedd. Handb. Ferns Brit. India 260. 1883. *L. membranifolia* Bedd. Ferns South. India t. 102. 1865.

Distrib. : S. India, Sri Lanka, Burma, Malay Islands, Polynesia.

C. fuscipes (Wall. ex Bedd.) Ching, Bull. Fan Mem. Inst. Biol. Bot. 8 : 313. 1938. *Aspidium fuscipes* Wall. List. no. 361. (*ex parte*), 1827 (*nom. nud.*) ex Bedd. Ferns Brit. India Suppl. 15. t. 366. 1876. *Lastrea fuscipes* Moore, Ind. Fil. 82. 1858 ; Bedd. Handb. Ferns Brit. India 243. 1883 (*pro parte*) *Tectaria fuscipes* C. Chr. Contr. U. S. Nat. Herb. 26 : 290. 1931. *Nephrodium membranifolium* Hook. Sp. Fil. 131. t. 261. 1862 (*pro parte*) ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 534. 1880. *N. membranifolium* var. *typica* Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 534 t. 75 A. 1880. *N. membranifolium* var. *dimorphous* Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 534. t. 75 BC. 1880. *Aspidium membranifolium* Bedd. Handb. Ferns Brit. India 225. 1883. Suppl. 48. 1892 (*excl. syn.*).

Distrib. : India (Eastern India and Eastern Himalayas), Burma, Annam.

C. ingens (Atk. ex Clarke), Ching, Bull. Fan Mem. Inst. Biol. 8 : 320. 1938. *Nephrodium ingens* Atk. ex Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 526. t. 73. 1880. *Lastrea dissecta* var. *ingens* (Atk. ex Clarke). Bedd. Handb. Ferns Brit. India 261. 1883.

Distrb. : India (Himalayas), Bhutan.

C. sagenoides (Mett.) Ching, Bull. Fan Mem. Inst. Biol. Bot. 8 : 312. 1938. *Aspidium sagenoides* Mett. Farnatt. Pheg. u. Aspid. 139. n. 269. 1858. *Lastrea melanopus* Bedd. Ferns Brit. India t. 38. 1865. *L. fuscipes* Bedd. Handb. Ferns Brit. India 261. 1883 (*pro parte*).

Distrb. : India (Eastern India & Eastern Himalayas), Burma, Siam, China, Cochin-China, Malaya & its islands, Java, Sumatra, Philippine Islands.

Cyrtomium Presl, Tent. Pterid. 84. t. 2. f. 19. 1836 ; C. Chr. Amer. Fern. Journ. 20 (2) : 41. 1930.

Lectotype : **C. falcatum** (L. f.) Presl, Tent. Pterid. 86. 1836.

C. caryotideum (Wall. ex Hook. et Grev.) Presl, Tent. Pterid. 86. t. 2. f. 26. 1836 ; Bedd. Ferns South. India t. 119. 1865. *Aspidium caryotideum* Wall. List. No. 376. 1828 (*nom. nud.*) ex Hook. et Grev. Icon. Fil. t. 69. 1828. *Cyrtomium falcatum* var. *caryotideum* (Wall. ex Hook. et Grev.) Bedd. Handb. Ferns Brit. India 211. t. 106. 1883. *Aspidium falcatum* Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 512. 1880 (*pro parte*).

Distrib. : Mountainous regions of India. Nepal, Bhutan, Sri Lanka, China, Japan, S. Africa, Sandwich Islands, Madagascar.

C. falcatum (L. f.) Presl, Tent. Pterid. 86. 1836 ; Bedd. Handb. Ferns Brit. India 211. 1883; C. Chr. Amer. Fern. Journ. 20: 47. 1930. *Polypodium falcatum* L. f. Suppl. Sp. Pl. 466. 1781.

Distrib. : Himalayas, China, Japan, Formosa, Asia.

C. fortunei J. Sm. Ferns Brit. & For. 286. 1866.

Distrib. : Himalayas, China, Japan.

C. hookerianum (Presl) C. Chr. Ind. Fil. Suppl. 1 : 101. 1913. *Lastrea hookeriana* Presl, Tent. Pterid. 77. 1836. *Aspidium caducum* wall. ex. Hook. et Grev. Icon. Fil. t. 171. 1829. (non H B K 1815). *Cyrtomium caducum* (Wall. ex Hook. et Grev.) Moore, Ind. Fil. 276. 1861 ; Bedd. Ferns Brit. India t. 45. 1865 ; Handb. Ferns Brit. India 211. 1883.

Distrib. : India (Himalayas), China, Japan, Australia.

C. macrophyllum (Makino) Tagawa, Acta Phytotax. Geobot. 3 62. 1936. *Aspidium falcatum* var. *macrophyllum* Makino, Bot. Mag. Tokyo 16 : 90. 1902.

Distrib. : Himalayas, China, Japan.

Didymochlaena Desv. Gen. Naturf. Freunde. Berlin Mag. 5 : 303. 1811.

Type : **D. sinuata** Desv. Ges. Naturf. Freunde Berlin Mag. 5 : 303. t. 7. f. 6. 1811.

D. truncatula (Sw.) J. Sm. Journ. Bot. 4 : 196. 1841. *Aspidium truncatum* Sw. Schrad. Journ. Bot. 1800 (2) : 36. 1801. *Didymochlaena lunulata* sensu Bedd. Ferns Brit. India t. 15. 1865 : Handb. Ferns Brit. India 199. t. 99. 1883 ; Suppl. 42. 1892.

Distrib. : E. India, Burma, Malaya Peninsula, Tropical America, East African Islands ; Malay Islands, Fiji.

Lastreopsis Ching, Bull. Fan Mem. Inst. Biol. Bot. 8 : 161. 1938.

Type : *L. tenera* (R. Br.) Tindale, Vict. Nat. 73 : 181. 1957.

L. tenera (R. Br.) Tindale, Vict. Nat. 73 : 181. 1957. *Nephrodium tenerum* R. Br. Prod. Fl. Nov. Holl. 149. 1819. *Lastrea recedens* J. Sm. ex Moore. Gard. Chron. 708. 1855 ; Bedd. Fern South. India t. 98. 1865 (*pro parte*) ; Handb. Ferns Brit. India 260. 1883.

Distrib. : S. India, Sri Lanka, Philippines.

L. wattii (Bedd.) Tagawa, Journ. Jap. Bot. 22 : 163. 1949. *Lastrea wattii* Bedd. Handb. Ferns Brit. India Suppl. 49. 1892.

Distrlb. : India (Manipur).

Pleocnemia Presl, Tent. Pterid. 183. 1836.

Type : *P. leuzeana* (Gaud.) Presl, Tent. Pterid. 184. t. 7. f. 12. 1836.

P. winitii Holtt. Reinw. 1 : 181. 1951. *P. leuzeana sensu* Bedd. Handb. Ferns. Brit. India 228. 1883 (*descr. pro parte, excl. t. 117*).

Distrib. : India (Assam) Thailand, Annam, China.

Pteridrys C. Chr. et Ching, Bull. Fan Mem. Inst. Biol. 5 : 129. 1934.

Type : *P. syrmatica* (Willd.) C. Chr. et Ching, Bull. Fan Mem. Inst. Biol. Bot. 5 : 131. t. 11, 17. 1934.

P. cnemidaria (Christ) C. Chr. et ching, Bull. Fan Mem. Inst. Biol. Bot. 5 : 136. t. 12, 18. (8-9), 20 (19). 1934. *Dryopteris cnemidaria* Christ, Bull. Acad. Geogr. Bot. 140. 1910. *Nephrodium syrmaticum* Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 ; 518. 1880. (*pro porte*).

Distrib. : India (Assam, Sikkim), Burma, China, Bhutan, Cochin-China.

P. zeylanica Ching ex C. Chr. et Ching, Bull. Fan Mem. Inst. Biol. Bot. 5 : 135. t. 20 f. 16-18. 1934. *Loatrea spectabilis* Bedd. Ferns South. India t. 108. 1863. *Nephrodium syrmaticm sensu* Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 519. 1880. *Lastrea syrmaticum sensu* Bedd. Handb. Ferns. Brit. India 243. 1883.

Distrib. : India (Assam, South India), Sri Lanka.

Quercifilix Copel. Phillip Journ. Sci. 37 : 408. 1928.

Type : **Q. Zeylanica** (Houtt.) Copel. Philip. Journ. Sci. 37 : 408. 1928.

Q. Zeylanica (Houtt.) Copel. Philip. Journ. Sei. 37 : 408. 1928.
Ophioglossum Zeylanicum Houtt. Nat. Hist. 14 : 43. 1783.

Distrib. : South India, Sri Lanka, Malesia to Polynesia, China.

Tectaria Cav. Anales. Hist. Nat. 1 (2) : 115. 1799 ; Ching, Sinensis 2 (2) : 9. 1931.

Type : **T. trifoliatum** (L.) Cav. Descr. 249. 1802.

T. artinexa (Clarke) Ching, Sinensis 2 : 15. 1931. *Nephrodium artinexum* Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 536. 1880. *Pleocnemia clarkei* Bedd. Ferns Brit India, Suppl. 15. t. 368. 1876 ; Handb. Ferns Brit. India 227. 1883.

Disirib. : India (Eastern Himalayas).

T. chattagramica (Clarke) Ching, Sinensis 2 (2) : 35. 1931. *Polypodium chattagramicum* Clarke, Trans. Linn. Soc. Lond. II. Bot. 1. 594. t. 81. 1880. *Dictyopteris chattagramica* Bedd. Handb. Ferns Brit. India 255. t. 155. 1883.

Distrib. : India (Meghalayas), Bangladesh, Burma.

T. cicutaria (L.) Copel. Philip. Journ. Sci. 2C. 410. 1907. *Polypodium cicutarium* L. Syst. Nat. ed. x. 2 : 1326. 1759. *Aspidium cicutarium* (L.) Sw. Schrad. Journ. 1800 (2) : 36. 1801 ; Bedd. Handb. Ferns Brit. India 221. 1883. *Sagenia coadunata* J. Sm. Journ. Bot. 4 : 1841 ; Bedd. Ferns South. India t. 81. 1863.

Distrib. : Throughout the Indian region from the planis upto 5000 ft. also tropics throughout the world.

T. coadunata (Wall. ex Hook. et Grev.) C. Chr. in Contr. U. S. Nat. Herb. 26 : 331. 1931. *Aspidium coadunatum* Wall. ex Hook. et Grev. Icon. Fil. 2 : 202. 1831. *Sagenia variolosa* Bedd. Ferns South. India t. 170. 1823. *S. gigantea* Bedd. Ferns South. India t. 80. 1863. *S. coadunata sensu* Bedd. Ferns South. India t. 81. 1863. *Nephrodium cicutarium* Hook. & Bak. Syn. Fil. 299. 1874 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 532. 1880. *N. cicutarium* var. *coadunatum* Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 540. 1880.

T. decurrens (Presl) Copel. Elmers Leaflets Philip. Journ. Bot. 1 : 234. 1907. *Aspidium decurrens* Presl, Rel. Haenk. 1 : 28. 1825 ; Bedd. Ferns South. India t. 245. 1870 ; Handb. Ferns Brit. India 219. 1883. *Sagenia pteropus* Moore, Ind. Fil. 89. 1858 ; Bedd. Ferns South. India t. 82. 1863. *Nephrodium decurrens* Hook & Bak. Syn. Fil. 299. 1874 ; Clarke. Trans. Linn. Soc. Lond. II. Bot. 1 : 339. 1880.

Distrib. : Eastern to Southern India, Sri Lanka, China, Malay Peninsula, Malay Islands, Polynesia, Formosa, Bangladesh.

T. dubia (Clarke et Bak.) Ching, Sinensis 2 : 23. 1931. *Nephrodium cicutarium* var. *dubia* Clarke et Bak. Journ. Linn. Soc. 24 : 417. 1888. *Aspidium dubium* Bedd. Handb. Ferns Brit. India Suppl. 47. 1892.

Distrib. : India (Assam).

T. gigantea (Bl.) Copel. Philip. Journ. Sci. Bot. 2 : 410. 1907. *Aspidium giganteum* Bl. Enum. Pl. Jav. 159. 1828.

Distrib. : South India, Sri Lanka, Java, Malaya.

T. multicaudata (Wall. ex Bedd.) Ching, Sinensis 2 (2) : 20. 1931. *Aspidium multicaudatum* Wall. ex Bedd. Handb. Ferns Brit. India 222. 1883. *Nephrodium multicaudatum* Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 540. t. 77. 1880. *Aspidium griffithii* Bedd. Ferns Brit. India Suppl. 15. 1876. *Sagenia griffithii* Bedd. Ferns Brit. India t. 377. 1880.

Distrib. : South-East India, Burma, China, Formosa.

T. heterosora (Bak.) Ching, Sinensis 2 : 29. 1931. *Nephrodium heterosorum* Bak. in Syn. Fil. 504. 1874; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 537. 1880. *Sagenia hetero'ora* (Bak.) Hook. Syn. Fil. 504. 18 ; Clarke, Trans. Linn. Soc. Lond II. Bot. 1 : 537. 1880.. *S. heterocarpa* Bedd. Ferns Brit. India t. 47. 1865. *Aspidium heterocarpum* (Bedd.) Bedd. Ferns Brit. India Suppl. 16. 1876 : Handb. Ferns Brit. India 219. 1883. *A. heterosorum* (Bak.) Bedd. Handb. Ferns Brit. India Suppl. 46. 1892.

Distrib. : E. India, Sikkim, Bangladesh.

T. leuzeana (Gaud.) Copel. Philipp. Journ. Sci. 2 C : 417. 1907. *Polypodium leuzeanum* Gaud. Freye. Voy. Bot. 361. t. 6. 1827. *Pleocnemia leuzeana* Presl, Tent. Pterid. 184. 1836 ; Bedd. Ferns Brit. India t. 134. 1886 ; Handb. Ferns Brit. Innia 228. 1883. *Nephrodium leuzeanum* Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 535. 1880.

Distrib. : E. India, Darjeeling & Sikkim Himalayas, Bangladesh, Burma, Malaya Peninsula, Malaya Islands, China, Polynesia, N. Australia.

T. macrodonta (Fée) C. Chr. Ind. Fil. Suppl. III. 181. 1934. *Sagenia*

macrodonta Féé. Gen. Fil. 313. t. 24 A. f. 1. 1852. *Aspidium cicutarium* (L.) Sw. Schrad. Journ. Bot. 1880. (2) : 36. 1801 ; Bedd. Handb. Ferns. Brit. India 220. 1883. *Nephrodium cicutarium* Hook. & Bak. Syn. Fil. 299. 1874 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 539. 1880.

Distrib. : Throughout India and also tropics of the world.

T. melanocaulis (Bl.) Copel. philip. Journ. Sci. 2C : 416. 1907.
Aspidium mclanocaulon Bl. Euam. Pl. Jav. 161. 1828.

Distrib. : India (Andaman & Niobor Islands) Malay Islands, Philippines, New Zealand.

T. polymorpha (Wall. ex Hook.) Copel. Philip. Journ. Sci. 2 C : 413. 1907. *Aspidium polymorphum* Wall. List. no. 382. 1828. ex Hook, Sp. Fil. 4 : 54. 1862. ; Bedd. Ferns South, India t. 116. 1865 ; Handb. Ferns. Brit. India 218. 1883. *Nephrodium polymorphum* Bak. Syn. Fil. 297. 1874. *N. polymorphum* Bak. in Hook. Syn. Fil. 297. 1867 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 537. 1880. (excl. vars.). *Aspidium polymorphum* Bak. in Hook. Syn. Fil. 297. 1867 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 537. 1880. (excl. vars.). *A. polymorphum* Wall. List. no. 382. 1828 ; Bedd. Handb. Ferns Brit. India 218. 1883.

Distrib. : S. India, N.W. Himalayas, Bangladesh, Burma, Sri Lanka, Malay Islands, Philippines.

T. polymorpha var. *cuneifolia* R. Bonap. Notes pterid. 14 : 50. 1923.

Disirib. : India (North Andamans), S. China. Burma, Malaysia, Malay peninsula, Indonesia, Philippines.

T. simonsii (Bedd.) Ching. Sinensis 2 : 32. 1931. *Aspidium simonsii* Bedd, Ferns Brit. India Suppl. 15 t. 367. 1876 ; Handb. Ferns Brit. India 48. 1883. *Nephrodium polymorphum* var. *simonium* Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 537. 1880.

Distrib. : India (Eastern & Sikkim Himalayas), Bangladesh, Bhutan.

T. subconfluens (Bedd.) Ching, Sinensis 2 : 27. 1931. *Aspidium subconfluens* Bedd. Ferns Brit. India Suppl. t. 364. 1876 ; Handb. Ferns Brit. India 215. t. 109. 1883. *Nephrodium subconfluens* Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 536. 1880,

Distrib. : India (Meghalaya), China, Bangladesh.

T. subtriphylla (Hook. & Arn.) Copel. Philp. Journ. Sci. Bot. 2 C : 410. 1907. *Polypodium subtriphyllum* Hook. & Arn. in Bot. Capt. Beech. Voy. 256. t. 50. 1848. *Sagenia subtriphylla* Bedd. Ferns South. India t. 242. 1863.

Distrib. : South India, China, Formosa.

T. ternifolia (v. A. v. R.) C. Chr. Ind. Fil. suppl. III. 185. 1934.
Aspidium ternifolium v. A. v. R. Bull. Jard. Brit. no. 11 : 3. 1913.

Distrib. : India (S. Andamans) Malaya, N. Sumatra.

T. trimeni (Bedd.) C. Chr. Ind. Fil. Suppl. 3 : 185. 1934. *Aspidium trimeni* Bedd. Handb. Ferns Brit. India 224. t. 114. 1883. *Sagenia gigantea sensu* Bedd. Ferns South. India t. 80. 1863 (*pro parte*).

Distrib. : S. India, Sri Lanka, Malesia.

T. variolosa (Wall. ex Hook.) C. Chr. Contr. U. S. Nat. Herb. 26 : 289. 1931. *Aspidium variolosum* Wall. ex Hook. Sp. Fil. 4 : 51. 1862 ; Bedd. Ferns Brit. India Suppl. t. 365. 1876, (non t. 170. 1878) ; Handb. Ferns Brit. India 216. t. 111. 1883 ; Suppl. 45. 1892. *Aspidium zollingerianum* Bedd. Ferns Brit. India t. 251. 1868.

Distrib. : East Bengal plains, abundant from Assam to Chittagong ; Burma, Malaya Peninsula, Formosa.

T. vasta (Bl.) Copel. Philip. Journ. Sci. 2 C : 411. 1907. *Aspidium vastum* Bl. Enum. Pl. Jav. 142. 1828 ; Bedd. Ferns Brit. India Suppl. 15. 1876 ; Handb. Ferns Brit. India 213. t. 108. 1883. *Nephrodium vastum* Hook. & Bak. Syn. Fil. 296. 1874 ; Clarke, Trans Linn. Soc. Lond. II. Bot. 1 : 536. 1880. *Sagenia alata* Bedd. Ferns. Brit. India t. 169. 1866.

Distrib. : E. India, Bangladesh, Malaya Peninsula and Malay Islands.

T. wightii (Clarke) Ching, Sinensis 2 : 28. 1931. *Nephrodium wightii* Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 538. 1880. *Aspidium polymorphum* var. *macrocarpum* Bedd. Ferns South. India t. 177. 1865.

Distrib. : S. India, Assam, Bhutan.

PERANEMTACEAE (Presl) Ching

Acrophorus Presl, Tent. Pterid. 93. t. 3. f. 2 : 1836.

Type : **A. nodosus** Presl, Tent. Pterid. 94. t. 3. f. 2. 1836.

A. paleolatus Pic. Ser. Webbia 31 (1) : 252. 1977. *Acrophorus stipellatus* auct. *pro parte* quoad pl. asiat. (non T. Moore, 1854). *Leucostegia nodosa* sensu Bedd. Ferns Brit. India t. 93. 1865 ; Handb. Ferns Brit. India 54. t. 26. 1883 (*pro parte*).

Distrib. : India (Himalayas), Asia.

Diacalpe Bl. Enum. Pl. Jav. 241. 1828.

Type : **D. aspidioides** Bl. Enum Pl. Jav. 241. 1828.

D. aspidioides Bl. Enum. Pl. Jav. 241. 1828 ; Bedd. Ferns South. India t. 257. 1865 : Handb. Ferns Brit. India 18. t. 7. 1883. *Peranema aspidioides* Bl. Enum. Pl. Jav. 241. 1828.

Distrib. : India (Himalayas), Nepal to Bhutan, Burma, China, Malay Islands, Philippines, New Guinea, Sri Lanka.

Lithostegia Ching, Sinensis 4 : 2. 1933.

Type : **L. foeniculacea** (Hook.) Ching, Sinensis 4 : 2. 1933.

L. foeniculacea (Hook.) Ching, Sinensis 4 : 2 : 1933. *Aspidium foeniculaceum* Hook. Sp. Fil. 4 : 36. t. 237. 1862. *Diacalpe foeniculacea* (Hook.) Clarke, Trans. Linn. Soc. Lond. II Bot. 1. 434. 1880 ; Bedd. Handb. Ferns Brit. India 18. 1883. *Lastrea foeniculacea* Bedd. Ferns Brit. India t. 36. 1865.

Distrib. : India (Eastern Himalayas), Burma, China.

Peranema D. Don, Prod. Fl. Nepal 12. 1825.

Type : **P. cyatheoides** D. Don, Prod. Fl. Nepal 12. 1825.

P. aspidioides (Bl.) Mett. Fil. Lechl. 2 : 33. 1850. *Diacalpe aspidioides* (Bl.) Enum. Pl. Jav. 241. 1828 ; Bedd. Handb. Ferns Brit. India 18. t. 7. 1883.

Distrib. : Eastern India and Eastern Himalayas, Nepal, Bhutan, Burma, Sri Lanka, Vietnam, Sumatra, Java, Borneo, New Guinea, Philippines.

P. cyatheoides D. Don, Prod. Fl. Nepal 12. 1825 ; Bedd. Ferns South. India t. 73. 1863 ; Handb. Ferns Brit. Indta 22. 1883.

Distrib. : Himalayas and S. India, Nepal, Bhutan, Formosa.

DRYOPTERIDACEAE (Ching) Ching

Arachniodes Bl. Enum. Pl. Jav. 2 : 241. 1828 ; Sledge, Bull. Brit. Mus. (N. H.) Lond. 5 : 39. 1973.

Type : **A. aspidioides** Bl. Enum Pl. Jav. 2 : 242. 1828.

A. amabilis (Bl.) Tindale, Contr. N. S. Wales Nation Herb. 3 : 90. 1961. *Aspidium amabile* Bl. Enum. Pl. Jav. 165. 1828. *Lastrea amabilis* Moore, Ind. Fil. 85. 1858 ; Bedd. Ferns South. India t. 109 ; Handb. Ferns Brit. India 229. 1883.

Distrib. : Throughout India in hilly regions, Sri Lanka, Nepal, S. China, Philippines.

A. aristata (Forst. f.) Tindale, Contr. N. S. Wales Nation Herb. 3 : 89. 1961. *Polypodium aristatum* Forst. Prod. 82. 1786. *Lastrea aristata* Moore, Ind. Fil. 85. 1858 ; Bedd. Ferns South. India t. 101. 1863 ; Handb. Ferns Brit. India 229. 1883. *Aspidium aristatum* Clarke. Trans. Linn. Soc. Lond. II. Bot. 1 : 511. 1880. (*excl. vars.*).

Distrib. : S. India, Burma, Sri Lanka, China, E. Australia, Polynesia.

A. assamica (Kuhn) Ohwi, Journ. Jap. Bot. 37 : 76. 1962. *Aspidium assamicum* Kuhn, Linnaea 36 : 108. 1869.

Distrib. : India (Himalayas, Assam), Burma, China, Japan, Tonkin, Thailand.

A. carvifolia (Kze.) Ching, Acta Bot. Sin. 10. 256. 1962. *Aspidium carvifolium* Kze. Linnaea 24 : 292. 1851.

Distrib. : India (Himalayas), China.

A. coniifolia (Moore) Ching, Acta Bot. Sin. 10 : 257. 1962. *Lastrea coniifolia* Moore, Ind. Fil. LXXXVIII. 1857.

Distrib. : India (Eastern Himalayas,) China.

A. nipponica (Rosenst.) Ohwi, Jap. Bot. 37 : 76. 1962. *Polystichum nipponicum* Rosenst. in Feddes Repert. 13 : 130. 1914.

Distrib. : India (Himalayas), Burma, China, Japan.

A. simulans (Ching) Ching, Acta Bot. Sin. 10 : 259. 1962. *Rumohra simulans* Ching, Sinensis 5 : 54. t. 8. 1934.

Distrib. : India (Himalayas), China.

A. speciosa (D. Don) Ching, Acta Bot. Sin. 10 : 259. 1962. *Aspidium speciosum* D. Don, Prod. Fl. Nepal 5. 1825. *Lastrea affinis* Wall ex Bedd. Handb. Ferns Brit. India 230. 1883.

Distrib. : India (N.E. Himalayas), Nepal, China, Tibet.

A. spectabilis (Ching) Ching, Acta Bot. Sin. 10 : 259. 1962. *Rumohra spectabilis* Ching, Sinensis 5 : 58. t. 11. 1934.

Distrib. : India (Himalayas), W. China, Burma, Thailand.

A. tripinnata (Goldm.) Sledge, Bull. Brit. Mus. (N.H.) 5 : 41. 1973.
Polystichum tripinnatum Goldm. Nova Acta Acad. Gesar. Leop. Carol. 19. Suppl. 1 : 463. 1843. *Lastrea coniifolia* sensu Bedd. Ferns Brit. India t. 261. 1868. (excl. fig., non *Aspidium coniifolium* wall. ex Mett.) ; Handb. Ferns Brit. India 230. f. 118. 1883.

Distrib. : South India, Java, Sarawak, Philippines.

Dryopteris Adanson, Fam. Pl. 2 : 20. 551. 1763 ; Ching Bull. Fan Mem. Inst. Biol. Bot. 8 : 363. 1938. *nom. cons.*

Type : **D. filix-mas** (L.) Schott, Gen. Fil. ad. t. 9. 1834.

D. acuto-dentata Ching, Bull. Fan Mem. Inst. Biol. Bot. 8 : 432. 1938.
Nephrodium kingii Hope, Journ. Bomb. Nat. Hist. Soc. 12 : 621. t. 9. 1899; 14 : 731. 1903 (non *Dryopteris kingii* C. Chr. 1905).

D. angustifrons (Moore ex Bedd.) O Kzte. Rev. Gen. Pl. 2. 812. 1891.
Lastrea angustifrons Moore, ex Bedd. Ferns Brit. India t. 226. 1867 : Handb Ferns Brit. India t. 262. 1883. *Nephrodium splendens* var. *angustifrons* Hook. Sp. Fil. 4 : 126. 1862. *N. angustifrons* Bak. Syn. Fil. 283. 1867.

Distrib. : India (Sikkim,) Nepal, China.

D. apiciflora (Wall. ex Mett.) O. Knutze, Rev. Gen. Pl. 2 : 812. 1891.
Aspidium apiciflorum Wall. ex Mett. Pheg. 4. Asp. 54. n. 128. 1858. *Lastrea apiciflora* (Wall. ex Mett.) Presl, Tent. Pterid. 76. 1836 ; Bedd. Handb. Ferns Brit. India 249. 1883.

Disirib. : India (Himalayas), Formosa.

D. approximata Sledge, Bull. Brit. Mus. (NH.) Bot. 5 : 11. f. 1. 1973.

Distrib. : S. India, Sri Lanka.

D. assamensis (Hope) C. Chr. et Ching, Bull. Dept. Biol. Coll. Sci. Sun. Yatsen Univ. 6 : 4. 1933. *Lastrea filix-mas* var. *assamensis* Hope, Journ. Bot. 1890 : 327. 1890.

Distrib. : India (Assam).

D. atrata (Wall. ex Kunze) Ching, sinensis 3 : 326. 1933. *Aspidium atratum* Wall ex Kunze, Linhaea 24 ; 279. 1851. *Nephrodium gamblei* Hope, Journ. Bomb. Nat. Hist. Soc. 12 : 533, t. 7. 1895. *Dryopteris gamblei* (Hope) C. Chr. Ind. Fil. 267. 1906. *Lastrea hirtipes* Bedd. Handb. Ferns Brit. India 232. 1883 (excl syn.) *Nephrodium hirtipes* Hook. Sp. Fil. 4 : 115. t. 249. 1862. (excl. Syn. *Asp. hiritipes* Bl.) ; Clarke, Trans. Linn. Soc. Lond. II. Bot. I : 513. 1880 (*pro parte*).

Distrib. : India (Himalayas), Nepal, Sri Lanka, Burma, China, Formosa, Japan, Tibet.

D. barbigera (Hook.) O. Ktze. Rev. Gen. Pl. 2 : 812. 1891. *Nephrodium barbigerum* Hook. Sp. Fil. 4 : 113. 1862 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 522. 1880. *Lastrea barbigera* Moore ex Bedd. Ferns Brit. India t. 227. 1867 ; Handb. Ferns Brit. India 246. 1883. *Nephrodium barbigerum* Hook. Sp. Fil. 4 : 113. 1862 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 522. 1880. *Lastrea falconeri* Bedd. Ferns Brit. India t. 41. 1865.

Distrib. : India (Himalayas), China, Formosa.

D. basiaurita Ching, Bull. Fan Mem. Inst. Biol: Bot. 8 : 405. 1938.

Distrib. : India (Kashmir).

D. blanfordii (Hope) C. Chr. Ind. Fil. 254. 1905. *Nephrodium blanfordii* Hope, Journ. Bomb. Nat. Hist. Soc. 12 : 624. t. 11. 1899. *Nephrodium remotum* Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 523. 1880. *Lastrea spinulosa* Bedd. Ferns Brit. India 336. 1870 (*non* Presl 1836). *Lastrea spinulosa* var. *remota* (Clarke) Bedd. Handb. Ferns Brit. India 252. 1883 ; Suppl. 59. 1892.

Distrib. : India (North-West to Darjeeling Himalayas).

D. chingii Nair, Indian For. 94 : 169. 1968. *Nephrodium filix-mas* (L.) Rieh. var. *fibrillosa* Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 520. t. 70. 1880. *Dryopteris fibrillosa* (Clarke) Hand-Mazz. Akad. Anz. Akad. Wien. no 72. 1922, (*non* Christ 1905.) ; 15 : 131. 1964. *Lastrea filix-mas* (L.) Presl var. *parallelogramme* (Kunze) Hook. Subvar. *fibrillosa* (Clarke) Bedd. Handb. Ferns Brit. India 250. 1883.

Distrib. : India (North-West Himalayas).

D. chrysocoma (Christ) C. Chr. Ind. Fil. 257. 1905. *Aspidium filix-mas* var. *chrysocoma* Christ, Bull. Herb. Boiss. 6 ; 966. 1898. *A. chrysocoma* Christ, Bull. Acad. Géogr. Bot. II : 253. 1902. *Nephrodium filix-mas* var. *schimperiana* Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 ; 520. t. 69. f. 2. 1880. *Lastrea intermedia* Bedd. Ferns, South. India t. 113. 1863 (*non* presl, 1836). *L. filix-mas* var. *elongata* Bedd. Handb. Ferns Brit. India 250. 1883 (*pro parte*). *L. filix-mas* var. *schimperiana* Bedd. Handb. Ferns Brit. India Suppl. 58. 1892. *Nephrodium schimperianum* Hope, Journ. Bomb. Nat. Hist. Soc. 14 : 733. 1903 (*non* *Aspidium* Hochst. 1841).

Distrib. : India (Himalayas, South India), China, Nepal, Bhutan.

D. chrysocoma (Christ). C. Chr. var. **major** Ching, Bull. Fan Mem. Inst. Biol. Bot. 8 : 438. 1938.

Distrib. : India (North-West Himalayas), China.

D. Clarkei (Bak.) O. Ktze, Rev. Gen. Pl. 2 : 812. 1891. *Nephrodium Clarkei* Bak. Syn. Fil. 497. 1874 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 518. 1880. *Lastrea filix-mas* var. *clarkei* Bedd. Ferns Brit. India Suppl. 17. t. 371. 1876 ; Handb. Ferns Brit. India 250. 1883.

Distrib. : India (Sikkim).

D. coeruleata (D. Don.) C. Chr. Ind. Fil. 258. 1905. *Nephrodium coeruleatum* D. Don, Prod. Fl. Nepal 6. 1825 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 521. 1880. *Lastrea filix-mas* Presl. var. *S. coeruleata* (D. Don.) Bedd. Handb. Ferns Brit. India 250. 1883.

Distrib. : India (Himalayas), Burma, Malaya Peninsula, China, Tonkin, Nepal.

D. costalisora Tagawa, Act. Phytotax. Geobot. 3 : 88. 1934.

Distrib. : India (Himalayas), Formosa

D. deparioides (More) O. Ktze. Rev. Gen. Pl. 2 : 812. 1891. *Diclisodon deparioides* Moore, Ind. Fil. XCVI. 1857. *Lastrea deparioides* Bedd. Ferns South. India 104. 1865 ; Handb. Ferns Brit. India 257. 1883.

Distrib. : S. India, Sri Lanka, Australia.

D. deparioides (Moore) Ktze, var. **concinna** C. Chr. Ind. Fil. Suppl 3 : 84. 1934 ; *Lastrea concinna* Bedd. Ferns Brit. India t. 207. 1864 (*non* *L. concinna* (Willd.) Moore).

Distrib. : S. India, Sri Lanka.

D. filix-mas (L.) Schott, Gen. Fil. ad. t. 9. 1834. *Polypodium filix-mas* L. sp. pl. 2 : 1090. 1753. *Lastrea filix-mas* Presl, Tent. Pterid. 76. 1836.

Distrib. : Cosmopolitan.

D. filix-mas var. **kingii** C. Chr. Ind. Fil. 265. 1905,

Distrib. : North-West Himalayas, Sikkim, Tibet.

D. fructuosa (Christ) C. Chr. Ind. Fil. 267. 1905. *Aspidium varium* var. *fructuosum* Christ, Bull. Soc. Fr. 52. Mem. 1 : 38. 1905.

Distrib. : India (Sikkim), China (Yunnan).

D. hasseltii (Bl.) C. Chr. Ind. Fil. 269. 1905. *Polypodium hasseltii* Bl. Fl. Jav. 195. t. 92. 1829.

Distrib. : Eastren Himalayas, Tonkin, China, Formosa, Ryukyu, Philippines, Malaya Islands, New Guinea.

D. hexagonopetra (Michx.) C. Chr. Ind. Fil. 270. 1905. *Polypodium hexagonopetrum* Mishx. Fl. Bor. Am. 2 : 271. 1803.

Distrib. : India (N. W. Himalayas), America.

D. hypophlebia Hayata, Icon. Pl. Formosa 4 : 154. f. 96. 1914.

Distrib. : India (Himalayas), Formosa.

D. gracilis (Moore) Ching, Bull. Fan Mem. Inst. Biol. 8 : 455. 1938. *Lastrea gracilis* Moore ex Bedd. Ferns Brit. India t. 198. 1866. *Aspidium thomsonii* var. *gracilis* Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 508. 1880. *Polystichum thomsonii* var. *gracilis* Bedd. Handb. Ferns Brit. India 207. 1883.

Distrib. : India (Sikkim).

D. gracillima Ching, Bull. Fan Mem. Inst. Biol. Bot. 8 : 452. 1938. *Nephrodium sparsa* var. *minor* Bak. Syn. Fil. ed. 2. 458. 1874 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 524. 1880. *Lastrea filix-mas* var. *minor* Bedd. Handb. Ferns Brit. India 254. 1883.

Distrib. : India North-West Himalayas. Sri Lanka.

D. kawakamii Hayata, Mater, Fl. Formos. 416. 1911.

Distrib. : India (Himalayas), Formosa.

D. lachoongensis (Bedd.) Nayar. & Kaur, Comp. to Bedd. Handb. 61 1974. *Lastrea filix-mas* var. *lachoongensis* Bedd. Handb. Ferns Brit. India Suppl. 58. 1892.

Distrib. : India (Sikkim).

D. lepidopoda Hayata, Icon. Pl. Formos. 4 : 161. f. 101. 1914.

Distrib. : India (Himalayas), Formosa.

D. macrocarpa Stewart, Bull, Torrey Bot. Club. 72 : 406. 1945. *Lastrea filix-mas* var. *elongata* sensu Bedd. Handb. Ferns Brit. India 250. 1883. non *D. elongata* (Wall. ex Hook.) Kze.

Distrib. : India (Himalayas, S. India), Sri Lanka.

D. marginata (Wall ex Christ) Christ, Philip. Journ. Sci. 2 : 212. 1907. *Aspidium marginatum* Wall. List No. 391. 1828. (*pro parte*) ex Christ. Bull.

Soc. Bot. France 52. Mém. 1 : 39. 1905. *Nephrodium filix-mas* var. *marginata* Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 521. t. 71. 1880. *Lastrea filix-mas* var. *elongata* Bedd. Ferns. Brit. India 250. 1883 ; Handb. Suppl. 56. 1892.

Distrib. : India (Eastern Himalaya and Eastern India), Sri Lanka, Malaya Peninsula, Formosa, China.

D. nigra Ching, Bull. Fan Mem. Inst. Biol. Bot. 8 : 430. 1938.

Distrib. : India (Meghalaya), China, Tibet.

D. odontoloma (Moore) C. Chr. Act. Hort. Gothob. 1 : 59. 1924.

Lastrea odontoloma Moore, Ind. Fil. 90. 1858 ; Bedd. Ferns. South. India 39. t. 114. 1863. *Nephrodium odontoloma* (Moore) Hope, Journ. Bomb. Nat. Hist. Soc. 14 : 736. 1903. *Nephrodium filix-mas* var. *2 normale* Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 519. t. 68. f. 2. 1880. *Lastrea filix-mas* var. *odontoloma* Bedd. Handb. Ferns Brit. India Suppl. 55. 1892.

Distrib. : India (North-West Himalayas), Indo-China, Tibet, China.

D. patentissima (Franch.) Panigr. & Basu Indian Journ. For. 3 (3) : 266. f. 1. t. 2. 1. & 4. 1980. *Aspidium patentissimum* Franch. Nouv. Arch. Mus. II. 10. 118. 1887, non Wall. 1829, nom. nud. *Nephrodium filix-mas* var. *khasiana* Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 519. t. 69. f. 1. 1880. *N. parallelogramma* subsp. *Khasiana* (Clarke) Hope, Journ. Bomb. Nat. Hist. Soc. 14 : 729. 1903. *Lastrea filix-mas* var. *parallelogramma* Bedd. Handb. Ferns. Brit. India 249. 1883. p. p. et suppl. 56. 1892, *pro parte* quoad. var. *Clarkei* Bedd.

Distrib. : India (Sikkim, Meghalaya, Tamil Nadu) China, Tibet.

D. pulvinulifera (Bedd.) O. Ktze. Rev. Gen. Pl. 2 : 213. 1891. *Lastrea pulvinulifera* Bedd. Ferns Brit. India t. 333. 1870 ; Handb. Ferns Brit. India 255. 1883 ; Suppl. 60. 1892. *Lastrea pulvinulifera* var. *zeylanica* Bedd. Ferns Brit. India Suppl. 17, 1876. *Nephrodium pulvinuliferum* Hook. & Bak. Syn. Fil. 500. 1874. (*pro parte*) ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 525. 1880 (*pro parte*). *Nephrodium sparsum* var. *squamulosa* Clarke, Trans Linn. Soc. Lond. II. Bot. 1 : 524. 1880 (*pro parte*). *Lastrea sparsa* var. *zeylanica* Bedd. Handb. Ferns Brit. India. 254. 1883. *Dryopteris harae* Ito in Hara's Fl. East. Himal. 476. t. 26. 1966.

Distrib. : N. E. Himalayas, Sri Lanka.

D. ramosa (Hope) C. Chr. Ind. Fil. 266. 1905. *Nephrodium ramosum* Hope, Journ. Bot. 1896. 126. 1896, sensu Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 523. 1880. *Lastrea rigida* Bedd. Handb. Ferns Brit. India 251. 1883 ; Suppl. 59. 1832 (non presl 1836).

Distrib. : North-East India, Afghanistan, Sri Lanka.

D. reflexosquammata Hayata Icon. pl. Formos. 4 : 176 f. 114. 1914.

Distrib. : India (Himalayas), Formosa.

D. redactopinnata Basu & Panigr. Indian J. For. 3 (3) : 270. t. 2. f. 3, 6. 1980.

Distrib. : India (Darjeeling & Sikkim Himalayas), Uttar Pradesh, (Kumaon), Kashmir.

D. rhodolepis (Clarke) C. Chr. Ind. Fil. 288. 1905. *Nephrodium rhodolepis* Clarke, Trans Linn. Soc. Lond. II. Bot. 1 : 526. t. 72. 1880. *Lastrea blumei sensu* Bedd. Ferns South. India t. 249. 1883 ; Handb. Ferns Brit. India 259. 1883.

Distrib. : India (Himalayas), Sri Lanka, Japan. Malaya Islands, Polynesia.

D. rosthornii (Diels) C. Chr. Ind. Fil. 289. 1905 *Nephrodium rosthornii* Diels, Engl. Jahrb. 29 : 190. 1900.

Distrib. : India (Eastern Himalayas), China.

D. rubra Ching, Bull. Fan Mem. Inst. Biol. Bot. 2 : 198. 1931. *Nephrodium urophyllum* var. *khasiana* Clarke, Trans. Lim. Soc. Lond. II. Bot. 1 : 547. 1880.

Distrib. : India (Assam, Meghalaya, Sikkim), China.

D. scottii (Bedd.) Ching, Bull. Dept. Biol. Col. Sci. Sun. Yatsen Univ. 6 : 3. 1933. *Polyodium scottii* Bedd. Ferns Brit. India t. 345. 1870.

Distrib. : India (Himalayas), Tonkin, China, Formosa.

D. serrato-dentata (Bedd.) Hayata, Icon. Pl. Formos. 4 : 179. 1914. *Lastrea filix-mas* var. *serratodentata* Bedd. Handb. Ferns. Brit. India Suppl 55. 1892. *Nephrodium serrato-dentatum* Hope, Journ. Bomb. Nat. Hist. Soc. : 12 : 622. t. 10. 1899. *Lastrea filix-mas* var. *odonotoloma* Bak. in Hook. et Bak. Syn. Fil. 498. 1874 (excl. fig. of Bedd. cited and the var. *nidus*). *Lastrea odontoloma* Bedd. Ferns Brit. India t. 115. 1865 ; Handb. Ferns Brit. India 258. 1883. *Nephrodium odontoloma* Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 521. 1880. *Lastrea odontoloma* sensu Bedd. Handb. Ferns Brit. India 248. t. 128. 1883 (*non* Moore, 1858).

Distrib. : India (N. E. Himalayas.) Bhutan, China, Formosa.

D. sikkimensis (Bedd.) O. Ktze, Rev. Gen. Pl. 2 : 813. 1891. *Polystichum sikkimensis* Bedd. Ferns Brit. India t. 127. 1866 ; Ferns Brit. India Suppl. 14. 1876. *Aspidium sikkimensis* Bak. in Hook. Syn. Fil. 256. 1867. *Lastrea sikkimensis* Bedd. Handb. Ferns Brit. India. 259. 1883.

Distrib. : India (Eastern Himalayas), China.

D. sino-fibrillosa Ching, Bull. Fan Mem. Inst. Biol. 10 : 180. 1940.

Dryopteris fibrillosa (Clarke) Hand-Mazt. in Akad. Ang. Akad. Wiss. Wien 1922 : 2. 1922. *Nephrodium filix-mas* var. *fibrillosum* Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 520. t. 70, 1880. *Lastrea filix-mas* var. *parallelogramma* subvar. *fibrillosa* (Clarke) Bedd. Handb. Ferns Brit. India 250. 1883.

Distrib. : Himalayas to S. W. China.

D. sparsa (D. Don) O. Ktze, Rev. Gen. Pl. 2. 613. 1891.

Nephrodium sparsum D. Don, Prod. Fl. Nepal 6. 1825 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 523. 1880 (excl. vars.). *Lastrea sparsa* (D. Don.) Bedd. Ferns South. India t. 103. 1863 (excl. parte) ; Bedd. Handb. Ferns Brit. India 252. 1883 (excl. parte). *Nephrodium sparsum* var. *latisquamm* Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 524. 1880.

Distrib. : India (Himalayas), Sri Lanka, Burma, Malaya Peninsula, Malaya Islands, New Guinea, Tibet, China, Formosa, Japan, Philippines, Formosa.

D. sparsa var. *nitidula* (Wall. ex Bedd.) C. Chr. Ind. Fil. 293. 1905.

Lastrea filix-mas var. *nitidula* Wall. ex Bedd. Ferns Suppl. t. 374. 1876; Handb. Ferns. Brit. India 254. 1883.

Distrib. : India (Himalayas).

D. spinulosa (Muell) O. Ktze, Rev. Gen. Pl. 2 : 813. 1891.

Polypodium spinulosum Muell. Fl. Fridrichid. 193. f. 2. 1767 ; Bedd. Handb. Ferns Brit. India 252. 1883. *Lastrea spinulosa* Presl, Tent. Pterid. 76. 1836.

Distrib. : India (Himalayas) Europe, Asia, America.

D. splendens (Hook.) O. Ktze. Rev. Gen. Pl. 2 : 813. 1891.

Nephrodium splendens Hook. Sp. Fl. 4 : 126. 1862 (excl. parte) ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 527. 1880. *Lastrea splendens* Bedd. Ferns Brit. India t. 42. 1865 ; Handb. Ferns Brit. India 262. 1883.

Distrib. : India (Himalayas), S. W. China, Malacca.

D. subimpressa Loyal, Nova Hedw. 16 : 467. 1978.

Distrib. : Eastern Himalayas.

D. submarginata Loyal, Nova Hedw. 16 : 467. 1968.

Distrib. : Eastern Himalayas.

D. subtriangularis (Hope) C. Chr. Ind. Fil. 296. 1905. *Nephrodium*

subtringulare Hope, Journ. Bot. 1890 : 327. 1890. *Lastrea filix-mas* var.

substriangularis (Hope) Bedd. Handb. Ferns Brit. India Suppl. 56. 1892.

Distrib. : India (Assam), China.

D. uropinna Price, Gard. Bull. Singapore 30 : 244. f. 1. A-C. 1977.

Distrib. : Himalayan region to S. Japan, Luzon.

D. varia (L.) O. Ktze, Rev. Gen. Pl. 2 : 814. 1891. *Polypodium varium* L. Sp. Pl. 2 : 1090. 1753. *Lastrea varia* Moore, Ind. Fil. 107. 1858 ; Bedd. Handb. Ferns Brit. India Suppl. 50. 1892.

Distrib. : India (Assam), China, Hong Kong, Japan, Formosa, Luzon Philippines.

D. wallichiana (Spreng.) Hyl. Bot. Notis. 392. 1953. Basu, Indian Journ. For. 3 (3) : 269. t. 2. f. 2 & 5. 1980. *Aspidium wallichianum* Spreng, Syst. veget. 4 (1) : 104. 1827. *Lastrea patentissima* Bedd. Ferns Sou h. India 39 t. 111. 1864. *nom superfl.* *Nephrodium patentissimum* Clarke, Journ. Linn. Soc. 15 : 156. 1876, incl. descr. in part, excl. type. *Nephrodium filix-mas* var. *patentissimum* (Clarke) Clarke Trans. Linn. Soc. London II Bot. 1 : 520. 1880. *pro parte* quoad sp. *Wallich* 340. *Dryopteris filix-mas* subsp. *patentissima* (Clarke) C. Chr. Ind. Fil. 265. 1905. *pro parte*, excl. *Nephrodium filix-mas* var. *fibrillosa* Clarke, 1880. *Lastrea filix-mas* var. *parallelogramma* Bedd. Handb. Ferns Brit. India 249. 1883. *pro parte* et suppl. 56. 1892 ; *pro parte* quoad subvar. *Clarkei* Bedd.

Distrib. : India (Himalayas), Sri Lanka, China, Burma, Java, Borneo, New Guina, Philippines, Formosa.

D. yunnanensis (Christ) Copel. Gen. Fil. 122. 1947. *Aspidium yunnanense* Christ, Bull. Herb. Boiss. 6 : 965. 1898.

Distrib. : India (Himalayas), China.

Nothoperanema (Tagawa) Ching, Acta Phytotax. Sin. 11 : 25. 1966.

Type : *N. squamisetum* (Hook). Ching, Acta Phytotax. Sin. 11 (1) : 27. 1966.

N. hendersonii (Bedd.) Nakaike, Enum Pterid. Jap. 239. 1975. *Lastrea hendersonii* Bedd. Ferns Brit. India. Suppl. 17. t. 377. 1876. *Nephrodium spectabile* Clarke, Trans. Linn. Soc lond. II. Bot. 1 : 526. 1880. *Lastrea spectabilis* (Clarke) Bedd. Handb. Ferns Brit. India. 257. 1883.

Distrib. : India (Himalayas) Nepal China, Taiwan, Thailand,.

N. squamisetum (Hook.) Ching, Acta Phytotax. Sin. 11 : 27. 1966.
Nephrodium squamisetum Hook. Sp. Fl. 4 : 140. t. 268. 1862. *Lastrea buchanani* Bedd. Handb. Ferns Brit. India 255. 1883.

Distrib. : India (Himalayas), Africa.

Polystichum Roth, Tent. Pterid. Fl. Germ. 3 : 31. 269. 1799 : Khuller & Gupta, Nova Hedw. 63 : 41. 1979 ; 64 : 428. 1980.

Type : **P. lonchitis** (L.) Roth. Rom. Mag. 2 (1) : 106. 1799.

P. acanthophyllum (Franch.) Christ, Bull. Soc. Brot. France 52 Mem. 1 : 30. 1905. *Aspidium acanthopoyllum* Franch. Bull. Soc. France 32 : 23. 1885. *Polystichum aculeatum* var. *acanthophyllum* (Franch.) Bedd. Handb. Ferns Brit. India Suppl. 43. 1892.

Distrib. : India (Himalayas), China, Nepal.

P. aculeatum (L.) Schott, Gen. Fil. ad. t. 9. 1834. Bedd. Handb. Ferns Brit. India 207. 1883. *Polypodium aculeatum* L. Sp. Pl. 2 : 1090. 1753. (*pro parte*).

Distrib. : Cosmopolitan.

P. anomalum (Hook. & Arn.) J. Sm. Ferns Brit. For. 151. 1866 ; Bedd. Ferns South. India t. 219. 1869 ; Handb. Ferns Brit. India 209. 1883. *Aspidium anomalum* Hook. & Arn. Sp. Fil. 4 : 27. 1862.

P. anomalum var. **travancoricum** (Bedd.) Sledge, Bull. Brit. Mus. (N.H.) 5 : 33. 1973. *P. aculeatum* var. *travancoricum* Bedd. Handb. Ferns Brit. India 209. 1883. *P. travancoricum* (Bedd.) Nayar & Kaur, Comp. to Bedd. Handb. 49. 1974.

Distrib. : South India, Sri Lanka, Tonkin, Thailand, China.

P. atkinsoni Bedd. Ferns Brit. India Suppl. 14. t 362. 1876 ; Handb. Ferns Brit. India 203. 1883. *Aspidium atkinsoni* Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 506. 1880,

Distrib. : India (Sikkim), Bhutan.

P. auriculatum (L.) Presl, Tent. Pterid. 83. 1836 ; Bedd. Ferns South. India t. 120. 1866 ; Handb. Ferns Brit. India 203. t. 102. 1883 (*excl. vars.*) *Polypodium auriculatum* L. Sp. Pl. 2 : 1088. 1753.

Distrib. : South India, Sri Lanka.

P. bakerianum (Atkinson) Diels, Nat. Pflanzn. 1 (4) : 191. 1899. *P. aculeatum* var. *bakerianum* Clarke, Trans. Linn. Soc. Lond. II. Bot. t. 66. 1880 ; Bedd. Handb. Ferns Brit. India 210. 1883.

Distrib. : India (Sikkim).

P. biaristatum (Bl.) Moore, Ind. Fil. 86. 1856; *Aspidium biaristatum* Bl. Enum. Pl. Jav. 164. 1828. *Aspidium aculeatum* var. *biaristatum* Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 510. 1880. *P. aculeatum* var. *biaristatum* (Clarke) Bedd. Handb. Ferns Brit. India 209. 1883.

Distrib. : (S. India, Meghalaya), Sri Lanka, Malaya Peninsula, China, Burma, Thailand, Java, Celebes, Formosa.

P. castaneum (Clarke) Nayar & Kaur, Comp. to Bedd. Handb. 50. 1883. *P. aculeatum* var. *castaneum* Clarke, Trans Linn. Soc. Lond. II. Bot, 1 : 509. 1880 ; Bedd. Handb. Ferns Brit. India 210. 1883.

Distrib. : India (Sikkim).

P. duthiei (Hope) C. Chr. Ind. Fil. 581. 1905. *Aspidium duthiei* Hope, Journ. Bomb. Nat. Hist. Soc. 12. 532. 1899.

Distrib. : India (Himalayas), China.

P. discretum (D. Don.) Diels, Nat. Pflanzn. 1 (4) : 199. 1899. *P. aculeatum* var. *setosum* Wall. ex Bedd. Handb. Ferns Brit. India 209. 1883.

Distrib. : India (Himalayas).

P. eximum (Mett. ex Kuhn) C. Chr. Bull. Dept. Biol. Coll. Sci. Sun yatsen Univ. no. 6 : 8. 1933. *Phegopteris exima* Mett. ex Kuhn in Linnaea 36 : 107. 1869.

Distrib. : India (Himalayas), China, Japan, Taiwan, Sri Lanka.

P. garhwalicum Nair & Nag, Journ. Jap. Bot. 51. 138. t. 1. 1976.

Distrib. : India (North-West Himalayas).

P. heteropalaceum Nair & Nag, Journ. Bomb. Nat. Hist. Soc. 75 : 153. 1977.

Distrib. : India (Meghalaya).

P. indicum khuller & Gupta, Nova Hedw. 32 : 425. 1980.

Distrib. : India (North-West Himalayas).

P. lachenense (Hook.) Bedd. Ferns Brit. India t. 32. 1865; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 506. 1880 ; Bedd. Handb. Ferns Brit. India 203. 1883. *Aspidium lachenense* Hook. Sp. Fil. 4 : 8. 1862.

Distrib. : Himalayas to China, Formosa, Japan.

P. lendum (D. Don) Moore, Ind. Fil. 86. 1858. *Aspidium lendum* D. Don, Prod. Fl. Nepal 4. 1825. *Polystichum auriculatum* var. *lendum* (D. Don) Bedd. Handb. Ferns Brit. India 204. 1883. *P. auriculatum* var. *sub-pinnatum* Bedd. Ferns Brit. India t. 136. 1867.

Distrib. : India (Himalayas), Nepal, Bhutan.

P. lendum (D. Don) Moore var. *bifurcatum* Biswas & Basu, J. Econ. & Tax. Bot. 3 : 961. f. 1. 1982.

Distrib. : India (Sikkim).

P. levingei Nair, Bull. Bot. Surv. India 11 : 192. 1869.

Distrib. : India (West Bengal : Darjeeling).

P. longipinnulum Nair, Amer. Fern Journ. 64 : 15. 1974.

Distrib. : India (Eastern Himalayas).

P. lobatum (Huds.) Presl, Tent. Pterid. 83. 1836. *Polypodium lobatum* Huds. Fl. Angl. 459. 1762. *P. aculeatum* var. *lobatum* (Huds.) Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 509. 1880 ; Bedd. Handb. Ferns Brit. India 207. 1883.

Distrib. : India (Himalayas).

P. lonchitis (L.) Roth, Rom. Mag. 2 (1) : 106. 1799 ; Bedd. Ferns Brit. India t. 128. 1867 ; Handb. Ferns Brit. India 203. 1883. *Polypodium lonchitis* L. Sp. Pl. 2 : 1088. 1753.

Distrib. : India (Himalayas), Japan, Europe, N. America.

P. mucronifolium (Bl.) Nayar et Kaur, Comp. to Bedd. Handb. 49. 1974. *P. aculeatum* var. *mucronifolium* Bl. Enum. Pl. Jav. 64. 1828 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 509. 1880 ; Bedd. Handb. Ferns Brit. India 210. 1883.

Distrib. : India (Meghalaya, Assam).

P. neolobatum Nakai, Bot. Mag. Tokyo 39 : 118. 1925.

Distrib. : India (Himalayas), Bhutan, Nepal, Taiwan, Japan.

P. nepalense (Spreng.) C. Chr. Ind. Fil. 84. 1905 ; 585. 1906. *Aspidium nepalense* Spreng. Syst. 4 : 97. 1827. *Aspidium marginatum* Wall. List no 366. 1828 (*nom. nud.*) ; Hope, Journ. Bomb. Nat. Hist. Soc. 14 : 459. t. 17. 1902. *Polystichum auriculatum* var. *marginatum* Bedd. Handb. Ferns Brit. India 201. 1883. *Aspidium auriculatum* var. *marginata* Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 126. 1880.

Distrib. : India (Himalayas), China, Formosa.

P. nepalense var. **subbipinnatum** C. Chr. Contr. U.S. Nation Herb. 26 : 284. 1931.

Distrib. : Eastern Himalayas Bhutan.

P. nigropalaceum (Christ) Diels, in Engl. & Prant. Nat. pfl. 1 (4) : 191. 1899. *P. aculeatum* var. *nigropalaceum* Christ, Ber. Schweiz. Bot. Ges. 3. 1893. *P. setiferum* var. *nigropalaceum* (Christ) Sledge, Bull. Brit. Mus. (N.H.) Bot. 5 : 37. 1973. *P. aculeatum* var. *angulare* Bedd. Handb. Ferns Brit. India 207. 1883. pp. (*non Aspidium angulare* Wall.). *Aspidium angulare* sensu Hope, Journ. Bomb. Nat. Hist. Soc. 14 : 472. 1902. (*non Willd.* 1810).

Distrib. : India (North-West Himalayas, S. India). China Sri Lanka.

P. obliquum (D. Don) Moore, Ind. Fil. 87. 1858. *Aspidium obilquum* D. Don, Prod. Fl Nepal 3. 1825. *Aspidium caespitosum* Wall. List no. 367. 1828. (*nom. nud.*). *Aspidium auriculatum* Sw. var. *caespitosa* Clarke, Trans. Linn. Soc. Lond. II. Bot. I : 507. 1880. *Polystichum auriculatum* var. *caespitosum* (Clarke) Bedd. Handb. Ferns Brit. India 204. 1883.

Distrib. : India (Himalayas), Bhutan.

P. prescottianum (Wall. ex Mett.) Moore, Ind. Fil. 101. 1858 ; Bedd. Handb. Ferns Brit. India 210. t. 105. 1883. *Aspidium prescottianum* Wall. ex Mett. in Abh. Senckneb. Ges. Frank. 2 : 48. 1858.

Distrib. : India (Himalayas). China and Formosa.

P. punctiferum C. Chr. Contr. U.S. Nation Herb. 26 : 288. t. 17. 1931.

Distrib. : India (Eastern Himalayas) Bhutan, Burma, China.

P. semifertile (Clarke) Ching, Lingnan. Sci. Journ. 15 : 398. 1938. *Aspidium aculeatum* var. *semifertile* Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 509. 1880.

Disirib. : (Eastern Himalayas, Tonkin,) Thailand, China.

P. setiferum (Forssk.) Moore ex Woynar Mitt. Naturw. Ver. Steierm. 49 : 181. 1913. *Polypodium setiferum* Forsk. Fl. Aeypt. arab. 185. 1775. *Polystichum aculeatum* var. *angulare* Presl ex Bedd. Ferns South. India t. 121. 1865 ; Handb. Ferns Brit. India 207. 1883.

Distrib. : North & South India.

P. setiferum var **crenatum** Nair, Amer. Fern Journ. 64 : 15. 1974.

Distrib. : India (Punjab).

P. setosum (Wall. ex Bedd.) Khullar & Gupta, Nova Hedw. 32. 420. 1980. *P. aculeatum* var. *setosum* Wall. ex Bedd. Handb. Ferns Brit. India 209. 1883.

Distrib. : India (North-West Himalayas, Sikkim).

P. squarrosum (D. Don) Féé, Gen. Fil. 278. 1850-52. *Aspidium squarrosum* D. Don, Prod. Fl. Nepal 4. 1825. *P. aculeatum sensu* Bedd. Ferns South. India t. 121. 1865. *P. aculeatum* var. *rufobarbatum* Wall. ex Bedd. Handb. Fern Brit. India 207. 1883.

Distrib. : India (Himalayas) to China.

P. stenophyllum Christ, Bull. Soc. Bot. France 52. Mém. 1 : 27. 1905.

Distrib. : India (Himalayas) to China, Formosa.

P. stimulans (Kunze ex Mett.) Bedd. Ferns Brit. India t. 31. 1865. *Aspidium stimulans* Kunze ex Mett. Abh. Senckneb. Ges. Frank. 2 : 327. 1858. *A. ilicifolium* D. Don, Prod. Fl. Nepal. 3. 1825. *Polystichum ilicifolium* (D. Don) Moore, Ind. Fil. 94. 1858. (*non* Féé 1850); Bedd. Handb. Ferns, Brit. India 206. 1883.

Ditstrib. : India (Himalayas) to W. China (Yunnan).

P. tacticopterum (Kunze) Moore, Ind. Fil. 105. 1858. *Aspidium tacticopterum* Kunze, Linnaea 24 : 210. 851.

Distrib. : S. India, Sikkim, Meghalaya.

P. thomsoni (Hook. f.) Bedd. Ferns Brit. India 1 : t. 126. 1865. Handb. Ferns. Brit. India 206. 1883. *Aspidium thomsoni* Hook. f. in Hook. 2nd cent. Ferns t. 25. 1860; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 508. 1880.

Distrib. : India (Himalayas) to China, Formosa.

P. tsus-simense (Hook.) Smith, Journ. Hist. Fil. 219. 1875. *Aspidium tsus-simense* Hook. Sp. Fl. 4 : 16. t. 220. 1862.

Distrib. : India (Himalayas).

LOMARIOPSIDACEAE Aloton

Leptogramma J. Sm. Journ. Bot. 4 : 51. 1841.

Lectotype : **L. totta** (Schlecht.) J. Sm. Journ. Bot. 4 : 52. 1841.

L. totta (Schlecht.) J. Sm. Journ. Bot. 4 : 52. 1841 ; Bedd. Handb. Ferns Brit. India 377. t. 215. 1883. *Gymnogramma tottum* Schlecht. Adumb. 13. t. 6. 1825. *Grammitis totta* Presl, Tent. Pterid. 209. t. 9. f. 4. 1836 ; Bedd. Ferns South. India t. 49. 1863. *Thelypteris totta* (Schlecht.) Nayar & Kaur, Comp to Bedd. Handb. 90. 1974.

Distrib. : Throughout India in mountainous regions, Nepal, Bhutan, Sri Lanka, China, Japan, Malesia.

BOLBITIDACEAE (Pic. Ser.) Ching

Bolbitis Schott, Gen. Fil. t. 14. 1834 ; Hennipman, Fl. Males. ser. II. 1 (4) : 314. 1978.

Lectotype : **B. serratifolia** (Mart. ex Kaulf.) Schott, Gen. Fil. 3 : t. 14. 1834.

B. angustipinna (Hayata) Ito, Journ. Jap. Bot. 14 : 443. 1938. *Leptochilus angustipinnus* Hayata, Icon. Pl. Fern. 5 : 297. f. 119. 1915. *Acrostichum crispatulum* Clarke var. *contaminans* Clarke. Trans. Linn. Soc. Lond. II. Bot. 1 : 580. t. 84. f. 2 A, C. 1880.

Distrib. : India (Himalayas), Sri Lanka, China, Taiwan, Thailand, Philippines.

B. contaminans (Wall.) Ching C. Chr. Ind. Fil. Suppl. 3 : 47. 1934. *Acrostichum contaminans* Wall. List. no. 22. 1828. *Gymnopteris contaminans* Bedd. Ferns Brit. India Suppl. 27. 1876 ; Handb. Ferns Brit. India 435. 1883 (*pro parte*). *Paecilopteris repanda* Bedd. Ferns South. India t. 202. 1865.

Distrib. : South India, Burma, Sri Lanka.

B. costata (Wall. ex Hook.) Ching, C. Chr. Ind. Fil. Suppl. 3 : 47. 1934. *Acrostichum costatum* Wall. ex Hook Sp. Pl. 5 : 262. 1864. *Gymnopteris costata* Bedd. Ferns Brit. India Suppl. 27. 1876 ; Handb. Ferns Brit. India 438. 1883.

Distrib. : North-East India, Nepal, Bangladesh.

B. crispatula (Wall.) Ching. C. Chr. Ind. Fil. 102. 1905.
Acrostichum crispatulum Wall. List no. 24. 1828. *Gymnopteris contaminans* Bedd. Ferns Brit. India 435. 1883 (*pro parte*).

B. deltigera (Wall ex Clarke) C. Chr. Ind. Fil. 3: 48. 1934. *Meniscium deltigerum* Wall. List no. 59. 1828 ex Clarke, Trans Linn. Soc. Lond. II. Bot. 1 : 572. 1880. *Gymnopteris flagellifera* Bedd. Ferns Brit. India Suppl. 27. 1876 ; Handb. Ferns Brit. India 433. 1883.

Distrib. : North-East India, Sikkim, Bangladesh, Burma, Japan, Formosa, China, Malesia.

B. presliana (Fée) Ching, C. Chr. Ind. Fil. Suppl. 3 : 49. 1934.
Acrostichum preslianum Fée, Acrost. 46. 1845. *Gymnopteris presliana* (Fée) Bedd. Handb. Ferns Brit. India 439. 1883.

Distrib. : South India, Philippines.

B. simicordata (Bak.) Ching in C. Chr. Ind. Fil. Suppl. 3 : 50. 1934.
Acrostichum semicordatum Bak. Syn. Fil. 422. 1868.

Distrib. : India, Australia.

B. sinuata (Presl) Henmann, Blumea 18 : 148. 1970. *Polyodium sinuatum* Presl. Rel. Haenk. 1 : 21. 1825.

Distrib. : India (Nicobar Islands), Thailand, Malesia.

B. subcordata (Copel.) Ching, C. Chr. Ind. Fil. Suppl. 3 : 50. 1934. *Campilum subcordatum* Copel. Philip. Journ. Sci. 37. 369. 1928.

Distrib. : India (Eastern Himalayas), China, Indo-China, Japan.

B. subcrenata (Hook. & Grev.) Ching, C. Chr. Ind. Fil. 3 : 50. 1934. *Acrostichum subcorenatum* Hook. & Grev. Icon. Fil. t. 110. 1829. *Gymnopteris subcrenata* (Hook. & Grev.) Bedd. Handb. Ferns Brit. India 437. 1883.

Distrib. : South India, Western ghats, Sri Lanka.

B. undulata (Wall.) Ching, C. Chr. Ind. Fil. Suppl. 3 : 50. 1934.
Notholaena undulata Wall. List no. 140. 1828.

Distrib. : India, Cambodia.

B. virens (Hook. & Grev.) Schott var. **compacta** Hennipman,
 Blumea 18 : 149. 1970.

Distrib. : India (Nicobar Islands), Vietnam. Thailand. Malesia.

Egenolfia Schott, Gen. Fil. t. 16. 1834 ; Ching, Bull. Fan Mem. Inst. Biol. Bot. 2 : 302. 1931 ; Nair & Kaur, Bull. Nation. Bot. Gard, Lucknow no. 100. 17. 1964.

Type : *E. hamiltoniana* Schott, Gen. Fil. t. 16. 1834.

E. appendiculata (Willd.) J. Sm. Ferns Brit. & For 111. 1866. *Acrostichum appendiculatum* Willd. Sp. Pl. 5 : 114. 1810. *Polybotrya appendiculata* (Willd.) J. Sm. Journ. Bot. 4 : 150. 1841 ; Bedd. Ferns South. India t. 194. 1865 ; Handb. Ferns Brit. India 424. 1883.

Distrib. : Throughout India, Philippines. Hongkong.

E. appendiculata var. **major** (Bedd.) Nayar & Kaur, Comp. to Bedd. Handb. 99. 1974. *Polybotrya appendiculata* var. *major* Bedd. Handb. Ferns Brit. India 424. 1883.

Distrib. : India (Sikkim).

E. appendiculata (Willd.) J. Sm. ssp. **vivipera** (Hook.) Hennipman var. **neglecta** (Hennipman) Dixit comb. nov. *Bolbitis appendiculata* (Willd.) Iwats. ssp. *vivipera* (Hook.) Hennipman var. *neglecta* Hennipman, Blumea 18 : 147. 1970.

Distrib. : India (Andaman & Nicobar Islands), Malesia.

E. asplenifolia (Bory) Fée, Gen. Fil. 358. 1852. *Acrostichum asplenifolium* Bory, Bél. Voy. Bot. 2 : 21. t. 3. 1833. *Acrostichum wightianum* Wall. List no. 2163. 1829 (nom. nud.). *Polybotrya asplenifolia* Presl, Tent. Pterid. 231. 1836 ; Bedd. Ferns South. India t. 195. 1865. *Polybotrya appendiculata* var. *asplenifolia* (Bory) Bedd. Handb. Ferns Brit. India 424. t. 235. 1883.

Distrib. : S. India, Common in Western Ghats, Sri Lanka.

E. bipinnatifida J. Sm. Hist. Fil. 132. 1875. *Acrostichum appendiculatum* var. *costulata* Hook, Sp. Fil. 4 : 252. 1860. Bedd. Ferns Brit. India t. 110. 1865. *Polybotrya appendiculata* var. *costulata* (Hook.) Bedd. Handb. Ferns Brit. India 426. t. 257. 1883.

Distrib. : Eastern India, Burma.

E. helferiana (Kze.) C. Chr. Contr. U.S. Nat. Herb. 26. 292. 1931 ; *Polybotrya helferiana* Kze. Farnkr. 2 : 35. t. 114. 1848. *Polybotrya helferiana* var. *hamiltoniana* Wall. ex. Bedd., Ferns Brit. India t. 111. 1855 ; Handb. Ferns Brit. India 426. t. 256. 1883.

Distrib. : India (Meghalaya. Assam. Andaman Islands), Burma.

E. nodiflora (Bory) Fée, Gen. Fil. 48. 1850-52. *Polybotrya nodiflora* Bory, Bél. Voy. Bot. 2 : 17. 1833. *Acrostichum hamiltoniana* Wall. List no. 29. 1929. (*nom. nud.*). *Polybotrya appendiculata* var. *subintegra* Bedd. Ferns Brit. India t. 3. 1865.

Distrib. : India (Assam, Meghalaya, South India), Burma, Bangladesh, Siam, Cambodia.

E. keralensis (Nayar & Chandra) Nayar & Kaur, Bull. Nation. Bot. Gard. Lucknow no. 94 : 4. 1964. *Bolbitis keralensis* Nayar et Chandra, Amer. Fern Journ. 54 : 9. 1964.

Distrib. : S. India.

E. sinensis (Bak.) Maxon, Proc. Biol. Soc. Wash. 36 : 173. 1923. *Acrostichum sinense* Bak. Kew Bull. 1906 ; 14. 1906. *Polybotrya appendiculata* var. *ludens* Bedd. Ferns Brit. India Suppl. 26. 1876.

Distrib. : Eastern India (Himalayas), Nepal, S. China, Japan,

E. vivipera (Hook.) C. Chr. Ind. Fil. Suppl. 3 : 102. 1934. *Polybotrya vivipera* Hook. Exot. Fl. 2 : t. 107. 1825.

Distrib. : Eastern India, Cambodia.

ELAPHOGLOSSACEAE Pic. Ser.

Elaphoglossum Schott ex J. Sm. Hook. Journ. Bot. 4 : 148. 1841 ; Sledge, Bull. Brit. Mus. (N.H.) Lond. Bot. 4. 81. 1967 ; Ghosh & Biswas. J. Econ. Tax. Bot. 4(3) : 995. 1980.

Type : **E. conforme** (Sw.) Schott, Gen. ad. t. 14 : 1834.

E. angulatum (Bl.) Moore, Ind. Fil. V. 1857. *Acrostichum angulatum* Bl. Enum. Pl. Jav. 101. 1828 ; *Acrostichum conforme* sensu Hook. Sp. Fil. 5 : 98. 1864. (*pro parte*) ; non Sw. *Elaphoglossum marginata* sensu Thw. Enum. Pl. Zeyl. 380. 1864. (non Wall. ex Fée). *E. laurifolium* sensu Bedd. Ferns South. India 67. t. 200. 1864 (*non* T. Moore). *E. latifolium* sensu Bedd. Handb. Ferns Brit. India 416. t. 248. 1883 (*non* J. Sm.). *E. krajina* Biswas, Misc. Inf. Kew 1939 : 240. f. 2. 1939.

Distrib. : S. India, Sri Lanka, Java.

E. beddomei Sledge, Bull. Brit. Mus. (N.H.) Bot 4 (2) : 85. 1967.
E. stigmatolepis sensu Bedd. Ferns South. India 67. t. 197. 1864 ;

Handb. Ferns Brit. India 418. 1883 (*non* T. Moore). *Acrostichum stigmatolepis sensu* Bak. in Hook. et Bak. Syn. Fil. ed. 2 : 51. 1874.

Distrib. : South India.

E. ceylanicum Krajina ex Sledge, Bull. Brit. Mus. (N.H.) 4 (2) : 95. 1967.

Distrib. : India (Sikkim), Sri Lanka.

E. cherrapunjii Ghosh et Biswas, J. Econ. Tax. Bot. 4 (3) : 996. t. 2. 1983.

Distrib. : E. India (Meghalaya).

E. commutatum (Mett.) v. A. v. R. Malaya Ferns Suppl. 1 : 427. 1917. *E. laurifolium sensu* Moore, Ind. Fil. 359. 1862 (*pro parte*) quoad specim. ex India. Ceylon, Java). *Acrostichum commutatum* Mett. apud. Kuhn in Ann. Mus. Bot. Lugd. Bat. 4 : 292. 1869.

Distrib. : India, Malaya Peninsula.

E. jowaiense Biswas & Ghosh, Journ. Bomb. Nat. Hist. Soc. 79 (3) : 650. f. 1 5. 1982. *E. viscosum sensu* Bedd. Handb. Ferns Brit. India 40. 1883.

Distrib. : India (Meghalaya, Sikkim), Nepal.

E. marginatum (Wall. ex Féé) Moore, Ind. Fil. 89, 1857. *Acrostichum marginatum* Wall. List no. 17. 1829. *nom. nud.* ex Féé Mém. Fam. Foug. 2 : 31. 1845 (excl. syn. Bl.). *Elaphoglossum conforme sensu* Bedd. Handb. Ferns Brit. India 416. 1883 (*pro parte*), excl. f. 247. non J. Sm.

Distrib. : South-East India, Eastern Himalayas, Nepal, Bhutan, Sri Lanka, Java.

E. nilgiricum krajina ex Sledge, Bull. Brit. Mus. (N.H.) 4 (2) : 94. 1967. *E. squamosum sensu* Bedd. Ferns South. India 67. t. 197. 1864 ; Handb. Ferns Brit. India 420. f. 25. 1883, excl. Specim. xe Ceylon, *non* J. Sm.

Distrib. : South India.

E. prainii Ghosh & Biswas, J. Econ. Tax. Bot. 4 (3) : 995. t. 1. 1983.

Distrib. : E. India (Nagaland).

E. simonsianum Ghosh & Biswas, J. Econ. Tax. Bot. 4 (3) : 1001. t. 4. 1983.

Distrib. : E. India (Meghalaya).

E. stelligerum (Wall. ex Bak.) T. Moore ex Alston et Bonner in *Candollea* 15 : 216. 1956. *Acrostichum stelligerum* Wall. ex Bak. in Hook. et Bak. *Syn. Fil.* ed. 2. 521. 1874. *Elaphoglossum viscosum sensu* Bedd. *Ferns South. India* 67 t. 196. 1864 ; *Handb. Ferns Brit. India* 40. 1883 ; (*pro parte*).

Distrib. : S. India, China, India-China.

E. stigmatolepis (Fée) Moore, *Ind. Fil.* XVI : 15. 1857. *Acrostichum stigmatolepis* Fée, *Mém. Fan. Foug.* 2 : 62. t. 24. f. 2. 1845. *Elaphoglossum conforme sensu* Bedd. *Ferns South. India* 67. t. 19. 1864 ; *Handb. Ferns South, India* 416. f. 247. 1883 (*pro parte* ; non J. Sm.) *E. ballardianum* Biswas, *Bull. Misc. Inf. Kew* 1939 : 239. t. 1. 1939.

Distrib. : South India.

E. thomsonii Ghosh & Biswas, *J. Econ. Tax. Bot.* 4 (3) : 996. t. 3. 1983.

Distrib. : India (Sikkim).

NEPHROLEPIDACEAE (Ching) Pic. Ser.

Arthropteris J. Sm. ex Hook. *Bot. Antarct. Voy. II. Fl. Nov.-Zel.* 2 : 43 : 1854.

Type : **A. tenella** (G. Forst.) J. Sm. ex Hook. *Bot. Antarct. Voy. II. Fl. Nov. Zel.* 2 : 43. 1854.

A. palisotti (Desv.) Alston, *Biol. Soc. Brot.* 30 : 6. 1954. *Aspidium palisotti* Desv. *Ges. Naturf. Berl. Mag.* 5 : 320. 1811. *Nephrolepis ramosa* Moore, *Ind. Fil.* 102. 1858 ; Bedd. *Handb. Ferns Brit. India* 284. t. 145. 1883.

Distrib. : South India, Sri Lanka, Malaya, Australia, W. Africa, Thailand to South China.

Nephrolepis Schott, *Gen. Fil.* t. 3. 1834.

Lectotype : **N. exaltata** (L.) Schott, *Gen. Fil.* t. 3. 1834.

N. biserrata (Sw.) Schott, *Gen. Fil. ad. t. 3.* 1834. *Aspidium biseriatum* Sw. *Schrad. Journ. Bot.* 1800 (2) : 32, 1801. *Nephrolepis acuta* Presl. *Tent. Pterid.* 79. 1836.

Distrib. : India (South India, North-East India), Bangladesh, Sri Lanka, Tropical Africa.

N. cordifolia (L.) Presl, Tent. Pterid. 79. 1836 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 541. 1880 ; Bedd. Handb. Ferns Brit. India 282. t. 144. 1883 ; *Polypodium cordifolium* L. Sp. Pl. 2 : 1089. 1753. *Nephrolepis tuberosa* Presl, Tent. Pterid. 79. 1836 ; Bedd. Ferns South. India t. 92. 1865.

Distrib. : Tropics and subtropics of the world.

N. delicatula (Decne) Pic. Ser. Webbia 23 : 181. 1968. *Nephrodium delicatulum* Decne in Jacq. Voy. Ind. Bot. 4 : 178. t. 179. 1844. *Nephrolepis paucifrondosa* d' Almeida, J. Indian Bot. Soc. 5 : 51. f. 1-5. 1926.

Distrib. : India (Himalayas) Nepal, Burma, Thailand, China (Yunnan).

N. exalatata (L.) Schott, Gen. Fil. t. 3. 1834 ; Bedd. Ferns South. India t. 93. 1865 ; Handb. Ferns Brit. India 283. 1883.

Disirib. : Tropics and subtropics of the world

N. falcata (Cav.) C. Chr. Ann. Dansk. Bot. Ark. 9 : 15. 1937. *Tectaria falcata* Cav. Descr. 250. 1802.

Distrib. : South India and tropics of the world.

N. hirsutula (Forst.) Presl, Tent. Pterid. 79. 1836. *Polypodium hirsutulum* Forst. Prod. 81. 1786.

Distrib. : India, Japan, Thailand, New Guinea.

N. multiflora (Roxb.) Jarrett in Morton, Contrib. U.S. Nat. Herb. 38 : 309. 1974 ; Manickam & Ninan, New Botanist 1 : 15 : 1976. *Davallia multiflora* Roxb. ex Griff. Calc. Journ. Nat. Hist. 4 : 515. t. 31. 1844.

Distrib. : S. India.

N. undulata (Afz. ex Sw.) J. Sm. in Curtis Bot. Mag. 72 : 35. 1846. *Aspidium undulatum* Afz. ex Sw, Schrad. Journ. Bot. 1800 (2) : 32. 1801.

Distrib. : India, China Japan.

OLEANDRACEAE Ching ex Pic. Ser.

Oleandra Cav. Anales Hist. Nat. 1 (2). 115. 1799 ; Pic. Ser. Webbia 20 (2) : 745. 1965.

Type : **O. neriformis** Cav. Anales Hist. Nat. 1 (2) : 115. 1799.

O. musaefolia (Bl.) Presl, Epim Bot. 42. 1849 ; Bedd. Handb. Ferns Brit. India 287. 1883. *Aspidium musaefolium* Bl. Enum. Pl. Jav. 141.

1828. *Oleandra neriformis sensu* Bedd. Ferns South. India 32. t. 91. 1863. (*non* Cav.).

Distrib. : South India, Sri Lanka, Java.

O. pistillaris (Sw.) C. Chr. Ind. Fil. Suppl. 3 : 132. 1934. *Aspidium pistillare* Sw. Schrad. Journ. Bot. 1800 (2) : 30. 1801. *Oleandra neriformis sensu* auct. Bedd. Ferns. Brit. India t. 264. 1868 ; Handb. Ferns Brit. India 285. t. 146. 1883 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 541. 1880.

Distrib. : Eastern Himalayas, Nepal, Bhutan, Malaysia, Java.

O. undulata (Willd.) Ching, Lingnan Sci. Journ. 12 : 565. 1933. *Polypodium undulatum* Willd. in Linn. Sp. Pl. 5: 155. 1810. *O. cumingi* var. *longipes* Hook. Sp. Fil. 4 : 15. 1860 ; Bedd. Ferns Brit. India t. 135. 1866. *O. cumingii* Hook. et Bak. Syn. Fil 303. 1868 (*non* J. Sm.) ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 542. 1880 ; Bedd. Handb. Ferns Brit. India 288. 1883.

Distrib. : Eastern India, Burma, Malesia, China, Taiwan.

O. wallichii (Hook.) Presl, Tent. Pterid. 78. 1836 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 542 ; 1880. Bedd. Ferns Brit. India t. 263. 1868. Handb. Ferns Brit. India. 287. t. 147. 1883. *Aspidium wallichianum* Hook. Exot. Fl. 1. t. 5. 1823.

Distrib. : India (Himalayas, Punjab to Assam), Nepal, Bhutan, Burma, Indo. China, Malay Peninsula, China, Formosa.

DAVALLIACEAE Mett. ex. Frank

Araiostegia Copel. Philip. Journ. Sci. 34 : 240. 1927.

Type : **A. hymenophylloides** (Bl.) Copel. Philip. Journ. Sci. 34 : 240. 1927.

A. beddomei (Hope) Ching in Chien et Chun Fl. Republ. Pop. Sin. 2 : 288. 1959. *Davallia beddomei* Hope, Journ. Bomb. Nat. Hist. Soc. 12 : 527. 1899.

Distrib. : Eastern India.

A. clarkei (Bak.) Copel. Philip. Journ. Sci. 34 : 341. 1927. *Davallia clarkei* Bak. Hook. & Bak. Syn. Fil. ed. II. 91. 1874. *leucostegia clarkei* (Bak.) C. Chr. Contrib. U.S. Nat. Herb. 26 : 294. 1931.

Distrib. : India (Himalayas), Tibet, S.W. China.

A. dareiformis (Hook.) Copel. Univ. Calif. Publ. Bot. 12 : 398. 1931.
Polypodium dareiformis Hook. 2nd Cent. Ferns. t. 24. 1860. *Davallia dareiformis* Levinge ex Clarke in Trans Linn. Soc. Lond. II. Bot. 1 : 443. 1880.

Distrib. : Himalayas from Nepal to Assam, Sri Lanka, Indo-China, S.W. China.

A. delavayi (Bedd. ex Clarke et Bak.) Ching, in Fl. Reip. Pop. Sin. 2: 289. 1959. *Davallia pulchra* var. *delavayi* Bedd. ex Clarke et Bak. Journ. Linn. Soc. 2 : 410. 1888. *Leucostegia delavayi* (Bedd. ex Clarke et Bak.) Ching in C. Chr. Ind Fil. Suppl. 3 : 120. 1934. *Leucostegia pulchra* var. *delavayi* sensu Bedd. Handb. Ferns Brit. India Suppl. 13. 1892.

Distrib. : Meghalaya, Sikkim, Nepal, China (Yunnan).

A. hookeri (Moore ex Bedd.) Ching in Fl. Reip. Pop. Sin. 2 : 291. 1959. *Acrophorus hookeri* Moore ex Bedd. Ferns Brit. India t. 95. 1865. *Leucostegia hookeri* (Moore ex Bedd.) Bedd. Handb. Ferns Brit. India 52. 1883.

A. hymenophylloides (Bl.) Copel. Philip. Journ. Sci. 34 : 241. 1927. *Aspidium hymenophylloides* Bl. Enum. Pl. Jav. 172. 1828. *Leucostegia hymenophylloides* (Bl.) Bedd. Ferns Brit. India Suppl. 4. 1876 ; Handb. Ferns Brit. India 54. 1883.

Distrib. : South India.

A. perdurans (Christ) Copel. Univ. Calif. Publ. Bot. 12 : 400. 1931. *Davallia perdurans* Christ, Bull. Herb. Boiss. 6 : 970. 1898.

Distrib. : India (Himalayas), W. China.

A. pseudocystopteris (Kunze) Copel. Phil. Journ. Sci. 34 : 241. 1927. *Leucostegia pseudocystopteris* Kunze in Bot. Zeit. 68. 1850 ; Bedd. Ferns Brit. India t. 92. 1865 ; Handb. Ferns Brit. India 54. 1883. *Davallia pulchra* var. *pseudocystopteris* Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 444. 1880.

Distrib. : India (Himalayas), Nepal.

A. pulchra (D. Don) Copel. Philip. Journ. Sci. 34 : 241. 1927. *Davallia pulchra* D. Don, Prod. Fl. Nepal. 2. 1825 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 444. 1880. *leucostegia pulchra* (D. Don) J. Sm. Lond. Journ. Bot. 1 : 426. 1842 ; Bedd. Handb. Ferns Brit. India 52. t. 25. 1883. *Acrophorus pulchra* sensu Bedd. Ferns South. India t. 10. 1863.

Distrib. : Throughout India in mountainous regions, Nepal, Bhutan, Sri Lanka, Burma.

Davallia Sm. Mém. Acad. Turin 5 : 414. 1793.

Lectotype : **D. canariensis** (L.) Sm. Mém. Acad. Sci. Turin. 5 : 414. t. 9. f. 6. 1793.

D. bullata Wall. ex Hook. Sp. Fil. 1 : 169. t. 50 B. 1846 ; Bedd. Ferns South. India t. 17. 1863 ; Handb. Ferns Brit. India 61. t. 31. 1883 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 445. 1880.

Distrib : Throughout India in mountainous regions, Nepal, Bhutan, Sri Lanka, Burma, Malaya Peninsula. Japan, China, Malay Islands.

D. denticulata (Burm.) Mett. ex Kuhn, Fil. Deck. 27. 1867. *Adiantum denticulatum* Burm. Fl. Ind. 236. 1768. *Davallia elegans* Sw. Schrad. Journ. Bot. 1800 (2) 87 : 1801 ; Bedd. Ferns South. India t. 18, 1863 ; Handb. Ferns Brit. India 59. 1883.

Distrib. : South India, Sri Lanka, Malaya Peninsula.

D. divaricata Bl. Enum. Pl. Jav. 237. 1828 ; Clarke Trans. Linn. Soc. Lond. II. Bot. 1 : 445. 1880 ; Bedd. Handb. Ferns Brit. India 60. 1883. *Davallia polyantha* Hook. Sp. Fil. 1 : 168. t. 59 A. 1846.

Distrib. : North-East India, Burma, Malaya Peninsula, Malay Islands. China.

D. solida (Forst.) Sw. Scharad. Journ. Bot. 1800 (2) : 87. 1801 ; Bedd. Ferns Brit. India t. 104. 1865 ; Handb. Ferns Brit. India 468. 1966. *Trichomanes solidum* Forst. Prod. 86. 1786.

Distrib. : India (Eastern Himalayas), Burma, Malaya Peninsula, Malay Islands, Polynesia.

D. trichomanoides Bl. Enum. Pl. Jav. 238. 1828.

Distrib. : India (Eastern Himalayas) Malaya Peninsula, Malay Islands, New Guinea.

Humata Cav. Desc. 272. 1802.

Lectotype : **H. ophioglossa** Cav. Descr. 273. 1802.

H. assamica (Bedd.) C. Chr. Contr. U.S. Nat. Herb. 26 : 293. 1931. *Acrophorus assamicus* Bedd. Ferns. Brit. India t. 94. 1866. *Davallia assamica* (Bedd.) Bak. Syn. Fil. 452. 1868 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 443. 1880. *Leucostegia assamica* (Bedd.) J. Sm. Hist. Fil. 84. 1875 ; Bedd. Hand. Ferns Brit. India 51. 1883.

Distrib. : India (Himalayas) S.W. China.

H. griffithiana (Hook.) C. Chr. Contr. U.S. Nat. Herb. 26 : 293. 1931.
Davallia griffithiana Hook. Sp. Fil. 1 : 168. t. 493. 1846 ; Bedd. Ferns Brit. India t. 106. 1865 ; Handb. Ferns Brit. India 60. 1883.

Distrib. : India (Meghalaya) Bhutan, Burma, S. China, Formosa.

H. heterophylla (Sm.). Desv. Prod. 323. 1825 ; Thothathri : et al. in Journ. Bomb. Nat. Hist. Soc. 74 : 251. 1977. *Davallia heterophylla* Sm. Mém. Acad. Turin 5 : 415. 1793.

Distrib. : India (Great Nicobar Islands) : Malesia, Polynesia.

H. repens. (L.f.) Diels, Nat. Pfl. : 1 (4) : 209. 1999. *Adiantum repens* L. f. Suppl. 446. 1781. *Davallia pedata* J. Sm. Mém. Acad. Turin 5 : 415. 1793 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 442. 1880. *Humata pedata* J. Sm. Journ. Bot. 3 : 416. 1841 ; Bedd. Ferns South. India t. 12. 1863 ; Handb. Ferns Brit. India 48. t. 23. 1883.

Distrib. : South-East. India, Sikkim ; Malay Peninsula, Malay Islands, Australia, Sri Lanka, N. Australia, China, Japan, Mascareen Islands.

Leucostegia Presl, Tent. Pterid. 94. t. 4. f. 11. 1836.

Type : **L. immersa** (Wall ex Hook.) Presl, Tent. Pterid. 95 t. 4. f. 11. 1836 ; Bedd. Ferns South. India t. 11. 1863 ; Handb. Ferns. Brit. India 51. 1883. *Davallia immersa* Wall. ex Hook. Sp. Fil. 1 : 156. 1846 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 443. 1880.

Distrib. : Throughout India in mountainous regions, Indo-China, W. China, Formosa, Malay Peninsula, Philippines, Malay Islands, New Guinea.

Paradvalloides Ching, Acta Phytotax. Sin. 11 : 18. 1966.

Type : **P. multidentatum** (Wall. ex Hook. et Bak.) Ching, Acta Phytotax. Sin. 11 : 18. 1966.

P. membranulosum (Wall. ex Hook.) Ching, Acta Phytotax. Sin. 11. 1966. *Davallia membranulosa* Wall. ex Hook. Sp. Fil. 1 : 159. t. 53 A. 1846 ; Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 443, 1880. *Leucostegia membranulosa* J. Sm. Hist. Fil. 84. 1875 ; Bedd. Ferns Brit. India t. 98. 1865. Handb. Ferns Brit. India 50. 1883.

Distrib. : E.W. India, Nepal, China.

P. multidentatum (Wall. ex Hook. et Bak.) Ching, Acta Phytotax.

Sin. 11 : 18. 1966. *Davallia multidentata* (Wall. ex Hook. et Bak.) *Lecuostegia multidentata* wall. ex Hook Syn. Fil. 91, 1867. Bedd. Handb. Ferns Brit. India 50. 1883. *Microlepia pteropus* Bedd. Ferns Brit. India t. 313. 1869.

Distrib. : India (Himalayas) S.W. China.

GYMNOGRAMMITIDACEAE Ching

Gymnogrammitis Griff. Icon. Pl. Asiat. 2 : 129. 1849.

Type : **G. dareiformis** (Hook.) Ching, ex Tard. et Christ, Notul Syst. Paris 6 : 2. 1937.

G. dareiformis (Hook.) Ching ex Tard. et Christ, Notul. Syst. Paris 6 : 2. 1937. *Polypodium dareiformis* Hook. 2nd Cent. Ferns t. 24. 1860 ; Bedd. Handb. Ferns Brit. India 316. t. 169. 1883. *Acrophorus hookeri* sensu Bedd. Ferns Brit. India t. 95. 1865.

Distrib. : India (Himalayas).

BLECHNACEAE (Presl) Copel.

Blechnidium Moore, Octava Nat. Print. Brit. Ferns 2 : 210. 1860.

Type : **B. melanopus** (Hook.) Moove, Octava Nat. Print. Brit. Ferns 2 : 210 : 1860.

B. melanopus (Hook.) Moove, Octava Nat. Print. Brit. Ferns 2 : 210. 1860. Bedd. Ferns Brit. India t. 50. 1865 ; Handb. Ferns Brit. India 133. t. 67. 1883. *Blechnum melanopus* Hook. Sp. Fil. 3 : 64. t. 161. 1860.

Distrib. : India (Meghalaya).

Blechnum L. Sp. Pl. 2 : 1077. 1753.

Lectotype : **B. occidentale** L. Sp. Pl. 2 : 1077. 1753 (*Orientale* ex err.) ed. II. 2 : 1524. 1763.

B. orientale L. Sp. Pl. 2 : 1077. 1753 ; Clarke Trans. Linn. Soc. Lond. II. Bot. 1 : 474. 1880 ; Bedd. Ferns South. India t. 29. 1863 ; Handb. Ferns Brit. India 132. t. 86. 1883.

Distrib. : Throughout India, Sri Lanka, China, Malaya. Peninsula, Malesian Islands. Polynesia, Australia.

Brainea J. Sm. Cat. Kew Ferns Gard. 5. 1856.

Type : **B. insignis** (Hook.) J. Sm. Journ. Bot. 5 : 237. t. 2. 1856.

B. insignis (Hook.) J. Sm. Journ. Bot. 5 : 237. t. 2. 1856 ; Bedd. Ferns Brit. India t. 139. 1867 ; Handb. Ferns Brit. India 396. t. 230. 1883. *Bowringia insignis* Hook. Journ. Bot. 5 : 237. t. 2. 1853.

Distrib. : India (Meghalaya) Burma, Malay Peninsula, Hong Kong.

Doodia R. Br. Prod. Fl. Nov. Holl. 151. 1810.

Lectotype : **D. aspera** R. Br. Prod. Fl. Nov. Holl. 151. 1810

D. aspera R. prod. Br. Fl. Nov. Holl. 151. 1810 ; Bedd. Ferns Brit. India t. 222. 1865 : Handb. Ferns Brit. India 137. t. 69. 1883.

Distrib. : India (Tamil Nadu), Sri Lanka, Java.

Woodwardia Sm. Mém. Acad. Sc. Turin 5 : 411. t. 9. f. 3. 1793.

Lectotype : **W. radicans** (L.) Sm. Mém. Acad. Sci. Turin 5 : 412. 1793.

W. orientalis Sw. Schrad. Journ. Bot. 1800. (2) : 76. 1801.

Distrib. : India (Himalayas), Tonkin, China, Formosa, Japan, Philippines.

W. unigemmata (Makino) Nakai, Bot. Mag. Tokyo 39 : 103. 1925. *W. radicans* (L.) Sm. Var. *unigemmata* Makino, Journ. Jap. Bot. 2 : 7. 1818. *W. redicans* sensu Bedd. Handb. Ferns Brit. India 135 t. 68. 1883.

Distrib. : India (Himalayas), China, Japan, Tonkin, Formosa, Philippines.

STENOCHLAENACEAE Ching

Stenochlasna J. Sm. Journ. Bot. 3 : 401. 1841.

Lectotype **S. scandens** (Sw.) J. Sm. in Hook. Journ. Bot. 3 : 401. 1841.

S. palustris (Burm.) Bedd. Ferns Brit. India Suppl. 26. 1876 ; Handb. Ferns Brit. India 421. t. 253. 1883. *Polypodium palustre* Brum. Fl. Ind. 234. 1768. *Acrostichum palustre* (Burm) Clarke, Trans. Linn. Soc. Lond. II. Bot. 1 : 577. 1880. *Stenochlaena scandens* (Sw.) J. Sm. in Hook. Journ. Bot. 3 : 401. 1841 ; Bedd. Ferns South. India t. 201. 1867.

Distrib. : India (Western Ghats, West Bengal), Sri Lanka, China, Queensland, Fiji.

AZOLLACEAE Wett.

Azolla Lam. Encycl. Méth. 1 : 343. 1783.

Type : **A. filiculoides** Lam. Encycl. Méth. Bot. 1 : 343. 1783.

A. imbricata (Roxb. ex Griff.) Nakai, Bot. Mag. Tokyo 39 : 185. 1925. *Salvinia imbricata* Roxb. ex Griff. Calc. Journ. Sci. 4 : 470. 1844.

Distrib. : Throughout India, Nepal, China, South Korea, Formosa, Ryukyu, Japan.

A. pinnata R. Br. Prod. Fl. N. Holl. 167. 1810.

Distrib. : Throughout India, Japan, Asia, Australia, Africa.

SALVINIACEAE Seguier

Salvinia Séguier, Fl. Veron. 3 : 52. 1754.

Type : **S. natans** (L.) All. Pedem. 2 : 269. 1785.

S. auriculata Aubl. Hist. Pl. Guian 2 : 969. t. 367. 1775 ; Kamathy, Sci. & Cult. 34. 396. 1968.

Distrib. : India (Kerala), Trop. America.

S. cucullata Roxb. ex Bory, Bél. Voy. Bot. 2 : 6. 1833 ; Roxb. ex Griff. Calc. Journ. Nat. Hist. 4 : 470. 1844.

Distrib. : Eastern India, Western Australia.

S. molesta Mitch. Br. Fern Gaz. 10. 251. 1973.

Distrib. : India (Kerala).

S. natans (L.) All. Fl. Pedem. 2 : 289. 1785. *Marsilea natans* L. Sp. Pl. 2. 1099. 1753.

Distrib. : Throughout India, Warm regions of Europe and S. E. Asia.

INDEX TO GENERA

A

- Acrophorus* Presl 145
Acrorumohra (H. Ito) H. Ito 137
Acrostichum L. 73
Actiniopteris Link 68
Actinostachys Wall. ex Hook. 58
Acystopteris Nakai 124
Adiantum L. 74
Ampelopteris Kunze 102
Amphineuron Holtt. 103
Anemia Sw. 59
Angiopteris Hoffm. 24
Anisocampium Presl 125
Anogramma Link 77
Antrophyum Kaulf. 80
Arachniodes Bl. 146
Araiostegia Copel. 168
Arthromeris (Moore) J. Sm. 34
Arthropteris J. Sm. ex Hook. 166
Asplenium L. 116
Athyriopsis Ching 125
Athyrium Roth 125
Azolla Lam. 174

B

- Belvisia* Mirb. 35
Blechnidium Moore 172
Blechnum L. 172
Bolbitis Schott 161
Botrychium Sw. 21
Brainea J. Sm. 173

C

- Cephalomanes* Presl 90
Ceratopteris Ad. Brongn. 83
Ceterach Willd. 123
Ceterachopsis (J. Sm.) Ching 123
Cheilanthes Sw. 63

- Cheilanthopsis* Hieron. 124
Christella Lev. 103
Christensenia Maxon 26
Christopteris Copel. 36
Cibotium Kaulf. 92
Colysis Presl 36
Coniogramme Féé 77
Cornopteris Naki 130
Coryphopteris Holtt. 107
Crepidomanses Presl 91
Crypsinus Presl 37
Cryptogramme R. Br. 66
Ctenitis (C. Chr.) C. Chr. ex Tard.
et C. Chr. 137
Ctenitopsis Ching ex Tard. et C.
Chr. 139
Ctenopteris Bl. 31
Cyathea Sm. 92
Cyclogramma Tagawa 107
Cyclosorus Link 108
Cyrtomium Presl 139
Cystopteris Bernh. 130

D

- Davallia* Sm. 170
Dendroglossa Presl 37
Dennstaedtia Bernh. 94
Diacalpe Bl. 146
Dicranopteris Bernh. 30
Dictyocline Moore 108
Dictyodroma Ching 131
Didymochlaena Desv. 140
Diphasiastrum Holub 10
Diplaziopsis C. Chr. 131
Diplazium Sw. 131
Dipteris Reinw. 57
Doodia R. Br. 173
Doryopteris J. Sm. 61

- Drymoglossum* Presl 38
Drymotaenium Mak. 38
Drynaria (Bory) J. Sm. 58
Dryoathyrium Ching 135
Dryopteris Adanson 148

E

- Egenolfia* Schott 163
Elaphoglossum Schott 164
Emidopteris Ching et Wu 95
Equisetum L. 19

G

- Glaphyropteridopsis* Ching 108
Gleichenia Smith 29
Goniophlebium Presl 38
Gonocarmous v. d. Bosch 85
Grammitis Sw. 32
Gymnocarpium Newm. 135
Gymnogrammitis Griff. 172

H

- Helminthostachys* Kaulf. 20
Hemionitis L. 78
Histiopteris (Agardh) J. Smith 95
Holcosorus Moore 39
Humata Cav. 170
Huperzia Bernh. 7
Hymenophyllum Sm. 85
Hypodematum Kunze 137
Hypolepis Bernh. 97

I

- Idiogramma* Ghosh 79
Isöetes L. 18

K

- Kuniwatsukia* Pic. Ser. 136

L

- Lastreopsis* Ching 141
Lemmaphyllum Presl 39
Lepidogrammitis Ching 39
Lepisorus (J. Sm.) Ching 39
Leptochilus Kaulf. 43
Leptogramma J. Sm. 161
Leucostegia Presl 171
Lindsaea Dryand. ex Smith 98
Lithostegia Ching 146
Loxogramme (Bl.) Presl 33
Lunathyrium Koidz. 136
Lycopodiastrum Holub ex Dixit 10
Lycopodium L. 9
Lygodium (L.) Sw. 59

M

- Macrothelypteris* (Ito) Ching 109
Marattia Sw. 26
Marginaria Bory 44
Marsilea L. 84
Matteuccia Todaro 123
Mecodium Presl ex Copel. 86
Meringium Presl 87
Metapolypodium Ching 44
Metathelypteris (Ito) Ching 109
Microgonium Presl 88
Microlepia Presl 95
Microsorium Link 44
Microtrichomanes (Prantl) Copel. 88
Mildella Trev. 62
Monachosorum Kunze 97

N

- Neocheiropteris* Christ 47
Nephrolepis Schott 166
Notholaena R. Br. 62
Nothoperanema (Tagawa) Ching 155

O

- Oleandra* Cav. 167
Onychium Kaulf. 67
Ophioderma (Bl.) Endl. 22
Ophioglossum L. 22
Oreopteris Holub 110
Osmunda L. 27

P

- Palhinhaea* (L.) Franco et Vasc. 10
Paradavalloides Ching 171
Paraleptochilus Copel. 47
Parathelypteris (H. Ito) Ching 110
Pellaea Link 62
Peranema D. Don 146
Phegopteris (Presl) Fée 110
Phlebodium (R. Br.) J. Sm. 48
Phlegmariurus (Herter) Holub 9
Phymatopteris Pic. Ser. 48
Phymatosorus Pic. Ser. 50
Pityrogramma Link 79
Plagiogyria (Kunze Mett. 27
Platycerium Desv. 57
Pleocnemia Presl 141
Pleuromanes (Presl) Presl 89
Pneumatopteris Nakai 110
Polypodiastrum Ching 51
Polypodioides Ching 51
Polypodium L. 52
Polystichum Roth 156
Pronephrium Presl 32
Prospitia Presl. 32
Pseudocyclosorus Ching 111
Pseudodrynaria C. Chr. 52
Pseudophegopteris Ching 113
Psilotum Sw. 20
Pteridium Gled. ex Scop. 98
Pteridrys C. Chr. et Ching 141
Pteris L. 68
Pyrrosia Mirf. 53

Q

- Quercifilix* Copel. 142

S

- Salvinia* Ség. 174
Schizaea Sm. 59
Scleroglossum v. Ald. v. Ros. 32
Selaginella P. Beauv. 11
Selenodesmium (Prantl) Copel. 89
Sinephropteris Mickel 123
Sphaerocionum Presl 89
Sphaerostephanos J. Sm. 114
Sphenomeris Maxon 102
Stenogramma Bl. 114
Stenochlaena J. Sm. 173
Syngamma J. Sm. 79

T

- Taenitis* Willd. ex Spring 83
Tapeinidium (Presl) C. Chir. 102
Tectaria Cav. 142
Thelypteris Schmid. 114
Thylacopteris Kunze ex J. Sm. 57
Triblemma (J. Sm.) Ching 136
Tricholepidium Ching 57
Trigonospora Holtt. 115

V

- Vandenboschia* Copel. 89
Vittaria Sm. 81

W

- Woodsia* R. Br. 124
Woodwardia Sm. 173

X

- Xiphopteris* Kaulf. 33