

TYPE COLLECTIONS IN THE CENTRAL NATIONAL HERBARIUM

VOLUME 2

U. P. SAMADDAR



BOTANICAL SURVEY OF INDIA


**TYPE COLLECTIONS IN THE
CENTRAL NATIONAL HERBARIUM**

VOLUME 2

TYPE COLLECTIONS IN THE CENTRAL NATIONAL HERBARIUM

VOLUME 2

U. P. SAMADDAR



BOTANICAL SURVEY OF INDIA

© Government of India, 1991

Date of Publication : August 1, 1991

Price :

Published by the Director, Botanical Survey of India, P-8, Brabourne Road, Calcutta-700 001 and Printed at Prabartak Printing & Halftone Ltd., 52/3, B. B. Ganguly Street, Calcutta-700 012.

AUTHOR'S NOTE

This second attempt in the series on the enumeration of Type Collections in the Central National Herbarium (CAL) forms yet another significant contribution in the pursuit of developing a National Data Base as envisaged in the review of the functions and organisation of the Botanical Survey of India.

The progress of the scientific activity of floristic survey in the Indian Subcontinent revealed an increase in finding of new species, sub-species and varieties in addition to the vast collections made earlier and deposited in CAL. In the present book Types of both Cryptogams and Angiosperms are enlisted. The information here are supplemented with nomenclature, synonyms, references, localities, altitudes, collection dates, and numbers which forms this treatise.

Apart from new species, sub-species and varieties, the plant genera (Angiosperms) new to science included here (inter alia) are : *Asraoa*, *Eskemukerjea*, *Kanjarum*, *Kurziodendron*, *Neodriessenia*, *Parakaempferia*, *Pauella*, *Pseudodissochaeta*, *Janakia* and *Santapaua*.

New taxa of cryptogams described (inter alia) include-mosses : *Andreaea kashyapii*, *Hydrogonium mussoorianum*, *Habrodon kashmirensis*, *Plagiothecium dehradunense*, *Lescuraea darjeelingensis*, *Haplocladium microphyllum* ssp. *capillatum* var. *bhutanicum*, *Hygrohypnum nairii*, *H. chophrae*, *Brachythecium chakratense*, *B. garhwalense*, *Pseudoleskeia laevifolia* var. *vasudharensis*, *Thuidium contortulum* var. *mussoriensis*, *T. squarrosum* var. *scabrisetum* and *Rhynchostegium calderii*; Pteridophytes: *Polysticum levingi*, *P. lanatum* var. *bifurcatum*, *Thelypteris multilineata* var. *bhutanica*, *Plagiogyria meghalayensis*, *P. elongata*, *Pteris silent-valleyensis*, *Elaphoglossum jowaiense*, *E. cherapunjii*, *E. thomsonii*, *E. prainii* and *E. simonsianum*; Lichens: *Graphina darjeelingensis* and *G. awasthi*.

Among the names of the botanists mentioned in this book for their valuable contribution in discovering new taxa are Nair, Vohra, Panigrahi, Dixit, Ghosh, Roy Chowdhury, Nayar, Jain, Deb, Thothathri, Majumdar, Babu, Rakshit, Bhattacharyya, Mukerjee, Subbarao, Banerjee, T. Chakraborty, H. Chakraborty, Malick, Shetty, Sivarajan, Santapau, Rhaghavan, Joseph, Ansari, Basappa, Muniyamma, Ellis, Rolla Rao, Henry, Billore, Hajra, Ramamurthy, A. S. Rao, Sastry, Balakrishnan, Subramaniam, Chandrasekharan, Sanjappa, Purohit, Vasudeva Rao, Sur, Gangopadhyay, Giri, Maiti, Dutta, T. Paul, Singh, etc.

Types of some species described earlier by workers like J. D. Hooker, Watt, Roxburgh, King, Kurz, Clarke, Sprengel, D. Don, O. Kuntze, Gage, Edgeworth, Mattfield, Prain, Thwaites, C. de Candolle, A. de Candolle, Hoogland, Wright, Koehne and Stapf are also included.

The vast collections in the Central National Herbarium (CAL) represent the Flora of India and many other countries of the world. This priceless botanical treasure plays a valuable role in teaching and research activity in the country in identifying botanical specimens and in the efforts of revisionary accounts of Indian plants with the help of its authentic materials.

It is further hoped that this attempt to enlist the Type collections for building the National Data Base on the Flora of India also helps in effective and speedy progress on the taxonomic revisions of the different flowering plant families being worked out in the Botanical Survey of India.

I consider it is a great privilege to record my indebtedness to Dr. M. P. Nayar former Director, BSI, for his suggestions and constant advice. In the preface to the earlier publication on the Type Collections in *CAL* his emphasis on the urgency of listing all the type materials gave boost to the author in continuing the work. I am grateful to Dr. B. D. Sharma, the present Director of the institution for his supervising the work. I thank Shri A. R. K. Sastry, Scientist SE, In-charge, Publication Section, Shri S. C. Pal, Publication Officer, Shri Pankaj Kr. Dutta, Shri Haramohon Mukherjee, Shri Samiran Roy of Publication Unit for speedy completion of the work and Shri Swapan Kumar Das for rendering help with typing the manuscript.

Type Collections in the Central National Herbarium

Abelmoschus tuberculatus Pal et Singh var. **deltoidefolius** Paul et Nayar (Malvaceae).

Bull. Bot. Surv. India 24(1-4) : 215. 1982 (1983); Fasc. Fl. Ind. 19 : 79. 1988.

Holotype : India : Rajasthan: Jhalawar, Sept. 19, 1964, *B. M. Wadhwa et D. M. Verma* 7581.

Paratype : Ibid., Sept. 21, 1964, *B. M. Wadhwa et D. M. Verma* 7524.

Abutilon neelgherrense Munro ex Wt. var. **fischeri** T. K. Pal et Nayar (Malvaceae).

Bull. Bot. Surv. India 25 (1-4): 183-184. 1983 (1985); Fasc. Fl. Ind. 19 : 90. 1988.

Holotype : India : Tamil Nadu : Coonoor river beach, Coimbatore Dist., 12000 ft. (363 m), Jan. 10, 1910, *C. E. C. Fischer* 1581.

A. ranadei Woodr. et Stapf (Malvaceae).

Kew Bull. 99. 1894 ; Fasc. Fl. Ind. 19: 95. 1988.

Isotype : India : Maharashtra : Ratnagiri, Ambaghat, *N. B. Ranade s. n.*

Acer acuminatum Wallich ex D. Don (Aceraceae).

Prodr. Fl. Nepal 249. 1825 ; Fasc. Fl. Ind. 9 : 4. 1982.

Type : India : Kashmir, *Wallich* 1225 p. p.

A. caesium Wallich ex Brandis (Aceraceae).

For. Fl. N. W. & C. Ind. III. tab. 21. 1874 non *A. caesium* (Reinward ex Bl.) Kostermans 1965 ; Fasc. Fl. Ind. 9 : 4. 1982.

Type : N. W. Himalaya : 7-10,000 ft., *Thomas, Thomson s. n.*

A. campbellii Hooker f. & Thomson ex Hiern var. **campbellii** (Aceraceae).

Hook. f. Fl. Brit. Ind. 1 : 696. 1875 ; Fasc. Fl. Ind. 9 : 5. 1982.

Type : Sikkim Himalaya : *Griffith* 926.

A. campbellii Hooker f. & Thomson ex Hiern var. **serratifolium** Banerji (Aceraceae).

J. Bombay Nat. Hist. Soc. 58(1) : 306. 1961; Fasc. Fl. Ind. 9 : 6. 1982.

Type : India : Sikkim, *J. H. Lace* 2250.

A. caudatum Wallich (Aceraceae).

Pl. Asiat. Rar. 2 : 4. 1830 & 2 : 28. t. 132. 1831; Fasc. Fl. Ind. 9 : 8. 1982.

Type : Kashmir : *Wallich* 1225 p.p.

Acer hookeri Miquel var. **hookeri** (Aceraceae).

Arch. Neerl. Sci. Nat. 2 : 471. 1852; Fasc. Fl. Ind. 9 : 9. 1982.

Type : Sikkim : 8-10,000 ft., *Hooker* 3.

A. hookeri Miquel var. **majus** Pax (Aceraceae).

Bot. Jahrb. 7 : 216. 1886; Fasc. Fl. Ind. 9 : 10. 1982.

Type : India : Sikkim : Rirhi to Rinchinpong, 2-5000 ft., Oct. 1, 1862, *Anderson* 407.

A. laevigatum Wallich (Wall. Cat. No. 1223. 1829 *nom. nud.*) (Aceraceae).

Pl. Asiat. Rar. 2 : 3-4. t. 104. 1830 ; Fasc. Fl. Ind. 9 : 10. 1982.

Type : Nepal : *Wallich* 1223.

A. oblongum Wallich ex DC. var. **oblongum** (Aceraceae).

Prodr. 1 : 593. 1824 ; Fasc. Fl. Ind. 9 : 12. 1982.

Type : Nepal : *Wallich* 1222.

A. oblongum Wallich ex DC. var. **membranaceum** Banerji (Aceraceae).

Phytologia 9(5) : 265-266. 1963 ; Fasc. Fl. Ind. 9 : 14. 1982.

Type : India : Uttar Pradesh : Mussoorie, Jharipani, March 29, 1957, *Banerji s. n.*

A. oblongum Wallich ex DC. var. **microcarpum** Hiern (Aceraceae).

Hook. f. Fl. Brit. Ind. 1 : 693. 1875; Fasc. Fl. Ind. 9 : 12. 1982.

Type : India : Arunachal Pradesh, *Griffith* 935.

A. osmastonii Gamble (Aceraceae).

Bull. Misc. Inforin. Roy. Bot. Gard. Kew. 446. 1908; emend Delendick in Brittonia 30(4) : 473-476. 1978 ; Fasc. Fl. Ind. 9 : 14. 1982.

Type : India : W. Bengal : Darjeeling, Salombong, Oct. 27, 1903, *B. B. Osmaston s. n.*

A. pectinatum Wallich (Wall. Cat. No. 1226. 1829. *nom. nud.*) (Aceraceae).

Nicholson in Gard. Chron. 1881(1) : 365. t. 69. 1881 nom. Illegit. ex Pax in Bot. Jahrb. 7 : 249. 1886; Fasc. Fl. Ind. 9 : 15. 1982.

Type : Nepal : Gossain Than on the Himalayas, Aug. 1821, *Wallich* 1226.

A. pentapomicum J. L. Stewart ex Brandis (Aceraceae).

For. Fl. N. W. & C. Ind. 111. 1874; Fasc. Fl. Ind. 9 : 15. 1982.

Syntypes : Jhelum Valley: 3-5000 ft., *J. L. Stewart* 648; Hazara, Khajan Valley, 4-5000 ft., *J. L. Stewart* 541, 542.

***Acer sikkimense* Miquel var. *sikkimense* (Aceraceae).**

Arch. Neerl. Sci. Nat. 2 : 471. 1852 ; Fasc. Fl. Ind. 9 : 16. 1982.

Type : India : Sikkim, 7-9000 ft., Hooker 4.

***A. sikkimense* Miquel var. *serrulatum* Pax (Aceraceae).**

Bot. Jahrb. 7 : 215. 1886; Fasc. Fl. Ind. 9 : 16. 1982.

Type : India : Arunachal Pradesh: Mishmee Hills, Griffith 936.

***A. stachyophyllum* Hiern (Aceraceae).**

Hook. f. Fl. Brit. Ind. 1: 694. 1875 ; Fasc. Fl. Ind. 9 : 16. 1982.

Type : India : Sikkim, 9000 ft., Hooker 5.

***A. sterculiaceum* Wallich (Wall. Cat. No. 1224. 1829. *nom. nud.*) (Aceraceae).**

Pl. Asiat. Rar. 2 : 3. t. 105. 1830 ; Fasc. Fl. Ind. 9 : 17. 1982.

Type : Nepal: 1921, Wallich 1224.

***A. thomsonii* Miquel (Aceraceae).**

Arch. Neerl. Sci. Nat. 2 : 470. 1867 ("Thomsonii") ; Fasc. Fl. Ind. 9 : 18. 1982.

Type : India : Sikkim, Thomson s. n.

***Aconitum gammiei* Stapf (Ranunculaceae).**

Kew Bull. 1907 : 56.

Holotype : India : Sikkim Himalaya, Chola Range, 12500 ft., Sept. 27, 1892, Fls. Blue, G. A. Gammie 1310, Acc. No. 7853.

***Aconogonium kuttienne* Maiti, Dutta and Babu (Polygonaceae).**

J. Bombay Nat. Hist. Soc. 77(2) : 303. 1980.

Holotype : India : Uttar Pradesh : Kuttii, Sept. 22, 1968, R. M. Dutta 277A.

Isotype : Ibid., R. M. Dutta 277B.

Paratypes : Ibid., 14-15000 ft., July 31, 1886, J. F. Duthie 5929; Kuttii Valley, way to Samzurkchan Glacier, 3 km from Kuttii village, C. 4000 m, Oct. 24, 1976, G. G. Maiti 790.

***A. pangianum* Pal et Maiti (Polygonaceae).**

Bull. Bot. Surv. India 26 (1 & 2) : 95. 1984 (1985).

Holotype : India : Arunachal Pradesh : Subansiri Dist., Pange, 3000 m, April 19, 1980, G. D. Pal 77734.

Isotype : G. D. Pal 77734,

***Acroblastum ambavanense* Reddi (Balanophoraceae).**

Willdenowia 5 : 389. 1969.

Holotype : India : Maharashtra : Poona Dist., in the Kate pani forest near Ambavane village, on the Western Ghats, 1050 m, growing on the hill slopes on the roots of *Carissa congesta* Wight and among bushes of *Carvia callosa* (Nees) Brem, Oct. 26, 1964, Reddi 99494a.

***Acronema johrianum* Babu (Apiaceae).**

Brittonia 25 : 159-161. 1973.

Holotype : Nepal : Ghopte, Gosainthan, 4290 m, flowers white, open place, Sept. 9, 1965, Shrestha & Sakya 3740.

***Actinocarya acaulis* (W. W. Smith) I. M. Johnston (Boraginaceae).**

Journ. Arnold Arb. 21 : 53. 1940 ; Ibid., 52 : 354. 1971.

Holotype : India : Sikkim : Naku La, Llonakh, top of Pass, 17,500 ft., Aug. 3, 1909, Smith & Cave 2811.

***Adiantum indicum* Ghatak (Adiantaceae).**

Bull. Bot. Surv. India 5(1) : 71. 1963.

Holotype : India : W. Bengal: Belgharia, a suburb of Calcutta (22°34'N, 88°22'E), 9 m, March 11, 1957, J. Ghatak J. 301.

A. *malesianum* Ghatak (Adiantaceae).

Bull. Bot. Surv. India 5(1) : 73. 1963.

Holotype : Malaya Peninsula, Oct. 4, 1958, J. Ghatak J 350.

A. *ramyam* Nayar et Geevarghese (Adiantaceae).

Bull. Bot. Surv. India 28 : 133. 1986 (1988).

Type : South India : Vellamunda (Wynad), Feb. 14, 1982, B. K. Nayar et P. V. Madhusudan K 11637.

***Adinandra gallatlyii* Paul (Theaceae).**

Ind. J. For. 2(1) : 20. 1979.

Holotype : Burma : 1877, Geo. Gallatly 671A.

Isotypes : Ibid., Geo. Gallatly 671B-C.

***Agapetes subansirica* Pal et Thoth. (Vaccinaceae).**

Bull. Bot. Surv. India 26(3 & 4) : 239. 1984 (1986).

Holotype : India : Arunachal Pradesh : Subansiri Dist., Begi, 190 m, April 24, 1980, G. D. Pal 78214.

Agrostis munroana Aitch. et Hemsley ssp. **indica** Sunanda Bhattacharya et Jain (Poaceae).

Bull. Bot. Surv. India 25(1-4) : 205. 1983 (1985).

Holotype : India, Himachal Pradesh: Karcham-Brue road, nearly 2 km from Baspa Valley, Aug. 27, 1973, *Janardhanan* 52862A.

Paratypes : India : Jammu & Kashmir: Gulmarg, 3000 m, Aug. 18, 1958, *Nanda* 1599 ; Sangam Valley above Liddarwat 3500-4000 m, Aug. 10, 1893, *Duthie* 13535 ; Yamhara Pass, 4000-4300 m, Aug. 13. 1893, *Duthie* 13582; Himachal Pradesh: Chamba, 3200 m, Nov. 9, 1898, *Lace* 1856; Uttar Pradesh: Kumaon, Dhaul Valley, 2500-3000 m, Aug. 7, 1886, *Duthie* 6231.

A. tungnathii Sunanda Bhattacharya et Jain (Poaceae).

Bull. Bot. Surv. India 25(1-4) : 204. 1983 (1985).

Holotype : India : Uttar Pradesh: Chamoli, on way to Tungnath, 4000 m, Aug. 29, 1978, *Bhattacharya* 1524A.

Agrostistachys hookeri (Thw.) Benth. (Euphorbiaceae).

Benth. & Hook. f. Gen. Pl. 3 : 303. 1880; J. Bombay Nat. Hist. Soc. 67(2) : 305. 1970.

Sarcoclinium hookeri Thw.

Enum. Pl. Zeyl. 4 : 279. 1861.

Type : Ceylon: *Thwaites* C. P. 3429 ABC.

A. indica Dalzel var. **longifolia** Muell.-Arg. (Euphorbiaceae).

DC. Prodr. 15(2) : 726. 1866; J. Bombay Nat. Hist. Soc. 67(2) : 303. 1970.

A. longifolia (Muell.-Arg.) Kurz

Prelim. Rep. Veg. Pegu, App. A : 111. APP. B : 79. 1875.

A. gaudichaudii Hook. f.

Fl. Brit. Ind. 5 : 406. 1887.

A. indica ssp. *longifolia* (Muell.-Arg.) Pax & Hoffm.

Engl. Pflan. 57 : 107. 1912.

Type : Burma : Tenasserim, *Helper* 4921.

A. meeboldii Pax & Hoffm. (Euphorbiaceae).

Engl. Pflan. 57 : 100. 1912 ; J. Bombay Nat. Hist. Soc. 67(2) : 304. 1970.

Sarcoclinium longifolium Wight

Ic. Pl. Ind. Or. 5 : 24. t. 1887-1888. 1852.

A. longifolia (Wight) Benth. ex Hook. f.

Fl. Brit. Ind. 5 : 407. 1887 (non Kurz 1875).

Type : Tamil Nadu : Courtallam, *Wight* 2612.

Aithyrium andersoni (C. B. Clarke) Panigrahi et S. K. Basu (Aithyraceae).
 Proc. 72nd Session; Ind. Sci. Cong. Lucknow, Sec. Botany, Abst.:
 14. 1985; Bull. Bot. Surv. India 27(1-4) : 115. 1985 (1987).

Asplenium andersonii Clarke
 J. Linn. Soc. 15 : 154. 1876. P. P. excl. var.
Syntypes : Sikkim : 1862, *Anderson* 73, 1511.

Alysicarpus roxburghianus Thoth. et Pramanik (Fabaceae).
 Bull. Bot. Surv. India 21 : 189. 1979 (1981).
Holotype : India : Shevaroi Hills, *Perrottet* 272A.
Isotypes : *Perrottet* 272 B & C.

A. vasavadae Hemadri (Fabaceae).
 Indian For. 97(1) : 65-67. 1971.
Holotype : India : Maharashtra : on the open slopes of Shivneri Hill,
 Junnar, Poona, Nov. 1, 1964, *Hemadri* 104350A.
Isotype : *Hemadri* 104350F.

Anaphalis cavei Chatterjee (Compositae).
 Bull. Bot. Soc. Beng. 3(1) : 17. 1949.
Isotype : India : Sikkim: Naku Chu, 5300 m, Sept. 25, 1915, *G. H. Cave*
 1486.

Andreaea kashyapii Dix. ex Vohra et Wadhwa (Andreaeaceae).
 Bull. Bot. Surv. India 6 : 321. 1964 (1965).
Isotype : Western Tibet : Manasarowar, July 18, 1926, *Kashyap* s. n.

Andrographis nallamalayana Ellis (Acanthaceae).
 Bull. Bot. Surv. India 8 : 362. 1966 (1967).
Holotype : India : Andhra Pradesh : Ahobilam, Nallamalais, Karnool
 Dist., Aug. 29, 1965, *Ellis* 25546A.

A. neesiana Wt. var. **rotundifolia** Sreem. (Acanthaceae).
 Bull. Bot. Surv. India 8 : 91. 1966 (1967).
Holotype : India : Tamil Nadu : Boluvampatti Hills, Coimbatore
 Dist., 1100 m, Sept. 24, 1963, *Sreemadhavan* 862A.

Aneilema glanduliferum Joseph et Rolla Rao (Commelinaceae).
 J. Indian bot. Soc. 47(3-4) : 367. 1969.
Holotype : India : Arunachal Pradesh : (NEFA), Tipi, North Bhalukpong, 1250 m, on way to Krishna, along wet rocky slopes near
 bank of river Bharali, Kameng Dist., 92°38'E., 27°1'N., Sept. 10, 1964,
Joseph 39707A.

Anemone rauii Goel et Bhattach. (Ranunculaceae).

J. Bombay Nat. Hist. Soc. 79(3) : 641. 1983.

Holotype : India : Uttar Pradesh : Tehri Dist., Gangi, open grassy slopes, 3000 m, Aug. 12, 1978, A. K. Goel 64419.

Angelica sikkimensis (Clarke) P. K. Mukh. (Umbelliferae).

Bull. Bot. Surv. India 24 (1-4) : 43. 1982 (1983).

Peucedanum sikkimensis Clarke

Hook. f. Fl. Brit. Ind. 2 : 710. 1879.

Syntypes : Sikkim : Lingdam, 12. 10. 70, C. B. Clarke 13065 H; Rishee, 5.10.75, C. B. Clarke 244987B.

Anisadenia pubescens Griff. (Linaceae).

Itin. Notes 54. 1848; Fasc. Fl. Ind. 13 : 2. 1983.

Type : India : Meghalaya : Khasia, Herb. Griffith s. n.

A. sexatilis Wall. ex Meissn. (Linaceae).

Pl. Vasc. Gen. Comment. 96. 1838; Fasc. Fl. Ind. 13 : 3. 1983.

Type : Sheopore, Aug. 1821.

Anoectangium kashmiriense Aziz & Vohra (Pottiaceae).

Bull. Bot. Surv. India 25 (1-4) : 239-240. 1983 (1985).

Holotype : India : Kashmir : Masjid Valley, July 25, 1892, Duthie 14398.

Anoectochilus nicobaricus Balak. et Chakrab. (Orchidaceae).

Bull. Bot. Surv. India 20 : 80. 1978 (1979).

Holotype : India : Great Nicobar Island, 6 km on East-West Road, Shaded places in inland forests, Dec. 1, 1975, in flower, P. Chakraborty 3226A.

Anthoxanthum borii Jain et D. C. Pal (Poaceae).

J. Bombay Nat. Hist. Soc. 72(1) : 90. 1975.

Holotype : India : Tamil Nadu : Palnis, Pambar stream near Shen-thadikana Dec. 6, 1898, Bourne 1954.

A. hookeri (Griseb.) Rendle (Poaceae).

J. Linn. Soc. (Bot.) 36 : 380. 1904 ; J. Bombay Nat. Hist. Soc. 72(1) : 94. 1975.

Hierochloe hookeri (Griseb.) Clarke ex Hook. f. Fl. Brit. Ind. 7 : 223. 1896.

Type : India : Sikkim, 2700-3600 m, J. D. Hooker s. n.

***Anthoxanthum sikkimense* (Maxim.) Ohwi (Poaceae).**

Bull. Tokyo Sci. Mus. 18 : 8. 1947; J. Bombay Nat. Hist. Soc. 72(1) : 95. 1975.

Hierochloe gracillima Hook. f.

Fl. Brit. Ind. 7 : 223. 1896; Bor. 166. 1940.

Type : India : Sikkim, 3300-3600 m, *J. D. Hooker s. n.*

***Antistrophe caudata* King et Gamble (Myrsinaceae).**

J. Asiat. Soc. Beng. 74 (Part II) : 154. 1906; Bull. Bot. Surv. India 18(1-4) : 83. 1976 (1979).

Isosyntypes : Malaya : Perak : Gunong Batu Pateh, 1050 m, *Wray* 456; *Wray* 3242.

***A. oxyantha* (Wall. ex A. DC.) A. DC. (Myrsinaceae).**

Ann. Sci. Nat. 2. ser., 16 : 84. 1841 et in DC., Prodr. 8 : 92. 1844; Bull. Bot. Surv. India 18(1-4) : 81. 1976 (1979).

Isotype : Bangladesh : Silhet, *Wallich* 2275.

***A. oxyantha* (Wall. ex DC.) DC. var. *bhutanica* Nayar et Giri (Myrsinaceae).**

Bull. Bot. Surv. India 18(1-4) : 81. 1976 (1979).

Holotype : Bhutan, alt. 1100 m, *King s. n.*

***A. solanoides* (King et Gamble) Nayar et Giri (Myrsinaceae).**

Bull. Bot. Surv. India 18(1-4) : 83. 1976 (1979).

Isotype : Malaya : Perak, Larut, 150-250 m, *King's Collector* 4254.

***Apama barberi* Gamble (Aristolochiaceae).**

Kew Bull. 1924 : 386. 1924 et Fl. Pres. Madras 2: 840. 1957 (Repr. ed.); Bull. Bot. Surv. India 12(1-4) : 1. 1970 (1972).

Type : Tamil Nadu : Tirunelveli, Kannikatti, May 1901, *C. A. Barber* 3093.

***Apluda blatteri* Sur. (Poaceae).**

Bull. Bot. Surv. India 28 : 193. 1986 (1988).

Holotype : India : Rajasthan : Mount Abu, Oct. 13, 1916, *Father E. Blatter* 2594.

***Aponogeton satarensis* Sundararaghavan, Kulkarni & Yadav (Aponogetonaceae).**

Kew Bull. 36(4) : 687. 1982.

Holotype : India : Maharashtra : Satara Dist., Mavashi Plateau, 1200 m, *S. R. Yadav* Y-15.

***Aporusa acuminata* Thw. (Euphorbiaceae).**

Enum. Pl. Zeyl.: 288 (1861); Bull. Bot. Surv. India 26 : 42. 1984 (1985).

Type : Sri Lanka : Ambagamowa Dist., near Balangodde, 610-915 m, Thwaites C. P. 2530.

***A. dioica* (Roxb.) Muell.-Arg. (Euphorbiaceae).**

DC. Prodr. 15(2) 472 (1866); Bull. Bot. Surv. India 26 : 43. 1984 (1985).

Alnus dioica Roxb.

Fl. Ind. 3 : 580 (1832).

Icono-type : Roxb. I. C. t. 2378.

A. villosula Kurz

For. Fl. 2 : 362 (1877).

Syntype : S. Andaman, N. Corbyn's cove, Kurz s. n.

***A. fusiformis* Thw. (Euphorbiaceae).**

Enum. Pl. Zeyl. : 288 (1861); Bull. Bot. Surv. India 26 : 42. 1984 (1985).

Type : Sri Lanka : Central Province, 1220-1830 m, Thwaites C. P. 30.

***A. latifolia* Thw. (Euphorbiaceae).**

Enum. Pl. Zeyl. : 288 (1861); Bull. Bot. Surv. India 26 : 42. 1984 (1985).

Agyneia latifolia Moon.

Cat. Ceyl. Pl. : 65 (1824), *nom. invalid, sans descript.*

Type : Sri Lanka : Thwaites C. P. 3433.

***A. villosa* (Lindl.) Baill. (Euphorbiaceae).**

Et. Gen. Euphorb : 645 (1858); Bull. Bot. Surv. India 26 : 45. 1984 (1985).

A. glabrifolia Kurz

J. Bot. 4 : 330 (1875); Fl. Brit. Ind. 5 : 349. 1887.

Type : Nicobar Islands: Kamorta, tropical forest, Feb. 1875, Kurz s. n. (3 sheets).

***A. wallichii* Hook. f. (Euphorbiaceae).**

Fl. Brit. Ind. 5 : 350. 1887; Bull. Bot. Surv. India 26 : 44. 1984 (1985).

Isolectotype : Wall. Num. List No. 8019.

***Ardisia dawnensis* Nayar et Giri (Myrsinaceae).**

Bull. Bot. Surv. India 27 (1-4) : 222-224. 1985 (1987).

Holotype : Burma : Below Mulai-yit peak, Dawna Range, Amherst Dist., Jan. 27, 1912, 3500-6000 ft. (900-1825 m), J. H. Lace 5624.

Ardisia meghalayensis Nayar et Giri (Myrsinaceae).

J. Bombay Nat. Hist. Soc. 72(3) : 825. 1975.

Holotype : India : Meghalaya : Jaintea Hills. J/M No. 903, CNH Acc. No. 279310.

A. neglecta Nayar et Giri (Myrsinaceae).

J. Econ. Tax. Bot. 3(1) : 281. 1982.

Holotype : Burma : Tenasserim : Rocky River Bank, 126 miles down- from Tavoy, March 1885, Dr. King's Collector 115.

A. russellii Nayar et Giri (Myrsinaceae).

J. Econ. Tax. Bot. 3(1) : 281. 1982.

Holotype : Burma : Tavoy, Heinze head waters, 608 m, Nov. 25, '21, P. T. Russell 2211A.

Isotypes : Ibid., P. T. Russell, 2211 B-E.

Paratype : Tavoy, headstream, The Ta aingya chaung, 550 m, Feb. 5, 1919, A. T. Gage 48.

Arenaria bhutanica Majumdar & Babu (Caryophyllaceae).

J. Arn. Arb. 50 : 626-628. 1969.

Holotype and Isotype : Bhutan : Chila, on way to Paro, 3630-4125 m, Aug. 24, 1963, G. Sengupta 721.

A. curvifolia Majumdar (Caryophyllaceae).

Blumea 26(2) : 446. 1980.

Holotype : J. E. Duthie 3858A.

Argyreia arakuensis Bal. (Convolvulaceae).

Bull. Bot. Surv. India 3 : 163. 1961 (1962).

Holotype : India : Andhra Pradesh : Araku Valley, Visakhapatnam, Dist., 800 m, Sept. 14, 1961, Balakrishnan 540A.

Isotypes : Balakrishnan 540 B-F.

Arisaema caudatum Engl. (Araceae).

DC. Monogr. Phan. 2 : 559, 1879. emend. descr. Rolla Rao et Ahuja; Bull. Bot. Surv. India 11 : 450. 1969 (1971).

Neo-type : India : Maharashtra : Panchgani Plateau near Mahabaleswar, Ansari 105095A.

A. setosum A. S. Rao et D. M. Verma (Araceae).

Bull. Bot. Surv. India 11 : 196. 1969 (1971).

Holotype : India : Arunachal Pradesh : Sirang to Guizing, Siang Dist., Nov. 20, 1958, R. S. Rao 17907A, ♀

Aristolochia gourigangaica N. C. Nair (Aristolochiaceae).

Bull. Bot. Surv. India 10 : 332. 1968 (1969).

Holotype : India : Uttar Pradesh : Gouriganga Valley, 2800 m, May 4, 1965, at Radgadi (Near Muniari), Kumaon Himalayas, *N. C. Nair* 35654.

Artobotrys nicobarianus D. Das (Annonaceae).

Bull. Bot. Surv. India 11 : 194. 1969 (1971).

Holotype and Isotype : India : Great Nicobar, from the forest from Galathea Bay to Pulobaha Bay, March 26, 1966, *K. Thothathri & S. P. Banerjee* 11524.

Arthraxon deccanensis Jain (Poaceae).

J. Bombay Nat. Hist. Soc. 68(1) : 297-299. 1971.

Holotype : India : Maharashtra : Sinhagad, Poona Dist., Nov. 1956, *R. M. Patil* 7825.

Arthraxon junnarensis Jain et Hemadri (Poaceae).

J. Bombay Nat. Hist. Soc. 68(1) : 300-301. 1971.

Holotype : India : Maharashtra : Warsubai Plateau, 16 km, West of Junnar, Poona Dist., Oct. 6, 1965, *K. Hemadri* 106849 A/I.

Isotypes : *K. Hemadri* 106849 A/II and A/III (Mounted on same sheet).

Arthromeris jarrettii Sastry et Chowdh. (Polypodiaceae).

Bull. Bot. Surv. India 11 : 442. 1969 (1971).

Holotype : India : Arunachal Pradesh : Amjee-Begi, Subansiri Dist., Nov. 30, 1964, *A. R. K. Sastry* 40966A.

Arundinella cannanorica V. J. Nair, P. V. Sreekumar and N. C. Nair (Gramineae).

J. Bombay Nat. Hist. Soc. 80 (2) : 396. 1983.

Holotype : India : Cannanore Dist., Bela (in Between Kumbla and Bedukka), ±100 m, Nov. 23, 1981, *P. V. Sreekumar* 71822.

A. setosa Trin. var. **nilagiriana** Subba Rao et Kumari (Gramineae).

J. Bombay Nat. Hist. Soc. 72(3) : 827. 1975.

Holotype : India : Tamil Nadu : Near Koilbetta, Ebanad, Nilgiri Dist., 1850 m, Sept. 11, 1970, *Subba Rao et Kumari* 19795A.

Asarum himalaicum Hook. f. & Thoms. var. **bhutanicum** W. W. Smith (Aristolochiaceae).

Rec. Bot. Surv. India 4 : 271. 1913; Bull. Bot. Soc. Beng. 19(1) : 37. 1965.

Type : Bhutan, *White* 166.

Asemanthia maingayi (Hk. f.) Ridley var. **hirsuta** (King) A. K. Sinha et B. Mitra (Rubiaceae).

J. Bombay Nat. Hist. Soc. 78(2) : 427. 1981.

Syntypes : Perak, *King's Collector* 225; Malacca, *Derry* 253.

A. maingayi (Hk. f.). Ridley var. **montang** (Ridley) A. K. Sinha et B. Mitra (Rubiaceae).

J. Bombay Nat. Hist. Soc. 78(2) : 428. 1981.

Type : Malacca : Mount Ophir, Padang Batu, *Ridley* 3215.

Aspidopterys balakrishnanii Srivastava (Malpighiaceae).

J. Econ. Tax. Bot. 4(3) : 1003. 1983.

Holotype and Isotype : India : N. Bengal : (W. Duars), Lower Tendu forests, *Haines* 561.

Asraoa triandra J. Joseph (Arecaceae).

Bull. Bot. Surv. India 14 : 144. 1972 (1975).

Holotype : India : Arunachal Pradesh : Hayuliang, Lohit Dist., Sept. 25, 1969, *A. S. Rao* 48105A.

Atalantia racemosa Wight var. **bourdillonii** Narayanan et Nayar (Rutaceae).

Indian J. Forestry 13(1) : 69. 1990.

Holotype : India : Kerala : Calar toopurha, Quilon.

Dist., 60 m, Feb. 27, 1899, *V.F. Bourdillon* 160, Acc. No. 76452.

Bambusa cacharensis R. Majumdar (Gramineae).

Bull. Bot. Surv. India 25(1-4) : 237-238. 1983 (1985).

Holotype : India : Assam : Cachar, Lakhimpur, Mar Bastee, *Majumdar* 74265A.

Begonia scutta Wall. ex DC. (Begoniaceae).

Prod. 15(1) : 328. 1864; J. Econ. Tax. Bot. 13(1) : 247. 1989.

Type : Nepal : Gossiantham, 1821, *Wallich* 3986.

Bhidea fischeri P. V. Sreekumar & B. V. Shetty (Gramineae).

Kew Bull. 42(3) : 683. 1987.

Holotype : India : Kerala : Cannanore Dist., Mukkarikandam, ± 50 m, Oct. 18, 1981, *P. V. Sreekumar* 71754.

Biermannia jainiana Hedge et Nageswara Rao (Orchidaceae).

Bull. Bot. Surv. India 26(1-2) : 97. 1984 (1985).

Paratype : India : Arunachal Pradesh : West Kameng Dist., Sessa, 1100 m, May 11, 1982, *Hedge* 4194.

Blastania fimbriostipula (Fenzl.) Kotschy & Peyr. (Cucurbitaceae).

Pl. Tinn. 15 : t. 7. 1865; Fasc. Fl. Ind. 11 : 12. 1982.

Type : Sudan : Nubia, *Kotschy* 205.

Blumea venkataramanii Rolla Rao et Hemadri (Asteraceae).

Curr. Sci. 42(19) : 693-694. 1973.

Holotype : India : Maharashtra : collected around cultivated fields near Vadgaon on Poona Bombay Road, Poona Dist., Dec. 12, 1968, *Hemadri* 118174A.

Boehmeria tirapensis Deb et Dutta (Urticaceae).

Indian For. 90(11) : 782. 1964.

Holotype : India : Tirap Frontier Division, Kothong, June 25, 1961, *Deb* 26081A.

Boerreria eradii N. Ravi (Rubiaceae).

J. Bombay Nat. Hist. Soc. 66(3) : 539. 1969.

Paratypes : India : Kerala : Punalur, June 20, 1968, *N. Ravi* 2372 B-D.

Bolbitis nagalandense R. R. Rao & N. S. Jamir (Lomariopsidaceae).

Reinwardtia 10(2) : 119. 1985.

Holotype : India : Nagaland : Changki, Mokokchung Dist., 1000 m, *Jamir* 7112.

Bothriochola jainii Despande et Hemadri (Poaceae).

Indian For. 97(10) : 593. 1971.

Holotype : India : Maharashtra : on the open top of Durga Kahilla, ± 1200 m, about 30 km west of Junnar, Poona Dist., Oct. 29, 1964, *Hemadri* 104241A.

B. parameswaranii Sreekumar, Malathi and Nair (Poaceae).

J. Bombay Nat. Hist. Soc. 85(1) : 163. 1988.

Holotype : India : Kerala : Idukki, Eravikulum National Park, ± 2100 m, Feb. 14, 1981, *P. V. Sreekumar* 71858.

Brachiaria brizantha (Hochst. ex A. Rich.) Stapf var. *ciliata* Basappa et Muniyamma (Poaceae).

Proc. Indian Nation. Sci. Acad. B, 49(4) : 379. 1983.

Holotype : India : Karnataka : Dept. Garden, Manasagangotri, introduced (Fiji Origin, USDA, USA : 355712, Sept. 11, 1980, *Basappa et Muniyamma* 1004.

Brachiaria chennaveeraiana Basappa et Muniyamma (Poaceae).

Proc. Indian Nation. Sci. Acad. B, 49(4) : 378. 1983.

Holotype : Rajasthan : Near Gomukh, Mount Abu, Oct. 15, 1980, *Basappa & Muniyamma* 2195.

B. cruciformis (J. E. Smith) Grieb. var. **divaricata** Basappa et Muniyamma (Poaceae).

Proc. Indian Nation. Sci. Acad. B, 49(4) : 379. 1983.

Holotype : India : Karnataka : near Mysore, Bogadi, Aug. 12, 1979, *Basappa et Muniyamma* 1451.

B. hybrida Basappa et Muniyamma (Poaceae).

Proc. Indian Nation. Sci. Acad. B, 49(4) : 379. 1983.

Holotype : Karnataka : Dist. Shimoga, near tunga Dam, Gajanur, Sept. 10, 1979, *Basappa et Muniyamma* 2851.

B. munae Basappa (Poaceae).

Proc. Indian Acad. Sci. Pl. Sci. 93(1) : 53. 1984.

Holotype : India : Tamil Nadu : Madurai, Madurai Kamraj University Campus, Dec. 20, 1979, *Basappa* 3001.

B. ramosa (L.) Stapf var. **pubescens** Basappa et Muniyamma (Poaceae).

Proc. Indian Nation. Sci. Acad. B, 49(4) : 380. 1983.

Holotype : India : Karnataka : Shimoga Dist., Benakanaholli, Aug. 20, 1980, *Basappa et Muniyamma* 2171.

B. reptans (A.) Gard. et C. E. Hubb. var. **hispida** Basappa et Muniyamma (Poaceae).

Proc. Indian Nation. Sci. Acad. B, 49(4) : 380. 1983.

Holotype : India : Karnataka : Shimoga Dist., Govinakovi, Oct. 10, 1980, *Basappa et Muniyamma* 2302.

B. semiundulata (Hochst.) Stapf var. **intermedia** Basappa et Muniyamma (Poaceae).

Proc. Indian Nation. Sci. Acad. B, 49(4) : 380. 1983.

Holotype : India : Karnataka : near Doopada Bore, Bedaguli, Aug. 8, 1979, *Basappa et Muniyamma* 2602.

B. semiundulata (Hochst.) Stapf var. **lantana** Basappa et Muniyamma (Poaceae).

Proc. Indian Nation. Sci. Acad. B, 49(4) : 380. 1983.

Holotype : India : Karnataka : near Doopada Bore, Bedaguli, Mysore, Aug. 8, 1979, *Basappa et Muniyamma* 2603.

Brachiaria setigera (Retz.) C. E. Hubb. var. **albistyla** Basappa et Muniyamma (Poaceae).

Proc. Indian Nation. Sci. Acad. B, 49(4) : 380. 1983.

Holotype : India. Tamil Nadu : Coimbatore, Agricultural College Campus, Oct. 25, 1980, *Basappa et Muniyamma* 2716.

B. stapfiana Basappa et Muniyamma (Poaceae).

Indian Nation. Sci. Acad. B, 49(4) : 379. 1983.

Holotype : India : Karnataka : near Ugra Narasimha Temple, Hampi, Bellaey, *Basappa et Muniyamma* 2100.

B. villosa (Lamk.) A. Camus var. **glaberrima** Basappa et Muniyamma (Poaceae).

Proc. Indian Nation. Sci. Acad. B, 49(4) : 381. 1983.

Holotype : Near Monkey point, Kasauli, Siwalik Hill, Chandigarh, Nov. 26, 1981, *Basappa et Muniyamma* 2911.

Brachystelma elenaduensis M. B. S. Char (Asclepiadaceae).

Curr. Sci. 47 : 965. 1978.

Holotype : India : Karnataka : Tumkur Dist., Elenadu (altitude : of the village bench mark, 780, 33 m of the contour, 800 m), sandy grassy land, April 2, 1969, *M. B. S. Char* 1.

Brachythecium chakratense Vohra (Brachytheciaceae).

Bull. Bot. Surv. India 22 : 118. 1980 (1982).

Holotype : India : Garhwal, Churani to Chakrata, 2135 m, Mar. 25, 1963, *Vohra* 1804.

Paratypes : India : Himachal Pradesh : Nachar Bari Forest, 2135-2440 m, Oct. 11, 1960, *Vohra* 468, 486; Rainsukh, Jaunsar, Oct. 1894 (Present under the name *Brachythecium emodi-glareosum* Broth.).

B. garhwalense Vohra (Brachytheciaceae).

Bull. Bot. Surv. India 22 : 122. 1980 (1982).

Holotype : India : Jaunsar Bahar, Deoban, 2135-2440 m, Sept. 28, 1879, *Duthie* (Present under the name *Brachythecium brachycladulum* C. Muell.

Brachytome russellii Deb & Gang. (Rubiaceac).

Candollea 42 : 353. 1987.

Holotype : Burma : Tavoy : Chaung, 1600 ft. (480 m), 19.3.1919, *P. T. Russell* 21.

Paratype : Valley of Talaingza, Chung, 1200 ft. (360 m), 8.4.1919, *P. T. Russell* 186.

Brachytome scortechinii King & Gamble (Rubiaceae).

J. Asiat. Soc. Beng. 72(4) : 198. 1903; Candollea 42 : 353. 1987.

Isolectotype : Perak (Two sheets).

Paratypes : Taiping, March 1884, *Scortechini* 14; Larut, 900-1015 m, July 1881, *King's Collector* 2136; April 1882, *King's Collector* 2930; Lower camp Gumong, Batu, Putch, 1020 m, L. Wray 950; L. Wray 2812; Perak, Maxwells Hill, L. Wray 2999.

Brassaiopsis andamanica R. N. Ban. (Araliaceae).

Indian For. 94(10) : 775-777. 1968.

Holotype : India : South Andaman black creek, *Kurz s. n.* Acc. No. 193177.

Isotype : Ibid., *S. Kurz* 193178.

Brucea mollis Wall. ex Kurz (Simaroubaceae).

J. Asiat. Soc. Beng. 42 : 64. 1873; Fasc. Fl. Ind. 4 : 9. 1980.

Type : Burma : Martaban, *Kurz* 2009.

Bulbophyllum brachypodium A. S. Rao et Balak. var. *parviflorum* A. S. Rao et Balak. (Orchidaceae).

Bull. Bot. Surv. India 10 : 350. 1968.

Holotype : Eastern Bhutan : Gumdrithang, 2000 m, March 16, 1965, flowered at National Orchidarium, Shillong, Aug. 23, 1965. *Balakrishnan* 43004A.

B. leopardinum Lindl. var. *tuberculatum* Balak. & Chowdhury (Orchidaceae).

Bull. Bot. Surv. India 9 : 90. 1967 (1968).

Holotype : Eastern Bhutan: Nyoth on tract between Shali and Tshiyangtsi ($90^{\circ}34'E$ & $27^{\circ}32'N$) 1800 m, March 13, 1965, collected in vegetative condition, flowered at Shillong on May 3, 1965, *Balakrishnan* 41971A.

B. odoratissimum Lindl. var. *racemosum* Balakrishnan (Orchidaceae).

J. Bombay Nat. Hist. Soc. 75(1) : 157. 1978.

Holotype : E. Bhutan : 1800 m, Oct. 24, 1965 (in Veg. condition), flowered at National Orchidarium, Shillong, Meghalaya State, in Pot No. 3614, *Balakrishnan* 43070A.

B. reptans (Lindl.) Lindl. var. *acuta* Malhotra et Balodi (Orchidaceae).

Bull. Bot. Surv. India 26 : 110. 1984 (1985).

Holotype : India : Uttar Pradesh : Gorapatta (Gori Valley), Pithoragarh, 1500 m, March 10, 1965, M. A. Rao 35340.

Bupleurum andhricum M. P. Nayar et R. N. Ban. (Umbelliferae).

Bull. Bot. Surv. India 12 : 255. 1970 (1972).

Holotype : India : Andhra Pradesh : Palakond, Godavari Dist., 762 m, Oct. 1, 1920, *V. Narayanaswami* 370A.

Burmannia candelabrum Gagnep. (Burmanniaceae).

Bull. Soc. Bot. Fr. 54 : 462. 1907; Fasc. Fl. Ind. 19 : 7. 1988.

B. candida Griff. ex Hook. f.

Fl. Brit. Ind. 5 : 665. 1888. *p.p.*

Type : East Bengal: *Griffith* 5598.

B. championii Thw. (Burmanniaceae).

Enum. Pl. Zeyl. 325. 1864; Fasc. Fl. Ind. 19 : 9. 1988.

Isotype : Ceylon (Sri Lanka) : *Thwaites* 2735.

B. coelestis D. Don (Burmanniaceae).

Prodr. Fl. Nepal 44. 1825; Fasc. Fl. Ind. 19 : 9. 1988.

Type : Nepal: *Wallich* 9005.

B. nepalensis (Miers) Hook. f. (Burmanniaceae).

Fl. Brit. Ind. 5 : 666. 1888; Fasc. Fl. Ind. 19 : 12. 1988.

Cyanotis nepalensis Miers et Wall.

Wall. Cat. No. 9006. 1840, *nom. nud.*

Type : Nepal: *Wallich* 9006.

B. pusilla (Wall. ex Miers) Thw. (Burmanniaceae).

Enum. Pl. Zeyl. 325. 1864 Excl. var. B; Fasc. Fl. Ind. 19 : 13. 1988.

Type : India : Concan, *Wallich* 9008.

B. stricta Jonkar (Burmanniaceae).

Meded. Bot. Mus. Utrecht 51 : 156. 1938; Fasc. Fl. Ind. 19 : 14. 1988.

Type : India : Cochin, *Meebold* 12337.

Calamus pseudoseanus S. K. Basu (Palmae).

J. Econ. Tax. Bot. 13(1) : 133. 1989.

Holotype : India : Tamil Nadu: S. Coimbatore, 17.10.1905, DFO Reg 23503, Acc. No. 494007.

Isotypes : Ibid., Acc. Nos., 493999, 494001, 94006.

Callophyllum kunstleri King (Guttiferae).

J. Asiat. Soc. Beng. 59 : 174. 1891; Bull. Bot. Surv. India 2 : (1 & 2) : 141. 1960.

Callophyllum apetalum Blanco.

Fl. Filip. (ed. 2) 429. 1845; et (ed. 3) 415. 1879 (non Willd. 1811).

C. pulcherrimum F. Vill.

Novis. app. 17. 1880 (non Wall. 1828).

Syntypes : Perak : Larut, *King's Collector* 5328, 5374, 5459.

Callophyllum polyanthum Wall. (Guttiferae).

Wall. Cat. 4844, 1828, nomen; Bull. Bot. Surv. India 2(1 & 2) : 144. 1960.

Isotype : Bangladesh : Sylhet, *Wallich* 4844.

C. trapezifolium Thwaites (Guttiferae).

Enum. Pl. Zeyl. 51. 1858 ; Bull. Bot. Surv. India 2(1 & 2) : 144. 1960.

Isotype : Ceylon : Hunasgiria Dist., Central Province, 4-5000 ft., *Thwaites* 2446.

Camellia siangensis T. K. Pal et Nayar (Theaceae).

Bull. Bot. Surv. India 27 (1-4) : 92. 1985 (1987).

Holotype : India : Arunachal Pradesh : Siang F.D., Minguing-Takepokong, 1150-1775 m, Nov. 18, 1958, *R. S. Rao* 17781-A.

Isotypes : Ibid., *R. S. Rao* 17781 B, C, D.

Campanula wallichii C. R. Babu (Campanulaceae).

J. Bombay Nat. Hist. Soc. 65(3) : 808. 1968.

Isotype : *Wallich* 1289.

C. wattiana Nayar et Babu (Campanulaceae).

J. Indian Bot. Soc. 49(1-4) : 183. 1970 (1971).

Holotype and Isotype : India : Jammu & Kashmir : Bagi, 2970 m, Sept. 23, 1888, *Watt* 8609.

Cantanthera lysipetala F. v. Muell. (Melastomataceae).

Journ. Bot. 24 : 289. 1886; Reinwardtia 10 : 52. 1982.

Hederella lysipetala (Muell.) Nayar

Kew. Bull. 20 : 237. 1966.

Isolectotype : New Guinea : Papua, Sogeri region, *Forbes* 419.

Medirilla anomala Cogn.

DC. Monogr. Phan. 7 : 1185. 1891.

Isotype : Ibid., *Forbes* 451.

Canthium andamanicum Gang. et T. Chakrab. (Rubiaceae).

J. Econ. Tax. Bot. 13(1) : 85. 1989.

Holotype : India : S. Andamans, April 1890, *King* s. n.

Capparis acutifolia Sweet ssp. **viminea** Jacobs (Capparaceae).

Blumea 12(3) : 429. 1965.

C. viminea Hook. f. & Th.

Hook. f. Fl. Brit. Ind. 1 : 179 (1872).

C. membranifolia Kurz

J. Asiat. Soc. Beng. 42 ii (1874) 70.

Type : *C. membranifolia* : Burma : Pegu, Kurz 1826.

C. assamica Hook. f. & Th. (Capparaceae).

Hook. f. Fl. Brit. Ind. 1 : 177. 1872; Blumea 12(3) : 434. 1965.

C. gallatlyi King

Ann. Gard. Calc. 5(1896) 118. t. 133B.

Type : *C. gallatlyi* : Burma : Tenasserim, Moolar Plain, 1877, *Gallatly* 880.

C. burmanica Coll. & Hemsl. (Capparaceae).

J. Linn. Soc. Lond. Bot. 28(1890) 19. t. 2; Blumea 12(3) : 439. 1965.

Type : Upper Burma : Shewenoungbu, fl., Collet 495.

C. cucurbitina King (Capparaceae).

J. Asiat. Soc. Beng. 58. ii (1889) 395; Blumea 12(3) : 446. 1965.

Type : Malesia : King's Collector 10027.

C. flavicans Kurz (Capparaceae).

J. Asiat. Soc. Beng. 39. ii (1870) 62; Blumea 12(3) : 451. 1963.

Type : Southwestern Siam : Teijsmann HB 5931.

C. floribunda Wight (Capparaceae).

IC. Pl. Ind. Or. 2(1840) 35. t. 14; Blumea 12(3) : 453. 1965.

C. andamanica King

Ann. Gard. Calc. 5(1896) 119. t. 137.

Type : *C. andamanica* : S. Andaman : January 1, 1892, King's Collector s. n.

C. fusifera Dunn (Capparaceae).

Kew Bull. (1914) 377; Blumea 12(3) : 454. 1965.

Type : S. India : Udumanparai, Anamalais, April 30, 1903, Barbar 5726.

C. grandis Linn. f. (Capparaceae).

Suppl. (1781) 263; Blumea 12(3) : 456. 1965.

C. grandis var. *auricans* Kurz

J. Asiat. Soc. Beng. 43 ii (1874) 70.

Capparis auricans (Kurz) Craib.

Kew Bull. (1922) 169.

Type : Burma : Kurz 1827.

Capparis nilgiriensis Subba Rao, Kumari and Chandrasekaran (Capparaceae).

J. Bombay Nat. Hist. Soc. 78(1) : 146. 1981.

Holotype : India : Tamil Nadu : Chinnacoonoor, 950 m, March 16, 1972, Subba Rao 40259A.

C. pachyphylla Jacobs (Capparaceae).

Blumea 12(3) : 476. 1965.

Type : Kingdom-ward 11183.

C. pubiflora DC. (Capparaceae).

Prod. 1(1824) 246; Blumea 12(3) : 479. 1965.

C. pubiflora var. *perakensis* Scort. ex King

J. Asiat. Soc. Beng. 58 ii (1889) 394.

C. perakensis (Scort. ex King) Ridl.

Fl. Mal. Pen. 1(1922). 124.

Type : Malesia : Scortechini 1784.

C. scortechinii King (Capparaceae).

J. Asiat. Soc. Beng. 58 ii (1889) 394; Blumea 12(3) : 488. 1965.

Type : Malaya : Perak, Batang, Padang Dist., Fl., 1885, King's Collector 8083.

C. sepiaria Linn. (Capparaceae).

Syst. Nat. 10th ed. 2(1759) 1071; Blumea 12(3) : 489. 1965.

C. glauca var. *angustifolia* Coll. & Hemsl.

J. Linn. Soc. Lond. Bot. 28(1890) 19.

Type : *C. glauca* var. *angustifolia* : Burma, Collett 538.

C. siamensis Kurz (Capparaceae).

For. Fl. Burma 1(1877) 63; Blumea 12(3) : 493. 1965.

Type : Teijsmann HB 5927.

C. sikkimensis Kurz (Capparaceae).

J. Asiat. Soc. Beng. 43 ii (1875) 181; Blumea 12(3) : 496. 1965.

Type : India : Sikkim, Kurz s. n.

C. tenera Dalz. (Capparaceae).

Hook. J. Bot. Kew Gard. Misc. 2 (1850) 41; Blumea 12(3) : 497. 1965.

Capparis disticha Kurz

J. Asiat. Soc. Beng. 43 ii (1874) 70. name.

Type : *C. disticha* : S. Burma, Kurz 20.*Capparis trinervia* Hook. f. & Th. (Capparaceae).

Fl. Brit. Ind. 1(1872) 175; Blumea 12(3) : 500. 1965.

C. kunstleri King

J. Asiat. Soc. Beng. 58 ii (1889) 396.

Type : Malesia : King's Collector 8337.

C. versicolor Griff. (Capparaceae).

Not. Pl. As. 4 (1854) 577; Blumea 12(3) : 501. 1965.

C. larutensis King

J. Asiat. Soc. Beng. 58 ii (1889) 393.

Type : *C. larutensis* : Malaya : King's Collector 5103.**C. zeylanica** Linn. (Capparaceae).

Sp. Pl. 2nd. ed. (1762) 720; Blumea 12(3) : 505. 1965.

C. crassifolia Kurz

J. Asiat. Soc. Beng. 42. ii (1874) 227.

Type : *C. crassifolia* : Lower Burma : Pegu, Kurz 1825.*C. polymorpha* Kurz

J. Asiat. Soc. Beng. 42 ii (1874) 227.

Type : *C. polymorpha* : Lower Burma : Pegu, Kurz 1828.**Caralluma nilagiriana** Kumari et Subba Rao (Asclepiadaceae).

J. Bombay Nat. Hist. Soc. 73(1) : 194. 1976.

Holotype : India : Tamil Nadu : From Garden of Botanical Survey of India, Southern Circle, Coimbatore (from cultivated material raised from wild plants collected in vegetative condition, growing in rocky areas on way from Anaikatty to Ebanad) 700 m, in Nilgiri Dist., Tamil Nadu, Aug. 20, 1974, Subbarao et Kumari 39262A.

Cardanthera uliginosa Buch.-Ham. var. **birbhumensis** Guha (Acanthaceae).

Bull. Bot. Soc. Beng. 21(1) : 47-48. 1967.

Holotype : India : Bengal : Nanur, Birbhum, Jan. 6, 1965, B. P. Guha 131.

Carex nandadeviensis Ghildyal, Bhattacharyya et Hajra (Cyperaceae).

Ind. Journ. Forestry 9(1) : 90-91. 1986.

Holotype : India : Uttar Pradesh : Chamoli, Bhojgara-Ramni, Nanda Devi National Park, July 5, 1982, P. K. Hajra 73974A.

Casearia insularis Vasad. & T. Chakrab. (Flacourtiaceae).

J. Econ. Tax. Bot. 5(4) : 991. 1984.

Holotype : India : Andaman-Nicobar Islands, North Andaman Island, Saddle peak range, Lamia Bay slope, March 31, 1977, *Balakrishnan* 5389A.

C. rubescens Dalz. var. **gamblei** N. Mukherjee (Flacourtiaceae).

J. Bombay Nat. Hist. Soc. 69(2) : 393. 1972.

Holotype : South India, evergreen forest of Pirmed 3500 ft., *T. F. Bourdillon* 181.

C. sikkimensis N. Mukherjee (Flacourtiaceae).

J. Bombay Nat. Hist. Soc. 69(2) : 392. 1972.

Holotype : India : W. Bengal : Kurseong, Dhobijhwa, 6000 ft., April 1882, *Gamble* 10316.

Cassia floribunda Cav. var. **pubescens** Singh (Leguminosae).

J. Econ. Tax. Bot. 10(2) : 326. 1987.

C. laevigata Gamble

Fl. Madras Part 3 : 399. 492. 1919 *pro parte* non Willd. 1809.

Type : Tamil Nadu : Nilgiri, 1800 m, June 1885, *Gamble* 16287.

Catanthera lysipetala F. v. Muell. (Melastomataceae).

Journ. Bot. 24 : 289. 1886; Reinwardtia 10 : 52. 1982.

Hederella lysipetala (Muell.) Nayar

Kew Bull. 20 : 237. 1966.

Isolectotype : New Guinea : Papua, Sogeriregion, *Forbes* 419.

Medirilla anomala Cogn.

DC. Mongr. Phan. 7 : 1185. 1891.

Isotype : Ibid., *Forbes* 451.

Centrantherum sengaltherianum B. M. Narayana (Compositae).

Curr. Sci. 50(6) : 279. 1981.

Holotype : India : Tamil Nadu : Tirunelveli Dist., 1100 m, in rock crevices, rare, March 18, 1978, *B. M. Narayana* 175A.

Ceropégia ciliata Wight (Asclepiadaceae).

Ic. Pl. Ind. Or. 4. t. 1262. 1848; Fasc. Fl. Ind. 16 : 13. 1984.

Type : India : Tamil Nadu : Nilgiri Hills, *Wight* 1927.

C. hookeri Clarke ex Hook. f. (Asclepiadaceae).

Fl. Brit. Ind. 4 : 73. 1883; Fasc. Fl. Ind. 16 : 17. 1984.

Type : Sikkim : 8-10,000 ft., *J. D. Hooker* 23.

***Ceropegia huberi* Ansari (Asclepiadaceae).**

Bull. Bot. Surv. India 10(2) : 219. 1968 (1969); Fasc. Fl. Ind. 16 : 17. 1984.

Holotype : India : Maharashtra : Ambaghat, Ratnagiri Dist., Aug. 29, 1967, *Ansari* 105001A.

Isotype : Ibid., *Ansari* 105001E.

Paratype : Ibid., 105033A.

***C. jainii* Ansari et Kulkarni (Asclepiadaceae).**

Bull. Bot. Surv. India 22 : 221. 1980 (1982).

Holotype : India : Maharashtra : Ratnagiri Dist., Ambolighat, Aug. 13, 1977, *Kulkarni* 121885A.

Isotype : Ibid., *Kulkarni* 121885D.

***C. kachinensis* Prain (Asclepiadaceae).**

J. Asiat. Soc. Beng. 69(2) : 170. 1900; Fasc. Fl. Ind. 16 : 19. 1984.

Holotype : Burma : Kachin Hills, Shaik Mokim 's. n.

Isotype : Burma, Shaik Mokim s. n.

***C. maccannii* Ansari (Asclepiadaceae).**

Bull. Bot. Surv. India 22(1-4) : 227. 1980; Fasc. Fl. Ind. 16 : 22. 1984.

Holotype : India : Maharashtra : Sinhagadh Hills, Pune, Dist., Aug. 7, 1964, *M. Y. Ansari* 97574A.

Isotype : Ibid., *M. Y. Ansari* 97574D.

***C. mahabalei* Hemadri et Ansari (Asclepiadaceae).**

Indian For. 97(2) : 105. 1971; Fasc. Fl. Ind. 16 : 24. 1984.

Holotype : India : Maharashtra : Ralegaon Hill, ± 1000 m, about 10 km west of Junnar, Poona Dist., Sept. 25, 1968, *Hemadri* 118070A.

Isotype : Ibid., *Hemadri* 118070G.

***C. metziana* Miq. (Asclepiadaceae).**

Analecta Bot. Ind. 3 : 11. 1852; Fasc. Fl. Ind. 16 : 25. 1984.

Type : India : Tamil Nadu : Devala (Nilgiri Dist.), Gamble 15601.

***C. noorjahaniae* Ansari (Asclepiadaceae).**

J. Bombay Nat. Hist. Soc. 69(1) : 250. 1972; Fasc. Fl. Ind. 16 : 26. 1984.

Holotype : India : Maharashtra : Wai-Panchgani Ghat, Satara Dist., *Ansari* 104880A.

Isotype : Ibid., *Ansari* 104880E.

***C. panchganiensis* Blatter et McCann (Asclepiadaceae).**

J. Bombay Nat. Hist. Soc. 36 : 534. 1933; Bull. Bot. Surv. India 22 (1-4) : 199. 1980; Fasc. Fl. Ind. 16 : 27. 1984.

Neotype : India : Maharashtra : Lingmala near Mahabaleshwar, Satara Dist., *Ansari* 105090A.

Ceropegia pussilla Wight et Arn. (Asclepiadaceae).

Wight, Contrib. 31. 1834; Fasc. Fl. Ind. 16 : 28. 1984.

Isotype : India : Tamil Nadu : Nilgiri Hills, Herb. *Wight* 1932.

C. raizadiana Babu (Asclepiadaceae).

Herb. Fl. Dehradun 305-306. 1977.

Holotype : India : Uttar Pradesh : Dehra Dun, Bindal, not uncommon, on ravine slopes, Fls. July-Oct., *C. R. Babu* 32215.

C. rollae Himadri (Asclepiadaceae).

Bull. Bot. Surv. India 10(2) : 123. 1968; Fasc. Fl. Ind. 16 : 29. 1984.

Holotype : India : Maharashtra : Dhak Khilla, about 27 km west of Junnar, Dist. Poona, ±1300 m, Sept. 29, 1965, *Himadri* 107472A.

Isotype : Ibid., *Himadri* 107472F.

Paratype : On the open top of Durga Khilla, ±1200 m, about 30 km west of Junnar, Oct. 1, 1965, *Himadri* 107547C.

C. sahyadrica Ansari et Kulkarni (Asclepiadaceae).

Indian For. 97(12) : 688. 1971; Fasc. Fl. Ind. 16 : 29. 1984.

Holotype : India : Maharashtra : Ambolighat, Ratnagiri Dist., Aug. 30, 1967, *Kulkarni* 108643A.

Isotype : Ibid., *Kulkarni* 108643D.

C. santapaui Wadhwa et Ansari (Asclepiadaceae).

Bull. Bot. Surv. India 10(1) : 95. 1968; Fasc. Fl. Ind. 16 : 30. 1984.

Holotype : India : Maharashtra : Mahad Ghats, near about milestone No. 30/7 from Mahabaleshwar, Satara Dist., ± 1160 m, Aug. 19, 1966, *Wadhwa* 109640A.

Isotype : Ibid., *Wadhwa* 109640H.

Paratype : Ibid., Sept. 13, 1966, *Wadhwa et Ansari* 109651A.

C. sootepensis Craib (Asclepiadaceae).

Kew Bull. 1911. 420.

Isotype : Siam: Chiengmai, in open deciduous Jungle on Doi Sootep, 450 m, *Kerr* 695.

Chlorophytum bharuchae Ansari, Raghavan et Hemadri (Liliaceae).

Indian For. 96(4) : 304. 1970.

Holotype : India : Maharashtra : from Shivneri Hill, Junnar, Poona, July 25, 1963, *Ansari* 88712A.

Choerospondias articulata Doli Chandra (Anacardiaceae).

J. Bombay Nat. Hist. Soc. 75(2) : 457. 1978.

Holotype : India : Uttar Pradesh : Mussoorie, May 15, 1899, P. W. Mackinnon s. n., Acc. No. 98623.

Chonemorpha pedicellata Seshagiri Rao (Apocynaceae).

J. Indian Bot. Soc. 32 : 44. 1953.

Type : N. W. India, Hb. Royle.

Chrysophyllum asraoii Rakesh et Thoth. (Sapotaceae).

Indian J. Forestry 13(1): 61. 1990.

Type : India: Maharashtra: Between shoutally and Kundally, 7 miles north West of Somwarpet, Oct. 16, 1963. A. S. Rao 94720.

Chrysopogon pseudozeylanicus K. G. Bhat. & C. R. Nagendran (Poaceae).

Reinwardtia 10(Part 2) 128. 1985.

Holotype : India : Karnataka : South Kanara Dist., Charmadi Ghat, about 15 km from Kotigehar on way to Dharmasthala, Nov. 24, 1977, K. G. Bhat. 550A.

Cleidion bishnui T. Chakrab. et Gang. (Euphorbiaceae).

J. Econ. Tax. Bot. 12(2) : 473. 1988.

Holotype : Burma : Katha Dist., Kadu Hills, July 11, 1911, J. H. Lace 5343.

C. javanicum Bl. var. **longipedicellatum** T. Chakrab. et Gang. (Euphorbiaceae).

J. Econ. Tax. Bot. 12(2) : 491. 1988.

Holotype : India : Sikkim : Munsong, April 26, 1909, Kari 1161, Acc. No. 411202.

Isotype : Ibid., Kari 1161, Acc. No. 411208.

Cleistanthus sankunianus Sivarajan & Bachachandran (Euphorbiaceae).

Kew Bull. 40(1) : 121. 1985.

Holotype : India : Kerala : Kottakkal Arya Vaidya Sala Herbal Garden, original source not known, Indu 1486.

Clematis smilacifolia Wall. ssp. **andamanica** Kapoor (Ranunculaceae).

Kew Bull. 19(2) : 331-333. 1965.

Holotype : India : Andaman Islands: Naumnagar, Sept. 27, 1890, King's Collector, Sheet No. 147.

Paratypes : Andaman: King's Collector s. n. Sheets Nos. 145, 146, 148, 149.

Clematis zemuensis W. W. Smith (Ranunculaceae).

Rec. Bot. Surv. India 4(5) : 166. 1911.

Holotype : Sikkim : Lower Zemu, 9000 ft., *Smith & Cave* 2671, Acc. No. 333.

Isotypes : Zemnu Valley, 9000 ft., *Smith & Cave* 2671, Acc. Nos. 331, 332, 339.

Cleome viscosa L. var. **nagarjunakondensis** Sundararaghavan (Capparaceae).

Bull. Bot. Surv. India 28 : 187. 1986 (1988).

Holotype : India : Andhra Pradesh : Nagarjunakonda Hills, Nalgonda Dist., 200 m, July 13, 1961, *K. Thothathri* 6616A.

Isotypes : Ibid., *K. Thothathri* 6616B-D.

Cnesmone javanica Bl. var. **glabriuscula** Balakr. & N. G. Nair (Euphorbiaceae).

Gdns. Bull. Singapore 31 : 49. 1978.

Holotype : India : South Andaman Island, Guptapara, Nov. 23, 1973, *Balakrishnan* 641A.

Paratype : Ibid., Oct. 22, 1975, *Balakrishnan & Nair* 2287.

Cocculus prainianus (Diels) Pramanik et Thoth. (Menispermaceae).

Kew Bull. 42(3) : 706. 1987.

Pericampylus prainianus Diels

Engl. Pflanzenr. Menisp. iv. 94 : 221. 1910.

Holotype : India : Naga Hills, Belegan, May 1899, *Prain's Collector (M. A. Hock)* 968.

Coelachne perpusilla (Arn. ex Steud.) Thw. var. **nilagirica** Ved Prakash et Jain (Poaceae).

Bull. Bot. Surv. India 24 : 187. 1982 (1983).

Holotype : India : Tamil Nadu : Nilgiri Dist., Parthimund, 2300 m, July 11, 1970, *J. L. Ellis* 34627.

Coelogyné corymbosa Lindl. (Orchidaceae).

Fol. Orch. Coelog. 7. 1854; Fasc. Fl. Ind. 5 : 9. 1980.

Syntype : India : Khasi Hills, *Hooker s. n.*

C. cristata Lindl. (Orchidaceae).

Coll. Bot. Sub. t. 33 (Textus). 1821; Fasc. Fl. Ind. 5 : 9. 1980.

Type : Nepal : Toka, March, 1821, *Wallich s. n.*, Wall. Num. List No. 1958/1.

***Coelogyné flaccida* Lindl. (Orchidaceae).**

Gen. & Sp. Orch. 39. 1830; Fasc. Fl. Ind. 5 : 10. 1980.

Type : Nepal : Toka, Apr. 1821, *Wallich s. n.*, Wall. Num. List No. 1961.

***C. fuscescens* Lindl. var. *fuscescens* (Orchidaceae).**

Gen. & Sp. Orch. 41. 1830; Fasc. Fl. Ind. 5 : 13. 1980.

Type : Nepal : Toka, March 1821, *Wallich s. n.*, Wall. Num. List No. 1962A.

***C. glandulosa* Lindl. var. *bournei* Das et Jain (Orchidaceae).**

Bull. Bot. Surv. India 18 : 241. 1976 (1979); Fasc. Fl. Ind. 5 : 14. 1980.

Holotype : India : Pulney Hills, Poombarai, May 25, 1899, *Bourne* 2941.

***C. glandulosa* Lindl. var. *sathyanarayanae* Das et Jain (Orchidaceae).**

Bull. Bot. Surv. India 18 : 241. 1976 (1979); Fasc. Fl. Ind. 5 : 14. 1980.

Holotype : India : Tamil Nadu : Kodaikanal, March 10, 1960, *Saldanha* 5211.

***C. griffithii* Hook. f. (Orchidaceae).**

Fl. Brit. Ind. 5 : 838. 1890 & I. c. Pl. t. 2108. 1894; Fasc. Fl. Ind. 5 : 14. 1980.

Syntype : India : Manipur: Khonghi Valley, 4-5000 ft. (1200-1500 m), April 27, 1882, *Watt* 6780.

***C. hitendrae* Das et Jain (Orchidaceae).**

Orch. Rev. 86. 195. 1978.

Holotype : India : Nagaland : Pulebadje, *Kataki* 60202.

***C. longipes* Lindl. (Orchidaceae).**

Fol. Orch. Coelog. 10. 1854; Fasc. Fl. Ind. 5 : 16. 1980.

Syntype : India : Khasi Hills, 4-5000 ft. (1200-1500 m); *J. D. Hooker* & *T. Thomson* s. n.

***C. nitida* (Wall. ex Don) Lindl. (Orchidaceae).**

Gen. & Sp. Orch. 40. 1830 [Excl. Syn. *Cymbidium nitidum* Roxb.]; Fasc. Fl. Ind. 5 : 19. 1980.

Type : Nepal: *Wallich s. n.*, Wall. Num. List No. 1954. p.p.

***C. occultata* Hook. f. var. *uniflora* Balakrishnan (Orchidaceae).**

J. Bombay Nat. Hist. Soc. 75(1) : 159. 1978.

Holotype : E. Bhutan : Nyoth Forest, 2000 m, Oct. 22, 1965, on moss covered tree trunk and rock boulders, *Balakrishnan* 43041A.

Coelogyne odoratissima Lindl. (Orchidaceae).

Gen. & Sp. Orch. 41. 1830; Fasc. Fl. Ind. 5 : 20. 1980.

Type : Herb. *Heyne s. n.*, Wall. Num. List No. 1960.

C. prolifera Lindl. (Orchidaceae).

Gen. & Sp. Orch. 40. 1830; Fasc. Fl. Ind. 5 : 23. 1980.

Type : Nepal : 1821, *Wallich s. n.*, Wall. Num List No. 1956.

C. punctulata Lindl. forma **brevifolia** (Lindl.) Das et Jain (Orchidaceae).

Fasc. Fl. Ind. 5 : 25. 1980.

Syntype : India : Khasi Hills, *J. D. Hooker & T. Thomson s. n.* CAL. 454607!.

C. raizadae Jain et Das (Orchidaceae).

Proc. Ind. Acad. Sci. 87B (5) : 119. 1978; Fasc. Fl. Ind. 5 : 25. 1980.

Holotype : India : Khasi Hills, Mawsmai, *Das* 55419.

C. schultesii Jain et Das (Orchidaceae).

Proc. Ind. Acad. Sci. 87B (5) : 119. 1928; Fasc. Fl. Ind. 5 : 28. 1980.

Holotype : India : Meghalaya: Khasi Hills, Cherrapunjee, *Das* 60256.

C. uniflora Lindl. (Orchidaceae).

Gen. Sp. Orch. 42. 1830; Fasc. Fl. Ind. 5 : 30. 1980.

Type : Nepal: *Wallich s. n.*, Wall. Num. List No. 1956.

Corallorrhiza anandae Malhotra et Balodi (Orchidaceae).

Bull. Bot. Surv. India 26(1-2) : 108. 1984 (1985).

Holotype : India : Uttar Pradesh : Gori Valley, growing on slopes in the shade of *Betula* trees, Martoli Bugyals, 4500 m, June 15, 1958. *T. A. Rao* 6851A.

Coriaria nepalensis Wall. (Coriariaceae).

Pl. Asiat. Rar. 3 : 67. t. 289. 1832; Fasc. Fl. Ind. 1 : 3. 1978.

Type : India : Uttar Pradesh : Kumaon, *Blinkworth-Wall.* CAT. 6817C.

C. terminalis Hemsley var. **xanthocarpa** Rehder & Wilson (Coriariaceae).

Sarg. Pl. Wilson 2 : 171. 1916; Fasc. Fl. Ind. 1 : 4. 1978.

Type : India : Sikkim : Lachung, 2745 m, *Gammie* 353.

Corybas purpurens Joseph et Yoganarasimhan (Orchidaceae).

Indian For. 93(12) : 815-817. 1967.

Holotype : India : Meghalaya : Elephant Falls, Shillong, ±1500 m, united Khasi & Jaintia Hills Dist., June 27, 1966, Joseph 36892A.

Paratype : Ibid., Oct. 16, 1965, Joseph 36891A.

Cotoneaster aitchisonii Schneider (Rosaceae).

Ill. Handb. Laubholz. 1 : 749. 1906; Bull. Bot. Surv. India 28 : 66. 1986 (1988).

Isotype : Afghanistan : Kuram-Tal, 2500 m, 1880, Aitchison 277.

C. cavei Klotz (Rosaceae).

Bull. Bot. Surv. India 5(3 & 4) : 213. 1963 (1964); Ibid., 28 : 67. 1986 (1988).

Type : India : Sikkim Himalaya: Prope, Kalapokhri, 3200 m, May 28, 1963, G. Klotz s. n.

C. duthieanus (Schneider) Klotz (Rosaceae).

Bull. Bot. Surv. India 5(3 & 4) : 211. 1963 (1964).

Isotype : India : Kumaon, Birch forest near Naffi in the Kutti Valley, 12-13000 ft., Sept. 11, 1884, Leg. J. F. Duthie.

C. falconeri Klotz (Rosaceae).

Wiss. Zeits. Univ. Halle. Math.-Nat. 15 : 535. 1966; Bull. Bot. Surv. India 28 : 68. 1986 (1988).

Holotype : N. Pakistan : Chitral, Jambatai, May 5, 1895, S. A. Harris 16109.

C. gilgitensis Klotz (Rosaceae).

Wiss. Zeits. Univ. Halle. Math.-Nat. 15 : 539. 1966; Bull. Bot. Surv. India 28 : 69. 1986 (1988).

Holotype : Kashmir, Nittar Valley, North of Gilgit, 3300-3600 m, Aug. 4, 1892, J. F. Duthie 12385.

C. kaganensis Klotz (Rosaceae).

Wiss. Zeits. Univ. Halle. Math.-Nat. 15 : 541. 1966; Bull. Bot. Surv. India 28 : 70. 1986 (1988).

Isotype : Pakistan: Hazara, Kagan Valley, Bhorji, July 12, 1897, Duthie's Collector vide Duthie 21276.

C. khasiensis Klotz (Rosaceae).

Bull. Bot. Surv. India 5(3 & 4) : 209. 1963 (1964).

Types : India : Meghalaya : Khasia, Laitlyngkot, 1650 m, May 16, 1886, Leg. C. B. Clarke No. 43, 911,

Cotoneaster nagensis Klotz (Rosaceae).

Bull. Bot. Surv. India 5 : 212. 1964 & 28 : 76. 1986 (1988).

Paratypes : Naga Hills, near Chukka forest, 2196 m, Dec. 1886, *D. Prain s. n.*; Khoima, Naga Hills, Dec. 1886, *D. Prain s. n.*

C. obovatus Wall. ex Dunn. (Rosaceae).

Misc. Infra. Kew Bull. 1921 : 119 (1921); Bull. Bot. Surv. India 28 : 76. 1986 (1988).

Isolectotype : Chenab Valley, Cherer near Sinhe, 2440 m, June 3, 1881, *R. Ellis* 1192.

C. sandakphuensis Klotz (Rosaceae).

Bull. Bot. Surv. India 5 : 213. 1964 & 28 : 78. 1986 (1988).

Holotype : Sikkim-Himalaya, Prope Sandakphu, infruticetis via, Rimbick, Oct. 9, 1941, *K. Biswas* 5721.

C. stracheyi Klotz (Rosaceae).

Wiss. Zeits. Univ. Halle. Math.-Nat. 15 : 854. 1966; Bull. Bot. Surv. India 28 : 79. 1986 (1988).

Type : India : Kumaon, Dwali, 2500 m, May 1848, *R. Strachey & Winterbottom* 1.

C. wallichianus Klotz (Rosaceae).

Wiss. Zeits. Univ. Halle. Math.-Nat. 15 : 541. 1966; Bull. Bot. Surv. India 28 : 79. 1986 (1988).

Holotype : Sikkim, Tankra, 3000 m, Aug. 5, 1892, *G. A. Gammie* 634.

Craibiodendron mannii W. W. Sm. (Ericaceae).

Notes R. bot. Gdn. Edinb. 24 : 159. 1912; J. Bombay Nat. Hist. Soc. 79(1) : 223. 1982.

Type : India : Meghalaya : Juudmaka Pahar, 6000 ft., Aug. 1891, *G. Mann s. n.*

Crotalaria beddomeana Thoth. et Ansari (Leguminosae).

Bull. Bot. Surv. India 20 : 180. 1978 (1979).

Isotype : India : Pulney Hills, 1958, *Beddome* 2.

C. decasperma Naik (Leguminosae).

Indian For. 92(12) : 760. 1966.

Holotype : Ghatangri forest, 4 km north of Osmanabad Town, Nov. 15, 1964, *Naik* 976A.

Isotypes : *Naik* 976B-C.

Crotalaria khasiana Bal. ex Thoth. et Ansari (Leguminosae).

Bull. Bot. Surv. India 20 : 180. 1978 (1979).

Isotype : India : Meghalaya : Khasia, 1520-2128 m, *J. D. Hooker & T. Thomson s. n.*

C. khasiana Bal. ex Thoth. et Ansari var. **macrophylla** Thoth. et Ansari (Leguminosae).

Bull. Bot. Surv. India 20 : 180. 1978 (1979).

Holotype : Burma : Peinnegon Yegyauny, Maymyo Dist., 1200 m, Dec. 6, 1924, *Ingkan* 592.

C. kodaiensis Debberm. et Biswas (Leguminosae).

J. Ind. Bot. Soc. 56 : 59-60. 1937.

Type : India : Madras : Kodaikanal Hills, 360 m, Jan. 1916, *C. Tomtand* 1605.

C. madurensis Wt. var. **kurnoolica** Ellis et Swamin. (Leguminosae).

J. Bombay Nat. Hist. Soc. 66(1) : 227. 1969.

Holotype : India : Andhra Pradesh : Srisailam, Karnool Dist., Oct. 21, 1964, *Ellis* 22144A.

C. paniculata Willd. var. **nagarjunakondensis** Thoth. (Leguminosae).

Bull. Bot. Surv. India 6 : 66. 1964 (1965).

Holotype : India : Andhra Pradesh : from the Hills surrounding the Nagarjunakonda Valley, Nalkonda Dist., 300 m, Nov. 26, 1961, *Thothathri* 9764A.

Isotypes : *Thothathri* 9764B-E.

C. sulphizzi Thoth. et Ansari (Leguminosae).

Ind. J. For. 2(3) : 282. 1979.

Type : Burma : Pegu (without precise locality) Jan. 4, 1871, *S. Kurz* 2508.

Croton caudatus Geisel. var. **obovoideus** Balakr. & T. Chakrab.

(Euphorbiaceae).

Bull. Bot. Surv. India 25(1-4) : 190-191. 1983 (1985).

Paratypes : India : Kerala : Travancore, April 26, 1896, *Bourdillon* 826; Ibid., no date, *Calder & Ramaswami* 1345; Ibid., Achankoil to Makona, Sept. 5, 1913, *Rama Rao* 1594; Tamil Nadu: Coimbatore Dist., Anaimali Hills, Aug. 30, 1913, *Fischer* 3638; Tirunelveli Dist., Shencottah to Aryankavu, Sept. 8, 1913, *Calder & Ramaswamy* 685.

Croton robustus Kurz var. **serratus** T. Chakrab., Balakr. & Gupta (Euphorbiaceae).

J. Econ. Tax. Bot. 9(2) : 251. 1987.

Holotype : Burma : Tavoy Valley of Sedi chamilf., alt. 2800 ft., ca 850 m, March 12, 1919, P. T. Russell 31.

Cryptocarya insularis Vasud. & T. Chakrab. (Lauraceae).

J. Econ. Tax. Bot. 6(2) : 446. 1985.

Isotype : India : Andaman Nicobar Islands: South Andaman Island, Dhanikhari, Feb. 6, 1974, N. G. Nair 852.

Curcuma vamana Sabu et Mangaly (Zingiberaceae).

J. Econ. Tax. Bot. 10(2) : 307. 1987.

Type : India : Kerala: Palghat Dist., Monsoon forest, July 20, 1984, Sabu et Mangaly Cu 37343B.

Cyanotis arcotensis Rolla Rao (Commelinaceae).

Blumea 14 : 345. 1967.

Holotype : India : Tamil Nadu: North Arcot Dist., Tippukadu reserve forest, 160 m, Nov. 10, 1963, fl., fr., Joseph 89886A.

C. cerifolia Rolla Rao et Kammathy (Commelinaceae).

J. Linn. Soc. Bot. London 59 : 306. 1966.

Holotype : India : From the experimental garden, Botanical Survey of India, Poona (from cultivated material raised from the cuttings of wild plants growing in weverly Estate, Anamalai Hills, Coimbatore Dist., Tamil Nadu), Aug. 8, 1962, Kammathy 77785A.

Cyathea balakrishnanii Dixit et Tripathi (Cyatheaceae).

Bull. Bot. Surv. India 26(3-4) : 170. 1984 (1986).

Isotype : India : Madhya Pradesh: Pachmari, Dhupgarh, ± 1000 m, Oct. 16, 1984, Dixit 41231A.

C. holttumiana R. R. Rao et N. S. Jamir (Cyatheaceae).

J. Econ. Tax. Bot. 6(1) : 207. 1985.

Holotype : India : Nagaland : Changki, Mokokchung Dist., Jamir 7996A.

C. nicobarica Balakr. et R. D. Dixit (Cyatheaceae).

Indian Journ. Forestry 10(1) : 43. 1987.

Paratype : India : Nicobar Island : Katchal Island, ± 160 m, Nov. 24, 1864, s. n., Acc. No. 1290.

***Cynandrium guttatum* Stapf (Melastomataceae).**

Hook. IC. Pl. 25 : t. 2419. 1895; Bull. Bot. Surv. India 10(3 & 4) : 335. 1968 (1969).

Isotype : *Haviland* 1549.

***Cynodon barberi* Rang. et Tad. (Gramineae).**

J. Bombay Nat. Hist. Soc. 24 : 846. 1916; Bull. Bot. Surv. India 8 : 204. 1966 (1967).

Lectotype selected : India : Agricultural College, Central Farm, Coimbatore, Aug. 1916, Madras Herbarium, S. India Flora No. 13715.

***C. barberi* Rang. et Tad. f. *longifolium* Jain (Gramineae).**

Indian For. 92(11) : 699. 1966.

Holotype : India : Orissa : Puri, 1889, *J. H. Walsh s. n.*

Paratypes : India : W. Bengal : Bakura, Adhkata, July 20, 1955, *Ganguly s. n.* CAL 1798; Bihar : Chaibasa, Rains, 1898, *s. l., s.n.*; Andhra Pradesh : Nagarjunkonda Valley, July 15, 1961, *Thothathri* 9683; Madras ; Madras Presidency, Precise locality ?, *Ramaswami* 541.

***Cynometra iripa* Kostel (Leguminosae).**

Allg. Med.-Pharm. Fl. 4 : 1341. 1835; Reinwardtia 10 : 66. 1982.

C. mimosoides Wall.

Type : *Wallich* 5817A.

***Cyperus atkinsoni* C. B. Clarke (Cyperaceae).**

J. Linn. Soc. 21 : 109. 1884.

Isotype : India : Kashmir : Aug. 1874, *Atkinson* 24196.

***C. atroglumosa* Govind. (Cyperaceae).**

Proc. Indian Acad. Sci. 81B (5) : 187. 1975.

Paratype : Karnataka : Nalur Oct. 7, 1961, *Govindarajalu* 4823.

***C. lurida* Govind. (Cyperaceae).**

Proc. Indian Acad. Sci. 81B(5) : 189. 1975.

Isotype : India : Tamil Nadu : Akkamalai, Valparai, 5400-6200 ft., Jan. 3, 1971, *Govindarajalu* 10742F.

***C. macer* Clarke (Cyperaceae).**

J. Linn. Soc. 21 : 160. 1884.

Isosyntypes : Burma : Pegu : *Kurz* 671; Bangladesh : Chittagong, *C. B. Clarke* 19779C.

Cyperus pentabracteatus Govind. et Hemadri (Cyperaceae).

Proc. Indian Acad. Sci. 82B(6) : 205. 1975.

Holotype : India : Maharashtra : Poona, Durga Khilla, Malavandara, 17-18 miles west of Junnar, Poona, *Hemadri* 107562A.

C. polyanthelus Govind. (Cyperaceae).

Proc. Indian Acad. Sci. 88 : 229-242 (1979).

Isotype : S. India : Manalar, Vattaparai, Feb. 1, 1969, *E. Govindarajalu* 4286B.

C. sanguinolentus Vahl subsp. **sacer** Koila. (Cyperaceae).

Bull. Bot. Surv. India 9 : 235. 1967 (1968).

Holotype : India : W. Bengal : Sodepur, July 3, 1964, *Korlaahalli* 232A.

Isotype : *Korlaahalli* 232B.

C. stricticulmis Govind. (Cyperaceae).

Proc. Indian Acad. Sci. 81B (5) : 194. 1975.

Isotype : India : Tamil Nadu : Balparai, Oct. 7, 1968, *Govindarajalu* 9042.

Cyrtandra burttii Balakr. (Gesneriaceae).

Notes R. bot. Gdn. Edinb. 35(1) : 115. 1976.

Holotype : India : Great Nicobar Island : 17 km from Campbell Bay to Alexander River, alt. ± 75m, above s.l., shaded stream sides in ever green hill forests, Aug. 21, 1975, *Balakrishnan* 3025A.

Isotype : Ibid., *Balakrishnan* 3025B.

C. occidentalis Balakr. et Burtt (Gesneriaceae).

Notes R. bot. Gdn. Edinb. 37(1) : 153. 1978.

Holotype : India : Great Nicobar Island : Campbell Bay to Chengappa Bay, 25 m, June 11, 1977, in flower, *Balakrishnan* 5700A.

Cyrtomium beddomei S. R. Ghosh (Dryopteridaceae).

J. Econ. Tax. Bot. 5(2) : 495. 1984.

Type : India : Tamil Nadu : Coonoor, Nilgiris, *Beddome* 19.

Dalbergia bhutanica Thoth. (Leguminosae).

Bull. Bot. Surv. India 14 : 189. 1972 (1975).

Holotype : Bhutan : Narchu Valley, Engo forest, Nov. 1898, *Prain* s. n.

Isotype : *Prain* s. n.

D. burmanica Prain (Leguminosae).

J. Asiat. Soc. Beng. 66 : 448. 1897 emend. Thoth.; Thoth., Tax. Revi Dalbergieae. P. 131. 1987.

Lectotype designated : Upper Burma: Ruby mines, March 1892, *Abdul Huk* 7A.

Syntype : Ibid., *Abdul Huk* 12.

Dalbergia cana Grah. (Wall. Cat. 5859. 1832, nomen) ex Kurz (Leguminosae).

J. Asiat. Soc. Beng. 42 : 70. 1873; Thoth., Tax. Revi. Dalbergieae P. 172. 1987.

Isotype : Moulmein, 1827, Wall. Cat. 5859.

D. candenatensis (Dennst.) Prain (Leguminosae).

J. Asiat. Soc. Beng. 70 : 49. 1901; Thoth., Tax. Revi. Dalbergieae P. 187. 1987.

D. torta Grah.

Type : *D. torta* : Penang : Singapore, 1832, Wall. Cat. 5873A.

D. clarkei Thoth. (Leguminosae).

J. Jap. Bot. 46(3) : 73. 1971; Thoth., Tax. Revi. Dalbergieae P. 114. 1987.

Isotype : India : Assam : Maoksandram, Khasia, 133 m, Jan. 1886, Clarke 42875B.

D. confertiflora Benth. (Leguminosae).

Miq. in Pl. Jungh. 1 : 225. 1852 et J. Linn. Soc. 4 (Suppl.) 41. 1860; Thoth., Tax. Revi. Dalbergieae P. 102. 1987.

Isolectotype : Bangladesh : Sylhet, Wall. Cat. 5448 I.

Syntype : Oudh, Wall. Cat. 5848E.

D. confertiflora Benth. var. *listeri* Thoth. (Leguminosae).

Bull. Bot. Surv. India 17 : 66. 1975 (1978).

Holotype : Bangladesh : Chittagong, Feb. 1876, *Lister* 8.

D. coromandeliana Prain (Leguminosae).

J. Asiat. Soc. Beng. 70 : 60. 1901 ; Thoth., Tax. Revi. Dalbergieae P. 126. 1987.

Lectotype : S. India: Shevagiri Hills, Tinnevalley Dist., *Wight* 798.

D. cultrata Grah. (Wall. Cat. 5861. 1832, nomen) ex Benth. (Leguminosae).

In Miq. Pl. Jungh. 1 : 254. 1852 et J. Linn. Soc. 4 (Suppl.) 39. 1860; Thoth., Tax. Revi. Dalbergieae P. 91. 1987.

Lectotype designated : Prome, 1826, Wall. Cat. 5861A.

Dalbergia cultrata Grah. ex Benth. var. **cultrata fusa** (Pierre ex Prain)

Thoth. (Leguminosae).

Thoth., Tax. Revi. Dalbergieae P. 94. 1987.

Holotype : Cochin China, Baochinah, Bin Hoa, 1879, *Pierre* 1706.

D. cultrata Grah. ex Benth. var. **maymyensis** Thoth. (Leguminosae).

Bull. Bot. Surv. India 17 : 65. 1975 (1978).

Holotype : Upper Burma : Maymyo plateau, 1200 m, July 1908, *J. H. Lace* 4141.

D. cultrata Grah. ex Benth. var. **tavoyensis** Thoth. (Leguminosae).

Bull. Bot. Surv. India 17 : 65. 1975 (1978).

Holotype : Burma : Tavoy : Tenasserim, Sept. 1901, *Manson* 71.

D. duarensis Thoth. (Leguminosae).

Bull. Bot. Surv. India 14 : 192. 1972 (1975).

Holotype : India : Mahakalaguri, Alipore Duars, Oct. 1891, *Heawood* 104.

D. gardneriana Benth. (Leguminosae).

J. Linn. Soc. 4 (Suppl.) : 42. 1860 ; Thoth., Tax. Revi. Dalbergieae P. 81. 1987.

Isolectotype : Nilgherries, *Hohenacker* 1591.

D. hemsleyi Prain (Leguminosae).

J. Asiat. Soc. Beng. 66 : 450. 1897 et Ann. Roy. Bot. Gard. Calc. 10(1) : 94. 1904.

Lectotype selected : Burma : Myingyin, Nov. 1890, *Prazer s. n.*

D. horrida (Dennst.) Mabberly (Leguminosae).

Taxon 25 : 538. 1977; Thoth., Tax. Revi. Dalbergieae P. 105. 1987.

D. multiflora Heyne

Wall. Cat. 5848. B, 1832 (nomen).

Isotype : Courtallum, Wall. Cat. *Heyne* 5848B.

D. horrida (Dennst.) Thoth. var. **concanensis** Thoth.

Thoth., Tax. Revi. Dalbergieae P. 110. 1987 (Leguminosae).

Holotype : India : Karnataka : Concan, Collector? *s. n.*, Acc. No. 130711.

D. horrida Thoth. var. **glabrescens** (Prain) Thoth. et Nair (Leguminosae).

Taxon 30 : 46. 1981; Thoth., Tax. Revi. Dalbergieae P. 109. 1987.

Lectotype designated : South India : Nagercoil, Nov. 1893, *Lawson* 218.

Dalbergia kingiana Prain (Leguminosae).

J. Asiat. Soc. Beng. 67 : 289. 1898 et *Ibid.* 70 : 46. 1901 et Ann.
Roy. Bot. Gard. 10(1) : 61. 1904; Thoth., *Tax. Revi. Dalbergieae* P. 78. 1987.
Holotype : Upper Burma : Myitkyina, Kachin Hills, Nov. 1987, *Shaik Mokim s. n.*

D. kurzii Prain (Leguminosae).

J. Asiat. Soc. Beng. 66 : 450. 1897 et *Ibid.* 70 : 50. 1901; Thoth.,
Tax. Revi. Dalbergieae P. 175. 1987.

Lectotype designated : Burma : Pegu, *Kurz* 2603.

Syntypes : *Ibid.*, *Kurz* 2608; Pegu Yomah, *Kurz* 1780; Pegu, *Maclelland s. n.*; Thykuolah, *Brandis* 1170; upper Burma, 1894, *Prazer s. n.*; Shan State, Nov.-Dec., 1887, *Alpin s. n.*

D. lanceolaria Linn. f. **ssp. paniculata** (Roxb.) Thoth. (Leguminosae).

Bull. Bot. Surv. India 25 (1-4) : 171. 1983 (1985).

D. paniculata Roxb.

Corom. Pl. 2 : 8 t. 114. 1798.

D. nigrescens Kurz

Pegu Rep. App. A. 48. B. 45. 1875.

Lectotype selected : Plate 114 of Roxb. *Corom. Pl.* 1798.

D. lanceolaria Linn. f. **ssp. paniculata** var. **hemsleyi** (Prain) Thoth. (Leguminosae).

Bull. Bot. Surv. India 25 (1-4) : 172. 1983 (1985).

D. hemsleyi Prain

J. Asiat. Soc. Beng. 66 : 450. 1897 et Ann. *Roy. Bot. Gard. Calc.* 10 (1) : 94. 1904.

Lectotype selected : Burma : Myingyin, Nov. 1890, *Prazer s. n.*

D. lanceolaria Linn. f. **ssp. paniculata** var. **maymyensis** (Craib) Thoth. (Leguminosae).

Bull. Bot. Surv. India 25 (1-4) : 172. 1983 (1985). ; Thoth., *Tax. Revi. Dalbergieae* P. 153. 1987.

D. maymyensis Craib

Kew Bull. 390. 1912.

Lectotype selected : Burma : Maymyo Plateau, 1200 m, July 1908, *Lace* 4134.

Syntypes : *Ibid.*, *Lace* 5793 (April 1912), 5848 (July 1912), 3113 (April 1908).

Dalbergia oliveri Gamble ex Prain (Leguminosae).

J. Asiat. Soc. Beng. 66 : 451. 1893 emend. Thoth.; Thoth., Tax. Revi.

Dalbergieae P. 134. 1987.

Lectotype designated : Upper Burma: Wautho, Feb. 1892, *Oliver s. n.*

D. ovata Grah. (Wall. Cat. 5854, 1832, nomen) ex Benth. (Leguminosae).

Miq. Pl. Jungh. 1 : 254. 1852 et Journ. Linn. Soc. 4 (Suppl.). 40. 1860; Thoth., Tax. Revi. Dalbergieae P. 58. 1987.

Isotype : Burma : Moolmyn, 1827, Wall. Cat. 5854.

D. ovata Grah. ex Benth. var. **glomeriflora** (Kurz) Thoth. (Leguminosae).

Bull. Bot. Surv. India 25 (1-4) : 170. 1983 (1985).

D. glomeriflora Kurz

J. Asiat. Soc. Beng. 42 : 70. 1873.

Type : Burma : Prome, Kurz 211.

D. ovata Grah. ex Benth. var. **puberula** Kurz (Leguminosae).

J. Asiat. Soc. Beng. 45 : 280. 1876 emend. Thoth.; Thoth., Tax. Revi. Dalbergieae P. 59. 1987.

Holotype : Pegu : Irawaddy Dist., *Brandis s. n.*

D. peguensis Thoth. (Leguminosae).

J. Jap. Bot. 50 (2) : 55. 1975.

Holotype : Burma : Pegu, Yomah, Mar. 1871, *Kurz* 2604.

Paratype : Ibid., Jan. 1868, *Kurz* 1781.

D. pinnata (Lour.) Prain var. **acaciaefolia** (Dalz.) Thoth. (Leguminosae).

Bull. Bot. Surv. India 25 (1-4) : 170. 1983 (1985).

D. acaciaefolia Dalz.

Kew Journ. Bot. 2 : 37. 1850.

D. tamarindifolia Roxb. var. *acaciaefolia* (Dalz.) Baker

Hook. f. Fl. Brit. Ind. 2 : 235. 1876.

Isotype : Concan: *Law s. n.*

D. prainii Thoth. (Leguminosae).

J. Jap. Bot. 50 (2) : 52. 1975.

Holotype : Upper Burma : Kalay and Chin Hills, June, 1894, *J. C. Prazer* 39.

D. pseudo-sissoo Miq. (Leguminosae).

Fl. Ind. Bat. 1 : 128. 1855; Thoth., Tax. Revi. Dalbergieae P. 67. 1987.

D. rostrata Grah. (Wall. Cat. 5867, 1832, nomen) ex Prain

J. Asiat. Soc. Beng. 70 : 45. 1901 et Ann. Roy. Bot. Gdn. (1) : 60. 1904.

Isotype : *D. rostrata* : Singapore, Wall. Cat. 5867.

Dalbergia pseudo-ovata Thoth. (Leguminosae).

J. Jap. Bot. 50 (2) : 52. 1975.

Holotype : Burma : Kangyi reserve, 24 m, Tharawaddy Dist., Feb. 1912, *Lace* 5707.

D. reniformis Roxb. (Hort. Beng. 53. 1814. nomen) (Leguminosae).

Fl. Ind. 3 : 226. 1832; Thoth., Tax. Revi. Dalbergieae P. 194. 1987.

D. flexuosa Grah.

Type : Sylhet, 1832, Wall. Cat. 5875.

D. rimosa Roxb. var. *foliacea* (Wall. ex Benth.) Thoth. (Leguminosae).

Bull. Bot. Surv. India 25 (1-4) : 170. 1983 (1985).

D. foliacea Wall. ex Benth.

Miq. Pl. Jungh. 1 : 255. 1852.

Lectotype selected : Amherst and Maulmyne, 1927, Wall. Num. List. No. 5856A.

D. rimosa Roxb. var. *laevis* Thoth. (Leguminosae).

Bull. Bot. Surv. India 17 : 64. 1975 (1978).

Holotype : Burma : Nahmedaung, Henzada, Feb. 1903, *Shaik Mokin* 1355.

D. spinosa Roxb. (Hort. Beng. 98. 1914, nomen) (Leguminosae).

Fl. Ind. 3 : 233. 1832 ; Thoth., Tax. Revi. Dalbergieae P. 193. 1987.

Type : Chittagong : *Roxburgh s. n.*

D. stipulacea Roxb. var. *kurzii* Thoth. (Leguminosae).

Bull. Bot. Surv. India 17 : 67. 1975 (1978).

Holotype : Burma : Martaban, *Kurz* 1774.

D. stipulacea Roxb. var. *mogkokensis* Thoth. (Leguminosae).

Bull. Bot. Surv. India 17 : 68. 1975 (1978).

Holotype : Burma : Mogkok, 1334 m, Ruby mines division, May 1910, A. Rodger 332.

D. stipulacea Roxb. var. *stipulacea* f. *puberula* Thoth. (Leguminosae).

Bull. Bot. Surv. India 17 : 68. 1975 (1978).

Holotype : Burma : Mandalay Dist., 1200 m, May 1915, G. Rogers 968.

D. tinneveliensis Thoth. (Leguminosae).

Ceylon Journ. Sci. (Bio. Sci.) 12 : 47. 1970.

Holotype : India : Tamil Nadu: Tinnevelly Dist., Jan. 1917, Fischer 4045.

Dalbergia thomsonii Benth. (Leguminosae).

J. Linn. Soc. 4 (Suppl.) : 33. 1860; Thoth., Tax. Revi. Dalbergieae P. 171. 1987.

Lectotype designated : India : Khasia, 600-1250m, *Hooker & Thomson s. n.*

D. velutina Benth. (Leguminosae).

Miq. Pl. Jungh. 1 : 225. 1852 et in J. Linn. Soc. 4 (Suppl.) 43. 1860; Thoth., Tax. Revi. Dalbergieae P. 99. 1987.

Isolectotype : Bangladesh : Sylhet, Wall. Cat. 5868B.

Syntype : Moolmyne, 1827, Wall. Cat. 5868A.

D. velutina Benth. var. *maingayi* Prain (Leguminosae).

J. Asiat. Soc. Beng. 66 : 117. 1897; Thoth., Tax. Revi. Dalbergieae P. 100. 1987.

Lectotype designated : Malacca, 1867-68, Malaya, *Maingay* 612.

Syntypes : Singapore, Malaya, 1894, *Ridley* 5923, 6086.

D. velutina Benth. var. *velutina* f. *burmanica* Thoth. (Leguminosae).

Bull. Bot. Surv. India 17 : 70. 1975 (1978).

Holotype : Burma : Mergui, April 1909, *Lace* 4787.

D. volubilis Roxb. var. *assamica* Thoth. (Leguminosae).

Bull. Bot. Surv. India 17 : 67. (1975) 1978.

Holotype : India : Meghalaya: Myantang Valley, Khasia and Jaintea Hills, Jan. 1916, *U. Kanjilal* 6574.

D. wattii Clarke (Leguminosae).

J. Linn. Soc. 25 : 17. 1889; Thoth.; Tax. Revi. Dalbergieae P.177. 1987.

Holotype : Moneypoor, Mayung, 1200 m, Nov. 1885, *Clarke* 42034.

D. yunnanensis Franch var. *collettii* (Prain) Thoth. (Leguminosae).

Bull. Bot. Surv. India 25 (1-4) : 171. 1983 (1985).

D. collettii Prain

J. Asiat. Soc. Beng. 66 : 445. 1897 P.P. et Ann. Roy. Bot. Gard. Calc. 10(1) : 53. 1904.

Dalbergia sp. Collett & Hemsley

J. Linn. Soc. 28 : 50. 1891.

Lectotype selected : Kawlo, 1666 m, Shan Hills, April 1988, *Collett.* 591.

Dalechampia burmanica Mukherjee (Euphorbiaceae).

Bull. Bot. Soc. Beng. 3(1) : 9. 1949.

Type : Burma : Myaukhaing reserve, July 12, 1949. *D. R. Pokhant* 562.

Decaisnea insignis (Griff.) Hook. f. & Thoms. (Lardizabalaceae).

Proc. Linn. Soc. 2 : 350. 1854; Fasc. Fl. Ind. 19 : 31. 1988.

Slakia insignis Griff.

Itin. Notes 187. 1848.

Type : Bhutan: *Griffith* 1838.

Decaschistia crotonifolia Wt. & Arn. (Malvaceae).

Prod. Fl. Pen. Ind. Or. 1 : 52. 1834; Fasc. Fl. Ind. 19 : 104. 1988.

Isotype : *Wight* Cat. No. 215.

D. cuddapahensis T. K. Paul et Nayar (Malvaceae).

Geobios New Reports 2 : 156-157. 1983; Fasc. Fl. Ind. 19 : 105. 1988.

Holotype : India : Andhra Pradesh : Swamipadhalu-Kodur, Cuddapah Dist., 550 m, July 23, 1962, *J. L. Ellis* 14332.

Delphinium malabaricum (Huth.) Munz var. *ghaticum* Billiore (Ranunculaceae).

Indian For. 99(7) : 436-439. 1973.

Holotype : India : Maharashtra : on the exposed slopes of Dhamni Hill, ± 700 m, near Mokhada, Vihigaon forest range, Oct. 19, 1967, *K. V. Billiore* 112948A.

Dendrobium assamicum Chowdhury (Orchidaceae).

Kew Bull. 43(4) : 667. 1988.

Holotype : India : Assam : Nowgong Dist., Outskirt of village Nellie, 45 miles from Guwahati towards Nowgong, 26° 15' N & 92° 30' E, March 31, 1969, *S. Chowdhury* 354.

D. diodon Reinb. f. ssp. *kodayarensis* Gopalan et Henry (Orchidaceae).

J. Econ. Tax. Bot. 12(2) : 487. 1988.

Holotype : India : Tamil Nadu: Kannya Kumari, Upper Kodayar, *Gopalan* 81452.

Derris andamanica Prain (Leguminosae).

J. Asiat. Soc. Beng. 66(2) : 104. 1897; Fasc. Fl. Ind. 8 : 7. 1982.

Lectotype : India : South Andaman : North corbyn' scove hill jungle, Jan. 1895, *King's Collector* s. n.

D. benthamii (Thw.) Thw. var. *benthamii* (Leguminosae).

Enum. Pl. Zeyl. 413. 1864; Fasc. Fl. Ind. 8 : 9. 1982.

Isotype : Ceylon : Central Province, *Thwaites* 2925.

Derris benthamii (Thw.) Thw. var. **wightii** (Baker) Thoth. (Leguminosae).

J. Jap. Bot. 51(5) : 145. 1976; Fasc. Fl. Ind. 8 : 11. 1982.

Isotype : India : Shevagherry Hills, Aug. 1936, *Wight* 834.

D. canarensis (Dalz.) Baker (Leguminosae).

Hook. f. Fl. Brit. Ind. 2 : 246. 1878; Fasc. Fl. Ind. 8 : 12. 1982.

Neotype : India : Karnataka : Kassarleh, Kanara, May 1853, *Dr. Ritchie* 1720.

D. cuneifolia Benth. var. **cuneifolia** f. **cuneifolia** (Leguminosae).

Miq. Pl. Jungh. 1 : 253. 1852 et Journ. Linn. Soc. 4 (Suppl.) : 112. 1860; Fasc. Fl. Ind. 8 : 13. 1982.

Lectotype : Nepal : Noakote, 1820, Wall. Num. List. No. 5887.

D. cuneifolia Benth. var. **cuneifolia** f. **assamica** Thoth. (Leguminosae).

Bull. Bot. Surv. India 3(2) : 192. 1961 (1962); Fasc. Fl. Ind. 8 : 15. 1982.

Holotype : India : Assam : Bishnupur, Cachar Dist., Dec. 1914,

U. Kanjilal 4835.

D. cuneifolia Benth. var. **longipedicellata** Thoth. (Leguminosae).

Bull. Bot. Surv. India 3(2) : 191. 1961 (1962); Fasc. Fl. Ind. 8 : 15. 1982.

Holotype : India : Sikkim : Sivoke, April 1908, *Ribu* 760.

D. elegans Benth. f. **andamanensis** Thoth. (Leguminosae).

Bull. Bot. Surv. India 3(2) : 184. 1961 (1962); Fasc. Fl. Ind. 8 : 16. 1982.

Holotype : India : Manipur Hill Jungle, South Andaman, Feb. 1894, *King's Collector s. n.*

D. elliptica (Wall.) Benth. var. **elliptica** (Leguminosae).

J. Linn. Soc. 4(Suppl.) : 111. 1860; Bull. Bot. Surv. India 3 : 194. 1961; Fasc. Fl. Ind. 8 : 16. 1982.

Isotype : Amboina, cultivated in Botanic Garden, Calcutta. Wall. List No. 5881A.

Paratype : Burma : Attran, 1827, Wall. List. No. 5881B.

D. elliptica Benth. var. **chittagongensis** Thoth. (Leguminosae).

Bull. Bot. Surv. India 3 : 195. 1961 (1962).

Holotype : Bangladesh: Chittagong: Kodla, in Hills, May 1887, *Badul Khan* 450A.

Isotypes : *Badul Khan* 450 B-E.

Derris feruginea (Roxb.) Benth. (Leguminosae).

Miq. Pl. Jungh. 1 : 252. 1852 et J. Linn. Soc. 4(Suppl.) : 109. 1860;
Fasc. Fl. Ind. 8 : 18. 1982.

Isolectotype : Bangladesh : Sylhet, Wall. Num. List. No. 5885A.

D. henryi Thoth. (Leguminosae).

J. Jap. Bot. 45(1) : 6. 1970.

Holotype : China : Yunnan : 1897, *Henry* 10893A.

Isotype : Ibid., *Henry* 10893B.

D. heyneana (Wt. & Arn.) Benth. (Leguminosae).

Miq. Pl. Jungh. 1 : 252. 1852 et J. Linn. Soc. 4 (Suppl.) : 110. 1860;
Fasc. Fl. Ind. 8 : 19. 1982.

Isolectotype : India, without precise locality, Western Peninsula,
Wight 920.

D. lushaiensis Thoth. (Leguminosae).

Bull. Bot. Surv. India 12 : 104. 1970 (1972).

Holotype & Isotypes : India : Valley near Fort Lungleh, 1000-2000m,
South Lushai Hills, Assam, March 1899, *A. T. Gage* 23A-C.

D. marginata (Roxb.) Benth. (Leguminosae).

Miq. Pl. Jungh. 1 : 252. 1852, in adnot; Fasc. Fl. Ind. 8 : 22. 1982.

Lectotype : Bangladesh : Sylhet, H.B.C., Wall. Num. List. No. 5909A.

D. monticola (Kurz) Prain (Leguminosae).

J. Asia. Soc. Beng. 66(2) : 361. 1897; Fasc. Fl. Ind. 8 : 23. 1982.

Type : Burma : Nattoung, 2200-2400 m, Pegu, *Kurz* 1776.

D. ovalifolia (Wt. & Arn.) Benth. (Leguminosae).

Miq. Pl. Jungh. 1 : 252. 1852 *ex parte* et J. Linn. Soc. 4 (Suppl.) : 115.
1860; Fasc. Fl. Ind. 8 : 25. 1982.

Isotype : India : Western Peninsula, *Wight* 917.

D. pseudo-robusta Thoth. (Leguminosae).

Bull. Bot. Surv. India 3 : 181. 1961 (1962).

Holotype : India : Arunachal Pradesh : Kheti-Tincha, Tirap F. D., Aug.
1958, *Panigrahi* 14550A.

D. robusta (Roxb. ex DC.) Benth. var. **robusta** (Leguminosae).

J. Linn. Soc. 4(suppl.) : 104. 1860. *pro parte*; Fasc. Fl. Ind. 8 : 26.
1982.

Isolectotype : Bangladesh : Sylhet, Wall. Num. List No. 5849B.

Derris robusta (Roxb. ex DC.) Benth. var. **assamica** Thoth. (Leguminosae).

Bull. Bot. Surv. India 3(2) : 180. 1961 (1962); Fasc. Fl. Ind. 8 : 27. 1982.

Holotype and Isotype : India : Upper Shillong, 1933 m, June 1920, Carter 1696A&B.

D. scandens (Roxb.) Benth. var. **scandens** (Leguminosae).

J. Linn. Soc. 4(Suppl.) : 103. 1860; Fasc. Fl. Ind. 8 : 27. 1982.

Lectotype : Plate 192 of Roxb. Pl. Corom. 2 : 49. 1798.

D. sinuata Benth. ex Thw. (Leguminosae).

Enum. Pl. Zeyl. 93. 1859; Bull. Bot. Surv. India 3 : 195. 1961 (1962).

Pongamia sinuata Wall.

Wall. Cat. 5911. 1831-32, *nom. nud.*

Deguelia sinuata (Thw.) Taub.

Engl. & Prantl. Nat. Pfl. Fam. 3 (2) : 345. 1891.

Isotype : Ceylon Province: *Thwaites* 1491.

D. thothathrii Bennet (Leguminosae).

J. Indian Forestry 1(1) : 23. 1978; Fasc. Fl. Ind. 8 : 29. 1982.

Type : India : Concan, *Stocks, Law* 10.

Dichanthium foulkesii (Hook. f.) Jain et Despande (Poaceae).

Bull. Bot. Surv. India 20 : 134. 1978 (1979); Fasc. Fl. Ind. 15 : 12. 1984.

Syntype : India : Pykora, Aug. 1878, *King s. n.*

D. glabrum (Roxb.) Jain et Despande var. **glabrum** (Poaceae).

Bull. Bot. Surv. India in 20 : 134. 1978 (1979); Fasc. Fl. Ind. 15 : 13. 1984.

Type : Bengal: *Roxburgh s. n.* (Roxb. Ic. 1194).

D. glabrum (Roxb.) Jain et Despande var. **punctatum** (Roxb.) Despande (Poaceae).

Fasc. Fl. Ind. 15 : 14. 1984.

Type : India: *Roxburgh s. n.* (l. c. 892).

D. jainii (Desh. et Hem.) Despande (Poaceae).

Bull. Bot. Surv. India 21 : 198. 1979 (1981); Fasc. Fl. Ind. 15 : 17. 1984.

Holotype : India : Maharashtra : Pune, on the open top of Durga Khilla, ± 2200 m, about 30 km west of Junnar, *Hemadri* 104241A.

Dicliptera nasikensis Naras. et Sharma (Acanthaceae).

J. Econ. Tax. Bot. 7(2) : 481. 1985 (1986).

Holotype : India : Maharashtra : From fallow land of Dangsaundane round of Satna range, Nasik District, alt. of *ca* 675 m above *m.s.l.*, May 20, 1983, *Narasimhan* 163977A.

Dicraea filifolia Ramamurthy et Joseph (Podostemonaceae).

Bull. Bot. Surv. India 6 : 333. 1964 (1965).

Holotype : India : Kerala : Parambikulam, Submergible area, Trichur Dist., Nov. 22, 1962, *Ramamurthy* 15325A.

Isotype : Ibid., *Ramamurthy* 15325B.

Paratype : Ibid., June 19, 1964, *Ramamurthy* 20018A.

Dicranopteris linearis Underw. var. *wattii* Panigr. et Dixit (Gleicheniaceae).

Bull. Bot. Surv. India 13 : 162. 1971 (1973).

Holotype : India : Manipur : Kala Naga Hills, 195 m, May 1882, G. Watt 7497.

Digitaria pennata (Hochst.) T. Cooke var. *shettyana* Pandey, Parmar et Vyas (Gramineae).

J. Econ. Tax. Bot. 5(2) : 475. 1984.

Holotype : India : Rajasthan : Jalore, from rocky habitat near Sareh Mandir, Sept. 21, 1978, R. V. Shetty 6659A.

Dillenia andamanica Parkinson (Dilleniaceae).

Indian For. 61 : 452. t. 29. 1935; Hoogl. in Blumea 7(1) : 127. 1952; Fasc. Fl. Ind. 2 : 5. 1979.

Type : India : Middle Andaman: Bomlungta, April 12, 1916, *Parkinson* 1185.

D. bolsteri Merr. (Dilleniaceae).

Philip. J. Sc. Bot. 7. 1912. P. 305; Blumea 7(1) : 54. 1952.

D. cauliflora Merr.

Philip. J. Sc. Bot. 9. 1914 (1915). P. 517.

Isotype : *D. cauliflora* : Samar, April 1914, *Ramos* Bs 1695.

D. bracteata Wight (Dilleniaceae).

Ic. Pl. Ind. Or. 2 : 4. t. 358. 1840 (as *bractiata* in tab.); Blumea 7(1) : 84. 1952; Fasc. Fl. Ind. 2 : 6. 1979.

Type : India : Balaghaut Mountains near Madras, *Wight* 13.

Dillenia excelsa (Jack) Gilg (Dilleniaceae).

Engl. & Prantl. Nat. Pfl. Fam. 3. 6. 1893. P. 123; Hoogl. in Blumea 7(1) : 86. 1952.

D. magnoliaefolia Martelli.

Becc., Malesia 3. 1886. P. P. 155.

Isotype : *D. magnoliaefolia* : Sumatra, Padang, Aug. 1878, Beccari P. S. 548.

D. eximia Miq. (Dilleniaceae).

Fl. Ind. Bat. Suppl. 1860. P. 620; Hoogl. in Blumea 7(1) : 75. 1952.

Wormia scortechinii King

J. Asiat. Soc. Beng. 58. II. 1889. P. 365.

Isotype : *D. eximia* : Sumatra, Teysmann 4402 HB.

Type : *W. scortechinii* : Malacca, March 1883, Scortechini 1864.

D. indica Linn. (Dilleniaceae).

Sp. Pl. 535. 1753; Blumea 7(1) : 108. 1952.

D. elongata Miq.

Fl. Ind. Bat. 1. 2. 1859. P. 12.

Isotype : *D. elongata* : Lubu Alung, Sum, W. Coast, Teysmann 453.

D. mansoni (Gage) Hoogl. (Dilleniaceae).

Blumea 7(1) : 83. 1952.

Wormia mansoni Gage

J. Asiat. Soc. Beng. N.S. 2. 1906. P. 73.

Holotype : Burma : Banks of Yunzalin Riv. above its junction with Salween, May 1905, Manson s. n.

D. marsupialis Hoogl. (Dilleniaceae).

Fl. Mal. 1. 4. 1951. P. 162. descr. angl.; Blumea 7(1) : 66. 1952.

Isotype : Ramos & Edano BS 75247.

D. megalantha Merr. (Dilleniaceae).

Philip. J. Sc. Bot. 9. 1914 (1915). P. 519; Blumea 7(1) : 61. 1952.

D. mindanaensis Elm.

Leafl. Philip. Bot. 7. 1915. P. 2611.

Type : *D. mindanaensis* : Philippine : Mindanao, Sept. 1909, Elmer 11883.

D. monantha Merr. (Dilleniaceae).

Philip. J. Sc. Bot. 9. 1914. P. 321 ; Blumea 7(1) : 102. 1952.

Type : Tatay, Palawan, April 1913, Merrill 9237.

Dillenia ovalifolia Hoogl. var. *sericea* Hoogl. (Dilleniaceae).

Fl. Mal. I. 4. 1951. P. 159. Descr. angl.; Blumea 7(1) : 34. 1952.

Isotype : *de Hann* 1715.

D. pentagyna Roxb. (Dilleniaceae).

Pl. Corom. I. 1795. P. 21. t. 20. Blumea 7(1) : 117. 1952.

D. augusta Roxb.

Hort. Bengal 1814. P. 43. *nomen nudum*.

Isotype : *D. augusta* : Wallich 948.

D. philippiensis Rolfe (Dilleniaceae).

J. Linn. Soc. Bot. 21. 1884. P. 307 Hoogl. in Blumea 7(1) : 63. 1952.

D. catmon Elm.

Leafl. Philip. Bot. 7. 1915. P. 2610.

Isotype : *D. catmon* : Philippine: Cabadbaran (Mt Urdaaneta), Agusan Prov., Mindanao, Aug. 1912, Elmer 13564.

D. pulchella (Jack) Gilg (Dilleniaceae).

Engl. & Prantl. Nat. Pfl. Fam. 3. 6. 1893. P. 123; Blumea 7(1) : 113. 1952.

Wormia pulchella Jack

Mal. Misc. 2. 7. 1822. P. 70.

Type : *W. pulchella* : Singapore, Oct. 15, 1892, Ridley 3965.

D. reifferscheidia Villar (Dilleniaceae).

Nov. App., 1880. P. 3. based on *Reifferscheidia speciosa* Presl; Blumea 7(1) : 68. 1952.

D. reifferscheidia var. *rosea* Elm.

Leafl. Philip. Bot. 8. 1919. P. 3079.

Isotype : *D. reifferscheidia* var. *rosea*: Luzon, Laguna Prov., June-July 1917, Elmer 18025.

D. reticulata King (Dilleniaceae).

J. Asiat. Soc. Beng. 58. II. 1889. P. 367; Blumea 7(1) : 78. 1952.

Holotype : Malacca : Dipong. March 1883. Scortechinii 1840.

D. sibuyanensis (Elm.) Murr. (Dilleniaceae).

Merr. Philip. J. Sc. Bot. 9. 1914. P. 332 : Blumea 7(1) : 99. 1952.

Isotype : Magallanes, Isl. of Sibuyan, March 1910, Elmer 12078.

D. suffruticosa (Griff.) Martelli (Dilleniaceae).

Becc, Malesia 3. 1886. P. 163; Blumea 7(1) : 70. 1952.

Wormia subsessilis Miq.

Fl. Ind. Bat. Suppl. 1860. P. 619.

Isotype : *W. subsessilis* : Teysmann 3203.

Dimeria borii P. V. Sreekumar, V. J. Nair et N. C. Nair (Poaceae).

J. Econ. Tax. Bot. 3(2) : 657. 1982.

Holotype : India : Kerala : Calicut Dist., Kanjeerakkadavu near Kakkayam, ± 900 m, Nov. 26, 1981, *P. V. Sreekumar* 71855.

D. keralae N. C. Nair, P. V. Sreekumar and V. J. Nair (Poaceae).

J. Bombay Nat. Hist. Soc. 80 (3) : 626. 1983.

Holotype : India : Kerala : Paramba, Cannanore Dist., on way to Bandudka ± 150 m, Oct. 16, 1981, *P. V. Sreekumar* 71717.

D. kollimalayana M. Mohanan et A. V. N. Rao (Poaceae).

J. Bombay Nat. Hist. Soc. 80 (3) : 615. 1983.

Holotype : India : Tamil Nadu : Kollimalai, Salem Dist., 1200 m, Feb. 19, 1982, *M. Mohanan* 56208.

Dimorphocalyx balakrishnanii T. Chakrab. & Premanath (Euphorbiaceae).

J. Econ. Tax. Bot. 4(3) : 1013. 1983.

Holotype : India : Andaman-Nicobar Islands : Havelock Island, without locality, ± 40 m, May 15, 1974, *Ansari* 1368A.

Paratypes : India : South Andaman Island : Goplakabang, 1890, *King's Collector s. n.*, Acc. No. 412226; without locality, 1900, *Prain's Collector* 60.

D. dilipianus Balakr. & T. Chakrab. (Euphorbiaceae).

J. Econ. Tax. Bot. 4(3) : 1017. 1983.

Holotype : India : Andaman Nicobar Islands : South Andaman Island, Carbyn's cove, ± 20 m, Oct. 10, 1973, *Nair* 498A.

Diplocisia glaucescens (Bl.) Diels var. *kunstleri* (King) Pramanik et Thoth.

(Menispermaceae).

Kew Bull. 42(3) : 705. 1987.

Cocculus kunstleri King

J. Asiat. Soc. Beng. 58 : 384. 1889.

Syntypes : Malaya : Perak : ulu Bublong, *King's Collector (Kunstler)* 4417, 10282.

Diplospora andamanica Balakr. et N. G. Nair (Rubiaceae).

Bull. Bot. Surv. India 24 : 31. 1982 (1983).

Holotype : India : North Andamans : Saddle peak, ca 600 m, Dec. 1, 1976 *Balakrishnan & Nair* 6762A.

Diplospora majumdarii Gang. et T. Chakrab. (Rubiaceae).

J. Econ. Tax. Bot. 13(1) : 86. 1989.

Holotype : Perak, Malaya Peninsula, Gunonginas, 5000 ft., L. Wray Jr. 4086, Herb. Acc. No. 207289.

Isotype : Ibid., L. Wray Jr. 4086, Acc. No. 207290.

Paratype : Ibid., Maxwell's Hills, June, 1893, N. H. Reddy.

Dipterocarpus andamanicus (King) Tewary & Sarkar (Dipterocarpaceae).

Indian Journ. Forestry 10(1) : 63-64. 1987.

D. turbinatus Gaertn. f. var. *andamanica* King

J. Asiat. Soc. Beng. 62(2) : 92. 1893.

Lectotype : South Andaman : Hill Jungle, Dhanikhari, April 11, 1891, Dr. King s. n., Acc. No. 50541.

Drepanostachyum jainiana (Das et Pal) R. Majumdar (Gramineae).

Bull. Bot. Surv. India 25(1-4) : 235. 1983 (1985).

Chimonobambusa jainiana Das et Pal

J. Econ. Tax. Bot. 4(3) : 1023-1024. 1983.

Holotype : India : Kalimpong : Hazra 12178.

Drypetes balakrishnanii T. Chakrab. et Gang. (Euphorbiaceae).

J. Econ. Tax. Bot. 12(2) : 393. 1988.

Holotype : Burma : Bhamo division, Bhamo, 1909, GEC Cubitt 31776.

Paratype : Bangladesh : Sylhet Dist.: Malycharow Hill, 1905. M. Abu Hussain 128.

Dunbaria mamyoensis Satyan. et Thoth. (Fabaceae).

Bull. Bot. Surv. India 27(1-4) : 152-153. 1985 (1987).

Holotype : Burma : Ani Sakau, Mamyo Plateau, ± 1160 m, Oct. 15, 1911, J. H. Lace 5494.

Elaphoglossum cherapunji S. R. Ghosh et A. Biswas (Elaphoglossaceae).

J. Econ. Tax. Bot. 4(3) : 996. 1983.

Holotype : India : Meghalaya : side of Chera valley, below Cherapunji, ± 300 m, Dec. 20, 1850, Dr. J. Fayer s. n., CAL 25228.

E. jowaiense Anjali Biswas et S. R. Ghosh (Elaphoglossaceae).

J. Bombay Nat. Hist. Soc. 79(3) : 650. 1982.

Holotype : India: Jowai, 1500 m, Aug. 1892, Dr. King's Collector s. n., (CAL 25226).

Paratype : India : Jowai: Punji, 1500 m, G. Mann s. n. (CAL 25248).

Elaphoglossum prainii S. R. Ghosh et A. Biswas (Elaphoglossaceae).

J. Econ. Tax. Bot. 4(3) : 995. 1983.

Holotype : India : Nagaland : Palan Bazar, Naga Hills, Sept. 1886, D. Prain s. n. (CAL 25233).

E. simonsianum S. R. Ghosh et A. Biswas (Elaphoglossaceae).

J. Econ. Tax. Bot. 4(3) : 1001. 1983.

Holotype : India : Meghalaya : Kalapani Hills, Sept. 1850, Simons s. n. (CAL 25244).

E. thomsonii S. R. Ghosh et A. Biswas (Elaphoglossaceae).

J. Econ. Tax. Bot. 4(3) : 996. 1983.

Holotype : India : Sikkim : West of Sikkim, Nov. 11, 1857, T. Thomson s. n. (CAL 25245).

Elymus harsukhii Dubey et Dixit (Gramineae).

J. Econ. Tax. Bot. 12(1) : 227. 1988.

Holotype : India : Kashmir : on way to Amarnath, 3800 m, 14th Aug. 1983, Murugessan 65618A.

Epilobium semiamplexicaule Chowdhery et S. Singh (Onagraceae).

Indian Journ. Forestry 11(1) : 80-82. 1988.

Holotype : India : Himachal Pradesh : Dist. Kinnaur, Rakcham, K. P. Janardhanan 52679A.

E. spitianum Chowdhery et Murti (Onagraceae).

Indian Journ. Forestry 12(3) : 244. 1989.

Holotype : India : Himachal Pradesh : Spiti Dist., Gette, 4300 m, Aug. 3, 1972, U. C. Bhattacharyya 49272.

Epipogium sessanum Hegde et Rao (Orchidaceae).

J. Econ. Tax. Bot. 3(2) : 597. 1982.

Paratype : India : Arunachal Pradesh : West Kameng Dist., 1100 m, S. N. Hegde 3945.

Eragrostis santapaui K. G. Bhat. & C. R. Nagendran (Poaceae).

Reinwardtia 10. Part 2. 127. 1985.

Holotype : India : Karnataka : Coorg Dist., Mercara, growing in moist soil, Dec. 18, 1980, K. G. Bhat 794.

Eria connata Joseph, Hegde et Abbareddy (Orchidaceae).

Bull. Bot. Surv. India 24 : 114. 1982 (1983).

Holotype : India : Arunachal Pradesh : Kameng Dist., Sessa, 1100 m, Aug. 7, 1980, N. R. Abbareddy 62391A.

Eria muscicola (Lindl.) Lindl. var. **brevilinguis** Joseph et Chandrasekaran (Orchidaceae).

Bull. Bot. Surv. India 15 : 267. 1973 (1976).

Holotype : India : Kerala : Western slopes of Agastyamalai, 1500 m, ($77^{\circ}14' - 77^{\circ}15'E$, Long. and $8^{\circ}36' - 8^{\circ}37'N$, Lat.), Trivandrum Dist., Oct. 6, 1973, Joseph 44630A, BSI/SC Acc. No. 86140.

E. muscicola (Lindl.) Lindl. var. **ponmudiana** Mohanan & Henry (Orchidaceae).

J. Econ. Tax. Bot. 8(2) : 425. 1986.

Holotype : India : Kerala : Trivandrum Dist., Ponmudi, \pm 900 m, Aug. 16, 1980, M. Mohanan 69213.

Eriocaulon kasaragodense R. Ansari, V. J. Nair et N. C. Nair (Eriocaulaceae).

J. Econ. Tax. Bot. 11(1) : 235. 1987.

Holotype : India : Kerala : Kasaragod Dist., Beckal, 175 m, abundant in wet marshy places near paddy fields, June 21, 1981, R. Ansari 70984.

E. keralense R. Ansari, V. J. Nair et N. C. Nair (Eriocaulaceae).

J. Econ. Tax. Bot. 11(1) : 235. 1987.

Holotype : India : Kerala : Kasaragod Dist., Bela, 200 m, in the crevices of open base rocks, Oct. 1, 1982, R. Ansari 74420.

E. nairii M. Chandrabose et V. Chandrasekaran (Eriocaulaceae).

J. Bombay Nat. Hist. Soc. 79(1) : 165. 1982.

Holotype : India : Tamil Nadu : Konalar, Anamalai, Coimbatore Dist., Feb. 2, 1980, Chandrabose 65821.

Eskemukerjea nepalensis Malick et Sengupta (Polygonaceae).

Bull. Bot. Surv. India 11 : 433. 1969 (1971).

Holotype : Nepal : Jumla, 3000 m, Sept. 1963, Itoh & Rajbhandari 1175.

Eugenia assamica Biswas et Purkayastha (Myrtaceae).

Kew Bull. 262 : (1938) 1938.

Type : India : Assam : Passighat, June, 1936, C. S. Purkayastha 13491.

E. memecylifolia Talb. (Myrtaceae).

J. Bombay. Nat. Hist. Soc. 11 : 236. 1897.

Type : India : Karnataka : Talbot 3127.

Eulalia fimbriata (Hack.) O. Ktze. (Poaceae).

Rev. Gen. Pl. 2 : 775. 1891; J. Econ. Tax. Bot. 6(1) : 179. 1985.

Pollinia fimbriata Hack.

DC. Monogr. Phan. 6 : 164. 1889; Hook. f. Fl. Brit. Ind. 7 : 112. 1896,

Type : Peninsular India, Malabar, and Gonoan, *Hook. f. et Thomson s. n.*

Eulalia manipurensis Bor (Poaceae).

Grass. Burma, Ceylon, India & Pakistan 156. 1960; J. Econ. Tax. Bot. 6(1) : 181. 1985.

Type : India : Manipur : Palel, 720 m, *N. L. Bor* 17736.

E. mollis (Griseb.) O. Ktze. (Poaceae).

Rev. Gen. Pl. 2 : 775 (1891); J. Econ. Tax. Bot. 6(1) : 181. 1985.

Erianthus mollis Griseb.

Nachr. Ges. Wiss. Goettingen 92. 1868.

Syntype : India : Himalaya borealiocid, 1600-2000 m, *Hook. f. et Thoms.* "Erianthus No. 3".

E. phaeothrix (Hack.) O. Ktze. (Poaceae).

Rev. Gen. Pl. 2 : 775. 1891; J. Econ. Tax. Bot. 6(1) : 183. 1985.

Pollinia phaeothrix Hack.

DC. Monogr. Phan. 6 : 168. 1889; Hook. f. Fl. Brit. Ind. 7 : 112. 1896.

Syntype : Sri Lanka, *Thwaites* 959.

E. quadrinervis (Hack.) O. Ktze. (Poaceae).

Rev. Gen. Pl. 2 : 775. 1891 ; J. Econ. Tax. Bot. 6(1) : 184. 1985.

Pollinia quadrinervis Hack.

DC. Monogr. Phan. 6 : 158. 1889; Hook. f. Fl. Brit. Ind. 7 : 110. 1896.

Type : India : Sikkim : Rimchingpong, *Anderson* 1365.

E. quadrinervis (Hack.) O. Ktze. var. *hirtifolia* (Hack.) P. Sur (Poaceae).

J. Econ. Tax. Bot. 6(1) : 185. 1985.

Pollinia hirtifolia Hack.

DC. Monogr. Phan. 6 : 165. 1889 ; Hook. f. Fl. Brit. Ind. 7 : 111. 1896.

Type : India : Himachal Pradesh : Simla, *Hugal* 758.

E. hirtifolia (Hack.) A. Camus

E. speciosa (Debeaux) O. Ktze. var. *velutina* (Hack.) Bor (Poaceae).

Grass. Burma, Ceylon, India & Pakistan 157. 1960; J. Econ. Tax. Bot. 6(1): 186. 1985.

Pollinia velutina Hack.

DC. Monogr. Phan. 6 : 169. 1889; Hook. f. Fl. Brit. Ind. 7 : 113. 1896.

Isotype : India : Meghalaya : Khasia, 1600-2000 m, *Hook. f. et Thomson s. n.*

Eulalia staintonii Bor (Poaceae).

Kew Bull. 12 : 411. 1958; J. Econ. Tax. Bot. 6(1) : 186. 1985.

Type : India : Uttar Pradesh : Kumaon, 7-8000 ft., Sept. 17, 1884, *J. F. Duthie* 3554.

E. thwaitesii (Hack.) O. Ktze. (Poaceae).

Rev. Gen. Pl. 2 : 775. 1891; J. Econ. Tax. Bot. 6(1) : 187. 1985.

Pollinia thwaitesii Hack.

DC. Monogr. Phan. 6 : 163. 1882; Hook. f. Fl. Brit. Ind. 7 : 111. 1896.

Isotype : Sri Lanka: *Thwaites* C. P. 949.

E. trispicata (Schult.) Henr. var. **hookeri** P. Sur (Poaceae).

Curr. Sci. 44(10) : 362. 1975; J. Econ. Tax. Bot. 6 (1) : 190. 1985.

Holotype : Nepal : Kukhure Dol (Phul), 2300 m, Oct. 5, 1967, *Mrs Pradhan* 7472.

E. wightii (Hook. f.) Bor (Poaceae).

Grass. Burma, Ceylon, India & Pakistan 158. 1960; J. Econ. Tax. Bot. 6(1) : 190. 1985.

Isotype : India : Tamil Nadu : Pulney Hills, *Wight* 3390.

Eulaliopsis duthiei P. Sur (Poaceae).

J. Bombay Nat. Hist. Soc. 72(3) : 815. 1975.

Holotype : India : Uttar Pradesh: Ganga Valley (Tehri-Garwal), June 11, 1883, *J. F. Duthie* 14.

Eulophia hirsuta Joseph et Vajravelu (Orchidaceae).

Bull. Bot. Surv. India 17(1-4) : 192. 1975.

Holotype : India : Kerala : Palghat Dist., Karasuryamalai-Anamada area, Nemaia division, March 5, 76, Alt. 1150 m, between N. Lat. 10°30'-10°35' and E. Long. 76°45'-76°50', *Vajravelu* 46201A.

E. nicobarica Balakr. & N. G. Nair (Orchidaceae).

Bull. Bot. Surv. India 15 : 271. 1973 (1976).

Holotype : India : Car Nicobar, June 15, 1974, *N. G. Nair* 1647A.

E. ucbii Malhotra et Balodi (Orchidaceae).

Bull. Bot. Surv. India 26(1 & 2) : 92. 1984 (1985).

Holotype : India : Uttar Pradesh : Pithoragarh, Gori Valley, Garjia, April 1984, *Bipin Balodi* 75561A.

Euodia lunu-ankenda (Gaertn.) Merrill var. ***tirunelvelica*** Henry et Chandr. (Rutaceae).

Bull. Bot. Surv. India 15 : 114. 1973 (1976); J. Econ. Tax. Bot. 13(1) : 199. 1989.

Holotype : India : Tamil Nadu : Agastyamalai, Tirunelveli Dist., July 2, 1964, *Henry et Chandrabose* 19224A.

E. parkinsonii Narayanan et Nayar (Rutaceae).

J. Econ. Tax. Bot. 13(1) : 201. 1989.

Holotype : India : Andamans, 1918, *Parkinson* 922.

Euphorbia linearifolia Roth var. ***nallamalayana*** Ellis (Euphorbiaceae).

Bull. Bot. Surv. India 8 : 345. 1966.

Holotype : India : Andhra Pradesh : Chelama, Na'llamalais, Kurnool, Dist., July 6, 1963, *Ellis* 16774A.

E. santapaui Henry (Euphorbiaceae).

Bull. Bot. Surv. India 6: 329. 1964 (1965).

Holotype : India : Tamil Nadu : Agastyamalai Hills, Tirunelveli Dist., 1650 m, Aug. 27, 1963, *Henry* 17353A.

E. tavoyensis N. P. Balakrishnan (Euphorbiaceae).

J. Bombay Nat. Hist. Soc. 73(1) : 189. 1976.

Holotype : Burma : Heinye headwaters, Tavoy \pm 650 m, Nov. 25, 1921, *P. T. Russell* 2216C.

Isotype : Ibid., *P. T. Russell* 2216A, B.

Paratypes : Ridge between Kyong Pyu chaung and Talaingya, Tavoy \pm 600 m, Feb. 1, 1919, *A. T. Gage* 26 A-E.

Eurya arunachalensis Chauhan (Theaceae).

Indian Journ. Forestry 13(1) : 76. 1990.

Holotype : India : Arunachal Pradesh : Perila, 9180 ft., April 2, 1957, *G. Panigrahi* 6375A.

Evrardia asraoa Joseph et Abbareddy (Orchidaceae).

Bull. Bot. Surv. India 25(1-4) : 232-234. 1983 (1985).

Holotype : India : Meghalaya : Khasi Hills, from Pynursla, ca 1500 m, *Joseph* 73566A.

Exacum wightianum Arn. var. **uniflorum** A. N. Henry et M. S. Swaminathan (Gentianaceae).

J. Bombay Nat. Hist. Soc. 80(2) : 459. 1983.

Holotype : India : Tamil Nadu : Agastyamalai Hills, Tirunelveli Dist., May 27, 1963, *Henry* 16335.

Festuca nandadevica Hajra (Poaceae).

Indian Journ. Forestry 6(1) : 79. 1983.

Holotype : India : Uttar Pradesh : Nanda Devi National Park, Deodí Ramni, in *Rhododendron-Betula* forest, Aug. 25, 1981, *P. K. Hajra* 73285A.

Fimbristylis crystallina Govindarajalu (Cyperaceae).

Proc. Indian Acad. Sci. 79B(4) : 166. 1974.

Paratype : India : Tamil Nadu : Coimbatore, Italiyar-Anamalaiyar, Valparai, Jan. 4, 1971, *Govindarajalu* 10825A.

F. multinervia Govindarajalu (Cyperaceae).

J. Bombay Nat. Hist. Soc. 69(1) : 162. 1972.

Type : Burma : Pegu : Karen Country Hills, Jonkeyeghat, *S. Kurz* 637.

Paratypes : Burma : Pegu : Jonkeyeghat, 7 Pagodas, *S. Kurz* 628, 636; Pegu: *D. Brandis* 1018; without collector's name 2; Upper Burma : *J. C. Prazer s. n.*; India: Nagaland: Leve Kohima, 1300 m, *Dr. D. Prain* 226.

F. pustulosa Govindarajalu (Cyperaceae).

Proc. Indian Acad. Sci. (Plant Sci.) 90(5) : 444. 1981.

Isotype : India : Tamil Nadu : Coimbatore, on the way to Italiyar and Anamalaiyar, occurring in marshy places 3000-3800 ft., *Govindarajalu* 12519D.

F. rugosa Govindarajalu (Cyperaceae).

Proc. Indian Acad. Sci. 78(2) : 48. 1973.

Isotype : India : Tamil Nadu : Kolhaiyar, Oct. 6, 1964, *Govindarajalu* 7615D.

F. scabrisquama Govindarajalu (Cyperaceae).

Proc. Indian Acad. Sci. 91(1) : 43. 1982.

Isotype : India : Tamil Nadu : Venniyar to Varaiyattumottai, Highways Mts., Madurai, occurring in grass slopes, *Govindarajalu* 12110A.

F. semidisticha Govindarajalu (Cyperaceae).

Proc. Indian Acad. Sci. 78(2) : 51. 1973.

Paratype : Mysore State : Bababudan Hills, *Govindarajalu* 8869B.

Fimbristylis tortifolia Govindarajalu (Cyperaceae).

Proc. Indian Acad. Sci. (Plant Sci.) 91(1) : 46. 1982.

Isotype : India : Tamil Nadu : Madurai, Kulikad, Highways, occurring in wet open grass rocks, Dec. 12, 1973, *Govindarajalu* 12000A.

F. unispicularis Govindarajalu et Hemadri (Cyperaceae).

Proc. Indian Acad. Sci. 82B (6) : 205. 1975.

Holotype : India : Maharashtra : Durga Khilla Plateau about 30 km west of Junnar, Poona, 1100 m, Oct. 1, 1965, *Hemadri* 107528A.

Isotype : Ibid., *Hemadri* 107528E.

Folioceros palisiformis Singh (Anthocerotaceae).

Bull. Bot. Surv. India 29(1-4): 176. 1987 (1989).

Holotype : India : Arunachal Pradesh : West Kameng Dist., Prope Tipi, 3000 m, April 16, 1982, *D. K. Singh* 11671.

Friesodielsia khoshooi Vasud. et T. Chakrab. (Annonaceae).

J. Econ. Tax. Bot. 6(2) : 435. 1985.

Holotype : India : Andaman-Nicobar Islands: Great Nicobar Island, 44 km, on east-west road, April 27, 1984, *T. Chakraborty* 10193A.

Gagea bulbifera (Pall.) Salisb. (Liliaceae).

Konig & Sims. Ann. Bot. 2 : 557. 1806; J. Bombay Nat. Hist. Soc. 83(1) : 88. 1986.

Ornithogalum bulbiferum Pall.

Reise 2(2) : 736. t. Q. f. 2. 1776.

Isotype : USSR : Astrakhan: *Pallus s. n.*

G. chitralensis Dasgupta & Deb (Liliaceae).

Candollea 38(2) : 478. 1983.

Paratype : Tashkent : *Vyedensky* 53.

G. persica Boiss. (Liliaceae).

Diagn. Pl. Or. Nov. 1(7) : 108. 1846 *pro parte*; J. Bombay Nat. Hist. Soc. 83(1) : 85-87. 1986.

Type : Iran : Persepolis: *Kotschy* 237.

Galium confertum Royle ex Hook. f. var. *laceanum* Nayar et Malick (Rubiaceae).

Bull. Bot. Surv. India 11 : 429. 1969 (1971).

Holotype : N. W. Himalaya: Bashahr Rogi Cliffs, 3000 m, Aug. 2, 1890, *J. H. Lace* 469.

Garcinia affinis Wall. ex Pierre (Guttiferae).

Fl. For. Coch.-Chine. 16. t. 78C, 79G. 1883; Bull. Bot. Surv. India 6(2-4) : 122. 1964 (1965).

G. cornea Choisy

Guttif. Ind. 53. 1851.

Type : Bangladesh: Sylhet, *Wallich* 4854.

G. andamanica King var. **andamanica** (Guttiferae).

J. Asiat. Soc. Beng. 59 : 170. t. 160. 1890; Bull. Bot. Surv. India 6(2-4) : 112. 1964 (1965).

Syntypes : India : Andaman Islands, *King's Collector* 224; *Kurz s.n.*; *Helper* 872.

G. andamanica King var. **pubescens** King (Guttiferae).

J. Asiat. Soc. Beng. 59 : 170. 1890; Bull. Bot. Surv. India 6(2-4) : 112. 1964 (1965).

Type : India : Andaman Islands: Hill's rocky place, Port Blair, *King's Collector* 136.

G. anomala Planch. & Triana (Guttiferae).

Ann. Sci. nat. Ser. 4. 14 : 329. 1860; Bull. Bot. Surv. India 6(2-4) : 117. 1964 (1965).

G. monosperma Berg et Schmidt.

Offiz. Gew. 4 : t. 33 d. 1863.

Syntypes : East Bengal: *Griffith* 848; Khasia, *Hooker f. & Thomson* 14.

G. brevirostris Scheff. (Guttiferae).

Tijdschr. Nederl. Ind. 31 : 353. 1870 et Flora 53 : 241. 1870; Bull. Bot. Surv. India 6(2-4) : 120. 1964 (1965).

G. eugenifolia Wall.

(Wall. Cat. 4873, 1831, *nom. nud.*) ex T. Anderson

Type : *G. eugenifolia* : Penang, *Wallich* 4873.

G. cambogia Desr. var. **conicarpa** (Wight) T. Anderson (Guttiferae).

Fl. Brit. Ind. 1 : 262. 1874; Bull. Bot. Surv. India 6(2-4) : 129. 1964 (1965).

G. conicarpa Wight

Icon. t. 121. 1839 et Illustr. 1 : 125. 1840.

Type : Shevagherry Hills : *Herb. Wight* 142.

G. echinocarpa Thwaites var. **monticola** Mahesh. (Guttiferae).

Hook. Kew Journ. 6 : 71. 1854 et Enum. Pl. Zeyl. 1 : 49. 1858; Bull. Bot. Surv. India 6(2-4) : 126. 1964 (1965).

Type : Ceylon: Central Province, *Thwaites* 335.

Garcinia kurzii Pierre (Guttiferae).

Fl. For. Coch.-Chine. Fnum. 14. t. 78A. 1883; Bull. Bot. Surv. India 6(2-4) : 123. 1964 (1965).

G. speciosa Kurz

Type : Andamans : Kurz 24.

G. microstigma Kurz (Guttiferae).

J. Bot. 13 : 324. 1875 et For. Fl. Brit. Burma 1 : 91. 1877; Bull. Bot. Surv. India 6(2-4) : 125. 1964 (1965).

Type : S. Andaman : *S. Kurz s. n.*

G. pedunculata Roxb. (Guttiferae).

(Hort. Beng. 42. 1814, *nom. nud.* et) Fl. Ind. 2 : 265. 1832; Bull. Bot. Surv. India 6(2-4) : 119. 1964 (1965).

Neotype proposed : Assam : Goalpara, *Wallich* 4860.

G. speciosa Wall. (Guttiferae).

Pl. Asiat. Rar. 3 : 37. t. 258. 1832; Bull. Bot. Surv. India 6(2-4) : 123. 1964 (1965).

Type : Burma : Amherst, *Wallich* 4855.

G. stipulata T. Anderson (Guttiferae).

Fl. Brit. Ind. 1 : 267. 1874; Bull. Bot. Surv. India 6(2-4) : 116. 1964 (1965).

Type : Sikkim : *Hooker f. & Thomson* 17.

G. talbotii Raizada ex Santapau (Guttiferae).

Rec. Bot. Surv. India (ed. 2) 16 : 14. 1960; Bull. Bot. Surv. India 6(2-4) : 113. 1964 (1965).

G. ovalifolia var. *macrantha* T. Anderson

Fl. Brit. Ind. 1 : 269. 1874.

G. malabarica Talbot

J. Bombay Nat. Hist. Soc. 11 : 234. t. 1. 1897 et For. Fl. Bomb. Pres. 1 : 96. 1909.

Type : India : North Kanara: Gairsoppah Ghat, *W. A. Talbot* 3713.

Garnotia schmidii Hook. f. (Poaceae).

Fl. Brit. Ind. 7 : 242. 1897; Fasc. Fl. Ind. 3 : 11. 1979.

Syntype : India : Tamil Nadu : Nilgiri Hills, *Schmid s. n.*

G. scoparia Thw. (Poaceae).

Enum. Fl. Zeyl. 363. 1864; Fasc. Fl. Ind. 3 : 12. 1979.

Type : Ceylon (Sri Lanka) : *Thwaites C. P.* 943.

***Gastrochilus corymbosus* A. P. Das et Chanda (Orchidaceae).**

J. Econ. Tax. Bot. 12(2) : 401. 1988.

Holotype : India : W. Bengal : Darjeeling, Jalpahar, 2200 m, Oct. 29, 1982, A. P. Das 823.

***Gaultheria seshagiriana* Sub. et Kum. (Ericaceae).**

Bull. Bot. Surv. India 10(2) : 223. 1968 (1969).

Holotype : India : Arunachal Pradesh : Siang Frontier Division, Nov. 19, 1958, R. S. Rao 17863A.

Isotype : Ibid., R. S. Rao 17863B.

***Geniosporum tenuiflorum* (L.) Merr. var. *longiracemosum* (Ramamurthy et Sebastine) Sant. & Jain (Labiatae).**

Indian For. 92 : 643. 1966.

G. prostratum (L.) Benth. var. *longiracemosum* Ramamurthy & Sesastine.

Bull. Bot. Surv. India 6 : 325. 1964 (1965).

Holotype : India : Tamil Nadu : Kudiraimoli Teri R. F., Tiruchendur taluk, Tirunelveli Dist., Nov. 29, 1961, Sebastine & Ramamurthy 13701A.

Isotype : Ibid., Sebastine & Ramamurthy 13701B.

***Gentiana tetrasepala* Biswas (Gentianaceae).**

Hook. f. Ic. Plant. 4 : ser. 5. t. 3359. fig. 1-9. 1938 ; J. Econ. Tax. Bot. 3(3) : 1000. 1982.

Holotype : India : Kumaon, Ratam (Ralam) Valley, 13000-14000 ft., (\pm 4000-5000 m), Aug. 26, 1884, Duthie 3166.

***Geranium acotinitifolium* L'. Herit. var. *album* Ghosh et Bhattacharyya (Geraniaceae).**

J. Econ. Tax. Bot. 10(2) : 312. 1987.

Holotype : India : Himachal Pradesh : Thumla, Spiti Valley, 4200 m, Aug. 1, 1972, U. C. Bhattacharyya 49241A.

***Geum aequilobatum* Purohit et Panigrahi (Rosaceae).**

Bull. Bot. Surv. India 21 : 205. 1979 (1981).

Holotype : India : Uttar Pradesh : Tehri Garhwal, on way to Ghuttu, Jan. 8, 1972, B. D. Naithani 48206.

***Gigantochloa ligulata* Gamble (Poaceae).**

Ann. R. Bot. Gard. Calc. 7 : 67. 1896; Reinwardtia 10(Part 3) : 361. 1987.

Isotype : Malaya : Perak, J. Wray Jr. 845.

Gigantochloa scorchedinii Gamble (Poaceae).

Ann. R. Bot. Gard. Calc. 7 : 62. 1896; Reinwardtia 10. Part 3. 341. 1987.

Isotype : Malaya Peninsula: Perak, *Father Scortechini s. n.*

Globba rubromaculata J. Lal et D. M. Verma (Zingiberaceae).

Bull. Bot. Surv. India 29(1-4) : 26. 1987 (1989).

Holotype : India : Arunachal Pradesh : Ganga Lake, 450 m, Itanagar, Oct. 2, 1986. *J. Lal* 2201A.

Glochidion airyshawii Balakr. et T. Chakrab. (Euphorbiaceae).

Bull. Bot. Surv. India 25(1-4) : 220-222. 1983 (1985).

Holotype : India : Andaman & Nicobar Islands: Middle Andaman Island, Lataw near Mayabunder, ± 20 m, May 4, 1974, *Balakrishnan* 1337A.

G. bilobulatum Vasud. et T. Chakrab. (Euphorbiaceae).

J. Econ. Tax. Bot. 5(4) : 936. 1984.

Holotype : India : Andaman-Nicobar Islands : North Andaman Island, Saddle Peak Range, Lamia Bay slope, March 31, 1977, *Balakrishnan* 5426A.

G. hongkongense Muell.-Arg. var. **puberulum** T. Chakrab. et Gang. (Euphorbiaceae).

J. Econ. Tax. Bot. 13(3) : 712. 1989.

Holotype : Upper Burma, Jan. 1892, *Abdul Huk s. n.*, Acc. No. 403944.

G. insulare Hook. f. var. **brevistylum** T. Chakrab. et Gang. (Euphorbiaceae).

J. Econ. Tax. Bot. 13(3) : 715. 1989.

Holotype : Burma, Mergui, Mar. 20, 1924, *C. C. Calder* (R. N. Parker) *s. n.* Acc. No. 419320.

G. subsessile Balakr. et T. Chakrab. ssp. **birmanicum** T. Chakrab. et Gang. (Euphorbiaceae).

J. Econ. Tax. Bot. 13(3) : 716. 1989.

Holotype : Burma : Kalay Hills, July 6, 1894, *J. C. Prazer* 96, Acc. No. 403959.

Isotype : Ibid., *J. C. Prazer* 96, Acc. No. 403955.

Glycine wightii Verdcourt subsp. **wightii** Verdcourt var. **coimbatorensis** Ajita Sen (Leguminosae).

J. Bombay Nat. Hist. Soc. 74(2) : 330. 1977.

Paratypes : India : Kerala: Bolampatti, South Fuel Working Circle, Dec. 7, 1909, *C. E. C. Fischer* 1508; Kamalapores, Sept. 1910, *A. Meebold* 11285.

Gomphogyne macrocarpa Cogn. (Cucurbitaceae).

Engl. Das Pflanzenr. 4 : 40. 1916; Fasc. Fl. Ind. 11 : 52. 1982.

Neotype : India : Arunachal Pradesh: Tirap Fd., Langsang, Kothang, Deb 26151A.

Gomphostemma keralensis Vivek, Gopalan & Ansari (Labiatae).

Kew Bull. 38(2) : 189. 1983.

Holotype : Kerala : Iddiki, 950 m, Oct. 10, 1976, *Vivekananthan* 48658.

G. nayarii Chauhan (Labiatae).

Bull. Bot. Surv. India 28 : 182. 1986 (1988).

Holotype : India : Meghalaya : Khasi Hills, Nongstoin, Sept. 5, 1985, A. S. Chauhan 81993A.

Grammitis pilifera Joseph (Grammitidaceae).

J. Bombay Nat. Hist. Soc. 76(2) : 348. 1979.

Holotype : India : Kerala: From Ponmudi Hills, Trivandrum Dist., Nov. 13, 1975, 750 m, *Ravi* 5711A.

Graphina awasthii C. G. Dharne et K. N. Roychoudhury (Graphidaceae).

Bull. Bot. Surv. India 10 : 267. 1968.

Type : India: W. Bengal: Darjeeling, Sukhna Reserve Forest, 300-400 m, on the bark of trees, Jan. 4, 1967, *C. G. Dharne* 544.

G. darjeelingensis Dharne et Roychoudhury (Graphidaceae).

Bull. Bot. Surv. India 10 : 267. 1968 (1969).

Type : India : W. Bengal: Darjeeling, Manibhanjan, on the bark of trees, 2500 m, Dec. 30, 1966, *C. G. Dharne* 221.

Grona filicaulis Kurz (Leguminosae).

Bull. Bot. Surv. India 17(1-4) : 205. 1975.

Holotype : Pegu, Dec. 29, 1870, *S. Kurz* 2549.

Gueldenstaedtia santapaui Thoth. (Leguminosae).

Bull. Bot. Surv. India 10 : 391. 1968.

Holotype : Bhutan : Chin-la-Pass, Paro, 4200 m, Aug. 20, 1963, *Balakrishnan & Sengupta* 619.

Gymnema lushaiense Rahman et Wilcock (Asclepiadaceae).

J. Econ. Tax. Bot. 13(1) : 181. 1989.

Type : India: Lushai Hills, Prope Lunglah, 3-4000 ft., *A. T. Gage* 74.

Gymnosporia bailadillana Narayanaswami et Mooney (Celastraceae).

J. Indian Bot. Soc. 20(4) : 191. 1941.

Type : India : Bailadilla Hill, Bastar State, H. Mooney 890.

Habrodon kashmiriensis Vohra (Fabroniaceae).

Bull. Bot. Surv. India 11 : 70. 1969 (1971).

Holotype : India : Kashmir: Tangmarg, N. W. Himalaya, 2000 m, Sept. 9, 1961, Vohra et Wadhwa 567.

Haplocladium microphyllum (Hedw.) Broth. ssp. **capillatum** (Mitt.) Reim.

var. **bhutanicum** Vohra (Thuidiaceae).

Bull. Bot. Surv. India 20(1-4) : 152. 1978 (1979).

Holotype : Bhutan : Dochula, 3050 m, on tree trunk, Sept. 22, 1964, Vohra 2003.

Paratype : Bhutan : Aug.-Sept. 1963, Wadhwa 185.

Harrisonia perforata (Blanco) Meir. (Simarubaceae).

Philip. J. Sci. Bot. 7 : 236. 1912; Fasc. Fl. Ind. 4 : 12. 1980.

Type : Philippines : Mandaloyon, Rizal Prov., Luzon, April 20, 1914, Merrill Species Blancoanae 433.

Hedychium calcaratum A. S. Rao et D. M. Verma (Zingiberaceae).

Bull. Bot. Surv. India 11 : 120. 1969 (1971).

Holotype : India : Meghalaya ; Jowai Bodorpur Road, 96 km from Shillong, Khasia & Jaintea Hills, July 22, 1957, G. K. Deka 10134A.

H. dekianum A. S. Rao et D. M. Verma (Zingiberaceae).

Bull. Bot. Surv. India 11 : 122. 1969 (1971).

Holotype : India : Meghalaya; Khlie Hriat Road from Jowai, Khasia & Jaintea Hills, July 15, 1966, G. K. Deka 35605A.

H. gracillimum A. S. Rao et D. M. Verma (Zingiberaceae).

Bull. Bot. Surv. India 11 : 124. 1969 (1971).

Holotype : India : Meghalaya: Experimental Garden, Botanical Survey of India, Woodlands, Shillong, Khasia & Jaintea Hills, July 1, 1967, Introduced from Pynursla, Khasia & Jaintea Hills, D. M. Verma 35650.

H. longipedunculatum A. R. K. Sastry et D. M. Verma (Zingiberaceae).

J. Bombay Nat. Hist. Soc. 65(2) : 293. 1968.

Holotype : India : Arunachal Pradesh : Subansiri Dist., Amjee, 1220 m, May 22, 1966, A. R. K. Sastry 45509A.

Paratypes : India : Naga Hills: July 1844, Anon. s. n., Acc. No. CAL-466858, in Flower; Konoma Hill, 2500 m, May 19, 1895, G. Watt 11609, in flower.

Hedychium radiatum A. S. Rao et P. K. Hajra (Zingiberaceae).

Bull. Bot. Surv. India 16 : 163. 1974 (1977).

Holotype : India : Arunachal Pradesh : Nyertsephu-Wana Camp Road, Kameng Dist., *A. S. Rao* 56655A.

H. robustum A. S. Rao et P. K. Hajra (Zingiberaceae).

Bull. Bot. Surv. India 16 : 166. 1974 (1977).

Holotype : India : Arunachal Pradesh : Jamiri, Kameng Dist., *A. S. Rao* 56654A.

H. rubrum A. S. Rao et D. M. Verma (Zingiberaceae).

Bull. Bot. Surv. India 11 : 126. 1969 (1971).

Holotype : India : Meghalaya: Experimental Garden, Botanical Survey of India, Woodlands, Shillong, Khasia & Jaintia Hills, Aug. 12, 1965, *D. M. Verma* 34599A.

Hedyotis bourdillonii (Gamble) Rolla Rao et Hemadri (Rubiaceae).

Indian For. 99 : 378. 1973; J. Bombay Nat. Hist. Soc. 83(2) : 466. 1986.

Oldenlandia bourdillonii Gamble

Kew Bull. 1919 : 404. 1919.

H. silent-valleyensis Vajravelu et al.

J. Bombay Nat. Hist. Soc. 80(2) : 402. 1983 (1984).

Holotype : *H. silent-valleyensis* : India : Kerala : Palghat Dist., Silent Valley, Kothipuzha dam-site, May 27, 1966, *Vajravelu* 27674.

H. devicolamensis D. B. Deb et Ratna Dutta (Rubiaceae).

J. Bombay Nat. Hist. Soc. 82(3) : 619. 1985.

Holotype : India : Kerala : Kottayam Dist., Devicolam, Sept. 11, 1968. *D. B. Deb* 30480.

Isotype : *D. B. Deb* 30481.

Paratypes : Umaiya Malei, 2025 m, Devicolam, April 17, 1966, *B. V. Shetty* 27311; Rajamallai, 1950 m, Feb. 2, 1970, *B. V. Shetty* 31783; Devicolam, Jan. 25, 1964, *K. M. Sebastian* 18458.

H. erecta Manilal et Sivarajan (Rubiaceae).

Notiser, 129(2) : 191. 1976.

Isotype : Thiruvangad, 50 MSI, Sept. 8, 70, *Sivarajan* 491.

H. santapaui Shetty et Vivek. (Rubiaceae).

Bull. Bot. Surv. India 11 : 447. 1969 (1971).

Holotype : India : Kerala: Umaiyamalai, Devicolam, Kottayam Dist., Nov. 17, 1965, *Shetty* and *Vivekananthan* 26471A.

***Hedyotis tavoyensis* Balakr. (Rubiaceae).**

J. Bombay Nat. Hist. Soc. 71(1) : 168. 1974.

Holotype : Burma : Tavoy: Talaingya Chaung, 900 m, Feb. 7, 1919,
A. T. Gage 68A.

Isotype : Ibid., A. T. Gage 68B.

***Hemiadelphis polyspermus* (Roxb.) Nees var. *joshianus* Rao et Biswas
(Acanthaceae).**

Indian For. 94(9) : 657-658. 1968.

Holotype : India : Uttar Pradesh : Chandmari, Varanasi, Varanasi Dist., 85 m, S. P. Singh 2558A.

***Heritiera kanikensis* Majumdar et Banerjee (Sterculiaceae).**

Bull. Bot. Surv. India 27(1-4) : 150-151. 1985 (1987).

Holotype : India : Orissa : Bhitarkanika, Aug. 7, 1973., L. K. Banerjee 10295A.

Isotypes : Ibid., 10295B-C.

***Hibiscus beddomei* Rakshit et Kundu (Malvaceae).**

Sci. & Cult. 27 : 192. 1961; Fasc. Fl. Ind. 19 : 126. 1988.

Holotype : India : Without precise locality, Beddome 92.

***H. fragrans* Roxb. (Malvaceae).**

Roxb. [Hort. Beng. 97. 1814. *nom. nud.*] Fl. Ind. ed. Carey 3: 195. 1832 ; Fasc. Fl. Ind. 19 : 13. 1988.

Neotype : Bangladesh : Sylhet, Wall. Cat. No. 1911A.

***H. hoshiarpurensis* T. K. Paul et Nayar (Malvaceae).**

Bull. Bot. Surv. India 25(1-4) : 188-189. 1983 (1985); Fasc. Fl. Ind. 19 : 131. 1988.

Isotype : India : Punjab : Dholbah (Hoshiarpur Dist.), Sept. 22, 1970, O. P. Misra 41888B.

***H. scandens* Roxb. (Malvaceae).**

Roxb. [Hort. Beng. 51. 1814. *nom. nud.*] Fl. Ind. ed. Carey 3 : 200. 1832 ; Fasc. Fl. Ind. 19 : 150. 1988.

Neotype : H. B. C., Wall. Cat. No. 1910.

***H. surattensis* L. f. forma *immaculata* Kurz ex Rakshit (Malvaceae).**

Sci. & Cult. 87(4) : 194. 1961.

Type : Burma : Pegu : Yomah, Kurz 1248.

Hibiscus talbotii (Rakshit) T. K. Paul et Nayar (Malvaceae).
 Bull. Bot. Surv. India 22 : 197 (1980) 1982; Fasc. Fl. Ind. 19 : 156. 1988.
H. hirtus var. *talbotii* Rakshit
 Sci. & Cult. 27(4) : 193. 1961.
Holotype : India : North Canara : *W. A. Talbot* 68.

Hiptage thothathrii Balakr. et Srivastava (Malpighiaceae).
 J. Econ. Tax. Bot. 4(3) : 985. 1983.
Holotype : India : N. Andamans : Saddle Peak, 600-700 m above
 m. s. l., *Balakrishnan* 5377.

Holboellia khasiana T. K. Paul et Nayar (Lardizabalaceae).
 Bull. Bot. Surv. India 27(1-4) : 105. 1985 (1987).
Holotype : India : Meghalaya : Khasia, Cherra, 5000 ft., (1520 m),
Griffith s. n., Acc. No. 16830.

H. latifolia Wall. var. *Iatifolia* (Lardizabalaceae).
 Tent. Fl. Nep. 24. t. 16. 1824; Fasc. Fl. Ind. 19 : 32. 1988.
Type : Nepal Himalaya : Wall. Cat. No. 4950.
 var. *angustifolia* (Wall.) Hook. f. & Thoms.
 Fl. Brit. Ind. 1 : 108. 1872.
H. angustifolia Wall.
 Tent. Fl. Nep. 24. t. 17. 1824.
Type : Nepal : Wall. Cat. No. 4951.

Hoya kanyakumariana Henry et Swaminathan (Asclepiadaceae).
 J. Bombay Nat. Hist. Soc. 75(2) : 462. 1978.
Holotype : India : Tamil Nadu : Vallachithodu, Lower Kodayar,
 Konyakumari, 600 m, Sept. 6, 1976, *Henry* 48243A.

H. manipurensis Deb (Asclepiadaceae).
 J. Indian Bot. Soc. 34(1) : 50. 1955.
Holotype : India : Manipur : Litan, 3000 ft., Sept. 1, 1953, epiphytic
 on *Ficus glomerata* Roxb., *Deb* 1081.

Hugonia belli Sedgwick (Linaceae).
 Indian For. 46 : 424. 1920; Fasc. Fl. Ind. 13 : 4. 1983.
Type : Solepoli, Coorg, *Sedgwick* & *Bell* 5457.

Humboldtia unijuga Bedd. var. *trijuga* Joseph et Chandrasekaran
 (Caesalpiniaceae).
 J. Bombay Nat. Hist. Soc. 81(3) : 729. 1984.

Holotype : India : Kerala: Koviltherimalai, near Bonaccord Estate, Trivandrum Dist., Oct. 2, 1973, *Joseph* 44511.

Huodendron biaristatum (W. W. Sm.) Rehder (Styracaceae).

J. Arn. Arb. 16 : 345. 1935.

Styrax biaristatus W. W. Smith

Notes Roy bot. Gard. Edinburgh 12 : 233. 1920.

Iso-syntype : China : Yunnan, Nov., 1919. *G. Forrest* 18833.

Hydnocarpus kurzii (King) Warb. ssp. *australis* Sleum. forma *latifolia*

N. Mukherjee (Flacourtiaceae).

J. Bombay Nat. Hist. Soc. 69(2) : 394. 1972.

Holotype : Burma : Pegu, *Kurz* 532.

Hydrogonium mussoorianum Vohra (Pottiaceae).

J. Bombay Nat. Hist. Soc. 63(2) : 464. 1966.

Holotype : India : N. W. Himalaya : Mussoorie, 2000 m, March 22, 1963, *Vohra* 453A.

Hygrohypnum choprae Vohra (Amblystegiaceae).

Bull. Bot. Surv. India 22(1-4) : 117. 1980 (1982).

Isotype : India : W. Bengal : Darjeeling, 2400 m, July 1935, *Chopra* 115 (Present under the name *Hygrohypnum angustirete* Dix. mss.).

H. nairii Vohra (Amblystegiaceae).

Bull. Bot. Surv. India 22(1-4) : 117. 1980 (1982).

Holotype : India : Uttar Pradesh : Kumaon, Pithoragarh, Kalamuni Pass, 3200 m, March 30, 1966, *Nair* M 12.

Hymenandra narayanaswamii Nayar et Giri (Myrsinaceae).

J. Bombay Nat. Hist. Soc. 72(3) : 819. 1975.

Holotype : Burma : Tavoy : *P. T. Russell* 2105.

Hypecoum leptocarpum Hook. f. et Thoms. (Papaveraceae).

Fl. Ind. 1 : 276. 1855 et Hook. f. Fl. Brit. Ind. 1 : 120. 1872; Fasc. Fl. Ind. 17 : 43. 1984.

Type : Tibet : Temp. region, 3050-3660 m, *T. Thomson* s. n.

Hypericum assamicum S. N. Biswas (Hypericaceae).

Webbia 25(2) : 671. 1971.

Holotype and Isotype : India : Assam : Nowgong, March 1848, *Simons* s. n.

***Hypericum bengalensis* S. N. Biswas (Hypericaceae).**

Bull. Bot. Surv. India 29(1-4) : 53. 1987 (1989).

Holotype : India : W. Bengal : Kalimpong, 1060 m, June 2, 1959, *S. K. Mukerjee* 5094.

Isotypes : Ibid., *S. K. Mukerjee* 5094A and 5094B.

***H. burmanicum* S. N. Biswas (Hypericaceae).**

Bull. Bot. Surv. India 29(1-4) : 54. 1987 (1989).

Holotype : Burma : Shan Hill, Fort Stedman, alt. 1215 m, Aug. 1892,

H. H. Hilldebrand s. n.

Paratype : Burma : Shan States, an erect shrubs, 5-6 ft., with yellow fls., March 5, 1932, *A. Long* 10374.

***H. griffithii* Hook. f. et Thoms. ex Dyer (Hypericaceae).**

Fl. Brit. Ind. 1 : 253. 1874. emend. ; Bull. Bot. Surv. India 13 : 161. 1971 (1973).

Isotype : Bhutan : 1861-62, *Griffith* 833.

***H. hookerianum* Wt. et Arn. var. *dentatum* S. N. Biswas (Hypericaceae).**

Bull. Bot. Surv. India 25(1-4) : 195-196. 1983 (1985).

Holotype : India : Tamil Nadu : Perumalmalai, Kodaikanal, ±2000 m, July 27, 1965, *K. M. Sebastian* 24992.

Paratype : Sikkim Himalaya : Below Changu, 3040 m, Sept. 22, 1926, *J. C. Dawa* 167.

***H. humifusum* L. ssp. *suborbiculatum* S. N. Biswas (Hypericaceae).**

Bull. Bot. Surv. India 29(1-4) : 53. 1987. (1989).

Holotype : Nepal, Near Jagat, alt. 8500 ft., by small stream in grass, flowers yellow, July 5, 1954, *Stainton, Sykes & Williams* 3374.

Paratype : India : W. Bengal : Dist. Darjeeling, *S. K. Mukerjee & party* 645.

***Hypocybe boriviliensis* Sharma, Sandhya Deshpande & Pradhan (Ord. Agaricales).**

Bull. Bot. Surv. India 28 : 171. 1986 (1988).

Holotype : India : Bombay, Borivili, National Park, Aug. 22. 1985, *Sharma* 169123 (Cryptogamic Unit).

***Hypoestes andamanensis* Thoth. (Acanthaceae).**

Reinwardtia 7 : 1-3. 1965.

Holotype : India : North Andamans : From forests of Austin II, 30 m, Feb. 1, 1959, *Thothathri* 9157A.

Isotypes : Ibid., *Thothathri* 9157B-H.

Hypoestes thothathrii Vasud. & T. Chakrab. (Acanthaceae).

J. Econ. Tax. Bot. 5(4) : 989. 1984.

Holotype : India : Andaman-Nicobar Islands: South Andamans, Rutland Island, Barakhari (N-S Road, 14 km onwards), Jan. 21, 1982, *Vasudeva Rao* 8601A.

H. triflora R. & S. var. **pedunculata** Balakr. (Acanthaceae).

Bull. Bot. Surv. India 16 : 172. 1974 (1977).

Holotype : India : Meghalaya : Raliang, Jowai Dist., Nov. 24, 1969, shaded moist places in subtropical forests, 1000-1500 m, *Balakrishnan* 50252A.

Ilex longicaudata Comber (Aquifoliaceae).

Bull. Bot. Surv. India 17(1-4) : 199. 1975.

Isotype : China : Yunnan : Shweli-Salween divide, Lat. 25°20' N, 3300 m, July 1917, *G. Forrest* 15657 (Male).

Illigera gammiei Nayar et Giri (Hernandiaceae).

Bull. Bot. Surv. India 25(1-4) : 249-251. 1983 (1985).

Holotype : India : Assam : Dihing River, March 19, 1894, *G. A. Gammie* 142A.

Isotypes : Ibid., *Gammie* 142B-E.

Impatiens chandrasekharanii M. Chandrabose (Balsaminaceae).

J. Bombay Nat. Hist. Soc. 75(3) : 901. 1978.

Holotype : India : Tamil Nadu : Akkamalai, Anamalai in Coimbatore Dist., July 23, 1978, *Chandrabose* 57269A.

I. konalarensis M. Chandrabose, V. Chandrasekaran and N. C. Nair (Balsaminaceae).

J. Bombay Nat. Hist. Soc. 81(3) : 676. 1984.

Holotype : India : Tamil Nadu : Coimbatore Dist., Konalar, Anamalai, Nov. 18, 1980, *Chandrabose* 69013.

Indigofera bella Prain (Fabaceae).

Reinwardtia 10. Part 2. 216. 1985.

Lectotype selected : Upper Burma : Kalay Hills, Sept. 9, 1894, *J. C. Prazer s. n.*

I. bracteata Grah. ex Baker var. **khasiana** Sanjappa (Fabaceae).

J. Bombay Nat. Hist. Soc. 81(2) : 449. 1984.

Holotype and Isotype : India : Meghalaya : Khasia Hills, 5-6000 ft., *Hooker f. & Thomson s. n.*

***Indigofera chitralensis* Sanjappa (Fabaceae).**

Bull. Bot. Surv. India 28 : 228. 1986 (1988).

Holotype : Pakistan : Chitral : June 20, 1985. Western Himalayas, S. A. Harris 16069.

I. *gangetica* Sanjappa (Fabaceae).

Bull. Bot. Surv. India 28 : 225. 1986 (1988).

Holotype and Isotype : India : Uttar Pradesh : Kumaon, Wallich 5480 ex parte.

I. *maymyoensis* Sanjappa (Fabaceae).

Reinwardtia 10. Part 2. 227. 1985.

Holotype : Burma : Maymyo plateau, 3500 ft., July, 1909, Lace 4846.

Paratypes : Nov. 5, 1911, Lace 5519 & June 21, 1913, Lace 6227.

I. *megalayensis* Sanjappa (Fabaceae).

Reinwardtia 10. Part 2. 227. 1985.

Paratypes : Burma : Mt. Victoria, Cat. 8000 ft., Chhin Hills, 1939, Dikason 8534; unknown locality, Chhin Hills, 1985, Dunn 82.

I. *minbuensis* Gage (Fabaceae).

Rec. Bot. Surv. India 3 : 40. 1904; Reinwardtia 10. Part 2. 229. 1985.

Lectotype : Burma : Chhin Hills, Minbu Dist., 4000 ft., Jan. 1903, S. Mokim 1158A.

I. *nigrescens* Kurz ex King & Prain (Fabaceae).

J. Asiatic Soc. Beng. 67 : 286. 1898; Reinwardtia 10. Part 2. 231. 1985.

Lectotype : Upper Burma : Maymyo, Oct. 1888, B. Khan 203.

I. *scabrida* Dunn. (Fabaceae).

J. Linn. Soc. 35 : 487. 1903; Reinwardtia 10. Part 2. 233. 1985.

Lectotype : China : Yunnan : 4600-6000 ft., A. Henry 9686 (Photo & Type number).

I. *sesquipedalis* C. B. Clarke ex Sanjappa (Fabaceae).

J. Bombay Nat. Hist. Soc. 81(2) : 445. 1984.

Holotype : India : Meghalaya : Shillong, 5000 ft., June 1, 1885, C. B. Clarke 38232.

I. *squalida* Prain (Fabaceae).

J. Asiatic Soc. Beng. 66 : 355. 1897 et Ann. Roy. Bot. Gard. Cal. 9 : 21. t. 27. 1901; Reinwardtia 10. Part 2. 235. 1985.

Lectotype selected : Upper Burma : Fort Stedmann, June 5, 1893, King's Collector 442.

***Indigofera thothathrii* Sanjappa (Fabaceae).**

J. Jap. Bot. 59(9) : 272. 1984.

Isotype : India : Uttar Pradesh : Bharaich Dist., Nishangara, near Forest Rest House, Sept. 11, 1956, *H. Pirson* 1412.

***I. tirunelvelica* Sanjappa (Fabaceae).**

J. Bombay Nat. Hist. Soc. 79(3) : 647. 1982.

Holotype : India : Tamil Nadu : Tirunelveli Dist., along roadside between Kalakkad and Sengalteri, 300 m, Dec. 18, 1980, *M. Sanjappa* 717A.

Isotypes : Ibid., *M. Sanjappa* 717B-F.

***I. trita* L. f. var. *marginulata* (Grah. ex Wt. & Arn.) Sanjappa (Fabaceae).**

Bull. Bot. Surv. India 26 : 118. 1984 (1985).

I. marginulata Grah. ex Wt. & Arn.

Type : India : Penen : Wall. Cat. No. 5467.

***Indofeveillea khasiana* Chatterjee (Cucurbitaceae).**

Kew Bull. 1947 : 119; Fasc. Fl. Ind. 11 : 62. 1982.

Type : India : Khasia, *G. Mann* s. n.

***Indotristicha tirunelveliana* Sharma, Karthik. et Shetty (Podostemaceae).**

Bull. Bot. Surv. India 16 : 157. 1974 (1977).

Holotype : India : Tamil Nadu : Thulukkanparai, Tirukkurungudi, Tirunelveli Dist., 850 m, Feb. 17, 1972, *Karthikeyan* 40159A.

***Iphigenia magnifica* Ansari et Rolla Rao (Liliaceae).**

Bull. Bot. Surv. India 20 : 162. 1978 (1979).

Holotype : India : Maharashtra : Dhule in Dhule Dist., Oct. 5, 1969, *Pataskar* 118218A.

Isotype : Ibid., *Pataskar* 118218D.

***I. sahyadrica* Ansari et Rolla Rao (Liliaceae).**

Bull. Bot. Surv. India 20 : 163. 1978 (1979).

Holotype : India : Karnataka : Hulical in Shimoga Dist., Aug. 25, 1963, *Raghavan* 90209A.

***Isachne albens* Trin. (Poaceae).**

Sp. Gram. 1. t. 85. 1828; Fasc. Fl. Ind. 14 : 13. 1984.

Type : Nepal : Sanko : Aug. 1821, *Wallich* s. n., Wall. Num. List No. 8658.

***Isachne angladei* Fischer (Poaceae).**

Kew Bull. 1932 : 323; Fasc. Fl. Ind. 14 : 13. 1984.

Paratypes : India : Palni Hills, Shembagnur, ± 1650 m, *Vannalderen* 1283; Anaimalai, 1200-1650 m, May 1917, *Blatter & Hallberg* 182.

***I. borii* Hemadri (Poaceae).**

Ind. For. 97 : 223-225. 1971; Fasc. Fl. Ind. 14 : 14. 1984.

Holotype : India : Maharashtra: Pune Dist., Dhak Khilla, ± 27 km west of Junar, Sept. 22, 1968, *Hemadri* 117968A.

Isotype : Ibid., *Hemadri* 117968E.

***I. bourneorum* Fischer (Poaceae).**

Kew Bull. 1932 : 324; Fasc. Fl. Ind. 14 : 15. 1984.

Isotype : India : Palni Hills; Kodaikanal, ± 1950 m, June 14, 1898, *A. G. & Lady Bourne* 2491.

***I. dimyloides* Bor (Poaceae).**

Kew Bull. 1949: 96; Fasc. Fl. Ind. 14 : 18. 1984.

Isotype : India : Sikkim Tarai, Dulkajhar, ± 1500 m, Oct. 16, 1884, *C. B. Clarke* 36764.

***I. gracilis* C. E. Hubb. (Poaceae).**

Kew Bull. 1927 : 77; Fasc. Fl. Ind. 14 : 25. 1984.

Isotype : India : Karnataka; Bababudan Hills, Santaveri, ± 1200 m, Oct. 1908, *Meebold* 10781.

***I. himalaica* Hook. f. (Poaceae).**

Fl. Brit. Ind. 7 : 23. 1897; Fasc. Fl. Ind. 14 : 27. 1984.

Syntype : India : Meghalaya; Khasi Hills, ± 1200-1500 m, *J. D. H. & T. T.*, s. n.

***I. meeboldii* Fischer (Poaceae).**

Kew Bull. 1932 : 323; Fasc. Fl. Ind. 14 : 31. 1984.

Isotype : India : Karnataka; Shimoga, ± 600-900 m, Oct. 1908, *A. Meebold* 10747.

Paratypes : *A. Meebold* 10746; Kumsi, ± 600-900 m, Oct. 1908, *A. Meebold* 10745.

***I. mysorensis* Raghavan (Poaceae).**

Indian For. 97 : 304-307. 1971; Fasc. Fl. Ind. 14 : 33. 1984.

Holotype : India : Karnataka : Shimoga Dist., Kundaguda near Agumbe, Aug. 19, 1963, *Sundararaghavan* 90008A.

***Isachne scabrosa* Hook. f. (Poaceae).**

Fl. Brit. Ind. 7 : 23. 1897; Fasc. Fl. Ind. 14 : 34. 1984.

Type : India : Khasi Hills, 4500-5000 ft., C. B. Clarke s. n.

***I. swaminathanii* Ved Prakash et Jain (Poaceae).**

Proc. Indian Acad. Sci. (Plant Sci.) 92 : 19-22. 1983; Fasc. Fl. Ind. 14 : 36. 1984.

Holotype : India : Maharashtra : Satara Dist., Mahabaleshwar, 1372 m, Oct. 9, 1979, *Ved Prakash* 337A.

Isotypes : *Ved Prakash* 337B-E.

***I. veldkampii* Bhatt. & Nagendran (Poaceae).**

Curr. Sci. 52 : 258-259. 1983; Fasc. Fl. Ind. 14 : 36. 1984.

Holotype : India : Karnataka : South Kanara Dist., Manipal, 150 m, Oct. 29, 1977, *K. G. Bhatt* 468A.

***Ischaemum calicutensis* Sreekumar, V. J. Nair et N. C. Nair (Poaceae).**

J. Econ. Tax. Bot. 4(3) : 1007. 1983.

Holotype : India : Kerala : Calicut Dist., Pokkunnamalai, near Nanminda, ± 850m, Oct. 29, 1981, *P. V. Sreekumar* 71803.

***I. kannanorensis* Sreekumar, V. J. Nair & N. C. Nair (Poaceae).**

J. Econ. Tax. Bot. 4(3) : 1009. 1983.

Holotype : India : Kerala : Cannanore Dist., Paramba, on the way to Bendukka, ± 150 m, Oct. 16, 1981, *P. V. Sreekumar* 71713.

***I. nairii* V. J. Nair et P. V. Sreekumar (Poaceae).**

J. Econ. Tax. Bot. 5(5) : 1205. 1984.

Holotype : India : Kerala : Calicut Dist., Kanjeera Kadavu, near Kakkayam, ± 900 m, Nov. 26, 1981, *P. V. Sreekumar* 71843.

***I. raizadae* Hemadri et Billiore (Poaceae).**

Indian For. 96(4) : 318. 1970.

Holotype : India : Maharashtra : Sadryaghat, near Harishchandragarh, Thana Dist., Nov. 16, 1968, growing on rock and rocky crevices along the stream, *K. V. Billiore* 115450A.

Paratypes : *K. V. Billiore* 115984R-S.

***I. tadulingamii* N. C. Nair & P. V. Sreekumar (Poaceae).**

Blumea 30(2) : 385-387. 1985.

Type : India : Kerala : Idduki Dist., Eravikulam National Park, 2200 m, April 7, 1980, *Sreekumar* 71863.

Ixonanthes khasiana Hook. f. (Ixonanthaceae).

Fl. Brit. Ind. 1 : 416. 1874; Fasc. Fl. Ind. 13 : 14. 1983.

Type : India : Khasia Mts., *F. de Silva*.

Janakia arayalpathra Joseph & Chandrasekaran (Periplocaceae).

J. Indian Bot. Soc. 57(4) : 308. 1978.

Holotype : India : Kerala : Trivandrum, Kursumalai *ca* 875 m, near Bonacord Estate ($77^{\circ}10' - 77^{\circ}11'$ E Long. and $8^{\circ}41'$ N Lat.), Aug. 23, 1975, Joseph 46503A.

Jasminum andamanicum Balakr. et N. G. Nair (Oleaceae).

Bull. Bot. Surv. India 21 : 214. 1979 (1981).

Holotype : India : S. Andaman : North Bay Hill jungle, $92^{\circ}45'$ E & $11^{\circ}43'$ N, Jan. 5, 1894, *Dr. King's Collector s. n.*, Acc. No. 286074.

Isotypes : Acc. Nos. 286073, 286075, 286076.

J. cordifolium Wall. ex G. Don ssp. **andamanicum** Srivastava & Kapoor (Oleaceae).

Bull. Bot. Surv. India 25(1-4) : 217-219. 1983 (1985).

Paratype : South Andaman : Chiriatapu, Jan. 20, 1916, C. E. Parkinson 891.

J. multiflorum (Burm. f.) Andr. var. **nicobaricum** Thoth. (Oleaceae).

Bull. Bot. Surv. India 5 : 99. 1963 (1965).

Holotype : Car Nicobar Island : March 30, 1959, *Thothathri* 9345A.

Isotypes : Ibid., *Thothathri* 9345 B-F.

J. simonsii Sinha, Maiti et Giri (Oleaceae).

J. Bombay Nat. Hist. Soc. 77(2) : 305. 1980.

Holotype : Without any precise locality, *Simons s. n.* [*J. attenuatum* Roxb. ex DC., C. E. C. Fischer, Aug. 12, 1936 (Acc. No. 285972)].

Isotype : Ibid., Acc. No. 285972A.

J. unifoliolatum Balakr. et N. G. Nair (Oleaceae).

Bull. Bot. Surv. India 24 : 33. 1982 (1983).

Holotype : India : North Andaman : Saddle Peak, 400-700 m, scrub forests, on red lateritic soil, Dec. 1, 1976, in flower, *Balakrishnan et Nair* 4766A.

Paratype : Ibid., March 27, 1977, in fruits, *Balakrishnan* 5160.

Jatropha maheshwarii Subr. et Nayar (Euphorbiaceae).

Bull. Bot. Surv. India 6 : 331. 1964 (1965).

Holotype : India : Tamil Nadu : Kanyakumari, at sea level, April 5, 1956, *Nayar* 107A.

Isotypes : *Ibid.*, *Nayar* 107B-I.

Jatropha villosa Wight var. **ramnadensis** Ramamurthy (Euphorbiaceae).

Bull. Bot. Surv. India 9 : 278. 1967 (1968).

Holotype : India: Tamil Nadu : In Melamadam forest, Tirupathur Taluk, Ramnad Dist., Aug. 15, 1964, *Ramamurthy* 20937A.

Paratype : *Ibid.*, Sept. 15, 1965, *Ramamurthy* 25824A.

Juncus rohtangensis Goel et Aswal (Juncaceae).

Indian Journ. Forestry 10(4) : 262-263. 1987.

Isotype : India : Himachal Pradesh: Lahul-Spiti, Rohtang Pass, 4000 m, July 27, 1979, *B. S. Aswal* 10554E.

Justicia gingiana Seb. et Ramamurthy (Acanthaceae).

Bull. Bot. Surv. India 6 : 99. 1964 (1965).

Holotype : India : Tamil Nadu : In Gingee Reserve Forest, South Arcot Dist., March 22, 1961, *Sebastine et Ramamurthy* 12360A.

Kalanchoe cherukondensis Subba Rao & Kumari (Crassulaceae).

Bull. Bot. Surv. India 17 : 177-178. 1975 (1978).

Holotype : India : Tamil Nadu : Coimbatore, Garden of Botanical Survey of India (from cultivated materials raised from wild plants collected in vegetative condition, growing in rocky areas on Cherukonda at, 1100 m, in Visakhapatnam Dist., Andhra Pradesh, 17th April, 1974, *Subba Rao et Kumari* 39300.

Kanjarum palghatense Ramamurthy (Acanthaceae).

Bull. Bot. Surv. India 13 : 153. 1971 (1973).

Holotype : India : Kerala : Palghat, Kanjarumpuzha Dist., Oct. 27, 1964, *Sebastine* 22312A.

Isotype : *Ibid.*, *Sebastine* 22312B.

Katherinea navicularis Balakr. & Chowdhury (Orchidaceae).

Bull. Bot. Surv. India 8 : 316. 1966 (1967).

Holotype : India: Near Narfong, on the road to Tashigang, ($91^{\circ}26'$ E & $27^{\circ}7'N$) 1560 m, March 29, 1965, collected in vegetative condition, flowered, under cultivation at Shillong on Sept., 24, 1965, *Balakrishnan* 43014A.

Knoxia plantaginea Wall. (Rubiaceae).

Pl. Asiat. Rar. 1(1830) 271; J. Econ. Tax. Bot. 6(1): 81. 1985.

Lectotype : Wall. Cat. No. 818.

K. platycarpa Arn. var. **foliosa** Thw. (Rubiaceae).

Enum. (1864) 152; J. Econ. Tax. Bot. 6(1) : 81. 1985.

Syntype : Sri Lanka, C. P., 1370-2134 m, *Thwaites* C. P. 51.

K. roxburghii (Spreng.) Rau ssp. **brunonis** (Hook. f.) Bhattacharjee et Deb var. **brunonis** (Rubiaceae).

J. Econ. Tax. Bot. 6(1) : 85. 1985.

Isotype : Kumaon, *Blinkworth* s. n. in Wall. Cat. No. 822.

Spermacoce brunonis Blinkworth Wall. Cat. No. 822. *nom. nud.*

K. microcarpa Kurz

J. Asiat. Soc. Ser. 2. 46. 1877. 138.

Syntypes : Burma : Pegu, Dec. 27, 1870, *Kurz* 3083; Irawaddy, Sittang Valley, *Kurz* 1433.

K. roxburghii (Spreng.) Rau ssp. **brunonis** (Hook. f.) Bhattacharjee et Deb var. **glaucia** (W. W. Sm.) Bhattacharjee et Deb (Rubiaceae).

J. Econ. Tax. Bot. 6 (1) : 85. 1985.

Hedyotis glauca W. W. Sm.

Kew Bull. 1911 : 344.

Syntypes : Burma : Kachin Hills, near Myit Kyina, *Shaik Mokim* 86; India : Assam : Darrang Dist., Charadi near Bishnath, *Burkill* 32470.

Kobresia gammiei Clarke (Cyperaceae).

Kew Bull. Add. ser. 8 : 68. 1908; J. Econ. Tax. Bot. 8(2) : 469. 1986.

Isotype : India : Sikkim : Tongsup, near Gongri, *Gammie* s. n.

Kurziodendron viridissimum (Kurz) Balakr. (Euphorbiaceae).

Bull. Bot. Surv. India 8 : 68. 1966 (1967).

Sabia viridissima Kurz

J. Asiat. Soc. Beng. 41(2) : 304. 1872.

Type : *Kurz* s. n., Acc. No. 97543.

Lactuca kashmiriana S. K. Mamgain et R. R. Rao (Asteraceae).

J. Bombay Nat. Hist. Soc. 83(2) : 406. 1986.

Holotype : India : Kashmir : Degwan, *B. M. Wadhwa* 67012A.

L. lahulensis Mamgain et R. R. Rao (Asteraceae).

Bull. Bot. Surv. India 27(1-4) : 120. 1985 (1987).

Holotype : India : Himachal Pradesh : Lahul-Kyelong, (3300 m), *U. C. Bhattacharyya* 40772.

Lagenandra lancifolia (Sch.) Thw. (Araceae).

Enum. Pl. Zeyl. 334. 1864; Engl. DC. Monogr. Phan. 11 : 622. 1879;
Bull. Bot. Surv. India 9(1-4) : 296. 1967.

Iso-syntype : Sri Lanka : *Thwaites* C. P. 3174.

L. nairii K. Ramamurthy et R. Rajan (Araceae).

J. Bombay Nat. Hist. Soc. 80(3) : 613. 1983.

Holotype : India : Kerala : Near the base of Adira Pally water falls as fully submerged patches in fresh running water on the bank of the Chalakudi River, Trichur Dist., 300 m, March 17, 1982, *Rajan* 73052.

Paratype : Ibid., March 22, 1981, *Ramamurthy* 66279.

L. thwaitesii Engl. (Araceae).

DC. Monogr. Phan. 11 : 621. 1879; Engl. Das Pflanzenr. 73. 230-231. 1920; Bull. Bot. Surv. India 9(1-4) : 296. 1967.

Iso-syntype : Sri Lanka : *Thwaites* C. P. 3173.

L. undulata Sastry (Araceae).

Bull. Bot. Surv. India 9 : 294. 1967 (1968).

Holotype : India : Arunachal Pradesh : From Amjee, about 1220 m, Subansiri Dist., May 21, 1966, *A. R. K. Sastry* 45489A.

Lasianthus andamanicus Hook. f. var. **ciliatus** Deb et Gangopadhyay (Rubiaceae).

J. Bombay Nat. Hist. Soc. 84(2) : 458. 1987.

Holotype : India : South Andaman : Beadnabad, ± 50 m, Nov. 30, 1973, *N. P. Balakrishnan* 654.

Paratypes : Little Andaman : Hut Bay, Sea-level, Sept. 1, 1976, *N. Bhargava* 4339; North Nicobar : Mildera, ± 30 m, May 4, 1977, *P. Chakraborty* 5600.

L. burmanicus Gang. et T. Chakrab. (Rubiaceae).

J. Econ. Tax. Bot. 13(1) : 117. 1989.

Holotype : Burma : Myit-Kyina Dist., Kadu, Ba-Bnlaung, Nov. 9, 1930, *Badal Khan* 11836.

L. lucidus var. **caudisepalus** Deb et Gangopadhyay (Rubiaceae).

J. Bombay Nat. Hist. Soc. 84(2) : 458. 1987.

Holotype : Arunachal Pradesh : Subansiri Dist., Apa Tani Valley, Hapolli, 1590 m, April 26, 1965, *Cox et Hutchinson* 499.

Lasianthus meeboldii Deb et Gangopadhyay (Rubiaceae).

J. Bombay Nat. Hist. Soc. 84(2) : 412. 1987.

Holotype : Burma : Tenasserim, Yaundan, March, 1911, *A. Meebold* 14790.

Paratype : Mergui, Sandawut Reserve, 50 ft., Jan. 31, 1919, *C. G. Rogers* 440 m.

L. wallichii Wt. var. **glabriusculus** Deb et Gangopadhyay (Rubiaceae).

J. Bombay Nat. Hist. Soc. 84(2) : 462. 1987.

Holotype and Isotype : Burma : March 1, 1849, *Falconer* 885.

Laurembergia agastyamalayana Henry (Haloragaceae).

J. Bombay Nat. Hist. Soc. 62(3) : 604. 1965.

Holotype : India : Tamil Nadu : Agastyamalai Hills, Tirunelveli Dist., 1400 m, Aug. 26, 1963 m, *Henry* 17328A.

Leontopodium simbrilligerum J. R. Drumm. (Compositae).

Kew Bull. 1910 : 76. 1910; Bull. Bot. Surv. India 14 : 171. 1972 (1975).

Holotype : W. Himalaya, Chumbi Valley, 3350 m, *King's Collector.*

s. n.

Lescuraea darjeelingensis Vohra (Musc.)

Bull. Bot. Surv. India 20 : 150. 1978 (1979).

Holotype : India : W. Bengal : Darjeeling, Tongloo, Batashi Road, 3060 m, June 17, 1961, *Wadhwa* 166.

Paratype : Bhutan : Chi La-Ha, 3000-4000 m, Aug. 20, 1963, *Wadhwa* 32.

Leucas lavandulifolia Rees var. **nagalapuramiana** Chandr. et Srin. (Labiatae).

Bull. Bot. Surv. India 17 : 165. 1975 (1978).

Holotype : India : Andhra Pradesh : Chittoor Dist., Nagalapuram Hills, Oct. 4, 1974, *Chandbose* 45228A.

L. mollissima Wall. var. **sebastiana** Subba Rao et Kumari (Labiatae).

Bull. Bot. Surv. India 11 : 452. 1969 (1971).

Holotype : India : Andhra Pradesh : Cherukonda, 900 m, Visakhapatnam, Dist., Jan. 4, 1968, *Subba Rao et Kumari* 29743A.

L. mukerjiana Subba Rao et Kumari (Labiatae).

Bull. Bot. Surv. India 10 : 358. 1968.

Holotype : India : Andhra Pradesh : Cherukonda, 1300 m, Visakhapatnam Dist., Aug. 29, 1966, *Subba Rao* 28182A.

Leucas vestita Benth. var. **devicolamensis** Shetty & Vivek. (Labiatae).

Bull. Bot. Surv. India 10(2) : 236. 1968 (1969).

Holotype : India : Kerala : Kottayam Dist., Upper Vagavurrai, Devicolam, Aug. 11, 1967, *Shetty & Vivekananthan* 28372A.

Limnopoia meeboldii (Fischer) C. E. Hubb. (Poaceae).

Hook. Ic. Pl. t. 3432. 1943; Fasc. Fl. Ind. 14 : 38. 1984.

Isotype : India : Kerala : Cochin, Chalakudi, Nov. 1910, *A. Meebold* 12520.

Lindernia crustacea (L.) F. Muell. var. **clarkei** Giri et Majumdar (Scrophulariaceae).

J. Econ. Tax. Bot. 3(1) : 253. 1982.

Holotype : India : Bihar : Noatolea, Lohardugga, Chotanagpore, 2000 ft., Nov. 12, 1883, *C. B. Clarke* 34173.

Liparis tigerhillensis A. P. Das et Chanda. (Orchidaceae).

J. Econ. Tax. Bot. 12(2) : 403. 1988.

Holotype : W. Bengal : Darjeeling, Tigher Hill, 2500 m, July 16, 1982, *A. P. Das* 1051.

Isotypes : Ibid., 1051 A & B.

Listera nandadeviensis P. K. Hajra (Orchidaceae).

Bull. Bot. Surv. India 25(1-4) : 181-182. 1983 (1985).

Holotype : India : Uttar Pradesh : Chamoli Dist., Himgoli, Nandadevi National Park, *P. K. Hajra* 73202A.

Litosanthes wardii (C. E. C. Fischer et Kaul) Gang. et T. Chakrab. (Rubiaceae).

J. Econ. Tax. Bot. 13(1) : 119. 1989.

Lasianthus wardii C. E. C. Fischer et Kaul

Kew Bull. 1940 : 292. 1941.

Type : Upper Burma : Nam Tamai, 3000-3500 ft., 1200-1350 m, 3.5 . 1926, *F. Kingdon Ward* 6679.

Litsea lakshmanniana Keshav. (Lauraceae).

Curr. Sci. 56(8) : 371. 1987.

Isotype : India : Karnataka : Coorg Dist., Shola forests of Tadiandamol, Oct. 28, 1983, 1550 m, *K. R. Keshavamurty et al.*, 4414C.

Lobelia clorata Wall. (Lobeliaceae).

Pl. Asiat. Rar. 2 : 42. 1831; Fasc. Fl. Ind. 19 : 45. 1988.

Type : India : Pundua, *F. De Silva*, Wall. List No. 1307.

Lobelia erectiuscula Hara (Lobeliaceae).

Journ. Jap. Bot. 4 : 328. 1965; Fasc. Fl. Ind. 19 : 49. 1988.

L. erecta Hook. f. & Thoms.

Journ. Linn. Soc. Bot. 2 : 28. 1958, non De Vries 1845; Clarke, Fl. Brit. Ind. 3 : 426. 1881.

Type and Isotype : Sikkim, Chula, 10-12000 ft., Nov. 6, 1849, *J. D. Hooker s. n.*

L. microcarpa Clarke (Lobeliaceae).

Fl. Brit. Ind. 3 : 424. 1881; Fasc. Fl. Ind. 19 : 52. 1988.

Isotype : "Tenassserim and Andamans": *Helper Kew Dist.*, n. 3447.

L. terminalis Clarke var. **terminalis** (Lobeliaceae).

Fl. Brit. Ind. 3 : 424. 1881; Fasc. Fl. Ind. 19 : 56. 1988.

Isotype : W. Mymensingh, Bhyadanga, Nov. 20, 1868, *C. B. Clarke* 8044.

Lonicera laceana Nayar et Giri (Caprifoliaceae).

J. Bombay Nat. Hist. Soc. 80(2) : 385. 1983.

Holotype : India : N. W. Himalaya : Kalatop Forest, 2128 m April 15, 1896, *J. H. Lace* 1316.

L. magnibracteata Nayar et Giri (Caprifoliaceae).

J. Econ. Tax. Bot. 3(2) : 593. 1982.

Holotype and Isotype : India : Sikkim Himalaya : Patan La, 10,000 ft., (3040 m), July 16, 1877, *G. King* 4195.

Luffa hermaphrodita Singh et Bhandari (Cucurbitaceae).

Baileya 11(4) : 133. 1963; Fasc. Fl. Ind. 11 : 73. 1982.

Type : India : Rajasthan : Jodhpur, *Bhandari* 1527A.

Macaranga nicobarica Balakr. et Chakrab. (Euphorbiaceae).

Gard. Bull. Singapore 31 : 57. 1978.

Holotype : India : Nicobar : Katchal Island, Mildera, 11th Aug. 1974, *Chakraborty* 2041A.

Paratypes : Ibid., 2584C-D.

Maesa andamanica Kurz var. **longipedicellata** Nayar et Giri (Myrsinaceae).

Bull. Bot. Surv. India 27 (1-4) : 263. 1985 (1987).

Holotype : S. Andaman : Without precise locality, Nov. 15, 1901, *R. L. Heinig* 451.

Maesa kanjilalii Majumdar et Giri (Myrsinaceae).

J. Bombay Nat. Hist. Soc. 78(1) : 149. 1981.

Holotype : India : Assam : Gouhati Road, 6-7 miles from Shillong, Oct. 14, 1930, *P. C. Kanjilal* 8684.

M. macrophylla Wall. var. **magnidentata** Nayar et D. Das (Myrsinaceae).

Indian For. 95(6) : 400. 1969.

Holotype : India : Naga Hills, April 1896, *Dr. King's Collector s. n.*

M. truncata A. R. K. Sastry (Myrsinaceae).

J. Sen Mem. Vol., pp. 433-436. 1969.

Holotype : India : Arunachal Pradesh: Subansiri Dist., Palin vicinity, about 1600 m., May 17, 1966, *A. R. K. Sastry* 45387A.

M. wardii Nayar & Giri (Myrsinaceae).

Bull. Bot. Surv. India 17 : 182-184. 1975 (1978).

Holotype : India : Nagaland : Naga Hills, *F. Kingdon Ward* 11241.

Mallotus subramanyamii Ellis (Euphorbiaceae).

Bull. Bot. Surv. India 25(1-4) : 199-201. 1983 (1985).

Holotype : South India : On the western slopes of Nilgiri Hills, Moolakadu, Ouchterlony, July 29, 1970, *Ellis* 34963A.

Paratype : Northern slopes of Nilgiris, no date given as (*M. muricatus* Rottlera). Anon., *s. n.*

Manisuris divergens (Hack.) O. Kuntze var. **hirsuta** (Fischer) Jain (Poaceae).

Bull. Bot. Surv. India 12 : 12. 1970 (1972).

Type : India : Karnataka (Mysore) : Bababudan Hills at Kalhatti, 1800 m, Oct. 1908, *Meebold* 10559.

M. forficulata Fischer (Poaceae).

Kew Bull. 1933: 353. 1933. emend. Jain; Fischer in Fl. Mad. 1761. 1934 *pro parte*; Bull. Bot. Surv. India 12 : 12. 1970 (1972).

Paratypes : India : Maharashtra : Khandala, alt. 600 m Sept. 1907, *Meebold* 8850; Mysore : Belgaum, Amboli Ghat, Oct. 6, 1900, *Talbot* 4305A; Kerala : Kavalay Cochin, 900 m, Nov. 1910, *Meebold* 12221.

Type : India : Maharashtra : Mahabaleshwar, Sept. 26, 1891, *Lisboa* *s. n.*

M. goaensis Rolla Rao et Hemadri (Poaceae).

Bull. Bot. Surv. India 10 : 106-109. 1968.

Holotype : India : Goa : Verna Village on Cortalim-Madgoa Road, Nov. 8, 1962, *Rolla Rao* 84474A.

Paratype : India : Goa : Provorium, Nov. 8, 1963, *Rolla Rao* 92850A.

***Manisuris mysorensis* Jain et Hemadri (Poaceae).**

Bull. Bot. Surv. India 10 : 280-282. 1968; Ibid. 12 : 15. 1970 (1972).

Holotype : India : Mysore : Castle Rock, Oct. 25, 1902, *Gammie* 15643A.

Isotypes : Ibid., 15643B and 15643C (A-C mounted on one sheet).

Paratypes : Mysore : Amboli Ghat, Oct. 6, 1960, *Talbot* 4305 D-E *pro parte*.

***M. ratnagirica* Kulkarni et Hemadri (Poaceae).**

Indian For. 100(4) : 250. 1974.

Holotype : India : Maharashtra : On the open rocky plateau, near Chaukul along the Amboli Ghat in Ratnagiri Dist., ± 650 m, Oct. 12, 1970, *B. G. Kulkarni* 121638A.

***M. santapaui* Jain et Deshpande (Poaceae).**

Bull. Bot. Surv. India 10 : 277-279. 1968; Ibid. 12 : 15. 1970 (1972).

Holotype : India : Maharashtra : Ratnagiri, in exposed pools on laterite hill top, Sept. 15, 1961, *C. Saldanha* CS 7130A.

Isotypes : Ibid., c-d.

Paratype : Ibid., *C. Saldanha* CS 7132B.

***M. talbotii* (Hook. f.) Bor (Poaceae).**

Grass. Burm. Ceyl. Ind. & Pak. 192. 1960; Bull. Bot. Surv. India 12(1-4) : 16. 1970 (1972).

Isotype : India : Goa : Marmagoa, Oct. 15, 1891, *Talbot* 2572.

***Marsdenia tirunelvelica* A. N. Henry et K. Subramanyam (Asclepiadaceae).**

J. Bombay Nat. Hist. Soc. 73(1) : 186. 1976.

Holotype : India : Tamil Nadu : Agastymalai Hills, Tirunelveli Dist., 1400 m, April 25, 1972. *Henry* 8421A.

***Mastixia tetrandra* (Thw.) Clarke**

Fl. Brit. Ind. 2: 745. 1879; Fasc. Fl. Ind. 19 : 27. 1988. *pro parte*.

var. *tetrandra* Matthew (Cornaceae).

Blumea 23 : 78. 1976.

Isotype : Ceylon : *Thwaites* C. P. 2441.

***Meconopsis bella* Prain (Papaveraceae).**

J. Asiat. Soc. Beng. 63. 2 : 82. 1894 et 64. 2 : 321. 1896; Fasc. Fl. Ind. 17 : 14. 1984.

Lectotype : India : Sikkim : Peykeangla, Jongri, July 1887, *Dr. King* s. n.

Meconopsis grandis Prain (Papaveraceae).

J. Asiat. Soc. Beng. 64. 2 : 320. 1895 et in Ann. Roy. Bot. Gard. Calcutta 9. 1 : 2. tab. 2. 1901; Fasc. Fl. Ind. 17 : 17. 1984.

Syntype : India : Sikkim : Jongri, 3900 m, June 23, 1892, *G. A. Gammie* 199.

M. horridula Hook. f. et Thoms. (Papaveraceae).

Fl. Ind. 1 : 252. 1855 et Hook. f. Fl. Brit. Ind. 1 : 118. 1872; Fasc. Fl. Ind. 17 : 19. 1984.

Type : India : Sikkim : Alpina, 4300-5100 m, *J. D. Hooker s. n.*

M. lyrata (Commins and Prain) Fedde (Papaveraceae).

Engl. Pflanzenr. 4. 104 : 246. 1909. *nomen synonymum* ex Prain.

Bull. Misc. Inf. Kew 1915 : 142. 1915; Fasc. Fl. Ind. 17 : 21. 1984.

Lectotype : India : Sikkim : Phullot, 3600 m, Aug. 1887, *Dr. King s. n.*

Syntypes : Tankra, Aug. 2, 1892, *Gammie* 415; chianie Nepal Frontier, 3000 m, Aug. 1888, *Dr. King s. n.*

M. napaulensis DC. (Papaveraceae).

Prodr. 1 : 121. 1824; Fasc. Fl. Ind. 17 : 22. 1984.

Isotype : *Wallich* 8121.

M. paniculata Prain (Papaveraceae).

J. Asiat. Soc. Beng. 64. 2 : 316. 1896; Fasc. Fl. Ind. 17 : 23. 1984.

Syntype : Nepal : Gossain Than, *Wallich* 1823B.

M. robusta Hook. f. et Thoms. (Papaveraceae).

Fl. Ind. 1 : 253. 1855 et Hook. f. Fl. Brit. Ind. 1 : 118. 1872 *pro parte*; Fasc. Fl. Ind. 17 : 25. 1984.

Isolectotype : In Himalaya temperate, Kumaon, 2400 m, *Wallich* 8124.

M. simplicifolia (D. Don) Walp. (Papaveraceae).

Repert. Bot. Syst. 1 : 110. 1842; Fasc. Fl. Ind. 17 : 26. 1984.

Holotype : Nepal : Gossain Than, *Wallich* 8125.

M. sinuata Prain (Papaveraceae).

J. Asiat. Soc. Beng. 64(2) : 314. 1896 (excl. var. *pratti*); Fasc. Fl. Ind. 17 : 28. 1984.

Syntype : India : Sikkim : Jongri, Pey-king-la, Aug. 1987, *Dr. King s. n.*

M. superba King ex Prain (Papaveraceae).

J. Asiat. Soc. Beng. 64. 2 : 317. 1896; Fasc. Fl. Ind. 17 : 30. 1984.

Holotype : Bhutan Himalaya : Ho-Ko-Cha, June 16, 1884. *Dungboo* 280.

Meconopsis villosa (Hook. f. ex Hook.) G. Tayl. (Papaveraceae).

Acc. Genus Meconopsis 28. 1934; Fasc. Fl. Ind. 17 : 30. 1984.

Isotype : India : Sikkim : *J. D. Hooker s. n.*

Melocalamus indicus R. Majumdar (Gramineae).

Bull. Bot. Surv. India 25(1-4) : 236-237. 1983 (1985).

Holotype : India : Assam : Cachar, Bhuban Hill, *Majumdar* 73083.

Melothria angulata Chakravarty (Cucurbitaceae).

J. Bombay Nat. Hist. Soc. 50 : 899. 1952; Fasc. Fl. Ind. 11 : 78. 1982.

Type : South India : Gomata, *Malcolmpeeth* 81.

M. assamica Chakravarty var. **scabra** Chakravarty (Cucurbitaceae).

Rec. Bot. Surv. India 17(1) : 145. 1959.

Type : India : Assam : Goalpara, Chirang Duar, *King's Collector s. n.*

M. maderaspatana (L.) Cogn. (Cucurbitaceae).

DC. Monogr. Phan. 3 : 623. 1881; Fasc. Fl. Ind. 11 : 83. 1982.

Type : N. W. India : *Royle s. n.*

M. mucronata (Blume) Cogn. var. **umbellata** Chakravarty (Cucurbitaceae).

Rec. Bot. Surv. India 17(1) : 150. 1959; Fasc. Fl. Ind. 11 : 85. 1982.

Type : South India : Without precise locality, *Rodriguez* 1955.

Memecylon collinum Balakr. et N. G. Nair (Melastomataceae).

Bull. Bot. Surv. India 24 : 30. 1982 (1983).

Holotype : India : North Andaman : Saddle Peak, 500-700 m, open scrub jungles in lateritic soils, Dec. 1, 1976, in flower, *Balakrishnan & Nair* 4764A.

M. subramanii Henry (Melastomataceae).

J. Bombay Nat. Hist. Soc. 77(3) : 492. 1980.

Holotype : India : Tamil Nadu : Kannikatti, Tirunelveli Dist., 765 m, April 22, 1972, *Henry* 8407.

Merrillioanax cordifolia A. R. K. Sastry (Araliaceae).

Blumea 15(2) : 297. 1967.

Holotype : India : Arunachal Pradesh : Subansiri Dist., Begi-Amjee, ±1540 m, May 12, 1966, in flower, *A. R. K. Sastry* 44893A.

Mesua ferrea Linn. subsp. **pulchella** Vesque (Guttiferae).

DC. Monogr. Phan. 8 : 631. 1893 ; Bull. Bot. Surv. India 5(3 & 4) : 339. 1963.

var. **pulchella**

Mesua pulchella Planchon & Triana

Ann. Sci. Nat. Ser. 4. 15 : 307. 1861.

M. coromandeliana Thw.

Enum. Pl. Zeyl. 50. 1858. *non* Wt. (1839).

Type : Ceylon: Central Provinces, forests of Galle and Ratnapoora, *Thwaites* 3404.

var. **coromandeliana** (Wt.) Maheshwari

M. coromandeliana Wt.

Icon. t. 117. 1839.

M. ferrea Wt. & Arn.

Prodr. 1 : 102. 1834. *non* Linn. (1753).

Type : Courtallum, *Wight* 132.

Mesua ferrea Linn. subsp. **thwaitesii** Vesque (Guttiferae).

DC. Monogr. Phan. 8 : 632. 1893; Bull. Bot. Surv. India 5(3 & 4) : 339. 1963.

M. thwaitesii Planchon & Triana

Ann. Sci. Nat. Ser. 4. 15 : 305. 1861.

Type : Ceylon: *Thwaites* 2675.

Microstegium borianum P. R. Sur (Poaceae).

J. Bombay Nat. Hist. Soc. 79(3) : 652. 1982; J. Econ. Tax. Bot. 6(1) : 168. 1985.

Holotype : India : Meghalaya : Khasia and Jaintea Hills, Lowlyngdoh, Sept. 26, 1937, *G. K. Deka* 15683.

M. vagans (Nees ex Steud.) A. Camus (Poaceae).

Ann. Soc. Linn. Lyon, 68 : 200. 1921; J. Econ. Tax. Bot. 6(1) : 173. 1985.

Pollinia vagans Nees ex Steud.

Sym. Pl. Glum. 1 : 410. 1855; Hook. f. Fl. Brit. Ind. 7 : 115. 1896.

Type : Nepal, *Wallich* 8807B.

M. vimineum (Trin.) A. Camus (Poaceae).

Ann. Soc. Linn. Lyon, n. s. 62 : 201. 1921; J. Econ. Tax. Bot. 6(1) : 174. 1985.

Andropogon vimineus Trin.

Mem. Acad. Sci. Petersb. ser. 6. 2 : 268. 1832.

Pollinia imberbis Nees ex Steud.

Sym. Pl. Glum. 1 : 410. 1855; Hook. f. Fl. Brit. Ind. 7 : 117. 1896.

Type : Nepal: *Wallich* 8832.

***Microula duthiei* Ban. (Boraginaceae).**

Indian For. 92(10) : 644. 1966.

Holotype : India : Sikkim : On Teesta Valley above Thangu, 4114 m, July 6, 1903, *F. E. Younghusband* 1538A, Acc. No. 308324.

Isotypes : *F. E. Younghusband* 1538 B-D, Acc. Nos. 308325, 308326, 308327.

***Minuartia ebracteolata* Majumdar & Giri (Caryophyllaceae).**

Candollea 38(1) : 345. 1983.

Holotype : India : Himachal Pradesh : Lahul, Batal, 4000 m, Sept. 11, 1961, *N. C. Nair* 16867.

***M. kashmirica* (Edgew.) Mattfeld (Caryophyllaceae).**

Bot. Jahrb. 57(Beibl. 126) : 32. 1921; Candollea. 38(1) : 346. 1983.

Type : India : Kashmir : Regiotemp, 5000-6000 ped. (1520-1820 m), *T. Thomson s. n.*

***Mitrophora andamanica* Thoth. ex Das (Annonaceae).**

J. Bombay Nat. Hist. Soc. 64(3) : 430. 1967.

Holotype : India : South Andaman : Baratang Island, April 30, 1964, *K. Thothathri* 10840A.

Isotypes : Ibid., *Thothathri* 10840 B-F.

***Moghania gracilis* Mukerjee (Leguminosae).**

Bull. Bot. Soc. Beng. 6(1) : 22. 1952.

Type : India : Western Ghats : Castle Rock, 600 m, Sept. 1918, *Bell* 4347.

***M. praecox* (Clarke ex Prain) H. L. Li var. *robusta* Mukerjee (Leguminosae).**

Bull. Bot. Soc. Beng. 6(1) : 19. 1952.

Type : Peninsular India : Flowering in January, Chanda, Yenkatapur, *J. F. Duthie* 9408A.

***M. rollae* Billote et Hemadri (Leguminosae).**

J. Econ. Tax. Bot. 3(2) : 617. 1982.

Holotype : India : Maharashtra : Ahmadnagar Dist., in association with *Carvia callosa* (Nees) Bremek. on a hill-slope (\pm 1200 m) of Panshet Forest, Bari near Kalsubai Hill, Oct. 9. 1970, *B. M. Wadhwa* 128313A.

***Molineria grandis* J. N. Mittra (Hypoxidaceae).**

J. Indian Bot. Soc. 43(3) : 487-489. 1964.

Holotype : India : Duphla Hills, Camp. 3, Nov. 15, 1875, *J. Lister* 337.

Molineria prainiana Deb (Hypoxidaceae).

Bull. Bot. Surv. India 6 : 77. 1964 (1965).

Holotype : India : Arunachal Pradesh : From Janakmukh, Abor Hills, Jan. 23, 1912, *I. H. Burkill* 36471.

Momordica macrophylla Gage (Cucurbitaceae).

Rec. Bot. Surv. India 3 : 61. 1908; Fasc. Fl. Ind. 11 : 94. 1982.

Type : Burma : Minbu, Pomin, riverside, *Seikh Mokim* 253.

Syntype : Tenasserim, Taepo, *Gallatly* 80.

Mycetia javanica (Bl.) Reinw. ex Korth. (Rubiaceae).

Netherl. Kruid. 118. 1841; Bull. Bot. Surv. India 28 : 118. 1986 (1988).

Bertiera javanica Bl.

Bijdr. 987. 1826.

Type : Penang, 1822, Wall. Cat. No. 6282.

Adenosacme javanica Wall. (Cat. 6282, 1832, *nom. nud.*) ex Ridley.

Fl. Mal. 2 : 63. 1923.

M. malayana Wall. ex Ridley

Fl. Mal. 2 : 63. 1923.

Type : Penang, 822, Wall. Cat. No. 6282.

M. listeri Deb (Rubiaceae).

Blumea 14(1) : 241. f. 1. 1966; Bull. Bot. Surv. India 28 : 120. 1986 (1988).

Type : India : Arunachal Pradesh : Daphla Hills, 1874, *J. L. Lister s. n.*

M. mukerjiana Deb et Dutta (Rubiaceae).

Ind. For. 91(5) : 272. t. 1. 1965 ; Sci. & Cult. 33 : 60. 1967; Bull. Bot. Surv. India 28 : 123. 1986 (1988).

Holotype : India : Assam : Lakhimpur, Makum Hills, *G. K. Deka* 16961.

M. rodgeri Deb et Mondal (Rubiaceae).

Bull. Bot. Surv. India 28 : 127. 1986 (1988).

Holotype : Burma : Ruby Mines Division, Mogak, July, 1910, *A. Rodger* 148A.

Isotype : Ibid., 148B.

M. stipulata (Hook. f.) O. Ktze. (Rubiaceae).

Rev. Gen. Pl. 1 : 289. 1891; Bull. Bot. Surv. India 28 : 129. 1986 (1988).

subsp. stipulata

Adenosacme stipulata Hook. f.

Fl. Brit. Ind. 3 : 95. 1880.

Isotype : East Nepal : 600-1500 m, *J. D. Hooker s. n.*

Mycetia stipulata (Hook. f.) O. Ktze. subsp. **macrostachya** (Hook. f.) Deb (Rubiaceae).

Bull. Bot. Surv. India 28 : 129. 1986 (1988).

Adenosacme macrostachya Hook. f.

Fl. Brit. Ind. 3 : 96. 1880.

Isotype : Arunachal Pradesh : Mishmi Hills, *Griffith s. n.*

Neanotis monosperma (Wt. & Arn.) Lewis var. **tirunelvelica** A. N. Henry & M. Chandrabose (Rubiaceae).

Bull. Bot. Surv. India 17(1-4) : 188. 1975.

Holotype : India : Tamil Nadu : Tirunelveli Dist., Manjanamparai in Singampatti R. F., Sept. 4, 1963, *Henry* 17454A.

N. sahyadrica Billore et Mudaliar (Rubiaceae).

J. Econ. Tax. Bot. 3(1) : 321. 1982.

Holotype : India : Maharashtra : On the slopes of Taramati dongar Harishchandragarh, Thana Dist., Sept. 18, 1968, *K. V. Billore* 115664A.

Paratype : India : Maharashtra : From higher slopes of Ajoba dongar, Washala forest range, Thana Dist., Oct. 17, 1967, *K. V. Billore* 111944 C.

Neodriessenia scorpioidea (Stapf) Nayar (Melastomataceae).

Bull. Bot. Surv. India 16(1-4) : 25. 1974 (1977).

Driessenia scorpioidea Stapf

Hook. f. Ic. Pl. 25. t. 2414. 1895; Merrill in J. Str. Br. Roy. As. Soc. 440. 1921.

Compimia scorpioidea (Stapf) Ridley

J. Str. Br. Roy. As. Soc. 57 : 40. 1911.

Isotype : Borneo: *Haviland & Hore* 3222.

Neolitsea balakrishnanii T. Chakrab. & Goel. (Lauraceae).

J. Econ. Tax. Bot. 6(2) : 449. 1985.

Holotype : India : North Andaman Island : Saddle Peak, March 31, 1977, ±700 m, *Balakrishnan* 5379A.

N. nicobarica Goel. & T. Chakrab. (Lauraceae).

J. Econ. Tax. Bot. 6(2) : 450 (1985).

Holotype : India : Nicobar Islands : *sine loc. exact.*, April 4, 1884, *King's Collector* 491A.

Isotypes : Ibid., *King's Collector* 491B, 491E ♂.

Paratypes : Kamorta, Feb. 1875, *Kurz s. n.* ♀; *sine loc. exact.*, *Exped. Novara, Jelinek* 137. ♀.

Neoluffa sikkimensis Chakravarty (Cucurbitaceae).

J. Bombay Nat. Hist. Soc. 50 : 895. 1952; Fasc. Fl. Ind. 11 : 98. 1982.

Type : India : Sikkim Himalaya : King s. n.

Nothaphoebe nicobarica T. Chakrab. et Vasud. (Lauraceae).

J. Econ. Tax. Bot. 6(2) : 443. 1985.

Holotype : India : Andaman-Nicobar Islands : Great Nicobar Island, 40 km on East-West Road, Feb. 22, 1980, Dwivedi 7893A.

N. panduriformis (Hook. f.) Gamble var. **paucinervia** T. Chakrab. & Vasud. (Lauraceae).

J. Econ. Tax. Bot. 5(4) : 997. 1984.

Holotype : India : Andaman-Nicobar Islands : Great Nicobar Island, 25 km on East-West Road, Oct. 17, 1979, D. K. Hore 7222A.

Nothopégia beddomei Gamble var. **wynaadica** Ellis & Chandras. (Anacardiaceae).

Bull. Bot. Surv. India 12 : 257. 1970 (1972).

Holotype : India : Kerala : Cannanore Dist., Chandanathode, Peria, April 18, 1966, Ellis 27133A.

Oberonia seidenfadeniana Joseph et Vajravelu (Orchidaceae).

Bull. Bot. Surv. India 13 : 344. 1971 (1973).

Holotype : India : Tamil Nadu : From Andiparai Shola, Eastern side of Anamalai Hills, 1350 m, Sept. 22, 1962, Joseph 17476A.

O. sulcata Joseph et Chowd. (Orchidaceae).

J. Bombay Nat. Hist. Soc. 63(1) : 54. 1966.

Holotype : India : Arunachal Pradesh : Selari Forest, 9 km N. E. of Bomdi La, Kameng Frontier' Dist., ±2700 m, about Lat. N. 27°19' and Long. E. 92°25', Sept. 28, 1964. Joseph 40358.

Ophiorrhiza caudipetala Deb et Mandal (Rubiaceae).

Kew Bull. 37(3) : 483. 1982.

Holotype : India : Meghalaya : ±1830 m, June 4, 1886, Clarke 44072C.

O. chandrasekharanii G. V. Subba Rao et G. R. Kumari (Rubiaceae).

J. Bombay Nat. Hist. Soc. 81(1) : 156. 1984.

Holotype : India : Andhra Pradesh: Vandachinta, Visakhapatnam Dist., June 1, 1968, Subba Rao 30049.

***Ophiorrhiza infundibularis* Balakr. (Rubiaceae).**

Bull. Bot. Surv. India 24 : 61. 1982 (1983).

Holotype : India : Great Nicobar, 31-32 km on East-West Road, ca 100 m, shaded places near streams, June 14, 1977, *Balakrishnan* 5745A.

***O. nairii* Ramamurthy et Rajan (Rubiaceae).**

J. Bombay Nat. Hist. Soc. 82(1) : 174. 1985.

Holotype : India : Kerala : Idduki Dist., Adimali Reserve Forest, 1800 m, March 28, 1980, *Ramamurthy* 66561.

***O. nicobarica* Balakr. (Rubiaceae).**

Reinwardtia 9(4) : 411-414. 1980.

Holotype : India : Great Nicobar Island, 17 km from Campbell Bay to Alexandra river, ca 75 m above s.l. Aug. 21, 1975, *Balakrishnan* 3027A.

***O. seshagiriana* Sikdar et Maiti (Rubiaceae).**

J. Bombay Nat. Hist. Soc. 78(1) : 143. 1981.

Holotype : India : W. Bengal : Buxa, Duar Forest Range, Jalpaiguri Dist.. on the way to Sinchu from Buxaduar, ±1800 m, May 16, 1976, *J. K. Sikdar* 4681A.

Isotypes : Ibid., *J. K. Sikdar* 4681 B-E.

***O. tirunelvelica* Henry et Subr. (Rubiaceae).**

Bull. Bot. Surv. India 12 : 277. 1970 (1972).

Holotype : India : Tamil Nadu : Tirunelveli Dist., near upper Godaiyar, Sept. 2, 1963, *Henry* 17419A.

***Ormosia assamica* Yakovlev (Leguminosae).**

J. Arn. Arb. 52(4) : 691. 1971.

Holotype : India : Aka Hills, Dec. 1890, *Badul Khan* 10.

***Orophaea katschallica* Kurz emend. Thoth. (Annonaceae).**

J. Bot. (Lond.) 4 : 323-324. 1875; Indian For. 92(8) : 530. 1966.

Type : Nicobar. Katchall, *Kurz s. n.*

***Oryza indandamanica* Ellis (Gramineae).**

Bull. Bot. Surv. India 27(1-4) : 225-227. 1985 (1987).

Holotype : India : Andamans : Rutland Island, July 26, 1986, *Ellis* PBL 12248.

***Osbeckia abrahamii* Giri et Nayar (Melastomataceae).**

J. Bombay Nat. Hist. Soc. 81(2) : 434. 1984.

Holotype : India : Kerala : Travancore, *Narayanaswami* 1379.

Osbeckia arunkumarensis Nayar et Giri (Melastomataceae).

J. Bombay Nat. Hist. Soc. 81(2) : 436. 1984.

Holotype : India : Sikkim : Eastern Himalaya, *Griffith* (Kew Distrib.) No. 2248.

O. darjeelingensis Giri et Nayar (Melastomataceae).

Bull. Bot. Surv. India 25(1-4) : 241. 1983 (1985).

Holotype : India : W. Bengal : Jhenaikuri, Darjeeling, Dec. 1879, *Gamble* 7597A.

O. lineolata Gamble (Melastomataceae).

Kew Bull. Misc. Inf. 241. 1819; Bull. Bot. Surv. India 25(1-4) : 244. 1983(1985).

Syntype : S. India : Pulney Hills, *Wight s. n.*, Kew Distrib. 1090 (K, CAL, Lectotype chosen at CAL).

O. lineolata Gamble var. **anamalayana** Giri et Nayar (Melastomataceae).

Bull. Bot. Surv. India 25(1-4) : 244. 1983 (1985).

Holotype : India : Tamil Nadu : Anamalai Hills, Konalar, 1950 m, Nov. 17, 1980, *Chandrabose* 57793.

O. nayarii Giri (Melastomataceae).

J. Econ. Tax. Bot. 4(2) : 609. 1983.

Holotype : India : Meghalaya : Khasia, 2000 ft., Sept. 14, 1886, *C. B. Clarke* 44776A.

Paeonia emodi Wall. ex Royle (Paeoniaceae).

Ill. Bot. Himal. 57. 1834; Fasc. Fl. Ind. I : 5. 1978.

Type : Shalma Mountain, Kumaon, Wall. Cat. No. 4727.

Panax assamicus Ban. (Araliaceae).

Bull. Bot. Surv. India 10 : 23. 1968.

Holotype : India : Eastern Himalaya, Khasia Mountains, *J. D. H. et T. T. s. n.*, Acc. No. 191778.

P. pseudoginseng Wall. (Araliaceae).

Trans. Med. Rhys. Soc. Cal. 4 : 117. 1829; Bull. Bot. Surv. India 10(1) : 23, 1968.

Lectotype : Nepal : Central Himalaya, Sheopore, June 1821, Wall. Cat. No. 3730, Acc. No. 191766.

Panax sikkimensis Ban. (Araliaceae).

Bull. Bot. Surv. India 10 : 21. 1968.

Holotype : India : Sikkim Himalaya, July 25, 1909, Smith & Cave 2781.

Panicum garadei Raghavan et Karthik. (Gramineae).

Bull. Bot. Surv. India 24 : 146. 1982 (1983).

Holotype : India : Karnataka : Dharwar, Sept. 16, 1907, L. D. Garade s. n.

P. jainii Karthik. et Raghavan (Gramineae).

Bull. Bot. Surv. India 24 : 145. 1982 (1983).

Holotype : Pakistan : Sind : Mirpurkas, Dec. 12, 1909, R. K. Bhide s. n.

P. nehruense Jauhar et Joshi (Gramineae).

Bull. Bot. Surv. India 8 : 97. 1966 (1967).

Isotype : *P. P. Jauhar* 9B (the type was isolated from a composite seed lot of *Panicum coloratum* obtained from Jodhpur where it is reported to have been introduced from Australia. The specimen has been named as *Panicum nehruense* and was reported of being arrived at CAL).

Paraboea nagalandiana Deb et Dutta (Gesneriaceae).

J. Bombay Nat. Hist. Soc. 85(1) : 168. 1988.

Holotype : India : Nagaland : Narum, 1200-1500 m, Dec. 1907. A. Meebold 7394.

Parakaempferia synantha A. S. Rao et D. M. Verma (Zingiberaceae).

Bull. Bot. Surv. India 11 : 206. 1969 (1971).

Holotype : India : Assam : Along Kana River, Chaldhowa, N. Lakhimpur, May 22, 1966, D. M. Verma 46585A.

Parardisia gamblei Nayar et Giri (Myrsinaceae).

Bull. Bot. Surv. India 28 : 250. 1986 (1988).

Holotype : Borneo, near Kuching, March 24, 1893, G. D. Havilland & C. Hore 608A.

Isotypes : Ibid., G. D. Havilland & C. Hore 608 B-E.

P. involucrata (Kurz) Nayar et Giri comb. nov. (Myrsinaceae).

Bull. Bot. Surv. India 28 : 248. 1986 (1988).

Ardisia involucrata Kurz

J. Asiatic Soc. Beng. 40(2) : 68. 1871.

Type : E. Himalaya, Herb. Griffith Kew Distrib. No. 3560.

Parvatia brunoniana (Wall.) Decaisne (Lardizabalaceae).

Arch. Mus. Hist. Nat. Paris 1 : 190. t. 12A. 1839; Fasc. Fl. Ind. 19 : 36. 1988.

Stauntonia brunoniana Wall.

Wall. Cat. No. 4952; 1824 *nom. nud.*

Type : Montes Silhet, Wall. Cat. No. 4952.

P. elliptica (Hemsl.) Nayar et Paul (Lardizabalaceae).

Fasc. Fl. Ind. 19 : 36. 1988.

Stauntonia elliptica Hemsl.

Kew Bull. 9, t. 2844. 1909.

Type : Assam : Gauhati, *Simons s. n.*, Acc. No. 16832.

Pauella sivagangana Ramam. et Seb. (Araceae).

Bull. Bot. Surv. India 8 : 348. 1966 (1967).

Holotype : India : Tamil Nadu : Esaniforest, Sivaganga, Ramnad Dist., Aug. 14, 1964, *Ramamurthy* 20794A.

Paratype : Ibid., Sept. 12, 1965, *Ramamurthy* 24685A.

Pauia belladonna Deb et Dutta (Solanaceae).

Indian For. 91(6) : 363-366. 1965.

Holotype : India : Arunachal Pradesh, Tirap Frontier Division, way to wakka, about 6 km away at about 2,100 m alt., July 2, 1961, *Deb*, 26422.

Pavonia procumbens (Wall. ex Wight & Arn.) Walp. (Malvaceae).

Rep. Bot. Syst. 1 : 301. 1842 non Casaretto 1842; Fasc. Fl. Ind. 19 : 194. 1988.

Isolectotype : India : *Wallich* 2688.

Pennilabium proboscideum A. S. Rao & J. Joseph (Orchidaceae).

Bull. Bot. Surv. India 10(2) : 231. 1968 (1969).

Holotype : India : Meghalaya : Between Umran and Umsaw besides the Gauhati-Shillong Road, July 23, 1966, *A. S. Rao* 45622.

Peristrophe bicalyculata (Retz.) Nees var. **subaequibracteata** S. S. R. Bennet (Acanthaceae).

J. Bombay Nat. Hist. Soc. 66(1) : 229. 1969.

Holotype : India : W. Bengal : Shyampur, Howrah Dist., Jan. 2, 1965, *Bennet* 1065A.

Isotype : Ibid., *Bennet* 1065B.

Phanera nicobarica Balakr. & Thoth. (Caesalpiniaceae).

Bull. Bot. Surv. India 17(1-4) : 201. 1975.

Holotype : India : Great Nicobar : 15 km on East-West Road, ±100 m, Aug. 23, 1975, *Balakrishnan* 3043A.

Isotypes : Ibid., *Balakrishnan* 3043B-C.

Paratypes : 18 km on North-South Road, ± 25 m, July 17, 1976, *Balakrishnan* 3824 A-B; on the way from Galathea Bay to Pulobaha Bay, ± 150 m, March 26, 1966, *Thothathri & Banerjee* 10661.

Pholidota convallariae var. **breviscapa** Deori et Joseph (Orchidaceae).

Bull. Bot. Surv. India 20 : 159. 1978 (1979).

Holotype : India : Arunachal Pradesh : From 43-camp, ± 1666 m, Kameng, *A. S. Rao* 50902A.

P. imbricata Lindl. var. **sessilis** Hook. f. (Orchidaceae).

Fl. Brit. Ind. 5 : 864. 1890; Bull. Bot. Surv. India 9(1-4) : 92. 1967 (1968).

Type : India : Naga Hills : Kohima, 1000-1500 m, *Prain s. n.*

Phrynum paniculatum Balakr. (Marantaceae).

Blumea 24 : 185-187. 1978.

Holotype : India : Great Nicobar Island, 8 km on East-West Road, Aug. 19, 1975, *Balakrishnan* 2979.

Paratypes : Ibid., 31-32 km on East-West Road, June 14, 1977, *Balakrishnan* 5758; on path to Shompenhouses, ex 36.8 km on East-West Road, June 17, 1977, *Balakrishnan* 5827.

Phyllanthus andamanicus Balakr. et N. G. Nair (Euphorbiaceae).

Bull. Bot. Surv. India 24 : 35. 1982 (1983).

Holotype : India : North Andaman ; Saddle peak ca 550 m, Dec. 1, 1976, *Balakrishnan & Nair* 4765A.

Pimpinella clarkeana Watt ex Banerji (Apiaceae).

J. Bombay Nat. Hist. Soc. 50(1) : 88. 1951.

Type : India : Manipur : Chingsow, 8000 ft., April 18, 1882, *G. Watt* 6556.

Paratype : Nepal : Arun Watershed, 10,000 ft., May 21, 1948, *Banerji* 446.

P. diversifolia DC. var. **sarmentifera** Goel et Bhattacharyya (Apiaceae).

J. Econ. Tax. Bot. 6(1) : 213. 1985.

Isotypes : India : Uttar Pradesh : Garhwal, Gangi, 3000 m, Aug. 10, 1978, *A. K. Goel* 46313 B.-C.

Pimpinella rollae Billiore et Hemadri (Apiaceae).

Indian For. 108(11) : 712. 1982.

Holotype : Raised from the seeds of *K. V. Billiore* 115489A-J, July 15, 1969, *K. V. Billiore* 115986A.

Isotypes : *K. V. Billiore* 115986 F-G.

Paratype : *K. V. Billiore* 115489A. The plant was observed in drying condition in the shade of a steep, overhanging rock on way to Kedarnath Temple (Popularly known as Harishchandragarh, Thana Dist., Maharashtra, India) from Panchnai, Nov. 17, 1968. Seeds were collected and shown in June 1969 in the experimental plots of BSI, Poona. A large number of herbarium specimens were prepared from sowing of fruits in addition to the live materials.

P. tongloensis P. K. Mukherjee (Apiaceae).

Bull. Bot. Surv. India 12 : 78. 1970 (1972).

Holotype : India: Sikkim : Oct. 14, 1868, *S. Kurz s. n.*, Acc. No. 188365.

Isotype : *S. Kurz s. n.*, Acc. No. 188367.

P. urceolata Watt ex Banerji (Apiaceae).

J. Bombay Nat. Hist. Soc. 50(1) : 89. 1951; Bull. Bot. Surv. India 12 (1-4) : 77. 1970 (1972).

Type : India : Sikkim : Tonglo, Rinchinpong and Gassing, 3500 ft., Oct. 2, 1862, *Anderson* 623.

Paratypes : Tonglo, without collector's name, Oct. 1, 1857 (two sheets); Tonglo? Oct. 14 & 21, 1868, *Kurz s. n.* (three sheets); Tonglo, 9000 ft., July 8, 1882, *Gamble s. n.*

Pinda concanensis (Dalzell) Mukherjee & Constance (Apiaceae).

Kew Bull. 1986. 41(1) : 226.

Heracleum concanense Dalzell

Hooker's Journ. Bot. Kew Gard. Misc. 2 : 260 (1850).

Isotype : *Heracleum concanense* : India : Bombay Presidency, in *Colibus concanensis*, Bombay Presidency, *Dalzell*.

Pittosporum eriocarpum Royle (Pittosporaceae).

Ill. Bot. Himal. 1 : 77. 1834; Fasc. Fl. Ind. 6 : 7. 1980.

Type : From above Sahasradhara to Mussoorie from 3000-6700 ft. of $30\frac{1}{2}^{\circ}$ Northern latitude, *Royle s. n.*

P. humile Hook. f. & Thoms. (Pittosporaceae).

Fl. Brit. Ind. 1 : 198. 1872; Fasc. Fl. Ind. 6 : 8. 1980.

Type : India : Khasi Hills, Banks of Borpani River, 5000 ft., *Hooker f. & Thomson*.

Pittosporum podocarpum Gagnepain var. **angustatum** Gowda (Pittosporaceae).

J. Arn. Arb. 32 : 295. 1951; Fasc. Fl. Ind. 6 : 11. 1980.

Type : China : Yunnan : Menglze, 7000 ft., *Henry* 10545.

P. reticosum Ridl. (Pittosporaceae).

Fl. Mal. Penins. 1 : 137. 1922.

Isotype : Pahang, 5-6000 ft., July 7, 1905, *H. C. Rabinson & L. Wray* 5444.

P. viridulum Nayar, Giri et Chandrasekaran (Pittosporaceae).

Fasc. Fl. Ind. 6 : 11. 1980.

Holotype : India : Tamil Nadu : Nilgiri, Nadugani forest, Kozhipalam, 825 m, Sept. 19, 1980, *Chandrasekaran* 69264.

Paratype : Ibid., *Chandrasekaran* 64100.

Plagiogyria elongata Dixit et Das (Plagiogyriaceae).

Proc. Indian Acad. Sci. (Plant Sci.) 90(4) : 377. 1981.

Paratypes : India : Assam : Shora, ±1400 m, Aug. 15, 1872, *Clarke* 18853; Meghalaya, ± 2000 m, *J. D. H. and T. T. s. n.*, Acc. No. 9467.

P. meghalayensis Dixit et Das (Plagiogyriaceae).

Proc. Indian Acad. Sci. (Plant Sci.) 90(4) : 378. 1981.

Holotype : India : Meghalaya : Shillong, Elephant Falls, March, 1979, *Rao* 4866.

P. triquetra Wall. ex Mett. (Plagiogyriaceae).

Abhandl. d. Senckenb. naturf. Ges. B. 2. 274. 1858; *Acrostichum* Wall. List No. 23. 1828 (*nom. nud.*); Proc. Indian Acad. Sci. (Plant Sci.) 90 (4) : 376. 1981.

Isotype : Nepal : 1921, *Wallich* 23, Acc. No. 9466.

Plagiothecium dehradunense Vohra (Plagiotheciaceae).

Bull. Bot. Surv. India 16(1-4) : 139. 1974 (1977).

Holotype : India : Uttar Pradesh : Dehra Dun, Rajpur, 900 m, Oct. 24, 1968, *C. L. Malhotra* 558.

Planchonella kingiana Rakesh et Thoth. (Sapotaceae).

J. Econ. Tax. Bot. 13(1) : 47. 1989.

Type : India : South Andaman, near Port Blair, Aug. 9, 1984, *Dr. King's Collector* 409.

P. kingiana Rakesh et Thoth. var. **andamanica** (Sapotaceae).

J. Econ. Tax. Bot. 13(1) : 47-49. 1989.

Type : India : Andaman : Herb. Hort. Calcuttensis, Dec. 1900, Dr. Prain's Collector 64.

Pogostemon travancoricus Bedd. var. *devicolamensis* Shetty et Vivekananthan (Labiatae).

Bull. Bot. Surv. India 15 : 155. 1973 (1976).

Holotype : India : Kerala : Lower Vagavurrai, Devicolam, 1800 m, Feb. 5, 1970, *Shetty & Vivekananthan* 33420A.

Polycarpaea corymbosa (L.) Lam. var. *longipetala* Srinivasan et Narasimhan (Caryophyllaceae).

J. Econ. Tax. Bot. 10(2) : 347. 1987.

Holotype : Pondicherry, on the road to Marakanam, Jan. 11, 1959, *Subramanyam* 7603.

Polygala angustifolia (Chodat) R. N. Ban. (Polygalaceae).

Bull. Bot. Surv. India 26(1) : 2. 1984 (1985).

P. arillata Buch.-Ham. ex D. Don var. *angustifolia* Chodat
Mem. Soc. Phys. Hist. Nat. Genev. 31(2) : 95. 1893.

Type : India : Sikkim (Darjeeling), Northside of Sinchal, Aug. 1862, 7200 ft., *Anderson* 738.

P. angustifolia (Chodat) R. N. Ban. var. *scandens* (Mukerjee) Giri et R. N. Ban. (Polygalaceae).

Bull. Bot. Surv. India 26 (1) : 3. 1984 (1985).

P. arillata f. *scandens* Mukerjee

Cent. Souv. Madras Herb. 49. 1954 ; Bull. Bot. Soc. Beng. 12 (1 & 2) : 31. 1958.

Holotype : India : Nagaland : Kohima, Naga Hills, 3500-6500 ft. (1060-1980 m), July 1886, *Prain s. n.*

P. arillata Buch.-Ham. ex D. Don var. *chartacea* (Mukerjee) Giri (Polygalaceae).

Bull. Bot. Surv. India 26(1) : 5. 1984 (1985).

P. arillata f. *chartacea* Mukerjee

Cent. Souv. Madras Herb. 50. 1954; Bull. Bot. Soc. Beng. 12 : 32. 1958.

P. arillata f. *meeboldii* Mukerjee Ibid., *pro parte*.

Holotype : India : Kerala : Travancore, 4500 ft. (1500 m), *Bourdillon s. n.*

Polygala arillata Buch.-Ham. ex D. Don var. *purpureescens* Clarke ex Mukerjee (Polygalaceae).

Cent. Souv. Madras Herb. 51. 1954 ; Bull. Bot. Soc. Beng. 12 : 32. 1958; Bull. Bot. Surv. India 26(1) : 4. 1984 (1985).

P. arillata f. *sikkimensis* Mukerjee

Cent. Souv. Madras Herb. 49. 1954 ; Bull. Bot. Soc. Beng. 12 : 31. 1958, *pro parte*, *Syn. nov.*

Holotype : India : Meghalaya : Khasia, Mamloo, 4000 ft., 1220 m, Clarke 45444.

P. arillata Buch.-Ham. ex D. Don var. *revoluta* (Mukerjee) Giri (Polygalaceae).

Bull. Bot. Surv. India 26(1-2) : 5. 1984 (1985).

P. arillata f. *revoluta* Mukerjee

Cent. Souv. Madras Herb. 50. 1954 ; Bull. Bot. Soc. Beng. 12 : 32. 1958.

Holotype : India : Tamil Nadu : Ootacamund, Nov. 1899, Watt s. n.

P. chinensis Linn. var. *hirsuta* Mukerjee (Polygalaceae).

Bull. Bot. Soc. Beng. 12(1-2) : 40. 1958.

Type : Bangladesh : Above kasalong, Chittagong Hill tracts, J. L. Lister s. n.

P. globulifera Dunn (Polygalaceae).

J. Linn. Soc. 35 : 486. 1903; Bull. Bot. Surv. India 26(1) : 6. 1984 (1985).

Isotype : China : Yunan : Forest South and South east of Szemao mts. 1220-1820 m, 1899, Henry 12805A.

P. globulifera Dunn var. *kachinensis* (Mukerjee) R. N. Ban. Comb. et. stat. nov. (Polygalaceae).

Bull. Bot. Surv. India 26(1) : 6. 1984 (1985).

P. arillata f. *kachinensis* Mukerjee

Cent. Souv. Madras Herb. 49. 1954 ; Bull. Bot. Soc. Beng. 12 : 30. 1958.

Holotype, Isotype : Burma : Kachin Hills, Nov. 1897, Shaik Mokim s. n.

P. hyalina Wall. ex Hassk. var. *pusilla* Mukerjee (Polygalaceae).

J. Bombay Nat. Hist. Soc. 51 : 524. 1953 (Emended); Bull. Bot. Soc. Beng. 12(1-2) : 35. 1958.

Type : Burma : Mree Hill, Maym, 1000 m, Sept. 12, 1915, A. Rodger 170.

Polygala jacobii Chandr. (Polygalaceae).

Bull. Bot. Surv. India 9 : 288. 1967 (1968).

Holotype : India : Tamil Nadu : From the banks of the Noyil river, Coimbatore town, 467 m, May 9, 1965, *Chandrabose* 28777A.

P. kyoukmyoungensis Mukerjee (Polygalaceae).

Bull. Bot. Soc. Beng. 12(1-2) : 48. 1958.

Holotype : Upper Burma : Kyaukmyoung, July 1891, *Abdul Huk s n.*

P. pellucida Lace (Polygalaceae).

Kew Bull. 1915 : 395. 1915 ; Bull. Bot. Soc. Beng. 12(1-2) : 35. 1958.

P. carnosa Mukerjee

J. Bombay Nat. Hist. Soc. 51 : 524. 1953.

Isotype : Burma : Maymyo Plateau, alt. 1050 m, *Lace* 4844.

P. ramaswamiana Mukerjee (Polygalaceae).

J. Bombay Nat. Hist. Soc. 53 : 55. 1955; Bull. Bot. Soc. Beng. 12(1-2) : 38. 1958.

Type : S. India : Peermade, *Meebold* 12858.

P. ramaswamiana Mukerjee f. **devicolamensis** Mukerjee (Polygalaceae).

J. Bombay Nat. Hist. Soc. 53 : 56. 1955 ; Bull. Bot. Soc. Beng. 12(1-2) : 38. 1958.

Type : S. India : Devicolam, *A. Meebold* 13185.

var. palinsensis Mukerjee

J. Bombay Nat. Hist. Soc. 53 : 56. 1955; Bull. Bot. Soc. Beng. 12 (1-2) : 38. 1958.

Type : S. India : Kilavare, Upper Palnis, 2000 m, Sept. 20, 1911, *C.E.C. Fischer* 3026.

P. raoi R. N. Ban. et L. K. Ban. (Polygalaceae).

Proc. Indian Acad. Sci. 82B (6) : 218. 1975.

Holotype : India : Tamil Nadu : Vivekanandapuram, flowering and fruiting June 28, 1971, *T. A. Rao* 8216.

Polygonum sarbhanganicum Subba Rao (Polygonaceae).

Bull. Bot. Surv. India 6(1) : 106. 1964 (1965); Bennet, Name Chang. Flowri. Pl. Ind. & Adj. Reg.

P. glandulosum Subba Rao

Bull. Bot. Surv. India 5 : 257. 1964, non Poir 1804, nec. R. Br. 1810.

Isotype : India : Assam : Goalpara, Sandy Banks of the river Sarbhanga about 1200 m, April 24, 1957, *R. S. Rao* 7162B.

Polypogon monspeliensis (Linn.) Desf. var. *indicus* Sunanda Bhattacharya et Jain (Poaceae).

Bull. Bot. Surv. India 25(1-4) : 208. 1983 (1985).

Holotype : India : Punjab : Ferozepore, s. l., s. n.

Paratype : Afghanistan, Herb. *Griffith* 6750.

Polystichum lentum (D. Don) Moore var. *bifurcatum* Biswas and Basu (Dryopteridaceae).

J. Econ. Tax. Bot. 3(3) : 961. 1982.

Holotype : India : Sikkim (East Dist.) : Gangtok, below Honuman top, Nov. 20, 1980, *P. Chakravarty* 818.

Isotype : *P. Chakravarty* 818A.

P. levingei N. C. Nair (Dryopteridaceae).

Bull. Bot. Surv. India 11 : 192. 1969 (1971).

Holotype : Bhutan : Dochula, 3250 m, Sept. 22, 1964, *Sengupta* 436.

Paratypes : Koongma, 3750 m, *Levinge* 14593; W. Bengal : Darjeeling, Oct. 15, 1880, *Levinge* 14594, 14595, 14596; Bhutan : Phullot, 3437 m, *Levinge* s. n. & 8541.

Polyzygus tuberosus Dalzell ex Walpers (Umbelliferae).

Ann. Bot. Syst. 2 : 715. 1852; Kew Bull. 41(1) : 228. 1986.

Paratype : India : Maharashtra : Bombay, Malwan, *Dalzell* s. n.

Pomatocalpa bhutanicum Balakrishnan (Orchidaceae).

J. Bombay Nat. Hist. Soc. 75(1) : 161. 1978.

Holotype : E. Bhutan : Tashiyangtsi, 1870 m, flowered in National Orchidarium, Shillong, June 24, 1965, *Balakrishnan* 41993A.

Portulaca pilosa Linn. ssp. *pilosa* (Portulacaceae).

Sp. Pl. (1753) 445; Blumea 17(2) : 295. 1969.

P. javanensis var. *grisea* Poelln.

Rev. Sudamer. Bot. 7(1943) 273.

Isotype : *P. Javanensis* var. *grisea* : Siam : Pedang, *Yapp* 289.

Potentilla panigraiana Dikshit (Rosaceae).

Bull. Bot. Surv. India 27(1-4) : 107-108. 1985 (1987).

Holotype : Kashmir : Bhurjile, Deosai Plain side, 4575 m, July 31, 1940, *R. R. Stewart* 20137.

Primula flagellaris W. W. Sm. (Primulaceae).

Rec. Bot. Surv. India 4 : 219. 1913; J. Econ. Tax. Bot. 11(1) : 178. 1987.

Holotype : Sikkim : Zemu Valley, *Smith et Cave* 2631.

P. subansirica G. D. Pal (Primulaceae).

J. Bombay Nat. Hist. Soc. 82(3) : 617. 1985.

Holotype : India : Arunachal Pradesh : Subansiri Dist., Begi, alt. 2000 m, April 24, 1980, G. D. Pal 78219.

Prunus bracteopadus Koehne (Rosaceae).

Fedde's Repert. spec. nov. 9 : 33. 1910/1911; Fasc. Fl. Ind. 18 : 27. 1984.

Isotypes : India : Jowai, 1524 m, Oct. 27, 1892, (Sphalm. 93) Dr. King's Collector; Burma Superior, Kachin montes, Sadan, 1981 m, 1900, Shaik Mokim s. n.

P. cerasoides D. Don var. *cerasoides* (Rosaceae).

Prodr. Fl. Nepal 239 (Oct.) 1825; Fasc. Fl. Ind. 18 : 21. 1984.

Isotype : Nepal : Narainhetty, *Hamilton* s. n.

P. phaeosticta (Hance) Maxim. (Rosaceae).

Bull. AC. Imp. Sc. St. Petersb. 3 : 29. 110. 1883; Fasc. Fl. Ind. 18 : 35. 1984.

Isolectotype : Hong Kong: Verticem montium Pakwan, *Hance* 6015.

P. pygeoides Koehne (Rosaceae).

Engl. Bot. Jahrb. 52 : 297. 1915; Fasc. Fl. Ind. 18 : 36. 1984.

Isotype : India : Bihar : Summit of Parusnath, 1220 m, Nov. 15, 1950.

T. Anderson s. n.

P. rufa Steud. ex Hook. f. var. *rufa* (Rosaceae).

Fl. Brit. Ind. 2 : 314. 1878. p. p.; Fasc. Fl. Ind. 18 : 23. 1984.

Isotype : Nepal : Gossainthan, Aug. 1821, Wall. Num. List No. 721.

P. rufa Steud. ex Hook. f. var. *trichantha* (Koehne) Hara (Rosaceae).

J. Jap. Bot. 37 : 99. 1962; Fasc. Fl. Ind. 18 : 24. 1984.

Lectotype selected : Eastern Himalaya : Sikkim, Temperate region, 3000-4000 m, J. D. Hooker & Thomson, p.p.

Pseudodissochaeta assamica (C. B. Cl.) Nayar (Melastomataceae).

J. Bombay Nat. Hist. Soc 65(3) : 559. 1968.

Isotype : India : Muneypore (Manipur), 225 m, Nov. 30, 1885. C. B. Clarke 42323.

Pseudoleskea laevifolia (Mitt.) Jaeg. var. **vasudharensis** Vohra (Leskeaceae).

Bull. Bot. Surv. India 22 : 122. 1980 (1982).

Holotype : India : Uttar Pradesh : Chamoli Dist., Vasudhara, 3500 m, Oct. 7, 1959, *Vohra* 29.

Paratypes : India : Himachal Pradesh : Nachar, Panchhotekhad, 2600 m, Oct. 10, 1960, *Vohra* 448; Tehri-Garhwal, Kidarkanta, 2440-3050 m, Oct. 1879, *Duthie* (Present under the name *Pseudoleskea fuscifolia* C. Muell.).

Pseudostellaria heterophylla (Miq.) f. **nepalensis** Majumdar (Caryophyllaceae).

J. Bombay Nat. Hist. Soc. 64(3) : 589. 1967.

Holotype : Nepal : Nomcha-Thanboche, 3450 m, *Banerjee et Shakya* 5655.

Psychotria balakrishnii Deb et Gang. (Rubiaceae).

Bull. Bot. Surv. India 25 (1-4) : 211. 1983 (1985).

Holotype : North Andaman : Saddle peak, ± 650 m, Dec. 2, 1976, *N. P. Balakrishnan & N. G. Nair* 4785A.

Paratype : Middle Andaman : Tugapur, ± 15 m, July 31, 1974, *N. Bhargava* 1949.

P. bangladeshica Gang. et T. Chakrab. (Rubiaceae).

J. Econ. Tax. Bot. 13(1) : 115. 1989.

Holotype : Bangladesh, *Griffith s. n.* (K. D. No. 3034).

Paratype : India : Assam, *Master s. n.*

P. beddomei Deb et Gang. (Rubiaceae).

J. Econ. Tax. Bot. 5(2) : 479. 1984.

Paratypes : Cheimunjee (?) 1200 m, (4000 ft.) April 9, 1898, (?) 969 ; 900 m (3000 ft.), April, 11, 1898, (?) 960.

P. bhargavii Gang. et T. Chakrab. (Rubiaceae).

J. Econ. Tax. Bot. 13(1) : 113. 1989.

Holotype : India : S. Andamans, Hobdaypur, Sept. 6, 1890, *King's Collector s. n.*, Acc. No. 212666.

Paratypes : Ibid., *King's Collector s. n.*, Acc. Nos. 212667, 212684; Anikhet-Hill Jungle, Sept. 3, 1892, *King's Collector s. n.*, Acc. No. 212611.

P. burkillii Deb et Gang. (Rubiaceae).

Bull. Bot. Surv. India 25 (1-4) : 213. 1983 (1985).

Holotype : India : Arunachal Pradesh : Siang Dist., Tuting to Koppu, 2000-2400 ft. (600-720 m), Nov. 7, 1958, *R. S. Rao* 17415A.

Isotype : Ibid., *R. S. Rao* 17415 B-C.

Paratype : Arunachal Pradesh : Subansiri Dist., Lalik Valley, 2000 ft. (600 m), March 2, 1912, *I. H. Burkhill* 38166.

Psychotria burmanica Deb et Gang. (Rubiaceae).

J. Bombay Nat. Hist. Soc. 81(2) : 439. 1984.

Holotype : Burma : Tavoy : Head waters of Sedi chung 900 m, May 15, 1920, *P. T. Russell* 1806A, in flower.

Isotype : *Ibid.*, *P. T. Russell* 1806B.

Paratypes : *Ibid.*, June 1, 1920, *P. T. Russell* 1832 A, B, C, in fruit, *D. Sterile*; Heinze No. I Camp., 510 m, April 18, 1921, *P. T. Russell* 2027 A,B,C, in flower.

P. calderi Deb et Gang. (Rubiaceae).

J. Econ. Tax. Bot. 5(2) : 477. 1984.

Holotype : Burma : Thabawlich, Cinchona Camp, March 21, 1924.

C. C. Calder s. n. A.

Isotype : *Ibid.*, *C. C. Calder s. n. B.*

Paratypes : *Ibid.*, March 30, 1924, *C. C. Calder s. n.*; Tenasserim, 1929, *K. Biswas* 126.

P. fulva Buch.-Ham. ex Hook. f. var. **brevi-bracteata** Gang. et. T. Chakrab. (Rubiaceae).

J. Econ. Tax. Bot. 13(1) : 116. 1989.

Holotype : India : Meghalaya; Jowai, Khlie Riat Area, N. of Jowai, May 26, 1957, *G. Panigrahi* 2566.

P. kurzii Deb et Gang. (Rubiaceae).

Taxon. 31(3) : 546. 1982.

P. polyneura Kurr.

J. Bot. 4 : 327. 1875.

Type : S. Andaman, *Kurz s. n.*

P. meeboldii Deb et Gang. (Rubiaceae).

J. Bombay Nat. Hist. Soc. 81(2) : 442. 1984.

Holotype : Ceylon (Sri Lanka) : Colombo : March, 1905, *A. Meebold* 2323.

P. nudiflora W. & A. var. **latifolia** Deb et Gang. (Rubiaceae).

Bull. Bot. Surv. India 25(1-4) : 215. 1983. (1985).

Holotype : India : Tamil Nadu : Kanniyakumari Dist., Vallachithodu, Lower Kodayar, ± 550 m, Oct. 7, 1980, *A. N. Henry* 68941.

Paratypes : Tamil Nadu : Kilaviarumalai, ± 500 m, Aug. 29, 1976, *A. N. Henry* 48121; Muthukuzhivayal, ± 1400 m, Aug. 6, 1977, *A. N. Henry* 49621; Tirunelveli Dist., Kannikatti, 830 m, Nov. 10, 1959, *K. M. Sebastian* 9627.

***Psychotria russellii* Deb et Gang. (Rubiaceae).**

J. Bombay Nat. Hist. Soc. 81(2) : 444. 1984.

Holotype : Burma : Tavoy : Heinze No. 1 Camp, 510 m, April 5, 1921, *P. T. Russell* 1943A (in flower).

Isotype : Ibid., *P. T. Russell* 1943B.

Paratypes : Ibid., Nov. 23, 1921, *P. T. Russell* 2186 (in flower); Ibid., April 28, 1921, *P. T. Russell* 2077 (in flower); Tenasserim, March, 1911, *A. Meebold* 14700 (in fruit).

***P. sekharana* K. Ramamurthy et R. Rajan (Rubiaceae).**

J. Bombay Nat. Hist. Soc. 80(3) : 621. 1983.

Holotype : India : Kerala : On the banks of Kannankuzhi river side, Trichur Dist., 250 m, March 22, 1980, *Ramamurthy* 66275.

***P. silhetensis* Hook. f. subsp. *tomentosa* Deb et Gang. (Rubiaceae).**

J. Bombay Nat. Hist. Soc. 80(2) : 466. 1983.

Holotype : India : Assam : North Lakhimpur, Makum, Nov. 21, 1911, *I. H. Burkhill* 35726.

Paratype : North Lakhimpur, Dulong Reserve forest, Nov. 21, 1957, *G. Panigrahi* 11251.

***Pteris silent-valliensis* S. R. Ghosh et R. K. Ghosh (Pteridaceae).**

J. Bombay Nat. Hist. Soc. 79(2) : 385. 1982.

Holotype : India : Kerala : Way to Kuntiphuzha river, Silent Valley, Palghat, April 21, 1980, *Vohra et Ghosh* 56301.

Isotype : Ibid., *Vohra et Ghosh* 56301.

***Pterocarpus macrocarpus* Kurz (Leguminosae).**

Journ. Asiat. Soc. Beng. 43 : 187. 1874; Thoth., Tax. Revi. Dalbergieae P. 214. 1987.

Lectotype designated : Pegu : Tonkyeghat, at 7 Pagodas, *Kurz* 1772.

***P. marsupium* Roxb. subsp. *acuminatus* (Prain) Thoth. (Leguminosae).**

Thoth., Tax. Revi. Dalbergieae P. 227. 1987.

Lectotype designated : Karnataka : N. Kanara, Yellapore, Oct. 1882, *Talbot* 4.

***Pyrenaria burmanica* T. K. Paul et Nayar (Theaceae).**

Journ. Jap. Bot. 64(1) : 8. 1989.

Holotype and Isotype : Burma : Tenasserim, Humatee, 5000 ft., April 10, 1877, *Geo Gallatly* 695.

Raddia angustifolia Soderstrom et Zuloaga (Gramineae).

Brittonia 37(1) : 32. 1985.

Isotype : Brazil, Bahia, 210 m, April 4, 1976, T. R. Soderstrom, G. F. Russell & J. Hage 2110.

Rhamnus collettii M. M. Bhandari et A. K. Bhansali (Rhamnaceae).

J. Econ. Tax. Bot. 5(4) : 913. 1984.

Type : Punjab Himalayas, 1892, H. Collett s. n.

Rhodiola imbricata Edgew. var. **lobulata** Singh et Bhattacharyya (Crassulaceae).

Bull. Bot. Surv. India 25(1-4) : 246-248. 1983 (1985).

Holotype : India : Himachal Pradesh : Sach Pass, 4535 m, July 23, 1964, N. C. Nair 32763A (Male plant).

Rhododendron santapaui A. R. K. Sastry, S. K. Kataki, Peter Cox, Patricia Cox and P. Hutchinson (Ericaceae).

J. Bombay Nat. Hist. Soc. 65(3) : 744. 1968.

Holotype : India : Arunachal Pradesh : Begi, c 1540 m, Subansiri Dist., May 23, 1966, A. R. K. Sastry 45720.

Rhynchoglossum lazulinum A. S. Rao et Joseph (Gesneriaceae).

Bull. Bot. Surv. India 9 : 280. 1967 (1968).

Holotype : India : Arunachal Pradesh : At Krishna, 30 km from Bhallukpong, 1250 m, on the way to Sessa, Kameng Dist., in shady moist places near the way side, on black humus soil, Sept. 10, 1964, Joseph 39716A.

Paratypes : N. L. Bor 18179 Aka Hills, 1936; G. K. Deka s. n. (B. S. I. Acc. No. 29477) Balipara F. T. 13.11.51; A. R. K. Sastry 40738 on way from Palim to Sayata, Subansiri district, N. E. F. A. on hilly slopes of damp forest floor and along small streams with Impatiens, on black clayes soil, 19.11.1964.

Rhynchosia fischeri Satyan. et Thoth. (Fabaceae).

Bull. Bot. Surv. India 28 : 241. 1986 (1988).

Holotype : India : Tamil Nadu : Dimbam, ±1165 m, Coimbatore Dist., Nov. 15, 1906, C. E. C. Fischer 1374

R. hainesiana Satyan. et Thoth. (Fabaceae).

Bull. Bot. Surv. India 28 : 243. 1986 (1988).

Holotype : India : Orissa : Angul forests, March 1917, H. H. Haines 3995.

R. jacobii Chandr. et Shetty (Fabaceae).

Bull. Bot. Surv. India 15 : 139. 1973 (1976).

Holotype : India : Tamil Nadu : Thulukkamparai, Eastern slope of Mahendragiri Hills, Tirunelveli Dist., 425 m, July 25, 1966, Shetty & Chandrabose 27989A.

Rhynchosia meeboldii Satyan. et Thoth. (Fabaceae).

Bull. Bot. Surv. India 28 : 243. 1986 (1988).

Holotype : India : Naga Hills, Shibong, ± 1000 m, Dec. 1907, A. Meebold 7582.

Rhynchostegium calderii Vohra (Brachytheciaceae).

Bull. Bot. Surv. India 22(1-4) : 124. 1980 (1982).

Holotype : Eastern Himalaya, Lalkuti, 2-3 miles on Sonada-Mungpu Road, 2500 m, June 29, 1926, Calder 1102A.

Rinorea longiracemosa (Kurz) Craib (Violaceae).

Fl. Siam. Enum. 90. 1925; Fasc. Fl. Ind. 12 : 7. 1983.

Neotype : Martaban, S. Kurz 41.

Rosa clinophylla var. **parvifolia** (Lindl. ex Prain) Ghora et panigrahi (Rosaceae).

Journ. Jap. Bot. 60(2) : 59. 1985.

R. involucrata var. *parvifolia* Lindl. ex Prain

Journ. Asiat. Soc. Beng. 73(2). No. 5. 202. 1904. p.p.

Syntypes : N. W. Himalaya : Kumaon, Wall. Num. List No. 696; Behar : Chotonagpur, J. J. Wood s. n.

R. hirsuta Chhabbi Ghora et Panigr. (Rosaceae).

Bull. Bot. Surv. India 28 : 117. 1986 (1988).

Holotype : India : Uttar Pradesh : Kumaon, Milanglacier, 3812 m, Aug. 28, 1848, Strachey et Winterbottom 5, Acc. No. 151642.

Isotype : Ibid., Acc. No. 151641.

R. lyellii Lindl. (Rosaceae).

Ros. Monogr. 12. t. 1. 1820; J. Jap. Bot. 60(2) : 59. 1985.

Type : Wallich 682A, Acc. No. 151437.

Rostellularia vahlii (Roth) Nees var. *ruplicola* Ellis (Acanthaceae).

Bull. Bot. Surv. India 11 : 435. 1969 (1971).

Holotype : India : Andhra Pradesh : Srisailam, Nallamalais, Kurnool Dist., July 12, 1963, Ellis 16842A.

Rubia aitchisonii Deb et Malick (Rubiaceae).

J. Bombay Nat. Hist. Soc. 63(3) : 781. 1966.

Holotype : Afghanistan : Badgis : April 29, 1885, Aitchison 338A.

Rubia albicaulis Boiss. var. **stenophylla** Boiss. (Rubiaceae).

Fl. Or. 3 : 19. 1876; Bull. Bot. Surv. India 10(1) : 12. 1968.

Lectotype designated : Schiras, June 17, 1842, Kotschy 543.

R. albicaulis Boiss. var. **stenophylla** Boiss. f. **infundibularis** (Hemsl. & Lace)

Deb et Malick (Rubiaceae).

J. Bombay Nat. Hist. Soc. 63(3) : 783. 1966.

Lectotypes : Baluchistan, Ziarat, J. H. Lace 3945, 3870.

R. angustissima Wall. ex G. Don (Rubiaceae).

Gen. Syst. 3 : 643. 1834; Bull. Bot. Surv. India 10(1) : 5. 1968.

R. charaeifolia Wall. ex G. Don

Lectotype designated : Taong Dong, Nov. 26, 1826, Wall. Cat. 6207.

R. cordifolia Linn. var. **cordifolia** f. **strigosa** Deb et Malick (Rubiaceae).

J. Bombay Nat. Hist. Soc. 63(3) : 782. 1966.

Holotype : Bhutan : Chumbi, Ta esiu Doom, June 17, 1884, G. King 482A.

Isotypes : G. King 482 B & C.

Paratype : India : Arunachal Pradesh : Jeaon to Poom, Kameng F. D., May 21, 1958, G. Panigrahi 16006A.

R. florida Boiss. (Rubiaceae).

[In Kotsch. Pl. Pers. Exs. Feb. 1845 &] Diagn. Pl. Or. Nov. 1(6) : 70. 1845; Bull. Bot. Surv. India 10(1) : 5. 1968.

Lectotype designated : Kotschy 245.

R. sikkimensis Kurz (Rubiaceae).

J. Roy. Asiat. Soc. Beng. 43(2) : 188. 1874; Bull. Bot. Surv. India 10(1) : 12. 1968.

Lectotype designated : India : E. Himalaya : Gunkabaree, Oct. 68, S. Kurz s. n.

R. tibetica Hook. f. (Rubiaceae).

Fl. Brit. Ind. 3 : 204. 1881; Bull. Bot. Surv. India 10(1) : 4. 1968.

Lectotype designated : Western Tibet : 3000-4000 m, Thomson s. n.

Rubus ghanakantae Rolla Rao et Joseph (Rosaceae).

Bull. Bot. Surv. India 12 : 261. 1970 (1972).

Holotype : India : Arunachal Pradesh : Kameng Dist., along the moist soil slopes leading to the ascent of Bomdila, at about 2160 m altitude, nearly 4-5 km from Rahung, June 5, 1957, Rolla 8086A.

Sabia campanulata Wall. var. **kingiana** Nayar et Majumdar (Sabiaceae).

Bull. Bot. Surv. India 24 : 219. 1982 (1983).

Holotype : India : Sikkim Himalaya : G. King s. n., Acc. No. 97399.

Sadiria boweri Dunn (Myrsinaceae).

Bull. Misc. Inform. Roy. Bot. Gard. Kew 1920 : 111. 1920 ; Bull. Bot. Surv. India 16 : 147. 1974 (1977).

Type : India : Arunachal Pradesh : Abor Summit of Bapu, 2080 m, March 7, 1912. I. H. Burkhill R.E.P. 36929.

S. eugenifolia (Wall.) Mez. (Myrsinaceae).

Engl. Pflanzenreich 9 (iv. 236) : 182. 1902; Bull. Bot. Surv. India 16(1-4) : 146. 1974 (1977).

Ardisia eugenifolia [Wall. Cat. No. 2276. 1830] ex A. DC.

Trans. Linn. Soc. 17 : 120. 1834; DC. Prodr. 8 : 130. 1844.

Pimelandra eugenifolia Hook. f. Benth. et Hook. f.

Gen. Plant. 2 : 647. 1876; Hook. f., Fl. Brit. Ind. 3 : 530. 1882.

Tinus eugenifolia O. Ktze.

Rev. Gen. 2 : 974. 1891.

Type : India : Meghalaya : Khasia, Wallich 2276.

S. eugenifolia (Wall.) Mez. var. **burmanica** Nayar et Giri (Myrsinaceae).

Bull. Bot. Surv. India 16 : 146. 1974 (1977).

Holotype : Burma : Myitkyina Dist., 1100 m, May 11, 1919, C. Gilbert Rogers 852.

S. griffithii (Clarke) Mez (Myrsinaceae).

Engl. Pflanzenreich 9(iv. 236) : 183. 1902; Bull. Bot. Surv. India 16 (1-4) : 146. 1974 (1977).

Pimelandra griffithii Clarke

Fl. Brit. Ind. 3 : 531. 1882.

Tinus kurzeana O. Ktze.

Rev. Gen. 2 : 973. 1891.

Ardisia eugenifolia Kurz (non Wall.).

For. Fl. 2 : 111. 1877 et J. Asiat. Soc. Beng. 2 : 225. 1877.

Type : Burma : Sine loc., Griffith 3598.

S. subsessilifolia Nayar et Giri (Myrsinaceae).

Bull. Bot. Surv. India 16 : 144. 1974 (1977).

Holotype : India : Arunachal Pradesh : Abor, Rengging, Feb. 20, 1912, I. H. Burkhill R.E.P. 36657.

Sageraea elliptica (A. DC.) Hook. f. & Thoms. (Annonaceae).

Fl. Ind. 1 : 93. 1855; Fasc. Fl. Ind. 10 : 6. 1982.

Type : Burma : Tavoy : Wall. Num. List No. 6470.

S. laurifolia (Grah.) Blatter (Annonaceae).

J. Bombay Nat. Hist. Soc. 34 : 294. 1931; Fasc. Fl. Ind. 10 : 7. 1982.

Type : Malabar: Concan, Stocks & Law s. n.

S. listeri King var. **andamanica** Chatterjee et Mukerjee (Annonaceae).

Proc. Linn. Soc. 154 : 268. 1941-42; Fasc. Fl. Ind. 10 : 9. 1982.

Type : R. E. P. Regd. No. 12585, Andaman Islands, March 28, 1900.

Santapaua madurensis Bal. & Subr. (Acanthaceae).

J. Indian bot. Soc. 42(3) : 411. 1963.

Holotype : India : Tamil Nadu : Madurai, Nallakulam, Alagar Hills, 200 m, Feb. 12, 1958, K. Subramanyam 5286A.

Isotypes : Ibid., K. Subramanyam 5286D-F.

Sarcanthus williamsonii Reichb. f. (Orchidaceae).

Hamb. Gartenzeit. 21 : 333. 1865 et in Gard. Chron. 1865 : 674; Bull. Bot. Surv. India 9(1-4) : 93. 1967 (1968).

Type : Moulmein, Tenasserim, Parish s. n.

Sauropolis varierii Sivarajan & Indu Balachandran (Euphorbiaceae).

J. Econ. Tax. Bot. 5(4) : 917. 1984.

Holotype : India : Kerala : Malappuram Dist., Kottakkal Arya Vaidya Sala Herbal Garden, Indu AVS 1579. (Original locality of collection not known).

Saussurea sudhansui Hajra (Compositae).

Indian For. 109 (2) : 77. 1983.

Holotype : Uttar Pradesh : Nanda Devi National Park, on the open hill slopes near Dharasi, 4800 m, Aug. 31, 1981, Hajra 73466A.

Saxifraga poluniniana H. Smith var. **mucronata** Bhattacharyya et Viswanathan (Saxifragaceae).

Bull. Bot. Surv. India 15 : 269. 1973 (1976).

Holotype : India : Uttar Pradesh : Kukinakhel, 3200 m, Eastern Chamoli, Garhwal, Western Himalaya, May 1967, U. C. Bhattacharyya 37230.

Schefflera bhutanica R. N. Ban. (Araliaceae).

Bull. Bot. Surv. India 8 : 100. 1966 (1967).

Holotype : Bhutan : 1220 m, Nov. 27, 1876, A. B. Acc. No. 192442.

Isotype : Acc. No. 192430.

S. chandrasekharanii K. Ramamurthy et R. Rajan (Araliaceae).

J. Bombay Nat. Hist. Soc. 79(1) : 163. 1982.

Holotype : India : Kerala : In dense evergreen forest, along, Munnarkumili Road, Idukki Dist., 2000 m, March 26, 1980, *Ramamurthy* 66388.

S. pushpangadanii T. Chakrab. (Araliaceae).

J. Econ. Tax. Bot. 6(2) : 421. 1985.

Holotype : India : Great Nicobar Island, 38 km on East West Road, Feb. 13, 1980, *Dwivedi* 7886A.

Schima notatum (Hack.) A. Camus (Gramineae).

Bull. Mus. Hist. Nat. Paris 27 : 373 (1921) ; Bor, Grass Ceyl. Ind. & Pakis. 218 (1960); J. Econ. Tax. Bot. 12(1) : 77. 1988.

Ischaemum notatum Hack.

DC. Monogr. Phan. 6 : 246 (1889).

Syntype : India : Bunagiri, above Dharahat, 2000 m, *J. F. Duthie* 5057.

S. sulcatum (Hack.) A. Camus (Gramineae).

Bull. Mus. d' Hist. Nat. 27 : 373. 1921; J. Econ. Tax. Bot. 12(1) : 78. 1988.

Ischaemum sulcatum Hack.

DC. Monogr. Phan. 248 (1889).

Type : India : Central Province, Narsingpur, 330 m, *O. Kuntze* s. n.

Schizachyrium sudhansui Singh (Poaceae).

J. Indian. bot. Soc. 60 (3-4) : 359. 1981.

Holotype : India : Karnataka : Raichur-Hyderabad Road, 19th km, C 400 m, Raichur Dist., Nov. 15, 1975, *Singh* 141725A.

Scleroglossum sulcatum (Kuhn) Ald. v. Ros. (Grammitaceae).

Bull. Jard. Buit. Ser. 2.7 : 39. 1912; Bull. Bot. Surv. India 26 : 112. 1984 (1985).

Type : Sri Lanka : Central Province: *Thwaites* 3807.

Sedum phariense H. Ohba (Crassulaceae).

J. Jap. Bot. 48(11) : 328. 1973.

Holotype : Tibet : Phari, July 1879, *Dungboo* s. n., Acc. No. 159421.

Selaginella ganguliana R. D. Dixit (Selaginellaceae).

Bull. Bot. Surv. India 26 (1 & 2) : 104. 1984 (1985).

Holotype : India : Kerala : Trivandrum Aruvikara, 40 m, Dec. 29, 1973, N. C. Nair 51560, Plant numbered-1.

Paratype : Kerala : Kuttikanam, Oct. 22, 1968, N. C. Nair 40008.

S. jainii Dixit (Selaginellaceae).

Bull. Bot. Surv. India 25(1-4) : 225. 1983 (1985).

Holotype : India : Madhya Pradesh : Bilaspur, Siang, Feb. 22, 1972, Panigrahi 16838A (Plant No. 1).

Isotype : Ibid., the rest seven specimens of the sheet, Panigrahi 16838A.

S. keralensis Dixit (Selaginellaceae).

Bull. Bot. Surv. India 27 (1-4) : 123. 1985 (1987).

Isotype : India : Kerala : Kuttikanam, Oct. 22, 1968, Nair 40195.

S. nairii R. D. Dixit (Selaginellaceae).

Bull. Bot. Surv. India 26 (1 & 2) : 106. 1984 (1985).

Holotype : India : Orissa : Jeypore, 830 m, Sept. 17, 1970, N. C. Nair 40637, Plant numbered-II.

Isotype : N. C. Nair 40637A.

Paratype : Orissa : Devighat, 900 m, Sept. 22, 1970, N. C. Nair 40692.

S. nayarii Dixit (Selaginellaceae).

Bull. Bot. Surv. India 27(1-4) : 123. 1985 (1987).

Isotype : India : Kerala : Kuttikanam, Jan. 2, 1971, Nair 818.

S. panigrahii Dixit (Selaginellaceae).

Bull. Bot. Surv. India 25(1-4) : 226. 1983 (1985).

Holotype : India : Madhya Pradesh : Bastar-Kutumsar, Kanger Valley, Feb. 19, 1963, Panigrahi 1119, Acc. No. 20678 (Plant No. I).

Isotypes : Ibid., the rest five specimens of Sheet No. 20678.

Senecio jowaiensis Balakr. (Asteraceae).

Indian For. 101 (11) : 703. 1975.

Holotype : India : Meghalaya : Jowai, Jarain, in shaded moist situation, 1300-1500m, flowers in October, Balakrishnan 47400A.

Isotype : Ibid., Balakrishnan 47400B.

S. pedunculatus Edgew. var. **albus** Ghosh et Bhattacharyya (Compositae).

J. Econ. Tax. Bot. 10(2) : 311. 1987.

Holotype : India : Himachal Pradesh : Lahul, Upper Chenab, Rappe, 2600 m, Aug. 15, 1971, U. C. Bhattacharyya 45329A.

Seshagiria sahyadrica Ansari et Hemadri (Asclepiadaceae).

Indian For. 97(3) : 126. 1971.

Holotype : India : Maharashtra : On the slopes of Sinhagadh Hill, Haveli Taluka, Poona Dist., 1000 m, June 28, 1963, *Ansari* 87750A.

Isotype : *Ansari* 87750D.

Sida mysorensis Wight et Arn. (Malvaceae).

Prodr. Fl. Pen. Ind. Or. 59. 1834; Fasc. Fl. Ind. 19 : 209. 1988.

Isolectotype : India : Mysore, *Wight* 183.

S. rhombifolia Linn. ssp. *rhombifolia* var. *scabrida* (Wight & Arn.) Mast. (Malvaceae).

Fl. Brit. Ind. 1 : 324. 1874; Fasc. Fl. Ind. 19 : 216. 1988.

S. scabrida Wight & Arn.

Prodr. Fl. Pen. Ind. Or. 57. 1834.

Type : Peninsular India, *Herb. Wight* 166.

S. tiagii Bhandari (Malvaceae).

Ann. Arid Zone 16 : 455. 1977; Fasc. Fl. Ind. 19 : 220. 1988.

Holotype : India : Rajasthan : Common on open sandy ground, near Mungari, Barmer Dist., Aug. 28, 1962, *Bhandari* 1570.

Silentvalleya nairii V. J. Nair, P. V. Sreekumar, Vajravelu and P. Bhargavan (Poaceae).

J. Bombay Nat. Hist. Soc. 79(3) : 654. 1982.

Holotype : India : Kerala : Palghat Dist., Silent Valley Dam site, ± 850 m, Oct. 5, 1979, *N. C. Nair* 64268.

Skimmia arunachalensis Goel et Mehrotra (Rutaceae).

J. Bombay Nat. Hist. Soc. 86(2) : 226. 1989.

Paratypes : India : Arunachal Pradesh : Bomdila Camp, April 13, 1957, *G. Panigrahi* 6885A-D.

S. laureola (DC.) Sieb. et Zucc. ex Walpers var. *wattii* Narayanan et Nayar (Rutaceae).

J. Bombay Nat. Hist. Soc. 85(3) : 652. 1988.

Holotype and Isotype : India : Manipur : Sirohipurar, 2425m, April 1882, *Watt* 6459.

Smithia agharkarii Hemadri (Fabaceae).

Indian For. 97(1) : 67. 1971.

Holotype : India : Maharashtra : On the open plateau of Dhak Khilla, ± 1150 m, 27 km west of Junnar, Poona Dist., Sept. 29, 1965, *Hemadri* 107486A.

Smythea velutina (Ridl.) S. P. Ban. et P. K. Mukh. var. *velutina* (Ramnaceae).

Bull. Bot. Surv. India 10(3 & 4) : 251. 1968 (1969).

Ventilago velutina Ridl.

Fl. Mal. Penin. 1 : 467. 1922.

Type : Perak: *Scortechini s. n.*, Acc. No. 87930.

Smythea macrocarpa Hemsl. var. *pubescens* King

J. Asiat. Soc. Beng. 65 : 382. 1896, *pro parte*.

Lectotype : Perak: *Scortechini* 2110.

S. velutina (Ridl.) S. P. Ban. et P. K. Mukh. var. *kingiana* S. P. Ban. et P. K. Mukh. (Ramnaceae).

Bull. Bot. Surv. India 10(3 & 4) : 251. 1968 (1969).

Type : Larut : Perak, 244-305 m, 1885, *King's Collector* 7726.

Sonchus jainii M. Chandrabose, V. Chandrasekaran and N. C. Nair (Asteraceae).

J. Bombay Nat. Hist. Soc. 81(1) : 149. 1984.

Holotype : India : Tamil Nadu : in Konalar, Anamalai, Coimbatore Dist., Nov. 18, 1980, *Chandrabose* 69015.

Sonerila cannanorensis Giri et Nayar (Melastomataceae).

Bull. Bot. Surv. India 27 (1-4) : 86. 1985 (1987).

Holotype : India : Kerala : Cannanore Dist., Chandanathode \pm 825 m, Dec. 18, 1979, *Ramchandran* 65288.

S. devicolamensis M. P. Nayar (Melastomataceae).

J. Bombay Nat. Hist. Soc. 71(3) : 632. 1974.

Holotype : India : Kerala : Devicolam, *A. Meebold* 13771.

S. elegans Wight var. *beddomei* Giri et Nayar (Melastomataceae).

Bull. Bot. Surv. India 27(1-4) : 72. 1985 (1987).

Holotype : India : Kerala : Palghat Hills, 1878, *Beddomei* s. n.

S. gamblei Giri et Nayar (Melastomataceae).

Blumea 31(1) : 235. 1985.

Holotype : India : Tamil Nadu : Nilgiri, *Gamble* 13454.

S. kanniyakumariana Gopalan et Henry (Melastomataceae).

J. Bombay Nat. Hist. Soc. 86(1) : 82. 1989.

Holotype : India : Tamil Nadu: Kannyakumari Dist., Upper Kodayar, March 20, 1984, *R. Gopalan* 77180.

Sonerila neglecta Nayar (Melastomataceae).

Indian For. 95(1) : 28. 1969.

Holotype : Burma : Tavoy : Heinize head waters, 666 m, Nov. 30, 1921, Acc. No. 173299.

S. sahyadrica Giri et Nayar (Melastomataceae).

Bull. Bot. Surv. India 26(3 & 4) : 175. 1984 (1986).

Holotype : India : Kerala : Quilon Dist., Pamba to Analhode, 1050 m, Nov. 30, 1976, Vivekanantran 48344.

S. talbotii Giri et Nayar (Melastomataceae).

Journ. Jap. Bot. 61 (11) : 344. 1986.

Holotype : Karnataka : N. Canara, Aug. 1983. W. A. Talbot 561.

Spatholobus purpureus Benth. ex Prain (Leguminaceae).

J. Asiat. Soc. Beng. 66(1897) : 414; Reinwardtia 10 (2) : 192. 1985.

Butea purpurea (Benth. ex Prain) Blatter

J. Ind. Soc. 8 (1929) : 137.

Type : Talbot 1630.

Sphaerostephanos wynadensis Nayar et Geevarghese (Thelypteridaceae).

Bull. Bot. Surv. India 28 : 137. 1986 (1988).

Type : S. India : Wynad, Attappady, Feb. 13, 1980, 850 m, K. K. Geevarghese CU 29250.

Sphyrantha airyshawii T. Chakrab. & Vasud. (Euphorbiaceae).

J. Econ. Tax. Bot. 5(4) : 959. 1984.

Holotype : India : Andaman-Nicobar Islands : North Andaman Island, Saddle peak, May 10, 1982, Vasudeva Rao 8951A.

Spiraea arunachalensis Panigr. et Purohit (Rosaceae).

Bull. Bot. Surv. India 26(1 & 2) : 83. 1984 (1985).

Holotype : India : Arunachal Pradesh : Kameng Dist., Morsing, 1500m, May 16, 1958, G. Panigrahi 15720.

S. darjeelingensis Panigr. et Purohit (Rosaceae).

Bull. Bot. Surv. India 26(1 & 2) : 83. 1984 (1985).

Holotype : India : W. Bengal : Darjeeling, 2135 m, April 3, 1877, fr., J. S. Gamble 4108C.

S. nayarii Purohit et Panigr. (Rosaceae).

Bull. Bot. Surv. India 26(1 & 2) : 78. 1984 (1985).

Holotype : India : Sikkim : Tiatong above Tallum Samdong, 3660 m, Aug. 1885, Fl., G. King's Collector, Gyalu s. n., Acc. No. 145139,

***Spiraea rhamniphylla* Panigr. et Purohit (Rosaceae).**

Bull. Bot. Surv. India 26 (1 & 2) : 76. 1984 (1985).

Holotype : India : Uttar Pradesh : Kumaon, W. Boogur, 2135 m, May 1848, Fl., R. Strachey & J. E. Winterbottom 3, Acc. No. 145271.

***S. subdioica* Purohit et Panigri. (Rosaceae).**

Bull. Bot. Surv. India 26 (1 & 2) : 86. 1984 (1985).

Holotype : India : Arunachal Pradesh : Kameng Dist., Bomdila, 2780 m, Sept. 14, 1964, fl., J. Joseph 39895.

***S. subrotundifolia* Panigr. et Purohit (Rosaceae).**

Bull. Bot. Surv. India 26(1 & 2) : 88. 1984 (1985).

Holotype : India : Sikkim : Tiampong, Jongri, Aug. 1887, fl., King's Collector s. n. (♂ + ♀ on the same sheet).

Isotypes : Acc. Nos. 145166 & 145168.

***S. tangensis* Purohit et Panigr. (Rosaceae).**

Bull. Bot. Surv. India 26 (1 & 2) : 80. 1984 (1985).

Holotype : India : Sikkim : Tangu, 3965 m, July 1, 1903, fl., F. E. Younghusband s. n., Acc. No. 145145.

***Stellaria nepalensis* Majumdar et Vartak (Caryophyllaceae).**

J. Indian bot. Soc. 50 : 215. 1971.

Holotype : W. Nepal : Pokhra alt. 970 m, May 25, 1968, Vartak 462.

***Sterculia khasiana* Debbarman (Sterculiaceae).**

Fl. Assam 1(1) : 154. 1934; Red Data Book of Indian Pl. 1 : 354. 1987.

Types : India : Meghalaya : Khasi Hills, Oct. 1877, C. B. Clarke 21884 A-D.

***Stigmatodactylus serratus* (Deori) Nageswara Rao (Orchidaceae).**

J. Econ. Tax. Bot. 9(2) : 255. 1987.

Pantlingia serrata Deori

Bull. Bot. Surv. India 20 (1-4) : 175-176 (1978) 1979.

Holotype : India : Meghalaya : Shillong peak, ca 2033 m, Deori 51262A.

***Stixis manipurensis* Deb et Rout (Capparaceae).**

J. Bombay Nat. Hist. Soc. 86(1) : 86. 1986.

Holotype : India : Manipur, Tengnoupul, 1500 m, July 30, 1987, Deb 3549A.

Stixis nayarii Sundararaghavan (Capparaceae).

Bull. Bot. Surv. India 28 : 191. 1986 (1988).

Holotype : Burma : Borwa, Tenasserim Dist., *Meebold* 15406.

S. ovata (Korth.) Hall. f. ssp. *fasciculata* (King) Jacobs (Capparaceae).

Jacobs, Fl. Mal. 1. 5 (1960) 97. f. 29 (map : 2a); Blumea 12(1) : 8. 1963.

Roydsia fasciculata King

Ann. Bot. Gard. Calc. 5. 2 (1896) 121. t. 140A.

S. fasciculata (King) Gagn.

Fl. Gen. I-C. 1 (1908) 201.

Type : *Gallatly* 499.

S. scandens Lour. (Capparaceae).

Fl. Cochinch. (1790) 295; Blumea 12 (1) : 10. 1963.

S. parviflora (Griff.) Pierre.

Bull. Soc. Linn. Paris 1 (1887) 655.

Type : *Griffith* (Burma and Malaya Peninsula) 175.

S. scandens Lour. (Capparaceae).

Fl. Cochinch. (1796) 295; J. Econ. Tax. Bot. 13 (1) : 20. 1989.

Roydsia parviflora Griff.

Not. Pl. As. 4 : 578. 1854.

Isotype : *R. parviflora* : Burma, without locality, *Griffith* s. n.

S. scortechinii (King) Jacobs (Capparaceae).

Fl. Mal. 1.5 (1960) 99. f. 29. 3 (map); Blumea 12(1) : 9. 1963.

Roydsia scortechinii King

J. Asiat. Soc. Beng. 58 ii (1889) 397.

Type : *King's Collector Kunstler* 8464.

Sunipia jainii Hynniewta et Malhotra (Orchidaceae).

J. Indian bot. Soc. 57(1) : 31. 1978.

Holotype : India : Nagaland : Tseninyu, 1200-1500 m, Nov. 14, 1973.

T. M. Hynniewta 56060A.

Swertia sikkimensis Burkill (Gentianaceae).

J. Asiat. Soc. Beng. n. s. 2 : 322. 1906; J. Bombay Nat. Hist. Soc. 81(3) : 731. 1984.

Syntypes : India : Sikkim : Giagong, Sept. 1903, *Prain* s. n.; Himachal Pradesh : Kunawar, 1831, *N. Vicary* s. n., Bhutan : Kungment, Aug. 3, 1884, *Dungboo* 295.

Symplocos nairii A. N. Henry, R. Gopalan & M. S. Swaminathan (Symplocaceae).

J. Bombay Nat. Hist. Soc. 81 (1) : 169. 1984.

Holotype : India : Tamil Nadu : At Muthukuzhivayal, Kanniakumari Dist., 1400 m, Sept. 27, 1980, *Henry* 68830.

S. pulchra Wight subsp. **coriacea** Gopalan et Henry (Symplocaceae).

J. Econ. Tax. Bot. 12 (2) : 481. 1988.

Holotype : Tamil Nadu : Kanniakumari Dist., 11 Forest, from Kodayar to Nalumukku, 1250 m, March 25, 1984, *Gopalan* 81443.

Syzygium chandrasekharanii Chandrab. et Chandras. (Myrtaceae).

J. Bombay Nat. Hist. Soc. 78(2) : 354. 1981.

Holotype : India : Tamil Nadu : Anamalai Hills, Coimbatore Dist., Feb. 15, 1980, *Chandrabose* 65811A.

S. parameswaranii Mohanan et A. N. Henry (Myrtaceae).

J. Bombay Nat. Hist. Soc. 84(2) : 408. 1987.

Holotype : India : Kerala : From the western slopes of Agastymalai, Trivandrum Dist., Alt. ± 1600 m, March 5, 1980, *M. Mohanan* 66051.

Tacca integrifolia Ker-Gawl. (Taccaceae).

Curtis, Bot. Magaz. t. 1488. 1812; Bull. Bot. Surv. India 15(3 & 4) : 189. 1973 (1976).

T. choudhuriana D. B. Deb.

Indian For. 90 : 241-243. 1964.

Type : *T. choudhuriana* : India : Arunachal Pradesh : Bandardia, Subansiri Frontier Division, Sept. 7, 1963, *A. K. Choudhuri* 36457A.

Tadehagi triquetrum (L.) Ohashi subsp. **andamanicum** Balakr. et N. G. Nair (Fabaceae).

Bull. Bot. Surv. India 24 : 29. 1982 (1983).

Holotype : India : North Andaman : Saddle peak, 650-720 m, above m.s.l., scrub forests on open hill slopes, Dec. 2, 1976, *Balakrishnan & Nair* 4798A.

Taeniophyllum andamanicum Balakr. & Bhargava (Orchidaceae).

Bull. Bot. Surv. India 20 : 154. 1978 (1979).

Holotype : India : South Andaman : Baratang Island, Nilambur, ± 25 m, on *Tectona grandis*, July 17, 1977, *Bhargava* 5932A.

Tarennia agumbensis Sundararaghavan (Rubiaceae).

Bull. Bot. Surv. India 10 : 341. 1968.

Holotype : India : Karnataka : Barakana, Agumbe, Shimoga Dist., May 19, 1960, Sundararaghavan 62732A.

Tectaria periya Nayar et Geevarghese (Dryopteridaceae).

Bull. Bot. Surv. India 28 : 136. 1986 (1988).

Type : South India : Wynad, 850 m, Feb. 15, 1982, B. K. Nayar & Periya, K. 11691.

Teramnus hookerianus Ajita Sen (Fabaceae).

J. Bombay Nat. Hist. Soc. 73(1) : 187. 1976.

Holotype : India : W. Bengal : Manbhumi, Purulia Dist., 1866-67, V. Ball s. n., Acc. No. 125331.

Paratypes : India : Madras : M. S. Rameswami 1015, Acc. No. 125305; Bengal: Bankura, Bishnupur, Koch Birai Canal, M. M. Sanyal 532.

Tetracera akara (Burm. f.) Merr. (Dilleniaceae).

Philip. J. Sci. 19 : 366. 1921; Hoogl. in Reinwardtia 2 : 208. 1953.

T. sylvestris Ridl.

J. Str. Br. R. A. S. 54 : 8. 1910.

Isotype : *T. sylvestris* : Ridley 6179.

T. arborescens Jack (Dilleniaceae).

Mal. Misc. 1(5) : 45. 1820; Reinwardtia 2 : 217. 1953.

T. Laevigata Miq.

Fl. Ind. bat. 1(2) : 8. 1859.

Isotype : *T. laevigata* : Sumatra : Teysmann 457 HB.

T. asiatica (Lour.) Hoogl. (Dilleniaceae).

Fl. Mal. Ser. 1. 4 : 143. 1951; Hoogl. in Reinwardtia 2 : 193. 1953.

Delima sarmentosa var. *glabra* Hook. f. & Thoms.

Fl. Ind. 1 : 61. 1855.

Isolectotype : *D. sarmentosa* var. *glabra* : Bangladesh : Sylhet, Feb. 1829, Wallich 6632.

T. fagifolia Bl. (Dilleniaceae).

Bijdr. 1 : 4. 1825; Hoogl. in Reinwardtia 2 : 219. 1953.

T. sumatrana Miq.

Fl. Ind. bat. Suppl. 1 : 618. 619. 1861.

T. obliquinervia Elm.

Leafl. Philip. Bot. 7: 2621. 1915.

Isotype : *T. sumatrana* : Sumatra : Teysmann 4454 HB.

Isotype : *T. obliquinervia* : Mindanao, Elmer 13862.

Tetracera korthalsii Miq. (Dilleniaceae).

Ann. Mus. bot. Lugd. Bat. 4 : 75. 1868; Hoogl. in Reinwardtia 2 : 213. 1953.

T. subrotunda Elm.

Leafl. Philip. Bot. 5 : 1771. 1913.

Isotype : *T. subrotunda* : Elmer, Palawan, 13048.

T. macrophylla Wall. ex Hook. f. & Thoms. (Dilleniaceae).

Hoogl. in Reinwardtia 2 : 215. 1953.

T. macrophylla Wall. ex Hook. f. & Thoms.

Fl. Ind. 1 : 63. 1855.

T. grandis King

J. Asiat. Soc. Beng. 58(2) : 363. 1889.

Isotype : *T. macrophylla* : Wallich 6628.

Holotype : *T. grandis* : Perak : Scortechini s. n.

T. maingayi Hoogl. (Dilleniaceae).

Fl. Mal. Ser 1. 4 : 144. 1951; Hoogl. in Reinwardtia 2 : 199. 1953.

Delima laevis Maing.

J. Asiat. Soc. Beng. 58(2) : 362. 1889, non *T. laevis* vahl

Holotype : Malacca : April 10, 1867, Maingay 1570.

Thelypteris multilineata (Wall. ex Hook.) Morton var. **bhutanica** N. C. Nayar (Thelypteridaceae).

Bull. Bot. Surv. India 11 : 193. 1969 (1971).

Holotype : Bhutan : Nov. 1964, Deb. 179.

Themeda sabarimalayana Sreekumar et Nair (Poaceae).

Bull. Bot. Surv. India 29(1-4) : 127. 1987 (1989).

Holotype : India : Kerala: Pathanamthitta Dist., on way to Sabovi-mala, ± 450 m, Dec. 23, 1980, P. V. Sreekumar 69433.

Thrixspermum muscaeflorum A. S. Rao & Joseph (Orchidaceae).

Bull. Bot. Surv. India 11 : 204. 1969 (1971).

Holotype : India : Between umran and umsaw, beside the Gauhati Shillong high way, July 23, 1966, Joseph 45621A.

Thuidium contortulum (Mitt.) Jaeg. var. **mussooriense** Vohra (Thuidiaceae).

Bull. Bot. Surv. India 22 : 124. 1980 (1982).

Holotype : India : Uttar Pradesh : Mussoorie, 1830-2135 m, Aug. 1895, Duthie s. n.

Thuidium squarrosulum Ren. et Card. var. **scabriisetum** Vohra (Thuidiaceae).

Bull. Bot. Surv. India 22 : 124. 1980 (1982).

Holotype : Bhutan : Lumichawa, 2500 m, Sept. 2, 1965, *Vohra* 2435.

Paratype : India : W. Bengal : Darjeeling, Lloyd's Botanic Garden, 2000 m, May 18, 1972, *Vohra* 3746.

Thymus serpyllum Linn. var. **album** Ghosh et Bhattacharyya (Labiatae).

J. Econ. Tax. Bot. 10(2) : 310. 1987.

Holotype : India : Himachal Pradesh : Lahul, Chetru, 3500 m, July 20, 1972, *U. C. Bhattacharyya* 48766A.

Tinomiscium nicobaricum Balakr. (Menispermaceae).

Bull. Bot. Surv. India 24 : 56. 1982 (1983).

Holotype : India : Great Nicobar : 25 km on East-west Road, near Galathea river, inland hill forests, ca 75m above m.s.l., June 16, 1977, *Balakrishnan* 5787A.

Tribulus rajasthanensis Bhandari et Sharma (Zygophyllaceae).

Bot. Not. 129. 367. 1976; Bull. Bot. Surv. India 24 : 162. 1982 (1983).

Type : India : Rajasthan : Jodhpur Dist., Oct. 3, 1959, *Bhandari* 537.

T. subramanyamii P. Singh, Giri et V. Singh (Zygophyllaceae).

Bull. Bot. Surv. India 25(1-4) : 197-198. 1983 (1985).

Holotype : India : Tamil Nadu : Coimbatore, on the way to var-apalayam, 666 m, July 25, 1956, *K. Subramanyam* 390.

Isotype : Ibid.

Trichocolea udarii Singh (order-Jungermanniales).

Bull. Bot. Surv. India 25(1-4) : 177-180. 1983 (1985).

Holotype : India : Meghalaya : Jarain, Jaintea Hills, ca 1450 m, April 24, 1981, *D. K. Singh* 11012.

Trichodesma decurrens Banerjee (Boraginaceae).

Indian For. 92(1) : 56. 1966.

Holotype : Burma : Zeaing, Jan. 1905, *Shaik Mokim*, 1112A.

Isotype : *Shaik Mokim* 1112B.

Trichosanthes bracteata (Lamk.) Voigt var. **tomentosa** (Heyne ex Clarke) Chakravarty (Cucurbitaceae).

Rec. Bot. Surv. India 17 (1) : 47. 1959; Fasc. Fl. Ind. 11 : 110. 1982.

Type : Peninsula Indiae Orientalis, *Herb. Wight* 1134.

Trichosanthes dicaelosperma Clarke (Cucurbitaceae).

Fl. Brit. Ind. 2 : 609. 1879; Fasc. Fl. Ind. 11 : 113. 1982.

Type : India : Meghalaya : Khasia, *J. D. Hooker & Thomson s. n.*

T. himalensis Clarke var. **glabrior** Clarke (Cucurbitaceae).

Fl. Brit. Ind. 2 : 608. 1879; Fasc. Fl. Ind. 11 : 115. 1982.

Type : India : Meghalaya : Khasia, *J. D. Hooker & Thomson 9.*

T. himalensis Clarke var. **sikkimensis** (Kundu) Thoth. (Cucurbitaceae).

Bull. Bot. Surv. India 2(1 & 2) : 169. 1960; Fasc. Fl. Ind. 11 : 116. 1982.

Type : India : Sikkim : *Clarke 36925A.*

T. listeri Chakravarty (Cucurbitaceae).

J. Bombay Nat. Hist. Soc. 50 : 895. 1952; Fasc. Fl. Ind. 11 : 117. 1982.

Type : Bangladesh : Chittagong hill tracts, *J. L. Lister 349.*

T. tomentosa Chakravarty (Cucurbitaceae).

J. Bombay Nat. Hist. Soc. 50 : 894. 1952; Fasc. Fl. Ind. 11 : 121. 1982.

Type : India : Nagaland : Kohima to Nerhema, *Watt 11640.*

Trigonostemon aurantiacus (Kurz ex Teijsm. & Binnend.) Boerl. var.

rubriflorus Balakr. & T. Chakrab. (Euphorbiaceae).

J. Econ. Tax. Bot. 5(1) : 169. 1984.

Holotype : India : Andaman-Nicobar Islands : Havelock Island, without locality, ±25 m, May 15, 1974, *Ansari 1361A.*

T. birmanicus T. Chakrab. & Balakr. (Euphorbiaceae).

J. Econ. Tax. Bot. 5(1) : 175. 1984.

Holotype : Burma : Upper Chindwin Dist., Numpakom drainage, April 29, 1911, Collector uncertain (Chin?) 5849.

T. nicobaricus T. Chakrab. (Euphorbiaceae).

J. Econ. Tax. Bot. 5(1) : 203. 1984.

Holotype : India : Andaman-Nicobar Islands : Great Nicobar Island, 9 km on East West Road, May 6, 1981, *Dwivedi 8521A.*

T. sunirmalii T. Chakrab. et Balakar. (Euphorbiaceae).

J. Econ. Tax. Bot. 5(1) : 179. 1984.

Holotype : Burma : Tenesserim, Nimchaung, ± 60 m alt., March 31, 1930, *Biswas 22.*

Paratypes : Lepokechung, Feb. 11, 1931, *Biswas 842*; Mergui, Cinchona Camp, March 21, 1924, *Calder s. n.*, Acc. No. 411835.

Trigonostemon viridissimus (Kurz) Airy Shaw var. **chatterjii** (Deb et Deka)

Balakr. et T. Chakrab. (Euphorbiaceae).

J. Econ. Tax. Bot. 5(4) : 967. 1984.

Holotype : India : Meghalaya : Jaintea Hills Dist., Dawki, May 4, 1943, *Deka* 19A.

T. viridissimus (Kurz) Airy Shaw var. **confertifolius** Balakr. et N. G. Nair (Euphorbiaceae).

Bull. Bot. Surv. India 24 : 36. 1982 (1983).

Holotype : India : North Andaman : Saddle peak, 500-700 m, open scrub forests, Dec. 1. 1976, *Balakrishnan & Nair* 4773A.

Trigonotis caespitosa Ban. (Boraginaceae).

Bull. Bot. Surv. India 8(3 & 4) : 322. 1966 (1967).

Holotype : India : Sikkim : Goraphu chu, Llonakh, 4785 m, Aug. 6, 1909, *Smith & Cave* 2220, Acc. No. 308558.

T. hookeri Benth. ex Clarke (Boraginaceae).

Fl. Brit. Ind. 4 : 173. 1883; Bull. Bot. Surv. India 8(3 & 4) : 327. 1966 (1967).

Isotype : India : Mount Khasia, 1829 m, *J. D. Hooker & T. Thomson s. n.* Acc. No. 307884.

T. microcarpa (DC.) Benth. ex Clarke (Boraginaceae).

Fl. Brit. Ind. 4 : 172. 1883; Bull. Bot. Surv. India 8(3 & 4) : 320. 1966 (1967).

Eritrichium microcarpum DC.

Prodr. 10 : 123. 1846.

T. peduncularis var. *microcarpa* (DC.) Brand

Engl. Pflanzenr. 97 : 198. 1931.

Isotype : Nepal : *Wallich* 928.

T. multicaulis (DC.) Benth. ex Clarke var. *cavei* Ban. (Boraginaceae).

Bull. Bot. Surv. India 8(3 & 4) : 321. 1966 (1967).

Holotype : India : Sikkim : Eumtso La (Yumchhola), 4267 m, July 17, 1909, *Smith & Cave* 1607A.

Isotype : Ibid., *Smith & Cave* 1607B.

T. ovalifolia (Wall.) Benth. ex Clarke (Boraginaceae).

Fl. Brit. Ind. 4 : 172. 1883; Bull. Bot. Surv. India 8(3 & 4) : 326. 1966 (1967).

Myosotis ovalifolia Wall.

Roxb. Fl. Ind. ed. Carey & Wall. 2 : 9. 1824.

Eritrichium ovalifolium (Wall.) A. DC.

DC. Prodr. 10 : 123. 1846.

Isotype : Nepal : Gossainthan, *Wallich* 927.

Trigonotis rotundifolia Benth. ex Clarke (Boraginaceae).

Fl. Brit. Ind. 4 : 172. 1883; Bull. Bot. Surv. India 8(3 & 4) : 324. 1966 (1967).

Myosotis rotundifolia Benth.

Royle III. Bot. Him. 395. 1836.

Eritrichium rotundifolium (Benth.) DC.

Prodr. 10 : 123. 1846.

Isotype : India : Kumaon, *Wallich* 930.

T. smithii Ban. (Boraginaceae).

Bull. Bot. Surv. India 8 : 325. 1966 (1967).

Holotype : India : Sikkim : Llonok, 4419 m, July 31, 1909, *Smith & Cave* 1811A, Acc. No. 308538.

Isotypes : *Smith & Cave* 1811 B-D, Acc. Nos. 308553, 308554, 308557.

T. tibetica (Clarke) Johnst. (Boraginaceae).

Contr. Gray Herb. Harv. 75 : 48. 1925; Bull. Bot. Surv. India 8(3 & 4) : 323. 1966 (1967).

Isotype : Tibet: *Hook. f. & Thoms.*, *Eritrichium* sp. n. 17.

T. tibetica var. **minor** (Clarke) Johnst. (Boraginaceae).

Contr. Gray Herb. Harv. 75 : 48. 1925; Bull. Bot. Surv. India 8(3 & 4) : 324. 1966 (1967).

Isotype : India : Sikkim : Alpine Sikkim, 4572 m, *Hook. f. & Thoms.*, *Eritrichium* sp. n. 15, Acc. No. 307585.

Tripogon anantaswamianus Sreekumar, V. J. Nair et N. C. Nair (Gramineae).

Bull. Bot. Surv. India 25 (1-4): 185-187. 1983 (1985).

Holotype : India : Kerala : Eravikulam Sanctuary, Idukki Dist., 2000 m, Nov. 15, 1980, *P. V. Sreekumar* 69432.

Triumfetta tungarensis Billore (Tiliaceae).

J. Econ. Tax. Bot. 3(2) : 621. 1982.

Holotype : India : Maharashtra : In the forest undergrowth of Mazivilli R. F., near Tungar Hill Mandvi range, Thana Dist., Nov. 23, 1968, *K. V. Billore* 115873A.

Trivalvaria dubia (Kurz) Sinclair (Annonaceae).

Gard. Bull. Singapore 24 : 47. 1955; Fasc. Fl. Ind. 10 : 11. 1982.

Type : India : S. Andaman, *S. Kurz s. n.*

T. kanjilalii D. Das (Annonaceae).

Bull. Bot. Surv. India 10 : 263. 1968; Fasc. Fl. Ind. 10 : 11. 1982.

Type : India : Meghalaya : Khasia Hills, wah-Maw-long, Oct. 20, 1917, *U. N. Kanjilal* 7280.

Urginea indica (Roxb.) Kunth (Liliaceae).

Enum. Pl. 4 : 333. 1843; J. Bombay Nat. Hist. Soc. 84(2) : 411. 1987.

Scilla indica Roxb.

Fl. Ind. 2 : 148. 1832.

S. coromandelliana Roxb.

U. govindappae Boraiah et Fatima

Bull. Bot. Surv. India 12 : 128. 1970.

Type : *U. govindappae* : *Boraiah et Fatima* 601.

U. nagarjunae Hemadri et Swahari (Liliaceae).

Ancient Sci. Life 2 : 105. 1982; J. Bombay Nat. Hist. Soc. 84(2) : 411. 1987.

Holotype : *Hemadri* 3001A.

Paratype : *Hemadri* 2925A.

U. rajii (Ansari) Deb et Dasgupta (Liliaceae).

J. Bombay Nat. Hist. Soc. 84(2) : 412. 1987.

Drimia rajii Ansari

J. Bombay Nat. Hist. Soc. 78(3) : 572. 1981.

Holotype : *D. rajii* : India : Maharashtra : Pune Dist., Dive Ghat, March 15, 1970, *Ansari* 104878A.

Isotype : Ibid., *Ansari* 104878D.

Utricularia khasiana Joseph et Mani (Lentibulariaceae).

Bull. Bot. Surv. India 25(1-4) : 192-194. 1983 (1985).

Holotype : India : Meghalaya : Shillong, Khasi Hills, 1496 m, Oct. 1, 1980, *Joseph* 76944A.

U. malabarica Janarthanan et Henry (Lentibulariaceae).

J. Bombay Nat. Hist. Soc. 86(1) : 84. 1989.

Holotype : India : Kerala, Kasaragod, Mulleriya, Aug. 21, 1985, *M. K. Janarthanan* 82924.

***Utricularia nayarii* M. K. Janarthanan et A. N. Henry (Lentibulariaceae).**

Bull. Bot. Surv. India 28 : 195. 1986 (1988).

Holotype : India : Kerala : Idduki Dist., Rajamallay near Munnar, Dec. 12, 1985, *M. K. Janarthanan* 82963.

Paratype : Ibid., Lockhart gap near Munnar, 1675 m, Oct. 12, 1963, *K. M. Sebastine* 17543.

***U. polygaloides* Edgeworth (Lentibulariaceae).**

Proc. Linn. Soc. 1 : 351. 1847; Basak in Taxon 25 : 189. 1976.

Neotype : India : W. Bengal : Burdwan, Jan. 1837, *Mrs. Skepwith s. n.*, Acc. No. 330152.

***U. sampathii* Subr. et Yog. (Lentibulariaceae).**

J. Indian bot. Soc. 60 : 123. 1981.

Holotype : India : Karnataka : Dhyan Ashram, by the side of small stream near Bannerghatta, 12°50'N & 77°45'E, about 10 kms from Bangalore, 900 m, Nov. 3, 1976, *Subramanyam* and *Yoganarasimhan* 123.

***U. tayloriana* Joseph et Mani (Lentibulariaceae).**

Bull. Bot. Surv. India 24 : 108. 1982 (1983).

Holotype : India : Meghalaya : 20th km on the Shillong-Pynursla Road, ca 700 m, Khasi and Jaintia Hills, Aug. 26, 1980, *Joseph* 73568A.

***Uvaria andamanica* King (Annonaceae).**

J. Asiat. Soc. Beng. 61 : 2 : 21. 1892 et Ann. Roy. Bot. Gard. Cal. 4 : 29. t. 25. 1893; Fasc. Fl. Ind. 10 : 13. 1982.

Lectotype : India : Andaman : Port Blair, April, 1884, *King's Collector* 207.

***U. eucinata* Bedd. ex Dunn (Annonaceae).**

Kew Bull. 1914 : 182. 1914; Fasc. Fl. Ind. 10 : 15. 1982.

Type : Russelkonda Hills, *Beddome* 50.

***U. hamiltonii* Hook. f. & Thoms. (Annonaceae).**

Fl. Ind. 1 : 96. 1855 et Hook. f., Fl. Brit. Ind. 1 : 48. 1872; Fasc. Fl. Ind. 10 : 15. 1982.

Type : Mongir Hills, *Hamilton ex Wallich* Num. List 6485E.

***U. hamiltonii* Hook. f. & Thoms. var. *kurzii* King (Annonaceae).**

Ann. Roy. Bot. Gard. Cal. 4 : 17. 1893; Fasc. Fl. Ind. 10 : 16. 1982.

Type : South Andaman : July 9, 1884, *King's Collector* 338.

***Uvaria hookeri* King (Annonaceae).**

Ann. Roy. Bot. Gard. Cal. 4 : 28. t. 22. 1893; Fasc. Fl. Ind. 10 : 16. 1982.

Type : South India : Wall. Num. List 6473C.

***U. lurida* Hook. f., & Thoms. var. *lurida* (Annonaceae).**

Fl. Ind. 1 : 101. 1855; et Hook. f., Fl. Brit. Ind. 1 : 50. 1872; Fasc. Fl. Ind. 10 : 17. 1982.

Type : Northern slopes of Khasia Hills, 2000 ft., *Hook. f. Thomson s. n.*

***U. lurida* Hook. f. & Thoms. var. *sikkimensis* King (Annonaceae).**

Ann. Roy. Bot. Gard. Cal. 4 : 29. t. 23. 1893; Fasc. Fl. Ind. 10 : 17. 1982.

Type : Sikkim, Mongpoo, Peak, 1981, *King s. n.*

***Vanda jainii* Chauhan (Orchidaceae).**

J. Econ. Tax. Bot. 5(4) : 977. 1984.

Holotype : India : Meghalaya : West Khasi Hills, Sonapahar, March 25, 1981, *Chauhan* 70887A.

***Vernonia anaimudica* Shetty et Vivek. (Compositae).**

Bull. Bot. Surv. India 12 : 266. 1970 (1972).

Holotype : India : Kerala : Kottayam Dist., Rajamallay, Devicolam, Feb. 2, 1970, *Shetty & Vivekananthan* 31786A.

***V. peninsularis* (Clarke) Clarke ex Hook. f. var. *kodayarensis* Henry et Gopalan (Compositae).**

J. Econ. Tax. Bot. 11(1) : 233. 1987.

Holotype : India : Tamil Nadu : Kannyakumari, Muthukuzhivayal, Feb. 18, 1983, *Henry* 77074.

***V. andamanica* Balakr. et N. G. Nair (Rubiaceae).**

Bull. Bot. Surv. India 24 : 32. 1982 (1983).

Holotype : India : North Andaman : Saddle peak, 500-700 m, open scrub jungles, Dec. 1, 1976, *Balakrishnan & Nair* 4768A.

***Viola betonicifolia* J. Sm. ssp. *jaunsarensis* (W. Beck.) Hara (Violaceae).**

J. Jap. Bot. 49(5) : 133. 1974; Fasc. Fl. Ind. 12 : 15. 1983.

Holotype : India : Konain, Jaunsar, *J. F. Duthie* 12963.

***V. canescens* Wall. (Violaceae).**

Fl. Ind. 2 : 450. 1824; Fasc. Fl. Ind. 12 : 19. 1983.

Isotype : Nepal : Wall. Num. List No. 1442.

Viola fedtschenkoana W. Beck. var. **muzaffarabadensis** W. Beck. (Violaceae).

Beih. Bot. Centralbl. Abt. 2. 40 : 59. 1923; Fasc. Fl. Ind. 12 : 22. 1983.

Holotype : India : Jammu & Kashmir : Muzaffarabad, *Inayat* 21166.

V. glaucescens Oudem. (Violaceae).

Miq. Ann. Mus. Bot. Lugd. Bat. 3 : 74. 1867; Fasc. Fl. Ind. 12 : 22. 1983.

Isotype : India : Sikkim : *J. D. Hooker s. n.*

V. hossei W. Beck. (Violaceae).

Beih. Bot. Centralbl. Abt. 2. 34 : 257. 1916; Fasc. Fl. Ind. 12 : 25. 1983.

Isotype : China : Yunnan, *Henry* 11768.

V. indica W. Beck. f. **barbata** (W. Beck.) Banerjee et Pramanik (Violaceae).

Fasc. Fl. Ind. 12 : 26. 1983.

Holotype : India : Matiana, 2700 m, Simla Hill State, *J. H. Burkhill* 28658.

V. jordani Henry [Cat. Pl. var. 169. 1853] var. **falconeri** (Hook. f. & Thoms.) W. Beck. (Violaceae).

Beih. Bot. Centralbl. Abt. 2. 34 : 380. 1917; Fasc. Fl. Ind. 12 : 27. 1983.

Isotype : India : Kashmir : *Falconer s. n.* EIC 193.

V. kunawarensis Royle (Violaceae).

111. Himal. Bot. 74. t. 18. f. 3. 1839; Fasc. Fl. Ind. 12 : 27. 1983.

Isotype : N. W. India, *Hb. Royle s. n..*

V. rupestris F. W. Schmidt. var. **himalayensis** (W. Beck.) Banerjee et Pramanik (Violaceae).

Fasc. Fl. Ind. 12 : 33. 1983.

Holotype : India : Kashmir : *Falconer s. n.* sub. EIC 191.

V. sikkimensis W. Beck. (Violaceae).

Beih. Bot. Centralbl. Abt. 2. 34 : 260. 1916; Fasc. Fl. Ind. 12 : 34. 1983.

Type : Sonada, 2500 m, *J. S. Gamble* 10318.

V. thomsonii Oudem. (Violaceae).

Miq. Ann. Musc. Bot. Lugd. Bat. 3 : 74. 1867; Fasc. Fl. Ind. 12 : 35. 1983.

Isotype : India : Sikkim, *J. D. Hooker s. n.*

Viola yunnanensis W. Beck. et Boiss. (Violaceae).

Bull. Herb. Boiss. 8 : 740. 1908; Fasc. Fl. Ind. 12 : 37. 1983.

Isotype : China : Yunnan, A. Henry 10685.

Vittaria arunachalensis R. D. Dixit (Vittariaceae).

J. Econ. Tax. Bot. 3(3) : 959. 1982.

Holotype : India : Arunachal Pradesh : Kameng Dist., Dirang, April 7, 1973, R. S. Rao 53927.

V. elongata Sw. var. **angustifolia** Holttum [Rev. Fl. Malaya 2 : 614. 1955.

descr. angl. nom. illeg.] ex Balakr. (Vittariaceae).

Bull. Bot. Surv. India 22 : 137. 1980 (1982).

Holotype : N. Nicobars; Car Nicobar Isl., N. G. Nair 2896A.

Wallichia mooreana S. K. Basu (Palmae).

Taiwania 28 : 146. 1983.

Holotype : S. China : Yunnan : Ponesee, March 17, 1868, D. J. Anderson s. n., Acc. No. 492466.

Paratypes : S. China : Yunnan : Szemao Mts., 1899, A. Henry 12331, Acc. Nos. 492505, 492507.

Wendlandia scabra Kurz var. **tenasseriensis** Nayar et Malick (Rubiaceae).

J. Bombay Nat. Hist. Soc. 75(1) : 156. 1978.

Holotype : Burma : Young Zalum River, Tenasserim, March 27, 1877, Gallatly 496A.

Isotypes : Gallatly 496B-E.

Zanonia indica L. var. **pubescens** Cogn. (Cucurbitaceae).

DC. Monogr. Phan. 3 : 927. 1881; Fasc. Fl. Ind. 11 : 126. 1982.

Type : India : Khasia Hills, Griffith 2521.

Zanthoxylum burkillianum Babu (Rutaceae).

Bull. Bot. Surv. India 16 : 55. 1974 (1977).

Holotype : India : Meghalaya : Abor Hills, Rotung on hill side over Dihang, 396 m, Burkhill 37598A.

Isotype : Ibid., Burkhill 37598B.

Z. pseudoxiphyllum Babu (Rutaceae).

Bull. Bot. Surv. India 16 : 66. 1974 (1977).

Holotype : India : Manipur : Sirohfurai, 2420 m, Watt 6452A ♂.

Isotype : Ibid., Watt 6452 B ♂.

Zanthoxylum tomentellum Hook. f. (Rutaceae).

Fl. Brit. Ind. 1 : 493. 1875 emend. Babu ; Bull. Bot. Surv. India 16 : 68. 1974 (1977).

Fagara tomentella (Hook. f.) Hand.-Mazz.

syn. sin. 7 : 624. 1933.

Lectotype : *Griffith* 743 ♀

Zenkeria jainii N. C. Nair, P. V. Sreekumar and V. J. Nair (Poaceae).

J. Bombay Nat. Hist. Soc. 78(2) : 352. 1981.

Holotype : India : Kerala : Eravikulam Sanctuary, ± 2000 m, Idukki Dist., Aug. 20, 1980, *P. V. Sreekumar* 68419.

Z. sebastinei Henry & Chandr. (Poaceae).

Bull. Bot. Surv. India 15 : 142. 1973 (1976).

Holotype : India : Tamil Nadu : Agastyamalai, Tirunelveli Dist., Aug. 26, 1963, *Henry* 17325A.