

References for *Colocasia* Schott

028.009

Genus ID : A-

- 1.Sharma AK, Mukherjee D.1959.Further investigations on the cytology of some members of Aroids.Proc. 46th Indian Science Congress .Part-3.356.
- 2.Cao LM, Long CL.2003.Colocasia bicolor (Araceae), a new species from Yunnan.Ann. Bot. Fennici.40.283-286.
- 3.Cao LM, Long CL.2004.Chromosome numbers of eight Colocasia taxa and karyotypes of five species occurring in China.Acta Botanica Yunnanica.26.310-316.
- 4.Wang GY, Zhang XM, Qian M, Hu XY, Yang YP.2017.Chromosome number and genome size variation in Colocasia (Araceae) from China .J. Plant Res.DOI 10.1007/s10265-017-0959-8.
- 5.Yen DE, Wheeler JM.1968.Introduction of Taro into the Pacific: The indications of the chromosome numbers.Ethnology.7.259-267.
- 6.Marchant CJ.1971.Chromosome variations in Araceae II. Richardieae to Colocasieae.Kew Bulletin .25.27-56.
- 7.Hotta A.1971.Study of family Araceae. General remarks.Jap. J. Bot.20.269-310.
- 8.Hsu CC.1972.Preliminary chromosome studies on the vascular plants of Taiwan (V). Cytotaxonomy on some monocotyledons.Taiwania .17 .48-65.
- 9.Sastrapradja S, Rijanti AM.1972.On the cytology of Javanese Colocasia.Ann. Bogor.5.173-177.
- 10.Ramachandran K.1978.Cytological studies on South Indian Araceae.Cytologia .43.289-303.
- 11.Kuruvilla KM, Singh A.1981.Karyotypic and electrophoretic studies on taro and its origin.Euphytica.30.405-413.
- 12.Zhang G, Yang Z.1984.Studies on chromosome numbers of main cultivars of Colocasia esculenta in China.Acta Horticulturae Sincae.11 (3).187-190.
- 13.Tanimoto T, Matsumoto T.1986.Variations of morphological characters and isozyme patterns in Japanese cultivars of Colocasia esculenta Schott and C. gigantea Hook.Japanese Journal of Breeding.36.100-111.
- 14.Coates DJ, Yen DE, Gaffey PM.1988.Chromosome variation in Taro, Colocasia esculenta : Implications for origin in the Pacific.Cytologia.53.551-560.
- 15.Sreekumari MT, Mathew PM.1989.Karyomorphology of six cultivars of taro (Colocasia esculenta Schott).New Botanist.16.127-135.

- 16.Okada H, Hambali GG.1989.Chromosome behaviors in meiosis of the inter-specific hybrids between *Colocasia esculenta* (L.) Schott and *C. gigantea* Hook. f.*Cytologia*.56.389–393.
- 17.Kuruvilla KM, Dutt B, Roy RP.1989.Karyomorphological investigations on aroids of North-Eastern Hills.*Journal of Cytology and Genetics*.24.13-22.
- 18.Zhang G, Zhang D.1990.The relationship between geographic distribution and ploidy level of taro, *Colocasia esculenta*.*Euphytica*.47 .25-27.
- 19.Sreekumari MT, Mathew PM.1991.Karyomorphology of five morphotypes of taro (*Colocasia esculenta* (L.) Schott).*Cytologia*.56.215-218.
- 20.Sreekumari MT, Mathew PM.1991.Karyotypically distinct morphotypes in taro (*Colocasia esculenta* (L.) Schott).*Cytologia*.56.399-402.
- 21.Sreekumari MT, Mathew PM.1995.Karyotype distinction in Indian taros.*Journal of Cytology and Genetics*.30(2).143–149.
- 22.Ivancic A, Lebot V.1999.Botany and genetics of New Caledonian wild taro, *Colocasia esculenta*.*Pacific Science*.53.273–285.
- 23.Kokubugata G, Konishi T.1999.Implication of a basic chromosome number of $x = 14$ in seven cultivars of two varieties of *Colocasia esculenta* by fluorescent in situ hybridization using rDNA probe.*Cytologia* .64.77–83.
- 24.Chaudhuri JB, Sharma A.1978-1979.Chromosome studies in certain members of Araceae.*Genet. Iber*.30-31.161-187.
- 25.Sharma AK, Das NK.1954.Study of karyotypes and their alternations in Aroids.*Agtron. Lusit*.16.23-48.
- 26.Alam SS, Deen SS.2002.Karyotype and Isozyme analysis in three forms of *Colocasia esculenta* (Araceae).*Bangladesh J. Bot.* .31.95–98.
- 27.Simmonds NW.1954.Chromosome behavior in some tropical plants.*Heredity*.8.139-145.
- 28.Probatova NS, Seledets VP, Rudyka EG.2008.IAPT/IOPB chromosome data 5.*Taxon*.57 (2).559.
- 29.Parvin S, Kabir G, Ud-Deen MM, Sarker JK.2008.Karyotype analysis of seven varieties of taro *Colocasia esculenta* (L.) schott. from Bangladesh.*J. Bio-sci.* .16.15-18.
- 30.Zhang CS.1998.A preliminary study on making plant chromosomal specimens using peppermint oil compound as pretreatment agent.*Journal of Wuhan Botanical Research*.16 (3).280–282.
- 31.Sreekumari MT, Mathew PM.1991.Effect of colchicine treatment in a triploid variety of taro [*Colocasia esculenta* (L.) Schott].*New Botanist*.18.211-215.

32. Huang SF, Zhao ZF, Chen ZY, Chen SJ, Huang XX. 1989. Chromosome counts on one hundred species and infraspecific taxa. *Acta Botanica Austro Sinica*. 5. 161-176.
33. Subramanian D, Munian M. 1988. Cytotaxonomical studies in south Indian Araceae. *Cytologia* . 53. 59–66.
34. Udengwu OS, Okafor NA. 1999. Karyomorphological studies of five edible *Colocasia* and *Xanthosoma* species. *Nucleus (Calcutta)* . 42 (1,2) . 39–44.
35. Jos JS, Magoon ML. 1970. Pollen mitotic studies in some aroides. *Chromosome Inf. serv.* 11. 4-6.
36. Jos JS, Bai KV. 1978. Supernumerary gametes in the pollen tube of *Colocasia*. *Curr. Sci.* 47. 281-282.
37. Strauss MS, Stephen GC, Gonzales CJ, Arditti J. 1980. Genetic variability in taro, *Colocasia esculenta* (L.) Schott (Araceae). *Ann. Bot.* 45. 429-437.
38. Lebot V, Aradhya KM. 1991. Isozyme variation in taro (*Colocasia esculenta* (L.) Schott) from Asia and Oceania. *Euphytica*. 56. 55-66.
39. Matthews P, matsushita Y, Sato T, Hirai M. 1992. Ribosomal and mitochondrial DNA variation in Japanese taro (*Colocasia esculenta* L. Schott). *Japan J. Breed.* 42. 825-833.
40. Irwin SV, Kaufusi P, de la Pena R, Cho JJ. 1998. Molecular characterization of taro (*Colocasia esculenta*) using RAPD markers. *Euphytica*. 99. 183-189.
41. Lakhanpaul S, Velayudhan KC, Bhat KV. 2003. Analysis of genetic diversity in Indian taro [*Colocasia esculenta* (L.) Schott] using random amplified polymorphic DNA (RAPD) markers. *Genetic Resources and Crop Evolution* . 50. 603–609.
42. Kreike CM, Van Eck HJ, Lebot V. 2004. Genetic diversity of taro, *Colocasia esculenta* (L.) Schott, in Southeast Asia and the Pacific. *Theor. Appl. Genet.* . 109 . 761–768.
43. Okpul T, Mace ES, Godwin ID, Singh D, Wagih ME. 2005. Evaluation of variability among breeding lines and cultivars of taro (*Colocasia esculenta*) in Papua New Guinea using ISSR fingerprinting and agro-morphological characterization. *Plant Genetic Resources Newsletter* . 143. 8-16.
44. Caillon S, Quero-Garcia J, Lescure JP, Lebot V. 2006. Nature of taro (*Colocasia esculenta* (L.) Schott) genetic diversity prevalent in a Pacific Ocean island, Vanua Lava, Vanuatu. *Genetic Resources and Crop Evolution* . 53. 1273–1289.
45. Singh D, Mace ES, Godwin ID, Mathur PN, Okpul T, Taylor M, Hunter D, Kambuou R, Rao VR, Jackson G. 2008. Assessment and rationalization of genetic diversity of Papua New Guinea taro (*Colocasia esculenta*) using SSR DNA fingerprinting. *Genet. Resour. Crop. Evol.* 55. 811–822.
46. Beevi PN, Sreekumari MT, Kumar V. 2011. Study of genetic diversity in South Indian taro (*Colocasia esculenta* (L.) Schott.) using Random Amplified Polymorphic DNA Markers. *Journal of Root Crops*. 37 (2). 162-167.

- 47.Sardos J, Noyer JL, Malapa R, Bouchet S, Lebot V.2012.Genetic diversity of taro (*Colocasia esculenta* (L.) Schott) in Vanuatu (Oceania): an appraisal of the distribution of allelic diversity (DAD) with SSR markers.Genet. Resour. Crop. Evol.59.805–820.
- 48.Macharia MW, Runo SM, Muchugi AN, Palapala V.2014.Genetic structure and diversity of East African taro [*Colocasia esculenta* (L.) Schott].African Journal of Biotechnology.13.2950-2955.
- 49.Chair H, Traore RE, Duval MF, Rivallan R, Mukherjee A, Aboagye LM, Van Rensburg WJ, Andrianavalona V, Pinheiro de Carvalho MAA, Saborio F, Sriprana M, Komolong B, Lawac F, Lebot V.2016.Genetic diversification and dispersal of Taro (*Colocasia esculenta* (L.) Schott).Plos One.<http://dx.doi.org/10.5061/dryad.b8262>.
- 50.Nunes RSC, Aguila EMD, Paschoalin VMF, da Silva JT.2016.DNA barcoding assessment of the genetic diversity of varieties of taro, *Colocasia Esculenta* (L.) Schott in Brazil.ISBN:978-1-922227-355.Breeding and Genetic Engineering: The Biology and Biotechnology Research.iConcept Press.
- 51.Begum KN, Alam SS.2009.Karyotype analysis in three morphological forms of *Colocasia fallax* schott.Cytologia.74 (2).209-214.
- 52.Larsen K.1969.Cytology of Vascular Plants III. A study of Thai Aroids.Studies in the Flora of Thailand.Danish Botanical Society.
- 53.Long CL, Liu KM.2001.*Colocasia lihengiae* (Araceae: Colocasieae), a new species from Yunnan, China.Bot. Bull. Acad. Sin.42.313-317.
- 54.Ito T.1942.Chromosomen und Sexualitat von der Araceae. I. Somatische Chromosomenzahlen einiger Arten.Cytologia.12.313-325.
- 55.Nakajima G.1933.Chromosome numbers in some angiosperms .Jap. J. Genet.9.1-5.
- 56.Sharma AK, Sarkar AK.1963.Cytological analysis of different cytotypes of *Colocasia antiquorum*.Bulletin of the Botanical Society of Bengal.17.16-22.
- 57.Sharma AK.1956.A new concept of a means of speciation in plants .Caryologia.9.93-130.
- 58.Kurakubo Y .1940.Uber die chromosomenzahlen von Araceae-Arten.Bot. Zool. (Tokyo).8.1492.
- 59.Rattenbury JA.1957.Chromosome numbers in New Zealand angiosperms.Transaction of the Royal Society of New Zealand.84 (4).936-938.
- 60.Rattenbury JA.1956.Cytotaxonomy and the migration of primitive peoples.Nature .178.545-546.
- 61.Pfitzer P.1957.Chromosomenzahlen von Araceen.Chromosoma.8.436-446.
- 62.Sarkar AK.1960.Chromosomal biotype in *Colocasia antiquorum*.Proc. 47th Indian Science Congress.Part 3.367.

63. Krishnan R, Magoon ML, VijayaBai K.1970.Desynapsis in *Colocasia antiquorum* Schott.Genetica.41.170-178.
64. VijayaBai K, Magoon ML, Krishnan R.1971.Meiosis and pollen mitosis in diploid and triploid *Colocasia antiquorum* Schott.Genetica.42.187-198.
65. Maeda Y.1932.Cytological studies in *Colocasia*.Proc. Crop. Sci. Soc. Japan.4.315.
66. Kawahara T.1978.Chromosome number of taros in Nepal and India.Chromosome Information Service.24.4-5.
67. Chakraborty BN, Bhattacharya GN.1984.Cytotaxonomy of Araceae: III. A study of chromosomes of *Colocasia antiquorum* Schott in relation to its ecological adaptation.Cell and Chromosome Research.7.17–19.
68. Sarkar AK.1991.Studies on the cyto-ecological correlations of some araceous taxa occurring in different habitat.Proceedings of the Indian Science Congress Association.78 (3,VIII).136–137.
69. Yang Z, Yi T, Li H, Gong X.2003.A cytological study on three species of *Colocasia* (Araceae) from Yunnan.Caryologia.56 (3).323-327.
70. Rao NS.1947.A note on the chromosome number in *Colocasia antiquorum* Schott.Current Science.16.229.
71. Mookerjee A.1955.Cytology of different species of aroids with a view to trace the basis of their evolution.Caryologia.7 (2).221-291.
72. Krishnan R, Magoon ML, VijayaBai K.1970.Meiosis in a desynaptic triploid *Colocasia antiquorum* Schott.Genet. Agrar.24.325-334.
73. Chakraborty BN, Bhattacharya GN.1984.Desynapsis as well as inversion heterozygosity in the natural population of triploid *Colocasia antiquorum* Schott.Cytologia.49.739-743.
74. Cusimano N, Bogner J, Mayo SJ, Boyce PC, Wong SY, Hesse M, Hetterscheid WLA, Keating RC, French JC.2011.Relationships within the araceae: comparison of morphological patterns with molecular phylogenies.American Journal of Botany.98 (4).654-668.
75. Henriquez CL, Arias T, Pires JC, Croat TB, Schaal BA.2014.Phylogenomics of the plant family Araceae.Molecular Phylogenetics and Evolution.75.91-102.
76. Nguyen VX, Mai DTT, Anh NT.2015.Genetic diversity and phylogenetic relationships among *Colocasia* species (Araceae) in Vietnam using SSR markers.Aroideana.38 (1).177-185.