

References for *Lasia* L.

Genus ID : A-028.014

- 1.Hore P, Tanti B.2014.Karyomorphological studies of two morphotypes of *Lasia spinosa* (Lour.) Thwaites available in Assam, India. *Annals of Plant Sciences*.3 (8).792-796.
- 2.Pfitzer P.1957.Chromosomenzahlen Von Araceen. *Chromosoma* .8.436-446.
- 3.Sharma AK, Mukherjee D.1959.Further investigations on the cytology of some members of Aroids. *Proc. 46th Indian Sci. Congr. .Part 3*.356.
- 4.Larsen K.1969.Cytology of vascular plants. III. A study of Thai Aroids .27.39-69. *Studies in the Flora Of Thailand*.Danish Botanical Society.Denmark.
- 5.Hotta M.1971.Study of family Araceae. General remarks .*Jap. J. Bot.*20.269-310.
- 6.Ramachandran K.1978.Cytological studies on South Indian Araceae. *Cytologia*.43.289-303.
- 7.Sultana S, Rahman RB, Alam SS.2006.Karyotype analysis in three morphological forms of *Lasia spinosa* (L.) Thwaites (Araceae). *Cytologia*.71 (4).359–364.
- 8.Alam SS, Munira S, Habib MA, Sultana SS.2012.Karyotype and RAPD analysis in common and rare forms of *Lasia spinosa* (L.) Thwaites. *Cytologia* .77 (4). 499–505.
- 9.Duvall MR, Learn GH Jr., Eguiarte LE, Clegg MT.1993.Phylogenetic analysis of *rbcL* sequences identifies *Acorus calamus* as the primal extant monocotyledon. *Proc. Natl. Acad. Sci.*90.4641-4644.