

References of *Vigna* Savi

Genus ID : A-140.002

- 1.Bairiganjan GC, Patnaik SN.1989.Chromosomal Evolution in Fabaceae.Cytologia.54.51 - 64.
- 2.Shamurailatpam A, Madhavan L, Yadav SR, Bhat KV, Rao SR.2012.Chromosome diversity analysis in various species of *Vigna* Savifrom India.Nucleus.55.107 - 114.
- 3.Shamurailatpam A, Madhavan L, Yadav SR, Bhat KV, Rao SR.2015.Heterochromatin distribution and comparative karyo- morphological studies in *Vigna umbellata* Thunberg, 1969 and *V. aconitifolia* Jacquin, 1969(Fabaceae) accessions.Comparative Cytogenetics.9.119 - 132.
- 4.Rao SR, Raina SN.2004.Studies on male meiosis in cultivated and wild *Vigna* species.Plant Biotechnology and Molecular Markers.331 - 345.
- 5.Shamurailatpam A, Madhavan L, Yadav SR, Bhat KV, Rao SR.2014.Heterochromatin characterization through differential fluorophore binding pattern in some species of *Vigna* Savi.Protoplasma.doi: 10.1007/s00709-014-0708-y.
- 6.Shi CW, Jiang XH, Ou LJ, Liu J, Long KL, Zhang LH, Duan WT, Zhao W, Hu JC.2015.Molecular cytogenetic characterisation and phylogenetic analysis of the seven cultivated *Vigna* species (Fabaceae).Plant Biology.17.268 - 280.
- 7.Simon MV, Benko- Iseppon AM, Resende LV, Winter P, Kahl G.2007.Genetic diversity and phylogenetic relationships in *Vigna* Savi germplasm revealed by DNA amplification fingerprinting.Genome.50.538 - 547.
- 8.Pardhe DD, Satpute RA.2011.A Comparative Analysis of Genetic diversity in Genus *Vigna* Savi genotypes using ISSR.International Journal of Pharm Tech Research.3.464 - 470.
- 9.Gupta SK, Bansal R, Vaidya UJ, Gopalakrishna T.2012.Development of EST- derived microsatellite markers in mungbean [*Vigna radiata* (L.) Wilczek] and their transferability to other *Vigna* species.Indian Journal of Genetics.72.468 - 471.
- 10.Phansak P, Taylor PWJ, Mongkolporn O.2005.Genetic diversity in yardlong bean (*Vigna unguiculata* subsp. *sesquipedalis*) and related *Vigna* species using sequence tagged microsatellite site analysis.Scientia Horticulture (Amsterdam).106.137 - 146.
- 11.Venoral G, Blangiforti S, Cremonini R.1999.Karyotype analysis of twelve species belonging to Genus *Vigna*.Cytologia.64.117 - 127.
- 12.Goel S, Raina SN, Ogihara Y.2002.Molecular evolution and phylogenetic implications of internal transcribed spacer sequences of nuclear ribosomal DNA in the *Phaseolus* - *Vigna* complex.Molecular Phylogenetics and Evolution.22.1 - 19.
- 13.Chen HK, Mok MC, Shanmugasundaram S, Mok DWS.1989.Interspecific hybridization between *Vigna radiata* (L.) wilczek and *V. glabrescens*.Theoretical and Applied Genetics.78.641 - 647.

- 14.Parida A, Raina SN, Narayan RKJ.1990.Quantitative DNA variation between and within chromosome complements of *Vigna* species (Fabaceae).*Genetica*.82.125 - 133.
- 15.Joseph LS, Bouwkamp JC.1978.Karyomorphology of several of *Phaseolus* and *Vigna*.*Cytologia*.43.595 - 600.
- 16.Han OK, Kaga A, Isemura T, Wang XW, Tomooka N, Vaughan DA .2005.A genetic linkage map for azuki bean [*Vigna angularis* (Willd.) Ohwi & Ohashi].*Theoretical and Applied Genetics*.111.1278 - 1287.
- 17.Isemura T, Kaga A, Tomooka N, Shimizu T, Vaughan DA.2010.The genetics of domestication of rice bean, *Vigna umbellata*.*Annals of Botany*.106.927 - 944.
- 18.Kaga A, Ohnishi M, Ishii T, Kamijima O.1996.A genetic linkage map for azuki bean constructed with molecular and morphological markers using an interspecific population (*Vigna angularis* x *V. nakashimae*).*Theoretical and Applied Genetics*.93.658 - 663.
- 19.Xu HX, Jing T, Tomooka N, Kaga A, Isemura T, Vaughan DA.2008.Genetic diversity of the azuki bean (*Vigna angularis* (Willd.) Ohwi & Ohashi) gene pool as assessed by SSR markers.*Genome*.51.728 - 738.
- 20.Zong XX, Kaga A, Tomooka N, Wang XW, Han OK, Vaughan DA.2003.The genetic diversity of the *Vigna angularis* complex in Asia .*Genome*.46.647 - 658.
- 21.El- Hayd EA, Haiba AA, El- Hamid NR, Al- Ansary AE, Mohamed AY.2010.Assessment of genetic variations in some *Vigna* species by RAPD and ISSR analysis.*New York Science Journal* .3.120 - 128.
- 22.Kaewwongwal A, Kongjaimun A, Somta P, Chankaew S, Yimram T, Srinives P.2015.Genetic diversity of the black gram [*Vigna mungo* (L.) Hepper] gene pool as revealed by SSR markers.*Breeding Science*.65.127 - 137.
- 23.Xu RQ, Tomooka N, Vaughan AD, Doi K.2000.The *vigna angularis* complex: Genetic variation and relationships revealed by RAPD analysis and their implications for in- situ conservation and domestication.*Genetic Resources and Crop Evolution*.47.123 - 134.
- 24.Wang L, Kikuchi S, Muto C, Naito K, Isemura T, Ishimoto M, Cheng X, Kaga A, Tomooka N.2015.Reciprocal translocation identified in *Vigna angularis* dominates the wild population in East Japan.*Journal of Plant Research*.128.653 - 663.
- 25.Shamurailatpam A, Wahlang DR, Madhavan L, Yadav SR, Bhat KV, Rao SR.2016.Further observations on chromosome diversity analysis in wild species of *Vigna* from India .*Nucleus*.doi: 10.1007/s13237- 016-0157- 4.
- 26.Bachulkar MP, Yadav SR, Dixit GB.1992.Karyotype studies in *Vigna khandalensis* (Sant.) Raghavan and Wadhwa.*Proceedings of the Indian Science Congress Association*.79.85 - 86.
- 27.Venora G, Saccardo F.1993.Mitotic karyotype analysis in the *Vigna* genus by means of an image analyser.*Caryologia* .46.139 - 149.

- 28.Galasso I, Pignone D, Perrino P .1993.Cytotaxonomic studies in *Vigna*. II. Heterochromatin characterization in *Vigna unguiculata* and three related wild species.*Caryologia* .46.275 - 282.
- 29.Galasso I, Pignone D, Perrino P .1992.Cytotaxonomic studies in *Vigna*.I. General technique and *Vigna unguiculata* C- banding .*Caryologia* .45.155 - 161.
- 30.Galasso I, Schmidt T, Pignone D, Heslop- Harrison JS.1995.The molecular cytogenetics of *Vigna unguiculata* (L.) Walp: the physical organization and characterization of 18s- 5.8s - 25s rRNA genes, 5s rRNA genes, telomere- like sequences, and a family of centromeric repetitive DNA sequences.*Theoretical and Applied Genetics*.91.928 - 935.
- 31.Galasso I, Saponetti LS, Pignone D.1998.Cytotaxonomic studies in *Vigna*. IV. Variation of the number of active and silent rDNA sites in *Vigna unguiculata* populations.*Caryologia* .51.95 - 104.
- 32.Galasso I, Harrison GE, Pignone D, Brandes A, Heslop - Harrison JS.1997.The distribution and organization of Ty1- copia - like retrotransposable elements in the genome of *Vigna unguiculata* (L.) Walp. (cowpea) and its relatives.*Annals of Botany*.80.327 - 333.
- 33.Guerra M, Kenton A, Bennett MD.1996.rDNA sites in mitotic and polytene chromosomes of *Vigna unguiculata* (L.) Walp. and *Phaseolus coccineus* L. revealed by in situ hybridization.*Annals of Botany*.78.157 - 161.
- 34.Soliman MI, Rizk RM, El - Eraky NM.2008.Cytogenetics comparison of cultivated and wild relatives of Genus *Vigna* in Egypt.*Catrina*.3.61 - 70.
- 35.Vasconcelos EV, de Andrade Fonseca AF, Pedrosa- Harand A, de Andrade Bortoleti KC, Benko- Iseppon AM, da Costa AF, Brasileiro- Vidal AC.2015.Intra - and interchromosomal rearrangements between cowpea [*Vigna unguiculata* (L.) Walp.] and common bean (*Phaseolus vulgaris* L.) revealed by BAC- FISH.*Chromosome Research*.23.253 - 266.
- 36.Lavania UC, Lavania S.1982.Chromosome banding patterns in some Indian pulses.*Annals of Botany*.49.235 - 239.
- 37.Bortoleti KCDA, Benko- Iseppon AM, De Melo, NF, Brasileiro - Vidal AC.2012.Chromatin differentiation between *Vigna radiata* (L.) R. wilczek and *V. unguiculata*(L.) Walp.(Fabaceae).*Plant Systematics and Evolution*.298.689 - 693.
- 38.Venora G, Padulosi S.1997.Karyotypic analysis of wild taxa of *Vigna unguiculata*(L.) Walpers.*Caryologia* .50. 125 - 138.
- 39.Menendez CM, Hall AE, Gepts P.1997.A genetic linkage map of cowpea (*Vigna unguiculata*) developed from a cross between two inbred, domesticated lines.*Theoretical and Applied Genetics*.95.1210 - 1217.
- 40.Fatokun CA, Danesh D, Menancio- Hautea DI, Young ND.1993.A linkage map for cowpea [*Vigna unguiculata* (L) Walp] based on DNA markers (2n= 22).6256 - 6258.O'Brien JS . In *Genetic maps 1992*. A

compilation of linkage and restriction maps of genetically studied organisms.Cold Spring Harbor Laboratory Press,Cold Spring Harbour.New York.

41.Lucas MR, Diop NN, Wanamaker S, Ehlers JD, Roberts PA, Close TJ.2011.Cowpea- Soyabean synteny clarified through an improved genetic map.The Plant Genome.4.218 - 225.

42.Ouedraego JT, Gowda BS, Jean M, Close TJ, Ehlers JD, Hall AE, Gillaspie AG, Roberts PA, Ismail AM, Bruening G, Gepts P, Timko MP, Belzile FJ.2002.An improved genetic linkage map for cowpea (*Vigna unguiculata* L.) combining AFLP, RFLP, RAPD, biochemical markers, and biological resistance traits.Genome.45.175 - 188.

43.Ubi BE, Mignouna H, Thottappilly G.2000.Construction of a genetic linkage map and QTL analysis using a recombinant inbred population derived from an intersubspecific cross of cowpea (*Vigna unguiculata* (L.) Walp.) .Breeding Science.50.161 - 172.

44.Fang JG, Chao CCT, Roberts PA, Ehlers JD.2007.Genetic diversity of cowpea [*Vigna unguiculata* (L.) Walp.] in four West African and USA breeding programs as determined by AFLP analysis.Genetic Resources and Crop Evolution.54.1197 - 1209.

45.Gupta SK, Gopalakrishna T.2010.Development of unigene - derived SSR markers in cowpea (*Vigna unguiculata*) and their transferability to other *Vigna* species.Genome.53.508 - 523.

46.Li CD, Fatokun CA, Ubi B, Singh BB, Scoles GJ.2000.Determining Genetic similarities and relationships among cowpea breeding lines and cultivars by microsatellite markers.Crop Science.41.189 - 197.

47.Ba FS, Pasquet RS, Gepts P.2004.Genetic diversity in cowpea [*Vigna unguiculata*(L.) Walp.] as revealed by RAPD markers.Genetic Resources and Crop Evolution.51.539 - 550.

48.Zannoul A, Kossoul DK, Ahanchede A, Zoundjihékpon J, Agbicodo E, Struik PC, Sanni A.2008.Genetic variability of cultivated cowpea in Benin assessed by random amplified polymorphic DNA.African Journal of Biotechnology.7.4407 - 4414.

49.Diouf D, Hilu KF.2005.Microsatellites and RAPD markers to study genetic relationships among cowpea breeding lines and local varieties in senegal.Genetic Resources and Crop Evolution.52.1057 - 1067.

50.Gillaspie Jr AG, Hopkins MS, Dean RE.2005.Determining genetic diversity between lines of *Vigna unguiculata* subspecies by AFLP and SSR markers .Genetic Resources and Crop Evolution.52.245 - 247.

51.Mignouna HD, Ng NQ, Ikea J, Thottapilly G.1998.Genetic diversity in cowpea as revealed by random amplified polymorphic DNA.Journal of Genetics and Breeding.52.151 - 159.

52.Panella L, Kami J, Gepts P.1993.Vigna diversity in wild and cultivated taxa of *Vigna unguiculata* (L.) Walp. (Fabaceae).Economic Botany.43.371 - 386.

- 53.Asare AT, Gowda BS, Galyuon IKA, Aboagye LL, Takrama JF, Timko MP.2010.Assessment of the genetic diversity in cowpea (*Vigna unguiculata* L. Walp.) germplasm from Ghana using simple sequence repeat markers.*Plant Genetic Resources: Characterization and Utilization*.8.142 - 150.
- 54.Fall L, Diouf D, Fall- Ndiaye MA, Badiane FA, Gueye M.2003.Genetic diversity in cowpea [*Vigna unguiculata*(L.) Walp.] varieties determined by ARA and RAPD techniques.*African Journal of Biotechnology*.2.48 - 50.
- 55.Li CD, Fatokun CA, Ubi B, Singh BB, Scoles GJ.2001.Determining genetic similarities and relationships among cowpea breeding lines and cultivars by microsatellite primers.*Crop Science*.41.189 - 197.
- 56.Nkongolo KK.2003.Genetic characterization of Malawian cowpea (*Vigna unguiculata* (L.) Walp.) landraces: diversity and gene flow among accessions.*Euphytica*.129.219 - 228.
- 57.Ogunkanmi LA, Ogundipe OT, Ng NQ, Fatokun CA.2008.Genetic diversity in wild relatives of cowpea (*Vigna unguiculata*) as revealed by simple sequence repeats (SSR) markers.*Journal of Food Agriculture and Environment*.6.263 - 268.
- 58.Badiane FA, Gowda BS, Cisse N, Diouf D, Sadio O, Timko MP.2012.Genetic relationship of cowpea (*Vigna unguiculata*) varieties from Senegal based on SSR markers.*Genetics and Molecular Research*.11.292 - 304.
- 59.Dikshit HK, Singh D, Singh A, Jain N, Kumari J, Sharma TR.2012.Utility of adzuki bean [*Vigna angularis* (Willd.) Ohwi & Ohashi] simple sequence repeat (SSR) markers in genetic analysis of mungbean and related *Vigna* spp.*African Journal of Biotechnology*.11.13261 - 13268.
- 60.Singh A, Dikshit HK, Jain N, Singh D, Yadav RN.2014.Efficiency of SSR, ISSR and RAPD markers in molecular characterization of mungbean and other *Vigna* species.*Indian Journal of Biotechnology*.13.81 - 88.
- 61.Samarajeewa PK, Nanayakkara HL, Ekanayake EM, Sumanasinghe VA.2002.RAPD analysis of genetic relationships of wild and cultivated *Vigna* species .*Annals of Sri Lanka Depart. Agriculture*.4.215 - 226.
- 62.Patil DM, Sawardekar SV, Gokhale NB, Bhave SG, Sawant SS, Sawantdesai SA, Lipne KA, Sabale SN, Joshi SN.2013.Genetic diversity analysis in cowpea [*Vigna unguiculata* (L.) Walp.] by using RAPD markers.*International Journal of Innovative Biotechnology and Biochemistry*.1.15 - 23.
- 63.Karuppanapandian T, Karuppudurai T, Sinha PB, Haniya AMK, Manoharan K.2006.Phylogenetic diversity and relationships among cowpea (*Vigna unguiculata* Walp.) landraces using random amplified polymorphic DNA markers.*General and Applied Plant Physiology*.32.141 - 152.
- 64.Nagalakshmi RM, Kumari RU, Boranayaka MB.2010.Assessment of genetic diversity in cowpea (*Vigna unguiculata*).*Electronic Journal of Plant Breeding*.1.453 - 461.

- 65.Kumari S, Sahai G, Pandey SK, Pandey HC.2012.Identification of putative hybrids of cowpea (*Vigna unguiculata* (L.) Walp.) and their genetic evaluation using RAPD marker.Agricultural Science Research Journals.2.506 - 511.
- 66.Coulibaly S, Pasquet RS, Papa R, Gepts P.2002.AFLP analysis of the phenetic organization and genetic diversity of *Vigna unguiculata* L. Walp. reveals extensive gene flow between wild and domesticated types.Theoretical and Applied Genetics.104.358 - 366.
- 67.Ajibade SR, Weeden NF, Chite SM.2000.Inter simple sequence repeat analysis of genetic relationships in the genus *Vigna*.*Euphytica*.111.47 - 55.
- 68.Dikshit HK, Jhang T, Singh NK, Koundal KR, Bansal KC, Chandra N, Tickoo JL, Sharma TR.2007.Genetic differentiation of *Vigna* species by RAPD, URP and SSR markers.*Biologia Plantarum*.51.451 - 457.
- 69.Alam SS, Tasmin R, Jahan I, Habib MA, Sultana SS.2013.Fluorescent banding and RAPD analysis of five cultivars in *Vigna unguiculata* subsp. *sesquipedalis* (L.) Verdc.*Cytologia*.78.73 - 79.
- 70.Xu P, Wu X, Wang B, Liu Y, Ehlers JD, Close TJ, Roberts PA, Diop NN, Qin D, Hu T, Lu Z, Li G .2011.A SNP and SSR based genetic map of asparagus bean (*Vigna unguiculata* subsp. *sesquipedalis*) and comparison with the broader species .*PLoS ONE*.6.doi: 10.1371/journal.pone.0015952.
- 71.Adetula OA.2006.Comparative study of the karyotypes of two *Vigna* sub species.*African Journal of Biotechnology*.5.563 - 565.
- 72.Tosti N, Negri V.2002.Efficiency of PCR- based markers in assessing genetic variation among cowpea (*Vigna unguiculata* subsp. *unguiculata*) landraces.*Genome*.45.268 - 275.
- 73.Sen O, Bhattacharya S.1988.Cytomixis in *Vigna glaberscens* TTKI (Wild).*Cytologia*.53.437 - 440.
- 74.Sen NK, Bhowal JG.1960.Cytotaxonomic studies on *Vigna*.*Cytologia*.25.195 - 207.
- 75.Sonnante G, Spinoso A, Marangi A, Pignone D.1997.Isozyme and RAPD analysis of the genetic diversity within and between *Vigan luteola* and *V. marina*.*Annals of Botany*.80.741 - 746.
- 76.Galasso I, Saponetti LS, Pignone D .1996.Cytotaxonomic studies in *Vigna*. III. Chromosomal distribution and reacting properties of the heterochromatin in five wild species of the section *Vigna*.*Caryologia* .49.311 - 319.
- 77.Rao SR, Chandel KPS.1991.Karyomorphological studies in the cultivated and wild *Vigna* species in Indian Gene Centre.*Cytologia*.56.47 - 57.
- 78.Alam SS, Mahbub MN.2007.Karyotype comparison in two varieties of *Vigna mungo* L. after staining with orcein and CMA.*Bangladesh Journal of Botany*.36.167 - 170.
- 79.Mannan MA, Kabir G, Shaheen NN, Akhter FN.1991.Meiotic studies in seven pulse crops of Bangladesh.*Cytologia*.56.511 - 515.

- 80.Gupta SK, Soufmanien J, Gopalakrishna T.2008.Construction of a genetic linkage map of black gram, *Vigna mungo* (L.) Hepper, based on molecular markers and comparative studies.Genome.51.628 - 637.
- 81.Chaitieng B, Kaga A, Tomooka N, Isemura T, Kuroda Y, Vaughan DA.2006.Development of a black gram [Vigna mungo (L.) Hepper] linkage map and its comparison with an azuki bean [Vigna angularis (Willd.) Ohwi and Ohashi] linkage map.Theoretical and Applied Genetics.113.1261 - 1269.
- 82.Ghafoor A, Sultana T, Rizvi ZF.2012.Genetic diversity in black gram [Vigna mungo (L.) Hepper] for randomly amplified polymorphic DNA (RAPD) markers .Pakistan Journal of Botany.44.473 - 478.
- 83.Ghafoor A, Sharif A, Ahmad Z, Zahid MA, Rabbani MA.2001.Genetic diversity in black gram (*Vigna mungo* L. Hepper) .Field Crop Research.69.183 - 190.
- 84.Shafique S, Khan MA, Nisar M, Rehman S.2011.Investigation of genetic diversity in black gram [Vigna mungo (L.) Hepper].Pakistan Journal of Botany.43.1223 - 1232.
- 85.Soufmanien J, Gopalakrishna T.2004.A comparative analysis of Genetic diversity in black gram genotypes using RAPD and ISSR markers.Theoretical and Applied Genetics.109.1687 - 1693.
- 86.Soufmanien J, Reddy KS.2015.Denovo assembly, characterization of immature seed transcriptome and development of Genic - SSR markers in black gram [Vigna mungo (L.) Hepper].PLoS ONE.10.doi: 10.1371/journal.pone. 0128748.
- 87.Sivaprakash KR, Prashanth SR, Mohanty BP, Parida A.2004.Genetic diversity of black gram (*Vigna mungo*) landraces as evaluated by amplified fragment length polymorphism markers.Current Science.86.1411 - 1416.
- 88.Kanimozhi M, Jayamani P, Nadarajan N.2009.Genetic diversity as assessed by ISSR markers in black gram (*Vigna mungo* (L.) Hepper) .Electronic Journal of Plant Breeding.1.12 - 17.
- 89.Shanthi P, Jebaraj S, Manivannan N.2006.Genetic diversity in Urdbean (*Vigna mungo* L. Hepper) .Legume Research.29.181 - 185.
- 90.Tantasawat P, Trongchuen J, Prajongjai T, Thongpae T, Petkhum C, Seehalak W, Machikowa T.2010.Variety identification and genetic relationships of mungbean and black gram in Thailand based on morphological characters and ISSR analysis.African Journal of Biotechnology.9.4452 - 4464.
- 91.Ganguly PR, Bhat KV.2012.Study of the pattern of variation for microsatellite markers in black gram, *Vigna mungo* (L.) Hepper germplasm.DHR International Journal of Biomedical and Life Sciences (DHR - IJBLs.3.2278 - 8301.
- 92.Srivastava P, Pandey A, Sinha DP .2011.Genetic diversity analysis in different varieties of black gram using RAPD markers .Journal of Plant Breeding and Crop Science.3.53 - 59.
- 93.Mahbub MN, Rahman ANMRB, Alam SS.2007.Development of marker chromosomes in three varieties of *Vigna radiata* L. (Fabaceae).Cytologia.72.221 - 225.

- 94.Khattak GSS, Wolny E, Saeed I.2007.Detection of ribosomal DNA sites in chickpea (*Cicer arietinum* L.) and mungbean (*Vigna radiata* (L.) Wiltz) by fluorescence in situ hybridization.Pakistan Journal of Botany.39.1511 - 1515.
- 95.Humphry M, Konduri V, Lambrides C, Magner T, McIntyre C, Aitken E, Liu C.2002.Development of a mungbean (*Vigna radiata*) RFLP linkage map and its comparison with lablab (*Lablab purpureus*) reveals a high level of colinearity between the two genomes.Theoretical and Applied Genetics.105.160 - 166.
- 96.Lambrides CJ, Lawn RJ, Godwin ID, Manners J, Imrie BC.2000.Two genetic linkage maps of mungbean using RFLP and RAPD markers.Australian Journal of Agricultural Research.51.415 - 425.
- 97.Isemura T, Kaga A, Tabata S, Somta P, Srinives P, Takehiko S, Uken J, Vaughan DA, Tomooka N .2012.Construction of a genetic linkage map and genetic analysis of domestication related traits in mungbean (*Vigna radiata*).PLoS ONE.7.doi: 10.1371/journal.pone. 0041304.
- 98.Tangphatsornruang S, Somta P, Uthaipaisanwong P, Chanprasert J, Sangsrakru D, Seehalak W, Sommanas W, Tragoonrung S, Srinives P.2009.Characterization of microsatellites and gene contents from genome shotgun sequences of mungbean (*Vigna radiata* (L.) Wilczek.BMC Plant Biology.9.doi: 10.1186/1471- 2229-9-137.
- 99.Wang LX, Elbaidouri M, Abernathy B, chen HL, Wang SH, Lee SH, Jackson SA, Cheng XZ.2015.Distribution and analysis of SSR in mung bean (*Vigna radiata* L.) genome based on an SSR-enriched library.Molecular Breeding.35.1 - 10.
- 100.Sangiri C, Kaga A, Tomooka N, Vaughan D, Srinives P.2007.Genetic diversity of the mungbean (*Vigna radiata*, Leguminosae) genepool on the basis of microsatellite analysis.Australian Journal of Botany.55.837 - 847.
- 101.Singh S, Jawali N.2009.Development and characterization of trinucleotide microsatellite loci from *Vigna radiata*.International Journal of Plant Breeding.3.71 - 75.
- 102.Singh S, Reddy SK, Jawali N.2012.Genetic diversity analyses of mungbean (*Vigna radiata* [L]. Wilczek) by ISSR.International Journal of Plant Breeding.6.73 - 83.
- 103.Gwag JG, Chung JW, Chung HK, Lee JH, Ma KH, Dixit A, Park YJ, Cho EG, Kim TS, Lee SH.2006.Characterization of new microsatellite markers in mung bean, *Vigna radiata* (L.).Molecular Ecology Notes.6.1132 - 1134.
- 104.Somta P, Musch W, Kongsamai B, Chanprame S, Nakasathien S, Toojinda T, Sorajjapinun W, Seehalak W, Tragoonrung S, Srinives P.2008.New microsatellite markers isolated from mungbean (*Vigna radiata* (L.) Wilczek).Molecular Ecology Resources.8.1155 - 1157.
- 105.Kumar SV, Tan SG, Quah SC, Yusoff K.2002.Isolation of microsatellite markers in mungbean, *Vigna radiata*.Molecular Ecology Notes.2.96 - 98.

- 106.Pandey A, Kumar A, Ramya.2011.Genetic diversity in green gram accessions as revealed by STMS markers.African Journal of Biotechnology.10.17081 - 17087.
- 107.Karuppanapandian T, Karuppudurai T, Sinha PB, Haniya KA, Manoharan K.2006.Genetic diversity in green gram [*Vigna radiata* (L.)] landraces analyzed by using random amplified polymorphic DNA (RAPD).African Journal of Biotechnology.5.1214 - 1219.
- 108.Santalla M, Power JB, Davey MR.1998.Genetic diversity in mung bean germplasm revealed by RAPD markers.Plant Breeding.117.473 - 478.
- 109.Palaniappan J, Murugaiah S.2012.Genetic diversity as assessed by morphological and microsatellite markers in green gram (*Vigna radiata* L.).African Journal of Biotechnology.11.15091 - 15097.
- 110.Bhat KV, Lakhampaul S, Chadha S.2005.Amplified fragment length polymorphism (AFLP) analysis of genetic diversity in Indian mungbean (*Vigna radiata* (L.) Wilczek) Cultivars.Indian Journal of Biotechnology.4.56 - 64.
- 111.Lakhampaul S, Chadha S, Bhat KV.2000.Random amplified polymorphic DNA (RAPD) analysis in Indian mungbean (*Vigna radiata* (L.)Wilczek cultivars.Genetica.109.227 - 234.
- 112.Saini A, Reddy SK, Jawali N.2004.Evaluation of long primers for AP - PCR analysis of mungbean [*Vigna radiata* (L.) Wilczek]: Genetic relationships and fingerprinting of some genotypes .Indian Journal of Biotechnology.3.511 - 518.
- 113.Prasad S, Reddy KS, Jawali N.1999.Abundance of randomly amplified hybridizing microsatellites in mungbean *Vigna radiata* (L.) Wilczek.Asia Pacific Journal of Molecular Biology and Biotechnology.7.1 - 11.
- 114.Betal S, Chowdhury PR, Kundu S, Raychaudhuri SS.2004.Estimination of genetic variability of *Vigna radiata* cultivars by RAPD analysis.Biologia Plantarum.48.205 - 209.
- 115.Lavanya GR, Srivastava J, Ranade SA.2008.Molecular assessment of genetic diversity in mung bean germplasm.Journal of genetics .87.65 - 74.
- 116.Kumar SV, Tan SG, Quah SC, Yusoff K.2002.Isolation and characterization of seven tetranucleotide microsatellite loci in mungbean, *Vigna radiata*.Molecular Ecology Notes.2.293 - 295.
- 117.Ranade R, Gopalakrishna T.2001.Characterization of blackgram (*Vigna radiata* (L) Hepper) varieties using RAPD.Plant variety and seeds.14.227 - 233.
- 118.Chattopadhyay K, Ali MN, Sarkar HK, Mandal N, Bhattacharyya S.2005.Diversity analysis by RAPD and ISSR markers among the selected mungbean (*Vigna radiata* (L.) Wilczek) genotypes.Indian Journal of Genetics.65.173 - 175.

- 119.Singh R, van Heusden AW, Yadav RC.2013.A comparative genetic diversity analysis in mungbean (*Vigna radiata* L.) using inter- simple sequence repeat (ISSR) and amplified fragment length polymorphism (AFLP) .*African Journal of Biotechnology*.12.6574 - 6582.
- 120.Islam S, Mir JI, Kudesia R.2015.Evalution of genetic diversity in *Vigna radiata* L.) using protein profiling and moleculr marker (RFLP).*International Journal of Plant Breeding and Genetics*.9.238 - 246.
- 121.Datta S, Gangwar S, Kumar S, Gupta S, Rai R, Kaashyap M, Singh P, Chaturvedi SK, Singh BB, Nadarajan N.2012.Genetic diversity in selected Indian mungbean [*Vigna radiata* (L.) Wilczek] cultivars using RAPD markers.*American Journal of Plant Sciences*.3.1085 - 1091.
- 122.Tripathy SK, Sardar SS, Dash S, Pal S, Acharya T, Mohapatra BR, Das AB, Rout GR.2015.RAPD markers for genetic characterization in mungbean [*Vigna radiata* (L.) Wilczek].*Legume Research*.38.279 - 288.
- 123.Saini M, Singh S, Hussain Z, Yadav A.2010.RAPD analysis in mungbean (*Vigna radiata* (L.) Wilczek): I. Assessment of genetic diversity.*Indian Journal of Biotechnology*.9.137 - 146.
- 124.Undal VS, Thakare PV, Chaudhari US, Deshmukh VP, Gawande PA.2011.Estimination of genetic diversity among wild Vigna species revealed by RAPD markers .*Annals of Biological Research*.2.348 - 354.
- 125.Adetula OA, Fatokun CA, Obigbesan G.2005.Centromeric banding pattern of mitotic chromosomes in *Vigna vexillata* (TVnu 73).*African Journal of Biotechnology*.4.400 - 402.
- 126.Ogundiwin EA, Thottappilly G, Akenova ME, Pillay M, Fatokun CA.2005.A genetic linkage map of *Vigna vexillata*.*Plant Breeding*.124.392 - 398.
- 127.Wang L, Do Kim K, Gao D, Chen H, Wang S, Lee S, Jackson SA, Cheng X.2016.Analysis of simple sequence repeats in rice bean (*Vigna umbellata*) using an SSR- enriched library.*The Crop Journal*.4.40 - 47.
- 128.Tian J, Isemura T, Kaga A, Vaughan DA, Tomooka N.2013.Genetic diversity of the rice bean (*Vigna umbellata*) genepool as assessed by SSR markers.*Genome*.56.717 - 727.
- 129.Wang LX, Chen HL, Bai P, Wu JX, Wang SH, Blair MW, Cheng XZ.2015.The transferability and polymorphism of mung bean SSR markers in rice bean germplasm.*Molecular Breeding*.35.doi: 10.1007/s11032- 015- 0280-y.
- 130.Chen H, Chen X, Tian J, Yang Y, Liu Z, Hao X, Wang L, Wang S, Liang J, Zhang L, Yin F, Cheng X.2016.Development of gene - based SSR markers in rice bean (*Vigna umbellata* L.) based on transcriptome data.*PLoS ONE*.11.doi: 10.1371/journal.pone. 0151040.
- 131.Muthusamy S, Kanagarajan S, Ponnuswamy S.2008.Efficiency of RAPD and ISSR marker systems in accessioning genetic variation of rice bean (*V. umbellata*) landraces.*Electronic Journal of Biotechnology*.11.1 - 10.

- 132.Osuji JO, Edeoga HO.2005.Karyological studies on *Vigna subterranea* (L.) Verdc. From Nigeria.Cytologia.70.167 - 169.
- 133.Ntundu WH, Bach IC, Christiansen JL, Andersen SB.2004.Analyses of genetic diversity in bambara groundnut [*Vigna subterranea* (L.) Verdc.] landraces using amplified fragment length polymorphism (AFLP) markers.African Journal of Biotechnology.3.220 - 225.
- 134.Somta P, Chankaew S, Rungnoi O, Srinives P.2011.Genetic diversity of the bambara groundnut (*Vigna subterranea* (L.) Verdc.) as assessed by SSR markers.Genome.54.898 - 910.
- 135.Massawe FJ, Dickinson M, Roberts JA, Azam - Ali SN.2002.Genetic diversity in bambara groundnut (*Vigna subterranea* (L.) Verdc) landraces revealed by AFLP markers.Genome.45.1175 - 1180.
- 136.Amadou HI, Bebeli PJ, Kaltsikes PJ.2001.Genetic diversity in bambara groundnut (*Vigna subterranean* (L.) Verdc) germplasm revealed by RAPD markers.Genome.44.995 - 999.
- 137.Odongo FO, Oyoo ME, Wasike V, Owuoche JO, Karanja L, Korir P.2015.Genetic diversity of bambara groundnut (*Vigna subterranea* (L.) Verdc.) landraces in Kenya using microsatellite markers.African Journal of Biotechnology.14.283 - 291.
- 138.Aliyu S, Festo JM.2013.Microsatellites based marker molecular analysis of Ghanaian bambara groundnut (*Vigna subterranean* (L.) Verdc.) landraces alongside morphological characterization.Genetic Resources and Crop Evolution.60.777 - 787.
- 139.Olukolu BA, Mayes S, Stadler F, Ng QN, Fawole I, Dominique D, Azam Ali SN, Abott GA, Kole C.2012.Genetic diversity in bambara groundnut (*Vigna subterranea* (L.)Verdc.) as revealed by phenotypic descriptors and DArT marker analysis.Genetic Resources and Crop Evolution.59.347 - 358.
- 140.Rungnoi O, Suwanprasert J, Somta P, Srinives P.2012.Molecular genetic diversity of bambara groundnut (*Vigna subterranea* (L.) Verdc.) revealed by RAPD and ISSR marker analysis.SABRAO Journal of Breeding and Genetics.44.87 - 101.
- 141.Tun, Tun Y, Yamaguchi H.2007.Phylogenetic relationship of wild and cultivated *Vigna* (Subgenus *Ceratotropis*, Fabaceae) from Myanmar based on sequence variations in non- coding regions of trn T-F.Breeding Science.57.271 - 280.
- 142.Doi K, Kaga A, Tomooka N, Vaughan DA.2002.Molecular phylogeny of genus *Vigna* subgenus *ceratotropis* based on rDNA ITS and atpB-rbcL intergenic spacer of cpDNA sequences.Genetica.114.129 - 145.
- 143.Raveendar S, Lee JR, Park JW, Lee GA, Jeon YA, Lee WH, Cho GT, Ma KH, Lee SY, Chung JW.2015.Identification of genus *Vigna* using ITS2 and matK as a two- locus DNA barcode .Plant Breeding and Biotechnology.3.153 - 159.

- 144.Kumar AV, Saini A, Jawali N.2010.Phylogenetic analysis of subgenus *Vigna* species using nuclear ribosomal RNA ITS: evidence of hybridization among *Vigna unguiculata* subspecies.Journal of Heredity.101.177 - 188.
- 145.Faris DG.1965.Origin and evolution of the cultivated forms of *Vigna sinensis*.Canadian Journal of Genetics and Cytology.7.433 - 452.
- 146.Valliancourt RE, Weeden NF .1992.Chloroplast DNA polymorphism suggests Nigerian center of domestication for the cowpea, *Vigna unguiculata* (Leguminosae).American Journal of Botany.79.1194 - 1199.
- 147.Marechal R, Mascherpa JM, Stainier F.1978.Etude taxonomique d'un groupe complexe d'espèces des genres *Phaseolus* et *Vigna* (Papilionaceae) sur la base de données morphologiques et polliniques, traitées par l'analyse informatique.Biossiera Geneve.28.1 - 273.
- 148.Baudoin JP, Marechal R.1988.Taxonomy and evolution of the genus *Vigna*.2-12.Proceedings of the Second International Mungbean Symposium.Asian Vegetable Res. Dev. Center, shanhua, Taiwan.
- 149.Egawa Y, Nakagahra M, Fernandez GCJ.1988.Cytogenetical analysis of tetraploid *Vigna glabrescens* by interspecific hybridization involving diploid Asian *Vigna* species .200 - 204.Proceedings of the Second International Mungbean Symposium.Asian Vegetable Res. Dev. Center, shanhua, Taiwan.
- 150.Maréchal, R.1969.Données cytologiques sur les espèces de la sous-tribu des Papilionaceae-Phaseoleae-Phaseolinae Première série (Cytological data on species of the subtribe Papilionaceae - Phaseoleae - Phaseolinae First series).Bulletin du Jardin botanique National de Belgique (Bulletin of the National Botanic Garden of Belgium).39.125-165. Botanic Garden Meise.
- 151.Maréchal, R.1970.Données cytologiques sur les espèces de la sous-tribu des Papilionaceae-Phaseoleae-Phaseolinae. Deuxième série (Cytological data on species of the subtribe Papilionaceae - Phaseoleae - Phaseolinae. Second series).Bulletin du Jardin botanique National de Belgique (Bulletin of the National Botanic Garden of Belgium).40.307-318+320-349. Botanic Garden Meise.
- 152.Frahm-Leliveld, J.A.1965.Cytological data on some wild tropical *Vigna* species and cultivars from cowpea and asparagus bean.Euphytica.14.251-270.
- 153.Vijaykumar A., Saini A. and Jawali N.2011.Molecular characterization of intergenic spacer region of 5S ribosomal RNA genes in subgenus *Vigna*: extensive hybridization among *V. unguiculata* subspecies.Plant Systematics and Evolution.294.39-55.Springer Vienna.
- 154.Patel O. P. and R. C. Verma.1999.Cytomorphology of induced tetraploidy in *Cajanus cajan* and *Vigna unguiculata*.Cytologia.64.29-36.
- 155.Jahan B., Vahidy A. A. and Ali S. I.1994.Chromosome numbers in some taxa of Fabaceae mostly native to Pakistan.Annals of the Missouri Botanical Garden.81.792-799.Missouri Botanical Garden Press.

- 156.Kongjaimun, A., Kaga, A., Tomooka, N., Somta, P., Shimizu, T., Shu, Y., Isemura, T., Vaughan, D.A. and Srinives, P.2012.An SSR-based linkage map of yardlong bean (*Vigna unguiculata* (L.) Walp. subsp. *unguiculata* Sesquipedalis Group) and QTL analysis of pod length.Genome.55.81-92.
- 157.Tosti, N. and Negri V.2002.Efficiency of three PCR-based markers in assessing genetic variation among cowpea (*Vigna unguiculata* subsp. *unguiculata*) landraces.Genome.45.268-275.
- 158.Coulibaly, S., Pasquet, R.S., Papa, R. and Gepts, P.A.F.L.P.2002.AFLP analysis of the phenetic organization and genetic diversity of *Vigna unguiculata* L. Walp. reveals extensive gene flow between wild and domesticated types.Theoretical and Applied Genetics.104.358-366.
- 159.Gillaspie, A.G., Hopkins, M.S. and Dean, R.E.2005.Determining genetic diversity between lines of *Vigna unguiculata* subspecies by AFLP and SSR markers.Genetic Resources and Crop Evolution.52.245-247.