

G-011.002

1. Gulline H F . 2003 . In: R. J. E. Wiltshire & W. D. Jackson, Index of chromosome numbers of Tasmanian spermatophytes . Pap. & Proc. Roy. Soc. Tasmania . 137 : 39–53.
2. Hair JB . 1968 . The chromosomes of the Cupressaceae. 1. Tetraclineae and Actinostrobeae (Callitroideae). . New Zealand Journal of Botany . 6 : 272-284.
3. Khoshoo TN . 1961 . Chromosome numbers in gymnosperms . Silvae Genetica . 10 : 1-9.
4. Li L C, Cong B, Liu C, Liu YQ,Weng RF . 1994 . Karyotype analysis of 3 species of Callitris (Cupressaceae) in Australia and its phylogenetic significance . Acta Bot. Yunnan . 16 : 349–353.
5. Mehra P N . 1988 . Indian Conifers, Gnetophytes and Phylogeny of Gymnosperms. . Panjab University.
6. Ohri D, Khoshoo TN . 1986 . Genome size in gymnosperms . Plant Systematics and Evolution . 153 : 119–132.
7. Zonneveld BJM . 2012 . Conifer genome size of 172 species covering 64 of 67 genera range from 8-72 picograms. . Nordic Journal of Botany . 30 : 490-502.
8. Hizume M, Fujiwara M . 2016 . Fluorescent chromosome banding patterns of several species in the Cupressaceae sensu stricto . Chromosome Botany . 11 : 1-8.
9. Gadek P A, Alpers DL, Heslewood MM, Quinn CJ . 2000 . Relationships within Cupressaceaesensulato: a combined morphological and molecular approach . American Journal of Botany . 87 : 1044-1057.
10. Piggin J, Bruhl J . 2010 . Phylogeny reconstruction of *Callitris* Vent. (Cupresseceae) and its allies leads to inclusion of *Actinostrobus* within *Callitris* . Australian Systematic Botany . 23 : 69-93.

