

1. Bou Dagher-Kharrrat M, Grenier G, Bariteau M, Brown S, Siljak-Yakovlev, Savoure A . 2001 . Karyotype analysis reveals interspecific differentiation in the genus *Cedrus* despite genome size and base composition constancy . *Theoretical and Applied Genetics* . 103 : 846-854.
2. Bou Daghar-Kharrrat, MB, Mariette S, Lefevre F, Fady B, Grenier-de March G, Plomion C, Savoure A . 2007 . Geographical diversity and genetic relationships among *Cedrus* species estimated by AFLP . *Tree Genetics & Genomes* . 3 : 275-285.
3. Hizume M . 1988 . Karyological studied in the family Pinaceae. . *Memoirs of the Faculty of Education Ehime University Series 3* . : 1-108.
4. Khoshoo TN . 1961 . Chromosome numbers in gymnosperms . *SilvaeGenetica* . 10 : `1-9.
5. Li L C, Fu YX. . 1995 . Studies on the cytotaxonomy and the historical phytogeography of *Cedrus* (Pinaceae). . *Acta Bot. Yunnan* . 17 : 41–47.
6. Mehra, P. N. . 1998 . Indian Conifers, Gnetophytes and Phylogeny of Gymnosperms . Panjab University .
7. Mehra PN, Khoshoo TN . 1956 . Cytology of conifers I . *Journal of Genetics* . 54 : 165-185.
8. Qiao CY, Ran JH, Li Y, Wang XQ . 2007 . Phylogenetics and biogeography of *Cedrus* (Pinaceae) inferred from sequences of seven paternal chloroplast and maternal mitochondrial DNA region. . *Annals of Botany* . 100 : 573-580.
9. Zonneveld BJM . 2012 . Genome sizes of 172 species, covering 64 out of the 67 genera, range from 8 to 72 picogram . *Nordic Journal of Botany* . 30 : 490-502.