

1. Gu Z J, Zhou Q X, Yue Z S . 1988 . A karyomorphological study of Cephalotaxaceae . Acta Phytotax. Sin . 36 : 47–52.
2. Khoshoo TN . 1961 . Chromosome numbers in gymnosperms . Silvae Genetica . 10 : 1-9.
3. Mehra P N . 1988 . Indian Conifers, Gnetophytes and Phylogeny of Gymnosperms . Panjab University.
4. Mehra P N, Khoshoo T N . 1957 . Cytology of conifers II . Journal of Genetics . 54 : 181-185.
5. Zonneveld B J M . 2012 . Conifer genome sizes of 172 species, covering 64 of 67 genera, range from 8 to 72 picogram . Nordic Journal of Botany . 30 : 490–502.
6. Leitch IJ, Hanson L, Winfield M, Parker J, Bennett MD . 2001 . Nuclear DNA C-values complete familial representation in gymnosperms . Annals of Botany . 88 : 843-849.
7. Cheng Y, Nicolson RG, Tripp K, Chaw SM . 2000 . Phylogeny of Taxaceae and Cephalotaxaceae genera inferred from chloroplast matK gene and nuclear rDNA ITS region . Molecular Phylogenetics and Evolution . 14 : 353-365.
8. Khoshoo TN . 1957 . Partial failure of meiotic spindle in Cephalotaxus drupacea var. pedunculata . Cytologia . 22 : 80-89.
9. Hizume M, Kaneko K, Miyake T . 2014 . A method for preparation of meiotic chromosomes of conifers and its application . Chromosome Botany . 9 : 83-88.