

1. Fang Y X . 1988 . Karyotypic analyses of *Glyptostrobus pensilis*, *Metasequoia glyptostroboides* and *Taiwania flousiana* . J. Shanghai Norm. Univ., Nat. Sci. Ed . 3 : 48–55.
2. Hizume, M . 1989 . Karyomorphological studies in twelve species in the Taxodiaceae with special reference to cytotaxonomical position of *Sciadopitys verticillata* . Educ. Ehime Univ., Ser. 3, Nat. Sci . 9 : 7–32.
3. Hizume M, Kondo T, Shibata F, Ishizuka R . 2001 . Flow cytometric determination of genome size in the Taxodiaceae, Cupressaceae sensu stricto and Sciadopityaceae. . Cytologia . 66 : 307-311.
4. Khoshoo TN . 1961 . Chromosome numbers in gymnosperms . Silvae Genetica . 10 : 1-9.
5. Ohri, D, Khoshoo T N . 1986 . Genome size in gymnosperms . Plant Systematics and Evolution . 153 : 119–132.
6. Schlarbaum S E, Johnson L C, Tsuchiya T . 1983 . Chromosome studies of *Metasequoia glyptostroboides* and *Taxodium distichum*. Botanical Gazette . 144 : 559–565.
7. Stebbins GL . 1948 . The chromosomes and relationships of *Metasequoia* and *Sequoia* . Science . 108 : 95-98.
8. Zonneveld BJM . 2012 . Conifer genome size of 172 species covering 64 of 67 genera range from 8-72 picograms . Nordic Journal of Botany . 30 : 490-502.
9. Hizume M, Abe KK, Tanaka A . 1988 . Fluorescent chromosome bandings in the Taxodiaceae . La Chromosomo . 50 : 1609-1619.
10. Hizume M, Kaneko K, Miyake T . 2014 . A method for preparation of meiotic chromosomes of conifers and its applications . Chromosome Botany . 9 : 83-88.