

1. Bai C, Alverson WS, Follansbee A, Waller DM . 2012 . New reports of nuclear DNA content for 408 U.S. plant species. *Annals of Botany* . 110 : 1623-1629.
2. Baranec T, . 1979 . Karyologickéštúdiumniektorýchdrohovrodu *Pinus* L . *Biologia* (Bratislava) . : 3-13..
3. Baranec T, Murín A . 2003 . Karyological analyses of some Korean woody plants. *Biologia* (Bratislava) . : 797-804.
4. Bogunic F, Muratovic E, Brown SC, Siljak-Yakovlev S. 2003 . Genome size and base composition of five *Pinus* species from the Balkan region . *Plant Cell Reports* . 22 : 59-63.
5. Cai Q, Zhang D, Liu ZL, Wang XR . 2006 . Chromosomal localization of 5S and 18S rDNA in five species of subgenus *Strobus* and their implications for genome evolution of *Pinus* . *Annals of Botany* . 97 : 715-722.
6. Chagne D, Lalanne C, Madur D, Kumar S, Frigerio JM, Krier C, Decroocq S, Savoure A, Bou-Dagher-Kharrat M, Bertocchi E, Brach J Plomion C . 2002 . A high density genetic map of maritime pine based on AFLP's . *Ann. For. Sci.* 59 : 627-636.
7. Chang HF, Li MX . 1984 . Study of karyotypes of three species of *Pinus*.*DongbeiLinyeYanjouShuoXuebao*, . Liaoning (J. N.-E. Forest.Inst.). 15 : 115–120.
8. Chen T H, Wu RJ, Lai HL, Li KY . 1992 . A preliminary study on cytology taxonomy and system evolution of pine. *J. Fujian Coll. Forest*. 12 : 437–444.
9. Cardona, M. A. & J. Contandriopoulos. 1983 . In IOPB chromosome number reports LXXIX . *Taxon* . 32 : 323–324.
- "10. Costa P, Pot D, Dubos C, Frigerio JM, Pionneau C, Bodenes C, Bertocchi E, Cervera M-T, Remington DL, PlomionC . 2000 . A genetic map of Maritime pine based on AFLP, RAPD and protein markers.  
. *Theoretical and Applied Genetics* 2000 . 100 : 39–48.. "
11. Devey, M.E.; Sewell, M.M.; Uren, T.L.; Neale, D. B. 1999 . Comparative mapping in loblolly and radiata pine using RFLP and microsatellite markers . *Theoretical and Applied Genetics* . 99 : 656-662.
12. Dhillon SS. 1980 . Nuclear volume, chromosome size and DNA content relationships in three species of *Pinus* . *Cytologia* . 45 : 555-560.

13. Dhillon SS. 1987 . DNA in tree species. In Cell and tissue culture in forestry (Bouga JM, Deuzan DJ, eds) . : 298-313.
14. Dhillon SS, Berlyn GP, Miksche JP. 1978 . Nuclear DNA content in populations of *Pinus rigida*. American . Journal of Botany . 65 : 192-196.
15. Doudrick RL, Heslop-Harrison JS, Nelson CD, Schmidt T,NanceWL,Schwarzacher T . 1995 . Karyotype of Slash Pine (*Pinus elliottii* var. *elliottii*) using patterns of fluorescence in situ hybridization and fluorochrome banding . Journal of Heredity . 86 : 289-296..
16. Druskovic, B. & M. Lovka. 1995 . IOPB chromosome data 9. Int. Organ. Pl. Biosyst. Newslett. (Zurich) . 24 : 15–19.
17. Echt C, Saha S, Krutovsky K, Wimalanathan K, Erpelding J, Liang C . 2011 . Anannotated genetic map of loblolly pine based on microsatellite andcDNA markers. BMC Genet . 12 : 17.
18. Funabiki K, Seido K, Takahashi K, Sato T . 1970 . A study on chromosome morphology of *Pinusthunbergii* and *P. densiflora*. Rep. For. Fac. Agr. Niigata Univ. 5 : 11-16.
19. Greilhuber J. 1986 . Severely distorted Feulgen DNA amounts in *Pinus* (Coniferophytina) after non-additive fixations as a result of meristematic self-tanning with vacuole contents. Canadian Journal of Genetics and Cytology . 28 : 409-415.
20. Grotkopp E, Rejmanek M, Sanderson MJ, Rost TL . 2004 . Evolution of genome size in pines (*Pinus*) and its life-history correlates: supertree analysis. Evolution . 58 : 1705-1729.
21. Gurzenkov N N . 1973 . Studies of chromosome numbers of plants from the south of the Soviet Far East . Komarov Lectures . 20 : 47–61.
22. Hall SE, Dvorak WS, Johnston JS, Price HJ, Williams CG. 2000 . Flow cytometric analysis of DNA content for tropical and temperate New World pines. Annals of Botany . 86 : 1081-1086.
23. Hizume, M. 1988 . Karyomorphological studies in the family Pinaceae. Memoirsof the Facultyof Education Ehime University Series 3 . Natural Science . 8 : 1-108.
24. Hizume M, Ishida F, Murata M. 1992 . Multiple locations of the rRNA genes in chromosomes of pines, *Pinus densiflora* and *P. thunbergii* . Japanese Journal of Genetics . 67 : 389–396.
25. Hizume M, Ohgiku A, Tanaka A . 1989 . Chromosome banding in the genus *Pinus* II. Interspecific variation of fluorescent banding patterns in *P. densiflora* and *P. thunbergii*. Botanical Magazine (Tokyo) . 102 : 25–36..
26. Hizume M, Shibata F, Matsusaki Y, Garajova Z . 2002 . Chromosome identification and comparative karyotype analysis of *Pinus* species. Theoretical and Applied Genetics . 105 : 491-497.

27. Hizume M, Shibata K, Maruyama Y, Kondo K . 2000 . Chromosomal localization of telomere sequence repeats in five gymnosperm species . *Chromosome Science* . 4 : 39-42 .
28. Hizume M, Shibala F, Murayama Y, Kondo T . 2001 . Cloning of DNA sequences localized on proximal fluorescent chromosome bands by microdissection in *Pinus densiflora* Sieb. Zucc. *Chromosoma* . 110 : 345-351.
29. Ho RH, Owens JN. 1947 . Microsporogenesis and pollen formation in lodgepole pine. *Canadian Journal of Botany* . 52 : 1669-1674.
30. Islam-Faridi MN, Nelson CD, KubisiakTL . 2007 . Reference karyotype and cytogenetic map for loblolly pine (*Pinus taeda* L.) . *Genome* . 50 : 241-251.
31. Jacobs MD, Gardner RC, Murray BG . Cytological characterization of heterochromatin and rDNA in *Pinus radiata* and *P. taeda* . *Plant Systematics and Evolution* . 223 : 71-79.
32. Joyner KL, Wang XR, Johnston JS, Price HJ, Williams CG . 2001 . DNA content for Asian pines parallels New World relatives. *Canadian Journal of Botany* . 79 : 192-196.
33. Kaya Z, Ching KK, Stafford SG . 1985 . A statistical analysis of karyotypes of European black pine (*Pinus nigra* Arnold) from different sources . *Silvae Genet* . 34 : 148–156.
34. Kedharnath S, Upadhyaya LP. 1967 . Chiasma frequency in *Pinus roxburghii* S arg. and *P. elliottii* Engelm. *Silv.Genet*. 16 : 112-113.
35. Khoshoo TN . 1961 . Chromosome numbers in gymnosperms . *SilvaeGenetica* . 10 : 1-9.
36. Komulainen P, Brown GR, Mikkonen M, Karhu A, Garcia-Gil MR, O'Malley D, Lee B, Neale DB, Savolainen O . 2003 . Comparing EST-based genetic maps between *Pinus sylvestris* and *Pinus taeda*. *Theoretical and Applied Genetics* . 107 : 667–678.
37. Kormuták A, . 1975 . Karyological structure of some *Pinus* species. *Biologia (Bratislava)* . : 545-550.
38. Kormuták A, Matúová R, Szmidt A, Lindgren D. 1993 . Karyological, anatomical and restriction fragment length polymorphism characteristics of the interspecific hybrid *Pinus banksiana* X *Pinus contorta*. *Biologia (Bratislava)* . 48 : 95–100.
39. Kim YY, Choi HS, Kang BY. 2005 . An AFLP-based linkage map of Japanese red pine (*Pinus densiflora*) using haploid DNA samples of megagametophytes from a single maternal tree. *Mol Cells*. 20 : 201-209.
40. Kumar S, Bansal HC, Singh D, Natarajan AT . 1966 . Consistency of karyotypes and classification and identification of chromosomes in the genus *Pinus*. *Ind. Jour. Genet. & Pl. Br.* 26 : 311-316..
41. Kuroki Y . 1969 . Karyotype studies on important conifers . *Bull. Miyazaki Univer.* : 103(Japanese).

42. Löve A, Kjellqvist E . 1972 . Cytotaxonomy of spanish plants. I. Introduction. Pteridophyta and Gymnospermae . *Lagascalia* . 2 : 23-35.
43. Love A, Love D . 1980 . In Chromosome number reports LXVIII . *Taxon* . 29 : 535.
44. Macpherson P, Filion WG . 1981 . Karyotype analysis and the distribution of constitutive heterochromatin in five species of *Pinus* . *Journal of Heredity* . 72 : 193–198.
45. Mariette S, Chagné D, Lézier C, Pastuszka P, Raffin A, Plomion C, Kremer A . 2001 . Genetic diversity within and among *Pinus pinaster* populations:comparison between AFLP and microsatellite markers . *Heredity* . 86 : 469–479.
46. Marie D, Brown SC . 1993 . A cytometric exercise in plant DNA histograms, with 2C values for 70 species. *Biology of the Cell* . 78 : 41-51.
47. Mehes-Smith M, Nkongolo KK, Michael P . 2011 . A comparative cytogenetic analysis of five pine species from North America, *Pinus banksiana*, *P. contorta*, *P. monticola*, *P. resinosa* and *P. strobes*. *Plant Systematics and Evolution* . 292 : 153-164 .
48. Mehra, P. N. 1988 . Indian Conifers, Gnetophytes and Phylogeny of Gymnosperms. Panjab University . : .
49. Mehra PN, Khoshoo TN . 1956 . Cytology of conifers I . *Journal of Genetics* . 54 : 165-185.
50. Miguel M de, Maria N de, Guevara MA, Diaz L, Sáez-Laguna1 E . 2012 . 2012RsAnnotateds genetic linkage maps of *Pinus pinaster*Ait. from a Central Spain population using microsatellite and gene based markers . *BMC Genomics* . 13 : 527.
51. MoromizatoS . 1972 . 1972Karyological studies on the pine species having two or three needles in a facule belonging to the genus *Pinus*. *Sci. Bull. Coll. Agr. Ryuku Univ.* 22 : 591-678 (Japanese)..
52. Muratova. 2005 . Caryological and cytogenetic studies of coniferous plants of Siberia and Far East. *Sibirskii Èkol. Žur.* 12 : 573–582..
53. Muratova. 2000 . Karyotypic variability and anomalies in populations of conifers from Siberia and the Far East. Pages 129–141 in H. Guttenberger, Ž. Borzan, S. E. Schlarbaum& T. P. V. Hartman (eds), *Cytogenetic Studies of Forest Trees and Shrubs---Review, Present Status, and Outlook on the Future* . Arbora Publishers . : 129–141 .
54. Murín A. &Májovský J., . 1978 . Löve Á. (ed.), IOPB chromosome number reports LXI . *Taxon* . 27 : 4.
55. Natarajan, G. 1978 . In Löve, A. (ed.), IOPB chromosome numbers reports LXII]. *Taxon* . 1.125 : 526-531.
56. Naydenov, K., F. Tremblay & P. Ganchev. 2003 . Karyotypic diversity in European black pine (*Pinus nigra* Arn.) from Bulgarian Provenances . *Phyton (Horn)* . 43 : 28-Sep.

57. Nelson C D , Nance WL, Doudrick RL, . 1993 . A partial genetic linkage map of slash pine (*Pinus elliottii*Engelm. var. *elliottii*) based on random amplified polymorphic DNAs . Theoretical and Applied Genetics . 87 : 145-151..
58. Neves G L, Davis JM, Barbazuk WB, Kirst M . 2014 . A high-density gene map of loblolly pine(*Pinus taeda* L.) based on exome sequence capture genotyping . G3 (Bethesda) . 4 : 29-37.
59. Nkongolo KK, Kim NS, Michael P . 2004 . Detection and physical mapping of the 18S-5.8S-26S rDNA and the pKF1660 probe with microsatellite sequences drived from the rice blast fungus (*Megna porthe grisea*) in conifer species . Hereditas . 140 : 70-78 .
60. Nunes JD, Torres GA, Davide LC, de Campos JMS . 2009 . Chromosome banding and DNA content in tropical *Pinus* species. ScientiaForestalis, . Piracicaba . 37 : 213-218..
61. O'Brien IEW, Smith DR, Gardner RC, Murray BG. 1996 . Flow cytometric determination of genome size in *Pinus*. Plant Science . 115 : 91-99.
62. Oda, Y., K. Kawabata & K. Nagano. 1982 . Karyological studies in some species of Pinaceae. Proc. Fac. Agric. Kyushu Tokai Univ . 1 : 7-16..
63. Ohri, D. & T. N. Khoshoo . 1986 . Genome size in gymnosperms. Plant Systematics and Evolution . 153 : 119–132..
64. Pederick LA . 1970 . Chromosome relationships between Pine species . Silvae Genet . 19 : 171-180..
65. Pellicer J, Estiarte M, Garcia S, Garnatje T, Penuelas J, Sardans J, Valles J. 2010 . Genome size unaffected by moderate changes in climate and phosphorus availability in Mediterranean plants. African Journal of Biotechnology . 9 : 6070-6077.
- "66. Plomion C, Bahrman N, Durel CE, O'Malley DM . 1995 . Genomic mapping in *Pinus pinaster* (maritime pine) using RAPD and protein markers.  
. Heredity . 74 : 661–668. "
- "67. Ritter E, Aragones A, Markussen T, Achere V, Espinel S, Fladung M, Wrobel S,  
Faivre-Rampant P, Jeandroz S, Favre JM  
. 2002 . Towards construction of an ultra  
high density linkage map for *Pinus pinaster*.  
. Annals of Forest Sciences . 59 : 637–643.. "
68. 68. Rake AV, Miksche JP, Hall RB, Hansen KM . 1980 . DNA reassociation kinetics of four conifers. Canadian Journal of Genetics and Cytology . 22 : 69-79.

69. Remington D, Whetten R, Liu BH, O'malley D . 1999 . Construction of an AFLPgenetic map with nearly complete genome coverage in *Pinus taeda*. *Theoretical and Applied Genetics* . 98 : 1279–1292.
70. Renfroe MH, Berlyn GP . 1984 . Stability of nuclear DNA content during adventitious shoot formation in *Pinus taeda* L. tissue culture. *American Journal of Botany* . 71 : 268-272..
71. Ribeiro M, Mariette S, Vendramin G, SzmidtA, Plomion C, Kremer A . 2002 . Comparison of genetic diversity estimates within and amongpopulations of maritime pine using chloroplast simple-sequence repeatand amplified fragment length polymorphism data. *Molecular Ecology* . 11 : 869–877.
72. Santamour FS . 1960 . New chromosome counts in *Pinus* and *Picea*. *Silvae Genetica* . 9 : 87-88.
73. Sax K, Sax HJ . 1933 . Chromosome number and morphology in conifers . *Journal of Arnold Arboretum* . 14 : 356-375.
74. Saylor, LC . 1972 . Karyotype analysis of the genus *Pinus* - Subgenus *Pinus* . *Silvae Genetica* . 21 : 155-163.
75. Saylor, L C . 1983 . Karyotype analysis of the genus *Pinus* subgenus *Strobus* . *Silvae Genet.* 32 : 119–124.
76. Shepherd M, Cross M, Dieters MJ, Henry R . 2003 . Genetic maps for *Pinus elliottii* var. *elliottii* and *P. caribaea* var. *hondurensis* using AFLP and microsatellite markers . *Theoretical and Applied genetics* . 106 : 1409-1419.
77. Shibata K,Ogoshi Y, Nakata G . 1956 . Cytological studies on coniferae I. chromosome numbers of some species of *Chamaecyparis*, *Cryptomeria* and *Pinus*. *Kromosomo* . 29 : 1025-1028 (Japanese)..
78. Siljak-Yakovlev S, Pustahija F, Šolic EM, Bogunic F, Muratovic E, Bašić N, Catrice, O, Brown SC. 2010 . Towards a Genome Size and Chromosome Number Database of Balkan Flora: C-Values in 343 Taxa with Novel Values for 242 . *Advanced Scienc* . .
79. Styles BT, Khosla PK . 1975 . Reports [In Löve, A. (ed.), IOPB chromosome numbers reports XLVII] . *Taxon* . 24 : 145.
80. Tanaka R, Hizume M . 1980 . C banding treatment for the chromosomes of some gymnosperms . *Botanical Magazine* . 93 : 167-170.
81. Taylor RL, Mulligan GA . 1968 . Flora of the Queen Charlotte Islands. Part 2.Cytological aspects of the Vascular Plants. Queen's Printer, Ottawa . : 148.
- "82. Temesgen B, Brown GR, Harry DE,Kinlaw CS, Sewell MM, Neale DB . 2001 . Genetic mapping of expressed sequence tag polymorphism (ESTP) markers in loblollypine (*Pinus taeda* L.). . *Theoretical and Applied Genetics* . 102 : 664–675. "

83. Teoh SB, Rees H. 1976 . Nuclear DNA amounts in populations of *Picea* and *Pinus* species. *Heredity* . 36 : 123-137.
84. Terasaka, O. 1982 . Nuclear differentiation of male gametophytes in Gymnosperms . *Cytologia* . 47 : 27-46.
85. Toda Y,Sotoyama S . 1972 . Variants of woods . *Iden* . 26 : 50-54.
86. Tominaga Y, Oga M. 1970 . Cytological studies on the calli of *Pinus densiflora* in vitro. *Bull. Hiroshima Agr.College* . 4 : 8-14.
87. Vendramin G, Anzidei M, Madaghiele A, Bucci G . 1998 . Distribution of genetic diversity in *Pinus pinaster*Ait. as revealed by chloroplast microsatellites. *Theoretical and Applied Genetics* . 97 : 456-463.
88. Westbrook JW, Chhatre VE, Wu LS, Chamala S, GomideNeves L, Muñoz P, Martínez-García PJ, Neale DB, Kirst M, Mockaitis K, Nelson CD, Peter GF, Davis JM, Echt CS . 2015 . A Consensus Genetic Map for *Pinus taeda* and *Pinus elliottii* and Extent of Linkage Disequilibrium in Two Genotype-Phenotype Discovery Populations of *P. taeda*. *G3 Genes Genomes Genetics* . : doi: 10.1534/g3.115.019588.
89. Wu R Q,Zeng X, Jing Y, Ren CH, Wu PS. 1995 . Karyotype analyses and fuzzy clustering on Masson pine and Japanese black pine of different seed sources. *J. Fujian Coll. Forest*. 15 : 164-169.
90. Wakamiya I, Newton RJ, Johnston JS, Price HJ. 1993 . Genome size and environmental factors in the genus *Pinus*. *American Journal of Botany* . 80 : 1235-1241.
91. Williams CG, Joyner KL, Auckland LD, Johnston S, Price HJ. 2002 . Genomic consequences of interspecific *Pinus* spp. hybridization. *Biological Journal of the Linnean Society* . 75 : 503-508.
92. Wyman J, Laliberte S, Tremblay MF. 1997 . Nuclear DNA content variation in seeds from 22 half-sib families of jack pine (*Pinus banksiana*, Pinaceae). *American Journal of Botany* . 84 : 1351-1361.
93. Yim KB . 1963 . Karyotype analysis of *Pinus rigida* . *Hereditas* . 49 : 274-276.
94. Zinnai I . 1952 . Tetraploid plants of Japanese red , *Pinus densiflora* S. et Z. discovered in transplant beds. *Jour. Jap. For. Soc.* . 34 : 185-187.
95. Zonneveld BJM. 2011 . Pine nut syndrome: a simple test for genome size of 12 pine nut-producing trees links the bitter aftertaste to nuts of *P. armandii* Zucc. exEndl. *Systematics and Evolution* . 297 : 201-206.
96. Zonneveld BJM . 2012 . Genome sizes of 172 species, covering 64 out of the 67 genera, range from 8 to 72 picogram . *Nordic Journal of Botany* . 30 : 490-502.
97. Hizume M, Kaneko K, Miyake T . 2014 . A method for preparation of meiotic chromosomes of conifers and its applications . *Chromosome Botany* . 9 : 83-88.