

LOUDON'S HORTUS BRITANNICUS .

CATALOGUE

ALL THE PLANTS

INDIGENOUS, CULTIVATED IN, OR INTRODUCED TO

BRITAIN.

PART I.

. THE LINNEAN ARRANGEMENT,

IN WHICH NEARLY 30,000 SPECIES ARE ENUMERATED :

THE SYSTEMATIC NAME AND AUTHORITY,
ACCENTUATION, DERIVATION OF GENERIC NAMES, LITERAL ENGLISH OF SPECIFIC NAMES,
SYNONYMS SYSTEMATIC AND ENGLISH or BOTH GENERA AND SPECIES,

HABIT, HABITATION IN THE GARDEN, INDIGENOUS HABITATION,

POPULAR CHARACTER, HEIGHT, TIME OF FLOWERING, COLOUR, OF THE FLOWER,

Alode of Propagation, Soil,

NATIVE COUNTRY, YEAR OF INTRODUCTION, AND REFERENCE TO FIGURES:

^

PRECEDED BY

AN INTRODUCTION TO THE LINNEAN SYSTEM,

PART II.

THE JUSSIEUAN ARRANGEMENT,

OF NEARLY 4,000 GENERA,

WITH

AN INTRODUCTION TO THE NATURAL SYSTEM, AND A GENERAL DESCRIPTION •
AND HISTORY OF EACH ORDER.

Second Edition with Additional Supplement to February, 1832.

EDITED BY J. C. LOUDON, F.L., H., G., & Z.S.

LONDON:

PRINTED FOR LONGMAN, REES, ORME, BROWN, GREEN, & LONGMAN,
PATERNOSTER-ROW.

PREFACE.

The titlepage of this Catalogue indicates how much it is in advance of all that have *^to published ; but there are **MOM Improvement!** in it not there stated, and till l' it may be requisite to give some explanation.

The enumeration of the species in the Linnaean Arrangement, and the use of figures in designating the varieties, are adopted with a view to facilitate the number them in gardens, of dried specimens in herbariums, of drawings, or of seeds. For w we have given an explanation of the Seton mode of cutting tallies (*u. xxi.*)

Emplest and best for cutting with • kaitf OH number sticks. The genera are numerately for the same purposes, and for more ready reference to the Unneau **Anad'** I the Jussieuan, and to both Arrangements from the General Index. A star *e either the number of a species, or that of a genus, indicates ih;it next in alliavu is an additional species or genu?, or several additional species or genera, in tit .ni (p. 467, to p. 490). A section (\$) indicates that the name has been changed, genus or species has undergone some alteration in the **Supplement**

In the Natural Order two numbers are placed, as totals of species (e. g. sp. 4. —) first indicates the number of species in the Catalogue, the second the total number described by botanists. An accurate idea is thus given of what additions are made to the British Homis, in any genus of plants.

be sign* used for the habits of plants (col. 3.), and those of their habitation and due le garden [col. 4.]; explained in p. vii. viii.. are improvements in botanical description by the Editor, first described in the *Enegdgpifia of Gardening* in 1822, and applied in *a fia of Plant** (Kt edit 18S9). The twenty-three varieties of habit are indicated by figures of the plants themselves ; as a tree for a tree, a shrub for a shrub, a ladder for plants of these descriptions, a grass for a grass, a bulb for a bulb, a plant firmer for an aquatic, &c, to recollect which requires no exertion of memory. A pel Mieateil by ; triangle, A, the sign of the Trinity, and therefore connected with perfection or eternity, instead of the old sign, if ; an annual remains a circle as before, c ^•, -uiiong other reasons, gardeners sow patches of annual flowers in circles ; and a high doobill circle, Q, instead of the old sign, £. The bark stove is a parallelogram, •, be considered as representing the section of a hot-house closed on all sides to make a degree of boat; the dry stove, three sides of a parallelogram, —], to main degree of heat; the green-house two and a half sides of a parallelogram, i_J, re may be considered as the section of a green-house; and the frame two sides, ss^I fillclogmn), —j, which may be supposed to resemble the section of a frame or pit. Unation will assist the reader in recollecting these signs. By combining the situation with those of habitation, G5J, 2Sk S3D U1> && one column is made to represent of two. Thus, with these natural signs of habit, which amount to twenty-four, nation and habitation, a high amount to nineteen, we have extended the power of classification of abridged botanical description from ten, the greatest number of signs entirely arbitrary, that, we believe, has hitherto been used in botanical works, and these all natural or characteristic, the number employed in this Catalogue.

The system:lic names are accented on a simple principle, which is explained in viii. The derivations of the genera are given, and the specific systematically translated, any explanatory words accompanying such translation being prior. Those names, whether of genera or species, which are commemorative, as B. honour of Sir Joseph Banks, are distinguished by having the subjoined letters in where the rest of the word is in Human, and in Roman where the rest of the word tiunksia.; those which have been applied to plants by the classic writers of antiquity distinguished by having the initial letter in Italic, as Pyrus, where the rest of the word is in Roman, anil in Roman where the rest of the word is in Italic, as Pyrm. All words, lecific, of unknown derivation, or aboriginal names, are wholly in Italic or wblan, according to the letter in which the preceding or following matter may occur. This mode of accentuation, which occurs KdruH in J82G, was first exemplified in the second volume of the *< Mag* with the intole of accentuation noted in the Catalogue, is continued in that **mag** in the AfogosiKt' *i)f Xatiiral History* not only in the scientific names of plants, but of animals and minerals,

Introductions are given to the Jnnican and Jussieuan Systems,

PREFACE.

engravings; and this circumstance, together with the important one of all the species names being literally translated, will, by giving the meaning of almost nil the terms used in botanical description, in a great measure supersede the necessity of a grammar of botany to the young gardener. The literal translation of the specific names may be considered to a certain extent, teaching him the Latin language, and the etymologies of the generic names will give him the meaning of a number of Greek words. The species of tor genus, where numerous, are subdivided into sections and subsections, which are still defined by specific characters; and so copious are the descriptive particulars after species, that we will venture to assert that the genus to which any plant belongs is known, the specific name, in a majority of cases, may be discovered by this Catalogue with the aid of a. *Sjxcw\$ PiaiUarum.*

In the popular descriptions of the Natural Orders, the medicinal properties and medical uses are slightly noticed; and the soil, propagation, and general treatment of different groups indicated; so that this part of the work, in our opinion the most valuable collection of botanical facts that has ever been brought into so small a space, may be considered an epitome of the history, uses, and culture of the whole vegetable kingdom. ^T*\\ ever has a sufficient theoretical and practical knowledge of plants and of vegetable culture to generalise on these subjects, will find in this Natural Arrangement the rudimental thing that can be said or written on botany, gardening, and agriculture.

K numerations of those species which are adapted for culture in the open air in Britain or in corresponding climates with an estimate of the quantity of ground which each tribe would occupy, are given in the Natural Arrangement, with a view to the formation of arboreturns or botanic flower-gardens; but, for an explanation of this improvement and of its value to the practical gardener, we must refer to the body of the work. (p. 19) The same proportionate enumerations will prove useful in contriving the size of the drawers, divisions, boxes, volumes, or pages, required for preserving a hortus si ecu * collections of drawings, of engravings, of seeds, or of specimens of woods.

The Supplement to the Linnean Arrangement (p. 467.) includes all the plants introduced to Britain, and figured in English botanical works, up to February, 1830; and *Additional Supplements* in continuation are intended to be printed from time to time, and sold singly, or done up with new impressions of the body of the work.

No further explanation appearing necessary in this Preface, it remains for the Editor to state that the Linnean Arrangement audits Supplement are entirely the work of Mr. GKC DON; the Natural Anangement is managed on Unit of Professor LISDLET, with additions and shemions rendered necessary by the plan of this Catalogue; and Dr. VILLK, the profound cryptogamist, assisted in arranging the Cryptogamia. Those who are acquainted with the present state of botany, and with the botanists of this country, will agree that three gentlemen better adapted for cooperating in producing a Catalogue like that submitted to the public are not to be found. All that the Editor claims for himself is a plan of the work, which he certainly considers much superior to any tiling that has preceded it. The literal translation of the specific names, the derivations of the generic names indications of derivations, and, above all, the general descriptions of the Natural Order will, he confidently expects, be found not only of great value in (Introducing the names of plants, and in recognising them at sight, both individually and in masses; but in assisting the botanical student and practical gardener to acquire some knowledge of their structure, physiology, analogies, affinities, properties, uses, and culture.

Jayswater, May, 1830.

J. C. L.

THE first Additional Supplement, enumerating all the new plants introduced to Britain and made known there, with the scientific changes in their names, &c., up to February, 1830 is now (August, 1832) published. In it, the genera are arranged alphabetically, for ready reference from the body of the work, and from the first Supplement (p. 467.); and I hope to save the necessity of a second index. All the genera in the body of the work, and in the first Supplement (p. 467.), to which there are any species added, or in which any alterations are made, in the first Additional Supplement (p. 577. to p. 602.), are marked with a star (*); and the same will be done, as future Additional Supplements are published.

Whoever wishes to ascertain the additions and improvement! made subsequently to the last Additional Supplement may consult, the *Gardenia** Magazine*; in which, under article which will be henceforth contained in every number, until we, "Floriculturist." Botanical Notices of New Plants, and of Old Plants of Interest, *Supplementary* to the Editions of the *Encyclopaedia of Plants*, and of the *Hortus Salmia*, will be found the name of every plant newly introduced or originated, and of every recent improvement in botanical nomenclature.

CONTENTS.

„ „, «nd General Introdm.-	- Page til	'List of Books referred to
• intw used in the I/mmean Arrangement	vii	Method* of naming and numbering Plant*
I/littlf» fo' Pronouncing HnUinieal Names	viii	Drying Plant* and forming a Herbarium
, borttie* f< Generic and Specific Name*	ix	Drawing Plants, Flowers, Fruits, to

LINNEAN ARTIAN REMEMT.

Introduction to the Linnean Arrangement Table of Linnean Classes.

	specie*	1	ia, genera and specie*	m
	ries	4	ia, genera and specie*	
	specie*	B	ti, genera nnd specie*	
		12	nutria, genera and 6pcie*	
		13	TAOCLrau	296
			ectindriit genera and tpe es	318
		14	Palpifulliritt, genera and specie*	331
		23	OKNKN	
			JBqyutis, geHera and »ccica	aae
			SupJrtun, genera and *ivcie*	sn
			Frusltransa, genera and species	as
		37	ifeeatMa, genera atid specie*	
			SegT't, genera and »jocie»	
			M<mdndr/a, genera and specie*	
		t3	THi'mdria, genera and specie*	373
		US	Ht-iiitdrta, genera and specie*	37+
	Lurnandape		CTrIA	
	iera and species		• uiliia, genera and species	37*
			, j:eu(Ta imd species	375
	VNraandje*		'frifiinlliit, gtoen and species	376
	a, gwieM		Telrutultia, genera and specie*	378
	nets and species		Pettlndria, genera and specie*	380
	j, genera and specie*		Heilnuria, genera anil specie*	381
	Jl, genera and species		Oetandria, genera and »Hxie*	Saa
	rin, genera and tpccie		Icotndria, gynra nnd specie*	
	fa, gaoen ami species		Pvlyt'iulria, gcnra and specie*	389
	, genera and specie*		AIIK tdiiph ia, gei icra a nd species	BBS
	ma, genera and upctiei		DUE CIA	387
			•, genu* and specie*	393
	nia, genera and species		Di'iidria, gtnera and species	
	fttii, genera and specie*		Tritiinrfia, genera ami specie*	
	genera and attecie*		indr, genera and sjecie*	
	nia, genera and species		j'ttiiiiilrin, genera nnd sp'ies	89S
	Munwf/niti, genera and species		Ilxritulria, genera and specie*	
	ynia, genets iud «pecie*		Qett'mlria, gentu anj specie*	
	nera and specie*		thsiitii^ genera and specie*	
	• HI A		Deci'mdria, genera and species	
	•>aynia, genera and i		Ti, gHK'a and specie*	
	Trigfinla, genera aid i		Jeotii'ilrin, >T)iera and specie*	
	.iftiiUu, genera suid specie*		ia, genera anil apectei	
	iRragynia, genera unil »i's		MaiusdvipMa, genera and species	403
	■H-HIA			
	Monogijnia, genera and species		Manofcia, genera and species	- 405
	mid specie*		CtoVfe, genera and *pecie*	
	i-iicra and specie*		FVieei	
	: -ra and specie*		PalifspodiUcm, genera and Bpcries	+17
	- fd species		Otmumt. ecu , Kf- ³ⁿ - ³ⁿ -d *p*cl ei	
			nera arid spi-i-10*	
			Dj, genera mid <MHM	
			PuroptJridct, Ktnera and spevies	
			:i,.., ind -p. II[at	
			iMCoptxtntir, genera and *pecie*	
			ilartilchaea, genera and species	
			Musci	
			A'ilomi, genera and specie*	
			Gytonositmi, genera and species	
			Apioperthshni, genera and specie*	
			H, Kenyra and species	
			Schutocurjn, Ktnera nnd dMCC*	
			'AtKir, genera and specie*	
			At'gm	
			Mathtne>T, genera and qweic*	
			Kostoeikina, genns and specie*	
			*ifen>aid*ri, genera and ipeciet	
			ice*, genera and specie*	
			Tloridea, genera and specie*	
			FucaUU'C, genera and specie*	
			LtekM	
			Idiotktlam, genera and *pecl«	
			•itithUmi, genera and *pecie*	
			Ih.mothiitiam, goner* and *pecies	
			AtfitUarni, genus and specie*	
			goDera and specie*	
			-iiumtnomycifes, genera and tpecies	
			PyreiumiyiAtes, genera and specie*	
			ÜttercomueiUs, genera and specie*	
			Coiuottcyctei, genera and specie*	

CONTENTS.

NATURAL ARRANGEMENT.

Explanatory Introduction	- 491
Introduction to the Juiucuean System	- 41S

VASCULARE &

UICOTYLEUD.VEjK.		<p>Combret&ceffi VtwhysietD- Khizo i:oreic Lopbirrn? Onugriiria; Hafot'tgeto M LyOirariw or Sili-iirim Tun*ri*dne<t 111-List oin i c e a Alinagi&ui PliiliittOliihca Mvrilicate CucurUUoac Passiforcni IjOlLieie Tiinieri&eise PorMi)4cC9</p> <p>K III U M V C T H I I . u CraMiiAoeM mr OpuntiJwetc</p> <p>Gross uLtrita; Kscail..iiic</p> <p>Cuuoni&ccff Unibell irerie Arnlift'ose t iipriiYiliEiwe Lor&niheu;</p> <p>!i]Mjirt^ Calysternt i si tit</p> <p>Stylicda; (iaodcnivtic iiiulikeoa: i</p> <p>triceie JVnciLcce Lvp'UTulfEE liloctiiaeai</p> <p>Styracinea; MyinofB Sap6tex Kltoniivs Ilri-xidx Oteiiffi</p> <p>Ja*mineat <tr^cltnca! Apocjnete Aiclepltdwa CietitiincK Bipooniftcon:</p> <p>Pcdaltnaa Seakinen Pol Diu on i c e * HydroU-Jt-en; CiuvutvulAcoe</p> <p>Roragiilics Cord&xowg Hvilropliflea:</p> <p>Scrophulfrina! Labl^Ue "VcrbcniccK</p> <p>5)3 Myop6rins 513 &csntMoeM 513 Ornlaaichem 513 I^milinlaria; B Primuaccir (nlobuSirire 514 I'lumbaKfneB 514 I'dllt:tV'IIK13 6W Nytiitginac 514 Amarantli^cea; 515 Phjtnlftnia 515 Illi-noptrtOK HogonAcsc Polygbnea Lauriott 516 I>ri>iic<as 511J ProtV-pa? 516 Ttymelje** Hfi ^/ajrriueit SantalilicWd; i mic</p> <p>517 Cythlen I'lijihnrlijVccoi 617 ADtdtaBKfl 617 UJmftccan SIS I'pfriirfaj Sit> 519 tteeat MI> II.iatielidi> ComftTb Erapctrwc</p> <p>330 ,i Ci)TYLEUHINEC</p> <p>Cycildra" Hydrochartideas Ali-riiict'iie Itufbtneae</p> <p>ass</p> <p>M (Jreliidwe Scitf nu ncie t'itiica; llus&ceo; Iriilliu BM litt'iiioiJorieea. 5M Hypoxidvse 524 AinHrylidcc H cmeroCilHdoe 524 Dioscoritx- 585 T..mea 5f5 smil'icea 525 Asphodeiete 525 Tuhpacea; me Mt'laiitlifftw ass BromelibcAee Poutedtres sen Comtelinec 52* Hera TJphinos</p> <p>1</p> <p>* f</p> <p>5 1</p> <p>50J</p> <p>7j</p> <p>^ '4</p> <p>54JI</p> <p>5^p</p> <p>* ^ A B</p> <p>54^p</p> <p>54 fl</p> <p>5411</p> <p>541 {</p> <p>54J 1</p> <p>541</p> <p>541</p> <p>544 AFITV'LLX.</p> <p>A1<e</p> <p>Iliction<</p> <p>Fingi</p>
FoiuVut.		<p>LyeopoJfnwt ^irj-ilticffl</p> <p>Miisci</p> <p>Hcpbtic*</p> <p>Genera) Index</p> <p>First Additional Supplemcit</p>
FiUet		<p>54t</p> <p>544</p> <p>544</p> <p>544</p> <p>546</p>
Equitetice*		<p>544</p> <p>544</p> <p>544</p> <p>544</p> <p>577</p>

CELLUta'RES,

FoiuVut.		<p>LyeopoJfnwt ^irj-ilticffl</p> <p>Miisci</p> <p>Hcpbtic*</p>
FiUet		<p>54t</p> <p>544</p> <p>544</p>
Equitetice*		<p>544</p> <p>544</p> <p>544</p>
	1	<p>AFITV'LLX.</p> <p>A1<e</p> <p>Iliction<</p> <p>Fingi</p>
	1	<p>544</p> <p>545</p> <p>*** I</p>

TABLE OF ABBREVIATIONS AND CHARACTERS.

COLUMN 2. English Name.		COLUMN & Height.		bar. he. barren heaths, bar. pa. barren pastures, ba. s. p. barren sandy places, bog. h. boggy heaths, bog. pi. boggy places.	
bast	bastard,	ftd	fruited.	fit floating.	
hg	bearing.	hyb.	hybrid.		
bran.	branched.	lk	like.		
brct	brauted.	lvd	leaved.		
brd	berried.	sctd	scented.		
cal	calyxed.	shpd	shaped.		
eld	colored.	spkd	spiked.		
dble	double.	sund	spined.		
flwd	flowered.	stkd	stalked.		
flwg	flowering.	stmd	stemmed.		
fmd	formed.	stmd	stamened.		
fr.	fringed.	wgd	winged.		
COLUMN 3. Habit.		COLUMN 7. Time of Flowering.		all sea wet w	
3f	Deciduous tree.		ja January,	s September.	
2	Evergreen tree.		f February,	o October.	
1	Palm tree.		mr March,	n November.	
* Deciduous shrub.			ap April	d December.	
<i> Evergreen shrub.			my May.	•P Spring.	
Jt deciduous undcr.shrub.			in June,	su Summer,	
tt. Evergreen under-shrub.			jl July,	aut Autumn.	
\$_ Deciduous twiner, ligneous or herbaceous.			au August	Winter.	
% Evergreen twiner, lig. or herb.					
X Deciduous climber, lig. or herb.					
fi Evergreen climber, lig. or herb.					
-* Deciduous trailer, lig. or herb.					
JU Evergreen trailer, lig. or herb.					
^ Deciduous creeper, lig. or herb.					
Xn Evergreen creeper, lig. or herb.					
^ Deciduous herbaceous plant -					
£ Evergreen herbaceous plant					
nb Grass,					
tf Bulbous plant					
ft Fusiform-rooted plant					
A Tuberous-rooted plant					
* Aquatic.					
E* Epiphyte.					
COLUMN 4. Duration and Habitation.		COLUMN 8. Color of the Flower.		bar. he. barren heaths, bar. pa. barren pastures, ba. s. p. barren sandy places, bog. h. boggy heaths, bog. pi. boggy places.	
A" Perennial		Ap apetalous.	Li lilac.	bo. me. boggy meadows,	
Q> Bicnn'l.		JERug a>rugi.	Lu lurid.	bor. fi. borders of fields,	
O Annual.		nous.	0 orange.	br. branches,	
• Bark, or moist, stove.		B blue.	Och ochraceous	bu. fl. bushy fields,	
C Dry stove.		Bd blood.	01 olive.	bu. hi. bushy hills,	
I_J Green-house.		Bh bluish.	Oliva olivaceous.	bu. pl. bushy places,	
I Iraape.		Bk black.	P purple.	cal.ba. calcareous banks,	
GH Bark stove perenniaL		Bksh blackish.	Pa pale.	cal.ro. calcareous rocks,	
Z'! Dry stove perenniaL		Br brown.	Pk pink or	ch. ba. chalky banks,	
iA] Green-house perenniaL		Bri brick-co-	rose.	ch. cl. chalky cliffs,	
.AJ Frame perenniaL		lored.	PI pellucid,	ch. fi. chalky fields.	
IS Bark stove bienniaL		Brownish.	R red.	ch. hil. chalky hills.	
j Hi Dry stove bienniaL		Bsh bluish.	Ro rosy.	ch. pa. chalky pasture!.	
' OI Green.house bienniaL		Bt bright	Rsh reddish,	ch. so. chalky soil.	
HI Frame bienniaL		C crimson.	Ru rufous.	ch. wo. chalky woods.	
ED Bark stove annual		Css caesious.	Rus russet.	clov. fi. clover fields.	
22 Dry stove annuaL		Ch chestnut	Rust ruaty-co-	clt. gr. cultivated ground.	
iQ] Green-house annual		Cin cinereous	lored.	cor. fi. corn fields.	
.Q] Frame annual		Cop copper-co.	rose.	dit. ditches.	
		Crea cramcam-co-	PI pellucid,	dit. ba. ditch banks.	
		lored.	R red.	d. m. pL dry mountainous places.	
		D dark.	Ro rosy.	dr. co. dry commons.	
		Din dingy.	Rsh reddish,	dr. fl. dry fields.	
		DI dull	Ru rufous.	dr. he. dry heaths.	
		Dp deep.	Rus russet.	dr. pa. dry pastures.	
		F flesh.	I Tan tan-color-	dr. wo. dry woods.	
		Fer ferruginousl	rent.	d. st pL dry stony places.	
		Fi fiery.	Taw tawny.	d. st w. dry stony woods.	
		Fla flame-co-	(Test testaceous.	dung. dunghills.	
		lored.	I Tran transparent.	ed. of d. edges of ditches.	
		Ful fulvid.	I Umb umber-co-	gra. ba. gravelly banks.	
		Fus fuscous.	I lored.	gra. he. gravelly heaths,	
		G green.	V violet.	gra. pa. gravelly pastures,	
		Gl glaucous.	Va variegated.	gra. so. gravelly soil,	
		Go golden.	Ve vermillion.	hea. heaths,	
		Gsh greenish.	Vy veiny.	hea. w. heaths and woods,	
		Gy grey.	W white.	hed. hedges,	
		Hoa hoary.	Wsh whitish.	hed. b. hedge banks,	
		L light	Y yellow.	highlv. Highland valleys.	
		La lake.	Y yellowish.	hil. pa. hilly pastures,	
		Ld livid.	I Yah yellowish.	ir. bog. Irish bogs,	
		Lem leraon-cld.	I	ir. mo. Irish mountains,	
				ir. roc. Irish rocks,	
				ir. sho. Irish shores,	
				ir. thL Irish thickets,	
				lak. lakes,	
				m. alp. moist alpine places.	
				mar. marshes.	
				mar.la. margins of lakes.	
				m. a.w. moist alpine woods.	
				m. c. h. moist chalky hills.	
				m. ch.s. moist chalky soil.	
				mea. meadows.	
				me. pa. meadows and pastures.	
				m. hed. moist hedges.	
				mic. ro. micaceous rocks.	
				m. me. moist meadows.	
				moi. fi. moist fields.	
				moi.gr. moist grounds.	
				moi. h. moist heaths.	
				mou. pi. moist places.	
				moi.ro. moist rocks.	
				xnol. w. moist woods.	
				mo. pi. mountainous places.	
				mos. b. mossy bogs.	
				moun. mountains.	
				m. pas. moist pastures.	
				ms.pas. mountainous pastures.	
				m. r. h. mountainous rocky heaths,	
				mrif. r. maritime rocks.	
				m. r. tr. moist rocks and trees.	
				m. s. pi. moist shady places.	
				m. thi. mountainous thickets.	
				m. wa. mountainous woods.	
				mudd. d. muddy ditches.	
				mud. s. muddy shores.	
				n. of e. north of England.	
				n. of s. north of Scotland.	
				old. w. 7. ¹¹ wa. t.	
				old. wa. j. os. hol. osier holts.	
				pas. pastures.	
				pea. d. peaty ditches.	

RULES FOR PRONOUNCING BOTANICAL NAMES.

riv.ba,	river banks,	■ so. co.	south coast	■ COLUMN 11. <i>Propagation.</i>
rivil.	rivulets,	■ s. of s.	south of Scotland,	■ B by budding.
ro. kid.	road sides,	■ p. bo.	spongy bogs,	■ C cuttings.
rub.	rubbish,	■ sta. wa.	stagnant water,	■ D division of the plant
s&hea.	sandy heaths,	■ tin w.	stones in water,	■ O grafting.
sal. m.	salt marshes,	ato. hi.	stony hills,	■ I inarching.
sa. ma.	sandy marshes,	sta pa.	stony pastures,	■ L layers.
san. fi.	sandy fields,	sto. pi.	stony places,	■ Ls leaves.
san.gr.	sandy ground,	sta wa.	stones and walls.	■ O offsets.
•an. pi.	eandy places,	sun.U	sunny hills,	■ R division of the root
san. sh.	sandy shorea	sun./a	sunny rocks,	■ S seeds.
sa. pas.	sandy pasture,	thick,	thickets,	■ Sk suckers.
saw. d.	salt-water ditches.	tr.	trees,	
saw. p.	Bandy wet places.	tr. & st	trees and stones,	
sc. alp.	Scottish alps.	tru. tr.	trunks of trees,	■ COLUMN 12. <i>Soil.</i>
scbog.	Scottish bogs.	tur. bo.	turfy bogs,	■ aq. watery places,
sc.Ul.	Scottish islands,	tur.he.	turfy heath,	■ ca common garden soil,
s. cliffs	sea cliffs.	unc. gr.	uncultivated ground,	■ cp. common peat, or bog,
sc. ma.	Scottish marshes.	unc. pi.	uncultivated places,	■ h. heavy rich clay.
sc. ma.	Scottish mountains.	wa. gr.	waste ground,	■ hL heavy loam.
sapas.	Scottish pastures.	w.al. h.	wet alpine heaths,	■ L loam.
sc. roc.	Scottish rocks.	w. alp.	Welsh alps,	■ Lp. loam and peat, most loam.
sa. sh.	Scottish shores.	watco.	watery commons,	■ It light vegetable soil
scthi.	Scottish thickets.	wat pL	watery places,	■ itL light loam.
gawds.	J Scottish wood*.	w. bog.	Welsh bogs,	■ m.s. moist soil.
sc. wo.		w. co.	wet commons,	■ p. peat
sea co.	sea coast	w. gr.	wet ground,	■ p.!. peat and loam, most peat
sea sh.	sea shore.	w. lak.	Welsh lakes,	■ r. rich garden soil,
sev. isl.	Severn isles.	w. roc.	Welsh rocks,	■ r.m. rich mould,
sha.ba.	shady banks.	w.s.a. p.	wet sandy places,	■ ru. rubbish,
Bha.bo.	shady bogs.	w. s. gr.	wet shady ground,	■ s. sand.
sha. la.	shady lanes.	w.sh. p.	wet shady places,	■ s.I. sandy loam.
•ha. pl.	shady places,	wy.sh.p.	watery shady places,	■ s.p. sandy peat
sh. roc.	shady rocks.	w. thi.	wet thickets.	■ sp.L sand, peat, and loam.
s. m. pL	shady moist places.			

Accents.—In the first sixty-two pages the acute (') is used to denote accented syllables merely; after p. 62, it is restrained to such vowels as are sounded short, while the grave (") is placed over those sounded long, as M&rtha, Mary. The systematic names are distinguished as classical, i. c. names applied to plants by the ancients, by the first letter being in Italic, as *jfibies*; as commemorative, by the terminating letter or letters being in Italic, as Banks/a; and as aboriginal, or of uncertain derivation, by the whole word being in Italic, as *JF/rua*. All the other names are formed, in almost every case, from the Greek and Latin.

RULES FOR PRONOUNCING BOTANICAL NAMES.

BYLLABLER

IN classical words there are as many syllables as there are vowels; except when *u* with any other vowel follows *g*, *g*, or *u*, and when two vowels unite to form a diphthong. The diphthongs are *<e*, *ce*, *at*, *ei*, *oi*, *ui*, *au*, *ru*, and *on*. These seldom coalesce in final syllables. *oo*, *ee*, *ca*, and other combinations which never occur as diphthongs, in classical words, follow, in commemorative names, the pronunciation of their primitives, as *Teddfi*, *Wd&dsio*.

VOWELS.

In this work, with the exception of the first sixty-two pages of the Linnean Arrangement, the sounds of the accented vowels are indicated by the mark placed over each, the long sound by a grave accent ("), the short by an acute ('); but, as in the abovementioned sixty-two pages the acute accent alone is employed, the following observations will be found useful in showing when the vowel is to be sounded long, and when short:

Every accented penultimate vowel is pronounced long, when followed by a vowel or a single consonant, as *Ach'iWka tomenthsa*; but it is shortened when followed by two consonants or a double one, as *56rbus*, *Taxus*; except when the first consonant is a mute and the second a liquid, as *AHws*.

Every accented antepenultimate vowel, except *u*, is pronounced short, as *Jelleborus*, *Jftimulus*; but when succeeded by a single consonant, followed by *e* or *i* and another vowel, it is lengthened, as *Stellaria*; except 1, which is short, as *7ilia*.

A unaccented, ending a word, is pronounced like the interjection *ah*, as *SMcta /ah/*.

f final, with or without a consonant preceding, always forms a distinct syllable, as *Sflelio*, *Aide*; also when the vowel is followed by a final consonant not *Trichoma-nes*, not *Tricho-mane*.

/ unaccented, if final, sounds as if written *eye*, as *Spica vehti /eye/*; but, when it ends a syllable not final, it has the sound of *e*, as *A/cspilus* (*Mespclus*), *Smith//* (*Smith#eye*).

y is subject to the same rules as *f*.

The diphthongs *er* and *a?* conform to the rules *fore*; *ei* is generally pronounced like *eye*; the other diphthongs have the common English sounds.

In addition to the primary accent, every word of more than three syllables contains a secondary accent, which is regulated by the same rules. The secondary accent must always be at least two syllables before the primary accent, as in *Cheflidbnium*; for its place the ear is a sufficient guide, and even were it entirely omitted, still, however inharmonious, the pronunciation would not be incorrect.

CONSONANTS.

- *C* and *a* arc hard before *a*, *o*, and *u*, as *Cbrnus*, *Calium*; soft before *e*, *1*, and *y*, as *Cetraria*, *Citrus*.

T *s* and *c*, before *ia*, *ie*, *ii*, *to*, *hi*, and *eu*, when preceded by the accent, change their sounds, *t* and *c* into *sh* as *bletuz*, *ficia*; and *s* into *zh*, as *Bfas/a*; but, when the accent is on the first diphthongal vowel, the pre-celine consonant preserves its sound, as *aurantiacum*.

Ch before a vowel, is pronounced like *k*, as *Chelulonium* (*kel*), *C6lchicum* (*kolkekum*); but in commemorative names it follows their primitives, as *Richardshnto*, in which the *ch* is soft.

Cm, *en*, *ct*, *em*, *en*, *mn*, *tm*, *ps*, *pt*, and other uncombinable consonants, when they begin a word, are pronounced with the first letter mute, as *Ptens /teris*, *t'nlcus /tukus*, *Gmellna /melina*, *6nidia /nidia*; in the middle of a word they separate as in English, as *lap-sana*, *L^ra-na*.

Ph, followed by a mute, is not sounded; but, followed by a vowel or a liquid, sounds like /, as /'hReun (*ficum*).

Sch sounds like *sk*, as &hoe'nus (*skenus*); in // and zm both letters are heard.

S, at the end of a word, has its pure hissing sound, as *Dactylis j* except when preceded by *e*, *r*, or *n*, when it sounds like %, as *Rtbes* (*ex*).

X, at the beginning of a word, sounds like 2, as *Xanthium*; in any other situation it retains its own sound, as *ftxus*, *Tfcnarix*. (*Gardener's Magazine*, vol. v. p. 333.)

LIST OF AUTHORITIES
FOR
GENERIC AND SPECIFIC NAMES.

A	<i>Albertini and Schwebit'z.</i> Writers upon Fltingi.	<i>Bb'hm.</i> <i>Boltmer.</i> A German botanical writer.
<i>Abel.</i>	<i>Abel.</i> A traveller in China, and author of a Notice of Chinese plants.	<i>Boissieu la Martiniere,</i> who accompanied La Peyrouse.
<i>Ach.</i>	<i>Acharius.</i> A Swedish professor, and writer upon lichens.	<i>Bojer.</i> A professor of botany in the Isle of France.
F	<i>F. Adams.</i> A Russian botanist, who travelled through Arctic and Eastern Siberia.	<i>Bolton.</i> An English writer on F&ngi.
<i>Adan.</i>	<i>Adanson.</i> A French systematic botanist.	<i>Bonpland.</i> A French traveller in South America, and botanist
	<i>Afzcltui.</i> A Swedish professor.	<i>Boreckhausen.</i> A writer upon the Flora of Hesse Darmstadt
<i>Ag.</i>	<i>Agardh.</i> A Swedish professor, and writer upon ijigie, &c.	<i>Bory de St. Vincent.</i> A French traveller and botanist *
<i>Ait.</i>	<i>Aiton.</i> The superintendant of the King's garden at Kew.	<i>Bosc.</i> A French botanist, and traveller in North America.
<i>Alb.</i>	<i>Albertini.</i> A writer upon Fangi.	<i>Boucher.</i> A writer upon the French Flora.
<i>All.</i>	<i>Allioni.</i> An Italian botanist	<i>F. A. G. Boucher.</i> A French botanist
<i>Anun.</i>	<i>Ammann.</i> An old Russian botanist.	<i>J. Bowie.</i> A collector of plants for Kew Gardens.
<i>Andcr.</i>	<i>Anderson.</i> A London merchant j published a paper on peonies.	<i>B. R.</i> <i>Botanical Register.</i> By Ker and Lindley.
<i>Andr.</i>	<i>Andrews.</i> A famous botanical draughts man.	<i>Br.</i> <i>Patrick Browne, M. D.</i> Author of Civil and Natural History of Jamaica. An Irish botanist
<i>Andrx.</i>	<i>Andrxjowski.</i> A Russian botanist.	<i>B. Rep.</i> <i>Botanical Repository.</i> By Andrews and others.
<i>Ard.</i>	<i>Arduini.</i> An Italian botanist.	<i>Brad.</i> <i>Bradley.</i> An old English writer upon sue culent plants.
<i>Asso.</i>	<i>Asso.</i> A Spanish botanist.	<i>Bredcmeyer.</i> A German.
<i>Aub.</i>	<i>Aul^L</i> A French traveller in Guiana.	<i>Bridel.</i> A Gennan writer upon mosses.
<i>And.</i>	<i>AudWrt.</i> A French cultivator.	<i>Brig.</i> <i>J. Brignoli.</i> Professor at Verona.
<i>Balb.</i>	<i>Balm.</i> A French professor of botany.	<i>Brm.</i> <i>Burmann.</i> A Dutch editor of other people's works.
<i>Bac.</i>	<i>Bade.</i> A German botanist ?	<i>Broegelmann.</i> A German botanist
<i>Baldw.</i>	<i>Baldwin, M. D.</i> of Savannah in Georgia A botanist	<i>Brong.</i> <i>A. Brongniart.</i> A French botanist
F	<i>Bartling, M. D., and Wendland,</i> of Gottingen, botanists.	<i>Brot.</i> <i>Brotero.</i> A Portuguese botanist
<i>BanJcs.</i>	<i>Bunks.</i> A great traveller and patron of science.	<i>Brou.</i> <i>Broussonet.</i> A French botanist, and traveller in Barbary.
<i>Jiarrl,</i> <i>Bart.</i>	<i>Barrelier.</i> A French botanist of 1714.	<i>Buc.</i> <i>Buchanan.</i> An English physician, and traveller in Nepal.
<i>Barlr.</i>	<i>Barton, M. D.</i> Formerly a professor at Philadelphia.	<i>Bui.</i> <i>Bulliard.</i> A French writer upon Fungi.
<i>Bast.</i>	<i>Bartram.</i> Formerly a nurseryman at Philadelphia.	<i>Burc.</i> <i>Burchell.</i> An English botanist, and traveller at the Cape of Good Hope.
<i>Batsch.</i>	<i>Bastard or Batard.</i> A writer upon the Flora of France.	<i>Burg.</i> <i>Burgsdorf.</i> A German botanist
<i>Batt.</i>	<i>Batsch.</i> A writer upon FGngL	<i>But.</i> <i>Buxbaum, M. D.</i> A Russian botanist; travelled through Armenia.
	<i>Battarra.</i> Published a History of Fungi, 1789.	<i>Cas.</i> <i>Casalpnum.</i> A famous old Italian botanist
	<i>J. C. Baumgarlen,</i> M. D. of Schasburgh, In Transylvania. A botanist	<i>Cam.</i> <i>Catnpna.</i> An Italian cultivator.
<i>Bauh.</i>	<i>Bavhin, brothers;</i> professors of medicine, published MRO, 1650.	<i>Campb.</i> <i>Campdera.</i> A Spanish botanist
<i>Bautn.</i>	<i>J. C. Baumgartcn, M.D., of Schasburgh, in Transylvania.</i> A botanist	<i>C.S.R.</i> <i>Castagne and Robillard.</i> French botanists ?
3 C	<i>Botanical Cabinet</i> By Loddiges and Sons.	<i>Carey.</i> <i>W. Carey, D. D., of Serapmore.</i>
<i>Beaup.</i>	<i>Beauprc.</i> A French botanist, mentioned in Dec. Systema.	<i>Capt. I.</i> <i>Carmichael.</i> A Scotch botanist
<i>Beauv.</i>	<i>Palisot dc Beauvois.</i> A French traveller and botanist	<i>H. Casfini.</i> A French botanist
<i>Bedf.</i>	<i>Duke of Bedford.</i> A great promoter of botany.	<i>L. Castagne.</i> A French botanist ?
<i>Bel.</i>	<i>Bellardi.</i> An Italian botanist	<i>M. Catesby.</i> A botanist and traveller in North America.
<i>Bcnth.</i>	<i>Bcutham.</i> An English botanist, secretary to the Horticultural Society, London.	<i>Cav.</i> <i>Cavamilles.</i> A Spanish professor and botanist
<i>Berg.</i>	<i>Bergius.</i> A Swedish writer upon Cape plants.	<i>C. Bauh.</i> <i>Caspar Bauhin.</i> A celebrated botanist of the sixteenth century.
<i>Berger.</i>	<i>J. Bergeret, M. D.</i> A French botanist	<i>Cer.</i> <i>Vicente Cervantez.</i> A Spanish botanist and professor.
<i>Bcrnh.</i>	<i>ftin/tardi.</i> A German botanist	<i>Chaix.</i> <i>Chair.</i> A French botanist and ecclesiastic.
<i>Bert.</i>	<i>Necan.</i> A writer upon the Flora of Italy, the Crisan ProfeM or, resident m	<i>Cham.</i> <i>Chamiso.</i> A German traveller round the world.
<i>Bes.</i>	<i>Bieberstein.</i> A Russian botanist of great note.	<i>Choi.</i> <i>Choisy.</i> A Swiss botanist.
<i>Bieb.</i>	<i>Bivona.</i> A Sicilian botanist	<i>Clair.</i> <i>Clairville.</i> A French botanist
<i>Bfo.</i>	<i>Rlume, M: D.</i> A Dutch botanist	<i>Clar.</i> <i>J. Clarion.</i> A French botanist
<i>Biutne.</i>	<i>Botanical Magazine.</i> By Curtis, Sims, &c.	<i>Clem.</i> <i>S. Clemente.</i> A S>>am&h agriculturist
<i>B.U.</i>	<i>Boerhaavc.</i> An old Dutch botanist	<i>Clutius.</i> An old French botanist and tra vller.
		<i>Co. a</i> <i>ColrWe's Catalogue.</i> Plants cultivated in Colville's nursery, Chelsea.
		<i>Colb.</i> <i>Colebrooke.</i> A celebrated English writer r rW*1UDian Plant*
		<i>^ ,</i> <i>J. F. Colladon.</i> A Genevese botanist.
		<i>Coll.</i>

AUTHORITIES FOR GENERIC AND SPECIFIC NAMES.

Com.	Commelin. A Dutch garden botanist	Gled.	Gleditsch. A German botanist
Corr.	Corria de Serra. A Portuguese botanist and diplomatist	Glox.	Gloxin. A botanist of Strasburgh.
Cr.	Crantz. An Austrian botanist	Gm.	Gmelin. A Russian botenist, and traveller in Siberia.
Cun.	A. Cunningham. A collector of plants for Kcw Gardens.	Goch.	Gochnat. A German botanist
Cur.	Curtis. An English writer upon plants.	Gold.	Goldie. A nurseryman at Ayr, in Scotland.
Cits.	Cusson. A Swiss writer upon Umbeliferse, whose wife burnt his herbarium.	Goldd.	Goldhack. A Russian botanist.
Cyr.	Cyrilli. An Italian botanist	Good,	Goodenough. Bishop of Carlisle An English botanist.
Dan.	Danthuine. A French botanist	Gou.	Gouan. A French botanist
Dat.	H. Davies, D. D. A Welsh botanist	Grah.	Graham, M. D. Regius professor of botany at Edinburgh. An assiduous botanist
D.Don.	David Don. Librarian to the Linnean Society.	Greu.	Greville. An English botanist, and writ" upon cryptogamic plants.
Deb.	Debry. A botanist of Frankfort	Gro.	Gronovius. A botanist of Leyden.
Dee.	Decandolle. A celebrated French system, atic botanist	Guer.	Guersent. A French botanist.
Del	Delile. A French professor, and traveller in Egypt	Gal.	GULdenstacdt. A Russian botanist
JDelan.	Delany. An English artist.	GUnt.	GUnther. A Silesian botanist
Delcss.	Delesscr. A French botanist	Habl.	Hnbltzl. A Russian botanist
Desf.	Desfontaines. A French botanist, and traveller in Barbary.	H<e.	Hanks. A German botanical writer.
Dcsps.	Dcsportes. A French botanist.	Hal.	Holler. A Swiss botanist
Desr.	Desrousseaux. A French botanist	Hal./.	Holier the younger.
Deso.	Desvaux. A French professor of botany.	Ham.	Hamilton. A Scotch botanist and traveller in the East Indies.
Deth.	Detharding. A botanist of Rostoch.	H.SfB.	Humboldt and Bonpland. Famous travellers and botanists.
Dick.	Dickson. An English cryptogamic botanist	H.QG.	Hooker, L.L.D., and Greville, L. L.D. English botanists resident in Scotland.
Die.	Dietrich. A German gardener.	H.QT.	Hooker and Taylor. English botanists.
Dil	Dillwyn. An English writer upon Con- fervas.	Har.	Har tig. A writer on the culture of forest trees.
Dit.	Ditmari. A Dutch botanist	Haw.	Haworth. An English writer upon succulent plants.
Domb.	Dmnbry. A French traveller in South America.	Hay.	Hayue. A German botanist
Dor.	Dorthes. A French botanist.	Hber.	Hortus Beroltnensis.
Don.	Douglas. A collector of plants.	H. C.	Chelsea botanic garden.
Dow.	Dowine. Not known.	H.Cels.	Catalogue of the plants cultivated in the gärden of M. Cels.
Dry.	Dryander. A Swedish botanist	Hedw.	Hedwig. A German cryptogamic botanist
Düb.	Dubois. A French botanist	Hei.	Heister. A German botanist.
Duch.	Duchesne. A French botanist or horticulturist	H.er.	Hortus Erfurtiensis. Catalogue of the plants cultivated in the botanical garden of Erfurt
Dufri.	Dufresne. A French writer upon valerians.	Her. am.	Herb. amat, by Delaune.
Duh.	DuhameL. A celebrated French physio- logical botanist	Herb.	Herbert. An English divine ana ootanist
Dum.	Dumont Courset. A writer upon French garden plants.	Herit.	VHeritier. A French botanist
DundL	Dundal. A French botanist	Herm.	Hermann. Professor of botany at Leyden.
Durand.	Durand. A French botanist	Heyne.	Heyne. A writer on East Indian plants.
Durb.	Durborough. Not known.	Hit.	Auguste St. Hilaire. A French botanist, and traveller in Brazil.
Duroi.	Du Rot. A German writer upon plants.	Hill.	Hill. An English compiler of botanical matters.
Dut.	Dutour. A Belgic botanist	U.K.	Hortus Kewensis. A catalogue of the plants growing in the King's garden at
Duval	Duval. A French botanist	H.L.	Liverpool botanic garden.
E.B.	English Botany. By Sowerby and Smith.	Hofg.	Hoffmannsegg. A botanist of Dresden.
Eddy.	Eddy. An English gardener.	Hofin.	Hoffmann. A German writer upon Um- bellifers, &c.
E. F.	English Flora. By Sir. J. E. Smith.		Hogg. A nurseryman it New York.
Ehren.	Ehrenberg. A German traveller in Arabia, &c.		Hbkert. A Swedish botanist.
Ehrh.	Ehrhart. A German botanist		Holtwell. An English botanist
El.	ELUs. London merchant and botanist		Holmskiold. A Danish botanist
Etl.	Elliot. An American botanist		Holmskiold. A Danish botanist
Exp.	Espt'r. A German writer on Fungi.		Hooker. An English botanist, and professor at Glasgow.
Et.	Ettlinger. A German writer on Salvia.		Hoppe. A German botanist, and collector of plants.
F.m.	Faucoult. A French botanist?		Hornemann. A Danish botanist and pro- fessor.
Feu.	FeuUlee. A Chilian botanist.		Homschuch. A German botanist and pro- fessor.
Fis.	Fischer. A Russian botanist		Hort. Of the gardens.
Fl.br.	Flora Briannica. By Sir James Edward Smith.		Host. An Austrian writer upon grapes and European plants.
Ft dan.	Flora Danica. By Oeder, Hornemann, and others.		Houttuyn. A botanist of Amsterdam.
Fi.gr.	Flora Gneca, by Sir J. E. Smith.		Of the Paris garden,
Fl.Ion.	Flora Londinensis. By Curtis and Hooker.		Hortus Taurinensis. Catalogue of the plants cultivated in the botanical garden at Turin.
Ft. per.	Flora Pcruviona. By Ruiz and Pavon.		Transactions of the London Horticultural Society.
Flu.	Flligge. A German writer upon grasses.		Hudson. An English writer upon British plants.
Forsk.	Forskahl. A Danish naturalist, and traveller in Arabia.		Humboldt. A celebrated Prussian traveller and philosopher.
	Forster. A traveller in the South Seas with Captain Cook.		Hunter. An English botanist
Fro.	Frazcr. A gardener, and collector of plants in North America.		Hartmann. A botanist of Stockholm,
Fries.	Fries. A Swedish botanist, and writer upon fungi		Jves. An American botanist
Fro.	Frolich. A German writer upon Gentiana.		Jussicu. A celebrated French systematic botanist.
Funk.	Funk. A German cryptogamic botanist		Jacquin. An Austrian traveller in South America, and botanist
Gae.	Gartner. A celebrated German carpolo-		Jackson. An English botanist
G.SsA.			Junghans. A botanist of Halle.
Gnu.			
Gay.			
G. J.			
Ger.			
Gilib.			
Gill.			
Gin.			

AUTHORITIES FOR SPECIFIC AND GENERIC NAMES. xi

<i>Jon.</i>	<i>Jones.</i> An accomplished writer upon Indian matters.	<i>Mayer.</i> Several German botanists of this name.
<i>Katn.</i>	<i>Kamjiffer.</i> A traveller in Japan. Not known.	<i>Medicus.</i> A German botanist of the last century.
	<i>Kitabel.</i> A Hungarian botanist	<i>Menz.</i> Menxies. A Scotch botanist, and traveller round the world with Vancouver.
	<i>KnighVs Protoc&B.</i>	<i>Merat.</i> A French botanist.
<i>Koel.</i>	<i>Koch.</i> A professor at Erlang.	<i>Mert.</i> Mertens. A French Botanist
	<i>Ifoeler.</i> A writer on German and French grasses.	<i>Mey.</i> Meyer. A German botanist
<i>Kbille.</i>	<i>Kofle.</i> A botanist of Erlang.	<i>Micheli.</i> Micheli. A Florentine botanist
<i>Kon.</i>	<i>Kbnig.</i> Several German naturalists of this name.	<i>Miers.</i> Miers. A South American collector.
<i>Kth.</i>	<i>Kunth.</i> A Prussian botanist.	<i>Mican.</i> Mi-can, A German writer on Brazilian plants.
<i>Kunxe.</i>	<i>Kunze.</i> A German cryptogamic botanist	<i>Miller.</i> Miller. An English gardener and botanist
<i>X*.</i>	<i>Linnlaus.</i> The celebrated Swedish re-former of natural history.	<i>Mirbel.</i> Mirbel. A French physiological botanist
<i>I.ab.</i>	<i>Labillardiere.</i> A French botanist	<i>Mocino.</i> Mocino. A Mexican botanist
<i>Lag.</i>	<i>Lagasca.</i> A Spanish botanist and professor.	<i>Moench.</i> Moench. A German botanist
<i>Lai.</i>	<i>La Lave.</i> A Mexican botanist	<i>Mohr.</i> Mohr. A German cryptogamic writer.
<i>Lam.</i>	<i>Lamarck.</i> A French botanist	<i>Molina.</i> Molina. An Italian writer upon the natural history of Chile.
<i>Lamb.</i>	<i>A. R. Lambert,</i> V. P. L. S. The most liberal botanist in England. Nothing in botany can well be completed without his assistance.	<i>Montin.</i> Montin. A Swedish botanist
<i>Lat.</i>	<i>Langsdorff.</i> A Russian botanist	<i>Morison.</i> Morison. An old writer on plants.
<i>Lap.</i>	<i>La Peyrouse.</i> A French writer upon the plants of the Pyrenees.	<i>Mordant de Launy.</i> Mordant de Launy. A French botanist
<i>Lar.</i>	<i>LMTLoche.</i> A French botanist.	<i>Moretti.</i> Moretti. An Italian botanist.
<i>Law.</i>	<i>Lawrence (Miss).</i> An English flower-painter.	<i>Mougeot.</i> Mougeot. A German cryptogamic botanist.
<i>ix.</i>	<i>Laxmann.</i> A German writer on Siberian plants.	<i>Muhlenberg.</i> Muhlenberg. A North American botanist
<i>Lee.</i>	<i>Leconte.</i> An American botanist	<i>Murray.</i> Murray. A German botanist
<i>Lech.</i>	<i>Lechenault.</i> A French botanist.	<i>Mussin-Puschkin.</i> Mussin-Puschkin. A Russian botanist and traveller in Caucasus.
<i>Led.</i>	<i>Ledebour.</i> A botanist and traveller in Siberia.	<i>Mutis.</i> Mutis. A Spanish botanist, resident in New Grenada.
	<i>Lee.</i> A nurseryman at Hammersmith.	<i>Michaux.</i> Michaux. A French botanist, and traveller in North America.
<i>Leers.</i>	<i>Leers.</i> A German botanist	<i>Nees and Martius.</i> Nees and Martius. German botanists.
<i>Leh.</i>	<i>Lchmann.</i> A German botanist and professor at Hamburg.	<i>Necker.</i> Necker. A German writer upon botanical affairs.
	<i>Lejeune.</i> A French botanist	<i>Nets v. Escnbeck.</i> Nets v. Escnbeck. A German botanist
<i>Lent.</i>	<i>Leman.</i> A botanist mentioned in Dec. Fl. Franc.	<i>Nestler.</i> Nestler. A botanist of Strasburgh.
	<i>Lepchin.</i> A Russian botanist	<i>Niven.</i> Niven. Once a collector of plants at the Cape.
<i>L*</i>	<i>Lepelletier.</i> A French botanist	<i>Nocca.</i> Nocca. A professor of Pavia.
<i>Lepel.</i>	<i>Lewis.</i> An American traveller.	<i>Noisette.</i> Noisette. A French nurseryman.
<i>Lewis.</i>	<i>Leysser.</i> A botanist of Halle.	<i>Noronha.</i> Noronha. A Spanish botanist who visited Madagascar.
<i>L.J.U.</i>	<i>Linnlaus the younger.</i> The son of the great Linnus.	<i>Nuttali.</i> Nuttali. A North American botanist
<i>Lich.</i>	<i>Lichtenstein.</i> A German botanist and professor.	<i>Oeder.</i> Oeder. A Danish botanist.
<i>Light.</i>	<i>Jightfoot.</i> A writer upon the Scottish	<i>Olivier.</i> Olivier. A French botanist and traveller in Persia.
<i>P.HJeb.</i>	<i>Liljeblad.</i> A professor at Upsal.	<i>Ortega.</i> Ortega. A Spanish botanist
<i>Lind.</i>	<i>Lindern.</i> A French botanist	<i>Oth.</i> Oth. A French writer in Decandolle's Prodromus.
<i>Lima.</i>	<i>Lindley.</i> An English botanist, and professor in London.	<i>Otto.</i> Otto. A Prussian gardener.
	<i>Livinston.</i> A Scotch botanist.	<i>Pallas.</i> Pallas. A Russian traveller and naturalist
	<i>Link.</i> A Prussian botanist	<i>Parmentier.</i> Parmentier. A French nurseryman.
<i>Lo. c.</i>	<i>Link and Otto.</i> Prussian botanists.	<i>Patnn.</i> Pattn. A Russian traveller.
	<i>Lobel.</i> An old writer upon plants.	<i>Perpenti.</i> Perpenti. A female Italian botanist
	<i>Loddiges* Catalogue.</i> Catalogue of the plants cultivated in Loddiges' nursery at Hackney.	<i>Persoon.</i> Persoon. A French botanist
<i>Lock.</i>	<i>Lock/tart.</i> Superintendent of the Trinidad botanical garden.	<i>Petagna.</i> Petagna. A Neajlwanian botanist
<i>Lod.</i>	<i>Loddiges.</i> English nurserymen and botanists.	<i>Pursh.</i> Pursh. A Prussian botanist, and traveller in North America.
	<i>Loefling.</i> A Swedish botanist and traveller.	<i>Plumier.</i> Plumier. A French botanist, and traveller in the West Indies.
<i>Ul</i>	<i>Loiseleur Dcslongchamps.</i> A French botanist	<i>Pohl.</i> Pohl. A German botanist
<i>Lon.</i>	<i>Londes.</i> A botanist of Gottingen.	<i>Poir.</i> Poir. A French botanical compiler.
<i>Lo.s.</i>	<i>Loddiges* Supplement.</i>	<i>Pott.</i> Pott. Poiteau. A French botanist and draughtsman.
<i>Lou.</i>	<i>Loureiro.</i> A Portuguese traveller in Cochim China,	<i>Pot.</i> Pot. Pollini. A professor at Verona.
<i>London.</i>	<i>J. C. Loudon.</i> Auth. of various agricultural, horticultural, and botanical works.	<i>Poll.</i> Poll. Pollich. A German writer on the plants of the Palatinate.
	<i>Linnean Society's Transactions.</i>	<i>Pop.</i> Pop. Poppig. A German botapist
<i>Lum.</i>	<i>Lumnitxer.</i> A Gorman botanist	<i>Pou.</i> Pou. Pourret. A French botanist.
<i>Lunan.</i>	<i>Lunan.</i> A writer on Jamaica plants.	<i>Presl.</i> Presl. A Bohemian botanist
<i>Lyell.</i>	<i>Lyell.</i> A Scotch botanist	<i>Raddi.</i> Raddi. An Italian cryptogamic botanist, and traveller in Brazil.
<i>Lyng.</i>	<i>Lyngbye.</i> A Danish writer upon cryptogamic matters.	<i>Raeuschel.</i> Raeuschel. A German botanist
<i>Ma. C.</i>	<i>Lyon.</i> A collector of American plants, tents <i>catalogue,</i> Cat. <i>atalogue,</i> of the at Clapton.	<i>Rafim'sque Schnialz.</i> Rafim'sque Schnialz. A modern writer upon botanical matters.
	<i>Mocino and Sess4.</i> Mexican botanists.	<i>Ram.</i> Ram. Ramand. A French botanist
	<i>Marcgraav.</i> A traveller in Brazil.	<i>Ruiz and Pavon.</i> Ruiz and Pavon. Spanish botanists, and travellers in Peru and Chile.
<i>Marsh.</i>	<i>Marshall.</i> A writer on American trees.	<i>Rbmer and Schultics.</i> Rbmer and Schultics. German editors of Linnus's Species Plantarum.
<i>Mart.</i>	<i>Martius.</i> A Bavarian botanist, and traveller in Brazil.	<i>Rau.</i> Rau. A German botanist
<i>Mas.</i>	<i>Masson.</i> A collector of plants at the Cape, and elsewhere.	<i>Robert Brown.</i> Robert Brown. A distinguished English botanist, and traveller in New Holland.
		<i>Reichenbach.</i> Reichenbach. A German botanist
		<i>Redout*?.</i> Redout*. A French botanical draughtsman.
		<i>Redowski.</i> Redowski. A Russian botanical collector
		<i>Reichard.</i> Reichard. A botanist of Frankfort
		<i>Reinwardt.</i> Reinwardt. A professor at Leyden.
		<i>^clhan.</i> ^clhan. A writer upon the Flora of Cheshire.
		<i>Req.</i> Req. Requien. A French botanist
		<i>Retx.</i> Retx. Retxius. A German botanist

<i>Reyn.</i>	<i>Reynier.</i> A botanist of Lausanne.	<i>Stier.</i>	<i>Suter.</i> A Swiss botanist.
<i>Rh.</i>	<i>Rheede.</i> Author of <i>Hortus Malabaricus</i> .	<i>Swert.</i>	<i>Swertia.</i> An old writer on plants.
<i>Rich.</i>	<i>Richard.</i> A French botanist.	<i>Swt.</i>	<i>Sweet.</i> An English botanist
<i>Risso.</i>	<i>Risso.</i> An Italian writer upon Oranges	<i>Swx.</i>	<i>Swartz.</i> A Swedish botanist, and traveller in the West Indies.
<i>Riv.</i>	<i>Rivinus.</i> A German botanist.	<i>Symes.</i>	<i>Symes.</i> A writer on English plants.
<i>R. It.</i>	<i>Redout&s Liliace'es.</i>	<i>Ten.</i>	<i>Tenore.</i> A Neapolitan botanist
<i>Robs.</i>	<i>Robson.</i> An English botanist	<i>Thib.</i>	<i>Thibaud de Chanvalon.</i> A French botanist
<i>Roch.</i>	<i>RocheL</i> Superintendant of the garden at Pest.	<i>Thomas.</i>	<i>Thomas.</i> A Swiss collector of plants.
<i>Roem.</i>	<i>Roemer.</i> A German botanist	<i>Thore.</i>	<i>Thore.</i> A French botanist
<i>Rokde.</i>	<i>Rohde.</i> A botanist of Bremen.	<i>Thory.</i>	<i>Thory.</i> A French botanist
<i>Rol.</i>	<i>Rolander.</i> A Swedish botanist	<i>Thou.</i>	<i>Du Petit Thouars.</i> A French botanist, and traveller in Madagascar.
<i>Roll.</i>	<i>Rollinson.</i> A nurseryman near London.	<i>Thouin.</i>	<i>Thouin.</i> A French botanist
<i>Rose.</i>	<i>Roscoe.</i> An English botanist, and writer upon Scitamme.	<i>Thuill.</i>	<i>Thuillier.</i> A French botanist
<i>Roth.</i>	<i>Roth.</i> A German botanical writer.	<i>Thun.</i>	<i>Thunberg.</i> A Swedish botanical traveller.
<i>Rox.</i>	<i>Roxburgh.</i> An Indian botanist	<i>Timm.</i>	<i>Timm.</i> A German botanist
<i>Rtb.</i>	<i>Rottboll.</i> A Danish botanist	<i>Tineo.</i>	<i>Tineo.</i> A professor at Palermo.
<i>Rtl.</i>	<i>Rattier.</i> A German missionary.	<i>Tode.</i>	<i>Tode.</i> A German writer on Flungi
<i>Rud.</i>	<i>Rudge.</i> An English writer upon botanical subjects.	<i>Torrey.</i>	<i>Torrey.</i> An American botanist
<i>Rudol.</i>	<i>Rudolph.</i> A German botanist	<i>Tou.</i>	<i>Tournefort.</i> An old French botanist, and traveller in Greece and Asia Minor.
<i>Rum.</i>	<i>Rumph' ms.</i> Author of <i>Herbarium Am. boinensis</i> .	<i>Trot.</i>	<i>Trattinik.</i> A botanist of Vienna
<i>Rus.</i>	<i>Russel.</i> A botanist of Aleppo.	<i>Trent.</i>	<i>Trentepohl.</i> A German botanist
<i>Sab.</i>	<i>Sabine.</i> An English amateur of botany.	<i>Trev.</i>	<i>Treviranus.</i> A German botanist
<i>Sal.</i>	<i>Salisbury.</i> An English botanist	<i>Trin.</i>	<i>Trinius.</i> A writer on grasses.
<i>Saint.</i>	<i>The Prince of Salm Dyck.</i> A noble German amateur.	<i>Turn.</i>	<i>Dawson Turner.</i> An English writer on sea weeds.
<i>S. \$ TV.</i>	<i>Schroder and JVettlrand.</i> German botanists.	<i>Turp.</i>	<i>Turp.</i> A French botanist and draughtsman.
<i>Santi.</i>	<i>Santi.</i> An Italian botanist	<i>Turra.</i>	<i>Turra.</i> An Italian botanist
<i>SavL.</i>	<i>SavL</i> An Italian botanist	<i>Tus.</i>	<i>Tussac.</i> A French writer on the Flora of the Antilles.
<i>Savig.</i>	<i>Savigny.</i> A French botanist	<i>Uc.</i>	<i>Ucria.</i> A botanist of Palermo.
<i>Schaff.</i>	<i>Schriffer.</i> A German writer upon F6ngi.	<i>Urv.</i>	<i>D*Urville.</i> A Captain in the French navy.
<i>Schlk.</i>	<i>Schkuhr.</i> A German writer upon grasses and ferns.	<i>Vail.</i>	<i>Vaillant.</i> A French botanist and traveller.
<i>Schl.</i>	<i>Schleicher.</i> A Swiss plant collector.	<i>Van.</i>	<i>Vandelli.</i> A Portuguese botanist.
<i>Schlee.</i>	<i>Schlechtendahl.</i> A German botanist	<i>Ven.</i>	<i>Ventenat.</i> A French botanist
<i>Schmidt.</i>	<i>Schmidt.</i> A Bohemian botanist	<i>Vest.</i>	<i>Vest.</i> An Styrian botanist
<i>Schott.</i>	<i>Schott.</i> A traveller in Brazil.	<i>Vig.</i>	<i>Vigvier.</i> A writer upon Poppies.
<i>Schous.</i>	<i>Schousboe.</i> A writer upon the Flora of Morocco.	<i>Vil.</i>	<i>Villars.</i> A French botanist
<i>Sehr.</i>	<i>Schroder.</i> A German botanist	<i>Vit.</i>	<i>Vitmann.</i> A German botanist
<i>Schrank.</i>	<i>Schrank.</i> A Bavarian botanist	<i>Viv.</i>	<i>Viviani.</i> An Italian botanist
<i>Schreb.</i>	<i>Schreber.</i> A German botanist	<i>W.</i>	<i>Willdenow.</i> A German botanist
<i>Schu.</i>	<i>Schubler.</i> A professor at Tubingen.	<i>Wahl.</i>	<i>Wahlenberg.</i> A Swedish botanist
<i>Schult.</i>	<i>Schultes.</i> A Bavarian botanist	<i>Wai.</i>	<i>Wallich.</i> Superintendant of the botanical garden at Calcutta.
<i>Schulx.</i>	<i>Schiltz.</i> A German botanist.	<i>Wald.</i>	<i>Waldstein.</i> A noble German patron of botany.
<i>Schum.</i>	<i>Schumacher.</i> A Danish botanist	<i>Wallr.</i>	<i>Wallroth.</i> A German botanist
<i>Schw.</i>	<i>Schweigger.</i> A German botanist, author of <i>Flora Erlangensis</i> .	<i>Walt.</i>	<i>Walter.</i> A writer on the Flora of Carolina.
<i>Schwa.</i>	<i>Schweegrichen.</i> A German cryptogamic botanist	<i>Wan.</i>	<i>Wangenheim.</i> A German botanist
<i>Schwein.</i>	<i>Schweinitz.</i> An American botanist		<i>Waldstein and Kitaibel.</i> Authors of the Flora of Hungary.
<i>Sco.</i>	<i>Scopoli.</i> An Italian botanist	<i>Wat.</i>	<i>Weber and Mohr.</i> German botanists.
<i>Seb.</i>	<i>Sebastiani.</i> An Italian botanist.	<i>Web.</i>	<i>Watson.</i> An English writer upon trees and shrubs.
<i>Sessi.</i>	<i>Sessi.</i> A Mexican botanist	<i>WeieL</i>	<i>Weber.</i> A Gennan cryptogamist
<i>Sib.</i>	<i>Sibthorp.</i> An English botanist, and traveller in Greece.	<i>Wethe.</i>	<i>Weigel.</i> A German botanist.
<i>Sirb.</i>	<i>Sieber.</i> A botanical collector.	<i>Wein.</i>	<i>Welthe.</i> A German writer on <i>Rbh</i> .
<i>Sims.</i>	<i>Sims.</i> An English garden botanist	<i>Weis.</i>	<i>Weinmann.</i> A German gardener, superintendant of one the royal gardens in Russia.
<i>Sm.</i>	<i>Sir J. E. Smith.</i> An English botanist, and purchaser of the Linncan herbarium.	<i>W. en.</i>	<i>Weis.</i> A German botanist
<i>Silt.</i>	<i>Sinclair.</i> Nurseryman at New Cross.	<i>West.</i>	<i>Willdenow's Enumeration of the Plants in the Berlin Garden.</i>
<i>Sol.</i>	<i>Solander.</i> A Swedish botanist, and companion of Sir Joseph Banks in Cook's voyage round the world.	<i>Wibel.</i>	<i>West.</i> An English botanist
<i>Sole.</i>	<i>Sole.</i> A writer on mints.	<i>Wig.</i>	<i>Wibel.</i> A German botanist
<i>Son.</i>	<i>Sonnarat.</i> A French botanist, and traveller.	<i>Wik.</i>	<i>Wiggers.</i> A German botanist
<i>Sow.</i>	<i>Sowerby.</i> An English botanical draughtsman.	<i>Wlk.</i>	<i>Wikstrom.</i> A Swedish botanist
	<i>Sparmann.</i> A Swedish travelling botanist.	<i>With.</i>	<i>Withering.</i> An English botanist
	<i>De Spin.</i> A botanist of Turin.	<i>Wnl.</i>	<i>Hendlund.</i> A German garden botanist.
	<i>Sprengel.</i> A German botanist	<i>Woods.</i>	<i>Woods.</i> An English writer on roses.
	<i>Stackhouse.</i> An English botanist	<i>Woodw.</i>	<i>Woodward.</i> An English botanist.
	<i>Sicchmann.</i> A writer on Artemisia.	<i>Worm.</i>	<i>Wonnshield.</i> A Norwegian botanist
	<i>Stephan.</i> A Russian botanist	<i>Wre.</i>	<i>Wredow.</i> A German divine and botanist
	<i>Sternberg.</i> A noble German botanist	<i>Wrede.</i>	<i>Wrede.</i> A German botanist.
	<i>SteudeL</i> A Gennan botanist	<i>WuL</i>	<i>Wulfen.</i> A German botanist
	<i>Steven.</i> A Russian botanist	<i>Y.</i>	<i>Young.</i> A writer in the Linnean Transactions.
	<i>Sturm.</i> A Gennan botanical draughtsman.	<i>Zea.</i>	<i>Zea.</i> A Spanish botanist
		<i>Zey.</i>	<i>Zeyher.</i> A German gardener.
		<i>Zuc.</i>	<i>Zuccagni.</i> Su)crmtcndant of the botanical garden at Florence.

NAMES OF BOOKS REFERRED TO.

AJ. China.	Abel (C.). A Voyage to China. 4to.	Ard. sag.	Arduini (Pietro), in Saggi dell' Accad. de Padova. 2 vols.
Ab. ins.	Abbott's Natural History of the Insects of Carolina. Folio.	Ard. sp.	<i>Id.</i> Animadversionum Botanicarum Specimen. 4to. Patavii, 1759.
Ac. boh.	Abhandlungen einer Gesellschaft in Boehmen, zur Aufnahme der Mathematik, der Vaterlandsdichen Geschichte und der Naturgeschichte. 8vo. Prags, 1775—1789.	As. res.	Asiatic Researches, or the Transactions of the Society instituted in Bengal 4to. Calcutta, 1788, &c.
Ac. bon.	Dc Bononiense Scientiarum, et Aratum Institute, Commentarii. 4to. Bononiae?. 7 vols. 1748—1791.	Asso ar.	De Asso (Ignatius). Synopsis Stirpium indigenarum Arragoniae. 4ta Marsilis, 1779.
Ac. not.	Commentarii Societatis Regiae Scientiarum Goettingensis. 4to. 1751. 1754. 1769. 1816.	Aub. gui.	Aublet (Fusee). Histoire des Plantes de la Guiane Franchise. 4 vols. 4ta London, 1773.
Ac. haf.	Acta Literaria Universitatis Hafniensis. 1 vol 4to. Hafnls, 1778.	Bal. mis.	Balbis (Joh. Baptist). Miscellanea Botanica. 4to. 1804.
Ac. hel.	Acta Helvetica Physico-mathematico-botanico-modica. 8 vols. 4to. Basiles, 1757—1777.	BaLst	<i>Id.</i> Horti Academicci Taurinensis Stirpium minus cognitarum aut forte novarum Icones et Description ea. Taurini, 1810.
Ac. hoi.	Kongl. Svenska Vetenskaps Akademii's Hafdiingar. 8vo. Stockholm, 1739—1816.	Bal. tic.	<i>Id.</i> Flora Ticincensis. 2 vols. 8vo. Ticin, 1816—1821.
Ac. par.	Actes de la Société d'Histoire Naturelle de Paris. 1 vol. fol. Paris, 1792.	Banks K.	Banks (Joscpus). Icones selectae Plantarum quas in Japonia collegit et delineavit E. Ksmpfer. Fol London, 1791.
Ac. pet.	Commentarii Academis Scientiarum Imperialis Petropolitans. 14 vols. 4to. Petropoli, 1728—1751.	Bar. fl.	Barton (W. P. C.). Flora of North America. 4to.
Ac. st	Kongl. Vetenskaps Akademis Handhagar. Stockholm, 1739. 8vo.	Bar. ic.	Barrclier (Jacobus). Plants per Gal. Ham, Hispaniam, et Italianam observatae. Fol. Parisiis, 1714.
Ac. tou.	Histoirc et Memoires dc l'Academie Royale des Sciences, &c, de Toulouse. 3 vols. 4to. Toulouse, 1782. 1784.178a	Bartr. it	Bartram (Will.). Travels through South and North Carolina. 1 vol. 8vo. Philadelphia, 1791.
Ac. up.	Acta Literaria et Scientiarum Upplandiae publicata. 8vo. 1720—1816.	Bast j.	Bastard (T.), in Schrader's Journal fur die Botanik V
Ag. sven. bot	Aghardt, in Svensk Botanik. U vols. 8vo. Stockholm, 1804 to the present time.	Batarra	Batarra (John Ant.). Fungorum Agri Arimincns Historia. 4to. F. venti©, 1759.
Al. au.	AUioui (Carolus). Auctuarium ad Floram Pedemontanam. 1 fasc. 8vo. Taurini, 17f&.	Batsch fun.	Batsch (Aug. Joh. Georg. CarL). Elenchus Fungorum. 4to. Hale, 1783—178a
Alb. & Schw.	Albertini (J. B. dc) and Schweinitz (L. D. de). Conspectus Fungorum in Lusatia Superioris Agro Nisiensi crescentium. 1 vol. 8vo. Lipsia, 1809.	Bauer h. h.	Bauer (Ferdin.). Illustrationes Florae Novae Hollandie. Fol. 1813.
Alp. sg.	Alpinus (Prosper). De Plantis JEgypti liber. 4to. Venetiis, 1592.	Ban. his.	Bauhin (Johannes). Historia Plantarum universalis. Fol. 1651.
ALped.	AUioni (Carolus). Flora Pedemontana 3 vols. fol. Taurini, 1785.	Bau. pin.	Bauhin (Caspar). Pinax Theatri Botanici. 4to. Basilea?, 1623.
Alp. ex.	<i>Id.</i> De Plantis Exoticis libri duo. 4to. Venetiis, 1629.	Bau. pr.	<i>Id.</i> Prodromus Theatri Botanici, in quo plants supra 600 proponuntur. 4to. Franc. Mcen. 1620.
Altau.	<i>Id.</i> Miscellanea Philosophico-mathematica Societatis private Tauri.	Beau. Ow.	Allioni. Palisot de Beauvois. Flore des Royaumes d'Oware et de Benin.. 2 vols. fol. Paris, 1805.
Am. ac.	Linnei Amcenitates Academics, seu Dissertations antehac seorsim editis. 10 vols. 8vo. Holmiae et Lipsiffi, 1749, et seq.	Beauv. gr.	<i>Id.</i> Essai d'une nouvelle Agrostographic. 1 vol. 8vo. Paris, 1812.
Amans s.	Amans. Obs. in Recueil des Travaux de la Societe d'Agriculture d'Agens. 1vol.	Beauv.in acam.JdL	Beauvain. Transactions of the American Philosophical Society. 4to, 3 vols. Philadelphia, 1774—1793.
Am. rut	Ammann (Johan.). Stirpium rariorum in Imperio Rutheno sponte provenientium Icones et Descriptions. 4to. Petrop. 1739.	BeL tau.	Bellardi (Ludovico). Memoires de l'Academie Royale des Sciences dc Turin. 4to. 1782.
An. bot	Annals of Botany, by C. Kdnig and J. Sims. 2 vols. 8vo. London, 1805 and 1806.	Berg. c.	Bergius (Peter Jonas). Descriptiones Plantarum ex Capite Bons Spei. 8yo. Stockholm, 1767.
And. ger.	Andrews (Henry). Coloured Engravings of Geraniums. Fol. London.	Berg. ho.	<i>Id.</i> in Kongl. Vetenskaps Academiens Handlingar. 8vo. Stockholm, 1739.
And. heaths.	<i>Id.</i> Coloured Engravings of Heaths, with botanical descriptions. 3 vols. London, 1802—1809.	Berg. ph.	Bergeret*. Phytonomatotechnie universelle. 3 vols. fol. Paris.
And. rep.	<i>fol.</i> London, 1802—1809.	Ber. mag.	Martini (Fried. Henr. Wilh.). Berlinisches Magazin, Oder ge. sammlte schriften. 4 band, 8vo. Berlin, 1765—1767.
An. mu.	<i>Id.</i> The Botanist's Repository for new and rare Plants. 10 vols. 4to. London, 1797, et seq.	Ber. ph. tr.	Bergius (P. J.), in Philosophical Transactions, vol. for 1772. London.
LL	Annales du Muséum d'Histoire Naturelle. 24 vols 4ta Paris, 1802 A to the present time.	Bes.eys.	Beelder (Basilius). Hortus Eystettensis. 2 vols. fol. Nuremberg, 1612.
An. wett.	Annalen der Wetteranischen Gesellschaft fur die gesammte Naturkundc.	Bib. it	Memorie di Matematica e tisica della Societa Italiana. 4to, 1 vol. Verona, 1782 and 1816.
Ard. m.	Arduni (Pietro). Memoria di Observazioni e di Sperienze sopra la Cultura e gli Usi di varie Piante. 4to. Padova, 1766.	Bieb. cen.	Bieberstein (Marschall von). Centuris Plantarum rariorum Rossis Meridionalis. Charkovia?. Fol
		Biv. cen.	Bivona Bernardi (Anton.). Sicutarum PUntarum Centuria prima. 8vo. Panormi, 1806.
		Biv. sic.	<i>la.</i> stirpium rariorum minusque cognitarum in Sicilia provenientium Descriptiones. 4to. 1813.

NAMES OF BOOKS REFERRED TO.

Black.	Blackwell (Eliz.). A curious Herbal containing 600 cuts of the use. Ail plants. 2 vols. to). London, 1737.	Bur. am.	Burmann (Johannes). Plants Africaneae a C. Plunder detectes et A J. Burmanno editis. Fol. Amstelod. 1755.
Boc. mu.	Bocccone (Paolo). Museo di Piante rare delta Sicilia, Malta, Corsica, Italia, Piemonte, e Germania. 4to. Venetia, 1697.	Bur. in.	<i>Id.</i> Flora Indica. 1 vol. 4to. Lugd. Bat 1768.
Boc. sic.	<i>Id.</i> Icones et Descriptiones rariorum Plantarum Siciliic, Melits, Gallis, et Italia?. 4to. Londini, 1674.	Bur. zey.	<i>Id.</i> Thesaurus Zeylanicus. 4to. Amst. 1737.
Boer.	Boerhaave (Hermannus). Index Plantarum quie in Horto A cad. Lugd. Bat recipiuntur. 8vo. Lugd. Bat. 1710.	Cam. ep.	Buxbaum (Joh. Christ). Plantarum minus cognitarum Centuria? quinque. 4to. 1728.
Bolt. fl.	Bolton (James). Filices Britannia. 4to. London, 1785—1790.	Cam. h.	Camerarius (Joachim). De Plantis Epitome utilissima. 1 vol. 4ta Fmncof. M ^o n. 1586.
Bolt pil	<i>Id.</i> Geschichte der Merkwürdigsten Pilze. Berlin, 1795	Cat car.	<i>Id.</i> Hortus Medicus et Philosophicus. 4to. Franc Ms ⁿ . 1588.
Bonam.	Bonamy (Franc). Flora? Nannetensis Prudromus. 1 vol. 12mo. Nantes, 1782. "	Cav. djs.	Catesby (Marsh). The Natural History of Carolina, Florida, &c. 4 vols. fol. London, 1741—1744.
Bon. mon.	Bonpland (Aime). Monographic des Melastomes et Rhexes et autres Plantes de cet Ordre. Fol. Paris, 1809.	Cav. ic.	Cavanilles (Ant. Jos.). Monadophila? Classis Dibsertationes. 10 vols. 4to. Paris, 1785—1789. Madriti, 1790.
Bon. nav.	<i>Id.</i> Description des Plantes rares cultives à Malmaison et X Navarre. Fol. 1813.	C. g.	<i>Id.</i> Icones et Descriptiones Plantarum qua; aut sponte in Hispania crescunt aut in Hortis hospitantur. 6 vols. foL Madrit 1791—1800.
Bon. rhex.	<i>Id.</i> Monographic des Melastomes et Rhexes et autres Plantes de cet Ordre. Fol. Paris, 1809.	Chan. cam.	Commentarii Societatis Regie Scientiarum Göttingensis. 4to. 1751—1816.
Bory.	Bory de St Vincent (J. I. M.). Voyage en Espagne a ^r travers les Royaumes de Galice, Leon, &c 8vo.	Chois. hyp.	Chandler's Camellias. London.
Bot cab.	The Botanical Cabinet. By Conrad Loddiges and Sons. 14 vols. 12mo and 4to, published monthly.	CL h.	Choisy (J. D.). Prodromus d'une Monographie de la Famille del Hypocrincéas. 4to. 1821.
Bot mag.	The Botanical Magazine. 8vo. 55 vols. London, 1787—1810.	Cl.pan.	Clusius (Carolus). Rariorum Plantarum Historia. Fol. Antwerp, NiOL.
Bot mis.	Hooker (Will. Jackson). Botanical Miscellany. 2 parts, 8vo. London, 1811.	Col. cas.	<i>Id.</i> Rariorum aliquot Stirpium per Pannoniamboserv. Historia. 1 voL 8vo. 1583 (Hal), 1584 (His.)
Bot reg.	The Botanical Register. 8vo. 14 vols. London, 1815—1830.	Col. ec.	Colladon (Frédéric). Histoire Naturelle et Médicale des Casse. 1 voL 4to. Montpellier, 1816.
Bot rep.	The Botanist's Repository for new and rare Plants. 10 vols. 4 to. London, 1797, et seqq.	Col. h. rip.	Columna (Fabius). Minus cognitarum Stirpiuin lēphras: 4to. Koxaas, 16W.
Braam.	Bream's Icones Chinenses. Not published.	Col. ph.	Colla. Hortus Ripulensis. 4to. Turin, 1829—1887.
Brad. sue.	Bradley (Richard). Historia Plantarum Succulentarum. 4to. London, 1716—1727.	Com. h.	Columna (Fabius). Phytobasanot. 4to.
Bra. saL	De Braune (Franz. Ant). Salzburgische Flora. 8vo. 1791.	Com. pet	Commelyn (Caspar). Horti Medici Amstelodamensis rariorum Plantarum Descriptio et Icones. 9 vols. fol. Amst 1703.
Brew. jour.	Brewster's Philosophical Journal. Edinburgh, published every three months.	Com. pr.	Commentarii Academiae Scientiarum Imperialis Petropolitans. 14 vols. 4ta Petrop. 1728—1751.
Brey. c	Breyneus (Jacobus). Exoticarum Plantarum Centuria. Fol. Gedani, 1678.	Com. r.	Commelyn (Caspar). Præludia Botanica ad publicas Plantarum Demonstrations. 4to. L. Bat. 1703.
Brey. pr.	<i>Id.</i> Prodromus Fasciculi rariorum Plantarum in Hortis Hollandie observatarum. 2 vols. 4to. Gedani, 1680^1689.	Cook.	<i>Id.</i> Horti Medici Amstelodamensis Plants rariores et exotics. 4to. L. Bat 1706.
Brid. mus.	Bridel (Samuel Elias). Muscologia. 4to. Goths et Parisiis, 1797—1803.	Cor. ca.	Cook (James). Voyage round the World (2d). 2 vols 4to. 1777.
Br. jam	Browne (Patrick). The Civil and Natural History of Jamaica. Fol. London, 1756.	Cr. au.	Cornuti (Jacob.). Canadenium Plantarum aliarumque nondum odita rum Historia. 4to. Paris, 1635.
Brot. lus.	Brotero (Felix Avicular). Flora Lusitanica. 2 vols. 8vo. Ollyssip. 1801	Crypt brit.	Crantz (Henr. Joh. Nepoin.). Stirpes Austriae. 8vo. 1762.
Brot. ph.	<i>Id.</i> Phytographia Lusitanise selection FoL Ollyssip. 1801.	Cup. pa.	Greville (Charles Kaye). The British Cryptogamic Flora. 5 vols. 8vo.
Br. rera.	Brown (Robert). General Remarks, Geographical and Systematical, on the Botany of Terra Australis. 4to. 1814.	Cup. sic.	Cupani (Francieus). Pamphylum Siculum, scu Historia Plantarum Sicilis. 1 vol. Fol.
Bruce tr.	Bruce (James). Travels to discover the Source of the Nile. 5 vols. 4to. Edinburgh, 1790.	Cur. Ion.	<i>Id.</i> Catalogus Plantarum Sicularum noviter detectarum. Panormi. 1652.
Bry. hist	Bryant (Charles). Flora Dietetica, or History of Esculent Plants. 1 vol. 8vo London, 1783.	Cyr. ne.	Curtis (William). Flora Londinensis. Fol. London, 1777, continued.
Buch.	Buchoz (Pierre Joseph). Collection des Flours. Fol. Paris.	Dalech.	Cyrillo (Domcnico). Plantarum rariorum Regni Ncariolitani S ^r ecimen. Fol. NeapoL 1788—1792.
Bui her.	Bulliard. Herbarie de la France. Fol. 1780, &c.	Dec as.	Dalechamps (Jacques). Hibtoriagencalis Plantarum. 2 vols. fol. Lugd. 1586—1587.
Bui ph.	Bulletin des Sciences, par la Société Philomathique de Paris. 1791—1816.	Dec. bis.	Decandolle (Augustus Pyramus). Astragalologia. 4to, ctfol. 180 ^r .
Bur at.	Burmann (Johannes). Rariorum Africarum Plantarum Decades 10. 4to. Amstelodami, 1738—17*19.	Dec. cac.	<i>Id.</i> Monographie des Biscutelles, in Annales du Museum, vol. 1& 1811. 4to.
		Dec. diss.	<i>Id.</i> Dissertation on Cacti, in Mémoire* du Museum, vol 17.
			<i>Id.</i> Dissertations difTerens sur ** Botanique. Various fascicle** 4to.

NAMES OF BOOKS REFERRED TO.

- iieuandolle (Augustus Pyramus). *Plantes rares du Jardin de Geneve.* Fasc. 1, 2. fol. Geneva, 1825.
Id. *Icones Plantarum Gallicariorum.* 4to. Paris, 1808.
Id. *Memoires sur les Légumineuses.* 4to.
Id. *Catalogus Plantarum Horti Botanici Monspicensis, addito observationum circa Species novas aut non satis cognitas Fascicule 8vo.* 1813.
Id. *Aitiales du Museum d'Histoire Naturelle.* 4to. 1802.
Id. *Plantarum Historia succulentarum.* Fol. and 4to. Paris, 1799—1830.
 Delile (Aline Rafeneau). *Mémoires Botaniques extraits de la Description de l'Egypte.* Fol. Paris, 1813. *Et Flora Egyptiaca illustrata.* Fol. Paris, 1813.
 De la Roche (Francois). *Eryngiorum necnon Generis novi Alepideae Historia.* Fol. Paris, 1808.
 Delessert (Benj.). *Icones selectae Plantarum in Systema universi descriptarum.* 4to. 1820.
 Watson (P. W.) *Dendrologia Britannica.* 1 vol. 1825.
 Description de l'Egypte, ou Recueil des Observations et des Recherches faites pendant l'Expedition de l'Armee Francaise. 4to, et fol. 1810.
 Desfontaines (Rend Louiche). *Mémoires dc 1^e Académie Royale des Sciences.* 4to. 1666.
Id. *Flora Atlantica.* 2 vols. 4to. Paris, 1798—1799.
Id. *Choix des Plantes du Corollaire des Institute de Touniefort* 1 vol. 4to. Paris, 1808.
Id. *Icones picta; Hort. Par. A Series of Drawings of Plants grown in the Jardin des Plantes, Paris.* Not published.
 Desvaux. *Journal de Botanique.* 8vo. 1808.
 Sturm (Jacob). *Deutschlands Flora.* 2 vols. 4to. Nurnberg, 1793, &c.
 Dickson (James). *Plantarum Cryptogamicarum Britannia Fasciculi.* 3 vols. 4to. London, 1785—1793.
 Dillenius (John J?c.). *Hortus Elthamensis.* 2 vols. fol. Londini, 1732.
Id. *Historia Muscorum.* 4to. Oxonii, 1741.
 Dodart (Denys). *Memoires pour servir a l'Histoire des Plantes.* Fol. Paris, 1676.
 Dodoneus or Dodoens (Rambrot). *Stirpium Historiae Pemptades vi.* Fol. Antwerp, 1583.
 Dufresne* (Pierre). *Histoire Naturelle et Medicale de la Famille des Valerianees.* 4to. 1811.
 Duhamel du Monceau (Henri Louis). *Traite des Arbres et Arbustes qui se cultivent en France en pleine terre.* 2 vols. 4to. Paris, 1755.
Id. *Traict des Arbres fruitiers.* 2 vols. 4to, Paris, 1768; 3 vols. 8vo, Paris, 1782; fol. Paris, 1808, et seq.
Id. See Dun. ar. A new edition, by Michel. 5 vols. folio. Paris, 1801—1816.
 Echal (Michel Felix). *Monographie des Anonacées.* 4to.
Histoire Naturelle, Medicale, et Economique, des Solanum et Genres qui ont été confondues avec eux. 4to. Montpellier, Du Roi (Joh. Phil.). *Die Harbkesche wilde Baumzucht.* 2 vols. 8vo. Braunschweig, 1771—1772.
 Dillwyn (Lewis Weston). *Synopsis of the British Conifers.* 4to. 1802—1814.
 Si-Jour. P. Jour. 3 The Edinburgh Philosophical Journal. Edinburgh. 8vo. Published quarterly.
- Ehr. pic. Ehret (George Dion.). *Plantw et Papillones rariores.* Fol. London, 1748—1759.
 Encyc. pi. Loudon (J. C. L.). *Encyclopaedia of Plants.* 8vo. London, 1830.
 Eng. bot English Botany, by Sir James Edward Smith, and Mr. James Sowerby. 36 vols. 8vo.
 Esper fuci. Esper (Eug. Joh. Christ.). *Icones Fucorum. Abbildungen der Fange.* 4ta Nurnberg, 1797—1799.
 Ex. bot Exotic Botany, by Sir J. E. Smith. 2 vols. 4to London, 1804—1808.
 See Hook. ex. fl. See Hook. ex. fl.
 Ferrari (Joh. Baptist). *Hespérides, sive de malorum aureorum culture et usu.* Fol. Roma?, 1646.
 Feuillee (Louis). *Journal des Observations Physiques, Mathématiques, et Botaniques, faites dans l'Amérique Meridionale, &c.* 4to. Paris, 1714—1725.
 Field (Baron.) *Geographical Memoirs of New South Wales.* 1 vol. 8vo. London, 1825.
 Collection of Drawings sent by Fischer to Deca'ido. Not published.
 De Tussac iF. R.). *Flora Antillarum.* Fol. Paris, 1808, et seq.
 Gmelin (Carol. Christ.). *Flora Baudensis-Alsatica.* 8vo. 1808.
 Schmidt (Franc. Wilib.). *Flora Bohemica.* 2 vols. fol. Pragae, 1793, 1794.
 Morris (Richard). *Flora Couspicua.* 1 vol. 8vo. London, 1826.
 Flora Danica, sive *Icones Plantarum sponte nascentium in Illegni Dauie et Norvegise, &c.* 9 vols. fol. Hafnia, 1761—1829. By Vahl, Horneman, and Muller.
 Sibthorp (Joh.). *Flora Gracca.* ed. J. E. Smith. 6 vols. fol. London, 1806—1829.
 Linneus (Carolus). *Flora Lapponica.* 8vo. Amstelodami, 1757.
 Curtis (William). *Flora Londinensis** Fol. London, 1777, continued.
 Sesse and Mochto. *Flora Mexicana icon. ined.* Iconibus ab ipso Sesse et Cervantesio curatis nondum editis usus est Candolleus.
 Magröff(W.). *Flora Monacensis.*
 Tenore (Michel). *Flora Neapolitana.* 4vols. fol. Neap. 1811, &c.
 Gunnerus (Joh. Enn.). *Flora Norvegica.* 2 vols. fol. Nidrosie, 1766 j Hafnias, 1772.
 Ruiz (HippoL) et Pavon (Jos.). *Flora Peruviana et Chilensis.* 5 vols. fol. Madriti, 1798—1799.
 Hofmansegg and Link. *Flore Portugaise.* Fol. Rostock and Berlin* 1806, &c.
 Picot de la Peyrouse (Philippe). *Figures de la Flore des Pyrenees.* Fol. Paris, 1795—1801.
 Lightfoot <John). *Flora Scotica.* 2 vols. 8vo. London, 1777.
 Pollini (Cyrus). *Flora Veionensis.* 3 vols. 8vo. Verona, 1822—1824.
 Forskaal (Petrus). *Flora iEgyptiaco. Arabica.* 4to. 1775. *Or, Icones Rerum Nat.* 4to. Hafnia, 1776.
 Commentarii Societatis Regiae Scientiarum Gottingensis. 4to. Gottingen, 1751, to the present time.
 Forster (George). *Characteres generuin Plantarum quas in Itinere ad Insulas Maris Australis, &c.* 4ta 1776.
 Fraser (John). *A short History of the Agrostis Cornucopia?* Fol. London, 1789.
 Fries (Elias). *Observationes Mycologicae.* 12mo. 2 vols. Hafnia, 1815—1818.
 Fuchs (Leonhard). *De Historia Stirpium Comroentarii insignes.* Fo., Basilis, 1542—1545.
 Gaertner (Josephus). *De Fructibus et Seminibus Plantarum.* 2 vols. 4to. Lipsia, 1788—1791.
 Garidel (Pierre Joseph). *Histoire des Plantes qui naissent aux Environs d'Aix.* 2vo. fol. Aix, 1715.

NAMES OF BOOKS REFERRED TO.

Gard. mag.	Loudon (J. C. L.). <i>Gardener's Magazine</i> . Vols. 1. to 6. London, 1826—1830, continued.	H. mon.	Decandolle (Aug. Pyr.). <i>Catij Horti Monspeliensis</i> , 8va spehi, 1813; 8vo, Gottingen, 1813.
Gaydfl.	Gay (John). <i>Fragment d'une Monographie des Buttneriacees</i> . 4to. Paris, 1823.	H. n. h.	Hotton (Petrus). <i>Rei herbaria toria et fatto sermo Acad. Lugd. 1695.</i>
Ger. em.	Gerard (John). <i>The Herbal, or General History of Plants</i> . 1 vol. fol. London, 1797. Enlarged by Johnson. 1 vol. fol. 1633.	Hof. cryp.	Hoffmann (George Franz.). <i>Historia Cryptogamica</i> . 4to. M 1787—1790.
Ger. gal.	Gerard (Ludovic). <i>Flora Gallo-provincialis</i> . 8vo. Paris, 1761.	Hof. ger.	<i>Id. Flora Germanica</i> .
Gil. euu	Gilibert (Joh. Em.). <i>Histoire des Plantes de l'Europe, ou Elemenrs de Botanique Pratique</i> . 2 vols. 8vo, Lyons, 1798; 3 vols. 8vo, Lyons, 1806.	Hof. got	<i>Id. Programma Hortus Fol. 179a</i>
Gm. it	Gmelin (J. G.). <i>Reise durch Sibirien</i> . 4 vols. 8vo, Gotting. 1751—1753.	Hof. ph.	<i>Id. Phytographische Blatter</i> . 1 Gottingen.
Gm. si.	<i>Id. Flora Sibirica</i> . 4 vols. 4to. Petropol. 1747—1769.	Hof. sa.	<i>Id. Historia Salicium Iconibus trata. Fol. Lipsis, 1785.</i>
Gou. iL	Gouan (Antoine). <i>Illustrations Botanice</i> . Fol. Tiguri, 1773.	Hof. urn.	<i>Id. Plantarum Umbelliferarum</i> 8vo. 1816.
Gou. m.	<i>Id. Hortus Regius Monspeliensis</i> . 8vo. Lugduni, 1762.	Hook. ex. fl.	Hooker (W. J.). <i>The Exotic</i> 3 vols. 8vo. Edinb. 1825—f
Grev. cryp.	Greville (Robert Kaye). <i>The Scottish Cryptogamic Flora</i> . 4 vols. 8vo.	Hook. jung.	<i>Id. Monograph of the British germanis. 4to. London, 1816.</i>
Gr. wo.	Sinclair (George). <i>Hortus Gramineus Woburnensis</i> . 1 vol. 8vo. 1825.	Hor. ph.	<i>Hors Physics Berolinenscs</i> . Fi* lin.
Hac. ca.	Hacquet (Balthazar). <i>Plantar Alpins Carniolics</i> . 4to. Vienna, 1782.	Hort. tr.	<i>Transactions of the Horticultural Society of London</i> . 7 vols. 1782.
Hag. lich.	Hagen (C. G.). <i>Tentamen Historis Lichenum prssertim Prussicorum</i> . 1 vol. 8vo. Regiomonti, 1782.	Host'gr.	Host (Nicol Thorn.). <i>Icores scriptiones Graminum rum</i> . 3 vols. fol. Vindob. 1803,
Halh.	Haller (Albert). <i>Historia Stirpium indigenarum Helvetiae</i> ; 3 vols. fol. Berne, 1768.	H. pfl.	Houttuyn (Martin). <i>Des r Linne! Pflanzensystem Anlcitung des Ho Werks</i> . 14 vols. 8vo. 1777—1788.
H. an.	Hortus Anglicus.	Ic. h. k.	Bauer (Francis). <i>Delineations of Plants cultivated in the Garden at Kew, published by J. Aiton</i> . Fol. London, 1791.
H. & B.	Von Humboldt (Alexand. Fr. Henr.) et Bonpland (Aime). <i>Plantes Equinoxiales</i> . Fol. 1808.	Is. ac. pa.	Isnard. See Ac. par.
H. & B. n.	Humboldt, Bonpland, and Kunth. <i>Nova Plantarum Genera et Species</i> . 7 vols. 4to. Paris, 1813 to 1825.	Jac. am.	Von Jacquin (Nicolas Jo) piura Americanarum Fol. 1761.
H. & G.	Hooker (W. J.) and Greville (R. K.). <i>Icones Filicum</i> . Fol. London, 1827.	Jac. am. pic.	<i>Id. The same work with plates, no date.</i>
Hay. ter.	Hayne (Frid. Gottlieb). <i>Termini Botanici Iconibus illustrati</i> . 4to. Berlin, 1799.	Jac. au.	<i>Id. Flora Austriae Icones</i> . fol. Vindob. 1773—1778.
Hay. us.	<i>Id. in Usteri's Annalen der Botanik</i> . 6 vols. 8vo. Zurich, 1791—1793.	Jac. c.	<i>Id. Collectanea ad Botanii spectantia</i> . 5 vols. 4to. 1786—1796.
H. ben.	Roxburgh (Will). <i>Hortus Bengalensis</i> . 1 vol. 8vo. Calcutta, 1814.	Jac. ec.	<i>Id. Eclogi Botanies</i> . Fol. 181
H. ber.	Willdenow (Car. Lud.). <i>Hortus Berlinensis</i> . Fol. Berlin, 1806—1810.	Jac. fr.	<i>Id. Fragmenta Botanica</i> . ***—1809.
H. cl	Linnaeus (Carolus). <i>Hortus Cliffbrtianus</i> . Fol. Amst. 1737.	Jac. gram.	<i>Id. Eclogi Botanies</i> . Fol. 181
Hed. cryp.	Hedwig (Johannes.). <i>Theoria Generationis et Fructificationis Plantarum Cryptogamicarum</i> . 1 vol. 4ta Petropoli, 1784.	Jac. h. vin.	<i>Id. Hortus Botanicus Vindobonensis</i> . 3 vols. fol. Vindob. 170
Hed. mus.	<i>Id. Species Muscorum Frondosorum</i> , 4to. Lipsiae, 1801.	Jac. ic.	<i>Id. Icones Plantarum rariorum</i> . fol. Vindob. 1781—1793.
Her. am.	Courset (Dumont de), or Delaune. <i>Herbier de V Amateur</i> . Paris.	Jac. m.	<i>Id. Miscellanea Austriae tantiam, &c, spectantia</i> . 4ta Vindob. 1778—1781.
Herb. ap.	Herbert (William). <i>A Botanical Arrangement of Bulbous Roots</i> . 8vo 1821.	Jac. ob.	<i>Id. Observations Botanies</i> . * 1764—1771.
Her. ger.	L'Heritier (Charles Louis). <i>Geranologia</i> . Fol. Paris, 1787, 1788.	Jac. ox.	<i>Id. Oxalis Monographia illustrata</i> . 4to. Vindob.
Her. lug.	Hermann (Paulus). <i>Horti Lugduni Batavi Catalogue</i> . 8vo. Lugd. Bat 1687.	Jac. sc.	<i>Id. Plantarum rarioiu Horrei Schcenbrunensis</i> . 4 v Vindobonen. 1797—1804.
Her. par.	<i>Id. Paradisus Batavus</i> . 4to. Lugd. Bat 1798.	Jac. st.	<i>Id. Stapelis cults</i> . Fol. 1806—1815.
Her. ser.	L'Heritier (Charles Louis). <i>Sectum Anglicum, sive Plants rariores</i> . Fol. Paris, 178a	Jo. h. n.	<i>Journal d'Histoire Naturelle</i> . 4to. Paris, 1792.
Her. st.	<i>Id. Stirpes novas aut minus cognite</i> . Fol. Paris, 1784—1785.	Jo. sc.	<i>Journal of Science</i> , edited at the Institution. 8vo. ***
H. er. wo.	Hortus Ericus Woburnensis. 4to. 1 vol. London, 1824.	Jun. ic.	<i>Junghans (Phil. Carp.). Iconaturum ad Vitam i Hals, 1787.</i>
H. flor.	Passus (Crispinus). <i>Hortus Floridus</i> . Fol. Arnheim, 1614.	Juss. 6ci.	Jussieu, in <i>Memoirs de l'Academie Royale des Sciences</i> . 1740.
H. eot	Hortus Gottingensis. 2 fasc fol Gottingae, 1809—1813.	Ksm. am.	Ksmper (Engelbert). <i>Exotics</i> . 4to. Leipzig, 1791.
H. has.	Moench (Conrad). <i>Enumeratio Plantarum indigenarum Hassiae</i> . 1 vol. 8vo. Cassel, 1777.	Ksm. ic.	<i>Id. Icones selectas Plantarum</i> . London, 1791.
H. kew.	Aiton (William). <i>Hortus Kewensis</i> . Ed. 1. 3 vols. 8vo. London, 1789.	Keraabr.	Kerner (John Sim.). <i>A und Beschreibungen Welche in Wurteni Wachsen</i> . 4 heft. 4to. 1783—1786.
H. mat	Lagasca (Mariano). <i>Catalogus Plantarum qua in Horto Regio Madritensi colebantur anno 1815</i> . 8vo. Madriti, 1818.	Ker rev.	Ker (John Bellenden). <i>tarum hucusque in Botanicorum ~</i> . London, 1801.

NAMES OF BOOKS REFERRED TO.


xvii

Ker strel ic.	Ker (John Bellenden). Coloured figures of the Genus Strelitzia. Fol.	Lob. ic.	Magazin der Gesellschaft Natur-iutchunden Freunde.
Kno. dcL	Knorr (Georg. Wolfgang). Deliciffi Natura's selecta. 2 vols. fol. Nurnberg, 1766—1767.	Lob. ob.	De Lobel or Lobelius (Mathias). Stir-pium Icones. 4to. Antwerp, 1601.
Kna th.	<i>Id.</i> Thesaurus Rei Herbariae Hortensis universalis. 2 vols. fol. 1770—1772.	Lcef. it	<i>Id.</i> Observations Botanies. Loeffling (Peter). Iter Hispanicum. 8vo. Stockholm, 1758.
Kr.siL	Krockcr (Ant Jon.). Flora Silesiaca renovata. 2 vols. 8vo. Vratislavis, 1787—1790.	Loe. pi.	Locslius (John). Flora Prussica. 4ta Hegiomonti, 1793.
Kth. miin.	Kunth (Car. Sigism.). Mimoses et autres Plantes Leguminosae du Nouveau Continent Fol. Paris, 1819, &c.	Loi. gal	Loiseleur Deslongchamps (J. L. A.). Flora Gallica, 2 vols. 12mo. Paris, 1806—1807.
Kun. nov. g.	<i>Id.</i> Nova Genera et Species Plantarum. 7 vols. 4to. Paris, 1825.	Loi. no.	<i>Id.</i> Notices sur les Plantes a ajouter & la Flore de France. 1 vol. 8vo. Paris, 1810.
Lab. n. h.	Labillardiere (Jac. Jul.). Novae Hollandiae Plantarum Specimen. 2 vols. fol. Paris, 1804—1808.	Loi. sup.	<i>Id.</i> Supplement to Flora Gallica. 8vo. Paris, 1807.
Lab. syr.	<i>Id.</i> Icones Plantarum Syris rariorum. Fol. Paris 1791—1812.	Loud. cnc. pi	Loudon (J. C. L.). Encyclopaedia of Plants. 8vo. London, 1850.
Lab. vo.	<i>Id.</i> Relation du Voyage à la Recherche de la Peyrouse. Paris, 1798. 2 vols. 8vo.	Loud. g. m.	<i>Id.</i> Gardener's Magazine. Vols. 1. to a London, 1826—30, continued.
Lamb. ci.	Lambert (Aylmer Bourke). Description of the Genus Cinchona. London, 1797.	Loud. m. n. h.	Loud. m. n. h. <i>Id.</i> Magazine of Natural History. Vols. 1. to 3. London, 1828—31, continued.
Lamb. pin.	<i>Id.</i> A Description of the Genus Pinus. Fol. London, 1803. New ed. 1829.	Lyng. hyd.	Lyngbye (H. C.). Hydrophytologia Danica. 4to. Copenhagen.
Lam. ic. in.	Lamarck (Jean Baptiste Monet de la). Icones Plantarum incd.	Mag. ber.	Martini (Henr.). Berlinisches Magazin oder Gesammelte Schriften. 8vo. Berlin, 1765—1767.
Lam. il.	<i>Id.</i> Illustration des Genres. 4to. Paris, 1791, &c.	Mag. h.	Magnol (Petrus). Hortus Regius Monspeliensis. 8vo. Monsp. 1697.
Lamour. dis.	Lamouroux (Justin). Dissertations sur plusieurs Espèces de Fuchs. 4to. Agcn, 1805.	Mag. mo.	<i>Id.</i> Botanicum Monsplicense. 12mo. Monsp. 1686.
T & Q. ab. ^{i c ~}	Link (Henr. Frid.) et Otto (FrM.). Abbildungen und Beschreibungen von selteuer Ptianen in Berlin Sarten. 4to. Berlin, 1821, and following years.	Mae. nat hU.	Mae. nat hU. Loudon J.C.I. Magazine of Natural History. Vols. 1. to 3. London, 1828—30, continued.
*- & O. dm.	/rf. Ueber die Gattungen Melocattus und Echinocactus in Hort Soc. Trans. Berlin, vol. 2. 4to.	Marcg. bra.	Marcgravius (Gcorg.). Historia Rerum Naturalium Brasiliae. Fol. L. Bat.
T o — ^{{ sp. -}	W. Hors Physics Berolinenses. Fol. Lapeyrouse (Philip Picot). Monographia Saxifragarum. Cujus partem facit Toulouse.	Mart. br.	Martius (C. F. P.). Nova Genera et Species Plantarum quas in Itinere per Brosiliain ann. 1817—1820. 4to.
{ sp. - ber. ^{sax.}	Le Roche (Daniel). Specimen Botanicum inaugurate sistens Descriptiones Plantarum aliquot novarum. 4to. 1766.	Mart. ccn.	Mart. ccn. Mart (John). Historia Plantarum riariorum. Cent 1. dec. 1—5. Fol. London, 1728.
Lar. dis.	De La Roche (Daniel). Specimen Botanicum inaugurate sistens Descriptiones Plantarum aliquot novarum. 4to. 1766.	Mart. med.	Von Martius (Carl. F. P.). Specimen Matris Medicis Braniilicnsis. Monach. 1824. Fol.
Lawr. pass.	Lawrence (Miss). Six numbers of coloured figures of Passion Flowers. Fol. London.	Mart palm, ic	Mart palm, ic <i>Id.</i> Genera et Species Palmarum quas in Itinere per Brazilian collegit. Monach. 1824, Fol.
Lawr. ros.	<i>Id.</i> A Collection of Roses from Nature. Fol. London, 1799.	Mass	Masson (Francis). Stapclis novae. Fol. London, 1798.
L. D. fr.	Ixuseleur's Duhamel, &c. A new edition by Loiseleur Deslongchamps. See Duh. fr.	Mat. com.	Mathiolus (P. Andr.). Commentarii in Libros Dioscoridis de Medica Materia.
Lee. pas.	Leconte's Pastures.	Mat. op.	<i>Id.</i> Opera Omnia. Fol. Francof. 1598; Basiles, 1674.
Lee. gr.	Leers (Joh. Dan.). Flora Herbor. nensis. 8vo. Colon. Allobr. 1789.	Mee. ic.	Meerburg (Nicol). Plantarum selectarum Icones pictæ. Fol. L. Bat 1798.
Leers her.	Lehmann (J. C. GX). Plants Asperi. folis Nucif. 2 vols. 4to. Berlin, 1818.	Mém. m.	Mémoires du Museum d'Histoire Naturelle. 4to. Paris, 1815—1819.
Leh. as.	<i>Id.</i> in Berlinisches Magazin, &c.	Mém. m&sc.	Mémoirs de la Societe Imp'riale des Naturalistes de Moscow. 6 vols. 4to. Moscow, 1811, to the present time.
Leh.b.	<i>Id.</i> Monographia Generis Primularum. 4to. Lipsiae, 1817.	Men. pu.	Menzel (Christ.). Pugillus Plantarum Tariorum. Fol. 1682.
Leh. m.	<i>Id.</i> Nicem Nicotianarum Historia. 4to. 1818.	Mer. sur.	Merian (Maria Sybilla). De Metamorphosis Insectorum Surinamensium. Fol. Hags, 1726.
Leh. nic.	<i>Id.</i> Monographia Generis Potentillarum. 4to. Hamburg, 1820.	M. fl. vi.	Moretti (Gius.). Notizia supra diverse Piante da aggiungiarsi alia Flora Vicentina. 4to. 1813.
Leh. pot.	<i>Id.</i> Monographia Generis Potentillarum. 4to. Hamburg, 1820.	M.h.	Morison (Robert). Plantarum Historia Universalis Oxoniensis. 8 vols. fol. Oxon. 1680.
L. f. dec.	Linnsus (Carolus, filius). Plantarum riariorum Horti Upsalicnsis Decas. Fol. Stock. 1762—i	Mic. am.	Michaux (Andr.). Flora Boreall-Americanæ. 2 vols. 8va Paris, 1803.
L. f. fa.	<i>Id.</i> Plantarum riariorum Horti Upsaliensis fasciculus 1. Fol. Lips. 1767.	Mic. ar.	Michaux (Andre' Franc, fils). Histoire des Arbres forestiers d'Amerique Septentrionale. \$ crova. 4to. Paris, 1810—1813.
Lind. col.	Lindley (John). Collectanea Botanica. Fol. London, 1821.	Mic. gen.	Micheli (Peter Ant.). Nova Plantarum Genera. Fol. Florence, 1719.
Lind. dig.	<i>Id.</i> Digitalium Monographia. Fol. London, 1821.	Mic. quer.	Histoire des Plantes d'Amérique Seirten.
lhd. ns.	<i>Id.</i> Rosarum Monographia. 8vo. London, 1820.	Mil. d. ed.	Milier (Philip). Gardener's Diction-
Lin. er.	Linnsus's Liceæ.		
Lin. tr.	Transactions of the Linnean Society of London. 4to. London, 1791—1829.		
Lk. obs.	Link (Henr. Frid.). Observations in Ordines Plantarum Naturales (Fungos tantum), (isa. L et 2. in		

Mil. ic.	Miller (Philip). Figures of Plants described in the Gardener's Dictionary. 2 vols. fol. London, 1760.	Pal. ros.	Pallas (Peter Simon) Flora Petrop. 1784—1788.
M. in.	Mémoires de l'Institut National des Sciences et des Arts. 1 vol. 4to. Paris, 1796, et seq.	Par. Ion.	Salisbury (Rich. Ant.) Londiniensis. 2 vols. 4to. 1805—1808
MissChalm, ic.	Figures of Bulbs, by Miss Chalmers. Not yet published.	Par. p.	Parkinson (John). A Pleasant Flowers. FoL 1629.
M. msq.	Usher, in Memoirs of Natural History of Moscow. See M6m. Mosc	Parry's app. ic.	Parry (William). A Voyage to cover the North West Paff 4to. London. Appendix.
Moen. wei.	Moench (Conrad). Verzeichniss AuslandUcher Baume des Lustschlosses Weissenstein. 8vo. Frankf. 1785.	Par.th.	Parkinson (John). Theatrum BcW cum. FoL London, 1760.
Ma gr.	Monti (Josephus). Catalogi Stirpium Agri Bononiensis Prodromus Gramina et Affinia complectens. 1 vol 4to. Bononis, 1719.	Patters.	Patterson (William). A Narrative of Four Journeys into the Country of the Hottentots. 4to. London, 1777.
Muhldan.	Muller(Otfrid.). Flora Danica Vols. 4. and 5. The rest by Vahl and Horneman. 5 vols. foL Havnis, 1566—1782.	Pen. dtsp.	Persoon (Christ Henr.). Tcnit*J Dispositionis methodic* FoL Trum. 8vo. Lijw. 1710.
M. um.	Morrison (Robert). Plantarum Urn. belliferum Distributio nova Fol. Oxon. 1672.	Pers. ie. pict	Id. Icones picti Specie rari Fungorum. 4to. Paris and S. 1803.
Munt. ph.	Munting (Abrah.). Phytographia curiosa 1 vol. fol. 1702.1713.1727.	Pet brit	Id. Synopsis Plantarum, seu Eridion Botanicum. 2 vols. Paris, 1805—1807.
Mur.	Murray (Joh. Andr.). Prodromus Designationis Stirpium Gottingen- sium. 8vo. Goffing, 1770.	PetfiL	Petiver (James). Herbarii Brita Catalogus. FoL London, 1704.
Muse. brit.	Hooker and Taylor. Muscologia Bri- tannica 8vo. London, 1818.	Petg.	Id. Pterigraphia Americana London.
Muse. hib.	Turner (Dawson). Muscologis Hibern- ics Specimen. 12mo. London, 1804.	Petm.	Id. Gazophylacium Naturis et FoL London, 1702—1704.
N. a p.	Nova Acta Academis Scientiarum Imperialis Petropolitans.	Ph. am.	Id. Mussi Petiveriani Centur* 8vo. London, 1795.
Nat ber.	Willdenow (Lud.). Der Gesellschaft Naturforschender Freunde zu Berlin neue Schriften. 4to. Berlin, 1795, and following years.	Ph.tr.	Pursh (Frederick). Flora Ameri- don. Septentrionalis. 2 vols. 8vo. 1814.
N. c g.	Novi Commentarii Societatis Regis Scientiarum Göttingensis. 4to. 1751, to the present time.	Pic. h. p.	Transactions of the Royal Socy London. Many vols. 4to. 1665, to the present time.
N. C p.	Novi Commentarii Academis Imperialis Petropolitans.	Pis.br.	Picciuoli (Giuseppe). Hortus Pan- ticus 4to. Firenze, 1715.
N. cur	Nova Acta Physico-medica Academis Cossarea Leopoklino-Caroiins Naturae Curiosorum. 4to. Continued to the present time.	Pis. lib.	Piso (Gulielm.). Historia Natu- Brasili. FoL Amsterdam, 1704.
Nees h. b.	Nees von Esenbeck (C. G.). Hors Physics Berolinenses collects ex Symbolis Virorum doctorum Link, Rudolphi, &c. fol. 1820.	PI. am.	Id. De Indis utriusque Re nitt et medica libri 14. FoL lx:S. Burmanno edits. FoL 1755.
Neespilze.	<i>Id.</i> Das System der Pilze und Schwamme. 4to. Wurtzburg, 1817.	Flan.h. b.	Id. Traite des Fougères de rique. FoL Paris, 1705.
Nees r. g.	<i>Id.</i> Assisted by Aug. Weihe. Rubis Germanicis cdendis,&c. 4to. Rome, 1823.	PLed.B.	Id. Nova Puntarum Americanan Genera. 4ta Paris, 170J
Men pot	Nestler (C. G.). Monographic de Potentilla 4to. Parish's et Argentorati, 1816.	PLic.	Id. Plantarum Americanarum 10, continentis Plantas olim C. Plumierius detexit. depinxit Edidit Johannes mannus. FoL Amst, 1755.
N. h.	Acta Literaria Universitatis Haf- niensis. 4to. Hafhis, 1778, and following years.	Plk.ic.	Plenck (Jos. Jac.). Icones PI ^ A medicinalium. Fasc. fol. IS ^o seq.
N. m. r.	Nees and Martius, in Nova Acta Phy- sico.medica Academis Cossarea Leopoldino-Carolins Naturae Cu- riosorum Bonns. 4to. From 1757, to the present time.	Pluk. al.	Plukennet (Leonard). Almagest Botanicum, sive Phyt Onof- con. 1 voL 4to. 179&.
Nocca tr.	Nocca (Dominicus). Ticinensis Horti Plants selects. 1 fasc. foL Pa- pis, 1814.	Plulcam.	Id. Amaltheum Botanicum. 1 vol. 1705.
NoL jar.	Noisette (Louis). Le Jardin Fruitier. 4to. Paris, 1813.	Plulcm.	Id. Almagesti botanici Mantissa. 1 4to. 1700.
Nov.com.got.	See N. eg.	Pluk. ph.	Id. Phytographia, seu Stirpium triorum, &c. 4vols. 4ta Lo- 1691-170&
Or. d.	De Ortega (Cas. Gomez). Novarum aut rariorum Plantarum Hor. R. Madritensis Decades. 4to. Ma- drid, 1797,1798.	Poc. crat	Pococke. Cratogl
Osb. it	Osbeck (Peter). A Voyage to China, &c. 8vo. London, 1771.	Pohl. fl. bo.	Pohl. (Imman.). Versuch einer rs*T Bohmens. 2 vols. 8vo. PTM 1810—1814.
Pa it ger.	Pallas (Peter Simon). Reise durch Verschiedene Provinzen des Russischen Reichs. 3 vols. 4to. Petrop. 1771—1776.	PtetT.il.	Poiteau et Turpin. Flore Paris* FoL Paris, 1808, &c
Pal. as.	<i>Id.</i> Species Astragolorum descriptis et Iconibus illustrate. FoL Lips. 1800.	Purt fl.	Purton (Th.). Midland Flora 8vo. Stratford upon Avon.
PaL ill.	<i>id.</i> Illustrationes Plantarum minus coonitarum. FoL Lipsia, 1801.	Qahis.	Quer (Martinez). Flora 4 vols. 4to. Madrid, 1764.
PaL it'	<i>Id.</i> Voyages dans PEmpire Russe 8 vols. 8vo.	Raddi m.	Raddi (Joseph). Memoir Observations in Atti vol. 9, and in Memorie di vols. 18. and 19.
PaL p.	<i>Id.</i> In the Pittsburgh Transactions. See Ac. pet	Rad. die	Radius (Justus). Dc Pyrola maphila Specimen. 4ta 1821.

NAMES OF BOOKS REFERRED TO.

six

Rauw hod.	7 Rauwolf (Leoni.). Aigentliche Be-	Sc. v.	Schrader (H. A.). <i>Commen-</i>
Rauw. it.	j schreibung der Raiss in den Mor-	Sc. verb.	<i>Uto de</i>
	ganlander. 4to. Laugangen, 158*	Sch. ag.	<i>Veronicis spicatis.</i> 8vo. Gottinga;
	—1583.	Sch. bs.	1803.
Ray sy.	Ray (Joh.). <i>Synopsis Stirpium Britan-</i>	Sch.ce.	<i>Id. Monographia Generis Verbasci</i>
Rchb. ac.	<i>nicarum.</i> 1 vol. 8vo.	Schaeff.	4to. Gottinge, 1813.
Reichenbach (Lodov.). <i>Monoeraphia</i>		Schleca.	<i>Scheuchzer (Johan.). Agrostographia,</i>
	<i>Generis Aconiti et Delphinii.</i>	Schk.cryp.	<i>sive Graminum, Juncorum, Cyper-</i>
Rchb. bot	Fol. Lipsiae, 1820.	Schk. fil.	<i>orum, iisque Affinium. Histon.</i>
	<i>Id. Hortus Botanicus.</i> 4to. Lipsiae,	Schk. han.	1 vol. 4to. Tigari, 1775.
Reb. neo.	1824.	Schk. us.	<i>Schwenck (Mart. W.). Beschryving</i>
	<i>Rebcnftisch (Joh. Frider.). Prodromus</i>	SchL ran.	<i>der Gewassen. Gravhage, 1760.</i>
	<i>Flora? Neomarchicffi.</i> 8vo. Bero-	Schm. ar.	<i>Schreber (John ChristJ. Beschreibung</i>
Red. lil.	lini.	Schou.	<i>der Gnesser. Fol. Leipsic, 1769 and</i>
	<i>Redout? (P. J.). Les Liliacees.</i> 8 vols.	Schm. ic.	1810. 3 fasc.
Red. ros.	fol. Paris, 1802—1816.	Schm. ic.	<i>Schaafft (Jac. Christ). Fungorum</i>
Rwi. spec.	<i>Id. Les Roses.</i> Fol. Paris.	Schm. ic.	<i>Bavaris et Palatinatus Icones.</i>
Retz.	<i>Rneaulme (Paul.). Specimen His-</i>	Schm. ic.	4 vols 4to. Ratisbonae, 1702—1770.
R-Hou.	<i>toric Plantarum.</i> 4to. Paris, 1611.	Schm. ic.	<i>Schkuhr (Christ.). Histoire des Carex</i>
	<i>Retzius (And. Joh.). Observations</i>	Schm. ic.	<i>ou Laiches.</i> 8vo. Lcipsig, 1802.
Riv. mon.	<i>Botanies.</i> 4to. London, 1774—1791.	Schm. ic.	<i>Id. Vier und zwanzigste Klassedes</i>
Riv. tct	<i>Reliquiae Houstoniana?</i> 4to. Lwidon,	Schm. ic.	<i>Linneschen Pflanzensystems.</i> 4ta
R-mal.	17H.	Schm. ic.	<i>Wittenberg, 1809.</i>
Rob. ic.	<i>Rivinus (Aug. Quirinus.). Ordo Plan-</i>	Schm. ic.	<i>Id. Abbildungen dor Farrnkrauter.</i>
Rw. ar.	<i>tarum Flore irregulari monopetalo.</i>	Schm. ic.	<i>Id. Botanisches Handbuch.</i> 3 vols. 8vo.
Rw. ac.	Fol. Lips. 1690.	Schm. ic.	<i>Wittenberg, 1791—1803.</i>
	<i>Id. Ordo Plantarum Flore irregulari</i>	Schm. ic.	<i>Id. in Usteri's (Paul) Annalen der</i>
	<i>tetrupetalo.</i> Fol. Lips. 1699.	Schm. ic.	<i>Botanik.</i> 6 vols. Zurich, from
	<i>Van Rhede (Henricus.). Hortus In-</i>	Schm. ic.	1791 to 1793.
	<i>dicus Mulabaricus.</i> 12 vols. fol.	Schm. ic.	<i>Schlchtendahl (A.). Animadversiones</i>
	1678—1703.	Schm. ic.	<i>in Ranunculeas Candolii.</i> 4to.
	<i>Robert (Nicolas.). Icones Plantarum.</i>	Schm. ic.	<i>Berlin, 1819, 182a In 2 fasciculi.</i>
	Fol. Paris, 1701.	Schm. ic.	<i>Schmidt (Franz.). CESTreichs Allge-</i>
	<i>Roemcr (Jac. Job.). Archiv fur die</i>	Schm. ic.	<i>meine Baumzucht</i> 2 vols. fol.
	<i>Botanik.</i> 3 vols. 4to. 1796—1799.	Schm. ic.	<i>Vienna, 1792—1794.</i>
	<i>Id. Scriptores de Plantis Hispanicis,</i>	Schm. ic.	<i>Schmidel (Cas. Christ). Icones Plan-</i>
	<i>Lusitanicis, et Brasiliensisbus.</i>	Schm. ic.	<i>tarum.</i> Fol. Norimb. 1762.
Rose, scit	1 vol. 8vo. 1796.	Schm. ic.	<i>Schneevoght (G. Voorhalm). Icones</i>
Rost. J.	<i>Roscoe (W.). Figures of the Order of</i>	Schm. ic.	<i>Plantarum rariorum.</i> 2 vols. fol.
	<i>Scitainimcs or Monandrous Plants.</i>	Schm. ic.	<i>Haarlem, 1793.</i>
	1 vol. fol. Liverpool.	Schm. ic.	<i>Schousboe (P. K. A.). Iagtagelser</i>
	<i>Rostkow (Fr. Gul. Ih.). Monogra-</i>	Schm. ic.	<i>over vaextriget i Marocco.</i> 4to.
	<i>phia Generis Junci.</i> 8vo. Berolini,	Schm. ic.	<i>Kiobenh. 1800.</i>
Roth abh.	1801.	Schm. ic.	<i>Schrank (Fr. von Paula). Plants</i>
	<i>Roth (Alb. Wilh.). Botanische Ab-</i>	Schm. ic.	<i>rariores Horti Monacensis.</i> Fol.
ttox. cor.	<i>handlungen und Beobachtungen.</i>	Schm. ic.	<i>Munich, 1817—1819.</i>
	4to. Nuremberg, 1787.	Schm. ic.	<i>Von Schreber (Joh. Christ Dan.).</i>
	<i>Roxburgh (William.). Plants of the</i>	Schm. ic.	<i>Icones Plantarum minus cognita-</i>
	<i>Coast of Coromandel.</i> 2 vols. fol.	Schm. ic.	<i>rum. Decas 1.</i> Fol. Hals, 1766.
Rtb. gr.	London, 1795—1798.	Schm. ic.	<i>Id. Beschreibung der Grsscr.</i> 2 vols.
	<i>Rottboll (Christ. Friis.). Descriptiones</i>	Schm. ic.	<i>fol. Lips. 1769—1779.</i>
	<i>et Icones Plantarum—rariorum.</i>	Schm. ic.	<i>Schwegrichen (Frid.). Species Mus-</i>
Ru. am.	Fol. Hafnie, 1773.	Schm. ic.	<i>corum Hedwig Supplementum.</i>
Rud. gui.	<i>Rumphius (George Everh.). Herba-</i>	Schm. ic.	4to. Lips. 1811.
	<i>rium Amboinense.</i> 6 vols. fol.	Schm. ic.	<i>Scopoli (Joh. Ant.). Flora Carniolica.</i>
	Am*st 1750.	Schm. ic.	8vo. Vienna?, 1760.
Rus. al.	<i>Rudge (Edward.). Plantarum Guians</i>	Schm. ic.	<i>Id. Deliciae Florae Insubricis.</i> 3 vols.
	<i>rarioiri Icones et Descr.</i> Fol.	Schm. ic.	fol. Ticini, 1786—1788.
	London, 1805.	Schm. ic.	<i>Seba (Alb.). Locupletissimi Rerum</i>
	<i>Russell (Alex.). Natural History of</i>	Schm. ic.	<i>Naturalium Thesauri Descriptio.</i>
	<i>Aleppo.</i> 1vol. 4to. 1756. Ed. 2.	Schm. ic.	4 vols. fol. Amst 1734—1765.
Sab. rom.	1794.	Schm. ic.	<i>Sebastiani (Ant.). Romanarum Plan-</i>
	<i>Sabbati (Liberatus.). Hortus Romanus.</i>	Schm. ic.	<i>tarum.</i> 2 vols. 4to. Rome, 1813
	7 vols. fol. Rome, 1772—1784.	Schm. ic.	—1815.
Sal. st	<i>Salisbury (Richard Anthony.). Icones</i>	Schm. ic.	<i>Seguier (Joh. Franc.). Plants Vero-</i>
	<i>Stirpium rarioiorum.</i> Fol. London,	Schm. ic.	<i>nscens.</i> 3 vols. 8vo. Verona, 1745.
San. vi.	1791.	Schm. ic.	<i>Seringe (N. C.I. Musec Helvltique</i>
	<i>Santi (George.). Viaggi al Mont Amiata</i>	Schm. ic.	<i>d'Histoire Naturelle.</i> 4to. 1818—
	<i>e per la Toscana.</i> 3 vols. 8vo.	Schm. ic.	1820.
Savi d.	Pisa, 1795—1806.	Schm. ic.	<i>Shaw (Thomas). Travels in Barbary</i>
Savi med.	<i>Savi (Gajetanus.). Dissertation Phaseolus.</i>	Schm. ic.	<i>and the Levant.</i> 1 voL fol. 1738.
avi ph.	<i>Id. Materia Medica vegetable Tos-</i>	Schm. ic.	<i>Supplement, 1746.</i>
Savi pis.	<i>cana.</i> Fol. Firenze, 1805.	Schm. ic.	<i>Sloane (Hans). A Voyage to Madeira,</i>
	<i>Id. Memoir on Phaseolus.</i>	Schm. ic.	<i>Barbadoes, Nevis, St Christopher's,</i>
	<i>Id. Flora Pisana.</i> 2 vols. 8vo. Pisa,	Schm. ic.	<i>and Jamaica.</i> 2 vols. fol. London,
Sc b. j.	<i>Schrader (H. A.). Journal flir die</i>	Schm. ic.	1707.
	<i>Botanik.</i> 10 fasc. 8vo. Got-	Schm. ic.	<i>Smith (J. Edward). Exotic Botany.</i>
Sc. dansk.	<i>Id. Det Konglige Danske Landhuis</i>	Schm. ic.	1804—1808.
	<i>holdings-sclkabs Schrifter.</i> 4 vols.	Schm. ic.	<i>Id. Plantarum Icones hactenus in-</i>
Sc. gen. nov.	8vo. Kiobbenhavn, 1776—1794.	Schm. ic.	<i>editte fasc. 1—3.</i> 1789—1791. Fol.
Sc. ger.	<i>Id. Nova Plantaxum Genera.</i> 1 fasc.	Schm. ic.	<i>Id. Icones picti Plantarum rari-</i>
	Fol. 1797.	Schm. ic.	<i>orum.</i> Fol. London, 1790—1793.
Sc. hrl.	<i>Id. *ijwGermanica.8vo. Gotting*</i>	Schm. ic.	<i>Id. A Specimen of the Botany of</i>
		Schm. ic.	<i>New Holland.</i> 4to. London, 1793.
Schan.	<i>Id. De Halophytis Pallasii.</i> 4to.	Schm. spic.	<i>Id. Spicilegium Botanicum.</i> Fol. Lon-
	<i>Id. Sertum Hanoverianum.</i> Fol.	Schm. spic.	<i>don, 1791—1792.</i>
Sc. phil.	Götting. 1795—1796.	Schm. spic.	<i>Id. Tracts relating to Natural His-</i>
Sesam.	<i>Id. Dissertation on the Genus Phi-</i>	Schm. spic.	<i>tory.</i> 1 vol. 8vo. 1798.
	<i>ladelphus.</i> Not published.	Schm. spic.	<i>See Mlm. mosc.</i>
	<i>Id. Systematische Sammlung Krypt-</i>	Sow.	
	<i>ogamicher Gewachse.</i> 2 fasc.	Sow.	
	«vo. Gfettingen, 1796,1797.	Sow.	

Spar. h.	Sparrman (Andr.). Kongl. Svenska Vetenskaps Academiens Handl. gar. 8vo. Stockholm, from 1739 to 1816.	Th. up.	Thunberg (Car. Petr.), in Acta Literaria et Scientiarum Upsaliac aut ab Academia Uxatalensi pubheata, 1750—181&
Spr. b. m.	Sprengel (Curt), in Berhnisches Magazin.	Til. pis.	Tilli (Mich. Aug.). Cataiogus Horti Pisani. Fol. Florence, 1723.
Spr. cryp.	<i>Id.</i> Introduction to the Study of Cryptogamous Plants, translated from the German. 8vo. 1813.	Todefun.	Tode (Hcnr. Jul.). Fungi Mecklenburgenses selecti 4to. Luneburg, 1790.
Spr. end.	<i>Id.</i> Neuc Endeckungen un Ganzen Umfang der Flanzenkunde. 8vo. Lipsia, 1820.	To. it	Tournefort (Pitton de). Relation d'un Voyage au Levant 2 vols. 4to. Paris, J717.
Spr. hal.	<i>Id.</i> Flora Halensis Tentamen novum. 8vo. Hals Sax. 1806.	Tou. ins.	<i>Id.</i> Institutions Rei Herbariae. 3 vols. 4to. 1717.1719.
Spr. urn.	<i>Id.</i> Plantaruin Umbellifrarum Prodromus. 4to. Halsr, 1813.	Tourn. vo.	See To it
Stack, fuel	Stackhouse (John). Nereis Britannica. Fol. Bath, 1795—1707.	Toz. mu. d. fir.	Tozzetti. Musco di Firenze.
Stcr. sax.	Sternberg (Graf. Casp.). Recvibio Saxi fragarum Icombus illustrata. 1 vol. fol. 1810.	Trat ar.	Trattinnick (Leop.). Archiv der Gewachskunde. 4to. Vienn. 1811 —1812.
Steer, stra.	Stercck (Ant.). Libellua de Stramonio, Hyosiamo, Aconito. 8vo. 1763.	Trat. au.	<i>Id.</i> Flora Austriaca sitca. Fol. 1792.
Stur. deu.	7 Sturm (Jacob). Deutschlands Flora. 4to. Many volumes. Nuremberg, 1798—1829.	Trat tab.	<i>Id.</i> Observations Botanies Tabularium Rei Herbaria illustrantes. 4to. 1811.
Stur. ger.	<i>Id.</i> j. Cistinace. The Natural Order of Cistus, or Rock-Rcs. 1 vol 8vo. London, 1830.	Trat th.	<i>Id.</i> Thesaurus Botanicus. Fol. Vicn. 1819.
Sw. au. ic. in.	Sweet (It.) Flora Australasica. 1 vol 8va London, 1837,1828.	Tr. ehr.	Trew (Christ Jac). Plants select@ ab Ehret picte. Fol. 1750—1773.
Sw. cis.	<i>Id.</i> Cistinace. The Natural Order of Cistus, or Rock-Rcs. 1 vol 8vo. London, 1830.	Trev. b.	Trevirinus (Lud. Christ), in Berlin. isches Magazin.
Swert fl.	. Swertia (Einan.). Florilegium tractans de varii Floribus. 1 vol fol. 1613.	Trev. del.	<i>Id.</i> De Delphinio et Aquilgia Observations. 4to. 1817.
Sw. fl. gar.	Sweet (R.). The British Flower Gar. den. 8vo. London, published in monthly numbers.	TrL ob.	Triumfetti (Joh. Bapt.). Observations dc Ortu ct Vegetatione Plantarum. 4to. Rome, 1685.
Sw. fl. gar. ic. in.	<i>Id.</i> British Flower Garden. 1823 to the present time, 3 vols. of the old series; and a part of a volume of the new series, 1830.	Tr. pi.	Trew (Christ Jac). Plantae rariores. Ed. J. C. Keller. Fol. 17G3.
Sw. ger.	<i>Id.</i> Geraniac. 8vo. Published in monthly numbers.	Turn. fuci.	Turner (Dawson). Historia Fucorum. 3 vols. fol. London, 1802, &c
Swz. b. m.	Swartz (OU) in Berlinisches Magazin.	Us. an.	Ustcri (Paul). Neue Annalen der Botanik. 8vo. 1794.
Swz. fil	<i>Id.</i> Synopsis F^licum, earum Genera et Species complectens. 8vo. Kilis, 1806.	Valil cc.	Vahl (Martinus). Elogac Americana Fol. 1796.
Swz. fl.	<i>Id.</i> Flora India* Occidentalis. 3 vols. 8vo. ErlangsR. 1797. 1800. 1806.	Vahl ic.	<i>Id.</i> Icones Plantarum in Elogis de scriptarum. Fol. 1798.
Swz. i c	<i>Id.</i> Icons Plantarum Indiae Occidentalnis. 1 fasc. Fol. 1794.	Vahl sy.	<i>Id.</i> Symbols Botanies. Fol. Hafh. 1790—1794.
Swz. ob.	<i>Id.</i> Observations Botanies. 8vo. Erlanga*, 1791.	Vail, it	Vaillant (Sebastien). Botanicon Parisiense, 1 vol. folio. Leid. 1727.
Swz. pr.	<i>Id.</i> Prodromus Descriptionum Vegetab. Indiae Occidentalnis. 8vo. Holms, 1758.	Vail pa.	<i>Id.</i> Botmicon Parisiense, Operis majoris Prodromus. 8vo. L. Bat 1723.
Tab. ic.	TaLernemontanus (Jac Theod.). Ei. cones Pldiaturam, cur. N. Bassso. 4to. Frankf. 1590.	Vau. con.	Vaucher (Jean Picr.-e). Histoire des Confreres d'Eau douce. 4to. Genve, 1803.
Tab. kr.	<i>Id.</i> Krauterbuch. Fol. IJ88.	Vcn. eels.	Ventenat (Etienne Pierre). Description des Plantes nouvelles ou peu connues du Jardin de J. M. Cels. Fol. Paris, 1800.
Tau. h. c	Tausch (Ign. Fred.). Hortus Canalius. Fol. part 1. Prague, 182a	Ven. ch.	<i>Id.</i> Choix des Plantes. Fol. Paris. 1803—1808.
Th. act haf.	Thunberg (Car. Petr.), in Acta Literaria Universitatis Hafniensis. 1 vol 4to. Hafn, 1770.	Ven. mal.	<i>Id.</i> Jardin de la Mulmaison. Fol. 1803—1805.
Th. gar.	<i>Id.</i> Dissertatio de Gardenia. 4to. Upsal, 1780.	Ven. tiL	<i>Id.</i> Monographic du Genre Tilleul. 4to. Paris, 1802.
Th. h.	<i>Id.</i> in Kongl. Svenska Vetenskaps Acadeniicns Handlingar. 8vo. Stockholm, 1759—1816.	Vib. ce.	Viborg on the Cerealia.
Thore j.	Thore (Jean). Observations in Journal de Botanique.	Vig. his.	Viguier (L G. A.). Histoire Naturelle des Pavots et des Arjmoncs. 4to. Montplicr, 1814.
Thou. af.	Du Petit Thouars (Aubert). Histoire des Vg^taux recueillies dans les lieux Australes d'Afrique. 4to. Paris, 1806.	ViL deL	Villars (D.) Histoire des Pinntos du Dauphinl. \$ vols. 8vo. Grenoble, 1786—1788.
Thou. or.	<i>Id.</i> Histoire des Plantes Orchid'es recueillies sur les trois Isles A us. trals d'Afrique. 8vo. Paris, 1822	Vil. s. j.	<i>Id.</i> in Schrader's Journal für die Rotanik. 5 vols. 8va Gottingen, 1803—1810.
Th. p.	Thunberg (Car. Petr.), in Novi Com. mentari Academic Imperialis Pctronolitans. 20 vols. 4to. 1720 —1776.	ViL stras.	<i>Id.</i> Catalogue m'thodique du Jardin de Strasbourg. Svo. 1807.
Thun. dra.	<i>Id.</i> Diss. Dracaena, 2 vols. 4to. Up- sal, 1780.	Viv. an.	Viviani (Dom.). Aunali di Botanica. 2 vols. 8vo. Gnova, ItO2, 1803.
Thun. er.	<i>Id.</i> Dissert de Erica. 4to. Upsal, 1785.	Viv. ft.	<i>Id.</i> Flors Italics Fragmenta. 4to. Genoa, 18C8.
Thun. fie.	<i>Id.</i> Diss. Ficus Genus. Upsal, 1786.	Viv. lib.	<i>Id.</i> Flors Libycae Specimen. Fol. Genoa, 1824.
Thun. ir.	<i>Id.</i> Dissertatio de Iris. 4to. 1782.	Voi nor.	Volckamer (Joh. Georg.). Flora Noribergensis. 4ta Noribergs, 1700.
Thun. jap.	<i>Id.</i> Flora Japonica. 8vo. Lips. 1784.	W. ach.	Willdenow (Car. Lud.). Tractatus de Achillei* ct Tanaceto. 8vo. Hals Magd. 1789.
Thun. pr.	<i>Id.</i> Dissertatio de Protea. 1781.	Wa. fl. hcl.	Wahlenberg (Georg.). De Vegetatione Helvetica. Svo. Furic 1813.
Thun. v.	<i>Id.</i> Voyage au Japon, ■■■ Svo.	Wa. lap.	<i>Id.</i> Flora Lapponica. Svo. Berlin. 1812.

Lie.	Wallicti (Nmh.). Planfer rnrjorca Asiatic*. 2parU,fal. !,oiiion.1830.	Won, er.	Woniland (Job. Clirist). Ericarum 1 cones et Dracriptlanffl. 4ti>, III, &c.
Jrf.	Asiatic Researches, or Trans- -;ii'tiins of ihu Society instituted. in Rwihi, -Ito. 1788.	Wen. tian.	Id. Sct-tum Hanoveratmrn. 4 faic. FoL
WaJtca.	Walter (T'lom.). Flora Caroliniana. Rvo. I'mlon, 1788.	Won. bar.	Id. Hfrttu Herrenhusanus. Fol. 1798.
Walt h.	Woltber; Atijr.Krt!..). ncstpnatio Plan. tarum Hrti cjs Bvo. UH 1731	Wen, ob.	/rf. BotnuiJirhc BcobnchtuiiEen. 1 voL 4in. I Lumver,
	WaUrteta (Franc.) et Ktaibd T-my. Doseriploiotes ct tc>m's l'anliiruin rnrlorum Hungnrite. 3 vols. l'ul. VIenn. 1803—1813,	W. etf	"Willtonow (Cnr. Lnd.i. In Acts Ac*. ileinO! electaralii MnguntinK Sd- entisrura UHIIHUI qon E^fordiat. 2 vols. BIOL Kriorif ct Gotbi J7ffl i the r^< in 4tu, ll'i:> and 177.
Wang.	Von Wangenheim (Fried Ad. JuL). Aiiplliuniung North Ametkani. char Holwtsn, 1 vol. folio. OottingK J787.	Wern, tr.	Triiins.icLioni: r>r Memoirs of tie Wer- ii-ty. 8vn. Edinburgh, . to the jivpsfit time,
Wi b. m.	Willilcuow (Car. Lurt.). Historic Amu- rtithor, Pol. Turid, IJKK lit. In tterJntsch: Mngzin exler GC wminleu; ifchriten, &c 4ta 1765 Ti7.	W. 1). b.	WilUienow tCar. Lud. I lortus Bero. liuctntii. Fol. Berlin, 18U6—1810.
Weig. ob».	^1 (Christ. Ehrntj.). Observa- tiones Botanicit>. 4<, Gryphia?, 177S.	W. n. her	Id. in Ni-wen Schriften <> Hi?rlin- k-hM Gcsetldfhad naturfoi den (reutirtc. 6 vols. Svo. Berlin, 1780-
Wein.	titMnn (J. Gull. Phytantiowi Ic(Hior;juJuci. 4vol& lol. ilegemh. 1737—nis, [Juh. Chriit.). Collectio l'plantaruin Uin exoticarum fju.mi iniigtarum. 4U. Haliovtt-ric, 1805, &c.	Wood.	WoodTUielWUliua}. Medical Botanj, 3 vols. «o. ITKL
		W. 11.	Willtk-imw (Car. Lud.). Phytopraphnia. 1-1. Krlmgj, 17i7.
		Za.	Zanoo (Antonio). Istoria Botanic*. Fol. Ilolognn, K75,
		Ztt	Zorn (Barthol!). Icone* PlsntaniM 'nnium. Bvo. Nurcmrx 17U —1784.

METHODS OF NAMING AND NUMBERING PLANTS.

In gardens or garden scenery where accuracy and intelligence are aimed at, all the less common plants, ligneous and herbaceous and tvli the variotios of fruits, should h:tvc ijTfir names affixed to them ; with the addition of llivir native ouintry, year of introduction if exotic, and natural order. Larch, oak, and yeiv are the most drirulje timbers for forming tallies en which to write or paint these names. The cheapest and best description of cast-iron tally is that of Mr. Stuart Murray (Caret. Mag. vol. Hi..p. 29.), the next bust thn used in the garden of the Horticultural Society. (Garil Mag, vol. vi. p. 407.) For trees, shrill^, and large plants, tho names may be painted on the ends of bricks, as in M-T-oddiges's arhuretiiii • or, what is belter, there are i>inctled bricks in itnition of Smrt Miii-ray's tally, and earthenware tallies <ith j.;inel, and with the names impressed by types on the clay before bum ing, manufactured by Patkeof TonstaQ near NewtastU under Lyne, iiml by Allurdyec of Clay Hills near hbmdtma. [Gard. Hag. 'o, vi, p. 399.]

Where plants cannot be named, they should hsve number* affixed to Uiem referring to L cataloguej and that theaanmmben maybe eenendlj ondCTstood, the common nun should ue made use of. TbaaB may be painteo on woud, iron, or earthenware tallies. But as ihe common numerds require Ih* nte of the pea or pencil, gardeners have de^is^d si for them which may be readily cut wiUi the knife on slicks.

SETOJJS MODE Or CUTTINO NU 31 BE US ON STICKS

Is much the simplest, and at the same time the most comprehensive, The signs made use of are as under. —



The advantages of these murks over those- in common use are, that they ar (and more easily made by sin^ly distinct cuts; und that no number, however high, reniirw more marks Ihaao it would require figures written with a pun. The only way En winch the memory is apt to misgive, in this scheme, is by confounding / and \, si ami K, A and y, with each other, u a cliild would confound the figures G and 9; but this slight inconvenience will be remedied by die following Uy, which may bt' easily b>niu in the mind. Li-t us recollect Out, in writing, wo naturally draw a stroll- ttora the ri^lit at top, to the at bottom, thus /, and not in the opposite direction, thus \ : now, in all the above numbers which differ from tach other in the direction »f tfo diagonal line, that wbith is in the direction usual in writing precedes the other, thus / \ sy N /, ,\; the other two, £ 3 6 1 8 9

A and V i will not be confounded, on recollecting that V >s the usual numeral letter for five.

As an example of application, suppose we take species 341, variety 8, the tally cut thus,—



The long **cut** (—) between the number of the species and the number of the variety is used to separate the **one** from the other, and is to be considered as the sign of separation. By the use of this **sign** several numbers may be cut on one stick; it is a fixed rule that the number still always read from the root or sharpened end of the tally, because it is necessary for the operator to hold that end in his hand in cutting the marks. Names written on tallies, on the contrary, should always read from the top, by which means the letters are kept at the greatest practicable distance from the soil.

DRYING PLANTS AND FORMING A HERBARIUM.

THE best mode for gardeners, and such as have little spare time to dry specimens of plants, is to gather them when perfectly free from exterior moisture, place them between paper, and press them with bag* of hot sand. In this way they dry of their natural color in a few hours. (*Mag. Nut. Jist.* vol. i, p. 196.) Or they may be dried in sand (*Ibul.* vol. iii, p. 459.)

To fill up a space that would otherwise remain an unseemly blank, we shall then observe, that every **gardener** ought to dry plants and form them into a herbarium. The **herbarium** may be a folio volume of stout thick paper, white, if he can afford it; it's not, brown; allowing in binding the thickness of a leaf for every page, in order that, when filled with specimens, it may shut close and preserve its form. Mr. **Toward**, who has formed one of the handsomest gardener's herbariums we have anywhere seen, in four thick folio volumes, uses cartridge paper, binding the back of each leaf a leaf of brown blotting paper, and round the margin of both surfaces of the double leaf so formed strips of cartridge paper. These strips rather more than compensate for the thickness of the dried specimens; so that when the leaves are bound up, their edges cut, and the books shut close, the external air is excluded, and the appearance as neat as that of any printed volume. (*Gard. Mag.* vol. iv. p. 4:16.) The specimens, if dried in sand, or between loose leaves of paper or in any old book, may be fixed to the cartridge side of the leaves of the herbarium book, according to any arrangement which may be preferred,

DRAWING PLANTS, FLOWERS, FRUITS, &c

EVERY gardener ought also to be able to draw **plants**; and this he may acquire himself by copying with a pen or pencil, without the aid of colors (for all the essential objects of drawings may be effected by lines), the outlines of the ribs, veins, and other inner lines of the different parts of the dried **specimens** as they lie flat before him on the paper. Having practised this for some time, he may then draw plants, flowers, and fruits from nature; beginning by washing one side of a pane of glass over with gum, holding that when dry between his eye and the plant or other object to be drawn, and then tracing the object with the point of a nail in the gum, or on it with the point of a black pencil. (*Gard. Mag.* vol. vi. p. SOS.) The outline of hard fruits, such as the apple and pear, may be correctly taken by dividing them into two parts, unless that of soft fruits if not too ripe. (*Gard. Mag.* vol. iv. p. 230.) Every gardener ought also to be able to draw landscapes, figures, and animals, all manner of games, pictures, and plans of gardens and maps of estates. This he may acquire **atmself** by copying the various engravings in our *Encyclopaedia of Gardening and Agriculture*; and for this, and landscape-scenery more particularly, by copying the elementary plate (Pl. I. sold by itself, under the title of *Elementary Detail for Pictorial Map-drawing*, for 3s. Gd.) of our *Illustrations of Landscape-Planning and Garden Architecture*. He may afterward* draw figures and landscapes from nature, first through a gummed pane of glass, and next on paper; and finally, measure and draw from nature implements and buildings, and map gardens, parks, and farms.

On correctly applying, pronouncing, and spelling the names of plants; on drawing plans neatly, and being able to produce; off hand a sketch of any thing; on being able to measure and value all kinds of country work; above all, on good penmanship, spelling, and writing a correct and intelligent letter, much of the success in life of a gardener depends. Without possessing these qualities, by a regular course of education at school, or determining to labour hard at self-instruction till he has possessed them, no young man in the present improving age should attempt the profession.

HO'RTUS BRITANNICUS.

PART I.

LINNEAN ARRANGEMENT.

of plants. For this purpose some one organ, common to plants in general, is fixed on: and, according to certain condition* in which this organ is found, individual species are referred to their places in the system, as word, by their Initial letters, are referred to their places in *U* alphabetical dictionary.

as words, by their initial letters, are referred to their places in C alphabetical dictionary.

II) to proffers of artificial syiisms dille/t-t/n organs have been filed on by diilinul Waniit*; but those which have been most extirrably employed are tie *eonBai* by Tonrnclfort, and the stamens and pistil* by Unnsiu. The »stem of I onrndrbt »;* been i RG+H deal employed in France, and may be comldeed as the artificial system or that country ; that of Lfimirus has been empWed in most other countries, and is justly esteemed by far the most perfect artificial syiem which has hitherto been produced. It b, therefore, adopted In this work.

wervw, " will soon parcel re that it is to be understood merely as a dictionary, to make nut any plant that inay fall in hia WHy." (*Gram. of Bot.*) " If we examine," says Decandol 1 e, " the first lici:d systems which hava been hitherto devized, we shall find the n^{ext} celebratc Of them, that which was propo^{ed} by Linnaus, to posse^s a dcclid superiority over all others, not only because it is consistently derived from one siupl^{er}eple, but also because the author of it, by means of n new nomenclature, has given to his terms the gi
«*in*tinctns» of meanin^g." (*Elements of the Philos. of Linnaas, by UcaimhUc <in> Sprengel*) V
11 of subsequent advance* in science may enable ix:tamMs to dispense with the Liimmen system altogether, it
"not for *ui* to aftlm ; but in the mean time nothing can be inure certain than that the Linnean system is the
nest leading arrangement for such a work as the present, in Uie existing »tale of botanical knowledge in
on tain. *

^aAccording to the Linnean »slom all plant* are furnished with flower*, either conspicuous OT loottupteuoui. ine plant!! with contpukuoi flowers are arranged according to the number and position ol ilicir stamens and P^{TTM*}; those with incontpicuoi Bowen areananged awctnliig to I lie situation of the flowers on the plant, or according to other circutTis-tanL-c* in the plant ilMIC

To discover the unnic of > plant by the Linnean system, therefore, all that it
iry for a beginner is to possess a specimen of It in flower, and to be able
to know it! different parts by the m n n given them by botanists. To d&korer
the CIOM, order, ami genus of a plant, it is only necessary to be able to distinguish and
name the (lif'eretit part* of the (lower. TheM [Kirts are: the cabx or cup /J'tg. 1.a),
which in that leaf, or thane leaves, by which the flower is usually enclosed when in a
hud, ami which, when the flower Is expamtwl, appear under it. The corolla
(corona, a crown) is the colored leaf, oi leaves, of a flower /Jtg. 1. It). Ike Btamt-n
(or fwlt principle of any thing) is the thread-like proews, or processes, Imme-
ili.itly within the leaves or the corolUi (Jtg. 2.): It cotwlfils of two parts, the Slfment
of umad /a) and the anther /> i ibis nnher contains what is called liejtollen, or
Iviiik meal \. In the centre ill the flower is the pistil /Jtg. Sj * it conaists of
thmM parts, the y,t-t- nu of the fruii

The pistil and utamen are the essential parts of a flower. The corolla or the calyx may be wanting, and yet the flower will live, though perfect, because the absence of those parts is no obstacle to reproduction. Even the style would be absent without preventing the fruit from ripening or the fruit from the flower; and there are many flowers which the author has seen which have no corolla, yet without a style, and the stigma to the germin without a style; but the style, the germin, and the style are essential.

The scil is contained in the pericarp, or seed-vessel, which is the germen when j^own to maturity, name of the seed-~~vi~~-••• VLirism-rnrling to tu form, substance, &c.; but the word pericarp (''''> about, *karpon*, fruit; is aptneite (o all its vari : the base ur medium lvieli connects the other parts of the fructification. (*Mazimiuu-srf Nnlurat History*, vol. I, p. 618A)

The degree of knowledge conveyed by the following Table, init the preceding observations, will enable a "alnncr to discover at least the das* and ordex of any plant wbleh he may find in flower.

Fittsr GkAKD DIVIHIU*.*- *Plants with conspicuous Flowers (Phaneroglintia).*

n illc larnd (lover).
'Malt :nd l'in,i... organs ell .rim¹.
not un.tcd elihiT atom or Ivlow,

	t' 1. Monihadria (<i>nvMfuj one</i>)
Two,	4. <i>Oiifr, n iiiiui</i> , <i>Ki.unplc.</i> 1 <i>aman).</i>
Thht,	f 5. <i>Trlandrla</i> [<i>tru. Uiritt, L uner, lIMI</i>]. <i>I'm ton.</i> 14
Four,	f 1. <i>Tclrnixtria</i> (<i>Mm, bar, fl^w^, II m^ii L i</i>)
Fltip,	f 5. <i>PomaElUiiA</i> (<i>ifififi*, atw, nrnnn</i>). <i>i.j</i>
Sin,	r 6, <i>Hi nJndria</i> (<i>At^ (Ii, mirr, man.) icila b>ioll.a</i>) s <i>I 1)6</i>
Seven,	t \ <i>atter, q man), ^hcultur</i> t <i>Hl^HJciAtalium.</i> 143
	V B> <i>OctdiuWa</i> (<i>mW>, i-lchl,</i> I <i>qfed<i><i.</i></i> 144
Mm,	f 9. <i>EnntSmrlrlft</i> (<i>mnM, nipe, lf)</i>
Ten,	(10. <i>ihcHMdirI</i> <i>iMy, ten, uiwr, a man)</i> <i>Saff&u'B umlrbta. lStt</i> V "• <i>Ilodccrfncrbi</i> (<i>rfttlaw,</i> rum, 1S7
Miinr, <i>fnqiimtl(ji)</i> <i>Iji,coaandrlB(Ti(vn', twrnij, aner, amin), fraia*</i> IVL <i>utacitlnl></i> <i>lothii?, lslij,</i> <i>Oijacincnlh.</i> 1U4	j (P
II'M	
... <i>h'fli II h'fliETII</i>	
Tva lorm, ami two ihurt,	f 14. <i>DicJnjnJa</i> (<i>iiii, tvite, duwtfluu, paw'a; *upi rtM Wy of twiglatiniKii</i> t <i>TsflerwMi Iceld.</i> IH
Knur Long and ibort.	ITDJ
	ttu ultf ill.
united,	
lie t)jmcenn, rinlo ono bottj,	I 1G. <i>Moiiad^nhla</i> (<i>irictHM onr. WIWIJKH. lit</i> I <i>ttlUra frierant.</i>
Into two bodi»,	
into many bodlos,	I-s. P,
(hi onthrt* or 1 <i><i</i> *, f *•* Into A ollmkcr,	- j
(itwIMni), >t-1 40. <i>fynnifTidri*</i> <i>tfiVIi » <*</i> ilkLuillin'uiwii, U <i>man, our, a</i> 11 <i>inn</i> SS6	
uinc plant.	R i mpn^*, mpn home). 6ticdr^jita
remt >Unu,	f 21. <i>DSirVia</i> (<i>rfij, ivfct.nUH hcucic), fannabl u</i>
mtordWiiTTntplimiTV •	
Hili hennatibtxUM-i	

SKCOKDGEAND DIVISION.—*Plants with inconspicuous Flowers {CryptogAuna}*

9. F/l<<aOUiri i'>B (*^"«^ t>r.'.


HO'RTUS BRITA'NNICUS.

PART I.

LINNEAN ARRANGEMENT.

CLASS I. ORDER 1.

MONA'NDRIA MONOGY'NIA.



Kistematic Name and Authority.	EnRlish Name.	Native Country.	Reference to FfeUKS.
• 1. CA'NNA L.	INDIAN SHOT.		(<i>Canna</i> , a cane or mat, <i>Celt.</i>) 34.-35.
1 patens <i>Rote.</i>	Indian	Rio Jan.	1778. R r.m Bot.rcg.576
2 indica L.	spotted	India	1570. R r.m Red HI. 201
2 maculiflta	scarlet	India	... R r.m Hook, ex fl.53
3 coccinea Rose.	yellow	& Amer. 1731. R r.m Bot. mag. 452	
4 h'ltea Rose.	Lambert's	E Indies 1629. R r.m BoL mag. 2085	
5 LamMrti B. M.	gigantic	Trinidad 1819. R r.m Bot.rcg.470	
6 gigantia Desf.	western	S. Amer. 1809. R r.m Bot. rcg.20fi'	
7 occideitalis Rose.	woolly	W. Indies 1822. R r.m Bot.rcg.772	
8 lanuginosa Rose.	flesh-colored	Maranh. 1823. D r.m Hose. scit. ic.	
9 carnæ Rose.	wild	Brazils 1822. D r.m Rose, wit ic.	
10 sylvestris Rose.	orange	S. Amer. 1820. D r.m Rose, scit ic.	
U aiirantara Rose.	multitorm	Brazil 1824. D r.m Rose, scit ic.	
12polymorpha/o.C.	blody	S. Amer. 1825. D r.m	
13sanguinealo.C.	esculent	S. Amer. 1822. D r.m	
Hesculentia L.o.C.	lanceolate	Brazil 1825. D r.m	
14 lancolata L.o.C.	bordered	Brazil 1818. R r.m Botreg.771	
16 limbata Rose.	variable	India 1822. R r.m	
17 variabilis W.	red	W.Indies 1820. R r.m	
18 rubra IV.	red-stemmed	 1821. R r.m
19 rubncaulis Lk.	eatable	Peru 1820. R r.m Bot.reg.775	
20edtli5.i1.	showy	S. Amer. 1820. tt r.m Bot mag. 2317	
21 speciosa B. M.	lottv	Brazil 1820. R r.m Bot cab. 743	
22 excisa Lod.	broad-leaved	Brazil 1820. R r.m BoLcab. 834	
23 latifolia Lod.	narrow-leaved	S. Amer. 1824. R r.m Pis.br. ic.	
24 angustifilia L.	apmy S	f. Indies 1820. R r.m Row. scit Ic.	
24 orientalis Rose.	oriental		
chinenis IV. en.	4 my.jn H.Y		
2Idva	yellow	E. Indies 1820. R r.m Rose, scit ic.	
SGcompacta Rose.	compact	E. Indies 1820. R r.m Rose, scit ic.	
27 denudata Rose.	naked	Brazil 1818. R r.m Rose, scit ic.	
2 latifolia	pruineous	Brazil 1818. R r.m Rose, scit. ic.	
28 pallida Rose.	yulc-flowering	W. Indies 1820. R r.m Rose. scit. ic	
2 latifolia	broad-leaved	W.Indies 1820. R r.m Rose, scit ic	
29 crocea Lac.	safr-on-colored	 1823. R r.m
30 pedunculata B. M.	stalked	O. Indies 1820. R r.m Bot mag. 2323	
<i>jinicea</i> Retz.	rushy	S. Carol. 1788. R r.m Sal. st. 3.2	
32 flaccida Rote.	flaccid	S. Amer. 1732. R r.m Ex. bot. 2.102	
33 glaficaL.	prdfa	S. Amer. ... R r.m Bot mag. 2302	
34 iridiflora/7.*r.	rufoous	Peru 1816. R r.m Bot mag. 1968	
* 2. MARANT/f L	irid-flowered		
« arundinacea L.	ARROW ROOT.	(<i>B. Maranti</i> , a Venetian physician, d. 1554.) <i>Cunnea</i> . 14. — 20.	
% Indica 7W.	rccd-like	S. Amer. 1732. k s.l Bot. mag. 2307	
37 gracilis Rud.	Indian	E. Indies 1800. D s.l SI jam. 149	
# divaricata Jtos.c.	slender	Guiana 1B21. D s.l Rud. gui. 3	
2purpurascens	divaricate	E.Indies 1718. R r.m Rose. scit. ic.	
T Pteal Jlf.r.	purpurascens	Brazil 1823. R r.m Rose. scit. ic.	
41 angustifilia B. M.	oblique	Guiana 1803. R s.l Rud. gui. 2	
42 Tonchat Aub.	yellow	Caracas 180U. R 61 Jac.12. 201	
43 gibba Sm.	harrow-leaved	W. Indies 1820. R 81 BotTnag. 2398	
44 sylvatica Rose.	wood	E. Indies 1819. R 81 Ru.am.4.7	
« variegata X.o	petiolated	E. Indies 1818. R s.l Kosc. scit. ic.	
47 blcolor JCf.r.	variegated	Malacca 1818. D s.l	
« malaccensis W.	two-colored	E. Indies 1800. D s.l	
	Malacca	G. 2 jnjl Y.W E. Indies 1820. D 81 Ru. am. 5.71	
* 3. CALATHEA ML? CALVTHEA (<i>Kata/hot</i> , a basket; from the form of the stigma.) <i>Cdnex</i> . 7.—9.	violet-colored		
Wzebrina Undl.	£ El or 2 year	R.Y Brazil 1815. H s.p Botreg.385	
▪ MardntozebrinaAJtf.			
«) viucca Lindl.			

MONANDRIA MONOGYNIA.

CLASS 1.

51 <i>flavescens Lindl.</i>	flavescant	£ El >r	4 au	Y	Brazil	1822.	D si	Bot. reg. 932
52 <i>Allouya Jac.</i>	Allouya	£ El >r	8	W	Trinidad	1824.	D s.l	
53 <i>longibractata B.R.</i>	long-bracted	£ El or	jl.au	P	Brazil	182a	D s.l	Bot. reg. 1020
54 <i>RiW Lo. C.</i>	Row's	£ El or	apjl	—	Brazil	1825.	D s.l	
55 <i>villosa Lo. C.</i>	villous	£ El or	apjl	—	Brazil	1825.	D s.l	
4. THA'LM L.	THALIA.	(J. <i>Thalius</i> , a German physician, d. 1588.)					<i>Cdnne. 2.—</i>	
56 <i>dealbata Rose.</i>	whited	*i AJ or	4 jl.au	B	S. Carol.	1791.	R p.l	Bot. mag. 1690
57 <i>geniculata Rose.</i>	jointed	£ El or	2 aus	B	W. Indies	1823.	D p.l	Pl. gen. 26.108
• 5. PHRY'NIUM L.	PHRTNIOM.	(P/irynos, a frog; a plant of marshes.)					<i>Cunnea. 14.—</i>	
58 <i>capitatum W.</i>	headed	£ El or	5 jl.au	W. P.	E. Indies	1807.	R s.l	As. res. 11. 3
59 <i>imbricatum Rox.</i>	imbricated	■ El or	my.jl	Pa.R	E. Indies	1818.	D s.l	
60 <i>dichotomum Rox.</i>	forked	£ El or	jl.au	W	E. Indies	1810.	R s.l	
61 <i>iarviflorum Rox.</i>	small-flowered	£ El or	jl.au	Y	E. Indies	1820.	R s.l	Rose, scit ic
62 <i>grandiflorum Rose.</i>	great-flowered	£ El or	jl.au	Y	Brazil	1823.	R 81	Rose, scit. ic.
63 <i>spicatum Rox.</i>	spiked	£ El or	jl.au	Y	E. Indies	1835.	R 81	Rose, scit ic
64 <i>Myrotha Rose.</i>	Myrrh-scented	■ El or	2 jl.au	W	S. Amcr.	1820.	R r.m	Rose, scit ic
	<i>Myrosma cannaefolia L.Jil.</i>							
65 <i>comosum Rose.</i>	tufted	£ ES or	3 jl.au	Y	Surinam	1812.	R r.m	Rose, scit ic.
	<i>Maranta comosa L.Jil.</i>							
66 <i>Parkeri Rose.</i>	Parker's	£ El or	2 jl.au	Y	Grenada	1823.	R r.m	Rose, scit ic.
67 <i>Cas&po Rose.</i>	Casupo	£ El or	13 jl.au	Y	S. Amcr.	1820.	R r.m	Rose, tscit. ic
	<i>Maranta Cas&po Jac.</i>							
68 <i>ellipticum Rose.</i>	elliptic	£ E) or	lijl.au	Pa.Y	Guiana	1822.	R r.m	Rose, scit ic
	<i>Maranta spicata Aub.</i>							
69 <i>virgatum Rox.</i>	twiggy	£ El or	8 jn.jl	W	EL Indies	1818.	D s.l	
70 <i>obliquum Lo. C.</i>	oblique	tf	tAlor	II my.jl	Y	E. Indies	1824.	D s.l
71 <i>angustifolium Lo. C.</i>	narrow-leaved	£ El or	1J my.jl	Y	E. Indies	1824.	D s.l	
• 6. HEDY'CHIUM Klin.	HEDYCIHUM.	(Hedys, sweet, chion, snow; flower.)					<i>Scitaminear. 20.—</i>	
72 <i>coronarium Rox.</i>	garland	£ El or	5 jh.s	Y	E. Indies	1791.	R p.l	Bot. mag. 708
73 <i>angustifolium Rox.</i>	narrow-leaved	£ El or	5 jn.s	S	E. Indies	1815.	R s.l	Rose, scit ic.
74 <i>aurantiacum Rose.</i>	orange-colored	£ El or	5 jl.au	O	E. Indies	1812.	D 81	Rose, scit ic.
75 <i>coccineum Rose.</i>	scarlet	£ El or	6 jl.au	S	E. Indies	1815.	D s.l	Bot. reg. 157
	<i>angustifolium B. R.</i>							
76 <i>gracile Rox.</i>	slender	£ EI or	3 jn	W	Bengal	1823.	R s.l	
77 <i>carneum Carey</i>	flesh-colored	£ El or	4 au	Pk	E. Indies	1H23.	D 1	Bot. mag. 2657
78 <i>longitònum Rose.</i>	long-leaved	£ El or	6 jn.jl	R	E. Indies	1819.	D 81	Rose, scit ic
79 <i>flavum Rox.</i>	yellow	£ El or	3 jn.au	Y	Nepal	1822.	II s.l	Bot. cab. 604
80 <i>flavescens Lod.</i>	flavescant	£ OS or	6 jn	Y	India	18S2.	R s.l	Bot. cab. 723
81 <i>hetromallum B. R.</i>	variable-haired	£ El or	3 jn.au	Y	India	1822.	R 81	Bot. reg. 767
82 <i>speciosum Wai.</i>	showy	£ El or	8 aus	Pa.Y	E. Indies	1823.	D s.l	
83 <i>maximum Rose.</i>	largest	£ El or	8 au	W	E. Indies	1820.	D s.l	Rose, scit ic
84 <i>c&tum R.Br.</i>	tall	£ El or	5 jn.d	Pa.R	K Indies	1818.	R s.l	Bot. reg. 526
85 <i>Gardnerianum Wai.</i>	Gardner's	£ El or	7 jn.au	Y	E. Indies	1819.	R r.s	Bot. reg. 771
86 <i>villosum Wai.</i>	villous	£ El or	3 jl.au	Crea	E. Indies	1823.	D 81	Rose, scit ic.
87 <i>glaucum Rose.</i>	sca-gie-en	£ El or	44jl.au	W	E. Indies	1822.	D s.l	Rose scit. ic.
88 <i>ellipticum Sm.</i>	elliptic	£ El or	5 JU	W	E. Indies	1804.	D 81	Rose, scit ic.
89 <i>acuminatum Br.</i>	acuminate	£ El or	4 jn.au	W	E. Indies	1820.	D s.l	Rose, scit ic.
90 <i>spicatum B. M.</i>	spiked	£ El or	3 jn	Y	India	1810.	R co	Bot. mag. 2300
91 <i>thyriflora Sm.</i>	thyrsiformed	£ E 3 or	4 jl.au	W	Nepal	1018.	D 81	Rose scit ic
7. ROSC(yEl Sm.	RO9COBA.	(W. <i>Roseoe</i> , the historian of the Medici.)					<i>Scilamince. 5.—</i>	
92 <i>purpurea Sm.</i>	purple	■ El or	jl.au	■	Nepal	1820.	K s.l	Ex. bot 97
93 <i>gracilis Sm.</i>	slender	■ El or	jl.au	■	Nepal	1821.	D s.l	
94 <i>elatior Sm.</i>	taller	■ El or	jl.au	■	Nepal	1820.	D s.l	
95 <i>spicata Sm.</i>	spiked	■ El or	jl.au	■	Nepal	1820.	D 81	
96 <i>capitata Sm.</i>	capitate	■ El or	jl.au	■	Nepal	1819.	D 81	
• 8. ALPI'NI! L.	ALPIMIA.	(P. <i>Alpini</i> , an Italian physician, d. 1615.)					<i>Scitamine. 24.—</i>	
97 <i>comosa Jac.</i>	tufted-spiked	£ El or	1 ap,jn	W	Caraccas	1752.	R s.l	Jac. ic. v. 3
98 <i>Galdnca Swz.</i>	Galanga	£ El dt	6 pf	W.T	R Indies	...	R 81	Ru. am. 5.63
99 <i>raccmosa Rose.</i>	racemose	£ El or	5 jls	W	W. Indies	1752.	R s.l	Pl. ic. 11.20
100 <i>occidentalis Swz.</i>	occidental	£ El or	6 jl	W	Jamaica	1793.	R s.l	
101 <i>ndtans Rose.</i>	nodding	£ El or	13 ap,jn	Rk	E. Indies	1792.	R 81	Ex. bot. 2.106
102 <i>diff'issa Rose.</i>	two-clft	£ El or	6 apny	P.B.Y	E. Indies	1818.	D s.l	Rose, scit. ic.
103 <i>auricul&ta Rose.</i>	eared	£ El or	4 apny	R.Y	E. Indies	1814.	D s.l	Ro&c. scit ic.
104 <i>exalt&ta Mey.</i>	lofty	£ El or	20 my,jn	R.Y	Surinam	1820.	D 81	Rose, scit ic
105 <i>cifrua B. M.</i>	drooping	£ El or	3 Pj	Pk	E. Indies	1804.	R 81	Bot. mag. 1900
106 <i>calcarata Rose.</i>	spur-flowered	£ El or	3 S	W	E. Indi's	1800.	R 81	Bot. rep. 421
107 <i>malaccensis Rose.</i>	Malayan	£ El or	5 ap-my	W	E. Indies	1799.	R 81	Bot. reg. SSS
108 <i>mOtica Rox.</i>	spurless	£ El or	5 w	W	E. Indies	1811.	R 81	Rose scit. ic.
109 <i>bracteata Rox.</i>	bracteate	£ E3 or	3 my,jn	W	E. Indies	182 K	D s.l	
110 <i>Antillidrum R. % S.</i>	Antilles	£ El or	4 my,jn	F	W. Indies	1826.	D s.l	Sl. jam. 105.2
111 <i>penicillata Rose.</i>	pencilled	£ iAJ or	3 my,jn	Pk	China		D s.l	Rose, scit ic
112 <i>spicata Rox.</i>	spiked	£ El or	2 ...	—	Sumatra	1822.	R r.m	
113 <i>striata Lk.</i>	striated	£ El or	4	—	E. Indies	1818.	D s.l	
114 <i>Roscocara R. & S.</i>	Roscoe's	£ El or	3 my	■	E. Indies	182a	D s.l	
	<i>bracteata Rose, non Rox.</i>							
115 <i>Cardamomum Rox.</i>	cardamom	£ El clt	8 au	W.p	E. Indies	1815.	R B.1	R. mal. 11.4.5
116 <i>tumuliflora B. R.</i>	tubular	£ El or	2 jl.au	R	Demerara	1820.	R s.l	Bot. reg. 777
117 <i>puncea Rox.</i>	scarlet	£ El or	6 mr	S	E. Indies	1820.	D 81	
118 <i>media Spr.</i>	mediate	£ El or	6 jl.au	R	E. Indies	1815.	D s.l	Lin. tr. 10.4, *
	<i>Cardamomum medium Rox.</i>							
119 <i>ingua?formis Box.</i>	tongue-formed	£ El or	6jl.au	R	E. Indies	1820.	D s.l	
120 <i>ALMghas Rose.</i>	Allug.ofCe>lonf	£ El or	2 fmr	R	E. Indies	1796.	R s.l	Bot. rep. 501
*9. HELLE'NI! W.	HELLENIA.	(C. N. <i>Hellenius</i> , Professor at Aba)					<i>ScitaminecE. 2.—5.</i>	
121 <i>cpfilea R. Br.</i>	blue-berried	£ El or	4 jn.jl	W	N. Holl.	182a	D s.l	Rose, scit ic,
122 <i>chinenis. W.</i>	China	£ El or	3 jn.jl	W	China	1825.	D s.l	
10. ZINGIBER Gae.	GINGER.	(The original Indian name.)					<i>ScitaminecE. 2.—14.</i>	
123 <i>panduriflum Rox.</i>	fiddle-lipped	£ El or	3 my-jn	Pk	E. Indies	1812.	R s.l	
124 <i>Mioga Rosa</i>	Mioga	A iAJ or	2 my.jn	Pk	Japan	1796.	R r	ra
125 <i>officinale-flocc.</i>	official	S El clt	2 jn.au	R	E. Indies	160b.	R r,m	Jac vin. 1.7*

126 <i>Zerumbet</i> Rose.	Zerumbet		Y.o	E. IndieB	1690.	n.r.m Ex. hot 2.112
127 <i>Casum&nar Rox.</i>	Casumunar		W.v	E. Indies	1807.	It r.m Bot mag. 1426
128 <i>purpureum</i> Rose.	<i>purple-bracted</i>			E. Indies	1796.	R s.l
129 <i>rosicum</i> Rox.	rosy			E. Indies	1822.	R s.l Box. cor. 2.196
130 <i>ribens</i> Rox.	red			E. Indies	1822.	R s.l Rose, scit ic
131 <i>ligulatum</i> Rox.	ligulate			E. Indies	1823.	1) s.l
132 <i>squarrosum</i> Rox.	<i>rough-spiked</i>			E. Indies	1822.	R al
133 <i>chrysanthum</i> Rox., ; ; <i>capitatum</i> Rose.	gold-flowered capitate			E. Indies	1821.	R s.l Rose, scit ic
135 <i>elitum</i> Rox.	lofty			E. Indies	1825.	II s.l Rosa scit ic.
• 11. CO STUS Rose.	Comm			E. Indies	1820.	R s.l
1% <i>arabicum</i> L.	Arabian					
137 <i>spicatus</i> IV.	spiked					
138 <i>speciosus</i> Rose.	snowy					
139 <i>ncpalensis</i> Rose.	Nepal					
• <i>peciosus</i> /3 <i>angustifolius</i> B. R.						
140 <i>afer</i> B. R.	African					
141 <i>maculatus</i> Rose.	spotted					
a Yer & <i>hirsutus</i> B. R.						
142 <i>spiralis</i> Rox.	spiral					
Alp. <i>spiralis</i> Joe.						
143 <i>Pisonia Lindl.</i>	Pison's					
144 <i>villossimus</i> Jac.	most villous					
145 <i>lanatus ho. C.</i>	woolly					
M2. KJEMPFERL* L.	GALANGALE.	[E. Kiemfifer, a German naturalist, d. 1716.)	Seitaminea.	7. - 8.		
146 <i>rotunda</i> L.	<i>round-rooted</i>	or	jlau	R. w	E. Indies	1761.
147 <i>Galanga</i> L.	<i>Galanga</i>				E. Indies	1728.
148 <i>anguillifolia</i> Jac.	<i>narrow-leaved</i>				E. Indies	1797.
149 <i>pandurita</i> Rox.	<i>Rddleshapett</i>				E. Indies	1797.
150 <i>latifolia</i> Bonn.	<i>broad-leaved</i>				E. Indies	1803.
151 <i>ovalifolia</i> Rox.	<i>oval</i>				Malacca	1822.
152 <i>marginata</i> Carey	<i>margined</i>				E. Indies	1822.
* 13. AMYUM Rose.	AMOMUM. (A. privative, <i>mo-mos</i> , impurity; & counter-poison.)	Seitaminea:	13. - 20.			
153 <i>Aftelia</i> Rose.	Afzelius's	^ 123 or	my.jn	Pk	S. Leone	1795.
154 <i>grandiflorum</i> Sm.	large-flowered	E or	3 jnjf	W	S. Leone	1797.
la5 <i>anguifolium</i> Rox.	narrow-leaved	E or	8 jnjl	R	Madagascar	...
156 <i>dealbatum</i> Rox.	whieed	E or	3 xmr.ap	W	Bengal	1819.
157 <i>sylvestre</i> Sum.	wild	E or	1 mr.ap	W	W. Indies	1819.
158 <i>tardamumura</i> L.	<i>gftia/Zcardamom</i>	E or	4 my.jn	pa.Br	E. Indies	1820.
159 <i>aculeatum</i> Rox.	prickly	E or	10 ap.my	O	Hindu*	1819.
160 <i>latifolium</i> Afx.	broad-leaved	E or	4 ap.jn	P.Y	& Leone	1824.
161 <i>subulatum</i> Rox.	subulate	E or	3 mr.ap	Y	Bengal	1822.
162 <i>Grana Paradisi</i> W.	grain of parad.	E or	5 fmr	R	Madagascar	...
163 <i>mlimum</i> ROT.	greatest	E or	5 jn	W	E. Indies	...
164 <i>aroiriticum</i> Rox.	aromatic	E or	3 my.jl	P.Y	E. Indies	182a
165 <i>sericum</i> Rox..	silky	E or	6 jnj.jl	W	E. Indies	1819.
• 14. CURCUMA L.	TURMERIC.	(Kurkum, the Arabic naracj	Seitaminea.	20. —		
166 <i>Zedoaria</i> Rose.	Zedoary	^ E3 or	3 ap.ap	R	E. Indies	1797.
167 <i>aromatica</i> Sal.	aromatic	2 jnj.l	Y	E. Indies	1804.	
168 <i>Idtifa</i> Rose.	broad-leaved	or	12 my.jn	Y	E. Indies	1880.
169 <i>Zcr&nbet Rox.</i>	Zerumbet	E or	3 ap.au	Y	E. Indies	1807.
170 <i>aruginosa</i> Rox.	rusty	E or	5 ap.au	RY	E. Indies	1807.
171 <i>rubescens</i> Rox.	rubescent	E or	3 my.s	Y	E. Indies	1805.
172 <i>cassia</i> Rox.	grey	E or	1 apjn	Y	Bengal	1819.
173 <i>comosa</i> Rox.	<i>tufcid-flowcred</i>	E or	2 my	RY	R Indies	1819.
174 <i>clata</i> Rox.	tall	E or	3 my	C	E. Indies	1819.
175 <i>ferruginea</i> Rox.	ferruginous	E or	4 my	Y	E. Indies	1819.
176 <i>leucorhiza</i> Rox.	white-rooted	E or	5 my	RY	E. Indies	1819.
177 <i>xanthorrhiza</i> Rox.	yellow-rooted	E or	6 my	R	Amboyna	1819.
178 <i>rubricaulis</i> Lk.	red-btemmed	E or	7 my	R	E. Indies	1822.
179 <i>angustifolia</i> Rox.	narrow-leaved	E or	8 my	Y	E. Indies	1822.
180 <i>vindiflora</i> Rox.	green-flowered	E or	9 my	Y.o	Sumatra	1822.
181 <i>petiolata</i> Rox.	petiolated	E or	10 my	B	Pegu	1822.
182 <i>Amada</i> Rox.	<i>Ama.da.ginger</i>	E or	11 my	RY	Bengal	1819.
183 <i>long-rooted</i>	long-rooted	E or	12 ail	B	E. Indies	1759.
184 <i>montana</i> Rox.	mountain	E or	13 my.jn	R.w	E. Indies	1824.
185 <i>reclinata</i> Rox.	rechnate	E or	14 ap.my	Pk	E. Indies	1824.
15. GLO'BA Rose.	GLOBRA.	(SO named in the Moluccas.)	Seitaminea.	8. — 9.		
186 <i>marantina</i> Rose.	maranta-like	4 (E or	1* jlau	Y	E. Indies	1800.
187 <i>bulbifera</i> Rox.	bulbiferous	E or	2 jlau	Y	E. Indies	1829.
188 <i>onx-snsis</i> Rox.	Orixian	E or	2 jlau	Pk	E. Indies	1819.
189 <i>Careyana</i> Hox.	Dr. Carey's	E or	3 jlau	Y	Pegu	1822.
190 <i>orecta</i> Dec.	upright	E or	4 jlau	W	E. Indies	1820.
191 <i>racmosa</i> Sm.	racemose	E or	5 jlau	Y	Ceylon	1812.
<i>libra</i> Rox.		E or	6 jlau	Y		
192 <i>pendula</i> Rox.		E or	7 jlau	Y		
193 <i>sessiliflra</i> B. M.		E or	8 jlau	Y		
		E or	9 jlau	Y		
		E or	10 jlau	Y		
		E or	11 jlau	Y		
		E or	12 jlau	Y		
		E or	13 jlau	Y		
		E or	14 jlau	Y		
		E or	15 jlau	Y		
		E or	16 jlau	Y		
		E or	17 jlau	Y		
		E or	18 jlau	Y		
		E or	19 jlau	Y		
		E or	20 jlau	Y		
		E or	21 jlau	Y		
		E or	22 jlau	Y		
		E or	23 jlau	Y		
		E or	24 jlau	Y		
		E or	25 jlau	Y		
		E or	26 jlau	Y		
		E or	27 jlau	Y		
		E or	28 jlau	Y		
		E or	29 jlau	Y		
		E or	30 jlau	Y		
		E or	31 jlau	Y		
		E or	32 jlau	Y		
		E or	33 jlau	Y		
		E or	34 jlau	Y		
		E or	35 jlau	Y		
		E or	36 jlau	Y		
		E or	37 jlau	Y		
		E or	38 jlau	Y		
		E or	39 jlau	Y		
		E or	40 jlau	Y		
		E or	41 jlau	Y		
		E or	42 jlau	Y		
		E or	43 jlau	Y		
		E or	44 jlau	Y		
		E or	45 jlau	Y		
		E or	46 jlau	Y		
		E or	47 jlau	Y		
		E or	48 jlau	Y		
		E or	49 jlau	Y		
		E or	50 jlau	Y		
		E or	51 jlau	Y		
		E or	52 jlau	Y		
		E or	53 jlau	Y		
		E or	54 jlau	Y		
		E or	55 jlau	Y		
		E or	56 jlau	Y		
		E or	57 jlau	Y		
		E or	58 jlau	Y		
		E or	59 jlau	Y		
		E or	60 jlau	Y		
		E or	61 jlau	Y		
		E or	62 jlau	Y		
		E or	63 jlau	Y		
		E or	64 jlau	Y		
		E or	65 jlau	Y		
		E or	66 jlau	Y		
		E or	67 jlau	Y		
		E or	68 jlau	Y		
		E or	69 jlau	Y		
		E or	70 jlau	Y		
		E or	71 jlau	Y		
		E or	72 jlau	Y		
		E or	73 jlau	Y		
		E or	74 jlau	Y		
		E or	75 jlau	Y		
		E or	76 jlau	Y		
		E or	77 jlau	Y		
		E or	78 jlau	Y		
		E or	79 jlau	Y		
		E or	80 jlau	Y		
		E or	81 jlau	Y		
		E or	82 jlau	Y		
		E or	83 jlau	Y		
		E or	84 jlau	Y		
		E or	85 jlau	Y		
		E or	86 jlau	Y		
		E or	87 jlau	Y		
		E or	88 jlau	Y		
		E or	89 jlau	Y		
		E or	90 jlau	Y		
		E or	91 jlau	Y		
		E or	92 jlau	Y		
		E or	93 jlau	Y		
		E or	94 jlau	Y		
		E or	95 jlau	Y		
		E or	96 jlau	Y		
		E or	97 jlau	Y		
		E or	98 jlau	Y		
		E or	99 jlau	Y		
		E or	100 jlau	Y		
		E or	101 jlau	Y		
		E or	102 jlau	Y		
		E or	103 jlau	Y		
		E or	104 jlau	Y		
		E or	105 jlau	Y		
		E or	106 jlau	Y		
		E or	107 jlau	Y		
		E or	108 jlau	Y		
		E or	109 jlau	Y		
		E or	110 jlau	Y		
		E or	111 jlau	Y		
		E or	112 jlau	Y		
		E or	113 jlau	Y		
		E or	114 jlau	Y		
		E or	115 jlau	Y		
		E or	116 jlau	Y		
		E or	117 jlau	Y		
		E or	118 jlau	Y		
		E or	119 jlau	Y		
		E or	120 jlau	Y		
		E or	121 jlau	Y		
		E or	122 jlau	Y		
		E or	123 jlau	Y		
		E or	124 jlau	Y		
		E or	125 jlau	Y		
		E or	126 jlau	Y		
		E or	127 jlau	Y		
		E or	128 jlau	Y		
		E or	129 jlau	Y		
		E or	130 jlau	Y		
		E or	131 jlau	Y		
		E or	132 jlau	Y		
		E or	133 jlau	Y		
		E or	134 jlau	Y		
		E or	135 jlau	Y		
		E or	136 jlau	Y		
		E or	137 jlau	Y		
		E or	138 jlau	Y		
		E or	139 jlau	Y		
		E or	140 jlau	Y		
		E or	141 jlau	Y		
		E or	142 jlau	Y		
		E or	143 jlau	Y		
		E or	144 jlau	Y		
		E or	145 jlau	Y		
		E or	146 jlau	Y		
		E or	147 jlau	Y		
		E or	148 jlau	Y		
		E or	149 jlau	Y		
		E or	150 jlau	Y		
		E or	151 jlau	Y		
		E or	152 jlau	Y		
		E or	153 jlau	Y		
		E or	154 jlau	Y		
		E or	155 jlau	Y		
		E or	156 jlau	Y		
		E or	157 jlau	Y		
		E or	158 jlau	Y		

MONANDRIA DIGYNIA.

CLASS I. ORDER II.

19. BOERHAA'V/JI L.	HOQWBED.	(The celebrated Dutch physician, d. 1758.)	<i>Njctagine*</i> .	9.-36.
S04 erecta X.	upright	£ (23 or Ifjl.s	W India	1733. S co Jac. vin. 1.5,6
205diffusa W.	spreading	£. 123 or 1 au.8	C India	1690. S co
206 procumbens Roxb.	procumbent	* 123 or 1 jl.n	R E. Indies	1800. ■ co R.mal.7.56
207 hirsCita X.	hairy	JU • or 1 my.au	S Jamaica	1733. ■ co Jac. vin. 1.1
208 plumbaginca Cav.	Plumbago-like	£ (23 or \ my.au	R Spain	1820. ■ co Cav. ic. 2.112
209 discolor Kth.	two-colored	£ (23 or 1* s	R Guayaquil	1825. S co
S10 viscosa Lag.	clammy	£. (23 or 3 ap.8	S Peru	1821. C co
211 scandens X.	climbing	CD or 6 ap.8	G Jamaica	1691. S co Jac. vin. 1.4
212 cxcHsa W.	tall	nHor 5 ap.n	R S. Amer.	1820. C co
20. CENTRANTHUSXVfc	CRNTBANTHUS.	(Kenton, a spur, an/hos, a flower)	<i>ValeriiLnece</i> . 3.-4.	
213 rdber Dec.	red	J A or H my.jl	C Britain	mea. R co Eng. bot 1531
2fl6reairo	white-flowering	J A or 11 my.au	W Britain	gard. D co
214 angustifolius Dee,	narrow-leaved	t A or U my.jl	C & Europe	1759. R co FLgr.29
215 Calcitrapa Dtftr.	caltrop-kaved	O or 1 my.jl	P Portugal	1683. S co Fl. gr. 50
* 21. VOCHY'IA Poir.	VOCHTSIA.	(The original name of the Caribes.)	<i>Vochysiee</i> .	1.—15.
216 excelsa Lam.	lofty	I CD or 30 au.s	Y Trinidad	1825. C s.p Aub.gui.18.6
22. POLLICHLI W.	POLLICBIA.	IJ. A. Pollich, a German botanist	<i>Chenopbde*</i> .	1.
217 campestris W.	field	OJ or 6 8	Ap C. B. S.	1780. C co Sm. spic. 1.1
23. SALICCRNIA X.	GLABSWORT.	(Sal, salt, cornu, a horn; salt-wort)	<i>Chenophdete</i> .	6.—18.
218ai*bica W.	Arabian	#i Jor 1 jnj.l	Ap Arabia	1758. C s.l M.h.2.33.7
219 fruticosa L.	shrubby	J ■ 1 au.8	Ap Britain	sea sh. S 8 Eng. bot. 2467
220 perennans IV.	perennial	J ■ \ au.8	Ap Siberia	1823. S s Pal it 1.1. 1
221 radicans E. R	rooting	J ■ \ 1 8	Ap Britain	sea sh. S s Eng. bot. 1691
222 herbacea L.	herbaceous	O clt 1 au.8	Ap Britain	sea sh. S s Eng. bot 415
223 procumbens E. R	procumbent	JU O clt \ au	Ap England	sal. m. S 1 Eng. bot 1691
24 vulgaria L.	MARETAIL.	(Hippos, a mare, oura, a tan.)	<i>HippwM**</i> .	1.-2.
	common	J "A " 1 my.jn	Ap Britain	dit R co Eng. bot 763
25. ZOSTE'RA L.	WBACKORAA s.	(Zoster, a xiband; referring to the leaves.)	<i>Fluviales</i> .	1.
225 marina X.	marine	* A " au.8	Ap Britain	sea sh. S ■ Eng. bot 467
26. CHLORA'NTHUS Sun.	CHLOBANTHUS.	(Odoros, green, onthos, a flower.)	<i>Chlor&nthetae</i> .	3.-4.
226 inconspicuus Sun.	obscure	I L R C U 1 ap.s	Ap China	1781. C co
227 monfetachys Lindl.	one-spiked	jg E2 cu 1 fmy	Ap China	1819. C co land. coL 17
228 monander R. Br.	one-anthered	t L p c u Ijjn	Ap China	1817. C co

CLASS I. ORDER 2.

MONA'NDRIA DIGY'NIA.

27. CORISPE'RMUM X.	TICKSEBD.	(Kori, a bug or tick, sperma, a seed.)	<i>Chawpbdea</i> .	11.—16.
229 Aywopifolium X.	hyssop-leaved	O un li jl	Ap Euroiie	1739. S co FLgr.1.1
230 squarrifolium X.	Tough-spiked	O un 1 au.s	Ap Russia	1759. S co Pal. ros. 2.99
231 RedowakV Fis.	Redowsky's	O un fjl.au	Ap Siberia	1822. S co
232 intermedium Schw.	intermediate	O un \ jlau	Ap Poland	1822. S co
£33p6ngens Vahl	pungent	O un ijl.au	Ap Siberia	1818. S co Bun. c 3.56
234 uitidum R.&S.	clear	O un 1 jlau	A p Hungary	1820. S co
235 cartescens w.S/JK.	canescent	O un 1 jlau	Ap Hungary	1824. S co
236 patens &&	spreading	O un 1 jlau	Ap	1824. S co
237 tenué Lk.	slender	O un 1 jlau	Ap	1825. S co
238 glomeratum Lag.	glomerated	O un 1 jlau	Ap	1824. S co
239 Manchalinii Stev.	Marschall's	O un 1 jlau	Ap Caucasus	1820. S co
*28. CALLITRICHE W.	WATER STARWOBT.	(Kallas, beauty, thrix, hair.)	<i>Hakragete</i> .	2.
240 verna X.	spring	£ o w \ ap.o	W Britain	dit. S aq Eng. bot 722
aquatica E. R				
2 aquatica E. FL	water	£ O w iap.o	W Britain	dit. S aq CoL ec. 1.316
41 auiumnalis X.	autumnal	* O w \ jn.s	W Britain	dit S aq Eng. bot 722.2
aquatica y E.R				
29. SLITUM X.	STRAWBERRY BUTE.	(Bliton, insipid, Gr. and Celt.)	<i>Chenopbdear</i> .	5.
242 capitiflum X.	berry-headed	O or 2 my.au	Ap Austria	*1633. S ru Pi&T.fl. 1.2
243 virgositum X.	twiggy	O or 2 my.s	Ap S. Europe	1680. S ru Bot mag. 276
244 chenopodioides Lam.	chenopod..like	O un 21 my.au	Ap Crimea	1797. S co M.h. i32.11
245 petiolare X*.	petiolate	O un ljjl.au	Ap	1824. S co
246 maritimum NuL	sea-side	O cu 1 jn.au	Ap N. Amer.	1825. S co
SO. MNIA'RUM Font.	MNIARUM.	(Mttiaros, mossy.)	<i>Chenopbdecc</i> .	1—2.
247 biflrum Font.	two-flowered	£ [AJ cu i jn.jl	W N. Zeal.	1823. S s.p For. co. goet 1.1
3L ASPICA'RPA Rich.	ASPICABPA.	(Aspis, a shield, karpas, a fruit)	<i>Malpighidcem</i> .	1.
£48 urens Rich.	stinging	J ■ 5J jn.jl	Ap & Amer.	1821. C co Mem. m. 2. IS
32. MONE'RMA R.Br.	MONEBMA.	(Monos, one, herma, a support j bne glume.)	<i>Graminae</i> .	1.
249 monandrum R. Br.	monandrous	m ^ cu I jl	Ap Spain	1804. S co Cav. ic. 39.1
Psillifrus nardoidea Trin.	Rottboellia monfindra Cav.			
33. CI'NNA X.	CINNA.	(Ancient name of a grass growing in Cilicia)	<i>Grammfar</i> .	1. — 2.
250 arundinacea X.	reedy	Jk A cu 3 jn.s	Ap Canada	1799. S co Schreb.gr. 49

CLASS II. ORDER 1.

DIA'NDRIA MONOGY'NIA.

298	glaucum	<i>H. K.</i>	glaucous	I	3	au	W	C.G.H.	1774.	C	l.p	Sal. st. 8			
299	acandens	<i>Vahl</i>	climbing	I	10	jl	W	£. Indies	1820.	C	co				
300	bracteatum	<i>Box.</i>	large- <i>h</i> Tacted	I	30	ap.my	W	EL Indies	1818.	C	81				
301	latifolium	<i>Rox.</i>	broad-leaved	I	20	Jnjl	W	E. Indies	1817.	C	81				
302	arborcscens	<i>Box.</i>	arborecent	I	12	...	W	E. Indies	1824.	C	r.m				
303	trinfrve	<i>Vahl</i>	three-nerved	I	20	...	W	E. Indies	1804.	C	l.p				
SOI	simplicifolium	<i>Fortt.</i>	simple-leaved	I	3	jnjl	W	S. Seas	1800.	C	r.m	Bot. mag. 980			
305	gracile	<i>R. Br.</i>	slender	I	3	year	W	Norfolk	1.1791.	C	s.p	Hot. rep. 127			
JGJ	elongat	<i>tum L</i>	elongated	I	10	ap.my	W	E. Indies	1820.	C	r.m	Ber ph. tr. 1772.			
Sff7	acumin&tum	<i>R. Br.</i>	acuminate	I	10	jn.au	W	N. Holl	1820.	C	l.p				
308	paniculatum	<i>Box.</i>	pnncled	I	5	ja	W	China	1818.	C	r.m	Bot. cab. 469			
309	undulatum	<i>Vahl</i>	wavy	I	5	ja	W	China	1819.	C	r.m	Bot. reg. 4-ii			
310	auriculatum	<i>Vahl</i>	auriculated	I	10	my.s	W	E. Indies	1790.	C	r.m	Bot. reg. L'ltt			
SII	campanulatum	<i>Lk.</i>	campanulate	I	4	...	W	...	1822.	C	r.m				
312	flexile	<i>Vahl</i>	flexible	I	10	ap.my	W	E. Indies	1825.	C	r.m				
313	capense	<i>fhun.</i>	Cape	I	8	apjl	W	C.G. H.	1816.	C	r.m				
314	wylamcuci	<i>Ma. C.</i>	Ceylon	I	6	jl.au	W	Ceylon	1823.	C	r.m				
315	tortu&sa;ura	<i>W.</i>	twisted	I	6	my.jl	W	C.G.H.	1818.	C	r.m	Jac. sc. 4.490			
		flxile Jac.													
316	dispensum	<i>Wa.</i>	two-seeded'	I	li	lor	8	...	W	Nepal	1825.	C	l.p		
317	divaricat&tum	<i>A. Zhd.</i>	divaricate	I	i	or	10	ap.jl	W	N. Holl.	1819.	C	r.m	Bot reg. 12W	
318	heterophllum/for.		variable-leaved	I	»	J	or	14	jnjl	W	Nepal	1820.	C	r.m	
319	azoricum	<i>L.</i>	Azorian	I	I	J	or	5	ap.n	Y	Madeira	1724.	C	r.m	Bot. reg. 89
320	frfiticans	<i>L.</i>	sprouty	S	or	3	apo	Y	S. Europe	1570.	C	r.m	Bot mag. 461		
321	humile	<i>L.</i>	low	I	k	or	3	jns.s	Y	S. Europe	1656.	L	co	Bot reg. 350	
322	odoratissimum	<i>L.</i>	sweetest-scent.	J	t	J	or	3	my.n	W	Madeira	1656.	C	r.m	Bot mag. 285
323	revolutum	<i>B. R.</i>	revolute-/füMi	£	i	J	or	12	mr.o	Y	E. Indies	1812.	C	r.m	Bot reg. 178
324	officinalc	<i>L.</i>	officinal	&	or	15	jno.	W	E. Indies	1548.	C	co	Bot mag. 31		
•	2 fol.	argenteis	silver-leaved	A	or	15	jno.	W	E. Indies	...	C	co			
3	fol.	aftreis	golden-leaved	T	or	15	jno.	W	E. Indies	...	C	co			
4	flribus	plenis	double-flowered^	or	15	jno.	W	E. Indies	...	C	co				
325	grandiflorum	<i>L.</i>	lurge-flowered	£.	i	or	15	jno.	W	E. Indies	1629.	C	r.m	Bot reg. 91	
326	stenopetalum	<i>Lindl.</i>	narrow-petalcd	«	CD	or	+	■	W	E. Indies	1824.	C	co		

44. LER'TA'NDRA Nut. LEPTANDRA. (*Leptos*, slender, *aner*, an anther.) *Scrophulariaceae*. 2.—

•45. VERO'NICA L. SPEEDWELL. (Name of a princess, or altered from Betonica.) *Scrophyllutirue*. 134. — 170

		A or	3 ja.	B	Hungary	1805.	D e o	W. & K. 2.102
329 foliosa Kit.	leafy	^ A or	3 ja.	B	S. Europe	1814.	D co	Hof. ph. E. 3
330 crenulata Vahl	notch-flowered	II	II	B	Sweden	1570.	D co	Sc. v. 29.1.1
331 maritima L.	marine	II	II	B	Germany	1819.	D co	
332 H6stiiR. &a	Host's	II	II	B	Persia	1821.	D co	
333 Stephanidna R. & S.	Stephan's	II	II	B	Siberia	1822.	D co	
334 angustifolia Fit.	narrow-leaved	H	H	B	Siberia	1731.	D co	Gme1.it 1.39
335 spiria Schr.	spurious	H	H	B	Russia	1797.	D co	
336 paniculata Pall.	paiified	H	H	B	S. Europe	1812.	D s.l	Hof. ph. E. 4
337 complicata ligfg.	complicate-/-^	H	H	B	1821.	D co	
338 azifira Schott	sky-blue	H	H	B	1821.	D co	
339 polystachya Lk.	many-spiked	H	H	B	S. Europe	1804.	D co	Sc.v.25.1.4
340 glabra Schr.	smooth	H	H	B	Siberia	1817.	D co	
2 alba	white	H	H	B	S. Europe	1812.	D co	
341 leucantha Fis.	white-flowered	H	H	B	S. Europe	1808.	D co	
342 amethystina W. en.	amethyst	H	H	B	1822.	D co	
343 elatior Ehrh.	taller	H	H	B	S. Europe	1812.	D co	Sev.22.2.2
344 acuta Mart.	acute	H	H	B	1820.	D co	
345 argita Schr.	sharp-notched	H	H	B	Europe	1817.	D co	
346 melancolica Desf.	melancholy	H	H	B	1817.	D co	
347 nltida Lk.	clear	H	H	B	1818.	D co	
348 longibracteata Lk.	long-bracted	H	H	B	Germany	IH04.	D co	Sc.v.23.1.2
2 latifolia	broad-leaved	H	H	B	1823.	D co	
349 media Schr.	mediate	H	H	B	1817.	D co	
350 periclitella Schott	peach-leaved	H	H	B	S. Europe	1812.	D co	Sc.v.24.2.3
351 pradita H. C.	very high	H	H	B	S. Europe	1731.	D co	Sc.v.24.2.1
353 austriaca Schr.	southern	H	H	B	—	—	D co	
353 longifolia L.	long-leaved	H	H	B	Crimea	1821.	D co	
2 incarnata	flesh-colored	H	H	B	Sweden	1823.	D co	
3 alba	white	H	H	B	—	1820.	D co	
354 grossa Mart.	gross	H	H	B	Siberia	1797.	D co	W.&K.3.244
355 ambigua Mart.	doubtful	H	H	B	Russia	1759.	D co	Hof. got 15.6
356 falcata Mart.	falcate-leaved	H	H	B	Siberia	1826.	D co	
357 neglecta Vahl	neglected	H	H	B	Greenland	1819.	D co	
358 inclata L.	hoary	H	H	B	1823.	D co	
359 grindia Fis.	grand	H	H	B	S. France	1822.	D co	
360 Wormskioldia R. & S.	Wormskiold's	H	H	B	1822.	D co	Bot. mag. 2975
361 rlgens Mart.	stiff*	H	H	B	England	ch. pa.	D co	
382 elegans Dee.	elegant	H	H	B	—	1800.	D co	
363 brevifolia Lk.	short-leaved	H	H	B	Hungary	1822.	D co	Eng. bot 2
364 spicata L.	spiked	H	H	B	1817.	D co	
365 incarnata G. Don	small flesh-cld	H	H	B	1819.	O co	
366 Cltis' Schott	Elcuse's	H	H	B	Austria	1823.	D co	
367 Waldsteinia Schott	Waldstein's	H	H	B	S. Europe	1833.	D co	
368 confusa R. & S.	confused	H	H	B	Europe	1819.	I) co	Bot. mag. 2210
369 menthaefolia Schott	mint-leaved	H	H	B	1824.	D co	
370 Barrelieria Schott	Barrelier's	H	H	B	1817.	D co	
371 orchidea Crz.	orchis-flowered"	H	H	B	England	moun.	D co	Eng. bot. 673
372 longitora R. & S.	long-rowered	H	H	B	Europe	1622.	D co	
S73 corymbosa Hort.	corymbose	H	H	B	Russia	1821.	D co	
374 h^brida L.	hybrid	H	H	B	Pyrenees	1822.	D co	
375 cassifolia Kit.	thick-leaved	H	H	B	Gou. il 1.1			
376 ruthenica Joe.	Russian	H	H	B				
977 F6na? Gou.	Pona's	A or	3 ja.	B				

378 <i>villösa</i> Schr.	villoua	^ A or	U	■ B	& Europe	1804.	D co	Sc.v.31.1 3
379 <i>pinnata</i> L.	pinnate-tawed	k A or	1	jn.au	Siberia	1775.	D co	Hof.got. 15.10
380 <i>incisa</i> // K.	<i>cut-leaved</i>	k A or	2	jn.au	Siberia	1779.	D co	JJotcab.UD7
381 <i>lacu&ta</i> H. K.	jagged leaved	ifc A or	■	jn.au	Siberia	1780.	D co	Jun. ic. 2
382 <i>parviflora</i> Vahl	small-flowered	« i or	■	jnjl	New Zeal.	1822.	C s.p	
383 <i>gentianina</i> /fe* Vahl	gentian-like	* A or	■	myjn	D.B	Levant	1748.	D co
384 <i>gallida</i> Horn.	Dale	A or	1	iny.jn	B	Tauria	1821.	D co
385 <i>ellidioides</i> L.	dawy-like	i A or	■])	B	SwitzcrL	1775.	D co
386 <i>fruticulosa</i> L.	shrub-like-rtAd	a. or	■	jn.au	Scotland	sc. alp.	D co	Hal. h. 15.1
387 <i>saxatilis</i> L.	rock	* A or	■])	Scotland	sc. alp.	D co	Eng. bot 1028
388 <i>nummularia</i> GOM	moneys wort-lv	Jc A or	..])	B	Pyrenees	1820.	D co
389 <i>alpina</i> L.	alpine	Jc A or	..])	B	Europe	...	D co
2 <i>Integerrifolia</i> R.S/S.	entire-leaved	* A or	ft	my.jn	B	Silesia	1814.	D co
3 <i>pumila</i> All.	dwarf	Jc A or	ft	my	B	Piedmont	1819.	D s.p
4 <i>obtusifolia</i> R. 8/S.	obtuse-leaved	Jc A or	ft	my	B	Scotland moun.	D s.p	Al пед. 22.1
5 <i>heterophylla</i> /& \$.	variable-leaved*	A or	ft	my	B	Europe	...	Eng. bot 484
6 <i>rotundifolia</i> R. \$S.	round-leaved	J: A or	ft,	my.	B	Europe	1816.	D s.p
390 <i>depauperata</i> Kit.	impoverished	k A or	ft,	ap.jl	B	Hungary	1823.	D s.p
391 <i>scrupulifolia</i> L.	Serpulhum-lvd	5 ^ or	ft	ap.jl	B	Britain me. pa.	D co	W.&K.3.245
2 tenella ScAm.	tender	Jc A or	i	ap.jl	B	Europe	—	Eng. bot 1075
3 neglecta ScAm.	neglected	Jc A or	ft	ap.jn	B	France	1820.	D co
4 quaternata	quaternate	* A or	J	my.jn	B	Crete	1819.	D co
5 <i>nuiMularifolia</i>	moneyswort-lvd	Jc A or	ft	my.jn	B	Hungary	1822.	D s.p
6 humifusa	humifuse	Jc A or	ft	my.jn	B	Falkl. I.	1776.	C r.m
392 <i>tenella</i> All.	tender	Jc A or	ft	my.jn	B	Italy	1775.	D co
393 <i>cretica</i> Pall.	Cretan	Δ-Δ pr	ft	my.jn	B	Britain rivul.	D co	Seg.ver.1.3.2
3H <i>microphylla</i> Kit.	small-leaved	Δ ^ or	ft	my.jn	B	Carolina	1821.	D co
395 <i>decu8sdta</i> //.IC.	<i>cxosi-leuvcd</i>	» J or	ft	my.jn	B	Britain mar.	D co	Eng. bot 655
396 <i>aphl ill</i> L.	<i>leaAenstalled</i>	EL A or	ft	my.jn	B	mar.	D co	
397 <i>Bccab&ng</i> L.	Brook-lime	£ A clt	2	myjn	B	France	1820.	D co
398 <i>carol imana</i> Pair.	Carolina	& A or	1	jnjl	B	England	—	AI пед. 22.1
399 <i>fnaKalhs</i> L.	Pimpernel-ftfc	£ A or	ft	my.jn	B	Europe	—	
400 <i>Bcutellata</i> L.	saucer-leaved	£ A or	2	jl	F	France	1820.	D co
401 <i>parmula</i> 7Vo/.	small-shielded	Δ A *r	2	jn.au	R	Crete	1819.	D co
402 <i>onentalis</i> //. /C	eastern	Jc A or	1	jlau	L.B	& Europe	1824.	D co
403 <i>tai'incs</i> W.	Taunan	Jc A pr	1	jlau	B	Levant	1748.	D co
404 <i>tenuifolia</i> Sfw.	fine-leaved	Jc B	1	jlau	B	Siberia	1820.	R LI
405 <i>trichodrpa</i> R. \$S.	hairy-capsuled	Jc B	1	jlau	B	Iberia	1821.	D co
	piloz irpa Lk	Jc B	1	jlau	B	Levant	182a	D 8.1
406 <i>caucasica</i> liicb.	Caucasian	Jc B	ft	jnjl	Pa. R	Caucasus	1816.	D co
2 <i>latifolia</i> G. Don	broad-leaved	Jc B	I	jrijl	Pa. R	Caucasus	1820.	D co
407 <i>Jacquinii</i> Schott	Jacquin's	Jc B	I	my.jn	B	Austria	1748.	D co
408 <i>pectinata</i> Vahl	pectinated	Jc B	I	myjn	B	Italy	1819.	D co
409 <i>Schmidti</i> *R.&S.	Schmidt's	Jc B	I	myjn	B	Bohemia	1820.	D co
410 <i>australa</i> Jac.	Austrian	Jc B	I	myjn	B	Austria	1748.	D co
411 <i>multifida</i> L.	much-cut	Jc B	ft	my jl	L.B	Siberia	1748.	D co
412 <i>pmnatiida</i> W.	pinnatifid	Jc B	1	myjn	B	Europe	1817.	D eo
413 <i>Alli6n</i> * Vil.	Alliom's	Jc B	ft	jn.au	L.B	& Europe	1748.	1) co
414 officin^lis L	officinal	Jc B	4	apjl	B	Britain bar.gr.	D co	Al. ped. 1. 46. i
415 <i>Tournefortii</i> Vil	Tournefort's	Jc B	ft	myjn	B	France	1821.	D co
416 <i>scltgera</i> D. Don	bristled	Jc B	ft	apjl	B	Scotland raed.	D co	Eng. bot 765
	hirsita Hopk.	Jc B						
417 <i>MullenVina</i> R.&S.	Muller'a	Jc B	1	jnjl	■	Syria	1825.	D B.p
418 <i>Chaixa</i> Lap.	Chaixa's	Jc B	ft	jnjl	■	S. Europe	1825.	D f1. I
419 <i>prostrata</i> L.	trailing	Jc B	1	myjn	■	Germany	1774.	D co
420 <i>plicata</i> Pohl	plaited	Jc B	2	myjn	■	Bohemia	1817.	D co
421 <i>dentSta</i> Schr.		Jc B	1	myjn	■	Europe	1818.	D co
422 <i>micrantha</i> Hofg.	small-flowered	Jt A or	ft	myjn	W	Portugal	1822.	D co
423 <i>Jamifolia</i> Hayne	dead-nettle-lvd	^ A or	ft	myjn	W	...	1825.	D co
424 <i>melissifolia</i> Pair.	balm-leaved	ji A or	1	myjn	W	...	1826.	D co
425 <i>latifolia</i> Z.	broad-leaved	ji A or	1	myjn	W	Austria	1748.	D co
426 <i>Tedcum</i> L.	Germander-/vd	^ A or	2	jn.au	L.B	Germany	1596.	D co
427 <i>polymorpha</i> W. en.	multiform	^t A or	1	jn.au	B	...	1817.	D co
428 <i>pilos</i> L.	pilos	^ A or	1	myjn	B	Bohemia	1819.	D co
429 <i>acutiflora</i> Lap.	acute-flowered	ji A or	1	myjn	B	France	1821.	D co
430 <i>crinita</i> Kit.	hairy	ji A or	1	jlau	B	Hungary	1822.	D
431 <i>Chamaedrys</i> L.	Germander	ji A or	ijl.ju	H	Britain m.pas.	D co	Eng. bot. 623	
432 <i>triticxfoha</i> L.	Circea-like	ji A or	I jnjl	L.B	Austria	1776.	D co	
433 <i>circvoVrs</i> G. Don	nettle-leaved	ji A or	ft myjn	11	Switzerl.	1820.	D eo	
434 <i>monfana</i> L.	mountain	ji A or	ft myjn	B	Britain moi.w.	D co	Jac.au. 1.51	
435 <i>peduncularis</i> Bicb.	pedunculated	ji A or	myjn	B	Caucasus	1826.	D co	
436 <i>petra/a</i> »?»		ji A or	myjn	B	Caucasus	1821.	D co	
437 <i>pcrfhota</i> R. Br.	rock	ji A or	jl.au	B	N. S. W.	1815.	D r.m	
438 <i>labiata</i> R. Br.	perfoliate	ji A or	apjl	L.B	N. Holl.	1802.	C r.m	
439 <i>plebäa</i> R. Br.	labiated	ji A or	jjnl	B	N. S. W.	1820.	D co	
«0 <i>sracilis</i> /I. Br.	slender	ji A or	myjn	B	N. S. W.	1825.	D co	
Mildistans R.Br.	distant	ji A or	myjn	B	N. S. W.	1823.	D Lp	
%4+2p61hra G.Don	fair	ji A or	f jn.au	B	Caucasus	1824.	S co	
443 <i>maxima</i> 6Yev.	largest	ji A or	1	myjn	B	Britain san.fi.	S	Bux. c. 1.34
444 <i>v6rna</i> L.	vernal	ji A or	ft ap.my	B	Britain	1805.	S co	
445 <i>digitata</i> TaA/	digitated	ji A or	ft jl	B	S. Europe	1805.	S co	
446 <i>triplylos</i> L.	three-leaved	ji A or	ft ap.my	B	Britain	san.fi.	Eng. bot 26	
447 <i>Adera>falia</i> 7M	ivy-leaved	ji A or	1	inr.jn	B	Britain	clt.gr.	Eng. bot 784
448 <i>ymbalaria</i> -/vd	Jc O or	2	ap.my	W	S. Europe	1821.	S co	
449 <i>percgrina</i> L.	outlandish	O cu	ft mr jn	W	N. Europe	1680.	S co	
450 <i>filiformis</i> 5w.	filiform	Jc O or	ft my	L.B	Levant	1780.	S co	
451 prs'cox All.	early	O or	ft txnr	B	& Europe	1775.	S co	
452 <i>acinifolia</i> L.	acinus-leaved	O or	O pr	B	S. Europe	1788.	S co	
453 <i>arvensis</i> L.	corn field	Jc O w	ft ap.my	B	Britain	oldw.	P. & T.fl.L.S3	
454 <i>agrestis</i> L.	rural	Jc O w	nirjl	B	Britain	cltgr.	Eng. bot 734	
2 versicolor	various-colored	O w	ft mrjl	B	Britain	fields.	Eng. bot 783	
3opáca	opaque	O w	ft mrjl	B	Britain	fields.	S co	
455 amccna Stev.	pleasant	O pr	ft ap.ray	R	Iberia	1824.	S co	
456 bUoba ToA/	two-lobcd	O pr	ft ap.my	B	Iberia	1819.	S co	
457 pulchUa Bernh.	pretty	O pr	ft ap.my	W	—	1819.	S co	

DIANDRIA MONOGYNYA.

CLASS 11*

458 pfrstea Pohr.	Persian	Opr	I ap.my	B	Persia	1818.	S co
459 Buxnami Ten.	Buxbaum's	O pr	i apjn	W	S. Europe	1800.	S co Fl. nap. 7.1
460 wrort/c^forme/^A.veronica-formed-*	DIPLOPHYLLUM Leh.	DIPLOPHYLLUM. (<i>Diploos</i> , double, <i>phyton</i> , a leaf.)	<i>Scrophdaniue</i> . 1.				
Veronica CYsta galli Stev.							
47. MICRA'NTHEMUM Mx.	MICRANTHEMUM. (<i>Mikros</i> , small, <i>anthos</i> , a flower.)	<i>Prtmulacece</i> . 1.—2.					
48 orbiculatum itf*.	round-taiwwf	JL AJ pr	} niy.jn	W	Carolina	1826.	D s.p
• 48. GALIPEA* Aub.	GALIPEA. (The name in Guiana.)	<i>Rutacece</i> . 1.					
4G2 trifoliata Aub.	three-leaved	• o r 4?	..	G	Guiana	1803.	■ p.l Aub. gui. 269
49. SCHWE'NCKM L.	SCHWE'NCKIA. (J. T. Schvoenck, Prof. Med. at Jena.)	<i>Scrophuldrince</i> . 1. — 7.					
463 americana L.	American	£ 33	1 au.s	Li	Guiana	1781.	C s.p Sch.bs.328.1
60. HEMI'MERIS L.	HEMIMERIB (Hemni, half, merit, a p:irtj a half flower.)	<i>Scrophuldrive</i> . 1.					
434 montana /..	mountain	■ ■ ■ -	- jn.au	-	C. G. H.	1816.	C p.l Lam. il. 538
51. HO'STvi Jac. .	HOSTA.	(N. T. Host, a German botanist.)					
465 cierdlea Jac.	blue	* Q or 6	jn.au	B	Mexico	1733.	C p.l Bot mag. 2611
Curn'itia punct&ta W.							
466 latifolia Hum.	broad-leaved	■ or 6	jn.au	B	Mexico	1824.	■ p.l
467 longifolia Hum.	long-leaved	■ or 6	jn.au	B	Mexico	1826.	■ ■
52. CORNU'TJ^ L.	COBNUTIA. (<i>J. Cornutus</i> , a physician of Paris.)	<i>Verbendceee</i> . 1.					
468 pyramidata L.	pyramidal	• o r 6	jl	B	Mexico	1733.	C p.l Lam.il. 541
53. GRATI'OLA L.	HEDGE HVSSOP. (<i>Gratia</i> , grace of God; as a purge.)	<i>Scrophuldriture</i> . 5.—35.					
469 officinalis L.	officinal	■ my.au	L.B	Europe	1568.	D co Fl. dan. 363	
470 virginica L.	Virginian	■ au	Y	Virginia	1759.	D co	
471 aurea Ph.	golden	■ my.jn	Y	N. Amer.	1820.	D co Bot cab. 1399	
472 latifolia R Br.	broad-leaved	■ jl au	W	N. Holl.	1822.	D p.l	
473 quadndentata Mx.	four-toothed	■ my.au	W	N. Amer.	1821.	D co Lam.il. 16.2	
54. BONNA'Yi< Lk.	BONNAYA. (<i>Ronnay</i> , a German botanist)	<i>ScrophidariKT</i> . 3. —12.					
474 brachiata Lk.	brach late	O pr 1	au	'W	Brazil	1823V	S r.tn L. &O. a.b. g.ll
475 veronicetef Wa Spr.	speedwell-lvd	■ ijn.s	Pk	£. Indies	1798.	C co R.mal.9.59	
Gratiola veronacifolu. Arias.							
476 reptans Spr.	creeping	fr ES3 or	jn.jl	B	E. Indies	1820.	D & p
* 55. SCHIZ A'NTHUS Ft. per.	SCIIIZANTHILS. (<i>Schizo</i> , to cut, <i>anthos</i> a flower.)	<i>Scrophutdrinte</i> .					
477 pinnatus Ft. per.	pinnate	O or 2	jn.o	WP	Chile	1822.	■ Lp Hook. ex. fl. 73
478 pömgens Hook.	spreading-sto/fced	O or 2	jn.o	W.P	Chile	1822.	■ Hook. ex. fl. 86
56. ELYTRA'RIA Mx.	ELYTRARIA	(<i>Elytron</i> , an envelope: its scaly stem.)					
479 virg&ta Mx.	twiggy	■ ■ ■	-		Carolina	1813.	D Bp Mic. am. 1.1
480 crenata Vahl	crcnate	■ ■ ■	-		E. Indies	1820.	D s.p Rox. cor. 127
Justice acatilis Rox.							
481 margin&ta Bcauv.	marginate	■ ■ ■	-		Guinea	1823.	■ 8.p
482 imbricata Pers.	imbricated	■ ■ ■	-		W. Indies	1824.	■ s.p
483 caulescens Led.	caulescent	■ ■ ■	-		E. Indies	1823.	■ s.p
484 ramosa Hum.	branched	■ ■ ■	-		Mexico	1825.	■ co
485 lyrata Vahl	lyrate	■ ■ ■	-		E. Indies	1825.	S co
57. HYPOESTES Sol.	HVPOESTES. (<i>Hypoestet</i> , an inner garment; the involucrum.)	<i>Acanthacea</i> . 4. —11.					
486 involucrata Rox.	involucred	• o r ljjl.au	W	E. Indies	1811.	C p.l	
487 purpurea R. Br.	purple	jg E J or 2	my.jn	Pu	China	1822.	C p.l
488 fastuosa Sol.	sumptuous	• or 2	my.jl	R	E. Indies	1818.	C p.l
489 serpens JL Br.	creeping	ArOSor	ijl.au	...	Australia	1820.	D p.l
*58. JUSTI'CId L.	JUSTICIA. (J. Justice, an eminent Scotch hort and bot)	<i>Acanthacea</i> . 58. —150.					
490 bicalyculata Vahl	two-calyxed	■ or 3	au	Li	E. Indies	1755.	■ 81 Cav. ic. 1.71
491 hispacia W.	hispida	£ E) or	iap.my	Y.w	S. Leone	1824.	D p.l
492 Echium L.	ex lulling JK	• or 3	mr.au	B	R Indies	1759.	C pi Bot mag. 1847
493 thrysiflora Rox.	thyrsiflower'd	• or 8	ap.jl	S	E. Indies	1812.	C p.l
494 vitelina Rox.	yolk-like	n. or 1	my.jn	Y.B	E. Indies	1818.	C p.l
495 alba liox.	white	ft • or 2	ap.jl	W	E. Indies	1816.	C p.l
496 corinca Aub.	scarlet tt.	■ or 5	f	S	S. Amer.	1770.	C p.l Bot mag. 432
497 t'chiofes L.	Echium-like j£ Z3	1	ap.my	R	E. Indies	1820.	C p.l
48 quadrifida Vahl	four times cut tt.	• or 3	mr.s	S	Mexico	1795.	C p.l Par. Ion. 50
499 pumila Die.	dwarf	«L or 1	mr.jn	R	S Amer.	1820.	C p.l
600 nigrans Lou.	blackish	«L J or 6	mr.8	W.R	China	1819.	C p.l
501 nitida Jac.	clear	tL QH or 4	mr.s	W	W. Indies	1790.	C pi Bot. rep. 57
502 bractolata Jac.	small-bracted	tL or 6	jl.au	P	Caracas	1823.	C p.l Jac. ic. 215
503 picta L.	painted	tt or 8	jl.au	C	E. Indies	1780.	C p.l R.mal.6.60
2 lrido-sanRUineae	lurid blood-lvd	■ or 8	jl.au	C	E. Indies	1780.	C p.l Bot mag. 1870
504 paniculata Vahl	paniced	*.Qor	jl.jl.au	Pk	E. Indies	1811.	S p.l R.mal.9.56
505 variegata Aub.	stuffed flowered*	2	my.jn	R	Guiana	1825.	C p.l Aub. gui. 1.4
506 alicina Vahl	vi\ovi-leaved M	li	au.s	R	Peru	1816.	C s.p
507 polystroma Rox.	many-seeded Jk	{	jl.au	P.B	E. Indies	1818.	S p.l
508 secunda Vahl	side-flowering J*	3	jn.jl	R	W. Indies	1793.	C p.l Bot mag. 2000
509 geniculata B. M.	jointed m	2	jn.jl	P	W. Indies	1820.	C p.l Bot mag. 2487
510 ciliaris L.	ciliated	1	jl.au	W	W. Indies	1780.	S 81 Jacvin.2 104
511 lufida Vahl	shining-leaved M	or 3	jl.au	S	W. Indies	1795.	C p.l Bot mag. 1014
512 elongata Vahl	elongated	■	myjn	R	E. Indies	1812.	C sp
513 cuspidate Vahl	cuspidate	■	ij.	S	Arabia	1820.	C sp
514 salvisflora Hum.	sage-flowered	■	jl.au	S	Mexico	1824.	C p?
£15 Gendarissa L.	Gendarussa	■	jn.jl	Li	E. Indies	1800.	C p.l Botreg. 635
516 carthaginensis Jac.	Carthagensa	■	jl.jl	P	Carthag.	1792.	C s.p. Botreg. 797
517 nemorosa Sun.		2	my.jn	W	W. Indies	1825.	C s.p. Botreg. 797
518 pedunculosa Mx.	grove	U	jl.au	Li	N. Amer.	1759.	C ■ 1 Bot mag. 2367
519 polystachya Lam.	pedunculated	2	jn.jl	P	Guiana	1821.	C p.l
520 procumbens L.	many-spiked	ijl.jl	au	Pk	E. Indies	179a	C s.p. Pluk. al. 56.
521 Guinguangelxia Ktne	procumbent	i	jl.au	P	E. Indies	1820.	C p.l
822 diffusa W.	diffuse	1	my.jn	W	K Indies	1826.	C p.l
523 htimilis Mx.	dwarf			W	N. Amer	1318.	C p.l
524 comata L.	comate			jl.au	Jamaica	1795.	R sp SI.jam. 1.103.2
525 eustachiflora Jac.	Eustachian			au.1	St. Eustac.	1799.	C s.p. Jac. am. pic. 15

ORDER I.

PIANDRIA MONOGYNIA.

Vahl D	\rightarrow	xipn8BA.	(D«, double bellklo, to shut: fruit of 2 valves)	Acanthacea: 11.—92.
49 resupinata Vahl	six angled	or	2 JI R	Pluk.al 279.6
550 biyalvis Vahl	resupinute	rm or	W.P	Cav. ic. 3.203
554 two-valved	■ or	i	S. Amer. 1805. S p.1	Cav. am. 6. 22.1
554 martinicensis Vahl	Martinique	tt-CDor	E. Indies 1818. C p.1	Jac. am. 5. 6
gr J ^ a n a Vahl	Peruvian	tt,	P W. Indies 1818. C s.1	Fl. per. 1.16. a
553 verticillaria Vahl	vrhrl-flowered	CD or	2 my.au P Peru 1818. C p.1	Bur. in. 8.64.1
554 chinensis Vahl	China	jC	my.jn P C. G. H. 1826. C p.s	Rox. cor. 2.153
555 pectiniflora Vahl	pectinated	tt,	alor Pa.B E. Indies 181& C s.1	K. Hou. a 1
556 scorpioides Vahl	Hcorpiou-like	tt,	alor B E. Indies 1798. C p.1	Br. jam. 2.1
557 audrigens Void	asHiengrt	tt	leu 2 jl.au R W. Indies 1818. S p.1	Rot. cob 721
558 rectusa Vahl	retuea	ig	2 rar.an R W. Indies 1818. S p.1	

R. Br. NELSONIA.	(D. Nelson, a bot who accompanied Capt. Cook.)	<i>Acanthacea.</i>	1.—5.
hairy	CD un	jil, au	R. E. Indies 1818. C. p.l
61. ERA'NTHEMUM It. Br. ERANTHEMUM.	(Ear, spring, antbos, a flower.)	<i>Acanthacea?</i>	10. — 24.
560 spumosum <i>b.</i> Br. thorny	tt-CD or r	jLaa	W. Indies 1733. C
561 pulchellum 2T. li. pretty	tt-CD or	ja.o	B. E. Indies 17N. C
*B variabile A ^r. variable	tt-L-Jor	jnjl	P. N. Holl. 1820. C
563 bicolor 2T Af. two-colored	tt. CD or	in.y.au	Luconia 18JZ. C
56* strictum llox. strict	■ Q44	mr.ap	Nepal 1818. C
5a? crenulatum Wai. crenulate	n-CDpr	o	Nepal 1824. C
566 capense L. Cape	tt. (L) or	apiny	E. Indies 1818. C
567 elegans J'aув. elegant	■ jnjl	Sc	Guinea 1824. C
568 roseum R. Br. rose-colored	tt-CDpr		E. Indies 1820. C
«J9 ambiguum R. \$ S. ambiguous	tt,CDor	&	1821. C
		R	C. s.p

^W WULFENIA (F. X. Wulfen, a German botanist.) *Scrophylidin**. 1.—
Carinthian O pr lijl.au B Carinthia 1817. S co Jac. ic. 1.2
2.

					Italy	1824.	D	s.p	Jac.au.	5.39
0<* pinnaca L.		Ang Jeaved							Scrophularidire. 6	55.
575 scabiosifolia L. & S.	Scabiosifolia	Ivd	t ^ Kor	2	m.v.o	Y	Chile	1773.	S. s.p	Bot. mag. 41
576 rugosa L. per.		wrinkled	tt .Jor	2	au.s	Y	Chile	1822.	C co	Bot. mag. 2405
577 integrifolia L.	entire-leaved	<. Jor	2	au.s	Y	Chile	1822.	C co	Hook: ex. fl. 99	
578 coriacea Cavanil.	F ory pose	Alor	1	myjn	Y	Chile	1822.	C co	Bot reg. 744	
		A .	1	ft myv.u	Y	Falkl. I.	1777.	S. co	Bot. reg. 723	
								D In	Bot. mag. 348	

581 <i>vulgaris</i> L.	Portugal	+	+	+	jn.jl	Li		bogs	D bog	Eng. bot	145
5 ² <i>alpina</i> L.	common	+	+	+	Jray	V	Britain	bogs	D bog	Eng. bot	70
« <i>S. grandiflora</i> Lam.	alpine	+	+	+	i ap	W	Germany	1794.	D bog	Fl. dan.	453
5 ³ <i>liffl. tea</i> Walt.	large-flowered	+	+	+	i apmy	B	Britain	ir. bog.	D bog	Eng. bot.	1128
»» <i>edentula</i> Hook.	yellow	+	+	+	* jn.jl	Y	Carolina	1816.	S bog	Bog Bot reg.	126
	toothless	+	+	+	i ap	Y	N. Amer.	1S23.	D bog	Hook. ex. fl. 1	
585 v. ^{C. f.} L A R I A	⁷ <i>omis</i> SS	DEDMDI	if ⁰ 1	£	^ Vittletottle; w. toroo <u>i</u>)	w. te. 31	-63.				
gffminor!.	IS?2	S	£	o	* Jn.jl X	2 ritaln	"ta.wa.Daq	Eng	bot		253
588 <i>Intermedia</i> Hay "	KSStrn	S	SS	Scu	* £. Sy. K	Y S ^ a ! S E w a S Q Bq3	Egg	Spok	&		

591	urticifolia	B. M.	nettle-leaved	or	2	my.jl	B	S. Amer.	•	S	p.1	Jac. ob. 4.86
592	umbrifera	Hum.						S. Amer.	•	S	p.1	Bot. mag. 1848
592	jamaicensis	Vahl	Jamaica		2	jn.s	B	W. Indies	1714.	C	p.1	Bot. mag. 1860
593	prismatica	Vahl	prismatic		2	my.jn	B	W. Indies	1699.	C	p.1	Jac. ic. 2.208
594	cayennensis	Vahl	Cayenne		3	my.jn	B	Cayenne	1892.			
595	hirsutissima	Lk.	most hairy		1}	mr.ap	B	Brazil	1822.			
596	Borubicanus		Orubrian		3	jn.au	V	Panama	1G99.			
597	mutabilis				3	mr.e	O	S. Amer.	1801.	Lp	Ehr.pic.5.1	

600 australis R. Br.	southern				W	N. Hull.	1823.	D co	
601 virginicus L.	Virginian				Wp	Virginia	17G0.	D m.8	
2 ffucrificus	Oak-leaved				W	N. Amer.	17XX	D co	
602 intermedium Schr.	intermediate				W	Europe	1820.	D m.s	
70. AMETHYSTEA L.	AMETHYSTEA. (<i>Amethystos</i> , the amethyst; color of flower.)				O pr	1J jn.jl	B	Labidhe. 1.	
603 cserilea L.	blue-flowering					Siberia	1759.	S p.l	Bot mag. 2448
71. ZIZIPHORA L.	ZIZIPHORA. (<i>Zizi</i> , the Indian name, and <i>phoreo</i> , to bear ?)				Jk A rk	t jl.au	II	Labiatee. 10.—13.	
604 ocinoldes L.	Thyme-like				O rk	!in	Pk	Siberia	1786. D 81
605 dinopodioides Xnm.	Basilicuicke								Ac. pet. 1810.11
icryllacea H. K.	headed				O or	JjLau	R	Syria	1752. FL gr. 1. IS
606 capitata L.	Spanish				O or	i jn	R	Spain	1759. Lam. il 1& 1
607 hispanica L.	spiked				O or	ijjn	P	Spain	1826. Lam. il. 18.2
fi 68 spicata Lag.	slenderer				O or	*"jn.jl	Li	levant	1752. Lam. il. 18.2
609 teniflor L.	Taurian				O or	Jjl.S	II	Tauria	1816. Bot. mag. 906
610 tafrica L.	Serpentillum-like*				A rk	* jl.au	R	Caucasus	1801. Bot. mag. 906
611 *eryllacea Bieb.	mediate				t. A rk	4 jn.au	R	Caucasus	1822. Bot. mag. 1093
612 media LK.	thick-flowered				JU rk	} jn.au	Ii	Siberia	1803. Bot. mag. 1093
613 dasyletha liib.									
Pouskhni H. K.									
72. CUNILA L.	CIINILA. (<i>Konos</i> , a cone, or <i>Cunila</i> , the name of a town ?)				Labidte. 2. — 5.				
614 marifina L.	Maryland				A or	1 jls	It	N. Amer.	1759. M. h.3.19.7
615 coccinea Nut.	scarlet				£ IAJ I*	s	S	N. Amer.	1823. C p.l Hook. ex. fl. 163
7a ZJEDEO'MA Pers.	HKDEOMA. (<i>Hedcoma</i> , a Greek name for mint.)							Lubidtic. 2.-3.	
fl16pulegiofdes Ph.	Pennyroyal-like				O or	Jnau	B	N.Amer.	1777. S co
617 /hyomofdes Pars.	Thyme-like				O or	£ jn.au	R	France	1699. S co M.h.3.19.6
74. MONARO ROrf L.	MONARDA. (<i>N. Monardext</i> a physic, of Seville, in 16th cent.)							Labidte. 17.—	
618 fistulosa L.	hoWow-stalked							N. Amer.	1656. D r.m Mil. ic. 122.2
619 media W. en.	mediate							N. Amer.	1656. 1) r.m
620 m6lis L.	soft							N. Amer.	1656. D r.m
621 oblongata JL.	obvoig. feaved							N. Amer.	1771. D r.l
622 chnopodia /.,	Wild-BasiWw*							N. Amer.	1789. D r.l Bot. mag. 145
623 purpurea Ph.	purple							N. Amer.	1821. D r.l
634 altissima W.	tallest							N. Amer.	1821. R pl
625 affinis Lk.	allied							N. Amer.	1821. R pl
626 rugosa H. K.	wrinkled							N. Amer.	1761. D r.l
627 Kalmirfa Ph.	Kalm's							N. Amer.	1813. D Pl Ph. am. 1
628 didyma L.	twin, or <i>Qswege tea</i>							N. Amer.	1752. D r.l Bot. mag. 546
629 Russellifna Sims	illuscl's							N. Amer.	1823. R pl Bot. mag. 2513
§630 ciliata /.,	ciliated							N. Amer.	1798. D r.l Pluk.al. 164.3
631 dtriodra Cav.	lemon-scented							Mexico	1825. II pl
1632 hirsfita L.	hairy							N. Amer.	1798. D r.l
633 gracilis Ph.	slender							N. Amer.	1798. D r.m
634 punctata L.	dotted							N. Amer.	1714. S s.p Bot reg. 87
75. FLOSMARI'NUS	L. ROSEMARY. (<i>lios</i> , dew, <i>marfnis</i> , of the sea; on sea-shores.)							Labidta. 2.	
635 officinalis /..	official				or	4 ja.ap		S. Europe	1548. FLgr.1.14
2 latifolius	broad-leaved				il	or	12 ja.ap	S. Europe	1548.
3 fol. argenteis	silver-leaved				•	or	4 ja.ap	S. Europe	...
4 fol. aOreis	gold-leaved				•	or	12 ja.ap	S. Europe	1548. C
636 chilensis MoL	Chile				il	or	4 jl	Chile	1795. C
76. SA'LVIA L.	SAGE. (<i>Salvus</i> , safe; medical qualities.)							Labidtee. 124.— 230.	
637 pomifera L.	apple-barked				tt.	or	2 jl.au	B	Candia 1699. C p.l
638 calycina Sm.	large-calyxed				tt.	or	1 jl.au	Pk	Levant 182.1 C co
639 canariensis L.	Canary				4 jns	P		Canaries 1697. C l.p	Tr.pl. 2.19
640 atrea L.	go'dcn-flwd				LJ or	3 ap.n	Y	C. G. H. 1731. C p.l	Bot mag. 182
641 dentata H. K.	tooth-leaved				1 Jor	* dja	W	C. G. H. 1774. C p.l	
642 interfifpta Schous.					or	1 apss	W	Barbary 1798. C s.l	Schou.fi. 1
643 pilanthes Lk.	hairy-flowered				or	2 jl.au	B	1822. C co	
644 pinnata L.	vr'ingcf-leaved				Q or		P	Levant 1731. C s.l	Boer. 1.167
645 Hablitz/Ytma W.	Hablitz's				or	iiii	B	Siberia 1795. C co	Bot mag. 1429
646 scabiostefolia Lam.	Scabious-leaved				J<	or	1 au	W	Siberia 1818. C co An.mu. 15.15
647 vulnerariaxifolia IV.	Vulneraria-lvd				or	jl	W	Armenia 1820. C co	
648 ltnccolata Cav.	lanceolate				O or	1 my.s	B	W. Indies 1813. S co	Jac. cc. 2.12
649 hirsuta Jacq.	hirsute				O or	1 my.jn	B	1801. S co	Jac. sc. 3.252
650 angustifolia Cav.	narrow-leaved				jl	jl	W	Mexico 1806. C co	Cav. ic. 317
651 leuca'ntha Cav.	white-flowercd				PI	IAI or	2	Mexico 1825. C co	Cav. ic. 1.24
652 azurca Lam.	azure-flowered				5. A or	6 au	B	Carolina 1806. C co	Bot. mag. 1728
653 ameth^fetina Sm.	Amethyst				yg	1Alpr	2 a.us	B	Colombia 1817. C 6.1 Sra. ic. 2.27
654 psetiduccoccinea Jinc.	bastard-scarlet				(23	3 jn.au	S	S. Amer. 1797. C s.p	Jac. ic. 2.209
655 Boosia TM Jac.	Boos's				1Alor	2 mr.ap	B	Peru 1821. C co	Jac.cc. 1.47
S. amce'na Bot. reg. 446.									
656 rosea Vahl	rosy				tt.	jl.or	W	E. Indies 1816. C s.l	
657 tubiflora Cav.	tube-flowered				tt.	1Alor	S	Mexico 1820. C r.m	
658 /amititola Jac.	Lamium-lvd				tt.	or	ji	S Amer. 1821. C p.l	Jac. sc. 3.318
659 mexicana L.	Mexican				£	1Alor	S	Mexico 1724. C p.l	Cav.ic.1.26
660 chamaedryoides Cav.	Germander-liken.				LAJ or	4 mv.jl	B	1795. C p.l	Bot mag. 308
661 <corodoniae61iaD^	Scorodonia-lvd				tt.	jl.au	W	1825. C co	
662 ce'sia IV. en.	grey				tt.	1Alor	S	1813. C p.l	
663 involucrata Cav.	involucrate				tt.	jl.au	P	1825. C s.l	Cav. ic. 2.105
664 purpurea Cav.	purple				g	1Alor	2) my.jl	P	1825. C 6.1 Cav. ic. 2.106
665 hispanica L.	Spanish				O or	1J jn.au	P.B	Spain 1739. S p.l	Bot reg. 359
666 ser6tina L.	late-flowering				g	1Alor	B	Ohio 1803. C s.l	Jac. ic. 1.3
667 dominica TV.	Dominica				El or	f' jl	W	W. Indies 1759. C sp	Swz. on. 18.1.1
668 Miae6lia Vahl	Lime-tree-lvd					4 jn.au	B.c	S. Amer. 1793. C p.i	Jac. sc. 3.254
669 polystachya Or.	many-spiked				LAJ	3 o.d	B	Mexico 1822. C co	Jac.sc. 3.318
670 micrantha Vahl	small-flowered				tAlor	1 my.jn	B	Cuba 1823. C co	
671 formosa W.	beautiful				» or	4 ap.o	S	Peru 1783. C p.l	Bot. mag. 376
672 coccinea L.					Al	2 ap.o	S	S. Amer. 1774. C p.l	Mur. 177a 1
673 pulchella Dec.	pretty				1Alor	2 o.f	S	S. Amer. 1821. C co	Col. h. np. 16
674 amarissima Or.	bitterest				1Alor	2 jlau	B	Mexico 1803. C fl.p	Bot. reg. 34'
675 glutinosa L.	glutinous				A or	3 jn.s	Y	Germany 1796. C co	M.h.3.ia18

762 scabriuscula H. K.	roughish oval Jeaud		E. Florida 177(1) D p.l
763 oviflisis Ph.			Carolina 1812. D p.l
764 tuberosa Mx.	tuberous		Carolina 1806. R p.l
765 anadUB. M.	Anise-scented		Carolina 1806. D p.l Bot. mag. 1213
•78. CATA'LPA J.	CATALPA. (Indian name.)		E. Florida 177(1) D p.l
	synnaga-tvd or 20 jn.au	W	N. Amer. 1726. S p.l Bot. mag. 1094
767 longissima H. K.	longest-podded	CD or 20	W. Indies 1777. L s.p PL ic. 57
768 microphylla Spr.	small-leaved	Dor 15	Hispaniol. 1820. C p.l PL ic. 55
Bignonia microphylla Lam.			
79. GHI'NII Schreb.	GIINIA. (Ghini, an Italian botanist.)	< Verbenaceae. 1.-2.	
769 spinosa Schreb.	thorny-bracted	O cu 2 au P	W. Indies 1733. S s.l R. Hou. 2
80. FONTANE'S/i<	Lab. FONTANESIA. (M. Desfontaines, professor of botany at Paris.)	Oleaceae. 1. — 2.	
770 phillyreae-ofdesla. a6.	phillyrea-like	tt-Jbpr 18 aw Y	Syria 1787. € 84' Lab. syr. 1.1
81. LINOCIE'R^ Swz.	LINOCIERA. (G. Linocier, a French physician.)	O&ifue. 3.-7.	
771 compacta R. Br.	compact	• o r 10	W W. Indies 1793. C Lsp Jac.c. 2.6.1
772 /igustrina Swz.	Privet-like	D o r 12	W E. Indies 1820. C ls
773 cotinifolia Vahl	Cotinus-leaved	sh CD or 12	W Ceylon 1818. C s.l
•82. ACA/NA L.	ACJENA.	(Akaina, a thorn; the berjeaj	SanguisSrbia? 10. — 16.
774 latebrosa H. K.	latebrose	AI 1 ap.jn G	C. G. H. 1774. C l.p.s
775 pinnatifida Fl. per.	pinnatifid	my.jn	Chile 1822. D l.p.s Fl. per. 1.104
776 ovahföha Fl. per.	oval-leaved	my.jn	Peru 1802. n. l.p.s Ven. eels. 6
777 adscendens Vahl	ascending	my.jn	Magellan 1822. u l.p.s
778 magellana Vahl	Magellan	my.jn	Magellan 1823. D co Lam.il. 22.2
779 Sanguin'irba Vahl	BurneUleaved	jn	N. Zéal. 1796. D l.p.s Lam. il. 12.1
780 Mcida Vahl	shining	i my.jn G	Falkl. I. 1777. D l.p.s Lam.il. 22.3
781 argentea Fl. per.	silvery	2 my.jn G	Chile 1822. D l.p.s Fl. per. 1.103
782 lsvigata 11. K.	smooth	4jn.au G	Magellan 1790. D l.p.s
783 ovina Cun.	sheep	* jn.jl G	N. Holl 1818. D co
84. O'RNU'S Pert.	FLOWERING! Asn.	(Qren, Heb. oreijas. Gr. ornus, Lat. a wild ash.)	Olinna. 5.
784 europe'a Pen.	European	Y or 30 my.jn W	Italy 1810. (f co FLgr. 1.4
785 american'a Ph.	American	y or 30 my.jn W	N. Amer. 1812. G co
786 rohindifolia Pen.	round-leaved	Y or 30 ap W	Italy 1697. G co W.b.m. 2.1
787 striata Swt.	striated	\$ or 30 ap.my W	N. Amer. 1818. G co
/Växinus Strifta Spr.			
788 flonbiunda Wai.	bundle-flowered^	or 30 .■ W	Nepal 1822. G co
84. MoaiNA.	(L. Morin, a French botanist)	Dipsæete. 1.	
789 pērsica L.	Persian	AI or 3 jl.au R w Persia	1740. C s.p FLgr. 1.28
85. CIRC JEM L.	ENCHANTER'S NIGHTSHADE.	(The enchantress Circe; damp sha, pi)	Onagrdrhv. 3.
790 lutetiina L.	Parisian	^L A or 1 jn.au R	Britain sha.pl. D co Enp. bot. 1056
791 intermedia Ehrh.	intermediate	3i A or ijn.au R	Europe 1821. D co FL dan. 256
792 alpina L.	mountain	^ A or i jn.s R	Britain moun. D co Eng. bot 1057
86. FE'DIA Moen.	FEDIA. (A name of Adonis, probably without meaning.)	Valentine*. 1.—2.	
793 cornucopia? Vahl	horn of plenty	O or 1 jn.jl R	S. Europe 1796. S co FL gr. 32
*87. PIMELE'A Forst.	PIMELEA.	(Pimele% fat?)	Thymdriet. 12.—32.
794 /iniflha Sm.	flax-leaved	Lj 2 f.au W	N. S. W. 1793. C s.p Bot. mag. 89
795 collina It Br.	hill	.. i J pr 2 ap.my W	N. Holl. 1824. C s.p
796 gladca R. Br.	glaucous	2 my.jn W	N. Holl. 1824. C s.p
797 /igistrina Lab.	privet-like	2 f.au	N. Holl. 1823. C s.p
798 /inoides Cun.	flax-like	2 ap.s W	N. Holl. 1826. C s.p
799 decussata R. Br.	decussate	a-Lj pr 3 my.jn R	N. Holl. 1824. C s.p
800 rōsea R. Br.	rosy	tL i lor 3 my.jn R	N. Holl. 1800. C s.p Sw.au. 8
801 filamentosa Rud.	thready	tl J lor 1	N. Holl. 1826. C s.p Bot. mag. 1458
802 drupacea Lab.	cherry-fruited	a. j or my W	N. Holl. 1817. C s.p Lin. tr. 10.14.1
803 pauciflora Schr.	few-flowered	H i lor my W	V. Di. L. 1812. C s.p Bot. cab. 540
804 spicita R. Br.	spiked	tl J or 2 mrs W	N. Holl. 1824. C l.p. Bot. cab. 179
805 incana R. Br.	hoary	tL i J cu 2 ap.my W	N. Holl. 1824. C s.p Lin.tr. 10.14.
88. CLA'DIUM Schr.	CLADIUM.	(Klados, a branch or twig.)	Cyperaceee. 5.—14.
806 germāicum Schr.	German	ilk A un 3 jl.au Ap	Britain mar. D m.8 Eng. bot 950
807 occidental Schr.	Western	M EJ cu 3 my.jn Ap	Jamaica 1820. D Lp
808 glomeratum IL Br.	glomerated	M iAI cu 1 jl.au Ap	N. Holl. 1816. D aq
809 /mceum R Br.	rushy	Jk uAJ cu 1 jl.au Ap	N. Holl. 1820. D aq
810 *cho-noJdes R. Br.	Schochnus-like	Jr iAJ cu + jlau Ap	N. Holl. 1823. D aq Lab. n.h. 1.18
Schas'nus acQthus Lab.			
89. GU'NNERJ L.	GUNNERA.	[E. Gunner, bishop of Drontheim, a botanist)	Urticae. 1.—3."
811 perp^nsa L.	well considered^	iAicu l jlau P	C. G.H. 1688. D m.s Bot mag. 2376

CLASS II. ORDER 2.

DIA'NDRIA DIGY'NIA.

9a ANTHOXA'NTHUM L.	SPRING GRASS. (Anthos, a flower, xanthos, yellow; spikes.)	Gramitwce.
812 odor&tum L.	sweet-scented	m Ag L my Ap Britain me pa. S h.l Eng. bot 647
21&xum	loose	JL A cu my Ap Britain me.pa. S co
3 pub^scens	pubescent	M A cu my Ap Britain me.pa. S co
4 ramoBum	branched	m A cu ST Ap Britain me.pa. S co
81S amarum Brot.	bitter	M A cu ST Ap Morocco 1810. S co
814 ovatum Lag.	ovate	M O CU i ji Ap Spain 1821. S co
815 gracile BivT	slender	Jm A cu lap Ap Sicily 1820. S co Biv. sic M
91. CORYCA'RPUS Zea.	CORYCAHPUB.	[Korys, a helmet, karpot, a fruit.)
816 arundinaccus Zea.	recl-like	Ji ^ c u S ap.jl Ap N. Amer. lbo. S co Mic.am. iō Fcstuca diandra Mx.

92. SCHMIDT Trat SCIIMIDTIA. (*Schmidt*, a German botanist) *Gramine**. 1.
817 subtilw Trat. small i Q cu i my.jl Ap Bohemia 1820. S p.1 Prat. au. 451
Colcauthua subtilis RSS.

CLASS II. ORDER S.

DIAN'DRIA TRIGYNIA.

93. PITER L. PEPPER. (<i>Pippul</i> , [REDACTED]ese, long pepper; or [REDACTED]i/o.todig>t)	<i>Pipers**-</i>	<i>^irji</i> ⁶
818 peltatum L. peltated « 2	Ar	[REDACTED]
819 umbellatum L. umbelled	W. Indies 1748.	C r.m PL am. 56.4
820 macrophyllum L. broad-leaved CL	W. Indies 1748.	C l.p
821 geniculatum W. swoUen/jomtcd «L	W. Indies 1800.	C r.m Aub.gui.23
882 aduncum L. hooked'	Jamaica 1823.	C r.m
883 hispidum 7/. Sf.B. hid&pidUeaved	Jamaica 1748.	C r.m Jac ic. S. S10
824 hirsutum Sws. hairy	S. Amer. 1793.	C r.m
825 Amaldeo L. Amalago	W. Indies 1793.	C r.m
83G M L Betle	Jamaica 1759.	C r.m SL jam. 1. 87.1
827 nigrum L. black	E. Indies 1804.	C r.m R.mal. 7.15
828 tuberculatum Jac. tuberculate	E. Indies 1790.	C r.m Lam. U.79.53
829 triocicum Itox. trioccious	S. Amer. 1816.	C r.m
830 discolor Swx. two-colored	E. Indies 1818.	C r.m Jac. ic. 8. £12
831 reticulatum /.	W. Indies 1821.	C r.m Bot cab. 610
832 decumanum L. netted	W. Indies 1748.	C r.m PL am. 57.75
833 Siriboa	Carthag. 1768.	C r.m Jac. ic. 2. 215
834 longum L. long	E. Indies 1768.	C r.m Ku. am. 5.117
835 glabrum Mil. smooth	E. Indies 1788.	C r.m Ru.am.5. lift 2
836 aurifolium Mil. Laurel-leaved	Catnpeac. 1768.	C r.m
837 tomentosum Mil. downy	W. Indies 1768.	C r.m
838 nildum Swz.	W. Indies 1768.	C r.m
839 racemosum Mil. great racemose	Jamaica 1793.	C r.m
840 acutifolium R.A.S. sharp-leaved	Campeac. 1768.	C r.m
841 colubrinum Lk. snake	Peru 1823.	C r.m Fl. per. 1.61. a.
842 mediate	Brazil 1820.	C r.m
843 marginatum Jac. marginate	W. Indies 1820.	C r.m Jac.ic.8. 8
844 glaucococcum Jac.	S. Amer. 1811.	C r.m Jac. ic. 2.15
	Peru 1822.	C r.m Jac. ec. 1.76
94. PE-EBCMI A R 4 P. PEPEHOMIA. (Analogous to <i>Piper</i>) <i>Piperdeca</i> . 41.—134.		
845 brathyphylla W. short-leaved £ 23 cu jns. Ap	S. Amer. 1818.	C r.m
846 amplexirau Hs&wz. stem-clasping £ ZJcu jns. Ap W.	Indies 1793.	C r.m
847 magnolia/ta Vahl Magnolia-lvd £ 23 cu ja.mr Ap	W. Indies 1743.	C r.m Jac. ic 2. 213
848 obtusifolia IV. obtuse-leaved £ 23 cu apjl Ap	W. Indies 1739.	C r.m Tr. chr. 54. 9
849 c'ma/gfia Jac. Clusia-leaved Jw 22 cu myjn G	S. Amer. 1K17.	C p.1 Jac. ic. 2.212
850 cuncifolia W. en. wedge-leaved £ 23 cu jn.jl Ap	Caracas 18&	C r.m Jac. ic. S. 214
851 macrostachya Rich, large-spiked ^ 23 cu myjn G	Trinidad 1K-4	C p.1
852 tenella Swz. slender	Jamaica 1820.	C pi
853 tristachya Kth. three-spiked	W. Indies 1826.	C p.1 H. & B.n. 1.6
854 reniflora Hook. reniform	St. Vine 1824.	C p.1 Hook. ex. fl. 164
855 ovalifolia Hook. oval-leaved	St. Vine. 1834.	C p.1 Hook. ex. fl. 165
856 alpita Vahl winged	S. Amer. 1812.	C r.m Fl. per. 3/48
857 acuminate W. en. acuminate	W. Indies 1812.	C r.m Hot.mag. 1882
858 distachya Pers. two-rowed	S. Amer. 179a	C r.m I'l. am. 51.67
859 conacea Vahl leathery	E. Indies 1815.	C r.m Bot. cab. 128
860 discolor it. * S. two-colored £ Sf cu jlau Ap	S. Amer. 1821.	C r.m Bot cab. 610
861 maculosa Vahl spot-stalked £ 23 cu i s Ap	StDomin. 1790.	C r.m Pl. am. 60.66
862 polystachya Kth. pellucid	S. Amer. 17+8.	C r.m Pl. am. 54.72
863 rotundifolia Kth. round-leaved £ 23 cu jlau Ap	S. Amer. 1822.	C r.m Pl. am. 69
864 hispidula R. & S. little-hispid	Jamaica 1818.	C r.m Swz. ic. 4
865 nummularifolia Kth. moneyw.-lvd £ 23 cu myjn Ap	Jamaica 1818.	C r.m
866 pubescens Haw. pubescent £ 23 cu myjn Ap	S. Amer. laoy.	C r.m
867 hifimilis Vahl low £ 21 cu myjn Ap	W. Indies 1708.	C r.m
868 trifolia Pers. three-leaved £ 23 cu jn.jn Ap	S. Amer. 1802.	C r.m Pl. am. 52.68
869 pulchella IV. fair	Jamcka 177H	C r.m Hot. cab. 574
870 pernikiifolia Kth. Pereskia-lvd £ 23 cu myjn Ap	S. Amer. 1820.	C r.m Hook. ex. fl. 67
871 Manilla Kth. bland £ 23 cu myjn Ap	Caraccas 1802.	C r.m Hook. ex. fl. SI
872 rubricating Nees red-stemmeil £ ZJCU myjn Ap	... 1822,	C r.m Nccah.b.8
873 polystachya W. many-spiked £ 23 cu jn.jl Ap	Jamaica 1775.	C r.m Hook. ex. fl. S3
874 quadrifolia Kth. four-leaved £ 23 cu jn.jl Ap	S. Amer. 1818.	C r.m Hook. ex. fl. 22
875 ina?qualifolia RAS. unequal-leaved £ 21 or jlau Ap	Peru 1800.	C r.m Fl. per. 1.46. a.
876 stellata Pers. starry £ 23 cu myjn Ap	Jamaica 1802.	C r.m Jac. vin. 2.217
877 incana Haw. great-dovtny £ 7Z cu f Ap	Brazil 1815.	C r.m Bot. cab. 503
878 tahnifolia Kth. Talimum-Wd ^ 23 cu myjn G	W. Indies 1820.	C p.1
879 tricarinata Haw. three-keeled j^ 23 cu myjn G	W. Indies 1818.	C p.1
880 amplexiflora Lk. stem-clasping £ 23 cu fap Ap	S. Amer. 1823.	C r.m
881 subtropunda Haw. roundish/^ £ 23 cu f Ap	... 1812.	C r.m
882 serpent JL 23 cu myjn G	Jamaica 182a	C p.1
883 epensis KIA. creeping % 7\$ cu my.au Ap	S. Amer. 1823.	C r.m
884 rubella Haw. small red £ 23 cu mr.ap Ap	W. Indies 1820.	C r.m Hook. ex. fl. 59
885 capensis L. Cape . * ,23 cu myjn G	G.G.H. 18ft	C r.l

CLASS III. ORDER 1.

TRIA'NDRIA MONOGY'NIA.

95. VALERIA'NA L.	VAMHIAN.	{Valere, to be in health: or, Celtic supine Lavender TM*	or	W my.jn Pk h	or, Valerius, a king? Va.lerHb.neeB. SO.—70.
886 c&tica L.		Celtic supine Lavender TM*	or	W my.jn Pk h	Switzerl. 1748. D dac. c. 1. 1
887 supina Vahl		Celtic supine Lavender TM*	or	W my.jn Pk h	S. Europe 1822. D co Jac. m. 2.17. 2
888Sahntca All.		Celtic supine Lavender TM*	or	W my.jn Pk h	R France 18*1. D co Al. ped. 1. 70.1
889 saxatilis L.		Celtic supine Lavender TM*	or	W my.jn Pk h	Austria 1748. D co Jac.au. 3.267
890 elongata Jac.		Celtic supine Lavender TM*	or	W my.jn Pk h	Austria 1812. D co Jac. au. 3. 219
891 montana L.		Celtic supine Lavender TM*	or	W my.jn Pk h	Switzerl. 1748. D co Bot. cab. 317
892 alliariaefolia Vahl		Alharia-lvd	or	R.w Caucasus	EGL D s.p
893 tuberosa L.		tuberous-rooted &	or	R.w Caucasus	S. Europe 1629. D co M. h. 3.15.20
894 asarifolia Dufr.		Asarum-leaved	or	R.w Caucasus	Crete 1824. D sp
895 diof ca L.		diocious	or	R.w Caucasus	Britain mar. D co Eng.bot.fi28
896 Phu L.	Phu	& A or	3 my.jl	R.w Caucasus	Germany 1597. D co Black. 250
897 tripterus L.		three-winged	1 mr.my	R.w Caucasus	Jac. au. 3.268
898 heterophylla Lois,		various-leaved	1 jn.jl	R.w Caucasus	Lof gal. 1.2
899 pyrenaica L.		Pyrenean	2 my.jn	R.w Caucasus	Eng. bot. 1591
900 officinalis L.		officinal	5 A m	R.w Caucasus	Eng. bot. 698
901 «ambuct(tolia Mile.		Elder-leaved	2 jn.jl	R.w Caucasus	
902 «symbifl5lia Vahl		Sisymbrium lvdifc	Q> 1 jn.jl	R.w Caucasus	
903 intermedia Vahl		intermediate	1	R.w Caucasus	
904 Cardamines 2a?6.		Cardamine-lvd^	A or	Pk	
905 capensis Thunb.	Cape	A or	1	Pk	
•96. VALERIANELLA	Dufr.	LA-MD'S LETTUCE.	(A diminutive of Valerian.) Valerianeof.	18. — 24.	
906 echinata Dec.		pricMy-capsuled	O cul	L jl.au Pk	S. Europe 1807. S co Col ec. 1.206
907 olit6ria Dec.		salad	O cul	*ap. my Bk	Britain compi. S eo Eng. bot 811
908 dentata Dec.		toothed	O un	1 ap.jn	Britain corn fl. S co Eng. bot 1370
909 vesicaria Dufr.		bladdery	O un	4 ap.my	Candia 1739. S co Fl.gr. 1.34
910 coronata Dec.		crowned	O un	4 ap.jn	Portugal 1731. S co Colec. 1.209
911 discoidea Dufr.		discoid	O un	Japjl B	Italy 1731. S co M.h.3.16.29
912 carinata Dec.		keeled	O un	4 ap.my	France 1819. S co
913 eriocarpa Dec.		woolly-fruited	O un	1 apmy Li	France 1821. S co M.h.3.16.33
914 radiata Vahl		radiated	O un	1 ap.my	N. Amer. 1821. S co
915 dasycarpa Bieb.		shaggy-fruited	O un	1 ap.my Li	Crimea 1821. S co
916 uncinata Bieb.		hook-fruitcd	O un	1 my.jn Li	Tauria 1822. S co
917 auricula Dec.		eared	O un	I my.jn Pk	S. France 1824. S co
918 pdmila Dec.		dwarf	O un	4 ap.my	& Europe 1826. S co
919 mixta Vahl		mixed	O un	1 my.in Pk	Provence 1818. S co
920 platyloba Dufr.		broad-lobed	O un	4. my.jn Pk 1825. S co
921 excarpa Stev.		stemless	O un	4 my.jn Pk	Siberia 1826. S co
		three-toothed	O un	1 my.jn Pk	Tauria 1821. S co
97. OXY'B APHUS R.	S P. UMBRELLA-WORT.	(Oxys, acid, baphe, dyer's color.) Hyctaginece. 12.—13.			
923 viscous Herit.		viscid	jp	6 my.s P	C-Lp. Bot. mag. 434
924 aggregatus Cav.		aggregate	tf	1 jl.au Pk	N. Spain 1811. S 81 Cav. ic. 437
925 glabfolius W. en.		smooth-leaved	tf	1 jl.au Pk	N. Spain 1811. C s.l Cav. ic. 379
926 Cervantes's Lag.		Crvantes's	jf ^ATCU	2 jl.jl P	Mexico 1823. S r.m Sw. fl. gar. 84
927 expansus R.S/P.		expanded	JU^A Jcu	2 jl.au P	Peru 1819. S r.m Fl. per. 1. 75. a.
928 ovatus Vahl		ovate	L Alcu	2 jl.s P	Peru 1820. S s.p FL per. 1.77. b.
929 nyctaginea Nut.		nocturnal	-4 T cu	1 jls P	Missouri 1823. S s.p
All6nia nyctaginea		A/x.			
930 pilosus Nut.		pilose	-* A cu	1 ■■■	Missouri 1812. S fl.p
All6nia pilosa PA.					
931 hirsutus Nut.		hairy	-4 A cu	1 jl.s	P Louisiana 1812. S s.p
932 angustifolius Nut.		narrow-leaved	-A A cu	1 jl.s	
Calymenia angustifolia AW.		Alhonia linearis Ph.			
933 decumbens Nut.		decumbent	-* A *	2 jl.s	Missouri 1818. S s.p
98. BOLDO'jf Lag.	BOLDOA.	(D. Soldo, a Spanish botanist)			Nyctaginece. S.—5.
934 purpurascen8 Cav.		purpurascen	£	1 myjn V	Cuba 1820. C sj Cav.h.m.7
935 fanceolata Lag.		Iwnce-leaved	tt	3 jn.jl P	W. Indies 1824. C 81 H. & B. 1. 44
Salpianchus arcuarius	//. § B.				
99. LOEFLI'NG/^ L.	LOSFUNGIA.	(P. Lafpling, a Swedish botanist)	G	■■■ e*	2.-3.
936 hispanica L.	Spanish	O cu ijin G	Spain 1770. S. I	Cav. ic. 1. 94	
937 pentandra Cav.	pentandrous	O cu * jn.jl G	S. Europe 1820. S	r.m Cav. ic. 2.148.3	
100. HIPPOCRATES L.	HIPPOCHATEA.	(Hippocrates, father of phys.) HippocratedceA*.	5. — 24.		
938 obcordata Lam.	obcordate	A. CU or 10 ...	G.w	W. Indies 1819. C p.l	Lam. U. 1. 28.1
scandens Jac.					
939 ovata Lam.	ovate	4. CD or 10 ...	G w	& Amer. 1793. C p.l	Lam. U. 1. A S
voldbilis L.					
940 indica W.	Indian	10	G.w	E. Indies 1818. E. Rox. cor. 2.130	
941 obtusifolia Rox.	obtuse-leaved	Qor 14	G.w	E. Indies 1818. E. Rox. cor. a 205	
942 arborea Rex.	tree	20	G.w	E. Indies 182a E. Rox. cor. 2.130	
101. TONSELLA Vahl.	TONSBLLA.	(Ravoua-tontelle, a Caribe name.) Hippocrateatee. 2.—10.			
943 scandens Vahl	climbing	(LCD or 6 aus.s G	Guiana 1824. C s.l	Aub.guLL. 10	
944 pyriformis H. T.	pear-shaped	Q or 5 ... G	Guiana 1825. C 31		
102. A'NTHODON B. S.P.	ANTHODON.	{Anthos, a flower, oifofi, a tooth.) Hippocratealce*. 2.—9.			
945 paniculatum Mart.	panicled	* C or 12 ■■■ Y.u	Rio.Jan. 1888. C p.l		
elliptic		* C or 12 ■■■ Y.G	Rio Jan. 1818. C p.l		
103 JO/HNW ROX.	JOHNIA.	{Rev. Dr. John, of Tranquebar.) Hippocrateetica. 2.			
947 salacioides Rot.	Salacia-like	Dor 4 ... Y.G	E. Indies 1822. C p.l		
948 coramdeliana Aojr.	Coramdel	Dor or 10 ■■■ Y.a	K Indies 1820. C p.l		
104. O'LAX L.	OLAX.	{Olax, a furrow; the flower.) Olacinete. 2.^3.			
949 scandens Rox.	climbing	L D or 8 year W	E. Indies 1820. C l.p Rox. cor. 2.		
950 imbricate Rox.	imbricated	i. CZ or 8 year W	E. Indies 1820. C l.p		

					Oiacinete. 1.—3
					Trebintdceee. 2.
					Terebintdcete. 3.—4.
					..8. C.p.l. SI.jam.2.222.1
WW dentata Jac. S56ilicif6haStuz.	tooth-leaved holly-leaved	• tm 30 jl • tm 15 R			C.p.l. Lam. il. 27 W. Indies LTH. C.p.l. Jac. am. 13.173.4
					Caribbee Is. 1789. C.p.l. Plum. ic. 118.1
					Legumindsee. 1.
957 bifolia Aub.	VOUAPA.	(ACaribe name.)			Guiana 1823. C.s.l. Aub. gui. 1.1
■109. OUTEA Aub.	JOUTAY.				Legnmindsee. 2.—3.
■58 bijuga Dc.	two-balled	(A Caribe name.)			E. Indies 1823. C.s.l. Lin.tr. 12.17
Macrolobium bijugum Cab.					
959 guanensis Jfffr.	Guuiua				
110. XYRTS L.	XYRIS.	1 O or 20			
960 operculata Lab.	covered	(Xyros, acute.)			Guiana 1825. C.s.l. Aub. gui. 1.9
961 brevifolia Afr.	short-leaved	£ 123 pr 1 jnjl Y			JUMcete. 7.*—27.
962 Ue's R.Br.	smooth	£ A pr jn.au Y			fc. S. W. 1804. S bog Bot mag. 1158
9G5bractea"ta2J?r.	bracted	£ E2 pr li jn.au Y			Carolina 1812. S bog
964 juncea J.J. tfr.	rush-like	£ iAJ pr I jn.au Y			N. Holl. 1819. S bog
W5 gr&chis R. Br.	slender	£ LAI pr 1 jn.au Y			N. Holl. 1825. S bog
•66 americana fw6.	American	£ LAI pr i jn.au Y			N. Holl. 1821. S bog
		£ A pr i jls B			Guiana 1825. D bog Aub. gui. 1.14
111. CALLISIA. L.	CALLISIA.	(Kalo. pretty.)			Commelinea. 1.—3.
»7 repens L	creeping	**E3pr ijnjl B			W. Indies 1776. R Bp Jac. am. 11.11
»» angustifolia L.	narrow-leaved	4 j- au B			CaiMina 1824. D p.l
»69 dianthifolia Dec.	pink-leaved	jnjl B			Guiana 1823. R s.p Red. lil. 7.990
vJO cajemnenu Sic/t.	Cayenne	jnjl B		 1818. D ps
971 dubia Dec.	doubtful	1 my.jl B			N. Amer. 1820. D ps Red. m. 6.359
yj2 hirtilla Vahl	hairy	B			Virginia 1779. R s.p
973 virginica L.	Virginian	B			Caracas 1804. R s.p
974 mollis Jac.		B			Mexico 1820. R s.p Plk.al. 135.174.4
975 obliqua Vahl		B			Lima 1813. D ps
976 tuberosa L.		B			Bengal 1794. R s.p Bot rep. 399
£77 cicestis W.		B			P.B. 1824. R s.p Bot mag. 1695
978 fa*ciculata JZ %P.	fascicled	B			America 1732. R s.p Red. m. 107.6
979 bengalensis Forsk.	Bengal	B			China 1818. D ps Ksm. 888
980 parviflora Zee	small-flowered	B			Brazil 182J. C. Bot. mag. 2644
981 communitis L.	common	B			Caracas 1806. R s.p Jac. ic. 2.294
Jg Pol'gama Ao/A	polygamous	B			America 1732. D r.m. W.h.b. 2.87
?J3deficiens/fer?	deficient	B			Trinidad 1820. D ps W.h.b. 2.87
JJ longicafis Jsic.	long-stalked	B			
985 caroliniana Walt.	Carolina	B			
986 p511hd H*A	pale blue	B			
	upright	B			
988 carpensis H. & B.	Caripe	Ift jnjl B			Trinidad 1826. D ps
989 cyanea R. Br.	bright-blue	1 jlau B			N. Holl. 1820. R s.l
990 africana L.	African	1 my.o B			CG.H. 1759. R r.m. Bot mag. 1431
					Commelinee. 11.—49.
W» uiuira n. JUT.	two-flowered				
992 ambigua Beauv.	ambiguous	£ • or V			N. Holl. 1820. R co
993 sinica Ker	Chinese	£ IA) or 1 my.jn V			S. Leone 1822. D r.m Beau. ow. 15
994 acuminata R. Br.	acuminate	j^ [23 or I jl.au P.B.			China 1820. I) r.m Bot reg. 659
»95 nudiculis R. Br.	naked-stemmed	JU E] or B			N. Holl. 1822. D s.l
996 Bipartita L.	8'piral	%! 1AI or ft jl.au B			E. Indies 1818. D s.l R.maL9.63
997 affinis R. Br.	long-leaved	Xr IA) or 1 jl.au B			E. Indies 1783. I) s.l
998 longiflora Hook.	naked-flowered	£ (A) or 1 au B			N. Holl. 1820. D e.p
999 nudinaria L.	serrulated	£ OI or ft jn.jl B			Mozambi. 182*. C p.l Hook. ex. fl. 004
1000 serrulata Vahl	equinoxial	ft in.jl B			E. Indies 1824. D ps l'luk. al. 27.4
1001 sequinotidic Beauv.		ft in.jl B			Trinidad 1824. D ps
		Guinea 1820. D ps Beau. ow. 15			
114. ORTEGII Jii L.	ORTEGIA.	(C. O. de Ortega, a Spanish botanist.)			CaryophijletB. 2.
1002 hispanica L.	Spanish	A cu I jn.jl Ap	Spain 1768. D l.p Cav. ic. 1.47		
1003 dichotoma All.	forked	A cu Jau.s Ap	Italy 1781. D i.p Al.tau.3.4.1		
					; joints.)
Pall.	recurved	O cu			S. s.l. Jac.au. 4. S65
Pall.	Siberian	O cu			
i009 inalacoph?Uum4be6.	opposite-leaved	O cu			
	hard-seeded	O cu			
	soft-leaved	O cu			
in***11. CROCUS L.	Caucua.	(Krokin, Chaldee.)			
1010 vernus L. B.		A or } fap P	England 1820. S s.l		
2 neapolitanus Ten.	Neapolitan	A or if.ap P.B.	Naples 1826. S s.l		
3 obovatus? &	obovate	A or fm P.S.	Europe 1826. S s.l	Gm. si. 23.1	
1011 albifl6rus Xrt.	white-flowered	A or fm P.W.	Austria 1629. O co		
1012 minimus Red	smallest	A or fm P	S. Europe 1629. O co	Bot. mag. 8G0	
1013 versicolor H. K.	party-colored	A or fm P	Crimea 1629. O co	Bot. mag. 2240	
1014 biflorus Mil.	two-flowered	A or fm P	W. Crimea 1629. O co		
1015 pusillus Ten.	little	A or if.mr P.Y.	Naples 1824. O co	Bot. cab. 1454	
1016 susi?nus Kit.	Suaian	A or * fm Y	Turkey 1G05. O ro	Bot. mag. 652	
1017 reticulatus Stev.	netted	A or fm B	Crimea 1820. O co		
1018 striatus lit.	channelled	A or fm W 1820. O co		
1019 sulphureus H. K.	Caucua.	A or i fm Y	S. Europe 1629. O co	Bot. mag. 1110	
		A or i frar P.Y.	Crimea 1629. O co	Bot. mag. 845	
1020 luteus Lam,	sulphur-cofomi	A or fm P.Y.	Naples 1824. O co	Bot. cab. 1454	
1081 lacenafl6rus Sal.	polyantha yellow	A or fm Y	Turkey 1629. O co	Bot. mag. 45	
	bottle-flowered	A or if.mr D.Y.	Greece 1820. O co	Fl. gr. 1.35	
	yellow	A or Cmr P.Y.	Greece 1820. O co	Bot. mag. 1111	

123. TRITOTJIA Ker	THITONIA. (<i>Trito</i> ηn, a weathercock; in allusion to the stamens.) <i>Irideaf.</i> 22.—2a
1092 crispa Ker	<i>curivd-leaved</i> tf LAI or I ap my Y C. O. H. 1787. O s.p.l Bot. mag. 678
DWJ aiigozanthaflura5w./Anigosan.-fl. tf JSUR	iili,jl Y C. G. H. 1825. O s.p.l Sw. fl. gar. ic.
1094 vlnidis Ker	<i>green-flouuered</i> tf iAlor . C. G. H. 17H8. O s.p.l Bot. mag. 1275
1095 rosen Ker	<i>rosy</i> tf iAlor . C. G. H. 179i. O s.p.l Hot. mag. 618
DSW capensis Xier	Cape tf iAlor au.o W C. G. H. 1811. O s.p.l Bot. mag. 1531
1097 lonKiflora Jfc-r	<i>long-flowered</i> tf iAlor ap.jn W C. G. H. 1774. O s.p.l Bot. mag. 216
1098 tenuiflora Fahl	<i>slender.flwd</i> tf iAlor ap.jn Y C. G. H. 1811. O s.p.l B.m.l502.f.maj.
1099 Luncolor Su>.	<i>self-colored</i> tf iAlor ap.jn Y C. G. H. 1811. O s.p.l B.m.l502.f.min.
1100 Uroch6nris4L .V	De la Roche tf iAlor au Y C. G. H. 1806. O s.p.l Bot. mag. 150J
1101 pthlddjfr	<i>pate-flowered</i> tf LAI or au W C. G. H. 1825. O s.p.l
1102 pectinata Ker	<i>pectinated</i> tf iAI or myjn Y C. G. H. 1825. O s.p.l
ImUK	<i>lined</i> tf iAI or my Va C. G. H. 1774. O s.p.l Bot. mag. «7
Ker	<i>striated</i> tf iAI or myjn B C. G. H. 1825. O s.p.l Jac. ic. 22GD
1105 sejurgera Ker	<i>axe-bearing</i> tf LAI or i " l v . C. G. H. 1774. O s.p.l Bot. mag. oH3
IIOOflavaAVr	<i>yellow</i> tf iAlor f.mr Y Ru C. G. H. 1774. O s.p.l Bot. reg. 747
1107 squaTula tor	<i>squalid</i> tf iAlor f.mr Ru C. G. H. 1801. O s.p.l BoL mag. 704
1108 fenestrata AVr	<i>windowed</i> tf iAlor ful C. G. H. 1758. O s.p.l Bot. mag. 184
1109 crocata Ker	<i>saffroned</i> tf iAlor ful C. G. H. 1795. O s.p.l Bot. mag. 609
III Odencsta Ker	<i>blasted</i> tf iAlor my ful C. G. H. 1825. O s.p.l Red. III. 3. 124
1111 miniata Ker	<i>vermilion-/w<2</i> tf iAlor au ful C. G. H. 1815. O s.p.l Bot. reg. 135
1112 xanthrispila Ker	<i>ycl low-spotted</i> tf iAI or I my,jl RY C. G. H. 1815. O s.p.l Bot. reg. 135
1113 refracta Ker	<i>refracted</i> tf iAlor myjn Y
• 124. VTATSfyVIA Ker	WATBONIA. (IV. Watson, a London apothecary.) <i>Iridea.</i> 13.—
UMspicelUAtfer	<i>spiked</i> tf iAlor my Pk C. G. H. 1791. O s.p.l Bot. mag. 523
1115 jDlantaginosa Ker	<i>Plantain</i> tf iAlor 2-jnl W C. G. H. 1774. O s.p.l Bot. mag. 525
1116 punctata/i.7i/-.	<i>dotted-flowered</i> tf iAlor 1 ap.my Pk C. G. H. 1800. O s.p.l Bot. rep. 177
1117 roseo-alba Ker	<i>red and white</i> tf iAlor 1 jlau Pk C. G. H. ... O s.p.l Bot. mag. 537
2 variegata	<i>variegated</i> tf iAlor 1 jlau Va C. G. H. ... O fl.p.l Bot. mag. 1193
1118 marginata Ker	<i>margined</i> tf LAI or 3 jn Pk C. G. H. 1774. O s.p.l Bot. mag. 608
2 inimur	<i>leaser</i> tf iAlor 3 o Pk C. G. H. 1812. O 8.ixl Bot. uwg. 1530
1119 strictiHora ATr	<i>strict-flowered</i> tf iAlor 1 jn K C. U. H. 1810. O s.p.l Bot. mag. 1406
1220 rose Ker	<i>rosy</i> tf iAlor 2 jlau P C. G. H. 1803. O s.p.l Bot. mag. 1U72
1121 brevifolia Ker	<i>short-leaved</i> tf iAlor 1 my Pk C. G. H. 1794. O s.p.l Bot. mag. HO1
1122 iridifolia Jac.	<i>Iris-leaved</i> tf iAlor 2 my F C. G. H. 1795. O s.p.l Jac. ic. 2.234
II2 Kjulda&cZ.	<i>bright</i> tf iAlor 4 my R C. G. H. 1795. O s.p.l Bot. mag. NO
1123 Menana Ker	<i>Menan's</i> tf iAlor 1 my,jn F C. G. H. 1750. O s.p.l Bot. m. 418.1194
1125 hifimilis Ker	<i>humble</i> tf iAlor 2 my,jl La C. G. H. 1754. O s.p.l Bot. m. 631.1193
1126 al caldi Jfe	<i>Aletna-like</i> tf 'Al or my,jl S C. G. H. 1774. O s.p.l Bot. mag. 441
	<i>vanegetad-^wd</i> tf iAlor 1 *my,jl Va C. G. H. 1774. O s.p.l Bot. mag. 533
125. BABIA'NA Ker	BABIANA. (<i>Babiancr.</i>) Dut.; the roots the food of baboons.) <i>Irideee.</i> 17.
1127 Thunblrji Ker	Thunberg's tf iAlor 1 ap P C. G. H. 1774. O s.p.l
1128 ringons Ker	<i>guining-flowered</i> tf iAlor 1 myjn P T. G. H. 1752. O e p Bot cab. 1006
1129 tubiflora Ker	<i>tube-flowered</i> tf iAlor 1 ijin Y.R C. G. H. 1774. O sp Bot. mag. 847
Ilfti tubata IF.	<i>long-lubei</i> tf iAlor 1 jn W.a C. G. H. 1774. O s.p Bot. mag. 680
1131 spathacea Ker	<i>heathy</i> tf iAlor 1 jii LB C. G. H. 1801. O s.p Bot. mag. 638
1132 *ambucina Air	<i>Hadar-scented</i> tf iAJ °f L ap.my B C. G. H. 1799. O s.p Bot. mag. 1019
liajdlsctica Ker	<i>two-ranked</i> tf iAlor 1 *jh,jl B C. Li. H. 1774. O s.p Bot. nug. 626
1134 plictata AVR	<i>folded</i> tf iAlor 1 h'iny,jn Pu C. G. H. 1774. O s.p Bot. mag. 576
1135 >strkt:aA'Dr	<i>strict</i> g iAlor 1 uiy,jn B.w V. G. H. 1757. O s.p Bot. mag. 621
1136 angust i folia A'er	<i>narrow-leaved</i> t iAlor 1 my,jn Va C. G. H. 1757. O s.p.l Bot. mag. 637
'xia striata A 3/.	
1137 tenuiflora Swt.	<i>slender-flwd</i> tf iAlor \ myjn P C. G. H. 1825. O s.p.l Sw. fl. gar. ic.
1138 sulphiroja Ker	nu\hur-flu>d tf iAlor i myjn Y C. G. H. 1715. O s.p Bot. mag. 1053
1139 mucronata Ker	<i>mucromate</i> tf iAI or 4 myjn P C. G. H. 1885. O s.p.l Jac. ic. 2. 253
1140 villosa Ker	<i>hairy</i> tf iAI or *au Dn C. G. H. 1778. O s.p Bot. mag. 583
1141 obiusfolia Ker	<i>obtuse-leaved</i> tf iAlor L my,jn B C. G. H. 1825. O »-p.Uac. ic. 2.284
J'xia villosa Jac. not H. K.	
1142 purpi rea Ker	<i>purple</i> tf iAlor < myjn P C. G. H. 1806. O s.p.l Bot. mag. 1052
1143 rdbro-cyanea Ker	red and blue tf iAlor ap.jn B.B C. G. H. 1794. O s.p Bot. mag. 410
1144 n'ina Spr.	<i>dwarf</i> tf iAlor ap.my B C. G. H. 1807. O s.p.l And. rep. 137
Cradolous ninus And. Bep.	
* 126 LAPEYROU'S/itf Ker	LAPEYROISIA. (La Peyrou's, the French navigator.) <i>Iridea.</i> 8.
1145 rorymboea Ker	<i>corymbose</i> tf Alor i my,jn U C. li. H. 17?I. O s.p Bot. mag. 595
1146 Cfalcata AVR	<i>falcate</i> tf tAlor myjn B C. G. H. 1b2J. O s.p.l Thun. ir.4.1.3
1147 fasciculata Ker	<i>fasciculate</i> tf iAlor i my,jn W C. G. U. 1825. O s.p.l Jac. ic. 2. 291
Galaxia plicata Jac.	
1148 fisMfolia Ker	<i>cloft-kenved</i> tf iAlor au.8 V C. G. H. 1809. O a p Bot. mag. 1246
1149 Janeps Ker	<i>two-edged</i> tf ^Jpr i b li C. G. H. 1824. II s.l Sw. fl. gar. 143
1150 aculata Swt.	<i>prickly</i> tf iAlor i my,jl B.Y C. G. H. 1825. O s.p.l Sw. fl. gar. icin.
1151 Ailemfdes Ker	<i>Catchfly-like</i> tf iAJor Jmy,jl V C. G. H. 1822. O s.p.l Jac. ic. 2.270
1152 Fabricic Ker	<i>Fabricius's</i> tf iAlor * my,jl B C. G. H. 1825. O s.p.l
*** MELASPHIE'RULA Ker	MELASPH.CBIXA. (>/fff<, black, juAfira, globule; bulbs.) <i>Irideaf.</i> 3.
1153 ^raminca Dec.	<i>Grass-leaved</i> tf iAI - 1 ap.au G C. G. H. 1787. O ap Red. hi. 163
1154 Intermtdia Swt.	<i>Intermediate</i> tf iAI or 1 ap.au Y.c; C. G. H. 1787. O s.p.l Bot. mag. 615
Aframica B. R. not	<i>Bed.</i>
1155 Sindil6ha Dec.	<i>Iris-leaved</i> tf iAJ or 1 ap.au Y o C. G. H. 1787. O s.p Jac. ic. 2.236.
• 128. GLADI'OLUS L.	Conv FLAG. (<i>Gladius</i> a sword; the leaves.) <i>Iridea.</i> 37.-46.
1 lo» Wzspnus Thuu.	Watson's tf iAlor 1 faid D.R C. G. H. 1791. O s.p.l Botmag.4RUe9
" 2 variegatus	<i>variegated</i> tf yAl or 1 m.ap R.w C. G. H. 1801. O s.p.l Bot mag. 5G9
" 1157 viperitu* Ker	<i>viper</i> tf iAlor i ap my G.Y C. G. H. 1787. O s.p.l Bot mag. 688
1158a)atus L.	<i>Vfinged-flwd</i> tf iAlor ■ myjn S.v C. G. H. 1795. O s.p.l Bot mag. 586
2 algooensis Herb.	<i>Algoa May</i> tf iAlor \$ jl O C. G. H. 1824. R-lp Bot. mag. 2iOS
1159 namuq6nsis Ker	<i>Namaqua</i> tf iAlor 5 myjn O.G C. G. H. 1800. O fl.p.l Bot. mag. 592
1160 permeabilis Lar.	<i>permeable</i> tf iAI or f myjn C. G. H. 1825. O s.p.l Lar. dis. 2
1161 brevifilius Jac.	<i>short-leaved</i> tf iAI or 1^ Ujn O £ 2- 5 ! ? ! • 2 s*p Bot.mag.727.992
1162 hinutus L.	<i>hairy</i> tf iAlor i japjn Pk C. G. H. 1795. O s.p.l Bot. mag. 574
1163 versicolor Ker	<i>various-colored</i> tf iAl or U myjn Pk C. G. H. 1794. O s.p.l Bot. mag. <1042
2 binervis B. M.	<i>two-nerved</i> tf iAl or 1 myjn Br C. G. H. 1806. O s.p.l Bot. mag. >1042
3 tentior Ker	<i>slenderer</i> tf iAlor 1 myjn Pk C. G. H. 1779. O s.p.l Bot. mag. 55d
1164 ednlis Ker	<i>eatable</i> tf iAI cul myjn Va C. G. H. 1816. O s.p.l Bot reg. 169
Ilias hastatus Tfit/n.	<i>hastate</i> tf iA ■ l" ap.my W C. G. H. 1810. O s p 1 Bot mag. 1564

1166 tristis L.	sad	■ iAI or 1 myjn	Br.Y C. G. H.	1745. O * .p.l Bot. mag. 272
1167 cōncolor Sal.	self-colored	* • Alor 1 myjn	Y C. G. H.	1790. O s.p.l Par. Ion. 8
1168 hyahnus Jac.	glassy	■ Alor 1 myjn	Y.R C. G. H.	1825. O fl.p.l Jac. ic. 2.242
1169 tenellus Jac.	tender	■ Alor 1 myjn	Va.Y C. G. H.	1825. O s.p.l Jac. ic. 2.248
1170 Colvillia Swt.	Colville's	! IAIpr 1 injl	S.Y hybrid	1824. O l.Sw. fl. gar. 155
1171 trichonemifolius Ker	Trichonema.lv	■ Alor 1 injl	Y C. G. H.	1810. O 8.p.l Bot. mag. 1483
1172 gracilis Jac.	small	■ Alor 1 mr.ap	B.w C. G. H.	1800. O s.p.l Bot. mag. 562
1173 debilis Sims	weak	5 LAJ pr 1 myjn	W C. G. H.	... O p.l Bot. mag. 2585
1174 recbrvus H. K.	recurved	■ Alor S ap.my	Str C. G. H.	1758. O s.p.l Bot mug. 578
1175 carnēus Jac.	fleath-colored	■ Alor 1 myjn	F C. G. H.	179a O s.p.l Bot mag. 591
1176 cuspidatus L.	pointed	■ IAI or 1 ap.my	W.BrC. G. H.	1795. O s.p.l Bot. mag. 582
1177 blandus H. K.	bland	■ IAI or 1 injl	L.P C. G. H.	1774. O s.p.l Bot. mag. 625
1178 dibidus Vahl	whitish	■ Alor 1 myjn	C. G. H.	1774. O B.pl Bot. mag. 648
2 pictus B.M.	painted	■ Alor 1 myjn	R.W C. G. H.	1794. O s.p.l Bot. mag. 1665
1179 trimaculatus Vahl	three-spotted	■ IAI or 1 myjn	R.W C. G. H.	1794. O s.p.l
1180 campanulatus B. R.	beW-flowered	■ Alor 1 myjn	L.P C. G. H.	1794. O s.p.l Bot. rep. 188
1181 angustus L.	nanovr-icaved	■ Alor 1 myjn	L.Y C. G. H.	1757. O s.p.l Bot mag. 602
1182 involutus Lar.	involute	■ IAI or 1 myjn	Pk C. G. H.	1757. O s.p.l Lar. dis. 2. 3
1183 flexuosus Thun.	flexuous	■ Alor 1 myjn	O e. G. H.	1825. O s.p.l Thun. ir. 1. 1
1184 undulatus Jac.	xvæle-Jtowered	■ IAI or 1 ap.my	Pk e. G. H.	1760. O s.p.l Jac. ic. 2.242
2 pálodus	pale	■ Alor 1 ap.my	P.Pk C. G. H.	1760. O s.p.l Bot. mag. 647
3 carnēus	flesh-colored	■ Alor 1 ap.my	F C. G. H.	1760. O s.p.l Bot. mag. 538
1185 floribndus Jac.	bundle-flwd	■ Alor 1 myjn	Ci C. G. H.	1788. O s.p.l Bot mag. 610
1186 Mfllen Ker	Miller's	■ IAI or 1 ap.my	V C. G. H.	1751. O s.p.l Bot mag. 632
1187 cardinalis B. M.	cardinal	■ Alor 1 jl.au	D.B C. G. H.	1789. O s.p.l Bot mag. 135
1188 byzantinus Mil.	Byzantine	Q A or 5 H.	R Turkey	W29. O s.p. Bot. mag. 874
1189 communis L.	common	■ A or 1 injl	II S. Europe	1506. O s.p.l Hot. mag. 86
2 c&rneus	flesh-colored	■ A or 1 injl	F S. Europe	1596. O r.m Bot mag. 1575
1190 tcnuis Bicb.	slender	■ A or 1 myjn	R Tauria	1823. O co
1191 imbric&tus I.	imbricated	■ A or 1 myjn	K Russia	1820. O lp
1192 segetum Ker	corn field	■ Alor 1 injl	P S. Europe	1596. O B.pl Bot. mag. 719

129. SYNNOT/!* Swt	SYNNOTIA.	(W. <i>Synnot</i> , who collected many plants at the Cape)	Iridee. 3.
1193 variegata Swt.	variegated	tf iAI or f ap.jn	Va C. G. H. 1825. O s.p.l Sw. fl. gar. 150
1194 bfcolor Swt.	two-colored	tf LAJ or \$ mr.ap	B.v C. G. H. 1786. O s.p.l Bot mag. 548
/xiabicolor AA/I	Sparaxiabicolor Avr.	(iliadiolus bicolor W.	
1195 galeatus Swt.	helmcted	tf iAI or 1 mr.ap	Y C. G. H. 1825. O s.p.l Jac. ic. 2. 258
		Gladiolus galcatus Jac.	

* 130. ANOMATHE'CA Ker	ANOMATHECA.	(Anomos, singular, theca, a capsule.)	Iridea. 1.
1196 juncea Ker	rushy	g 1Alor ap.my	Li C. G. H. 1791. O B.pl Bot mag. 606

131. ANTHOLY'ZAL.	ANTHOLYZA.	(Anthos, fl., lyssa, rage; fl. like a mouth ready to bite.)	Iridea>. 3—4.
1197 sthiopica L.	Ethiopian	tf iAJ or 3 myjn	S.o C. G. H. 1759. O a.p.l Bot. mag. 561
1198 prs&ita Dec.	very tall	tf iAI or 2 ja.f	O C. G. H. ... O s.p.l Bot mag. 1172
ethiopica fi vitfgera B. M.			^
1199 montna Ker	mountain	tf j^d fra 1 jn	Br C. G. H. 1822. R s.p Bot cab. 1022
132. ANISA'NTHUS Swt.	ANISANTHUS.	(Anisns, unequal, anthos, a flower.)	IrtletE. 3.
1200 splēndens Swt.	splendid	tf JAJ ^r li myjn	S C. G. H. 1805. O s.p.l Sw. fl. gar. ic.
1201 Cunōnvi Swt.	Cunon's	tf iAJ or 2 myjn	S C. G. H. 1756. O s.p.l Bot. mag. 843
AntholyzaCunon/rt B.M.	GladiolusCunonia H. K.		
1202 quadianguliris Swt.	quadrangular	tf iAJ ^f mr.ap	Y C. G. H. 17C0. O s.p.l Bot mag. 567
Gladiolus quadrangularis 2f.Af abbreviatus And. rep.			136

133. XIPHI'DIUM Aubl.	XIPHIDIUM.	(Xiphos, a sword; the leaves.)	Ilcomodordcece. 2.
1203 album W.	white	£ [3]or \ W	W W. Indies 1787. U s.p
1204 caerGleum Aubl.	blue	£ EJ or \ B	Guiana 1793. II s.p Aub.gui. 1.11

134. LEPTA'NTHUS Mx.	LEFTANTHUS.	(Leptos, slender, anthos, a flower.)	Pontedrenee. 1.
1205 gnmlneu Mx.	grassy	■ A 1 jn.au	Y N. Amer. 182a D aq Hook. ex. fl. 94
135. HETERANTHE'RA Beauv.	HETERANTHERA.	(Heteros, variable, aner, anther.)	Pontedrenee. 3.-7.
1206 reniformis % £/P.	kidney-leaved	^ (£) or fltl.au	B S. Amer. 1824 Sk aq Fl. per. 1.71
1207 aciita JE.	acute kidn.-lyd	^ LAJ or flt jnjl	W Virginia 1812. Skaq Beauv.in ac. am.
Leptanthus reniformis Mx. not R. %P.			[4. 173. ic]
1208 limsda Vahl	bog	A A or fltl.au	B N. Amer. 1822. D m.8 Mic.am.5.1

136. WACHENDCKRF/^ L.	WACHENDORFIA.	(E. J. Wachendorf, a Dutch bot.)	Hamodoraee. 8.—a
1209 thysiflora L.	thyrsflowered	£ LAJ or 2 myjn	Y C. G. H. 1759. D r.m Bot. mag. 1060
1210 paniculata Thun.	panicled	£ iAI or 2 f Y	C. G. H. 1700. D r.m Bot mag. 616
1211 Hibberti Swt.	Hibbert's	^ Alor 2 fjn Y	C. G. H. 1823. D r.m Bot mag. 2610
paniculata/3 B.M.			
1212 hirsuta Ker	hairy	■ • Alor Ujn V	C. G. H. 1687. D r.m Bot mag. 614
1213 brevifolia Ker	short-leaved	■ Alor 1 mr.ap P	C. G. H. 1795. D r.m Bot. mag. 1166
1214 Brcynia/R. & S.	Brcynius's	■ Alor 1 mr.ap	C. G. H. 1825. D r.m Brey.c.37
1215 graminea Thun.	grass-leaved	■ IAI or 1 jn V	C. G. H. *** D r.m
1216 tencilla Thun.	tender	■ Alor 1 jn Y	C. G. H. 1816. D r.m

137. HJEMODO'RUM Sm.	HAMODORUM.	(Haima, blood, doron, gift; flower.)	Hamodordceg*. 2.-6.
1217 planifolium R. Br.	flat-leaved	£ 1 or 1} jl.n O	N. S. W. 1810. S s.p Bot. mag. 1610
1218 teretifolium R. Br.	round-leaved	£ \ 1 jn O	N. Holl. 1822. D s.p

138. ARFSTEA L.	ARISTEA.	(Aristea, a point, or beard; the leaves.)	Iridece. 5.-7.
1219 cyanea//.A\	bright blue	£ LAJor i apjn B	C. G. H. 1759. S s.p Bot mag. 458
1220 capitata H. K.	headed	£ LAI or 3 jl.au B	C. G. H. 1790. C s.p Bot mag. 605
1221 spiralis Vahl	spiral-Jfcnueraf	£ IAJ or 1 ap.my Pa.B	C. G. H. 1795. C s.p Bot mag. 520
1222 melaletica Ker	black and white	IAJor 1 myjn Pa.B	C. G. H. 1796. C s.p Bot mag. 1277
1223 pusilla Ker	little	£ LAJ or i jnjl B	C.G.H. 180& C s.p Bot mag. 1231

139. DI'LATRIS L.	Di LATHIS.	(Dis, two, later, a tile.)	Heemodordcece. 3.—
lgJymbosarA ■	corymbose	■ iAI or 1 my P.r C. G. H.	179a S s.p Ex. bot L16
1225 visc6sa Thun.	clammy	■ iAI or 1 my C. G. H.	1795. S s.p Lam. U. 34
1226 paniculata Thun.	panicled	■ iAI or 1 my,jl B	C. G. H. 1825. S s.p
1227 tinctoria EU.	dyer's	£ iAI dy 1^jl.au Pk	N. Amer. 1812. S s.p Mic. am. 1.4

IXlatris Heritiera Pcrs. EMLatris tinctoria Ph.

1305 tuberfa <i>L.</i>	tuberose	tf A or	mr.ap	G.B	Levantf	1597.	O s.p	Bot. mag. 531	
1306 alfita <i>Poir.</i>	winged	tf A or	B	Algiers	1801.	O co	Desf.at. 1.6		
1307 AiphiumA.	Xiphia	tf A or	B.Y	Spain	1596.	O co	Bot. mag. 686		
1308 Jiphiodes <i>Ehrh.</i>	Xiphium-Hke	tf A or	B.Y	vSpain	1571.	O co	Bot. mag. 687		
1309 lusilánica <i>Ker</i>	Portuguese	tf A or	2 ap	B	Portugal	1796.	O co	Bot. mag. 679	
1310 tenuiloha <i>Pall.</i>	slender-leaved	tf A or	limy	L.B	Jfeuria	1796.	D co	Pa), it. 3.2	
1311 pē>aicaL.	Persian	tf A or	mr.	B.Y	Persia	1629.	O co	Bot. mag. 1	
1312 reticulata <i>Bieb.</i>	netted	tf A or	mr.ap	B	Iberia	1821.	R co		
1313 caucayica <i>Hqfm.</i>	Caucasian	tf A or	mr	Y	Caucasus	1821.	It co		
•U3. MOUE'AL									
1314 anftista <i>Ker</i>	MORJSA.	(R. Moore, a botanist of Shrewsbury.)			Iridett.	20.—93.			
1315 edulis <i>Ker</i>	narrow Jeaved	tf IAJ or	f my.jn	Ful	C. G. H.	1740.	D s p	Bot. mag. 1276	
1316 odifra <i>Sal.</i>	eatable	tf TAI cul	4 my.jn	Li	C. G. H.	1792.	I s.p	B.mag.hu 1-238	
1317 loiiKiftia <i>Andr.</i>	sweet-scented	tf IAJ or	2 my.jn	Li	C. G. H.	1792.	D s p	Par. Ion. 10	
1318 longiflora <i>ILK.</i>	long-leaved	tf iAJ or	3 my.jn	Y	C. G. H.	1803.	D s p	And. rep. 45	
1319 setifera <i>Ker</i>	long-flowcred	tf iAJ or	my.jn	Y	C. G. H.	1801.	D s p	Bot. mag. 712	
1320 tristis <i>Ker</i>	setaceous	tf iAJ or	i jn jl	Y	C. G. H.	182/5.	I s.p	Thun. ir. 1.1	
1321 crispa <i>Ker</i>	curled	tf LAJ or	my.jn	B	C. G. H.	1718.	I s.p	Bot. mag. 577	
1322 bituminosa <i>Ker</i>	bituminous	tf iAJ or	j ap my	Y	C. G. H.	18(3).	D s p	Bot. mag. 1284	
1323 polyntachya <i>Ker</i>	m.iny-spiked	tf iAJ or	1 jn jl	Y	C. G. H.	1825.	I up		
1324 viscana <i>Ker</i>	vi.cid	tf iAJ or	1 jn	Li	C. G. H.	INCH.	D s p	Bot. mag. 587	
1325 ramosa <i>Ker</i>	ramose	tf LAJ or	3 my.jn	Y	C. G. H.	17H.	I s.p	Bot. mrig. 771	
1326 plumaria <i>Ker</i>	feathered	tf LAJ or	1 my.jn	Y	C. G. H.	18*5.	I s.p		
1327 siBryinchium <i>Ker</i>	SUyrinclusm	tf A cul	\ myjn	B	S. Europe	1597.	I Bp	Bot. mag. 1407	
1328 Tenorca/tta <i>Swt</i>	Tenorc's	tf A or	1 ap.my	P	Naples	1624.	K l.p	Sw.fl.gar.110	
1329 mindta <i>Ker</i>	small	tf iAJ or	\ jnjl	B	C. G. H.	1825.	I s.p		
1330 papilionacea <i>Ker</i>	butterfly	tf LAJ or	\ myjn	Va	C. G. H.	1795.	U s.p	Bot. mag. 750	
1331 cihāta // K.	fingud-leaved	tf iAJ or	\ apjn	P	C. G. H.	1587.	D e.p	Bot. mag. 1061	
1332 barbigra <i>Sal.</i>	bearded	tf iAJ or	i ap.my	Y	C. G. H.	1587.	D s.p	Bot. mag. 1012	
* 144. M A'R CA Schreb. MA MCA. (Maraino, to become flaccid; the flowers.)	(Maraino,				I idea:	7.			
1333 Northiana <i>Ker</i>	North's	■ 123 or	4 ap.au	YB	Brazil	1789.	D s.p	Bot. mag. 654	
1334 cerifca <i>Ker</i>	blue	■ iAJ or	2 myjn	B	Brazil	1818.	I co	Bot. nff. 713	
1335 Sablni Lindl.	Capt. Sabine's	■ T	2 ap	P	St. Thorn.	1822.	R pI	Hort. tr. fi. 1	
1336 martinic6nsia <i>Ker.</i>	Martinico	tf [23 or	2 jn	Y	Martinico	1782.	I s.p	Bot. mag. 416	
1337 hfmuiilis <i>R&S.</i>	low	tf iAJ or	J "my.jl	Y	Brazil	1823.	D p.l	Bot. cab. 685	
1338 palud6sa <i>W.</i>	marsh	A (23 or	1 jl.au	W	Ciukiiia	1792.	Sk s p hot. mig.	646	
1339 semi-aperta <i>Lod.</i>	half-open	jf LAJ or	1J my.jn	Y	Brazils	18*0.	D co	Bot. cab. 685	
• 145. SISYRINCIUM//. SISYRINCHIUM M. (Sys ig or hog, rhygchos, a snout)					I? idea:	14.—23.			
1340 plicatum <i>Spr.</i>	folded	■ (23 or	2 f.mr	W	M Indies	1779.	Sk s.p	Bot. mag. 655	
Marica plica'ta <i>B. M.</i>									
1341 jaalmifolium //.	palm-leaved	tf (23 or	■ f.mr	W	Brazil	1823.	D p.l		
1342 bermudianuni //.	Bermuda	tf tAI or	jnjl	B	Bermudas	1732.	D co	Bot. mag. 94	
Marica triodioides <i>B. M.</i>									
1343 Nuttall* <i>Swt.</i>	Nuttall's	tf A or	1 jn jl	■	N. Amer.	1823.	D s.p		
bermudia'nuiu Nut.									
1344 fineps /..	two-edged	tf A or	1 jnjl	B	N. Amer.	1693.	I co	Bot. mag. 461	
1345 mucronatum <i>Mx.</i>	mucronate	tf A or	\$ Jn'j	B	N. Amer.	1812.	J pi		
1345 convoluitu <i>Noc.</i>	convolute	jf ^AJ or	J my.jn	Y	S. Amer.	1816.	I co	Red. HI. 47	
1347 tenuifoliu <i>J/i.m.</i>	slender-leaved	tf AJ or	4 my.jn	Y	S. Amer.	1816.	D co	Red. hi. 275	
1348 Idtcum /..	yellow	tf iAJ or	myjl	Y	S. Amer.	1823.	D fl.p		
1349 caftornicum <i>Spr.</i>	Californian	£ AJ or	my s	Y	California	1796.	Sks.p	Bot. mag. 983	
1350 micranthus tW	small-lloered	tf iAJ o"	1 Jd-J	Y	S. Amer.	1815.	D co	Cav.dis. 191.2	
1351 tridilohum <i>Hum.</i>	Iris-leaved	tf AJ or	i myjl	Y	S. Amer.	1822.	D s.p	Bot reg. 646	
MARica tridit'olia A	7.								
1352 lixum <i>Lk.</i>	loose	tf AJ or	j my.jl	Y	S. Amer.	1818.	D s.p		
1353 striatum <i>CW.</i>	channelled	^ A or	2 ap.s	Y	Mexico	1788.	Ske.p	Bot. mag. 701	
146. BOBA'RT/y L. BOB A tin/v. (Jacob Bobart, professor of botany at Oxford, 1669.)					Iridear.	3			
1354 gladiäta <i>Swt.</i>	gladiated	tf iAi or	2 jnjl	Y	C. G. H.	1816.	D s.p	Bot. reg. 229	
Muricagladi&aJfor, Mor'Vz gladiata IV.									
1355s)Kitliacea//.	sheathed	tf iAJ or	J jnjl	■	C. G. H.	1798.	D s.p	Thun. ir. 1	
Mor'rt spathacca <i>Ker.</i>									
1356 aurantaca <i>Zuc.</i>	orange	tf A or	f f.ap	O	Belgia	1827.	D s.p		
147. VIEUSSEUX'iiLar. VIEI»SEUXIA. (M. Vteusseux, a physician of Geneva.)					Iuleee.	11.			
1357 tripetalos <i>Wes Dec.</i>	3-petaled-like	tf iAJ or	■ ap.my	V	C. G.H.	1802.	O fi.p	Bot. mag. 70S	
Mor'rt tripetalos <i>Ker.</i>									
1358 pavonica <i>Dec.</i>	peacock	tf A or	1 myjn	R.B	C. G. H.	1790.	O s.p	Bot mag. 1247	
Mor'ri pavonia <i>B.M.</i>									
1359 villosa <i>Spr.</i>	villous	tf AJ or	1 ap.my	P	C. G. H.	1789.	D s.p	Bot. mag. 571	
/ris villosa <i>Ker.</i>									
1360 tricuspis <i>Fit.</i>	three-pointed	tf AJ or	■ J	G	C. G. H.	1776.	D s.p	Bot. mag. 696	
1361 /ris tricuspis <i>Thun.</i>	More'a tricuspid Ker, Ferrftri/i trictipos JA.	tf Jn'j	■						
1362 Belleneften <i>Swt.</i>	grey-ed	tf J^j or	1 myjl	R.H	C. G. H.	1776.	D s.p	Bot mag. 168	
Bellen den's									
1363 tenuis It. #S.	f&ortc'a tricuspis y	Kitea It. M.	tf TA or	1 jnjl	Y	C. G. H.	1803.	D i.p	Bot mag. 772
Mor'& tenuis // K.	slender								
1364 spiribif R. Sf S.	spiral	tf iAl or	1 myjn	P	C. G. H.	1807.	D s.p	Bot mag. 1047	
1365 unguiculata R. ft S.	soft-clawed	tf iAlor	■ myjn	W	C. G. H.	1824.	D s.p	Iar. dis. 5	
Morce'a unguiculata H.K.									
1366 liurida <i>Swt.</i>	lurid	tf iAJ or	1 jnjl Cr	C. G. II.	1817.	D Bp	Bot. reg. 312		
Morte'a Urilda B. R.									
1367 fugax <i>Lar.</i>	fugacious	tf LAJ or	1 jnjl	P	C. G. H.	1825.	D s.p	Lar. dis. ic.	
148. RENEA'LMM R. Br. RBNEALMIA. (P.&M. L. Renealmc, first a Fr. phys., otherabot) Iridceee. 3—4.									
1368 grandiflora R. Br.	large-flowered	tf LAJ or	H ap	W	N. Zeal.	1822.	R p.l	Sw. fl. gar. 64	
1369 paniculata/t.fr.	panirled	tf iA or	\ ap.au	W	N. Hull.	182J.	D p.i		
1370 pulchella A. Br.	pretty	tf LS or	1 ap.jl	W	N. Holl.	1823.	D p.l		
149. PARDA'NTHUS Ker	PARDANTHUS. {Pardos, a leopard, anthos, a flower.)				Iidete.	fi			
1371 chindnsis Ker	Chinese	tf A or	2 jn jl	O	China	1759.	R p1	Bot mag. 171	
1372 nepalcnris Swt	Nepal	tf A or	2 Jn jl	O	Nepal	1823.	D p.l		

150. SCHCE'NUS L. Boo Rrsu. (*Schoinus*, a cord; made into cordage.) *Cyperacete.* 5.—C8.
 1373 mucronatus L.. pointed Jb A un 1 ap.my Ap S. Europe 1781. 1) bog M. gr. 1. 43
 1374 nigricans /., blackish Jb & A_ww f 1 jl Ap Britain sp. bo. D bog Eng. bot. J121
 1375 imberbis R Br. beardless Jb iAI □ 1 ap.my Ap N. Holl. 1822. D bog
 1376 melanostachys.fl.flr. black-spiked Jlb Alcu 1 il au Ap W. Indies 1H22. 1) bog
 1377 stellatus W. star-headed JlitiAlpr fs.'d Ap
 151. CH^TCKSPORA R. Br. CIATOSPORA. (*Chafe*, a bristle or awn, *spora*, «»ed-l *Cyperaceat.* 2—23.
 1378 ferruginea Hum. rusty Jk A un i ap.my Ap Europe 1821. D bog Sc. *—. 1.1.4
 Schce'nus ferrugineus L.
 1379 turbinata R. Br. turbinated Jb iAI cu i jl Ap N. Holl. 1820. D bog
152. LEPIDOSPE'R'M A Lab. LEPIDOSPERMA. (*Lepis*, a scale, *sperma*, a seed.) *Cyperacef.* 1. -- SI.
 1380 gladiata U7f.A. gladiated JIULACU 4jl.au Ap N. Holl. 1819. D bog Lab.n. h. 1.12
- Isa DUU'CHIUM Pets. Dnuciiuv. (The name of an island, where found.) *Cyperaceat.* L--8.
 138J spathaceum Pw*. spathaci-ous Jb A cu ... jl.au Ap N. Amer. 1818. D bog Pluk. <1. 301.1
154. RHYNCHO'SPORA Vahl RIIVNCHOSPORA. (*Rhygchos*, a beak, *spora*, a seed.) *Cypcracc**. 3.-38.
 1382 alba Vahl wh'te-headed Jb A w 1 au Ap Britain bogs. D bog Eng. bot 985
 1383 fusca Vahl brown-headed Jb A w 1 au Ap Britain bogs. D bog Eng. bot. 1575
 1384 comata Lk. tult-headed Jlib(ZQcu If' au Ap Brazil 1820. D bog
155. FIMBRI'STYLIS Vahl FIMDRISTYMS. (*Fimbria*, a fringe, *stylis*, a style.) *Cypeiacea?* 4.-65.
 1385 dichotoma ftiM riichotomous Jl O un 1 jn.jl Ap E. Indie* 1819. 1) bog Ktb. gr. 1.1
 1386 hamnum R. &S. annual Jl O un jn.jl.ap S. Europe LsO. S bog Host gr. 3.63
 1387 puberula Vahl mossy Jli A un 1 jl.au Ap N. Amer. 1820. 1) bog
 1388 dipliylla Vahl two-leaved M A un 1 jn.jl Ap Nepal 1822. S bog
- III-1 ABILDGAA'RD/i4 Vahl ABILDGAARUIA. (*Prof. Abilgaard*, of Copenhagen.) *Cyperdccee.* 8.-4.
 138) moixMfchava Vahl oncsipked dliaIcu ijn.au Ap N. Holl. 1819. D bog Sl.jam. 1.7ft 2?
 1300 trfstachys Vahl three-spiked JlAI cu 1 jn.au Ap N. Holl. 1824. D bog
- 157 TSO'LEPI'S R.Br. ISOLEPI'S. (*Ios*, equal, />&, a scale; flower of equal scales.) *Cyperacea.* 8.—48.
 1391 fltitans R. Br. floating & A cu fit jl au Ap Britain dit. D aq Eng. bot 216
 6'cirpus fltitians E. B.
 1392 setacea R. Br. bri.stle-Mr Jl O cu \ jl.au Ap Britain bogs. S bog Eng. bot. 1693
 1393 tenuissima D Don slenderest Jb O cu i jn.jl Ap Nepal 1821. S bog
 .Vcirkpusdensus IVal.
 1394 nodosa R. Br. knotted JUI LAI cu 1J jl.au Ap N. Holl. 1820. D bog Un. tr. 10.13.2
 A'cirpus gracilis Rud.
 1395 complanata R.S/S. flattened Jl iAI cu ... jl.au Ap E. Indies 1823. D bog
 1396/ołosche' nusA\$5. whole Schoenus Jb A cu A jl Ap England scash. Skbog Eng. bot. 1612
 1397 romana RffS. Roman Jb A cu 3 jl Ap Austria ... Sk bog Jac. au. 5.448
 1398 australis R \$S. southern Jl A cu 3 jl Ap S. Europe ... Sk bog Pluk. ph. 40.5
168. FCI'KPUSAAR. CUB RUSH. (*Cirs*, rushes, *Celt.*) *Cyperdea.* 17.—120.
 1399 cespitifl. B's L. tufted Jlb A ■ l jl Ap Britain tur. he. Sk bog Eng. bot. 102
 1400 pdci florae E. B. flowered Jlb A w i n u Ap Britain bgs. m. Skbog Eng. bot. 1122
 1401 elongittus Ham. elongated Jlb A un 1 s Ap Nopal 1825. S bog
 1402 lacurtris L. lake J• A ec 6 jlau Ap Britain riv*rs. Sk bog Eng. bot. 635
 1403 RitiruUitf M. jointed Jlb EM un 2 a.s Ap Europe 1821 S bog R. mal 12.71
 1404 glafem E. B. glaucous Jlb A w 2 jl au Ap England sal. m. Sk bog Eng. bot. 2321
 1405 triquetor /, triangular Jlb A W 3 au Ap England mar. Sk bog Eng. bot. 1694
 1406 murcronatus jr. shnqi-pointed Jlb A un 2 an Ap Kur. Asia ... Sk bog
 1407 carnarius E. B. keeled JUI A un 3 jl.au Ap England riv. ba. Sk bog Eng. bot. 11\$)
 1408 carcinus 5m. Carox-like Jlb A CU 1 np.my Ap Bnfa.n bogs. D bog Eng. bot. 1010
 .^choBluscompresso E. B.
 1409 rufus E. F. rufous Jb A cu I jl Ap Scotland bogs. D bog Eng. bot. 71l
 Schee'mis rufis E. B.
 1410 maritimus L. salt-marsh Jb A w ■ ls Ap Britain sal. m. Sk bog Eng. bot. 542
 J411 radicans Srk. rooting Jl A un ■ lau Ap Europe 1800. D bog Sellk. us. 4.1
 1412 atrovirens MM. dark-green Jlb A un ■ lau Ap N. Amer. 1H18. 1) bog
 1413 Luzufce L. Luzula Jb I un ■ ls Ap R Indies 1776. Sk bog Pluk. m. 27.417.3
 1414 Hylyaticus L. wood Jb A " jl.s Ap Britain m. n. p. Sk bog Eng. bot 919
 1415 quinquanguUrusr*<iAlvc.angled* Jb A cu ■ Jn.jl Ap . . . 1820. D bog
159. ELEO'CHARIS rt. Br. SPIKE RUSH. (*Elos*, a marsh, *ehalro*, to delight.) *Cyperacete.* 9.—28.
 1416 aedta R Br. acute-Avz/af Jb IAIcu * jl.au Ap N. Holl. 181Q. i>log
 1117 geniculata R. Br. jointed Jb IAI cu Ap W. Ind. 1822 D bog Sl. jam. 1.83. 3
 1418 palustris Roth. marsh Jb A " Ap Britain mar. Sk bog Eng. bot 131
 JSI*ovia Roth ovate Jk A un ■ ijl Ap Germany 1818. Sk bog
 i-raj'milticaatl Sm. Many-stemmed Jb A cu ■ jl.au Ap Britain bogs. D bog Eng. bot. 1187
 <circpi' multicaulis E. B.
 T421 aric-ul5ris R. Br. needle Jlb A w f jl Ap Britain mar. Skbog Eng. bot. 74l>
 "c(nuiaiciuUrii E. B.
 }22 obtuse-scaled Mi A cu {jl.au Ap N. Amer. 1818. D bog
 1421 glaucescens R S/S. glauccscnt Jlb A cu I jl.au Ap N. Amer. 1830. D bog
 1424 tenulis R.S/S. slender Jlb A cu 4jl.au Ap N. Amer. 1824. D bog
160. ELY'NA Schr. ELA'NA. (*Elyo*, to cover.) *Cyperdccee* 1.
 14y> spicata Schr. spiked ■ A cu { jn.jl Ap Europe 1819. S bog Vil.del.2.6
- Carex Bellardi All. myosuroides Vil.
 1H1 ERIOPHORUM L COTTON GRAM. (*Eriion*, wool, *phoreo*, to liear; seeds in wool.) *Cyperdc**. 7.-9.
 3JS? vaginatum L. sheathed JUI A P' ' mr.ap Ap Britain moors. D bog Eng. bot 873
 JS^[10]? s¹ chyf S?*■ mny-spiked Jb 2 fir ■ jn.jl Ap nntain bog*. 1) bog Eng. bot. 563
 14H pubescens Eng. Ft. downy m □ or ■ my.jl Ap England bogs. 1) bog
 1426 angustifolium Reich. narrow-leaved m A P' ■ an Ap Britain bogs. D bog Eng. bot. 564
 14M virk-nicum L. Virginian m A P' ■ my.au Ap N. Amer. IHFO. D bog Pluk. al. 29Q.4
 1431 fracile Ruth slender Jl A P' ■ jl.au Ap Scotland ac. mo. D bog Eng. bot. 2402
 1432 capitulum E. B. headed Jl A P' f au.s Ap Scotland sc. mo. D bog Eng. bot. 2387
- *162. TRTCHOOPHORUMMeR. TRTCHOOPBORITM. (TORfe, hair*, pAftrq, to bear.) *Cyperdcceat.* 2
 1433 cypriinum Pen. Cyperus-Uke Jb A cu 2| my.s Ap N. Amer. 1802. D bog Pluk. m. 419.3
 1434 alpinum Pers. alpine Jb A cu ■ Ap Scotland bogs. D bog Eng. bot 311

• 163. CYPETUJS L.	CYPEHI'S. (<i>Cynris</i> , a name of Venus; roou aphrodisiacal.)	<i>Cyperdete.</i> 50.— 300.
1435 Aituni R. & S.	Aiton'8 Ji iAJ cu H jn.jl Ap C. G. H. -	S bog
1436 articulatus L.	jointed ■ O - au Ap Jamaica 1817. S bog SI. jam. 81.1	
1437 niloticus Forsk.	Nilotic ■ O cu au Ap Egypt 1810. S bog	
1438 viscosus H. K.	clammy ■ my.au Ap Jamaica 1781. Sk bog Jac. ic. 2.295	
1439 compactus Retz.	compact ■ jl.O un jl.au Ap Chile 1819. S bog	
1440 tenellus Vahl	slender ■ uOJcu 5 Ap C. G. H. 1819. S bog Pluk. al. 300.4.5	
1441 mucronatus Rtb.	mucronate ■ QJ un 5 Ap W. Indies 1819. S bog Rtb. gr. 8.	
1442 pannonicus L.	Pannoman jut O cu Ap Hungary 1781. Skbog Host gr. 3.20	
1443 briza>'us Rich.	Briza-Hke ■ El un 1 jl.au Ap S. Amer. 1817. S bog	
1444 alternitifolius L.	alternate-lvd ■ El cu 1 jnjil Ap Madagascar 1781. Sk bog Jac. ic. 2.998	
1445 dubius Rtb.	doubtful ■ El cu 2 fmr Ap E. Indies 1802. S bog Rtb. gr. 20.4.5	
1446 conglomeratus L.	conglomerated ■ my.s Ap Arabia 1818. S bog Rtb.gr. 15.7	
1447 compresus L.	compressed ■ jl Ap Jamaica 1820. S bog SI. jam. 77.1	
1448 vegetus W.	froh M A cu > jl Ap America 1790. Sk bog Jac. vin. 3.12	
1449 ligularia L.	g ■ O un 1 jl.au Ap Jamaica 1817. S bog Rtb.gr. 11.2	
1450 Linzula W.	Luzula m EJ cu 2 my.s Ap W. Indies ... Skbog Rtb.gr. 13.3	
1451 paniculatus Vahl	panicled ■ El cu 1 my.jn Ap E. Indies 1804. D bog	
1452 pStens	spreading ■ O un 1 jl Ap W. Indies 1821. S bog	
1453 rigidus	rigid ■ El un 2 ■ Ap W. Indies 1826. S bog	
1454 pallescens	pallescens ■ A un 2 jn.jl Ap S. Europe ... D bog Desf.atl.9	
1455 flavescens L.	flavescnt ■ G cu 1 n.s Ap Germany 1776. S bog Host gr. 3.72	
1456 fuscus L.	brown ■ O cu JL Li Ap England mar. S bog Host gr. a 73	
1407 conglomerates Rtb.	conglomerated ■ El cu my.s Ap Arabia 1820. D bog	
1458 erubescens Lk.	erubescent ■ LAI cu my.jn Ap & Europe 1820. D bog	
1459 austri&lis ScAn	southern ■ O un my.jl Ap & Europe 1824. S bog Seg. vcr. a.2.2	
1460 esculentus L.	eatable A cul Ap & Europe 1507. Skbog Hobtgr.3.75	
1451 hydra A/j.	hydra ■ A un Ap N. Amer. 1812. S bog	
1462 Pangorei A/	Pangore's JyfaEJ un Ap Ceylon 1824. S bog Rtb. gr.7.3	
1463 b&dios L.	brown JB LCU Ap Algiers 180X. Skbog Desf.at. 1.7.2	
1464 pātulus Host	spreading jib O un 1 ■ Ap S. Europe 1824. S bog Host gr. 74	
1465 longus L.	long Mi A cu 3 jI' Ap England mar. Sk bog Eng. hot. 1309	
1466 tenuiflorus Rib.	slender-flwd M A cu 1 Lau Ap S. EurojMi 1818. S bog Jac. ic. 2.29G	
1467 Mōnū L. fil.	Monti's ■ O un 1 s ■ Ap Italy 1834. S bog Mo. gr. 12.1. 2	
1468 paramattensis X*.	Paramatta ■ O un 1 my.jn Ap Paraniata 1826. S bog	
1469 Strigosus L.	strigose ■ JifaElcu ■ Ap W. Indies 1786. Skbog Rtb.gr.40. 11.3	
1470 J'nā L.	Ira jifaEJcu ■ Ap E. Indies 1802. Skbog R.mal. 12.56	
1471 distans Vahl	distant ■ u\w cu 2 jl.au Ap W. Indies 1820. D bog Jac. ic. 299	
1472 fastigiatus Rtb.	fastigiate ■ El cu 1 my.au Ap E. Indies 1800. Skbog Rtb.gr.32.7.2	
1473 Alopcurodes A/A.	Foxtail-like ■ LCU 2 my.au Ap C. G. H. 1804. Skbog Rtb.gr.38.8.2	
1474 crugntus Rtb.	bloody ■ A un 1 jnjil Ap S. Europe 1823. S bog Rtb.gr. 5.1	
1475 glomeratus L.	clustered ■ k o cu 2 my.au Ap Italy 1804. S bog	
1476 polycēphalus Lk.	many-headed ■ QJ un ■ Ap S. Europe 1820. S bog	
1477 polystachys 7/6.	many-spikid ■ QJ un ■ Ap E. Indies 1819. S bog Rtb.gr.11.1	
1478 fascicularis Lam.	parcelled ■ O un ■ Ap S. Europe ... S bog Lam.ii.38.	
1479 leucocephalus	white-headed Mi A un au Ap W. Indies 1819. S bog	
1480 carcinus D. Don	Carcx-hke ■ A un aus Ap NeNil 1826. S bog	
1481 tristilus D. Don	three-furrowed ■ A un n Ap Nepal 1824. S bog	
1482 diffonis L.	two-formed Mi O un jnjil Ap E. Indies ... S bog	
1483 prostratus ttfrrt.	prostrate ■ O un ■ Ap Egypt 1825. S bog Rtb.gr. 9.2	
1484 dives Del.	rich ■ A un ■ jnjil Ap Egypt 1824. S bog Del. »g. 5.3	
• 164. PAPY^RUS Lk.	PAPYRUS. (Zfotair, Syrian; hence Papirus, Egyptian paper.) <i>Cyperdete<e.</i> 4.-6.	
1485 antiquorum Lk.	ancient ■ EJor 10 jl.s Ap Egypt 18d3. D b?i Mic. gen. 44.19	
1486 odoratus W.	sweet-scented ■ El ■ 10 jl.au Ap W. Indies 1819. D bog SI. jam. 74.10	
Cyprius odoratus Brtn.		
1487 laxiflora Spr.	lax-flowered ^ E l - ■ jl.au Ap Madagasc.1822. D bog	
1488 elegans G. Don	elegant ^ IAJ or ■ jl.au Ap W. Indies 1820. D bog SI. jam. 1.75.1	
Cypinus elegans L.		
165. KYLLINGI L.	KVLLINGA. (P. Kylline, a Danish botanist, d. 1696.) <i>Cyperdete.</i> 6.—14.	
1489 monocephala L.	one-headed JbEfunk jnjil Ap India 1793. Skbog Rtbgr.4.4	
1490 intermedia R. Br.	intermediate Jb El un 4 jnjil Ap N. Holl. 1822. S bog	
1491 polycēphala Lk.	many-headed ■ un 1 jl.au Ap Brazil 1820. 1) bog	
1492 uncinata Lk.	hooked ■ un 1 jl.au Ap Brazil 1820. D bog	
1493 triceps L.	three-headed ■ l u ■ i ■ Ap India 1776. Skbog IUh.gr.4.G	
1494 brevifolia Rib.	short-leaved ■ El un 1 my Ap W. Indies 1817. D bog Rtb. gr. 4.3	
166. Af ARI'SCUS Vahl	MARISCUS. (<i>Mar</i> , a marsh, <i>Celt.</i> ; growing in marshes.) <i>Cyperdete.</i> 7.— 28.	
1495 umbellatus Vahl	umbelled Mi El cu 14 jn.au Ap E. Indies 1789. Skbog Rtb.gr.4.2	
1496 clatus Vahl	tall ■ Et CU ■ Ap E. indies 1803. Sk bog Jac. ic. 2.500	
1497 confluxus Lk.	conHexed ■ El CU ■ Ap Brazil 1819. D bog	
1498 aggregatus W.	aggregated ill El cu T jnjil Ap 1822. D bog	
1499 Jianleus Vahl	ranic-xro<<<	
1500 laevis R. Br.	smooth ■ lA lun 2 jl.au Ap S. Amer. 1820. D bog Rtb.gr. 6.1	
1501 ovulare Vahl	egg-seeded ■ Et un 1 jl.au Ap N. Holl. 1818. S bog	
	Carolina 1818. S bog Pluk. al. 91.4	
167. REMIRE'A Aub.	REMIHEA. (The native name in Guiana.) <i>Gramknee.</i> 1.—2.	
1502 maritima Aub.	sea ■ A cu Ajl.au Ap Florida 1822. Dseas. Aub. gui. 16	
168. LY'GEUM L.	LYCEUM. (Lygeo, to bend; its flexibility for ropes, baskets, etc.) <i>Granuñcte.</i> 1.	
1503 spartini L.	Broom ML A oe 1 myjn Ap Spain 1776. D co Cl. h. 2. S	
169. COITNUCOTIJE L.	CORNUCOPIA (<i>Cornu</i> , a horn, <i>cōma</i> , plenty; the spike) <i>Grantmes.</i> 1.	
1504 cucullatumL.	hooded ■ O cu j au Ap Levant 1788. S co Fl.gr. 1.51	
170. CE'NCHRUS L.	CENCHRUS. (<i>Kegchros</i> , the Greek name of the Millet.) <i>Graminece.</i> 4.-12.	
1505 cchinatus L.	echinated ■ O cu 2 aud. Ap W. Indies 1691. S co Cayic.39.C2	
1506 iribuloides L.	Tribulus-like ■ O cu 1 my.au Ap N. Amer. 1818. S co CaV.ic.5.461	
1507 splenifex Cav.	thorn-producing ■ A cu 1 myjn Ap S. Amer. 1820. S co Cav. ic. 5.461	
1508 auatrifilis R. Br.	southern ■ O cu 1 jn.au Ap N. Holl. 1822. S co	
171. FUIRE'Nil Rtb.	FUIRENA. (<i>G. Fuirena</i> , a Danish botanist.) <i>Cyperacet.</i> 1. — 9.	
1509 umbellata Rtb.	umbelled ■ k El cu 1 myjn Ap W. Indies 1825. D bog Rtb. gr. 17.	
172. HYPCKLYTRUM Vahl	HYPOLVTRIUM. (<i>Hypos</i> , underneath, <i>elytron</i> , involvrum.) <i>Cyperacea.</i> 1.—9>	
1510 senegalense Rich.	Senegal JI 23 cu 2 jl.au Ap Senegal 1824. S bog	

- | | | | | |
|---------------------------|--|---|-------------|------------------------------|
| 173. PENNISETUM Rich. | PENNISETUM | (<i>Penna</i> , a feather, <i>seta</i> , a bristle.) | G | ■ ■ ■ |
| 1£11 scoturga Rich. | setose | Jb O un 2 jl.au Ap | Brazil | IHI7. S co Ru. am. fi. 7.2.6 |
| 1512 polyst&chymum R/S/S. | many-spiked | Jb O un 2 jl.au Ap | E. Indies | IS04. S co Ru.am.6.7.2 |
| 1513 violaceum Lam. | " | Violet Jb O un 1 . . . Ap | Senegal | 1810. S co |
| 1514 orient&le Rich. | oriental | m Q un 1 injill Ap | Guiana | 1817. S co |
| 1515 compressum R. Br. | compressed | m E3 un 1 jl.au Ap | N. Holl. | 1810'. S co |
| 1516 Alocoides A # S. | Holcus-like | M O un 1 jlau Ap | £. Indies | 1816. S co |
| Panicum Alocofodis Rox. | | | | |
| 1517 cenchroides Rich. | Cenchrus-like | M Ol cu U my.au Ap | C.G.H. | 1777. S co |
| 1518 barbatum R. SrS. | bearded | Jk O un 1 jlau Ap | E. Indies | 1823. S co |
| 1519 dichotomum Del | dichotomous | Jit Q) un 2 jl.au Ap | Egypt | 1833. S co |
| 1520 nepalense Spr. | Nepal | Jb O un 2 jlau Ap | Nepal | 1822. S co |
| 174. SPARTINA Schreb. | SPARTINA. | (<i>Spartina</i> , a rope made from broom.) | Gramine* | 8.—10. |
| 1521 striata Roth | strict | m cu 1 au Ap | Britain | sal. m. D co |
| 1592 cynosuroides Roth | Dog's-tail-like | Jb cu 3 au.s Ap | N. Amer. | 1781. D co |
| 1523 alterniflora Xoi. | alternate-flwd | Ji cu 6 au.s Ap | France | 1819. D co |
| 1524 polyst&chya Mhl. | many-spiked | Jb cu 6 au.s Ap | N. Amer. | 1781. D co |
| 1525 puraila Roth | dwarf | M cu i aus.s Ap | N. Amer. | 1826. 1) co |
| 1526/uncca A/A. | rushy | Mi cu 11 jlau Ap | N. Amer. | 1781. D co |
| 1527 patens A/A. | spreading | ilk A cu 14 jlau Ap | N. Amer. | 1781. S rm |
| 1528 gemulata ifoitv. | jointed | Jh lAI cu 2 jl.au Ap | Java | 1822. S co Bur. in. 12.2 |
| 175. DIPLA'ZIA Rich. | DIPLAZIA. | (<i>Diplazo</i> , to double; the glumes.) | Cyperdcete. | 1. |
| 1529 karattio Wa. Rich. | Karutas-leaved | Jb El un 3 au s Ap | Guiana | 1835. 13 co |
| 176. tfa'RDU'S L. | MAT GRASS. (<i>Nardos</i> , Gr., <i>nard</i> , Hub, having a peculiar perfume.) | Gramine* | 1.—2. | |
| 1530 striata L. | strict | Jb A cu i jn.l Ap | Britain | molh. D bog Eng. bot 290 |
| 177. ORYZO'PSIS Mx. | ORYZOPSIS. | (<i>Oryza</i> , rice, <i>opsh</i> , appearance.) | Graminece. | 1. |
| 1531 asperifolia Mx. | rough-leaved | Jit cu 3 jl.au Ap | N. Amer. | 1822. D co |

CLASS III. ORDER 2.

TMA'NDRIA DIGYNIA.

- | | | | | | |
|-------------------|------------------------------|-------------------------|---|-----------------------|--|
| 1532 | setacum Mr. | setaceous | Jb O un | \ j.lau* | Ap N. Amer. 1818. S co |
| 1533 | ciliatifolium Mx. | ciliated-leaved | Jb QS un | 1 jLau | Ap S. Amer. 1820. S co |
| | pubescens W. | | | | |
| 1534 | débil Poir. | weak | Jb O un | J | Ap Asia 1817. S co |
| 1535 | fibiforme Swz. | lillorin | Mi O un | J | Ap W. Indies 1817. S co |
| | <i>SwdTzidium</i> Flug. | | | | |
| 1536 | vaginatum Swz. | sheathed | Jb O un | I au | Ap W. Indies 1820. S co |
| 1537 | coujugatum Berg. | conjugate | Jb O un | 1 jnjl | Ap S. Amer. 1817. S co |
| 1538 | not Hum Flug. | marked | Jb O un | 1 jl.au | Ap W. Indies 1817. S co |
| 1539 | inaequile Lk. | unequal | Jb O un | 1 jl.au | Ap Manilla 1819. S co |
| 1540 | Kora Forst. | Kora | Jb ES or | 1j jLau | Ap E. Indies 1810. S co |
| | littoralis JL Br. | | | | |
| 1541 | gl&brum Poir. | smooth | Jb (SI un | 1 jl.au | Ap W. Indies 1816. S co |
| 1542 | pusillum Ven. | weak | Jfa GEJ un | J jl.au | Ap W. Indies 1822. S co |
| | orbicul&tum Poir. | | | | |
| 1543 | diffirme R. & S. | two-formed | Jb O un | 1 jljau | Ap N. Amer. 1818. S co |
| 1544 | la'vis Mx. | 6mooth | Jb O un | 2 jnl.jl | Ap N. Amer. 1820. S co |
| 1545 | quadrifarium Lam. | four-rowed | Jb O un | 1 jnjl | Ap Trinidad 1820. S co |
| 1546 | dubium Dec. | doubtful | Jb O un | 1 jl.au | Ap . 1818. S co |
| 1547 | villosum Thun. | villous | Jb O un | 1 jl.au | Ap Asia 1824. S co |
| 1548 | distichum L. | two-ranked | Jb OQ cu | 11 jl | Ap Jamaica 1776. S co |
| 1549 | serotinum Flug. | evening | Jb O cu | Ijl.jl.au | Ap N. Amer. 1804. S co |
| | Digitaria pildsa W. | | | | |
| 1550 | diw^ctup//. | dissected | Jb O un | J | Ap S. Amer. 1818. S co |
| 1551 | scrobiculatum L. | furrowed | m d c u | n jjs | Ap E. Indies 1778. S co |
| 1552 | tencillum lte</> | tender | * O u n | lijl.au | Ap . 1817. S co |
| 1553 | pancul&tum L. | panidied | Jt ED cu | 8 jls | Ap Jamaica 1782. S co |
| 1554 | stoloniferum Bosc | stoloniferous | Jb iAI cu | 2 JLB | Ap Peru 1794. S co |
| 1555 | virg&tum L. | twiggy | Jb O un | 1 injl | Ap W. Indies 1820. S co |
| 1556 | elegans Flug. | elegant | Jb O un | 14 jl | Ap Brazil 1816. S co |
| | | | | | |
| | | AupuiPoaN. | (<i>Amphi</i> , round, <i>pogon</i> , a beard.) | Graminece. | 1—5 |
| | strict | Jb TAJ cu | ... my.jn | Ap N. Holl 1823. S co | |
| | | | | | |
| 12 ^o P | AXONOPUS Beauv. | AXONOPUS. | (<i>Axon</i> , axis, <i>pous</i> , a foot.) | Graminece. | 3. |
| 1558 | cimicinus Beam. | cimexian | Jb O cu | I jl a | Ap India 1788. S co |
| 1559 | digitatus Beauv. | digitate | Jb (Q) un | J | Ap W. Indies 1820. S co |
| law | aQreus Beauv. | golden | Jhfrqun | J | Ap W. Indies 1821. S co |
| | | | | | H. & B. 1.27 |
| | | MILLET GRASS. | { <i>Milr.</i> , a thousand, seeds.) | Graminece. | 7.—20. |
| | | common | Jt A * | 3 jnjl | Ap Britain m. s. p. S m.s Eng. bot. 1106 |
| | | crowded | Jb A un | 1 jnjl | Ap Switzerl. 1825T. S co |
| | | velvety | Jb O un | lijl.au | Ap Mexico 1823. S co |
| | | Gallician | Jb O un | 2 jn.jl | Ap Galicia 1824. S co |
| | | vernal | Jb O un | 2 jn.jl | Ap Tauria 1824. S co |
| | | small-leaved | Jb O un | 1 jnjl | Ap N. Spain 1825. S co |
| | | frutescent | Jb A cu | 1 ... | Ap Crimea 1822. S co |
| | | | | | |
| 182. | CERE'S/4 Pers. | CBRESIA. | (<i>Ceres</i> , a goddess, inventress of tillage.) | Graminece. | 1.*. 2. |
| 1568 | elegans Pers. | elegant | Jf IAlor | 2 jl.au | Ap Peru 1816. S eo Jdc. ff. 80 |
| | i*aspalum membranaceum Flug. | | | | |
| | | | | | |
| 18a | PIPTATHETRUM Beauv. | PIPTATHIERUM. | (<i>Pipto</i> , to fall, other, an awn.) | Graminece. | 3. |
| 1569 | paradoxum Beauv. | paradoxical | M A cu | 3 jh.jl | Ap France 1771. S co Host gr. 3.23 |
| | Jfilum paradoxum L. | Arachne virescens Trin, | | | |
| 1570 | caerulescens Beauv. | ceruleolescent | Jb A cu | 2 jn.jl | Ap Barbary 1819 S co Desf. at.1.12 |
| | Jfilum cruseleciifH Schous. | | | | |

TRIANDRIA DIGYNIA.

- 1655 tenacissima R.\$\$. most tenacious JBkO ■ 2 jl.s Ap Brazil 1819. S co
 1656! emisatum R.\$\$. Pennisetum jw O ■ 2 jl.au Ap S. Amer. 1821. S co
 ld? scändens Spr. scandent Jk O ■ 3 JL Ap 1820. S co
 it&S auricoma Lk. golden-hairca Jj O un ■ JL Ap IK IS. S co
 IC'9 aspera Lk. rough Ja A un ■ jl.au Ap C. G. H. 1820. S co
 litij muncata Lk. prickly Jj O ■ JL Ap N. Amer. 1818. S co
 1661 cenchroides Cenchrus-like i O " " JL.ap Cayenne 1821. S co
188. ECHINO'CHLOA JRMKW. PRICKLY GRASS. (*Echinos*, a hedgehog, *chloea*, grass.) *Graminea*. 8.—
 1662 ttagnata Uettuo. stagnant Jlt O un J jUu Ap" Hungary 1802. S & Host gr. 3.51
 1663 commutata R. #S. changed Jj O un Ujl.au Ap
 IMicum stavninum Host
 1664 intermedia It. Si S. intermediate Jj O un li jl.au Ap 1818. S co
 1665 crythropurpurea R.Sfs. red-seeded Jj O un 1 jl.au Ap E. Indies 1781. S co
 1666 crua-edri Beauv. crow's-foot Jj O un 1 jl.au Ap Britain xnoif. S co Eng.bot.876
 1667 <rus.galli Beauv. cock's-foot Jj O un li jl.au Ap
Pdñxum crus-galli E. li.
 1668 echinata It. 4 & prickly Jj O ■ 1 jl.au Ap N. Amer. 1818. S co
 1669 setigera It. % S. setigerous Jj O ■ 1 jl.au Ap E. Indies 1820. S co
189. ORTHOPCVGON R. Br. ORTHIOPOGOV. (*Orftos*, straight, *pō'ōn*, »JjJ!!». *Graminea*. 6.-28.
 1670 interitus R. Br. hairy Jj OB as 1 jnjl Ap W. Indies 1791. S co
 1671 uidulatitibus Spr. wavy-leaved Jj A un M Ap S. Euro/c 1795. S co Host gr. S.52
 1672 liurmum It. Br. Hurman's THun Jj Jl Ap E. Indies 1800. S co Hur. in. 12.1
 1673 Aromoides R. #S. Bromus-like Jj ■ un jl.au Ap Trinidad 1820. S co Ru.ain.6.5.2
ollaccus Spr.
 1674 compaditus R. Br. compound Jj [13 un Jj Ap Ceylon 1818. J J H. pfl. 12. 96.1
 1675 africanus G. Don African Jj E3 un Jj Ap Africa 1824. J J Beau. ov. 2.67.1
Oplismenus atricaius Beauv.
190. MOKA'CHNE Beam
 1676 unilateralis is Beauv. one-sided A/lium villusum Lk.
- 1678 spedita W. spiked Jj O ■
- V.Y2. [SA'CHNE R. Br. IS.VCHNE. (Isos, equal, *achne*, a gume.)
 1679 australis R. Br. southern Jj A un I jnjl Ap N. Holl. 1820. S bog
193. ZAPPA'GO Schreb. LAPPOAC, (L < M, the burdock; with rough prickly flowers.) ^ J Jf Tsf
 1680 racemosa Schreb. racemose m 6 cu Jj.l.a Ap S. Europe 1771. S co Hostgr. 1.S6
194. STIPPA L. GBASS. (*Stippa*, a silky or feathery substance.) *Graminea*. g.—
 1681 pennita L. feathered ilk A or Q jl.au Britain al.roc. 1 s.l Eng. bot.
 itiHShtmili Cav. low Jj O cu Jj.l.a Ap S. Amer. 1802. S co Cav. ic. 5. 466.1
 H-8Jt6rcilis Delt. twUted Jj A un 1 jl.au Ap Greece 1890. D pi Fl.gr. 86
 1681 Lagta* It & S. Lukasch Jj A un U jnjl Ap Spain 1820. D pi Dcif. at. 1.28
 liR1 tcn.irissanna L. most tenacious M A ■ Jj ■ Suiiii 1817. D co Desf. at. 1.30
 liH capillata L. long-h.ired ML A ■ E jl.au Ap Europe 1815. D co Hoat gr. 3.5
 1687 Jfincea L. Jj ■ ■ Jj A CU Ap France 1772. D co Fl.gr. 1.85
 1688 gigantea Lag. giant Jj A cu 3 jl.au Ap Spain 1823. D 8U
 1689 anstilla/,. nwhed Jj A un 1 jl.au Ap S. Europe 1819. D pi Fl.gr. 87
 1690 bicolor Vahl two-colored Jj A un 2 jl.au Ap K. Amer. 1820. D Fp Cav. ic. 5.466.2
 1691 con ferta R.%S. crowded Jj A cu 2 Jl.l.a Ap 1819. P Lp
 1692 sibirica Lam. Siberian Jj A cu 3 jl.au Ap Siberia 1777. D co Gm. si. 1.22
193. ARISTIDAZ. Ai:isTin\.
 1693 vagans R. Br. wandering Jj AJ cu 2 jl.au Ap N. Holl. 1816. D pi Cav. ic. 5.471
 1694 stipoidea R. Br. Stipa-hkc Jj Ub_AJcu 1|jnjl Ap N. Holl. 1826. S pl
196. CH/ETA'RIA Beauv. CIIJEXAKIA (*Maria*, "an arrow or blistle") *Graminea*. 4.-21.
 1695 adscensis Beauv. ascension Jj Ut(Z)un 1 Jl.l.a Ap W. Indies 1820. S co SL. jam. 1.2.56
 Aristida adscensis W.
 1696 h^stmx Beauv. i>orcup»^ Jj [S] un 2 jl.au Ap E. Indies 1820. S co Pluk.al. 191.5
 1697 divaricata Hum. ci"»-spate Jj AJ un 2 jl.au Ap S. Amer. 1818. S co Jar. ec. 6
 1698 cwrulacencis R. \$ * esrulescnt Jj O un 2 jl.au Ap Canaries 1818. S co Dcsf. at. 1.21.2
197. CH*TP'*US Lk. CiiRTiiRdB. Chate, a head of hair, oura, a tail; silky panicles.) *Graminea*. 1.—2.
 1699 fasciculus Lk. bundled Jj O un i Jlf Ap Spain 181a S co
- 198 LAGUR'RUS L. HARE'S-TAII. Gnxss. (I-agos, a hare, oura, a tail.) *GravMnea*. 1.
 1700 ovatus L. ovate-sptfr Jj O cu 1 jn Ap Guernsey bor. fi. S co Eng. bot 1384
199. ECHINOPOGON Beauv. ECIUNOPOGON. (*Echinos*, prickly, *poeon*, a beard.) *Graminea*. 1.
 1701 ovitava Beauv. ovate Jj JU un 1 jl.au Ap N. Holl. 1820. S pfl Lab. n. h. 1.21
200. HIRRO'CHLOE Gm. HIBROCHILOE. (*Micros*, holy, *chloe*, grass.) *Graminea*. a—7.
 1702 borealis Illy. northern Jj A cu 1 my,jn Ap Scotland moun. D co Gr. wo. 1G7. ic.
JLACUH borealis Schr. odoratus L.
 1703 ariRalis R.fS. southern Jj A cu 1 jnjl Ap S. Europe 1777. D co
 1704 frjtrans R A.S. sweet-scented Jj A cu 1 jnjl Ap N. Amer. 1777. D co
21. POLYPO'GON Delti POLYPOGOX. (Poly, many, *pagan*, a beard.) *Graminea*. 1704
 1705 monspelitica Delt. montpellier Jj O pr Jj.l.a Ap Britain way s. S co Eng. bot 1704
 1706 littoralis H.S.n. Khorc Jj O pr Jj.l.a Ap Britain sea s. D co Eng. bot 1251
 ifKrostii htoralis E. B.
 1707 maritimus IV. sea-side M O un 1 jl.au Ap 8. Europe 1817. S co
 1708 tativifl. Fis. Tarturian Jj O un 1 jl.au Ap Tartary 1820. S co
 ltO) glomeratus IV. glomerated Jj O un li jl.au Ap N. Amer. 1820. 3 co
- 202 GASTRI'DFUM Beauv. GASTRIDIVM. (*Gadridion*, a little swelling.) *Graminea*. 2.
 1710 australis Beauv. southern Jj O ag i jl.au Ap Britain san. IS co Eng. bot 1107
 JVHm lendgerum E. B.
 1711 maticum Spr. beardless M O un linjl.au Ap Sicily 1819. S co

* 203. <i>JGR Cy STIS L.</i>	<i>m</i>	BENT GRASS. (The Greek name for all grasses. <i>Agros</i> , a field.) <i>Graminete.</i> 40.—110.
1712 <i>stolonifera L.</i>		creeping <i>Fiorin</i> j ^ ag 1 jl Ap Britain moi.m. C h.l Eng. bot 1532
2 <i>latifolia Sin.</i>		broad-leaved - <i>Jt A ag</i> 1 jl Ap Britain mo. pa. S co Gr.wo.225 ic.
3 <i>angustifolia Sin.</i>		narrow-leaved * <i>A ^ g</i> 1 jl Ap Britain mo. pa. S co Gr.wo.346.ic.
4 <i>aristata Sin.</i>		awned <i>WX A ag</i> 1 jl Ap Britain mo. i». S co Gr.wo.345.ic.
5 <i>ncmorialis Sin.</i>		grove <i>4 A a g</i> I JJ Ap Britain woods. S co
6 <i>pallidris Sin.</i>		marsh * <i>A ag</i> 1 jl Ap Britain mar. S co
1713 <i>diffusa Host</i>		diffuse <i>Jb A un</i> 1 jl.lau Ap Austria 1818. I) co Hostgr.4.55
1714 <i>neglecta Schr.</i>		neglected <i>Jb A un</i> 1 jl.lau Ap Pannonia 1820. D co
1715 <i>capillaris Sm.</i>		capillary <i>Jb A un</i> Jjl.lau Ap Caucasus 1825. D co Sm. ic. in. 3.54
1716 <i>tenella R. & S.</i>		slender <i>Jt A un</i> 1 jl.lau Ap Tyrol 1821. D e o Sc. ger. 1.5.1
1717 <i>spina R. & S.</i>		spurious <i>Jb A un</i> 1 jl.lau Ap Pannonia 1826. D e o Hostgr. 4 51
1718 <i>decumbens Host</i>		decumbent <i>Jb A un</i> 1 jnjl Ap Pannonia 1819. D e o Hostgr. 4.54
1719 <i>Elba L.</i>		white <i>Fiorin</i> ^ A ag Ap Britain mar. S m.s Gr.wo.342.ic.
2 <i>purpureascens</i>		<i>purpura8ccnt</i> Wf A ag Ap Britain mar. S co Eng. bot. 118
3 <i>vivipara</i>		viviparous wk A ^ g Ap Britain mar. S co
1720 <i>dulcis Sm.</i>		sweet <i>Jb A un</i> jn.jl Ap Greece 1818. D co
1721 <i>tenuifolia Bieb.</i>		fine-leaved <i>Jb A un</i> jl.lau Ap Caucasus 1820. D co
1722 <i>pätna G. u.</i>		spreading <i>Jb A un</i> jl.lau Ap Switzerl. 1818. I) co
1723 <i>valentina R. k S.</i>		<i>Valentia</i> <i>Jb A un</i> jl.lau Ap France 1819. D co
1724 <i>gigantea Roth</i>		gigantic <i>Jb A un</i> jl.lau Ap Caucasus 1820. D co Sc. ger. 1.11. 2
1725 <i>dubia L.</i>		doubtful <i>Jb A un</i> jl.lau Ap Germany 1818. D co Lee. her. 4. 4
1726 <i>frondosa Ten.</i>		frondose <i>M A un</i> jn.jl Ap Italy 1824. D co
1727 <i>vulgaria E. B.</i>		common <i>Jk A w</i> jl.lau Ap Britain me. pa. S 1 Eng. bot. 1671
1728 <i>ambigua A. & S.</i>		ambiguous <i>Jb A un</i> 1 Jl Ap Europe 1819. D co Host gr. 56
<i>lMJuies&L Lk.</i>		<i>Juresc</i> <i>Jb A un</i> 1 jnjl Ap Germany 1820. D co
1730 <i>Juixira Mx.</i>		rough <i>Jb A un</i> 1 jnjl Ap N. Amer. 1820. D co
1731 <i>capensis Tte&D.</i>		Cape <i>Jb A un</i> Jjl.lau Ap C. G. H. 1818. D co
1732 <i>virginica W.</i>		Virginian <i>Jb A un</i> Jjl.lau Ap N. Amer. 1820. I) co Lab. n. h. 1.23
1733 <i>maritima Lam.</i>		sea-side <i>Jb A un</i> Jjl.lau Ap France 1818. D co
1734 <i>versicolor Stev.</i>		party-colored <i>Jb A un</i> Jjl.lau Ap Caucasus 1820. D co
1735 <i>debilis Poir.</i>		weak <i>Jb A un</i> Jjl.lau Ap France 1818. D co
1736 <i>Frsteri R. & S.</i>		<i>Forster's</i> <i>Mi A un</i> 1 jnjl Ap Austria 1821. D co
17J7 <i>hispida W.</i>		hispida <i>Jb A un</i> Jjl.lau Ap Europe 1805. S co
1738 <i>sylvatica L.</i>		wood <i>Jb A un</i> Jjjnjl Ap Britain woods. S m.s Lee. her. 4.3
1739 <i>varia Host</i>		variable <i>Jb A un</i> Jnl Ap Germany 1817. D co
1740 <i>latifolia 0<<</i>		broad-leaved <i>Jb A un</i> Jnl Ap Germany 1826. D co
1741 <i>Ten6r Otto</i>		Tenore's <i>Jb A un</i> Jnl Ap Italy 182a D co
1742 <i>plebeia A. ifr.</i>		plebeian <i>Jb A lin</i> 1 jl.lau Ap N. Holl. 1806 1) co
1743 <i>aemula A. ffc</i>		rival <i>Jb A lin</i> 1 jl.lau Ap N. Holl. 1822. I) co
1744 <i>Billardieri R. Br.</i>		<i>La Billardier's</i> <i>M A un</i> 1 jl.lau Ap N. Holl. 1806. D co Lab. n. h. 1.31
1745 <i>verticillata W.</i>		<i>vrhoTl-Jtotuer Mi O un</i> 1 jnjl Ap S. Europe 1800. S co
1746 <i>spica-venti W.</i>		<i>vrhoTl-Jtotuer Mi O un</i> 1 jnjl Ap Enclaid san. fi. S s.l Eng. bot 951
1747 <i>interrupta L.</i>		long-awned <i>Jb A cu</i> 1 jl.lau Ap S. Europe 1820. S co Host gr. 48
1748 <i>pallida Dec.</i>		<i>windward-bpkd Jb O cu</i> 4 jnjl Ap France 1819. S co
1749 <i>purpurca Griff.</i>		interrupted <i>Mi O un</i> 2 jl.lau Ap Vallesia 1821. S co
1750 <i>pauciflora It.&S.</i>		pale <i>Jb O un</i> jnjl Ap Switzerl. 1824. D co
1751 <i>retrofracta IV.cn.</i>		purple <i>Jb O un</i> 1 jnjl Ap N. Holl. 1806. S s.l
204. TRICHXCHLOA <i>Dec IRKHOCHLOA</i> <i>Un</i> (<i>Thrix</i> , hair, <i>chloa</i> , grass.) <i>Graminete.</i> 6.—30.		
1752 <i>longiseta Trtn.</i>		long-awned <i>Jb A cu</i> 1 jl.lau Ap Amer. 1820. S co
ylgrfstis temufl6ra <i>Muhl.</i> <i>Chma tenuiflora Lk</i>		
1753 <i>mexicana Trin.</i>		Mexican <i>Mi A cu</i> J. I Ap Mexico 1780. S co Wil.h.b.12
ifgrfstis mexicana L.		<i>Chma mexicana Beauv.</i>
1754 <i>fohosa Ttin.</i>		leafy <i>Jb A cu</i> 1 jl.lau Ap N. Amer. 1819. S co
ifurotis filiformisW.		<i>Clinia nTiformrisLA.</i>
1755 <i>sobolifera Trtn.</i>		sobohfero <i>M A cu</i> 1 jl.lau Ap N. Amer. 1819. S ro
1756 <i>microsperma Dec.</i>		small-seeded <i>Jb O cu</i> 1 jl.lau Ap Mexico 1820. S co
<i>AgTlMu microsperma Lag.</i>		
1757 <i>capillaris Dec.</i>		capillary <i>Jb A cu</i> jl.lau Ap Carolina 1818. S co
805. TRICHO'DIUM Mx. TRICHO'DIUM. <i>[square]</i> hair, <i>eidos</i> , like; inflorescence.) <i>Graminete.</i> 12.—18.		
1758 <i>caninum Schr.</i>		dog <i>Jb A w</i> 1^ jl.lau Ap Britain nns. S co Eng. bot. 1856
1759 <i>alpinum II.S.S.</i>		alpine <i>Jb A un</i> Ap Europe f821. D co Sc. ger 1. 3.4
1760 <i>rupestre R. & S.</i>		rock <i>Jb A un</i> Ap S. Europe 1815. S co Sc. ger. 1 3.5
1761 <i>nvgluctum R. & S.</i>		neglected <i>Jb A un</i> jnjl Ap Pannonia 1819. D co
1762 <i>flavescens Host</i>		flavescent <i>Jb A un</i> jnjl Ap Europe 1817. D co Host gr. 4.52
1763 <i>setaceum R. & S.</i>		bristly <i>Jb A un</i> jl.lau Ap Britain dr. he. S co Eng. bot 1188
1764 <i>rubrum Schult.</i>		red <i>Jb A un</i> jl.lau Ap Lapland 1820. D co
Z765 <i>hybridum Schult.</i>		hybrid <i>Jb A un</i> jnjl Ap Switzerl. 1821. D co
1766 <i>diffusum Lk.</i>		diffuse <i>M A un</i> J Ap 1816. S co
1767 <i>decumbens Mx.</i>		decumbent <i>Jb A ag</i> jnjl Ap N. Amer. 1786. S co Fras. mon. ic.
1768 <i>laxiflorum Mx.</i>		loose-flowered <i>Jb Q un</i> jl.lau Ap N. Amer. 1818. S co Mic. am. 1. 8
1769 <i>elegans Thore</i>		elegant <i>Jb O un</i> jnjl Ap France 1818. D co Thore j. 2
206. TRTSTEGIS Nees. TRISTEGIS. (Tress, three, <i>stege</i> a covering; three glumes.) <i>Graminea</i> 1		
1770 <i>glutinosa Nees</i>		clammy <i>M A CU</i> * jnjl Ap 1822. S co Nees h. b.7
207. SPORCBOLUS R Br. SPOROBOLUS. (<i>Svoros</i> , a seed, <i>ballo</i> , to cast forth.) <i>Graminee</i> 5.—10.		
1771 <i>indicus 72 Br.</i>		Indian <i>M O cu</i> 2 au.o Ap India 177a S co SLjamVai
1772 <i>purpurascens Beauv.</i>		<i>purpurascens JlbOcu If aus.</i> Ap W. Indies 1806. S co
ifgrfstis purpurascens Stvx.		
1773 <i>tenacissimus Beauv.</i>		most tenacious <i>Jb EJ cu</i> i aus. Ap E. Indies 1801. ■ □ Jac. ic. 16
1774 <i>elongatus R. Br.</i>		elongated <i>Jb O cu</i> i aus. Ap N. Holl. 182a ■ □
1775 <i>diandrus Beauv.</i>		diandrous <i>Jb iQI cu</i> i aus. Ap E. Indies 1820. ■ □
Ag6tis diandrus Retz.		
208. AIRO'PSISD^v. AIROPSIS <i>Aira</i> , hair grass, <i>opsis</i> , like.) <i>Graminea</i> 4.-6.		
1776 <i>globosa Desv.</i>		globose <i>Mi A un</i> 1 jnjl Ap France 1826. D co Thore j. 1.7.3
1777 <i>obtusata Desv.</i>		obtuse-valved <i>Jb A un</i> 1 jnjl Ap Carolina 1826. D co
1778 <i>t'amlölei Desv.</i>		Decandolle's <i>Jb A cu</i> 1 jnjl Ap S. Europe 1820. S co Dec. ic. 1
— <i>Hastidea Dec.</i>		<i>Aira /rgrostidea Lois.</i>
1779 <i>brevidolia R. % s.</i>		short-leaved <i>Jb A cu</i> 1 jnjl Ap Missouri 1818. S co
<i>ira brevifolia Ph.</i>		
Trtn. PBMBALU. (<i>Perilallo</i> , to encompass.) <i>Graminete.</i> 1.		
1780 <i>hispanica Trtn.</i>		Spanish <i>Jb O un</i> 1 jn Ap Spain 1820. S co Cav. Ic 44 1
<i>Aira involucrata Cav.</i>		

210. PSA'MMA Beauv.	MAT GRASS.	(<i>Psammos</i> , sand; Its place of growth.)	<i>Graminea</i> . 1—2.
1781 areñaria Beauv.	sand	Já A cu 2 jnjl Ap Britain sea & co. S 8 Eng. bot. 520	
Brando arcnaria E.B.			
211. CRY'PSIS H. K.	CRYPTIS. / <i>Krypto</i> , to conceal; the flowers concealed in (heleaves.)	<i>Gromline</i> *. 3.—&	
1782 aculeata H. K.	prickly	V o u n *au Ap S. Europe 1783. S co Host gr. 1.31	
1783 alopecuroides Sehr.	Foxtail-like	M Q un ij.l.au Ap Germany 1816. S co Host gr. 1.29	
Heleocholo Alopecuroides Host.	Phalaris crypsoides Urv.		
1784 chaenoides Lam.	Rush-like	JU O un J au Ap & Europe 1783. S co Host gr. 30	
212. ILOPECURUS L.	FOXTAIL GRASS.	(<i>Alopex</i> , a fox, <i>aura</i> , a toil.)	<i>Gramine</i> *. 15—22.
1785 bulbosus L.	bulbos	1 A w ' l'jl Ap England sal. m. S las Eng. bot. 1249	
1785 arundinaceus Pair.	Icoid-like	Jk A ag 2 jnjl Ap Europe 18C6. D co	
1787 macrōstachya Pair.	long-spiked	Já A cu 1 jn.au Ap Burbary 1826. S co	
1788 colobochnoides R. S(S)	Colobache-lkjUfa	A cu 1J jn.au Ap Siberia 182a S co	
1789 brahytachys Bieb.	short-spiked	1 A □ A □ Jn.jn.ap Siberia 1820. S co	
1790 pratensis L.	nicaillow	Já A ag 2 my Ap Britain mea. S hl Eng. bot. 759	
1791 nigricans Horn.	blackish	M A ag 4 jnjl Ap Europe 1815. S co Jacec. 13	
1792 alpinus E. B.	Alpine	Já A cu i my.jn Ap Scotland sc.mo. S s.l Eng. bot. 1126	
1793 agrestis L.	wild	M O w M jlau Ap Britain ro. aid. S s.l Eng. bot. 848	
1794 geniculatus L.	jointed	A A w 1 myau Ap Britain mea. S m. Eng. bot. 1250	
1795 fulvus E. B.	bladdered	M A w 1 jn Ap England ponds. S m.s Eng. bot. 1467	
17% utricularius Per*.			
1797 ramosus Poir.	branching	A O un 1 jlau Ap Italy 1777. S co Host gr. 3.7	
1798 ventricosus Pen.	ventricose	Já A un 1 jnjl Ap Carolina 1819. D co	
1799 cchinatus Thun.	hedgehog	JB tfl un 1 jnjl Ap France 1821. D co	
		C G. II. 1825. S co	
213. PHLE'UM L.	CAT'B-TAIL GRASS.	'Greek name for <i>Typha?</i> , applied by L. to this plant.)	<i>Graminea</i> . 9.—
1800 pratense L.	meadow	Jb A ag 2 jl Ap Britain me. pa. S m.s Eng. bot. —	
1 majus Sin.	larger	M A ag 2 jl Ap Britain me. pa. S co Gr. wo 195. ic.	
2 minus Sin.	smaller	B A □ 2 jl Ap Britain me pa. S co Gr. wo. 197. ic.	
1801 nodosum L.	knotted	JU A w Ujl.s Ap Britain Wales. S co Fl. dan. 380	
1802 Bertolini M R. & S.	Bertolini's	M A un 1 jnjl Ap Italy 1826. O co	
	stolonif'rum Bert. ?		
1803 alpinum	Alpine	iU A cu 1 jl Ap Scotland sr. alp. S hl Eng. bot 519	
1804 Micheli W. en.	MicheU's	M A ag 1 jn.jl Ap Scotland al. roc. S co Eng. bot 2265	
1805 commutatum Gau.	changed	M A un \ jnjl Ap Valesia 1820. D co	
1806 cfliuum Sm.	cat's	JC Q un 1 jl Ap Greece 1819. S co	
1807 cchituum Host	hedgehog	M Q un jnjl Ap S. Europe 1818. S co Host gr. 3.11	
1808 Gerard All.	Gerard's	M A un 1 jnjl Ap France 1820. D co Jac. ic. S01	
214. ACHNODONTON Beauv.	ACHNODOTON.	(<i>Ach?</i> w, & chaff or husk odous,& tooth.)	<i>Grammitce</i> . 2.
1809 Bellardi Beauv.	Bellardi's	Já A un i jnjl Ap Spain 1798. S co	
Phalaris Bell&xdf W.			
	slender	Já O un 1 jnjl Ap Mesopota 1804. S co Bar. ic. 14.1	
215. CHIL&CHLOA Beauv.	CHILOCHLOA.	(<i>Cnigos</i> , fodder, <i>chloa</i> , grass.)	<i>Gramlnete</i> . 5.—&
1811 Boehmcri Beauv.	B<rhmer's	Já O w T Jl Ap England plains. S co Eng. bot. 479	
Phl&uai Bd'hmeri E. B.			
1812 cuspulata R. S.S.	cuspidate	Já A un j jnjl Ap Austria 1821. D co Host gr. 4.20	
1813 aspora R. % S.	rou^h-	Jlk O w 1 jlau Ap England hea. S. co Eng. bot 1077	
Pnl'cum paniculiflum E. B.			
1814 annua R. % S.	annual	Já O un J jnjl Ap Caucasus 1821. S co	
Phl'cum finnuum Bieb.			
1815 arenaria Schr.	s.md	Jáfa O w ijl.au Ap England seaxo. S co Eng. bot 222	
Phdlaris arenaria E. B.			
216. PHA'LARIS L.	CANARY GRABS.	(<i>Phalaros</i> , brilliant; having shining seeds.)	<i>Graminete</i> . 13.—23.
1816 canariensis L.	Canary	Já O ag 2 jn.au Ap Britain unc. pi. S r.m Eng. bot 1310	
1817 aquatica H. K.	water	+ OUn Ujjj.jl Ap Egypt 1778. S co Host gr. £ 38	
1818 minor Retz.	smaller	Já O un 1 Jjl Ap S. Europe 1823. S co	
1819 commutata R. S/S.	changed	Já O or jnjl ^JP Italy 182a S co Bar. ic 700.2	
1820 cap'nui8 Thun.	Cape	Já O un jnjl AP C G. H. 1804 S co	
1821 caerulea Dcft.	circulscsent	Já Q un jnjl AP Spain 1818. S Bux. c. 4.53	
1822 nitida Presl	shining	Já Q un jnjl ^P S. Europe 1820. S CO	
1823 quadifolius Lag.	four-valved	Já O un jnjl AP Spain 1824. S CO	
1821 appendiculata R. S/S.	appeiidiculate	Já O or jnjl AP Egypt 1820. S CO	
1825 paradoxia L.	paradoxical	M O un jnjl AP Levant 1687. S CO Host gr. 2.40	
1826 microstachya Dec.	small-spiked	Já Q un 1 jnjl AP N. Amer. 1826 S CO	
1827 bulbosa Cav.	bulbous	Jlk A un 1 jnjl AP Spain 1798. S CO Cav. ic. 1.64	
1828 nodosa Bieb.	knotty	Jáfa A un H jn.jl AP S. Europe 1820. D co	
SI7. DE'GRAPHIS Trm.	DEGRAPHIIS	(Dis, twice, grapho, to mark)	<i>Graminea</i> . 2.—
1829 arundinacea Trin.	Reed-like	jáfa A w 3 jl Ap Britain riv. ba. S co Eng. bot 402	
Artido colorata E. B.			
IQO: 2vari Ka ta	variegated	Já A or 3 jl Ap Britain gard. ■ co	
i-30 americana Ell.	American	Já A um 3 jl Ap N. Amer. 1820. ■ co	
IQO: 19 hflarisarundiniflora Mx.			
IB31 canescens Beauv.	canaeacent	Já A w ij.l.au Ap England san.sh. S s.l Eng. bot. 1190	
ilira cantescens i5. A			
1832 articulatus Beauv.	jointed	Mi A cu i jl.au Ap S. Euro ic 1816. S co Deaf, at 1. IS	
219. ^I'HA L.	HAIR GRASS.	(The Greek name of our Loliom.)	<i>Graminea</i> . 12.—18.
1833 /ilchel)a W.	pretty	Já O un ij Ap Spain 1820. S s	
1834 flexuosa L.	Hexuous	Já A w 1 jlau Ap Britain hea. S s.l Eng. bot 1519	
WV caryophylla L.			
HJ18 versicolor R. S.S.	Cltivc-pink./ii</	Já O w Ap Britain sa. pas. S s.l Eng. bot 812	
1837 atropurp <i>></i> iroa IVahl.	dark-purple	Já A un Ap Spain 1821. I) co	
1838 linduicra Lag.	maggot-bearing	Jlk O un i jn.jl Ap Finnmark 1820. S co FL dan. 961	
1839 humifus Bieb.	dwarf	Jáfa A 1 jn.jl Ap Spain 1826. D co	
1840 refracta Lag.	refracted	Jáfa A un 1 jn.jl Ap Caucasus 1826. D co	
3841 glabrata Brot.	smooth	Jlk A un 1 jnjl Ap Spain 1825. D co	
1842 pallens Sjt.	pale	Jáfa A un 1 jl.au Ap Portugal 1820. D co	
1H43 arundinacea L.	Reed-like	Jáfa A cu 3 jl.au Ap N. Amer. KLO S co	
1844 seminedutra Kit.	half-neuter	Jáfa A cu 1 jl.au Ap Cumana 1817. S co	
J*hflaris seminectra R. S S		Hungary 1812. S co	

220. CATABRO'SA Beauv. CATABBOSA. (<i>Katabrosis</i> , food.)	Gramlnca. 2.—	
1815 aquatica Beauv. aquatic & A cu 1\$ my.jn Ap England rivu). D aq Eng. bot 1557		
<i>Aim aquatica E. B.</i>		
1846 viridula R.#S.	greenish £ A cu 1 jnj.l Ap —— 1816. D aq	
•231. DESCHA'MPS/^\ Beauv DESCHAMPSIA. (AT. <i>Deschamps</i> , M.D. a French botanist.) Graminece. 11.—14.		
1847 capitiflora Beauv. tufted jlk A wo jnj.l Ap Britain m.sp. S co Eng. bot. 1453		
ATA caspita L.		
2 vivipara viviparous m A cu 3 jnj.l Ap Britain m. sp. S co		
1848 glomerata G. Don glormerated JIU A cu 3 jnj.l Ap Scotland moun. S co		
1849 la>vigita RS/S. smoth-leaved Jb A cu 1 jlau Ap Scotland sc. alp. S co Eng. bot 2102		
<i>Aim Isvigatj Sm.</i>		
1850 nlnina R S/S. Alpine M A un Ap Denmark 1819. S □ Fl. dan. 961		
1851 bottmica Trin. Itottmica M A cu 3 Jnj.l Ap Bottnia 1816. S □ Wa. lap. 1.4		
Air a. Irrttmu JVah.		
1852 moitiina G. Dan ATA montiflta L.	mountain Jlb A un 2 jnj.l Ap Lapland 1817. S co Stel.gr. 4	
/Sr,3 ambifl. Beam. ambiguous jlb A un 2 jnj.l Ap N. Amer. 18*1. S co		
IHJ4 ilicocar. It. # S. two-colored jif A un 1J jnj.l Ap France 1817. S co		
IH-JO'a'rcia Beauv. Kush-like jlb A un 2 jnj.l Ap S. Europe 1819. S co		
1856 media It. # S. mediate juk A un 1 jnj.l Ap Montpel. 1817. S co		
292 EIUA'CHNE 7i.Br. ERIACHNB. (<i>Eriion</i> , wool, achm, a glume.) Graminece. 1.—8.		
1857 obtfoa R. Br. obtuse Jb A cu 1 jnj.l Ap N. Ainer. 1819. S co Ac. pet 11.7		
jfira truncata Mid.		
22.J. ilVE'NA L.	OAT GRASS. (Aten, from r./kn, to eat, Cel.) Gramneat. 17.—40.	
1858 pra-cox Beano. Praj'cox E. B.	early Jib O w i my.jn Ap Britain hea. S co Eng. bot 1216	
<i>Aim praj'cox E. B.</i>		
1859 brevin Roth short □ O un f jnj.l Ap Germany 1804. S co Host gr. 3.42		
1860 filba Dec. white □ O un 1 jnj.l Ap France 18*10. 1) co		
1861 oricntalis Schreb. eastern □ O *B 3 jnj.l Ap 17*8. S co Host gr. 3.44		
1862 sativa L. cultivated □ O ag 3 jnj.l Ap —— rm Host gr. 2.59		
1 molanospferma black-seeded □ O ag 3 jnj.l Ap —— co		
2 leucosperma white-seeded □ O ag 3 jnj.l Ap —— co		
IMS ndda L. naked □ O <2 2 jnj.l Ap —— rm Host gr. 3.43		
1867 semifleryrens Vil. insipid □ O w 4 au Ap Britain cor. fi. co Host gr. 2.58		
IHC8 fallax Itee. barren □ O cu 4 jlau Ap Barbary 1640. □ co Host gr. 2.57		
18ti9setacea Pi. hirsute □ O un 3 jus Ap Barbary 17*8. S co		
1870 pratensis Z. evergreen □ A un 1 jnj.l Ap Cierniau 1820. 1) co Vil. del. 2.5		
1871 gromodfes Gou. doubtful □ A un 1 jnj.l Ap France 1824. 1) co		
1872 pl. imculinis Schr. setaceous □ A un 1 jnj.l Ap France 1825. D co Vil. del. 2. 5		
1873 vernedlor Ti. meadow □ A ag 1 Jajl Ap Britain me. pa. S hi Eng. bot. K04		
1874 pumila Detf. flus talked □ A un 2 jnj.l Ap France 1820. D co		
party-colored □ A un 1 jnj.l Ap Germany 18*1. 1) co Sc.gcr.6.2		
dwarf □ A un □ Ap S. Euro>E 1819. 1) co Vil. del. 2.4.5		
224. TRISE'TUM Pers. TRISBTLM. (Treis, three, seta, a brist. Ic.) Graminea. 20.—35.		
1875 stnfitum Per*. striated Jlb O un li jl.au Ap S. Europe 1804. S co Lee. her. 9. 3		
<i>Avena striata Lam.</i>		
1876 tenui R. \$ S. slender M O un - jnj.l Ap S. Europe 1820. S co Host gr. 2.55		
1877 L6'flmKf <th>num R. & S. Ixclflng's M O un - jnj.l Ap Spain 1770. S co Cav. ic. 1.45.1</th>	num R. & S. Ixclflng's M O un - jnj.l Ap Spain 1770. S co Cav. ic. 1.45.1	
1878 pensylvaniun J3craft; PcmnHylyvanian Jlb O un □ Ap N. Amer. 1785. S co		
1879 neglectum R % S. neglected M O un jnj.l Ap Italy 1826. S co Savi pis. 1.1.4		
1880 incranthum /f. 8j S. small-flowered JIU. O un □ Ap .18 .fi. S co		
1881 parviflorum Pers. small-flowered Jlb O un □ Ap Italy 1820. S co Desf.at. 1.32		
18H-yllopecurus li. % S. Fox-tul M O un 2 jnj.l Ap .1816. S co		
1883 coiuensatun R. \$ S. condensed Jlb O un H jnj.l Ap Sicily 1819. S co		
1884 pilosum Bicb. pilose Jlb O un 2 jnj.l Ap Iberia 1818. S. co		
1885 pub^scens Jl. \$ S. downy Jlb A w lijl.au Ap Britain ch. pa. D s.l Eng. bot 1640		
Av-na pub&scens E. B. Av-na pub&scens E. B.		
1886 rigidum Bicb. rigid Jlb A un 1 jnj.l Ap Caucasus 1819. D co		
1887 alptuum R. \$ S. alpine M A w □ Ap Britain sc. alp. D co Eng. bot 2141		
Avini planiculmis E. B. alpina E. F. Carpathian Jlb O un □ jnj.l Ap Austria 1824. S co		
1888 rarp.tu.u R. SF.S. flavescent Jlb A w □ jnj.l Ap Britain ... S co Host gr. 4.31		
1889 flavor en-. ft. ,V.N. Avena flavescens E. B.		
1890 alp^tre Beauv. mountain Jlb A un 1 jnj.l Ap Europe 1819. D co Host gr. 3. 39		
1891 brevitilium R. 8s S. short-leaved Jlb A un 1 ju-jl Ap S. Europe 1818. S co Ho-t gr. fi. 40		
1892 argenteum R. \$ S. silvery Jlb A un 1 } jnj.l Ap Switzrl. 1819. S co Host gr. 2. 53		
1893 distichophyllum Vil. distich-leaved M A un 1 jnj.l Ap Switzrl. 1796. D co Host gr. 2.53		
1894 airodes Beauv. Aira-like Jlb O un * jnj.l Ap Switzrl. 1800. S co Host gr. 2.43		
225. TRICHfl/TA Beauv. TRICHJETA. (Trels, three, chcete, a bristle.) Graminee. 1.		
1895 ovata Beauv. ovate Jlb A un ... jnj.l Ap S. Europe 1824. S co Cav. ic. 6.591.2		
2fr6mus ovdtus Cav. Trisetum ovatum Pers. Avena ovata Spr.		
226. DANTHO'N/y* Dec. DANTIIONIA. (Af. <i>Danthoine</i> , a French botanist) Graminete. 10.—18.		
1896 longil'lia R. Br. long-leaved Jb iAI un 2 jl.au Ap N. Holl. 1823. □ co		
1897 sennannuldris R. Br. semi-annualr Jb iAI un 2 jl.au Ap N. Holl. 1823. □ co Lab. n. h. 1.3		
jfrtndo semiannularis Lab.		
1898 pilosa R. Br. pilose Jb LAI un □ jl.au Ap N. Holl. 1823. □ co		
1899 provincialis is Dec. Provence Mi Q un □ jnj.l Ap S. Europe 1821. □ co Vil. del. 2.2.9		
yvleña provincialis Poir.		
1900 *tng6sa Beauv. strigose Jb O w 3 jnj.l Ap Britain hed. S co Eng. bot. 1266		
Av&w* strig6sa E. B.		
1901 spicata Mil. spiked Jb O un 2 jnj.l Ap N. Amer. 1824. S co		
A vlna glumacea Mx.		
1902 serlera Nut. silky Jb LAI un 2 jnj.l Ap N. Amer. 1826. S co		
1903 macrfintha R % S. long-flowered Jb O un 2 jnj.l Ap C. G. H. 1810. S co		
1904 curvitöna R.8/S. curved-leaved JU iAI un 1 jnj.l Ap C. G. H. 1816. S co		
1905 calycina A \$. large-cdyxed Jb LAI un 2 jnj.l Ap C. G. H. 1818. S co		
227. GAUDI'SIA Beauv. GAUDINIA. (3f. <i>Gaudiit</i> , a Swiss botanist.) Graminece. 1.		
1906 fraglhs Beauv. brittle Mi A un U jnj.au Ap Sijain 1778. D co Host gr. 2.54		
At. frigUi* L.		

228. CENTOTHE'CA Desu. CEVTOTHEICA. (*Kenteo*, to prick, *tkeea*, a sheath.) *Graminea*. 1.
1907 /appacoa Dcsy. Burdock *M* O cu 1 jl. Ap E Indies 1773. S co Beauv. gr. 14.7
Cenchrus/appaccus L.
229. PHILAGMITES *Trin* REED. (*Phragmos*, a hedge j. forming hedges.) *Gramineee*. 1.
1908 commfinis *Trin*, common fe A ec 6 jl.s Ap Britain dit D aq Eng. hot. 401
^ri'mdoPhragmftes *E. B.*
230. ^RU'NDO With. WATER REED. (*Arundo*, a reed, *Lat.*, *aru*, water, *Celt.*) *Graminea*. a—9
1909 IMmax W. Donax Ja A ec 10jl.au Ap S. Europe lo48. S co [J>tgr.4.38
2.verscolor stried M A or 3 jl.au Ap S. Europe 1648. S co M. h. 3.8.9
1910 tcnax Vahl tough Ja A un 3 jl.s Ap S. Europe 18Ei D co Cy. ne. 2.12
VIII festuccea W.cn. Fescue-like Ja A un 2 jl.s Ap Germany 1823. Deo
231. CAL AM A GRO'STIS Adan. CALAMAGROSTIS. (*Kalamos*, a reed, *agrostis*, grass.) *Graminea*. 18. --30.
1912 epigejos Roth terrestrial Ja A w S jl.au Ap Britain moi. pi. S bog Lng. bot 40J
jfrAmio epigejos *E. B.*
- 1913 stricta Spr. strict Jb A w Ujl.au Ap Scotland mar. S bog Eng. bot. 21G0
i4r6udo stricta *E. B.*
- 1914 htoreá Dec. shore Ja A un ■■■ Ap Europe 1818. D bog Sc. ger. 1.4. 2
1915 effusa Gm. effuse Ja A un 1J jn.jl Ap Siberia 1818. D bog
- 1916 laxa //OA/ loose Ja A un ■■■ Ap Europe 18m D bog Hostgr.4. 43
1917 Haller/V/m Dec. Haller's Ja A un 2" jnjl Ap Germany 1821. D bog be. ger. 1.4.3
^ri'ndo Pseudophragmit Schr.
MIR LaiKsd6rf/ Spr. Langsdorfs Jlfe A un 4 jnjl Ap 182a D bog
l> glauca Horn. glaucous Jlfa A un lftjnjl Ap N. Amer. 1819. D bog
l'M) btrigosa Wahl. rnt Jlfa A un ■■■ * jnjl Ap Finnmark 1817. D bog Wa.lap.ll. a. b.
l> ai vana Host variable 4Kb A un 1 jnjl Ap Alp. Eur. 1819. 1) bog
IH2 acutiflora R.S/S. acute-flowered a ^ un 1 jnjl Ap Europe 1820. D bog
1923 sylvatica Dec. wood Ja A un 2 jl.au Ap Germany wo. S bog Host gr. 4.49
Calamagrostis arundinacea Roth, A ^Tostis arundinacea L.
1921 confinis Spr. similar Ja A un 2 jnjl Ap N. Amcr. 1819. D bog
1923 Hostii R. & S. Host's Ja A un lijn.jl Ap Alp. Eur. 1819. D bog Host gr. 4.48
sylvatica Host
- 192G lanceolata Roth lanceolate Ja A w 2 jl.au Ap Britain m. s. s. D bog Eng. bot. 2159
ArhnAo C'alainagr6stis E.B. jfrgr6stis lanceolata B. & S.
- 1927 tenuila Host tender Ja A un f jn.jl Ap Bohemia 1824. D bog Host gr. 4. 50
jfrfindo htmiilis R SfS.
- 1928 niontana Host mountain Ja A un 1 jnjl Ap Germany 1820. D bog Hostgr.4.46
1929 speciosa Schr. showy Ja A un 2 jnjl Ap Germany 1818. D bog
232. CHRYSURUS Mx. CHRVSURUS. (*Chrysos*, gold, *oura*, a tail; the flowers.) *Graminea!* 4.
1930 auricus Beauv. goUan-sjiketl Jfc O uif jJl Ap Levant 1770. S co Host gr. 3.4
Cynosirus adreus L. cynoburolden Pers.
- 1931 echinatus Beauv. hedgehog Mi O w 2 au Ap England san. fl. S ■ Eng. bot 1333
19J2 cfVisus Beauv. cfUse M O or 1 jl.au Ap S. Europe 1824. S ■
1933 elegans Beauv. elegant Jb O or 1 jl au Ap S. Europe 1816. S ■ Desf. at. 1.17
Cynosurus elegans Del.
233. SESLE'RJ.4 Sco. SESLERIA. (*Sesler*, a physician and botanist of the ISth century.) *Graminea!* 8.—11.
1934 elongata Host elongated Ja A un ■■■ Ap Germany 1805. S ro Hostgr. 2.97
1935 cylindrica Dec. cylindrical Ja A un ■■■ j.l.au Ap S. Europe 1821. D co
1936 tenuifolia Schr. fine-leaved Ja A un ■■■ j.l.au Ap Germany IK20. D co Sc. ger. 1.6.4
19J7 caerulea Ant blue Ja A w ■■■ my.jn Ap Britain fields. S co Eng. bot. Ibl3
E. B.
1938 nkida Ten. shining Ja A un 1 jl.au Ap Italy 1826. D co
19J9 tenella Host tender Ja A R' j ap.my Ap Switzerl. 1819. S co Host gr. 2.100
1940 diaticha Pers. two-rowed Ja A un J jn.ju Ap (Germany 1876. D co Jac. ic. 19
1941 sphacerocephala Ard. round-headed Ja A pr f ap.my Ap Switzerl. 1819. S co Hostgr. 2.99
234. CYNOSURUS L. DOG'S-TAIL GRASS. (*Kyon*, a dog, *oura*, a tail.) *Gramineee*. I.—8.
1942 cristatus L. crested M A >g 2 au Ap Britain pas. S s.l Eng. bot. S16
235. KCELK'MA Pers. KOILERIA. (A. Kohler, professor of natural history at Mayenc.) *Graminecar*. 15.—20.
1943 cristata Pers. crested Ja A un 1 jn.ju Ap Europe 1800. S co Hostgr. 2. ~.~
Sglubra tt,fs. smooth Ja A or 1 jn.ju Ap Britain bus. S co Eng. bot. 643
Aim cinst&tta E. B.
- 1944 gracillbra R. & S. large-flowering Ja A un 1 jn.ju Ap S. Europe 1824. S co
l)~t glauca Dec. glaucous Ja A un J jn.jl Ap Denmark 181S. D co
- 1946 jx-nhylianica Dec. Pennsylvanian Ja A un 1 jn.jl Ap N. Amer. 1817. D co
1W lobata Bich. lohatc Ja A un * jn.jl ■ Caucasus 1825. D co
- 1948 macrantlia li.&S. long-flowered Ja A un 1 jn.ju Ap Russia ls1'4. S co
IW vallesiacra Dec. Vallais Ja A un ■■■ Ap Vallais JK17. D co
IDJO tuberosa Pers. tulicrous Ja A un ■■■ j.l.au Ap Europe 1802. ■ co Lam. il. 45.4
1951 imbecens Beauv. pubescent Ja A un ■■■ jnjl Ap S. Europe 1800. S co Ger. gal. 1
1952 liibQta Dec. hairy Ja A un ■■■ jnjl Ap Vallesia IS2I. D co
1953 villosa Dec. villous Ja A un ■■■ jnjl Ap Sweden 1818. D co
l!54 Dhleoides Pers. Cat's-taiMikc Ja O ■■■ j.l.au Ap Portugal 1802. S co Deaf. at. 1.23
1955 hispida Dec. hispid Ja o un ■■■ jfjl.au Ap Mediterr. 1819. S co Savi pis. 1.5
1966 brachystachya Dec. short-spiked Ja A un ■■■ jnjl Ap S. France IH2L. D co
1957 aegyptiaca Jac. ? Egyptian Ja O un ■■■ i jnjl Ap ■■■ W26. 8 co
- * 236. WA'CTYLIS L. COCK'S-FOO-J GRAHS. (*Dactylos*, a finger j. finger-grass.) *Gtamlnete*. 10.—20.
195H flonuTata L. glomcerated Ja A og ■■■ jnjl Ap Britain mea. S h.l Eng. bot. 335
lf;Jl Klaiu'tsons W. en. glaucescent Jui A un ■■■ jn.jl Ap Switzerl 1818. D cu
19H1 littorata IV. shore Ja A un ■■■ jn.ju Ap S. Kuropc 181H. S co Lam.il. 45.5
1961 h.spamra W. en. Spanish Ja A un ■■■ Ap Spain 1814. S co
1962 glauca Roth glaucous Ja A un ■■■ Ap S. Xony 1800. S co
l!51 littaritima Schr. sea Ji A un ■■■ Ap Cinnmany 1820. D co
l!52 rupens Dcsf. creeping 4A<>n #(!!:ji Ap liarbary 18i.l. S co Desf. at 1.15
liU'S adseriflens R. SfS. ascending Ja I A un ■■■ Ap C. G. H. 18^3. S co
1966 cynosuroides Roth Cynoborus like Ja O un ■■■ jnau Ap E. Indies 182a S co
1967 patens // K. spreading Ja A un ■■■ au.s Ap N. Amer. 1781. S co
237. GLYCE'RIA B. Br. GLYCERIA. (*Glycros*, sweet; spike of flowers.)
1968 flcitanis R. Br. floating ■■ A w 1} my.au Ap Britain iwnds. S aq Eng. bot. 150
Festica fluitans L.

	FESCUE GRASS.	(<i>Festu-</i> % pasture cr food, <i>Celt.</i>)	<i>Graminea.</i> 47.-68.
238. <i>FESTUCA</i> L.	tender	<i>m</i> O un f jlau	Ap N. Amer. 1804. S co
1969 <i>tenella</i> Ph.	sheep's	Jb A ■■■ J jn	Ap Britain dr.pa. S sl Eng. bot 565
1970 <i>ovina</i> /.	Hordeum-like	Jb A ■■■	Ap Britain „ S co Gr. wo. 159. ic.
1971 <i>Aordciiformis</i> Sin.	fine-leaved	■■■ A ■■■	Ap Britain dr.pa. S co
1972 <i>tenuifolia</i> L.	viviparous	A ■■■	Ap Britain sc.mo. S 8.1 Eng. bot 1355
1973 <i>vivipara</i> E. B.	long-haired	Jb A a g ■■■	France 1820. D co
1974 <i>capillata</i> Lam.	curved-leaved	Jb A un ■■■	Switzerl. 1825. D co
1975 <i>curvula</i> Gau.	alpine	Jib A un ■■■ jnjl	Switzerl. 1821. D co
1976 <i>alpina</i> GAU.	mountain	Jb A un ■■■ jnjl	Switzerl. 1819. D co Host gr. 4.63
1977 <i>alpstris</i> Host	Haller's	jut A ■■■	S France 181. D co
1978 <i>Halleri</i> Vil.	amethystine	Jb A un ■■■ jl	S Europe 1804. S co
1979 <i>amethystina</i> Host	grey	Jib A w ■■■ 1 jnjl	England bar. he. S co
1980 <i>cassia</i> E. B.	strict	Jib A un ■■■ 1 jnjl	Austria 1818. D co
1981 <i>strista</i> Host	intermediate	jut A u n ■■■ jnjl	Switzerl. 1K19. D co
19H2 <i>intermedia</i> R. & S.	hairy	jdk A u n ■■■ k	Germany 1818. S co
198J <i>hirsuta</i> Host	pale	Jb A "S ■■■	Switzerl. 1819. D co
1984 <i>pallens</i> Host	Vallais	jut A i ■■■	Vallais 1820. I) co
1985 <i>valleHiaca</i> Gau.	yellow	Jb A un ■■■ 1 ■■■	Switzerl. 1820. D co
19H6x <i>Enthiga</i> 11% S.	glomerated	Jb A un ■■■ 1 jnjl	France 1826. D co
1987 <i>glomerata</i> Rus.	fallacious	Jb A un ■■■ 1 jnjl	France 1826. D co
1988 <i>fallax</i> Thuil.	variable	Jb A un ■■■ 1 jnjl	S. Europe 1823. S co
1989 <i>vaiii</i> Host	flavescens	M A ■■■ f jnjl	Savoy 1804. S co
1990 <i>flavescens</i> Bel.	sheathed	Jb A ■■■ ■■■ 1 jnjl	Hungary 1804. S co
1991 <i>vaginata</i> Kit.	Pannonian	jib A ■■■ 1 jnjl	Hungary 1804. S co
1992 <i>pannonica</i> Wul.	bulbous	jib A ■■■ 1 jnjl	Italy 1820. D co
1993 <i>bulbosa</i> Biv.	ciliated	jib A ■■■ i jlau	Portugal 1802. S co
1994 <i>cihata</i> Lk.	nodding	Jb A un ■■■ 3 jnjl	N. Amer. 1805. S co
1995 <i>ndtans</i> Ph.	downy	Jib A un ■■■ 3 jnjl	Hungary 1822. S co
1996 <i>pubescens</i> Lk.	various-leaved	Jb A un ■■■ 3 jnjl	France 1812. S co
1997 <i>heterophila</i> Jl	hardish	Jb A ■■■ 1 jnjl	Britain me.pa. S
1998 <i>duriuscula</i> L.	thicket	Jb A un ■■■ 1 jnjl	Europe ... S co
1999 <i>dumetorum</i> Jl	three-flowered	Jb A un ■■■ 1 jnjl	S. Europe 1819. S co
2000 <i>triflora</i> Zte.	red	Jub A ■■■ 1 jnjl	Britain me. pa. S
2001 <i>rubra</i> L.	Welsh	Jb A "S ■■■ 1 jnjl	Wales ... S co
2002 <i>cimbrica</i> Sin.	smooth	Jb A ■■■ 1 jnjl	Britain - S co
2003 <i>glabra</i> 5/n.	Fa ⁿ nas	Jib A un ■■■ 1 jnjl	Spain 182a D co
2004 <i>Elliⁿna</i> L/ie.	Cynosurus-like	Jb A un ■■■ 1 jnjl	S. Europe 1820. S co
2005 <i>cyno</i> Kuroldes Desf.	large-flowered	Jb A un ■■■ 1 jnjl	N. Amer. 1812. S co
2006 <i>grandiflora</i> Lam.	Yatc-Jlotvering	Jb A un ■■■ 1 jnjl	Austria 1819. D co
2007 <i>scrifta</i> Jfcf	smooth	M 1 ag ■■■ 1 jnjl	Russia 1806. D co
2008 <i>loj'vis</i> Fis.	reddish	Jb A ■■■ 1 jnjl	S. Europe 1776. S co
2009 <i>rubens</i> Po«.	glaucous	Jb A a g ■■■ 1 jnjl	S. Europe ... S co
2010 <i>glatica</i> Lorn.	Mexican	Jib A un ■■■ 1 jnjl	Mexico 1805. S co
2011 <i>mexicana</i> Zio/tm	Barbaric	Jib A un ■■■ 1 jnjl	Minorca 1825. D co
2012 <i>balcarica</i> Zuc.	loose	Jib A un ■■■ 1 jnjl	Switzerl. 1819. D co
2013 <i>laxa</i> Host	scabrous	JL A un ■■■ 1 jnjl	Austria 1821. 1) co
2014 <i>scabra</i> faA/	rough	Jb A un ■■■ 1 jnjl	France 1818. D co
2015 <i>aspera</i> Pair.			
239. <i>SCHEDONOC'RUS</i> Beauv.			(<i>Sciedion</i> , near to, <i>a</i> , <i>ros</i> , <i>a</i> , <i>mountain</i> .) <i>Gramine*</i> . 17.—36.
2016 <i>pratensis</i> A Jt	meadow	■■■ A ag If jnjl	Britain m.pas. S co Eng. bot 1592
	<i>Festuca pratensis</i> E. B.		
2017 JhoMioides R. & S.	Phcenix-like	Jib A <■■■ 1 jnjl	Switzerl. 1819. S co
2018 clatio II. & S.	taller	Jib A ■■■ ■■■	Britain m. me S co Eng. bot 1593
	<i>Festdca elatinum</i> E. B.		
2019 <i>camilccens</i> R. % S.	cifirulescent	Jib A un 1 jnjl	Barbary 1818. ■ co
2020 <i>spadiceus</i> R. & S.	sheathy	Jib A <■■■ If ap.my	Italy 1775. ■ co Hostgr. 3.20
	<i>Frstdca spadicea</i> L.		
2021 <i>inermis</i> R. \$ S.	unarmed	Jb A un 2 jnjl	Germany 1794. S co Hostgr. 1.9
	<i>ifromus uiernis</i> E. B.		
2022 <i>calanatus</i> R. & S.	Reed-1 like	Jb A un 3 jlau	Scotland m.me. S co Eng. bot 1005
	<i>Festuca calamaria</i> E. B.		
2023 <i>deciduus</i> R. & S.	deciduous	Jb A ag If jnjl	Britain m.me. S co Eng. bot 226b
	<i>Festdca decidua</i> E. B.		
2024 <i>sylvaticus</i> R. & S.	wood	Jb A ag 2 jnjl	Germany 1804. S co Host gr. 2.78
	<i>Festdca sylvatica</i> Host		
2025 <i>oliaceus</i> R. & S.	loliaceous	^ A ag 1 ■■■	Britain mo. pa. S co Eng. but 1281
	<i>FestOca oliacea</i> E. B.		
2026 Scheutzl II. & S.	Schutzer's	Jb A un 1 ■■■	S Europe 1816. S
	<i>Fes/Uca Scheutzeli</i> Gau.		
2027 <i>ndtans</i> II & S.	nodding	Jb A un 1 jnjl	Hungary 1825. S co Hostgr. 4.61
	<i>FestUca n 6 tans</i> Ph.		
9028 <i>nigrescens</i> R. if S.	nigrescent	Jib A ag 1 jnjl	Switzerl. 1819. S co
	<i>FestUca nigrescens</i> Lam.		
9029 <i>violaceus</i> R. \$ S.	violaceous	Jib A un 1 jnjl	Europe 1817. S co
	<i>Fes/Uca violarpa</i> Gau.		
2Q3Q poset'Ormla R. \$ S.	Poa-form	Jb A ag 1 jnjl	Switzerl. 1819. S co Hostgr. 2.81
	<i>FestUca pistorinis</i> Host		
9031 <i>pdmilus</i> R. \$ S.	dwarf	Jb A un ■■■	S. Europe 1820. ■ co
9032 <i>tendillus</i> R. & S.	tender	Jib A un ■■■	N. Amcr. 1804. ■ co Hostgr. 2.91
	<i>Festica tendia</i> Ph.	Brachypodium/tv/tfcoldes L	
240. <i>MYGALU'RUSZ.</i> *.	MOUSE-TAIL.	(<i>Mygafr</i> , a field mouse, <i>oura</i> , a tail.) <i>Graminc*</i> . 6.	
2033 caud&tus Lk.	tailed	Jb Ow ijn	Ap Britain ways. S co Eng. bot 1412
	<i>FestUca Mydrus</i> E. B.		
2034 ffromoides Lk.	Bromus-like	Jb O w imyjn	Ap Britain walls. S co Eng. bot 1411
	<i>Festica fromoides</i> E. R.		
2035 stipoldes Lk.	Stig-a-like	Jb O un 1 jnjl	Majorca 179a S co Bar. ic. 76.1
	<i>Festica stipoldes</i>		
2036 delicate Lk.	delicate	Jb Q un <jnjl	Spain 1817. S co
	<i>Festica delicate</i>		
9037 uniglumis Lk.	single-husked	Jb O w ■■■	Britain seaco. S co Eng. bot 1430
	<i>Festica uniglumis</i> E. B.		
2038 Alnpuccroides Lk.	Foxtail-like	Jb A un ■■■	S. Europe 1819. S co
	<i>Festica tendia</i> Ph.		
241. BBOME GRASS.		(<i>BBOM</i> , a Greek name for a sort of wild oat)	<i>Gramlnete.</i> 47.—66.
2039 <i>secalinum</i> L.	Rye-like	Jb Ow 2 jnjl	Ap England corn n. S co Eng. bot. 1171
2040 elongitus Gau.	elongated	Jib O un 2 jnjl	Ap Switzerl. 1819. S co

<i>R S T R A R I A</i> 7n. <i>pubescens</i> <i>Trin.</i>	<i>Festuca intermedia</i> M A un T jn.j Ap 1820. D co.	<i>ROSTRARIA.</i> (<i>Rostrum, a beak.</i>)	<i>Grantme*</i>
<i>festucoides</i> <i>Trin.</i>	<i>pubescens</i> <i>Trin.</i>	<i>J. O un T jn.au</i>	<i>Ap Europe</i> 1820. S co.

		BHA? IIYlion»*.	(<i>Brachys, short, pouss a foot</i>)	Gramline*.	22.-2a			
		J U n A W	3 j n a U	AP	Britain	hea	S co	K.g.bot730
	<i>jirumus</i> pmnataui							
2088	rupestre A * & Zfrdmus rupestris <i>Host.</i>	rock	Jb A un	1 jnjl	Ap	SL Europe	18ia	D eo Hostgr. 4.17
2089	brevicostum R. A.S.	short-bristled	Jb A un	1 jnjl	Ap	pp.....	1825.	D co
2090	strigosum R. S. S.	strigose	Jb A un	1 jnjl	Ap	Tauria	1819.	8 ro
2091	cffispitosum R. < f. S.	tufted	Jb A un	1 jnjl	Ap	S. Europe	1820.	S co
2092	ramdum li & S.	branched	M A un	2 jnjl	Ap	S. Europe	1818.	8 ro
2093	Jdjhsnicpfds R. & S.	Micenic-like	Jt A un	1 jnjl	Ap	Germany	1821.	D ro
	»M obtumfolium Lk.	blunt-leaved	Jb M un	1 jnjl	Ap	Germany	1821.	Deaf, at 1.24.1
2095	Venorianum It & S.	Tenore*B	Jk A un	1 jnjl	Ap	Spain	1818.	8 co
2096	longifolium R. y S.	long-leaved	Jb A un	2 jnjl	Ap	Italy	1823.	S co
2097	sylvaticum Jis(S.)	wood	Jb A w	2 jnjl	Ap	Barbary	ie18.	S co
	IVrdmus sylvaticum B. R.			2 jnjl	Ap	Britain	bed.	
						S co	Eng. bot	729

SRASIDB OAT. {*Unus*, one, alone; united glumes.) Graminea. 1—7.
broad-leaved. I^b A un 4 inil Ap N. Amer. 180P.

broad-leaved	JUB	A	JUN	Ap	N. Amer.	1793.	S co			
pamfrld	JUB	A	JUN	Ap	N. Amer.	1793.	8 co Cat car. 1.32			
spiked	M	vw	li	Ap	N. AmL.ri	HIM	S co			
		in	f	jnjl	Ap	N. Amer.	1789.			
Quinquefida	Bcauv.	five-cleft	Ib	A	S	injl	Ap	N. Amer.	1820	S co 1..a
					(Tret's,	three,	cuspis,	a point.)		

847. CER ATCCHLOA Beauv. HORN GRASS. (*Keras*, a horn, *chloa*, grass; the seeds.) *Graminea*. 1. — 2L
 2115 gnioloides Beauv. Uniola-like *M Q un* *lft jl* Ap N. Amer. 1788. S co H. ber. 1.3
Festuca tenuioides JV.

248. SCHISMUS JfcnwL Sen ISM us. (*Schisme*, a cleft; outer palea.) *Graminete*. I.
 2115 marginatus Beauv. margined *Mi O un* *ijn.jl* Ap Spain 1781. S co Lam.il. 46.1
Festuca calycina L.

849. TKIO'DI A R. Br. TRIODI*. (*Treis*, three, *odous*, teeth; the palea.) *Graminea*. 1. — 10.
 2117 decumbens R. Br. decumbent *Jii A w* *1 jl.au* Ap Britain ... S co Eng.bot.7C2
Poa decumbens L.

250. BECKMANNA Host. BRCKMANNIA. (The author of the History of Inventions.) *Graminete*. 1.
 2118 erucaeformis Host Eruca-form *Mi Q un* *2 jl* Ap Europe 1773. S co Host gr. a 6
Cynusurus erucaeformis W.

251. ME'LICA /,. MELIC GRASS. (*Mel*, honey; the name in Italy for the great millet.) *Graminete*. 9. — 24.
 2119 cilifata/. ciliated *Jh A or* *3 jl* Ap Europe 1771. S sl Host gr. 2.12
Bauhinia All. *Jic A un* *2 in.il* Ap Italy* ... S co Host gr. 4.24!
 2121 niftans L. nodding *Jh A or* ... Ap Britain moun. S sl Eng. bot 1039
 2122 uniflora Retz. one-flowered *Jlh A P* *my.jn* Ap Britain groves. S m s Eng. bot. 1058
 2123 pyramidalis Lam. pyramidal *Jlh A* ... Barbary 1804. S co Bar. IC. 1.6.1
 8124 speciosa Mhl. snowy *Jlh A* ... N. Amer. 1820. S co M.h. 3.8.7.51
 2125 glabra Ph. smooth *Jlh A* *jnjl* Ap N. Amer. 1812. S co If. li. SL7.51
 2126 altUsima L. tallest *Jlh A* *jl.au* Ap Siberia 1770. S Ht>btgr.2.9
 2127 sylvatica Lk. wood *Jlh A* *jnjl* Ap Germany 1824. D co
un

* 252. MOLIN'NZ/f Schrank. MOLINIA. (?*J Molina*, who wrote on the plants of Chile, in 1782.) *Graminete*. 1.
 2128 cajrulea Schrank blue *Mi A w* *1 an* Ap Britain bogs. S p.m Eng. bot. 750
Melica esculca E. B.

253. /TM'ZA L. QUAKING GRASS. (*Brizo*, to nod; trembling spikolets.) *Graminea*. 9.
 2129 minor /. smaller *Jlh O* or *ftjl.au* Ap England corn fl. S co Eng. bot. 1316
 2130 virens L. green *M O* or *lft jl an* Ap Spain 1800. S Hay. ter. 25. 6
 2131 hifmilis Afefc humble *Jtt O un* *jl.jnjl* Ap Caucasus 1825. S Sc. ger. 4. 10
 2132 media L. mediate *ill A* or *II my.jn* Ap Britain pas. Eng. bot. 340
 2133 Cltisii If & a *Clusius's* *Jlh ^ or* *myjl* Ap S. Europe 1820 ... Thorej.3.24.2
 2134 gemulata Thun. jointed *m o or* *jlau* Ap C. G. H. 1816. Fl.gr. 1.74
 2135 elatior Sm. taller *Q* or *jl* Ap Greece 1817. S Host gr. 2.30
 2136 maxima £. greatest *o or* *ij* Ap S. Europe 1633. S
 2137 rubra Lam. red *Jlh O pr* *1 jnjl* Ap S. Europe 1H20. S

254. PO'A /,. MISADOW GRASS. (*Poa*, grass or herbage.) *Graminea*. 72. — 143.
 2138 alpina L. Alpine *Jn A cu* *I jnjl* Ap Scotland sclnp. S 81 Eng. bot. 1003
 2139 collina Host hill *Ji A un* *1 jnjl* Ap Austria 1826. D co Host gr. 2. 66
P. thermalis Pen.

2140 coucina Gnu. neat *Jlh A* ... S. France 1817. D co

2141 flexuosa E. B. zigzag *Jlh A* ... Scotland scalp. S h.l Eng. bot. 1123

2142 distichophylla Gail. distiuh-leaved *Jlh A* ... Austria 1819. D co Host gr. 4.26

2143 laxa Hoe. *joosc-spikcd* *Jlk A un* *fjn.il* Ap Germany 1800. S ro Host gr. 3.15

2144 Hallcrirto R. & S. *llaller's* *Ji A un* *ft jnjl* Ap Vallcsia 1821. D co

2145 cic'sia JR. grey *Jlh A w* ... Scotland scino. S sl Eng. bot 1719

2146 urachiphylla PA. short-leaved *Jh A un* *1 J>jl* N. Amer. 1816. S co

2147 viviparous viviparous *Jh A un* *ijnjl* Switzerl. 1800. S co Fl. dan. 807

2148 trivialis /. common *Jh A ag* *H jnj.ap* Britain me |>. S h.l Eng. bot. 107?

2149 pratensis L. meadow *Mi A ag* *1 my.jn* Ap Britain me pa. S sl ling. bot. 17>*

2150 angustifolia W. narrow-leaved *Jlh A ag* *2 jnj.ap* Germany ... S cp l'ee her. 6 3

2151 humile E. B. humble *Jlh A w* *ft my.jn* Ap Britain mc.pa. S si Kng. Lot. J004

2152 annua L. annual *Jh O w* *mr.o* Ap Bnta.n pas. S hi Eng. bot 11H

2153 coespitosa Forst. tufted *Jh A un* *1 jnjl* Ap N. Zeal. 1822. S co

2154 badensis //AA Baden *Jh A un* \ Ap Baden 1800. S co Host gr. 2. (II)

2155 sudetica Hae. Swedish *Jh A un* *3 jl.au* Ap Germany 1802. S co Host gr. 3. Id

2155 hybrida C.M. hybrid *Jh A un* *WJ* Ap Europe 1819. D co

2156 Gmelini Koch. Gmehn's slender *Jh A un* *i jnjl* Ap Germany 1818. D co

2157 gracilis Desf. slender *Jh A un* *i jnjl* Ap Switzerl. 1826. D co

2158 cunisia All. Mount Cenis *Jh A un* \ Ap Mt. Ccnis 1791. S co Host gr. 3.1

2159 rufa L. yellow *Jh A un* *Ujl.au* Ap N. Amer. 1604. S co

2160 hirsuta Aff. hairy *Jh A un* *1 jnjl* Ap N. Amer. 1819. D co

2161 serotina Ehrh. late-flowering *Jh A un* *2 jl.s* Ap Germany 1800. S co Lee. her. 6. X

2162 effusa Kit. marthi *Jh A un* *1 jl.au* Ap Pannoma Ib22. D co Vil. stras. 2. ff

2163 vridis Mhl. open *Jh A un* *1 jnjl* N. Amer. 1812. S co

2164 tv/ifOffo rmis Host green *Jh A ag* *5 jnjl* AP Dalmatia 1800. S co

2165 atrovirens liesf. Festuca-form *Jh A un* *2 jl B* S Kurope 181< 1) co Host gr. 3.17

2166 curvula Hum. dark-green *Jh A un* *1 my.jn* Ap Kurope 181! 1) co Dest. at. 1.17

2167 abyssinica // K. iiin vel utc *Ji o un* *1 my.jn* Abyssinia 1775. 1) CO

2168 capularis /. AbysHimn *Mi o un* *4 au.o* N. Amer. 1781. S co JAC. ic. 1.17

2169 Poiretta K. & S. hait-panicled *Jh A un* *4 jnjl* Europe 1821. S co M. h. 3.6.33

2170 chinensis Retx. Poiret's China *Jh o un* *1 jnjl* China 1820. D co

2171 austriensis R. Br. Australian *Jh Al un* *1 jnjl* N. Holl. 18Z2. S co Bur. in. 11. 3

2172 plebeian Lab. *Jlh A un* *1 jnjl* S co Lab. n. h. 1. !*

2173 mucronata Beauv. plebeian *Jlh 1A1 un* *1 jnjl* Ap N. Holl. 1892. S co

2174 Molineri Balb. mucronate *jut* (23 un) || jnjl Ap Guinea 1824. S co Beau. ow. 1.4

2175 supina Schr. Moliner's *M A un* *1' jnjl* Ap Italy 1807. S co Bal. mis. 5. 1

2176 mulicea Kit. supine *Jlh A un* *1 jnjl* Ap Switzerl. 18V2. D co Host gr. 4.27

2177 spicata L. Millet *Jlh A un* *2" jnjl* Ap Europe 1821. 1) co

2178 Ciudfni R. & R. spiked *Jlh A un* *1 jnjl* Ap Spain 1826. 1) co

2179 Rterilis Bub. barren *Jlh A un* *1 jnjl* Ap Switzerl. 1818. D co

2180 angustifolia R. Br. narrow-j/v&ai *Mi A un* *1 jnjl* Ap Tauria 1821. S co

2181 tcnax Lk. tough *Jlh A* *2 jl.au* Ap Melv Isld. 182a S co

2182 comprefisa L. *Ji O* *1 jnau* Ap 1817. S co Eng. bot. 505

2183 glauca E. B. glaucous *Jlh A* *2 jn.au* Ap Britain walla. S 8.1 Eng. bot. 505

2184 nomorahs L. wood *J> A* *1 iti* Ap Britain moun. S B.I Ebn. bot. 17

2 angustifolia Sin. narrow-leaved *Jlh A* *1 mr.jn* Ap Britain woods. S h.l iing. bot. J2tf

9185 depauperita Kit. starved *Jlh A* *1 jnjl* Ap Pannonia 1819. 1) co (Jr. wo. 182. ic.

£251 domingensis Lk. 2252 filiformis Beauv. <i>P6a chinensis L.</i>	St. Domingo filiform	JukJZDpr M O pr	3 Jn 2 jl.au	Ap Ap	W. Indies China	1820. 1820.	S co S co	Jac. ic. 22 Jac. ec 4	
2253 cynosuroides A. AS. 2254 prdcera J?.&	Cynosurus-like tell-	M .Qj pr JkESp	ft 4 jl	my.jn Ap	E. Indies Brazil	1824. 1823.	S co S GO		
263. CY'NODON Pers. 2255 Dáctylon Pers. Panicum Dáctylon E.B. 2indica	CTNODOW. {Kuon, a dog, odtnu, a tooth.)				Graminea?	5.—10.			
2256 linearis W. en. 2257 stellatus W. en. 2258 virgatus W. en. 2259 pr ^c ox JL*&	Dactylon linearis stellate twiggy early	4 A w M O un Jk O un Jk O un	1 jl 1 jnjl 1 jnjl 1 jnjl	Ap	England EL Indies St.Helena E. Indies	... 1796. 1819. 1819.	S co S co S co S co	Epig. boL850	
264. ACTINO'CHLOA W. 2260 procumbens Desv. Chondrosium procumbens Desf.	ACTINOCHELLOA. procumbent	Jk O un	{Aktin% a ray, chlo, grast.)	Ap	Philippine	1816.	S co	Jac. ec. 12	
2261 tenuis W. Ghadrismum tenuis Kth.	slender	Jk O un	1 jLau	Ap	Mexico	182a	S co		
2262 prostrata Lag. 2263 hirsuta Lag.	prostrate hairy	Jk O un Jk O un	JjLau 1 jl.au	Ap Ap	Mexico Mexico	1820. 182a	S co S co	H.&B.1.57	
265. DINE'BRA Beauv. 2264 arabica Beauv.	DINBBRA. Arabian		(Its Arabic name.)			Gratiniae.	1.—5.		
2265 Lima Trin. Dinebra Lima Beauv.	Lima	in O w	JjLau	Ap	Spain	1776.	S co	Cav. ic. 1.91	
267. ECHINA'RIA Ztou 2266 capitata Desv.	ECIUXARIA. headed	Mi Q cu	} my.au	Ap	S. Europe	1771.	S co	Graminea. 1.—	
	Cenchrus capitatus Sm.	Sesleria	echinata Host					Host gr. 3.8	
268. PAPPO'PHORUM Schreb. 2267 nigrans R Br. 2268 phleoides Lag.	PAPOPHORUM. black	Jl • Al un	2 jl.au	Ap	N. Holl.	1822.	S p.l	Graminece. 2.—7.	
	Phleum-like	M C3 un	2 jl.au	Ap	S. Amer.	1822.	S p.l		
269. TRI'TICUM X. 2269 sstivum L. 2270 hybrlrum L. 2271 sibericum W. 2272 compositum L. 2273 Linneendum Lag.	WHEAT. summer	Jlk 0 ag	■	{Tritum, rubbed; ground into flour.)	Baschkiros	■	S rm Host gr. 3.26	Graminete. 30.—	
	winter	Jlk 0 ag	4 jnjl	A p			S rm Host gr. 3.20		
	Siberian	M O Clt	5 jnjl	A p	Siberia	1800.	S co		
	composite	Jvk O ag	3 jnjl	Ap	Egypt	1799.	S co		
	Linexan	Jlk O Clt	3 jlau	A p	Spain	1820.	S rm M.h.ai.7		
	turgid	Jlk 0 ag	3 jnjl	A p			S co		
	fastubsum Lag.	Mi O clt	3 jlau	Ap	Spain	1820.	S co		
	proud	Mi O Clt	3 jlau	Ap	Spain	1824.	S rm Host gr. 3.28		
	Gxrtnerifinum Lag.	M O Clt	3 jlau	Ap	Spain	1821.	S co		
	platysachyum Lag.	M O Clt	3 jlau	Ap	Spain	1820.	S co		
	broad-spiked	M O Clt	3 jlau	Ap	Spain	1820.	S co		
	cochlearie Lag.	aheW-seeded	M O Clt	3 jlau	Ap	Spain	1820.	S co	
	d6rum Desf.	stiff	Ml O Clt	3 jlau	Ap	S. Europe	1820.	S co	
	T. vill6sum Host								
2280 sicutum A. * S. 2281 Aordeffirme Host	Sicilian	Jlk 0 Clt	3 JLau	Ap	Sicily	182a	S co	Host gr. 4. 6?	
2282 compactum Host	Barley-form	Jlk 0 Clt	S jlau	A P	Austria	1819.	S co		
2283 Cevalloa Lag.	compact	Jlk 0 clt	3 jlau	A P	Austria	1819.	S co	Host gr. 4.5	
2284 polonicum L.	Cevallos	Jdk O Clt	3 jnjl	A p	Austria	1819.	S co	Host gr. 4.7	
2285 SpMa L. Zea Host	Polish	Jlk 0 ag	4 ■	A P		162.	S co		
2286 monococcum L.	Spelt	Jlk 0 ag	3 jnjl	Ap			S rm Host gr. a 21		
2287 atratum Host	one-grained	Jlr O ag	3 jnjl	A P			S rm Host gr. 3.30		
2288 Cienfuegos Lag.	dark-headed	Jlk O Clt	3 jlau	A p	Austria	182a	S rm Host gr. a 33		
2289 Bauhn's Lag.	Cienfuegos	Jlk 0 dt	3 jlau	A p	Spain	1821.	S co	Host gr. 4.8	
2290 dicoccum SchU. 1 album 2rufum	Bauhn's	Jlk 0 clt	3 jlau	Ap	Spain	1821.	co		
2291 tricoccum SchU. 9292 venulicum Ser.	two-seeded	M O clt	3 jnjl	A P			co		
2293 vill6sum Beauv. Sec&le villdsum L.	white-seeded	Jlk O dt	3 jnjl	Ap			co		
2294 squarr6sum Roth	rufous-seeded	Jlk O dt	3 jnjl	A p			co		
2295 hispanicum W. 2296 creticum R.SS.	three-seeded	Jlk O dt	3 jajl	Ap			co		
2297 campstre Kit.	small, veined	Jlk O Clt	3 jnjl	Ap	Egypt	1800.	co	Sch. ce. 17.3	
	villous	juk O clt	3 jnjl	Ap	S. Europe	1819.	co		
							FLgr. 1.97		
270. AGROPY'RUM Beauv. 2298/dnceum Beauv.	AOROPYBUM. rushy	M f ■	■ f ■	{Agrof, a field, pyros, wheat.)	Gramine*. 26.—69.				
7Yticum panicum L.		M f ■	■ f ■	Ap	England sea-sh.	S co	Eng. bot 814		
2299 rtpens R. \$ S. Triticum r6pens L.	creeping	Mi A w	2 jlau	Ap	Britain	rub.	S m.s Eng. bot 909		
1 ar'vense R. & S. 2 subulatum R. if 5.	corn field	Mi ■	■ jlau	Ap	Britain		co	Trat tab. 5(tf g)	
3 dumet6rum#.*S.	subulate	Mi A W	1 jlau	A p	Britain		co	Schreb. gx. 'n&'	
4 VaillantM>i/mR. & S.	thicket	Jlk A w	jljl au	A p	Britain		co	Schreb.gr.iii.*;	
5 Leerudnum R. & S.	Vaillant's	Jlk A w	jljl au	A p	Britain		co	Srhrch. gr. 2*	
6 capillare<.*S.	Leers's	Jlk A w	1 jlau	A p	Britain		co	Lee. her. 12.4	
2300 dasyanthus R.SS.	capillary	Jlk A w	1 jlau	A p	Britain		co		
2301 distichum R. \$ S.	thick-flowered	Jlk A w	jljl au	Ap	Russia	1820.	co		
2302 acu turn R. \$ S.	two-rowed	Mi A w	jljl au	A p	Egypt	1816.	co		
2303 Rlaucum R.fyS.	acute-flwoered	Mi A un	jljl	A p	France	1818.	co		
2304 obtusiflorum R. Sf S.	glaucous	Jlk A un	jljl	Ap	France	1819.	co		
2305 densefloridum R. S.	obtuse-flowered	B A un	1 jnjl	Ap	France	1823.	co		
2306 densiflorum R. \$ S.	dense-flowered	Mi A un	1 injl	Ap	Siberia	1823.	co		
£307 muriratum R. \$ S.	muricated	M A un	Ujnjl	Ap	Siberia	1826.	co		
£308 sibircum R. Sf S.	Siberian	Jlk A un	1 my.jn	AP	Siberia	1820.	co		
£308 prostratum R. AS.	prostrate	* A un	1 my.jn	Ap	Tauria	1819.	co		
2309 angustifolium A. * S.	narrow-leaved	Mi A un	2 my.n	Ap	Siberia	1824.	co	N. c.g. 14.1*	

Host gr. 4.10

ORDER II.

TRIANDRIA DIGYNIA.

271. LO'LIUM L.	RYE GRASS.	{ <i>loloa</i> , <i>Celt.</i> , <i>lolium</i> , I ^a (at)}	Gramine*	7.-11.
XB4 perenne L.	perennial	<i>M</i> A ag [1] my.jn Ap	Britain	me pa. S
<i>lvigileRAS\$.</i>	common	<i>jib</i> A ag U myjn Ap	Britain	co en <i>v.</i> , bot. 315
2 tenu <u>A</u> -S.	slender	<i>M</i> A ag If myjn Ap	Britain	co Host gr. 1.25
3 compositum R. A.S.	composite	<i>Jk</i> A a# [1] myjn Ap	Britain	co Gr. wo. 211. ic.
4 ramosum R. A.S.	branchy	<i>m</i> A ag [1] i myjn Ap	Britain	co Sen. ag. 1.7. D.
0 <i>RwsleUdnum</i> Sin.	Sin. Russell's	<i>jd</i> A *A IJ myjn Ap	Britain	co Lee. gr. 12.1
6 <i>Whitw6thiSin.</i>	Whitworth's	<i>M</i> A ag <i>Uljmyjn</i> Ap	Britain	co Gr. wo. 216. ic.
<i>J Stickneydnum</i> Sin.	Stickney's	<i>M</i> A a? U myjn Ap	Britain	co
>paniculatu <i>m</i> Sin.	paniculate	<i>M</i> A *f Uinyjn Ap	Britain	co
9 monstruoso Sin.	monstrous	<i>M</i> A ag [14] myjn Ap	Britain	co
10 viviparous R.A.S.	viviparous	<i>M</i> A [1] myjn Ap	Britain	co
11 multiflorum Pers.	many-flowered	<i>M</i> A myjn Ap	Britain	co
oqoe ¹² h ? m l e K	humble	<i>A</i> A *i myjn Ap	Britain	co
l ¹³ multi <u>fl</u> rum Lam.	many-flowered	<i>JU</i> A un 2 myjn Ap	France	182& S
*326 speciosum Stev.	showy	<i>M</i> A un 2 myjn Ap	Tauria	1824. S
2327 tenuu L.	slender	<i>M</i> A un 3 <i>jla</i> Ap	S. Europe	1591. S
23 ³ temul'tnum L.	drunken	<i>Darnel M</i> OpP 4 <i>jla</i> Ap	Britain	cor. fi. S
2329 arveus With.	corn-field	<i>Mi</i> OpP 2 Jl Ap	England	cor. ft S
2330 rigidum Gau.	rigid	<i>jii</i> [1] myvn Ap	SwiftierL	1820. S 00

LVME GBASS.	(According to Linnaeus, <i>etc/o</i> , to cover)	Graminece.	23.—25.
sand	A ag 4 apjn Ap	Britain seaco. S s Eng. bot. 1872	
jointed	A ^a apjn kjl Ap	England sea sh. S g Eng. bot 1286	
sandy	JU A un kjl AP	Siberia 1806. S co	
gigantic	Jib A or ^a au Ap	N. Amer. 1790. S co	
Mexican	Jib A cu 3 ^a au Ap	Mexico 1833. S co	Jac ec. 19
Siberian	Jib A un 6 injl Ap	Siberia 1758. D co	Schreb.gr.2.21.1
racemose	Jib A un Sii ^a jl Ap	Siberia 1826. S co	
tender	1 A un 3 injl Ap	Siberia 1801. S co	
Philadelphia	JH At A un 4 jl.au Ap	N. Amer. 179a S co	
Canadian	M A un 4 jl.au Ap	N. Amer. 1699. S co	M. h. 3.2.10
glaucous-lvd	Jib A un 4 injl Ap	N. Amer. 1819. D co	
Virginian	Jib A un 3 injl Ap	Virginia 1781. S co	
striated	JVb A un 3 injl Ap	N. Amer. 17W. S co	
villous	^ A un Ap	N. Amer. 1802. S co	
Eurojean	Jib A un Ap	England woods. S.s.1 Eng. bot 1317	
long-awned	Jiu O cu Ap	Smyrna 1806. S co	Schreb.gr. 24.3
intermediate	Jtt O un Ap	Caucasus 1823. D co	Bux. c. 1.52
Medusa's head	Jib O un Ap	Portugal 1784. S co	Schreb. gr. 24.2
Rush-like	JU A un Ap	Siberia 1806. S co	M. irop. 1.45
Barley-like	Jib O un 3 my. injl Ap 1822. D co	
hystrnxZ.	MI A CU 2 ^a jl Ap	Crimea 177a S co	Jac. ic. 2.303
^a P ^a Iahf, trix, etc TM .		N. Amer 181a S co	
P pseudo-h. fitrix R S	bastard poror* Q cu 1 injl Ap		

273. SECALE L. 2353c ^{er} alez.	RYE. corn winter	(An ancient name derived from <i>sreō</i> , to cut)	Gramineae. S.— al Host gr. 2.48 S co Black 424 S co Host gr. 2.48
		jib@ clt 3 jnj! Ap Crimea	
		Jik CD Clt 3 jnj! AP Tauria	
		jib O clt 3 jnj! Ap Tauria	
		jib O clt 2 W A.P. Tauria	
		M O clt S jnj! Ap Tauria	
		S jnj!	
originate If. Triticum orientale	oriental <i>Pisak</i>	jib O clt 3 jnj! Ap Levant	1807. S co W. n. ber. 2.4.3

	THTKAFOGON.	{Tetra. four, pogon, a beard.)	Graminis.	1.	
ous	Jb O cu	1 Jl.au	Ap	Barbary 181& S co	Desf. at a 255
		(According to Bodjeus a Slapel, <i>hordus</i> . heavy.)			
giant	Jb O clt	3 jl	Ap	Sicily ... S	r.m Host gr. a 34
celestial		4 Jl	Ap	Levant ... S	co
black			Levant	... S	co Vib. ce. 1
six. rowed			& Europe 1818.	S	co
two-rowed			Tartary	—	r.m Hostgr. asft
beardless			Tartary	—	r.m Hostgr.3.36
naked			Tartary	—	co
Spelt Barley			S. Europe 1819.	S	co Ard. sag. 2. 4
flattened			Europe	...	r.m Host gr. 3.37
Rye-like			Nepal 1817.	S	
Nepal			Italy 1770.	S	co Mo.gr. 60.96
bulbous			S Europe 1824.	S	
strict			Britain sal. m. S	S	Pl.gr. 1.98
wall			T G. H. 1817. S	S	co Desf. at 1.37
cape			Britain m.me. S	S	co Eng. bot 1971
pratense	Aid				h.l Eng. bot. 400

- 2372 marftimum *With.* Mi O W 1 jn.jl Ap Britain sal. m. S m.8 Eng. bot. 1205
 2373 hfstrix *Roth* porcupine M O CU 1 jn.jl Ap Spain 1821. S* co
 2374 juba' tumZ., maned M Q> CU 1 jl.au Ap N. Amer. 1782. S co
276. MICRCCCHLOA *R. Br.* MICBOCHLOA (*Mikros*, small, *chloa*, grass.) Graminea. I.
 2375 Betticea *R. Br.* setaceous M Q c*v*i i jl Ap E. Indies 1806. S co Box. cor. 132
277. JEGOPCKGON/Jiim. JECOPOGON (*Aix*, a goat, *pogon*, a beard.) Graminea!. 2.-5.
 2376 puHillis *Beauv.* small Jb O cu 4jl.au Ap S. Amer. 1822. S co Cav. h. m. 1.5.2
 2377 trisetus *Lag.* three-bristled M O cu 1 jl.au Ap S. Amer. 1822. 8 co Cav. h. m. 1.5.3
278. OPHIU'RUS *Beauv.* HARD GRASS. (*Ophis*, a snake, *oura*, a tail.) Graminea. 3.-4.
 2378 incurv&us *Beauv.* incurvate i Q c u f jl Ap Britain sea co S ms Eng. bot 761
 Rottboellia incurvata L.
 8379 filiformis *Beauv.* filiform Jb O cu | jl Ap Portugal 1800. S co Bar. ic. 117.1
 RottbocWo filiformis *Roth*
- 2380 subultus *Beauv.* subulate Jk O cu 1 jl Ap S. Europe 1806. S s.1 Bar. ic. 5
 Leptfrus cylindricus *Trin.* Rottbocilia cylindrica W. Moncma subulfita Beauv.
279. ROTTBOE'LLL* L. ROTTBOELLIA. (*C. Rottboett*, a Danish botanist) Graminea. 2.—
 2381 exalt&t*a* L. exalted M EJ cu 8 ... Ap E. Indies 1806. S co Rox. cor. 2.157
 2382 perfor&a Rox. perforated Jb ES cu 3 ... Ap E. Indies 1822. S co Rox. cor. 2.182
280. PHOLIU'RUS *Trin.* PiroLiuaua (*Pholis*, a scale, *euro*, a tail.) Graminea?. 1.
 2383 pannbmcus *Trin.* Pannonian Jb O cu T jl Ap Hungary 1804. S co Host gr. 1/24
 Ophiurus pannonicus Beauv.
281. ANTHE'PHORA *Schreb.* ANTHIEPHORA. (*Anthos*, a flower, *phoreo*, to bear.) Graminea. 1.—2.
 2384 elegans *Schreb.* elegant JlibOlcu 2 au.s Ap Jamaica 177& S pl Schreb.gr. 41
 Tripsacum hermaphroditum L. Cenchrus laevigatus *Trin.*
282. MANISU'RUS *L.* MANISURUS. (*Manis*, the scaly lizard, *oura*'a tail; spikes.) Graminea. 1.—2.
 2385 granularis *Sum.* grained MU^cu 2 jl.au Ap E. Indies 1821. D co Rox. cor. 2.118
283. PE'ROTIS *H. K.* PEROTIS. (*Peros*, deficient; flower.) Graminea. I.—2.
 2386 latifolia *H. K.* broad-leaved 1 O c u 2 au.s Ap E. Indies 1777. S s.p R. mal. 12.62
284. SA'CCHARUM *L.* SUGAR CANS. (*Soukar*, its Arabic name.) Graminea. 10.—32
 2387 officin&rnum *L.* officinal ■■■■■ 12 -- Ap India 1597. Sk r.m Sl. jam. 1.66
 2388 violaceum *Tus.* violaceous ■■■■■ clt 10 jl.au Ap W.Indies 1824. D co R a n t 1.24
 2389 sinense *Rox.* Chinese ■■■■■ clt 6 ... Ap China 1822. D r.m
 2390 prbcerum *Rox.* tall ■■■■■ 10 ... Ap E. Indies 1822. D r.m
 2391 contractum *Pair.* contracted Jb ES cu 4 n.jl Ap Trinidad 1823. D co
 2392 caudatum *Mey.* caudate ■■■■■ 4 ... Ap W. Indies 18KL D r.m Sl. jam. 1.70.1
 8393 polystachyum *Swz.* many-spiked JllhAicu 2 jl.au Ap Trinidad 1823. D co
 8394 mexicachum *Brou.* Mexican ■■■■■ 1 jl.au Ap Mexico 1820. S co
 8395 dftium *H. if.* doubtful Jib ES cu 2 jl.au Ap Trinidad 1826. D co
 8396 Mnja *Rox.* Munja ■■■■■ 4 ... Ap E. Indies 1805. D r.m
- 285..ERIA'NTHUS Mr. ERIANTHUS. (*Erion*. wool, *anthos*, a flower.) Graminea. 2.-34.
 2397 saccharoides *Mx.* Sugar-cane-like Jib A P* 5jl.au Ap N. Amer. 1826. 1) Lp
 Saccharum gigantum Pers.
 2398 brevibarbis *Mx.* shorUbcarded ilk A cu 4 jlau Ap N. Amer. 1822. D pl
 Saccharum brevib&rbre Pers.
286. TRIC HOLJE'N A R. % S. TRICHOL.*NA*. (*Thrix*, hair, *lana*, a cassock.) Graminea. 1.
 2399 micrantha *R. % S.* small-flowered Jb iAl pr 1 jn.jl Ap Teneriffe 1825. S pl Jac.ec. 34
 S&ccharum Tenerifte L. Pinxcum Teneridie Spr.
287. IMPERATii Cyr. (*Ferrante Imijerati*, a Neapolitan botanist of the 16th century.) Graminea. 2.—
 2400 arundinacea *Cyr.* reedy Jb A ec 2f jl.au Ap S. Europe 1817. S co Cyr. ne. 2.11
 2401 spontanea *Beauv.* spontaneous JlliEJpr 3 jl.au Ap Africa 1824. D co R.mal 12.46
 Saccharum spont'ncum L.
288. LEF/RSL4 Swz. LEERSIA. (J. D. Leers, a German botanist.) Graminea. 3.—
 2402 oryzoides *Swz.* Rice-like Jb A cu 2 jlau Ap Levant 1793. S co Host gr. 1.35
 2403 virginica *W.* Virginian M A CU 1 J₂ jl.au Ap N. Amer. 1770. S co Jac. ic. 2.305
 2404 austriulis *R. Br.* Australian M iAJ un 1¹ jl.au Ap N. Holl. 1819. D pl
289. ARUNDINA'RIA Mx. CANE-BRAKE. (Altered from *arundo*, a reed.) Graminea, 2.
 2405 macrospérma *Mx.* long-seeded Jib A or 10 jn Ap N. Amcr. 1809. S co
 2406 glaucescens R\$S. glaucous M E cu 10 ... Ap E. Indres 1818. D pl
 Lud61ifa glaucescens W. Panicum glaucescens Lam.

CLASS III. ORDER S.

TRIA'NDRIA TRIGY'NIA.

- * 290. //OLCSTEUM *L.* HOLOSTEUM. (*Holos*, all, *osteon*, a bone; by antiph. no bone.) CarttonhHttea 2—4
 £407 umbellitum *L.* umbellated O P' i jlau Pk "England old wa. "S co" Eng. bot S?
 2408 diandrum *Swz.* diandrous O u n i jn.jl W Jamaica 1824. S co Swx. pr. 7
291. POLYCA'RPON *L.* ALL-SEED. (*Polys*, many, *karpos*, seed.) Paronychiece. 2.-3.
 2409 tetnphf *Hum L.* four-leaved O cu i jl W England san. pi. S co Eng. bot. 1031
 2410 diphylum *Cav.* two-leaved O cu i jn.jl W Spain 1821. S co Cav. ic. 151.1
- 292 LE^CHEif *L.* LECHEA. (G. Leche, a Swedish botanist, died 1764.) Cistinea. 4.-6.
 2411 major *Ph.* greater k APT 1 jl.au W Canada 1780. D co Lam.il. 1.52. 2
 viju^ga&/i. minor *L.* lesser ■■■■■ ijl.au W Canada 1802. D co Lam. il 52.1
 2412 minor *Ph.* Thyme-leaved ■■■■■ * jl.au W N. Amer. 1812. D s.p
 2413 thymifolia *Mt.* minor *Wai.*
 2414 racemulosa *Mx.* racemulose ■■■■■ A pr iil.au W N. Amer. 1816. D s.p Lam. il. 3.2H1
- * 893. ERIOCAU'LON Lam. PIPEWORT. (*Erion*, wool, *kaulos*, a stem.) Eriocatlecc. 3.-34.
 8415 septanguUre E.B. aeven.angied * A cu % W Scotland bogs. D aq Eng. bot 775

- | | | | | | | | |
|------------------------------|-----------------|---|---|---------------------|-----------|-------------|-------------------|
| 2416 austral* <i>Jl. Br.</i> | Australian | I Al | 1} jn | W | N. Holl. | 1820. | D bog |
| 2417 fascicul&um <i>Lam.</i> | fascicled | O c u | i au.8 | W | Guiana | 1825. | S bog Lam. il. 50 |
| 294. MO'NTII! L. | CHICKWEED. | { <i>Joseph Monti</i> , professor of botany, at Bologna.) | <i>Portulaceae</i> . 2. | | | | |
| 2418 fontana L. | fountain | £ O w | 1 ap. ray | W | Britain | springs. S | aq Eng. bot 1206 |
| 2419 rivulari&um Qm. | Irook | * O un | i jh.jl | W | Labrador | 1823. | D aq |
| 295. AFOLLU'GO L. | MOLLUGO. | (A name in Pliny, retained by Linnaeus.) | <i>CaryopkQUE*</i> . 5.—14. | | | | |
| 2420 verfracta L. T. | whorled | O un | I jl | Ap | Virginia | 1748. S co | Ehr.pic.6 |
| 2421 triphylla Lk. | three-leaved | O un | jl.au | W | Brazil | 1821. S ms | |
| Linkii Dec. | | O un | jn.jl | W | E. Indies | 1818. S co | Pluk.al.21.75.6 |
| 2422 oppositifolia L. | opposite-leaved | O un | jl.au | W | Brazil | 1826. S co | Sen. mo. 6 |
| 2423 Schrankfi Scr. | Schrank's | O un | jn.jl | W | | | |
| dichotoma Schrank | | O un | i jn.jl | W | C G. H. | 1824. S co | |
| 2424 hirta Thun. | hairy | O un | i jn.jl | W | | | |
| 296. MINUA'RTJML. | MINUARTIA. | { <i>Minuart</i> , a Spanish apothecary.) | <i>Paronychite</i> . 3. | | | | |
| 2425 dichotoma L. | dichotomous | O cu | i jn.jl | Ap | Spain | 1771. S co | Ac. st. 1758.1.2 |
| 2426 campestris L. | field | O cu | lin jn.jl | Ap | Spain | 1806. S co | Ac. st. 1758.1.3 |
| 2427 montana L. | mountain | O cu | jn.jl | Ap | Spain | 1806. S co | Ixef.it. 1.4 |
| " 297. QUE'Rit L. | QUERIA. | { <i>Don J. Quer y Martinez</i> , a Spanish surgeon.) | <i>CaryopkOllea</i> . 1.—2. | | | | |
| 2428 hispanica L. | Spanish | O cu | lin my.s | Ap | Spain | 1800. S co | Qu. his. 6.15.2 |
| J g 8. KGBNI'OLF L. | KCENIOIA. | { <i>Samuel Katnig</i> , a Swiss mathematician^ | <i>Polygfmeae</i> . 1. | | | | |
| 2429 islfodica L. | Iceland | O cu | ftap | Ap | Iceland | 1773. S co | Lara. il. 51 |
| 299. PROSERPINA'CA L. | PROSRPINACA. | { <i>Pnaero</i> , to creep.) | <i>Hatoilgete</i> . 2. | | | | |
| 2430 pattstria L. | marsh | & O cu | 1 jl | W | Canada | 1818. D aq | |
| 2431 pectinata Ph. | pectinated | & O cu | jl.au | W | N. Araer. | 1822. S aq | |
| *300. ELODE'A Mx. | ELODEA. | { <i>Elodes</i> , growing in marshy places.) | <i>Fluvidles</i> . 1—3. | | | | |
| 2432 guyanensis Mt. | Guiana | -O c u | ijl.au | W | Guiana | 1820. S aq | M. in. 1811.4.1 |
| 301. HOUTTU'YNLf | Thun. | HOUTUTNIA. | { <i>Dr. Houttuyn</i> , a virtuoso of Amsterdam.) | <i>Aroidae</i> . 2. | | | |
| 2433 cordata Thun. | | £ CS3 | 1 my | jl Y.o | Japan | 1820. D s.l | Bot mag. 2731 |
| 2434 fce'tida Hart. | foetid | £ E3 cu | my.au | Y.o | Japan | 1800. D s.l | |

CLASS IV. ORDER 1.

TETRA'NDRIA MONOGY'NIA.

- | | | | | | | |
|---|----------------|--|--|--------------------|----------------------------|-----------------------------|
| 302. PETRCPHILA | <i>R.Br.</i> | PETROPIITLA. | { <i>Petros</i> , a stone, <i>phileo</i> , to love.) | <i>Proteacca.</i> | 8.—10. | |
| 2435 tercifflia <i>R. Br.</i> | round-leaved | • L.J or | 4 jl.au W | N. Holl. | 1824. S s.p | |
| 2 « filiofia Jl Br. | thread-leaved | H u Jor | 5 jn.jl W | N. Holl. | 18'4. S s.p | |
| 2437 rigida <i>R. Br.</i> | rigid | • L.J or | 5 jn.jl W | N. Holl. | 182a S 8.p | |
| 2438 pulchella Jl Br. | neat | ■ L.J or | 5 jl.au W | N. S.W. | 1790. S s.p Bot mag. 796 | |
| Pr6tea <u>wcfis<i>lia</i></u> <i>Sal.</i> | | | | | | |
| 2439 dveref <i>lia</i> <i>R. Br.</i> | various-leaved | ■ i_Jor | 5 ... | N. Holl. | 1803. S s.p | |
| Z440 pedunculata <i>R. Br.</i> | pedunculated | ■ L.I or | 4 jl.au | N. Holl. | 1824. S ip | |
| 2441 fastigiata Jl Br. | fustigiate | ■ u.J or | 5 jl.au | N. Holl. | 1820. S s.p | |
| 2442 trifida Jl Br. | tntid | ■ J ■ | 4 jl.au | N. Hull | 1820. S s.p | |
| 303. ISOPOGON Jl. Br. | ISOPOUPON. | { <i>Ios</i> , equal, <i>pogon</i> , a beard.) | | <i>Proteadete.</i> | 12.— | |
| 2443 teretifolis Jl. Br. | round-leaved | • L.J or | 4 my.jn Pa | N. Holl. | 1823. S s.p | |
| 2444 anethifolius <i>R. Br.</i> | Dill-leaved | • L.J or | 5 my.jn Pa | N. Holl. | 1796. S s.p Cav.ic.6.549 | |
| 2445 formbus <i>R. Br.</i> | handsome | * L.J or | 4 mr.jn Pk | N. Holl. | 1805. S «p» | |
| S446 anemonifidhus <i>R. Br.</i> | Anemone-lvd | il L.J or | 5 jl.au Y | N. Holl. | 1791. S s.p Bot mag. 697 | |
| 2447 axillaris <i>R. Br.</i> | axillary-Jltm* | il L.J or | 4 ap.my Pa | N. Holl. | 1824. S s.p | |
| 2448 ceratophyllus <i>R. Br.</i> | Buckshorn-lvd | • L.J or | 4 my.jn Pa | N. Holl. | 1824. S s.p | |
| 2449 polycyphalus <i>R. Br.</i> | many-headed | • L.J or | 3 my.jn Pa | N. Holl. | 1824. S s.p | |
| 2450 prolnquus <i>Cun.</i> | allied | • i.J or | 3 my.jn Pa | N. Holl. | 1824. S s.p | |
| 2451 divaricatus <i>R. Br.</i> | divaricate | • l.J or | 3 my.jn Pa | N. Holl. | 1824. S e.p | |
| BA <i>Pr</i> 6tea divaricata <i>Andr.</i> | | | | N. HolK | S s.p Bot rep. 465 | |
| 2452 trilobus <i>R. Br.</i> | three-lobed | ■ ■ ■ | 4 my.jn Pa | N. Holl. | 1803. S fl.p | |
| 2453 attenuata Jl Br. | attenuate | ■ ■ ■ | 4 ap.jn Pa | N. Holl. | 1822. S s | |
| 2454 longitius Jl. Br. | long-leaved | ■ i_Jor | 3 jn.jl Y | N. Holl. | 182a C p.l Bot.reg.900 | |
| 304. PROTEA ⁱ L. | PROTEA. | { <i>Proteus</i> , a self-transforming sea god: diversity) | | <i>Proteacea.</i> | 47.—5a | |
| 2455 cynaroides L. | Artichoke-like | • i.J or | U mr.n P | C. G. H. | 1774. <i>Bot</i> mag. 770 | |
| 2 * 56 <i>ti66hJl</i> ■ | broad-leaved | ti _ or | 7 jls. P | C. G. H. | 180a S s 1 Bot. mag. 1717 | |
| — cocinea | scarlet | * i.J or | 7 jls. S | C. G. H. | 180a C s1 | |
| Q ^a em 3 viridiflra | green-flowered | Mi _ or | 7 jls. G | C. G. H. | 1806. C s1 And. rep. C46 | |
| 2457 villiferal/wf. | villiferous | L.J or | 7 jls. P | C. G. H. | 1800. C s1 Bot rrg. 1023 | |
| 2458 compacta Jl. J. | compact | i.J or | 6 jl.au | C. G. H. | 1810. C Up | |
| 2459 longiflora Lam. | long-flowered | i.J or | 7 ja.mr | P | C. G. H. | 1795. C s.l pEx.bot.2.81 |
| 2460 speciosa L. | splendid | i.J or | 2 mr.jn | P | C. G. H. | 1786. S s1 Bot. mag. 1183 |
| 2461 coccinea Jl. Br. | scarlet | N.J or | 5 mr.au | S | C. G. H. | 1824. S s1 |
| 2462 obtusa Kn. pr. | obtuse | L.J or | 10 mr.ap | H | C. G. H. | 1786. C s.l p Bot. rep. 110 |
| speciosa B. M. | | | | | | |
| h^lla R. Br. | large-leaved | ■ i_Jor | 8 mr.au W | C. G. H. | 1824. S 8.1 | |
| ■ JS me alefica R. Br. | handsome | ■ i_lor | 6 my.jn R | C. G. H. | 1789. S p.l Bot mair. 1711 | |
| 2466 hifwita W. en. | blk. & white | ■ J or | 6 mr.jl D.P | C. G.H. | 1786. C alp Bot rSff03 | |
| g J pinensi K.cn. | hairy | ■ J or | 4 mr.au Pa | C. G.H. | 1819. S sd *** | |
| S** lepidocarpum R. Br. | pale | ■ J or | 4 mr.au Pa | C. G.H. | 1819. S a 1 | |
| 2469 ncriifolia R. Br. | scaly-fruited | ■ J or | 6 | f | Bot.wpLaDL : | |
| Olcander-lvd | | ■ J or | 6 | 5 | s.lpBot.rcg.i'08 | |

StfOpulchella it Br.	neat ft	i_J or	3 mr.au R	C. G. H.	1795.	L s.l	Bot. reg. 80
1 ciliata	ciliated ft	i_J or	3 mr.au R	C. G. H.	1795.	C s.l	And. rep. 870
2 glabra	smooth ft	i_J or	3 mr.au R	C. G. H.	1795.	C s.l	And. rep. 877
3 glieci6sa	splendid ft	[J or	3 mr.au R	C. G. H.	1795.	C 8.1	And. rep. 442
2471 pftens R. Br.	spreading ft	L_J or	2 mr.jn W.p	C. G. H.	1789.	C 6.1	Bot. rep. 543
8472 magnifica Kn. pr.	magnificent ft	i_J or	6 mr.jn W	C. G. H.	1789.	S 8.1	Bot. rep. 4d8
8473 longifolia R. Br.	long-leaved ft	i_J or	2 mr.au P	C. G. H.	1798.	S 8.1	Bot. reg. 47
8474 incompta 11. Br.	untrimmed ft	i_J or	3 ap.my W	C. G. H.	1822.	C s.l.p	
8475 umbonellata Kn. pr.	embossed ft	i_J or	7 mr.au W.bk	C. G. H.	1798.	C s.l.p	Bot. rep. 14*
8476 ligulifolia Kn.pr.	strap-leaved ft	i_J or	7 mr.au P	C. G. H.	1798.	C al	Bot. rep. 133
8477 mellifera Than.	honey-bearing ft	[J or	6 my.d Pa.Y	C. G. H.	1774.	S 8.1	Bot. mag. 346
2 alba	white fti	[J or	6 my.d W	C. G. H.	1795.	C s.l	And. rep. ic.
8478 grandiflora A. -Br.	great-flowered ft	i_J or	8 my.jn W	C. G. H.	1787.	S p.l	Dot. mag. 2447
8 marginata	margmated ft	L_J or	6 my.jl W	C.G.H.	1795.	C 8.1	Bot. reg. 539
8479 Scolymus Thun.	Golden thistle fti	[J or	3 myjn P	C. G. H.	1780.	C 8.1	Dot. mug. 698
8480 mucronifolia Sal.	mucronate-lvd fti	[J or	3 jld W	C. G. H.	1803.	C 8.1	Bot. mag. 933
8481 nana A Br.	dwarf fti	[J or	8 apjl Pk	C. G. H.	1787.	C s.l.p	Ex. bot. 144
2482 pmiula if. flr.	pendulous ft	i_J or	2 mr.jn V'	C. G. H.	1806.	C 1p	
8485 tenax A 22/.	tough fti	[J or	2 fmy Pk	C. G. H.	1801.	■	Par. lond. 70
8484 canaliculata R. Br.	channel-leaved ft	i_J or	3 fd P	C. G. H.	1800.	■	Bot. rep. 437
8483 acuminata B. M.	sharji-pointed ft	i_J or	3 mr.jn P	C. G. H.	1809.	■	Bot. mag. K194
8486 acaulis 7%**	stemless ft	[J or	1/ my.s P	C. G. H.	1802.	■	Bot. mag. 2065
8487 lorea R. Br.	thonged ft	[J or	1/ mr.au W	C. G. H.	1824.	S.1	
8488 elongata R. Br.	elongated ft	[J or	4ijlau P	C. G. H.	1820.	■	
8489 angustata R. Br.	naxrovr-leaved ft	i_J or	1 mr.au P	C. G. H.	1824.	■	
8490 revoluta if. Br.	rcvolute-leaved ft	i_J or	1f mr.au P	C. G. H.	1795.	■	
8491 tenuifolia A. jfr.	fine-leaved ft	i_J or	1 my.m ...	C. G. H.	1816.	■	
2492 glaucophylla ifopr.	glaucous-leavedft	i_J or	1 ap.my G	C. G. H.	1806.	■	Par. Ion. 11
2493 Uevis R. Br.	smooth-leaved ft	[J or	1 myjn G	C. G. H.	1803.	■	Bot. mag. 8439
	longifolia Sal.						
8494 scabra R. Br.	rough-leaved ft	_J or	1 my.jl Br	C. G. H.	1809.	■	
8495 ripens Thun.	creeping ft	i_J or	1 jn.au ...	C. G. H.	1800.	■	Wein. 897. a.
8496 turbinate R. Br.	turbinate-flwd ft	i_J or	1 ap.my Pk	C. G. H.	1803.	■	Par. Ion. 108
8497 Scopolendrium R. Br.	Hart's-tongue ft	i_J or	1 -	C. G. H.	1802.	■	
8498 cordiflta 7%ML	heart-leaved ft	L_J or	1/ mr.my P	C. G. H.	1790.	■	Bot. rep. 889
8499 amplexicaulis R. Br.	stem-clasping ft	i_J or	4 ja.mr	C. G. H.	1784.	S p.5	Bot. rep. 469
8500 homilis R. Br.	low-flowering ft	i_J or	1 jn.au Br	C. G. H.	1802.	■	Par. Ion. 67
8501 acer6sa R. Br.	chaffy fti	[J or	3 mr.my Bk	C. G. H.	1803.	■	Bot. rep. 532
	grandiflrum it Br.	great-flowered ft	[J or	4 my.jl Y	G.G.H.	1800.	p.l Par. Ion. 116
	Leucodendron grandiflorum Sal.						
8510 ptherberum it. Rr.	iXovny-leafed	[J or	8 myau Y	C. G. H.	1774.	C s.1	
8511 p&tuliu ii. Br.	spreading	[J or	2 jn.au Y	C. G. H.	1823.	S s.p	
2512 spatulatinu R. Br.	spatulate	[J or	2 jn.au Y	C. G. H.	1825.	S sf	
2513 tomentisum Kn. pr.	cottony	[J or	2 aua Y	C. G. H.	1789.	S p.?	
2514 parile Kn. pr.	matched	[J or	2 a.us Y	C. G. H.	1789.	C fp	
2515 candicans B. R.	whitish	[J or	2 a.us Y	C. G. H.	1790.	S p.l	Bot. rep. 894
	Prteacandicans Andr.						
8516 hypophyllum R. Br.	under-leaved	ft i_J or	1 - .. ■	C.G.H.	1787.	S pl	Pl. man. 440.3
	306. MI METES R.Br. MIMETES.	(Mimetes, a mimic.)			Proteacea.	8. — 13	
8517 hlrta R. Br.	hairy	[J or	3 jn.au R	C. G. H.	1774.	C 8.1	Wein. 4.899. a.
8518 pauciflora R. Br.	few-flowered	[J or	3 jn.au R	C. G. H.	1818.	S Lp	
8519 pah'istris Kn.pr.	marsh	[J or	3 jn.au P	C. G. H.	1808.	C l.p	Boer. 2.194
8520 cucullata it. Br.	cucullate-Zirf	[J or	2 ... P	C.G.H.	1789.	S 8.1	Pluk.al.212.304.6
	Prtea cucullata L.						
8521 Hartogii R. Br.	Hartoge's	:■[J or	5 jnau	C. G. H.	1824.	■ 8.1	Wein. 4.906
8522 divaricata R. Br.	divarirate	[J or	2 jn.8 W	C. G. H.	1795.	■ 8.1	
8523 vaccini6lia Swt.	Vaccinium-lvd	[J or	3 ...	C. G. H.	1800.	■	
	Deastella vaccinifolia Kn. pr.						
\$32& purpurea R. Br.	purple	i j o r	2 nd P	C. G. H.	1789.	C s.l	
	307. SERRUR'M R. Br. SEKRURIA.	(Dr. J. Serrurier, prof. bot. at. Utrecht).			Proteaceae.	34.-46.	
8525 glaberrima R. Br.	very smooth	[J i lor	1 jn.au P	C. G. H.	1825.	S si	
5256 diffVisa R. Br.	diffuse	J ll or	3 jn.au P	C. G. H.	1803.	S s.1	
2527 abrotanifolia Kn. pr.	Southernw.-lvd	:■[J or	4 jnau Pk	C. G. H.	1803.	C l.p	Bot. rep. 122
2528 millef6lia Kn. pr.	thousand-lvd	<■[J or	4 jnau P	C. G. H.	1803.	C Up	Bot. rep. 337
2529 rtremisw/J//Kn.pr.	Wormwood-lvd	:■[J or	5 jn au P	C. G. H.	1789.	C l.p	
2630 pinnata R. Br.	pinnate	<■[J or	1 jn.au Pk	C. G. H.	1803.	S p. I	Bot. rep. 512
8531 arenaria R. Br.	sand	<■[J or	1 jn.au P	C. G. H.	1803.	C s.p	
8538 cyanoides R. Br.	Cyanus-like	'■[J or	1^ jn.au P	C. G. H.	1803.	S pl	Pluk.m.61.345.6
2033 nitida R. Br.	shining	<■[J or	2 jn.au P	C. G. H.	1823.	S s.1	
8534 pedunculata R. Br.	>edunculate	>■[J or	7 in.a P	C. G. H.	1789.	C p.l	Botrep.264
8535 scoparia R. Br.	Broom	■[J or	3 jn.au P	C. G. H.	1809.	C p.l.s	
2536 Niveti R. Br.	Niven's	■[J or	1 jn.au P	C. G. H.	1800.	S 8.1	
	Prtea decumbens Andr.				C sp.1	Bot rep. 349	
2537 villosa R. Br.	villous	■[H or	2 jnau P	C. G. H.	1883.	S si	
2538 teniculacea R. Br.	fennel-leaved	ii lor	2 jn.au P	C. G. H.	1880.	S 8.1	
2539 ciliata R. Br.	ciliated	■[n or	2 jn.au P	C. G. H.	1803.	C 8.1	
2540 congata R. Br.	heaped	■[l or	2 jnau P	C. G. H.	1820.	S 8.1	
2541 saurarr6si R. Br.	squarerose	■[l or	2 jnau P	C. G. H.	1810.	S 8.1	
SH2 shylcoides it? Br.	Phyllica-like	■[l or	3 jn.au P	C. G. H.	1788.	■ p.1	Bot rep. 507. *
2543 se'mula R. Br.	rival	■[l or	3 jn.au P	C. G. H.	1803.	C l.p	
544 flonda R. Br.	flowery	■[l or	2 jn.au P	C. G. H.	1824.	S al	Thun. pr. 1.1
8545 pteris Kn. pr.	matched	■[l or	2 jn.au Pk	C. G. H.	1803.	C p.l	Bot rep. 507
2546 odorata Swt.	sweet-scented	■[l or	2 jn.au Pk	C. G. H.	1803.	C p.l	Bot. rep. §45
	Vrolea abrotaniCSlia odorata Andr.						

ORDER 1.

TETRANDRIA MONOGYNIA.

2547 ematginsta <i>Suit.</i>	emarginated	LJ or	• 2	jn.au	Fk	C.G.H.	1800.	C p.1	Bot rep. 536	
» 4 Serrurill arenaria <i>Kn. pr.</i>	decumbens	Ljor	1	jn.au	P	C. G.H.	1818.	S s.1	Thun. pr. 1.1	
2549 adscindens A Br.	ascending	Ljor	2	jn.au	P	C.G.H.	1819.	S 81		
2550 rubricates A Br.	red-stemmed	Ljor	2	jn.au	P	C.G.H.	1818.	S 81		
2551 flagellaris A Br.	whip	Ljor	2	jn.au	P	C.G.H.	1816.	S s.1		
2552 glomerate A Br.	glomerated	Ljor	3	jn.au	P	C. G. H.	1789.	S p.1	Bur. af 99.2	
2553 decpiens A Br.	deceptive	Ljor	4	jn.au	P	C. G. H.	1806.	C Lp		
2554 Roxburghii R. Br.	Roxburgh's	Ljor	3	jn.au	W	C. G. H.	1806.	C l.p		
2a55 Burmanni tt. Br.	Burmann's	Ljor	2	jn.au	P	C.G.H.	1786.	C Lp	Bur. af. 99.1	
2a56 tritennata A Br.	triternate	Ljor	7	jn.au	P	C. G. H.	1802.	S p.1	Bot rep. 447	
2a57 elongata R Br.	elongated	Ljor	1	jn.au	P	C.G.H.	1800.	C l.p		
2558 crithmifolia A Br.	Samphire-lvd	Ljor	2	jn.au	R	C. G. H.	1818.	S 81		
n. 308. NIVE'NW R. Br.	NIVBNIA,	(J.Niven, a collector of Cape plants.)								
2559 sceptrica A Br.	sceptre/-	LJ or	2	jn.au	Pa.F	C G. H.	1790.	S p.1	Protedeea. 7.—12.	
2560 spathula & A Br.	spatulae	LJ or	2	jn.au	P	C. G. H.	1790.	C s.1	Spar.h. 1777.55.1	
2a61 parvifolia A Br.	small-leaved	u Jor	1*	jn.au	P	C.G.H.	1823.	S p.1	Thun.pr.41	
otto P'tea spatulata W.										
2562 spicata A Jr.	spiked	Ljor	2	jn.au	P	C.G.H.	1786.	S p.1		
2563 crenulifolia A Br.	Samphire-lvd	LJ or	2	jn.au	Pa.P	C G. H.	1797.	S p.1	Bot rep. 243	
& 64 media It. Br.	middle	Ljor	6	jn.au	P	C. G. H.	1803.	C s.p	Bot.rep.234	
*>6a Lagopus A Br.	Hare's-foot	LJ or	4	jn.au	P	C G. H.	1810.	S s.1		
•ws* 5 ? R E P H A L U S A Br.										
2567 setacea A Br.	SOROCKPHALIIS. {Soros, a heap, kenhale, a head.)	Proteace*. 7.—lft								
2567 imberbis A Br.	setaceous	Ljor	2	jn.au	P	C. G. H.	1823.	S Lp		
2ab8 spaUoides A Br.	beardless	Ljor	3	jn.au	P	C G. H.	1806.	C s.p		
£2*fenuiflora R.Br.	Spatalla-like	i Jor	3	jaau	P	C. G. H.	1803.	C s.p		
25701anata R.Br.	slender-leaved	Ljor	3	jn.au	P	C. G. H.	1802.	C l.p		
2571 l'mbricata A Br.	woolly	Ljor	2	jn.s	P	C. G. H.	1790.	C l.p	Thun. pr. SO. 3	
2572diversifolia R.Br.	imbricated	Hujor	3	ap.jl	P	C. G. H.	1794.	S l.p	Bot. rep. 517	
various-leaved	Ljor	4	ap.jl	P	C. G. H.	1803.	C l.p			
or?W- SPATA'LLA A Br.	SPATALLA.	(Spatala, wantonness.) Proteace. 2.—1&								
2573m61lis A2fr.	soft	Ljor	2	ap.au	P	C.G.H.	1826.	S p.1		
2574 nWca A Br.	white	Ljor	2	ap.au	P	C G. H.	1806.	S 81		
2g5 bracteata R Br.	bracteate	Ljor	3	ap.au	P	C. G. H.	1806.	S 81	Thun.pr. 1.2	
2576 ramlusa R Br.	ramulose	Ljor	3	au.s	P	C. G. H.	1787.	C Lp		
o2I proifera A Br.	proliferous	Ljor	21	myjn	P	C G. H.	1789.	S s.1	Bot rep. 429	
2578 Inctriva A Br.	Fncurved-leaf	Ljor	2	ap.au	P	C. G. H.	1812.	S p.1	Thun. pr. 2.3	
J579 caudata A Br.	caudate	si uJor	3	myjn	P	C. G. H.	1806.	C Lp		
2580 Thunbergii R. Br.	Thunberg's	Ljor	1	my.au	P	C. G. H.	1823.	S p.1		
2581 brevifolia A Br.	short-leaved	short-leaved								
9. SI DENA'NTHOS LoA.	ADENANTHOS.	(Aden, a gland, anthos, a flower.) Proteace. 3.-5.								
2582 apov/gadap.	obovate	LJ or	5	my.au	R	N. Holl.	1826.	S s.p	Lab.n.h. 1.37	
2583 cuneata R. Br.	vreAge-Jeaved	LJ or	—	R	N. Holl!	1824.	C 8!p.ILab.n.h.i!36			
2584 sericea R. Br.	silky	LJ or	—	R	N. Holl.	1824.	C s.p.1 Lab. n. h. 1.38			
*JJ2 CONOSPERMUM	A Br.	CONOSPERMUM.	(Konot,&cone,sperma,seed.) Proteace. 5.-9.							
»85 ellipticum A Br.	elliptic	LJ or	3	my.au	W	N. Holl.	1822.	C ap		
»86 cricifilium R. Br.	Heath-leaved	Ljor	3	jaau	W	N. Holl.	1820.	C s.p	Lin.tr. 10.17.1	
»87 taxifolium Sm.	Yew-leaved	Ljor	3	jaau	W	N. Holl.	1824.	C s.p	Bot.mag.2724	
»88 longitilium Sm.	long-leaved	Ljor	4	jaau	W.v	N. Holl.	1824.	C s.p	Sm.ex.bot.2.82	
»89 capitatum J2. Br.	capitate	LJ or	3	jaau	B	N. Holl.	1824.	C s.p		
		* raceme, keras, a hora) Proteacea. 1. NO								
*3J4. PERSOCN/^ Sm.	PERSOONIA.	(C. H. Persoon, of Dutch origin, a celeb, bot.) Proteacea. 15.—25.								
*JJ1 Pimf6lia A Br.	Pine-leaved	LJ or	4	jnjl	Y	N. S. W.	1822.	I, s.p		
*92 uniperina Lab.	Juniper-hke	LJ or	4	jnjl	Y	N. S. W.	1856L	C s.p	Lab.n.h. 1.45	
2593 linearis R Br.	Ihear-Zeaved	LJ or	4	jnjl	Y	N. S. W.	1794.	L s.p	Bot mag. 760	
2594 flexifolia A Br.	bent-leaved	Ljpr	2	jnjl	Y	N. Holl.	1824.	C p.1	Bot cab. 922	
2595 teretifolia R Br.	round-leaved	Ljor	4	jnjl	Y	N. Holl.	1822.	C s.p		
SS96 Chama'pitrys Cun.	Ground-pine	c Jor	4	—	Y	N. Holl.	1824.	C l.p		
2597 nutans R Br.	noddins	Ljor	—	—	Y	N. Holl.	1824.	C Lp		
2598 hirsuta R Br.	hairy	Ljor	4	myjl	Y	N. S.W.	1800.	C Lp	Bot cab. 327	
21 P'uinoM Cun.	frosty	Ljor	3	jnjl	Y	N. S.W.	1824.	C Lp		
2600 m6llis R Br.	soft	Ljor	3	jnjl	Y	N. S.W.	1826.	C l.p		
2601 scifra R Br.	scabrous	Ljor	4	jnjl	Y	N. S.W.	1826.	C l.p		
28W latifolia Andr.	broad-leaved	Ljor	4	jnjl	Y	N. S. W.	1795.	C p.1	Bot rep. 4289	
»>4 ferruginea AM.	rusty	Ljor	3	jnjl	Y.K	N. S. W.	1828.	C s.p	Sm.ex bot 283	
2605 lanceolata Andr.		Ljor	4	jnjl	Y	N. S. W.	1791.	III	Bot rep. li	
		A Jr. AGASTACHYS. {Agastos, admirable, staehys, a spike.) Proteacea. 1.								
J316. GREVIXLE^ H.	Br. GREMLLEA^ (C. F. GreviUe, a patron of natural science.) Proteacea. 31—4*.	sweet-scented								
<607 sericea A Br.	silky	uJor	6	aps	Pk	N. S. W.	1790.	S s.p	Motmaff. 86S	
2> £ punicea A Br.	scarlet	Ljor	—	ap.au	P	N. S. W.	1822.	C l.p	Sm.n.h.9. 5	
2609 dffbia A Br.	doubtful	Ljor	3	aps	V	N. S. W.	1K20.	C l.p		
2610 acanthifolia Cun.	Acanthus-lvd	Ljor	4	aps	P	N. S. W.	1824.	C l.p	Y. n. h. ic.	
2611 concinna R Br.	neat	IJor	4	ap.i	P	N. S. W.	1824.	C l.p		
2612 juniperina R Br.	Juniper-like	Ljor	4	aps	Pk	N. S. W.	1892.	C l.p	Bot cab. 1003	
2613 aspera A Br.	rough	Ljor	3	ap.«	Pk	N. S. W.	1824.	C l.p		
2614 linearis R Br.	linear-fcauof	Ljor	6	aps	W	N. S. W.	1790.	S s.p	Bot rep. 272	
2 incarnaata R M.	flesh-colored	Ljor	4	aps	F	N. Holl.	1790.	C p.1	Bot mag. 2661	
Sálbaorf.	strict	UJ or	4	aps	W	N. Holl.	1790.	C s.p	Bot. cab. 1003	
J615 strcta A ffr.	UJ or	4	ap.8	Pk	N. S. W.	1820.	C s.p			
616 rosmarinifolia dm.	Rosmary.lvd	uJor	4	aps	R	N. S. W.	1824.	C s.p		
617 ripana A Br.	river-side	LJ or	4	aps	Pk	N. & W.	17ij.J.	C l.p		
2618 Dlndersii Cun.	Flinders's	Ljor	3	aps	P	N. S. W.	1824.	C l.p		
W9 arenaria A 2?.	sand	Ljor	5	ap.8	Pk	N. S. W.	1803.	C In		
ruminata A Br.	acuminate	Ljor	4	aps	R	N. & W.	1805.	S l.p		
	long-styled	Ljor	n	aps	Pk	N. S. W.	1809.	C al		
		I)	4							

TETRANDRIA MONOGYNIA.

CLASS IV.

2822 mucronulata R. Br.	mucronulate	H J or	4 ap.8	Fk N. S. W.	1809. C l.p
£623 sulphurea Cun.	sulphur-c/d	■ J or	4 ap.s	Fa.Y N. S. W.	1884. C l.p
2624 Bauern' R. Br.	Bauer's	H \ J or	4 ap.au	R N. Holl.	1824. C s.p
2625 pubescens Hook.	pubescent	H J or	4 ap.s	R N. Holl.	1822. C s.p Hook. ex. fl. 216
8686 cinerea R. Br.	cinerous	H J or	4 ap.8	Pa.R K. S. W.	1822. C l.p
OC&I monfaria R. Br.	mountain	■ J or	4 ap.8	V N. S. W.	1842. C s.p
2628 Auxifolia R. Br.	Bo-leaved	■ J or	6 f.s.	Pk N. S. W.	1790. S i.p Bot. rep. 218
8629 mucromifolia Ctm.	mucronate-lvd	■ J or	3 ap.s	V N. S. W.	1824. C s.p
8630 phylloids R. Br.	Phylica-like	■ J or	3 my.au	R N. Holl.	1823. C s.p
8611 podalyria-tfa Swt	Podaluria-lvd	■ J or	5 my.au	R N. Holl.	1821. C s.p
8632 ferfer-folia Surf.	Berberis-leaved	H \ J or	4 my.au	R N. Holl.	1821. C s.p
2633 podocarpifolia Swt.	Podocarpus-lvd	H J or	4 my.au	R N. Holl.	1824. C e.p
2634 collina Kn.pr.	hill	■ J or	4 mrjl	Pk NSW.	1802. C l.p
2635 asplenifolia R. Br.	Asplenium-lvd	1 J or	5 mrjl	Pk N. S. W.	1806. C l.p
2636 heterophylla Host	variable-leaved	H J or	4 ap.au	R N. Holl.	1821. C s.p
8637 trifurcate Art.	■ J or	3 ap.au	R N. Holl.	1821. C s.p	
317. HAKEA. (Baron Hake, a German patron of botany.)	HAKEA.			Proteacea?	45.—58.
2638 pugioniformis R. Br.	dagger-formed	• \ J or	6 myjn	W V.D.LL.	1796. S s.p Bot cab. 353
8639 pinnis Kn. pr.	matched	H J or	6 myjn	W	1796. C l.p
8640 obliqua R. Br.	oblique-?UMi	H \ J or	6 ...	w	1803. C fl.p
2641 gibbosaa Cav.	gibbosus-rufa/	• J or	7 my.jn	W	1790. C Bp Cav. ic. 6.534
8642 aciculari* R. Br.	neet-e-leaved	H J or	3 myjn	W	N. SLW. 1790. C fl.p Ven.mal.111
2ri43 suavolens R. Br.	sweet-smelling	H L J or	4 ap.s	W	N. Holl. 1803. C s.p
8644 pectinata Coll.	pectinated	H J or	4 myjn	W X. Holl.	1810. C s.p Col. h. rip. 11
2f45 microcarpa R. Br.	small-fruited	H J or	4 myjn	W	V.DLL. 1819. C s.p Botreg/475
2645 propinquia Cun.	allied	* LJ or	4 myjn	W	N. S. W. 1824. C s.p
2647 sulcata R.Br.	mlcatc-leaved	H i J or	4 myjn	W	N. Holl. 1820. C fl.p
2648 subulate Cun.	mbulate-leaved	H J or	4 myjn	W	N. S. W. 1824. C fl.p
2649 varia R. Br.	variable	■ J or	3 jnau	W	N. Holl. 1825. C fl.p
2650 Klfcina Cun.	Furze-like	■ J or	4 myjn	W	■ ■ ■ 1824. C s.p
2651 lanigera Ten.	woolly	■ J or	3J myjn	W	N. Holl. 1880. C fl.p
2612 epiglottis Lab.	epiglottis	■ J or	4 my.jn	W	N. Holl. 1819. C s.p Lab. n.h. 1.40
2153 longifolia Dum.	long-leaved	■ J or	3 myjn	W	N. Holl. 1825. C s.p
8R54 trifurcata R. Br.	three-forked	■ J or	5 ■	W	N. Holl. 1824. C fl.p
2655 flexilis Bucr.	pliant	■ J or	4 ...	W	N. Holl. 1824. C s.p
2R56 florula R. Br.	flowery	■ J or	5 myjn	W	N. Holl. 1803. C s.p Bot mag. 2579
207 lineifirs R. Br.	linear	III-J or	■ ■ ■	W	N. Holl. 1884. C s.p
2658 ilicifolia R. Br.	Holly-leaved	• L J or	4 jl.8	W	N. Holl. 1803. C s.p
2659 nitida R. Br.	glossy	■ J or	5 njnl	W	1803. C s.p Bot. mag. 2246
8660 amplexicaulis R. Br.	stem-clasping	■ J or	2 ...	W	N. Holl. 1803. C 8.p
8661 prostra& R. Br.	prostrate	» J or	■ -	W	N. Holl. 1803. C B.p
2662 ceratonth Da R. Br.	horn-leaved	■ J or	■ ■ ■	W	N. Holl. 1803. C B.p
2663 Lambert* Swt	Lambert's	■ J or	4 jnjl	W	N. S. W. 1825. C s.p
SIG carduifolia Lo. C.	Thistle-leaved	■ J or	4 myjl	W	N. Holl. 1825. C 8.p
2665 cerasophylla Lo. C.	Buckshorn-lvd	■ J or	4	W	N. Holl. 1824. C s.p
2666 acanthophylla Lk.	prickly-leaved	H J or	S -	W	N. S. W. 1821. C 8.p
2667 undulata H. Br.	wave-leaved	■ J or	3 ■	W	N. Holl. 1803. L B.p
2668 oleifolia R. Br.	Olive-leaved	■ J or	5 njnl	W	N. Holl. 1794. C s.p
2669 saligna R. Br.	yyUm.leaved	■ J or	7 mr'jl	W	N. Holl. 1791. C s.p Bot rep. 215
2670 cincarea R. Br.	circneous-lmf	■ J or	5 njnl	W	N. Holl. 1803. S fl.p
2671 cane&cens Lk.	canescent	■ J or	3 jnau	W*	N. Holl. 18<9. C fl.p
2B72 dactyloides Cav.	Dactyhs-like	■ J or	7 jn.au	W	N. S. W. 1790. C s.p Cav. ic. 6.535
2673 elliptica Lab.	elliptic	■ J or	4 jn.au	W	N. Holl. 1794. C fl.p
8674 ruscifolia Lab.	Ruscus-leaved	■ J or	4 jn.au	W	N. Holl. 1824. C fl.p Lab.n.h. 1.39
2675 clavata R. Br.	clavate	H J or	5 jn.au	W	N. Holl. 1824. C <.p Lab. n. h. 1.41
2676 marginata A. Br.	marginate	H t or	4 jn.au	W	N. Holl. 1824. C s.p
2677 repanda Cun.	repand	H \ or	■ myjl	W	N. Holl. 1824. C fl.p
8678 cucullata Cun.	cucullate	■ J or	■ myjl	W	N. Holl. 1824. C fl.p
2679 ferruginea Cun.	rusty-leaved	■ J or	■ myjl	W	N. Holl. 1825. C fl.p
2680 latifolia Lo. C.	broad-leaved	■ J or	■ myjl	W	N. Holl. 1825. C fl.p
2681 petrophyllodea Ma.C.	Petrosylla-lk	H J or	■ ■	W	N. Holl. 1825. C s.p
2682 echinata Ma. C.	hedgehog	H J or	■ ■	W	N. Holl. 1824. C e.p
2G83 angustitoha Ma. C.	narrow-leaved	H J or	■ ■	W	1824. C s.p
318. STENOCA'R'PUS R. Br.	STENOCARPUS.	(Stcnos, narrow, karpos, fruit)		Proteacea?	1.—2.
2684 lafgnus R. Br.	Willow-teawrf	H i J or	5 njnl	G N. Holl.	1819. C fl.1 Bot.reg.441
• 319. LAMBERT/i* Sm.	LAMBERTIA. (A. B. Lambert, Esq., F. R. S., Pres. L. S., etc.)			Proteacea.	■
2685 uuiflora R. Br.	one-flowered	H i J or	3 jn.au	... N. Holl.	1884. C s.p
8686 echinata R. Br.	hedgehog	■ J or	3 jn.au	... N. Holl.	1824. C s.p
2687 longiflora Lod.	long-leaved	H J or	4 jn.au	R N. Holl.	1826. C s.p
8688 formosa \$m.	handsome	■ J or	4 jn.au	R N. S. W.	1788. C fl.p Bot rep. 69
2689 intermis R. Br.	unarmed	■ J or	3 . . .	N. Holl.	1824. C B.p
320. XYLOM'LUM Sm.	XYLOMELUM.	(Xyltm, wood, melon, an apple.)		Proteacea.	1.
250»yrif6rme Sm.	pyrifonn	i L j or	14 . . .	N. S. W.	1789. S Bp Cav. ic. 6.536
H Skea pyrifommis Cav.					
* 321. TELO*PEA R Br.	WARRATAR.	(Telopos, seen at a distance; its brilliant blossoms.)		Proteacea.	1.—2.
2C91 speciosissima R Br.	most splendid	f ull or	10 myjl	S N. & W.	1789. C sp "Bot. mag. 1128
322. LOMAT'TIA R. Br.	LOMATIA.	(Loma, a border; winged edge of the seeds.)		Proteacea.	5.—8.
2692 silafloria R. Br.	Silaflm-leaved	■ J or	8 jn.au	O N. S. W.	1792. C s.p Bot. mag. 1272
8693 tinrtfria R. Br.	dyer's	■ J or	2 . . .	N. Holl.	1822. C s.p Lab. n. h. 1.42,43
2694 longifolia R. Br.	long-leaved	H J or	8 jn.	G N. S. W.	1816. C l.p Botreg.442
2695 flicifolia R. Br.	Holly.leaved	■ J or	3 jn.au	... N. Holl.	1824. C e.p
2696 denULta R. Br.	toothed	■ J or	3 . . .	Chil.	1824. C l.p Fl. per. 1.94. a.
* 323. RHOPAL'A R. Br.	RHOPALA.	(Roupala, its aboriginal name.)		Proteacea.	3.—16.
2697 mfeia R. Br.	mediate	J L or	10 myjn	... Guiana	1823. C l.p Aub.gui. 1.32
Roupala montana Aubl.					
\$698 dentata R. Br.	tooth.Jeaved	ti J or	10 myau	G S...Amer.	1802. C l.p
2G99 sessilifolia Rich.	sessile-leaved	\$ L or	10 . . .	G Guiana	1803. C l.p Rud. gui. 1.31
* 34. KNIGHT'A R. Br.	KNICHTIA.	(T. A. Knight, Esq. F.R.S., Pres. H. S., etc.)		Proteacea.	1.
2700 exc'lsa R. Br.	lofty	f L j or	10 . . .	F A. Zeal.	1824. C s.p Lin. tr. 10.2

* 325. BA'NKS/jf R. Br.	BANKSIA. neaUflowered	32.-36.
2702 sphaerocarpa R. Br.	round-fruited	
2703 nifitans R. Br.	nodding-JftM*	
2704 eridifolia/.	Heath-leaved	
2705 spinulosa R. Br.	spinulose	
2706 collina A. i?r.	hill	
2707 clatior R. Br.	taller	
2708 occidentalis R. Br.	western	
2709 littoralis R. Br.	sea-side	
2710 marginata Cav.	marginate	
2711 depresso R. Br.	depressed	
2712 australis R. Br.	southern	
2713 insularis R. Br.	island	
2714 intcgriifolia R. Br.	entire-leaved	
2715 c6mpar R. Br.	well-matched	
2716 verticillata R. Br.	verticillate	
2717 coccinea R. Br.	sc-drct-Jflowercd	
2718 paludosa R. Br.	marshy	
2719 media R. Br.	mediate	
2720 oblongifolia Cav.	oblong-leaved	
2721 latifolia R. Br.	broad-leaved	
2722 marcescens R. Br.	marcescent	
2723 attenuata R. Br.	attenuated	
2724 serrata R. Br.	s&vf-leaved	
2725 82'mula R. Br.	rival	
2726 dryandro0UfttR.Br.	Dryandra-like	
2727 queratifolia if. Br.	Oak-leaved	
2728 dentiflta/..	toothed	
2729 speciosa R. Br.	showy	
2730 grandis W.	great-flowered	
2731 prostrata R. Br.	prostrate	
2732 repens Lab.	creeping	
• 326. DRY ANDRI< R Br.	DRVANDRA. (<i>Jonas Dryander</i> , a distinguished botanist.)	Lsp. Lab. ya. 15.23
2733 florib&ndia R. Br.	many-flowered	
2734 cuneata R. Br.	wedge-Aviwd	
1 brevifolia	short-leaved	
2 longifolia	long-leaved	
2735 armata R. Br.	armed	
2736 falcata R. Br.	falcate	
2737 formosa R. Br.	handsome	
2738 mucronulata R. Br.	mucronulated	
2739 plumosa R. Br.	feathered	
2740 obtusifl R. Br.	obtuse-leav?d	
2741 nivea R. Br.	white-leaved	
2742 longifolia R. Br.	long-leaved	
2743 Baxteri R. Br.	Baxter's	
2744 tenuifolia R. Br.	fine-leaved	
2745 teridif6lia R. Br.	Pteris-leaved	
2746 olechnifolia R. Br.	Blechnum-lvd	
2747 nervosa R. Br.	nervose	
327. STRUTHI'OLA	L. STRUTMOLA. (<i>Struthion</i> , a little sparrow; seeds beak-like.)	np*gf" 15/-18-
■'48 virgata L.	twiggy	
2749 ciliata Andr.	ciliated	
2750 lateri fibra Horn.	side-flowered	
2751 longiflora Lam.	long-flowered	
2752 pubescens H. K.	pubescent	
2753 juniperina W.	Juniper-leaved	
2754 likens Pair.	shining	
2755 angustifolia Lam.	parrot-leaved	
2756 erecta W. en.	upright	
stricta Donn		
2757 glabra L.	smooth	
2758 incana Lod.	hoary	
2759 imbricata H. K.	flve&-leaved	
2760 striata Lam.	striated	
2761 tomentosa H. K.	woolly-leaved	
2762 ovata Thun.	ovate-leaved	
328. OPERCULA'RIA	L. STRUTMOLA. (<i>Struthion</i> , a little sparrow; seeds beak-like.)	np*gf" 15/-18-
2763 umbellata Gae.		
2764 aspera Gae.		
2765 hfsida Spr.		
aspera J. diph^lla Gae.		
2766 sessiliflra J.	sessile-flowered	
8767 ocyinif6lia J.	Basil-leaved	
	vAl un	
	Ed cu	
	W	C. G. H. 1824. S. sp. An. mu. 4.70.2
	W	E. Indies 1824. S. sp.
^329. CRYPTOSPE/RMUM Pers.	CRYPTOSPERMUM. (<i>Kryptos</i> . hidden, <i>sperma</i> , seed.)	OpereularU*. 1.
2768 Yofingif Pers.	Young's	a j w 4 jl.au
Opercularia palcat'a Y.		N.S.W. 1793. C co Lin.tr. 3.5
330. PATRIN'N/i J.	PATRINIA. (Af. <i>Patrin</i> , a Siberian traveller.)	Valerihntai. 7. —
2769 sibirica J.	Siberian	17SL B co Bot.mag.2325
Valeriu'nithenica W.	sibirica L.	
2770 rupistris J.	rork	
Valeridna sibirica W.	Fidia rupestris'	
2771 scabiosae6lia Fis.	Scabi6sa-leaved	
2772 serratulissfolia Fis.	Serratula-lvd	
2773 intermedia Fis.	intermediate	
2774 coronata Fis.	crowned	
2775 nudiflscula Fii.	nakedish	
* 331. PO'THOS L.	Pomos.	(<i>Potha</i> , its name in Ceylon.)
8776 scandens L.	climbing	Aroidets. 27. — 39.
8777 officinalifl Rox.	officinal	% ES cu 4 myjn Ap E. Indies 1821. Sk s.p. Uu. am. 5.1.3
	t E tu 20 apjl Ap E. Indies 1820. Sk p. 1	

TETRANDRIA MONOGYNIA.

CLASS IV.

2778 <i>Pipila</i> Rox.	Peeplee	■ ap.jl	Ap	E. Indies	1820.	Skp.l	
2779 <i>gigantea</i> Box.	giant	■ ap.jl	Ap	E. Indies	1824.	Skp.1	
2780 <i>pertusa</i> Box.	perforated	■ ap.jl	Ap	E. Indies	1824.	Skp.l	
<281 <i>aca&lis</i> L.	stemless	1 ap.jl	Ap	W. Indies	1790.	Sks.p	
2782 <i>lancolata</i> L.	<i>lanceolated</i>	3* ap.jl	Ap	Barbadoes	1790.	Sks.p	
2783 <i>crenata</i> L.	crenate	J my.jn	Ap	W. Indies	1823.	Sks.p	
2784 <i>gr&cihs</i> Bud.	slender	1 my.jn	Ap	Trinidad	WX.	Sk s.p	
2785 <i>violacea</i> Sum.	<i>violet-fruited</i>	2 ap.jn	Ap	Jamaica	1793.	Sks.l	
2786 <i>cannifolia</i> H. K.	Canna-leaved	3 ap.my	Ap	W. Indies	1789.	Sks.p	
2787 <i>crassinervis</i> Jac.	thick-nerved	2 ...	Ap	S. Amer.	1796.	Sks.p	
2788 <i>cordata</i> L.	<i>heart-leaved</i>	3 ap	Ap	America	1770.	Sks.p	
2789 <i>sagittata</i> B. M.	<i>arrow-leaved</i>	3 au	Ap	W. Indies	1800.	Skp.l	
2790 <i>angustata</i> H. k A	narrow	my.jn	Ap	Trinidad	1823.	Sk s.p	
2791 <i>macrophylla</i> Swz.	large-leaved	3 my.jn	Ap	W. Indies	1794.	Ska.p	
grandifolia Jar.						Jac. ic. 3.610	
2792 <i>obtusifolia</i> H. K.	blunt-leaved	■ my.jn	Ap	Barbadoes	1790.	Skp.l	
2793 <i>digitata</i> Jfc.	digitate	■ my.jn	Ap	W. Indies	1820.	Sks.p	
2794 <i>conacca</i> tif-a.a.	coriaceous	■ jn	Ap	Brazil	1824.	Sk Bp	
2795 <i>Harr.Wi</i> Grah.	Harris's	■	Ap	Brazil	1814.	Ske.p	
2796 <i>rubrine'ria</i> LA.	red-nerved	■	Ap"	S. Amer.	1810.	Skp.l	
2797 <i>P'matd</i> L.	natinated	■	Ap"	S. Amer.	1803.	Skp.l	
2798 <i>Laaia</i> /fax	Lasia	■	Ap	E. Indies	1819.	Skp.l	
2799 <i>pentaphl</i> Ila FT.	five-leaved	■	Ap	Cayenne	1803.	Skp.l	
2800 <i>pinnata</i> L.	pinnated	■	Ap	E. Indies	18%.	Sk p.l	
2801 <i>decunfa</i> Box.	running	■	Ap	E. Indies	1822.	Skp.l	
2802 <i>heterophylla</i> Box.	variable-leaved	■ TELcu	20	■ Ap	E. Indies	1824.	Skp.l
332 SYMPLOCA'RPUIS	Nut.	SYMPLOCARPUIS	(<i>Symploke</i> , connection, larpes, fruit.)			Aroidee. 1.	
2803 fœ'tidus Nut.	fœtid	* AJ cu	1 mr.jn	Ap	N. Amer.	1735.	Sk aq
Pothos fœ'tidus B. M.	Jrac6ntium fte'tidum L.					Bot mag.a36	
333. GYMINOSTACHYS R Br.	GYMINOSTACHYS	R Br.	(<i>Gymnos</i> , naked, <i>stachys</i> , a spike.)			Aroidete. 1.	
2804 <i>anceps</i> B. Br.	two-edged	■ iAI ° r	1 Jn.jl	... N. Holl.	1820.	Sk p.l	
• 334. RIVI'NI* L.	RIYIXA.	(A. Q. <i>Rivinus</i> , a botanist of Saxony, d. 1722.)				Chenopdeet. 6.-8.	
2805 <i>hdmlis</i> L.	humble	■ ID or	2 Ja o	W W. Indies	1699.	S r m Bot mag. 1781	
2 cantees W.	canescent	n. CD or	2 my.au	W W. Indies	1804.	C Ip	
2806 <i>purpurascens</i> Schr.	purpurascens	ft. CD or	■ my.au	Pk W. Indies	1815.	c t.p	
2807 <i>lae</i> VM L	smooth	n. CJ or	■ f.s	Pk IV. Indies	1733.	S r.m Bot. mag. 2J33	
2808 <i>lancolata</i> W.	lanceolate	■	■ my.jl	... Brazil	1815.	C Lp	
2809 <i>brasiliensis</i> L.	Brazilian	■	■ jn.jl	G Brazil	1790.	C b	
2810 <i>octandra</i> L.	octandrous	H.Qor	■ my.jn	W W. Indies	1752.	C p.l Br. jam. 149.23.2	
335. CAMPHOROSMA L.	CAMPHOROSVA.	(<i>Camphora</i> , % I'at, camphor, <i>osmc</i> , Gr. a smell.)				Chenopdete. 3.-5.	
2811 <i>mqnspellaca</i> L.	Montpeher	a. L cu	14 aug	Ap S. Europe	KSIQ.	C p.l Suhk. hsn. 1.26	
2812 <i>acuta</i> L.	a.acute-leaved	O un	■ au	Italy	1820.	S s.p I' bic.404	
2813 <i>ovata</i> W.tft.	ovate-leaved	O un	■ au	Ap Hungary	1820.	B.p W.&K.63	
336. ANISACA'NTHA B. Br.	B. Br.	AM SACANTHA.	(<i>Anisos</i> , unequal, <i>akantha</i> , a.ipine.)			Chenopdlear. 1.	
2814 <i>divaricata</i> B. Br.	divaricate	M il cu	2 .	N. Holl.	18^4.	C p.l	
337. ALCHEMILLA L.	LADIES' MANTLE.	(<i>Alkemelyeh</i> , its Arabic name.)				Sanguitdrbea. 11.	
2815 <i>vulgäris</i> L.	common			Britain	me. pa.	D co Eng.bot597	
2 varivgata	variegated			Britain	gard.	D co	
3 glabra Dec.	smooth			Pyrenees	1817.	I co	
2816 <i>hybrida</i> Dec.	hybrid			Britain	moun.	D co MiLic. 18	
montana W.en.							
2817 <i>pubescens</i> L.	pubescent			Caucasus	1813.	D co H.ber.2.79	
2818 <i>scrifica</i> W.en.	silky			Caucasus	1813.	D co	
2819 <i>alpina</i> L.	alpine			Britain	rocks.	D co Eng.bot.244	
2820 <i>pcntaphilla</i> L.	five-leaved			Switzerl.	1784.	D co Buc. mu. 1.1	
2821 <i>fissa</i> Sent.	cleft-leaved			Switzerl.	18.6.	I co	
2822 <i>SvhbaMia>felia</i> Dec.	Sibbaldia-lvd			Mexico	1823.	D p.l.sH.&B. n.6.561	
2823 <i>capensis</i> Thun.	Cape			C. G. H.	1818.	I) pl I-am. ll. Sri. 2	
2824 <i>A'phanes</i> Leers	Parsley-picrt			Britain	-	S co Eng. bot 1011	
arvénensis Sco.	A'phanes arvénensis L.						
2825 <i>corliocopiooides</i> Lag.	Cornucopia-like			Spain	1826.	S co	
• 338. SANGUISORBA L.	GREAT BLRNET.	(<i>Sanguis</i> , blood, <i>sorbeo</i> to absorb.)				Sanguisdrbete. 7.	
2826 <i>oficinalis</i> L.	officinal			Britain	me. pa.	S co Eng. Iwt. 1312	
2 auricul&t;a	eared			Italy	-	Boc. nu. 19. 9	
2827 <i>cárnea</i> Fis.	flesh-colored			-	1823.	D co Sch. mo. 69	
2828 <i>mauntanica</i> Desf.	Mauritanian			Mauritan.	1810.	D co M. h. 8.18.4	
2829 <i>tenuifolia</i> Fis.	fine-leaved				1820.	D co	
2830 <i>média</i> L.	middle			Canada	1785.	I) Za.h. 1811SB	
8831 <i>caiatlcnais</i> L.	Canadian			Canada	li&J.	D co Cor. ca. 174	
2832 <i>ncgl^ntu</i> G. Don	neglected			Eurofie?	1800.	O co	
* 339. DORSTEN'I'il L.	DORSTENIA.	{T. Dorsten, a German imtanirI)		Vrticcw. ft	-16.		
2833 <i>cordifolia</i> Lam.	heart-leaved	EJ cu	my.au	G W. Indie*	1822.	D co Swz. ob. 7.2	
2834 <i>brasilensis</i> Lam.	Brazilian	EJcu	ap.au	G S. Amer.	1792.	R s.l	
2835 <i>Houston</i> L.	Houston's	EJ cu	jn.jl	G S. Amer.	1747.	R s.l Bot mag. 2017	
2836 <i>An(dia</i> Lam.	Arum-leaved	EJCU	my.jl	O Brazil	IM&	R s.l Bot mag. 2476	
2837 <i>Cot/rajrva</i> L.	Contrajerva root	12S TM	mr.jn	G S. Amer.	1748.	Skp.l Jac. ic. a 614	
2&S Draku'na L.	Draksmu contra	jeu	py.au	Vera Cruz	1818.	U pi Bot cab. 677	
2839 <i>ccratosanthes</i> Lod.	Buckshorn-flwdjg	El cu	\$ mr.jn	G S. Amer.	1826.	D 6.p Bot cab. 1215	
340. ISNA'R'Dii L.	ISNARDA.	{A. T. D. d*Isnard, a French botanist.)		Onagrarue. 1. — 6.			
2840 <i>palustris</i> L.	marsh	j , ^ * i j		Britain	pools s bog		
341. ELJEAGNU'S L.	OLEASTER.	{Elaia, an olive, agnos, the chaste tree.)		ELtuignear. 7.—12.			
2841 <i>angustifolia</i> L.	narrow-leaved	or 15 jl	Y	S. Europe	1633.	C co Pal.ros.1.4	
2842 <i>argentea</i> Ph.	silvery	or 10 jl.au	Ap	N. Amer.	1813.	C co	
2843 <i>orientalis</i> L.	Oriental	1 J or 10 jl.au	Ap	Levant	1748.	L pi Pal. ros. 1.5	
2844 <i>arborea</i> Box.	tree	1 J or 30 ...	Ap	Nepal	1819.	C co	
2846 <i>latifolia</i> L.	broad-leaved	CD or 4 Jl.au	Ap	£. Indies	1712.	L p.l Bur. icy. 59.2	
2846 <i>acuminata</i> /Jr.	acuminated	I or 4 ...	Ap	-	... C co		
2847 <i>conferta</i> Rox.	crowded	or 10 ...	Ap	Nepal	1825.	C s.p	

342. GLOBULA'RIA L.	GLOBULARIA.	(The flowers in globose heads.) <i>Globuldrhue.</i> 9.—13.
2848 longifolia L.	long-leaved	ft. pr. 3 Jlau W Madeira 1775. L pl Botreg. 685
ralicina Lam.		
2849 Alypum L.	Alypo	<-i_Jpr 2 a.u.s Pa S. Europe 1640. L p.l Gar. aix42
2 integrifolium	entire-leaved	<, »J or 2 a.u.s Pa S. Europe ... L p.l
2850 vulgáris L.	common	£ A P myjn B Europe 1640. E p.l Bot mag. 2256
2851 /influa Lam.	Flax-leaved	£ A or f myjn B Spain 1818. D p.l
vulgaris Brot.		
2852 spinosa MU.	prickly-leaved	£ i_J pr I myjn B Spain 164a C Lp
2853 cordifolia L.	heart-leaved	£ A pr *jnjl B Germany 163a C Lp Jac. au. 3.245
2854 nana Lam.	dwarf	£ A pr i B France 1824. C Lp Dec. ic. 1. 3
2855 nudicaulis L.	naked-stalked	£ A pr i B Germany 1629. C p.l Jac au. 3. 230
2856 Acellidifolia Ten.	Daisy-leaved	£ A pr R Italy 1825. C sp
•343. HOUSTONI <i>n</i> J	L. HOUSTONIA. (W. Houston, M. D. a British botanist, d 1730)	
2857 cerflea £..	blae-floivered	£ A pr * my.au L.B N. Amer. 1785. D sp
2858 purpurea L.	^purple-flowered	£ A pr * my.au P N. Amer. 1800. D co
344. DI'PSACUS L.	TEASEL. (<i>Dipsao</i> , to thirst j leaves hold water.) <i>Dipsacea.</i> 7.—10.	
2859 fulignum All.	fullers'	■ P Britain hedg. S 1 Eng. bot 2080
2860 sylvestris Mil.	wood	■ P Britain m.hed S m.s Eng. bot 1032
2861 laciniatus L.	cut-leaved	■ P Germany 1683. S m.s Jac. au. 5. 403
2862 ferox Lot.	fierce	■ P S. Europe 1818. S co
2863 GmeUni Bieb.	GmeUn's	■ B Caucasus 1820. S m.s
2864 inermis Wai.	unarmed	■ W Nepal 1823. S m.*
2865 pilosus L.	pilose	■ W Britain moi.pl. S m.s Eng. bot 877
345. CEPHALA*RIA Schr.	CEPHALARIA. (<i>Kephale</i> , a head; the flowers in round heads.) <i>Dipsaceaef.</i> 20.	
2866 alpina Schr.	Alpine	■ j or 3 jn.jl LY Switzerl. 1570. D co Bes. eys. 8.1
Succisa alpina Spr.		
2867 albescens R. & S.	albescent	■ Siberia 1804. D co
2868 grisea R. & S.	Greek	■ S. Europe 1819. D co
Scabi6sadecurrrensL.		
2869 rigida R. & S.	stiff-leaved	■ C.G.H. 1731. S p.l Com. h. 2.93
2870 attenuata R. & S.	alender-leaved	■ C.G.H. 1774. S Lp
2871 scabra R.S/i	scabrous	CO. H. 1825. C co
2872 transylvanica R. % S.	Transylvanian	Transylv. 1699. S co Jac. vin. 2.111
2873 syriaca R.Ses.	Syrian	Syria 1633. S co M. h. 3.14.14
2874 leucantha R. & S.	white-flowered	France 1739. D co Ger. cm. 721.8
2875 tatarica A.* &	Tartarian	Russia 1759. S co Ac. up. 1744.1
2876 uralensis Mur.	Uralian	Siberia 178a S co C. g. 1782.4
2877 Uevigita W.&K.	smooth	Hungary 1805. D co Vf.&K. 3.230
Succisa centaurioides Spr.		
2878 corniculata R. & S.	small-horned	Hungary 1801. D co W. & K. 13
2879 leucanithema R. & S.	white-flowered	France 182a S co
Lepicephalus leucantha Lag.		
2880 coriacea W. en.	coriaceous	Tauria 1813. D co
Lepicphalus coriacea Lag.		
2881 cretacea R. & S.	chalky	Caucasus 1818. D co
S 2882 Vaill&pntf u. & S.	Vaillant's	W Aleppo 1822. S co
Pterocephalus VaiUfntV Lag.		S. Europe 1597. S co
5 2883 papposa R. & S.	downy-headed	P Spain 1823. S co
ft 2884 diffracta R.S/S.	diandrous	B Spain 1819. S co Gae.fr. 86.4
ft 2885 plumosa R. A. S.	plumose	Scabi6sa WdUchi' Lk.
* 346. SC ABICSA Vail SCABIOUS.	leprosy; medicinal qualities.) <i>Dipsacaz.</i> 68.—103.	
2886 dichotoma CTc.	forked	■ 1 jn.au Pk Sicily 1804. S co Boc. mu. 120
h 2887 Succ sa L.	Devil's Bit	■ 1 au.o V Britain pas. D co Eng. bot 878
Succ sa pratlnsis Moen.		
2888 glabrata Schott	smooth	■ A or 1 jn.jl P Austria 1819. D co
Succisa glabrata Spr.		
2889 australis Wul.	southern	■ k A pr 2 5n.. P Styria 1820. D co
2890 p>eud.austraUsA. % S.	bastard australis	■ A or 2 jn.jl P Russia 1825. D co
2891 stricta W. K.	strict	■ A or 2 jn.jl R Hungary 1820. S co W.&K.2.138
ndrica W. en.		
2892 integrifolia L.	entire-leaved	■ R France 1748. S co
2893 hybrida All.	hybrid	■ R S. France 1819. S co
2894 Saicéd Lag.	Salceu's	■ W Spain 1823. D co
2895 arvensis L.	corn field	■ P Britain cor. fl. S co Eng. bot. 659
2896 collina Req.	hill	■ P Switzerl. 1819. S co
hirsuta Lap.		
2897 montana Bieb.	mountain	■ W Caucasus 1820. D co
2898 sylvatica L.	wood	■ P Austria 1633. D co Jac au. 4.362
HS Pubiscns Kit.	pubescent	■ W Hungary 182a D co
2901 carpatica Fis.	Teasel-leaved	■ W Germany 1818. D co
SKW hirsuta J.Ait.	Carpathan	■ W Prussia 1819. D co
SUM diversifolia Bourn,	long-leaved	■ Li Hungary 1802. D co W. & K. 5
i ££* JIII&ta SP-	various-lea	■ R TranBylv. 1826. K co
\ VMS dumetorum Brou.	ciliated	■ R Germany 1802. D co
Ftrocophalus dumetorum Coul.	doubtful	■ W a Europe 1820. C co
I ?v c^nescens Kit.	canescent	■ 1 jl.au Li Hungary 1802. D co W.&K. 53
7 ^ gram6ntia L.	Gramont	■ 1 jlau LB & Europe 1597. D p.l Gcr. h.582. 2
2908 setiforma L.	bristle-bearing	■ 2 jn.jl W 8. France 1826. R co
mdlhs W. en.	long-haired	■ « jn.jl 1820. S co
2910 rupestris Bieb.	rock	■ P Caucasus 1824. D co
2911 commutata R. Sc S.	changed	■ B Siberia 1826. S co
field		
2913 amoe'na Jac.	pleasant	■ P Hungary 1818. S co W. & K. 3.204
	 1820. S co
g j * Pyrenica AIL	Pyr>nean	■ P S. France 1819. S ALped.1.25.2
g } columbaria L.	pigeon-house	■ P Britain dr. pa. S Eng. bot 1311
2916 micrinth	small-flowered	rk Armenia 1825. S An. mu. 11.25

TETRANDR1A MONOGYNIA.

k 2917 grandiflora Sco.	great-flowered	9	+	+	+	Barbary	1804.	S	co	Scop. d. 3.14		
I 2918 lucida VU.	shining	+	-	-	+	Dauphinyl	1800.	D	co			
I 2919 5ilcnif61ia W.&K.	Silene-leaved	+	-	-	+	Hungary	1826.	S	co	W.&K.2,157		
& 2920 lutea Fis.	yellow	+	-	-	+	Russia	1820.	S	co			
(2921 sicula L.	Sicilian	+	-	-	+	Sicily	1783.	S	co	Jac. vin. 1.15.		
i 2922 saxatilis Cav.	rock	+	-	-	+	Spain	1827.	S	co	Cav. ic. 2.184		
I 2923 brachiatia Sm.	branched	+	-	-	+	Greece	1823.	S	co	Fl.gr. 109		
Pterocephalus brachiatia Spr.												
I 2924 legionensis Lag.	Leon	+	-	-	+	Spain	1820.	S	co			
I 2925 rotata Bieb.	rotate	+	-	-	+	Iberia	1823.	S	co			
I 2926 Biebersteinii R. & S.	Bieberstein's	+	-	-	+	Iberia	1820.	S	co			
I 2927 intermedia R. 4 S.	intermediate	+	-	-	+	S. Europe	1824.	S	co			
I 2928 rutajolia Vahl	Rue-leaved	+	-	-	+	Sicily	1804.	D	co	Boc. sic. 52		
I 2929 maritima L.	K?..	+	-	-	+	Italy	168a	S	co	M. h. 6.15.29		
I 2930 Webbiana D. Don	Webb's	+	-	-	+	Mt.Ida.	1818.	t>	co	Bot.reg.717		
I 2931 crenata Cyr.	crenate	+	-	-	+	Italy	1825.	S	co	Cyr. n.é. 1.3		
coronopifolia Sm.												
4 2932 holoscrcea Bert.	all-silky	+	-	-	+	Pyrenees	1818.	D	co			
4 2933 tomentosa Cav.	woolly	+	-	-	+	Spain	1827.	S	co	Cav. ic. 2. 183		
I 2934 mollissima Dec.	softest	+	-	-	+	Italy	1820.	S	co			
\ 2935 stellata L.	starry	+	-	-	+	Spain	1596.	S	co	Cl. h. 2.1. ic.		
laxiflora LA:												
4 2936 simplex D?<.	simple	+	-	-	2	jl.au	W	S. Europe	1820.	S	co	Desf. at 1.39.1
\ 2937 pectinata Z.fl^.	pectinate	+	-	-	1	jjl.au	V	Arabia	1824.	S	co	
ty 2938 prolifera /.	proliferous	+	-	-	1	jl.au	Y	Egypt	1683.	S	co	Her. par. 125
\ 2939 atropurpurea L.	dark purple	+	-	-	+	jl.s	Br	1629.	S	co	Botmag.247
2 carnea	flesh-colored	+	-	-	+	jl.au	F	E. Indies	1629.	S	co	
3 rdsea	rose-colored	+	-	-	+	jl.au	R	E. Indies	1629.	S	co	
4 variegata	variegated	+	-	-	+	jl.au	Va "	E. Indies	1629.	S	co	
5 alba	white	+	-	-	3	jl.au	W	E. Indies	1629.	S	co	
6 prolifeia	proliferous	+	-	-	3	jl.au	P	E. Indies	1629.	S	co	
§ 2040 argentea L.	silvery	+	-	-	2	jn.o	W	Levant	1713.	D	co	An. mu. 11.24
§ 2941 urceofata Jfrtf.	jagged	+	-	-	3	jl.au	Y	Barbary	1804.	S	co	M. h. 6.13.24
I 2942 africana L.	African	+	-	-	6	jl.o	W	Africa	1690.	S	p.l	Her. par. 219
\ 2943 altissima Jrtc.	tallest	+	-	-	5	my.jn	L.B	Africa	1819.	S	co	Jac. vin. 2.185
africana Hort. indurata L.												
\$ 2944 nitens R. & S.	glittering	+	-	-	-	-	Azores*	1779.	D	co		
Scabiosa lycida H.K.												
ft 2945 cretica L.	Cretan	+	-	-	P	Crete	1596.	S	p.l	Mor.h.3.15.31		
{ 2946 graminititia L.	Grass-leaved	+	-	-	B	SwitzerL	1683.	D	p.l	Bot. reg 835		
i 2947 caucaea Bieb.	Caucasian	+	-	-	B	Caucasus	1803.	D	p.l	Bot mag. 886		
§ 2948 elegans Spr.	elegant	+	-	-	L.B	S. Europe	1813.	D	co			
ft 2949 lyrata Forsk.	lyre-faifov	+	-	-	P	Turkey	1799.	S	81			
\$ 2950 palacisfina L.	Palestine	+	-	-	Ci	Palestine	1771.	S	s.l	Jac. vin. 196		
Astrocephalus palestina Spr.												
I 2951 isetensis L.	Isetsk	+	-	-	W	Siberia	1801.	S	s.l	Gm. si. 2.88		
I 2952 ucrainica L.	Ukraine	+	-	-	L.Y	Ukraine	1795.	C	s.l	Gm. si. 2.87		
I 2953 ochroclaca L.	yellowish white	+	-	-	Y	Germany	1597.	D	s.l	Jac. au. 5.439		
I 2954 bann&tica Kd.	Bannatic	+	-	-	Pk	Hungary	1800.	D	co	W.&K. 10.12		
Collfim^ Ten.												
2955 Scopoli* Lk.	Scopoli's	+	-	-	Str	S. Europe	1818.	D	co			
2956 negl^cta Horn.	neglected	+	-	-	R	Germany	1825.	S	co			
2957 incana Schult.	hoary	+	-	-	R	Europe	182a	D	co			
2958 ceratophylla Ten.	Buckshorn-lvd	+	-	-	R	Italy	1826.	S	co			
pilosa Lag. ?												

347. KNAUTIA/T/ 4 L.
9959 orientalis L.
2960 propontica IV.

KNAUTIA.	(C. Knaut, physician at Halle, in Saxony, died 1694)	Dipsu <u>cea</u> .	2.
Oriental	O or 1 jn.s	R	Levant
Propontic	^ Q or 2 jn.au	P	Levant
			1713. S co Sc. han. 1.22
			Til. pis. 15a 48
• 348. GA'LIUM L.	BED-STRAW.	(Gala, milk; used for curdling milk.)	Rubid <u>cea</u> *. 75.—16a
2961 rubrifoldes L.	Madder-like	1 jl	W
2962 diffVisum Sehr.	diffuse	1 jjl	S. Europe 1775. D co Bux. c.2.29
2963 valmtioides Bieb.	Valantia-like	1 jjnl	Germany 1820. D co
2964 palfistre L.	marsh	jjnl	Caucasus 1819. D co
2965 trifidum L.	trifid	W	S. Europe m me. D m8 Eng. bot 1B57
2966 montanthum L.	mountain	W	Denmark 1826. D co
2967 Villarsii Gau.	Villars's	W	France 1819. D co Fl. dan. 48
2968 catureif6liura Trev.	Savory-leaved	W	France 1823. D co Vil. del. 2. 7
9969 pubescens Sehr.	pubescent	W	Carpathia 1817. D co
2970 lithospermif6liumFis.	Lithosper-.lvd	W	Germany 1819. D co
2971 frutic6sum W.	shubby	W	Russia 1826. D co
2972 Hallerii R. & &	Haller's	W	Crete 1819. C co
vernum Sco.		W	SwitzerL 1819. D co
2973 Withering'E. B.	Withering'g	W	England hea. D <p Eng. bot 2206
2974 tfeble Hofm.	weak	W	Portugal 1824. D co
2975 campestre Schous.	field	W	Barbary 1820. S co
2976 divaricatum Lam	divaricate	W	France 1818. S co Dec.ic.24
2977 megalosperrum All.	large-seeded	W	S. France 1826. D co Alped.79. 4
2978 glabrum Thun.	smooth	W	C G. H. 1825. D co
2979 uligin6sum L.	marsh	W	Britain mar. D m.% Eng. bot 1972
S(H)I aufltriacum Jac	Austrian	W	Europe 1804. D co Jac. au. 90
C98J Jlorri mi All.	Buccome'a	Pk	Eurolic 1801. D co Hoc. mu. 1*5.101
9SM2 obliquum VU.	oblique	W	& France 1826. D co Vil. del. 8. 8
S9H3 ht6de8 Lk.	marsh	W	Spain 1825. S co
2P84 rdbrum L.	red	P	Italy 1597. D co Ger. h. 967.3
2985 v6rum L.	true cheese rennet	Y	Britain bu. pL D m.s Eng. bot 660
2986 tyrolpse IV. en.	Tyrolean	3r	Tyrol 1801. D co
2987 iplKigo L.	great hedge	Mollugo 3	Britain hedg. D co Eng. bot 1673
2988 erectum E. B.	upright	W	Britain m. pass. D m.s Eng. bot. 2sJ.7
2P89 lucidum All.	shining	W	Iberia 1819. D co Al пед.1.77.2
2990 pusillum E. R	little	i lau	England moun. D 81 Eng. bot 74
2991 alp&tre Gau.	Alpine	i jjnl	SwitzerL 1819. S co Vil.deLS.7
2992 anisophyllum VU.			
2992 fr4gile Poir.	brittle	O un	S. Europe 1824. S co

2993 <i>Iffl've Thuil.</i>	smooth	■ ■ ■ ■ ■	jn.	■ ■ ■ ■ ■	S. Europe 1820. D co	Vil. del. 2.7
299* <i>supinum Lam.</i>	supine	■ ■ ■ ■ ■	■ ■ ■ ■ ■	■ ■ ■ ■ ■	Europe 1819. D co	
2995 <i>tenuifolium All.</i>	fine-leaved	■ ■ ■ ■ ■	■ ■ ■ ■ ■	■ ■ ■ ■ ■	S. France 1824. D co	
2996 <i>baldcnse Spr.</i>	Mount Baldo	■ ■ ■ ■ ■	■ ■ ■ ■ ■	■ ■ ■ ■ ■	S. Europe 1819. D co	
2997 <i>tenuissimum Bieb.</i>	slenderest	■ ■ ■ ■ ■	■ ■ ■ ■ ■	■ ■ ■ ■ ■	Tauria 1818. S co	
2998 <i>Jinifilum L.</i>	Flax-leaved	■ ■ ■ ■ ■	■ ■ ■ ■ ■	■ ■ ■ ■ ■	S. Europe 1759. D co	Bar. ic. 583
2999* <i>nigdum Vil.</i>	rigid	■ ■ ■ ■ ■	■ ■ ■ ■ ■	■ ■ ■ ■ ■	1778. D co	
3000 <i>cimbricum All.</i>	cinereous	A un	■ ■ ■ ■ ■	■ ■ ■ ■ ■	S. France 1821. D co	Al. ped. 774
3001 <i>sylvaticum L.</i>	wood	A un	■ ■ ■ ■ ■	■ ■ ■ ■ ■	S. Europe 1658. D co	Fl. dan. 609
3002 <i>aristitum L.</i>	awned	A un	■ ■ ■ ■ ■	■ ■ ■ ■ ■	ItaW 1699. D co	* Boc. mu. 83.75
3003 <i>campanul&tum Vil.</i>	campanulate	A pr	■ ■ ■ ■ ■	■ ■ ■ ■ ■	S. Europe 1821. D co	Jac. au. 1.81
	glaucum <i>Jac.</i>					
3004 <i>purpureum L.</i>	purple	3t	■ ■ ■ ■ ■	1 jnjl	■ ■ ■ ■ ■	Switzerl. 1731. D co
3005 <i>anglicum E. B.</i>	English	£	■ ■ ■ ■ ■	i jnjl	England Wales. D s.l	Eng. bot. 384
3006 <i>helveticae Weigel</i>	Swiss	£	■ ■ ■ ■ ■	I jnjl	Switzerl. 1819. D co	Ac pel. 1714.15.1
	sax&title <i>Dec.</i>					
3007 <i>pumilum Lam.</i>	dwarf	■ ■ ■ ■ ■	■ ■ ■ ■ ■	■ ■ ■ ■ ■	S. Europe 1820. D co	Lam. il. 60.2
3008 <i>scabrum Jac.</i>	rough	■ ■ ■ ■ ■	■ ■ ■ ■ ■	■ ■ ■ ■ ■	Austria 1819. D co	Jac.au. 5.422
3009 <i>saxatile L.</i>	rock	■ ■ ■ ■ ■	■ ■ ■ ■ ■	■ ■ ■ ■ ■	Britain hea. D <p	Eng. bot. 815
3010 <i>microstperm Def.</i>	small-seeded	■ ■ ■ ■ ■	■ ■ ■ ■ ■	■ ■ ■ ■ ■	S. Europe 1819. S co	
3011 <i>microstperm Vahl</i>	small-fruited	■ ■ ■ ■ ■	■ ■ ■ ■ ■	■ ■ ■ ■ ■	Europe 1819. S co	Dec. ic. 26
3012 <i>parisiense Lk.</i>	Parisian	■ ■ ■ ■ ■	■ ■ ■ ■ ■	■ ■ ■ ■ ■	France 1817. S co	Kng. bot. 1641
3013 <i>tricorne Sm.</i>	three-horned	■ ■ ■ ■ ■	■ ■ ■ ■ ■	■ ■ ■ ■ ■	Britain hhdg. S co	Eng. bot. 2173
3014 <i>verrucosum E. B.</i>	waited	■ ■ ■ ■ ■	■ ■ ■ ■ ■	■ ■ ■ ■ ■	Britain 1819. D co	
	saccharatum <i>All.</i>					
3015 <i>Cucullaria R. S. Valantini</i>	Cucullaria	■ ■ ■ ■ ■	■ ■ ■ ■ ■	■ ■ ■ ■ ■	Valantini Cucullaria L.	
		my.jl	■ ■ ■ ■ ■		Levant 1780. S co	Bux. c. 1.19 2
3016 <i>Hpinium E. B.</i>	spurious	■ ■ ■ ■ ■	■ ■ ■ ■ ■	■ ■ ■ ■ ■	Britain cor. fi. S co	Eng. bot. 1871
3017 <i>boreale L.</i>	northern	■ ■ ■ ■ ■	■ ■ ■ ■ ■	■ ■ ■ ■ ■	Britain moun. D co	Eng. bot. 105
3018 <i>brevifolium Stev.</i>	short-leaved	■ ■ ■ ■ ■	■ ■ ■ ■ ■	■ ■ ■ ■ ■	Caucasus 1819. I) co	
3019 <i>rotundifolium L.</i>	round-leaved	■ ■ ■ ■ ■	■ ■ ■ ■ ■	■ ■ ■ ■ ■	Europe 1821. D co	Jac. au. 94
3020 <i>ellipticum L.</i>	elliptic	■ ■ ■ ■ ■	■ ■ ■ ■ ■	■ ■ ■ ■ ■	1819. D co	
3021 <i>suavolens IVahl.</i>	sweet-scented	■ ■ ■ ■ ■	■ ■ ■ ■ ■	■ ■ ■ ■ ■	N. Europe 1821. W. h.b. 66	
3022 <i>minimum il. ^5.</i>	least	■ ■ ■ ■ ■	■ ■ ■ ■ ■	■ ■ ■ ■ ■	S. Europe 1818. D co	Al. ped. 77.1
	mur&le <i>All.</i>					
3023 <i>verticill&tum Dan.</i>	whorled	■ ■ ■ ■ ■	■ ■ ■ ■ ■	■ ■ ■ ■ ■	France 1824. S co	Loi. no. 2.1
3024 <i>capillare Cav.</i>	capillary	■ ■ ■ ■ ■	■ ■ ■ ■ ■	■ ■ ■ ■ ■	Spain 1820. S co	Cav. ic. 2.191.1
3021 <i>pilosum H. K.</i>	hairy	■ ■ ■ ■ ■	■ ■ ■ ■ ■	■ ■ ■ ■ ■	N. Amer. 1778. D co	
3026 <i>maritimum L.</i>	maritime	■ ■ ■ ■ ■	■ ■ ■ ■ ■	■ ■ ■ ■ ■	France 1826. D co	
307* <i>caucasicum Lag.</i>	Caucasian	■ ■ ■ ■ ■	■ ■ ■ ■ ■	■ ■ ■ ■ ■	Caucasus 1824. S co	
■ ■ ■ ■ ■	villous	■ ■ ■ ■ ■	■ ■ ■ ■ ■	■ ■ ■ ■ ■	S. France 1824. D co	Bar. ic. 82
3029 <i>tau>icum JR.*S.</i>	Taurian	■ ■ ■ ■ ■	■ ■ ■ ■ ■	■ ■ ■ ■ ■	Tauria 1818. D co	
	Gällum reflellum <i>Presl.</i>	■ ■ ■ ■ ■	■ ■ ■ ■ ■	■ ■ ■ ■ ■		
3050 <i>parine l.</i>	Cleavers	■ ■ ■ ■ ■	■ ■ ■ ■ ■	■ ■ ■ ■ ■	■ ■ ■ ■ ■	Eng. bot 816
3031 <i>infestito W. * K.</i>	troublesome	■ ■ ■ ■ ■	■ ■ ■ ■ ■	■ ■ ■ ■ ■	S. Europe 1817. ■ ■ ■ ■ ■	W. & K. 3.202
3032 <i>scaberrimum Vahl</i>	roughest	■ ■ ■ ■ ■	■ ■ ■ ■ ■	■ ■ ■ ■ ■	Egypt 1821. ■ ■ ■ ■ ■	
	hispidum <i>W. en.</i>					
3033 <i>gras'cum L.</i>	Grecian	■ ■ ■ ■ ■	■ ■ ■ ■ ■	■ ■ ■ ■ ■	Candia 1798. D co	Alp. ex. 167.166
		A cu	■ ■ ■ ■ ■	■ ■ ■ ■ ■		
349. JIU'BIA L.	MADDER.	(Iluber, red; the roots as a dye.)			Rubidace. 9.-22	
3034 <i>tincturum L.</i>	dyer's	-* A ■ ■ ■ ■ ■	Y	■ ■ ■ ■ ■	S. Europe 159a D s.l	Lam. il. 60.1
3035 <i>perigrina L.</i>	foreign	Jc A w	■ ■ ■ ■ ■	■ ■ ■ ■ ■	England bu. pi. D co	Eng. bot. 851
3036 <i>cordifolia L.</i>	heart-leaved	-* A cu	■ ■ ■ ■ ■	■ ■ ■ ■ ■	Siberia 1783. D pi	Pal. it 3.2.1
3037 <i>Boccone Pet</i>	Bocccone's	Si A or	■ ■ ■ ■ ■	■ ■ ■ ■ ■	Italy 1823. D sp	
3038 <i>lucida L.</i>	bright	B. un	■ ■ ■ ■ ■	■ ■ ■ ■ ■	Majorca 1762. C lp	Fl. gr. 142
3039 <i>fruticosa Jac.</i>	shrubby	«.i un	■ ■ ■ ■ ■	■ ■ ■ ■ ■	Canaries 177* C PJ	Jac. ic. 1.25
3040 <i>sptndens</i>	shining	■ or	■ ■ ■ ■ ■	■ ■ ■ ■ ■	Spain 1812. C Lp	
			■ ■ ■ ■ ■	■ ■ ■ ■ ■		
3041 <i>galoides Poir.</i>	Galium-like	«.» Jor	2 jn.au	■ ■ ■ ■ ■	■ ■ ■ ■ ■	Canaries 1816. ■ ■ ■ ■ ■
3042 <i>angustifolia L.</i>	narrow-leaved	«.» LJ or	2 jn.au	■ ■ ■ ■ ■	■ ■ ■ ■ ■	Spain 1772. ■ ■ ■ ■ ■
						Lam. il. 60.2
* 350. ASPE'RULA L.	WOODHOOF.	(Diminutive of asper, rough; leaves.)			Rubiacete. 24. —31.	
3043 <i>Isevg&ta L.</i>	smoothed	■ ■ ■ ■ ■	■ ■ ■ ■ ■	■ ■ ■ ■ ■	S. Europe 1775. I) s.l	M. h. 21.4
3044 <i>taurina L.</i>	bull	■ ■ ■ ■ ■	■ ■ ■ ■ ■	■ ■ ■ ■ ■	Italy 1739. D s.l	
3045 <i>pyrenaica L.</i>	Pyrcean	■ ■ ■ ■ ■	■ ■ ■ ■ ■	■ ■ ■ ■ ■	Spain 1821. D co	
3046 <i>longiflora W.ScK.</i>	long-flowered	■ ■ ■ ■ ■	■ ■ ■ ■ ■	■ ■ ■ ■ ■	Hungary 1821. D co	W. & K. 2.150
3047 <i>aristata L.</i>	awned	■ ■ ■ ■ ■	■ ■ ■ ■ ■	■ ■ ■ ■ ■	S. Europe 1823. D co	
3048 <i>supina Bieb.</i>	supine	■ ■ ■ ■ ■	■ ■ ■ ■ ■	■ ■ ■ ■ ■	Caucasus 1821. D co	
3049 <i>brevifolia Ven.</i>	short-leaved	■ ■ ■ ■ ■	■ ■ ■ ■ ■	■ ■ ■ ■ ■	Europe 1823. D co	Ven. eels 63
3050 <i>cynanchica W.</i>	Cynanche	■ ■ ■ ■ ■	■ ■ ■ ■ ■	■ ■ ■ ■ ■	England ch.hil. D s.l	Eng. bot 33
3051 <i>alpina Spr.</i>	alpine	■ ■ ■ ■ ■	■ ■ ■ ■ ■	■ ■ ■ ■ ■	Caucasus 1820. D co	
3052 <i>rígida Sm.</i>	rigid	■ ■ ■ ■ ■	■ ■ ■ ■ ■	■ ■ ■ ■ ■	Greece 1819. D co	Fl. gr. 118
3053 <i>crassifolia L.</i>	thick-leaved	■ ■ ■ ■ ■	■ ■ ■ ■ ■	■ ■ ■ ■ ■	Levant 1775. D s.l	
3054 <i>tomentosa Ten.</i>	tomentose	■ ■ ■ ■ ■	■ ■ ■ ■ ■	■ ■ ■ ■ ■	S. Europe 1817. D co	Fl.gr. 119
	inc&naSm.					
3055 <i>scabra Presl</i>	rough	■ ■ ■ ■ ■	■ ■ ■ ■ ■	■ ■ ■ ■ ■	Italy 1824. D co	
3056 <i>tincturia L.</i>	dyer's	■ ■ ■ ■ ■	■ ■ ■ ■ ■	■ ■ ■ ■ ■	Europe 1764. D s.l	Tab. ic. 733.1
3057 <i>mont&ta Kit.</i>	mountain	■ ■ ■ ■ ■	■ ■ ■ ■ ■	■ ■ ■ ■ ■	Hungary 1801. D co	
3058 <i>arvensis W.</i>	corn-Held	■ ■ ■ ■ ■	■ ■ ■ ■ ■	■ ■ ■ ■ ■	Europe 1596. S co	Lob. ic. 801.8
3059 <i>arcadiensis B. M.</i>	Arcadian	■ ■ ■ ■ ■	■ ■ ■ ■ ■	■ ■ ■ ■ ■	Arcadia 1819. D co	Bot mag. 2146
3061 <i>hirta Ram.</i>	hairy	■ ■ ■ ■ ■	■ ■ ■ ■ ■	■ ■ ■ ■ ■	Pyrenees 1817. D co	
3061 <i>hiruuta Brot.</i>	hirsute	■ ■ ■ ■ ■	■ ■ ■ ■ ■	■ ■ ■ ■ ■	Portugal 1819. D co	
3012 <i>rhizoides Bieb.</i>	Galium-like	^ A or	4 jn.au	■ ■ ■ ■ ■	S. Europe 1710. D co	Jac. au. 1.81
Crifium Klaficum L.						
3063 <i>humiflisa lies.</i>	trailing	-* A or	i jn.au	■ ■ ■ ■ ■	Caucasus 1818. D co	
Galium humifusum						
3064 <i>longifolia Sm.</i>	long-leaved	§ A or	1 jl.au	R	S. Europe 1820. D co	Fl. gr. 117
3065 <i>Aparine Bes.</i>	Cleavers	§ A or	2 jl.au	W	S. Europe 1818. D co	Fl.gr. 1.117
3066 <i>odor&ta L.</i>	sweet-scented	^ A or	J my.jn	W	Britain woods. D s.l	Eng. bot 755
351. SHERA'RD/il L.	FIELD MADDBB. {W. Sherard, a celebrated British botanist.)				Rubidete. 2.	
3067 <i>arvensis L.</i>	corn field	O w	\ ap.g.	B	Britain cor. fi. S co	Eng. bot 891
3068 <i>muralis L.</i>	wall	O un	ijn.au	Y	Italy 1805. S co	Al.ped.77.1
	Gällum murale All.					

352. SPERMACCKCE L.	BUTTON-WEED.	(<i>Sperma</i> , seed, a <i>Anote</i> , a point; capsule.)	Rubiaceae. 22.—68-
3069 glabra Mx.	smooth	O or 1 jn.ju W	KAner. 1823. S P.1
3070 tuodina Mx.	Diodia-like	O un * jn.jl W	Jamaica 1810. S co
3071 ancēdens TV.	ascendent	Q un 1 in.jl V	W. Indies 1820. S co
3072 hispida L.	bristly	im un 2 jn.jl W	E. Indies 1741. S S.1 Mur.3.6
3073 latifolia Aubl.	broad-leaved	OB un 1 jl. W	Guiana 1803. S 8.1 Aub. gui. 19.1
3074 strigosa B. M.	meagre	Q un 1 jl.au W	W. Indies 1760. S 8.1 Bot mag. 1558
3075 cornifolia Fis.	Dogwood-leaved	■■■ un 1 my.jn R	Brazil 1819. S S.1
3076 stylosa Lk.	long-styled	■■■ un 1 my.jn W	Manilla 1819. S 8.1
3077 suffruticosa Jac.	sum-uticose	tL 1 lun i.jn.au W	1824. Si Jac. sc. 322
3078 radicans Aubl.	rooting	jE 2 un 1 jl. W	Guiana 1803. S S.1 Aub.gui. 1.20.4
3079 las'vis Lam.	smooth	Q un 1 my.jl Str	W. Indies 1820. S co Sl. jam. 1.94 2
3080 sporula Aubl.	rough	Q un 1 my.jl W	Trinidad 1819. S co
3081 teuijor L.	slender	Q or 1 jl.au Pk	W. Indies 1732. S co
3082 lanceolata Lk.	lanceolate	O or 1 jl.au 9 ...	W. Indies 1818. S p.1
3083 imoTia Vahl	Flax-leaved	■■■ un 1 my.jl W	W. Indies 1818. S co
3084 scabra TV.	rough	■■■ un 1 my.jl W	E. Indies 1818. S co
3085 hirta L.	hairy	O un 1 my.jl W	W. Indies 1Hia S co
3086 villosa Sum.	villous	■■■ un 1 my.jl W	W. Indies 1810. S co
3087 rfiba Joe.	red	£ ■■■ un 1 my.jn p	1804. S 8.1 Jac. sc. 256
3088 Ffischer* Lk.	Fischer's	■■■ un 1 my.jn W	Jamaica 1821. S 8.1
3089 mucronata Nees	mucronata	tE 1 un 2 my.jl W	Jamaica 1822. D 8.1
3090 Roxburgh** Spr.	Roxburgh's	t • a or 1 jl.au W	E. Indies 1818. S P.1
353. DKVDIA L.	DIODIA.	(<i>Diodos</i> , a journeying, a way-side plant.)	Rubiaceae. 4.—10.
3091 virginica L.	Virginian ^	* A cu J.jn.jl W	Virginia 1820. S co Jac. ic. 129
3092 verticillata PhM	verticillate	i* EJcu i. jn.jl W	W. Indies 1821. S co
3093 prostrata Suit.	prostrate	JU Q cu 4 jn.jl W	W. Indies 1818. S co
3094 sarmcntosa Sun.	twiggy	JL ■■■ I jn.au W	W. Indies 1821. C sp
354. KN01LM L.	KNOXIA.	(<i>Robert Knox</i> , an English traveller in 1681.)	Rubiaceae. 3.
3095 zeylanica L.	Ceylon	n. O or 1 jl.au W	Ceylon 1826. S co Bur. in. 13
3096 corymbosa TV.	corymbose	ttQor I jn.au W	E. Indies 1818. S co Lam.IL59.2
Spermacoce sumatrensis Retz.			
3097 umbellata Sol.	umbellate	tL Q or } jn.au W	E. Indies 1820. S co
Spermacoce teres Rox.			
355. BIGELCV/ii Spr.	BIGELOVIA.	(<i>Prof. Biegelow</i> , of North America.)	Rubiaceae. 3.—11.
3098 verticillata Spr.	whorled	- Q o r lijn.au W	Africa 1732. S s.p DLel.277.358
Spermacoce verticillata L.			
3099 commutata Spr.	changed	tf FAlor i. jn.jl W	W. Indies 1818. D s.l
Spermacoce verticillata Host			
3100 stricta Spr.	strict	OJ pr f jn.jl W	E. Indies 1820. S s.l
Spermacoce stricta L.			
356. CRUCIANE'LLA L.	CHOSSWORT.	(Diminutive of crux, a cross j leaves.)	Rubidex. 13.—16.
3101 angustifolia L.	narrow-leaved	O cu i. jn.jl Y	France 165a S co Ex. bot 2.109
3102 latifolia L.	broad-leaved	O cu i. jn.jl G	France 1633. S co Bar. i c 520
3103 seytiaca L.	Egyptian	Q cu i. jn.jl Y	Egypt 1800. S co
3104 pátula L.	spreading	8 cu i. jn.jl Y	Spain 1798. S co
3105 pubescens TV.	pubescent	■■■ cu i. jn.jl P	Candia 1799. C l.p
3106 glomerate Bieb.	glomerated	A CU 1 jl.au Y	Iberia 1824. S co
3107 ciliflu Lam.	ciliated	■■■ cu ijl.au Y	Levant 1805. S co
3108 tuberculosa Cau.	tuberclose	O cu 1 il.au Y	Spain 1826. S co
3109 maritima L.	maritime	■■■ cu 1 jl.au Y	France 1640. C l.p
3110 amricana Mil.	American	■■■ cu 1 jl.au Y	S. Amer. 1780. S co
3111 monspelica L.	Montpelier	■■■ cu ijl.au Y	France 1791. S ls.p
3112 molluginoides Bieb.	Mollugo-like	k A cu 1 jl.au G	Caucasus 1800. D co Bux.c.2.30.1
3113 anomala Bieb.	anomalous	■■■ cu 1 jn.au Y	Caucasus 1820. S co
357. CALLICA'RPA L.	CALLICARPA.	(<i>Knlos</i> , beautiful, <i>karpos</i> , fruit.)	Ferbenaceat. 12. — 24.
3114 americana L.	American	ft i. or 6 jn.jl R	N. Amer. 1724. C s.p Cat car. 247
3115 &naL.* ^♦	hoary	* or 3 ... P	E. Indies 1799. C s.p Bot. mag. 2107
tomentosa Lam.			
3116 lanata L.	woolly	■■■ or 4 jn.jl P	E. Indies 1788. C l.p
3117 incifta Box.	hoary	■■■ or 4 jn.au R	E. Indies 1800. C l.p
3118 macrophylla Vahl	long-leaved	■■■ or 6 - - Pk	India 1808. C i.p Vahl By. 3.53
3119 lanceolana Rox.	lanceolate	■■■ or 4 jn.au P	E. Indies 1M& C l.p
3120 ferruginea Sun.	rusty	■■■ or 2 jn.jl B	Jamaica 1794. C l.p
3121 longitolia Lam.	long-leaved	Ljp ■■■ 3 mr.ap W	China 1825. C co Botreg. 864.
3122 reticulata Swx.	netted	i. lor 4 jn.au R	Jamaica 1820. C l.p
3123 rubella B. B.	reddish	■■■ or 2 my. Fk	China 1822. C co Bot. reg. 883
3124 purpurea J.	purple	■■■ or 3 jl.n P	E. Indies 1822. C l.p
3125 arborea Rox.	tree	■■■ or 12 jl.n P	RIndies 1820. C l.p
358. AQUA'RTIA Jac.	AQUARTIA.	(<i>B. Aquart</i> , Jacquin's companion in America.)	Solohce. 1. — 2.
3126 tomentifusa Jac.	woolly	• D or 4 ... W	S. Amer. 1819. C s.l Jac.am. 15
359. WITHERI'NG/i*	Herit	WITHERINGIA.	Soldwar. 6.—16.
3127 wlanacea Herit.	solanaceous	£ 2 2 or 1 my.s Y	S. Amer. 1742 D l.p Her. ser. 33.1
3128 stramonifolia Klh.	Stramon.-lvd	MCJor 3 my.au Y	Mexico 18J3. C l.p
3129 crassifolia Dunal	thick-leaved	• L J or 2 my.au Y	C. G. H. 1706. C l.p Di. el 273.332
3130 montana Dunal	mountain	■■■ IAJ or 1 my.au W	Peru 1822. C 1 p Feu. ob. 46
3131 phyllantha Dunal	leaf-flowered	O or 1 jl.au Y	Peru 1822. C l.p Cav. ic. 4. ST>91
3132 pinnatifida Dunal	pinnatifid	£ iAJ or 1 jn.au B	Peru 1822. C l.p FLper.2.170.6
360. ^GI'PHILA L.	JEGIPHILA.	(Aix, a goat, philos, dear; liked by goats.)	Verbendceee. 7.—14.
3133 martinic^nsis L.	Martinique	• Qor 6 n . W	W. Indies 178a S p.1 Jacob 2.27
3134 arborescens W.	arborescent	• o r 10 - -	W. Trinidad 1823. C l.p Aub. gui. 1.24
3135 elata Swx.	tall	■■■ 11 - - Pa.Y	W. Indies 1823. C p.1
3136 foetida Sun.	fetid	■■■ 11 Li	W. Indies 1800. C l.p
3137 trifida Sun.	trifid	■■■ 11 W	Jamaica 1826. C p.1
3138 diffisa Andr.	diffuse	■■■ 11 W	W. Indies 1804. C l.p Bot. rep.578.1
8139 oboviflora Andr.	obovate	■■■ 11 Y	W. Indies 1804. C l.p Bot. rep.5/S.2

3221 brasiliensis Jac.	Brazilian	* D o r	10	■	O	Brazil	1822.	C	l.p	■ ■ ■ ,	
3222 salviifolia Lam.	Sage-leaved	* i .Jor	3	au.8	C	C.G.H.	17(50).	C	8.1	Jac. so. 1.28	
3223 jalicifolia Vahl	Willow-leaved	H' O or	6	S. Amer.	1823.	C	l.p		
3224 jraligna W. en.	Willow-like	«i .Jor	6	au.8	W	C. G. H.	1816.	C	l.p	Jac. sc. 1.29	
3225 madagascariensisraA/Madagascar	• • o r	10	-	-	O	Madagascar	1824.	G	Lp	Lam.iL 1.69.3	
3226 paniculata Wai.	paniced	* O ° r	14	Cap	W	Nepal	1823.	C	l.p		
3227 diversifolia B. & S.	various-leaved	SO or	6	-	...	Java	1823.	C	l.p		
3228 americana L.	American	■ ■ ■	10	au	Y	Mexico	188G.	C	l.p		
366. E'XACVM L.	EXACUM.	'	(Ex, out of, ago, to drive; expels poison.)			Gettijanea?	2.—14.				
3929 tetragnum Rtb.	four-angled	O pr	1^ Jls	B	Nepal	1820.	S	p.1			
3250 pulchellum Ph.	pretty	O pr	i au	Pk	N. Jersey	1826.	S	p.1			
367. CO UTO VBEA	Aubl.	COUTOUEBA.	(IU name in Guiana.)			Gentihneea.	3.				
3231 rambobum Aubl.	branchy	O or	1J jl.au	W	Brazil	1824.	S	p.1	Aub.gui.	1.28.1	
Exacum ramifsum Void											
3232 spicatum Aubl.	spiked	£ IDS or	1J jl.au	W	Maranh.	1823.	S	p.1	Aub. gui.	1.27	
JS'xacum spic&tum Vahl											
3833 verticillatum G.Don	whorled-headtd	tf Ql	1	jl.au	W	Trinidad	1818.	S	p.1		
J^xacum verticillatum W.	Gentidina	verticillata L.									
368. MICROCA'LEZ*. MICROCALE.		(Mikros, small, kalos, pretty.)				Gentia*ea».	1.				
3234 filiforme Lk.	filiform	O pr	Jjn.jl	Y	Britain	sa.ma.	S	i.i	Eng.bot	235	
fx acum filifirme E. B.											
369. PLA'DERA Boar.	PLADER*.		{Pladaro, abounding in juice.)			Gentihneetc.	2.-5.				
3235 virgite Box.	twiggy	OJor	1 jl.au	K	E. Indies	1820.	S	p.1			
3236 decussate Box.	decussate	OJor	1 jls	W	W. Indies	1816.	S	p.1			
370. SEBM'A R. Br.			(A. Seba, an apothecary of Amsterdam.)			Gentihneea.	5..-8.				
3237 ovate B. Br.	ovate	uQjpr	Jau	H	N. Holl.	1820.	S	p.1	Lab. n. li.	1.52	
3838 albens it. Br.	whitish	Opi	i a u	W	C. G. H.	1820.	S	s.l	Bur. at.	74.4	
3239 cordate R. Br.	hezrt-leaved	O or	üJau	Y	C. G. H.	1815.	S	co	Bur. af.	74.5	
3240 afirea R.8/S.	golden	■	jnau	Y	C. G. H.	1824.	D	s.1	Pluk. al.	275.3	
3241 guianensis Spr.	Guiana	■	ijJau	Br	Guiana	1825.	S	p.m.s	Aub. gui.	1.26.1	
Ä'xacum guianensis Aubl.											
371. MITRASA'CME R Br.	MITRASACME.	(Mitra, a mitre, akme, a point)				Gentihnete.	3.—20.				
3242 polymorpha B. Br.	multiform	f 1ai pr	i jn.jl	W	X. Holl.	182G.	S	s.1			
3243 canescens B. Br.	cancscens	£ IAI P'	i jn.au	W	N. Holl.	1824.	D	s.1			
3244 wrpyUifolia R. Br.	Thyme-leaved	£ ^ pr	ijn.au	W	N. Holl.	1826.	D	s.1			
372. Fna^SER^ Walt	FRASEKA.	(J. Fraser, a collector of plants in North America.)				Gentiahear.	1.				
3245 carolinensis Walt	Carolina	£ Q or	4 jl.au	G.Y	Carolina	1795.	S	co			
' Walter^ Mx.											
■ 373. PENJEM L. PEHSA.	(P. Peru, auth. with Lobe\ of Adversaria Botan., 1570.)					Peutaceat.	7.—14.				
3246 myrtoides L.	Myrtle-like	■ i_J or	2 my.jn	R	C. G. H.	1816.	C	p.1			
3247 fruticulata L.	fruticulose	■ J or	1 my.jl	K	C. G. H.	1822.	C	p.1			
3248 mucronata L.	mucronata	■ or	8 jn.jl	Y	C. G. H.	1787.	S	p.1	Ven. mal.	87	
3249 squamosa L.	scaly	■ or	1 jn.jl	R	C. G. H.	1787.	S	p.1	Bot. reg.	106	
3250 sarcocolla Thun.	thick-necked	■ T or	1 jn.jl	R	C. G. H.	1825.	S	p.1	Pluk. m.	446.6	
3251 marginata L.	marginated	M I or	li my.jl	R	C. G. H.	1816.	C	p.1			
3252 lateriflora Thun.	side-flowering	si t_J or	1 jn.jl	R	C. G. H.	1825.	S	p.1	Mag.ber.	1.3.2	
374. BLJE'RIA L.	BLKRIA.		R.S., physician at Boston, Line.)			Ericete.	11..18.				
3255 ericoides L.	Heath-hke	■	2 au.o	P	C. G. H.	1774.	C	s.p	Pet g.	471.2.10	
3254 muscosa H. K.	mossy	■	1 jn.au	P	C. G. H.	1774.	C	l.p			
3255 fasciculata W.	fascicled	■	2 jn.jl	...	C. G. H.	1812.	C	s.p			
3256 paucifolia Wnl.	few-leaved	■	2 jn.jl	...	C.G.H.	1812.	C	s.p	Wen. c.	2.43	
3257 articulata L.	jointed	M or	2 my.jn	Pk	C. G. H.	1795.	C	s.p	Lam. a	78	
3258 scabra W.	scabrous	• or	2 my.jn	P	C. G. H.	1824.	C	a.p			
3259 depresso Lich.	depressed	H i or	1 jn.jl	? ...	C. G. H.	1816.	C	s.p	Wen.c.	1.31	
3260 purpurea L.	purple-covered	• L_ or	2 my.jn	P	C. G. H.	1791.	C	s.p			
3261 dumosa Wnl.	bushy	• LJ or	2 jn.jl	...	C. G. H.	1806.	C	s.p			
3262 glab^ua Wnl.	smoothish	• I or	2 jn.jl	...	C. G. H.	1816.	C	s.p	Seb. thes.	1.20.2	
3263 ciliaris L.	ciliated	• i or	2 jn.au	W	C. G. H.	1795.	C	s.p	Wen. c.	2.49	
375. SYMPIE'ZA Lich.	SYMPIEZA.	(Sympiezo, to press together.)				Ertce*.	1.				
3264 capitellata Lich.	small-headed	sH _Jor	fijn.au	...	C. G. H.	1812.	C	s.p	Wen. c.	2.37	
Blae'ria bracteata Wnl.											
376. CHOME'LLf Jac.	CHOMELIA.	(J. B. Chomel, a Fr. bot, phys. to Louis XV.)				Rubidcear.	2.—3.				
3265 spinosa Jac.	spiny	• D < »	12	...	W	W.Indies	1793.	C	p.1	Jac.am.	18. 13
Ix6raspinosa Lam.											
3266 fasciculata Sim.	fascicled	* • or	5	-	W	W. Indies	1825.	C	p.1		
Ix6rafasciculata Sum.											
377. ADI'NA Sal.	APINA.	(Adinos, clustered; its flowers in heads.)				Rubideaa.	1.—2.				
S267 globiflora Sal.	globe-flowered	is CD or	3 jl.au	W	China	1804.	C	s.l.p	Par. Ion.	115	
NaucleaAdina Sm.											
378. BOUVA'RDL* Sal.	BouVARnu.	(Dr. Bouvard, a superin. of the Paris garden)				Rubiucee.	3.—7.				
3268 triphilla Sal.	three-leaved	* _ or	2 ap.n	S	Mexico	1794.	C	s.p	Par. Ion.	88	
Jacqini Hum.	Houstonia coccinea B. R.	H i or	2 ap.n	S	Mexico	1794.	C	s.p	Bot. reg.	107	
2 pubescens	pubescent	*M or	2 ap.n	S	Mexico	1794.	C	s.p			
3 glabra	smooth	A i or	2 Jls	R	S. Amer.?	1814.	C	Lp	Bot. reg.	245	
3269 versicolor Ker.	various-colored	A i or	2 Jls	R	S. Amer.?	1814.	C	Lp			
3270 longiflora Kth.	long-flowered	■ ■ ■	2	...	W	Mexico	1827.	C	s.p	Cav. ic.	572.1
379. IXWRA L.	IXORA.	(A Malabar idol, to which its flowers are offered.)				Rubidceee.	19.—SO.				
3271 grandiflora Ker.	great-flowered	* O or	4 au	O	E. Indies	1814.	C	l.p	Bot reg.	154	
coccinea L.											
9S7S Bandhda Rox.	Bandhooka	• Q or	3 jl	F	E. Indies	1815.	C	p.l	Bot. reg.	513	

			5Jn.au	S	E. Indies	1823.	C p.l	Ru.am.4.46
							M	Botmag.169
	saffron-colored ^colored ft	• • or • or white	3 aus 2 au 4 au	O R W K	Indies 1822. E Indies 1822. C p.l	Bot. reg. 782 Rot cab.1048 Bot. reg.100		
3282 P. IS* ^	wedge-leaved twice-branched	• • or HQor	4 jn.au 2 A.	W E. Indies E. Indies	1768. 1822.	C fl C p.l	BoL reg. 648	
3281 £11.22 U*	Pavetta	M a or	3 jn.au	W E. Indies	1800. 1800.	C p.l C p.l	Bot. cab.009 And. rep. 78	
3281 C23.2	small-flowered	• • or	20 au.o	W E. Indies	1800 1823.	C p.l C p.l	Vahl.3.11.52	
	bearded	• • or	12 jnjl	W E. Indies	1823. 1820.	C p.l C p.l	Bot.mag.2505	
		*Dor	4 jn.au	W P Bengal	1819. 1819.	C p.l C p.l	Bot.rcg.540	
l^firoa^i				K Indies	181&	C Si		
3288flavaL.c. *	yellow	iQot	3 - Y	E. Indies	1825.	C p.l		
							Aub.gui.158	

WI-C ATESBJS'il W. LILT THORN. (*Si Catesby*, author of the Nat. Hist. of Carolina.) *Rubidacee.* 3
 ①Si<pin6sa W.
 '292 Idūfflia Lindl. spinous J C o r 12 my.s Y I. Provid. 172A C s.p Bot. mag. 131
 <S*3parvifloraPars. broad-leaved » D o r 41 jnjl . Y W. Indies 1823. C p.l Bot reg. 858
 small-flowered H Q o r 2 jnjl W Jamaica 1810. C Lp SLjam.2.207.1

PAVETTA. (The name in Malabar.) *Rubidacee.* 3.
 Indian • • o r 4 au.o W E. Indies 1791. C p.l Bur. in. 13.3
~~Lam. Pav4tta Rox.~~
 • i j o r 4 mr.o W China 1799. C p.l Bot. reg. 198

SUN. ERNODEA. (*Ernodes*, branching) in allusion to habit of plant i *Rulwccce* 1—3

COCCOPTSELM. (To***, fruit W ^ a vase; berries.) <i&irfc^. L 71
 creeping E 3 un l my P W. Indies 1793. D s.p Br. jam. 6.1
 COCCOPTSELM. W. IESij*** an Engl.Uhman who travelled in Virginia.) fl<Wrfee*. 1.
 creeping or i j n W N. Amer. 1761. L, s.p Cat. car. 1.20

SJEU MADDER. 7?et, <<an <'; leaves like ears.) *Rybidee.* 7.-
 * J n , f u w E-Indie. 1821. S Vp

milk.*fe O u n J U u W E. Indie, S S. S. S.

Profef-rofbotanyatFlom.ee.)

A<lmater<c4ntharom. C i.p Her. at 1.1

(m).0<f,tonnderofBotMv,etc..d.1799L) CWo ...
 i tm30 ... Fa C. G. rf 1775. C ,1 Bur. af 235.83

'telea ovata fan. SerInSSorata adagaw-ttM. C i.p Lam. il. 910

2.60

TETRANDHIA MONOGYNIA.

A5* IV.

»397. AMMA'NSJJ L.

3318 rotiitnil *Vilia Hum.*
3319 intifilia J.
3330 hi'imilw Mr.
SU'll rmni'isior L.
3*!2 verticil! its *Lam.*
SS&TbixsiHam
3355 pentambla *Box.*
- octandra *L.JU.*
»anuinolenta *Stoz,*
multipli.lira /for.
3330 racemuwi *Roth*
83) mriculii »'.
3132 bMciftm /,.
3333 vridis *Horn.*
3934 c&tptici *Hitti.*
3335 <>yl>tlaca W.

round.lenvcd
buad-leaved
hiini.li;
brancilicr
wlriori *Jlutuered*
red
ivi. it.
pcitandrmu
octandmn
bltwdy
itiflu*
many-flowered
eared
berry-bearing
Egyptian

J. jlnwMitn. dr*
O un J Jlau J'k Nepal 1819.
O un ijL.lau W W. Indie 1823.
O mn 3 jlau R N. Amer. 1823.
[fl CU 1 jlau W Guineia 18W.
K Nepal IRIS.
S s.i
P E. Indies 1778.
P K. Indies 1823.
It li Indies 1820.
R Jamaica 1803.
W. isaa S s.1
K U ii 8 i.l
P ism. S BO
l Egypt ISM. S co W,h.b.7
Ap. S co 8 .l Lam.ll77.4
P Caspian S eo
Ap Egs,t S »l W.li.b.6

Sallicube. 18.—'22.

398. FA*ii*.VRA

333fi Pten'.Li L.
Ptoriita l*ti'roia
3337 FiperUn h.
3S38 /ragddei

(Arabic name used by Avicenna.)

f • t m S U au.s	G	Jbautct	1768.	C p.1	Dr.Jarn I
					Japan 177^
					W. Indies 175).
					1835.
					Kasm. 833
					Jac uni. SJL 14

399. ZIF.TUA Sin.

3340 Smihtf Andr.
ia.iiciwhu.tu /(. Br.
3341 macrophilla lionp,
3342 Ifevigatn AM.
g34fl inicruph j-lla Amp.
33H niluia /U/.
3S4S Binun Vee.
3346 obcorritii ('un.
SHI revali'tii ('un.
3348 octandrn Stvt.

ZIEHM.

Snu III's
long, leaved
nruyoth./riuvlt
glaucous
creelung
Huufitciulose
tguHTE-ttaited
Bfarw
Smilnx-liku
ft. CJ
ovatu
antariic
glandular

{M. Zier, a loused botanic.)

L_J or 3 u>jl	W	N. S. W.	1808.	C »p	Bot mag. 1395
!_or 4 myJj	W			18CQ.	C
my,jl	W	N. HCII.	1822.	C	
_or 3 myJl	W	N. M.II.		C tp	
!_or 4 myJl	W	N. Hoi I.		C i.v Lin. Ir. 1U. 17.2	
I or 1 myJl	W	N. Hull.	18SSL	C	
I or 4 myJl	W	N. Hull.		C	
I or 4 myJl	W	N. Hull.		C i	
I(or 4 myJl	a	N. Hull.	1825.	C Sw. au. ic. in.	

*nutate**, ft—12:

400. CI'SSUS L.

3349 iritiglncia L.
3351 admits /for.
3351 latilitifl /fun.
335° glafsta Roz.
3353 cassia 4fr
3354 rejH-ii* Lam.
SiM puncUciflisa Mr.
3356 qintraueulariB /,.
3357 *Ieyoldw L.
3338 nniljcina A1/i.
3359 ovata Lnm,
3340 aiiNirutira Fen.
iiM-ruis »'.
SjG> trilhilii Lnm.
33S4 aunculala /toi.

Cusui.

Vine-sdrung
DdliliC
Urnii tl caved
glaucous
creelung
Huufitciulose
tguHTE-ttaited
Bfarw
Smilnx-liku
ft. CJ
ovatu
antariic
glandular

(Tlic Greek name or theivy.)

20 Q	India	E. — BS.
A mi 8 ... G	Indiut 18S1	Pluk. m.27.337.i
& -n ... G	E. Indie* ISC*	Ku. am. 5. IC4. i
fl_a ta 10 ... G	1818. C	K-maLV. 11
15 ... G	S. I'eone	D
!_or 4 myJj		18SL C
2 ... (;		R.mal.7.48
W ...	Careone	I"
30 ... (f	E. Indies	C pl In. am. K -It
B Dor W ... 0		C «.p Sl.jain. 1. 1U.
10 ... a	W.	co
10 ... G		18M. C El
ill) jn.au G	N. a w.	I7S0. C sj Ilr. j:iin. I 1,1
30 ... G	E. IndiM	1819. I) TO
12JLU1 G	C. G. rL	1792. C S.p 1
E. Indies	E. Indies	182J. C co B. nial. 5 45
10 ... G	K. iii-ites	Ififlt C > J:te. «c. I. 33
ii ... G	Ji m. lira	C co Jtu, ntii
A. - un 11 ... G	E. Indie*	C ll1 Sl.jatn, '
J II no 6 ... G	K. III.ik'	C co PL ic. S
winsed 6 ... 0	S. Amer.	182S. C co Jac. tun. 183,10
five-leaved B ip.i G	Japan	nSA C «.p
elmiKated Li irr 10 ... G	K. initii.'S	r.in
v:iriou»-coved th or 10 ... a	c. a 11	Big V l.p
oriental J_ un 8 ... G	F. Indies	taa u CO
fi-lcavcd Brazil 4. 1 un 10 Jl	Bralil	Ijim. 1
		BiL ni^f. iW

E. — BS.

Pluk. m.27.337.i

Ku. am. 5. IC4. i

K-maLV. 11

R.mal.7.48

Ilr. j:iin. I 1,1

But.mng.24SB

El

sj

co

CLASS IV. ORDER 2.

TETRA'NDRIA DIGY'NIA.

w. falua	w. P ¹⁷ "TM". slender-leaved	(Count de Btibfn^he celeb. French naturalist.)	C <i>onj</i> jUe*. 2. - 4 .
	O cu *jn	W England sea ca S co 2ng. bot 1313	
	perennial	Jjn.jl W France 1817. S p.l Lam.il. 87.2	
rrophylla PA.	T (» « «) - with, melon, apple; fr. accompany fl.)	llamamelUlc*. 2.	
	Virginian	or 10my.n Y NAmer.1726. L pl Dub. ax.11114	
	long-leaved	or 15 my.n Y N. Amer. 1812. L p1	
	HYPECOUM. {Hypecheo, to rattle j noise of seeds in pods.)	Papaverceai. 3. - 6 .	
procumbent	-A O or 1 jn.jl Y	S. Europe 1596. S co Schk. ban. 1.27	
pendulous	* O or § jn.jl Y	& France 1>40 S co Par. th. 372.2	
erect	O or i my.n Y	Siberia 1759. S co Am. rut. 58.9	

CLASS IV. ORDER 3.

TETRA'NDRIA TETRAGY'NIA.

* uragbga Jac.							
diuretic	■ or Myrtle-leaved	4 aus. or 4 my.au	P & Amer. W	1790. N. Amer.	L p.1 1818. L p.1	Jac. am. 24.16	lihammete. 5.—10.
Rhacomia	■	4 ... 4 ap.my 4 ...	W W W	Jamaica 1798. W. Indies 1795. Martiniq. 1826.	L p.1 C s.p. C s.l.	Jac. ic. 2.311 Fl. per. 84.6	
34 UJ 3495 STM 4. P ¹ ^A » 15 integrifolia Lam.	W LEX						
44 16 ^ 6							
... vwuvvs ^{il} n5ta Desf.							
SJWbaU^ricaJ!)^ IJ19 Pertido H. £							
Minorca	I or	10 myjn	W	Minorca 1815.	G co		
Perado	I or	10 ap.my	Pk	Madeira 1760.	G s.l	Mce. ic. 2.6	
Canary Islands	I or	16 jnjl	W	Canaries 1820.	G		
opaque	or	30 myjn	W	Carolina 1744.	G p.1	Mce. ic. 2.5	
saffron-colored	«Jtm	30 ..	W	C. G. H. 1794.	G s.p		
loose-flowered	or	20 apjn	W	Carolina 1811.	G ap	Lam. il. 355	
Chinese	M I or	10 jl	W	China 1814.	G s.f	Bot. mag. 2043	
Dahoon	A J or	6 my.jn	W	Carolina 1726	G s.l		
Cassine	Dahooon £	12 au	W	Carolina 1726.	G 81	Cat car. 1. 31	
narrow-leaved	m J or	6 myjn	W	Carolina 1H06.	G s.l		
Cassine-hke	Ml or	6 jl.au	W	.. 1820.	G ro		
emetic	t or	10 jl.au	W	Florida 1700.	G p.1	Cat. car. 2.57	
Willow-leaved	* or	5 myjn	W	Mauritius 1818.	C p.1	Jac. c. 4.2.2	
Myrtle-leaved	m or	6 jl.au	W	W. Indies 1806.	C p.1		
Paraguay Tea	m	18	W	Paraguay 1823.	C p.1	Lamb. pi. 12. ic.	
PLANERA.							
Richard's	il	(J. J. Planer, a German botanist.)			Ulm &c.e.		
K.	or	12 ap.my	Br	N. Amer. 1760.	G co		
Gmelin**	fl						
	or	12 ap.my	Br	N. Amer. IJ&	G eo		
COLDBNIA.							
trailing	J E JF ^{Ulm} k* North American botanist)						
	-* O or	k jl.au	W	E Indies 16W	Boragine*. 1.—2.		
		E 2			co Lam. iL89		

PENTANDRIA MONOGYNIA.

CLASS V.

•414. POTAMOOETON L.	POND-WEED. (<i>Potamot</i> , a river, <i>geiton</i> , near; situation.)	<i>Atismdce*</i> .	14.—44,
3436 natans L.	natant	^ A cu ft au	G Britain riv. D co Eng. bot. 1822
3437 fluitans Roth	floating	£ A cu ft jl.au	R Britain dit. D co Eng. bot. 1286
3438 heterophyllum Schreb.	vanous-lvd	* A cu ft jl.au	G Britain dit. D co Eng. bot. 1285
3439 iyrfoliatum I.	perforate	■ A cu ft jl.au	P Britain riv. D co Eng. bot. 168
3440 densum £.	close-leaved	A cu ft my.jl	G Britain dit. D co Eng. bot. 397
3441 luccns L.	shining	■ A cu ft jn.jl	G Britain dit. D co Eng. bot. 376
3442 crifspum L.	curled	■ A cu ft jn.jl	R Britain rivul. D co Eng. bot. 1012
3443 rompt'ssum L.	■ ■ ■ ■ ■	■ A cu ft jn.jl	G Britain rivul. D co Eng. bot. 418
3444 cuspidatum Schr.	cubipdate	■ A cu ft jn.jl	Br Britain rivul. 1) co Loe. pr. 66
3445 pectinatum /.	pectinate-fai	■ A cu ft jn.jl	Oi Britain dit. D co Eng. bot. 323
3446 lancolatuum J5. A	apeax-leaved	■ A cu ft jl.au	Oi England w.lak. D co Eng. bot. 1185
3447 gramineum E. B.	Grass-like	■ A cu ft jl.au	G Britain dit. D co Eng. bot. 2253
3418 pusillum L.	small	■ A cu ft jl.au	G Britain dit. D co Eng. bot. 215
3449 sct&ceum i.	■ ■ ■ ■ ■	■ A cu ft jl.au	G Britain ... D co
415. RU'PPM L.	RUPPIA.	(H. B. Ruppi, a German botanist.)	<i>Fluvidles</i> . 1.—2.
3450 maritima L.	sea	■ A cu ft jl	G Britain sa. w. d. S s.l. Eng. bot. 136
•416. SAGI'NA L.	PEARLWORT.	(<i>Sagina</i> , something nourishing, <i>Lat.</i> ; sheep-food.)	<i>CaryopMleas</i> . 6.—8.
3451 procumbens /..	procumbent	Θ w ■ my.s	W Britain " " Eng. bot. 880
3452 cer.tstdc Sm.	Cerastium-like	Θ w ■ 'jn.ji	W Britain " " Eng. bot. 169
Mocnch'a cerastoldes Spr.			
3453 tillorinus Pou.	filiform	○ cu * jn.jl	W Pyrenees 1824. S co
3454 aptala L.	apetalous	○ w / my.jn	W Britain rub. S co Eng. bot. 881
3415 maritima Don	sea-side	○ w ft my.au	W Ireland sc. alp. S co Eng. bot. 2195
3456 ercta L.	upright	○ w ft ap my	W Britain ... S co Eng. bot. 609
Mochnch'a quaternella Ehrh.			
417. IXIAJE'A L.	TILLJS.A.	(M.A. Tilli, an Italian botanist, d. 1740.)	<i>Crassuldcea</i> . 2.—4.
3467 muscosa L.	mossy	O cu ft jn.o	Pi England sa. he. S co Eng. bot. 116
3478 aquatica R. \$ S.	water	^ O cu ft jn.au	P. a N. Europe 1816. S l.p. Us. an. 6.1
4 ia BULLIA'R'Dii Dec.	BULLIARDA.	(M Bulliard, a French botanist.)	<i>Crassuldaw</i> . 1.—2.
3459 Vaillant* Dec.	Vaillant's	^ O cu ft jn.au	W S. Europe 1825. S l.p. Dec. pL 74
419. RAD'OI.A Dil.	RADIOLA.	(<i>Radiohis</i> , a little ray : radiated capsule.)	<i>CaryopMlece</i> . 1.
3460 millegiana Sm.	all-seed	O cu ft jl.au	W Britain san. pi. S bog Eng. bot 893
420. GONOCA'RPUS Thun.	GOWOCARPUS	(<i>Gonia</i> , an angle, <i>karpos</i> , fruit; seeds.)	<i>Salicdriar</i> . 1.—9.
3461 micranthus Thun.	small-flowered	o cu 1 jl.au	W China 1806. S co Thun. jap. 69.15

CLASS V. ORDER 1.

PENTA'NDRIA MONOGY'NIA.

421. MIRA'BILIS L.	MARVEL OF PEHU.	(<i>Mirabilis</i> , wonderful: flowers.)	<i>KydagInete</i> . 5.—
3462 dichotoma L.	forked	■ ■ ■ jl.au	Y Bjexico 1640. R co Mart. ce. 1.1
3463 Jaldpa L.	Jalap	■ ■ ■ 2 jn.s	R W.Indies 1596. R r.m Bot. mag. 371
1 flavia	yellow-flowered	■ ■ ■ 2 jn.s	Y W. Indies 1596 R r.m
2 alba	white-Itotvrred	■ ■ ■ 2 jn.s	W W. Indies 159a R r.m
3 rubro-alba	red and white	■ ■ ■ 2 jn.s	R.W. W. Indies 1596. R r.m
4 rubro-fl&va	red and yellow	ft ■ ■ 2 jn.s	R.Y. W. Indies 1596. R r.m
3464 hibrida I*pel.	hybrid	ft ■ ■ 2 jn.s	R 1813. R r.m
3465 longiflora L.	long-flowered	ft ■ ■ 2 jn.s	W Mexico 1759. R r.m Ex. bot 1.23
3466 suaveolens Kth.	sweet-scented	ft A or lijl.au	W Mexico 1824. R r.m
•422. ABRO'NIA J.	ABROMA.	(<i>Abros</i> , delicate, the involucrum.)	<i>Nyctaglnete</i> . 1.
3467 umbellata Lam.	'umbelled	JU A el ft ap.my	R California 1823. D s.p. Hook. ex. 194
Tricratus admirabilis Her.			
*423. PLUMB A'GO L.	LEADWOHT.	(<i>Plumbum</i> , a disorder in the eyes: med. qual.)	<i>Plumbagines</i> . &—12.
34C8 eurojMEA L.	European	^ A or 3 e.o	B S. Europe 1596. C p.l. Bot. mag. 2139
3469 /apathifolia W.	Dock-leaved	Ji jn.jl	Iberia 1822. Sk s.p
3470 zeylanica L.	Cingalese	■ ■ ■ 2 ap.s	W E. Indies 1731. Sk s.p R mal. 10. 8
3471 rosca L.	rosy	tL ■ or lft mr.jl	R E. Indies 1777. Sk r.m Bot. mag. 230
3472 »ändens-L.	climbing	LCD or	3 jn.jl W W. Indies 1699. Sk s.p SL jam. 1.133-1
3473 occidentalis Sco.	West Indian	■ ■ ■ 3 apa	W W. Indies 1817. C l.p
zeylanica 0 R. % S.	scamlecs Host noikV.		
3474 tristis // K.	dark-Jotwra/	tL i_J or lft my.jn	Br C. G. H. 1792 F l.p
3475 capensis Thun.	Cape	n. _J or lftn	B C. G. H. 1818. F l.p Bot. reg.417
424. //ELIOTRO'PIUM L.	TURNSOLE.	(<i>Helios</i> , the sun, <i>trope</i> , a turning; flowers.)	<i>Boraclnece</i> . 22.—79.
3476 peruvianum L.	Peruvian	JU J or 2 my.s	Li Peru ~ 1757. C r.m "Bot. mag. 141
2 hybndum	More's hybrid	tt 1_J or 3 my.s	Li hybrid 1815. C r.m
3477 corymbosum R S(P).	corymbose	SU i_J or 4 my.s	Li Peru 1808. C r.m Bot. mag. 1609
grandiflorum Sehrank			
3478 linifolium Lch.	Flax-leaved	n-L-Jor	lftjn.8 W C. G.H. 1815. C s.p
Myosotis Irutieaa TV.			
3479 brvifolium Wai.	short-leaved	Is D or	■ ■ ■ W.Y Nepal 1824. C pi
3480 zeylamicum Lam.	Ceylon	« O or	Jl.au W Ceylon 1818. S r.m Bur. in. 16.2
S481 humilis Lam.	humble	■ D or	my.jn W S. Amer. 1752. C si PLic.227. 2
3482 maroccanum Lch.	Morocco	■ L _J or	jn.ji W Morocco 1823. F sl
3483 undulatum Vahl	crispum Qef.	■ Jl.au	LB N. Africa 1820. F s.p Desfat41
European			
3484 europium L.	European	O or I jn.o	W & Europe 1562. F ■ Jac. au. a 207
S445 commutatum R. 4> S. changed		O or	W S. Europe 1800. F ■
S451 supinum L.	supine	* Q or	W S. Europe 1640. F ■ Gou. m. ■ ■ ■

432. ilNCHU'SA L.	BUGLOSS.	(<i>Agehousta</i> , paint for the skin; use.)	<i>Boraginea</i> . 28.—50.
3560 Barrelieri Gm.	Barretter's	☒ A or 2 my.jn B	S. Europe 1820. D co Bot. mag. 2349
Myosotis obtusa W. \$ K.			
3561 myosotidin Vira Leh.	ikyfiosotis-flwd	☒ A or 1 jn.8 B	Tauria 1825. D co
Jlfyfiosotis macrophylla Bieb.			
3562 naniculata H. K.	panicled	☒ 3 my.jn B	Madeira 1777. C p.l Fl. gr. ltd
3563 Imec'na Gar.	pleasing	4 jn.jl B	S. Europe 1817. S co M. h. 28. 7
3564 asperrima Del.	roughest	14 my.jn B	Egypt 1817. S co
3565 Miller* W. en.	Miller's	1J my.jn Pk	Ilevant 1713. S co
3566 stylosa Bieb.	stylose	1 my.jn B	Tauria 1820. S co Bieb.cen.23
3567 prceera Bet.	tall	3 my.jn B	Gachia 1824. S co
3568 tinctoria L.	dyer's	1*jn.o P	MontpeL 1596. D co Bot. rep. 576
3569 hispida Forsk.	bristly	1* my.jn B	Egypt 1817. S co
3570 (endilla Horn.	delicate	1 f my.jn W	Ceylon 1820. S co
zeylanica Pat.			
3571 officinalis L.	officinal	P	Britain sea co. D co Eng. bot. 662
3572 ochroleuca Bieb.	yellowish-white^	Pa.Y	M.Caucas. 1810. D co Bot mag. 1608
2 italica FT.	Italian	R.p	S. Europe 1597. S co Bot. reg. 483
3573 incarnita Schr.	flesh-colored	T	S. Europe 1816. S co
3574 latifolia XeA.	broad-leaved	Q) or 2 my.jn B	1826. S co
3575 angustifolia L.	narrow-leaved	A or 2 my.jn P	S. Europe 1640. D co Fl. gr. 164
3576 leptophylla K. \$ S.	slender-leaved	A or 2 Jl.o P	Europe 1640. S co Bot. mag. 1897
angustifolia B. Af.			
3577 longifolia IAMI.	long-leaved	2) jn.au B	Italy 1819. S co
3578 /ycopoides Bes.	Ly <i>ll</i> ik k	li jn.au B	Russia 1824. S co
3579 maculata //urn.	spoltuin-leaved	1J my.jn B	1824. S co
3580 undulata L.	waved-leaved	2 jn.au B	Spain 1752. D co Bit mag. 2119
3581 Gmelini Led.	Gmelin's	2 my.jn B	Podolia 1817. S co
3582 caplnsis Thun.	Cape	1 jl B	C. G. H. 1800. S p.l Bot. rep. 336
3583 hifbrida Ten.	hybrid	1) jn.au W.B	Italy 1820. S co Fl. nap. 11
3584 Agardhi Leh.	Agardh's	I jl.s B	Siberia 1820. S co
3585 verrucosa Lam.	warty	i jn.au P.v	S. Europe 1821. S co Jac.vin.3. 21
3586 rupestris R. Br.	rock	1 jl B	Siberia 1802. D co PaLit.3.71. E.3
3587 sericea f. Sf S.	silky	1 jn.au B	Siberia 1802. D co
433. SY'MPHYTUM L.	COMFRET.	(<i>Sumphuo</i> , to make unite;	healing qualities.) <i>Boragineie</i> . 10.—
3588 otticinale L.	officinal	myjl W	Britain " wet pl. D co Eng. bot 817
2 patens Sib.	open	my.jl B	Britain wet pl. D co
3589 bohemicum Schott	Bohemian	my.jl K	Bohemia 1810. D co
3590 tuberosum L.	tuberous	my.o Y	Scotland m s.p.l. D co Eng. bot 1502
3591 orientate L.	eastern	my.jl W	Turkey 17*2 D co Bot. mag. 1912
3592 caucasicum Bieb.	Caucasian	P	Caucasus 1820. H co
3593 tauncum W. en.	Taurian	my.jl W	Tauria 1806. H co Bot. mag. 1787
3594 bullatum Jac.	b <i>ll</i>	2) my.jl W	Caucasus 1818. H co Jac. ec. 81?
3595 cordatum Kit.	heart-leaved	2 my.jl Y	Transylv. 1813. H co W. & K. 7
3596 aspemnum H. K.	roughest	6 my. s R.B	Caucasus 1799. H co Bot. mag. 929
2 hybridum	hybrid	4 my.s R.B	hybrid 1825. H co
3597 peregrinum Led.	foreign	2 my.au	Podolia 1816. H co
434. ONOSMCDIUM Mx.	ONOSMODIUM.	(<i>Onosma</i> , <i>Onosma</i> , <i>eidos</i> , like; similarity.) <i>Boraglnct.</i> 2. - 3.	
3598 hispidum Mx.	bristly	^ A or 1 jn Y	N. Amer. 1759. D s.l. M.h.3.11.^8.3
PrlshuTi hispida Leh.			
3599 m6Ue Mx.	soft	^ i jn.au W	N. Amer. 1812. D 8.1 Mic am. 15
Purshta nullis Leh.			
435. CYNOGLO'SSUM L^ HOUND'S-TONGUE. (At/on, a dog, <i>eloua</i> , a tongue; leaves.) <i>Boraglnete</i> . 19.—			
3600 officinale L.	official	2 jn.jl B.p	<i>Bohemia</i> 1819. S 68 Eng. bot 921
3601 Haenkei Schult	Haenke's	3 ju.jl B.p	Britain aha. la. S co
3602 sylvaticum E. B.	wood	2 jn.au P	France 1820. S co Eng. bot 1642
3603 Dioscoridis Vil.	Dioscorides's	2 au L.B	Madeira 1658. H co
3604 pictum H. K.	painted	2 jn.au W.p	Germany 1820*. S co Bot. mag. 2li>4
3605 bicolor W.	two-colored	2 jn.au F	— 1819. H co
3606 elongitum Horn.	elongated	2) jn.au Br	Spain 1821. H co Des. at. 1.44
3607 clandestinum Desf.	clandestine	2 jn.au B	N. Amer. 1812. D p.l
3608 amplexicaule Mx.	stem-clasping	2 my.jl R	
virgin leum L.			
3609 apenninum L.	Apennine	apil R	Italy 1731. H co Col.ec. 1.70
3610 magellense Ten.	Magellan	my.jl P	Naples 182a H co
3611 australis B. Br.	southern	my.jl P.R	N.Holl. 1820. H co s.p
3612 hirsutum Thun.	hairy	jl.au LB	C. G. H. 1806. H co
3613 canescens IV.	cancscnt	jnau B	E. Indies 1819. H co Jac. 8C. 489 .
hirsutum Jac.			
3614 Coltim? Ten.	Columna's	jn.au P	Apennin. 1825. S co Col. ec. 1.178
3615 holoscrecum Stev.	velvety	jn.au V	Siberia 1821. S co
3616 umbellatum Kit.	umbellated	my.au P	Hungary 1817. S co
3617 glomeratum Fra.	clustered	my.au N. Amer. 1812. S co	
3618 c/wirtohum L.	Wallflower-lvd	Levant 1596. S co	
^nchfisa lanSta L.			
436. RINDR. Pal.	RINDKRA.	{ <i>Kinder</i> , dean of medicine in Moscow.)	<i>Boraglnete</i> . 1. —
3tiV laivluhtn Jt.	Silofitl	A « r 1 Jnjl K	Siberia 1818.
tetrafpis JPal.	fynoglou«um U — M		
437. OMPH ALCDES Leh.	VENUS' NAVEL-WORT. (<i>Omphalos</i> , a navel, <i>eidos</i> , like; seeds)		<i>Boraginea</i> ? 7.—
3620 nitida I*.	shining	£ A or £ apjn W	Portugal 1480. III eo Bot. mag. 2589
Cynoglossum nitidum if. Af.	Picutia nitida B.&S.		
3621 verna Leh.	spring	tf A el j mr.ap B	S. Europe 1633. D co Bot. mag. 7
Cynoglossum Omphalodes L.	Picotia v^rna R.S.F.S.		
3622 semmervirens D.	evergreen	£ A or 2) my.jl B	Britain rub. D co Eng. bot 45
yfnehtsa sempervirens L.			
3023 amplexicaulis Leb.	stem-clasping	Q or 1 jn.au W	Spain 1823. S co Leh. b. 93.6
Picotta Arassicsfoliat \$.	Cynoglossum Arassicsfolum Lag.		
3624 hitoralu Leh.	sand	O pr 1 jn.au W	France 1826. H co
3625 scorpioides Leh.	scorpion-like	* O pr 1 jn.au B	Bohemia 1825. H co Leh. b. 98.7
Cynoglossum scorpioides Joe			
3626 /inifolia Leh.	Flax.l	O or 1 jn.au W	Portugal 1648. S co
Cynoglossum /inifolium L.			

• 446. TOURNEFO'R/tif L.	TOIIRNEFORTIA. (J.	P. de Tournefort, the celeb.	Fr. bot.)	Boragineee. 11.—48.
3731. Musser8chmijliaR.Br.	Messrschmidt'st i J or	6 jn.o G	Canaries	1779. C 61 JJot recg. 464
3732. fruticosa R. Br.	shrubby	4 jnjl W	Caniaries	1800. C pi
3733. angustifolia R.fyS.	narrow-leaved	5 jn.n W.o	Canaries	1820. C p.l
3734. argentea L.	silvery	8 jnn W	E. Indies	IK22 C p. Ru. am. 4.55
3735. foetidissima TV.	most fetid	9 s Pk	Jamaica	1739. F p. Pluk.ic.226.230
3736. cymosa L.	cymose	9 jl Pk	Jamaica	1777. C p. Jac. ic. 1.31
3737. scabra Lam.	scabrous	<> Dor	4 jnjl W.R	W. Indies 1700. C p. p.
3738. bicolor Swz.	two-colored	• • Or	6 Jnjl G	Jamaica 1812. C p. SI. jam. 2.162.4
3739. suttruticosa R. Br.	suffruticose	• I. 1 or	4 jnjl W	Jamaica 1759. C p.
3740/auriflora fun.	Laurel-leaved	• I. Or	6 jnjl Y.v	S. Amer. C p.
3741. mutabilis Ven.	changeable	• I. Jor	4 jn.au W.G	Java C p. Yen. ch. 3
3742. Argertia if Br.	Arguzia	• I. I	i jn.o W	Siberia 1780. C 81
Mehrschmidtf. Arguzia L.				

447. MESSERSCHMID <i>d</i> /i L.	MESSERSCHMIDIA. (D. <i>Messerschmid</i> , a Ger. bot.)	<i>Boraainete</i> .	4.—12.
3748 <i>hirsutissima R.Sf.</i>	<u>hairiest</u>	<i>f</i> O or 10 myjl G.r W. Indies 1818.	C p.l SL jam. 2.212.1
Tournefort <i>M</i>	<i>hirsutissima Swz.</i>		
3744 <i>volutilis R.</i>	<u>s.</u> twining	t. CD or 10 jLau G.T Jamaica 1752.	C p.l SI.jam. 1.143.2
Tournefort <i>M</i>	<i>volutilis L.</i>		
3745 <i>scandens R</i> # S.	climbing	L O or 10 jl.au G.v Peru 1816.	C p.l Fl. per. 2.148
Tournefort <i>R</i>	<i>scandens W. volobilis IT AS.</i>		
3746 <i>Zaun</i> <i>folia R.</i> 8f. S.	Laurel-leaved	<i>fl</i> IZD ^{or} 12 JI Y W. Indies 1819.	C p.l Ven.ch.2
Tournefort <i>Zaunfolia Fen.</i>			

*448. NOLA'NA L. S747 prostrate <i>L.</i> 3748 paradoxia Lindl.	NOLAVA. trailing paradoxical	(<i>Nola</i> , a little bell; corolla.) 8 or 1 j l s B .lk el 1 au B Chile	Soldrwa. 2.—a 1761. C p. Bot. mag. 731 1825. S co. Bot. reg. 865
---	------------------------------------	---	--

•449. ARETM L. ARBTIA. (*B. Aretius*, a Swiss, prof. in univ. of Berne, d. 1574.) *Primuldecar.* 4. — 9. — 3749 helvetic^a L. Swiss £ A P^r i myjn W Switzerl. 1775. D s.p Schk. han. 1. ^ i^ndrsace Aretin Lan.

3750 <i>alpina</i> L.	alpine	£ A pr	\ myjn	Pk	Switzerl.	1775.	D s.p	Bot cab. 297
3751 <i>Vitaliiflora</i> L.	Vital's	A 1»	I " yjn	Y	Pyrenees	17H7.	I) s p	Par. Ion. 107
3752 <i>argentea</i> Gae.	silvery	j£ ^J pr	i myjn	W	Switzerl.	1826.	D s.p	
■ 450. /INDRÖ SACE L.	ANDROSACE.	Miwrr.aman, Jflflow, a buckler; calyx.)	Primulace. 183.—S&					
3753 <i>maxima</i> L.	g —	O	i nw-jn	W	Austria	1797.	S p.l	Tan. AH 4e 331
3754 <i>elongata</i> L.	elongated	O p	i ap.my	W	Austria	1776.	S p.l	Jac. au. 4.330
3755 <i>septentrionalis</i> L.	northern	O P'	i ap.my	W	Russia	1755.	S p.l	Bot mag. 2021
3756 <i>nifna</i> Horn.	dwarf	O P*	I ap.my	W	Denmark	1803.	S pi	nnnaO
3757 <i>lactiflora</i> Fls.	inilk-flowered	5> O>	Jjn.s	W	Siberia	1806.	D sp	Bot mag. 20.»
2Z ⁵⁸ 1, 1, 2, 011168 Horn	Alisma-Lke	O> ?r	i jns	W	Siberia	1820.	S l.p	
3759 <i>filif</i> < Srmis Retx.	filiform	Opr	* myjn	W	Siberia	1820.	S co	Gm.si.4. Ag. 44.*
3760 <i>ChainsBjanw</i> WvL	Bastard-jasmine*	JM pr	ft jn.au	Pk	Austria	1768.	D sp	Bot. cab. 232
3761 <i>villfoa</i> L,	vinous	^ j	ft jn.jl	Pk	Pyrenees	1790.	D s.p	Bot mag. <g
3762 <i>laetca</i> L,	m\ V.-whUe	S ^j pr	ft jn.au	W	Austria	1752.	D sp	Bot. mag. 8W
3763 c&rnea /	...		ft jl au	F	Switzerl	17S	n s.p	Dot cab. 4U

2 odor&tum	scented	At Alor	\ f.ap	H.w	Cyprus	1731.	S	p.l
3 albiflorum	whiteflowered	A LAI or	i f.ap	W	Cyprus	1731.	S	p.l
3821 Acderefulum n. K.	Ivy-leaved'	A A or	\ »P	P	Britain	banks.	S	p.l
europium B.M. & E. B.						Eng. bot	548	
1 purpurascens	purpurascent	A A or	ijn.au	P	Britain	gard.	S	p.l
2 albidum	VrlitisMuui	A A or	\ jn.au	W	Britain	gard.	S	p.l
3822 repandum Sin.	repand	A A or	\ ap.my	R	Greece	1816.	S	pi
3823 neapolitanum Ten.	Neopolitan	A A or	\ ap.my	R	Italy	1824.	S	p.l
3824 lalifolium 8m.	broad-leaved	A A or	\ ap.my	R	S. Europe	1824.	S	p.l
3825 linearifolium Dec.	linear-leaved	A A or	\ ap.my	P	S. Europe	1824.	S s.p	Fl. gr. 18'
456 HOTTCN/i! L.	WATER VIOLET. (<i>P. Hotton</i> , prof, in the univ. of Leyden, d. 1709.) <i>Primylacete</i> . 1.—A							
3826 palustris L.	marsh	* A or 1	jLau	F	England	dit	S aq	Eng bot S64
* 457. LYSIMA'CHIA L.	LOOSE STRIFE. (<i>Lysis</i> , dissolution, <i>mncfie</i> , strife.) <i>Primulacea</i> . 18—29.							
3827 * ulgaris L.	common	A or 3	il s	Y	Britain	watsh.	D co	Eng. bot 761
3828 Ephemerum L.	Ephemerum				Spain	1730.	D pi	Bot.mag. 2346
3829 atropurpurea L.	dark-purple				Levant	1820.	S.C p.l	Hook ex. fl. 180
3830 dñbia H. K.	doubtful				Levant	1759.	S p.l	Mur. 1782.1
3831 stricta H. K.	strict				N. Amer.	1781.	D p.l	Bot. mag. 104
3832 thysiflora L.	thyrs-flowered	A or			England	lakes	D co	Eng. bot 176
3833 capitate Ph.	headed				N. Amer.	1813.	D co	
3834 quadrifolia L.	four-leaved				N. Amer.	1794.	D pi	Lam. il. 1.101 8
3835 longitflia Ph.	long-leaved				N. Amer.	1798.	D p.l	Bot. mag. 660
3836 quadriflora H. K.								
3836 verticillate Pall.	whorled				Crimea	1820.	D co	Bot. mag. 2293
3837 ciliata L.	ciliated				N. Amer.	1732.	D m.s	Walt h. 12
3838 punctata L.	dotted				N. Holl.	1658.	D co	Jac. au. 4.366
3839 angustitolia Mx.	narrow-leaved				N. Amer.	1813.	D p.l	
3840 hybrida Mx.	hybrid				N. Amer.	1820.	D co	
3841 Lfnum stellatum L.	Flax-star				Italy	1658.	S s.l	Mag. mo. 162
Aster6linon stellatum Lk.								
3842 nemorum L.	wood	k A or	imy jl	Y	Britain	m.s.pl.	D m.s	Eng. bot 527
3843 maculata R. Br.	spotted	\ iAI or		Y	N. Holl.	1825.	D p.l	
3844 Nummulana L.	Moneywort	IU \ A or		Y	Britain	m.me.	D m.s	Eng. bot 528
*458. JNAGA'LLISL.	PIMPEINEL. (<i>Anagelao</i> , to laugh; removing despondency.) <i>Pritnuidcea</i> . 9.—14.							
3845 arvensis L.	PIMPEINEL. (Anagelao, to laugh; removing despondency.) <i>Pritnuidcea</i> . 9.—14.							
phecnicea Lam.								
3846 carnea Schl.	fleshy	O or	i jn.s	F	Switzerl.	1819.	S co	
3847 caerulea Schreb.	blue	-* O w	ijn.s	B	Britain	corn fi.	S co	Eng. bot. 1823
3848 indica Suit.	Indian	O pr	1 my.s	B	Nepal	1824	S co	Sw. fl. gar. 132
3849 fmitcosa Fen.	shrubby	jg OI or	3 my.jl	Ve	Morocco	1803.	C p.l	Bot mag. 831
collina Schous.								
3850 Iatilolia L.	broad-leaved	\ iQ or	1 my.jl	P	Spain	1759.	C p.l	Mee. ic. 1.22
3851 Morfelli L.	Monelh's	\ LAI or	1 my.s	B	Italy	1648.	C pi	Bot. mag. 319
3852 /inif6ha L.	Flax-leaved	\ OI or	1 jn jl	B	Portugal	1796.	C s.l	Bot. mag. 2389
3853 tenglla L.	delicate	WK A or	i au.s	Pk	Britain	bog.pl.	D l.p	Eng. bot 530
459. .DIAPE'NSIA £.	DIAPENBIA. (<i>Diapente</i> , composed of five; flowers 5-cleft) <i>Hydrate</i> *. 1.							
3854 lapponica L.	Lapland	\ A or	J f.mr	W	Lapland	1801.	D s.l	Bot. mag. 1108
460. PYXIDANTHE'RA Mx. PYXIDANTHERA. (<i>Pyxis</i> , a box, <i>anthera</i> , an anther.) <i>Hydrdea</i> . 1.								
3855 barbulata Mx.	small-bearded	JU_Alor	,jl	W	Carolina	1806.	D Lp	Mic.am.17
Diapensia cuneifolia Ph.								
461. CO'RIS L.	Corns.	(A Greek name used by Dioscorides.) <i>Primulacea</i> . 1.						
3856 monspheensis L.	Montpelier	jf O or	\ jn.jl	Li	S. Europe	1640.	S s.p	Bot. mag. 2131
462. GA'LAX L.	GALAX.	(Gala, milk; whiteness of the flowers.) <i>Saxifraga</i> . 1.						
3857 aphylla L.	leafless	^ A or	i jn.jl	W	N. Amer.	178a	D e.p	Bot. mag. 754
Blandfurda cordata B. Rep.								
463. 3/ENYA'NTHES L.	BUCK BEAN. (<i>Men</i> , month, <i>anthos</i> , flower; time in blossom.) <i>Genithaea</i> . 1. — 2.							
3858 trifoliolate L.	three-leaved	^ A or	1 jl	W	Britain	moi.pl.	C co	Eng. bot 495
3359 americana G. Don	American	\ A or	1 jl	W	N. Amer.	1818.	D co	
trifoliolate Mx.								
*464. VJLLA'SR/4 Ven.	VILLARIA. (VULars, a distinguished French botanist)							
3860 nymphoides Ven.	Water-lily-like ^ A or	fit jn.jl	Y	England rivers.	Skp.1	Eng. bot. 217		
A/cnyanthes nymphoides L.								
3861 cordata EU.	cordate		or fit jn.jl	Y	N. Amer.	1820.	O p.l	
3862 indica Ven.	Indian		or fit my.au	W	C.G.H.	1792.	Skp.1	Bot mag. 658
ii/cnyanthes indica L.								
3863 ovata Ven.	ovate-leaved	^ iAI or	fit my.jl	O	C. G. H.	1786.	Sk p.l	Bot mag. 1909
3864 Iacunosa Ven.	pitted	A iAJ or	fit jn.jl	W	N. Amer.	1812.	Skp.1	Ven.ch.9
3865 *armentosa B. M.	twiggy	JII iAI or	1 jn.jl	Y	N. Holl.	1806.	Sk bog	Bot. mag. 1323
3866 /mrnassifolia Jl. Br.	Parnassia-lvd	\ iAI or	8>jn.o	Y	N. S. W.	1825.	Sk bog	Bot. mag. 1029
3867 remfdrmis It. Br.	kidney-form	\ iAI or	1 jl.au	Y	N. Holl.	1824.	Sk bog	
• 465. CHIRO'N/if L.								
3868^asminoides Thun.								
3869 /ychnoides Thun.								
3870 nudicaulis L.								
3871 /inoides L.								
3872 tetragona L.								
3873 baccifera L.								
3874 angystifolia H. K.								
3875 frutescens L.								
2 albiflora								
3876 decussata Ven.								
^Sfi.-E^STOMA Sal.	ELJSTOMA..	(<i>Eustoma</i> , beautiful mouth; flower)						
3877 silemiolum 5a/.	Sdene-leaved							
3878 Canteffium Pp. ^{RRH.}	ERYTHILKA.	(<i>Erythraia</i> , red: flowers.) <i>Gentianete</i> . 10.—30.						
Chir6nw Cnteürum E. B.	O or i JLau	Pk	Britain	heaths.	S s.1	Eng. bot 417		

M73. NEMO'PHILA Bart. *Nemophila*. {*Nemos*, a grove, *phileo*, to love.) *Hydrophyllacee.* 2.
 3937 phaceoides Bart. Phacelia-like ij or 1 jl.au B N. Amer. 1822. S co Bot mag. 2373
 3938 pamculata Spr. panicled & \$ my.jn P.B N. Amer. 1813. D p.l
Hydrophyllum appendiculatum Mx.

479. RAMON'Dil Mx. RAMONDA. (Af. *L. Ramond*, a French botanist.) *Soldneic.* 1.
 3939 pyrenaica Mx. Pyrenean J* A or 1 my p Pyrenees 1731. D 81 Bot mag. 23ft
/xrbascum Myc6ni L. Chaixa Mycon L.

480. FERBA'SCUM L.		MULLEIN.		(Altered from <i>barbascum</i> , bearded.)		<i>Soldnete.</i> 67.-74.		
3940 Tiliapsus L.	Shepherd's-club	6	jl.au	Y	Britain	ro. sid.	S co Eng. bot 549	
3941 phapifonnc Schr.	thick-leaved	2	jl.au	L.Y	Europe	1817. S	co	
3942 crassiiflum Haffg.	pointed	3	jl.au	Y	Spain	1818. S	co Fl. por. 1.26	
3943 cuspidatum Schr.	woolly	4	my.jn	Y	Vienna	1817. S	co Sch. mo. 1.1	
3944 gossypiflum Bieb.	snow-white	4	jl.au	Y	Caucasus	1820. S	co	
3945 niveum Ten.	whitest	3	jl.au	P.v	Naples	1823. S	co Fl. nap. 22	
3946 candollissimum Dec.	Thapsus-like	4	jl.au	Y	Montpel.	1823. S	co	
3947 /hapsoides Htfg.	long-flowered	3	jl.au	Y	Spain	1819. S	co Sch. mo. 5.2	
3948 macr.inthum Lk.	southern	3	jn.jl	Y	Portugal	1820. S	co 11. por. 27	
3949 australis Schr.	Phlomus-like	4	jl.au	Y	S. Europe	1815. S	co Sch. mo. 2	
3950 jriilomoiries L.	close-flowered	3	jn.jl	Y	Italy	1739. S	co H. has. 170.4	
3951 condensatum Sc.ir.	grove	4	jl.au	Y	Austria	1820. S	co Sc. verb. 3	
3952 nemorosum Schr.	mountain	5	jl.au	Y	Austria	1820. S	co Sc. verb. 1.2	
3953 montanuin Schr.	hill	3	jl.au	Y	France	1819. S	co H. got. 2.12	
3954 collinuin Sc.*r.	inverted-flwd	3	jl.au	Y	Germany	1820. S	co Sc. verb. 5.1	
3955 versiflum Schr.	branch-bearing	6	jl.au	Y	S. Europe	1824. S	co	
3956 ramigerum /Jr.	Bastard's	6	jl.au	Y	France	1824. S	co	
3957 Bastardii R. & S.	mucronate	4	jl.au	Y	Greece	1814. S	co	
3958 mucronatum Lam.	waved	2	jl.au	Y	S. Europe	1570. S	co Fl.gr. 227	
3959 sinuatum L.	pinnatifid	1	my	Y	Archipcl.	1788. S	p.1	
3960 pinnatifidum TA/I/	waved	1	jl.au	Y	S. Europe	1819. S	co Fl. gr. 226	
3961 undulatum Lam.		1	jl.au	Y	Tauria	1813. S	p 1 Bot mag. 1777	
3962 bipinnatifidum B. M.	bipinnatifid	1	jl.au	Y	S. France	1821. S	Vil. del. 2.13	
3963 Chaix's	Hornemann's	1	jl.au	Y	Europe	1821. S		
3964 Bocraavij L.	long-stalked	1	jl.au	Y.p	S. Europe	1731. S	Mil. ic. 2.273	
3966 elongatum IV. en.	crowded	1	jl.au	Y	Europe	1813. S	co	
3967 com>actin lieb.	3	jl.au	Y	Tauria	1820. S	Til. pis. 50?		
3968 /maphalodes Bleb.	Gnaphaliu-ik	2	jl.au	Y	Caucasus	1825. S	co	
3969 ZychnitisZ.,	Lychnitis	3	jn.au	C	Britain	ro. sid.	S co Eng. bot 58	
3970 austriacum Schott	Austrian	5	jl.au	Y	Austria	1818. S	co	
3971 pulverulentum E. B.	powdered	3	jn.au	Y	England	bor. fi.	S co Eng. bot. 487	
3972 lanatum Schr.	woolly	3	jl.au	Y	Italy	1825. S	co	
3973 Stevenii Fis.	Steven's	5	jl.au	Y	Siberia	1821. S	co	
3974 inale Dec	hog	5	jl.au	Y	Montpel.	1817. S	co	
3975 pyramidatum Bieb.	pyramidal	3	jl.au	Y	Caucasus	1804. S	co Sw. fl. gar. 31	
3976 ovatum Schr.	ovate	3	jl.au	Y	Europe	182*. S	co H. got 15	
3977 rotundifolium Ten.	round-leaved	4	jl.au	Y	Italy	1823. S	co Fl. nap. 23	
3978 floccosum Kit.	woolly	3	jn.jl	Y	Hungary	1805. S	co W. & K. 59	
3979 leptostachyum Dec.	slender-spiked	3	jl.au	Y	S. France	1825. S	co	
3980 angustifolium Ten.	narrow-leaved	3	jl.au	Y	Naples	1824. S	co	
3981 formosum Bieb.	handsome	2	jl.au	Y	Russia	1818. S	co Bot. reg. 558	
3982 spectabile Bub.	showy	2	jn.jl	Y.P	Tauria	1820. S	co	
3983 longifolium Ten.	long-leaved	3	jl.au	Y	France	1824. S	co	
3984 nigrum L.	black, rooted	■ my.au	Y	Naples	1824. S	co Fl. nap. 21		
3985 orientale Bieb.	eastern	■ my.au	Y	England	ch. so.	S co p.1 Eng. bot 59		
3986 iflopecfirus Thuil.	Foxtail	■ my.au	Y	Caucasus	1821. S	co		
3987 ramosissimum Poir.	braiu-hiest	■ my.au	Y	France	1820. S	co		
3988 hasmorphoidal U.K.	haemorrhoidal	2	jn.au	W.P	Madeira	1777. S	co	
3989 speciosum Schr.	comely	1	jl.au	Y	Austria	1818. S	co H. got 16	
3990 ovalifolium H. K.	oval-leaved	1	jl.au	Y	Caucasus	1804. S	co Bot mag. 1037	
3991 grandiflorum Schr.	great-flowered	1	jl.au	Y	Europe	1820. S	co H. got. 15 •	
3992 lattana L.	Moth	1	jl.au	Y	Britain	gra. pi.	S co Eng. bot 393	
2£iba	white	1	jl.au	W	Britain	gard.	S co	
3993 Alattariofdes Lam.	Blattaria-like	1	CD or	Y	France	1819. S	co	
3994 lyratum Lam.	lyrate	1	O or	jl.au	Y	Spain?	1819. S	co
3995 repandum W.en.	repand	1	O or	jl.au	Y	Europe	1813. S	co
3996 Osbeckii W.	Osbeck'a	1	Q or	jl.au	G	Spain	1752. S	To. it 2.83
3997 glabrum W.en.	smooth	3	jl.au	Y	Europe	1805. S	p.1	
3998 virgatum With.	twiggy	1	Q or	jl.au	Y	Britain	gra. pi.	Eng. bot. 550
3999 cyprium B. At.	copper-co/orai	5	jl.au	Y	Caucasus	1798. S	co Bot mag. 1226	
4000 pheonicium L.	Vur'ic-Jtowered\$	1	A or	my.au	P	P. S. Europe	1716. S	Bot. mag. 885
4001 punicum Schr.	yellow	1	A or	my.au	P	S. Europe	1820. S	H. got. 14
4002 rubiginosum W.&K.	rusty	1	jl.au	Y.B	Hungary	1817. S	p.1 W.&K. 197	
4003 ferrugineum H. A.	ru^t-colored	1	jl.au	Y.B	S. Europe	16PI. S	Bot. rep. 162	
4004 triste Lk.	dark	2	jl.au	Y.R	S. Europe	1788. S	And. rep. 162	
ferrugineum And. rep.	not of others.							
4005 totomiflum Dcsf.	Betony-leaved	2	jn.au	Y	Armenia	1825.	An. mu. 11.4	
4006 spinobum L.	thorny	1	my.jn	P	Crete	1824.	Alp. ex. 36	

481. DATU'R'A L.		THORN APPLE.		(Alteration of the Arabic name, <i>tdtSrah.</i>)		<i>Solunea.</i> 9.—10.	
4007 trox L.	fierce	O or	3	jl.s	W	China	1731. S
4008 Stramonium L.	Stramonium	O clt	3	jl.s	W	England	rub. S sl Eng. bot 1288
4(49) muncata Lk.	rouncated	O or	2	jl.s	W	■ 1820. S	r.m
4010 Tdula L.	Tatula	O or	3	jl.s	B	N. Amer. 1&9.	S sl Mee.ic.2.13
4011 faMu6sal.	proud	O or	3	jl.s	P	Egypt	1629. S
4012 Mite/L	Metel	O or	2	jn.s	W	Asia	1596. S
4013 guayaquilnsis Kth.	Guayaquil	O or	2	jl.s	'W	Guayaquil	1826. S
4014 laevis L.	smooth-fruited	O or	2	jn.s	W	Africa	1780. S
4015 ceratocalon Or.	horn-stalked	O or	2	jl.s	W	S. Amer. 1805. S	r.m Jac. vin. 3.82
482. BRUGMA'NS/if Pers.	BRUGMANSIA.	(Prof. S. J. Brugmant, Ger. bot author.)					Jac. sc. 3.339
4016 suaveolent IV. en.	sweet-scented	1 • or	IS	au.s	AV	Peru	1733. C
Datura arborea Hort.							I.p
4017 Candida Pers.	wh'te, stalked	J • or	10	au.s	W	Peru	1813. C
Dattta arborea W. en.							I.p FL per. 3 128

- | | | | | |
|---|---|--|-------------------|-----------------------------------|
| 483. LISIA'NTHUSJSr. | LISIANTHIJS. | (<i>Lis</i> , smooth, <i>anthot</i> , a flower.) | Gentianete. | 10.—31. |
| 4018 Iongifolus L. | long-leaved | ill- lor | y | Jamaica 1793. |
| 4019 glaucifolius Jac. | glaucous-leaved* | • or | p | Br. jam. 9.1 |
| 4020 exsArtiu Swz. | drawn out | ■ | Y | Jac. ic. 1.33 |
| 4421 latifolius Swz. | broad-leaved | or | 4 jl.s | W. Tndies 1793. |
| 4022 umbellatus Swz. | umbelled | • or | 6 jl.au | Jamaica 1821. |
| 4023 cordifolius L. | heart-leaved | • or | 2 | Jamaica 1822. |
| 4024 acutangulus R. S. P. | acute-angled | El or | jlau | Jamaica 1816. |
| 4025 grandiflorus Aubl. | great-flowered | ly | jl.au | Peru 1820. |
| 40.6 angustifolius Kth. | narrow-leaved | a Q or | jl.au | Trinidad 1818. |
| 4>27 alatus Aubl. | winged | • | jl.au | S. Amer. 1824. |
| | | | | Mexico 1884. |
| 484. SPIGE'L U L. | WORM GRASS. | (<i>A. Spigelia</i> , prof. at Padua, d. 1625.) | Gentiane'a. | 2.—5. |
| 4028 Anthelmia /,, | Anthelmia | Ow 14 | G.B | W. Indies 1759. |
| 4029 maniändica L. | Maryland | ^ A or | jl.au | S. Amer. 1694. |
| 485. NIC'A'NDIU J. | NICANURA. | { <i>Nicander</i> , a Greek phys. flourished about | 150 B. C.) | <i>Soldne</i> *. 2. |
| 40.30 physaloides Pers. | Physalis-like | O or 2 jLs | P | <i>Peru</i> 1759. |
| A'tropo physaloides L. | | | | D s.l. Bot. mag. 2458 |
| 4031 Indica R.%S. | Indian | O or | 2 jLi | B R Indies 1824, |
| Phjrsals indica Lam. | | | | S co Ru. am. 6.26.1 |
| 486. ANI'SODUB Lk. | ANISODUS. | { <i>Anisos</i> , unequal, <i>odous</i> , a tooth; calyx.) | Solunee. | 2. |
| *32 stramonifolius (r. Don & tramonium-lvd^ | A or | 5 au.s | Y.G | Nepal 182a |
| Whitleya str. tinoiutola Svo. | | | | R co Sw. fl. gar. 125 |
| 4033 Ifridus Lk. | lurid | ^ ^ o r | 3 au.o | Lu Nepal 1824. |
| Nicajdra anomala Lk. | | | | R co |
| *487. J/YOSCY'AMUS L | HENBANE. | (<i>Hys</i> , a hog, <i>hyamot</i> , a bean; eaten by swine.) | Soli'mete. | 13.— |
| 4034 niger L. | black | Q O | 3 jnjl | Y.p Britain rub. |
| 2 finnuus | annual | ■ | 1 jnjl | S. s.l Eng bot. 591 |
| 4015 albus L. | white | ■ | 2 jl.au | S. s.l Hot mag. 2S94 |
| 4036 reticulatus L. | netted | ■ | 1 jl.au | S. co Black. III |
| 4037 Senecio'nis IV. en. | Groundsel | ■ | lin ro | S. co Com. h. 77. 22 |
| 4038 atiros L. | golden | ■ | mro | C. s.l |
| 4<39 can.tnchensis Ker | Canary | ■ | H year | S. r.m Bot. mag. 87 |
| 4'40 pusillus L | small | * | Y | C. s.l Bot reg 180 |
| + 41 physaloides L. | Physalis-like | ■ | I | S. s.l Pluk. al. 37.5 |
| 4<45 agrestis Kit. | field | ■ | 1 mr.ap | D p 1 Bot. mag. &2 |
| 4<>13pa'lidus Kit. | pale | ■ | 1 ap.iny | S. pl. Sw. fl.gar. 27 |
| 4<>44 muticus /,, | awnless | ■ | 1 ap.iny | Egypt 1822. |
| *45 orientalis Bleb. | eastern | ■ | 1 mr.ap | Iberia 1821. |
| 4H6 auriculatus Ten. | eared | II ap | P.Y | D Bot mag. 2414 |
| | | | | Italy 182J. S |
| *488. VICOTIA'NA L. | TOBACCO. | (<i>J. Nicot</i> , of Nismes, procured seeds from a Dutchm.) | Soldnect. | 24.—27. |
| 4047 fhticōM L. | shrubby | tt i or | 4 jl.au | Pk China 1699. |
| fW* chinensis Fis. | Chinese | O or | 6 jls | Pk China 1819. |
| *49 macrophyllid W.en. | long-leaved | O or | 6 jl.au | Pk America ... |
| 40.80 Tabucum L. | Tabacum | O clt | 4 jlau | Pk America 1570. |
| 1 attenuftum R. S(S. | attenuated | O clt | 4 jlau | S. Amer. 1570. |
| 2 macropolyllum 7{#& long-leaved | O clt | 7 jlau | Pk S. Amer. 1570. | |
| 3 pallescens A * S. | pallescent | O clt | 4 jlau | S. Amer. 1570. |
| 4 alines R. # V. | wing-stalked | O clt | 4 jlau | S. Amer. 1570. |
| 5 seritnum R. S/S. | late | O clt | 4 jlau | S. Amer. 1570. |
| 6 gracilis R. \$ S. | slender-stalked | O clt | 4 jlau | S. Amer. 1570. |
| 7 V6rd.inij SS. | Verdan | O clt | 4 jlau | S. Amer. 1570. |
| 3P* S.P*111 R. k & | tongue-leaved | O clt | 4 jlau | S. Amer. 1570. |
| 4051 dilatata Lk. | dilated | O or | 3 jl | Pk IK20. S. r.m |
| 4GJ2 ybarfensis It \$ B. | Ybarra | O or | 2 jls | Pk S. Amer. 1823. |
| 4054 angustifolia | narrow-leaved | O or | 4 jls | Pk Chile 1819. |
| *15 viscasa Uh. | Buenos Ayres | O or | 3 jl.s | Pk B. Ayres 1821. |
| 4056 pusilla L. | clammy | O or | 3 jl.au | Pk B. Ayres 1824. |
| 4007 glutinosa L. | small | jy III or | 3 au | S. r.m Mil. ic. 2 185.2 |
| 4008 rfistica L. | clammy | O or | 4 jl.s | S. Peru 1759. |
| 4059 hñmialis Desf. | rustic | O clt | 3 jls | S. r.ra Bot rep. 484 |
| <<ri0 paniculata L. | humble | O or | 1 il.s | S. co Black. 437 |
| 4062 Langsdorffii Nees | paniced | O or | 3 ■ | S. r.m |
| 4<<3 repanda W. | Honeywort-like | O or | 2 au | S. r.m Fl. per. 2.129 |
| 4if4 if dictreni Ag. | repanad | O or | 5 au | S. co Leh. nic. 2 |
| *fi5 plumbaginifl. halif.cn. | decurrent | O or | 2 jnjl | S. co Bot. mag. 222 |
| undulata R. Br. | Plumbago-lvd | O or | 2 jl.s | S. co Bot. mag. 2484 |
| *W undulata R. Br. | wav#_Txf | I Alelt | 2 my.Jn | America 181(5. |
| | | | | D r.m Jac ff. 84 |
| *7 suaveolens Uh. | Vinca-flowred | ^ \ or | 2 jl.s | N. S. W. 1800. |
| *S nevroleura Las. | four-valved | O or | 1 jl.au | D r.m Bot. mag. 673 |
| *JS multivalvis B. R. | many-valved | O or | 2 jl au | S. r.m Bot reg 1057 |
| 4070 nina B R. | dwarf | O cu | jn | co Bot. reg. 883* |
| *Pf TWNJA J. | PETUNIA. | { <i>Petun</i> , the name for tobacco in Brazil.) | Solare*. 1.—2. | |
| *1/1 Pvctaginiflora J. | Mary.-Ptru-flwd | ^ or | 4 jn.s | C co Sw. fl.gar. 119 |
| bicotida nyctaginiflora Leh. | | | | |
| 4070. S S. 9. P O / L M Jac. | STM***. (G. A. Scopoli, celeb, prof. of botany d | 1789.) | SoMne*. 1. | |
| *X carniflca Jbf. | Carniolian. | .U or | 1 ap.my | D e o Bot mag. 1126 |
| * H. & S. jyyoscyamus ScopoUa L. | | | | |
| | IPOMGSA. (Ips, bindweed (error of Lin.), <i>homoios</i> , similar.) | | Convolvldceee. | 111.—193. |
| QuamocMVf. | Quamodit | i O o r | 6 jls | S. r.m Bot.mag. 244 |
| 2 flore albo | white-flowered | jj O or | 6 jl. | E. Indies Ifi29. S. r.m |
| 4014 dissecta IV. | cut-leaved | £ El or | 10 jns | W R Indies IR> S. r.m |
| 2105 quinqueloba Vahl | five-lobed | El un | 8 jls | Tropics 1813. C <P W.ph. 1.2.3 |
| J<<6 Kinuata Or. | tcoUofit-leaved | f. iAl or | 6 jl.au | W. indies 1810. C co |
| | various-leaved | & El un | 10 jl.au | Cuba 1817. S co Jac. vin. 159 |
| | digitated | JL El or | 10 aus.s | W. Indies ... s r.m Jac. fr. 42.4 |
| | tubercled | J. • or | 6 jl au | Bourbon 1818. C pl. PI. ic. 92.1 |
| | | | | 1. n. Jac sc. 2. 199 |

4060 <i>Carolina L.</i>	Carolina	\$ O or	10	Jlau	P	Carolina	rm	Di.el.84,98
4081 <i>venrta Val.</i>	veiny	ft_IZ2 or	10	Jlau	P	Mauritius	1820.	C I P
4082 <i>tuberosa L.</i>	tuberous-roofed	A I23 or	10	...	Pa.Y	W. Indies	1731.	C s.p Bot. reg. 768
4083 <i>paniculiflta L.</i>	panicld	A I23 m	20	jns	Pk	E. Indies	1799.	C B.p Bot reg. 62
4064 <i>enosperma Beauv.</i>	woolly-seeded	A IS! or	10	Jlau	Pk	Guinea	1823.	S r.m Beau. ow. 2
4085 <i>senegalensis Lam.</i>	Senegal	A (2)2gr	10	Jlau	Pa.P	Guinea	1823.	S r.m
4086 <i>marrorrhiza L.</i>	long-rooted	A 23 or	10	Jlau	Pk	Guiana	1824.	S r.m Jac. am. 90.1
4087 <i>splendens G. Don</i>	shining	A (SI or	18	jn.jl	Pk	Guinea	1823.	S r.m
4088 <i>pentaphylla Jac.</i>	five-leaved	iOor	20	au.i	W	W. Indies	1739.	S s.p Jac. ic 2.319'
4089 <i>pilosa Crv.</i>	Cavanillea's	ft_CDor	10	aus.s	■ ■	Mexico	1815.	C p.l Cav. ic. 3.256
4089 <i>Cavanillesii R. & S.</i>	armed	■ ■ ■	6	jls.	■	Mexico	1824.	S p.l Cav. ic. 5.478.«
4090 <i>armata71.\$P.</i>								
4091 <i>muncata Cav. not Jac.</i>	thick-seeded	J O or	10	aus.s	P	E. Indies	1815.	C Lp Bot.reg.86
4092 <i>tuberculata B. R.</i>	pendulous	ft_t_	10	my.o	Pk	N. S.W.	1808.	R l.p Bot. rep. 613
4091 <i>ternata Jac.</i>	ternatleaved	tDun	3	Jlau	W	Brazil	1824.	S r.m Jac. 6C.37
4094 <i>parviflra Vahl</i>	small-flowered	JiQJun	4	Jlau	P	W. Indies	1818.	S r.m SI. jam. 97.1
4095 <i>pes-tigris L.</i>	tiger's-isot	£ O or	6	au	R	E. Indies	1732.	C s.p I.i. cl. 318 411
4096 <i>cafni Swt.</i>	Cairo	ft_CD or	8	jn.au	R	Egypt	1680.	C r.m Bot. mag. 699
	<i>Convolvulus cairicus B.M.</i>	<i>Ipomoc'a palmate Forsk.</i>						
4097 <i>platcnis Ker</i>	Plata	ft_E3or	10	jns	V	S. Amer.	1817.	S r.m Bot reg. 333
4098 <i>chryseides Ker</i>	golden	ft (23 or 4	jns	Y	China	1817.	S r.m Bot reg. 270	
4099 <i>caerulea Ker</i>	blue	1 O or	■ ■	L.B	E. Indies	1818.	S r.m Bot. reg. 276	
4100 <i>scotsa Ker</i>	bristly	■ ■ ■	■ ■ ■	P	Brazil	1817.	S r.m Bot reg. 335	
4101 <i>scabra Gm.</i>	rough	■ ■ ■	■ ■ ■	W	S. Amer.	1804.	S r.m	
4102 <i>Tnrpithum L.</i>	Turbcth	■ ■ ■	■ ■ ■	W	Ceylon	1759.	S r.m Bot. mag. 2093	
4103 <i>lutæola Jac.</i>	yellowish	■ ■ ■	■ ■ ■	S	Carolina	1759.	S r.m Bot. mag. 221	
4104 <i>coucinia L.</i>	scarlet	■ ■ ■	■ ■ ■	D.R	W.Indies	1713.	S r.m Bot. rep. 99	
4105 <i>Iacun6sa L.</i>	pitted	■ ■ ■	■ ■ ■	W	N. Amer.	1640.	S r.m Di. cl. 87. 102	
4106 <i>Gossypn6ha W.</i>	Gossypium-lvd	ft_CD or	15	■ ■ ■	P	—	—	C s.p Bot reg. 75
insignia B. R.								
4107 <i>bhia-nox L.</i>	good-night	■ ■ ■	■ ■ ■	■ ■ ■	■ ■ ■	W.Indies	1773.	S s.l Bot. mag. 752
4108 <i>noctiluca Herb.</i>	night-shiniug	■ ■ ■	■ ■ ■	■ ■ ■	■ ■ ■	E. Indies	1820.	S co Bot. reg. 889
lutih6ra B. R.								
4109 <i>sanguinca Vahl</i>	Wood-flowered	■ ■ ■	■ ■ ■	fn	D.R	W. Indies	1812.	C 81 Bot reg. 9
4110 <i>mutauilis B.R.</i>	changeable	■ ■ ■	■ ■ ■	my.s	P	S. Amer.	1812.	C p.l Bot. reg. 39
4111 <i>candicans B. M.</i>	whitish	* IAI or	15	jn.au	W	N. Amer.	1776.	R p.l Bot mag. 1603
4112 <i>cwrulcscens Rox.</i>	cacrulescnt	ft_ (23 or 10	jn.au	B	E. Indies	1820.	■ ■ ■	
4113 <i>tutir'cha Swt.</i>	Vine-leaved	ft_ • or	10	jn.au	Y	E. Indies	1817.	■ p.l Bur. in. 18.1
	<i>Convolvulus vitifolius L.</i>							
4114 <i>Jalnpa B. M.</i>	Jalap	■ ■ ■	10	aus.s	R	America	173a	C r.m Bot reg. 621
	<i>Jaldpa fi rascal? A</i>	<i>Conv6lvulus Jaldpa B.M. 1572</i>						
4115 <i>Micnauxi? Swt</i>	Michaux's	A CD m	10	aus.s	W	Georgia	1815.	C p.l Bot reg. 342
	<i>Jaldpa a. B.R.</i>	<i>macrorhiza Mx.notL.</i>						
4116 <i>henaticiflia L.</i>	Hepatica-lvd	JO or	10	aus.s	P	E. Indies	1759.	S co Bur. in. 50. 20. S
4117 <i>sagittate Pair.</i>	sagittate	i A or	6	Jlau	P	S. Europe	1821.	D co Cav. ic. 2.107
4118 <i>plataniflia Vahl</i>	Plane-leaved	ft_ 123 or	8j1.ju	W	W. Indies	1820.	I co	
4119 <i>barbate Roth</i>	bearded	J O or	10	Jlau	P	N. Amer.	1824.	S co Di.eL80.92
4120 <i>v&ria/fo/A</i>	various	fc.CEjun	10	Jlau	P	America	1810.	S r.m
4121 <i>acuminata Vahl</i>	acuminate	ft_ D3 un	6	Jlau	P	W. Indies	1818.	S rin
4122 <i>angula'ris W.</i>	angular	f. CD or	■ ■	Jlau	S	E Indies	1823.	■ Pi
4123 <i>communita R. Sf S.</i>	changed	\$_ O un	■ ■	Jlau	P	N. Amer.	J8IU	S r.m Cat car. 87
4124 <i>triquetra Vahl</i>	triangular	ft_ CD or	■ ■	Jlau	P	W. indies	1824.	S rm
4125 <i>longiflora H.%B.</i>	long-flowered	ftEJor	6	Jlau	P	S. Amer.	1824.	S rm
4126 <i>Battdas Poir.</i>	Batatas	A IS] clt	6	—	P	Tropics	1&7.	R p.l R.mal.7.50
	<i>Conv6lvulus Battdas L.</i>							
4127 <i>punctate if. £ S.</i>	spottcti-calyxed	JL ■ or	6	jls.	P	■ ■ ■	■ ■ ■	■ ■ ■
4128 <i>fattigiat Swt.</i>	fastigiat	* ■ or	10	jn.jl	P	E. Indies	1816.	C Lp
	<i>Conv6lvulus fastigatus Rox.</i>							
4129 <i>bicolor Swt.</i>	two-colored	ft_ (23 or	6	aus.s	Y.p	E. Indies	1816.	S p.l
	<i>Convolvulus bicolor Rox.</i>							
4130 <i>pandurate B. R.</i>	fiddle-shaped	\$_ A or	12	jls.	W.p	N. Amer.	1732.	D p.l Bot. mag. 1939
	<i>Conv6lvulus panduratus B. M.</i>							
4131 <i><olaniflia L.</i>	Nightshade-lvd	J O or	8	Jlau	Pk	America	1759.	C s.p PL ic. 94.1
4132 <i>campanulata L.</i>	be\Jtuwered	j j O or	8	aus	P.w	E. Indies	1800.	S s.I R.mal. 1156
4133 <i>multiflra Mil.</i>	many-flowered	J; OJ or	6	jn.jl	P	Jamaica	—	S Lp Pluk. ph. 167.1
4134 <i>malabanca L.</i>	Malabir	ft_ O or	10	—	P	E. Indies	1823.	C Lp K. mal. 2.51
4135 <i>violacea L.</i>	■ ■ ■	■ ■ ■	8	aug	P	S. Amer.	1732.	S s.I PL ic. 93.1
4136 <i>carpa L.</i>	fleshy	J E Q or	10	aus.s	F	S. Amer.	1799.	S s.I Jac. am 26. 18
4137 <i>repanda L.</i>	repand	jOor	10	au	Sc	W.Indies	1793.	C s.p Par. Ion 81
4138 <i>sibinia L.</i>	Siberian	£ O or	8	Jlau	F	Siberia	1779.	S co Pal. it. 3.723. K
4139 <i>speciosa L.fil.</i>	showy	ft_ I"or	8	Jlau	P	E. Indies	1778.	S p.l Bot. mag. 2446
4140 <i>purpurea Lam.</i>	great puq>le	J O or	—	n.s	D.P	America	1629.	S co Bot. mag. 113
*	2 incarnata	flesh-colored	J O or	—	F	America	1629.	S co
*	3 v&ria	various	J O or	n.s	St	America	1629.	S co Bot. mag. 1682
4	4 leiu&nttha	white-flowered	£ Q or	n.s	W	America	1629.	S r.m
4141 <i>discolor Jac.</i>	sundry-colored	j; O or	20	n.s	B.w	—	—	S co Bot mag. 1005
4142 <i>triloba L.</i>	three-lobed	10	n.jl	V	W.Indies	1752.	S s.I	
4143 <i>Aiilerifolia L.</i>	Ivy.leaved	iOor	10	—	V	S. Amer.	1773.	S si Pl.ic.82 93.2
4144 <i>Nit. IerH</i>	Kit	—	10	ls	L.B	America	1097.	S s.p Bot. mag. 188
4145 <i>Aedcraccia L.</i>	Ivy-like	iOor	10	jls	B	N. Amer.	1729.	S s.p Bot. reg. 85
	<i>barbata Roth</i>							
4146 <i>barbfgera Swt.</i>	bearded	J O pr	3	jns	P	N. Amer.	1824.	S co Sw.flgar.86
4147 <i>cuspidate Pers.</i>	sharp-pointed	J O or	10	Jl.jl	L.B	S. Amer.	1732.	S s.p Fl.per.2.lift*
4148 <i>Dilfenti Desf.</i>	Dillemus's	J OI or	A	Jl.au	B	Guinea	...	S p.l Di. cl. 84 9H
4149 <i>/amnifolii L.</i>	Tannus-leaTcd	O or	10	jl	B	Carolina	1732.	S co Di. el. 318.410
4150 <i>grand i flora L.</i>	great-flowered	ft_ • or	8	s	W	E. Indies	1802.	S s.I Bot rep. 188
4151 <i>muru-ata Jac.</i>	muncate	& O ■	8	Jlau	P	E. Indies	1777.	S co Jac.sc 3.323
...	<i>purpurascens Bot. reg. 290</i>							
4152 <i>obscura L.</i>	obscure	■ ■ ■	8	jn.au	W	E. Indies	1732.	S s.p Bot. reg. 239
4153 <i>involucrita Beauv.</i>	involucrate	■ ■ ■	4	jn.jl	R	Guinea	182a	S s.I Beau. ow. ** w
4154 <i>hlspida Vahl</i>	hispid	■ ■ ■	10	jl.au	P	E. Indies	1820.	S r.m
4155 <i>capit&te Vahl</i>	headed	■ ■ ■	10	jl.au	B	K Indies	1823.	S s.I

ORDER L.

			4	jl.s	Pk	E. Indies	1815.	S	s.1
4156 pilosa Swt.	P>loM	—	4	jl.s	Pk	S.Amer.	1817.	S	r.m Jac. ic ₄ 2.318
Chenopodium pilosus	Box.	white-flowered	^ O or	6 jl.au	W	W. Indies	1825.	S	r.m Jac. vm.3.69
4157 leucantha Jac.		broad-flowered	■ ■ ■ ■ ■ ■	■ ■ ■ ■ ■ ■	W	E. Indies	1818.	S	r.m .
4158 latiflora Desv.					P	W. Indies	1823.	S	r.m Jac am. pic. 26
4159 strigilata Roth	*)?K ^{oae}		■ ■ ■ ■ ■ ■	■ ■ ■ ■ ■ ■	& Amer ¹	1810.	C	s.1 Pl. am. 102	
4160 filiformis Jac.	filiform			20 jl.s	W	W. Indies	1823.	S	r.m J.am.89^2
4161 panthaea B.SfS.	Auricula-?!!!				W	W. Indies	1822.	S	co Bot reg. 999
Convolvulus umbellatus L.		3 or	2 jl.au		Pk	E. Indies	1812.	C	pl
4162 corymbosa Forst.	corymbose	■ ■ ■ ■ ■ ■	6 au.a			K Indies	1820.	C	pl
4163 pudibunda B.B.	blushing					E. Indies	1816.	C	pl
4164 gangetica Swt.	Gangetic			12 my.au	Pk	E. Indies	1817.	C	pl
Convolvulus gangeticus Box.				12 jl.au	W	E. Indies	1816.	C	pl
4165 blanda Stirf.	**ft				R	E. Indies	1817.	C	pl R. mal. 11. S3
Convolvulus blandus Rox.		• or		10 jn.au	R	Carolina	1819.	S	co Bot reg. 437
4166 Kochia frigida Swt	Roxburgh s	J. ■ ■ ■ ■ ■ ■		8 jn.au	R	Kindles	1778.	S	co Botreg.317
multiflora Rox.					P	Mexico	1732.	R	s.p Dk el. 87.102
4167 sepiana Rox.	hedge	L.	3 jn.s		Y	India	1800.	S	s.l Jac. ic. 317
4168 sagittifolia Ker	sagittate-leaved [^]	A	6 jLau		Pk	E. Indies	1816.	S	co
4169 denticulata A. J.Pr.	denticulate [*]	■ ■ ■ ■ ■ ■	6 myjl		Pk	Mexico	1824.	C	pl
Convolvulus Mddium W.		ED or	6 jl.au		P	E. Indies	1778.	S	s.p R. mal. 11.65
4170 glaucifolia L.	glaucous-lvd	■ ■ ■ ■ ■ ■	6 au.o		P	Mexico	1823.	S	r.m Cav.ic. 3.250
4171 angustifolia Jac.	narrow-leaved		6 jl.au		V	E. Indies	1822.	C	pl
4172 spherocephala Swt.	round-headed		6 jl.au		Vi	E. Indies	1816.	S	pl
Convolvulus spherocephalus Rox.			10 jl.au		P	Cayenne	1824.	C	pl Bot mag. 2645
4173 hast! geratA.	spear-beaning	■ ■ ■ ■ ■ ■	6 jl.au		P	E. Indies	1770.	S	s.p Bot reg. 319
4174 tridentatdL.	three-toothed	■ ■ ■ ■ ■ ■	10 my.jl		P	S.Amer.	1724	R	s.p PL am. 89.104
4175 stans Cav.	standing				P	E. Indies	1800.	C	pl
4176 aurifolia Swt.	Laurel-leaved	■ ■ ■ ■ ■ ■	4 jn.au		Pk	R Indies	1806.	D	p.l B. maL 11.
Convolvulus auritolioides Bo ¹		or	■ ■ ■ ■ ■ ■	■ ■ ■ ■ ■ ■	P				
4177 jalicifolia Rox.	Willow-leaved		■ ■ ■ ■ ■ ■	■ ■ ■ ■ ■ ■	P				
4178 amplexicaulis Sims	Bi-pinnate like		■ ■ ■ ■ ■ ■	■ ■ ■ ■ ■ ■	P				
4179 brasiliensis L.	sea-side		■ ■ ■ ■ ■ ■	■ ■ ■ ■ ■ ■	P				
4180 stipulacea Swt.	Brazilian		■ ■ ■ ■ ■ ■	■ ■ ■ ■ ■ ■	P				
Convolvulus stipulaceus Box.	stipulaceous				P				
4182 reptans L.	creeping	^ E3 or	i jn.au		P				

* 492. CONVOLVULUS *L.* BINDWEED. {*Convolvere*, to entwine; stems Britain

- 4236 tricolor L. three-colored O or 3 jl.au St S Europe 1629. S co Bot mag. 27
 2 albiflbrus white-flowered O or 1 jl.au W S. Europe 1629. S co
 4237 filiranlu Vahl thread-stmd JU[A-jun 1 jn.jl Y Guinea 1824. S sl
 4238 Sibth⁶puii R. & S. Sibthorp's - A or 1 jns - W France 1823. R co Fl. gr. 193
 4239 ebracteatus Detf. ebrai-teate - A or 1 jn.s W 1819. R co
 4240 multifidus Thun. many-cleft - O un 2 jns P C. G. H. 1821. S s.l
- *493. ARGYREIA Lou. SILVER-WEED. (*Argyreios*, silvery; texture of leaves.) *Convolvulaceae*. 7.—12.
 4241 splendens Swt. shining 1. CD or 1 n Fk E. Indies 1820. C co Bot mag. 2088
 Lettsbina splendens Box.
 4242 speciosa Swt. showy % * or 10 jl.au R E. Indies 1818. C r.m Bot mag. 2446
 Ipomoea a speciosa Bot. mag. 2239. C. nervosus Brm. Lettsbina nervosa Box.
 4243 cuveata Ker wedde-leaved fc. * or 10 au.s P E. Indies 1822. C s.l Bot. reg. 661
 Lettsdmania cuneata Box.
 4244 cym&a Swt. cymose fc. * or 10 ... Pk E. Indies 182a C r.m
 Lettbinfo cymosa Box.
 4245 bdna-nox Swt. good-night t. O or 20 ... W E. Indies 1799 C r.m
 Lettsinjia bona-nox Box.
 4246 umflora Swt. one-flowered % CD or 30 ... W E. Indies 1817. C r.m
 Lettedinia uni flora Box.
 4247 ornata Swt. adorned fc_E2or 10 ... W E. Indies 1824. C r.m
 Lettsbina ornata Box.
494. CALYSTEGIA BBr. BEARBIND. (*Kalyx*, a calyx, *stege*, a covering.) *Convolvulaceae*. 10.—11.
 4248 septum B. Br. great hedge j A or 6 jns W BriUin m. hed. R s.l Eng. bot 313
 Convolvulus bdp. []
 2 incarnata red-flowered A or 6 jns R N. Amer. ... R co Bot mag. 732
 4249 sylv&tms B. & S. wood A or 18 jns W Hungary 1815. R co
 4250 spithamea Ai Ph. span A or 1 jl W N. Amer. 1796. R co Hook. ex. fl. 07
 Convolvulus spithamea L.
 4251 paradoxa Ph. paradoxical T A un 3 jns W N. Amer. 1810. R s.l
 4252 Catesbofria Ph. Catesby's T A un 3 jns P N. Amer. 1810. R s.l
 4253 dahurica Fis. Dahurian A pr 1jjl Pk Dauria 1823. S co Bot mag. 2609
 4254 tomentosa A. SS. woolly I A or ... jnjl - N. Amer. 1818. D p.l
 Convolvulus stans Mx.
 4255 Soklanella B. Br. Soldanella JU A or 1 jnjl F Britain seash. R s.l Eng. bot 314
 Convolvulus Soldanella E. B.
 4256 reniformis B. Br. kidney-shaped - A un T I I F N. Holl. 1817. R s.p Sw. fl. gar. 181
 4257 marginata B. Br. margin ate " - A Pk N. Holl. 1824. R r.m
495. MORENOS Lai. MOREXOA. (*P. Moreno*, a Mexican patriot) *Convolvulaceae*. 3.—5.
 4258 patula Lai. spreading % El or 6 ... S Mexico 1826. C p.l
 4259 grandiflora Lai. large-flowered i Eflor fi ... S Mexico 1826. C p.l
 4260 globo&a Lai. globose % LSI or 6 ... S Mexico 1827. C p.l
496. PORA^NA Brm. PORANA. (*Poreuo*, to journey; extension of branches.) *Convolvulaceae*. 1.—3.
 4261 volubilis Brm. twining fc. * or ^50 jl.au W K Indies 1820. S p.l Bur. in. 21
497. DINETTUS Swt. DINETUS. (*Dineios*, that may be twined.) *Convolvulaceae*. 2.
 4262 racemosa Swt. racemose - f O P* 12 au. 8 " Nepal 1823. S co Sw.fl.gar.127
 Porkna racemosa Box.
 4263 paniculata Swt. panicled J_ * or 10 au.s W E. Indies 182a C p.l
 Porana paniculata Box.
498. COB^T^ Cav. COBJSA. (*B. Cobo*, a Spanish botanist.) *Cobtehcete*. 1. — 2.
 4264 scandens Cav. climbing fl_iAJor 20 my.o P Mexico 1792. S p.l Bot mag. 851
- *499. GI'LIL B.&P. GILIA. (*P. S. Gilio*, a Spanish botanist) *Polemoniaceae*. 3. — 6.
 § 4265 coronopifolia Pen. Buckhorn-lvd ^ O | or '3 au.s S Carolina 1726. C l.p Ex. bot. 1.13
 Cdtnti coronopitdia W. Ipomoea elegans Mx.
 4266 parvillora Spr. small-flowered O or 2 s.n B America 1793. S co Ex. bot. 1.14
 llKJinopsis mconspicua 5m. Cdtnta parviflora Ph.
 4267 capitata Hook. capitate Q or 1 jl.au B Colombia 1826. S p.l Bot. mag. 2698
500. CALDA^S7A W. CALDASIA. (*J. Caldas*, a Bogotan naturalist) *Polemoniaceae*. 1.
 4268 heterophylla W. variable-leaved O or lijn.au B N. Spain 1813. S s.p Bot.rdg.96
 Bonphindia geminiflora Cav.
501. HOITZIA J. HOITZIA. (*Hoitzit*, its name in Mexico.) *Polemoniaceae*. 3. — 8.
 4269 cocinea Cav. scarlet i j o r 3 ... S Mexico 1824. C r.m Cav. ic. 6.365
 4170 crathea Cav. blue * i j o r 1 ... Pa.B Mexico 1824. C r.m Cav. ic. 6.366
 127i glandulosa Cav. glandulous * i _Jor 2 ... Pa. R Mexico 1825. C r.m Cav. ic. 4.367
502. RETZI/f Thun. RBTZIA. (*A. J. Retzius*, prof. nat. hist at Lund, in Sweden.) *Convolvulaceae*. 1.— ■
 4272 spicata Thun. spiked <t i_J cu 4 my.jn Br C. G. H. ... C l.p Lam. il. 103
- *503. LUBI^N'il Com. LUBINIA. (*M. drSt. Lubin*, a French officer.) *Primulaceae*. 1.—3.
 4273 atropurpuriflra Lk. dark-purple JU iAJ cu 2 ... P C. G. H. 1820. C l.p H.ber.27
504. "E'PACRIS Forsk. EPACRIS. (fx, upon, akro**t* the top; habitation.) *Epacridae*. 14.— 20.
 4274 purpurascens B. Br. purpurascent ft iJor 3 ja mr P N. S. W. 1803! C sp Bot.mag.84*
 2 rObra rod-flowered ft iJor 3 ja.mr R N. Holl. 1K03. C s.p Hot. cab. 876
 4275 pulchilla B. Br. neat ft iJor 4 apjn Pk N. S. W. 1804. C 8.p Bot. mag. 1170
 4276 microphilla B. Br. small-leaved ft iJor 2 . . . - N. S. W. Ih22. C l.p
 4277 grandiflora B. Br. great-flowered ft iJor 3 ja.jn S N. S. W. 1803. C sp Bot.mag.982
 4278 0i^osin^oflora Cun. Onosma-flwd ft iJor 2 apjl R N. Holl. 1823. C s.p
 4279 rusciflbia li. Br. Ruscus-leaved ft iJor 2 apjn - N. Holl. 1824. C s.p
 4280 sparsa B. Br. scattered . iJor 2 apjn N. Holl. 182&. C s.p
 4281 apiculata Cun. small-tufted ftLJor 2 apjn N. Holl. 1825. C s.p
 4282 obtusiflba B. Br. blunt-leaved ft iJor 3 apjn W N. S. W. 1804. C s.p Ex. bot. 140
 4283 imprecsaa Lab. impressed . iJor 3 apjl C N. Holl. 1824. ■ s.p Sw. au. 4
 4284 heterondoma Lab. vanous-stmd ft iJor 3 apjl W N. Holl. 1823. C s.p Lab.n.h.56
 4285 paludbsa R. Br. marsh . iJor 3 apjl Pa R N. Holl. 1825. C s.p
 4286 exsertd R. Br. drawn-out ft iJor 2 apjn W Vi. Di. L. 1812. C l.p
 4287 mucronulata B. Br. small-pointed ft LJor 3 apjl R N. Holl. 1824. € s.p
505. STYPHE^NIA B Br. STYPHKLIA. (*Syphyllos*, rigid; compact habit) *Epaerideae*. 8.— 9.
 4288 tubiflora Sm tube-flowered ft iJor 6 my.au C N. S. W. 1802. S s.p Sm. n. h. 1+
 4289 longifolia B. Br. long-leaved ft iJor 3 apjn G N. S. W. 1807. C l.p Bot. reg.**

- 42901 «a7l. 7?r.
W.H adscemlens R. Br.
429* viridiflora R. Br.
4i⁴ triflbra B. Rep
4204 epacrioldc* Hort.
4295 latiflria A Tir.
- fruitful ■ U or 5 ... G N. Holl. 1829. C s.p
ascend. ng ■ U^of 3 " G N. Holl. 1822. C s.p
green-flowered ■ U or 4 apjn G N. S. W. 17^U. C a p Bot. rep. 312
three-flowered ■ U or 1 my au Pk N. S. W. 1796. C s.p Bot mag. 129?
Epacriis-like ■ U or 6 w C N. Holl. 1823. C s.p
broad-leaved ■ U or 4 apjl Pk N. Holl. 1823. C s.p
506. MELICHRUS it Br. MELICIBUS. (*Mrtichros*, honey-colored; gl mds of flowers) *Epaeridete*. 2.—&
4296 rotatus it Tir. rotate « U o r lijp.au S N. Holl. 18^a4. C sp Cav. ic. 4.349.1
4247 m&dius Cm. middle 1 y o r 2 ap.my S N. Holl. 1824. C s.p
007. LISSANTHE R. Br. LISSANTUB. (*Lissos*, smooth, *anthos*, a flower.) *Epaeridex*. 5.—6.
4299 sapida R. Br. Savoury ■ i L J or 4 jnjil W N. Holl. 1824. C s.p
4299 subul&tta R. Br. subulate ■ U or 2 apjl W N. Holl. ■ C s.p
4 XX strigula R. Br. strigose ■ U or 3 myjl W N. Holl. C s.p
4 Jul daphnooides R. Br. Daphne-like ■ U or 3 jnjil W N. HU. 1818. C s.p
4302 ciliaU R. Br. Ciliated ■ U or 3 myjl W N. Holl. 1825. C s.p Bot.cab.4fib¹
- 508 ASTROLOWTA U Br. ASTBOLOMA. (*Astron*, a star, *loma*, a fringe; flower.) *Epaertdece*. 2.—6.
4W3 humifusum R. Br. trailing i j o r 1 myo S N. S. W. 1H07. C s.p Bot. mag. 1439
4XH denticulatum 71 Tir. denticulate « U o r 1 my au Pa R N. Holl. 1824. C s.p
509. SPRENGFLM Sm SPBBNGBLU. (C. C. Sprengel, of Spandau, in Br.mdenburgh.) *Epacrldete*. 1—2.
4:05uidcrnataSm. fLesh-cototed ft LJ or 2 apjn F N. S. W. 179J. C s.p Bot. mag. 1719
510. ANDERSONI^N Ni⁴ R. Br. ANDERSOVU (Messrs. W. A, IV., mAG. Anderson.) *Epacrider*. 1.—6.
K5 sprengeliirc R. Br. Sprengeha-like ft LJ or 2 mrjl Pk N. Holl. 180J. C s.p Bot. mag. 1o45
511. LYSINEWIA R. Br. LVSIVEMA. (*Lysis*, a sep-iration, *nema*, a stamen.) *Epaeridete*. 4.—fi.
J J⁷ pcampetalum/R^Br. five-petaled ft U^o 2 fmpy Pk N. Holl. 1813. C 1.p
4J08 consicuum R. Br. conspicuous ■ i L J or 2 fmpy Pk N. Holl. ■ C ip
4JUy pi'mgens R. Br. pungent ■ U or 2 flap⁷ W N. S. W. 1804. C Lp Bot. m.-g. 844
2 rhbrum red ■ U or 2 tap R N. S. W. 1814. C Lp Bot. mag. 1199
F/pacry rosaa Lod.
4310 attenuatium Lk. attenuated « L j o r 2 f.ap W N. S. W. 1812. C 1p Bot cab. 38
4311 lasianthum Tt Tir. hairy-flowered ft LJ or fi fmy Pk N. Holl. 1820. C 1p
- 512. DRACOPHYLLUM 71 Tir. DBACOPHYLLUM. (*Dmkos*, a dragon, *phyllon*, a leaf.) *Epacrdete*. 3.—&
4312 scc.'ndum it Tir. side-flowering ft U or 2 ap au W N. Holl. 1823. S s.p Sw. au. ic. in.
4J13 longitklum TZ Tir. long-leaved i j o r 2 ap au W N. Holl. 18^a4. C s.p
44314 gracileTZ Tir. slender ft j o r 2 ap.my W N. Holl. 182J. C s.p Bot. mag. 2678
513. TROCHOCA'RPA 71.Tir. TBOCHOCABPA. (*Trochos*, a wheel; *karpos*, fruit; form.) *EpacrMea*. 1.
4315tefirina R. Br. LaurcUtte ft U o r 4 ap.au W N. Holl. 182J. C s.p Lin.tr.8.9
Styphitia cornifolia Rud.
514. ACRO'TRICHIE R. Br. AcaoTKiniE. (*Akros*, a point, *thrix*, hair: coroHO *Epacrldef*. 3.—8.
4316 corditaT? Tir. heart-m-rf ft U o r 1 apjl W N. Holl. 1823. C s.p Lab. n. h. 1.63
4317 ovalifolia R. Br. oval-leaved « U o r Jap.au W N. Holl. 182J. C s.p
4318 divaricata 7%. Tir. divaricate ft • or J my W N. Holl. 1824. C s.p
515. MONOTOCOA R. Br. MOXOTOCOA. (*Monos*, one, *tokos*, birth; *germen* one-seeded.) *Epaeridete*. 4.—0.
•319elliptica R. Br. elliptic ft i (or 8 my au W N. S. W. 1H02. C 1p
4J20)meata7J. Tir. lined ft U or 6 my.au W V. Di. L 1804. C s.p Lab. n.h. 1.61
- 14 ■■■■■ R.Br. Broom ■ i J or 5 my.au W N. Holl. 1825. C s.p
■■■■■ R.Br. white ■ LJ or 6 my.au W N. Holl. 1824. C s.p
516. PONCELETTW R. Br. PONCELETIA. (*M. Poneelet*, a French botanist.) *Epacridea*. 1.
«*sprengelwid* R. Br. Sprengeha-hke ft U o r 1 myjn ... N. Holl. 182G. C pJ
- Jj1⁷ COSMEALIA R. Br. COSMBLIA. (*Kosmeo*, to adorn; beauty.) *Epaeridete*. 1.
*4 rubra Tt Tir. reti-stotvered ft U o r i ... It N. Holl. 1826. C s.p
- J J⁰ ^ EUCOPO^GON R. Br. LsticoponoM. (*Isukos*, white

ogon, beard; flowers.) *Epaeridete*. 13—48
F&E ^ ceolutus R. Br. lanceolate ft U or 12 my.au W N. S. W. 1790. C s.p Bot. rep. 287
JSM HLL R. Br. Kiehle's ■■■■■ 5 my.au W M. Holl. 1812. C 1.p Lab.n.h. 1.66
■■■■■ Kiehle Kichei Lab.
*3s7 obovktus 71 Tir. obovate ■■■■■ 1 my.au W N. Holl. 1824. C Lp Lab.n.h.lfi?
Styphelia obovka Lab.
4328ericoides It.Br. H^oonth iakwt A. in* 6 ... W K.S.W. 1815. C lp Cav.ic.4.347.1

SSSI TM «C
JTS SSI"iSf fln ■■■■■ StJS ■■■■■ StJS
8temrasping ft U OJ 3 ... X9 KSW. 181^a8. C lp Lin.tr.8.8

J J⁰ ^ JJ¹ small-leaved ft LJ or 2 myjl W K Holl. 1818. C lp Cav.ic.4.849.
■■■■■ R. Br. Brialied ft LJ or 3 my.au "W N. Holl. 1823. C l.p
■■■■■ R. Br. pressed togetherft \ or 3 my.au W N. Holl. 181m C l.p
Juniperinus it Br. Junipe>r/^ve< ft t lor 3 apjn W N. S. W. 1804. C Lp Bot. cab. 447
scigorous ft i or 2 ... W N. Holl. 1884. C l.p

W ■■■■■ CYATHODES. (A>/A^w, ac like; nectary) *Epaeridete*. 3.—7.
glaucous 1 U o r 20 apjn V.Di. 1. 1818. C ep Lab.n. h. 1.6
Pnckly Cedar ft LJ or 6 apjn V. Di. 1. 1822. C s.p Lab. u. h. 1.6
4110 Whchaa Ox^cedrus Lab chalTy ■■■■■ U o r 8 ap.au W N. Holl. 1823. C s.p Gae.flr.2.94

■■■■■ NTHERA R.Br. STENANTHEBA. (*Stenos*, narrow, *anthera*, an anther.) *Epaai-de**. 1.
■■■■■ Br. Pino-leaved ft U o r 6 myjl S N. S.W. 1811. C s.p Botreg.218

521. AZA'LEA L. AZALEA. (*AzaJeos*, dry, arid; habitation.) *Rhodordrea*. 1.1—
■■■■■ indica L. Indian ft LJ or 4 mr my S China 1808. C p1 Bot. mag. 1480
2 purpurea pldna double-purple ft \ or 4 mr my P China
3 variegata variegated ft i or 4 mr my St China
4 ilba pure-white ft I or 4 mr.my W China 1819. C p1 ■■■■■
5 aurantla orange ft I or 4 mr.my O China 1²². C ii.l
■■■■■ Pontic ft or 3 my p China lh24. C p1
Chinese ft or 3 my Y China ■■■■■ U p Bot. cab. 885
Pontic ft or 6 myjn Y Turkey ■■■■■ L Bp Bot.ni4g.433
glaucous Sk or 6 myjn Y Turkey ■■■■■ L s.p
white-flowered ft or 6 myjn W Turkey ■■■■■ L sp Bot.mag.2::ai

PENTANDRIA MONOGYNIA.

4 <i>cuprea</i>	<i>copper-colored</i>	<i>Sk</i>	or 0	inr.my Cop	Turkey	—	L s.p
5 <i>pallida</i>	<i>three-colored</i>	<i>Sk</i>	or (i)	mr.my Pa.Y	Turkey	—	L s.p
6 <i>tricolor</i>	Marygold-like	<i>ft</i>	or 6	mrmY Y.R	Turkey	—	L s.p
4344 <i>calendulaeca Mx.</i>	BaftVon-colored	<i>ife</i>	or 4	my.jn O	N. Amer. 1806.	L s.p	
1 <i>crtaea</i>	fine-golden	<i>*</i>	or 4	my.jn Saf	N. Amer. ...	L p.l	Bot mag. 172
2 <i>chrysocarpa</i>	large-flowered	<i>a*</i>	or 4	my.jn Y	N. Amer. ...	L s.p	
3 <i>grandiflora</i>	<i>copper-colored</i>	<i>Sk</i>	or 4	my.jn O	N. Amer. ...	L s.p	
4 <i>cuprea</i>	shining	<i>ft</i>	or 4	my.jn Cop	N. Amer. ...	L p.l	
5 <i>splendens</i>	flame-colored	<i>ft</i>	or 4	my.jn It	N. Amer. 1812.	L s.p	Bot reg. 145
6 <i>flimmea</i>	triumphant	<i>ft</i>	or 4	my.jn O	N. Amer. ...	L p.l	
7 <i>triumphans</i>	ignescens	<i>ft</i>	or 4	my.jn R	N. Amer. ...	L p.l	
8 <i>igneous</i> •	canescens	<i>ft</i>	or 3	my.jn R	N. Amer. 1812.	L s.p	
4345 <i>canescens Mx.</i>	showy	<i>ft</i>	or 4	my.jn S	N. Amer. ...	L p.l	Den. br. 116
4346 <i>speciosa W.</i>	larger-fcarfe/	<i>ft</i>	or 4	my.jn S	N. Amer. ...	L p.l	Bot cab. 634
1 <i>major</i>	curled	<i>ft</i>	or 4	my.jn S	N. Amer. ...	L s.p	
Scrfspa	orange	<i>ft</i>	or 4	my.jn S	N. Amer. ...	L s.p	
3 <i>aurintia</i>	oblique	<i>ft</i>	or 4	my.jn —	N. Amer. ...	L s.p	
4 <i>obliqua</i>	Plum-leaved	<i>ft</i>	or 4	my.jn —	N. Amer. ...	L s.p	
5 <i>prunifolia</i>	hooded	<i>ft</i>	or 4	my.jn —	N. Amer. ...	L e.p	
6 <i>cucullata</i>	revolute	<i>ft</i>	or 4	my.jn —	N. Amer. ...	L s.p	
7 <i>revoluta</i>	ciliated	<i>ft</i>	or 4	my.jn —	N. Amer. ...	L s.p	
8 <i>ciliata</i>	acute-leaved	<i>ft</i>	or 4	my.jn —	N. Amer. ...	L s.p	
9 <i>acutifolia</i>	<i>wavcA-leaved</i>	<i>ft</i>	or 4	my.jn —	N. Amer. ...	L s.p	
10 <i>undulata</i>	twisted-leaved	<i>ft</i>	or 4	my.jn —	N. Amer. ...	L s.p	
11 <i>tortifolia</i>	naked-flowered	<i>ft</i>	or 3	my.jn S.Pk	N. Amer. 1734.	L s.p	
4347 <i>nudiflora L.</i>	Ai/y-white	<i>ft</i>	or 4	my.jn W	N. Amer. ...	L s.p	
1 <i>alba pldna</i>	double-white	<i>ft</i>	or 4	my.jn W	N. Amer. ...	L s.p	
2 <i>bella</i>	soft	<i>ft</i>	or 4	my.jn Bh	N. Amer. ...	L p.l	
4 <i>carnea</i>	fleshy	<i>ft</i>	or 4	my.jn P.R	N. Amer. 1734.	L p.l	Bot reg. 120
5 <i>caroliniana</i>	Carolina	<i>ft</i>	or 4	my.jn S	N. Amer. ...	L p.l	
6 <i>coccinea</i> -	scarlet	<i>ft</i>	or 4	my.jn S	N. Amer. ...	L p?	Bot mag. 180
7 <i>corymbosa</i>	corymbose	<i>ft</i>	or 4	my.jn S	N. Amer. ...	L s.p	
8 <i>crispa</i>	curled	<i>ft</i>	or 4	my.jn Pk	N. Amer. ...	L s.p	
9 <i>cumulata</i>	bundled	<i>ft</i>	or 4	my.jn S.Pk	N. Amer. ...	L s.p	
10 <i>discolor</i>	various-colored	<i>ft</i>	or 4	my.jn W.s	N. Amer. ...	L s.p	
11 <i>fastigita</i>	pyramidal	<i>ft</i>	or 4	my.jn Pk	N. Amer. ...	L s.p	
12 <i>floridea</i>	many-flowered	<i>ft</i>	or 4	my.jn Pk	N. Amer. ...	L s.p	
13 <i>globosa</i>	globose	<i>ft</i>	or 4	my.jn Pk	N. Amer. ...	L s.p	
14 <i>glomerata</i>	glomerated	<i>ft</i>	or 4	my.jn Pk	N. Amer. ...	L s.p	
15 <i>incana</i>	hoary	<i>ft</i>	or 4	my.jn Pk	N. Amer. ...	L s.p	
16 <i>incarnata</i>	<i>tLcsh-colored</i>	<i>ft</i>	or 4	my.jn t	N. Amer. ...	L s.p	
17 <i>irinabilis</i>	wonderful	<i>ft</i>	or 4	my.jn S	N. Amer. ...	L s.p	
18 <i>magnifica</i>	magnificent	<i>ft</i>	or 4	my.jn S	N. Amer. ...	L s.p	
19 <i>montana</i>	mountain	<i>ft</i>	or 4	my.jn S	N. Amer. ...	L s.p	
20 <i>pallida</i>	<i>pdc-flowered</i>	<i>ft</i>	or 4	my.jn Pa.R	N. Amer. ...	L s.p	
21 <i>paludosa</i>	marsh	<i>ft</i>	or 4	my.jn Pk	N. Amer. ...	L s.p	
22 <i>papilionacea</i>	butterfly	<i>ft</i>	or 4	my.jn Str	N. Amer. ...	L s.p	
23 <i>partita</i>	five-parted	<i>ft</i>	or 4	my.jn W.R	N. Amer. ...	L s.p	
24 <i>parviflora</i>	small-flowered	<i>ft</i>	or 4	my.jn ...	N. Amer. ...	L s.p	
25 <i>prolifera</i>	proliferous	<i>ft</i>	or 4	my.jn ...	N. Amer. ...	L s.p	
26 <i>pamila</i>	dwarf	<i>ft</i>	or 4	my.jn W	N. Amer. ...	L s.p	
27 <i>purpurascens</i>	purpurascens	<i>ft</i>	or 4	my.jn P	N. Amer. ...	L s.p	
28 <i>purpurea</i>	purple	<i>ft</i>	or 4	my.jn P	N. Amer. ...	L s.p	
29 <i>purpurea plena</i>	double-purple	<i>ft</i>	or 4	my.jn P	N. Amer. ...	L s.p	
30 <i>rosea</i>	roseate	<i>ft</i>	or 4	my.jn R	N. Amer. ...	L s.p	
31 <i>rubriflora</i>	reddest	<i>ft</i>	or 4	my.jn D.R	N. Amer. ...	L s.p	
32 <i>rubescens</i>	rubescent	<i>ft</i>	or 4	my.jn R	N. Amer. ...	L s.p	
33 <i>rubicunda</i>	ruddy	<i>ft</i>	or 4	my.jn R	N. Amer. ...	L s.p	
34 <i>rubra</i>	red	<i>ft</i>	or 4	my.jn R	N. Amer. ...	L p.l	Bot cab. 51
35 <i>rutilans</i>	shining-red	<i>ft</i>	or 4	my.jn D.R	N. Amer. ...	L p.l	
36 <i>scmiduplex</i>	semidouble	<i>ft</i>	or 4	my.jn W	N. Amer. ...	L p.l	
37 <i>staminea</i>	taw-stamened	<i>ft</i>	or 4	my.jn R	N. Amer. ...	L p.l	
38 <i>stellata</i>	starry	<i>ft</i>	or 4	my.jn R	N. Amer. ...	L p.l	
39 <i>tricolor</i>	three-colored	<i>ft</i>	or 4	my.jn S.W	N. Amer. ...	L p.l	
40 <i>vanabilis</i>	variable	<i>ft</i>	or 4	my.jn R	N. Amer. ...	L p.l	
41 <i>variegata</i>	variegated	<i>ft</i>	or 4	my.jn R,w	N. Amer. ...	L p.l	
42 <i>versicolor</i>	party-colored	<i>ft</i>	or 4	my.jn R,w	N. Amer. ...	L p.l	
43 <i>Violacea</i>	violet-colored	<i>ft</i>	or 4	my.jn V	N. Amer. ...	L p.l	
4548 <i>bicolor Jt.</i>	Cohu rail's	<i>ft</i>	or 4	my.jn S	N. Amer. ...	L p.l	
4347? <i>lwidixi A.</i>	two-colored	<i>ft</i>	or 4	my.jn St	N. Amer. 1734.	L s.p	Tr.clir. 48
1 <i>odorifera</i>	dummy	<i>ft</i>	or S <i>il.au</i>	W	N. Amer. J7J4.	L s.p	Mce.ic.IL 9
2 <i>crispia</i>	* ^{1*2*3*4*5*6*7*8*9*10*11*12*13*14*15*16*17*18*19*20*21*22*23*24*25*26*27*28*29*30*31*32*33*34*35*36*37*38*39*40*41*42*43*44*45*46*47*48*49*50*51*52*53*54*55*56*57*58*59*60*61*62*63*64*65*66*67*68*69*70*71*72*73*74*75*76*77*78*79*80*81*82*83*84*85*86*87*88*89*90*91*92*93*94*95*96*97*98*99*100*101*102*103*104*105*106*107*108*109*110*111*112*113*114*115*116*117*118*119*120*121*122*123*124*125*126*127*128*129*130*131*132*133*134*135*136*137*138*139*140*141*142*143*144*145*146*147*148*149*150*151*152*153*154*155*156*157*158*159*160*161*162*163*164*165*166*167*168*169*170*171*172*173*174*175*176*177*178*179*180*181*182*183*184*185*186*187*188*189*190*191*192*193*194*195*196*197*198*199*199*200*201*202*203*204*205*206*207*208*209*2010*2011*2012*2013*2014*2015*2016*2017*2018*2019*2020*2021*2022*2023*2024*2025*2026*2027*2028*2029*2030*2031*2032*2033*2034*2035*2036*2037*2038*2039*2040*2041*2042*2043*2044*2045*2046*2047*2048*2049*2050*2051*2052*2053*2054*2055*2056*2057*2058*2059*2060*2061*2062*2063*2064*2065*2066*2067*2068*2069*2070*2071*2072*2073*2074*2075*2076*2077*2078*2079*2080*2081*2082*2083*2084*2085*2086*2087*2088*2089*20810*20811*20812*20813*20814*20815*20816*20817*20818*20819*20820*20821*20822*20823*20824*20825*20826*20827*20828*20829*20830*20831*20832*20833*20834*20835*20836*20837*20838*20839*20840*20841*20842*20843*20844*20845*20846*20847*20848*20849*20850*20851*20852*20853*20854*20855*20856*20857*20858*20859*20860*20861*20862*20863*20864*20865*20866*20867*20868*20869*20870*20871*20872*20873*20874*20875*20876*20877*20878*20879*20880*20881*20882*20883*20884*20885*20886*20887*20888*20889*208810*208811*208812*208813*208814*208815*208816*208817*208818*208819*208820*208821*208822*208823*208824*208825*208826*208827*208828*208829*208830*208831*208832*208833*208834*208835*208836*208837*208838*208839*208840*208841*208842*208843*208844*208845*208846*208847*208848*208849*208850*208851*208852*208853*208854*208855*208856*208857*208858*208859*208860*208861*208862*208863*208864*208865*208866*208867*208868*208869*208870*208871*208872*208873*208874*208875*208876*208877*208878*208879*208880*208881*208882*208883*208884*208885*208886*208887*208888*208889*2088810*2088811*2088812*2088813*2088814*2088815*2088816*2088817*2088818*2088819*2088820*2088821*2088822*2088823*2088824*2088825*2088826*2088827*2088828*2088829*2088830*2088831*2088832*2088833*2088834*2088835*2088836*2088837*2088838*2088839*2088840*2088841*2088842*2088843*2088844*2088845*2088846*2088847*2088848*2088849*2088850*2088851*2088852*2088853*2088854*2088855*2088856*2088857*2088858*2088859*2088860*2088861*2088862*2088863*2088864*2088865*2088866*2088867*2088868*2088869*2088870*2088871*2088872*2088873*2088874*2088875*2088876*2088877*2088878*2088879*2088880*2088881*2088882*2088883*2088884*2088885*2088886*2088887*2088888*2088889*20888810*20888811*20888812*20888813*20888814*20888815*20888816*20888817*20888818*20888819*20888820*20888821*20888822*20888823*20888824*20888825*20888826*20888827*20888828*20888829*20888830*20888831*20888832*20888833*20888834*20888835*20888836*20888837*20888838*20888839*20888840*20888841*20888842*20888843*20888844*20888845*20888846*20888847*20888848*20888849*20888850*20888851*20888852*20888853*20888854*20888855*20888856*20888857*20888858*20888859*20888860*20888861*20888862*20888863*20888864*20888865*20888866*20888867*20888868*20888869*20888870*20888871*20888872*20888873*20888874*20888875*20888876*20888877*20888878*20888879*20888880*20888881*20888882*20888883*20888884*20888885*20888886*20888887*20888888*20888889*208888810*208888811*208888812*208888813*208888814*208888815*208888816*208888817*208888818*208888819*208888820*208888821*208888822*208888823*208888824*208888825*208888826*208888827*208888828*208888829*208888830*208888831*208888832*208888833*208888834*208888835*208888836*208888837*208888838*208888839*208888840*208888841*208888842*208888843*208888844*208888845*208888846*208888847*208888848*208888849*208888850*208888851*208888852*208888853*208888854*208888855*208888856*208888857*208888858*208888859*208888860*208888861*208888862*208888863*208888864*208888865*208888866*208888867*208888868*208888869*208888870*208888871*208888872*208888873*208888874*208888875*208888876*208888877*208888878*208888879*208888880*208888881*208888882*208888883*208888884*208888885*208888886*208888887*208888888*208888889*2088888810*2088888811*2088888812*2088888813*2088888814*2088888815*2088888816*2088888817*2088888818*2088888819*2088888820*2088888821*2088888822*2088888823*2088888824*2088888825*2088888826*2088888827*2088888828*2088888829*2088888830*2088888831*2088888832*2088888833*2088888834*2088888835*2088888836*2088888837*2088888838*2088888839*2088888840*2088888841*2088888842*2088888843*2088888844*2088888845*2088888846*2088888847*2088888848*2088888849*2088888850*2088888851*2088888852*2088888853*2088888854*2088888855*2088888856*2088888857*2088888858*2088888859*2088888860*2088888861*2088888862*2088888863*2088888864*2088888865*2088888866*2088888867*2088888868*2088888869*2088888870*2088888871*2088888872*2088888873*2088888874*2088888875*2088888876*2088888877*2088888878*2088888879*2088888880*2088888881*2088888882*2088888883*2088888884*2088888885*2088888886*2088888887*2088888888*2088888889*20888888810*20888888811*20888888812*20888888813*20888888814*20888888815*20888888816*20888888817*20888888818*20888888819*20888888820*20888888821*20888888822*20888888823*20888888824*20888888825*20888888826*20888888827*20888888828*20888888829*20888888830*20888888831*20888888832*20888888833*208888888}						

4**?" ALL. AMA'NDI* L. ALLAMANDA. (Dr. F. AUamand, prof. at Leyden.)	<i>Apodine*</i> . 1.-2.
*>9 cathartic L. cathartic H Q or 12 jn.jl Y Guiana 1785. C r.m Bot. mag. 338	
ZEK THEOPHRA'ST^ L. THEOPHRABA. (Tkcophrastus, father of nat. hist.)	<i>Apoepnete</i> . 1.-2.
uf Lindl. Jussiaeus's * i_J or 3 ... W Hispanio. 1818. S r.m lahd. col. 26	
.527. CLAVIM Rper. CLAVIJA. (J. Clavijo Paxardo, a Spanish naturalist)	<i>Myroxinece</i> . 1. - 4.
*1 macrophyila Ft. per. long-leaved & D or SO ... W S. Amer. 1816. C r.m	
.528 n'NCA /,. PERIWINKLE. (Tinea/urn, a band; twining shoots)	<i>Apoepnete</i> . 5. - 6.
J>62 hertcea J3f. herbaceous JU ■■■■■ ■■■■■ Hungary 1816. D s.l Bot. mag. 2002	
*>oj nrnor L. lesser JL, or 1 mr.s V Britain bu. pi. D co Eng. bot 917	
I>H o argenteo-variegata white-jfotsraf JL or 1 mr.s W Britain gard. D co	
43G1 Jfotgo variegata double-striped JL or 2 mr.s V Britain gard. D co	
variegated JU or 2 mr.s V England groves. D co Eng. bot. 514	
4365 pusillaZ. greater JU or 2 mr.s B England gard. co	
amall-flowered O or au B E. Indies 1778. S s.l Mur.2.1	
a 2 alba 3 ocellata	B
roseate n. CD or 1 mr.o R.w E Indies 1756. C r.m Bot. mag. 248	
vrwhite-flowered «. or 1 mr.o W E. Indies 1756. C r.m	
rerf-eyed tt-Qor 1 mr.o St E. Indies 1756. C r.m	
529. ^E%RIU L. NERIUM. (Neros, humid; habitat.) <i>Apoepnete</i> . 3.-5.	
4367 Oleander L. Oleander ■■■■■ Jor 8 jn.o R S. Europe 1596. L r.m Lam. il. 174	
2 filbum ■■■■■ Jor 8 jn.o W S. Europe 1596. L r.m Bot. cab. 700	
3 splendens ■■■■■ Jor 7 jn.o R & Europe 1814. L r.m	
* variegatum ■■■■■ Jor 8 jn.o St S. Europe ... L r.m Bot. cab. 666	
*368 odirum H. K. sweet-scented ■■■■■ Zor 6 jn.au Pa.R E. Indies 1683. C r.m R.mal. 9. 2	
2 cernicum ■■■■■ Zor 6 jn.au Pk E. Indies 1683. C r.m	
3 plenum ■■■■■ Zor 6 jn.au Pa.R E. Indies 1683. C r.m Bot reg. 74	
*369 flavescens Spin flavescent ■■■■■ Zor 6 jn.au Pa.Y. . . . 1816. C r.m	
R.Br. ALSTONIA. (Dr C. Alston, prof. of medicine, Edinburgh.) <i>Apocynete</i> . 2. - 4.	
poisonous H Q p r 6 my.jn W E. Indies 1823. C p.l	
school » r zor 8 mr.my W E. Indies 182*. C s.p R.mal. 1.45	
oleandraf&lia Lod. Cerium tinctorium Hort.	
Lo.C. CRTPTOLEPIS. (Kryptos, hidden, lepis, a scale.) <i>Apocfinect</i> . 1.-2.	
elegant tQor 6 Brazil -1824. C p.l	
ALVXIA. (Alyxis, anxiety; gloomy appearance.) <i>Apocpine*</i> . 1.-8.	
Richardson's ■■■■■ Jor 4 apjn W N. Holl. 1823. C p.l Bot.reg. ic. in	
musciLeaved ■■■■■ Jor 5 ap.jl W N. Holl. 1820. C s.p	
H'DIAKth. THENANDIA. (M. Thenard, a disting. French chemist.) <i>Apocpnec</i> . 1. - 2.	
In Kth. bundle.floweredi.Qor 10 ... B Mexico 1823. C p.l H.&B.3.240	
Br. PRESTOVIA. (C. Preston, M. D., a correspondent of Ray.) <i>Apocpnec</i> . 2.-6.	
woolly ft CD or 8 my.au W? Brazil 1820. C p.l	
smoothed S. or 8 my.au W? S. Amer. 182*. C p.l	
(Dr. W. Wright, a Scotch physician.) <i>Apocpnat</i> . 44.-6.	
dysenteric f. or 10 ... W E. Indies 1778. C r.m R.mal 1.47	
coccinea A.V. sra	
cerium cocclineum Wat.	
large-flowered ■■■■■ el 20 jn W Pi Indies 1820. C p.l Bot. reg. 911	
long-leaved ■■■■■ or 20 ... W E. Indies 1818. C r.m	
Ecnites. (Echis, a viper; smooth twining shoots.) <i>Apocpnec</i> . 22. - 60.	
umbelled % O or 15 jl W Jamaica 1733. C p.l Jac. am. 30.22	
twin-flowered k, n-n, o o 20 jl W W. Indies 1793. C p.l Jac am. SO. £1	
St. Domingo ■■■■■ or 10 jn.lau Y W. Indies 1820. C p.l Jac. ic. 53	
suberct ■■■■■ or 10 jn.au Y Jamaica '1759. C p.l Bot. mag. 1064	
twisted ■■■■■ CD or 10 jn.au Y Jamaica 1778. C p.l Jac. am. 33.27	
Heynes's ■■■■■ CD or 5 jn.au Y E. Indies 1818. C p.l	
twin-spined ■■■■■ CD or 5 jn.au Y C. G. H. 1795. C p.l	
succulent ■■■■■ Jcul 1 jln. Pk C. G. H. 1810. C s.p N. a. p 14.9.2	
tuberous ■■■■■ Br C. G. H. 1813. C s.p	
clove-leaved ■■■■■ Oor ft o ... Pa.Y R Indies 1812. C p.l Bot mag. 1919	
Roth antidyseentric ■■■■■ C. J or 6 ... Pk K Indies 1823. C p.l	
frutescent ■■■■■ 10 jn.jl E. Indies 1821. C p.l	
netted ■■■■■ 6 jn.lau Y E. Indies 1816. C p.l	
Malabar ■■■■■ G jn.lau R Malabar 1822. C p.l R.mal.9.12	
woolly ■■■■■ 3 jn.lau Y Guiana 1824. C p.l	
red-stemmed ■■■■■ 6 jn.lau Y Guiana 1824. C p.l	
long-flowercd ■■■■■ 6 my.au W? Brazil 1816. C p.l	
panicled ■■■■■ 10 jn.lau Y S. Amer. 1823. C p.l	
two-formed ■■■■■ 8 jl P.v Carolina 1806. C p.l	
Richard's ■■■■■ 3 jn.lau Y Guiana 1824. C p.l	
erguliria glabra W. ■■■■■ 10 ... - Trinidad 1826. C p.l	
(Vollo, to enclose; used for fences in Java.) <i>ApocQnea</i> . 1—2	
J. D. or 10 ... W E. Indies 1818. C p.l Itu. am. 5. 29'	
L? bundle-floweredi. O or 6 jn.au . . . Jamaica 182a C p.i	
monbndna Sivx.	

564. VARROWS L.	VARRONIA. (<i>M. T. Varro</i> , a celeb. Roman, lift years A. C.)	<i>Corditice</i> *.	11-.; U.
4531 angustifolia liesf.	narrow-leaved f O or 15	- W Santa Cruz 1808.	C 8.1
curass&vica Lam.	<i>Curia angustifolia</i> It.		
4532 aluifflia Horn.	Aider-leaved	— 1 ft SO	W Mc\co 1818. C co
4533 martincensis Jac.	Martinique	6	W Martinico 1795. C 8.1 Jac. am. 41.32
4534 Cordia martinicensis R. 8f S.	blistered	12	W W. Indies 1823. C p.l Jac. am. pic. 4J
4535 monoucrma Jac.	one-seeded	1 D 3	W Caracas 18.0. C p.l Jac. sc. 1.39
4536 Klopiisa Lam.	globose	m Qor 6	W Jamaica 1818. C si SI jam. 2.194. £
4537 lineata X.	lined	• OR 4	W W. Indies 1793. C s1 Br.jam. 13.2
4538 mirabiloides Jae.	Mirabilis-like	f I or 12	W Hispanio. 1798. C 8.1 Jac. aro. 41.33
4539 alba Desv.	White	30	W Trinidad 1820. C 8.1 Com. h. 1.80
4540 mōlijs Dttf	soft	4	W W. Indies 1821. C 8.1
4541 dichotoma R. OP. R. %S.	forked	4	Peru 1818. C s.p Fl. per. 2.145. a
565. PATAGO'NULA L.	PATAGONULA. (<i>Patagonia</i> , its native country.)	<i>Boraginece</i> . 1.	
4542 americanica I.	American	\$ D o r 20 jn.au	W S. Amer. 1732. C s.p Lam.il.96.
Cordia Patagonula If. K.			
566. EHRETT/il L.	EHRETIA. (D. G. Ehret, a celeb. German bot. draughtsman.)	<i>Cordiūcete</i> . 11.—28.	
4543 J/inifolia L.	Tinus-leaved	tm 30 jn.jl	W Jamaica 1734. C p.l Tr.ehr.4.25
4544 intembidis Herit.	internoded	I or 6	W Antilles 1819. K1 Her. st. 1.24
4545 aspera Rox.	rough-leaved	I or 10	W E. Indies 1795. p.l Rox. cor. 1.55
4546 senfta Rox.	serrated	I or 6	W E. Indies 1823. C
4547 laeVis Rox.	smooth	I or 12	W E. Indies 1823. Rox. cor. 1.56
4548 laxa Jac.	loose	I or 6	W Bourbon 1826. Jac. sc. 41
4549 microph. jrla Lam.	small-leaved	I or 10 jn.au	W E. Indies 1818. Pluk. ph. 31.1
4550 Auxif/ia Rox.	Box-leaved	• OR 8	W E. Indies 1823. Rox. cor. 1.57
4551 acuminata R. Br.	acuminate	I J or 15 jl	W N. Holl. 1820.
4552 d tibia Jac.	doubtful *	I or 20	W Jamaica 1825.
4553 divaricata Dec.	divaricate	I 15	W Havannah 1820. s.1 Dec. flfon. 4
567. BOURRE^R/i Gae.	BOURRERIA. (<i>Bourrer</i> , an apothecary at Nuremberg.)	<i>Cordiaceae</i> . 2.	
4554 succiflta Jac.	succulent	• tm 45	W W. Indies 1758. C s.l Jac. ob. 2. 2.26
Ehr'tia Bourrferia Lam.			
4555 exsiccata Jae.	dry-fruited	f • or 15	W W. Indies 1804. C s.l Jac. am. 173.17
568. ELLI'S/il L.	ELLISIA. (J. Ellis, F.R.S., an eminent English naturalist.)	<i>Hydrophyllae</i> . 1—	
4556 Nyctelea L.	Nyctelea	O cu \ jl.au	W Virginia 1755. S co Lam. il. 97
569. SERSALI'S/il R.Br.	SERSALISIA. (J. B. Sersalis, a Neapolitan ecclesiastic.)	<i>Sapbtete</i> . 1.—2.	
4557 sericea R. Br.	sttly-leaved	i j o r 6	W N. Holl. 1772. C p.l
Sideroxylon sericeum H. K.			
570. MANG'LLA Jus.	MANGLILLA. (Its name in PeruO	<i>MyrAne</i> *. 2. — II.	
4558 Milleria Ma Pers.	Miller's	j L J tm S O j n j l	W C. G.H. ... Ce o Bot mag. 1858
Sideroxylon mifte L.	A/ynine mitis Spr.		
4559 canariensis W.	Canary	\ tm 40	— Teneriffe 182a C s.p
Sideroxylon canariense IV.	3/ytscne canariensis Spr.		
* 571. ARDrSIA Sum.	AH nisi A.	(Ardis, a spear point; segments of corolla.)	<i>Myrsinea</i> . 2a— 32.
4560 acuminate IV.	acuminated	i or → jl.au..	Guiana 1803. C p.l Bot. mag. 1678
leacbrea guianensis Aub.			
4561 finifolia Swz.	Tinus-leaved	I I or 20 ll.au	R W. Indies 1820. C Sl. jam. 2.105
4562 paniculate Rox.	panicled	I or 12 jl.au	R E. Indies 1818. C Bot reg. 638
pyramidalis Roth			
4563 colorata Lk.	colored	I or 10 jl.au	R E. Indies 1816. C 8.1 Bot cab. 465
4564 corikcea Swz.	leathery	I or 7	S Antilles 1824. C 8.1
4565 «olanacea Rox.	Nightshade-like	I or 10 jn.s	R E. Indies 1798. Pi Bot. mag. 1677
4566 pyramidMis Cav.	pyramidal	I tm 25 jlau	R SantaCruzS18. s1 Bot. cab. 448
4567 mil mil is Vahl	humble	I or 6 jlau	K Ceylon 1820. s1 Bur. xcy. 103
4568 htoralis B. R.	sea-side	I 4 jlau	E. Indies 1809. C Bot. rep. 630
umbellate Rox.			
4569 lanceolate Rox.	lanceolate	*d) 6 jl.au	K E. Indies 1820. C p.l Bot rep. 49
4570 elegans Andr.	elegant	10 jl.au	R E. Indies ikOD. C p.l Bot rep. 49
crenata Rox.			
4571 punctate L.	dotted	I 10 jnau	W China 1823. C s.1 Bot reg. 887
4572 lenticigintaa Ker	speckled	• or 6 year	W China 1814. C B.1 Bot reg. 533
crenulata B. C crenata B. M.			
4573 macrocarpa I.al.	long-fruited	I 5 ...	F Nepal 1824. C
4574 complinata IVal.	levelled	I 6 ...	Pk Pinang 1824. C p.l
4575 thyrsiflora D. Don	thyrsiflwd	I 5 ...	Pk Nepal 1824. C p.l
4576 excelsa H. K.	tall	I J 80 ji.au	Pk Madeira 1784. il Gae. fr. 1.77
Hebrdencia R. & S.			
4577 canariensis Zey.	Canary	I 10 jl au	R Canaries 1820. C
4578 crenulata Ven.	crenulated	I 10 jns	R W. Indies 1809. C
4579 serrulata Swx.	saw-leaved	I 3 jlau	R W. Indies 1820. C
4580 lateriflora IF.	side-flowering	I 6 ..	W. Indies 1793. C
4581 canaliculata Lo. C.	channelled	I 6 jl.au 1821. C
4582 pubescens Lo. C.	pubescent	I or 6 jl.au 1820. C Bot cab. 1083
572. EMBE^SLIA Bnn.	EMBELIA.	(/Embilla, its name in Ceylon.)	<i>Myrstnea</i> . 1.—6.
4583 robusta Rox.	robust	I or 20 ...	W.a E. Indies 1823. C p.l
573. ARDUPNA L.	ARDUINA.	(P. Arduini, curator of the econom. garden of Padua.)	<i>Apocynace</i> . I. rr
4584 bispinosa L.	\ two-spined	i J cu 2 mr.au	W C. G. H. 1760. C p.l But. cab. 387
574. STRYCHNOS L.	STRYCHNOS.	The Greek name of the sola num.	<i>Apocineis</i> . 6. —11*
4585 Nux-vdmica L.	Vomit-nut	I P G.W	E. Indies 1778. S p.l Roxcor. 1.*
4586 coluurtna L.	snake	I Im	E. Indies 1820. S li mal. 7. J
4587 potatōrum L.	drinkers'	I m	E. Indies 1794. P Rox. cor. 1.
4588 madagascarenensis Thou.	Madagascar	> n m	Madagas. 1823. s.p
4589 spinosa Lam.	spiny	I K	Madagas. 1818. C
4590 axillaris T'nib.	axillary	I K	E. Indies 1824. P
* 575. FAGRJ&A Thun.		CJ. T. Fagretrus, M.D., a Wend of Thunberg.)	<i>Apocfi*f*</i> [W]
4591 zelánica Thun.	Ceylon	T Q or Vi ...	W Ceylon I*16 C *v.l Wti 1782,
Willuirhbñia zelanic* Srr.			

ORDKR I.

4657 edulis B. M.	eatable	£ EJcul	2 Jl.au	Y	S. Amer.	1773.	C co	Bot. mag. 1068
4658 chenopodiifolia Lam.	Goosefoot-lvd	j(A un	2 jl.au	Y	Peru	17 ⁸ *	S s.l	
4659 tuberosa IV. en.	tuberous	A un	2 jl.au	W		1815.	I e.p	
4660 pubescens L.	pubescent	O un	2 jl.au	Y	America	1640.	S s.l	Feu. ob. 3.1
4661 micranthus Zar.	small-flowered	O un	26 jl.au	Y	America	1826.	S co	
4662 nodosa Law*.	knotty	O '» Zi	jl.au	Y	America	181ft	S co	
4663 angulata L.	an <i>E</i> Uar-branched	O un	2 jns	W	India	17.32.	S s.l	Di. el. 13.12.12
4664 phludelphica Mr.	Philadelphian	O un	1 jl.au	Y	Philadelph.	1800.	S co	
4665 atriplicifolia Jac.	Atriplex-leaved	O un	2 jl.au	Y	£. Indies	1820.	S co	Jac. fr. 85. a
4666 chenopodiifolia HI	Roth's	O un	1 jl.au	Y	E. Indies	1826.	S co	
4667 uquata Jirie.	even	O un	IJjn.jl	Y		182c.	S co	
4668 barbadensis Jac.	Birbadoes,,	O un	2 jl.au	Pa. Y	W. Indies	1798.	S s.l	Jac. ic. 1. 89
4669 minima X.	smallest	O un	Ujl.ju	Pa.Y	E. Indies	1759.	S s.l	ILmal. 10.71
4670 pruinosa L.	frosty	O un	1 jl.au	I'a.Y	America	1726.	S s.l	Di. el. 10. 9.9
4671 prostrata Herit.	trailing	O un	1 jl.au	L.H	Peru	178 ¹ .	S s.p	Bot rep. 75
4672 g g A f.	smaU-Howered	O un	Hjl.ju	N. Holl.	1820.	S s.p		
4673 g g A f.	L	O un	1 jl.au	N.	Spain	1824.	S s.p	
4674 dubia L.	doubtful	O un	2 jl.au	F	Brazil	1821.	S s.p	
4675 feedidissima Lag.	mobt letid	O un	H jl.au	N. Spain	1820.	S s.p		

fl87. SA'RACIL* FLper.	SARACHA.	(J. Saracha, a Spanish botanist.)	Soltate*.	2.—3.
4675 procumbens fl per.	procumbent	Jc O or 3 jn.jl	Pa.Y Peru	IX22. D co Fl. per. 2. 180

A'tropa procumbens Cao.

4677 umbellata Jac.

588. LY'CIUM L.	BUX-TUBVB.	(Lucia in Asia Minor, its original country.)	Sotunee.	18.—28.
4678 afraum L.	African	or 10 jn.jl	V	C. ii. H. 1712. I p.l
4679 tenui Lk.	slender	or	T	C. G. H. 1819. I p.l
4580 rigidum Thun.	rib <ul style="list-style-type: none">ul	or	T	V. G. H. 1793. I p.l
4681 tetraiidrum Thun.	tetrandrous	or	T	C. G. H. 1810. I p.i
4G2 carnosum Dull.	fleshy	HK	T	C. G. H. 1795. I p.i
4683 rithemcuin AJur.	small-leaved	or	T	C. G. H. 1795. I p.i
am buñbarum /.	Russian	L	T	Siberia 1804. I p.l
4684 turbinatum Per*.	Barbary	or	V	Barbary 1606. I co
4685 europium L.	European	EL J or	V	Den. br. 9
4689 lanceolatum Pair.	apeur-leaved	or	V	Shaw bar £49
4690 chipense Mil.	Chinese	EL J or	V	Duh. no. 119. 31
4691 cinerquin Thun.	cinerous	or	V	Mic. gen. 105.1
4692 Trewianutu Duh.	Trew's	or	V	Duh. no. 32
4693 hürnduin Thun.	very prickly	I L J or	V	Den. br. 8
4J94 boerhavu/ ^v /ium L.	Boerliaavia-lvd	tl gr	V	Duh. no. 30
4695 carolininuni Pit.	Carolina	or	V	Her. st. 45.23

589. LYCIOSERISSA R 8/S. LVCIOSERISSA. (Lycium, lye, serissa, ser; serissa-like lycium.) Sotine*. 1.

4695 Scapensis A 4E. Cape m LJ un 2 jn.au G C. G. H. 1820. C p.l

590. LYCOPERSICUM Ton. LVCOPERSICUM. (Lykos, wolf, persikon, peach j aphrodisiac.) Sotinea. 9. — ltt

4697 pimpiliifidhum/u/OT/	l'impincilla-lvd	O clt	3 my.jl	G Peru	... S r.m
Solānuin pimpinellifiliun L.					
4698 regulare Dunal	reRular-leaved	O clt	3 my.jl	G	... S r.m Dun. so. 2.81
4699 peruviaium Dunal	Peruvian	^ iAJ clt	3 my.jn	Y Peru	1823. D co Feu. ob. 3. 25
4700 Humboldti W.	Humboldt's	O clt	3 jl.s	Y	S. Amer. 1822. S co W. h.b.27
4701 pynf6rme Dutial	pear-sha ied	O clt	3 jl.s	Y	1823. S co Dun. so. 26
4702 cerasiforme Dunal	cherry,bhaped	O clt	3 jl.s	G	1800. S r.m Jac. \ in. 1.11
5blānum Pseudo-lycopersicinu Jac.					
21ātqum ycWow,fruited		O clt	3 jn.s	G	Peru 1800. S r.m
4703 esculentum Dunal	eatable apple	O clt	3 jl.s	G	S. Amer. 1590. S r.m Ru. am. 5.1*34.1
Solātum Lycopersicum L.					
1 erythrocarpum	red-fruited **	O clt	3 jl.s	G	S. Amer. 1596. S r.m
2 chrybOi-arpum	yellow-fruited	O clt	3 jl.s	G	S. Amer. 1596. S r.m
3 leucocarpum	white-fruited	O fit	3 jl.s	G	S. Amer. 1596. S r.m
4704 proctihibens Dunal	prucumbent	-* O clt	1 jl.s	Crea	S. Amer. 1700. S r.m
4705 commutatum Spr.	changed	O clt	3 jl.s	Y	S. Amer. 1818. S r.m

* 591. SOLA^NUM L. NK.IITSHADE*. (Solor, to comfort; soothes by stupefying.) Soltince. 143. — 370.

4706 tuberbiuum L.	tuberous	A A ag	2 jn au	W	Peru 15-17. R r.m Bau. pr. 89 89
2 Commsoni Poir.	Commson's	A A clt	2 my.o	W	S. Amer. 1822. R co Hort.tr. 5.10
4707 appcticulatūin//,\$7.appemli<uljeu*1 rjJ or	3 jn.ju	W	Mexico Ib23. R co Dun. bo. 2. 84		
4708 Seaforthianum Aiulr.	Seaiorth's	JL O or	20 jl.s	Pk	Barbadoes1804. C l.p Bot. rep. 504
4709 ^etacum Cav.	HeuU/eavcd	4 jl.jl	Pk	S. Amer. 1803. C l.p Bot. rep. 511	
4710 muricatum // K	prickly	Qor	3 jl.jl	V	Peru 1785. C l.p FFau ob 772. 15
4711 diverbiflum II. ^B.	various-leaved	H ! lun	3 jn.jl	Pk	Caraccas 1825. C r.m Dun. so.2.88
4712 oliganthum Lk.	few-flowered	1 i lor	3 .	W	1824. C co
4713 laciniatum R. Br.	cut-leaved	LIL Jor	3 jl.au	V	N. Holl. 1772. S s.p Bot. mag. 349
2 herbaceum	herbaceous	£ El	3 jn.au	V	V. Di. L. 1772. R si
4714 fuercifolium L.	Oak-leayed	A or	in jl	V	Peru 1787. C r.in Feu. ob. 772.15
4715 radkans L.	jootoing		jl.au	P	Peru 1771. I s.p 1*1 dec. 1.10
4716 corymbosum Jac.	fcorymbose		jl.au	V	Peru 1786. D co Jac. ic. 1.40
4717 Dulcamara L.	Bitter-sweet	I P	jl.s	V	Britain hed. C si Eng. bot.5*5
2 álba	white-flowered	P	jl.s	W	Britain hed. C co
3 vanegata	variegated	P	jl.s	V,w	Britain hed. C co
4718 angulatum RSJS.	angular	Dot	jl.jl	W	Lima 1823. C pi Dun.so. 2.95.1
quitense Lam.					
4719 pginatum Cav.	dwarfish	O un	B	Bonaria 1819. S co Cav. ic. 5 439. *	
4720 Teghre Aub.	Tcgoré	*D or	2 jn.jl	B	Guiana 1823. C p 1 Aub. gui. 1 84
4721 htoriile Lk.	sea-side		4 my.jl	W	France 1819 C co
4722 macrncarpum L.	long-fruited		1 my.s	B	Peru 175». C sp Mil. ic. 2.294
4723 calycinum Dunal	large-calyxed		1 jl.jl	B	Mexico 1820. S co Dun. so. i'0
4724 pentadactylw G.Don	five-fingered	» D c u	3 jn.au	B	Trinidad 1818. C r.m
4725 racenuflurum Dunal	cluster-riwd	£ Ol or	jl.s	P.B. S. Amer. 1818.	
4726 iethiopicum /.	Ktuopidii	O or	Mils	W Ethiopia C p.l Jac.sc. 3.33	
4727 Zuccagmiftu/nDunal	Zuccigm's	O or	U J>jl	W	S Ip Ja'l vin. J.*
					S <6 Dim. M. II

4788 triquetrum Cav.	three-cornered	• or	2 }njl	W	N. Spain	1820.	C p.l	Cav. ic. 3.259
4729 Pseudo-ca>sicuin L.	Bart capsicum	■ or	4 jn.s	W	Madeira	1596.	S r.m	Sab. rom. 59
4730 nodiflorum Jac.	joint-flowering	T un	10 jnjl	W	I. France	1822.	S co	Jac. ic. 2.326
4731 Dilleiu* Schult.	Dilleni's	O un	2 jn.a	W.	.	1818.	S co	Di. el. 275.355
4732 guineinse Lam	Guinea	O p	4 jn.s	G	Guinea	1817.	S s1	Di. el. 274.354
4733 mclanocerasumff.fn.	black cherry	P	2 jn.s	W	Virginia	1820.	S ■	Di. el. 275.356
4734 nigrum L.	Wack-berricd	■	3 jn.s	W	Britain	nib.	S s1	Eng. bot. 5G6
4735 pterocaule Dunal	wing-stemmed	O un	2 jn.s	V	America	1H20.	S co	Di. el. 275.256
4736 rubrum Mil.	red-berried	O un	2 jnjl	W	S. Amer.	1817.	S co	
4737 flavum Kit.	yellow-berried	O un	2 jnjl	W	Brazil	1817.	S co	
4738 olerdequin Rich.	jiotherb	O un	i jn.au	W	S. Europe	1816.	S co	Pis. lib. 4. 3
4709 judaicum Shul.	Jewish	O un	1 jn.au	Pa.V	E. Indies	1800.	S co	
4740 triangulare Lam.	triangular	O un	4 jn.au	W	S. Europe	IH23.	S co	
4741 mimatum Bernh.	red-beet ricd	O un	4 jnjl	W	S. Europe	1823.	S co	
4742 humile Bernh.	humble	O un	1 jnjl	W	S. Europe	1823.	S co	
4743 villosum IV. en.	villous	O un	5 jn.s	W	Barbades	1820.	S s1	Di. cl. 274.353
4744 hirsutum Dunal	hairy	O un	2 jn.s	W	Egypt	1817.	S co	
4745 Kitaibel'a	Kitaibel'a	O un	1 jn.s	W	Pannonia	1822.	S co	
4746 Be'ssen Wein.	Besscr's	O un	1 jn.s	W	America	1820.	S co	
4747 patulum Pers.	spreading	O un	4 jn.s	V	India	1818.	S 81	Di. el. 275.355
4748 suffruticosum&ft0iu.	half shrubby	O or	4 my.s	W	Barbary	1804.	C l.p	
4749 crispipi F. per.	curled	O or	18 jn.jl	W	Chile	1824.	C co	Fl. per. 2.158
4750 longiflorum Vahl	long-flowered	J Jor	3 jn.au	V	Cayenne	1823.	C p.l	Dun. so. 9
longifolium Dunal								
4751 bomblinc Jac.	Bombay	I-Jor	12 jn.jl	W	Mexico	1822.	C co	Dun. so. 101
4752 /aurifl.ijum Dunal	Laurel-leaved	CD or	8 jnjl	... S. Amer.	1820	C p.l	Dun. so. 8	
4753 Cervantes' Lag.	Cervantes's	C Jr	4 myjn	W	Mexico	1818.	C co	
4754 verbasifolium L.	Mullein-leaved	■ or	7 jnjl	W	W. Indies	1749.	S p.l	Jnc. vin. 1.13
4755 puhecsen IV.	pubescent	■ or	5 jnjl	P	K. Indies	1820.	C s1	W. ph. 1.3
4756 platamfuliu Sims	Plane-leaved	■ un	3 JI	B	S. Amer.	1813.	S co	Bot mag. 1618
4707 argenteum Dunal	silvery	I lor	jnjl	...	Brazil	1824.	C pa	Dun. so. 3!)
4758 auric ul< H. K.	ear-leaved	■ or	V	Madagascar	1773.	S p.l	Scop. d. 3. H	
4759 diph^lluin L.	two-leaved	■ or	JnV	W	W. Indies	1699.	S s1	Jac. ic. 2.322
4760 triste Jac.	sad	■ or	6 jnjl	V	W. Indies	IHSu	C p.l	Jac. am. pic. 49
4761 arboreum H. S/B.	tree	f or	40 jnjl	W	Cumana	1819.	C p.l	Dun. so. 2.194
4762 havanense L.	Havannah	* or	5 Jlau	B	W. Indies	1793.	C co	Jac. am. 49.35
4763 aggregatum Jac.	clustered	* or	6 jnjl	P C.	G. H.	1821.	C p.s	Jac. ic. H. 323
4764 yciooides W.	Lycium-like	* or	4 myjn	Pa.B	Peru	1791.	C s.p	Jac. ic. 1. 46
4765 uniflorum L.	one-flowered	* or	3 myjn	B	N. Spain	1820.	C co	
4766 neglectum Dunal	neglected	* or	4 jnjl	V	W. Indies	18-J4.	C p.a	PI. am. 245.4
4767 monanthum It S(S.	one-flowered	* or	2 jnjl	B	N. Spain	1818.	C p.l	
uniflorum Lag.								
4768 tkgax J.ic.	fleeting	• or	5 myjl	W	Caracas	1816.	C pl	Jac. ic. 324
4769 stellatum Jac.	starred	• I-Jor	6 jnjl	B	...	1822.	C co	Jac. ic. 1.325
4770 virgatum Ijum.	twiggy	• I-Jor	4 jls	V	W. Indies	1820.	C p.s	Dun. so. 40
4771 scandens L fit.	climbing	I aor	10 jlau	...	Surinam	1820.	C p.l	
4772 conaceum H+k.	coriaceous	• or	4 jlau	P.B.	...	1820.	C p.l	Bot. mag. 2708
4773 lenthum Cav	pliant	LCD or	5 jn.s	P.V	N. Spain	1823.	C p.l	Cav. ic. 4. %8
4774 elagnifolium Cav.	Oleaster-lvd	• or	6 jnjl	B	Chile	182a	C co	Cav. ic. 3.2K!
4775 stelliferum R. Br.	star-bearing	»L or	3 jn.au	Pa.P	N. Holl.	182a	C p.l	Sm. ex. bot. 2.88
4776 racemum L.	racemose	• or	4 jlau	W	W. Indies	1781.	C co	Jac. am. 50.3b'
4777 igneum L.	firc-spined	• or	3 mm	W	S. Amer.	1714.	C s.p	Jac. vin. 1.14
4778 atum Dunal	half-armed	• or	6 myjn	W	...	1820.	C co	
4779 cuneifolium Dunal	Bahama	i-jor	6 jnjl	V	Bahama	1732.	S p.l	Di. el. 271.350
4780 mexicanum M. SS	netted	• or	3 jnjl	W	W. Indies	1820.	C p.l	Dun. so. 48
4781 jamalencense Stiff.	shaggy	• or	3 jnjl	W	Trinidad	1821.	C p.l	Vahl ic. 3.21
4782 brasiliense Dunal	tall	• or	■ Jnjl	W	...	1820.	C p.l	
4783 lanceolatum Jac.	soft	• or	■ Jnjl	B	Trinidad	1818.	C s1	Dun. so. 180
4784 cuneifolium Dunal	hybrid	* i or	2 Jnjl	P	C. G. H.	1662.	C p.l	Boc sic 8.5
4785 cuneifolium Dunal	starlet	* or	2 Jnjl	B	Guinea	1815.	C p.l	Jac. vin. 2.11J
4786 cuneifolium Dunal	Cestrum-lvd	all or	2 Jnjl	W	...	1810.	C p.l	Jac. ic. 43
4787 jamaicensis Stiff.	Jamaica	41 • or	4 jn.jl	W	Jamaica	1818.	C p.l	
4788 heterochonchum Z) i/n/i	variable-haired	fl or	2 jnjl	W	S. Amer.	1824.	C p.l	Dun so. 20
4789 cuneifolium Dunal	wedge-leaved	* or	2 jnjl	W	Guiana	ISIS.	C p.l	Sl. jam. 141.3
4790 mexicanum M. SS	Mexican	tk I or	3 jnjl	W	Mexico	1825.	C p.l	
4791 mexicanum M. SS	twining	% or	8 jnjl	V	W. Indies	1823.	C p.l	Pluk. m. 4.32
4792 brasiliense Dunal	Brazilian	m or	2 jnjl	B	Brazil	1830.	C p.l	Pl. am. 454. 4
4793 lanceolatum Jac.	lance-leaved	A • or	10 jlau	W	W. Indies	1816.	C co	Jac. ic. 2.239
4794 umbrussum Dunal	shady	• or	3 jnjl	W	Trinidad	1825.	C p.l	Dun. so. 126
4795 ubiginosum Vahl	rusty	m CD or	3 jnjl	W	Guiana	1821.	C p.l	Vahl ic. 13
4796 opnumnum W.	peaked	• il I or	2 4 jnjl	Pa.B	S. Europe	1818.	C p.l	Dun. so. 16.2
4797 bonariense L.	clammy	ft or	4 jnjl	W	...	1810.	C p.l	Dun. so. 2.54.
4798 opnumnum W.	Buenos Ayres	ft \ or	10 jn.s	W	B. Ayres	1727.	C II	DieL364.272.351
4799 celatum Cav.	autumnal	CD or	A jn.jl	B	...	1820.	C p.l	Dun. so. 16
4800 ganteum Jac.	star-like	I or	2 jnjl	W	W. Indies	1818.	C p.l	
4801 vnum Sivz.	half-unarmed	ft or	7 jlau	B	W. Indies	1752.	C Iq	Jac. am. 40. 3
4802 >ugineum Jac.	lanceolate	I or	7 jnjl	Pa.B	Mexico	1800.	C s.l	Bot. mag. 2173
4803 tonaccum B. R.	giant	f \ or	15 jnjl	V C	G. H.	17S<	C s.p	Bot. mag. 1921
4804 ovigorum Dunal	grim	ft or	5 jlau	Pa.B	W. Indies	1816.	C p.i	Pluk. al. S55.3.
4805 Mffongena L.	ferruginous	ft i or	4 jlau	Pa.B	W.	1816.	C p.i	Jac. M. a. 134
4806 violaceum	saponaceous	ft i or	4 jlau	W	Chile	1825.	S p.i	Bot. reg. 2097
4807 insanum L.	egg-bearer	iQ cul	4 jnjl	B	Africa, &c. l597.	S lp	Pluk. ph. 2G6.2	
4808 esculetum Dunal	r\o et-frui/ed	iQ cul	5 jnjl	B	Tropics	1597.	S r.m	
4809 Br6wn^n Dunal	white-fruited	O cul	5 jnjl	B	Tropics	1597.	S r.m	
4810 xiolaceum R. Br.	je\ow-fruited	iQ cul	5 jnjl	B	Tropics	1597.	S r.m	
4811 meum L.	red-fruited	iQ cul	5 jnjl	B	Tropics	1597.	S Km	
4812 W09 insanum L.		iQ cul	5 au.s	B	E Indies	1815.	S l.p	Pluk. aL 22a 3
4813	Brown's	■ L J or	3 Jn au	V	N. Holl.	1820.	C p.l	
4814	Sodom	■	■	■	Africa	■	■ r.m	Her. lug. 575
4815	hoary	■	■	■	Ceylon	JSS-	■ p.l	Scop. d. 1.1
4816	Indian	■	■	■	India	1732.	S p.l	Di. el. 270.549

4814 undatum <i>Lam.</i>	waved	111	or	3	jnjl	V	Malabar	1820.	C	s.p	R.mal.2.37
4815 <i>Anequivi</i> <i>Lain.</i>	Madag. potato	A	GS	un	3	my.s	W	India	1818.	L	r.m
4816 <i>cgagulans</i> <i>Forsk.</i>	curdling			3	jil	W	Arabia F.	1802.	C	s.p	Jac. sc. 4. 469
4817 <i>sanctum</i> <i>Dunal</i>	holy		L	or	3	jnjl	P	Egypt	1818.		s.p
4818 <i>marginatum</i> <i>IV.</i>	white-edged		L	or	4	jns	P	Africa	1775.		s.p
4819 <i>campachinense</i> <i>W.</i>	Campeachy		L	un	2	jil	V	America	1732.		s.p
4800 <i>Trdnym</i> <i>Poir.</i>	Trongum	RII	un		jnjl	W	E. Indies	1816.		co	Ru. aro. 5.86 1
4821 <i>congense</i> <i>Lk.</i>	Congo		Q	or	3	jnjl	B	Guinea	1821.		co
4822 <i>fuscatum</i> <i>L.</i>	dusky		Q	or	1	jnjl	S	S. Amer.	1817.		Jac. ic. 1.42
4823 <i>acutealatumsum</i> <i>Jac.</i>	prickliest		Q	or	3	apjl	Pa.B	S. Amer.	1816.		Jac. ic. 1.41
4824 <i>polyacanthum</i> <i>Lam.</i>	many-spined		•	or	4	jns	R	W. Indies	1821.		Cav. ic. 3.236
	pamfliftrum <i>Cav.</i>										
4825 <i>mynacanthum</i> <i>Dunal</i>	inany-spined		•	or	3	jl.au	P		1822.	C	p.l
4826 <i>mamin6sum</i> <i>L.</i>	teated		Oor	4	jl.au	Pa.B	W. Indies	1699.	S	s.p	Pluk. al. '226.1
4827 <i>stramonifolium</i> <i>Jac.</i>	Stramonium-lvd*		•	or	6	jns	P	W. Indies	1778.	C	s.p
4828 <i>flavescens</i> <i>Dunal</i>	flavescent		■	or	3	jnjl	B	Trinidad	1826.	C	s.l
4829 <i>ftrox</i> <i>L.</i>	fl'ree		jg	EJun	2	a.u.s	P	E. Indies	1795.	C	l.p
4830 <i>cainqi»nulatum</i> «J?r.	campanulate		Q	or	1	jns	W	N. Holl.	1819.	S	p.l
4831 <i>armatum</i> <i>R. Br.</i>	armed		•	Jor	2	a.u.a	•	N. Holl.	1819.	S	p.l
4832 <i>pungentium</i> <i>R. Br.</i>	pungent		Q	or	1	a.u.s	•	N. Holl.	1823.	S	co
4833 <i>cinfreum</i> <i>R. Br.</i>	ashy		•	Jor	1	a.u.s	P	N. Holl.	1823.	S	p.l
4824 <i>Mildri</i> <i>Jac.</i>	Miller's		•	Jor	3	jl.au	W	C. G. H.	1762.	C	61
4835 <i>rigescens</i> <i>Jac.</i>	rigescents		•	Jor	ljnjl	V	C. G. H.	1823.	C	81	
4836 <i>ciliatum</i> <i>Lam.</i>	ciliated		•	O un	2	jnjl	W	—	1820.	C	81
4837 <i>trilobatum</i> <i>L.</i>	threc-lobed		•	or	12	au	V	India	1759.	C	s.p
4838 <i>acetosifolium</i> <i>Lam.</i>	Sorrel-leaved	I	cu	1	jlau	W	E. Indies	1759.	S	p.l	Bur. in. 57.22.2
4839 <i>carolinense</i> <i>L.</i>	Carolina	iQ	un	2	jl.s	Pa.B	Carolina	1732.	S	p.l	Bur. in. 22.2
4840 <i>vol&ceum</i> <i>Jac.</i>	violet	•	or	4	jnjl	B	E. Indies	1817.	C	s.p	Jac. ic. 2.331
4841 /Vracanth'a 5m.	Ftre-thom	•	or	4	a.u.s	P	Madagascar	1789.	C	r.m	Ex. bot. 2.64
2 inermis	unarmed	•	or	4	au.o	P	Madagascar	1789.	C	r.m	Bot mag. 2547
4842 <i>spinosisimum</i> <i>Lo. C.</i>	spiniest	•	or	3	jnjl	P	•	1820.	C	p.l	
4843 <i>melanoxylum</i> <i>Lk.</i>	black-wooded	•	or	3	jnjl	V	•	1821.	C	p.l	
4844 <i>virginianum</i> <i>Pers.</i>	Virginian	O	un	1	my.au	V	Virginia	1662.	S	s.p	Di. el. 267.346
4845 <i>Jacquin'i</i> <i>W.</i>	Jacquin's	fm	un	2	s.n	B	E. Indies	1804.	S	s.p	Jac. ic. 2.332
4846 <i>xanthocarpon</i> <i>S.Sf/W.</i>	yellow-fruited	O	un	1	ijnjl	B	Ethiopia	1824.	S	co	Sc. han. 1.2
4847 <i>Balbis</i> <i>Dunal</i>	Balbis's	: L	Jor	4	aps	B	S. Amer.	1816.	C	co	Bot reg. 140
4848 <i>tectum</i> <i>Pers.</i>	covered	: L	Jor	3	apa	Y	Mexico	1824.	C	co	Cav. ic. 4.309
592. NYCTE* RIUM <i>Fen.</i>	NYCTERIUH.	(Nycteros, nocturnal;					time of flowering.)	<i>Solanace.</i> 7.—S.			
4840 <i>cordifolium</i> <i>Ven.</i>	heart-leaved	• LJ or	2	ap.my	P		CaiLlsL	1779.	C	co	Ven. maL 85
Solanum vespertilio <i>H. K.</i>											
4850 <i>amazonium</i> <i>B. M.</i>	Amazonian	• • or	3	jn.au	P	Mexico	1800.	C	■	Bot reg. 71	
Lamberts* Swt Solanum nmaztiauumA 2f.											
4851 <i>lobatum</i> <i>Swt.</i>	lobed	O	or	2	jl.au	Y	Louisiana	1813.	S	co	Ph. am. 8.7
Solanum heterandrum <i>Ph.</i>											
4854 <i>Fontanesianum</i> <i>Dunal</i>	Desfontaine's	O	or	2	jl.s	Y	Brazil	1813.	S	co	Bot reg. 177
4853 <i>corn'utum</i> <i>Lk.</i>	horned	• • or	1	jn.au	Y	.Mexico	1823.	S	SD	An. mu. 3.9	
4854 <i>rostratum</i> <i>Lk.</i>	beaked	O	or	2	jn.au	Y	Mexico	1823.	S	s.p	Dun. so. 24
4855 <i>heterodoxum</i> <i>Dunal</i>	variable	O	or	2	jn.au	B	Mexico	1820.	S	s.p	Dun. so. 25
593. CA'PSICUM <i>L.</i>	CAPSICUM.	(<i>Kapto</i> , to bite; pungent qualities.)					<i>Sol&neat.</i> 23.-28.				
4856 <i>Jinnuum</i> <i>L.</i>	annual	O	cul	1	jnjl	■	S	r.m	Kno. th.2.6		
4857 <i>longum</i> <i>Dec.</i>	long,fruitcd	Q	cul			■	S	r.m	Fuchs733		
4858 <i>cordiforme</i> <i>Mil.</i>	heart-shaped	O	cul		jnjl	W	India	1548!	S	r.m	
4859 <i>tetragramum</i> <i>Mil.</i>	four-angled	O	cul		jnjl	W	India	—	S	r.m	
4860 <i>angulfussum</i> <i>Mil.</i>	angular/rufuted	O	cul		jnjl	W	India	—	S	r.m	
4861 <i>sphaericum</i> <i>JV.cn.</i>	g\6b\ af-fruitedn.	• cul	2	ap.jl	W	•	1807.	C	r.m		
4862 <i>ovatum</i> <i>Dec.</i>	ov&te-fruited	it	• cul	3	jns	W	•	1824.	S	r.m	
4863 <i>cönicum</i> <i>Lam.</i>	conical	O	cul	1	jnjl	W	Guiana	1820.	S	r.m	
4864 <i>pendulum</i> <i>IV. en.</i>	pendulous	FL	• cul	2	ap.jl	W	•	1804.	C	r.m	
4865 <i>fluteum</i> <i>Lam.</i>	yeUow-fruited	«L	• cul	2	jn.au	E. Indies	1820.	S	r.ra		
4866 gr6ssum <i>W.</i>	large	n.Ocul				W	India	1759.	S	r.m	Bes. cys. 1.11.1
2 globusum	g\6b\ fruited	M	clct			W	E. Indies	...	S	r.m	
3 luteum	ycMovr-fruited	n-Qclt				W	E. Indies	...	S	r.m	
4867 conoldes <i>Mil.</i>	cone-like	r~l				W	India	1750.	C	r.m	
4868 pyramidale <i>Mil.</i>	pyramidal	■				W	Egypt	1750.	C	r.m	
4869 <i>cerasifirme</i> <i>IV.</i>	cherry-shaped	• 9				Pa.Y	W.Indies	1739.	S	r.m	
4870 <i>globiferum</i> <i>Mey.</i>	globe-bearer	■				W	Guiana	1824.	S	r.m	
4871 <i>MiWcri</i> <i>R. & S.</i>	Miller's	■				W	W. Indies	...	S	r.m	
4872 <i>fructescens</i> <i>L.</i>	frutescent	■				Pa.Y	India	1656.	C	r.m	Ku.a.M.o.88
2 <i>toruliflum</i> <i>Jac.</i>	sub-twisted	■				W	E. Indies	1820.	S	r.m	
4873 <i>bicolor</i> <i>Jac.</i>	two-colored	■				W	W. Indies	1804.	C	r.m	Bot mag. 1835
4874 <i>baccatum</i> <i>L.</i>	berried	■				W	•	1731.	C	r.m	SL. jam. 1.146.2
4875 <i>sinense</i> <i>IV.</i>	Chinese	■				W	China	1807.	C	r.m	Jac. vin. 3.67
4876 <i>microrarpon</i> <i>Dec.</i>	small-fruited	■				W	•	1820.	C	r.m	
4877 <i>cerasiflrum</i> <i>Lk.</i>	cherry-flwd	■				W	•	1823.	C	r.m	
4878 <i>micranthus</i> <i>Lk.</i>	smalUflowered	n. cul	3	myjn	W	Braill	1824.	C	r.m		
594. LEE\4 L.	LEEA.	(<i>J. Lee</i> , father, son, and grandson, esteemed bot and cult)					<i>Melihcece.</i> 6.—				
■ W	Eldcr-leaved	■				Y	E. Indies	1790.			Cav.dis.7.
4880 <i>squaU</i> <i>L.</i>	even	■				G	E. Indies	1777.	Lp		
4881 <i>crispia</i> <i>L.</i>	curled	■				W	C. G. H.	1767.	Lp	Bot rep. 355	
4882 <i>macrophylla</i> <i>Box.</i>	long-leaved	■				W	K Indies	1806.			
4883 <i>hirta</i> <i>Rox.</i>	hairy	■				G	E. Indies	182a			
4884 <i>robusta</i> <i>Rox.</i>	robust	■				O	E. Indies	182a			
* 595. SPERM ADI'CYON <i>Rox.</i>	sweet-scented	• • or	4	o	W	E. Indies	ISia	C	lp	Bot reg. 348	
596. DENTE'LLA <i>Font.</i>	DENTELLA.	(Dim. of <i>dens</i> , a tooth; corolla)									
4886 rtpens For ^z .	creeping	• • or	jjl		W	N. HoU.	1802.	S	co	Lam.il. 118	
Oldenfindio reiens i.											
597. MACROCNETMUM <i>L.</i>	MACROCNEUM.	(<i>Makros</i> , long, <i>kneme</i> , a leg; fl. stalks)					<i>Rubihece</i> *. S. — M				
4887 <i>jamaicense</i> Z.,	Jamaica	\$ D or	14	...	W	Jamaica	180a	C	p.l	Swa.ob.6&3.	
4888 strictum <i>Rox.</i>	strict	• • « :	10	...	W	E. Indies	1804.		p.l		
4889 tinctorum <i>H.tfb.</i>	dying	ICII or	30	s.o	B	Trinidad	1820.		p.l		

- | | | | |
|-----------------------------------|---|--|--|
| 598. EXOSTE'MM A Rich. | UXOSTEMMA (f.J., without, stemtn,a crown.) | RuhthauB. | 4. — lti. |
| 4890c. iribajxumlf. | Caribscan | t or 20 jns. | W. Indies 1780. C j.p Bot. rep. 481 |
| 4891 flonbundum W. | many-flowered | J · tm 40 | W. Indies 1794. Lamb. ci. 27.7 |
| 4882 longiflbrum R. & S. | long-flowered | f · or JO jnjl | Caracass 1890. p. Lamb. ci. 12 |
| 48113 brachycarpum IL%S. | short-fruited | Q o r 30 jnjl | Jamaica p.1 Lamb. ci. 8 |
| 599. HYMENODICTYON Wai. | HYMENODICTYOK. (<i>Hymen</i> , a membrape, <i>diktyon</i> , a net) <i>Rubiace</i> *. 2.-4. | | |
| 4894 thyrsiflbrum Wai | thyrsse-flowered* CD or 15 jnjl | G.Y E. Indies 1819. C Lp | |
| Cinchona thrysiflra Rox. | | | |
| 4895 excelsum IVal | tall | ■ □ □ 30 jl.au | G.Y E. Indies 182a C Lp Rox. cor. 2.106 |
| Cinchona excelsa Rox. | | | |
| 600. CINCHO ^N J « L. | CINCHIO.VA. | | |
| 4896 officinalis L. | official | ■ m 18 jl.au | Cured the Countess of dnchon of a fever.) Muse*. 2.—20. It Peru 1810. C Lp Lamb.ci.1 |
| 4897 scabra Lod. | scabrous | ■ m 6 | 1820. C Lp |
| 601. BURCHE'LLL* R. Br. | BURCHILLIA. | (<i>W. Burchell</i> , a traveller in Africa.) RuMdeea?. 2.— | |
| 4898 bubanza R. Br. | buffalo | • · or 3 my.jn S | C. G. H. 1818. C r.m Bot. reg. 899 |
| 4899 Sap&isisi. JR | Cape | i l Q e l 3 mr | S C. G. H. ... C p.l Bot. reg. 466 |
| 602. RONDELET/il L. | ROXDELETIA. | (<i>W. Rondet</i> , a celebrated physician.) Rubi(tee)e. 9. — 31. | |
| 49(X) americana L. | Americau | • or 10 au | W. Indies 1752. C s.p FLic.213.1 |
| 4901 Ucvigkta J7.it. | smooth-leaved | H · or 12 jl.au | W. Indies 1790. C s.p . |
| 4902 hirta Swz. | hairy | ■ ■ 10 jn.au | Pk Jamaica 1776. G f.B Bot.cab.3-in |
| 4903 /anfulnia Sun. « | Laurel-leaved | ■ ■ 5 jl.au | Jamaica 1824. G \$ B |
| 4904 hirauta Swz. | hairy | ■ ■ 5 jl.au | Jamaica 1824. G \$ B Br.jam.2.2 |
| 4905 thyrsoides Swx. | thyrsoid | ■ ■ 5 jl.au | Jamaica 1819. C s.p |
| 4906 racembsa Sivz. | racemose | ■ ■ 6 jl.au | Jamaica 182a C s.p Br. jam 2. \$ |
| 4907 tomcmbsa Swz. | tomontose | ■ ■ 6 jl.au | Jamaica 181J. C s.p |
| 4908 paniculat Rox. | panicked | ■ ■ 6 jn.au | E. Indies 1820. C s.p |
| 603. COUTARPA Aubl | COUTAREA. | (<i>Coutari</i> , its name in Guiana.) Rubiaceaf. 1. | |
| 4909 sucibsa Aubl. | beautiful | ■ ■ 12 ... P | Guiana 1803. C s.p Aub.gui. 122 |
| Portland/a hexandra W. | | | |
| 604. PORTLAND/^ L. | PORTLANDIA; | (Duchess of Portland, faiaw patroness of b ot ^.) A i M a «» | |
| 4910 grandiflbra L. | great-flowered | f Cjor 12 jn.au | W. Jamaica 1775. C s.p Botmag.28*, C sp |
| 4911 coccinea Swz. | scarlet | ■ ■ 5 ... S | Jamaica 1802 C sp |
| * 605. ADENOCHORAFFis | ADENOPMORA. | (■ ■ a gland, phoreo. to bear.) Campandhcece, 14.— | |
| 4912 marsupiiflra Fis. | purse-flowered | ■ A or ■ ■ Pa.B | 1818. R co Bot. reg. 149 |
| Campanula coronata B. R. | | | |
| 4913 Gmelin Fis. | Gmelin's | ■ my.jl | Pa.B Siberia 1820. D co Gm. si. 3.33 |
| 4914 coronopiflta Fis. | Buckshorn-lvd | ■ A pr 1 jn.jl | Dahuria 1822. R co Sw. IL gar. 104 |
| Campanula coronopiflbia R. Sf.S. | | | |
| 4915 denticulata Fis. | denticulated | ■ ^ or 1 jl.au | B Siberia 1817. D p.l Sw.fl.gar. 116 |
| Campanula tricuspidMa R. & S. | | | |
| 4916 comtinis Fis. | common Siberian | Jt A or 4 jl.s | Pa.B Siberia 1810. D p.l |
| Campanula commfncis R. Sf.S. | | | |
| 2 suavolens Hort. | sweet-scented | ■ ■ jnjl | Pa.B Siberia 1816. D co |
| 3 hybrida R. A S. | hybrid | ■ ■ jjjl | Pa.B Siberia 1818. D co |
| H917 /lhfbila G. Don | Lily-leaved | ■ ■ 2 my.s | Pa.B Siberia 1784. D pi Bot reg. 236 |
| Campanula /lhfbila W. | | | |
| 4918 Kabelaisuna G. Don | Rabcluis's | ■ A or A jn.au | B Siberia 1823. D p.l |
| Campanula Kabelaisuna Ana R. & S. | | | |
| 4919 Lamarkidna Fis. | Lamark's | ■ A or lijn.au | Pa.B Siberia 1824. D Lp |
| Campanula Lamarkidna R. & S. | | | |
| 4920 stylbsa Fis. | long-styld | ■ A or 1 my.jn | Pa. B Siberia 1820. R co Gm.si.3.27 |
| Campanula stylbsa Lam. | | | |
| 4921 intermedia Svt. | intermediate | ■ A or ■ ■ Pa.B | Siberia 1819. D p.l |
| Campanula intermedia R. & S. | | | |
| 4922 eriopiflbia Fis. | Periopio-lvd | ■ A or ■ ■ 8 | Pa.B Siberia 1824. D p.l |
| Campanula eriopiflbia Lam. | | | |
| 4923 Fischer* G. Don | Fischer's | ■ A or 2 jl.s | Pa.B Siberia 1819. D p.l |
| Campanula Fischer R. & S. | | | |
| 4924 veretkijfb/ia Fis. | Pereskia-lvd | ■ A or liji ■ | Pa.B Siberia 182L D p.l |
| Campanula /texcskue/bia R & S. | | | |
| 4925 verticillata Fis. | whorled | ■ ■ ■ 2 jn | L.B Siberia 1783. D s.1 Pal. it 3. G.I |
| Campanula verticillata W. | | | |
| * 606. WAHLENBE'RGIASchr. | WAHLENBERGIA. | (G.Wahlenberg, M.D., until 1811 Lapp.) <i>Campanddior</i> *. 5.— | |
| 4926 grandiflbra Schr. | great-flowered | ■ A or 1 jn.au | B Siberia 1782. D p.l Bot. mag. 252 |
| Campanula grandiflbra L. | | | |
| * 27 pendula Schr. | pendulous | O or ■ jl.au | W.p Madeira 1777. ■ s.l |
| 4928 elongata Schr. | elongated | O or ■ jl.au | B — 1819. ■ co |
| Campanula elongata W. | | | |
| * 607. CAMPANULA L. | BELL-FLOWER. | (Dimin. of catnpna, a bell; corolla.) <i>CampantUdceat</i> . 129.—220. | |
| T* ²⁹ cenisia L. | Mt. Cenis | ■ A or i jnjl | Switzerl. 1775. R co Al. ped. 1. a 2 |
| J930 uniflbra L. | one-flowered | ■ A or ijnil | Lapland 1815. R sp Fl. lap. 9.5,6 |
| 4931 diffusa Vahl | diffuse | ■ A or i Jju | Alps,Italy 1826. R co Cyr.ne. II. 2 |
| 4932 muophph(flaJ&7. | small-leaved | ■ A or 4 jnjl | B Hungary 1820. R co |
| J933 Uell&rdi All. | Bellardi's | S A or 4 jn.jl | Italy 1813. R co AL ped. 1.85.5 |
| 4334 pflia L. | russet | ■ A pr 1 jnjl | Austria 1779. R co Bot. cab. 554 |
| 4935 Zol'sii Wul. | Zoys's | ■ & or 1 jn.au | D.B Carniola 1813. D co Jac. ic 2. 334 |
| 4936 carpatica L. | Carpathian | ■ A or 4 jn.au | Carp. Alps 1774. D p.l Bot mag. 117 |
| 4937 rotundiflbia L. | round-leaved | A or 4 jn.au | Britain hea. D p.l Eng. bot. 866 |
| 2 tibre &bo | white-flowered | A or 1 jn.au | Britain woods. R co |
| 4938 exc^sa Schl. | cut off | A or i myjn | B Switzerl. 1820. R co |
| L usilla Hae. | diminutive | A or i jl.au | Pa.B Switzerl. 1821. R co |
| ca-spithsa Sco. | tufted | A or i jn.au | Austria 1819. R co Bau. pr. 34.51 |
| Pfimila B. M. | dwarf | A or t jn.au | Switzerl. ■ D p.l |
| dujo, 2flbralbo | white-flowered | A or t jn.au | R co Bot mag. 512 |
| gg puMsens Schmidt | pubescent | A or 1 ap au | B Bohemia 1813. D co |
| erminii Lk. | slender | QJor 1 ap au | N. S. W. 1794. R co |
| Hermelin's | | Jin.au | Portugal 1823. S co Fl.nor.2.72 |

PENTANDRIA MONOGYNIA.

CLASS V,

4945 quadrifida R. Br.	four-cleft	ft	Jn.au	B	N. Holl.	1820.	S co	
4946 littoralis Lab.	sand	ft	ap.au	B	N. Holl.	1820.	S p.l	Lab. n.h. 1.70
4947 Scheilchzeri Vil.	Scheuchzer's	f	jn.au	B	Europe	1813.	D co	Bot cab. 485
4948 angustifolia Lam.	narrow-leaved	1	jn.au	B	France	1818.	D co	Mag. ma 46
4949 virgata Raji.	twiggy	:	1	jn.au	N. Amer.	1823.	D co	
4950 lanceolata Lap.	lanceolate	1	jn.au	B	France	1819.	R co	Al. ped. 47. a
4951 heterodoxa Vest	heterodox	ft	jn.au	B	Siberia	1818.	K co	
4W52 Waldsteinia It. & S. Waldstein's	flexibsa Kit.	ft	jn.au	B	Hungary	1824.	R co	W. & K. 136
4953 /inifolia Hue.	Flax-leaved	f	jn.au	B	SwitzerL	1819.	D co	Vil. del. 1.10
4954 rigescens Pall.	rigescens	ftjn.au	B	Siberia	1820.	R co		
4955 spatulata Sm.	spatulate-/w*	1	jl.au	B	Greece	1817.	S p.l	Fl.gr. 203
4956 ramosissima Sm.	branchiest	ft	jl.au	B	Greece	1820.	S co	Fl.gr. 204
lobevio des W.								
4957 ptilula A.	spreading	1	jl.au	V	Britain	pas.	S p.l	Eng. bot. 42
4958 neglecta R. & S.	neglected	2	m au			1818.	S co	
4959 calycina Bieb.	-	1	jl.au	B	Tauria	1820.	R co	
4960 Broussonetiina A. \$ & Hroussonet's		1	jl.au		Mogadore	1820.	S co	
41*61 Locflingy Brot.	Loefling's	1	jl.au		S. Europe	1818.	S co	
4962 infundibuliflum Vest	funpidwd	2	jl.au		Siberia	1825.	D co	
4963 Aapunculatum L.		3	jl.au		Britain	hed. b.	S r.m	Eng. bot 283
4*64 elegans R. & S.	elegant	^ ^ o r	1	jl.au	Siberia	1821.	R ro	
siccibsa W.								
4965 persicifolia L.	Peach-leaved	3t	A or	3	jl.a	B	Europe	159ft D p.l Fl.dan. 1087
2 maxima	largest peach-lyd'i	A or	3	jl.a	B	Europe	1596.	D <p.l Bot. mag. 397
3 flora pleno	double A/uoflw^t	& or	3	jl.a	B	Europe	1596.	D co
4 fibre albo	white-flowered	^ A or	3	jl.s	W	Europe	1596.	D co
5 flwo albo pleno	dble white-flwd^f	A or	3	jl.s	W	Europe	1596.	D co
6 grand is	Urge-flowering	^ A or	3	jl.s	B	Europe	1596.	D co
4966 pyramidal is L.	pyramidal	5	A or	4	jl.s	Pa.B	Carmola	1596. D p.l
2 fibre albo	white-flowered	3	A or	4	jl.s	W	Europe	... D p. I
4967 versicolor Sm.	various-colored	£. A or	4	jl.s	St	Greece	1788.	D a1 Bot.repi396
planiflora W. en.								
4968 obnqua W.en.	oblique							
4969 amenckna L.	American							
4970 planiflora Lam.	flat-flowered							
4971 acumpanata Mx.	acuminate							
4972 Ottontina R. & S.	Otto's							
4973 nitida ii. Jic	shining							
2 fibre caeruleo	blue-flowered	^ A or						
4974 aurea L.	golden-flowered*	^ \ or	3	jl.s	Y	Madeira	1777.	S s.p Bot. reg. 47
2 latiflbia R. \$ S.	broad-leaved	St	2	jl.s	Y	Madeira	1777.	S s.p Iuh. cd. nov. 41
3 angustiflbia R. & S.	narrow-leaved	M	or	2	jl.a	Y	Madeira	1777. S s.p Jac. sc. 4.472
4975 rhomboidea L.	rhomboid	ifc	A or	2	jl.	Pa.B	Switzerl.	1775. D p.l Bot. cab. 603
4976 Alpine Lam.	Alpinus	A	or	1	jn.jl	B	Italy	1800. D p.l
4977 latiflbia L.	broad-leaved	A	or	4	jl.	P	Britain	s. m. p. S s.p Eng. bot. 302
2 fibre albo	white-flowering	^ A or	3	jl.au	W			
4978 macrantha F.*	long-flowered	^ A or	3	jn.jl	P	Russia	1822.	D co
4979 eriocarpa Bieb.	woolly-fruited	^ A or	2	jn.jl	P	Caucasus	1823.	R co
4980 urticiflbia Schmidt	Nettle-leaved	^ A or	3	au	V	Germany	1800.	D co
4981 Trachelium L.	Throatwurt	& A or	4	jn.au	V	Britain woods.	D p.l	Eng. bot. 12
2 fibre pleno	double A/t/e-flwd^	A or	3	jl.au	B	Britun	g.trd.	D co
3 fibre albo	white-flowered	^ A or	3	jl.au	W	Britain woods.	D co	
4 fibre pleno albo	dble white-flwd^t	A or	3	jl.au	W	Britain	gard.	D co
4982 stellidiflbia Lap.	Daisy-leaved	^ Q) or	2	jl.au	B	Pyrenees	1823.	S s.p
4983 crenata W.	crenatiel-leaved	& A or	2	jl.au	B	Russia	1810	R co
neglecta Bes.								
4984 rapunculoides L.	Rampion-like		3	jn.jl	B	England	woods.	D p.l Eng. bot. 1369
4985 infundibuliformis Sims	funnel-shaped		2	jl	P	Siberia	1822.	D co Bot mag. !&>
4986 trachcliodes Bieb.	Throat wort-like		3	jl.au	B	Caucasus	1817.	R co
4987 sarmatica B. R.	Sarmatian		2	jn.au	Pa.B	Siberia	1803.	D co Bot. reg. 2SI
gummifera R. % S.								
4988 bonomensis L.	Bononian		2	au.s	B	Italy	1773.	D co M. h. 2.5.4.38
4989 simplex Dec.	simple-stmd		2	jl.au	B	S. France	1819.	R co
4990 obhquiflbia Ten.	oblique-leaved		2	jl.au	B	Italy	1824.	R co Fl. nap. 17
4991 trichocalytina Ten.	hairy-calyxed				B	Italy	1823.	R co Fl. nap. 16
4992 ruthénica Bieb.	Russian		2	jn.au	B	Caucasus	1815.	D co
4993 iprēta R. & S.	despised		2	jl.au	Pa.B	Siberia	1825.	R ro
4994 graminiflbia L.	Grass-leaved		1	jn.jl	B	Italy	1816.	D s.p W. & K. 15?
4995 tenuiflbia Kit.	line-leaved		1	jl.jl	V	Hungary	1817.	R pi W. & K. 155
4996 glomerata L.	clustered		2	my.s	V	Britain	ch. pi.	D p 1 Eng. bot. 90
3 fibre albo	white-flowered		1	my.s	W	Britain	ch. pi.	D co
4 fibre>filo fillo	dble wlnte-flwi		1	my.s	W	Britain	gard.	D co
4997 cephalantha Horn.	head-flowered		1	jl.i	B	Russia	1817.	R co
4998 cephalitis Horn.	roimcf-headed		1	jn.jl	B		1818.	D co
4999 speciosa Horn.	showy		2	my.jn	P	Siberia	1824.	R co Bot reg. 690
glomerata /3 dahurica B IL								
5000 nica*ensis If. \$ S.	Nice		*	jl.jl	p	Piedmont	182a	D co AL ped. 39.1
glomerata All. petra>*A L.								
5001 congesta R. # 5.	crowded		1	jl.au	B	France	1823.	R co
5002 elliptica Kit.	ellipticul		1	jl.au	B	Hungary	1826.	R co Boc. mu. 58
5U03 foehsia Ten.	leafy		1	jl.au	B	Italy	1826.	R co Fl. nap. 18
5004 Rafneri Perp.	Rainer's		1	jl.au	B	Italy	182a	R co Bib. it 13. 1817
5005 Cerviciana L.	Thruatwort		3	jl	L.P	Germany	1708.	S s.p Bot. cab. 452
5006 macrostachya W.	long-spiked		1* jn.jl	B	Hungary	1814.	S co	
multiflbra Kit								
5007 cervicurioides R. \$ S.	Cervicaria-like		1	jlau	B	Italy	1822.	R co
■ ■ ■ ■ ■ Bieb.	hill		1	jl.au	B	Caucasus	1803.	I) p.l Bot. mag. 527
5009 aziirea Sal.	azure		Hjn.jl	L.P	Switzerl	1778.	D p.l Bot.. mag. A31	
5010 Lorek Pol.	Lore's		lf jn.jl	P	Italy	1824.	S co Bot. mag. fcdtf	
baldensis Balb.								
5011 lactiflria Bieb.	milk-flowered		6	jl.s	W	Siberia	1814.	C &p Bot. reg. 241
5012 thyrsoidae..	crowded-jlfM*		2	ils	Pa.B	Bavaria	1817.	C sp Bot. cab. 515
5014 velutlna Desf.	thyrsoid		2	jiu.au	B	SwiterL	1785.	S s.p Bot. mag. 129V
	velvety		ft	my.jl	B	S. turope	1826.	R s.p Fl.gr.210

£086 triquetra L.	triangular	■	1	my.au	B	C. G. H.	1774.	C	B.p	
5087 fenestralis Cav.	windowed	■ uOJ ^{or}	1	jn.s	B	Mexico	1824.	S	p.m	
5088 longiflora L.	long-flowered	£ [1	my.au	W	Jamaica	1752.	S	sp	
5089 tomcnidaa L.	woolly	^ LAI «P	£	my.*	B	C. G. H.	1821.	D	s.p	
50M scundin L.	side-flowering	j£ IAJ ^{or}	j,	my.au	W	C. G. H.	1790.	S	s.p	
50MI acuminatea Sivz.	acuminated	JU E3 or	2	jn.au	G.v	W. Indies	18<A	C	p.l	
50112 mcotiaurfldia Roth	Tobacco-leaved	£ OJ or	2	jn.jl	-	Indies	1822.	C	p.l	
5093 Clayton/ana Mx.	Clayton's	3t A ^{or}	2	jn.jl	B	N. Amer.	1824.	D	s.p	
	goodniahls W.	spicata l.am.							W.h.b.1.30	
5094 nssfurgens L.	nssurgent	■ ■	3	jno	S	W. Indies	1787.	C	sp	
5095f ilgcns/.4J?.	fulgent	3t AJ r	3	my.s	S	Mexico	1809.	C	B.p	
5096i oerbascifolia Swi.	Mullein-leaved	3t A or	6	my.jn	R	Nepal	1822.	D	rm	
5097 pyramidalis Walt.	pyramidal	f iAI ^{or}	4	ft	P	Nepal	1822.	D	co	
509B cardinalis B.J.	cardinal flower	■ ■	3	my.8	S	Virginia	1722.	C	sp	
	shining		3	my.s	S	Mexico	1814!	C	sp	
50O debilis L.	feeble	■ ■	1	jl.au	B	C. G. H.	1774.	S	sp	
5101 alatail. 2?r.	vringed-stalked	■ ■	11	my.au	B	N. S. W.	1804.	S	sp	
51U2 Tipu L.	Tupa	8 b.o	8	b.o	S	J. Fernan.	IK24.	It	CO	
5103 sphyliitic* L.	sphyliitic	■ ■	2	au.o	L.B	Virginia	1665.	C	ap	
5104 sunnemannis L.	Surinam	■ ■	2	jal	O	W. Indies	1786.	C	sp	
2 rubra			2	jal	It	W. Indies	1820.	C	*.p	
5105 gracilis R. Br.	red	■ ■	1	jl.o	D.B	N. S. W.	1801.	S	sp	
5106 purpurascens R. Br.	slender	■ ■	1	jn.au	B	N. S.W.	1809.	D	sp	
5107 inflata L.	purpurascent	■ ■	lftjlau	Pa.B	N. Amer.	175ft.	S	ap	Ac. up. 1741,1	
5108 Cliffrtiana L.	inflated	■ ■	JJ	jlau	Pk	N. Amer.	1733.	S	sp	
5109 Michauxii Nut.	Clifford's	■ ■	1	il.au	II	N. Amer.	1800.	S	co	
Cliffr/Wmrt. Mx. not Nut.	Michaux's									
5110 trialata D. Don	three-winged	£ LAJCU	ijl.au	B	Nepal	1822.	S	s.p	Hook. ex. fl. 44	
micrantha Hook.										
5111 tircnsl.	burning	■ ■				England	hea.	S	L	
6112 serrulata Schott	serrulated	O or	f	jn.jl	B	Spain	1820.	S	ap	
5113 puberula Mx.	mossy	k A ^{or}	I	jn.jl	Pa.B	N. Amer.	1800.	D	s.p	
5114 adia* na Mx.	pleasing	& A or	3	jn.au	B	N. Amer.	1812.	D	SI	
5115 minima L.	small	g Alcul	■ ■		W	C. G. H.	1772.	R	sp	
5116 minima B. M.	least	a iAi ^{or}	■ ■		W	C. G. H.	1800.	D	sp	
5117 laurcntian L.	Laurentian		*jl		B	Italy	1778.	S	B.p	
5118 tenella Biv. »	delicate	iQlor	■ ■	my.jl	P.y	Sicily	1821.	D	co	
5J19 campainoldes Thun.	Campanula-Ik	■ ■	■ ■	my.au	W	China	■ ■	I	co	
6120 £nnus L.	Erinus	■ ■	■ ■	jn.8	B	C. G. H.	1752.	S	8.p	
5121 campanulata Lam.	campanulate	** IDI or	■ ■	jn.jl	B	C. G. H.	1821.	S	sp	
5122 finnoldia L.	Erinus,likc	£ OJ or	■ ■	jn.au	B	C. G. H.	1759.	R	sp	
5123 bicolor H. K.	two-colored	iQJ or	■ ■	jn.au	Pa.B	C. G. H.	1795.	C	sp	
5124 ilicifolia AVr	Holly-leaved	£ LAJ or	■ ■		Pk	C. G. H.	1815.	D	sp	
5125 anceps L.	two-edged	id or	■ ■			C. G. H.	1818.	S	B.p	
5126 pubescens H. K.	pubescent	£ i^or	■ ■			C. G. H.	1780.	R	&p	
5127 zylanica L.	Ceylon	£ iAI ^{or}	■ ■			E. Indies	1821.	D	sp	
5128 lutea L.	yellow	£ iAI ^{or}	■ ■			C. G. H.	1774.	S	8.p	
5129 hirsuta L.	hairy	£ iAI ^{or}	■ ■			C. G. H.	1751.	C	Bur. af. 40.2	
5130 varhfbilia A3f.	vanous-leaved	£ iAJ or	■ ■			C. G. H.	1812.	C	sp	
5131 coronopifolia L.	Coronopus-lvd	£ vA ^{or}	■ ■			C. G. H.	1752.	S	sp	
5132 umbellata Vest.	umbellate	£ iAJ ^{or}	■ ■			C. G. H.	1818.	D	sp	
5133 Thunbergia Swt.	Thunbergia's	ff <A ^{or}	■ ■		c. G.H.	1812.	D	s.p		
roronopitula Thun. not B. M.										
5134 crenata Thun.	notched-leaved	£ iAJ or	■ ■	an.my	B	C. G. H.	1794.	C	"P	
5135 dentata Cav.	toothed	£ iAI ^{or}	1	ji.jl	B	N. Holl.	1824.	D	s.p	
5136 inundata R Br.	overflowed	A A or	1	ji.jl	B	N. Holl.	1821.	D	Cav. ic a 522	
5137 carulea B. M.	blue	A A or	U	ji.jl	B	C. G. H.	1824.	D	s.p	
5138 Sims's Swt.	Sims's	tf	g Alor	I o.n	B	C. G. H.	1819.	D	pL Rot mag. 2701	
pedunculata B.M. not R. Br.										
5139 cdoncolor R.Br.	self-colored	< lAlor	1	my.jn	B	N. Holl.	1819.	D	co Bot mag. 2251	
5140 globosa R. Br.	globose	£ LAT ^{or}	1	my.jl	B	N. Holl.	1824.	D	p.1	
5141 jenecioides Cun.	Senecio-like	f .Aior	1	jl	B	N. Holl.	1824.	D	p.1	
Is6toma axillaris L'tnd.									Bot reg. 964	
5142 pedunculkta A.2?r.	pedunculated	tf	LAJ or	1	o.n	B	N. Holl.	1819.	D	co
5143 corymbosa B. M.	corymbose	£ iAJ or	§	jn.au	R	C. G. H.	■ ■	D	p.1	
610. MONOTYSIS Sal.	MONOPSIS.	(Monos, one. opsis, a face.)				Lobelihcce.	2. — M			
5144 conspicua Sol.	conspicuous	O of	ijl.au	B	C. G.H.	1812.	S	s.p	Bot mag. 1499	
Lobelia Speculum B. M.										
5145 inconspicua Sal.	inconspicuous	O or	\jl.au	P	C. G. H.	1812.	S	s.p		
611. PHYTEITMA L.	RAMPION.	(A name adopted from Dioscorides.)				Campamdhcea	28.-96			
5146 pauciflorum L.	few-flowered	■ ■	my.jn	B	Switzerl.	1823.	D	p.l		
5147 globularifolium Ster.	Globularia-lvd	■ ■	my.jl	B	S. France	1820.	D	p.l		
5148 Scheuchzeri All.		i nw.jn	■ ■		Switzerl.	1813.	D	co	Bot. mag. 1797	
5149 sco zomn'folium Vil.	Scorzonera-lvd	i jl.au	■ ■		Alps	1819.	D	p.l		
5150 sibiricum Vest.	Siberian	i jn.au	■ ■		Siberia	1817.	D	p.l		
5151 Michlit AH.	Micheli's				Switzerl.	IS22.	D	p.l		
5152 hemisphaericum L.	hemispherical				Switzerl.	1752.	D	p.l	Jac.ia2.SSS	
5153 hispid um Schl.	hispida				Switzerl.	1825.	D	p.l		
5154 comsum Wul.	tufted				Austria	1752.	S	s.l	Jae, ap. 5. \$Q	
5155 orbiculare L.	round-headed				Austria	1752.	S	s.l		
5156 Sieberw. Spr.	Sieber's				England	ch.pa.	D	p.1	Eng. bot 1**	
5157 Charm&ii Vil.	Charmeli's				Pyrenees	1826.	D	p.1		
5158 humile Schl.	humile				Pyrenees	1823.	D	p.1	Vil. deL 2.1. "a	
5159 ina?qtatum Kit.	levelled		1	my.jl	B	Switzerl.	1825.	D	p.1	
5160 ellipticum VU.	elliptic		1	jn.au	B	Austria	1820.	D	p.1	
orbiculare W.			1	jn.au	B	Switzerl.	1817.	D	p.1	
f If? & jm Schmidt	short-leaved		1	jn.au	B	Switzerl.	1819.	D	p.1	
5163 cordatum B. M.	black		1	jn.au	D.B	Bohemia	1820.	D	a.p	
5164 reticulifolium Vil.	Betony-leaved		2	jl.au	B	Hungary	1804.	D	co	
5165 spicatum L.	Bipkd		2	jn.jl	Pa.B	S. Europe	1818.	D	p I	
Slon Hillerl. All.	Hallcr's		2	jn.au	W	Europe	1597.	D	p.1	
5167 ovatum IV.	ovate-spiked		2	jn.au	D.V	Europe	1814.	D	Bot. mag. 2 ⁵ 7	

✓VH^purjirirea Rox. (5347 Utintblr Hun, 5348 undulata IIW.	Cfdamlia puqili: •waved, leaved	m lor 60 O E. Innn 5349 violicen Sun. viditi-Aerrifrf <i>Tapagamia</i> violecea Sure.	lor 10 P Itox. cor. 1.5* CD or 10 P.B.R. C r.m. 1 lor 80 Y i: ludiM C r.m.	E. Indies C r.m. I.iii.tr. & SJ! C r.m.
650. CEPHAE*LIS Su*. CaMtun 5349 violicen Sun.	(Kephale, a heat1: flower*.)	RubiAcryx. 8.-26.		
5350 tuinenlosa fAM ii^il punleca Vtihl	*. ESjor 1 Jn-jl W W. Indie* 181S.	C r.m. Aub. gui I. tw		
tall	a 1-i or 4 jns Br Trinidad 1835.	C r.m. /Vul. gui. 1. 55		
5353 win'ureA IK.	tt 11 or & jtl.S w Jamuka 1793.	C r.m. l.p.		
<i>Tapabmin</i> piip&re1 Aith	* Zi of I ap.jn p Trinidad 18JEL.	C r.m. Aub. gui. 1. G33		
5354 involucrifta W. BSS5 pedunculifl. SaL	o lor 5 jni. W Guiana 18JR	C r.m. Aub. gui. 1.64		
5356 asillarii Sws. calycin L.	* 11 or S f W S. Leane 1d'ii.;	C r.m. Par. luu. 103	ft Und. col. £1	
OJtillfly	(* 1 or 4 ap.my W Uraii	C		
651. GEOTHILA D Don. GenrnLA £357 ia>ifstSrmii D. Don kidney-ihaped	(Gf, the earth, phileo, tolove.1 llub&ccte. 1.	U 1P Jac. am. 4fi		
Psycho tria herbicea L.	fAlor i apjn r.a W. Inuiei 17:			
659. SARCOCE'PHALUS^/i G ns 5358 esculentut Af%,	Puca. (Sax, Besh. keptiaU, a head; flower*.) RKB&cft. 1.			
eatable	a • fl 15 jl i'k S-Lcone	C P-I Hort. tr. 5.18		
053. HIHITE'I.LA W. ESnentLta.	(HirAi*, li airy; young branches.) Chry&oliaWinesr. 2.—13L			
5359 racembsa Lam. racemose	? 1 lor SS ... V W. [iidie] 1782.	C Lp Aub. gui. 1.98		
ameriduia Aub,	Irinndrou*			
5300 triundit Swx. ameriduia Joe. not Aub.	» 11 or SO ... W Jamaica 1816.	C Lp Jac. am. iic II		
paiiicu&ta				
K54. TIUPH.VSIA Lou, Twraisu. SStil auraiaital Mm.	{Triphatiot, triple; Sower* and leave*.)			
Llmunui IriloDaia It,	a 1 Jfr s JnJl W China 17W.	c r.m But- rep. 143		
635. HTIS Z.	Tan. (Giyd			
538 vinifeia L.	wine-bearing 1 fr jn.jl G Various 1648.	C r.m. Amfx-t&a*. ia—23.		
5381 dent'ita Lk.	crumpled 1 ft 20 jnjl G	C C Sdim, ic. 34 8		
5385 Wallfchtf Dec.	tooDuil 1 or 10	C r.m.		
5386 gflfbrJla Koih	Wnllich'a 1 or W	C r.m.		
5387 Indlea i.	i moot tied 10 ***% *	E. 1 rj.lii-s 1H19.		
5388 enrieba Dec.	Indian 1 or 10	C .1 R.mnl.7.6		
tmica Swx.	Caribc 1 or 10	C r.m. SL jam. £.210.*		
S369 jabrtisca L. 2b&ca* albii	WiJd Vine 1 fr 10 .. G N. Amer.	L s.p. Jac.sc.42(!		
5370 jntivllii Ajr. \\ IIIIII. JV.	white-borrjed 1 lr 10 .. G H. Aratr.	L C *P Jac tc 425		
lummer	or 10 .. G N. Amer.	WS		
5371 Purlnf Ham.	tnualc.!,yivfd 1 or 10 .. G N. Amer.	C r.m.		
5372 cordiftilia Mr.	Fur.ut's 1 or 10 .. G Nepal	r.m.		
5373 rip&ria Hx.	cord a U'-It.; *«! 1 or 10 .. G N. Amer. 18061	C C Joe. <. 427		
5374 rotundifolia Mi.	river.bank T or 'M mjj,Jn	C C <p Hot, mag. 2429		
roundle>aved	J. or 90 .. G N. Amer. 1606.	B-p		
65ft AMPELOTSJSJit.	Ampelopfil	[Ampetot, a vine, rwemblanco.)	Amveh'dca: 4.-7.	
5375 cordita Mr.	lieatt-/iv)tv(/	i or H ap.my N. Amer. 1803.	C co	
Cltius Ampeluplis Pen.				
5376 biplnniU AT*.	bi pi mi ate	or 15 Jl.au Fa K. Amer. noa	C co	
5377 AeuBfftem A.r.	IvJ-ltke	J or OO JllJI Ro N. Amer. ma	C co	Cor. ca.JO0
CIMU* quinquef&liui	Mr.	or 60 up. my Pa K. Amer. (80S	C co	
5378 hiwtita Dunn	hairy			
657. BHA'MNUS L.	BI'CKTHOBN.	(Ram, a tult of tranche*, Cell.)	J u met f. sa-6i..	
5379 jflatenius L.	bast.-lml.- A la tut	a or 40 apjn G S. Ktirop 1620.	J u co	
1 balericra		or 40 apjn G B. Eurtif* Balearic	J u co	
Sgftbra	Sponinh	or 40 apjn G S. Europe	I. co	
5 maculita	cmooth	or 40 op. n G S. Europe	I. co	
6 a'irea	entire-leaved	or 40 ap. D G S. Europe	I. co	
7 v.-irrg^ta	nj nt ted	or 40 II G S. Europe	I. co	
8 argL-itteaa	golden -riged	or 40 np! II O. S. Europe	I. co	
9 ljiuinijra	TjiriegaU-d	or 40 up. n O. S. l-nrujkj	I. eo	
£380 hybridus Htril.	Mwr-edged	or 40 II G S. Kuri.ju.	I. co	
£381 CtlM W.	JiiKgtel-ZtdwJ	or 40 II G BL T-ii "j!"	I. co	
5S88 glanduliwu.W,jTt.	Cluuijt'i	or 30 my.jn a S. Europe 102ft	L co	Hcr.ur.S
5383 integrifliliu* Dec,	glandukir	or 15 jn.jl G Caauiel 17K5.	C PJ \m. ii	
5384 surnahf n&is Sen.	in tire-leaved	or 1 or 3 G Y Tenerift 18SJ2.	C r.m. Jie-. mon i"	
5385 kinRifblids Dettf	Surinam	or 1 or 1 .. G Y Surinam 1SJ2	C rm Scop,i	
	Inngt.ctived	or 6 .. G	L co	
5388 ^ n o M e V /frtS.	Winttr-lwr.-lk	a 1 J or ll au.i Y C G. H. 1778.	C p. I Her. trr 9	
5387 feltifluu* Thuk	C Iltis-leavcd	ft 1 or SO ap.jn G.T C .1 Hur. af. wj <	C i. co LJIk. boL I?"	
5388 catharticiu I.	purgung	or 15 my.jn G.V >iig!and hod. L r.m.		
£389 virgutis /iflr.	dyc-r»	or 8 in.jl y issa L r.m.		
5390 tincliriuM W. S/IL	dyer's	or 5 my.jn PaV Uillt.,ny L r.m.		
5391 ii. (Wt dries f.	rock	or 6 jn.jl G.V S. Europe 1884.	I. co W. & K	
S395 mxiiiius /..	Olive-lifco	or 1 my.jn G.Y Europe 1755.	C co ArLi m. 'sJ	
tiS95 olcold^ /j.	Box-leaved	or 4 JnJl G.Y Spain 171SL	C co Jai!. su. ».	
W'l ftuxifluua J'ojr.	pUbOM'Cltt	or 6 my.jn Pa, Kr.mce 1817.	I. co rtuk.al.™	
5395 pub'tcciu JP<.	crenulnte	or 4 my.jn OY Iteetttb pa, E. r.m.	C P-I	
539A crmulktuf J/ A*	V, & liter II. h.k c	or 8 mr G.Y Spain 1754.	L co AF, IC 2. TM	
5397 feclodtum L.	A i nurad	or H t.d. G.Y Sjai! 1754	L co	
	dwarf	or 6 t.d. G.Y Stbera 1BB3.	L co Scnp-e^	
53feErythT<ixylon />aa	lic-d.UODd	or 6 ll.au. Y.w Canniola	L co	
5399 piimilu* I.		or 9 ll.au. Y.w		

665. SEKA.XIA Com.	SENAZIA.	(Senac, a French physician.)	Pittosporea.	2.—
5467 undulata Lam.	wave-leaved	Loor 12 ... W	Bourbon 1785. C	Lp
5468 nepalensis Dec.	Nepal	Loor 10 ... W	Nepal 1820. C	Lp
Celastrus verticillatus Box.				
* 666. EVO'NYMUS Tou.	SPINDBL TBBB.	(Having afoed name, by antiphrasis; fetid.)	Celastrinece.	17.—30.
5469 europeus L.	European	> or 15 my.jl W	Britain 'bed.	S B1 Eng.bot.362
1 pdmplus	dwarf	or 4 my.jl W	—	S 8.1
2 leucocarpus Dec.	white-fruited	or 12 my.jl W	—	L r.m
5470 verrucosus Sco.	warted	ill or 6 my.jn G	Austria 1763.	L p.1 Schm. ar. 72
5471 latifolius Bauh.	broad-leaved	da or 10 jn.jl G	Austria 1730.	L s.1 Bot. mag. 2384
54-12 hanus Bieb.	dwarf	ft or 3	Caucasus 1825.	C r.m
5473 atropurpureus Jac	dark purple	ft or 6 jn.jl	N. Amer. 1756	L p.1 Schm. ar. 73
5474 americanus Jac	American	or 6 jn.jl Pk	N. Amer. 1683.	L s.p Schm. ar. 75
5475 sarmentosus Nut.	sarmentose	or 3 my.jn Y	N. Amer. 1824.	L p.1
scändens Hort.				
5476 angustifolius Ph.	narrow-leaved	ft or 6 jn.jl Y.B	N. Amer. 1806.	L p.1
5477 obovatus Nut.	obov&te-leaved	ft or 3 my.jl Pk	N. Amer. 1820.	L p.1
5478 japonicus Thun.	Japan	M i_J or 6 jn.au Pk	Japan 1804.	C p.1 Ka?m. ie. 8
5479 lcid us D. Don	shining	ft or 7	Nepal 1820.	C s.p
5480 micranthus D. Don	small-flowered	ft or 8	Nepal 1820.	C s.1
5481 chinensis Lou.	Chinese	ft J or 4 my.jn W	China 1820.	C co
5462 echinatus Wai.	prickly	J or 10 my.jn W	Nepal 1824.	C co Bot mag. 2767
5483 grandiflora Wai.	large-flowrcd	J or 8 my.jl W	E. Indies 1824.	C s.p
5484 grissus Wai.	thick	J or 6 my.jn W	Nepal 1824.	C co
5485 Hamiltonidni< Wai.	Hamilton's	J or 20 mr W	Nepal 1825.	L r.m
•667. COLLETTA* Com.	COLLTIA.	(Collet, a French botanist)	Rha'mnear.	4.—7.
5486 spinosa Lam.	spiny	D or 2 my.jl Ap	Peru 1823.	C Lp Lam. il. 2.129
horrida W.				
5487 aeratifolia Ven.	saw-leaved	■■■■■ my.jl Y	Peru 1822.	C l.p Ven. eels 15
5488 obcordata Ven.	obcordate	■■■■■ G.v	Peru 1822.	C l.p Ven. eels 92
(5489'Ephedra Ven.	Ephedra-ft'Ae	J or ■■■■■ G	Chile 1823.	C co Ven. ch. 16
*668. CEANO ^X THUSL.	RED WOOD.	(Keo, to prick; name used by Theophrastus.)	Bhdnnete.	SSL —
5490 lasigatus Dec.	smooth-leaved	■■■■■ G.v	W.Indies 1818.	C l.p
.Ahāmnus laevigatus	Vahl			
5491 capensis Dec.	Cape	*1_J or 3 my.jl W	C. G.H. 1823.	C l.p
.fīhamnus capensis Thun.				
5492 zeyl&nicus Roth	Ceylon	D or 3	Ceylon 1818.	C s.1
Celastrus zeyl&nicus Schult.				
\$5493 asiaticus L.	Asiatic	i_J or 12 jl.au Pa.Y	Ceylon 1091.	C p.1 Cav. ic. 5. 440.I
5494 sphaerocarpus Dec.	round-fruited	■■■■■ 15 ... G.v	Jamaica 1824.	C p.1
Jthāmnus sphaeropermus Swz.				
5495 infestus // & B.	troublesome	■■■■■ or 4 ...	Mexico 1824.	C Lp H.&B.n.7.61*
(5436 colubrinus Lam.	snake-wood	■■■■■ tm 20 my.n G	Bahama 1762.	L co Jac. vin. 3.50
iZhamnus colubrinus L.				
(5497 reclinatus Hen.	reclineat	■■■■■ 5 au G	Jamaica 1758.	L co Br. jam. 29.2
iithamnus ellipticus H. K. •				
5498 mystacinus Dec.	bearded	CDor 13 n W.o	Africa 1775.	S p.1
/Zhamnus mystacinus H. K.				
(5499 cubensis Lam.	Cuba	Qor 5 ... C	Cuba 1820.	C p.1 Jac.vin.49
jRhamnus cubensis L.				
5500 azureus Desf.	blue	LJ or 10 ap Pa.B	Mexico 1818.	C p.1 Bot. reg. 291
ca&rfleid Lag.-				
5501 tardiflorus Horn.	late-flowering	ft or 3 s.n W	N. Amer. 1820.	L s.1
5502 americanus W.	American	ft or 2 jl.o W	N. Amer. 1713.	S p.1 Bot. mag. 1*79
5503 macrocarpus Cav.	long-fruited	Dor 3 jl.o Y	N.Spain 1824.	C l.p Cav. ic. 37270
5504 nepalensis Wai.	Nepal	ft or 10 jn.o ... Y	Nepal 1820.	C p.1
5505 ovatus Desf.		ft or 3 jn.o W	N. Amer. 1818.	L s.1
5506 perenni Ph.	perennial	A or 2 jn.s W	Carolina 1822.	L s.1
5507 intermedium Ph.	intermediate	or 2 jnjl W	N. Amer. 1812.	C Lp Pluk. al. 28. &
5508 sanguineus Ph.	blood-stalked	or 2 my.jl W	Missouri 1812.	C Lp
5509 microphylus Ius Mr.	small-leaved	ft or 1*jn.jl W	N. Amer. 1806.	L p.1
(5510 afriicanus W.	African	I or 6 mr.ap B	C. G. H. 1712.	C p.1 Pluk. ph. —
5511 Mocinianus Dec.	Aiocino's	J or ■■■■■ ... Mexico	1824.	C p.1
Gf9. CORYNOCA'RPU'S Font.	CORYNOCARPUS.	{Koryne, a club, karpos, fruit.)	Myrsine*.	1.
5512 Isvigatus Forst.	smooth	± i_J or 20.	N. Zeal. 1823.	L r.m Font 16
• 670. POMADE'RRI'S Lab.	POMADEBRISz	(Poma, a lid, derris, a skin; berries.)	Rhahncc.	
5513 phillyreoldes Sieb.	Phillyrea-like	■■■■■ 2 ap.my Pa.Y	N. Holl. 1818.	C s.p
5514 Lanigera B.M.	woolly	■■■■■ 3 ap.my Pa.Y	N. Holl. 1806.	C s.p Bot. mag.
5515 elliptica Lab.	ova-leaved	■■■■■ 10 my.jl Pa.Y	N. Holl. 1805.	C s.p Bot. mag.
5516 acuminate Lk.	acuminate	■■■■■ 8 my.jl Pa.Y	N. Holl. 1816.	C s.p
5517 discolor Ven.	two-colored	■■■■■ 5 my.jn Crea	N. Holl. 1814.	C p.1 Vent mal
5518 intermedia Sieb.	intermediate	■■■■■ 4. . . .	N. Holl. 1825.	C p.1
5519 aptlata Lab.	apetalous	■■■■■ 7 myjn Pa.Y	N. Holl. 1803.	C p.1
5520 aspera Sieb.	rough	■■■■■ 6 myjn Crea	N. Holl. 1825.	C s.p
5521 globulosa G. Don.	globulose	■■■■■ 6 ap.jn E	N. Holl. 1803.	C D1
Ceanothus globulodus Lab.				C p.1 Lab.n.h.
5882 capsularis Ct. Don	capsular	M_J or 3 ap.jn F.Y	N. Holl. 1820.	C p.1
Ceanothus capsularis Forst.				
5523 Wendlanditifl Swt.	Wendland's	LJ or 6 ap.jn P.Y	N. Hull. 1810.	C p.1 Sw. a"-.ic*
Ceanothus Wendlanditifl's R. & S.				
6524 spatulata Ct. Don	spatulate	L_J or 4 ap.jn P.Y	N. Holl. 1826.	C p.1 Lab. n. »»!
Ceanothus spatulatus Lab.				
6526 phycifolia Lad.	Phylica-leaved	ft i_J or ■■■■■ ap.jn Pa.Y	N. Holl. 1819.	C s.p Bot. cab. *
6526 fedifolia OM.	Leddim-leaved	ft i_J or ■■■■■ ap.jn Pa.Y	N. HoU. 1824.	C s.p
Sm. APPLB-BEBBV.	(J. J. LabiUardiere, a French botanist.)	Pittosporea-		
climbing	4 LJ or 12 jn.au	C	N. S. W. 1790.	S B p Bot.
changeable	J or 8 jn.s	P	N. S. W. 1795	S sp Bot.
long-flowered	4 J or 20 jn.s	C	V. Di. L. 1810.	S sp Bot.
spindle-shaped	Jj t_j or 8 jn.au	B	V. Di. L. 1823.	S sp Lab.n.
narrow-leaved	1 j or 12 jn.au	Crea	N. HoU. 1820.	C sp
small-flowered	1 j or 12 jn.au	R	N. Holl. 1825.	C s.p

(*iMa*, an olive, *dendron*.ftxag. *Celatirtne**. 5.-8.
« U ... G.v Mauritius 1771. C p.l Jac.ic.1.48
or 3jn.au G.w N. Holl. ... C s. ... Gw N&w 179&c J
? f. ? c N h 2 4 1
c xylocarria Kt. CD or 3 F T Antilles 1816. C s.l.p Vén.ch. 23

DIOSMA.	(Diw, diTine, ome, smellO	Rytdeee.	25.-42.
opposite-leaved * <u>m</u> <u>i</u> <u>j</u> or	3 mrjl	W C.G. H. 1752.	C p1 Com. r. 1.1
succulent <u>W</u> <u>u</u> <u>l</u> <u>j</u> or	3 apjn	W C.G. H. 1752.	C p1 Wen. c. 1.1
Cypress-leaved «L L J or	lijnjl	Pk C.G. H. 1790.	C p1 PI. al. 279.2
liticear-leaved M- i- Jor	mr jl	W C.G. H. 1800.	C p1
pectinated «-l- or		B C.G. H. 1812.	C p1
subulate »	apjn	W C.G. H. 1818.	C p1
red » L Jor	mr jl	W C.G. H. 1800.	C p1
four-angled * L Jor	fmy	R C.G. H. 1752.	C p1 Bot. rep. 451
slenderest »	jlau	W C.G. II. 1789.	C p1
hairy-leaved L Jor	apjn	W C.G. H. 1820.	C p1
twiggy V Jor	apjl	Pk C.G. H. 1731.	C p1 Com. r. 3.3
Heath-like L Jor	mr jl	W C.G. H. 1820.	C p1
capitate * L Jor	mr jl	W C.G. H. 1756.	C p1 Bot mag. 2338
long-leaved * L Jor	2 myjn	D C.G. H. 1790.	C p1
dotted * L Jor	2 mr jl	W C.G. H. 1823.	C p1 Wen. c. 1.19
heart-shaped CL i- or	Hapl. W	W C.G. H. 1823.	C p1
scaly tt- i- or	1 apjl	W C.G. H. 1823.	C p1
corymbose tt- i- or	1 st apjl	W C.G. H. 1818.	C p1
most fetid • L Jor	2 myjl	W C.G. H. 1818.	C p1
hybrid ft. L Jor	2 mr jl	W C.G. H. 1824.	C p1
Broom ft. L Jor		C.G. H. 1823.	C p1
delicate «-l- or	i»51!	W C.G. H. 1823.	C p1
Furze-like		C.G. H. 1823.	C p1
	1 apjl		
	(Dicha, without, osme, smell)		Ruthcex. 1.
* LJ or	2 apjl	W C. G. H.	- C p1 Jac. col. 3.20.1

pretty	L	f.s	P	C O H	S	P-!	B-o-t-mag.	1357
proacMeaved	L	jLau	W	r ^ «	J2?*	P ¹	Bot rep.	33
	L	apjn	P	CG.a	174.	F	p.1 Bot cab!	210
	L			co a	IT?!"	r.P!	y<-*i>	
illiated	CZ		P	CG H	S *	Wen.		
proliferous	L	2 apjm	W	CG.H.	1790.	C	p.1 Wen.c.3.77	
wrinkled	L	2 apjn	W	C. O. H.	1790.	C	p.1	
ambiguous	L	2 apjn	W	C G. H.	1810.	C	p.1 Bot cab.	461
cuminate	L	2 apjn	V	C G. H.	1812.	G	p.1 Wen. c.	1.28
bearded	L	2 apjn	W	C G. H.	1810.			
imbricated	L	3 apjm	Pk	C G. H.	1774.	8	jjl	Wei.c. 1.9
clothed	L	2 apjn	W	C G. H.	1824.			
fix-leaved	L	2 apjn	W	C G. H.	1825.	C	p.1	
	L			C G. H.	1826.			

- | | | | | | | | | |
|--|--|--|-----------------------|---------|-----|-----------|------------------------|----------------------------------|
| S Venenatidna R. & S. Venenat's | ft | LJ or | 2 | apjn | E | C. G. H. | 1794. | C Lp |
| S exsiccata Dec. | dried up | * i_J or | 2 | apjn | V | C. G. H. | 1794. | C Lp |
| 5598 hispida W. | hispid | tt. Jor | 1 | jn.au | V | C.G. H. | 178a | C p.l |
| 5599 brevifolia Lam. | short-leaved | ft u.J or | 2 | apjn | V | C. G. H. | 1818. | C Lp |
| 5610 reflexa Lk. | reflexed-teaved | * i_I or | 2 | apjn | V | C. G. H. | 1820. | C Lp |
| 5601 Brun'ades Lk. | Brunia-like | ft Jor | 2 | apjn | V | C. G. H. | 1820. | C Lp |
| 678. SP1RANTHE ^X RA St. Hel. | SPIRANTHERA. | (<i>Speira</i> , a spiral, <i>anthera</i> , an anther.) | Ruthcea. | 1. | | | | |
| 5602 odoratissima St. Hel. | most odorous | * O or | 6 | ... | W.R | Brazil | 1823. | C p.l N. M.r. 11.31 |
| Terpn&tliusjasminiodbrus Nee* | | | | | | | | |
| *679. PITTO'SPORUM Ban. | PITTOSPORUM. (<i>Pitloo</i> , to pitch, iporof, feed; in pitchy fluid.) | Pittospre* | 13—17. | | | | | |
| 5003 coriaceum H. K. | coriaceous | ft tJor | 10 | my | B | Madara | 1787 | L P.p |
| 5604 viridiflorum B. At. | green-flowered | ftt Jor | 6 | myjn | G | C. G. H. | 1806. | C p.l Bot mag. 1(81) |
| 5605 Tobria H. K. | Tobiru | fti Jor | n' | mr.au | W | China | 1804. | C p.l Bot. mag. 131 ⁶ |
| 5606 undulatum B. Rep. | waved-leaved | ft tJor | | fjn | W.v | N. S. W. | 1789. | C e.p. Bot reg. 16 |
| 5607 revolutum H. K. | revolute | fti tor | | f.ap | Y | | 1795. | C s.p. Bot. reg. 186 |
| 5608 fulvum Rud. | fulvid | * tJor | | ap.my | Y | N. a W. | 1820. | C r.m Lin. tr. 10. i.0 |
| 5609 tomentosum Bonp. | woolly | ftt Jor | | ap.o | Y | N. Holl. | 1824. | E r.m Bon. nav. 21 |
| 5610 ferrugineum H. JC | rusty-leaved | ftQor | | f.my | Y | N. Holl | 1787. | G s.p |
| 5611 hirtum W. en. | hairy | » Jor | | my | Y | Guiana | 1822. | C r.m |
| hirsutum Lk. | | | | | | Canaries | | |
| 5612 Ander&nfis Fis. | Anderson's | ft iJor | L | myjn | — | N. Holl. | 1820. | C r.m |
| 5613 tenuifuhum Gae. | thin-leaved | ft Jor | L | myjn | — | N. Holl. | 1820. | C r.m Lam. il. 143.1 |
| 5614 expense Nois | expensive | ft^ Jor | L | myjn | — | | 1820. | C r.m |
| 5615 mauritiuum Lo. C. | Mauritius | ftt Jor | L | myjn | — | Mauritius | 1825. | C r.m |
| 680. LASIOPETALUM Sm. | LASIOPETALUM. (<i>Lasiost</i> , hairy, <i>petalon</i> , a petal; flower.) | Byttneridce. 2. | | | | | | |
| 5616 parviflorum Sm. | small-flowered | fti Jor | 3 | apjl | W | NN. Holl. | 17910. | C Ch.pri. Hamtr. 1019. 2 |
| 5617 lerrugineum Bud. | rusty | ■\J or | 4 | apjl | W | N. Holl. | 1791. | C s.p Bot mag. 1766 |
| *681. THOMA ^S ^ Gay | THOMASIA. | (<i>M. Thomas</i> , a collector of Swiss plants) | Byttnerietcea. | 5. | | | | |
| 5618 purpurea Gay | purple | * Lj or r | 3 | apjl | P | N. Holl. | 1803. | C a.p. Bot mag. 1756 |
| Lasiopetalum purpureum B. M. | | | | | | | | |
| 5619 foliosa Gay | leafy | #i Jor | | apjl | Ap | N. Holl. | 182a | C s.p. Gay.dis.7 |
| 5620 aolankeea Gay | Solanum-like | il Lj or | | apjl | W | N. Holl. | 1803. | C s.p. Bot. mag. 1486 |
| 5621 triphylla Gay | three-leaved | * Ljor | | apjl | Ap | N. Holl. | 1824. | C s.p. Lab. n. h. 1.88 |
| Lasiopetalum triphyllum Lab. | | | | | | | | |
| 5622 guerciflia Gay | Oak-leaved | ■■■ or | 3 | apjl | P | N. Holl. | 1803. | C s.p. Bot mag. 1485 |
| 682. SERI'NG/il Gay | SERINGIA. | (itf. <i>Seringe</i> , a Swiss botanist.) | Byttnerihcea. | 1. | | | | |
| 5623 platyphyl lla Gay | broad-leaved | * LJ or | 12 | apjl | W | N. Holl. | 1802. | C Bp Meth. m. 7 |
| Lasiopetalum arborescens H. K. | | | | | | | | |
| 683. BYTTNE'R/i Loe. | BYTTNERIA. | (<i>B.S.A. Buttner</i> , prof. bot. at Gottingen.) | Byttneridce*. 8.—2a | | | | | |
| 5624 dasypylla Gay | thick-leaved | * i_J or | 3 | jnjl | W | V.Di.L. | 1780. | C p.l Bot rep. 603 |
| Commerb ^{ns} s dasypyphylla B. Rep. | | | | | | | | |
| 5625 hermanni/oo/o Gay | Hermannia-lvd | L J or | | | | N. Holl. | 1823. | L p.l |
| 5626 herbaria Rot. | herbaceous | ff (Z2 — | | jnjl | | E. Indies | 1820. | p.l Box. cor. 1. 9/d |
| 5627 grandifolia Dec. | large-leaved | • • un | | | | E. Indies | 1823. | p.l |
| 5628 cafts/jevelia Jac. | Catalpa-leaved | * • un | | | | Caracas | 1823. | L p.l Jac. sc. 1.46 |
| 5629 cordata Lam. | cordate | * CJ un | | | | Peru | 1824. | p.l Gay dis. 5. ISO. |
| 5630 scabra L. | rough-leaved | tt. cu | | | P | W. Indies | 1793. | p.l Oav.dis.5.148.1 |
| 5631 microphylla L. | small-leaved | tt. CD cu | | | W.p | S. Amer. | 1816. | L p.l Cav.dis.5.148.8 |
| 684. A YEN ■ E. | A YENIA. | (The Duke D'Ayen, of Koailas.) | Malvacee. 2.—6. | | | | | |
| 5632 pusilla L. | small | £ O cu | 1 | jl.s | P | Jamaica | 1756. | C r.m Mil. ic. 79.118 |
| 5633 laevigata Surx. | smooth | a. • cu | 2 | ... | S | Jamaica | ... | C r.m |
| 685. CALODE'NDRON Thun. | CALODENDRON. | (<i>Kalos</i> % beautiful, <i>dendron</i> , a tree.) | Ruthcee. 1. | | | | | |
| 5634 capense Thun. | Cape | 2 LJ or | 40 | ... | Pk | C. G. H. | 1789. | C Lp H. n. h. 4.22 |
| 686. TODDA ^S LIA J. | TODDALIA. | (<i>Kaka Toddali</i> , its Malabar name.) | Terebinetvdetv. 3.—4. | | | | | |
| 5635 aculeata Pers. | prickly | ft O or | 6 | ... | W | E. Indies | 1790. | C s.p. Lam.il. 139 |
| Scopoliart aculeata Sm. | Paullinia asia'tica L. | | | | | | | |
| 5636 paniculata Lam. | panicled | ft | • un | 6 | ... | W | I. France | 1819. |
| Scopolia incrmis Sm. | | | | | | | C s.p. Lam. il. 139. > | |
| 5637 angustifolia Lam. | narrow-leaved | ft | • or | 6 | ... | W | Mauritius | 1824. |
| Rubinia angustifolia Spr. | | | | | | | C p.l | |
| * 687. ESCALLO ^N /ymlutisEacAtLOMA. | (<i>Ecatton</i> , a Spaniard, and American traveller.) | Eseallonuz. £—* | | | | | | |
| 5638 discolor Mytis | sundry-colored | ft CD'or | 6 | ... | W | S. Amer. | 1820. | C Lp Veil. ch. 5m |
| 5639 floribunda //, \$ B. | flowery | ft O or | 6 | ... | W | S. Amer. | 1827. | C Lp |
| 688. RITYSCHL4 Jac. | UUYSCIIA. | (<i>F. Ruy*ch</i> , M. D., a celebrated Dutch anatomist) | Maregravietcea. I. | | | | | |
| 5640 clusup/o/ia Jac. | Clusia-leaved | ft O or | 4 | ... | P | W. Indies | 1823. | C p.l Jac. am. 51. < |
| * 689. BURSA ^R RIA Cau. | BURSARIA. | (<i>Buna</i> , a pouch.) | Pittospbre. | 1. | | | | |
| 5641 spinosa Cav. | thorny | » L jor | 10 | aud | W | N. S. W. | 1793. | C s.p. Botmag.J7e/ |
| 63a CEDREXA L. | BASTARD CEDAR. | (<i>Cedrus</i> , the cedar tree: aromatic resin.) | Cedritete. 3-T | | | | | |
| 5642 odorata L. | sweet-scented | f a t S O | | | Pk | W. Indies | 1739. | C l.p. |
| 5643 Todna Rox. | Toona | | | | | E Indies | 8823. | C a |
| 5644 velutina Dec. | velvety | | | | | | | |
| 691. HOV&SIA Thun. | HOVENIA. | (ID. <i>Hoven</i> , a senator of Amsterdam.) | Rhdmnete. 2. | | | | | |
| 5645 dlcis Thun. | sweet | li_jfr | 8 | jl.au | W | China | 1812. | C pJ Bot mag. a- |
| acerba Lindl. | | | | | | | | |
| 6646 ina^ualis Dec. | unequal | LJ fr | 10 | — | W | Nepal | 1823. | C p.l |
| dulcis D. Don and Rox. | | | | | | | | |
| pubescens Swt. | | | | | | | | |
| * 692. BRXTN/il L. | BRUNIA. | (C. <i>Brun</i> . a traveller into the Levant and Russia.) | Brynnitec. | 24.—" | | | | |
| 5647 JieVis Thun. | smooth | — Jpr | 2 | jn au | W | C6.H. | 1822. | Pj Brey. cl. 22.° |
| 5648 nodiflora L. | knot-flowered | ft Jpr | 6 | jlau | W | C. G. H. | 1786. | Pj |
| 5649 globosa Thun. | globose | ft L Jpr | 2 | J n a u | W | C. G. H. | 181&. | p. |
| 5650 deiista Thun. | hurnttipred | tt. i_J or | 1 | jn.au | W | C. G. H. | 1804. | pi |
| (5651 lanuginosa L. | woolly | ft L Jpr | 3 | jn.au | W | C. G. H. | 1774. | pi |
| 5652 comsda Thun. | tufted | • LJ pr | 3 | jlau | W | C. G. H. | 1820. | p. |
| 5653 verticillata L. | whorled | ft uJ or | 3 | jn.au | W | C. G. H. | 1794. | Pj |
| 5654 squarrose Thun. | squarrose | ft J J pr | 2 | jl.au | W | CG.H. | 1820. | Pj |

PENTANDRIA MONOGYNIA.

CLASS V.

5730 uliginosa Bes.	swamp	I ap.my P	Carinthia 1823.	D co
5731 rotundifolia Mx.	round-leaved	\ ap.jn P	N. Amer. 1817.	D pl
573M asarifolia Ph.	Asarum-leaved	i ap.jn W	N. Amer. 1820.	D pl
5733 dentata Ph.	toothed	t ap.jn B	N. Amer. 1817.	D pl
5734 clandestina Ph.	clandestine	i my Br	Pensylv. 1800.	D pl
5735 blanda Nut.	charming	ft my.jl W	N. Amer. 1802L.	D pl
5736 ovata Nut.	ovate-leaved	i ap.jn P.D	N. Amer. 1783.	D pl
5737 primulifolia Ph.				W. h. b. 24'
5738 pnummifolia Gro.	(not of Pur*)	Priraroae-lvd	3t A or	I ^P i^n p.a.B
5739 Selkirk* GoL	Alleghany	A	1 ap.jn B	N. Amer. 1824.
5740 hirta L.	Selkirk's hairy	A B'	* juijl B	N. Amer. 1823.
5741 humilis ILS/S.	humble	A or	i ap.my B	England ch. so.
5742 hirsuta R.S/S.	hirsute	Alor	& ap.jn W	Mexico 1824.
5743 collina 2te*	hill	A or	i ap.jl B	Bohemia 1820.
5744 campestris Bieb.	field	A or	1 mr.my B	Poland 1822.
5745 palustris Smi.	marsh	A or	i ap.my P	Tauria 1824.
574H Schmidtia Schult	Schmidt's	A or	4 my.jn B	Britain mos.b.
5747 pyrenaica Ram.	Pyrencean	A pr	£ my.jn B	Austria 1821.
5748 odorata L.	sweet-scented	A	ap.jn B	Pyrenees 1817.
1 purpurea			mr.my P	Britain sh.apl.
2 alba			mr.my P	Britain ch.pl.
3 corylea			mr.my P	Britain d.p.
4 purpurea plena	hluc-flowered	mt A '/*	mr.my B	Britain gard.
5 alba plena	double-purple	9n A fra	mr.my P	Britain gard.
6 corylea plena	double-white	ft? A fra	j mr.my W	Britain gard.
7 pallida plena	double-blue	4U A fra	£ mr.my B	Britain gard.
5749 niplina Jac.	Alpine	Ar A fra	i mr.my Fa.B	Britain gard.
5750 japonica Lan.		*r A or	*my.jn D.P	Austria 1823.
odorata Thun.		SW A	* ap.jn B	Japan 1818.
5751 palmaris Ham.	palm	A or		Jac. au. 242
5752 cantha L.	dog's	A w	ap.jl Y	Nepal 1825.
5753 sylvestris Kit.	wood	A or	\ ap.jn B	Britain hca.
5751 Kx6keri Gm.	Kroker's	A or	i my.jn Pa. R	Hungary 1820.
5755 neglecta Schmidt	neglected	A or	1 my.jn P.B	Siberia 1820.
5756 Alhuni Pio	Allioni's	A or	i my.jn B	Crimea 1821.
arenaria Dec.	dwarf		femynj Pa.B	France 1823.
5757 phimia Vil.				D p.1 Piovi. 1. 2
lancifolia Thore, ericetorum Schr.				Vil.stras.5
5758 glauca Bieb.	glaucous	ifc	i my.jn P.B	Poland 1822.
5759 sarmentosa Bieb.	twiggy	%	* juijl B	Caucasus 1824.
5760 lactea E. B.	milk-colored	jfc	, i my.jn £	England moi. h.
5761 Kuppit All.	Ruppius's	A pr	i my.jn B	Italy 1822.
5762 HornemannsAnaR.&S.	Hornemann's sjt	A or	i my.jn B 1819.
stricta Horn.			1 my.jn Crea	Germany 1817.
5763 joersicifolia Roth	Peach-leaved	st A or	1 my.jn L.B	AI.ofEur. 1683.
5764 montana L.	mountain	i ^ o	i juijl Pa.B	Baltic 1820.
5765 littorkhs Spr.	shore	5t A ot	1 juijl B	Siberia 1823.
5766 Langsdorfii Fis.	Langsdorf's	^ A ot	li my.jn B	Croatia 1820.
5767 stagn^na Kit.	bog	3t A or	... Teneriftc	Teneriftc 1817.
5768 Broussonetiiana R. & S.	Broussonet's	At or		D p.1
5769 hastata Mxen.	hastate	^	my.jn	Carolina 1823.
5770 Nuttallit Ph.	Nuttai's	^ A or	my.jn	Missouri 1812.
5771 triparita Cita Ell	three-parted	^ A or	ij	N. Amer. 1823.
5772 debilis Mx.	weak	^ A or		N. Amer. 1820.
5773 valderia TV.cn.	Valderian	& A or		Mt. Cenis 1805.
5774 cenisia All.	Mount Cenis	^ A or		Mt. Cenis 1759.
5775 nummularifolia Alt.	Moneywort-lvd	^ A or		S. France 1820.
5776 canadensis L.	Canadian	^ A or		N. Amer. 1783.
5777 striata II. K.	streaked	^ A or		N. Amer. 1772.
5778 ochroleptica Lewis'	yellowish-white	^ A or		N. Amer. 1800.
5779 pubescens H. K.	pubescent	^ A or		N. Amer. 1772.
5780 mir-4bilis W.	wonderful	je A		Germany 1732.
5781 radicans Dec.	rooting	%f A		Carolina 1823.
5782 biflora L.	two-flowered	^ A		Al.ofEur. 1752.
5783 uniflora W.	one-flowered	^ A		Siberia 1774.
5784 decumbens Thun.	derumbent	je		C. G. H. 1819.
5785 arboriscens W.	arborescent	a		Spain 1779.
5786 saxatilis R. % S.	rock-inhabiting	^		Portugal 1824.
saxatilis Hqfg.				D p.1
5787 tricolor L.	thre-colored			Britain corn fi. S co Eng. bot 1287
5788 barbatica Kit.	Banatian			Germany 1820. S co
5789 arvntis Sib.	lorn.rield			Britain S co
5790 eriocarpa Dec.	woolly-fruited			N. Amer. 182a D p.1
5791 prostrata A *5.	prostrate			Tcenenff 1824. D p.1
5792 declinata F. AX.	turncl aside			Pannonia 1818. D p.1
5793 Kitaibclicina K. & S.	Kitaibcl's			Pannonia 1820. D p.1
5794 Selkirk/7 GoL	Selkirk's			N. Amer. 1822. D p.1
5795 bicolor Ph.	two-colored			N. Amer. 1818. D p.1
5796 altaica Ker	Altaic			Siberia 1805. D co
2 purplea	purple			Bot.reg.54
5797 rothomagensis De*.				France 1783. D co
5798 saxatilis Schmidt	Rouen			Bot mag. 1498
lutea Bes.	rock			Bohemia 1810. D co
5799 sudetica W. en.	Sudetic			D p.1
S800 lutea.E.A	yff			Germany 1805. D eo
5801 grandiflora L	great-flowered			Britain m. pas. D p.1
5802 ortades Bieb.	oreads			Switzerl. ... D p.1
5803 Pallas, Fis.	Pallas's			Tauria 1818. D p.1
5804 chrysanthia Fis.	yellow-flowered			Siberia 1818. D p.1
5805 amecnia Sumes	pleasing			Germany 1820. D p.1
5806 Zdys/i Wtl.	Zoys's			Scotland tnoun. D p.1
5807 calcarcta L.	spurred			Carinthia ... D co
				Jac. c.4 11*1
				Switzerl. 1752. D p.1

ORDER I.

PENTANDRIA MONOGYNYIA.

5808 corn&ta <i>L.</i>	honed	■ A or ■ A or	I myjn B	Pyrenees 1776. D p.1	Bot.mag.791
5809 Vmarstan <i>i</i> R. & S.	Villars's	■ A or calcarata <i>Vil.</i>	i j n j l B	Vallesia 1830. D p.1	
5810 gracilis <i>Sib.</i>	slender	■ A or	J j n j l P	Greece 1817. D p.1	Fl.gr.S2S
702. ERPE ^X TION <i>Swt.</i>	SPURLESS VIOLET.	(<i>Erpetos</i> , trailing, ton, a violet)		<i>Vwlani*</i> . 2.-3.	
5811 reniWrmis <i>Suit.</i>	reniform	J^ Alpr	i m y . o P3	N. Holl. 1823. D sp.1	Sw. fl. gar. 170
nolaAederacea&fla <i>finesZ</i> w.					
581S//ederaceaG.i)o»	Ivy-leaved	KW-Alpr	ft my.o P.B	N. Holl 1823. D sp.1	Lab.n.h. 1.91
Hola Aederacea <i>Lab.</i>					
703. SG?LEA Ging.	SOLBA. (TV. Sole, author of an essay on the genus <i>Mentha</i> .) <i>natorite</i> . I.				
5813c6ncolorGinff.	self-colored	£ A or	1 j n j l G	NT Amer. 1788. D p.1	Lin.tr.28
"tola cõcolor Forst.					
704. POMB'A'lil Van.	POMBALIA. (<i>Marquis de Pombal</i> , a Portuguese statesman.) <i>Fiolarieat</i> . 1. --S.				
5814 Itiba Ging.	Itulu	S E l m 1 jl	W Brazil	1822. S co	Bot mag. 2453
Ionfdium <i>Ipecacuânhaf</i> B M.					
705. ION'DIUM Ven.	IONIDIUM.	(<i>Ion</i> , a violet, <i>eidos</i> , like; similarity.)		<i>Violar&a.</i> 2.—30.	
5815 capcnsc B. \$ S.	Cape	■ j ■ 1 myjl W	C. G. H. 1824. C l.p		
Hola capensis <i>Thun.</i>					
wlSstrictum <i>Ven.</i>	strict	■ QS or	1 myjl W	W. Indies 1824. S p.1	
3>17 Sprengelcicum R. & S. Sprengel's		■ A) or	1 myjl W	Pensylva. 1818. S p.1	
S&dea atricta Spr.					
°°18 polygalasfbium Tun.	Polygala-lvd	■ LJ or	1 ap.au G.Y	S. Amer. 1707. C Lp Vcn. mai. 87	
KDL» stiff verticillata Spr.					
•819 verbcnâccum //.*JR Vervain-like		■ EJ or	i my.jl Pa.B	Mexico 1823. S Lp H. & B. n. 5.407	
706. KOISSETTFA H. % R	NOISETTIA. (IS. C. Noisette, nurseryman at Paris.)			<i>Violariiee.</i> 1.—3.	
5820 longifilia II. \$ A	long-leaved	tt.CJor 1 ...	Cra Cayenne	1824. S p.1	H.&B.n.5.429.11
Ionidium longif&lium B. # S.	<i>Vola</i> longiflia <i>Poir.</i>				
* 707. CALY'PTRION Ging.	CALYPTBION. (<i>Kalyptra</i> , a veil, <i>ion</i> , a violet)			<i>Violariea.</i> 1.—6.	
5821 Aubletij Ging.	Aublet's	L CD or	ti ...	Crea Guiana 1823. S p.l	Aub. gui. 2,319
■ ^ H o l a Hybanthus iteA.	Ionfdium Hybaanthus	ft £S.		" * . - .	
"822 pyrifolius Afar*	Pyrus-leaved	tt.[Z]or 3	Brazil	1822. S p.1	Martbr.)C.
708. ALSODE'A Thou.	ALSODEA. (<i>Alsodes</i> ,leafY.)			<i>Violarticc.</i> 2.-7.	
■ Eucliflora Thou.	few-flowered	* C J or 6 ...	W Madagas.	1824. C p.1	Thou. af. 17
tifulia Thou.	broad-leaved	* Q - 6 ...	W Madagas.	1824. C p.1	Thou. af. 18.8
709. CERANTHE ^X RA Beam.	CERANTHERA. (<i>Keras</i> , a horn, <i>anthera</i> , an anther.) <i>VitOariee.</i> 1.—ft				
■ »5 subintegritolia Jtfl<<. sub-entire-lvd ft • or	6 jnjl W	Guinea	1824. C 61	Beau. ow. 2. G6	
a ■ ■ PHY'LICA L.	PHYLICA. (<i>Phyllikos</i> , leafy j copious evergreen foliage.)			<i>Bhdmneaf.</i> 45.-52.	
■ ■ ramostssima Dec.	branchiest	i t J or 16	my.jn W	St.Helena 1810. C p.1	
\$27 stipularis L.	stipular	ftuJor 3	my.s W	C. G. PI. 1786. C p.1	
■ J28 parviflora i.	small-flowered	ft LJ or	ap.jl W	C. G. H. 1790. C p.1	
ggJericoldesL.	Heath.like	ft i J or	3 ap.s	C. G. H. 1731. C p.1	
USOglabratatmn.	smooth	ft i J or	2 my.jn W	C. G. H. 1817. C p.1	
5831ace,t-a■	chaffy	ft LJ or	3 myjn W	C. G. H. 1820. C p.1	
■ ■ ■ Serg.	wool-bearing	ft i J or	3 n W	C. G. H. 1774. C p.1	
anceolata Thun.	lance-leaved	ft i J or	3 ap.my W	C. G. H. 1790. C p.1	
blcolor Z.	two-colored	ft i J or	2 myjl W	C. G. H. 1817. C p.1	
■ Linda 27<>.	side.flowering	ft uJ or	2 myjn W	C. G. H. 1817. C p.1	
imwrbisL.	beardless	ftLjor SJmyjl W	C. G. H. 1824. C p.1		
a M	southern	■ i J or 6	apjl W	C. G. H. 1818. C p.1	
dwarf	dwarf	■ i J or 1	myjn W	C. G. H. 182a. C p.1	
■ hairy	hairy	■ i J or 3	apjl W	C. G. H. 1820. C p.1	
villus	villus	■ i J or 2	my W	C. G. H. 1790. C p.1	
papillose	papillose	■ i J or 3	myjl Pa.Y C. G. H. 1820. C p.1	Seba.2.49.5	
Rosemary-lvd	Rosemary-lvd	■ i J or 3	W C. G. H. 1815. C p.1		
cylindrica	cylindrical	■ i J or 2	ap.au W	C. G. H. -	
Exeglaria WnL	lofty	I u J or 16	my.jn W	C. G. H. 1823. C p.1	Wen. c. 1
Horizontalis Vcn.	horizontal	■ i J or 2	myjn W	C. G. H. 1830. C p.1	Ven. mal. 57
f&HSplum&saJ*.	feathered	■ i J or 2	mr.my W	C. G. H. 1752. C p.1	
Wsquarrosa^u.	square	■ i J or 2	au.n W	C. G. H. 1800. C p.1	
Commelin's	pubescent	■ i J or 1	myjn W	C. G. H. 1800. C p.1	
headed	headed	■ i J or 2	f.ap W	C. G. H. 1774. C p.1	
globosa	globosa	■ i J or 1	my.au W	C. G. H. 1800. C p.1	
callous-fcatvrf		■ i J or 3	myjl W	C. G. H. 1800. C p.1	
		■ i J or 1	mr.ap W	C. G. H. 1774. C p.1	
4)803 *Dicata L.	spiked	ftLjor 2	n.d W	C. G. H. 1774. C p.1	Lam.iL127.3
gSJcordatai.	heart-leaved	■ i J or 2	myjn P.Y	C. G. H. 1789. C p.1	Com. rar. 62. IS
L.	Myrtle-leaved	■ i J or 3	:: D.Y	C. G. H. 1816. C p.1	
eduies Desf.	Box-leaved	■ i J or 2	my.s W	C. G. H. 1759. C p.1	
>ichta Lod.	Lcdum-leaved	■ i J or 1	S myjl W	C. G. H. 1820. C p.1	
ot.	Olive-hkc	■ i J or 1	S myjn W	C. G. H. 1800. C l.p	
rosmarinifolia W.en.	diccios	ft i J or 2	my.s W	C. G. H. 1817. C p.1	
Orientalis LK.	axillary./fwd	■ i J or 2	S myjn W	C. G. H. 1812. C p.1	
redinata WnL.	oriental	fti Jor 3	myjn C. G. H.	1820. C p.1	
W.	recline	ftl Jor 3	myjn C. U. H.	18123. C p.1	
Thun.	acuminate	■ i J or 2	myjn C. G. H.	1819. C p.1	
	paniculate	■ i J or 3	myjl C. G. H.	1817. C p.1	
	trichotomous	ft i J or 3	myjl C. G. H.	1818. C p.1	
	Thyme-leaved	ftL Jor 3	my jl C. G. H.	1820. £ Pl	
	tree	■ i J or 16	myjl Tris.d'Ac. 1817.		
	Pine-leaved	ftl Jor 2	my W	1789L ▷ p.1	
	racemose	ftujor 2	my.s C. G. H.	1790. C p.1	
	imbricated	-	1 au.u W	C. G. H. 1801. C p.1	

711. CRYPTANDRA Sm. CYPTANDRA. (<i>Kryptos</i> , hidden, <i>oner</i> , a man; concealed stamens.) <i>Rhdtnnece</i> . 4. BSJlmcifbYiaSm.	Heath-leaved «* J»	Ijmy.jn	W	N. Holl. 1821.	C sp	Lin. tr. 10.18.1
5872 ambra Sm.	bitter	* i _ J cu	14	my.jn	W	N. Holl. 1821.
5873 obovata Sieb.	obovate	* i _ J cu	1 J	my.jn	W	N. Holl. 1825.
5874 spin&cens Sieb.	spinescent	* i _ J cu	1*	my.jn	W	N. Holl. 1824.

712. PLECTRCPNIA L. PLECTRONIA. (<i>Plectron</i> , a cock's spur; tree with large spines.) <i>Bhdmneat</i> . L — 2.						
5875 corymbosa L.	corymbose	i u.J or 20	...	W.a	C. G. H. 1816.	C p.l Bur. af. 94

714. CONOCA'RPUIS Jac.	BUTTON TREK.	(<i>Konos</i> , a cone, <i>karpoo</i> , a fruit.)				
5876 crectus Jac.	upright	tm 10	...	Pa.Y	Jamaica 1752.	C p.l Cat. car. 2, S3
5877 procumbens Jac.	procumbent	or 1	...	Pa.Y	Cuba 1730.	C p.l Jac. am. 52.2
5878 racemibus L.	racemose	• Dor 10	■	W	S. Araer. 1820.	C p.l Jac. am. pic. 79
5879 acutifilius //.	acute-leaved	• 10	—	W	S. Amer. 1824.	C p.l

714. CY'PHIA Res.	CVPIIIA.	(<i>Kuphos</i> , curved; stigma.)				
5880 volubilis W.	twining	Oor 1	—	P.B	C. G. H. 1795.	D l.p

5881 bulbosa Berg.	bulbous	}au.s	P.B	C. G. H. 1791.	D l.p	Bur.af.38.1
5882 Cardamine W.	Cardamine	H ISI pr 3	jl.au	...	C. G. H. 1823.	S l.p
5883 incisa W.	cut-leaved	lQIpr	ijlaii	Pa.R	C. G. H. 1819.	S s.p
5884 />hyteuma W.	Hampton	■ A	Iff	Pk	C. G. H. 1822.	D l.p Bot.reg.fi25

* 715. LIGHTFOOT/jf Herit. LIGHTFOOTIA. (<i>Rev. J. Lightfoot</i> , author of first Fl. Scot.) <i>Campanulaceae</i> . 3.-6.						
5885 oxycooldes W.	Oxycoos-like n.	Aor i jl	B.w	C. G. H. 1787.	C sl	Ex. bot. 2. 69
5886 tenilla Lad.	slender tt.	i J el %	jn.jl	B	C. G. H. 1822.	C p.l Bot. cab. Iu38
5887 subulata /fen/.	awl-/?tfi/f tt.	LJ or £ au	B	C. G. H. 1787.	D 81	Her. ser. 4. 5
Campanula fruticosa JL interrupta Pers.						

■ 716. SHEEP'S SCABIOUS.	(Applied by Pliny to a wild potherb.)					
5888 montana L.	mountain	O pr 1	Jn.jl	B	Britain sap.a	Eng. bot. 882
5889 perdnmis Lam.	perennial	£ A P» 1	jn.jl	B	France 1787.	D co Bot mag. 2198

717. LAGCE*CIA L.	CUMIN.	{ <i>Lagos</i> , a hare, <i>oikas</i> , a dwelling; j seeds in wool)				
5890 cuminoides£.	Cumin-like	Q cu 1	jn.jl	G.Y	Levant 1640.	S co Lam. il 242

718. HE'DERA L.	IVY.	(<i>Hedra</i> , cord, <i>Celt</i> , j the shoots.)				
5891 Rrlx L.	common	or 40 o.n	G	Britain woods.	L co	Eng. bot. 1267
^2 poëtica		or 20 o.n	G	Britain woods.	L co	
"3 arborea	poet's	or 4 ap.my	G	Britain	...	C r.m
4 chrysocarpa	tree	or 8 o.n	G	Greece 1815.	L co	
5892 cananensis IV.	yellow-berried	or 30 o.n	G	Canaries	...	C co
5893 plndula Swz.	Canary	or 20 o.n	G	Canaries	...	Mag. ber. 2.5.1
	pendulous	or 12 ...	Pa.G	Jamaica 1824.	C p.l	Swz. fl. 9
* 719. RRBES L.	CURRENT.	(An acid plant, mentioned by the Arabian physicians.)				
5894 rubrum L.	red	* fir 4 ap.my	G	Britain riv.ba.	C r.m	Eng. bot 1289
2 album	white	I fir 4 ap.my	G	Britain	—	C r.m
3 sylvestre	wild	ft or 4 ap.my	G	Britain	...	C r.m
5895 carpaticum Kit.	Carpathian	A or 4 ap.my	G	Carpathia 1818.	C co	
5896 petra>im L.	rock	Sk or 4 my	R	England moun.	C co	Eng. bot. 705
5897 multiflurum Kit.	many-flowered	tt or 5 ap.my	G	Hungary 1822.	C co	Bot. mag. 2368
5898 spicatuin Robs.	spiked	or 4 ap.my	G	England m.w.	C co	Eng. bot. 1290
5899 trifidum Mx.	trifid	or 4 ap.my	P.Y	Quebec 1823.	C co	Bot mag. 2368
5900 procumbens Pall.	trailing	fr J my jn	P	Oahuaria 1804.	L xns	Palros. 2.b5
5901 rigens Mx.	stiff	or 6 ap.my	G	N. Amer. 1812.	C co	
5902 prostratum Herit.	prostrate	or J ap.my	Y	N. Amer. 1777.	L 81	Her. st. 1.2
5903 alp'num L.	Alpine	or 3 ap.my	G	Britain woods.	C co	Eng. bot 704
5904 tristc Pall.	sad	or 3 ap.my	Y.P	Siberia 1820.	C co	Pal. pet. 10
5905 albincrvx Mx.	white-nerved	ff 4 ap.my	G	N. Amer. ...	C co	
5906 sangufneum/%.	blood-flowered	ft 3 ap.my	Bd	N. Amer. 1817.	C co	
5907 aureum Ph.	golden	or 8 ap.my	Y	Missouri 1812.	C r.m	Bot reg. 125
I fructu Albo	white-fruited	or 8 ap.my	Y	N. Amer. ...	C co	
S fructu rrbro	red-fruited	or 8 ap.my	Y	N. Amer. ...	C co	
5908 missouriensis Hort.	Missouri	or 6 ap	Y	Missouri	...	C co
5909 rrigrum L.	black	fr 5 ap.my	G	Britain m.hed.	C r.m	Eng. bot. 1291
5910 floridum Herit.	florid	or 4 ap.my	Y	N. Amer. 1729.	C co	Di. el. 244.315
5911 laxittorum Ph.	loose-flowered	or 4 ap.my	Y.o	N. Amer. 1812.	C co	
5912 resinum Ph.	gummy	or 3 ap.my	Y.o	N. Amer. 1800.	L co	Bot mag. 1583
5913 glandulifsum R. &c P.	glandular	or 8 ap.my	Y	Peru 1820.	C co	Fl. per. £33. b.
5914 viscosissimum Pit.	most viscid	or 4 ap.my	Y	N. Amer. 1826.	C co	
5915 glaciMc Wai.	frozen	or 4 ap.my	Y	Nepal 1823.	C co	
5916 hirtellum Mx.	roughish	or 3 ap.my	Y.G	N. Amer. 1812.	L s.l	
5917 gracile Mx.	slender	or 4 ap.my	Y.o	N. Amer. 1812.	L r.l	
5918 trifl&rum W.	three-flowered	or 4 ap.my	G.Y	N. Amer. 1812.	L r.l	W.h.b.1.61
	stamineum Horn.					
5919 saxatile Pall.	rock	ft my.jn	G	Siberia 1817.	C co	
5920 oriente Desf.	eastern	or my.jn	G.Y	Syria 1824.	C co	
5921 Diacantha L.	two-spined	or my.jn	G.Y	Siberia 1781.	L r.l	Schm. ar. 97
5922 reclitratum L.	recline	or ap.my	P.a	Germany 1683.	L co	
5923 macracanthum Lo.C.	long-spined	or my.jn	G	C co		
5924 Grossulana L.	Rough gooseb.	Sk fr mr.ap	G	England hed.	C r.m	Eng. bot. 12°
5925 UVa-crispa L.	Smooth gooseb.	ft ft mr.ap	G	England hed.	C r.m	Eng. bot. 2057
5<26 oxyacanthodesL.	Hawthorn-like A	or 3 ap.my	W.Y	N. Amer. 1705.	L co	Di. el. 139.16 ⁶
5927 grossularioides Mx.	Rough gooseb.-lkScf	fr my.jn	G	N. Amer. —	C co	
5928 rac6stre Poir.	lake	or ap.my	Y.G	N. Amer. 1812.	C p.l	

Cyn&bat L.	Dog bramble	A	or 4 ap	G	Canada 1759.	C el Schm. ar. 98
KWO caucasicum Adams	Caucasian	■ fr	4 my.jn	G	Caucasus 1819.	C co
coif* G 5 O N 9 N VI ^ L.	GRONOVI.	(J. F. Gronovius, a botanist of Leyden.)				
5931 scandens L.	climb-ne	LD&cu 6	in if	G.Y	Jamaica 1731.	C pl JAC. id
5932 HumboldtV. &> R.&S. Humboldt's	fl. O cu 8	jl.ad	Y	S. Amer. 1820.	S s.p	

IN THE S		£ ACHYRANTHES. (<i>Achyron, chaff, anthos, (I; envelopes.) Amaranthacee.</i>)		13.—28.
silver	■ ■ ■	1 my.o	& c u y	1713. C ls Boc.sic.16.9
rough	■ ■ ■	3 my.o	Pk India	1751. C ls MiLic.1.11.2
curled	tt. i CU	2 my.o	W.a S. Amer.	1810. S co
twiggy	tt. i leu	2 my.o	W.G W.Indies	1817. S co
branched	i Qcu	W	E. Indies	1824. S co Pluk. al. 334.5
obtuse-leaved	■ ■	2 my.n	Pk.	C l.p.
extending	■ Qcu	6 ap.au	P . . .	C r.m Bot. nag. 830
snowy	tt. J CU	2 my.jl	W Canaries	1780. C r.m
shrubby	tt. CU	6 my.jl	P E. Indies	1820. C r.m
pubescent	tL J CU	U ap.jl	Pk.	1821. C r.m
southern	«. CU	2 Jn.jl	W N. Holl.	182a S co
dichotomous	O cu	1 Jn.jl	G N. Amer. . .	S co
whorled	a. I leu	1 il.au	W C. G. H.	1820. S co

A M A I H . A M I S O 1 H - & B - CHAMISSOA. (*M. Chamisso, who accompanied Kotxebue.*) *Amaranth&ce*. I.—2.*
*W6al«88jina.«A tallest *C Jor 5jl.au Y Jamaica 1816. C co H.&B.n.2.125
Achyranthes altissima L.

» ffl!1!2P^L2? ^ET₁U^R? 2? *^{PHI}!-0XRaus. (*Phitos*, a lover, *xeros*, arid j soIL) *Amaranthace**. 3.-7.
 ffl!S? ¹₁A¹₁18R, ^A₁ checkedered ^{JU}. El cu * jl.o Pk & Amer. ... C r.m Her. par. 15
 ^ /leecrumbe vermicularie /.,
 fQta? ²⁸!!1!1⁸? ^EBrazilia! » tt-Qcu ³ jl.o W Brazil 1790. C r.m Jac. ic 2.346
 »» aggregate if. \$2?. aggregate JUElcu * jl.s W Trinidad 1820. S s.p

24. DESMOCH/ETA Dec.	DESMOCHATA. (<i>Desmos</i> , a bond, <i>chaite</i> , hair.)	<i>Amaranthacea</i> .	a—12.				
<i>atropurpurea</i> Dec.	dark-purple	■■■■■	1 au.o	P	E. Indies	1759.	C Lp R.maL10.59
<i>Achyranthes</i> /appacea J.							
<i>navescens</i> Dec.	flavescens	iQJ cu	1 jl.a	Y	E. Indies	1823.	S co Pluk. al. 83. 2
<i>micrantha</i> Dec.	small-flowered	O cu	1 jn.s	Y	E. Indies	1820.	S co Pluk. al. 82.2
<i>wosstrata</i> Dec.	prostrate tt.	■■■■■	2jl.au	G.p	E. Indies	1793.	D l.p Ru.am.6.11
<i>omentosa</i> R. \$ s.	woolly tt.	O cu	2 jl.s	W	E. Indies	1818.	S co
"	hook-seeded	O cu	1 jl.s	...	E. Indies	1820.	S co
"	globose	iQj or	1 jl.s	Pa.R	R Indies	1817.	S co
"	spreading)£ El cu	3 au.o	W	E. Indies	1823.	C Jp Pluk. al. 260.1
<i>Achyranthes pfitzera</i> L.							

Bo⁵ k. CI¹ ADO¹ ST¹ ACHYS D. *Don* CLADOSTACHYS. (*Klados*, a branch, *stachys*, a spike.) *Amaranthaceaf.* 9.
*** ftutescens *D. Don* frutescens tt. • cu 4 au.n G E. Indies 1777. C sp¹ Ru. am. 5. 83
floen Achyranthes muricata *L.* Chamissoa muricata *Spr.*
»¹ 9 alterniſhuha *Svet.* alternate-leaved OJ cu 1 jLn P E. Indies 1789. S p.l Pluk. al. 260.1
Achyranthes alterniſhuha *W. Desmochs* ² a alterniſhuha *Dec.*

<i>CEBRUM L.</i>	<i>KNOTGRASS.</i>	<i>(<i>Illecebra</i> of Pliny, from <i>Ulico</i>, % to allure.)</i>	<i>Paronychietie.</i>	<i>7.—15.</i>
<i>scillitum L.</i> ***	<i>whorled</i>	<i>M d R 4 il</i>	<i>England</i>	<i>lmts-al S p.l Eng.bot.895</i>
<i>cymosum VU.</i>	<i>sprouty</i>	<i>* O pr</i>	<i>S. Europe</i>	<i>18.U S p.l FLgr.245</i>
<i>chia cymbosa Dtr.</i>				
<i>Poir.</i>	<i>prickly</i>	<i>Jc O pr I jl</i>	<i>G Barbary</i>	<i>1821. S p.l Boc. sic. 20.3</i>
<i>Lich.</i>	<i>white-ended</i>	<i>Jc O cu ^jn.jl</i>	<i>W C. G. H.</i>	<i>1817. S s.p</i>
		<i>O cu \jn.jl</i>	<i>W Peru</i>	<i>1810. S s.p</i>
	<i>Gomphrena-lk</i>			
	<i>clustered</i>	<i>)£ E Jc u i jn.jl</i>	<i>W Brazil</i>	<i>1820. S s.p</i>
	<i>diffuse</i>	<i>if AJ cu 1 jn.jl</i>	<i>W Trinidad</i>	<i>1817. S fl.n</i>

727. ALTERNANTHE	RA Br.	ALTERNANTHERA.	(Stain, alternately fert. and bar.)	<i>Amaranthaceae</i>	E. 10.-25.
coldestf. 26..		Chaff-flower	£ El cu	1 jn.au	W Buenos A. 1732. D Lp Di.cl. 8.7.7
pinbsa Horn.		Polygonum-like	£ iAJ cu	1 jn.au	W America 1731. C r.m Her. par. 17
axillaris Dec.		frutescent	s. i_ or	1 jn.au	W Peru 1820. S s.p Iler.-st. 1.37
		Caraccas	jg OSJu	1 jn.au	W Trinidad 1819. S s.p
		canescent	jg El cu	1 jl.au	W Cumana 1825. S s.p
		silky	£ El cu	2 jn jl	W Quito JK20. S s.p
		procumbent	£ EJ cu	4 jn jl	W Brazil 1818. S s.p
		sessile-flowered	jg OJ cu	} jl.o	Br E. Indies 1778. S r.m R. xnal. 10.11
		fg. \ke-leaved	j£ El cu	1 jn.jl	G S. Amer. 1821. S r.m Jac. am. 60.4
		spinous	O cu	1 my.jn	Y 1823. S r.m

TOM. headed	PARONYCHIA.	(Maladies of finger-nails; mod. qual.)				Paronychia.		12.—20.			
		J	n	a	u	w	S p a i n	1 6 8 3 .	D	p	1 0 b p i . 4 2 0 . 1
		1	jn.au	W	Spain	1812.	D	s.l			
		i	my.jl	W	Brazil	1820.	S	co			
		3	Jn.au	W	Spain	...	D	51	Scop. d.	13	
		1	jn.au	W	Spain	1683.	I	s.l			
		§	injl	W	Italy	1821.	S	s.p			
		i	jnjl	W	Spain	1816.	S	sp	Vil.del.	1.16	
		i	jn.au	W	Pyrenees	1820.	S	s.p			
		1	jn.au	W	Pyrenees	1820.	S	sp	Vil. a. j.	1801, Ic	
		i	l'	W	France	1818.	S	up	Jac. ic.	8.2.1	
	O cu	1	jn.au	W	Bengal	1820.	S	s.p			
						1817.	S	s.p			

{*Chen, a goose, leia, prey.*) Chenopbde*. 1.

tt. i_J cu 1 au.8 G C. G. H. 1758. C r.m

Mx. ANYCHIA.
Mx. forked
Mx. *canadensis* L. ^{led from it's affin' y to} *Paronychia*.) *Paronychiae* ^{* O} *O* *r* *I* *s* *T* ^{N. Amer. 1806.}

<i>JE'RUA</i> Forsk. <i>janata</i> J.	woolly	DDcu Arabic name	E. Indie, 1\$%"-,*
Javanese	« . a c u 2 a . . a u W	£ Indie. 1761 C r m Bur. in 65 "	

732. LESTIBUDE**Sfi* R.Br. LESTIDUDESIA. (*F. J. Latiboudois*, a Flemish bot.) *Amaranthikcete*. 3. —
£993 paniculate *R. Br.* paniced 233 cu 3 jn.s P.Y Jamaica 1733. C r.m SL.jam. 1.91.1

5994 trifgyna *R. Br.* three-styled ITBleu au.o W Senegal 1777. r.m Jac. vin. 3.15
5995 virgata it. *Br.* twiggy .1 leu au.o G — 1815. r.m Jac. ic. 2.339

733. DEERI'NG/i* R. Br. DEERLVGIA. (*Dr. C Deering*, an Eng. bot. author.) *Amaranth^cece*. 2.
5995 celosioides *R. Br.* Cock's-coimb-lk D8 cu 6 au.o W E. Indies 1804. S s.l Bot mag. 2717
5997 indica *Spr.* Indian iQI pr 2 au.n W E. Indies 1804. S p.l
Celosia baccata Retx.

' 734. TRIANTHEMA L. TRIANTHEMA. (*Treis*, three, *anthemon*, flower; disposition.) *PortuMcae*. 3. — 6.
5998 monogyna L. monogynous O cu 1 my.jn P.o Jamaica 1820. S co Her. par. 2.213
5999 decandra £. decandrous O cu 1 jn.jl G E. Indies 1824. S s.p Bur. m. 31.3
6000 obcordata Box. hcnrUcavcd fc* LJ CU i jn.jl G E. Indies 1816. S s.p

735. CELOSIA L. COCKS-COMIL. (*Kelos*, burnt; appearance.) *AmarantMcete*. 14. -i-22.
6001 argentea L. linear-leaved Ty-spikcd rmTorr jn.s0 J-china 1740. ■ r.m Mart cen. 1.7
2 lineans Qlor 1 J-Jl F E. Indies 1714. S r.m It inal. 10.39
6002 cristataX. created mi or 2 jns D.R Asia 1570. S r.m Lam. il. 168.1
1 elata tall QJor 2 jn.s DK Asia 1570. S r.m Kno.del.I.H.5,6
2 compacta compact Qlor 2 jn.s D.R Asia 1570. S r.m
3 flavescent flavescent Qlor 1 jn.jl Y Asia 1570. S r.m
6003 comosa *Retx.* tufted Dj or 1 jn.s Pk E. Indies 1802. S r.m
6004 coccinea L. scarlet O or 5 jn.s P China 1597. S r.m
6005 cernua B. Rep. drooping O. or 3 jl.au P E. Indies 1809. S r.m Bot rep. 635
6006 castrensis L. Cdtns mi or 2 jl.s P E. Indies 1739. S r.m Bar. ic. 1195
6007 Monsnii L. Monson's rm or 3 jl.s W E. Indies 1778. S r.m Pluk.all 1.334.4
6008 nodiflora L. knotted-flwd fmor 2 jl.s G E. Indies 1780. S r.m Jac. vin. 1. 98
6009 pyramidalis Brm. pyramidal rmcu 1 jn.s W E. Indies 182a S s.p Bur. in. 25. 1
6010 margaritacea L. pearly tt-l eu 2 jl.s Y W. Indies 1817. S s.p R. mal.1038
argentea Hort.

6011 nitida Vahl shining O cu 1 JU P Malabar 1706. S SI.jam. 1.91.5
6012 dichotoma Roth dichotomous O cu 1 jn.au Y E. Indies 1824. ■ Wen. her. 2
6013 glauca Wni. grey tt. \ cu 1 jn.au W C. G. H. 1818. S 3.1
6014 echinata/.R hedgehog O cu 1 jn.au P Oroncko 1821. S 8.1

736. OPLOTHE*CA Kut. OPLOTHECA. (*Oplon*, armour, *theca*, a sheath; capsules.) *Amaranthice*<v.2.—3.
6015 floridana Nut. Florida £ A or 3 s W N. Amer. 1824. K p.l Bot. mag. 2603
6016 interrupta Nut. interrupted JU Qfl or 2 jn.au G W. Indies 1733. C r.m Jac. ic. 1.51
Gomphr&nma interrupta W.

737. GOMPHRE'NA L. GLOBE AMARANTH. (*Gop'nphos*, a club; flowers.) *Amaranthdce**. 7. — 26.
6017 gibbosa L. round Oor my.ø Pw India 1714. S r.m R. mal. 10.37
«alba white 0 or 1 jn.o W India 1714. S r.m
6018 perennis L. perennial £ Csl or 2 jl.o P.Y 8. Amer. 1732. C r.m Di. el. 24. 20.29
6019 arborescens X. arborescent n. O or 3 jl.o W S. Amer. 1802. C r.m
6020 lactea Dec. milky white L cu 2 jn.au W S. Amer. 1823. S s.p Dec. mon. 93
6021 lanata R. Br. woolly 01 cu 1 jn.au W N. Hull. 1824. S s.p
6022 amaranthoides Roth Amaranth-like OJ c^u 2 jn.a^u R fde^s 1826. S P Mexico 1826. S s.p Jac. sc. 482
6023 decumbens Jac. decumbent IBI cu i jn.au P

738. LAHA^Y^ R. & S. LAIIAYA. (*M. Lahaye*, a botanical gardener.) *Paronychie**. 10. — 15.
6024 diffusa R & S. diffuse IOJeu ijJL.au W Canaries 1779. S Lp W.h.b..ll
Mellia diffusa W. /Uecbrum divaricatum H. K.
6025 aristata R. 8f.S. awncd s. LJ cu i jn.jl W Canaries 1780. C l.p
M6UPA anstata H. K.
6026 latifolia R. # 5. broad-leaved tt. LJ cu \ jn.au W Tenerife 1810. S s.p
/Uecbrum latifolium W.
6027 stellata R. % S. stellate tt. leu \ jn.au W Guinea 1820. ■ s.p
6ms tenuiflora R. if S. line-leaved i jn.au W S. Leone 1817. ■ s.p
6029 corymbbsa R. # S. corymbose tt. f- cu I jn.au W Ceylon 1823. ■ s.p Bur. zey. 65. S
6030 ^naphalioides R. 8f.S. Gndphahum-lkflL t_J cu W N. Africa 1818. ■ i.p
6031 polycarpoides it. 8f.S. Polycaron-lk tAI cu jn.au P.W Sicily 1817. ■ opp
Hayea polycarpoides Biv.
6032 filsinifbha «& 5. Chickweed-lvd -* O cu ijmau W S. Europe 1817. S s.p Boc. sic. 38
Hayea alsinitha Biv.
6033 minuarto/c5 It & & Minuartia-like Jk Q cu \ jn.au W Spain 1826. S s.p

739. GLAUOT L. BLACK SALTWORT. (*Glavkos*, grey; leaves.) *Saliciirte*. 1.
6034 maritima L. ea IL A ray.jn F Britain salt m. S s.l Eng. bot 13

740. THE^IUM L. BASTARD TOAD-FLAX. (*Thcs*, a labouring servant; mean habit.) *Santaldee*. 6. — S.
6035 /inophyllum L. Flax-leaved ^ A cu i jn.jl W England ch. pa. D p.l Eng. bot. 247
6036 intermedium Schr. intermediate ^ A cu J jn.jl W Germany 1818. S p.l Schk. han.51
6037 montanum Ehrh. mountain ^ A cu i jn.jl W Europe 1817. S p.l Sc.b. j. 119
6038 alpinum L. Alpine £ A cu £ jn.jl W Germany 1814. D p.l Jac.au. 5.416
humifissum Dec.

6039 ebractektum Hay. bractless A cu ijn.jl German 1814. D p.l Sc. b. j. 1800.7
6040 amplexicatle L. stem-clasping IJ cu 4 W C. G. H. 17b7. C s.l

741. COMA'NDRA Nut. COMANDRA. (*Kovie*, hair, *aner*, an anther.) *SantalUceee*. 1.—2.
6041 umbellata Nut. umbellate ^ A cu 1 jn G N. Amer. 1782. JJ p.l Pluk. m. 342.1
Thesium umbellatum L. Hamiltunia umbellata Spr.,

742. LEPTOME^RIA R. Br. LEPTOMERIA. (*Leptos*, slender, *merit*, a part; habit) *Santalucete*. 2. —
6042 acidaliJr. acid «. LJ CU 1 ... W N. Holl. 1823. S p.l
6043 Billardi^rj R. Br. LabiHardiere's ttjeu 1 ... W N. Holl. 1823. S p.l Lab. n. h. 1 J^o
Thesmm drupkeem Lab.

«IS" DAR^INGTOXN/^ Dec. DARLINGTONIA. (*Dr. Darlington*, an American bot) Z,T Dec. leg.
0044 brach^loba Dec. short-podded £ A or 1, s.o W N. Araer. 1803. D p.l
• Acacia brach^loba TV.
6045 glandulosa Dee. glandulous * A or H 5.0 W N. Amer. 1806. Dp.l Vcn.ch.
Mimosa glandulosa Mx. Acacia glandulosa IV.

6018 <i>Bihai Swz.</i>	serrata W.	serrated	• or 12	... Y	Malabar 1824.	C p.l R. mal. 5.48
6046 <i>serrata W.</i>	entire-leaved	J Q o r	12	... Y	Guiana ...	C p.l
745 HELICO^NIAX.	HELICONIA. (<i>Helicon</i> , mountain of the Muses; affinity to <i>Musa</i> .)	<i>Mushcece</i> .	10.—12.			
6018 <i>Bihai Swz.</i>	<i>Bihai</i>	j.g El or	12 jlau	O	W. Indies 1786.	Sk s.p Swz. oh. 96.5.2
6049 <i>caribea Lam.</i>	Caribcan	£ El or	12 jlau	O	W. Indies 1798.	Sk s.p Pl. gen. 59
6050 <i>BAUiia Rich.</i>	Ballia	£ El or	20 jls	O	Maranhao 1823.	Sk s.l Mer. sur. 54
6051 <i>hfamilis Jac.</i>	dwarf	£ El or	6 jlau	S	Caraccas 1798.	D s.p Jac. sc. 1. 48.49
6052 <i>braziliensis Hook.</i>	Brazilian	jg IS spl	8 au	■	Brazil 1820.	D p.l Hook. ex. H! 190
6053 <i>psittacorum W.</i>	weerot-bedked	jg • or	4 au	O	W. Indies 1797.	Sk s.p And. rep. 124
6051 <i>Swartzia R & S.</i>	Swartz's	£ E) or	4 jlau	Y	Jamaica 1800.	Sk s.p Bot mag. 502
6055 <i>psittacorum/3 B.M.</i>	Indian	£ 123 or	4 . . .		Madagasc. 1818.	Sk.s.l Ru. am. 5.62.2
6055 <i>Indica Lam.</i>						
6056 <i>buccinata Box.</i>	whited	■ ■ ..				Sk s.p
6056 <i>dealbata Lo. C.</i>	hairy-flowered	El or	5 jlau	O		& Amer. 1800. Sk s.l

* 746. <i>L. PLANTAIN TREE.</i> (Altered from <i>Maux</i> , Egyp., in honor of <i>Antonius Musa</i> .)	<i>Mushcece</i> .	9.—14.				
paradisaca L.	common paradise	El clt	20 o.d.	Pk	Tropics 1690.	Sk r.m Tr. ehr. 3.18.20
6059 <i>sapi^ntum L.</i>	wise men's Banana	El clt	20 mr o	Pk	Tropics 1729.	Sk r.m Tr. ehr. 4.21.23
6060 <i>nepalensis TV.</i>	Nepal	TClor	6 fmr	Y	Nepal 1823.	Skr.m
6061 <i>ornata Box.</i>	ornamented	El or	5 mr.d	O	E. Indies 182a	Sic r.m
6062 <i>glaucafar.</i>	glaucous	EJor	10 ...	Pk	E. Indies 1824.	Skr.m
6063 <i>inaequata Jac.</i>	spotted	EAior	10 ...	Pk	Mauritius 1818.	Skr.m
GOB* <i>superba Box.</i>	superb	El or	14 jl	Pk	E. Indies 1820.	Skr.m
6064 <i>rosacea Jac.</i>	rosaceous	El or	15 f.my	P	Mauritius 1805.	Sk r.m Bot reg. 706
6066 <i>coccinea Andr.</i>	scarlet	El or	4 mr.d	S	China 1792.	Sk r.m Bot. mag. 1559

* 747. STRELITZ <i>/i*</i> H. K. STRELITZIA. (Queen of Geo. III. . house of Mecklenburg-Strelitz.)	<i>Musuteec</i> .	8.				
august	El or	IS f.my	W	C. G. H. 1791.	Sk p.l Kcr strel. ic.	
queen's	El or	8 ap.my	Y	C. G. H. 1773.	Sk p.l Red. hi. 77, 78	
ovaXe-leaved	El or	8 f.ap	Y	C. G. H. 1777.	Sk p.l Bot.mag. 119,120	
humble	El or	6 my.jn	Y	C.G. H. ...	S p.l	
tneay.stalked	El or	5 fmr	Y	C.G. H. 1795.	S p.l Ker strel. ic.	
narrow-leaved	El or	6 myjn	Y	C. G.H. 1778.	S p.l Ker strel. ic.	
small-leaved	El or	6 my.jl	Y	C. G.H. 1796.	S p.l Ker strel. ic.	
Rush-leaved	El or	6 my.jn	Y	C. G.H. ...	S p.l Bot reg. 516	

CLASS V. ORDER 2.

PENTA'NDRIA DIGY'NIA.

748. MELODrNUS Forst.	MEODINU&	(<i>Melon</i> , an apple, <i>dineo</i> , to turn round; fruit.)	<i>ApocynneG. t.</i>			
6275 <i>scandens Forst.</i>	climbing	%Q • pr I35 jlau	'W N.Cald. 1775.	C s.p Itm. il. r79		
wo xnonogynus Box.	monogynous	i. • or 10 jl	W	E. Indies 1820.	C r.m Bot reg. 834	

- 749. PERIPLOCA £.	PERIPLOCA.	(<i>Periploke</i> , an intertwining; habit)	<i>Asclemidiae.</i> 4.—13.			
77 gr ^o ca L.	Grecian	J or JO jlau	Br Syria	1597.	K d Botreg. 803	
J S t J t	smooth	JL I—I or 6	... G.Y	Canaries 1779.	C p.l Cav. ic. 3 217	
1079 angustifolia/fla.	narrow-leaved	£ or 6	... Y	S. Europe 1800.	C p.l Dec. ic 2.7	
naauntiana Poir.	Mauritian	A_O° r 10	... Br	Malabar 1823.	C p.l R. mal. 9.11	

W f i a S ? ^ 0 1 ? 0 ^ 1 1 - ^ CiYPToTEGA.	(<i>Kryptos</i> , concealed, <i>stcge</i> , a covering.)	<i>Asclepiudece.</i> 2.				
W f i a S ? ^ 0 1 ? 0 ^ 1 1 - ^ CiYPToTEGA.	large-flowered * or 6 jn.jl	Pk India* 1818.	C r.m Bot reg. 435			
W f i a S ? ^ 0 1 ? 0 ^ 1 1 - ^ CiYPToTEGA.	ascanensis Lo. C.	Madagascar a_Q or 10 jlau	... Madagascar 1826.	C p.l		

R.Br. HEMIDEBMUS.	(<i>Hemisus</i> , half, <i>demos</i> , a tie: filaments.)	<i>Aselepihdece.</i> 1.—2.				
Indian	i. • or 6	... G	Ceylon 1796.	C l.p Bur. zey. 83. 1		

W f i a S ? ^ 0 1 ? 0 ^ 1 1 - ^ CiYPToTEGA.	(<i>Asclepidae</i> , Arabic.)	<i>Asclepidae.</i> 3.—5.				
W f i a S ? ^ 0 1 ? 0 ^ 1 1 - ^ CiYPToTEGA.	6 jl	W Egypt	1752.	C 81 Alp. a# 134		
W f i a S ? ^ 0 1 ? 0 ^ 1 1 - ^ CiYPToTEGA.	W India	1816.	C f.l.p W. ph. 1.5.2			

W f i a S ? ^ 0 1 ? 0 ^ 1 1 - ^ CiYPToTEGA.	... N. Holl.	1824.	C p.l			
W f i a S ? ^ 0 1 ? 0 ^ 1 1 - ^ CiYPToTEGA.	... L. J. C. U. 3 J. U. u	W L. J. L. 111 3 up	W. Ph. 1.5.2			
W f i a S ? ^ 0 1 ? 0 ^ 1 1 - ^ CiYPToTEGA.	... L. J. C. U. 3 J. U. u	W L. J. L. 111 3 up	W. Ph. 1.5.2			

W f i a S ? ^ 0 1 ? 0 ^ 1 1 - ^ CiYPToTEGA.	(Mrs. Harrison, of Aighburgh, near Liverpool.)	<i>Atclep&de*</i> .	1.			
or 6 jlau	S Brazil	1825.	C s.l Bot mag. 2699			

W f i a S ? ^ 0 1 ? 0 ^ 1 1 - ^ CiYPToTEGA.	(<i>Stephanos</i> , a without, <i>Stephanos</i> , crown; stamens.)	<i>Aselepidetv.</i> 2.—C				
6091 trillbrus R. Br.	linear	J IAJ or 4 jlau	W C. G. H. 1816.	D p.l		
6092 linearis B. Br.		J IAJ or 4 jlau	W C. G. H. 1816.	D p.l		

W f i a S ? ^ 0 1 ? 0 ^ 1 1 - ^ CiYPToTEGA.	Cynfinchui viminale L.	I. C r.m Alp. ag. 190				
6094 Swartz/dhi/m IL & S. Swartz's	t Ocu 6	W Jamaica 1820.	C s.l SI.jam. 1.131.1			
ylsleplias viminahs Swz.						

W f i a S ? ^ 0 1 ? 0 ^ 1 1 - ^ CiYPToTEGA.	hasrttum	Th. p.M.9.6				

6151 amplxldcaftls Mx.	stem-clasping	8	jLau	R	N. Amer.	1816.	D r.m	Ab. Ins. 7
6152 acuminata Ph.	acuminate	2	jLau	RW	N. Amer.	1826.	D r.m	
/aurifdha Mx.								
6153 cinfcrea Walt.	cinereous	2	jLau	Br	N. Amer.	1325.	D r.m	
6154 incarnata L.	flesch-caforaf	2	jLeu	P	N. Amer.	1710.	D p.l	Bot reg. 250
6155 pulchra Ehrh.	fair	2	jLau	P	N. Amer.	1710.	D p.l	
6156 variegata L.	variegated	A or	4	jLau	W	N. Amer.	'ft	Bot. rang. 1182
6157 curassifica L.	Curassoa	JAI	3	jns	S	S. Amer.	S r.m	Bot. reg. 81
2 alba	white		3	jns	W	S. Amer.	1692.	S r.m
6158 eitrifolia Jac.	citron-leaved		1	jLau	W	S. Amer.	isis.	D r.m Jac. ic. 343
6159 nfvica L.	snowy		3	jLs	W	N. Amer.	1730.	D p.l Bot mag. 1181
6160 parviflora J. AT.	small-flowered		4	jLO	W	N. Amer.	1774.	C r.m Jac. cc. 28
6161 polytachya Walt.	many-spiked		4	jLau	W	N. Amer.	1825.	D r.m
6162 cibilia Mx.	weak		2	jLau	W	N. Amer.	1818.	D r.m
6163 verticillata /.	whorl-leaved		3	jLau	WG	N. Amer.	1751.	D pi Pluk. m. 336.3
6164 quarinfolia Ph.	four-leaved		1	jLau	W.R	N. Amer.	1818.	T r.m Jac. ob. 33
6165 mexicXna Chit.	Mexican		3	jLau	W	Mexico	1821.	C s.p Cav. ic. 1.58
6166 rusea //, & B.	roseate		1	jnau	R	Mexico	1824.	S r.m
t167 angustifolia Lk.	narrow-leaved		3	jLau	W	Mexico	1817.	D r.m
MM virgata Balb.	Flax-leaved		3	jLau	W	Mexico	1820.	C s.p
6169 /infeli i la*	Willow-leaved		2	jLau	W	Mexico?	1817.	C p.l Bot cab. 272
6170 sahifMia Cod.			8	jLau	W	Ceylon	1810.	C p.l W. ph. 2.1795
(>171 asthmatica L.								
Cynanchum Ipecacudintha W.								
6172 alexia Jae.	antidotal		8	--	O	Ceylon	181&	C p.l R. mat. 9.13'''
6173 longifolia Ph.	long-leaved	3	2	jLau	Pa.P	N. Amer.	1816.	I) p.l
6174 tuberosa L.	tuberous-rooted		2	jLs	O	N. Amer.	1690.	I) s.l Bot. reg. 76
6175 rbra L.	red		1	jLs	R	Virginia	1825.	D p.l
6176 decumbent IV.	decumbent	A A or	2	jLau	O	N. Amer.	1731.	D p.l
6177 Linaria Cav.	Toad-flax-tod	E iAl or			W	Mexico	1802.	I) s.l Cav. Ic. 1.57

772. STYLA'NDRA. Nut. STYLANDRA. (*Stylos*, a column, *aner*, a man; *anthers*.) *Asclepidde*, L
6178 pumila Nut. dwarf ^ AJ or \ jn.au G.v N. Amer. 1824. D r.m
Podostigma pubescens Spr. Jsclep'u pedicellata Ph.

•77a GONO'LOBUS Mx.	GONOLOBW.	(<i>Gonia</i> , an angle, <i>lottos</i> , a pod.)	<i>Asclepihde</i> .	19.—28.
6179 maritimus R. Br.	sea-shore	f. CD or	6 jn.jl	G Carthage 1823. C p.l Bot. reg. 931
Cynanchum maritimum L.				
6180 suber'sus R. Br.	cork-barked	Q o i	J.s	G America 1733. C p.l Di. el. 229.296
6181 crispiflbrus R. Br.	curled-flowered	L or	jLau	W.a S. Amer. 1741. C s.p Pl. ic. 216.1
CilHU planiflorus A Br.	flat-flower red	F Tio r	jn.au	G Trinidad 1818. C p.l Jac. am. pic. 81
6183 grandiflcr* R. Br.	large-flowered	S CD or	jl.au	G Trinidad 1826. C p.l Bot. reg. 1053
6184 rostflitus R. Br.	beaked			G Trinidad 1824. C p.l
6185 n'igr R. Br.	black			M Mexico 1825. C p.l Cav. ic. 2.159
Cynanchum nWrumb				
6186 racemosus R. fir.	racemose			Caraccas 1820. C p.l Jac. am. pic. 80
6187 diadematus Ker	crowned			M Mexico 1812. C s.p Bot reg. 252
6188 echinatus Lo. C.	hedge-hog			
6189 carolinianus R. Br.	Carolina			P Carolina 1824. C p.l Bot. cab. 365
hirsutus B. C. not of others				
6190 hirsutus Mx.	hairy			P N. Amer. 1806. C l.p Sw. fl. gar. I
6191 ls'vis M.	smooth			B N. Amer. IBOii C l.p
6192 macropodus Nut.	large-leaved	M or	jn.jl	Y N. Amer. 1822. C p.l
6193 discolor R. S. S.	various-colored	J A or	jl.au	G N. Amer. 1809. C p.l Bot mag. 1273
Cyn&chum discolor B. if.				
6194 obnqus R.Br.	oblique	A or	jl.au	Carolina 1818. D p.l Jac. ic. 2.341
6195 Nuttallirinw Spr.	NuttaU's	J -	Jlau	G Mississippi 1822. D p.l
viridiflors Nut.				
6196 prostratus R. Br.	prostrate	JU LAI or	Jlau	G Mexico 182& C p.l Cav. Ic. 1.7
6197 uniflorus H.SfB.	single-flowered	-\$ • or	W	Mexico 1825. C p.l H.&B.n.S.23B

•774. PERGULA^RIA L.	PEBUULARIA.	(<i>Pergula</i> , trrelliswork; fit for arbours.)	<i>AsclepiUdece</i> .	3.—~6.
6198 odoratissima 8m.	sweetest-scent.	% \ fra	15 jn.jl	G K Indies 1784. C r.m Hot. rep. 185
6199 minor H. K.	amalcr	f	Ira	8 my.au Y.u £ Indies 1790. C r.m Bot mag. 755
6200 sanguinolenta Lindl.	bloody	cu	6 jl.au	G V & Leone 1822. C r.m Bot. mag. 2532

775. MARSDFN/i! R. Br.	MARSDENIA.	(W. Marsden, Esq., auth. of Hist. of Sumatra.)	<i>Asclepitidccc.</i>	2.—8.
6201 erecta R. Br.	upright	fra	jl.au	W Syria 1097. 81 Jac. vin 1.38
6202 suavcolcn R. Br.	sweet-scented	lira	jl.au	W N.S.W. 1816. s1 Bot reg. 489
•776. H O ^ R. Br.	HOYA. (T. Hoy, F. L. S., gardener at Syon House, d. 1821.)			<i>Asdepiphdtt.</i> 5.—7.
6203 carn&sa R. Br.	fiathyleaved	JL'dor	10 jlau	Pk Asia 1802. L r.m Bot mag. 788
%624 lanceolata Hort.	lanceolate	Q or	2 ...	W E. Indies 1815. C r.m
g05 crassiflbia Haw.	thick-leaved	Q or	10 ...	China 1821. C r.m
go6 P6tt8ii Hort	Pott's	ft. CD or	10 ... Y	China 1824. C r.m
trinervis Horri.	three-nerved	ft. CD or	10 ... W	China 1824. C r.m Bot. cab. 936

777. TYLO'PHORA R. Br.	TYLOPHIORA.	(<i>Tylos</i> , a wart, <i>phono</i> , to bear; lvs of corolla.)	<i>Asclrpidddea.</i>	a—5.
6908 grandiflbra R. Br.	great-fibiventid*	J J = 10 jn.jl	... NN.Hidl. 1882. C p.l	
6209 barbita R. Br.	bearded	f J or	100 jn.jl	N. Holl. 1822. C p.l
6210 cxill* Colb.	slender	f Q or	10 jn.jl	Pa.P E. Indies 1823. C p.l Lin. tr. 12.16

•778. CEROPIGTIA L.	CBROPBGIA.	wax, pege, a fountain.)	<i>AsclrpUdraf.</i>	10.—12.
6211 bulbba Rox.	bulmus	2 aps	Ka E. Indies 1821. K p.l	Rox. cor. 1. 7
6212 tuberctma Rox.	tuberous		R.a E. Indies IKJL R p.l	Rox. cor. 1.9
6213 acuminkta Rox.	acuminate		P Command. 1820. R p.l	Rox. cor. 1.8
6214 junccea Rox.	rushy		Y E. Indies 1822. C s.l	Rox. cor. 1.10.
6215 aphylla Haw.	leafless		W E. Indies 1817. C p.l	
6216 dichotoma Haw.	dichotomous		Y E. Indies 1804. C s.l	
f117 africkna Lod.	African		Y E. Indies 1823. C s.l	Bot cab. 906
6218 iinurja Par.	sinuated		Pa.R C G. H. 1818. C p.l	Bur. af. 15
1219 australis Lo. C.	southern		... N. Holl.? 1820. C p.l	
Haw.	Stapelia-form		P C.G.H. 1824. C p.l	

■779. STAPBLTLW L.	STAPELIA.	(Dr. J. B. ii Stapel, Dutch editor of Theophrastus.)	<i>AselerAMea.</i>	36.
wi grandifl,ra Mas.	great-flowered	tt. ID cu 1 s.d.	D.P C. G. H. 1795. C s.l	Mass. 11

•384 quinquerenifrif Haw.	fivo-nrvcd	tt. — leu	1 jn.8	Y. St	C. G. H. 1800.	§ 1.1
6295 Wendlandtina R. & S. Wendkml's tt.	ZD cu	jn.s	Y. s	C. G. H. 1818.	§ 8.1	Wen. c. 2.52
Stapelia rugtoa Wnl.						
6296 marmorkta R. % S.	marbled	tt. ZD cu	1 in.s	Y. St	C. G. H. 1820.	C s.1
6297 refusa Haw.	reticule	tt. HI cu	* jl.au	Y. St	C. G. H. 1820.	C s.1
6296 Woodfordiana Haw.	Wooriford's tt.	ZD cu	-	C. G. H. 1810.	C s.1	
6299 hiobra Haw.	scentless	ZD cu	-	Y. St	C. G. H. 1788.	C 8.1
6300 clypeata Haw.	shielded	tt ^ cu	-	C. G. H. 1812.	C 8.1	
Stapelia clypeata Jac.						

785. OBE'SIA Haw.	OIIESIA.	(Obesus, fat; flowers.)	Asclepihdece. &
6301 geminata Haw.	Iw'injowerett	* my n P.St C. G. H. 1795.	C 8.1 Bot mag. 1326
Stapelia geminata Mas.			
6302 decora Haw.	graceful	tt. — I cu	i my.n Y. St C. G. H. 1795.
6303 serrulata Haw.	serrulate	tt. — cu	jn.au P C. G. H. 1805. F Mass. 26
Stapelia serrulata Jac.			Jac. st. ic.

786. DUVA'LL if Haw.	DUVALLIA.	(A. Duval, a French botanist.)	AsdcpiMca.: 12.
6304 reclinata Haw.	recline	— — jls	C. G. H. 1795. C 8.1 Mass. 28
Stapelia reclinata Mas.			
6305 elegans Haw.	elegant	— —	C. G. H. 1795. C s.1 Bot. mag. 1184
6306 caspitbsa Haw.	tufted	— —	C. G. H. 1790. C s.1 Mats. 29
6307 hirtella Spr.	roughish	— —	C. G. H. 1804. C 8.1 Jac. st. ic.
Stapelia hirtella Jac.			
6308 radiata Haw.	rayed	— —	C. G. H. 1795. C 8.1 Bol. mag. 619
6309 Jacquiniidta R. & S.	Jacquiu's	— —	C. G. H. 1802. C 8.1 Joc.st. ic
radiata Jac.			
6310 replicata Spr.	folded back	— — cu	jls P C. G. H. 1812. C s.1 Jac. ft. ic.
Stapelia replicata Jac.			
6311 tuberculata Haw.	tuberculate	n. — eu	i jls Br C. G. H. 1808. C s.1
6312 Itcig^ta Haw.	smooth	n. — eu	Br C. O. H. — C 8.1
6313 glomerata Haw.	glomerated	tt. — leu'	i — Br C. G. H. 1804. C 8.1
6314 compacta Haw.	compact	tt. — leu	Br C. G. H. 1840. — s.1
6315 mastbdes Spr.	large-teated	tt. — leu	i jl.au Br C. G. H. — s.1

787. CARUNCULA'RIA Haw.	CARUNCULARIA.	(Caruncula, fleshy protuberance; fl.)	AsclepiMecB. 1.
6316 pedunculata Haw.	pedunculate	tt. H] cu	jn.n Br.P C. G. H. P9U. C s.1 Bot mag. 793
Stapelia pedunculata Mas.			

788. PECTIN A'RI A Haw.	PECTLVARIA.	(Pfc/<i>a comb.)	Asdepialear. 1.
6317 articulata Haw.	jointed	tt. ^ cu	jl P C.G.U. 1800. C s.1 Mass. 30
Stapelia articulate Mas.			

789. PIARA'NTHUS R.Br.	PIABANTHUS	(Piar, fatness, anthos, a flower.)	AsctepiMcce. 3.-4.
6318 pGllug R. Br.	dark	tt. ^ cu	* aus.s D.P C. G. H. 1774. C 8.1 Bot. mag. 1648
Stapelia pulla B. M.			
£319 punctatus A 2r.	dotted	tt. 3 cu	JjLn D.P C. G. H. 1795. § 8.1 Mass. 24
6380 parviflbrus Spr.	small-flowered	tt. 3 cu	J au Y.G C. G. H. 1795. § 8.1 Mass. 22.35
Stapelia parviflbra Mas.			

790. HUE'RNM R.Br.	HUBRNIA.	(J. Huernius, an early collector of Cape plants.)	Asclepiadea. 11—12.
^wl reticulata R. Br.	netted	CLZHu	i jl.au P.St C. G. H. 1793. C s.1 Bot. mag. 1662
Stapelia reticulata B. M.			
J>22 campanuKita R. Br.	bcll.shaped	■ !13.■ Jjlo	Y. St C. G. H. 1795. C 8.1 Bot. mag. 1227
5?23 venusta R. Br.	graceful	tt.H]cu i jnjl	Y. St C. G. H. 1795. C 8.1 Mass. 3
≤324 lentiginbsa R. Br.	freckled	a. leu ijjl	Y. St C. G. H. 1795. C 8.1 Bot. mag. 506
f'B5 KUTtkta R. Br.		tt. "Z3 cu \$ au.n	Y. St C. G. H. 1795. C 8.1 Mass. 4
J>22 V. Br.	humble	tt. — Iru \$ au.n	Y. St C. G. H. 1795. C 8.1 Mass. 5
J>22 R. Br.	Xubc-flowered	tt. — leu 2 au.n	Y. St C. G. H. 1805. C 8.1 Bot cab. 225
J>22 R. Br.	bearded	tt. — leu i au.n	w. St C. G. H. 1795. C 8.1 Mass. 7
J>22 * Haw.	curllet	tt.U]cu	* C. G. H. 1795. C 8.1 Jac. st. ic.
J>22 clavSgora //aw.	club.bearing	tt. Z3cu *Jl.n"	Y. St C. G. H. 1795. C 8.1 Jac. st. ic.
J>22 ocellata Jac.	occilated	tt-ZDcu J au.n	Y. St C. G. H. — C 8.1 Jac. st. ic.
Stapelia ocellata Jac.			

*J91. BRACHYSTE'LMA if. Br.	BRACUVSTELMA.	{Brachys, short, stelma, a crown.)	AtclejAddct. 1.
6332 tuberbum 7f.Hr.	tuberous	X ID cu	1* jn.jl P C. G. H. 1821. C s.1 Botreg. 722

*792. CARALLITMA R. Br.	CARALLUMA.	(IU Indian name.)	AsclcpUdece. 2.
6333 adscendens R. Br.	ascending	tt.Z3cu	2 jl P E. Indies 1804. C s.1 Rox. cor. 1.30
Staplia adscendens Rox.			
6304 umbellata Haw.	umbellel	tt-Z3cu. Pk E. Indies 1804. C s.1

fts^KW nere^n» L.	SWE/R T/i < L.	FELWORT.	/E. Swert, a celebrated cultivator in Holland.)
KW nere^n» L.	perennial	or 1	Alp. Eur. 1596. D p.l Mil. ic. 139
JIJfnrncticulitaGm.	small-horned	A pr	Switzerl. 1817. D p.l
JIJfnrncticulitaGm.	Michaux's	24 jn.jl	Ypp. Switzerl. 1817. D p.l
JIJfnrncticulitaGm.	corniculata Ph.	Y pp	Bot rep. 117

GENTIAN.	(Virtues first experienced by Gentivs, king of Illyria.)	Geutntnea M.—W.
yellow	5t A or 4 jn.jl	Y Alp. Eur. 1596. D p.l Mil. ic. 139
hybrid	A pr	Switzerl. 1817. D p.l
purple	3 jn.jl	D p.l
white-flowered	B	D p.l
Pannonian	W	D p.l
two-lobcd	W	Jac. au. 2 136
Burser's	A or 2 jl au	Jac. au. 5.28
campanulate	Y	Dec. ic. 15
seven-cleft	L.B	HJO. D p.l
Spotted	Levant	Trat. au. 3.10
Swallow-wort-lk?	B	Pal. ros. 2.96
long-leaved	AustRIA	Jac. au. 29
crossed	D.B	Bot. mag. 1229
ascending	Siberia	Bot. mag. 1410
decumbent	Jnl	Bot. mag. 1078
frigid	B	Pal. ros. 2.96
cold	Siberia	Bot. mag. 705
Jrigida //»: «gida Pall.	W	Bot mag. 723
	W	Trat. au. 7.1
	W	Pal. ros. 2.95

6353 gelida Bieb.	ice-cold	P.Y	Siberia	1807.	D	p.l
6354 Pneumonanthe L. 2 guttata	Wind-flower spotted	B	England	moi.li.	D	p.l
6355 Pae&do-pnmonan̄. the R. & S.	C'B ibtard	B.	.	.	D	p.l
6356 linearis Fro.	I. wind-flower	I	N. Amer.	1800.	D	p.l
6357 triflora Pall.	Mnesr-leaved	J	Carolina	1816.	D	p.l
6358 Saponaria l.	three-flowered	J	Siberia	1807.	D	p.l
6359 Cateab'i Walt.	Soapwort-/i/d	J	N. Amer.	1776.	D	p.l
Ssiō intermliui Jt. M.	Catesby's	J	N. Amer.	1803.	D	p.l
IVkil ochrolucu Frit.	intermediate	J	N. Amer.	1820.	I)	p.l
6362 incarnata S. M.	five-flowered	J	N. Amer.	180J.	D	p.l
6363 quinqueflora Lam.	golden^	Pa.B	N. Amer.	1824.	S	p.l
afarnelloides A/r.			Norway	182a	D	p.l
6361 aorca L.	umbellate	fmy.Jn	Caucasus	1823.	D	p.l
qumqefbia Fl. dan.	steinless^	i mr.my	Wales	walls.	D	p.l
6365 umbellata Bieb.	narrow-leaved	i mr.my	Alp. Eur.	1819.	D	p.l
6366 acatilis /.	short-leaved	i ap.my	Switzcr.	1819.	I)	Bar. ic. 103. 2
2 anii(Uit?)lia	alpine	i .my.jl	Alp. Eur.	1817.	D	p.l
6M7 brachyphilla Fit.	Altaian	i ap.my	Siberia	1824.	I)	Pal. ros. 97. 1
63K sibylma Fit.	Pyrencean	i ap.my	Pyrenees	1825.	D	p.l
6359 alliaca Pall.	dwarf	i iip.my	Switzcr.	1817.	I)	W. & K. 207
6370 pyrenitca L.	spring	i jn.jl	England	moun.	D	p.l
6371 jM.mila L.	summer	i jl	Austria	1818.	I)	Hacc. 4.2.3
6372 verna L.	Bavarian	i jl.au	Germany	1775.	D	p.l
6373 awtva R. AS.	imbricate	i ap.jn	Switzerl.	1819.	D	p.l
6374 bavdrica, L.	bladdery	P S	Altai	1824.	I)	Bar. ic. 101
6375 imbricata Fro.	snowy	i au	Europe	1822.	I)	
6376 angulbsa Bieb.	humble	ap	Scotland	scalp.	D	81
6377 utnculosa Z.	studded*	...	Caucasus	1824.	D	p.l
6378 nrl.vls L.	German	i au	Germany	1818.	S	p.l
6379 humiliis Scw.	Amarella	i au	Britain	ch.pa.	S co	Fl. mon. 278
6380 clavata A A/.	field	i au	Britain	gra.pa	S co	Eng. bot. 236
6381 germijinica W.	obtuse-leaved	i jl.au	Switzerl.	1826.	S	p.l
6382 amarietra II.	meadow	i jl.au	Siberia	1817.	S	p.l
6384 obtufl folia IV.	Caucasian	i jl	Caucasus	1804.	S	p.l
6385 pratinus Fro.	narrow-leaved	i jl.au	N. Amer.	1812.	D	p.l
6386 caucWa B. M.	icy	i jl.au	Alp. Eur.	1819.	S	p.l
6387 nnKjstifolia Mx.	Carinthian	i jlau	Switzerl.	1817.	S	p.l
6388 gl.ctiaca MJ.	ciliated	f au.a	Germany	1739.	D	p.l
6389 canthiaca Fro.	fringed	tau.s	Caucasus	1818.	D	p.l
6390 cihata Z.	bearded	i au.a	Siberia	1764.	D	p.l
6391 cihata W.	long-haired	J jn.jl	N. Amer.	1804.	S	p.l
6392 barbata Fro.		L.B	Bot mag. £031			
ciliata 2 O/.						
6393 crinuq Fro.						
fimbriata A Arp.						
795. HYDRCOLEA Z.	HYDROLEA. (<i>Hydor</i> , water, <i>elaia</i> , oil; situation, and oily.)		<i>Convotvulaceat.</i> 2. - 6.			
6394 npinbgn L.	thorny	£\ I or	1 jn.jl	P.n	S. Amer.	1791.
6395 quadrivalvis	four-valved	£ IAJ or	1 jLau	B	Carolina	1824.
caroliniāna Mx.					S bog	
' 796. EVCLVULUS L.	EVLVULUB. (<i>Evolo</i> , to roll out; not twining, opp. <i>Convolv.</i>)		<i>Convovul&ce*</i> . 12.-22.			
f13% nummpularia L.	Money-wort					
tlk#7 Rangiticus L.	Ganges	* O pr	1 jl.au	B	Jamaica	1816.
am flsinoids R. Br.	Chickweed-like*	O pr	i jn.au	B	E-Indies	1820.
6399 hiniut Lain.	hairy	Jc III	jn.au	B	E. India	1817.
6400 villbsus R. # P.	villous	i. TAlpr	J jl.au	B	Trinidad	1818.
6401 sericeus Sw.	silky	i. TAlpr	J jl.au	W	S. Amer.	1810.
6402 tncanus Pers.	hoary	i jn.au	B	W. Indies	1816.	
serices R. & P.	Flax-leaved	Jt O pr	2 au.s	B	S. Amer.	1810.
6403 /inifolius /.	Nuttallia	Jt O pr	Jn.au	B	N. Amer.	1824.
6404 Nuttalliji/tf R. & S.	emarginate	Jopr	1 s	B	E. Indies	1816.
6405 latifolius Ker	broad-leaved	Jopr	« jn.jl	W	Brazil	1819.
FALKIA.	(J. P. Folk, a Swedish botanist, died 1774)		<i>ConvovulacecB.</i> 1.			
6407 repens L.	creeping	JW IAJ or	i my.au	Pk	C. G. H.	1774.
798. CU'SCUTA L.	DODDER.		<i>Convoloulucetv.</i> 7. - 23.			
6408 europea B'a L.	Euro/jean	A cu		Britain	hea.	D par Eng. bot. 378
6409 Apifnymum Mur.	True Dodder	A cu		Britain	hea.	D par Eng. bot. 55
6410 chinensis Lam.	Clunrae	A cu		China	1803.	D par
6411 australis R. Br.	southern	A cu		N. Holl.	1818.	S par
6412 chilensis B. M.	Chilian	A cu	year	Chile	1821.	D par Bot. reg. 603
6413 verrucosa Swt.	warted	A cu	apo.o	Nepal	1821.	D par Sw. fl. gar. ff.
6414 Horterk Swt.	Hooker's	A cu	s.o	E. Indies	1821.	D par H. Hook. ex. ff. 150
reflxia var. verrucosa Hook.	American	£, E3 cu ... jlo	W. Y. S. Amer.	1816.	D par SI. jam.	1.128.4
6415 americana L.						
799. DICHO'NDRA Font.	DICUONDRA. (D/«, twice, <i>chondros</i> , grain ; form of caps.)		<i>Convolvulcea.</i> 5. - 7.			
6416 r&cius Font.	creeping	% IAJP1	1J jnau	W	N. S. W.	1803.
6417 sericea Sum.	silky	i wE2cu	Ijjnau	W	Jamaica	1793.
6418 carol inensis Mi.	Carolina	Irr-AJcu	& jn.au	W	Carolina	1810.
repens fi Poir.					D	
6419 argentea IV.	silvery	JWIAJcu	§ jn.au	W	W. Indies	1800.
6420 rotundifolia Lk.	round-leaved	J^iAJcu	jn.au	W		1819.
., 800. CRESS'A L.	CREBSA.	(Cressa, a native of Crete, Lat.: abounds there.)	<i>Convolvulcea.</i> 1. - 3.			
6421 microtek A Sun.	CrCun	Jt O cu i	jl.au	P	Levant	1822.
6422 debilla Stvx.					S s.p	Lam. fil. 183
6423 maypurena G. Don	weak	O cu	jn.jl	W	Jamaica	181d
AncistrocerebrjBB	Maypurena	O cu	ijn.au	W	Trinidad	1817.
Kth.					S co	H. AB.n.8.1

802. <i>VELVZIA</i> W. 6484 <i>rigida</i> W.	VBLBZIA. rigid	<i>{C. Vein,</i> physician and botanist at Madrid, j O cu * jl W.P Spain 1633. S co	<i>Caryopk Alea.</i> 1. <i>Banic.</i> 1018
801. <i>LINC&NIA</i> L. 6425 <i>alojecuroldea</i> L.	LINCO.VIA. Fox-taiMikk	(Uncertain; probably a man's name) • i Jor 2 myjn. WW (C.G.H. 1616. C pi Swz.b.m. 4	<i>Brunfoeea.</i> 3. - 4.
6426 <i>flymifblia Sun.</i> <i>Diosma defasta Thun.</i> 6427 <i>cupidkta Sun.</i> <i>Diosma cuspidata Thun.</i>	Thyme-leaved cuspitate	M Jor 2 myjiiWW CCGGHH.1825. C pi Swz. b. m. 85. 4 ft _Jor 2 myjn W C. G. H. 1825. C pi SwYb.mi. 5.884. £7.1	
804. <i>BUMA'L Dil Thun.</i> 6428 <i>trifolia Thun.</i>	BUMALDA. three-leaved Staphylta Bumalda Dec.	<i>(J. A. de Bumalda, a botanist of Bologna.)</i> ■ LJCU 2 jn.s ... Japan 1812 S co	<i>Bruntoce*.</i> 1.
•905. <i>HEITCHEIM</i> L. 6429 <i>americana</i> L.	HEITCHEBA. American	<i>(J. H. Heucher, prof. med. at Wittcmberg.)</i> <i>Saxifrage*</i> . 7. - 8.	
G430 <i>hispida Ph.</i> 6431 <i>glabra Pail.</i> 6432 <i>pubescens Ph.</i> 6433 <i>villcea Mx.</i> 6434 <i>caule&cens Ph.</i> 6435 <i>macrophylla La. C</i>	hispid smooth pubescent villous caulescent long-leaved	• or 1 myjl P N. Amer. 1616. D s.l Pluk.al.58.3 • or 3 my P N. Amer. 1826. D p • or 1 myjn Pk N. Amer. 1824. I) co • or 1 myjl Pk.v N. Amer. 1812. D Lp • or 4 myjl W N. Amer. 1812. I) l.p • or 1 myjl W N. Amer. 1812. D l.p • or 1 myjn Pk N. Amer. 1810. D co	
806. <i>CUSSO*N/i*</i> L. 6435 <i>thy mi flora L.</i> 6437 <i>tripterU Coll.</i> 6438 <i>spicata L.</i>	CIJSSONIA. (<i>A. Custon</i> , a celebrated French liotanist.)	<i>Aralidcet.</i> a thyrs-flowered * i Jor 6 ... G C. G. H. 1795. C l.p Th. up.3.12 three-winged ft T Jor 4 ... G C. G. H. 1816. C l.p Col. h. rip. 26 spike-flowcrcd ft LJ or 6 ... G C. G. H. 1789. C s.l Th.up.3. 13	
807. <i>ANA'BASIS</i> JL 6439 <i>famarisciflra L.</i> 6440 <i>aphf Ha /.</i> <i>Salsola articulata</i> Font. 6441 <i>flrida Bieb.</i> 6442 <i>oppositiflora Schr.</i>	ANABASIS. Tamarisk-lvd leafless flowery opposite-flwd	<i>(Anabahlo, to ascend; climbs trees.)</i> <i>Chenopbdeee.</i> 4. — 9. a. i cu Jn.jl G Spain 1753. C l.p Cav. ic.3.2R3 a. + cu Jn.jl G Asia Min. 1817. C l.p Cav.ic.3.284	
808. <i>SALSO'LAZ.</i> 6443 <i>KM L.</i> 6444 <i>TYagus L.</i> 6445 <i>fomarisciflra Pall.</i> 6446 <i>crissa Bieb.</i> 6447 <i>rofacea L.</i> 6448 <i>oppositiflora Desf.</i> 6449 <i>brachiatra Pall.</i> 6450 <i>glatica Bieb.</i> spicata Pall. 6451 <i>vermiculata L.</i> 6402 <i>rigida Pall.</i> 6453 <i>microphylla Cav.</i> 645* <i>verrucosa Bieb.</i> 6455 <i>folidsa &/<.</i> <i>Anabasis foliosa L.</i> 6456 <i>laniflra Z.</i> 6457 <i>sattva L.</i> 6458 <i>Sorb. L.</i>	SALTWOBT. Kali Tragus Tamarix-like fleshy rosy opposite-leaved tt. branched glaucous tt.	<i>(Salsus, salt; saline properties.)</i> <i>Chenopbdite.</i> 16.—51. O ec 1 jl.au F Britain sea. sh. S s.l Eng. bot 634 O cu 2 jl.au W S. Europe 1817. S s.l Pal. ill. 2.29 O cu 2 jl.au Y Tauria 1820. S a.l Pal.il2.25 cu 1 jl.au W.v Caucasus 1818. S p.l Bux.c.1.14.2 8 cu 1 jl.au Pk Asia 1759. S il Schk. han. 1.57 Opposite-leaved tt. O cu 1 jl.au -- Sicily 182a C 81 Cav. ic 245 branched O cu 1 jl.au Br Tauria 1818. S <p Pal. III. 2.22 glaucous tt. -J cu 4 jl.au Y.o Caucasus 1821. C s.p Pal. ill. 2.19	
— 809. <i>KO^CH/i*</i> Schr. 6451 <i>Ayssopiflra JlwB</i> Jlw <i>trigyna £*.</i> Jlw <i>prostrata Schr.</i> 6462 <i>arenaria Both</i> Chcnnpodium arenarium Kit.	KocniA. Hyosop-leaved trigynous trailing sand	<i>(A. Koch, a German botanist)</i> <i>Chenopbdea.</i> 9. — 12. O cu Ifjn.jn au G Siberia 1801. S co Pal. it. 14;l. PL O cu 3 jl.au G Spain 1804. S s.l Cav. ic. 3.289 Jlw 2 Jn.jn au G S. Europe 1780. C s.l Jac.au. 3.294 O cu 1 myjn W.o Hungary 1822. S s.l W. & K. 1.78	
5463 <i>edoTdes %Ar.</i> H464 <i>muricata fcAr.</i> B erjphora Schr. ><66 <i>coparia & Ar.</i> G467 <i>dasy&ntha ScAr.</i>	Stonecrop-like muncatc wool-bearipg broom Cypress thick-flowered	O cu fi Jn G Crimea 1821. ■ s.l Pall. ill. 36 O cu 1 jl.au G Egypt J77& ■ si Al. tau. a 4.2. O cu f Jn.jl G Spain 1823. ■ si Sr.hal.3 O cu 3 jn.s G Greece 16K. ■ co Sc. hal. 1.1. O cu 1 jl.au G Caucasus 182J. ■ sp Pal ill. 1.10	
810. <i>CHENOPD' DIUM</i> L. 6478 <i>Bfmu Henuku*</i> L. 6480 <i>Arbicium L.</i> g**" " " L. X <i>Jiobrum</i> .. **72 chrysomelanospeY- mum Bulb. J73 cnthraginivK Zue. J74 puicturrum Sco. Jga Kuuu>nmc Jac. r<>7 Qumba W. <H8 rhoibifrlum Mhl. 6J79 serotinum L.	GOOSEFOOT. Good Heqry city. Atriplex-like red Cyclow and "J Jblakw seeded J farthagena small-dotted Guinea wall green Quinoa red Quinoa rhomb-leaved late fig-leaved white rounilinh-leaved green entire-leaved thick-leaved Opulus-leaved	<i>{Chen,</i> a goose, pout, a foot; leaves) <i>Chenopbdeete.</i> 34. — 72. Good Heqry Aw 1 myau G Britain rub. 1) co Eng.bot. 1033 city. O w 1 au G Britain dungh. S co Eng. bot. 717 Atriplex-like O un 3 au.s S China 17H0. S co Jac. vin.3.80 red O w 2 auH R Britain dungh. S co Eng. bot. 1721 Cyclow and "J O un 1 au.8 G Galicia 1820. S co Jblakw seeded J O un 1 au.8 G Galicia 1820. S co farthagena O un 1 jl.au G Carthag. 1817. S co small-dotted O un 1 jl.au G Europe 1826. S co Scon.d. 1.11 Guinea O un 2 au.s G Guinea 1790. S co Jac. ic.i.345 wall O w 14 au.a G Britain rub. S co Eng. bot 1732 green Quinoa O cul 3 jl G Peru 1822. S co red Quinoa O cul 3 jl G Peru 1822. S co Feu. ob. 10 rhomb-leaved O un 1 jl.s G N Amer. 1807. S co late O un 2 jl.s G Spain 1821. S co fig-leaved 8 w \$ au.s G Britain dungh. S co Eng. bot 1724 white O w iffjl.S G Britain rub. S co Eng. bot. 1723 rounilinh-leaved O w 2 jl.s G Britain rub. S co green O w 2 jl.s G Britain rub. S co entire-leaved O w 2 jl.s G Britain rub. S co thick-leaved O w 2 jl.s G Britain rub. S co Opulus-leaved O un S jl.au G Europe 1810. S co Bast.j. 1814,3	
(*81 <i>Album L.</i> 2 subrotundum 3 vlide 4 integriflum 5 crasiflum ^WS opuliflum Schr. J70 <i>litoides Lej.</i> Schradera/inum R. ft S. Schrader's fa lidum Schr. not of others J71 <i>urn Poir.</i>	pctioled Blitum-like Schraderainum R. ft S. Schrader's fa lidum Schr. not of others cuUeaved	O un 1 jl.au G S. Amer. 1824. S co O un 1 jl.au G France 182a S co O un 3 jl.au G Europe 1817. S co O >n 1} jl.au J Europe 1821. S co	H 2

C487 humifusum Zuc.	ground-spread	O un I jl.au	G Europe 1810.	S co
6488 giganteum D. Don	gigantic	O un 6 au.s	G Nepal 1818.	S oo
6489 hibriderum L.	hybrid	S W II au.s	G Britain rub. S	Eng. bot. 1919
6490 Bitrya L.	Botrys	O un 1 jn.s	G S. Europe 1548.	S co Fl. gr. 243
6491 Aotryoides Sm.	BotrysJJte	O w 1 jn.au	II Britaiu sea. sh. S	Eng. bot. 2247
6493 fœtidum Lam.	fetid	O un 4 jn.au	G Europe 1800.	S co Fl. dan. 1152
6493 multifidum L.	many-deft	■ ■ 2 jn.ø	G Buenos A. 1732.	D Di. el. 60.77
6494 ambrosioides L.	Ambrosia-like	fir if jt.0	G Mexico 1640.	S eo M.h.5.35.8
6495 sutructosum tV.	suffruticose	«L fun 2 jl.o	G Mexico 1819.	S co
6496 anthelminticum L.	anthelmintic	«.i fun 3 jl.ju	G America 1732.	C co Di. eL fiC. 76
6497 graveolens W.	heavy-umelling	O un 4 jl.au	G Mexico 1823.	S co
6498 gluticum JL.	glaucous	O w ljl.ju	G England rub. S	Eng. bot 1454
6499 marginatum Spr.	margined	O un 2 jl.au	G Europe 1819.	S co
6500 iutulum 7to/A	spreading	O un 1 jl.au	G E. Indies 1817.	S co
6501 crassifolium Zteff.	thick-leaved	O un 2 jl	G Europe 1802.	S co
6502 6lidiun Sm.	rank-bmulling	O w 1 jl.au	G Britain rub. S	Eng. bot 1034
6513 polyspermum L.	many-seeded	O w 1 jl.au	G Britain rub. S	Eng. bot 1480
6504 ualifilum Jar.	caudate	O un 1 jl.au	G Guinea 1806.	S co Jac. ic. 2.344
(J505 lateralq // K.	lateral	O un 2 jl.au	G Europe 1781.	S co
6506 lancolatum 3/A.	spoor-leaved	O un 1 au.s	G Pennsylv. 1809.	S co
(5507 ariatum L.	awned	O un 1 jn.s	G Virginia 1771.	S co Gm. si. 3.15.1
(VK8 scipium Mayer	hedge	O un 2 jn.jl	G Moravia 1823.	S co
6509 acuminatum W.	acute-leaved	O un 1 jl.au	G Siberia 1810.	S co W.n.ber.2.5.2.
6510 auctiflum E. B.	sea.	O w 5 au	G Britain unc. gr. S	Eng. bot. 1481
6511 maritimum L.	small-leaved	O un 2 jl.au	G Britain sal. m. S	Eng. 1st. 633
6512 parviflum It. \$ S.	11. & S. Biebrsteiniiflum	O un 3 jl.au	G Caucasus 1825.	S co PaL ill. 3.44
6513 Biebrsteiniiflum 11.	garden	It un 3 jl.au	G Caucasus 1824.	S co
6514 hortense R. \$ S.	Salsola divulgens Forr.		G S. Europe ... S	co
6515 hirshtum Bleb.	hairy	O un 3 jl.au	G Euro] 1791.	S co Fl. dan. 187
6516 fruticbum Schr.	shrubby	tt. ec 2 au.s	G England sea sh. C	co Eng. bot. 685
Salsola fruticbsa E. B.				
6517 altissimum Bich.	tallest	O un 6 jlau	G Italy 1775.	S co Sc. hal. 1.3
6518 salsum/t*.4r&	Saltwort	O un 1 au.s	G Astracan 1782.	S co Jac. vin. 383
Salsula salsa L.				
6519 setgerum Dec.	bristle-bearing	O un 2 jnj.l	G S. Europe 1822.	S co
6520 radiatum Schr.	rayed	O un 1 jl.au	G N. Amer. 1820.	S co An. mu. 2.34
6521 VuUasianum R.&S.	Pallas's	O un 2 jl.au	G Persia 1826.	S co Bux.c. 1.31.1
6522 spic/tum/t.#&	spiked	O un 1 jl.au	G Spain 1821.	S co Cav. ic. 3.290
Salsola sifla Cav.				

811. BETA L.

6523 vulgiris L.	BEET.	(BetL red. in Celtic.)	Chenopbdite. 7.
1 viridis	common	cu 4 au	G S. Europe 1548.
2 rufra	green	dt 4 au	S. Europe ... S r.m Pl. ic. 169
31utea	red-rooted	clt 4 au	G S. Europe ... S r.m
6524 macrorhiza Stev.	long-rooted	clt 4 au	G S. Europe ... S r.m Kern. ab. 235
6525 pâtula H. K.	spreading	clt 6 au	G Caucasus 1820.
6526 chela L.	Sicilian	un 1 au	G Madeira 1778.
6527 crfspa Trat.	curled	cul 6 au	G Portugal 1570.
6528 trfgyna Kit.	three-styled	or 3 jl.au	G S. Europe 1800.
6529 maritima L.	sea	i cul 1 au	G Hungary 1/9G. S r.m Trat.-av."ll7

812. BO^SEif L.

6530 Yervambra L.	GOLDEN ROD.	IE. G. Bosc, a botanist of taipsic.)	Chenopbdete. 1.—2.
	Yervamora	i ll_jor 8 ... Ku Canaries 1728.	C p.l Walt. h. 24.10
813. HERNIA*RIA L.	RUPTURE-wottT.	(Hernia, rupture; supposed med. qual.)	Paronychiar. 8.—12.
6531 glabra L.	smooth	* A cu £jl	England sa.gr. S co Eng. hot KM
6532 hirsuta L.	hairy	Jc A cu \ jl.au	England sa.gr. S co Eng. bot. 1379
6533 fruticosa L.	shrubby	tt. cu i my.au	Spain 1814. C p.l Lob.ic.85
6534 polygonoides Cav.	Knotgrass-like	tt. (cu i my.au	G S. Europe 1702. C l.p Cav. ic. 2. 131
6535 incana Bieb.	hoary	* cu * jn.au	G S. Europe 1822. S l.p Pluk. al. 5.J..!
hirsuta Bieb.			
6536 alpina Vil	alpine	■ A cu * my.au	G S. Europe 1822. S lp
6537 annua Lag.	annual	Jk O cu ijlau	Spain 1824. S co
6538 cinerea Dec.	cinerous	Jk O cu i jnjl	G Montpel. 1823. S co

•814. U'MUSL.

6539 carapestris L.	English field	tm 80 ap.my Br	Britain hed. L co Eng. bot. 1886
6540 subcrbsa Afoen.	cork-barked	tm 40 ap.my Br	Britain hed. L co Eng. bot. 2161
6541 fruticosa tV.	shrubby	m or 8 ap.my Br	Europe ... G co
6542 glabra E. R.	smooth	tm 60 ap.my Br	Britain hed. L co Eng. bot. 22 l\$
6543 major & A	greater	tm 40 ap.my Br	... G co Eng. bot. 2542
6544 montina E. B.	mountain	tm 40 ap.my Br	Britain hed. S co Eng. bot. 1887
6545 effusa W.	spreading.^wr/ \$	tm 40 ap.my Br	Britain ... L co Schk. han. 57
- cihata UArA			
6546 americanæ Ph.	white American'	tm 40 ap.my Br	N. Amer. 1752. G co
6547 alâta Mx.	winged	tm 30 ap.my Br	N. Amer. 1820. G co Mic. ar. 3.5
6548 Alba Xi.	white Hungar.	tin 30 ap.iny Br	Hungary 1824. G co
6549 hñmllis Amm.	low	or 6 ap.iny Br	Siberia ... G co
6550 crispa ff.	curled	or 20 apmy Br	N. Amer. ... G co
6551 fulva Ph.	deep-yellow	tm 60 ap.my Br	N. Amer. ... G co Mic. ar. 3. 6
pñndula W.			
6552 scabra Mil.	scabrous	£ tm 40 ap.my Br	N. Amer. ... L co
6553 piimila W.	dwarf	-a or 2 ap.my Br	Siberia 1771. L p.l Pal. ros. 1.43
6554 parvithilia Jac.	small-leaved	A or 12 my lir.	1822. G co Jac. sc. 262
6555 microphylla Pers.	small-leaved	■ or 10 ap.my Br	Siberia ■ L coPal.ros.l.48.A.B.C
6556 chinensis P*.	China	» ior 3.	China ■ ■ L
6557 intcgncfia Rox.	entire-leaved	f or 40 ... Br	E. Indies 1822. G s.f Rox. cor. 1.78

815. CHAILLE'r'il Dec. CHAILLETIA. (If. Chaillet, a Swiss botanist.) Chaillete*. 1.—6.

6558 Toxicaria G. Don	Rat poison	* L J P 5 my.jl W	S.Leone 1824. C p.l
816. PHY'LLIS L.	BASTARD HARE'S EAR. (Phyllon, a leaf: its chief beauty.) Rubihcea. 1. ^		
■S9N■ ■ L.	Noble	m cu 3 jh.jl G	Canaries 1699. C r.m DL el. 299.386

- | | | | | |
|------------------------------|-----------------|--|----------------------------|------------------------------|
| 817. CORIA'NDRUM | /., CORIANDER. | (Koris,* bug; smell of leaves.) | <i>Umbellifer*</i> . | I—J- |
| 6560 sativum L. | cultivated | O clt 2 jn | W England | fields. S co Eng. bot 67 |
| 818. BIFORIS Spr. | BIFORIS. | (Bis. double, <i>forts</i> , a flap; fruit.) | <i>UmbeUiferte</i> . | S. |
| 6561 testicuhitum Bieb. | twin-fruited | O Tun 2 jn .j1 W | S. Europe | 1640. S co Pluk-al. 169.2 |
| Cbnandrum testiculMum L. | | O un ljjn.jl W | Caucasus | 1817. S co |
| 6562 radians Bieb. | radiant | | | |
| Conundrum testiculitum Bieb. | | | | |
| 819. SCA'NDIX L. | SCANDIX. | (Greek name for an eatable plant) | | |
| 6563 Pcten L. | Veaus's Comb | O ■ 1 jn.jl W | Britain | co. fl. Eng. bot. 1317 |
| 6564 austrMis L. | southern | O un 1 my.jn W | S. Europe | 1713. co Col.ec. 1.90 |
| 6565 pinnatifida Ven. | wing-cleft | O un limyjn W | Persia | 1805. co Ven.eels 14 |
| 6566 falcatu Lon. | falcate-seeded | O un 1 my.jn W | Crete | 1817. co Hof.um.2.2 |
| W^lia radians Hofm. | | | | |
| 6567 grandiflora L. | large-flowered | O un If my.jn W | Iberia | 1818. S co Hof. um.2.3 |
| W^lii grandiflura Hofm. | | | | |
| 6508 ibérica Bieb. | Iberian | O un 1 jnjl W | Iberia | 1823. S co Hof. urn. 2.4 |
| Wylia ibérica Hofm. | | | | |
| *820. jfINTHRI'SCUS Pera. | Rouon CHERVIL. | (By Pliny to a plant like Scandix.) | | <i>UmbclUfera</i> . & — 9. |
| 6569 vulgXris Per*. | common | O w 1 J my.jn W | Britain | he. ba. S co Eng. bot. 813 |
| Scandix ifinfrhiscus L. | | | | |
| 6570 tuberculata it. &j S. | tuberculate | O un Ift my.jn W | Syria | 1810. S co |
| 6571 taurica Fis. | Taurian | O un 1 my.jn W | Tauria | 1820. S co |
| 6572 hispida Dec. | hispid | O un 1 my.jn W | France | 1819. S co |
| 6573 nodtta Pen. | knotted | O un 1 my.jn W | Sicily | 1656. S co Jac.vin.a25 |
| 6574 trichosperma R8/S. | hairy-seeded | O un li ... W | Hungary | 1818. S co |
| 6575 nemorosa Spr. | grove | A un 2,1 my.jn W | Russia | 1810. D co Gm. si. 1.49 |
| 6576/umarijoldes Spr. | Fumaria-like | A un 1 my.jn W | Hungary | 1810. D co W.&K.3.224 |
| *821. CHJEROPHYLLUM L. | CHERVIL. | (Chain, to rejoice, <i>phyllon</i> , leaf; <i>mfl.</i>) | <i>UmUBifetw.</i> 8*—9B. | |
| % 6577 sylvestre L. | wood | A w & my.jn W | Britain | hed. D co Eng. bot. 752 |
| 6578 nltidum Vahl | shining | A un 3 my.jn W | Europe | 1820. D co |
| 6579 hfbridum Ten. | hybrid | A un 1 my.jn W | Italy | 1822. D co |
| (5580 divaricatum Poir. | divaricate | A un 1 my.jn W | Europe | 1810. I) co |
| 6381 monogonum Kit. | one-angled | A un 1 my.jn W | Hungary | 1810. D co |
| angulatum Kit. | | | | |
| ^8582 satvum Pers. | cultivated | O cul li my.jn W | England | he.ba. D co Eng. bot. 120 |
| Scandix Cerefulium L. | | | | |
| 6583 nroctumbens Lam. | procumbent | Jfc O uu 1 jnjl W | Virginia | 1699. D co M. h. 9.11. ult. |
| 6584 bulbosum L. | bulbosous | & A un li jnjl W | Europe | 172i. D co Jac.au. 1.63 |
| Afrrhis bulbusa Spr. | | | | |
| 6585 temuJumL. | giddy | Q w 3 jl.au W | Britain | hed. D co Eng. bot. 1521 |
| 6586 hirsutum L. | haxtyUcaved | A un 24 jnjl Pk | Switzerl. | 1759. U co Jac. au. 2.148 |
| 6587 Cicutana ViL | Cicutn-leaved | A un li my.jn HW | S. France | 1810. U co |
| 6588 roseum Bieb. | rosy | A un 1 my.jn R | Caucasus | 1817. I) co Hof. urn. 1.1.33 |
| 6589 hüm lie Bieb. | humble | A un 1 my.jn W | Iberia | 1&3. D co |
| 6590 coloratum L. | colored | A un 1} jl.au Y | Illyria | 1806. D co |
| 6591 aureum L. | golden | A un 2 jl.au Pk | Scotland b. of ti. | D co |
| 6592 maculatum Horn, | spotted | A un 3 my.jn W | S. Europe | 1810. D co |
| 6593 torquatum Dec. | torquate | A un 3 my.jn W | Piedmont | 1819. I) co |
| '6594 Bieberstein M K. & S. | Ilioersterain's | A un 2 my.jn W | Caucasus | 1825. D co |
| 6595 tenuifolium Poir. | fine-leaved | A or 1 my.jn W | S. Europe | 1818. D co |
| 6596 capense Thun. | Cape | A un 1 my.jn W | C. G. H. | 1810. S co |
| 6597 aromiticum L. | aromatic | A or 3 jn.jn W | Germany | 1726. D co Jac.au. 2.150 |
| 6598 angelicafolium Bieb. | Angelica-lvd | A un 3 my.jn W | Caucasus | 1820. D co |
| 6599 canadensePrrr. | Canadian | A or 1} jl.au W | N. Amer. | 1699. D co M. h. 9.11 |
| ^ Staon canadense L. | | | | |
| ^6600 Claytoni Pers. | Clayton's | A or 2 jl.au W | N. Amer. | 1806. D co Spr. un. 3.6 |
| Jl/yrhisi Clayton&ni Mx. | | | | |
| _ 822. JERY'NGIUM L. | ERYNGO. | (Eryggano, to belch: expels wind) | <i>UmbeUifcra:</i> 34.—58. | |
| 6601 aquaticum L. | aquatic | | N. Amer. | 1699. D bog Botreg.372 |
| 6602 longifolium Cav. | long-leaved | | Mexico | 1820. D s.l. Cav. ic. 6.555 |
| 6603 bromelto/diM7W Lar. | Pine-apple-lvd | | NewSpain | 1824. D □ Lar. cr. 28 |
| 6604 ebracteatum Lam. | bractless | | B. Ayres | 1817. D □ Lar. cr. 32 |
| 6605 pl3num L. | RaUeaved | | Europe | 1596. I) s.l. Jac. au. 4.391 |
| 6606 aspenfolium Lar. | rough-leaved | | | 18J0. I) 81 Lar. er. 11 |
| W&D dichotomuin Desf. | dichotomous | | S. Europe | 180. I) 81 Desf. at 1.55 |
| & K^Kntoum Bieb. | giant | | Caucasus | 18J0. I) SI |
| U^o kracile Lar. | slender | | N. Spain | 1824. I) 61 Lar. er. 24 |
| ^M^A uspidatum L. | tntid | | Spain | 1699. D 81 Lar. er. 9 |
| ^C^A meum Pers. | blue-floicred | | Caspian | 1816. I) SI M.h.7.37.13 |
| ^C^A tum L. | sea-holly | | Britain | seash. D s.l. Eng. bot.7IH |
| ^C^A tum Lam. | alpine | | Switzerl. | 1597. D s.l. Bot. mag. 9*22 |
| ^C^A tum | stiff | | France | 1816. D 81 Vil. del. 17 |
| firi-i" --- Porm | small | | Spain | 1640. D 81 Lar. er. hi |
| ^C^A cornicuntum Lam. | Gahum-like | | Portugal | 1810. D si Bot. maff. 14S7 |
| ^C^A mucronat Cav. | small-horned | | Portugal | 18(13. I) 81 Lob. ic. 2.23 |
| 15* virginknum Lam. | slender | | Spain | 1824. D 81 Spain |
| 7^1 virgittum Lam. | HaWy-leaved | | Spain | 1816. D s.l. Lar. er. 10 |
| 7^1 virgittum Lam. | Virginian | | N. Amer. | D s.l. I*r. er. 19 |
| 7^1 virgittum Lam. | virgate | | N. Amer. | Amer. 1810. D 81 W. Indies |
| 7^1 virgittum Lam. | fetid | | 1714. D 81 Her. lug. 237 | |
| 7^1 virgittum Cav. | serrated | | NewSpainl(MX). | D si Cav. ic. 6. (S4 |
| TM*& Cervantesii Lar. | Cervantes's | | Mexico | D si liar. er. 18 |
| ^ campfote L. | field | | Britain | D 81 Eng. bot. 57 |
| J25 Bourgghti Gou. | Bourgati's | | P.B. S. France | 1731. D 81 Gou il. 7.3 |
| ^C^A lepidostylinum L. | amethystine | | L.B. Styria | 164H. D s.l. M. h. 7.35. 2 |
| 7^1 lepidostylinum L. | triangular | | B. Europe | 1824. D 81 Desf. at 1.01 |
| 7^1 lepidostylinum L. | dilated | | B. Portugal | 1821. D s.l. Lar. er. 4 |
| Kii. Lar. | Anderson's | | B. | 1800. D s.l. |
| Kii. Lar. | sheep | | W. N. Holl. | 1824. D 81 |
| Kii. Lar. | tufted | | N. Spain | 1818. D s.l. Lar. er. 7 |

6633 azitrepum <i>Daf.</i>	azure	A A or 2	jl.au	B	S. Europe 1790.	D s.1	
6634 glomeratum <i>Lam.</i>	glomerate	£ A or 1	jl.au	B	S. Europe 1826.	D s.1 Lar. er. 6	
823. SANI'CULA <i>L.</i>	SANICLB.	<i>(JSano, to heal; supposed healing qualities.)</i>		<i>Umbellifere. a</i>			
6635 europs*a <i>L.</i>	European	£ A w	i*!njj	yH	Britain woods.	D s.1 Eng. bot 98	
6636 canadensis <i>L.</i>	Canadian	5 A un	2 jn.jl	W	Canada 1800.	D s.1	
6637 manländica <i>L.</i>	Maryland	5t A un	1*jn.jl	W.G	N. Amer. 1765.	D s.1 Jac ic. 2.348	
834. ECHINO'PHORA <i>L.</i> SEA PARS.VEP. (<i>Echinos</i> , a hedgehog, <i>phoreo</i> , to bear j flowers.) <i>UmbeUlfer*</i> . 3.							
6638 spinhsa <i>L.</i>	prickly	£ A cu	f jl	W	England sea co.	D s Eng. bot 2413	
6639 tenuifolia <i>L.</i>	thin-leaved	£ A cu	1 jl.au	W	Apulia 1731.	D s 1 M. h. 9.1.2	
6640 trichophylla <i>Sm.</i>	hair-leaved	£ A cu	1 jl	W	Levant 1820.	D co	
825. DAU'CUS <i>L.</i>	CARROT.	<i>(Daio, to separate; discusses flatulence.)</i>		<i>UmbeUlferaf. 15.—21.</i>			
6641 Carbeta <i>L.</i>	common	i Q) m	3 jnj	W	BriUin b. offi.	S s.1 Eng. bot. 1174	
2 hortensis	yellow garden	5 Q) cul	3 apjl	W	Europe gard.	S r.m	
3 aur&itia	/oil-orange	CD cul	3 myjl	W	Europe gard.	S r.m	
4 prolixox	early Ziorf	Q cul	3 my jl	W	Europw gard.	S r.m	
6642 mauritidianus <i>L.</i>	Mauntanian	Q un	3 jnj	W	Spain 1768.	S s.1 Al. ped. 2.61.1	
6643 polygamus <i>Gau.</i>	polygamous	Q un	3 jnj	W	Spain 1817.	S co Jac. vin. 3.78	
6644 maritimus <i>Lam.</i>	sea	Qun	3 jnj	W	Britain Cornw.	S s.1 Eng. bot. 25iH)	
6645 gimmerifer <i>Lam.</i>	gum-bearing	Q	3 jnj	W	Italy 1H18.	S co Bou. mu. 20	
6646 Gingidium <i>L.</i>	Chervil	Q	3 jnj	W	France 1722.	S 81 M. h. 9.9.10	
6647 lucidus <i>L.</i>	shining	Q	3 jnj	W	S. Europe 1807.	S BI M h. 3. 9. 14 4	
16648 muncutus <i>L.</i>	Pk	3 jnj	W	Barbary 1683.	S s.1 M. h. 9.14.4		
6649 aurcus Zoff	golden-flowered	Y	3 jnj	W	& Europe 1823.	S co Desf. at 61	
6650 littoralis <i>Sm.</i>	shore	Y	3 jnj	W	Greece 1820.	S co Fl. gr. 271	
6651 hlpidus i)*T.	hispida	Pk	3 jnj	W	Barbary 1804.	S s.1 Desf. at 63	
6652 meifMius <i>Brot.</i>	Meum-leaved	Pk	3 jnj	W	Spain 1817.	S co Brot. ph. 1.4	
6653 parvifl5rus <i>Detf.</i>	small-flowered	Y	3 jnj	W	S. Europe 1824.	S co Desf. at 1.68	
6654 orinUus .Offf.	long-haired	Y	3 jnj	W	Barbary 1804.	S 61 Desf. at 62	
6655 montevideensis <i>Spr.</i>	Monte Videan	Q un	3 jnj	W	Monte Vi. 1827.	S co	
826. CAU'CALIS <i>L.</i>	BUR PARSLEV.	<i>(Greek name used by Thcophraitius.)</i>		<i>Umbellifera?. 12.—27.</i>			
46636 grandiflora <i>L.</i>	grcat-flowered	O un	1 jl.au	W	S. Europe 1648.	S co Jac. au. 1.54	
6657 latifolia <i>L.</i>	broad-leaved	O w	3 jl.au	R	England ch. ti.	S co Eng. bot. 198	
6658 ffaucoides <i>L.</i>	Carrot-like	O w	11 jn	R	England ch. R.	S co Eng. bot. 197	
16659 platycarpus <i>L.</i>	broad-seeded	O un	* jl.au	W	S. Europe 1800.	co M. h. 9.14.2	
6660 mauntanica <i>L.</i>	Mauntanian	O un	1 my jl	W	Mauri tan. 1818.	co	
6661 leptoph?lla <i>L.</i>	flncleaved	O un	1 jl.au	Pk	Europe 1739.	co Jac. vin. 195	
6662 pbmila IV.	dwarf	O un	1J jl.au	Pk	S. Europe 1640.	co Cav. ic 2.101	
5 6653 maritima <i>Bauh.</i>	sea	O un	i my jl	H	France 1810.	co M. h. 9.14	
\$6664 orientklis <i>Bux.</i>	oriental	O un	4 jn.jl	W	Levant 1699.	co M. h. 9.14.5	
2 pulch?Trima <i>W. en.</i>	molt beautiful	O un	2 jn.jl	W	Caucasus 1816.	co Bux. c. 3.33	
6665 hispida <i>Desf.</i>	hispida	O un	1 my jn	W	Italy 1818.	co	
6666 glkbra <i>Forsk.</i>	smooth	O un	i my jn	W	Egypt 1820.	co For. eg. 23 2,3	
(6667 htorklis <i>Bteb.</i>	shore	O un	1 my jn	W	Caucasus 1819.	co Hof. urn. ed.2 1.	
C&chrys htorialis <i>Spr.</i>						[1.36]	
827. TORPLIS <i>Gae.</i>	TURILIS.	<i>(Toreyo, to carve; seeds.)</i>		<i>UmbeUlferaf. 6.—9.</i>			
6668 ilnhrscus <i>Gm.</i>	Anthnscus	O w	2ijl.au	R	Britain hcd.	S co Eng. bot. 987	
CaArahs ilnhrscus <i>Sco.</i>		O w	1 jl.au	Y	Britain co. fl.	S co Eng. bot. 1314	
6669 infesta H. K.	troublesome	O w	1 my jl	W	Britain co.fi.	S co Eng. bot. 199	
Ca6calis arveisis <i>Hud.</i>	helvetica Gm.	Jl	O w	■ myjl	W	Syria 1817.	co
6670 noritoa <i>Gae.</i>	knotted	O un	■ myjl	W	Egypt 1824.	co	
6671 tuberculata <i>Poir.</i>	tuberculate	O un	■ myjl	W	Austria 1817.	S co Jac.au.29	
6672 tnchouperma <i>Spr.</i>	hair-scseeded	O un	■ myjl	W			
Scandix trichosperma <i>L.</i>		O un	■ myjl	W			
6673 neglecta <i>Spr.</i>	neglected	O un	1 jnj	W			
828. OLIWRIA Vcn. OLIVE HI A.	TURILIS.	<i>(G. A. Olivier, a French botanist.)</i>		<i>Utnbclllferar. 1.</i>			
6674 decumbens Veil.	decumbent	—* O cu	1 myjl	P	Bagdad 1816.	S co Ven.cch.21	
829. LEDEBITI// Lk. LEDEBUBIA.	<i>(A. Ledebure, a botanical author.)</i>		<i>UmbeUtfcree. 1.</i>				
6675 pimpinelloides Lk.	Pimpinella-hkc	■ L un	2 jnj	W	Tauria 182a	S co	
TYgium tauncum <i>Led.</i>							
830. 3/Y'RRHIS Mor.	MYRRH.	<i>(A name used by Dioscoride*)</i>		<i>Umbelliferaf. 1.</i>			
6676 odorata <i>Sco.</i>	sweet-sueted	Jl	£ cc	2} myjn	W	Britain m. pas.	D co Eng. bot 697
831. JIPNINIUM L.	EARTH NIJT.	<i>(Bounos, a hill; habitation.)</i>		<i>UmbeWer*. 4.</i>			
6677 flexufisun 5/i.	flexuous	■ A w	2 myjn	W	Britain past	D co Eng. bot. 988	
Bulboc&Htanuin E. B.							
6678 pyrene un LoL	Pyrenean	A A un	1 jnj	I W	Pyrenees 1826.	D co	
6679 Bulbocaitanum L	common	A A un	i jnj	I W	Portugal 1818.	D co Hof. urn. 1.1 4	
6680 mmno'ides Lk.	Ammi-like	A A un	1 Jnj	jl	Portugal 1820.	D co	
flexuifsum <i>Brot.</i>							
832. WALLRO'TH/jtf Spr.	WALLROTHIIA.	(K WaUroth, M. D., a German botanist)		<i>UmbeWfene. 2.</i>			
6681 tuberosa Spr.	tuberous	A A un	ft myjl	W	Hungary 1823.	D co W. & K. 2.182	
Iftnium alplnum Kit.	Ligusticunj alpliuon Spr.						
6682 spl6ndens Spr.	shining	A A un	1 myjl	W	Pyrenees 1825.	D eo Fl. pyr.85	
Zigdsticum splendens Lap.							
833. (ES A'NTHE L.	WATER DAOPWORT.	(Oinos, fine, anthos, a flower; odor.)		<i>Vmbcllfere. 18.—23.</i>			
6683 fistulosu L.	fstulous	k A P	2 jn.au	F	Britain dit.	D ms Ping. bot 363	
fi6B4 pimpinelloides L.	Pimpinella-like^	A un	jn.au	W	England sal m.	D ms Eng. bot. 347	
6685 gymnorhiza Brig.	naked-rooted	■ jn.jl	W	Baltic	1817. D co		
Excedanifilia IVre.	pimpinelloides						
6686 approximate Merat	approximate	■ ■ ■	■ ■ ■	W	France 1817.	D eo	
Man pimipelloides Thuil.							
6687 involucrul Cav.	involute	■ ■ ■	■ ■ ■	■	Spain 1818.	D co	
fi6B7 involucrul Cav.	Peucedan.-lvd	■ ■ ■	■ ■ ■	■	Europe 18J0.	D aq Lob. ic. 729	
6689 llaifihaBteb.	Silaiimleavcd	■ ■ ■	■ ■ ■	■	Britain dit	D aq Eng. bot 348	
jcucedanifolia E. B.	not Pall.						

6690 <i>virgata</i> Pair.	twiggy	■ ■ ■	jn.au	W	Spain	1820.	D co
6691 <i>prolifera</i> L.	proliferous	■ ■ ■	n.au	W	Italy	1739.	D co Jac. vin. 3.62
6692 <i>globulbsa</i> L.	<i>globe-headed</i>	■ ■ ■	n.au	W	Portugal	1710.	I co Gou.il. 18.9
6698 <i>crocata</i> L.	saffron-colored	■ ■ ■	n.au	W	Britain	dit.	I m.s Eng. bot 2313
«34 <i>apiifolia</i> Brot.	Parsley-leaved	■ ■ ■	n.au	W	Portugal	1806.	D co Sub. rom. 84
6695 <i>interrGpta</i> 7%un.	interrupted	■ ■ ■	jn.jl	W	C. G. H.	1818.	S co
6696 <i>int-brians</i> Thun.	inebriating	■ ■ ■	au.s	W	C. G. H.	1816.	S co
6697 <i>tenuifolia</i> 7%».	fine-leaved	■ ■ ■	jn.jl	W	C. G. H.	1819.	S co
«698 <i>austrklis</i> Wul.	southern	■ ■ ■	jn.jl	W	& Europe	1817.	I co
6699 <i>gluf.ca</i> Dec.	glaucous	■ ■ ■	jn.jl	W	S. Europe	1823.	D co
6700 <i>nodifl&ra Schous.</i>	knot-flowered	■ ■ ■	jn.au	W	N. Africa	1820.	S co
<i>Cbnium dichotomum Brou.</i>							
834. CRITHMUM L.	SAMPHIRE.	(Krithe, barley; similarity of seeds.) <i>UmbelVferaf.</i> 8.—4.					
6701 <i>maritimum</i> L.	sea	■ A cul	1	jl.s	W	Britain	s. cliffs. D r.m Eng. bot 819
6702 <i>latifolium</i> L.	broad-leaved	■ iAJ cu	li	jl	Y	Canaries	1780. D r.m
Tendria canariensis Spr.							
•835. ilTHAMA'NTA L.	SPICKNEL.	(Found upon Mount Athamas in Sicily.) <i>Umbellifera.</i> 13.—14.					
\$6703 <i>Libanitis</i> L.	Rosemary	■ Aw	2	jn.jl	W	England	ch.pa. S co Eng. bot. ISX
6704 <i>Cervaria</i> L.	Cervaria	■ A un	4	jl.au	Pa.P	Europe	1597. S co Jac. au. 1.69
Ligubicum Cervaria Spr.							
6705 <i>sibirica</i> L.	Siberian	■ A un	■	jLau	W	Siberia	1771. ■ co Om. ri. 1.40.1,8
6706 /macifolia Spr.	Panax-leaved	■ A uu	■	jl.au	W	Spain	1823. ■ co Boc. sic. 1
Cachrys panacifolia Vahl							
6707 condens&ta L.	<i>close-headed</i>	■ A or	■	jl.s	W	Siberia	1773. ■ co Gou.il 83.26
6708 <i>inchna</i> W.	ho<try	■ A un	■	Jau	W	Siberia	1802. ■ co
6709 <i>sicula</i> L.	Sicilian	■ A un	■	jnjl	W	Sicily	168a ■ co Za. h. 70.48
6710 <i>Matthloli</i> Wal.	Mattluoli's	■ A or	■	jl.jl	W	Carniola	1802. ■ co Jac. le. 1.57
6711 <i>retensis</i> L.	Cretan <i>Carrot</i>	■ A un	■	jl.jl	W	Austria	1596. ■ co Jac.au. 1.62
6712 &nnua	annual	O un	■	jl.jl	W	Candia	1731. ■ co
6713 <i>lasiantha</i> IV.	hairy-petaled	■ A un	■	jn.au	W	Spain	1818. ■ co
6714 <i>tortuba</i> Spr.	twitted	■ I cu	■	jn.au	W	S. Europe	1826. ■ s.p Desf. at. 1.73
6715 <i>carwfolia</i> Step.	Caraway-leaved	■ A un	■	jl.jl	W	Siberia	1819. ■ co
836. PIMPINE'LLA L.	BURNET SAXIFRAGE.	(Bipinnula, twice pinnate j leaves.) <i>UtnbeUifene.</i> 10.					
6716 <i>Saxifraga</i> L.	Saxifrage	■ 1	jn.au	W	Britain	dry pa. S co Eng. bot. 407	
6717 l&tea Desf.		■ 4	jn.au	W	S. Europe	182a S co Desf. at 76	
6718 <i>intermedia</i> Nut.	intermediate	■ li	jn.au	W	Europe	1827. S co	
6719 <i>cretica</i> Poir.	Cretan	■ 1	jn.au	W	S. Europe	1826. S co	
6720 <i>rigras</i> W.	<i>huck-rooted</i>	■ 1	jn.au	W	Germany	1683. S co Boc. mu. 80	
6721 <i>magna</i> L.	great	■ 4	jn.au	W	England	woods. S co Eng. bot. 408	
2 dissecta	<i>cut-leaved</i>	■ 4	jn.au	W	Europe	1800. S co M. h. 9.5	
6722 <i>media</i> Hafm.	mediate	■ 4	jn.au	W	Europe	1817. S co Jac. au. 397	
6723 <i>dissecta</i> Hrtz.	<i>cut-leaved</i>	■ li	jn.au	W	England	dry pL S co	
6724 <i>rotundifolia</i> Spr.	round-leaved	■ 2	jn.au	W	Caucasus	1817. S co Spr. um. 2.4.1.4	
6725 <i>tenuis</i> Sieb.	slender	■ A un	1	jn.au	W	Crete	1820. S co
837. 7HA'NIUM Spr.	TRAGIUM.	(Tragos, a goat; odor.) <i>UmbeUfer<e.</i> 7.—10.					
6726 <i>jntaum</i> L.	Anise	O cul	1	jn.au	W	Egypt	1551. S co
Sison ^n'sum Spr.							
6787 Colfmmne Spr.	Columna's	■ A un	2	jn.au	W	S. Europe	1810. S co Col. ph. 17
Pimpinellii TYagium Lam.							
6728 <i>depresum</i> Sieb.	depressed	■ A un	li	jn.au	W	Crete	1822. ■ co
6729 <i>aromaticum</i> Spr.	aromatic	Q) un	1	jn.au	W	Caucasus	1820. ■ co
6730 Briteri Spr.	Brotero's	■ A un	li	jn.au	W	Portugal	1822. ■ co
Pimpinella ftubonoldes Brot.							
6731 <i>villosum</i> Desf.	villous	■ A un	2	jn.au	W	Morocco	1817. S co
Schousboei Spr.							
6738 <i>pcregrinum</i> Spr.	foreign	■ A un	8	jn.au	W	Italy	1640. S co Jac. vin. 8.131
838. PHELLA'NDRIUM L.	WATER HEMLOCK.	(Phelco, to deceive, aner, a man.) <i>Umbellifcne.</i> 1.					
6733 <i>aquaticum</i> L.	aquatic	k Q> P	3	jnjl	W	Britain	rivul. D aq Eng. bot. 684
839. DO'NDII* Spr.	DONDIA.	(Dondie Dupre'e, a French botanist) <i>UmbeUfrene.</i> 1.					
6734 Epipactis Spr.	Epipactis	■ A pr	4	mr.ap	Y	Alps	1823. D pl Jac.au. 5. U
Astrandia Epipactis L.							
840. TRACHYSPE'RMIUM Lk.	TBACIYSPERMUM.	(Traehys, rough, sperma, seed.) <i>Umbelliferte.</i> 1.					
6735 cōpticum Sp.	Egyptian	O un	2	jnjl	W	Egypt	1773. S co Jac vin. 2.196
A mm\ cōpticum L.							
6736 rigens G. Don	stiff* tt.	i_J cu	li	jnjl	W	C. G. H.	1787. S co
Ctinum rigens L. 2ftnum rigens Spr.							
Wl. i<'MMI L.	AMMI.	I Ammos, sand; habitation.) <i>UmbellUferaf.</i> 8.					
22 ^y H»*K* Lam-	Visnaga	Q un	2	jn.au	W	S. Europe	1596. S co Gae. fr. 1.107
SiiX "ne'tnifMium Lam.							
6740 Joeberi Hok.	Fennel-leaved	Q un	■	jnjl	Y	S. Europe	1551. S co Black. 447
Ul'crulaBfolium Lag.	Bocher's	Q un	■	jnjl	W	Caucasus	1818. ■ co
G74B acatilis Spr.	Ferula-leaved	Q un	■	jnjl	W	Crete	1817. ■ co
iftinnum aca6le Bieb.							
6743 glauriflum Lap.	glaucous-lvd	J O un	li	jnjl	W	Chile	1822. ■ co
0744 ttauciflum Seo.	Carrot-leaved	5 A un	2	jl.au	Pa.Y	& Europe	1820. ■ co
I Ammos, sand; habitation.) <i>UmbellUferaf.</i> 8.							
842. mTBON L.	BUDON.	(Boubon, the groin: supposed medical virtues.) <i>UmbellUfer*</i> . 5.-7.					
V>745 maced6nicum W.	Macedonian	CD cu	2	jn.au	Pa.Y	Greece	1596. S co Black. 382
i^th^manta maced6nica Spr.	stitHeaved	3	jn.au	Pk	Sicily	1710.	Boc. mu. 8.76
(6746 rigidum Spr.	gum-bearing	3	jl.s	Pk	Crimea	1804.	Sm. ex. bot 120
2 gummiferum Sm.		21	jn.au	W	Siberia	1816.	
46747 buchtornense Fis.	Buchtorn						
ilthamaftha rlgida Horn.							
*6748 glaucum Spr.	glaucous	■ o	■	lijnjl	W	Tauria	1817. S co
Seseli iictra'um Bieb.							
g49 siculum Spr.	Sicilian	■ o	■	W	Sicily	1820.	M. h. 9.
% TOO dich6tomum Spr.	dich otomous	■ o	■	W	Caucasus	1818.	Bot. mag. 8073
Ressell dich6tomum Bkb.							

- | | | | | |
|--------------------------------------|------------------------|--|--|--------------------------------------|
| an. CICITTA L. | COWDANE. | (A name of doubtful meaning used by Pliny.) | <i>Umbelliferae.</i> | 4 |
| (815 virusa L. | poisonous | ^ m 3 jl | W Britain ditch. | D.m.s Eng. bot 479 |
| 6616 maculata L. | | A P jjjl.au | W N. Amer. 1759. | D e o Pluk. al.*76.1 |
| 6817 daurica Fl.*. | Daurian | A un 3 jl au | W Siberia 1817. | D co Gm. *1.47 |
| 6818 bulbifera L. | bulbiferous | A un 1 jl.au | W N. Amcr. 1810. | D co |
| • 852. CO ^N IUM L. | HEMLOCK. | (<i>Konao</i> , to whirl round) | causes giddiness and death) | <i>VmbeUlfera.</i> 4. |
| 6819 maculatum L. | spotted | Q) m 5 jn jl | W Britain hcd. | S co Eng. bot. 1191 |
| 6820 croaticum J.«. | Croatian | Q r a 6 jn.jl | W Hungary 1818. | S co |
| 6821 <i>Armaadacha</i> Hnok. | Arracacha | ** G3 cul | G.w S. Amer. 1823. | I) r.m Hook. ex. fl. 152 |
| 6822 moschatum //, \$ B. | musk | ■ jnjl | W S. Amcr. 1824. | D r.m H.&B.n. 5.420 |
| 853. CAPNOPHYLLUM G<>. | (&) cul | (<i>Kapnos</i> . smoke, <i>phyllon</i> , a leaf.) | <i>UmbeUlfera.</i> I. | |
| 6823 africanum Gwr. | African CAPNOPHYLLUM. | 3 jn.s W C.O.H. 1759. | S s.l Jac vin. | 2. Id* |
| Conium africanum L. | Rhvdxa capensis Lk. | | | |
| *854. SMY'RNIUMi. | ALEXANDERS. | (<i>Smyrna</i> , myrrh; similar qualities.) | <i>UmbeWfere.</i> 9. — 1 | |
| 6824 Dioscoridii Spr. | Dioscoride's | CD or 2 myjn Y | & Europe 1700. | S co W. & K. 1. |
| perfoliatum mot. | | | | |
| 6825 Dodona <i>^i</i> Spr. | Dodonus's | Q) or 2 my.jn | Spain | — S co Lob. ic. 790 |
| perfoliatum L. | | | | |
| 6826 Oluskrum L. | Olusatrum | 3(Q) cul | 4 myjn | G Britain sea co. S s.l Eng. bot 230 |
| 6827 apuūtiHun JV. | Smallage-lvd | ^ A or 1 myjl | P.Y Camria | 1731. I) s.l |
| 6828 cicutarium Bkb. | Cicutn-like | ^ A or 2 myjl | G.w Caucasus | 1827. D co |
| \$6829 nudicaulc Bieb. | naked-stmd | ^ A or 3 myjn | G Caucasus | 1817. D co Spr. um.4.7 |
| 6830 cordatum Walt. | hearUfeated | ^ A or 1 jnjl | D.P N. Amer. 1597. D s.l | |
| trifoliatum Bart. | Thapsia tñfolida L. | golden | 2 my.jn Y | N. Amcr. 1699. D r.m |
| ^6831 attreum L. | | | | |
| Sson afircum Spr. | | | | |
| 6832 integrerrimum L. | cntireJcavcd | £[A un 1} jn | Y | N. Amer. 1759. D r.m |
| 5ison integrerrimum Spr. | | | | |
| 855. iTPIM L. | PARSLEY. | (<i>Apon</i> , water, <i>Celt. i</i> habitation.) | | |
| 46833 Petrosellum L. | True Parsley | L Q cul 3 jn jl | L.Y Sardinia | 1548. S r.m |
| 6834 gravēlons L. | heavy-smelling Celery* | Q cul 4 jn.au | W Britain ditch. | S m.s Eng. bot. 1210 |
| 6H35 prostratum Lab. | prostrate | £ iAI un ij,jl | W V. Di. L. | 1826. S co Ven. mal. 81 |
| 6836 romanum Zuc. | Roman | Q) un 1 jn.jl | W Italy | 1820. S co |
| 6837 fractoph [^] llum Hofm | broken-leaved | CD un 1 jn.jl | — 1826. S co | |
| 6838 ternatum Pall. | ternate | ^t A un 1 jn.jl | Siberia 1824. S co | |
| 856. JEGOPOPDIDIUM L. | GOOT-WEED. | (<i>Aix</i> , a goat, <i>podion</i> , a little foot; leaves.) | <i>Umbrillfcrtv.</i> 1. | |
| 6839 Podagraaria L. | Podagria, or com. | Am 2 my.jl | W Britain sh. pi. | D m.s Eng. bot 9M |
| 857. METUM Tou. | BAWD-MONEY. | (Afcion, smaller; delicacy of leaves.) | <i>Umben'fene.</i> 4. | |
| 6840 /ftinius Jac. | Square Parsley | O un 1 jl | P 1778 | Jac. vin. 2.1.W |
| M | Mutellina | ^ A un | | |
| 6841 Mutellina Gae. | Mutellina | 1 jla | P Austria 1774. D e o Al.ped.G.0.1 | |
| Phellandnum Mutellina L. | | | | |
| 6842 athamanticum Joe. | Athamanta-lk | 3t A or | 1\$ ap.jn P.Y Britaj me. pa. I) m.s Jac. au. 4.303 | |
| yitham&nta A/eum L. | | | | |
| 6843 sibiricum 5pr. | Siberian | ^ A un | 6 jn.jl Y Siberia 1818. D co | |
| Zig6sticum arenarium Lk. | | | | |
| • 6844 graveolens L. | DILL. | { <i>Ano</i> , upward, <i>theo</i> , to run; grows quick.) | <i>Uml*IHfcne.</i> 5. - 6. | |
| 6845 setigenum L. | heavy-smelling | Q cul 3 jnjl | Y Spain* 1570. S r.m | |
| 6846 Sow Rox. | corn | O un fijn.jl | Y Portugal 1796. S co | |
| 6847 Foeniculum L. | Sowa | O m 1 jn.jl | Y E. Indies 1810. S co | |
| Jl/eum Foeniculum Spr. | Fennel | ^t A cul 6 jl.au | England ch. pi. S B.1 Eng. bot 1208 | |
| 2 dtice | sweet | A cul 4 jl.au | Italy 1824. S co | |
| 6848 pijierktum Bert. | peppered | Q cul 6 jnjl | Italy 1824. S co | |
| • 059. CA% RUM£. | CARAWAY. | (A native of <i>Caria</i> , in Asia Minor.) | <i>Umbellifern.</i> 8. | |
| Wto Cdruil, | common | & CD dt 2 my.jn | W Britain me. pa. S B.1 Eng. bot 1503 | |
| 6850 simplex W. | | & CD un 1 my.jn | W Siberia 1816. S s.l | |
| *860. CNI'DIUMCtt*. | CNIDIUM. | (The ancient name of an orach.) | <i>Umbelliferar.</i> 7. — 10. | |
| 6851 Monnier Cus. | Monnier's | O un If jl.au | W S. Europe 1771. S co Jac. vin. 1.62 | |
| 6852 Silatis Spr. | SilaUs | A 2 jn.au | Y England mca. D co Eng. bot 2142 | |
| Peucedanum SilaUs L. | | | | |
| ^6853 alsaticum Spr. | Alsatian | A un 6 jn.au | L.Y Austria 1774. D co Jac. au. 1.70 | |
| /cuedanum alsaticum L. | | | | |
| 6854 pyren'sum Spr. | Pyrencean | A un 1 jnjl | W Pyrenees 1731. D co Gou.il. 11.5 | |
| 6855 canadense Spr. | Canadian | A un 2 jn.au | W Canada 1817. D co | |
| SeUnum canadonse Mx. | | | | |
| 7856 atropurpureum Spr. | dark-purple | A un 2 . . . | Philadelp. 1810. D co | |
| Snymium atropurpureum Lam. | | | | |
| 6857 Ffschcri Spr. | Fischer's | A un 2 jn.au | W Siberia 1820. D co | |
| Angélica Ffschcri R. & S. | ^thamantia dehuata | Fijn.au | W Siberia 1820. D co | |
| 858. JJUPLEimUM L. | HARE'S-EAR. | an ox, <i>pleuron</i> . a rib; leaves.) | <i>UmbeUlfera.</i> 34. - 37. | |
| w ⁸ rotunoifohum L. | round-leaved | 2 jn.jl | Y England co. fl. S co Eng. bot 99 | |
| TS ubovatum Lk. | subovate | imy.jl | Y Spain 1819. S co | |
| j ¹⁰ heterophyllum Lk. | various-leaved | 1 my.jl | Y Egypt 1818. S co | |
| Wil longitium W. | long.leaved | 3 my.jl | G.v Switzerl. 1713. D co Cam. h. 38 | |
| J ¹² afireum Fis. | golden | 1 my.jn | Y Siberia 1820. D co | |
| 6K13 angulosum L. | angulose | i myjl | G.Y Switzerl. 1759. D co | |
| W6 pyrenaicum Gou. | Pyrcnean | 1 my.jl | G.Y Pyrenees. 1814. D co Gou. it 4.1.2 | |
| @65 graminifolium Vahl | Grass-leaved | I myjl | G.Y Switzerl. 1768. D co Jac. ic. 1.56 | |
| 6861 petra [^] um L. | rock | M myjl | G.Y Switzerl. 1762. D co Pluk. ph. 50.5 | |
| 6867 stellatum L. | starry | 1 my.jl | G.Y Switzerl. 1775. D co Hal. h. 771.18 | |
| 8 ⁸ falcatum L. | falcate | J m y ? | G.Y Germany 1739. D s.l Jac. au. 2.158 | |
| S5 ⁹ rtgidum L. | rigid | 2 my.jl | G.Y France 1820. S co M.h.9.12 | |
| ISL9 P ⁹ yp'Huin Led. | many-leaved | 2 my.jl | G.Y Caucasus 1823. S co Za.h.40 | |
| ggl baldense Tou. | Mount Bakio | 1 my.jn | G.Y Mt. Baldo 1817. S co | |
| E42 sibiricum Vest. | Siberian | 2 il | G.Y Siberia IH26. S co | |
| 70 ³ exalthum Bkb. | exaltctl | | Tauria 1807. D s.l | |

6874 n&dura <i>H. K.</i>	<i>naked-stalked</i>	I AJ pr 1} o	G.Y	C.G.H.	1778.	C Lp		
6875 tranunculoides <i>W.</i>	Crowfoot-bke	A Pr i Jl*au	G.Y	Pyrenees	1790.	D Lp	Par.th.7	
6876 gnricifolium <i>W.</i>	Carex-lcaved	A Pr i Jl*au	G.Y	France	1817.	S co		
6877 Buncridnum <i>W.</i>	Burser's	A Pr i Jl*au	G.Y	1817.	S co		
6878 paniculatum <i>Brot.</i>	paniculate	pr li jn au	Y	Spain	1824.	S co		
6879 trifidum <i>Ten.</i>	tnfid	pr 2} jn.au	Y	Italy	1824.	S co		
6880 d*inceum <i>L.</i>	R.	Jl*au	G.Y	S. Europe	1772.	S co		
6881 Pollichit <i>R. & S.</i>	Pollich's	Jl*au	G.Y	Palatine	1818.	S co		
6882 nodiflbrum <i>Sm.</i>	knot-flowering		G.Y	Greece	1823.	S co	Fl.gr. 260	
6883 oppositifolium <i>Lap.</i>	opposite-lvd		G.Y	Pyrenees	1819.	S co	Fl.pyr.73	
6884 lanciflrium <i>Horn.</i>	lance-lcavcd		Y	Tauna	182a	S co		
6885 OdontUes <i>L.</i>	Odontites		G.Y	Italy	1749.	S co	Jac. vin.3.91	
6886 semicompitum <i>Xr.</i>	semi-compound		G.Y	Spain	1778.	S co	Gou.il. 7.1	
6887 grfcile <i>Bieb.</i>	slender		G.Y	Caucasus	1820.	S co		
6888 glaucum C. *R>	glaucous		G.Y	S. Europe	1819.	S co		
6889 tenuissimum <i>Z.</i>	slenderest		G.Y	England sea sh.	S s	Eng. bot 478		
6890 Gerard Jac.	Gerard's		G.Y	S. Europe	1804.	S co	Jac.au. 3 256	
6891 lancifolium /forn.	lance-leaved		G.Y	1823.	S co		
862. TENO'R/!* Spr.	TENORIA.	(A. Tenore, professor of botany at Naples.)				Umbellifertr.	8.	
6892 arborescens Spr.	Arborescent	I l-Jpr 1 aus.	G.Y	C. G. H.	1810.	C ls		
Apledrum arborescens <i>Thun.</i>								
6893 fructictom Spr.	shrubby		G.Y	S. Europe	1595.		Den.br. 14	
6894 corftcum Spr.	coriaceous	i-Jpr	G.Y	Gibraltar	1784.		Jac. ic. 23"II	
6895 firuticescens Spr.	fruticetcent		G.Y	Spain	17&		Cav. ic. 2.105	
6896 canescens Schous.	canscent	i-Jpr	G.Y	Barbary	1809.			
6897 plaitaflncuni Spr.	Plantain-wi		G.Y	N. Africa	1810.		Desf. at 1.57	
6898 spinbusm Spr.	thorny		G.Y	Spain	1752.		Gou.il. 8.2.3	
6899 diffurm Spr.	two-formed		G.Y	C. G. H.	1752.			
* 863. HYDROCOTYLE L. PENNYWORT. (<i>Hydro</i> , water, <i>kotyle</i> , a cavity; leaf.)						Umbellifene.	18 — 60.	
6900 vulgaris L.	common		imy.jn	R	Britain	wa. pi. 1)	c.p Eng. bot. 751	
6901 natans Cyr.	nodding		jn.au	R	Italy	1819.	D p Cyr. ne. 1.6	
6902 amencana L.	American		i my.au	G	N. Amer.	17X.	D P Spr. um. 2.3	
6903 peduncularis R. Br.	pedunculate		my.au	G.Y	N. Holl.	1823.	D 8	
6904 umbellata L.	umbell'd	nH A un	ijl.au	G	N. Amer.	1795.	D p Spr. um. 1	
6905 bonariensis Lam.	Buenos Ayres %, LAJ un		G	Peru	1810.	D p	Cav. ic. 5.438	
6906 nepalensis Hook.	Nepal		G	Nepal	1820.	D co	Hook. ex. fl. 30	
6907 nitidula Rich.	rather shining ft* E2 un		my	G	Java	1820.	D co	Hook. ex. fl. 29
6908 villtaa L.	villous & LAI un		1 au	G	C. G. H.	1795.		
6909 crnifrmis Thun.	kidney-shaped ijE iAJ un		ijl.au	G	C. G. H.	1820.		
6910 triloba Thun.	three-lobcd		LAI un	i jn.au	G	C. G. H.	1810.	
6911 chinensis L.	Chinese ft,	LAJ un	ijn.au	G	C. G. H.	1818.	B5 Th.dis.2ded.3	
6912 asiatica L.	Asiatic (&, LAI un	1 jl.au	G	C. G. H.	1690.	R. mal. 10.36		
6913 plebba R. Br.	plebeian ft,	LAI un	imy.au	G.Y	N. Holl.	1823.	D co	
6914 nirsikta Sun.	hairy £, E) un	i jlau	G	W. Indies	1810.	D p		
spicata Lam.								
6915 sibthorpeida Lam.	Sibthorpia-like Jk E3 un	\ my.jl	W	I. France	1800.			
6916 ranunculus L.	Ranunculus-Ik jfc, -AJ un		W	N. Amer.	1800.		Spr. um. 1.2.	
6917 repanda Per*.	repand ft,	-AJ un	1 jn.jl		N. Amer.	1806.	D p	Spr. um. 2. 4.
864. BOWLES'S/J R. & a BOWLESEA.	(TV. Bowles, an Irish botanist.)					Umbellifero.	I—5.	
6918 tenera Spr.	tender JW -AJ cu	i jn au	G.Y	M. Video	1827.	D co		
865. SPANA'NTHE Jac.	SPANANTHE.	(Spanos, rare, anthon, a flower; umbels.)				UmbeWfer*	1—a	
6919 paniculata Jac.	panicled	jg Q un	2 jl.au	W	Caracas	1795.	S s.l Jac ic. 2.350	
HydrocotyleSpana'nthe IV.								
866. DRIP&f Doc.	DRUSA.	(M. Le Dru, a botanist.)				Umbelliferar.	1.	
6920 oppoaitifolia Dec.	opimsite-lcaved^ iAJ un	ijl.au	W	Tencrife	1824.	D e o	An. mu. 10.38	
*867. TRACHYMETNE Rud.	TnAcnYMENE	(Trachys, rough, hymen, membrane; involuc.)				Umbellifera> 2-5.		
6921 fimbriis Spr.	linear-fibwd		L JCU a 5* jnau	W	XN. Holl.	1824. CC sp	Cav. ic. 5.485	
6922 indsa Rud.	cut-leaved		I Jcu	—	N. Holl.	1819!	C fl.p Lin.tr." 10.21.2	
868. ULOSP'E'RUMM Lk.	BROADSEED.	(Oulos, curled, sperma, a seed.)				UmbeWferte.	1.	
6923 dichotomum Lk.	dichotomous	O cu	1 jn.jl	W	Barbary	1800.	S co Desf. at 1.66	
Cbnium dichotomum Desf.								
869. OTHITSA L.	FOOL'S PARSLEY.	(Aithusso, to make hot; acrid.)				VmbelMeree.	4.	
6924 Cytilipium L.	Dog poison, or cam.	O Jl*	>X	Britain	co.fi.	S co Eng. bot.	1192	
& Ys cynapioides Bieb.	Cynapiuin-like	O lijl.au	W	Caucasus	1817.	S co Hof. um. 1. 9		
6925 elcta Hafn.	superb	O U jl.au	W	1820.	S co Hof. um. 5.6.5		
6927 fatua H. K.	insipid	O 2 Jls	W	—	1781.	S co		
870. IMPERATCTRIA L.	MASTERWORT.	(Reputed imperial virtues in medicine.)				Umbellifera?.	3.	
6928 Ostruthium L.	Sparrow, or ctn.^i A m	2 my.jl	Pk	Scotland	m.al p.	D co	Eng. bot 1380	
6929 angustifolia Bel.	narrow-leaved ^ & un	2 jl.au	P	S. Europe	1819.	D co	Lob. ic. 700	
6930 mexickna // C.	Mexican	5 A or 4 jl.au	G.Y	Mexico	1818.	S co		
871. CALLISACE Fis.	CALLIBACE.	{Kallas, beauty, sakos, a buckler; seeds.)				Umbellifero.	L	
6931 dahurica Fis.	Dahurian		A un 2 jlau	G	Siberia	1816.	D co	
872. THYSSELNUM Hafm.	TJIYSBELINUM.	(Thyo, to burn, selinon, parsley; acrid juice.)				UmbeUlfene.		
6932 palastrc Hafm.	marsh	& A un	3 jl.au	W	Britain	mars.	D m.s Eng. bot. 229	
Scfnum palustre L.								
6933 Plinii Lob.	Pliny's		A un 3 jl.au	W	Italy	1800.	D ee	
£cnum Plinii Hal.								
*873. SELVSUML.	MILK PARSLEY.	(Selene, the moon; nfgricans Gau.				UmbeWfera.	23.—	
\$6934 montanum W. en.	mountain		A un 3 jl.au	W				
nfgricans Gau.								
vn&* austrlacum Jac.	Austrian		2 jl.au	W	Austria	1804.	D co Jac au. 1.71	
6136 cartijlia L.	Caraway-leaved		2 jl.au	W	Austria	1774.	D co Jac au. 1.16	
\$6937 Chubr&N Jac.	Chabra's		lijl.au	W	Austria	1791.	D co Jac.au. 1.72	
Imperatbra Chabrss' Spr.								
\$6938 Segucri L.	Seguier's		4 jl.au	W	Italy	1774.	D s.l Jac. vin. 1.61	
Imperatbra Seguierj Spr.								

6939 latifolium <i>Bieb.</i>	broad-leaved	fr A un 2 il.au	W	Caucasus 1816.	D s.l	
46940 Oreosennum <i>Sco.</i>	Mountain Parsley	A un 2 ij.l.au	W	Europe 1810.	O s.l	Jac. an. 1.68
^thaminta Oreosennum <i>L.</i>						
46941 dccccinns <i>Wnl.</i>	deceiving	i A un 2 jn.jl	W	Madeira 1787.	D s.l	Sc. nan. a 1.13
6942 sih ricum <i>Rctz.</i>	Siberian	U A un 2 jl.au	W	Siberia 1810.	D co	
6943 polymorphum <i>Spr.</i>	multiform	Si A un 2 jl.au	W	Russia 1810.	D co	
46944 baicalense <i>IV.</i>	Baikal.	U A un 2 in.au	W	Hungary 1820.	D co	
46945 lineare <i>Schum.</i>	linear-Arc	vaf S A u:i S jl.au	W.R	Europe 1817.	D co	FL dan. 330
Ferula rablense <i>WuL</i>						
6946 terebinthaceum <i>L.</i>	turpentine	■ A un 2 jl.au	W	Siberia 1816.	D co	
46947 poucedanoides <i>Detf.</i>	Peucedanum-lk	■ A un 2 jl.au	W	& Europe 1810.	D co	
6948 ven&lum <i>Spr.</i>	Venetian	■ A un 2 jn.au	W	S. Europe 1810.	D co	
6949 rigidulum <i>Viv.</i>	rigidish	■ A un 1 jl.au	W	Italy 1826.	D co	Viv. fr. 1.15
6950 scabrum <i>Lap.</i>	rough	■ A un 1 jn.au	W	Pyrenees 1824.	D co	FL pyr. 77
6951 podoticum <i>Bes.</i>	Podolian	■ A un 1 jn.au	W	Podolia 1817.	I> co	
6952 elegans <i>Bnlb.</i>	elegant	■ A un 1 jl.au	W	Europe 1817.	D co	Bal.st4
6953 Bellardi <i>Balb.</i>	Bellardi'a	■ A un U jn.au	W	S. Europe 1817.	D co	
6954 pyrenaicum <i>Gou.</i>	Pyrencean	■ A un 2 ij.l.au	W	Pyrenees 1800.	D co	Gou. il. 3.5
Angelica pyrenaea <i>a Spr.</i>						
6955 caucasicum <i>Bieb.</i>	Caucasian	■ A or ij.lau	Y	Iberia 1823.	D co	
Imperatona caucasica <i>Spr.</i>						
6956 angchastrum <i>Lk.</i>	Angelica-like	■ A u:i Iij1.au	W	Pyrenees 1817.	D co	Gou.il. 5.5
Angelica lancifolia <i>Spr.</i>	A'cseh pyrenarum <i>L.</i>					
•874. ANGE'LICAZ.	ANGELICA.	(Supposed angelic virtues.)		UmbeUifera. 9.—10.		
46957 Archangelica <i>L.</i>	archangel	Q cul 4 jn.au &	England	wa.pl. S m.s Fl. dan. 206		
46938 atropurpurea <i>£.</i>	dark-purple	Q un 6 jl.au	Canada	1759. S co Cor. ca. 199		
6959 lucida <i>L.</i>	shining	Q un 2 jl.au	W	Canada 1640.	S co Jac. vin. 3.24	
6960 triquintae <i>Mx.</i>	tnquinate	A un ij.lj.au	W	Canada 1810.	D p Spr. um. 6.2	
6961 sylvestris <i>L.</i>	wood	Aw 6 in.au	F	Britain m.wo.	D in.s Eng. bot. 1128	
69H Razoul'i Gou.	Razoul's	A un 2 jn.au	Pa,P	Pyrenees 18 Iti. 1) co Gou. il. 13. fi		
6963 verticillata <i>L.</i>	whorled-jfrvd	A un 6 1 G	Italy	1683. D co Jac. vin. 2.130		
Imperathria verticillaris <i>Dec.</i>						
6964 pratensis <i>Spr.</i>	meadow	■ A un 2 jn.au	F	Galicia 1818.	D co	
Imperatona palustris <i>Bes.</i>						
6965 flavescens <i>Bes.</i>	flavescens	■ A un 2 jl.au	Y	Galicia 1818.	D co	
875. ZIGU'STICUMZ LOVAUB.	(Growing in great abundance in Liguria.)	Umbelliferof. 14.—20.				
46996 Levisticum <i>L.</i>	common	■ ■ cul 6 jn.jl P.Y	Italy	1596. D co		
6967 scAticum <i>L.</i>	Scotch	■ ■ cul 2 jn.jl W	Britain	sc. sh. D co	Eng. bot. 1207	
^6968 nodiflbum <i>Vil.</i>	knot-flowering	■ ■ 3 jl.au G	Alp. Eur.	1810. D co	Al. ped. 72	
Smyrnium nodiflbum <i>All.</i>						
6969 peloponnestmuni <i>L.</i>	Peloponnesian	■ ■ 4 my.jl P.Y	Switzerl.	1596. D co	Jac.au. 5.13.	
6970 Sprengtli <i>Bieb.</i>	Sprengel's	■ ■ 2 jl.au	W	Italy 1823. D co		
6971 alatum <i>Spr.</i>	winged-dwarf	■ ■ 5 jl.au R.W	Caucasus	1816. D co		
y^thamfanU alata <i>Bieb.</i>						
6972 othamantoldes <i>Spr.</i>	Atharaanta-lk	■ ■ A un 2 ij.l.au	IV	Pyrenees 1817.	S co	Jac. vm. 2.197
A thamanta pyrenaica <i>Jac.</i>						
6973 pyrenaicum <i>L.</i>	Pyrenean	■ ■ A un 3 jl.au	W	Pyrenees 1804.	S co	Gou. il. 14.10
Grindium pyrenaicum <i>Spr.</i>						
6974 cendicans // K.	whitish	■ ■ 2 jl.au P.Y	—	178a S co	Jac. vin. 3.18	
6975 peregrinum <i>L.</i>	foreign	■ ■ 2 jn.jl L.Y	Portugal	1633. S co		
6'tson peregrinum <i>Spr.</i>						
6976 baleanicum <i>W.</i>	Balearic	■ ■ 1 jn.jl Y	Minorca	1804. D co		
6977 longiflbum <i>L.</i>	long-leaved	■ ■ 3 jn.jl Pa,P	Siberia	1804. D co	M.h.9.15.1	
6978 actsiflbum <i>Mx.</i>	Alder-leaved	■ ■ 3 jl.au W	Canada	1810. D co		
6979 obtusiflbum <i>Horn.</i>	obtuse-leaved	■ ■ 1 jl.au W	N. Africa	1824. D co		
876. PHYSOSPE'RUM Cus. PHYSOSPERMUM.	(to inflate, sperma, seed.)	UmbclUfene** I.				
6980 commutatum, <i>Spr.</i>	altered	^ A un li jl.au W	England	bu. fi. D co	Eng. bot. 683	
.Ligisticum cornubiense <i>L.</i>						
877. PLEUROSP'E'RUM llqfnu PLEUROBPEBMUM.	(Pleuron, a rib, sperma, seed.)	UmbelUfers. 1.				
6981 austriacum <i>Hgfm.</i>	Austrian	^ A un 2 jl.au W	Austria	1597. D co	Jac. au. 2.151	
/ig&sticum austriacum <i>L.</i>						
878. H ASSELQUI'ST/J L HASSELQUISTIA. (F. Hasselquist, M.D., its discoverer, d. 1752.) Umbeliteree. 2.						
6982 aegyptiacum <i>L.</i>	Egyptian	f) un U jl W	Egypt	1768. S co	Jac. vin. 1.37	
6983 cordate <i>L.</i>	heart-leaved	O un 2 J jl W	1787. S co	Jac. vin. 2.193	
879. ARTE'DM L.	ARTEDIA.	(P. Artedi, a Swedish naturalist, d. 1735.)		Umbelliferat. 1.		
■ ■ ■ ■ ■	scaly	O un li jl W	Levant	1740. S co	Lam. il 193	
8H0. FE'RULA L.	GIANT FENNEL.	(Fen. to jstake; used Europe) UmbelUfers. 1.19. +26.				
6985 commfinis <i>L.</i>	common	■ ■ 4 jn.jl	Siberia	1816. D 8.1	PaLit2.Rpp.N	
698G sibirica <i>Spr.</i>	Siberian	Y				
Peucedanum sibiricum <i>W.</i>						
■ ■ ■ ■ ■	glaucous	■ ■ ■ ■ ■ P.Y	Italy	1596. D s.l		
■ ■ ■ ■ ■	thyrse-flvrcrd	■ ■ ■ ■ ■ W	Crete	1825. D co	Fl. gr. 280	
■ ■ ■ ■ ■	capillary	■ ■ ■ ■ ■ Y	Spain	1820. D co	M. h. 9.15	
■ ■ ■ ■ ■	knotted-flwd	■ ■ ■ ■ ■ Y	S. Europe	1596. D s.l	Boc. mu. 76	
J9ubon rigidius <i>L.</i>						
■ ■ ■ ■ ■	Ferulago	■ ■ ■ ■ ■	S. Europe	—	D co	Jac.au. 5
nodi flora <i>Jac. not L.</i>						
«2 strcta <i>Spr.</i>	strict	■ ■ ■ ■ ■ 2 jn.jl Y	C. G. H.	1818. S co		
JN3 longiR.Ua <i>Fis.</i>	long-leaved	■ ■ ■ ■ ■ 4 jl.au Y	Siberia	1820. D co		
WW weoldeu <i>L.</i>	Meum-like	■ ■ ■ ■ ■ 3 jl.au Y	Levant	1810. I> co		
915 orientalis <i>Tou.</i>	eastern	■ ■ ■ ■ ■ 3 jl.au Y	Levant	1759. D s.p	To. it. 3.239	
8&G ptaica IV.	Persian	■ ■ ■ ■ ■ 6 jlau Y	Persia	1782, S s.l	Bot. mag. 2096	
Assafotida B. M.						
6%7 Assafe'tida <i>L.</i>	Assafotida	■ ■ ■ ■ ■ Y	Persia	—	S r.m	
4%8 caspica <i>Bieb.</i>	Caspian	■ ■ ■ ■ ■ Y	Caucasus	1819. D co	Spr. um. 7.14	
6999 nudia <i>Spr.</i>	naked	■ ■ ■ ■ ■ Y	Siberia	1821. D co	Spr.um. 7.15	
ZJf tingiUna <i>Herm.</i>	Tangier	■ ■ ■ ■ ■ Y	Barbary	liso. S s1	Her. par. 165	
«ui obtusifolia <i>Spr.</i>	blunt-leaved	■ ■ ■ ■ ■ G	Greece	1819. D co	FL gr. 277	
-Pfucedanum obtusifolium <i>Sm.</i>						
%& pubescens <i>Pall.</i>	pubescent	■ ■ ■ ■ ■ Y	Siberia	1820. D co		
'003 villosa <i>Walt.</i>	villous	■ ■ ■ ■ ■ W	N. Amer.	1824. D co		

•881. /ASERPITIUM L.	LASERWORT. (<i>Laser, its gum, pis, pitch; smell</i>)	<i>UmbeUiferte.</i> 22.—26.
7004 latifolium L.	broad-leaved	£ un 3 jn.jl W Europe 1640. D co Jac. au. 2.146
7005 trilobum Crx.	three-lobed	£ un 5 my.jl W Levant 1640. D co
\$7006 aquilegiforme W.	Columbine-lvd	5 un. my.jl W Austria 1716. D co Jac. au. 2.147
Siicr aquilegiforme Gae.		
7007 /.ibanicum Lam.	Rosemary	^t A un 1J jn.jl p Germany 1800. Deo Lob. ic. 705
pub&cens Lag.		
7008 marginatum Kit.	margined	& A un 2 jlau P Hungary 1823. D co W. & K. 2.109
7009 Archangelicum Joe.	archangel	£ A un 4 jlau P Carniola 1800. D co Jac. ic. 1.58
7010 Aller L.	Withy	5 A un 3 my.jl W Austria 1G40. D co Jac. au. 2.145
7011 gallicum Tou.	French	[E A un 3 jn.jl Y S. Europe 1F88. D co liuk. ph. 198. 6
7012 triquetrum Ven.	three-cornered	£ A un 3 jn.jl pa.Y Constant 1816. D co Ven. eels 97
Cnchrys triquetra Spr.		
7013 angustifolium L.	narrow-leaved	& A un 2 Jn.jl Pk S. Europe 1738. D co M.h.9.19.9
7014 pruthenicum L.	Prussian	£ A un 3 jlau W Germany 1759. D co Jac. au. 2.153
7015 ^thamfinic Spr.	Athamanta-lik fr	Q un 3 jlau W Siberia 1817. S co
*7016 *ilaffi-lum w.	Silaura.lcaved	5 A un 2 jn.jl Pa.Y Italy 1791. D co Jac. au. 44
Z.ig6sticum apiftdes Lam.		
7017 peucedanoides L.	Peucedanum-lkdt	A un 2 jn.jl Pk Italy 1816. D co Jac ic 2.350
&7018 Illicidum W.	shining	5 Q un 1 :! Pa.P Switzerl. 1775. S co
Ligisticum/crulacum All.		
7019 caucasicum Bkb.	Caucasian	^ A un 1 jn.jl W Caucasus 182a S co
6'uidium caucasicum Bieb.		
7020 Oaireum W.	Olden	■ A un 2 jlau Y Levant 1820. S co Bux.c. 1.43
7021 muoldes Desf.	Suum-like	■ A un 2 jlau Y & Europe 1818. S co Desf. at 1.69
7022 pilifolium W. en.	pilose	Q un 2 jlau Pa.Y 1759. S co
7028 hinfitum Lam.	hairy	■ A un 1 :! Pa.Y Alps 1759. S co M.h.9.15
Hallcri All.		
7024 scabrum Cav.		
7025 hifspidum 2ftf.	hii	■ G un 3 jlau W Spain 1800. S co Cav. ic. 2.190
		■ G un 4 jn.jl Caucasus 1817. S co
•882. PEUCE'DANUM L. SULPHUR-WORT. (<i>Peuke, a pine, danot, burning; pungent</i>) <i>UmbelUferte.</i> ■-■		
7026 ottinile L.	officinal	^ A m 6 my.jl Y Enxland salt m. D c.l Eng. but. 1767
7027 album Hoi.	white	■ A un 6 my.jl Y N. Kuroplc 1819. S co
7028 paniculatum Lol.	uanicled	■ A un 6 my.jl Y France 1818. S co
7029 gallicum Tou.	French	■ A un 6 my.jl Y 1800. S co Gin. si. 1.41
pariHicns Dec.		
7(W aren^num Kit.	sand	■ A un 5 jn.jl Y Hungary 1816. ■ c.l W.&K.1.20
7031 ifilicum Pen.	Italian	■ A un 8 my.jl Y Italy 1810. ■ c.l Lob. ic. 781
7032 all>atre L.	Alp	■ A un 8 my.jl Y France 1739. ■ c.l
7033 ruthenicum Bkb.	Russian	■ A un 2 my.jn Y Russia 1818. ■ co
7034 tauncum Bkb.	Taunan	■ A un 2 my.jn Y Tauria 1820. ■
7030 loiKifT>limm Kit.	loiKifK-lcaved	■ A un 5 jlau Hungary 1823. D co W. & K. 251
7036 Mathiol. Spr.	Mathiolus's	■ A un 3 jn.jl Italy 1818. D co
tenui folidum Poir.		
7037 id'tensc Spr.	Isetsk	■ A un 5 jn.jl W.Y Siberia 1824. ■ co Gm. si. 1.42,43
7038 tenuiflum Thun.	fine-leaved	■ A un 8 my.jl C. G. H. 1810. ■ co
7039 rupstre Hojin.	rock	■ A un 8 my.jl Spain 1818. ■ co
7040 siblicum W.	Siberian	■ A un 8 my.jl Siberia 1804. ■ c.l W. & K. 1.60
7041 aArcum //, K.	golden	■ A un 8 my.jl Canaries 1779. ■ co
7042 Vucontibrum Spr.	vocuntians*	■ A un 8 my.jl DauphinylbOO. S co Vil. del. 2.14
//eracium pdmilum Vil.		
•883. PASTINA'CA L. PARSNEP. (<i>Pastinum, a forked planting tool; resemblance</i>) <i>UmbellUferte.</i> 10.—II.		
7043 lbcida Gou.	ghinmg.leafed	4 Q u n 1J jn.jl Y S. Europe 1771. S s.1 Jac. vin. 2. 199
7044 satWa L.	cultivated	■ C cul 4 jlau Y England ch. pi. S u.l Eng. bot 556
7045 opaca Bernh.	opaque	■ Q un 7 jlau Y Europe 1800. S co
7046 gravtolcs Bieb.	heavy-scented	■ Q un 2 jlau Y Tauria 1817. S co Hof. urn. 1. B. 6
Malabalala gravtalens Hofm.		
7047 pnpnclipifolia Bteb.	Uurnet-leaved	■ Q un 2 jlau Y Caucasus 1818. S co Buz. c 3.37
Malabalala pnpncliploides Ififm.		
7048 disscta Ven.	cuUcaved	^t Q un 6 jn.jl W Levant 181fi. S co Ven. eels 78
47049 Opopanax L.	Oi>opanax	% A m 6 jn.jl W S. Europe 1G40. D co Gou. U. 19.13.14
Ferula Opopanax Spr.		
7050 nudicaalis Spr.	naked-shmd	^t A un 1jl.au W N. Amer. 1817. D co
Sm^rmum nudicaule Ph.		
7051/wniculafcea Spr.	Fennel-like	3t A un 1jl.au W N. Amer. 1820. D co
Ferula /cenicufacea Nut.		
(7052 rfgida Spr.	rigid	■ A un 2 jlau W Virginia 1774. D s.p M.h.9.7.1
S\xm rigidius L.		
884. HERACLE* r/M L. Cow PARSNEP. (<i>Heracles, Hercules; sacred to him.</i>) <i>UmbcWfene.</i> 28.-34.		
7053 Sphondylium L.	Sphondylium	■ A myjn W Britain hcd. D co Eng. bot 939
7054 flav-Bccns W.	flavescens	■ A myjn W Austria 1789. 1) co Jac. au. 2.173
7a% clcgaua Jac.	narrow-leaved	■ A myjn W Britain w. pla. D co Jac. au. 2.174
7057 longiflbum Jac.	elegant	■ A myjn W Austria 1810. 1) co Jac. au. 2. 175
7058 sibfricum L.	long-leaved	■ A myjn W Austria 1800. D co Jac. au. 2.374
7059 7'anace L.	Siberian	■ A myjn W Siberia 1768. D co Gm. si. 1.50
True Cow Parsley		
7060 austriacum L.	Austrian	■ A flau W Siberia 1&96. D co Lob. ic. 701
7061 alpinum L.	alpine	■ A un 2 jn.jl W Austria 1752. D co Jac. au. 1.61
7062 pyrenicum Cus.	Pyrcnacan	■ A un 3 jn.jl W Switzerl. 173!. D co Bar. ic. 55
7063 iisperum Bieb.	rough	■ A un 3 jn.jl W Pyrenees 171>. S co H. her. 53
7064 pubescens Bieb.	pubescent	■ A un 4 jn.jl W Caucasus 1818. S co
7065 villsum Fis.	villous	■ A un 4 jn.jl W Caucasus 1823. S co
7066 caucasicum Stcv.	Caucasian	■ A un 4 jn.jl W Siberia 1826. S co
7067 amphiflbum Lap.	large-leaved	■ A un 6 jn.jl W Caucasus 1818. S co
7068 gummiferum W. en.	gum-bearing	■ A un 4 jn.jl W Pyrenees 1818. S co Fl.pyr.79
7069 Biueibsum Hofm.	showy	■ A un 5 jn.jl W 1819. S co W.h.b. 53,54
7070 murginktum Fis.	large-margined	■ A un 5 jn.jl W Siberia 1617. S co
7071 decipiens Fis.	deceptive	■ A un 5 jn.jl W Siberia 1817. S co
7072 giganteum Fis.	giant	■ A ox 8 jn.jl W Siberia 1820. S co
7073 lanatum Mx.	woolly	■ A ox 4 jn.jl W N. Amer. 1810. ■ co
7074 tomentiflbum Sm.	tomentose	■ A ox 4 jn.jl W Greece 1820. ■ co
7075 subvillidum W.	subvillous	■ Q or 6 jn.jl W Siberia 1817. ■ co Fl. gr. 281

7076 basviçum <i>Lam.</i>	Basvic	Q> un 8 jn.jl	W	Siberia	1818. S	co
7077 lacinatum <i>Fit.</i>	laciniated	CD un 8 jn.jl	W	Siberia	1819. S	co
7078 tauricum <i>Fis.</i>	Taurian	O) un 5 jn.jl	W	Tauria	1820. S	co
7079 vorruçusuin <i>Stev.</i>	warted	Q) un 7 jn.jl	W	Tauria	1820. S	co
7080 /igisticif Shum <i>Bieb.</i>	Lovage-leaved	Q> cu 2 jn.jl	W	Tauria	1816. S	co
•885. 7X)RDY'LIUM L. HARTWORT. (<i>Tornos. a lathe, iUo, to turn; seeds.</i>) <i>Umbc Ulfene.</i> 8.-9.						
7081 syriacum <i>Bio.</i>	Syrian	Q un *Jl	W	Syri	1597. S	co
\$7082 officinale L.	officinal	Q w 2jl	F	England	corn fi.	S co
7083 peregrinum L.	foreign	O un 2 jl	W	Levant	159a. S	co
t7084 lipulum <i>Bio.</i>	Apuhan	O un lfj	W	Italy	1739. S	co
7085 infimum L.	greatest	O w ?	W	England	corn fi.	S co
7086 lusitanicum IV. en.	Lusitanian	O un ? Lau	W	Spain	1818. S	co
17087 hfamile Dcsf.	humble	O un ? Lau	W	S. Europe	1823. S	co
7088 tiibrium <i>Seo.</i>	Siium-leaved	O un H n.jl	R	Carmola	1816. S, co	Scop. car. 194. 8
886. ASTRA'NTIAZ. MAOTERWORT. (<i>Astron. A star;</i>) <i>Umbc Ulfen*</i> . 6.						
7089 mdixim B. M.	greatest	■ ■ ■ ■ ■ ■ ■	D s.p	Bot. mag.	1553	
7090 major L.	Aelleborifbha Sal.	heturopylla W.				
7W1 pauciflora Bert.	few-flowered	5 A or 1 jn.jl	W	Alp. Eur.	1596. D p.l	Ex. bot. 2.76
7092 nrinor L.	smaller	£ A pr 1 myjn	Pk	Sicily	1820. D co	Hoc. sir. 5
7093 Jcarni6lica Jac.	Caruiolan	5 A pr 1 myjn	St	Switzerl.	1G86. D p.l	Bot cab. 93
7094 caucasica W.	Caucasian	k A or \ jnjl	Pk	Carniola	1812. 1) pi	Jac. au. 10
intermedia Bieb.				Caucasus	1818. D co	
887. ZOSVIA Hofm. ZOBIMIA. (//, N. and Z. Zosima, disting. editors of Greek classics.) <i>Vmbelliffrce.</i> 1.						
7095 ahsinthitlia Hofm.	Womwood-lvd	O w 2 jl.au	W	Persia	1816. S co	Ven. ch. 22
ileracfetim absinthiflum Ven.						
888. IV'MA Hofm. UDMIA. Taurian (<i>Rumia, the goddess who presided over suckling.</i>) <i>Utnbe/lferaf.</i> 1.						
7096 Uifrica Hofm.		£ A cu 1 Jl	W	Crimea	1819. T> co	
889. CAXJHRY'S L. CACHIRVB.						
7097 Libaritis L.	Rosemary	3 jl.au	Y	Sicily	1570. S	co Schk. nan. 1.65
7098 Morisfts AIL	Monson's	Y	S. Europe	1710. S	co M. u.m.a.i	
7099 kesclodei Bieb.	Seseli-hke	W	Tauria	1820. S	co Hof.um. 1.4	
7100 athainantoldes Bieb.	Athamanta-llk	W	Caucasus	1818. S	co	
7101 niicrocypria Bieb.	small-fruited	W	Caucasus	1810. S	co M. h. 9.1. 3	
7102 odontalKica Pall.	toothache	Y	Siberia	1819. S	co Pa.it. 3. ap. 9	
7103 crétla Lam.	Cretan	W. Y	Greece	1823. S	co An.mu.11.2	
7104 ta6nca W.	Taurian	Y	Tauria	1816. S	co	
7105 alpina Bieb.	alpine	W.Y	Tauria	1826. S	co	
7106 alita Bieb.	winged	W	Levant	1752. S	co To. it. 2.121	
Xascrpitium /erulaceum L.						
890. iGASY'LLIS Spr. AGASYLIS. (Greek name of the gum ammoniac plant) <i>UmbeWfne.</i> 1.						
7107 caucasica Spr.	Caucasian	‡ Q) un 3 jl.au	W	Caucasus	1818. S	co
Cachrys latifolia Bieb.	Slier caucasicum Spr.					
891. HIPPOMA'RATHRUM Lk. HIPPOMARATHRUM. (<i>Hippos, horse, marathron, fennel.</i>) <i>Umbellifercc.</i> 1.						
7108 siculum Lk.	Sicilian	^ A cu 3 jl.au	V.	Sicily	1640. De o	Boc. sic 18

CLASS V. ORDER S.

PENTA'NDRIA TRIGY'NIA.

892. nBU'RNUM L.	VIBURNUM.	(<i>Vieo, to tie; pliability.</i>)	Ca	ns.	28.-36.
7109 Tnu L.	Laurestinc	* or 5 mr.d	W	S. Europe	1597. S co Bot mag. 38
2 hirta	hairy	A or 5 mr.d	W	Kurope	— L co
7110 strictum	strict	» or 4 mr.d	W	S. Europe	... L co
2 virgatum	twiggy	it or 4 mr.d	W	S. Europe	... L co
7111 finolites Mutis	Tinus-hkc	t Q o r 1 ...	W	S. Amer.	1820. C P.1
7112 lucidum Mil.	ehwning-leaved	A or 6 mr.d	W	Spain	1596. L co Cl. h. 49
7113 rugifsum iVrs.	rough-Awin	* i Jor 4 ...	W	Canaries	1796. L co Bot mag. 2082
7114 villidum Swx.	villous	A O or 6 ...	W	Jamaica	1824. L co
7115 prumfilum L.	Plum-leaved	A or 8 myjn	W	N. Amer.	1731. L co Den. br. 23
7116 odoratissimum Kcr	bweetcsUscent	A L-J or 8 f	W	China	1818. L co Bot reg. 456
7117 iquamatum MM.	scaly	■ or 6 jl	W	N. Amer.	1822. L co Den. br. 24
7118:yrif&lium Lam.	Pear-leaved	A or 6 myjn	W	N. Amer.	... L co Den. br. 22
7119 Lentago L.	Lentago	A or 8 jl	W	N. Amer.	1761. L co Den br. 21
7120 fircum Pall.	naked	A or 8 myjn	Pa.Y	N. Amer.	1752. L co Bot. maa 2381
7122 obovatum Walt.	Daurian	A or 2 jnjl	W	Dahuria	1785. L co Pal. ros. 1.38
7123 cassim<ics L.	obov&te-lpaved	A or 2 myjn	W	N. Amer.	1812. L co
7124 IttvigAtum II. K.	Pumca-leaved	A or 2 myjn	W	N. Amer.	1812. L co
7125 nitidum H. K.	Cassine-like	• or 3 jnjl	W	N. Amer.	1761. L co
7126 dent'um L.	smoothed	A or 10 jl.au	W	N. Amer.	1724. L co MiL v 1.83.1
7127 ubescens Ph.	ihihiitK-leaodc	A or 2 myjn	W	N. Amer.	1758. L co Den. br. 25
7128 laitanooldes Mr.	tooth/eaved	A or 5 jnjl	W	N. Amer.	1736. L co
7129 Lant^na L.	pubescent	A or 3 jnjl	W	N. Amer.	17J6. L co
7130 tomentuum Thun.	Lantana-hkc	A or 5 injjn	W	N. Amer.	... L co Eng. bot 331
7131 molle Mr.	Wayfaring tree	A or 10 iuy.jn	W	Britain	hed. L co
Z132 acerifbium L.	tomentose	M L_ or 6	W	Japan	C p.1
2r6sea vancata Pall.	soft	Hk or 6	M	'W	N. Amer. 1812. L co
Rose	Maple-leaved	A or 4 jn.jl	W	N. Amer.	1736. L co Den. br. 118
variegated-wl	Guelder Rose	A or 10 myjn	W	Britain	moi.w. L co Eng. bot 332
oriental	Snowball tree	A or 14 myjn	W	—	L p.1
Cranberry-7Av	A or 10	R	Britain	—	L p.1
Ph.	A or 12 jl	W	Caucasus	1827. L p.1	
	A or 12 jl	W	N. Amer.	... L p.1	
	A or 12 jl	W	N. Amer.	1812. L p.1	

PENTANDMA TRIGYNIA.

893. SAMBITCUS L.	ELDEB. <i>I</i> sambuca. a musical instrument made of this tree.)	<i>Caprifoliat</i> 8.—11.
7137 <i>Twlus L.</i>	Dwarf	" Wspk-Britain wa.gr. R co ting. bot 475
7138 <i>humilis Mil.</i>	humble	2 jn.jl W.a R co
7139 <i>chinensis Lindl.</i>	Chinese ,	4 s.o. W China 1823. R co
7140 <i>nigra L.</i>	black	ec 15 my.jl Britain hed. C co Eng. bot 476
3 variegata	variegated	or 10 my.jl Britain - co
3 adrea	<i>goldcn-rdged</i>	or 10 my.jl Britain - h.l
4 pulverulc'nta	powdered	or 10 my.jl Britain - co
5 monstrosa	monkrous	or 10 my.jl Britain - co
6 viridii	<i>green-berried</i>	or 10 my.jl Britain - co
7 ilbida	<i>white-berried</i>	or 10 my.jl Britain - co
7141 <i>laciniata Retz.</i>	lacinated	SI or 8 my.jl W N. Amer. 1761. C co Schm. ar. 144
7142 <i>canadensis* L.</i>	Canadian	H" or 6 jn.au W N. Amer. 1812. C co Schm. ar. 142
7143 <i>pubens Mx.</i>	downy	SI or 12 my.jn G.T S. Europe 1596. C co Jac. icl. 59
7144 <i>raceinisa L.</i>	racemose	SI or 12 my.jn P S. Europe 159ft C co
2 purpurea	Spiric	SI or 12 my.jn Y.G & Europe 1596. C co
3 flavescens	*	
• 894. <i>RHV'S Tou.</i>	avescent	
7145 <i>Cotinus l.</i>	SUMACH. (<i>Shudd, red, Celt.; fruit</i>) <i>Terelndtuce*</i> . 74.-88.	
7146 <i>Aetopium L.</i>	Wild Olive	S Europe 1616. L Jec. au. 3. 210
7147 <i>Coriana L.</i>	Metopium	W. Indies 18tf. C p.1 Br.jam. 13.3
7148 <i>typhlna l.</i>	Coraria-vtf	* S. Europe 156. L co Den. br. 136
2 Irutescens	fever	5 N. Amer. 1629. S co Den. br. 17,18
7149 <i>javanica IV.</i>	frutscsent	S S co
7150 <i>scmialita Mur.</i>	Java	*
7151 <i>glabra L.</i>	half-winged	Java" 1799. S p.1 Den. br. 15
7152 <i>elcgyns IV.</i>	smooth	Macao 1780. L p.1 Mur. 6.3
7153 <i>vmdifltoa Poir.</i>	elegant	N. Amer. 1726. L p. Di el. 243.3*4
7154 <i>ptimila Afx.</i>	green-flowered	* N. Amer. 1726. p. Den.br. 16
7155 <i>whinoldes W.</i>	<i>dwarfpoisonous**</i>	<i>dwarfpoisonous**</i>
7156 <i>chinlndis Mi.</i>	Schinus-like	S. Europe 1806. p.
7157 <i>pauciflora L.</i>	Chinese.	Brazil 1824. p.
7158 <i>afata Thun.</i>	few-flowered	China 1800. p.
7159 <i>copalina l.</i>	winged	C. G. H. ... p.
7160 <i>trijuga Poir.</i>	gum copal	C. G. H. 1824. p.
7161 <i>obllqua 7%Kn.</i>	thrc-i>air-/Mf	N. Amer. 1688. p. Jac. sc. 3.341
7162 <i>Commersiaitf Poir.</i>	oblique	il Brazil 1825. p.
7163 <i>auminata Dec.</i>	Commerson's	H C. G. H. 1825. p.
7164 <i>succedanca L.</i>	acuminate	Jl Brazil 1810. p.
7165 <i>vernix l.</i>	succedancous	M Nepal 1820. p.
venenata Dec.	varnish	*
7168 <i>Mcku-Amida Rox.</i>	Bucku-Amela	It China p.1 Kaem. 795
7167 <i>yraxinifltha D. #<m</i>	• "jor	
7168 <i>obscura Iitcb.</i>	Ash-leaved	G.Y Nepal 1823. S co
7169 <i>vermcifera Dec.</i>	obscure	
Juglandifolia <i>Wai.</i>	varmsh-bearinga L_J or	Iberia 1820. L p.1
7170 <i>Toxicodendron L.</i>	Poison-Oak	G.Y Nepal 1823. S co
7171 <i>radicans l.</i>	rooting	N. Amer. 1640. L co
2 volubilis	twinning	N. Amer. ... L co Bot. mag. 1806
3 microcarpa	*-small-fruited	N. Amer. - co DuB. no. 48
7172 <i>glaAca Desf.</i>	glaucous	N. Amer. co Di.el.375
7173 <i>Imeatiflha Or.</i>	hneatcd-leaved	C. G. H. 1821. p.
7174 <i>Thunbergiana ILSeS.</i>	Thunberg's	C (J. H. Cuba 1818. p.
7175 <i>Ibcida L.</i>	shining-taived	C. G. H. 1697. Com. h. 1.93
7176 <i>Cavanillesii Dec.</i>	Cavanilles's	Mexico 1816. Cav. ic. 2.135
liicina Cav.		
7177 <i>BurmAnni Dec.</i>	Bunnann't	N. Amer. 1897. C p.1 Bur. af. 91.2
liicina H. K.		
7178 <i>nervtoa Poir.</i>	ncrovae	C. G. H. 1800. p.
7179 <i>concfna Burc.</i>	neat	C. G. H. 1820. p.
7180 <i>villbaa L.</i>	hairy	C. G. H. 1714. Pluk. al. 219. 8
7181 <i>pyroldes Burc.</i>	Pear-like	C. G. H. 1816. p.
<i>In</i> pubocens L.	pubescent	C. G. H. 1800. p.
7183 <i>atomaria Jac.</i>	atomed	C. G. H. 1800. p.
7184 <i>Albida Schous.</i>	white-famed	C. G. H. 1800. p. Jac. sc. 343
7185 <i>excisa Thun.</i>	<i>cut-leaved</i>	Mogadore 1816. p.
7186 <i>macronota Thun.</i>	macronate	C. G. H. 1816. p.
7187 <i>elliptica Thun.</i>	elliptic	C. G. H. 1824. p.
7188 <i>pendufina Jac.</i>	penrulous	C. G. H. 1818. p.
elongate Jac.	<i>phahed-leaved</i>	C. G. H. 1758. C p.1
7190 <i>viminallis Vahl</i>	green-twiggly	G.Y C. G. H. 1774. p. Jacsc & 344
7191 <i>l&ncea L.</i>	<i>l&nce-leaved</i>	C. G. H. 1810. p.
7192 <i>fragns L.</i>	awceUscented	C. G. H. 1824. p.
7193 <i>rltida Mil.</i>	rigid	C. G. H. 1700. p.
7194 <i>cihata Licit.</i>	ciliated	C. G. H. 1816. p.
7195 <i>trid&ctylis Burc.</i>	three-fingered	C. G. H. 1816. p.
7196 <i>angustifolia L.</i>	narrow-leaved	C. G. H. 1714. p. Pluk. al. 219. 6
7197 <i>rosmanitolia Vahl</i>	Rosemary-lvd	C. G. H. 1800. p. Bur. af 91. 1
7198 <i>sernefTilia Burc.</i>	saw-leaved	C. G. H. 1816. p.
7199 <i>iintulfia Jac.</i>	w&eA4eaved	C. G. H. 1816. p. Jac. K. 3.346
<i>thèsans Hort</i>		
7200 <i>dioica Brou.</i>	dificious	*LJor Mogadore 1825. p.
7201 <i>Oxyacantha Schous.</i>	» or	Barbary 1823. p.
7202 <i>Oxyacanthoides Dün.</i>	Hawthorn-like	Burbary 184. p.
7203 <i>lobata Poir.</i>	SI or	Teneritte 1800. p.
7204 <i>sinuate Thun.</i>	<i>lobei</i> Leaved	C. G. H. 1816. p.
7205 <i>cuneiflia Desf.</i>	<i>Binuate-leaved</i>	G.Y C. G. H. 1816. p.
7206 <i>Stridentitez..</i>	wedge-leaved	C. G. H. 1816. sp.
7207 <i>crefate Thun.</i>	three-toothed	C. G. H. 1816. p.
7208 <i>micrantha Thun.</i>	crnated	C. G. H. 1818. p.
7209 <i>disscta Thun.</i>	small-flowered	C. G. H. 1820. p.
7210 <i>tomentos* L.</i>	dissected	C. G. H. 1691. p. Com. h. 1.92
7811 <i>inrisa L.</i>	<i>wooWyJcaved</i>	G.Y C. U. H. 1789. p.1
	<i>cut-leaved</i>	2

7818 dentate <i>Thun.</i>	toothed	C C U 8 ...	C G. H. 1798.	C p.i
spicata <i>Thun.</i>	piked	C J or 3 ...	C 6. H. 1819.	C p.l
7213 pentaphylla <i>Deft.</i>	fiv-leaved	C J or 4 ...	Barbary 1816.	C p.l Deaf, at 1.77
7214 zizypmna <i>Tinco</i>	Ziziphus-hke	C J or 4 ...	Sicily 1800.	C p.l
7216 aroinatica <i>H. K.</i>	aromatic	SB or 8 my	N. Amer. 1759.	L p.l An. mu. 5.30
7216 suaveolent <i>H. K.</i>	sweet-smelling	dh or 6 my	N. Amer. ...	L p.l
7217 heterophylla <i>Dec.</i>	various-leaved	A L I or 6	C. G. H. 1800.	C p.i
7218 digitate <i>L.</i>	digitate	H J or 10 ...	C. G. H. 1820.	C p.i
895 CASSINE L. CASSINE (Its name amongst the Indiana of Florida.) <i>Cddtritiae. 7.</i>				
7219 Maurocenia L. Maurocenia Hotcher	tt. L or 5	W C. G. H. 1690.	C sp.1 Di. el. 121.147	
7220 capensis L. Cape <i>Phillyrea</i>	tt. L or 1	jl.au W C. G. H. 1629.	C sp.1 Bur. af. 85	
7221 Colpodn Thun.	COMKKM tree	* i. J or 6 ...	W C. G. H. 1791.	C sp.1 Bur. ac 86
7222 ethiApica <i>Thun.</i>	Ethiopian	* i. J or 2	jl.au W C. G. H. -	C sp.1
7223 barbara L. barbarous	m J or 8	jl.au W C. G. H. 1818.	C sp.1	
7224 excelsa Rox.	tafl	* i. J or 18	Nepal 1820.	C sp.1
7225 oppositi folia MU.	opposite-leaved	H J or 5	Jamaica 1824.	C sp.1
896. SPATHE^LIA L. SPATHIELIA. {Spatha, a palm tree; similarity.) <i>Tercbinthcea. 1.—8.</i>				
7226 simplex L.	simple	± • tm 40 ...	R Jamaica 1778.	S sp Bot. reg. 670
897. STAPHYLE^AZ. BLADDER-NUT. {Staphyle, a bunch; disposition of fructification.) <i>Ccltufr'itut. 3.-5</i>				
7227 pinnata L.	pinnate-feawd	Sk or 6 ap.jn	England hed. L co Eng. bot. 1560	
7228 trifolia L.	three-leaved	A or my.jn	W N. Amer. 1640. L co Schm. ar. 80	
7229 occidentalis Sun.	western	IU2OT	W Jamaica 1824. C p.l Sl. Jam. 2.220.1	
898. STACKHOU^Sii It Br. STACKHOUSIA. (Af. Stackhouse, a British botanist.) <i>StackhmUUe. 1.—2.</i>				
7830 liliifolia Cun.	Flax-leaved	i. Jor 2	N. Holl. 1821	C s.l.p
899. 7W/MARIXJL TAMARISK. (Tamarisci, a people of the Pyrenees; habitat) <i>Tamaraseinete. 6.—18.</i>				
72J1 gallica L.	French	or 12 my.o	F England so. co. C s.l Eng. bot. 1318	
7232 articulate Vahl	jointed	D or 30	Pk E. Indies -	C l.p Vah.sy.2.38
7233 dioica Rox.	dioecious	6	- K Indies 1823. C p.l	
7234 Indica W.	Indian	O 6	F E. Indies ... C fp	
7235 germanica L.	German	or 8 jn.s	Pk Germany 1582. C ms Mil. ic. 862.8	
7236 dav&nca L.	Dahurian	or 6	Pk Dahuna 1818. C l.p	
900. TU'RNEIU L. TURNERA. <i>JJV. Turner, M.D., died 1568.) Turntrihcea. 11.—84</i>				
7237 tiliifolia L.	Elm-leaved	H 3 jn.s	Y Jamaica 1733. C p.i H.cl. 10	
2 angusti folia B. M.	narrow-leaved	3 ap.s	Pa.Y Jamaica 1733. C s.p Bot mag. 881	
7238 elegans Otto	elegant	3	Pa.Y Brazil 1821. C s.p	
7239 cuneiformis J.	wedge-form	JnjV	Y S. Amer. 18.Jl. S. s.p	
7240 /riomflora Sims	Trionium-flwd	year	Pa.Y Brazil 1812. C s.p Bot. mag. 8106	
7241 brasiliensis W.	Brazilian	Y	Brazil 1810. S fl.p	
72(2) Pumil&a L.	Dwarf	Y	Jamaica 1796. S s.l Sl. jam. 1.187.6	
7243 cistoides L. **	Cistus-like	Y	America 1774. S s.l PI. ic. 150. 1	
7244 rupestris Auh.	rock	Y	Guiana 1824. S Bp Aub. guil. 113.1	
7245 guianensis Aub.	Guiana	Y	Guiana 182. S s.p Aub. gu. 1.114	
7246 hirta IV.	hairy	Y	Brazil 1818. S sp	
7247 racemosa Joe.	racemose	Y	Siberia? 1789. S 81 Jac.vin.3.94	
<i>Piriqua racemosa Suit.</i>				
901. DRY'PSISL. DRYPSIS. (Dryptot to tear; armed leaves.) <i>Caryoptijllct. 1.</i>				
7248 spinosa L.	prickly	pr jn.jl	Pa.B Italy 1775. S s.l Bot mag. 8816	
902. fLSrN CHICKWEED. (Algos, a shady place; habitation.) <i>Caryoph\$le*. 5.-6.</i>				
7249 media L.	mediate	Ow 1 jls	W Britain fields. S co Eng. bot. 537	
7250 Segetalis L.	Sword-grass	O cu 1 jl.9	W France 1805. S co	
7251 moJugheea Lag.	Mollugo-like	O cu * jls	W Spain ISifi. S co	
7252 puWscens Lk.	pubescent	O cu i jl.s	1810. S co	
7253 mucronata L.	mucronate	O cu 1 jn.jl	W & Europe 1777. S co Fl.gr. 293	
*903. DRYMA^RIA W. DRYMARIA. (Drymos, a forest; habitation.) <i>Caryoph\$li*e. 1. — 5.</i>				
7254 cordata IV.	cordate	3pr ijn.au	W Jamaica 1800. D co Lam. it 51.2	
904. TELFSVHIUM U ORPINS. (Sup. to heal old ulcers, like that of <i>Telcphus</i> .) <i>Portuldcat. 1.—2.</i>				
7855 Imperau L.	Imperato's	-* A w ijn.au	W S. Europe 1658. D s.l Lam.il. 213	
905. COR RIG POL A L. STRAPWORT. (Dim. in. of <i>corrugia</i> , a leather thong.) <i>PorhditceG. 3.</i>				
725G Utteralis L.	6hore	Jk O or i jl au	W England so. co. S s Eng. bot. 668	
7257 capensis Thun.	Cape	J; O pr i jl.au	W C. (i. H. 1819. S co	
7258 tQlevbiiblia Pou.	Orpine-leaved	Jt A cu e jl.au	W S. Europe 1822. S	
*906. PHARKAXEUM W. PHARNACBUM. (Pharnaccs, king of Pontus, first used it.) <i>CaryopkSUea.W.-^.</i>				
7859 Acclidifolium Poir.	Daisy-leaved	O cu i jl.au	W W. Indies 1823. S s.l PL am. 21.1	
7859 ~ MoMfgo Adhdif&lia Dec.				
7260 cordiflum L.	heart-leaved	O f u I jn.au	W C.G.II. 1823. 6.1 Jac. sc. 349	
7261 dichotomum Thun.	forked	i. cu i jl.s	W C.G.H. 1783. 8.1	
275* K ^{lom} cr. lrum Thun.	giomcrateil	iD cu i jl.au	W C.G. H. 1817. s.p	Pluk. m. 331.4
7263 Cervikna L.	Cerviana	O cu I jn.	W Russia 1771. co	Gin. si. a 2a 2
475* spergulaWo* Poir.	Spergula-like	O cu i jn.au	E. Indies 1819. Ldm.il 214.8	
7265 lineare Thun.	hncar-leaved	tt. Jcuy i my.o	W C. G. H. 1795. 8.1 Bot rep. 326	
7W6 umheUdtum Forsk.	tt. Ocu i jl.s	W Egypt 1820. 8.1		
7267 iVolh.go L.	Mollugo	tt. Ocu i jl.au	W E. Indies 1752. 8.1 Bur. in. 5.4	
7268 incanum L.	hoary	tt-uJpr 1 my.o	W C. G. H. 1782. s.1 Bot mag. 1883	
7269 pruinaBum Haw.	frosty	<-l-Jpr i jn.au	W C. G. H. 1817. s.1	
JSPI P°?ULACA^RIA Joe. PURSLANE TREK. (A Por/Waco-llke plant) <i>Poriuttce. 1.</i>				
7270 afra Jac.	African	a. j c u 3 ...	P Africa 1/32. C r.m Jac. c. 1.88	
908. BASELLA L. MALAHAR NIGHTSHADE. (Its Malabar name.) <i>Chenapodett. 8.—10.</i>				
red				
black	i. OD cul 8 jls	Pk E. Indies 1731. S r.m		
white	fl_Ocul 3 jls	W China 1822. S r.m		
shining	fl_Ocul 8 jln	W E. Indies 1688. S r.m Pluk. aI 63.1		
heart-leaved	fl_Ocul 6 jln	W E. Indies 1802. S r.m		
branchy	fl_Ocul 6 jln	Pa.P R Indies 1802. S r.m		
	 S r.m		

- 7277 tuberata //, & B. tuberous & Elor 6 jl.n S. Amer. 1824. S r.m
 margined %_ Oleu 4 jn.au — Mexico 18J4. S r.m

909. PORANTHE'RA Rud. PORANTHEBA. (*Poros, a pore, anthcra, an anther.*) 1.
 7S79 ericiflta < Af. <i. Heath-leaved a. L. cu ijn.au W N. Holl. 1824. C s.1 Lin.tr. 10. ©. 2

CLASS V. ORDER 4.

PENTA'NDRIA TETRAGY'NIA.

910. PARNA'SSI A L. GRABS OP PARNASSUS. (*Mt. Parnassus*: elegance, and sup. liab.) *Droserheete.* 4.-7.

- | | | | | | | | | |
|-------------------------------------|-----------------------|--------|----------------|---|----------|-------|-------|----------------|
| 7280 <i>halfiiflirB</i> <i>L.</i> | <i>mar*!*</i> | k A dr | i jLau | W | Britain | bogs. | D bog | Eng. not. 82 |
| 7281 <i>parviflerra</i> <i>Dec.</i> | <i>small-flowered</i> | ^ A pr | <i>I</i> my.jl | W | N. Amer. | 1820. | I) | bog |
| 7282 <i>caroliniana</i> <i>Mx.</i> | <i>Carolina</i> | £ A P* | <i>i</i> my.jn | W | N. Amer. | 1802. | D bog | Bot. mag. 1459 |
| 7283 <i>usaritolia</i> <i>Ven.</i> | <i>Asarura-leaved</i> | ^ A pr | <i>ijl.au</i> | W | N. Amer. | 1812. | D bog | Ven. mal. 90 |

CLASS V. ORDER 5*

PENTA'NDRIA PENTAGY'NIA.

- | | | | |
|---|------------------|--|--|
| •911. ARABIA L. | ARAUA. | (<i>Ara</i> , annoyance; spinel to travellers.) | <i>AraUlcce.</i> 16.—34. |
| 7284 spinosa L. | thorny | or | W Virginia 1688. R p.l Dcn.br. 116 |
| 7285 hispida Mx. | bristly | or | W N. Amer. 1799. R p.l Bot mag. 1085 |
| 7280 Multilobed Bergia R.&S. Muhlenberg's t | naked-stalked | or | W N. Amer. ... R i.p. |
| 7289 frágans G. Don | sweet-scented | | |
| Panax fráRrans Rox. | | | |
| 7290 aculeátum G. Don | prickly | IC2OT | — W Nepal 1820. C a.p. |
| /federa aculeata Ham. | | | |
| 7291 umbaculifera Aor. | shady | | W E. Indies 1818.. C 8.1 |
| 7292 pentaphylla Thun. | five-leaved | | Japan 1810. C 8.1 |
| 7293 ferruginea // . * B. | rusty | | Trinidad 1826. C 8.1 |
| 7294 digitata Rox. | digitate | | W E. Indies 1817. C 8.1 |
| 7295 imcans // ft B. | glittering | | Trinidad 1828. C s.p. |
| 7296 pubescens Ztec. | pubescent | | W. Indies 1818. C s.p. H.mon.45 |
| 7297 arborea L. | tree | | G Jamaica 182a C r.m Jac.BC. 51 |
| 7/dera arborea Sti*. | | | |
| 7298 capitata Sun. | capitate | | G W. Indies — C r.m Bot cab. 946 |
| /federa capitata Sun. | | | |
| 7299 cochlate Lam. | elated | elated | W E. Indies 182a C s.p Ru. am. 4.31 |
| 912. SCIODAPHYLLUM Br. | | | |
| 7300 Bréwini Spr. | Brown's | SCIODAPHYLMJM. (<i>Skiodes</i> , shady, <i>phyllon</i> , leaf.) | <i>Aruliccar.</i> 7.—8. |
| Aralia Sciodaphyllum Sun. | | TODPH | Jamaica 179a C sp Br.jam. 19.1 |
| 7301 digitatum G. Dan | finger-leaved | f • or 20 | — W |
| Actinophyllum digitatum Wai | | * | G E. Indies 1820. C s.l |
| 7302 pedicellatum Poir. | pedicellate | • or 10 | — P Peru ... C s.l Fl per. 308 |
| 7303 cénicum Poir. | conical | I Dor 10 | — Pa.R Peru ... C s.l Fl. per. 509 |
| Actinophyllum cénicum R. & P. | | JLChor 10 | — Pa.R Peru ... C s.l Fl. per. 509 |
| 7304 acuminatum Poir. | acuminate | O or 10 | — Y Peru ... C 8.1 Fl. per. 310 |
| 7305 pentandrum Poir. | pentandrous | • or 18 | — Pa.R Peru 1820. C B.1 Fl. per SI 1 |
| 7906 anomalous G. Don | anomalous | • or 50 | — W.o Trinidad 1817. C si |
| CaroHea insignis Hort. | | | |
| •913. RO'CHRISTDec. | HOCIRIA. | (<i>M. De la Roche</i> , a botanical author.) | <i>CrassuRice*</i> . 2. |
| 7307 falcate Dec. | tickle-leaved | si L J or 3 | jns. R C. G. H. 1795. C 8.1 Bot. mag. 2035 |
| Crassula falcata L. | | | |
| 7308 perfoliata Haw. | perfoliate | • i_J or 4 | jl.au S C. G. H. 1700. C 8.1 Dec. pi. 13 |
| 2 SO* Haw. | white-flowering | • i_J or 4 | jl.au W C. G. H. 1800. C 8.1 |
| 914. KALOSANTHES Haw. | KALOSANTHES | (<i>Kalos</i> , beautiful, <i>anthos</i> , a flower.) | <i>Crassul&cae.</i> 9. |
| 7309 coccinea Haw. | scarlet | tt or 1 | jn.au S C. G. H. 1710. C s.l Bot mag. 495 |
| Crassula coccinea L. | | | |
| 2 flor albo Brad. | white-flowering | tt i_J or ! | 1 jn.au W C. G. H. 1811. C 8.1 Brad. sue. 50 |
| 7310 mfredia Haw. | mediate | n. i_J or 1 | Jnl R C. G. H. 1810. C 8.1 |
| 7311 versicolor Haw. | changeable | tt i_J or 2 | mr.s R C. G. H. 1817. C 8.1 Bot reg. 320 |
| 7312 bicolor Haw. | two-colored | tt i_J or 1 | Jnl YB C. G. H. 1800. C 8.1 |
| 7313 odoratissima Haw. | sweetest-scented | tt i_J or 1 | jrjl Pk C. G. H. 1793. C 8.1 Bot rep. 26 |
| 7314 ffava Haw. | yellow | tt i_J or 1 | aus Y C. G. H. 1802. C 8.1 Pluk.aL314. 2 |
| 7315 cymbia Haw. | cymose | tt i_J or 4 | au R C. G. H. 1800. C 8.1 |
| 7316 biconvexa Haw. | double-convex | tt i_J or " | " C. G. H. 1823. C 8.1 |
| 7317>asminea Haw. | Jasmine-like | tt i_J or \$ | ap.my W C. G. H. 1815. C 8.1 Bot mag. '2078 |
| 915. CRA'SSULA L. | CRASSULA. | (Dim. of <i>crassus</i> , thick; fleshy leaves and stems.) | <i>CrassvlactB.</i> 52.—83. |
| 7318 ramosa JKK. | branched | 2 Jl.au | Pk C. G. H. 1714. C s.l |
| 7319 tetragona /* | four-angled | 2 au | W C. G. H. 1711. C 8.1 Dec. pi. 19 |
| 7320 acutifolia Lam. | acute-leaved | tt or | W Greece? 1795. C 8.1 DecpLS |
| 7321 biplana Haw. | biplanate | tt lor | W C. G. H. 1823. C B.1 |
| 7322 rotundifolia/fow. | round-leaved | tt or | aus W C. G. H. 1800. C 8.1 |
| 7323 bullulata Haw. | small-studded | tt or | au.B Y C. G. H. 1800. C 8.1 Martcen. ic. |
| 7324 scabrella Haw. | small-scabrous | tt or | W C. G. H. 1810. C 8.1 |
| 7325 filicifolia Haw. | thread-stmd | tt or | VV' C. G. H. 1820. C 8.1 |
| 7326 revolvens Haw. | revolving | tt or | aus W C. G. H. 182a C 8.1 |
| 7327 arbor&cens W. | arborescent | tt J or 3 | my.jn Pk C. G. H. 1739. C s.l Bot mag. 384 |
| 7328 obliqua H. K. | oblique-teavif | BH J or 4 | ap.my R C. G. H. 175a C s.l Dec. ic.79 |
| 7329 lactea H. K. | milk-colored | tt or f.s.o | W C. G. H. 1774. C s.l Bot mag. 1771 |
| 7930 albiflora Dec. | white-flowered | tt or | C. G. H. 1800. C 8.1 Bot mag. 23P1 |

7331	ciliata L.	ciliated	tt.i	lor	i jl.au	Y	C. G. H.	1732.	C s.1	Dec. pi. 7	
2	média Haw.	mediate	tt.i	lor	I jlau	Y	C. G. H.	1818.	C 8.1		
3	minor Haw.	smaller	tt.	or	ijjlau	Y	C. G. H.	1818.	C 8.1	Jac. sc. 433	
7-8	dejecta Jae.	dejected	tt.	or	1 jl.au	W	C. G. H.	J820.	C 8.1		
7332	conclina Haw.	neat	tt.	or	-	-	C. G. H.	1818.	C 8.1		
7-8&1	lingularifolia Haw.	latchet-leaved	tt.	or	-	-	C. G. H.	1818.	C 8.1		
7335	obtusu Haw.	obtuse	tt.	or	i aus	W	C. G. H.	1823.	C 8.1		
2	major Haw.	two-bracted	LJ	or	iau.s	W	C. G. H.	1823.	C 8.1		
7337	squamulif^a //aw.	larger	tt.	or	S	W	C. G. H.	1817.	C s.1		
7338	coccinea*/aw.	squamulose	L	or	Pa.R	C	C. G. H.	1823.	C s.1		
7339	telephio idis Haw.	small scarlet	L	or	t jn.jl	P Y	C. G. H.	1818.	C 6.1	Di. ek. 99.117	
7340	si ilbra /.	Telephium-likott.	L	or	1 iau	W	C. G. H.	174.	C 8.1		
7341	biconvexa Haw.	rough-leaved	LJ	or	1 jnjl	W	C. G. H.	1800.	C 8.1		
7342	ramuliflora Lk.	double-convex	LJ	or	1 jlau	W	C. G. H.	1822.	C s.1		
7343	corymbulif. 1 Lk.	branchlet-flwd	LJ	or	1 jnjl	W	C. G. H.	1822.	C 8.1	Bur. af. 9	
7344	columbaris /.	corymbulose	L	or	I*jnif	W	C. G. H.	1789.	C s.1		
7345	imbricata // K.	columnar	L	or	1 apau	W	C. G. H.	1820.	C 8.1		
7346	mcoides Haw.	imbricated	L	or	2 jlau	P.v	C. G. H.	1732.	C 8.1	Scop. d. 3.6	
7347	rosularia Haw.	Heath-like	tt.	or	1 jns	Pk	C. G. H.	1760.	C s.1		
7348	perfoliata Sco.	small-rosy	tt.	or	* Jli	W	C. G. H.	1819.	C 8.1		
7349	punctita L.	threaded	tt.	or	i my.au	Pk	C. G. H.	1783.	C ill.		
7350	marginalia // JK	dotted	tt.	or	1 in	W	C. G. H.	1739.	C 8.1		
7351	pulicaria /.	marginal	tt.	or	1 myjn	Pk	C. G. H.	1774.	C s.1		
7-8>	Kpathulara W.	pellucid	tt.	or	1 au	W	C. G. H.	1774.	C 8.1	Di. el. 100, 119	
7J53	cordisca H. K.	spathulated	tt.	or	1 myjn	Pk	C. G. H.	1774.	C 8.1	Dec. pi. 49	
7354	Cetylomonis L.	ho, tr-lt'av'd	L	or	1 myjn	Pk	C. G. H.	1774.	C 8.1	Bot cab. 539	
7355	orbicularis* L.	Cotyledon-!/*/ \$	A	or	1 my	W	C. G. H.	1774.	C 8.1		
7356	retroflexa L.	orbicular	jg	A	1 my	Pk	C. G. H.	1788.	C 6.1		
7-17	lineolata H. K.	bunt bick	O	or	1 in	Y	C. G. H.	1774.	C 8.1		
7358	contauroides H. K.	small-lined	f	£2)	1 myjn	Pk	C. G. H.	1774.	C 8.1	Bot mag. 17-ii	
7359	glomerata L.	Centauria-like	£	Ol	1 myjn	Pk	C. G. H.	1774.	C 8.1	Her. lug. 553	
7360	Magnifi Dec.	glomerated	O	pr	1 myjn	Pk	C. G. H.	1774.	C 8.1		
7361	subulata L.	Magnol's	O	o	1 myjn	W	C. G. H.	1774.	C 8.1		
7362	pulchilla // Jfc	subulate	O	pr	1 myjn	W	C. G. H.	1800.	C 8.1		
7363	glabra Haw.	pretty	O	or	R**	W	C. G. H.	1810.	C 8.1	Her. lug. 552	
7364	glomeretta 0 H. K.	smooth cluster	O	P	1 my	W	C. G. H.	1774.	C 8.1		
7365	ribbens L.	annual red	O	pr	I myjn	Pk	Italv	1749.	C 8.1		
7366	verticillaris /.	VfhorUflowered	O	pr	* Jli	Pk	S. Kuroj	1748.	C 6.1		
7-8	expand // K.	expanded	O	pr	Jinjl	W	C. G. II.	1774.	C 8.1		
7-8>	spana H. K.	s	;	21	pr	% jnjl	Pk	C. G. H.	1774.	C 8.1	
7JS	difflisa // K.	diffuse	O	pr	Opr	W	C. G. H.	1774.	C 8.1		
7-8j	moschata Forst.	mubby	O	pr	* my.n	W	N. S. W.	1794.	C 8.1		
*916.	TURGCysIA Haw.	TURGOSIA. (Err. for Pyrgosia: pyrgos, tower; clusters of flowers.) <i>Crassuldeete.</i> 7.									
7370	linguefolia Haw.	tongue-leaved	£	iAI or	1 au	W	C. G. H.	1803.	C s.1		
7-8	linguefolia Haw.										
7-8	tomentosa W.										
7-8?	pertusa //w.										
7-8>	turrata Haw.										
7-8	2 rusea Haw.										
7-8	obovata Haw.										
7-8	aloidea Haw.										
7J76	capitella Haw.										
917.	GLOBITLEA Haw.	GLOBULEA. (<i>Globulus</i> , a globule; stems.) <i>Crassullcea.</i> 16. — 29.									
7-8	cultivata Haw.	sharp-leaved	«Ll_Jor		1 jlau	W	C. G. H.	1732.	C s.1	Bot. mag. 1940	
7-8	Crassula cultMa L.										
7-8	atropurpurea Haw.	dark-purple	tt.	J or	\ au	W	C. G. H.	1823.	C &1		
7-8	radicans Haw.	tf#r#/t-oolingtt.	tt.	or	1 jno	W	C. G. H.	1823.	C 8.1	Com. pr. 2	
7-8	ngua Haw.	tongue	tt.	or	* jnjl	W	C. G. H.	1823.	C 8.1		
7-8	« a »aw.	little-tongue	a.	\ or	% jnjl	W	C. G. H.	1823.	C 8.1		
7-8	capitata Ma Haw.	capitate	£	IAj or	I	W	C. G. H.	1819.	C 8.1		
7-8	obvalvata Haw.	guarded-leaved	a.	j or	i jlau	W	C. G. H.	J795.	C 8.1	Dec. pi. 61	
7-8***	canescens Hiw.	canescent	a.	IAj or	4 jlau	W	C. G. H.	1800.	C si		
7-8>	pumiculata Haw.	panirulate	jg	IAj or	i jnjl	W	C. G. H.	1823.	C 8.1		
7380	impressa Haw.	impressed	tt.	or	\ aus	W	C. G. H.	1820.	C si		
2	2 minor Haw.	smaller	tt.	or	i alor	W	C. G. H.	1820.	C s.1		
7387	sulcata Haw.	sulcate	tt.	or	ij!18	G''	C. G. H.	1818.	C 8.1	Dec. pi. 55	
7-88	nudicaulis Haw.	nakod.stemmed	tt.	or	i my.8	W	C. G. H.	1732.	C s.1		
7-8>	hsipida Haw.	hispida fiat-lvd	tt.	i Jor	4 aus	W	C. G. H.	1823.	C s.1	Dec. pi. 133	
/	muKembryanthodes				1 aus	W	C. G. H.	1820.	C 8.1		
7-8	altior Haw.				4 aus	W	C. G. H.	1820.	C 8.1		
7-8	mollis Haw.	soft	tt.	or	1 au	W	C. G. H.	1774.	C 8.1		
7-8	subpicana Haw.	huaryish	tt.	or	4 au	W	C. G. H.	1823.	C s.1		
7-8	erecta Haw.	erect	tt.	or	1 au	W	C. G. H.	1823.	C 8.1		
7n6	URTGVGYNE Haw.	GURTOQINNE. (<i>Kurtos</i> , curved, jiy/u?, a style.) <i>Crassulacea.</i> S.									
739J	undata Haw.	waved	tt-Ljor	*	a u	W	C. G. H.	1818.	C s.1		
7J10J	undulata Haw.	small-waved	tt.	J or	au	W C. G. H.	1797.	C s.1			
7-8	VAU A'NTHES Haw.	VAUANTiEa (<i>Fau</i> , the letter V, <i>anthos</i> , a flower.) <i>Crassulacea.</i> 1.									
7-8	chloriflora Haw.	Chlora-flowered	njor	fjl	Y	C. G. H.	1774.	S s.1	Dec. pi. 67		
7-8	Crassula dichotoma L.										
7-9	GISETKA4 L.	GISEKIA. (<i>P. D. Gische</i> , a Danish botanist.) <i>Phytolococe*</i> . *									
73J6	phsLrnaceoldes L.	Pharnaceum-like	(P. O cu	1 jn	Pa.G	E. Indies	1783.	Rox. cor. 2. 18			
7-9	zrNUM Bauh.	FLAX.	(TMn,	a thread,	Cell.)		CaryopIrlleec.	43. — 61.			
7-8	afireum W. \$ K.	French	O or	I jlau	Y	France	177.	S co	Ger. gal. 15.1		
7-8		golden	O ox	^jnjl	Y	Hungary	1820.	S p.l	W. & K. 2.177		
7-8		yellowish	O or	I jnjl	Y	Tauria	1820.	S p.l	Bux c 5.59		
7-8		knotted-flwd	O or	ijjlau	W	Italy	1759.	D s.1	M. h.f. 5.1U1		
7-8		strict	O or	1 myjl	Y	S. Europe	1759.	S pl			
7-8		stiff-leawd	O or	1 jl	Pa.Y	Missouri	1807.	S co			
7-8		Virginian	Q or	1 jl	Y	N. Amer. IK07	S GO				

7404 dahūricura/J.S.S.	Dahurian	1 jnjl Y	Dahurla 1816. D pi	Jac. vin. 2.154
7405 marttiinum L.	sea	2 jl.au Y	S. Europe 1890. D co	Bot. mag. 234
7406 arbucum L.	tree	2 my.au Y	Candia 1788. C pl	
7407 campanulatum L.	campanulated	1 jnjl Y	Euroojic 1705. D pl	
7408 tauricum W. en.	Taurian	ljin.jl Y	Tauria 1818. C pl	
7409 capitatum R.%8.	capitate	1 jnjl Y	Austria 1816. D pl	
7410 flicvum L.	yellow	\$ jnjau Y	Austria 1793. D pl	Bot. mag. 312
7411 trigynum Aw.	three-styled	2 ja.o Y	E. Indies 1799. C pi	Lot. mag. 1100
7412 africanum L.	African	1 jnjl Y	C. G. H. 1771. C pl	Bot. mag. 403
7413 quadrifolium L.	four-leaved	2 myjn Y	C. (%. H. 1787. C pl	JSolmag. 431
7414 viebsum L.	clammy	2 jnjau P	S. Europe 1818. D co	
7415 Apcrifibrium&i&	St John's-wt-lvd hairy	i jnjil P	Caucasus 1807. D co	Bot. mag. 1048
7416 mesothum L.	Ascyrum-Ivd	ljin.jl au B	Austria 1751. D co	Jac. au. 1.31
7417 ascyrinthum B. M.	fibrous	1 jl.au W	Portugal 1800. D co	Bot. mag. 108
7418 nervotoum W. Sf K.	Narbonne	liSnjl B	Hungary 1822. D co	W. & K. 105
7419 narbonense L.	most used	2 jnjl B	S. France 1759. D co	
7420 usitatissimum L.		2 jnjl B	Britain co. li. S co	Eng. bot 1367
7421 marginatum Poir.	angustifolium W. en.	tvAit'-marginced ^ A 1810. S co	
7422 reflexum //, K.	reflex-leaved			
7423 squamuollium Schr.	squamulose		S. Europe 1777. D co	
	austriacum Bieb.		Tauria 1819. D co	
7424 dittusum SchuU.	diffuse		B. 1823. S p.!	
7425 Lewis's P.	narrow-leaved		B N. Amer. 1820. S pl	
7426 angustifolium Hud.	Siberian		P England sa.pa. S co	Eng. bot. 381
7427 sibiricum Dec.	austriacum B. M. not L. perenne L.		B Siberia 1773. S pl	Bot. mag. 1086
7428 Anglicum Mil.	English	jt A or 2 jnjl B	England ch. pi. S pl	Eng. bot 40
perenne E. B.				
7429 austriacum L.	Austrian	k A or 1 jnjl B	Austria 1775. D co	Jac. au. 418
7430 roontanum Schl.	mountain	1 jnjl B	Switzcr. 1817. S pl	Scop. car. 11
7431 bicolor Schous.	two-colored	5£ or ljin.il Y.R	Morocco 1820. S co	
tenuifolium (3 bicolor Pers.				
7432 alphum L.	alpine	£ A or i jl.au B	Austria 1739. S co	Sw. fl. gar. 17
7433 decumbens Desf.	decumbent	£ A or li jnjl R	N. Africa 1817. S co	Desf. at. 1.79
angustifolium £ decumbens Pers.				
7434 grandiflrum Desf.	large-flowering	■ A or * Jnjil B	S. Europe 1820. ■ ■	Desf. at. 1.78
7485 tenuifolium i.	slender-leaved	5 A or II jnjl Pk	Eurojk; 1759. ■ ■	Jac. au. 3. 215
7436 salsoloides Lam.	Salsola-like	n. or 1 jnjl Pk	S. Europe 1810. ■ ■	Bar. ic. 795?
7437 suffruticosum L.	fluff ruticose tt.	LJ or 1 au Pk	Sluin 1759. ■ ■	Cav. ic. 2.108
7438 catharticum L.	purging	O w i jnjau W	Britain dr. pa. ■ ■	Eng. bot. 382
922. DRCVSERA L.	SUNDEW. (Drosos)	dew, dew-like glands on leaves.)	Droscritcear. 4.—32.	
7439 pauciflora Banks	few-flowered	I jnjau W	C. G. H. 1823. S bog	
7440 acatilis Thun.	Btcmless	Ijnjau W	C.O. H. 1823. S bog	
7441 rotundifolia L.	round-leaved	A i jlau W	Britain tur.bo. S bog	Eng. bot. 817
7442 americana W.	American	A i jnjau W	N. Amer. 1810. S bog	
7443 longifolia L.	long-leaved	A i jlau W.H	BriUin tur.bo. S bog	Eng. bot. 868
interini:dia Desv.				
7444 anglica Hud.	English	* A Pr i i u u w - R	England tur.bo. S bog	Eng. bot 869
7445 lincaria Got.	Knew. leaved	£ A P IJ a u F	N. Amer. 1818. S bog	
7446 filiform Rafi.	filiform	£ A pr i "Win P	N. Jersey 1811. S bog	
7447 binata Lab.	binate-faztvi	£iAJI>r ii>a u W	N. Holl. 1821. S bog	Lab. n.h. 1.105
923. FTBLIS Sal	BYBLTS. (<i>Bt/lilis</i> , the daughter of Miletus.)		Droscritceea. 1.	
7448 tiniflira 5a/.	Flax-flowcrcd	£ iAI pr i myjn B	N. Holl. 1800. S bog Par. Ion. 95	
924. SIBBA'LD/iJ L.	SIHBALDIA. (Sir Robert Sibbald, prof. of phys., Edin.) Rosnccat. 4.—V.			
7449 procumbens L.	procumbent	tA i jnjau Y	Britain sc.al. I) s. Eng. bot 897	
2 helvetica G. Don	Swiss	i jnjau Y	Switzerl. 1819. D s1	
3 amricaia G. Don	American	i Jnjau Y	N. Amer. 1820. D s1	
7450 parviflra W.	small-flowered	i jnjau Y	Cappadocia... D s1 W.n.bcr.nnjl,5.4	
7451 ericta W.	upright	1 jnjau Pk	Siberia 1806. D 81 Am. rut. 112.13	
	polygynous	i jnjau Y.a	Siberia 1824. D s1	
925. ALDROVAWDii L.	ALDROVANDA. (Ulysses Aldrovandus, an Ital naturalist)		Droscritceee. 1.	
7453 vesiculosa L.	vesiculose	* A cu ijnjau W	Italy 182a D i p Ac.bon.2.12	
926. COMMERSCPN/ii W.	COMMERSONIA. (V. Cotmnerson, Fr. traveller and bot)		Bi/tncrtcear. 2.—3* k	
	broad-leaved	* O or 3 jnjl W	Moluccas 1806. C i.p Bot. mg. H9	
	echinatcd	• or	Moluccas 1820. C p.i Ru. am. 3.119	
927. RUL'NG/iJ R. Br.	RULINUIA. (J. P. Ruling, author of Essay on Nat Ord.)		Bytnrcrtceaa. 4.	
7430 pannosa R.Br.	nggei Ueard «L J or ? my	W N. Holl. 1819. C Lp	Bot. mag. 2191	
7457 coryUolia Cunn.	Hazel-leaved	i j o r 3 ... W	N. Holl. 1824. C Lp	
7458 hermanni Vrhl/iaSwt	He r m a n n i-a lvdi jor	2 ... W	N. Holl. 1819. C s1	
7459 cistiflria Cun.	Cistus-leaved	MI J or 2 ... W	N. Holl. 1824. C Lp	
928. ARME'RIA W.en.	THRIFT. (<i>Armtria</i> , swoctlwlliam.In.)		Plumbagince. 19.—20.	
7460 vulgaris W. en.	common	A or i jnjau R	Europe ... D co Schk. han. 87	
7461 maritima W. en.	sea-side	A or i myjl R	Britain sea co. D s1 Eng. bot. 226	
751tice Armtria L.				
7462 nplna W. en.	alpine	A or i iny.au P	Carinthia ... D s1	
7463 arcnfria Pers.	sand	A or i my.au Pk	France ... D s1	
7464 littoralis W. en.	shore	A or 1 my.au Pk	S. Europe ... D s1	
7465 dianthoides Horn.	Pink-like	A or i my.jn Pk	Europe 1810. D s1 Sw.fl.gar. ic *	
7466 montana G. Don	mountain	A or i my.jn Pk	Scotland moun. I) 61	
7467 hiimilis Lk.	dwarf	A or i my.jn Pk	S. Europe 1817. D s1	
7468 hirta W.	hairy	A or i my.jn Pk	N. Africa 1820. D s1	
2iEphaUtes//. A.	rounf-headed	A or 1 my.jn Pk	Europe 1800. D s1	
7470 pongna R. & S.	Sungent	A or i my.jn Pk	Sinin 1818. D s1	
7471 juniuierif6ia W.		A or 1 my.jn Pk	Spain 1818. D s1 Qu. his. ■ ■	
7472^iniflra7ia7. & S.	unipcr-Icavcd	A or 1 my.jn Pk	Portugal ... D s1	
7473 rtlihcea W.	Pine-leaved	A or 1 my.jn VV	Spain 1718. D s1 Civ. ic. 2.109	
7474 denticullata Bert.	GzMc-lcavcd	j£ & or ljin.yjn F	Naples 1810. D s1	
7475 plantaginea W.	toothed	A or 1 my.jn R	S. Europe 1818. D s1	
	Plantain./wf	ff		

7476 <i>rcorgofimeftilla Lk.</i>	Bcorzonera-lvd £ A or 1 my.jn S	8. Europe 1816. D Bl
7477 <i>latifolia W. en.</i>	broad-leaved £ A or 2 my.jl L.R	Algarbia 1740. D s.p Jac. fin. 1.42
7478 <i>fasciculata W. en.</i>	bundled tt. (J or 2 ap.au F	Portugal ... D s.l .Ven.cels3S
r 929. <i>STA'TICE L.</i>	SEA LAEOT)ER. (Statixo, to stop; astringency.)	<i>Plutnbaeinca.</i> 61.—74.
7479 <i>fframinifolia H. K.</i>	Grass-leaved £	Siberia 1780. I) s.l
7480 <i>Limonium L.</i>	marsh Wild Beet £	England mud.s D r.m Eng. bot 102
7481 <i>carolinika Walt.</i>	Carolina £	Carolina 1820. B
7482 <i>Gmchni V.</i>	Gmclin's £	Siberia 17'6. I) &l Gm. si. 2. 90
7483 <i>Scopkria Pall.</i>	Broom £	Siberia 1796. I) 81
7484 <i>latifolia Sm.</i>	broad-leaved £	Siberia 1791. D pi
7485 <i>oleifolia Pou.</i>	Olive.leaved £	Italy 1688. I) &l
7486 <i>auriculatolia Vahl</i>	Auricula-lvd £	Barbary 1781. D s.l Scop. d. 1.10
7487 <i>cmarginata W. en.</i>	cmarginate £	Gibraltar ... D s.l
7488 <i>cordkra L.</i>	"i my.jl	S. Europe 1752. D s.l Bar. ic. 805
7489 <i>scabra Thuru</i>	f	C. G. H. 1788. S r.m
7490 <i>virgata W. en.</i>	1 my.jl	Spain ... D s.l
7491 <i>roticulata L.</i>	1} jn.au	England mud.s D s.l Eng. bot 328
7492 <i>tetragona Thun.</i>	i	C. G. H. 1820. D 61
7493 <i>dichotoma Caw.</i>	2 jn.au	S. Europe 1810. D B.l Cav. ic. 1.50
7494 <i>altaica Hort.</i>	i jn.au	Siberia 1820. D s.l
7495 <i>viminea &Ar.</i>	1 jn.au	... 1818. D 81
7496 <i>caspia W. en.</i>	i jn.au	Pa.B Caspian Sea ... S s.l Gm. si. 2.89.2
7497 <i>echioides L.</i>	1 jn.au	Pa. B S. Europe 1752. S s.l Fl.gr. 298
7498 <i>spathulata D ^.</i>	Echium-like £	Barbary 1804. D s.l Bot. mag. 1617
7499 <i>foillidifolia 5m.</i>	I	Greece 1810. D r.m FEigr. 293
7500 <i>greVsia Poir.</i>	Daifly-leaved £	Greece 1810. D r.m
7501 <i>Willdenovtma Pois».</i>	(Greek Willdenow's	France 1800. D r.m W.h.b. 63
	spatulata W. en.	
7502 <i>specifsa L.</i>	showy £	Russia 1776. D p.l Bot. mag. 656
7503 <i>conspicua A M.</i>	conspicuous £	Ajor jn.au Russia 1804. D s.l Bot. mag. 1.29
7504 <i>tatarica X.</i>	Tartarian £	Pk Russia 1731. I) p.l Sw. fl. gar. 37
7505 <i>ovalifolia Poir.</i>	oval-leaved £	Pk Russia 1816. I) s.p
7506 <i>flexuosa W.</i>	zig-zag £	Canaries 1816. I) s.p
7507 <i>purpurea L.</i>	purjilcd £	Siberia 1791. D p.l
7508 <i>minita L.</i>	small £	P C. G. H. 1800. D p.l
7509 <i>pubeficens Drc.</i>	pubescent £	R Mediterr. 16/8. D p.l Plu.al. 200.3
7510 <i>articulata Lot.</i>	jointed £	R S. Europe 1824. D s.p Boc. sic. 13
7511 <i>pectinata H.K.</i>	inclined £	R S. France 182a D s.p Loi. gal. 2.6
7512 <i>fechvus JKdf.</i>	hedgehog £	B Canaries 1780. D p.l
7513 <i>suffruticosa L.</i>	suffruticos £	R Caucasus 1813. C s.l Fl.gr. 300
7514 <i>monoptala L.</i>	one-petaled £	B Siberia 1779. D r.m Gin. si. 2.88.2,3
7515 <i>cinerea Poir.</i>	Iey	P Sicily 1731. C r.m Boc. sic. 16,17
7516 <i>pruinosa L.</i>	Fennel-leaved £	B C. G. H. 1810. I) s.l
7517 <i>erulacea Z.</i>	scollop-leaved £	W S. Europe 1823. D 81
7518 <i>ginuata L.</i>	Thouin's £	Y Siberia 1796. D s.l Plu. al. 28 4
7519 <i>Thouinw Viv.</i>	spiked £	P.Y Levant 1629. S r.m Bot. mag. 71
7520 <i>spicata W.</i>		B N. Africa 1700. S s.l Mart. ccn. ^i
lyrata Bieb.		I jn.au W Caucasus 1819. S 81 Gm. si. 2.91.2
7521 <i>australis Sjw.</i>	southern £	N. Holl. 1823. S s.l
7522 <i>Taxanthema austrikis It. Br.</i>		
7522 <i>alata IV. en.</i>	winged £	least O cu ±ap.my Y Britain cor.fi. S co Eng. bot. 435
7523 <i>mucronata L.</i>	mucronate £	1 jn.au P.Y 1806. D s.l
7>24 <i>globularinfolia' Desf.</i>	Globularia/-wj	ljin.au R Barbary 1784. C r.m Her. st18
incana L.		1 my.s W Sicily 1822. D r.m Bar. ic. 793
7526 <i>macrophilla TV.</i>	hoary £	1 jn.au Pk Egypt 1823. D r.m
Z^7 egyptica Pers.	long-leaved £	2 my.jn W Canaries 1824. C r.m
7528 <i>auriculata Bes.</i>	Egyptian £	I my. W Egypt 1823. D r.m Bot. mag. 2563
?529 dotalis S».	eared £	i jn.au B Galicia 1817. D 81
7530 <i>cuneata // T.</i>	tall £	1 jn.au B Siberia 1820. D s.l
	wedge-leaved, £	ijn.au B Siberia 1820. D s.l

CLASS V. ORDER 6.

PENTA'NDRIA POLYGY'NIA.

930. MYOSITRUSX. MOUSE-TAIL. (*Mys.a mouse, oura, a tail; spiked receptacle/*) *Ranunculheea.* 1.
531 minimus L. least O cu ±ap.my Y Britain cor.fi. S co Eng. bot. 435

 Moen. CEHA TOCEPHALUS. (*Krasa, a horn, kepha/c, a head; fl.*) *Ranunculcea.* 2.
orthoceras Dec. straight-horned O cu i my Y S. Europe 1739. S co Jac. au. 48
AanUnculus falcatus Bieb.

 Parsley-leaved * or 3 fap (Xanthos, yellow, rhixa, a root) Banunculcear. 1.
HerU YELLOW or 3 fap P.Y N. Amer. 1766. Sks.p Bot mag. 1736

CLASS VI. ORDEK].

HEXA'NDRIA MONOGY'NIA.

• IV3. ATARCISSUS /..	NARCISSUS.	(Narke, stupor; effects of smell.)	Amarylidea.: 77.-
7535 pofticua L.	poet's ff	A or I my W	S. Kuro>c O co Par. p. 70*
7536 recfrvus Haw.	recurved-taHfdtf	A or I my	S. Eurojic O co
7537 patcfariB Sal.	bwin-flwd tf	A or I my W	England O co Eng. bot. 275
<i>iracticus E. B.</i>	<i>majMis B. M.</i>		
2 exsbertus Haw.	drawn out tf	A or I my W	England gard. O co
3 pl&mus Haw.	double tf	A or I my W	England gard. O co Par. p. 75.1
7538 aiiBUHffolius B. M.	narrow-leaved	tf A or ap.my W	S. Europe 1570. O co liot mag. 193
<i>radufloruh Sal.</i>			
7539 biflrrus B. M.	two-flowered tf	A or I ap.my W	Britain mea. O co Eng. tot. 27G
7540 tenuior // K.	slenderer tf	A or I my L,Y	1789. O co Bot mag. 379
7541 crenulatus B.M.	crenulated tf	A or I mr.ap W	SnaV*** O co
7542 Eremalhus B.M.	Trew's tf	A or I mr.ap W.v	Spain O r.m Bot. mag. 940
7543 homomus Sch.	bundle-flu d tf	A or I mr.ap W.v	Spam O r.m
7544 fiotulmus /aw.	hollow-stalked tf	A or I 2 ap W.v	O r.m
7545 cerninus Haw.	wax-colored tf	A or I 1 ap W	O r.m
7546 Taxiflora L.	Tazetta, or <i>Polyanthus</i> tf	A or I mr.ap yr	Spain 1759. O s.l Par. p. 81.3
7547 multiflribus Haw.	tall many-flwd tf	A or I ap Y	— O s.l Bot mag. 925
<i>Tazetta B. M. not L.</i>			
2 atricus Haw.	golden many-flwd^ A or	I ap Y	— O s.l Bot. mag. 1026
7548 obslMus Haw.	obsolete tf	A or I au.s W	Spain 1819. O s.l Par. p. 89.4
7549 latiflbus Haw.	broad-lvd. orange tf	A or I ap.my W	— O s.l
7550 Macattia Lindl.	Mac Leay's tf	A or I ap.my Crea	Smyrna 1815. O s.l Bot. reg. 987
7551 orientMis /«	oriental tf	A or I mr.ap W	Levant O co Bot. mag. >48
7552 papyracus B. M.	paper tf	A or I mr.ap W	O co Bot mag. 947
<i>unicolor Ten. nfcvius L.</i>			
2 Jasmincus Haw.	Jasmine-like tf	A or I ap W	— O s.l
7553 Lilna Haw.	Luna tf	A or I ap W	— O s.l
7554 grandiflbrus Haw.	great-flowered tf	A or I ap W.Y	— O s.l
7555 citrinus Haw.	Citron-colored tf	A or I ap W.Y	— O 0.1 Bot. mag. 946
<i>oricon&lis y B. M.</i>			
7556 italicus B. M.	Italian tf	A or I mr.ap Pa.Y	S. Europe — O co Bot mag. 1188
<i>pYa;cox Ten.</i>			
2 scimpilenus	semi-double tf	A or I mr.ap Crea	Italy — O s.l
3 plenus	double tf	A or I mr.ap Crei	Italy — O p.l
7357 tereticafthis Haw.	round-stalked tf	A or I mr.ap Pa,Y	Spain — O co Bot mag. 1298
7558 compressus Haw.	flat-stalked tf	A or I mr.ap L,Y	Spain — O co
7559 blfrons B. M.	two-faced tf	A or I mr.ap Y	S. Europe — O co Bot. mag. 1186
7560 primulhus Haw.	Covi&Wp-citpped tf	A or I mr.ap Y	— O co Bot mag. 1299
7561 Jonquilua L.	Jonquil tf	A or I 5 ap.my Y	Spain 1596. O r.m Bot. mag. 15
2 fibre pldno	double tf	A or I 1 ap.my Y	Spain 1596. O r.m
7562 Hmlis Haw.	similar Jonquil tf	A or I «P Y	— O s.l
7563 gracilis Lindl.	slender tf	A or I 1 ap.my Y	— O co Bot reg 816
7564 hcininillB Haw.	lesser curled cup tf	A or I np Y	— O s.l Par. p. <3.4
7565 viridiflbrus Schous.	green-flowered tf	A or I au.o G	Barberry 1629. O r.m Bot mag. 1687
7566 scrotinus L.	late-flowered tf	A or I au.o Pa.Y	Barberry 1629. O r.m Cl. h. 252
7567 trilobus L.	three-lobed Jonquil tf	A or I 1 ap.my Y	S. Europe HB9. O r.m Bot mag. 78
calathinus Sal.	odurus B. M. tmparttus Horn.		
7568 interjeutus Haw.	interjected^ A	I ap Y	1810. O &1
7569 odhrus L.	sweet-scented A, or	I ap.ny Y	& Europe 1629. O 8.1 Botmag.<)Sl
calathina et B. M.			
7570 pusillus G. Don	small Jonquil tf	A or I ap E	— 1818. O s.1
Phylgyne minor Haw.			
2pknu8 double Q. Anne's Jonquil tf	A or	I lap T	1810. O 8.1
7571 rugulbus Haw.	rugulose cup tf	A or I ap T	1818. O s.1
7572 nifins Haw.	nodding tf	A or I mr.my Y	S. Europe 1789. O 8.1 Bot. mag. 945
trilobus B. M.			
7673 concolor Haw.	self-col. sulphur tf	A or I ap Y	1820. O 8.1
7574 infundibulkris Sal	funnel-flowered^ A or	I mr.my Y	— O 8.1 Par. p. 92
7575 pulchelluk Haw.	pretty tf	I mr.my Y	Spain — O s.1 Bot mag. 1262
7576 stri&tuhis Haw.	small-channelled tf	A or I *ap *Y	— O s.1
7577 triandrus L.	triandrous tf	A or I *ap.my W	Portugal 1629. O 8.1 Par. p. 93.2
7578 cernuus Sal.	pale drooping tf	A or I ap Y	1818. O 8.1 Bot. mag. 48
<i>triandrus B. M. not L.</i>			
7579 capax Sal.	capacious tf	A or I ap Pa.Y	— O 8.1 Red. Ill. 177 ?
7580 montanus B. R.	mountain tf	A or I ap.my W	Portugal ... O s.1 Bot reg. 183
7581 galanthiflbus Haw.	Snowdrop-lvd tf	A or I «my W	— O 8.1 Par. p. 73
7582 aUicans Haw.	whitening tf	A or I mr.ap Pa.Y	— O 8.1 Par. p. K16.
7583 Bulbocodium /.	Bulb, or Hoop-pet. tf	A or I mr.ap Y	Portugal 1629. O 8.1 Bot. mag. 88.
7584 inflatus L.	inflated tf	A or I mr.ap Y	— O 8.1
2 minor Haw.	lesser swln tf	A or I mr.ap PaSu	Spain 1629. O 8.1 Par. p. 107.8
7585 lobulMus Haw.	small-lobed tf	A or I ap.my Y	— R s.l
7586 tenuiflulus Sal.	slender-leaved tf	A or I 5 ap.my W	— O si Sw. fl. gar. 11*
7587 incomparabiis B.M.	incomparable tf	A or I ap.my Y	Portugal 1629. O co Bot mag. 121
7588 gibus Haw.	incomparable white tf	A or I ap W	— O 8.1
2 pl&mus Haw.	double orange Plicnixti A or	I ap Su	— O 8.1
7589 semipatius Haw.	clove-cupped tf	A or I mr Su	— O 8.1
7590 Haw6rthi G. Don Ha worth's tf	A or	I ap Y	1700. O 8.1
Quclth c6ncolor Haw.			
2 pfenus sulphCireus Haw.	dble-sulph. tf	A or I ap Su	1629. O 8.1
7591 aurantiu6 Haw.	orange mixed tf	A or I mr Y	1629. O 8.1
2 planss Kitew au-	C double yellow"	I mr Y	1629. O 8.1
rantius Haw.	£ and orange S	I mr Y	1629. O 8.1
7592 tortoKus Haw.	tw&sted-petaled tf	A or I Hap W	Spain 1629. O co Par.p.101-*
moschktus L.	musk tf	A or I mr.ap W	Spain 1759. O co Bot mag. 9BJL
7594 patulus Sal.	uioschatus & B.	I mr W	Spain — O co Bot. mag. 130"

2 <i>dlbens</i>	flaw,	whitish <i>Nor/oikfl</i>	A or	I n r	W	Spain	-	O co
7595 <i>serratus</i> <i>Haw.</i>	serrated	tf A or	S mr.ap	Pa.Y	S. Europe	-	O co	
2 <i>suavis</i> <i>Haw.</i>	sweet	tf A or	mr.ap	Pa.Y	S. Europe	-	C) co	
7596 <i>spbrius</i> <i>J/aw.</i>	spurious	tf	ap	Y	England	-	O co	
7597 <i>obscissus</i> //aw.	dipt trunk	tf	my	Y	Spain	-	O co	
7598 <i>Telambnius</i> <i>Haw.</i>	Tciamon's	tf A or	ap	Y	-	-	Par. p. 107	
2 <i>pi en us</i> //atu.	double <i>Daffodil</i>	tf & or	ap	Y	-	-	O co	
3 <i>grandifipenus</i> <i>Haw.</i>	great double	tf A or	ap	Y	-	-	O co	
7599 <i>Pseido-iVarcissus L.</i>	Daffodil	tf A or	mr.ap	Pa.Y	England woods	O co	Par. p. 101.1	
festMis //aw.							Eng. bot. 17	
2 <i>seditionis</i> //aw.	Scotch <i>Daffodil</i>	tf A or	mr	Y	Scotland gard.	O co	*	
3 <i>pl&nus</i> //aw.	double <i>Ihyffodil</i>	tf A or	my	Y	France	...	O co	
4 <i>plenissimus</i>	most double	tf A or	mr.ap	Y	-	O co	Par. p. lul.6	
7600 <i>tubiflbrus</i> <i>Sal.</i>	tube-flowered	tf A or	mr.ap	W.v.	-	O i'o		
7601 <i>lorifolius</i> <i>Haw.</i>	strap-lvd two-cld\$	A or	mr	Y	-	O co	Bot. mag. 1187	
bfcolor <i>B. M. not L.</i>								
2 <i>breviflos</i> <i>Haw.</i>	short-flwd two-cldf	A or	mr	Y	-	O co		
3 <i>anceps</i> <i>Haw.</i>	two-edged two-cld\$	A or	mr	Y	180)	O co	
7602 bcolor /.	two-colored	tf A or	ap.my	W. T	Spain	1629.	O co	
7603 Sablni Lindl.	Sabine's	tf A or	ap.my	Y	-	O co	Bot. reg. 732	
7604 nfveus IV. en.	snowy	tf A or	my	W	S. Europe	...	O co	
7605 obvalfris <i>Sal.</i>	trenched round	tf A or	mr.ap	Y	Spain	...	O co	
7606 major 2*. <i>M.</i>	larger	tf A or	mr.ap	Y	Spain	1629.	O co	
7607 propfnnus <i>Sn.</i>	allied	tf A or	mr.ap	Y	Spain	1629.	O co	
7608 n6bili8 //aw.	noble	tf A or	mr.ap	Y	-	O co	Botmag.1301.su.	
7»» A'jax Sal.	Ajax	tf A or	mr.ap	Y	-	O co		
7610 phmilius <i>Sal.</i>	dwarfish	tf A or	mr.ap	Y	Spain	...	H. flor. 8	
7G11 minor <i>L.</i>	smaller	tf A or	mr.ap	Y	Spain	1629.	O s.l Bot inag 6	
*934. PANCRATIUM L. PANCRATIUM. (<i>Pan</i>, all, <i>kratys</i>, potent; med. qual.)								
7612 <i>zylfinicum</i> <i>L.</i>	Ceylon	tf E or 1	jnjr	W	Ceylon	1752.	Sk r.m Bot. reg. 47!	
7613 <i>longiflrum</i> <i>Box.</i>	long-flowered	tf ftflor	2 jnjl	W	E. Indies	IH10.	O r.m	
7614 <i>vere6ndum</i> <i>Sol.</i>	ruddy	tf fA or	lljn.au	W	E. Indies	1776.	Sk r.m Bot reg. 413	
7615 <i>marlitnum</i> <i>L.</i>	sea	tf A or	2 my jl	W	S. Europe	ISW7.	Sk s.p Bot. reg. 161	
7616 <i>caroliniknum</i> <i>L.</i>	Carolina	tf A or	2 jnjl	W	Carolina	1751.	Sk r.m Cat. car. 3.5	
7617 <i>canariense</i> <i>Ker</i>	Canary	tf IAJOT	ljnjnjl.	W	Canaries	1815.	Sk r.m Bot reg. 174	
7618 <i>ill^ricum</i> <i>L.</i>	Illyrian	tf A or	1 my,jn	W	Europe	1615.	Sks.p Bot. mag. 718	
7619 <i>undufatum</i> <i>Kth.</i>	v/nvo-leaved	tf CAT or	1 Jnjl	W	S. Ainer.	-	Sk r.m	
7>20 <i>expansum</i> <i>B. M.</i>	expanded	tf (25 or	2 jnjl	W	W. Indies	IH20.	O r.m Bot mag 1941	
7621 <i>litorMe</i> <i>Jac.</i>	sea-side	tf El or	2 my,au	W	S. Amer.	1758.	Sk r.m Bot. mag. 1879	
7622 <i>DryAndri</i> <i>Ker</i>	Dryander's	tf /Alor	2 my,au	W	-	-	Sk r.m Bot. mag. 8>5	
7623 <i>angustum</i> <i>Ker</i>	nariow-leaved	tf 2S,Jor	lmy,au	W	-	-	Skr.m Bot. reg. 221	
7624 <i>dlstichum</i> <i>Herb.</i>	two-ranked	tf GS,Jor	1* my,au	W	S. Amer.	-	O r.m	
7t25 <i>crassifbVmn</i> <i>Herb.</i>	thick-leaved	tf ft a	1jmy,au	W	S. Amer.	-	O r.m	
7626 <i>rotatum</i> <i>Ker</i>	wheel-crowned*	tf AJ or	1 Jsj	W	Carolina	1803.	Sk r.m Bot mag. 1082	
7>27 <i>mexichnum</i> <i>L.</i>	Mexican	tf - 123-	1 au	W	Mexico	1732.	Sk r.m Pi. el 2ift. 289	
7028 <i>acutiflbum</i> <i>Swt.</i>	sharp-leaved	tf I&J or	2 my,au	W	Mexico	1824.	O r.m Bot reg. 910	
<i>mexicnum</i> <i>B. R.</i> not of others, //emero ^c allis								
7629 <i>gianense</i> <i>Ker</i>	Guiana	tf El or	n	W	Guiana	1815.	Sk r.m Bot reg 265	
7J30 <i>at ilof</i>	spreading	tf El or	■ jl,au	W	W. Indies	IK22.	Sk r.m lied. lil. & 380	
?<31 pediMeLoA	foot (<i>lonffislwd</i>)	tf El or	■ au	W	Brazil	IH20.	Skr.m Bot cab. 8>?	
7632 <i>hragrans</i> <i>Sal.</i>	fragrant	tf El or	■ my,au	W	W. Indies	IS1<.	Sk r.m Bot cab 834	
7f33 <i>billiflorum</i> (w.	two-flowered	tf El or	■ my,au	W	E. Indies	1820.	Sk r.m	
7M4 <i>cariba?um</i> <i>L.</i>	Caribbean	tf El or	1jmy,au	W	W. Indies	1730.	Sk r.m Bot. mag. 826	
76to <i>declinatum</i> <i>Jac.</i>	declined	tf El or	2 my,jl	W	W. Indies	1825.	O r.m Bot cab. 558	
736 <i>amafranum</i> <i>Sal</i>	handsome	tf El or	1 my,au	W	Guiana	1790.	Sk r.m Bot. mag. 1467	
56J7 ovatum Jlfpt.	ovate-leaved	tf El or	1 my,au	W	W. Indies	...	Sk r.m Bot. reg. 43	
42 P 16 Sal.	showy	tf El or	11 my,au	W	W. Indies	1759.	Sk r.m Bot mag. 1453	
americanum <i>Mil</i>	American	tf ■■■■■	2" jnjl	W	Jamaica	1820.	O r.m	
*935. ISMETNJS; Sal.								
7640 nMans <i>Herb.</i>	ISMENE.	(The daughter of Cfidipus and Jocasta.)					<i>Amaryllfdec.</i>	3.
7641 <i>ancrJitum</i> <i>calathnum B. Af.</i>	nodding	tf C3 or 2} jnjl,ji	W	Brazil	1800.	Sk r.m BoL mag. 1561		
?WI <i>calathium</i> <i>Herb.</i>	cu <i>v</i> -Jtowered	tf Elor	2 h	W	Brazil	1800.	Sk r.m Bot. reg. 215	
?>42 <i>Andncaes</i> <i>Herb.</i>	Amancaes	tf Elor	■■■■■ Y	Peru	1804.	Sk r.m Bot reg. 600		
<i>Pancratium</i> <i>Amdncaes</i> <i>B. M.</i>								
& *: EUCRO5IA Ker								
**3 bcolor Ker	EUCHOSIA.	{Eu, well, krossos, a fringe; cup of stamens.)					<i>AmaryUidece.</i>	1.
	two-colored	tf El or	1 ap-my	O	CapeHor.	1816.	O ltl Bot. reg. 207	
7f7* EU R CYLES Sal. (<i>Eurus</i> , broad, <i>kleio</i> , to close up; perianth tube by stamens.)								
○* <i>amboinlsn</i> <i>Sal.</i>	Amboyna	tf El or	2 my,jn	W	Amboyna	1759.	O ltl Bot. mag. 1419	
KA an cratium araboifnus L.	Arabian	tf El or	1 my,au	W	Arabia	1759.	Sk r.m Bot. reg. 215	
** nervisa G. Don	nrvycd-leaved	tf El or	1 my,jn	W	E. Indies	1822.	O ltl	
7f. winum nerv5sum llor.	showy	tf El or	11 my,au	W	W. Indies	1759.	Sk r.m Bot. reg. 43	
** australasia G. Don	Australasian	tf El or	1 my	W	N. Holl.	1821.	O ltl Bot. reg. 715	
•rancratium australasicum Ker.	australis	tf El or	1 my	W	N. Holl.	1821.	O ltl Bot. reg. 715	
7641 F AIOSTE MMA R Br. CALOSTEMMA. (<i>Kolas</i> , beautiful, <i>temma</i> , a crown; flower.)								
7642 H ln K,r	ye"ow	tf tf M,or	1 n	Y	N. Holl.	181b.	O sl Bot reg. 421	
7643 ! »! Mirro in R. *r.	purple	tf tf LAJor	1 n	P	N. Holl.	1810.	O pl Bot. rcf. 422	
! L album R. Br.	fragrant	tf tf iAJ or	1 my,jn	Y	N. Holl.	1824.	O pl	
CHLIDA'NTHUS Herb.	white	CULIDANTH.I.S.	(<i>Chlidcios</i> * delicate, <i>anthoa</i> , a flower.)		<i>Amaryllide</i> *.	I.		
Agrans Lndl.	fragrant	tf iAJ or	1 my,jn	Y	B. Ayres	1820.	O ltl Lmd.col.34	
Pancratium Ihtoum	Pair.							
CHRXSIPHrALA Ker								
S J V a * r -v	veHow	tf ^or	f my	O	Peru	1820.	O ltl Bot reg. 778	
ato steno x8on flavum Herb.								
Z Paucifl&ra Lindl.	few-flowered	tf uA7 or	i my	O	Peru	1822.	O ltl Hook. ex. fl. 132	
tel cur* lcnt.ta Herb.	curve-toothed	tf LAJ pr	1 o	Y	Peru	1825.	O pl Bot reg. ■■■	
** crucea Red.	saffron-colored	tf tAJ or	1 my	Cop	S. Amer.	1820	■■■	
re Ker	LOPIHOLA.	d'yim of lophia, a crest; crest of the sepals.)						
i-onostyhs americana Ph.	golden	* A »r	» my,il	Y	N	AUKT 1811.	D p.] Bot mag. 1596	

- | | | | |
|---|---|---|--------------------------------------|
| 12. ARGOLA'SIA. | ARGOLABIA. | {Argos, white, <i>lasios</i> , woolly; outside of calyx.) | <i>Hatnodoracear.</i> |
| 7656 plumusa J. | feathery | £ ij or 1* | W C. G. H. 1787. & i.l.p |
| Lanaria plumtaa H. _% K. | | | |
| 9*3. ANIGOZA'NTHOSA. Br. ANIGOZANTHOS. (Anigozo, to expand, <i>anthos</i> , flower.) | <i>Hcemodorrhcecc.</i> 2. | | |
| 7657 flavidia R. Br. yellowish-gr. _{<} VR-/jIM _% IAI or 3 my.s | IAI or 3 my.s | Y N. Hou. 1808. D s.p Bot. mag. 1151 | |
| Schwagrichenfa illvida Spr. | | | |
| 7658 rufa Lab. | rufous | £ iAI or 1\$ my.s | Ru N. Hou. 1824. D l.p Lab. vo. 1.22 |
| 944. BARBAC&XZA Van. BAHDACSNA. (2&&IC<KI, govt. of MinasGeraes.) | <i>Harmottordcete.</i> l.—l. | | |
| 7659 purporea Hook. purple | £ E J or 1 j1 | P Brazil 1825. D s.l Bot mag. 2777 | |
| 945. CONO'STYLIS R. Br. CONOSTYTILIS. (Konos, a cone, <i>stylos</i> , a style.) | <i>Hcemodoritccce.</i> 3. - 4. | | |
| 7660 aculeata R. Br. prickly | £ LAJ or 1 fo. | ifon. 1820. D i p | |
| 7661 stigera R. Br. stigerosus | £ IAJ or 1 | N. Hou. 1825. D s.p | |
| 7662 serrulata R. Br. serrulated | £ IAJ or 1 | N. Hou. 1824. D s.p | |
| 946. URA ^X NIA Schreb. URANIA. (Ouranios, sublime; stateliness.) | <i>Musdcea.</i> 1. | | |
| 7663 speciosa W. showy | £ El or 20 ... | II Madagasc. ... Skp.l Jac. sc. 1.93
<i>Javentula madagascariensis</i> Sonn. | |
| * 947. BUONAPA'RTE if Fl. per. BUONAPARTBA. (Napoleon Buonaparte.) | <i>Bromeliaceof.</i> 1. | | |
| 7664 juncea Fl. per, Ku'h4eaved | jj E3 or + ... | B Peru 1800. S s.l Fl. per. 3.242 | |
| • 948. AGA ^V E L. . . 1 AOAVB. | (Altered from <i>agauos</i> , admirable.) | | <i>Bromeliacea.</i> 11. |
| 7665 americana L., common 2 variegata | American variegated Amer. c | £ S. Amer. 1640. Skr.m Ilot.rcp.4S8 | |
| 7666 Milleri Haw. Miller's | £ | S. Airier. 1600. Sk r.m | |
| 7667 flaccida Haw. flaccid | £ | 170U Skr.ir | |
| 7668 lunda Jac. lurid | £ | S. Amer. 1790. Skr.m | |
| 7669 angustifolia Haw. narrow-leaved | £ | VeraCruz 1731. Sk r.m Bot mag. 1528 | |
| 7670 yoi&ftlia W. Yucca-leaved | £ | 1790. Skr.m | |
| 7671 Karaite Mil. Karatto | £ | 1819. Skr.m | |
| 7672 mexicana Haw. Mexican | £ | S. Araer. 1768. Skr.m | |
| 7673 vivipara L. viviparous | £ | Mexico 1817. Skr.m | |
| 7674 polyacanthina flata many AMrs spnd | £ | & Amer. 1731. Sk s.p Com. pr. 15 | |
| 7675 virginica L. Virginian | £ | 1800. Skr.m | |
| | | N. Amer. 1765. Sk r.m Bet mag. 1157 | |
| 949. FURCREM Yen. FURCRQSA. (M. Fourcroy, a celebrated chemist.) | <i>BromeUhcea.</i> 7. | | |
| 7676 gigantea Ven. gigantic | ■ 23 or 20 ja.s | G S. Amer. 1690. Sk r.m Bot mag. 2205 | |
| Agave foytida L. tuberous | ■ | O S. Amer. 1739. Skr.m | |
| Ififi tubercifia Haw. Cuba | ■ | G S. Amer. 1739. Sk r.m Jac. am. 260. 25 | |
| 7678 cubensis Haw. rigid | ■ | S. Amer. 1768. Sk r.m | |
| 7679 rigida Haw. southern | ■ | N. Holl. 1811. Skr.m | |
| 7680 austriiis Haw. Cantula | ■ | China 1818. Sk r.m | |
| 7681 Cintula Wai. Madagascar | ■ | Madagascar. 1825. Skr.m | |
| 950. LITT-ffIM Brig. LITHBA. (Duke of Lptta, near Milan.) | <i>Bromelidcea?</i> 1. | | |
| 7683 geminifera Brig. twinflowered | £ A or 15 ja.au | G America 1810. Skr.m Jo.sc.3.1 | |
| Agave geminiflora Xer. Yer. Buonapartba Juncea Haw. and Hort. | | | |
| • 951. BROME^L/l.IL. BROMBLIA. (Olafus Bromel, a Swedish botanist) | <i>Bromelidcea.</i> 13.—18. | | |
| 7684 semiscirrata IV. en. half-sawed-forf | £ Ej or 3 year | G S. Amer. ... Sk r.m | |
| 7685 Pinguin L. Pinguin | £ El ec 3 HIAP | R W. Indies 1690. Sk r.m Jac. am. pic. 91 | |
| 7686 sylv&tris W. wild | £ QS or 3 JI | C S. Amer. 1820. Sk r.m Bot. mag. 2392 | |
| 7687 fastuosa Lindl. proud | £ (23 or 4 au.s | P S. Amer. 1815. Sks.p Lind.col. 1 | |
| 7688 KarMus L. Karatas | £ (23 or 8 Pk | W. Indies 1739. Sk r.m Jac. vin. 1.31,32 | |
| 7689 chrydintha Jac. golden-flowered | £ (23 or 2 Y | Caracas 1819. Sk s.p Jac. sc. 1.55 | |
| 7690 hngulka L. tongue-leaved | £ (23 or 1 J my.jn | Y S. Amer. 1759. Sk r m Pi. ic 64.1 | |
| 7691 bractea;xta Swx. red-bracted | £ (23 or 2 8.0 | Pk Jamaica 1785. Sk r.m Par. Ion. 40 | |
| 7692 Acringa L. Acanga | £ El or 2 | Brazil 1822. Sk s.p Pis. br. 91 | |
| 7693 exstidans Lod. sweating | £ fgt T^ or 2 s.o.. | W. Indies 1820. Sk r.m Bot cab. 801 | |
| 7694 paniculigera Sam. panicle-bearing | £ El or 2 ap.my | W. Indies 1822. Skr.m | |
| 7695 htumilis Jac. low | £ (A)or 1 mr | Pk 1789. Skr.m Jac.ic. 1.60 | |
| 7696 sessiliiflora Lo. C. sessile-flowered | £ El or 3 jl.au | S. Amer. 1826. Sk r.m | |
| 952. AfIANA'SSA Thun. PINE-APPLE. (Nonas, the name in Guiana.) | <i>Bromelidcete.</i> 4. | | |
| 7697 bracteata B. R. bracted | £ El ft 3 mr | C.p Brazil 182a Sk r.m Bot reg. 1081 | |
| 7618 debiliu B. R. weak | £ El ^ 3 mr | C Skr.m | |
| 7699 Ificida B. R. shining | £ El fir 3 year | Pk S. Amer. ... Sk r.m Di. el. 25.21.22 | |
| BromeliYi Ificida W. cultivated | £ [23 rf 3 year | P S. Amer. 1690. Sk r.m Bot mag. 1554 | |
| 7600 sativa B. R. Bromelia Ananas L. | | | |
| 953. GUZMA'N.N/i! Fl. per. GUZMANNIA. (A. Guzman, collector of natural history.) | <i>Bromeliacete.</i> 1* | | |
| 7701 tricolor Fl. per. three-colored | £ El or A my | G.s S. Amer. 1820. Sk r.m Lind. col. 8 | |
| | | | |
| 954. PITCANtN/* Herit FITCAIRNIA. (W. Pitcairn, M.D., of London.) | <i>Bromeliacce.</i> 16.—19. | | |
| 7702 bToiac Uatulfia H. K. Pine-apple-lvd | £ (23 or 2 jn | S Jamaica 1781. Sks.p Bot. reg. 1011 | |
| 7703 angustifolia H. K. narrow-leaved | £ I or 2 year | SantCruz 1777. Sk *p Bot. mag. 1547 | |
| 7704 integrifolia B. M. entire-leaved | £ I or 2 au | K W. Indies 1800. Sks.p Bot.inag. 1463 | |
| 7705 latifolia H. K. broad-leaved | £ I or 2 au.s | S W. Indies 1785. Sk s.p Bot mag. H56 | |
| 7706 flammea B. R. dame-colored | £ 2 n F | It Janeiro 1825. Sk s.p Uotreg. Ju92 | |
| 7707 bracteata H. X. tar^e-bracted red | £ Q3 or 2 apamy | W. Indies 1799. Sks.p lied. UL 7, 7* | |
| latifolia Red. | | | |
| 7708 furfuracea W. en. fufuraceous | £ El or 2 ja.au | R a Amer. 1816. Sk r.m | |
| 7709 staminea B. M. /ov-stamined | £ El or 2 ja | S. Amer. 1823. Sk r.m Bot mag. 114 | |
| 7710 Jchil6nsisZo. C. Chile | 1 ja | Chile 1820. Sksp | |
| 7711 hfilmis Hort. low | 1 ja | S. Amcr. 1820. Ske.p | |
| 7712 intermedia Hort. intermediate | 2 ja | S. Amer. 182a Sk ap | |
| 7713 mdiflbra Hort. Iris-flowered | 2 ja | S. Amer. 1820. Skr.m | |
| 7714 media Hort. mediate | 1 ja | S. Amer. 1820. Skr.m | |
| 7715 sulphurea B. Ren. 2 ja | 2 ja | W. Indies 1797. Sks.p Bot mag. 1*g | |
| 7716 albinos Herb. sulphur-JfuKi if | 3 s | Brazil 1824. Skp.l Bot. mag. 2tJ | |
| 7717 suaybolena B. R. white-flowered | 2 ja | Bnuul 1824. Sk r.m But mag. !!!! | |

* 955. POURRET/v* R. & P. POURRRTIA. (A Pourret, a French botanist) *BromelBcea:* 3.-6.
 7718 pyramidata R. 8f P. pyramidal £ El or 1 jnjl Y Peru 1822. Sk s.p Kl. per. 3.257
 7719 coarctata R. 8f P. coarctite £ El or lftmyjn Y Chile 1822. Sk s.p Feu. ob. 3.39
 7720 magnifspatha Coll. large-spathe £ El or 5 my.jn G.w S. Amer. 1820. Sk s.l Col. h. rip. 19

* 956. TILLA'NDSM L. TILLANDSIA. (*Ellas Tillandfius*, professor of physic at Abo.) *Bromeliacec.* 28.-40.
 7721 usneoides L. Usnca-like £ El or 6 1 W. Indies 1833. Skp ■■■■■
 7722 Bartramia EIL Bartram's £ El cu ■■■■■ Carolina 1825. Sk s.p
 7723 recurvata L. recurved-Jurf £ El or ■■■■■ Jamaica 1793. Sks.p SI.jam. 1.121.1
 7724 xiphoides B. R. Xiphium-hkc /? El or ■■■■■ W B. Ayrcs 1810. Sks.p Bot. reg. 105
 7725 scrata L. auw-lcaved j| El or ■■■■■ Y Jamaica 1793. Sk s.p PL ic. (ff. 75.1
 7726 setacea Swx. setaceous £ El or ■■■■■ B W. Indies 1824. Sks.p
 7727 stricta B. M. strict £ El or ■■■■■ B Brazil 1810. Sk s.p Bot mag. 1529
 7728 bulbosa Hook. bulbous £ El pr I au B Trinidad 1823. Sk r.m Hook. ex. fl. 173
 7729 pAlhra Hook. fair £ El pr 0 fn Pk Trinidad 1823. Skr.m Hook.ex.fl. 154
 7730 obscura Lo. C. obscure £ El or 2 jn.au ■■■■■ S. Amer. 1820. Sk s.p
 7731 aloifolia Hook. Aloc-lcaved jSfAicu 1 n.d Pk Trinidad 1824. Sk s.p Hook. ex. fl. 205
 7732 nitida floo*. shining £ El cu 2 n.d B Jamaica 1823. Sks.p Hook.ex.fl.218
 7733 utriculata L. bladdered gfAlor 2 ... P.v S. Amer. 1793. Sk s.p
 7734 tenuifolia L. slender-leaved £ ITlor ■■■■■ B W. Indies 1825. Sks.p
 7735 flexubsa Stuz. 2 pallida flexuose /| El or 1 ■■■■■ B W. Indies 1790. Sks.p Jac.am.63
 7736 ineps Lod. pale /j El or 1 jnjl Y W. Indies 1815. Sks.p Bot reg. 749
 7737 nbtans Swz. two-edged £ El or ■■■■■ B W. Indies 1820. Sks.p Bot cab. 771
 7738 fasciculata Swz. nodding £ El or 2 au B Jamaica 1793. Sks.p
 7739 canescens Swz. fascicled £ El or 1 jnjl B W. Indies 1820. Sks.p
 7740 paniculata L. canescnt £ El or ■■■■■ B W. Indies 1824. Sks.p
 7741 ramosa GUL panicld £ fAI or jnjl B W. Indies 1820. Sks.p
 7742 rigida GUI. ramos £ (AJ or jnjl ■■■■■ Chile 1823. Sks.p
 7743 polystachya Spr. rigid £ V±or jnjl ■■■■■ Chile 1823. Sks.p
 7744 bractata Hook. many-spiked £ El or ■■■■■ W S. Amer. 1825. Sk s.p
 7745 coarctata Gill. bractod £ El or Jnjl ■■■■■ W. Indies 1824. Sks.p
 7746 compressa Gill. straitened £ El or jnjl ■■■■■ Chile 1823. Sks.p
 7747 gracilis Gill. compressed £ El or [njl ■■■■■ Chile 1823. Sks.p
 7748 angustifolia Swz. slender £ El or [njl ■■■■■ Chile 18U3. Sks.p
 narrow-leaved £ El cu ijU' B W. Indies 1822. Sks.p

* 957. BILLBERGZrf Thun. BILLBEROIA. (J. G. Billberg, a Swedish botanist.) *Bromelidce. 6.*
 7749 iridifolia B. R. Iris-leaved £ El or 1 mr SBY R.Janciro 1825. Sks.p Bot reg. 1068
 Brom&lia tridifolia N. & M.
 7750 amoeba Lindl. pleasing £ El or 2 jn ■■■■■ T S. Amer. 1817. Sks.p Bot reg. 344
 Brom&lia pallida B. H. Tillandsia amw^ia B. C.
 7751 pyramidalis Lindl. pyramidal £ El or 2 frar L R. Janeiro 1817. Skr.m Bot.reg.203
 Bromclia pyramidalis B. M. nudicafni B. R.
 7752 nudicaulis Lindl. naked-stemmed £ El or S apjl C Trinidad 1822. Skr.m Hook.ex. fl. 143
 g. Brom'lfa nudicaulis Hook.
 7753 clavata Lindl. club-shaped £ C51 or limy B Trinidad 1824. Sk r.m Bot. reg. 766
 BromMm melanantha B. R.
 7754 zebrina Lindl. zebxe^streaked £ E3 or Ilia M. S. Amer. 1820. Sk r.m Bot mag. 9686
 Brom&ua zebra B. M.

958. CARAGUA'TA Plu. CARAGUATA. (Its name in South America.) *Bromeliacea. 1.*
 7755 lingulataLindl. tongue-leaved ■■■■■ 2 jnjl Y Jamaica 1795. Sks.p Jac. am. 82
 Tillandsia lingulata Jac.

* 959. PONTEDE'R/^ L. PONTBDERIA. (*Julius Pontedera*, prof. bot. at Padua.) *Ponledrcee. 6.—10.*
 756 azitca Kny. fine-blue ■■■■■ 1 jnjl B Jamaica 1824. O ■■■■■
 757 crassipes Mart. thick-pctioled & El or fs.o B Guiana 1825. ■■■■■ Mart. br. 4
 758 cordataL. heaxUeaved * A o 2 jn.au B N. Amer. 1759. ■■■■■ Bot mag. 1156
 759 angustifl.Mia PA. narrow-leaved ■ A or 2 jn.au B N. Amer. 1806. ■■■■■
 760 dilatata/faw. spreading £ fT^or 2 my B KIndies 1806. ■■■■■ Bot rep. 490
 761 lanceolataL. lanceolate * El or 2 au B N. Amer. 1815. ■■■■■ Bot cab. 613

960. H J E M A ' N T H U S L. BLOOD-FLOWER. (*Haima, blood, anthos, a flower; flowers.*) *Amart/lidceec.* 21.—
 762 coccineus L. scarlet tf Alor au.o R C G. H. 1629. O r.m Bot mag. 1075
 763 coarctatus Jac. straitened tf Alor f.mr Pk C G. H. 1795. O r.m Bot reg. 181
 764 tigrinus Jac. tiger-spotted tf Alor f.d F C G. H. 1790. O r.m Bot mug. 1705
 765 rncalutus Jac. ByotteA-U-aved tf Alor ap.au C. G. H. 179a O s.l.p
 766 quadrivalvus Jac. tour.valved tf Alor 8.0 F' C. G. H. 1790. O r.m Bot. mag. 1523
 767 crassipes Jac. thick-pctioled tf Alor ■■■■■ R C. G. H. 1820. O 1
 768 hydrocarpus Jac. glass-fruited tf Alor ■■■■■ R C. G. H. 1822. O s.l.p Jacsc. 409
 769 rotundifolius B.M. round-leaved tf IAI or ■■■■■ S C. G. H. 1790. O s.l.p Bot mag. 1618
 770 orbicularis Fis. orbicular tf IAI or jn.o W C G. H. 1820. ■■■■■
 771 lanceolataL. scarlet tf LAJ or jnjl S C. G. H. 1722. O r.m Bot mag. 1315
 W*htunXH>Jae. low tf ^ or my.B S C. G. H. 1825. O r.m
 772 pumilio Jac. dwarf tf fAIor s.n Pk C. G. H. 1789. O s.l.p Jac. sc. 1
 773 pub^tceens L. pubescent tf fAIor a.us W C G. H. 1774. O r.m Bot.rab.703
 774 albiflos W. white-flowered tf fAIor au W C G. H. 1791. O r.m Bot. mag. 1239
 775 multiflorus L. many-flowered tf fAIor ap.au S S. Leone 178a O r.m Bot mag. 961
 776 lancifl.Mia Jac. siifar-leaved tf fAIor my.s C G. H. 1794. O r.m Jac. sc. 1.60
 777 cann'tus W. kcul-lcaved tf fAIor s.o Pk C G. H. 1759. O r.m
 778 sanguinicus L. bloody tf fAIor a.us C C G. H. 1820. O r.m
 779 mosciatus Jac. mut,kscnted tf fAIor JU R C G. H. 1816. O r.m
 780 amaryllisJac. Amaryllis-like tf fAIor a.us Pk C G. H. 1825. O r.m
 781 amarylloWtu Jac. Aestucolorcd tf fAIor f a.us Pk C G. H. 1819. O s.l.p Botreg.509
 782 carnus Ker Aestucolorcd tf fAIor I jnjl

783 rnfalls L. snow (Gala, milk, anthos, a flower; color.) *Amarylidcar. 2.*
 784 plicatus Bieb. plaited A or } ja.rar W Britain mca. O co Eng. bot. 19
 A or ija.ap W Crimea 1818. O co Bot reg. 545

* 962. LEUCOVUM L. SNOW-FLAKE. (*Leukos, white, ion, a violet*) *Amaryllidccc. 5.-7.*
 785 vernum L. spring A or Jap.mr W Germany 1596. O s.l Botmag.46
 786 carriaghicum B. M. Carpathian A or Jap.ap W Casp.M 1816. O s.l Bot mag. 173
 787 trichophyllum Brot. hair-leaved A or ija.f W Barbary 1820. O s.l Bot. reg. 544
 788 n^>fl&rum Red summer tf A or II ap.my W England m.mc. O s.l Eng. bot. 6121

2 m&jus <i>Herb.</i>	larger	tf E1 or	3 il.au	R.w	S. Ainer.	1789.	O r.m
3 minus <i>Herb.</i>	smaller	tf ED or	H jl.au	R.T	S. Amur.	1789.	O r.m
4 viridifolium	green-leaved	5 123 or	3 jl.au	W	Demerara	1819.	O r.m
5 berbicense	Iterbice	tf EJor	2 jlau	W	Berbicc	1819.	O r.m
78&9 hybridum <i>B. M.</i>	hybrid	tf EM or	2 jl.au	Pk	hybrid	1820.	O r.m Bot mag. 2336
7860 pheatum <i>Hort.</i>	plaited	tf LAJor	2 jn.au	W	China	181&	O r.m
* 965. CYRTA'NTHUS H. K. CYRTANTHII'S.	(Kttrtos, curved, anthos, a flower.)	Amaryllidee.	8.—10.				
7861. anRustiblius II. K.	narrow-leaved	tf tAJor	1 my.jn	O	C. G. H.	1774.	O r.m Bot. mag. 271
7862. collinus B. R.	lull	tf ^J or	% my.au	G	C. G. H.	1816.	O r.m Hot. reg. 162
7863 spirilis B. R.	spiraWcatvd	tf iAl or	1 my.au	S	C. G. H.	17XX.	O r.m Bot. reg. 167
78W ventricibus W.	bellied	tf iA] or	1 my.au	R?	C. G. H.	1770.	O r.m Jac. sc. 1.76
78W obnqus H. K.	ohWque-leaved	tf iAl or	2 my.au	G.o	C. G. H.	1724.	O r.m Bot. mag. 1133
7866 stratus <i>Herb.</i>	striated	tf IAJ or	\$ jl	O	C. G. H.	...	O r.m Bot. mag. 25J4
7K7 odorus AT	sweet-scented	tf iAl or	i my.il	C	C. G. H.	1818.	O r.m Bot. reg. 503
7868 pallidus Sims	pale	tf LAI or	1 my.jl	Pk	C. G. H.	1822.	O r.m Bot. mag. 2471
966. GASTRONEWA <i>Herb.</i> GASTRONEMA. (Gaster, a belly, ncma, a filament)	Amarylfxdete.	1.					
7869 clavatum <i>Herb.</i>	clavate	tf ■■■	\ my.jn	W.a	C. G. H.	181&	O r.m Bot reg. 168
Cyrtanthus uniflrdus B. R.							
* 967. BRUNSVI'G/J Hei. BRUNSVICIA.	(Hie noble family of Brunswick.)	Amaryllidee.	12.				
7870 lificiria <i>Herb.</i>	shining	tf LAJor	1 au	Pk	C. G. H.	1818.	O r.m Botrcg^J?
AmaryMij latfcomia B.R. Nertnc latfcomia Ker	smaller	tf iAJ br	jl.au	Pk	C. G. H.	1822.	R s.l Botreg.a^i4
7871 minor Lindl.			1 jn.au	S	C. G. H.	1814.	O nn Red. HL 370.372
7872 Josephine? R. L.	Josephine's	tf iAl or	1 jn.au	S	C. G. H.	1814.	O r.m Bot. reg. 192.193
2 minor B. R.	smaller	tf iAJ or	1 jn.au	S	C. G. H.	1823.	O r.m Bot. mag. >78
3 stnara R AT.	striated	tf iAJ or	1 jn.nu	S	C. G. H.	1752.	O r.m Bot. mag. UH9
7873 multifloM Hei.	many-flowered	tf iAJ or	1 jn.au	R	C. G. H.	1752.	O r.m Bot. mag. UH9
Amarylhf nrientMis L							
7874 marginMa H. K.	raZ-margined	tf iAl or	1 s.o	S	C. G. H.	1795.	O r.m Jac. sc. 1.65
7875 radula Jnc.	rasulcaved	tf iAl or	ap.au	R	C. G. H.	1790.	O r.m Jac. sc. 1.68
7876 striata Jac.	striated	tf iAJ or	i s.o	Pk	C. G. H.	1795.	O r.m Jac. sc. 1.70
7877 falckta Ker	sickle-lcavcd	tf iAl or	f my.jn	R	C. G. H.	1774.	O r.m Bot. inag. 1443
Ammocharis falctn Herb.							
7878 toxicchia Ker	*poison-&if%	tf iAJ P	1 s.o	Pk	C. G. H.	1774.	O rjn Bot. reg. 567
Hsemuiithid toxicarius L.							
7879 coránica Kcr	Coranic Poison-bulbj	iAJ P	1 8.0	Pk	C. G. H.	1815.	O r.m Bot reg. 159
Ainm6chrnis cordnica Herb.							
7880 ciliiris Ker	fringed	tf iAJ or	1 jl.o	Bk	C. G. H.	1752.	O r.m Brey. c. 39
HwniHithus ciU^ris IV.							
7881 dltidlii Swt.	two-rowed	tf LAJ or	..	It?	C. O. If.	1823.	O r.m PaLitl
Ainar^lli* dlsticha Pat.	Bbphanc disticha Kcr						
968. NERPNA <i>Herb.</i>	NEHINE.	(Nerine, the daughter of Nereus.)					
7882 curvifolia <i>Herb.</i>	curve-leaved	■ iAJ or	1 my.s	P	C. G. H.	1777.	O r.m Bot mag. 725
Amarillis curvifolia Jac.							
7883 con'isca <i>Herb.</i>	glittering	tf iAJ or	1 jl.au	S	C. G. H.	1809.	O r.m Bot. mag. 1069
7884 sarmYnsis Kcr	Guernsey Lily	tf iAl or	1 s.o	K	Jaijan KK9.	O r.m Bot. mag. 204	
7885 venista <i>Herb.</i>	lieautiful	tf LAI or	1 jn.jl	S	C. G. H.	17XX.	O r.m Bot.mag.10UO
7886 flexioislaHerb.	flexuous	tf LAJor	1 8.0	Pk	C. G. H.	17XX.	O r.m Botreg. 172
7887 hfmihis <i>Herb.</i>	low	tf LAJor	2 jn.jl	R	C. G. H.	17<5.	O r.m Bot. inag. 7^6
7888 unduKita <i>Herb.</i>	waved-flowered	tf LAI or	I my.jn	Pk	V. d. H.	17..7.	O r.m Bot. mag. &M
7889 aurca Swt.	golden	tf LAI or	1 aus.s	Y	China	1777.	O r.m Bot. mag. 409
Lycorus aurca <i>Herb.</i>							
7890 radiata	radiated	tf laJor	1* jn.jl	Pk	China	1758.	O r.m Bot rep. 95
Lycorus radiate <i>Herb.</i>							
7891 röbca Swt.	roseate	tf iAI or		Pk	C. G. H.	1818.	O r.m Bot mag. 2124
7892 pulchella <i>Herb.</i>	pretty	tf LAJor		Pk	C. G. H.	1820.	O un Bot. m jg. 241/7
7893 versicolor <i>Herb.</i>	various-colored	tf iAl or	I JL0	Va	hybrid	1815.	O r'm Herb. ap. it*
* 969. AMARY'LLISK AMARYLLIS.							
7894 pumilio Spin	dwarf	tf ■■■ n	Pk	C. G. H.	1774.	O F.m	
7895 piunca Ker	modcbt	i my.jl	Pk	C\ G. H.	17V>	O nn Kerrov.8 R	
7896 blanda Ker	charming	tf H my.jn	Pk	C. G. H.	1704.	O r.m Bot. mag. 1450	
7897 Belladonna L.	Belladonna Lily	tf 2 jl.s	F	W. Indies	1712.	O p.l Bot mag. 733	
2 pálhda	pale-flowered	tf 2 jl.s	F	Madeira	...	O pi Red. ll.470	
7898 pallidn <i>Herb.</i>	pale	tf 1 jl.au	Pa.R	V. G. II.	1777.	O r.in Bot reg. 714	
7899 reticulata H. K.	netted-veined	tf 1 apiny	P	Brazil	1777.	O r.in But. leg. G57	
7KX HtriatiPolia Swt.	channeled-lvd	EJor	1 ap.my	P	Brazil	1815.	O r.m
I'm brevifMia Swt.	short-leaved	fAlor	2 ap.my	R	hybrid	1S21.	O r.m
JUC suK-rbe Swt.	superb	I AJor	1 ap.jn	St	hybrid	181L.	O rm
TM grandiflora Swt.	large-flowered	IZSor	2 my.jn	St	hybrid	1821.	O r.m
4U4 atrortibeni Swt.	dark, red	USlox	1 my.jn	R	hybrid	1821.	O r.in
7WQ Qlnhra Swt.	obscure	2.3 or	1 my.jn	St	hybrid	1821.	O r.m
AW intermedia Lindl.	intermediate	[SI or	1 my.jn	St	Brazil	1821.	O r.in
7o7 sulcMa Swt.	furrowed	IS]	1 np.jn	St	hybrid	1821.	O r.m
7XJ recurred			1 ap.jn	St	hybrid	18*1.	O r.m
Wells's			1 Jjlau	St	hybrid	1821.	O r.m
Vallet's			1 year	P	hybrid	1821.	O r.m
7JL ruw scens Swt.	rul)escent	If ap.my	R	hybrid	1821.	O r.m	
7/2 purpur&scens Swt.	purpurascnt	2 year	P	hybrid	1821.	O r.m	
7913. p^ten8 Swt.	Hreading	1 year	Va	hybrid	18*J1.	O r.m	
7914. vencolor Swt.	various-colored	1 year	Va	hybrid	1821.	O r.m	
7915 costata Swt.	ribbed	1 year	St	hybrid	1821.	O r.m	
7916 dioica Swt.	dio?cious	1 year	R-p	hybrid	1821.	O r.m	
7917 amce na Swt.	pleading	1 year	R	hybrid	1821.	O r.m	
7D18 rugosa Swt.	rugose	1 year	R	hybrid	1821.	O r.m	
7D19 def.ra Swt.	comely	1 my.au	St	hybrid	1821.	O r.m	
7D20 pale	1 jn.jl	Pa.O	hybrid	1821.	O r.m		
7D21 variegated	1 year	P.R	hybrid	1821.	O r.m		
7D22 nerve-icaved	1 year	St	hybrid	1821.	O r.m		
Johnson's	2 my.jn	St	hybrid	1810.	O r.m		
Anderson's	1 jnjl	R	hybrid	1821.	O r.m		

HEXANDRIA MONOGYNYIA.

CLASS VI.

7925	Colvilia Swt.	tf	El or	1*	Jn,jl	R.P	hybrid	1821.	O	r.m	
7926	unduliflora Swt.	tf	El or	1	year	P.w	hybrid	1821.	O	r.m	
7927	palliflora Swt.	tf	El or	2	year	R.w	hybrid	1821.	O	r.m	
7928	discolor Swt.	tf	El or	1	year	St	hybrid	1821.	O	r.m	
7929	expansa Swt.	tf	EJ or	1	year	D.E	hybrid	1821.	O	r.m	
7930	patentibima Swt.	tf	El or	1	au.s	C	hybrid	1821.	O	r.m	
7931	bractata Swt.	tf	El or	1	year	R	hybrid	1821.	O	r.m	
7932	Annesleyana Swt.	tf	El or	1	my,jl	P	hybrid	1823.	O	r.m	
7933	striatiflora Swt.	tf	El or	1*	year	St	hybrid	1821.	O	r.m	
7934	consobrina Art.	tf	El or	1	my,jn	Pk	hybrid	1823.	O	r.m	
7935	phoenicea Swt.	tf	El or	1	year	U.my,jl	P	hybrid	1821.	O	r.m
7936	sternantha Swt.	tf	El or	2	my,jn	R	hybrid	1821.	O	r.m	
7937	reginae L.	2	my,jl	S	America			1725.	O	r.m	
	pulverulenta Lod.	tf	El or	If	my,jn	S	hybrid	1820.	O	r.m	
	3 vittoa // Tr.	tf	El or	If	my,jn	K	hybrid	1820.	O	r.m	
7938	rubricunda Swf.	tf	El or	1	year	P.	hybrid	1823.	O	r.m	
7939	pallescens Swt.	tf	El or	1	year	D.P	hybrid	1821.	O	r.m	
7940	accidens Surf.	tf	El or	1\	year	hybrid		1821.	O	r.m	
7941	coccinea Swt.	tf	El or	2	year	S	hybrid	1821.	O	r.m	
7942	ringens Swt.	tf	El or	1	year	K.w	hybrid	1821.	O	r.m	
7943	carnescens Surf.	tf	El or	1	year	P.w	hybrid	1821.	O	r.m	
7944	rusea?iba Swt.	[AT or	El or	2	year	R.w	hybrid	1821.	O	r.m	
7945	spathacea Stu.	tf	El or	If	year	11?	hybrid	1821.	O	r.m	
7946	imbuta Swt.	tf	El or	2	year	R	hybrid	1821.	O	r.m	
7947	compacta Surf.	tf	El or	1	year	R	hybrid	1821.	O	r.m	
7948	consanguinea Sw.	tf	El or	1	year	O	hybrid	1821.	O	r.m	
7949	principis Salm	tf	El or	1	jn,jl	Brazil		1823.	O	r.m	
7950	vittata // K.	Landed	I.AJ or	1	ap,my	St	C. G. H.	1769.	O	r.m	
	2 major Lnf//.			2	my,jl	St	C. G. H.	1774k	O	r.m	
7951	lineata Q.	lined	tf IAJ or	\	ap,au	R.w	'S.Amer.	1820.	O	r.m	
	vittata y Harrison*	B, R.									
7952	p&chra Swt.	fair		\	ap,au	P.w	hybrid	182a	O	r.m	
7953	variabilis Swt.	variable	tf	I.A or	1	ap,au	R.w	hybrid	1823.	O	r.m
7954	Forbesii Lindl.	Forbes's	tf	.Alsol	I.Jl.au	PkW	C. G. H.	1823.	O	s.l	
	2 purpurea LindL	purple	tf	I.Jl	If	jlau	P	C. G. H.	1823.	O	s.l
7955	vittifera Swt.	riband-bearing	tf	IAlor	Ilfmy,jl	S	Chile	1807.	O	r.m	
7956	advena Ker	stranger	tf	IAlor	* my,jn	P.v	Chile	1821.	O	r.m	
	2 cerina Lindl.	wax-colored	tf	IAlor	S my,jn	S	Brazil	1815.	O	r.m	
7957	rutila Ker.	fiery-red	tf	IAlor	I ap,my	S	hybrid	1816.	O	r.m	
7958	vittata Hort.	banded	tf	IAlor	1	ap,my	S	hybrid	1820.	O	r.m
7959	spillndens Herb.	shining	tf	IAlor	1	my,jn	S	hybrid	1822.	O	r.m
7960	attenuata Swt.	attenuated	tf	IAlor	1	year	Pa.R	hybrid	1822.	O	r.m
7961	punctata Swt.	dotted	tf	IAlor	1	year	Pa.It	hybrid	1822.	O	r.m
7962	fuliginea Ker	fulgid	tf	IAlor	% ap,my	Ve	Brazil	1810.	O	r.m	
	2 vittata Hort.	banded	tf	IAlor	f ap,my	V	hybrid	1820.	O	r.m	
7963	subbarbata B. M.	slightly bearded	tf	E or	1	year	Va	Brazil	1833.	O	r.m
7964	mutabilis Swt.	changeable	tf	E or	1	year	R.Saf	hybrid	1822.	O	r.m
7965	rubro-crocea Swt	reddish saffron	tf	EAl or	1	year	Cop	hybrid	1822.	O	r.m
7966	cuprea Swt.	copper	tf	E or	1	year	Ve	Brazil	1815.	O	r.m
7967	crockta Ker	saffron-colored	tf	E or	1	ap,my	Ve	Brazil	1815.	O	r.m
	2 regime Hort.	queen's	tf	E or	1	my,jn	S	hybrid	1810.	O	r.m
7968	stylifera B. M.	stylbse	tf	EAl or	1	my,jn	R	Maranhao	1821.	O	r.m
	marañensis B. R.										
7969	pulvrcultenta Herb.	powdery	tf	EAl or	1	ap,my	O	Brazil	1819.	O	r.m
7970	venusa Swt.	veiny	tf	I AJ or	1	jn,o	St	hybrid	1822.	O	r.m
7971	latcritia Swt.	brick-co/ored	tf	I Al or	\	jn,o	Bri	hybrid	1822.	O	r.m
7972	fulva Swt.	tawny	tf	I AJ or	1	jn,o	Br	hybrid	1822.	Cv	r.m
7973	glaucocentus Swt.	glaucous	tf	LAJO*	1	jn,o	R	hybrid	1822.	O	r.m
7974	Hoodia Swt.	Hood's	tf	IAlor	1	jao	S.w	hybrid	1822.	O	r.m
7975	macrantha Swt.	long-flowered	tf	IAlor	1	jn,o	R.R	hybrid	1822.	O	r.m
7976	breviflora Swt.	short-flowercd	tf	IAlor	1	jn,o	R	hybrid	1822.	O	r.m
7977	cleftata Swt.	dehifflful	tf	IAlor	f	jn,o	D.11	hybrid	1822.	O	r.m
7978	aiKibiUs Atta	lovely	tf	I AJ or	11	jn,o	Va	hybrid	1822.	O	r.m
7979	principes Atta	princely	tf	I AJ or	1	jn,o	S	hybrid	1822.	O	r.m
7980	Imperialis Surf.	imperial	tf	I Al or	1	jn,o	S	hybrid	ISM.	O	r.m
7981	augusta Surf.	august	tf	EAl or	2	jn,o	S	hybrid	1822.	O	r.m
7982	ensipiflora Swt.	curled.flowercd	tf	EAl or	2	my,jl	S	hybrid	1822.	O	r.m
7983	tortuliflora Swt.	twibted-flwd	tf	EAl or	2	my,jl	S	hybrid	1822.	O	r.m
7984	igneaens Swt.	ignescent	tf	EAl or	1	my,jl	Fla	hybrid	1822.	O	r.m
7985	tricolor Swt.	three-colored	tf	EAl or	2	my,jl	Cop	hybrid	1822.	O	r.m
7986	equestris Atte.	equestrian	tf	EAl or	1	jl,o	S	W. Indies	1710.	O	r.m
	2 major	larger	tf	EAl or	\	jl,o	S	W. Indies	1710.	O	r.m
	3 plena	double	tf	EAl or	If	my,s	S	W. Indies	1801.	O	r.m
7987	miniata 71% P.	vermilion	tf	EAl or	1	jn,jl	F	Peiu	1825.	O	r.m
		flame	tf	EAl or	1	ap	Fla	Peru	1825.	O	r.m
			tf	EAl or	P.v	S. Amer.		Fl. per. 3.286.6			
7988	solandrayiflora Lindl.	red-clianched	tf	EAl or	If ap,jn	R.w	S. Amer.	1820.	O	r.m	
	2 rubro-striata	purpurascens	tf	EAl or	If ap,jn	P.w	S. Amer.	1820.	O	r.m	
7989	3 purpurascens	banded	tf	EAl or	2	jl,au	Va	hybrid	1819.	O	r.m
7990	vittata Lindl.	calyptrata	tf	EAl or	If my,au	G	Brazil	1816.	O	r.m	
7991	calyptrata Ker	courty	tf	EAl or	If my,au	G.s	Brazil	1810.	O	r.m	
7992	auhca Ker	broad-putaled	tf	EAl or	If my,au	O	Brazil	1824.	O	r.m	
	2 platyptala B. JR.	parrot	tf	EAl or	If my,au	G.s	Brazil	1816.	O	r.m	
7993	psittacina Ker	netted	tf	EAl or	If ap,jn	S	W. Indies	1822.	O	r.m	
7994	retinervia Swt.	twisted	tf	EAl or	n. jn,o	G.s	hybrid	1822.	O	r.m	
7995	tortuosa Swt.	four-colored	tf	EAl or	U	jn,o	St	hybrid	1822.	O	r.m
7996	quadricolor Swt.	many-channelled	tf	EAl or	11	jn,o	St	hybrid	1822.	O	r.m
7997	multistrata Swt.	channeled	tf	EAl or	If jn,o	S	hybrid	1822.	O	r.m	
7998	canaliculata Swt.	recurved-flwd	tf	EAl or	If jn,o	R	hybrid	1822.	O	r.m	
7999	recurviflora Swt.	six-spotted	tf	EAl or	If jn,o	Cop	hybrid	1822.	O	rnt	
8000	sexmaculata Swt.	campanula tc	tf	EAl or	2	my,jl	BB	hybrid	1822.	O	r.m
8001	multistrata Swt.	net-flowcrd	tf	EAl or	f my,au	U.II	N. Amer.	1658.	O	r.m	
8002	rticiflora Swt.	handsomest	tf	EAl or	1	my	R?	Mexico	1825.	O	r.m
8003	formoissima L.	ncrvose	tf	EAl or	f my	B	Siberia	1822.	O	r.m	
8004	ncrydsia Hum.	Tartarian	tf	EAl or	1	my					
«005	Utarica PaU.										

MissChapm. ic.

Bot mag. 47

- * 970. PHYCE'LLA Lindl. PHYCELLA. (Dim. *otphykos*, red alkanet; color.) *AmarylUdea*. a
 8006 *fgnea Lindl.* fiery JE J or 1 n S Chile 1824. O r.m Bot reg. 809
Amaryllis fgnea Lindl.
- 8007 *cyrtantholdes Lindl.* Cyrtanthus-lk V (29 or f Jn C Chile 1824. O r.m Bot mag. 2399
Amaryllis Uicrytantholdes B. M.
- 8008 *corusca Lindl.* glittering tfEJor 1 • S Coquimbo 1825. O i.p.l
971. VALLOTif Sal. VALLOTA. (*Pierre Pallet*, a French botanist) *Amaryllidae*. \
 8009 *purpurea Herb.* purple tf iAJ or 1\$ my.jn S C G. H. 1774. O p.s Bot. reg. 553
Amaryllis purpurea H. K.
 2 minor Herb. smaller tf iAJ or 1 my.jn S C. G. H. 1774. O p.i Bot mag. 1430
972. GRIFFI'N/if Ker GRIFFINIA. (*William Griffin*, Esq., of South Lambeth.) *Amaryllideaf.* &
 8010 *hyacinthina It. Br.* violet-colored tf E or 1 in.s B S. Amer. 1815. O r.m Bot reg. 163
 8011. *intermedia Lindl.* intermediate 9 (23 or fmpr.ap B Brazil 1815. O p.l Bot reg. 990
 8012 *pariflora Ker* small-flowered ? [Z2or | jns Pa.P S. Amer. 1815. O r.m Bot reg. 511
973. STERNBERG/ RG/^ Kit STKRNBERQIA. (*Count Caspar Sternberg*, a celeb, bot) *Aman/Uide**. 4.—6.
 8013 *colchiciflra W. 8c K.* Coichicum-flwd tf A or I aus. Y Hungary 1816. O r.m IV. & K. 2.157
 8014 *ClixBlana Ker* Eluse's ↑ A or *aus. Pa.Y Constant. -- O r.m Cl. h. 1.163
 8015 *lutea Ker* yellow A or | aus. Y S. Europe 1596. O r.m Bot. mag. 290
 2 angustifolia narrow-leaved A or 4 au Y S. Europe 1595. O co
 8016 *cixigua Ker* small tf A or * aus. Y N. Africa 1820. O r.m
- * 974. ZEPHYR'A'NTHES Herb. 2EFIIYRANTHES. (*Zephyros*, west wind, *anthe*, flower.) *Amarylldcaf*. 9.
 8017 *tubiflatha Herb.* tube-sputhe tf QS or i my.jl W S. Amer. ... O r.m Bot mag. 1586
Amaryllis tubiflatha Hent.
- 8018 *Candida B. M.* white tf A or I 8 W Peru 1822. O r.m Bot reg. 724
Ainarylliic Candida B. R.
- 8019 *Atamasco Herb.* Atamasco Lily tf A or i my.jn W N. Amer. 1629. O r.m Bot mag. 239
Amaryllifl Atamasco L.
- 8020 *grandiflbra Lindl.* large-flowered tf AJ pr I my.jn Pk Mexico 1824. O 81 Bot reg. 902
 8021 *carinata B. M.* keeled tf AJ pr I my.jn Pk Mexico 1824. O s.I Bot mag. 2594
 8022 *vereunda Herb.* blushing tf AJ pr I ap.my Pa.It Mexico 1824. O s.I Bot mag. 2583
 8023 *striata Herb.* channelled tf Al pr I ap.my W Mexico 1824. O s.I Bot mag. 2593
 8024 *rifosa Lindl.* rosy tf AJ or I my.jn It Havannahl823. O r.m Bot. reg. 821
 8025 *chloroleuca Herb.* greenish-white tf A or imy.au Pa.G O r.m Ker rev. 8.2
- * 975. HABRA'NTHUS Herb. HADRANTHUS. (*Habros*, delicate, *anihos*, flower.) *Amarylldca?*. 6.
 8026 *versicolor Herb.* changeable ttf A to I Pk S. Amer. 1821. O r.m Bot mag. 2485
 8027 *loriflthus B. M.* strap-leaved tf AJor 4-11 S. Amer. 1821. O r.m
 8028 *graciliflbus Herb.* slender-leaved tf A'' W S. Amer. 1821. O r.m Bot. mag. 2464
 8029 *anftubus Herb.* narrow tf A'pr \s R Brazil 1822. O p.l Bot. mag. 2639
 8030 *bifidum Herb.* two-cleft tf AJ or 5 jn Pk B. Ayres 1823. O 81 Bot mag. 2597
 8031 *spathaceus Herb.* spatheaceous tf Al ij - B. Ayres 1823. O r.m
- 97a DORYA'NTHES Corr. DORYANTHES. (Dory, spear, *anthe*, flower; stem.) *Amaryllidea?*. 1.
 8032 *excelsa Corr.* gigantic tf. iAi or 20 jl.au Crea N. S. W. 1800. Sk s.p Bot mag. 1685
977. GETHY'LLIS Z. GETHYLIS. (*Getheo*, to rejoice; perfume of flowers.) *Amaryllidete*. 5.
 8033 *spiralis L.* *pvrj-leaved I kJ or % jij,jj | WV C. G. H. 1780. O s.p Bot. mag. 1088
 8034 *ciliaris L.* fringed I kJ C. G. H. 1788. O s.p Jac. ac. 1.79
 8035 *villbsa L.* villous I kJ C. G. H. 1787. O s.p
 8036 *lanceolata L.* siic&r-leaved I kJ C. G. H. 1790. O s.p
 8037 *Afra B. B.* African I kJ C. G. H. 1820. O e.p Bot. reg. 1016
978. POLIA'NTHES L. TUBEROSE. (Polis, city, *anthe*, flower.) *Hemcrocallidecc*. S.
 8038 *tuberosa L.* tuberous A iAJ or 3 aus. W E. Indies 1629. O r.m Bot. reg. 63
 2 Mure pleno double A iAi or 3 aus. W O r.m
 8039 *gracilis Lk.* slender AIZSor 3 aus. Pa.Y Brazil 1822. O r.m
- * 979. ALSTR(EMEPR/14 L. Pelegrina. (Baton C. Alstremer, a Swed. bot) *Amaryllitiear*. 10.—33.
 8040 *Pelegrina L.* Pelegrina or 1 Jna St Peru 1753. 11 r.m Hot inng. ISO
 8041 *Littifl F** Littifl or 1 finr S Peru 1778. It 1>p Hot maf. H
 8042 *FIOH Martini Ker* St. Martin* fl. Ijin w.p.y Chile 1822. S L

Bot. reg. 731
 8043 *pulchella Sims* pretty A E 3 or 3 in 8 Chile 1822. S L

Hook ex. fl. J/H
 8044 *roseata Hook.* roseate A ES or 1 Jn.jl Pk Peru 1824. R L

Hook ex. fl. 181
 8045 *SalsUla L. Sal* sill a 4123 clt 6 jnjil G.c S. Amer. 18U6. R l.s.p Bot mag. 1613
 8046 *ovata Cav.* ovate ** A or 4 jnjil RUY Chile 1824. It Ls.p Cav. ic. 1.7G
 8047 *eclulis J.* eatable IZQ 4 jl.au R Trinidad 1820. R l.s.p Fl. ant. 14
 8048 *pauciflura Kth.* few-flowered 4 C3 or O.u Caracas 1823. 11 l&p
 8049 *lurtdla Kth.* rather hairy J A or 4 jl H.Y.G Mexico 1824. S p.l Sw. fl. gar. 228

* 80. CONANTHEFRA Fl. per. CONANTIERA. (*Konos*, a cone, *anthera*, an anther.) *Asphodhlca*. 3.—5.
 v8050 *campanulata//w/fc.* beU-flowcrat J J or I mt B Chile 1823. O Up Hook. ex. fl. 214
 8051 *bifolia Fl. ptr* two-leaved tf i-mr TAB Chile 1823. O "sd" Bot cab. 904
 8052 *Slmsii Swt* Sims's tf I mr.my B Chile 1823. O r.m Bot mag. 241%
 bifbliaifeA/.notF/.jw.

MI- HYPO'XISZ.
 8053 *junccea Sm.* HYPOXIS. (Hypo, beneath, oxys, sharp; base of capsule.) *Hypoxlfia?* 22.—2ft
 XK hygrometrica Lab hygrometric A'or Y Carolina 1787. O l.p Sm. epic. 15. lb
 «MS ilba L. white iAJ or jnjil Y N. Holl. 1820. O l.p Lab. n. h. 1.8
 Kd*i obtifa B. R.* obtuse tf tAlor i'n Y C. G. H. 1806. O p.i Jac. c. 4.2.1
 8107 *linoiris // Jtcp.* star-flowercd tf iAJ or i ap.my Y C. G. H. 1816. O p.i Bot. reg. 159
 W.->H ifellata L. elegant II".Alor I apjn W.D C. G. H. 1792. O Lp Bot rep. 171
 8079 *elegans Andr.* G&w-leaved tf tAlor I apjn W.B C. G. H. 1752. O s.p Bot mag. (iti2
 8060 *serifata L.* Bshort-scaped tf iAlor \ my.jn Y C. G. H. 1788. O l.p Bot mag. 709
 8111 *breviflspa Kth.* ovate tf iAlor Y Brazil 1823. O p.l
 8012 *ovata W.* Veratrum.Ivd A or Y C. G. H. 1806. O e.p Bot mag. 1010
 "03 veratriflbia Is. Carolina tf iAlor i jnjil Y Carolina 1822. O Lp
 8013 *caroliensis Mt.* decumbent tf tAlor Jjn.s Y Jamaica 1755. O p.l Mil. i c \ 3i£
 Tmp. stell pilis Ker starry-haired tf iAlor 1 jl Y C. G. H. 1821. O s.p Bot reg. 663
 2HS g. obtiqua Jac. oblique-leaved tf iAlor i jnjil Y C. G. H. 1795. O p.l Bot rep. 195
 S > n "s R. Br. meadow 5 iAlor i mr.jn Y N. Holl. 1824. O p.l
 bra Lod. rough "ff^Jor i aus. Y C. G. H. 182a R p.l Bot cab. 970

8170 erecta L.	upright	£ A or ft J ⁿ i ^l	Y	N. Amer.	1752.	O p.l	Bot. mag. 7J0
8071 Bobolifcra Jac.	shookbearing	giAJor	Jjn.s	Y	C. G. H.	1771.	O p.l
8072 villosa L.	villous	g^tAJ or	fjn.s	Y	C. G. H.	1774.	O pi
8073 aquatica L.	aquatic	j^Aor	ijn.jl	Y	C. G. H.	1787.	O p.l
983. CURCITLIGO	Gae. CuacLiao.	<i>(Curculio, a weevil; seeds.)</i>			Hypoxfdeec.	6.-9.	
8074 Sumatr'na Rox.	Sumatran	£ El or	3 jl	Y	Sumatra	1818.	O 1.p
8075 phcata Z)m	plaited-foivat	£ iAJ or	1f jn.jl	Y	C. G. H.	1788.	O 1.p
2 glabra	smooth	* IAJ or	Ij jn.jl	Y	C. G. H.	17H8	O 1.p
8076 breviflora Rox.	Orchis-like	£ [A3 or	J jn.jl	Y	E. Indies	1800.	O 1.p
8077 breviflora Dry.	short-leaved	jg E2 or }	my.jl	Y	E. Indies	1804.	O 1.p
8078 latiflora Dr.	broad-leaved	£ K3 or	1f my.au	Y	Poole Pin.	1804.	O 1.p
8079 recurvata Dry.	recurved-/caww/£ OS	1 ...	Y	Bengal	1805.	O 1.p	
983. MOLINE^RM Coll. MOLINEHIA. <i>(Igititio Molinerio, direct, of bot gar. at Turin.)</i>					Hypoxidac.	1.	
8080 plicate Coll.	plaited	£ 123 or	1 au	S.Y	Java	182a	D p.l
984. BAMBLTSA Schr.	BAMHOO CANE. <i>(Sambos, the Indian name)</i>				Grammeee.	8.	
8181 flrmidinacea W.	Reed-like	spjlliQee40	- -	Ap	India	1730.	Ski Rox.cor. 1.79
8082 verticillata W.	v/horl-fjowcrec	IP cu 20	- -	Ap	India	180* Sk 1	
8083 spiniba Ham.	spin'y	dlili P or 20	- -	Ap	K Indies	1820.	Sk 1
8084 stricta Rox.	strict	jlbb • or 20	- -	Ap	R Indies	1824.	Sk 1,
8085 glauca ho. C.	glaucous	jlbb • or 20	- -	Ap	E. Indies	1H26* Sk 1	
8086 iirgra Lo. C.	black	jlbb l or 20	- -	Ap	R Indies	1825.	Sk 1
8087 aristata Lo. C.	awned	jlbb r-or 20	- -	Ap	E. Indies	1812.	Sk 1
8088 puWscens Lo. C.	pubescent	jlbb rnor 20	- -	Ap	E. Indies	1826.	Sk 1
985. MELOCA'NNA Tr'm. MELOCAXNA. <i>(Melon, apple, kanna, reed.)</i>					Graminete.	1.—2.	
8089 bamhuso Vcs, Trin.	Bambusa-like	Xk • or 20	... Ap	E. Indies	1818.	Sk 1	
Jiumbusa baccitera Rox.							
986. NA'&TUS J.	NASTUS. <i>(Nattos, the Greek name for a kind of reed)</i>				Graminete.	4.	
8090 latiflora Spr.	broad-leaved	Jb • or 20.			Cumana	1818.	Sks.p
Bambusa latifolia H. \$ li.							
987. CA'LAMUS L.	CALAMUS. <i>(Kalam, a reed, Arab.)</i>				Pdtoue.	6.—13.	
8091 verus JJOU.	true	ce 20	... G	Cochinch.	1824.	S s Ru.am. 5.54	
mvn nlger W.	black	ce 20	G	Moluccas	1824.	s Ru.am. 5.54	
8093 riidcnuin Lou.	cable	ce 20	G	E Indies	1812.	S 81	
80L4 Zalatca Gae.	Zalat-ca	ce 20	G	f. Indies	1812.	S 81 Gae. fir. 2.139.2	
80U5albus/Vr».	white	ce 20	G	E. Indies	1812.	S r.m Ru. am. 5.52	
8096 draco FK	dragon	ce 20	G	E. Indies	1819.	S r.in»Ru. am. 5.58.1	
988. EHRHA'RIM Thun. EIIRHAIITA. <i>(F. Ehrhart, a Swiss naturalist)</i>					Graminccv.	4.—14.	
8077 piincca Sm.	Panic-grass	JUUMAICU	my.jl	Ap	C. U. H.	1790.	S co Sin. ic. in. 1.9
film distichiph'lla Lab.	distich-leaved	JUUTAJun	jnl	Ap	N. Holl.	1820.	1) r.m
mm gigantia Ehrh.	giant	JUATJUUI	jn.jl	Ap	C. G. H.	1818.	D s.p
8100 calycina Sm.	/	JBciAlun	jn.jl	Ap	C. G. H.	1818.	1) s.p Sm. ic. in. 1.33
Alta. capensis L.							
989. if'CORUSL.	ACORUS.	<i>(A, priv., kore, pupil of the eye; med. qual.) Arold ear. 3.</i>					
8101 Calamus L.	Sweet Flag	^ A >n 2 jn.jl	Ap	Britain	pools.	D m.s Eng. bot. 356	
8102 terrestris Spr.	terrestrial	£ A ^r 1 ju.jl	Ap	China	1822.	1) co	
8103 /rainlnicii // K.	Uras^jeaud	£ /^ cu if	Ap	China	179G.	D s.p Sm.epic. 15.	
990. ORO'NTIUMZ.	OnONTIUM.	<i>(A Greek name adopted by Linnaeus.)</i>			AroUte c	I. —4.	
8104 aquaticum L.	aquatic	■ A cu 4.jn	Ap	N. Amer.	1775.	D t.p Hook. ex. fl. 19	
991. RO%HDE4 Roth	ROHOEA.	<i>(Mich. Bohde, of Bremen, in Germany.)</i>			Aroideee.	1.	
8105 japonb-a Roth	Japan	^ A ce 2 ja.ap	Ap	Japan	1783.	D i p Bot mag. 898	
Or6ntium japonicum	Thun.						
• 992. TUPI'STRA Ker	TUPISTRA.	<i>(Dimin. of typis, a mallet; flower.)</i>			AroU&e.	1.	
8106 squalida Ker	squalid	£ QS cu 2 ap	Ld	Amboyna	1810.	R l.p Bot reg. 704	
993. TA'CCA Forst	TACCA.	<i>(Its Malay name.)</i>			Aroideee.	&	
8107 pinnatifida Forst.	pinnatifid	ec 2 ... P		R Indies	1793.	R l.p Bot cab. 692	
8108 >aspacra/fox.	rough	tf E cu 2 jl.au	Br	E. Indies	1816.	R l.p	
8109 iffVis Rox.	smooth	tf EScu 2 jl.au	Br	E. Indies	1820.	R l.p	
8110 guineensis G. Don	Guinea	tf E3cu 2 jl.au	Br	R Indies	1823.	R l.p	
8111 phalferia ?o?.	crest-bearing	tf G3cu 4 jlau	Br	Mauritius	1826.	R l.p	
8112 integriflora ACT	entire-leaved	ftmy.ii P		R Indies	1810.	Bot mag. 1488	
• 994. JU'NCUS L.	RUSH. <i>(Jung9» to join; first ropes made of rushes.)</i>				Juhccce.	37. —	
8113acutus //.	great sharp sea	Jb A cc 6 jl.au	Ap	Britain	sea co.	S m.s Eng. bot. 1614	
8114 maritimus Lam.	lesser sharp sea	Jb A - 4 au	Ap	Britain	sal. in.	S m.s Eng. bot. 1725	
8115 conglomerate L.	conglomerated	Jb A - 2 jn.jl	Ap	Britain	moi.p.	S m.s Eng. bot. 835	
8116cffTiSU8//.	effuse	Jb - 3 my.au	Ap	Britain	moi.p.	S m.s Eng. bot. 836	
8117 g'n6cus Ehr.	glaucous	Jb ec 2 jl	Ap	England	moi.p.	S m.s Eng. bot. 665	
8118 balticus DM	Baltic	Jb ec 1 jl	Ap	Britain	seaco.	S m.s	
deludes Lk. arcticus Hook.							
8119ErticusL.	retic	Jb A un 1 my	Ap	Norway	1822.	S las Fl. dan. 1094	
8120 Aifitaniil L.	filiform	Jb A un a u	Ap	Britain	tur.bo.	S m.s Eng. bot. 1175	
8121 trflidus L.	three-leaved	jut A un	Ap	Scotland	scalp.	S m.s Eng. bot. 1482	
8122 monanthos Jac.	one-flowered	Jb A un	Ap	Europe	1820.	D m.s	
8123 squarrus L.	squirrosc	Jb A w	Ap	Britain	sa.hea.	S m.s Eng. bot. 933	
8124 Gpsncr Sm.	Gesncr's	Jlb A cu	Ap	Scotland	scalp.	S m.s Eng. bot. 2174	
Rf&clis E. B. tlhuU Hook.							
8125 tenuis W.	slender	Jb A un 1 jl.au	Ap	N. Amer.	1818.	S m.s Rost. j. 1.3	
8126 capitatus Weigel	headed	O un ft JKau	Ap	Europe	1823.	S s	
py^my		Jb A un i jn.au	Ap	France	1817.	S m.s	
8128 planiflthus It. Br.	flat-leaved	Jb iAI un 1 jl.au	Ap	N. Holl.	1822.	D co	
8129 pauciflurus R. Br.	few-flowered	Jb iAI un 1 jl.au	Ap	N. Holl.	1822.	D co	
8130 vaginatUK R. Br.	sheathed	M iAI un 2 jl.au	Ap	N. Holl.	1822.	I co	
8131 lampocirpus Ehrh.	shining-fruited	Jb A w 2 jn.au	Ap	Britain	moi.p.	S m.s Eng. bot. 2143	
8132 acutiflorus Ehrh.	sharp-flowered	Jb A w 2 jl.au	Ap	Britain	moi.p.	S m.s Eng. bot. 238	
8133 obtusiflorus Ehrh.	blunt-flowercd	Jb A w 1 jau	Ap	Briuui	mar.	S m.s Eng. bot. 2144	

- | | | | | | | | | |
|---------------|------------------------------|--|-------------------------------------|--------------------------|----------------------|-----------|---------------------|------------------------|
| 8134 | valvatus L.*. | valved | mu A ■■■■■ | Ap | Spain | 1820. | S | m.s |
| 8135 | polycephalus D. Don | lnany-headed | Jb A ■■■■■ | Ap | Scotland | moun. | D | m.s |
| 8136 | uliginosus Roth | bog | Mi A ■■■■■ | Ap | England | tur.hc. | S | m.s Eng. bot. 801 |
| 8137 | supinus G. Don | supine | Jb A ■■■■■ | Ap | Scotland | moun. | D | m.s |
| 8138 | aristMus Mx. | bearded | Jb A un 1 | Ap | N. Amer. | 1823. | S | |
| 8139 | plebeian R. Br. | plebeian | Mi LAI un ? | Jajl | N. Holl. | 1816. | D | co |
| 8140 | subverticillatus IVul. | bulb-whorled | Jb A un 1 | jl.au | Europe | 1821. | S | m.s |
| 8141 | bulbus L. | bulbous | Jb A w 1 | jl.au | Britain | ... | S | m.s Eng. bot 934 |
| | compressoes Jae. | | | | | | | |
| 8142 | botticus Wold. | Bothnian | Jb A un | ■■■■■ | N. Europe | 1820. | S | m.s |
| 1143 | COMOSUS Bieb. | mud | Jb A un | 1" jlau | N. Holl. | 1816. | D | eo |
| B144 | bufiinus L. | toad | Jb O w | ■■■■■ | Britain | w.s.gr. | S | m.s En. bot. 802 |
| 8145 | Tenageia Ehrh. | Tenageia | 1 O un | & jnau | France | 1817. | S | ms Vait.pa.HD.1 |
| 8146 | trigluminisf. | three-flowered | Jb A un | £ jl | Britain | bgs.m. | S | ms Eng. bot. 899 |
| 8147 | biglumis L. | two-flowered | Jh A un | ■■■■■ | Britain | bgs.m. | S | ms Eng. bot. 898 |
| 8148 | cas&neus Sm. | chestnut | Mi A un | 1 jl | Scotland | sc.alp. | S | ms Eng. bot. 900 |
| 8149 | Jacqrini L. | Jacquin's | Jb A un | ijjl.au | Europe | 1817. | S | ms Jae. au. 3.221 |
| | biglumis Jae. | | | | | | | |
| •995. | LITZULA Do; | LIZULA. | (Gramen Lvzula of Bauhin, | glowworm grass; shines.) | Junceee. | 17.—31. | | |
| 8150 | vernalis Dec. | venial | Jb A un | i ap | Europe | 1820. | D | m.s |
| 8151 | pilosa Sm. | hairy | Jb A " | i ■■■■■ | Britain | groves. | D | m.s Eng. bot. 736 |
| 8152 | flav&cens Dec. | flavescent | Jb A un | ■■■■■ | Europe | 1820. | S | ins Hostgr.3.94 |
| 8153 | Fdrsteri Dec. | Forster's | ju. | A w | England | woods. | S | m.s Eng. bot. 1293 |
| 8154 | glabrata Desv. | smooth | M A un | ■■■■■ | Norway | 1819. | D | m.s |
| 8155 | albida Dec. | Vfht-e-hcaded | Jb A un | ■■■■■ | Switzerl. | ... | S | m.s Lee. her. 13.6 |
| 8156 | littea Dec. | yellow | Jb A | ■■■■■ | Switzerl. | ■ | S | m.s |
| 8157 | spadicea Dec. | light red | Jb A | ■■■■■ | S. Europe | 1821. | S | m.a Vil. del. 2. 6 |
| | Juncus spadiceus TV. | | | | | | | |
| 8158 | nfyca Dec. | snowy | Jlli A un | ■■■■■ | Switzerl. | 1770. | S | m.s Schrb.gr. 7.7 |
| 8159 | maxima Dec. | greatest | JHla A w | ■■■■■ | Britain | woods | S | m.s Eng. bot 7J7 |
| 8160 | arcuata Stvz. | arched | Jllb A un | ■■■■■ | Scotland | moun. | S | m.s |
| | Juncus arcuatusftm | | | | | | | |
| 8161 | spicMa Dec. | Bluked | xik A un | ■■■■■ | Scotland | sc.alp. | S | m.fl Eng. bot. 1176 |
| 8162 | pedifurmis Dec. | foot-formed | Jllb A un | ■■■■■ | Europe | 1824. | S | m.s |
| 8163 | J confesta Lejun | close-headed | Jlb A W | ■■■■■ | France | 1805. | S | m.s Host gr. a 97.5 |
| 8164 | sudeticus Dec. | Sudetic | M A up | ■■■■■ | Silesia | 1817. | S | ins Kr.sil.52 |
| 8165 | campbris Dec. | field | J A | ■■■■■ | Britain | bar.pa. | S | m.s Eng. bot 672 |
| 8166 | multiflbris Le.J. | many-flowered | Jb A un | ■■■■■ | France | 1818. | S | r.m |
| | | | | | | | | |
| 996. | CO'RYPHA L. | FAM PALM. | (Koruphe, the summit; | | leaves only at top.) | Pnlmte. | 7.—13. | |
| 8167 | umbraculifera L. | umbrella-bearing | ■■■■■ 100 | ... | E Indies | 1742. | S | r.m R. mal. 3.1.12 |
| 8168 | glaucescii Hort. | glaucescent | Jf for 100 | ... | E Indies | 1820. | S | r.m |
| 8169 | effta Rox. | tall | ID or 150 | ... | E Indies | 1825. | S | r.in |
| 8170 | Ustan Lam. | Utan | ID or 50 | ... | Moluccas | 1825. | S | r.m |
| 8171 | Uectorum//,\$& | roof | J! or 15 | ... | W.Indies | 1825. | S | r.m |
| 8172 | Pdmos H. & B. | Pumos | J! or 20 | ... | Cuba | 1824. | S | r.m |
| 8173 | australis 11. Br. | southern | J! or 50 | ... | N. Holl. | 1824. | S | r.m |
| | | | | | | | | |
| 997. | TALIETRA Mart | TALI EH A PALM. | (Its aboriginal name.) | | Pdlnue. | 1.—7. | | |
| 8174 | bengalensis Spr. | Bengal | iQ or 10 | — | E Indies | 1823. | S | r.m |
| | | Corypha Taliera Rox. | | | | | | |
| 998. | LICUATLA Thun. | LICUALA. | (Its Macassar name.) | | Pidmc. | 2. | | |
| 8175 | spinosa Thun. | spiny | J Q e c 6 | ... | W.u | E. Indies | 1802 | S r.m Ru.am. 1.9 |
| 8176 | peltata Rox. | peltate | JtCIlor 6 | ... | W.Y | E. Indies | 1825. | S r.m |
| | | | | | | | | |
| oi999. | THRPNAX L.JU. | THRIXAX. | (Thrinax, a fan; form of leaves.) | | Liilnue. | 1.—■ | | |
| 8177 | parviflora Sun. | ismall-flowered | f O ee 15 | ... | W.u | Jamaica | 1778. | S r.m |
| | | | | | | | | |
| 1000. | TRADESC A'NT/4 L. | SPIDER WORT. (John Tradescant, gardener to Charles I.) | Commetinea. 9.1.—99. | | | | | |
| 8178 | fuscita Ker | browned | f Ej or | ■■■■■ | S. Amer. | 1824. | L | r.m Botreg.482 |
| 8179 | malabarica L. | Malabar | f El or | 1 jlau | P | E Indies | 1776. | Skr.m R. mal. 9.63 |
| 8180 | tuberosa Rox. | tuberous | A El or | 14 jl au | B | E. Indies | 1817. | R p.l |
| 8181 | 8jecisca L. | showy | f IJU or | 1 jl.au | B | Mexico | 1825. | D r.m |
| 8182 | discolor H. K. | various-colored^ | ■■■■■ | 1 ap.s | VV | & Amer. | 1784. | Sk&p Bot mag. 1192 |
| 8183 | cordifolia Sivz. | heart-leaved | f, El | ■■■■■ | B | Jamaica | 1819. | D r.m |
| 8184 | crecta Cao. | upright | 2 jlau | ■■■■■ | B | Mexico | 1794. | S r.m Bot. mag. 1340 |
| 8185 | uudata Hum. | waved | 1 jn,jl | ■■■■■ | B | Trinidad | 1819. | D r.m |
| 8186 | diuretica Mart. | diuretic | Jjn,jl | ■■■■■ | B | Brazil | 1825. | D r.m |
| 8187 | polchellia Hum. | neat | 1 jlau | ■■■■■ | B | Mexico | 1825. | D e.p |
| 8188 | subaspera Ker | roughish | 1 my.o | ■■■■■ | P | N. Amer. | 1812. | D r.m Bot. mag. 1597 |
| 8189 | crassifolia Cav. | thick-leaved | 3 jlo | ■■■■■ | B | Mexico | 1796. | L B.p Bot. mag. 1598 |
| 8190 | virginica L. | Virginian | \ my.o | ■■■■■ | B | N. Amer. | 1629. | D p.l Bot mag. 105 |
| | 2affa | white | my.o | ■■■■■ | B | N. Amer. | 1629. | I) r.m |
| | 3 rfabia | red | my.o | ■■■■■ | R | N. Amer. | 1629. | D r.m |
| | 4 plena | blue and white | my.o | ■■■■■ | B | N. Amer. | 1629. | D r.m |
| | 5 caerulea fflbida | pilose | my.o | ■■■■■ | B | N. Amer. | 1629. | D co |
| | 6 tiffia Swz. | rose-flowered | my.o | ■■■■■ | Pk | Carolina | 1802. | D r.m Bot cab. 370 |
| | procfin.bens IV. | many-flowered | i jn,jl | ■■■■■ | B | Jamaica | 1820. | D r.m |
| | multiflbra Jae. | trailing | * jn,jl | ■■■■■ | B | Trinidad | 1824. | D r.m Jae. ie. 2.355 |
| | | | | | | | | |
| | small-flowered | f El or | 1 au s | ■■■■■ | B | Peru | 1822. | L r.m Fl. per. 3.272.8 |
| | broari-icavcd | * O or | 1 Jj s.n | ■■■■■ | B | Lima | 1816. | S r.m Fl. per. 3.272.1 |
| | divaricata | f El or | i injl | ■■■■■ | B | Trinidad | 1818. | D r.m Aub gui. 1.12 |
| | knotted | f El or | 1 jlau | ■■■■■ | B | W.Indies | 1783. | L s.p Jae.am.64 |
| | panicled | ,QJ or | 1 au | ■■■■■ | B | E. Indies | 1816. | S pi |
| | | | | | | | | |
| 1001. | CAMP^LIA Rich. | CAMPELIA. | ■■■■■, a bending, helias, the sun.) | | Commelinete. | 1. | | |
| Zanonia Rich. | Zanonia.ll/f | Tradscant/a Zanonia L. | ■■■■■ | | W. Indies | 1759. | S r.m Red. Id. 192 | |
| | | | | | | | | |
| | CYANO TIS D. Don | CTANOHS. | (Kyanos, blue, ous, an ear.) | | Commfdnear | 3.-4 | | |
| | | | | | | | | |
| | ■■■■■ D. Dw* | axillary | OJor i jj;; | ■■■■■ | E. Indies | 1822. | S r.m R. mal lo! 13 | |
| | ■■■■■ stata(Z).i)o> | crested | OJor 1 jl s | ■■■■■ | Ceylon | 1770. | S r.m Botmai 1435 | |
| | | | | | | | | |
| | Bono ^ adescndta cristata L. | | | | | | | |
| | ■■■■■ barbnta D. Don | bearded | A or 1 jU | ■■■■■ | Nopal | 1824. | D rm | |

1003. CARTONETMA R. Br.	CARTONEMA.	(<i>Kartos</i> , shorn, <i>nema</i> , a filament) <i>Comtnlinear</i> . 1.
8203 spicatum AJ3!r.	spiked	£{ZSor 1jl.au D N. Holl. J822. D i p
•1004. DICHORIZA'NDR A il/i*. DICHORIZANDRA.	(S** twice, cAomo, to part, anrr, anther) <i>Comme'nea</i> . 4.-5.	
8204 thrysiflra A/**,	thyrs-flowered £ ED or	4 au II Brazil 1822. R r.m Bot. reg. 682
8205 puberula Mart.	downy £ ES or	3 au s 11 Brazil 1823. D s.p
8206 oxytiwala fe-■	sharp-petaled ■ A gr 2 jl.au R Brazil 1810. Ex.r.m Bot. mag. 2721	
8207 gracilis Jlfar.	slender ■■■■■ 1_a.uu B Brazil — D r.m	
* 1005. AGAPANTHUS HerU.	AFRICAN LILV.	(<i>Agape</i> , love, <i>anthos</i> , flower.) <i>Hemerocattxete</i> . 3.
8208 umbellatus Ucri.	umbellate tf JAior	3ja.au B C. G. H. 1692. R r.m Bot. mag. 500
2 variegatus	Variegated tf tfl Alor	2 ja.au B — R r.m
8209 praecox W. en.	early tf iAl or	4 jn.au P.B C. G. H. — R r.m
8210 minor hod.	tf iAl or	2 ja.au B C. G. H. — D r.m Bot. cab. 42
l**cr		
1006. BLANDFCRD W Sm.	BLANDFORIMA.	(George, Marquis of Blandford.) <i>Hemerocallideie</i> . 2.
8911 nabilis Sm.	noble *iAlor	2jl.au O N. S. W. 1803. S s.l.p Ex. bot. 1.4
8212 grandiflora JZ 2?r.	large-flowered £ iAlor	2 jl.au C N. S. VV. 1812. S s.l.p
1007. i/EMEROCA'LLIS/.. DAY LILY. (<i>Hemera</i> , day, *i/o* beauty; duration.) <i>Hemeroallidcat</i> . *		
8213 /raminea // JC	grassy-icavcU	£ A or 1 jn.jl LY Siberia 1759. R s.l Bot. mag. 87.
8214 flava //,	yellow £ A or 2 jn Y Siberia IWrt. R s.l Bot. mag. 19	
8215 disticha Donn	two-rowed £ A or 2 my.jl O China 1798. R s.l Sw. fl. bar. 28	
8216 f&lva L.	fulvous £ A or 4 jn.au Ful Levant 1596. R s.l Bot. mag. 64	
• 1008. FU'NKL* Spr.	FUNKIA.	(<i>Henri</i> /Funk, a German crytogramist) <i>Hemerocallidcat</i> . 2.-3.
8217 subcordata Spr.	subcordate-vrf £ A or 1 au.s W Japan 1790. R p.l Bot. mag. 1433	
J/microdillia japonica IV.		
8218 ovata Spr.	ovate-leaved £ A or 1J my.jl B Japan 1790. R p.l Bot. mag. 894	
/temerocallis crsrlca Andr.		
1009. RHIPIDODE'NDRON TV. RIHIPIDODEDROV. (<i>Rhipis</i> , a fan, <i>dendron</i> , a tree.) <i>Hemeroallidcat</i> . L		
8219 platicile Haw.	plaited *i_Jgr 8 jn.jl R Africa 1723. C s.l Bot. mag. 457	
distichum W.	if'loii phatilis // K.	
2mäjus	larger •t-Jgr 10jn.jl R Africa 1723. C s.l	
1010. PACHIDE'NDRON W. PACIWENDROV. (<i>Pachys</i> , thick, <i>dendron</i> , tree.) <i>Hemeroallidcat</i> . 8.		
8220 afriquetum Haw.	African *i_J gr 8 jl R C. G. H. 1819. Sk s.l Bot. mag. 2517	
A'löe afric&ma Haw.		
8221 pseudoo-africAlium Haw.	bast. African * J gr 6 mr n O C. G. H. 1731. Sk s.l Bot. mag. 1322	
A'löe psuedo-africflna HerU.	africana /2angustior B. M.	
8222 principis Haw.	prince's * i_J g* 5 mr.n Y C. G. H. 1821. Sks.1	
jillo principis Haw.		
8223 ferox Haw.	hedgehog * LJ gf 6 ap.my Y C. G. H. 1759. Sks.1 Bot. mag. 1975	
A'löe ferox Dec.		
8224 pseudo-ferox Haw.	bast, hedgehog ** i_J gr 4 my.jn O C.G.H. 1820. Sks.1	
A'löe psuedo-ferox Salm.	subferox Spr.	
8225 angustifolium Haw.	narrow-leaved * L_J 8T 8 jl.au R C. G. H. 1806. Sks.1	
if'loo angustiflaria Haw.		
8226 supralae vc Haw.	above smooth M LJ gr 5 ... Ö C. G. H. 1731. Sks.1 Com. pr. 71.20	
1011. if'LOE L.	ALOE.	(<i>A'Uothetis</i> Arabic name.) <i>Hemeroallidcat</i> . 49.—L0.
\$92? dichotoma L.	dichotomous ■■■■■ 18	C. G. H. 1780. Ls s.1
8228 purpurascens Haw.	purpurascens ■■■■■ * 12jl.o	C. G. H. 1789. C s.l Bot. mag. 1474
8229 socotrina Dec.	Socotrine ■■■■■ 12 f.ap	C. G. H. 1731. C s.l Bot. mag. 472
8230 spicata /,	spike-flowered ■■■■■	C. G. H. 1795. Sk s.p
8231 pluridens Haw.	many-toothed ■■■■■ 2 jl.au	C. G. H. 1823. Sk s.1
8232 arborescens Dee.	artwrecessent ■■■■■ 12 mr.n	C. G. H. 1731. C s.l Bot. mag. 1306
8233 gracilis Haw.	slender ■■■■■ 3 jn.jl	C. G. H. 1822. Sks.1
8234 fiavispina Haw.	yellow-spined ■■■■■ 5 au	C. G. H. 1793. Sk s.1
8235 mitroformis L.	mitre-formed ■■■■■ 6 au	C. G. H. 1732. Sk s.l Bot. mag. 1270
8236 n'ibilu Haw.	noble ■■■■■ 5 au	C. G. H. 1800. Sk s.1
8237 distans Haw.	distant ■■■■■ 6 au	C. G. H. 1732. Sk s.l Bot. mag. 1362
2 refl'xa Salm.	reflexed ■■■■■ 4 jlcu	C. G. H. 1820. Sk s.1
3 depreßsa Haw.	depressed ■■■■■ 4 jl.au	C. G. H. 1820. Sks.1
8238 Commelin's W.	Commelin's ■■■■■ 3 ...	C. G. H. 1819. Sk s.1
mitrifurmis Salm		
8239 xanthacanthia TV.	yellow-spined ■■■■■ 3 jn O	C. G. H. 1817. Sk s.1
8240 vulgariifl Dec.	common ■■■■■ 2 my.au Y Levant 1596. Sks.1 Dec.pi.27	
barbadensis Haw.		
8241 chinensis Haw.	Chinese ■■■■■ 3 jn Y China 1821. Sks.1	
8242 cassia Salm	canious ■■■■■ 2 jn O C.G.H. 1818. Sks.1	
2 clatiior 7<kw.	taller ■■■■■ 9 njl R C. G. H. 1821. Sks.1	
8243 frutescens Salm	frutescens ■■■■■ 5 R C. G. H. 1819. Sks.1	
8244 albisp^na Haw.	white-spined ■■■■■ 7 S C. G. H. 1796. Lss.1	
8245 glauca 7/aw.	glauca ■■■■■ 7 as R C. G. H. 1731. Sks.1	
2 rhodacantha B. M.	rhodacantha B. M. ■■■■■ red-spnd a. i fer 4 jas R C. G. H. 1731. Sks.1	Bot. mag. 1278
8246 spiniflorus /four.	spinier ■■■■■ 4 jas R C. G. H. 1820. Sks.1	
8247 lineata Haw.	lined ■■■■■ 4 jas S C. G. H. 1789. Sk s.p	
2 glaucescens Haw.	glaucescent ■■■■■ 5 ... S C. G. H. 1789. Sks.1	
8248 depressa //AW.	flaUcaved ■■■■■ 6 au O C. G. H. 1731. Lss.1	Bot. mag. 1332
8249 brevifolia Haw.	short-leaved ■■■■■ ijnjl O C. G. ft 1810. Sks.1	Bot. mag. 996
8250 proli. flra Salm	proliferous ■■■■■ ijnjl O C. G. G. 1819. Sks.1	
2 major Salm	greater ■■■■■ 2 mr.jn O C. G. H. 1819. Sks.1	
8251 serrata Dec.	sav-leaved ■■■■■ 2 mr.jn O C. G. H. 1818. Sks.1	
8252 paniculata Joe.	paniced ■■■■■ S C. G. H. 1795. Sks.1	Jac. fr. 62
striatafluiw..		
8253 striatula Haw.	■■■■■ i J gr 3 jajl R C. G. H. 1821. Sks.1	
ciliaris Haw.	fringe/oiva/ ■■■■■ 5 jn.jl R C. G. H. 1821. Sks.1	
8255 tentuor //au.	slender ■■■■■ *L i J gr 2 jl.au Y C. G. H. 1821. Sks.1	
8256 albocincta Haw.	white-edged tt ■■■■■ 3 jn O C. G. H. 1812. Sks.1	
8257 serrulata A. K.	aaw.leaved tL ■■■■■ 7 jl.au R C. 6' H. 1789. Sks.1	
obscira /fotv.	obscure tt ■■■■■ 2 jl.au R C. G. H. 1819. Sks.1	Bot. mag. 1323
nirta Tire.		

S glaficior <i>Haw.</i>	more glaucous	SL iAI gr	2 jl.au	R	C. G. H.	1819.	Sks.1
3 magnidens <i>Haw.</i>	large-toothed	«L IAI gr	3 jnj.l	R	C. G. H.	1820.	Sks.1
8259 palliscens <i>Haw.</i>	Pallescent	tLAiAltr	3 jnj.l	R	C. G. H.	1820.	Sks.1
8260 latifolia <i>Haw.</i>	road-lvd-soap	SL J gr	6 jlau	S	C. G. H.	1795.	Sks.1
8261 saponaria <i>Haw.</i>	common soap	*L J gr	4 jlau	R	C. G. H.	1727.	Sks.1
2 Qteo-stnata <i>Haw.</i>	yellow-striped	(L) gr	4 jlau	It	C. G. H.	1821.	Sks.1
8262 micracantha JX M.	small-spined	VLA gr	3 jl	Pk	C. G. H.	1819.	Sks.1
8263 virens <i>Haw.</i>	green	III gr	3 au.s	Y	C. G. H.	1790.	Sks.1
8264 subcreta <i>Haw.</i>	subcrect	£ IAJgr	5 mrjn	S	C. G. H.	1789.	S s.1
2 semiguttata /tow.	half-spotted	£ IAJgr	3 mrjn	S	C. G. H.	1819.	Sks.1
8265 acuminata <i>Haw.</i>	acuminata	£ IAJgr	3 mr.my	O	C. G. H.	1795.	Sks.1
hilmilis B. M.							Bot mag. 757
8266 echinMa W.	cchinata	£	6				
8267 tuberculata ■■■■■	tubcrc.-hedgeh.	LAJ gr	2 mr.my	O	C. G. H.	1821.	Sks.1
8268 tenuifolia <i>Haw.</i>	thin-leaved	£ IAJ gr	1 jnj.l	O	C. G. H.	1821.	Sks.1
8269 vanegata IV.	variegated	£ LAJ gr	4 mr.s	Pk	C. G. H.	1720.	Sks.1
8270 aristata <i>Haw.</i>	bearded many4vd	£ iAI gr	2 myjn	O	C. G. H.	1801.	Sk&l
8271 suTuberculata <i>Haw.</i>	suTuberculated?	LAJ gr	2 jnj.l	O	C. G. H.	1620.	Sks.1
8272 hiimilis Thun.	humble	£ IAJ gr	1 mazjn	O	C. G. H.	1731.	Sks.1
8273 incitvra <i>Haw.</i>	incurred	£ IAJ gr	1 mr.jn	O	C. G. H.	1796.	Sks.1
hiimilis (3 inctivra B. M.							Bot. mag. 828

1012. BOWIES *Haw.* BOWIEA. (Jltouife, a collector of plants for Kew gardens.) *HemcrocaU'idae.* 8.
8274 myriacantha *Haw.* inany-spined ■■■■■ apjn R.G C. G. H. 1823. Sks.p
8275 atricana *Haw.* African ■■■■■ o R.o C. G. H. 1823. Sk s.p

1013. GASTEMtLA-Háti. GASTERIA. (Gaster, a belly; base of flowers.) <i>Hemerocallidiace.</i> 42.							
8276 pfilchra <i>Haw.</i>	fair	«. L gr	3 jn.au	S.B	C. G. H.	1759.	C ■■■■■ Bot. mag. 765
8277 maculata <i>Haw.</i>	spotted	«-i J gr	2 jlau	SR	C. G. H.	1759.	Sks.1 Bot mag. 979.
2 ftflax <i>Haw.</i>	fallacious	«. J	2 jlau	S.B	C. G. H.	1820.	Sks.1
8278 picta <i>Haw.</i>	painted	■■■■■	3 jlau	S.B	C. G. H.	1820.	Sks.1
8279 formbsa <i>Haw.</i>	beautiful	■■■■■	2 jlau	S.n	C. G. H.	1820.	Sks.1
8280 fasciata <i>Haw.</i>	banded	■■■■■	1 jnj.l	R	C.G.H.	1820.	Sks.1
nf gricata y fasciata	Saltn						
2 laxa <i>Haw.</i>	loose	■■■■■	2 jlau	S.B	C.G. H.	1820.	Sks.1
8281 retata <i>Haw.</i>	netted	£ LAJ gr	2 jlau	SR	C.G. H.	1820.	Sks.1
8282 nfricans <i>Haw.</i>	dark	£ IAJgr	2 jnj.l	R	C.G. H.	1790.	Sks.1
2 marmoritta <i>Haw.</i>	marbled	fr Al ■■■■■	1* jnj.l	R	C.G. H.	1820.	Sks.1
8283 crassifolia <i>Haw.</i>	thick-leaved	■■■■■		R	C.G. H.	1820.	Sks.1
8284 brcvif.Mia <i>Haw.</i>	short-lvd tonguef	LAJ gr	14 jnj.l	R	C.G. H.	1809.	Sks.1
2 pveridis <i>Haw.</i>	very green	£ IAJ gr	3 jlau	S.B	C.G. H.	1820.	Sks.1
8285 obtusi folia <i>Haw.</i>	blunt-leaved	£ ■■■■■	jlau	R	C.G. H.	1796.	Sks.1
8286 mollis <i>Haw.</i>	soft muddy-lvd f	■■■■■	ymjl	R	C.G. H.	1823.	Sks.1
8287 subnigrans <i>Haw.</i>	blackish	■■■■■	1-u	S.B	C.G. H.	1820.	Sks.1
2 glabrior	smoother	■■■■■	jlau	S.B	C.G. H.	182a	Sks.1
8288 dtsticha <i>Haw.</i>	two-rowed	je ■■■■■	jlau	S.B	C.G. H.	1820.	Sks.1
2 major <i>Haw.</i>	larger	£ ■■■■■	jlau	SR	C.G. H.	1820.	Sks.1
8289 consprucata <i>Haw.</i>	dirtyd	£ ■■■■■	mr.s	R	C.G. H.	1796.	Sks.1
8290 angulMa <i>Haw.</i>	angled	£ IAJgr	2 mr.n	R	C.G. H.	1791.	Sks.1
longifolia <i>Haw.</i>							Bot. mag. 1322
2 minor <i>Haw.</i>	smaller	f tAlgr	liau	R	C.G. H.	1820.	Sk*.1 -
8291 sulcata Uaw.	sulcated	£ tAlgr	2 jlau	SE	C.G. H.	1820.	Sks.1
8292 excavata <i>Haw.</i>	excavated	tflAlgr	U ...	R	C.G. H.	1824.	Sks.1
2 obllqua <i>Haw.</i>		■■■■■	li jn.au	R	C.G. H.	1759.	1'8.1
8293 angusti folia <i>Haw.</i>	narrow-leaved	■■■■■	1} mrs	R	C.G. H.	1731.	Sks.1
2 longifolia <i>Haw.</i>	long-leaved	■■■■■	1} mrs	R	C.G. H.	1796.	Sks.1
8294 lwVig <i>Haw.</i>	smooth	■■■■■	limr..	R	C.G. H.	1820.	Sks.1
8295 subveruebsa <i>Haw.</i>	subverueose	IAK	■■■■■	R	C.G. H.	1820.	Sks.1
2 narvipunct'ita	small-dotted	■■■■■	2 !!!	R	C.G. H.	1731.	Ski.1
J296 verrucosa <i>Haw.</i>	warty	£ IAJ gr	2 mr.n	R	C.G. H.	1790.	Sks.1
0*97 intermedia <i>Haw.</i>	middle tonguef	■■■■■	2 mr.n	R	C.G. H.	1820.	Bot. mag. 837
2 asierrima <i>Haw.</i>	roughest leavediz	IAI gr	2 mr.s	R	C.G. H.	1820.	Bot. mag. 1322.1
3 lie vior <i>Haw.</i>	smoother	je ■■■■■	jlau	S.B	C.G. H.	1820.	Sk&l
4 lóngior <i>Haw.</i>		■■■■■	jlau	SR	C.G. H.	1820.	Sks.1
8298 ripens <i>Haw.</i>	crecyng-t ootedz	IAJK	1 jl	R	C.G. H.	1821.	Sks.1
8299 parva <i>Haw.</i>	small	■■■■■	2 jlau	S.B	C.G. H.	182a	Sks.1
8300 declpiens <i>Haw.</i>	deceiving	■■■■■	2 jlau	SR	C.G. H.	1820.	Sks.1
nfgricans <i>Haw.</i>							
8301 carinata <i>Haw.</i>	£ IAJgr	2 jnj.l	R	C.G. H.	1731.	Lss.1 Bot. mag. 1331	
8302 strigata <i>Haw.</i>	ridged	£ IAler	2 jlau	S.R	C.G. H.	1820.	Sks.1
8303 ltEtipunctata <i>Haw.</i>	lively-spotted	£ IAJgr	2 jlau	S.R	C.G. H.	1820.	Sks.1
2 denticuKita	denticulated	£ IAler	2 jlau	S.H	C.O. H.	1822.	Sks.1
''04 subcarinata <i>Haw.</i>	obbcure-kecled	£ Al ■■■■■	2 jnj.l	S.G.H.	1818.	Ske.1	
2 viridior <i>Haw.</i>	greener	£ IAJgr	2 jlau	O	C.G. H.	1820.	Sks.1
3 lie vior <i>Haw.</i>	waved	£ LAJgr	2 jlau	S.R	C.G. H.	1796.	Sks.1
4 lóngior <i>Haw.</i>	smooth-keeled	f IAJgr	3 jnj.l	S.R	C.G. H.	1820.	Sks.1
8298 ripens <i>Haw.</i>	smaller	■■■■■	2 jlau	R	C.G. H.	1820.	Sks.1
8299 parva <i>Haw.</i>	shining	£ IAJgr	1 jlau	S.R	C.G. H.	1822.	Sks.1
8300 declpiens <i>Haw.</i>	large-dotted	£ IAler	1 jlau	R	C.G. H.	1820.	Sks.1
nfgricans <i>Haw.</i>	triangular	£ IAJgr	1 \$jn.au	R	C.G. H.	182a	Sks.1
8301 carinata <i>Haw.</i>	obtuse	■■■■■	1 jn.au	R	C.G. H.	1820.	Sks.1
8302 strigata <i>Haw.</i>	scimitar-lvd	■■■■■	mr.s	R	C.G. H.	1819.	Sks.1
8303 ltEtipunctata <i>Haw.</i>	smaller	£ LAJgr	jlau'	O	C.G. H.	1820.	Sks.1
2 denticuKita		£ IAJgr	jlau	S.R	C.G. H.	1820.	Sks.1
''04 subcarinata <i>Haw.</i>	many-dotted	£ S gr	jlau	S.R	C.G. H.	1820.	Sks.1
2 magnipunc^ta	large-dotted	£ IAI gr	jlau	S.R	C.G. H.	1823.	Sks.1
■■■■■ ensifolia <i>Haw.</i>	sword-loaved	£ IAJ B*	jlau	S.R	C.G. H.	1823.	Sks.1
J215 candicans <i>Haw.</i> rough mat £.white		f*		R	C.G. H.	189&	Sk s.1
8ii5 S nita ■■■■■	smeared	£ LAJ K	jlau	R	C.G. H.	1820.	Sks.1
8ii7 bicolor <i>Haw.</i>	two-colored	£ IAJ 8*	jlau	S.R	C.G. H.	1824.	Sks.1
J216 HAWORTHIA. (A. H. HawoMh, F.L.S., disting. bot.) <i>Hetcrocattfdcar</i> 64	clammy	£ LAJgr	H jnj.l	Gy	C.G. H.	1727.	Sks.1 Bot mag! 814
viscosa L.							

8319 <i>indurata</i> Haw.	hard <i>branchy</i>	£ iAI gr	*jn.Jl	Gy	C. G. H. 18S0. Sk 8.1
8320 <i>pseuido-tortuosa</i> SoAft	subtwist. <i>truing</i> .	^ IAJgr	1 jl.au	Gy	C. G. H. 1818. Sk s.l
8321 <i>tortuosa</i> Haw.	collared	I AJgr	1 Jls		C. G. H. 1823. Sk s 1
8322 <i>concinna</i> Hava.	neat	I AJgr	s ...	Gy	C. G. H. 1818. Sk s.l
8323 <i>cordifolia</i> Haw.	heart-leaved	I AJgr	ft myjl	Gy	C. G. H. 1817. Sk s.l
8324 <i>aspcruiiscula</i> Aatv.	roughish	I AJgr		Gy	C.*K. H. 18ia Sk HI
8325 <i>tortutaa</i> //w.	twisted		1 my.s	Gy	C. TJ. H. 1794. Sk s.l
^{^Mœ} <i>rigida</i> B. M.					
8326 <i>côrta</i> //aw.	<i>abort-twisted</i>	■ IAlgr	H ~		C. G. H. 1876. Sks.l
8327 <i>tortella</i> Haw.	little-twisted	* IAJgr			C. G. H. 1817. Skb.l
8328 <i>nltidia</i> Salm	shining	IF IAJgr			C. G. H. 1825. Sks.l
8329 <i>setta</i> //ww.	bii8tle-tv//r<	* IAJgr			(O. H. 1818. Sks.l
8330 <i>expansa</i> Haw.	xi-anded	£ IAJgr			C. G. H. 1795. Sks.l
rigida Itec.					
8331 <i>hbrida</i> Salm	hybrid	tf IAJgr	1 j^n	Gy	C. G. H. 1821. Sk s.l
8332 <i>scabra</i> /aw.	rough	I AJgr	i jn.jl	Oy	C. G. H. 1818. Sk 81
8333 <i>rfrida</i> Haw.	sordid	JP IAJgr	jl.au	Gy	C. G. H. 1820. Sk s 1
8334 <i>venhsa</i> Haw.	veiny	j£ IAlgr		Oy	C. G. H. 18. U Sk 8.1
A'Xoc vndosa Lam.					
8335 <i>recfirva</i> Haw.	recurve-feawr	£ iAJ gr	1 au	Gy	C. G. H. 1795. Sks.l
8336 <i>nfgicans</i> //aw.	granuiat. black	£ iAJ gr	i an	Gy	C. G. H. Skel.
8337 <i>pifira</i> //iw.	small	if IAJ gr	*my.jn	Gy	C. G. H. 1821. Sks.l
8338 <i>teusellita</i> Haw.	rftr/r-checkeredij	LAJgr	4 jnau	Gy	C. G. H. 1823. Sks.l
8339 <i>multifaria</i> //aw.	many-sided	£ iAI gr	■ jnjl	Gy	C. G. H. 1824. Sks.l
8340 <i>retusa</i> IK	recluse cushion	\$ LAJ gr	1 my.jl	Gy	C. G. H. 1720. Sk s.l
8341 <i>mltica</i> //iw.	blunt cushion	f IAJ gr	ft JnJI	Gy	C. G. H. 1870. Sks.l
8342 <i>mrahilis</i> Haw.	admirable <i>cuvionz</i>	£ iAJ gr	f my.au	Gy	C. G. H. 1795. Sks.l
8343 <i>turgida</i> Haw.	turgid cushion	lg IAJ gr	ft au	Gy	C. G. H. 1819. Sks.l
8344 <i>tarte</i> vireiw //aw.	lively green	£ iAJ gr	ft "ii	Gy	C. G. H. 1819. Sks.l
8345 pp. 2	papillose	if IAJ gr	1 my.au	Gy	C. O. H. 1820. Sks.l
2 <i>ticmipapillsa</i>	naif-papillose	i IAJ gr	ft au	Gy	C. G. H. 1820. Sks.l
8346 <i>crecta</i> Haw.	erect pearl	£ IAI gr	i ...	Gy	C. G. H. 1838. Sks.l
8347 <i>fasrfata</i> Haw.	banded-Mrar	£ iAJ gr	I jl.au	Gy	C. G. H. 1818. Sks.l
2 mäjor	larger	£ IAJ gr	* I jl.au	Gy	C. G. H. 1821. Sks.l
8348 <i>granata</i> Haw.	grained	je IAI gr	i jn au	Gy	C. G. H. 1735. Sks.l
A'lo'i margaritifera y minima Dil.					
8349 <i>brevia</i> Haw.	short pearl	£ IAJ gr	i jn.jl	Gy	C. G. H. 1810. Sk s.l
Aloe margaritifera y minima // M.					
8350 <i>minor</i> Haw.	lebser ;rtir/	£ AI gr	1 my.au	Gy	C. G. H. Sks.l
8351 <i>inargaritifera</i> // K.	pearl-bearing	£ IAJ gr	1 my.s	Gy	C. G. H. 1739. Sks.l
8352 <i>Haworthi</i> Hnrt.	Haworth's	^	1 au	Gy	C. O. H. 1801. Sks.l
8353 <i>sciniglubrMa</i> //aw.	hall'-sinoothed	£ LAJgr	■ my.au	Gy	C. O. H. 1811. Sks.l
8354 <i>râdula</i> Haw.	■				1805. Sk s.l
1 UeVior Haw.	smoother	£ IAJ gr	1 au	Gy	C. a. H. 1825. Sks.l
2 pluriperlta	many-peared	tf IAlrr	1 au	Gy	C. G. H. 1820. Sks.l
3 asptrior Haw.	rougher	£ LAJ gr	1 au	Gy	C. G. H. 1821. Sks.l
8355 <i>attenuata</i> //w.	attenuated-f*ar/f	LAJ gr	1 my.au	Gy	C. G. H. 1790. Sks.l
ii'loc râdula if. Af.					Bot mag. 1345
8356 Reinwarf Salm	Reinwart's pearl	£ LAJ gr	i jn.jl	Gy	C. G. H. 1820. Sk s.l
Hawi'jrth/a					
8357 <i>semimargaritifera</i> //aw.	semi-pearl-bg	£ iAI gr	1 mr.s	Gy	C. G. H. 1819. Sks.l
2 major //aw.	larger	£ IAJ gr	1 mr.s	Gy	C. G. H. 1819. Sks.l
3 minor Haw.	smaller	£ IAJ gr	1 mr.s	Gy	C. G. H. 1819. rik s.l
4 multiplerata //aw.	many-peared	£ LAJ gr	1 mr.s	Gy	C. G. H. 1819. Sks.l
8358 <i>coarcata</i> Haw.	compressed	£ LAJ gr	4au	Gy	C. G. H. 1821. Sks.l
&359 <i>albicans</i> Haw.	white-rr/crf	£ IAJ gr	1 Jl	Gy	C. G. H. 1715. Sks.l
8360 la~vis //aw.	smooth whde-edxvd	£ LAJ gr	x au	Gy	C. G. H. 1820. Sks.l
8361 <i>ramifcra</i> Haw.	branch-bearing	£ IAI gr	1 jl.au	Gy	C. G. H. 1821. Sks.l
8362 <i>viresentis</i> Haw.	vircscnt	£ LAJ gr	1 a.us	Sy	C. G. H. 1819. Sks.l
2 minor	smaller	£ IAJ gr	ft a.us	Gy	C. G. H. 1819. Sks.l
8363 ararnoldcs Haw.	cobweb-like	£ IAJgr	1 au	Gy	C. G. H. 1727. Ls.s.l
2 minor Haw.	smaller colnveb	£ IAJ gr	a.us	Gy	C. G. H. 1819. Sks.l
8364 transitions H. K.	transparent	£ LAJ gr	\$ my.au	uv	C. G. H. 1795. La s.l
8365 <i>setata</i> Haw.	bfvsMis-leavcd	£ LAJ gr	1 jl.au	Gy	C. G. H. 1820. Sks.l
2 nfgicans Haw.	blackish	£ LAI gr	1 jl.au	Gy	C. G. H. TSM. Sks.l
3 media//aw.	mediate	■	1 jl.au	Gy	C. G. H. 1820. Sks.l
4 major Haw.	larger	£ LAJgr	1 jl.au	Gy	C. G. H. 1820. Sk s.l
8366 pâl ida //aw.	palcirrarn	£ IAJgr	ft Jn.jl	Sy	C. G. H. 1820. Sks.l
8367 atroyrens //aw.	dark-green	£ IAJgr	1 my	Gy	C. G. H. 182a Sks.l
A'lo'6 arachnoldcs	pumila B M.				Bot mag. 1361
8368 <i>pumila</i> //aw.	dwarf cobweb	£ LAJgr	1 my	Gy	C. G. H. 1752. Sks.l
8369 <i>floracâitha</i> //aw.	gretm-pñned	If LAJgr	I au	Gy	C. G. H. 1820. Sks.l
8370 <i>angulifjlia</i> //w.	narrow-leaved	£ IAlgr	ft jn	Gy	C. G. H. 1884. Sks.l
8371 <i>reticulata</i> Haw.	netted	JgiIAJgr	4 my.au	Gy	C. G. H. 1794. Skfl.l
8372 <i>cymbifñrmis</i> Haw.	boat-formed	£ LAJgr	f my.au	Gy	C. G. H. 1795. Sks.l
concava Haw.					Bot mag. tf'4 Hot. mug. bl/3
8373 altilineti Haw.	ridge-lincil	£ LAJgr	i JUu	Gy	C. G. H. 1824. Sks.l
8374 obtfsls //aw.	small blunt	£ iAJ gr	ft jn	Gy	C. G. H. 1N24. Sk s.l
8375 planifflmia Haw.	flat-leaved	£ IAJgr			C. G. H. 1824. Sk s.l
8376 mucronata //aw.	vnenibr. pointed	£ LAJ gr			C. G. H. 1820. Sks.l
8377 limpida Haw.	limpid	£ LAJgr	} au		C. J. H. 1K9. rik 81
8378 aristata Haw.	awned	£ IAlgr	1 jl	Gy	C. G. H. 1820. Sks.l
8379 cuspidata //aw.	cuspidate	£ IAJ gr	\ au	Gy	C. I*. H. 1819. Sks.l
8380 denticulate Haw.	denticulate	^ LAJgr	au	Gy	C. G. H. 1819. Sks.l
1015. A'PICRA W.					
8381 spirMis Haw.	APICRA. (Apikros, not bitter; unlike bitter aloes.)				Hemerocall'dc*. \ 1-
8382 spirElla Haw.	gn-a spiral	M. gr	1 am a	Gy	(gr. 9. 1. 199A SkB.l
8383 spirElla Jac.	small spiral	n. gr	1 au	Gy	C. G. H. 1808. Sks.1'
8384 icentragNiia Haw.	little-blistered	tL gr	1 my.jn	Gy	C. G. H. Sk 81
'2 tñtia	five-angled	«. J gr	1 jn.jl	Gy	17.VL Sksl
8385 rig'tla II.	twisted	tL f gr	1 jn.jl	Gy	C. G. H. 1800. Sksl
pseudo-rigida Sa/m	rigid	«. J gr	f ap.my	Gy	C. G. H. 1820. Sks.l
8386 aj.pgra Haw.	rough	tL J gr	I jn	Gy	C. G. H. 1795. Sks.1
8387 asj>erula //aw.	pale roughish	tL i_J gr#	I jn.jl	Gy	C. G. H. 1824. Sks.1

8188 bicarinata <i>Haju.</i>	double-keeled	tt-i ₁ tc ₁	1	jin	C. G. H.	1820.	Sks.1
83K9 <i>Tagra Haw.</i>	ro?igh-b'ack	tL\ gr	1	■	C. G. H.	1823.	Sks.1
83D0 imbricata <i>Haw.</i>	imbricated	tL\ T gr	■	■	C. G. H.	1731.	Sks.1
^locspirklis Aaf.							Bot. mag. 14A5
8391 foholbsa <i>Haw.</i>	small-leafy	a. I ₁ gr	1	jn.au	Gy	C. G. H.	1795.
* 1016. J.I'LIUMZ.	LILY.	*	(<i>Li</i> ,	whiteness.	Celt.	flowers.)	Tulipuccte. 26.-28.
8392 candid urn <i>L.</i>	white	A or	3	Jn.jl	W	Levant	1556. O r.m
2 bistratium	striated-/tv<tf	A or	4	Jn.jl	W		Bot. mag. 278
3 "fcipatum	spiked	tf ^ or	4	Jn.jl	W		
4 vaneg.iltum	variegated	tf A or	4	Jn.jl	W		
H.W3japonicum <i>Thun.</i>	Japan	tf A or	■	Jl.au	W	China	1804.
83D4 longiflbrum <i>Thun.</i>	long-flowered	tf A or	2	myjn	W	China	1820.
83? 3' " nepalense <i>Wai.</i>	Nepal	tf A or	3	Jn.jl	W	Nepal	O r.m Bot reg. 560
fcJl/G carolinianum Mr.	Carolina	tf A or	2	Jl'au	O	N. Amer.	1825. O p.l
autumnMe 7ord							Bot reg. 580
8C97 bulbiferum <i>L. com.</i>	orange, orbulb-bgtf	A or	3	jn.jl	O	Italy	1596.
2 umbclfttum	umhe\ -jtwd orange	tf A or	3	jn.jl	O	Italy	1596.
838 latifolium ZA:	broad-lvd-gii/6i/.	tf A or	2	Jn.jl	O	Europe	1820.
8J99 lanciflum <i>Thun.</i>	lance-leaved	tf A or	4	Jl.au	O	Japan	18J0. O r.m
8100 spectabile <i>Fis.</i>	showy	tf A or	2	jn.jl	L.0	Dauria	1754.
pennsylvanicum <i>B.M.</i>	daCircum <i>Ker</i>					O p.l	Bot mag. 872
8401 unicolor <i>Sal.</i>	self-colored	tf A or	2	Jl	R	China	1806.
8402 Catesbeatt Cur.	Catesby's	tf A or	1	Jl.au	S	Carolina	1787.
8403 philadelphicum L.	Philadelphian	tf A or	5	Jl.au	S	N. Amer.	1757.
SW4 amrtnum M*.	Andes	tf A or	4	Jl.au	S	N. Ainer.	1819.
8405 cuiuilemce L.	Canadian	tf A or	4	Jl.au	L.0	N. Amer.	1629.
2 riourum	rciUjtawcrcd	tf A or	4	Jl.au	Q	N. Amer.	1822. O p.l
8406 pendulifluru'm Red.	pendulous-flwd	tf A or	1	Jn.jl	Q	N. Amer.	1820. O r.m
8407 superbum <i>Lam.</i>	biptcrb	tf A or	6	Jn.au	■	N. Amer.	1727. O p.l
8408 Murtagon <i>L.</i>	Martngon	tf A or	3	Jl.au	P	Germany	1596.
2 duber>cns	pubescent	tf A or	3	Jn.au	O	Germany	1506.
8409 glabrum <i>Snr.</i>	smooth	tf A or	4	Jn.au	O		O r.m
8410 croccum <i>JBernh.</i>	saffron	tf A or	3	Jl.au	Y		
8411 chalcedumcum L.	Chalcedonian	tf A or	2	Jl.au	S		1596. O co
8(12) pyrenacum <i>Gou.</i>	Pyrenean	tf A or	2	Jl.au	D.0	Levant	1596. O p.l
8-11) Pomphnium L.	Scarlet Pom pom	tf A or	2	myjn	R	Pyrenees	1596. O p.l
8414 monadllphum Bleb.	monadclphous	tf A or	2	Jn.jl	Y	Siberia	1629. O r.f
8-15 tigrinum <i>Ker</i>	tiger-spotted	tf A or	6	Jl.s	O	Caucasus	1600. O r.l
8410 pumilum <i>Bed.</i>	dwarf	tf A or	1	jn	S	China	1804. O r.l
8417 tenuiflbum <i>Bieb.</i>	fine-leaved	tf A or	2	Jn.jl	S	Dauria	1816. O r.l
* IC17. TULIPAL.	TULIP.					Caucasus	1820. O p.l
8418 sylv6strisL.	wild, Russian	tf A 8F	1	ap.my	Y		
8819 repens <i>Fis.</i>	creeping	tf A or	2	ap.my	Y	England	ch.pit O sl Eng. bot 63
84J0 tfircica Both	Turkish	tf A or	2	ap.my	St	Russia	1819. O r.m
84*1 Cehidna Schr.	Ccls's	tf A or	lijn.jl	Y	■		O sl Sw. fl. gar. 186
8422 cornbta Bed.	horned	tf A or	2	my	St	Levant	1816. O r.m Bot. mag. 71
acuminata Vahl							
8423 altaica Pall.	Altaian	tf A or	1	ap.my	Y	Levant	
8424 biflora Pall.	two-flowered	tf A or	1	ap	Y	Altai	...
8*25 tricolor Sm.	three-colored	tf A or	1'	ap.my	S	Russia	1806. O r.m Bot reg. 535
8126 Aulinus Sm. St. Am.	sun's-eyc	tf A or	1	ap	R.D	Russia	1817. O r.m
**W7 Gesme/Vna L.	Gesner's	tf A or	2	ap.my	St	Italy	1816. O s.l Bot. reg. 204
2 versicolor	party-colored	tf A or	2	ap.my	Va	Levant	1577. O r.m Bot mag. 1135
3 plena	double	tf A or	1	ap.my	Va		
4 laciniata	cut-sepaled	tf A or	2	ap.my	Va	Levant	1803. O r.m
5 lutea	yellow	tf A or	11	ap.my	Y	Levant	160.'. O r.m
7428 montana B. R.	mountain	tf A or	1	jl	S	Levant	1603. O r.m
J4J1 sujvcolens Roth	sweet-scented	tf A or	1	ap.my	Y	Persia	1826. O r.m Bot. reg. 1106
2*10 pnc'cox Ten.	eaxy,flowering	tf A or	1	ap.my	S	S. Europe	1M3. O r.m Bot. mag. 8J9
JUL puhescens W.	pubescent	tf A or	1	ap.my	R	Italy	1825. O r.m Sw. fl. gar. 157
JJ32 Clissu:/rt Red.	CIUSUH'S	tf A or	1	jn	W.p	ISL4. O r.m Sw. fl.gar. 78	
84J3 btellata B. M.	starred	tf A or	2	mr.ap	V	Sicily	1636. O r.m Bot. mag. 1390
* 1018. FKITILLA^RIA	L. FHITILLART.					Kumana	1827. O r.m Bot. mag. 2762
8*34 imperialis L.	Fritillaria	tf A or	4	mr.ap	D.Y		
1 ratora	Crown Imperial	tf A or	4	mr.ap	Persia	1596. O co	
2 flvka	red-Jlowered	tf A or	4	mr.ap	R	Persia	1596. O co Bot. mag. 194
8435 fersica L.	yeWofv-Jlowered	tf A or	4	mr.ap	Y	Persia	1596. O co Bot. mag. 1215
2 minimia Swert.	Persian	tf A 8F	1\$	ap.my	Br	Persia	1596. O co Bot. mag. 1537
J<6 Obliqua B. M.	least Persian	tf A or	1	ap.my	Br	Persia	1596. O co Bot. mag. 962
2+* tulipiflair Bieb.	oblique-toivn/	tf A or	1	ap	Br.P	Caucasus	1596. O co Bot. mag. 857
8) > vertinirita W.	Tulip-leaved	tf A or	1	my	Br.P	Crimea	1822. O cq
5*8 ruthenica Wik.	whorled	tf A or	1	ap'	P	Crimea	1823. O co
8*80 pyrenaica L.	Russian	tf A or	1	my	P	Caucasus	1826. O l.p
Pyrencean		II	myjn	D.P	1605. O co	Bot mag. 953
8*81 n ^g ra B. M.	black	tf A or	1	my	Y.p	Pyrenees	1596. O co Bot mag. 664
piet>I'yrenäica J2.3.							
9H2 v. en.	nervcd-leavcd	tf A or	1	my	D.P	Caucasus	1826. O co Bot. mag. 853
minor B. M.							
11Mow-flowered	tf A or	1	ap.my	Y	Caucasus	1812. O co	Bot mag. 1538
Lusitanian	tf A or	1	myjn	Br.P	Italy	1825. O co	
11LT broad-leaved	tf A or	1	ap.my	R	Spain	18J5. O co	
11J7 meleagris L.	guinea-fowl	tf A or	1	mr.my	P	Caucasus	1604. O co
g^ meleagris 3 alba Hort.	early white	tf A or	1	mr.my	W	Britain	moi. p. O co
g^ meleagris 3 alba Pers.	Melcagris-like	tf A or	1	apmy	P	Europe	- O co
11W nvclcggo Ules Fis.	slender	tf A or	1	my	P		
11*0 tenella Bieb.							
racemosa a. minor B. M.	white-flwd. Russian	tf A or	1	ap.my	W	Siberia	1824. O l.p
>ll 'eucantha Us.	white American	tf A or	1	ap.my	W	Siberia	1822. O l.p
Eir? ?	spear-leaved	tf A or	f my	D.P	N. Amer.	- O l.p	
11J lanceoCita Ph.					Karaschat	1759. O co	Lin. tr. 10.11
84JU fli^junc kamschatc6nsc L.	chaste	tf A <r	1	apmy	py	o	, p
o** pudica Sm.						p n a m . ,	

1019. CHARLWOOT>/i4 Swt. CHAHLWOODIA. (<i>G.Ciarlzoood</i> , F.L.S., an enthusiastic Eng.lmt.) <i>Asphodellea</i> . 4.											
8455 conglsta <i>Swt.</i>	crowded	I	CDor	mr	P.B	N. Holl.	am.	X	Sw. au	18	
8456 stricta <i>Swt.</i>	upright	I	CDor	mr	B	N. Zeal.	am.	X	Bot. reg.y56	1820.	
Dracaenastricta <i>B.M.</i>											
8457 indivisa <i>G. Don</i>	undivided	I	d or	10	—	N. Zeal.	—	C	p.l		
Dracunivisa <i>Font.</i>											
8458 australis <i>G. Don</i>	southern	I	CDor	10	—	B.W	N. Zeal.	1823.	C	p.l	
Dracaena australis <i>Forst.</i>											
•1020 DRACENA L. DRACJENA. (Drufczina, she-dragon; juice like dragon's blood.) <i>Asphodellear</i> . 18.—20.											
8459 DT\CO /,. Dragon, or common	f or 10	—	W	E. Indies	1640.	C	p.l	Black.	538		
8460 cnstifflia Jfeur.	sword-leaved	I	Q or	4	—	W	—	—	1800.	C	p.l
8461 arborea Lk.	tree	I	CDor	—	—	W	—	S. Leone	1800.	C	p.l
8462 rffigrans Ker	sweet-scented	I	CDor	—	—	W	—	Africa	1768.	R	p.l
8463 tessellata W.	tessellated	I	CDor	—	—	W	—	Madagascar	1816.	C	p.l
inarginata Lam.											
8464 cernua Jac.	drooping	I	CDor	10	my	W	Mauritius	...	C	p.l	Jac. sc. 1. 96
8465 rflexa Lam.	reflexed-leaved	I	CDor	4	myjl	W	Madagascar	1819.	C	p.l	
8466 ferrca K/.	iron	I	CDor	8	mr.ap	W	China	1771.	II	p.l	Bot mag. 2053
8467 teriminilis Jac.	terminal	I	CDor	10	jnjl	W	E Indies	1820.	C	p.l	Jac. ic. 2. 448
8468 purpurea Lo. C.	lur>lc-leaved	I	CDor	15	jnjl	W	E. Indies	1820.	C	p.l	
8469 umbrarulifera Jo*	umbrella-bearing	I	CDor	10	—	W	Mauritius	1788.	C	p.l	Bot cab. 289
8470 ovata B. M.	ovate	I	CDor	2	—	W	S. Leone	1798.	C	p.l	
8471 interrupta Haw.	interrupted	I	CDor	2	jnjl	W	S. Leone	1798.	R	p.l	Bot mag. 1180
8472 leoiicitis Lo. C.	Sierra Leone	I	CDor	3	jnjl	W	& Leone	1824.	C	p.l	
8473 notosa Lo. C.	knotted	I	CDor	4	ap.my	W	C. G. H.	IK20	C	p.l	
8474 triatia Lo. C.	striated	I	CDor	4	ap.my	W	Mauritius	1825.	C	p.l	
8475 mauritiana Lam.	Mauritian	I	CDor	6	—	W	C. G. H.	1816.	R	p.l	
8476 undulata L.	waved	I	CDor	4	jl.au	B	N. Holl.	18'0.	C	p.l	
8477 nūtans Cun.	nodding	I	CDor	4	jl.au	B	N. Holl.	18'0.	C	p.l	
1021. PHYLLO ^X MA Haw. PHYLLOMA. (<i>Phillion</i> , leaf, <i>loma</i> , fringe; cld-toothed edges.) <i>AsphodelUce</i> . 4.											
8478 noltiform Haw.	Aloe-flowered	f or 10	ap	O	Bourbon	1766.	R	p.l	Bot. mag. 1585		
Dracaena marginata //, K.	Lomatophyllum borbonicum	W.									
8479 Jacquinii G. Don	Jacquin's	i.J or	4	jn	O	—	—	Sk p.l			
jj'loz hexaiHjtala Sal.	lomatophyllum	Jacquinii Spr.									
8480 rufocinctum AW.	rufous-edged	i.J or	3	jn	O	E. Indies	1818.	Sk.s.l			
A'oil rufocincta Haw.											
8481 m'cum Lk.	thin-leaved	s> _Jor	3	jn	O	Mauritius	1817.	Sk.s.1			
'A'loc macra Haw.											
1022. ALE ^{vnt} ISL.	ALETRIS.	(Aleton, meal; covered with powder)				<i>Hemerocallidea</i> . 2.					
8482 farinosa L.	mealy	I	A or	i	jn	W	N. Amer.	1768.	R	s.p	Bot mag. 1418
filba Mx.											
8483 afirca Mx.											
1023. TRI'TOMA Ker	TRITOMA.	(Treis, three, temno, to cut; angles of leaves.) <i>Hemerocallidete</i> . S.									
8484 Uvaria Ker	Uvaria	* JVJ or	2	a.u.s	O	C. G. H.	1707.	R	p.l	Bot. imag. 758	
8485 media Ker	mealy intermediate	tf	JJ or	2	year	O	C. G. H.	1789.	R	p.l	Bot. mag. 744
8486 piimila Ker	dwarf	je	AJ or	1	s.n	O	C. G. H.	1774.	R	p.l	Bot. mag. 764
1024. VELTHEIM' M/i1 Gled. VELTHIEIMIA. (F.A. de Veltheim.a. German botanist) <i>Hemcrocallide</i> *. 3.-4.											
8487 viridiflora Jac.	green-leaved	tf	LAJ or	2	ap.n	F.w	C. G. H.	1738.	O	r.m	Bot. mag. 501
8488 intermedia Hort:	intermediate	tf	iAl or	1*	ap	F.w	C. O. H.	1800.	O	r.m	
8489 glaCica Jac.	glaucous	tf	LAJ or	2	j&ap	F.o	C. G. H.	1781.	O	r.m	Bot. mag. 1091
1025. SANSEVIE ^N JU Thun. SANBBVIERA. (<i>Sunsevter</i> , a Swedish botanist) <i>Hemeroallidca</i> . 14.											
8490 glaCica Haw.	glaucous	je	EI cu	2	—	W.o	—	—	Sks.l		
8491 stenoph? ila Lk.	narrow-leaved	je	EI cu	3	—	—	—	1818.	Sks.l		
8492 polyphylla Haw.	many-leaved	je	E3 cu	2	—	W.o	—	—	Sks.l		
8493 gumeeflisis IV.	Guinea	je	E) cu	2	jn.n	G	Guinea	1600.	Sk s.p	Bot mag. 1180	
8494 longiflora Sims	long-flowered	je	J2J or	2	Jl	W	Africa	1824.	Sk co	Bot mag. 2634	
8495 he'te v'irens Haw.	lively-green	je	E3 cu	2	—	W.o	—	—	Sks.p		
8496 fulvochneta Haw.	fulvous-edged	H	E3 cu	1	—	—	Brazil	1820.	Sks.p		
8497 spicata Haw.	spiked	je	Ef cu	2	—	W.o	E. Indies	1790.	Sks.p	Cav. ic. 3. 246	
8498 zeylanica IV.	Ceylon	je	ES cu	2	jn.n	W.o	Ceylon	1731.	Sks.p	Bot. reg. 160	
8499 ensifolia Haw.	sword-leaved	je	Ef cu	1	—	T	E. Indies	—	Sks.p		
8500 lanuginea W.	woolly	je	E1 cu	2	—	W.o	E. Indies	—	Sks.p	R.maLII.42	
8501 grandiflora Haw.	large-pointed	je	C3 cu	3	—	W.o	—	—	Sks.p		
8502 ptmilia Haw.	dwarf	je	Ef cu	1	—	W.o	CG. H.	1796.	Sks.p		
8503 cjmica Andr.	sessiflora B. M.	je	Ef cu	1	—	W.o	China	1792.	Skl.p	Bot. rep. 361	
1026. TULBA [%] GH/J4 L. TULBA«HIA. (<i>Tulbagh</i> , a Dutch governor at the C. G. H.) <i>Hemeroallide</i> ee. 3.											
8504 alliacea L.	Garlic-scented	tf	iAJ or	1	myjl	Br	C. G. H.	1774.	O	r.m	Bot. mag. 806
8505 ainnis Lk.	allied	tf	LAJ or	1	my jl	Br	C. O. H.	1820.	O	r.m	
8506 cepacea L.	Onion-scented	tf	iAl or	J	ap	Br	C. O. H.	1795.	O	r.m	
• 1027. YU'CCA L. ADAM'S NEEDLE. (<i>Yuca</i> , its na in St Domingo.) <i>Tulipaceat</i> . 19.— 22.											
8507 gloriiflora L.	glorious	li	or	4	jl.au	W.o	Aiaocica	1596.	Sk r.l	Bot mag. 1260	
8508 acuminata Swt.	acuminate-/scud	m	or	6	au.8	W	—	1800.	Sk r.l	Sw.fl.gar.ic.in*	
8509 aloifolia L.	Aloe-leaved	m	Ll or	2	a.u.s	W.o	a Amer.	1696.	Sk r.l	Bot mag. 1700	
8510 temufloria Haw.	slender-leaved	m	or	1	—	W.o	Malta	1817.	Sk r.l		
8511 draennis L.	dragon	m	LJ lor	8	au.s	W.o	& Amer.	1732.	Sk r.l	Di. eL 324. 117	
8512 concava Haw.	homMow-leaved	je	A or	lftau	W	o	1816.	Sk r.l		
8513 oUqua Haw.	obWque-leaved	je	A or	4	—	W.o	1808.	Sk r.l		
2 major T	larger	je	A or	4	—	W.o	1808.	Sk r.l		
8514 flaccida Haw.	flaccid	je	A or	2	—	W.o	1816.	Sk r.l		
8515 fierrulkta Haw.	scrurlatc-edged	m	Ll or	10	—	W.o	Carolina	1808.	Sk r.l		
8516 recurva Sal.	recurve-Jeaved	m	or	3	au	W.o	Georgia	1794.	Sk r.l	Par.Ion.31	
8517 supērba Haw.	superb	m	LJ lor	10	au	W.o	—	—	Sk r.l	Bot. rep. 473	
8518 glaucescens Haw.	glaucescent	m	or	2	jl.au	W.o	N. Amer.	1819.	Sk r.l	Sw. II. gar. 53	
8519 filamentosa L.	thready	m	A or	2	S.0	W.o	Virginia	1675.	Sk r.l	Bot mag. 900	
8520 riifo-cfncta Haw.	rufoso-edged	m	A or	Jll	W.o	—	—	1816.	Sk r.m		
8521 stricta Ker	upright	m	or	Jl	W.o	Carolina	1817.	Sk r.m	Bot mag. 22»		
8522 conspicua Sal.	conspicuous	a	LJ lor	3	—	W.o	—	1818.	Sk r.m		
8523 angustifolia Nui.	narrow-leaved	m	or	2	jl.au	W.o	Missouri	1811.	Sk r.l	Bot. mag. & ^{rtBIA}	
8524 crenplata Haw.	crenulate	m	or	2	—	W.o	—	1818.	Sk.s.L		
8525 arctata Haw.	bowed	m	or	1	jl.au	W.o	—	1817.	Sk.s.1		

- | | | | | |
|---------------------------------|----------------------|---|----------------------------|--|
| 1042. EITCOMIS Herit. | Encovia | (<i>Eukomes</i> , beautiful-haired; tufted crown of fl. spike) | <i>Asphodèle</i> | 7. |
| 8680 nana H. K. | dwarf | jp IAI <w f frly | Br C. G.H. 1774. | O r.m Bot mag. 1405 |
| 8581 <i>purpuracafilis</i> Dry. | purple-stalked | jp IAI or 2 | G.n C. G. H. 1794. | O r.m Bot. rep. 569 |
| 8582 <i>hirta</i> Jac. | two-leaved | ^ AAJ or | Iap.my L.G. C. G. H. 1792. | O r.m Bot. mag. 840 |
| 8583 rigida JT. | royal | AAJ or | 2 mr.ap G C. G. H. 1702. | O r.m Di. el 92.109 |
| 8584 <i>uiululata</i> H. K. | Vfjavcd-leafed | AAJ or | 2 mr.ap G C. G. H. 1760. | O r.m Hot. mag. 1083 |
| 8585 punctata /ferrf. | ((Kitted | AAJ or | 2 j.l G.B. C. G. H. 178a. | O r.m Bot mag. 913 |
| 8586 stnata H. K. | striated | AAJ or | 2 year G C. G. H. 1790. | O r.m Bot mag. 1539 |
| 1043. PELIOSA'NTHES Andr. | PELIOSANTHES. | (<i>Peltos</i> , livid, <i>anthos</i> , flower.) | <i>Melanthacecc.</i> | 2. |
| & 8687 hemis Andr. | dwarf | jf (2S cu \ my.jn | E. Indies 1809. | D r.l Bot. mag. 1532 |
| 8588 <i>Teta</i> B. M. | Teta | f ES' cu \ ap | B.G. E. Indies 1807. | Sk s.p Bot mag. 1302 |
| 1044. APHYLLA'NTHES L. | LILY PINK. | (<i>Aphyllos</i> , leafless, <i>anthos</i> , flower.) | <i>Asphodelé</i> * | 1. |
| 8589 monspeliens JB L. | Montpelier | jf Ad pr 1 jnjl | R France 1791. | R s.p Bot. mag. 1132 |
| 1045. SOWERBYEM Sm. | SOWERDAA. | (James Sowerby, F.L.S. H.S., an emin. bot. artist.) | <i>Asphodelea</i> | 1 |
| 8590 Junccea Sm. | Rush.leaveU | jf IAI br 1 myjl | Pk N. S. W. 1792. | R s.p Bot. mag. 1104 |
| * 1046. /TILLIUM L. | GARLIC. | (iflf, hot, Cxw.) | <i>Asphodelle</i> | 150. |
| 8591 sativum L. | cultivated | tf A cul *i J^n jl | W Sicily 1548. | O r.m M. h. 4.15.9 |
| 8592 Ophioscorodon CDon | Gar. Rocambole | tf A clt 4 a.us | Pa.R Greece ... | O r.m M. h. 2.15.10 |
| Ecordobrasum fi L. | | | | |
| 8593 controyrsuin Schr. | contrary-strnd | ■ A pr 1 Jnjl | P | 1816. O co |
| 8594 arenriuin L. | sand | ■ A pr Jn | P | Britain moi w. O co |
| 8595 ScorolopraHim L. | Rocambole | ■ A cul Jl | L.P | Denmark l>96. O co |
| 8590 littoreum Bert. | sea-Hide | ■ A cu Jhau | P | Italy 1818. O co |
| | W.cn, | | | |
| 8597 vine.ile L. | vine | ■ A pr | P | Britain mea. O co |
| 8598 confertum Pers. | crowded | ■ A cu | P | Europe . O co |
| campestre Schl. | | | | M. h. 2.15.3 |
| 8599 Ptnhfl G. Don | Pursh's | tf A or 2 au | Pk.P N. Amer. 1818. | O co |
| vineale Ph. | | | | |
| 8600 proliferum Schr. | proliferous | tf A cu 3 jl.o | | 1820. O co Bot. mag. 1469 |
| 8601 t'pa L. | common Onion | tf Q cul 3 jnjl | | ... S r.m |
| 2 aggregatum G.Don | aggregated | tf A cul 1 jnjl | | ... () co |
| 3 paucituruin Ttev. | few-flowered | tf A cu 2 jnjl | | O co |
| | Ixck | tf Q cul 2 a p. my | | |
| 8602 PtTrum L. | Vine Leek | ■ A cu 2 j.lau | | Switzerl. 1562. S r.m Black. 421 |
| HUH rofundum L. | round | ■ A pr 14 jl | | England sun.hi. O co Eng. bot 1657 |
| 8C05 HfiUcri (x. Don | Haller's | ■ A pr 2 j.lau | | S. Europe 1820. O co |
| SKHJ rubicindiu Niu. | reddish | ■ A pr 1 j.lau | | P.R S. Europe 1818. O co Fl. gr. 344 |
| tiiUJU Waldstcini G. Don | Waldestein's | ■ A or 2 Jnjl | | C. G. H. 1804. O co Bot. mag. 1560 |
| ^rnpelipfr.isum W. 8f K. | | | | Hungary 1826. O co W. & K. 82 |
| 8608 exuterm G. Don | drawn out | tf A pr 3 jn.au | | |
| 8J19 crectum G. Don | upright | tf -Alpr 3 jnjl | | |
| 8flO ncutiflorum Los. | acute-flowered | tf A pr" i j | | |
| 8611 Synnot's Gi. Don | Synnot's | tf -A pr" U J^n au | | |
| 8112 verrubrum G. Don | w <rty-scpa/ed | -A pr" 14 J | | |
| HGLI multilldrum Drcf. | many-flowered | tf A pr 14 J | | |
| 8<il4 desceulmens /,, | d d d d d | tf A pr | | D.P Switzerl. 179a co Bot mag. 251 |
| 8il5 bpha. r(Mephalon L. | round-headed | tf A pr | | R Europe 1759. co Bot. mag. 1764 |
| 8616 pnmuin IJc. | frosted | tf A pr | | R Spain 1819. co |
| 8517 carneum IV. en. | ilioAi-colored | tf A pr | | R Odessa 1819. co |
| W>18 guttMum Steu. | spotted | tf A pr | | V Palestine 1548. L r.m M. h. 4.14. 3 |
| 8519 ascal6mcum L. | Askalon, or Shallot | tf A cul V | | S. Europe 1804. O co Mat. op. 420. ic. |
| 2 mifus | greater, or Scallion | tf A clt | | F Britain m. pas. O r.m Eng. bot 2438 |
| 8620 Schanoprasum L. | Chives | tf A cul V myjn | | |
| tenuiflum Sal. | r | | | |
| 8621 sibiricum W. | Siberian | tf A pr 1 jl-au | | |
| glaucum Dcsff palustre Sal. | | | | |
| 8622 acut um Spr. | acute | tf A pr 1 Jl | | |
| 8623 folibusu tied. | leafy | tf A pr | | |
| 8624 fUltibium L.* | fistulose | tf A cul lft ap.my | | |
| 8M5 ccapaform G. Don | Ccpa-form | tf A cul 2 jl.s | | |
| obHquum L. | obWque-leaved | tf A pr Ujnjl | | |
| 8f27 rainosum G. Don | branchy | tf A or 2 Jn | | |
| 8628 PalKiaii Mur. | Pallas's | tf A pr | | |
| 8i29 pUKilm Cilr. | small | tf A pr | | |
| 8630 rubellum Bieb. | reddish | tf A pr * Jl.au</td <td></td> <td></td> | | |
| 8b31 reticulatum Fra. | xQ,roasted | tf A pr 1 Jnjl | | |
| BtcKjpm /3 lbum B.M. | angulifloraPA. | | | |
| 8V>2 oxyptalmum G. Don | sharp->etuled | tf A or S au | | |
| 8633 intermedium Lam. | intermediate | tf A or 2 au | | |
| paniculMum VU. | | | | |
| 8634 oleraccum L. | oleraceous | tf A pr 1 Jl | | |
| viri'scens Lam. | | | | |
| 8635 carinatum L. | keeled | ■ A pr i myjn | | |
| 8636 ffexum G. Don | twisted | ■ A pr J | | |
| foytidum W. en. | | | | |
| 8637 violaceum W. en. | violet | ■ A pr 1 J | | |
| 8i28 aspermum G. Don | rough | ■ A or J | | |
| 8639 longspathum Red. | long-spathed | ■ A pr 3 jl.au | | |
| 8640 ffscum Kit. | brown-flowered | tf A pr 2 jl.au | | |
| 8641 paraculatum L. | pamled | tf A pr 2 jnjl | | |
| 8642 pällens L. | Vd-c-Jlowered | tf A pr ? J J.l | | |
| 8643 tenuiflum Ten. | slender-flwd | tf A Pr 1 J | | |
| 8644 pulchellum G. Don | pretty | tf A pr J | | |
| paniculMum lied. | | | | |
| 8645 flavum L. .. | yellow | ■ A pr 1 Jnjl | | |
| 8646 caucasicum Bieb. | Caucasian | ■ A pr Jnjl | | |
| ff Z ^V?*™ Mb. | rock | ■ A pr R | | |
| 8648 globosum Bieb. | globose | ■ A pr 1 au.s | | |
| 8650 montanum Sm. | Bmally-flowered | ■ A br 1 jnjl | | |
| mountain | | ■ A pr R | | |
| | | | | nift |
| | | | | Fl. gr. 319 |

J. Jwlmoschatum Z.	musky	5 A pr	iau.s	w.p	S. Europe	1823.	O co	W. & K. J. 68	
8652 saxatile Bieb.	rocky	1	j.lau	W	Crimea	1823.	O co	"M	
8053 strlctum Schr.	strict	tf Ap	1	Pk	1821.	O co		
... deflexum W.									
... lincare L.	linear-fe<w>s	tf Apr	1 jn.jl	W.ii	Siberia	1753.	O co	Gm. si. 1.13	
of 5 nutans L.	nodding	S A pr	1 j.lau	R	Siberia	1785.	O co	Bot. mag. 1143	
... senescens L.	senescent	tf A pr	Jjnjl	W	Germany	1596.	O co	Gm. si. I. 11. 2	
... Andershs G. Don	Anderson's	tf A pr	1 jn.jl	P	Siberia	1818.	I) co		
...>8 spinum G. Don	spurious	J A Dr	1 jn.jl	P	Siberia	1820.	I) co		
oos Klciuem Schr.	glaucous	tf A or	* Jjnjl	Pk	Siberia	1800.	D co	Bot mag. 1150	
bicalense W. en.	spirale IV. en.	senescent Br.M.							
... angulsum K.	anguiose	tf A or		L.P	Germany	1739.	O co	Bot. mag. 1149	
... laxum G. Don	loose	tf A pr	1 jn.jl	R	Siberia	1817.	O co		
wfS acut&ngulum IV. en.	acute-angled	tf A pr		R	1816.	O co		
f GJ danubile S>r.	Danube	tf A pr		R	Germany	1818.	O co		
<t4 seritum G. .	late-flowring	tf A pr		R	Switzerl.	1820.	O co		
J*15 illidum AfA	whitish	tf A pr		W	Crimea	1K20.	O co		
806f denudatum Dec.	stripped	tf A pr		W.u	Siberia	1819.	O co	Red. lil. 357	
flavocenso.		tf A Pr							
8667 rikbiu Schr.	red-rooted	tf A pr	1 jn.jl	P	Germany	1805.	O co	Bot. mag. 1381	
bisulcum B. M.									
8G88 prostratum Trev.	trailing	tf Apr	i jn.jl	R	Siberia	1823.	O co		
9ID congustum G. f»«	crowded J-wrf	tf A or	1 my.jn	P	Siberia	1818.	O co		
JIM narcissi fltrum Vit.	Narcissus-flwd	tf	1 j.lau	P	France	O.r.m Vil. del. 2. G		
TM71 pedemontaium IK.	Picdmontsc	tf	1 j.lau	R	Piedmont	1817.	O co Al. ped. US. 1		
8C72 sunvt-olcna J*	weet-smelling	tf	1 j.lau	W	Austria	1801.	O co Jac. JC. «. 364		
- nrnhf'iuuih hm. Graminifolium Pow. meitiruin Thore									
2*73 oehrolciicum W.S/K.	yellowibh white	tf A P'	1 jl	Crica	Hungary	181(1	O co	W. & K. 186	
oeJ *Cerilius II <th></th>		drooping	tf A br	1 jn	R	N. Ainer.	1804.	O co	Bot. mag. J324
TM5 t'aroliuMiniTn Dec.	Carolina	tf A P'	% J>	U	N. Ainer.	1818.	O co	lied lil. 100	
<>70 stellatum B. M.	starred	tf A pr	1 ap	W	N. Amur.	1811.	O co	Bot. mag. 1570	
8G77 para<16xum G. Don	wonderful	tf A P'			Caucasus	1823.	O co		
Scilla paradisiaca Bieb.									
8>79 mafolifin m'fligie, Df. Strozzera, sifif A op'r	roseate	tf A br	1 jn.jl	Y	S. Europe	1820.	O co	Red. lil. 2.99	
8680 cahadencs L.	Canadian M	tf pr	1 jn.jl	G.w	Austria	1518.	O co	Bot. mag. 1148	
8681 mutabile Mx.	changeable	tf A P'	1 jn.jl	P	N. Amer.	173a.	O co		
8681 incarnatum Horn.	ticxh-co/orcd	tf A l'r	1 ll	R	N. Amer.	1824.	O co	Red. lil. 240	
rftscum £. B. It. ambfguum Sm.				R	Greece	1812.	O co	Bot. reg. 978	
8H83 ainc^nun G. Don	plouising	tf A pr	1 my.jn	R	France It.	1821.	O co	Fl.nap.28	
carneum Bert.									
2*4 rtaeum L.	roseate	tf A br	1 jn	Pa.P	France	1752.	O co	Red. lil. 4.213	
2*3 longifolium Kth.	long-leaved	tf A l'r	1 j.lau	UP	Mexico	1826.	O sl	Bot. reg. 1034	
JGJ illyricup Jac.	Ilynan	tf A or	1 jn	P	Austria	1820.	O co	Jac. ic. 2. 365	
*67 subhirsutum L.	rather hairy	tf A P'	1 my	W	S. Europe	1596.	O co	Bot mag. 774	
cildtum Cyr.									
aS8 Clusianni Retz.	Clusius's	tf A pr	\$ jn.au	W	S. Europe	1803.	O co	Cl h. 1.192	
2*9 Chamsmluy L.	Dwarf Moly	tf A Pr	1 jn	W	S. Europe	1774.	O co	Bot. mag. 1203	
5*w brachyslemon Dec.	short-stamencd	tf A °r	1 my.jn	W	Europe	1819.	O co	Red. hi. 7.374	
S? H? Ten.				W	Italy	1824.	O co	Fl. nap. 28	
SP2 caspium Bieb.	Caspian	tf A br	1 ap	W	Crimea	1822.	O co		
.... Amarjih* caspia L.									
ewe Cowkni Lindl.	Cowan's	tf A Pr	I jn	W	Chile	1823.	O co	Bot. reg. 758	
RISS "ob>num Cyr.	Neapolitan	tf A P'	1 Jl	W	Naples	1823.	O co	Sw. fl. gar. 201	
triquetrum L.	triangular-f/aff	tf A Dr	I my.jn	W	Spain	1759.	O co	Bot mag. 869	
Endurmum Ten.	drooping	2u>tf	f ap my	W	Italy	1825.	O co	Fl. nap. 31	
G.Don	intermediate	tf A °r	1 my.jl	W	Hungary	1820.	O co		
	b'ack-ovaried	tf A pr	fil.au	W	Barbary	1818.	O co		
	dark-purple	tf A pr	M	V	Hungary	1821.	O co	W. & K. 1.17	
	three-seeded	tf A l'r	Jl	P	S. Europe	lii4.	O co	Bot mag. 499	
2J 1 tncjecum H. K.				W	N. Amur.	1770.	O co		
2J < ursim L.	bear	tf A br	ap.my	W	Britain woods.	O co	Eng. bot. 122		
0703 Victionis L.	Victorialis	tf A cu	1 my	G.v	Austria	1739.	O co	Bot mag. 1222	
ftTn 2 aiugistif-Mium	narrow-leaved	tf A cu	1 my	GY	Scotland	1819.	O co	Cl. pan. 224	
°01 odbrum L.	sweet-scented	tf A Pr	Jn	W	S. Europe	1820.	O co	Bot mag. 1142	
tataricum L.									
2005 uncifolium G. Don	Rush-leaved	tf A pr	Jmy.jn	W	Chile	1826.	O co		
001 striatum Jac.	striated	tf A br	1 my.jn	W	Carolina	1800.	O co	Bot. mag. 1035.1524	
W7 striatellum Limll.	small-striated	tf A pr	lj ap.my	W.v	Chile	1823.	D co	Bot. mag. 2419	
Ornithogalum graminicum B. M.									
ans Vcn.	fragrant	tf A pr	1*8.0	W	W.Indies	1822.	O co		
J0>7 ALBITCA L.	ALBUCA.	(Aldus, white: color of flowers.)							
09 a 1 J	tallest	tf A l or	4 ap'm'y	W	C.G. H.	1789.	O r.m	Jac. ic. 1.36	
Dr	greater	tf A or	3 ap.my	GY	C.G. H.	1759.	O r m	Bot. mag. 804	
J nYmor L.	smaller	tf A l or	1 my.jn	Y	C.G. H.	1768.	O si	Bot. mag. 7d0	
S 3 flaccida Jac.	flaccid	tf l A l or	2 my.jn	Y	C.G. H.	1791.	O r.m	Jac. ic. 2.144	
g 2 virlift J	green-flowered	tf A l or	1 my.jn	G	C.G. H.	1798.	O r.m	Bot. mag. 1656	
dry.	compresso	tf A l or	1 my.jn	Y	C.G. H.	1774.	O r.m		
iastiki Ma Dry.	peaked	tf A l or	1 my.jn	W	C.G. H.	1774.	O r.m	Bot. rep. 450	
RaudHU Jac.	tailed	tf A l or	1 my.jl	W	C.G. H.	1791.	O r.m	Jac. ic. 2.442	
illibia Ker	throAd-leaved	tf A l or	1 my.jl	W	C.G. H.	1820.	O r.m		
asetosa Jac.	bristly	tf A l or	1 my.jl	G	C.G. H.	1795.	O r.m	Bot mag. 14G1	
p vittata Ker	banded	tf A l or	1 U u	Y.c	C.G. H.	1802.	O sp	Hot. mig. 1329	
physodes B. M.	flatulent	tf A l or	1 my.jl	W	C.G. H.	1804.	O r.m	Bot. mag. 1046	
exuviate B. M.	adder's skin	tf	1 my.jl	W	C.G. H.	1795.	O r.in	Bot. mag. 871	
Anthridium exuviatum Jac.									
orea Jac.	golden	tf	Alor	G.Y	C.G. H.	1818.	O r.m		
cc.	Abyssinian	tf	vtor	au	W	Abyssinia	1818.	O r.m	
?gax Ker	sweet-scented	tf	or	ju.jl	Y.G	C.G. H.	1791.	O s.p.Uac. sc. 1.81	
872r Anthuncum fragrans Jac.	fleeting	tf	or	my.jl	Y.o	C.G.H.	1791.	O r.m Bot. reg. 311	
8727 Anthuncum fragrans Jac.	dammy-Zt'lWfd	tf	IAI or	1 my.jn	W.a	C.G.H.	1779.	O r.m Jac. ic. 2.445	
"Piftis L.	smral-leaved	tf	IAI or	K	C.G.H.	1795.	O s.p	Jac. ic. 2.439	

1048. XANTHORRH(E\A Sm. XANTHOMUKEA. (*Xanthos*, yellow, *rheo*, to flow; gum.) *Aphodxilce*. 6.—7.
8728 australis R. Br. southern n. i.) or 3 ap.my W Holl. 182*. Sk s.p
8729 htmilis R. Br. dwarf £ iAI or 2 ap.my W N. Holl. 188⁵ Sk fl.p

pumilio R. ltr. spear tt.L—|cu 4 ap.my W N. S. W. 1803. Sk.s.p
8730 hastihs t< lit. intermediate tt. i_|or 2 ap.my W N. Holl. — Skl.p
8731 media £. 2fr. smaller ■ IAJ ■ 2 ... W N. S.W. 1804. Sk.s.p
8732 minor R. Br. braeted ■ IAJcu 2 ... W N. S.W. 1810. Sk.s.p
8733 bracteata R. Br.

1049 LAXMA'NN/i* R. Br. LAXMANNIA. (Laxmann, a Siberian traveller.) 'AsphodZlear. 1.—8.

8734 gracilis ii. -Br. slender £ iAI or £ jujl P.w N. Holl. 1824. D s.p

*1(J50 THYSANO*rUS.B.2&\ THYSANOTIH. (*Tbyisanotos*, fringed; three inner sepals.) *Asphodel** 4.—21.

8735 iilnceus A. 5f. Rush-/-** tf iAI pr 4 a.us P N. S. W. 18(H. O s.p Bot reg. 656

8736 elatior R. Br. taller A iAI or 1 au P N. Holl. 1823. O s.p

8737 isanthrus R. Br. cycn-anthercd A LAJ pr 4 a.us P N. S. W. 1822. O s.n Bot. reg. 655

8738 tuher+su* R. Br. tuberous A IAJ el 1 jn P N. Holl. 1825. O s.l

1051. EKIOSPE'LLUM Jac. ERIOSPERMOM. (*Erión*, woo], *sperma*, seed; envelope of seed.) *Asphodllear*. 8.

8739 latifohum Juc. hroned-leaved tf iAI cu 1 jn au L.D C. G. H. 1800. Sk s.p Jac. IC. 420

8740 Ueliendeni Suit. Bellenden's tf LAI or 1 jl.au L.B C. G. H. 1800. Sk s.p Bot. mag. 1002

lahtuliuin a.B. M. not Jac. pubescent tf LA(CU ■ W.G C. G. H. 1820. Sk.s.p Bot reg. 578
8742 launginfusin Jac. woolly tf IAI or ■ Jl an W.O C. G. H. 1820. Sk.s.p Jac. sc. 3.265
8743 luncctfMiuni Jac. spear-leaved tf IAI(CU 1 jn.au LB C. G. H. 1795. Sk s.p Jac. ic. 2. 421
8744 parviflMiunJirfc. buial-leaved tf IZU cu 5 jn.au DB C. G. H. 1796. Sk s.p Jac. ic. 2. 422
8745 foliolifercur Ker le.ift-bearing tf IAJ cu fijn.au Y.O C. G. H. 1806. Sk.s.p Bot reg. 795
8746 paradox um Jac. wonderful tf IAJor 1 jn.au ■ C. G. H. 1825. Skip Jac. c. sup. 81.1

Ornithogalum parad6xum Jac.

* 1052. GA^XGEA Sal. GAGEA. (Sir Thoma* Gage, a bot amateur.) *AtpJtod&e** 16.

8747 uniflora G. Don one-flowered tf A or 4 my.jn Y Siberia 1781. O s.p N.c.p.18.6.3

Ornithogalum uniflorum L.

8748 stnata Swt. striated tf A or i my.jl Y Europe 1826. O si

Ornithogalum striatum W.

8749 pygina G. Don pygmy tf A or 4 ap Y Spain 1825. O co

Ornithogalum Jygmac un IV.

8750 bobemich G. Don Bohemian tf A or 4 ap Y Bohemia 1825. O co Ac. boh. 2.121.10

Oriuthigalum bohemicum Schmidt

8751 fasciculiris Sal. hwnUc-Jlwdf. tf A ■ | mr.ap^Y Britain woods. O s.p Bot. mag 1200

late B.M. Ornithogalum futcumf.

8752 sylva'tica IV. en. wood tf A P' | mr.ap Y Europe ... Os.p Us. an. 5.1.1

8753 villofa Bish. villous tf A or | mr.ap Y Caucasus 1S25. O pl

8754 stellaris Sal. starry tf A A|pr | my Y Sweden 1759. O e.p Us. an. 15.1

Ornitigalum minimum Roth. arvense Pert.

8755 Sternberg/ Hoppe Sternberg's tf A or \ mr.ap Y Switfrl. 1826. O pl

8756 bracteolaris Sal. bractolar tf A or ft ap Y Europe 1817. O co Sw. fl. gar. 151

Ornitigalum luteum L. not E. B.

8757 MMihaca IV. fihedthed tf A pr A my Y Germany 1759. O s.p Hay. us. 15.1

8758 thlorAntha Bieb. yellow-flwd tf A or *ap Y Siberia 1819. O co

8759 gUtica Swt. glaucous tf A or *ap Y Switzcr. 1825. O co Sw. fl. gar. 176

8760 pusilla Schmidt small tf A or *ap Y Bohemia 1823. O co

8761 circinMa //. rounded ■ A pr A my.jn Y Siberia 1789. O s.p Pal. it. D 2

8762 ser6tina B. M. late tf A ■ Wales ... O s.p Eng.bot.793

Anthericum ser6tinum E. B.

* 1053. ORKITHO'GALUM L. STABOF BETHLEHEM/ (*Ornis*, a bird, *gala*, milk.) *AphodUece*. 48.-56.

8763 bulbiferum Pall. bulb-bearing tf A or | p. W Russia 1821. O r.m Pal. it. 2. ap. 106

8764 umbellatum L. tnnbclled tf A or 1 apjn W England mo p. O co Eng.bot130

8765 refractum W. refracted tf A or J ■ W Hungary 1820. O pl

8766 timbratiuum W. fringed tf A or £ f.mn W Crimea 182⁸. O co Lind. col. 28

8767 exscapum Ten. stem less tf A or 4 my.jn W Italy 1824. O r.m Fl. nap. 34

8768 montimum Ten. mountain tf A or i my.jn W Italy 1824. O pl

8769 comsum L. tufted tf A or IJn.au W Austria 181. O pl Jac. ic. 2.426

8770 canticum Jac. conical tf A or 1 my.jn W C. G. H. 1823. O r.m Jac. ic. 2.408

8771 nfveum Sol. snowy tf A|or 1 jnjl W C. G. H. 1774. O r.m Bot. reg. 235

8772 ciliatum Thun. ciliated tf IAlor △my.jn W G C. G. H. 1819. O r.m

6773 crenulkatum Thun. crenulated tf ^Jor } ap.my W C. G. H. 1816. O r.m

8774 maculkatum Thun. spotted tf A) or A my.jn W C. G. H. 1813. O rm

8775 tenillum Jac. delicate tf LAJor *jn W C. G. H. J81H. O r.m Jac. ic. 2.427

8776 scdndum Jac. side-flowering tf LAJor 4 nu W C. G. H. 1826. O r.m Jac. ic. 2.433

8777 pilosquin Thun. pilose tf IAlor 1 ap.my W C. G. H. 1826. O r.m

8778 revolt.turn Jac. revolute-JfriMf tf IAlor 1 mr.jn W C. G. H. 1795. O r.m Bot. mag. 653

8779 dātum, Andr. tall tf AJor 3 mr W Egypt 1804. O r.m Bot. rep. 523

8780 coryinbōsum Fl. per. corymbose tf T 14 my W Chile 1823. O r.m Bnt. reg. 906

8781 tcuifolium Red. fine-leaved tf IAI or 1 ap.my W G. G. H. 1819. O r.m Red. lll. 312

8782 pfasinum Ker green+-ouvrfrf tf IAlor i m W C. G. H. 1816. O r.m Hot. reg. 158

8783 sauvcolens Jac. sweet-scented tf IAI & 1 my.jl W C. G. H. 1826. O r.m Jac. ic. 2.431

8784 barbtoum Jac. bearded tf IAI or 1 my.jl W C. G. H. 1795. O r.m Jac. sc. 1.91

8785 polyph^llum Jac. many-leaved tf IAJ ec A J W C. G. H. 1824. O r.m Jac. ic. 2.430

8786 ovatum Thun. ovate tf IAJ or 1 my.jn W C. G. H. 1824. O r.m

8787 rupestre L. rock tf IAJ or 4 my.au W C. G. H. 1795. O r.m

8788,unciflium Jac. Rush-leaved tf AJ or f jl.au W C. G. H. 1794. O r.m Bot.mag. 97⁷

8789,fuscatum Jac. dull tf IAlor 4 Jnjl L.Y C. G. H. 1820. O r.m Jac. ic. 2.429

8790 odoratum Jac. sweet-scepted tf IAlor U my.jn F C. G. H. 1795. O r.m Bot. rep. 260

8791 nyramdale L. pyramidal tf A or 2 jn.jl W Spain 1752. O r.m Jac. ic. 2.425

8792 longibracteatum Jac. long-bracted tf IAlor 4 my W C. G. H. 1817. O r.m Jac. sc. 1.29

8793 seilloides Jac. Squill-like tf IAlor 4 jnjl W C. G. H. 1795. O r.m Jac. sc. 1.888

8794 latifohum L. broad-leaved tf Alor 4 jnjl W Egypt 1629. O r.m Bot. mag. 87⁷

8795 narbon^usc L. Narbonne tf A or 2 jn.jl W S. Europe 1810. O co Bot. mag. <5W

8796 pyrenäicum I*. Pyrcnean tf A or 24.ap.il G T C. G. H. 1771. O co Ren. spec. fl.

K7/ «tach»oides H. K. Stachys-like tf A or 1 jn.jl W C. G. H. 1796. O r.m Bot. mag. Iij⁷

H798 lacteum Jac. milk-white tf IAI or 1 jn.jl W C. G. H. 1774. O r.m Bot. mag. 8W

8799 caudatum II. K. caudate tf IAJ or 3 f.au W o C. G. H. 1774. O r.m Bot. mag. 8W

Jf/0 nutans L. nodding ■ A or 4 jn.jl W Britain me.pa O ro King.bot POT

8801 conrctatum Jac. compressed-/fufv if LAI or 14 Injl W o C. <. H 1804. O r.m Jar. ic. 43d

8802 atireum <i>Cur.</i>	golden	tf IA.Jor	f jn.jl	Y	C. G. H. 1790.	O	r.m	Bot. mag. 190
8803 flavidum <i>Jac.</i>	yellowest	tf iAJ or	1 jnj	Y	C. G. H. 1804.	O	r.m	Jac. ic. 436
8801 thyrsifolius <i>Jac.</i>	thyrsifolius	tf iAlor	i jnj	W	C. G. H. 1757.	O	r.m	Rot. mag. 1164
2 flavescens	flavescens	tf iAlor	i jnj	W	C. G. H. 1800.	O	r.m	Bot. reg. 305
8807 arabicum <i>L.</i>	Arabian	tf iAlor	i mr.ap	W	Egypt 1529,	O	r.m	Bot. mag. 7&8
8806 ixioidea // K.	Ixia-like	tf iAlor	i my.jn	W	California 1795.	O	s.p	
8807 hispidum <i>Horn.</i>	hispid	tf iAlor	i	W	C. G. H. 1824.	O	r.m	
8808 vires <i>Lamil.</i>	greenish	tf iAlor	i jnj	W	Del. Bay 1823.	O	co	Bot. reg. 814
8809 umbratum <i>B. M.</i>	one-leaved	tf iAlor	i jnj	G	Gibraltar 1805.	O	r.m	HoLmag. 915 953
8810 Squill <i>B. M.</i>	afflinal Squill	tf iAlor	i ap.my	W	S. Europe 1829.	O	r.m	Bot. mag. 918
Scilla maritima <i>L.</i>								
•1054. SCILLA L.	SQUILL.	(<i>Skyllo</i> , to injure; roots poisonous.)			Asphodillae.	22.—32.		
8811 italicica <i>L.</i>	Italian	tf A or	i ap.jl	B	Switzerl.	1H05.	O pi	Bot. mag. 663
8812 peruviana <i>L..</i>	Peruvian	tf A or	i my	D.B	Spain	1607.	O r.m	Bot. mag. 749
2 alba	white	tf A or	i my	W	S. Europe	1607.	O co	
8813 lusitanica <i>L.</i>	Lusitanian	tf	i	L.B	Portugal	1777.	O Pl	Bot. mag. 1999
8814 Lilio-J/yacinth* <i>L.</i>	Lily-Hyacinth	tf A or	i my.jl	B	S. Europe	1597.	O co	Red. HI. 205
8815 amce. na <i>L.</i>	pleasing blue	tf A of	i mr.ap	I	Ixiant	1596.	O pl	Bot. inag. 341
8816 sibirica <i>H. K.</i>	Siberian	o A or	J fmr	B	Siberia	1796.	Oji.1	Bot. mag. 1025
8817 prasW <i>W.</i>	c>\-flowering	tf A or	i mr.ap	D.B		1790.	I	
8818 verna <i>L.</i>	vernal	tf A or	J apmy	B	Britain	rocks.	O si	Eng. bot 23
8819 unifolia <i>L.</i>	one-leaved	tf A or	i my.Ju	W	Portugal	-	O 81	
8820 ryacinthoides <i>L.</i>	Hyacinth-like	tf iAlor	4 au	B	Madeira	1585.	O r.m	Bot. mag. 1140
8821 autumnalis <i>L.</i>	autumnal	tf A or	4au.s	Pk	England dr. pa.	O pl	Eng. bot 78	
8822 bifoliaL.	two-leaved	tf A or	* cap	B	England woods.	O pl	Eng. bot. 24	
2 alba	white	tf A Pr	i Jap	W	S. Europe	... O co		
3 rrhra	red	tf A pr	i	It	S. Europe	... O co		
8823 ptimila <i>BroL</i>	dwarf	tf A >r	1 ap.my	B	Spain	1821.	O r.m	
8824 umbellata <i>Ram.</i>	uinbellcd	tf A or	\ ap	B	Pyrenees	1822.	O pl	Brot ph. 41.8.6
8825 cfrnua <i>Red.</i>	drooping	■ ■	\ ap.my	Pk	Spain	1815.	O pl	Bot. mag. 1461
8826 Indica <i>Rox.</i>	Indian	tf ES3 or	i	K Indies	1816.	O pl		
8827 campanula <i>H. K.</i>	campanulate	tf A or	1 my.jn	D.P	Spain 81683 p.	O	Pl Bot. mag. 127.1102	
8828 non scripta <i>L.</i>	undescribed	tf A or	f mr.jn	B	Britain woods.	O co	Eng. bol 377	
n titans Sm.								
2 caruca	flesh-catered	tf A or	% mr.jn	Pk	Britain woods.	O 81	Bot. mag. 1641	
3 alba	white	tf A or	\$ mr.jn	W	Britain woods.	O s.l		
8829 brevifolia tor	short-leaved	tf iAl or	\ ja	Pk	C. G. H. 1811.	O s.l	Bot. mag. 1648	
8830 maurita'nica Schous.	Mauntanian	tf A or	\ ap.my	B	Mauritan. 1819.	O r.m		
8831 wojymbia <i>B. M.</i>	corymbose	tf iAl or	\ aud.	Pk	C (5 H 1793.	O s.l	Botmag. 1748.IL	
8832 esculenta <i>B. M.</i>	esculent Quamash	tf A cc	1 my.jl	W	N. Amer. 1811.	O 81	Bot. mag. 1574	
1055. BELLEVIA ChIA Lap. BELLEVALIA. (P. R. Bellevall, a French botanist.) Asyfiodi/ear. 1.								
8833 operculata <i>Lap.</i>	lid-covered	tf A or	5 my	W	Italy 1596.	O co	Bot. mag. 936	
ficilla romana <i>L.</i> //yacinthu« romanus <i>L.</i>								
1056. BARNA'R'DJil Lindl. BARNABDIA. (E. Barnard, F.L.S., Vice-Sec. Hort. Soc. London.) Asphodile*.	1.							
834<cioldcesAiS.	Scilla-hke	tf .Al or	i jl.au	P	China 1826.	O pl	Bot. reg. 1029	
1057. PUSCHKI'N// Adams PUSCUKINIA. (Count M. Pouschkin, a Russian bot.) Asyphodilea: 1.								
88& scilloides Adams	Scilla-like	tf jf or	± my.jn	P.B	Siberia 1819.	O 81	Lind. col. 24	
Adamsia willoldes W.								
1058. MASSON/N/il L.	MASSONIA.	(Mr. F. Masson, author of Stnpclim Nova;)			Asphodilea.	13.		
28,(*) laffia-lia <i>L.</i>	broad-leaved	tf LAJ cu	i mr.ap	W	C. G. H. 1775.	O 81	Bot. mag. 848	
28,7 cordita <i>Jac.</i>	heart-leaved	tf iAl or	A my.jn	W	C. G. H. 1826.	O r.m		
wj A ionRiffha <i>Jac.</i>	long leaved	tf LAJ cu	i mr.ap	W	C. G. H. ...	O B1	Jac. sc. 4.457	
22 » Candida Burc.	white	tf iAl cu	i mr.ap	W	C. G. H. ...	O s.l	Bot reg. 694,	
840 grandiflora <i>Lindl.</i>	large-flowered	tf iAl cu	i o	W.G	C. G. H. 1825.	S s.l	Bot reg. 958 ,	
88* muricata <i>Ker</i>	Yrckly-leaved	tf iAl cu	L ap.my	W	C. G. H. 1790.	O 81	Bot. mag. 559	
**& scabra // K.	rough	tf iAlcu	i ja.ap	W	C. G. H. 1790.	O a 1	Bot rep. 220	
particular								
cvnikiita <i>L.</i>	prckly-leaved	tf IAICU	\ y	W	C. G. H. 1790.	O 81		
I i i uuciflora // K.	few-flowered	tf IAICU	\ my	W	C. G. H. 1790.	O 81		
I- iungustiflbia <i>L.</i>	narrow-leaved	tf IAICU	\ mr.ap	W	C. G. H. 1775.	O 81	Bot mag. 736	
Ui uiuulata Thun.	waved-leaved	tf IAICU	\	W	C. G. H. 1791.	O 81		
8847 easifMia Ker	sword-leaved	tf IAICU	*	Li	C. G. H. 1790.	O Si	Bot mag. 554	
8818 violacea Amir.	yuAut-colored	tf iAlor	i myjn	-	C. G. H. 1800.	O r.m	And. rep. 46	
1059. EREMITRUS Bieb.	EHIEMURUS.	(Eretros, solitary, oura, a tail; spikes.)			Asphodileaf.	1.		
»4U spectabilis Bieb.	beautiful	A or	2 my.jn	Y	Siberia 1800.	O 81	Sw. fl. gar. 188	
ilspodelus alfaicus Pall.								
1060. ZJULBrNE IV. en.	BULBISE.	{Bolbos, a bulb.)			Asphodileee.	19.—24.		
<w0 hispida Sut.	hispida	■ ■	1 my.jn	G.w	C.G. H. 1774.	Sk s.l	Jac. ic. 2.409	
Anthericum hispidum <i>L.</i>		£ iAlor	1 jl	Y	N. Holl. 182a	Sks.l	Bot. cab. 320	
»1 semiarbhta Spr.	semi-bearded	tf LAJ or	1 jnj	Y	N. Holl.	1890.	O s.p	
SBro Anthericum semibarbatum It. Br.		tf A or	In	Y	N. Holl.	1823.	O s.l	
»* « austriklis Spr.	southern	tf O or	my.jn	Y	C. G. H. 1731.	S s.p	Bot. mag. 1451	
Anthericum bulbiflum R. Br.	bisulcate	tf iAlor	1 ap.au	Y	C. G. H. 1759.	Sk r.m	Bot. mag. 1339	
annua W. en.	annual	tf 2 jn.au	W	C. G. H. 1759.	O r.m	Jac. vin. 181		
Anthericum unnuum L.		tt i or	S mr.au	Y	C. G. H. 1812.	C s.l	Jnc. ic. 2.403	
8g longscapa IV. en.	loig-scaped	«L» ior	2 mr.au	Y	CO H 1702.	C s.l	Bot. mag. 816	
0856 aaphodeloldes Spr.	Asphodelus-lk	ip iAlor	1 ap.jn	Y	C. G. H. 1793.	Sks.l	Bot. mag. 1454	
DO.. Anthericum aaphodeloldes L.	beaked	ti iAlor	1 my.au	Y	C. G. H. 182a	O r.m		
£.7 rostrata W. en.		tf 1Alor	1 jnj	Y	C. G. H. 1825.	Sk.s.p		
JM frutescens W. en.	frutescent				C.G. H. 1823.	S 8.p		
2 ^ pugioniformis LAr.	dagger-formed				C.G.H. 1818.	O r.m	Jac. ic. 2. 406	
ori triquetra Spr.	three-sided							
uqz Anthericum triquetrum Thun.								
scabra Spr.	rough							
Anthericum sebrura Thun.	ciliated							
ciliat. J.								
praBmfsra Spr.	praemorse							
Anthericum prsm6rsu Jac.								

1071. DIANJS'LLJ* Ker	DIANELLA. (<i>Diana</i> , sylvan goddess j <i>habits woods.</i>)	<i>Asphodellee. 9.—11.</i>
8926 laiVis R. Br.	smooth AiAJor 12 au B N. Holl. 1822.	Sk s.p
8926 longifolia R. Br.	long-leaved A 1AI or 2 au B N. Holl. 1822.	Sk s.p Bot. reg. 734
8327 strumusd Ker	strumous AiAJor Ifmr B N. Holl. 1822.	Sks.p Bot reg. 751
888 revoluta R. Br.	revolute A 1AJor 2 my.au B N. Holl. 1821.	Sks.p
8929 congehta A. Br.	crowded AiAJor 2 my.au B N. Holl. 1820.	Sks.p
8930 incinorbsa f<>.	grove AiAlor 2 au B t. Indies 1731.	Sks.p Ru. am. 5.73
8931 ensifolia B. Al.	sword-leaved A 1SI or Ijau W E. Indies 1731.	Sk p.l Bot. mag. 1404
Dracaena ensi folia II. K.		
8932 racifera R. Br.	blue AiAJor 2 my.au B N. Holl. 1705.	Sks.p Bot.inag. 505
8933 divaricata R. Br.	divaricated AiAlor 3 jl.au B N. S.W. 1805.	Sks.p

1072. CORDYLPNE R. Br.	CORDYLINE. (<i>Kordyle</i> , a club.)	<i>AspiwdZlece. 2.</i>
8934 hemichryta Com.	half-golden m or 2 . . . Bourbon	1823. Sk veg
8935 cannefilia R. Br.	Canna-leaved * LJ or 4 . . . N. Holl. 1820.	Sk p.l

• 1073. EITSTREPHUS R. Br.	EIJSTREPHIUS. (<i>EU</i> , well, <i>strrpho</i> , to twine; habit)	<i>Asphodilete. 2.</i>
8936 latifolius R. Br.	broad-leaved % _J or 3 jn.jl Pa.P N. S. W. 1800.	C sp Bot.mag. 1245
8937 angustifolius R. Br.	narrow-leaved fc. i_ or 3 jl Pa.P N. S. W. 1820.	*C s.p

■ 1074. ^SPA'RAGUS L.	ASPARAGUS. (A. intens. <i>sparasso</i> , to tear: <i>pickles.</i>)	
8938 officinalis L.	officinal	England seaco. S r.m Eng. bot. 339
8939 longifolius Fis.	lggg-leaved	Siberia 1827 S co
8940 amirius Dec.	bitter	France 1824. S r.m
8941 sylvestris W. #A.	wood	Hungary 1819. S r.m W. & K. 3.201
8942 verticillaris L.	whorUcaved	Caucasus 1752. S r.m Bux. cen. 5.37
8943 declinatus L.	declined	C. G. H. 1759. R s.p
8944 maritimus Pall.	maritime	Caspian 1823. R s.p
8945 decumbens Jac.	dcumul)cnt	C. G. H. 1792. R s.p Jac. sc. 1.97
8946 scandens Thun.	climbing	C. G. H. 1755. R Mp
8947 dahuricus Fts.	p.iuhan	Dauria 1823. R s.p
8948 falcifolius L.	&ck>lcaved	E. Indies 1792. R s.p Bur. zejr. 13. a
8949 racemosus JV.	racemose	E. Indies IN08. R s.p
8950 Broussonet's Jac.	Broussonet's	Canaries 18 22. R s.p
H9il retrofract IK	retrofracted	Africa 1759. R s.p Pluk. al. 375. J
8952 asiaticus L.	Asiatic	Asia 1759. R s.p Pluk. al. 15.4
8953 sethiopicui L. •	Ethiopian	C. G. H. 1816. *K s.p
8954 Albus L.	white	Spain 1540. R ap M. h. 1. L 3
8955 stipulaceus Lam.	stipulaceous	C. G. H. 1821. R s.p
*8956 acutifolius L.	acute-leaved	Spain 1640. R s.p Par.th.3
8957 flexuosus Thun.	flexuous	C. G. H. 1800. R s.p
8958 aphyllus L.	leafless	S. Europe 1040. R s.p M.h.1.1.2
8959 dropendens Thun.	drooping	C. G. H. 1819. R s.p
8960 horridus L."	horrid	S. Europe 1800. R s.p Cav ic. 2.136
8961 subulatus thun.	awl-leaved	C. G. H. 1811. R s.p
8962 caudatus L.		C. G. H. 1621. R s.p Jac. sc. 3.266
8963 aarmamentus L.	twiggy	W. o Ceylon 1710. R r.n R. nul. 10.10

1075. HERRETRI A R. & P.	HERRERIA. (C. A. de Herrera, a Spanish agriculturist)	<i>Asphodilete. 2.</i>
8964 parviflora B. R.	small-flowered % C3 or 8 jn.jl G.Y Brazil 1824.	C r.m Bot. reg. 1042
8965 stellata/ Sf P.	starred t_E3 cul 8 jn.jl G.v Chile 1825.	C r.m Fl. per. 3. 303. a

* 1076. LUZURIA>i<Fl. per.	LUZURIAGA. (D. Jgn. M. R de Luxuriago, M.D., Span.bot)	<i>AsphotdetE. 2.-3.</i>
8966 cymosa R. Br.	cymose % ILJ pr 3 my.jn G N. Holl. 1825.	C p.l
8967 montana R. Br.	mountain EAIAto . . r G G NN.Holl. 1820.	C s.p

* 1077. DRI'MIA Jac.	DRIMIA.	<i>Asphod>hac. 10.</i>
8968 altissima Ker	tallest tf iAn	If au.s VV.G C. G. H. 1791. G s.p Bot. mag. 1074
8969 elata Jac.	tall tf LAJ or 2 o.n R.u C. G. H. 1799. O s.p Bot. mag. 822	
8970 cili-Xris Jac.	ciliated tf IAJ or Ifs P.w C. G. H. 1800. O s.p Bot. mag. 1444	
8971 pusilla Jac.	little tf IAJ or * my.jn G C. G. H. 1793. O s.p Jac. ic. 2. 374	
8972 lanceolifolia Ker	tpcar-leaved tf IAJ or \ so P C. G. H. 180a O s.p Bot. mag. 643	
8973 undulata Jac.	undulated tf IAJ or \ aus G C. G. H. 1819. O s.p Jac.'ie. 2.376	
8974 purpurascens Jac.	purpurascens tf LAJ or / aus P C. G. H. 1818. O s.p	
8975 revoluta B. M.	revoluta tf LAJ or / au G C. G. H. 1774. O s.p Bot. mag. 1380	
lancculifolia (B. M. //yacintu*) revolutus // K.	Lachnantha reflexa Andr.	
8976 mtdia Jac.	intermediate tf LAJ or / au W C. G. H. 1820. O s.p	
8977 acuminata Lod.	pointed tf IAJ or au Br C. G. H. 1829. O p.l Bot. cab. 1041	

9.J? - MI'LILJil Cav.	MILLEA. (<i>JulincaMula</i> , chief gardener, royal garden, Madrid.)	<i>Asphodelets. 1.</i>
'978 biflbra Cat.	two-flowered tf [SI or 1 my.jn W Mexico 1826.	O s.l Cav. ic. 2.196

1079. UROPETALON Ker	UROPETALOY.	<i>(Ouria, a toil, petaton, a petal.) AspAodtoef. 6.</i>
8180 crispum Burc.		tf 1 jl.au G C. G. H. 1816. O l.p Bot. reg. 156
8970 "or"tinu" Ker	late-flowering tf J'v or ijn'iù G.R Spain 1625.	O l.p Bot. mag. 859
QJB^ fulvum Swt.	tawny tf Alor 1 jn.au G.R Mogadore 1808.	O l.p Bot. mag. 1185
Sc\yacintu(3B.M.		
8983 longifolium Lindl.	long-leaved tf 2 au P.B Mozambi. 1825.	O p.l Bot. reg. 974

■ 1084 amethystinus	hyacintus	<i>(A boy killed by Zephyrus) Asphodite*. 2—8.</i>
8985 orientalis L.	oriental tf A or i ap.ay B S. Europe 1759. O l.p Sw. (l. gar. 135	
2 rdber	tf A or f inr.ap B Levant ISM. O r.m Bot reg. 995	
3 flvus	tf A or f mr.ap R 1596. O r.m	
4 albus	tf A or \$ mr.ap Y 1596. O r.m	
H semiplnuB	white tf A or f mr.ap Va 1596. O r.m	
6 multiplex	semi-double tf A or % mr.ap Va 1596. O r.m	
	manifold.	

ZTJCCA'GNII Thun. ZUCCAGMA. (A. Zuccagni, gard. raperin, at Florence.)

vjridisTctin. green tf IAJor f au G ^G.H. 1774. O 1 — <

"yacintu vridis L. LachenMw viridis fV.

* 1083. MUSC AMU Detf. GRAPE HYACINTH.		(<i>Moschos</i> , musk; smell.)	<i>Atphodilece.</i> 7. — 10.
8987 glaticum B. 21.	' glaucous	■ A or % my P.o Persia	1825. O s.l Bot. recg. 1085
8988 moschatum B. M.	musk	■ A or fap my B.Y Levant	1596. O 61 Bot.mag.7J4
//yacinthuj Muscari L.			
8989 macrincarpum Swt.	long-fruited	tf A or i ap.my G.Y Levant	1596. O s.l Boi. mag. 1565
moschatum /3 fiavum B. M.			
8990 nfttum Ker	ciliated	tf A or 1 my Br.p Crimea	1822. O s.l Bot. reg. 394
8991 cuinr.bium W.	tufted	tf A or 4 np.my B S. Europe	1822. O 81 Bot. mag. 133
2 mon-trssum	monstrous	tf A or \ ap.my B S Europe	1593. O s.l M. h. 4.11. S
8992 p. illens Fis.	Sillid	tf A or 4 my Pa.B Crimea	— O 81
8993 fotrydickt IV.	Sotrys-like	tf A or ap.my B Italy	1596. O 81 Bot. mag. 157
2 pallium	pale	tf A or ap.my Pa.B Italy	1596. O co Sw. fl. gar. 15. e
3 album	white	tf A or i ap.my W Italy	1596. O co Sw. fl. gar. 15. b
8994 racemusum W.	racemose	tf A or I ap B Europe	1780. O 81 But. mag. 122
2 minus	smaller	tf A or i ap B Europe	1780. O 81
• 1083. LACHENA ^{SLM} Jac.	LACHEVALIA.	(<i>W. de la Chenal</i> , bot auth.)	<i>Asphodilets.</i> 35. —
89'tf pendulu H. K.	i^>^>^>^>^> \$*AIP'	f mr.niy R.Y C. G. H. 1789. O 81 Bot. mag. 590	
2 maculata	sptt^>^>^>^>^> AJ pi	^ mr.my ILv C. G. H. 1789. O 81	
8906 punctata Jac.	dotted	tf Al or my.jn P C. G. H. 1824. O 81 Jac. ic.2. 597	
8997 ruin da Jac.	TtiA-dotted Jtwd	tf Al pr f.s.o R C. G. H. 18a). O 81 liot. mag. 993	
8998 quadricor Jacob.	four-colored	tf fffAAlpr mr.ap S.Y C. G. H. 1774. O S1 Bot. mag. 588	
2 colorata	colored-leaved	tf LAlpr 4 mr.my S.Y C. G. H. 1774. O 81 Bot. mag. 1097	
8919 scriptina Jac.	late	tf -Alpr au Pk Spain 1820. O 81	
9000 tricolor Jac.	thrc-e-roled	tf LAlpr 1 ap.my R.Y C. G. H. 1774. O 81 Jac. ic. 1.61	
9001 lut&ola Jac.	yellowish	tf LAlpr 1 ap.my Y.R C. G. H. 1774. O S1 Bot. mag. 82	
2 maculata	■■■■■fc ■■■■■	tf LAlpr 1 ap.my Y.R C. G. H. 1774. O S1 Bot. mag. 1704	
9002 flava Antr.	yellow	tf IAJr J my.jn Y C. G. H. 1790. O sl Bot. rep. 77	
9003 Hcsitlura Andr.	sestislt^nWered	tf iAl pr I my.jn R C. G. H. 1804. O 81 Hot. rep. 460	
9004 isopetala Jac.	equal-pcf filed	tf ■■■■■ f my.jn W.p C. G. H. 1804. O 81 Jac. ic. 2.401	
9005 umftlia Jac.	one-icaw "I	tf LAlpr 4 mr.ap W.B C. G. H. 1795. O sl Bot. mag. 766	
9006 pusilla Jac.	small	tf IAJor ■■■■■ W C. G. H. 18'5. O 81 Jac. ic. f 385	
9007 bifltha Ker	two-leaved	tf LAlpr mr.ap Pk C. G. H. 1813. O 81 Bot. mag. 1611	
9008 flsea Andr.	roseate	tf ufelpr 1 ap my Pk C. G. H. 1800. O 81 Bot. rep. 296	
9009 containiata H. K.	contaminated	tf IAJpr * f..r Pk C. G. H. 1774. O S1 Bot. mag. 1401	
9010 angufnca Swt.	serpent	tf LAlpr ap.jl W C. G. H. 1825. O 81 Sw. fl. gar. 179	
9011 /Inflbra Jcc	Lily-flowered	tf LAlpr ap.jl W C. G. H. 1825. O 81 Jac. c. 2.287	
9012 (Jrtw/ma Jac.	glaucous	tf LAlpr mr.ap G.w C. G. H. 1795. O 81 Jac. ic. 2.391	
9013 mutabilis Lad.	■■■■■	tf LAJel ■■■■■ B C. O. H. 1825. O S1 Bot. cab. 1076	
9014 orchidifl. // K.	Orchhuulike	tf iAJ pr f.ap G.Y C. G. H. 1752. O 81 Bot. mag. 1269	
9015 paliwla // A.	LAI pr	tf mr.ap PA.B C. G. H. 1782. O 81 Bot. recg. 287	
9016 hy-dcinthoides Jac	Hyacutji-like	tf iAl pr ** mr.ap W.R C. G. H. 1812. O S1 Jac. ic. 2.382	
9017 angustifilia Jac.	narrow-leave-l	tf iAl pr 1 ap.my W C. G. H. 1793. O si Bot. mag. 735	
9018 patula Jac.	spreading-yiwi	tf LAlpr J ap.my W.pk C. G. H. 1795. O 81 Jac. ic. y. 384	
9019 purpurea Jac.	purple	tf iAJ or mr.my P C. G. H. 1826. O 81 Jac. ic.2.393	
9(20) lancieflilia Jae.	lance-leaved	tf LAlpr mr.my W.o C. G. H. 1818. O si Jac. ic. 2.402	
9J21 glnucum B. R.	glauconous	tf A or f my P.o Persia 1825. O si Bot. reg. 1085	
9022 frv-irans Jac.	sweet-scented	tf LAlpr mr.my W.R C. G. H. 1806. O si Bot. mag. 1-73	
9023 unicolor B. M.	self-colored	tf LAlpr my.jn Pk C. G. H. 1798. O si Bot. mag. 1372	
9014 lucirid Ker	g/o&sy-tcaved	tf LAJpr mr.my Pk C. G. H. 1811. a.s1 B xt. mag. 1517	
9025 raccinosa A. M.	race-mose	tf iAJ pr my W.o C. G. H. 1790. () 81 Bot. mag. 817	
9026 piiKtulXta Jnc.	bliktervd	tf LAlpr 1 ja.ap W.o C. G. H. 1798. O 81 Bot. mag. 745	
9027 purpuro-caerulea Jac.	purple-blue	tf LAlpr 1 up.my B.P C. G. H. 1810. O si Bot. mag. 1497	
9028 nrcyflui Ker	nervetUcavcd	tf ■■■■■ \$jn Pk C. G. H. 1795. O 81 Jac. ic. 2.394	
9029 violacea Jac.	vio/ct-colored	tf ■■■■■ 1 mr.ap L.B C. G. H. 1795. O 81	
1084. PHO'RMIUM L.	FLAX LILY.	(<i>Phormos</i> , a basket; U.S.C in New Holland.)	<i>Aspliodelea?</i> . 1.
9030 tenax Forst.	tough	y■ A ec 6 au G.w N. Zeal. 1798. H Up Cook 2.96	
•1085. CYANE'LLAZ.	CYANELLA.	(Dimin. odkij&wpWdjudlfkulfw flowers.)	<i>Asphodlra?</i> . 5.
90-31 capensis L.	Cape	tf LAI br 1 jl.au B C. G. H. 1768. O s.p Bot. mag. 568	
9032 K.tea T/ian.	yellow	tf LAJ pr 1 jl au Y C. G. H. 1788. O s.p Bot mag. 1252	
9033 orchidifòrmis Jac.	Orchis-formed	tf LAI or 1 au B C. G. H. 1826. O s.p Jac. ic. 2.447	
9034 filba Thun.	white	tf AJ or 1 jl.au W C. G. H. 1819. O s.p Th.h.7.2	
9035 lineata Jitore.	lined	tf LAI pr 1 jl.au St C. G. H. 1816. O s.p	
1086. LEO'NTICE L.	LEONTICE.	(The Greek name for wild chervil.)	<i>Berberideee.</i> 4.. 5.
9036 ChrysKonum L.	A Alcu	1 mr.jn Y Levant 1740. R s.l.p M. h. 3.15.7	
9037 al-/Nci Pall	A. AJ or	i ap.my Y Siberia 1818. R s.l.p Lam.il. 254.2	
9K3 Leontoptitalon L.	Altaijan	1 ap.my Y Levant 1597. R n.l.pM. h.3.15.3	
90J9 veaicuna Pall.	Lion'-8-loaf	A AJ cu 1 ap.my Y Siberia 1821. R s.l.pPal. it.3.5.2	
1087. CAULOPHY'LLOM A*/. CAULOPHYLLUM. (<i>Kaulos</i> , a stem, <i>phyllon</i> , a leaf.)	Berberideat.	1.— 8.	
9040 (halirtridoides Mx.	Thahctrum-Ik	A A cu 5 my Y.o N. Amer. 1755. R up Mic. am. 1.21	
Z.cöntice /halictroldes L.			
1088. DIPHYLLE'IA Mx.	DIPHYLLEIA.	(Dis. twice, <i>phyllon</i> , leaf; always two.)	<i>Bcrher'dece.</i> 1.
9041 cymosa Mx.	cymose	^ Wx \$ my.jn W N. Amer. 1812. D ip Bot mag. 1668	
1089. 7>RrNOS L.	WINTER BERRY.	(<i>Prion</i> , a saw j serrated leaves.)	<i>Ixdmnce.</i> 11.— 13.
9042 deciduous Dec.	deciduous	11 or 4 jnjjl W Virginia — L Ls	
/Nex primoides H.K.			
9043 verticillatus L.	verticillated	■ sa or 6 jl.au W N. Amer. 1736. L s.p Den. br. 30	
9044 amblguus Mx.	ambiguous	A or ■ — W Carolina 1812. L Its Den. br. 29	
9045 lsvigut Ph.	smooth	tt or 4 jl.au W N. Amer. ... L Its Den. br. 28	
9046 lanceolktus Ph.	lanceolate	St or 4 jnjjl W Carolina 1811. L Its	
9047 glaber L.	smooth	■ or 1} jl.au W Canada 1759. L Its	
9048 atumkrus Nut.	atomed	or, 2 jl.au W N. Amer. 1822. L Its	
9049 jtrunifolius Desf.	Plum-leaved	■ or — W 1810. L Its	
»»50 corikceus Ph.	leather-leaved	■ or 3 jnjjl W N. Amer. 18'0. L Its	
9051 lucidug // K.	shining	m or 2 jnjjl W N. Amer. 1778. L Its	
..... ^ xi « can'Mc'iss R.	mountain	il ■ or — W W. Indies 1820. C Its	
9052 mont inua Swx.			
* 1090. BE'RBERIS L.	BERBERRY.	(<i>Berberys</i> , its Arabic name.)	<i>Berberideat.</i> 15.— 8.
9053 vulgaris /«	common	" Si fr 8 ap.my Y England bu. pi. L co Eng. bot. 49	
2 violacea	violet-/ui7crJ	* fr 8 ap.my Y L co	
3 alba	white fruited Sk	fr ti ap.my Y L co	

4 Ihtea Dec.	yettovr-fruited	■	fr	6	ap.my	Y	Europe	... L co
5 purp&rea Dec	purple-fruited	■	fr	6	ap.my	Y	Europe	... L co
6 rtgra Dec.	black-fruited	■	fr	7	ap.my	Y	Europe	... L co
7 asperma Dee.	seedless	■	fr	6	ap.my	Y	Europe	— L co
9054 daGrica Lo. C.	Daunan	■	or	8	apjn	Y	Dauna	1818. L co
VO55 provincialis Lo. C	Provence	■	or	8	ap.my	Y	France	1821. L CO
9056 ibérica Fts.	Iberian	■	or	5	ap.my	Y	Iberia	1818. L r.m'
vulgtois fi ibérica Stev.								
9057 cunad&niss Mil.	Canadian	■	fr	8	ap.my	Y	Canada	1759. L co
9058 cmargmata W. en.	emarinate	■	or	3	ap.my	Y	Siberia	1790. G co
9059 sincnsis Detf.	Chinese	fik	or	4	ap.my	Y	China	1815. L co Den. br.86
9010 anstkt Dec.	awncd	■	or	6	ap.my	Y	Nepal	1820. Hook.ex.fl. 98
rhitria R li.								
9061 erotica L.	Cretan	■	or	6	ap.my	Y	Candia	1759. L co
9062 sibrica Pall.	Siberian	Sk	or	2	jnjl	Y	Siberia	1790. L co Bot. reg. 487
9053 ilicifolm Furst.	Holly-leaved		or	4	jl.au	Y	T.delFue.	1791. L r.m
9064 asiatica Itox.	Asiatic	■	or	4	...	Y	Nepal	1823. L r.m
9065 WallichV/m Dec.	Wallich's	jor	4	apjn	Y	Nelial	1820. L r.m DelBS. 2.1	
9066 rusculia Lam.	Ituscu-leaved	■	or	5	my.jn	Y	B. Ayres	1821. L r.m
9067 heterophyllt J.	various-leaved	■	or	4	ap.iny	Y	Magellan	1805. L r.m Lam. il. 253
* 1091. MAHOWA Nut	MAHONIA.	[Bernard WMahon, of N. Amer., lover of bot.]	co	Berberidcte. 14.				
9068 nervosa Nut.	nervose							
Jt'iluris nervosa Ph.		fit	-J or	10	...	Y	N. Amer.	1826. G r.m l.h. am. 1. 5
9069 s/guifolium Nut.	Ho/ly-leaved	■	or	J	...	Y	N. Amer.	182k. G r.m Ph. am. 1.4
Berberis i4/guifolium Ph.	bundled	il	or	10	ap.my	Y	California	1819. L co Bot mag. 2396
9070 lusricul.Hriu Dec.								
Bitberi* pinnkta Lag.								
1092. NANDPNA Thun.	NANDIMA.	(Nandin, name in Japan.)		Berberidcte. 1.				
9071 domfotica Thun.	domestic	■	i_J or	6	jnjl	GMr China	1804. C pl Bot mag. 1109	
* 1093. COSSI'GN/iJ.	COSSIONIA.	(M. de Cossigny, a French naturalist)		Saptndlcete. 1.				
9072 borbonica Dec.	Bourbon	■	or	JO	...	Mauritius	1824. C pl	
pinnkta Lam.								
1094. HI'LL/if L.	HILLIA.	(Sir John Hill a voluminous botanical author.)		Rubiaceaj. 2.				
9073 longiflora Swx.	lon^flowered	■	n Q or	1	ljfmr	W W. Indies	17H9. C Bp Bot mag. 721	
9074 tetrandra Swx.	tetrandrous	■	<-CZ or	1	jnjl	W Jamaica	1793. C s.p	
1095. RICHARDSO^N/^ Kth.	RICUARDSOMA.	(Richard Richardson, an Eng. bot.)		Rubiillcett. 8.				
9075 scabra Kth.	rough	a. • W	Sa	W VeraCruz	... C l.p	Sw. fl. gar. 91		
Uichárdim scabra L.								
9076 pilcesa Kth.	pilose	- * O w	i jl.au	W Brazil	1814. S co			
109R SCHRA'DEIM Vdh	SCBRADBRA.	(Henr. A. D. Schroder, a German botanist)		Bubiaceta. 1—2.				
9077 cephalotes W.	round-headed	* • or	4 jl.au	K Jamaica	1820. C Lp			
Fuchsia involucrata Swx.								
1097. ISE'RT/^ Schreb.	IBERTIA	(P. E. Isert, a Tanish surg. on coast of Guinea.)		Rubiace. 1.—2.				
9078 roccinea Vahl	scarlet	«O spl	10 jl.au	S Guiana	1820. C 81 Aub. gui. 1.123			
Guccarda coccina AUBL.								
1098. CANARPNA L.	CANARINA.	(A native of the Canaries.)		Cavipanulhceee. 2.				
W79Campinula L.	Campanula	■	iAI or	3	jamr	O Canaries	1696. 11 r.m Bot. mag. 444	
9080 isvigKta G. Don	smoothed	■	AI	3	ja.ap	O Canaries	1825. R r.m	
1099. FRANKEWLrf L.	SEA HEATH.	(John Franckinus, prof. bot. at Upsal.)		Frankeniace. 9.—16.				
9081 lajvis L.	smooth	■ fr A cu	ijl.au	F EnRland sal. m. D s.l	Eng. bot. 205			
»82 Nöftria Thun.	Nothria	■ f	ijn.au	F C. G. H. 181H. 1) s.l	Berg. c 171.1.8			
'083 nodiflora Lam.	knot-flowered	■ f	ijjl.jl	F C.G.H. 1818. D s.l	Lain. il. 262.4			
9084 hirsfta Sib.	hairy	■ cu	i jnjl	L.B Siberia	1789. D s.l Fl.gr 343			
hlsptida Dec.								
9085 intermédia Dec.	intermediate	!L ^ cu	■	W S. Europe	1817. D s.l	Mic. gen. 22.2		
hirshta Dec. not Sm.								
9086 pulvern'lénta L.	powdery	■	■	R England c. co. S s.l	Eng. bot. 2222			
tmi corymbusa Desf.	corymbose	■	■	R Barbary 1819. D BI	Debt at. 1.93			
IXI88 mollis Bieb.	soft	■	■	R Caucasus 1824. D s.l				
9089 mcifolia Sm.	Heath-leaved	■	■	R Canaries 1816. D s.l				
•1100. PE^PLISZ.	WATER PURSLANE.	(Greek name of purslane.)		Saiicarke. 2. - 4.				
9090 Portula L.	common	O < ! U	^ jl.s	P Britain wa.pl. S aq	Eng. bot. 1211			
9091 altermflbia Bieb.	alternate-leaved	O cu	\ jl.au	P Siberia 1816. S co				
*1101. STEPHAW.4 Dec.	STEPHAMADA.	(F. Stephana professor at Moscow.)		Cappardice. 1.				
9092 clcomoldes Dec.	Cleomc-like	*C2 or.	.	. Caracas 1823. C l.p Jae.sc. III				
Capparis paradiba Jac.								

HEXA'NDRIA DIGY'NIA.

1102. ORY'ZAZ.	It ICE.	(Eruz, its Arabic name.)		Gramline*. 3.
9093 sattva L.	cultivated	M O ag 2 jl	Ap	E. Indies 1596. S aq Cat. car. 1 14
9094 latiflia Desf.	broad.leaved	O ag 2 au	Ap	S. Amer. 18'-0. S aq
9095 ncpalensis G. Don	Nepal	iQJag 2 au	Ap	Nepal 1818. S aq
1103. iiTRAPHA'XIS Z.	ATRAPHAXIS.	(A, privative, trepho. to nourish.)		Polwbae 2.
? 9095	prickly	«t _ J cu 2 au	Ap	Levant 1732. L l.p DoTbr 119
L.	ww&Ueaved	«t _ cu 2 jnjl	Ap	C. G. H. 1732. L i p DL d L & SS

1104. OXY'RIA Hill 9098 rcnifunii Sm. 7tumex digynus L. S americ;ia renifrmis Ph.	MOUNTAIN SORREL. kidney-shaped ^ A cu ft jn.jl (Osys, acid.) G Britain alp. pa. D p.l Eng. bot. 910 American £ A cu I jn.jl G N. Amer. ... D p.l	Polygbyttea?. J.
---	--	------------------

CLASS VI. ORDER 8.

HEXA'NDRIA TRIGYNIA.

1105. FLAGELLA^RIA L. FLAQELLARIA. (FlagCuum, rod; long pliant branches.) Mncece. 1 1. 9099 Indica £. Indian L O c u 7 jn.jl W India 1782. Skp.l Red. UL 257	
1106. SCHEUCHZE^RJ/f L. SCHIEUCHZERIA. (John and James Scheuchzer, German bot.) Juncaglne*. 1. 9100 palAstris L. marsh j£ ^ cu f myjji Br England sp. bo. O bog Eng. bot 1801	
1107. TRIGLCPCCHINZ., ARROW GRABS. (TreU, three, glochin, a point; capsule.) JuncagUic*. 6.—12. 9101 palAstro L. marsh 1 jl.au G Britain wa.mc. 1) bog Eng. bot. 366 2 Dirksonu Dickson'a 2 jl.au G England mar. 1) bog 9102 derpiens R. Br. deceiving 1 jl.au G N.HoU. 1820. D bog 9103 Barrchéri Loi. Barrcher's 1 jl.au G Italy 1820. 1) bog 9104 inarltimum L. sea 1 my.au G Britain sal. in. D bog Eng. bot. 255 9105 ellitum Nut. tall 1 my.au G N. Amer. 1818. D bog 9106 bulbubuin Z. bulbous-roofed 1 o P C. G. H. 1806. O bog Bot. mag. 1445	
1108. LICHTENSTEI'N/i* W. LICHTENSTEINIA. (M. von Lichtenstein, a Prus. trav.) Melanthacac. 2. 9107 lievigata IV. smoothed £ iAI pr 1 ... B C. G. H. 1824. O BI Cyniitton lujvigatum Spr. 9108 undulata W. wave-leaved £ iAJ or 1 jl.au B C. G. H. 1814. O s.l	
1109. MYRSIPH'LLUM W. en. MYRSIPHYLLUM. (Myrsine, a myrtle, leaf.) Asphodelccc. 2. 9109 asparagoides W. en. Asparagus-like 4 iAJ cu 6 imr.o G.w C. G. 1702. R Bp Her. lug. 681 Malbola asparagoides L. 9110 angustiflhum W. narrow-leaved \$ iAl cu 6 mrd G.w C. G. H. 1752. R s.p Til. pis. 17.12. 2	
1110. TOFJE^LIXM Hud. TOFIELDIA. (Mr. Tqfield, a friend of Hudson the bot.) Melanthheeee. 5.—8. 9111 paKists Sm. marsh j£ A cu 1 jl.au G Britain bgs. m. D m.s Eng. bot. 536 9112 alpina Sm. alpine • jg A, CU i jl.au G N. Europe ... D m.s Red. hi. £56 Yt.ll(ix,trs)Dec. 9113 stenopetila Sm. narrow-petaled £ A cu i my G N. Amer. 1820. D m.s 9114 pubtbeens Dry. pubescent £ A cu * ap.my W N. Amur. 1790. D m.s Pluk. m.342.3 9115 glutinosa Ph. glutinous £ A pr I ... W N. Amer. 1825. D p.l	
1111. 3/ELA'NTHIUM L. MELANTHIUM. (Melas, dusky, antlos, flower.) Melanthhcece. 8—12. 9116fframlincum Cav. grassy tf tAI cu 1 myjn W Mogadore 1823. O s.l Cav. ic. 587.1 9117junceum IV. liu»/eaved tf tAI cu Jjn.n Pk C.G. H. 1781. O s.p Bot. mag. 558 9118sucimlinum Lam. suie-flowering tf tAI cu 1 jn.n W C.G. H. 1812. O s.p Lam.ih26s).2 9119 umflbrum Jac. one-flowered tf tAI cu 1 myjn L.Y C.G. H. 1787. O s.p Bot. mag. 767 Tiliipa Breyniana L. 9120 capense W. Cape tf iAJ cu 4 myjn Br.v C. G. H. 1768. O 8p 9121 ciliatua Thun. ciliated tf iAJ or i jn.jl P.T. C. G. II. 1810. O s.p 9122 sibinicum L. Siberian tf A, or 1 Siberia 1823. O 8. 9123/ihalangioldes IV. Phalangium-lk tf A or 1 jn.jl W Carolina 1810. O s.l Jac. ic. 2.419	
1112. ANGUILLA'IUA R. Br. ANUUILLARIA. (AnguiUa, an eel; twisted seed.) Metanthdce*.3.—5. 9124 indica R. Br. Indian £ AJ or imy,jl D.P. Tranqueb.1818. O s.p 9125 biglamulosa R. Br. biglamulous £ AJ or \ myjn P N. Holl. 1818. O s.p 9126 dioica if. Br. dioecious £ AJ or i myjn P N. S. W. 182a O s.p	
1113. ORNITHOGLO'SSUM &/. BIRD'S TONGUE. (Ornis, bird, glossa, tongue; petals.) Melanthhcete. 2. 9127 undulatum Suit. waved tf iAJ or au.o G C. G. H. 1825. O s.p Sw. fl. gar. 131 9128 viride Sal. green-flowerd tf iAI or io.n. G C.G.H. 1788. O s.p Botmag.994 Jtfelanthium viride Thun.	
1114. MEDEVJ./4 L. MKHEOLA. (Medea, the sorceress; med. qual.) Smilukeee. 1. 9129 virginica L. Virginian £ A cu \$ jn Y. Virginia 1759. li Bp Bot. mag. 1316 Gyrbnia virginica Nut.	
1115. XEROPHY'LLUM Mx. XEROPITLLUM. (Xeros, dry, on, a leaf; leaves.) McLanthhceee. 3. 9130^raminctim M^/. grassy j£ A or 2 myjn N. Amer. 1812. R sp Bot mag. 1599 Hultnias ^raminica Ker 9131 tf-nax Nut. • tough-frawrf j£ A or 1) myjn W N. Amer. 1811. R s.p Ph. am. 1.9 HcK>nia< tf-nax Ph. 9132 asphodeloides Nut. Asphodel-like £ A or li myjn W N. Amer. 1765. R s.p Bot. mag. 748 setifolium Mx. Helunias asphodeloides L.	
1116. WU'RMB^ii Thun. WURMBEA. (F. Fan Wurmbs, Sec. Ac. of Sc. at Batavia.) Melanthilcecc. 4. 9133 longiflira W. long-flowered tf iAI cu \ myjn W C. G. H. 1788. O 8.1 Bot mag. Vf/J/ 9134 purpurea Dry. purple tf iAJ cu 1 myjn P C. G. H. 1788. O 8.1 Bot.mag.fJ't Afelanthium spicktum Hout. 9135 camnanufata W. cam pa nu late tf LAI or \ my.jl W C. G. H. 1819. O s.l A/elinthium monopetalum L. 9136 phmila w. dwarf tf .Al or \ myjn W C. G. H. 1800. O s.p	
1117. ANDROCY'MBIUM L. ANDROCYMBU'M. (Aner, anther, kymbion, a saucer; stam.) Melanthuete. 3. 9137 euromoldes W. Eucomis-like tf iAI cu \ mr.my G C. G. H. 1794. O s.p Sw. fl. gar. 165 Azclinthium eucomoldes Jac. 9138 melanthioides W. Melanth.-like tf iAI cu \ jn.jl W C. G. H. 1823. O s.p 9139 volutare Burc. tolloA-lcaved ^ iAJ or ap.my W C. G. H. 1816. O s.p	

*1118. TRI'LLIUMZ.	TRILLIUM. (<i>Trilix</i> , tissue of three threads; triple-leaved calyx.)	<i>Melanthhcct.</i>	14.
9140 sessile /.	sessile-leaved A A or f ap.my Br	N. Amer. 1759. R gp	Bot mag. 40
9141 netio Kitum PA.	petiolate A A or Jap.my Br	N. Amer. 1811. R s.p	
9142 crythrocarpum Mx.	red-fruited A A or i my.jn K,w	N. Amer. 1811. 11 s.p	Sw.fl.gar.S13
pictum Ph.			
9143 ovatum Ph.	ovate ■ A or i my.jn Pa.P	N. Amer. 1812. R s.p	
9144 p&milum Ph.	dwarf ■ A or A my.jn R	Carolina 1812. R s.p	
pusillum Mx.			
9145 cernuum L.	drooping-jftMt A A or 1* ap.my \W	N. Amer. 1758. R s.p	Bot mag. 954
9146 erectum L.	erect A A or i ap.my Br	N. Amer. 1759. R s.p	Bot. mag. 470
rhomboideum Afr.	purp&rcum Ker		
2 Albun	white-Jtimicri'd A A or ap.my W	N. Amer. 1800. R ap	Bot mag. 1027
9147 obovatum Ph.	ohovatc-leaued A A or i ap my R	Canada 1810. R Bp	
9148 undulatum IV.	vraeui-petaled A A or ap.my R	N. Amer. 1818. R s.p	
9149 pendulum W.	pendulous A A or i ap.my W	N. Amer. 1805. R s.p	W.h.b.1.35
erectum § Bot.mag.IWn			
5150 Catesbe' Ell.	Catesby's A A or * my R	Carolina 1820. R fl.p	Cat car. 1.45
9151 ncrovosum Ell.	ncrvose A A or 4 ap.my R	Georgia 1820. R s.p	
9152 stylidsum Nut.	stylose A A or ap.my II	Carolina 1823. R s.p	
9153 grandiflbrum Sal.	large-flowered A A or * ■■■ W	N. Amer. 1799. R s.p	Par. Ion. 1
crythrocarpum B. Af.			

1119. CO'LCHICftn L.	MEADOW SAFFOV.	(<i>Colchis</i> , its native country.)	<i>MelatUhhcte.</i> 9.
M5 autumibile L.	Byzantine tf A or i so P	Levant 1629. O s.p	Bot mag. 1122
2 album	autumnal. tf A or i so P	Britain mead. O s.p	Eng.bot. 133
3fuhisvariegatis	vhfite-flowered tf A m i so W	Britain mead. O s.p	
4 flbra pfeno	variegated-lvd 3' A m S.O P	Bnt tin gard. O s.p	
01.% crociflirum Sims	double-flwd tf A or P	Britain gard. O s.p	
9157 alptnum Dec.	Crocus-flwd tf A or : au.s P	S. Europe 1820. O s.p	Bot. mag. 2673
9158 arenarium Kit.	alpine tf A or P	Apciniin. 1820. O s.p	
9159 umbrsbm Fis.	sand tf A or ■■■ P	Hungary 1816. O s.p	"W.&K.9.179
D160 variegatum L.	shaded tf A or Pk	Crimea 1819. O pi	Bot. rej. 541
9161 tessellatum Mil.	chckcred-jftMtf A or * au.o P	Greece 1629. O pi	Bot. mag. 1028
9162 mont^nun L.	tessellated tf A or * au P	S. Europe 1600. O s.p	
Mere/ultra mont^na Ram.	mountain tf A or i au P	O pl Al пед. 1.74.2	
		S. Europe 1819., O p.	

1120. MERENDE'RA Bieb.	MERBNDBRA.	(A name given to <i>Colchicum</i> by the Spaniards.)	<i>Melanthoncete.</i> 1.
916J caucasica Bkb.	Caucasian tf A or \ au	P Caucasus 1823. O Bp	
Bulbocodium trigynum Adams,	C61chicum caucasicum Spr.		

1121. HELO'NIAS L.	HELONIAB.	(<i>Helos</i> , a marsh; habitation.)	<i>MeUmtkacete.</i> 3.-4.
9164 bullata W.	boss-garnished ■ A or 1 ap.my P	N. Amer. 1758. R s.p	Bot mag. 747
9165 erythrosperma Afr.	red-seeded ■ A or J j" "W	N. Amer. 1770. R s.p	Bot. mag. 803
9166 angustifolia Mr.	narrow-leaved f A or 1 my.jn "W	N. Amer. 1823. R s.p	
1122. KOIVXA Mx.	NOMVA.	(P. C Noñy, an American botanist)	<i>MelaniMcav.</i> 1.
9167 georgitaa Mx.	Georgian ■ A or Si jl.au W	Georgia 1812. R s.p	Pluk. m. 342.1
1*halangium virgatum Lam.			

1123. APONOGE^TON Thun.	APONOGETON.	(<i>Apom</i> , water.Gr//, <i>gciton</i> , neighbour, Gr.\ habitat.)	<i>Fluvitiles.</i> 4.
9168 monostKchyon /.	siinpLe-spiked ^ E3 cu { au.o Pk	E. Indies 1803. O pl	Bot rep. 406
9169 distkehyon H. K.	two-spiked * LAI cu \ rayjl W	C. G. H. 1788. O pl	Bot mag. 1293
9170 angustiflbum H. K.	narrow-leaved ^ LAI cu ft ap.s W	C. G. H. 1788. O pl	Bot. mag. 1268
"171 cribrum Thun.	curled-leaved ^ QS cu i jl.s W	Ceylon 1820. O pl	

9172 Adans&ni B. M.	Adanson's i 122 or 6 jn.au G	Florida 1810. S 81	Bot mag. 1434
C6rypha inhuor Jac.	Chuma^rops aca6lis Mx.	Rh?phis arctis W.	
9173 Blackburni)RA Lo,C.	Blackburn's f O or 6 ...	G tropics 1825. ■ s.l	
9174 grainimifia Lo. C.	Grass-leaved f CU or 6	G S. Amer. 1825. ■ s.l	
9175 umbraculifera Lo.	C. umbrella-bear. f O o r	Jamaica 1825. ■ s.l	
9176 PalmJU Lo. C.	Small Palm f O or 6	Georgia 1823. ■ s.l	

1125. L*VISTO^NIAA Br.	LIVISTONIA.	(Patrick Murray) of <i>Levistone</i> , near Edinburgh.)	<i>Pdlnue.</i> 2.
9177 inermis R. Br.	unarmed f or 10 - -	N. Hol 1824. S r.m	
9178 htmiiis R. Br.	humble f i J or 6 - -	N. Hol 1824. S r.m	

9179/7ydrolpathum Hud.	Water Dodk	* A ■■■ jl.au G	Britain riv.ba. R co Eng.bot. 2104
aqu^ticus Sm.			
9180 britannicus L.	British * A m 2 jn.jl G	N. Amer. ... R co	
»J81 m^ximus Schreb.	largest A un 6 jl.au G	Europe 1800. R co	
9182 glomeratus Schreb.	glomerato A un 2 jlav G	Europe 1818. R co	
9183 divaricatus IV.	divaricated A un 2 ll.au G	Italy 179J. S co	TIL pis. 37.2
9184 Ncmoldapatnum&ArA.	Grove Dock A un 3 jl.au G	S. Europe 1810. R co	
9185 nemortous Schr.	grove A un 2 1 G	Germany 1810. R co	
9186 condylbdcis Bieb.	knobbed A un 3 jl G	Caucasus 1816. S co	
9187 crlspus L.	curled A w 2 jn.jl G	Britain jrubble. R co	Eng. bot 1998
9188 Patientia /..	Patience A cul 4 jn.jl G	Italy 157a R co	
Sjj-J confertus %\	cose-headed A cul 4 jn.jl G	Europe 17J.W. R co	
SJS »»»»»»»»»»»»»»»»»»			
9191 •eninlftris /..	Brazilian Umy G	Brazil IK22. R co	
9192 spatulatus Thun.	whorled 2 jn.au G	N. Amer. 1820. R co	
carefiliflus Camp.	spatulatc 1 jn.au G	C. G. H. 1820. S co	
9193 acutus L.	*h*rp.leaved ■■■ G	Britain wa.pl. R co	Eng.bot724
9194 laevigatus W.	smoothed ■■■ G	Europe 1810. R co	
9195 sylvestris Wallr.	wood ■■■ G	Germany 1800. R co	
9196 heteroprh 11 us Schulz	variable-leaved ■■■ G	Germany 1818. R co	
9197 uncatus Trev.	hooked ■■■ G	Prussia 1826. R co	
9198 nepalensis Spr.	Nepal ■■■ G	Nepal 1817. R co	
IM- tuberosus Rox.			
»199 obtusif&lius L.	obtuse-leaved ■■■ 3 jn.au G	Britain rubble. R co	Eng.bot 1999
»^00 pilcher L.	fair ■■■ 2 jn.au G	Britain gra.pa. R co	Eng. bot. 1576

CLASS VI. ORDER 4.

HEXA'NDRIA POLYGY'NIA.

- | | | |
|---|------------------------|--------|
| 1129. WENDLA*NDIA W. WENDLANDIA. (J. C. Wendland, a German botanist) | <i>Metdspirhiee.</i> | 1. |
| 9248jDopuliflvia W. Poplar-leaved f. Jor 10 jnjl W Florida 1759. C co | | |
| Cucullus connatus Dec. | | |
| 1130. DAMASO^NIUM ScArct. DAMABOMUM. (<i>Damazo</i> , to subdue; med. qual.) | <i>Hydrocharfdeet.</i> | S. |
| 9249) Indicum W. Indian ^ [73 or 1 jl s W E. Indies 1800. S aq Bot mag. 1201 | | |
| 9250 ovalifolium R. Br. oval-leaved S. iAj or 1 jijl W N. Holl. 1824. S aq | | |
| 1131. ACTINOCA'RPU\$ R. Br. ACTINOCARPUS. (<i>Aktin</i> , ray, <i>karpos</i> , fruit; radiated.) | <i>Alismhcece.</i> | 2.-4. |
| 9251 minor R. Br. smaller A. A. my.au W N. S. W. ... S aq | | |
| 9252 Damastrum R. Br. Damasonium ^ A. jn.au W England dit. S aq Eng. bot. 1615 | | |
| illisma Damabonia L. Damasonium bellatum Pcrs. | | |
| 1132. JLI'SMA i. WATER PLANTAIN. (<i>Alis</i> , water, <i>Celt.</i> : habitation.) | <i>Alismhcece.</i> | 9.—11. |
| 9253 Plantago L. Plantain ^ A or 1* Jnjl P.w Britain pools. S aq Eng. bot 837 | | |
| 9254 lanceolata With. apcar-leafed f= A or H jnjl P.w Britain pools. S aq | | |
| 9255 parviflora Ph. small-floweredd A. A or 1 jl.au W N. Amer. IK 16. S aq | | |
| 9256 trivialis Ph. trivial A. A or li jnjl W N. Amer. 1816. Sk aq | | |
| 9257 cordifolia L. heart-leaved ^ E3pr 1 jl.au W W. Indies 1810. D aq M.h.15.4.6 | | |
| 9258 parnassifolia L. Parnassia-lvd ^ -Ajpri i jnjl W Italy 1820. D aq | | |
| 9259 natanis L. floating f= A or i jl.au W Wales allak. S aq Eng. bot 775 | | |
| 9260 ranunculoides L. Ranunculus-lk de A or i jau P. Britain tur.bo. S aq Eng. bot. 326 | | |
| 9261 repens Cav. creeping ik A F i jl.au W.p Wales&Eng.dit. D aq Cav. ic. 1.55 | | |

CLASS VII. ORDER 1.

HEPTANDRIA MONOGY'NIA

1133. TRIENTA^XLISX. WINTER GREEN. (*Trkns*, one third; humble stature.) *Primulacea*. 9.
 9262 europea^{ta} L. European £ A cu i my.jn W Britain m.wo. R s.p Eng. bot 15
 9263 americana Ph. American £ A cu ft jl.au W N. Araer. 1816. R s.p

1134. DISA'NDRA L. DISANDRA. (DJW, difficult, a ner, anther J number variable.) *Scrophularinaria*. 1.
 9264 prostrata L. trailing JUiAlor | my.au Y Madeira 1771. R P-1 Bot. mag. 218

• 1135. FISO ^S NIA L.	PISONIA.	(IV. <i>Piso</i> , a physician at Amsterdam.)	<i>Nyctaginea</i> .	11.—16.
9265 aculeata L.	prickly	IQcu 10 mr.ap G Jamaica 1739. C pl Lam.il. 861		
9266 fragrans Desf.	fragrant	• Dcu 3 1823. C pl		
9267 macroph Ua Lk.	long-leaved	H.I.leu 3 182J. C pl		
9268 nigrans Suns.	blackish	H1.leu 3 G.w W. Indies 1806. C pl		
9269 obovata Lk.	obovate	H1.leu 3 182a C pl		
9270 mexicana Lk.	Mexican	* Lleu 4 Mexico 1824. C pl		
9271 nitida W. en.	shining	• Dcu 3 Madagasc. 1824. C pl		
9272 grandis R. Br.	large	iLlor 12 N. Holl. 1805. C pl		
9273 subcordata Swt.	subcordate-wf	Tlror 30 Jamaica 1821. C pl		
9-74 obtusata Swt.	obiauw-leaved	Tlror 4 W. Indies 1824. C pl		
9275 hirtella Kth.	rather hairy	* 1lor 4 Mexico 1825. C pl		

1136. PETIVETRLf L. PBTIVERIA. (*J. Petiver*, F.R.S., apothecary, London.) *ChenojAdete*. 2.
 9276 galliacea L. Garhc-scented i t Q cu 2 jn.jl W Jamaica 1759. C pi Tr.chr. 33.67
 9277 octandra Jae. octandrous n-CJeu S jn.jl W W. Indies 1737. C pl PI. ic. 2J3.219

' 1137. ^E'SCULUS J.	HORSECHEST.VUT.	(<i>Esca</i> , food ; in fabled Golden Age.)	<i>Hippocastnea</i> .	5.
9278 HippoLast.uura L.	Horsech., or com. ^	tm 20 ap.iny W Asia 16-29. S co Schro. ar. 1.38		
2 vancgktuin		or If np.my W Asia 1629. S co		
9279 glabra IV. en.	Bmooih-leaved	or 12 my.ju G.Y N. Amer. 1812. G co		
9280 oh'u'nis Mx.	Ohio	or SO ap.my W N. Amer. ... S co		
9281 rubicunda Dec.	ruddy	or 12 jn> R N. Amer. 1820. G co Den.br. 121		
catnea Wat.				
9282 pallida IV. en.	pale-flowered	% or 12 jn G.T N. Amer. 1812. G co		

1138. VA^VIA Boer.	PAVIA.	(Pierre Paw, professor of botany at Leyden.)	<i>Hippocas/dnete</i> .	7.
9283 rufra Lam.	rod-flowered	it or 6 my.jn S N. Amer. 1711. G s.l Den. br. 120		
jE'sculus Pavia L.				
2 arguta	short-notched	■ or 4 ... R Europe 182a G co Bot.reg. 993		
9284 camea B R.	(Lcsh-colored	■ or 16 ... R 1820. G co Bot. reg. IUS6		
9285 hhmilis 6. Don	humble	■ or 4 my.jn R ■ ... G co Bot. reg. 1018		
JE'SCUIUB liumilis Lod.				
9286 h^briria Dec.	hybrid	■ or 4 my R Y Georgia 1812. G s.l		
jK'sculus discolor Ph.				
9287 fl?iva Dec.	yellow	■ or 20 my.ju Y N. Amer. 1764. G s.l Den.br. 163		
jEfrculus flXva //, K.				
9288 ncglcta G. Don	neglected	■ or SO my.jn Pa.Y 1823? G co Bot. reg. 1009		
JE'aculy* neglcta Lindl.				
0289 macrostachya Dec.	long-spiked	■ or 6 jn.jl W N. Amer. 1820. G co CoL h. rip. 19		
parviflra Walt.				
i&sculus macrostachya Mx.				

1139. JONE^SJi W. JOVBSIA. (*Sir William Jones*, a celebrated scholar and bot) *Leewntnb**. &
 9290 Asba Rux. Asoca f • or 20 ... O E. Indies 1796. C pl R.mal.5.5!)

9291 scandens llox. scandent |_CDor 20 -- O E. Indies 1820. C pl

1140. DRACO'NTIUM //-. DiUGOAr. (Draaron, a dragon; stems.) *Aroideite*. 2.-6.

9292 polyuhllum L. many-leaved L. 2 mrjn Ap India 17-39. R ltl Bot rep. 700

9293 spinosum L. prickly E. ap.my Ap Ceylon 1759. R ltl

1141. CALLA L. CALLA. (*Kallos*, beauty.) *Arode. 4.-5.*

9294 palft&tris L. marsh i ^ or jUu W N. Amer. 1768. 1) p Bot. mag. 1831

9295 aromatica Rox. aromatic j? iA) or 2"jl WW China 1813. D r.m Bot mag. 2279

9296 pertusa Kth. jwforated ^ El cu 6 ap.jn W W. Indies 1752. R p.l Jac.sc.SJ. 184.5

jRachnitum pertusum L.

9297 occulta Lou. hidden-spiked tf 1Alor my.jn W China 1817. D r.m Bot. cab. 12

1142. HICHA'RD/il Kth. RICHARDIA. (L. C. Richard, an eminent French botanist) *Aroidece*. 1.

9298 athiopica Kth. Ethiopian £ufijor 3 ja.my W C G. H. 1731. Sk r.m Bot mag. 832

Calla athiopica L.

1143. PARINATRIUM J. PARINARIUM. (*Parinar*L the Guiana name of the genus.) *Chrysobattne**.3.—b\

9299 excUjm G. Don tall f CD fr 60 ... W S Leone 1822. C 1

9300 macropylum G. Don long-leaved *• fir 3 ... W S. Leone 1822. C r.l

9301 campestre Aub. field iQor 6 ... WV Guyana 1824. C S.1 Aub.guLS06

Petrocarpa camp&tris W.

CLASS VII. ORDJCR 2.

HEPTANDRIA DIGY'NIA.

1144. il'MEUM L. LIVEUM. (*Loimos*, a pest; a poisonous plant) ' PorhOhcea 1—4.

8302 AfricānumX. African fciAIP f jnjl W C. G. H. 1774. K tip

CLASS VII. ORDER 3.

HEPTA'NDRIA TETRAGY'NIA.

1145. SAURUMtUS L.	LIZARD'S-TAIL. (<i>Saura</i> , a lizard, <i>oura</i> , a tail; flower.spikc.) <i>Piperhcete. X</i>
9303 cérnuus L.	drooping * A cu 2 s Ap Virginia 175a D s.p Ijim. ll. 1
9304 l&cidus Jae.	•hining fe A cu 1 s Ap N. Amer. 1791. 1 Jac ec. 1.18
9305 chintasis Jlott.	Chinese £3 AJ cu 1J ... Ap China 1819. D 1

1140. ASTRA'NTHUSIMU.	ABTHANTHOS. (<i>Aitron</i> , star, <i>anthot</i> , flower; segments.) <i>Eomnlineee. 1.—2.</i>
9306 cochinchinensis Lou.	Cochinchinese * LJ CU 4 jn,jl W* China 1823. C r.m Bot mag. 894

CLASS VII. ORDER 4.

HEPTA'NDRIA HEPTAGY'NIA.

1147. SE'PTASL.	SEPTA*. (<i>Septem</i> , seven ; all parts of flower divided into seven.) <i>Sempervfvea. 3.—4.</i>
9307 capensis B.	Cape £ TAI cu J au.s W C. G. H. 1774. II s.p Hot. rep. 90
9308 gloniiflora B. M.	globe-flowered £ LAJ cu \$ mr.ap W C. G. H. 1809. R s.p Bot. mag. 1472
9300 umbellata Haw.	umbel £ iAJ cu } jl W.G C. G. H. 1800. R s.p Jac. vin. 352

CLASS VIII. ORDER 1.

OCTA'NDRIA MONOGY'NIA.

* 1148. TROPJE'NOLUM L.	INDIAN CRIBS. (<i>Tropantyattropy</i> ; leaf and flower.) <i>Tropeble*</i> . 8.—13.
9310 nrtinus L.	smaller Jc Θ or 1 jn.o O.Y Peru 1596. S s.l Bot. mag. 98
2 ffra pleno	double-flowered! iAJ or J jn.o O.Y Peru 159ft. C 8.1
9311 maju* L.	greater £ O cul J jn.o O.Y Peru 1680. S 8.1 Bot. mag. 23
2 ffra píeno	double-flowered! LAJ or J jn.o O Peru Ifitti. C r.m'
9312 adduncum Sm.	hooked ISIGJor J jn.o O Peru 1775. S r.m Bot. mag. 1351
peregrinuin Jac>	
9313 phimutum Amtr.	pinnate-/faud + - J 2 jn.o P.o Peru -- C r.m Bot. rep. 535.
9314 hybndum /.	hybrid + - 4 jn.au O -- C rm Berg.ho. 32.1
9315 pcregilnum L.	strange + - 2 J.s O N. Grana. 1810. C si Jac.sc. 1.68
9316 pcntph^llum Lam.	five-leaved + - O M. Video 1824. C s.l Lam. il. 277
9317 polyn^llum Cav.	many-leaved + - 3 jn.Jl O.Y Chile 1827. C s.I Cav. ic. 4. 365

1149. ROXBUR'RGH [^]	Jon. ROXBOBGHIA. (JV. <i>Boxburgh,M.D.,6TCct.</i> of bot gar. Calcutta.) <i>Asphodylets. f L</i>
9318 gloriosoides Hog.	Gloriosa-like fl. (29 or 6 ap Pk.v R Indies 1803. Sk pl Bot mag. 1500

1150. GR'L'SLE [^] L.	GRISLEA. (ff <i>Gtixfey</i> , a Portuguese surgeon and author.) <i>Salicurke. 2.</i>
9319 tomentosa Jiox.	downy ■ prp 3 my.jn R K. Indies 1804. C s.p Bot reg. 30
9320 secunda L.	side-flowering ■ L or 4 — ... Cumana 1820. C p.l

1151. CASTE'Lif Turp.	CASTELA. (M. <i>Castel</i> , author of a poem upon plants.) <i>Bhdmea. 1.—2.</i>
9321 erecta Turp.	erect • Dor 4 . . . W.Indies 1825. C p.l

1152. BORONI'A Sm.	BORONIA. IF. <i>Borone</i> , a servant of Dr. Sibthorp.) <i>RutiHeeee. 10.—14.</i>
9322 /cdifolia Gay	Lcdum-leaved * LJ or 11 mr.ap W N. S. W. 1814. C s.p Ven. mal. 59
9323 pinniflta Sm.	pinniflta LJ or uJ 2 mr.ap N. Holl. 1824. C s.p Hot. reg. 58
9324 tt'traulra Lab.	tutrandrous A or 3 mr.jn N. Holl. 1825. C s.p Sw. au. ic. in.
9325 alata Sm.	winged a 1 or 3 mr.jn N. Holl. 1824. C s.p
9326 anemmesefolia Cun.	Anelliione-lvd H L or jnjl N. S.W. 1816. C B.p Bot reg. 842
9327 berrulata Sm.	serrulated tt J or mr.jn N. Holl. 1824. C s.p Sm.tr. 7
9528 /wlygal-a-folia Sm.	Polygal-leavctil 1 or mr.jl N. Holl. 1826. C B.p Lab. n. h. 1. 126
9329 pilosima Lab.	hairy-stamened H i_J or mr.jl N. Holl. 1823. C s.p Bot reg. 1000
9330 denticulata Sm.	denticulate • u.1 or mr.jl N. Holl. 1825. C s.p
9331 parad6xa Dec.	wonderlul il 1 or N. Holl. 1825. C *p

1153. TETRATHECA Sm.	TETRATHECA. (<i>Tetra</i> , four, <i>theke</i> , a cell; cells of anther.) <i>Tremndre*</i> . 6.
9332 juncea SOT.	rushy «. 1 J pr 2 jl.au P N. S. W. 1803. S A B. Sm. n. h. 1.2
9333 tfricasfolia Sm.	heath-leaved «. 1 J pr 2 jl.au P N. Holl. 1820. S sp Sm.ex.bot. 1.20
9334 pifosa Lab.	pilose IL i_J Pr 1 jl.au P V. Di. L. 1823. S sp Lab. ll. h. 1.122
9335 hymifolia Sm.	Thyme-leaved a. L pr 1 jl.au P N. Holl. 1824. S sp Sin. ex. bot. 22
9.136 glandulosa Lab.	glandulose «. 1 J pr 1 jl.au P V. Di. L. 1822. S sp Lab. n. h. 1. lJ3
9337 rubioides Cun.	Rubia-like tt. 1_T°r jl.au P N. Holl. 1825. S B.p

MVO [^] C ORR A A Sm.	ConnGA. (J. <i>Coma de Serra</i> , a celebrated Portuguese botanist) <i>Butucetc. 5.</i>
9338 alba B. Bon	white * \ J or fi apjl W N. S. W. 1793. C s.p Bot. rep. 18
9339 rdfa Gat.	reddish - Q or 6 apjl IV W N. Holl. 1821. C B.p I.Lab. v2 1.17
9340 speciosa A Bon.	showy * i J or 3 apjl S N. S. W. 180&. G sp Bot. reg. 26
9341 pulchella A. Ar.	pretty m i J or 5 apjl S N. Holl. 1824. G fl.p Sw. au. 1
9S42 virens Sm.	green^/owr<MI m i J or 6 my.n G N. S. W. 1600. C S.p Bot. reg 3

1155. PORLIE?IM R. St P. POHLIEQA. (*P. Antoine Porlier, a Spanish promoter of botany.*) *Zygophyllite.* 1.
 9343 hygrometrica ft. per. hygrometric a. • or - . . Peru 1820. C p.i s F.l. per. pr. 7
- * 1156. MIMU^SOPS L. MIMUSOFS. (*Mimo, an ape, ops, the face; flowers.*) *Sapote* <e. 5. — 6.
 9344 ENngi L. Elengi t. or 15 ... W E. Indies 1796. C p.l Rox. cor. 1.14
 9545 pamfölia R. Br. small-leaved t 1.J or 10. . . . N. Holl. 1821. C p. I
 9316 dissecta Spr. dissected J LJ fr 15. . . . S, Sea Itl. 1804. C p.l
/Pchras dissecta Font.
 9347 Kadki L. Kauki T. or 10. . . W E. Indies 1796. F p.l Ru. am. 3.8
 - 9348 hex&ndn Roz. hexandrous T. or 10. . . Y.W India 1804. F p.l
1157. SCHMIDE^LJi4 L. SCUMIDBLIA. (*G. C. Schmidel, prof, hot, acad. of Erlang.*) *Sapindtkcete.* 3.—18.
 9349 aerflata Dec. naw-leaved T. or 12. . . W E. Indies 1804. C p.l Rox. cor. 1.61
Ornitroph serrata Rox.
- 9350 racemosa L. racemose T. 15 my.jn W E. Indies 1820. F p.l Bur. in. 32.1
 9351 Cominia Swz. Cominia T. 15 my.jn W Jamaica 1759. F p.l SLjam. 2.208.1
Ornitrophe Cominia W.
- * 1158. CUPA^Nlif L. CUPANIA. (*Father Franc. Cupani, an Italian capuchin.*) *Sapindhcca.* 6.—13.
 9362 tomentifsa Swz. tomentose t. or 15. . . W W. Indies 1818. C p.l Pluk.am. 110,
 9353 exctlsa Kth. lofty T. 15 my.jn W Mexico 1824. C p.l
 9354 glabra Sun. smooth T. 15 my.jn W Jamaica 1822. C p.i
 9355 saponarioides Sun. Saponaria-like T. 15 my.jn W W. Indies 1810. C p.i
 9356 canescens Pert. canescens T. 15 my.jn W E. Indies 1818. C p.l
Mollnia canescens Rox.
 9357 dentata Sessi toothed T. 15 my.jn W Y
- IDor — — W Mexico 1824. C p.l Fl. raex. ic.in.
1159. EUPHORIA Com. EnpnoBiA. (*Euphoros, fertile; fruit.*) *Sapindhcete.* 4. — 5.
 3858 Litchi Desf. Lee-Chce T. C J f r 15 my.jn W China 1786. L r.m Lam.il. 306
Dimocarpus LitckKi Lou.
 3359 Long^na I'am. Longan £ C 3 f r 15 my.jn W China 1786. L r.m Buc.99
Dimocarpus Longdn Lou.
 9360 Acpelhelium Dec. Burdock T. 20 my.jl W E. Indies 1809. C r.m Lam. il. 764
Aepheleum /appaceum W. Scyfalia Ramboottan Rox.
 9361 rertillata B.R. whorled i 4fr 15 Sjnjau W.R EL Indies 1820. C r.m Bot. reg. 1059
1160. MELICO^CCAZ. HONET BEHHY. (*Mell, honey, kokkos, berry; flavour.*) *Sapinddcie.* 4. — 7.
 9362 bfjuga L. two-paired T. fr 16. . . Y Jamaica 1778. C lt.l Jac. am. 108.72
 9363 paniculata J. paniced T. fr SO. . . St.Domin. 1820. C c Mém. in. 3.5
 9364 olvys?ffnnis Dec. Olive-shaped T. fr 20. . . Y W. Indies 1824. C p.l
 9J65 trfjuga J. three-paired T. 20. . . . Ceylon 1820. C s.p M6m.HL 3.8
1161. BLVGHIA H. K. AKEE THEC (Capt. Bligh first carried bread-fruit to the W. Ind.) *Saphuticete.* 1.
 9366 sfpida H. K. savoury i O f fr SO. . . W Africa 1793. S r.m An.bot.2.16,17
1162. MATA^YBA Aub. MATAIBA. (*Vernacular name in French Guiana.*) *SapindHcetc.* 2. — 5.
 9307 guianensis Aub. Guiana S CD tm 60. . . W Guiana 1801 C p.l Aub. guil. 1. 128
- 9368 Patrotona Dec. Patris's ± Qtm 50. — W Guiana 1825. C p.l
1163. TALI'SIA Aub. TALIBIA. (*Toulichia, its name in Guiana.*) *SapindHcete.* 1. *— 4.
 9369 guianensis Aub. Guiana T. Q or 8. . . R Guiana 1824. C p.l Aub. gui. 1. 136
1164. THOU'I'N/4 Poit. THOUINIA. (*And. Thouin, of the Jardin des Plantes, Paris.*) *Sapindacce.* 1. — 8.
 9370 pinnata Turp. pinnate i O o r 8. . . W N. Spain 1823. C p.l An. mu. 5.26
1165. VITMX'KmA Vahl VITMANNIA. (*F. Vitmann, professor at Milan, 1792.*) *Malpighi&cete.* 1.
 9371 elliptica Vahl elliptic ± D or 20. . . Y Ceylon 1817. C 81 Gac. fr. 2.156. 3
1166. KÖLREUTE^R/^ Lax. KOLREUTEBIA (J. G. Kolreuter, a clcb. Ger. bot. and nat.) *Sapindiacea.* 1.
 9372 pppnchilita Lax panidled \$ or Jjau Qhina HG3- R co Blt.tre3.3^
1167. GUA^REA L. GI'AREA. (*Guara, its name in Cuba.*) *Mel&cece.* 3. — 5.
 9373 grandiflora Dec. great-flowered £ O tin 20 my.jn W S. Amer. 17552 C r.m Cav.dis.7.210
trichilioides L.
 9374 Swartzia Dec. Swartz's ± O tm 30 mr.jn W Jamaica 1818. C pi SI.jam.2. 170. i
trichilioides Sun.
 9375 ramiflora Ven. branch.flwd J • or 20. — W PortoRico 1824. C p.l Ven.ch. 41
1168. ylMY^RIS L. AMVRIS. (A. intensive, myrrha, myrrh; perfume of gum.) *Terebinthhceee.* 10. — 20.
 9376 Lunkiw Spr. Lunan's f • or 12 jn.jl W Jamaica 1820. C f.
 9377 sylvatica Jac. wood CJ or 16 Jn.jl 'W Carthag. 1793. f. v.
 9378 brasiliensis Spr. Brazilian E or 20 jn.o W Brazil 1818. f. v.
 9379 Plunieri Dec. Plunder's E or £0. . . W W. Indies 1820. f. v.
clemifura Dec.
 9380 heptaplifila Rox. seven-leafed £ or 20. . . W R Indies 1823. f. v.
 9381 toxifera W. poison-bearing £ or 20. . . W W. Indies 1818. f. v.
balsam! fora L.
 9382 acuminata Rox. acuminate £ or 20. . . W E. Indies 1823. f. v.
 J3S2 nána Uox. dwarf £ or 20. . . W E. Indies 1822. f. v.
 9384 maritima Jac. sea £ or 20. . . W S. Amer. 1810. f. v.
 '38] Tecomaca Dec. Teeomaca £ or 20. . . W Mexico 1827. f. v.
1169. BALSAMODE^NDRON Kth. BALPAMODENDRON. (*Balsamon, balsam, dendron, tree.*) *Tereblnt.* 1.—5.
 9386 zeyldnicum Kth. Ceylon f CD m 30. . . . Ceylon ... C p.l
Amfxn zelanica Retx.
- 1 im H K D W! O M Sw. HEDWIGIA. (*John Hedwig, a cryptogamist, died 1799.*) *Terebinthhcete.* 1.—2.
 9387 balsamifera Swz. balsam-bearing ± CD or 40. . . W W. Indies 1820. C Lp Gae. fr. 2. 2
1171. XIME^NJ^ L. XIMENIA. (*F. Ximenes, a Spanish naturalist*) *Olacinete.* 2. — 4.
 9388 americanæ IV. American f or 15. . . W W. Indies 1759. C s.p Jac. am. pic. 107
 9389 inermis L. unarmed f or 40. . . . W. Indies 1810. C Lp
- * 1172. BIE'CKIA L. BACKIA. (*A. Back, a Swedish physician.*) *Myrmcete.* S. — 9.
 9390 fruticescens L. frutescent £ Jpr 3 s.d W China 1806. L s.l Obbl.
 g^ virgata Ker slender £ Jpr 3 au.o W N. Caled. 1806. C s.p Bot reo 508
 SS' Kralus Cun. branliest £ Jpr 3 ... W N. Holl. 1826. C s.l p
 J3 ramostoma Cun. close-leaved £ Jpr 3 au.o W N. S. W. ... C s.p

9455 octagyna <i>Lo. C.</i>	eight-angled ft t J or	2	C G. H.	1816. C s.p
9456 purp&rea <i>Andr.</i>	purple-flowered ft J el	2	C G. H.	1789. C s.p
9457 elata <i>Andr.</i>	tall ft J or	2	C G. H.	1790. C s.p
9458 sulphurea <i>Lod.</i>	sulphur ft J el	2	C G. H.	1805. C s.p
9459 laniflora <i>Wnl</i>	woolly-floweredft J or	2	C G. H.	1790. C s.p
sordida <i>Andr.</i>	tube-flowered ft J de	2	C G. H.	1775. C s.p
9460 tubiflora <i>L.</i>	bucciniformis&J trumpet-form-J:wif	2	C G. H.	1818. C s.p
9461 simpliciflora <i>Donn</i>	simple-flowered ft J or	2	C G. H.	1774. C s.p
9463 Archeriana <i>Andr.</i>	Lady Archer's ft J or	2	C G. H.	1796. C s.p
9464 tpiria <i>Andr.</i>	spurious ft J or	2	C G. H.	1796. C s.p
9465 Hibbertiana <i>Andr.</i>	Hibbert's ft L J spl	2	C G. H.	1800. C fi.p
9466 conspicua // K.	conspicuous ft t J or	2	C G. H.	1774. C s.p
9467 linneoue Bedf.	Linnse'a like ft J or	2	C G. H.	1812. C s.p
9468 curviflora <i>L.</i>	curve-flowered ft J or	2	C G. H.	1774. C s.p
2 rrhra	red-flowered ft J or	2	C G. H.	1800. C s.p
9469 triphylla <i>Lk.</i>	three-leaved ft J spl	2	C G. H.	1822. C s.p
9470 monadelpha <i>B. M.</i>	monadelphous ft J or	2	C G. H.	1789. C s.p
9471 conclina // K.	neat ft J de	2	C G. H.	Bot. mag. 1370
9472 pellcida <i>Andr.</i>	pellucid ft J or	2	C G. H.	C s.p
9473 pellucidooides <i>Bedf.</i>	Pellucida-like ft J or	1	C G. H.	And. heaths 3
pellucida fibra <i>Andr.</i>	pellucida fibra	1	C G. H.	1800. C s.p
9474 viridiflora <i>Hort.</i>	rarfat-green-flwd ft L J or	2	C G. H.	1810. C s.p
9475 Lannuina Bedf.	Linnauis's fti J de	2	C G. H.	179a C fl.p
Linnaelz <i>Andr.</i>	perspScua /3 // K.			And. heaths 8
9476 clavata //or/.	clubbed green-flwdft u J or	n	C G. H.	1800. C s.p
9477 Bedfordina G. Don D. p. Bedford's ft J or	2 ap.my	2	C G. H.	And. heaths 2
Linnsdna superba 2fe<#			C G. H.	1800. C s.p
9478 hirsuta <i>Lod.</i>	hairy ft L	1	C G. H.	1800. C s.p
9479 pannota <i>Sal</i>	cloth.flowered ft L	1	C G. H.	Bot cab. 751
9480 erubescens <i>Andr.</i>	erubescenft L	1	C G. H.	C s.p
9481 Lechno H. K.	Lee's ft J	2	C G. H.	1800. C s.p
9482 colorans <i>Lod.</i>	coloring ft J or	2	C G. H.	1788. C s.p
9483 onosnuclora <i>Sal</i>	Onosma-flwd ft J el	1	C G. H.	1817. C s.p
9484 viridis <i>Andr.</i>	green-covered ft J cu	2	C G. H.	Botcab.224
9485 sanguinea <i>Lod.</i>	bloody ft J el	1	C G. H.	1789. C s.p
9486 longifltha <i>Dowm</i>	long-leaved ft J or	2	C G. H.	1800. C s.p
longifltha 2 Sal. pinca Wnl.			C G. H.	1815. C s.p
9487 ministriflbia Roll.	Pinaater-lyd m i lor	1	C G. H.	1808. Bot cab. 8d
2 rrhra	red ft J or	1	C G. H.	Ic.h.k.4
9488 jf nea Thun.	Pine-leaved m i J or	2	C G. H.	1810. C s.p
2 discolor	various-colored ft J or	2	C G. H.	179a C s.p
	spiral ft J or	2	C G. H.	C s.p
4 favoldes	honeycomb-lk ft J or	2	C G. H.	C s.p
5 echioïdes	Echium-like ft L J or	2	C G. H.	C s.p
6 pulchella	pretty ft J or	2	C G. H.	C s.p
7 purpurea	purple ft J or	2	C G. H.	C s.p
J ["] 9/moides JVWi	Flax-like ft J or	2	C G. H.	C s.p
9490 longiflora <i>Sal</i>	long-flowered ft J or	2	C G. H.	1806. C s.p
^ttl pitophyta Spr.	Pine-leaved ft u J or	2	C G. H.	1808. C s.p
piniflbia <i>Andr.</i>		2	C G. H.	1810. C s.p
2 spirkis	spiral ft i or	2	C G. H.	And. heaths 3
t. Ann 3 discolor	various-colored ft i or	2	C G. H.	1800. C s.p
9192 afirea >*irfr.	go\& colored ft i el	2	C G. H.	1810. C fl.p
9193 elongate Lad	elongated ft u J de	1	C G. H.	1799. And. heaths 2
S12 lanata WW.	woolly ft i or	1	C G. H.	Bot. cab 738
J12 Boweofti Lod.	Bowie's ft i el	1	C G. H.	Wen.er.5.5.ic
SH% coccinea L.	scarlet/fouN*raf ft i or	1	C G. H.	1822. Bot. cab. 842
9*97 && dans Zof	sweating ft J cu	1	C G. H.	1783. And. heaths 1
MW ostrina 7/Jf.	purple ft i or	1	C G. H.	1810. Bot. cab. 287
	unsavoury ft i or	1	C G. H.	1820. Bot. cab. 1218
Mas66ni Thun.	Masson's ft i gr	2	C G. H.	1800. C s.p
2	smaller ft i or	2	C G. H.	1808. C s.p
^J gemmifera <i>Lod.</i>	gem-bearing ft J spl	2	C G. H.	Bot cab. 457
J02 bicolor ^ndr.	two-colored ft i or	2	C G. H.	1802. C s.p
J03 extirgens <i>Andr.</i>	rising ft J spl	2	C G. H.	7 C s.p
2 major	larger ft J or	2	C G. H.	Bot cab. 835
3 grandiflira	great-flowered ft i or	2	C G. H.	1792. C s.p
4 carnea	flesh-colored ft i or	2	C G. H.	1800. C s.p
Spilliida	pale ft i or	2	C G. H.	1800. C s.p
g ^{ra} 04vesftta77kM ^m .	clothed ft J spl	2	C G. H.	1810. C s.p
1flba	white ft L el	2	C G. H.	1789. C s.p
2 incarflata	tLesh-colored ft i J spl	2	C G. H.	1808. C s.p
3 purpurea	purple ft i spl	2	C G. H.	1800. C s.p
	rosy ft i de	2	C G. H.	1800. C s.p
5 f61gida	bright ft J spl	2	C G. H.	1789. C s.p
6 coccinea	scarlet ft J spl	2	C G. H.	1800. C s.p
7lhcta	yellow ft i spl	2	C G. H.	1789. C s.p
8 mutabilis	changeable M L or	2	C G. H.	1789. C s.p
	elegant ft i snl	2	C G. H.	1800. C s.p
	roseate ft i el	2	C G. H.	1810. C s.p
Niveniana Andr.	Niven's ft J spl	2	C G. H.	1798. C s.p
q ^{ca} 2 minor	smaller ft i lor	1	C G. H.	Bot. cab. 782
507 aspera <i>Andr.</i>	rough m L	1	C G. H.	1799. And. heaths 2
	my.jn Y	1	C G. H.	1812. C s.p
		1	C G. H.	1802. C s.p
		1	C G. H.	And. heaths 3

II. VBrnicox. — Flowers much inflated.

Lady Monson's M. lei	2 fs	P.o	C G. H.	179a C s.p
Red Nightshades J or	4 ap.s	W	C G. H.	1787. C s.p
	1 my.au Y	C G. H.	1780. C s.p	Bot. mag. 1915
woolly *	lija.a	P.r	C G. H.	And. heaths 2
	2 my.au W	C G. H.	1803. C fl.p	And. heaths 3
Dickenson's ft J or	2 my.au W	C G. H.	1809. C s.p	
white ft J or	2 my.au W	C G. H.	1809. C s.p	
red * J or	2 my.au R	C G. H.	1809. C s.p	
smooth ft i J de	1 my.au W	C G. H.	1820. C s.p	
Honeywort-lk	4 roy.n D.S	C G. H.	1774. C *p	Bot mag. 230

2 major Snana	larger dwarf	H lor	4 my.jn	S	C. G. H.	1800.	C Lp	
9515 pectinifolia <i>Sal.</i>	pectinated-lvd	L or	1 my.jn	K	C. G. H.	1800.	C s.p	
9516 pripceps <i>Andr.</i>	princely	L or	H m i	S	C. G. H.	1800.	C s.p	
2 carnea	<i>tiesh-colorcd</i>	L or	1 my.jl	F	C. G. H.	1804.	C s.p	
9517 Swauhbn* <i>Andr.</i>	Swainson's	L or	2 JLlo	ILP	C. G. H.	1794.	C s.p	
9518 Sprengelia <i>Swt</i>	Sprengel's	L or	2 jn	Y.p	C. G. H.	1806.	C s.p	
tricolor <i>Spr.</i>							And. heaths 3	
9519 tricolor <i>Nuts.</i>	three-colored	M lor	2 jn.jl	R.o	C. G. H.	1810.	Rp	
2 minor	nailer	M lor	2 jn.jl	Ru	C. G. H.	1810.	+	
3 major	larger	L or	2 jlau	R.G	C. G. H.	IK10.	s.p	
9720 Templedn/i <i>Hort</i>	charming	L JW	i ap.s	L.P	C. G. H.	J800.	P	
9521 blanda <i>Andr.</i>	inflated		14 my.s	W.a	C. G. H.	1800.	And. hnntlisS	
9522 ferruginea <i>Andr.</i>	rusty	L or	1 my.jl	Ii	C. G. H.	1798.	C s.p	
9523 metutefibra <i>B. M.</i>	ninepin-flwd	M Ljel	1 jn.au	O	C. G. H.	1798.	C s.p	
9524 tumida <i>Ker</i>	tumid	L or	U my.s	S	C. G. H.	1812.	C s.p	
9626 tistuliefibra <i>Sal.</i>	pipe-flowered	L or	2 s	W	C. G. H.	1800.	Hot reR. tift	
9521 ubtilta <i>Andr.</i>	bottle	L or		W.E	C. G. H.	1791	And. heaths 3	
9528 acuminate <i>Andr.</i>	pointed-fcatwrf	L or	U	C. G. H.	1791	C s.p		
Spallida	pale	T jor	U jnjl	Pa.R	C. G. H.	IK20.	And. heaths 2	
							Bot. cab. 2 Jo	
III. — Flowers narrowed upwards, with a spreading border.								
9529 Lawsbni <i>B. M.</i>	Lawson's	ft	LJ el	N p.jn	F	C. G. H.	1802.	C +
9530 ventricosa <i>Thun.</i>	bellied	ft	i J iu	N ap.s	F	C. G. H.	1787.	C Bot mug. 35u
2 coccinea	scarlet	ft	LJ or	N ap.s	S	C. G. H.	-	C s.p
3 stellifera	star-bearing	ft	LJ or	N ap.s	F	C. G. H.	-	C b.p
4 c&neea	flesh-colored	ft	i J or	N ap.s	F	C. G. H.	-	C Bp
5 alba	white	ft	LJ or	N ap.s	W	C. G. H.	-	C s.p
6 supgrba	superb	* LJ or	N ap.s	S	C. G. H.	-	C s.p	
7 erecta	erect	ft	LJ or	N ap.s	F	C. G. H.	-	C s.p
8 nana	dwarf	ft	LJ or	N ap.s	F	C. G. H.	-	C Bp
9 hirsute	hairy	* LJ or	N ap.s	F	C. G. H.	-	C Bp	
9531 praCgnans <i>Andr.</i>	swelled	ft	LJ or	1 my.jl	K	C. G. H.	1796.	C s.p
9532 glutinosa <i>Berg.</i>	glutinous	ft	LJ or	jl.o	P	C. G. H.	1787.	C s.p
2 drosorooides <i>Lam.</i>	Sundew-like	ft	LJ or	jl.o	P	C. G. H.	1787.	C s.p
9533 gracilis <i>Lod.</i> not <i>Wnl</i>	slender	ft	LJ or	l.my	P.E	C. G. H.	1812.	C b.p
9534 vernici flua <i>Sal.</i>	varnish-flowing	ft	i J or	mr.s	P	C. G. H.	1804.	C s.p
9535 tetragona <i>Thun.</i>	quadrangular	ft	LJ or	N Y	L.Y	C. G. H.	178a	And. heaths 3
pugionifolia <i>Sal</i>								
9536 dense <i>Andr.</i>	dense	L or	1 my.au	R	C. G. H.	1810.	C s.p	
9537 libyuwa <i>Andr.</i>	Irby's	L or	jn.o	W.a	C. G. H.	1800.	C s.p	
9538 asminiflora <i>Andr.</i>	Jasmine-flwd	L or	jn.n	W.pk	C. G. H.	1794.	C t.p	
2 alba	white	L or	2 jn.n	W	C. G. H.	1790.	C s.p	
9539 ampullacea <i>Cur. flask</i>	Ampullacea-ik	ft	LJ or	2* myjl	R.Y	C. G. H.	1800.	C Bp
9540 ampullaciodes <i>Bed.</i>	Bandon's	ft	LJ or	2 jl.au	P	C. G. H.	1810.	C e.p
9541 Bandontina <i>Andr.</i>	Lady Shannon's				W.p	C. G. H.	1806.	C s.p
9542 Shannonitina <i>Lod.</i>	Rus&ell's		11	my.jn	Pk	C. G. H.	1820.	C s.p
9543 Russellfmi <i>Lod.</i>	recurred-fcaivdft	ft	LJ or	my.au	Pk.w	C. G. H.	1787.	C s.p
9544 rectirta <i>Thun.</i>	Blemler-flwd	ft	LJ or	ap.in	L.Y	C. G. H.	1800.	And. heaths 4
9545 tenuiflora <i>Andr.</i>	white	ft	LJ or	ap.in	W	C. G. H.	1818.	C s.p
2 alba	Flax-leaved	ft	LJ or	jl.jl	P	C. G. H.	1796.	C s.p
9546 finifolia <i>Sal.</i>	slender	ft	LJ or	jl.au	R	C. G. H.	1800.	C s.p
9547 tenuis <i>Sal.</i> not <i>W.</i>	emarginate	ft	LJ or	—	P	C. G. H.	1802.	C np
9548 emarginata <i>Andr.</i>	Lady Clifford's	ft	LJ or	ap.my	W	C. G. H.	1812.	And. heaths ic
9549 Chifordiana <i>Lod.</i>	excelling	ft	LJ or	jn.n	W	C. G. H.	IK10.	C s.p
9550 pnc stans <i>Andr.</i>	Hyacinth-like	ft	LJ or	jn.au	Pk	C. G. H.	1798.	And. heaths 3
9551 AyacinthoulM <i>Andr.</i>	peaked	ft	LJ or	H my.s	W	C. G. H.	1797.	C s.p
9552 fastigata <i>L.</i>	WalkerMna <i>Andr.</i>						Bot. cab. 207	
infundibuliflora <i>Lk.</i>	funnel-shaped	ft	LJ or	1 ap.my	P	C. G. H.	1821.	C s.p
infundibuliflurmis <i>Andr.</i>	funneUhpd	ft	LJ or	2 a <u>n</u> s	Pa.R	C. G. H.	1802.	C s.p
Aitonidna <i>Andr.</i>	Alton's	ft	LJ or	2 jn.s	W.p	C. G. H.	1790.	C s.p
9556 lutea <i>L.</i>	yellow	L or	ff.my	P.Y	C G. a	1774.	C s.p	
2 alba	white	L or	4 ap.au	P.Y	C G. a	1810.	C s.p	
9557 comosa <i>L.</i>	tufted	L or	f ap.au	R	C. G. H.	1787.	Ic. h. k. 18	
1 rabra	rea-flowered	L or	f ap.au	R	C. G. H.	1787.	Wen. er. 127 ic,	
2 alba	vrihle-flowered	L or	f ap.au	W	C. G. H.	1787.	C Bp Ami. heaths 2	
9558 Musckri <i>Andr.</i>	Muscari	ft	LJ or	1 mr.jl	W	C. G. H.	1790.	C s.p
9559 muscanoldes <i>Hort.</i>	Muscari-like	ft	LJ or	2 jn.jl	C	C. G. H.	1800.	C s.p
9560 rfaphneflibra <i>Sat.</i>	Daphne-flwd	ft	de	It ap.my	Pa.P	C. G. H.	1791.	Bot. cab. 543
9561 carnL-old <i>Lod.</i>	Daphne-flwd	ft	de	It ap.my	Pa.P	C. G. H.	1810.	C s.p
9562 rfaphnoides <i>Lod.</i>	Bub-flesh-CM	If	jl	Pk	C. G. H.	1800.	C s.p	
mirabilis <i>HorL</i>	Daphne-like	L or	2 myjn	W	C. G. H.	1800.	Bot. cab. 154	
9563 decolbrans <i>IV.</i>	discoloring	ft	LJ or	myjl	P	C. G. H.	1812.	C Bp
9564 Parmentiendntf <i>Lod.</i>	Parmentier's	L Jde	jl.au	Pa.P	C. G. H.	1810.	C fl.p	
2 rbsea	rose-colored	L Jor	jl.au	R	C. G. H.	1810.	C Bp	
9565 Bonplandtini <i>Lod.</i>	Bonpland's	L Jor	mr.s	Pa.Y	C. G. H.	1812.	C s.p	
9566 struthiolajflora <i>Lod.</i>	Struthiola-flwd	ft	LJ or	—	C. G. H.	—	C s.p	
9567 Humenda <i>Lod.</i>	Sir A. Hume's ft	LJ or	—	Pk	C. G. H.	1808.	C s.p	
9568 denticulata <i>L.</i>	denticulated	• L-Jor	ap.my	P	C. G. H.	1821.	C s.p	
9569 radiata <i>Andr.</i>	radiated	ft LJor	ap.my	C	C. G. H.	1798.	C s.p	
9570 aristata <i>Andr.</i>	awned	• L-Jor	11 mr.au	D.P.W	C. G. H.	1801.	And. heaths 1	
9571 aristella <i>Bed.</i>	small-awned	ft LJor	1 jn.jl	R	C. G. H.	1806.	And. heaths 3	
aristata mUior <i>Hort.</i>								
9572 tricolor <i>Swt.</i>	three-colored	* L - J or	...ny.au	P.Y.O	C. G. H.	—	And. heaths 3	
aristata minor <i>Andr.</i>								
9573 primulodes <i>Andr.</i>	Cowslip-like	ft	LJ or	4 apjl	P.R	C.G.H.	1802.	C s.p
9574 mdndula /,«*.	neatish	L or	2 fo	P	C. G. H.	1810.	C s.p	
9575 moschata <i>Lod.</i>	larger.	ft LJor	2 fo	P	C. G. H.	1810.	C s.p	
9576 conckva <i>Lod.</i>	musk-scented	L or	If myjl	G	C. G. H.	1805.	Bot. cab. 614	
9577 Coventryana <i>Lod</i>	concave	L or	f mr.ap	Pa.P	C. G. H.	lh08.	Bot. cab. LM	
9578 Jiroadleydna <i>Andr.</i>	Lord Coventry's	pr	1 jn.jl	Pk	C. G. H.	1808.	C s.p	
9519 wluSmM <i>Lqd.</i>	Broadley's	A L-J or	2 ...	R	C. G. H.	1800.	And. heaths ic	
	admirable	ft	LJ or	1 my.jn	P	C. G. H.	1800.	C f.p

gig^o efosa *Lod.***S**9583 n^obilis *Ro C.*
9584 notabilis *Wnl.*
9585 armata⁵ar.
9586 trissula *Lod.*
S riibra

fgnawed

quadrangular

July

noble

notable

armed

spruce

red

calycina W.

9589 calycinoides *Bed.*calycina capitata *Hort.*glomerate *Andr.*

large-calyxed

• i_J or

2 my.jl

P

C. G. H.

1799.

C s.p

And. heaths 3

Bot mag. 1250

9591 andromeda²/ZdrA *Andr.*

Androm.flwd

Hi_I pr

2 mr.jn

Pk

C. G. H.

1803.

C fl.p

9592 cumulaeflora *Sal.*

hecp-flowered

■ ■

1 my.s

P

C. G. H.

1801.

C s.p

9593 /?enistsfulia 5/a.

Genista-leaved

■ ■

1 jn.d

P

C. G. H.

—

C s.p

9594 elegans *Andr.*

elegant

■ ■

J el

ft

mm

G

C. G. H.

1799.

C 8.p

Bot. mag. 966

9595 triflora *L.*

three-flowered

■ ■

J or

1 mr.jn

W

C. G. H.

1774.

C ap

Wen. er. 12.13

9596 flagellaris *Lk.*

twiggy

■ ■

J pr

1* my

P.Y

C. G. H.

1820.

C s.p

9597 bracteata *Thun.*

nrrf-bracted

■ ■

J or

f my.jn

II

C. G. H.

1800.

C s.p

9598 turgida *Lk.*

turgid

■ ■

J el

1 apjl

P

C. G. H.

1821.

C s.p

9599 lachnaBfflia*<*irfr.

Lachnca-leaved

dtt

J dc

Iftmyjl

W

C. G. H.

1793.

C s.p

black-tipped

*■ Jcu

f mr.jn

R

C. G. H.

1790.

C s.p

9600 varia *Bed.*

various

*■ J or

1 ji

PY

C. G. H.

1810.

C s.p

^ Cassbnii *Hort*9602 salax *Sal.*9<3 biccans *L.*P<>4 flagax *Sal.*sWb triflmphans *Lod.*9606 phylcodes *W.*9007 incurva *Wnl.*9K8 tenuifolia *L.*9%9 aptrta *Spr.*9610 Thunbergia *W.*9611 faxifolia //, *K.*9612 petiolata *Thun.*9613 vesieuPins *Sal.*9614 imbrickta *L.*9>5 microphylla £, *o£*96(6) velleriflora *Sal.*9617 bruniddro *L.*9f18 capitata *L.*

9619 nodiflbra 5/a/.

9620 patens *Andr.*9621 fimbriata *Andr.*

9622 melanthera 7%v.

9f33 flaccida *Lk.*9624 Solundm *Andr.*9625 humifbsa *Sal.*9R26 spirlis *Nois.*9H7 sexfaria //, *Jf.*9f128 pachyphilla⁵a/.9629 coryd U8 *Sal.*9630 pallida *Wnl.*

<< 2 rrbra

Saeswaf9633 oppositifolia *Andr.*

2 rubra

9634 biflbra *Lk.*9615 obcordata *Nois.*

9636 spumbsa i.

9637 vulgaris *L.*Callfnia vulgäriss *Sal.*

2 alba

3 fluc pit-no

4 decumbens

5 tomcntbsa

G variegkta

7 coccinea

ai.

ra Sal.

K&rida Andr.

K mfind Sal.

<< SJ Pe H¹c¹B¹o¹ Sal.

Bfr.

9648 S,S⁵ S,⁵ * ii⁵ Brf.

TM*Y cupped

gown^{ed}

obnqua Thun. not W.

9650 chl&iAydiflbra *Sal.*

cloak-flowered

9651 canaliculata *Andr.*

channeled

callous

canaliculata minor *Andr.*

horizontal-wf

globeflowred

IV.

Gnaphalium-lk

carpenter's

Pink-leaved

s⁵ dianthiflbia *Sal.*

Calycina-like

il i_J or

2 jlS

R.P

C. G. H.

1789.

C s.p

And. heaths 4

IV. CALVIN*. — Flowers enclosed in the inflated calyx. —

conical

M\| or

i my.au

P?

C. G. H.

1830.

C s.p

Bot. cab. 1179

Coris-leaved

* L_I pr

1* au.d

Pa.P

C. G. H.

1774.

C BP

Bot. mag. 423

IV.

calycina W.

9589 calycinoides *Bed.*calycina capitata *Hort.*glomerate *Andr.*

large-calyxed

• i_J or

2 my.jl

P

C. G. H.

1799.

C s.p

And. heaths 3

Bot mag. 1250

9590 calycina W.

9591 andromeda²/ZdrA *Andr.*

Androm.flwd

Hi_I pr

2 mr.jn

Pk

C. G. H.

1803.

C fl.p

9592 cumulaeflora *Sal.*

hecp-flowered

■ ■

1 my.s

P

C. G. H.

1801.

C s.p

9593 /?enistsfulia 5/a/.

Genista-leaved

■ ■

1 jn.d

P

C. G. H.

1821.

C s.p

9594 elegans *Andr.*

elegant

■ ■

J el

ft

mm

G

C. G. H.

1799.

C 8.p

Bot. mag. 966

9595 triflora L.

three-flowered

■ ■

1 mr.jl

Y

C. G. H.

1820.

C s.p

9596 flagellaris Lk.

twiggy

■ ■

J pr

1* my

P.Y

C. G. H.

1820.

C s.p

9597 bracteata Thun.

bracteated

■ ■

J or

f my.jn

II

C. G. H.

1800.

C s.p

9598 turgida Lk.

turgid

■ ■

1 apjl

P

C. G. H.

1821.

C s.p

9599 lachnaBfflia 5/a/.

Lachnca-leaved

dtt

J dc

Iftmyjl

W

C. G. H.

1793.

C s.p

9600 varia Bed.

various

*■ J or

1 ji

P.Y

9658 lavandulieffbia <i>Sal.</i>	Lavender-lvd reddish short-leaved	m. lor ■ ■ ■ or	2 my.au S jn. 1 ja.au	F ^{...} Fk -	C.G. H. C.G. H. C.G. H.	1795. 1814. 1800.	C fl.p C s.p C s.p	Bot mag. 2165
V. GLOBOS&FLO^RM.—Flowers small, globose.								
9661 ardens <i>Andr.</i>	glowing	■ lor ■ ■ or	2 ap.jn 2 jl. ^o	S W	C.G. H. C.G. H.	1800. 1800.	C s.p C s.p	Bot.reg.115
9662 nftida <i>Andr.</i>	bright	■ lor ■ ■ or	2 jn.a	P	C.G. H.	1810.	C s.p	And. heaths 3
9663 nltens <i>Hort.</i>	shining	■ lor ■ ■ or	If mrjl	W	C.G. H.	1788.	C s.p	Bot. mag. 443
9664 physodes <i>L.</i>	flatulent	■ lor ■ ■ or	I	-	C.G. H.	1810.	C e.p	Bpt. cab. 153
9665 orbiculkris <i>Lod.</i>	orbicular	Z ■ ■ or	3 my.au	G.p	Portugal	-	C s.p	Lin. cr. 9. ic.
9666 viridiprophroa <i>W.</i>	green and purple!	■ lor ■ ■ or	S f.jn	W	S. Europe	1658.	C s.p	Fl.gr. 45
9667 arborea <i>L.</i>	tree	iN^lor ■ ■ or	5 f.jn	W	S. Europe	1658.	C s.p	
2 stylosa <i>Andr.</i>	long-styled	■ J pr ■ ■ or	4 f.jn	W	Europe	1800.	C s.p	
3 squarrusa	squarrose	■ lor ■ ■ or	1 my.au	O	C.G. H.	1803	C s.p	Bot mag. 1139
9668 resinosa <i>B. M.</i>	resinous	ll ujor ■ ■ or	1 my.de	W	C.G. H.	1800.	C s.p	And. heaths 2
9669 Lanibert* >>> <i>Andr.</i>	Lambert's	• ujde ■ ■ or	1 my.au	R	C.G. H.	1791.	C s.p	
©670 guttaeflora <i>Stil.</i>	spot-flowered	»L lor ■ ■ or	1 my.au	R	C.G. H.	1791.	C s.p	And. heaths ic.
incarnata <i>Thun.</i>								
9671 incamata <i>Andr.</i> not <i>Thun.</i>	flesh-cW	ill-Jor ■ ■ or	1 f.jn	Pa.R	C.G. H.	1790.	C s.p	And. heaths 1?
9672 ritbens <i>Thun.</i>	rcA.flowered	*l-lor ■ ■ or	1 jn.a	D.li	C.G. H.	1798.	C s.p	Bot. cab. 557
9673 minima // <i>Ccl.</i>	least	ill-Jor ■ ■ or	1	R	C.G. H.	1810.	C s.p	
9>74 I fbula <i>Lk.</i>	button	Ht-Jor ■ ■ or	1	P	C.G. H.	1823.	C s.p	
9675 axillaris <i>Thun.</i>	ax-flowered	• LJor ■ ■ or	1	Pk	C.G. H.	1798.	C s.p	
9676 strigifsa // <i>K.</i> not <i>Sal.</i>	meagre	m-Jor ■ ■ or	1	W	C.G. H.	1798.	C s.p	And. heaths ic.
axillariH <i>Andr.</i>								
9677 margantacca <i>Thun.</i>	pezr-Jllovcrcd	*l-Jcl ■ ■ or	1 my.s	W	C.G. H.	1775.	C s.p	And. heaths 1
9678 tomentusa <i>Sal.</i>	tomentose	*l-Jor ■ ■ or	1	P	C.G. H.	1798.	C s.p	
9679 totta <i>Thun.</i>	Hottentot	• l_Jor ■ ■ or	1	P	C.G. H.	1810.	C s.p	
ferox <i>Sal.</i>								
9680 pendula <i>IVnl.</i>	pendulous	m Jor ■ ■ or	1J jalau	P	C.G. H.	1791.	C s.p	Wen.er.10.13.ic.
9681 latralis <i>W.</i>	sulc-Jhwerec	*L_Jcu ■ ■ or	ll mrjl	R	C.G. H.	1791.	C s.p	And. heaths 1
9682 pedunculata <i>H. Cel.</i>	pedunculate	HI-Jor ■ ■ or	1 on	P	C.G. H.	1818.	C s.p	And. heaths 4
ciliata <i>Hurt.</i> not <i>L..</i>								
9683 filipctrifolia <i>L.</i>	Crowberry-lvd	ill-Jor ■ ■ or	1} ap.jn	P	C.G. H.	1774.	C s.p	Bot. mag. 447
9684 incirvita <i>Andr.</i>	incurred	*l-J de ■ ■ or	1 mr.my	W	C.G. H.	1802.	C s.p	And. heaths ic.
9687 claginifolia <i>Sat.</i>	Selago-leafed	ill-I or ■ ■ or	1 fjn	-	C.G. H.	1801.	C s.p	
9688 planul'lin <i>L.</i>	flat-leaved	il I or ■ ■ or	2 jl.s	P	C.G. H.	1795.	C s.p	Wen.er. 8 7. ic.
9687 A-erpyllifolia <i>Lod.</i>	Thyme-leaved	*l pr ■ ■ or	JnJl	W	C.G. H.	1810.	C s.p	Bot cab. 744
9685 mantlia <i>Thun.</i>	Marum-leaved	• l-J or ■ ■ or	2 my.jn	P	C.G. H.	1773.	C s.p	And. heaths 1
9689 purMnH <i>Hurt.</i>	fiery	il J or ■ ■ or	1 s.o	P	C.G. H.	1810.	C s.p	
9690 helianthincifolia <i>Sal.</i>	Helianth.-lvd	■ Jor ■ ■ or	2 f.ap	W	C.G. H.	1796.	C s.p	
9691 hwpialula <i>Thun.</i>	sub-bristly	<totrfi ■ ■ or	1 jn.au	P	C.G. H.	1791.	C s.p	
9692 (linusa <i>Andr.</i>	bushy	■ Jor ■ ■ or	1 ap.my	P	C.G. H.	1812.	C s.p	And. heaths ic.
9693 Schnlliiffla <i>Lod.</i>	Scholl's	■ Jcu ■ ■ or	1 my.jn	P	C.G. H.	1810.	C s.p	Bot cab. 538
9694 mollissima <i>Lo. C.</i>	softest	■ Jor ■ ■ or	1-my	W	C.G. H.	-	C s.p	
9695 Blaudorfifma <i>Andr.</i>	■ BlandfordV	■ Jor ■ ■ or	U mr.jn	Dk.Y	C.G. H.	1803.	C s.p	And. heaths 3
2lfsca	brown	■ Jor ■ ■ or	14 mr.jn	Crea	C.G. H.	1803.	C e.p	
9696 sanguinolinta <i>Lod.</i>	bloody	■ Jpr ■ ■ or	S my.jl	P	C.G. H.	1818.	C s.p	Bot. cab. 468
9697 pygmy* <i>Andr.</i>	pygmy	■ Jor ■ ■ or	i jl.o	R	C.G. H.	1806.	C s.p	And. heaths 4
9698 Suvil'Vam <i>Andr.</i>	Savile's	■ Jor ■ ■ or	2 dn	R.P	C.G. H.	1810.	C s.p	And. heaths ic.
9K99 carncolata <i>Nois.</i>	sub-Acah-cfcf	■ J or ■ ■ or	2 au.a	P.B	C.G. H.	1802.	C s.p	
9700 glilia <i>Andr.</i>	pipMn-Jlowrcd	m J or ■ ■ or	2 jl.o	P	C.G. H.	1817.	C s.p	And. heaths 4
9701 subufta <i>IVnl.</i>	subulate-ZeaMrd	»J or ■ ■ or	1* jn.jl	P	C.G. H.	1800.	C s.p	
9702 hnica <i>Spr.</i>	unique	il I or ■ ■ or						
pedunculata <i>Sal.</i>								
9703 aggregate <i>WnL</i>	clustered	m I I el ■ ■ ■ or	■ ■ ■	T	C.G. H.	1&D7	C s.p	
2 alba	white-Jfouwrai	il I-J el ■ ■ ■ or	■ ■ ■	■ ■ ■	C.G. H.	1822.	C s.p	
9704 perlata <i>Hort.</i>	very broad	M I-J or ■ ■ ■ or	■ ■ ■	■ ■ ■	C.G. H.	1810.	C s.p	
9705 stellifera <i>Andr.</i>	star-bearing	m IJ or ■ ■ ■ or	■ ■ ■	■ ■ ■	hybrid	1800.	C s.p	And. heaths ic.
9706 conglbla <i>Wnl.</i>	closc/eeded	a l-J de ■ ■ ■ or	■ ■ ■	■ ■ ■	C.G. H.	1820.	C s.p	
9707 stellaris <i>Nois.</i>	starry	il J or ■ ■ ■ or	■ ■ ■	■ ■ ■	C.G. H.	1810.	C s.p	Bot. cab. 893
9708 paniculata <i>L.</i>	panicld	sll-I or ■ ■ ■ or	■ ■ ■	■ ■ ■	C.G. H.	1774.	C s.p	Bot. cab. 1194
2 alba	white-Jloiuered	■ J or ■ ■ ■ or	■ ■ ■	■ ■ ■	C.G. H.	1774.	C s.p	
9709 flagelliformis <i>Andr.</i>	twig-fbrmed	■ l-J of ■ ■ ■ or	jaau	P	C.G. H.	1800.	C s.p	And. heaths 4
9710 suavéolens <i>Lod.</i>	sweet-scented	■ Jel ■ ■ ■ or	au	Pk	C.G. H.	1800.	C s.p	Bot. cab. H*
9711 tuberculifus <i>So.</i>	tuberculate	• L J or ■ ■ ■ or	f.au	P	C.G. H.	1790.	C s.p	
9712 amoAia <i>WnL</i>	pleasing	sVLJor ■ ■ ■ or	mrjl	P	C.G. H.	1795.	C s.p	Wen.er.17.73.ia
plumbosa <i>Andr.</i>								
9713 faVvis <i>Andr.</i>	smooth	Ai-Jde ■ ■ ■ or	my.jn	W	C.G. H.	1821.	C s.p	
2 alba	white	■ l-J or ■ ■ ■ or	my.jn	W	C.G. H.	1801.	C s.p	
9714 PcsUa <i>Lod.</i>	Pesiza	*L Jde ■ ■ ■ or	1 mr.s	W	C.G. H.	1812.	C s.p	Bot. cab. 265
9715 gracilis <i>Wnl.</i>	slender	• Ujor ■ ■ ■ or	f.jn	W	C.G. H.	1794.	C s.p	Wen.er. 8.9. ic
9716 Impccilla <i>Surf.</i>	weak	• L J or ■ ■ ■ or	1 Jjl	p	C.G. H.	1793.	C s.p	
gracilis <i>Sal.</i>								
9717 interyalaris <i>Sal.</i>	distant-iatwf	ill-Jor ■ ■ ■ or	my.n	p	C.G. H.	1800.	C s.p	
9718 nidularia Z<ML	nestling	• L Jpr ■ ■ ■ or	mr.ap	Pk	C.G. H.	1809.	C s.p	Bot. cab. 764
9719 preYox /or/	early	• l J or ■ ■ ■ or	ja.mr	P	C.G. H.	1820.	C s.p	
9720 Persdltuo Z^rf	Garland-flower	M J or ■ ■ ■ or	1J Em	P	C.G. H.	1774.	C s.p	Bot. mag. 342
2 alba	white	il J or ■ ■ ■ or	1 f.my	W	C.G. H.	1800.	C s.p	
3 rubra	red	il J or ■ ■ ■ or	1 f.my	R	C.G. H.	1800.	C s.p	
9721 grandinbsa lorf.	hailstone	*L J or ■ ■ ■ or	1 f.d	P	C.G. H.	1810.	C s.p	Bot. cab. 627
9722 pubescens <i>L.</i>	pubescent	■ J or ■ ■ ■ or	lf.d	R	C.G. H.	1790.	C s.p	
2 minor	smaller	■ i J or ■ ■ ■ or	lf.d	R	C.G. H.	1802.	C &p	
3 pubescéntior	more pubescent*	i J or ■ ■ ■ or	1 f.d	R	C.G. H.	1802.	C s.p	
9723 auriculhris 5a/.	eared	si LJ or ■ ■ ■ or	2 ap.au	P	C.G. H.	1800.	C s.p	
9724 modesta <i>Sal.</i>	modest	il LJ or ■ ■ ■ or	1 ap.my	P	C.G. H.	1800.	C s.p	
9725 m61hs Bedf.	soft	■ J or ■ ■ ■ or	?* my	P	C.G. H.	179a	C s.p	H. er. wo. 3.
9726 hirtiflora <i>B. M.</i>	hairy-flowered	■ cu ■ ■ ■ or	C.G. H.	1790.	C s.p			Bot. mag. 481
puhescens <i>Andr.</i>								
9727 mitrnf6rmis <i>Sal.</i>	mitre-formed	il LJ or ■ ■ ■ or	2 jn.jl	R	c. G. H.	1800.	C &p	
9728 Thun. not <i>Andr.</i>								
SJS *1810?1^a *.*	Cistus-leaved	il i J cu ■ ■ ■ or	1 my.jn	W	C. G. H.	1823.	C s.p	
2JS EZFiPV^	Constantia	* £j or ■ ■ ■ or	2 mr.au	P	C. G. H.	1810.	C fl.p	
9730 decunata <i>Hort.</i>	declined	il H or ■ ■ ■ or	1 s.o	P	C. G. H.	182a	C s.p	
97J1 mucosa <i>L.</i>	mucous	« LJ el ■ ■ ■ or	1 f.au	R	C. G. H.	17B7.	C s.p	And. heaths 1

9732 ramentacea L.	ramentaceoui	■ ■ ■	D.R.	C. G. H.	1786.	C s.p	And.heaths 1
9733 melittifera Lk.	honey-bearing	■ ■ ■	I	ap.my	P C	G. H.	1820. C Bp
9734 micrtstoma Berg.	small-mouthed	■ ■ ■	I	au.n	P C	G. H.	1810. C s.p
9735 odorala Andr.	perfumed	■ ■ ■	I	apjl	W C	G. H.	1904. C s.p Bot. cab. 633
9736 cruciformis Andr.	cruciform	■ ■ ■	I	ap.n	P C	G. H.	1800. C s.p And. heaths ic.
9737 canescens Andr.	cancsent	■ ■ ■	Ij	my.au	Pk	C. G. II.	1790. C s.p And. heaths 2
9738 para Ld.	pure	■ ■ ■	I	L-Jpr	f	aus.	W C. G. H. 1807. C s.p Bot. cab 72
9739 racemosa 7%tm.	racemose	■ ■ ■	I	Jel	li.ap.my	Pk	C. G. H. 1793. C s.p Wen.er.IO.3.ic.
9740 absmthoides L.	Wormwood-Ik	■ ■ ■	I	L-Jor	1A	mrjn	P C. G. H. 1792. C s.p
9741 scarita Thun.	scarious	■ ■ ■	I	Jor	I	jnjl	P C. G. H. 1800. C s.p Bot cab. 477
9742 campanulata Wnl.	be-flowered	■ ■ ■	I	Jel	I	ap.au	Y C. G. H. 1791. C s.p And. heaths 1
9743 scoparia L.	broom	■ ■ ■	I	i-pr	6	ap.my	G C. G. H. 1770. C s.p Lin. er. 14. ic
2 minima	least	■ ■ ■	I	Jor	3	ap.my	G C. G. H. 1800. C s.p
9744 triceps Lk.	three-headed	■ ■ ■	I	de	1	myjn	W C. G. H. 1820. C s.p Bot cab. 962
9745 montana Bedf.	mountain	■ ■ ■	I	Jor	2	o	P C. G. H. 1818. C s.p
9746 crossata Spr.	pot-flowered	■ ■ ■	I	Jor	2	myjn	P C. G. H. 1816. C s.p
9747 nteva Ld.	snowy	■ ■ ■	I	Jor	2	ap.my	W C. G. H. 1801. C s.p
9748 coarcftta Wnl.	crowded	■ ■ ■	I	Jor	1	myB	P C. G. H. 1822. C s.p
9749 if ctiefa Lk.	Actson	■ ■ ■	I	Jpr	1	myjn	Pa.P C. G. H. 1800. C s.p And. heaths 2
9750 conferta Andr.	crowded-Jltm?	■ ■ ■	I	de	II	fap	W C. G. H. 1780. C s.p
9751 glomeratesod. not Andr.	penicilhfltria Sal.	■ ■ ■	I	Jor	2"	fap	P C. G. H. 1792. C s.p Wen. er. 4.5. ic
9752 penicilhfltria Sal.	wAse-pencilled	■ ■ ■	I	T-CU	2	my.au	P.W.BrC. G. H. 1800. C s.p
calculata Wnl.							
9753 villosa Andr.	villous	■ ■ ■	J	L-J C U	2	fjn	W C. G. H. 1800. C s.p And. heaths 3
9754 taraeflura Andr.	turban-flwd	■ ■ ■	J	J pr	2	my.au	R C. G. H. 1800. C s.p And. heaths 3
9755 mutabilis Andr.	mutable	■ ■ ■	J	J or	§	fo	< C. G. H. 1798. C s.p
9756 obliqua W.	ohMque-leavcd	■ ■ ■	J	J or	1	au.0	P C. G. H. 1789. C s.p And. heaths 1
9757 flava Ld.	thrc-lvd yellow*	■ ■ ■	J	J el	1	ap.s	Y C. G. H. 1795. C s.p Bot. cab. 882
9758 folibsa Andr.	leafy	■ ■ ■	J	J or	2	Y	C. G. H. 1800. C s.p And. heaths 4
9759 flava H.K.	yellow	■ ■ ■	J	J or	2	my.o	Y C. G. H. 1705. C i.p And. heaths 2
2 imbricate B.M.	imbricated	■ ■ ■	J	J or	2	my.o	Y C. G. H. 1795. C sp Bot. mag. 1815
9760 ovata Ld.	ovate-flowered	■ ■ ■	J	J or	1	myjn	P C. G. H. 1811. C s.p Bot cab. 417
9761 ovaliflora Bo. C.	oval-flowered	■ ■ ■	J	J or	1	...	P C. G. H. 1817. C s.p
9762 decora Andr.	graceful	■ ■ ■	J	Jel	2	ja.n	P C. G. H. 1790. C s.p And. heaths 3
9763 hispida Andr.	hispid	■ ■ ■	J	Jor	2	jl.au	P C. G. H. 1791. C s.p Arid, heaths ic.
9764 cordata Andr.	heart-leaved	■ ■ ■	J	Jde	fapjn	W C. G. H. 1799. C s.p And. heaths 3	
9765 Passorina W.	Sparrow-wort	■ ■ ■	J	J or	2	my.n	W C. G. H. 1800. C s.p Pet g. 3.7
9766 calyculata Pers.	calyculate	■ ■ ■	J	J cu	2	myjl	P C. G. H. 1800. C s.p
9767 filiformis Sal.	filiform	■ ■ ■	J	J or	2	fjn	... C. G. H. ... C s.p
9768 setXcca Andr.	brMe-leaved	■ ■ ■	J	J or	Iff.ap	W C. G. H. 179a C s p And. heaths 1	
9769 tenuissima IVnL	Blendcrest	■ ■ ■	J	Jpr	1	fau	R C. G. H. 1803 C s.p Wen. cr. 6. 9. ic
9770 verecinda Sal.	reddish	■ ■ ■	J	J or	2	aps	R C. G. H. ... C s.p And. heaths ic.
cfrnua Andr.							
9771 floribunda Ld.	bundle-flwd	il	I-J cu	1	myjn	Pa.P C. G. H. 1800. C s.p Bot cab. 176	
VI OVATIFLO - Flowers small, not globose.							
9772 carduifolia 5aL.	Thistle-leaved	■ ■ ■	I	my.au	P	C. G. H.	1806. C Bp
9773 austriahs L.	southern	■ ■ ■	I	mr jl	P	Spain	1769. C s.p
9774 cinerca // K.	grey	■ ■ ■	I	jns	P	Britain	he. L s.p
2 alba	white-flowered	■ ■ ■	I	ec	1	jns	W Britain hca. L s.p
3 atropurpurca	dark-purple	■ ■ ■	I	or	1	jns	D.P Britain mo.he. L co
4 r&ura	red	■ ■ ■	I	or	1	jns	R Britain mo.he. L co
5 c&rnca	Aesh-colored	■ ■ ■	I	or	1	jns	F Britain mo.he. L co
9775 stricta Donn	6trict	■ ■ ■	I	or	2	au.n	P S. Europe 1765. C s.p And heaths S
9776 ramulosa Viv.	small-branchy	M'	or	2	au.n	P S. Europe 1800. C s.p	
2 rfcbra	red	M'	or	2	jnjl	R S. Europe 1800. C s.p	
9777 reflfexa Lk.	reflexed	il	I-J cu	1	myjn	W C. G. R. 1820. C s.p	
9778 cernua L.	drooping-Jlw	■ ■ ■	I	aud	P C. G. H. 1791. C Bp Bot cab. 823		
9779 lancolMa Pers.	epeax-leaved	il	I-J or	1	jnd	W C. G. H. 1791. C s.p Wen. er. 8.13.ic	
9780 leucanthera Andr.	white-antheredil	il	I-J or	Jja.my	W C. G. H. 1803. C s.p		
9781 infercians Lod.	blackish	il	I-J or	1	jna.u	C. G. H. 1816. C s.p	
9782 Tlralix L.	Tetralix	il	ec	1	jna.u	F ... Britain moi. h. L Bp Eng. bot 1014	
2 alba	yrhite-flowered	M	or	1	jna.u	W ... *~... L s.p	
3 cärnca	flesh-colored	M	or	1	jna.u	F Britain mo.he. L co	
9783 tewrifolia Spr.	Teucrium-lvd	41	I-J or	2	myB	- C. G. H. 1812. C s.p	
9784 cineraifcens IV.en.	cincrascent	■ ■ ■	I	Jor	1	ap.my	P C. G. H. 1810. C s.p
9785 complanata Nois.	Rat-flowered	■ ■ ■	I	J or	2	myjl	Y C. G. H. 182a C s.p
9786 approximata ScM.	approximate	■ ■ ■	I	J or	1	myjl	R C. G. H. 1823. C s.p
9787 vrs;ta IVnL.	twiggy	■ ■ ■	I	J or	1	myjl	P C. G. H. 1818. C s.p
9788 urceolaris Thun.	small-beaked	■ ■ ■	I	J or	1	myjl	W C. G. H. 1778. C s.p
9789 rottsttl Bed.	fortunate	■ ■ ■	I	J or	2	apjn	W C. G. H. 1810. C s.p
9790 fadsU Sal.	cube-flowered	■ ■ ■	I	Jde	1	apjl	Pa.P C. G. H. 1795. C s.p
9791 cubtea L.	larger	■ ■ ■	I	J or	1	apjl	P C. G. H. 1790. C s.p And. heaths 1
«,«, 2 mk J or	drawn out	■ ■ ■	I	J or	2	ap.s	P C. G. H. 1811. C s.p
9792 exnr6mpa Spr.	hairy-leaved	■ ■ ■	I	J or	2	my.au	P C. G. H. 1816. C s.p
9793 Masophylla Spr.	rising	■ ■ ■	I	Jde	1	myjn	W C. G. H. 1821. C s.p
9794 a88fingensL.	spiked	■ ■ ■	I	J or	1	f.o	W C. G. H. 1802. C s.p
9795 cana Andr. not L.	naked-flowered	■ ■ ■	I	J or	1	s.o	W C. G. H. 1800. C Bp And. heaths 1
2 spicata	hoary	■ ■ ■	I	J or	2	jl.au	D. Y C. G. H. 1783. C s.p Sin. ic. in. 3. SI
9796 nudifl&ra L.	reA-flowered	■ ■ ■	I	J cu	1	jna.u	W C. G. H. 1810. C s.p
9797 incana Wnl.	milk-flowered	■ ■ ■	I	J or	2	jns	R C. G. H. 1810. C Bp Bot cab. 901
2 ritbra	milk-colored	JH	or	2	jno	W C. G. H. 1818. C s.p	
9798 lacti flora Ld.	regerminating	■ ■ ■	I	J or	1	my.au	R C. G. H. 1791. C s.p
9799 lactiflor Bo. C.	white.	■ ■ ■	I	J el	1	myjn	W C. G. H. 1805. C s.p
9800 regermans W.	roughish	■ ■ ■	I	J el	1	jnjl	W C. G. H. 1800. C s.p
2 lls	small-bracted	■ ■ ■	I	J or	1	mrjl	R C. G. H. 1810. C s.p
9801 scabriuscula Lk.	crest-flowered	■ ■ ■	I	J or	li	myjn	Pa.P C. G. H. 1803. C s.p
9802 bracteofafis Lam.	protruding	■ ■ ■	I	J de	1	ap.my	W C. G. H. 1805. C s.p
9803 enstfflfra Sal.	ziz-zag	■ ■ ■	I	J or	ljapjl	W C. G. H. 1792. C s.p And. heaths 1	
9804 prottfidens Lk.	umbelled	■ ■ ■	I	J or	3	myjl	P Portugal 1782. C sp Bot cab. 1217
i*05 flexutis Andr.	reflexed-stmnd	*L-J or	S	Jns	R	C.G.H. 1799. C s.p	And. heaths 3
M divaricata WnU							
9806 umbellate L.							
'07 stamlnca Andr.							

9606 <i>caftia Sal.</i>	grey	iLJ or	8	mr.Jl R	C. G. IT.	1800.	C s.p		
9809 <i>latifolia Andr.</i>	broad-leaved	iLJ or	2	my.au R	C. G. H.	1800.	C s.p	And. heaths f	
9810 <i>carnica L.</i>	<i>ficah-coloured</i>	iLJ or	1	ja.np Pa.P	Germany	1763.	L s.p	Bot. mag. 11	
2 <i>herbacea Wnl.</i>	herbaceous	iLJ or	4	mr.my P	Portugal	1648.	C s.p	Bot. mag. 471	
9811 <i>mediterranea</i>	Mediterranean	iLJ or	2	ap.jn R	Sicily	1819.	C s.p		
9812 <i>sicula Cus.</i>	Sicilian	iLJ or	1	f.au R	C.G.H.	1810.	C s.p	Bot. cab. 843	
9813 <i>arbùscula Lod.</i>	little tree	iHJ>r	1	ll.au R	Cornwall	hea.	L s.p	Eng. bot. 3	
9814 <i>varans L.</i>	wandering	*iLJ or	1	ll.au W	Europe	—	L s.p		
2 <i>alba</i>	white-flowering	*iLJ or	1	jlau R	C. G. H.	1805.	C s.p		
3 <i>tenella</i>	delicate	*iLJ or	1	jlau Pk	C. G. H.	1800.	C s.p	Bot. cab. 103	
9815 <i>longipedunculata X.</i>	long-stalked	iLJ or	1	jlau P	C. G. H.	1800.	C s.p	And. heaths 4	
9816 <i>magnifica Andr.</i>	magnificent	iLJ or	2	av.in P	C. G. H.	1800.	C s.p	And. heaths 4	
9817 <i>Sainsburyana Andr.</i>	Salisbury's	iLJ or	2	jns. P	C. G. H.	1800.	C s.p	And. heaths 4	
9818 <i>Comptoniana Andr.</i>	Compton's	iLJ or	2	my.jl P	C. G. H.	1802.	C s.p	And. heaths ic.	
9819 <i>ciliaris L.</i>	ciliated	iLJ or	1	jlau P	Cornwall	hea.	C s.p		
9H20 <i>recurvata Bedf.</i>	curcurvate	iLJ or	i	apjh P	C. G. H.	1810.	C s.p	Bot. cab. 1093	
9821 <i>pilosa Lod.</i>	pilose	iLJ or	1	jlau G	C. G. H.	1800.	C s.p	Bot. cab. 606	
9822 <i>A'phanes Spr.</i>	Parsley-piert	iLJ or	2	mr.jn P	C. G. H.	1820.	C s.p		
9823 <i>arctata Lo. C.</i>	narrow	iLJ or	2	aps P	C. G. II.	1820.	C s.p		
5824 <i>Siberia W.</i>	whitish	mLJ or	4	mr.au W	C. G. H.	1789.	C s.p	Bot. mag. 440	
1825 <i>prop6ndens Andr.</i>	forward-hanging*	iLJ or	It	jlau P	C. G. H.	1800.	C fl.p	And. heaths 2	
1826 <i>intertexa Lod.</i>	interwoven	iLJ or	U	jl.nj Y	C. G. H.	JH18.	C s.p	Bot. cab. 1034	
m 27 <i>jubata Lod.</i>	maned	iLJ or	2	jn.o P	C. G. H.	1816.	C s.p		
9828 <i>stylaris Spr.</i>	/fliff-styled	uLJ or	2	my.au ...	C. G. H.	1812.	C s.p		
9829 <i>acutangula Nois.</i>	acute-angled	KU nit	2	mr.ap W	C. O. H.	1860.	C s.p		
9830 <i>pyramidalis B. M.</i>	pyramidal	iLJ or	1	fm.y Pk	C. O. H.	17h7.	C s.p	Bot. mag 36G	
9831 <i>denexa Hort.</i>	deflexed	iLJ or	2	jl W	C. G. H.	1810.	C s.p		
9832 <i>pyramidaliformis Wnl.</i>	pyramid-form	mLJ or	2	Hd P	C. G. H.	1K18.	C s.p		
9833 <i>halictraeflora llort.</i>	Thalictraeflora	iLJ or	II	mr.my Y	C. G. H.	—	C s.p		
9834 <i>nola-flora Sal.</i>	bell-flowered	iLJ or	n	W	C. G. II.	1818.	C e.p		
gloinera llort.									
9835 <i>bryanthia Thun.</i>	very flowery	*LJ or	1	ji.Jl W	C. G. H.	1812.	C s.p		
9836 <i>icenuflora Andr.</i>	Echium-flwd	*iLJ el	S	C. G. H.	17fw.	C s.p	And. heaths 3		
2 sup&rb	superb	*LJ or	i	fin S	C. G. H.	1800.	C s.p		
3 purpurea	purple	sLJ or	1	fjn S	C. G. H.	1800.	C s.p		
4 coccinea	scarlet	illJ or	1	fjn S	C. G. H.	1812.	C fl.p		
9837 <i>argentiflora Andr.</i>	silver-flowered	HIJ or	1	apjl W	C. c. H.	1800.	C s.p	And. heaths ic.	
9838 <i>filamcrosis Andr.</i>	filamcnos	HIJ or	2	year p	C. G. H.	1800.	C s.p	Bot. reg. 6	
9339 <i>tubiflcula Bed.</i>	small-tubed	iLJ or	2	ja.8 R	C. G. H.	1800.	C s.p	And. heaths 2	
pubescens minima Andr.									
9840 <i>pulchilla Thun.</i>	pretty	iLJ or	R	C. G. H.	1792.	C s.p	Thun. er. 24.4		
9841 <i>bella Spr.</i>	beautiful	iLJ or	1	ja.o R	C. G. H.	1800.	C s.p	Bot.oab.307	
pulchdla llort. not Thun.									
9842 <i>visc*riat.</i>	clammy-fluid	iLJ d	2	mr.jl R	C. G. H.	1774.	C s.p	Ic.h.k.l	
9843 <i>flexicafilis H. K.</i>	crook-stalked	iLJ or	I	jammy	C. G. H.	1800.	C s.p	And. heaths 2	
glandulosa Andr.									
9844 <i>tenella Andr.</i>	delicate	illJ or	my.au	P	CG.H.	1791.	C s.p	And. heaths 2	
9845 <i>alopecuroldes Wnl.</i>	Foxtail-like	iLJ or	1	au.d R	1810.	C s.p	Bot. cab. 874		
9846 <i>stirfurtaea Sal.</i>	scurfy	iLJ or	2	jnii F	France	1731.	C fl.p	And. heaths 1	
9847 <i>multiflora L.</i>	many-flowered	iLJ or	2	jn.n W	Europe	—	C fl.p	And. heaths 2	
2 alba	white	iLJ or	2	jn.au Y	C. G. H.	1789.	C <p	And. heaths 2	
9848 <i>depressa L.</i>	depressed	iLJ or		rn.y					
rufiflora Andr.									
9849 <i>adenophora Spr.</i>	gland-bearing	*uJ or	2	my.jl	CG.H.	1810.	C s.p		
9850 <i>nkna Sal.</i>	dwarf	m iJ el	fm.y.au Y	...	C. G. H.	1792.	C s.p		
9851 <i>ambigua Spr.</i>	ambiguous	tk iJ or	2	ap.jny P	C. G. H.	179a	C >p	And. heaths Ic.	
cylindrica Andr.									
9862 <i>pallistris Andr.</i>	marsh	il LJ or	1	my.o F	C. G. H.	1799.	C s.p	And. heaths 2	
9853 <i>formosa Thun.</i>	handsome	A uJ or	2	jns R	C. G. H.	1795.	C s.p	Thun. er. 82.3	
2 alba	vrhite-flowered	*iLJ or	2	jns W	C. G. H.	1795.	C s.p		
3 rrhra	red-flowered	il LJ or	2	jnfi R	C. G. H.	1795.	C fl.p		
9854 <i>florida Thun.</i>	florid	ill or	1	my.au R	C. G. H.	1803.	C fl.p	Thun. er. 64.6	
2 mosenkta	musk	il LJ or	1	mr.au R	C. G. H.	—	C s.p		
9855 <i>antheria Spr.</i>	many-Hovered	JtlJ or	2	apjl R	CG.H.	1800.	C s.p	Bot. cab. 234	
flrida llort.									
9856 <i>muconhita Andr.</i>	mucronated	il LJ or	1	rny.au P	C. G. H.	1800.	C s.p	And. heaths 3	
9857 <i>Solandriiflora Andr.</i>	Solndr's	il LJ el	1	mr.s Fk	C. G. H.	1800.	C s.p	And. heaths 2	
9858 <i>aedta Andr.</i>	noined-cupped	M LJ de		my.jl	C. G. H.	1799.	C s.p	And. heaths 2	
9859 <i>mpetroides Andr.</i>	Empetrum-like	il LJ or	1	my.au LF	C. G. H.	1788.	C s.p	And. heaths 2	
9860 <i>pulverulenta Lo. C.</i>	JMWderd	il LJ or	1	jn.au P	C. G. H.	1820.	C fl.p		
ytkil turrigera Sal.	turret-bearing	il LJ or	1	jn.s R	C. G. H.	1796.	C fl.p		
9862 <i>procumbens Hort.</i>	procumbent	A C or	t	jn.Jl ,P	CG.H.	1800.	C fl.p		
cupressina llort.									
9863 <i>Berguma L.</i>	Bergius's	iiLJ cu	1	ap.au P	C. G. H.	1787.	C fl.p	And. heaths 2	
quadriflora Andr.									
9864 <i>riibens Thun. not Andr.</i>	reddish	iiuJ or	1	ap.o P	C. G. H.	1810.	C s.p		
peduncularis Sal.									
9S65 <i>quadriflora Sal.</i>	four-flowered	• Ljor	1	mr.au ...	C. G. H.	1800.	C fl.p		
9866 <i>barpata Andr.</i>	bearded	illJ or	1	myau W	C. G. H.	1799.	C s.p	And. heaths 2	
2 major	larger	iLJ or	1	mr.au R	C. G. H.	1800.	C fl.p	And. heaths 2	
3 minor	smaller	*LJ or	1	mr.au R	C. G. H.	1800.	C fl.p	And. heaths 2	
9867 <i>retroflexa Wnl.</i>	retroflexed	iLJ or	1	jl.s W	C. G. H.	1787.	C s.p*	Wen. er. 8.7.1c.	
pule*lla Andr.									
9868 <i>equiseticilia Sal.</i>	articuläris Thun.	Equisetum-lvd	iLJ or	2	jl	C.G.H.	1800.	C s.p	
articuläris Hort. not B.M.									
9869 <i>hyminiflia Andr.</i>	Thyme-leaved	iLJ	my.au P	C. G. H.	1789.	C fl.p	And. heaths 2		
9870 <i>exserta Bedf.</i>	exserted	iLJ or	2	fn. P	C. G. H.	1810.	C s.p	And. heaths 4	
petitata Andr.									
9871 <i>fucata Wnl.</i>	painted	il LJ or	1	ap.my P	CG. H.	1819.	C s.p		
9872 <i>oxyccifolia Sal.</i>	Oxycoccus-lvd	JU iJ or	Jf.my R	...	CG. H.	1791.	C s.p		
9R73 <i>ctnus Wall.</i>	slendcMtN*	* iJ el	l" f o Pk	C	C. G. H.	1790.	C a.p		
<S74 <i>liaria Thun.</i>	hairy-leaved	M iJ or	2 apjn Rpk	CG. H.	1795.	C fl.p	Thun. er. 56		
9875 <i>strig'sa H. K.</i>	A iJ or	1 mir.ap Pa.lt	CG. H.	1775.	C s.p				
9876 <i>coufertiflora tVnl.</i>	crowded-flwd	il J or	M n y j r w	CG. H.	1818.	C s.p			
9877 <i>confertiflta Wnl.</i>	crowded-leaved	* uJ or	2 my.au ...	C.O. H.	—	C s.p			

- | | | | | | | | | |
|--------------------------------|--------------------|-----------|---------|----|----------|-------|-------|---------------|
| 9878 mollearis <i>Sal.</i> | <i>soft-leaved</i> | 4F L J de | 1 ap.o | R | C. G. H. | 1803. | C s.p | Schn. ic. IT |
| 9879 racemifera <i>Andr.</i> | raceme-bearing | ■ I J or | 1 ap.jn | R | C. G. H. | 1803. | C s.p | And. heaths 3 |
| 9880 pilulifera <i>L.</i> | pellet-bearing | 4F I ICU | 1 ap.my | R | C. G. H. | 1789. | C s.p | |
| 9881 catervosfolia <i>Sal.</i> | huddled-leaved | 4I \ I or | 1 ap.jn | R | C. G. H. | 179a | C s.p | |
| 9882 sicaefolia <i>Sal.</i> | dagger-leaved | 4I \ I or | 1 f.au | - | C.G.H. | -- | C s.p | |
| 9883 tardiflora <i>Sal.</i> | slow-flowering | 4I: I C U | 1J mr.8 | F | C.G.H. | 1790. | C s.p | Botmag.480 |
| ↑ pubescens <i>B. M.</i> | | | | | | | | |
| 9884 canescens <i>Wnl.</i> | canescent | ■ I J or | 2 mr.o | R | C G . ft | 1802. | C s.p | And.heath12 |
| ↑ pubescens minor <i>Andr.</i> | | | | | | | | |
| 9885 parvi flora <i>Sal</i> | small-flwd downy | 4I \ I pr | 1 mr.8 | Pk | C. G. H. | 1790. | C s.p | |
| 9886 exfuga <i>Sal</i> | small downy | 4i I J pr | 1 mr.8 | Pk | C. G. H. | 1790. | C s.p | |

VII. DIBBLE.—*Doubtful to what section they belong*

(Stephan FMiot, a North American botanist.) *Ericece.* 1—2.

JJ76. CHLO^RA L.	racemose	H	or	2 my.jl	W	Georgia	...	L.s.p.
^ "imperfornata L.	YELLOW-WOBT. (<i>Chloros</i> , green : dried flowers of <i>C. perfoluta</i> .)	Genthinca: 2.-3.						
"	perfoliate	O	or	1 jn.jl	Y	Britain	sh. sc. S. s.l	Eng. bot. 60
"	imperfoliate	O	or	1 jn.jl	Y	Italy	1833. S	S & I

W⁷⁷ MICHAU'X/i Herit. MICIIAUXIA. {A. Michaux, botanist to Louis XVI.) Campanulaceet. 2.
 ' ^-{Fnanuloides/fcr//. Campanula-Ik or 4 jn.au L.B Levant 1787. S r.J Bot. mag. 219
 ' ^-{F Fen. smooth-Btnd 5i or 3 jl.au L.B Persia 1820. S r.l Ven.cels81
 JEFFERSO^NZ/I Bart. JEFFERBONIA. (T. Jafffrson, President of United States.)
 phbllum dipbllum L. < * my ' W N. Amer. 1792. D L

• 1179. DODO*LE*V* L.	DODONAA.	(<i>R. Dodonaei</i> , a celebrated botanist)	Terebitdhhcece. 18.—22.
9953 viscosa L.	clammy	il IJum 6 jnjl G	S. Amer. 1690. C pl Cav. ic. 4. 327
9545 jamaicensis Dec.	Jamaica	D c u 5 jnjl G	Jauhaica 1810. C fl.p Cavic. 1327
viscosa <i>Cav. angustifolia Sum.</i>			
9955 bialata Kth.	two-winged	Q un 4	S. Amer. 1822. C co
9956 /aurifolia Sieb.	Laurel-leaved	Q cu 5	N. Holl. 1823. C s.p
9957 Bunnanatina Dec.	Burmann's	Q cu 5 jnjl	C pl Bur. zey. 23
9958 micrurarpa Dec.	small-fruited	cu 4 jnjl	C pl Lam. il 304.2
9959 jalicitbia Dec.	Willow-leaved	I Jcu 4 jnjl	C p.1
angustifolia Lam.			
9960 dioica Rox.	dioecious	I Q cu 4 jnjl G	E. Indies 1819. C p.1 Ru. am. 4.50
heterophylla Hort.			
9961 oblongifolia Lk.	oblong-leaved	I I un 4	G.P N. Holl. 1823. C co Bot. reg. 1051
9962 triquetra Andr.	three-sided	I I un 5	... N. S. W. 1790. C S.D Bot. rep. 230
9963 cuncta Sm.	vicAgc-It'aved	I cu 3	... N. Holl. 182a. C Pj Lin. tr. 11.19
9964 truncata Lod.		LJCU 3	... N. Holl. 1820. C p.1
9965 asplinifolia Rud.	Fern-leaved	I cu 3 jnjl	N. Holl. 1820. C p.1 Lin. tr. 11.20
9966 filiformis Lk.	filiform/twiwrf	I Jcu 3 M	... 1820. C p.1
9967 angustissima Hort.	narrow-leaved	I cu 3 jnjl	... 1823. C p.1
9968 «la;agnoids Rud.	Elaugnus-like	I cu 5 jnjl	& Domin. 1800. C p.1
9969? pinnata Sm.	pinnate-leaved	I cu 4 jnjl	N. Holl. 1824. C p.1
9970 itcabra Lo. C.	scarious	I I Jcu 4 jnjl	N. Holl. 1820. C p.1
9971 attenuata Cun. •	attenuate	I I Jcu 4 jnjl	N. Holl. 1820. C p.1
1180. LAWSONI*N/i* L.	LAWSONIA. (<i>J. Lawson</i> , M.D., author of a voyage to Carolina.) <i>Salichrice</i> . a—6.		
9972 inermi L.	unarmed	O cu 10	Egypt 1752. C s.p Rauw. ic. 60.7
9973 spinosa L.	prickly	I O c u 18	E. Indies 1759. C s.p R. mal. 1.40
9974 purpurea Lam.	purple	\$ • cu 12	E. Indies 1820. C s.p
* 1181. OSBECK/ff L.	OSDECKIA. (<i>P. Osbeck</i> , a Swedish c		Melastomifcete. 4.—
9975 cylindrica L.	Ceylon	« Q pr 2 jlau	1799. C s.p Bot. reg. 565
9976 chinensis L.	Chinese	tt-Opr	1818. C s.p Bot reg. 542
Mel&btoma osbeckidides Kcr			
9977 stellata Don	starred	«-[Z]pr 1 jn.au	Nepal 1820. C p.1 Bot. reg. 674
9978 neilensis Hook.	Nepal	I tt.(ZZ)pr U Jn	Nepal 1821. C p.1 Hook. ex. fl. 31
* 1182. AHEOCIA L.	RHEXIA. (<i>Rhexis</i> , a rupture;		Melastomiacea. 6.—8.
9979 mariana L.	Mariland	A f jn.au	N. Amer. 1759. D s.p Bot. cab. 366
2 rubella Ph.	reddish	i jn.au	N. Amer. 1823. D s.p Sw. fl. gar. 41
9980 angustifolia Lam.	narrow-leaved	I jn.au	N. Amer. 1812. D s.p
marianna y exfibida	Mx.		
9981 chit>sa Mx.	ciliolate	I Jn 1 jn.au	Carolina 1812. D p.1 Ph. am. 1.10
9982 virginiana L.	Virginian	f jn.au	N. Amer. 1759. D p.1 Bot mag. 968
9983 Hypericoides W.	Hypothecum-like	O or 1 jkau	Guiana 1820. S s.l Aub. gu. 1.161
9984 versicolor B. R.	changing-cld	«L • or f s	Brazil 1825. C s.p Bot. reg. 1066
• 1183. FINOTHERETRA L.	CEVOTHERA. (<i>Oinos</i> , wine, <i>thera</i> , a catching; acquired smell.) <i>Onagr&rue</i> . 55.—60.		
9985 biennis L.	biennial	CD or 4 Jns	N. Amer. 1629. S co Fl. dan. 446
9986 salicifolia Detf.	Willow-leaved	CD or 2 jn.au	1794. C co
9987 globularis H. C.	globular	CD or 3 jn.au	1824. C co
9988 gaurodes Horn.	Gaura-like	Q or 4 jl.au	1824. C co
9989 hpectabilis Bernh.	showy	Q or 2 jn.au	1810. C co
9990 albicans Lam.	whitish	Q or 8 jn.au	Mexico 1820. C co
9991 grandiflora H. K.	great-flowered	Q or 8 jn.au	Peru 1823. C co
suaveolens Pen.			N. Amer. 1778. C co Bot. mag. 2068
9992 parviflora L.	small-flowered	I Jn 4 jn.au	N. Amer. 1757. S co Mee. ic. 1.34
9993 muricata Mur.	rickto-stalked	I Jn 3 jl.au	N. Amer. 1789. S co Mur. 6.
9994 longiflora Jac.	long-flowered	I Jn 3 jl.s	B. Ayres 1776. S co Bot mag. 365
9995 odorata Jac.	sweet-scented	I ap.au 2 ap.au	& Amer. 1790. S co Bot reg. 147
undulata Dry.			
9996 nocturna Jac.	night^meMng	£ Q or 2 ap.au	Y C. G. H. 1790. S co Jac. ic. 3.455
strata Led.			
9997 villosa W.	villous	I O or 2 jl.au	Y C. G. H. 1791. S co
9998 dentate Cav.	toothed	I O or i jn.au	Peru 1818. S co Lind. coL 10
ChamissSs & Lk.			
9999 serrulata Nut.	serrulated	A pr 1 jn.	Y N. Amer. 1824. S p.1 Hook. ex. fl. 140
10000 cheiranthifolia Hort.	Cheiranth.-lvdtt.	I pr or Ift jn.au	Chile 1823. S.l Bot reg. 1040
10001 minima Ph.	least	i jn.au	N. Amer. 1825. co
10002 8crftina Hort.	atc-Jlowering	I Jn.aus	... 1820. co Sw.fl.gar.184
10003 cruciflata Nut.	cruciate-yfu>a	3 jn.au	N. Amer. 1824. co
10004 tetragntha Roth	four-appled	U jn.au	N. Amer. 1820. co
10015 corymbigra Lam.	corymbose	3 s	Mexico 1816. co Bot mag. 1974
10005 stricta Led.	strict	Ift jnjl	... 1822. co
10007 mtdia Lk.	intermediate	S J U u	N. Amer. 1823. P.1
10008 sinuata L.	Bcttop-leavcd	Jl.s	N. Amer. 1770. S co
10009 tcnuifolia Cav.	flnc-leaved	Jl.s	Peru 1824. s.l Mur. 5.3
10010 micr&ntha Horn.	small-flowered	myjl	California 1823. S co
hirta Lk.			
10011 mollissima L.	softest wave-lvd	Q or	Y B. Ayres 1732. S co Schk. han. 1.10 ⁵
10012 pubescens* W.	pubescent	Q or	W S. Amer. 1825. S co
10013 pinnatifida Nut.	pinnatifid	I Jn.au	W N. Amer. 1816. S co
10014 purpurifca Cur.	purple-flowered	I Jn.au	P N. Amer. 1794. S co Bot mag. 352
10015 rtoeo-filia Bernh.	red and white	my.au	P Nepal 1827. S co
10016 Roianitzuvi Led.	Romanzow's	my	P N. Amer. 1817. S co Bot. reg. 562
pleasing red		R.w	P N. Amer. 1825. S co
10017 amce^na Hort.	trailing	I Jn.au	P Florida 1824. S co
10018 humiflora Nut.	delicate	I Jn.au	P Chile 1822. S co
10019 tcnilla Cav.	white-stalked	ap.au	W N. Amer. 1811. S co Bot mag. 24x4
10020 albicaulis Fra.	shrubby	I my.au	D.Y N. Amer. 1737. D s.p Bot mag. <03>
10021 fruticosa L.	ambiguoust	I Jn.au	1 jn.au Y N. Amer. 1818. D co
10022 ambigua Nut.	hoary	I Jn.au	2 jn.au Y N. Amer. 1820. S co
10023 incknia Bart.	hybrid	I Jn.au	1 jn.au Y N. Amer. 1813. D co
10024 hybrida Mx.	dwarf	imy.s	D.Y N. Amer. 1757. D p.1 Bot. mag. S55
10025 ptimila L.	small	I Jn.au	Y N. Amer. 1817. S co
10026 pusilla Mx.			
10027 Uncaria Mx.			

JJ028 macrocarpa Ph.	long-fruited	1 jn.jl	N. Amer. 1811. R s.p	Sw.fl.gar.5
10029 missour'nsis B. M.	Missouri	1 jn.au	N. Amer. 1818. R af	Bot mag. 1592
alata Nut.				
10030 glauca Mr.	glaucous	2 my.o	N. Amer. 1812. D s.p	Bot. mag. 1606
10031 Frasen' Ph.	Eraser's	11 my.o	N. Amer. 1811. D s.p	Bot. mar. 1674
100J2 triloba Nut.	three-lobed	1 my.s	N. Amer. 1822. S co	* ><
rhizocarpa Spr.				
■ mrt*egH.K.	rosy-flowered	1 myau	Peru 178a S pl	Bot mag. 347
10034 tetraptera Cav.	four-winged	1 jnau	Mexico 1796. S s1	Bot mag. 468
10035 clavita Kth.	clubbed	1 jl.au	Mexico 1827. S co	
10036 nervosa /torn.	nervose	2 jl.au		1827. S co
1000 capitoza 2? Af.	turfy	1 Jnjl	N. Amer. 1811. R p.l	Bot mag. 1593
10038 acanthus Cav.	stemless	1 my.s	Chile 1821. S co	Bot. reg. 7s3
10039 scippiosa Nut.	showy	1 mr.s	N. Amer. 1821. S co	Hook. ex. fl. 80
10040 twiggy	Ujn	P.w	Peru 1823. S co	FL per. 315
* 1184. GAU'RA L.	GAIIRA.	(Gauros, superb; flowers.)	Onagrdrie. 6.-7.	
10041 biennis L.	biennial	CD or 5 au.o	R.w N. Amer. 17h2 S pl	Bot mag. 389
10042 mutabilis Cav.	changeable	H]pr 2 jl.au	Y N. Amer. 1795. S s1	Bot mag. 388
10043 trnotherelfbra Zuc.	(Enothera-flwd	Q> or 11 jl.au	P S. Amer. 1816. S s1	
10044 coccinea Fra.	scarlet	: A r lau.o	S Louisiana 1811. S s1	
10015 fruticosa Jac.	shrubby	“ pr 3 ...	R.w S. Amer. 1815. S s1	Jac. ic. 3. 457
10046 tripetala Cav.	three-petaled	O cu 11 au	Pk Mexico 1804. S s1	Cav. ic. 4.396.1
1185. CLA'RK/ii Ph.	CLARKIA. (Capt. Clark, accomp. Capt Lewis to Rocky Mountains.)	Onaghrke. 1.		
10047 pulchella Ph.	pretty	O P' Ift Jn	P N. Amer. 1826. S p.l	Bot. reg. 1100
* 1186. EPILECTBIUM L.	WILLOW HERB. (Epi, upon, lobos, a pod; flower.)	Onagrdrie. 21.—25.		
10048 angustifolium L.	narrow-leaved	3t A or	Britain mea. D cc	Eng. bot 1947
2 album	white-flowered	5 A or	4 jl.au W Britain gard. D co	
10049 spicatum iMtn.	spiked	5 A or	4 jl.au P N. Amer. D co	
KJ050 angustissimum/T.	JC narrowest-/^ A or	2 jl.au P Alp. Eur. 1775. D ms	Bot mag. 76	
10051 Dodon<>i AIL	Dodocns's	“ t A or	lljl.au P France 1700. D co	
HAlleri Retz.				
10052 rosmarinifolium Ph.	Rosemary-lvd	k A or	2 jl.au P N. Amer. 1800. D co	
squamiflum Nut.				
10053 latiflbum L.	broad-leaved	^ A or	2 jl.au R N. Europe 1779. D co	Par. Ion. 58
10054 hir8ixtumL.haiTy,OT CodlinSfcCream^	A or	4 jl.au P Britain watpl. D co	Eng. bot 838	
1X55 tomenfusum Fen.	tomentose	^ A or	3 jlau P Asia 1818. D co	Ven. eels 90
10056 parvifl&rum E.B.	small-flowered	3t A pr	3 jlau P Britain watpl D co	Eng. bot 795
m61le Lam.				
10057 villsum Thun.	villous	I A Jor	2 jl.au P C. G. H. 1799. O co	
10058 mont'mun. Z.	mountain	A w	2 jnjl P Britain woods. D co	Eng. bot 1177
10059 strictum MM.	strict	A or	“ lau P Pennsylv. 1817. D co	
JCHO rfeum ScArrfi.	rose-colored.	A w	“ 1 Pk England mar. D ms	Eng. bot 693
100K1 alsinifolium Vil.	Chick weed-lvd	A w	“ 1 Pk Britain sc. al. D ms	Eng. bot 2000
10082 tetragrnum L.	square-stalked	A w	“ 1 Pk Britain mar. D ms	Eng. bot 1948
10063 coloratum 3/A.	colored	A or	“ P N. Amer. 1805. D lp	
100G4 al/itstre Schmidt	alpine	“ 3 injl	P Switzerl. 1820. S lp	
100E4 dahitricum FM.	Daurian	“ p	“ W Dauria 1822. S co	
100G5 jwifistre L.	marsh	A w	“ jl P Britain mar. D co	Eng. bot 346
JNXJ7 obscfirum Schr.	obscure	A or	“ jl.au P Europe 1815. S co	
J0066 lanceolatum 5e*.	lance-leaved	A or	“ P Italy 1810. D co	
10069 alplnum L.	alpine	A w	“ Jlau R Britain aLriv. D sj	Eng. bot 2001
TnJIS ⁷ GELA Lou	GELA.	(Gelao, to shine; leaves?)	(Marine*. 1.	
10070 lanceoUta	lanceolate	*1 Jor	4 ... Y Cochinch. 1820. P 8.p	
Silas lanceolata Spr.				
JSS2 ¹ cocinea L.	FUCHSIA. (Z. Fuchs, a celebrated German botanist)	Onogrhrke. 13.—19.		
10072 gracilis Lindl.	scarlet	“ lor	6 my.o S.p Chile 1788. C p.l Bot mag. 97	
decussata B. M.	slender	“ tLjor	8 my.o S.p Chile 1823. C p.l, Bot reg. 847	
2 mul <i>ifl</i> ra	many-flowered	“ l-Jel	6 my.o S.p Chile 1824. P Bot reg. 1052	
10073 tenella Hort.	delicate	“ l-ei	8 my.o S.p Chile 1824.	
10074 virgita Swt.	twiggy	“ h or	4 my.o S.p Mexico 1825.	
10075 cónica B. B.	conic	“ jn.o	4 my.o S.p Chile 1825.	Bot reg. 1062
10076 macrostfeinon Fl. per.	long-stamened*	I Jor	3 JLs S.p Chile 1823.	Bot cab. 1062
10077 venusta Kth.	beautiful	“ l-Jel	6 o P Mexico 1825.	
[W78] ycioides Andr.	Boxthorn-like	“ l Jor	4 ap.o R Chile 1796.	Bot. mag. 1024
[Wfr parviflora B. R.	small-flowered	“ i Jor	4 my.o R Mexico 1824.	Bot. reg. 1048
Jij80 arborescens Moc.	arborescent	J i or	16 o Pk Mexico 1824.	Bot. reg. 943
JU81 sk <i>mr</i> cr <i>ss</i> excortidta L.	barked	“ ill-lor	S jn.o G.P N. Zeal. 1824.	Bot. reg. 857
“ “ r>da Swt.	hybrid	“ ll J or	3 my.o S hybrid 1825.	
a) etala R. % P.	apeLulous	II J or	10 s.o P Chile 1825. P FL per. 3.322	
IrJi?* CYMINO'SMA Gae.	CYMINOSMA. (Kyminon, cumin seed, o\$me, smell)	But&clear. 2.-4.		
XU084 pedunculMa Dec.	peduncled	m • cu 6 fd	G E. Indies 1800. C It.l	
inno* ^{3m} bolffera pedunculata Lou.				
*U085 odorata Dec.	sweet-scented		China 1818. C Itl	
Jambolifera odorata Lou.				
JL ² L V P ² RIUM Jac.	ELAPHBIUM f	(Elaphros, contemptible; wood.)	Rutdceaf. I.—J.	
glabrum Jrtc.	“ moth	“ 20 ... W Carthag. 1818. C p.l Jac.am.71.4		
Faglra Elaphrium IV.				
-1191. MELFCOPE Font. MBLICOPB. (Mell honey, hope, incision; nectary of notched glands.) Ruthcae. 1.				
'0087 ternata Forst.	ternate	• t11 or	N. Zeal. 1222. C p.l Lam.il. 294	
*J192. STADMA'NNLI Lam.	STADMANNIA. (M. Stadmann, a botanical traveller.) Sapindacece. 1.—ft,			
'0088 australis R.Br.	southern	I L J O F 30 . . .	N. Holl. 1823. C alp	
JL ² OXYCO'CCUS Pen.	CRANBERRY.	(Oxy\$, acid, kokkos, fruit)	fnee^.	4.
Palustris Ph.	marsh	ft*	fir imy.jn Pk Britain tur.bo. L p	Eng. bot 319
JacciniumOxyceducus X.				

- 10090 *macrofirpifl Pers.* long-fruited JU fr f my.jn Fk N. Araer. 17G0. L bog Den. br. 122
/acclinium macrocarpum H. K.
- 10091 *erythrocarpus Pers.* red-fruited n. fr 2 my.jn Pk N. Amer. 1806. I. bog Den. br. 31
ericetus Ph.
- 10092 *hispidulfiflvr**. rather hiRpid JU fr i ap.my W Huds. Bay 1815. L bog Ph. am. 13
Gaultheria <epyllumia PA. vaccinium hispidulum L. ^rbatus tilifurmis Lam.
- * 1194. *FACCI'NIUM L.* WHORTLK BERRY. (Perhaps from *bacca*, a berry.) *Erictce.* 43.—57.
- 10093 *jfrctOKtaphylos L.* Bear's Grape Y or 10 npju W Caucasus 1800. L l.p
- 10091 madeirens Lk. Madeira - fr 6 jn.au ILlo Madeira 1777. L p And. rep. 30
yfrctostaphyloa Andr. padifilum Sm.
- 10095 diff. asum H. K. diffuse \$ or 20 my.jl Pk Carolina 1785 L p
- arbbreum Mx.
- 10096 dumtoum Andr. bushy II or 3 my.jn W N. Amer. 1774. L p Bot mag. 1106
hirtaium H. K. frondosum Mx.
- 2 hhmile 5^{in!}?5. 5 or 2 my.in W N. Amer. 1774. L p.l Den. br. 32
- 10097 venustum A. K. beautiful A or 3 my.jn W N. Amer. 1775. L p
- 10098 frondosum L. frondose it or 3 my.jn W.o N. Amer. 1761. L p Bot. rep. 140
glaucum Mx.
- 10099 pallidum //, JE pale A or 2 my.jn W N. Amer. 1772. L p
- 10100 clevatum /o. C. elevated A or 3 my.jn W N. Amer. 1772. L p.l
- 10101 stamineum L. long-stained A or 2 my.jii W N. Amer. 1772. L p.l Pluk.al.339.3
- 10102 mucronatum £. mucronate A or 4 my.jn W N. Amer. 1772. L p.l
- 10103 Album Z. white-flowered A or 2 my.jn W N. Amer. 1772. L p Bot rep. 263
stamineum ^/ufr.
- 10104 lignistrinum /,, Privet-leaved A or 3 my.jn P N. Amer. ... L p
- 10105 rvsnosum H. K. resinous A or 4 my.jn P.o N. Amer. 1772. L p Wang. 30.69
- 10106 elevatd A or 3 my.jn Y.u Canada 1772. L p
- 10107 1 videscens viridescent-JfuM/A 3 my.jn Pk N. Amer. 1772. L p
- 2 rubescens rubrscnt-tud A or 3 my.jn Br N. Amer. 1804. L p
- 10108 parviflorum Andr. small-flowered A or 7 my. W N. Amer. 1806. L p Bot. rep. 123
corymbTisum L.
- 10109 corymbisphrum Mx. corymbose A or 3 my.jn Pk N. Amer. 1772. L p
- 10110 disomphrum var. Mx. disomphrum var. Mx. A or 6 my.jn W N. Amer. 1765. L p Bot. rep. 138
- 10111 virgatum H. K. twiggy A or 3 ap.my Pk N. Amer. 1767. L p
- 2 angustifolium ?Pal. narrow-leaved A or 3 ap.my W N. Amer. 1767. L p.l Den. br. 34
- 10112 marianum Wat. Maryland A or 2 my.jn W N. Amer. 1812. L p.l Den. br. 124
- 10113 grandiflrum Wat. great-flowered A or 2 jl.au W N. Amer. 1812. L p.l Den. br. 125. A
- 10114 elongatum Wat. elongated A or 2 jl.au W N. Amer. 1812. L p.l Den. br. 125. B
- 10115 minutiflrum Wat. sm.ll-flowered A or 2 jl.au W N. Amer. 1812. L p.l Den. br. 125. C
- 10116 glabrum Wat. smooth A or 2 jl.au Pk N. Amer. 1812. L p.l Dcn.br. 125. D
- 10117 galézansMx. Gne-leaved A or 2 my.jn W N. Amer. 1806. L p
- 10118 formosum Andr. darkened A or 2 my.jn Pk N. Amer. 1770. L p Bot.reg.302
Tuscatum/A".
- 2 angustifolium narrow-leaved A or 2 my.jn Pk N. Amer. ... L p
- 10119 tcniflum H. K. delicate A fr 1| my.jn Pk N. Amer. 1772. L p
- pennsylvanicum Mx. pennsylvanicum Mx.
- 10120 MytililiH L. Myrtle Bilberry* fr If ap.jn R Britain hea. L p Eng.bot.456
- 2 fructu albo white-fruited A fr 1J ap.jn G Britain moors. L p
- 10121 uliginosum L. bog Blealtei ry A fr 2 ap.my Pk Britain ... L p Eng.bot.581
- 10122 angUtitohum //, K nar.-lvd Blueit's A or 2 ap.my Pk N. Amer. 1776. L p
- myrtillodes Mx.
- 10123 whCinum Lod. Willow-faitm! A or 1jmyjn W N. Amer. ... L p
- 10124 ca?pitum Mx. turfy JH or 1 myjn W Huds. Bay 1823. L p
- 10125 fuis ldff'a L. Cowberry tt. \$ ap.jn Pk Britain Skp Eng.bot.598
- 2 major larger tt. p* f ap.jn Pk N. Amer. ..^ Skp
- 3 maxima largest tt. f ap.jn Pk N. Amer. ... Skp
- 10126 haUertftMun Lo. c7l1alfcria-lvd A or 14 myjn W N. Amer. ... L p
- 10127 myrtiflum Mx. Myrtle-leaved JU pr 1 myjl W Carolina 1812. L p
- 10128 crasbifilum Andr. thick-leaved tt. pr 1 jn.jl Pk Carolina 1787. L p Bot mag. 1152
- 10129 nitidum PA. meridional . or 2 mrjn W.G Jamaica 1778. L p
- 2 decumbens decumbent JU pr 1 myjn Pk Caracas 1825. C Cp
- 10130 lauertfntes Mx. lanceolate tt. pr 1 myjn Pk Carolina 1794. L p Bot. reg. 480
- 2 lanceolktus obtuse tt. pr 1 gmyjn Pk Florida ... L p Bot mag. 1550
- 3 obtusus obtuse ft pr 1* myjn Pk Carolina ... L p
- 10131 prunifolium Lo. C. Plum-leaved A or li myjn Pk N. Amer. ... L l.p
- 10132 Auxifdlum Andr. Box-leaved JU cu 1 myjn W N. Amer. 1794. L p Bot. mag. 923
- 10133 ovateni RA. » ovate * fr 2 my Pk N. Amer. 1826. L p.l
- * 1195. JI/E^TE'CLYON L. MEMECLTON. (Greek name of the fruit of the arbutus.) *Melasnncaj* 2—6.
- 10134 capitellktum L. small-headed : or 10 jl E. Indies 1796. L p.l Bur. .e y ^
- 10135 edule Box. edible J : ft 20 ... P.c Ceylon 1820. C p.l
1196. LAGE'TTA J; LAGETTA. (Lagefto, its name in Jamaica.) Thymetafar. 1.
- 10136 lintcaria J. linen * D c u 6 year W Jamaica 1793. C l.p Lam. il. 2B
- Daphne Lagitia Swz.
1197. DA'PHNE L. DAPHNE. (Daio, to burn, phone, noise, crackles when burning.) TAymeleaf. 22.-33.
- 10137 Mexercum L. Mezcercon A m 4 f.ap. Pk England woods. C p 1 Eng. bot 1381
- 1 redbrum red-flowered A or 4 f.ap. Pk England woods. C p 1
- 2 album whxte-Jiowered A or 4 f.ap. W ... C p.l
- 3 autunnMe autumnal A or 4 au,o K Europe ... C s.l
- 10138 Laureola L. Spurge Laurel ■ or 6 ja.mr G Britain woods. S.s.l Enp. bot. 119
- 10139 pontical L. Politic * ■ or 4 ap.my G Y Pontus 175<. C s.l Bot. mag. 1282
- 2 rtbra Hort. red A | or 4 ap.my R hybrid 1827. C s.l
- 10140 rinifMia5u. Tinus-leaved * | or 6 ... Jamaica 1773. C l.p
- 10141 Gnidium J. Gnidium tt or 2 jn.au W Spain 1597. C s.l Bot. rah. 150
- 10142 od^ra Thun. sweet-scented A i_J or 3 mr.d P.w China 1771. C r.m Bot. mag. 1587
- 2 variegau variegated A i_J or 4 ap.my W Japan 1800. C l.p
- 10143 indica L. Indian . i_I or 4 myjl W China 1800. C p.l
- 10144 papyracea Wai. paper * uj or 4 mr.au W Nepal 1824. C s.l As.res.13.315.ift
- cannabina Wat.

CLASS VIII. ORDER 2.

OCTA'NDRIA DIGY'NIA.

1805. GALETNW W. GALEMIA. (*C. Galenus*, a celeb, physician of Pergamus.) *Chenopbdeat*. 1. —3.
 10212 africana W. African *n. i_* cu 2 jn.au W C. G. H. 1752. C p.l Lam. il 314
1206. APHANA'NTHE Lk. APIANANTHE. (*Aphanes*, obscure, *anthos*, flower.) *AmaranihUceoj*. 1.
 10213 celosioides Lk. Cockscomb-lik IL U) W.G Brazil 1813. C p.l
1307. WEINMA'NN;/! I* WEINMANNIA. W. Weinmann, a German botanist) *Cunonihceaj*. 5.—IS.
 10214 glabra L. smooth • Dor 6 my.jn W Jamaica 1815. C r.m Lam. 11.313.1
 pinnata L.
 10215 hirta Stvz. hairy *m* lor 6 my.jn W Jamaica 1820. C r.m
 10216 ovata Can. ovate-leaved *Ml* or 6 my.jn W Peru 1824. C r.m Cav.ic.6.566
 10217 elliptica Kth. elliptic >*l* or 4 my.jn W & Araer. 1824. C r.m
 10218 trifoliata L.fil. three-leaved *inor* 4 my.jn W C. G. H. 1820. C r.m Lam. il. 313.2
1208. MCEHR'NGJM L. MQSIIKINGIA. (P. H. G. Mahring, a celebrated physician.) *Caryophylleee*. 2. - 3.
 10219 muscosa L. mossy 3 A cu 1 jn.jl L.P S. Europe 1775. D >l Schk. han. 1.10!
 10&0 wdifilia W. Sedum-leaved 1 A cu 1 jn.jl W.s S. Europe 182a D 6.1 W. b. in. 3.23
1209. AREMCTNIA Dec. AREMONIA. (Change from *Agrimonia*.) *Rosace**. 1.
 10221 agrimonoides Dec. Agrimony-like A or 1 jLau Y Italy 1739. D co Col.ec. 1.144
 Agnmonia agrimonoides L.

CLASS VIII. ORDER 3.

OCTA'NDRIA TRIGY'NIA.

- * 1210. POLY'GONUM L. POLTGOMJM. (*Polys*, many, *gonu*, a joint; stems.) *Polygbnea*? 68. —102.
 10222 ramosissimum Mx. branchiest my.jl N. Amer. 1810. S co
 10223 gracile It. Br. slender jn.au N. Holl. 1822. D m.s
 10224 amphibium L. amphibious jn.au Britain dit. D 81 Eng.bot.436
 2 hirshtum
 10225 coccineum MhL scarlet jn.au N. Amer. 1819. D
 10226 niloticum Leh. Nile jl.au Africa 1824. D m.s
 10227 ocreatum L. booted in.s Siberia 1780. D s.l Gm.si.3.8
 10228 plebeium R. Br. plebeian jl N. Holl. 1822. S m.s
 10229 virginianum Bieb. Virginian 3 au.s N. Amer. 1640. D 81 Par. th. 857.6
 10230 nmt Pers. *mild Water Peppers** or 1 jn.au N. Amer. 1800. S m.s
 10231 salsugineum Bieb. briny Q pr 1 jn.au Caucasus 1817.
 10332 Persicaria L. Persicaria w 1 j.l.au Britain dit S V.l Eng. bot 756'
 10333 incanum Schmidt hoary un 2 j.l.au Germany 1804. S s.l Pet. brit 3.8
 10234 /apathifolium L. Dock-leaved w 1 jn.s England dungh. S 81 Eng. bot 1382
 10235 persicarioides Kth. Persicaria-like or 1 jn.au Mexico 1816. S co
 10236 senegalensis Mil. Senegal or * jl.au Guinea 1825. S m.s
 10237 Hydr. Apipcr L. Water Pepper Q w 1 j.l.a Britain wat pi. S s.l Eng. bot 989
 10238 glabrum Box. smooth or 1 jn.au E. Indies 1810. S co
 10239 m'inus Hud. smaller w 1 au.s England wat.co. S s.l Eng. bot. 1013
 10240 arifolium L. Arum-leaved O cu 6 my.o N. Amer. 1816. S s.l PL am. 398.3
 10241 australe Spr. southern cu 1 jl.au N. Holl. 1818. S m.s
 10242 barbitale L. bearded A cu 1 J^a China 1819. S
 10243 tinctorium Lou. dyer's Q dy 2 jl.au China 1776. C 8.1
 10244 decipiens R. Br. deceiving A or 2 jl.au N. Holl. 1822. S m.8
 10245 flonbundum Schle. bundle-flowered or 2 jl.au Siberia 1818. S co
 10246 orientale L. oriental or 2 jn.au E. Indies 1707. S co Bot mag. 213
 2 album
 10247 pennsylvanicum L. Pennsylvanian or 4 jLo N. Amer. 1800. D co
 10248 setosum Jac. bristly or 1 in.a1 AsiaMin. 1817. D co Jacob. 3.57
 10249 LaxmfinJ Lep. Laxmann's A or Dauria 1800. D co
 10250 horniarioides Del. Horniaria-like ^ cu Egypt 1827. D co
 10251 erectum L. upright un P.R. N. Amer. 1792. D 8.1
 10252 acetosum Bieb. sour un Crimea 1820. S co
 10253 maritimum L. sea-side A un S. Europe 1820. D s.1
 10254 glaucum Nut. glaucous A or N. Amer. ... D co
 10255 equisetiforme Sm. Equisetum-fmd3 O un S. Europe ... S co
 10256 BeUrdi AIL Bcellardi's O un S. Europe ... S co
 10257 littoriale Lk. sea-shore Jc A un S. Europe ... S co
 flagellaris Bert.
 10258 arenarium W. \$ K. sand my.au Hungary 1807. W.&K.67
 10259 Dryandri Spr. Dryander's E. Indies 1800.
 elegans Dry.
 10260 patulum Bieb. spreading H jn.s W Tauria 1820. S co
 10261 aviculare L. small birds' 3 ap.o G Britain rubble. S co Eng. bot 125
 10262 articulatum L. jointed 1 jn.au W N. Amer. 1817. S co
 10263 acclustum IV. en. sourish 2 jnjl Wo Siberia 1816. D 8.1
 10264 allgnum TV. en. Willow-like 4 my.au W.G Siberia 1816. D 8.1
 10265 divaricatum L. divaricating 3 jl.au W.G Siberia 1759. D co Gm. si. 3.11.1
 10266 diffusum Pall. diffuse jl.au W Siberia 1817. D co
 10267 undulatum Mur. vraye-leaved 3 jnjl W.G Siberia 1789. D co Gm. si. 3. 10
 10268 alpinum All. alpine 2 ray.au W Switzerl 1816. D s.1 Al.ped.68.1
 10269 elegans Ten. elegant 2 ap.u W.G Naples 1824. D co
 10270 chinense L. Chinese dy 6 jl.au W.o China 1795. D co Bur. in. 30.3 *
 10271 patens D. Don spreading 3 A or 1 jl.au ... Nepal 1823. S s.l
 10272 sagittatum L. arrow-leaved 1 O cu 6 jl.au W.G N. Amer. 1759. D co H. cl. 12

10273 cilinbde Mr.	hairy-jointed	4	j.l.i	Pk	Canada	1800.	S	co	
10274 adpressum <i>R. Br.</i>	compressed	2	my.au	R	N. Holl.	1822.	L	i.p	Lab. n. h. 1.127
10275 dumetorum <i>L.</i>	thicket	12	my.s	W	S. Europe	1803.	S	co	Fl. dan 759
10276 Convolvulus <i>L.</i>	Woodbine	3	my.s	W.a	Britain	corn fi.	S	si	Eng. bot 941
10277 tataricum <i>L.</i>	Tartarian	2	jl.au	W.pk	Siberia	1759.	S	si	Gm.si.3.13.1
10278 emarginatum <i>Roth</i>	notch, leaved	2	jlau	Pk	China	1796.	S	si	Bot. reg. 1065
10279 cymatum <i>Trev.</i>	cymose	2	jl.au	Pk	Nepal	1827.	S	co	
10280 Fagopyrum <i>L.</i>	Brk-Wheat	2	jl.au	Pk	England	corn fi.	S	bi	Eng. bot. 1041
	American climbing	4	jl.s	Pk	N. Amer.	1749.	D	co	Pluk. al. 177.7
10281 scandens <i>Pall.</i>	silky	ijl.au	W	Siberia	1810.	D	co	Gra. si. 3.9	
10282 sericeum <i>Huds.</i>	hirsutissim	1	my.s	W	N. Amer.	1817.	D	LP	
10283 Walt.	hairy	1	my.s	W.G	Britain	al. pas.	D	si	Eng. bot 669
10284 viviparum ^{/*}	viviparous	2	my.jl	Pk	Siberia	1807.	D	s.p	
10285 ellipticum <i>W.</i>	elliptic	1	my.s	Pk	Britain	me. pa.	D	co	Eng. bot 509
10286 Bistorta <i>L.</i>	Snakeweed	1	my.s	P	Nepal	1820.	D	co	
10287 petiolatum <i>D. Don</i>	petiolated	2	my.au	P	Nepal	1820.	D	co	
10288 macrophyllum <i>Hum. D.</i>	Don long-leaved	1^	my.au	P	Nepal	1820.	D	co	
10289 affine <i>D. Don</i>	kindred	1	my.au	R	Nepal	1822.	D	co	

1211. TRAGOPY^RUM Bieb. GOAT'S WHEAT. (*Trogos, agat, pyros, wheat.*) *Polygynece.* 3.—&
10290 lanceolatum Bieb. lanceolate tit or 2 jl.au Pkr Siberia 177a L.s.l Bot.reg.255
Balzomous fructus prosp. R. R.

10891	<i>Polygonum frutescens B. R. buxifolium Bieb.</i>	Box-leaved	A	or	lijl.au	Pk	Siberia	1800.	L	s.l	Bot mag.	1065
10298	<i>Polygonum crispulum B. M. cauc&sicuyn Hgfg. polygamum Spr.</i>	polygonous	fit	or	2 jLau	Pk	Carolina	1810.	L	p.l	Ven. eels	63

1212. COCCOCLOBA L.	SEASIDE GRAPH. ' (Kokkos, a berry, lobos, a lobe; fruit.)	Polypbtete. 18. — 23.
10293 uvifera L.	grape-bearing	W.a W. Indies 1690.
10294 pubescens L.	pubescent	W.a W. Indies 1690.
10295 orbiculafaris Lo. C.	round-leaved	W.a S. Amer. 1825.
10296 excoriata L.	barked	W.a W. Indies 1733.
10297 nivea Jac.	white-leaved	W.a Jamaica 1818.
10298 punctita L.	spotted	W.G W. Indies 173a.
10299 diversifolia Jac.	various-leaved	W.a St. Dom. 1818.
10300 /auritifolia Jac.	Laurel-leaved	W.G Caracas 1822.
10301 flave-sceng Jac.	flavescnt	C.r.m Jac. sc. 3. 267
10302 obtusifolia Jac.	obtuse-leaved	W.G St. Dom. 1820.
10303 microstachya W.	small-spiked	W.o Carthag. 1822.
parviflora Poir.		W.G W. Indies 1824.
10304 obovata Kth.	obovate-leaved	W.G & Amer. 1824.
10305 barbadensis Jac.	Barbadoes	W.G Barbados 1790.
10306 brasiliensis Spr.	Brazilian	W.G Brazil 1825.
10307 latifolia Lam. .	broad-leaved	W.G S. Amer. 1812.
10308 acuminata Kth.	acuminate-wi	W.o N. Grenada 1&0.
10309 tenuifolia L.	narrow-leaved	W.a Jamaica 1820.
10310 longifolia Lk.	long-leaved	W.G W. Indies 1810.
		C.r.m

1213. BRUNNI'CHM Gae. BBUNWICHIA. *If. Brunnich, a Danish naturalist.*) *Polygbnea. 1.*
 '0311 cirrhbsa Gat. tendrilled H—I^r 6 ... Pk Carolina 1787. Lp Gae.fr. 1.45.2

J0313	pinnata L.	vr\ngtx\leaved	H— or	f0	W.o	W. Indies	1752.	C	r.ra	Jac. ob. 3.62.12
1<<14	tetragona Aub.	quadrangular-rtwd	— or	20	W	Guiana	1825.	C	lp	
10315	Vespirilio Sun.	bat	i—Q or	20	W	S. Christo.	1823.	C	s.p	Ac. haf. 3.11.1
J0316	meliastiflia J.	Melia-leaved	A—CD or	12	W.G	Brazil	1819.	C	s.p	Hook. ex. fl. 110
10317	senegalensis B.J.	Senegal	LCD or	16	W	Guinea	1822.	C	r.m	
10318	Cupam's Kth.	Cupam's	— or	20	W	Trinidad	1818.	C	s.p	
10319	pscrmdrpa Rich,	round-fruited	A—CD or	26	W	Guiana	1824.	C	lp	
10,320	curassavica L.	Curassoa	< D or	18	W.G	S. Amer.	1739.	C	r.m	Jac. ob. & 61.8
10&21	barbadensis Jac.	Barbadoes	—	16	W.a	W. Indies	1786.	C	s.p	Jac. ob. 3.62. P
10,323	polyphylla Schunt.	many-leaved	—	20	W.o	W. Indies	1739.	C	s.p	Jac. ob. 3.61.10
10,323	cartnaginnsis Jac	Carthagena	—	16	W	Carthag.	1818.	C	s.p	Jacob. 3.61.6
10,324	caribae' Jac.	Caribsan	^— or	10	W.a	W. Indies	...	C	s.p	Jac. ob. 3.62.7
10,325	pubescens Kth.	pubescent	iD or	16	W	S. Amer.	1820.	C	sp	
10,326	hfswida Jac.	hispid	—	20	W	Trinidad	1825.	C	lp	
10,327	bipinnata Poir.	bipinnate	LQ or	20	W	Brazil	1816.	C	r.m	Jac. sc. 3.268
10,328	cauliflra Jac.	stem-flowering	H— or	18	W	Caraccas	1822.	C	sp	Jac. ic. 3.450

,1215. S E R I A L Plu.	SERIANA. (<i>Phtis</i> [Serjeant, a French friar and botanist])	<i>Sapindacea.</i>	5.—21.
10329 sinuata Schunu	sinuata-fairvfl fl	Hor 15	... W.a S. Amer. ... C co Jacob. 3.61.2
J0330 divarickta SCAtm.	divaricate fl	Qor 20	— W.G Jamaica 1824. C r.m Ac. haf. 3. 12.2
10331 caracastaa W.	Caraccas	4 CD OT 15	■■■■ Caraccas 1816. C co Jac.sc 199
\ 3.32 mexicana W.	Mexican	fl 1	■■■■ W.a Mexico 1823. C r.m Ac. haf. 3.11.3
10333 triterpnata W.	Mexican	I or 20	■■■■ W.a S. Amer. 1739. C r.m Jacob. 61.10

~~Urville triterne 77.~~ triterne *J.C. 3gr* ²⁰ URVILLEA. *CD. d' Urville*, a French navigator and botanist. *Sapindace**. 2.
Urvillea ^{or} ²⁰ W Caraccas 1824. C *
rusty ^{26 myjl} W Brazil 1823. C s.p. Bot. reg. 107

* 1217. CARDIOSPE'RUM L. HEART SEED. (*Kardia*, a heart, *spetna*, seed) *Saphdace**. 6.—L2.

10336 *Crataegus* *lanceolata* L. Winter Cherry 1 E 3 cu 4' - 10' 10' - 20' 10' - 20' 10' - 20' W.a India 1594
 10337 *Crataegus* *lanceolata* L. var. *flavescens* S. G. long-flowered 1 E 3 cu 4' - 10' 10' - 20' 10' - 20' 10' - 20' W.a Jamaica 1817

10535	<i>grandiflorum</i>	<i>Sun.</i>	large-flowered	»	• or 20	J.E.Lau	W	Jamaica	1817.	C.C.
10538	<i>coulteroides</i>	<i>Kth.</i>	Colutea-like	»	• or 10	J.I.Iau	W	Caracas	1818.	8.1

10339 *hirsutum* W. hairy t. or 15 Jl au W. Guinea 1822 L C e1
 10340 Corundum L. Indian heart ■■■■■ 4 Jl au W.a Brazil 1750. S co

10341 *pubticens* *Lag.* *subpubescent* 6 Jan S N. Spain 1823. S co

I: SAPI'NDUS L. SOAP BERET, (tow-soap, *Indus*, Indian; fruit; cleanses *vener.*) *Sapindacea.* 11. - & 20.

Saponaria L. Saponaria edged (t)e c 20 ... Wa W. Indies 1697. CC pd Com. h. 94
V. [ao r 20 ... W.o Carolina ... CC pd

Forsyth'ka > aor 20 W Granada ... C p.l. 00X6Q-251

**rtWfQ o r 25jLs
frutescent - g or 10 ..**

Aub-gUL 1,159

10318 /aurifMius Vahl	Laurcl-leaved		W.o	E. Indies	1820.	p.l R. mal. 4.
10349 lonjifMius Vahl	lonjj-leavcd		W.u	E. Indies	IHQ	co
10350 emnrginatus Vahl	emargmated		W.o	E. Indies	1822.	J.m
10352 Indicus Pair.	rusty Indian		W.o	E. Indies	1821.	Pi

CLASS VIII. ORDER 4.

OCTA'NDRIA TETRAGY'NIA.

1219. KALANCH&E Adan.	KALANCHOB.	(Chinese name.)	Semperv'ivea.	7.-8.
10353 lacimata Dec.	cut-leaved	n. Z3 cu	2 Jl.au	£. Indies 1781. Ls &l Dec. pl. 100
10354 crenfcta Dec.	crenated	tt. ZD cu	2 JLs	S. Leone 1793. Ls s.l Hot. mag. 1436
Verea crenata Andr.				
10355 acutiflora Haw.	acute-flowered	tt. ^] cu	2 jl.s	W £. Indies? 1806. Lss.1 Bot.rep.560
Verea acutiflora Andr.				
10356 egyptica Dec.	Egyptian	«L Z] C U	2 JLau	Y Egypt 1820. La i.l Dec. pl. 63
10357 spatula Dec.	spatulate	ajcu	2 jl.Lau	Y China 1820. Lss.1 Dec.pl. 64
10358 rotundiflia Haw.	round-leaved	«. ZD cu	2 jl.au	W C. G. H. 1820. Lss.1
Crassula rotundiflia Haw.				
10359 ceratophylla Haw.	horn-leaved	«. Z3 cu	2 jl.au	Y hina 1820. Lss.1

1220. BRYOPHY'LLUM Sal. BRYOPHYLLUM. (*Biyo*, to grow, *phyllon*. a leaf; leaf plant) *Semperiveo* of I.
10360 calycinum Sal. /ar/,
cupped «L Z2 cu 2 ap.jl G.p E. Indies 1800. Ls s.l Par. Ion. 3
Cotyledon calycinum Lar.

1221. PA^RIS L.	PARIS	{Par, equal; regularity of parts.)	Asphodelear.	3.-4.
10361 quadrifolia L.	four-leaved	& A cu £ my.jn G	Britain woods. D pl Eng. bot. 7	
10362 verticillata Bieb.	verticillate	5 A cu J ap.my G	Caucasus 1825. D veg	
10363 polyphylla Sm.	many-leaved	^ A cu ap.jn G	Nepal 1826. D veg	

1222. ADCKXA L.	ADOXA.	{A, privative, doxa, glory; not showy.)	Saxifrdgea?	1.
10364 Moschellina L.	Moschatellina	A A cu \ mr.my G.v	Britain woods. D s.p Eng. bot 453	
1223. £LATINE L.	WATER-WORT.	{Elate, afrm, Greek.)	Caryoph/lletg.	2..4.
10365 Hydr6piper L.	Water Pepper	* O cu i au W	S. Europe ... S s.l Vail. pa. 2.2	
10366 tripétala Sm.	thrc-pctaled	i O cu 1 au F	Britain mar.Ian. S s.l Eng. bot 955	
	triandra Schk.	hexāndraD^c. Hydrōpiper E.B.		

* 1221. HALORA'GIS Font. HALORAGIS. (*Hals*, the sea, *rhex*, berry of grapes.) *Halordgear.* 2.-5.
10367 Cerfdia H.K. tail-like tt.i Jcu 2 ap.s G.R N. Zeal. 1772. C s.p Jac.ial.69
10307 prostraU Font. prostrate J U L J C U ftmy.1 G.» N. Zeal 1820. C s.p

CLASS IX. ORDER 1.

ENNEA'NDRIA MONOGY'NIA.

1225. CINNAMCTMUM jIL Br.	CINNAMON.	{Kinamon, cinnamon, Arab.)	Lattirue.	11.—15.
10309 vērum Swt.	true	Q m 40 jns.	W E. Indies	1768. C s.p Bot mag. 1636
/atinu Cinnamōnum L.	Cässia B. M.			
10370 nitidum Hook.	shining Cinnamon^	O m 20 ... G	EL Indies	1823. C p.l Hook. ex. fl. 176
Zafrus nitida Rox.				
10371 Cassia D. Don	Cassia	t D m SO my.s	G.v Ceylon	1763. C s.p Bot rep. 51%
*Lafrus Cassia L.	Cinnamomum B. Rep.			
10672 J/alabatruin G. Don	Malabar Leaf	± or SO ... G.v	E. Indies	1805. C s.p R. mal. 5. S3
/aurus j/lalabatrum L.				
10375 dñlee G. Don	Camphor tree	£ • m 20 mr.jn G.w Japan	1727. C s.p Jac.sc.4.3.2	
L&lriu Cthnphora L.				
10374 glaucum G. Don	glaucous Camphor	• or 20 ... P.v Japan	1800. C Lp	
Xaurus glaeca Thun.				
10375 dñlee G. Don	sweet cinnamon^	• or 40 ... W. Y E. Indies	1820. C Lp	
/atirus dtlcs Rox.	BunndnniNecs			
10376 Beiolgta Swt.	Bciolgota	CD or 40 ... Y.G E. Indie* 1818. C Lp		
Zafirus Beiolgta Ham.				
10377 grâcile Swt.	slender	• or 20 ... Y.G E. Indies	1820. C Lp	
Xatirus grâciliis Hort.				
10378 montanum G. Don	mountain	D or 40 ... W.o W.Indies	1810. C Lp	
Lauras montana Swz.				
10379 Culidban Swt.	Culilaban	• or 20 ... G.T E. Indies	1823. C s.l Ru. am. 2.14	
ZaiSrus Culilaban L.				

* 1228. LAU'RUS L.	LAUREL.	(Blaur, green, Celt.)	LatirHue.	25.—70.
10380 spl\$ndens Hort.	shining	or 40 ... Y.o	E. Indies	1800. C l.p
Jl&2A »acea Sivz.	leather-leaved	or 50 ... W.p	W. Indies	1810. C l.p
— miir	Willow-leaved	CD or 20 ... Y.G	W. Indies	1826. C l.p
10384 Chlor6xylon L.	thyrs-flowcrcd	O or 30 ... Y.o	Madagasc.	1810. C l.p
10385 floribunda Swz.	Cogwood tree	tm 60 ... G.w	Jamaica	177a C s.p Br.jam.7
	bundlc-flowered	• or 40 ... Y.o	W. Indies	1800. C l.p

CLASS IX OADEB ?

ENNEANDRIA TRIGY'NIA

L.	HHITBARB. (<i>Rha</i> , the Wolga; first brought from its banks.)	<i>Polygbnea?</i>	14.—15.
anticum L.	Uhapontic	ft A m 4 my.jn W.o Asia	1573. R co Sab. rom. 1. 34
nts Fis.	wave-leaved	■ A cul 4 my.jn W.a China	1734. R co Am. ac. 3. 4
J. Mricuin Hort.	Siberian	■ A cul 8 my.jn W Siberia	1800. R.r.m
^ m IF. \$	windowed	■ A cul 6 myjn W Siberia	1800. R.r.m
astrMe D. Doh	hybrid	■ A cul 5 myjn W.a Asia	1778. R co Mur. 1
* & W.	southern	K A ■ 8 — ■ Nepal	1823. R.r.m
caspium W.	palmated	ft A m 5 ap.my W.a Bucharia	1763. R co I, f fa. 7.4
<*<\$<2 crtpum H. C.	compact	ft A m 3 myjn W Tartary	1758. R co Mil. it 2.218
H33 austrUcum H. C.	Caspian	■ A cul 6 myjn W Russia	1817. R.r.m
artSircium L.	curled	ft A cul 5 myjn W	1800. R.r.m
	Austrian	■ A cul 5 myjn W Austria	1800. R.r.m
	Tartarian	I A no 3 myjn W.o Tartary	1793. R co
	Ribes	■ A cul 2 myjn W.a Levant	1724. R co An. mu. 1. 49
183a PLEEA Mx.	PLERA.	(<i>Pleias</i> , the seven stars; flowers.)	
''36 tenuifolia My.	slender-leaved	I B I A cul ft my.s Br Carolina	Jwiceat. 1. 1824. D p.1 Bot mag. 1966

CLASS IX. ORDER 3.

ENNEA'NDRIA HEXAGY'NIA.

1234. JKTTOMUS L. FLOWKBINO RUSH, (*nous, an ox, tcmmo, to cut; cropped by oxen.*) *Bubmeee.* 9.—
 10437 umbellatus L. umbelled [^]A el 2 jn.jl Pk Britain dit D r.l Eng. bot. 651
 10438 latifdhus D. Don broad-leaved * A el 1 jn.jl W Nepal 1823. D r.l

CLASS X. ORDER 1.

DECA'NDRIA MONOGY'NIA.

1235. EDWA'RDS/4 Sal. EDWARDSSIA. (5. Edwards, a celeb, bot. draughtsman.) *Leg. Pap. Soph.* 5.—&
 10430 grandiflora Sal. large-flowered ^{Sc} or 12 my.jn Y N. Zeal. 1772. C s.p Bot. mag. 167
 10440 chrysopillya Sal. golden-leaved ^{Sc} or 12 my.jn Y N. Zeal. — C s.p Bot. reg. 738
 10441 nitida Dec. shining * Q or 8 — Y Bourbon 1820. C s.p
Sophbra *scricea* *Hit.*
 10442 microphylla Sal. small-leaved 4 I or 6 my.jn Y N. Zeal. 1772. E s.p Bot. mag. 14 IS
 10443 minima Lo. C. least A or 4 my.jn Y N. Zeal. 1818. E s.p

- 1236 MYROSPERMUM H. %B. MYROSPERMUM. (*Myron, balsam, sperma% seed.*) *Leg. Pap. Soph.* 1.— 4.
 10444 toluiferum Mx. tolu-bearing i • m. . . . S. Amer. ... C p.l Wood. 3. 193

- * 1237. SOPHCTRA L. SOPHORA. (Altered from *Sophera*, the Arabic name.) *Leg. Pap. Soph.* 12.—15.
 10445 japonica L. Japanese [■] Japan 1763. ■ Bot. reg. 585
 2 fuliginea variegkritis varicgated-lvd [■] Japan
 3 plndula drooping Japan [■] Japan
 10446 chinensis Lo. C China [■] China 1818.
 10447 glauca Uch. glaucous [■] E. Indies 1818.
 10448 havanensis Jac. Havannah [■] Havannah 1823. p.l Jac. am. 173.1
 10449 tomentosa L. woolly [■] India 1690. p. Tr. ehr. 59
 10450 crassifolia Hit. thick-leaved [■] Guinea 1818. p.l
 10451 secundiflora Lag. side-flowering [■] N. Spain 1820. Df Cav.ic.5.401
 10452 littoralis Schr. shore [■] Brasil 1820. Df
 10453 flavescentes H. K. flavescent [■] Siberia 1785. D r.l
 10454 galgooides Pall. Galgo-like [■] Siberia 1817. D r.m Pal. as. 88
 10455 olopecuroides L. Foxtail-hke [■] Levant 1731. D r.l PaLas.87
 10456 sericea Nut. silky [■] N. Amer. 1820. D r.m
Astragalus carndus Ph.

1238. ORMCTSIA Jack. ORMOSIA. (*Ormos, a necklace; seeds worn in necklaces.*) *Leg. Pap. Soph.* 2.—*
 10457 dasycarpa Jack. thick-fruited t or 20 in.jl B W. Indies 1793. C lt.1 Lin. tr. 10.26
 10458 coccinea Jack. acalet-seeded j Q or 20 jl.au B Guiana 1823. C s.l Lln.tr. 10.27
Robinia coccinea Aub.

- * 1239. ANAGVRIS Tou. BEAN TREFOIL. (*Ana, backward, gyros, a circle; pod.*) *Leg. Pap. Soph.* 4.
 10459 fo'tida L. fetid ft [—] or 9 ap.my Y Spain 1170. C p.l Bot cab. 740
 10460 glacea Hort. glaucous [■] or 6 ap.my Y S. Europe 1800. C p.l
 10461 latifolia W. en. broad-leaved [■] or 10 ap.my Y Tenenffe 1815. C p.l
 V10462 indica Wai. India * or 8 jl Y Nepal 1821. D p.l Hook.ex.fl.1⁵¹
Baptisia nepalensis Hook. Thermopsis /burnifolia D. Don

1240. MACROTROPUSDrc. MACROTROPA { *Makros, lóng, ip, Qi*, a key flower. } *Leg. Pap. Soph.* 1
 10463 fce'tida Dec. fetid * L_1 or 6 a Pm Y China
jfnagryis ftc'tida Lou. sinensis Stev. chinensis Spr.
 1241. THERMOPISISA. [^] THIERMOPSISA. (*Thermos, lupine, obris, resemblance.*) *Leg. Pap. Soph.* 4
 10464 rhombifolia Nut. rhomb-leaved [■] A or 2 jn.jl N. Amer. 1811. D r.m
Cytigus rhombifolius Ph
 10465 abacea Dec. Fabo-like [^] A or 2 jn.jl ■ Siberia 1818. D r.m Pal. as. 90.
Sophbra fa.Yacea. Pall. /upinoldes var. a. L.
 10466 lancolata R. Br. spear-leaved [^] A pr 1 Jn.jl ■ Siberia 1779. D lt.1 Bot. mag. I⁸⁹
Podalyria /upinoldes IV. Sophbra tapinofdces Pall.
 10467 corganis 2)rc. Corgan [—] A or 1 jl ■ Altai 1820. D r.m Pal. as. 90.1
Sophbra alpina Pa//. Podalyria alpina W. en.

1242. VIRG'LII Lam. VIRGILIA. (*Publius Virgilii Maro, prince of Latin poets.*) *Leg. Pap. Soph.* 6."?♦
 10468 robiniofcta Dec. Robinia-like * L_J or 8 au.s T C. G. H. 1818. C ap
Robinto capensis Brm.

- 10469 Uiteail* [—] vc Wavr-flowercd * f or 15 jn.jl Y N. Amer. 1812. E p.l Mic. ar. ie.
 10470 sylvatica Dec. wood * i_J or 4 au.o ■ C. G. H. 1816. E s.p

- Sophbra sylvatica Bare.*
 10471 affrea Lam. golden-flowered * |_J or 6 jl Y Abyssinia 1777. E ■ Her.sU.7⁵
 10472 intrbsa R. Br. intruded ti I or 8 my.au Y.w C. 6. H. 1790. C p.l Bot. mag. I⁵⁹⁰
 10473 capensis Lam. Cape * \ or 5 jl.au P.w C. G. H. 1767. E ■

- * 1243. CYCLOCTPIA Fen. CYCLOPIA. (*Kyklos, a circle, posus, a foot; base of pods.*) *Leg. Pap.* [■]
 JS*74.joni8toldestf.X. Genista-like • I J or 8 jlau Y C. G. H. 1787. C pi Bot.mag.g[^]
 10475 ^aloides Dec. Galium-like * \ or 4 jl.au Y C. G. H. 1820. C p.l Fluk.aL41*

- ^{4AM} ^{7P%} & ^{4all} l^{tea} Berg, ^spalathus callosa Brm.
 10476 latifolia Dec. broad-leaved m i_j or 5 jl.au T C. G.H. 1820. C pi
Genista ftxifolia Brm.

* 1244. BAPTISIA Ven.	BAPTISIA. (<i>Bapto</i> , to dye; blue tincture from leaves.) <i>Leg. Pap. Soph.</i> 8.—10.
10477 perfolita R. Br.	perfoliate  pr 3 au Y Carolina 1732 D c Di.cl. 102.122
10478 lanceolata Ell.	lanceolate S A or 1 jl.au Y N. Amer. 1818. D s.p
Podalyria uni flora Mx.	
10479 australis R. Br.	southern  N. Amer. 1758. D c Bot mag. 509
10480 mrtillis Nut.	soft  N. Amer. 1824. D s.p
10481 villosa Ell.	villous  N. Amer. 1811. D c
<WS tinctoria L. Br.	dyer's  N. Amer. 175a D c
J048. exaltata Sivt.	exalted  N. Amer. 1816. D c Bot mag. 1099
10484 alba R. Br.	vwhite-Jlouiered  N. Amer. 1724. D c Bot. mag. 1177
1245. PODALY'R'M Lam.	PODALYHIA. (<i>Podalarius</i> , a son of JEsculapius.) <i>Leg. Pap. Soph.</i> 13.—14.
10485 Burchillii Dec.	Burchell's *  or 6 my.jl P C. G. H. 1816. E p.l
10486 hirsuta W.	hairy  or 2 jl.au P C. G. H. 1774. E p.l
10487 sericea R. Br.	silky  or 6 ja.o P C. G. H. 1778. E p.l
KH88 cuneifolia Ven.	wedge-leaved  6 myau W C. G. H. 1804. E p.l
10489 myrtillifolia W.	Myrtle-leaved  6 ap.jl P C. G. H. 1795. E p.l
10490 ftuxifolia W.	Box-leaved  2 my.jl B C. G. H. 1790. E p.l
10491 oleaculifolia Sal.	Olive-leaved  4 my P C. G. H. 1804. E p.i
10492 styracifolia B. A.	Styrax-leaved  6 my.jl Pk C. G. H. 1792. E p.i
calyprata W.	
10493 Rlaúca Dec.	glaucous  6 my.jl P C. G. H. 3810. E Lam. il. 327.4
10494 argentea Sal.	silvery  6 f.jn W E C. G. H. 1789. E Bot. mag. 753
bittia B. M.	
10495 liparioides Dee.	Liparia-like  f.jn F C. G. H. T
J0496 subbiflorata Dec.	subbiflowered  f.jn P C. G. H. T
10497 cordata B. Br.	heatx-leaved  my.jl B C. G. H. 1794. E
1246. CHORI'ZEMA Lab.	CHIORIZEMA. (<i>Choros</i> , dance, <i>xema</i> , drink.) <i>Leg. Pap. Soph.</i> 4.—5.
10498 Ucifolia Lab.	Holly-leaved  ^Jor 2 mr.o Y.H N. Holl. 1803. S s.p Lab. vo. 181
10499 nana B. M.	dwarf  LJ or ^Jmr.o Y.B N. Holl. 1803. S s.p Bot. mag. 1032
1p500 rhimbæa R. Br.	rhomb-leaved  i.J or 2 ap.jn Y N. Holl. 1803. S <>p
J0501 Henchmanni IL Br.	Henchmann's  spl 2 ap.jn S N. Holl. 1824. C s.p Botreg. 986
1247. PODOLO^BIUM R. Br.	PODOLOBIUM. (<i>POUS</i> , foot, <i>tobas</i> , pod; with footstalks.) <i>Leg. Pap. Soph.</i> 5.—7.
?5LJ2trilobatum<^r.	three-lobed *  J < 2 ap.jl Ys N. a.W. gI S 8J got. TM*^TM^n
10503 stauroph^ilum Dec	croas-leaved *  i.J pr 2 mr Y N. Hull. 1803. S s.p
^quifblum ho. C.	
10504 heterophyllum Cun.	various-leaved M J or S ap.jl  N. Holl. 1824. E s.p
10505 scandens Dec.	climbing A. L J or 3 ap.jn  N. Holl. 1824. E 8.p
Chordzema scandens Sm.	
2 humifusum Dec.	trailing >^i.Jor 1 ap.jn T N. Holl. 1823. C s.p
Davcaria humifusa Sieb.	
10506 cortaeum Dec.	leatherUeaved * i.J or 2 ap.jn Y N. Holl. 1824. C Bp
1848. OXYLO^BIUM Andr.	OXYLOBIUM. (<i>Oxys</i> , sharp, <i>lobos</i> , pod;  intedl  N. Holl. 1824. E s.p
Arborescens * s.t.	leaved * i.Jsp 1 2 ap.my O  N. Holl. 1824. E s.p
10488 arborescens ZJ. J7r.	arborescent * J or 6 ap.jn Y V.D.L. 1805. C s.p Bot. reg. 392
10509 ellipticum R. Br.	ellipticoidMT* * J or 3 my.s Y V.D.L. 1805. C s.p Lab. n.11. 135
Upio cordifolium /wrrf.	hearUeaved  i.Jor 3 aps Y N. S. W. 1807. C a.p Bot rep. 492
spinosum Dec.	spiny  or 2 ap.jn Y N. Holl. 1825. C s.p
Cirugineum Lo. C.	fusly  or 2 ap.jn Y N. Holl. 1820. C s.p
obtUBiRilium Swt.	blunHeaved  i.J or 2 mr D.O N. Holl. 1824. C s.p Sw.au. 5
1<514 Pulten* Dec.	Pulteney's  S Ven. CALLiSTACiwa /Kilos, beautiful, <i>stachus</i> , spike.) <i>Pap. Soph.</i> 8.
1p249. CALLFSTACHYS lanceoVata Ven.	spear-leaved * LJ or 4 jn.au Y N. Holl. 1815. C s.p Bot. reg. 216
ovate-leaved * LJ or 3 jn.au Y N. Holl. 1815. C s.p Bot. mag. 1925	
imJZ<<eata Sm.	wedge-leaved m \ J or 3 jn.au T N. Holl. 1824. C s.p
10518 UnariajMia Ma. C.	Toad-flax-Ivd * i.J or 6 jn.au T N. Holl. 1824. C s.p
J250. BRACHYSETMA R. Br.	BRACHYSEMA. (<i>Brachys</i> , short, <i>sema</i> , standard.) <i>Leg. Pap. Soph.</i> 2.—3.
J0510 Utifolium it. Br.	broad-leaved ft. i.J or 3 apil C N. Holl. 1803. C s.p Bot. reg. 118
10520 undulatum Ker	wave-leaved JU uJ cu 3 mr.ap G N. & W. 1820. C s.p Bot reg. 642
* 1251. GOMPHOLO^BIUM Sm.	GOMPHOLOBIUM.  Saf'MT ^{1^&''**1*! ''} s.p
10o21 bartigerum Dec.	bearded-keeled M LJ or  apal
fimbriatum Sieb.	
10522 grandiflurum Sm.	large-flowred  S mr.s Y N. S.W. 1803. C s.p Bot. reg. SI
2 scitifolium Sm.	bristle-leaved  2 mr.s Y N. & W. 1803. C s.p
10523 latiflrium J. K.	broad-leaved S mr.s Y N. S. W. 1803. C s.p Ex. bot 58
i Sm.	
twiggy  R. Br.	edged M de 1 mr.jl Y N. Holl. 1820. C s.p
<> pedunculare Dec.	pedunculate n. i CM or i ap.jl Y N. Holl. 1803. C s.p
T tetraetheroldes Sieb.	Tetrathcca-likctt. J or 11 apil T N. Holl. 1824. C s.p
i vdThniodes Ma. C.	Merbelia-like n. i lor Wall Y N. Holl. 1823. C s.p
IK lvmorphum R. Br.	multiform il i.J de 2 mr.au Y N. Holl. 1803. C s.p Bot mag. 1533
lanktum Cun.	woolly tL or 1 mr.in Y N. Holl. 1824. C s.p
fetculatum Ma. C.	netted-leaved n. or 14 apil Y N. Holl. 1824. C s.p
V ci iomentu8um Lab.	delicate tt ^ Jor 11 mr.ap Y N. Holl. 1894. C s.p
J0534 glabratum DeC.	tomentose LJ de 3 apil Y N. Holl. 1803. C s.p Lab. n.h. 1.134
40545 venustum R. Br.	smooth <1 Jor 1) ap.au Y N. Holl. 1820. C s.p
40546 pinnatum Sm.	beautiful *1 Jde 2 ap.3? N. Holl. 1803. C s.p
<0537 glaucescens Cun.	glauceacent <. J or h ap.jl T N. Holl. 1820. S s.p
Goinholbbium minus Sm.	
BURTO*N/y R. Br.	BUHTONIA. (<i>D. Burton</i> , a collector for Kew garden) <i>Leg. Pap. Soph.</i> 3.—4.
rougb./>Md tL J pr i my.jl Y N. Holl. 1813. C s.Lp	
BT isidin ucc.	scsile-leaved n. J or * my.jl Y N. Holl. 1824. C s.n
m^norD^c.	smaller n.1 Jor * rarjl Y N. Holl. 1812. C s.p
Goinholbbium minus Sm.	
TL Br. JACKSOVIA. (G. Jackson, librarian to A. B. Lambert.) <i>Lea. Pan. Snh</i> 4 — c	
Broom/Jr<<	*i Jpr 2jn.au Y N. S. W. 1803. c*^n^BotSb 427
spinous	* L J p r 2 ap.s Y N. Holl. 1803. C s.p Lab. n. f. 1.136

- 10543 furcellata Dec. fork-branched A u J ap.s Y N. Holl. 1824. C <.l.p Bon. nay. 11
 10544 reticulata Dee. netted «|_J or ap.s Y N. Holl. 1820. C 8-lp
 Davies/a reticulata Sm.
1254. VIMIN' A'RIA Sm. Rum BROOM. (Vimen, a twig; habit) Leg. Pap. Soph. 2.
 10515 denudata&n. stripped ft i_J or 3 jn.s Y N. Holl. 1789. C sp Bot mag. 1190
 10546 lateriflora Zir. side-flowering * LJ or 3 jn.s Y N. Holl. 1824. C s.p
 1255. SPHJEROLO^BIUM Sm. SPIAROLOBIUM. (Sphaira, sphere, lobos, pod; shape.) Leg. Pap. Soph. 2.
 10547 vimneum Sm. twiggy • I_J or 2 my.au Y N. Holl. 1802. C fl.p Bot. mag. 969
 III n us Lab.
- 10548 medium R. Br. mediate * i_J or 3 jn.au R N. Holl. 1803. C fl.p
1256. AO^TUS Sm. AOTUB. (A, privative, out, ear; ear-like appendages to calyx wanting.) Leg. Pap. Soph. 3.
 10549 villosa Sm. villous ft S ap.jn Y N. Holl. 1790. C s.p Bot. mag. 949 .
 lericoldeH Heath-like ft or 1 ap.jn Y N. Holl. 1810. C s.l.p Ven.mal. 35
 2ferruginica rusty C^or i* ap.jn Y N. Holl. 1820. C s.l.p Lab. n. h. 1.132
 10550 inehna Ma. C. hoary ft |_J or 2 ap.jn Y N. Holl. 1824. C s.l.p
 10551 virgina Dee. twiggy ft or 1 ap.jn Y N. Holl. 1824. C s.l.p
1257. DILLWY'N/i* Sm. DILLWYMA. (TV. Diliwyn*) celebrated botanist) Leg. Pap. Soph. 11.—12.
 10552 floribunda Sm. bundled wd or 2 ap.jl Y N. S. W. 1794. C s.l.p Ex. bot. 26
 10553 Tiriulia Sm. Heath-leaved ft or 2 mr.jl N. S. W. 1794. E s.l.p Ex. bot. 25
 10554 glaberrima Sm. smoothest » L_J or 2 mr.jl N. S. W. 1800. E s.l.p Bot. mag. 944
 10555 parvifolia R. Br. small-leaved ft or 2 mr.jl N. S. W. 1800. E s.l.p Bot. mag. 1527
 10556 decidue-leavcd nccde-leavcd ft or 1J ap.jn N. Holl. 1826. - Rnl
 10557 rufis Sieb. rustic ftl or 2 ap.jn N. Holl. 1824. - Blp
 2 brevifolia short-leaved fti or 2 ap.jn N. Holl. 1824. E sp.l
 3 tercifolia round-leaved «|_J or 2 ap.jn E N. Holl. 1824. E sp.l
 4 hispidula hispidish ft uJor 2 ap.jn N. Holl. 1824. -
 10558 phylloclades Cun. Phyllica-like ft L_J or 2 ap.jn N. Holl. 1824. C STP
 10559 tenuifolia Sieb. fine-leavcd ft i l or 1J ap.jn N. Holl. 1824. C s.l.p
 10560 scricca Cun. silky L_J or 1J ap.jn N. Holl. 1822 k C s.l.p
 10561 Juniperina Lod. Juniper-featwrf Ljpr 2 ap.my Y V. D. L. 1818. C s.l.p Bot. cab. 401
 10562 cincrascent R. Br. cincrascent ft or 2 mr.jl Y N. S. W. 1819. C s.l.p Bot. mag. 2247
1258. EUTA'XIA R.Br. EDTAXIA. (Eutaxia, modesty; appearance.) Leg. Pap. Soph. 2.
 10563 wyrtifolia R. Br. Myrtle-leaved L_J pr H au O N. Holl. 1803. C s.p Bot. mag. 1274
 10564 pungent pungent LJ or 4 ap.jn O N. Holl. 1825. C p.l Sw. au. 28
DiUfnia pfmgns fta.C.
1259. SCLERO'THA'MNUS R. Br. SCLEROTHAMNUS. (Skleros, hard, thamnos, shrub.) Leg. Pap. Soph. 1.
 10565 microphyllus R. Br. smalUleaved ft L_J or 2 my.jn Y N. Holl. 1803. C Blp
1260. GASTROLCPBIUM it. Br. GASTROLODIUM. (Gaster, belly, lobos, pod; shape.) Leg. Pap. Soph. 1.—3.
 10566 bilobum R. Br. two-lobed ft i_J or 2 mr.my Y N. Holl. 1803. C s.l.p Bot. reg. 411
1261. EUCHPLUS R. Br. EIJCHILUS. (Eu, well, cheilos, a lip; well lipped.) Leg. Pap. Soph. 1.
 10567 obcordatus R. Br. obcordate-fw* L_j or 2 mr.jn Y N. Holl. 1803. C s.l.p Bot. cab. 60
1262. PULTEN^ii Sm. PULTKVEA. (IV. Pulteney, M.D., a botanical author.) Leg. Pap. Soph. 43.
 10568 /aphnoldes Sm. Daphne-like ftl Jor 2 ju.jl Y N.S. W. 1792. C s.l.p Bot. mag. 1391
 10569 obcordiat Andr. obcordate/w/ or 1 my.jl Y V. D. L. 1808. C s.l.p Bot. rep. 574
 10570 biloba It. Br. two-lobed ft uJor ap.my Y N. S. W. 1817. C fl.Kp Bot. mag. 2091
 10571 sc2bra H. K. rough-leaved ft or my.jl Y N. S. W. 1803. C s.l.p
 10572 incurvata Cun. incurved fti or my.jn Y N. Holl. 1823. C s.l.p
 10573 ferruginea Rud. rusty ft or 2 ap.jn Y N. Holl. 1810. C s.l.p Lin. tr. 11. S3
 10574 raccmulosa Dec. raccmulose ft or 2 ap.jn Y N. Holl. 1820. C s.l.p
 10575 parviflora Sieb. small-flowered ft or 2 ap.jl Y N. Holl. 1824. C s.p.l
 10576 rtricta B. M. strict or 2 ap.jn Y N. S. W. 1803. C s.p.l
 10577 elliptica Sm. elliptic ft or 2 ap.jn Y N. Holl. 1810. C Blp Bot. mag. 1588
 10578 plumosa Sieb. feathery »|_J or 1* ap.jn Y N. Holl. 1824. C Blp Lin. tr. 11.24
 10579 hymifolia Sieb. Thyme-leaved ft or 1 ap.jn Y N. Holl. 1810. C 8.l.p
 10580 hypolempira Sieb. brightish ft or 2 ap.jn Y N. Holl. 1824. C s.l.p
 10581 villifera Sieb. hair-bearing ft or 2 ap.jn Y N. Holl. 1817. C s.l.p
 10582 xilygalifolia Rud. Polygala-lvd or 2 ap.jn Y N. Holl. 1801. C s.p.l Lin. tr. 11.25
 10583 flexilis Sm. flexible ft or 1 ap.jn Y N. S. W. 1820. C fl.F.p
 10584 cuch'ila Dee. Euehilus/Z/Ae retusus ft or 1 ap.jn Y N. Holl. 1789. C s.p.l
 10585 retusa Sm. retusus ft i lor 1 ap.my Y N. S. W. 1824. C l.s.p Bot reg. 373
 10586 cunca Lo. C. v/rdgc-leaved fti or 1 ap.jn Y N. Holl. 1824. C s.p.l
 10587 oxahdifoia Cun. Oxalis-leaved ft or 2 ap.jn Y N. Holl. 1826. C Bpl
 10588 capitellata Sieb. sm.il I-headed ft or 1* ap.jn Y N. Holl. 1823. C Bpl
 10589 /inophylia Sm. Flnx-leaved ft or 2 my.jl Y N. S. W. 1789. C fl.p.l
 10590 microphylla Sieb. 0mall-lra/cd ft or 1 ap.jn Y N. Holl. 1810. C s.p. Sc. nan. 3.18
 10591 stipulans Sm. stipular ft or 2 ap.jl Y N. S. W. 1792. C Bpl
 10592 paleocea Sm. chatty il ap.jn Y N. S. W. 1824. C fl.p Bot. mag. 435
 10593 aristata Sieb. awned ap.jn Y N. Holl. 1823. C s.l.p Bot.cab.291
 10594 cchinula Sieb. small-prickled ft or clothed ft ^Jor 3 ap.jl Y N. Holl. 1803. C sp.l
 10595 vestita R. Br. canescens ft or 1 npmy Y N. Holl. 1822. C sp.l
 10596 canescens Cun. Polium-leaved ft or 2 ap.jn Y N. Holl. 1820. C fl.l.p
 10597 polifolia Cun. ft or 2 ap.jn Y N. Holl. 1824. C s.l.p
 10598 dentata Lab. ft or 2 ap.jn Y N. Holl. 1824. C s.l.p
 10599 Juniperina IMb. thin-leaved ft or 1 ap.my Y N. S. W. 1817. C s.p.l Lab. n. h. 1.131
 10600 tonuitifolia It. Br. villous ft or 2 ap.my Y N. S. W. 1790. C s.l.p Lab. n. h. 1.1 &
 10601 vilif.sa Sm. thick-leaved fti or 2 ap.jn Y N. S. W. 1824. C s.l.p Bot mag. 2086
 10602 crassifolia Lo. C. flesh-colored ft or 1 ap.jn Pa.R N. Holl. 1824. C s.l.p
 10603 incarnata Ma.C. silvery ft or 1 ap.my Y N. Holl. 1823. C s.l.p
 10604 argentea Cun. procumbent ft or 1 ap.my Y N. Holl. 1824. C s.l.p
 10605 proctimbens Cun. C. Rosemary-lvd ft or 2 ap.jn N. Holl. 1824. C Blp
 10606 rosmarinifolia Ma. 10307 aaiwra Sieb. rough ft or N. Holl. 1822. C s.l.p
 10607 comiva Sieb. tufted ft or N. Holl. 1825. C s.p.l
 10608 squarrtsa Sieb. avarrose ft or 5 ap.jl N. Holl. 1822. C Bpl
 10609 phytoco/des Sieb. alylica-hke ft or 1 ap.jn N. Holl. 1825. C Bpl
 10610 cundida Lod. white-leaved ft or liap.jn N. Holl. 1825. C Bpl
C Bpl Bot. cab. ISSo

DAVIETSL	Sm.	DAVIBSIA.	(Rev. H. Davies, F.L.S., a Welshbotanist)	Leg. Pap. Soph.	13.
w ^s 12 racemulosa Dec.	racemulose	i or 2) jn.au	N. Holl. 1823.	C s.p. I	
*0013 umbellulata Sm.	small.umbellcd	i or 2) ap.jn	N. Holl. 1818.	C sf.p	
JL104 inrrauata S/n.	thick-leaved	i or 2) iny.jl	N. Holl. 1820.	C s.p.	
aciculari* Sm.	ncclv.e-leaved	it or 2) injnl	N. S. W. 1804.	C s.l.p	
ggttina Sm.	Vunc-leauott	t J or 3 ap.au	N. S. W. 1712.	C s.l.p Bot. cab. 1234	
JJ109 juniprira lad.	Junipor-/ite	L_J or 2 my.jl	N. Holl. 1825.	C s.l.p Bot. cab. 4i	
10618 corymbftsa S/n.	corymbose	i or 2 my.au	N. S. W. 1804.	C s. l.p	
I0619 mimojoldes R. Br.	Mimos-like	L_J or 2 jn.au	N. S. W. 1809.	C s.l.p Bot rep. 526	
glifica Led.	broad-leaved	i or 3 my.au	N. S. W. 1805.	C s.l.p Bot. mag. 1757	
JJ620 latifolia A. Br.	squarese	A or 21 my.jl	N. Holl. 1824.	C s.l.p	
JUB21 squarrosa S/».	cordite-leaved	i or 3 jn_jl	N. Holl. 1824.	C s.l.p Bot. reg 1005	
10622 cordata Sm.	winged	ti or 3 my.au	N. S. W. 1818.	C s.l.p Bot. reg. 728	
10523 alataS/«.		L_J or 2 my.jl	M. Holl. 1823.	C s.l.p	
Ofi24/uncea Sm.					
1264. MIRBE'L/yf Sm.	MIR BELT A.	(C. F. B. MirbcL a celebrated French physiologist.) Leg. Pap. Soph. fi.			
10625 reticulata Sm.	reticulated	L_J or 3 my.au	N. S. W. 17JK.	C s.l.p Bot mag. 1211	
10626 dilatata R. Br.	dWatlm-leaved	i or iny.au	N. Holl. 180i.	C s.l.p Bot. reg. 1041	
10627 speciosa Steb.	showy	i or myjl	N. Hull. 1824.	C s.l.p	
lp^e püngens Cun.	pungent	i or 2 myjl	N. Holl. 1824.	C s.l.p	
10619 grandiflora B. M.	large-flowered	L_J or 2 myjl	N. Holl. 1825.	C s.l.p Bot. mag. 2771	
10630 Baxter's Ma. C.	Baxter's	JU L_J or 2sp <i>suau</i> Yaa	N. Holl. 1828.	IL spplBot mag. 1434	
1265. CE'RCIS L.	JUDAS THEE.	(Kerkis, a name of Theophrastus's.) Leg. Cats. Cass. 2.			
10631 Siliquastrum	*£ or 20	my.jn R	S. Europe 1596.	L co Bot. mag. 1138	
2 florc albo	white-flowered	\$ or 20	my.jn W	L r.m	
3	small-flowered	\$ or 20	my.jn P	Buchana 1827.	L co
AU632 canadensis L.	Canadian	J 18	my.jn Pa.R	N. Amer. 1730.	Lt co
1266. SCHOT^T/iff Jac.	SCHIOTIA, (R. van der Schot, Jacquin's fellow-travel, in Amer.) Leg. Cats. Cass. 4—5.				
10633 speciaja Jac.	showy	• or 5 jld	C C G. H. 1759.	C Tp Hot. rej. 348	
10634 nwirtri/dfolia Afz.	Tamarind-lvd	tt or 6 my.i	C C G. H. 1795.	C lp Bot. mag. 1153	
10635 stipulata H.K.	/or^stipuled	> G or fi iny.s	C C G. H. 1794.	C Lp	
10636 alitff Thun.	winged	L_J or 5 jii.jl	C C. G. H. 1816.	C lp	
1267. OMPHALCTBIUM Jac.	OMPHIALOBIUM.	(Omphalos, the navel, lobos, a pod.) Ixg. Qes. Cass. 1.			
10637 Schftcia Jac.	Schot's	H L 6 my.jl	Pa.P C. G. H. 1810.	C r.m Jac fr. 15.4	
Schbt/a latifolia Jac.					
1268. BAUVXIA Plu.	MOUNTAIN EBONY.	(J. and C. Bauhin, two celeb. botanists.) Leg. (Ues. Cass. 45.-68.			
JO638 divaricata L.	divaricata	• or 5 jn.s	W W.III.II. 1718.	C It1 H. cl. 15	
10639 Lamarkiana Dec.	Lamark's	• or ■	W W.Amer. 1818.	C a1	
*X ⁸⁰ rctbsa Pair, divaricata Lam. and Delf.	spathacaous	• or 6	W Mexico 1823.	C It1 Fl. mex. ic. in.	
subrotundiflida Cau.	roundish-lvd	• or 6	W Acapulca 1820.	C It1 Cav. ic. 5.406	
10642 lunkria Cav.	haff-xnoow-lvd	• or 6	W Acapulca 1820.	C It1 Cav. ic. 5.407	
H. H. H. / / ^	long-cared	■	W Jamaica 1756.	C It1 Mil. ic. 1.61	
10634 Porrecta Svat.	stretched out	■	W Indies 1737.	C It1 Bot mag 1708	
latiflaria Cav.	broad-leaved	pr	W. Indies 1818.	C It1 Cav. ic. 5.405	
H.K.	white	• or 6	my.jn W E. Indies 1777.	C It1	
647 Jcuinianata L.	acuminated	• or 8	W E. Indies 1808.	C It1 R. mal. 1.34	
J06W Pauletti Pers.	Pauletia	• or 4	W Panama 1820.	C It1 Cav. ic. 410	
lrgo-flowered	spathecious	• or 4	W Peru 1820.	C It1	
lesteuler-))ctaled	roundish-lvd	• or 5	W NwSpain 1818.	C It1	
aculcata L.	pr/cly-stal/kcd	I or 5	(* jn.au W. Indies 1737.	C It1 PL ic. 44.1	
	lver-leaved	• or 6	W Brasil 1823.	C It1	
	emarKinata	Q C	W Carthag. 1700.	C It1	
0654 annata Otto	armed	Q C	W Brazil 1824.	C It1	
0655 incrims Pers.	unarmed	6	Y.B. Acapulca 1810.	C It1 Cav. ic. 5.409	
JUSSII rufescens Lam.	rufescens	Oor	Pd R Africa 1810.	C It1 Lam. il. 239. S	
JJ657 parviflora Vahl	smdll.flowered.	6	E. Indies 1808.	C It1	
OK58 madagascarisDt/F.	Madagascar	4	W Madagasc. 1826.	C It1	
10659 dngula Jac.	clawed	5	Carccas 1817.	C It1 Jac. fr. 15.1	
aculekta Jac. am.	VI.2	5	W		
variegata L.	variegated	• or 6	St E. Indies 1690.	C It1 R. mal. 1.32	
chiensis Dec.	China	• or 6	K China 1800.	C It1 Braam 26	
*0612 tomentosa L.	tomentose	• or 6	Y.w E. Indies 1808.	C It1 R. mal. 1.35	
JJ663 pubescens Dec.	pubescent	• or 4	W Jamaica 1823.	C It1	
J0664 multinervia Dec.	many-nerved	• or 5	W Caracas 1817.	C It1	
Klaueccens Dec.	glaucescent	• or 6	W Cumana 1817.	C It1	
Vahl	racemos	£ sp 20	E. Indies 1790.	C It1 Vahl sy. 3.62	
Rox.	purple	• or 6	E. Indies 1778.	C It1 R. mal. 1.33	
corymbosa Rox.	recluse	• or 7	W E. Indies 1820.	C It1	
anguina ROT.	corymbose	• or 6	E. Indies 1818.	C It1 Dec. leg. 13.70	
scandens L.	small Upd/ climbing	« f 30	W.Y E. Indies 1790.	C It1 IL mal. 8.29	
U «lian « "818 Aub.	Ullana	f 20	W Guiana 1820.	C It1 Aub. (jui. 1.145	
		6	W Caracas 1K24.	C It1 Kth. mini. 46	
		15	W Carthag. 1810.	C It1 Jac. am. 17J. 3	
		6	W S. Amer. 1800.	C It1 Her. am. 315	
		6	W E. Indies 1817.	C It1	
		8	W E. Indies 1820.	C It1	
10076 semibifida Rox.	semibifld	ar. 4. CD or	W E. Indies 1817.	C It1	
W77 ferruginea Rox.	rusty.leaved	10	W E. Indies 1820.	C It1	
10378 malantrica Rox.	Malabar	15	W E. Indies 1810.	C It1	
10579 triandra Rox.	triandrous	CD or 15	W E. Indies 1823.	C It1	
dica Lo.C.	Indian	CD or 6	W E. Indies 1820.	C It1	
Hort.	small-leaved	o 6	W a Amer. 1817.	C It1	
	showy	10	W 1820.	C It1	
HYMENIE"-4 L.	LOCUST THEE.	(Hymen, god of marriage, leaves joined.) Lee. Can Cma i R			
loSt erru	Courbaril	t f i or 20	W. Indies 1688.	C pi Lam il 330"R	
**^ CandoHcirt Kth	warty-podded	TCJ or SO.	Madagasc. 18(8).	C pl p.l 330"2	
	Dccandolle's				

DECANDRIA MONOGYNIA.

1270. CYNOME^TRA L.	CYNOMETOA. (& «*» dog, mctr., matrix; pods.)	Leg. Cas. Can. SL—4.
10686 caulinflora L.	stem-flowering t • or 30	RR E. Indies 1804. C s.l.p Lam. f. 331
10687 polyandra tfo*.	polyandrous I • or 20	R E. Indies 1822. C s.l.p Rox. cor. 28C
• 1271. CA'SSIA L.	CASSIA. (<i>Katsa</i> , to tear off, <i>Arab.</i> , bark stripped from tree). /.-<. Ors. Cass.. 106. — 204.	
10688 bacillaris L.	rod ■•• or 3	Y B. Indies 1782. C ro Bot. reg. 881
10689 speciosa Schr.	showy ■•• or 5	jnjl Y Brazil 1816. C l.p
10690 corymbosa Lam.	corymbose ■•• i Jor 3	Y B. Ayres 1796. C s.p Bot mag. 633
10691 floribunda Cav.	bundle-flwd ■•• or 4	jnjl Y New Spain 1818. C l.p
10692 lasigkta W. en.	smooth ■•• pr 3	my.au Y — — C UI
10693 pendula W. en.	pendulous ■•• or 3	Jl.au Y S. Atmer. 1820. C p.1
10694 torifera Cav.	torose ■•• or 6	jnjl Y China 1816. C s.l Jac. ic. 73
10695 chinensis Lam.	Chinese « LJ or 4	jnjl Y China 1807. ■ P.1
10896 coromandeliana Jac. I.	Coromandel i r i o r 8	jnjl Y Coroman. 1823. ■ 8.1 Jac. fr. 100
not IV en.	3	
10697 Sophora L.	Sophora • • or 4	Jls Y E. Indies 1658. ■ It.1 R. mal. 2.52
10698 bracte-ita £.	bractcatc • I lor 6	ap.d Y W. Indies J822. ■ co
10699 alata, L.	winged-leaved • I lor 12	... Y W. Indies 1731. C p.1 Jacob. 245.2
10700 obovata Coll.	obovate-lvd Senna O m	li jl.au Y Egypt 1610. ■ It.1 Jac. ec 1.87
St/nna Lam.		
10701 ligustrinoides Schr.	Ligustrina-like tt. • un	I jl.au Y Arabia 1800. C It.1
10702 lanceolata Forsk.	lanceol. true Senna*. • m	I jl.au Y Levant ... C It.1 Del. asg. 27.1
orientals Pcrs.	acutuoba Dec.	
10703 italica Lam.	Italian Senna O or 3	jnjl Y S. Europe — S co M. h. 2.24.2
10704 Tardian Kth.	Tarantan ■•• or 1	ljjl.au Y Cumana 1817. C 8.1
10705 Richardia Vna Kth.	Richard's «••• m 1	jl.au Y Cumana 1823. C 8.1 Kth. mim. 42
10706 sericea Swz.	silky-/wirf O tin	Jl my.au Y Jamaica 1731. S s.p Jac. ic. 3. 459
10707 obtusifolia L.	blunt-leaved ED un 2	jl.au Y Jamaica 1732. S Di. el. 62.72
10708 ciliata Hafm.	ciliated a. n. m	I Y Cuba 162a S
10709 humilis Coll.	humble lf ITTm	■ Y S. Amer. 1810. S 8.1 Pl. ed. B. 76.2
10710 Tbra L.	Tora ■•• un	au Y E. Indies 1693. S r.m Di. el. 6a 73
10711 ciliaris Coll.	ciliated-/-wrf jf EH m	E. Indies J817. S ■
10712 pentagibbia Mil.	five-angled If Blm	jnjl Y Peru 1700. S 8.1
10713 Tagera Lam.	Tagera ■■■■■	jnjl Y E. Indies 1803. S p.1 R. mal. 2.53
10714 viminea L.	twiggy ■•• pr	Y W. Indies 1786. C p.1 Sl. jam. 2.189.67
10715 macranthra Dec.	long-anthered ••• or	Y Brazil 1824. C s.l Col. cas. 8
10716 quinquangularis Rich.	five-angled ••• or	Y Cayenne 1818. C s.l
10717 scindoides Jac.	Senna-like ■■■■■	3 jnjl Y E. Indies 1808. C s.p Jac. ic. 1. 70
10718 Apoucouita Aub.	Apoucouita ••• or	8 jl.au Y Surinam 1820. C s.p Aub. gui. 1.146
acuminata IV.		
10719 bicapsularis L.	two-capsuled ••• or	■ my.jn T W. Indies 1739. ■ p.1 Pl. ic. 76.1
10720 mollissima H. \$B.	softest-leaved ••• or	■ my.jn T S. Amer. 1816. ■ p.1
10721 aTtem inoides Gau.	Artemisia-like ■ I or	■ T N. Holl. 1820. ■
10722 glutinosa Dec.	clammy ■ I or	■ T N. Holl. 1818. ■
10723 dispar W. en.	unequal ■•• or	■ S. Amer. 1824. ■ Bp
10724 Bértcri Coll.	Barter's ■•• or	■ W. Indies 1824. ■ It.1 Col. h. rip. 24
10725 acapulcensis II. Sf/B.	Acapulca ■•• or	■ Acapulca 182a ■ s.l
10726 glauca Lam.	glauous ■•• or	■ E. Indies 1818. ■ L
10727 sulphurea Dee.	sulphur ■•• un	■ E. Indies 1800. ■ p.1 R. mal. 6.9,10
arboriscus I'ahl		
10728 biflora /.	two-flowered ■■■■■	6 ap.d ■ W. Indies 1766. C ■ Bot. mag. 810
10729 geminiflora M. & S.	twin-flowered ■■■■■	5 jnjl ■ Mexico 1824. C ■ CoL cas. 3
10730 pallida Vahl	Silc-yellow ■■■■■	6 jnjl ■ S. Amer. 1818. C ■
10731 Bardayna Swt	Sarclay's ■■■■■	4 jnau ■ N. Holl. 1827. C ■ S.p
10732 mexicana Jac.	Mexican ■■■■■	5 jnjl ■ Mexico 1824. C ■ s.l Jac. sc. 203
10733 tomentosa Lam.	tomentose ■■■■■	15 Jls ■ S. Amer. 1822. C ■ co Jac. ic. 1.72
multiglandulosa Jac.		
10734 stipulacea H. K.	/ff/gc-stipuled ■■■■■	■ Chile 1786. ■ It.1 Feu. ob. 3.42
10735 cupidata W. en.	cupidate ■■■■■	■ S. Amer. 1820. ■ co
10736 auriculata L.	small-cared ■■■■■	■ E. Indies 1777. ■ It.1 Pluk. al. 314. 4
10737 fastigiata Vahl	peaked ■■■■■	■ E. Indies 1818. ■ 8.1
10738 frondosa H. K.	frondose ■■■■■	■ W. Indies 1769. ■ It.1 Jac. ic. 1.74
crista Jac.		
10739 polypilla Jac.	many-leaved ■■■■■	■ Q 4 jnjl ■ W. Indies 1816. C 8.1 Jac. ic. 460
10740 marginata W. en.	marginate ■■■■■	■ 3 my.jl ■ Surinam 182a C co
10741 hirta W. en.	hairy ■■■■■	■ 3 jl.s ■ S. Amer. 1820. C co
10742 longisiliqua L.	long-podded ■■■■■	■ 6 jnjl ■ W. Indies 1800. C s.1
10743 icerptiaca W. en.	Egyptian ■■■■■	■ 3 my ■ Egypt 1822. C co
10744 occidentalis L.	western ■■■■■	■ W. Indies 1759. C p.1 Bpt. reg. 83
10745 planisiliqua L.	flat-podded ■■■■■	■ W. Indies 1822. C co Pl. am. 77
10746 purpurea Rox.	yur\c-stemmc4* ■•• or	■ myjl ■ E. Indies 1821. C co Bot. reg. 856
10747 hirsuta L.	hairy ■•• or	■ jn.au ■ America 1778. C s.p Jac. sc. 270
10748 linckris Mt.	Wucax-podded ■■■■■	■ jn.jl ■ Carolina 1800. C 8.1
10749 p^tula H. K.	spreading ■■■■■	■ aus ■ W. Indies 1778. C It.1
10750 pubescens Jac.	pubescent ■■■■■	■ S. Amer. 1818. C 8.1 Jac. fr. 57
10751 filiflata Dec.	furrowed-branched ■■■■■	■ S. Amer. 1820. C 8.1 CoL cas. 6
C^rnua Balb.		
10752 ruscif?ia J/ic.	Buscus-leaved ■■■■■	■ Madeira 1816. C It.1 Jac. ic. 1.71
10753 ligustrina L.	VrvicUcaved ■■■■■	■ Baham. Is. 1726. C p.1 Bot. reg. 109
10754 rohtoioides W. en.	Robinia-hkc ■■■■■	■ S. Amer. 1823. C CQ
10755 manlandica L.	Maryland ■■■■■	■ N. Amer. 182a D s.p Di.cL26O.3S0
10756 nigrans Vahl	blackish ■■■■■	■ Egypt 1817. C 8.1
10757 elliptica H. & B.	elliptic ■■■■■	■ Trinidad 1818. C 8.1
10758 emarginata L.	-notch-leaved ■■■■■	■ Jamaica 1759. C p.1 Sl. jam. 2.180.1>
10759 atomaria L.	atomed ■■■■■	■ S. Amer. J810. C 6.1
10760 fl6rida Vahl	large-flowered ■■■■■	■ E. Indies 1820. C s.1
10761 reticulata FT. en.	netted ■■■■■	■ S. Amer. 1821. C co
10762 spectabilis Dec.	showy ■■■■■	■ Caracas 1820. C 8.1 CoL cas. 7 no
J.152 aversiflora Herb.	contrary-flwd ■■■■■	■ Brazil 1820. C co Bot. mag. 2638
10764 AIMUB L.	Absus ■■■■■	■ India 1777. S It.1 Bur. zey. 97
10765 Thonningii Dec.	Thonning's ■■■■■	■ Guinea 1824. S s.1
J.066 /otoldps J. \$B.	Lotutlike ■■■■■	■ Trinidad 1820. C 8.1
10767 diphlla Lam.	two-leaved ■■■■■	■ W. Indies 1781. ■ It.1 Cav. ic. 5. 600
	2 myjl	

10768 bifoliata <i>Dee.</i>	bifolate	H Q o r	2	jn.jl	Y	Brazil	1820.	S	81	Col. caa. 9i3	
10769 gracilis <i>Kth.</i>	slender	• • or	2	jn.jl	Y	Orinoco	1817.	S	81	Kth. mim. 36	
10770 tenifolia <i>H.8(B.</i>	weak	• • or	2	jn.jl	Y	Orinoco	1820.	S	s.l		
10771 hneata <i>Stvz.</i>	lincated	it Q o r	1	jn.jl	Y	Jamaica	1818.	S	s.l		
10772 prostrata <i>W. en.</i>	prostrate	1. Eg or		\ my.au	Y	S. Amer.	1819.	S	co		
10773 pilosa <i>L.</i>	pilose	tf E3 pr	1	jn.jl	Y	Jamaica	1818.	S	s.l		
10774 trifurca <i>Jac.</i>	three-flowered	Opr	1	jn.jl	Y	W. Indies	1816.	S	s.l	Jac. sc. 480	
10775 Hurnmanni/ Dec.	Burmann's	.Q pr	1	jn.jl	Y	C G. H.	1810.	S	s.l		
flexuös.i B>m.											
10776 vir'ta Sun.	twi _{BKV}	«*-Opr	1	jn.jl	Y	W. Indies	1810.	S	si		
10777 nictitans <i>L.</i>	twinkling	O pr	2	jl	Y	N. Amer.	1800.	S	lt.l	PI al. 314.5	
10778 Chamnecrista <i>L.</i>	Ground Crista	tt. • pr	1	jns	Y	America	1699.	S	r m	Bot. mag. 157	
10779 glandulifosa <i>L.</i>	glandulous	O pr	4	au.o	Y	W. Indies	1822.	S	co		
10780 mimosoldes <i>L.</i>	Mimosa-hke	O pr	2	jns	Y	Ceylon	1806.	S	lt.l		
10781 cipensis <i>Thun.</i>	Cape	*LL_Jor	1	jn	Y	C. G. H.	1816.	S	Itl	Bot. cab. SI 1	
10782 piinula <i>Lam.</i>	dwarf	- * O pr	1	jn.jl	Y	E. Indies	1816.	S	s.l	Pluk. al. 120.1	
procumbens <i>L.</i> not F											
10783 procumbens <i>W.</i>	procumbent	- * O un	1^	jn.jl	Y	N. Amer.	1806.	S	lt.l	Com. pet 11	
10784 vEschynom6menac Dec.	vEschynomene	ED or	1	jn.jl	Y	"W. Indies	1810.	S	lt.l	Col. caff. 17	
10785 Parkeria/na Dec.	Parker's	tt.	• or	1	aus	Y	Demerara	1817.	S	lt.l	
10786 äspera <i>Mhl.</i>	rough	O or	1	jl.au	Y	Georgia	1818.	S	lt.l		
10787 angustissima <i>L.</i>	narrowest-/^	O or	1	jl.au	Y	E. Indies	1820.	S	lt.l	Ku. am. 6.67.1	
10788 Wallichdmi Dec.	Walch'h'a	O o r	1	jn.jl	Y	Nepal	1817.	S	Itl		
dimidiata <i>Rax.</i>											
10789 flexuba <i>Z.</i>	flexuous	O	or	1	jl.au	Y	Brazil	1810.	S	lt.l	Brey. c 23
10790 microphf lla <i>W.</i>	small-leaved	O pr	1	jns	Y	San. Cruz	1810.	S	lt.l		
10791 arcñaria //, k. B.	sand	tt.	• or	1	jnau	Y	Maypures	1819.	S	r.m	
10792 australis <i>B. M.</i>	southern	* i	• or	3	jl.au	Y	N. Holl.	1824.	C	Itl	Bot. mag. 2676
10793 pulchella.8\$.	pretty	• [J	or	3	jl.au	Y	Mauritius	1825.	C	UI	

1272. CATHARTOCA'RPU\$ Per*. CATHARTOCARPUS. (*Kathairo*, to purge, *karpos*, fruit.) Leg. Ctes. Cass. 8-13.
10794 brasilianus Lam. Bréhílian IQhr 30 -- Y "S. Amer. 1820. C l.p. Brey. c. 21
Glossy Catbird. See also V. II.

127a PARKINSON [%]N/J4 L. PARKINSONIA. (*J. Parkinson*, apothecary, and bot author.) *Leg.Cees.Cass.* 1.
10802 aculeata L. prickly * Q o r 12 ... Y w. Indies 1739. C Lp Jac am. 80

* 1274. POINCLITJY <i>i4</i>	L. FLOWBR FENCE.	(<i>M. de Pomici</i> , governor of the Antilles.)	Leg. Oes. Cass. 3.
1080J pulchrērnma JL	very fair	• or od 10 jn.s. Rk. o E. Indies 1691.	C r.m. Bot mag. 995
<i>Utillea</i>	yellow	• Qor 10 jn.jl Y W. Indies 1691.	C r.m.
J0004 insfgns Kth.	noble	■ Qor 15 ... Cop S. Amer. 1823.	p.l Kth. mim. 44
10805 el4ta JL	tall	■ Qor 15 ... Y E. Indies 177a	r.m.

J. 1275 CJESALPI[^]/if Blu. BRABILETTO. (C. (*ketalpbetus*, phys. to Pope Clem. VIII.) Leg. Cas. Cast. 20.-32.

J061y, ^rtH.K.	Nuga	< D c c	10	...	Y	E. Indies	1801	S	p.l	Ru. am. 5.50
J087 pamculata <i>Desf.</i>	panicled	D	1	..	Y	Malabar	1817	S	p.l	R.mal.6.19
**J08 acclandis <i>Roth</i>	scandent	D	1	..	Y	E. Indies	1800	S	p.l	
J0809 brasili^nsis <i>L.</i>	Brazilian	D	1	..	O	Jamaica	1739	S	p.l	
J0810 Sippan <i>L.</i>	Sat{>}an	D	1	..	Y	E. Indies	1773	S	p.l	Rox. cor. 1.16
J0811 mimosoldcs <i>Lam.</i>	Mimosa-like	D	1	..	Y	E. Indies	1806	S	p.l	R. mal. 6.8
J0812 2 b f 1 5	two-paired	D	1	..	Y	Jamaica	1770	S	p.l	Sl.jam.S.181.2.23
J0813 bahamensis <i>Lam.</i>	Bahama	D	1	..	W	Bahama	1820	S	p.l	Cat car. 2.51
J0814 Crista <i>L.</i>	Criata	D	1	..	W.T	Jamaica	1800	S	p.l	PI. gen. 68
J0815 Cacalaco <i>H. & B.</i>	Cacalaco	D	1	..	Y	Mexico	1824	S	p.l	H.&B.2.137
J0816 casbioldcs <i>W.en.</i>	Cassia-like	D	1	..	Y	& Amer.	1821	S	p.l	
J0817 mucronata <i>W.en.</i>	mucronate	D	1	..	Y	Brazil	1823	S	p.l	
J0818 punctata <i>W.en.</i>	dotted	D	1	..	Y	Brazil	1820	S	p.l	
J0819 enneaijhilla <i>Rox.</i>	nine-leaved	D	1	..	Y	Kindles	1816	S	p.l	
J0820 oleopcpnna <i>Rox.</i>	oil-seeded	D	1	..	Y	E. Indies	1820	S	p.l	
J0821 chiin^nsis <i>Rox.</i>	China	D	1	..	Y	E. Indies	1310	S	p.l	
J0822 tortu?sa <i>Rox.</i>	twisted	D	1	..	Y	E. Indies	1820	S	p.l	
J0823 subekSa <i>Lo. C.</i>	cork-barked	D	1	..	Y	E. Indies	1823	S	p.l	
JPJ24 vesicaria <i>Lam.</i>	bladdered	D	1	..	Y	E. Indies	1820	S	p.l	
J0825 prbceria <i>Pop.</i>	tall	D	1	..	t nor	Cuba	1824	S	p.l	

1276. REICHA'K'DIA Roth RRICUARDIA. (J. J. Reichard, a celeb. French botanist.) *Leg.Cas.Cast.* I.—■
10826 hexapetalia Roth six-petaled A Q o r 10 ... Y E Indies 1824. S p.f
Cmaloxis ligulata Hawne.

Kth. COULTBBIA. (Dr. Coulter, a botanical author.) Leg. Cat. Cass. 2.-5.
 horrid • o r 15 ... O Carthage 1824 S p.l H.&B.569
 10828 tinctoria/>». dyer+B • o r 12 ... O Carthage 1822. S p.l H.&B.568
 Poinciana tdrria R. & p. Ciscalpinio tinctoria Domb.

1278. GUILANDFNI J. NICKER FREE... (A). *Quilauan*, a Prussian trav. in Africa). Leg. Cat. Class. 2-7. 10829 Bandue L. Bondue ... * o r 12 ... Y India 1640. C s.p Lam. il. 336

1279. *MORINGA* Dec. HORseradish Tree. (Name in Malabar.) Leg. Cat. Cast. 1—4

■ 831 pterygosperma Dec. winged-seeded i • cu 15 ... Y E. Indies 1759. C pl Jac. ic. 3.461
Hyperanthera Horingia Vahl

Hoffmane Ggw Cav. Hoffmannibggia (J. C. a distin. nat) Lee. Cas. Cats
F. acaria Cfl.; iickle-tetffrfr • • c u 2jl.au e 18W. C sip cJv^{te} 398

1281. ADENANTHETRA L. ADENANTHERA. (*Aden,* a gla;id, *anthera,* an anther.) *Leg. Mimbsce.* 2. - 5.
 10833 pavorina L. peacock MO or 5 my.au Y.w E. Indies 1759. C s. l.p Jac. c. 4.23
 10834 falcata L. falcate t Q or 5 ... Y E. Indies 1812. C s.l.u Ru. am. 3.11
1282. CA^DIA Forsk. CADIA. (*Qadhy,* its Arabic name.) *Leg. Cas. Cass.* 1.
 10835 purplea W. puiple * d) c u 6 ja.jl W.p Arabia 1775. C R.t.l Pic. h. p. 9. ic.
varia Herit.
1283. JPROSO'PISL. PROSOPIS. (*Prosopw^a* a mask; by Dioscorides to *Arctium Zappa.*) *Legum. Mimbsea.* 6.
 10836 spidgera /, snake-bearing f Q e c 20 ... Y.a E. Indies 1812. S r.m Rox. cor. 1.63
 2 Adenanthera gland-anthercd \$ • ec 20 ... Y.a E. Indies 1820. S s.p
Adenanthera aculeata Rox.
 10837 dfilicis Kth. sweet T D e c SO ... W.G N. Spain 1818. S p.1 Kth.mim.34
Ifcacia edulis IV. en.
 10838 cumanensis Kth. Cumana T O or 20 ... W.a Cumana 1822. S p.1
Acacia cumanensis W.
 10839 dominensis Dec. St. Domingo f • or 30 ... Y.a StDomin. 1818. S p.1
 10840 juliflora 7Jr. July flower f O c 30 jl ... Y Jamaica 1800. S p.1
h ifcacia falcata Dctf. Mimosa piliflora Sun.
 10841 pAllida Kth. pale Algaroba f • ec 20 ... W.o Peru 1825. S co
1284. LAGONY'CHIUM Bieb. LAGONYCHIUM. (*Lagos,* a hare, *onychion,* a little nail) *Legum. Mimbsea.* 1.
 10842 Stephanira Bieb. Stephan's tt ____ or 10 jl.au Y Persia Casp. 1816. S pi Bux. c. 48
Mimosa micrantha Vahl. /fcacia Stephani/w Bieb.
1285. ERYTHROPHLETUM Afz. RED WATER TREE. (*Erythros. red, phleos,* a water plant.) *Leg. Mim.* 1.
 10843 guineensis G. Don Guinea i • P 100 ... Pa.Y & Leone 1793. S p.1
Afeelsa grandis Hort.
1286. GAGNEBfNA Neck. GAOVEBINA. (Of no meaning.) *Leguminbsa Mimbsea.* S.
 10844 femartscina Dec. Tamarisk-like si • or 6 ... Y Mauritius 1824. S p.1 Dec. leg. 12.64. B
Acacia femarf scina Lam.
- 10845 axillaris Dec. axillary i Q w 6 ... Y Mauritius 1824. S p.1 Dec. leg. 12.64.A
Mimosa pterocarpa Lam.
1287. HiEM ATCXYLON L. LOGWOOD. (*Hahna,* blood, *xylon,* wood; color of wood.) *Leg. Cats. Cass.* 1.
 10846 campechiannum L. Campechy J D d y S O ... Y S. Amer. 1724. S r.m Cat.car.2.66
1288. COPAI'FERAL. BALSAM OF CAPEVI. (*Copaiba,* Brazilian name, *fero,* to bear.) *Leg. Cas. Cass.* 2—14.
 10847 officinalis L. officinal t CD m 20 ... W S. Amer. 1774. C s.T Jac. am. 86
 10848 guianensis Desf. Guiana j D m 90 ... W Guiana 1826. C' p.1 Mém. mu. 7.13
1289. TACHYQA^LIA Aub. TACHYGALIA. (*TachygaU,* its name in Guiana) *Leg. Cas. Cass.* 1.—3.
 10849 bfjuga Dec two-paired I Q or SO ... Y Brazil 1822. C p.1
1290. HARDWI'CKM Rox. HARDWICKIA. (*Major-Gen. Hardwick, E. I. C. Artillery.*) *Leg. Cas. Cass.* 9.
 10850 pinnata Rox. pinnate t D o r 40 ... Y E. Indies 1818. C p.1
 10851 binata Rox. binate J • or 40 ... Y E. Indies 1820. C p.1 Rox. cor. 209
1291. BA^PHJA Afz. BAPMA. (*Baphe, a dye; use)* *leguminbsa Swartxtea.* 1.
 10852 nltida Lod. shining ± D or 30 jn.n W S.Leone 1793. C s.p Bot. cab. 367
1292. TRICHI'LIAZ. TRICHILU. (*Tricha,* ternary; capsule 3-celled and 3-valved.) *Meliicea.* 5.—19.
 10853 hirtal. hairy f O or 15 jn.jl W Jamaica 1800. C p.1 Sl. jam. 2.220.1
 10854 tpondidiles Suix. Hog-plum-like t O or 20 jn.jl W Jamaica 1800. C p.1 Jac.sc. 1.103
 10855 havandnsis Jac Havannab \$ • or 20 jn.jl W Havan nan 1824. C p.1 Jac. am. 175.38
glabra L.
 10856 odor Ma B. Rep. sweet-scented f O or 15 jn.jl W W. Indies 1801. C l.p Bot rep. 637
 10857 glandulosa Sm. glandulous J • or 20 jn.jl W N. Holl. 1824. C l.p
- * 1293. JffETLIA L. BEAD TREE. (The Greek name of the manna ash.) *Meliacea.* 8.—10.
 10858 Axedardach L. Azedarch Y J or 45 jn.au B S.ya 1656. C p.1 Bot. mag. 1066
 10859 robusta Rox. robust H or 30 jn.au W.R E. Indies 1820. C l.p
 10860 sumatrensis G. Don Sumatra Lilac S • or 20 jn.au W.R E. Indies 1824. C l.p
 10862 semperv^rens W. evergreen T L or 30 jn.au B Jamaica 1656. G 8.i Bot. reg. 613
 10863 composita W. compound^vd t t or 20 jn.au W.R Tencriff. 1810. C l.p
 10864 excelsa Dec. lofty L.J or 40 jl.au W.R E. Indies 1819. C l.p
 10865 Axadirchta W. Azadirchta J • or 60 jn.au W E. Indies 1759. C q.ii Cav. dis. 7.208
1294. SANDO'RICUM Cav. SANDAL TREE. (*Santoor,* the aboriginal name.) *Melidcea.* 1.
 10866 Indicum Cav. Indian 1 • tm 40 ... W E. Indies 1820. C l.p Ru. am. 1.61
- * 1295. TURRJEV4 I* TURRSA. (*George Turraa,* professor of botany at Padua.) *Melidcea.* 2.—8.
 10867 vkrens L. green • or 20 ... W E. Indies 1820. C p.1 Sm. ic. in. 1.10
 10868 rlgida Yen. rigid J • or 20 ... Y Mauritius 1816. C p.1 Yen. ch. 48
1296. QUIV'SIAC&v. QUIVISIA. (*Bois de Quivi,* name in Isle of France.) *Meliicea.* 1.—4.
 10869 heterophylla Cav. vanous-loaved t Q c u 16 ... W la France 1822. C p.1 Cav. dis. 213
1297. SWIETE^NLJ L. MAHOGANY TREE. (*Baron van Swieten,* celeb, phys. and bot.) *Melidcea.* 2.—3.
 10870 Mahgongi L. Mahogany f O tm 80 ... R.Y W. Indies 1734. C pi Cav.dis.7.209
 10871 febrifuga Rox. febrifuge J • m 60 ... W.v E. Indies 1796. C p.1 Rox. cor. 1.17
1298. CHLOROOXYLON Dec. CHILOROXYLON. (*Chloros,* green, *tylon,* wood.) *Meliicea.* 1.
 10872 Swietinia Dec. Van Swieten's ± • tm 100 ... W E. Indies 1820. C l.p Rox. cor. 64
Swietenia Chloroxylon Rox.
- * 1299. FLINDE'RS/J R. Br. FLINPBRSIA. (*Capt. M. Flinders,* a celebrated voyager.) *Meliacqa.* 1.—*.
 10873 australis R. Br. southern £ LJ tm 60 ... W N Holl. 1623. C l.p Br. rem. 1
1300. CARAPA Aub. CABAPA. (*Carape,* its name in Guiana.) *Meliicea.* 4. —
 10874 guianensis Aub. Guiana J D o r SO ... Y Guiana 1824. C l.p Aub.gui.S87
 10875 guineensis G. Don Guinea t O ec 20 ... Y & Leone 1793. C l.p
Afzelta splendens Hort.
- 10876 moluccensis Lam. Moluccas t CZI or 20 ... Y E. Indies 1820. C l.p Ru. am.*®¹
 10877 procra Dec. tall f O or 40 ... Y W. Indies ... C l.p

EKEBE'RG/i* Spar.	EKEDEBGIA.	(C. G. Ekeberg, a Danish naturalist.)	MeUdce*. i.
MS Spar.	Cape	t L Jor 20 jLau W C. G. K. 1789. C p.l Lam il. 358	
IS ^{tr} <Jus ^S Rox.	HBYNEA.	(Dr. B. Heyne, a German botanist)	Melidcear. S.
three-paired	J or 20	W Nepal 1812. C	
JWJ80 quinquejuga Rox.	J or 20	.. W E. Indies 1816. C)p Bot mag. 1738
1303. GUAPACUM. LIGNUM MTJR THEE.	(Guatac, its name in Guiana.)	Zusophfjilee. 3.-5.	
10881 officinale /.	t r-lm 40 Jls	B W. Indies 1694. C ip Lam. il. 542	
10882 arboreum Dec.	J l-m 30 ...	B Trinidad 1816. C p.l Jac. am. 83	
Zygophyllum arboreum Jac.			
10883 verticale Or.	vertical	arior 8 — B W. Indies 1820. C p.l	
1301. ZYGOPHYLLUM L. BBAN-CAPER. (Zygos, a pair, pkyllon, a leaf; lvs. in pairs.)		Uea. 13.-18.	
10884 simplex L.	g imple t E) or jl.au Y	St Jago 1825. S	For. eg. 12
10885 cordifolium L.	heart-leaved • J cu 6 o	O C. G. H. 1774. C	
10886 Fab %o L.	Fabago ♀ A cu 4 jl.s	O.w Syria 159G. C	Lam.il. 545.1
10887 foetidum &Ar.	fetid * J pr 4 jn.au	O.T C. G. H. 1790. C	Sc. han. 17. 9
2 insuave J9. Al.	unpleasant * J pr 4 jn.au	C. G. H. 1790. C	Bot mag. 372
10888 fruticulsum Dec.	fruticulose Jl or 1 jl.au	Y N. Holl. 1820. C	
10889 maculatum //, K.	Jl pr 4 o.n	C. G. H. 1782. C	
10890 mleophi Humilium L.	small-leaved Jl or jl.au	Y C. G. H. 1816. C	
10891 Morisina L.	Morogona * J cu 3 my.s	Y C. G. H. 1712. C	
10892 fulium L.	essile-leaved m \ cu 3 jl.au	Y C. G. H. 1713. C	Bot. mag. 2184
10893 spinosum L.	spiny a. J or 1 jl.au	Y C. G. H. 182a C	Bur. af. 5.2.2
10894 coccineum L.	scarlet • J or 3 ...	S Egypt 1823. C	For. sg. 87.11
10895 album L.	white * ZZ or 2 on W	Canaries 1779. C	L. f. dec. 1.6
WB96 protitratum TAUH.	prostrate JU iA or ft jl.au	Y C. G. H. 1810. C	
in ³⁰⁵ . F A G O W J * Tou.	FAONIA.	(Mons. Fagon, a great patron of botany.)	Zygophyllum. 3.—10.
10897 cratica L.	Cretan tt. L-J cu ~ if'jn au P	Candia 1739. 8 it. Bot. mag. 241	
10898 arabica L.	Arabian tt. Jcu 2 jn.au P	Arabia 17C9. S it.	
10899 glutinosa Dec.	glutinous JU Jor 1 iny.au R	Egypt 1820. S p.l De1.aBg.2a2	
J206. RHIBULUS TVw.	CALTROPS.	(Tribu, to tear; prickly capsules.)	Zygophyllum. 7.—18.
10900 pistoides L.	Cistus-like —* IQJ or J	Y S.Amer. 1752. % ill	Bot.reg.791
10901 white	white -* O cu "1 jn.Jl	W Guinea 1826. S	
10902 earth	nearly unarmed Jc (QJ or	Y S. Europe 1596. S co Lam. il.346.1	
10903 woolly	woolly -* IQJ or J	Thibet 1820. S 8.1	
10905 maximus L.	greatest ** O pr J	Y E. Indies 1822. S.1	Bur. ley. 106.1
10906 trijugatus Nut.	three-paired -* O or J	Y Jamaica 1728. 6.1	Jac. ic. 3.462
* 1307. LA'RREJ Cav.	LARREA.	(J. A. H. de Larrea, a Spanish botanist)	Zygophyllum. 1.—a
10907 nltida Cav.	shin'ng4eaved n. J or 2 jnjl	Y & Amer. 1823. S	Cav ic. 6.559
* 1308--DICTAMNUS i.	FRAXINELLA.	(A Cretan city; grows on mountains near it)	Rutaceae. 2.
10908 Fraxinella Pert.	A or 3 my.jl P	Germany 1596. S p.l	Jac.au. 5.428
10909 albus L.	white k A or 3 my.jl W	Germany 1596. S p.l	
10909 filibus Lk.			
1309. RUTA Tou.	RUE.	(Rute, Gr., and the same in most languages.)	Rutaceae. 16.—25.
10910 Clus.	vrlngal-leaved n. J un 2 mr.au	G.v Canaries 1780. C r.m	
10911 Pen.	mountain tt. J un 2 aus.s	G.Y S. Europe 1596. C co Jac. ic. 1.76	
10912 L.	Chalco tt. J un 2 jns.s	G.Y Africa 1722. C r.m	
10913 OeO8a Dec.	heavy-scented tt. m 3 jns.s	G.Y S. Europe 1562. C co Dull. ar. 2.61	
10914 divaricata Ten.	bracted tt. — un 1) myjn	Y.a Sicily 1819. C co	
10915 angustifolia Pers.	divaricate tt. — un 2 my.jl	Y.o Italy 1820. C co	Lam.il. 345.1
10916 chalepensis Vil.	narrow-leaved tt. J un 2 jns.s	G.Y France 1722. C r.m	Bot mag. 2311
10917 macrippilla Sol.	long-leaved tt. J un 3 jl	Africa 1820. C r.m	Bot mag. 2018
10918 albiflora Hook.	white-flowered tt. J pr 2 jl.au	G.Y Nepal 1823. C r.m	Hook. ex. fl. 79
10919 c<5rica Dec.	Corsican tt. J un 1 jns.s	W Corsica 1818. C eo Hoc. mu. 59	
10920 patav'nai.	Paduan Aj ± un li jn.jl	Y.o Italy 1819. C r.m Mic. gen. 19	
10921 pubescens IT. en.	pubescent 5 A un 1) my.au	G.Y Spain 1816. C co	Lam. il. 345.2
10922 villbsa Bieb.	villous a...J un 1 jnjl	Y.o Caucasus 1818. C co	Bux. c.2.28.2
10923 lifolia L.	Flax-leaved tt. — un 1) jn s	G.Y Spain 1752. C r.m	
10924 s ^c /infifolia grandiflora B. M.	sweet-scented tt. J un 2 jns.s	Y.a Tauria 1800. C r.m	Bot. rep. 565
U0925 dahurica Dec.	Dahurian tt. J un If my.jl	Y.a Dahuria 1820. C co	Bot. mag. 2254
1310. CROWWE Sm.	CROWEA.	(J. Crowe, of Norwich, a celebrated British botanist.)	Rutaceae. 2.
10926 Julia Lo. C.	Wittow.leavcd m \ Jor 3 jl.d	P N. S. W. 1790. C s.l.p	Bot. mag. 989
broad-leaved <J Jor 3 jl	P N. S. W. 1825. C fl.p		
M ¹ : ERIOSTEMON Sm.	EHIOSTEMON. (Erion _{krood} , stemon, stamen; fringed Mamcnts.)	Rutaceae. 9	
10928 Auxilius Sm.	Box-leaved H Jor 2 my	Pk N. Holl. 1824. C p.l	
10929 calicifluius Sm.	Willow-leaved H or 3 ap.jl	R N. Holl. 1824. C s.p.l	Lin. tr. 11.86
Job's rtus CIIH.	Myoporum-like H or	R N. Holl. 1824. C fl.p	
10932 cuspisatus Citn.	obovatn H or ^ ap.jj	R N. Holl. 1824. C s.p.l	
10933 micfolius Mn. C.	cuspidate H J or 3 ap.jl	R N. Holl. 1824. C s.p.l	Bot cab. 1S47
10934 linearifolius Dec.	Heath-leaved J 3 ap.au	R N. Holl. 1824. C s.p.l	
10935 lancifolius Gae'	lincnr-leavcd J 3 aji.au	R N. Holl. 1824. C s.p.l	
10936 squameus Lab.	lanceolate J 3 ap.au	R N. Holl. 1824. C s.p.l	
scaly J 2 ap.au	R N. Holl. 1822. C s.p.l		
PIEBALIUM. (Phibalee, a myrtle appearance.)			
squamulose * L Jor 2 ap.jj	Y N. Holl. 1824. 8.I.p	Ven. mal. 102	
oldiei » J or 6 ap.jl	Y N. Holl. 1822?	C s.l.p Fl. n. h. ic.	
Lachnea-like J or 3 ap.jl	W N. Holl. 1824. C s.l.p		
tall J or 10 ap.jl	W N. Holl. 1825. C i.l.p		
linear J or 3 ap.au	W N. Holl. 1825. C s.p.l		
Willow-leaved J or 3 ap.au	W N. Holl. 1825. C fl.p		
CHOISY^ H. & B.	CHOIBYA.	(J. D. Choisy, a Genevese botanist)	Rutaceae. 6.51S
ternate Q or 6 jlau	W Mexico 1825! C 1 p u.EB.		

1314. CO'DON FT. CODON. (*Kodon*, a little bell; corolla.) *Solanacaj.* 1.
10944 Royeū* W. Royen's £ Ol cu 1 i W C.G.H. 1801. S Itl BoL rep.325
1315. GO'MPHI A Schreb. BUTTON-FLOWER. (*Gomphos*, a club; german.) *Ochnhcece.* 6.-25.
10945 zeylanica Dec. Ceylon ft D or 4 ... Y Ceylon 1823. C s.l Bur.zey.56
O'chna zeylanica Lam.
- 10946 obtusifolia Dec. obtuse-leaved ft D el 3 ... Y Jamaica 1803. C BI An. mu. 8
10917 JabottJpitaSwz. Jabotapita ft • or 5 ... Y Jamaica 1820. C BI Lam. il 472.2
10948 nftida Swx. glossy-leaved ft O el 4 ... Y Jamaica 1803. C s.l An.mu.13
10949 laevigata Lo. C. smooth-leaved ft O or 4 ... Y. 1820. C al
10930 feurifolia Swx. Laurel-leaved ft O or 5 ... Y Jamaica 1823. C s.l Dec mu. 17.15
1316. QUA'SSM W. QUASSIA. (*Quassi.a* negro slave of Surinam, first used it med.) *Stharubdceaf.* 1. — 4.
10951 amara W. bitter 2 Dm 20 jn.jl R Guiana 1790. C p.l Bot mag. 497
1317. SIMARIPBA Aub. SIM ARUBA. (*Simarouba*, its name in Guiana.) *Simarhbear.* 2.-3.
10952 excelsa Dec. lofty I Q m 40 ... Y.w Jamaica 1818. C l.p Ac. haf. 1788.8
Quftssia cxcisla Swx.
- 10953 officinalis Dec. officinal ft am 10 ... Y.w W. Indies 1789. C l.p Aub.gui.2.33
Quassia SimariUta L.
1318. SIMATBA Hil. SIMABA. (Its name in Guiana.) *Simarhbeat.* 2.-7.
10954 orinocensis/i:^2r. Orinoco ft a m 10 ... W Orinoco 1818. C Lp H &B.514
10955 guianensis Aub. Guiana ft D or 10 ... W Guiana 1826. C Lp Aub. gui. 1.153
Zwingenz amara W.
1319. HEISTETR/ii L. Bois PERDRIX. (*Lawrence Heister*, prof, bot Halstadt, d. 1758.) *Olactneat.* 1. — 3.
10956 coccinea Jac. scarlet ± D o r 20 ... S W.Indies 1822. C Lp Jac.am. 81
- * 1320. LIMO'NIA L. LIMOXIA. (*Limoun*, the Arabic name of the Citron.) *Aurantihcea.* 4. — 11.
10957 crenulata Rox. crenulate ft D or 4 ... W E. Indies 1808. C r.m Rox. cor. 1. 86
10958 parviflora B. M. small-flowered ft i | fr 6 year W China 1800. C r.m Bot mag. 2416
10959 scindens Rox. scandent fl i | or 6 jn.jl W China 1800. C r.m
10960 citrifolia TV. Orange-leaved ft L J fr 4 year W China 1800. C r.m
1321. BE'RGERI* Kon. BKRGKRA. (*C.J. Berger*, professor of botany at Kiel.) *Aurantihcea.* 3.
10961 Konfg/f Rox. Konig's ft O cu 3 jn.jl W E. Indies 1820. L pi Bot cab. 1019
10962 intcgerrima Rox. very entire-tod ft D cu 4 jn.jl W W. Indies 1823. C s.p
1322. ATALA'NT/jf Corr. ATALANTIA. (*Atalanta*, the daughter of Schceneus.) *Aurantiheete.* 1.
10963 monophylla Dec. one-leaved ft D or 4 jn.au W E. Indies 1777. C r.m Itox. cor. 1.83
Limbium monophyllum Rox.
1323. GLYCO'SMIS Corr. GLYCOSMIS. (*Glykys*, sweet, osme, smell: perfume.) *AurantiUcece.* 2.
10964 pentaphylla Corr. five-leaved £ D or 20 jn.jl W E. Indies 1790. C r.m Rox. cor. 1. 64
Limbnia pontaphylla Retz.
10965 arborea Corr. tree ID or 20 my.au W E. Indies 1796. C r.m Rox. cor. 1.85
1324. CLAUS&NA Brm. CLAUSENA. (Not explained.) *Aurantidcece.* 1—2.
10966 pentaphylla Drc. five-leaved f D o r 20 jn.au W Coroman. 1800. C r.m
Limbnia pentaphylla Herb. Lam. not Box.
1325. FEROWI! Corr. FERONIA. (Goddess of forests.) *Aurantiaccw.* 1. — 2.
10967 elephantum Corr. elephant apple H Q o r 4 ... W E. Indies 1804. C r.m Rox. cor. 2.141
1326. MURRA'YI! Kon. MURRAYA. (*J. A. Murray*, a Swede, prof. bot. Gottingen.) *Aurantiacea?* Q.
10968 exotica L. exotic * D or 8 au.s W E. Indies 1771. C Itl Bot. reg. 434
10969 paniculata WaL paniced ft D or 8 jl W E. Indies 1823. C r.m Hook. ex.fl.79
1327. COO'K/4 Sona WAMPEE TREE. (*Copt. Cook*, celeb, Eng. circumnavigator.) *Aurantihcea.* 1.—2.
10970 punctata Relz. dotted £ • fr 15 ... W China 1795. C Jtl Jac. sc. 1.101
1328. G/ERTNETRJ Rox. GARTNERA. (J. Gartner, M.D., F.R.&, a celebrated bot) *MalpighiUcece.* 2.
10971 racembsa Rox. clustered L D or 15 mr.ap W.Y E. Indies 1796. C p.l Bot rep. 600
Hf ptage mandablbta Gae.
- 10972 obtusifolia ROT. obtuse-leaved 4. D o r 20 ... W China 1810. C p.l
Hiptage obtusifolia Dec.
1329. BUNCHO'SIA J. BUNCHOSIA. (*Bunchos*, coffee, A rob.; similar seeds.) *Malpighiucta?* U. — 16.
10973 glandul&sa Dec. glandulous ft D or 10 ... Y W. Indies 1804. C p.l Cav. dis. 8.239.2
Malpighi glandulSsa Cav.
10974 glandulifera //.* B. gland-bearing ft D or 10 mr.my Y W. Indies 1806. C p.l Jac. ic. 3.469
Malpighia glandulifera Jac.
10975 mtdia Dec. intermediate ft • or 10 mr.my Y W. Indies 1790. C p.l
Malpighi mtdia H. K.
10976 polytachya Dec. many-spiked ft D or 10 mr.my Y W. Indies 1806. C p.l Bot. rep. 604
Malpighia polystachya Andr.
10977 tuberculata Dec. tuberculate ft • or 6 my.au Y Caracas 1806. C p.l Jac. sc. 1.104
Malpighia tuberculata Jac.
10978 argentea Dec. silvery ftd)or 10 jl.au Y Caracas 1810. C p.l Jac.fr. 83
10979 nitida Dec. shining • D o r 10 jl.au R St.Domin.1800. C p.l Cav.dis.&239.1
Maipghia nitida L.
10980 paniculata Dec. paniced ft D o r 10 my.au P Jamaica 1820. C p.l
10981 cumane'nsis G. Don. Cuinana ft O or 10 s Y Cumana 182a C p.l
10982 odorata Dec. scented ft • or 10 myau Y Carthage 1806. C p.l Jac. am. 177. **
10983 canescens Dec. canescnt ± D o r 20 JLau Y W.Indies 1742 C p.l
Malpighia canescens H. K.
1330. MONOTROPA L. YELLOW BIRD'S-NEST. (*Monos, onc, trepo*, to turn; fls. one way.) *Monotrhpce.* T
10984 uniflbra Mx. one-flowered ^ A cu i W N. Amer. 1824. S s.p Hook. ex.
10985 Hypatips L underpines 5 A cu *jn.jl W Britain woods. S s.p Eng. bot 69
1331. DION; E\4 L. DioNiEA. (One of the names of Venus.) *Brosrhcect.* 1.
10986 Musripula /.. Venus's Flytrap j£ tAJ cu i jl.au W Carolina 1768. S bog Bot mag. 785

- | | | | |
|---------------------------------------|--|---|---|
| 1332. GARJTOA Rox. | GARVGA. | (Garugo, its Telinga name.) | Terbinthacea. 1. |
| 10987 pinnate Rox. | mng & Ueaved | I Q o r 20 ... Y E. Indies | 1808. C p.l Rox.cor.3.208 |
| 1333. BOS WE'LL/* | Rox. OLIBANUMTREB. | (Dr. John Boswell, of Edinburgh.) | Terebinthaceae. 2. - 3. |
| 10988 serrata Stack. | serrated | t Om 20 ... Fa.Y E. Indies | 1820. C Lp As. res. 9,377. ic. |
| 10989 glabra Box. | smooth | J Q o r 30 ... Pa.Y Coroman. | 1823. C Lp Itox. cor. & 207 |
| 1334. ICICA Aub. | ICICA. | (Its name in Guiana.) | Terebinthaceae. 4. — 15. |
| 10990 heterophylla Dec. | various-Jeaved | ■ CD or 20 ... W | Guiana 1826. C l.p Aub. gui. 1. 133 |
| iflmryis hctrophylla W. | | | |
| 10991 guttiferens Aub. | Guiana | f D o r 30 .. W | Maranh. 1823. C l.p Aub. gui. 1.131 |
| iflmryis guianensis W. | | | |
| 10992 enca-ndra Aub. | nine-stamened | t o r 20 .. W | Maranh. 1822. C l.p Aub. gui. 1. 134 |
| 10993 Tacamahaca Kth. | Tacamahac | i Q o r 20 .. W | Trinidad 1819. C l.p |
| 1335. KA'LMM L. | KALMIA. | (P. Kalm, professor at Abo, in Sweden.) | Bhodorrhce. 6. |
| 10999 latifolia L. | broad-leaved | ■ or 8 my.jl R | N. Amer. 1734. L s.p Hot mag. 175 |
| 10995 angustifolia L. | narrow-leaved | ■ or 5 my.jl | N. Amer. ■ L s.p Bot. mag. 331 |
| Stribra | red-flowered | ■ or 5 my.jl | N. Amer. ■ L s.p Bot. cab. 502 |
| 3 fulris variegatis | variegated-lvd | ■ or 5 my.jl | N. Amer. ■ L s.p |
| 4 minima | least | ■ or 5 my.jl | N. Amer. ■ L s.p |
| 5 nana | dwarf | ■ or 5 my.jl | N. Amer. ■ L s.p |
| 6 rbsa | rose-colored | ■ or 5 my.jl | N. Amer. ■ L s.p |
| 7 phmila | dwarf | ■ or 5 my.jl | N. Amer. ■ L s.p |
| 8 ovita Ph. | ovate | ■ or 5 my.jn | N. Amer. ■ L s.p |
| 10996 cunctata Mx. | cuneate-leaved | ■ or 5 my.jl | W.R N. Amer. 1820. L s.p |
| 10997 glauca L. | glaucous | ■ or 2 ap.my | P N. Amer. 17<7. L s.p Bot. mag. 177 |
| 10998 rosmarinifolia PA. | Rosemary-lvd | ■ or 2 ap.my | R N. Amer. 1812. L s.p |
| 10999 hirsuta Walt. | hairy | —Jor | R N. Amer. 1786. L s.p Bot mag. 138 |
| 1336. ZEDUM L. | LABRADOR TEA. | (Leden, Greek name of the Cistus.) | Rhodorecete. 4. |
| 11000 palustre L. | marsh | ■ or 2 ap.my | W Europe 1762. L s.p Bot. cab. 560 |
| 2 decumbens | decumbent | ■ or 1 ap.my | W Huds'. Bay 1762. L s.p |
| 11001 latifolium // K. | broad-leaved | ■ or 3 ap.my | W N. Amer. 1763. L s.p Bet. cab. 584 |
| 11002 canadense Lod. | Canadian | ■ or 11 jn | W Canada ... L s.p Bot. cab. 1049 |
| 11003 greenlandia Beta. | Greenland | ■ or 2 tip.my | W Ireland bogs. L s.p |
| *1337. AMMYRSPNE Ph. | AMMVISINE. {Ammos, | sand, myrsine, a myrtle j habitat) | Rhodorecea. J. |
| 11004 Auxifilia Ph. | Box-icved | —L or 1 ap.my | W N. Amer. 1736. L s.p Bot reg. 531 |
| Xedum duxifolium | L. LeiohprUum ihmifolium Spr. | | |
| 133a AHODO'RA L. | RHODORA. | (il*o-fow, a rose; color of flower*) | Rhodordceee. 1. |
| 11005 canadense L. | Canadian A | or 3 ap.my | P N. Amer. 1767. L p.l Bot mag. 474 |
| *1339. iZHODED'NDKON L. | RHODODENDRON. (Rhodon, a rose, dendron, a tree.) | RAodorilcete. 23.—24. | |
| 11006 maximum Z. | largest | ■ spl SO jn.au | Pk N. Amer. 1736. h s.p Bot mag. 951 |
| 11007 album Ph. | white | ■ or 20 jii.au | W N. Amer. 1811. L s.p |
| 11008 purpCircum PA. | purple | ■ spl 25 jn.au | P N. Amer. ... L s.p |
| 11009 ponticum L. | Pontic | ■ spl 12 my.jn | P Gibraltar 1763. L 6p Bot mag. 650 |
| 1 album | white | ■ or 8 my.jn | W gardens — L s.p |
| 2 angulifolium | narrow-leaved | ■ or 6 my.jn | P gardens — L s.p |
| 3 ■ fi ■ | Cassine-leaved | ■ or 6 my.jn | P gardens ... It S.p |
| 4 ceruleascens | ceruleaceous | ■ or 10 my.jn | B gardens — L s.p |
| 5 contortum | twisted-petaled | ■ or 8 my.jn | P gardens — L s.p |
| 6 fibre pleno | double-flowered | ■ or 12 my.jn | P gardens ... L s.p |
| 7 fblis argenteif | silver-leaved | ■ or 6 my.jn | P gardens ... L s.p |
| 8 fblis aftreis | golden-leaved | ■ or 6 my.jn | P gardens ... L s.p |
| 9 frondibsum | leafy | ■ or 5 my.jn | P gardens ... L s.p |
| 10 intermedium | intermediate | ■ or 12 my.jn | P gardens ... L s.p |
| 11 macrophyllum | long-leaved | ■ or 12 my.jn | P gardens ... L s.p |
| 12 ovatum | ovate/eaved | ■ or 6 my.jn | P gardens — L s.p |
| 13 rbuscum | rose-flowered | ■ or 10 my.jn | R gardens — L s.p |
| 14 rubescens | rubescens | ■ or 10 my.jn | R gardens — L s.p |
| 15 alicitiflum | Willow-leaved | ■ or a my.jn | P gardens — L s.p |
| 16 nepalense | Nepal | ■ or 5 ap.my | P Nepal 1817. L s.p |
| 11010 cinnabmeum TTa. | Cinnamon | ■ spl 20 jn | P Nepal 1820. L ps Den. br. 168 |
| 11011 obtusum Wat. | obtuse | ■ spl 12 myjn | P Gibraltar 1763. L s.p Ex. bot 6 |
| 11012 arboreum Sm. | tree | ■ spl 120 ... | S Nepal 1820. L s.p |
| 2 album | white | ■ spl 120 — | W Nepal 1818. L s.p |
| 11013 punctatum Andr. | | ■ or jn.au | Pk N. Amer. 1786. L s.p Bot rep. 36 |
| 2 majus | | ■ or 6 jn.au | Pk N. Amer. 1786. L s.p Bot reg. 37 |
| 11014 azaleoides Desf. | larger dat u Ulvd 41 | ■ spl 3 jn.au | Pk 1820. L s.p Bot. rep. 379 |
| 2 odoratum | Azalea-like | ■ or 3 jn.au | Pk 1820. L s.p |
| 11015 hibridum B. R. | sweet-scented | ■ or 3 jn.au | Pk 1820. L s.p |
| H016 dat Ticum L. | Herbert's hybrid* | ■ 3 jn.au | Pk 1820. L s.p |
| 2 atrovlrcns | Daurian A | or 3 mr.d | P Siberia' 1780. L s.p Bot reg. 195 |
| 3 alticum Fis. | dark .green | or 3 f.ap | P Siberia' 1780. L s.p Bot. mag. 636 |
| JJ017 myrtiflbum Lod. | Altaic | or 3 mr.ap | P Altai 1810. L s.p Bot reg. 194. |
| 11018 ferrugineum L. | Myrtle-leaved | ■ spl 12 my.jn | P Gibraltar 1763. L s.p |
| 11019 bicajum L. | rusty/eaved | ■ spl 12 my.jl | S Switzerl. 1752. L s.p Bot. cab. 908 |
| | hairy.leaved | ■ or 1 my.jl | S Switzerl. 1656. L s.p Bot. cab. 65 |
| | vanegatcd-lvd | ■ or 1 my.jl | 1800. L s.p Bot mag. 1853 |
| | yellow.flwd | ML or \ jn.jl | Y Siberia' 1796. L s.p Par. lon. 80 |
| 11021 caucasicum Pall | Caucasian | ■ au | P Caucasus 1803. L s.p Bot mag. 1145 |
| 11022 campanulatum B | D r | ■ ap.my | Nepal 1817. L s.p |
| 11023anth>bgdn. D. | | ■ ap.my | Nepal 1820. L s.p |
| aronviticum Wat. | | | |
| 11024 scutsum D. Don | setose | ■ d or | V Nepal 1825. L s.p |
| 11025 catawbiense Mx. | Catawba | ■ 4t or | P N Amer. 1809. L s.p Bot. mag. 1671 |
| JJO16 Catogium Howt.. | Latesby's | ■ 4k or | P N. Amer. 1810. L s.p |
| 11027 camchtamicum Pall | Kamchatka | ■ 4t or | P Kamtsch. 1802. L s.p Pal ros. 1.33 |
| 11028 Chamaecestus L. | Ground Cistus | ■ tt or | P.A Austria 1786. C s.p Bot. mag. 488 |
| 11029 lapponicum Wahl. | Lapland | ■ L or | R Lapland 1810. L s.p |
| J340. L | PIGAA. | (Epi, upon, grtia, earth; the stems on ground.) | Rhodoriacea. 1. — 2. |
| | creep'ng | IU pr | Jjl.a u W N. Amer. 1736. L s.p Bot reg. 201 |

• 1341. ANDROMEDIF L.	ANDROMEDA.	(The virgin <i>Andromeda</i> .)	<i>Ericat.</i> 27.—44.
11031 <i>tetragona</i> L.	four-angled	tt. — J or	W Lapland 1810. L s.p
11032 <i>Aypnoldia</i> L.	Moss-like	tt. — W. E Inland	L s.p IH. dan. 10
11033 <i>calyculata</i> L.	small-calyxed	or 1} f.ap	W. N. Amer. 1748. L s.p Pal ros. 2.72.1
1 ventricosa	ventricose	or	Kussia 1748. L s.p Dot mag. 1286
2 latifolia	broad-leaved	or	Newfoun. 1748. L s.p HoL cab. 530
3 nana	dwarf	cu	L B.p Uot.cab. 862
11034 <i>angusti folia</i> PA.	narrow-leaved	or	N. Amer. 1748. L s.p
11035 <i>crispa</i> Dc.jj.	curled	or	N. Amer. 1824. L s.p
11036 <i>foliolata</i> L.	Poliom-leaved	or	— 1» B.j*
1 latifolia	or	my.s	N. Amer. — L s.f*
2 media	middle, orb. ^{Wid. Rovetary}	or	my.8
3 subulata	a.w.-leaved	tt.	my.s
4 minima	smallest	tt.	ap.my Pk
5 oleifMia	Olive-leaved	tt.	ap.my Pk
11037 <i>glaucophylla</i> Lk.	glaucous-lvd	tt.	my.s Pk
J.Dolifolia/3 latifolia lvd.			N Amer. ... L s.p Bot cab. 546
11038 <i>ro>marinifflia</i> Ph.	Hobmerry-lvd	tt. or 1	mys Pk
11039 <i>jamaicensis</i> Swx.	Jamaica	• • O r 6	N Amer. ... L e.p PaLros.2.70. a
11040 <i>marianna</i> L.	Maryland	tt. or 2	my.jl W
1 ovkis	ovaUeaved	tt. or 2	my.il W
2 oblonga	oblong-leaved	tt. or 2	my.jl W
11041 <i>speciosa</i> Mx.	showy	tt. or 3	jns W
1 nltida	Bhiring.Icaved	tt. or 3	jns W
2 pulvrellnta	duby/<iv.a'	tt. or 3	jns W
SglAca Wat.	glaucous	tt. or	— — L s.p Den. br. 26
11042 dealbata <i>Linell.</i>	whitened	tt. or	2 ap.my* Pk
11043 fasciculata AUBB	bundled	tt. or	— — L s.p Bot reg. 1010 _m
11044 aalicifolia Wat.	Willow-leaved	ttQ or 20	jn W
11045 Auxifolia Lam.	Box-leaved	ttl or 2	Pk
11046 coriacea H. K.	thick-feavef	tt. or 2	jn.au Pk
nitida Mx. Iticida Lam.			N Amer. 1765. L Bp Bot. mag. 970
2 rikbra. Ld.	red-flowered	* or 3	jn.au R
11047 acuminate //, K.	acuminate	tt. or 3	au W
Iticida Jac. fl. <i>populifl.</i> la Lam.	reticulata Watt.	tt. or 2	Jauxina Mx.
11048 <i>Catesba</i> Ti Walt,	Catesby's	or' 2	jn.jl W
spinulifl. Ph.			N Amer. 1793. Sk s.p Bot mag. 1955
11049 spickta Walt.	spiked	tt. or	jn W
11050 axillaris H. K.	axil-flowering	tt. or	my.au W
2 longiflilia	long-leaved	tt. or	my.au W
Walteri W.			N Amer. ... Sk ■ Bot mag. 2357
11051 canescens Lo. C.	cancsrent	tt. or	— W
11052 racemusa L.	branchy	fra	N Amer. ... L s.p
paniculata Walt.			N Amer. 1736. L s.p
2 latifolia	broad-leaved	tt. or 3	jl.au W
Sstrcta	strict	tt. or 4	jl.au W
11053 jai>6nica Than.	Japan	tt. or 3	... W
11054 rubjeinosia Pert.	rubiginous	tt. or 10	... W
pubescens Pair.			W. Indies ... C s.p
11055 arboreal*.	Sorrel tree	tt. or 40	jl.i W
11056 florblinda Ph.	bumilc-flwd	tt. or 3	my.jn W
11057 avuhCAa JVaU.	oval-leaved	i or 20	Ncii.il 1825. L. D. As.rea.13.391.ic.
11058 pilulifera/#.	pellet-bearing	tt. or 3	jn.jl W
			— ... L s.p
1342. LYON ^o Ni4 Nut.	LVOMA.	(J. Lyon, an American collector of plants.)	<i>Erfcete.</i> 6.
11059 ferruginea Nut.	ferruginous	tt or 3	jn.jl W
Andromeda ferruginea Wait.			N Amer. 1784. L s.p Ven. mal. 80
11060 rigida Nut.	rigid	tt or 20	ap.my W
Andromeda ferruginea //, K.			K Amer. 1774. L Bp Bot cab. 430
11061 paniculata Nut.	pamceled	tt. or 3	my.jn W
Andromeda paniculata L.			N Amer. 1748. L s.p Den. br. 57
11062 multiflora Wat.	many-flowered	tt. or	— W
11063 capra'tbia Wat.	tendnl-leaved	tt. or	— W
leafy	tt. or	my.jn W	N Amer. 1812. F. sp Dcn.br. 128
11064 frondosa Nut.			N Amer. 1806. F. sp Dcn.br. 127
Andrmcdia frondosaPn.			
1343. ENKIA'NTHUS IMU.	ENKIANTHIB	(Egkuos, pregnant, anthos, flower.)	<i>Ericete.</i> 2.-3.
11065 quinqueflora Lou.	five-flowered	■ L j el 3 fs	Pk China 1812. C s.l.p Bot mag. 161!)
11066 reticulata Lindl.	netted	■ L Jcl 3 ja.f	Pk China 1822. C p.l Bot. rcg. 885
1344. GAULTHE'R/4 L.	GAULTHERIA. (M. Gauthier, M.D. a Fr. writer on maple sugar.)	<i>Ertccca.</i> 4.—18.	
11067 procumbent L.	procumbent	pr Jjl.fl	W N. Amer. 1762. Sksp Bot. rep. 116
11068 SAnZoZ Ph.	Shallon	tt or 4	my W N. Amer. Jl'26. L s.p Ph. am. ic.
LOWHragrans J. Don	fragrant	tt v-J or 6	... P.R. Nepal 1824. I. s.p As.res.13.297.ic.
11070 antipoda Forst.	antipodal	H J o r 6	... W N. Zeal. 1820. C b.p
* 1345. ARBUTUS L.	STRAWBERRY TREE. (<i>Arboise</i> , austere bush, <i>Celt.</i> , rough fruit.)	<i>Ertceee.</i> 10.—17.	
11071 tA>edoE.	Unedo	or 10 s.d	W Ireland ir.ro. S co Eng. bot 2377.
2rdber	rcA-flowered	tt or 10 s.d	Pk *L co
3 plenus	double-Jfouuroa	tt or 5	s.d. W.o L co
4 schizopetalus	entire-leaved	11 or 6	s.d. Pk L co
5 crispus	cut-petalcd	tt or 7	s.d. W. L co
6 crispus	curled	tt or 8	s.d. W. L co
7 alicifilius	Willow-leaved	tt or 6	s.d. W. L co
11072 canarfensis Lam.	Canary	tt l or 8	my.jn W.G Canaries 1796. L co Bot mag. 1677
11073 andrachno'ides Lk.	Andrachnc-lk	tt or 8	fmy W.G L s.p Bot. reg. 619
h^bnda D. It.			
11074 serratiflilia Nois.	serrated-leaved	tt \ or 6	fmr W.6 F. ■ ■ ■ ■ ■
11075 Andr&chnc L.	Andrachnc	tt or 10	nir.ap W.G Levant 1724. G p.f Bot. rcg. 113
11076 auriflha L.	Laurel-leaved	t or 20	... W Mexico 1825. L B.p
11077 denBilla Kih.	dense-flowered	J or 20	... W Mexico)82. L s.p
11078 phillyraiflBa Pro.	Phillyreal-levd	tt-i Jor i	... W Peru 1812. L s.p
11079 tomentoflilia Ph.	wooWy-leaved	tt or 6	... W N. Amer. 1826. L p.1
	dwarf	tt J or 4	W Magellan 1220. L s.p

- | | | | |
|--|---|---|---------------------------------------|
| 1357. MICOW,* R.&P. MICONIA. | (<i>D. Mieon</i> , M.D., a Spanish botanist.) | <i>Melattomheea.</i> | 15.— |
| 11144 <i>grandifolia</i> D. Don large-leaved | ... W | Trinidad | 1890. C I.p. |
| <i>Melastoma grandifolia</i> //, C. | | | |
| 11145 <i>rubescens</i> D. Don rubescent | - W | S. Amer. | 1818. C Lp |
| 11146 <i>tenuifolia</i> D. Don fine-leaved | - W | S. Amer. | 1818. C Lp |
| 11147 <i>angustata</i> D. Don narrow | - W | Trinidad | 1820. C Lp |
| 11148 <i>impedita</i> Kins. <i>D. Don aca&ile-leaved</i> | - W | W. Indies | 1822. C Lp Bon. mon. 29 |
| 11149 <i>decussata</i> D. Don decussate | - W | Guiana | 1818. C Lp Aub.gui. 1.156 |
| 11150 <i>semicrenata</i> D. Don half-crenated | - W | Guadalou. | 1817. C Lp |
| 11151 <i>purpureascens</i> D. Don purpurascant | - W | Jamaica | 1817. C Lp |
| 11152 <i>longifolia</i> G. Don long-leaved | - W | Guiana | 1817. C Lp |
| 11153 <i>prasina</i> G. Don lock-green | - W | Guiana | 1817. C Lp |
| <i>Melastoma lsvigia</i> Aub. | | | |
| 11154 <i>trinervia</i> D. Don three-nerved | - W | Jamaica | 1795. C Lp |
| <i>Mel&toma trinervis</i> W. | | | |
| 21155 <i>tetraandra</i> D. Don turandrous | S1Q or 2 | ... | Jamaica 1815. C Lp |
| <i>Melastoma tetsindra</i> Swz. | | | |
| 11156 <i>Acinodendron</i> ZJ.Mw* <i>Acinodendron</i> • or | 6 | P | Jamaica 1804. C Lp PL ic. 142.2 |
| <i>Melastoma Aoinodendron</i> L. | | | |
| 11157 <i>purpureascens</i> > J.Mw* <i>purpurascant</i> | Si or 8 | P | Guiana 1804. C Lp Aub. gui. 1.154 |
| <i>Melastoma purpura</i> IV. <i>purpureascens</i> Aub. | | | |
| 11158 <i>losigata</i> D. Don smooth | S1Q or 6 | W.o | a Amer. 1815. C s.p Botreg.663 |
| <i>Melastoma loevigata</i> L. | | | |
| 1356. CHITO^NI A D. Don CHITONIA. (<i>Chiton</i> , a coat of mail; calyx scaly.)^ <i>Melastomhca.</i> 5.— | | | |
| 11159 <i>macrophylla</i> D. Don long-leaved | W | Trinidad | 1820. C Lp |
| 11160 <i>pyramidalis</i> G. Don pyramidal | W | Trinidad | 1817. C Lp |
| <i>Melastoma pyramidalis</i> Desf. | | | |
| 11161 <i>fflipcans</i> D. Don white-leaved | « Q or 10 | W | Mexico 1815. C Lp |
| <i>Mel&toma albicans</i> Swz. | | | |
| 11162 <i>Fothergilla</i> I. Don <i>Fothergill's</i> ± • or 20 | ... | P | a Amer. 1815. C s.p Aub. gui. 1.175 |
| <i>Melastoma Fothergilla</i> Rich. <i>Fothergilla mirabilis</i> Aub. | | | |
| 11163 <i>Tamonia</i> G. Don <i>Tamonia</i> i or 12 | ... | P | W. Indies 1815. C s.p |
| <i>Melastoma Tamhnia</i> Swz. <i>Swartzia</i> Spr. | | | |
| 1359. CLIDI?MI* D. Don CLIDEMIA. (<i>Cidemii</i> , an ancient Greek botanist) | | | <i>Melastomdeete.</i> 12.— |
| 11164 <i>legans</i> D. Don elegant | Si or J^jl.au | W | Maranh. 1822. C s.l Aub. gui. 1.167 |
| <i>Melastoma elegans</i> Aub. | | | |
| 11165 <i>spicata</i> D. Don spiked | ... | | Guiana 1823. C p.l Aub. gui. 1.165 |
| 11166 <i>agrestis</i> D. Don field | ... | | Guiana 1822. C p.l Aub. guL 1.166 |
| 11167 <i>cernua</i> D. Don drooping | ... | | Peru 1820. C p.l |
| <i>Melastoma cernua</i> Bonp. | | | |
| 11168 <i>microphylla</i> D. Don small-leaved | Si CD or 1 jl.au | W | a Amer. 1817. C p.l |
| 11169 <i>holosericea</i> D. Don velvety | A Q or 4 jl.au | W | Brazil 1820. C p.l Pink. am. 252.2 |
| 11170 <i>alifta</i> G. Don winged | s1^ or 4 jl.au | W | Maranh. 1819. C p.l Aub. gui. 1.158 |
| <i>Melastoma alata</i> Aub. | | | |
| 11171 <i>nfvea</i> D. Don snow-white-/wfl • or 4 jl.au | W | S. Amer. 182a. C p.l Bon. mon. 44 | |
| 11172 <i>aggregata</i> 2. D. Don aggregate | SI or 1f jl au | W | Peru 1820. C p.l |
| 11173 <i>velutina</i> D. Don velvety | S1Q or 2 jl.au | W | W. Indies 1816. C p.l |
| <i>Melastoma vcluAna</i> W. | | | |
| 11174 <i>hrlta</i> D. Don hairy | H Q or 6 s.d | P | Jamaica 1740. C s.p Pluk.al. 264.1 |
| <i>Melastoma hrlta</i> L. | | | |
| 11175 <i>rdhra</i> G. Don red | • si or 6 my.jn | P | Guiana 1793. C s.p Aub. gui. 1.161 |
| <i>Melastoma rtbra</i> L. | | | |
| 1360. PLERCPMA D. Don PLEROMA. (<i>Pleroma</i> , fulness; the cells of capsule.) | | | <i>Melastotndcea.</i> 5.— |
| 11176 <i>heteromalla</i> D. Don <i>heleromallous</i> • o r 4 jls. | P | Brazil | 1819. C Lp Bot.reg.644 |
| <i>Melastoma heteromalla</i> B. R. | | | |
| 11177 <i>granulosa</i> D. Don granulous | il • or 10 aus. | P | Brazil 1819. C p.l Bot.reg.G71 |
| <i>Mdastoma granulosa</i> Dcso. 7?hexia | | | |
| 11178 <i>vimfnea</i> D. Don twiggy | Fontanesii Kth. il CJ or 6 jn.au | P | Brazil 1821. C s.p Bot.reg.664 |
| <i>JRhexia vimfnea</i> B. R. | | | |
| 11179 <i>holosericea</i> D. Don velvety | si • or 10 jl | B | Brazil 1816. C p.l Bot.rcg.32S |
| <i>Ahcxia holosericea</i> Bonp. | | | |
| 11180 <i>glomerata</i> G. Don headed | * Q or 1jl | B | W. Indies 1818. C p.l Bot.cab.334 |
| <i>Rhtxia. glomerata</i> Lod. | | | |
| 1361. TOCO^CA Aub. | TOCOCA. | (<i>TOCOCO</i> , its name in Guiana.) | <i>Melastomhcee.</i> 2. |
| 11181 <i>AubRtii</i> D. Don Aublet's | * • or 3 au.s | Pa.K | Guiana 1826. C p.l Aub. gui. 1.174 |
| <i>guianensis</i> Aub. | | | |
| 11182 <i>TnnMaD</i> . <i>Don</i> woolly | • or 3 au.s | Yf | Trinidad 1817. C p.l |
| <i>Melastoma lanata</i> H. C. | | | |
| * 1362. ACIO^riS D. Don ACIOTIB. (<i>Akis</i> , a point, <i>ous</i> , an ear; petals.) | | | <i>Melastomhcete.</i> 8. |
| 11183 <i>discolor</i> D. Don various-colored | * • or 1 jl.s | W.R | Trinidad 1816. C p.l PI. ic.48 |
| <i>Melastoma discolor</i> L. | | | |
| 1U84 <i>aquatica</i> D. Don aquatic | .. pr f jn.jl | W.R | a Amer. 1793. C bog Aub. gui. 1.169 |
| <i>JZnexia aquatica</i> L. | | | |
| <i>Melastoma aquatica</i> Aub. | | | |
| 1363. MERIA^N^ Swz. | JAMAICA ROSE. | (M. S. Merian, authoress of a work on insects.) | <i>Melastomheex.</i> 3. |
| 11185 <i>leucantha</i> Swz. | white-flowered | il • or 6 ... W | Jamaica 1825. C p.l |
| 11186 <i>purpurea</i> Swz. | purple | i l ^ or 6 ... P | Jamaica 1825. C p.l |
| 11187 <i>coccinea</i> Swz. | scarlet | • or 6 ... S | Jamaica 1818. C p.l |
| % 1364. PETALON^MA Swz. | PETALOMA | (<i>Petalon</i> , petal, <i>lama</i> , fringe; petals in calyx teeth.) | <i>Melattomhce*</i> 9. |
| U1188 <i>myrtilloides</i> Swz. | Bilberry-like | • • < * 10 ... W.Y | W. Indies 1823. C p.l Si.jam. 187. 3 |
| (11189 <i>Mourvi</i> Swz. | Mourin | * Q or 3 jl.s | W. Guiana 1817. C p.l Aub. gui. 1.180 |
| <i>Mouriri guianensis</i> Aub. | | | |
| 1365. ACISANTHE^RA J. ACISANTHIERA. | (<i>Akis</i> , point, <i>anthera</i> , anther; anthers pointed.) | <i>Melastoma</i> | |
| 11190 <i>quadrtka</i> Pcrs. | squared | <i>siricu</i> 3 | Jamaica 1804. C p.l Br. jam. 22.! |
| <i>Ahxia. Acisanthiera</i> L. | | | |
| 1366. DA^IS L. | DAIB. | (Derivation unknown.) | <i>ThymelaP*</i> . 1.— 3. |
| 11191 <i>cotinifolia</i> Z. | Cotinus-leaved | si. I or 10 in il | W.o C G.H. 1776. C s.l Bot mag. m |

1367. BUCPDA L. OLIVE BARK TREE, (<i>Sous</i> , an ox; fruit like an ox'a horn) <i>Combrethfa</i> . 1.—4.			
11192 &ceras L.	Ox's horn	j p e c 25 au.s	Y.w Jamaica 1793. C l.p Lam.il 356
,1368. SAMRDA L.	SAMVDA.	(Greek name of the birch; resemblance.)	<i>Samydetv.</i> 8.—15.
11193 glaberrata Sivz.	smooth	• Dor 6 jl.i	W W. Indies 1800. C l.s
JJ194 villtaa Sura.	villous	• • or fi JLi	W W. Indies ltfll. C l.s
11195 decurrens Spr.	decurrent	• • or 5 ...	G Brazil 1820. C s.l
11196 pubescens L.	pubescent	• Dor 4 my.au	W W. Indies 1793. C l.p Jac. am. 132
JJ197 serrulata L.	serrulate	■ 3 jl	W W. Indies 1723. C s.p Jac. c. 2.17.1
11198 raea H.K.	rose-colored	■ 4 jn.jl	Pk W. Indies 1793. C s.p Bot.mag. 550
11199 nitida L.	glossy	■ ■ 4	W-o W. indies 1793. C l.p Br.jam. 23.3
11200 macrophylla TV.	long-leaved	■ ■ ■ ■	E. Indies 1820. C s.l
viridiflora Pair.			
iiIS ¹ C fts EA «?W I	CASEARIA	(«Co*caniM,cooperatorin Hortus Malabaricus8.)	<i>Samjdeof.</i> 6.—23.
11201 ramiflora Vahl	branchflwd	» Q o r 4 ...	Y.o Guiana 1824. C s.l Aub. gui. 1.127
Irouetia guianensis Aub..			
11202 parviflora JF.	sm.Jl-flowered	« • or 6 ...	Y.o S. Amer. ISI& Martiniq. 1827. C s.l Sl.jam. 2.211.2
11203 parviflora IV.	small-leaved	« D o r 6 ...	Y.a Jac. am. 85
Decandra Jac.	wood	*CZ)or 8 ...	Jamaica 1823. C s.l
J!Xf Hirsuta Swz.	hairy	i i Q o r 8 ...	Jamaica 1825. C 8.1
J!2f serrulata Swz.	serrulate	i i Q o r 6 ...	Jamaica 1818. C s.l
iiIS ^{0*} AQUILA'RIA L. AQUILARIA. (<i>Aquila</i> , an eagle; called Boisd'Aigle by Fr.colonists.)	<i>Samjdea</i> .	1.—2.	
11207 malaccensis Lam.	Malacca	*C j o r 6 ...	W.o Malacca 1823. C l.p Cav. dis. 7.224
ovata Cav.			
iii? ¹ CERATOPETALUM Sm. CERATOPETALUM. (<i>Keras</i> , a horn, <i>petaton</i> , a petal.) <i>CunoniUcear</i> . 1.			
11208 gummiferum Sm.	gum-bearing	£ L_J or 50 ...	Y N. Holl. 1820. C s.l Sm. n. h. 1.3
iJ? ² - BARWI'N/J4 Rud.	DARWINIA. (<i>Dr. Darwin</i> , author of the Botanic Garden, a poem.)	<i>Myrtacea</i> . 2.	
J1209 fasciculito Bud.	fascicled	*i J o r 29 my.jl ...	N. Holl. 1820. C s.p Lin. tr. 11.22
H210 foxifolia Cun.	Yew-leaved	• i J o r 29 my.jl ...	N. Holl. 1824. C s.p.l

CLASS X. ORDER 2

DECA'NDRIA DIGY'NIA.

,1373. ROYE ^o Ni! L.			
11211 flucida L.			
J1212 villbsa L.			
}13paffens7%im.			
JJ14 »iyrtiflia IV.cn.	Myrtle-leaved	■ Jor	W C.G. H. 1800. C Up
J1215 glabra Z.	smooth	■ Jor	4 s W C. G. H. 1731. C l.p Com. h. 1.155
H216 pubescens W. en.	pubescent	■ Jor	4 jl.au W o CG.H. 1752. C l.p Bot.reg.500
110...cuneSita Poir.			
J27 hirsuta L.	hairy-leaved	■ ■ 7 jl.au	W CG.H. 1752. C p.1 Lara.il 370.2
J28 angustiflia TV.	narrow-leaved	■ ■ 7 jl.au	W C.C. H. 1789. C p.1
Jgiglrlfella W.ai.	broad-leaved	■ ■ 5 jn.jl	W CG.H. 1816. C s.p
"221 polyandra W.	ambiguous	■ ■ 6 jn.jl	W CG.H. 1815. C p.1 Ven.mal.17
	polyandrous	■ ■ J or 6	W CG.H. 1774. C p.1
lisX ⁴ , T1*1A'NTHEMAZ. TRIANTHEMA. (<i>Treis</i> , three, ant, <i>Jicmon</i> , fl.; flowers by threes.) <i>Portuldeear</i> . 3.-12.			
11201 mon6\$y ^{na} '-	monogynous	- * O w 2 jl.au	G Jamaica 1710. S co Dec pi. 109
lioG;fdecandraZb	decandrous	- * ED w lijl.au	G India 1762. S co Bur. in. 51.3
'''attobcorrutta/fox.	obcordate	- * O cu 1 jl.au	G EL Indies 1820. S co
1 POSI ⁵ SCLEK A'NTHUS L. KNAWEL. (<i>Skleros</i> , hard, <i>anthos</i> , flower; dry, juiceless calyx) <i>Paronychiilar</i> . 3—5.			
J{**5annus2.,	annual	O un jl.au	G Britain s.a. fl. S co Eng. bot 351
i iSC ⁷ perennis L.	perennial	- * A, un i a.us	G Britain sa. hea, D co Eng. bot. 352
"287 pungent A. A*.	pungent	O cu ij1.au	G N. Holl. 1826. S s.l
PACHYNE ⁸ M A B. Br. PACHYNEMA. (<i>Pachys</i> , thick, <i>nema</i> , a filament; stamens.) <i>Dillenidcea</i> ? 1.			
comptenhtum li.Br. fat-branched tL _J or 1).			N. Holl. 1825. C s.l Del. ess. 1.73
Cy*TN L. CUNOXIA. (J. C. <i>Cuno</i> , of Amsterdam, in 1750.) <i>Cunoniheca</i> . 1.—2 ?			
capensis L.	Cape	!1 J or 20 au	W CG.H. 1816. C co Botreg.826
L. HYDRANGEA. (<i>Hydor</i> , water, <i>aggos</i> , a vessel; seed-vessel.) <i>Saxifragef</i> 6.—8.			
11251 arborescent A or 6 il.au	W.a Virginia 1736. L p.1 Bot. mag. 437		
heat-leaved tt or 8 jl.au	W.G. Carolina 1806. L p.1 Den. br. 42		
enovf-white-leavedSH or 5 jl.au	W.o Carolina 1786 L p.1 Den. br. 43		
no ^o heteromallous	■ or 4 ...	W.o Nepal 1821. C	
U235 mornensis AM. Oak-leaved	■ or 4 jn.s	W.o Florida 1803. C	Bot. mug. 975
Hortensia opuloldcs Lam.	■ or 3 ap.s	Pk China 1788. C	Bot. mag. 438
alternifl / ^ alternatclwd	A cu i ap.my Y	Britain w.sli.p. 1) bog Eng. bot 54	
opposiUfohuini.	opposite-leaved	i ap.my Y	Britain w.sh.p. 1) bog Eng. bot 490
nepaleiisc D. Don	Népal	{ ap.my Y	Nepal 1820. D bog
GOULDEN SANHAREE (Chrysos, gold, spfen, spleen; med. <i>qunU</i> Saxfrügea)? 3.			
ligulata Wai.	ligulate	£ ^ or i ap.jn	W.R Nepal 1821. D p.1 Bot cab. 747
Mogasea cihiata Haw.			
crassifolia L.	thick-leaved	jf A or 1 mr.myP	Siberia
	heart-leaved	£ ^ or 1 mr.my P	Siberia
		■ ■ ■ ■	1/9. D s.1

11242 media Haw.	intermediate		1 apjn P	Britain	moun. D co		
11243 hirsute L.	hirsute		1 myjn F	Ireland	ir.mou. I) s1	Eng. bot. 2322	
11244 Geum	Geum		1 jnj! W	Ireland	ir.mou. I) s1	Eng. bot. 1561	
11245 dentata /.*.	toothed		myjn W	Britain	moun. D co		
11246 crenata /tow.	crenated		myjn W	 1796. D co		
11247 spheroidea Haw.	spheroidal		myjn F	 1800. D co		
11248 Sumburusa	shady London pride	£ A or	apjn F	Britain	moun. D s1	Eng. bot. 663	
11249 punctata L.			myjn W	Siberia	1699. D s1	M.h.3.9.17	
11250 polita Z.*.	polished	£ A or	myjn W	Ireland	moun. D co		
11251 cuneifolia L.	wedge-leaved	£ A or	nj! W.a	Switzcr.	1768. I) s1		
2 davurica D. Don	Dahurian	£ A or	W	Siberia	1809. D ps		
11252 hifrida Lk.	hybrid	£ A or	W	Piedmont	1810. D co		
11253 stellaris L.	starry		W	Britain	al.riv. D s1		
2 Schielcheri D. Don	Schielcher's	£ A or	W	Switzerl.	1819. D p.s	Eng. bot 167	
11254 dissimilis O. Don	dissimilar	£ A or	B1jnj!	BenNcvls al.rlv.	D s1		
11255/eucanthemifolia Atc.	Stock-leaved	A or f	Jn	W	N. Amer. 1812. D ps		
11256/sarmentowii L.	twiggy	JU Al or	1 jnj!	W.R	China 1771. D s1	Bot mag. 92	
11257 cuscuthf6Tmi» Lod.	Dodder.formed	JU AJ pr	i jnj!	W	China 1815. D s1	Bot. cab. 186	
11258 critaaFh.	gnawed		W.Y	N. Amer. 1812. D s1			
2 lurstria D. Don	hairy		W.Y	N. Amer. 1800. D s.p			
11259 rotundifolia L.	round-leaved		W.R	Austria 1596. D s1	Bot mag. 424		
2 repanda D. Don	repand		W	Caucasus 1800. D s.p	Stcr. sax. 17.5		
11260 granulata L.	grain-rooted		W	Britain mepa. D il	Eng. bot 500		
2 plena			W		D s1		
11261 bulbifera L.	bulb-bearing		W	S. Europe 1819. D ps	Col. cc. 1.317		
11262			W	Siberia 1802. D al	Stcr. sax. 25		
11263 cfrnual	Siberian		W	Scotland sc. alp. I) 61	Eng. but. 661		
11264 rivularia Z.	drooping		W	Scotland sc. alp. D s1	Eng. bot 2275		
11265 Acicracea L.	brook		W	Levant 1752. S 81	Fl. gr. 379		
11266 HirculusL.	Ivy-leaved		Y	England tu. bo. I) s1	Eng. bot. 1009		
WilTI ilagelarlis Ster.	Hirculus		Y	Greenl. 1819. I) s.p	Ster. sax. 6		
11268 flizodesZ.	rod-like		W	Britain al.riv. D s1			
11269 aizooneZ.	A izoon-like		W	Britain ... D s1	Eng. bot. 39		
11270 teWib.	autumnal		Y				
11271 bronchiMisL.	smooth		W	Canary 1826. D s1			
11272 sispralaL.	bronchial		W	Crea Siberia 1819. D s.p	Gm. si. 4.65.8		
11273 tenella Wul.	rough		W	Switzcr. 1752. D 81	Jac. au. 5.31		
H274 gryoldesZr.	delicate rough		jnau	W	Cariuthia 1819. 13 s.p	Jac. c. 3.17	
11275 Aieracifi.haJH/.	Bryum-like		ijnjl	Switzerl. 1752.	Jac. m. 2.5.1		
11276 semipubescent Swt.	Hawkweed-lvd	£ A or	myjn	Hungary 1789. D s1	W. & K. 1.18		
11277 pcamtlyvanica L.	semi-pubescent	£ A or	myjn	Lu N. Amer. 1800. I) s.p			
	PennsVlvania	£ A or	1) myjn	G.Y N. Amer. 1732. D sf	Di.cL253.328		
	smooth	£ A or	2) apjn	G.Y N. Amer. 1732. I) s.p			
11278 virginensis Mx.	Virginian	£ Al or	i myjl	G.v N. Amer. 1790. I) 81			
11279 congesta Haw.	heat-ed-Jlw	£ AI or	e myjl	W N. Amer. 1812. D s1	Bot. mag. 1664		
nivMis Ph.			W				
11280 nivMis L.	snowy		W	Britain scalp. D 81	Eng. bot. 440		
11281 Cotyledon L.	Cotyledon		W.n	Alp. Eur. 15% D s1	Fl. dan. 241		
11282 lingulata BeL	tongue-leaved		W	Switzerl. 1800. D s1	Lap. sax. 11		
2 inertia D. Don	intermediate		W	Carniola 1800. D s1	Ster. sax. 1. a		
11283 crustacea Vest	crustaceous-rrfgf	£ A or		Switzerl. 1800. U s1	Ster. sax. 1. b		
11284 intermedia Haw.	intermediate	£ A or		W.R			
11285 inteta IV. em.	untouched	■ A or	1 myjl	W Tyrol	H.ber.2.75		
recta Lap.							
S minor Haw.	smaller	£ A or	1 myjn	W Alp. Eur. ... D 81			
3 parviflora Haw.	small-flowered	£ A or	1 myjn	W Alp. Eur. ... D s1			
11286 Atzbon Mur.	Aizoon	£ A or	1 myjl	W Alp. Eur. 1731. D 61	Jac. au. 5.438		
11287 mutataL.	changed	£ A or	ft jnj!	L.Y Switzerl. 1779. D s1	Bot. mag. 351		
11288 oppositifolia L.	opposite-leaved	£ A or	ft mr.ap	P Britain al. roc. D s1	Eng. bot. 9		
11289 binora All.	t*ro-fift& red	JL A or	ft myjn	P Switzerl. 1820. D ps	Al. ped. 21.1		
11290 retusa Gou. .	rectuse	JL A or	ft myjn	P Piedmont 1826. D ps	Lap. sax. 18		
11291 cassia L.	grey	£ A or	i myjn	P.y Switzerl. 1752. D 81	Bot. cab. 421		
11292 diapensioidez Bel.	Diapensia-like	£ A or	i apjn	W Switzerl. 1825. D s.p	A I. ped. app.3		
11293 aretioides Lap.	Arctia-like	£ A or	ft myjl	Y Switzerl. 1826. D p.s	Lap. sax. 13		
11294 Burscridna L.	Burscr's	£ A or	1 mrjn	Crea Carniola 1826. D p.s	Ster. sax. 10.1		
11295 androsacea L.	Androsacc/vtf	£ A or	i myjn	W Austria 1792. D 81	Jac. au. 4.389		
11296 sedolodes L.	Sedum-like	£ A or	i myjn	Y Europe 1820. D p.s	Jac. m. 2. 21.22		
11297 tenera Suter	tender	£ A or	ft myjn	Crea Switzerl. 1819. D p.s	Al. ped. 61.2		
11298 geranioides Z.	Crane's-bill-lk	£ A or	1 jnj!	W Pyrenees 1770. D si	Lap. sax. 43		
11299	plashy	Q> or	ft myjn	Tauria 1817. S ps	Bieb.ros.2.73		
11300 pedatifida Ehrh.	pedatifid	£ A or	ft myjn	Scotland scalp. D s1	Eng. bot. 2278		
11301 quinquifida Haw.	five-cleft	£ A or	i myjn	Britain moun. D s1			
11302 visebsa Haw.	clammy		i myjn	Pyrenees JHS. D s1	Lap. sax. 40		
			ft myjn	Spain 1804. D s1	Bot. mag. 1651		
			1 myjn	... 1825. D si			
11506 ajugifolia L.	Bugle-leaved		1 jnj!	Pyrenees 1770. D 81	Lap. sax. 31		
11307 affinia D. Don	kindred		ft myjn	— — D s1			
laevis Bonn							
11308 hifra Donn	hairy	£ A or	1 jn	Scotland scalp. D s1	Eng. bot. 2291		
11309 platypetala Sm.	broad-petaled	£ A or	1 jn	Scotland alroc. D s1	Eng. bot. 227C		
11310 decipiens Ehrh.	deceptive	£ A or	ft myjn	Wales wal.p. D 81	Eng. bot. 455		
palmata Sm.							
11311 Sternbergii W.cn.	Steinberg's	£ A	1 my	W Germany ... D s1	Ster. sax. 24		
11312 incurvifolia D. Don	incurved-lvd	£ A	\ myjn	Ireland rocks. D s1			
11313 denudata D. Don	stripped	£ A	i myjn	Scotland cmo. D s1			
11314 pulchilla D. Don	pretty	£ A	ft myjn	Germany 1818. D s1			
11315 tridentata D. Don	three-toothed	£ A	ft myjn	W Pyrenees ... D s1	Lap. sax. 37, #		
11316 moschata Wul.	musky	£ A	ft myjn	L.Y Pyrenees ... D s1			
musco*des Ster.	Moss-like	£ A	ft myjn	Pa.Y Switzerl. 1819. D s1	Lap. SAX. 35		
11317 muscoles Wul.							
moschata With.							
JJ3J8 tricuspidata Rib.	tricuspidate	£ A	ft myjn	Greenl. 1824. D s1	Fl. dan. 976		
11919 pygmaea Haw.	pygmy	£ A	ft myjn	W.Y Scotland scalp. D s1	Eng. bot. S2J*		
moschata&i?			O w	\ ap.my W Britain walls. S s1	Eng. bot 501		
11320 tridaactylites L.	three-fingered						

11321 contraversa <i>Ster.</i>	contrary-turned	O or	4	myjn	W	S. Europe	1824.	S	s.l	Ster. sax. 16	
U322 petn' aZ.	rock	O or	* ap.my	W	Norway	1752.	8	s.l	Jac. ic. 1.81		
11323 adscendens <i>Vahl</i>	ascending	£ ^ or	i	ray	W	Pyrenees	1752.	D	s.l	Ster. sax. lft 1,8	
aqu.itica <i>Lap.</i>											
11324 luiceofata <i>Haw.</i>	upcar-bracted	f ^ or	4	myjn	W	Europe	1800.	D	s.l		
2 obtusa <i>Haw.</i>	obtuse	£ A or	i	myjn	AV	Europe	1820.	D	s.l		
11325 leptophylla <i>Pen.</i>	fine-leaved	£ A or	4	myjn	W	Wales	moun.	D	s.l		
2 angustifida/D/Jon	narrow-cleft	£ A or	4	myjn	W	Wales	moun.	D	s.l		
11326 condensata <i>Gm.</i>	dense	£ A or	4	myjn	VV	Scotland	c. mo.	D	s.l	Fl.bad.2.3	
11327 ketevirens i). Uojt	lively green	£ A or	4	yj	w	Scotland	c. mo.	D	e o		
2 viscusa <i>Hort.</i>	Moss-like	£ A or	4	ap.jn	W	Britain	d.roc.	D	co	Eng. bot 45*	
3 angustifiliu <i>Hort.</i>	clammy	£ A or	4	ap.jn	w	Scotland	moun.	D	co		
4 muscosa <i>D. Dun</i>	mossy	£ A or	4	myjn	W	Scotland	moun.	D	co		
5 pulchella <i>D. Don</i>	pretty	£ A or	4	myjn	W	Scotland	inoun.	D	co		
11329 cespitosa <i>L.</i>	turfy	£ A or	4	myjn	C	Wales	w.alp.	D	si	Eng.bot.794	
11330 exarata <i>VUL.</i>	engraved	£ A or	4	myjn	W	S. Europe	1S18.	D	BI	VH.del.4.45	
11331 granlandia <i>H. K.</i>	Greenland	£ A or	4	j	W	Pyrenees	1732.	D	81	Lap. sax. 19	
11332 elongella <i>Sm.</i>	longish-stalked	£ A or	1	ap.my	W	Scotland	sc. alp.	D	s.l	Eng. bot 2277	
1381. TIARELLA <i>L.</i>	TIARELLA.	{Tiara, a Persian diadem;				mitred capsule.)	Saxifrdge.	3.—J.			
11333 cordifolia <i>L.</i>	heart-leaved	£ A or	* ap.my	W	N. Amer.	1731.	D	s.p	Hot. mag. 1589		
11334 Menzisii <i>Ph.</i>	Mcnzies's	£ A or	1	ap.my	W	N. Amer.	1812.	1)	s.p		
11335 polyphylla <i>D. Don</i>	many-leaved	£ A or	1	ap.my	W	Nepal	1U20.	D	p.l		
* 1382. MITE'LLA X.	MITELLA.	(Dimin. of mitra, a mitre;				(capsule.)	...&*&**&*	*			
11336 diphylla <i>L.</i>	two-leaved	£ Apr	4	ap.my	W	N. Amer.	1731.	D	p.l	Bot.	
11337 cordifolia <i>Lam.</i>	heart-leaved	5 Apr	4	ap.my	W	N. Amer.	1812.	D	lam.il.	373.3	
11338 niidaL.	naked	£ Apr	4	jn.au	W	N. Amer.	1758.	D	p.l	Lam. IL 373.2	
" reniformis <i>Lam.</i>											
11339 prostrata <i>Ms.</i>	prostrate	— or	4	ap.my	W	N. Amer.	1818.	D	p.l		
1383. ASTILBE <i>Ham.</i>	ASTILBB.	(A intensive, stilbe, brightness.)				Saxifrdge.	1.—2,				
11340 decandra <i>D. Don</i>	decandrous	■ A or	2	myjn	W	Carolina	1812.	D	s.p	Ven.mal.54	
Tiarella biaternata <i>Ven.</i>											
•1384. GYPSO'PHILA <i>L.</i>	Struthium <i>L.</i>										
11341 Struthium <i>L.</i>	peaked										
JJ342 fastigiatä <i>L.</i>	hill	2 jl.au	W	Podolia	1821.	S	p.l				
{W43cdinii* <■	sand	14jl.au	W	Hungary	1801.	S	p.l	W. & K. 41			
J1344 arenkna <i>Mur.</i>	clammy	14 jnjl	W	Levant	1773.	S	p.l	Mur. 3			
J345 ten'iu <i>Poir.</i>	delicate	1 il.au	st	Europe	1816.	S	p.l				
JJ347 altissima <i>L.</i>	tallest	14jl.au	W	Siberia	1759.	S	p.l	Gm.si.4.60			
J1348 grandiflora <i>Poir.</i>	great-flowered	2 jl.au	F	Spain	1800.	S	p.l				
}) «PerfoliataL.	perfoliate	2 jl.au	W	Iberia	1732.	S	p.l	Di. el. 276			
11350 sabulata <i>Stev.</i>	sandy	2 jl.au	W	'lauria	1817.	S	p.l				
„ icorzonneridYui Delf.	perfoliata Bieb.	14jlau									
{351 acutifolia <i>Fa.</i>	acute-leaved	3 jl.au	W.G	Siberia	1820.	S	co				
11352 paniculata <i>L.</i>	pamcled	4 jnjl	W	Siberia	1759.	S	p.l	Jac. au. 5. ap. 1			
J1353 adscendens <i>Jae.</i>	ascending	2 jl.au	W	S. Europe	IWK	S	p.l	Jac.vin.2.138			
J1354 glifica <fw.	glaucous	W	Caucasus	1822.	S	co					
J1355 Steveni <i>FIB.</i>	grassy	3	1 jl.au	R	Greece	1810.	S	p.l			
J1356 Steveni <i>FIB.</i>	elegant	1 jn.s	W.pk	Crimea	1828.	S	co	Seh. rao.21			
Jjagrtensi.	Steven's	2 jl.au	W	Iberia	1822.	S	co				
J1359 cretica 5m.	creeping	1 jl. ⁹	St	Siberia	1774.	S	p.l	Bot mag. 1448			
J1360 Cretaria Bieb.	Cretan	1 jl.au	W	Crete	1810.	S	p.l	FLgr.384			
J1361 fine-leived		1 jl.au	R	Caucasus	1824.	S	p.l				
J1362 prostrata <i>L.</i>	doulifful	1 my.s	W		1815.	S	p.l				
} «3 muralia <i>L.</i>	trailing	1 Jls	R	Siberia	1759.	S	p.l	Bot mag. 1281			
11364 serdina <i>JV.</i>	wall	*jn.o	F	Germany	1739.	S	s.l	Lam. IL 375.1			
iiHiS glomerata <i>Pall.</i>	Ute-flowering	a.u.s	W	Europe	1818.	S	p.l				
11366 A'xifraga <i>L.</i>	glomerate	1 jl.au	Pa.R	Tauria	1818.	S	p.l				
H367 rigida X.	Saxifrage	1 jl.au	Pk	Germany	1774.	S	p.l	Ex. bot S. 90			
	rigid	4 jii.au	Pk	France	1769.	S	B.1				
* 1385. SAPONA'XRIA <i>L.</i>	SOAPWOBT.	(Sapo)	mucilaginous sap.)			Caryopftillece.	10.—17.				
II68 VaccariaZ..	Cow-herb	O or	2 jl.au	Pk	Germany	1596.	S	s.l			
11309 perfoliata <i>Rot.</i>	perfoliate				KIndies	1800.	S	co			
11370 officinalis <i>L.</i>	olicimal				England	hed.	D	co	Eng. bot 1060		
2 pltna	double						D	co			
11371 orymoldes <i>L.</i>	Basil-like										
JJ72 glutinosum <i>Bieb.</i>	clammy				France	1768.			Bot. mag. 154		
\J7i plurigenes <i>L.</i>	stretching	1 jl.au			Caucasus	1817.	■	IF	Bieb. cen. 2.66		
\J74 orientalis <i>L.</i>	eastern	1 jn.au			Levant	1680.	■		Jac. vin. 2.109		
J1375 crspifusa <i>Dec.</i>	turfy	1 jl.au			Levant	1712.	S	sp	Di. el. 167.204		
J1376 ICetea <i>L.</i>	yellow	4 jn.au			Pyrenees	1824.	D	s.p			
HJ77 6	Daisy-leaved	1 JUu			Switzerl.	1804.	D	s.p	Al пед. 23.1		
					Italy	1825.	D	b.p	Boc.mu.62.1		
■ ■ ■ DIA'NTHUS <i>L.</i>	PINK. (DKW, divine, anhot, fl.; magnificence, fragrance.)										
J1378 prostrata <i>L.</i>	urolierqualiferous n or	i jl.au	Pk	Eneland	Espana	SE	pf	Eng. bot. 956			
H'2rl ciminiijit! » L>	smal\flowered				S. Europe	1771.	S	p.l			
11380 Armeria <i>L.</i>	Armeria				England	grapa.	S	p.l	Eng. bot 317		
11381 armcriodes <i>Raf.</i>	Armeria-like				New Jersey	1826.	S	co			
11382 pubescens <i>Sm.</i>	pubescent	O or			Greece	1820.	S	s.l	Fl.gr. 397		
H'383 Pseud-Armeria <i>Bieb.</i>	False Armeria	£			Crimea	1820.	S	p.i	Bot mag. 2288		
11384 discolor <i>Sims</i>	various-colored				Caucasus	1803.	S	s.l	Bot mag. 1168		
11385 barbatus <i>L.</i>	bearded	SweetwUiam			Germania	1573.	S	r.m	Bot mag. 207		
11386 dentatus <i>Ft*.</i>	toothed				Siberia	1826.	S	p.l			
iiS aggregatus <i>Poir.</i>	aggregate					1817.	S	B.1			
T'588 latifolius <i>IV.</i>	broad-leaved						C	s.l	Sw. 6. gar. 2		
1) S? J*p6nicus <i>Thun.</i>	Japanese						.1823.	C	p.l	Thun.Jap.83	
cephalotes <i>Ser.</i>	headed							Caucasus	1822.	C	p.l
^1 capitatus <i>Dec.</i>	headed										
atrorubeni <i>Bieb.</i>											

11392 polymorphus <i>Bieb.</i>	multiform	£ A or	1 jn.o	R	Crimea	1823.	C p.l
2 dibtimi <i>B. Kit.</i>	day.flowering	£ A or	1 Jn.o	R	Hungary	1816.	C p.l
11393 Balbii <i>Ser.</i>	Balbii's	£ A P*	I jl	R	Genoa	1817.	C s.l
glaucocephalus <i>Horn.</i>							
11394 ferrugineous <i>L.</i>	rusty	£ Q* or	I I jl.s	Ur	Italy	1750.	S p.l Mil. ic. 1.81.1
11395 Carthusianorum <i>L.</i>	Carthusians'	if ^ or	JJ jl.au	R	Germany	1573.	C 81 Loe.pr.7
11396 atrorubens <i>All.</i>	dark-red	£ A or	1 JU	C	Italy	1802.	C Bl Jac. ic. 3.467
11397 arbifascia <i>B. II.</i>	little tree	<L Jpr	Ijlj.au	R	China	1824.	C p.l Bot.reg.103G
11398 arboreus <i>L.</i>	tree	n.-	li jn.au	Pk	Greece	1820.	C s.l Bot.cab.459
11399 fruticosus <i>L.</i>	shrubby	■	li jjn-s	Pk	Greece	1815.	C r.m To. it. 1.9
11400 suffruticosus <i>TV.</i>	half-shrubby	■	li jjnl	Pk	Siberia	1804.	C p.l
11401 carolinianus <i>Walt.</i>	Carolina	£ AJ or	I jn.s	P	N. Amcr.	1811.	C r.m
11402 rupicola <i>Bieb.</i>	rock-inhabiting	£ A or	I jjnl	11	Italy	1820.	C 81 Fl.nap.39
Bisigniuit Ten.							
11403 caryophylliids <i>Schult.</i>	Clove-like	■ ■ ■	I	R	1817.	C s.l
11404 asper <i>W.</i>	rough-stated	■ ■ ■	I	Pk	SwitzerL	1822.	C 81
scaler <i>Sent.</i>							
11405 coinnus <i>IV.ickK</i>	hill	■ ■ ■	fjl.e	W	Hungary	1800.	C 81 Par. Ion. 62
11406 scrutis <i>Lap.</i>	serrated	■ ■ ■	I jl.s	Pk	Pyrenees	1827.	C s.l
11407 campistris /%£.	field	■ ■ ■	I jlau	W.R	Tauria	1815.	C s.l Rot. mag. 1876
11408 alpestris <i>Balb.</i>	alpine	■ ■ ■	ij.ijl	R	Ebrope	1817.	C 81 Bal.mis.7.1
11409 niftidus <i>IV.Sjk.</i>	shining	■ ■ ■	I jl.au	K	Carpath.	1822.	C 81
11410 diffusus <i>Sib.</i>	diffuse	■ ■ ■	Ujl.au	I	Cyprus	1820.	C 81
11411 attenuates <i>Sm.</i>	attenuated	■ ■ ■	§ jjn.jl	II	Spain	1822.	C 81 Brotph.2.73
11412 hirtus /#.	hairy	■ ■ ■	I jl.au	U	Trance	1821.	C 81
11413 guttatus <i>Bieb.</i>	spotted	■ ■ ■	I jl.8	R	Caucasus	1816.	C s.l
11414 pallidiflorus & r.	pale-flowered	■ ■ ■	I jjnl	P	Siberia	1817.	C s.l
pál lens <i>Bieb.</i>							
11415 vrsicolor /%£.	changable-cld	£ ■ H	■ ■ ■	RY	Russia	1823.	C s.l
11416 pratins <i>Bieb.</i>	meadow	£ ■ H	I jl.3	W.v	Crimea	1820.	C 61
ochroleucus <i>Lie.</i>							
11417 chinensis /.	China	£ <3 or	1 jls	R	China	1713.	S r.m Bot mag. 28
11418 montanus <i>Bieb.</i>	mountain	£ A or	* jn.s	R	Caucasus	1803.	C 81 Bot. mag. 1162
11419 Ciryophyllus <i>L.</i>	Clove	■ A or	2 jn.au	F	England walls.	C r.m Eng. bot. 214	
2 flore pleno	double Carnation	£ A or	2 jn.au	C	England	-.	C r.m Bot. mag. 39
3 fructicosus	shrubby Carnation	£ A or	3 jn.au	C	England	...	C r.m
4 imbricatus	imbricated Wheat-car	£ A or	li jn.au	F	England	...	C r.m Bot. mag. 1662
11420 glaucophyllus i./C.	glaucous-lvd	£ A or	I	R	1827.	L co
11421 sylvestris <i>Jac.</i>	wild	£ A or	I JSjl	R	S. Europe	1732.	C 81 Bot. mag. 1740
virginicus <i>Sims</i>							
11422 longicatilis <i>Ten.</i>	long-stcmmed	£ A or	I jjnl	W	Italy	1820.	C 81
U42J suavefolcs Spr.	sweet-smelling	£ A ■	3nj	W	1820.	C s.l
■ ■ ■	Poirt's	£ A or	I jn.jl	P	1816.	C s.l
2 fibre pltno	double-flwd	£ A or	I	P	Greece	1820.	C s.l
11425 Liposchitztinia <i>Ser.</i>	Liboschitz's	£ A or	I Ijnjl	W	Tauria	1817.	C s.l
11426 monadelphous <i>Ven.</i>	monadelphous	£ A or	I jjnl	W.Pk	Levant	...	C s.l Ven.cels39
procumbens <i>Pers.</i>							
11427 sylvaticus <i>Hoppe</i>	wood	■ A or	Iijn.s	R	Ratibus	1815.	S p.l
11428 pomeridikus <i>L.</i>	afternoon	■ A or	I jl.jl	Y	Levant	1804.	C s.l
11429 icetoictuluB <i>W.</i>	finc-prtalcd	■ A or	Ijjl	W	Caucasus	IK14.	C s.l Par. Ion. 57 Bot. mag. 1739
11431 divaricatus <i>Urv.</i>	pungent	■ A or	I au.o	Pk	Spain	1781.	C s.l
11432 bicolor/&&	divuncate	■ A or	I jl.jl	P	Greece	1822.	C s.l
11433 furcXtus <i>Balb.</i>	two-colored	■ A or	I jl.jl	Pk	Tauria	1816.	C s.l
11434 virginicus <i>L.</i>	forked	■ A or	I jl.jl	P.*	Piedmont	1619.	C s.l Bal.mis.7.2
pCingens <i>Poir.</i> rupestris <i>L.</i>	true virgin	■ A or	I jl.jl	R	Montpel.	1816.	C s.l Di. el. 385
11435 dclto^ds /.	deltoid	£ A or	I jn.o	F	Britain gra. pa.	C 8.1	Eng. bot 61
114% glaucus <i>L.</i>	glaucous-/avai	£ A or	I jn.o	W	Britain gra. pa.	C 6.1	Di. el. 29a 348
11437 Ayssopifolius <i>Hort.</i>	Hyssop-like	f Apr	ft jn.o	Pk	Europe	J810.	S co
11438 crcnatus <i>Thun.</i>	crenated	£ ^A or	I au	F C. G. II.	1817.	C s.l	Bot reg. 256
11439 marginatus <i>Poir.</i>	bordered	£ A or	I jjnl	W	Europe	1820.	C s.l
11440 rigidus <i>Bieb.</i>	rigid	£ A or	I	R	Casp. Sea	1802.	C s.l
11441 clavatus <i>Spr.</i>	clavate	£ A or	I jn.o	Pa.Pk	C s.l
11442 sulavis <i>W.</i>	sweet	£ A or	I jn.o	F	Britain rocks.	C 8.1	Eng. bot. 62
11443 caeViis <i>Sm.</i>	grey	£ A or	* jnjl	R	Austria	1759.	C s.l Bot mag. 1205
11444 alpinusL.	alpine	£ A or	I jjnl	R	S. Europe	1820.	C s.l
11W5 glacialis <i>Hae</i>	icy	£ A or	I'M				
neglectus <i>Lok</i>							
11446 tiner <i>Balb.</i>	tender	£ A or	I	R	Alp. Eur.	1817.	C s.l Bal.mis.7.3
Hornemajin* <i>Ser.</i>	Ilornemann's	£ A or	I	R	Italy	...	C 81
ture-Mus <i>Horn.</i>							
11448 Sternb^rgii <i>Sib.</i>	Sternbrcg'a	■ A or	I jjnl	R	C s.l	
11449 petrus'us VV.&K.	rock	£ A or	I jl.au	W	Hungary	1804.	C s.l W.&K.3.222
" uribus majoribus	Dee. larger.flwd	£ A or	I jjnl	Pk	1804.	C s.l Bot mag. 1204
11450 Petrus'us W. M.	Iberian	£ A or	I jjnl	P	Iberia	1817.	C al
Willdcnbvii Lk.							
11451 ruthenicus <i>loem.</i>	Russian	£ A or	I jjnl	P	Russia	1816.	C s.l
11452 gallicus <i>Pers.</i>	French	£ A or	I jjnau	P	S. France	...	C s.l
11453 albens H. K.	whitish	■ A or	f au	W	C. G. H.	1787.	C p.l Bot. reg. 256
11454 plumanus <i>L.</i>	feathered	£ A or	Ijn.au	W.p	Caucasus	1629.	C s.l
moachatus Deaf, d	>ius Horn,						
11455 hortensis <i>Schr.</i>	garden	£ A or	I jl.au	St	Hungary	1805.	C r.m
11456 sax&tis <i>Pers.</i>	Rock	£ A or	I jjnl	W	& Europe	1816.	C s.l
11457 squarrosus <i>Balb.</i>	squarerose	£ A or	I jn.au	W	Tauria	1817.	C p.l Bieb. cen. 1.33
11458 Mussini Horn.	Mussini's	£ A or	I jjn.au	W	Caucasus	1823.	C p.l
11459 caedsicus <i>Sims</i>	Caucasian	■ A or	I jn.s	P	Caucasus	1803.	C p.l
11460 fragrans Art.	fragrant	£ A or	I jjn	W	Austria	1804.	C 8.1 Bot. mag. 795
11461 punctatus <i>Spr.</i>	dotted	£ A or	I jjn	Pa.Ii	C r.m Bot. mag. 2067	
11462 neritinus W. ft K.	J&tc-Jtowersng	£ A or	I jl.s	P	Hungary	1804.	C r.m Hot. cab. 896 ^
11463 arenarius L.	sand	£ A or	t my	P	Europe	C r.m
11464 prostrata <i>Jac.</i>	prostrate	■ A or	* jjnl	R • C. G. H.	1821.	C sl W.&K.2J2J	
11465 limbriatus <i>Bieb.</i>	fringed	£ A or	I jjn.au	Li	Iberia	1815.	C sl Bot mag. 2036
orion&lia <i>Suns</i>							C sl Jac.sc. 3.271
							C s.l Bot mag. 100

J25 P'um'flus Spr.	feathery	* A or	Jftjls	W.Li M. Bald.	... C s.l
J. monspessulanus L.	Montpôlicr	£ A or	1 jlau	R Montpcl.	1764. C p.l
superb		£ A or	2 jl.s	VV Europe	1516. C 81
JjM Fischer's	Fischer's	£ A or	1 jn.jl	II Russia	1820. C si Bot. mag. 897
JJ*70 pulchellus Ehfh.	pretty	ik A or	1 jnjl	W.R Siberia	1827. C up

CLASS X. ORDER 3.

DECA'NDRIA TRIGY'NIA.

CAMPION.		(A name adopted from Pliny.)		<i>CaryopJifflcte.</i> 1.	
H471 bacciferJ.,	berry-bearing	i ^ w	1 jn.jl	W England	hed. D co E>g. bot 1577
*1388. SILE^NEZ.	CATCHFLY. (<i>Sialuti</i> , saliva; viscid frothy moisture of stalks)			<i>Caryophlllew.</i> 158.—227.	
11472 acalis L.	stemless	^ pr	ft jn.au	Pk Britain sc. alp. D p.l	Eng. bot. 1181
2 alba	w\ t-flower ing	£ pr	ftjn.au	W Britain sc. alp D s.p	
3 frscapa	seamless	£ pr	ft jn.au	R Switzerland 1819. D s.p	Al ped. 79, 2
11473 pumilio 6/lrfm	dwarf	^ pr	ft jn.au	P Germany 1822. D co	Stur.deu. 1. U
11474 fimbrikta Sims	fringed fluid	& pr	2ft" my.au	W Caucasus 1803. D s.l	Bot mag 90S
11475 lacera Sims	torn	3 pr	Iftmy.au	W Caucasus 1818. S co	Bot mag. 2255
111^« CUCuinalM lacerus Bieb.					
JHE ?tel!^ita w_*	•tw-towni	■ pr	1 jn.au	W N. Amer. 1696. D co	Bot. mag. 1107
J177 inflMa A'm	inflated	■ cu	1 mys	W Britain co. fl. D co	Eng. bot. 164
Jit^Tu*WolR Sbhrank	narrow-lvd	or	1 jn.jl	W Europe 1817. I) co	Fl. nap. 37
{ 479 ovSt Ph.	ovate	or	1 myjl	W N. Amer. 1820. D co	
?W> maritima D<r.	sea	cu	f a.u.s	W Britain sea sh. D s.l	Eng. not. 957
J S1 - n a If. K.	Faba-leavcd	cu	fjnjn.au	W Sicily 1731. S co	Hoc. in. 133. 92
JX *** L-	Behen	un	2 jn.jl	W Crete J713. S co	Di. el. 397.409
J83 indica/far.	Indian	un	2 jn.jl	p Nepal 1823. C co	
J4 viscaginoMcs Horn.	Virago-like	cu	* jn.jl	Pk Dauria 1824. D co	
JM frainifoliaO//j	Grass-leaved	or	2 jn.jl	W Altaia 1819. I) co	
Si ffrainifolia Mur.	procumbent	cu	f jn.jl	Pk Siberia 1825. D co	
JUS? libilia Bl'b.	small-red	ft	myjn	F Portugal 1732. S co	Di. el. 314.406
J Si r o ^ a Moon.	petalless	ft	jn.jl	A.P. 1801. S co	
JF92 hispanica Othh	Spurrey-like	■	1 jn.jl	W Armenia 1824. D co	
J Pannora Z.	Gypsophila	■	P. 1822. D co		
J Ofites Per*.	fleshy	ft	Jnjl	Crea Spain 1819. S co	
J Other	Spanish	A cu	1 jlau	Crica England gra. so. D co	Eng. bot 85
J Volga	Volga	A pr	1 jlau	UT Volga 1824. D co	
J small-flowered	small-flowered	A pr	1 jlau	H Hungary 179fi. D co	
J effuse	effuse	A pr	1 jlau	Volga 1823. D co	
J Siberian	Siberian	A >u	Iftjn.au	Siberia 1773. D co	Hof.gotl.150.ic.
J Russian	Russian	A or	2 jn.jl	Russia 1820. S co	
J many-flowered	many-flowered	Q cu	1 jn.au	H Hungary 1794. S ■	W.&K.1.5G
J tall	tall	A or	3 jn.jl	Tauria 1819. S ■	
J Tartarian	Tartarian	je II	2 jnau	Russia 1769. D co	
J gigantic	gigantic	je II	3 jnjl	Africa 1738. S s.l	Walt. h. 11
J clammy	clammy	3 jnjl		levant 1739. S co	To. it 2.361
J conic	conic			England sun. fi. S s.l	Eng. bot. 932
J conoid	conoid			& Europe 1683. S s.l	M.li. 5. 36.6
J cylindrical-flwd^	cylindrical-flwd^	1 jnjl		S. Europe 1824. S co	
J vravc-ivealed	vravc-ivealed	ft au		C. G. H. 1775. S p.l	
J English	English	*	jn.jl	Britain san. fi. S co	Eng. bot 1178
J Portuguese	Portuguese	1 jn.jl		Portugal 1732. S co	Di. el. 311.401
J three-toothed	three-toothed	A myjn		Barbary 1823. S 81	
J French	French	1 myjn		France 1683. S s.l	Di. el. 310.399
J compressed	compressed	1 jnjl		Valencia 1825. S co	
J Basil-like	Basil-like	1 mr.jn	 1823. S co I	
J two-rowed	two-rowed	Iftjnji	 1817. S s.l	Sch. mo. 39
J Cerastium-like	Cerastium-like	f lau	W S. Europe 1732. S s.l	Di. cl. 309.397	
J five-wounded	five-wounded	1 jn.au	Bd England san. fl. S co	Eng. bot. 86	
J Canary	Canary	ft jnjl	W.R Madeira 1822. S co	Spr. end. 3	
J night	night	2 jn.au	Br S. Europe U183. S 81	Di. el. 310.400	
J grey	grey	W	N. Africa 1818. S co		
J halncest	halncest	O or	R Spain 1&1. S co		
J Holn.	fat	Q. nr	X	Br.R S Europe 1818. S co	
J enclosed	enclosed	ft jnjl	R 1820. S co	
J various-colored	various-colored	ftfjl	R	Greece 1817. S co	Fl.gr. 410
J reflexed	reflexed	1 jlau	Br	S. Europe 1726. D co	Mag. mo. 171.ic
J small, petaleil	small, petaleil	U^n	R 1821. S co	
J minutc-flowred	minutc-flowred	U^n	II	Portugal 1823. S co	
J cauesciit	cauesciit	R	R	Naples 1822. D co	
J tlichotonuiis	tlichotonuiis	ft	W	Hungary 1791. S s.l	
J		W	Iberia 1823. S co		
J552 vespertina Retx.	cveifniT	ft jn.jl	Pk 1815. S co	
J 533 sabulettum Lk.	gr J1Sjt	ft jn.jl	R 1794. S 81	Jacvin. 3.81
J 534 cheiranthiflia Schott	WallOvd	ft jn.jl	R	Barbary 1796. S co	Bot mag. 677
J 535 obtusifolia W.	obtuse-leaved	1 jlau	P. 1818. S co	
J 522 colorata Schous.	colored	1 jnjl	P. 1820. S co	
J 515 imbricata Desf.	imbricated	£ i	P.	San Rocco 1821. S co	
H520 hirsutissima	thick-leaved	1 jlau	P. 1819. S co	
J	lender	O pr	P. 1818. S co	
J	Jcpisee	1 jlau	C. G. H. 1774. S p.l		
J	ciliated	1 jnjl	W. 1823. S co	
J	pendulous	O or	Pk 1804. S s.l	
J		myjl	R	Sicily 1731. S s.l	
J				T,ot mag. 114	

11543 <i>hfspida</i> Desf.	hispid		Jn.jl		Barbary	1817.	S	co	
11544 <i>laxiflora</i> Biot.	loose-flowered				Spain	1820.	S	co	
U5V> <i>dechmbeens</i> Bernh.	decumbent				Spain	1823.	S	co	
11546 <i>secundiflora</i> Oth	side-flowering				Syria	1820.	S	co	
11547 <i>viscosissima</i> Ten.	clammiest				Italy	1824.	S	to	
11548 <i>longicaulis</i> Pou.	long stemmed				Spain	1818.	S	co	
11549 <i>quadrifolia</i> Dec.	four-toothed				Alps	1822.	D	co	
11550 <i>pusilla</i> W. & J.C.	weak h d				Hungary	1804.	D	co	
11551 <i>alpestris</i> Jac.	alpine	4 A un	1 my.jl		Austria	1774.	D	s.l	
11552 <i>rupestris</i> L.	rock	& Q un			SwitzerL	1774.	S	s.l	
11553 <i>glaucaulis</i> Lag.	glaucous-leaved	Q or			Spain	1820.	S	co	
11554 <i>parviflora</i> O/A	small-leaved	CD or				1817.	S	co	
11555 <i>mapertia</i> L.	unopcn-Jtawcreed	O un	2 jn.jl		Madeira	1732.	S	81	
11556 <i>clandestina</i> Jac.	hidden-flowered	O un	1 jn.jl		C. G. H.	1801.	S	co	
11557 <i>portensis</i> L.	Oporto	O cu	1 jl.au		Portugal	1759.	S	s.l	
11558 ,Antirrhinum.	Snapdragon	O un	1 jn.jl		N Amer.	1712.	S	p.l	
11559 <i>gemmiflora</i> IV.	twin-flowered	O cu	1 jn.jl			1816.	S	co	
11560 <i>flavescens</i> W. -K.	flavescens	i A P ^r	1 jn.jl		Hungary	1804.	D	p.l	
11561 /imphlia W.	Flax-leaved	O pr	1 jl.au			1817.	S	81
11562 Heretical..	Cretan	O pr			Candia	1732.	S	s.l	
11563 <i>divaricata</i> Clem.	divaricate	O or			S. Europe	1818.	S	co	
11564 , <i>edoldes</i> Jac.	Scdum-like	O pr			Crete	1804.	S	co	
U565Suxifraga L.	Saxifrage	t A cu			France	I(40).	D	61	
11566 <i>petra</i> AFK# AT	rock	5 A P ^r			Hungary	1822.	I)	co	
11567 <i>ctmirnula</i> ftr*	heU-flowered	\$ A P ^r			Piedmont	IH2J.	D	co	
11568 <i>viscida</i> Spr.	viscid	5 3 or			Carniola	1820.	S	co	
11569 <i>longipetala</i> Ven.	long-petaled	O pr	1 jn.au		Barbary	1822.	S	co	
11570 <i>nuntans</i> L.	nodding	3t A un	2 jn.jl		Britain	cal. ro.	D	co	
11571 <i>saxatilis</i> Sims	stone	^ A cu			Siberia	1800.	D	s.l	
11572 <i>livida</i> W.	livid	& A pr	1 jn.jl		Carniola	1816.	D	al	
11573 <i>tenuis</i> IV.	slender	ik A pr			Buical	1816.	D	p.l	
11574 <i>quadriplda</i> O-A	four-cleft	^ Q or			Verona	1818.	S	co	
11575 <i>viridiflora</i> L.	green-flowered	Q cu			Spain	1739.	S	co	
11576 <i>chlorantha</i> W.	pink-flowered	^ A cu			Germany	1732.	D	si	
11577 <i>nicaeensis</i> All.	Nice	O or			Piedmont	1820.	S	co	
11078 <i>ramosissima</i> De{	branched	3l Q or			Barbary	1820.	S	co	
11579 <i>diversifolia</i> O/F	different-leaved	O or				1820.	S	co
USHO Psammithis /.*.	Psammithis	^ Q or				1818.	S	co
11081 <i>cath6hea</i> Oth	universal	5 A cu			Italy	1711.	D	co	
11582 <i>elcgans</i> Errf.	elegant	O pr			Portugal	1819.	S	co	
11583 <i>spatulata</i> Zitt.	spatulate	Q or			Caucasus	1823.	S	co	
11584 <i>tenuifolia</i> GrtA	fine-leaved	O or			Dahuna	1820.	S	co	
11585 <i>ripens</i> Pat.	creeping	O pr			Siberia	1823.	D	co	
11586 <i>virginica</i> L.	Virginian	^L A or			N. Amer.	1783.	D	p.l	
11587 <i>Catesbe</i> Walt	Catesby's	3t A or			Carolina	1810.	D	ps	
11588 <i>strcta</i> L.	upright	O pr			Spain	1802.	S	co	
11589 <i>muscinula</i> Z.	flytrap				Spain	1596.	S	p.l	
11590 <i>noctiflora</i> L.	night-flowering				England	san. fi.	S	s.l	
11591 <i>oriflata</i> H.K.	ornamented				C.G. H.	1774.	S	p.l	
11592 <i>Corsica</i> Dec.	Corsican				Corsica	1820.	D	co	
11593 <i>egyp'tica</i> L.	Egyptian				Egypt	1800.	S	81	
11594 <i>sctXcca</i> All.	silky				S. Europe	1801.	S	8.1	
11595 <i>pubescens</i> Loi.	pubescent				Corsica	1818.	S	co	
11596 <i>picta</i> P ^r *	painted				1822.	S	co	
11597 <i>laevigata</i> Sm.	smooth-leaved				Greece	1817.	S	co	
11598 <i>bicolor</i> 7%ore	two-colored				France	1820.	S	Dec. ic. 42	
11599 <i>rambsa</i> Oth	branchy				Crete	1820.	S	Fl.gr. 4125	
11600 <i>reticulata</i> Desf.	netted	O							
11601 <i>pennsylvamca</i> Mx.	Pennsylvanian	k A			Barbary	1804.	S	p.l	
11602 <i>vallesia</i> Z*	Vallesian	5 A			N. Amer.	1806.	I)	Bot. rec. 247	
11603 <i>fruticosa</i> L.	shrubby	J			SwitzerL	1765.	D	s.l	
11604 <i>caspica</i> Pers.	Caspian				Sicily	1629.	C	p.l	
11605 <i>Requienii</i> Oth	Uequien's				Caucasus	1823.	I)	Com. h. 33	
IlhO6 amo?na L.	pleasing				Corsica	1823.	D	co	
11607 <i>supina</i> Bieb.	supine	# jn.au	Pk		Tartary	1779.	D	p.l	
11608 <i>deprensa</i> Bkb.	depressed	ijn.jl	W		Caucasus	1804.	D	s.l	
11609 i-rad6xa X.	paradoxical				Iberia	1816.	S	co	
11610 <i>chloracifolia</i> Sm.	Chloro-leaved	1 jl	Pk		Europe	—.	D	p.l	
11611 <i>itifilia</i> D ^r .		1 au.s	LY		Armenia	1796.	D	Bot. mag. 807	
11612 pAtula Detf.	spreading	Hmy,jn	W		Italy	1759.	S	Jac. ob. 4.7D	
11613 <i>polyphilla</i> L.	many-leaved	1 myjn	W		Barbary	1823.	D	co	
11614 <i>ncromalis</i> W.8/K.	grove	1 jn.jl			Germany	...	D	Cl. h. 1.290	
11615 <i>longiflora</i> Ehrh.	long-flowered	Q cu			Hungary	1852.	S	co	
11616 <i>Auplcuro</i> ^desL.	Bupleurum-lik				Hungary	1793.	D	W. & K. 1.8	
11617 <i>mollissima</i> /*.	softest	A cu			Persia	1801.	S	To. it 154	
11618 <i>cana</i> CMA	hoary	A or			Italy	1739.	D	W. & K.3.248	
11619 <i>régia</i> Sims	royal	A or	U my.au	C	N. Amer.	1811.	D	Bot. mag. 1724	
11620 <i>Mociniña</i> Dec.	Mocini's	Al or	I jn.jl	P	Mexico	IH27.	S	FL mxx. ic in*	
11621 <i>afccndns</i> Lag.	ascending	O cu	J myau	Pk	Spain	1822.	S	co	
11622 <i>cespiflora</i> &ftf.	turfy	A pr	1 my.au	Pk	Caucasus	1824.	D	co	
11623 <i>cordifolia</i> AIL	cordate-leaved	A or	i jn.jl	Pk	Piedmont	1819.	S	AL ped. 2a 3	
11624 <i>iltacion</i> Mur.	Atocion	O pr	{ my.jl	Pk	Levant	1781.	S	Jac. vin. 3. &	
11625 <i>Psebdo-iJtacion</i> Desf.	Bast-Atocion	O or							
11626 <i>Armèria</i> L.	Sweetwilliam	O or		Pk	N. Africa	1820.	S	co	
2'alba	white-flowered			Pk	England	cor. fl.	S	s.l	
11627 <i>compacta</i> TU.	compact	O or	Ijjls	Pk				44nn	
11628 <i>congesta</i> Sm.	crowded	O or	1 jn.jl	Pk	Caucasus	1823.	S	co	
11629 <i>perfoliata</i> Oth	perfoliate		J	R	Greece	1818.	S	co	
Cucubalus chonetbus Par.					& Europe	1817.	S	co	

V9 STELLARIA L. STITCH WORT. (*Stella*, a star; starlike flowers.) *Caryophylle**. &> -T^{2/3}

11630 *majorum*L. grove k Aw 1 ap.jn W Britain moi.wo. 1) co Eng. bot

11631 *latifolia*Pers. broad-leaved ^ A un lijn.au W Germany 1816. D co

J <i>fi32dich6tomaL.</i>	forked	O pr	1} year	W	Britain	clt. gr. S	Eng. bot 537
-i <i>Kuumuoosa wui.</i>	bulbous	O pr	1 year	W	Cannthia	182J. I)	Jac. ic. 3. 4ii8
J <i>•6 H viscula Bieb.</i>	clammy	O pr	1 year	W	Hungary	1820.	W. & K. 22
J <i>M J/olestea L.</i>	Holosteae	O pr	1 apjn	W	Britain	woods 1)	Eng. bot 511
J <i>63xi Laxmanni Fis.</i>	Laxmann's	O pr	1 apjn	W	Siberia	1823. S	
J <i>•37 velutina & r.</i>	velvety	O pr	1 myjn	W	Baical	1824. I)	
J <i>i38^ ramineaL.</i>	Grau-leaved	O pr	1 apjn	W	Britain	hed. b. D	Eng. bot 803
J <i>L639 glauca With.</i>	glaucous marsh	O pr	f my.au	W	Britain	moi. m. I)	Eng. bot 825
J <i>(140crassifolia&ArA.</i>	thick-leaved	O pr	1 my.au	W	Germany	...	D co
J <i>1641 uliginosa Cur.</i>	bog	O pr	1 jnjl	W	Britain	rivul.	S bog Eng. bot 1074
J <i>aquatica Pott.</i>							
J <i>11642 crastidids L.</i>	Cerastiura-like	3i	ft Jnjl	W	Scotland	scalp. D	bog Eng. bot 911
J <i>11643 humif&a Swz.</i>	trailing	3	ft myjn	W	Norway	181& S co	Fl. dan. 978
J <i>11644 arenariaL.</i>	sand	O pr	ft jn	W	Spain	1799. S co	
J <i>11645 scaplerga W.</i>	scape-bearing	O pr	i jnjl	W	Scotland	scrivu. D co	Eng. but 1266
J <i>11646 i multicaulis Jac.</i>	many-stemmed	5	jnjl	W	Carmthia	1816. S co	Jac.c. 1.19
J <i>11647 dahinea IF.</i>	Dahunan	£	1 jnjl	W	Dahuria	1818. S co	
J <i>U54 murMis Lk.</i>	wall	O pr	ft jn.jl	W	Candia	1824. S co	
J <i>11649 elegans Ser.</i>	elegant	O pr	1 myjn	W	Siberia	180t. S co	
J <i>11650 ttnigipes Gol.</i>	long-stalked	O pr			N. Araer.	1820. S p.l	
J <i>*19° ARENA^RIA L.</i>	SANDWORT.						
J <i>purple</i>		(Arena, sand; habitation.)					
J <i>red</i>	O pr	i jl	P	Spain	182a S	s.l	
J <i>marine</i>	O pr	i jnaj	P	Britain	san.fi. S	s.l	Eng. bot 852
J <i>152rubraJ.</i>	A pr	ft jnjl	P	Germany	1795. D	s.l	
J <i>H633 marina Roth</i>	salt-pit	O pr	ft jnjl	P	Bohemia	1820. S s.p	
J <i>marginata Dec.</i>	mediate	O w .	J jl	P	Britain	scaco. S co	Eng. bot 958
J <i>654sapina&r.</i>	glandular	O or	i jnjl	P	Europe	1820. S co	Jac. sc. 355
J <i>H5 media,,</i>	Canada	O pr	* jnjl	R	N. Amer.	1812. S p.l	
J <i>i... marina & B.</i>	Grass-leaved	A Pr	ft jnjl	W	Siberia	1817. D e o	H. got 5
J <i>666glandul.sj Jac.</i>							
J <i>1168 adensis Pers.</i>	smoothest	A or	jnjl	W	Caucasus	1816. D co	
J <i>1678 gramimflua Schr.</i>	long-leaved	O pr	jjl	W	Siberia	1823. D co	Gm. si. 63.2
J <i>Bifflia Bieb.</i>	Dahurian	O pr	1 jnjl	W	Dahuria	1824. D sl	
J <i>2 glaberrima Dec.</i>	handsome	O pr	i jnjl	W	Dahuria	1824. D sl	
J <i>graminiflva IV. en.</i>	Nardus-leaved	O pr	jnjl	W	Siberia	1827. D co	
J <i>g. dittof</i>	Otitcs-like	O pr	4 jnjl	W	Siberia	1820. D co	
J <i>ah&rica Fts.</i>	stiff	O pr	ft jnjl	W	Siberia	1823. D co	
J <i>Fis.</i>	Pine-leaved	O pr	i jl.au	W	Caucasus	1823. D s.p	
J <i>Adams</i>	subulate	O pr	i jl.au	W	Caucasus	1822. D s.p	
J <i>P&gens Step.</i>	viscosa Fa.	O pr	* jnjl	W	Siberia	1820. D s.l	
J <i>capilldns Pair.</i>	capillary	£ A					
J <i>na</i>	Juniperus	O pr	jnjl	W	Armenia	1800. D s.p	Sin. ic. in. 1.35
J <i>A*</i>	strict	O pr	"myjn	W	N. Amer.	1812. D s.p	
J <i>toriciflMia/..</i>	Larch-leaved	O pr	i au	W	Britain	...	Jac. au. 3.272
J <i>Si imb" cata Bleb.</i>	imbricated	O pr	i jnjl	W	Caucasus	1820. D &l	
J <i>22 procea Snr.</i>	tall	O pr	ft jnjl	VV	Siberia	1820. D s.l	
J <i>673rostrata U'.&K.</i>	rostrate	O pr	i au	W	Hungary	1816. D s.p	Vil. del. 4.47.1
J <i>1674striata Vtl.</i>	striated	O pr	ft jnjl.au	W	Switzerl.	1683. D s.p	Vil. del. 4.47.6
J <i>167a austriaca Jac.</i>	Austrian	A pr	ft jn.6	W	Austria	1793. D s.p	Jac.au. 3.270
J <i>Villirsi/Balb.</i>							
J <i>76cap>Ilkeea</i>	capillacous	O pr	4 jnjl	W	Piedmont	1819. S	Al-ped.89.2
J <i>677trifl.r-iA</i>	three-flowered	A pr	i ap.jl	W	Europe	1816. D s.p	Cav. ic. 3.249. 8
J <i>1178grandiflral.</i>	great-flowered	A pr	i jn.s	W	Switzerl.	1783. D s.p	ALped.10
J <i>JS: X</i>	Helm's	A pr	i jnjl	W	Siberia	1826. D s.l	
J <i>680 macrocarium Ph.</i>	long-fruited	A pr	i jnj.j	W	N. Amer.	1810. D s.l	
J <i>681 canescens Vahl</i>	cancscnt	A pr	* jnjl	W	1817. D s.l	
J <i>inacrocarpa Horn.</i>							
J <i>11682 verna L.</i>	vernal	£ A pr	imyau	W	Britain	mount D	Eng. bot.51S
J <i>11683 ramostssima TV.</i>	branchiest	£ O pr	ft jnjl	W	Hungary	1816. S co	
J <i>H681 cassiputia&ArA.</i>	turfy	£ A or	i jnjl	W	Switzerl.	1826. D s.l	
J <i>11685 Gerard's</i>	Gerard's	£ A Pr	i myau	W	France	1822. D s.p	Jac. au. 445
J <i>11686 saxatilis L.</i>	rock	A pr	ft jljau	W	Germany	1732. D s.p	Gm. si. 4,63.8
J <i>JS7pcndula W. ^K.</i>	pendulous	A pr	» M	W	Hungary	1816. D co	W. & K. 2.87
J <i>11688 tenuifoha L.</i>	uncleaved	O pr	* jnjl	W	England	aan.fi. S co	Eng. bot 210
J <i>S Barrelieri Vil.</i>	Barrelier's	O pr	ft jnjl	W	S. France	1K20. S co	Bar. ic. 580
J <i>4 viscidula Vil.</i>	hybrid	O P'	ft jnjl	W	S. France	1827. S co	ViLdel.4.47
J <i>thandrou</i>	viscidish	O pr	ft jnjl	W	France	1818. S co	
J <i>JS7 Mediterranean</i>		O pr	i jnjl	W	1817. S t,l	Sch. mo. 30
J <i>Mediterranean</i>	O cu	* jnjl	W	Barbary	1816. S co		
J <i>hairy</i>	O pr	i jnjl	W	Mediterr.	1823. S co		
J <i>calyculate</i>	A Pr	i jnjl	W	Alps	1822. D co	Jac c. 16	
J <i>setacea Thui.</i>	setaceous	A Pr	i jnjl	W	Caucasus	1880. D ■	
J <i>heteromalla Pers.</i>		A Pr	i jnjl	W	Hungary	1817. D s.l	
J <i>11696 fasciculata Gou.</i>	fascicled	O cu	jn	W	France D co	
J <i>fastigiata £m.</i>					Scotland	scmo. D s.p	Eng. bot. 1744
J <i>filifolia Forsk.</i>	thread-leaved						
J <i>JG9 glomerata Bieb.</i>	glomerated				Arabia	D s.p	Vahl sy. 1.12
J <i>11699 uliginosa Schl.</i>	marsh				Tauria	1818. S co	
J <i>(1700polygono'ides Wxd.</i>	Knotgrass-like				Switzerl.	1819. D si	Dec.ic.46
J <i>Hed verticillata W.</i>	whorled				Switzerl.	1822. D co	Al. ped.64.4
J <i>IZ^2 tetrAquadra L.</i>	square				Armenia	1823. C s.p	
J <i>4S lnceolau AU.</i>	lanceolate				Pyrenees	1731. D s.p	AL ped. 2. 89.1
J <i>montaia L.</i>	Cherleria-like				Switzerl.	1823. D s.l	ALped.26,5
J <i>mountain</i>	dense				France D s.p	
J <i></i>					Hungary	1824. D s.l	
J <i></i>					France	1800. D s.p	Bot. mag 1118

11707 <i>ferryllifolia L.</i>	Thyme-leaved	O w	<i>i</i>	Jn.jl	W	Britain	walls.	S	co	Eng. bot 923
11708 <i>coimbricensis Brat.</i>	Coimbra pubescent	O pr	<i>1</i>	jn.jl	W	Portugal	IHH9.	S	■	Brot.ph.73
11709 <i>pubescens Dec.</i>	A pr	<i>£</i>	jl		W	Archifwl.	1820.	S	co	
11710 <i>brevicaulis Ster.</i>	short-stalked	A pr	<i>A</i>	jn.jl	W	Alp. Eur.	1823.	I)	co	
11711 <i>sritbra Poir.</i>	rough	A br	<i>i</i>	jnjn	W	Alp. Eur.	1822.	D	co	
11712 <i>ciliata L.</i>	ciliated	A br	<i>i</i>	mr.au	W	Ireland	ir mou.	I)	s.p	Eng. bot 1745
11713 <i>marginatafl? 26. in.</i>	margined	3i A Br	<i>1</i>	jn.jl	W	Caucasus	1818.	D	s.p	
11714 <i>multicadhs Wul.</i>	many-stalked	k A pr	<i>1</i>	jl.jau	W	Europe	1794.	D	s.p	Jac. c. 1.17.1.
11715 <i>trincris L.</i>	three-nerved	O w	<i>5</i>	myjn	W	Britain	woods.	S	s.p	Eng. bot 1483
11716 <i>balefifica L.</i>	Balearic	A pr	<i>i</i>	mr.au	W	Majorca	1787.	D	s.p	Her. st. 1. 15
11717 <i>ncplaidcs L.</i>	Fepis-like	A pr	<i>i</i>	my.jl	W	Britain	sea sh.	I)	s.p	Eng. bot 189
11718 <i>procumbens Vahi</i>	prucumbent	4 Alpr	<i>1</i>	jl.jau	W	Egypt	1801.	D	s.p	Vahsy.2.33
1391. CHEKLEWI* Hal. CHURLERIA. (<i>John H. Charier</i> , assistant to Jno. Bauhin.) <i>Caryophyllece</i> . 1.—a										
11719 <i>mloldes L.</i>	Seduni-like	£ A or	<i>1</i>	ji.au	Y.w	Scotland sc. alp.	D s.l	Eng. bot 1212		
1392. GARIDE'LLI* Tou. GA RID ELL A. (P. Garidel, M.D., of Provence, bot. auth.) <i>Ranunculacear</i> . 1.—2.										
11720 <i>Nigellastrum L.</i>	Nigella-feawca*	O or	<i>1</i>	£ jn.jl	B.o	France	1736.	S	co	Bot. mag. 1266
1393. MALPI'GHL* I* BARBADOES CHERRY. (<i>M. Malpighi</i> , prof. med. Bologna.) <i>Malmghuilece</i> . 12.—19.										
11721 <i>fucata Ker</i>	painted	■ ■ ■	—	—			1814.	C	p.f	Bot. reg.189
	<i>macrophylla Deaf.</i>									
11722 <i>urens L.</i>	stinging	■ ■ ■	—	—	Pk	S. Amer.	1737.	■	p.l	Bot teg. 94
IV123 <i>cuñenias H. & B.</i>	Cuba	■ ■ ■	—	—	Pa.R	Havannah	1824.	■	p.l	
11724 <i>Angustifolia L.</i>	narrow-leaved	■ ■ ■	—	—	Pk	W. Indies	1737.	■	p.l	Bot cab. 321
11725 <i>quitulum L.</i>	<i>Holy-leaved</i>	■ ■ ■	—	—	Pk	S. Amer.	1759.	■	I	Cav.dis.8.L'X)'2
11726 <i>coccifera L.</i>	berry-ben ring	■ ■ ■	—	—	Pk	W. Indies	1733.	■	p.l	Bot. reg. 5M
11727 <i>glabra L.</i>	smooth-leaved	■ ■ ■	—	mr.jl	It	W. Indies	1757.	■	p.l	Bot. mag. 813
11728 <i>biflora Poir.</i>	two-flowered	■ ■ ■	or	10 jl.jau	Pa. II	S. Amer.	1810.	■	p.l	Cav. dis. 8.234.8
	<i>jnnicifbia Cav.</i>									
11729 <i>punicifolia L.</i>	Pomegranate-lvd	ft O or	12	...	Pk	W. Indies	1690.	■	PL ic.	16G. 2
11730 <i>aginea Sum.</i>	<i>Beech-leaved</i>	sH O or	10	jl.jau	Y	S. Amer.	1820.	■	p.l	
11731 <i>dbbia Cav.</i>	doubtful	i D o r	10	jl.jau	Y	Jamaica	1820.	■	p.l	Cav. dis. 8.242
11732 <i>nlida A/t.</i>	glimsy-leaved	* CD or	6	mr.au	Pk	W. Indies	1733.	■	Cav. dis. 8.239.1	
1394. BYRSO'NIMA Rich. BVRSONIMA. (<i>Bt/rsa, a hide; used for tanning</i>) <i>Malpghidicear</i> . 13.—31.										
11733 <i>verbascifolia Dec.</i>	Verbascum-lvd	* CD or	g	jl.jau	Pa.R	Guiana	1810.	C	p.f	Aub. gui. L184
	<i>Malpighia verbascifolia L.</i>									
11734 <i>/aurifolia Jl. \$B.</i>	Laurel-leaved	■ ■ ■	10	jl.jau	Y	Cumana	1824.	■	H	
11735 <i>nervosa Dec.</i>	nervose	• D or	8	jl.jau	Pa.Y	Brazil	1820.	■	H	
	<i>Malpighia macrophylla J.</i>									
11736 <i>altissima Dec.</i>	talc&t	t • or	60	jl.jau	W	Guiana	1820.	C	p.l	Aub. gui. 1. 181
11737 <i>crasBif uha Dec.</i>	thick-leaved	f CD or	20	jl.jau	Y	Guiana	1793.	C	p.l	Aub. gui. 1.183
	<i>Malpighia crassifolia L.</i>									
11738 <i>MoureUa Dec.</i>	Mourcila	J CD or	20	au	■	S. Amer.	1823.	C	p.l	Aub. gui. 1.183
	<i>Malpighia MourUa Aub.</i>									
11739 <i>chrysophyila H.S/Ji</i>	golten-leaved	• D or	10	au	Y	Orinoco	1823.	C	p.l	
11740 <i>spicata Dec.</i>	spiked	6	jl.jau	Y	Antilles	1810.	C	ps.1	Cav. dis. 8.237	
11741 <i>Ibida Dec.</i>	shming-leaved	■ C or	6	my.au	Pk	W. Indies	1759.	C	p.l	Bot mag. 2462
	<i>Malpighia lucida Sivz.</i>									
11742 <i>conScea Dec.</i>	AcAthercy-Icaved	2 CD or	30	my.au	W	Jamaica	1814.	C	p.l	SL jam. 2.163.1
	<i>Malpighia enriacea Swx.</i>									
11743 <i>reticulata Dec.</i>	netted	ftQor	10	jl.jau	P.v	Cayenne	1823.	C	pi	
11744 <i>volubilis Dec.</i>	twining	ICZ)or	10	aus.s	Y	W. Indies	1793.	C	p.l	Bot mag. 809
11745 <i>nlidifl Dec.</i>	pale	• • or	4	...	Pa	Cayenne	1820.	C	p.l	
	<i>Malpighia pallida Poir.</i>									
1395. BANISTEMt/^ L. BANISTERIA. (<i>lieu. J. Banister, a curious botanist</i>) <i>Malpighiticea</i> *. 12.—47.										
11746 <i>acunulata Cav.</i>	auriculate	LCD or	10	—	Y	Brazil	1820.	C	ps.1	Cav. dis. 255
11747 <i>ciliata IV.</i>	ciliated	■ ■ ■	—	—	Y	Brazil	1796.	C	s.l	Cav. dis. 9-254
11748 <i>spfendens Dec.</i>	shining	■ ■ ■	10	—	Y	S. Amer.	1812.	C	8.1	Cav. dis. 253
	<i>heterophylla W. fulgens Lam.</i>									
11749 <i>Humboldt'swa Dec.</i>	Huaiboldt's	■ ■ ■	10	—	Y	S. Amer.	1824.	C	ps.1	
11750 <i>sericea Cav.</i>	silky	■ ■ ■	10	—	Y	Brazil	1810.	C	s.l	Cav. dis. 9. 259
11751 <i>tomentifl Desf.</i>	tomentose	■ ■ ■	10	jnau	Y	S. Amer.	1820.	C	ps.1	
11752 <i>faurifolia L.</i>	Bay-leaved	■ ■ ■	10	jl.jau	Y	Jamaica	1733.	C	s.l	Bot reg. 937 ..
11753 <i>ferruginea Cav.</i>	rusty	■ ■ ■	10	—	Y	Brazil	1820.	C	ps.1	Cav. dis. 9. 213
11754 <i>periplocafoha Desf.</i>	Periuloca-Wd	■ ■ ■	10	jn.au	Y	PortoRicoll818.	C	ps.1		
11755 <i>folgens L.</i>	shningfi-ited	■ ■ ■	6	—	Y	W. Indies	1759.	C	r.m	
11756 <i>zanzibarica Boj.</i>	Zanzibar	■ ■ ■	10	—	Y	Zanzibar	1825.	C	ps.1	
11757 <i>ovata Cav.</i>	ovate-leaved	■ ■ ■	or	6 jn.au	Y	St.Domm.1810.	C	ps.1	Ven. ch. 51. A	
1396. TRIO'PTERIS L.	TRIOFTERIS.	(Tris, three, pteron, a wing; capsule.) <i>Malpighihcca</i> . 2.-8.								
11758 <i>jamaicensis L.</i>	Jamaica	L Q o r	10	...	Y	Jamaica	1822.	C	p.f	
11759 <i>sericea Lo. C.</i>	silky	LCDoR	■ ■ ■	—	S. Amer.	1823.	C	p.l		
1397. TETRA'PTERIS Cav. TETRAFTERIB. (<i>Tetra, four, awing; capsule.</i>) <i>Malpighiiicete</i> . 2.-7.										
11760 <i>Mixifolia Cav.</i>	Box-leaved	* CD or	6	...	■ ■ ■	StDomin.1822.	C	p.f	Cav. dis. 262.1	
	<i>Tnóptericis duxifdia W.</i>									
11761 <i>citriflifl Pers.</i>	Orange-leaved	il CD or	6	—	■	Cayenne	1818.	C	p.l	Cav. dis. 262.9
	<i>Tri6pteris citrifolia Sun.</i>									
*1398. HETERO'PTERIS/f.>J9. HBTEROPTERIB. (<i>Heterus, various, pteron, wing; caps.</i>) <i>Malpighiæct</i> . 8.-J;										
11762 <i>purpiircia//. \$B.</i>	purple	LCDor	10	...	P	W. Indies	1759.	C	p.l	Tav. dis. 9. « #
	<i>Bamsfertfl purptirea L.</i>									
11763 <i>parvifolia Dec.</i>	small-leaved	LCDoR	10	—	P	W. Indies	1820.	F	■	Ven.ch.51
11764 <i>brachifta H. \$B.</i>	crw*-branched	L CD or	10	—	Y	W. Indies	1759.	F	■	
	<i>Banist&ria brachi& L.</i>									
11765 <i>chrysophylla L.</i>	golden-leaved	L O or	10	—	■	Brazil	1793.	C	r.m	Jac. sc. 105
	<i>Banisteria chrysophylla Lam.</i>									
11766c!Dr{ilea//.^A	buc-Jiowering	LCDoR	10	—	B	Havannah	1823.	C	p.l	Pl.ed. B ₂ J ² *
U767 <i>nlitida Kth.</i>	glossy	• • or	10	—	Y	S. Amer.	1809.	C	8.1	Cav. dis. 244
	<i>Banisteria nitida Lam.</i>									
HZ51 <i>P Kita Dec.</i>	fea <i>feati</i>	■ ■ ■	10	—	Y	S. Amer.	1809.	C	s.l	Botreg.!S°
11769 <i>flonbOnda//. QB.</i>	appendiculat	■ ■ ■	or	10	—	Y	St. Vine.	1820.	C	p.l
	<i>bundle-flwd</i>	■ ■ ■	or	10	—	B	Mexico	1824.	C	p.l

- | | | | | | |
|-------|----------------|--------|---|----------------------------------|--------|
| 1399 | <i>HIR^TA</i> | Jac. | <i>HIRJEA.</i> (J. N. de la Hire, a French physician, d. 1727.) | <i>Malplghiiucc*</i> . | 5.—19. |
| 11770 | rcdninta | Jac. | reclined ^ Q o r 10 ... Y w. Indies ... C s.l Jac am. 146 | | |
| 11771 | odorata | W. | sweet-scented j _Q or 8 Guinea 182J. C s.l | | |
| 11772 | irritans | Rox. | nodding a Q or 10 jn.au ... E. Indies 1820. C p.l | | |
| 11773 | indica | JZo* | Indian i. or 10 jn.au ... E. Indies 1820. C s.l Rox. cor. S. 1G0 | | |
| 11774 | glauca&scens | Lo. C. | glaucescent Q or 10 jn.au ... E. Indies 1823. C s.l | | |
| 1400 | ERYTHRCXYLON | L. | <i>ERVTHIROXVLO.V.</i> (<i>Erythros</i> , red, <i>xylon</i> , wood.) | <i>Erythrox&lece.</i> a — SS | |
| 11775 | /ypericifblum | Lam. | Hypcricum-lvdv • tm 40 ... Y.G Mauritius 1818. C s.l Cav. dis. 8.230 | | |
| 11776 | havanensis | Jac. | Havannah \$ o r 50 ... Y.u Havannah 1822. C 81 Jac. am. 87.2 | | |
| 11777 | /auriflum | Lam. | Laurel-leaved \$ O tm 50 ... Y.a Mauritius 1823. C 81 Lam. dis. 226 | | |
| 1401 | SETHM | Kth. | <i>SETIIA.</i> (S. Sethi, author of a work on culinary vegetables.) | <i>Erythroxyl<i>b</i></i> . | 1. |
| 11778 | Indica | Dec. | Indian £ • tm 40 ... Y E. Indies 1824. C 81 Rox. cor. 1.88 | | |
| | | | Erythr6xylon monogynum Rox. | | |
| 1402 | SPERGULA'STRUM | Mx. | <i>SPBRGULACTRUM.</i> (From similarity to <i>Spergula</i> .) | <i>Caryophyllece.</i> 1. — 3. | |
| 11779 | lanuginosum | Mx. | woolly £ A cu J.jn.jl W.p N. Araer. 1821. S s.p | | |
| | | | Microptalon lanuginfrsum Pers. | | |

CLASS X. ORDER 4.

DECA'NDRIA PENTAGY'NIA.

- | | | | |
|--|-----------------------------|--|------------------------------------|
| 1403. CNE'STIS J. | CNESTIS. | <i>Iknœo</i> , to scratch; prickly capsules.) | <i>Connaracete</i> . 3-9. |
| 11780 <i>gfabra</i> <i>Lam.</i> | smooth | ■ CD or 10 ... W.a | Mauritius 182i C s.l Lam.il. 387.1 |
| 11781 <i>poiphylla</i> <i>Lam.</i> | many-leaved | ■ L3 or 6 ... P | Mauritius 1823. C s.p |
| 11782 <i>cornicuta</i> <i>Lam.</i> | small-horned | ■ CD or 10 ... P | Guinea 1793. C s.p |
| 1404. AVERRHOA^ L. | AVERRHOA. | (<i>Averrhoes</i> , of Corduba, in Spain.) | <i>Oxaldece</i> . 2. - 3. |
| 11783 <i>Bilimbi</i> <i>L.</i> | Bilimbi tree | i [j f r 8 au.s R.Y. E. Indies? 1791. C s.l Cav. dis. 7.219 | |
| 11784 <i>Carambbla</i> <i>L.</i> | Carambola free* O ft | ■ - G.R. Ceylon 1733. C s.l Cav. dis. 7. 220 | |
| _ 1405. SPCNDIAS L. | Hoa PLUM. | (Greek name of a kind of wild plum.) | <i>Terebinthhcete</i> . 4. |
| 11785 <i>pup&rea</i> <i>L.</i> | purple | + • fr 30 ... W.u W.Indies 1817. C s.p Sl. jam. 2.219 | |
| 11786 <i>littea</i> <i>L.</i> | yellow | £ • fr 30 ... Y.a W.Indies 1739. C s.p Mor.sur. 13 | |
| Myrobaianus <i>L.</i> | | | |
| 11787 <i>dilicis</i> <i>Forst.</i> | sweet <i>Otaheite apple</i> | • fr 50 ... Y.a Society Is. 1793. ♀ Lam. il. 384 | |
| 11788 <i>mangifera</i> <i>Pers.</i> | mango-bearing | 1 • fr 50 ... w E. Indies 1820. ♀ | |
| Mangifera pinnata <i>L.</i> | | | |
| 1406. BUCHANA*NU Rox. BUCHANANIA. (<i>Francis Buchanan</i> , now Hamilton, M.D.) <i>Tercbtuthhcett</i> . 2. | | | |
| 11789 <i>latifolia</i> flat, | broad-leaved | t Q f r 30 ... W. E. Indies 1820. C s.l.p | |
| U790 <i>angustifolia</i> <i>Aox.</i> | narrow-leaved | f1 Z1fr 30 ... W. E. Indies 1820. C s.l.p | |
| Mangifera axillaris <i>Lam.</i> | Sp6ndias axillaris //. | Bcng. | |
| 1407. POUPART/ii Com. POUPARTIA. (Called <i>Bois de Poupart</i> by inhab. of Bourbon.) <i>Tt</i> , | | | |
| 11791 <i>borb6nica</i> <i>Lam.</i> | Bourbon | ± • tm 40 ... P Bourbon 1825. C s.Lif | |
| 140a ROBE'RG/J Schreb. ROBERGIA. (<i>Laurent Roberg, yrof.</i> med. at Upsal.) <i>Terebinthfce</i> < g. \—& | | | |
| 11792 <i>frutescens</i> <i>W.</i> | frutescenc | * • or 6 ... W Guiana 1823. C s.l.p Aub. gui. 187 | |
| •1409. COTYLE^DON L. NAVELWORT. (<i>Kofyle</i> , a cavity; cuplike leaves.) <i>Orassnlacea</i> . 41—45. | | | |
| 11793 <i>orbiculata</i> <i>Haw.</i> | romd.leaved | t L I CIU 2 il.au R C G. H. 1798. C s.l | |
| 11794 <i>ovata</i> <i>Haw.</i> | ovate-leaved | ■ I ru 2 jl.o K C G. H. 1789. C s.l Bot mag. 321 | |
| 11795 <i>ppilaria</i> <i>Thun.</i> | papillary | ■ i cu 1 a.u R C G. H. 1822. C 81 | |
| 11796 <i>Wobblonia</i> <i>Haw.</i> | | ■ i cu 2 jl.s K C G. H. 1690. C s.l | |
| 11797 <i>craosifolia</i> <i>Haw.</i> | thick-icaved | t L or ■ C. G. H. 1824. C 1.1 | |
| 11798 <i>trHbra</i> <i>Thun.</i> | thrc-flowered | tt. ■ or *mys Pk.WC.G. H. 1821. C s.l | |
| 11799 <i>alternans</i> IV. | alternate | n. ■ or ■ jnau C. G. H. 1816. C s.l | |
| 11800 <i>ramusa</i> <i>Haw.</i> | branchy | M- or ■ jnjl C. G. H. 1748. C s.l M. h. 12.7.39 | |
| ramosissima <i>Mil.</i> | | | |
| 11801 <i>ramobllsima</i> <i>Salm</i> | branchiest | t L i lor 1 my.jn — C. G. H. 1816. C 81 | |
| 11802 <i>fascicul^ris</i> <i>H. K.</i> | cluster-/ai;af | t L ■ neu 1 jl. — C. G. H. 1759. C 81 Bur. af. 18 | |
| 11803 <i>hemisphaerica</i> <i>L.</i> | hemispherical | ti i J IU 2 my. W.p C. G. H. 1731. C s.l DeepL87 | |
| 11804 <i>vlridis</i> Uftv. | green | t L \ or ■ — C. G. H. 1824. C 81 | |
| 11805 <i>rotundifolia</i> <i>Haw.</i> | round-leaved | t L \ or ■ — C. G. H. 182(5. C s.l | |
| 11806 <i>cristata</i> <i>Haw.</i> | crested | ■ t Al or ■ Va C. G. H. 1820. C 81 | |
| 11807 <i>clavifl.lia</i> <i>Haw.</i> | club-leaved | tf ■ lor ■ P C. G. H. 1824. C s.l | |
| 11808 <i>spuma</i> <i>L.</i> | spurious | «■ L Jcu 1 jl.au — C. G. H. 1731. C s.l Com. r. 83 | |
| 11809 <i>undulata</i> /law. | waved | t L i lor 1 jnjl C. G. H. 1818. C s.l Bur. af 19.2 | |
| muconatona <i>Lam.</i> | | | |
| 11810 <i>asminalifera</i> <i>Salm</i> | Jasmine-flwd | tt. \ or ■ jl.au C. G. H. 1818. C s.l | |
| 11811 <i>datta</i> <i>Haw.</i> | tall powdered | <. \ or ■ jnjl C. G. H. 18J6. C s.l | |
| 11812 <i>cujeita</i> 7% II». | wedgelike | «. \ or ■ my C. G. H. 1818. C s.l | |
| 11813 <i>corugeana</i> <i>Haw.</i> | glittering | tt. \ or ■ jn C. G. U. 1818. C s.l Bot. cab. 1030 | |
| 11814 <i>curviflra</i> <i>Sim</i> | curve-flowered | tt. \ cu \ o C. G. H. 1818. C s.l Bot. mag. 2044 | |
| purphrea <i>Haw.</i> | | | |
| 11815 <i>decussata</i> <i>Smf</i> | croii-leaved | t L J cu 1 au S C. G. H. 1819. C s.l Bot. mag. 2518 | |
| papifarii <i>Haw.</i> | | | |
| 11816 <i>tricupidata</i> //, u. | tricuspidate | tt-l. or ■ C. G. H. 1823. C s.1 | |
| 11817 <i>cacalurina</i> <i>Thun.</i> | Cacalura-like | ■ C. G. H. 1818. C s.1 | |
| 11818 <i>rhombifolia</i> <i>Haw.</i> | rhomb-leaved | ■ L \ or ■ C. G. H. 1823. C s.1 | |
| 11819 <i>raamml.Hvls</i> -in / nun. | mammillary | tt. \ or ■ my. W.p C. G. H. IS!! C s.1 | |
| 11820 <i>canaliculata</i> <i>Uta</i> <i>Uaw.</i> | small-channelled | tt. \ or ■ jn-jl C. G. H. 1818. C s.1 | |
| 11821 <i>ungulata</i> <i>Lam.</i> | clawed | tt. \ or ■ my.jn C. G. H. 1818. C s.1 | |
| 11822 <i>maculata</i> <i>Sal.</i> | spotted | tt. \ or ■ my.jn W.p C. G. H. 1H18. C s.1 | |
| 11823 <i>coccinea</i> <i>Cav.</i> | scarlet | tt. JCU S jnjl S C. G. H. 1818. 1 | |
| 11824 <i>tuberculosa</i> <i>Lam.</i> | tuberculose | tt. \ or 1 jn.au D C. G. H. 1800. C U | |
| 11825 <i>gracilis</i> <i>Haw.</i> | slender | tt. \ or 1 jn.au ... C. G. H. 1800. C U | |

\$11826 cespitsba <i>Haw.</i>	turfy	£ iAI cu	1 jn.au	Y	California 1796.	C s.1	
Unguef6rmis <i>H. K.</i>							
11827 dichotoma Jstw.	fork-spined	£ AJ	1 *au	W	C. G. H. W18	Gm. si. 67.2	
\$11828 spinbsa <i>L.</i>	spiny	£ Q> cu	ft jnjl	K	Siberia 1790.		
Sedum spindsum <i>W.</i>							
11829 serrata <i>L</i>	notch-fcauof	£ A cu	1 in.jl	R	Siberia 1732.	D s.l	
Spanish		£ Q> cu	ft jnjl	K	Spain 1796.	S 8.1	
						Dec. pi. 122	
navel Penny-wort	£ A pr		Y	Britain	sha.roc. D	s.l	
Mucizoni's	£ A pr		Y	Portugal	182a D	s.l	
yellow	£ A pr		Y	England	moi.ro. Sk s.1	Eng. bot. 1522	
11833 Malacoph? <i>Pall.</i> soft-leaved	O pr		P.Y	Davuria	1815. S s.1	Pa. it. 3. ap. G.I	
11832 Orteat? <i>Chrys.</i> Malacoph? <i>Hum Fit.</i>							
11833 M t f							
•U10. SE'DUM <i>L.</i>	STONECROP. (<i>Sedere</i> , to sit; manner of growth.)				Craauldeca. 61 — 8 ■		
11834 maximum <i>Lk.</i>	greatest purple	£ A or	2 jls	W	Spain	1791. D s.l	
latifbluin <i>Bert.</i>							
11835 albicans <i>Haw.</i>	white	£ A or	2 jls	W	Europe	1794. D s.l	
11836 verticillMum <i>L.</i>	v/horUeaved	£ A or	1 jls	Pk	S. Europe	... D s.1	
11837 tripltfllum <i>Haw.</i>	three-leaved	£ A or	1 jl.au	Pk	S. Europe	... D s.l	
11838 argitnum <i>Haw.</i>	sharp-toothed	£ A or	Iftjl.au	P	Carpathia	1820 D s.l	
11839 paucidens <i>Haw.</i>	few-toothed	£ A or	A jnjl	W	1818. D s.1	
11840 pectinatu <i>Haw.</i>	pectinate	£ A or	4 ap.au	W.o.	1818. D s.1	
11841 TQlvvium <i>L.</i>	common Orpine	£ A or	2 jls	P	Britain	bor. fi. D s.1	
11842 7b1epht>d1c* <i>Mx.</i>	Tclphiuin-like	£ A or	1 jls	P	N. Amer.	1810. D s.l	
11843 y/nucumpscros <i>L.</i>	Auacamp&cros a.	£ A or	4 jl.au	P	France	15JK 1) s.1	
11844 livulum JK	hyd	£ A or	ijn.au	W.o.	1816. D s.1	
11845 ^izuon <i>L.</i>	Aizoon	£ A "or	1 jls	Y	Siberia	1757. D s.1	
11846 joopulifiMium/.	Poplar-leaved	■ A or	1 jl.au	W	Siberia	1780. D s.l	
11847 opijositofliolum? A,	opposite-leaved	£ A or		W	Caucasus	... 1) s.1	
11848 spurium <i>Bleb.</i>	spurious	■ A or		Pk	Caucasus	1816. D s.l	
11849 dent a turn <i>Haw.</i>	toothed	■ A or		P	1810. 1) s.1	
11850 hfbridium <i>L.</i>	hybrid	■ A or	my.jl	Y	Siberia	1766. D s.l	
11851 ternutum <i>Mx.</i>	ternate	■ A or	ftjlau	W	N. Amer.	1789. D s.1	
portulacotrics <i>MM.</i>						Bot. reg. 142	
11852 stellutum <i>L.</i>	starred	O or	i jnjl	W	S. Europe	1640. D s.l	
11853 divaricatum <i>H.J.C.</i>	spreading	n. i or	I jn.jl	Pk	Madeira	1777. C s.1	
11854 Ccpofa <i>L.</i>	Ccpfa	O or	1 jl.au	W	France	HU0. S s.l	
11855 spatulatum <i>Kit.</i>	spathulate	£ Q) or		W	Hungary	1815. S s.l	
11856 ffcmprvividcesifet	Scmperviv.-like	£ A or		R	Iberia	1823. D 8.1	
11857 ochrolchum <i>Sin.</i>	yellowish-white	£ A or		W	Greece	1818. D si	
11858 altissimum <i>Pen.</i>	tallest	£ A or	1 jl.au	P.Y	S. Europe	1739. D s.1	
Jacqta Haw. SempervWum «diffirme Jac.						Jac. vin. 1.81	
11859 cerulcsccns <i>Haw.</i>	cuerlescent	£ A or					
11860 riasphyllum <i>L.</i>	thick-leaved	£ A pr		Y	1820. D s.1		
11861 reflexum <i>L.</i>	reflex-leaved	£ A or		W	walls	1) s.1	
11862 septangulXrc <i>Haw.</i>	seven-angled	£ A or		Y	Britain	walls. D s.1	
11863 glaucin Sm.	glaucus	£ A or		J>0l	Eng. bot. 695	
11864 collnum IV. en.	hill	£ A or		Y	1795. D s.1	
11865 irinius <i>Haw.</i>	smaller shoit-t-ludz	£ A or	4JLau	Y	England	bar sa. U s.1	
reflexum Dec. not of others.			1 jn.au	Y	Eng. bot. 2477	
11866 spirMe <i>Haw.</i>	spiral	£ A or	ft jlau	Y	J815. D 8.1	Dec. pl. 115	
sexangijare Dec. not E B.					Europe	1810. D 8.1	
11867 recurvatum IV.	recurred green	£ A or	ft jnjl			Dec. pi. 116	
11868 anopetalin Dec.	upward-lvd grcen	£ A or					
rupestre VU.							
H aurantiacum <i>Sal.</i>	Orange	£ A or	i Jj	O	& France	1820. D s.1	
11869 virens // K.	green	£ A or	I jnjl	Y	Portugal	1774. D s.1	
11870 rupestre L	rock	£ A pr	i jlau	Y	England	rocks. D s.1	
11871 virbccc* IV. en.	viresent	£ A or	1 jn.au	G.Y	Eng. bot. 170	
11&72 VonXaridnum Sm.	Forster's	£ A or	il	Y	Wales	w.roc. D si	
11873 quadrifidum // K.	four-cleft	■	*i jlau	W	N. Asia	1800. 1) s.1	
11874 nCidum // K.	naked-branchedn.	i J pr	* jnjl	W.a	Madeira	1777. R s.1	
11875 quinquefidum W.	five-clett	■		Pa.Y	IIH0. D si	
11876 hexapetalum <i>Haw.</i>	six-petaled	■		W	Caucasus	1816. S s.1	
11877 s'xflidum Btob.	annual six.cleft	■		W	S. Europe	18K. D s.1	
11878 monregalcnseAi^.	Montreal	■		R	Switzerl.	1826. D sp	
11879 ^pens Dec.	creeping	■		P.T	Spain	1752. D si	
11880 hispanicum L.	Spanish	■		W	England	rocks. D s.1	
11881 album //.	white	■		Pk	Britain	m.al.p. D s.1	
11882 villbsum L	villous	■		I au	W	N. Europe	17-m. S s.1
11883 Annuum L.	annual	■		iau	B	Italy	1795. S s.1
11884 atratum L.	dark annual	■		B	Africa	1822. S s.1	
11885 neriliuem Vahi	pale-h\Q	■		B	England	walls. D s.1	
11886 sexangulari..	*ix-angled	■		Y	Eng. bot. 520	
11887 acre L.	acrid	■		Y	England	walls. D s.1	
11888 saxatile W.	rock	■		Y	Eng. bot 1646	
11889 pallens Yen.	pa'e-glaucous	Qox	ft jnjl	W	& Europe	1830. D s.1	
11890 Andersunii G. Don	Anderson's gl.	£ Q) or	ft jnjl	W	S. Europe	1816. S s.1	
glaficum W. Sc K.	not E. B.					W. & K. 2.181	
11891 pallidum Bt&b.	pale	Q or	i in.jl	Pa R	Caucasus	1817. S s.1	
11892 oblongum <i>Haw.</i>	oblong	£ A or	i jlau	W	Britain	rocks. D s.1	
11893 aiglicum L.	English	£ Apr	ft jlau	W	Britain	rocks. D s.1	
2 microphllum	small-leaved	£ A or	I jnau	W	Britain	rocks. D 8.1	
3 hib^namicum	Irish	£ A or	ft jn.jl	W	Ireland	rocks D s.1	
1411. PE'NTHORUM W. PBNTHORUM. (<i>Pente</i> , five, <i>horos</i> , column; capsule.)						Craauldece. 1.	
11894 seddles W.	Sedum-like	£ A cu	1 jl.au	G.Y	Virginia	1768. D s.1	
						Lam. il. 390	
1412. GRIITLUM L.	GBIELITM.	(Grcios, old; aspect^				Rosuete. 3.	
11895 tenuifolium L.	slender-leaved	2 ap.my	Y	C. G. H.	1790. R p.l	Sw. fl. gar. 171	
11896 humifsum Thun.	trailing	1 apjl	Y	C. G. H.	1825. R p.l	Bur.af.34. 1	
11897 laciniatum Gae.	jagged	£ or	ft au	Y	C. G. H.	1825. R p.l.s Sw. A. gar. 306	
1413. BICPHYTUM Dec. BIOPHYTUM (<i>Bios</i> , life, <i>pkyton</i> , plant; pods open when touched.)	Oxalie. 1.						
Ib98senitvumDec.	sensitive	O pr	ft jls	Y	China	1823. S s.1	
O'xalis sensitiva L.						Jac,ox.7a4	

- * 1414. CXALIS L. WOOD SORREL. (*Orys*, sharp, or sour; qualities.) *Oxalidea?* 97.—160.
.. 1. HEDRSAROI'DR*. — *Peduncles many-flowered*; stems leafy: leaves trifoliolate, middle one petiolated.
- 11899 Plumiért Jac. Plunder's *-C* or 2 year Y S. Amer. 1K23 C p.l Bot. reg 810
11900 Barretters Jac. H. Barretters *Cipr* 1st year Pa.R Caracas 1824 f. II Jac. ox. 9
H901 frutirima Bad. shrubby *JP'* 1 year Y Ilio Jan. 1817. f. II
- II. CORNICILIAVA. — *Peduncles 1,2, or many, flwd; stems leafy, lvs. palmate-trifol., leaflets sessile.*
- 11*02 pergrannas Haw. perennial ■ LAI 2 my.s Y N.S.W. ... O s.p
HlyQ3bil² fl. rida So. Dilleniuss's O pr 2 iny.au Y America 1798. S s.p DI.cl.221
11904 stricta L. florid O pr 2 my.au Y America 1798. S s.p
11905 Lygin Ph. strict A pr II jn.o Y N. Amer. 1658. O s.p Jac. ox. 4
11906 cormiculata L. Lyon's LAI pr 1 jl.o Y N. Amer. 1816. O s.p
11907 mlnrophil. Pair. small-homed O pr ... Britain sh.roc.O s.p Eng.bot.1720
riihens Haw. small-leaved iQlpr + N.S.W. — S s.p
- 11908 ripens Thun. creping-stkd ■ 1 mr.ap Y C.G.H. 1793. O »p Jac. ox. 78.1
11909 rfsca Jic. rosry iAJp 1 mr.ap Chile 1823. O s.p Bot mag. 2415
11910 flonftinda Lch. bundle-flwd ■ IAJor 1 jn.au R S. Amer. ? 1827. 1) sf Bot reg. 1123
11911 latenflura Jic. lateral-flowered ■ iAJ pr 4 mr.ap P C. G. H. 1824. O s.p Jac.C.204
- III. SE8SIMF6L1E. — *Peduncles axillary, one-flwd; stems elongated, leafy: leaves trifoliolate, sessile.*
- 11912 macroBtlys Jac. long-styled ViAlor \an P C. G. H. 1793. () s.p Jac. ox. 9
11913 tubiflura Jac. tube-flowered tf LAIor 1 o.n Pk C. G. H. 1790. O i.p Jac.ox. 10
11914 canébeens Jac. canescent tf IAlpr 1 ja.s P C. G. H. 1821. O sp Jac. ox. 11
11915 secfinda Jac. side-flowering tf LAI or § o.n Id C. G. H. 1790. O s.p Jac. ox. 12
J1916 hirta L. haT-stalked tf LAI or f o.n Li C. G. H. 1787. O sp Jac. ox. 13
J1917 hirtella Jac. small hairy tf LAI pr 4 mr.ap R C. G. H. 1821. O sp Jac. ox. 16
J1918 multiflora Jac. many-flowered tf IAlor f fm.r Li C. G. H. 1789. O s.p Jac. ox. 15
J1919 rubella Jac. branching small red tf iAJ or 4 s.n Pk C. G. H. 1791. O s.p Hot. mag. 1031
J1gDrosacea Jac. rmccolored 4 IAlpr | s.n Pk C. G. H. 1793. O fl.p Bot. mak. 1698
J1921 fitigida A.A. fulgid tf IAJpr i s.n C C. G. H. 1820. (> s.p Bot. reg. 1073
- IV. CAULIFLORJB. — *Peduncles axil, one-flwd i stem* elongated, few-leaved* (ff/fc *3--5? ■■■■■
- J1922 repatria Jac. creeping-rooted i " LAI i " d " F" C. G. H. 1798. O s.p Jac. ox. 20
J1923 virginea Jac. virgin < iAI dr < ja.au W C. G. H. 1820. O s.p Jac. sc. 3.275
J1924 disticha Jfc. two-rowed tf iAI pr % ja.s Pa.Y C. G. H. 1818. O s.p Jac. ox. 18
" S? incarnata L. flesh-colored tf IAlpr * ap.jn V C. G. H. 1719. O s.p Jac. ox. 71
veiny tf S * * o.n V.V C. G. H. 1823. O s.p
11927 furckta Loo*. folk-leaved tf S dr J s R C. G. H. 1823. O s.p Bot. cab. 1056
- V. CAPRI'MJR. — *Stems; peduncles 1,2, or many, flowered; leaflets 3 or many, stalked.*
- J122? niv? " Ait. Bowie's' tf LAI pr ft sin' C C. G. H. 1823. O s.p
H929 sericea L. silky tf iAI or 4 ap.my Y C. G. H. 1754. O s.p Jac. ox. 77.1
liaH Burmann's tf IAJpr i o.n Y C. G. H. 1820. O s.p Bur. af. 29
J1931 tetrphylla Cap. four-leaved tf El pr i Jn.jl P Mexico 1823. O 8.p Cav. ic. 3.237
**932 violacea L. violet-co/oreo* tf A or i my.jn L.P N. Amer. 1772. O s.p Jac. ox. 80.8
!S? Pmctata B. M. dotted tf A or 4 ap.jn P .O s.p Bot. reg. 2781
52? <Prtna L. goava-foot tf LAJor 4 mr.jn F C. G. H. 1757. O s.p Jac. ox. 76.1
remua Thun. drooping tf iAI or 4 f" y Y C. G. H. 1757. O s.p Jac. ox. 6
- > tr? compressed tf LAI or 4 year Y C. G. H. 1794. O s.p Jac. ox. 78. 3
silentXta Jac. toothed tf LAI or 4 " d F C. G. H. 1793. O s.p Jac. ox. 7
! mirata Jac. purpled tf iAJ pr 4 o.n F.W. C. G. H. 1822. O s.p Jac. sc. 356
HYZ prophylia Horn, long-leaved tf LAI pr 4 J*-8 Pa.R C. G. H. 1820. O s.p
JJ*O livida Jac. livid tf LAI or 4 o.n F C. G. H. 1793. O s.p Jac. ox. 8
H941 lobata Sims lobed tf iAI or 4 o.n Y C. G. H. 1823. O s.p Bot mag. 8386
- VI. SIMPLICIFOLJE. — *Stemless; leaves simple; scapes one, or many, flowered.*
- **42 monophylla L. one-leaved tf iA)pr i o.n Y C. G. H. 1774. O s.p Jac. ox. 79.3
H943 lepida Jac. pretty Cape tf IAlpr 4 ja.n W C. G. H. 1823. O s.p Jac. ox. 21
H944 rostrata Jac. beaked tf iAI 4 o.n P.v C. G. H. 1795. O s.p Jac. ox. 22
- VII. PTEROPODS. — *Stemless; leaves bisfoliate, or trifoliolate; petioles winged; scapes one-flowered.*
- JJStScrispa Jac. curled tf iAJpr 4 an W G G. H. 1798. O s.p Jac. ox. 23
J SHfilep'ina Jac. hare's-rarrd tf IAlpr 4 on W G G. H. 1795. O s.p Jac. ox. 25
J1947 asurina Jic. assWarof tf iAI pr 4 n.d Y G G. H. 1792. O s.p Jac. ox. 24
J1948 lanceafflia Jac. spear-leaved tf IAlpr 4 o.n Y G G. H. 1795. O s.p Jac.ox.26
J1949 abffifolia Jac. Bean-leaved tf IAJ pr i o.n Y G G. H. 1794. O s.p Jac. ox. 27
- VIII. AcTogELLE. — *Stemless; leaves trifoliolate, petioled; scapes one-flowered.*
- 11950 /abumiflba Jac. Laburnum-lyd tf LA)pr 4 8.0 ■ G.G. H. 1793. O s.p Jac. ox. 28
11951 sangufneia Jac. bloody/fawt/ tf IAlpr iod V C. G. H. 1795. O s.p Jac. ox. 29
11952 rbbro-flava Jac. red and yellow tf IAlpr R.T C. G. H. 1823. O s.p Jac. ox. 50
1 H³3 tricolor Jac. threecolored tf iAI or * o.d W.a C. G. H. 1794. O e.p Jac. ox. 47
11354 cilikris Jac. bowed tf IAlor i o.n P C. G. H. 1713. O s.p Jac. ox. 50
HYSarcuAta Jac. flacid tf IAlpr 4 o.n V C. G. H. 1795. O s.p Jac. ox. 31
1 W6facqila Jac. runty tf IAI pr 4 jn jl W G G. H. 1820. O s.p Jac. sc. 3.274
ambiguou tf IAJpr is.d W G G. H. 1790. O s.p Jac. ox. 43
y/Avc-leaved tf IAJpr I o.n W C. G. H. 1795. O s.p Jac. ox. 44
brown-spotted tf LAJ pr ft my.jn Y GG. H. 1795. O ap Jac. ox. 45
sulphur-colored tf iAI pr 4 o.n P.v G G. H. 1795. O s.p Jac. ox. 63
specious tf LAI pr is.n V C. G. H. ■ O s.p Jac. ox. 60
variable tf LAI or 4 o.d W.R C. G. H. 1795. O s.p Jac. ox. 58
great-flowered tf LAI or 4 o.d W C. G. H. 1790. O Bp Jac. ox. 54
Sims's tf LAJ or 4 o.d W C. G. H. 1790. O Bp Bot mag. 1683
purple tf IAJpr 4 o.n P C. G. H. 1812. O s.p Jac. ox. 56
small convex tf iAI or ft Jan Pk C. G. H. 1789. O s.p Jac. ox. 55
smooth O pr ijnil P 1818. S co
gran-edged tf IAJpr 4 sd W C. G. H. 1812. O co Jac. ox. 68
11968 pulchella Jac. i A el 4 o.n W C. G. H. 1795. O co Jac. ox. 69
11969 obtusa Jac. pretty S iAlor 4 o.n W C. G. H. 1812. O s.p Jac. ox. 79.1
11970 lankta L. XAunUleaved tf IAlor ion W C. G. H. 1791. O s.p Jac. ox. 77.2
11971 lufeola Jac. vrooWy.leaved tf iAlor i jas Y C. G. H. 1823. O s.p Jac. ox. 65
yellowish tf iAlor 4 ap.my W Britain grov. O co Eng.bot.762
little sorrel or com. tf A cul i ap.my VV N. Amer. — O co
11973 americana Dec. American tf A pr i ap.my Li G.G. H. 1791: O co
delicate tf LAIor 4 8d W ■ ■ ■ ■ O s.p Jac. ox. 19
floating * Al or 4 o. V C.G.H. 1815. O s.p Jaesc.2.205
thread-atmd tf E) or

11977 blfida <i>T</i> .	c\ovcn-leaved	tf IAlor	\$ s.o	V	C. G. H.	1791.	O	s.p	Jac. ox. 70.4
11978 cuneifolia <i>Jac.</i>	wedge-leaved	tf IAlor	I ap. my	W	C. G. H.	1793.	O	s.p	Jac. ox. 41
11979 Hnearis <i>Jac.</i>	linear-shaped	tf IAlor	V	C. G. H.	1795.	O	s.p	Jac. ox. 32	
11980 reclinata <i>Jac.</i>	reclining	tf IAlor	Je.n	Pk	C. G. H.	1796.	O	s.p	Jac. ox. 34
11981 cunepta <i>Jac.</i>	cuncate	tf iAlpr	J ja.s	Y.w	C. G. H.	1822.	O	s.p	Jac. ox. 40
11982 minuita <i>Jac.</i>	venmlion	tf IAlpr	i ja.s	Ve	C. G. H.	1819.	O	Bp	Jac. ox. 35
11983 pusilla <i>Jac.</i>	weak	tf IAlpr	I ja.s	Pa.R	C. G. H.	1823.	O	s.p	Jac. ox. 42
11984 PiCthe CoL	Piotta's	tf IAlpr	i ja.s	O	C. G. H.	1816.	O	s.p	Col. h. rip. 1
11985 tdnera <i>Spr.</i>	tender	tf IAlpr	i my	Y	Brazil	1826.	O	s.p	Bot. reg. 1045
11986 carnosa <i>Mol.</i>	fleshy	tf IAlpr	Y	Chile	1826.	O	8.p	Bot. reg. 1063	
IX. ADENOPHT'LUS. - Stems leafy I <i>lvs petioled</i> , 3—5-foliate; leaflets linear underneath, at top glandular.									
11987 gKbra Thun.	smooth	tf iAl or	§ myjn	P	C. G. H.	1795.	O	s.p	Jac. ox. 76.3
11988 versicolor L.	various-colored	ff IAlor	I jamr	C	C. G. H.	1774.	O	s.p	Bot mag. 155
11989 gracilis <i>Jac.</i>	slender	tf IAlpr	i ja.rar	*s.o	C. G. H.	1820.	O	s.p	Jac. ox. 33
11990 elongata <i>Jac.</i>	elongated	tf IAlor	I	W	C. G. H.	1791.	O	s.p	Jac. ox. 37
2 amoëvia	flue red	tf IAlpr	J so	R	C. G. H.	1810.	O	s.p	Jac. sc. 206
11991 tenuifolia <i>Jac.</i>	flnc.lcaved	tf IAlor	io.n	W.R	C. G. H.	1790.	O	s.p	Jac. ox. 38
11992 polyph Ha <i>Jac.</i>	many-leaved	tf IAlor	I	Pa.P	C. G. H.	1791.	O	s.p	Jac. ox. 39
11993 Hifolia <i>Jac.</i>	thread-leaved	tf IAlor	I	Pk	C. G. H.	1822.	O	s.p	Jac. sc. 273
11994 pentaphylla Sims	five-leaved	tf IAlor	I	Pk	C. G. H.	1800.	O	s.p	Bot. mag. 1549
X. PALM An FOLIJE. - Stemless; leaves petioled palmately and peltately. 5-l&foliate; scapes one-flwd.									
11995 upinifolia Jac.	Lupine-leaved	tf LAI pr	i o.n	Y	C. G. H.	1791.	O	s.p	Jac. ox. 72
11996 Hava /..	yellow	tf AJ pr	f mr.ap	Y	C.G.H.	1775.	O	s.p	Bot. reg. 117
11997 pectinata <i>Jac.</i>	pectinated	tf iAlpr	\ s.n	Y	C. G. H.	1790.	O	s.p	Jac. ox. 7a
11998 flabclifolia <i>Jac.</i>	fan-leaved	3 IAJ pr	I s.n	Y.B	C. G. H.	1789.	O	s.p	Jac. ox. 74
11999 tomentosa Z,	dov/hy-leaved	tf iAl pr	i ap.my	W	C. G. H.	1791.	O	s.p	Jac. ox. 81
• 1415. LY'CHNIS L. - LVCIINIS. (<i>Lychnis</i> , a lamp; down of leaves as wicks.) <i>Caryopkfiliee</i> . 20 — 93.									
12000 Visçaria L.	clammy	■ A or	1 myjn	Pk	Britain	rocks. I)	co	Eng. bot 788	
2 plena	double	£ A or	I myjn	H	Britain	gard.	D	co	
12001 neglecta G. Don	neglected	£ A or	I	W	—	—	D	co	
Visçaria alba // or/.		£ Apr							
12002 alpina L.	alpine	£ A or	I ap.my	Pk	Scotland	scroc.	D	p.l	Eng. bot. 2254
12003 helvetica G. Don	Swiss a(pine	£ A pr	I jn.jl	R	Switserl.	1814.	S	s.p	Bot mag. 394
alpina B. M. not E. B.									
12004 chaledonica L.	Chaledonian	■ A or	2 jn.jl	S	Russia	1596.	D	p.l	Bot. mag. 257
StElba	v/hitc-jiowcred	■ A or	2 Jiijl	W	Ilussia	—	D	p.l	
3 plena	■ A or	2 jn.jl	S	Russia	—	D	p.l		
12005 fulgens Fis.	shining	■	U jn.jl	S	Siberia	1822.	D	p.l	Bot. mag. 478
12006 grandiflora Jac.	great-flowered	■	W jn.s	II	China	1774.	C	p.l	Bot. mag. 223
coronata Thun.									
12007 Flos Jbyu L.	Flower of Jove	£ A or	I	R	Germany	1726.	■	co	Bot mag. 390
12008 coronaria Lam.	crown	jt A or	3 jn.s	R	Italy	1596.	■	co	Bot. mag. 24
Agrostemma coronaria L.									
2 alba	vrhxtcflowered	£ A or	3 jn.s	R	—	—	■	co	
3 plena	douhc-Itowercdz	A or	H jn.s	R	Levant	1713.	■	r.m	Bot mag. £95
12009 Ccc'i Rbsa Desf.	Hose of heaven	O or	1 jl.au	F					
Agrostemma Cucii	irosa L.								
12010 aptala L.	petallcss	■	■	■	Lapland	1810.	S	&d	Fl. lap. 12.1
12011 brachypetala Horn,	short-petaled	■	■	■	Siberia	1817.	S	co	
12012 diidra Sib.	day-flowering	■	■	■	Britain	—	D	co	Eng. bot. 1579
sylvi'btris Jloppe									
2 flora pleno	double-flwd	■	■	■	Britain	hedges.	D	co	
12013 vesijertla Sm.	evenmng.^ft^g-	■	■	■	Britain	wa.&ti.	D	co	Eng. bot. 1580
dioica L.									
S fibre pleno	double-flwd	■	■	■	Britain	fields.	D	co	
12014 lee\la // K.	joyful	■	■	■	Portugal	1778.	S	i.l	
12015 Corsica Loi.	Corsican	■	■	■	Corsica	1818.	S	s.p	
12016 sibirica L.	Siberian	■	■	■	Siberia	1817.	I)	co	
12017 pyrenica Berg.	Pyrenean	■	■	■	Pyrenees	1819.	D	s.p	
12018 Floscituh L.	Cuckoo-flower	■	■	■	Britain	m.me.	O	co	Eng. bot 573
2 flor pl&no	double-flwd	■	■	■	Britain	bogs.	D	p	
12019 Githago /..	Corn-cockle	■	■	■	Britain	com.fl.	S	co	Eng. bot. 741
Agrostemma Githago L.									
2 nicsdnis	Nicean	■	■	■	Italy	1794.	S	co	
Agrostemma nicaeensis IV.									
1416. CERA'STIUM L. - MOUSE-EAR CHICKWEED. (<i>Keras</i> , a horn; form of capsule.) <i>CaryophOttete</i> . 48.-79.									
12020 pauciflbrum Stev.	few-flowered	O un.	I jn.jl	W	Siberia	1816.	S	co	
12021 incnorMc Bieb.	grove	O un.	I jn.jl	W	Caucasus	1H18.	S	co	
12022 perfoliatum /..	per foliate	O un.	I jn.jl	W	Greece	1725.	S	co	Di.eL217.284
12023 caudicisum Fis.	Caucasian	O un.	I jn.jl	W	Caucasus	1820.	S	co	
1S024 duharicum Fit.	Dahurian	■ A un.	I my.s	W	Siberia	1815.	D	co	Bot mag. 1789
amplexicate B. M. //olosteum Fis.	greatest	O un.	2 Jiijl	W	Siberia	1792.	■	co	Gm. si. 4.62.2
12025 maximum L.	Stellaria-hke	O un.	*myjl	W	N. Amer.	1810.	■	co	
12026 stellarioideu Afoc.	forked	O un.	Jnjl	W	Spain	1725.	■	co	
12027 dichotomum /..	glandulous	O cu	W	Europe	1827.	■	co		
V2U'2V) inflatum H. C.	inflated	O cu	I jn.jl	W	Europe	1827.	■	co	
12030 rudcralc Bieb.	rubbmh	O un.	J myjl	W	Caucasus	1817.	■	co	
12031 tauricum Spr.	Taurian	O un.	I my.jl	W	Tauria	1820.	■	co	
12032 vulgkum L.	common	O w	* apjn	W	Britain	san pi.	S	co	Eng. bot 789
12033 ovalc Pers.	oval	O un.	I ajxn	W	Europe	1800.	S	co	
12034 viHcusum L.	cinnmy-leaved	O w	I ap.s	W	Britain	pas.	S	co	Eng. bot. 790
12035 Aolosteoldes Fries	Holosteum-like	A un.	i my jl	W	Sweden	1818.	D	co	
12036 barbulbum Lk.	small bearded	O un.	* * *	W	S. Europe	1820.	S	co	
12037 anomalam W.SfK.	anomalous	O un.	i myjl	W	Hungary	1817.	S	co	
12038 semidec&ndrum L.	semi-decadrous	O w	i mr.ap	W	Britain	walls.	S	co	Eng. bot. 1630
12039 tetrindrum H. K.	tetrandrous	O w	J my.jn	W	Scotland	san.sh.	S	co	Eng. bot. 166
12040 pent&ndrum L.	pentandrous	8 un.	i jn.jl	W	S. Europe	1821.	S	co	
1^041 gracile Desf.	slender	8 un.	i jnyjl	W	S. Europe	1818.	S	co	
12042 brdhyp&talumDf<	short-petaled	O un.	I ap.my	W	S. Europe	1816.	S	co	Dec ic. 44
12043 difltaum Pers.	spreading	£ A un.	*ap.s	W	Europe	1820.	D	co	
12044 segyllifolium W.	Thyme-leaved	O un.	I ap.s	W	Siberia	1817.	S	co	
12045 manticum L.	Mantic	O un.	I ■■■	W	Hungary	1801.	S	co	W. & K. 96

18046 rupestre <i>Us.</i>	rock	Q	un	Jnjl	W	Siberia	1820.	S	co		
12047 campanulatum <i>Viv.</i>	<i>beto.</i> flowered	O	un	ijll. jl	W	Italy	1824.	S	co		
12048 grandiflorum <i>W. & K.</i>	<i>longi</i> flowered	A	pr	4.. jl	W	Siberia	1818.	D	p.g		
12049 tomenttaum <i>L.</i>	woolly	A	un	* Jnjl	W	S Europe	1648.	D	co		
12050 Bicherteinff Dec.	Bichertoin's	A	un	* Jnjl	W	Caucasus	1820.	D	p.l		
12051 pusillum <i>Ser.</i>	weak	O	pr	4jnjl	W	Siberia	1824.	S	co		
12052 lajidum <i>Lam.</i>	woolly	O	pr	i	W	Alp. Eur.	1819.	D	p.l		
19053 alpinum <i>L.</i>	alpine	A	um		W	Britain w.alp.		D	co		
12054 ovatum <i>Hort.</i> *	<i>orate-leaved</i>	A	pr	£ Jnjl	W	Carinthia	1816.	D	co		
12055 lati folium <i>L.</i>	broad-leaved	A	pr	4 jnjl	W	Britain w.alp.		D	co		
12056 glaciMe <i>Gau.</i>	icy	A	pr	i jnjl	W	Switzerland	1819.	D	pi		
12057 lylaticum <i>Wf.</i> ■	wood	A	pr		W	Hungary	1820.	D	p.s		
12058 arvense <i>L.</i>	field	A	w	ft my.au	W	Britain corn ft*.		D	eo		
12059 atractum <i>L.</i>	strict	A	un	1 ray.il	W	Austria	1793.	D	co		
2 sufruticisum <i>W.</i>		sufruticose		A	un	1 my.jl	W	S. Europe	1796.	D	co
12060 dioclum //, <i>K.</i>	dioecious	A	un		W	Spain	1776.	D	co		
12061 pennsylvanicum <i>Horn.</i>	Pennsylvan.	A	un		W	Pennsylv.	1810.	D	co		
12062 pubescens <i>Gol.</i>	pubescent	A	un		W	N. Amer.	1821.	D	co		
12063 matrense <i>Kit.</i>	Matrensan	A	un	*. 3 \times jl	W	Pannonia	1817.	D	co		
12064 pilsum <i>Horn.</i>	pilose	A	un		W	1820.	D	co		
12065 Sprungclif Ser.	Sprengel's	Q	un		W	1819.	S	co		
12066 hirs&tum <i>Ten.</i>	hairy	A	un		W	Italy	1824.	D	co		
12067 Tenorethiwn Ser.	Tenore's	O	un	i'5J	W	Italy	1822.	S	co		
pilosum <i>Ten.</i>											

1417. LARBREM Hil. LARBREA.
12068 aquatica Hil. water
Stellaria aquatica Pers. Cer&sti

(*Abbi Larbre*, a botanical author.)

Caryophyllace. 1.

* 1418. SPE'RGULA L.	SPURRY.	(<i>Spargo</i> , to scatter; expels seeds.)	<i>Caryophylle</i> e.	8. — 14.
12069 arvénis <i>L.</i>	corn-field	O w s jlau	W Britain san.fl. S co Eng.bot 1535	
12070 pentandra <i>L.</i>	pentandrous	O s	w England gan.fl. S co Eng.bot 1536	
12071 pallida <i>Sal.</i>	pale	A pr	Pa.R C. G. H. 1810. D s.p.	
12072 nodosa <i>L.</i>	knotty	A w	ijlau W Britain san.hc. D co Eng. bot 694	
12073 /aricina <i>L.</i>	Larix-like	Apr	pjnl W Siberia 1820. D f.p. Lam.il. 392.1	
12074 saginoides <i>L.</i>	Sagina-like	A w	4jn.au W Scotland scalp. D co Eng. bot 210'	
12075 subulata <i>Sun.</i>	awl-shaped	A w	W Britain san.hc. D co Eng. bot 1082	
12076 glabra <i>W.</i>	smooth	■ ■ ■	Europe 1816. D s.p. Al пед. 64.1	
1419. BE'RGM L.	BERGIA. (<i>P. J. Bereius</i> , M.D. professor of nat. hist Stockholm.)	<i>Caryophyllea</i> 1. — 2.		
verticillata <i>W.</i>	whorled	O cu	/ jnjl W.R Egypt 1820. S fl.p Del. sg. 26.1	

CLASS X. ORDER 5.

DECA'NDRIA DECAGY'NIA.

1420. PHYTOLA'CCA <i>L.</i>	PHYTOLACCA.	<i>{Phytolacca, a plant, lacca, lac; crimson fruit.}</i>	*	
18978 octandra <i>L.</i>	octandrous tf	EJ or 6 jl.n	W.a Mexico	1732. C i.l.
12079 stricta <i>Hgfm.</i>	strict	Q or 3 jnj	W.o S. Amer.	1820. C i.p.
-?/ heptindra <i>Rct?</i> ?				
19080 abyssinica <i>W.</i>	Abyssinian	*	my.jn	W.a Africa 1775. R s.l. Hof. got.2
19081 bogotensis <i>Hum.</i>	Bogotan	f CSJor	3 ...	W Bogota 1824. C s.p.
12082 dodecadandra <i>W. en.</i>	dodecadrous	H E3 or	6 my.jn	R Africa ... C s.l.
12083 decadandra <i>L. decand.</i>	Virginian Poke \$	A cul	5 aus.s	L.W Virginia 1615. C s.p. Bot mag. 931
12084 mexicana <i>Suit.</i>	Mexican	jf QS or 3	jl.au	W Mexico 1824. C s.p. Bot mag. 2633
icosandra <i>B. M.</i>				
12085 icosandra <i>L.</i>	Icosandrous	a	3 JLn	W E. Indies 1758. C s.p. Mil. ic. 207
12086 dolcea <i>L.</i>	dioecious	\$	14 ...	W.o S. Amer. 1768. C s.p. Her. st 70

CLASS XI: ORDER 1.

DODECA'NDRIA MONOGY'NIA.

1421. if'SARUM L.	ASARAOACCA.	(A, privative, <i>seira</i> , a bandage; unfit for garlands.)	<i>Asdrinte</i> .	4. - 5.
12087 arifMium Mx.	Arum-leaved	A ij Br	N. Amer. 1823. D p.l Hook. ex. fl. 40	
12088 euro'Mc'um L.	Eurojean	A f my P	England woods. D p.l Eng. bot 1083	
12089 canad'nsC L.	Canadian	A cu fapjl Br	Canada 1713. D p.l Bot cab. 889	
12090 gramiliflum Lo.C.	large-leaved	A cu up. my Br	N. Amer. 1820. D p.l	
12091 vir^finicum L.	Virginian	A cu \$ ap.my Br	Virginia 1759. D p.l Sw.fl.gar. 18	
1422. BOCCO%N/^ L.	BOCCONIA.	(Paolo Boecone, M.D., a Sicilian monk.)	<i>Papaverrhcea</i> .	2. - 3.
12092 flerwens L.	frutescent	Celandine Q 10[0]ja.ap W.Y	W. Indies 1739. C r.m Bot. cab. 83	
12093 integrif6lia/T4-A	entire-leaved	M or 4 ja.ap W	Mexico 1820. C I.p H.&B. n.1.35	
1423. MACLETAY^ R. Br.	MACLEAYA.	(A. Macleay, secretary to the Lin. Soc.)	<i>Papaverhcete</i> .	1.
12094 cordkta It. Br.	cordate	jf A or 6 my.jl R.Y	China 1795. D r.m Bot. mag. 1905	
Bocc5nia cordkta L.				
1424. BA'SSU L.	BASSIA.	(F. BassL curator of the botanic garden at Bologna.)	<i>Sapbetae</i> .	3. - 4.
12095 longifolia L.	long-leaved	f O or 40 . . . E Indies	1811. C p.l Lam. il. 398	
12096 latifilia Box.	broad-leaved	t or 40 . . . Y Pi	Indies 1790. C p.l Rox. cor. 1. 19	
12097 butyrkcea Box.	butter-bearing	J or 40 . . . Nepal	1821 C s.l	
1425. BLA^KEi4 L.	BLAKEA.	(M. Blake, a promoter of useful knowledge.)	<i>Melastbmear</i> .	2. - 12.
12098 trinervia L.	three-nerved	• Q or 8 jnjil W	Jamaica 1789. L s.p Bot mag. 451	
12099 quinquenervia Aub.	five-nerved	• o r 10 jnjil W	Trinidad 1820. C s.p Aub. gui. 1.210	

)n>Di i w i m i A MOKOOL

On) I|*ui

a

ilur^Mr w»dur»»
W. inJw)7Ai. I

ten*

i.lqittt

Hook r t IM

m./mi.
I1IL 4

BM. m*t 1*17

1 au «

Aum

Aum

Bot. DM]. 11*1*

, our of tlic tM-n»in!"

KaMtdibr. S —J

* Wtt D f-l

Mrw, • crlrtwalwl lfotln t* > wi «n 1 W(eM«.

III. — .

s . 1

• Ifitt

[ja] [Wft] [H] [t] [z] [Cw ea. 1]

KIMIM IT*. S i|, It.maiytj

1730. Sfc., *Mur. Uf:n*

12218 uftglandulsa *Dec.* single-glanded O pr 1 jn.jl W.B Mexico 1818. S Bp Cav. ic. 4.306
Clevme uniglandulosa L.

1456. HELIOCA'RPUS t. HELIOCARPUH. (*Helios*, the sun, *karpos*, fruit; radiated.) *TUftce**. 1.—2.
12219 amricanus *L.* American HO or W ♦♦ P Vera Cruz 1733. C p.l Lam. il. 409

CLASS XI. ORDER, 2.

DODECA'NDRIA DIGY'NIA.

1457. CALLrCOMA *B.* Rep. CALLICOMA. (*Kalos*, beautiful, *kome*, hair; tufted heads of fl.) *Cunonihceae*. 1.
12220 senatifulia *B.* Rep. saw-leaved ft Jor 4 my.au Y N.S.W. 1793. C s.p Bot rep. 566

1458. AGRIMONIA <i>Tou.</i>	AGRIMONY. (From <i>Argemone</i> , a plant used in cataract of the eye.) <i>Roshceee</i> . \$.
12221 <i>Exmaloria</i> <i>L.</i>	Eupatoria k A™ 3 jn.jl Y Britain bor.fi. D co Eng. bot 1335
12222 <i>nepalensis</i> <i>D. Don</i>	Nepal ^ or 3 ji.au Y Nepal 1820. D co
12223 <i>dahanica</i> IV.	Dahurian Y Dahuria 1811. D co
12224 <i>odorata</i> <i>Lam.</i>	scented Y Italy 1640. D co
12225 <i>suaviolens</i> PA.	sweet-smelling Y Virginia 1810. D co
12226 <i>pilosa</i> <i>Led.</i>	pilosè Y Siberia 1819. D co
12227 <i>repens</i> /.	creeping Y Levant 1737. D co
12228 <i>parviflora</i> <i>H. K.</i>	small-flowered Y N. Amer. 1766. D co
12229 <i>striata</i> Mx	striated-frimf W N. Amer. 1812. D co

CLASS XL ORDER 3.

DODECA'NDRIA TRIGY'NIA.

1459. AESETDA <i>L.</i>	RESEDA. (<i>Resedo</i> , to calm, to appease; medical qualities.) <i>Reudhcea</i> . 22.-2.3.
12230 <i>jesamo'ides</i> <i>L.</i>	Sesamum-like O un I jl.au Ap France 1787. S 81 ALpcd.2.8H.3
12231 <i>lithifolia Horn.</i>	Flax-leaved O or I jlau Y.a S. Europe 1819. S co
12232 <i>glauaca</i> <i>L.</i>	glauous jg iAl un I my.jl Ap S. Europe 1700. S s1 Pluk. al. 107. S
12233 <i>dipetalia H. K.</i>	two-petaled £ iQl un I & C.G. H. 1774. S si
12234 <i>rcopana Brou.</i>	Brooin-/ste tL \ un au.s Ap Tenerife 1815. S si
12235 <i>Lutentia</i> .. little yellow Dyer's weed O ag I jn.jl Ap Britain wa.gr. C s1 Eng.bot.320	yellow Dyer's weed O ag I jn.jl Ap Britain wa.gr. C s1 Eng.bot.320
12236 <i>crispala Lk.</i>	curled O un E jn.jl Ap Portugal 1823. S co
12237 <i>virescens Horn.</i>	virescent O I jl.jl.ap Spain 1820. ■ s.i Cl.h. 1.295
12238 <i>canescens IV.</i>	canescent O I my.jl Ap Spain 1597. ■ co
12239 <i>chinensis Lou.</i>	China 2 jn.jl Y.a China 1819. ■
12240 <i>mediterranea L.</i>	Mediterranean Ijns Ap Palestine 1791. ■ s.l Lind. c. 22
12241 <i>odorata L.</i>	scented <i>Mignonette</i> 1 jn.o Ap Egypt 1752. ■ r.m Bot. mag. 29
2 frut'scens	frutescent <i>Mignon.</i> tL I or 2 jn.o Ap Egypt 1752. ■ r.m Bot reg. 227
12242 <i>Phyteimia L.</i>	Phytcumia l.jn.8 Ap S. Europe 1752. ■ s.i Jac.au. 2.132
12243 <i>ramosissima Pou.</i>	branshiest 2 jn.au Ap Spain 1816. ■ 81
12244 <i>sax&tilis Pou.</i>	rock O I jn.jl.ap Spain 1816. ■ s1
12245 <i>pruina Del.</i>	frosted Y I jn.jl.ap Egypt 1824. ■ s1
12246 <i>lutea L.</i>	yellow 3 jn.jl.ap Britain ch.so. S s1 Eng. bot. 321
12247 <i>undata L.</i>	wave-leaved 1 jn.au Ap Spain 1739. S 81 Bar. ic. 587
12248 <i>filba L.</i>	upright white 3 ray.o Ap S. Europe 1596. S 81 Lob. ic. 222
12249 <i>fruticul&sa L.</i>	small-shrubby a. I J or 1 s Ap Spain 1794. C 81 Jac. ic. 3. 474
12250 <i>bipinnata TV.</i>	bipinnate-Zvf B. I J or 2 jn.au Ap Spain 1816. C 81
12251 <i>myriophylla Ten.</i>	many-leaved jt Q or 2 jl.au Italy 1824. S CO

*1460. EUPHCTRBIJ L.	SPURGE. (<i>Euphorbus</i> , physician to Juba, king of Mauritania.) <i>EuphorUacet.</i> 213.
12252 <i>uncinata Dec</i>	hooked - Z gr 1 jn.au Ap C.G.H. 1794. C s1 Dec. pl 151
12253 <i>squarrosa Haw.</i>	squarrose a. I gr Ap C. 81
12254 <i>tngbna Haw.</i>	upright triang. • ZD gr 9 jl.au Ap E. Indies 1768. C 81
12255 <i>antiquorum L.</i>	antique * 13 gr 9 jl.au Ap E. Indies 1768. C s.p R.maL2.4S
12256 <i>lactea Haw.</i>	milk-colored * ID gr 4 jlau Ap E. Indies 1804. C s.p
12257 <i>gr&ndidens Haw.</i>	great-toothed * ZD gr 5 jl Ap C. 6. H. 1823. C s.p
12258 <i>canariensis L.</i>	Canary 1 ZH gr 4 jn.n Ap Canaries 1697. C 8.p Dec. pi. 140
12259 <i>stelleap'inia Haw.</i>	Btarry.spined * ZD gr 20 mr.ap Ap C. G. H. 1824. C si
12260 <i>csrulscens Haw.</i>	csrulcbcent * ZD gr 3 ' Ap C. G. H. 1824. C 8.1
12261 <i>tetragona Haw.</i>	four-angled il ZH gr 4 ... Ap
12262 <i>heptagona L.</i>	seven-angled A 3 gr 3 jLn Ap C. G. H. 1731. ■ ap Brad. sue. 2.13
2 ritbra Haw.	Tca-sjuned H Z3 gr 4 jn.n Ap C. G. H. 1731. ■ s.p
12263 <i>enneagyna Haw.</i>	nine-angled * ZD gr 3 jl.o Ap C. G. IL 1790. ■ B.p
12264 <i>mammillaris L.</i>	teated-angled • ID gr 2 jl.au Ap C. G. II. 1759. ■ s.p Com. pr. 9
12265 <i>ccreiformis L.</i>	Cereus-formed 1 ZD cu Ap C. G. H. 1731. ■ sp Bur.af.9.3
12266 <i>officinarum L.</i>	officinal ■ IN! Africa 1597. ■ s.p Dec. pi. 77
12267 <i>odontophylla IV. en.</i>	tooth-leaved ■ C. G. H. 1824. s.p
12268 <i>polygona Haw.</i>	many-angled ft ZD gr 6 jn.jl Ap C. G. H. 1790. ■ ..p
12269 <i>ertosia Haw.</i>	eroded * J gr 3 jls" Ap C. G. H. 1805. ■ s.p
12270 <i>ernifilia L.</i>	Oleander-lvd ft ZD gr 4 jn.jl.ap India 1690. ■ s.p Dec. pi. 46
12271 <i>toxiranana Afx.</i>	poisonous ft ZD gr 20 jl.au Ap Guinea 1793. ■ s.p
12272 <i>louhogyna Lam.</i>	crested-angled ft ZD cu 3 - Ap Madagasc. 1824. ■ s.p Dec. pi. ic^.
12273 <i>hystrix Jac.</i>	porcupine ft ZJ gr 6 jn.au Ap C. G. H. 1695. ■ 8.p Jac. 8C 2.207
Trcisia hystrix Haw.	
12274 <i>vkrians Haw.</i>	varying-stmd ft ZD gr 4 ... Ap E. Indies 1800. C s.p
12275 <i>grandifolia Haw.</i>	great-leaved ft ZJ cu 6 ... Ap S. Leone 1798. C s.p
18276 <i>cucumerina W.</i>	cucumber- ^{to} tL ZD gr Ap C. G. H. 1823. C s.p VaU.it 5
12277 <i>gminifimmaJXnv.</i>	large-teated ft ZD gr :: Ap Mexico 1823. C s.p
12278 <i>lanifera Haw.</i>	wool-bearing ft I gr 3 ... Ap Mexico 182a C s.p
12279 <i>geminispina Haw.</i>	double-spined ft ZD g' 3 ... Ap Mexico 1823. C 8.p
12280 <i>melotrichia v. maini</i> M>	maini? ft ZD g' f my.s Ap G. 6. H. 1774. C s.p Bot rep. 017
12281 <i>Caput McDbael</i>	great Med. Headit-3 gr 2 au Ap Africa 1731. C s.p Com. pr. 7
Med&sma, m'jor /foif	
2 haimlis Haw.	humble - 1 au Ap C. G. H. 1768. C s.p Bur.af.8

12365	feptophylla	Pers.	fine-leaved	1	jl.s	Ap	S. Europe	1817.	S	co			
12365	pterocedca	Brat.	winged-berried	1	jn.jl	Ap	Spain	1826.	S	co			
12367	prolffra	Horn.	proliferous	1	my.jn	Ap	Nepal	1820.	D	s.p			
12368	undulata	Bieb.	waved	14	jl.au	Ap	Caucasus	1818.	D	co			
12369	LáthyriB L		Cauer Spurge	3	my.o	Ap	England	d.at.pl.	S	co			
12370	valentlna	Or.	Valentian	1*	jl.au	Ap	Spain	1804.	S	8.1			
12371	terrariaZ.		Terracina	1J	jl.au	Ap	S. Europe	1810.	S	co			
12372	diffisa W.		diffuse	i	jl.au	Ap	Austria	1798.	S	8.1			
12373	^"pios L.		Veax-rooted	#	jnjil	Ap	Candia	159a	C	s.l			
12374	larta H. K.		joyous	1	Jn.jl	Ap	- -	1758.	C	s.l			
12375	/satis Pars.			A	cu	2	jlau	Ap	Spain	1820.	D	cq	
12376	£cniKtoldes L.		Genista-like	L	J cu	1	jl.au	Ap	C. G. H.	1808.	C	sq	
12377	corifolia Lam.		Cons-leaved	t	J cu	1	jl.au	Ap	C. G. H.	1810.	C	s.p	
12378	spinbsa L.		prickly	tt.	J cu	2	my.s	Ap	Levant	1710.	C	8.1	
12379	Iruticoso L.*.		shrubby	tt.	J cu	2	my.s	Ap	Sicily	1824.	C	8.1	
12380	nnumulariBlolia W. en.	Moneyw.-lvt.		L	J CU	1	my.jn	Ap	- -	180).	C	co	
12381	/Mithymoldes L.		Epithyraum-lk	£	A or	1	my.jn	Ap	Austria	1805.	D	co	
12382	villba IV.SfK.		willoes	^	A or	2	my.jl	Ap	Hungary	1820.	D	W. & K. 1.93	
12383	dulcis L.		sweet	J	or	1	my.jn	Ap	S. Europe	1759.	D	co	
12384	purprata Thuil.		purple	^	my.jl	Ap	France	1820.	D	co			
12385	ambigua Kit.		ambiguous	^	my.jn	Ap	Hungary	1817.	D	co			
12386	carrolica Jac.		Carmolan	5	au	Ap	Carmola	1796.	D	co			
12387	angulata Jac		angled	^	jl.au	Ap	Austria	18JL	D	co			
12388	flavicomia Dec.		yellow-haired	5	jl.jlau	Ap	S. France	1820.	D	co			
12389	'ithyiisa L.		Juniper-leaved	n.	jl.jlau	Ap	S. Europe	1741.	C	s.l			
12390	portandica L.		Portland	£	jl.jlau	Ap	Britain	sea sh.	C	s.l			
12391	saxatilis Joe.		rock	£	jl.jlau	Ap	Austria	1817.	C	s.p			
12392	Cordcriftaa Dec.		Corderi's	£	jl.jlau	Ap	S. Europe	1824.	C	8.1			
12393	Paráljas L.		Paralia	£	jl.s	Ap	England	seash.	C	8.1			
12394	2 suffriticosa		sufruticose	tt.	jl.s	Ap	Europe	- -	C	8.1			
12394	petalidldcs		Petelium-like	tL	S	jl.au	Ap	1795.	C	8.1		
12394	rlgida Haw. not Bieb.												
12395	aincea // K.		rushy	£	iAJcu								
12396	cong&cta W. en.		heaiid	^	A cu								
12397	aféppica L.		Aleppo	£	AJ cu								
12398	taunnenis W. en.		Turin	^A	cu								
12399	Mgfalia L.		corn	O	un								
12400	diversifolia Desf.		various-leaved	O	or								
12401	plumbosa Hort.		feathery	O	or								
12402	csspitusa Ten.		turfy	£	A or								
12403	biuidbelfata Poir.		double-umbell.	£	A cu								
12404	angustifolia Haw.		narrow-leaved	£	A su								
12405	multicorymbosa Haw.		Provence	A	un								
12406	provncialis W.		pubescent	O	cu								
12407	pubescens Desf.		Rush-like	O	un								
12408	juncoides Haw.		narrow-leaved	£	A su								
12409	helioscpbiaZ., sun-observing		Observing	Q	w								
12410	serrata L.		Jartwort	Q	w								
12411	trunçate Pen.		narrow sav-lydy	Q	w								
12412	créticatfaw.		truncate	Q	w								
12413	punicia Swx.		Cretan hoary	Q	w								
12414	veruebsa L.		scarlet Jowlered	Q	w								
12415	corollate £.		warty	Q	w								
12416	spathulifolia Haw.		coronate	Q	w								
12417	coralhoides L.		spatula-leaved	Q	w								
12418	androsemidlia Donn.	Tutsan-lvd	Coral-like	Q	w								
12419	19 pilbsa L.		hairy	Q	w								
12420	orientiblis L.		oriental	Q	w								
12421	Vaieriünte Lam.		Valcnan-feawd	Q	w								
12422	platyphilos L.		broad-leaved	Q	w								
12423	stricta E. B.		strict	Q	w								
12424	serrulata ThuiL		serrulated	Q	w								
12425	literate L.		blotch-leaved	Q	w								
12426	denticulate Lam.		denticulate	Q	w								
12427	atl&ntica Pers.		Atlantic	Q	w								
12428	ftpleuroules Desf.		marginalized	Q	w								
12429	marginata H.&B.		long-leaved	Q	w								
13430	longitubia D. Don		Esula	Q	w								
12431	E'sula L.		wood	Q	w								
12432	ayvla tica L.		Erythrina	Q	w								
12433	Erythrina Lk.		Ncapuheten	Q	w								
12434	neapothanta Ten.		gravel	Q	w								
12435	glarebsa Bieb.		two-winged	Q	w								
12436	bilatia Lk.		Ural	Q	w								
12437	urallnsis Fis.		small-flowered	Q	w								
12438	micrantha Bieb.		curled	Q	w								
12439	crispate Horn.		joint-fruited	Q	w								
12440	condylocarpa Bieb.		berry-bearing	Q	w								
12441	fragifera Lk.		Gérard's	Q	w								
12442	Gerardino Jac.		Cypress	Q	w								
12443	Cyparissias L.		twiggy	Q	w								
12443	virgata Kit.		Myrsinites	Q	w								
12444	il/yrsinifcs L.		reflexed	Q	w								
12445	renexa Spr.		rigid	Q	w								
12446	rigida Bieb.		imbricated	tt.	cu	aus.s							
12447	imbricate Pers.		Nicean	£	AI un	1*							
12448	mcaceensis VU.		marsh	£	A un	4	my.au	Ap	Caucasus	1818.	C	s.p	
12449	palustris L.		emarginate	£	A un	2	jl.au	Ap	Portugal	1804.	C	if	
12450	emarginata H. K.		Irish	£	A cu	1	my.jn	Ap	Spain	1809.	D	s.1	
12451	hibcmica W.		Willow-leaved	£	A un	2	my.jn	Ap	Sweden	1570.	D	8.1	
12452	sahcifolia Hort.		Almond-like	tt.	mr.jn	2	mr.jn	Ap	Italy	1758.	I	co	
12453	! S S y g. s		Characias	n.	pr	4	mr.jn	Ap	Britain	fields.	I	co	
12454	Characias L.		Venetian	tt.	J cu	1	mr.my	Ap	Hungary	1804.	D	co	
12455	venea L.		glaucescent	£	A un	1	mr.jn	Ap	England	woods.	C	co	
18456	glauceicens IF.								Europe	1820.	C	pi	
										1823.	D	co

Eng. bot 2255

Jac. ic. 1.88

Jac. ic. 1.87

Lob. ic. 361

Den. br. 45

Biv. cen. 1.35

Bot rep. 616

W. & K. 1.93

Jac. au. 3.213

W. & K. 135

Scop. car. 21

Jac. ic. 481

Boc. sic. 5

Eng. bot 441

Pa. it. ger. 1

Jac. sc. 1.107

Jac. sc. a 483

Eng. bot 883

Jac. ic. a 483

Gm. si. 2.93

Jac. au. 376

M.h. 10.3.3

Bot. cab. 390

Eng. bot. 333

Jac. ic. 3.482

Bux. c. 2.27,*

Des. at. 1.103

Eng. bot. 1399

Jac. au. 436

Bux.c.2.25

Jac. i.c. 3.4

Fl. dan. 866

Eng. bot \$?

Bot. cab. 973

Eng. bot ^

Eng. bot • »

CLASS XI. ORDER 4.

DODECA'NDRIA TETEAGY'NIA.

IOJIS³. CALLIGONUM L. CALLIGOXUM. (*Kolas*, beautiful, *gonu*, joint j joinU instead of Its.) ~~277~~ 12472 PaUasia H. K. Pallas's tt. or 4 au G.w Casp.Sea 1780. C s.l PaL KM. 2.77,78

CLASS XI. ORDER 5.

DODECA'NDRIA PENTAGY'NIA.

191J?> GLrNUS L 12473 /otoldes L	GLINHS. Lotus-like	(A name given by Theophrastus to the maple.)	FUoide*. 1.—5.
	iQJuf iijl	Y	S. Europe 1788. S 81 Boc.sic.2LII
IS? ? b,f ACKWE'LLW Com.	BLACKWEJXIA.	(Elizabeth BlackweU, bot. artist)	<i>Homaline*</i> 5.—8.
1245 ax H ₁ h r18 *-<>-	axillary	■ 4	W Madagasc.1824. C s.p Lam. il. 412.1
	spiral	■ or 4	W E Indies 182a C s.p
	panicled	■ or 6	W Bourbon 1820. C s.p
	entire-leaved	■ or 6	W Madagasc.1823. C s.p Lafn.il.412.2
K ham. grandiflora2. f>Mtf.	large-leaved	■ U or 5 au	W China 1824. C s.p
iolst+ G ASTO'N X/J. J. GASTONIA. (Gaston de Bourbon, son of Henry IV., patron of bot)	Araliacat. 1.—2.		
<<479 palmata Wai.	palmate	• o r 4 f.mr	W.o 1818. C p.l Botreg.894

CLASS XI. ORDER 6.

DODECA'NDRIA HEXAGY'NIA.

.1467. CEPHALOTUS soft. CEPHALOTUS. " (Kephalotos, headed; glandular-headed stam.) *RosaetB. I.*
18480 follicularis Lab. Collided £ L A] C U 1 ... W N. Holl. 1822. S bog Lab. n. h. 2.145

CLASS XI. ORDER 7.

DODECA'NDRIA DODECAGY'NIA.

J ^{AN} U ^{ARY}	V ^{OLUME}	L. HOUSELEEK.	<i>(Sempervivum to live for ever; tenacit;</i>	Crasmaceae.	29-32.
arb ^{rtum} L.	tree	J a	9 9mr.d d Y Y	Levant	1640. C s.l Bot. reg. 99
^482 cananense L.	variegated-wf	• 1 or	4 mr.d Y	Levant	1640. C s.l
12483 frutescens Haw.	Canary	«. 1 or	3+jn!!. Y	Canaries	1699. C 81 Dec. pL 141
2 ² 4g Jutini, sum H.Jjr.	frument	tt. i. Jor	; jlau	Teneriffe	1824. C s.l
S S ^ C	g)andulob-fMi tt. L or	; mr.my	Madeira	1777. C s.l Bot. mag. 1963	
i 2 hf bridum Haw.	ciliated	n. L J or	W	Madeira	1777. C s.l
i «87 cespitosum Spr.	hybrid	«. t lor	jlau	Teneriffe	1815. C s.l
19- barbktum Horn.	turfy	■■■■■ vap.s	Y	hybrid	1820. C 81
*m barbktum C. Sm.	ciliatum B. M.			Madeira	1815. C s.l Bot. mag. 1978
lot*** ^ineolkre Haw.	bearded	tt-Ljor	1 jlau	Canaries	1815. C s.l
ISSS Smithii Sims	Smith's	tt.Ljor	jlau	Teneriffe	1815. C s.l
*** ^a aereum C. Sm.	golden	tt.i or	jlau	Canaries	1815. C s.l Bot. mag. 1980
calyciforme Haw.					
8 ipiurium Haw.	spurious	tt-Ljor	1 jlau	Canaries	1820. C s.l

12401	drbicum	<i>C. Sm.</i>	city	St. i_	or	2	jn.au	Y	Teneriflē	1816.	S	s.l
12492	rvtūHum	<i>Haw.</i>	tow retuic	«-L-Jor		2	jnau	Y	Teneriflē	1824.	S	al
IMyitubilitufirinc	<i>Haw.</i>	table-shaped	e. <_J or			1	ll.ou	Pa.Y	Tencnlle	1817.	C	rl
12494	tortuutfum	//. K.	twisted	«.l J			J.lau	Y	Canaries	1779.	C	s.l
12495	villsnu	//. K.	villous	a. r.-l			Jjtijl	Y	Madeira	1777.	D	8.1
124M	Mxian	<i>Haw.</i>	los>c			i	jl uu	Y			
12417	stellatum	<i>Sm.</i>	starred	•Pl	or		i jl.au	Y	M. Baldo	1790.	D	s.l
	villoHuin	<i>B M.</i>									Seg.	ver. 2.17
12498	tectfrum	<i>L.</i>	roof	If.	A or	1	jn.s	F	Britain	roofs.	D	s.p
12499	of nctauni	<i>Haw.</i>	African	t	AJ				.C. G. H.	1768.	D	s.l
12500	dodrantalc	IF. en.	nine-inch			1	jl.au	Pk	Tencriffē	1815.	S	s.I
12501	flagellifonne	/5.	whl>fonned	£	A or		Jjlau	R	Siberia	1823.	C	s.I
12502	hirtmtn	<i>A.</i>	hairy	£	A or	1	injl	W	Italy	1804.	D	s.p
12501	SONHOHTUMI	# M.	Boholiforous	£	A or	1	injl	P.i.Y	Germany	...	I)	s.p
12504	Klubitvruin	/.	globiferous	f	A or			Y	Germany	1731.	I)	s.p
12505	KramilKTium	<i>Haw.</i>	great-Huwrcrd	£	A or			Pa.lt	Europe	1731.	D	s.l
	Klomllerterum	<i>B. M.</i>	not L. IV.								Bot.iuaK.JU7.2H5	
12506	inoit müui	<i>L.</i>	mountain	£	A or	i	jn.il	R	Switicrl.	1752.	1)	s.l
22507	vuadrihuim	<i>Haw.</i>	cubpiiatc	£	—	1	jnjl	R	Siberia	1810.	D	s.l
12508	pñnnilum	<i>Bieb.</i>	dwarf	£	A or	1	Jn.jl	Pa.R	Caucasus	1824.	1)	s.l
12509	arachnofdcuuii	<i>L.</i>	cobwebbed	£	A or	i	Jnjl	K	.Italy	1699.	D	s.p
1469.	MONA'NTHES	//aw.	MONANTHBB.						(Monot, one, anthos, a flower.)			Crassuticta?
12510	Mlyphylla	<i>Haw.</i>	many-leaved	tf	IAICU							1.
	ScinpcryWum	Monantho* <i>b.</i>		ft	jls			R	Canaries	1777.	D	s.l
											Bot.	mag. 93

CLASS XII. ORDER 1.

ICOSA'NDRIA MONOGY'NIA.

- | | |
|--|--|
| •1470. CA'CTUS L. MELON THISTLE. (A name applied by Theophrastus to a spiny plant) | <i>Opun/idcea*</i> . 12.- |
| 12511 Meloc&etus L. Melon Turk's CapfL | W. Indies 1188. O s.p. Dec.pl. T |
| 12512 macrac&ntha Haw. long-spined | S. Anicr. 1820. O s.p. |
| 12513 deprissu Dec. depressed | S. Amer. 1798. O s.p. |
| 12514 parvisp'ma Haw. small-spined | W. Indies 181#. O s.p. |
| 12515 uru/Acca Haw. hedgehog | 1818. O s.p. |
| 12516 HCUHH. Haw. old | Mexico 1823. O s.p. |
| S12517 latiflrina Haw. broad-spined | Mexico 1823. O s.p. |
| 12518 redicta Spr. restored | Mexico 1796. O fl.p |
| nAbilis Haw. | |
| U2519 n6bilis L. noble | 1796. O e.p |
| rrecarva Mil. | |
| 12521 corrugata GUI. twisted | I jn.au Anticua 1768. O s.p. |
| 12522 foliosa GUI. corrugated | Chile 1824. O fl.p |
| leafy | Chile 1624. O s.p. |
| •1471. MAMMILLAMUA/Jaw. MAMMILLA HI A. | {Mamma, the nipple; tubercles.) <i>Optntidcear.</i> 22.- |
| 12523 simplex Haw. simple red-spined n. "H ur | I jn.au W W. Indies 1688. O s.p. Dec. pl 111 |
| Cltcr* mammillaris L. | |
| 12524 prolifera Haw. white-mined prolific. n. *-Jpr | I jn.au S. Amer. 1800. O •p |
| 12525 discolor Haw. vanous-coloredtt. ZD K | t jn.au R 1800. O •p |
| Cactus deppressa Dec. | |
| 12526 straminca Haw. stnv/spinrid a. Igr | i jn.au . 1811. O s.p. |
| 12527 flavescens Haw. AavaccinUspined. 3 K | i jn.au Y . 1811. O •p |
| 12528 cónica Haw. conc-heedsd tt. 3 gr | i jn.au . 1808. O s.p. |
| 12529 parvimámma Haw. small-teatil tt. Igr | i Jn.au . 1817. O s.p. |
| 12530 mafriim&nuna/firjw. large-tea ted < " gr | my.jn Pk Mexico 1893. O ■ |
| 12531 stflttaii Haw. starred tt. ZD gr | S Amer. 1815. O s.p. Bot cab. 79 |
| Cactus scellata JK | |
| 12532 geminiHpina Haw. two-spined | R Mexico 1823. O s.p. |
| 12533 coronata Ma G Don | S. Amer. 1820. O s.p. |
| Cactus coronata W. | |
| 12534 coronaria Haw. garland | 5 Jn.au 6 S. Amer. 1817. O s.p. |
| 12535 vivipara Ph. viviparous | Louisiana 1811. O s.p. |
| 12536 lanifera/nw. | s.p. |
| Cactus glomeriflta Lam. | |
| ✓SSH missounensis Swt. Missouri | t ... II Mexico 1823. O s.p. |
| Cactus manunillaris Nut. | |
| 12538 pusilla Swt. | i jn.au .. Missouri 1818. O s.p. |
| Cactus pusilla Dec. | |
| 12539 ambigua G. Don | i jn.au . 1820. O s.p. |
| Cactus ambiffua Gill. | |
| 12540 spinida G. Don | spiny Chile 1827. O s.p. |
| Cactus spi n6sa Gill. | Chile 1827. O s.p. |
| 12541 quadrata G. Don | quadrata Chile 1827. O s.p. |
| Cfetus quadrata Gill. | |
| 12542 caudata G. Don | tailed Chile 1827. O •P |
| ioe^o " caudata Gill.///. | |
| 18543 coccinea G. Don | scarlet Chile 1827. O s.p. |
| mm Cactui ooccinea Gill. | |
| 12544 solitaria G. Don | solitary Chile 1827. O s.p. |
| Cactus solitaria UIC//. " | |

13618 <i>frigida</i> <i>Ua Nut.</i>	brittle	<..	J gr		-	-	N. Amer. 1814. C s.p
12619 <i>polyacantha</i> <i>Haw.</i>	many-spined				-	-	N. Amer. 1814. C s.p
Cactus ft-rox JV.K.							
12fi20 <i>undia</i> //».	middle	<i>many-spineilit-</i>			-	-	N. Amer. 1814. C ip
12W1 <i>tilia</i> -a <i>flnw.</i>	loafy	<..			Y		S. Amer. 1H17. C »p
12iJ <i>J</i> <i>pili</i> - <i>Unto</i> .	small	<..			Y		S. Amer. 1805. C s.p
li <i>OJ</i> <i>Hilophrca</i> <i>G. Don</i>	sulphur-co/ore/*/*				Y		Chile 1827. C s.p
6 <i>ctUM</i> <i>sulphrca</i> <i>Gill.</i>							
<i>scrifica</i> <i>tf. Don</i>	silky	<..			E	Chile 1827. C s.p	
Cactus <i>scrifica</i> <i>Gill.</i>							
* <i>Iffl. RHIP'SALIS Gae.</i> <i>RHIPSALIS</i> . (<i>/flips</i> , a willow branch : flexible branches.) <i>Cacti.</i> 6.							
12625 <i>Cahsutha Gar.</i>	<i>Cassutha</i>	tt.	Z3 cu I S		YY	VWVW <i>Indie</i> 728. C s.p	Hook. ex. fl. SI
Cactus <i>pendiila</i> <i>Stoz.</i>							
12626 <i>har.witruh</i> <i>Haw.</i>	iKirnsitic				Y	S. Amer. 1800. E	Dec. pi. 59
12627 <i>grandilK.rus</i> <i>Haw.</i>	large-flowered				W	& Amer. 1818. E	Bot. mag. 2740
<i>far'tus finiMn</i> <i>Stilm</i>							
12fi28 <i>F'sifirur>ra</i> <i>Htiw</i>	bundled				Gsh.W	Brazil 1817. E s.p	Rot mag .3079
r'C29 II <i>ognjidi's//III.</i> Saltwort-like					Y	E. In.lics 1817. E s.p	Bot. max. 24fil
12630 <i>iiwei!iniiryaiithoides//rtii;</i> <i>Mesem.-lktL</i>	cu				W	S.Amer. 1817. E s.p	But. mag. 3078
Cactus <i>portulacifolia</i> <i>L.</i>							
• 1476. PERE'SK/* Haw. BARB A DOES GOOSEHERRY. (N.F. <i>Pcireskius</i> . Provence, lover of bot.) <i>OjnnitacetB.</i> 4.							
126J1 <i>aculeita</i> <i>Haw.</i>	prickly	* 13	ft 5 o.n		W	W. Indies 1696. %M> Di. el. 2tf. 294	
6actUh <i>VerCukia</i> <i>L.</i>							
12& Krahil'Mia <i>Haw.</i>	large-loaved	* Z3 fr				Brazil 1818. c «P	
12 <i>23</i> <i>loiiKiNp.na</i> <i>Haw.</i>	long-Npinced	H Z3 "				S. Amer. 1808. C a.p	
19+& HJM<rtulaiwn)i.tif. Do// Portulaca-lvd" Z3 ft 3 ...						W. Indies 1820. C s.p	
Cactus <i>portulacifolia</i> <i>L.</i>							
* 1477. BAHTO*N/if Sims BARTON FA. (Dr. Z. S. Barton, professor of botany at Philadelphia.) <i>Lodsr*</i> . 9.							
12 <i>N5</i> <i>onuta</i> <i>Ph.</i>	ornamented	& OJ or	2 jl.s		W	Missouri 1811. S «P Bot. mag. 1487	
decapetalta B. M.							
196J6 <i>nula</i> <i>Ph.</i>	naked-seeled	^ OJ or	2 jl.s		W	Missouri 1811. S s.p	
1478. MEKTZVLU L. <i>MENTZBLIA.</i> (C. <i>Menzel</i> , of Brandenburg; bot author.) <i>Lothe*</i> . 3.							
12 <i>V7</i> <i>fispera</i> /.	rough	tL CD or	H jl.au	Y	America 1733. S co	Pl. ic. 174. 1	
12 <i>U8</i> <i>oliKospcrma</i> <i>Sims</i>	few-beedded	E LAJ or	2 my.jn	Y	Louisiana 1812. S s.l	Bot. mag. 1760	
12 <i>oJ</i> <i>hispida</i> >f.	hispid	LAJor	I jn.jl	Y	Mexico 1820. S s.l	Cav.ic. 1.70	
• 1479. i'HILADE'LPHUS L. <i>SYRINGA.</i> (Used by Athenams for a tree now unknown.) <i>PhtiaiUlpccce.</i> 6.							
12640 <i>coronarinH L.</i>	garland		or 8	my.jn	W	S. Europe 1596. L co	Bot. mag. 391
2 nafius Mil.	dwarf		or 2	my.jn	W		L co
3 fibre pltno	double-flwd		or 8	my.jn	W	S. Europe ...	L co
4 ftilis variegatis	variegated-lvd		or 8	my.jn	W	3. Europe ...	L co
12641 <i>grclilis</i> <i>Lo. C.</i>	slender		or 3	mr.jn	W	N. Amer. 1820. li co	
12641 <i>i</i> <i>nepalmis</i> <i>Wat.</i>	Nepal		or 6	jnjl	W	Nepal 1H22. L co	
Y2CAS. inod->ruh Mil.	scentless		or 2	jnjl	W	Carolina 17J8. J co	Bot. mag. 1478
ISM44 <i>granvittTua IV.</i>	large-flowered		or G	jnjl	W	Carolina 1H11. L co	Bot. reg. 570
12645 <i>hirsutus</i> <i>Nut.</i>	hairy		or 3	jn	W	N. Amer. 1820. L co	Dcn.br. 47
1480. LEPTOSPF/RMUM <i>Forst.</i> <i>LEPTOSPERMUM.</i> (<i>Leptos</i> , slender, sprma, seed.) <i>Myrtlcete.</i> 29.							
12646 <i>grandiflmuin Sm.</i>	large-leaved	ft i	I or 5	jijl	W	N. S. V. 1803. C p 1	Bot. mag. 1810
12647 grand iflorum <i>Loil.</i>	larKe-flowered		or	jijl	W	N. S. W. 1810. C p 1	Bot. cab. 514
12648 <i>cmarginatum</i> <i>Wnl.</i>	cinarginate		or	jijl	W	N. S. W. 1818.	
12649 sericeuin <i>Lab.</i>	silky		or	jnjl	W	N. S. W. 1818.	E P.
12650 marginatum <i>Lab.</i>	margined		or	jnjl	W	N. S. W. 1820.	E P.
12651 pubescenM <i>Sm.</i>	pubescent		or	jnjl	W	N. S. W. 1774.	E P.
12632 lanigerum <i>Sm.</i>	woolly		or	jnjl	W	N. S. W. 1774.	Bot Cab. 1192
12653 fcoparium <i>Forst.</i>	broom N Zeal. <i>Taitt</i>	or	jnjl	W	New Zeal. 1772.		Bot. rep. 622
12654 tuberculatum <i>Pair.</i>	tuberculatu		or	jnjl	W	N. S. W. 1816.	s.p
12655 attenuatuin <i>Stu.</i>	attenuated	ft I	J or	myjl	W	N. S. W. 17U5.	E
12 <i>56</i> ruhricafile <i>Vcn.</i>	red-btemmed	il LJ	or	jnjl	W	N. S. W. 1817.	E p.
12 <i>57</i> poroprh <i>Hum Cav.</i>	porous-leaved	ft	or	myjl	W	N. S. W. 1800.	E
12658 flexuobum <i>Lk.</i>	flexuose	ft	or	myjl	W	N. S. W. 1823.	p. Col. h. rip.app.l
12659 flavscens <i>Sm.</i>	flavcscent	il L J	or	myjl	W	N. S. W. 1787.	Bot. mag. 2695
12 <i>60</i> triloculK-re <i>Ven.</i>	trilocular	il I	or	jnjl	W	N. S. W. 1800.	Bot. cab. 791
12 <i>61</i> jimpennuin <i>Sm.</i>	iun\\eT-leaved	ft J	or	jnjl	W	N. S. W. 1790.	Vcn. mal. 89
12 <i>62</i> bacchini <i>Sm.</i>	herry-fruited	il J	or	jnjl	W	N. S. W. 1790.	Cav. ic. 4. 331. <
12663 ararhnofdcum <i>Sm.</i>	cobwebbed	il I J	or	myjl	W	N. S. W. 1795.	p. Gae. fir. 1.35
12 <i>64</i> pariflT>huin <i>Sm.</i>	small-leaved	si I	or	5 jnjl	W	N. S. W. 1789.	p.
12 <i>65</i> Helleatum <i>Cav.</i>	starred	sH LJ	or	jnjl	Y	N. S. W. 1790.	E P. Cav. ic. 4. 330
VJIM <i>i</i> imbricatum <i>Sm.</i>	imbricated	• i I	or	5 jnjl	W	N. S. W. 1823.	D
12 <i>67</i> obtitum <i>Lo. C.</i>	othtuasayn	il I J	or	jnjl	W	N. S. W. 1820.	s.P
J2668 multicafic <i>Cun.</i>	many-stcmmed	sH I	or	4 jnjl	W	N. S. W. 1824.	S.D
12669 /yminfdhum <i>Cun.</i>	Thyme-leaved	• I	or	5 jnjl	W	N. S. W. 1824.	
12070 amblguum 5m.	ambiguous	il t	or	3	I W	N. S. W. 1791.	E Pl. Ex. bot. 1.69
Metrosideros corifolius <i>Ven.</i>							
12671 ovatifolium <i>Swt.</i>	ovate-leaved	s* LJ or	5	■	■	N. Holl. C s.p.1	
I-abricit myrtifolia 1ia <i>Hort.</i> not <i>W.</i>							
12672 squarrösom <i>Sicb.</i>	squareose	* i	or	4 jn.au	W	N. Holl. C p.l	
12673 pendulum <i>Sicb.</i>	pendulous	ft i	or	4 jn.au	W	N. Holl. C p.l	
1481. FABRI'CM Gae. <i>FABRICIA.</i> (J. C. <i>Fabric</i> 'ms, the celebratil Danish entomologist.) <i>Nyrthcea.</i> &							
1&.74myrtifolia <i>Gae.</i>	Myrtle-leaved	i j or	3	... Y	N. Holl. ... C s.p	Gae. fr.355.4	
12675 hevigiita <i>Sm.</i>	smooth.leaved	* i J or	3	myjn Y	N. S. W. 1788. C sp	Bot. mag. 1304	
12676 sericca <i>Nats.</i>	silky	i j or	2	... Y	N. Holl. 1820. C s.p		
12677 stricta <i>Lod.</i>	strict	»i J or	3	ap.jn. ... N. Holl. 1827. C s.l		Bot. cab. 1219	
14R2. METROSONDE^ROS							
1SS? ^Irils lhill,							
WS79 hlsimlus <i>Sm.</i>							
12080 floribiius 5m.							
12 <i>61</i> costatiu <i>Uac</i>							
12 <i>62</i> glomiil'ircr <i>Sm.</i>							
12683 angustifolius <i>Sm.</i>							

12684 <i>fisperus Ma. C.</i>	rough	• i_J or 5	—	N. Holl.	1824.	C s.l
12685 <i>capiUtus Stn.</i>	capitate	M i_J or 5	—	Pk	N. Holl.	1824. C s.l
• 1483. CALLISTEWON R. Br. CALLISTRMON. (<i>Kallislot</i> , most beautiful, <i>stemon</i> , <i>stam.</i>) <i>Hfyrt&ce*</i> . 14.—18.						
ISitifl upccidus &v.' showy		*	—	10	nir.jn	C N. S. W. 1823. C s.l Hot mag. 1761
Metrodor roN spocioaus B. M.						
12687 <i>vnninalis G. Don</i>	twiggy	■ _Jor	10	mr.Jn	R	N. & W. 1800. C s.l Gae. ft. 1.34. 4
Metrosidéros virainalis <i>Gae.</i>						
12*188 <i>lancifftus Swt.</i>	spear-leaved	■ _Jor	10	Jn.n	C	N. S. W. 1788. C l.l Bot. mag. 260
Metrosid <i>-re</i> cltrinus <i>H. M.</i>						
12*58) <i>plungens G.Don</i>	pungent	■ H_Jor	6	my.jn	—	N. aw. 1827. C 8.1
199U <i>rigidus li. Br.</i>	rigid	■ uJor	5	np.my	—	IH00. C s.l Dot reg. 393
12<3 J Ine^ris. Vin.	linear	■ L'Jor	G	jn.jl	—	N. Holl. 1788. C s.l Wen. han. 19.11
12*52 <i>InicanMiu- lk.</i>	linear-leaved	■ (ir	10	mr.jn	K	N. S. W. 1820. C p.l
12*53 <i>vindifloriu. SW.</i>	green-flowered	■ Jor	5	jn.au	G	N. S. W. 1818. C s.l Bot. mag. 2602
K' ■ Pine-leaved		■ L'Jor	6»J.	G	N. Holl. —	C s.l Wen. c. 1.16
K' ■ margin'itu* Swt.	margined	■ L'Jor	5	—	P.Y	N. S. W. 1824. C s.p
■ forinmuis <i>Ma. C.</i>	handsome	■ L'Jor	5	—	—	N. S. W. 1824. C rp
■ microph^llus Mil. 6*	small-leaved	■ L'Jor	5	—	—	N. Holl. 1824. C rp
12*W <i>seinopforiens 7.0.C.</i>	ever-blowing	■ L'Jor	6	mr.jn	C	N. Holl. 1818. C p.l Bot cab. £23
12699 <i>riiKiiKius G. Don.</i>	small wrinkly	■ L'Jor	6	mr.jn	Pk	N. S. W. 1821. C p.l
Metrosidi'TOs rugulbus <i>W.</i>						N. S. W.
12700 <i>Inphanthiui 5m/.</i>	crest-flowered	■ i_Jor	6	jn.au	G.v	N. Holl. 1800. C s.l Sw.au. 29
MetrosiU ros lophanthus <i>Vent. salijnu8~2j. M.</i>						
12701 <i>leptostachys Sivt.</i>	sciider-bpked	M _ or	6	jn.au	G.v	N. Holl. 1820. C s.l
• 1484. PSI'DIUM L.	GUJAWA. (<i>Psilia</i> , a Greek name of the pomegranate.) <i>Myrtlece</i> . 16.—22.					
12702 <i>pyrifcrum L.</i>	pear-bearing	* C 3 f r	7	jn.jl	W	W. Indies Ih6. C r.m Bot. reg. 1079
12703 <i>LitifMuin Lk.</i>	broad-leaved	■ ■ ■	10	h.jl	W	S. Amor. 1800. C r.m
J2704 <i>poinireron L.</i>	apple-bearing	■ ■ ■	10	jn.jl	W	W. Indies IW2. C r.m II.mal.3.35
17or> h.ipidisiiuum <i>Jiic.</i>	most savoury	■ ■ ■	10	jn.jl	W	C r.m
1*70; J guineense <i>Swx.</i>	Guinea	■ ■ ■	8	jn.jl	W	Guinea 1812. C r.m
1-707 <i>arboindticum Aub.</i>	aromatic	■ ■ ■	5	...	W	W. Indies 1779. C r.m
12708 <i>grandiflurum Aub.</i>	great-flowered	■ ■ ■	8	jla	W	(Juiana IKX) C r.m Aub. gui. 1.191
W70> <i>guian^nae Rich.</i>	(■ ■ ■)	■ ■ ■	5	jla	W	Guiana 1822. C r.m
1*710 <i>fragrans Lo. C.</i>	fragrant	■ ■ ■	8	ap.au	W	Guiana 1823. C r.m
1/711 <i>conlMum B. M.</i>	cordate	■ ■ ■	5	my.jl	W	W. Indies 1811. C r.m
1'-712 <i>nionliuin Suas.</i>	mountain	■ ■ ■	—	W	W. Indies 1779. C r.m	
12713 <i>polyedrpum Andr.</i>	many-fruited	■ ■ ■	3	my	W	Trinidad 1810. C r.m
1-714 <i>CattkMs/Ms/ Lindl.</i>	Cattley's	■ ■ ■	20	mr.jn	S. Amer. 1818. C r.m Lind.col. 16	
12715 <i>wiyriffr.liuin Lo. C.</i>	Myrtle-leaved	■ ■ ■	<J	fjl	—	C r.m
12716 <i>oligosp^rmum Hort.</i>	few-seeded	■ ■ ■	10	...	W	1817. C r.m
12717 <i>chineike Lo. C.</i>	Chinese	■ ■ ■	6	...	W	1820. C r.m
						China
1485. NE'LITKIS Gae. NELITIIIS. (<i>Ne</i> , priv. <i>elytron</i> , seed-case; berry unpartitioned.) <i>Myrtheete</i> . 1.—3.						
12718 <i>JambostUa Gae.</i>	Jambosella	• • fr	10	...	W	Society Is. 1810. C s.p Gae.fr. 27.5
• Psidium decaspemnum <i>L.JU.</i>						
1480. MELANOPSI'DIUM Col. MELANOPSIDIIM. (<i>Melanos</i> , black, <i>psidiu</i> nt, <i>guava</i> .) <i>Rubiaceaf</i> . 1.						
12719 <i>nlgrum 6W.</i>	black	\$ • or	12	...	Crea.	...
						1828. C s.p Col. h. rip. 35
■ ■ ■	EUGENIA. (<i>Prince Eugene</i> , of Savoy, patron of botany.) <i>Myrtacat</i> . 25.—87.					
12720 <i>javánira</i>	Java	« D f r	10*	...	W	Java IH21. C s.p Hu. am. 1. 38.2
12721 <i>malactnsis L.</i>	Malay Apple tree	■ ■ ■	25	my.au	S	E. Indies 1768. C s.p Bot rep. 458
12722 <i>marrarpa Rox.</i>	long-fruited	■ ■ ■	20	...	W	E. Indies 1R22. C sp
12723 <i>glandulifera Rox.</i>	glanduliferous	■ ■ ■	10	...	W	E. Indies 1821. C s.p
12724 <i>Rox.</i>	pretty	■ ■ ■	10	...	W	E. Indies IHVtt. C Bp.
12725 <i>Rox.</i>	watery	■ ■ ■	20	...	W	E. Indies 1820. C s.p
12726 <i>in'at a Rox.</i>	acuteinate	■ ■ ■	10	myjl	W	E. Indies 1820. C s.p
12727 <i>temifMiji Rox.</i>	three-leaved	■ ■ ■	—	W	E. Indies 1822. C s.p	
12728 <i>obtusifolia Rox.</i>	obtu AC-leaved	■ ■ ■	—	W	E. Indies 1821. C 8p	
12729 <i>amplexicaths Rox.</i>	utoin-rlasping	■ ■ ■	myjl	W	E. Indies 1823. C sp Bot.reg. 1033	
12730 <i>ceraso^les Rox.</i>	Cherry-like	■ ■ ■	—	W	E. Indies 1822. C s.p	
12781 <i>sumairana Lo. C.</i>	Sumatra	■ ■ ■	—	W	Sumatra 1823. C s.p	
S12732 <i>Jambos L.</i>	Jainbos	■ ■ ■	fjl	Gv	E. Indies 1708. C sp Bot.mag. 1696	
12733 <i>baruensis Jac.</i>	Baruan	■ ■ ■	—	W	S. Amer. — C s.p Jac. ic. 3.486	
VSIM axillaris <i>W.</i>	axillary	■ ■ ■	—	W	Jamaica 1793. C s.p	
12735 <i>fragrans JV.</i>	sweet-scented	■ ■ ■	ap.my	W	Jamaica 17M. C s.p Bot. mag. 1212	
12716 <i>Mini Aub.</i>	Mini	■ ■ ■	—	W	Guiana 1803. C s.p Aub. gui. 1. 197	
12737 <i>elliptica Sm.</i>	elliptic-/; uifed	■ ■ ■	my.s	W	N. S. W. 1790. C s.p Bot. mag. 1872	
12738 <i>Myrtle-leaved</i>	i	■ ■ ■	apjl	W	N. Holl. 1818. C s.p Bot. reg. 627	
JV/rutus australis <i>B. M.</i>						
12739 <i>ig6strina IV.</i>	Vivet-leaved	■ ■ ■	8 au	W	Hispaniol. 1708. C s.p	
J*78 "niflaraL.	one-flowered	■ ■ ■	7 ja.mr	W	Brazil 1759. C s.p Bot. mag. 473	
*J2741 zycl&nica <i>W.</i>	Ceylon	■ ■ ■	10 jn.jl	W	Ceylon 17W. C s.p Bot. rep. 619	
12742 <i>biiUimica Wnl.</i>	balsam	■ ■ ■	—	W	— 1816. C s.p	
12743 <i>latitolia Aub.</i>	broad-leaved	■ ■ ■	—	W	Guiana 1793. C s.p Aub. gui. 1. 199	
12744 <i>ractita Raeu.</i>	bracteat	■ ■ ■	—	W	E. Indies 1820. C s.p	
12745 <i>Vincop^laLflm.</i>	long-leaved	■ ■ ■	—	W	E. Indies IHV0. C s.p	
JS1 J xl filia <i>W.</i>	Box-leaved	■ ■ ■	my.jn	W	W. Indies 1818. C s.p	
12747 <i>flombinda West</i>	bundle-flowered	■ ■ ■	—	W	W. Indies 1820. C sp	
12748 <i>obovata Poir.</i>	obovate	■ ■ ■	—	W	I. France 1823. C s.p	
12749 <i>glauca Fis.</i>	glaucous	■ ■ ■	—	W	— 1822. C s.p	
12750 <i>sincmarif nsis Aub.</i>	Sinemaria	■ ■ ■	myjl	W	Guiana 1823. C fl.p Aub. gui. 1.198	
12751 <i>auriculMa IM. C.</i>	small-cured	■ ■ ■	—	W	— 1825. C s.p	
12752 <i>pegalinom'ids Lo. C.</i>	Pctalamia-like	■ ■ ■	—	W	E. Indies 1825. C s.p	
12753 <i>gracilis Lo. C.</i>	slender	■ ■ ■	—	W	E. Indies 1825. C Bp	
12754 <i>rugfts Lo. C.</i>	wrinkly	■ ■ ■	—	W	E. Indies 1824. C s.p	
, 1488. WARXYOPHV'LLUS /... CLOVB THKE. (<i>Karyon</i> , nut, <i>phyllon</i> , & leaf; <i>lliids</i> of flowers.) <i>Myrtacc</i> . 1.						
12755 <i>aroraaticus L.</i>	aromatic	± • cul i	10	...	Y.w	Moluccas 1797. C l.p Dot.mag.2749.2750
* 1489. AfY'nTUS L.	MYRTLE. (<i>Myron</i> , sweet ointment; fragrance.) <i>Myrtuccce</i> . 14—55.					
* 12756 coinnfus L.	common	* LJ or	6	jl.au	W	S. Europe 1597. C r.m Huh.no. 43
1 romana	Roman	* LJ or	6	jl.au	W	S. Europe 1597. C r.m Mil. ic. 184 1

ICOSANDRIA MONOGYNIA.

CLASS XII.

fitarenfina	Tarentine	• L-J or	6	jLau	W	S. Europe	1597.	C	r.m
3 italicica	Italian	* LJ or	6	jLau	W	S. Europe	1597.	C	r.m
4ba?rica	Batic	* L-J or	6	jLau	W	& Europe	1597.	C	r.m Black. 114
5 lusitanica	Portuguese	* L-J or	6	jLau	W	S. Europe	1597.	C	r.m a. h. 11
6 belgica	broad-ivDutchm	i-J or	2	jLau	W	& Europe	1597.	C	r.m
7 mucronata	pointed-leaved	* t-J or	6	jLau	W	S. Europe	1597.	C	r.m
8 fibre pleno	double-flowered	L-J or	6	jLau	W	S. Europe	1597.	C	r.m
9 variegata	variegated	■■■■■	6*	jLau	W	a Europe	1597.	C	r.m
10 maculata	spotted	■■■■■	5	... P	W	N. Holl.	1824.	C	s.p
12757 tenuifolia Sm.	lnc-leaved	■■■■■	6 jn.jl	P	China	1776.	C	s.p	Bot mag. 250
12758 tomentosa H. K.	vroo Wy-leaved	■■■■■	fi in.jl	P	China	1823.	C	s.p	
12759 affinis Hart.	kindred	■■■■■	10 ap.my	W	Jamaica	1759.	C	s.p	Br.jam.25.3
12760 bifora L.	two-flowered	■■■■■	5 ...	W	N. Holl.	1824.	C	s.p	
12761 trinervis Sm.	thrc-ncrvcd	■■■■■	6 ...	W	Surinam	1793.	C	s.p	
12762 Ifacida L.	shining	■■■■■	3 jn.jl	W	W. Indies	1793.	C	s.p	
12763 dumrwsa FilAI	bushy	■■■■■	6 ...	W	Dominica	1776.	C	s.p	Gac. fr. 33.3
12764 Grtis Swz.	Greg's	■■■■■	6 jLau	W	Jamaica	1787.	C	s.p	Pl. ic. 208.1
12765 virgultosa Ais.	twiggy	■■■■■	10 my.jl	W	Jamaica	1759.	C	s.p	Pluk.al.155.3
12766 kens Sun.	Wild Clove	■■■■■	30 ...	W	Hispaniol.	1759.	C	s.p	Pl. ic. 208. 2
12767 conacea Vahl	Allspice-like	£ O or	20	my	W	W. Indies	... C	s.p	Bot. cab. 178
12768 6 8 t d X i	doubtful	ail	or	6 jn.ju	W	Maranh.	1828.	C	8.p Bot. reg. 1044
12769 obscura B. R.									
*1*9° CALYPTERA NTHERS Swz. CALYPTERANTUE3. (<i>Kalyptra</i> , veil, <i>anthos</i> , flower.), <i>Myrtacea</i> . 4.-9,									
12770 Zuzygium Sun.	Zuzygium	■■■■■ tm	20	my.jl	W	W. Indies	1778.	L	s.p Br.jam.7.2
VZniJamboland Vf.	Jambolan tree	t or	... W	W	E. Indies	1796.	L	s.p Ku. am. 1.42	
12772 Chytracilia Swz.	forked Chytraculaf	O or 20	mr.my	W	Jamaica	1778.	L	s.p Br. jam. 37.2	
412773 caryophylhdha W.	Clove-leaved	f or 20	... W	E. Indies	1822.	L	s.p Ilu. am. 1.41		
Eugenia caryophyllata liox.									

1491. *PIME'NTA* Lindl. PIMBNTA. (*Pimknto*, Indian pepper, *Span.*) *Uyrtheete*. 1.
12774 vulgaris Undl. common Allspice $\frac{1}{2}$ cul 30 my.jl W W. Indies 1723. C s.p Bot mag. 1236
Myrtus Paninta L.

1492. *OLY'NTHIA* LindL OLTNTRIA. (*Olynthos*, an unripe fig.) *Myrihcew.* 1.
12775 disticha Undl. two-rowed £ • or 2 ap.jl W Jamaica 1793. C s.p Bot mag. 867
it/yxtus disticha Swz.

1993. EUCALYPTUS Herit. EUCALYPTI:*	(EU, well, <i>kolypto</i> , to cover: calyx-lid)								
12776 rob&sta Sm.	robust Gum tree	■ Jtm 30	au.s	W	N. S. W.	1794.	L	1.p	Sm. n. h. 13
12777 reticulata Lk.	netted	■ Jor 30	-	W	N. Holl.	1823.	L	co	
12778 pilularis Sm.	pelleted	■ Jtm 30	-	W	N. & W.	1804.	L	lp	
12779 rostrata Cav.	beaked	■ Jtm 30	-	W	N. S. W.	1804.	L	Lp	Cav. ic. 4.342
12780 cornuta Lab.	horned	■ Jor 30	-	W	N. Holl.	1803.	L	s.p	Lab. vo. 1. 20
12781 glabulus Lab.	globuled	■ Jor 30	-	W	N. Holl.	1810.	L	s.l	Lab. n. h. 1.13
12782 orbicularis Lo.C.	orbicular	■ Jor 30	-	W	N. Holl.	1816.	L	8.1	
12783 incrassita Lab.	thickened	■ Jor 30	-	W	N. Holl.	1818.	L	Lp	
12784 marginata Sm.	thick-edged	■ Jtm 30	ap.jl	W	N. Holl.	1794.	L	Lp	
12785 longiloba Lk.	long-lcaved	i 1 or 30	-	W	N. Holl.	1823.	L	co	
12786 tereticornis Sm.	round-horned	■ Jtm 30	-	W	N. S. W.	1804.	L	Lp	
12787 resinifera Sm. resin-bg. red Gum tree	many-flowered	■ Jtm 30	ap.jl	W	N. S. W.	1788.	L	1.p	Bot rep. 400
12788 multiflora Sm.	small-headed	■ Jtm 30	-	W	N. Holl.	1820.	L	J	
12789 capitellata Sm.	\Viow-like	■ Jtm 30	-	W	N. Holl.	1804.	L	1.p	Sm. n. h. 42
12791 persicifolia L.	Peach-leaved	■ or SO jl	-	W	N. S. W.	1804.	L	Lp	Bot cab. 501
12792 media Lk.	intermediate	■ or SO	-	W	N. Holl.	1823.	L	co	
12793 mucronata Lk.	mucronate	■ or SO	-	W	N. Holl.	1823.	L	co	
12794 triantha Lk.	thrc-flowvrcd	■ or 30	-	W	N. Holl.	1883.	L	co	
12795 elonsitka Lk.	lengthicKd	■ Jor 30	-	W	N. Holl.	1883.	L	co	
12796 lotik-leaveil	Myrtle-lcared	■ Jor 30	Jn	W	N. Holl.	1823.	L	co	Bot. reg. 947
12797 microptiflra Lk.	small-lcaved	■ Jor 30	-	W	N. Holl.	1823.	L	co	
12798 stenoph'Aa Lk.	narrow-leaved	■ Jor 30	-	W	K. Holl.	1823.	L	co	
12799 angustifolia Lk.	narrow-leaved	■ Jor 30	-	W	N. Holl.	1810.	L	s.l	
12800 phillyreoldes Lo. C	Phillyrea-like	■ Jor 30	-	W	N. Holl.	1820.	L	8.1	
12801 viminalis Lab.	twiggy	■ Jor 30	-	W	N. Holl.	1810.	L	Lp	
12802 pulchella Lo. C.	pretty	■ Jor 30	-	W	N. Holl.	1820.	L	8.1	
12803 verruebsa Hort.	warty	■ Jor 30	-	W	N. Holl.	1820.	L	8.1	
12804 pulverulenta Lk.	hairy	■ Jor 20	-	W	N. Holl.	182a	L	co	
12805 purpurascens Dym.	Hypericuni-lyd	■ Jor 30	-	W	N. Holl.	182a	L	co	
12806 albicalyx Hort.	purpurascen	■ Jor 30	-	W	N. Holl.	1810.	L	&1	
12807 perfohata Hort.	white-stcninid	■ or 30	-	W	N. Holl.	1820.	L	s.l	
12808 heterophylla Swt	licrfuhate	■ Jor 30	-	W	N. Holl.	1820.	L	s.l	
12809 pulviflra Lk.	varius-leaved	■ Jtm 30	-	W	N. Holl.	1820.	L	s.l	
12810 cordata Lod.	powdery	■ Jtm 30	jn	W	N. Holl.	1816.	L	co	Bot. mag. 2087
12811 hirsute Lk.	small-leaved	■ Jor 20	-	W	N. Holl.	... L			
12812 microphylla Cun.	Cotinus-lcaved	■ Jor 30	-	W	N. Holl.	1824.	L	Bl	
12813 rotinotilia Lod.	powdered	■ Jor 30	-	W	N. Holl.	1818.	L	s.l	
12814 pulviflra Cun.	waved	■ Jor 30	-	W	N. Holl.	1824.	L	8.1	
12815 unduuta Hort.	winged	■ Jor 30	-	W	N. Holl.	1820.	L	8.1	
12816 alata Hort.	Botrys-like	■ Jtm 30	ap.jl	W	N. Holl.	1816.	L	8.1	Cav. ic. 4.341
12817 Atrotyodes Sm.	bloody-mouthed	■ Jtm 30	ap.jl	W	N. Holl.	1803.	L	1.p	
12818 haemastoma Sm.	rough	i 1 or 30	...	W	N. Holl.	1803.	L	Lp	
12819 scabra Nois.	Almond-leaved	■ or SO	...	W	N. Holl.	1820.	L	s.l	
12820 amygdalina Lab.	■ Jtm 30	-	W	N. S. W.	1788.	L	Lp		
12821 piperata Sm.	oblique-leaved	■ Jtm 100jlau	W	V.D. L.	1774.	L	1.p	Par. Ion. 15	
12822 corymbiflra Sm.	corymb-Jluid	■ Jtm 30	W	N.S.W.	1788.	L	Lp	Cav. ic. 4. 3	
?SS! P ^{an} culata Sm.	piniclcld	■ Jor 30	-	W	N. & W.	1804.	L	s.p	
12823 siovata Lab.	ovate	■ Jor 30	...	W	N. Holl.	1820.	L	s.l	
12824 small curved	■ Jtm 30	-	W	N. Holl.	... L				
12825 pauciflra Sieb.	few-flowered	■ Jtm 30	-	W	N. Holl.	... L			
12826 virgata Sieb.	twiggy	■ Jtm 30	-	W	N. Holl.	... L			
12827 stricta Sieb.	strict	■ Jtm 30	-	W	N. Holl.	... L			
12828 cugeiuordcs Sieb	Eugenia-like	■ Jtm 30	-	W	N. Holl.	... L			

1494. ALANGIUM.	ALANGIUM.	(<i>Alangt.</i> its Malabar name.)	<i>MyrtHceco.</i>	£.—3.	
12831 decap&alum <i>Lam.</i>	ten-petaled	H or 10 ... Pa.P	E. Indies	1779. C pl. R. mai. 4.17	
12832 hexapetalum <i>Lam.</i>	six-petaled	H or 15 ... P	E- Indies	1823. C pl. R. mal. 4.26	
1495. CALYTRIX Lab.	CALYTRIX.	(<i>Kalyx</i> , a calyx, <i>trixos</i> , triple.)	<i>Myrt&ce*</i> .	4.—&	
1248J glabra R. Br.	smooth	H or 4 ap.jn	W N. Holl.	1818. C s.l Bot cab. 586	
12833 rnoide* <i>Cun.</i>	Heath-like	H or 4 ap.jn	W N. Holl.	1824. C s.p	
12R15 floribGndia <i>Hort.</i>	bundle-flwd	H or 4 ap.jn	W N. Holl.	1820. C 8.p	
12836 pubescens* <i>Suit.</i>	pubescent	H or 4 ap.jn	W N. Holl.	1824. C s.p	
1496. FCETTDIA Com.	FOSTIDIA.	(<i>Fetidus</i> , fetid; odor of wood.)		— — — L	
128J7 xnauntiana <i>Lam.</i>	Mauritius	1 or 26 ... W	Mauritius	18*5. C <l	
1497. PITNICA W.	POMEGRANATE. (<i>Punicus</i> , Carthaginian; from N. Africa to Rome.)	<i>Granthea.</i>	8.		
12538 nana L.	dwarf	*i Jor '5 jls " R	W. Indies	1723. C r.m Bot. mag. 634	
12839 Graffatum L.	common	St fr 18 jns S	S. Europe	1548. C r.m Bot. mag. 1838	
8 album	vrhie-flowered	Sc or 10 jns W	China	— c.r.m Bot. rep. 96	
3 plenum	double-floweredH	or 10 jns S	S. Europe	— C r.m Tr.ehr.71.2	
4 flavum	yellow-/tow?rcrfSi	or 10 jn.i Y	S. Europe	— C r.m	
1498. jtfMY'GDALUS Tou.	ALMOND.	[■] to lacerate j fissured shell.)	<i>Roshcete.</i>	6..9.	
12840 nftna L.	common dwarf A	[■] rar.ap R	Russia	1683. B s.l Bot. mag. 161	
2 gefrigica Dec.	Georgian A	[■] mr.ap R	Georgia	1818. B p.l	
3 campestris Bes.	field	[■] mr.ap R	Podolia	1818. B s.l	
12841 inckna W.	hoary	[■] mr.ap R	Caucasus	— ■ s.l Pal. roll. 1	
12842 sibirica Lo C.	Siberian	A [■] mr.ap R	Siberia	1820. ■ fl	
1284J onentklis H. K.	oriental	[■] mrnp R	Levant	1156. ■ fl	
12844 commanis L.	common, or sweet	[■] mr.ap R	Barbary	1548. ■ nl	
2 amhra	bitter	[■] mr.ap R	Barbary	1548. ■ h.l Black. 195	
3 fragilis Dec.	brittle	[■] mr.ap 11	Barbary	... ■ h.l Noi.jar.7. 3.2	
4 macrocarpa Dec.	long-fruited	[■] mr.ap R	Barbary	■ hi Noi.Jar.7.3	
5 persicoldcs Dec.	Peach-like	[■] mr.ap R	Barbary	— ■ h.l Noi.jar.7.3.1	
12845 p&mila W.	double dwarf	[■] my.jn R	China	1683. ■ fl Dot mag. VYI6	
Prunus sinensis Per.*.					
1499. PE'RSICA Tou.	PEACH.	(Originally from Persia.)	<i>Roshceat.</i>	8.	
12846 vulgkris Mil.	common	3* fr 15 ap.my R	Persia	1568. B co Noi.jar.ic.	
AvdS gdalus Persica L.					
1 cfirne seccdente	parting-fleshed *£	fr 15 ap.my R	Persia	1562. B co	
8 c&rme adhaerente	adhering-fleshed \$	fr 15 ap.my R	Persia	1562. B co	
3 fl̄re pleno	double-flwd \$	or 15 ap.my ...	Persia	... B co	
12847 lsVis D^c.	smooth, or <i>Nectarine</i> 5	fr 15 ap.my R	Persia	1562. B co Noi.jar. 80. S3.	
^m^gdalus Pdrsicla Nectarina //, A'					
1 came secedente	parting,fleshed Y	fr 15 apmy R	Persia	1562. B co	
8 came adhsrente	adhering-fleshed J	fr 15 ap.my R	Persia	1568. B co	
1500. ^RMENPACA 3w.	APRICOT.	(Originally from Armenia.)	<i>Rotdeaf.</i>	4.	
12848 vulgans Lam.	common	% Prunus ylmenlaca L.	fr 15 f.mr W	Levant	1548. B co Lam.il. 431 n.8
1 ovalifolia Scr.	oval-leaved	Y	fr 15 f.mr W	Levant	154R B co Noi.jar.1.1.8.8.
2 cordifolia Ser.	heart-eaved	\$	fr 15 f.mr W	Levant	1548. B co Noi.jar. 1.3.8.3
12849 dasycarpa Pen.	thick-fruited	\$	fr 15 ap W	■ ■ 1800. B co Bot. cab. 1850	
Prunus dasycarpa Ehrh.					
2persicifoliaX.u.	Peach-leaved	*£	fr 15 ap W	— 1800. B co L.duh.fr.5.552.1	
18850 Biberica ft.	Siberian	&	fr 6 ap Pk	Siberia 1788. L r.m PaLros.1.8	
Pr&nus Riberica L.					
18851 brigantica Pers.	Brigantian	SI	fr 4 reir Pk	S. Europe 1819. L co L.duh. fr. 5. SB	
Prunus brigantica Vit					
1501. PIU^NUS Tou.	PLUM.	(<i>Prune</i> , its Greek name.)	<i>Rostiecaf.</i>	8.—10.	
1x852 spinii L.	spiny Sloe tree	B	or 15 mr.ap W	llritrin hod. S co ling. bot. 842	
i microcarpa IVaUr.	small.fruited Sk		or JO mr.ap W	Britain hed. S co	
3 macrocarpa JYallr.	long-fruited fib		or lu mr.ap W	Britain hed. S CO	
4 ovsita Ser.	ovate-fruited A		or 10 mr.ap W	Britain hod. S co Black. 494	
12853 ineititia L.	graftcl	Sk	fr SO ap W	— S CO Eng.bot.841	
12854 candicans Balb.	whitish	Sk	or 15 ap W	— 1820. S CO	
18855 Cocomilla Ten.	Cocomilla tt		or 20 ap W	Calabria 1894. L 8.1	
12856 maritima JKon.	sea	tik	or 4 my W	N. Amer. 1800. L co	
18K57 pubescensPoir.notPA.pultescent	fib		or 8 my W	fr 20 ap W England 1848. G r.m Eng. bot. 1783.	
18858 domestua L.	domestic Plum	■	fr SO ap W	— G co Duh.ff. 8.13	
8 Claudina Pen.	Petite Heine Claude^		fr SO ap W	— G co Duh.fr. 2.11	
SmyrobalaL.	Myrobalan		fr 20 ap W	— G co Dull. fr. 2.15	
4 dTamasca L.	Dumas Musque^		fr SO ap	— G co Duh.fr. 2. 20.3	
5 luroneinfisrr.	TufmPrunier Suisse*		fr SO ap Turin	— G co Duh. fr. 2 20.1	
6 Juliania L.	Siflw Julian's		fr 80 ap W	— G co Duh. fr. 2 20.4	
I V ^ n f S -	*Catherine's		fr SO ap W	— G co Duh.fr. 8.89.5'	
8 Aubertina Sor.	D/we Aubert's		fr SO ap W	— G co Duh. fr. 2.8.10,	
2 Seta Imperatrice viol.	pyramidalis JDic.	■	fr 20 ap W	— G co Duh.fr. 2.18	
18859 divaricata Led.	divaricate	■	or 10 ap W	Caucasus 1820. L co	
ASUS J. CHERRY.	(First brought from Cerasys, a town of Pontus in Asia.)	<i>Rosdce/e.</i> SQ.—42.			
18860 avium ifoen.	birds* Corone	+	or 50 ap.my W	England ... S co Black. 425	
1 sylvestris Ser.	wild	i	fr 50 ap W	Britain woods. S co	
8 macracarpa 5w.	to w purple Jd		fr 50 ap W	SwitzerL ... S co L.duh.fr. 1.4. D	
3 pallida Ser.	white and red	*	or 15 ap.my W	— B co L.duh. fr. 5.4.B	
.4 multiplex Ser.	death-flowered	*	fr 20 ap.my W	S. Europe — B ee L.duh.fr.5.18.D	
12861 dur&cina L.	hard		fr SO ap W	— B co L.duh. fr. 5.2.B	
1 cordigera Ser.	heart-bearing		fr SO ap W	— B co L.duh.fr.5.18.B	
2 oblonga Ser.	blunted		fr SO ap W	— B co L.riuh.fr.516.A	
3 mammillaris Ser.	mammillary	\$	fr 20 ap.my W	— B co L.riuh.fr.516.A	
xnacrophjrla Porr.					

12662 <i>Juliana</i> Dec.	<i>Saint Julian's</i>	20	ap.my	W	S. Europe	...	Q co	L. duh. fr. 5.19	
<i>U Heaumiana</i> Ser.	<i>Heaum's</i>	15	ap.my	W	—	—	B co		
3 pendula <i>Ser.</i>	<i>pendulous</i>	10	ap.my	W	S. Europe	...	G co		
12663 <i>Pseudo-cerasus</i> Lftuff.	<i>Bastard Cher.</i>	6	ap.my	W	China	1821.	G co	Bot. reg. 800	
12864 <i>capromiana</i> Dec.	<i>Capronian</i>	20	ap.my	W	S. Europe	...	•G co	Noi. jar. 5	
1 Montmorecydn <i>a</i> Ser.	<i>Montmorenc</i>	20	ap.my	W	—	—	G co	Duh. fr. 2.11	
2 pallescent <i>Ser.</i>	<i>pallicent ceranMe</i>	20	ap.my	W	—	—	G co	Duh. fr. 1.8.	
<i>Sgobbita</i> Ser.	<i>Gobbetia, white-fleshf</i>	20	ap.my	W	—	—	G co	Duh. ft. 4.3	
4 polygyna <i>Ser.</i>	<i>polyg. Cer. a bouquet</i> *	20	ap.my	W	—	—	G co	Duh. ft. 4.5	
5 multiplex <i>Ser.</i>	<i>doub'e-flowered%</i>	12	ap.my	W	—	—	G co		
6jversicifbra <i>Ser.</i>	<i>Peach-Howered</i>	20	apuny	W	—	—	G co		
7 vanegdita <i>Ser.</i>	<i>variegated</i>	10	ap.my	W	—	—	G co		
8 griditta <i>Ser.</i>	<i>Griotte</i>	20	ap.my	W	—	—	G co		
9 cordigera <i>Guigne</i>	<i>heart-bearing</i> J	20	ap.my	W	—	—	G co	L.duh.fr.5.I4.A	
2 sessilifl&ra <i>Ser.</i>	<i>sessile-flowered</i>	fr 20	ap.my	W	—	—	G co	Den. br. 131	
12866 <i>Chamajcerasus</i> Loi.	<i>Ground Cherry</i>	or 8	my	W	Austria	1587.	L s.l	Jac. ic. 1.90	
12867 <i>ncrsiciflia</i> Loi,	<i>Peach-leaved</i>	or 8	my	W	N. Amer.	—	G s.l	Mil. ic. 89.2	
12868 <i>pitnila</i> Mx.	<i>dwarf</i>	or 2	my	W	N. Amer.	17*56.	L s.l	Bot mag. 1117	
12869 <i>pygme/a</i> Loi.	e	or 4	my	W	N. Amer.	1823.	B co		
12870 <i>nigra</i> Loi.	<i>northern Choke</i>	or 20	ap.my	W	N. Amer.	1773.	G co		
12871 <i>borekhs</i> Mx.	<i>depress* Ph.</i>	or 4	my	W	N. Amer.	1822.	L co		
12872 <i>hyemalis</i> Mx.	<i>depressed Sand</i>	or 4	my	W	N. Amer.	1805.	L s.l		
12873 <i>prostrata</i> <i>Ser.</i>	<i>Winter Black Choke</i>	or 4	my	W	N. Amer.	1805.	L s.l	Bot reg. 136	
12874 <i>chicasa</i> <i>Ser.</i>	prostrate J*	J or 1	ap.my	Pk	Crete	1802.	L s.l		
12875 <i>pubescens</i> <i>Ser.</i>	<i>Chicasaw Plumtk</i>	or 6	ap.my	W	N. Amer.	1806.	L 8.1		
Prtnus pubescens PA	<i>pubescent</i> dk	or 1?	ap.my	W	N. Amer.	1820.	G co		
12877 <i>japonica</i> Loi.	J	or 2	mr.my	Pk	Japan	1810.	G co		
2 <i>multiplex</i> <i>Ser.</i>	<i>double</i>	ill	or 4	mr.my	Pk	Japan	1810.	G co	
12878 <i>pennsylvanica</i> Loi.	<i>Pennsylvanian</i>	or SO	my	W	N. Amer.	1773.	S co	W.ar.aS	
12879 <i>Mahaleb</i> Mil.	<i>Mahaleb</i>	or 20	ap.my	W	Austria	1714.	G co	Jac. au. 3. 227	
12880 <i>Padus1)»\`</i>	<i>Bird Cherry</i>	or 30	ap.my	W	Britain woods.	S co	Eng. bot. 1383		
1 <i>vulgaris</i> <i>Ser.</i>	<i>common</i>	or 30	ap.my	W	Britain woods.	L co	I. duh. fr. 5.1		
2 <i>parviflora</i> S<r.	<i>small-flowered</i>	or 30	ap.my	W	N. Europe	...	Fl. dan. 205		
3 <i>rubra</i> <i>Ser.</i>	<i>red Cornish Bird</i>	or 30	ap.my	W	Britain	...	W.ar.4.2		
Pr&nus rilbra W.									
4 bracteba <i>Ser.</i>	/on^-bracted	T	or SO	ap.my	W	Europe	L co		
12881 <i>virginiana</i> Mx.	<i>Virginian</i>	i	or 30	my.jn	W	Virginia	1724.	L co	W. nr. 5.9
12882 <i>canadensis</i> Loi.	<i>Canadian</i>	or 15	my.jn	W	Canada	1820.	L co	Pluk. aL 158.4	
12883 <i>nepalensis</i> <i>Ser.</i>	<i>Nepal</i>	-J or 2	...:	W	Nepal	1820.	G co		
12884 <i>serdfina</i> Loi.	late American Bird \$	or 30	my.jn	W	N. Amer.	1629.	L co	Den.br.48	
12885 <i>occidentalis</i> Zrt	Went /mtom K	D or 20	year	W	Jamaica	1784.	L co		
12886 <i>lusitanica</i> Loi.	Portugal Laurel \$	or 20	Jn	W	Portugal	1648.	S co	Mil. ic. 196.1	
12887 <i>Laurocerasus</i> Loi. com.	Laurel Ch. A	or 12	ap.my	W	Levant	1629.	C co	L. duh. ft. 5.1	
12888 <i>sphrocarpa</i> Loi.	round-fruited *	or 10	jn.jl	W	Jamaica	1820.	C ap	SI jam. 2.193,1	
12889 <i>carolininida</i> Mx. Carol. Evergr. Bird ±		30	my	W	Carolina	1759.	L co		
• 1503. CHRYSOBALANUS L. COCOA PLUM. {Chrysos, gold, balanas, acorn; fruit) Chrysobalhm*. 2.-4.									
12890 <i>Icaco</i> L.	Iceao West Indian* CD ft	15	...	W	W. Indies	1752.	L r.m	Jac.am. 94	
12891 <i>oblongifolius</i> Mx.	oblong-leaved *i_Jfr	3	my.jn	W	Georgia	1812.	C l.p	Bart. it. a ic.	

1504. LOPHTRA Banks LOPIHRA. {Lophos, a crest; bract at flower.) Lophirete. 1.
J2892 afitancafanfts African Scrubby Oak± mor 10 jn.jl W & Leone 1822. S s.l Gac.fr. 3.188

CLASS XII. ORDER 2.

JCOSANDRIA DI-PENTAGY'NIA.

1505. JIF'E'SPILUS £.	MEDLAR. {Mesos, half, piios, bullet; fruit resembling half a bullet) Rosace*. **								
12893 <i>germifinica</i> L.	Jeon — \$	fr ²	13my.jl	WW	England lied.	G ill	Eng. bot. 1523		
S stricta Dec.		fr	12	myjl	W	Europe	...	G co	
3 diffits Dec.		ft	12	myjl	W	Europe	...	Duh.ft. 1.3	
12894 Smiths Dec.		or	12	myjn	W	.	.	Ex. bot. 1.18	
grandiflora Sm.						1800.	L co		
• 1506. CRATIE^GUS I. HAWTHORN. (Kratost strength : hardness of wood.) Roster*. 49.-52.									
12895 <i>Pyracantha</i> Pert.	Evergr. Thorn	or 10	my	W	S. Europe	1629.	S s.l	Schm. ar. 90	
12896 <i>crenulata</i> Box.	crenulated	or	JO	my.jn	W	Nepal	1830.	B si	
12897 <i>Crtis-galli</i> L.	Cockspur	or	20	myjn	W	N. Amer.	1691.	BC Den. br 56	
1 splindcns ff. K.	shining	or	20	my.jn	W	N. Amer.	...	B co Pluk. al. 46.1	
S pyracanthifbliai)^.	Pyracant-lvd	or	20	my.jn	W	N. Amer.	...	B po	
3 #alificflia	'Willow-leaved	or	20	myjn	W	N. Amer.	...	B co	
4 linear is Dec.	linear	or	20	myjn	W	N. Amer.	...	B co	
Jlfspilus linekris Desf.									
12898 <i>punctata</i> H. X.	dotted-/r»^d	or	15	my	W	N. Amer.	1746.	B co	
12899 <i>elliptica</i> H. K.	elliptic-teawd	or	20	my	W	N. Amer.	1765.	B co	
12900 <i>irunifolia</i> Bosc	Plum-leaved	or	20	myjn	W	N. Amer.	1818.	B co	
12901 <i>pruneUXB Wa.</i> Bosc	Prunella-lvd	or	20	myjn	W	.	.	B co	
12902 <i>latifolia</i> Pers.	broad-leaved	or	20	my.jn	W	N. Amer.	1820.	B co	
12903 <i>parviflora</i> H. K.	smalMeaved	or	15	myjn	W	N. Amer.	1704.	B co	
12904 <i>ovalifolia</i> Horn.	oval-leaved	or	20	myjn	W	N. Amer.	1810.	B co	
12905 <i>apiifolia</i> Mx.	Parsley-leaved	or	15	myjn	W	N. Amer.	1812.	B co	
12906 <i>pyrifltha</i> H. K.	Pear-leaved	or	15	jn	W	N. Amer.	1765.	B co	
Jedulidr. fo r^* W.									
ISS [«] P <th>thulka</th> Alar.	thulka	glandulous	or	20	myjn	W	N. Amer.	175a	B co Botcab. ICIS
12895 P <th>thulka</th> Alar.	thulka	npatufoleaved	or	15	myjn	W	N. Amer.	1806.	B co Bot. cab. 126*
J8910cordataA/tf.	soarlet-/ml/rfr	or	20	ap.my	W	N. Amer.	16831	B co Den. br. 62	
J8910cordataA/tf.	hexauiiaved	or	SO	my	W	N. Amer.	1738	B co Den. nr. 69	

12911 lobata <i>Bosc</i>	lobed	or 15	my.jn	W	—	—	B co		
12912 purpurea <i>Bosc</i>	purple	or 15	my.jn	W	—	1822.	B co	Den. br. 60	
12913 florentina <i>Zuc.</i>	Florentine	or 15	my.jn	W	—	1800.	B co		
P ⁺ rus crategifolia	Lab.								
12914 rfigra IV. & K.	black	or 30	ap.my	W	Hungary	1819.	B co	Den. br. 64	
12915 pentagyna W. & K.	five-styled	or 15	my.jn	W	Hungary	1820.	B co		
12916 flava H. K.	yellow Pear-berried	or 20	my.jn	W	N. Amer.	1724.	B co	Den. br. 59	
12917 fissa <i>Bosc</i>	cleft	or 15	my.jn	W	—	1810.	B co		
12918 maroccana <i>Pers.</i>	Morocco	or 15	my.jn	W	Barbary	1822.	B co		
12919 Oxyacantha <i>L.</i>	com. Hawthorn	or 15	my.jn	W	Britain	hed.	S co	Eng. bot c. ic.	
2 rusea	roseate/ZuMf	or 15	my.jn	R	—	—	B co		
laciinata <i>Wattr.</i>									
3 major	greater-fruited	or 15	my.jn	W	—	—	B co		
4 pr&NtX	early	or 15	my.jn	W	—	—	B co		
5 plena	&ou)e-flowered	or 15	my.jn	W	—	—	B co		
6 aurea	gohlen-berried	or 15	my.jn	W	—	—	B co		
12920 oxyacanthoides TTtoJ.	Oxyacantha-lk	V	or 15	my.jn	W	France	1822.	B co	
12921 enocirpa Z/wt.	woolly-fruited	or 15	my.jn	W	Britain	woods	B co		
12922 monogyna <i>Pall.</i>	one-styled	or 15	my.jn	W	Siberia	—	B co	Pal. ros. 1.12	
12923 heterophylla <i>Flug.</i>	various-leaved	or 20	my.jn	W	N. Amer.	1816.	B co	An. mu. 12.38	
12924 Azarblus L.	Azarole	or 15	my.jn	W	S. Europe	1640.	B co	Bot. rep. 579	
J2925 Aronia <i>Bosc</i>	Aronia	or 15	my.jn	W	S. Europe	1810.	B co	Foe crat. 85	
12926 orientalis <i>Bosc</i>	oriental Axarole	\$	or 15	my.jn	W	S. Europe	1810.	B co	
12927 tanacetifolia Pers.	Tansy-Ivd Axarole	V	or 15	my.jn	W	Greece	1789.	B co	
2 taurica	Taurian	**	or 15	my.jn	W	Tauria	1800.	B co	
Celsiana Hort									
12928 odoratissima A. R.	sweetesUsctd A.%	or 15	my.jn	W	Crimea	—	B co	Bot. rep. 590	
12929 aciniMa. Uc.	fringed	or 15	my.jn	W	Sicily	1816.	B co		
12930 melanocarpw Bieb.	black-berried	or 15	my.jn	W	Tauria	1820.	B co		
12931 flabellata <i>Bosc</i>	fan-leaved	or 15	my.jn	W	S. Europe	—	B co	Bot cab. 1100	
12932 spinosissima <i>Lod.</i>	spiniest	or 15	my.jn	W	Europe	—	B co		
12933 mexickna 3/8c S.	Mexican	Jor 15	my.jn	W	Mexico	1823.	B co		
12934 viridis L.	green	or 15	my.jn	W	Carolina	1810.	B co		
12935 OlivienVina <i>Bosc</i>	Olivier*	or 20	my.jn	W	Asia Min.	1820.	B co		
12936 Poiret'sma Dec.	Poiret's	or 20	my.jn	W	N. Aincr.	1810.	B co		
12937 glabra <i>Lod.</i>	smooth	or 15	my.jn	W	Italy	—	B co	Bot cab. 248	
12938 alpina <i>Mil.</i>	alpine	or 20	my.jn	W	N. Amer.	—	B co		
12939 luciria <i>Mil.</i>	shining	or 20	my.jn	W	Siberia	1810.	B co		
12940 sangutnea <i>Pall.</i>	bloody	or 15	my.jn	W	Carp-it, mo....	B CQ			
12941 carpatica Lo. C.	Carpathian	or 20	my.jn	W	France	—	B co		
12942 corallina Thun.	coralline	or 15	my.jn	W	N. Amer.	—	B co		
12943 florida Lo. C.	Florida	or 20	my.jn	W					
12944 communis L.	PVRUS.	(Peren,	Celtic.)		Rosaceae.	40.—46.			
1 A ⁺ chras WaUr.	common Pear	V	or 20	ap	England woods.	G co	Eng. bot 1784		
1 A ⁺ chras WaUr.	Acras entire-lvd	5	fr 20	ap	G	co			
2 Pyraster IVallr.	Fyns.serrated-lod [^]	5	fr 20	ap	G co	Gae.fr. 87.2		
sylvestris Gac.									
3 satWa Dec.	garden	\$	fr 20	ap	W	—	G co		
12945 Polveraria L.	Polver's hooded-lvd	?	or 15	apjn	W	Germany	1786.	G co	
lorv. ^{Bo} »wyllenVnfl Dec.								Bot cab. 1009	
12946 *alviflaria Dec.	Sage-lvd Aurelian	\$	or 15	myjn	W	France	1806.	G co	
«947 amygdaliformis Tj/.	almond-formed	\$	or 15	myjn	W	S. Europe	1810.	G co	
lofuuo *allc «»ia Zoi.									
^s@48 daeagnibflia Pall.	Elaeagnus-lvd	\$	or 20	myjn	W	Siberia	1806.	G co	
orientalis Horn.									
WUlow-leaved	V	or 20	my.jn	W	Russia	1780.	G co	Bot.reg.514	
ta D.Don	woolly	or 15	my.jn	W	Nepal	1818.	G co		
CD enata D. Don	crenated	J	or 15	myjn	W	Nepal	1820.	G co	
W «da Muuhi* Mt. Sinai Medlar	snovr-white-lvd	£	ft 20	myjn	W	Levant	1820.	G co	Den. br. 49
953 niv&lia Z.	sour apple	%	or 6	my	W	Austria	...	G p.l	Jac. au. 2. 107
12944 acerba Hfc.			fr 20	ap.my	W	Europe	...	G co	Fl. dan. 1101
3/afusij3 sylvestris 27. rfnn.									
12955 italpus L.	Apple tree	fr 20	ap.my	W	Britain	woods.	G r.m	Eng. bot 179	
12956 astracanica Dec.	Astracan apple	fr £0	ap.my	W	Astracan	1810.	G co		
12957 floribunda B. R.	bundle-flwd	or 8	ap.my	W	China	1818.	G co	Bot reg. 1006	
12958 diolca W.	dioecious	cu 10	ap.my	W	China	1818.	G co		
12959 spectabilis U.K. showy Chinese apple	Ch.	ft 20	my	Pk	China	1780.	G co	Bot mag. 267	
12960 urunifolia W. Plum-lvd	Siberian crab	fr 20	ap.my	Pk	Siberia	1758.	G co	Mil. ic. £ 269	
12961 baccta L.	berried	or 15	ap.my	Pk	Siberia	1784.	G co	Den.br. 51	
12962 tomentbsa Dum.	tomentose	or 20	myjn	W	Eurojie	1810.	G co		
12963 coronifolia L. crown sweet-setd crab	blunt-leaved	or 20	my	Pk	Virginia	1724.	G co	Bot. mag. 9009	
annulifolia // K.	narrow-leaved	or 20	my	I'k	N. Amer.	1750.	G co	Den. br. 132.	
12965 ria Ehrh.	White Beam treef	tm 40	myjn	W	Britain	moi.w.	G co	Eng. bot. 1858	
1 obtusifolia Dec.	blunt-leaved	tm 40	my jn	W	Europe	...	G co	Fl. dan. 302	
2 acutifolia Dec.	sharp-leaved	tm 40	myjn	W	Europe	...	G co	Duh. no. 34	
aliiina W.en.									
12966 intermedia Ehrh.	intermediate	tm 40	ap.my	W	Sweden	1789.	G co	Fl. dan. 401	
1 latifolia Dec.	broad-leaved	tm 40	ap.my	W	Denmark	1789.	G co	Duh.ar.1. 80	
2 angusUCbia Dec.	narrow-leaved	fr 10	ap.my	W	—	—	G co		
edulis W.en.									
12967 terminalis Ehrh.	gripping Wild Serviced	tm 50	ap.my	W	England	woods.	S co	Eng. bot. 298	
Crats ^{gus} terminalis L.									
12968 tnlobata Dec.	three-lobed	or 20	myjn	W	S. Europe	1810.	G co	Lab. syr. 4.10	
12969 upsaliflisa Lo. C.	Upsal	or 20	myjn	W	Sweden	1824.	G co		
12970 auriculata Dec.	auriculated	or 20	my.jn	W	Egypt	1800.	G co		
12971 pinnatifida E.B.	pinnatifld	tm 40	myjn	W	England	rocks.	S co	Eng. bot. 2331	
Sorbus bybrida L.	small-fruited	or 10	myjn	W	N. Amer.	...	L co		
*2972 microcarpa Dec.									
iSibus microcarpa Ph.									
J12973 aucupariaGh??. fowler's MountainAsh		or 30	myjn	W	Britain	moi.w.	S co	Eng. bot. 337	
iSS amerrickna Dec.	American	or 15	myjn	W	Canada	1782.	G co	Den.br. 54	
*2975 Sorbus Gae.	True Service	fr 30	myjn	W	England	iroi.w.	S co	Eng. bot. 350	

13034 ralicifolia L.	Willow-leaved	■ or 5 Jn.au	Pk	Britain	mgLw. Skco	Eng. not 1468	
2 alpestris Patt,	alpine	■ or 5 Jn.au	W	Russia	1820. Sk co	Pal. ros. 1.22	
3 alba fArA.		■ or 5 Jn.au	W	N. Amer.	... Sk co	Mil. ic. 257.2	
paniculata W.							
4 latifolia JK	broad-leaved	■ or 5 Jn.au	W	Europe	--	Skco	
teA obovata RaJL not IK. tf #.							
J1030 carpifolia W. en.	Hornbeam-lvd	■ or 4 Jn.au	W	N. Amer.	... L co	Den. br. 66	
i*6 tomentosa L.	tomentose	■ or 5 au.s	Pk	N. Amer.	1736. Skp.l	Schm. ar. 1.51	
13037 sorbifolia L.	Sorbus-leaved	■ or 4 au	W	Siberia	1759. Skco	Schm. ar. 1.58	
2 alpina		■ or 3 au	W	Siberia	1817. Skco	Pal. ros. 1.25	
13038 Aruncus L.	Goat's-beard	A or 4 jnjl	W	Siberia	1633. D p.l	Pal. ros. 1.26	
2 americinus Mr.	American	A or 4 jnjl	W	N. Amer.	... D p.l		
13039 Filipendula L.	Dropwort	A or 5 Jn.au	W	Britain	m. pas. D co	Eng. bot 284	
5 plena	dewy	A or 5 Jn.au	W	Britain	gard. D co		
3 nftnor Campb.	smaller	A or 5 Jn.o	W	Europe	--	D co	
4 pubescens Campb.	oubescnt	A or 5 Jn.o	W	France	... D co		
13040 Ulm^ria L.	Meadow-sweet	A or 1 Jn.o	W	Britain	m. mea. D co	Eng. bot 960	
2 multiplex Dec.	double-flowered	A or 2 Jn.o	W	Britain	m.raea. D p.l		
3 tomentosa Campb.	woolly	A or 2 Jn.s	W	Britain	m.me. D co		
4 denudata Campb.	stripped	A or 2 Jn.s	W	Britain	m.me. D co		
5 variegata	variegated	A or 2 Jn.s	W	Britain	mmea. D co		
13041 lobata Mur.	lobed	A or 2 jlau	W	Siberia	1765. D p.l	Jac. vin. 1.88	
?3S2 Palmita Thun.	palmate	A or 2 jlau	R	China	1823. D p.l		
1JO43 digitata W.	digitate	A or 2 jlau	R	Siberia	1823. D p.l	PaL ros. 1.27	
			11				
,1516 GILL&NIA Moen.	GILLBNIA.					(Probably the name of some obscure botanist.) Rosacee, 2.	
13044 trifoliata Moen.	three-leaved	A or 2 jnau	R.w	N. Amer.	1713. D p.l	Bot. mag. 489	
Spiraea trifoliate L.							
13045 stipulacea Mi*.	Air/p-stipuled	A or 2 jn.au	R.w	N. Amer.	1805. D co		
Spiraea stipulacea W. «t.							
,1517. SESITVUM I*	SESUVIIM.		(Not explained.)		Ficoide*. 5.-7.		
1P046 pedunculatum !«.	peduncled	£ (23 or 2 jn.jl	R.W	W. Indies	1692. C r.m	Lam. il. 434.	
ij047 sessile Pers.	sessile-flowered	jg [23 or 3 jn.jl	R.W	W. Indies	... C r.m	Dec. pi. 9	
Portularastrum Dec.							
13048 revolutiflum Vahl	revolute-leaved	E3 pr 1J jl.au	R.W	S. Amer.	... C Lp	Bot mae. 1701	
13049 loDRifolum W.	long-leaved	O pr 1J jl.au	R.W	S. Amer.	1816. S l.p		
13050 rufepens W.	creeping	O pr 1 jl.au	R.W	E. Indies	1816. S Lp	Ru.am.5.72.1	
,1518. AIZOPOK L.	AIZOON.	{Aei, always, zoon, alive	O	Ficoide/e. 5.—16.			
13051 canariense L.	Canary	iQJ or 1 jl.au	Y	Canaries	1731. S r.m	Bot rep. 201	
13052 Glinus-like		1 jn.au	Y	C G. H.	1774. C r.m		
?3 perfoliatum L.	perfoliate	1 jn.au	Pk	C. G. H.	1818. C r.m		
?3 hispanicum L.	Spanish	tJUu	Ap	Spain	172a S r.m	Dec. pl 30	
WU55 lanceolatum Mur.	spear-leaved	f au	Pk	O G. H.	1752. S r.m		
,1519. TETRAGONIA i. TETRAGONIA.		{n, four.Amalf, an angle; fruit.)		Ficoidcir. 10.—16.			
13056 expansa//ff. expanded	N. Z. Spinach	O cu 1 6 ams	G	N. Zeal	1772. C s.l	Bot. mag. 2362	
13057 crystallina Ker	crystalline	iQJ un 2 jn	G	Peru	1788. ■ 8.1	Dec. pi. 34	
J1Q8 fruticosa L.	shrubby	tt. n. iJ un	2 jl.s	G	C. G. H.	1712. ■ 8.1	
13t19 dec&mbens Mil.	trailing	tL J un	1 jls	G	C. G. H.	1758. ■ 6.1	
J3060 Tetrapteris Haw.	trmngd-seededn.	✓ un	2 jl.s	G	C. G. H.	1795. ■ 8.1	
J14 spicata L.	spiked	a. □ un	1 jl	G	C. G. H.	179T. ■ 8.1	
J1 herbacea L.	herbaceous	Tf LAJ un	* jnjk	s	C. G. H.	1752. ■ 6.1	
J2033 chinkte // K.	hedgehog	f Olun	my.au	G	C. G. H.	1774. C 8.1	
J3064 line&rro JShni.	linear	B. Ljun	my.au	G	C. G. H.	1819. C 8.1	
J3065 obovata Haw.	obovate	*t uJ un	H s .	G	C. G. H.	1821. C 8.1	
*1520. MESEMBHY A'NTHEMUM L. Fia MARIGOLD. (<i>Mcgembria</i> , mid-day, anthenon, fl.) FicoUL 313.—350.							
I. ACAI/UA.—Stems none or very short, leaves large.							
13066 nsinutum Haw.	minute	£ IAJ cu	A s.n	Pk	C. G. H.	1795. ■ s.l	
^ 67 minimum Haw.	smallest	tf IACU	ft. s.d	Pa. Y C. G. H.	1776. ■	Bot. mag. 1376	
13068 nerpusilim Haw.	>ery small	j£ IAJCU	s.d	Pa. Y C. G. H.	1819. ■		
130G obconellum Haw.	small obcordate	jg LAJCU	i fo	W C. G. H.	1776. ■ s.l	Bot. mag. 1647	
J1P070 obconellum Haw.	small conical	j£ LAJ cu	ifo	W C. G. H.	1786. ■ 8.1		
13071 ficiflora Salm	fig-like	tf IAJ cu	fo	C. G. H.	1819. ■ 8.1		
J3072 truncatellum Haw.	small truncated	df IAJ cu	fo	Pa. Y C. G. H.	1795. ■ 8.1		
13073 fibuliforme Haw.	button-shaped	f IAI cu	fo	... C. G. H.	1795. ■ 8.1		
Sji UV88 Krone HaTM>	grape-like	< iAI cu	-	... C. G. H.	1820. ■ 8.1		
?3 gaufranne Haw.	nut-shaped	tf IAI cu	-	C. G. H.	1790. ■ si		
S3 testiculare Aft.	testicular	■ Si cu		W C. G. H.	1774. ■ BI	Bot. mag. 1573	
1J07 ph^llum Haw.	eight-leafed	f tAJcu	in	Y	C. G. H.	1819. C s.l	
s longisulcum	rather longer,/^	iAJ cu	in	Y	C. G. H.	1774. C s.l	
3 roseum	rosy	tf IAIcu	in	R	C. G. H.	1792. C s.l	
	obtuse/own	£ LAJCU	mr.ap	Pk	C. G. H.	1776. C s.l	
					CG. H.	1775. C s.l	
					CG. H.	1822. C 8.1	
	large-dotted				Y	C. G. H. 1K22. C 8.1	
					Y	C. G. H. 1822. C 8.1	
					Y	C. G. H. 1822. C 8.1	
					Y	C. G. H. 1822. C 8.1	
13083 canum Haw.	hoary	1AJcu	A -	Y	C. G. H.	1795. C 8.1	
13084 alo)dea Haw.	Aloe-like	LAJCU	i ...	Y	C. G. H.	1819. C 8.1	
13085 albintum Haw.	white-marked	IAJor	i s	Y	C. G. H.	1825. C 8.1	
13086 albipunctatum/faut.	white-dotted	IAJ or	is	Y	C. G. H.	1823. C s.l	
13087 cairinum Haw.	dog-chap	LAlg	i au.o	Y	C. G. H.	1717. C s.l	Dec. pi. 95
J1088 lupinum Haw.	wolf-cA <p></p>	IAlg	i ...	Y	C. G. H.	— C 8.1	
*30t0 vulpinum Haw.	fox-chap	LAlg	i au.o	Y	C. G. H.	1795. ■ 8.1	
J190 h^bridum Haw.	bastard			Y	C. G. H.	— C 8.1	
PSJ Wum L.	whitish	IAJgr	i*JL <i>ci</i>	Y	C. G. H.	1714. ■ 8.1	
3Q2 agn'mum Haw.	Vamb-chap	IAJor	imy.jn	Y	C. G. H.	1824. ■ 8.1	Bot. mag. 1824
2 minus Haw.,	less	IAJor	i myJn	Y	C. G. H.	1824. ■ 8.1	
3 cretisculum/Zhtv.	more erect-^rfjg	■ H	i myjn	Y	C. G. H.	1824. ■ 8.1	

13093 tigrinum Haw.	tiger-chap	£ IAJ S ^T	* in	Y	C. G. H.	1790.	C si	Bot reg. 260
13094 felinum Haw.	cat-chap	£ IAf ^r	i au.n	Y	CG. H.	1730.	C 8.1	Dec. pi. 152
13095 mustellinum Haw.	weasel-dtop	£ iAJ K ^r	I	... Y	C. G. H.	1820.	C 8.1	
13096 erminium Haw.	criminecttap	j£ iAJ or	*my.jn	Y	C. G. H.	1824.	C 8.1	
13097 murinum Haw.	mouse-chap	j£ IAJgr	t.	Y	C. G. H.	1790.	C 8.1	
13098 muscullnum Haw.	little mouse-*	£ IAI gr	ijl	Y	CG. H.	1824	C 8.1	
13099 dolabrilörme Haw.	hatchet-fonned	tt. J gr	i my.n	Y	C. G. H.	1745.	C 8.1	
13100 acapigrum Haw.	scape-bearing	j£ iA] cu	1 au.8	Y	C. G. H.	1823.	C 8.1	
13101 carWans Haw.	keeling	IAICU	- - -		C. G. H.	1818.	C 8.1	
13102 denticulatum Haw.	denticulate	■■■/or	*ap	Y	C. G. H.	1793.	C 8.1	
2 gildicum	glaucous	■■■/or	i ap	Y	C. G. H.	—	C 8.1	
3 cundidiasmus	whitest	■■■/or		Y	C. G. H.	—	C 8.1	
13103 rnb fistinum Haw.	robust	■■■/or		Y	C. G. H.	1795.	C 8.1	
13104 compactum // X.	compact	<t u.J or		Y	C. G. M.	1780.	C 8.1	
13105 quadrifidum Haw.	quadnfid	tt. i_J or		Y	C. G. H.	1795.	C 8.1	
13106 bifidum Haw.	bifid	■■■/or		Y	C. G. H.	1803.	C 6.1	
13107 bibrateatum //aw.	double-bracted	tt. i_ or		Y	C. G. H.	1732.	C 8.1	
13108 rostr.Muin L.	beaked	£ iA] or	p	Y	C. G. H.	1732.	C 8.1	Di. el. 186.229
13109 tuberculatum Mil.	pimpled	£ iA] or	ap	Y	C. G. H.	1732.	C 8.1	
13110 ramulbusum Haw.	brunihy	n. _ or		Y	C. G. H.	1791.	C 8.1	
13111 pisifirpe Haw.	pea-shaped	£ IAlor		W	C. G. H.	1796.	C 8.1	
13112 moniliforme Haw.	bracelet-shaped	£ iA] or	* mr.ap	W	C. G. H.	1791.	C 8.1	
13113 scalpratinum Haw.	great kmfe-s>hapeiff	£ A1 or	au.o	Y	C. G. H.	1714.	C 8.1	Di. el. 188. %A
13114 fragrans Salm	fragrant	£ iA] or	i	Y	C. G. H.	—	C 8.1	
13115 grandiflorum Haw.	great-flowered	£ iA1 gr		Y	C. G. H.	1824.	C 8.1	
13116 prapfngue Haw.	very fat	£ iA1 or	i jl	Y	C. G. H.	1792.	C 8.1	
13117 médium Haw.	intermediate	£ iAi or	{ au.o	Y	C. G. H.	—	C 8.1	
13118 cultatrum Haw.	cultrate	£ iA] or	£ au.o	Y	C. G. H.	1820.	C 8.1	
13119 Incidum Mil.	shining	£ iAlor	■ au.o	Y	C. G. H.	173.	C 8.1	
13120 adscéndens Haw.	ascending tongue	£ iA1 or	£ aun	Y	C. G. H.	1805.	C 8.1	
13121 pustulatum Haw.	blistered	£ iA1 or	au	Y	C. G. H.	1818.	C 8.1	
13122 longum Haw.	long tongue	£ iA1 or	f au.o	Y	C. G. H.	1725.	C 8.1	Dec. pi. 71
1 deprecsum B.M.	depressed	£ iAJ or	au.o	Y	C. G. H.	—	C 8.1	
2 dee live Haw.	sloping	£ iAJ or	\$ au.o	Y	C. G. H.	—	C 8.1	
3 angustius Haw.	narrower	£ iAl or	■ au.o	Y	C. G. H.	—	C 8.1	
4 purpurascens Haw.	purpurascen	£ iAJ or	f au.o	Y	C. O. H.	1819.	C 8.1	
5 uncatum Haw.	hooked	£ iAJ or	■ au.o	Y	C. G. H.	1819.	C 8.1	
6 attiblens Haw.	elevating	£ iAJ or	\$ au.o	Y	C. G. H.	1819.	C 8.1	
13123 linguaefdrmc Haw.	tongue-bhaped	£ iAJ or	mr.n	Y	C. G. H.	1732.	C 8.1	
2 rufescens Haw.	rufescent	£ iAJ or	g mr.n	Y	C. G. H.	1732.	C 8.1	
3 subrcuatiatum//aw.	subrcuatiac	£ iAJ or	\$ mr.n	Y	C. G. H.	1820.	C 8.1	
4 probratum Haw.	prostrate	£ iAJ or	\ mr.n	Y	C. G. H.	—	C 8.1	
5 liisurgens Haw.	rising	£ iAJ or	mr.n	Y	C. G. H.	1819.	C 8.1	
13124 latum Haw.	broad tongue	£ iAJ or	& mr.n	Y	C. G. H.	ff.20.	C Bl	Di. el. 184.225
2 brève Haw.	short	£ iAJ or	4 m m	Y	C. G. H.	1795.	C 8.1	
13125 depressoium Haw.	depressed tongue	£ iAJ or	\$ s.n	Y	C. G. H.	1802.	C 8.1	
2 lividum Haw.	livid	£ iAJ or	■ s.n	Y	C. G. H.	1819.	C 6.1	Bot. mag. 1866
13126 cruciatum Haw.	cross-leaved	ui_ I or	my.n	Y	C. G. H.	1792.	C 8.1	
13127 taurinum Haw.	bull's-Aorn	n. T_J or	>n	Y	C. G. H.	1795.	C 8.1	
13128 SUMii Haw.	Salm's	n. i_I or	a s.n	*	C. G. H.	1818.	C 8.1	
2 seraicruciatum fiz/m	half-crossed	n. i_I or	1 s.n	Y	C. G. H.	1818.	C 8.1	
3 angustifolium/faw.	narrow-leaved	<1 l or	1 Bn	Y	C. G. H.	1823.	C 8.1	
13129 surréctum Haw.	erect	ft J or	1 an	Y	C. G. H.	1819.	C 8.1	
2 brevifolium Haw.	short-leaved	tL i_J or	js.n	Y	C. G. H.	1819.	C 8.1	
13130 heterophllum Haw.	various-leaved	£ VJor	"\$ mr.o	Y	C. G. H.	1795.	C 8.1	
13131 angustum Haw.	narrow tongue	£ iA1 or	I mro	Y	C. G. H.	1790.	C 8.1	
2 pallidum Haw.	pale	£ iA1 or	mr.o	Y	C. G. H.	1790.	C 8.1	
3 heterophllumjac.	various-leaved	£ iA1 or	au	Y	C. G. H.	1732.	C 8.1	Bot. rep. 540
13132 diffbrme Haw.	deformed	tt. i_ or	£ au	Y	C. G. H.	1820.	C S.1	Di. el. 194.242
13133 bigiberatum Haw.	two-bunched	n. □ gr	1 au	Y	C. G. H.	1H18.	C s.1	
13134 bidentatum Haw.	two-toothed	tL L_J or	■ ■	Y	C. G. H.	1818.	C 8.1	
2 majus Haw.	larger	n. i_ or	f* mr.n	Y	C. G. H.	1732.	C s.1	Di. el. 194.241
13135 semicylindricum Haw.	semicylindriict.	□ or	■■■ R	Y	C. G. H.	1780.	C 6.1	
13136 gibbusum Haw.	gibbous	£ IAJor	■■■ R	Y	C. G. H.	1795.	C 8.1	
13137 luteovinide Haw.	yellowish-greennt.	J or	Jjamy	R	C. G. H.	1792.	C 8.1	
13138 pervlndc Haw.	very green	■■■ Alor	ja.my	R	C. G. H.	1702.	C 8.1	
13139 pubescens Haw.	pubescent	■■■ Alor	1 jls	W	C. G. H.	1717.	C 8.1	Dec pi. 5
13140 calamiforme L.	Heed-shaped	■■■ Alor	1 ...	W	C. G. H.	1796.	C 8.1	
13141 olisubulatum Haw.	obdubulatc	tt. □ or	1 f 8	R	C. G. H.	1792.	C 8.1	
13143 cylindricum Haw.	cylindrical	■■■ Alor	I f.s	R	C. G. H.	1794.	C 8.1	
13143 teratifolium Haw.	round-leaved	■■■ Alor	4 ...	R	C. G. H.	1794.	C 8.1	
13144 toretifisculum 77aw.	rather rounder	■■■ Alor	i ■■■	R	C. G. H.	1717.	C Bl	Di. el. 224.223
13145 tellidifbrum L.	Daisy.flowered	£ iAJ or	i jn.au	R	C. G. H.	1717.	C Bl	
2 subuKitum Mil.	awl-shaped	£ iAJ or	£ jn.au	R	C. G. H.	1717.	C Bl	
3 viride Haw.	pea-green	£ iAJ or	■■■ R	Y	C. G. H.	1793.	C 8.1	
13146 acutum Haw.	great acutclvd	£ iAJ or	*apn	R	C. G. H.	17W.	C 8.1	
13147 punctatum Haw.	dotted awl-lyd	£ iAlor	£ ap.n	R	C. G. H.	1789.	C 8.1	
13148 diminutum //aw.	diminished	■■■ Alor	4 ap	R	C. G. H.	1789.	C S.1	
2 cauliculatum Haw.	small-stemmed	IA or	i ap	R	C. G. H.	1789.	C S.1	

II. FASCICULAVA.—Leaves in bundles ; flowers polygynous ; calyx five-leaved.

13149 breum Dil.	thong.sta/ked	£ IAJor	* 1 s	Pa.Y	C. G. H.	1732.	C Bl	Di.eL200.255
2 congestum Haw.	crowded	■■■ Alor	1 s	Pa.Y	C. G. H.	1805.	C Bl	
13150 purp&reo-album/faw.	purplish white	£ IAJgr	Jau	W	C. G. H.	1824.	C si	
13151 diTersibium L.	various-leaved	£ iAlor	1 mr.o	Pa.Y	C. G. H.	1819.	C s.1	Di. el. 19a 253
2 glaucius Haw.	more glaucous	£ iAlor	1 au	Pa.Y	C. G. H.	1726.	C s.1	
3 brevifolium Haw.	short-leaved	£ iAlor	1 au	Pa.Y	C. G. H.	1726.	C s.1	
4 la?le vlyrens Haw.	bright green	£ iAlor	1 au	Pa.Y	C. G. H.	1726.	C s.1	
5 ^tMMrrens Haw.	dark-green	£ iAlor	1 au	Pa.Y	C. G. H.	1726.	C s.1	
■■■ piens/faw.	deceiving	■■■ Alor	1 au	Pa.Y	C. G. H.	18~0.	C 8.1	
■■■ Alor	doubtful	■■■ Alor	1 my n	Pa.Y	C. G. H.	1800.	C s.1	Brad. sue. 4.40
■■■ Alor	small-horned	■■■ Alor	1 mr.my	Pa.Y	C. G. H.	1732.	C s.1	
■■■ Alor	equal.leaved	■■■ Alor	1 mr.my	Pa.Y	C. G. H.	1732.	C s.1	Dec.pi. 108
■■■ Alor	procumbent	■■■ Alor	1 mr.my	Pa.Y	C. G. H.	1820.	C 8.1	
13156 tricolorum Haw.	three-colored	■■■ Alor	1 o	Y.R	C. G. H.	1794.	C 4.1	Bot. mag. 2144

13157 pugionifl. <i>Srmel.</i> <i>long</i> dagger, formed tt. i_J or	1 jl.s	Pa.Y C G. H.	1714.	C i.l	Di.el.269
2 <i>clavatum</i> <i>Haw.</i> <i>tiesb-colored</i> tt. L_J or	1 jl.s	Pk C G. H. tt	1714.	C B.1	
3 <i>purpureum</i> <i>Haw.</i> <i>purple</i> tt. L_J or	1 jl.s	Pk C G. H.	1714.	C s.l	
4 <i>bilinnc Jfew.</i> <i>biennial</i> tt. L_J or	1 jl.s	Pa.Y C. G. H.	1714.	C s.l	
13158 <i>capitatum</i> <i>Haw.</i> tt. i_J or	JU	Pa.Y C. G. H.	1717.	C L.	Bot reg. 494
2 <i>ramiifwrum</i> //J. <i>branch-bearing</i> tt. cu	jl.s	Pa.Y C. G. H.	1816.	C 8.1	
13159 <i>urevicaule</i> <i>Haw.</i> <i>short-stemmed</i> tt. (J or	jl.a	Pa.Y C. G. H.	1820.	C 8.1	
13160 <i>coruscans</i> //sw. <i>glittering</i> <i>leggg*</i> . L_J or	jl.s	Pa.Y C. G. H.	1791.	C s.l	
13161 <i>clongMum</i> <i>Haw.</i> <i>elongated tuberous</i> & L_J or	my	Pa.Y C. G. H.	1793.	C 8.1	Bot. reg. 493
2 minus <i>Jtow.</i> <i>smaller</i> {L_J"}	my	Pa.Y C. G. H.	1793.	C 8.1	
3 <i>fusif6nne</i> <i>Haw.</i> <i>fusiform</i>	my	Pa.Y C. G. H.	1793.	C 8.1	

III PROBTBA VA.—*Trailers; flowers polygynous; calyx five-leaved; leaves connate.*

13162 <i>geminatum</i> <i>Jac.</i> <i>twin</i> %* i or	-	Pk C. G. H.	1819.	C 6.1	Jac.fr. 50
13163 <i>simile</i> <i>Haw.</i> <i>similar</i> -	i	Pk C. G. H.	1819.	C t.1	
13164 <i>laxum</i> W. <i>loose</i> -	-	Pk C. G. H.	1820.	C 8.1	
13165 <i>sarmenobsum</i> <i>Haw.</i> <i>twiggy</i> JU(J or	-	Pk N. Holl.	1805.	C B.1	
13166 <i>rigidiflde</i> <i>Haw.</i> <i>stiff-stemmed</i> i v i lor	i my.jn	Pk C. G. H.	1819.	C 8.1	
13167 <i>viliqum</i> <i>Haw.</i> <i>robust</i> 1" my.jn	Pk C. G. H.	1810.	C 8.1		
13168 Scholl/i Salm Sell <i>oil's</i> Pk C. G. H.	1732.	C 8.1			Di. el. 212.273
13169 <i>filamentosum</i> <i>Haw.</i> <i>thready</i> £mr.ap	Pk C. G. H.	1795.	C 8.1		
13170 <i>serrulatum</i> <i>Haw.</i> <i>saw-leaved</i> i n.d	Pk C. G. H.	1802.	C 8.1		
2 <i>viridius</i> <i>Haw.</i> <i>greener</i> **. LJ or	[n.d]	Pk C. G. H.	1802.	C 8.1	
13171 <i>ruhriate</i> <i>Haw.</i> <i>red-stalked</i> JU i_J or	-	Pk C. G. H.	iaia	C 8.1	
2 <i>densius</i> <i>Haw.</i> <i>denser</i> *1 i or	Pk C. G. H.	1818.	C 8.1		
3 <i>subvirens</i> <i>Haw.</i> <i>tall</i> <i>greenish</i> JU LJ or	aus.s	Pk C. G. H.	1714.	C 8.1	
13172 <i>acinaeiforme</i> L. <i>Bcinfar-forrad</i> JU \ or	1 aus.s	Pk C. G. H.	216ngum	C 8.1	
216ngum <i>Haw.</i> <i>long</i> JU \ or	Pk C. G. H.	1811.	C s.1		
13173 <i>Isvig&tum</i> <i>Haw.</i> <i>polished</i> *au	Pk C. (S. H.	C 8.1			
13174 <i>rubrocinctum</i> <i>Haw.</i> <i>red-bordered</i> *au	Pk C. G. H.	1768.	C s.1		Dec. pi. 41
2 <i>compressum</i> <i>Haw.</i> <i>compressed</i> -	Pk C. G. H.	1804.	C s.1		
3 <i>tenerum</i> <i>Haw.</i> <i>tender</i> -	Pk C. G. H.	1835.	C s.1		
13173 <i>subpulchrum</i> <i>Haw.</i> <i>subul.</i> <i>Davy-ftwd</i> J~ i_J or	i)lau	Pk C. G. H.	1811.	C 8.1	Dec. pi 39
13176 <i>edule</i> L. <i>eatable</i> <i>Hottentot Fig</i> *~ i_J or	Pk V. D.L.	1820.	E 8.1		
\$13177 <i>dimidiatum</i> <i>Haw.</i> <i>halved Hot. Fig</i> J~ L_J or	Pk N. Holl.	1804.	C s.1		
13178 <i>glaucescens</i> <i>Haw.</i> <i>glaucescent</i> !U i or	Pk N. Holl.	1791.	C 8.1		
13179 <i>abbreviatum</i> <i>Haw.</i> <i>short-jointed</i> (U) i gr	Pk C. G. H.	1821.	C 8.1		
13180 <i>Kiss</i> <i>Haw.</i> <i>Ross's</i> -	Pk C. G. H.	1774.	C 8.1		
13181 <i>viridescens</i> <i>Haw.</i> <i>viresent</i> -	Pk C. G. H.	1824.	C 8.1		
13182 <i>equilaterale</i> <i>Haw.</i> <i>equal-sided</i> JU i or	Pk N. Holl.	1773.	C 8.1		
13183 <i>viridis</i> <i>Haw.</i> <i>upright-green</i> JU i or	Pk N. Holl.	1804.	C 8.1		
13184 <i>reptans</i> H. K. <i>creeping</i> -	Pk N. Holl.	1803.	C 8.1		
13185 <i>debile</i> <i>Haw.</i> <i>smooth weak</i> < *	Pk N. Holl.	1803.	C 8.1		
13186 <i>austrMe</i> <i>Haw.</i> <i>southern</i> i jl.au	C. G. H.	1727.	C 8.1		DL el. 201.257
13187 <i>crassifolium</i> L. <i>thick-leaved</i> JU(J or	imy.au	N. Holl.	1803.	F	
13188 <i>clavellatum</i> <i>Haw.</i> <i>small cUh-lvd</i> JU(J or	* jn jl	Pk N. Holl.	1810.	F	
2 minus/fotu.		Pk N. Holl.	1803.	C 8.1	
3 <i>aggregatum</i> <i>Haw.</i> <i>aggregate-wrf</i> 4 Jn.JJ	Vjl.au	Pk N. Holl.	1803.	C 8.1	

IV.

13189 <i>forficatum</i> L. <i>sensor-leaved</i> i_J ..	V	3L C. G. H.	1792.	C 8.1	1.26
J3H0 <i>geminatum</i> <i>Haw.</i> <i>twin-shoted</i> *~ i or	W	C. G. H.	1793.	C s.1	
13im <i>Inaninatum</i> <i>Haw.</i> <i>while-edged</i> i_J or	W.pk	C. G. H.	1820.	C s.1	
13192 <i>rostellatum</i> <i>Haw.</i> <i>little beak</i> <. i_J or	jn.au	Pa.P C. G. H.	1714.	C s.1	Di. el. 192.240
1319J <i>perfolidatum</i> Mil. <i>great perfoliate</i> i_J Jof	jn.au	Pa.P C. G. H.	1819.	C 8.1	Di.el.239
2 <i>monocanthum</i> Biad. <i>one-spined</i> <L or	jn.au	Pa.P C. G. H.	1725.	C 8.1	Dec pi 54
13194 <i>uncinellum</i> <i>Haw.</i> <i>small-hooked</i> << i_J or	au	Pa.P C. G. H.	... C. G. H.	C ■	
13195 <i>uncinatum</i> <i>Haw.</i> <i>hooked</i> << i_J or	au	Pa.P C. G. H.	1824.	C ■	
131 P7 <i>eindens</i> <i>Haw.</i> <i>one-toothed</i> -	W	C. G. H.	1792.	C 8.1	
13198 <i>viride</i> <i>Haw.</i> <i>green</i> petiolate' iL or	W	C. G. H.	1821.	C s.1	
13199 <i>acutangulum</i> <i>Haw.</i> <i>acute-angled</i> tt. L_J or	W	C. G. H.	1822.	C s.1	
13200 <i>curtum</i> <i>Haw.</i> <i>shorusheathed</i> < i_J or	W	C. G. H.	1823.	C s.1	
2 <i>majus</i> <i>Haw.</i> <i>larger</i> tt. L_J or	W	C. G. H.	1824.	C s.1	
3 <i>politum</i> <i>Haw.</i> <i>polished</i> tt. L_J or	W	C. G. H.	1825.	C s.1	
4 <i>minus</i> <i>Haw.</i> <i>smaller</i> tt. L_J or	W	C. G. H.	1826.	C s.1	
11201 <i>yaginiatum</i> <i>Haw.</i> <i>sheathed</i> tL L_J or	W	C. G. H.	1802.	C 8.1	
2 <i>pariflurum</i> <i>Haw.</i> <i>small-flowered</i> s. J_J or	W	C. G. H.	1800.	C 8.1	
13202 <i>pamflurum</i> <i>Haw.</i> <i>small-flowered</i> s. J_J or	W	C. G. H.	1793.	C 8.1	
13203 <i>rfgidum</i> <i>Haw.</i> <i>rigid</i> tt. L_J or	W	C. G. H.	1792.	C 8.1	
13204 <i>tenellum</i> <i>Haw.</i> <i>delicate</i> <i>perfoliate</i> <. L_J or	W	C. G. H.	1799.	C B.1	
13205 <i>imbrickatum</i> //, JL <i>imbricated</i> tt. L_J or	W	C. G. H.	1799.	C B.1	
2 <i>medium</i> <i>Haw.</i> <i>intermediate</i> tt. L_J or	W	C. G. H.	1802.	C B.1	
3 <i>viride</i> <i>Haw.</i> <i>green</i> tt. L_J or	W	C. G. H.	1803.	C B.1	
13206 <i>multiflorum</i> <i>Haw.</i> <i>many-flowered</i> tt. i or	W	C. G. H.	1792.	C 8.1	Pluk. ph. 117.1
2 <i>m'uis</i> <i>Haw.</i> <i>smaller</i> tL i or	W	C. G. H.	1793.	C B.1	
3 <i>rubrum</i> <i>Haw.</i> <i>rciUflowered</i> tt. J_J or	Pk	C. G. H.	1820.	C B.1	
4 pKtens W. <i>spreading</i> B- i_J or	W	C. G. H.	1820.	C B.1	
5 rtens <i>Haw.</i> <i>sinning</i> tL i_J or	W	C. G. H.	1727.	C 8.1	Di. el. 208.266
13207 URibeflatum <i>Haw.</i> <i>umbel-flowered</i> tt. J_J or	W	C. G. H.	1727.	C 8.1	
2 mnömluin W. <i>anomalous</i> tt. J_J or	W	C. G. H.	1802.	C 8.1	
13208 tumfdulum <i>Haw.</i> <i>rather tumid</i> tt. J_J or	W	C. G. H.	182a	C 8.1	
2 minus <i>Haw.</i> <i>smaller</i> tt. J_J or	W	C. G. H.	1802.	C 8.1	
13209 fohosutn <i>Haw.</i> <i>leafy</i> tt. J_J or	W	C. G. H.	1819.	C 8.1	
13210 lineolatum <i>Haw.</i> <i>small-lined</i> tt. J_J or	JU	Pk" CG. H.	181U.	C 8.1	
2 minus <i>Haw.</i> <i>smaller</i> tt. J_J or	JU	Pk" CG. H.	181U.	C 8.1	
3 le*ve Thun. <i>smooth</i> tt. J_J or	JU	C. G. H.	1819.	C s.1	
4 nltens <i>Haw.</i> <i>shining</i> tt. J_J or	JU	C. G. H.	1819.	C 8.1	

V DBLTJEFULIA.—Leaves more or less deltoid or hatchet-form.

J321 aerrXtum I k <i>puspworded</i> <- U or	2 in.jl	Pk "C. G. H."	1707.	C s.1	Di. el 191 \$38
J3213 Reteropetalumffaw. various-petldcd tt. U or	2 my"au	Pk C. G. H.	1794.	C 8.1	
1sor	glaucous tL ^ or	Pk C. G. H.	-	C B.1	
thick-leaved ii-LJor 1] jlau	Pk C. G. H.	-	C 8.1		

- | | | | | | | | | | | | | | |
|-------|-------------------|-----------------|--------------|----------|-----|----------|----------|-------|------|------|------|-----------|-----------|
| 13215 | mutifibilefotu. | changeable | tt Jor | lftjl.8 | Pk | C. G. H. | 1792. | C | 81 | Dec | pi. | 60 | |
| 13216 | inclafidens Haw. | encompassed | tL _ or | lft jn.s | Pk | C. G. H. | 1805. | C | B.1 | Bot. | rep. | 388 | |
| 13217 | caulescens Mil. | caulescent | delta-lvd*. | 1* my.jl | Pk | C. G. H. | 1731. | C | 81 | Di. | el. | 195.243-4 | |
| 13218 | deltoidium Haw. | great dclta-/ud | a. L _ or | lft my | Pk | C. G. H. | 1731. | C | 81 | Dec. | pi. | 58 | |
| 13219 | muricatum/ftwi. | muricated | delta-iudt*. | I _ or | Mmy | Pk | C. G. H. | 1731. | C | B.1 | Di. | el. | 195.245-7 |
| | 2 minus Haw. | less | tL__ or | lft my | Pk | C. G. H. | __ | C | s.1 | | | | |
| 13220 | microphyllum Haw. | small-leaved | tL _ or | i | Pk | C. G. H. | 1795. | C | 8.1 | | | | |
| 13221 | mucronatum Haw. | mucronated | __ or | __ - | Pk | C. G. H. | 1794. | C | 8.1 | | | | |
| 13222 | pygmae um Haw. | pygmy | tL _ (or | __ - | Pk | C. G. H. | 1805. | C | 8.1 | | | | |
| 13223 | pulchellum Haw. | pretty | tLLjor | __ ap | Pk | C. G. H. | 1793. | C | 8.1 | | | | |
| | canescens Haw. | | | | | | | | | | | | |
| | 2 revolutum Haw. | revolute | tL i Jor | __ ap | Pk | C. G. H. | __ | C | __ | __ | __ | __ | |

VI. THI'QUBTBA.—Leaves more or less three-cornered, distinct; calyx five-leaved, styles five.

- | | | | | | | | | | |
|-------|---------------------------|-----------------------|--------|-----------|----------|-----------|--------|----------|-------------------|
| 13224 | máximum <i>Haw.</i> | largest moon-lyd <L i | or | lft mr.au | Pk | C. G. H. | 1787. | s.1 | Bot rep. 358 |
| 13225 | luníttum <i>W.</i> | lunate n. | or | 1 Jl | Pk | C. G. H. | 1812. | | Di.eL 213. 275.6* |
| 13226 | falcatum <i>L.</i> | sickle-leaved tt. | or | 1 jn.au | Pk | C. G. H. | 1727. | | |
| 13227 | dec&embens <i>Haw.</i> | decumbent tt. | or | 1 my.o | Pa.R | C. G. H. | 1759. | | |
| 13228 | incirvum <i>Haw.</i> | incurred tt. | or | lftjnj | Pk | C. G. H. | 1802. | | |
| | 2 dilatans <i>Haw.</i> | spreading fw/tctftt. | or | lft jn | Pk | C. G. H. | — | | |
| | 3 pallidus <i>Haw.</i> | paler a. | or | 1 Jl | Pk | C. G. H. | — | | |
| | 4 densifolium <i>Haw.</i> | dense-leaved tL | or | lft jn | Pk | C. G. H. | 1819. | | |
| | 5 rubeum <i>JV.</i> | rosy tL | or | Jftjnl | Pk | C. G. H. | — | | |
| 13229 | confertum <i>Haw.</i> | crowded-leaved tL | i J or | lft BO | Pk | C. G. H. | 1805. | | |
| 13230 | falcifürme <i>Haw.</i> | sickle-shaped tt. | i J or | lft jl.au | Pk | C. G. H. | 1805. | | |
| 13231 | glomérfatum <i>L.</i> | clustered tL | uJ or | lft jnau | Pk | C. G. H. | 1732. | Bot cab | 1043 |
| 13232 | inflexum <i>Haw.</i> | inflexed tL | i J or | lft jn.au | Pk | C. G. H. | 1819. | | |
| 13233 | scabrum <i>L.</i> | scabrous tL | i J or | lft jl | P | G. H. | 1731. | | |
| | 2 purpureum <i>Haw.</i> | purple tL | i J or | 1 jl | Pk | G. H. | 17.11. | | |
| 13234 | versicolor <i>Haw.</i> | changeable-cld tt. | i J or | 1 my.au | Pk | G. H. | 1795. | | |
| 13235 | retroflexum <i>Haw.</i> | retroflexed tt-Ljor | | lft my.o | Pk | G. H. | 1724. | | |
| 13236 | imbricans <i>Haw.</i> | imbricating XL | or | 2 my.o | Pk | G. H. | 1818. | | |
| 13237 | deflexum <i>H. K.</i> | deflexed tL | or | 1 jl.o | Pk | G. H. | 1774. | | |
| 13238 | lcptálezon <i>Haw.</i> | slender tL | or | HJLo | Pk | G. H. | 1803. | | |
| 13239 | polyanthon <i>Haw.</i> | many-flowered tL | or | 1 au | .Pk | G. II. | 1870. | | |
| 13240 | flexile <i>Haw.</i> | Jit A IIC | ■■■ | lft au | Pk | G. H. | 1819. | | |
| 13241 | polypy'lum <i>Haw.</i> | many-leaved tt. | or | 2 jn.o | P | G. H. | 1820. | | |
| 13242 | violaceum <i>Dec.</i> | violet tt. | or | 2 jn.o | Pk | G. H. | 1732. | | |
| 13243 | emarginátum <i>L.</i> | notch-réouvrail tL | or | 2 jn.o | W | G. H. | 1820. | | |
| 13244 | dilatátum <i>Haw.</i> | dilated tL | or | 2 jn.au | Pk | G. H. | 1793. | | |
| 13245 | virgátum <i>Haw.</i> | twiggy tL | or | 3 jn.au | Pk | G. H. | 1774. | | |
| 13246 | bractéatum <i>Haw.</i> | bracted tL | or | 3 f.ap | Y | G. H. | 1811. | Bot cab. | 251 |
| 13247 | apicáps <i>Haw.</i> | two-edged tL | or | Ujl.o | Pk | G. H. | 1819. | | |
| | 2 pallidum <i>Haw.</i> | pale tL | i J or | lft s.o | P.Pk | C. G. H. | 1794. | | |
| 13248 | gracíle <i>Haw.</i> | slender-starry tL | J or | lft n | R | C. G. H. | — | | |
| | stellatulum <i>Haw.</i> | lft au.n | | | | | | | |
| | 2 gracilius <i>Haw.</i> | slenderer tL | i J or | 1 au.n | R | C. G. H. | — | C | s.l |
| 13249 | radiátum <i>Haw.</i> | rayed tL | i J or | 1 au.n | R | C. G. H. | 172. | C | s.l |
| 13250 | comproMSum <i>Haw.</i> | compressed tt. | i J or | <*JS" | D.R. | C. G. II. | 1792. | C | s.l |
| 13251 | platínum <i>Haw.</i> | spreading tL | i J or | Pk | C. G. H. | 1811. | C | s.l | |
| 13252 | aspercum <i>Haw.</i> | rough tL | i J or | lto.n | — | C. G. H. | 1818. | C | 8.1 |
| | 2 escrúscens <i>Haw.</i> | eaerúlescent tL | i J or | 1 au.8 | C | C. G. H. | 1820. | C | s.l |
| 13253 | formísum <i>Haw.</i> | beautiful tt. | i J or | 1 my.au | C | C. G. H. | 1787. | C | s.l |
| 13254 | siectábil <i>Haw.</i> | showy tt. | i J or | 1 8.o | C | C. G. H. | 1806. | C | s.l |
| 13255 | conspícum <i>Haw.</i> | conspicuous n. | J or | lftjnj | Pk | C. G. H. | 1810. | C | 8.1 |
| 13256 | bländum <i>Haw.</i> | bland tL | i J or | 2 jn | W | C. G. H. | 1818. | C | s.l |
| 13257 | curviflúrum <i>Haw.</i> | curve-flowered tL | i J or | 1 mr.o | Y | C. G. H. | 1750. | C | s.l |
| 13258 | affíreum <i>L.</i> | gohlcn-JlowcrediL | J or | 1 ... | C | C. G. H. | 1822. | C | 8.1 |
| | cymbífolium <i>Haw.</i> | boat-leaved tt. | J or | lftjnj.au | Y | C. G. H. | 17H3. | C | s.l |
| 13260 | aurantium <i>Haw.</i> | orange-flowered*- uJ | or | lft jn.jl | O | C. G. H. | 1696. | C | si |
| 13261 | glaticum <i>L.</i> | glaucous-leaved* i J | or | 3 | Y | C. G. H. | 1795. | C | Dec. pi. 148 |
| 13262 | strictum <i>Haw.</i> | strict tL | i J or | 1 | Y | C. G. H. | 1793. | C | s.l |
| 13263 | cymbíiforme <i>Haw.</i> | boat-shaped tL | i J or | 1 jl.aili | Y | C. G. H. | 1821. | C | 8.1 |
| 13264 | muconífórmehaw | sword-shaped tL | i J or | ft 8.0 | Y | C. G. H. | 1727. | C | Brad. sue. 2.20 |
| 13265 | granítifórmehaw | grain-shaped tL | i J or | 1 o | Pk | C. G. H. | 1774. | C | 8.1 |
| 13266 | m61le <i>H. K.</i> | wafí-leaved tt | LJ | | | | | | |

VII. FILICAU'LIA.—*Sums slender; leaves distinct, rounded, and dotted; flowers opening in the morning.*

- | | | | | | | | | | | | | |
|---|---------------|---------------------|----------------------|-------|---------|----------|-------|--------|--------|-------|----------------|--|
| 13267 | coccineum | Haw. | scarlet-flowered | tL | J or | lft my.s | S | CG. H. | 1696. | C si | Bot. mag. 59 | |
| 2 | actius | Haw. | acuter-calyxed | tL | J or | Jt my.s | S | CG. H. | ... | C 8.1 | | |
| 3 | minus | Haw. | smaller | tt. | J or | lft my.s | S | CG. H. | ... | C 8.1 | | |
| 13268 | bicolorum | L. | two-colored | tt. | J or | 1 my.s | O | CG. H. | 1732. | C B.1 | Di. el 202.258 | |
| 2 | patulum | Haw. | spreading | tt. | J or | 1 my.s | O | CG. H. | — | C s.1 | | |
| 3 | minus | Haw. | smaller | tt. | J or | ft my.s | O | CG. H. | — | C B.1 | | |
| 13269 | inajquale | Haw. | unequal-cifijipaitt. | tt. | J or | 1 my.s | O | CG. H. | 1716. | C 8.1 | Brad, sue 1.7 | |
| 13270 | tenuifolium | L. | slender-leaved | tt. | J or | I jns | S | CG. H. | 1700. | C s.1 | Dec. pi. 82 | |
| 2 | erectuum | Haw. | erect | tt. | J or | lftjn.8 | ■ | CG. H. | ... | C 8.1 | | |
| 13S71 | variabilis | Haw. | variable | tt. | J or | ft | jn.au | Y | CG. H. | ... | C 8.1 | |
| 2 | lseMus | Haw. | smoother | tL | V or | lftjn.au | Y | CG. H. | 1796. | C 8.1 | | |
| 13272 | spiniforme | Haw. | thorn-shaped | tt-L | J or | 1 | 8.0 | Pk | CG. H. | 1793. | C 8.1 | |
| 2 | subaduncum | Haw. | rather hooked | tt. | LJ or | 1 | 8.0 | Pk | CG. H. | ... | C 8.1 | |
| 13273 | curvifolium | W. | crooked-leaved | n. | LJ or | 1 | O | Pk | CG. H. | 1799. | C 8.1 | |
| 13274 | flexifiblixn | Haw. | bent-leaved | tt. | LJ or | Ifto | Pk | CG. H. | 1820. | C 8.1 | | |
| 2 | laet'e virens | Haw. | lively green | tt. | LJ or | lft | o | Pk | CG. H. | 1818. | C 8.1 | |
| 13275 | adfinicum | Haw. | hook-leaved | tt. | LJ or | 1 | f.mr | Pk | CG. H. | 1795. | C 8.1 | |
| 13276 | inconspicuum | Haw. | inconspicuous | tt. | LJ or | ft | jns | R | CG. H. | 1823. | C 8.1 | |
| 13277 | filicaule | Haw. | thread-stalked | tt. | J or | lft | a | Pk | CG. H. | 1800. | C s.1 | |
| 13278 | spinatum | L. | thorny | tL | J or | lft | jns | Pk | CG. H. | 1714. | C s.1 | |
| 13279 | stipulaceum | /.. | stipulaceous | tt. | J or | lft | my.jn | Pk | CG. H. | 1723. | C s.1 | |
| 13280 | cor&ellinum | Thun. | coral | tt. | J or | 1 | my.jn | Pk | CG. H. | 1820. | C si | |
| (%E) productum Haw. prolonged-calyxcdn. i | | | | | | | | | | | | |
| 1£ f. & L. P. ni | Haw. | pretty white-flwdt | uL | uj or | lft | a.us | W | CG. H. | 1823. | C s.1 | | |
| 13M3 | Haworth's | Donn | Haworth's | tt. | LJ or | 1 | jajn | Br | CG. H. | 1793. | C s.1 | |
| 13284 | 1*Te H. K. | smooth white-wooded | tt. | LJ or | lftjl.8 | . | . | CG. H. | 1774, | C 8.1 | | |

13285 verruculatum L.	small-warted	«Li lor	If ray.jn	Y	C. G. H.	1731.	C 8.1	Di.el.859
2 Candolff	Dccandolle's	a. i lor	I* my.jn	Y	C. G. H.	-	C 8.1	Dec. pi. 36
13286 insilltium TV.	grafted	a.i lor	1 au.o	P	CG.H.	1780.	C 8.1	
purpäreo-crbccum	Haw.							
2 flvo-crbceuin	Haw. yellow & safC	tt. or	1 au.o	Y	CG. H.	1816.	C 8.1	
3 minus	smaller	n. / or	au.o	Y	C. G. H.	-	C 8.1	
13287 luteum Haw.	yellow	a. or	1J my.au	Pa.Y	C. G. H.	1824.	C 8.1	
13288 luteolum Haw.	yellowish	a. i or	i my.au	Pa.Y	C. G. U.	1824.	C 8.1	

VIII. VERWJCO*SA. — Leaves and branches almost always more or less warty or papilous.

13289 crystallinum L.	crystalline	J± O or	1 my.au	W	Greece	1725.	S e.1	Dec. pl. 128	
13290 glaciXle Haw.	frozen	Jc o or	f my.au	W	Greece	-	S 8.1		
13291 pinnatifidum L.	wing-cleft	Jc O or	1 my.o	Y	C. G. H.	1774.	S 8.1	Bot mag. 67	
13292 sessiliflrum H. K.	sessile-flowered	Jc iQj or	\$ jl	Y	C. G. H.	1774.	S 8.1		
2 filbum Haw.	white	dk iQj or	f jl	W	C. G. H.	-	S 8.1		
13293 humifTisum H. K.	trailing	a. \ or	Jlau	W	C. G. H.	1774.	C 8.1		
13294 clandestine Haw.	clandestine	<.L-Jor	j my.au	W	C. G. H.	1822.	C 8.1		
13295 Aituni Jac.	Alton's	£ OJ or	J jn.o	Pk	CG.H.	1774.	S 8.1	Jac. vin. 3.7	
13298 lanceolatum Haw.	small-spear-A/f	O or	f my.au	W	C. G. H.	1795.	S 8.1		
2 rbscum Haw.	rosy	iQj or	my.au	Pk	C. G. H.	1813.	S 8.1		
13297 cordiflum L.	heart-leaved	ft i or	my.s	Pk	C. G. H.	1774.	C 8.1	Dec. pi. 102	
13298 pomeridianum L.	afternoon	■■■■■	1 jlau	Y	C. G. H.	1774.	S 8.1	Bot. mag. 540	
2 Andrëws ²	Andrews ² s	■■■■■	1 jlau	T	C. G. H.	-	S 8.1	Bot rep. 57	
J32S9 pUusUm Haw.	hairy yellow	■■■■■	f jii.au	Y	C. G. H.	1800.	S 8.1		
13300 calendulaceum/iit. Pot-marigold	fau	Y	C. G. H.	1819.	S 8.1				
13301 heliantholdcs H.K.	Sunflower-like	■■■■■	\$ au.o	Y	CG.H.	1774.	S 8.1	Dec. pi. 135	
Cand611ii Haw.									
13302 limpidum II. K.	clear	O or	1 j,	R	C. G. H.	1774.	S 8.1	Jac. ic 3.488	
13303 tricolor Haw.	three-colored	O or	R	C. G. H.	1795.	S 8.1	Bot. mag. 2144		
13304 rbscum Haw.	roseate	O or	■■■■■	Pk	C. G. H.	■■■■■	S 8.1		
3 linearc Thun.	linear	O or	■■■■■	W	C. G. H.	1819.	S 8.1		
4 album	vrhMe-Jlowering	iQj or	■■■■■	W	C. G. H.	1819.	S 8.1		
13305 villusum L.	villous tt.	iJ or	■■■■■	W	C. G. H.	1759.	C 8.1		
13306 caducum H. K.	deciduous	iQj or	jlau	■■■■■	C. G. H.	1774.	S 8.1		
13307 apetalum H. K.	metalless	O or	jlau	■■■■■	C. G. H.	1774.	S 8.1	Jac. vin. 3.6	
13308 nodiflrum L.	knot-flowered	iQj or	au.o	■■■■■	Egypt	1739.	S 8.1	Dec. pi. 88	
13309 cihatum H. K.	ciliated	O or	1 ...	W	C. G. H.	1774.	C 8.1		
13310 geniculiflorum L.	joint-flowering	iQj or	*jn.1	W	C. G. H.	1727.	C 8.1	Dec.pl. 17	
13311 Ttpbhum £.	Aiter-Icaved	i or	jn.o	Pa.Y	C. G. H.	1700.	C 8.1	Di. el. 179.220	
13312 expansum L.	expanted/vd	□ or	4jl.au	Pa.Y	C. G. H.	1705.	C 8.1	Dec. pi. 94	
13313 vriana Haw.	varying	tt.	i or	4^jl.o	Pa.Y	C. G. H.	1706.	C 8.1	Pet. g. 78.10
13314 tortubsum L.	twisted-leaved	□ or	*jn.o	Pa.Y	C. G. H.	1705.	C 8.1	Di. cl. 181. 222	
13315 pallens H. K.	jirte-floivcred	□ or	J j.lau	Pa.Y	C. G. H.	1774.	C 8.1	Dec. pi. 47	
13316 loratum Haw.	loratc	tt.	i or	flau	W	C. G. H.	1819.	C 8.1	
13317 relaxatum W.	relaxed	tt.	(or	I jlau	Pk	C. G. H.	1815.	C 8.1	
13318 crassicalyx Haw.	thick-stemmed	tt.	L or	1 jl.au	PILY	C. G. H.	1815.	C 8.1	
13319 anatbicum Haw.	skeleton-leaved	tt.	□ or	fjlau	W	C. G. H.	1803.	C 8.1	
2 fragile Haw.	brittle	tt.	i or	£ jlau	W	C. G. H.	1803.	C 8.1	
13320 rectum Haw.	straight	tt.	J or	fjl.au	W	C. G. H.	1819.	C 8.1	
13321 crassuloides Haw.	Crassula-like	tt.	J or	ijl.au	Pk	C. G. H.	1819.	C 8.1	
13322 incóptum Haw.	untrimmed	tt.	iJ or	4Jlau	W	C. G. H.	1819.	C 8.1	
13323 spfident L.	shining	tt.	LJ or	U jn.au	W	C. G. H.	1716.	C 8.1	
J3324 flexubsum Jfotu.	flexuous	tt.	J or	1jl.au	W	C. G. H.	1793.	C 8.1	DeepL85
13325 acumintum flinw.	acuminate	tt.	i-j or	2 au.s	W	C. G. H.	1820.	C 8.1	
13326 ainicaule Haw.	white-stemmed	a. i gr	1 au	W	C. G. H.	1819.	C 8.1		
13327 sulcatum //ow.	furrowed	tt.	J or	3 au.s	W	C. G. H.	1794.	C 8.1	
13328 fastigctum Haw.	peaked	tt.	LJ or	li ■■■■■	W	C. G. H.	1792.	C 8.1	
2 reflexed //kw.	reflexed	tt.	V or	U aus	W	C. G. H.	1820.	C 8.1	
13329 umbelliflrum » ^r .	uinbellate-flwd	tt.	□ or	■■■■■ aus	W	C. G. H.	1820.	C 8.1	
13330 paltsccns Haw.	pullescent	tt.	□ or	ii aas	W	C. G. H.	1820.	C 8.1	
13331 micranthon Haw.	small-flowered	tt.	iJ or	—	W	C. G. H.	1804.	C 8.1	
parviflrum Jac.									
13332 Juncum Haw.	Hush-leaved	tt.	i or	1 au.o	Pk	C. G. H.	1800.	C 8.1	
13333 granulicadle Haw.	granulated-stmd	tt.	□ or	—	... C. G. H.	1820.	C 8.1		
13334 Üne Haw.	slender	tt.	□ or	—	... C. G. H.	1819.	C 8.1		
JJ335 longispinulum Haw.	long-spinuled	tt.	J or	1 au.n	Pa.Y	C. G. H.	1820.	C 8.1	
13336 spinuliferum Haw.	suinulc-bearing	tt.	LJ or	1 jn.o	Pa.Y	C. G. H.	1754.	C 8.1	
13337 gr6ssum Haw.	thick	tLi lor	1 au.o	Pa.Y	C. G. H.	1774.	C 8.1		
13338 salmboicum Haw.	salmon-colored	tt.	iJ or	3 au.o	Pa.Y	C. G. H.	1819.	C 8.1	
13339 canaliculMum Haw.	small-chan-/vi	tt.	J or	2 JL	Pk	C. G. a	1794.	C 8.1	
13340 viridiflrum H. K.	green-flowered	tt. j or	2 JL	G	C. G. H.	1774.	C 8.1		
13341 tenuiflrum Jac.	slender-flwd	tt.i or	2 JL	Pk	C. G. H.	1820.	C 8.1		
13342 nitidum Haw.	bright	tt-1 or	2 jlLo	Y	C. G. H.	1790.	C 8.1		
J3343 brachiMum H. K.	brachiated	tt.	J or	1 jn.au	Y	C. G. H.	1774.	C 8.1	
13344 subincanum Haw.	rather hoary	tt.	J or	2 au.s	W	C. G. H.	1820.	C 8.1	
13345 testaceum Haw.	tWccotored	* LJ or	3 au.s	O	C. G. H.	1714.	C 8.1	DI. el. 264	
13346 tuberbus L.	tuberous-rao-dtt.	□ or	3 jn.o	O	C. G. H.	1714.	C 8.1		
13347 m i n s 8 H a w .	smaller	tt.	1*jn.o	O	C. G. H.	1824.	C 8.1		
JSS TMA ² K>rtaum Haw.	long-rooted	tt.	2 my.jn	W	C. G. H.	1714.	C 8.1		
13348 noctiflrum J.	night-flowering	tt.	2 jn.au	W.pk	C. G. H.	1732.	C 8.1		
JSS HAW. HAW.	st>w-colored	tt.	2 jn.au	Str	C.G.H.	1714.	C 8.1		
13349 datum Haw.	elate	n. i J or	3 jn.au	S	C. G. H.	1820.	C 8.1		
13350 defoliatum Haw.	fulvous	tt.	2 jn.au	Ful	C. G. H.	1820.	C 8.1		
13351 hontale Haw.	foliated	tt.	2 jn.au	W.pk	C. G. H.	1795.	C 8.1		
13352 speciosum Haw.	horizontal-^	tt.	2 jn.au	Str	C. C. H.	1793.	C 8.1		
13353 nrctans L.	specious	tt.	1} my.o	S	C. G. H.	1704.	C 8.1		
13354 maculMum Haw.	spotted-sealed	tt.	1} my.au	S	C. G. H.	1732.	C 8.1	Bot mag. 448	
13355 flXvum Haw.	small yellow	tt.	fj ...	8	C. G. H.	1820.	C 8.1		
13356 obriquum Haw.	oblique	tt.	4 au	Y	C. G. H.	1819.	C 8.1		
iIS ⁵ ? parviflrum Haw.	small-leaved	tt. J or	1 au	P	C. G. H.	1820.	C 8.1		
{£5 ² breviflrum H. K.	short-leaved	tt. J or	1'jl.O	R	C. G. H.	1795.	C 8.1		
{JgroBubglobsum Haw.	subglobose	tt.	1 jl.o	R	C. G. H.	1792.	C 8.1		
***Q pulverulentum Haw.	powdery	tt.	my	Pk	C. G. H.	1792.	C 8.1		

13361 <i>hispidum L.</i>	hispid	JL I_ or	f	my.o	P	C. G. H.	1704.	C	8.1	Di. el. 278
2 <i>platypetalum Haw.</i>	broad-petaled	JU_ or	f	my.o	P	C. G. H.	1820.	C	8.1	
13362 <i>hirtellum Haw.</i>	dwarf bristly	JL_ or	I	my.n	Pk	C. G. H.	1792.	C	8.1	
13363 <i>candens Haw.</i>	glittering	JU_ or	i	HjLäü	P	C. G. H.	1820.	C	8.1	
13364 <i>subcompreiumum//aw.</i>	subcompressed	JU i_J or				C. G. H.	1823.	C	8.1	
2 <i>minus Haw.</i>	smaller	tt. i_J or	1	jl.au	P	C. G. H.	182J.	C	8.1	
13365 <i>flonbitindum Haw.</i>	bundle-flwd	JUu or	I	my.o	Pk	C. G. H.	1704.	C	8.1	Di. el. 214.289
13367 <i>torquatum Haw.</i>	torqued	JL_ or	\$	my.o	Pk	C. G. H.	1820.	C	8.1	
13368 <i>stratum Haw.</i>	striated-bristly	JU_ or	f	my.o	Pk	C. G. H.	1819.	C	8.1	Di. el. 281
2 pollens	pale	Isi_ or	I	my.o	w	C. G. H.	1821.	C	8.1	Dec. pl. 138
13369 <i>attenuatum Haw.</i>	attenuated	JL_ or	I	my.o	w	C. G. H.	1818.	C	8.1	
13370 <i>hispidobum Haw.</i>	brUgly-leaved	JUL-Jor	\$	my.o	w	C. G. H.	1818.	C	8.1	
2 <i>roscum Haw.</i>	rosy	1* I_J or	i	my.o	Pk	C. G. H.	1818.	C	8.1	
13371 <i>echinktum H. K.</i>	hedgehog	tt. J_jor	^	jl.o	Y	C. G. H.	1774.	C	8.1	Dec. pi. 24
2 <i>album Haw.</i>	white	tt. I_J or	i	jl.o	W	C. G. H.	1774.	C	8.1	
13372 <i>strumbinum L.</i>	strumose	tt.i_lor	^	au	Pa.Y	C. G. H.	1820.	C	6.1	
13373 <i>barluituin L.</i>	trailing bearded*	J_ or	f	jn.au	Pk	C. G. H.	1705.	C	8.1	Dec. pi. 28
13374 <i>gtelligeruin Haw.</i>	star-bearing	tt.i_lor	£	my.o	Pk	C. G. H.	1793.	C	8.1	Bot. mag. 70
13375 <i>BtellMum Dec.</i>	Starry-bearded	tt.i_jor		S.O	Pk	C. G. H.	1716.	C	8.1	Di.el.235
iiiButuii jciWi										
13376 <i>densum //aw.</i>	densc-bearded	tt. i_lor	f	my.au	Pk	C. G. H.	1732.	C	8.1	Bot mag. 1220
13377 <i>bulbosum Haw.</i>	bulbous	*Li_ or	£	au	Pk	C. G. H.	1820.	C	8.1	
13378 <i>intonsum Haw.</i>	unshaved	tL_ or	i	jl	Pk	C. G. H.	1824.	C	8.1	
2 (<i>libum Haw.</i>)	white	tt_ or	£	£	Pk	C. G. H.	1824.	C	8.1	

1521. HYMENO'GYNE *Haw.* HVMBNOGYNB. (*Hymen, membrane, gyne, woman; style,*) *Flcoideee.* 1.
13379 *gftbra Haw.* smooth idJeu f jlLo Pa.Y C.U.H. 1787. S 8.1 Bot rep. 57
Mesembryanthemum glabrum *H. K.*

CLASS XII. OASSR 8.

ICOSANDRIA POLYGYNIA.

J522. JKTSA TOIL	ROSE. (Rhod, red, Celt; color of flower.)	Roidscts.	116.—150.
I. SIMPLICIFOLIA. — Leaves ample, cxs stipulat.			
13380 ferferifblia PalL	Berberry-lvd tt _J or 1} jnjI Y Persia	1790.	L r.m Par. Ion. 101
II. FERGCEB. — Very prickly ; branches clothed wUh persistent tomentum ; fruit naked.			
13381 ferox Law.	fierce tt or 3 jn.au R	Caucasus	1796. L co Dot. reg. 420
kamch&tica Red.			
13382 kamch&tica Ven.	Kamtschatka tt or 4 jl.au R Kamtsch. 1802. L co Bot. reg. 419		
2 nUens Lindl.	shining tt or 4 jl.au R 1822. L co Bot reg. 824		
III. BBACTEA^TJE.—Branches and frvU clothed ivth persistent tomentum.			
13383 involucrata Rox.	involucrated * I or 3 jl.au W R Indies	1818. L co Bot. reg. 739	
13384 bracteata Wnl.	bractcated J or 2 au.o W China	1795. C l.p Vcn. eels 28	
2 scabricatilis Lindl.	rough-stemmed■ or S au.o W China	- C l.p Bot mag. 1377	
IV. CIKNAM6MEJE.—Setigerous or unarmed bracteate ; leaflets lanceolate, glandless; disk slender.			
13385 nitida W.	glossy ■ or 2 jn.au R N. Amer. 1807. L co Lind. ros. 2		
13386 Alpa Base	Turnip ■ or 4 jn.au R N. Amer. ... L co Red. ros. 1.7		
13387 fucida Shrh.	hfxcht-leaved ■ or 2 jn.au R N. Amer. 1724. L co Di.eL245.316		
13388 pennsylvanica Lk.	Pennsylvanian tt or 5 jnjI Pk Pennsylvan.... L r.m		
13389 gemella W.	twin-flowering tt or 3 jl.au R N. Amer. 1800. L* co		
13390 laxa Lindl.	loose Carolina tt or 3 jl.au R N. Amer. L co Lind. ros. 3		
13391 parvi flora Ehrh.	small-flowered tt p* lijn.au F N. Amer. 1724. L s.p Lawr. ros. 3		
2 flure pr-no	double-flowered tt pr 1} jn.au F N. Amer. ... L co		
13392 taurica Bieb.	Taurian tt or 6 - Pk Tauria - L co		
13393 Wo6dNi Lindl.	Woods's tt or 3 mr.jn R N. Amer. ... L co Bot.reg.97C)		
13394 caroHna L.	Carolina tt or 6 jnjI R N. Amer. 1726. L s.p Lind. ros. 4		
2 Honda Donne	Florida tt or 5 jnjI R N. Amer. ... L s.p		
13395/raxiniflolia Bore.	Ash-leaved tt or 6 my.jn R Newfound. ... L co Bot. reg. 458		
13396 cinnamomea L.	Cinnamon tt or 6 my. P Europe ... L co Eng. bot 2388		
2 flora pleno	double-flwd tt or 5 my.jn P Europe ... L co Lind. ros. 5		
3 flora somipleno	scmidouble-flwdik or 7 my.jn R Siberia 1805. L co		
13397 pygmaea Bieb.	pygmy tt or 3 jnjI Pk Caucasus 1820. L r.m Bic.ben. 1.2		
13398 ctinophylla Red.	recluuing-lvd tt or 6 my.jl W 1820. L r.m Red. ros. 1.43.ic		
13399 fruticosa Bes.	shrubby tt or 6 jnjI Pk Vallesia 1817. L r.m		
13400 ambigua Wallr.	ambiguous tt or 6 jnjI Pk Germany ... L r.m		
13401 dahurica Pall.	Dahunian tt or 6 my.jl Pk Siberia 1824. L r.m		
13402 adenop^lla W.	gland-leaved tt or 4 jnjI Pk - - L r.m		
13403 majalis Retx.	May tt pr 3 my.jn Pk Europe - - L co Fh dan. 688		
13404 macrophylla Lindl.	long. leaved tt or 6 - Nepal 1822. L co Lind. ros. 6		
13405 Lyont Ph.	Lyon's tt or 4 my.au Pk - - L r.m		
V. PIMPINELLIFOJL.—Pückles equal, numerous, bractless, sepals connivent persistent; disk almost none.			
13406 alptaa L.	alpine ■ or 3 jnjI Pk Switzerl. 1683. L r.m Bot. reg. 424		
2 pyrenhica Gou.	Pyrcean ■ or 3 jnjI Pk Pyrenees L co Gou.il. 19		
« ^ M ?^ du^na L.	pendulous ■ or 5 my.jn P Switzerl. 1726. L co Lawr. ros. 91		
13407 suavis W.	sweet ■ or 4 Pk - - 1818. L r.m		
13408 balsamea Spr.	balsam ■ or 3 jnjI R - - 1825. L r.m		
13409 muricata *.	muricated ■ or 3 Pk - - 1820. L r.m		
13410 rubiflra Stn.	reddish ■ or 2 jnjI Pk England sea sh. L co Eng. bot 2521		
L ^M 2 melanop^lla Lindl.	black-fruited tt or 2 jnjI Pk N. Amer. . . . L co		
13411 stricta Carolina tt or 3 jn Pk N. Amer. . . . L co Lind. ros. 7			

13412 aciculkris <i>Lindl.</i>	aciculär	II	or	6	my.jn	Pk	Siberia	1805.	L co
2 pauciflora <i>Lindl.</i>	few-flowered	ft	or	6	my.jn	Pk	Siberia	1813.	L co
13413 sulphurea // K.	double sulphur	8k	or	3	jl"	Y	Levant	1829.	L 81 Hot. reg. 4*
13414 lutescens <i>Ph.</i>	lutescent	8k	or	3	my.jn	Pa.Y	Siberia?	1780.	L co liind, ros i)
13415 <i>Eglantaria</i> L.	Eglantine	8k	or	4	my.jn	Y	—	—	L co Hot. mag. 3>J
lutea unicolor <i>B.M.</i>	yellowish	Se	or	4	my.jn	Y	—	—	I- co Bot. mag. 1570
2 luttola <i>Thore</i>	—	—	—	—	—	—	—	—	—
hispida R M.	—	—	—	—	—	—	—	—	—
13416* flava <i>Wik.</i>	yellow	8k	or	4	jn.jl	Y	—	—	L r.m
13417 spinosissima L.	spinest Scotch	Je	or	2	jn.jl	W.B	Britain sa.hca. L pi	F.ng. bot. 197	—

Garden Varieties.

Blush, Anderson's Double	Marbled, Double Crimson	Two-colored, Small Double
Lady's	Marbled, Double Dark	White, Large Double
Blush, Double Lady's	Marbled, Double Light	White, Large Semi-double
Blush, Double Pink	Purple, Double	White, Small Double
Blush, Double Provinc	Purple, Small Double Light	White, Whitley's Double
Blush, Double Rose	Red, Double Dark	Yellow, Globe Double
Blush, Dutch Double	Red, Double Light	Yellow, Large Double
Blush, Princess Double	Red, True Double	Yellow, Pale Double
Crimson, Double	TVo-colorct, Large Double	Yellow, Small Double

2 reversa <i>Lindl.</i>	reversed	8k	or	1	my.jn	W	Siberia	1814.	L co Hot. reg. 451
3 saniculifolia <i>Lindl.</i>	Buract-lvdf	or	3	my.jn	W	—	—	L co	—
13418 grandi flora <i>Lindl.</i>	large-flowered	8k	or	4	my.jn	W	Siberia	1818.	L co Bot. reg. 888
13419 myriacantha Dec.	myriath-spined	ii	or	1	my.jn	W	S. France	1820.	L co Lind. ros. 10
13420 Bieberstefni <i>Lindl.</i>	Bieberstein's	8k	or	2	my.jn	W	Caucasus	1822.	L co
ferox <i>Bkb.</i>	—	—	—	—	—	—	—	—	—
13421 involuta Sm.	involted	8k	or	2	h.II	W	Hebrides	moun.	L co Eng. bot 2068
13422 revrsia W. & K.	reversed	■	or	5	iti-il	W.Y	Hungary	1816.	L co W. Ac K. 264
13423 alftica Red.	Altaiian	■	or	3	jii.ll	W	Altaiia	1818.	L r in Pal. ros. 75
spinoblossima y PalUiff <i>Lindl.</i>	—	—	—	—	—	—	—	—	—
13434 melanocarpa Lk.	dark-fruited	■	or	■	jii.jl	W	S. Europe	1817.	L r.m
13425 oxyacanthä <i>Bieb.</i>	sharp-spined	■	or	■	jn.jl	W	Caucasus	1820.	L r.m
13426 Sabmi Woods	Sabine's	■	or	8	my.jn	W.R	Britain woods.	L co	—
13427 Doniuma Woods	Don's	■	or	4	my.jn	Pk	Britain hetl	L co	—

VI. CE.VTIFOLIJR — Prickles unequal, braceate ; disk thick, surrounding the throats leaflets rugose.

IS428 damascena Mil.	Damascus	A	or	3	jn.jl	Pk	Levant	1573.	L co Lawr. ros. 38
----------------------	----------	---	----	---	-------	----	--------	-------	--------------------

Garden Varieties.

Agathe, Rouge	Brunswick	Hundred-leaved, Petite	Quatre Saisons, flesh-co-
Argentea	Cluster, Pale	Incomparable	lored
Auguste, Belle	Couronne, Belle	Mignonne, Favorite	Quatre Saisons, Francois
Aurora	Couronne, Petite	Monarque, Grande	Quatre Saisons Panachd.
Belgic, Blush	Damas Argenté	Monthly, Red	Quatre Saisons Pomponc
Belgique Carnée	Damas Pourpré	Monthly, White	Quatre Saisons sans spines
Belgique Violette	Damask, Hlsh	Pajstana	Quatre Saisons, semb
Bifra Carnea	Damask, lied	Paragon	double
Bifra de Naples	Damask, White	Parnassus	Royal, Great
Bifera grandiflora	Egyptian	Perpetual	Swiss
Blush, Early	Emirvor	Prolific	Valiant
Blush, Imperial	Felicite	Quatre Saisons	Versailles
Blush, Monthly	Goliath	Quatre Saisons blanche	York and Lancaster
Blush, Watson's	Gracieuse	—	Zealand

13429 centifolia L.	hundred.lvd Provins	■	or	3	jn.au	Pk	S. Europe	1596.	L r.m lied. ros. 1.1
provincialis Mil.	—	—	—	—	—	—	—	—	—

Garden Varieties.

Aunay, Belle d'	Constance	Petite Holland	Provins, Royal
Aurora	Cramois, Grand	Persian	Provins, Scarlet
Belgic, Red	Cumberland	Pompon, Gros	Provins, Semidouble
Blandford, or Kingston	Dragon	Pourpré Aimable	Provins, Shaillers
Blush, Royal	Duchesse d'Angouleme	Pourpré Favorite	Provins, Single
Bourbon	Duchesse de Berri	Pourpré Violette	Provins, White
Bright Crumpled	Elysian	Prolific	Rouge Superb
Cabbage, Blush	Emperor	Provins, Blush	Sans ptillea
Cabbage, Single	Juno	Provins, Cabbage	Souchet
Carmine	Louis XVIII.	Provins, Childings	Spongs
Carmine, Superb	Malta	Provins, Common	Striped Nosegay
Centfeuilles Anemone	Mfrc Gryone	Provins, Damask	Surpassantc
Centfeuilles de Bruxclles	Mottled Puqile	Provins, Dutch	Syren
Centfuillles de Hesse	Nea>olitau	Provins, Early	Trianon, Belle de
Centfeuilles Gaufrœ	(Eillet	Provins, Grand	Versailles
Chamois	One-sided	Provins, Imperial	Vilmoriu
Cluster	Pencilled	Provins, Invinciblo	—
2 muscbsa Mil.	moss	flft or 3 jn.jl Pk	— L r.m Red. ros. 1.8

Garden Varieties.

Moss, Blush	Moss, Dark	Moss, Single	Moss, White
Moss, Common	Moss, Prolific	Moss, Striped	—
3 pompnia Dec.	Pompone	■ or 2 jn.jl Pk	S. Europe L r.in Red. ros. 1.21

Garden Varieties.

De Meaux	Mignonne Charmante	Provins, Dwarf	Rheims, Dc
Dwarf Bagshot	Pompone	Provins, Small	St. Francis
Mosby de Meaux	. Pompone, Proliferous	—	—

14a 4bipinntaiJtafc	bipinnate ft	or 3 jn.jl R	— L co Red. ros. £. 4
13430 gtlila L.	French officinal St	or 2 Jnjl P	S. Europe 1596. L co Bot. reg. 4K

Garden Varieties.

Admirable	Fanny Bias	Mignonne	Prince
Aigle noir	Fiery	Mignonne, Blush	Prince William V.
Albanian	Flanders	Mignonne, Dark	Princess
Amaranth	Flemish	Mignonne, Favorite	Prolific
Antwerp	Formidable	Mignonne, Red	Promville
Atlas	Fringed	Mignonne, Semidouble	Proserpine
Belle Aurora	Garnet	Mignonne, Striped	Provins Pulmonaire
Beauté Aimable	Gay	Mirabelle	Purple, Blue
Beauté Rouge	Giant	Mogul	Purple, Bright
Beauté Supreme	Gloria Mundi	Montauban	Purple, Favorite
Bijou	Granaatappel	Morocco	Purple, Grand
Bishop	Grand Monarque	Mottled, Black	Purple, Light
Black Frizzled	Grand Sultan	Natalie	Purple, Royal
Blue	Henry IV.	Negrette	Pyramid
Bouquet Rouge Royal	Hermelin, Belle	Negro	Queen
Brunette	Hervi	Ninon de l'Enclos	Ranunculus
Brussels	Hollandc, Noir dc	Nonpareil	Ranunculus, Early
Buonaparte	Hundred-leaved, Blush	Nonsuch	Red and Violet
Burning Coal	Hundred-leaved, Dutch	Normandy	Roi de France
Cardinal	Hundred-leaved, Single-ton's	Officinal	Rose de Parade
Carmine	Imperatrice	Officinal, Blush	Royal Red
Carmine Brillante	* Incomparable	Officinal, Carmine	Royal Virgin
Carmine, Proliferous	Infernal	Orleans	Sable
Carnation	^ Invincible	Ornement de Parade	St. John's
Catalouian	Italian	Passtana	Sanspareil
Champion	Josephine	Panaché, Petite	Sceptre
Chancellor	Junon	Paradise	Shell
Changeable	King	Paragon	Spanish
Cherry	La Dauphine	Pavot	Stadholder
Clementine	L'Ombré Agréable	Perruque	Stepney
Coquette	L'Ombre Superbe	Phoenix	Striped Nosegay
Coulcur de Feu	Leyden	Plicate	Pluto
Cramoisié, Belle	Lisbon	Pomona	Sultana
Cramoisié, Grande	Lively	Pompadour	Superb Red *
Crimson, Dutch	Lurid	Pompon Bazard	Trafalgar
Crimson, Purple	Maiden	Poniutowsky	Triumphant
Crimson, Royal	Majorca	Poppy	Tuscany
Crown	Malabar	Porcelaine	Two-colored'
Cupid	Malta	Portland	Velvet, Double
Damask, Black	Mantau Royal	Pourprc, Belle	Velvet, Semidouble
Delicious	MarUed	Pourpr Bouquet	Velvet, Single
Dingy	Marbled, Dark	Pourpre Channante	Velvet, Striped
Due de Guiche	Marbled, Double	Pourpre de Tyr	Venetran
Duchesse d'Orléans	Marbled, Grand	Pourprc, Grande Belle	Victory
Dwarf Proliferous	Margaret	Pounce, Point	Violet, Dark
Enchanter	Matchless	Pourpres, Roi des	Violette, Belle
Enfant de France	Mauve	Pounce, Velours	Violette et Rouge
Eucharis			

13431 2 pumila L. dwarf wild officinal A or J R Austria 1810. L co Jac.au. 198
 marginata Wallr. margined Sk or 5 Pk Germany 1818. L r.m
 13432 pulchella Spr. neat Sk or 2 II France 1824. L r.m
 13433 parvitulifolia A. uruAUvd Burgundy ik or 1 P Europe -- L r.m Bot. reg. 452

VIT. VILLEOSAS.—Prickles straight; sepals conniv.persist, j disk thick, surround, throats branches upright
 13434 turbiflataf. K. turhmatc Frankfort Sk or 5 jn.au Pk 1629. L r.m Lawr. ros. 69
 lorbeissnica double Orbessandl or 4 inau Pk L r.m
 13435 coruscans Lk. glittering A or 6 jn.au Pk L r.m
 13436 gracilis Woods slender flk or 8 jnjl R Britain highl.v. L r.m Eng. bot. 583
 13437 vilKsa L villous ■ or 6 jnjl Pk Britain hedges. L co Eng. bot. 2459
 mollis Sm.
 13438 heterophylla Woods various-leaved ■ or 6 jnjl Pk Britain hedges L co
 13(39) scabnuscula E. B. rather rougher ■ or 6 jnjl Pk Britain hedges. L co Eng. bot. 1896
 13440 tomentosa Sm. ilownly-lvd Dog Sk or 6 jnjl Pk England hed. L co Eng. bot. 990
 2 resiliens Lindl. resinous Sk or 4 jnjl R Ireland . . . L co
 15441 cuspidata Lieb. cuspidate ■ or 6 jnjl Pk Tauria 1817. L r.m
 13442 montana Stev. mountain ■ or 6 jnjl R Tauria 1821. L r.m
 13443 molis Led. soft ■ or 6 jnjl R Caucasus 1818. L r.m
 13444 Wilidenfivii Spr. Willdenow's ■ or 4 jnjl W Siberia 1820. L r.m
 13445 alba L. single white ■ or 4 jnjl W Crimea 1597. L r.m Lawr.ros.37

Garden Varieties.

Agate	Due d'Yorck	Maiden's Blush, Great	Rbsea
Belle Aurorc	Eliza	Maiden's Blush, Small	Simonneville
Blanche à Coeur Vert	Feuille Fermée	Moraga la Favorite	Spineless Virgin
Blanche de Belgique	Grande Cuisse de Nymphe	Muscat Rouge	Thorniest*, Double
Blush, Double White	Hennette, Belle	Nova Cœfestis	TnangulVis
Bouquet Blanc	Joanne d'Arc	Nbva Plena	White, Double
Celestial	Maiden's Blush, Cluster	Petite Cuisse de Nymphe	White, Semidouble
13446 hibernica Sm.	Irish HI	or 2 jn.n Pk	Ireland ir.thi. Skco Eng.bot.S196
V111. RuBiGiNbsJE.—Prickles unequal; sepals persistent; disk thick ; branches arched ; leaflets glandular.			
13447 l&tca Mil.	single yellow tt	or 3 jn Y	Germany 1596. L r.m Bot mag. 363
2 punlca Mil.	scarlet Austrian A	or 3 jn Y.o	Germany 1596. L r.m Bot mag. 1077
13448 rubiginosa L.	rusty Sweet Briar it	or 5 my.jn Pk	Britain ch.ba. S co Eng. bot 991

Garden Varieties.

American, Single	Double	Monstrous	Scarlet
Blush	Dwarf, Semidouble	Mossy	Tree, Double
Clementine	Maiden	Petite Hessoise	White, Semidouble
Cluster	Mannines	Royal	Zabeth

2 <i>micrantha</i> <i>Sm.</i>	•mall-flowered flk	or	(5	my.jl	Pk	Britain	thick.	L	co	Eng.bot.S499	
3 <i>umbellata</i> <i>Iscrs</i>	umb. <i>Sweet Briar</i> <i>Sk</i>	or	4	my.jn	Pk	Germany	...	L	r.m	Lawr.ros.G-i	
4 <i>sepium</i> <i>Thuil.</i>	heilge	St	or	3	my.jn	Pk	Britain	thick.	L	co	
5 <i>inolitra</i> <i>As.</i>	scentless	3k	■	6	my.jn	Pk	Britain	hcd.	L	co	Cur. Ion. 117
13449 <i>Bdrynia</i> <i>Woods</i>	Horrors	3k	■	6	mrjl	Pk	Britain	hed.	L	co	Eng. bot. 2579
13450 <i>suavolens</i> <i>Rofi.</i>	sweet-smelling	3k	■	6	my.jl	Pk	N. Amer.	1800.	L	r.m	
13451 <i>pulverulenta</i> <i>Bicb.</i>	Caucas.	dusty	■	6	my.jl	Pk	Caucasus	1817.	L	r.m	
13452 <i>iberica</i> <i>Stev.</i>	Iberian	3k	■	6	my.jl	Pk	Iberia	18m	L	r.m	
13453 <i>floribunda</i> <i>Stcv.</i>	bundle-flwd	3k	■	6	my.jl	W	Tauria	1819.	L	r.m	
rubiginbsa <i>Bleb.</i>			■	6	my.jl						
13454 <i>pruniosa</i> <i>Lindl.</i>	frosted	3k	or	3	my.jn	Pk	Siberia	1818.	L	co	
13455 <i>glutinosa</i> <i>Sm.</i>	glutinous	<i>Cretan Sk</i>	or	2	my.jn	Pk	Candia	1821.	L	co	Red. ros. 1.47

IX. CANINJS.—Prickles equal, hooked; lvs glandless; sep. decid.; disk surround, throat j branches arched.

13456	sericea Lindl.	silky	3k	or	6	my.jn	Pk	Nepal	1822.	L.p.l	Lind. ros J2
13457	caucasea Lindl.	Caucasian	3k	or	20	jnj.l	K	Caucasus	1798.	L co	
13458	can't na L.	dog, or <i>Hip</i>	3k	or	8	jnj.l	Pk	Britain	hed.	L co	Eng. bot. 992
2 coll'ma, Jae.	hill	3k	or	8	jnj.l	Pk	Britain	hed.	L co	Eng. bot. 1895	
3 dumetorum Tftui/.	thicket	3k	or	8	jnj.l	Pk	England	hed.	L co		
4 cassia E. B.	grey	3k	or	6	jnj.l	Pk.w	Scotland	hed.	L co	Eng. btt. 2367	
13459	saxatilis Stev.	rock	3k	or	5	jnj.l	Pk	Tauria	1820.	L rm	
13460	baltica Roth	Baltic	3k	or	5	jnj.l	Pk	Baltic	[]	L r.m	
13461	leucodutha Bleb.	white-flowered	3k	or	10	jnj.l	W	Iberia	[]	L rm	Lind. ros. 11
13462	nuda ffow«	naked	3k	or	5	jnj.l	Pk	Britain	hedges.	L co	
13463	Sherardf Dav.	Sherard's	3k	or	6	jnj.l	Pk	Britain	-hedges.	L co	
13464	sarmientacea Sivz.	twiggy	-*	or	6	jnj.l	Pk	Britain,	hedges.	L co	Cur. Ion. 5.34
13465	br.ictescens Woods	bractescent	3k	or	6	jnj.l	Pk	Britain	hedges.	L co	
13466	Förmer* Sm.	Forater's	*	or	6	jnj.l	Pk	Britain	hedges.	L co	
13467	surculosa Woods	sponggy	**	or	4	jnj.l	Pk	Britain	hedges.	L co	
13468	rufribilia VU.	red-leaved	Hk	or	6	jnj.l	P	Europe	1814.	L co	Bot. rcg. 430
2 Uedutert Thory	Rertoute's	3k	or	3	jnj.l	P	1822.	L ro	Red. ros. 1. 38	
13469	Montezbne H.&B.	Montezuma's	[]	3	jnj.l	Pk	Mexico	1825.	C r.m	Red. ros. 1. 16	
13470	indica L.	blush	Indian	[]	20	year	F	China	1789.	C p.l	Lawr. ros. 2ti

Garden Varieties referable either to R. Indica or R. semperfibrans.

Xiba	Centifolia	Lhida	Purpurca		
Animating	Cerise éclatante	Major	Sanguinea		
Aéro-rilgra	Chiffonnée	Minor	Sans Epines		
Bengale à Bouquet	Cucullata	Monströsa	Subalba		
Bengale à Fl. Panaché	Elegans	Moonshine	Terneaux		
Bengale Blanche	Hirula	Nigra	Thisbe		
Bichitrina	Gigant ^a	Ochroleuca	Veloutec		
Carnescens	Lie de Vin				
2 odoratissima Swt.	sweetest-scented	i* — I or 3	f.au Pa.Pk China	1810. C p.l	Bot. reg. 804
S p&mild Jterf.	dwarf	» — J or 1	my.au Pk China	— C p.l	Red. ros. 1.42
4 Fongilpha W.	long> F'/low-lyd«	5 my.au	China	— C p.l	Red. ros. 2.12
5 Noisettidna	Noisette's	• — I or 10	my.s hybrid	— C p.l	Red. ros. 2.77. ic
13471 RoxWirgh Hort.	Roxburgh's	fl — I or 13	my.s China	1821. C p.l	
13472 Ayacanthina Hort	Hyacinth-scented	fl — I or 13	my.s hybrid	1821. C p.l	
13473 Boursotiltii Hort.	Boursoult's	fl — I or *13	my.s hybrid	1821. C p.l	
1J474 Fra-seridmi Hort.	Fraser's	m — I or 4	my.s hybrid	— C p.l	
1S475 purpurea Hort.	purple	* — I or 4	my.s hybrid	— C p.l	
13476 flavescens Hort.	flavescens China	* — I or 3	my.s Crea China	1821. C p.l	
13477 Liingii Swt	Lady Long's	ft — I or 20	jnl R hybrid	— L	
13478 nivea Dec.	snowy	T — I or 20	jl Pa China	1820. C Bp	
13479 semperfliuens Cur.	ever-blowing	» — I or 10	year C China	1789. C p.l	Bot. mag. 284
13480 Lawrencetia Swt.	Miss Lawrence's*	— I or 1	year R China	1810. C p.l	Bot. mag. 1762
13481 microphylla Box	small-leaved	• — I or 3	— Pk E. Indies	1823. C p.l	Bot. reg. 919

X. SYSTYLIE.—*Styles joined into an elongated column : stipules adnate.*

X. STYLISED.—Styles joined into an elongated column; stipules annual.									
13482	systylea <i>Desf.</i>	clustered-styled dk	6	my.jl	Britain	hed.	L	co	Eng. bot 1895
2	Monsonia <i>Lindl.</i>	Lady Monson's ik	3	my.jl	Britain	hed.	L	co	Red. ros. 3.67. ic
13483	arvensis <i>Hud.</i>	corn white dog	8	jnl	Britain	hcd.	L	co	Eng. bot 188
13484	h^brida <i>Schl.</i>	hybrid	4	my.jn	Switzercl.	—	L	co	
13485	semperfiriens <i>L.</i>	evergreen	fL	—	S. Europe	1629.	L	co	Bot reg. 465
	2 cnpcolkta <i>D. Don</i>	Vfjnvn%Ayrsrh.	—	—		—	—	1818.	L co
1348K	geminata <i>Rau.</i>	twin	3k	—	Switzercl.	1819.	L	r.m	
13487	Itussolima <i>Hort.</i>	Russel's cottage X	—	—		—	L	p.1	
13488	multiflora <i>Thun.</i>	many-flowered X	—	—	China	1804.	C	s.l	
	2 multiplex	double	—	—	China	1804.	C	s.p	Bot. mag. 1059
	3 alba	white	—	—	China	1810.	C	p.1	
13489	Grevillea <i>Hort.</i>	Greville's	—	—	China	—	C	p.1	
13490	Brununii <i>Lindl.</i>	Brown's	—	—	Nepal	1822.	C	co	Lind. ros. 14
13491	mochbata <i>Mil.</i>	musk	—	—	Barbary	1596.	L	r.m	Lawr. ros. 64
	2 tibre plj-no	double-flwd	—	—	Barbary	1596.	L	r.in	Lawr. ros. 53
	3 ncipalensis <i>Lindl.</i>	Nepal	—	—	Nepal	1822.	L	co	Bot reg. 829
	4 arburva <i>Pers.</i>	tree	—	—	Persia	1824.	L	co	
	5 nivea <i>Lindl.</i>	snow bush	—	—		—	L	co	
	evrgiff' atooBC	Ev ratine	—	—		—	—	1822.	L co
13492	rubifolia <i>R. Br.</i>	Bramble-lvd	—	—	N. Amer.	1800.	L	p.1	Bot reg. 861
	2 fenestrata <i>Lindl.</i>	windowed	—	—	N. Amer.	1800.	L	p.l	

XI. BANKSIA^NIE. — *Stipules narrow, nearly free, often deciduous; leaves often ternate; stems scaudent.*

13493	<i>sinica</i> Ait.	—	<i>Spathe</i> narrow, nearly free, often decurrent; leaves often pinnate terminalia stems scandent.
13494	<i>Banks/a?</i> R. Br.	three-lvd <i>China</i> £:	or 5 my. Jl W China 1759. L p. Lind. ros. 16
		Lady Banks's fl	or 20 jnjl W China 1807. L p. Bot reg. 397
	2 flwr luteo	yellow-flowered fl	or. Y China 1824. L p.
	3 multiplex		E. Indies 1824. L p.
13195	<i>microcarpa</i> Thun.	small-fruited	J or 20 jnjl Y E. Indies 1824. L p. Bot. reg. 1105
			W China 1822. L p. Lind. ros. 18

*lo23. .fllTBUS L.

BRAMBLE, E.

{*Rub*, red, *Celtic*; fruit)

Rosdccc*. 63.—116.

I. PiNNArirtuL — *Lvs pinnate, underneath smooth, from No. 13500. undern. clothed with white tomentum.*

13406	<i>rosneftulosa</i> 5m. 2 coronkriui	Rose-leaved garland	Jor ■LJor	3 ap.o 3 ap.o	W W	Mauritius 1811. Mauritius 1811.	X pl	Sra. ic. pic. 3.60 Bot. mag. 1783
-------	---	------------------------	--------------	------------------	--------	------------------------------------	---------	--------------------------------------

13497 <i>Eglantria</i> Trat	Eglantine	il L J or	3 ap.jl	W	N. Holl.	1835.	C p.l
13498 <i>pinnatus</i> IV.	pinneate	• I or	5 jn.jl	Pk	Madeira	1789.	C p.l
13499 <i>suberectus</i> Ander.	sub-upright	Sk or	4 jn.s	W	Britain woods.	Sk co	Eng. bot. 2572
ncmorusus Hay.							
1 J500 <i>micanthus</i> D.	Don email-flowered	Sk	fr 10	my.au	K Nopal	1822.	C co
paucitiorus Lindl.	distant	■ or	5 jl	W	Nepal	1818.	L co
13501 <i>dibitanus</i> D. Don	petalless	■ flor	6 jl.au	■	Is. France	1823.	C p.l
13502 <i>apitalus</i> Pair.	occidental	■ fr	5 my.jn	W	N. Amer.	1696.	Skco
13503 <i>occidentalis</i> Mis /.	rough	■ or	5 my.jn	W	Nepal	1821.	I pi
13504 <i>asperX</i> . Don	Raspberry	■ fr	5 my.jn	W	Britain m.wo.	Sk r.m	Kng bot. 2442
13505 <i>Idaeus</i> L.							
	II. PALMATIFLUII.	Leaves palmate, 3—5-foiate.					
13506 <i>laciniatus</i> W. en.	j-xg-leaved	Jk or	12 jn.s	W n			L co
ii'gjt tnvialis Ph.	trivial	Jk or	7 jn.jl	W	N. Amer.	1789.	Skco
13508 <i>caeViens</i> L.	grey Dewberry	Jk fr	2 jn.jl	W	Britain bor. fi.	Skco	Eng. bot. 826
13509 <i>foliolium</i> D. Don	small loathy	■ or	6 jn.jl	W	Nepal	1818.	L co
13510 <i>pedunculosus</i> Ztov/	pedunculated	■ or	5 jn.jl	W	Nepal	1818.	L co
13511 <i>flagellaris</i> IV. en.	rod	■ or	6 jn.jl	W	N. Amer.	1789.	Skco
13512 <i>inermis</i> IV. en.	unarmetl	Jk or	12 jn.jl	W	N. Amer.	1805.	Sk co
13513 <i>corylifolius</i> E. B.	Hazel-leaved	Jk or	10 jl	W	Britain lied.	Sk co	Eng. bot. 827
13514 <i>agrestis</i> IV. & K.	field	Jk or	6 jn.jl	W	Hungary	1820.	L co
13515 <i>plicatus</i> Weihe	folded	■ or	10 jn.s	W	Iaintain hcd.	li co	W. & C. 3.2fis
13516 <i>aftinis</i> Weihe	similar	Jk or	8 jn.s	W	Britain hedges.	L co	Necs r. g 3 •
13517 <i>Klimblius</i> Schoit	Elm-leaved	Jk or	10 jn.s	W	Spain	1823.	L co
13518 <i>liukiuensis</i> Ser.	Link's	■ or	10 jn.jl	W	Europe	1821.	I. co
paniculatus Schle.							
13519 <i>fruticosus</i> L.	common olirubby-*	or	10 jn.s	Pk	Bril.-in	hcd.	L co
2 illus	llite-fruited	■ or	10 jn.s	W	Britain	hcd.	L co
3 pltuus	double-flauK'redJk	or	6 jn.s	W	liritum	hcd.	L co
13520 <i>tomentifluis</i> Weihe	woolly-leared	■ or	10 jn.s	Pk	Germany	1818	L co
13521 <i>glandulibus</i> JV. en.	glandular	■ or	10 jn.s	W	Germany	1810.	L co
13522 <i>hirtus</i> W.QK.	hairy	Jk or	10 jn.s	W	Hungary	1816.	L vo
13523 b&nctus Schrcb.	holy	■ or	8 jn.jl	W	Palestine	1823.	L co
13524 canescens Dec.	canescent	Jc or	6 jn.s	W	Alp. Eur.	1820.	K co
13525 <i>fsistigiius</i> Weihe	fastigate	■ or	8 jn.s	W	Switzcr.	1819.	L co
13526 /iajiflMms Weihe	Lime-tree-lvd	Jk or	6 jn.jl	W	Germany	1819.	L co
13527 cordiflous HVMe	heart-leaved	■ or	8 jn.jl	W	Germany	1816.	L co
11528 rubraefolis Weihe	red-slemimud	Jk or	6 jn.jl	W	Germany	1818.	L co
13529 Menkii Weir.c	MenkV.	■ or	8 ju.jl	W	Germany	1816.	L co
13530 BeUeditandUilli	j gehtochlcii.tiaM^	or	10 jn.jl	W	Europe	1823.	L co
13531 Schleicheri Weihe	Schleicher's	Jk or	8 jl.jl	W	Germany	1818	L co
11532 horridus Weihe	horrid	■ or	6 jn.jl	W	Germany	1817.	L co
13533 Icu66tachys Sehl.	white-i-piked	Jk or	10 jn.jl	W	Britain hedges.	L co	SI. jam. 2.213.1
13534 rhammiflous Weihe	Buckthoru-lvd	■ or	10 jn.s	W	Britain	hed.	L co
1 i535 nitidus Weihe	shining	fa or	3 jn.s	W	Britain	thick.	L co
13536 argutus Lk.	&hi>-toothli-d Jk	or	3 jn.jl	W	N. Amer.	1823.	li co
13537 cuneiflous Ph.	wedge, leaved	■ or	3 jn.jl	W	N. Amer.	1811.	rflco
13538 sanguinolentus Lk.	blood-mf	■ i or	4 10 jn.s	Pk	I. France	1824.	C co
13539 Sprengelii* Weihe	Sprengel's	■ or	10 jn.s	Pk	Germany	1823.	L co
villusus/3 vulpinus							
Ser.	shaggy	or	3 jl.au	W	N. Amer.	1777.	Skco
13540 villusus J.K.	stngose	si or	3 jn.jl	W	N. Amer.	...	Skco
13541 strigulus Ph.	Jamaica	si lor	6 .	W	Jamaica	1822.	C co
13542 jamaicensis L.	bribly	2 .	3 au	W	Canada	1768.	Skco
13543 hspidus L	woolly	or	10 jn.jl	W	Siberia	1820.	U CO
13344 lanugnius Stev.	four-leaved	Jk or	10 jn.jl	W	N. Amer.	1819.	L co
13545 tetraphyllus H. K.	small-leaved	■ or	2 a.us	Pk	China	1818.	I# co
13546 parvifl <i>l</i> is I.*	Canadian	Jk or	3 jn.jl	W	N. Amer.	1811.	Skco
13547 canadiflous L.	stone	Jk A or	* i	W	Britain m.wo.	Sk p.l	Eng. bot. 2233
J3548 saxatilis /.	three-flowered	« A or	i in	W	Canada	-	Skpl
13549 triflorus Rick.	arctic	A fr	my.au	W	Scotland	aL ro.	Skpl
13550 arcticus L	close-styled	^ A or	i jn.jl	H	Labrador	1802.	Ex. bot. 2. 86
13551 pistillatus Sm.							
acatlis Mx.							
13552 sgopodioides Ser.							
19553 odorktus L.							
13554 reflexus Ker	sweet-scented	A or	7 jn.jl	R	N. Amer.	1700.	Skco
13555 moluccanus L.	reftxed	JL J or	3 jl.au	Il	China	1817.	Skco
13556 rugwtus Sfn.	Moluccas	JU. i or	3 jl.au	It	E. Indies	1810.	Skl.p
13557 stellatus Sm.	wrinkly	A y_Jor	3 jn.jl	R	S. Amer.	1819.	C co
13558 Chamaem&rus L.	starred	^ A or	i my.jn	W	N. Amer.	1824.	Sm. ic pin. 3,64
							Eng. bot. 716
1524. DALIBA'R'Dif L	Cloud herry.	^ ^ dr					
13559 iroholdes Mx.	Violet-like	A cu J my jn	W	N. Amer.	17G8.	D l.p Mic. am. 1. 27	
repens L.							
1526. COMARO'PSIS Rich.	COMAROFISM.	(<i>Comarum</i>) marsh cinquefoil, <i>opsis</i> , appearance.)					
13560 <i>Doniana</i> Dec.	Don's	f A or	i my.jn	Y	N. Amer.	1800.	D l.p Bot mag. 1567
Dalibarda fragario'ides B. M.	Waldstefni'a	<i>Uoniina</i> Trat					
13561 fragario'ides Dec.	Strawberry-Ik	J A cu i my.jn	W	N. Amer.	1803.	D 1 p Mic. am. 1. 28	
Dahbafrda fragarioides Mx.							
152R FRAGANRIA Toff.	STRAWBERRY.	(<i>Fragrans</i> , fragrant; <i>perfumed fruit.</i>)					
13562 vescia L.	edible	A fr	1 ap.my	W	Britain woodb.	Rs.s.l.	Eng. bot 1524
13561 caly^na Ijoi.	grandiflra	A fr	1 ap.jn	W	France	... Rs r.l	
grandiflra Tliuil.							
13564 monophlla W~	one-leaved	£ A fr	1 my.jn	W	1773.	Rss.l
13565 olfilla Ehrh.	hill Green Pine*	A fr	1 ap.n	W	Germany	1768.	Rs r.l
J?S VSPVP**Buch.	MajaufedChamp.	A fr	1 ap.my	W	France	...	Rs r.l
J^S B^ flin? w luch.	Bresling	^ A fr	1 ap.jn	W	France	...	Rs r.l
1?5S K^ vnoi^? &r~	Plane-like	^ A fr	M ap.my	W	N. Amer.	...	Noi. jar. 13. 2
11569 elatjor;/?rA.	taller	^ A fr	M ap.my	W	Britain woods.	R< r.l	E.g. bot. 2197
11570 ennaiUnsis MM.	Canada	*			N. Amer.	...	Rs r.l

13571 <i>virginiana</i> <i>Mil.</i>	scar/c/Virginia	£ / ^ fr	1 ap.my	W	N. Amer.	■ &	Rsr.l	Noi. jar.	12.2
13572 <i>grandiflora</i> <i>Ehrh.</i>	Pine great-flwd	£ A fr	1 ap.my	W	Surinam	1759.	Rsr.l	Mil. ic2.1SS	
J357J <i>ciliolata</i> <i>Ehrh.</i>	Chile	£ A f ^r	j myjn	W	S. Amrr.	1727.	Rs r.l	Noi. jar.	1% 1
13574 <i>bonariensis</i> <i>J.</i>	Buenos Ayres	je A ft ^r	2 apjn	Ap	B. Ayres	...	Rsr.l		
13575 <i>indica</i> <i>Andr.</i>	yellow Indian	J^ A or	1 my.o	Y	India	1805.	Rs s.p	Bot. reg.	61

1527. CO'MARUM *L.* COMA RUM. (*Komaron*, the arbutus; similarity of fruit) *Boisicar.* 1.
13 JG paKistre *L.* marsh *Cinquefoil* & cu Uin.il P Britain sp.bo. D p Eng.bot.172
Potentilla GSmarum Sco.

*1528. POTENTILLA *L.* CINQUBFOIL. (*Potens*, powerful; supposed med. qual.) *Roshecce.* 102.—126.

I. PORENTILLA'srauM. —	Petals yellow, rarely purple ; haves palmate, from No. 13G40. pinnate.								
13577 <i>fruticosa</i> <i>L.</i>	shrubby	ft	■ ■ ■ ■ ■	■ ■ ■ ■ ■	England	■ ■ ■ ■ ■	Eng. bot.	88	
13578 <i>floribunda</i> <i>Ph.</i>	bundle-flowered	ft	■ ■ ■ ■ ■	■ ■ ■ ■ ■	N. Amer.	1811.	if co	Den. br.	70
I-79 <i>nivea</i> <i>L.</i>	snowy	^	■ ■ ■ ■ ■	■ ■ ■ ■ ■	Siberia	1816.	D co	Bot. cab.	460
13580 <i>angustifolia</i> <i>ScAteC.</i>	many-leaved	5	■ ■ ■ ■ ■	■ ■ ■ ■ ■	Siberia	1824.	D co	Leh. pot	19
<i>leucophylla</i> <i>Fis.</i>									
13581 <i>grandiflora</i> /«.	great-flowered	fc	■ ■ ■ ■ ■	■ ■ ■ ■ ■	Siberia	1640.	D co	Bot. mag.	75
13582 <i>uniflora</i> L. ^f	one-flowered	■ ■ ■ ■ ■	■ ■ ■ ■ ■	£ my.jn	Dahuria	1819.	D co	Leh. pot	18
13583 <i>macrantha</i> <i>Lerf.</i>	long-flowered	■ ■ ■ ■ ■	■ ■ ■ ■ ■	i my.jn	Siberia	1820	I) co		
13585 <i>minima</i> <i>Half.</i>	smallest	■ ■ ■ ■ ■	■ ■ ■ ■ ■	i my.jn	Switzerl.	1818.	D co	Bot. cab.	480
13586 <i>glacialis</i> <i>Half.</i>	icy	5	■ ■ ■ ■ ■	■ ■ ■ ■ ■	Switzerl.	1811).	D co	Sor hel.	1.7
13587 <i>frigida</i> fit	frigid	i	■ ■ ■ ■ ■	■ ■ ■ ■ ■	Alp. Ddp	1819.	D co	Nes. pot.	10.3
13588 <i>velutina</i> <i>Leh.</i>	velvety	■ ■ ■ ■ ■	■ ■ ■ ■ ■	apjn	France	1819.	D co		
13588 <i>villosa</i> ■■■■■	villous	■ ■ ■ ■ ■	■ ■ ■ ■ ■	apjn	N. Amer.	1820.	D co	Leh. pot	16
13589 <i>speciosa</i> ■■■■■	Strawberry-like	■ ■ ■ ■ ■	■ ■ ■ ■ ■	apjn	S. Europe	1800.	I) co	Leh. pot	15
13591 <i>bithynica</i> <i>Horn.</i>	showy	<3t	■ ■ ■ ■ ■	apjn	Crete	tf.'l.	I) co	No*, pot.	11
13592 <i>decliniata</i> <i>Horn.</i>	dcclinatc-Aran	■ ■ ■ ■ ■	■ ■ ■ ■ ■	apjn	Bithynia	1817.	D co		
• 13593 <i>norvegica</i> /..	Norwegian	■ ■ ■ ■ ■	■ ■ ■ ■ ■	apjn	1817.	D co		
13594 <i>monspeliensis</i> <i>L.</i>	Montpelier	fc	■ ■ ■ ■ ■	apjn	N. Europe	1764.	S co	Fl. dan.	171
I-95 <i>elatior</i> <i>Schl.</i>	taller	<£	■ ■ ■ ■ ■	2 jn.jl	France	K80.	I) co	M. h.2.20.2	
13596 <i>subacaulis</i> <i>L.</i>	sub-stemless	31	■ ■ ■ ■ ■	■ ■ ■ ■ ■	Siberia	1824.	D co	Leh. pot.	14
<i>cincrica</i> <i>Chaix</i>					& France	1820.	D co	Scop. car.	1.23
13597 <i>reptans</i> <i>L.</i>	creeping	■ ■ ■ ■ ■	■ ■ ■ ■ ■	i jn.s	Britain	me.pa.	D co	Eng. bot.	862
13598 <i>huimifusa</i> <i>Nut.</i>	trailing	■ ■ ■ ■ ■	■ ■ ■ ■ ■	i jn.s	Missouri	1823.	D co		
13599 <i>flagellaris</i> <i>W.</i>	rod	■ ■ ■ ■ ■	■ ■ ■ ■ ■	i jn.s	Siberia	1824.	D co	Leh. pot	12
13601 <i>umbroia</i> <i>Stev.</i>	shady	■ ■ ■ ■ ■	■ ■ ■ ■ ■	i jn.s	Tauria	1818.	D co		
13601 <i>vinosa</i> <i>Lo. C.</i>	vinous	■ ■ ■ ■ ■	■ ■ ■ ■ ■	i jn.s	1818.	D co		
13602 <i>canadensis</i> <i>L.</i>	Canadian	■ ■ ■ ■ ■	■ ■ ■ ■ ■	i jn.s	N. Amer.	1800.	I) co	Nes. pot.	10.1
13603 <i>garmentosa</i> <i>W. en.</i>	twiggly	■ ■ ■ ■ ■	■ ■ ■ ■ ■	i jn.s	N. Amer.	180*	D co		
13604 <i>verna</i> <i>L.</i>	»P'ing	■ ■ ■ ■ ■	■ ■ ■ ■ ■	i jn.s	Britain	highl. p.	I) co	Eng. bot.	37
13505 <i>ojaca</i> <i>L.</i>	opaque	■ ■ ■ ■ ■	■ ■ ■ ■ ■	i jn.s	S. Europe	libO.	D co	Jac. ic.	1.91
13506 <i>crl.cea</i> <i>Leh.</i>	Sdfrron-colored	■ ■ ■ ■ ■	■ ■ ■ ■ ■	i jn.s	Switzerl.	1816.	D co	Jac. ic.	3.499
13607 <i>debilis</i> <i>Schl.</i>	sabanda Dec. filiform	■ ■ ■ ■ ■	■ ■ ■ ■ ■	i my.jl	Switzerl.	1819.	D co		
13607 <i>debilis</i> <i>Schl.</i>	weak	■ ■ ■ ■ ■	■ ■ ■ ■ ■	i my.jl	Scotland	sc. alp.	D co	Eng. bot.	561
13605 <i>atrea</i> <i>L.</i>	golden	■ ■ ■ ■ ■	■ ■ ■ ■ ■	i my.jl	S. Europe	182a	D co	Jac. ic.	1.490
2 Balisburgensis Ser.	Salisbury	■ ■ ■ ■ ■	■ ■ ■ ■ ■	i my.jn	Tauria	IK2I.	I) co	I-eh. pot	9
saliburgensis Hae.					Germany	1815.	D co	W. & K. 2.	199
13609 dtula IV.fy.K.	spreading	■ ■ ■ ■ ■	■ ■ ■ ■ ■	I my.jl	Armenia	1820.	D co	Nes. pot.	5.1
136K ^eranoides W.	Geranium-like	■ ■ ■ ■ ■	■ ■ ■ ■ ■	I my.jl	Switzerl.	1819.	D co	Hal. h.	21.4
13611 Hälleri Ser.	Haller's	■ ■ ■ ■ ■	■ ■ ■ ■ ■	I my.jl	N. Amer.	1810.	D co		
13613 pumila Pair.	dwarf	■ ■ ■ ■ ■	■ ■ ■ ■ ■	I my.jl	Italy	1822.	D co	Fl. nap.	1.44
13613 Thoia Vi Ten.	Thomas's	■ ■ ■ ■ ■	■ ■ ■ ■ ■	I my.jl	Switzerl.	1819.	D co		
I:J14 pctiiD'a Schl.	rock	■ ■ ■ ■ ■	■ ■ ■ ■ ■	I my.jl	Britain	gra. pa.	I) co	Eng. bot.	89
13613 ar^entca L.	silvery	■ ■ ■ ■ ■	■ ■ ■ ■ ■	I my.jl	Europe	1818.	I) co	I^*h. pot.	10
13616 Guntlheri Pohl	Guntlher's	■ ■ ■ ■ ■	■ ■ ■ ■ ■	I my.jl	It^dy	1820.	I) co	Fl. nap.	45
1-3617 ralckbra Ten.	Calabrian	■ ■ ■ ■ ■	■ ■ ■ ■ ■	I my.jl	Europe	1817.	I) co	Bot. cab.	706
13618 Weinmanni Lod.	Weinmann's	■ ■ ■ ■ ■	■ ■ ■ ■ ■	I my.jl	Alp.Delp.	1818.	D co	Vil. del.	3. 45
13619 inehnata ViL	inclinetl-A//?;i.	■ ■ ■ ■ ■	■ ■ ■ ■ ■	I my.jl	Europe	1817.	D co	Tau. h. c.	1.10
13620 can&cens Bcs-	cancessnt	■ ■ ■ ■ ■	■ ■ ■ ■ ■	I my.jl	Switzerl.	1786.	D co		
1-3621 intermedia /..	intermediate	■ ■ ■ ■ ■	■ ■ ■ ■ ■	I mys	S. Europe	1816.	I) co		
13622 collina VMM	hill	■ ■ ■ ■ ■	■ ■ ■ ■ ■	I my.jl	1818.	D co		
13623 inc'sa Desf.	cut-leaved	■ ■ ■ ■ ■	■ ■ ■ ■ ■	I my.jl	1820.	D co		
13634 virginita Leh.	twiggly	■ ■ ■ ■ ■	■ ■ ■ ■ ■	I my.jl	1820.	D co		
J3.3& tainca IV.	Taurian	■ ■ ■ ■ ■	■ ■ ■ ■ ■	I my.jl	Tauria	IK2I.	I) co	I-eh. pot	9
13626 thuringiaca Bernh.	Thuringian	■ ■ ■ ■ ■	■ ■ ■ ■ ■	I my.jl	Germany	1815.	D co		
13627 adscendens W. en.	ascending	■ ■ ■ ■ ■	■ ■ ■ ■ ■	I my.jl	Hungary	1806.	D co		
13:23 stipuldris L.	stipular	■ ■ ■ ■ ■	■ ■ ■ ■ ■	I lau	Siberia	1727.	I) co	Gm. si.	3.57.2
- 13629 obscura W.	obscure	■ ■ ■ ■ ■	■ ■ ■ ■ ■	I my.jl	Siberia	180(.	D co	Bal. tic.	10
13630 lacinifusa W.&K.	jagged	■ ■ ■ ■ ■	■ ■ ■ ■ ■	I my.jl	Hungary	1816.	I) co	Leh. pot	7
13631 pedMa W.	pedate	■ ■ ■ ■ ■	■ ■ ■ ■ ■	I my.jl	Kurojic	1819.	I) ro	Bot. cab.	579
I-3632 hirsata Mx.	hairy	■ ■ ■ ■ ■	■ ■ ■ ■ ■	I my.jl	N. Amer.	18W.	I) co		
13633 hrlta W.	hairy	■ ■ ■ ■ ■	■ ■ ■ ■ ■	I my.jl	S. Europe	1725.	D co		
13634 recta L.	upright	■ ■ ■ ■ ■	■ ■ ■ ■ ■	I my.jl	S. Europe	1648.	D co	Jac. au.	4.383
■■■■■ nstracanica Jac.	Astracan	■ ■ ■ ■ ■	■ ■ ■ ■ ■	I my.jl	Siberia	1787.	I) co	Jac. ic.	1.92
13635 atrusanguinea Lod.	dark-bloody	■ ■ ■ ■ ■	■ ■ ■ ■ ■	I my.jl	Nepal	1822.	D co	Bot. mag.	268)
1-637 formusa D. Don	beautiful	■ ■ ■ ■ ■	■ ■ ■ ■ ■	I my.jl	Nepal	1822.	D ro	Hook. ex. fl.	88
■■■■■ nepalensis Hook.	Russian	■ ■ ■ ■ ■	■ ■ ■ ■ ■	I my.jl	Siberia	1799.	D co	M. h.	20. 2
1V8 rutkenica W.	diffusc	■ ■ ■ ■ ■	■ ■ ■ ■ ■	I my.jl	Siberia	1817.	D co		
13639 difflsfa IV. en.	forked. leaved	■ ■ ■ ■ ■	■ ■ ■ ■ ■	I my.jl	Siberia	1773.	D co	Gm.it	1.27.1
1J640 bifurca L.	rather silky	■ ■ ■ ■ ■	■ ■ ■ ■ ■	I my.jl	Astracan	1827.	D co		
2 subsericea Ser.	trailing	■ ■ ■ ■ ■	■ ■ ■ ■ ■	I my.jl	Siberia	1696.	D co	Jac. au.	5.406
7S641 sup'ma L.	Burnet-like	■ ■ ■ ■ ■	■ ■ ■ ■ ■	I my.jl	Levant	1758.	D co	Bux.cen.	1.48
13542 pimpinelloides L.	Burnet-like	■ ■ ■ ■ ■	■ ■ ■ ■ ■	I my.jl	1827.	D GO		
13643 ornithopodioides Lan.	Cicutariaefoha W.	■ ■ ■ ■ ■	■ ■ ■ ■ ■	I my.jl	Galacca	1818.	D co	Nes. pot	2.2
13644 cicutariaefoha W.	Cicutaria-leaved	■ ■ ■ ■ ■	■ ■ ■ ■ ■	I my.jl	Dauria	1797.	D co	Nes. pot	36
13645 viscosa Donn.	clammy	■ ■ ■ ■ ■	■ ■ ■ ■ ■	I my.jl					
hspsida Nest.									
1-646 pennsylvanica L.	Pennsylvanian	■ ■ ■ ■ ■	■ ■ ■ ■ ■	I my.jl	N. Amer.	1725.	D co	Jac. vin.	2.189
hspsida W.en.									
13M7 inultifida L.	many-cleft	■ ■ ■ ■ ■	■ ■ ■ ■ ■	I my.jl	Siberia	1759.	D co	Ser. hel.	1.8.
13648 biflra W.	two-flowered	■ ■ ■ ■ ■	■ ■ ■ ■ ■	I my.jl	Siberia	1820.	D co	Leh. pot	20
1-649 verticillaris W.	whorled	■ ■ ■ ■ ■	■ ■ ■ ■ ■	I my.jl	Siberia	1818.	D co	Am. rut	16

13GBO <i>gedidea</i> BLu.	Geum-like		iJnJI	Y	Tauria	1820.	D co	Leh. pot. 2
13651 <i>Sprengelidwa</i> Lch.	Sprengel's		1 jnjl	Y	Siberia	1824,	D co	Leh. pot. 3
fragarioides W.								
13652 <i>Filipendula</i> Schl.	Dropwort		1 Jnjl	Y	Dahuria	1893.	D co	
13653 <i>Sanguisorba</i> Schl.	Burnet		1 myjn	Crea	Siberia	1826.	D co	Leh. pot. 5
13654 <i>agrimonioides</i> Bleb.	Agrimony-like		I myjn	Y	Caucasus	1817.	D co	
13655 <i>candidans</i> If. \$ B.	whitish		i myjn	Y	Mexico	1820.	D co	
13636 <i>Egedii</i> Worm.	p'gedius's		J myjn	Y	Denmark	1820.	D co	Fl. dan. 1578
13657 <i>serica</i> Z.	silky		my.s	Y	Siberia	1780.	D co	Leh. pot. 6
13658 <i>Anserina</i> £.	Goose <i>Tansy</i>		w i	my.jn	Britain	m. me. 1)	co	Eng.bot.861
13659 <i>hneariloba</i> A./\$ff.	linear-lobed		iAJ or	1 my.jl	Mexico	1824.	D co	Fl.mexc.iC in.298
13660 <i>splchndens</i> HelL	lineatū Trev.		fine jfc	A	Nepal	1822.	D co	Sw. fl. gar. 191

IT. FHACARIA'STRUM. — Petals white or red, sometimes yellow&h ; leaves pinnate, from No. 13664. palmate.

13661 <i>Saleş&vii</i> Step.	Salscovius's		ft or	2 jn.au	W	Siberia	1823	L p.)	Leh. pot. 1
13662 <i>ruprestis</i> L.	& A P'		5 my.s	W	England	al. roc.	D co	Kng. bot. 2058	
13663 <i>fragarioides</i> L.	Strawberry-Ik		I my.jn	W	Siberia	1773.	D co	Gm. si. 3.3* 2	
13664 <i>alba</i> ..	white		& A pr	i f.au	W	Wales w. tip.	I) co	Kng. bot. 13S4	
13665 <i>alchenilloulces</i> Lap.	Alchemilla-like		Cfe	A	Pyrenees	182J.	D co	Ac. tou.1.17	
13666 <i>micrantha</i> Rani.	small-flowered		A or	£ my.jn	W	Switzerl.	1820.	D co	Ser. hel.1.5
13667 <i>hybrida</i> IVallr.	hybrid		A or	4 myjn	W	Germany	18iO.	D co	
13668 <i>glabra</i> Lod.	smooth shrubby		or	2 jn.au	W	Dahuria	1818.	D co	Bot cab. 914
13669 <i>Fragaria</i> Pair.	Strawberry		A cu	£ my.jn	W	Britain banks.	D co	Eng. bot 1785	
13670 <i>caulescens</i> L.	caulecent		A or	1 my.jn	W. Y	Austria	1759.	D'co	Jac. au. 3.220
13671 <i>Clusiana</i> Jac.	Cluhius's		A pr	1 jl.au	W.Y	Austria	1806.	D co	Bot. mag. 1327
13672 <i>upmolades</i> JV.	Lupin-like		A pr	^jn.jl	W.Y	Alp. Eur. 1739.	1) co	Bot cab. 654	
13673 <i>multidentata</i> Ser.	many-toothed		A pr	g jnjl	W.R	Austria	1798.	D co	Jac. au. 5. ap. 25
nitida Jac.									
13674 <i>Valeria</i> L.	Valdert		A °r	1 my.jn	W	Piedmont	1825.	D co	Al.ped.24.1
13675 <i>nitida</i> L.	shining		A or	^nj.l.	W.R	Switzerl.	1816.	D co	Boc. mu. 9.4
13676 <i>appennina</i> Ten.	Apennine		A or	4 myjn	W	Apennines	1821.	D co	Fl. nap. 46
13677 <i>Boccuni</i> Nest.	Boccon's		A or	4 jnjl	W	Apennines	1821.	D co	Ncs. pot. 10. 2
13678 <i>tridentata</i> Sol.	three-toothed-toe		A	1.3	W	Scotland sc. alp.	D co	Eng. bot 2389	

* 1529. TO'RMENTILLIA L. SEPTFOIL. *XTormentum*, pain: supp. efficacy in tooth-ache.) *Rosacear.* 2.

13679 <i>reptans</i> L.	creeping		TA A " i jn.jl	V	Britain	me.pa.	Rs co	Eng. bot. 864
13680 <i>erecta</i> L.	erect		^ A w 1 my.o	Y	Britain	bar.pa.	D co	Eng. bot. 863

* 1530. GE'UM L.

13681 <i>canadense</i> Mur.	Canadian		to give a relish:	roots of <i>G. urbanum.)</i>	<i>Rosaceae.</i> 21.— 37.				
13682 <i>macrophyllum</i> W. en.	long-leaved		A or	2 jnjl	Y	Canada	1810.	D co	Mur. 5.4
13683 <i>heterophyllum</i> Desu.	various-lvd		A or	2 jnjl	W	Kamtsch.	1804.	D pl	
13684 <i>intermedium</i> Bes.	intermediate		A or	14 myau	Y	—	—	D co	
13685 <i>virginianum</i> Ph.	Virginian		A or	14 jl.au	W	Volhynia	1794.	D pl	
13686 <i>Album</i> Ph.	white-Jiflowered		A or	1 jl.au	W	N. Amer.	—	D pi	Her. par. III
canadense Jac.									
13687 <i>Portenschlagi</i> -1	Portenschlag's		A or	ljnjnl	Y	—	1820.	D co	
num Trat J									
13688 <i>urbanum</i> /.	city		A or	H my.au	Y	Britain woods.	D pl	Eng. bot 1400	
13689 <i>chiloinse</i> Balb.	Chile		A or	2 ...	Cop	Chile	1826.	D co	Bot. reg. 1088
coccineum B. R.									
13690 <i>ranunculoides</i> Ser.	Ranunculus-Ik		A or	1 jnjl	Y	—	1823.	D co	*
heterocarpifolium Hart.	not Desf.								
13691 <i>rivMe</i> L.	river		A or	1 jnjl	ILBr	Britain m. me.	D pl	Eng. bot. 106	
ii intermediate W.	intermediate		A or	14 myjn	Y	Germany	1794.	D pl	W. h. b. 1.69
13692 <i>nutans</i> Rafi.	nodding		A or	H jnjl	Y	N. Amer.	1825.	D pi	
13693 <i>hybridum</i> Jac.	hybrid		A or	R.Br	Europe	...	D pi	Jac. ic. 1.94	
13694 <i>pyrenicum</i> Ram.	Pyrenean		A or	h jnjl	Y	Pyrenees	1804.	D pl	Lam. il. 44
inchnatum Schl.									
13695 <i>rotundifolium</i> Lan.	round-leaved		A or	1 jnjl	Y	Russia	1820.	D co	
Adams'a <i>rotundifolia</i> Fls.									
13696 <i>brachypetalum</i> Ser.	short-petaled		A or	1 jnjl	Y	—	1818.	D co	
13697 <i>atlanticum</i> Desf.	Atlantic		A or	1 jnjl	Y	S. Europe	1810.	"D co	
sylvaticum Pou.	biflurum Brat.		strict	1 myjn	St	N. Amer.	1778.	D pl	Jac. id. 93
strictum Ph.									
13698 <i>strlctum</i> Ph.	Agrimony-Ik		A or	1 myjn	W	N. Amer.	1811.	D pl	
13699 <i>agrimonioides</i> Ph.	ciliated		A or	1 jnjl	W	N. Amer.	1818.	D co	
13700 <i>ciliatum</i> Ph.	radiated		A or	1 jnjl	Y	N. Amer.	1815.	D co	

1531. KWB.RIA Dec.

13702 <i>japonica</i> Dec.	KERRIA.	(W. Ker, a collector of plants for Kew Gardens.)	<i>Bosccea.</i> I.
Cörchorus japonicus L.	ft or	3 year V	Japan 1700. L co Bot mag. 129 <i>q</i>

1532. CALYCA'NTHUS L. ALLSPICE. (*Kalyx, calyx, anthos, flower; colored calyx.*) *Calycanthea.*

13703 <i>floridus</i> W.	florid		Sk fra 6 my.au Br	Carolina 1726. L l.p	Bot. mag. 503
13704 <i>frittilis</i> W.	fertile		A fra 3 my.au Br	Carolina 1726. L l.p	Bot reg. 404
13705 <i>lsevigatus</i> W. en.	smooth-feawrf		Si fra 3 my.jl Br	N. Amer. 1806. L l.p	Bot reg. 481
13706 <i>oblongifolius</i> Nut.	oblong-leaved		Sk fra 4 my.au Br	N. Amer. 1820. L l.p	
13707 <i>pennsylvanicus</i> Lo-C.	Pennsylvanian		SI fra 4 my.au Br	Pennsylv. 1820. L l.p	

1533. RHEE'D/4 L. RHEEDIA. (Henr. van Rheede win Draakenstein, patron of bot.) *Guttiferce?* 1.— .

13708 <i>javànica</i> Hort.	Java	<i>IDor.</i>	Java 1826. C s.l.p
-----------------------------	------	------------------------	--------------------

1534. CHIMONA'NTHUS Lindl. CHIMOXANTHUS. (*Cheimon, winter, anthos, fl.; time of fl.*) *Calycanthea.:J.*

13709 <i>fragrans</i> Lfmtt.	fragrant		A fra 6 df Y.R	Japan 1766. L l.p	Bot.mag.400
Calycanthus pra ⁿ cox L.					
SgrandiflorusLmd.	large-flowered		SI fra 8 df	Y.R China	... L e o Bot. reg. 451
31uteua	yellow		II fra 6 d.f.	Y Japan	1818. L l.p

13710 <i>octopeta</i> I.	DYRAS. (Having leaves like tne oak, which was sacred to the Dryads.) <i>Bosccefa.</i> ^
S amencana	American JU pr * jn.au W N. Amer. 1800. D s.p

13711 integrifolia Vahl entire-leaved ijn.au W Grcenland 1824. D s.p. Fl. dan. 1216
 13712 tenuella Ph. delicate i jn.au W Canada 1820. D i p. Hook. ex. fl. 220
 intcgrifolia Hook.

1536. COLITRIA R.Br. COLORIA. (Kolouros, deprived of a tail; seed.) Rosace*. 1.—3.
 13713 potentilloides R.Br. Potentilla-like L A pr i in O Siberia 1780. D p.1 Jac.vin.3.68
 Geum potentilloides L.

* 1537. SIEVER'S/i W. SIEVERSIA. Rosticeee. 6.
 13714 montana Spr. mountain Jac. au. 4.373
 13715 reptans Spr. creeping Jac. au. 5. ap. 22
 13716 anemoneoides IV. Anemone-like
 13717 glacialis Spr. icy
 Adamsii / glacialis Fis.
 13718 triflora R.Br. three-flowred N. Amer. 1826. D p.1
 Geum triflora Ph. T M T Y N. Amer. 1826. D p.1

1538. EUPOMAVTIA R.Br. EUPOMATIA. well, pomaxo, to close with a lid; fl. bud.) Anonhce*f 1.
 13720 /aurina R.Br. Laurel-like T or 4 .. N. Holl. 1824. G SJW

CLASS XIII. ORDER 1.

POLYANDRIA MONOGYNIA.

* 1539. CA'PPARIS L. CAPER TREE. (Kabar, capers, Arabic.) Capparidea?. ZJ.—118.
 13721 marianna Joe. Marianne Is. i D < w 4 ... W Timor 18^0. C 81 Jac.sc. 109
 13722 spirifosa L. common spiny * cul 3 my.au W S. Europe 1596. C s.l Bot mag. 29
 13723 ovata Desf. ovate A I or 3 my.au W S. Europe ... C s.l Boc. sic. 42.
 Fontantiss Dec.
 13724 horbacea W. herbaceous £ Al or 2 ... W Tauria 1818. S 81
 1372-3 aegyptia Lanu Egyptian • 0 or 3 ... W Egypt 1822. C sl Del. eg. 31.3
 13725 aphilla Roth leafless • CZD or 4 ... W E. Indies 1822. C L?
 13727 zeylanica L. Ceylon • i j or 6 ... W Ceylon 1819. C Braam 29
 13728 acuminata acuminate • 1 or 6 ... W E. Indies 1822. C Pluk.al.338.3
 13729 sepWma L. hedge • flor 4 ... W E. Indies 1823. C LP
 13730 Urassi Dec. Brass's • 1 or 4 ... W GoldCoast 1793. C ■
 13731 frondosa Jac. leafy • 0 or 7 ... W Cariag. bOO. C al Jac. am. 104
 13732 triflora Mil. three-flowered • 0 or 4 ... W S. Amer. ... C p.1
 13733 cynophallophora Z. Bay-leaved • 0 or 8 ... G J.W W. Indies 1752. C r.m Jac. am. 98
 13734 custachiana Jac. St. Eustach • 1 or 6 ... st St.Eustachilh22. C p.1 Jac. am. pic. 146
 13735 <alfrna f'all/ Wil lvt-leaved • 1 or 8 ... W Sant. Cruz 1807. C r.m
 13736 linear-leafed • 1 or 15 ... W W. Indies 1793. C r.m Jac. am. 102
 13737 verruedsa Jac. warty • 1 or 8 ... W Carthage. 18 JO. C s 1 Jac. am. 99
 13738 pulcherrima Jac. fairest • 1 or 10 ... W Carthage. 1700. C 81 Jac am. 106 •
 arborescens Mil.
 13739 amygdalina Lam. Amand-like • 0 or 6 ... W W. Indies 1818. C s.l PI. gen. 40.16
 13740 odoratissima Jac. sweetest-scented * 0 or 6 ... W Caracas 1814. C r.m Jac. sc. 1.110
 13741 tenuisiliqua Jac. slender-podded • 1 or G ... W Caraccas 1823. C s.l Jac. am. 105
 13742 ferruginea L. ferruginous • 1 or 4 ... W Jamaica ... C s.l Br.jam. 28.1
 octandra Jac.
 13743 Brcynia L. Breynius's • 1 or 11 ... W W. Indies 1752. C Lp Jac. am. 103
 13744 jatnacensis Jac. Jamaica • 0 or 4 ... W Jamaica 1793. C r.m Jac. am. pic. 101
 13745 torul&sa Swz. twisted • 0 or 6 ... W W. Indies 1822. C si
 uncata Lo. C.
 1-3746 undulata Lo. C. waved • 0 or 6 ... W C si
 13747 auriculkta Lo. C. eared • 0 or 6 ... W C s.l

1540. NIEBITHH/ii Dec. NIEBUHRIA. (Carslen Niebuhr, a traveller in Arabia.) Capparide*. '3.-7.
 13748 caffra Dec. Caffrarian I J or 4 ... W C. G. H. 1818. C ap

Crateva chitra Burc.
 JSS m! liaga8 (> anensis D<rc Madagascar • or 4 ... W Madagasc. 1822. C p.1
 13750 oblongiRilia Dec. oblong-leaved < i_ | or 4 ... W E. Indies 182*. C s.p
 Capparid heteroclita Rox.

RVAMEA. (John Ryan, M.D. F.R.S. a correspondent of Vuyl's.) Flacourtihncc.
 13751 speciosa Vahl showy I Q o 10 ... W Trinidad 1823. C s.l Vahl ec. 1.9
 Pdtrmnia pyrifera Rich.

H A S ? * ^ ! * H u - MARCGRAVIA. (G. Marggroaf, auth. of a voy. to Brazil.) Marcgraviaece. 2.-4.
 13752 umbellata Z. umbelled t a c u 2 0 ... W W. Indies 1792. C s.l p Jac.am.96
 13753 conacea Vahl leathery-iewarf. O cu 20 ... W Guiana 1820. C s.l p

1543. ANTHOLCTMA Lab. ANTHOLOMA. (Anthos, a flower, loma, a fringe.) Marceramhceo* 1.
 13754 montana Lar*. mountain • o r 10. . . . N. Holl. 1810. C i1 x £ v o < .

1544. NORANTEA Aub. NORANTEA. (Gonoro-Antcgr, its name in Guiana) 11
 15755 guianensis Aub. Guiana • or 4 ... V Guiana 1818. u s.l 11
 A'scyrum violaceum Aub.
 15756 brasiliensis Choi. Brazil • o r 4 . . . Brazil 1-TO G o

1545. ACTJPA L. ACT*:A. (*Aktaia*, an elm; rescmftance in foliage.) *Ranunciddee*. 3.—
 13757 spic&ta L. spiked *Bane-berry* & A or 3 ap.jn W Britain m.wo. K s.l Eng. bot. 918
 13758 alba Mil. white-berried & A or 2 ap.jn W N. Amer. ... 11 p.l Cor. ca. 77
 brachypetaia Dec. americkna Ph.  A or 1J ap.jn W N. Amer.  R p.l
 brachypéiala/3 rCbra Dec.

1546. MACRO'rYS Raft. MACROTYS. (*Makros*, long, otu, ear; capsule.) *Ranuncubets.* 1.—2.
 13760 racemosa Itaf. racemose  A or 2 ap.jn W N. Amer.  D p.l
Acts?& racemosa L.

*1547. SANGUINAMIA L. Puccox. (*Sanguis*, blood; blood-colored juice.) *Papaverdce*. I.
 13761 canadensis Z. Canadian *Bloodwort* jc A pr i mr.ap W N. Amer. 1680. R s.p Bot mag. 162
 • 1548. PODOPHYLLUM /* DUCK'S-FOOT. (*Potts*, foot, *phyton*, leaf; resemblance.) *PodophyMcea*. 1.—2.
 13762 peltatum L. peltate *May Apple* £ A cu | my R N. Amer. 1654. D s.p Bot mag. 1819

1549. CHELIDO^NIUM Bank. CELANDINE. (*Chelidon*, the swallow; time of flwg.) *Papaverhcea*. 3.—5.
 13763 m\ju₉ Mt. common larger £ A w 2 ap.o Y Britain shn.bx £ co Eng. bot 1581
 13764 grandiflora Dec. great-flowered ^ ^ or 2 ap.o Y Duhuria 1320. D co dahuricum?/₇?/.

13765 laciniatiuin Mil. jagged & A or 2 ap.o Y S. Europe ... D co Mil. ic. 1.92.1

1550. ROME'RL* Med. ROMEBIA. (J. J. fflfrnrr, prof. bot. at Landihut, died 1820.) *Papaverdcat*. 2.—3.
 13766 hybrida Dec. hybrid Q or 2 myjn P Britain hed. S co Eng. bot 201
Chelidofium hy'bridum L.

13767 refracta Dec. refracted O or 1 jnjl V Tauria 1818. S co Delcss.2.8

* 1551. GLAUCLUM Tou. HORN POPPY. (*Glaukos*, glaucous; color of plant) *Papaverdcete*. 4.—5.
 13768 flavura Dec. yellow Q or 2 jn.o Y Britain san.sh. S co Eng. bot. 8
UMcum H. K.

13769 tibicum Sm. tawny  Q or 2 s.u.8 O S. Europe 1802. S co Sw. fi. gar. 35

13770 phoeniceum 5m. purple O or 2 jnjl R England san.fi. S co Eng. bot. 1433
corniculatum L.
 2 flaviflorum yellow-flowered O or 1 jnjl Y.a Tauria 1823. S co

13771 rubrum Sm. red O or 1 jnjl R Greece 1818. S co Fl.gr. 488

* 1552. PAPA*VER Tou. POPPY. (*Papa*, pap, *Celt*; formerly used in children's food.) *Papaverhcete*. 21.—27.

I. HISPI'MCA'RPA.—*Capsules hispid.*
 13772 nudicatil L. naked-stalked Siberia 1730. S 
 1 Ibteum yellowr-flowered Siberia 1730. S 
 2 coccineum scarlet hybrid 182a S 
 3 glabratum Dec. smooth Siberia 1800. S s.l
 4 radicatum Dec. rooted Norway 1800. S s.l
 13773 rubro-auranficum F. red orange Dahuria 1822. S 
 13774 microcarpum Dec. small-fruited Kamtsch 1824. S 
 13775 pyrenaicum W. Pyrenean rjrcnees 1800. S 
aurantiacum Loi. Bar. ic. 764
 2 puniceum Dec. Tcd-Jlotttered 31 A or 1 jn.au R Pyrenees ■ S 
alpinum Lap.

13776 alpinum L. alpine  A or 1 jnjl W Austria 1759. S co Jac. au. 1.83
Biirsert Crz.
 13777 hybridum L. hybrid O or  England chai.fi. S co Eng. bot 43
 13778 Argemone L. Argemone O or  Britain corn fi. S co Eng. bot 643

II. GLABRICA/RPA.—*Capsules smooth.*
 13779 d&bium L. doubtful O or 2 jnjl Britain san.fi. S co Eng. bot. 644
 13780 /hoe'as L. Rhceas common Corn O or 2 jnjl S Britain corn fi. S co Eng. bot 645
 2 maculata *Bliottcd-Jtou>crcl* O or H jnjl Va Britain fields. S co Cur. Ion. 32
 3 coccinea scarlet O or  jnjl S Britain gard. S co Tab. ic. 1
 4 carnea flesh-colored O or  jnjl F Britain gard. S co Bar. ic. 763
 5 alba white O or  jnjl W Britain gard. S co Wein. 788. d.
 fi variegata variegated O or  jnjl Va Britain gard. S co Wein. 79a a.
 7 multiplex double o or  Va Holland 1818. S co Wein. 79a b.
 13781 trilobum Spr. thre-lobed O or  R Caucasus 1818. S co
 13782 laevigatum Bieb. smooth-leaved O or  R Caucasus 1820. S co
 13783 arenarium Bieb. sand O or 1 jn.jl R S. Europe 1818. S co
 13784 Roubi_x* Vig. Roubius's O or  R Levant 1815. S co
 13785 floribundum Detf. floribundum [^]Q or 1 jnjl S N. Holl. 1876. S co
 13786 horridum Dec. horrid O or  R S. Europe 1825. S co
 13787 setigrum Dec. brittle-bearing O or  W England corn fi. S co
 13788 somniferum L. somniferous O m 2 jlau W Britain gard. S co Lam. ic. 451
 1 nigrum Dec. black-seeded O m 2 jlau R Britain gard. S co Lob. ic. 274. 1
 2 flimbriflum fringed O or 4 jlau Va Britain gard.  Swert.fl. 2.^2.1
 3 rtibro-pleium double red O or 4 jlau Va Britain gard.  Wein. 795. c.
 4 album Dec. xwhite-seeded O m 4 jlau W Britain gard.  Eng. bot 2145
 5 albo-plenum double white-seeded O or 5 jlau Va Britain gard.  Swert fl. 2.22. 2
 6 variegum variegated O or 5 jlau Va Britain gard. 

13781 oriontMe L. oriental [^]A or 5 jlau R Levant 1714.  Bot mag. 57
 13790 bractatum Lindl. bracted  A or 4 jlau R Siberia 1818.  Lind. col. 23

pulcherrimum Fis. Caucasian O or  myjn R Caucasus 1813.  co Bot. mag. 1675

13791 caucasicum Bieb. Armenian O or  myjn Y Armenia 1815.  s.p.

1553. MECON'PSIS Vis. MECONOPSIS. (*Mekon*, a poppy, *opsis*, resemblance.) *Papaverbee*. 1.—4.

13793 cambria Vig. Welsh ffc A or 1 my.au Y England aLroc. S ap Eng. bot 66

Papaver cambricum L.

*1554. ARGEMO^NE Tov. ARGEMONE. (*Argema*, cataract of the eye; sup. med. qual.) *Papaveracea*. 4.

13794 mexicana L. Mexican O or  jlau Y Mexico 1592. S co Bot mag 243.

13795 albiflora Sims white-flowered O or 14 jlau W Mexico 1821. S co Bot mag. 23**

\$ 13796 Kulphraca Suit sulphur O or  jlau Su Mexico 1827. S co Sw. fl. gar. ic. m

17797 grandiflora Swt. great white-ftodk. A or  jlau W Mexico 1827. S co Sw.fl.gar.226

L SIDESADDLE FLOWER. (*Dr. Sarrax*, 'n, a French physician.) *Sarraci-nia*. 4.—6.

H 25 vanblanks Mr. pustuled  Alcu 1 jnjl Y N. Amer. 1752. R bog Bot. mag.  17w

adunca Ex. bot. 53

- 13800 *rhabra* Walt. red £ A1 cu 1 jn.jl P N. Amer. 1786* R bog Hook. ex. fl. 13
psittacula Afr.
- 13801 *purpurea* L. purple £ AJ cu 1 jn.jl P N. Amer. 1640. R bog Bot mag. 849
1556. OTMPHJt?A *Seek.* WATER LILY. (*Nymphae*, a water-nymph; habitation.) *Nymphydcae.* 18.—21.
I. CYANEA. —Anthers drawn out at ton; flmucrs blue or bluish; leaves peltate, entire, or bluntly sinuated.
- 13802 *acutifolia* Dec. shield-leaved *sQ5or flt jl.s B C. G. H. 1/92. R m.s And. rep. 197
oerilea Andr. and R M. not Savig.
- 13803 *cajrfilea* Savig. blue SED or flt jn.s B Egypt 1/92. R m.s Bot. mag. 2058
sieH'ita var. B. U.
- 13804 *stellata* JK starred-Jfwd £ E]or flt jn.s B E. Indies 1803. R m.s Botrep. 300
13805 *cyanea* Rox. Indian blue *(23 or flt jn.s B E. Indies 1800. R m.a
calUtra Doun
- II. LOTOS. —Flowers white or red; leaves peltate, often toothed, rarely entire, underneath pubescent.
- 13806 *edulis* Dec. eatable feGlor flt jn.s W E. Indies 1800. R m.s Par. Ion. 14
eaculenta ROT. CastMui edfilis Sal.
- 13807 *puWscns* W. pubesc. //WOTB *Lotus* = El or flt my.au Pk E. Indies 1803. R m.s Beau. ow. 2. 83
13808 *rufra* ROT. tea-flowered £ 123 or flt jl.au R E. Indies 1803. R m.s Bot mag. 1280
2 rosea B. M. roseate * CSJ or flt jl.au Pk E. Indies 1803. R m.s Bot mag. 1364
- 13809 *Lotus* L. Egyptian Lotus * IZ3 or flt jn.s Pk Egypt 1802. R m.s Del. sg. 60.1
CastMirt mystica Sal.
- 13810 *thermis* Dec. thermal Iluni. *Lotus* 2= 23 or flt jn.s W Hungary 1800. R m.s Bot mag. 797
Lotus B. M. not L.
- 13811 *ampla* Dec. ample/riv*/! * E3 or flt jn.s W Jamaica 1801. R m.s Pl.mss. 4
various-colored^ E2 or flt au Pk.w E. Indies 1807. R m.s Bot mag. 1189
- III. CASTAYI A. —Flowers white; leaves cordate, not peltate.
- 13813 *reniformis* Walt. kulney-shaped * A or flt jn.s W Carolina 1810. R m.s Dcless. 2. 5
13814 *alba* L. white i ^ or flt jn.jl W Britain riv.&c. R m.s Eng bot. 160
2canadensis Canadian % A or flt jn.jl W Canada 1820. R m.s
- 13815 *odorata* H. K. sweet-scented ^ A or flt jn.jl W N. Amer. 1780. R m.s Bot. mag. 819
13816 *lutea* Dec. smaller £^ ^ or flt jl W N. Amer. 1812. R m.s Bot. mag. 1602
odor<ta/3 *Ttiao* T B. M.
- 1S17 *nitida* R. M. shining cup-fiwd ^ A or flt jl.au W Siberia 1809. R m.s Bot. mag. 1359
1518 *pumila* Ni //. K. pygmy H A or flt my.s W China 1805. R m.s Botm.ig. 152J
blanda Mey. charming Guiana ** £E\ or flt jn.s W Trinidad 1810. R m.s
1557. XITPHAR Sm. NUPHAR. (*Neufar*, Arabic name for *Jymphæa*) *Nymphaeacear.* 5—6.
1J820 l&tea Sm. common yellow ^ A or flt jn.jl Y Brit. iwols.&c. II m.s Eng. bot 159
- 13821 *Kalmia* U. K. Kalm's ^ A or flt jl.au Y Canada 1807. R m.s Bot mag. 1243
15822 *sngittaefolia* Ph. arrow-leaved * ^ or flt jn.s Y N. Amer. 1824. R m.s
- 13823 *ptirala* Sm. dwarf yellow A A or flt jl.au Y Scotland al.lak.ll m.s Eng bot. 2298
minima E. B.
- 13824 *advena* H. K. stranger & A or flt jl.au Y N. Amer. 17/2. R m.s Bot. mag. 684
2Vympha a advena H. K.
- 15~. LIMIN' OCHARIS ton. LiMNOCHAiiis (*Limne*, pool, chairo, to delight; habitat) *HydrocharUea.* 1.—3.
- 13825 *l'uinitri* Rich. Plumicrt ^ IZ3 or 1* jn.n Y Brazil 1822. S m.s Bot mag. 2525
1559. TERNSTRCF///* L. TERNSM(EVIA. (*M. Ternström*, a Swedish naturalist.) *Ternstrocmulcev.* 4.—14.
- 13826 br'vipes Dec short-peduncled * CZ) or fi jl au II S. Amer. 1818. C s.p Col. h. rip. 38
J3827 peduncufris Dec. pedunculated ft(Z)or 6 jn au ... W. Indies 1S18. C pl
- 13828 punctiflora Sun. dotted ft(Z)or 6 jn.au G W. Indies 1820. C si Aub.gui. 1.228
13829 ventosa Spr. veiny ftQor 6 jn.au ... Brazil 182 fc C s.l
- ■ ■ ■ ■ Thun. CLEYERA. [A. Cleyer, M. D., a German botanist] *TernstreenUlicea.* 1.—2.
- 13830 *japonica* Thun. Japan ft^Jor 5. . . . Japan 1820. C s.} Ksm. 5.774. ic.
1519. FKEZIETRj Swz. FREZIEKA. (il. F. Fretier, a traveller in Peru and Chile.) *Ternstrcemidetra**. 1—7
13831 *theaodes* Swz. Tea-hke <i_J or Jamaica 1818. C s.l
- * 1062. LETTSOM/ A. R. & P. LETTBOMM. (*Cockley Lettsom*, an English naturalist.) *Ternstreenmidetra*. 1.
13832 *tomentosa* R.S/P. woolly ft i_J or 4. . . . Peru 1823. C s.1 Fl. per. 14
1503. BROWNLO'W/ il Rox. BROWNLOWIA. (The late L n ^ JB^tmrn/ow. dnugh. of Sir A Hume.) *TOihceee.l*
13833 el^ta Rox. tall £ • or 60 ... Y E. Indies 1830. C s.1 Rox. cor. ic.
- ■ ■ EURY^AL-E Sal. EURYALE. (*Euryne*, one of the Gorgons; thorny menacing habit.) *Kympteteete.* 1.
13834 *ferox* Sal. fierce & El or flt jl.s II India 1809. R m.s Bot. mag. 1447
Annesl&a spinosa B. R.
- 1.165. BI'XA L. AVOTTA. (US South American name.) Bixinc*. 3.
- 13835 *Orellana* Orellana IUJra 2) my.au Pk W. Indies 1690. S s.p Bot.mag. T466
JS.'5 Jmr. n^frea/for. P^Tle J a m 20jn.au P E. Indies 1817. S s.p
- 13837 *Urucuriuna* W. Urucu 2 a » « 20 jn.au Pk Brazil 1820. S s.p
- ■ ■ BE'RRYi* Rox. BERRYA. (Dr. Berry, who first introduced it to bot gard. of Calcutta.) *Tiliacete.* 1.
Amomilla Rox. Amomilla £ • or 20 jl.au ... E. Indies 18J0. C p.1
- ■ ■ ■ ■ (Not explained.) Bixinee. 1.—7.
- £££(oroc) ^ serraate ftQeu 4 jlau Y W. Indies 189& C s.n Bot reg. 97-2
VS2^rr^ ^ serraate ftQeu 6jn.au Y Montserrat 1823. C s.1
1J841 *Mimuraria* Dec. Tea-formed ftQeu 6 jnau Y Bourbon 1820. C s.1 Bory 2.24
Ludia ncterophyrlalla^/y
1568. LIU'DIA Lam. LuniA. (*Ludo*, to sport: in shape of lys. in young and old plant) *BUtncte.* 2—3
18<2 heterophylla lutm. various-leaved ft CD or 4 jl.au Y Mauritius 1823. C sp I*in.IL466. L2
13843 sessiliflora IMW. se aisle-flowered ft Q or 4jl.au Y Mauritius 1820. C s.p Jacsc 1.112
1569. itiSTIA L. LJETIA. (J. de Laet, of Antwerp, author of a history of America.) *Bixinee* 1.—fi
138M Thamnij Sun. Thamnia ft a or 4 jn.au W V. Indies 1824. C s.p Br.jajn.25?
1570. SLOANE^ L. SLOANEA. (Sir Hans Sloane, principal founder of British Museum) *Tilitaxm* 9—s
13845 dentata L. tooth-leaved t Q t m 4 0 ... W S. Amer. 1752. C p.1 PI. ic?244
3B4S *ineinaria^ngi* Aub. Sinemaria H Q or 15 jn.jl W Guiana 18-23. C s.p Aub. gui. 212

- | | | | |
|--|-------------------|--|--|
| 1571. APEVBA Marg. | APEIDA. | (Its name in Guiana.) | Tiliacea. 4.-8. |
| 13847 <i>Tibofirbou</i> Aub.
hirsute Lam. | Tibourbou | ■ • OR 7 ... Y | S. Amer. 1756. C p.l. Aub. gui. 1. 213 |
| 13848 <i>Petowno</i> Aub.
hispida Gae. | Petourao | ■ □ ■ 40 .. Y | S. Amer. 1817. C p.l. Aub. gui. 1. £15 |
| 13849 <i>aspera</i> Aub. | rough-capsuled | ■ • tin CO — Y | Cayenne 1792. C ■ Aub. gui. 1. 216 |
| 13850 <i>beta</i> Aub. | smooth-graved. | ■ • or 10. -- G | Cayenne 1817. C ■ Aub. gui. 1. 21+ |
| 1572. SPARRMA'NN/J L. | SPAHBMANNIA. | (A. Sparrmann, trav. in China and C. G. H.) | Tiliticce. 1. |
| 13851 <i>africana</i> L. | African | £ L-J or 10 mr.jl W | C. G. H. 1700. C p.l. Bot. mag. 516 |
| 1573. ENTELEPA R. Br. | ENTELEA. | (Enteles, perfect; all the stamens fertile) | Tilidcrv. 1. |
| 13852 <i>arbortscens</i> R. Br. | arborescent | f U or 20 ray W | N. Zeal. 1820. C p.l. Bot. mag. 2480 |
| 1574. MUNTI'NGJJ L. | MUNTXGIA. | (Abraham Hunting, prof, bot at Groningen, d. 1683.) | Tiliacea. 1. |
| 13853 <i>Calabhra</i> L. | Calabura | • • c u 3 jn.jl [V | Jamaica 1690. C p.l. Jac. am. 107 |
| 1575. GRE'W/y/l J. | GREWIA. | (Nehemiah Grew, M.D. F.R.S., physiological botanist) | Tiliacea*. 27. — 65. |
| 13854 <i>tomentosa</i> J.
hirsuta 5/. | woolly | ■ (or 10 - | Java |
| 13855 <i>hirsuta</i> Vahl | hairy-leaved | • • OR 10 ... | P E. Indies 1816. C p.l. |
| 13856 <i>bracteata</i> Roth | /arge-bracted | ■ ■ or 10 ... | P E. Indies 1820. C p.l. |
| 13857 <i>Mallococca</i> L.
Mallocucca crenata
Forst. | woolly-berried | ■ ■ or 10 aus. | Pa.P E. Indies 1792. C ftp Forst. 29 |
| 13858 <i>M/crocos</i> 77. X. | small-berried | • • OR 7 ... | G E. Indies 1799. C ■ 11. mal. 1.56 |
| 13859 <i>oppositifolia</i> Buc. | opposite-leaved | ■ ■ OR 10 ... | P Nepal 1818. C ■ |
| 13860 <i>umbellata</i> /Sox. | umbellate | ■ ■ OR 10 ... | P E. Indies 1816. C p.l. |
| 13861 <i>bicolor</i> J. | two-colored | ■ ■ OR 10 ... | W Guinea 1818. C ■ An. mu. 1. 50.2 |
| 13862 <i>Ruthii</i> Dec. | Roth's | ■ ■ OR 10 ... | W E. Indies 1819. C ■ |
| 13863 <i>jaivipa</i> Mia Roth | Sago-loaved | ■ ■ or 6 jl.s | W E. Indies 1818. C ■ |
| 13864 <i>flava</i> D.r. | yellow | ■ ■ OR 10 ... | Y C. G. H. 1819. C ■ |
| 13865 <i>ovalifolia</i> J. | oval-leaved | ■ ■ OR 8 jl.s | W E. Indies 1818. C ■ |
| 13866 <i>affinis</i> Lindl. | similar | ■ ■ or 3 ... | G China 1824. C CO |
| 13867 <i>paniculata</i> Box. | panicled | ■ ■ or 10 ... | E. Indies 1816. C ■ p.l. |
| 13868 <i>oriental</i> is L. | oriental | ■ ■ OR 10 jl.au | W E. Indies 1767. C ■ p.l. |
| 13869 <i>pildsa</i> Lam. | pilose | ■ ■ or 10 ... | W E. Indies 1804. C ■ p.l. |
| 13870 <i>scrutata</i> Ttec. | serrulated | ■ ■ or 6 jl.s | W E. Indies 1818. C ■ p.l. |
| 13871 <i>carpinifolia</i> J. | Hornbeam-lvd | ■ ■ or 10 ... | W Guinea 1823. C ■ p.l. |
| 13872 <i>occidentalis</i> L. | occidental | ■ ■ or 10 jl.s | p C. G. H. 1690. C ■ p.l. |
| 13873 <i>asiatica</i> X. | Asiatic | ■ ■ or 12 jl.au | p E. Indies 1792. C ■ p.l. |
| 13874 <i>subinsquamis</i> Dec. | rather unequal | tt • or 10 ... | W E. Indies 1816. C ■ p.l. |
| 13875 <i>aspera</i> 76/A | rough | • • or 7 jl.s | W E. Indies 1818. C ■ p.l. |
| 13876 <i>filiaef</i> Mia Vahl | Lime-trce-lvd | • • or 12 - | W E. Indies 1812. C ■ p.l. |
| 13877 <i>villosa</i> W. | villous | ■ ■ or 8 jl.s | EL Indies 1816. C ■ p.l. |
| 13878 <i>terebinthinae</i> Dec. | turpentine-*C | ■ ■ or 10 - | W 1820. C ■ p.l. |
| 13879 <i>obtusif</i> Mia W.en. | blunt-leaved | » ■ or 6 - | W C. G. H. 1818. C ■ |
| 13880 <i>ulmifolia</i> Box. | Elm-loaved | • • or 10 ... | W E. Indies 1816. C ■ p.l. |
| 1576. 7TL1A L. | LIME TREE. | (Plelea, the elm; resemblance?) | Tiliacea. 9.—10. |
| 13881 <i>microphylla</i> fen. | small-leaved | ■ tm 50 aus. | Y.o Britain woods. L co Eng. bot. 1705 |
| parvifolia Ehrh. | | tm 50 | |
| 13882 <i>intermedia</i> Hay. | intermediate | ■ tm 50 jn.au | Y.o Britain woods. L co Fl. dan. 553 |
| 13883 <i>riibra</i> Dec. | common red | ■ tm 50 jn.au | Y.G Britain woods. L co |
| conthhai-a Bose | | | |
| 13881 <i>platyphylla</i> Sco. | broad-leaved | *£ tm 50 aus. | Y.o Britain woods. L co Ven. til. 1.2 |
| grandiflha Ehrh. | cordifolia Bes. | | |
| 2 laciniata Mil. | fringed | ■ tm 50 au | Y.o Britain ... L co |
| 13885 <i>americana</i> L. | American | ■ tm CO jn.jl | Y.o N. Amer. 1752. L co Den. br. 134 |
| glabra Ven. canadensis Mx. | loose-flowered | Y tm 50 jn.jl | W N. Amer. 1820. L co |
| 13886 <i>laxiflora</i> Mx. | pubescent | £ tm 20 jl.au | Y.o N. Amer. 1726. L co Den. br. 135 |
| 13887 <i>pubescens</i> H. K. | thin-leaved | ^ tm 20 jl.au | Y.G N. Amer. ... L co |
| caroliniana 3//. | white | £ tm 30 ju.au | Y.G Hungary 1767 L co Den. br. 71 |
| 2 leptophylla Ven. | | | |
| 13888 <i>alba</i> W. § K. | various-leaved | *£ tm 30 jn.au | Y.o N. Amer. 1811. L co Yen. til. 5 |
| argentea Dec. | | | |
| 13889 <i>heterophylla</i> Ven. | | | |
| alba Mx. | | | |
| 1577. CO'RCHORUS L. | CORCHOBUS. | (Kore, pupil, horco, to purge; sup. med. qual.) | Tiliacea. 1. — 4. |
| 13890 <i>siliquiflus</i> L. | •iliquose | tt. ■ u 3 jn.au | R W. Indies 1732. C l.p. Jac. vin. 3.59 |
| 13891 <i>hirtus</i> L. | hairy | tt. ■ e u 2 jn.au | Y S. Amer. 1820. S p.l. P. ed. B. 103. 2 |
| 13891J <i>pilobolus</i> Lk. | hairy-podded | ■ ■ 2 jn.au | Y S. Amer. 1818. S p.l. |
| 13893 <i>triloculfans</i> L. | thrcellcelled | ■ ■ 1 jn.au | Arabia 1790. S co Jac. vin. 2.173 |
| 13894 <i>olitrius</i> L. | potherb | ■ ■ 2 in.au | India KMO S co Cam. h. 12 |
| 13S9J <i>ae/stuans</i> L. | Bcnri'hing | ■ ■ 2 jn.jl | S. Amer. 1731. S co Jac. vin. 1. 85 |
| 13896 <i>tifidens</i> L. | three-toothed | ■ ■ cu 2 jn.au | Senegal 1824. S p.l. Pluk. al. 127.4 |
| 13897 <i>acutangulus</i> Lam. | acute-angled | ■ ■ un 3 jn.jl | E. Indies 1816. C co Pluk. ph. 44.1 |
| 13898 <i>capsularis</i> L. | captular | O un lfj.jl | E. Indies 1725. S l.p. Ru. am. 5.78. * |
| 13899 <i>hirsutus</i> L. | hairy-capsuled | tt. un 3\$ jn.jl | Y S. Amer. 1752. S p.l. P. ed. B. 104 |
| 1578. GRPAS L. | AXCHIOVY PEAR. | (Grao, to eat; fruit.) | Guttiferis ajpnis. 1. |
| 13900 <i>cauhfdrla</i> L. | stem-dowering | £ • fr 50 - | W Jamaica 1768. C l.m Sl.jam.2.217.1. ■ |
| 1579. CALOPHY'LUM L. | CALOPHVLLUM. | (Kalos, beautiful, phyllon, a leaf.) | Guttifene. 4.—* |
| 13901 <i>Inophy'lum</i> L. | tibrous-leaved | t • tm 90 ... | W E. Indies 1793. C 81 It mal. 4.38 |
| 13902 <i>Tacamanalitca</i> W. | Tacainahaca | J • tm 30 ... | W Bourbon 18^2. C s.l. Pluk. al. 147.3 |
| 13903 <i>C'daba</i> Jic. | Cnlaba tree | t • tm 30 ... | W India J780. C s.l. Jac. am. 165 |
| 13y04 <i>spurmm</i> Dec | spurious | f • tm 30 ... | W Malabar 1800. C s.l. It. mal. 4.39 |
| <i>Cdlaba</i> I« | | | |
| 1580. MAMME'A L. | MAMMEE TBEE. | (Matney, its American name.) | Guttifene. 1.—3. |
| 13905 <i>americana</i> L. | American | £ ■ & (W ... | W Amer. 1737. C s.l. Jac. am. 182.8 |
| 1581. BONNE' TM | Schreb. BONNKIIA. | (Charles Bonnet, a Genevese naturalist.) | Guttifere ct. 1.— |
| 13906 <i>nobilis</i> Vahl | marsh | FOOR | R Trinidad 1819. C s.l. Aub. gui. 1. 2*? |

1582. O'CHNA Schreb. OCHNA. (Greek name of the wild pear tree; some resemblance.) *Ochnhcece.l.—M.*
 13907 obtusata Dec. blunted ft • or 4 jl.au Y E. Indies 1790. C Lp Rox. cor. 1.89
 squarrusa L.
 13908 lfcida Dec. bright ft • or 6 ... Y E. Indies 1819. C Lp Sc. dansk. 2. 6
 13909 nitida Thun. shining ft • or G ... Y C. G. H. 1815. C Lp Dec. mu. 17.2
 13910 multiflora Dec. many-flowered ft l-or 8 ... Y S. Leone 1820. C l.p Dec. mu. 17.3
 13911 mauntidna Lam. Mauritian HI— or 8 ... Y Mauritius 1822. C l.p Dec. mu. 17.5
 13912 arborea Bare. tree fQor 20 ... Y C. G. H. 1823. C l.p
 1o913 atropurpurea Dec. dark-purple JZuMff *L_Jor 4 ... P C 6. H. 1816. C l.p Pluk. aU 263.1,2
- * 158a DELPMA L. DELIMA. (*Delimo*, to shave off; leaves used for polishing.) *DtUenAcej. 1.-.6.*
 13914 s.irmontosa /,. sarmentose £. • or 10 ... Y Ceylon 1820. C Lp Bur. in. 37.1
 Tetracera sarmentosa Vahl
1584. DOLIOCA'RPU\$ Rai DOLIOCARPUS. (Dolio*, deceitful, Aaroo*. fruit; poisonous.) *DUIentocco*.l. — 4.*
 13915 Calini-a Gm. Calinea L O or 10 ... Y Guiana 1822. C Lp Aub. gui. 1.221
 Tetracera Calinéa W. *Calinéa scändens* Aub.
1585. ■■■■■ Corr. BENGAL QUINCE. (*Male*, one of the Hesperides.) *Aurantidcea. 1.—2.*
 13916 Marmelos Corr. Marmelos ftQfr 6 ... W.R E. Indies 1759. C 1 Rox. cor. 2.143
1586. BERTHOLE^T/^ H. & B. BBAZIL NOT. (*L. C. Bertholet*, a celebrated chemist.) *Lecythidc*. 1.*
 13917 cxcelsaJ7.421 tall 1 O P 100. . . . Para ... C pls
- 1587. LAGEItSTIU'E'MI^ L. LAGERSTIUMBIA. (*Magnus Lagerstram*, friend of Lin.) *Salicari**. 4. — 7.
 13918 indica L. Indian ft O or 6 au.o P E. Indies 1759. C s1 Bot. mag 405
 13919 regtaa Rox. queen's * Q or 12 ... R E. Indies 1792. C p1 Rox. cor. 1&>
 13920 graiidiflra Rox. large-flowered »• or 12 ... R E. Indies 1818. C p1
 13921 parviflbra Rox. small-flowered ftQor 12 ... W> E. Indies 1818. C p1 •
1588. LE'CYTHIS L. LECYTIHS. (*Lekythos*, an oil jar; eed-vessel.) *Leeyth\de**. 6. — H.
 13922 grandiflbra Aub. large-flowered f a or 60 ... Yw Trinidad 1824. C s1 Aub. gUL 2. 283
 13923 minor jac. smaller SB i i ur N ... Yw Carthage 1825. C s1
 13924 amdra Aub. bitter ft[Z]or 10 ... Yw Guiana 1825. C 81 Aub. gui. 2. 286
 13925 Adnthusn Aub. Adatimon ft a or 8 ... Yw Maranh. 1825. C 81 Aub. gui. 2. 289
 13926 pamflbra Aub. small-flowered ftj Jor 8 ... Y.w Trinidad 1825. C 81
 13927 bracteata Aub. bracted ■■■ or 10 ... Y.w Maranh. 1825. C 81 Aub. gui. 2. 282
- *1589. CTSTUS Tou. ROCK ROSE. (*Jdste, ahox*) capsule.) *Cistinece. S4.*
 I. ERYTHROCI'STUS.—Outer se p. narrow, inner with scarious margins; pet. rose-red or purple; caps. 5-locular.
 13928 purpureus Lam. purple 41 or 2 my,jl P — — — — Sw. cis. 17
 13929 heterophylus2)?tf. various-leaved * or 2 jnjl P Algiers — — — — s.p Sw. cis. 6
 13930 parviflorus Lam. small-flowered * or 3 jnjl Pa.R Crete 1800. — — — — s.i Sw. cis. 14
 13931 corapicatus Lam. complicated * or 3 jnjl R Spain 1818. — — — — 61
 13932 villus Lam. villous * or 3 jnjl P S. Europe 1640. — — — — Duh. ar. 1.64
 13933 créticus L. Cretan * or li in.au P Levant 1731. — — — — Jac. ic. 1.95
 2 tatinicus Dec. Taunan • or 2 jnjl P Tauria 1817. — — — —
 13934 incanus L. hoary • or 2 jn.au P S. Europe 1596. — — — — s.p Bot. mag. 43
 13935 DunalwMff* Swt. Dunal's * or 9 myjn ■ — — — 8.p
 undulatus Dunal
 13036 cribpus L. curled-leaved * or ■ jnjl P Portugal 1656. — — — — Sw.cis.22
 13937 filbidus L. white-leaved * or ■ jnjl Pa.P Spain 1P40. — — — — Par. th. 1
 13938 candidissimus Dunal whitest * or ■ Jn.jl Pa.R Canaries 1817. — — — — Sw. cis. 3
 13939 vaginatus H. K. sheathed * or ■ apjn Pa.P Tenenfife 1779. — — — — Sw. cis. 9
 13940 scirceus Vahl silky * or ■ jnjl R Spain 1826. — — — — Ef Bar. ic. 1315
 13941 platysepalus Swt. broad-sepaled A or ■ jnjl R — — — — Sw. cis. 47
 13942 canehcems Swt. canescens lk or ■ jnjl R — — — — C 81 Sw. cis. 45
 13943 cymdsus Dunal cymose * or ■ jnjl P — — — — C 61
11. LEDONI A — Sep. 5,2 outer ones largest s pet. white; stigm sub sessile, large capitate; caps. 5 orlO locular.
 13944 aalifiblius L. Sage-leaved * or 2 jnjl W S. Europe 1548. C s.p Sw.cis.5i
 13945 corboriensis Pou. Corbor ft ~ or 1/.my.jn. W Spain i ~ 165 & .C s.p Sw. cis. 8
 13946 obtusifolius Swt. obtuse-leaved ft or 3 jnjl W — — — — C s1 Sw. cis. 42
 13947 psilosépalus Swt. smooth-sepaled ft or 3 jnjl W — — — — C s1 Sw. cis. S3
 13948 florcntinus Lam. Florentine ft or 3 jnjl W Italy" 1825. — — — — 8.1 Sw. cis. 59
 13949 monspchenis L. Montpelier ft or 2 jnjl W S. Europe 1656. — — — — 8.p Jac. c 2.8
 13950 L'don Lam. Lcdon Gum ft or 1 jl.au W France 1730. — — — — s.p Duh. ar. 1.66
 13951 hirsutus Lam. hairy ft or 2 jnjl W Portugal 1656. — — — — Sw. cis. 19
 13952 laxus //, A. loose ft or 2 jnjl W Spain 1056. — — — — Sw. cis. 12
 13953 longiflMius lam. long-leaved ft or 4 jnjl S. Europe 16X. — — — — » Sw. cis.
 13954 popuhfilius Lam. Poplar-leaved ft or 3 myjn W Spain 1656. — — — — Sw. cis. 23
 13955 LitifMius Swt. broad-leaved • J or 4 jn W — — — — Sw. cis. 15.
 13956 eoiditfllius Swt. heart-leawl ft or 4 jnjl W — — — — s.p Sw. cis. ic'in.
 13977 /aurifi.lius L. Lnurel.leavcd ft or 4 jnjl W Spain 1731. — — — — B.p Sw. cis. 52
 13958 ladaniferus L. ladanum-bg ft or 4 jnjl W Spain 1629. — — — — 8.p Bot. mag. 112
 Salbflorus white-flwd gum ft or 4 jnjl W Spain 1700. — — — — Ef Sw. cis. 84
 fl< « 3m a 2 urat8 "potted ft or 4 jnjl W Spain 1700. — — — — S.I Sw. cis. 1
 13959 unduKvts Lie. waved ft or 4 jnjl W S. Europe 1800. C p1
- c^priu Lam. Cyprus ■ or 4 jnjl W Greece 1800. C p1 Sw.cis.39
 ladaniforus B. M. stcnohftlus Lk ■ or 3 jnjl W Spain 1810. C a.p Sw. cis. 32
 13961 Cluwi Dunal Clusius's ■ or 3 jnjl W Spain 1800. C a.p Sw. cis. 32
- *1590. HELIA'NTHEMUM Tou. SUN ROSE. (*Helios*, sun, *anthemon*, fl.; golden blos.) *Cistne**. 115—124.
 J. H[A]LIMUM.—Sepals 3, rarely 5; petals very often spotted at base; stigmas capitate, subtrilobed.
 13962 Libanotis IV. Rosmary-M II—J or 1 jn Y Spain 1752. C p1 Bar. ic. 294
 13963 urheatum Mil. wmbel-/u/ ft J or 2 jnjau W S. Europe 1751. W pi Sw.cis.5
 13964 CA scabrusum Pers. rough mt J or 3 jnjil Y Italy 1775. C p1
 13965 algarvi-nse Dunal ■■■■■ J or 3 jl.au Y Portugal 1H00. C p1 Bot. mag. 627
 13966 formosum Dunal ■■■■■ J or 4 myjl Y.Br Portugal 1780. C p1 Sw. cis. 50
 13967 filtripliciflMium W. beautiful m J or 6 jnjil Spain 1656. C p1 Bar. ic. 292
 13968 Aalimiflbum W. Orache-leaved ft J or 6 jnjil Y Spain 1656. — P-1 Sw. cis. 4
 13969 ocymo'ides Pers, Sea-Purslane-lydf. ft J or 4 jnjil Y Spain 1800. — B.p Sw. cis. 13
 ampuciflThum Cav. not MU. — — — — C — Vcn.ch.20
 13970 alyssoidcs Fen. Alysbum-like ft J or 3 m Y Spain 1800. C ■■■■■ Vcn.ch.20
 13971 rugdsum Dunal wrinkly si J or 3 jnjl Y Portugal 1800. C s.p

13972 <i>lasianthymum Pers.</i>	hairy-flwd	• J or	3 M	Y	Spain	1826.	sp	Bar. ic. 289	
13973 <i>involucratum Pers.</i>	involute	• I or	3 jn.jl	Y	Spain	1826.	s.p		
13974 <i>chiranthbides Pers.</i>	Stock-leaved	* I or	3 jn.jl	Y	Portugal	1818.	s.p		
13975 <i>candidum Swt.</i>	white-leaved	• or	3 jn.jl	Y	Spain	..	8.1	Sw. cis. 2J	
13976 <i>elongatum Pers.</i>	long-peduncled	* I or	2 jl	Y	Spain	1800.	P1		
II. LECIEOPDBS									
13977 <i>canadense Mx.</i>	Canadian	& A or	1 jn.jl	Y	N. Amer.	1799.	S s.l	Sw. cis. 21	
13978 <i>rosmanniflum Ph.</i>	Koseinary-lvd	or	1 jn.jl	PP.Y	Gandia	1823.	C sp		
13979 <i>glomeratum Lag.</i>	glomerate	tt. or	1 jn.jl	Y	Mexico	1820.	C s.p	Sw. cis. ic. in.	
13980 <i>ramuliflorum Mx.</i>	branchlet-flwd	A or	1 jn.jl	Y	Carolina	1823.	C s.p		
13981 <i>brasiliensc Pers.</i>	Brazilian	JL, & or	3 jn.jl	Y	Brazil	1823.	C s.p	Sw. cis. 43	
13982 <i>polygaUefohum Su//.</i>	Milkwort-lvd	it., pr	1 jn.au	Y	Carolina	18i3.	C s.p	Sw. cis. 11	
13983 <i>epipactis Thib.</i>	Carolina	larger	smaller;	pet. yellow	spotted at base	;	stigm. capitate.		
13984 <i>Tuberaria Mil.</i>	Tuberina	J. Ajor	i jn.jl	W	S. Europe	1752.	S s.p	Sw. cis. 18	
13985 <i>clantagineum Pers.</i>	Plantain-like	Jl A or	* jn.jl	Y	Spain	1823.	S s.p		
13986 <i>filigl6sumSui/.</i>	woody	tt. or	i jn.jl	Y	... MOI	1804.	S sp	Sw. cis. 46	
13987 <i>globularia>f.Mium Pers.</i>	Globul.-lvd	Jf Ajor	1 jn.jl	Y	Spain	1810.	S		
13988 <i>guttatum Mil.</i>	spottou/airf	O or	1 jn.jl	Y	England	san.pa.	S s.l	Eng. bot. 544	
13989 <i>serratum Pers.</i>	savr-petaled	O or	1 jn.jl	Y	Spain	180k	C s.p	Cav. ic. 2.175.1	
13990 <i>eriocaulon Dunal</i>	woolly-stemmed	O or	1 jn.jl	Y	Spain	1817.	S s.p	Sw. cis. SO	
13991 <i>punctatum W.</i>	dotted	O or	i jn.jl	Y	S France	1816.	S s.l	Sw. cis. 61	
13992 <i>niconspicuum Thib.</i>	inconspicuous	O or	i jn.jl	Y	Piedmont	1819.	S s.p		
IV. MACULANUA.—Sepals five, outer ones narrow; pet. often spotted at base; stigmas small subtrilobed.									
13993 <i>lunulatum Dec.</i>	lunulate	tt.	or	± jn.jl	Y	Span	1826.	S s.p	Al.au. 2.3
V. BRACLYPE TALOV.—Sep. 5, outer ones minute; pet. often spot, at base, small; stigm. simple: caps triquetrous.									
13994 <i>lepidolum Thib.</i>	Lithos-leaved	Q or	j jn.jl	Y	England	san.pa.	S s.l	Eng. bot. 2414	
13996 <i>niloticum Pers.</i>	Nile	Our	i jn.jl	Y	Egypt	1817.	S s.p	Sw. cis. ic. in.	
2 nūjus Dec.	larger	O or	1 jn.jl	Y	Europe	1817.	S s.p	Sw. cis. ic.in.	
3 proctrbmbs Dec.	prorumbeut	O or	\ jn.jl	Y	France	1811.	S s.p		
13997 <i>denticulatum Thib.</i>	denticulate	O or	1 jn.jl	Y	France	1818.	S s.p	Sw. cis. ic. in.	
caliciflum Gou.									
13998 <i>*alicitulum W.</i>	Willow-leaved	O or	4 jn.au	Y	S. Europe	1759.	S s.p	Cav. ic. 2.144	
LIDO <i>sanguincum Lag.</i>	bloody-stem'ned	O or	i jn.jl	Y	Spain	182&	S	Sw. cis. ic. in.	
14000 <i>ajgyptiacum Mil.</i>	Egyptian	O or	f jn.jl	W	Egypt	1161.	S if	Jac. ob. 3. 68	
VI. EKIOLAHPON.—Sepals five, outside silky, two outer ones minute; styles at base twisted: capsules pilose.									
14001 <i>canariense IV.</i>	Canary	£> L-J or	1 jn.jl	Y	Canaries	1790.	C p.l	Jac. ic. 197	
14G02 <i>lilppit Pers.</i>	Lippius'	«. or	J jn.jl	Y	Egypt	1820.	C s.p	Sw. cis. ic. in.	
14003 <i>kahfricum Del.</i>	Kahinan	tt. J or	j jn.jl	Y	Egypt	1820.	C sp	Del. leg. 31.2	
14004 <i>confertum Dunal</i>	close-flowered	tt. J or	J jn.jl	Y	Tenerife	...	C s.p		
VII. FUMA ^{NA} .—Cal before flwg. a HI tie twisted, 5-scalp, 2 outer ¹ ones small; styl. capiLfrinj. subtrilob.									
14005 <i>Fumana Mil.</i>	Fumana	tt.	J or	1 jn.jl	Y	France	171.	C s.p	Sw. cis. 1G
14006 <i>mitrofoles Dunal</i>	Erica-like	ttt. J or	J jn.jl	Y	S. Europe	...	C s.p	Cav. ic. 2.172	
14007 <i>proi-imbens Dunal</i>	procumbent	JL	or	J jn.jl	Y	S. Europe	...	C s.p	Bar. ic. 445
14008 <i>la/vipes W.</i>	smooth-footed	tt. J or	J jn.au	Y	France	1690.	C s.p	Sw. cis. 24	
14009 <i>arabicum Pers.</i>	Arabian	tt.	J or	J jn.jl	Y	S. Europe	1826.	C s.p	
viscidulum Stev.									
14010 <i>Barrel)eri Ten.</i>	Barrelier's	tt. or	J jn.au	Y	Italy	1820.	B.p	Bar. ic. 416	
14011 <i>la^veiVr*</i> .	smooth	tt. or	J jn.jl	Y	Span	182a	S.p	Cav. ic. 2. 145.1	
14012 <i>viride Ten.</i>	green	«L or	J jn.jl	Y	Italy	1825.	s.p		
14013/ <i>uniptrinuin iMg.</i>	Jumper-like	tt. or	J jn.au	Y	S. Europe	1800.	s.n	Bar. ic. 443	
14014 <i>glutin>suin Pers.</i>	clammy	tt. or	J my.s	Y	S. Europe	1790.	p.l	Cav. ic. 2.145.2	
14015 <i>fllyimflblum Pers.</i>	Thyme-Icaved	tt. or	14 jn.au	Y	Spain	1658.	s.p	Bar. ic. 444	
VIII. PsEuocniSTUS.—Sep. 5, outer ones minute; pet. small; styles at base twisted; stigm. capitate 3-lobed.									
14016 <i>origaniflum Pers.</i>	Marjoriam-lvd	tt. J or	1 jn.au	J	Spain	1795.	C s.p	...	
J4017 <i>mtile Pers.</i>	soft	* -> J or	1 jn.au	J	Spain	18J.	C s.p	Cav. ic. 2. 32f2	
14018 <i>dictiotomum Dunal</i>	dichotomous	tt. — or	1 jn.au	J	Spain	1826.	C sp	Cav. ic. 3. 263.1	
14019 <i>oalandicum Dec.</i>	(Eland	JL	or	J	Germany	1816.	C s.t)	Sw. cis. 2	
14020 <i>alpfotreDunal</i>	alp	JL	or	J	Germany	1818.	C s.l	Cr. au. 6.1	
14021 <i>itâhcum Pers.</i>	Italian	tt. —(or	1 jls	J	Italy	1799.	C s.p	Bar. ic. 366	
14022 <i>penicilUtum Thib.</i>	pencilled	JL	or	* jn.jl	Spain	1817.	B.p		
14023 <i>obovatum Dunal</i>	obovate	tt. —J or	1 jn.jl	J	Spain	1826.	s.p		
14024 <i>vineMc Pers.</i>	vine	JL	or	i jn.jl	Germany	1817.	E SD	8w. cis. 77	
14025e <i>AnnumDunal</i>	hoary	JL	or	g jn.jl	S. Europe	1772.	B.p	Sw. cis. 56	
14026 <i>maritillum Dec.</i>	Marum-leaved	JL	or	my.jn	Britain	al.roc	C sp	Eng. bot 396	
14027 <i>rotundiflumDtnff</i>	round-leaved	JL J or	} jn.au	X	Spain	JBJf.	C s.p	CAV. IC. 2.142	
14028 <i>pamecuKitum Dunal</i>	panicked	JL J or	jn.»u	X	Spain	1826.	C s.p		
14029 <i>cin&rum Pers.</i>	grey	tt. —J or	1 jn.au	J	Spain	1818.	C s.p	Cav. ic. 2.141	
14030 <i>crassifiliuin Pers.</i>	thick-leaved	tt. or	1 my.jl	J	Barbary	1818.	C 8.1		
14031 <i>squamatum Pers.</i>	scaly	tt. —J or	1 jn.jl	J	Span	1815.	C 8.1	Cav. Ic. 8.139	
IX. EIHELIANTHIEMUM.—Calyx 5-sepal3, outer ones smaller, cover. 2-valved 1-cell-ed capsule; stigm. simple.									
14032 <i>glauicum Pers.</i>	* glaucou3-wi tt.	or	2 jnaju	Y	Spain	1815.	C s.l	Cav?ic. 2.261	
14J33 <i>tomentosum Dunal</i>	tomentose	JL	or	fjl.au	Y	Svtbtm	scalp.	C 8.1 Eng bot 2208	
14034 <i>lavanduliflum Dm.</i>	Lavunder-lvd	tt.	or	J jn.jl	S. France	IS17.	C 8.p	Bar. ic. 288	
14035 <i>rtechadifT.lium Pers.</i>	Stoechas-lvd	tt.	or	J my.jl	Spain	1810.	C s.p	Sw.cis icin.	
14036 <i>scrylliflum Mil.</i>	Wild Tliymie-lvd	tt.	or	my.s	Austria	1731.	C pi	Sw. cis. 60	
1W37 <i>vulare Gae.</i>	common dwarf	JL	or	i my.s	Britain	gra.ban.	C 6.1	Eng. bot. 1321	
2 flore pKno	double-flwd	JL	or	* J	...	C s.l	Sw. cis. 64		
14038 <i>nudiculae Lag.</i>	naked-stmd	JL	or	J jn.jl	Spain	1826.	C sp		
14039 <i>Ieptophyllum2)/na/</i>	fine-leaved	JU	or	J jn.jl	Span	1818.	C s.p	Sw. cis. 20	
14040 <i>acunnn itum Pers.</i>	acuminat	JU	or	i jn.jl	S. Europe	1810.	C s.p	Viv fr. IS. 14.1	
14041 <i>ovatum Dunal</i>	ovate	JL	or	J jn.jl	Geneva	1818.	C s.p	Viv fr. 1.8. a	
14042 <i>obseCirum Pers.</i>	obscure	JL	or	i my.au	Europe	1816.	B.p	Sw. cis. ic. in.	
14043 <i>nummuririum Mil.</i>	Mone> wort-lvd	JL	or	jn.au	Spain	1752.	E 8.1	Cav. ic. 2.14->	
14044 <i>gymnophyllum Mil.</i>	Surrey	JL	or	J jn.jl	England	cli.hil.	C 8.1	Eng. bot. 2207	
14045 <i>axilliflum Ten.</i>	Hyssop-lvd	JU	or	t rny.jn	Italy	...	C s.l		
2 cyprium Swt.	copper-colored	JL	or	f my.jn	Cop	Naples	...		
14046 <i>hirtum Prrs.</i>	double-flwd	JL	or	* my.jn	Cop	Italy	...		
14047 <i>Ligulc>r Dunal</i>	hairy-calyxcd	tt. —J or	1 jn.jl	Y	Spain	175P.	C 8.1	Bar. ic. 488	
14048 <i>atrcum Pers.</i>	goldon	JL	or	S jn.au	Y	Spain	162d	L	
						Montytier	...		

14049 obtusifolium DIIMI/	blunt-leaved	tt.	or	1	jn.jl	Y	Cyprus	1739.	C	s.l		
14030 angustifolium Pers.	narrow-leaved	JL	or	f	[jn.jl]	Y		1800.	C	s.l	Jac. Tin. 3.53
14051 majoramifoliumta? Marjoram-lvd JU	or	1	lwyj]}	Y	w	France	1818.	C	s.p			
14058 enosēpalon Swt.	woolly-sepaled	1*	or			S. Europe	...	C	s.p	Sw.cis.76		
14059 grandiforum Dec.	large-flowered	1*	or	1	Jn.jl	Y	Italy	1800.	C	s.p	Scop. car. 25	
14054 asperum Lag.	rough	JL	or		jn.au	W	Spain	1827	C	s.f	Sw. cis. 49	
14055 pilosum Pef.	hairy	J-	pr	14	myjn	W	S. France	1731.	C	s.l	Bot. man. 1803	
14056 jampsuecumfrfV. Saminsuch.-lvd fU	or	I			W	France	...	C	p.l	Sw. cis. 2U		
14057 pulverulentum Dec.	powdered	JU	or	I		W	France	...	C	al	Eng. bot. 1322	
14058 polifilium Dec.	Palm-leaved	*.	or	I	wi	W	England downs.	C	s.p	Cav. ic. 2.147		
14039 violaccum Pers.	violet	tt.	or	1	jn.jl	W	Spain	1826.	C	s.p	Cav. ic. 3.263.2	
14020 strictum Pers.	strict	tt.	or	1	jn.jl	W	Spain	1820.	C	s.p	Sw. cis. 48	
14061 linetra Pers."	linear	tt.	or	1	j>jl	W	S. Europe	1818.	C	s.p		
14062 virgMum Pers.	twiggy	I	or	I	my.jl	W	S. Europe	1818.	C	sp	Jac. ic. 1. 98	
14063 fec'tidum Pcv*.	fetid	fl.	or	I	my.j.	W		1800.	C	s.p	Sw. cis. 62
14064 apem'mum Pers.	Awnmine	n.	or	i	my.jl	W	Italy	1731.	C	s.p	Sw. cis. ic. in.	
14065 papyraceum Swt.	↓\v>ct.white	I	or	2	jn.au	W	Europe	...	C	s.p		
14066 hispidum Dwfur/	tispid	I	or	I	my.jl	W	S. Europe	1816.	C	s.p	Sw. cis. 38	
14067 variegatum Swt.	variegated	I	or	*	my.jl	11.w	S. Europe	...	C	s.p	Desf.at.109	
14068 ciliatum i>m.	ciliated	I	or	1	my.jl	R	EuroKJ	182ft.	C	sp	Sw.cis.7	
14069 rhodanthum Dunal	red-flowered	I	or	ft	my.jl	R	Spain	1800.	O	s.p	Sw. cis. 51	
14070 canescens Swt.	canescent	JU	or	i	I	R	S. Europe	1800.	C	s.p		
rbscum Jar. via.			or	1	in'fl	R.W	S. Europe	1800.	C	s.p	Sw. cis. 26	
14071 versicolor Siot.	party-colored	tt.	or	4	jn.ll	R	S. Europe	1800.	C	s.p	Sw.cis.10	
14072 venustum Swt.	beautiful	JL	or	I	my.jl	R	S. Europe	I.K.	C	s.p		
2 fibre ple-no	double-flwd	JU	or	*	W!	Y	Spain	1800.	C	si>	Sw.cis.53	
14073 crocodim i>r\$.	conper-co/a/vd	JU	or	5	jhj	P.Y	France	1795.	C	81	Sw. cU.37	
14074 sulphuruum IV. en.	su'ph\T-colored	I	or	4	jn.au	RV	S. Europe	...	C	81	Jac. ic. 1.99	
14075 flut&b'l. Pers.	changeable	tt.	or	>i•I.	Pk	Spain	1815.	C	s.p	Sw. cis. 55		
14076 roseum Dec.	roseate	1*										
1591. SWA'RTZ/i l W.	SWARTZIA	(Olof Swartz, a celebrated botanist.)					Legum.	Swartxice.	3.—li.			
14077 pinnata IV.	* 3 o r 6 ... >V						Trinidad'	1820V	C	p.l	An. mu. 9.24.2	
14078 grandiflora W.	large-flowered	I	or	I	Y	Trinidad	1821.	C	p.l	Vahlcc.1.3		
14079 simplicifolia W.	simple-leaved	I	or	I	W	W. Indies	1818.	C				

CLASS XIII. OR>ER '2.

POLYAN'DRIA DIGYNIA.

1592. BAU'ER J* H.K. BAHBRA. (*Francis and Ferdinand Bauer, celeb. Ger.bot. draughtsm.*) *Cunoniaceae.* 2.
14080 rubisBfMia//. JC. Madder-leaved * i-J pr 1^jJU Pk N. S. W. 1793. (T s.p Bot. mag. 715
14081 htunilis Lk. dwarf «i_J or 1 jl.d R N. S. W. 1804. C sp Bot.cab.ll<J7

1593. FOTHERGI'LL^ L. FOTIIERGILLA. (*John Fothergill, M.D., a patron of botany.*) *Hamamalltea.* 4.
14082 alnifolia /,. Alder-leaved tt or 4 ap,jj W N. Amer. 1755. L s.p But. mag. 1341
14083 major B. M. larger-leaved *fe or 4 my.jn W N. Amer. 1765. L s.p Bot. mag. 1342
14084 Gardtni Jac. Garden's ft or 4 my.jn W K Amer. 1765. L s.p Jac. ic. 1.100
14085 serftina B. M. late a or 4 au W N. Amer. 1755. L s.p

1594. CURATE'LLA L. CURATELLA. (*Kureuo, to shave j leaves polish bows in Guiana.*) *DULenitcear.* 2.
14086 american'a L. American H O or 8 ... W S. Amer. ... L s.p Aub. gu'i. 1.232
14087 alata Ven. winged-^io/crfil Q or 8 ... W Guiana ... L s.p

1595. PLEURA'NDRA Lab. PLEURANOKA. (*Pleura, rib, aner, anther; stamens*) *DULenitce(e.6.—8S.*
14088 bracteata R. Br. bracteate IJ or 2 my.jn Y N. Holl. 1813. C s.p Dele&s. 1.78
14089 acicularis Lab. needle-leaved II or 2 my.jn Y N. Holl. 1822. C Bp Lab. n. h. 2.14+
14090 strleta B. Br. strict H or 2 my.jn Y N. Holl. 1826. C s.p
14091 Cheorum Can. Garland-flower II or 2 my.jn Y N. Holl. 1824. C s.p
14092 calycina Cun. /ar/rr-calyxed H or 2 my.jn Y N. Holl. 1826. C s.p
14093 ericifolia Cun. Heath-leaved IIJ or 2 my.jn Y N. Holl. 1824. C s.p

*1596. P&OTXIA L. REO.W. (*Paon, a physician who first used it in medicine.*) *Ranunculticce.* 19.
1. MOI/TAN. — Shrubby ; disk of A. expanded into a manbranaceous pitcher more or less inydvine carvels
14094 Moutan H. K. Chinese Tree U or 3 ap,jn P China 1789. C p.l
1 paJJaverdeca Papny,ftotvcrd 9a or 3 ap,jn W China 1789. C p.l Bot cab. 547
iBk Banks's Sk or S ap,jn P China 1789. C p.l Bot.mag. 1154
3 rosea roseate It or 3 ap,jn Pk China — p.l
4 Annels&i Annscley's or 3 ap,jn Pk China — p.l Hort tr. 6.7
IA h % iV* — herbaceous i disk of JL scarcely or not expanded, surrounding only so much of base of carpels.
14095 abiff?ra Pall. white.flowered ft A or 2 my.jn W Siberia 1784. R s.l
1 vestMis Ander. virgin ft A or 2 my.jn W Siberia ... R s 1 Bot. rep. 64
2 Candida Ander. white * A or 2 my.jn F Siberia ... R s.l
3 tartarica Tartarian X A or 2 my.jn F Siberia ... R si
4 sibirica Siberian ft A or 2 my.jn W Siberia ... R s.l
5 rubescens rubescent ft A or 2 my.jn Pk Siberia 1784. R s.l Bot. reg. 42
6 uniflra single-flowered ft A or 2 my.jn R Siberia ... R al
7 Whittleji Vlhittey's double whiter A or 2 my.jn Bh China 1784. R s.l Bot. rep. 612
8 Humei Hume's double crimson I. 2 my.jn R China 1784. R s.l Bot. mag. 1768
9 fragrans fragrant I. 2 my.jn R China 1784. II s.l Hort. tr. c. ic.
14096 triternMa Pall. triternate I. 3 my.jn P Siberia 1790. R 8.1 Bot. mag. 1441
dafrica Andr.
14097 corallina Retz. coralline I. 3 my.jn R England ... R si Eng. bot. 1513
14098 officinalis Retz. officinal 3 my.jn R Switzerl. 1548. R s.l Bot. mag. 1784
2 rosea roseate I. 3 my.jn R ... K 61
3 blanda bland I. 3 my.jn Pk ... R s.l
4 rubra double red I. 3 im.jn It ... tt s.l

5 carnescens	carnescent	J. or	3 my.jn	W	—	—	R 61
6 albicans	whitening	J. A or	3 my.jn	W	—	—	R si
14099 lobata Desf.	lobed	J. A or	2 my.jn	R	Spain	... R 8.1	Sw. fl. gar. 70
14100 anomala W.	anomalous	J. A or	1* my.jn	Pk	Siberia	178a R 8.1	Bot. mag. 1754
laciniata <i>Pal. ros.</i> 2.85							
14101 hibrida <i>Pall.</i>	hybrid	ft A or	2 my.jn	R	Siberia	1788. R 8.1	Pal. ros. 2.86
14102 tenuifolia L.	fine-leaved	t< A or	1 my.jn	R	Siberia	1765. R si	Bot. mag. 926
i. PUBESCB'NTES. — Same as the preceding, but with the leaves underneath pubescent, not smooth.							
14103 peregrina Mil.	foreign	ft A or	3 my.jn	D.P	Levant	1629. R 61	Bot. mag. 1050
2 Grévillei	Greville's	ft A or	2 my.jn	P		... R s1	
3 compacta	compact	ft A or	2 my.jn	P		... R 8.1	
14104 critica Lindl.	Cretan	ft A or	2 my.jn	P	Candia	... R 8.1	Bot. reg. 819
14105 paradoxia Ander.	paradoxical	t< A or	2 my.jn	P	Levant	... R s1	
2 fimbriata	double fringed	if A or	2 my.jn	P		... R 8.1	Sw. fl. gar. 19
tartarica Mil.		a					R SI
14106 mullis Ander.	soft	i(A or	H my.jn	P		... R SI	Bot. cab. 1263
14107 arietina Ander.	ram	i(A or	2 my.jn	P		... R 8.1	
14108 debrca Ander.	comely	i(A or	2 my.jn	P		... R 8.1	
1 Pali&ctf	Pallas's	i(A or	2 my.jn	P		... R SI	
2 elatior	taller	i(A or	2 my.jn	P		... R s1	
14109 htimilis Retz.	dwarf	i(A or	2 my.jn	P	Spain	1613. R 61	Bot. mag. 1422
14110 Rdssi Biv.	Ru&s's	i(A or	2 my.jn	C	Sicily	... R SI	Sw. fl. gar. 122
14111 plibens B. M.	downy	i(A or	2 my.jn	R		... R s1	Bot. mag. 2L/4
14112 villosa Swt.	villous	i(A or	2 my.jn	R	S. Europe	1816. R 8.1	Sw. fl. gar. Uo
acssiliflra Sims		a					

CLASS XIII. ORDER 3.

POLYANDRIA TRIGYNIA.

* 1597. HIBBE'R'T/i* Andr. HIUBEBTIA. (<i>George Hibbert, a disting. collect. of plants.</i>) <i>Dillenidcea*</i> . 9.—19.
14113 volubilis B. Rep. twining % \ or 8 my.o Y N. S. W. 1790. C s.p Bot. rep. 126
14114 grossiliar? fulu Sal. (gooseberry-lvd footed f J or 6 mr.au Y N. Holl. 1803. C s.p Bot. mag. 1218
14115 dentata It. Br. twiggy 3 jn.au Y N. Holl. 1816. C Bp Bot. reg. 282
14116 traltna if. Br. WiUow-leaved 3 jn.au Y N. Holl. 1823. C sp
14117 virgata T. J?r. twiggy 3 jn.au Y N. Holl. 1822. C e.p
14118 lineans R. Br. linear 2 jn.au Y N. Holl. 1821. C s.p
14119 obtusifolia R. Br. obtuse-leaved 2 jn.au Y N. Holl. 18124 C s.p
14120 pedunculata R. Br. pedunculated H 1_I or 2 my.au Y N. Holl. 1821. C s.p Bot. reg. 1001
corifolia Bot. mag. 2672
14121 cistifolia R. Br. Cistus-leaved * I_I or 1 my.au Y N. Holl. 1826. C s.p

1598. HOMA^LIUM Jac. HOMALIUM. (<i>Homa/os, equal; stamens.</i>) <i>Homallnea</i> . 1.—2.
14122 racemosum Jac. racemose * Q or 6 my jl W W. Indies 1816. C 8.1 Bot. cab. 261

*1599. DELPHI'NIUM Tou. LARKSPUR. (*Delphin, dolphinj supp. resembl. in nectary.*) *Ranuncul.* 45.—58.

I. CONH6LIDA. — Ovary one ; petals four joined into one. Annual.
14123 Oliverium Dec. Oliver's O or B S. Europe 1826. S co Deless. 1.51
14124 Ajhcis L. Ajax's O or U Pk Switzerland 157a S r.m
2 flr&rc pflno double-flowered Q or 1 jnjl Va Europe 1573. S co Cl. h. 2.2
14125 Consthda L. Coinfrey O or 2 jnjl B England sail. fl. S r.m Eng. bot. c. ic.
2 flr pK-no double-flowered Q or 2 jnjl Va England ... S co Bcs.eys. 2.14.1
14126 pubebcens Dec. pubescent O or 2 jnjl B Mediterr. 1816. S co Fl.gr. 504
ambiguum Lot.
14127 AcovC\ A. Aconite-like O or 1 jnjl P Levant 1801. ■ Vahl sy. 1.13
14128 peregrinum W. foreign O or 1 jnjl B Italy 1629. ■ Al.ped. 2.25.3
,,uncum Dec.

II. DELPHINE'LLUM. — Ovaries three j petals free, smooth. Annual
14129 ambiguum L. doubtful O or 2 jl.au B Barbary 1759. S P-I
14130 cardiopetalum Dec. heart-petaled O or 1 jnjl B Pyrenees 1818. S co Mat com. 557
14131 virgatum Poir. twiggy O or U jnjl B Syria 1823. S co Deless. 1.55
III. DELPHINA'BTRUM. — Ovar. 3 or 5; petals free, lower ones bearded in disk, entire, from No.HlCo. bifid.
14132 grandiflrum L. great-fldwered D.B Siberia 1816. D pl Bot. mag. 1686
2 fl&rc pflno double-flowered D.B Siberia D pl
3 album white-Jowlered W gardens 1816. D co
4 p&hdum pale-blue B gardens 1810. D co
14133 chmnae Pu. China D.B China 1818. D co Bot cab. 71
14134 cheilanthurum Fis. lip-flowered D.B Siberia 1819. D pl Bot. reg. 473
14135 pseudo-peregrinum Fis. bast, foreign R Siberia 18123. D co
14136 puniceum Pall. scarXet-flowered P Siberia 1785. D pl
14137 albilfltrum Dec. white-flowered * W Armenia 1823. D co Deless. 1.58
14138 ochrolucum Stcv. yellowish white 'W Iberia 1823. D pl
14139 h^bridum IV. hybrid B Siberia 1794. D pl
hirsbtum Pers.
14140 fissum W.&K. cleft ■ Hungary 1816. D co W.&K. 1.81
14141 velutinum Bert. velvety ■ Italy 1819. D co
14142 pentngnum Lam. five-styled ■ S. Europe 1819. D co Desf. at. 1.11*
14143 legans Dec. elegant ■ N. Amer. ... D co
2 fibre pleno common double-flwd ■ N. Amcr. 1741. D co Fl. con. 43
14144 amoëVium Stev. pleasing blue 2 jl.au Pa.B Siberia 1818. D co Gin. si. 4.77
14145 spuxium Fis. spurious 4 jl.au B Siberia 1810. D co
14146 Iric6rneJtf*. three-horned Jjl au B N. Amer. 1806. D pl Bot. cab. 506
14147 asfrcrum Mx. azure 6 jl.au L.B Carolina 1805. D pl
14148 exaltatum H. K. exalted 3 jl.au B N. Araer. 1758. D pi Mil. ic. 27tt 2
14149 urcadatum Jac. urceolate ■ ■ ■ 1801. D pl Bot mag. 17.1
14150 flexu&sum Bieb. flexuous ■ ■ ■ Caucasus 182a D pl
14151 revolutum Desf. revolute ■ ■ ■ ■ D pl

14152	mesoleftcum	<i>Lk.</i>	white-middle	^ A or	3	jinjl	B.w	—	1822.	D	p.l	Ficon.	£5
14153	palmitidum	<i>Dec.</i>	palmate-cleft	:fc A or	3	llau	B	Siberia	1824.	D	co	Gm.si.	4.79
	2	<i>glabellum Dec.</i>	Broothish	ik A	3	jl.au	B	Siberia	1817.	D	co	Gin.	si. 4.75
14154	intermedium	//. <i>K.</i>	intermediate	& A or	8	jl	B	Silesia	1710.	D	p.l		
1	<i>pilosissimum Dec.</i>	hairiest	^ A or	9	jl	B	Siberia	—	D	co	Gm.	8i. 4.80	
2	<i>leptostachyum Dec.</i>	slender-spiked	jg A or	6	jl	B	Pyrenees	■	D	co			
	intermedium	<i>Lap.</i>											
3	<i>ranunculiflbum Dec.</i>	Ranunc.-lvd	^ A or	6	jl	B	Pyrenees	...	D	co			
4	<i>luxum Dec.</i>	loose-spiked	^ A or	6	jl	B			D	co	Mil.	ic.	119
14155	eUktum	<i>IV.</i>	tall common Bee	^ A or	6	jns	B	Siberia	1597.	D	p.l	Schk.han.	2.145
14156	alp'inum	<i>IV.</i> \$ <i>K.</i>	alpine Bee	& A or	5	jlau	B	Hungary	1816.	D	co	W & K.	a 246
14157	cuneatum	<i>Stev.</i>	wedge-leaved	^ A r	4	jinjl	B	Siberia	1816.	D	co	Bot.rcg.	327
14158	villsum	<i>Stcv.</i>	villous	^ A or	4	jl.au	B	Caucasus	1818.	D	co		
14159	dictyocarpum	<i>Dec.</i>	netted-fruited	^ A or	4	jinjl	B	Siberia	1817.	D	p.l		
1	(160)	<i>laxiflrum Dec.</i>	loose-flowered	^ A or	3	jinjl	B	Siberia	...	D	pi		
14161	montanum	<i>Dec.</i>	mountain	31 A or	4	jl au	B	Switzerl.	1819.	D	co		
2	<i>bractcusum Dec.</i>	bractose	^ A or	8	jl.au	B	S. Europe	1816.	D	co			
14162	dasycarpum	<i>Stev.</i>	thick-fruited	^ A or	4	jlau	B	Caucasus	1819.	D	co		
14163	speciosum	<i>Bieb.</i>	showy	31 A or	4	jlau	B	Caucasus	1816.	D	co	Deless.	1.62
14164	triste	<i>Fis.</i>	sad	& A or	2	jlau	B	Dahuria	1819.	D	co		
IV. STAPHISA ^X GRIA —Ovaries 3 or 5 j petals free; spur short; capsules ventricose.													
14165	Requienii	<i>Dec.</i>	Requin's	^ CD or	4	my.jn	L.B	Majorca	1824.	S	co	Dleess.	1.63
14166	pictum	<i>W. en.</i>	painted	^ Q or	1	ap.apu	L.B	S. Europe	1816.	S	s.p	Wood	151
14167	Staphiskgria	<i>L.</i>	Stavesacre	^ Q or	2	ap.apu	L.B	S. Europe	1596.	S	s.p	FLgr.	508

***inOO.** *iiCONITUM Tou.* **ACOXITUV.** (*Plentiful about Acona in Bithynia.*) **Ranunculhcete.** **109.—**

I. A'NTIIORA. — *Sepals persistent; helmet conical or semicircular; ovaries 5; flowers cream-colored, sometimes variegated with blue; roots napiforme /, leavet palmately cut into linear lobes.*

14168	A'nthora L.	Anthora	"	A or	Ujn.au	Pa.Y	Pyrenees	1590.	R co	Jac. au. 4. 382
14169	grandiflorum Rchb.	largo-flowered	"	A or	11. jn.au	Pa.Y	Jura	1821.	R co	Rchb. ac. 1. B
14170	Decandellia Kchb.	Dcaidollc'i	"	A or	12. jn.au	Pa.Y	Siberia	1821.	R co	Rchb. ac. 3
14171	anthorofdeum/teAft.	Anthora-like	&	A or	13. jn.au	Pa.Y	Jura	1821.	R co	Rchb. ac. 4
	inchinatum Ser.									
14172	Pallas's/* Ilrbh.	Pallas's	i	A or	2. jn.au	Pa.Y	Siberia	1821.	R co	Rchb.ac. 6.A. B
14173	ciliophorum Rchb.	well-crested	k	A or	1f. jn.au	Pa.Y	Caucasus	1821.	R co	Rchb. ac. 5
14174	Jacquinet Rchb.	Jacquin's	t	A or	1J. jn.au	Pa.Y	Austria	1800.	R co	Jac. au. 4.382
14175	nemorosum Bieb.	grove	ff	A or	2. jlau	Pa.Y	Caucasus	1823.	R co	Rchb. ac. 6
14176	versicolor Stcv.	various-colored	^	A or	3. jn.s	B.v	Siberia	1820.	R co	Bot. cab. 794
II. LYC6CTONUM. — <i>Sepals deciduous; helmet conico-cylindrical; ovaries 3; flowers cream-colored or voluite, rarely purple or variegated; lobes of leaves cuneiform, pinnate, rarely bipinnate.</i>										
14177	lyctctonum E.	woofs-bane	"	or	3. juu	P	Alp. Euf.	1596.	D & co	Rchb.ac.5S
14178	Vulparia Rchb.	fox-fane	"		3. jlau	Pa.Y	Alp. Eur.	1821.	D co	Jac. au. 4. 38J
14179	cynictonium Rchb.	tall dog's-bane	"		3. jlau	Pa.Y	France	1820.	D co	
	altissimum Dec.									
14180	laxiflbrum Dec.	loose-flowered	£	A or	3. jlau	Pa.Y	Switzerl.	1823.	D co	
14181	rubicundum Fis.	reddish	&	A or	2ijl.au	P	Siberia	1819.	D co	
14182	carpathicum Ser.	Carpathian	£	A or	2i. jlau	P	Carpa.mo.	1810.	D co	Bot. mag. 2196
H183	auBtale Rchb	southern	£	A or	2ijl.au	P	Denmark	1821.	D co	Fl. dan. 123
14184	lupicidum Rchb.	wolPs bane	3f	A or	2. jlau	Pa.Y	Europe	1821.	D co	Rchb. ac. 54
14183	tragbctonum Rchb.	gnat's-bane	3l	A or	2} jn.au	Pa.Y	Switzerl.	1822.	D co	
14186	thriophorum JfcAA.	beast's-bane	3i	A or	2J. jn.au	Pa.Y	Europe	1824.	D co	-
14187	rectum Rchb.	upright	^	A or	3. jn.au	Pa.Y	Europe	1824.	D co	
1 st 188	pyrenaeicum L.	Pyrenean	JH	A or	4. injl	Y ,	Pyrenees	1739.	D co	Rchb. ac. 48
14189	album H. K.	white	ik	A or	4. jlau	W	Levant	1752.	D co	
14190	barbatum Pat.	bearded	3i	A or	2. injl	Pa.Y	Siberia	1807.	D co	Deless.1.61
14191	hispidum Dec.	hispid	^	A or	2. injl	Pa.Y	Siberia	1823.	D co	
14192	septentrionale TV.	northern	3t	A or	4. jlau	B	N. Europe/SOO.	D co		
14193	ochroleucum JV.	yellowish whitest	A	or	3. jn.au	L.Y	Caucasus	1794.	D co	
14194	macrophyllum/ior/.	large-leaved	k	A or	2. jlau	Pa.Y,			D co	

III. CA'MMARUM.—*Sepals deciduous; helmet conical, compressed; ovaries 3 or 5; flowers blue, white, or variegated, rarely flesh-colored; roots tuberous; lobes of leaves trapeziform pinnate.*

14195	<i>variegatum</i> L.	variegated	A A or	5 jn, au	P, w	S. Europe	1597.	R co	Ser.heM5.18.13
2	bicolor <i>Rchb.</i>	two-colored	A A or	4 jl.au	W, B	Switzerl.	1821.	R co	Rchb. ac.31
3	cieriMeum <i>Ser.</i>	blue	A A or	4 jl.au	B	Switzerl.	1819.	R co	
4	albiflurum <i>Ser.</i>	white-flowered	A A or	4 jl.au	W	Switzerl.	1819.	R co	
14196	<i>rychnanthum</i> <i>Rchb.</i>	bilFlowered	A A or	4 jl.au	P.B	Switzerl.	1821.	R co	
2	bicolor <i>Rchb.</i>	two-colored	A A or	4 jl.au	W.B	Switzerl.	1819.	R co	Cl. h. 2.98.1
14197	<i>hamktum</i> <i>Rchb.</i>	hooked	A A or	4 jl.au	P.B	Italy	1810.	R co	Ser. hel. 1.15.12
14198	<i>gibbusum</i> <i>Ser.</i>	gibbose	A A or	1 jl.au	B	Caucasus	1818.	R co	Ser. hel. 1.15.14,15
14199	<i>gracile</i> <i>Rchb.</i>	slender	A A or	1 jl.au	B	Switzerl.	1821.	R co	Rchb. ac. 7
14200	<i>Cammarum</i> <i>Jac.</i>	Cammarum	A A or	1 jn.s	P	Austria	1821.	R co	Rchb. ac. W
14201	<i>rostratum</i> <i>Bernh.</i>	beaked	A A or	1 jn.au	B	Switzerl.	1819.	R co	Bot cab. 203
2	<i>pilosasculum</i> <i>Ser.</i>	rather hairier	A A or	3 jn.au	P	Carpa.rao.	1800.	R co	
<i>AnM</i>									
	<i>Cammarum</i> <i>Wahl.</i>								
14202	<i>heburgynum</i> <i>Dec.</i>	blunt-styled	A A or	3 jn.au	P	Switzerl.	1819.	R co	Ser. heL 1.15.18,19
2	<i>multifidum</i> <i>Schl.</i>	many-cleft	A A or	3 jn.au	P	SwitzerL	1819.	R co	
14203	<i>nasatum</i> <i>Fis.</i>	great-nosed	A A or	3 jn.au	P	Siberia	1818.	R co	Rchb. ac. 10
J	paniceum	panicled	A A or	3 jn.s	Pa.B	France	1815.	R co	Rchb. ac. 32
	Rchb.	acuminate	A A or	3 jn.au	B	Switzerl.	1819.	R co	
14506	<i>lacimbum</i> <i>Rchb.</i>	jagged	A A or	3 jnau	B	Switzerl.	1820.	R co	
14207	<i>cernuum</i> <i>Rchb.</i>	drooping	A A or	3 jn.au	B	Switzerl.	1800.	R co	Rchb. ac. 33
2	<i>pacifilbrum</i>	few-flowered	A A or	3 jn.au	B	Switzerl.	1821.	R co	
3	<i>rambsum</i>	branchy	A A or	3 jn.au	B	Switzerl.	1821.	R co	
4	<i>flexicaale</i>	zigzag-stemmed	A A or	1 jn.au	B	SwitzerL	1819.	R co	
14208	<i>CGessi</i> <i>Rchb.</i>	<i>Clusius</i> s	A A or	1 jn.au	B	Switzerl.	1819.	R co	Cl h.2.97.1
14209	<i>Florkeanw</i> <i>Rchb.</i>	Flbrke's	A A or	3 jn.au	B	SwitzerL	1821.	R co	Stoer.stra.69.ic.
2	bicolor <i>Rchb.</i>	two-colored	A A or	3 jn.au	B.w	Switzerl.	1801.	R co	
14210	<i>japonicum</i> <i>Thun.</i>	Japan	A A or	6 ls	F	Japan	1790.	R co	
2	<i>csmMeum</i>	azure	A A or	4 jl.au	B	Japan '	1700.	R co	
14211	<i>uncinatum</i> <i>L.</i>	hooked	A A or	2 jl.au	B	N. Amer.	1768.	R co	Rchb. ac. 35
2	<i>Michauxidniwi</i> <i>Ser.</i>	Michaux's	A A or	2 jl.au	B	N. Amer.	1800.	R co	Bot. mag. 1119
14212	<i>Funkii</i> <i>Rchb.</i>	Funkie's	3 jnjl	B	Switzerl.	1825.	R co		

14213 Meyer; Rchb.	Meyer**	A A or	3. jn.j	B	Switzerl.	1825.	K co	
14214 nitidum Fis.	shining	A A or	3 jn.jl	B	Switzerl.	1825.	R co	
14215 plicatum Kolle	plaited	A A or	3 jn.jl	B	Switzerl.	1825.	K co	
14216 rigidum Ilchb.	rigid	A A P	3 jiujl	R	Switzerl.	1825.	R co	
14217 zöbconum Rchb.	poisonous	A A P _o	3 jn.jl	B	Switzerl.	1823.	R co	
14218 toxicum Rchb.	>< poisonous	A A or	3 jn.jl	B	Switzerl.	1821.	R co	Rchb. ac. 37
14219 umbrotum SchL	shady	A A or	3 jn.jl	B	Switzerl.	1825.	R co	
14220 albidum Rchb.	whitish	A A or	-	W	Europe	1824.	R co	
14221 oligocarpum Rchb.	few-podded	A A or	-	B	Europe	1823.	R co	
14222 neomontnum L.	new mountain	A A or	3 jl.au	B	Europe	1799.	R co	Bar ic. 610
14223 Bpeculus Otto	showy	A A or	3 jl.au	B	- -	1823.	R co	Deless. 1.54
14224 tortifsum IV. en.	twisting	A A or	6 jl.au	P.B	- -	1812.	R co	Rchb. ac. 24
14225 Gmerini lclib.	Gmelin's	A A or	4 jn.au	B	Siberia	1821.	R co	Rchb. ac. 40
14226 illinotum Rchb.	anointed	A A or	4 jl.au	B	- -	1821.	R co	
14227 Schlcicheri Rchb.	Schleicher's	A A or	2 jn.jl	B	Switzerl.	1821.	R co	Rchb. ac. 1. 1
	Napellus yy pubescens Dec.							
14228 Bernhoxiuum Rchb.	liernhardti's	A A or	1 jn.jl	B	Europe	-	R co	
14229 flacidum Rchb.	flacid	A A or	5 jl.au	B	Siberia	1822.	R co	Ser. hel.J. 15.30
14230 maximum Poll.	largest	A A or	6 jl.au	B	Kamtsch.	1823.	R co	Rchb. ac. 17
14231 productum Ilchb.	long.lipped	A A or	1 jl.au	B	Siberia	1821.	R co	Rchb. ac. 38
14232 inolle Rchb.	Soft	A A or	5 jl.au	B	- -	1820.	R co	Rchb. ac. 31
14233 exaltatum Bergh.	exalted	A A or	6 jl.au	B	Siberia	1819.	R co	Ser.hel.l. 1537, 1 st s
14234 intermedium Dec.	intermediate	A A or	5 jl.au	B	- -	1820.	R co	
IV. NAPE I.LI/9.—Sepals decubous; helmet semicircular, rarely navicular j ovaries 3,5, or 7, rarely pilose, + flowers blue and white or variegated; lobes of leaves cuneate bipinnate; roots tuberous.								
14235 Nancifus L.	Monk's-hood	A A ls ^r	myjl	■	Europe	15IXJ	R co	Rchb. ac. 2
2 fibbus	white-flowered	A A or	-	■	Switzerl.	1819.	R co	
3 rubellus	small red-Jlivd	A A or	-	■	Switzerl.	1819.	R co	
14236 K ^{DE} Vm/m Rchb.	Kelle's	A A or	-	■	S. Europe	1820.	R co	Rchb. ac. 11. 1
2 pygime'uin Rchb.	pxgmy	A A <r	-	■	S. Europe	1822.	R co	Rchb. ac. 12.1
14207 cominutivurn Rchb.	changed	A A or	-	■	S. Europe	1823.	R co	Rchb. ac. 18.3
14238 callybdtryon Rchb.	inne-racemed	A A or	-	■	S. Europe	... R co		Ilchb. ac. 16 1
14-39 Hällerj Rchb.	Haller's	A A or	-	■	S. Europe	1821.	R co	
2 bicolor Rchb.	two-colored	A A or	-	■	Switzerl.	1820.	R co	
14240 virgatum Rchb.	Iwiggy	A A or	-	■	S. Europe	1822.	R co	
14241 hians Rchb.	gaping	A A or	-	■	S. Europe	1823.	R co	
14242 amce'iium Rchb.	pleading blue	A A or	-	■	S. Europe	... R co		Rchb. ac. 14.1
14243 Ilöpppt Rchb.	Hopp's	A A or	-	■	Carinthia	1823.	R co	
14241 smigale-Mum Pall.	half-helmeted	A A or	-	■	Siberia	1818.	R co	Rchb. ac. 41
14245 inclutonum Rchb.	badger's-banc	A A or	-	■	- -	1821.	R co	
14240 formosum Rchb.	handsome	A s or	-	■	S. Europe	1824.	R co	Rchb. ac. 18.1
14247 acutum Rchb.	acute-leaved	S* A or	-	■	S. Europe	1821.	R co	Rchb. ac. 14.2
14248 neubergense Dec.	Neuberg	A A or	-	■	S. Europe	1822.	R co	Rchb. ac. 17.1
	strctrum Rchb.							
14249 laxum Rchb.	loose	A A or	-	■	S. Europe	1820.	R co	Rchb. ac. 15.4
14231 ls ^x tum Rchb.	joyful	A A or	-	■	S. Europe	1821.	R co	Rchb. ac. 13.2
14251 squarrifsum Rchb.	squarrose	A A or	-	■	Siberia	1822.	R co	
14252 elvitum Rchb.	tall	A A or	-	■	Europe	1822.	R co	
14253 angustifolium Rchb	narrow-leaved	A A or	-	■	Europe	1824.	R co	
14254 pubescens Moen.	pubescent	A A or	-	■	Europe	1824.	R co	
14255 eufcRhyon Rchb.	well-spiked	A A or	-	■	Europe	1824.	R co	Rchb. ac. 15. 3
14256 clphiniifiliumAc/#A.l)clphiniium-lvdA	A or	-	-	■	N. Amer.	1820.	R co	Rchb. ac. 42
14257 Kfhlen Rchb.	K'd'hler's	A A or	-	■	Europe	1824.	R co	
14258 vendstum Rchb.	beautiful	A A or	-	■	Switzerl.	1823.	R co	
14259 eriostemum Dec.	woolly-stamened	A A or	-	■	Europe	1821.	R co	
14260 WildcnöM Rchb.	Wildcnow's	A A or	-	■	Europe	1823.	R co	
142G1 rigiduim Rchb.	rigid	A A or	-	■	Switzerl.	1825.	R co	Rchb. ac. 7.1,2
2 grandiflurum Fis.	large-flowered	A A or	-	■	Siberia	1825.	R co	
14262 indinctum Koch	anointed	A A □	-	■	Europe	1829.	U	
14263 fminens Koch	eminent	A A or	-	■	Europe	1810.	R co	Ilchb. ac. 15.1
14264 strictum Dec.	strict	A A or	-	■	Siberia	1824.	R co	
14265 galactonum Rchb.	lizard's-bane	A A or	-	■	Hungary	1825.	R co	
1426G leucunthum Rchb.	white-flowered	A A or	-	■	1823.	R co		
14267 Ottoniunum Ilchb.	Otto's	A A or	-	■	Europe	1824.	R co	
14268 recñgnitum Rchb.	recognised	A A or	-	■	1830.	R co		
14269 Sprengelii Rchb.	Sprengel's	A A or	-	■	Europe	1824.	R co	
14270 amphiflurum Rchb.	large-flowered	A A or	-	■	Austria	1823.	R co	Rchb. ac. 8.1
14271 versicolor Rchb.	party-colored	A A or	-	■	Switzerl.	1819.	R co	Bot. cub. 794
14272 decorum Rchb.	neat	A A or	-	■	Pyrenees	1824.	R co	
14273 tauricum W.	Taurian	A A or	-	■	Tauria	1752.	R co	Rchb. ac. 15.4
laxum Rchb.								
14274 volQbile Pall.	twining	A A or	-	■	Siberia	1799.	R co	Rchb. ac. 25
14275 Brahnj Rchb.	Braune's	A A or	jl.au	■	Switzerl.	1821.	R co	
14876 bifl5rum Fis.	two-flowered	A A or	jnjl	■	Siberia	1817.	R co	Kch'. ac. 22

CLASS XIII. ORDER 4.

POLYA'NDRIA TETRAGY'NIA.

1601. CARYCTCAR L.	BUTTER NUT.	(Karyon. & nut: bearing nuts.) Rhnobblea. 3.-6.	
14277 nuciferum L.	nut-bearing	D fr 100" ... Rv Guiana 1820. C l.p Bot.mag. 17	
14278 tomentosum W.	woolly	ft 100 ... W Guiana 1820. C Bl Aub.guL23	
Pekea tuberculata Aub.	smooth	D fr 100" ... Guiana 1820. C s.l Aub.gni.240	
14279 gftbrum Pers.			
14280 WinteriForst^ Wintera aromatica Mur.	Winter's	(Drimys, acrid; bark acrid and aromatic.) I am ... W? Magellan 1827. C s.l.p *orst.42	

CLASS XIII. ORDER 5.

POLYANDRIA PENTAGYNIA.

1603. TRACHYTELLA Dec. TRAWII-TELLA. (*Trachytes*, roughness; leaves.) *Dittenihceet.* 1.—S.
 14281 ~~cta~~^{ta} a Dec. Acta » «. u.j cu 6 ... W China 1823. C p.l
1604. CIMI CI'FUG A L. BUQWOBT. (*Cimex*, a bug, *fueo*, to drive away; sup. qual) *Ranuncutece**. 4.—5.
 14282 fa'tida L. fetid £ A in 4 jn.jl L.Y Siberia 1/77. D p.l Lam. iL 487
^{cta} ~~ta~~ *Cimicifuga* L.
- 14283 americana Mx. American ■ A or 2 jn.jl W.Y Carolina 1824. D p.l Deless. 1.66
^{Acte} ^d podocarpa Dec.
- 14284 cordata Ph. heart-leaved ♀ A cu 3 jn.jl W.Y N. Amer. 1812. D p.l Bot. mag. 2069
- 14285 palmata Mx. palmated ■ A cu 4 j.lau W.Y N. Amer. 1812. D p.l Bot mag. 1630
- 1605. AQUILE^XGIA L. COLUMBINE. (*Aquila*, an eagle; claw-shaped nectaries.) *Ranunculcea*. 15.
 14286 vulgaris L. common ■ A or 2 my.jl B Britain fields. D co Eng. bot. 297
 2 corniculata Zto. smal. horn. doble ■ A or 2 my.jl B.w Europe ... D co Cl. h. 2. 204. 1
 3 inversa Dec. inverted double ^ A or 2 my.jl B.w Europe ... D co Cl. h. 2. 204. 2
 4stellata JDec. starred double ^ A or 2 my.jl B.w Europe ... D co Cl. h. 2. 205. 1
 5 degener Dec. degenerate double ^ A or 2 my.jl B.w Europe ... D co Cl. h. 2. 205. 2
 14287 viscosa Gou. clammy ♀ A or 1J my.jn P Montpel. 1752. D co Gou. il. 19.1
- 14288 sibirica Lam. Siberian ■ A or 1J my.jl B.W Siberia 1806. D co Deless. 1.47
^{vulgaris dahbrica} W.
- 14289 glandulosa Fis. glandular ■ A or 1J my.jl B Siberia 1822. D co
² concolor Dec. self-colored ■ A or 1J my.jl B Siberia 1822. D co
 14290 alpina L. alpine ■ A or 1J my.jl B Siberia 1731. D co Sw fl. gar. 218
- 14291 grandiflora Pat. large-flowered ■ A or 1J my.jl B Siberia 1818. D sp
- 14292 alpina f. grandiflora Dec. Pyrenean ■ A or 1 my.jl B Pyrenees 1818. D fl.p Trev.del.2
- 14293 canadensis L. Canadian ■ A or 1 my.jl R.o rf Amer. 1640. D i.p Bot mag. 246
- 14294 formba Fis. beautiful ■ A or 1 my.jl R.o Kantsch. 1822. D fl.p
- 14295 vindiflora Pall. green-flowered ■ A or 1 my.jl B Siberia 1780. D co Jacic 1.102
- 14296 daviniana Patr. Davunian ■ A or 1 my.jl B Siberia 1827. D co
- 14297 atropurpurea W. en. dark purple ■ A or 1 my.jl B Siberia 1827. D sp Bot reg. 922
² Fisher's Fis. Fisher's ■ A or 1 my.jl B Siberia 1827. D co
- 14298 parviflora Led. small-flowered ■ A or 1 my.jl B Siberia 1819. D 8D Gm.iL 4.74
- 11299 h. jbrida B. M. hybrid ■ A or 1 my.jl B Siberia — D co Bot mag. 1221
^{bicolor} Pen.
- 14300 anemone'ides IV. Anemone-hike ^ A or 1 my.jl P Altai 1827. D «.p
^{glandulosa} var. Fis.
1606. NIGE'LLA Tou. FENNEL FLOWEB. (Mferr, black; seeds.) *Ranunculcea*. 10.—11.
 ... I. EH6BATO9. —Stamens disposed in many series; sepals white or blue; carpels Joined.
- 14301 damascena L. damask O or 14 jn.s LB S. Europe 1570. S co Bot mag. 22
 2fl6repleno double-flowered O or 14 jn.s LB S. Europe 1570. S co CLh.2.208.1
³4302 ~~uarecta~~^{Xta} Gm. compressed O or 14 jn.s W.o S. Europe 1793. S co
- Stamens disposed in many series, collected into 8 or 10 bundles; sepals bluish.
 2 crictica Dec. cultivated O or 14 jn.s LB Egypt 1548. S.l¹ Zo. ic. 119
 3 citrina Dee. Cretan O or 14 jn.s Pa.B Crete ... S co Mat com. 581.4
^{citrina} citron-cld-seeded O or 14 jn.s Pa.B S. Europe ... S co M.h.3.12.18.5
 Indian O or 14 jn.s Pa.B E. Indies ... S co
 corn-field O or 14 jn.s B.w Germany 1683. S sl Schk. han. 2.146
 Spanish O or 14 jn.s B.w Spain 1629. S sl Bot. mag. 1265
^J4307 aniculiacea Dec. ~~J~~ennel-scented O or 14 jn.s B.w Tauria 1824. S co M. h. 3.12.18.6
^J4307 divanoata Seup. var. O or 14 jn.s B.w Egypt 1817. S co Deleas. 1.46
- III. NICELLA'STRUM. —Stamens disposed in one series; sepals yellow.
- 14308 orientalis L. yellow oriental O or 14 jn.s Y Syria 1699. S sl Bot. mag. 4264
- 14309 corniculata Dec. small-horned O or 14 jn.s Y ... 1820. S co
- U310 ciliaria Dec. ciliated O or 14 jn.s Y Levant 1827. S co Deleas. 1.45
- * 161/7. REAMITR.M L. REAMURIA. (Heine A. F. de Reaumur, an entomologist, d. 1757.) *Ficoideee* 1~2
 14311 Aypenc'des W. Hypericum-lk n. J pr 2 jl.o P Syria 1800. C a.p Bot reg. 845
^{/influtia} Sal. //ypericum alterniflum Lab.
1608. COLBE'R17/f Sal. COLBERTIA. (M. Colbert, a botanical patron.) *DULenihcece*. 2.
 *312 coramandellina Sal. Coramandel ■ or 15 mr.ap Y Coraman. 1803. C P.J Rox cor 20
 Dillenir pentagynia IV.
- *4313 scabrida D. Dom. roughish f • or 10 — Y Nepal 1820. C p.l
- IAJ??*-TE TRA'CERA L. TETRACERA. (Tetra, four, Tceras, a horn; capsules.) *Dillenibce**. 3.-23.
 *4314 potat'na Afz. drinking % a or SO S. Leoné 1822. C p.l
- obovata Dec.
 JSiS all? a V' Alder-leaved % • or 12 ... Y Guinea 1793. C p.l
 4316 Jvit? L¹ twining t O o r 12 ... Y S. Amer. 1818. C p.i Gae. ft. i. 69 3
- DiR?ma scandens Horn'A. *Olaus Wormius*, celebrated Danish phys. and nat.) *DiUenidce**. -i 4,
- . 1611. CARPODO'NTOS £«*. CARPODONTOB. (*Karpos*, a fruit, *odonof*, toothed.) *Hyperidne**. 1
 *318 lñcida Lab. shining ii_J or 20. ... N. Holl. 1820. C ?1 Lab. vo. 2.18
- . 1612. SAURAU'Jii W. SAURAUJA. (*SauratoO*, a Spanish botanist ?) *Ternstrcemicdeca* 2.—19
 4319 excelsa W. tall ft • or Caracci : Nepal & 8 Jl ^ *
- "nepalensis Dee. Nepal * D o r Nepal

CLASS XIII. ORDER 6-

POLYA'NDRIA POLGY'NIA.

1613. <i>NELU'MBIUMJ.</i>	SACRED BEAN	(<i>Nelumbo</i> , its name in Ceylon.)	<i>Nympheehceee.</i> 5. ~7.
14321 <i>speciosum W.</i>	showy	^ C 3 d'f flt jn.au Pk India	1787. R m.s Bot mag. 903
14322 <i>caspicum Fis.</i>	Caspian	#k El or flt : Pk Casp. Sea	1822. R mB
14323 <i>Tantira Rh.</i>	Tamara	* 2d or flt : Pk Malabar	1818. R m.s R. mal. 11.30
14324 <i>jamaicense Dee.</i>	Jamaica	* E] or lit : Pa.B Jamaica	1824. R m.i
14325 <i>Idicum W.</i>	yellow-flowered	[or flt : Y Carolina	1810. R r.o.s An.mu.13.29.42
1614 DILLETNM L. DILLEXIA. (<i>J. J. Dillenius</i> , prof. bot. Oxford, a famous cryptog.) <i>DULenihcece.</i> I.-ft			
1432fi <i>speciosa Thun.</i>	Bhoviy-flawercl f	CD tm SO ... Y E. Indies	1800. C p.l Ex. bot 1.2.3
1615 ILLUCIUM L.	ANISEED TREE.	(<i>Illicio</i> , to allure; perfume.)	<i>Magnolitcea?</i> 3.
14327 <i>floridum Ell.</i>	red Florida	sii : or 8 ap.jn R Florida	1766. C s.p Bot. mag. 439
14328 <i>flincktum L.</i>	Anise-scented	sii : or 6 my.jn R Japan	18.10. C p.l Cl. h 2.20.2.3
14329 <i>parviflurum Ms.</i>	small yellow-flwd	[or 6 my.jn Y Florida	1790. C p.l Yen. eels 22
1616. SMEATHMA'NN/i* Sol. SMBATHMANIA. (<i>Smeathmann</i> , an African traveller.) <i>Passifibre*</i> . 1.—3.			
14330 <i>lacwigta Sol.</i>	smooth-leaved	» D o r 6 f.m.r W S. Leone	1822. C s.p
1617. MROIODE'NDRONX. TULIP TREE.	(<i>Leirion</i> , a lily, <i>dendron</i> , tree.)	<i>Magnoliaceat.</i> 1.	
14331 <i>Tulipifera L. common tulip.bearing*jf</i>	or 60 jnjl Y.R	N. Amer. 1663. t s.l Bot mag. 275	
2 obtusiloba	obtuse-lobcd	\$ or 60 jnjl Y.E Pennsylv. 1663. ■■	
* 1618. MAGN(TL)4 L. MAGNOLIA. (<i>Peter Magnol</i> , M.D. prof. bot. Montpelier.)		<i>Magnoliacear.</i> 17.—SL	
14332 grand i flora L.	large-flowered	W Carolina 1734. L 1p	
1 elliptica	oval-ferruginous	spl 20 jn o W Carolina 1734. L 1p	
2 obovata	ovate broad-leaved	spl 20 jn.o W Carolina 1734. L 1p	
3 lancolata	Bpc&r-icaved	spl 20 jn.o W Carolina 1734. L 1p	
4 ferruginea B. M.	rusty	or 20 jn.o W N. Amer. — L 1p	
5 exoniensis	Exeter	or 20 jn.o W N. Amer. — Bot cab. 814	
6 rotundiflora	round-leaved	spl 20 jn W N. Amer. —	
7 pre'cox	early	or £0 my.o W N. Amer. —	
8 crispa	curled	spl 20 jn W N. Amer. —	
14333 glauca L.	glaucous decidu. swamp*	or 20 jn.s W N. Amer. 1688. F Bot. mag. 2164	
2 Gordonida Hort.	Gordon's dble	or 20 jn W 1750. F	
3 BurchellifVina Hort.	Bun-hell's dble	or 20 jn W —	h
4 Thomsonia/m Hort.	Thomson's	or 20 jn.o W hybrid" 1817. L 1p	
14334 longifolia/f if.	long-lvd cverg.swamp	or 80 jn.s W N. Amer. ... S p.1	
glauca & longii'lia Ph.			
14335 triptala L.	threc-petalcd	or 30 myjn W N. Amer. 1752. L s.l Mic.ar.5	
umbrella Lam.			
14336 acuminate L.	acuminated	or 60 my.jl Y.o N. Amer. 1736 L 8.1 Bot. cab. 418	
14337 auriculata Lam.	car-leaved	or 40 ap.my W Carolina 1780. L p.1 Bot. mag. 1206	
14338 pyramidata Bart.	pyramidal	or 20 ap.my W Carolina 1811. G p.1 Bot. rog. 407	
14339 macrophylla Mx.	long-icaved	or 30 jn.jl W N. Amer. 1800. S p.1 Bot. mag. 2189	
14340 maxima LotL	largest	or 30 jn.au W N. Amer. 1820. L p.1	
14341 cordata Mx	heirt-eaved	or 40 jnjl Y.w N. Amer. 1801. L s.1 Bot. cab. 474	
14342 conspicua Sat.	conspicuous	or 30 f.ap W China 1789. G p.1 Bot. mag. 1621	
gracilis Sal. Yiilan Doc.			
14M3 tomentuKn Thun.	tomentoce	—J or 20 mr.ap P China 1804. L p.1 Par. Ion. 87	
Kbbus Dec.			
14344 obovata Thun.	obovate	St J or 6 apjn P China 1790. ■■ Kami. 43	
2 discolor Dec.	various-colored	a — or 6 apjn P.W — 17D0. ■■ Bot. mag. 390	
14345 Soulaiigflna Hort	Soulange's ft	ft — or 5 apjn P.W hybrid 1826. ■■	
14346 fuscata Andr.	brown-stalked	ft i or 3 ap.my Br China 1789. ■■ Bot. mag. 1000	
14347 anomellYm Sal.	Anona-leaved	ft i or 3 ap.my Br China 1804. ■■ Par. Ion. 5	
14348 pumila Andr.	dwarf	ft — or 3 year W China 1786. ■■ Bot mag. 977	
1619. MICHE'LU L.	MICUELIA.	(P.A. Michelu the great Florentine botanist.)	<i>Magnoliacea.</i> L — ■
14349 <i>Champaca L.</i>	Champaca	J 0 t m 2 0 ... Y E. Indies 1779. C s.l 11. mal. 1.19	
* 1620. UVA'RIA/..	UNARIA.	(<i>Uva</i> , a grape; bunches of fruit like grapes.)	<i>Anonhcea.</i> 6. — 10-
14350 seyanica L.	Ceylon	r 20 ... R.o K Indies 1794. C p.1 R. mal. 2.10	
14351 velutina Dunal	velvety	» L J or 6 ... Br E. Indies IH23. c Pi	
14352 Gsertneria Dec.	Gsertner's	ft . or 6 ... Br E. Indies 1820* C p.1 Gae.fir.2.Hf/	
14353 futea liox.	yellow	• CD or 6 ... Br E. Indies 1822. C p.1 Rox.cor.1.3'	
14354 ltica Boj.	shining	ft . or 6 ... B Madalbar 1825. C p.1	
14355 tomentosa Box.	woolly	ft O or 6 ... Br E. Indies 1822. C p.1 Rox. cor. 1.■	
* 1621. UNO'NA L.	UNONA.	(<i>Uno</i> , to unite: stamens with germen.)	<i>Anandcea?</i> 7. -38.
14356 N&rum Dunal	Narum	L Dor 10. Br Mabalbar 1820. C p.1 R. mal. 2.9	
14357 fuscata Dec.	brown	• Dor 5 . Br Guiana 1818. C p.1	
14358 acuminate Dee.	acuminate	CDt 6 — Br Guiana 1820. C p.1	
14359 esculenta Dunal	esculent	10 — Br Madras 1818. C p.1	
14360 odorata Dunal	sweet-scented	O 5 — Br E. Indies 1819. C p.1 Lam.il. 495.1	
14361 longi folia Lam.	long-leaved	4 — Br Bengal 1820. C p.1 Son. it 4. 1.3 k	
14362 aromatic Dunal	aromatic	Qor 6 — Br Guiana 1820. C p.1 Aub.gui.l.	
1622. ANCPNA Adan.	CUSTARD APPLE,	(<i>Menona</i> , its Banda name.)	<i>Anonaceaf.</i> 17. —36.
14363 muricata L.	muricated Sour-sop	G.Y W. Indies 1656. C r.m Tac. ob. 1.5	
14364 fourifolia Dunal	Laurel-leu ved	Br W. Indies 1823. C r.m Cat car. 2. 67	
J4365 obtusiflvia Tus.	obtuse-leaved	Y.c W. Indies 1810. C r.m FID	
"366 palustris W.	marsh Cork-wood	Y W. Indies 1731. C r.m	
J4367 Aub.	long-leaved	Y.o Guiana 1820. C r.m Aub.gui.JIJ	
J55 paludosa A.*	spotted	Y.G Trinidad 1818. C r.m Aub.gui.;Uf	
J55 paludosa A.*	marsh	G Guiana 1803. C r.m Aub.gUL	
14370 squamosa L.	scaly Sweet-sop	W.o S. Amer. 1731. C r.m R.maL3.w	
14371 cinerea Dunal	t	Y.a W. Indies 1818. C r.m	

14372 <i>Cherimolia</i> Mil. tripetala H. K.	Cherimoyer	+	□	h	18	jl.au	Br	S. Atner.	1739.	C r.mTr. ehr.49
14373 <i>reticulata</i> L.	netted	†	•	fir†	20	...	W.a	S. Amer.	1690.	C r.m R. mal. 3.30.31
14374 <i>mucosa</i> Jac.	mucous	ft	•	fir	12	...	Y.G	E. Indies	1820.	C r.m Ru. am. 1.4.5
14375 <i>gibbra</i> L.	auiooth-fruited	±	•	or	16	jl.au	Br	Carolina	1774.	C r.m Cat. car. 2.64
14376 <i>amplexicaulis</i> Lam.	stem-clasping	■	□	or	12	...	Y.a	Mauritius	1824.	C r.m
14377 <i>asiatica</i> L.	Asiatic	■	□	or	12	...	Y.a	Asia	1816.	C r.m
14378 <i>mexicana</i> Lo. C.	Mexican	■	□	or	12	...	Y.a	Mexico	1823.	C r.m
14379 <i>senegalensis</i> Dunal	Senegal	■	Dor	10	...	Y.a	Guinea	1824.	C r.m Deless. 1.86	

* 1623. MONODCRA Dunal. MONODOBA. (*Monos*, one, *doron*, gift; fruit solitary.) *Anonhcecc.* 1.
14380 *Myristica* Dunal Amer. Nutmeg* • or 20. Jamaica ... C p.l Gae. ft. SL125.1
Anbna *Myristica* *Gae.*

1624. ARTABOTRYS R. Br. ARTABOTRYS. (*Artqo.* to suspend, *botus*, grapes; by tendrils.) *Anonhea*. 1.
14381 *odoratissima* fflbz. sweet-scented Q or 6 jnjl Br China 1758. C r.m Bot.reg.423
hexapetalia W.

1625. GUATTETRII R. & P. GUATTERIA. (J. B. Guattieri, prof. bot. at Parma.) <i>Anonhcecc.</i> 5.—MS
14382 <i>rufa</i> Dunal rufous ft Q or 3 jl.au Br China 1822. C r.m Bot. reg. 336
14363 <i>virgata</i> Dunal twiggy Lancewood ± tm 30 □ W Jamaica 1793. C p.l Dun. mon. 31
<i>Uvaria lanceolata</i> Swx.
14384 <i>ceraspides</i> Dunal Cherry-like HQor 16 — G E. Indies 1820. C p.l Rox. cor. 1.33
14385 <i>superba</i> Dunal cork-barked *dlor 8 — W E. Indies 1820. C p.l Rox. cor. 1.34
14386 <i>aurifolia</i> Dunal Laurul-leaved >Qor 8 — W Jamaica 1818. C p.l Dun. mon. 32

1626. ASIMINA Adan. ASIMINA. (A name of Canadian origin, meaning unknown.) *Anonhcecc.* 4.—5.
14387 *triloba* Dunal three-lobed-/W Se or 8 au Pa.P N. Amer. 1736. S p.l Cat. car. 2.83

Anbna *triloba* L.
14388 *parviflora* Dunal small-flowered Si cu 3 ap.my Br N. Amer. 1806 L pi Dunrmon. 9
14389 *pygmaea* Dunal pygmy SI cu 2 ... W N. Amer. 1812. L p.l Bart. it. 1
14390 *grandiflora* JMfl/ large-flowering SI or 3 ap.jl W Georgia 1820. L p.l Dun. mon. 11

1627. XYLOPIA L. XYLOPIA. (Altered from *Xylopicron*; *xylon*, wood, *pticros*, bitter.) *Anon&ceas.* 3.—9.
14391 *muricata* X. rough-/rw7-rf «O or 4 . . . W. Inaiesl/93. C p.l gr-f* r*
14392 *gKibra* L. smooth./mtorf t • tm 20 . . . Jamaica ... C p.l Pluk. al. S^ 4
I M i g f a f i . firutescent i o o r 4 . . . Guiana 1823. C p.l Aub. gui. 1.21)2

1628. HEPATICA Dil. HEPATICA. (<i>Hepatikos</i> , relating to liver; lobed leaves.) <i>Ranunculace</i> *. 4.—5L
14394 <i>ameru&na</i> B. R. com. American — — — — N. Amer. 1800. D 81 Bot. reg. 387
14395 <i>angulosa</i> Dec. angulose — — — — N. Amer. 1816. D sl
14396 <i>acutiloba</i> Dec. acute-lobed ^ww. — — — — N. Amer. 1818. D sl
14397 <i>triloba</i> Chaix common 3-lobed — — — — Europe 1573. D si Eng. bot 51
^nemuiia Hepatica L. 1 ceriilea blue — — — — D sl 2 cscr&leo-plt-na double blue — — — — D sl 3 rubra red — — — — D sl 4 rbbro-plena double red — — — — D sl 5 alba red-anthered wl — — — — D sl 6 nivea mow-white — — — — D sl

* 1639. i4NEMO%NE L. ANEMONE. (*Anemos*, the wind; growing in exposed places.) *Ranunculacea?* 38.—

I. PIJLSATILLA.—Seeds ending in long bearded tails; lvs of involv., sess. palmately parted into linear lobes.
14398 <i>vemalis</i> L. spring ap W Switzerland 1752. R s.p Sw. fl. gar. 205
2 flore HitM yellow-flowered *ap Y S. Europe ... R co
14399 <i>Halleri</i> All* Haller's drooping ap.my P Switzerland 1816. R s.p Bot cab. 940
JJJW <i>cernua</i> Thun. spreading I my.jn R.w Japan 1806. R s.p
14401 pMens L. spreading 1 jnjl L.Y Siberia 17^>2 R s.p Bot. reg. 61
2 ochroleuca B. M. yellowish white 1 jnjl V Siberia 1752. R PS Bot. mag. 1994
J4402 l>uhatilla L. com. Pulsatilla ap.my B England ch. pa. R s.p Eng. bpt 51
14403 acutipetaloida ScA. acute-petaloid i my D.P Switzerland 1819. R s.p
14404 pratinensis L. meadow i my D.P Germany 1731. R s.p Fl. dan. 611'
U40. oluofeta Sims obsolete 4 my P Germany ... R sp Bot mag. 1863
14406 albina Stev. Albana I my W Caucasus 1821. R s.p
14407 dahurica Fis. Dahurian I my F Dahuria 1819. R s.p

II. PUBONA'NTUUS.—Seeds ending in long bearded tails; leaves of involucrum trifid. petioled.

14408 alpina L. alpine * A *r jl W.p Austria 1058. R s.p Bot. mag. 2007
14409 sulphurea L. Bulphur-colored & or ap.my Su Europe 1816. R s.p Cam. ep. 393. ic. flpiK»lia W.
14410 micrantha Dec. malflowered ^ A or } jn.jl W.p Austria 1800. II s.p Jac. au. 1.85 alp'ma Jac.

III. ANEMOVA'NTHEA.—Seeds void, tailless; pedicles 1 or 2, leafless, l/fiwd.; leaflets of invol. sess. or petioled.

14411 coronaria L. garland A A or ap.my St Levant 1596. R Lp Bot mag. 841
2 plena double-flowered& A or f.ap.my St R r.l
14412 pavonina Dec. peacock-ry^ A A or ap.my R France : II < p M.h.4.25.10
2 flure plzno doubleflowrcrdA A or ap.my R Europe : R s.p Cl. h. 261. ic.
3 ftlgens Dec. shining A or ap.my R S. Europe 1818. R s.p
hortensis Thore, pavonia Lot. garden A A or I ap.my St Italy 1597. R s.p Bot mag. 123
14413 hurtensis L. small-flowered A A or I ap.my St Italy 1597. R s.p Bot mag. 123
stellata Dec. versicolor Sal. palmated A A or I my.jn Y Portugal 1597. R s.p Bot reg. 200
14414 parviflora 3fr. small-flowered A A or I ap.jn W N. Amer. 1824. R s.p An. mu. 3.21.1
cuneifolia J. caroliniana Walt. Carolina A A or I my.jn W Carolina 1824. R s.p
14416 caroliniana Walt. tenella Ph. Apcnnine A A or) mr.ap B England woods. R s.p Eng. bot 1063
14417 apennina L. blue A A or 4 my.jn B Siberia 182rf. R s.p Deless. 1.14
14418 caerulea Dec. Ural A A or i my B Siberia 18^4. R s.p
14420 baldensis L. Mount Baldo A A or W Switzerl. 1792. R *p Jac. ic. 1.103
fragifera Mur. grove A A or i mr.my W.R Britain woods. R s.p Eng. bot 355
14419 nemordia L. double-floweredA A or i mr.my W.R Britain woods. R s.p
j**f2 quinquefolia L. 5-lyd Amer. vwood& A or } mr.my W N. Amer. 1817. R s.p Bar. fl. S. 2
'4423 trifolia L. three-leaved A A or ^ ap.my W France 1597. R s.p M. h. 4.25.1

14424 lancifolia PA.	lance-leaved	A A or	ap.my	W	N. Amer. 1822.	R s.p	
14425 ranunculoides L.	Ranunculus-lk	A A or	4 mr.ap	Y	England woods.	R s.p	Eng. bot 1484
14426 reflexa Stev.	reflexed	A A or	t mr.ap	Y	Siberia 1818.	R s.p	Defess. 1.15
14427 sylvestris L*	wood. Snowdrop*	A or ft	ap.my	W	Germany 1596.	R s.p	Bot. mag. 54
14428 alba J. ochotensis Fts.	white	‡ A or	jn	VV	Siberia 1820.	R s.p	Bot. mag. 2167

IV. ANCMONOSPB' BMO9.—Seeds tailless compressed; pedicles from involucrum numerous, \flwd,							
14429 virginiana L.	Virginian	2L A or	‡ my.jn	W	N. Amer. 1722.	R s.p	Her. par. 18
14430 multifida Dec.	many-cleft	* A or	1 jn.jl	W	Magellan 1824.	R ap	Deless. 1.16
14431 pennsylvanica L.	Pennsylvania	‡ A or	1 my.jn	W	N. Amer. 1766.	R ep	
14432 dichotoma L.	forked	‡ A or	1 my.jn	Pa. W	N. Amer. 1768.	R s.p	L. f. dec. 2.15
V. OMALOCARPUS.—Seeds compr., oval-orbicul., smooth, without tails; pedicles numer. umbel, leafless, l-Jiwd.							
14433 narcissiflora L.	Narcissus-flwd	‡ A or	1 my	W	Siberia 1773.	R a p	Bot. mag. 1120
14434 sibirica L.	Siberian	3t A or	} jn	W	Siberia 1804.	R s.p	
VI. PULSELLATOI'DES.—Seeds very hairy; calyx 15 or SO-sepalled; leaves of involucrum 2 or 3, sessile, at top cut.							
14435 capSnsis L.	Clematis caplnsis	‡ tAI or	1 mr.ap	P	C G. H. 1795.	S pl	Bot. mag. 716

Clematis caplnsis Pmr.

1630. ilTRAGE^NE L. ATRAGENB. (A name given by Theophrastus to a species of Clematis.) <i>Ranunculus</i> . 5.							
14436 austriaca B. M.	Austrian	±	or 8	jn.jl	B.Y Austria	1792.	L co Bot. rep. 180
alptaa L.	Clematis alpina Mil.						
14437 sibirica L.	Siberian	1	or 12	jn.jl	W.Y Siberia	1753.	Le o Pal. ros. 2.76
Clematis sibirica Mil.							
14438 ochotensis Pall. Ochot&k, petaled	±	or 12	my.jl	W	Siberia 1818.	L co	
Cilmatis ochotensis Poir.							
14439 americana R. M. whorled American^	or 15	my.jn	P	N. Amer. 1797.	L s.p	Bot mag. 887	
Clematis verticillaris Dec.							
14440 occidentalis Horn, occidental	±	or 10	jn.jl	1818.	L ap
Clematis occidentalis Dec.							

*1631. CLEMATIS L. VIRGIN'S BOWER. (Klema, a tendril; petioles of old leaves.) *Ranunculaceae*. 39.-84.

I. FLAMMULA.—Seeds with long feathery bearded tails; flowers panicled; leaves pinnate or bipinnate, from No. 14450. ternate or bUern.; from No. 14459. peduncles trifid, 3.flwd, Ivs tern. from No. 14463. peduncles solitary, \flowered, leaves pinnate; and from No. 14470. peduncles solitary, \flowered, leaves undivided.

1 mi erecta All.	upright	k A or	3 jn.au	W	Austria 1597.	D pl	Jac. au. 3.291
2 hispanica	Spanish	‡ A or	3 jn.au	W	Spain 1800.	D co	
14442 Flammula L.	Flame	‡	or 20	jl.o	W	France 1596.	S co Kno. th. 2.9
1 rotundifolia	round4eavcd	±	or 20	JLo	W	France 1596.	L co
fragrans Ten.							
2 vulgtais	common	i	or 20	jl.o	W	France —	L co
3 maritima L.	sea	i	or 20	jl.o	W	S. Europe ...	L pl
14443 Massoniflora Dec.	Masson's	or 20	jn.8	C. G. H. 1731.	Sko Di. el. 119.145
14444 orientalis L.	oriental	LLjor	12	...	Y.w	Levant 1731.	Sko Di. el. 119.145
itMdglauca W.	glaucous	1	or 8	jl.o	Pa.Y	Siberia <	L co Den. br. 73
It4kJ chinensis Jtetz.	Chinese	±	or 12	ap	W.o	China 1820.	L co Retz. 2
14445 pamculata Thun.	panicled	‡ Jor	12	...	W	Japan 1800.	Li CO H.pfl.7.2
14446 Viulba L.	{ white vine, or "I" Traveller's Joy }	or 20	jl.au	W	England hed.	S co Eng. bot 612	
14447 virginiana L.	Caripe	fl	12	...	W	Trinidad 1820.	‡ Pl
14448 brasiliensis Dec.	Virginian	X	15	jn.au	G	N. Amer. 1767.	L sp Den. br. 74
14449 brasiliensis Dec.	Brazilian	fl	12	...	W	Brazil 1823.	L Deles*. 1.1
14450 dioica L.	dioicus	fl	15	my.jn	G,Y	W. Indies 1733.	L s.p SL. jam. 1.128.1
14451 americana Mil.	American	fl	12	...	W	S. Amer. —	L s.p
14452 glycinoides Dec.	Glycine-like	fl	10	...	—	N. Holl. 1826.	L pl
14453 conacea Dec.	leather-/<lawer/	‡	12	...	W	N. Holl. 1821.	L pl
australis IM. C.							
1 Haed aristata B. R.	wned-anthered	‡ J or	15	my.au	G.v	N. Holl. 1812.	L s.p Bot. reg. 238
14454 Adysariflbia Dec.	Hedysarum-lvd	‡	or 12	o	W	E. Indies 1819.	L co Bot. reg. 599
14455 triternata Dec.	internate	‡	or 12	...	W	... 1800.	L pl
14456 brachikta B. R.	brachiate	fl	1 I or	12	o.d	Y.G C. G. H. ...	L s.p Bot. reg. 97
14457 grandiflora Dec.	large-flowered	fl	CD or	12	f.my	Y.o S. Leone 1823.	L pl
14458 zansibarensis Boj.	Zanzibar	A. I—Jor	JO	—	—	Zanzibar 1820.	L pl
14459 dahurica Pers.	Dahurian	—	12	8	Y.G	Duhuria 1820.	L pl
14460 angustifolia Jac.	narrow-leaved	A or	4	my.s	W	Austria 1787.	D pl Den. br. 112
14461 lineariloba Dec.	linear-lobed	A or	4	my.jl	W	Carolina? 1823.	D pl Deless. 1.3
14462 diversifolia Dec.	various-leaved	A or	4	8.0	W	— D	Deless. 1.4
14463 Viurna L.	Traveller's Joy	‡	12	jn.8	P	N. Amer. 1730.	S s.p Di.el. II.8.1f*
14464 cylindrica A Af.	y	A or	3	jl.s	B	N. Amer. 1802.	L pl Bot. mag. lieu
14463 Simstii Swt.	Sims's	‡	or 8	jn.au	P	N. Amer. 1812.	L co Bot mag. 181°
cordata Sims							
14465 reticulata Watt.	netted	—	8	jn.8	P	N. Amer. 1812.	L s.p Den. br. 72
14466 integrafolia L.	entire-leaved	—	2	jn.au	B	Hungary 1596.	D pl Bot. mag. 65
2 angustiflora Y.	narr. cintre-lvd	—	2	jn.au	B	Hungary ...	D pl
3 elongata Trat.	elongated	—	2	jn.jl	B	Europe ...	Trat. tab. 4.1?
14471 ochroleuca H. K.	yellowish white	—	2	jn.jl	L.Y	N. Amer. 1767.	D pl Bot. cab. 661

II. VITICELLA.—Tails of seeds short, beardless; leaves ternate or decomound.							
14472 flpirida Thun.	florid	or 10	aps.	W.Y	Japan 1776.	L s.l	Bot mag. 834
2 fibre pleno	double-flowered	‡	or 10	ap.s.	W. Y. . . .	L s.l	
14473 Viticella L.	vine-bower	‡	or 20	jn.s	P	Spain 1599.	S co Bot mag. 5»
2 plena	double purple	‡	or 20	jn.s	P.	L co	ntd
14474 campaniflora Brot.	bell-flowered	‡	or 6	jn.s	P	Spain 1810.	L s.l Bot. cab. 987
14475 crispa L.	curled-flowered	‡	or 6	jl.s	Pa.P. N. Amer. 1726.	L pl Bot mag.	

III. CHEIR6PSIS.—Invol. calyciform, situated at top of peduncle underflower; tails of seeds bearded.							
14476 cirrhosa L.	tendnlcd	JL	or 12	mr.ap	W.G Spain	1596.	Le o FLgr.51
14477 pedicellata Swt.	pedicellate	fl	or 12	jnau	W.G Majorca	...	L co Bot.mag.**"
cirrhosa B. M.							
14478 Bemitrifolia Lag.	half three-lobed*	or 12	mr.ap	W.G Spain	... L co	05	
14479 baileynia J.H.C.A.	Balearic	‡ « J or	12	f.mr	Y.w AKnorea	1783.	Le o Bot mag. **"
calycina H. K.							
14480 ?Ray LIA Dec	NABAELIA.	L D J W	12	L Y	Ceylon	1796L	L s.p Rox.cor.-
Hyacinthina W.							
Jtr.igene zeylanica L.							

* 1633. THALI'CTRUM L. MEADOW RUB. (*Thallo*, to grow green; young shoots.) *Ranunculaceae*. 58.-67.

I. TRIPLE^HION.—*Fruit triquetrous, 'angles winged, substantiate.*

14519 grācīlis Dec.	slender	£ iAl or	1 mr.ap	Y.a	C. G. H. 1820. S p.l	Deless. 1.19
14550 hirsūte Dec.	hairy	£ IAJ or	1 a p. my	Y.u	C. G. H. 1823. S p.l	Bur. af. 51
Adunis hirsuta Poir.						
14551 daucifūlia Dec.	Carrot-leaved	£ iAl or	1		C. G. H. 1822 S p.l	
* 1636. FICA'Rl A. Dil.	PILEWORT. (<i>Ficus</i> , a fig; fig-like tubercles of root.)				Ranunculōcea. 1. — 2.	
14352 ranunculōdes, Voc. Ranunculus-like A	w i	mr.my	Y	Britain he.ba. R l.p	Eng. bot 584	
Ranunculus Ficaria L.						
2 plena	doubtfully-flowered&	A or	I	mr.my	Y	Britain he. ba. R l.p
* 1637. R ANU'NCULUS BaUh. CROWFOOT. (<i>Rana</i> , a frog; inhabiting humid places.)	Ranunculaceae. 49.—165.					
I. BATRA^CHIUM. — <i>Pericarp transversely striated : petals white ; claw yellow, marked with a nectarifer. hollow.</i>						
14553 Adcfacūs L.	Ivy-leaved	■ A P' fit	my.au	W	Britain wat. pi. D co	Eng. bot. 2009
14554 tripartitus Dec.	three-parted	■ ^ pr	ap.au	W	Europe dit. D co	Deless. 7.43
14555 obtusiflorus Dec.	blunt-flowered	■ A pr	flt ap.au	W	England dit. D co	Pet. brit. 39.1
14556 pantothrix Dec.	all-hairy	■ A pr	flt ap.au	W	Britain dit. D co	Bar. ic. 566
U fluvialis Wahl.	long-lvd river	■ ^ pr	flt ap.au	W	Britain dit. D co	Fl. dan. 376
weuccapafolius All.		■ A	or flt			
3 cespitosus ThuiL.	turfy	■ A pr	flt jl		Ap	Britain dit D eo
rigidus Pers. circinatus Sib.		■ A lt	flt ap.au	W	Britain dit D co	Bar. ic. 565
14557 aquatilis L.	water	■ A pr	fit ap.au	W	Britain dit D co	Bar. ic. 565
2 peletans Dec.		■ A	or			
II. RANUNCULASTRUM. — <i>Carpels smooth, disposed into a close spike ; roots grumose.</i>						
14558 bullatus L.	blistered	A A or	1 myjn	Y	S. Europe 1640. B co	M. h. 31.50
2 grandiflōris Dec.	large-flowered	A A or	myjn	Y	S. Europe 1640. B co	M. h. 31.50
3 flōra pleno	double-flwd	A A or	myjn	Y	S. Europe 1640. D co	Cor. ca. 95. ic.
14559 chsrophylloides L.	Chervyieaved	A A cu	myjn	Y	Portugal ... D r.m	Bar. ic. 581
14560 gregarius Dec.	gregarious	A A or	myjn	Y	Italy 1817. R co	
Thomas M Ten.						
14561 paludosus Poir.	marsh	A A un	myjn	Y	Barbary 1816. D co	
14562 millefoliatus Vahl	thousand-lvd	A A or	myjn	Y	Sicily 1820. D co	Vahl ey. 2.37
14563 grācīlis Dec.	slender	A A or	i myjn	Y	Archipel. 1818. D s.p	
14564 umariae-filius Desf.	Fumitory-lvd	A A cu	myjn	Y	— D co	Desf. pic. 74
14565 oxysperimus W.	sharp-seeded	A A or	my	Pa.Y	Caucasus 1822. D co	
14566 Hornemannii Schl.	Hornemann's	A A un	—		N. Amer. 1820. D co	
14367 cicutarioides Schl."	Cicutariae	A A or	—		Siberia 1818. D co	Schl. ran. 4 2
14568 pedatus W. * K.	pedate	A A or	myjn	X	Hungary 1806. D co	Bot. mag. 2229
14569 illyricus L.	Illyrian	A A or	myjn	X	S. Europe 1596. D co	Jac.au. 3.222
sericeus W.						
14570 monspeliacus L.	Montpelier	A A or	ap.my	Y	S. France ... D co	M. h. 30. 43
2 cuneatus Dec.	wedge-leaved	A A or	ap.jn	Y	S. Europe ... D co	Dec. ic. 50
apiifolius Desf.						
3 rotundifolius Dec.	round-leaved	A A or	1 apjn	Y	S. Europe — D co	
monspeliacus Gou.						
14571 asiaticus L.	Asiatic com. gard.	A A or	^ myjn	Va	Levant 1596. D r.m	Mil. ic. 2.216
2 sangueinus Mil.	blood-colored	A A or	f myjn	S	Syria — D r.m	Cl. h. 1.242. ic
3 tenuilobus Dec.	finc-lobed	A A or	\$ myjn	W	Greece — D r.m	Bau.hi. 3.362. ic
14572 cortusa/d/i<W.en.	Cortusa leaved	A A or	1 myjn	Y	Teneriffe 1826. D s.l	
14373 crticus L.	Cretan	A A	1 ap.my	Y	Candia 1658. D co	M. h. 31.48
III. TrifERA. — <i>Carpels smooth, subglobose ; roots grumose.</i>						
14574 Thbra L.	Thora kidney-lvd	A "A or	Y		Austria 1710. D co	Jac. au. 5.442
14575 scutatus W.JkK.	ahield-leaved	A A or	—		Hungary 1817. D co	W. & K. 2.187
14576 brevifolius Ten.	short-leaved	A A or	—		Naples 1824. D co	
14577 hybrida Biv.	hybrid	A A or	f myjn	Y	Austria 1820. D co	Stur. deu. ic.
IV. HECATUNIA. — <i>Carpels smooth, ovate, roundish, disposed into a round head; roots fibrous ; flowers white*</i>						
Ivs cut, from No. 1459U. leaves undivided, from No. 14595. yell. Ivs undivided, from 14603. Ivs dissected.						
14578 rotundifolius L.	Uue-leaved	3> A P*	i " ^ , 1 w		Austria 1759. D r.m	Jac. sc. 1.6.7
14579 tsopyrdites Dec.	Isopyrum-like	A or	myjl		Siberia 1818. a I) CO	
14580 Rläciklis L.	icy	■ jn.ju	W		Iaplnd 1775. D s.i	FL dan. 19
2 aconitoides Dec.	Aconitum-like	A P*	§ jn.ju	W	Switzcr. 1819. D s.i	
14681 Scquit-ri Vil.	Scffuic'a	A	ijnjl	W	Piodinnnt 1HIP. 13. 8. r	Vil. doL 4.40
I-UHX MCOMILLIUM /..	Afciiiltu-uuutl	A Or	1 myjn	Y	Alp. Kur. LOW. I> CO	
1 hiimilis Dec.	humble	A or	i myjn	W	Europe ... D r.m	M.h.3.2.1
2 crassicaulis Dec.	thick-stemmed	A or	1 myjn	W	Europe 1769. D r.m	
14583 plataniifolius L.	Plane-trcc-lvd	A >r	2 jnjf	W	Germany 1769. D co	Fl. dan. III
H fluc pltno	double-flowered	A or	1 myjn	W	Alp. Eur. 1596. D co	Bot. mag. 204
oconitifolius B. M.						
14584 alp&tris W.	alpine	A Pr	4 jn.au	W	Scotland al. riv. D co	Eng. bot. 9390
14585 crenatus IV.SfK.	crcnated	A or	* jnjf	W.R	Hungary 1818. D s.p	W. & K. 2.10
14586 npifolius Pers.	Apium-leaved	Q) or	2 jnjf		Konaria 1816. S m.s	Deless. 1.26
14587 lficerus Bel.	torn	A or	f myjn	W.A.Y	S. France IK21. D co	Bel. tau. 5.8
14588 frigidus Schrank	frigid	A cu	f myjn	W	S. Europe 1827. D i p	Sch.mo.57
Pyrenean	A or	1 myjn	W	Pyrenees 1807. I> s.p	Deless. 1.27.3	
14589 Bupleurum-Niusta/L	Bupleurum-lvd	A or	* myjl	W	Pyrenees 1818. D r.m	Jac. m. 1.18.1
14590 plantaginaceus All.	Plantain-lvd	A or	1 ap.my	W	Piedmont 1819. D Bp	Al.ped.76.1
pyrenaicais J. & A.	jolantagincus Dec.					
14591 angustifolius Dec.	narrow-leaved	0 0	1 ap.my	W	Granada 1822. D s.p	Deless. 1.27. A
14592 amplexicaulis L.	stem-clasping	0 0	—		Pyrenees 1633. D co	Bot. mag. 266
14593 parnassifolius L.	Parnassia-lvd	0 0	—		S. Europe 1769. D ro	Bot. mag. 386
14594 gramineus L.	grassy	0 0	—		Wales al.me. D e o	Eng. bot. 2306
Sphecenicifolius D^c	Phæne-leaved	0 0	—		Europe gard. D ro	M. h. 2. 4.30. »
3 fluc plno	double-flw cring	0 0	—			D r.m
14595 Lingua L.					Britain mud.d. D co	Eng. bot 100
14596 Flaramula L.	Flame Spearw.	0 0	—		Britain wapL D co	Eng. bot 387
2 serrata Dec.	serrated	0 0	—		Britain wa.pl. D co	M. h. 2.4.29. »
3 ovala Dec.	ovule-leaved	0 0	—		Britain wa.pl. D co	
14597 reptans Dec.	creeping Spearw.	& A cu	—		Britain wa.pl. D co	Fl. dan. 108
14598 fibiflrmis Mt.	thread-formed	j^ A	—		N. Amer. 1823. D s.l	
J <99 bonariensis Poir.	Buenos Ayrcs	—	—		N. Amer. 1817. S m.s	Delets. 1.29
14600 salsuginosus Pall.	salt	—	p		Siberia 1822. D co	Jac. vin. 31
J 100 symcharia	boat-leaved	—	jnjf		Siberia 1824. D s.p	Am. rut 13. *
14602 auricomus L.	golden-haired	—	ltap.my	Y	Britain woods. D co	Eng-bot. 62*
14603 cassubicus L.	Cassubian	—	2 jnjf	Y	Siberia 1794. D co	Bot. mag. 221
11604 abortivus L.	abortive	—	J my.au	Y	N. Amer. 1713. D co	

14605 hyperboreus <i>Box.</i>	northern		ap.my	Y	N. Europe	1820.	D co	E.J. dan. 331
14606 sceleratus <i>L.</i>	hurtful		2	my.jn	Y	Britain	wapl.	S co Eng. bot 681
14607 pygmaeus <i>Wold.</i>	pygmy		i	ap.my	Y	Lapland	1810.	D s.p Fl. lap. 3.3
14608 nivalis <i>Gou.</i>	snowy		i	jn.au	Y	Lapland	1775.	D s.f Fl. lap. 3.2
14609 montanus <i>IV.</i>	mountain		i	jn.au	Y	Lapland	1775.	D s.l Jac.a.u. 325,326
14610 Vill&rsii Dec.	Villars's		1	my.jl	Y	S. Europe	1819.	D co Cr. au. 2.4, 2
14611 Gouans W. pyrenseus <i>Gou.</i>	Gouan's	A A or	1	my.au	Y	Pyrenees	1818.	D co Gou. il. 17.1,2
14619 acris <i>L.</i>	acrid					Britain	me.pa.	D co Eng. bot 652
2 flore pleno	double-flwd					Britain	...	D co Bot. mag. 215
3 sylvaticus <i>Thuil.</i>	wood					France	...	D co
4 multifidus <i>Dec.</i>	many-cleft					Europe	...	Lob. ic. 686.1
polyfrithemos <i>Lob.</i>								
14613 brbitus <i>Ten.</i>	Bmtian					Italy	1823.	D co Fl. nap. 1.50
14614 Steven's Bes.	Steven's					Volhynia	1819.	D co
14615 caucasicus <i>B*eb.</i>	Caucasian					Caucasus	1820.	D co
14616 polyanthemoides <i>L.</i>	many-flowered					N. Europe	1796.	D co Lob. ic. 666
14617 hirtius <i>Banks</i>	hairy					New Zeal.	1820.	D s.p
14618 hispidus <i>Mx.</i>	hispid					N. Amer.	1810.	D co
14619 nemoribus <i>Dec.</i>	grove					Switzerl.	1810.	O co
polyanthemos <i>Suter</i>								
SpaefliflurusDec.	few-flowered					Switzerl.	1819.	D co Amans5
, aureus <i>Schl.</i> villosus <i>St. Am.</i>								
14620 crassicaulis <i>H. C.</i>	thick-stemmed					Europe	1827.	D co
14622 lanuginosus <i>L.</i>	woolly-leaved					S. Europe	1683.	D co Fl. dan. 597
14622 tuberbus <i>Lap.</i>	tuberous					Pyrenees	1820.	D co
14623 napelliflbius <i>Dec.</i>	Napellus-leavedjfc					Turkey	1822.	D co
14624 dissipatus <i>Bab.</i>	CUT-fe[TFE]f					Caucasus	1818.	D s.p
14625 marylandicus <i>Poir.</i>	Maryland		1	my.jl	Pa.Y	N. Amer.	1811.	D co
14626 repens <i>L.</i>	creeping		A	my.au	Y	Britain	me.pa.	D co Eng. bot 516
2 fibre pleno	double-flowered,*	A or	T	my.au	Y			
J15S Mre&tus // K.	pale hairy	O		jn.o	Y	England	rubble.	S co Eng. bot. 1504
J4628 pennsylvanicus <i>L.</i>	Pennsylvanian	A un	1	jn.il	Y	N. Amer.	1785.	S p.l Jac. ic. 1.105
14629 fascicularis <i>Mhl.</i>	bundled	A or	injl	Y	N. Amer.	...	D co Schl. ran. 2.30.2	
14630 tomentibus <i>Poir.</i>	woolly	A or	my.jl	Y	N. Amer.	1820.	D co	
14631 obtusifolius <i>Horn.</i>	obtuse-leaved	A un	my.jl	Y	Spain	1819.	D co	
14632 trifoliatus <i>Desf.</i>	trifoliate	A un	my.jl	Y	Siberia	1817.	D co	
14633 grandiflirus <i>L.</i>	large-flowered	A or	i my.au	Y	Capparicioa	...	D co Desf. ch. 44	
14634 appiceus <i>Sm.</i>	Burdock-like	LAJor	jn.jl	Y	N. Holl.	1822.	D s.p	
;4635 piebeusil.jfr.	plebeian	LAJor	my.jl	Y	N. Holl.	1820.	D co	
14636 bulbibus <i>L.</i>	bulbous		my.jn	Y	Britain	me. pa.	R co Eng. bot 515	
2 fibre pleno	double-flowered	A or	my.jn	Y			Lob. ic 666. S	
3 bractetus <i>Schl.</i>	bracted	A A or	my.jn	Y	Pyrenees	..	R co	
V. ECHINELLA — Carpels rough, from tubercles or prickles; leaves dissected or lobed, from No. 14647. entire.								
14637 Philonotis'ikte.	moisture-loving	O or	ijn.au	Y	S. Europe	1800.	S co Col. ec. 316.1	
14638 pifervulus <i>L.</i>	little upright	G w	jl.au	Y	England	...		
14639 tuberculous Kit.	tuberculated	O or	1 jn.au	Y	Tauria	1817	S 88	
14640 arvensis <i>L.</i>	corn-field	O un	1 jn.au	Y	Britain	cor.fi.	S co Eng. bot 135	
*4641 muncatus <i>L.</i>	p'r'c Yy-Seeded	O un	jl.au	Y	S. Europe	1683.	S co Lam. il. 498	
2 creticus <i>Dec.</i>	Cretan	O un	jljau	Y	Crete	...	S co M.h.2.4.29.24	
3 caroUnus <i>Dec.</i>	Carolina	O un	jjljau	Y	N. Amer.	...	S co Ven. eels 73	
echinatus <i>Ven.</i>								
*4642 ventribus <i>Ven.</i>	ventricose	O cu	jl.au	Y	Brazil	..	S co	
14643 muricatus <i>y brasiliensis Dec.</i>	small-flowered	O w	mir.jn	Y	England		Eng. bot 120	
iff* ³ P ^r »flbrus <i>L.</i>	Scio	O cu	jinjl	Y	Archipel.			
14644 chlus <i>Dec.</i>	sessile-flowered	O cu	jinjl	Y	N. Holl.			
J*4645 sessiliflbus <i>R. Br.</i>	thrc-lobed	O or	1 jn.jl	Y	Greece	1818.		
14646 trilobus <i>Desf.</i>	Snake'8-tong.-lvd	O um	jn.jl	Y	S. Europe	1823.		
14647 ophioglossiflbus <i>Iusrtf.</i>	knot-flowered	8 um	* i myjl	Y	Sicily	1714.		
14648 nodiflorus <i>L.</i>	toothed	8 um	i myjl	Y	Sicily		Bot. mag. 2171	
S dentatus <i>Dec.</i>								
•163a TRO'LLIUS <i>L.</i>	GLOBE FLOWEV.	(7Vof,a globe, Ger. s shape of flowers.; Ranunculaceae. 6.						
14649 europeus <i>L.</i>	European	2 ⁿ	my.jn	Y	Britain	groves.	D p.l Eng. bot 28	
2 h&milis <i>Dec.</i>	dwarf	1	my.jn	Y	Austria	1800.	D p.l	
14650 napelliflbius <i>Crx.</i>	Nupcellus-leaved		my.jn	Y	Europe	1800.	D p.l	
14651 asiticus <i>L.</i>	Asiatic		my.jn	Y	Siberia	1759.	D p.l Bot mag. 225	
2 intermedium	intermediate		my.jn	Y			D p.l	
3 hbridus	hybrid		my.jn	Y				
14652 caucasicus <i>Stev.</i>	Caucasian		my.jn	Y	Caucasus	1817.	D p.l	
14653 patulus <i>Sal.</i>	spreading		my.jn	Y	Siberia	1800.	D p.l Dclss. 1.44	
ranunculus Minus <i>Sm.</i>	American	■ A or	myjl	Y	N. Amer.	1805.	D co Bot mag. 1988	
14654 americanus <i>Mill.</i>								
14655 lumnoides <i>L.</i>								
14656 halictroides <i>L.</i>								
1639. JSOPY^RUM <i>L.</i>	ISOPVRUM. (<i>Iso\$ equal, yros, wheat; taste of seeds.</i>) Ranunculaceae. 2, -4.							
•4655/uinan6ides <i>L.</i>	Fumitory-like	O pr	jn	W.G	Siberia	1741.	S s.l Am. rut. 74.12	
14656 /halictroides <i>L.</i>	Thalictrum-lk	^ A P'	I mr.ap	W.o	Italy	175J.	D s.l Jac. au. 2. ia>	
• 1640. ER A'NTHIS <i>Sal.</i>	WINTER ACONITE. (<i>Erao, to love, anthos, flower; fl. lovely in winter.</i>) Ranuncu. 2.							
14657 hyemalis <i>Sal.</i>	common winter	A A or	\ ja.mr	Y	Italy	1596.	O co Bot. mag. 3	
7elleborus hyemalis <i>L.</i>								
14658 sibiricus <i>Dec.</i>	Siberian	A A or	\ mr.ap	Y	Siberia	1826.	O co	
1641. HELLE'BORUS <i>L.</i>	HELLEBORE. (<i>Helcin, to cause death, bora, food; poisonous.</i>) Ranunculaceae? 8.—JL							
14659 figer L.	black Christmas Rose.	A or	* jn.mr	Pk	Austria	1896.	D ram Bot mag. 8	
2 angustifolius	narrow-leaved	£ A or	1 mr.ap	Pk		1596.	I co	
14660 v/ridis <i>L.</i>	K ^{roen}	£ A or	2 mr.ap	G	Britain	woods.	D co Eng. bot 200	
14661 purpurascens <i>Kit.</i>	purpurascens	£ A or	1 mr.ap	P.G	Hungary	1817.	D s.l W.&K. 2.101	
14662 odorus <i>Kit.</i>	sweet-scented	£ A or	1 mr.ap	P	Hungary	1817.	D s.p W.&K.a271	
14663 atrorliensis <i>JKF.</i>	dark-red	£ A or	1 mr.ap	G	Hungary	1820.		
146M dumetorum <i>Kit.</i>	thicket	£ A or	1 mr.ap	G	Hungary	1817.	I s.l	
iKwifcc'tidusL.	fetid B*ar's-foot	A or	H f.ap	G	England cha.pa.	D co	Eng. bot 613	
		O	4					

- 14666 *lvidus H. K.* livid three-lvd ^{AJ} **H** 1 ja.my **T** Corsica 1710. D p.1 Bot. mag. 72
2 *integrilobus* entire-leaved **JF** **Alor** 1 ja.my **T** Corsica 1710. D p.1
14642. *COPTISSal.* COPTIS. (*Kopto*, to cut; leaves much cut.) *Ranunculaceae*. I.—2
14677 *trifolia Sal.* three-leaved **jf** A P^r 4 ap.ray Br N. Amer. 1782. D p.1 Bot. cab. 173
Jfelleborus trifclius L.
- 1643. CA'LTHA L. MARSH MARIGOLD. (A syncope *otkalathos*, a **Tablet**; **flower**.) *Ranunculceal.—II,*
14668 *palGstris L.* common marsh ^A **1** ap.my Y Uratian mar. D m.s Eng. bot. 506
2 flûre pleno double-flwd ^A or **1** up. my Y **1** m.s
- 14669 *radicans G. Don* rooting WK A or \ ap.my Y Scotland sc.ma. D m.s Lin. tr. 8.17
14670 *asarifolia Dec.* Asarum-leaved ^A or i ap.my Y Unilasch. 1824. D m.s
- 14671 *minor Mil.* smaller ⁱ A or i my **T** Britain moun. D m.s Tab. ic. 750.2
14672 *parnassifolia Rff.* Parnassia-lvd ^A or \$ap.my Y N. Amer. 1815. D m.s
ficario'ides *Ph*
- 14673 *flabellifl M /%* fan-leaved ^f or **1** ap.my Y N Amer. 1818. D m.s Ph. am. 2.17
14674 *natana Ph /* floating ^A or ^{fl} my-jn W Siberia 1816. D m.s Gm.si.4.82
1644. HYDROPELTIS Z. HYDROPELTIS. (*Hydor*, water, *peUe*, buckler j leaves.) *Hydropeleeee*. 1.
14675 *purpurea Mx.* purple ^{AJ} cu **jLau** R N. Amer. 1798. D m.s Bot mag. 1147
Braseta peltata Ph.
- 1645. HYDR A'STIS L. HYDIUSTIS. (*Hydor*, water; growing in humid places.) *Ranuncul&ce**. 1.
14676 *canadensis L.* Canadian ^{j^J} or 4 my.jn G Canada 1759. D m.1 Mil. ic. 2.285

CLASS XIV. ORDER 1.

DIDYNA^MIA GYMNOSPE'RMIA.

1646. A'JUGA L. BUGLE. ^{priv. xugon, a yoke j} calyx one.leaved.) **11. —17.**
14677 *Pva Schreb.* Iva **O cu** **4i' au** S. Europe 1759. S & I Fl. gr. 525
14678 *orientalis L.* oriental **A** or **14** my.jn Levant 1732. D s.p Di. el. 53. 61
14679 *pyramidalis L.* pyramidal **A** or **4** my.jn Britain sc.mo. D s.p Eng. bot 1270
14680 *integriflUa D. Don* entire-leaved **A** or **4** jl.au Nepal 1821. D s.l
14681 *alpina L.* alpine **4** my.England moun. D co Eng. bot 477
14682 *genevensis L.* Geneva **4** my. Switzerland 1606. D co Bui. her. 3H1
14683 *foliosa Trat.* leafy **4** my. Switzerl. 1826. D s.l
14684 *rupestrifl Schl.* rock **4** my.jl Switzerl. 1826. D s.l
14685 *rcptans L.* common creeping **4** my.jn Britain moi.w. D n.p Eng. bot. 489
2dlba **4** my. Britain moi.w. D s.p
3 rubra **4** my. Britain moi.w. D s.p
14686 *australis AJ?* southern **4** my. N. Holl. 1822. D s.l
14687 *Chamelpitya Schreb.* Ground Pine **4** ap.j England san.fi. S s.l Eng. bot. 77
1647. ANISO'MELES R. Br. ANISOMELES. (*Anisos*, unequal, **1** (member; anther.) **Labita. 4.-7.**
14688 *malabarica R. Br.* Malabar **n.** or **I*** jl.au V E. Indies 1817. C co R. maL 10.93
Ncpeta malabárica L.
- 14689 *ovata R. Br.* ovate-leaved **O** **1** jl.au Pk E. Indies 1823. S l.p **Bur. zey. 71.1**
14690 *furcata Lk.* forked ***** jl.au B Nepal 1824. C co
- 14691 *A'juga furepta L.* musk **2** jl.au P N. Holl. 1824. C s.l

* 1648. TEITCR/I/Af L. GERMANDER. (*Teucer*, a Trojan prince, first used it in medicine.) *Labihtce*. 50.—87.

- I. AXILLARIFLURA. — *Flowers axillary, solitary or in pain*,
- 14692 *frcticans L.* narrow-lvd shrubby * J or **3** ns V Spain 1640. C r.m Di. el. 284.366
14693 *latifoliu L.* broad-lvd tree **J** or **3** h.s V Spain 1640. C r.m Bot mag. 245
14694 *hetcophylum Jfertt.* various-leaved **J** or **2** jnjl P Madeira 1759. C r.m
14695 *Laxmanni Muir* Laxmann's k A or **1** jn.au Va Siberia 1800. C co W. & K. 1.G9
14696 *breviflum Schreo.* short-leaved **1** or **1** jnjl Pk Crete 1824. C co Fl.gr. 528
14697 *asiaticum L.* Asiatic **S** **1** or **2** n.o P 1777. C r.m Jac.vin.3.41
14698 *cubense L.* Cuba **jg** **D** or **14** my. P Cuba 1733. C co Jacob.2.30
14699 *nissolianum L.* Nissolian **1** **Vjl** P Spain 1752. D co M. h. 22.19
14700 *Pseodo-Chamaypitsl., Bast.Gr.P.* **n.** or **4** jn.jl P S. Europe 1820. C s.p Sab. rom.3.25
14701 *campanulatum L.* bellflowered **9t A o'** **1** jl.au W Levant 1728. D co

II. VEBTICILLFLORA. — *Flowers ivhorled, whorls for the most part divided in halves.*

- H702 *rcgium L.* royal **tL** **I** or **14** my.o P Spain 1699. C Tm Pluk. al 65.1
14703 *multiflorum L.* many-flowered **A** or **1** jl.s L.R Spain 1731. C co Boc.mu. 117
14704 *Iticidum L.* shining **5** A or **lijn.s** P S. Europe 1730. C r.m Mag. h. 52
14705 *ChamsMrys L.* **Germander** **A** or **i** my.au P England old w. C co Eng. bot. 680
14706 *5c6rdium L.* Water Germanander * A or **4** jl.au P England mar. D r.m Eng. bot 828
14707 *scordiflides Schreb.* Scordium-like **A** or **4** jnjl P Crete 1819. D s.l
14708 *spin&sum X.* thomy O or **4** my.jn W Spain 1640. S co Cav.ic.1.31
14709 *Butrys L.* Botrys O or **4** my.jn R S. Europe 1633. S co Ger. em. 525.*

III. TRIFLORUM. — *Peduncles axillary, S-flowered.*

- 14710 *trifidum Retz.* tri&d Jeaved tt- L J or **1J** jn.au P C. G. H. 179L C r.m
- IV. CAPITOPHYLLFLORA. — *Flowers capitate; leaves entire from No. 14717. crenated.*
J2J? ?! dwarf **2** **jhl** P Spain 1818. C co Bar. ic 1080
J2J? ?! dwarf **4** jl.au P Spain 1816. C co Bar. ic. 1092
14713 *Jhymifblium&Airk* Thyme-leaved «L **4** jn.o P Spain 1816. C co
14714 *supinum L.* **tupinc** tt **4** jn.o W Austria 1752. C co
14715 *montanum L.* **dwarfmountain».** **4** jl.o W S. Euroi>e 1710. C co Jac.au. 5.417
C co Fl.gr. 534

- | | | | | | | | |
|--|--|------------------------|--|-----------|-----------|-----------------------|-----------------------|
| 14785 | Nepctella L.
amethystina Poir. | small Nepete | ^ A cu | 1 jLau | R | S. Europe 1758. | D co |
| 14786 | gravoltens Vil.
patella All. | heavy-smelling | £ A cu | Ijjl.au | P | S. Europe 1804. | D co |
| 14787 | suaveolens Roem. | sweet-scented | ■ A or | 1} jn.au | B | 1817. | D co |
| 14788 | amethystina Desf. | amethystine | A or | \} jjl.au | B | S. Europe 1816. | O co |
| 14789 | /amifolia IV. en. | Lamium-lvd | A un | 1Jjl.au | P | Armenia 1806. | D co |
| 14790 | angustifolia Vahl
arragonensis Lam. | narrow-leaved | A cu | 2 jn.jl | p | Spain 1798. | D co |
| 14791 | Catrtia L. | | | | W | Britain ro. aid. | D co |
| 14792 | macroixa Led. | long-tailed | | | W,p | Siberia 1820. | D co |
| 14793 | ucknna L. | Ukraine | | | | Ukraine 1798. | D co |
| 14794 | nuda L. | naked | | | | S. Europe J713. | D co |
| 14795 | pannonica Jac. | Fannonian | ■ A cu | 4 au.o | R | Hungary 1683. | D co |
| 14796 | pannonica L.
paniculata Cr% | | | | | Jac.au. 2. 129 | |
| 14797 | camptlea H. K. | blue | ■ A cu | 1J my.jn | B | 1777. | D co |
| 14798 | violacea L. | violet-colored | A cu | 2 jjl.s | B | Spain 172a. | D co |
| | latif&Ha Dec. | broad-leaved | A or | 4 jlau | P | Pyrenees 1816. | D co |
| 14799 | grandiflora Lap. | | | | | Boc. mu. 36 | |
| 14799 | crispa W. | cmled-leaved | £ A cu | 2 jl.au | Pa.B | Levant 1800. | B co |
| 14800 | imbricata Lag. | imbricate | 5 A or | 2 jl.au | B | Spain 1820. | D sl |
| II. VEHTICILLIFLORJE.— <i>Flowers in whorls.</i> | | | | | | | |
| 14801 | teucriifolia W. en. | Teucrium-lvd | ^ A or | 14 jl.au | P | Armenia 1816. | Deo |
| 14802 | diffusa Fis. | diffuse | ik A or | Ijjl.au | P | Siberia 1824. | Deo |
| 14803 | teucrioides Lam. | Teucrium-like | 5 A or | 1J jl.au | W | S. Europe 1820. | Deo |
| 14804 | marrubiotdes IV. en. | Horehound-lk | 4 A or | Ijln.au | R | Italy 1640. | Dpl |
| 14805 | itálica L. | Italian | 5 A or | 1 jn.au | R.w | Algers 1817. | Deo |
| 14806 | multibracteatallo/. | many-bracted | * A or | 3 jlau | P | Caucasus 1816. | Deo |
| 14807 | suptaa Steu. | supine | 31 A or | 1 jlau | B | S. Europe 1774. | Deo |
| 14808 | lanata Jac. | woolly | 3f A un | Ijmy.jn | P | Spain 1683. | Deo |
| 14809 | tuberftsa L. | tuberous-rooted | * A un | 2 jn.au | V | Morocco 1801. | Deo |
| 14810 | reticulata Desf. | netted | 5 A un | 2 jl.au | P | N. Africa 1817. | Deo |
| 14811 | Scordbitis L. | Scordotis | 3t A or | Ijjn.au | | | Alp. ex. 283 |
| III. SPICATFLOCH.— <i>Flowers in spikes.</i> | | | | | | | |
| 14812 | multiflida L. | | | | | 1796. | D co |
| 14813 | dotyotdea UK. | | | | | Gin. si. 3.55 | |
| 14814 | malabárica Pers. | | | | | S co | Cav. ic. 1.49 |
| | | | | | | S.s.l | R. mal. 10.93 |
| 1655. ELSHO'LTZ/4 | | | | | | | |
| 14815 | ocymoldes Pers. | W. ELSHOLTZIA. | (J. S. Elsholtz, a Prussian botanist.) | | | Labihta. 3.-5. | |
| 14816 | cnstAta IV. | Basil-like | J^ O un | 1 jl | P | E. Indies 1824. | S co |
| 14817 | paniculata W. | crested | O or | II my.jl | Pk | Siberia 1789. | IS co |
| | Jtfysppus cristatus Lam. | large-panidied | ± QJ or | 3 jlau | Pk | E. Indies 1820. | S.s.l |
| 1656. LAVA'NDULA L. LAVENDER. (<i>Lavo, to wash j use of its distilled water.</i>) | | | | | | | |
| 14818 | Spka W. | common | 6pike | tt. | dt | Labiate. 11.-12. | |
| | 2 alba | | | | | C 81 Schk. nán. 2.157 | |
| 14819 | latifolia Ehrh. | white-flowered | tt. | or | lls | Li | S. Europe 1568. |
| 14820 | StoTchias L. | broad-leaved | tt. | dt | la | W | C s.l |
| 14H21 | viridis Merit. | Stoechas | tt. | J or | lls | Li | S. Europe 1568. |
| 14822 | dentata L. | green | tt. | — or | U my.jn | P | S. Europe 1562. |
| 14K23 | pinnata L.JU. | tooth-leaved | tt. | — or | U my.jn | Li | Madeira 1777. |
| 14824 | multiflida L. | pinnated | tt. | — or | apau | Li | Spain 1597. |
| 14825 | heterophylla Via. | many-cleft | IQJor | 1 jjl.s | Li | Madeira 1777. | |
| 14826 | formbsa Lk. | various-leaved | — or | 2 my.au | Li | Canaries 1597. | |
| 14827 | abrotanoides Lam. | beautiful | — or | 2 ap.au | Li | 1816. | |
| 14828 | cambsa W. | Southernw.-lk | tt. | — or | U jn.s | Canaries 1699. | |
| | | (toshly-leaved) | g Oil | 1 jn.jl | Li | E. Indies 1788. | |
| 1657. SIDERRTIS L. IRONWORT. (<i>Sideros, iron; sup. to cure wounds by iron weapons.</i>) <i>LabicUce. 26.—43.</i> | | | | | | | |
| I. EBRACTEAVA.— <i>Bracteas under calyx none.</i> | | | | | | | |
| 14829 | Canary | \$H I or | 3 my.au | Y | Canaries | 1697. | C r.m Jac. vin. 3. 30 |
| | whitish | cu | 3 ap.jl | Y.Br | Madeira | 1714. | C r.m Com. h. 2.99 |
| 14830 | Cretan | tt.-J | 1 jn.s | W | Candia | 182a | C co |
| 14831 | mountain | O or | 1 jlau | Y.Br | Austria | 1752. | S co Jac. au. 5.434 |
| 14832 | elegant | O or | li jl | V | | 1787. | S co Mur. 1.4. |
| 14833 | Roman | O or | 1 jn.au | W | Italy | 1740. | S co Cav. ic. 2.187 |
| 14834 | decumbens | -* O or | } jlau | W | C. G. H. | 1820. | S s.l |
| II. BnACTEA'r'a.— <i>Bracteate, bracteas entire.</i> | | | | | | | |
| 14835 | colored-calyxedin. | or | 1 jlau | Pa.Y | Russia | 1821. | D co |
| 14836 | Syrian | — or | 1J jn.s | Y | Levant | 1597. | C r.m Sab. rom. 3.40 |
| 14837 | Taurian | cu | Ji jn.s | Pa.Y | Tauria | 1822. | C co |
| 14838 | perfoliate | ^ A | au.n | Y | Levant | 1731. | C co |
| 14839 | white-flowered | tt. | 1 jlau | W | Spain | 1823. | D co Cav. ic. 30* |
| III. DEVTA'^IE.— <i>Bractcate, bracteas toothed.</i> | | | | | | | |
| 14840 | hoary | tt. | 1J jlau | ■ | Spain | 1752. | C co Cav. ic. 2.186 |
| | narrow-leaved | | 1 jlau | ■ | Spain | 1820. | D co |
| 14841 | spiny | | lijn.s | ■ | Spain | — | C co |
| 14842 | Holly-leaved | | j'-s | ■ | Levant | 1818. | C co |
| 14843 | Hyssop-leaved | | 1 jn.s | T | Spain | 1818. | D co |
| 14844 | Pyrenean | | 1 jn.n | L.Y | Pyrenees | 1597. | C co Schk.han.2.1 |
| | | | 1 jn.n | Y | Pyrenees | 1827. | C co |
| IV. EBRACTEABA.— <i>Bracteas under calyx none.</i> | | | | | | | |
| 14845 | alpine | ■ | 1 jnau | Y | a Franco | 1822. | D co |
| 14846 | fetid | ■ | 1 jn.n | Y | Spain | 1822. | D co |
| 14847 | Germander-lvd | ■ | 1 jn.au | Y | Spain | 1816. | D co |
| 14848 | Scorciium-hke | ■ | 1 au.n | Y | France | 1597. | C co Bar.ic.S< |
| 14849 | hairy | ■ | U jn.jl | Y | S. Europe | 1731. | C co Cav. ic. 4. SW |
| 14850 | curled-leaved | ■ | li jn.s | Y | Gibraltar | 1816. | C co |
| 14851 | woolly | ■ | 1 jn.au | V | Egypt | 1818. | S co Her.ser.i |
| | | | | | | | |
| <<♦ i lllcrit. BYSTBOPOOON. /Bu, to close, pogon, beard: throat of fl. <i>Lahiataf.</i> | | | | | | | |
| 14855 | plumiflous iferrf. | feathery, fjoirrreddt. | LJ or | li jn.jl | Pa.P | Canaries 1779. | C >1 Her.tfr. |
| 14856 | onganifolius Merit. | Onganumlyvd tt. | L J or | If ilan | Pa.P | Canaries 1815. | C fik Her.ser.i |

14857 canariensis Herit.	Canary dotted	tt.il —	or Ujn.au ■ ■	Pa.P	Canaries	1714.	C	p.l	Com. h. 2.65
14358 punctatus Nerit.		■ ■	jl.a	Pa.P	Madeira	1775.	C	p.l	Her.ser.7
* 1659. AfE'NTHil L. MINT. (<i>Mintha</i> , daughter of Cocytus, changed into this plant) <i>Labittte.</i> 52.-62.									
\$ 14859 quadrifolia Box.	four-leaved	Ljor	2 jl.au	P	Nepal	1830.	C	s.1	
14860 viridis L.	green spear	A cul	2 au	P	Britain	mar.	D	co	Eng. bot 2424
14861 levigata W. en.	polished	A or	jl.jl	P		—	D	co	
14862 tenuis Mx.	slender	A °r	1 jl.au	P	N. Araer.	—	D	co	
14863 crispata Schr.	crumpled	If	jl.au	P	1807.	D	co	
14864 nepetoides LeJ.	Nepeta-like	li	jl.au	P	Bilgio	1890.	D	co	
14865 niperita L.	pepper	2	aus.s	P	England	watpl.	D	co	Eng. bot. 687
14866 Aalsamica W. en.	Buteam-scented	jl.jl	au	P	Italy	1804.	D	co	
14867 ninaca Jac.	Nile	I	jl.au	P	Egypt	1796.	D	co	Jac. vin. 3.87
14868 sylvestris L.	wood	2	jl.au	Li	Britain	watpl.	D	co	Eng. bot 686
14869 pubescens W. en.	pubescent	li	jl.au	P	Spain	1800.	D	co	Lob. ic. 511
14870 canescens Both.	canescent	li	jl.au	P	Britain	—	D	co	FL dan. 484
14871 nemortaa W. en.	grove	2	jl.au	P	—	D	co	
14872 hirta W. en.	shaggy	11	au.s	P	—	D	co	
14873 gratissima W.	most grateful	n	jl.au	P	Germany	1799.	D	co	
14874 Auricularia L.	Indian eared	2	jl.au	P	E. Indies	1796.	D	co	Ru. am. 6.16
14875 divaricata Lag.	divaricate	2	jl.au	P	Spain	1824.	D	co	
14876 cocinea Box.	scarlet	1	jn.au	S	E Indies	1823.	D	s.p	
14877 crispa L.	curled	2	jl.au	P	Siberia	1640.	D	co	
14878 incana IV. en.	hoary	U	jl.au	P	1790.	D	co	
14879 undulata W. en.	wave-leaved	I*	jl.au	P	1816.	D	co	
14880 rotundifolia L.	round-leaved	2	aus.s	R	England	moi.pl.	D	co	Eng. bot. 446
2 variegata	variegated	2	aus.s	R	—	D	co	
14881 macrostachya Ten.	long-spiked	2	jl.au	P	S. Europe	...	D	co	
rotundifolia W. en.									
14882 rugusa Both.	wrinkly	2	jl.au	P	Germany	1800.	D	co	
14883 lavandulaKicea W. en.	Lavender-Jwf	1	jl.au	P	Spain	1823.	D	co	
14884 capensis T/tun. Cape		1	jl.au	P	C. G. H.	1816.	D	co	
II. CAPITIFLORJE. — Whorls of flowers by Mate.									
14885 hirsuta L.	hairy water	1*	*	■	Watpl.	D co	Eng. bot. 447		
aquatica JL									
14886 pyramidalis Ten.	pyramidal	■	■	P	Naples	1824	D	co	
14887 acutifolia Sm.	sharp-leaved	■	■	Li	Britain	wat pi.	D	co	Eng. bot. 2415
14888 borealis Mx.	northern	■	■	P	S. Amer.	1820.	D	co	
14889 odorata Sm.	scented Bergamot	^ A or	■	P	England	watpl.	D	co	Eng. bot. 1025
citrate Ehrh.									
III AXILLARIFLORA — Whorls of flowers axillary, remote.									
14890 arvensis H. K.	corn-field	■	■	■	Britain	cor. fl.	D	co	Eng. bot. 2119
... 2pravox Sole	early-flowering	■	■	■	Britain	...	D	co	Sol. min. c. ic.
14891 agrestis Ten.	toothed	■	■	■	Britain	m. fl.	D	co	Eng. bot 2120
J4892 dentate W. en.	Austrian	■	■	■	Germany	1816.	O	co	
14893 austriaca Jac.	gentile	■	■	■	Germany	1809.	D	co	
14894 gentMisL.	common red	■	■	■	Britain	pools.	D	co	Eng. bot. 2118
14895 rubra Am.	Baden	■	■	■	Britain	wat pi.	D	co	Eng. bot. 1413
14896 badensis Gm.	bland	■	■	■	Lapland	1810.	D	co	
14897 blfinda Wai.	northern	■	■	■	Nepal	1824.	D	co	
14898 borealis Mx.	Canadian	■	■	■	N. Amer.	1800.	D	co	
14899 canadensis L.	southern	■	■	■	N. Amer.	1801.	D	co	
J4900 australis B. Br.	Pennyroyal	■	■	■	N. Holl.	1820.	D	co	
J4901 Pulegium L.	stag Hyssop-lcr.ved	§ L A or	■	■	Britain	wet co.	D	co	Eng. bot. 1026
J4902 cervina L.	cultivated	^ A or	■	■	France	1648.	D	co	M.li.3.7.7
H9U3safrivaZ.	smoothed	■	■	■	England	1) co	D	co	Eng. bot 448
14904 glabrata W.	slender	■	■	■	Egypt	1802.	D	co	
14905 Kracilis Sm.	marsh	■	■	■	Britain	wat pi.	D	co	Eng. bot 449
14906 paldstris Sm.	villous	■	■	■	Britain	mar.	D	co	
14907 villosa Sm.	meadow	■	■	■	Britain	wat pi.	D	m.s	
14908 prafensis Sm.	river	■	■	■	Britain	mea.	D	m.s	Sol. min. 17
14909 rivalis Sm.	marsh	■	■	■	Britain	riv. sid.	D	m.s	
14910 paludosa Sm.					Britain	mar.	D	m.s	
1660. PERILLA L.									
14911 ocytoides W.	PERILLA.	O cu	ijl.au	W	India	1770.	S	s.l	Bot. mag. 2395
14912 fruticosa Box.	Basil-like	■>Ljor	6 jl	—	Nepal	1823.	C	p.l	
1661. HYPTIS Jac. HvFTis. (<i>Hyptios</i> , resupinate; limb of corolla turned on its back.) <i>Labitta-</i> 19.—29									
14913 obmacrata B. Br.	bractless	"	" JV ee	■	Pa.P	Wr Indies	1778.	I	Ap.mil.**.QO.2
14914 suay^olens Jac.	sweet-scented	£	JO un	■	W. Indies	1800.	D	co	An. mu. 7.39.2
14915 Pseud.-6'hmma>L>0ft. Bast.	Ger-lyde	OI un	■	—	W. Indies	1821.	D	co	An. mu. 7.31.1
14916 ChammMrys W.	Germander-/-^	OI un	■	■	Guiana	1821.	D	co	An. mu. 7.27.4
14917 radi^ta W.	radiated	^ iAI cu	■	■	Pa.P	Carolina	1690.	D	An. mu. 7.27.2
14918 capsita Jac.	ttcipitate	tf OI	■	■	Pa.P	W. Indies	1714.	I	An. mu. 7.27.1
14919 polyanthos Poit.	many-flowered	^ A	■	■	...	Peru	1819.	D	co
14920 inflate Spr.	inflated calyx	tf	■	■	...	Brazil	1823.	D	co
Marsypianthus hypothec Mart.	short-stalked	■	■	■	Li	S. Amer.	1822.	D	
14921 recurvata Poit.	recurved	■	■	■	Pa.P	Cayenne	1820.	D	
14922 polystachya Hum.	many-spiked	■	■	■	...	Mexico	1824.	D	
14923 canescens Hum.	canescent	■	■	■	...	Caraccds	1818.	D	
Teicrium rhombifolium W.	pectinate	■	■	■	Pa.P	W. Indies	1776.	D	An.mu.7.30
14925 pectinate Poit.	pectinate	■	■	■	...				
Arpeta pectinata L.									
14926 allida Kth.	whitish	tt. leu	■	■	W	Mexico	1824.	D	
14927 spicata Poit.	spiked	£ O)un	■	■	W	W. Indies	1820.	D	
14928 Plumieri Poit.	Plumier'8	jg Olun	■	■	W	W. Indies	1819.	D	
J4929 stachyldcs Lk.	Stachys-like	mcu	■	■	...				
JJigPersica W.	Persian	tL Jcu	■	■	Pa.P	Persia	1800.	I	Lin. (r. 6.12
1*931 scoparia Poit.	Broom	£ Olun	■	■	...	Maranh.	1825	I	An. nm. 7.31.2

*J£7°. ZIETEWIL GLETZ. ZIETENIA. (Probably the name of some obscure botanist.) *Labiatae.*
 15024 *lay&ndulifolia* Pen. Lavendercr.lvd TL or 4jl.au P Levant 1824. C co
 orientalis Glcd. Stachys lavandulifolia W

		STINKING HOREHOUND.	(<i>Balto</i> ,	to reject:	offensive odor.)	<i>Labiate.</i>	5 . - 7
502		common	3t A ^u	2 jl.au	P Europe	D s.l	
		black	^ A w	2 jl.s	P BriU.n	... bed.	D co
J5027 & Iba "Z" ***		white	* A *	2 jl.s	W Britain	... D	co
2 1505 cinerea D. Don		grey	k A ^m	2 J ^l .au	p Nepal	1824. C si	
Iau29diaticha W.		two-rowed	^ w	1 jl	B India	1828. S CO	
* 1672. A/ARRIPBIUM L.	HOHEHOUND. [Marrob, a bitter juice, Heb.]	<i>Labila</i> Ue.	20.— 81.				
	I. PEINTADOV. — <i>Calyx five-toothed.</i>						
?2S ^a Af Skum L.	Alyssum	* A ■ UJ ■	P Spain	1597. C 8J	Ger. h. 579.1		
j.-31 astracanicum Jac.	Astracan	■ ■ ■ ■ ■	Pa.P Levant	1816. D co	Jac. ic. 1.109		
?2S ^b peregrinum L.	foreign'	■ ■ ■ ■ ■	W Sicily	1640. I) co	Jac.au. 2.160		
\£? c ^t ticum Mil.	Cretan	■ ■ ■ ■ ■	Levant	D co			
JO34 candidissimum L.	whitest	■ ■ ■ ■ ■	Levant	1732. D s.p	DL el. 274.214		
i50.15 molissimum D. Don	softest	■ ■ ■ ■ ■	Nepal	1820. C p.l			
?S ^f a ^t > e Wont.	kindred	■ ■ ■ ■ ■	Siberia?	1822. D co			
?5037 supticum L.	supine	■ ■ ■ ■ ■	S. Kurope	1714. I) co			
?J/w8 catanaRilium & fr.	CatminUcaved	■ ■ ■ ■ ■	Levant	1819. n co	Boc. mu. 2.96		
?J/a9 leonurus Desr.	Lconurus-like	■ ■ ■ ■ ■	Caucasus	1819. D co			

				— Calyx ten-toothed.				
Vulgaris L. P. Ten. " Hispanicum L. 15047 orientale Spr. J9allula onentalis V15048 acutabellum&um L.	African common white Apulian hairy cinereous curled-Avzwrf Spanish Preoriental saucer-leaved	P W W Pa.P. Pa.b Pa.P & Europe P P	C. G. H. Britain Italy Spain Spain Spain S. Europe	1710. rubble. 1818. 1823. 1714. 1714. 1821.	D D D D D D C	p.l co co co co co p1	Com. h. 2.90 Eng. bot 411 Her. par. 600 Her. par. 201 Ber. jo. 129	

IJO4.)	Pseudodictamnus L.	Bast	Dictamnus.	J or	lfjl.ju	R	Candida	1596.	C	P-1	Lam.iL	508.2
1673.	LEONU ^A RUSi,	MOTHERWOBT.			a lion, oura, a tail; spike of flowers.)					Labiatet.	14.	
15050	Card ^A aca L.	cardiac	3		3 jl.au	R	Britain	gra ba. S	co	Eng. bot.	286	
15051	vill ^o rus Desf.	villus			3 jl.au	P	Tauria	1820. S	co			
15052	Marrubi ^A strum L.	Marrubiastrum			2 jn.au	P	Austria	1710. S		Jac. au.	5.405	
15053	crispus Mur.				2 jl.au	R	Siberia	1658. S		Mur.	8.4	
15054	supinus L.	supine			1 jn.au	R	Siberia	1816. S				
15055	tatancus L.	Tartarian			S au.o	F	Russia	1756. S				
15056	sibiricus /.	Siberian			S jn.au	R	Siberia	1759. S				
JS5	condensatui Horn.	condensed			3 jl.au	P	Siberia	1819. S				
15058	alt ^A etus Spr.	Altaian			2 jl.au	P	Altai	1824. S				
JS5»	> ⁸ ectus Schult.	neglected			3 jl.au	P	S. Europe	1818. S				
YEP	multidens Desf.	many-cleft			1 jl.au	P		1817. S				
J2Si	heterophyllus Swt.	various-leaved			4 jh.au	P	8. Amer.	1819. S				
15062	ltereru l.tndl.	torn			3 Jl.au	Pk	Nepal	1884. S	co			
15x	tanakuti Pcrs.	woolly	5 A		8 jn.au	Y	Siberia	1752. S	go	Gm.si.a	54	
		Moen' flajeta lanjia	or									
		Y	L	multifida								

15073 <i><alviB>folia Jac.</i>	Sage-leaved	*	or	3	jn.au	P	—	—	C co	Jac. BC. 3.359		
15074 <i>microphyllea Sieb.</i>	small-leaved	£	A or	3	jl.au	Pa.br	S. Europe	1827.	D pl			
15075 <i>orientalis Dec.</i>	oriental	jk	A or	3	Jnl	Y	Levant	1757.	D r.m	Mil ic. 2.204		
15070 <i>Nissolia W.</i>	Missole's	£	Al or	2	Jnl	Y.Br	N. Africa	1714.	D pl	Bot. mag. 1891		
15077 <i>samia L.</i>	Samian	£	A or	3	jnl	Br	Levant	1818.	D co			
15078 <i>lunaria Sm.</i>	Honesty-leaved^	A or		3	jn	Br	Levant	1821.	D r.m	Bot. mag. 2542		
16079 <i>Russellina Lag.</i>	Russell's	£	A or	3	jl	Br	Levant	1821.	D r.m	Bot. mag. 2542		
<i>lunaria JS Russellidae B. M.</i>												
15080 <i>cinnabarinum Cav.</i>	hair-bracted	k	A or	E		Pa.br	Spain	1820.	D co	Cav. ic. 3.247		
15081 <i>herba venti L.</i>	wind herb	£	A or	E		R	S. Europe	1594.	D co	Bot. mag. 2449		
15082 <i>pungens W.</i>	pungent	£	A or	E	jl	Br	Armenia	1820.	D co	Sw. fl. gar. 33		
15083 <i>tuberosa L.</i>	tuberous	£	A or	E	jn.o	L.P	Siberia	1759.	D co	Bot. mag. 1555		
15084 <i>alpina J.-att.</i>	alpine	£	A or	E	jl	P	Siberia	1812.	D s.l	Pal. p. 2.13		
15085 <i>laumata L.</i>	vaged4eaved	£	A or	E	jl	P	Levant	1731.	D co	Sw. fl. gar. 24		
1675. LEUCAS JI. Br. ^LEUCAS. (<i>Leukosy white; downy whiteness of flowers.</i>) <i>Labi&t*.</i> 11. — 27.												
15086 <i>chinensis R. Br.</i>	China	O o r	lfjn.o	W	China	1820.	S s.l					
/*hilmis chinensis Retz.												
15087 <i>zeylanica R. Br.</i>	Ceylon	O un	E	P	E. Indies	1777.	E	E. Indies	Jac. ic. 1. III			
15088 <i>/mifolia Spr.</i>	Flax-leaved	O or	E	P	E. Indies	1816.	E	E. Indies	s.l			
Philmis zeylanica Mur.												
15089 <i>Plukenetti Spr.</i>	Plukenett's	O o r	3	jls	W	E. Indies	1820.	S s.l				
Philmis Plukenetti Roth												
15090 <i>cephalotes Spr.</i>	round-headed	O or	2	jl.s	W	E. Indies	1818.	S s.l				
15091 <i>abera Spr.</i>	rough-leaved	fOI un	1	jl.au	W	Caramanian	1818.	S s.l				
15092 <i>indica R. Br.</i>	Indian	O un	Ujl.au	W	E. Indies	1789.	S	B.I				
15093 <i>urticifolia R. Br.</i>	Nettle-leaved	O un	Ujl.s	W	E. Indies	1810.	S	s.l				
15094 <i>flaccida R. Br.</i>	flaccid	iQJor	2	jl.s	W	N. Holl.	1823.	S	s.l			
15095 <i>martinicensis R. Br.</i>	Martinique	O	E	W	W. Indies	1781.	S	8.1	Jac. ic. 1.110			
Philmis martinicensis IV. caribaea Jac.												
15096 <i>biflora R. Br.</i>	two-flowered	£	OJ or	2	jl.s	W	Ceylon	1819.	S	s.l		
1676. LEONCTTIS R. Br. LION'S EAR. (<i>Leontis, a lion, eus, an ear; fanciful likeness.</i>) <i>LabidUe.</i> 4.												
15097 <i>Leonurus R. Br.</i>	Lion's tail	E	LJor	3	o.d	S	C. G.H.	1812.	C p.l	Bot. mag. 478		
.Philmis Leonbrus L.												
15098 <i>ovata Spr.</i>	ovate	E	!_J or	If	jn.jl	O	C. G. H.	1713.	C	p.l	Mil. ic. 2.162.1	
J>hilmis Leonuis Thun.												
15099 <i>intermedia Lindl.</i>	intermediate	H	E or	3	s.o	O	C. G.H.	1822.	C	p.l	Bot. reg. 850	
15100 <i>nepetifolia R. Br.</i>	Catmint-leaved	O	E or	3	s.o	O	E. Indies	1778.	S	s.l	Bot. reg. 281	
* 1677. MOLUCCELLA L. MOLUCCA BALM. (Brought from the Moluccas.) <i>Labiott.</i> 4.—7.												
h 15101 <i>spinosa L.</i>	prickly	O cu	1	jl.au	C	Levant	1596.	S	co	Lam. il. 510		
15102 <i>laevis L.</i>	smooth	O cu	1	jl.au	Pa.P	Syria	1570.	S	co	Bot. mag. 1852		
15103 <i>Marrubiastrum Step.</i>	Marrubiastrum	O or	1	jl.au	P	Siberia	182a	S	s.p			
15104 <i>tuberosa Pall.</i>	Juberous-roota/A	A cu	2	jl	Pa.P	Tartary	1796.	S	1.p	Pal. it. 3. T.		
' 1678. CLINOPODIUM L. WILD BASIL. (<i>Kline, bed, pou, foot j</i> fl. like turned bed feet.) <i>Labihle.</i> 3.—4.												
15105 <i>vulgare L.</i>	common	k	A or	1	Jn.au	Pk	Britain	gra.ba.	D	co	Eng. bol. 1401	
15106 <i>ffigyttaicum Lam.</i>	Egyptian	A A or	1	jn.au	P	Egypt	1759.	D	co			
15107 <i>onganitium Lab.</i>	Origanum-lvd	k A or	i	jn.au	Pk	S. Europe	1825.	D	p.l			
• 1579. PYCNA'NTHEMUM Mx. PVCNANTIIBMUM. (<i>Pyknos, dense, anthemos, flower.</i>) <i>LabihUe.</i> 7.—11												
15108 <i>incanum Mx.</i>	hoary	3	JLo	W	N. Amer.	1732.	D	e.o	Di. el.	74.85		
15109 <i>Monardisfo Mx.</i>	Small Monarda	E	2	jl.au	W	N. Amer.	?1816.	D	s.p			
15110 <i>anstatum Mx.</i>	awned	E	2	au	W	N. Amer.	1752.	D	co	Mic. am. 2. S3		
15111 <i>verticillatura Pers.</i>	whorled	E	2	jl.au	W	Carolina	1820.	S	s.p	Mic am. 2.31		
15112 <i>nfidum Nut.</i>	naked	E	2	JLau	W	Carolina	1824.	D	s.p			
15113 <i>/infidum Ph.</i>	Flax-leaved	E	Ujl.au	W	N. Amer.	1739.	D	co	Her. par.	218		
15114 <i>lanceolatum Ph.</i>	spenr.leaved	E	A or	1	jl.au	W	N. Amer.	1812.	D	co		
* 1680. ORIGANTJMZ. MARJORAM. (<i>Oros, a mountain, ganos, joy ; odor and habitation.</i>) <i>Labiate.</i> 20.—24												
15115 <i>vulgare L.</i>	common	A cul	2	jn.o	Pk	Britain	ch. w.	D	s.l	Eng. bot. 1143		
2 fibre albo	white-flowered	E				Britain	m.woo.	D	co			
15116 <i>hirtinile Poir.</i>	dwarf	E				1818.	D	co			
15117 <i>virens Lk.</i>	green	E				Spain	1824.	D	co			
15118 <i>nonnale D. Don</i>	normal	E				Nepal	1819.	D	co			
15119 <i>heraclicotum L.</i>	bast. Winter-sweet	E				W. Europe	1640.	D	s.l	Lob. ic. 492		
15120 <i>criticum L.</i>	Cretan	E				S. Europe	1596.	S	s.l	Schk. flan. 2.16*		
15121 <i>mogastachyum Lk.</i>	large-spiked	E				Pk	S. Europe	1823.	D	co		
15122 <i>smyrna'urn L.</i>	Smyrna,	E				W	Smyrna	1722.	C	r.m		
15123 <i>hirtum Lk.</i>	hairy	E				Pk	Levant	1823.	D	co		
15124 <i>oblongatum Lk.</i>	oblong	E				W	1823.	D	co		
15125 <i>Onych L.</i>	Onites	E				Pk	Sicily	1759.	D	co	Boc. mu. 38	
15126 <i>syriacum L.</i>	Syrian	E				Pk	Syria	1823.	S	co		
15127 <i>Majordna L.</i>	Sweet Marjoram	E				Pk	Portugal	1573.	S	r.m	M.h. 11.3.1	
15128 <i>Man L.</i>	Mastic	E				Pk	1823.	C	co	Bot. mag. 2005	
majoranoides W.												
15129 <i>Indicum Roth</i>	Indian	^ OJ or		jl.jl	W	E. Indies	1816.	S	r.m			
15130 <i>Tournefort'8 U. H. K.</i>	Tournefort'8	« or		au.s	P	Amorgos	1788.	C	co	Bot. rep. 537		
15131 <i>sip'leum L.</i>	Mount Sipylus	n.	or	jns	Pk	Levant	1699.	C	r.m	Her. lug. 463		
15132 <i>Dictamnus L.</i>	Dittany of Crete	—	or	jn.au	Pk	Candia	1551.	C	r.m	Bpt. mag. 290		
15133 <i>benghalense Gar.</i>	Bengal	«L	I or	jl.jl	Pk	Bengal	1820.	C	1.p	Bur. in. 38.3		
15134 <i>egyptiacum L.</i>	Egyptian	B.	fra	jn.au	Pk	Egypt	1731.	C	co	Alp. a*g 95		
• 1681. THYTTUSL. THYME. (<i>Thymos, courage; smell reviving.</i>) <i>LabihUe.</i> 33.—36.												
15135 <i>villus L.</i>	villous	f	jl.jl	P	Portugal	1759.	C	co	Fl. por. 1.14			
15136 <i>cephal&tes L.</i>	Numidian	f	jl.jl	P	Portugal	1759.	C	co	Fl. por. 1.13			
15137 <i>numfdicus Poir.</i>	Barbary	i	jl.au	P	Barbary	1816.	D	s.l				
15138 <i>Marinosci Ten.</i>	Marinosci's	tt.	or	ijl.au	R	Italy	1824.	D	s.l			
15139 <i>creticus Brot.</i>	Cretan	tt.	or	1 jl.au	P	Mediterr.	1817.	C	8.1			
<< M Zjynaus capitatus Lk.												
15140 <i>odoratissimus J^A.</i>	weetest-scentedtt.	or	E	jn.au	P	Tauria	1817.	C	s.l			
15141 <i>aciulkrisr*.</i>	needle-leaved	fU	or	E	jn.au	Hungary	1806.	C	co	W.&K. 8.1*7		
151B <i>hir<utu Bleb.</i>	hairy	«	or	E	jl.au	Spain	1821.	C	s.l			

15143	Zygis L.	Zygis	tL	or	1	au	P	Spain	1771.	S	r.m	Bar.	1c.	777	
15144	Marschaliciflum W.	Marshall's	tt,	or	jjn.au	P	Crimea	1817.	C	co					
15 HJ	glabratum Lk.	smooth	tt.	or	i	jn.au	P	S. Europe	1823.	C	co	Fl.	por.	15	
15145	angustifolius Schreb.	narrow-leaved**	or	ijnn.au	P					-	C	co			
15147	Serpulium L.	Wild Thyme	U	or	\	jn.au	P	Britain	heaths.	C	s.p	Eng.	bot.	1514	
15148	Ixserens Ehrh.	drawn out	JU	or	1	jn.au	P	Britain	heaths.	D	co				
15149	colonusite A.	hill	£*	or	l	il.au	P	Tauris	1820.	D	co				
15150	montanus Kit.	mountain	£*	or	■	■	St	Hungary	1806.	D	e.p				
15151	nummularius B'teb.	Moneywort	1*	or	i	jnjil	P	Crimea	1822.	C	co				
15152	pannonicus W. en.	Pannoman	■	or	\	jn.au	P	Hungary	1817.	C	co				
15153	lanuginosus Sehk.	woolly	■	or	i	jnau	P	Britain	Scot	C	co				
15154	citriod&rus Pert.	Lemon-scented	£,	or	i	jn au	P			C	co				
15155	lfcidus Ehrh.	nhming-leaved	tt.	or	1	jn.au	P			C	co				
15156	vulgaris L.	common garden*	cul	1	my.au	P	a Europe	154a	C	r.m					
	2 latifolius	broad-leaved	cul	1	my.au	P			C	co					
15157	Mastichina L.	Mastich	tt.	— or	1	jls	Pa.P	Spain"	1596.	C	co				
15158	elongatus Lk.	elongated	tt.	or	1	jls	P		—	—					
15159	tomentosus W. en.	tomentose	tt.	or	1	jn.au	W	Spain	1816.	C	co				
15160	lanceolktus Desf.	lanceolate	tt.	or	ijjla u	P	N. Africa	1823.	C	co					
15161	'Aagoriganum L.	Goat's Origan,	tt.	— or	1	my.jn	P	Candia	1640.	C	co				
15162	ericifolius Roth	Heath-leaved	tt.	or	ijjma	P	Spain	1806.	C	co					
15163	cro&licus Pert.	Croatian	tt.	or	1	jlau	P	Hungary	1802.	D	co				
15164	Piperella L.	Small Peppermints.	or	i	jl.au	P	Spain	1810.	D	s.l					
15163	filiformis W.	filiform	V.	—J	or	jn.jl	P	Minorca	1770.	C	co				
15166	virgatas Ten.	twiggy	■	or	1	jla u	P	Italy	1824.	D	s.l				
15167	fruticulbus Bert.	fruUculoise	or	1	ilau	P	Sicily	1822.	C	s.l					

1682. A CYNO9 Pert. ACT.VOS. (The Greek name of a balsamic plant) Labiate. 11.
 15168 vulgiris Pert. BatU-lud common O or $\ddot{\text{e}}$ jnau-⁹ Britain dryh. S co Eng. bot. 411

<i>Thymus acynoides</i>	<i>G.Don</i> , variable clvd.	O or	t	in.jl	F	Italy	1822.	S	co	
<i>Thymus acynoides</i>	<i>Ten.</i> , heterophylus	Poir.	k	■■■	jnau	F	S. Europe	1776.	s.1	Bot. mag. 2153
<i>Thymus acynoides</i>	<i>Pert.</i>	Paduan	k	■■■	jnau	F	Austria	1731.	s.1	Jac.au. 197
<i>Thymus acynoides</i>	<i>Pers.</i>	alpine	£	Ol or	in.i.8	P		1810,	11	
<i>Thymus acynoides</i>	<i>G. Don</i>	great-flowered	%.	Q>	i jlau	P				
<i>Thymus acynoides</i>	<i>Sm.</i>	apyrous								
<i>Thymus rotundifolius</i>	<i>Pert.</i>	round-leaved	tt.	or	■■■	P	Spain	1820.	■	co
<i>Thymus rotundifolius</i>	<i>Pers.</i>	jmpurascens	k	Q> or	■■■	P	Spain	1820.	■	co
<i>Thymus rotundifolius</i>	<i>G.Don</i>	herb baroni	tt.	or	1 jlau	P	Corsica	1820.	8.1	
<i>Thymus rotundifolius</i>	<i>Sm.</i>									
<i>Thymus suaveolens</i>	<i>G. Don</i>	sweet-scented	O or		1 jlau	P.R	Greece	1817.	S	fl.1
<i>Thymus suaveolens</i>	<i>Sm.</i>									
<i>Thymus villosus</i>	<i>Pers.</i>	villous	O or		ij nau	R	Germany	1817.	S	co
<i>Thymus villosus</i>	<i>Lk.</i>	heavy-scented	tt.	or	1 jn.au	P	Crimea	1820.	S	co

•163. CALAMI'NTHA Ph. CALAMINT. (*Kalos, bciutifUl, tmintha, mint.*) *LabiittB.* 8.-9.
 «179 grandiflora Pers. great-flowered ^ A or 1 jns R Italy 1596. D co Bot mag. 208

1517 grandiflora Pers.	green-flowered	A or	I jns	R			
1518a grandiflora B. M.							
1518a caroliniana 5w.	Carolina	k A or	1 jnjl	F	Carolina	1801.	D co Bot mag. 997
Thymus grandifl. A Af.							
15181 vulgaris Sw.	common	k A or	2 jl.au	BR	England	bor.fi.	D 81 Eng. bot 1676
Melissa Calamintha L.	Th^mus Calamintha E.B.						
15182 A epeta Ph.	Nepete	k A or	li JLo	B	England	ch.hil.	D co Eng. bot 1414
15183 Marum-leaved	A or	li jn.jl	P	Spain	1788.	D co	Cav.ic. 6.576
M petra marifulia Csv.	Th^mus mantMius IV. en.						
15184 critica P<<.	Cretan	n. or	Jjnjl	P	S. Europe	1596.	D r.m Bar. ic. 1166
15185 Iriticusa Pert.	shrubby.	< or	ls	P	Spam	1752.	C r.m
15186 dbla Lk.	white	£ A or	fil.au	W	Hungary	1818.	D p.l

1684. MELISSA.  {*Melissa*, a bee; gather abundance of honey from it.) Labiate. 5.-7.

13187	<i>officiniihs L.</i>	officinal		jn.o	W	S. Europe	1573.	D	co
	<i>S rom'jna</i>	Roman hairy		m.o	W	S. Europe	...	D	co
15188	<i>cordifolia Pert.</i>	heart-leaved		jn.o	W.P	Italy	1808.	D	co
15189	<i>altissima Sm.</i>	tallest		jl.au	W	Greece	1818.	D	s1
151U0	<i>pyrenkica Jae.</i>	Pyrcnacan		jlau	W	Pyrenees	1800.	D	81
	<i>Pyrcnacan L.</i>					Jac. vin.	2.183		

IS191 polyanthos *Lo. C.* many-flowered ■ A or T jiau w — 1820. D s.t

* 1685. DRACOCE'PHALUM L.	DRAGON'S HEAD.	(Drakon, dragon, <i>kephale</i> , head; fl.)	<i>Labioidæ</i> . 23. — 90
15192 <i>Virginianum</i> L.	Virginian	3t A or 3	jls.
15193 <i>speciosum</i> J.Swt.	showy	TV	Or 2 il au
17194 <i>denticulatum</i> L.	toothed	^	TM*
10115 <i>variegatum</i> <i>Veil.</i>	variegated	■	■
15196 <i>cordatum</i> <i>Nut.</i>	cordate-leaved	■	■
15197 <i>Ruyachidium</i> L.	Ruych's	■	■
15198 <i>aubracium</i> L.	Austrian	■	■
15199 <i>feotryoldea</i> Stev.	Botrys-like	■	■
15200 <i>palmatum</i> L.	palmated	■	■
15g1 canari'nsL	Canary <i>Balm qfGUead</i> *	or	
15202 <i>peregrinum</i> L.	foreign	&	
15203 <i>argunense</i> Fit.	Argun	5	or
15204 <i>grandiflrum</i> L.	great-flowered	\$	A or
15205 <i>altaianum</i> Lam.	Altaiian	5	A or 1
la20h' moldavicum L.	Moldavian	O	O or 2
2 albisflrum	white-flowered	O	O or 2
15207 <i>parviflorum</i> Nut.	small-flowered	k	& cr
I'iQiticaneescensZ..	canescent	O	or 2
<i>Him ibicum Step.</i>	Iberian	k	A or 1
15210 <i>petaltura</i> L.	peltate	O	or li
"211/hymiflorumi.	Thyme-flowered	O	ijns
15n?".	nodding	■	A or 1
g3chamiedryoWeJRI/*.	Germander-Ik.	AJ or	i
M ribricum L.	Siberian	A or	jDau
R N. Amer.	1683. D	p.l Bot mag.	467
Pk ^{ck} . Siberia	1822. D	co Sw. fl. gar.	93
St Carolina	1787. D	p.l Bot mag.	214
P Carolina	1812. D	co Ven.cels	44
P N. Amer.	1824. D	p.l	
B N.Europe	1(99. D	p.l Fl. dan.	121
B Austria	1097. D	p.l Jac. ic.	1.112
P Siberia	D	co	
P Siberia	1815. D	co	
Pa.P Canaries	1G97. C	rm Com. h.	2.41
B Siberia	1759. D	p.l Bot mag.	1084
B Siberia	1822. D	co Bot. cab.	797
B Siberia	1759. D	p.l Bot. mag.	1009
P Georgia	1787. D	co N.c. p.26.	3
B Moldavia	1596. S	co Lam. il	513.1
Moldavia	1596. S	co	
W N. Amer.	IBI'5. D	p.l	
B -Levant	1711. D	co Sw. fl. gar.	58
B Iberia	1824 D	p.l	
B Levant	1711. S	co Lam.il.	513.2
P Siberia	1752. S	co Gm.Bi.a50	
B Siberia	1731. D	co Botreg.	841
B —	1823. D	Up	
B Siberia	1760. D	p.l Bot mag.	5185

1686. MELITTIS L. BASTARD BALM. (*Meli*, honey; bees gather abundantly from.) *Labiatae*. 2.-4.
 15215 Melissophyllum L. Melissa-leaved ^ A or * my.jn F England woods. D co Eng. bot. 577
 2 alpinum ^ A or I my.jn F SwitzerL ... D co
 15216 grandiflora Sm. great-flowered ^ A r 1 my W!t England woods. D co Eng. bot 636

• 1687. (yCYMUM L. BASIL. (*Ozo*, a smell; powerful scent.) *Labiatae*. 20. — 31.
 15217 thrysiflorum L. thyrs-flowered jfc S un lijl.au W E. Indies 1806. S s.l Jac. vin. & 72
 15218 gratissimum L. most grateful n. S un lijl.au W Indies 1751. C s.l Jac. ic. 3.495
 15219 basileum L. common un 2 jl.au W Indies 1743. C s.l Black. 104
 15220 pilosum W. en. pilose O cul 1 jl.au W India . . . 1848. S r.m
 15221 micranthus W. en. small-flowered Q un 1 jl.s W 1816. S 8.1
 15222 viride IV. en. green tt. HHun 1 jl.au Pa.P. 1816. S 8.1
 15223 suave W. en. ariet-scented a. Ljun 3 jl.s W.a 1816. S 8.1
 J5224 canum 5ww hoary D un 3 jl.s W China 1822. CO Bot. mag. 2452
 15225 sanctum L. holy O un 1 jl.s W E. Indies 1758. R. mal. 70.92
 15226 febrifugum Lindt. febrifuge S un 1 jl.s P S. Leone 1822. CO Bot. reg. 753
 15227 album L. white lol or 3 jn.o W E. Indies 1816. S 8.1
 15228 amencanum L. American rm un 1 jl.s W W. Indies 1789. s.l Jac. vin. 3.86
 15229 caryophyllatum Rox. Clove-scented 2.3 or 2 jl.au T E. Indies 1817. S 8.1
 15230 villsum llox. villous CSJor 1 jl.s W E. Indies 1783. S 8.1
 15231 wienthoite L. Mint-like O un 1 jl.s W E. Indies 1823. S 8.1
 15232 inodruin /.am. scentless ttt or 1 jl.a W E. Indies 1822. S 8.1
 15233 cristatum Rox. crested O or 1 s.o Abyssinia 1802. S 8.1 Her. ser. 43
 15234 grandiflrum Herit. great-flowered 1 I un 1 jl.s W E. Indies 1573. r.m Schk.han.2.166
 15235 minimum L. least O cul i-Jun Mauntius? 1825. C 8.1
 15236 cordiflum J?q. heart-leaved i-Jun

* 1688. IAJMSVTZEHA Jac. LUMNITZERA. (*Stephan Lumnitzer*, a botanical author.) *Labiatae*. 5. - 9
 % 15237 ocyoimoides Jac. Ocyium-hke O or 1^ jl.s W 1823. S co

\ 15238 polyhachya Jac. many-spiked E Jun 1 jl.au W E Indies 1783. S 8.1 Mur.3.3
 O'cymum polycladum Lk.
 15239 polystachyon L. slender-flwd £ Q Jun 1 jlau Pa.P. E. Indies 1703. S 8.1 Ru am. 5.912
 415240 capitata W. headed O ec 1J jl.au W China 180G. S.1
 O'cymum capitellatum L.
 15241 mosuhfita Spr. musk iQJ or U jls W N. Holl. 1S23. S co
 Plectranthus moschata R. Br.

• 1689. PLECTR A'NTHUS Herit. PLECTRANTHUB. (*Plektron*, cock's spur, *anthos*, flower.) *Labiutue*. 23. — 38.

\ 15242 fruticosus Herit. shrubby tt. L or 3 jn.s B C.G.H. 1774. C r.m Her. eer. 85.41
 15243 scutellano'ides R.Br. Skullcap-like O or S jl.au B E. Indies 1764. S l.p But mag. 1446
 O'cymum scutellarioides H. A.
 15244 galeatus Vahl helmcodod O un 2 jl.au B Nepal 1820. S s.1
 15245 grav&olens R. Br. heavy-smelling tt. i. I un 2 au.s B N.Soll. 1826. C p.l
 15246 australis R. Br. southern O un 2 au.s B N. Holl. 1826. C p.l Bot reg. 1098
 15247 incknus Lk. hoary £ iAl or 3 jlau B 1822. D co
 15248 monachorum Spr. monks' O un 1 jlau W E. Indies 1796. S s.1
 O'cymum monachorum L.
 15249 mofhs Spr. soft iGJun 1 8 V E. Indies 1781. S 8.1
 O'cymum m6Ue H. K.
 15250 asper Spr. rough S un S au.s B E. Indies 1824. C 7
 | 15251 Forskoi(Urt T barbatus B. R.
 strobiliferous O " 1 jl.au B E. Indies 1804. S 8.1
 15252 secundus Rox. side-flowering O un 1 jl.au B R. Indies 1816. S 8.1
 15253 rotundifolius Rox. dotted jf O or 2 jamy B Africa 1775. r.m Her. ser. 87.41
 15254 punctatus Herit. round-leaved O un 1 jl.au B Mauritius 1823. S 8.1
 15255 rhymiflorus Spr. Thyme-flwd O un 1 au.s E. Indies 1826. S 8.1
 15256 viscius Spr. clammy tt. I I un 1 jaus E. Indies 1826. S 8.1
 15257 viscius Spr. small, flowered n. or 3 jn.s S Amer. 1805. W. h. b. 65
 15258 parviflrus W. en. heart-leaved tt. L or 3 jn.s Nepal 1823. S 8.1
 15259 cordifolius D. Don comose n. I pr 2 au Nepal 1821. CO Bot. mag. 2318
 15260 comitus Sims colored tt. L un 2 au.s Nepal 1820. S 8.1
 15261 colorktus D. Don Coetsa tf TA un 2 o Nepal 1823. S 8.1
 15262 Coetsa Ham. tematus Sims ternate Omime Plantk £ or 1 au Madagasc. 1821. D r.m Bot. mag. 2460
 15264 temifolius D. Don three-leaved tf i un 2 aus Nepal 1820. C p.l

1690. POGOSTE^MON Dcsf. POGOBTEMON. (*Pogon*, a beard, *stemon*, a stamen.) *Labiatae*. 1.

15265 plectranthodes Desf. Plectranth-lk t t j or 1 ... W. C ep

• 1691. TRICHOSTE^MA L. TRICHOSTBMA. (*Thrix*, hair, *sterna*, stamen; slender.) *Labiatae*. 2.—*
 15266 dichotomum L. dichotomous O or 1 jn jl B N. Amer. 1759. S s.l lam. il. 515
 15267 lineare MM. hncar-featwtf O pr 1 jn.au B N. Amer. 1759. S s.p Di. el. 285.369

1692. PROSTANTHE^RA Lab. PROSTANTHERA. (*Prostheke*, appendage, *anthera*, anther.) *Labiatae*. 1. — &

15268 denticulata R. Rr. toothed sh i-J or 4 jn.au ... N. Holl. 1824. C s.p
 15269 rhumbra R. Br. rhomb-far iwi S un jn.s N. Holl. 1823. S s.p
 15270 rotundifolia if. Br. round-leaved SH i-J or 3 jn.au N. Holl. 1824. S s.p
 15271 incisa R. Br. cut-leaved *L J or 3 jnau N. Holl. 1823. S s.p
 19T& vioKicca /{ Br. violet-colored *L J or 5 my.au V N. Holl. 1820. C s.p Bot. reg. 107*
 15273 linearis R. Br. linear-leaved *I J or 3 jn.au N. Holt. 1824. C s.p
 15274 lasianthos Lab. villous-flowered H i-J or 2 jnjl P.w N. & W. 1808. C s.p Bot. reg. I*³

* 1693. SCUTELLA^RIA L. SKULLCAP. (*ScuMla*, a small dish; figure of calyx.) *Labiatae*. 31.—⁴ ■

I. SOLITARIIFLORX. — Flowers axillary, solitary.
 L gal S un A I jn.s B Britain wat.pl. D co Eng. bot 523
 spear-leaved S un A I jn.s P Germany 1798. D co
 Scordium.lvd S un A I jn.s B Siberia 1817.
 less % jl.au Pk Britain xn.hed. D S¹ Eng. bot. 52*
 spring i jnjl B S. Europe 1821. D pi
 ambiguous I jnjl B N. Amer. 1824. D pi
 Havannah S un A I jn.s B Havannah 1793. D El Jac. ob. 2.29
 dwarf i jnjl B N. Holl. 1823. D P-i A 106
 very small S un B N. Amer. 1822. S p.l Hook.ex.fl^{iu} ^

15284 grandiflora Sims
15285 lateriflora L.

II. RACEMIFLORA—*Flowers in racemes, spicate, axillary.*

large-flowered	^ A or	lj.l.au	P.Y	Siberia	1804.	D	8.1	Bot mag. 635
side-flowering	£ A or	1 jn.a	B	N. Araer.	1752.	D	p.1	

15286 orientalis L.
15287 macrantha Fit.
15288 zlt&UnLebed.
15289 decumbens Sieb.
hfrta Sm.

III. TERMINALIFLORAL—*Racemes terminal, simple.*

oriental	k A or	1 jl.s	Y	Levant	1729.	D	pi	Bot. mag. 2120
long-flowered	1 A or	ijts	B	Siberia	1827.	D	co	
Altaian	^ A or	ft jn.jl	...	Siberia	1817.	D	p.1	
decumbent	5-Aior	^ jn.jl	B	S. Europe	1825.	D	p.1	

15290 alpina L.
15291 peregrina L.
15292 altimaria L.
15293 pilosida Zle&
15294 Columna AIL
15295 rubictincta Horn.
15296 nigrescens Spr.
15297 albida L.
15298 lupulina L.
15299 integrifolia L.
13300 Alyssopifolia £.
15301 purpurascens Sti.
15302 caroliniana Lam.
15303 nervosa Ph.

alpine	£ A or	f jn.o	B.w	Hungary	1752.	D	p.1	Sw. fl.ear.90
foreign	k A or	2 jn.o	V	Italy	1683.	D	co	W. & K. 2.125
tallest	£ A or	1 jl.au	D.P	Levant	1731.	D	p.1	Bot. mag. 2548
pale	k A or	2 jl.au	W	Crimea	1824.	D	co	Gm. si. 58
Columna's	k A or	1j jn.au	B	Italy	1806.	D	co	Sw. A. gar. 52
ruddy	k A or	2 jl.au	Pk		1823.	D	co	
nigrescent	* A P*	2 jl.au	Y	Candia	1826.	R	co	
whitish	k A or	lf jn.il	W.p		1771.	D	s.1	Sab. rora. 3. 29
wolf	k A or	I in.s	Y.w	Tartary	1739.	D	p.1	Schm. ic. 73
entire-leaved	jg A *	1 jn.s	B	N. Amer.	1731.	D	p.1	Pluk.aL441.6
Hyssop-leaved	k A or	1 jn.s	B	N. Amer.	1731.	D	co	
purpurascens	k A or	2 jn.jl	B	W. Indies	1830.	D	p.1	
Carolina	5 A or	lf jn.jl	B	Carolina	1811.	D	co	Lam. IL 515.3
fibrous	k A or	1 jl.au	B	Virginia	1826.	R	co	

15301 pirosa Afo.
15J05 serrata Andr.

IV. COMPOSITIFLORAL—*Racemes compound, terminal.*

pilose	k A or	1 jl.au	B	N. Amer.	1825.	D	p.1	
law-leaved	k A or	4 jn.s	B	N. Amer.	1800.	D	s.1	Bot rep. 494

* 1694. PR UN&LLA L. SELF-HEAL. (*Die breuna, disease of jaws, Ger.; supposed mod. qual.*) **Lobtite**, 13.

15306 vulgaris L.
,, 2ilba

common	^ A m	ft jl.au	Pk	Britain me. pa.	D	co	Eng. bot 961
white	4jlba	d	W	Britain me. pa.	D	co	

15307 pennsylvanica Jo308ovata W.

Pennsylvanian	Pa.B	N. Amer.	1801.	D	Pl	W.h.b.9
---------------	------	----------	-------	---	----	---------

15309 grandiflora X.
15310 latifolia Z.O. C.

great-flowered	P	America		S	fp	
----------------	---	---------	--	---	----	--

15311 longifolia P*r*. J5312 alba Pall.

broad-leaved	L.B	Austria	159a	D	p.1	Bot mag. 2014
--------------	-----	---------	------	---	-----	---------------

15313 intermedia Z*. J5314 2. er.

long-leaved		France	1800.	D	p.1	
-------------	--	--------	-------	---	-----	--

J5317 Alyssopifolia L.
15318 hirtetern A.

white		Siberia	1800.	D	p.1	
-------	--	---------	-------	---	-----	--

1695. CLEO^NIAL.
15319 lusitanica L.

intermediate		Portugal	1790.	D	6.1	Bot mag. 337
--------------	--	----------	-------	---	-----	--------------

irr^66 - PRA^SIUM L.
15321 m^nw L.

cut		1823.	I) co			
-----	--	-------	-------	--	--	--

15322 leptostachya L.

jagged		Austria	1713.	D	p.1	Lam.il. 516.
--------	--	---------	-------	---	-----	--------------

hastate		Spain	1818.	D	p.1	
---------	--	-------	-------	---	-----	--

Hysop-leaved		France	1731.	I) p.1	M.h.11.5.7
--------------	--	--------	-------	--------	------------

hairy		Europe	1812.	D	p.1	
-------	--	--------	-------	---	-----	--

--	--	--	--	--	--	--

CLEONM. (A Greek name employed by Theophrastus.) **Labtite**, 1.

Portugal O or 1 jn.jl L.B Portugal 1710. S co Mil. ic. 1.70

PRASIUM.	(The Greek name for horehound.)	LabiaUe.	2.
----------	---------------------------------	-----------------	----

greater Spanish*. J cu 2 jn.au w Spain J699. C r.m FLgr.S84

smaller Sicilian*. J cu 1 Jn.au W Sicily 175L C r.m			
---	--	--	--

--	--	--	--

PIIRVMA. (Meaning unknown.) **Jabiata**. 1.

slender-spiked ^ A cu lf au.s W.p N. Amer. 1802. D Lp Pluk.am.380.5

CLASS XIV. ORDER 2.

DIDYNAMIA ANGIOSPE'RMIA.

*1698. GESNE^RJ^L.

GEBNEBIA. (Conrad Gesner, of Zurich, the famous botanist.) **GesnerUte**. 11.—26.

15323 sckbra Sum.

rough tt.Qor 1 jn.au S Jamaica 1820. C p.1

15324 corymbosa Sun.

corymbose O-CZlor 2 jn.au S Jamaica 1822. C p.1

15325 hirsuta H. & B.

hairy ttDor 1 jn.au S Cumana 1826. C p.1

penduAna Lindl.

Elm-leaved jg EJ or 2 jn.au S S. Amer. 1820. C p.1 **Bot reg.** 1032

15327 calycma Swx.

ftzrgr-calyxed a. CD or lf t. — S Jamaica ... C p.1

15328 hilmills L.

humile it • or * S Cuba ... R p.1 PL gen. 1&3.2

15329 tubiflora Cav.

tube-flowered A El or 2 fmr S. Amer. 1815. R p.1 Cav. ic.584

15330 Douglasii Lindl.

Douglas's A El or lf s RT Rio Jan. 1826. R p.1

J5331 aggregMa Ker

aggregate J T 2 au S Brazils 1816. R p.1 Bot. reg. 329

15332 bulbataa Ker

bulbous AE or 2 my.jn S Brazils 1816. R p.1 Bot reg. 343

15333 acaulis L.

stemless AE or 1 ... S Jamaica 1793. R I.p SL jam. 1.102.1]

MJEP; COD O N 9 PHORA

£ «M» CODONOPHORA. (Kodon, a little bell, phoreo% to bear.) **Gesnerite**, 2.

Gesnre

laeolate tt-a or 2 myjl S & Amer. 1752. C p.1 Bot mag. 1023

15335 grandiflora Lindl.

great-flowered lf myjl S Brazil 1818. C p.1 Bot reg. 428

Gesnre praainata Ker

1702. GLOXI'N/i* Herit	GLOXINIA. (<i>? P. Gloxin</i> , of Colmar, a botanist) <i>Gesneriea?</i> 4.
15342 maculata Herit.	<i>gspoteA-stalked</i> £ E or 1. il.o P S. Amer. 1739. C s.p Botmag. 1191 ¹
15343 specioa B. R.	showy £ E2 or 4 jn.n P S.jmer. 1815. C s.p Bot. reg. 2J3
2 Alba	<i>vrlite-flowered</i> £ LSi or £ jn.n W S. Amer. ... D s.p
15344 caulescens B. R.	caulescent £ ES or £ jl.au P Pernamb. 1826. I. pl Bot. reg. 1127
15345 hirsuta B. M.	hairy £ GH'or * jn.au B S. Amer. 1824. D s.p Bot mag. 2690
1703. LINNJE\4 Gro.	LINVJEA. (Gar/ von Linni, most eminent of modern naturalists.) <i>Caprifl<i>l<i>aceaaf</i></i></i> . 1*
15346 borealis Gro.	northern IU A pr i my.au F Scotl. dry st.c. D Lp Eng. bot 433
2 amencana	American t, ^ or £ my.au F N. Amer. 1800. D Lp
1704. MELIA'NTHUS L.	HONEY FLOWER. (<i>Meli</i> , honey, <i>anthos</i> , flower; abounds in.) <i>Rutacee?</i> 3.
15347 minor L.	greater H i J or 10 my.jl Br C. G. H. 1688. Sk s.f Bot. reg. 45
1548 irffior L.	smaller H i J or 2 au Br C. G. H. 1696. Sk s.l Bot mag. 301
15340 coinjsws Vahl	tufted * i_j or 3 ... Y C. G. H. 1820. Sk.s.l Com. r. 4.4
1705 KRAMER^U/if L.	KRAMERTA. (J. G. II. and W. II. Kramer, German botanists.) <i>Polygplete</i> . 1.— 7.
15350 panciflora 8eni	few.flowered • or 4 ... Mexico 1824. C sp Fl. mex. ic. in.
* 1706. BIGSO'SIA L.	TRUMPET FLOWER. (<i>Abm Bignon</i> , libr. to Louis XIV.) <i>Bignoniacee</i> . 53.—82.
I. <i>Bix</i> *jE.—Leaves binate, tendriUcd; peduncles one or few ftwd,froni 15360. racemose, from 15368. paniced.	
15351 unguis L.	claw ft O or 10 ... Y W. Indies 1759. L. s.p PL. am. 94
15352 staminod.Lam.	toe-stamcned ft. I or 10 ... Y <i>Hibiscum</i> 1822. *C Lp PHIL SpeC. 56. 2
15353 capitoPita L.	tenchained mmml £± I or 10 ... Y N. Amer. 1710. G B.p Bot. mag. 864
15354 pub^cens L.	pubescent ft. • or 15 jn.jl Y Campach. 17. 9. C
15355 roquinctuMis L.	equinoctial ft. • cl 15 jn.jl Y Guiana 1768. C PL ic. 55.1
15356 Chainherplayn/i B.R.	Chainherlayne stft. O el 40 ap.o Y Brazil 1820. C tr Bot reg. 741
15357 s.ilic.a illia Kth.	Willow-leaved ft. • or 40 ap.o Y Trinidad ISL'4 C
15358 Bonp. Lam.	yellowish-whiteft. O or 10 ... Y.W. Amer. 1824. C ll
15359 R>cttal>ilis Vahl	ft. O or 10 jn.jl Y W. Indies 1790. C
ir.JCA plecta Kth.	showy ft. CD or 10 ... P W. Indies 1800. C l.p
15360 rigescens Jac.	painted ft. D or 10 ... Va S. Amer. 1823. C s.l
L'f[H] Licti flora Vahl	rigescut ft. • or 10 jn.jl Pk Caracas 1823. C s.p
15361 Inconamtha Lie.	mdk-flowered ft. D or 20 jn.jl W SantaCruz182J. C s.p
15362 airifttha Vahl	smaller-flwd ft. O or 20 jn.jl Pk N. Holl. ? ... B.p
15363 granditlli.i Jac.	L:ircd-leaved ft. ^ or 20 jn.jl Y* - Guiana 1804. C 1P
lo.X-7 latif?nia Rich.	large-loavcild ft. - or 60 ap.jl Y* - Caracas JSIo. C P.1
15368 elongata Vahl	broad-leaved ft. - or 10 - P Cayenne 182 i. C l.p
15369 mollissima Kth.	elongated ft. Dor 10 ... - S. Amer. 18J0. C l.p
15370 caiidif ans litch.	whitish ft. Dor 10 ... - Cayenne 1820. C l.p
15371 diversiflaria Kth.	various-leaved ft. • or 10 ... Mexico 1825. C l.p
15372 floribunda Kth.	buudle-flwd II lor 12 ... Mexico C l.p
II. TERNAV/E.—Leaves ternate, tendrilled, except Xo. 15382.	
15373 cruegera L.	cross-bearing Y.d S. Amer. 1759. L. s.p PL. ic. 58
15374 chirtre Aub.	Chiriri R.O Guiana 1824. C Lp Aub. gui. 2.260
heterophila 71:	
15375 yon'ista B. R.	lovely LOor 8.d O S. Amer. 1816. C Bot. reg. 249
15376 inc. irn.-ta Aub.	flesh-colored P.R Guiana 1820. C Aub. gui. 2.251
15377 m.Mlis Vahl	soft *-Hof 10 Guiana 1818. C Vahl ec. 2, 10
tomontii-a Rich, not Tinn.	
15378 pilliida Lindl.	vn\Q-flowered ft. Q 15 JI W W. Indies 1823. C r.m Bot reg. 965
iffTJ) litttirMis Kth.	shore ft. 10 ... P.R Mexico 184. C Lp
15380 iridiflora Lod.	green-flowered ft. • or 10 my G S. Amer. 18SS. C Lp
15381 apurensis Kth.	Apure ft. O or 10 ... Y Orinoco 1824. C Lp Bot reg. 1626
15382 triph?lla L.	three-leaved S O or 10 ... W S. Amer. 1733. R Lp
15383 cethin?na L.	bristly JrM/crf fl. I or 20 ... Pk Guiana 1804. C Lp Aub. gui. 2.264
Jacardinda echinata Spr.	
III. DICITAVA.—Leaves digitate.	
15334 fluviatilis Aub.	river ft. Q or 10 ... Guiana 1824. C Lp Aub. gui. 2.267
15385 Leuci'xylon L.	white-wooded ft. Q or 10 ... W. Indies 1759. C Lp Bot rep. 43. 4
Vf&6 rhysanthia Jac.	yclnw.flowered ft. Q or 10 ... Guiana 182a. C Lp Aub. gui. 2.211
15387 serratiflaria Vahl	saw-leaved ft. Q or 10 ... W. Indies 1822. C Lp
15388 osculiflaha Kth.	Horschestnut-lvd ft. Q or 10 ... Mexico C Lp
IV. PINNA'T/E.—Leaves simply pinnate.	
15389 bijuga Vahl	twin-leaved H Q or 6 ... Madagasc. 1822. C l.p
15390 Clematis Kth.	Clematis ./^ ft. O or 15. . . . Caraccas 1820. C l.p
V. COMPOSITE.—Leaves decomposed or bipinnate.	
15391 varilibilia Jac.	variable ft. 10 or 10 ... Y.W. Caracas 1819. C l.p Jac. scS. SIS
15503 indjea L.	Indian 18 40 ... W India 1775. C l.p R. mal. 1.43
15393 subcrbsa Box.	Milliungfinia hortensis /.
15394 Chica Hum.	Chica ft. • or 10 ... Orinoco 1819. C Lp
IS395^asminalia Kth.	Jasmine-leavedft. O or 10 ... Orinoco 1826. C Lp
15396 alba Aub.	white ft. CD or 8 ... W Guiana 1823. C l-p Aub. gui. S66
Jacardinda filba Spr.	
VI. SIMPLICIFOLIA.—Leaves simple, cordate, five-nerved.	
15397 tomentosa Thun.	woolly 2 O or 10. . . . Japan 1820. C pl Ksm. 860
Incarvillea tomentosa Spr.	
VII. DU^BIA.—Doubtful to which of the sections they belong.	
15398 dectpiens Lo. C.	deceiving ft. • or 10. . . . E. Indies ?1823. C
15399 crenata Lo. C.	crenated ft. CDor 10 E Indies? 1823. C
15400 lucida Lo. C.	shining ft. O or 10 E. Indies? 1823. C
15401 multifida Lo. C.	many-cleft J. I or 10 E. Indies? 1823. C p.
15402 quadrangulaTis Box.	quadrangular ft. O or 10 E. Indies? 1823. C p.
15403 purpurea Lo. C.	purple • Dor 6 & Amer. 1822. C Lp
1707. AMPHILO^BIUM Kth.	AMPHILOBIUM. (Amphi, around, lobos, a pod.) <i>Bienoniacet</i> . L- 15404 panicuKitum Kth.
panicled ft. • or 20 ... P W. Indies 1738, C Lp PL ic. 56.1	
■ paniculate W.	
*J708. SPATHO^DEA Beam.	SPATHODEA. (Spatha, a sheath; fl. sheathed at base.) <i>Bignoniacea</i> . 1-~
15405 unkta Spr.	hooked ft. Q or 10 Y Guiana 1804. C s.p Bot mag. J
Bigh&nia unkta B. M.	

15406 corymbosa <i>Ven.</i>	corymbose	iJ CD or 6	-	Y	Trinidad	182*	C l.p
15407 Rheedii <i>Spr.</i>	Rhecede's	£ or 16	-	Crea	E. Indies	1794.	C l.p R.mal.1.43
Bigndntospathacea	<i>Rctz.</i>						
15408 RoxbArgw <i>Spr.</i>	Roxburgh's	tCJor 12	-	Pk	E. Indies	1820.	C l.p
Bigndnco quadnlocularis	<i>Rox.</i>						
15409 kcVis Beam.	smooth	f Dor 12	-	P	Guiana	1825.	E l.p Beau. ow. 1. 29
15410 longiflora <i>Ven.</i>	long-flowered	\$ a or 12	-	R	E. Indies	1816.	L Lp R. mal 6. 29
Bignbni chelondldes <i>L.</i>							
15411/raxinifMia Kth.	Ash-leaved	LCD or 10	-		Caracas	1822.	C Lp
Bignonia/raxmifblia <i>Spr.</i>							

1709 FIELD/d. FIELDIA. (*Baron Field*, some time chief judge in N. S. Wales.) *Bignoniadgm.* 1.
15412 australis *Cun.* southern ft., i-j or 1 jl.au W N. Holl. 1826. C a.l Ex. fl. 232

• 1710. TECOMA J.	TECOMA.	(<i>Tecomaxochitl</i> the Mexican name.)	<i>Bignoniace.</i> 13.
15413 splcidida <i>Spr.</i>	splendid	Q or	Y Brazil 1820. C p.l
15414 pentaphylla <i>J.</i>	five-leaved	Q or	Y W. Indies 1733. C p.l Marcg.br. 118
Bignbni J. centaphylla <i>L.</i>		Q or	
15415 australis <i>R. Br.</i>	southern	A Ljor	O N.S.W. 1793. C s.p Botmag.865
Bigndnco Pandora Andr.			
15416 digitata <i>Kth.</i>	digitate	H Q or	Y S. Amer. 1818. C l.p
15417 mols <i>Hum.</i>	soil	S _j or	Y Mexico 1824. C p.l
15418 radicans <i>J.</i>	rooting	- or 30	O N. Amer. 1640. R s.p
Bignonia radicans <i>L.</i>			
1 major	greater Ash-lyd\	or 30	O N. Amer. 1640. R s.p Bot mag. 485
* 2 minor	smaller Ash-4vd	I or 20	S N. Amer. 1640. C Bp Cat. car. 1.65
15419 grandiflora <i>Suit.</i>	large-flowered	J J or 30	O China 1800. C r.m Bot. mag. 1598
Bignonia erandiflora <i>Thun.</i>	chinensis <i>Lam.</i>	Incarvillea grandiflora <i>Spr.</i>	
15420 capinsis <i>B. R.</i>	Cape	L or 8	Jls O C. G. H. 1823. C p.l Bot. reg. 1117
Bignui Mi cap^nsis <i>H. K.</i>			
15421 stans <i>J.</i>	standing	t D " 12 au	Y America 1730. S l.p Pl.ic.54
Bignofno stans <i>L.</i>			
15422 incisa <i>Svit.</i>	cut/cavcd	H • or 10 au	C S. Amer. 1850. C l.p
Bignofia inc^sa Co. C			
15423 fimbucifolia <i>Kth.</i>	Sambucus-lvd	A CD or	Y Peru 1824. C p.1
15424 rosafrlia <i>Kth.</i>	Rose-icavcd	• O or	Y Peru ... C p.1
15425*orbifolia <i>Kth.</i>	Sorbus-leaved	*D or	Y & Amer. ... C p.1

1711. CHILOPSIS D. Don CHILOFSIS. (*CheUos*, a lip, *opsis*, resemblance; flower.) *Rignonihcea.* I.
15426 salignaD. Don Willow-feawrf • L.J or 10 ... Pk Mexico 1825. C p.l Cav. ic. 3.269,
Bigninia l'mearis Cav.

V/Vi. JACARA/2IDA3.	JACARANDA.	^ QU name in Braijl	Braijl Guiana 1793. <i>Bignoniilce</i> *. 7.
15427 copia D. Don	Copaya	t D or 20	Braijl Guiana 1793. <i>Bignoniilce</i> *. 7.
procdra Spr.	Bignoniif procera W.	coputa Aub.	
JS42S tomentif sk R. Br.	woolly	J O ' ? : ! " "	S Brazil 1824. C l.p Bot reg. 1103
15429 bahamensis B. M.	Bahama	J _J or 10	jl.au B Bahamas 1724. C pd Cat car. 1.42
Bignbnir coerilea L.			
15430 mimosifolia I. Don	Mimosa-leaved	i C3 or 10	ap.my B Brazil 1818. C p.1 Bot. reg. 631
ovalifMia R. Br.		20	- Y Brazil 1820. - Lp
k*tl braillana Pers.	Brazilian	t LJ or 15	1825. F Lp
15432 pub6scens ho. C.	pubescent	I O or 15	- B W. Indies 1800. F 1P
15433/iicif7ha D. Don	Teni-leaved	i Q o r	
rhombifolia Mex.			

1712. DIDYMOCA'RPUIS Jac. DIDDMOCARPUS. (*Didymos*, double, *karoos*, ft.) *Didymocdrpe*s. 1. — 8.
i 15434 Rex/i Bow. Rex's DIDDMOCARPUS. \ jaau B C. G. E 182a D p.1 Hook. ex. fl. 227

• 1714. SALPIGLOSSIS R. g/P.	SALPIGLOSSIS.	(<i>Salpigx</i> , a tube, <i>elossa</i> , a tongue.)	<i>Solhnete.</i> 3.
15435 straminea Hook.	straw-colored	O or 1 jn.au	D.P Chile 1824. S p.1 Hook. ex. fl. 229
15436 sinuata R. H.P.	scalloped	ff	D.P Chile 1827. ■
15437 parviflora G. Don	small-flowered	Ejor	Y Chile 18L7. ■ P

*1715. ECCREMOCA'RPUIS A ft P. ECCREMOCARPUS. (*Eklcretnes*, pendent, *karpos*, fruit.) *Bienoniaeee.* 2. - 3.
15438 scarcer R. ft P.

15439 longifl & Hum. rough L u J p r 6 j u u O Chile 1824. C s.l.p Bot reg. 939

15439 longifl & Hum. long-flowered fl _ pr 6 jl.au O Peru 1825. C 8.1.p

- * 1718. CHELO*NE L. CHELONE. (*Chelone*, a tortoise; back of helmet of fl. compared to.) *Scrophular.* 4.-6.
 15468 glabra L. smooth ^ A or 4 au.o W N. Amer. 1730. I p.1 Tr. ehr. 83
 15459 obliqua L. oblique £ A or 4 au.o P N. Amer. 1752. D p.1 Bot. reg. 175
 15460 Lvööii Ph. Lyon's ^ A or 4 jl.s P N. Amer. 1812. D p.1 Bot. mag. 1864
 major B. M.
 15461 gentiano)des Hum. Gentian-like £ AJ or 3 jl.au Oa Mexico 1825. D p.1
 15462 barbata Cav. bearded ^ A or 3 jns O.s Mexico 1794. D p.1 Bot reg. 116
1719. TOURRETTIA. (*M. A. de la Tourrcette*, a friend of Rousseau's.) *Sehameete.* 1.
 15463 Jappacea W. Burdock] O c u 6 jn.au RG Peru 1788. S s.l Sal. st. 5.3
 Dombëya feppacea Hsrit.
1720. MARTY'N/4 L. MARTYNIA. (*John Martyn*, F.R.S., prof, bot, Cambridge.) *Sesdmeat.* 4.-5.
 15464 diandra Glox. two-anthered CD or lljl.au R NewSpain 1731. S s.l Bot rep. 575
 15465 prouoscldea Glox. proboscis-like O or | jn.au I*B America 1738. S r.m Bot mag. 1056
 annua L.
 15466 longiflora L. long-flowered iQjor 2 jl.au Pa.P C. 6. H. 1781. ■ s.l Mee. ic.1.7
 yellow fm or lf ■ Y -Brazil 1825. ■ co Bot reg. 934
 15467 lbtca Lindl.
1731. CRANIOLA'RIA L. CRANIOBLARIA. (*Kranion*, a skull]: sup. resemb. in capsules.) *Sesamea?* 1.
 15468 annua L. annual Ojor lljl.au ■ S.Amer. 1733. S s.l Jac. am. 110
 M.trjruia Craniolaria Swz.
1722. jtCA'NTHUSL. BEAR'S BREECH. (*Akantha*, a spine; some species spiny.) *Acanthiicear.* 7.-9.
 15469 mollis L. soft ■ A or S jl.8 P.w Italy 1548. D co Lam. il. 550
 15470 nigr Mil. black ■ A or 3 jl.s P.w Portugal 1759. D co
 15471 hispiticus Hort. Spanish ■ ■ 2 jl.8 W Spain 1700. D co
 15472 spinosus L. prck'y.leaved ■ ■ 3 jl.8 P.w Italy 1629. D co Bot mag. 1808
 15473 spinosissimus Pers. spiniest ■ ■ 3 jl.s P.w S. Europe 1629. D co
 15474 ilicifolius L. Holly-leaved ■ ■ - E. Indies 1759. C co R.mal.2.48
 15475 carduifolius Thun. Thistle-leaved ■ ■ jl.s B C. G. H. 1816. S p.1
1723. BLE'PHARIS J. BLEPIHARIS. (*Blepharis*, the eyelashes; bracteas of calyx.) *Acanthacete.* 5. — 10.
 15476 capensis Pers. Cipe. £ O or 1 jl.au B C. G. H. 1818. S l.p
Acanthus capensis Thun.
 15477 furcata Pers. (orkspined n.\ or 2 jn.au - C. G. H. 1816. S
 15478 procumbens Pers. procumbent t\ or 1 jn.au - C. G. H. 1825. S l.p
 15479 hoerhaavid/dja Pers. Hoerhaavia.lvd El or 2 jl.au E. Indies 1821. S l.p Bur. in. 42.2
 15480 linearifolia Pers. linear-leaved O or 2 jn.au ■ Guinea 1823. S l.p
1724. LEPIDA'GATHIS W. LEPIDAOATHIS. (*Lepis*, a scale, ogathis, a ball.) *AcanthiUe**. 1. — 2.
 15481 cristata W. crested £ ES or 2 . . . E. Indies 182a C s.l
- * 1725. B ARЛИTR/J L. BARLERIA. (Alt from *J. Barrelier*, a Domini. of Paris, a eel. bot) *Acanthicece.* 12. — 21.
 15482 flixifolia L. Box-leaved il C3 cu 1 . W E. Indies 1768. C Lp R.mal.2.47
 15483 jolanifolia L. Solanum-leavedft. * or ■ B W. Indies ... C Pl. gen. 43.2
 15484 PrionUie L. Prionitis il O or 3 jl.au O E. Indies 1759. C R. mal.9.41
 15485 longifolia L. long-leaved jf OS or 2 jl.s W E. Indies 1781. S l.p Pluk. al. 13a 4
 15486 crista L. crested * O or 2 jns B E. Indies 1796. C p.1 Bot. mag. 1615
 15487 longiflora L. long-flowered il • or 3 jn.a ■ E. Indies 1816. C p.1 Vahl sy. 1.16
 15488 flava Jac. yellow ■ ahle Eranthemum flavum W. E. Indies 1816. C p.1 Bot. reg. 191
 mUs B. R. Justicia flava ■ ahle Eranthemum flavum W. E. Indies 1816. C p.1 Bot. cab. 360
 15489 Silba Hort. white «D o r 3 jl.jl W N. Holl. 1815. C co Bot. cab. 344
 15490 purpurea Lod. purple * CI]pr 2 s P E. Indies 1814. C l.p Bot cab. 344
 15491 cssrbka Box. blue * or 2 jn.a B E. Indies 1823. C l.p
 15492 dichotoma Box. forked • or 2 jl.au P £ Indies 1823. C l.p
 15493 Briggsa W. strigose * Q or 2 jl.au B E. Indies 1820. C l.p
1726. PHAYLO'PSIS J. PIAYLOPSIS. (*Phaylos*, contemptible, opsis, appearance.) *Acanthdccc.* 3.—&
 15494 longiflora Sims long-leaved <- Q p r 2 ap.o W S.Leone 1822. S co Bot. reg. 2433
 15495 imbricata Swz. imbricated £ El or 1 mr.jl W Mauritius 1822. S m.s
 ^theiluma imbricata Spr.
 15496 glutintaa G. Don clammy j£ O or 1 mr jl W Guinea 1824. S m.s
- 1727. RUE'LL/iLL. RUELLIA. (*John RueUe* of Soissons, bot., phys. to Francis L.) *Acanthhcece.* 27. — 84*
 15497 ocymo'ides Cam Basil-like ■ ■ 1| jl.au B Mexico 1815. C Lp Cav. ic 5456*!
 15498 ciliata Horn. ciliated ■ ipr 2 jl. P E. Indies 1806. C l.p
 15499 austrodis CaiK southern ■ ■ 2 jl.au B N. Holl. 1824. D l.p Cav. ic. 6.586.1
 15500 te'tida Hum. fetid ■ ■ 2 n.au B S. Amer. ... C l.p
 15501 biflbra /,. two-flowered ■ ■ 1 ' Pa.B Carolina 1765. C l.p
 15502 vioKicca Aub. violet-colored ■ or Jjl.jl V Guiana 1820. D l.p Aub.gui.2.271
 15503 ralicifrla Vahl Willow-leaved ■ or 1 jl.au B E. Indies 1820. S l.p
 15504 cihoa Ph. ciliated ■ or 1 Lau B Georgia 1824. D l.p
 15505 rubricafilis Cav. red-stemmed ■ ■ 1 jl.au B Mexico 1823. C l.p Cav. ic. 5. 417
 15506 ovhta Cav. ovate-leaved ■ ■ 2 jl.au D.B Mexico 1800. I) In Cav. ic. JL 5254
 15507 pfitala Jac. spreading ■ n.CJpr 1 Jjl.jl D. B Mexico 1800. I) In Jac. ic. 1. 119
 15508 anuophilla Wai. unequal-leaved £ El or 2 jn.au Pa.V £. Indies 1774. C l.p Bot. cab. 1070
 persicifolia B. R. amygdalaeflha Hort. ■ ■ B KIndies? 1823. D l.p
 15509 formusa Andr. handsome tt-Dpr 2 jns S Brazil 1808. C s.p Bot mag. 1400
 elegans Poir.
- 15510 macropf lla Vahl long-leaved £ El pr 3 ... R & Martha 1824. C Lp Vahl sy. 2.39 .
 15511 tuberosa /,. tuberous-roofa ■ ■ 2 jl.au B Jamaica 1752. C Lp SI. jam. 1.95.
 15512 streps L. rattling tf Alpr 2 jl.au Pa.B N. Amer. 1726. D Lp Schk. han. 2.177
 15513 lactea Cav. milky j£ IAJ P' 2 jn.au Pa.V Mexico 1796. D Lp
 15514 clandestina L. hidden £ C2 pl 2 jl.au B Barbadoes172& C Lp Cav.ic. 3.255
 15515 paniculata L. panicled E)P* 3 au P W. Indies 1768. C Lp Di. el. 248.320
 15516 ftilgidd Andr. bri ■ ■ jLau S W. Indies 1804. C s.p SI. jam. 1.100.2
 waved £ (23 pr ■ ■ R Indies 1824. C Lp Bot rep. 527
- 15518 hlrta Vahl hairy ig EJ or 1 jl.au B'' E. Indies 1817. C Lp
 pubescent ig G3 pl 2 jn.au D.B C. G. H. 1823. C l.p
 15520 tetragona Lk four-cornered ig El pr 2 jl.jl B Brazil 1824. C Lp
 15521 longitilia Both long-leaved ig El or 2 jl.au B E. Indies 180'. C Lp
 15522 cerifua Aer. drooping ig El or 1 jlau B E. Indies 1816. D Lp
 15523 depndens Box. dependent £ El or lkjl.au B E. Indies 1816. D Lp

1728. HYGROTH1LA R. Br. HVGBOPHILA. (to love; habitat.) *Acanthdce**. 1.—3.

15524 *rfgens* R. Br. gaping t* El or Indies 1820. D bog
Ruellia ringens L.

1729. SLETHUM J. BLECHUM. (*Blechon*, Gr. name of a plant resembling marjoram.) *Acanthhcece*. 4.—R
15525 *Brtweui* J. Browne's * El or 2 jn W. Indies 1780. C jp SL jam. 1.100.1
15526 *laxiflorum* J. loose-flowered j. El or 1 jn.jl W Jamaica 1818. C lp
15527 *angustifolium* R.Br. narrow-leaved f El or 1 jn.jl B Jamaica 1824. C lp
15528 *brasiliense* Lo. C. Brazilian . f El or 1 jn.jl B Brazil 1824. C lp

1730. AMASOWJ L. AMAZONIA. (*Thomas Atnason*, an American traveller.) *Verbenhce**. 8.—4.
15529 *erecta* L. erect £ E pr 1* au.s Y Maranh. 1823. Sks.l Aub. gu. 2.252
Taligdla campastris Aub.

15530 *punctea* Vahl scarlet £ El pr 1) au.s Y Trinidad 1825. Sk s.l

1731. APHELA'NDRA R.Br. AVUELWDR A. (*Apheles*, simple, oner, a male; 1-celled anther.) *Acanthucce*. 1.
15531 *cristata* R. Br. crested il • spl 3 jn.s S W. Indies 1733. C Lp Bot. mag. lu78
Ruellia cristata B. R. Justfcfa pulcherrima L.

1732. GEISSOME^RIA B.R. GEISSOMERIA. (*Geisson*, penthouse, meris, part; calyx imbric.) *Acanth*.
15532 *longiflora* B. R. long-flowered m • el 3 jl.au S Brazil 1826. C Lp Bot reg. 104J

1733. CROSSA'NDRA Sal. CROSSANDRA. (*Krossos*, a fringe, aner, a male; anthers.) *AcantMceee*. 1.
15533 *undutefolia* Sal. wave-leaved il • spl lja.jn O.B E. Indies 1800. C p.l Bot reg. 69
Ruellia infundibuliformis B. R. Harrachto spicusa Jac.

1734. THUNBERGJW L. THIONBERCIA. (*Ch.P.Thunberg,T.H.S., F.M.L.S.*, celeb, bot & trav.) *Acanthhceat*. 8.
15534 *grandiflora* Rox. large-flowered t Q or 6 my.s B E. Indies 1820. C pl Bot mag SKR
15535 *Fragran Box*. fragrant — 4 my.s W E. Indies 1796. C pl Bot. mag. ISSI
15536 *confota Coll.* heart-leaved — 3 my.s Trinidad 1820. C pl Col. h. rip. 21
15537 *canensis Retz.* Cape — 3 my.r W C. G. H. 182*. u pl Lam. il. &c. 1
15538 *alata Lod.* winged — 4 my.s Y E. Indies 1823. C pl Bot cab. 1045
15539 *angulata* Fq/. angular — 4 my.s Y Madagasc. 1823. C pl Bot cab. 1044
15540 *coccinea Wai.* scarlet % • or 4 mys Nepal 1823. C pl Hook. ex. fl. 195
15541 Hawtayneina Wai. Hawtayne's taspiio ... S' Nepal 1826. C pl
S

1735. HEBENSTREIT/^ L. HEBENSTREITIA. (*J.E. HebenstrcU*, M.D., prof, bot, Ldpic.) *Verbendc*. 10.-12.
15542 *scabra Thun.* rough « I J or 1 jn.jl w C. G. H. 1824. C s.n
15543 *IeSaL* toothed E I pr 1 myss W C. G. H. 1739. S pppl Bot. mag. 483
15544 *integrifolia L.* entire-leaved — 1 my.jn W C. G. H. 1792. C pl Bot. rep. 252
av'real? Rep.
15545 *ciliata L.* ciliated <- I J pr 1 my.jl W C. G. H. 1815. C pl
15546 *capitata Thun.* capitate <- I J or 1 jn.jl W C. G. H. 1823. C s.p
15547 *albiflora Lk.* white-flowered <- I pr 1 my.s W C. G. H. 1822. C pl
15548 *fruticosa Thun.* shrubby tt. I or 1 jn.jl W C. G. H. 1816. C pl Bot mag. 1970
15549 *cno'ides Thun.* Erinus-like &* I J pr 1 my.n W C. G. H. 1816. C pl
15550 *chamaHryfilia Lk.* Germander-leaved I J pr 2 my.s W C. G. H. 1822. C pl
15551 *cordata L.* heart-leaved < t I J pr 1 jl.au W C. G. H. 1774. C pl

1736. GMELI'Nif L. GMELIVA. (*George Gmelin*, a celeb. Ger. naturalist and trav.) *Verbenhcece*. 5.-6.
15572 *speciosissima* D. Don showiest br 15 ... W Nepal 1823. C s.l
15573 *arborea Rox.* tree — or 15 ... Y E. Indies 1824. C s.l
15574 *villlosa Rox.* villous — or 10 — E. Indies 1818. C
15575 *asiatica L.* Asiatic — or 10 — Y E. Indies 1792. C pl Lam.il. 452
15576 *parviflora Rox.* small-flowered f Q or 10 — O E. Indies 1817. C Lp Rox. cor. 32

• 1737. PRE'MNA L. PREMNA. (Premium, a stump of a tree: trunk.) *VerbenhetB*. 2. — IS.
15557 *reticulata J.* netted t (HI or 10. Jamaica 1819. C s.l Gae.ff. 1.56
15558 *integrifolia L.* entire-leaved 1 • or 10. E. Indies 1824. C s.l Ru. am. 3.141

* 1738. LANTA^NA L. LANTANA. (An ancient name of Hburnum; similarity.) *Verbenace**. 28.-37.

I. ACULEAVB. — Stems prickly.

15559 <i>aculeata L.</i> prickly iit • or 10 ap.n R W. Indies 1693. C pl Bot. mag. 96
15560 <i>horrida Kth.</i> nornd — or 3 ap.n R Mexico 1824. C pl
15561 <i>purpurea Horn.</i> purple — or 2 jn.s P S. Amer. 1820. C pl
15562 <i>coccinea Lo. C.</i> scarlet — or 3 jn.s ♀ S. Amer. 1824. C pl
15563 <i>scabrida H. Jf.</i> roughish > I or 2 s W. Indies 1774. C pl Bot cab. 1T71
15564 <i>melissifolia H. K.</i> Balm-leaved * or 2 jl.s Y W. Indies 1732. C pl Di. eL57.66
15565 <i>nivea Fen.</i> white-flowered mi — or 3 jl.s W E. Indies ... C pl Ven. mal 8
15566 <i>Lockhartii G. Don</i> Lockhart's ii — or 8 jl.o W Trinidad 1820. C pl

II. INE'RMB& — Stems not prickly.

15567 <i>stricta Swz.</i> strict D o r 3 ... Pa.P Jamaica 1733. C pl SL jam. 195.4
15568 <i>radula Swz.</i> rasp-leaved — — — — W. Indies 1803. C pl
15569 <i>hispida Kth.</i> hispid — — — — Mexico 1824. C pl
15570 <i>(alvitefolia Jac</i> Sago-leaved — — — — C. G. H. 182a. C pl Jac. sc. a 285
15571 <i>pilosa Kth.</i> pilose — — — — Cuba IK2J. C pl
15572 <i>recta H. K.</i> upright — — — — Jamaica 1758. C pl Jac. sc. a 360
15573 <i>parvifolia Sal.</i> involucrate — — — — W. Indies 1690. C pl Pluk. al. 114 5
15574 <i>involuta L.</i> scented — — — — W. Indies 1758. C pl Pl. ic. 71.2
15575 <i>trifolia L.</i> three-leaved — — — — W. Indies 1733. C pl Bot. mag. 1449
15576 <i>Cannara L.</i> Cammara — — — — W. Indies Iffil. C pl Di. el. 56. 65
15577 <i>fuctoa Ker</i> painted — — — — S. Amer. 1823. C pl Bot. reg. 798
15578 <i>annua L.</i> annual — — — — S. Amer. 1733. C pl Bot. mag. 1022
15579 <i>lavandulacea W.</i> Lavender-like — — — — S. Amer. 1820. C pl
15580 <i>crucifera Lk.</i> copper-co/ortv/ — — — — Jamaica ISIS. C pl
15581 <i>violacea Desf.</i> violet-co/ored — — — — S. Amer. 1818. C pl
15582 <i>geminata Spr.</i> twin — — — — Trinidad 1819. C pl
15583 <i>braziliensis Lk.</i> Brazilian — — — — Brazil 1823. C pl
15584 <i>alba Mil.</i> white — — — — S. Amer. ... C pl
15585 <i>mixta W.</i> mixed — — — — W. Indies 1782. C pl Bot. cab. 68
15586 <i>hirta Grah.</i> hairy — — — — Mexico 1825. C pl

1789. ALOY'SM Or. ALOTSIA. (*Maria Louisa*, mother of Ferd. VIT. king of Spain.) *Verbenacea?* 1.—2.
 15587 citriodura Or. Lemon-scented Sc or 3 my.s Pa.P Chile 1784. C 1 Bot mag. 367
Ferbena triphylla Herit.

•1740. LI'PPLfL. LIPPIA. (*Augustine Liggi* a French phyg., traveller in Abyssinia.) *Verbenacea?* 1.—25.
 15588 purp&rea Jac. purple <& * or 3 jn.jl R Mexico 1823. C p.l Jac cc 85

1741. AFELAMPY^RUM L. Cow WHEAT. (*Melas*, black, *pyros*, wheat; seeds.) *Scrophulariaceae*. 5—7*
 15589 crisifatum L. crested ■ w 1J jl.au Y England com fi. S co Eng. bot. 41
 15590 arvencse L. purple corn-field ■ w jl.jn.jl Y England corn fi. S co Eng. bot. 53
 15591 nemorosum L. grove O or 11 jl.au X Europe . S Fl.dun 305
 15592 pratensc L. common meadow O w 1} jl.au X Britain woods. S ■ Eng. hot 113
 15593 sylvaticum L. wood O w 1} jl.au X Britain m.wo. S ■ Eng. bot. 804

* 1742. SELA^GO /,. SKLAGO. (*Set*, *Bight*, *jack*, salutary, *Celt.*; supposed meil. qualities.) *Verbenacea*. 15.—40
 15594 canescens Thun. canescent a. LJ pr 1f jl.n Pa.P C.O.H. 1812. C p.l
 15595 diffusa Thun. spreading n. \J pr 1J jl.au P C. G. H. 1807. C p.l
 15596 spf nea Lk. spiny <& i_J pr 3 ... P C. G. H. 1824. C p.l
diffusa Thun.

15597 ovata Thun. ovate-headed ■ Lip ■ D. P. C. G. H. 1774. C p.l Bot mag. 186
 15598 spicata Lk. spiked n. Lipr ■ P C. G. H. 1824. C p.l
 15599 ramulosa Lk. small-branched <& Lipr ■ W C. G. H. 1824. C p.l
 15600 bracteata Thun. bracted <& Lipr ■ C. G. H. 1812. C Lip Ven. mal. 10
lucida Ven.
 15601 dentata Poir. toothed <& 1J pr 1} jl.au W C. G. H. 1823. C p.l
teretifolia Lk.
 15602 joogalo'ides Thun. Milkwort-like ■ LIP ■ 1J.0 P C. G. H. 1807. C p.l
 15603 rapunculoides L. Rampion-like tL\Jpr 2 -- V C. G. H. 1824. C p.l
fulvo-maculata Lk.
 15604 fasciculata L. fasciculated <& ti_Jpr 1* jn.jl B C. G. H. 1774. C p.l Bot reg. 184
 15605 rotundifolia Thun. round-leaved 1 jn.jl P C. G. H. 1816. C p.l
 15606 corymbosa L. corymbose ■ 2 jl.au W C. G. H. 1699. C p.l Com. h. 240
 15607 spiria L. spurious ■ 2 jl.au V C. G. H. 1779. S p.l Bur. af. 42.3
 15608 heterophylla Thun. various-leaved <& i_J or 1 jl.au P C. G. H. 1823. C p.l Bur. af. 42.1

1743. rPTEX L. CHASTE TREE. (*Veo*, to bind; flexible twig9.) *Verbenacea*, 12.—28.
 I. MONOPHYLLA.—Leaves simple,
 15609 ovata Thun. ovate-leaved M\| or 4 jl.au P China 1796. C Lp

II. TRIPHYLLO.—Leaves ternate.
 15610 trifK.ra Vahl three-flowered ■ fi — P Cayenne 1819. C 1p
 15611 altissima L. tallest ■ 8 — P Ceylon 1802. C 1p
 15612 arborea Rox. tree ID or . — P
 15613 alata Roth winged . Dor 10 — P E. indies 1820. C 1p
 15614 bicolor IV. en. two-colored ■ 4 — P E. Indies 1810. C 1p
 15615 trifMia L. three-leaved ■ 4 — P E. Indies 1759. C p.l Bot mag. 218
 15616 Negundo L. Negundo ■ 4 — P E. Indies 1812. C 1p Ru.am. 4.19

III. PENTAPIHYLLI.*.—Leaves quinate
 15617 umbrosa Sum. shady f CJ < CO ... 1' Jamaica 1823. C 1p
 15618 A'gnus Castus L. true Chaste Trees or 6 B W.B Sicily 1570. C CO
21atiffia broad-leaved ill or 6 s W.B Sicily 1570. C CO
 15619 incisa Lam. cat-leaved M. \| or 4 jl.s P China 1758. C p.l Bot mag. 364
Negundo B. M.
 15620 Leuc6xylon L. white-wooded ■ LIP - - - ■ Ceylon 1793. C 1p
Wallrothta Leuc6xylon Roth

•1744. CHLOA'NTHES B. Br. CHLOANTHES. (*Chtoa*, greenness, *anthos*, flower.) *Verbenacea*. 2.—3.
 15621 Stee'chadis R. Br. StcechaWife n. LJ or 2 jn.au G.Y N. Holl. 1823. C s.p
 15622 glandulosa R. Br. glandulous n. \| or 2 jn.au G.Y N. Holl. 1824. C s.p

1745. ZAPAWL4J. ZAPANIA. (P. A. Zappa, an Italian botanist.) *Verbenacea*. 2.—10.
 15623 rtoechadifolia L. Stcechas-leaved JU CZ un 1 au.s P W. Indies 1732. C 1p Br. jam. 3.1
Lippia stcechadifoHa Kth.
 15624 nodiflVa Lam. knot-flowered fU LAJ un 1 jl.au P America 1664. C Lp Fl.gr. 553
Perbena nodiflora L. Lippia nodiflora Rich.

1746. PRFFA Adan. PRIVA. (Derivation unknown.) *Verbenacea*, 2.—7.
 15625 mexicana Pers. Mexican ■ E3 pr 2 au.s V Mexico 1726. C 1p DLel.302.3S9
hispida J. Verbena mexicana L.
 15626 /appufcea Pers. Burdock-like ■ (SI or 1) jn.au B W. Indies 1817. C s.l SL jam. 1.110-1
echinhta J.

1747. STRE'PTIUM Rox. STHEPTIUM. (*Streptos*, twisted; spiral tube of corolla.) *Verbenacea*. 1*
 15627 asperum Rox. rough ^ CS or 2 jLau V E Indies 1791. C 1p Rox. cor. 2.1⁴⁶
Priva leptostachya Pers. Toftula aspera IV.

1748. SPIELMA'NNJM Med. SPIELMANNIA. (J.fl.SmeZ/wann,prof.med.andbot.atStrasburg.) *Verbenacea*.*
 15628 africana W. African si LJ or 3 f.n W C. G. H. 1710. C r.m Bot mag. IK*

•1749. FERBEWA L. VEEVAIN. (*Ferfaen*, its Celtic name.) *Verbenacea*, 26.—■
 I. ■ ■ — Leaves undivided.
 15629 angustifolia Mx. narrow-leaved ■ 3 jn.au B N. Amer. 1802. D co
paniculata Lam. panicled ■ 3 jlau B N. Amer. 18X. D co Bot. reg. U
 ■ ■ ■ Buenos Ayres ■ 6 jl.o B B. Ayres 1732. S co Di el. 300. J
 ■ ■ ■ Carolina ■ 6 jn.s W N. Amer. 1732. D co Di cl. 301. J
 Mettle-leaved ■ 3 jl.s W N. Amer. HS3. C co Rob. ic. 26
 many-spiked ■ 4 jlau R Mexico 1820. D p.l
 diffuse ■ 3 jn.au B N. Amer. 1818. D p.l
KSi "T??" H ** a Veronica-leav ■ ijL.au B Mexico 1825. S co
 15637 stneta Ven. strict ■ 3 jl.au B N. Amer. 1802. D co Bot. mag. 1976
 15638 lasidstachys Lk. hairy-spiked ■ Q or 2 jl.au P California 1826. S co

		II. Ternate-leaved									
15639	trifida <i>Hum.</i>	trifid	II	1818.	S	•					
15610	bractcosa <i>Mx.</i>	bracteos	II	Mexico	1820.	S	8				
15641	Aubletfo. <i>Jac.</i>	Aublet's	II	N. Amer.	1774.	S	r.m	Bot. mag.	308		
15642	Lambert [#] <i>B. M.</i>	Lambert's	II	S. Amer.	...	D	co	Bot. mag.	2200		
15643	hastata <i>L.</i>	halberd/caved	II	Canada	1710.	D	co	Her. par.	242		
1564*	pinnatitida <i>Pk.</i>	pinnatilid	II	N. Amer.	181(1).	1)	p.I				
15645	seabra <i>Vahl</i>	scabrous	II	Mexico	1825.	D	r.in				
15646	officinalis <i>L.</i>	officinal	II	Britain	ro. sid.	D	co	Eng. bot.	769		
15647	bifida <i>D. Don</i>	sister	II	Nepal	1824.	S	d	Sw. fl. gar.	202		
15648	spikria <i>L.</i>	spurious	II	N. Amer.	1731.	S	—				
1:3649	sujuna <i>L.</i>	supine	II	Spain	1640.	S	co	Par. th.	675.2		
15650	pmitr'ita <i>It. Br.</i>	prostrate	II	N. Amer.	1794.	D	co				
15651	hiltida <i>A. P.</i>	hispid	II	Peru	1816.	S	s	Fl. per.	34. a.		
15652	caesceis <i>Hum.</i>	canescent	II	Mexico	1824.	S	co				
15653	elegans <i>Hum.</i>	elegant	II	Mexico	1818.	S	co				
15654	ennoldes <i>W.</i>	Erinus-like	II	Peru	1818.	S	co	FL per.	1. 33. C		
	multifida <i>R. & P.</i>										

1750. COLEBROCKKEJ Sm. COLEBROOKEA. (7. *T. Colebrookee*, an accomplished botanist) Verhenhtxa. 2.
 15655 nppoñitifia Sm. opposite-leaved * i.J or S ... W Nepal 1810. C s.l Sm.ex.boLl15
 15656 ternifia Rox. ternate-leaved ft il or 3 ... W E. Indies 1823. C l.p

1751. AVICE'NNJM L. AVICCINNA. (Acicennm, a celebrated Penian philos. and phys.) Mffopdrüue. 1.-3.
 15637 tomentosa L. downy-leaved ± u. 20 ... Pk India 1791. C l.p Beau. ow. 47

* 1752. CLEIIODE'NDIUM L. CTERODENnai'M. (Kleros, lot, dendron, tree j med. uncert) Verben. 38.—40.

		I. QUINQIFEDENTATA. — Calyx Jive-toothed, or almost now:									
15658	ligustrinum R. Br.	Vrwct-leaoed	3 au n	W	Mauritius	1789.	C	p.1	Jac c. sup. 5.1		
15659	pulvrescens Lindl.	pubescent H Q or	4 jl.au	W	W. Indies	ISL/4	C	fp	Bot reg. 1035		
15660	floribundum Lindl.	bundle-lwd	6 jl.au	Li			C	l.p			
15661	iñennic it. Br.	unarmed	4 au u	W	E. Indies	IG/2	C	p.1	Jac. c. sup. 4.1		
15662	angustiflum Spr.	narrow-leaved *	4	—	Mauriliis	1814.	C	s.l	Lam.il. 544.2		
15663	jairiflum Lo.C	Willow-leaved	4 o r	—	V. Indies	1H24.	C	s.l			
15664	coronandri umin Spr.	CoroiMndeHi • or	6	—	Mauritius	18JJ.	C	6.1			
	madagascariense Lo. C.	o'fatuiin Pair.									
15665	volutile Beam.	twining	1 O or	6	W	Guinea	1823.	C	p.1	Beau. ow. 1. 32	
15666	ftuxifbniim Sm.	Box-leaved	* D <	4 au	W	—	—	—	—	182U.	C p.1
	Volkamer*** Ainfolia W. en. *										
15667	heterophyllum R. Br.	various-lvd	■ or	3 aus.s	W	Mauritius	1805.	C	l.p	Bot. rep. 554	
	Volkamer/Vi angibifolia B. Rep.										
15668	macrmhylum B.M.	long-leaved	■ O or	■ jl	W.B	E. Indies	181.2.	C	p.1	Bot. mag. 2536	
15669	rierrktuin D. Don	serrated	■ i CD or	■	Nepal	1822.	C	s.l			
15670	lividum L.	livid	■'LJpr	n'"	China	1824.	C	co	Bot. reg. 945		
15671	cternuum Wai.	drooping	■ Q or	■	E. Indies	IH21.	C	s.l			
5672	natus Wai.	nodding	■ II lor	ilD"	W	E. Indies	—	C	8.1	Bot mag. 3049	
		II. QUINQIEPARTIVA. — Calyx 5-parted :		tm		W		L Indies		C Bur. —	
15673	Siphonanthus R. Br.	Tube-flower	* • or	6	—	W		L Indies	—	C	Bur. —
	Siphonanthus indica L.										
J5674	verticillatum D. Don	whorled-Zi/d	■ ■	6	—	Nepal	1818.	C	l.p		
15675	odoratunt Hum.	scented	■ ■	4	—	Nepal	1821.	C	8.1		
15676	feetidum D. Don	fetid	■ Q or	5	—	Nepal	1820.	C	l.p		
15677	trichiJtonmm Thun.	three-forked	■ i J	6	—	Japan	18<0.	C	l.p	Osb. 11	
15678	fortunatum L.	fortunate	■ CZjor	W	E. Indies	1784.	C	l.p			
15679	ternifilium D. Don	terrate-leaved	■ D	\$"...	—	Nepal	1K23.	C	8.1		
15680	attenuatum R. Br.	attenuated	■	6	—	N. Holl.	1824.	C	8.1		
15681	biquaintrum Vahl	K'iled	a	10. jn.s	S	China	17M.	C	s.p	Bot. reg. 649	
15682	tomcnfsum It. Br.	downy	■ t	5 tnr.ap	W	N. S. W.	1794.	C	s.p	Bot. mag. 1518	
15683	infortunatum /.	unfortunate	■ J or	6	—	E. Indies	...	C	l.p		
15684	costatum It. Br.	ribbed	■ D or	6	—	N. Holl.	1S2J.	C	l.p		
15685	phlomotdes L.	Phlomis-like	■ D or	4 aus.s	W	E. Indies	1820.	C	p.1	Bur. in 45.1	
15686	calamitus L.	calamitous	■ D or	4 jl.o	W	E. Indies	IHii.	C	8.1	Bur. in. 44	
15687	frkgrans Ven.	fragrant	m r~or	6 aud	W	W. R China	1791.	C	sp	Ven inaL70	
	2 flue pheo	double-flowered	H CD or	6 aud	W	W. R China	1790.	C	»p	Bot. mag. 1834	
15G88	viscosum Vat.	clammy	H or	6 niy.au	W	E. Indies	179H.	C	bp	Hot mug. 1805	
15G89	paniculatum L.	panicid	II or	6 jl.o	S	Java	1809.	C	s.p	Bot reg. 495	
15CU	glaucum Wat.	glaucous	II or			E. Indies	ISS-i.	C	8.1		
15ml	helianthifilum Wat.	Stiffflower	• or			E. Indies	1824.	C	8.1		
15692	violaccin L.	bloody-colored	■ 1 lor	+	—	R. Indies	182VJ.	C	p.1		
15693	MrticaefMiuinRox.	Kettle-leaved	III lor	—		R. Indies	1824.	C	8.1		
15694	hastatum WaU	hMxxA-leaved	t Q o r	—	W	E. Indies	18J.0.	C	8.1		
15695	Leucosceprium D. Don	White Spiked	H Q or			Nepal	1826.	C	8.1	Sm.ex.bot 2.116	
	Leucosceprium canuin Sm.										

1753. VOLKAME^RZd L. VOLKAMERIA. (J. G. Volkmer, a German botanist.) Verbence*. «.—a
 15696 aculeata L. nnckly * Q o r 4 au.o W W. Indies 17"», C p.1 Br. jam. 20.2

15697 japonica 2%
n. Japan 1 J or 50 ... P.w Japan 18-U C P.1

1754. HOLMSKIO LD/i RetB. HOLMSKMLDIA. (Th. Holmskiold, a Danish bot author.) Verbenete. 2.
 15698 sanguinea R.tz. bloody * • or 4 ... S Peru 1796. C p.1 Bot.reg. 792

Hastinga coccinea Sm.

15699 scandens Sitt. climbing | • or 5 ap.d S E. Indies 1824. C p.1

Hastiugiascandensfor.

1755. PE^V'RE^ L. PETREA. (Robert James Lord Petre, a botanical patron, d. 1742.) Verbendea? 3. - 6.
 15700 voltililis L. twisting I Q or 20 jl.au P VeraCruz 1733. C r.m Bot mag. 628

15701 erecta Hort. erect m Q or 10 ... B S. Amer. 1823. C r.m

15702 rugosa Hum. wrinkly « D o t 10 ... B Caracas 1824. C r.m

* 1756. JEOLLA'NTHUS Mart. JEoLLktrmvs. (Aiollo, to vary, antkos, flower.) Verbence* L

15703 imyholem Mart. sweet-scented O or 1 jl.au W BraxU 1825. S 81

1757. TAMONIA Aub.	TAMONIA.	(<i>Tamone, its name in Guiana.</i>)	Verbendcete. 2.
15704 curassavica Pen.	Curassoa	■ 1 jl.au B W. Indies 1823. S 81	
verbemacca Swx.	Ghinia spm&sa W,		
15705 mhtica Pers.	awnless	Olor 1 jl.au B Guiana 1820. S s.1	
Ghinia mhtica Swx.			
1758. CITHARE'XYLUM L.	FIDDLE WOOD.	(<i>Kithara, lyre, xylon, wood.</i>)	Verbenacea. 9.-15.
15706 cauditro L.	caudate	1 • or 20 ... W Jamaica 1763. C l.p Jac.ic.3.501	
erectum Jac.			
15707 villfrsum Jac.	villous	■ 10 ■ St. Domin. 1784. C p.1 Jac, i d . 118	
15708 subserratum Swx.	aubserrated	* 15 ■ W W. Indies 1820. C 81	
15709 dentatum Lo. C.	toothed	• 15 ■ £. Indies? 1824. C s.1	
15710 quadrangulare Jac.	square-stalked	1 CD tm 50 ■ W* Jamaica 1759. C p.1 Jac. vin. 1.22	
15711 cincrum L.	dark-colored	i J Q or 15 ■ W W. Indies 1739. C 81	
15712 mscle Jac.	soft	j Q or 15 ■ W W. Indies 1816. C 81 Jac. am. 118	
15713 pentandrim Fen.	pentandroua	II • or 6 ■ W Porto Rico 815. C 81 Yen. eels 47	
15714 sericeum Lo. C.	silky	I • or 15 ■ W £. Indies ?1824. C s.1	
1759. DURA'NIM L.	DIJRANTA.	{ <i>Castor Durantes, a phye, and bot, d. 1590.</i>)	Verbenacea. 7.—10.
W15 Plumieri L.	Plumier's	■ Dor 15 ♀ S. Amer. 1739. C 81	
15716 EUfsia U.	Ellis's	■ 6 au E W. Indies 1739. C P4 Bot mag. 1759	
microphylla Detf.			
15717 incrmis Mil.	unarmed	■ ■ ■ S. Amer. 1739. C p.1	
15718 xalapensis Kth.	Xalapa	■ ■ ■ Mexico 1K22. C p.1	
15719 macrorarpa Kth.	long-fruited	■ ■ ■ W. Indies 1818. C P4	
dentata Pen.			
15720 Mutisia L.	Mutis's	■ ■ ■ W. Indies 1820. C p.1	
15721 argentea Lo. C.	silvery	■ ■ ■ E. Indies ?1824. C p.1	
PEDA'LIMUM L.	PEDALIUM.	(<i>Pedalion, a rudder; dilated angles of fruit</i>)	Pedallnete. 1.
15722 murex L.	caXtop-fruUed	rm.cu \\\ au.s W.P E. Indies 1778. C l.p Lam. il. 538	
• 1761. MYCOPORUM Font.	MYOPORUM.	(<i>Myo, to shut, poros, a pore; leaves.</i>)	Myopofirue. 12.—18.
15723 elf pticum R. Br.	elliptictowo*	• i lor 2 ja.mr W N. S. W. 1789. C l.p Bot rep. 283	
Pogonias glabra Andr.			
15724 acuminatum R. Br.	acuminate	ft L-Jpr 3 ... W N. S. W. 1812. C 1.p	
15725 montanum R. Br.	mountain	ft i lor 2 ... W N. Holl. 1823. C l.p	
15726 parvifolium R. Br.	small-leaved	ft u Jpr 3 year W N. Holl. 1803. C l.p Bot mag. 1691	
15727 tuberculatum R.Br.	tubercl	ft L Jpr 3 - W N. Holl. 1803. C l.p	
15728 insulare R. Br.	insular	ft I or 3 ja.mr W N. S. W. 1800. C p.1	
15729 viscosum R. Br.	viscid	ft i Jpr 3 ... W N. Holl. 1803. C 1.p	
15730 crassifolium Font.	thick-leaved	ft I lor tt ... W N. Zeal. 1822. C l.p	
15731 débile R. Br.	weak	ft - - - W N. S. W. 1793. C l.p Bot mag. 1830	
15732 adscendens R. Br.	ascending	ft - - - W N. Holl. 1820. C l.p	
15733 diff&sum R. Br.	diffuse ft	3 f.au. W N. Holl. - C l.p	
15734 oppositifolium MTr.	opposite-leaved	ft i Jpr 3 year W N. Holl. 1803. C l.p	
• 1762. STENOCHRUS R. Br.	STENOCIHLUS.	(<i>Stenos, narrow, cheUos, a lip; Hower lip.</i>)	Myopdrinace. 1.—3.
15735 glaber R. Br.	naoOthUeaved	* i J or 2 year R N. Holl. 1803. C s.p Bot mag. 1942	
15736 maculatus Ker'	spotted	* LJ or 3 ap.my S N. Holl. 1820. C s.p Bot reg. 647	
15737 longidlius R. Br.	long.leaved	* L J or 4 ap.jl S N. Holl. 1825. C l.p	
1763. BO'NTL* L.	BONTIA.	(<i>J. Bant, M.D., a native of Batavia.</i>)	Myopdrbue. 1.
15738 caphnoides L.	Daphne-like	mcJor 6 jn Y.p W.Indies 169a C p.1 DLed. 49.57	
* 1764. OROBA'NCHE L.	BROOM RAPE.	(<i>Orobos, a vetch, agcho, to strangle.</i>)	OrobanchP*. 6.—31.
15739 mfijor L.	greater	A cu -1jn.jl Br Britain unc.pj. S s.1 Eng. bot. 421	
15740 clatior Suter	taller	■ -jl.au Y Britain clov.fl. S 81 Eng. bot 568	
15741 minor Sm.	smaller	■ I jl.au Y.W Britain clov.fi. S s.1 Eng. bot 422	
15742 rrbra Stn.	red	■ } au P Britain ir.roc. S 81 Eng. bot.c. ic.	
15743 csmMea Fit.	blue	■ V Britain sea co. S s.1 Eng. bot 423	
15744 ramdsia L.	branchy	■ a.us Br.P Britain hemp fi. S s.1 Eng. bot. 84	
1765. CRESCE'NTLI L.	CALABASH TREE.	(<i>Pietro Crescenti, of Bologna, agr. author.</i>)	Saldne<e. 2.-7.
15745 CujUe L.	Cujete	f • cu 20 ... G.w Jamaica 1690. C r.m Jac. am. III	
15746 acuminata Bonn.	acuminate	J • cu 20 ... G.w Cuba 1822. C r.m	
15747 cucurbitina L.	Gourd-fruited	\$ O cu 10 ... W W. Indies 1733. C r.m PI. ic. 109	
1766. TANATCIUM Sun.	TAMECIUM.	(<i>Tanaekes, stretched out; stem.</i>)	Soldnea?. 4.—10.
15748 pinnatum W.	pinnate	I D o r 50 ... R Mozambiq. 1824. C r.m Jac. c. a 18	
Crescentia pinnata Jac.	Tripinnaria africana Spr.		
1767. CASTILLEV.4 Mutis	CASTILLEJA.	(<i>Don Castilleja, a Spanish botanist.</i>)	Scrophuldrigue. 4.
15749 Integrifolia L.	entire-leaved	<.Dor - - - Amer. '1825. C l.p Sm. ic. in. 39	
15750 moranensis Kth.	Moran	a.nor - - Mexico 1825. C s.p	
15751 pfl ida Spr.	pale	^ or i jn.8 L.P Siberia 1782. D s.p Gin. si. 3.24	
Bartsfia pallida L.			
15752 septentrionalis Lindl.	northern	O pr U ■■■ W.o Labrador 1824. S p Bot. reg. 925	
1768. EUCHROMA Nut.	EIJCHROMA.	(<i>EU, well, chroma, color; bracteas.</i>)	Scrojjihuldrigue. 2.
15753 coccinea Nut.	scarlet-bracted	O or jlau Y N. Amer. 1787. S s.p Plukal. 102.5	
Bartsia coccinea L.	CastilK-ja coccinea Spr.		
15754 grandiflora Nut.	great-flowered	^ ^ or 1} jl.au P.T Louisiana 1811. D l.p	
Castilleja sessiliflora Ph.	grandiflbra Spr.		
1769. HALLE'RJI4 L.	HALLERIA.	{ <i>Albert Holler, a disting. bot and physician.</i>)	Scropkulttue. 1.— ^a
15755 liicida L.	shthning-leaved	m i J or 6 jn.au S C. G. H. 1752. C p.1 Bot mag. 1774	
15756 cinctica Thun.	elliptic	* i J or 6 ... S C. G. H. 1816. C p.1 Bur. af. 84.1	
.1770. LATHRÆ? A L.	TOOTHWORT.	(<i>Lathraios, concealed; growing in concealed places.</i>)	Orobuhcheai. F ¹ —
15757 Squamarial.	scaly	£ £ cu 1 ap G Britain dry wo. D co Eng. bot 50	
	YELLOW RATTLE.	(<i>Rhin, snout, anthos, flower.</i>)	Scrophulurite. 4. — W
	Cock's comb	Ow 1jn.a u V Britain mea.pa. S co Eng. bot 657	
	larger	O or 3 jlau Y Britain cornfi. S co	

15760 <i>yflectorblophus</i> <i>Poll.</i>	Cock's comb	O pr	lijn.au	Y	Europe	1880.	■ CO	
15761 <i>JYixigo L.</i>	Trixago	O p?	1 jn-a"	Y	Europe	1800.	■ CO	M.h.&24.8
	jlectorblophus itefi.	Bartsia	Trixkgo L.	TYixago rhinanthina	Zir.			
15772, BA'RTSM L.	BAHTSIA.	(John Bartsck.M.D., a friend of Linneus's)			<i>ScrophOdhrue.</i>	1—10.		
15702 viscbsa L.	viscous	O or	J.lau	Y	Britain	mar.	S m.s	Eng. bot 1045
15763 Odontites flm.	Odontites	O or	*jl.s	Pk	Britain	mea.pa.	S co	Eng. bot 1415
15764 alpna £.	alpine	O or	4jl.au	P	Britain	alp.nv.	S m.s	Eng. bot 361
15765 latifolia Sm.	broad-leaved	O or	1 jl.s	P	& Europe	...	S co	
15766 /infoliai.	EYEBRIGIT.	(<i>Euphraino</i> , to delight; sup. to cureblindnesB.)			<i>Scrophuldrlime.</i>	4.—31.		
15767 flcea L.	Flax-leaved	O or	{ jl.s	P	France	1826. ss	■ CO	
15768 alp'ina Lam.	yellow	O or	lfljl.1	Y	S. Europe	1816.	S co	Jac.au. 398
salisburgensis Funk	alpine	O or	† jl.8	■	Europe	1827.	S co	
15769 officinalia L.	officinal	O w	■ jl.s	W	Britain	past	S co	Eng. bot 1416
• 15770 <i>intirihnum i.</i>	SNAPDRAGON.	(Anti, like, rhin, nose; snout-like flowers.)			<i>Scrophul.</i>	10.—	■	
15770 <i>intirijus L.</i>	greater	jf	2	jn au	PK	England	oldw.	S co Eng. bot 129
2 coccneum	scarlet-/foutfra/£	H	2	jn.au	g	England	oldw.	S co
3 bfcolor	two-colored	f	2	jn.au	S.w	England	oldw.	S co
4 fltre pfeno	double-flowered	H	2	jn.au	F	England	oldw.	S co
15771 meonanthum Lk.	smaller.flwd	£	2	jns	Pk	S Europe	1817.	■ CO
15772 medium Lk.	intermediate	£	2	jns	Pk	Europe	1821.	S co
15773 angustiflbum Poir.	narrow-leaved	■	2	jns	Pk	Europe	1817.	S co
15774 aempcrvlns Lap.	evergreen	■	2	jns	Pk	Pyrenees	1821.	S co
15775 molle L.	soft-leaved	■	1	jl.o	W	Spain	1752.	C s.l
15776 Asarlna L.	Asarina	■	■	W	Italy	1649.	C r.m	Bot mag. 902
15777 sculm Uc.	Sicilian	■	1* jlau	W	Sicily	1804.	D lp	
15778 Grontium L.	Orontium	■	1 jl.s	F	Britain	san.fi.	S &l	Eng. bot 1155
15779 calyndrum Ven.	targe-calyxed	■	1 jl.au	K	■ am	1810.	S co	
IV 1775. LINANRIA Tou.	TOADFLAX.	(<i>Linum</i> , flax; similar leaves.)			<i>Scrophularitue.</i>	70.— 96.		
15780 fr&ticans Spr.	sprouting	I. FftimedSiE. — Small shrubs.	B-Ljor	1 jnjl	Y	C. G. H.	1822.	C s.p
Antirrhinum frticans Thun.								
15781 cokpria Brou.	broom	B - L J O I	1 jn.jl	Y	Teneriffe	1816.	C II	
15782 dalmaticaifo?*.	Dalmatian	a. _Jor	HJn.jl	Y	Levant	1731.	C ss	Bux.c.1.24
II. — Stems prostrate; leaves lobed, angular or toothed from No. 15791 subentire.								
15783 CymbalaYa W.	Cymbalaria	A or	my.n	Y	England	oldw.	1) s.t	Eng. bot 502
J.784 acutiloba Fis.	acute-lobed	i	mys	P	Siberia	1825.	D s.p	
15785 pubescens Presl	pubescent	I, A of	*my.n	Pa	Naples	1850.	D co	
15786 pil,sa Dec.	hairy	t A	i.jns	P	Pyrenees	1800.	D 81	Jacob. 2.48
15787 caufisica Mus.	Caucasian	JL A or	ijlau	Y	Caucasus	1818.	S	
15788 JH&tme Desf.	Elatine	■ O or	i jin	Y	England	corn fi.	S 8.1	Eng. bot. 692
15789 datinoldesi W.	Elafme-like	■ O or	■	Y	S. Europe	1821.	S CO	Deaf. at 2.132
J5790 cirrhosa IV.	tendrilled	■ O or	■	Pa.B	Egypt	177J.	S 8.1	Jac. vin. 1.82
J5791 sptria W.	spurious	■ O or	■	Y	England	corn fi.	S CO	Eng. bot. 691
15792 lanlera Desf.	wool-bearing	Ji O or	jns	Y	Portugal	1818.	S CO	
15793 dealbata Lk.	whited	** O or	i jl.s	Y	Portugal	182a	S co	
Antirrhinum lanferum Bro.								
III. ERE'CTA. — Flowering stems erect; lower leaves opposite or i>horledfram No. 15841 scattered.								
15794 latifolia Dec.	broad-leaved	O or	■	■	N Amer.	1800.	■ co	Desf. at. 2.134
15795 triphlla W.	three-leaved	O or	■	■	Sicily	1596.	s.l	Bot. mag. 324
J.796 re>xa Desf.	reflexed	O or	■	■	S. Europe	1810.	■	Al. tau. 2.1
J.797 origanifilia Bee.	Marjoram.lvd	■	■	■	S. Europe	1785.	■	Bar. ic. 598
Ja7M virgaSita Desf.	twiggy	■	■	■	N. Africa	1817.	■	Desf. at 2.135
JS79D cretXcea Fis.	cretaceous	■	■	■	Siberia	1827.	8.1	
L78(K) /hymitilla Dec.	Thyme-leaved	■	■	■	S. Europe	1818.	CO	
15801 fl^va Desf.	yellow	■	■	■	N. Africa	1820.	CO	Desf. at 2.136
15802 tranataptina Spr.	beyond the Tagus	■	■	■	Portugal	1810.	■ BI	
Antirrhinum lusitanicum Lk. pedunculatum Poir.								
15803 alsinifilia Spr.	Alsine-leaved	■	■	■	Corsica	1834.	S s.l	
15804 villbsa Spr.	villous	■	■	■	Spain	1786.	D 1P	Bar. ic. 597
IBNxi rubrifolia Dec	red-leaved	■	■	■	S. France	1826.	S 8.1	
15806 saxatilis Dec.	rock	■	■	■	Spain	1819.	S 8.1	
15807 hirta W.	hairy-leaved	■	■	■	Spain	1759.	S co	Jac. ic. I.II7
15808 triornithophoraSr.	tree-birds.bgs	■	■	■	Portugal	1710.	C s.p	Bot mag. 523
15809 bipartite W.	twb-parted	■	■	■	Barbary	1815.	D lp	Sw. fl. gar.SO
speciosa Jac.								
15810 minor Dec.	small	O w	■	■	England	san.fi.	S ■	Eng. bot. 2014
15811 hittoralis IV.	seaside	* O or	■	■	Austria	J820.	S 8.1	
15812 chalepensis Aff.	Aleppo	O or	■	■	Levant	1680.	S co	M.h.5.35.9
J5813 tristis Mil.	dark	£	-Alor	■	Spain	1727.	S 8.1	Bot. mag. 74
15814 versicolor Moen.	various-colored	O or	■	■	France	1777.	S 8.1	Jac. ic. I.116
15815 hlans S.	lepting	A or	■	■	S. Europe	1818.	D 8.1	
Antirrhinum larinorhiza Spr.								
15816 micrantha Spr.	small-flowered	* ■	■	■		1820.	S ■	Cav.ic. 1.69.3
15817 viscbsa Spr.	clammy	■	■	■	Spain	1786.	S 8.1	Bot. mag. 368
15818 purpCirea Mil.	purple	■	■	■	S. Europe	1648.	S co	Bot. mag. 99
15819 arenaria Dec.	sand	■	■	■	S. Europe	1823.	S 8.1	
15820 alpna Dec.	alpine	■	■	■	Austria	1570.	C 8.1	Bot. mag. 205
15821 lusitanica Tou.	Spanish	■	■	■	Spain	1819.	S 8.1	
15822 repens H. K.	Pyrenean	■	■	■	England ch. hil.	S CO		Eng. bot 1253
15823 pyrenaica Dec.		■	■	■	Pyrenees	1881.	S 8.1	
15824 Heelhva W.	Hslava	■	■	■	■	1803.	S co	
15825 Pelissenfyna Dec.	Pelisser's	■	■	■	Mropel640.	S 8.1	Bar. ic. 1162	
15826 proceraDr.	tall	■	■	■	S. Europe	■ D	co	
15827 ^alioides Ven.	Galium-like	■	■	■	S. France	■ D	co	
15828 monspessulana L.	Montpellier	■	■	■	S. France	■ D GO		
J5J29 amethystina Spr.	amethystine	O or	■ jn.jl	B.T	Spain	■ 13	0 w tI * »	
15830 suphna Dec.	supine	Jc Q or	■ jl	rY	Spain			

* 1777. ACME'S IA Fen. NEMESIA. (Applied by Dioscorides to a kind of Antirrhinum.) *Scrophularina*. S.—*. 15853 *Chamaelittula* Henr. *Chamaelitella* Lvd & LAJ or 2 ap s P C. G. H. 1787. D co

15854 fastens <i>Fen.</i>	stinking	■ iAl	2 ap.s	P	C. G. H.	1798.	Deo	Ven. mai.41
15855 bicornis <i>Pers.</i>	two-horned	O or	2 jl.au	P	C. G. H.	1774.	S ed	Bur. af. 75.3

1778. MAURA'NDYA Jac. MAURANDYA. (*Dr. Maurandy*, a bot. prof. at Carthagena.) *Scrophularin.* 3.
 15856 sem¹ cfrrlflorens Jac. flowering f. i_J or 10 year P Mexico 1796. C l.p Bot mag. 410
 15857 autirrhiniflora Jftiw. Snajdrugr-flwd fc i_J or 10 year P Mexico 1814. C l.p Bot mag. 1643
 15858 Barclayd/w B. R. Barclay's % _J or 10 year B.w Mexico 1825. C l.p Bot reg. 1108

1779. GERA'R'DJA L.	GERARDIA.	(John Gerarde, an old English botanist.)	Scrophuidirue.	6.-26.
15&59 rfcphinfiltia L.	Larkspur-lvd	& E2 or 2 jn.jl Pk	E. Indies 1800.	C l.p Rox. cor. 1 IK
158H0 purpurea L.	purple	1* jl.au	N. Amer. 1772.	S s.1 Bot mag. 2048
15S61 tenuifolia Vahl	slender-leaved	1 jl.au	N. Amer. 1812.	S s.1 Pluk.al. 12.4
15862 maritima Rafi.	seaside	2 jl.au	N. Amer. 1823.	S Lp
1586 S flvka L.	yellow	1jjl. au	D l.p	Pluk.am.389.3
15864 queriefolia Ph.	Oak-leaved	4 jl.au	N. Amer. 1796.	Ph. am. 19

1780. SEYME^XR/A Ph. SETMERIA. (*Henry Seymour, an English naturalist*) 2.-4.
15865* *tenuifolia Ph.* fine-leaved O I¹ J¹^{su} N Amer 1780 b 8 P

15873 <i>Resupinata</i> L.		resupinate	IL MU^TICJB.— <i>Helmet bluntnis/i, bUleatate; calyx bifid, from 15879. quinquefl.</i>						
15876	<i>paltuſtris</i> L.	marsh	2	jn.jl	P	Britain	bog.m.	S	Eng. bot 399
15877	<i>euph rasiokles</i> Stev.	Eyebright-like			P	Siberia	1816.	S	Gin. si. 3.43
15878	<i>canajensis</i> L.	Canadian	<i>U</i>	jl.au	Y	N. Amer.	1810.	S	Sw. fl. gar 67
15879	<i>sylvatica</i> L.	wood	1	my.jl	Pk	Britain	ra.he.a.	S	Eng. bot 400
15880	<i>myriophyllum</i> Pall.	myriad-leaved	1	my.jl	Y	Dauria	1816.	P	Pal. it. J. ap. 8. *
15881	<i>cinois</i> L.	tufted		jl.au	Y	Italy	1775.	P	Al. cpld. 1.4.1
15882	<i>Scutellaria</i> Caro Ufwmk.	circumcisus	5	au	P	Sweden		P	F. L dan. 26 ^
15883	<i>Scutellaria</i> Charles' Sc.					Austria	1787.	P	Jac. au. 3. 258
15884	<i>foliosa</i> L.	leafy			Y	Austria	1784.	P	Jac. au. 2. 139
15885	<i>verticillata</i> L.	whorled			my.jn	Austria	1790.	P	Jac. au. 3. 206
15886	<i>flammea</i> L.	flame			Y.8	Switzerl.	1775.	P	Hal. h. 8.3
15887	<i>versicolor</i> Wahl	party-colored			jn.au	Y.R	Switzerl.	1819.	

1782		ERINUS. (<i>Erino</i> Sy a wild fig tree; applied to this plant.) <i>Scrophulariace.</i> 5.—	
15889	<i>alpinus L.</i>	smooth alpine A or mr.ap B Pyrenees 1739. D s.l.	
15K89	<i>hisp&nicus Pers.</i>	hairy Spanish A or mr.ap R Spain 1739. D B.I.	
15890	<i>fragrans H. K.</i>	fragrant n. i. ^J or my.in Y.w C G. H. 1776. C 8.1 Bur. af. 4&J	
15H31	<i>Ljphilicla Thun.</i>	my.in Y.w C G. H. ... C 8.1 Bot. reg. 7<	
	<i>tristia Thun.</i>	Lyclinidca n. my.in P C G. H. 1825. C 8.1	
		<i>dark-flowered</i> n.	
* 1783.	3/I'MULUS L.	(<i>Mimo</i> , an ape; seeds like face.) <i>Scrophulariace.</i> 10.—	lf.,-
15893	<i>ringen* L.</i>	MONKEY FLOWER. or JIau L.B. N. Amer. 1759. D p.l. Bot mitt 2 »»	
15894	I ^w s ^p s Ph.	gaping A or f au Pa.P Missouri 1824. D p.l. Ph. am. 2	
15895	¹⁸ ¹⁹ ²⁰ ²¹ ²² ²³ ²⁴ ²⁵ ²⁶ ²⁷ ²⁸ ²⁹ ³⁰ ³¹ ³² ³³ ³⁴ ³⁵ ³⁶ ³⁷ ³⁸ ³⁹ ⁴⁰ ⁴¹ ⁴² ⁴³ ⁴⁴ ⁴⁵ ⁴⁶ ⁴⁷ ⁴⁸ ⁴⁹ ⁵⁰ ⁵¹ ⁵² ⁵³ ⁵⁴ ⁵⁵ ⁵⁶ ⁵⁷ ⁵⁸ ⁵⁹ ⁶⁰ ⁶¹ ⁶² ⁶³ ⁶⁴ ⁶⁵ ⁶⁶ ⁶⁷ ⁶⁸ ⁶⁹ ⁷⁰ ⁷¹ ⁷² ⁷³ ⁷⁴ ⁷⁵ ⁷⁶ ⁷⁷ ⁷⁸ ⁷⁹ ⁸⁰ ⁸¹ ⁸² ⁸³ ⁸⁴ ⁸⁵ ⁸⁶ ⁸⁷ ⁸⁸ ⁸⁹ ⁹⁰ ⁹¹ ⁹² ⁹³ ⁹⁴ ⁹⁵ ⁹⁶ ⁹⁷ ⁹⁸ ⁹⁹ ¹⁰⁰ ¹⁰¹ ¹⁰² ¹⁰³ ¹⁰⁴ ¹⁰⁵ ¹⁰⁶ ¹⁰⁷ ¹⁰⁸ ¹⁰⁹ ¹¹⁰ ¹¹¹ ¹¹² ¹¹³ ¹¹⁴ ¹¹⁵ ¹¹⁶ ¹¹⁷ ¹¹⁸ ¹¹⁹ ¹²⁰ ¹²¹ ¹²² ¹²³ ¹²⁴ ¹²⁵ ¹²⁶ ¹²⁷ ¹²⁸ ¹²⁹ ¹³⁰ ¹³¹ ¹³² ¹³³ ¹³⁴ ¹³⁵ ¹³⁶ ¹³⁷ ¹³⁸ ¹³⁹ ¹⁴⁰ ¹⁴¹ ¹⁴² ¹⁴³ ¹⁴⁴ ¹⁴⁵ ¹⁴⁶ ¹⁴⁷ ¹⁴⁸ ¹⁴⁹ ¹⁵⁰ ¹⁵¹ ¹⁵² ¹⁵³ ¹⁵⁴ ¹⁵⁵ ¹⁵⁶ ¹⁵⁷ ¹⁵⁸ ¹⁵⁹ ¹⁶⁰ ¹⁶¹ ¹⁶² ¹⁶³ ¹⁶⁴ ¹⁶⁵ ¹⁶⁶ ¹⁶⁷ ¹⁶⁸ ¹⁶⁹ ¹⁷⁰ ¹⁷¹ ¹⁷² ¹⁷³ ¹⁷⁴ ¹⁷⁵ ¹⁷⁶ ¹⁷⁷ ¹⁷⁸ ¹⁷⁹ ¹⁸⁰ ¹⁸¹ ¹⁸² ¹⁸³ ¹⁸⁴ ¹⁸⁵ ¹⁸⁶ ¹⁸⁷ ¹⁸⁸ ¹⁸⁹ ¹⁹⁰ ¹⁹¹ ¹⁹² ¹⁹³ ¹⁹⁴ ¹⁹⁵ ¹⁹⁶ ¹⁹⁷ ¹⁹⁸ ¹⁹⁹ ²⁰⁰ ²⁰¹ ²⁰² ²⁰³ ²⁰⁴ ²⁰⁵ ²⁰⁶ ²⁰⁷ ²⁰⁸ ²⁰⁹ ²¹⁰ ²¹¹ ²¹² ²¹³ ²¹⁴ ²¹⁵ ²¹⁶ ²¹⁷ ²¹⁸ ²¹⁹ ²²⁰ ²²¹ ²²² ²²³ ²²⁴ ²²⁵ ²²⁶ ²²⁷ ²²⁸ ²²⁹ ²³⁰ ²³¹ ²³² ²³³ ²³⁴ ²³⁵ ²³⁶ ²³⁷ ²³⁸ ²³⁹ ²⁴⁰ ²⁴¹ ²⁴² ²⁴³ ²⁴⁴ ²⁴⁵ ²⁴⁶ ²⁴⁷ ²⁴⁸ ²⁴⁹ ²⁵⁰ ²⁵¹ ²⁵² ²⁵³ ²⁵⁴ ²⁵⁵ ²⁵⁶ ²⁵⁷ ²⁵⁸ ²⁵⁹ ²⁶⁰ ²⁶¹ ²⁶² ²⁶³ ²⁶⁴ ²⁶⁵ ²⁶⁶ ²⁶⁷ ²⁶⁸ ²⁶⁹ ²⁷⁰ ²⁷¹ ²⁷² ²⁷³ ²⁷⁴ ²⁷⁵ ²⁷⁶ ²⁷⁷ ²⁷⁸ ²⁷⁹ ²⁸⁰ ²⁸¹ ²⁸² ²⁸³ ²⁸⁴ ²⁸⁵ ²⁸⁶ ²⁸⁷ ²⁸⁸ ²⁸⁹ ²⁹⁰ ²⁹¹ ²⁹² ²⁹³ ²⁹⁴ ²⁹⁵ ²⁹⁶ ²⁹⁷ ²⁹⁸ ²⁹⁹ ³⁰⁰ ³⁰¹ ³⁰² ³⁰³ ³⁰⁴ ³⁰⁵ ³⁰⁶ ³⁰⁷ ³⁰⁸ ³⁰⁹ ³¹⁰ ³¹¹ ³¹² ³¹³ ³¹⁴ ³¹⁵ ³¹⁶ ³¹⁷ ³¹⁸ ³¹⁹ ³²⁰ ³²¹ ³²² ³²³ ³²⁴ ³²⁵ ³²⁶ ³²⁷ ³²⁸ ³²⁹ ³³⁰ ³³¹ ³³² ³³³ ³³⁴ ³³⁵ ³³⁶ ³³⁷ ³³⁸ ³³⁹ ³⁴⁰ ³⁴¹ ³⁴² ³⁴³ ³⁴⁴ ³⁴⁵ ³⁴⁶ ³⁴⁷ ³⁴⁸ ³⁴⁹ ³⁵⁰ ³⁵¹ ³⁵² ³⁵³ ³⁵⁴ ³⁵⁵ ³⁵⁶ ³⁵⁷ ³⁵⁸ ³⁵⁹ ³⁶⁰ ³⁶¹ ³⁶² ³⁶³ ³⁶⁴ ³⁶⁵ ³⁶⁶ ³⁶⁷ ³⁶⁸ ³⁶⁹ ³⁷⁰ ³⁷¹ ³⁷² ³⁷³ ³⁷⁴ ³⁷⁵ ³⁷⁶ ³⁷⁷ ³⁷⁸ ³⁷⁹ ³⁸⁰ ³⁸¹ ³⁸² ³⁸³ ³⁸⁴ ³⁸⁵ ³⁸⁶ ³⁸⁷ ³⁸⁸ ³⁸⁹ ³⁹⁰ ³⁹¹ ³⁹² ³⁹³ ³⁹⁴ ³⁹⁵ ³⁹⁶ ³⁹⁷ ³⁹⁸ ³⁹⁹ ⁴⁰⁰ ⁴⁰¹ ⁴⁰² ⁴⁰³ ⁴⁰⁴ ⁴⁰⁵ ⁴⁰⁶ ⁴⁰⁷ ⁴⁰⁸ ⁴⁰⁹ ⁴¹⁰ ⁴¹¹ ⁴¹² ⁴¹³ ⁴¹⁴ ⁴¹⁵ ⁴¹⁶ ⁴¹⁷ ⁴¹⁸ ⁴¹⁹ ⁴²⁰ ⁴²¹ ⁴²² ⁴²³ ⁴²⁴ ⁴²⁵ ⁴²⁶ ⁴²⁷ ⁴²⁸ ⁴²⁹ ⁴³⁰ ⁴³¹ ⁴³² ⁴³³ ⁴³⁴ ⁴³⁵ ⁴³⁶ ⁴³⁷ ⁴³⁸ ⁴³⁹ ⁴⁴⁰ ⁴⁴¹ ⁴⁴² ⁴⁴³ ⁴⁴⁴ ⁴⁴⁵ ⁴⁴⁶ ⁴⁴⁷ ⁴⁴⁸ ⁴⁴⁹ ⁴⁵⁰ ⁴⁵¹ ⁴⁵² ⁴⁵³ ⁴⁵⁴ ⁴⁵⁵ ⁴⁵⁶ ⁴⁵⁷ ⁴⁵⁸ ⁴⁵⁹ ⁴⁶⁰ ⁴⁶¹ ⁴⁶² ⁴⁶³ ⁴⁶⁴ ⁴⁶⁵ ⁴⁶⁶ ⁴⁶⁷ ⁴⁶⁸ ⁴⁶⁹ ⁴⁷⁰ ⁴⁷¹ ⁴⁷² ⁴⁷³ ⁴⁷⁴ ⁴⁷⁵ ⁴⁷⁶ ⁴⁷⁷ ⁴⁷⁸ ⁴⁷⁹ ⁴⁸⁰ ⁴⁸¹ ⁴⁸² ⁴⁸³ ⁴⁸⁴ ⁴⁸⁵ ⁴⁸⁶ ⁴⁸⁷ ⁴⁸⁸ ⁴⁸⁹ ⁴⁹⁰ ⁴⁹¹ ⁴⁹² ⁴⁹³ ⁴⁹⁴ ⁴⁹⁵ ⁴⁹⁶ ⁴⁹⁷ ⁴⁹⁸ ⁴⁹⁹ ⁵⁰⁰ ⁵⁰¹ ⁵⁰² ⁵⁰³ ⁵⁰⁴ ⁵⁰⁵ ⁵⁰⁶ ⁵⁰⁷ ⁵⁰⁸ ⁵⁰⁹ ⁵¹⁰ ⁵¹¹ ⁵¹² ⁵¹³ ⁵¹⁴ ⁵¹⁵ ⁵¹⁶ ⁵¹⁷ ⁵¹⁸ ⁵¹⁹ ⁵²⁰ ⁵²¹ ⁵²² ⁵²³ ⁵²⁴ ⁵²⁵ ⁵²⁶ ⁵²⁷ ⁵²⁸ ⁵²⁹ ⁵³⁰ ⁵³¹ ⁵³² ⁵³³ ⁵³⁴ ⁵³⁵ ⁵³⁶ ⁵³⁷ ⁵³⁸ ⁵³⁹ ⁵⁴⁰ ⁵⁴¹ ⁵⁴² ⁵⁴³ ⁵⁴⁴ ⁵⁴⁵ ⁵⁴⁶ ⁵⁴⁷ ⁵⁴⁸ ⁵⁴⁹ ⁵⁵⁰ ⁵⁵¹ ⁵⁵² ⁵⁵³ ⁵⁵⁴ ⁵⁵⁵ ⁵⁵⁶ ⁵⁵⁷ ⁵⁵⁸ ⁵⁵⁹ ⁵⁶⁰ ⁵⁶¹ ⁵⁶² ⁵⁶³ ⁵⁶⁴ ⁵⁶⁵ ⁵⁶⁶ ⁵⁶⁷ ⁵⁶⁸ ⁵⁶⁹ ⁵⁷⁰ ⁵⁷¹ ⁵⁷² ⁵⁷³ ⁵⁷⁴ ⁵⁷⁵ ⁵⁷⁶ ⁵⁷⁷ ⁵⁷⁸ ⁵⁷⁹ ⁵⁸⁰ ⁵⁸¹ ⁵⁸² ⁵⁸³ ⁵⁸⁴ ⁵⁸⁵ ⁵⁸⁶ ⁵⁸⁷ ⁵⁸⁸ ⁵⁸⁹ ⁵⁹⁰ ⁵⁹¹ ⁵⁹² ⁵⁹³ ⁵⁹⁴ ⁵⁹⁵ ⁵⁹⁶ ⁵⁹⁷ ⁵⁹⁸ ⁵⁹⁹ ⁶⁰⁰ ⁶⁰¹ ⁶⁰² ⁶⁰³ ⁶⁰⁴ ⁶⁰⁵ ⁶⁰⁶ ⁶⁰⁷ ⁶⁰⁸ ⁶⁰⁹ ⁶¹⁰ ⁶¹¹ ⁶¹² ⁶¹³ ⁶¹⁴ ⁶¹⁵ ⁶¹⁶ ⁶¹⁷ ⁶¹⁸ ⁶¹⁹ ⁶²⁰ ⁶²¹ ⁶²² ⁶²³ ⁶²⁴ ⁶²⁵ ⁶²⁶ ⁶²⁷ ⁶²⁸ ⁶²⁹ ⁶³⁰ ⁶³¹ ⁶³² ⁶³³ ⁶³⁴ ⁶³⁵ ⁶³⁶ ⁶³⁷ ⁶³⁸ ⁶³⁹ ⁶⁴⁰ ⁶⁴¹ ⁶⁴² ⁶⁴³ ⁶⁴⁴ ⁶⁴⁵ ⁶⁴⁶ ⁶⁴⁷ ⁶⁴⁸ ⁶⁴⁹ ⁶⁵⁰ ⁶⁵¹ ⁶⁵² ⁶⁵³ ⁶⁵⁴ ⁶⁵⁵ ⁶⁵⁶ ⁶⁵⁷ ⁶⁵⁸ ⁶⁵⁹ ⁶⁶⁰ ⁶⁶¹ ⁶⁶² ⁶⁶³ ⁶⁶⁴ ⁶⁶⁵ ⁶⁶⁶ ⁶⁶⁷ ⁶⁶⁸ ⁶⁶⁹ ⁶⁷⁰ ⁶⁷¹ ⁶⁷² ⁶⁷³ ⁶⁷⁴ ⁶⁷⁵ ⁶⁷⁶ ⁶⁷⁷ ⁶⁷⁸ ⁶⁷⁹ ⁶⁸⁰ ⁶⁸¹ ⁶⁸² ⁶⁸³ ⁶⁸⁴ ⁶⁸⁵ ⁶⁸⁶ ⁶⁸⁷ ⁶⁸⁸ ⁶⁸⁹ ⁶⁹⁰ ⁶⁹¹ ⁶⁹² ⁶⁹³ ⁶⁹⁴ ⁶⁹⁵ ⁶⁹⁶ ⁶⁹⁷ ⁶⁹⁸ ⁶⁹⁹ ⁷⁰⁰ ⁷⁰¹ ⁷⁰² ⁷⁰³ ⁷⁰⁴ ⁷⁰⁵ ⁷⁰⁶ ⁷⁰⁷ ⁷⁰⁸ ⁷⁰⁹ ⁷¹⁰ ⁷¹¹ ⁷¹² ⁷¹³ ⁷¹⁴ ⁷¹⁵ ⁷¹⁶ ⁷¹⁷ ⁷¹⁸ ⁷¹⁹ ⁷²⁰ ⁷²¹ ⁷²² ⁷²³ ⁷²⁴ ⁷²⁵ ⁷²⁶ ⁷²⁷ ⁷²⁸ ⁷²⁹ ⁷³⁰ ⁷³¹ ⁷³² ⁷³³ ⁷³⁴ ⁷³⁵ ⁷³⁶ ⁷³⁷ ⁷³⁸ ⁷³⁹ ⁷⁴⁰ ⁷⁴¹ ⁷⁴² ⁷⁴³ ⁷⁴⁴ ⁷⁴⁵ ⁷⁴⁶ ⁷⁴⁷ ⁷⁴⁸ ⁷⁴⁹ ⁷⁵⁰ ⁷⁵¹ ⁷⁵² ⁷⁵³ ⁷⁵⁴ ⁷⁵⁵ ⁷⁵⁶ ⁷⁵⁷ ⁷⁵⁸ ⁷⁵⁹ ⁷⁶⁰ ⁷⁶¹ ⁷⁶² ⁷⁶³ ⁷⁶⁴ ⁷⁶⁵ ⁷⁶⁶ ⁷⁶⁷ ⁷⁶⁸ ⁷⁶⁹ ⁷⁷⁰ ⁷⁷¹ ⁷⁷² ⁷⁷³ ⁷⁷⁴ ⁷⁷⁵ ⁷⁷⁶ ⁷⁷⁷ ⁷⁷⁸ ⁷⁷⁹ ⁷⁸⁰ ⁷⁸¹ ⁷⁸² ⁷⁸³ ⁷⁸⁴ ⁷⁸⁵ ⁷⁸⁶ ⁷⁸⁷ ⁷⁸⁸ ⁷⁸⁹ ⁷⁹⁰ ⁷⁹¹ ⁷⁹² ⁷⁹³ ⁷⁹⁴ ⁷⁹⁵ ⁷⁹⁶ ⁷⁹⁷ ⁷⁹⁸ ⁷⁹⁹ ⁸⁰⁰ ⁸⁰¹ ⁸⁰² ⁸⁰³ ⁸⁰⁴ ⁸⁰⁵ ⁸⁰⁶ ⁸⁰⁷ ⁸⁰⁸ ⁸⁰⁹ ⁸⁰¹⁰ ⁸⁰¹¹ ⁸⁰¹² ⁸⁰¹³ ⁸⁰¹⁴ ⁸⁰¹⁵ ⁸⁰¹⁶ ⁸⁰¹⁷ ⁸⁰¹⁸ ⁸⁰¹⁹ ⁸⁰²⁰ ⁸⁰²¹ ⁸⁰²² ⁸⁰²³ ⁸⁰²⁴ ⁸⁰²⁵ ⁸⁰²⁶ ⁸⁰²⁷ ⁸⁰²⁸ ⁸⁰²⁹ ⁸⁰³⁰ ⁸⁰³¹ ⁸⁰³² ⁸⁰³³ ⁸⁰³⁴ ⁸⁰³⁵ ⁸⁰³⁶ ⁸⁰³⁷ ⁸⁰³⁸ ⁸⁰³⁹ ⁸⁰⁴⁰ ⁸⁰⁴¹ ⁸⁰⁴² ⁸⁰⁴³ ⁸⁰⁴⁴ ⁸⁰⁴⁵ ⁸⁰⁴⁶ ⁸⁰⁴⁷ ⁸⁰⁴⁸ ⁸⁰⁴⁹ ⁸⁰⁵⁰ ⁸⁰⁵¹ ⁸⁰⁵² ⁸⁰⁵³ ⁸⁰⁵⁴ ⁸⁰⁵⁵ ⁸⁰⁵⁶ ⁸⁰⁵⁷ ⁸⁰⁵⁸ ⁸⁰⁵⁹ ⁸⁰⁶⁰ ⁸⁰⁶¹ ⁸⁰⁶² ⁸⁰⁶³ ⁸⁰⁶⁴ ⁸⁰⁶⁵ ⁸⁰⁶⁶ ⁸⁰⁶⁷ ⁸⁰⁶⁸ ⁸⁰⁶⁹ ⁸⁰⁷⁰ ⁸⁰⁷¹ ⁸⁰⁷² ⁸⁰⁷³ ⁸⁰⁷⁴ ⁸⁰⁷⁵ ⁸⁰⁷⁶ ⁸⁰⁷⁷ ⁸⁰⁷⁸ ⁸⁰⁷⁹ ⁸⁰⁸⁰ ⁸⁰⁸¹ ⁸⁰⁸² ⁸⁰⁸³ ⁸⁰⁸⁴ ⁸⁰⁸⁵ ⁸⁰⁸⁶ ⁸⁰⁸⁷ ⁸⁰⁸⁸ ⁸⁰⁸⁹ ⁸⁰⁹⁰ ⁸⁰⁹¹ ⁸⁰⁹² ⁸⁰⁹³ ⁸⁰⁹⁴ ⁸⁰⁹⁵ ⁸⁰⁹⁶ ⁸⁰⁹⁷ ⁸⁰⁹⁸ ⁸⁰⁹⁹ ⁸⁰¹⁰⁰ ⁸⁰¹⁰¹ ⁸⁰¹⁰² ⁸⁰¹⁰³ ⁸⁰¹⁰⁴ ⁸⁰¹⁰⁵ ⁸⁰¹⁰⁶ ⁸⁰¹⁰⁷ ⁸⁰¹⁰⁸ ⁸⁰¹⁰⁹ ⁸⁰¹¹⁰ ⁸⁰¹¹¹ ⁸⁰¹¹² ⁸⁰¹¹³ ⁸⁰¹¹⁴ ⁸⁰¹¹⁵ ⁸⁰¹¹⁶ ⁸⁰¹¹⁷ ⁸⁰¹¹⁸ ⁸⁰¹¹⁹ ⁸⁰¹²⁰ ⁸⁰¹²¹ ⁸⁰¹²² ⁸⁰¹²³ ⁸⁰¹²⁴ ⁸⁰¹²⁵ ⁸⁰¹²⁶ ⁸⁰¹²⁷ ⁸⁰¹²⁸ ⁸⁰¹²⁹ ⁸⁰¹³⁰ ⁸⁰¹³¹ ⁸⁰¹³² ⁸⁰¹³³ ⁸⁰¹³⁴ ⁸⁰¹³⁵ ⁸⁰¹³⁶ ⁸⁰¹³⁷ ⁸⁰¹³⁸ ⁸⁰¹³⁹ ⁸⁰¹⁴⁰ ⁸⁰¹⁴¹ ⁸⁰¹⁴² ⁸⁰¹⁴³ ⁸⁰¹⁴⁴ ⁸⁰¹⁴⁵ ⁸⁰¹⁴⁶ ⁸⁰¹⁴⁷ ⁸⁰¹⁴⁸ ⁸⁰¹⁴⁹ ⁸⁰¹⁵⁰ ⁸⁰¹⁵¹ ⁸⁰¹⁵² ⁸⁰¹⁵³ ⁸⁰¹⁵⁴ ⁸⁰¹⁵⁵ ⁸⁰¹⁵⁶ ⁸⁰¹⁵⁷ ⁸⁰¹⁵⁸ ⁸⁰¹⁵⁹ ⁸⁰¹⁶⁰ ⁸⁰¹⁶¹ ⁸⁰¹⁶² ⁸⁰¹⁶³ ⁸⁰¹⁶⁴ ⁸⁰¹⁶⁵ ⁸⁰¹⁶⁶ ⁸⁰¹⁶⁷ ⁸⁰¹⁶⁸ ⁸⁰¹⁶⁹ ⁸⁰¹⁷⁰ ⁸⁰¹⁷¹ ⁸⁰¹⁷² ⁸⁰¹⁷³ ⁸⁰¹⁷⁴ ⁸⁰¹⁷⁵ ⁸⁰¹⁷⁶ ⁸⁰¹⁷⁷ ⁸⁰¹⁷⁸ ⁸⁰¹⁷⁹ ⁸⁰¹⁸⁰ ⁸⁰¹⁸¹ ⁸⁰¹⁸² ⁸⁰¹⁸³ ⁸⁰¹⁸⁴ ⁸⁰¹⁸⁵ ⁸⁰¹⁸⁶ ⁸⁰¹⁸⁷ ⁸⁰¹⁸⁸ ⁸⁰¹⁸⁹ ⁸⁰¹⁹⁰ ⁸⁰¹⁹¹ ⁸⁰¹⁹² ⁸⁰¹⁹³ ⁸⁰¹⁹⁴ ⁸⁰¹⁹⁵ ⁸⁰¹⁹⁶ ⁸⁰¹⁹⁷ ⁸⁰¹⁹⁸ ⁸⁰¹⁹⁹ ⁸⁰²⁰⁰ ⁸⁰²⁰¹ ⁸⁰²⁰² ⁸⁰²⁰³ ⁸⁰²⁰⁴ ⁸⁰²⁰⁵ ⁸⁰²⁰⁶ ⁸⁰²⁰⁷ ⁸⁰²⁰⁸ ⁸⁰²⁰⁹ ⁸⁰²¹⁰ ⁸⁰²¹¹ ⁸⁰²¹² ⁸⁰²¹³ ⁸⁰²¹⁴ ⁸⁰²¹⁵ ⁸⁰²¹⁶ ⁸⁰²¹⁷ ⁸⁰²¹⁸ ⁸⁰²¹⁹ ⁸⁰²²⁰ ⁸⁰²²¹ ⁸⁰²²² ⁸⁰²²³ ⁸⁰²²⁴ ⁸⁰²²⁵ ⁸⁰²²⁶ ⁸⁰²²⁷ ⁸⁰²²⁸ ⁸⁰²²⁹ ⁸⁰²³⁰ ⁸⁰²³¹ ⁸⁰²³² ⁸⁰²³³ ⁸⁰²³⁴ ⁸⁰²³⁵ ⁸⁰²³⁶ ⁸⁰²³⁷ ⁸⁰²³⁸ ⁸⁰²³⁹ ⁸⁰²⁴⁰ ⁸⁰²⁴¹ ⁸⁰²⁴² ⁸⁰²⁴³ ⁸⁰²⁴⁴ ⁸⁰²⁴⁵ ⁸⁰²⁴⁶ ⁸⁰²⁴⁷ ⁸⁰²⁴⁸ ⁸⁰²⁴⁹ ⁸⁰²⁵⁰ ⁸⁰²⁵¹ ⁸⁰²⁵² ⁸⁰²⁵³ ⁸⁰²⁵⁴ ⁸⁰²⁵⁵ ⁸⁰²⁵⁶ ⁸⁰²⁵⁷ ⁸⁰²⁵⁸ ⁸⁰²⁵⁹ ⁸⁰²⁶⁰ ⁸⁰²⁶¹ ⁸⁰²⁶² ⁸⁰²⁶³ ⁸⁰²⁶⁴ ⁸⁰²⁶⁵ ⁸⁰²⁶⁶ ⁸⁰²⁶⁷ ⁸⁰²⁶⁸ ⁸⁰²⁶⁹ ⁸⁰²⁷⁰ ⁸⁰²⁷¹ ⁸⁰²⁷² ⁸⁰²⁷³ ⁸⁰²⁷⁴ ⁸⁰²⁷⁵ ⁸⁰²⁷⁶ ⁸⁰²⁷⁷ ⁸⁰²⁷⁸ ⁸⁰²⁷⁹ ⁸⁰²⁸⁰ ⁸⁰²⁸¹ ⁸⁰²⁸² ⁸⁰²⁸³ ⁸⁰²⁸⁴ ⁸⁰²⁸⁵ ⁸⁰²⁸⁶ ⁸⁰²⁸⁷ ⁸⁰²⁸⁸ ⁸⁰²⁸⁹ ⁸⁰²⁹⁰ ⁸⁰²⁹¹ ⁸⁰²⁹² ⁸⁰²⁹³ ⁸⁰²⁹⁴ ⁸⁰²⁹⁵ ⁸⁰²⁹⁶ ⁸⁰²⁹⁷ ⁸⁰²⁹⁸ ⁸⁰²⁹⁹ ⁸⁰³⁰⁰ ⁸⁰³⁰¹ ⁸⁰³⁰² ⁸⁰³⁰³ ⁸⁰³⁰⁴ ⁸⁰³⁰⁵ ⁸⁰³⁰⁶ ⁸⁰³⁰⁷ ⁸⁰³⁰⁸ ⁸⁰³⁰⁹ ⁸⁰³¹⁰ ⁸⁰³¹¹ ⁸⁰³¹² ⁸⁰³¹³ ⁸⁰³¹⁴ ⁸⁰³¹⁵ ⁸⁰³¹⁶ ⁸⁰³¹⁷ ⁸⁰³¹⁸ ⁸⁰³¹⁹ ⁸⁰³²⁰ ⁸⁰³²¹ ⁸⁰³²² ⁸⁰³²³ ⁸⁰³²⁴ ⁸⁰³²⁵ ⁸⁰³²⁶ ⁸⁰³²⁷ ⁸⁰³²⁸ ⁸⁰³²⁹ ⁸⁰³³⁰ ⁸⁰³³¹ ⁸⁰³³² ⁸⁰³³³ ⁸⁰³³⁴ ⁸⁰³³⁵ ⁸⁰³³⁶ ⁸⁰³³⁷ ⁸⁰³³⁸ ⁸⁰³³⁹ ⁸⁰³⁴⁰ ⁸⁰³⁴¹ ⁸⁰³⁴² ⁸⁰³⁴³ ⁸⁰³⁴⁴ ⁸⁰³⁴⁵ ⁸⁰³⁴⁶ ⁸⁰³⁴⁷ ⁸⁰³⁴⁸ ⁸⁰³⁴⁹ ⁸⁰³⁵⁰ ⁸⁰³⁵¹ ⁸⁰³⁵² ⁸⁰³⁵³ ⁸⁰³⁵⁴ ⁸⁰³⁵⁵ ⁸⁰³⁵⁶ ⁸⁰³⁵⁷ ⁸⁰³⁵⁸ ⁸⁰³⁵⁹ ⁸⁰³⁶⁰ ⁸⁰³⁶¹ ⁸⁰³⁶² ⁸⁰³⁶³ ⁸⁰³⁶⁴ ⁸⁰³⁶⁵ ⁸⁰³⁶⁶ ⁸⁰³⁶⁷ ⁸⁰³⁶⁸ ⁸⁰³⁶⁹ ⁸⁰³⁷⁰ ⁸⁰³⁷¹ ⁸⁰³⁷² ⁸⁰³⁷³ ⁸⁰³⁷⁴ ⁸⁰³⁷⁵ ⁸⁰³⁷⁶ ⁸⁰³⁷⁷ ⁸⁰³⁷⁸ ⁸⁰³⁷⁹ ⁸⁰³⁸⁰ ⁸⁰³⁸¹ ⁸⁰³⁸² ⁸⁰³⁸³ ⁸⁰³⁸⁴ ⁸⁰³⁸⁵ ⁸⁰³⁸⁶ ⁸⁰³⁸⁷ ⁸⁰³⁸⁸ ⁸⁰³⁸⁹ ⁸⁰³⁹⁰ ⁸⁰³⁹¹ ⁸⁰³⁹² ⁸⁰³⁹³ ⁸⁰³⁹⁴ ⁸⁰³⁹⁵ ⁸⁰³⁹⁶ ⁸⁰³⁹⁷ ⁸⁰³⁹⁸ ⁸⁰³⁹⁹ ⁸⁰⁴⁰⁰ ⁸⁰⁴⁰¹ ⁸⁰⁴⁰² ⁸⁰⁴⁰³ ⁸⁰⁴⁰⁴ ⁸⁰⁴⁰⁵ ⁸⁰⁴⁰⁶ ⁸⁰⁴⁰⁷ ⁸⁰⁴⁰⁸ ⁸⁰⁴⁰⁹ ⁸⁰⁴¹⁰ ⁸⁰⁴¹¹ ⁸⁰⁴¹² ⁸⁰⁴¹³ ⁸⁰⁴¹⁴ ⁸⁰⁴¹⁵ ⁸⁰⁴¹⁶ ⁸⁰⁴¹⁷ ⁸⁰⁴¹⁸ ⁸⁰⁴¹⁹ ⁸⁰⁴²⁰ ⁸⁰⁴²¹ ⁸⁰⁴²² ⁸⁰⁴²³ ⁸⁰⁴²⁴ ⁸⁰⁴²⁵ ⁸⁰⁴²⁶ ⁸⁰⁴²⁷ ⁸⁰⁴²⁸ ⁸⁰⁴²⁹ ⁸⁰⁴³⁰ ⁸⁰⁴³¹ ⁸⁰⁴³² ⁸⁰⁴³³ ⁸⁰⁴³⁴ ⁸⁰⁴³⁵ ⁸⁰⁴³⁶ ⁸⁰⁴³⁷ ⁸⁰⁴³⁸ ⁸⁰⁴³⁹ ⁸⁰⁴⁴⁰ ⁸⁰⁴⁴¹ ⁸⁰⁴⁴² ⁸⁰⁴⁴³ ⁸⁰⁴⁴⁴ ⁸⁰⁴⁴⁵ ⁸⁰⁴⁴⁶ ⁸⁰⁴⁴⁷ ⁸⁰⁴⁴⁸ ⁸⁰⁴⁴⁹ ⁸⁰⁴⁵⁰ ⁸⁰⁴⁵¹ ⁸⁰⁴⁵² ⁸⁰⁴⁵³ ⁸⁰⁴⁵⁴ ⁸⁰⁴⁵⁵ ⁸⁰⁴⁵⁶ ⁸⁰⁴⁵⁷ ⁸⁰⁴⁵⁸ ⁸⁰⁴⁵⁹ ⁸⁰⁴⁶⁰ ⁸⁰⁴⁶¹ ⁸⁰⁴⁶² ⁸⁰⁴⁶³ ⁸⁰⁴⁶⁴ ⁸⁰⁴⁶⁵ ⁸⁰⁴⁶⁶ ⁸⁰⁴⁶⁷ ⁸⁰⁴⁶⁸ ⁸⁰⁴⁶⁹ ⁸⁰⁴⁷⁰ ⁸⁰⁴⁷¹ ⁸⁰⁴⁷² ⁸⁰⁴⁷³ ⁸⁰⁴⁷⁴ ⁸⁰⁴⁷⁵ ⁸⁰⁴⁷⁶ ⁸⁰⁴⁷⁷ ⁸⁰⁴⁷⁸ ⁸⁰⁴⁷⁹ ⁸⁰⁴⁸⁰ ⁸⁰⁴⁸¹ ⁸⁰⁴⁸² ⁸⁰⁴⁸³ ⁸⁰⁴⁸⁴ ⁸⁰⁴⁸⁵ ⁸⁰⁴⁸⁶ ⁸⁰⁴⁸⁷ ⁸⁰⁴⁸⁸ ⁸⁰⁴⁸⁹ ⁸⁰⁴⁹⁰ ⁸⁰⁴⁹¹ ⁸⁰⁴⁹² ⁸⁰⁴⁹³ ⁸⁰⁴⁹⁴ ⁸⁰⁴⁹⁵ ⁸⁰⁴⁹⁶ ⁸⁰⁴⁹⁷ ⁸⁰⁴⁹⁸ ⁸⁰⁴⁹⁹ ⁸⁰⁵⁰⁰ ⁸⁰⁵⁰¹ ⁸⁰⁵⁰² ⁸⁰⁵⁰³ ⁸⁰⁵⁰⁴ ⁸⁰⁵⁰⁵ ⁸⁰⁵⁰⁶ ⁸⁰⁵⁰⁷ ⁸⁰⁵⁰⁸ ⁸⁰⁵⁰⁹ ⁸⁰⁵¹⁰ ⁸⁰⁵¹¹ ⁸⁰⁵¹² ⁸⁰⁵¹³ ⁸⁰⁵¹⁴ ⁸⁰⁵¹⁵ ⁸⁰⁵¹⁶ ⁸⁰⁵¹⁷ ⁸⁰⁵¹⁸ ⁸⁰⁵¹⁹ ⁸⁰⁵²⁰ ⁸⁰⁵²¹ ⁸⁰⁵²²		

15899 floribundus <i>B. R.</i>	bundle-flowered	O or	L au	Y	N. Amer. 1826.	S co	Bot. reg. 1125
15900 likteus <i>L.</i>	yeUovrJowered^	^ pr	*jn.s	Y	Chile 1826.	D p.l	Bot reg. 1030
2rivuliiris	rivulet	J ^ pr	f jn.s	Y	Chile 1826.	D p.l	
15901 moschatus <i>Dou.</i>	musk	IT A pr	t jl.s	Y	Colombia 1826.	1> p.l	Bot reg. 1118
15902 lanatus <i>Ph.</i>	woolly	f ^ I p r	I *jn.jl	Y	N. Amer. 1826.	S p.l	
, 1784 HORNEMA'NN/^ W.en. HORNEMANIA. (<i>Prof. Hornemann, Copenhagen.</i>) <i>Scrophulfnrup. I.—2.</i>							
15903 bicolor <i>W. en.</i>	two-colored	O p r \	jn.s	B	E. Indies 1816	8 61	Grafiola goodcnUefblia Horn.
W85. TITTMIA'NN/4 Rchb. TITTMANIA. (<i>TUtnann, aman's name.</i>) <i>ScrophuUfrinar. 2.-8.</i>							
15904 ovata Rchb.	ovdt-leaved	JE O pr i	jn.jl	Pa. R	SocietyIsL 1816.	S s.p	Plan. h. b.9
Hornemanniia ovkta L.c.							
15905 viscosa Rchb.	clammy	- * O p r i	jnjl	...	E. Indies 182a	S s.p	Hornemanniia visciosa JV.
1786. MA*ZUS Lou. MAZUS. (<i>Mazes, a teat; tubercles closing mouth of corolla.</i>) <i>Scrophuldr'tm*.</i>							
15906 rugatu Lou.	wrinkly	-* .Q] pr }	my.s	Y.P	China 1780.	S s.l	Sw.fl.gar.36
15907 Limlernia japonica Thun.							
15907 pumilio R. Br.	dwarf	O pr *	jnjl	Pa.P	V. D. L. 1823.	S &p	
TORENIA. {Rev. Olgf Toren, a Swedish botanist.) <i>ScrophiUdr'irue. 2.-6.</i>							
15909 Asiatic diffuse	JUQSUN	JU ES] un	1 ji.au	P	E. Indies 1823.	S s.l	R. mal.9.53
15909 diffusa Ro'x.			1 jl.au	P	Nepal 1818.	S 81	
1788. ISOPLEXIS Lindl. IaoPLEXis. (JTOS, equal, plexis, H:jent; segments of cor. equal.) <i>Scropfiul. 2.</i>							
cinancensis Lindl.	Canary	* i_ or 4	jl.jl	Tr.O	Canaries 1698.	S p.l	Lind. dig. 27
Digitellis canariensis L.							
xcentrum Lindl.	sceptre	tt.	i_ or 2	jl.au	Br.O Madeira 1777.	S p.l	Lind. dig. 28
Digitalis sceptrum L.							
* 1789. DIGITALIS L. FOXGLOVE. (<i>Digitale, finger of a glove; form of flower.</i>) <i>Scrophular'ttue. 24.—2a</i>							
I. TRANSVERSJS.—Upper lip of ventricose corolla transverse.							
15912 purpurea L.	purple	4 jn.s	P	Britain hed.b.	S co	Lind. dig. 2	
22alba	white	4 jl	W	Britain hcd.b.	S co		
Th&psi L.	Thansi	14 my.au	P	Sjain 1732.	S co	lind. dig. 3	
MM lucescens W. & K.	woolly	3 jnjl	R	Portugal 1818.	S co		
XianiWguia Mur.	fuscous	2 jnjl	Hungary 1823.	S co			
ochroleuca Jac.	ambiguous	3 jl.au	L.Y	Switzerl. 1596.	S co	Lind. dig. 7.	
ava Lindl.	yellowish whitest	A or	4 jl.au	L.Y	Europe	S co	Lind. dig. 8
	tawny	^ A or	3 il.au	Br.	S co	Lind. dig. 9
II. PORRE'CUB.—Middle lobe of lower Up of corolla drawn out.							
smooth-leaved	'2 jl.au	Y	Hungary 1816.	S co	Lind. dig. 10		
iron-colored	4 jl.au	Br	Italy 1897.	S co	Lind. dig. 12		
woolly-spiked	3 jl.au	Br.Y	Russia 1827.	S co			
golden	3 jl.au	Br	Greece 1788.	S co	Lind. dig. 13		
grey	2 jl.o	W.Br	Greece 1789.	S co	Lind. dig. 14		
woolly	2 jnjl	W.Br	Hungary 1890.	S co	Lind. dig. 15		
orientalis Lam	eastern	ljjnjl	W	Levant 18LU	S co	Lind. dig. 16	
III. BIDENTAVJE.—Loiver lip of corolla bifid.							
smaller	i	Jnjl	P	Spain 17S9.	S s.l	Lind. dig. 6	
small-flowered	A or	1 jn.au	Br.	S co	Lind. dig. 17	
ismall-flowered	A or	2 jn.au	Y.Br	Switzerl. 1S17.	S co		
rigid	A or	1 jn.au	Y.a	S co	Lind. dig. 19	
purpurascens	A or	2 jn.au	Pk	Germany 1776.	S co	Lind. dig. 20	
tube-flowered	A or	2 jn.au	Y	S co	Lind. dig. 22	
small-yellow	A or	2 jl.au	L.Y	France 1629.	S co	Lind. dig. 23	
parviflora	lutescent	A, or	2 jl.au	Y	S co	Lind. dig. 21
15935 media Roth	obscure	A, or	1 jl.au	O	Spain 1778.	C pi	Lind. dig. 26
intermedia Pert.	intermediate	A or	2 jn.au	Y	Germany 1817.	S co	
* 1790. SCROPHYLAMIA L. FIG WORT. (Tuberous on roots like scrophulous tumours.) <i>Scroph. 45.-53.</i>							
1503R laminiifl. h~ Leaves simple, undivided, from No. 1394, lance-/Wd	lance-/Wd	i A un	3 jl au	G.Y Carolina 18LU	1) co		
lanceolata P.							
cretaceous	^ A un	2 jl	Br	Silicia 1S27.	S eo		
Uru.	Balm-leaved	2 jl	Br.P	Levant 18>3.	S co		
oriental	2 jl	Br	Levant 1710.	1) co	Schk.han.2 173		
rock shrubby	2 jl	P	Caucasus 1818.	1) co			
branch icst	2 jl	P	France 1821.	D co			
fruticosa Dec.							
15942 frutescens L.	frutescent		jn.au	D.P	Portugal 1768.	D co	Her. lug. 547
15943 adscondens IV. en.				Br 1816.	D co	
15914 marilandica L.	ascending			C.Br	N. Amer. 17VJ.	D l.p	
ISJ'i perigrina L.	Maryland		myjl	Italy 1640.	S co	Cam. h. 43	
15946 nimbosa L.	foreign		2 jn.au	Bil	Britain woods 1740.	D co	Eng. bot 1544
15H47 aquatica L.	knotty-rooted		2 myjl	Urtaian wat.pl.	1> co	Eng. bot. &H	
35D48 glabra // K.	water		2 myjl	Cnnries 1779.	S l.p	Jac.MC.2.209	
15949 Smithi* Horn.	emunth./eaeuii		np.my	P Canaries 1815.	D s.		
15950 biserrata W. en.	Smith's		ap.au	Br 1816.	I) co		
15951 arguta H. K.			2 ju.au	Br			
15952 vernalis L.	doubly-toothed		U mj,jn	Canaries 1778.	S co		
159.33 glandulosa Kit.	yellow spring		2 nif.iny	Britain m.sh.pl.	S co	Eng. bot 567	
15954 rugosa IV.	glandular		2 jn.s	Kr.P Hungary 1S06.	D co	W.&K.2H	
15955 Scopoli's Hoppe	rough		2 jn.s	Br.P Hungary 1816.	S co		
(S ? Scorodonia L.	Scopoli's		2 jn.au	Br Austria 1823.	D co	Scop. car. 32	
grandidentata Ten.	large-toothed		3 jl.au	P Britain wat.pl.	D co	Eng. bot. 2209	
altaica Mur.	Altaian		1 myjn	Pa Y Siberia 1786!	C co	Mur. 4.2	
Itonicifolia L.	Betony-leaved		jn.au	P Spain 1752.	D co	Bar. ic. 274	

15960 <i>hirsuta</i> Horn.	hairy	2	jl.au	Y	Europe	1816.	D	co			
15961 <i>appendiculata</i> Jac.	4	3	jl	D.P	Morocco	1805.	D	co	Jac. sc. 3.286		
15962 <i>Balbis</i> /j Horn.	■	2	jn.au	Y.o	Italy	1817.	D	co			
15963 <i>auricula</i> JL	Ibis'a	2	jl.au	Br	Spain	1772.	D	co	Lob. ic. 533		
15964 <i>trifoliata</i> L	en-leaved	2	in.y.s	R.v	Africa	1731.	S	co	Pluk. al. 313.6		
15965 <i>laciniata</i> Kit.	three-leaved	2	in.y.s	R.v	Africa	1731.	S	co			
15966 <i>verbenaefolia</i> Poir.	II.	—	—	Leaves cut or jagged.							
15967 <i>pinnatifida</i> Bro.	fag-leaved	3	jl	Br.P	Hungary	1806.	D	co	W.&K.2.170		
15968 <i>pinnatifida</i> Bieb.	Vervain-leaved	2	l	Br.P.	1816.	S	co			
15969 <i>ehuWfUa</i> Bieb.	pinnatifid	3	Lau	P.w	Span	1819.	S	co			
15970 <i>grandiflora</i> Dec.	variegated	2	n.au	Br.P	Casp. sho	1816.	D	co			
	Dwarf Elder-lvd	3	lau	Y.P	Caucasus	1819.	S	co			
	great-flowered	3	n.au	Y.p	S. Amer.	1820.	S	co			
15971 <i>lyrata</i> W. en.	lyrate-teawerf	3	jl.au	Br.P	Portugal	1816.	D	co	W.h.b.1.55		
15972 <i>mellifera</i> H. K.	honey-bearing	3	jl.au	Br.P	Barbary	1786.	D	co	Desf.at.2.143		
15973 <i>sambucifolia</i> L.	Elder-leaved	3	jl.8	R.u	Spain	1640.	D	co	Mil. ic. 2.231		
15974 <i>hispida</i> Desf.	hispid	2	il.au	Y.p	S. Europe	1824.	S	co			
15975 <i>tanacetifolia</i> W. en.	Tansy-leaved	3	jl.8	Br.P	Tauria	1804.	D	co	W. h. b. 1.56		
15976 canlna L.	dog	3	jn.au	Br.P	S. Europe	1683.	D	co	Lob. ic. 2.55		
15977 chrytinthemif&lia W. en.	Chrys.-lvd	3	jn.au	Br.P	Tauria	1816.	D	co	W. h. b. 1.59		
15978 ILICKU L.	ehming-leaved	2	in.au	Br.P	Levant	1596.	D	co	W. h. b. 57		
15979 ilicilolia Mil.	Fein-leaved	3	jt.au	P.w	Crete	1700.	S	co	Tou. it. 85		
15980 multifida IV. en.	many-clefWif	3	jn.au	Br.P	1816.	D	co	W.h. b.1.58		
1791. V ANDE'LLIii L. VANDEIXIA. {Dr. Louis VandeUi, prof. bot Coimbra, Portugal.) <i>Scrophul.</i> 1. — 2.											
15981 diffusa L.	diffuse	O cu	1	jl.au	W	S. Cruz	1824.	S	81	Marcg. br. 32.1	
1792. SI BTHO'RPI A L. SIRTHORFIA. {H. Sibthorp, M.D., prof. bot. at Oxford.) <i>Scrophuldrinhar.</i> 1.—2.											
15982 europa^a L.	European	&	A cu	ft	jl.au	y	England	w.sh.p.	D s.l	Eng. bot. 619	
1793. LTMOSE'LLA L. MUDWORT. (Limns, mud; habitation.) <i>Scrophularxne.</i> I.— 5.											
15983 aquatica L.	water	▲	O pr	4	jl.s	F	Britain	mud.pl.	S s.l	Eng. bot 357	
* 1794. BROWALLIJA L. BROWALLIA. (John Browallius, bishop of Abo.) <i>Scrophvlarhce.</i> 3.—4.											
15984 demissa L.	low	UQJor	f	jns	B	S. Amer.	1735.	S	si	Bot mag. 1136	
15985 efta L.	tall	iQJor	lj.jns	B	Peru	1768.	S	si	Bot mag. 34		
15986 elong^ta Kth.	elongated	iQJor	U jns	B.w.	S	81			
lactea Hort.											
• 1795. STEMCTDIA L. STEMOMA. (Stemon^jtamen dis , double; \$^anthered filaments.) <i>Scrophul.</i> 6.—14.											
15987 suffruticosa Bonp.	suffruticose	ILCD	—	Il^jl.au	R	Brazili	1820.	C	p.l		
ColGmnea tnfofata Lk.											
15988 viscosa ROT.	clammy	OD un)	jl.au	B	R Indies	1818.	■	p.1		
15989 vrtirillaris Lk.	whorled	O un	J	jl.au	P	Brazil	1825.	■	p.1		
15990 maritima L.	seaside	KB un)	jl.au	R	S. Amer.	1817.	■	■	Jac. am. 174.66	
15991 duranti/6/ja Swz.	Duranta-leaved	QS un	1	jl.au	W	W.Indies	1823.	S	p.l	SI.jam. 1.124.2	
Capraria duranti/oia L.											
15992 parviflora R. Br.	small-flowered	O un	i	jl.au	W	S. Amer.	1759.	S	p.l		
1796. TREVIRANA^Ni* W.en. TREVIRANA. (Dr. Treviranus, a German botanist) <i>Scrophuldritue.</i> 1.											
15993 coccinea W.en.	scarlet	£ E3	spl	^ au.s	S	Jamaica	1778.	R	l.p	Bot mag. 374	
Cyrilla pulchella Herit.											
1797. COUJMKEA L. COLIJMNEA. (Fabius Columna, of noble family of Colonna, in Italy.) <i>Scroph.</i> 4.— &.											
15994 scandens L.	climbing	fl	• or	6	au.s	S	W. Indies	1759.	C	s.p	Bot. reg. 805
15995 hirsute Sun.	hairy	s.	• or	2	au.n	Pa.P	Jamaica	1780.	C	s.p	Br. jam. 30.3
15996 rutilans Swz.	red-leaved	tt.	CJ or	2	au.s	P	Jamaica	1823.	C	s.p	
15997 tri(oli?ita)Lk.	three-leaved	», Q or	3	au.n	B	182a	C	s.p		
1798. RUSSE'L/i* L.	RUSSELIA. (A. Russel> M.D. F. R.S., auth. of nat. hist. of Aleppo.) <i>Scrophu.</i> 3. --&										
15998 multiflora B. M.	many-flowered	A Q or	4	jn.au	R	S. Amer.	1812.	C	s.p	Bot mag. 1528	
15999 floribunda Hum.	bundle-flwd	i Q or	4	...	R	Mexico	1824.	C	81		
16000 ternifolia Kth.	ternate-leaved	HO or	4	...	R	Mexico	1818.	C	81		
1799. DOD A'RT/4 L. DODARTIA. {F. Dodart, M.D. French bot. and acad.) <i>Scrophuldritue.</i> 1. — ^											
16001 onentahs L.	oriental	^ A or	^ jl.au	P	Levant	1752.	D	s.p	Sw. fl. gar. 147		
1800. LINDE'RN/il L. LINDEKNIA. {F.B. von Lindern, M.D of Strashburg,bot.author.) <i>Scrophul.</i> !...^											
16002 Pyxidaria L.	Pyxidana	O pr	1	jn.au	B	S. Europe	1789.	S	s.l	Lam.il. 522	
1801. HERPE'STIS R. Br. HBRPEOTIS. (Herpestes, any thing that creeps; habit.) <i>ScrophuldryMB.</i> 8. — ^											
16003 stricta Schr.	strict	£ (29 pr	1	au	B	1824.	D	bog		
16004 Monnier'a R. Br.	Monnier's	■ pr	jl.s	■.B	S. Amer.	177*	D	bog	Ehr. pic. 1* *		
16005 portulacacea B. M.	portulacaea	■	jl.au	B	S Amer.	1823.	D	bog	Bot. mag.		
16006 cuncifilia Ph.	wedge-leaved	i A pr	■	jl.au	B	N. Amer.	1812.	D	bog		
16007 dinglexiculis Ph.	stem-clasping	^ & un	\	jl.au	B	Carolina	1818.	D	bog		
16008 rotundifolia Ph.	round-leaved	A ^ un	\	jl.au	Pa.B	N. Amer.	1824.	D	bog		
16009 micrantha Ph.	small-flowered	■ AJ un	\	jlau	B	N. Amer.	1817.	D	bog	Swz. ic. 33	
16010 Brfwnii Pers.	Brown's	■ AJ un	i	jlau	B	Jamaica	...	D	bog		
1802. DIA'SCIA Lk. DIASCIA. {Dis, two, askion, little bladder; two protub. at base of cor.) <i>Scroph.</i> t*V											
16011 Bergiami Lk.	Bergius's	O un	i	jl.au	...	C. G. H.	1815.	S	61	Pluk. ph. 520-*	
1803. CAPRA^RIA L. CAPRARIA. ■, a goat: leaves liked by goats.) <i>Scrophuldritus.</i> 5. ^t X											
16012 biflbra L.	two-flowered	■ un	2	jf.au	W	S. Amer.	1752.	S	1.p	*	
16013 cuncta R. Br.	vrdge-leaved	■ un	2	...	W	S. Amer.	1759.	C	p.l	*	
16014 lanceolata JL	spear-leaved	■ un	3	...	W	C. G. H.	1774.	C	p.l		
16015 uudulata J..	wave-leaved	a. ■ un	2	mrjl	W	C. G. H.	1774.	C	p.l	Bot mag. 15 ⁵⁶	
16016 hiimihs W.	dwarf	■ un	1	jlau	.W.	E. Indies	1781	C	p.l		
■ Nut	COLLINSIA. {Zaccheus Collins, a vice-pres. of Ac. Nat. Sc. Philadcl.) ■										
■ verna JVX*	spring	O el	1	rnyjn	B p	N. Amer.	1826.	S	l.p.s	Sw.	
16018 parviflora B. R.	small-flowered	J c o o r	i	jn.jl	P.n	Colombia	1826.	S	l.p.s	Bot. reg.	
16019 grandiflura T.A.	great-flowered	O el	1	m y j l	Pk.B	Colombia	182fi.	S	l.p.i	Bot. reg.	

CLASS XV. TETRADYNAZIA.

1816. SCHIZOPETALON Sims SCHIZOPBTALON. (*Schho*, to cut, *petition*, a petal.) *Cruciferae*. 1.
16077 Wfilkeri Sims Walker's .QJeu 2 my.jl W Chile 1822. S p.1 Bot mag. 2379

1817. MATHROL;* R. Br. STOCK. (P. A. Matthioli, an Italian physician, d. 1577.) *Cruciferae*. 17.—28.
I PACIYNOVITM.—Prfrt* obovate; back iff stigma thick, not horned; silicle at top thick, not tricuspidate.

16078 incaria ML Br.	hoary <i>Gillyflower</i>	<u>tl</u> or	1 my.ii	P ₁	England	clifls.	S l.p	Eng. bot. 1935
2 coccinea	scarlet	<u>tl</u> or	1 my.n	S ₁	England	gard.	S l.p	
3 &iii	white branching*	J or	1 my.n	W	England	gard.	S l.p	
4 multiplex	double	<u>tl</u> or	1 my.n	Va	England	gard.	S l.p	
16079 annua Swt.	annual	O or	2 my.n	R	S. Europe	1731.	S l.p	Bau. his. 2.875.1
2 alta	white <i>Ten-week stock</i>	Q or	2 my.au	W	S. Europe	■	S l.p	Dalech. 802.1
3 flôre pleno	double	Q or	1 my.au	R	■	■	S l.p	
16Q80 siimpliciuulis Swt.	simple-staid	£ CD or	2 my.au	P	■	■	S l.p	Bes.eys.2.3.3
2 allia	white	£ Q or	2 my.au	W	■	■	S l.p	Bes. eys. 2.3.*
16081 glbrifci Dec.	smooth	tt. or	2 my.n	W	■	■	S l.p	M.h.3.8.8
2 purpurca	purple	tt. or	2 my. n	P	■	■	S l.p	
S flore pleno	double	tt. or	2 my.n	P.W	■	■	S l.p	
16082 grs\:a Swt.	Grecian <i>Wallflower</i> -lyd	Q or	2 my.n	W	S. Europe	...	S l.p	
IGISO tencstrMis R. Br.	window	£ O or	1 jl.au	P	...	1709.	S l.p	Jac. vin. £179
16084 siniiita R. Br.	Bco'ot.leaved	£ Q or	1 iny.au	V	England	sea sh.	S l.p	Eng. bot 462
laws acads Dec.	stemless	O or i	1 jn.jl	R	Egypt	1818.	S sp	

II. LUPE^SRIA.—Petals oblong, of a yellowish purple; back of stigma thick, not horned.

1G086 torulosa Dec.	twisted	"tt-Ljor	2 m'yll	P	C.G. fl.	1810.	C s.p	
1G'87 odoralissima R. Br.	swcrcst-setd	tt. J or	2 jn.jl	Ld	Persia	1797.	G r.m	Bot. mag. 17H
2 frigrans Fis.	fragrant	tL J or	2 jn.jl	Ld	Crimea	182.	C r.m	
16088 varia Dec.	various	tf AJ or	J jn.jl	Lfl	Levant	1821.	C r.m	Fl. gr. 636
IQIS'jtataricaiJrc.	Tartarian	£ Q or	1 jn.jl	RY	Tartary	1831.	S s.p	Pal. it. 1. ap. O
16U90 tristis R. Br.	d&rk-Jtuwered	tt.L J or	lj my.jl	Ld	S. Europe	1788.	C s.l	Bot. mag. 729

III. PittVaiA.—Petals yellowish purple; back of stigma horned.

10091 longiptala Dec.	long.petalcld	O or	1 ui.jl	R.Y	Bagdad	1818.	S s.p	Vcn. eels 93
16092 coronopifttha Dec.	Buckhoru-lyd	j£ Q or	1 jn.jl	P	Greece	1819.	S s.p	Fl. gr. 637

IV. ACINOVON.—Petals obovate; stigmas horned; silicles at top tricuspidate.

16093 tricuRipidata R. Br.	three-pointedWud	O pr	f jlau	P	Barbary	1739.	S el	Lam.il. 564.9
16094 parviflora R. Br.	small-flowered	O br	jjl.au	P	Morocco	1799.	S s.l	

181R CHEIRA'NTHUSZ. WALLFLOWER. (*Kheyry*, its Arabic name, *anthos*, flower.) *Crwifer**. 11.—li^L

I. CHEIR^XRI.—Styles scarcely any; seeds imwarginafe.

16095 Che)ri L.	Cheiri, or garden	tt. J or	2 ap. I	Φ	S. Europe	1573.	S r.m	Bui. her. 349
2 flôre pl&no	double-flwd	tt. or	2 ap.jl	Y	■	■	C r.m	Bes.eys.2.5.3
3 grandiflorus	large, flowered	tt. J or	2 ap.jl	Y	S. Europe	1573.	C co	Bes. eys. 2.5.*
4 mäximus	largest	tt. or	2 ap.jl	Y	S. Europe	1573.	C co	Bes. eys. 2.5.1
5 serratus	saw-flowered	tt. or	2 ap.jl	Y	S. Europe	1773.	C co	
6 p&tulus	(flffVe-spreadingt)	tt. J or	I	Se	S. Europe	1573.	C co	
7 ferrugineus	rwhty-flowered	tt. J or	I	Se	Br	S. Europe	1573.	C co
8 vbris	various	tt. J or	2 ap.jl	Va	S. Europe	17#.	C co	
9 gavescens	flavescent	tt. J or	2 ap.jl	Y	a Europe	1573.	C co	
Id thynoldes	tbyr>e.flowered	tt. J or	2 ap.jl	Bd	S. Europe	1573.	C co	
11 sanguincus	bloody	tt. or	2 ap.jl	D.Br	■	■	C r.m	
12 hsmänthus	double bloody.flwdtt.	tt. or	2 ap.jl	Bd	S. Europe	1573.	C co	
16096 fruticulhus L.	small shrubby	tt. i or	■	Y	Britain old wa.	S co	Eng. hot. 1934	
16097 arliorus Swt.	tree	sh □ or	■ ap.jl	Y	Egypt	1827.	C ■	
16098 alpinus L.	alpine	tt.i lor	Ainy.jl	Y	N.Europe	1810.	C sip	Wa. lap. lfcl
16099 firmus IV.	firm	tt. or	1 my.jl	Y	Europe	1816.	C s.p	
16100 ochrolencus Hal.	pale yellow	£ A or	£ ap.jl	Pa.Y	SwiUerl.	1820.	D co	HaLh. 449.1*
diibius Suter								

II. CHEIRO^VDES—Styles filiform; seeds marginate; silicles four-sided.

16101 tenuiflilius Herit.	fine-leaved	tt. \ J or	2 niyjn \	Madeira	1777.	C l.p		
16102 mutabilis Merit.	changeable	tt. i J or	■ mr.my	Y.p	Madeira	1777.	■ l.p	Bot. mag. ■
U longifolius Fen.	long-leaved	tt. i J or	3 s.d	W.p	Madeira	1815.	■ co	Ven. mal. ■
16103 jcparius W.	broom	tt. j-j or	3 my.o	W.p	Teneriffe	1812.	L r.m	
2 aeruginosus Dec.	rusty	tt. ■ or	3 my.o	Rust	Tenenffe	1812.	L r.m	
3 chama'Jeo Ker	chameleon	■ J or	3 my.o	Y.p	Tenenffe	1812.	C r.m	Bot reg. 219
16104 semixrlsrens Schous.	ever-blowing	tt. L J or	■ year	W	Barbary	1815.	C s.l	
2 frutscscnt Pcrs.	frutscscnt	tt. ■ or	■ mr.jl	W	Teneriffe	1815.	C co	
16105 riniflthus Pers.	Flax-leaved	tt. i J or	2 mr.au	P	Spain	1815.	C s.l	

*1819. tfASTU'RITIUM ML Br. NASTURTIUM. (*Nasus*, nose, *tortus*, tormented; effects.) *Crucif.* ■

I. CARDAMINUM.—Petals white, larger than calyx; silicles cylindrical; glands four at base of stamen*.

16106 officialeA.ér.	official	^ A cul	1 my.jl	W	Britain rivul.	D co	Eng. bot S&	
----------------------	----------	---------	---------	---	----------------	------	-------------	--



Sis[^]mbrum astiurtium L.

11. BRACHY'OBOS.—Petals yellow; silicles cylindrical; glands of receptacle small. ^{ot*} |

16107 sylvestre R. Br.	wood creeping	£ A w	1 jn.s	Y	Britain watpl.	D co	Eng. bot»	
Sisyrnbrium sylvestre L.								
16108 palnstre Dec.	marsh	^ O ^	1 J. ^l .	■	Britain watpl.	S co	Eng. bot I? ⁴⁷	
Sisfimbrium terrestre L.								
16109 bursiflum Dec.	Sheph. Purse-lvd	O un	i ■ ■	Crea	Kamtsch.	1820.	S co	
16110 coronopiflum Dec.	Buckhorn-lvd	O un	* jnjl	Pa.Y	N. Africa	J824.	S co	Dcsf. ap. ⁴⁸
Jfjl sagittatum A. r.	arrow-leaved	jg A un	■ myjn	Pa.Y	Siberia	1780.	D co	Jac. ic. *• * ⁴⁹
JJJf PP«ense Dec.	Lippa	j£ A un	■ myjn	Y	Carinthia	1775.	D co	
JJJf Pyrenicum R. Br.	Pyrenean	* A un	■ myjn	Y	Pyrenees	1775.	D co	
16114 amphibium R. Br.	amphibious	* A w	■ myjn	Y	Britain riv. ba.	D co		
1 nd jyjf»n Dec.	undivided-vrf	A & w	■ jn.jl	Y	Britain riv. ba.	D co		
2 variifolium Dec.	various-leaved	§b A w	1 jn.jl	Y	Britain riv. ba.	D co		

i<nr w ¹¹¹ - CLANDESTINA^RIA.	— Petals none or very small and white ; silkies cylindrical.
i5!5 ^b nghalgnse Dec.	Bengal O un 2jn.au W E. Indies ... S co
*jJg microsjwrmum 2tec. small-seeded	O un # jn.au W China 1820. S co
in 2)?) Indian	O un i jn.au Ap China — S co
ic? o ^o ; L EPTOCARP^A Dec.	LEPTOCARREA. (<i>Leptos, slender, karpas, fruit</i>) Crucifer. 1.
16118 LoesHj Dec.	Loesel's un 11au Y Germany 1683. S co Jac. au. 4.324
TurrUis Loeselis R. Br. Sisymbrium Loesel* L.	
1821. NOTO'CER AS R. Br. NOTOCERA*. (<i>Notos, the back, keras, a horn; back of pod.</i>) Crudfene. a	
Iffim ¹ - DIC ER^TIUM.	— Silkies dehiscent, two-horned's seeds compressed; flowers small; leaves entire.
*°iiy canariense R. Br. Canary	O un 4 au.s Y Canaries 1779. S co Jac. ec. III
Dec. Spanish	O un au.8 Y Spain 1821. S co
2? T ^E J* CERANRium	Silkies four-horned : flowers small, perhaps apetalous; leaves sinuately toothed.
U quartricorne Dec.	four-horned O un i jnjl W Siberia 1820. S co Deless.2.16
<ar/Bimum (ua <i>ri</i> c ⁵ nie step	— Tdst(irtiumquadricorne Dec.
? A R# * R. Br. WINTER CRESS. (Anciently called Herb of St. Barbara.) Cruel/era?. 4.—6,	
Dbris. s. /.- common	* A cul 1* my.au Y Britain rub. D co Eng. bot. 443
» Barbara L.	
pecox JZA.- early JSWfeste Cr**jfe A cul 1 ap.o Y	England brooks. D co Eng. bot. 1129
16121 <ca7Jtr Taurian	£ A un 1 ap.o M TT Tauria 1830. D co
>150 iSuna 7>c. Iberian	tf A un 1 my.au* Y Iberia 1816. D Jp
1612R Zi "ihi <i>m</i> Ricus FK M.	Plantain-like ^ A un ft jl.s Y Levant 1799. D co /
16121 jantagnnea Dec.	fismbrim Barbara L.
ifi?? ³ STEVE ^x NJi* Adams STEVENIA. (C. Steven, a Russian botanist.) Cruel/era.	
ifi <i>s</i> N ² Wes Fts.	Alyssum-like O un i myjl W Siberia 1824. S co Deless.2.20
*oi*8 cheirantholdes Dec.	Wallflower-Ik £ CD un 1 myjl W.p. Siberia 1818. S co Deless.2.21
16129 S ² R ^A * Y ¹¹ Ster. BRAYA. (Count Bray, a German nobleman.) ^Cruafer* 1.—2.	
aliAiia Ster. alpine	£ A cu j jn P Carinthia 1823. S p.i Deless.2.22
ifiKS pA'RRY^ R.Br. PARRY. (Capt. Edw. Parry, R.N., sent to discover N.W. passage.) Crucifero. I ^z	
retica R. Br. arctic	JQ cu ... P Melville 1.1820. S p.i Parry's app.
16 ¹ ^6v TURRI ^V ns2W'. TOWER MUSTARD. (Turris, tower; pyramidal growth.) Crucifer. 2.-3.	
fi?!!*L smooth long-podded	O w 1 my.jn W England gr.pa. S co Eng. bot 777
16132 ^iMiginEsa Dec*	marsh O un 1 jnjl W Siberia 1823. S co Pal. it 2. app. O
1827. -rf'RABISZ. WALL CRESS. (Originally from Arabia.) Crudferte. 49.-67.	
I. ALOMA^TIUM. — Seeds immarginate ; stem leaves cordate, clasping, from No 16W. sessile.	
X ^r m A ^r . vernal	O pr 1 my.jn P France 1710. S s.l Bar. ic. 476
Mair. Clusius's	JU A pr % mr.my W.Y Switzerl. 1596. D p.l Bot. mag. 226
Uhiuna Dec. dwarf	JU A or I mr.my W Pyrenees 1596. D co Cl.h.2.125.2
albulas/ whitish	JU A or £ mr.my W Switzerl. 1819. D co
IT. CTI. waved-leave	JU A pr f ja.o W Caucasus 1798. D s.l Jac. ec. 71
wave	SU A or L mrjl W S. Europe 1810. D co
long-leaved	SU A or myjl W Persia 1820. D co
soft	JU A or * myjl W Caucasus 1817. D co Delew.2.25
bow-leaved	£ CD pr 1' jl.au W Volga 182a S co
auriculata auricled	O un f my W S. Europe ... S co Al. ped.2.2
Ma Bieb. ^{abgera ii/.}	
2 recta JW. upright	O un fm.y W Europe ... S co Vil. del. 3.37
3 p ^u tula Dec.	O un spreading fm.y W Hungary 1818. S co W.&K.1.59
TurrUis p ^u tula fArA.	
4 dasycarpa Andrz. thick-capsuled	O un S my W Podolia 1820. S co
J6141 saxifiths All. stone	JU CD un fmy W Switzerl. ... S co VU. del. 37
16U2 crisjuta W. curled	JU A pr 1 my W Carniola 1818. D co
16143 sugittita Dec.	sagittate £ CD un 1 my.jl W S. France ... S co
2 suhogiabrata Dec.	j^ CD un 1 my.jl W Europe 1820. S co
3 ovMa Poir.	jt CD un 1 my.jl W N. Amer. 1818. S co
4 oblongiita Dec.	jf CD un 1 my.jl W N. Amer. 1827. S co
shnrtpotl	£ A w 1 my.jl W N.Europe 1824. S co
spatulata	jf CD or * my.jl W Britain rocks. D s.l Eng. bot 587
16147 Allionii Dec.'	£ CD un * jnjl W Bonaria 1823. S co
I ^{MO} Tu ^u , ^u stricta J.	£ A un 2 my.jn W Italy 1804. D co
is85SSViS ^{fe}	wall £ A un J my.jn W Italy 1824. D co
1G150 ciliata A. ^	upright £ CD un my Crea England rocks. D s.l Eng. bot. 614
TurrUis alpVna W hoary-rfa/ftal	ciliated £ CD pr I P. Ireland ir.sh. S s.l Eng. bot. 1746
16151 fincana Roth	
16152 Thalia TM L.	
16155 procfirrensW.* _k pro?S??nt	
16156 pra'cox W. & J?	early y. A or t* A un f jnjl W Hungary 1818. D co
J0157 SchiwerrecktaDec.	Schiwerreck's rock JU A or * jnjl W Austria 1826. D co
16158 petraea Lam,	hispida SU A or * my.jl W Austria 1820. D co
2 hispida	Shastula hastulate JU A or 4 myjl W Scotland rocks. D co Cr. au. 3.2
	ambiguous £ CD or 4 myjl W Britain rocks. D co fl. dan. 386
	lyrate O un P. Siberia 1824. D co Eng. bot 469
	purple sand O pr P. N. Arær. ... P co
Haller's	£ CD un 1 jn.jl Pk Germany 1798. S l Hook. ex. fl. SSI
	W Switzerl. ... S co W. ft K. 190

16163 stolonifera <i>Horn.</i>	shoot-bearing	IU A or ft my,jl W	Carniola 1818.	D co	Scop. car. 39
16164 ovine'ngis <i>Wul.</i>	Ovirian	t. A or ft jn,jl Pa.R	Carinthia 1824.	D eo	Jac. i d . 125
16165 cebennensis <i>Dee.</i>	Cevennes	£ O un lft jn,jl Pa.P	S. France 1820.	S co	
16166 Iaugloba <i>Lk.</i>	hairy-podded	£ O or 1J my,jl W	Mexico 1820.	S co	
II. LOMA'SPORA. — Seeds marginate; stem leaves clasping, from No. 16171. sessile.					
16167 Turrua <i>L.</i>	Tower Mustard	Q> w lft ap.my Su	England walls. S s.i	Eng. bot 178	
2 umbrusa <i>Stev.</i>	shades	j£ Q un 1 ap.my Su	Russia 1827.	■ co	
3 plindula <i>Desf.</i>	pendulous	£ Q> un 1 ap.my Su	Europe 1817.	■ co	
19168 pendula <i>L.</i>	pendulous	Q un 1 my,jl W	Missouri 1759.	■ 8.1	Jac. vin. 3.34
16169 Patnnwia Dec.	Patnn's	A un 1 my,jl W	Siberia 1818.	■ co	Deless. 2.27
16170 levigktia Dec.	polished	A un llmy,jn W	N. Amer. 1821.	D co	
16171 canadensis <i>L.</i>	Canadian	A un 2 my,jl W	N. Amer. 1768.	D s.l	PIuk.al.86.8
16172 nutans <i>W.</i>	nodding	A P' ft mrap W	Switzerl. 1658.	D co	Jac. au. 3.281
16173 fellidifolia <i>Jac.</i>	Daisy-leaved	A P* i my,jn W.Y	Switzerl. 1773.	D p.l	Jac. au. 3.280
16174 pumila <i>Wul.</i>	dwarf	£ P ft my,jl W	Austria 1816.	D co	Cr. au. 1.281
Sellidifolia Crz. ciliaris <i>W. en.</i>					
16175 casfalea <i>Wul.</i>	blue	f. A P ft jn,jl Pa.B	Switieri. 1793.	D co	Al. ped. 1.40.2
16176 stellula <i>Bert</i>	little-starry	A or ft my, W	Italy 1817.	D co	
16177 collina <i>Ten.</i>	hill	■ A un ■ W	Naples 1824.	D co	
III. DU'BIJG. — Doubtful to which of the sections they belong.					
16178 fucida <i>L.</i>	shining-leaved	£ A pr ft jn,jl W	Hungary 1790.	D p.l	
16179 stenopetalia <i>W.</i>	narrow-petaled	£ CD or ft jn,jl W	1818.	S co	
16180 alpestris <i>Schl.</i>	alpine	g-Tft or ft my,jl W	Switzerl. 1819.	S co	
16181 dasycarpa <i>Andrx.</i>	thick-fruited	£ * £ ♀ r ijn,jl W	Podolia 1827.	D co	
* 1828. MACROPO*DIUM Jl. Br. MACROPODIUM. (<i>Makros</i> , long, <i>pous</i> , foot; of seed-vessel.) <i>Cmcifera</i> . 1*					
16182 nivkle R. Br.	snowy	£ A P* 1 jn,s W	Siberia 1796.	D co	Pal it. 2. ap. U
Cardaraine nivkis Pall.					
' 1829. CARDAMINE L. LADY'S SMOCK. (Dim. of <i>kardamon</i> , water cress, Gr.s taste.) <i>Cruciffer<e.25.—&-</i>					
I. INDIVPS'. — Leaves nearly all undivided.					
16183 rhomboidea Dec.	rhomboid	£ A un ft jn,jl W	N. Amer. 1824.	D p.l	Pluk. al. 435.6
'6184 rotundifolia Mx.	round-leaved	g A un i my,jn W	N. Amer. 1823.	■	
16185 asarifolia L.	Asarum-leaved	£ A P' i Jn,jl W	Italy 1710.	D p.l	Bot. mag. 1375
16186 Aellidifolia L.	Daisy-leaved	£ A Pr i &Pjnj W	Scotland seal,	D s.l	Eng. bot 2355.
2 alpina Dec.					
tellidifolia Crz.	alpine	jg A or i apjn W	Austria 1658.	D s.l	Jac. mus. 1.17.*
II. TRI'LOAE. — Leaves for the most part three-lobed.					
16187 resedifolia L.	Reseda-leaved	O un £ jl W	Germany 1658.	S co	Al. ped. 1.57.8
16188 africkna Z.	African	tf iAI un 1 my jn W	C. G. H. 1691.	D co	Her. par. 208
16189 trifidia L.	three-leaved	£ A pr li mr.ap W	Switzerl. 1629.	D p.l	Bot. mag. 452
16190 chilensis Dec.	Chile	tf A un i mr.ap W	Chile 1825.	D co	
III. PINNATIPARTIVE. — Leaves for the most part pinnated.					
16191 granulosa All.	granular	* A un 1 ap.my W	Italy 1820.	D ms"	
16192 anfara L.	bitter	■ A P* i ap.my W	Britain wat.pl. D ms Eng bot 100		
16193 uliginosa <i>Bieb.</i>	bog	■ A pr ft apji W	Tauria 1819.	D ms	
16194 proptipens <i>Fis.</i>	creeping	* A un ft apmy W	Siberia 1821.	I) ms	
16195 pratensis L.	meadow Cuckoo fl.^	A pr 1 ap.my P	me. pa. D ms Eng. bot 776		
3 plena	&owhe-Jtowered	i A P' 1 ap.my L.P	... D mb		
16196 dentata <i>Schult.</i>	toothed	■ A un ft apjn W	Russia 1820.	D ms	
16197 pennsylvanica <i>Mhl.</i>	Pennsylvanian	■ A un 1 my,jn W	N. Amer. 1818.	D ms	
16198 hirsuta L.	hairy	O un year W	Britain mo.s.p. S ms Eng. bot. 492		
16199 sylvatica <i>Lk.</i>	wood	O un	Europe ... S ms Fl. dan. 735		
16200 umbrota <i>Andrx.</i>	shaded	O un	Europe 1824. S ms		
16201 paryiflora L.	small-flowered	O un "apamy W	France ... S ms Gm. si. 64		
16202 impatiens L.	impatient	O un ap.jn W	Britain al. roc. S ms Eng. bot. 80		
16203 latifolia <i>Vahl</i>	broad-leaved	■ A or ■ jn,au P	Spain 1710. S ms Her. par. 203		
16204 macrophytia W.	long-leaved	■ A or ap.au P	Siberia 1824. D ms Gm. si. 3.62		
16205 Chelidonia L.	Celandine-/Mi	■ A jn,au P	Italy 1739. D ms W. & K. 2.140		
16206 /halictroides All.	Thlasictrum-like	O jn,jl W	Piedmont 1818. S ms Vil. del. 3.38		
Plumieri Vil.					
16207 glauca Spr.	glaucous	£ A or ft my,jl W	Calabria 1827. D m.8 Deless.2.31		
1830. PTERONEITRON Dec. PTERONEURON. (<i>Pteron</i> , wing, <i>neuron</i> , nerve; pods.) ^ T W ^ ^ I IM					
16208 carnosum Dec.	fleshy	A A or ft jn,jl W	Hungary 1824.	D p.l	W.&K.**"
Cardimine carnsa W.S/K.					
16209 gTJE cum Dec.	Grecian	O un ft jn,jl Pa	S. Europe 1710. S co	Boc. BIC.***	
Cardaminic graeca L.					
183L DENTARIA L. DENTARIA. (<i>Dens</i> , tooth; tooth-like structure of roots.) Crudferee. 1					
16210 polyph'tlla W.S/K.	many-leaved	A A el 1 my,jn P	Hungary 1818.	D s.p	W. & K.
16211 cmneaphyl'la L.	nine-leaved	A A el 1 my,jn Pa,Y	Austria 1656.	D s.p	Jac. au.
16212 glandulosa W. % K.	glandulous	A A or 1 my,jn L.P	Hungary 1815.	D s.p	
16213 laciniata Mhl.	jagged	A A or 1 my,jn W	N. Amer. 1822.	D s.p	Pluk.m
16214 diph'j'la Mx.	two-leaved	A A el f my,jn W.p	N. Amer. ... D s.p Bot. mag.		
16215 maxima Nut.	largest	A A el 2 my,jn Pa,P	N. Amer. 1823.	D s.p	
16216 trifolia W.S/K.	three-leaved	A A el 1 my,jn W	Hungary 1824.	D s.p	
16217 digitata Lam.	digitate	A A el lft my,jn Pa,P	Switzerl. 1656.	D s.p	
pentaph'llos L.					
16218 pinnata Lam.	pinnate	A A el 1 my,jn Pa,P	Switzerl. 1683.	D s.p	Gar. aix. %
16219 quinquefolia <i>Bieb.</i>	five-leaved	A A or 1 my,jn P	Tauria 1820.	D s.p	
16220 bulbifera L.	bulbiferous	A A el lft ap.my P	England sha.pl. D s.p		
16221 tenuifolia Led.	fine-leaved	A A or 1 my,jn L.P	Siberia 1825.	D s.p	Gm.
1832. NEUROLO ^X MA Dec. NEUROLOMA. (<i>Neuron</i> , a nerve, <i>loma</i> , a fringe.) Crucifera?.					
16222 arabisiflrum Dec.	Arabis-flwd	£ A,A or ft ap,jl P	Siberia 1800.	D co	
^'rabis grandiflra L.	abid.				
JS?i ^{UN} ARTLE HONESTY. (Luna, the moon; broad silvery silicles.) Crucifer*.					
16223 rediviva L.	revived	A or 3 my,jn L.P	Germany 1596.	D co	Lar.
biennial ,					
16224 biannis Moen.	4 my,jn L.P	Germany 1570.	S co	Lar.	
annua L.					
2 albiflra Dec.	white-flowered	£ CD or 3 my,jn W	Germany 1570.	S co	

1834. *MCCOTTIA* L. RICOTIA. Egyptian Lunaria Dec. 
1835. *PARTHENOCARYA* L. *Turra* FABSETIA. (Philip Farsetti, a noble Venetian botanist.) *Crycifera**. 6.-7. co Desf. at. 2.160
- 16226 cheiranthoides R. Br. *Stocklike* tt. Jdu 1 jn.jl W.P Levant 1788. C co Ven. eels 19
- sgyptaca *Turra*, Cheiranthus Farsttia L. 10227 sutruficosa Dec. sub-shrubby tt. or 1 ap Y Persia 1823. C co Ven. eels 19
- 16228 lunanoidea ff. Br. Lunaria-like tt. or 1 jn.jl T Archipd. 1731. S co Tou. it 1.248
- Lunaria graeca IV. en. 
- 0229 criocarpa Dec. woolly-fruited tt. or 1 jn.jl Y Greece 1820. S s.p Deless. 2.34
- 10230 cleopatra B. Br. bwkla-podded & Q or 1 jn.jl Y S. Europe 1596. S co Dalech. 1141.1
- 16831 cheilanthifoli? Del. Cheiranthus lvd O or 1 jn.jl Y Levant 1818. S co -rily'ssum cheiranthsblum IV. 
- 16232 *{ERTERO\} Dec. BERTEHOA. (C. J. Bertero, a friend of Decandolle's ft) hoary Q or 1* Jl W Europe 1640. 
- 16233 mutabilis Dec. changeable £ A or " 1J jl au W.pk Levant 1802. S co Vcn. eels 85
- ifioo. * «*» mutabilis R. Br. oblique ■■■■■ or 1 jl W Sicily 1823. C co Fl.gr. 633
- 16234 obliqua Dec. oblique ■■■■■ Y botanical draughtsm.) *CruciferAi.* 3L
- iJ £ ? ^{AUBBIEV} W* Adan. AUDRIETIA. (M. Aubrkt, Leaut 1710. D p.l Bot. mag. 126
- 16235 deltoidea Dec. deltoid %> A ' ' \ »" Farsetia deltoidea R. Br. rily'ssum deltoideum L. 
- 16236 purpurea Dec. purple t- A Fr * ns-J" Greece 1820. D co Fl.gr. 643
-  4 «rjd P S. Europe 1823. D co Sw. fl. gar. icin. Druba Asperidiflora Lam.
- 16237 vesica RIA Law. VESICARIA. (*Fesica*, a bladder; inflated silicles.) *Cructferar.* 4. —12.
- utricularia Law. bladdered A or 1 apjn LY Levant 1739. S sl Botmag. 130
- utricularia A Af reticulata Lam. netted ■■■■■ A or 1 my.jn Y S. Europe 1700. S s.l Lam. il. 559.2
- uni Vesicularia L. sinukta P. < Cretan Q or ap.jn LY Spain 1595. ■■■■■ a h.2.134.1
- 10241 crdtica Poir. Cretan Q or my.au Y Crete 1739. ■■■■■ B.I Alp. ex. 118
- rily'ssum crdticum L.
- 16242 Andrz. SCHIVEBBCKIA. (Andr. Scrivereck, a Russian botanist.) *Crtctfene.* 1.
- Idolica Andrz. Podolian £ A or 4 my.jl W Podolia 1821. D s.p Sw. fl. gar. 77
- ifl'ssum podolicum Hort.
- *1840. ^LY'SSUMZ. MADWOHT. (A, priv., lyssa, canine madness; sup med. qual.) *Crnc'tfer<e.* 32.-^53.
- Seeds in each cell for 2, sometimes boraeredwUha wingi^mens toothed
- 16243 ADYBE^THIM. ISSaxatilc L. 
- y*4 gemoplnsc L. Gemona tt. <r 1 ap.my Y Candia 1710. C s.l Hot. mag. IMI
- 5*5 orientale Andr. oriental f or 1 ap.my Y Europe -- CCoo J ^ % « »
- spatulate or Jap.my Y Siberia 1818. C eo Deless. 275.1
- silvery Bertoloni's A or 1 ap.my Y Switzerl. 1823. D co
- wall A or 1 ap.my Y Switzerl. 1820. D co W. & K. 1.6
- obtuse-leaved or 1 ap.my Y Hungary 1828. C co Deless. 2.38
- attain icum / to n Atlantic or i ap.my Y Crete 1817. C co Desf. at. 2.141f
- *S wpyHfgleum itef. Thyme-leaved or i ap.my Y S. Europe 1822. C co
- ro? vem. < irn. vernal A cu 1 jn.jl 1819. C co Sob. mo. 96
- JgJ* tortifnum W.%K. twisted A cu I JAV Y Hungary 1804. D s.l W.&K.91
- JIS al. 'stre L. alpine A cu 1 jn.jl S. Europe 1825. D co Al. ped. 18.2
- 16246 Marschallidnum Andr. Marschall's A or J ap.my Y Caucasus 1820. D co
- alpfstre Art. 
- 5*7 montkn L. mountain £ A or ft.jl au Y 1713. D s.l Bot. mag. 419
- 16258 cuneif>lium Tefi. wedge-leaved £ A or 4 WJ Y Italy 1820. D co
- 16259 olympicum Hort. Mt. Olympus £ A or 1 jn.jl Y 1700. D co
- 16260 diffusum 7Vn. diffuse £ A or 1 jn.jl Y Italy 1820. D co
- 16261 WulfetitnumW.cn. Wulfon's £ A or 1 jn.jl Y Cannthia 1819. D co
- G22 rostratum Stcu. beaked O un 1 jn.jl Y Crimea 1823. C co Acs pet 3.15.1
- 16263 umbellatum Desv. umbellate O or 1 jn.jl Y Tauria 1821. C co
- 16264 mieroptalura Fis. small-petaled O un 1 my.jl Y Siberia 182a C co
- 1627 campestre L. field O un 1 jn.jl Y France 1768. B.p Bar. ic. 912. 2
- 1627 G hirsutum Bieb. hairy O or 1 jn.jl Y Tauria 1817. C co Delesca.2 40
- 1627 calycinum L. calyxed O un 1 jn.jl Y Austria 174a C co Jac. au. 338
- 16268 minimum IV. smallest O un 1 jn.jl Y Spain 1791. C co Trat. thi. 35
11. ANODONTEA. — Stamens toothless; silicles subinflated.
- 16269 edatatum W.Sf K. toothless £ Q un 1 jn.jl Y Hungary 1820. S co W. & K. 1.92
- 16270 dasycarpum Step, thick-fruited O or * jn.jl Y Siberia 1819. S co Mag.bcr.18IK.?
- 16271 macrocarpum Dec. long-fruited tt. or i my.jl W France 1823. C 61 Deless. 2.41
- 16272 rupestre Ten. rock tt. un 1 jn.jl W Naples 1825. C co Fl. nap. (X)
- 16273 Aqulifolium W. Purslane-lvd tt. un S jn.s W S. Europe 1820. C co Boc. mu. 39
- 16274 spinosum L. thorny tt. Jcu * jn.jl W S. Europe 1683. C s.l Bar. Ic. 808
- ff. Desy. WBNIOCUS (Mene, the moon, okkos, eye; seeds?) Cruciferte. I.
- ill'ssum Aniliuum Step. Flax-leaved O or i jn.jl Y Caucasus 1817. S co Deless. 2. 42
1842. ICONIC R. Br. KONIGA. (Charles Konig, F.R.S., British Museum.) *Crucifrc.* 1.
- 16276 maritima R. Br. seaside O or V mB W England sea co. S co Eng. Got. 1729
- A/ssum marltimum Lam. Adysetum maritimum Lk. 2 variegata variegated tt. i J or 1 jn.s W C B.I
- Cav. wool-beafhig 8 J* * y J Y S. Europe 1710. S co f" i 1 M!

184*. PELT A*RI AX. PELTARIA. (<i>Pelte</i> , a small buckler; form of silicie.)	<i>Crucifene</i> . 2.-3.
16279 <i>tfliliacea L.</i> GarWe-scented ^ A pr 1 my.jl W Austria 1601. D s.l Jac. an. 2. 1-	
16280 <i>glastifolia Dor.</i> Woad-leaved O or 1 my.jl W & Europe 1825. S s.p Deless. 2.43	
1845. PETUOCA'LLIS R. Br. PETROCALMS. (<i>Petros</i> , rock, <i>kallos</i> , beauty & beautifying rocks.)	<i>Cruc. I.</i>
16281 <i>pyrenaica R. Br.</i> Pyrenean £ A pr ft my.jn Pk Pyrenees 1759. D s.l Bot. mag. 713	
Dråba <i>pyrenaica L.</i>	
* 1846. Dli A*BA L. DRABA. (<i>Drabe</i> , acrid; quality.)	<i>Crucifene</i> . 36.—60.
I. AIZ6PSIS.— <i>Plants perennial; flowers yellow & scapes naked; leaves rigid, ciliated; styles filiform.</i>	
16282 <i>aizrtides L.</i> Aizoon-like £ A pr } fap Y Wales rocks. D al Eng. bot. 1271	
16283 <i>ciliaris L.</i> ciliate-leaved £ A ii * f ■ Y Switzert. 1731 D s.l Ger. gal. 1311	
16284 <i>brachystemon Dec.</i> short-stemmed £ A pr ft.ap Y Switzcr. 1819. D s.l.p Bot. mag. 170	
aizoides B.M.	
16285 <i>cuspidata Bieb.</i> cuspidate ■ Apr ft fap Y Iberia 1820. D s.p.l	
16286 <i>fryoldes Bieb.</i> Bryum-like ■ A pr ft Up Y Tauria 1820. D s.p.l	
16287 <i>bTunurfbilia Stev.</i> Brunia-leaved ■ Al>r ft jn.jl Y Caucasus 1820. I BF	
16288 <i>fricstbia Stev.</i> Heath-leaved * A pr ft jnjl Y Caucasus 1821. D s.l	
16289 <i>pilosa Adams</i> pilose ■ & pr ft jnjl Y Siberia 1825. D s.l	
16290 <i>Azoon Wahl.</i> Aizoon £ A pr ■ Y Carinthia 1823. D co	
II. CHIRYSODRA^BA.— <i>Plants peren.; lvs not rigid nor keeled; fl. yel.; styles almost nones silic. oval-oblong-</i>	
16291 <i>alpina L.</i> alpine £ A pr ft ap.my Y Lapland 1820. D oo Wa. lap. 11.4	
2 siliqua pür>sa hairy-podded £ A pr ft ap.s Y Greenland 1820. U co	
16292 <i>glaciärs Adams</i> icy ■ A pr ft ■ Y Siberia 1H26. D s.l	
16293 <i>rcpens Bieb.</i> creeping ■ & ft jnjl Y Siberia 1818. D s.l	
16294 <i>Gmelii Adams</i> Gmelin's ■ & Jnjl Y Siberia 182a D s.l Gm. si. 3.56	
16295 <i>muricella Wahl.</i> small-prickly ■ A pr ft jnjl W Lapland 1810. D s.l Wa. lap. 11.8	
hirta Fl. dan	
16296 <i>irrömpa Stev.</i> rough £ A pr ft fap Y Caucasus 1821. D s.p.l Deless. 2.44	
III. LEUCODRA^BA.— <i>Plants perennial; leaves not rigid; flowers white; petals obtuse or subemarginate*</i>	
16297 <i>oblongMa R. Br.</i> oblongate ■ A pr £ myjn W Baffin's B. 1823. D s.p	
16298 <i>corymbosa R. Br.</i> corymbose ■ A pr £ myjn W Baffin's B. 182a D s.p	
16299 <i>hirta L.</i> hairy ■ A pr I myjl W N.Europe 1823. D e o Wa. lap. II. 3	
16300 <i>rupestris R. Br.</i> rock £ A ■ i myjl W Scotland al.roc. D bl Eng. bot. 1339	
hirta E. B.	
16301 <i>nivalis W.</i> snow £ A pr ■ ar W Switzerl. 1824. ■ 81 Fl. dan. 142	
16302 <i>lapponica W.</i> Lapland £ A pr ■ ar W Lapland 1810. ■ 61 Wa. lap. 11. 5	
^ndrosheca Wahl.	
16303 <i>helvt'ica Sc!.</i> Swiss £ A pr ft myjl W Switzerl. 1819. ■ 81	
16304 <i>fladnisensis Wul.</i> Fladniso £ A pr ft myjl W Switzerl. 1819. ■ 81	
^ndrostaea W.	
16305 <i>tomentosa Qair.</i> woolly £ A pr ft myjl W Switzerl. 1819. ■ 81	
16306 <i>stellata Jac.</i> stellate £ A pr ft myjl W Pyrenees 1820. ■ co	
hirta Jac.	
10307 <i>sihquida Bieb.</i> siliquose £ Apr I myjn W Caucasus 1822. ■ 81	
16308 <i>austriaca Crz.</i> Austrian ■ A pr i myjl W Austria 1824. ■ Si	
16309 <i>lasiocarpa Bock.</i> woolly-fruited ■ A pr imyjl W 1820. ■ 81	
IV. HALA'BGES.— <i>Plants biennial; styles short & flowers white, rarely yellow.</i>	
16310 <i>incana L.</i> hoary £ Q>pr J my.jn W Britain aLroc. S 81 Eng. M>138B	
conferta Ehrh.	
16311 <i>confusa Ehrh.</i> confused £ O)pr % myjn W N. Europe .. S co Fl. dan. 130	
incna Fl. dan.	
16312 <i>cinerea Adams</i> grey £ G>pr ft. myjl W Siberia 1818. ■ co	
16313 <i>dahurica Dec.</i> Daurian £ Q)pr ■ ■ W Dauria 1824. ■ co	
16314 <i>aurea Vahl</i> £ QD ■ Y Denmark 1820. ■ co Fl. dan. 1460	
V. DRA'BEA.— <i>Plants annual; styles none; flowers small, yellow or white.</i>	
16315 <i>liitca Gilib.</i> yellow C pr ft jnjl & Europe 18>4 S ■ An. mu. H-■	
16316 <i>nemorahs L.</i> grove O pr ft myjn Y Europe 1759. S ■ H. sys. 4.60 1	
16317 <i>murahs L.</i> wall O pr f my W England moun. S ■ Eng. bot. 91	
1847. ERO'PHILADflr. EROPHILA. (<i>Ero</i> , spring, <i>phieo</i> , to love; <i>time off</i> flowering.)	<i>Ciquidaea. 17*</i>
16318 <i>vulgäris Dec.</i> common ^ O pr i mr.ap W Britain walls. S s.l Eng. bot. 5B6	
Draba verna L.	
16319 <i>americana Dec.</i> American O pr i mr.ap W N. Amer. 1816. ■ co	
16J20 <i>pra'cox Dec.</i> early O pr ft fap W Caucasus 1820. ■ co	
Dråba pretax Stev.	
1848. COCHLEA^RIA Ton. SCURVY GRASS. (<i>Cochlear</i> , a spoon; concave leaves.)	<i>Cructferai. ■</i>
I. KERNIORA.— <i>Sile. globose, at base subtetragonal; filam. 4, in middle jointed; styles short; stigmas wag*</i>	
16321 <i>saxatilis Lam.</i> rock it A un ft jnjl W Austria 1773. D s.l Jac. au. 2. i<&	
3/agrum saxatilis L.	
16322 <i>auriculata Lam.</i> auriculate ^ A unfit jnjl W S. Europe 1818. S co ■ h.3.140.»	
II. ARMORAVEA.— <i>Silices elliptic or oblong; styles short, filiform; stigmas capitate* neartdisciforme</i>	
16323 <i>Armoracca /.</i> Horseradish * A cul 3 my W England wat.pl. D s.l Eng. bot. <&>	
16324 <i>macrocarpa AW^SK</i> : long-capsuled * A un 3 jl W Hungary 1806. D i l W.&K.W*	
III. CÜCHLEAR.— <i>Silices variable inform, at top not emarginate; styles very short or none.</i>	
16325 <i>glastifolia L.</i> Woad-leaved ■ ft myjl W Germany 1648 S co	
16326 <i>SngliciaL.</i> English ft my W Britain sea sh. S co Eng. bot. 55	
16327 <i>officinalis L.</i> officinal ft ap.my W Britain sea sh. S co Eng. bot. 5a0	
■ ■ S rotundifolia Sm. smaller I ap.my W Britain moun. S co	
16328 <i>pyrenXica Dec.</i> round-leaved i ap.my W Pyrenees 1820. S co	
16J29 <i>grdnlandica L.</i> Pyrcnca , ft ap.jn W D ■ ■ ■	
Greenland ft myjn F Scotland scalp. S co Eng. bot. 6»>	
16330 <i>danica L.</i> Danish \ myjn W Britain sea sh. S co Eng. bot. 6»>	
16331 <i>fenestrata R. Br.</i> windowed ft my.jn W Greenl. 1820. S s.p	
16332 <i>integrifolia Dec.</i> entire-leaved I my.jn W Siberia 1822. S s.p	
IV. IONOPSI DIUM.— <i>Silices rounded, compressed, at top emarginate; flowers Mac.</i>	
16333 <i>acaulis Desf.</i> stellate £ A un ft jaap Li Portugal 1824. D co Jac. ec.	
pusilla Bro.	
■ ■ ■ ■ ■ SHEPHERD'S PURSE. (<i>Thlao</i> , to compress; seedf.)	<i>Cruafer*. 8-—17<</i>
16334 <i>broad-leaved</i> ^r A un 1 mr.ap w Crimea 1822. D co ■ ■ ■	
16335 <i>ceratocarpum Mur.</i> horn-fruited O w lftjl W Sib'ia 1779. S co	

<i>!S5^{ar} y.^{ena} L-</i>	corn <i>Penny Cress</i>	O w	i jnjil	W	Britain	corn fi.	S co	Eng. bot. 1659
{6337 <i>colpnum Bkb.</i>	hill	O un	1 jnjil	W	Caucasus	1818.	S co	
J6338 <i>alhaccum L.</i>	<i>Gar-Wc-scented</i>	O un	* myjl	W	S. Europe	1714.	S co	Jac. ic. 1.121
J6339 <i>perfohatum L.</i>	perfoliate	^ Q w	i apjl	W	England sto. pa.	D s.l	Eng. bot. 2354	
J6340 <i>moptanum Z.</i>	mountain		w	Austria	...	S s.l	Jac.au. 3.2J7	
16341 <i>alpcstre L.</i>	alpine		w	England m.pas.	D s.l	Eng. bot. 81		

i*12 ⁵⁰ CAPSE'LLA 1 f.m.	SHEPHERD'S PURSE.	f	A diminutive of <i>capsula</i> , a capsule.)	<i>Crudferce.</i> 1.				
16J42 <i>bursa pasturis Moen.</i>	common	Q w	■■■	W	Britain	ro. sid.	S co	Eng. bot 1485
Thlaspi <i>bursa pastbris L.</i>								
2 minor Dec.	smaller	O w	f.n	W	Britain	ro. sid.	S co	Tab. ic. 199
3 integriflolia Dec.	entire-leaved	O w	1 f.n	W	Britain	ra sid.	S co	M. h. 2.20.1
4 coronopifolia Dec.	Buckhorn-leaved	O w	1 f.n	W	Britain	hea.	S co	
5 aptala Afx.	peUlless	O w	1 f.n	W	Britain	fields.	S co	

1851. HUTCHI'NSIIL R. Br. HUTCHTNIA. (*Miss Hutchins*, an Irian cryptogaraist) *Crucifer**. 7.—11.

*IBERIDE'LLA—Styles filiform; leaves entire or somewhat toothed.

Jr. round-leaved	jfc	A pr	i ray.jl	W.p	SwitzerL	1759.	D co	Al. ped. 1.55.2
^								

^ styl&a Deg.	Cepaea-leaved	tf A pr	* myjl	Pk	Carinthia	1824.	D i p	Jac. m. 2.1
/beris stylosa Ten.	to/i-styled	t A pr	imyjl	W.pk	S. Europe	1824.	D e o	Bot mag. 2772

calydna Desv.	II. "8A&ruRTfoLVM. —Leaves pinnateflobcd.							
large-calyxed	‡ A p r	i ap.my	W	Siberia	1823.	D 8p		

alpine	■■■ pr	4 ap.jn	W	Germany	1775.	D co	Jac. an. 2.137
rock	■■■ i	mr.my	W	England	rocks.	S co	Eng. bot 111

"idium petraeum L.	procumbent	· □ ■ I	ap.jn	W	Spain	1823.	S Bp	
ffimbens Desv.	kepidium procumbens L.							

T E S D A L I A B. Br. TEESDALIA. (R. Teesdale, an English botanical author.) <i>Crucifer</i> . 2.								
nudicaulis R. Br. naked-stalked	O pr	* my jl	W	Britain	gra.pa.	S co	Eng. bot. 327	

ifi<†iberi [†] nudicaulisi.	regular	□ pr	J f.my	W	S. Europe	1824.	S co	
16351 regularise.	:							

lepidium Dec.	CANDYTUPT. (From the country called Iberia, now Spain.) <i>Crucifer</i> . 22.-25.							
ifa ^B ERIDI-ASTRUM	Radicile horizontal; seeds submarginate; septum nearly double? fructif. pedicels racemose,							
LQ5Z semperflorens L.	ever-flowering tt. i J fra 1\$ year	W	Sicily	1679.	C r.m Za. h.	165		

U' IBBRI'DIUM.	— Radicle descending; seeds bnmarginate; septum simple; fructiferous pedicels racemose;							
Gibraltar	from No. 16364. coiybose.							

rock	■■■ or	I ap.jn	W	Span	1732.	C eo	Bot. mag. 124
				S. Europe	1739.	C eo	Gar.aixOI

Coris-leaved	tt. or	■■■ apjl	W	S. Europe	1800.	C eo	Bot. mag. 1642
pubescent	tt. or	*ap.jn	Pa.V.			C eo	

evergreen	tt. or	i apjl	W	Candia	1731.	C eo	Riv. tet. 224.2
Garrax's	tt. or	i apjl	W	Piedmont	1820.	C eo	Al. ped. 40.3.45.2

crowded	tt. or	i apjl	W	Spain	1827.	C eo	
bitter	O or	jnjil	W	Englandchalfi.	S co	Eng. bot 52	

intermediate	jfc Q or	jnjil	W	France	1823.	S co	Bul. ph. 82. 21
pinnate-leaved	O fra	jnjil	W	S. Europe	1596.	S co	Lob. ic. 218

sweet-scented	O fra	jnjau	W	Geneva	1806.	S co	Cl.h. 132.1
umbellate	O or	jnjil	W	S. Europe	1596.	S co	Bot mag. 106

Lagasca's	O or	jnjil	W	Spain	1822.	S co	
Flax-leaved	f Q or	jnjau	W	S. Europe	1759.	S p. I	Gar. aix 105

ciliata	■■■ Q or	jnjau	W	Caucasus	1802.	S co	Bot mag. 1030
Taurian	£ O or	f mil	R	Caucasus	1823.	S co	

violet	O or	jnjil	W	P	1782.	S co	
dwarf	jfc Q or	ijn.jl	W.p	Dauphiny	1822.	S co	Bot mag. 2788

spatulate	O or	jnjil	P	Pyrenees	1823.	S r.m	Berg. ph. ic.
Tenore's	£ A or	jnjil	Pa.P	Naples	1823.	C co	Sw.fl.gar.88

16373 coitacta Pert.	contracted	a. or	i myjl	W	Spain	1824.	C p.l	

1. IONDRA ^{BA} .	— Calyx at base two-spurred; gla.							
	car-podded	O or	1 jnjil					

lf*75 engenfölia Dec.	Ericron-leaved	O or	1} jnjil					
16376 nfpida Dec.	hispid	O or	1\$ jnjil					

16377 cichoriifülia Lot.	Succory-leaved	3l Q or	1 jnjil					

II."*■■■								
■■■								

16381 cilata Dec.	ciliated							

IV.	depresso							
16383 microcaria Dec.	small-fruited							

1855. MEGACARPiEW Dec. MEOACARPJBA. (*Megas, great, karoos, fruit.*) *Crucifene.* 1.—2.
16396 laciniata Dec. jagged £ Q or 1 jn.jl Y Siberia 1818. S s.p Dec. bis. 3
Biscutilla megacarpa *Fis.*

1856. EUCLID' DIUM B. Br. EUCLIDIUM. (*Eu, well, kleidoo, to shut up; silicles.*) *Crudferee.* 8.
16397 syriacum 11. Br. Syrian O cu f jlau W Levant 1778. S co Jac. au. 1.6
z̄tunias syriaca *Gae.*
16398 tat&ricum Dec. Tartarian O or f jl.au W Tartafy 1821. S co Pal. it. 3. ap. 77.2
Fella tenuissima *Pall.*

1857. OCHTHO'DIUM Dee. OCIITHIODIUM. (*Ochthodes, waited; surface of silicles.*) *Cruciferar.* 1.
16399 egyptiacum Dec. Egyptian O cu f au Y Egypt 17b7. S co Jac. vin. 2.145
2/funias aegyptiaca L. /apistrum ajgyptiacum // K.

1858. ANASTATIC A L. ROSE OP JERICHO. (*Anastasis, resurrection; reviving in water.*) *Crucifene.* 1.
16W0 hierochuntina L. common HJcu | jn.au W Levant 1597. S co Jac. vin. 158

1809. CAKPLE Tou. C A RILE. (An Arabic word used for this plant.) *Crudfer**. 3.-4.
16401 roaritima Sco. Ben Rocket O or f jn.s P Britain seash. S s.l Eng. bot 231
16402 fegyptiaca W. Egyptian O or 1 jn.au P Egypt ... S co M. h. 2.3.7.20
mantima fi sinuatifulia Dec.
16403 americana Nut. American O or 1 jn.au P America 1823. S co Fl. ant. 1.17

1800. JIAP'I STRUMBauh. RAPISTRUM. (*Rapum, a turnip; similar leaves.*) *Crucifere.* 3.-5.
16404 perenne Berger. >crennial ^ Dun ij jl Y Germany 1789. D 61 Jac. au. 5.414
CnBle pr̄ennis Herit. il/yagram perenne L.
16405 rug&suin Berger. wrinkly O un 1f jn.jl Y S. Europe 1739. S «l Al. ped. 1.78
Cak)e rugbsa Herit. A/yagram rugosum L.
16406 orientMe Z)r. oriental O un 1 jl Y Levant 1795. S co Fl.gr. 612
A/yagram onentale L.

1861. CHORI'SPORA Dec. CHORISPORA. (*Choris, separately, spora, seed; separated.*) *Cntcfera.* 3.-4.
VQVJ ten&la Dec. slender O pr * jn.jl P Siberia 1780. S co Pal. it. 3. L. 3
iEaphanus tcnellus Pall.
Sarcuata bowed O pr | jn.jl P Siberia .. S co
iZiphanus arcuhtus W.
1H408 sibinca Dec. Siberian O un & jn.jl Y Altai 1823. S co Mur. 1775.48.11
1^&ibenca Itec. Iberian O un i jn.jl Y Iberia 1821. S co Deless.2.58
1852. MALCOTMTA R. Br. MALCOMIA. (IV. Malcolm, mentioned by Ray.) *Cructfce.* ft—16.
16410 africana R. Br. African O or f jn.jl P Africa 1747. S s.l Boc. sic. 42.1
//&peris africhna L.
1G411 /aruxacit'ulia Dec. Dandelion-leaved O or P P Siberia 1795. S l.p
ICA12 ISxa Dec. loose O or P P Siberia 1820. S co
1S413 cMa Dec. Chian dwarf-branching O or ifi P Chio 1732. S si Di. el 147.178
16414 mnrtima R. Br. seaside O or my.jn V S. Europe 1713. S s.l Bot. mag. 166
Cheiranthus maritimus L.
16415 arenaria Dec. sand O or | jn.jl V Algiers 1804. S s.l Desf. at 2.162
//&spcris arcnaria Dec.
16 H6 pa'viflora Dec. smallUflowered O or P V S. Europe 1823. S co Dec'ic.35
15417 lyr'ita Dec. lyrate O or ft P Cyprus 1820. S co Fl. gr. 635
Cheiranthus lyratus Sm.
13*18 littrea R. Br. shore O or 1 jn.n W.v S. Europe 1683. S s.l Lob. ic. 331.1

* 1863. //E'SPERIS L. ROCKET. (*Hesperos, evening; flowers more fragrant towards.*) *Crucifene.* 17.—23.

I. HESPEM'DIUM. — *Limb of petals linear, dark; silicles two-edged, with the valves keeled.*
18<9 tristis L. sad £ Q or 1 ap.jn D.P Austria 1629. S 61 Bot. mag. 730
16420 fragrans Fis. fragrant £ Q or 1 my.au D.P Siberia 1817. S p.l

II. DECLO'SMA. — *Limb of petals obovate; silkies cylindrical or subtetragonal.*
16421 laciniata All. fringed £ Q or li my.jn P S. France .. S co Al. ped. 82.1
16422 runcinata JV.S/F.K. . runcinate £ QD or U jjjf W.p Hungary 1804. S s.l W.&K. 2.200
16423 bituminbsa Sam. clammy £ Q or ij jn.jl W.P 1816. S 61
16424 matronalis L. matronly £ A or 4 mv.au P Italy 1597. D p.l Lam. II. 564.1
2 fibre albo white-flowered £ A or 4 my.au W. D p.l Tab. kr. 692.1
3 fibre pleno rhbro double-rcd-flwd £ A or 2 my.au R. D p.l Weia 572. C.
4 fibre plfcno albo double white-flwd £ A or 2 my.au W. D p.l
5 fibre plfcno varicgato var.dble-flwd £ A or 2 my.au W.R. D p.l Munt. ph. 186
6 folunora leaf-flwrcl £ A or 2 my.au G. D p.l Wcin. 572. A.
16425 inodora L. scentless £ A or 4 my.jn Pk Britain pas. D s.l Eng. bot 731
matronalis 3 sylvestris Dec.
164C6 siberica L. Siberian £ A or 3 my.au Pk Siberia 1800. D p.l
matronalis y siberica Dec.
16127 heterophylla Ten. various-leaved £ A or 4 myau R Italy 1823. D co
16428 grandiflora B. M. large-flowered £ A or 4 jl.au W.p 1820. D co Bot. mag. 2683
16429 etta Horn. tall £ CD or 4 jn.jl Pk Europe 1821. S co
16430 aprica Poir. exposed £ A or i my.jn P Siberia 1822. ■ co
16431 rp̄anda Lag. repand £ Q or 2 jn.jl R Spain ' 1821. ■ p.l
1/4.32 bicuspida Poir. two-pointed £ Q pr 1 jlau R Armenia 1827. ■ co
Cheiranthus bicuspida IV.
16433 pulchella Dec. neat O pr ■ jlau R S. Europe 1827. S co
16434 ramosissima Desf. branchiest O pr ■ jlau R Algiers 1816. S co Desf. at 2.161

1361. ANDREOSKIAf Dec. ANDREOSKIA. {(*Ant. Andryejowški, a Russian botanist*) *Crucifer**. &
1G435 integriflbia Dec. entire-leaved ■ Q un 1 my.jn W.R Siberia 1816. S co
Aiſymbrium intgriflbum L.
16436 eglandul.sa Dec. glandless O un 1 jl.au ■ Siberia 1818. S co
A'ktinbinus Sievers/Ett'i Rcdow.
16437 peothiat Dec. pectinated ^ QQ un 11 my.jn W Siberia 1827. S co
iſymbrumpotinatum Dec. i/espous pinnata Pers.
Spinatiffida W. pinnatifid ^ Q un 1 my.ap W Siberia 1857. S co
Cheiranthus pinnatifidu* I"

1865. SISY'MBRIUM L. SISVMBHIUM. (*Sisybos*, a fringe j fringed roots.) *Cruet/era.* 43. fil.

16502 firmum Swt.	firm	£ Q or	1 jl.au	Y	Switzerl.	1819.	S co
Cheiranthus f. firmus W. en.							
16503 collimum Andrx.	hill	£ Q> or	1 myjl	Y	Caucasus	1823.	S co
Cheiranthus collinus Bieb.							
16504 ictophyllum Uum Andrz.	fine-leaved	£ Q> or	1 myjl	Y	Iberia	1821.	S co
Cheiranthus leptophyllum Bieb.							
16905 pumilum Horn	dwarf	£ Q> or	1 myjl	Y	Switzerl.	1819.	S co
pätilum Horn	spreading	£ Q> or	1 myjl	Y	Europe	1820.	S co
16507 diffusum Ehrh.	diffuse	£ Q> un	1J myjl	Y	S. Europe	1731.	D e o Jac.au. 1.75
Ianceolatum B. Br. spear-leaved	£ Q un	1 myjl	Pa. Y	& Europe	1597.	S co	Jac.au. 1.74
Cheiranthus erysimoides L.							
16509 versicolor Andrx.	party-colored	£ CD or	1 myjn	Va	Caucasus	1825.	S co
Cheiranthus versicolor Bieb.							
16510 rheicum Dec.	Rhessian	£ Q> or	1 myjl	Y	Switzerl.	1819.	S co
Cheiranthus rheticus Schl.							
16511 longifl. Hiim Dec.	large-flowered	£ Q) or	1 myjl	Y	S. Europe	1823.	S co
grandiflorum Desf.							
16512 decumbens G. Don	decumbent	£ Q or	1 myjl	Y	Switzerl.	1819.	S co
Cheiranthus decumbens Scnl.							
16513 diffusum Dec.	doubtful	£ Q un	1} myjl	Y	1823.	S co
Cheiranthus dubius Horn.							
16514 asperum Dec.	rough	£ Q) un	1} myjn	Y	N. Amer.	1822.	S co
Redowskii Wein.	Uedowski's	£ Q) or	1 myjl	W	Siberia	1821.	S co
16516 strigatum Dec.	strigose	£ Q) or	1 myjl	Y	Siberia	1816.	S co
Cheiranthus strigatus Led.							

III. CORINGIA.—Styles scarcely any; petals erectish; leaves cordate, clasping the stem.

16517 ahfignum Baum.	alpine	£ A un If	myjn P.Y	Germany	1793.	D 81	Vil. del. 3.36
ifrassica alpina L.							
16513 perfoliatum Crx.	perfoliate	O or	1 jn.jl	W	Austria	1818.	S co
/frassica orientalis L.							

16519 austriacum Baum.	Aust Hare's ear	O un	1 jn	W	England	cliffs.	S co
ifrassica austriaca Jac.							

IV. DU*BIA.—Doubtful to which of the sections they belong.

1fi>20 altissimum LeJ.	tallest	£ Q) or	3 myjl	Y	Germany	1818.	S co
16521 bicolor Dec.	two-colored	£ Q) or	1 myjl	Y	Switzerl.	1819.	S co
Cheiranthus bicolor Horn.							

16522 quadricorne Step.	four-horned	O or	1 jn.jl	Y	Siberia	1821.	S co

1868. LEPTA ^N LEUM Drc.	LEPTALEUM.	{Leptaleos, slender: leaves.)			Cruciferar.	1.—2.
16523 filiforme Dec.	thread-leaved	O cu J jn.jl	Y	Siberia	1820.	S co Deless. 2.68
Sisymbrium filiforme W.						

1869. STANLEY Nut STANLEYA. {Edward Lord Stanley, F.R.S. vice-pres. of Lin. Soc.)	Crucifer. 1—3.
16524 pinnatifida Nut.	pinnatifid
Jf AJ or	1 jn.jl
	Y
Louisiana	1816.
S p.l	

Cleome pinnata Ph.

1870. CAMELPNA Crx.	GOLD OF PLEASURE. {Chamai, on ground, linon, flax; dwarf.)	Crucifer. 5.—6.
I. CHAMELPNUM.	—Silicles obovate, marginate; styles conic; stigmas simple.	

16525 saftva Crx.	cultivated	O ec	1 myjl	Y	Britain	corn fi.	S s.l Eng. bot. 12J4
Jtfagrum sativum L.							

1 pilosa Dec.	pilose	O ec	1 myjl	Y	Britain	...	S co M. h. 2.21.2
Cheiranthus sylvestris Wallr.							
2 glabrata Dec.	smooth	O ec	1 myjl	Y	Europe	1806.	S 81 Bau. his. "2, -"
Pers. tooth-leaved		O un	1 myjl	Y	Podolia	1820.	S co Deless. 2.G9

16526 dentata Pers.	tooth-leaved	O un	1 jn.jl	Y	Europe	1806.	S 81 Bau. his. "2, -"
16527 microcarpa Andr.	small-fruited	O un	1 jn.jl	Y	Podolia	1820.	S co Deless. 2.G9

II. PSEUDOLIVM'.	—SUicles globose, immarginate; styles filiform; stigmas capitate.						
16528 barbarorum/ia Dec.	Cress-leaved	£ A un	1 jn.jl	Y	Siberia	1818.	S co Deless. 2.70
16529 austriaca R. Br.	Austrian	£ A un	1 jn.jl	Y	Austria	1795.	S s.l Jac.au. 2.11*

187L KE'SLIA Desv.	NEBLIA.	(Not explained.)			Crucifer. 1.		
16530 paniculata Desv.	panicked	O un	lijl.a.u	Y	Europe	1683.	S co Gae.fr. 2. HI
A/ygram paniculatum L.							

1872. SENEBI'TRIL Poir.	WART CRESS. {J. de Senebier, of Geneva, aveg. physiol.)	Crucifer. 3.—4.			
16531 pinnatifida Dec.	pinnatifid	O w i jLau	W	England	rubble. S co Eng. 1

Chironopus didyma Sm.	Chironopus didymum L.	O w i jn.a.u	W	Britain	ro. si. S CO Eng. I
36532 Cbrnopus Poir.	Buckhorn	O w i jn.a.u	W	Britain	ro. si. S CO Eng. I

Cbrnopus Kuclii Gac					
16533 nilotica Dec.	Nilotic	O cu *	jn.jl	W	Egypt 1827. S co Del. scg 34.?

Cochlearia nilotica Del.					
1873. LEPKDIUM L.	PEPPBRWORT.	{Lepis, a scale; form of silicles.)			Crucifer. 31.—58.

I. CARDALMA.	—Silicles ovate-cordate; valves subturgid, wingless, terminated by the exserted filiform	"g{£
16534 Draba L.	Whitlow	O un 1 myjn W Europe 1596. D co Jac.au. 4.w

Cochlearia Drkba L.		
1874. ELLIPSA'RIA.	—Silicles elliptic; valves keeled, wingless; style filiform, long.	

16535 chalepense L.	Aleppo	O un 1 myjl	W	Aleppo	1718. S co
16536 lastifolium Desf.	Woad-leaved	O un 1} myjl	W	Barbary	1823. S co Desf. at 1A?
16537 ample*icaule IV.	stem-clasping	^t A un \ jn.jl	W	Siberia	1826. D co

II. BRADYPI'PTUM.	—SUicles elliptic; valves keeled, wingless; calyx persistent.
16538 coronopifolium Fts.	Buckhorn-lvd

16539 sativum L.	cultivated	Cress O cul	1} jn.jl	W	Persia 1548. S co
2 crispum Dec.	curled	O cul U	jn.jl	W	Persia 1548. S co Bau. nffg
3 latifolium Dec.	broad-leaved	O cul	1* jn.jl	W	Persia 1548. S co M.b. & "» ^

Y [»] LEPIA.—'Silicles suborbicular, emargin.; valves navicular, winged; wings adhering to the short style; cotyledons	
16540 campestris R. Br.	hoary field

Thlaspi campestris L.	
16541 hirtum Sm.	hairy

16542 spinaum /..	prickly
O un 1 s	W

	Levant	1787. S co
--	--------	------------

VI.	<i>Silicles subelliptic, at top emarginate; valves keeled, wingless; style very short.</i>
16548 <i>Subulitum L.</i>	<i>Virginian & V. leaved ft.</i> O un 1 Jn.jl W America 1713. S co Sciih. han. 2.183
Jia45 <i>ruderMe L.</i>	rubble O un 1 Jn.jl Britain sea co. S co Assoar. 6.3
1W46 <i>incisum Roth</i>	<i>cut-leaved</i> O un 1 Jn.jl Tauria 1818. S co
JMSZ <i>perfolitum A.</i>	bladdery O un 1 ap.au Crimea 1820. S co Bux. c. 1. 26
Cardamines L.	perforated O un 1 Austria 1640. S co Jac. au. 4. SMJ
Spanish Cress	Spanish Cress Q un 1 Jn.jl Spain 1789. S co Ard. sp. 1.18
divaricate tL	divaricate tL [un 1] my.au C.G. H. 1774. C p.l
Bona Jen e £.	Buenos Ayres O un 2ft my.jn S Amer. 1732. S co Di. el. 286.370
cuneif. Clum Zfcc.	wedge-leaved O un 1 Jn.jl NewCam. 1820. S co
Seel 2 um Foss.	Fish-poison O ec Is Societysl. 1779. S co
16554 <i>Ayslopif Wium Desv.</i>	Hyssop-leaved O un 1 jn.jl N. Holl. 1820. S co
VII. LISPIDIA'BTRUM.	<i>Silicles subeUiplic, very entire valves keeled, wingless; style very short.</i>
Nitaceum ffrst	pather W N. Zeal. 1824. S co
lyrate	lyrate S co
broad-leaved	broad-leaved Britain seaco. D co Eng. bot. 182
similar	similar Siberia 1819. D co
' thick-leaved	' thick-leaved Hungary 1820. D eo W. & K. 1.4
suffruticose	suffruticose Spain 1683. S co Cav. ic. 2.161.2
Cao.	Spain 1823. S co
J8561 line ire D.	linear Germany 1793. S co Lob. ic. 223
16562 /ben s/;	Iberis
riccimum Dec.	African *L 1 J un i w C. G. H. 1816. S co Deless. 2.73

VIII. DU'DIUM.—Doubtful to which section it belongs.
J6564 diffusum Dec diffuse O un 1 jn.jl W N. Amer. 1823. S co

BIVON*4.	<i>{Ant. Bivoni Bernardi, an Italian botanist.)</i> Cruciferae. 1.
Thlaspi ICiteum Biv.	O un ft Jn.jl Y Italy 1824. S co Cup. sic. 2.256

EUNO^MIA	EUNOMIA. (JSu, well, name, fodder; used for fodder.) Crucifera. 1.—3.
epijositifMia Dec.	opposite-leaved £ _£ dt 1 jn.jl W Syria 1827. S co Lab. syr. 5.9
£epidium opositifolium Lab.	

THIONE^MA R. Br.	JETUIONEMA. {Aitho, to scorch, nema, stamen; tinge.) Cruciferae. 1.—d.
8ax&tie Jl Br.	rock O pr * Jn.jl l' S. Europe 1759. S co Jac.au. 3.236
»P> saxifl L.	
ib 36y Juxbarmi Dec.	slender O pr i Jn.jl Pa.R Carniola 1823. S s.p. Boc.mu.70
IRP-A »P> arachicum Btcb.	Buxbaum's O cu jn.jl Pa.R Levant 1823. S co Bux. c 1.5.1
»70monospermumil.Br.	one-seeded ^ Q) cu ft jl.au Pa.P Spain 1778. S co

*1877. iSA^TIS JBbtA. WOAD. (Isazo, to render equal; supposed effect on rough skin.) Crucifera. 12.^11L

I. SAMERAMUA.—Silicles oval or orbiculate, indehiscent, girdled by a broad leafy wing.	
ArmeniaL.	Armenian O or Ifjtl.au Y Levant 1822. S co Desv. 3.25.6
mica Brot.	Portuguese O or 1 my Y Portugal 1822. S co

II. GLA'STUM.—Silicles oval oblong, nearly linear, subdehiscent j margin subcrouse.

Kit.	early	V	... liQf DD s.l Al ped. 86.2
Dec.	seaside		Hungary 1822. S co Trat. ar. 2.68
iia Fis.	blunt-fruited		Tauria 1823. S co Deless. 2.76
Stcv.	common dyer's field		Tauria 1824 S co Deless. 2.79
TM>19 banfatica Lk.	Bannatic		England corn fi. S 8.1 Eng. bot. 97
Ji80 orientMis IV.	oriental		Persia 1824. S co
16581 canescens Dec.	cancsent		Bann.it 1819. S co
2 ib'rca Stev.	Iberian		S. Europe 1818. S co
16582 alppica&x>.	Aleppo		S. Europe 1822. S co Bux. c. 1.5
			Ibeia 1823. S co
			Levant 1739. S 8.1 Scop. d. 216

16583 lasioc&pa Fa. hairy-podded O or ft jn.jl W Siberia 1824. S s.p. Cruciferae. 1.
16584 gymnocarpa Fis. naked-podded O or ft my jl W Siberia 1820. S 8.1 Fisch. & fil. in.

16585 perfolitum L. perfoliate O pr ft jn.jl Pa.Y France 1648. S co Schk.han. 2.178

16586 lith6phila Bieb. stone-loving £ CD un 2 jn.jl W Iberia 1824. S s.p. Deless. 2.80
Crambe macrocfirpa Bieb.

16587 tarMts Dec. smooth-leaved O or my.jl Pa. B Astracan 1827. S co Deless. 2.81
Aaphanus lsvigatus Bieb.

16588 tomentosum Dec. woolly W. ^ CD cu 1 ap.jl Y Astracan 1817. S s.l PaLit. 103.2
Chenanthus tomentosus W.
16589 torulosum Dec. twisted ^ Q) cu 1 ap.jl Y Iberia 1820. S co

16590 oleraceal. common oleraceous £ Q> cu 2 ap.jn Y England cliff*. \$ r.m Eng. bot 657

Gaiiien Varietiet.

4 *

I. nccjilala Dec.	
X vulgirU Dec. common, IUGA>Ar&C Cab. M.t.3.1.6	
3 jwordolia Dec. Oak-leaved i.m.	
4 Mhtllca Dec. Sabdlian, Borecole Lob. ic 2+7.1	
•Julia Dec. Pulm.leaved, Cfw Palmier	
6 covttai Dec. rilibed, Oiou dc Beaueais	

\ 2. cosQta Dee.

1 vulgSrls Dec. common, Ch. & gr. cStes I^h.ic 244.1	
3 oblonga oblong, Cow trmcchuda M. h. & L 4 t & bulltta Deo.	

5 mtijor D<:, grfater,

k, BntMsU sprouts

i L.

2 (ileffra iJtr.

3 pabuliri i

4 A'ipo-ftrissina L.

S (imiinilnis /Vc.

3 rutabaga 7J<,

Ili59f Jtp»

5 depressa 2Jcc.

3 obf'iiK'

4 illufern D.c.

npai /,

5 olclfcrtu Are.

3 esculentuH Dec.

Sflbtu /

S HAviii; /J--i,

p * A.

16596 clin&wls L.

ifoJG violJLceu L.

hoary

llulcaric

scmporvireni Schrank

Vi'm Jtrirni'n.i

HQO (imVuiv Ten.

innalftida ltsf.

MMS huillis A-c.

p y j

166W mtnlnxi /f/E.

16805 repii) ib Zfec.

II. E~~d~~Cfc8TtUM. — Silicic? south; terminated by a content bank at base, 1 or 2 nccdril.

JfflXi Riditr.-i Vil.

166i7 iiiontnsii //J..

lulu of Man

lftW Cheirfinthitis Vil.

16610 ebcinmthM^ni Dec. Rtnck.flwd

/i:i'hjinus choir.'inthifl^rn JV.

Ifi6j] Ttiundt'iirtit 6'uii.

Tonrfnefort's

16813 ftevigitiU /.'A'.

niuoniltJiviu-if

10613 valcndna D.rr.

Vnkntin

i'sisjmbrum valent'num L.

15fi14 Pfuudo-crucSitruB JTriv. BasURocktt

10515 fkvtlcuulut Cyr. ihrubby |£

III. MicdPo*D[tjM.—Siictfx rotted above receptacle on tkortpedietisi tlyla thort, tleuter, U-rminateH

copiate tUgma;

16616 clongMa Ehrh. ctong.itcd tf a* un ^ »<>> Y Hungary 1801. S L\ W.&K. 1.

16617 subulirU li>vt. (iwl Q uti I jn.jl Y Spain 1813. S co Bnjt,i>b,4J

ISM. SINATIS n>».

to hurt, opt, the eye; Criicifra; £8.—^

X

1W18 rCtgIA L. black.

Sfurlgda

Prrs.

genitTitau Detf.

— Sitidfi h-rete ar stibte/ragona/; style* dart, small, not rostrifarm.

4 liy.jn Y Itritain corn ft. S r.m Ens

tmsM 4 mygn V Britain enru tl. S r.m

tw-InU'd 4 myJK Y Europe Itili. S co .

VIIIILIS 4 my.jn Y i W27. S co

* in. (bulk* IS19. 5 co

lointad 1 Jn.jl Y co

bk1 2 ju.Jl Y 1818. S CO

II, CBII»TOBI.>(*"P|!I. — Sflices terminated hi) an enifitir conicui beak.

16621 levigaU L. tmooth

1fi652 h i i f ouricilatc

16fi23 rnmfisa /tor. branchy

i i l W.

L pube>cent

16fi12 tafricB FU. com Charlock

Tauruin

ISM3 nutipinnatfida Las. mtaini

Dn Kater

WB5 All., „, Jao. Alluni

MS3G t6,rBld« JDti. turgid

E, indies IS imy. s co

China 170. S co W.h.b. L.

China 17Ha S co An Up. IP»

China 1B0L S eo

Sicily 1) r.m Ard.

Levant com fi. s i.i irnK

Tmiria IRIS. S ca

Ivt.i.n B TO

S eo

1S13. S co

icre terete, filoc.; cell* generally Isceded; beak ovate, Utfceded, smooth, haaty.jointed f' Q Un 3 jl Y S. fitropc 17/1. S CO , variuut-le;iv<1 <f_ Q> un 1} my Jn Y Spain 19B2. S co

.....) v. LSUCOSINA^PIS. — <i>Silkies hispid or smooth, surmounted by an cnisiform beak; valvules torulose.</i>
<<39 &ta L. white O HF 3 inal Y Britain corn fl. S rm Eng. bot. KW
hsipid 2 *un 2 jnl Y Morocco 1804. S rm Schou.4
hsipid &tal<n. cut-leaved O un 1 mr.ap Y Spain ... S co
iM hastata De. hastate O U 2 jnjl Y N. Holl. 1817. S co
leafy O un 1 ap.my Y Levant 1820. S co
wW* Spula Ten. Apulian O un 2 jnjl Y Italy 1823. S co
i*x* *. DiBA'ocwa — <i>Calyx somewhat spreading, at base bisaccate.</i>
16645 frutescens f.J.C. shrubby O -LJC U ljjn.d Y Madeira 1777. C si
VI. DU^BIA. — <i>Doubtful to what section it belongs.</i>
naked-stemmudf £ un 1 jnjl Y Syain 1818. b co
iJS ⁵ - MORICA^NDJ/f Dec. MORICANMA. (<i>Stephen Moricand, an Italian botanist.</i>) Crucifer*. 1 - 3.
TM ***** Oec. corn Cabbage find £ ® un 1 jn V S. Europe 1739. C co Boc. sic. 25. 3,4

1886. DIPIOTA'XIS Dec. DIPLOTAXIS. (*Diptoos, double, tans, seriesj seeds.*) Crucifer*. 11 — 14.

ion! , CATO¹ «PUM. — Styles scarcely any; s'dicles pendulous, sessile, or often pedkeeled; stigmas biobed.

S ¹ S ¹ i a 1 ? C hick, leav ~	O un ^ j n j l Y Sicily 1818. S co
16 M9 Binduia 2 5 ! * pendulous	O un 1* f.mr Y Burbary 1823. S co Desf. at 156
1<... psymbrium pendulum Desf.	
<< u0 hispida Dec. hispid	O un J ap.my Y Egypt ... S co Deleu. 89
.a.,, - ANoCA'hPiw. — Styles conk, compressed, terminated ^h by a two-lobed stigma; silkies erect, sessile.	
16651 erucoides Dec. I&uca-like	O un 1 jnjl W.p S. Europe 1736. S r.m Jac. v.n. 2.170
16652 virgata i)<?c. twiggy	O un f jnjl Y Spain 1817. S co
<wr.. *j5>> virgata Cow.	
16653 catholia Dec. universal	O un 1 ap.my Y Spain 1822. S co
ICTRA ¹ b r i u m cathlum L.	
16654 tenuifolia Dec. fine-leaved	A un l.jl.o Y England walls. D s.l il..a. ^ >w
1R e r T o s y m b r i u m tenuiflum L.	
16655 muralis Dec. wall	O un l.jl.s Y England san.pl. S co Eng. bot. 10'J0
*-n * <u>Imbrioni murale</u> £.	
Barreltiri Dec. Barrelier's	O un ijn.jl Pa.Y S. France 1770. S co Baric 1016
viminica Dec. twiggy	O un l my Y S. France ... S co Boc. sic. 10
rock	A un l in Y S. Europe ... D co
ROCKET.	(Uro, to burn: biting qualities.) Crucifer. 3.
1665y aaKva Lam. cuUivated-Jlwd	O cul 11 Jl Pa. Y S. Europe 1573. S 81 Schk. lian. 2.18d
jBrassica Ertica L.	
Salbo-gUbra white smooth O cul H W S. Europe 1577. S co Black. 243	
3 ilpo-pilbsa white pilose O cul U W S. Europe 1577. S co	
4 flavo-gUbra yellow smooth O cul U Y S. Europe 1577- S co BuIher.SI3	
5 flavo-pilotia yellow pilose O cul U Y S. Europe 1577. S co FL gr. 64G, W7	
«ffrigida Pers.	
1 (Tuccides Horn	turgid O cul ll Pa.Y & Europe 1577. S co
8 /	eruca-like O cul ll Pa.Y & Europe 1577. S co
***>Ivesicaria Cav.	exotic O cul ljl Y & Europe 1577. S co
bladdery O cul l jl W Naples 1800. S co	
bladdery O un If* jl Pa.Y Spain 1820. S co Assoar.4	

, JS⁸ - E'LLA L. CRESS ROCKET. (*Latinised from vrvar, the Celtic name of the cress.*) Crucifera. 1.

<*> Pseudo-cftisus L. Bastard Cytisus. i_J or 3 ap.my Y Spain 1759. C co Cav. ic. 142

BO^LEUM Desv.	BOLEUM.	(Bolos, a ball j round pods.)	Crudfera. 1.
asperum Desv.	rough	tt_Jpr 1 myjl Crea Spain 182a S s.p Desv. 3.6	

CARRICUTFRA Adulam. CARRICUTBRA.			Cruciferae. 1.
Fella Dec.	annual Veils	O pr	Eng.botlUJ

1891 SUCCO^WW Med.	SUCCOWIA.		Crucifers. 1.
1G665 balearica Med.	Balearic	i.l Jac. vin. 2. U4	

1892 PSYCHPNE Desf. PsveniNE			Crucifene. X
16606 stylpsa Desf.	style	O un 1 jn.jl Y S. Europe 1818. S co Dt*f. at 2. i<	
2 thlaspi Psychme	W.		

1(G67myagroides Forsk.	Myagrum-like	a. J cu 2 mr Li Egypt 1822 C co Vcn.mal.16	
2 macrocrpa Spr.	long-fruited	n. J cu 1 mr Egypt 1820. C co Gne. fr. 2.142 2	

Corvln, Desv.	Corvin's	O cu l ap.ju w S. Europe 1815. S co Bntf. ph 1'	
Myagrum iberoWes Brot.			
** faAK*ut		(A Greek name for sea cabbage.)	O ^r f/mr. 11.-11

IfifOTmaritimaL — ^wrr i o 1,1 qfsVtcle depressed, thick; s/ i^wa *mife; larger filament

Ibi671 pinnatifida R. oriental A A un 1 jnjl W Siberia 1759. D 81 Jac. ic. 1.128

WdSL orientals L. Tartarian A A cul 3 ju.jl Levant 1752. D sl

1W73 tafurica Jac. rough A A " 1 my W Siberia 1789. D sl Jac. ic. 1.129

16774 aspera Bieb. heart-leaved A A or G my W Tauria 1820. D co

IG>75 cordifolia Stev. Caucasus 1822. D co

III. DEVDROCIUMBE.—Lower joint of *sUicle filiform*, short; stigma seated upon a short style.
 16679 fruticosa L. shrubby «. LJ un S_x my.n W Madeira 1777. C s.l
 16680 stngusa Uerit. ~~theagre~~ «. i_J un 1J my.jn W Canaries 1779. C s.l Jac. ic. 1. ISO

1897. R A'PH ANUS L, RADISH. (*Rha*, quickly, *phainomai*, to appear; rapid germination.) *Cruciferce*. 4.-9.

I. RA'PHANUS.—*SUiclefungose, bUoclar.*

16681 saflvus L	cultivated	ft O cul 3	my.jn W.p	China	1548. S r.m	Lam.il.566
2 radicula Dec.	small-rooted	ft O cul 3	my.jn W.p	China	1548. S r.m	
2 rotundus Mil.	round-rooted	ft O cul 3	my.jn W.p	China	1548. S co	Wein.860.c
3 oblongus Dec.	oblong	Turnip ft O cul 3	my.jn W.p	China	1548. S r.m	
i 4 oleiferus Dec.	oil-bearing	Q ec 3	my.jn W.p	China	1548. S r.m	
chuunis MU.						
3 niger Dec.	black Spanish	fft Q cu ³	my.jn jW ^W China	1548. S r.m		
2 vulgans Dec.	common	ft O cul 3	my.jn jW.p	China	1548. S r.m	M.h.2.3.15.2
3 rotundus Dec.	round	ft O cul 3	my.jn W.p	China	1548. S r.m	M.h.2.3. Ul
4 griscus Dec.	gr.	ft O cul 3	my.jn W.p	China	1548. S r.m	
5 albus Dec.	white	ft O cul 3	my.jn W.p	China	1548. S rra	
orbicularis MU.						

16682 eaudatus L. tailed ft O cu 11 my.au Pa.P Java 18J5. S co L.f. dec.3.10

II. UAPHANISTRC/M.—*Silicic terete, acuminate; at maturity moniliform, joints one-seeded.*
 16683 Raphanistrum L. Wild Radish Britain edra fi. S Eng. bot 856
 2 flure ilbo Dtr. white-flowered O w W Britain corn fi. S Wcin. 86S.a.
 3 purpurascens Ztec. purpurascent O w P Britain corn fi. S Wein. fU&b
 4 flure flavo/tec. yellow-flowered O w Y Britain corn fi. S Bau.his.2.844.ic
 16684 Landra Moret. Landra £ Q cu 5 Y Italy 1820. S co
 16685 maritiinus 5m. sea ft A w 3 my.jl Y Britain seaco. D co Eng. bot 1643

1898. DIDE'SMIS Desu. DIIDESMIS. (*Dis*, twice, *desmos*, band; silicles of two joints.) *Cructfera?*. 2.-4.

16686 bipinnatus Dec. bipinnate O un 1 jnj.l W N. Africa 1827. S co
Sinapis bipinnata Desf.
 16687 aigypnacous Deso. Egyptian O un 1 jnj.l Y Egypt 1818. S co Deless. 2.92
Jlyagruin tegyptiacum L.

1899. ENARTHROCA'RPUSEft. ENARTHROCARPUS. (*Ennea*, 9, *arthron*, joint; *happosoffi*). *Ceneff.* 2.—
 V & S arcuatus Lb. arched Y Grète 1820. 3 co Labssyr. 2
 16689 pterocarpus Dec. winged-fruited O un Y Egypt 1823. S co Deless. 2.93
/dphanus pterocarpus Pers.

1900. 27UWIASZ,. BUNIAS. (Bounos, a hill; habitation.) *Cruclfera*. 3.
 16690 Erucago L. Erucago O un 11 jnj.l Y Austria 1640. S co Jac.au. 4.340
 16691 asprca Retz. rough O un U jnj.l Y Portugal ... S co
 16692 onentalis L. oriental ■ A un Ij my.jl Y Levant 1731. D co Gm.si.a57

1901. ERUCANRIA Gae. ERUCARIA. (Uro, to burn; biting qualities.) *Cruclfera*. 3.—Z"5*a a
 16693 aleppica Gae. Aleppo Condyllocarpus lacivagatus W. O un 1 jljau Wp Levant 1660. \$ 81 Gae.flr. & 14*-
 16694 tenuifolia Dec. fine-leaved O un li jnj.l Y Spain 1817. S co
Sinapis hispanica L.
 16695 crassifolia Del. thick-leaved O un \$ Jnd W.p Egypt 1823. S co Dcl.aBg.34.1
irratfsica crassifolia Forsk.

1902. HELIO'PHILA L, HELIOPHILA. (*Helios*, the sun, *phUeo*, to love.) *Cructfene*. 22.-49.

I. CARPONE^MA.—*SUicles sessile, terete, acuminate, indehiscent, or scarcely dehiscent.*
 16696 filiformis L. filiform iQipr Ijj.lau Pa.P C. G. H. 1786. S s.p Lam. il. 56* »
 11. LEPTORMUS.—*SUicles sessile, subcompressed, very slender, submoniliform; joints ovate, oblong.*
 16697 dissecta T/ian. dissected O pr 1 jnj.l B C. G. H. 1819. S s.p
 2 pinnata Vahl pinnate O pr 1 jn.au W C. G. H. 1792. S s.p

III. ORMISCUS.—*SUicles sessile, much compressed, moniliform; Joints orbicular, one-seeded.*

16698 amplexicaulis L. stem-clasping pendulous O pr W:W C. G. H. 1794. S s.p Jac.fr. a57
 16699 pindula W. pinnata Fen. O pr W:W C. G. H. 1794. S s.p Jac.fr. a57

16700 trifida Thutu. trifid O pr C. G. H. 1819. S s.p
 16701 pinnata L. small O pr W C. G. H. 1824. ■ s.p Pluk.m.432.2
 16702 /epidioides Lk. Lepidium-like O pr W C. G. H. 1818.

IV. SITESECAKPJE.—*Silicles sessile, compressed, oval, or orbicular.*
 16703 diffesa Dec. spreading O pr \$ jnj.l W C. G. H. 1818. S s.p
 16704 peltariaDec. shield-podded O pr f jnj.l W C. G. H. 1820. S s.p

Peltaria capensis L.

V. ORTHUSELIS.—*SUicles sessile, unipinnately branched near with straight margins.*
 16705 pilosa Lam. pilose O pr 1 mys B C. G. H. 1768. if s.p

16706 arabotodes B. M. integrifolia L. Arabis-like O pr f jnj.l B C. G. H. 1768. S s.p Bot mag. 496

16707 stricta B. M. pilosa ^ incisa Dec. O pr f jnj.l B C. G. H. 182a ■ s.p Bot man.

16708 digit'a L. digitate O pr 1 mys B C. G. H. ■ s.p
 16709 pectinata Burc. pectinated O pr 1 jnj.l W C. G. H. 18*9. ■ s.p

16710 coronopifolia L. Buckhorn-lvd. O pr Jijnjl V C. G. H. 1778. »p Her. lug.

If711/tt-niculcea R. Br. Fennel-leaved O pr C. G. H. 1774. »p
 16713 platysihqua R. Br. Sampliirc-lcnvcl sprr j Jns p C. O. H. ism ■ p

16714 fcoparia Burc. broad-podded tt. OJ pr 1 jljau V C. G. H. 1774. ■ s.p Deless. 2.98

16715 linearifolia Burc. broom IL uJ pr 1 jnj.l p C. G. H. 1819. ■ s.p

linear-leaved n. i_ pr 1 jnj.l R C. G. H. 1826. ■ s.p

VI. PACHYSTYLM.—*SUicles sessile, subcompressed; linear, velvety; style thick, conical, short; with entire leaves.*

16716 incana H. K. hoary n. j.J pr 2 my.au P C. G. H. 1774. C s.p

VII. CARPOPODUM.—*SUicles pedicellate, linear.*

16717 deonuAdes Dec. Cleome-like M-Upr 1 jl C G. H. 1802. C co

Cheiranthus strictus L. Cleome capensis L.

~~■~~ SUBULA^RIA L. AWLWOET. (Subula, an awl₅ form of leaves) *Cructfene*. 1.

16718 aquatica L. water £ O cu E jl W Britain W-TSIW--

■ 1904. CLECTME W. CLBOME. (*Kleto*, to shut. First applied to a plant like *Shupis*.) *Capp.* 20.—53.

16722 heptaphilus L.	seven-leaved	jl.au	R	Mac. 1823.	
16723 spinosai.	prickly	1 jn.jl	W	Jamaica 1817.	
16724 pungent	pungent	ODpr	2 jl.au	"W. Indies 1731.	s.p Marcg. br. 34
				W. Indies 1812.	al. W. h. b. 1.18
16725 pubescens B. M.	pubescent	O pr	ft.jnjl	W 1815.	S s.1 Bot. mag. 1857
16726 <i>rosea</i> Vahl.	<i>rose-colored</i>	O pr	i jn.jl	Brazil 1823.	S co Bot. reg. 960
II. SILIQUANUA.—Torus small; thecaporum short or none; leaves simple, from No. 16728. 3 v. 1 r. <i>I. foliate.</i>					
16727 monophyl Ha L.	one* leaved	Opr	ft.jnjl	Y Ceylon 1759.	S s.1 R. mal. 9.34
16728 <i>zevylancia</i> Dee.	Ceylon		jn.jl	Y	Bur. xcy. 100.2
16729 procumbens Jfcl.	procumbent	Opr	4 jn.jl	Y W. Indies 1798.	S Bl Jac. am. 120
16730 violacea L.	vinet-co/ored	O pr	jnjl	P Portugal 1776.	S Schk.han. 2.189.b
16731 Dilleniflma Dec.	Dillenioides W.	O pr	jnjl	W.Y Levant 1732.	S 81 Di. cl 266.345
16732 arabica h.	Arabian	O pr	2 jnjl	Y Arabia 1794.	S s.1 L. f. fa. 8
16733 ibérica Dec.	Iberian	O or	i jnjl	W Iberia 1820.	S 81
16734 <i>polygamia</i> L.	twiggy	O or	jnjl	W Persia 1820.	S s.1 Bux.c. 1.9.2.
16735 aculeata L.	polygamous	cm or	2 jnjl	W W. Indies 1824.	S s.1 SL jam. 12+. 1
16736 <i>aculeata</i> L.	prickly	O or	f	W S. Amer. 1817.	s.1
16737 diffusa Bauh.	Houston's	Opr		W W. Indies 1730.	S s.1
	diffuse	O or	1 jnjl	W Brazil 1823.	S 81

-1905- PERITOMA Dec. PERITOMA. (*Peritome*, a cutting round about; base of calyx.) *Capparide**. 1.

16738 serrulata Dec.	serrulated	O or	ft jn.au	P Missouri 1823.	S s.1
----------------------	------------	------	----------	------------------	-------

Cleome serrulata Ph.

1906. GYNANDRC/PS1S Dec.	GYNANDBOPSIS.				<i>Capp. 7—11.</i>
16739 sessiliflora Dec.	sessile-flowered	O or	1 jn.au	Indies 1820.	S s.1
*6740 cleome triphyllo L.	Cleome triphyllo L.	iQJor	1 jl.au	R 1824.	S 81 Bot. mag. 2G56
*6741 candelabrum Swt.	chandelier				
*6742 cleome candelabrum B. M.		O or	1 jn.au	W W. Indies 1816.	S s.1 Her.lug. 565.ic.
16743 triphylla L.	three-leaved				
*6744 cleome triphylla L.		fQBor	3 jn.au	W Carthage. 1818.	S s.1 H.&B.5.436
16745 speciosa Dec.	showy	O or	2 jn.au	W f. Indies 1640.	S s.1 Bot. mag. 1681
*6746 cleome speciosa H.S/B.					
*6747 pentaphylla Dec.	five-leaved	O or	1 jnjl	W.p Maranh. 1825	S 81
16748 pentaphylla Lindl.	neat				
16749 glabrata R. Br.					
16750 occidentalis R. Br.	western				
16751 glauca R. Br.	glaucous				
16752 longiscapa Swt.	long scaped				
glaucia B. M.	glauca B. M.	iAI or			

CLASS XVI. ORDER 1.

MONADE'LPHIA TRIA'NDRIA.

1907. PATERSO Wil R. Br. PATERBONIA. (*Colonel William Paterson, an excellent botanist.*) *Iridacee.* 7.

16745 scribba R. Br.	silky	lftmyjl	N. S. W. 1803.	D s.p Bot mag. 1041	
16746 lanata R. Br.	woolly	l Alor	i my.jj	N. S. W. 1824.	D s.p Sw. au. 15
16747 loiriifolia R. Br.	long-leaved	tf iAlor	i my.jj	N. S. W. 1818.	D s.p
16748 mima R. Br.	intermediate	y: Alor	1 my.jl	N. S. W. 1816.	D s.p Sw. au. ic. In
16749 glabrata R. Br.	smooth		1 my.jl	N. S. W. 1814.	D s.p Bot reg. 51
16750 occidentalis R. Br.	western	l Alor	1 my.jl	N. a. W. 1824.	D s.p
16751 glauca R. Br.	glaucous	l Alor	1 my.jl	N. S. W. 1820.	D s.p Bot. cab 1182
16752 longiscapa Swt.	long scaped	l Alor	1J my.jl	N. S. W. —	D s.p Sw. au. 39

1908. ORTHROSA/NTHES Swt. ORTHIROSANTHES. (*Orthros, morning, anthos, flower.*) *Iridacee.* 1.

16753 multiflora Swt.	many-flowered	iAJ or	1 jnjl	B N. Holl. 1820.	D s.p Sw. au. 11
-----------------------	---------------	--------	--------	------------------	------------------

•1900. FERUAHtlif L. FERRARIA. (*J. B. Ferrari, an Italian botanist.*) *Iridacee.* 6.

16754 undulata L.	wavy	JiAor	ft mr.ap	G.Br C. G. H. 1775.	O sp Bot mag. 144
16755 anisotropa Swt.	narrow-leaved	3 iAlor	ft my.jl	Br C. G. H. 1835.	O s.p

16756 anthraca AIT Lorrato/r viridiflora li. Rep.

16757 obtusifolia Swt.	obtuse-leaved	tf l Alpr	ft my.jl	Br C. G. H. 1825.	O «.p Sw. fl. gar. 148
16758 uncinata Swf.	honked	tf l Alpr	ft my.jl	Br C. G. H. 1825.	O s.p Sw. fl. gar. Hil

16759 divaricata Swt.

divaricate		tf my.jl	Br C. G. H. 1825.	O s.p Sw. fl. «.ar. W2
------------	--	----------	-------------------	------------------------

^ S TIGEB FLOWER. (*Tigris, tiger, eidos, likeness; spotted flowers.*) *IrMear.* 2.

16760 pavonia Jac. peacock

tf rirria Tigridia B.M.	pavonia L.	tf A or	1 my.s	O.R Mexico 1796.	O s.p Bot. mag. 532
-------------------------	------------	---------	--------	------------------	---------------------

2 lvna Hort.

16761 conchuliflora Swt.	lion	tf A	»pl	1 my.s	D.Y Mexico 1823. O s.p
--------------------------	------	------	-----	--------	------------------------

shell-flowered

16762 ovata Thun.	GALAXIA.	tf Alor	1 my.s	Galaktiao, to abound in milk.
-------------------	----------	---------	--------	-------------------------------

16763 grandiflora B. R.

16764 mucronuliflora Sal.	great-flowered	tf Alor	ft my.s	D.Y C. G. H. 1799.
---------------------------	----------------	---------	---------	--------------------

16765 versicolor Sal.

16766 graminea Thun.	various-colored	tf Alor	ft my.g	P
----------------------	-----------------	---------	---------	---

GrsM-leaved

16767 galaxia Thun.	tf Alor	I jl.au	L.Y	C. G. H. 1799.
---------------------	---------	---------	-----	----------------

0 sp Bot. mag. 129

1912. CYPELLA Hto-6. CYPELLA. (*Ki/pellon*, a kind of cup; shape of flower.) Miter. 10.
 16767 Herbert* B. M. Herbert's ff iAl or ' 1 jn.au Ve B. Ayres 1823. O s.p Bot. mag. 2699
 Tigidia Herbert* B. M. Monc'i Herbert* B. M.

* 1913. HERBERTIA Swt. HERBERTIA. (/to*. # Rev. If. Herbert, an assiduous botanist.) Irdeat*. J.
 16768 pulchella Stv. neat tf .Al ... B.p Chile 1827. O s.p Sw.fl.gar.222

1914. HOMETR/il Ven.	HOMERIA.	(Homer, the father of epic poetry.)	Irideat. 10.
16769 spicata Swt.	spiked	tf LAI or 1) myjn Y	C. G. H. 1785. O s.p Bot. mag. 1283
Morefa spicata B. M.			
16770 collina Swt.	hill	tf iAl or 2 myjn P	C. G. H. 1768. O s.p Bot. mag. 1033
Mora'a collina B.M.			
16771 exaltata Swt.	exalted	tf iAl or 3 myjn Ve	C. G. H. 1768. O s.p Bot. mag. 1103
More'i collina /3 exaltata B. M.			
16772 lineata Surf.	lined	tf IAJor 1 myjn Ve	C. G. H. 1825. O s.p Sw.fl.gar. ic ip.
16773 />orrifolia Swt.	Leek-leaved	tf iAl or 2 myjn Ve	C. G. H. 1825. O s.p Sw.fl.gar. ic. in.
16774 elegans Swt.	clegajit		
Morce'a elegans Jac.			
16775 flaccida Swt.	flaccid	tf LAI or 1j myjn Ve	C. G. H. 1823. O s.p Jac sc. 1.2
MoffeV collina « miniMa minor B.			
16776 miniflora Swt.	vermilion	tf LAJor 1J myjn Ve	C. G. H. 1810. O s.p Bot. mag. 1612
Itl777 virgkta Swt.	twiggy	tf ■ or 2 myjn Ve	C. G. H. 1799. O s.p Sw.fl.gar. 152
Mora'a virgkta Jac.			
16778 flexubsa Swt.	flexuous	tf IAI 8F 1 myjn P	C. G. H. 1825. O s.p Jac ic. 2.228
'MoreVi flexubsa B. M.		1 myjn Y	C. G. H. 1823. O s.p Bot. mag. 695

CLASS XVI. OHDEK 2.

MONADE'LPHIA PENTA'NDRIA.

1915. WALTHERN/4 L.	WALTHP.RIA.	(A. F. Walther prof. med. at Leipsic)	Byttneriucae. 5,-12.
16779 americana L.	American	tf DD un 2 my.o Y	S. Amer. 1691. S. l.p Jac. ic. 1.130
16780 indica L.	Indian	* D un 9 jn.au Y	E. Indies 1799. C. p.l Bur. zey. 68
16781 elliptica L.	elliptic	■ I un 3 " " Y	E. Indies 1812. C. Bp Cav. dis. 6.171<
16782 gibra Pair.	smooth	■ - un 3 jl Y	Guadalou.1823. C. Bp Sch. mo. 55
ls'vis Schrank			
16783 microphylla Cav.	small-leaved	* Q u n 2 jnjl Y	E. Indies 1824. C. s.l Cav.dis. & 1702

1916. HERMANNN/1 L. HERMANNIA. {Paul Hermann, prof. bot., Leyden.) Byttmridcear. 38.-42.

I. TRIONE'IXA.	— Calyx inflated, but especially after flowering filaments much dilated.		
16784 althaeifolia L.	Althaea-leaved	ft or 2* mr.jl V~"~"CTG."H. 1728. C. l.p Bot. mag. ■	
16785 plicktta W.	plaited-/cui'f/ M.	■ or 3 n.d Y	C. G. H. 1774. C. l.p Jac. sc. 2.213
16786 candicans H. K.	whifish	* LJ or 3 apjn Y	C. G. H. 1774. C. l.p Jacsc. 1.117
16787 mollis L.	soil-leaved	M J or 2 myjn Y	C. G. H. 1814. C. l.p
16788 decdmibcns IV. en.	decumbent	41 L-J " 1 myjn Y	C. G. H. 1821. C. l.p
16789 Ayyosopifolia L.	Hyssop-leaved	ft or 7 apjn St	C. G. H. 1725. C. l.p Cav. dis. 6.1M-J
16790 trifoliolate L.	three-leaved	ft or 2 iny.au Y	C. G. H. 1752. C. l.p Cav. dis. 6.18&«
16791 triphylla Cav.	three-leaved	ft LJ or 2 my.jl Y	C. G. H. 1819. C. l.p Cav. dis. 6.178.3
16792 gUndulosa Lk.	glandular ft	L-J or 2 my.jl Y	C. G. H. 1822. C. l.p
10793 fragrans Lk.	fragrant ft	J or ■ -	C. G. H. 1822. C. l.p

II. HERMANNE'LLA. — Calyx not, or scarcely inflated; filaments not dilated.

16794 discrmet6lia Jac.	two-pointed-hvdft	LJ or 2 mr.ap Y	C. G. H. 1795. C. l.p Jac. sc. 1.1'i
16795 dfsticha Schr.	two-rowed	ft i or 2 my.au Y	C. G. H. 1789. C. l.p Jac. sc. 1.118
rotundiflolia Jac.			
16796 melochioides Burc.	Melochia-like	ft i lor 2 my.jl Y	C. G. H. 1818. C. l.p
16797 Ayyonifolia Burc.	Bryony-leaved	ft or 3 my.jl Y	C. G. H. 1818. C. l.p
16798 calviblia L.	Sage-leaved	ft or apjn Y	C. G. H. 1795. C. l.p Cav. dis. 6.119
16799 milcans Schr.	glittering	ft or 2 my.au Y	C. G. H. 1790. C. l.p Jac. sc. 1.119
latiflolia Jac.			
16800 involucrata Cav.	involucred	■ i-J or 2 myjn Y	C. G. H. 1794. C. l.p Cav. dis. 6.177-1
hisot wordifolia Jac.	Germnnder-lvd	■ i-J or 2 ap.n Y	C. G. H. 1794. C. l.p Jac. sc. 1.120
18802 denudata L.	stripped	■ LJ - 2 my.jl Y	C. G. H. 1774. C. l.p Jac sc. 1.122
VMM almfolia Jac.	Alder.lcaveil	ft J or 7 fm.y Y	C. G. H. 1728. C. l.p Btt m a g J
10804 runciflilia Jac.	wedge-lenvrld	A J or 2 nu.s Y	C. G. H. 1791. ■ Lp Jaesc. 1.1J*
Itl05 holosericea Jac.	velvet-teaved	41 L-J or 2 my.jn Y	C. G. H. 1792. ■ Lp Jac sc. 3.2ip
Mitt hirsute Jac.	hAry branched	* H i or 2 my.jn Y	C. G. H. 1790. ■ Lp Sc.han.1.t
16807 scifbra Cav.	rough-leafed	* i J or 3 mr.ap Y	C. G. H. 1789. ■ Lp Jaesc.] ■
15808 inulti flora Jac.	many-flowered	m \ or 3 mir.my Y	C. G. H. 1791. ■ Lp Jacsc. ia»
16809 fluminea Jac.	JL&m&jtowercd	* or 3 year O	C. G. H. 1794. ■ Lp Bot. mag. 1^
16810 anguKiris Jac.	angular	■ or 3 ap.my Y	C. G. H. 1791. ■ Lp Jacsc. Jp"
16811 tnfurcata L.	three-forked	■ i-J or 3 apjl P	C. G. H. 1789. ■ Lp Jac. sc. 1.1»
16812 odorkta // K.	sweet-scented	■ i-J or 3 fo Y	C. G. H. 1780. C. l.p
16813 lavandulifolia L.	Lavcndr-lvd	■ or lg my.s Y	C. G. H. 1732. C. l.p Bot mag. 50*
16814 velutlna Burc.	velvety	■ or 2 my.jl Y	C. G. H. 1818. C. l.p
16815 fihiblia L.	thread-leaved	■ lor Uiny.au Y	C. G. H. 1816. C. l.p Jacsc. 1.12
16816 procimbcns Cav.	procumbent	■ or U myjn Y	C. G. H. 1792. C. l.p Cav. dis. 6.1
16817 tenuitbia A M.	slender-leaved	■ or 2 jnjl Y	C. G. H. ... C. l.p Bot. mag. K
36818 incisa W.	cut-leaved	■ or 2 jnjl Y	C. G. H. 1806. C. l.p <-.
16819 pulverullnta B. Rep.	powdered	■ or 2 my.au Y	C. G. H. 1800. C. l.p Bot rep. *o*
KJ820 argentea Sm.	silvery	■ L-J or 2 jnjl Y	C. G. H. 1820. C. l.p
16821 coronopifolia Lk.	Buckhorn-lvd		C. l.p

1917. MAHE'RNI L.	MAITERNIA.	(An anagram of <i>Herhainls</i> ; affinity.)	Byttncriace**. -l^VTTII. *
16822 verticillata L.	whorled	n\ J or 2 jn.au Y	C. G. H. 1820. C. l.p Cav. dis. & "
1683 resetifolia Burc.	Reseta-leaved	ft i or 2 jn.au Y	C. G. H. 1816. C. l.p cfr
16824 binjnpta L.	biinnate	a. i or 2 jn.au Br	C. G. H. 1752. C. l.p Bot mag.
Jnr.	cuWrA	tt. i lor 2 ilau Y.w	C. G. H. 1792. C. l.p g. J S. f ^
	diffuse	tt-u-Jor 2 jn.au Y	C. (?). H. 1774. C. l.p Bot. cat. w

16886 hirctna Swt.	<i>goat-scented</i>	■ or 10	my.o	W.a	Brazil	-	C	pi	<i>Bot.reg.321</i>
fa/tida B. M. hirsuta	<i>painted</i>	■ 15	s	R	Brazil	1820.	Q	Lp	<i>Bot.reg.67S</i>
16887 picturata JCrr.	<i>wedge-leaved</i>	■ 15	jl.au	-	S. Amor.	1825.	n.L	Cav. ic. 12.292	
16888 cuneifolia Ca>	<i>eatable</i>	■ 30	jl.au	W	W. Indies	...	E	Lp	<i>Bot mag. 1989</i>
16889 edulis B. M.									
incarnata fi B. R.									
16890 incarnata L.	<i>flesh-colored</i>	* -A Jor	30	jl.au	Pk	N. Amer.	1629.	C r.m	<i>Bot. reg. 332</i>
16891 ciliata H. K.	<i>ciliated</i>	t CDor	6	jls	Pk	Jamaica	1783.	C p.l	<i>Bot. mag. 288</i>
16892 multiflora Jac.	<i>many-formed</i>	■ 10	1820.	C p.l	
16893 caracasana W.	<i>Caracas</i>	■ 15	jnjl	Pk	Caracas	1821.	C p.l		
rubricating Jac.									
16894 tripl.irtita fo-o*	<i>tripartite</i>	■ or 14							
16895 vitifolia U \$ B.	<i>Vine-leaved</i>	■ or 10	::				S. Araer.	1823.	E p.l
i. DUPPLICALYCES.— <i>Flowers naked; calyx double; leaves three-lobed.</i>									
16896 holosericea L.	<i>velvet-tea wd & d)</i>	or 10	myau	W.p	Vera Cruz	1733.	E p.l	<i>Bot reg. 59</i>	
16897 hirsata L.	<i>hairy</i>	D or	8	B	W	W. Indies	1778.	PI. am. 88. A	
16898 glauca // X.	<i>glaucous.fazwrig</i>	Q un	6	au.s	W	Cayenne	1779.	Lp	<i>Bot. reg. 88</i>
stipulata Aub.									
le*>9 lutea L.	<i>yellow</i>	■ cu	4	my.jn	Y.w	America	1714.	R p.l	<i>Bot. reg. 79</i>
l<0 punctata L.	<i>dot-dotted-leaved</i>	■ cu	6	my.jn	W.v	Peru	1784.	C p.l	<i>Bot. cab. 101</i>
16901 rotundifolia L.	<i>round-leaved</i>	■ cu	8	my.au	W	W. Indies	1779.	C p.l	<i>Cav. dis. 10.290</i>
16902 hemicycla Mey.	<i>half-round</i>	■ cu	10	...		Surinam	1817.	C p.l	
glabrata Kth.									
16903 racemosa Brot.									
2 alata cscrlea B. R.	<i>blue-winged</i>	■ cu	20	mr.o	S	Brazil	1815.	C p.l	<i>Bot mag. 2001</i>
3 princeps Hort.	<i>chief</i>	■ cu	15	apjl	Br	hybrid	1823.	C p.l	<i>Bot reg. 848</i>
ii. SIMPLICALYCES.— <i>Flowers naked; calyx simple; leaves three-lobed.</i>									
16904 gracilis Lk.	<i>slender</i>	O cu	6	au	W	1823.	O co	<i>Bot. reg. 870</i>	
16905 auberti Ra L.	<i>Cork.barked</i>	\$ CUCu	6	jns	W	W. Indies	1759.	C p.l	<i>Ex. bot. 1.28</i>
16906 Aederacea L.	<i>Ivy-leaved</i>	LDun	3	jnjl	W	W. Indies	...	1p	<i>Pl. am. 84</i>
16907 angustifolia Sco.	<i>narrow-leaved</i>	t Clcu	6	jns	W	W. Indies	1773.	1P	<i>Bot. reg. 188</i>
heterophylla Jac.									
16908 obscura Lndl.	<i>obscure</i>	LDcu	6	jl.au	W.a	W. Indies	1823.	E Lp	
16909 minima L.	<i>smallest</i>	LDcu	6	jl.au	W	Curasao	1690.	E Lp	<i>Bot reg. 144</i>
16910 peltata Cav.	<i>peltate</i>	LDcu	6	au.8	G	W. Indies	1778.	E Lp	<i>Bot reg. 507</i>
16911 maculata Bonn	<i>spotted</i>	LDcu	4	jno	Ap	N. Amer.	1812.	E p.l	
plata Ph. not of others.									
16912 iridi&ntum W.	<i>Maidenhair</i>	% r-l or	10	—		Norfolk	1.1820.	E p.l	
16913 ftryoniaefolia H.S.B.	<i>Bryony-leaved</i>	J. Q or	12	—		Mexico	1823.	E p.l	
16914 caculta L.	<i>common blue</i>	■ or	30	jno	W.B	Brazil	1699.	E s.p	<i>Rot. mag. 28</i>
2 ca?riMeo-racemosa	<i>blue-clusters</i>	■ or	30	jno	P	hybrid	J8J0.	E co	<i>Hort. tr. 4.9</i>
3 angustifolia	<i>narrow-leaved</i>	■ or	30	jno	W.B	hybrid	1820.	E co	
4 glauropa Jlla	<i>glaucous-leaved</i>	■ or	20	jno	B	Brazil	—	E co	
5 Colvillw Swt.	<i>Colvillw hybrid</i>	& or	12	jls	Va	hybrid	1824.	E co	<i>Sw. fl.gar. I¹</i>
16915 chinensis Swt.	<i>Chinese</i>	% or	30	jno	W.B	China	—	E co	
16916 filumentosa Cav.	<i>thready</i>	■ or	20	jno	W.B	America	1817.	C co	<i>Bot. reg. 584</i>
palmata B. C.									
16917 scribula L.	<i>saxr.leaved</i>	^ or	20	...	W.o	Martinq.	1800.	C Lp	<i>PL am. 79</i>
16919 digitrita G. Don	<i>digitate</i>	* D or	12	...	B	Trinidad	1820.	C p.l	
16919 ixifolia L.	<i>pedate</i>	S O or	15	...	W.a	W. Indies	17H1	C p.l	<i>PL am. 81</i>
16920 palmata Lk.	<i>palmate</i>	CD or	18	...	W	Brazil	1818.	C p.l	
16921 heterophylla Lam.	<i>various-leaved</i>	% or	15	...	W	St.Domin.	1817.	C p.l	<i>PL ic. 139,1</i>
V. Du^ffii A.— <i>Doubtful to what section it belongs.</i>									
16922 p&tula Wnl	<i>spreading</i>	\$ * or 11	—	—			C	E	
1924. MURUCIPIAJ.	<i>MUBUCUIA.</i>	(Murucyja, its name in Brazil)					Patsifihrece.	4.—10.	
16923 ocellata Pert.	<i>smnll-eyed</i>	j _ D fr	12	jl.au	S	W. Indies	1730.	C p.l	<i>Bot. reg. 574</i>
Passi flora Murucuia L.									
16924 perfoliata Spr.	<i>perfoliate</i>	t, • or	15	jl.au	P	W. Indies	1806.	C p.l	<i>Bot reg. 78</i>
Passiflora perfoliata L.									
\$16925 Herbertia Vina Swt.	<i>Herbert's</i>	J. • or GO	jl.s	G	N. Hol.	1821	C p.l	<i>Bot reg. 737</i>	
Passi flora Herberti(iae B. It.									
% 16926 adiantifolia Swt.	<i>Adinnumt-lvd</i>	E. • or 20	jn.au	O	Norfolk	1.1792.	C p.l	<i>Bot reg. 533</i>	
Passiflora tzdiantifolia B. R.									
1925. TACSONIA3.	<i>TACSONIA.</i>	(Tacsq, its name in Peru.)					Passiflbre*.	I.—1*..	
16927 peduncularis Pers.	<i>pedunculate</i>	t_ O or	10.	.	Peru	1815.	C Lp	Cav. ic. 5.4*	
Passi flora ieduncularis Cav.									
1926. TRIOONIA Aub.	<i>TRIOONIA.</i>	(Treis% three, gonia, an angle; fruit)					JUppocratehcea.	2.	
16928 villbsa Aub.	<i>villus</i>	■ O or	.	.			Cayenne	1820.	C s.l
16929 melliis Dec.	<i>soft</i>	■ Qor.	.	.	Brazil	1823.	C s.l	Aub. gui.	
* 1927. EKTOIUM Herit. HERON'S BILL. (<i>Erodios</i> , heron; fruit like head and beak.) <i>Gerantaceae?</i> 35.'^									
I. PINNATISK'TA. — <i>Leaves pinnately parted.</i>									
16930 petraAim W.	<i>rock</i>	^ A	i	jnjl	P	S\ Europe	164a	D e o	Gou.il. 21. ¹
16931 anthemidif Tilium Bibb.	<i>Anthc.-lvd</i>	^ A or	i	jnjl	P	Iberia	18W.	C Lp	
16932 glandulsum W.	<i>glandular</i>	■	■	i	P	Spain	1798.	S Lp	Fl pyr. 1. ¹
gravolens Lap.									
16933 Stephanianum W.	<i>Stephan's</i>	■ A or	i	jnjl	B	Dahuria	182a	S s.p	
multifidum Pat.									
16934 alp'num Lk.	<i>alpine</i>	k	Apr	myau	R	Italy	1814.	D co	Her.ger.3.
16935 crassifolium Desf.	<i>thick-leaved</i>	f	A Jor	imr.au	S	Cyprus	1788.	D r.m	Sw. ger. I ¹ .^
16936 laciniatum Cav.	<i>fringed</i>	f	A Jor	i mvau	R	Crete	1794.	J> r.m	Cav. dis.*.**.
16937 stylatum W.	<i>hairy</i>	3	A or	**	B	Egypt	1818.	S s 1	Jac. ec. J. ^
IVMII muricatum Hort.	<i>styled</i>	A or	**	P			1826.	S s.1	
IVMII cibinium W.	<i>prickly</i>	8	or	i jnjl	R				
16938 cibinium W.	<i>stork-beaked</i>	8	un	N jnjl	P	S. Europe	1711.	IK27.	S s i . i 18
16939 cicutarium Lem.	<i>Hemlock-leaved</i>	W	Japs.	P	Britain	ro.sid.	S co	Jac. vm. »^gj	
16940 cicutarium W.	<i>bipinnate</i>	O un	myn	P	Numidia	1803.			
JSS? bipinnatum W.	<i>Cav. Kurnet-leaved</i>	O or	f jnjl	P	S. Europe	1800.			
	<i>pilose</i>	O or	I jnjl	D.P	France	1800.	S co		
16945 cicutiflum Thuil.	<i>Cicuta-leaved</i>	O or	f jnjl	P	France	1816.	S co		

Sig^rm&numW'.	Roman	£ CDpr	\ my.jn P	Rome	1724	S CO	Eng. bot 377
>^r caucalifbium Swt.	Caucalis-leaved [^]	A pr	1 my.au P	France	1816.	S CO	Sw. ger. 6
tun *ipinum Desf.	musk	O pr	1 my.jl P	England	m.pas.	S 8.1	Eng. bot. 902
SSJ K S " ! *	Botrys	O or f	jnjl B	Italy	1818.	S CO	Cav. dis. 4. 90. 2
iBj. n^	II. INDIV' SA.—Leaves somewhat three-lobed or undivided.						
1659 T. num W.	Murcian	Q or	1 jnjl B	N. Africa	1818.	S CO	Cav. dis. 4.126.1
169592 J. serotinum Stev.	crane	un	i in.jl P	Crete	1596.	S 8.1	Cav. dis. 4. 88.2
M>J2 serotinum Stev.	late	& Q pr	Jl.s B	Siberia	1821.	D CO	Sw. ger. 137
inssa, S Uica Si 2Ar-	nitMplcum <*						
S J TM A	Chian						
S C U T. num Ton.	hymen-like	£ Q pr	1 jnjl P	Levant	1724.	S CO	Cav. dis. 4. 92.1
PSRS of. num w.	Gousson	£ A or	f year	Barbary	1789.	S r.m	Sw. ger. 23
16017 Sfeff. num Jac.	Mallow-like	O or	i my.jl B	Naples	1821.	D CO	
16058 nS A^& n^ Jac.	currant-leaved	O or	1 jnjl P	S. Europe	1596.	S CO	Cav. dis. 4. 91.2
16959 32 E * S miffa *	flesh-co/orcrf ft.	Q or	i my.jl F	C. G. H.	1787.	C r.ra	Jac. ic. 3 509
© KEX? R V illum **& glaucous-lvd	O un	i l.lau R	P	Egypt	1732.	S CO	Di. el. 124.150
2 ^ & \dvs V'	Malope-like	£ A or	% jnjl P	N. Africa	1810.	S CO	Cav. dis. 4.90.1
16061 12 2 S? m Len Z'	Corsican	△ A or	* jnjl P	Corsica	1817.	S CO	
16962 S a S? Mac black-sti Pna		△ Al or	1 jnjl P	1823.	S CO	
16963 SS? V? Ma		& A w	imy.S F	England	san.sh.,	D CO	Eng. bot. 646
files	Richard Vi	△ Al pr	I ap.s W	Minorca	1783.	C 8.1	Bot mag. 18
169641 JT^ nro des Herit.	Reichardi Mur.	& A un	* ap.s R	& Europe	182L	D CO	
**>* utt&reum Z*m.	shore						

CLASS XVI. ORDER 3.

MONADELPHIA HEXA'NDRIA.

LindL	GILLIESIA.	(Dr. Gillies, of Concepcion, in Chile.)	Gilliestite.	I.—2.
grassy	tf	J c u 1 aus	G Chile	1825. O 8.1 Bot mag. 2716

CLASS XVI. ORDER 4.

MONADE'LPHIA HEPTA'NDRIA.

16W% i	stalked leaves ; leaves oblong, entire, or lobed ; lobes entire, or scarcely toothed: from No. 16997.						
16967	cordate, 3-lobed, or at base appendic.	; from No. 16997. <i>Ivs pinnate, segments cut or multifid.</i>					
B rain Jac.	long-leaved	A IAJ pr	i my.jn Pk	C. G. H.	1812.	R r.m	Jac. ic. 3.518
ovahfbium Swt.	flowered	A iAl pr	f my.jn Y	C. G. H.	1812.	R r.m	Jac. ic. 3.521
16968	oval-leaved	C. AiAlpr	imy.jn W	H.	1820.	R r.m	Sw. ger. 106
16969	netted	A iAlpr	f my.jn Pk	C. G. H.	1820.	R r.m	Sw. ger. 91
16970	ciliated	A iAlpr	iapjn F	C. G. H.	1795.	R r.m	Her. ger. 7
16971	lar#-rooted	A iAlpr	t ap.my Y	C. G. H.	1794.	R r.m	Bot. rep. 60
16972	stul&tum Andr.	A iAlpr	* jnjr Y	C. G. H.	1802.	R r.m	Bot mag. 1718
16973	i Andr.	A iAlpr	i ap.my Y	C. G. H.	1795.	R r.m	Bot. rep. 152
16974	i Pers.	A iAlpr	i ap.my Y	C. G. H.	1795.	R r.m	Bot. rep. 282
16975	virgin	A iAlpr	i jn.au Y	C. G. H.	180L	R r.m	Bot. rep. 222
16976	rave-flowered	A IAJpr	Jmy.jl Y	C. G. H.	1795.	R r.m	Bot. rep. 317
16977	lino^re Pers.	A IAJpr	Y	C. G. M.M.	1795.	R r.m	Bot. rep. 292
16978	nfvrum Swt.	A LAJpr	* jnjl Y	C. G. H.	1795.	R r.m	Bot. rep. 193
16979	snow-	A LAJpr	* jnjl W	Y	1800.	R r.m	Bot. rep. 182
16980	Leebum Swt.	A IAlor	* jnjl W	Y	1821.	R r.m	Sw. ger. 3
16981	irrassatum Andr.	A IAlor	i my.jn W	C. G. H.	1823.	R r.m	Sw. ger. 1
16982	cirinatum Swt.	A IAlor	i myjn R	C. G. H.	1801.	R r.m	Bot. mag. 761
16983	ilprslssum Jac.	A IAlor	i myjn Y	C. G. H.	1812.	R r.m	Sw. ger. LISS
16984	depressed	A IAlor	^ myjn Y	C. G. H.	1812.	R r.m	Sw. ger. 1
16985	yellow	A IAlor	i myjn Crea	C. G. H.	1812.	R r.m	Jac. ic. 3.520
16986	i fiteum Andr.	A IAlor	* myjn Y	C. G. H.	1802.	R r.m	Bot. rep. 123
16987	i niimmarifolium&i. Nummul-lvdAit.	A IAlor	* myjn Y	C. G. H.	1801.	R r.m	Par.kmTsf
16988	erobifbium Swt.	Orobos-leaved	A IAJ pr	i my.jl Bd	hybrid?	1824.	R r.m
16989	undulcrflbrum Swt.	wave-flowered	A iAl or	i my.jl P	hybrid?	1824.	Sw. ger. 3M
16990	cong&tum Swt.	crowded	A iAI pr	* my.jl Li	C. G. H.	1821.	R r.m
16991	retusc	A iAI pr	i my.jl DC	C. G. H.	1824.	it	Sw. feftSW
16992	r liuteaflium Swt.	Colutea-leaved	A IAJ pr	i my.jl YR	hybrid?	1824.	R r.m
16993	li volituum Pers.	revolute	A iAI pr	* jlau P	hybrid?	1824.	11 rim
16994	" i uiricilatum IV.	car-leaved	A iAlpr	* dp.jn Pk	C. G. H.	1800.	Bot. rep. 354
16995	Urbamniflium Pers.	ja^-leaved	A IAJ pr	t myjn Pk	C. G. H.	r.m.	Jac. ic. 3.519
16996	nd-sorrel/crf	A IAlpr	i my.au Y	C. G. H.	1800.	R r.m	Bot. rep. 131
16997	nervifolium Jac.	nerved-leaved	A IAlpr	i my.au Va	C. G. H.	180L	R' r.m
16998	WMS triphyllum Jac.	thrcclcnved	A iAlpr	i my.au Y	C. G. H.	1812.	R r.m
16999	rtflxum Pers.	retex-fruved	A IAJ pr	ap.my Y	C. G. H.	1812.	Jac. ic. a 517
17000	roset-colored	A IAl pr	* jnjl W	C. G. H.	1812.	R r.m	Bot. rep. 224
17001	roset-colored	A IAl pr	I mr.my Pk	C. G. H.	1800.	R r.m	Bot. rep. 173
17002	rapaccum Jac.Rape	Funitory-flwd	A . Alpr	i apjn Pk	C. G. H.	1792.	R r.m
17003	nodding	A . Alpr	i apjn E	C. G. H.	1788.	R r.m	Bot.rep.2J9
17004	corydaliflbrum Swt.	Corydalis-flwd	A IAJ pr	Japjn	& T&SgSaS2!£&	1817	
17005	barbatum Jac.	bearded	A IAI pr	I Jlau			
17006	fissiflbum Pers.	clove-leaved	A IAJ pr	ap.au	Pk		
17007	set^sum Swt.	bristly	A IAI pr	t ap.au	Pk		
17008	AubomfTiliun Pers.	Bubo-leaved	A IAJ pr	I mr.jl W	P		
17009	viola?flbrum Swt.	Violet-flowered	A IAJ pr	I mr.jl W	P		
17010	floribfindum Ait.	bundle-flwd	A IAJ pr	i mr.my Pk	C. G. a	1810.	H r.m
17011	pilose	A IAJ pr	i my.jl Pk	C. G. a	1810.	it "	Bot. rep. 420
17012	winged-formed	A iAI pr	iniyjn Y	C. G. a	1800.	S rm	500 r ep. 259
				C. G. R	1800.	R r.m	Bot rep. 2

- | | | | | | | | | | | | | |
|-------|----------------------|--------------|----------------|----------|---|--------|-------|----|-------|--------|------------------|-----|
| 17009 | <i>purpureascens</i> | <i>Pers.</i> | purpuraacnt | AiAJp* | i | my.jn | P | C. | G.H. | 1800. | It r.m not rep. | 204 |
| 17010 | <i>hirsutum</i> | <i>Jac.</i> | hairy | A LAJ pr | I | mr | Pk | C. | G. H. | 17H8. | It r.m Hot. rep. | 317 |
| 17011 | <i>melananthum</i> | <i>Jac.</i> | black-flowered | A LAI P* | I | my.jn | D.Br. | C. | G. H. | 17L10. | It rm Sw.gr. | 73 |
| 17012 | <i>dioicum</i> | <i>Ait.</i> | dioecious | A LAI pr | I | jn.jl | D.Br. | C. | G. H. | 179**. | It r.m Hot. rep. | 209 |
| 17013 | <i>atrum</i> | /frnt | dark-frown | A LAJ pr | I | iny.jl | D.Br. | C. | G. H. | 1793. | K r.m Sw.gr. | 72 |

J. DIMA*CRIA. Petals 5, unequal, two upper connivent spreading at end: stamens shorter than sepals; 5 fert U& stemless herbs

III. CYNO'SBATA.—Petals oval, nearly equal, almost twice as long as calyx; stamens 10, erect, the 5 alternate ones bearing the anthers; stems shrubby, erect.

- 17035 lateritium W. brick-colored t*-Ljor lljn.au R C. G. H. 1800. C r.m Jac. cc. 1.97
 17036 cynosbatifolium W. Eglantine-lyd tt-Ljor 1J apjl D.R C r.m W. h. b. 2.78

IV. PERISTERA.—Petals nearly equal, as long as calyx, or a little larger; stamens 10; 5 longer, nearly equal* or one only occasionally abortive; 5 alternate, very short, sterile, tooth-like: herbs with stems, and with the appearance of *Erodium* or *Geranium*.
172. *Peristera* ^{flaviflora} Benth. sp. n.

- | | | | | | | | |
|---------------------------------|-----------------|---|---------|---|----------|-------|----------------------|
| 17037 colum/knum <i>Jac.</i> | dove's foot |  | £ jn.o | P | C. G. H. | 1795. | R r.m Jac. sc. 2.133 |
| 17038 procumbens <i>Pers.</i> | procumbent |  | 4 ap.my | P | C. G. H. | 1801. | S r.m Bot. rep. 234 |
| 17039 numifaim W. | trailing |  | i my.jl | P | C. G. H. | 1801. | S r.m Sw.ger.42 |
| 17040 chamaedryfulum/Jac. | Chaimedrys-lvd^ | LAJ pr | i my.jn | W | C. G. H. | 1812. | R r.in Jac ic.3. 528 |
| 17041 austral'e W. | southern | tt. | * my.au | R | N. S. W. | 1792. | S r.m Jac.cc. 1.100 |
| 17042 althaioides <i>Herit.</i> | Althaea-like | f OJ pr | 4 ap.jn | W | C. G. H. | 1724. | S r.m Jac. c. 4.21.2 |

i. *GRENVILLEA*. — Petals 5; 2 upper ones large, obliquely obovate, 3 lower ones small, spatulate: stamens V-**
4 fertile, ascendent, b sterile, short, subulate.

- | | | | | | | | | |
|---|---------------------|------------|------------|-----------|-----------|--------------|-------------|--|
| <i>Geranium Grenifloc</i> Andr. | | | | | | | | |
| V. OTT'DIA.—Petals oblong, linear, nearly equal, about twice as long as calyx, the two upper auricled *****, base on the upper side; stamens 10, erect, & fertile, 2 upper spatulate or subulate, Slower shorter; stew* shrubby, fleshy; leaves alternate pinnated, fleshy; flowers whitish,* | | | | | | | | |
| 17044 <i>laxum</i> Swt. | <i>Jooapanicled</i> | <i>tt.</i> | <i>i</i> | <i>or</i> | <i>pr</i> | <i>1</i> | <i>apin</i> | <i>W</i> |
| 17045 <i>ceratophyllum</i> Herit | horn-leaved | <i>tl.</i> | <i>-</i> | <i>or</i> | <i>1</i> | <i>my.jn</i> | <i>W</i> | <i>Africa</i> |
| 17046 <i>dasyactylon</i> Sims | thick-stemmed | <i>tt.</i> | <i>J</i> | <i>or</i> | <i>1</i> | <i>jl.d</i> | <i>W.p</i> | <i>C. G. H.</i> |
| 17047 <i>crithmifolium</i> Sm. | Samphire-lvd | <i>tt.</i> | <i>i</i> | <i>or</i> | <i>1</i> | <i>my.jn</i> | <i>W.p</i> | <i>C. G. H.</i> |
| 17048 <i>iltcrmans W.en.</i> | alternate Pars.-lvd | <i>tt.</i> | <i>»-1</i> | <i>or</i> | <i>1</i> | <i>my.au</i> | <i>W.p</i> | <i>C. G. H.</i> |
| 17049 <i>carnosum</i> Ait. | five-hy-stalked | <i>n.</i> | <i>—</i> | <i>or</i> | <i>1</i> | <i>in.au</i> | <i>W.p</i> | <i>C. G. H.</i> |
| | | | | | | | | <i>1821. S p.I Sw. ger. 196</i> |
| | | | | | | | | <i>1786. C r.m Bot. mag. 315</i> |
| | | | | | | | | <i>1795. C r.m</i> |
| | | | | | | | | <i>1790. C r.m Sm. ic. pic. 1&</i> |
| | | | | | | | | <i>1791. C r.m Wen. her. 2.9</i> |
| | | | | | | | | <i>1724. n. r.in Sw. ger. 98</i> |

V.1. POLYACTIUM. — Sepals nearly equal, revolute; petals 5, nearly equal, obovate : stamens 10 ; fertile, the lower long, subulate, upper broad, spatulate, reflexed at end ; the fertile ones shorter, incurved at end: petal with a very large dark brown spot, which is scarcely edged with yellow.

- 17050 multiradiatum W. en. many-rayed A IAJ pr 1 my.jn D.Br C. G. H. 1820. R r.m Sw. ger. 145

VII. ISOPE TALIJM.—Upper sepal ending in a honey pore and not in a tube: petals 5, equal: united in a very short cup; 5 or 6 fertile, spreading, incurved at end; sterile unequal, subulate, incurved: i with a fleshy stem.

- VIII. CAMPY'LIA. — Petals 5, unequal, two upper larger, with an ariculated claw: stamens 10, hairy or f^lcent; 5 fertile, erect; 5 alternate steele, of which the two upper are longer and hooked back: herbs at tip a little shrubby, branched: leaves stalked, ovate or tiblong, toothed or cut: petals with an appendage claw.

17052 <i>blatthrium Jac.</i>	Moth <i>Muleync</i> tt. j_ or woolly-stmnd tt.	j_ or or	jn au mr.ju	V W	C. G. H. C. G. H.	1720. 1794.	S.r.m Sw. ger. 88 C.r.m Jac. sc. 2.1*
17(W3 eriost-mon <i>Jac.</i>	velvety tt.			D.P.	C. G. II.	1820.	C.r.m Sw. ger. 75
170.H <i>holoscrecum Swt.</i>							

- | | | | | | | | | | | |
|----------------------------------|-------------------|--------|----|----|-------|--------|----------|----------|------------------|-------------------------------|
| 1705' OftioltuRc <i>Jar.</i> | CEnothera/A-c | £ | A | or | 1 | nirjii | Pk | C. G. H. | 1812. | S r.m J. ⁴ ic.3.ag |
| 1705' coronopifolium <i>Jac.</i> | IUickhorn-lvd | tt. | | or | 1 | jn.o | P.v | C. G. H. | 1791. | S r.m Bot. rep. 3» |
| 1705' c'uuniu <i>Pers.</i> | hoary | a. | | or | 1 | jn.o | Pk | C. G. H. | 1820. | S r.m Sw. ger. »• |
| 1708 carinatum <i>Swt.</i> | keeled | tt. | LJ | or | If | jn.o | W.p | C. G. H. | 1820. | S r.m Sw. ger. »• |
| 17059 verbascifl̄rum <i>Swt.</i> | Verbsjcum-flwdtt. | L J pr | | 1 | jn.au | Li | C. G. H. | 1811. | S rm Sw. ger. »• | |

i. PIYIMATA'NTHUS. — *The same as piycedding, hut with the upper petals warded above the rim, Subsp. very short, 5 fertile, recurved, spreading, 5 sterile, straight.*

- 17060 tricolor B. M.
17061 datum Swt.

IX. MYRRHIDIUM. — Petals* er rarely 5, the two upper very large, obovate, cuneate, usually branching lines; two or three lower much smaller, oblong, linear: stamens 10, with their tube straight, generally with nantheis scrouns 5 barren alternately, rarely *jeſatite*: *bhāntūlōor* *H* *H* *is* *i* rarely shrubby; stems round; leaves pinnate or temate, often multifid. From No. 17001, anthers o,*

- | | | | | | | | | |
|---|-----------------|---------|--------|----------|------|----------|--------|------------------|
| <i>rarely shrubby</i> , stems round, leaves pinnate or palmate, often matifolia. From No. 17061. 17064. anthers 5, p
from No. 17064. anthers 5, petals 5; from No. 170dJ. anthers 7, petals 4. | | | | | | | | |
| 17082 canariense JV. | Canary | tt. | [—] pr | 1^ jls | W.R | Canaries | 1802. | C r.m W.h.b 17 |
| 17163 mynhijdium Ait | Myrrh-leaved | tt. | [—] pr | 1 my.au | W.R | C. G. H. | 1696. | C r.m Jac. < |
| 17064 coriandrifolium jfct. | Coriander-lvd | £ OJ | pr | I mr.s | W.R | C. G. H. | 1721. | S r.m Sw. g. |
| 7.Oia lucerum Jac. | torn-leaved |) £ iAJ | or | 1 jn.au | Pk | C. G. H. | 1731. | S r.m Jacie- |
| VJ*** lucernif()hum Jac. | Anemone-lvd | ff Ol | or | 1 n.au | Pk | C. G. H. | | S r.m Jac ic 335 |
| 17067 eaucalift. myn. Jac. | Caucalis-leaved | OJ | pr | I mr.s | Pk | V. G. H. | 181*2. | S r.m Jac. ic |
| 17068 multicaule Jac. | matif-Rtalkei* | £ O' pr | * | jnau | Pa.V | C. G. H. | 1802. | S r.m Jacie |

S zonUum Swt.	zoned	tt.	uJ or	2 my.n	Pa.P C. G. H.	1787.	C r.m
3 roseum Swt.	rose-colored	tt.	i_I or	2 myn	R C. G. H.	1787.	C r.m
4 álbo-marginatum Ztec.	white-raarg.n.	i_	or	2 my.n	Pa.R C. G. H.	1787.	C r.m And. ger. ic.
5 peltatum Ait.	peltated	tt.	L_J or	2 jn.au	P C. G. H.	1701.	C r.m Bot. mag. 20
2 variegatum Andr.	variegated	-uJor	2 jn.au	P C. G. H.	1701.	C r.m And. ger. ic.	
iii. PLATVPE TALA.—Two upper petals broader and shorter, and very obtuse.							
17129 ovale Herit.	oval-leaved	tt. Jor	1\$ my.jl	p C. G. H.	1774.	C r.m Her. ger. 28	
17130 llegans W.	elegant	tt. ^J or	3 mr.Jn	W.R C. G. H.	1795.	C r.m Bot. rep. 28	
2 minus Surf.	araallery.Jtd	tt. L_J or	3 my.au	W.R C. G. H.	1795.	C r.m Sw. ger. 3. 232	
3 majus Swt.	larger-flowered	tt. L_J or	3 my.au	W.R C. G. H.	1795.	C r.m Sw. ger. 1.36.6	
iv. ANISOPB TALA.—Two upper petals longer and broader j stems shrubby.							
ü GLAUCESCE'NTIA.—Leaves smooth or nearly so, more or less glaucous.							
* Petals white, upper ones most often lined or spotted with red.							
17131 glauicura Herit.	glaucous-leaved*	tt. qor	3 jn.au	W.R C. G. H.	1775.	C r.m Sw. ger. 235	
17132 diversifolium	different-leaved	tt. qor	3 jn.au	W.R C. G. H.	1794.	C r.m	
17133 suspicuum Whl.	suspicious	tt. qor	3 jn.au	W.R C. G. H.	-	C r.m	
17134 Bororum H.,	sister	tt. -	3 jn.au	W.R C. G. H.	-	C r.m	
17135 laevigatum W.	smooth tcrn.-lvdn.	tt. -	3 ap.jl	W.R c. G."H.	-	C r.m Cav. dis. 4.121>1	
17136 grandiflorum W.	great-flowered	tt. -	3 my.au	W.R C. G. H.	1794.	C r.m Sw. ger. 29	
17137 variegatum W.	variegated-JhtM/t.	tt. -	3 ap.jl	W.R C. G. H.	1812.	C r.m Cav. dis. 4.118.3	
** Petals red or violet, upper ones striped with purple.							
17138 patulum Jac.	spreading	tt-i Jor	3 ap.jl	P.bd C. G. H.	1821.	C r.m Jac. ic. 3.541	
17139 saniculifolium W.	Sanicle-leaved n.	i_ el	3 jn.au	PR C. G. H.	1806.	C r.m Jac. ic. 539	
17140 fijscatum Jac.	clouded	tt. i_J el	3 ap.jl	P.R C. G. H.	1812.	C r.m Jac. ic. 3.540	
ii. LINEA^TA.—Flowers white, seldom red; two upper petals intensely lined with red; leaves ovate, cordate, or							
rentform, toothed, or undivided.							
17141 penicillatum W.	pencilled	tt. uJ el	3 jn.au	W.R C. G. H.	1794.	C r.m Bwt. lmag. 1148	
17142 Aetblinum AH.	Kirch-leaved	tt. i_J el	3 jn.au	W.R C. G. H.	1759.	C r.m Sw. ger. 215	
17143 formosissimum Pers.	handsomest	tt. (el	2 jn.au	W.R C. G. H.	...	C r.m Sw. ger. 3.215.6	
2 linettum Swt.	lineate-flowered*	tt. -	2 my.jl	W.R C. G. H.	1794.	C r.m	
m. TOMENTOSA.—Petals white, narrow; leaves cordate, soft, tomentose; stipules spreading.							
17144 tomenttaum Jac.	woolly	CL^Jor	3 jn.jl	W C. G. H.	1790.	C r.m Bot. mag. 518	
17145 rito/fulium Jac.	currant-leaved	tt. i_J or	3 my.jn	W C. G. H.	1798.	C r.m Jac. ic. a 538	
iv. PAPILIO.VAYEA.—Leaves cordate, flat, toothed i lower petals linear, upper ones lined with purple.							
17146 papilionaceum Ait.	butterfly	tt. LJ or	3 ap.jl	P.W C. G. H.	1724.	C r.m Sw. ger. 27	
17147 cordatum Ait.	heart-leaved	tt. i_J or	3 mr.jl	P.W C. G. H.	1774.	C r.m Bot. mag. 165	
17148 rubrofncutum Lk.	red-edged	tt. V_J or	3 mr.jl	P.w C. G. H.	1774.	C r.m	
17149 conduplicatum W. doublcd-hrt-lvdn.	i_J or	3 mr.jl	P.w	1774.	C r.m		
v. TuvrRASCEfSTia.—Leaves cordate or cuneate, toothed, undivided, or sublobatej lobes obtuse s flowers							
purple; lower petals oblong or obovate.							
* Leaves undivided, cuclulate,from No. 17156. lobed.flat.							
17150 culculatum Ait.	hooded-leaved tt.	i_ or	3 mr.jl	C. G. H.	3690.	C r.m Cav. dis. 4.106.1	
2 stratiiflorum Swt.	stratified-flwd	tt. -	3 mr.jl	C. G. H.	1810.	C r.m	
3 majus Swt.	greater Royal Geo.	tt. -	3 mr.jl	C. G. H.	1812.	C r.m	
4 grandiflorum Swt.	great-flowered	tt. -	4 mr.jl	C. G. H.	1818.	C r.m	
Sparviflorum Swt.							
17151 specibum W.	specious	tt. -	4 ap.jl	C. G. H.	1794.	C r.m	
17152 cochleatum W.	cochleate-frtf	M. I or	3 mr.jl	C. G. H.	-	C r.m	
17153 acenfbium Herit.	Maple-leaved	tt. -	3 ap.my	P. C. G!"H.	1784.	C r.m Her. ger. 21	
17154 angulsum Ait.	angulose	tt. uJ or	3 jl.s	C. G. H.	1724.	C r.m Cav. dis. 4.112.2	
17155 Barrington's* W.	Barrington's	tt. i_J or	3 mr.jl	C. G. H.	-	C r.m	
17156 Watsonia Lk.	Watson's	tt. i_J or	3 mr.jl	C. G. H.	-	C r.m Sw. ger. ISO	
17157 adulterinum Herit.	counterfeif	tt. J or	3 ap.jn	C. G!"H.	1785.	C r.m Sw. ger. 22	
17158 semirilobum Jtc.	half-thre-lobed	tt. -	3 ap.jl	C. G. H.	1800.	C r.m Jac. sc. 2. 1.239	
17159 vitifoliuum Ait.	Vine-leaved	tt. -	3 ap.au	C. G. H.	1724.	C r.m Cav. dis. 4.111.2	
17160 capitatum Ait.	CAnt&teRose-sctdLi	J or	3 ap.au	CG. H.	1690.	S r.m And. ger. c. ic.	
17161 rrbens W.	red-flowered	<. i_ or	3 my.jl	CG. H.	-	C r.m	
vi. CM'SPA.—Leaves lobed, lobes at top acutely toothed.							
17162 obtusiflbium Ait.	blunt-leavedtt.	tt. J	3 ap.au	P. -	-	C r.m Sw. ger. 8	
17163 tricuspidatum Herit.	three-pointed	tt. f or	3 my.au	W.p C. G. H.	1780.	C r.m Her. ger. 30	
17164 scabrum Ait.	rough wedge-lvd.	i_J or	3 ap.au	W.R C. G. H.	1775.	C r.m Jac. ic. 3.542	
17165 hermannia/o/litmJac.	Hermannia ia-lvdt.	J or	3 ap.jn	Pk C. G. H.	-	C r.m Jac. ic. 3.545	
17166 ericipum AU.	curl-leaved	n. i_ or	3 jl.n	P. C. G. H.	1774.	C r.m Her. ger. 32, 33	
17167 exstipulatum Ait.	exstipulate	tt. -	3 my.au	Vi C. G. H.	1779.	C r.m Her. ger. 35	
17168 pustulsum Swt.	pimply	tt. J or	3 my.au	W.pk C. G. H.	1820.	C r.m Sw. ger. 11	
17169 p&lidium IV.	pa)c-flowered	tt. J or	3 ap.au	Pk. -	...	C r.m	
17170 ternata	ternate	tt. J or	3 ap.au	Pk. C. G. H.	-	C r.m	
vii. RA'DULA.—Leaves beyond the middle parted; lobes toothed, phunntifld r flowers purplish or pale.							
17171 guercifbhum Ait.	Oak-leaved	tt. i_J or	3 mr.au	P. C. G. H.	1774.	C r.m Her. ger. 14	
2 bipinnatifidum Suit.	bipinnatifid	* J or	4 mr.au	P. C. G. H.	1774.	C r.m Her. ger. 15	
17172 graveolens Ait.	strong.rose-sctd	i_J or	3 mr.jl	P. C. G. H.	1774.	C r.m Her. ger. 17	
2 variegatum Swt.	variegated-vrf	LJ or	3 mr.jl	P. C. G. H.	-	C r.m And. ger. ic.	
Geranium capitatum var. Andr.							
17173 glutinosum Ait.	glutinous	tt. tt. d_J or	3 my.jn	Pk.w C. G. H.	1777.	C r.m Bot. mag. 143	
17174 nigrlscens Swt.	nigrescent	ft L_J or	4 my.jn	P. C. G. H.	1777.	C r.m	
17175 hlpidum W.	hispid	tt. J or	4 my.jl	P. C. G. H.	1790.	C r.m Cav. dis. 4.110.1	
17176 rädula Ait.	rasp-leaved	tt. L_J or	4 mr.jl	P. C. G. H.	1774.	C r.m Bot. mag. 95	
17177 balbameum Jac.	balsamic	tt. L_J or	4 jl.s	Pk C. G. H.	1790.	C r.m Jac. ic. 3. £43	
17178 aspernum IV.	rough multifid	J,J or	4 jl.8	Pk C. C. H.	1795.	C r.m Roth abh. 10	
17179 denticeffum Jac.	tooth-leaved	tt. L_J or	4 jnl.j	Pk C. G. H.	1789.	C r.m Sw. ger. 109	
17180 delphiniflthum W.	Larkspur-lv	J,J or	4 ap.jl	Pk C. G. H.	-	C r.m	
XIV. DU'BIA. —Doubtful to which of the sections they belong.							
17181 dscipes Haw.	disk-stalked	tt. i_ leu	5	Africa	1808.	C r.m	
17182 sprium W.	spurious	tt. -	2 ap.jl	V. -	-	C r.m	
17183 gratum IV.	grateful	Citron-satdn.	2 ap.au	Pk.vy	C r.m	
17184 pr thm IV.	spurious	tt. i_ or	2 ap.jl	Pk. -	-	C r.m	
{17185 consanguineum W.	kindred	tt. i_ or	2 ap.au	Pk. -	-	C r.m	
K 5 " ; <> <> *.	Wildenow's tt.	i_J or	2 my.au	W.vy C. G. H.	-	C r.m	
17187 unicolor&rum W.	one-colored	tt. -	2 my.au	C C.G.H.	-	C r.m	
17188 almfldum W.	Aidor IOBVI	tt. -	2 ap.jl	Pk.vy	C r.m	
17189 amplinhnum W.	IargSt	tt. -	2 ap.jl	P. -	-	C r.m	

17190 imbricatum <i>Swt.</i>	imbricated	*L i-J or	3	my.au	Li.p	C.G. H.	1806.	C	r.m	Sw. ger. 1.65		
17191 Aeracle#/i/iuwSwtCow.Parsnep-lvdig	J	or	1	my.s	Gy	C. G. H.	1H00.	C	r.m	Sw. ger. 3.211		
17192 pulverulentum <i>Sivt.</i>	powdery	£ i-	or	1	my.s	Gy.bdC.	G. H.	1822.	C	r.m	Sw. ger. 3.218	
17193 pdiciclkltum <i>Swt.</i>	pedicelled	£ i-	or	1	my.s	Gy.bdC.	G. H.	1822.	C	r.m	Sw. ger. a 250	
171*44 curdufolium <i>Swt.</i>	Thistle-leaved	n. LJ	or	3	my.s	Li.	C.G. H.	1816.	C	r.m	Sw. ger. 1.15	
17195 rugosum <i>Swt.</i>	wrinkly	tL	J	or	3	my.s	Pk.li	C. G. H.	1800	C	r.m	And. ger. ic.
17196 BeaufortiVrNtf/n Swt	Beaufort's	tL	J	or	3	my.au	Li	C. G. H.	..*	C	r.m	Sw. ger. 2.158
17197 principisse <i>Swt.</i>	princess	«. LJ	or	3	my.n	Dk.PC.	G. H.	1820.	C	r.in	Sw. ger. 2.139	
1719H citriodrum <i>Hurt.</i>	Citron-scented	tL	LJ	or	3	my.au	W	C. G. H.	1800.	C	r.m	And. ger. ic.
1719* uniflorum <i>Dec.</i>	one-flowered	tL\	or	3	my.au	..	C. G. H.	1800.	C	r.m		
17200 viscosissimum <i>Swt.</i>	clammiest	«. i-	J	or	3	rayjl	Li.W	C. G. H.	1820.	C	r.m	Sw. ger. 2.118
17201 itatentissimum <i>Wnl.</i>	most spreading	tL	i	or	3	my.au	Li.w	C. G. H.	1890.	C	r.m	Wen. c. ic.
17202 acugnatum <i>Cartn.</i>	Acunna	tL	—	or	3	my.au	...	Tris.d'Ac.18ia	C	r.m		
17203 quinquelobum <i>Col.</i>	five-lobed	tL	J	or	3	my.s	R	C.G. H.	1820.	C	r.m	CoL h. rip. S3

Garden Varieties.

1 abrotanifolium <i>Sw. ger.</i> 351	79 Daveyndum <i>Sw. ger.</i> 32	157 mod6stum <i>Sw. ger MA</i>
2 accroides <i>Sw. ger.</i> 242	80 Denpsidnum <i>Sw. ger.</i> 20	158 Mosfyn* <i>Sw. ger.</i> 10
3 acidum <i>Sw. ger.</i> 261	81 dentatum <i>And. ger. ic</i>	159 multiflbrum <i>Sw. ger.</i> 396
4 acutisobum <i>Sty. ger.</i> 184	82 depéndens <i>Sw. ger.</i> 195	160 multinérvo <i>Sw. ger.</i> 17
5 e'mulum <i>Sw. ger.</i> 160	83 différme <i>Sw. ger.</i> 105	161 Murrayndum <i>Sw. ger.</i> 164
6 affluens <i>Sw. ger.</i> 194	84 dilatatum <i>Sw. ger.</i> 378	162 mutâbile <i>Sw. ger.</i> 213
7 albinbtuin <i>Sw. ger.</i> 359	85 dimacraeflbrum <i>Sw.eer.</i> 220	163 nânun <i>Sw. ger.</i> 102
8 Alltafi <i>Sw. ger.</i> 229	86 diss&tum <i>Sw. ger.</i> 247	164 Nai rnii <i>Sw. ger.</i> 372
9 amoe^num <i>Sw. ger.</i> 121	87 diverlobum <i>Sw. ger.</i> 361	165 nervbsum <i>Sw. ger.</i> 47
10 anthriscifbhum <i>Sw. ger.</i> 233	88 Dobrednum <i>Sw. ger.</i> 253	166 Newshamdnui <i>Sw. ger.</i> 144
11 árdens <i>Sw. ger.</i> 45	89 dumbsum <i>Sw. ger.</i> 19	167 notátura Stu. <i>ger.</i> 208
12 ardescens <i>Sw. ger.</i> 831	90 elátum <i>Sw. ger.</i> J6	168 nummularifbium <i>Bot. rep.</i> 123
13 argfltum <i>Sw. ger.</i> 344	91 éléctum <i>Sw. ger.</i> 23K	169 oblatum <i>Stu. ger.</i> 35
14 armatum <i>Sw. ger.</i> 214	92 eiegans (L.) <i>Sw. ger.</i> 132	170 obovatum <i>Sw. ger.</i> 367
15 asperifolium <i>Sw. ger.</i> 169	93 Regans (11.) <i>Sw.eer.</i> SOS	171 obscrum <i>Sw. ger.</i> 89
16 atroftscum <i>Sw. ger.</i> 82	94 eiegans (VIII.) <i>Sw.eer.</i> 222	172 obtusifolium <i>Sw. ger.</i> 25
17 atropurpiireum <i>Sw. ger.</i> 152	95 errectum <i>Sw. ger.</i> 187	173 optabile <i>Sw. ger.</i> 63
18 atrosanguineum <i>Sw. ger.</i> 151	<M erioca&lon <i>Sw. ger.</i> 357	174 opulifbium <i>Sw. ger.</i> 53
19 aurantiacum <i>Sw. ger.</i> 198	97 eriophylum <i>Sw. ger.</i> 1H	175 ornatum <i>Sw. ger.</i> 39
20 AvToiulnum <i>Sw. ger.</i> 364	98 cnoslpalum <i>Sw. ger.</i> 371	176 Palkii <i>Sw. ger.</i> 224
21 haileyanum <i>Sw. ger.</i> 87	99 eximium <i>Sw. ger.</i> 26.	177 pannifbium <i>Sw. ger.</i> 9
22 Bakeridnutn <i>Sw. ger.</i> 240	100 exomåtum <i>Sw. ger.</i> 381	178 pdrticeps <i>Sw. ger.</i> 49
23 Barclayndum <i>Sw. ger.</i> 304	101 exp&nsum <i>Sw. not figured</i>	179 patens <i>Sw. ger.</i> 125
24 BamardVi/iuwi <i>Sw. ger.</i> 127	102 Fairfu? <i>Sw. ger.</i> 178	180 paucidentatum <i>Sw. ger.</i> 186
25 basilicum <i>Sw. ger.</i> 360	103 flaccidum <i>Sw. ger.</i> 337	181 pavonlnum <i>Sw. ger.</i> 40
26 Beadbnto <i>Sw. ger.</i> 191	104 flexubsum <i>Sw. ger.</i> 180	182 pectinithium <i>Sw. ger.</i> 66
27 Beaufort*?? IMn <i>Sw. ger.</i> 138	105 flocciflum <i>Sw. ger.</i> 129	183 pedunculatum <i>Sw. ger.</i> 346
28 bflllum <i>Sw. ger.</i> 60	116 flundum <i>Sw. ger.</i> 41	184 phocniccum <i>Sw. ger.</i> 207
29 Bcntinckianum <i>Sw. ger.</i> 350	107 folibsum <i>Sw. ger.</i> 310	185 pinguifbium <i>Sw. ger.</i> 52
30 biflbrum <i>Sw. ger.</i> 287	108 Folsjamboe <i>Sw. ger.</i> 312	186 planifbium <i>Sw. ger.</i> 219
31 biparfitum <i>Sw. ger.</i> 143	109 formbusum <i>Sw. ger.</i> 120	187 platanifbium <i>Sw. ger.</i> 326
32 Blandfordiánwra <i>Sw. ger.</i> 101	110 Francisii <i>Sw. ger.</i> 349	188 platyptalon <i>Sw. ger.</i> 116
33 blAndum <i>Sw. ger.</i> 4	111 ftulgens (XIII. i.) <i>Sw. ger.</i> 339	189 Potter* <i>Sw. ger.</i> 147
34 Boy'lca <i>Sw. ger.</i> 50	112 fuscatum <i>Sw. ger.</i> 210	190 Pfmcednum <i>Sw. ger.</i> 386
35 Breestinum <i>Sw. ger.</i> 64	113 glabnfblum. (ac.) <i>Sw. ger.</i> 363	191 principissas <i>Sw. ger.</i> 139
36 Brightinum <i>Sw. ger.</i> 227	1H/jauctifolium <i>Sw.eer.</i> 179	192 psiloph^Uum <i>Sw. ger.</i> 356
37 Broughtonte <i>Sw. ger.</i> 181	115 GowiniSw. <i>ger.</i> 333	193 pubescens <i>And. ger.</i> ic.
38 Brown/i <i>Sw. ger.</i> 146	116 grandidcntatum <i>Sw. ger.</i> 217	194 pulcherrimura <i>Sw.eer.</i> 134
39 Aryonifbium <i>Sw. ger.</i> 320	117 grandiflbrum (VIII. ii.)	195 ptlchrum <i>Sw. ger.</i> 107
40 BurnelJanittu <i>Sw. ger.</i> 369	118 Grenvillianitm And ger.	196 pulveréntum <i>Sw. ger.</i> 218
41 enloéphatan <i>Sw. ger.</i> 201	119 Gurneycum <i>Sw. ger.</i> 393	197 purum <i>Sw. ger.</i> 334
42 calycinum <i>Sw. ger.</i> 81	120 Hammersita? <i>Sw. ger.</i> 225	193 />yrethnfolium <i>Sw. ger.</i> 153
43 campythaeflrum <i>Sw. ger.</i> 251	121 Harwoiid/Ve <i>Sw. ger.</i> 383	199 quadntibrum <i>Sw. ger.</i> 321
44 campyloptpalim <i>Sw.eer.</i> 379	122 icdysarifbliumd.) <i>Sw. ger.</i> 355	200 ramigerum <i>Sw. ger.</i> 3T2
45 candidum <i>Sw. ger.</i> 128	123 Yo*rc<rfthrwm <i>Sw. ger.</i> 133	201 ramulbsum <i>Sw. ger.</i> 177
46 cardifolium <i>Sw. ger.</i> 15	124 HozTetnum <i>Sw. ger.</i> 80	202 recurvatuin <i>Sw. ger.</i> 223
47 cariescens <i>Sw. ger.</i> 388	V25 Hhrnci <i>Sw. ger.</i> 324	203 recurvifbium <i>Sw.eer.</i> 343
48 cartilaRincum <i>Sw.eer.</i> 382	125 Husseyrintwi <i>Sw. ger.</i> 92	204 rfegiuin <i>Sw. ger.</i> 368
49 caryophyll&ccum <i>Sw. ger.</i> 347	Vil imbricatum <i>Sw. ger.</i> 63	205 reticulatum <i>Sw. ger.</i> 143
50 cérinum <i>Sw. ger.</i> 176	128 imperiale <i>Sw. ger.</i> 365	206 llichidnum <i>Sw. ger.</i> 370
51 chserophyllum <i>Sw. ger.</i> 257	129 incanscens <i>Sw. ger.</i> 203	207 rigescens <i>Sw. ger.</i> 112
52 Charlwoodii <i>Sw. ger.</i> 380	130'incrufatum <i>Sw. ger.</i> 308	208 ringens <i>Sw. ger.</i> 256
53 chelidomfblum <i>Sw. ger.</i> 341	131 incrvum <i>Sw. ger.</i> 249	209 Robinbbni <i>Sw. ger.</i> 150
54 phenopodifbium <i>Sw. ger.</i> 328	132 inscription <i>Sw. ger.</i> 193	210 rotundilobum <i>Sw. ger.</i> 932
55 chrysanthemifbium <i>Sw. ger.</i> 124	133 insigmum <i>Sw. ger.</i> 300	211 rubescens <i>Sw. ger.</i> 30
56 Clinton*? <i>Sw. ger.</i> 392	134 intertextum <i>Sw. ger.</i> 135	212 rugbsum <i>And. ger.</i> ic.
57 darum <i>Sw. ger.</i> 366	135 involucratum maximum Do. 33	213 Hessel/dawn <i>Sw. ger.</i> 385
58 coarctatum <i>Sw. ger.</i> 70	136 Jenkins?/n <i>Sw. ger.</i> 154	214 sacpefibrens <i>Sw. ger.</i> 58
59 Colviblntim <i>Sw. ger.</i> 260	137 jonquillinum, <i>Sw. ger.</i> 241	215 salcbrsum <i>Sw.. ger.</i> 309
60 Colvilli Sw. ger. 86	138 Kntgi <i>Sw. ger.</i> 248	816 SaundersM <i>Sw.ger.</i> 205
61 Colvilli (I. <i>Sw. ger.</i> 260	139 labyrinthicum (L.) <i>Sw. ger.</i> 276	217 Scarborovitf. <i>Sw. ger.</i> 117
62 Comptnbue <i>Sw. ger.</i> 122	140 Lamberts <i>Sw. ger.</i> 104	218 scintillans <i>Sw. ger.</i> S8
63 cõmptum <i>Sw. ger.</i> 955	141 lancobatum And. ger. ic	219 scitulum <i>Sw. ger.</i> 3H
64 concavum <i>Sw. ger.</i> 237	112 lasiocafilon <i>Sw. ger.</i> 364	220 ScfWi <i>Sw. ger.</i> 264
65 concinnum <i>Sto. ger.</i> 108	143 latiflbium <i>Sw. ger.</i> 335	221 scutatum <i>Sw. ger.</i> 95
66 conclafisum <i>Sw. ger.</i> 305	144 latitbbum (VIII. ii.)	222 Befitum <i>Sw.eer.</i> 190
67 cdncolor <i>Sw. ger.</i> 140	145 ltilobum <i>Sw. ger.</i> 236	223 sclenifbium <i>Sw. ger.</i> S9
68 congestum <i>Sw. ger.</i> 325	146 lxiflbrum <i>Sw. ger.</i> 216	224 srratifsblum <i>Sw. ger.</i> 221
69 corbscans <i>Sw. ger.</i> 173	147 Lghkt?ck<ff <i>Sw. ger.</i> 377	225 Seymðurs <i>Sw. ger.</i> 37
70 Coamitnum <i>Sw. ger.</i> 189	148 lepidum <i>Sw. ger.</i> 156	226 sisymbiifbium(1.) <i>Sw.ger.</i> JJS
71 crenalbrum <i>Sw. not figured</i>	149 linearum <i>Sw. ger.</i> 116	227 Smithidntfdml. <i>Sw.ger.</i> 358
72 crenatum (XIII. i.) <i>Sw.ger.</i> 345	150 Lousadidnum <i>Sw. ger.</i> 44	228 SmithifSw. gcf. 110
73 crenatum And. ger. ic.	151 lbteam <i>Bot. rep.</i> 328	239 fmyrnifbium <i>Sw. not figured</i>
74 crenulatum <i>Sw. ger.</i> 162	152 macranthon <i>Sw. ger.</i> 83 •	230 soljibilo <i>Sw. ger.</i> 24
75 crispum <i>Sw. ger.</i> J83	153 magnistipulatum <i>Sw. ger.</i> 313	231 SouthqtoYiffrm <i>Sw. ger.</i> PAS
76 crithmifbium <i>Sw. ger.</i> 354	154 Mattocks/a/iuwi <i>Sw. ger.</i> 234	232 sppctabile <i>Sw. ger.</i> 136
77 cruentra <i>Sw. ger.</i> 170	155 mclisrinum <i>Sw. ger.</i> 5	233 sphsrocephalon <i>Sw. ger.</i> 317
78 cuneiflorum <i>Sw. ger.</i> 330	156 mixtum <i>Sw. ger.</i> 71	234 qihondylifMiinn <i>Sw.ger.</i> 24G

235 Spinii Sw. ger. 362	245 torrefactum <i>Su></i> . ger. 243	254 verbasciflrium Sw. ger. 149
236 Staplebmt Sw. ger. 212	246 tyrianthinum Sw. ger. 183	255 verccundum Sw. ger. 316
237 Stowartif Sw. ger. 353	247 Vanriëbe Sw. ger. 7	256 versicolor Sw. ger. 78
238 striatum Sw. ger. 1	248 varium Sw. ger. 1G6	257 vesperinum Sw. ger. 239
239 suffusum Sw. ger. ic. in.	249 veniferum Sw. ger. 322	258 villosum Sw. ger. 100
240 sulphbreum Sw. ger. 163	250 vcniflrum Sw. ger. 258	259 villousum (VIII. ii.) Sw. ger. notfigd.
241 Synnbtii (XL) Sw. ger. 342	251 venosum Sw. ger. 209	260 viscosissimum Sw. ger. 118
242 tanacctiflrium Sw. ger. 336	252 ventistum Sw. ger. 167	261 WcelluTinum Sw. ger. 175
243 Th\$neer Sw. ger. 74	253 verbasciflrum Sw. ger. 157	262 Youngii Sw. ger. 131
244 TibbitBtffUki Sw. ger. 158		

CLASS XVI. ORDER 5.

MONADELPHIA OCTA'NDRIA.

1930. AITO*NL4 L. fil. AITONIA. *{IV. Alton, head-gardener, royal gardens, Kew.}* MeMceee. 1.
17204 capensis L.J.U. Cape tt. i_J or 2 aps Fk C. G. H. 1777. G r.m Bot mag. 178

19*31. COMESPE'RMA Lab. COMESPERMA. (*Kome, hair, sperma, seed.*) Polygalece. 2.—10.
17205 tfricina Dec. Heath-like il i_J or 3 my.au P N. HoU. 1822. C s.p.l
17206 cordifbia Cun, heart-leaved *LJ or 3 my.au P N. HoU. 1822. C s.p.l

CLASS XVI. ORDER 6.

MONADE'LPHIA DECA'NDRIA.

* 1932. GERAWIUM *Hcrit. CRANE'S BILL.* (*Geranos, crane; capsule like head and beak*) Geraniacee. 58.-66.

I. UNIFLORA.— <i>Perennial; peduncles one-flowered.</i>		
17207 sibiricum L.	Siberian	3c A P* 1 jnj1 W Siberia 1758. D s.l Jac. vin. 1.19
17208 sanguineum L.	bloody	3r A P' 1 jns Bd Britain rocks. D s.l Eng.bot.272 .
2 villosissimum Dec.	hairiest	3 ^ o r } jns Bd Europe ... D co
17209 lancastrense With.	Lancaster	.* A P' I jns St Britain ... D 61 Cav.dis/4.76.3
prostKtum Lk. sanguinum /3 prostratinum Dec.		
17210 multifidum D. Don	multifid	H LAJ or 1 jn.n R C.G.H. 1817. C 81 Sw. ger. 3.245
II. BIPLORA.— <i>Perennial i peduncles two-flowered.</i>		
17211 incanum L.	hoary multifid	CTG.H. 1701. S r.m Cav. dis. 4.82.2
17212 canescens Herit.	canescent	C. G. H. 1787. S r.m Her. ger. 38
17213 argenteum L.	silvery-leaved	S. Europe 1699. D s.l Sw. ger. 59
17214 cinereum Cav.	grey	Pyrenees ... D s.l Her. ger. 37
vkrium Herit.		
17215 anemonefolum Herit.	Anemone-lvd	Madeira 1788. C r.m Sw. ger. 244
17216 macrorfizum L.	long-rooted	Italy 1576. D s.l BoL mag. 2420
17217 tuberbum L.	tuberous-rooted	Italy 1596. D r.m Sw. ger. 155
17218 gymnocalon Dec.	naked-attemded	1 jns B Iberia 1814. D 81
17219 ibricum Cav.	Iberian	H jns Levant 1802. D s.l Sw. ger. 84'
17220 nodbsum L.	knotty	Z 1 my.o P . England moun. D 81 Eng. bot. 1091
17221 angulatum Cur.	angular-stalked	1 myjn P . 1789. D 81 Bot mag. 203
17222 Wallichianum Swt.	Wallich's	fmy.au R Nepal 1819. D s.l Sw. ger. 90
17223 Vlassovianum Fis.	Vlassov's	1 my.au St Crimea 1821. D s.l Sw. ger. 228
17224 striatum L.	channeled	1 my.o St Italy 1629. D s.l Bot mag. 55
17225 refl^xum L.	rcfLeXrflowered	1} my.jn R Italy 1758. D 21 Cav. dis. 4. 81.1
17226 phae'um Lam.	dusky	Bk England m.thi. D s.l Eng. bot 322
17287 fuscum L.	brown	Br S. Europe 1759. 1) co
17228 lividum Herit.	livid-flowered	P Switzerl. 1775. I) s.l Her. ger. 39
17229 enostemon Fis.	woolly-stmnd	B Siberia 1822. D co Sw. ger. 197
2 pallidum Swt.	pale	Nepal 1822. D s.l Sw. ger. 197.6
17230 Aatrachioles Cav.	Batrachium-lk	B Europe 1817. D 81 Cav. dis. 4. 85.1
17231 Lambert* Swt.	Lambert's	R Nepal 1824. D s.l Sw. ger. 338
17232 sylvaticum L.	wood	Britain m.thi. D B.1 Eng. bot 121
17233 pratense /., meadow Crotfoot-lvd		B Britain me.pa. D B.1 Eng. bot 404
2 flrc Mho	white-flowered	W Britain gard. D co
3 flre variegato	vanegated-flwd	U my. ll Str Britain gard. D co
4 daturicum	Daurian	Ijmy.jl B Dauria 1818. D co
5 flore pleno	douhlc-flwd	2' my.ll B Scotland moun. D co
17234 longipes Dec.	long-stalked	1 my.jl Li . 1823. D co
LondOs'i Lk.		
17235 maculatum L.	spotted	f my.au P N. Amer. 1732. D s.l Cav. dis. 4.8a 2
17236 cojnnum Bieb.	hill	1 my.au P Siberia 1815. D co
17237 palustre L.	marsh	2 jn.au P Germany 1732. D B.1 Sw. ger. 3
17238 flconitiflrium Herit.	Aconite-leaved	li my.jn W Switzerl. 1775. D 81 Her. ger. 40
17239 dahbricum Dec.	Dahurian	1 my.jn P Dahuria 1820. D co
17240 caeruleum Pat.	blue	jns B Dahuria 1824. D s.l
17241 altaicum Hort.	Altaian	Pa.R Altaia 1818. D co
17242 pildsum Forst.	pilose	\$ my.au P N. Zeal. 1821. D co Sw. ger. 119
17243 parviflbrum W. en.	small-flowered	S my.jn P V. D. L. 1816. D co
17244 ncpalencs Swt.	Nepal	A my.au R Nepal 1818. D co Sw. ger. 12
17245 crisfatum Stcv.	crested	lijn.s R Iberia 1820. D B.1 Soc. mos. 4.5
albñum Bieb.		
17246 pyrenaicum L.	Pyrenean	my.au P Britain me. pa. S 8.1 Eng. bot 405
17247 nemoribus Ten.	grove	jns P Italy 1821. S co
pyrenaicum y nemoribus Dec		
17248 umbrbsum Kit.	shaded	J A pr 1 jn.s P Hungary 1804. S co W. & K. 2.144
III. A'NNVA.— <i>Annual; peduncles two-flowered.</i>		
17249 mblle L.		
17250 pusillum L.		

17251 rotundiflura £.	round-leaved	O w	I 3n.ll	Pk	England	gra.ba.	S	co	Eng. bot.	157
17252 pallens Bieb.	pale	O un	i J	Pa.R	Iberia	1820.	S	co		
17253 columbinum L.	dove	J g	8:-	PK	Britain	cha.ba.	S	co	Eng. bot	259
17254 dissectum L.				PK	Britain	was.gr.	S	co	Eng. bot	753
17255 carolinianum L.	Carolina	O un	j jlau	P	N. Amer.	1725.	S	co	Cav.dis.4.84.1	
17256 Villsum Ten.	villous	O un	ijn.jl	W.vy	Italy	1820.	S	co		
17257 bohemicum L.	Bohemian	O un	jn.au	P	Bohemia	1683.	S	co	Cav.dis.4.81.2	
17258 divaricatum Ehrh.	straddling	O un	jl.au	P	Hungary	1799.	S	co	W.&K.2.123	
17259 Ucidium L.	shining	O un	■ my.au	Pk	Britain	ston.pl.	S	co	Eng. bot	75
17260 Robertianum L.	flefft-Robert	O w	1 ap.o	R	Britain	ston.pl.	S	co	Eng. bot	1486
17261 purpureum Vtt.	purple	O w	1 ap.o	R	Switzerl.	1819.	S	co	VilDel.3.40	
17262 inodorum Ph.	scentless	O pr	1 ap.o	R	N. Amer.	?1800.	S	co		
17263 Briceanuw* Swt.	Miss Brice's	O pr	t ap.o	Pa.R	Bristol	rocks.	S	co	Sw. ger. icln.	
1933. PHILOTHE^CA Bud.	PHILOTHBCA, (<i>Psilos</i> (efr. <i>phiros</i>), Smooth, theke, sheath, tube of stain.)	Bud.	1.							
17264 australis Bud.	southern	si II	J or 2 ap	W	tf.	HolL	1824.	C p.1	Lin.tr.	11.21
	Eriostemum salsoloide Sm.									
1934. CHEIROSTETMON K'S/B.	CHEIROSTBMOW.	{Cheir, hand, stemon, stamen.)								
17265 platanoides #.\$•£.	Platanus-like	£ • or 30.								
1935. HUGCPNIA L.	HUGONIA. (<i>Dr. A. J. Hugo</i> , Hanover, friend of Haller's.)	<i>Chlieracea affine</i> .	2.-3.							
17266 ntfstax L.	mustachio	» D o r 10	jn.jl	Y	Ceylon	1818.	C s.l	R.mal.	2.29	
17267 serrate Lam.	serrated	■ I Q o r 8	jn.au	Y	Mauritius	1824.	C s.l	Cav. dia.	3.73.1	
1936. OMPH ALO^BIUM Gae.	OMPHALOBIUM (<i>Omphalos</i> , a navel, labos, a pod.)	<i>Connarfae.</i>	2.—12							
17268 indicum Gae.	Indian	* Q o r 8	...	Pa.R	Ceylon	...	C p.l	Bur.zey.	89	
Ahus Rxdece Hjhvel Mil.	Connarus asiaticus W.									
17269 africānum Dec.	African	« D < » 8	—	Pa.R	Guinea	1822.	C p.1	Cav.dis.7.221		
Connarus africanus Lam.										
1937. CONN A^RUS L.	CONWARUS. (The Greek name given to this kind of tree.)	<i>Connarhcece.</i>	3.-7.							
17270 paniculatus Box.	panicled,	i Q o r 8	...	W	Chittagong	1824.	C p.1			
17271 nitidus Box.	shining	MO or 8	...	W	Silhet	1824.	C p.1			
17272 pubescens Dec.	pubescent	i Q o r 6	...	W	Guiana	1822.	C p.1	Aub. gui.	1.187	
*1938. BRO'WNEJ Jac.	BROWNEA. (<i>Patrick Browne</i> , M.D. auth.of hist. of Jam.)	<i>Legum. Cafs. Geoff</i>	3.-5.							
17273 coccinea Jac.	scarlet	sH O spl 6	jlau	S	W. Indies	1793.	C r.l	Jac. am.	121	
17274 rfcasa Pers.	roseata	* Q o r 8	jn.au	S	Trinidad	1820.	C s.l	Ph.tr.	1771.8,9	
17275 latifolia Jac.	broad-leaved	i Q o r 6	...	S	Trinidad	1824.	C 8.1	Jac. fr.	16	
1939. PA'RK/i* R. Br.	NITTA TREB.	(<i>Mungo Park</i> , the celeb. African traveller.)	<i>Legumin. Mimbea.</i>	2.						
17276 biglobbsa B. Br.	biglobular	I O esc 30	mrap	Ve	Guinea	1822.	S r.m	Beau.ow.2LS0		
Jiiga biglobbsa W.		4 • esc 30	mrap	Ve	SL Leone	182a	S r.m			
17277 uniglobosa G. Don	uniglobular									
*194a. HO^VEK R.Br.	HOVEA. (<i>A. P. Hove</i> , a Polish botanist.)	<i>Leg. Pap. Lbtea?</i>	<i>GenUteaf.</i>	11.						
17278 longifolia B. Br.	long-leaved	■ H tor 3	jns	P	N. S. W.	1805.	S s.p	Bot.reg.614		
17279 linearis B. Br.	Wineax-leaved	■ ■ ■ 3	mr.jl	P	N. S. W.	1796.	S s.p	Bot.reg.463		
Poirëta linearis Sm.										
17280 rosmarinifolia Curt.	Rosemary-lvd	■ ll lor	S	mr.jl	P	N. Holl.	1824.	C s.p		
17281 lanceolata B. M.	spear-leaved	■ ^ J or 3	mr.jl	P	N. Holl.	1805.	C s.p	Bot mag.	1624	
17282 elliptica Dec.	elliptic-leaved	■ ■ ■ 3	mr.jl	P	N. Holl.	1817.	C s.p			
PoireUaclJpticas;».										
17283 latifolia Lod.	broad-leaved	■ ■ or 3	mr.jl	P	N. Holl.	1820.	C s.p	Bot cab.	30	
17284 purpurea Swt.	purple	■ ■ or 3	mr.jl	P	N. Holl.	1820.	C s.p	Sw. au.	13	
17285 Celsi >onp.	Celf's	■ ■ or 4	mr.jl	B	N. Holl.	1818.	C s.p	Bot.reg.280		
17286 mucronata Cun.	mucronate	■ ■ or 4	mr.jl	B	N. Holl.	1824.	C s.p			
17287 pannosa Can.		■ ■ 3	mr.jl	P	N. Holl.	1824.	C s.p			
17288 lanigera Lo. C.	wool-bearing	■ ■ 3	mr.jl	P	N. Holl.	1824.	C s.p			
19*1. PLATYLO^BIUM Sm.	FLAT PEA. (<i>Platys</i> , broad, lobos, pod.)	<i>Leg. Pap. Lbt. GenU.4.—5.</i>								
17289 formbsum 5m.	beautiful	* LJ or 4	jn.au	O	N. S. W.	1790.	S s.p	Bot. mag.	469	
17290 parviflrum Sm.	small-flowered	m J or 4	my.s	O	N. S. W.	1792.	S s.p	Bot. mag.	1520	
17291 ovatum Andr.	ovate	M J or 4	jns	O	N. Holl.	1825.	8 s.p			
17292 triangulare B. Br.	triangular-fed	« i J or 4	jns	O	V. D. L.	1805.	S s.p	Bot. mag.	1508	
1942. PLATYCHRUM Delau.	PLATYCHILUM.	(<i>Platys</i> , broad, cheilos, lip; corolla.)	<i>Leg. Pap. Lbt. GenU. L</i>							
17293 CelsWnun Her. am.	Cete's	§ » u j o t 3 J J ...	Y	N. Holl	L	1820.	C s.p	Her. am.	187	
Gompholbium Cehianum Hort										
1943. BOSSIzTil Ven.	BOSSI^A. (Af. Boissieu-Lamartiniere, accomp. La Perouse.)	<i>Leg. Pap. LbU Genii.</i>	14.							
	I. APIY'LLE.—Stems flat, leafless.									
17294 Scopolodrium Sm.	Plank-plant	* i J or 10	my.Ji	Y	N. S. W.	1792.	C 8.1P	Bot rep.	191	
17295 enskta Sieb.	sword	H LJ or 6	my.jn	Y	N. Holl.	1824.	C s.Lp	Sw. au. ic.in.		
VS96rtiSk.B.Br.	xufoui. Jflowered*	J or 6	jns	O	N. Holl.	1803.	C s.Lp			
	II. Foub&JB.—Stems flat, leafy.									
17297 /inophylla B. Br.	Flax-leaved	* [J or 3	jl.s	O	N. Holl.	1803.	C s.Lp	Bot. mag.	2491	
17298 heterophylla Ven.	various-leaved	* [J or 3	my.d	Y	N. S. W.	1792.	C a.Lp	Bot. mag.	1144	
17299 rotundophylla Dee.	round-leaved	* [J or 3	my.jn	Y	N. Holl.	1824.	C s.Lp			
17300 rhombifolia Sieb.	rhomb-leaved	* [J or 1	my.jn	Y	N. Holl.	1820.	C s.Lp	Sw. au. 9		
	III. TERETICAI/LUE.—Stems round, leafy.									
17301 cordifolia Swt.	heart-leaved	■ ■ J or 1	my.jn	Y	N. Holl.	1820.	C s.Lp	Sw. au.	20	
17302 microphylla Sm.	small-leaved	■ ■ J or 3	my.au	Y	N. S. W.	1803.	C s.Lp	Bot. cab.	656	
17303 /enticularis Sieb.	Jcint\leaved	■ ■ J or 3	my.jn	Y	N. Holl.	1823.	C s.Lp	Bot cab.	1238	
17304 cinerea B. Br.	grey sharp-lvd	■ ■ J or 3	my.jl	Y	V. D. L.	1802.	C s.Lp	Bot reg.	306	
17305 prostrata B. Br.	prostrate	■ ■ J or 1	JLi	Y	N. S. W.	1803.	C s.Lp	Bot mag.	1493	
17306 foliosa Cun.	leafy	■ ■ J or 4	my.jn	Y	N. Holl.	1824.	S s.Lp			
17307 ftuxifolia Cun.	Box-leaved	SU i J or 4	my.jn	Y	N. Holl.	1824	S s.Lp			
1944. GOCDLFR.Br.	GOODIA. (<i>Peter Good</i> , a collector of plants for Kew gardens.)	<i>Leg. Pap. Lbt</i>	6 n k a							
17308 ZotifoliaSm/.	Lotus-leaved	JL J or 3	ap.H	Y	V.D.L.	1793.	S s.p	W mag	958	
17309 pubescens Sims	pubescent	S J J or 2	ap.jl	Y	V.D.L.	1805.	S s.p	Bt	S'1310	
17310 polisperma Drc.	many-seeded	« i J or 2	ap.jl	Y	V.D.L.	1790.	C s.p	A		
Cftisus tomentbsui Andr.	in the gardens, under the name of Gooldia totifolia, frequently.									

- * 1945. SCOTTJi< R.Br. SCOTHA. (*Robert Scott, M.D., prof, bot, Dublin.*) *Leg. Pap. Lbt. Genis.* 1
17311 dentate *R. Br.* tooth-leaved **l*₁*J*or S jn.8 R.u N. Holl 1803. C s.p.
1946. T E M P L E T O W J* R.Br. TEMPLBTONIA. (*J. Templeton, excellent' Irish bot.*) *Leg. Pap. Lbt. Genis.* 8.
17312 retusa *R. Br.* retiae-leaved **u*₁ or 2 mr.jn K N.Holl. 1803. C s.p. Bot. mag. 2334
17313 glauca *B. M.* glaucous • LJ or 2 ap.my R N. Holl. 1818. C ssp Bbot reg. 359
1947. RA'FN/i4 Thun. RAFNXA. (*C. G. Rafn, Copenhagen a bot author.*) *Leg. Pap. Lit. Genis.* 5.—14.
17314 elliptica *Thun.* elliptic tt. t*J* or 2 jn.jl Y C. G. H. 1819. C 81
17315 cuncifolia *Thun.* wedge-leaved tt. i*J* or 2 jn.jl Y.p C. G. H. 1816. C s.l
17316 trifolia *Thun.* three-flowered & O or 3 jn.jl P C. G. H. 1786. S s.l Bot mag. 482
BorbnsTz cordata *B. Rep.*
17317 lancea *Dec.* lance-leaved tt. LJ or 2 jn.jl Y C. G. H. 1823. C s.l Th. h. 1800.4
17316 opposita *Thun.* opposite tt. LJ or 2 jn.jl Y C. G. H. 1824. C 61
Crotalaria opposite *L.*
1948. VASCOM Dec. VASCOA. (*Fascode Gama, a celeb. Portug. circumnav.*) *Leg. Pap. Lbt. Genis.* 2.
17319 amplexicaulis *Dec.* stem-clasping, at t-j or 4 jl.au Y C. G. H. 1816. € p.l
Uafn/Vi amplexicaulis *Thun.* Crotalaria amplexicaulis *L.*
17320 perfoliata *Dec.* perfoliata **i*₁*J* or 4 jl.au Y C. G. H. 1812. C p.l Seba 1. 24.5
Borbnnirt perfoliata *Thun.* Crotalaria amplexicaulis *Lam.*
- * 1949. BORBO*NJ/I L. BORDOVIA. (*Gaston Bourdon, son of Hen. IV., France.*) *Leg. Pap. Lbt. Genis.* S.-13.
17*21 barbate *Lam.* bearded ■■■ or 4 jl.au c. G. H. 1823. S p.l Lam.il.fil 0.2
17322 irincriva *L.* three-nerved ■■■ or 6 jl.au ■■■ C. G. H. 1759. S p.l Pluk. al. 297.4
17323 lanceoflate *L.* lance-leaved ■■■ or 5 jl.au ■■■ C. G. H. 1752. C p.l Jacsc. 2.217
17324 undulata *Thun.* wave-leaved ■■■ or 4 jl.au ■■■ C. G. H. 1812. C p.l
17325 cordata *L.* heart-leaved ■■■ or 6 jl.s ■■■ C. G. H. 1759. S p.l Jacsc. 2.218
17326 crenato *L.* notch-leaved ■■■ or 6 jn.au Y C. G. H. 1774. S p.l Bot. mag. 274
17327 ciliata *W.* ciliated ■■■ or 3 jl.au Y C. G. H. 1816. S p.l H.pfl.8f62.2
17328 ruscifolia *B. M.* Ruscus-leaved ■■■ or 3 jl Y C. G. H. 1790. S p.l Bot mag. 2128
1950. HA'LLIA Thun. HALLIA. (*Berger Martin Hall, pupil of Linnaeus.*) *Leg. Pap. Lbt. Genis.* 5.-7.
17329 alata *Thun.* winged ■■■ or 1 jl.au P C. G. H. 1818. S l.p
17330 flaccida *Thun.* flaccid ■■■ or 1 jl.au P C. G. H. 1789. S l.p
17331 cordata *Thun.* heart-leaved ■■■ or 2 au P C. G. H. 1787. S l.p Jac.sa 3.296
17332 asarina *Thun.* Asarum-like ■■■ or 1 jl.au P C. G. H. 1810. ■ l.p
17333 imbricate *Thun.* imbricated ■■■ or 2 II au P C. G. H. 1812. ■ 8.1 Bot mag. 1850
Hällia monophyilla Desv.
- * 1952. CROTALARIA L. CROTALARIA. (*Krotalon, a Castanet; seeds rattling.*) *Leg. Pap. Lbt. Genis.* 8A.—140.
- I. DECURRENTI&.—Leaves simple; stipules decurrent; racemes terminal or opposite the leaves; stems herbaceous; corolla yellow, smaller than calyx, o/i nearly equal.
- 17336 alata Jfcem. winged P or 1 jl.au Pa.Y Nepal 1818. S co
17337 stipularia Desv. /arge-stipuled ■■■ or 1 jl.au Y Cayenne 1823. S co
17338 platycarpa IM. broad-podded Olun 1 jl Y N. Amer. 1823. S co
17339 Purhii Dec. Pursh's ■■■ 1 jn.jl Y N. Amer. 1800. S co Pluk. al. 277. 2
ffrigida Ph.
17340 parviflora Roth small-flowered O un 2 au.s Y R Indies 1817. S co
17341 sagittalis L. arrow Oun 1 jnjl Y America 1731. S co Pluk. al. 169.6
17342 bialata Schrank two-winged O or 1 jnjl Y 1820. 8 co
17343 ovkis Ph. oval O or 1 jl.au Y Carolina 1810. S co
17344 rubigindsa W. rusty ED un 1 jl O R Indies 1807. S co
- II. ADBCURRENTUE.—Leaves simple J stipules not decurrent, often nones racemes terminal or opposite the lean,
- flowers variable.
- 17345 verucosa L. waited O or 1 jl.au B W. Indies 1731. S r.m Bot reg. 1137
cerulea Jac.
2 acuminate Dec. acuminate O or 1 jl.au B E. Indies 1731. S p.l Bot rep. 208
17346 semperfibrrens Fen. ever-blowing ■■■ un 3 mr.s Y R Indies 1816. C p.l Ven.cels 17
17347 retusa L. retiae-leaved ■■■ un 1 jnjl Y.p E. Indies 1731. S ■■■ Bot reg. 253
17348 spectabilis Roth showy ■■■ or 1 jn.jn au P. E. Indies 1820. S co
17349 pulcherrima Rox. very neat ■■■ ■■■ Y Mysore 1814. C co Bot mag. 2027
17350 benghalensis Lam. Bengal O un ■■■ Y E. Indies 1806. S 8.1 Pluk. al. 169.5
17351 jincea L. rush-stalked El un 1 jnjl Y E. Indies 1700. S p.l Bot rep. 422
17352 tenuifolia Rox. fine-leaved tt.r-lor 2 jnjl Y E. Indies 1816. S p.l Bot reg. 982
17353 fenestrata B. M. vridow-calyxed O un 2 jnjl Y E. Indies 1815. S p.l Bot. mag. 1933
17354 sericea Retz. silky [Q] un Hjnjl Y E. Indies 1807. ■■■
17355 Burmanni Dec. Burxmann's ED or 1 jl.au Y E. Indies 1800. ■ co Bur. in. 48.1
sericea Brm.
17356 hirsute W. hairy O or ■■■ jl.au Y R Indies 1818. ■ co
17357 paniculate W. panicled H or ■■■ jnjl Y Java 1820. ■ 8.1
chinensis Lam. not L.
17358 pithecia Andr. fair tt. O un 2 mr.myY E. Indies 1800. ■ pi Bot rep. 601
17359 Berteriina Dec. Berter's tt. O or 2 my,jl Y W. Indies 1818. ■ 8.1
17360 Nova; Hollandiaj Dec. New Holland jf ES or 2 my,jl ■■■ B.1
17361 paulina Schrank St. Paul's ■■■ 2 au.s ■■■ Brazil 1823. co Seh. mo. 88
17362 bifaria L. two-rowed ■■■ ■■■ 3 jl.au ■■■ R Indies 1817. co
17363 diffifta Lk. diffuse ■■■ ■■■ i jnjl ■■■ E. Indies 1790. ■■■ co
17364 bifora L. two-flowered ■■■ un ■■■ i jnjl ■■■ E. Indies 1824. ■■■ co
17365 glatica JV. Slaucous ■■■ ■■■ i jnjl ■■■ Guinea 1816. ■■■ co Bur. in. 48.2
17366 pumila Schrank Swarf Jcrm or ■■■ ■■■ ■■■ 1816. ■■■ co
square-stalked ■■■ O un 2 o.n E. Indies 1806. ■■■ S.1 Bot rep. 593
17367 tetragyna Andr. Flax-leaved tt.^ or ■■■ jl.au ■■■ Nepal 1820. ■■■ co
17368 inifilia L. furred tt.Qor ■■■ jl.au ■■■ Jamaica 1820. ■■■ co
17369 pellita Bert. acuminate Qor ■■■ jl.au ■■■ CG.H. 1820. ■■■ co
17370 acuminate Dec. Theban tt-i-Jor 2 my,jl ■■■ Egypt 1818. C B.1 Del. e.g. 37.1
17371 thebkica Dec.
- III. AxiiARiFLbRJE.—Leaves simple : stipules not decurrent or none; flower P^{d, inc*late} or *«& axillary..
- 17372 procumbent Af. * S. procumbent JUESJor 1 my,jl Y Mexico 1723. S s.l
17373 tuberosa Ham. tuberous A O or 1 jnjl P Nepal 1821. S ■■■

IV. ANTHYLLORDBS.—Leaves simple; stipules not decurrent or none; flowers lateral and terminal; calyx five-parted, very hispid, with two wing-formed lobes.

- 17374 anthylloides Lam. Anthyllis-like •••' or 4 aus Y E. Indies 1789. F s.1
 17375 calycina Schrank large-calyxcd O 1 Jn.jl B E. Indies 1816. F s.1 Sch. mo. 11
 17376 nepalensis Lk. Nepal O un 1 jnjl B Nepal 1823. F co
 17377 Roxburghia Dec. Roxburgh's n. • or 2 jn.jl Y EL Indies 1820. F s.1
 stricta Rox. anthylloides D. Don Chinese n. • or 2 jnjl Y China 1818. F 8.1
 17378 chinensis L. hairy O un \| jn.au Y E. Indies 1804. F co
 17379 prostrata Rtl. prostrate JiOuu \| jnjl Y E. Indies 1816. F p.1
 17380 hirta W. en. O un \| jn.au Y E. Indies 1816. F p.1

V. TRIFOILOIDS.—Leaves trifoliate; racemes terminal or opposite the leaves; stipules broad, leathery.

- 17381 arborescens Lam. arborescent • i_J un 6 jn.o Y C. G. H. 1774. S sp Jac. vin. 3.64
 incansecens L.

- 17382 turgida Her. am. turgid • o r 3 jl.au Y 1820. C 8.1

i. OPPOSITIFOLIADS.—Leaves trifoliate; racemes opposite the leaves; stipules setaceous or none.

- 17383 taburnifolia L. Laburnum-lvd £ Ol un 5 jls. Y E. Indies 1739. S co R. mal. 9. 27
 17384 pendula Bert. pendulous tt • or 5 jls. Y Jamaica 1820. C 8.1
 17J85 anagroides Hum. Anagyris-like • J' or 6 jls. Y Trinidad 1823. C 8.1
 17386 Brôneana Bert. Browne's .! lor 4 jl.au Y Jamaica 1816. C 8.1
 17387 br. cteata Rox. torge-bracted • or 4 jl.au Y E. Indies 1820. C 8.1
 17388 cyrtoscrtes Box. Cyttisus-like • J' or 3 jl.au Y E. Indies 1826. C 8.1
 17389 cajaniflora Wa. Hum. Cajan-Hke • or G jls. Y S. Amer. 1824. C 8.1
 17390 foliota TV. leafy • or 3 jn.jl Y E. Indies 1818. C s.1
 17391 orixensis W. en. Onxa O un 1* jl.au Y E. Indies 1816. S p.1
 17392 virgata Box. twiggy • or 3 jn.jl Y E. Indies 1816. C 8.1
 17393 cubensis Dec. Cuba O or 1 jl.au Y Cuba 1820. S 8.1
 hirta Lag.
- 17394 purpurascens Lam. purpurascens rm or 1 jl.au P Madagasc. 1824. S 8.1
 17395 p&mila Or. dwarf nmor Jjn.jl Y Cuba 182a S 8.1
 17396 incana L. hoary tol un 2 jnjl G.Y. W. Indies 1714. S 8p Bot reg. 377
 17397 nrtcans Lk. glittering J' Blun 2 jl.au Pa.Y 1820. S co
 17398 curta Lk. short-keeled m i un 1 ap.my Y S CO
 17J90 viteirna Ker. yolk-like J' un 2 TO* X Brazil 1819. C co Bot reg. 417
 17400 senegalensis Bac. Senegal nm or 1 jnjl Senegal 1819. S s.1
 17401 villosa Thun. villous 231 or 1 jnjl C. G. H. 1824. S 8.1
 17402 modicaginea Lam. Modicago-like rmor E. Indies 1816. S 8.1 K. cur. 4.1803.5
 Trifoliastrum W.
 17403 purpurea Ven. darf-purple • i J un 4 mr.my P C.G.H. 1790. F Bot reg. 128
 17404 microphylla Vahl small-leaved Jc [Q] or } jl.au Y Arabia 1820. F 8.1
 17405 micrantha Lk. small-flowered O un 1 jl.au Ceylon 1823. C CO
 17406 pillida H. K. pale-flowered O un 2 jnjl Pa.Y Africa 1775. F BD
 17407 argentea Jac. silvery <-i_J or 1 jnjl Y C. G. H. 1823. F 8.1 Jac. sc. 2.220
 17408 pulchella Andr. neat <_J un 3 Lau Y C. G. H. 1800. F P.; Bot. mag. 1699
 17409 SaXuna B. Rep. Salt's • un 3 jnjl Y Abyssinia 1810. F p.1 Bot. rep. 648
 17410 angustifolia Jac. narrow-leaved • i J un 1 J my.s Pa.Y C. G. H. 1815. F BD Jac. BC 2.219
 17411 obsecra Dec. obscure U-uJor 2 jnjl Y C. G. H. 1820. F 8.1

ii. AXILLA^RES.—Leaves trifoliate; flowers axillary.

- 17412 /otifolia L. Lotus-leaved * Q un 3 jnjl Y o Jamaica 1732. F Di.el. 102. 121
 17413 axillaris H. K. axil-flowered O un 1 jl.au Y Guinea 1781. F FL mex. ic. in.
 17414 triantha A. & S. three-flowered O or 2 jnjl Y Mexico 1824. F

VI. QUINQUEFOYIA.—Leaves 5-foliate.

- 17415 quinquef5ia Z# five-leaved O un 1 jnjl Y E. Indies 1792. C sp R. mal. 9.28

VI. D. J. BLE.—Doubtful to which of the sections they belong.

- 17416 angulata Mil. angular O or 1 jnjl Y S. Amer. 1706. F co
 17417 fruticosa MO. shrubby • or 2 jnjl Y Jamaica 1716. F co
 17418 Langsdorff* Fis. Langsdorf's O or 1 jnjl Y 1820. F co
 17419 filiformis Box. tawny O or 1 jnjl Y R Indies 1817. S co

1953. HYPOCALYPTUS Thun. HYPOCALYPTUB. (H under, kalypto, to veil.) Leg. Pap. Lbt. Gents. 1.
 17420 obcordatus Thun. obcordate • LJ pr Uj P C.G. H. 1823. Cfs-p
 Crotalaria cordifolia L.

1954. VIBCTRGIA Spr. VIBORGIA. (Eric Viborg, an acute Danish botanist) Leg. Pap. Lbt. Genis. 2. — &

- 17421 sertcea Thun. silky • J or 3 jl.au Y C. G. H. 1780. C l.p
 17422 obcordata Thun. obcordate • LJ O T 2 jl.au Y C. G. H. 1810. S l.p Bot cab. 509
 Crotalaria floribunda Lod. obcordata Berg.

1955. LODDIGETSL1 B.M. LODDICESIA. (Conrad Loddiges, nurseryman, Hackney.) Leg. Pap. Lbt. Genis. 1.
 17423 oxalidifolia B. M. Oxalis-leaved tt. LJ pr \| my.s Pa.P C. G. H. 1802. C p.l Bot.mag. 965

1956. LEBECKIA/ii Thun. LEBECKIA. (Lebeck, an obscure botanist.) Leg. Pap. Lbt. Genis. 6.—1L

- I. SIMPLICIFOLIJS.—Leaves simple.
 17424 subnftda Dec. sub-naked <L I J or \| ap.my Y C. G. H. 1824. C p.l
 17425 contaminata Dec. contaminated • i J or 5 ap.my Br C. G. H. 1787. S p.1
 17426 sepiaria Thun. hedge • i J or 3 ap.my Y C. G. H. 1820. C p.1 Pluk. al. 424.1

II. TRIFOLIA*T2E.—Leaves trifoliate.

- 17427 subternata Lk. subterete • i J or 3 jn.jl Y C. G. H. 1824. S co
 17428 scircea Thun. silky • i J or 2 ap Y C. G. H. 1774. S p.1
 17429 cytisoides Thun. CyUsus-like 1 J or 3 ap Pk C. G. H. 1774. S p.1 Com. b.2.107

1957. DICHPLUS Dec. DICHILUS. (Dis. twice, cheilos, lip; calyx.) Leg. Pap. Lbt. Genis. 1.
 17450 Lebeckioides* Dec. Lebeckia-like ■ i J or 2 ap.my Wv C. G. H. 1826. C p.s Dec. leg. & 35

1958. SARCOHYLLUM Thun. SARCOHYLLUM. (Sarcophyllo, flesh, phyllon, leaf) Leg. Pap. Lbt. Genis. 1.
 17431 carnifsum Th. Hesby-Jointed-lvd • uJ cu 3 my.au Y C. G. H. 1812. C s.1 Bot mag. 2502

195a JISPA'LATHUS L. ASPALATHUS. (A, priv., spaoo, to extract; thorns.) Leg. Pap. Lbt. Genis. 31.—86.

- I. FABCICUIAVE.—Flowers subsessile; leaves in fascicles.
 17432 spinbsaZ. spiny m \ot 2 jl.au Y C. G. H. 1824. C p.1 F
 17433 capitata L. capitate • LJ or 2 jl.au Y C. G. H. 1823. C p.1 PhX
 17434 astroUcs L. starry • LJ or 2 jl.au Y C. G. H. 181a C p.1 Sebale
 17435 Chenopoda L. Goosfoot • LJ pr 3 jl.au Y C. G. H. 1750. C p.1 Brey.c II

17436	álbena	<i>Thun.</i>	whitish	[■■■■■]	4	jl.au	W	C G. H.	1774.	C	p.1	Lam.il	620.1
17437	hfstrix	<i>L.</i>	porcupine	[■■■■■]	2	jl.au	Y	CG. H.	1824.	C	p.1	Pluk.al.	413.1
17438	tfymjfolia	<i>L.</i>	Thyme-leaved	[■■■■■]	2	jl.au	Y	CO. H.	1825.	C	p.1	Pluk.m.	413.6
17439	mcifolia	i.	Heath-leaved	[■■■■■]	2	jl.au	Y	C. G. H.	1789.	C	p.1		
17440	hfspida	7%in.	hispid	[■■■■■]	2	jl.au	Y	C. G. H.	1818.	C	p.1		
17441	asparagooides	<i>L.</i>	Asparagus-Ik	[■■■■■]	3	jl.au	Y	C. G. H.	1812.	C	p.1		
17442	multiflora	<i>Thun.</i>	many-flowered	[■■■■■]	2	jl.au	Y	C. G. H.	1818.	C	p.1		
17443	uniflora	X.	single-flowered	[■■■■■]	3	jl.au	Y	C. G. H.	1812.	C	p.1	Pluk.m.	414.7
17444	carnata	Z.	fleshy-leaved	[■■■■■]	3	my.jn	Y	C. G. H.	1795.	C	p.1	Bot mag.	1289
17445	crassifida	A Rep.	thick-leaved	[■■■■■]	2	jl.au	Y	C. G. H.	1800.	C	p.1	Bot.rep.	353
17446	affinis	7%ir.	kindred	[■■■■■]	3	nj.l	Y	C. G. H.	1822.	C	tp		
17447	genistoides	<i>L.</i>	Genista-like	[■■■■■]	2	jl.au	Y	C. G. H.	1816.	C	p.1		
17448	squarrosa	7%oo.	squarrose	[■■■■■]	2	jl.au	Y	C. G. H.	182a	C	p.1		
17449	lalio'riæa	/ferf.	Gallium-like	[■■■■■]	2	jl.au	Y	C. G. H.	1817.	C	p.1		
17450	subulata	<i>Thun.</i>	awl-leaved	[■■■■■]	2	jl.au	Y	C. G. H.	1789.	C	p.1		
17451	globulsa	J. /te/.	globular	[■■■■■]	3	nj.l	O	C. G. H.	1802.	S	p.1	Bot. rep.	510
17452	/aricina	Dec.	Lurchn-leaved	[■■■■■]	2	jl.au	Y	C. G. H.	1823.	C	p.1		
17453	aranebsa	<i>L.</i>	cobweb	[■■■■■]	2	nj.l	Y	C. G. H.	1795.	S	p.1	Bot mag.	829
17454	Indica	£.	Indian	[■■■■■]	1	lau	R	C. Indies	1759.	S	p.1	R.mal.	9.37
17453	ciliaris	<i>L.</i>	ciliated	[■■■■■]	1	lau	Y	C. G. H.	1799.	C	p.1	Bot. mag.	2233
17456	quinquefolia	Z.	five-leaved	[■■■■■]	1	jl.au	Y	C. G. H.	1816.	C	p.1	Pluk.al.	278.4
17457	candidans	H. K.	whitish	[■■■■■]	1	nj.l	Y	C. G. H.	1774.	C	lp		
17458 ¹	arglinea	X.	silver-leaved	[■■■■■]	2	jl.au	Y	C. G. H.	1759.	C	p.1		

II. TRIFOLIA^XUE.—*Turners subsessile*; leaves trifoliate, scarcely fasciculate,

17459 <i>sericea</i> Thun.	silky	sh. LH or	2	jlu	Y	C. G. H. 1816.	C. p.l.	Pluk.al 278.4
17460 <i>callosa</i> L.	callous	m. P'	3	J ^l au	Y	C. G. H. 1812.	c 1P	Bot mag. 2329
17461 <i>muconata</i> L.	muconate		3	injl	Y	C. G. H. 1796.		
17462 <i>pedunculata</i> Herit.	/on'-peduncled		6	J ^l au	Y	C. G. H. 1775.		Bot mag. 344

* 1960. CPLEX L.	FURZE.	[Ac. a point, <i>Celt.</i> ; leaves sharp.)	Leg.	Pap.	Lot.	Gents.	4.
17463 euro ^{MB} 'a L.	Kuropéan	<i>m</i> ag 6 f.ja Y Britain san.he. 8 co Eng. bot 742					
17464 nång Sm.]	dwarf	<i>m</i> or 2 aud Y Britain dry he. S co Eng. bot. 743					
17465 hibernica G. Don	<i>i</i> Irish	<i>SH</i> or 6 aud Y Ireland ... S co					
17466 provincialis Loi.	Provence	<i>m</i> or 4 jl.au Y S. Europe 1823. S co	Loi.	no.	6.2		

1961. STAUROCA'NTHUS Lk. STAUCANTHUS. (*Stauros*, cross, *akantha*, spine.) Leg. Pap. Löt. Genit. 1.
17467 *aphyllus* Lk. . . leafless * or 3 ... Y Spain 1823. S p.1
1st flex mitis *Horn* *genistoides* Brot.

1902. SPA'RTIUM L.	BROOBL (<i>Sparton. cordage</i> : use in early ages.)	Leg.	Pap.	Löt.	Gents.	1.
17468 <i>lündicum</i> L. 2f15repnfeno	<i>Rush or Spanish*</i> ec 6 jls. Y & Europe 1548. S co Bot mag. 85 double-flowered A or 6 jls. Y S. Europe 1548. S co					

1963. GENI'STA L. **GENISTA.** (*Gen*, a small bush, *Celt.*) **Leg. Pap. LbL Gents.** 48.-76.

I. INB'UMES.—Unarmed; leaves all or for the most part trifoliate.				Lg. eels	
		(Gen., a small bush, Cet.)		(Gen., a small bush, Cet.)	18. 70.
17469	paryflora Dec.	small-flowered	H	or 3 my.au	Y S. Europe 1817. S p.l Yen. eels 87
		Jto&rtium parviflormum	Ven.		
17470	clavata Poir.	clavate	■ i lor	3 my.au	Y Mogadore 1812. C p.l Ven.cels? 7
		Spartium sericeum	Ven.		
17471	canariensis L.	Canary	■ l Jfira	2 my.8	Y Canaries 1656. S s.l Bot.reg.217
17472	candidans L.	whitish	■ or	2 apil	Y Spain 1735. C 8.1 Den.br. 80
17473	patens Dec.	spreading	■ k or	2 apjl	Y Spam ... S s.p Cav.ic.2.170
		Spartium pátens Cav.			
17474	angulata Haji.	angulate	m or	3 my.jn	Y Maryland 1739. C p.1
		Spartium angulatum W.			
17475	/imflia L.	Flax-leaved	m or	3 jajn	Y Spain 1739. C p.1 Bot mag. 44
		Spt&rtium iinifolium Desf.			
17476	triquetra H. K.	triangular	* or	3 my.jn	Y Corsica 1770. C s.p Bot. mag. 31
17477	bracteolata Lk.	bracteolate	i j o r	2 mr.my	Y 1823. C co
17478	umbellata Poir.	umbellated	sH or	3 ap.jn	Y Barbary 1799. C p.1 Desf. at 2.18
		A'ptium umbellatum Desf.			
17479	radiata Sco.	rayed	■ or	lj jn.jl	Y Italy 1758. S s.p Bot mag. 22
		E'rtium radiatum L.			

II. Spixbaa.— <i>Spiny ; leaves trifoliate.</i>										
17480	hdrrida <i>Dec.</i>	horrid	1	or	3	myjl	Y	Pyrenees	1821.	L p.1 Oil. eu. 2.239.ic
17481	luroidica <i>L.</i>	Portuguese	*	or	2	mr.mn	Y	Portugal	1771.	C co Bot. rep. 419
17482	ftrrox <i>Poir.</i>	fierce	*	_J or	ljjnjl	Y	Barbary	1800.	C p.1 Bot.reg.368	
17483	Spartium ftrrox <i>Desf.</i>	The Barbed Spanish Broom	1	or	2	mlb	Y	Sardinia	C p.1	Bot. tab. 54

17483	<i>triacanthos</i>	<i>Brot.</i>	three-spined	•	or	2	myjl	Y	Spain	...	C	p.1	<i>Brot ph. 54</i>	
i. SIMPKICHTFOLLS.— <i>Spiny j leaves simple.</i>														
17484	<i>sylvestris</i>	<i>Sco.</i>	wood	*	or	2	jn.jl	Y	Hungary	1818.	L	p.1	Jac.ic.557	
17485	<i>Scorpius</i>	<i>Dec.</i>	Scorpion	A	or	4	mr.ap	Y	S. Europe	1570.	S	p.1	Den. br. 78	
	<i>Spärtfötter</i>	<i>Scorpius L.</i>												
17486	<i>hispanica</i>	<i>L.</i>	Spanish	•	or	2	jn.jl	Y	Spain	1759.	C	co	Cav. ic. 3.21	
17487	<i>itAlica</i>	<i>Lod.</i>	Italian	i	or	3	jn.jl	Y	Italy		L	i:o		
17488	<i>anglica</i>	<i>L.</i>	English	Petty Whin	*	or	2	my.jn	Y	Britain	moi.	he.	S co	Eng. bot 132
17489	<i>germanica</i>	<i>Z.</i>	German	i	or	2	in.au	Y	Germany	1773.	L	co		

- | | | | | | | | | | | | | |
|-------|--|---------------------------|---------|----|----------|-------|-----------|-----------|---------|-----|----------------|----------------|
| 17498 | sphrocr̄pa <i>Lam.</i> | round-podded | A i_Jor | 4 | jnjl | Y | S. Europe | 1731. | S | p.l | Ren. spec. 33 | |
| | <i>Spaxtium</i> sphrocr̄firpon <i>L.</i> | | | | | | | | | | | |
| 17499 | athme'nBis <i>Dec.</i> | -ffitna | A or | 3 | jnjl | Y | Sicily | 1816. | L | p.l | Bot. mag. 2674 | |
| | <i>Spartium</i> ethnense <i>Btv.</i> | | | | | | | | | | | |
| 17500 | tetragona <i>Bes.</i> | four-angled | -* | or | 1 | jn,jl | Y | Podalia | 1822. | L | p.l | |
| 17501 | scarius <i>Viv.</i> | srarious | il | or | 6 | jnjl | Y | Italy | 1821. | L | co | |
| | <i>januensis</i> <i>Viv.</i> genuīens Pers. | | | | | | | | | | | |
| 17502 | anxantica <i>Ten.</i> | Anxantic | m | or | 4 | jnjl | T | Italy | 1818. | L | 81 | Fl. nap. 2.66 |
| 17503 | tinctoria <i>L.</i> | dyer's <i>Green-weedm</i> | or | 3 | jn,au | T | Britain | dry. pa. | S | co | Eng. bot 44 | |
| 17504 | sibirica <i>L.</i> | Siberian | * | or | 2 | jn,au | T | Siberia | 1785. | L | co | Jac VIQ. 2.190 |
| 17505 | polygabefbla f.w. | Polygala-leaved* | or | 3 | jn,au | Y | Spain | 1820. | L | co | | |
| 17506 | ftoridalaL. | flowery | H | or | 6 | jn,au | T | Spain | 1752. | S | co | |
| 17507 | mlatrica <i>Pol.</i> | Mantic | m | or | 3 | jn,au | T | S. Europe | 1816. | L | co | Fl. ver. 2.4.2 |
| 17508 | ovata <i>Kit.</i> | ovateJeaved | A | or | 3 | jn,au | T | Hungary | 1816. | C | co | W. & K 83 |
| 17509 | pätnula <i>Bieb.</i> | spreading | A | or | 3 | jn,au | E | Caucasus | 1818. | L | co | |
| 17510 | triangularis <i>W.</i> | three-angled | A | or | 2 | myjn | Y | Hungary | 1815. | L | co | W. & K. 153 |
| 17511 | sagittalis <i>L.</i> | anovi,jointed | A | or | imy,jn | Y | Germany | 1570. | L | co | Jac. au. 3.209 | |
| 17512 | diffusa <i>W.</i> | diffuse | A | or | 3 | myjn | Y | Italy | 1816. | L | co | Jacic 3.553 |
| 17513 | procumbens <i>Kit.</i> | procumbent | JU | or | 1} jn,au | Y | Hungary | 1816. | L | co | | |
| | <i>prostrata</i> <i>Lam.</i> | | | | | | | | | | | |
| 17514 | dec&nibens <i>Durb.</i> | decumbent | & | or | J | myjn | Y | Burgundy | 1775. | L | p.l | Bot. cab. 718 |
| 17515 | pilosa <i>L.</i> | pilose <i>Green-weed</i> | A | or | 6 | myjn | Y | England | san.he. | S | co | Eng. bot 208 |
| 17516 | pilocarpa <i>Lk.</i> | hairy-fruited | m | or | 4 | jnjl | Y | | 182a | S | p.l | |

* 1964. CYTISUS L. CYTISUS. (First found at *Cythnus*, one of the Cyclades.) Leg. Pap. Löt. Gents. 34.-44.

- | | | | | | | | |
|-------|----------------------------|----------------|------------|---------|-----------|-------|----------------|
| 17517 | nubigenus Lk. | cloud-born | • u.j or 6 | my.jn W | Teneriffe | 1779. | C p.l |
| | Spärtium nublgenum/f. K. | | | | | | |
| 17518 | aums L.f. | white Portugal | A or 8 | my W | Portugal | 1752. | S co Duh.ar.23 |
| | Spärtium multiflorum H. K. | | | | | | |
| | Sincamatus | Qesh-colored | A or 8 | my F | Portugal | 1818 | Ş 8i |

II. LABU'NUM. — *Calyx campanulate; legumen many-seeded, not dilated at the upper sutures flowers yellow.*

- | | | | branches unarmed, leafy. | | | | | | |
|-------|--------------|---------------|---|--------|--------|--------|-----------|-------|--------------------|
| 17519 | Laburnum | <i>L.</i> | common Laburnum? | tm 15 | my,jn | Y | Switzerl. | 1596. | S co Bot mag. 176 |
| 17520 | alptnus | <i>Mil.</i> | alpine Scotch Laburn. | tm 30 | jn | Y | Europe | 1596. | S co |
| 17521 | nfricans | <i>L.</i> | black-roofed | A or 3 | jnjl | Y | Austria | 1730. | S 81 Bot reg. 802 |
| 17522 | sessiliflius | <i>L.</i> | com. sessilc-lvd | A or 6 | myjn | Y | Italy | 1629. | S 81 Bot mag. 255 |
| 17523 | triflorus | <i>Herit.</i> | three-flowered | A or 4 | jnjl | Y | Spain | 1640. | S 81 |
| 17524 | patens | <i>L.</i> | spreading | A or 4 | jnjl | Y | Portugal | 1752. | S |
| | | | Spartium pâtens <i>L.</i> not <i>Cao.</i> | | | | | | |
| 17525 | m61lis | <i>IV.</i> | en, | soft | A or | ■ jnjl | Y | | 1818. B.1 |
| 17526 | grandiflorus | <i>Dec.</i> | great-flowered | A or | ■ jnjl | Y | Portugal | 1816. | ■ CO |
| | | | Spartium grandiflorum <i>Brot.</i> | | | | | | |
| 17527 | *copânus | <i>Lk.</i> | common Broom | A or 6 | apjl | Y | England | com. | S co Eng. bot 1339 |
| | | | Spartium jcoparium <i>L.</i> | | | | | | |
| | 2£bus | | white-flouering | A or 6 | apjl | W | England | com. | S co |

III. CALYCC/TOME.—*Calyx campanulate, subbilabiate; legumen thickened towards the upper sutures branches spiny; flowjsrs yellow.*

Sp&rrnib spimbus Brot. not W. Calycotome villosa Lk. Spartium villosum Pour.

- | 17530 proterstum L. f. V. & K. | | white-flowered | | A | or | 4 | infl. | Pa. Y | Hungary | 1806. | S. 81 | Bot. reg. 121 |
|--------------------------------|--|-----------------|--|----|----|---|-------|-----------|-----------|-------|---------------|----------------|
| 17532 Andreia Dec. JV. | | white | | A | or | 4 | jn.jl | S. Europe | Austria | 1792. | S. 81 | Bot mag. 1438 |
| 17533 purpureus Sco. | | purple-flowered | | A | or | 3 | my.au | P | Austria | 1760. | S. 81 | Bot. mag. 1176 |
| 17534 albiflorus | | white-flowered | | A | or | 2 | my.au | W | Austria | 1760. | B1 | |
| 17534 biflorus Herit. | | two-flowered | | A | or | 3 | my.jn | Y | Hungary | 1760. | 8.1 | Bot reg. 308 |
| 17535 elongatus IV. \$ K. | | long-branched | | I | or | 4 | my.jn | Y | Hungary | 1804. | S. 81 | W. & K. 2.183 |
| 17536 multiflorus G. Don | | many-flowered | | I | or | 1 | Y | | Europe | 1800. | 8.1 | |
| elongatus Hort. not | | W. % K. | | - | | | | | | | | |
| 17537 falcatum IV.S(K.) | | Bickle-shaped | | A | or | 3 | jn.au | Y | Hungary | 1816. | S. 81 | Bot cab. 520 |
| 17538 austriacus L. | | Austrian | | A | or | 3 | jn.s | Y | Austria | 1741. | S. s.l | Jac. au. 1.21 |
| 17539 hirsutus L. | | hairy | | A | or | 5 | jn.au | Y | S. Europe | 1739. | S co | Jacob.4.96 |
| 17540 ruthenicus Fis. | | Russian | | A | or | 3 | jn.jl | Y | Russia | 1817. | S co | |
| 17541 sup'inus Jac. | | supine | | *- | or | 1 | my.au | S. Europe | 1755. | S. 61 | Jac. au. 1.20 | |
| 17542 capitatus Jac. | | capitate | | A | or | 3 | jn.jl | Y | Austria | 1774. | S. s.l | Bot. cab. 497 |
| 17543 ciliatus Wahl. | | ciliated | | A | or | 3 | jn.jl | Y | Carpathia | 1817. | S co | |
| 17544 pol'arichus Bieb. | | much-haired | | A | or | 1 | in.jl | V | Tauria | 1818. | S co | |

1734 por *richards Bleo.* much-haired A or I, jn. Y Tauria 1818 S co
 V. 'Lorran'^{Dt\$} — *Tube of calyx short, obconic; upper lip biparted, lower one 3-toothed; corolla scarcely longer than cal*
 •*mrA~ y* * many-stemmed decumbent undershrubs: flowers few, subcapitellate, yellow.

- | | | | | | | | | | |
|-----------------------------|---------------------|---|----|------|---|--------|-------|---|-----------------|
| 1754 <i>Argenteus</i> L. | Bilver-tan'd | A | or | 3 au | Y | France | 1739. | S | <i>If</i> |
| 1756 <i>calycinus</i> Bieb. | <i>larg-calyxed</i> | A | or | 2 au | Y | Tauria | 1820. | S | co Bot cab. 673 |

pauciflorus IV.
Tim manui Walt. dwarf
17548 *pygmaeus* W. pygmy or JSP¹ Levant 1816. Galacea Den. br. 81

- VI.**, *CHIBONA NTHUB*. — *Upper lip of calyx bifid, lower one trifid; lobes acute, length of tube & petals persist.*

legumen oval, much compressed, two-seeded.

- 49 onentahs *Loi.* eastern A or 3 S; Europe 1818. S co
 17550 glomeratus *Bo.J.* D. — Doubtful to what section it belongs.
 glomerated gor 3. Zanzibar 1826. S s.l.

1965. ADENOCA'RPU'S *Dee.*, ADENOCARPUS. 3 (*Aden,* gland, *karpos,* fruit.) Leg. Pap. Lot. Gents. 6.
 17551 hispanicus *Dec.* Spanish*. A or 3 jnjl Y Spain 1816. S co
Ciribus hispanicus *Lam.*

17555 foliosus Dec.	small-leafy	*i__ or	6 my.jn Y	Canaries	1629.	S co	Bot.mag.42
Cytisus fohosus H. K.							
17556 frankeniolifc* Choi.	Frankenia-like	* i_J or	2 apjl Y	Canaries	1815.	C s.l	
Genista viscosa IV.							
* 196a ONCTNIS L. REBTHARROW. (<i>Onos</i> , an ass, <i>onemi</i> , to delight j grateful to.) Leg. Pap.Lbt. Gen. St. —11L							
I. EUONOVII.— <i>Stipules adhering to the petioles.</i>							
i. NA ^V I BIX.— <i>Leaves simple or bipinnate; flowers axillary, pedunculate, yellow.</i>							
17557 criapa L.	curved	« J un	2 jn.au	Spain	1739.	S p.l	Mag. mo. 17
17558 hispanica L.	Spanish	*. J or	1 my.s	Spain	1799.	C p.l	Bot. mag. 2540
17539 vaginalis Vahl	sheathed	n. un	2 jn.au	Emp	1815.	C co	Vcn. eels 32
17560 Mtr.x Dec	Goat-root	n. un	1 my.s	S. Europe	1680.	S p.l	Bot mag. 329
17561 pfnguis L.	greasy	tt. un	If jl	& Europe	1739.	C co	
matrix fl Dec.							
17562 cuspidate Desf.	cuspidate	H un	1J my.s	Algiers	1818.		
17563 picta Deif.	painted	J A or	1 my.jn	Barbary	1820.		
17564 longifolia W.	long-leaved	a. i or	2 jn.au	Tenenne	1816.		
17565 falcate Via.	falcate	a. lor	1 jn.au	S. Europe	1819.		
17566 ramosissima Desf.	branchiest	n. or	i jn.jl	Sicily	1819.		
17567 areparia Dec.	sand	n. un	1	S. France	1819.		
17568 biflora Desf.	two-flowered	O un	I jlau	Barbary	1818.		
17569 visebsa L.	clammy	O un	I jlau	S. Europe	1759.		
17570 brachycarpa Dec.	short-podded	O or	i jn.jl	Spain	1823.		
17571 breviflora Dec.	short-flowered	O or	i jn.jl	S. Europe	1800.		
17572 sicula Cus.	Sicilian	O or	i jn.jl	Sicily	1817.		
17573 pubescens L.	pubescent	O un	1 jn.au	S. Europe	1680.		
17574 ornithopodioides L.	Bird's-foot-like	O un	1 i jlau	Sicily	1713.		
17575 pallida Horn.	pale	O or	I jn jl	P	1820.		
ii. NATR'DIUM.— <i>Leaves simple or trifoliate; flowers axillary, pedunculate, purplish, rarely white.</i>							
17576 rotundifolia L.	round-leaved	* or	2 myjl	Pk	Switzerl.	1570.	C 8.1 Bot. mag. 335
17577 tnbracteata Dec	thre-bracted	1 or	1 my.jl	Pk	S. Europe	1800.	
17578 fruticosa L.	shrubby	m pr	2 my.jn	Pk	S. France	1680.	
17579 tridentata L.	three-toothed	tt. i J or	1 jn.au	P	Spain	1752.	
17580 angustissima Lam.	narrowest	tt. or I	1 myjl	Pk	Spain	1825.	
17581 cenisia L.	Mount Cenis	A un	ijn.au	Pk	Italy	1759.	
17582 fce'tida Schous.	fetid	O or	* jnjl	Pk	Morocco	1818.	
17583 geminiflra Lag.	twin-flowered	O or	i jn.au	P	Spain	1817.	
17584 reclinata L.	recline	Jk O un	jn.au	St	S. Europe	1800.	
17585 Cherleria L.	Cherler's	O un	jn.jl	P	S. Europe	1771.	
17586 pendulosa Desf.	pendulous	O or	I jn.au	P	S. Europe	1818.	
iii. BUGRAVA.— <i>Leaves simple or trifoliate; flowers sessile or on very short peduncles, purple, rarely white.</i>							
17587 arborescens Desf.	arborescent	m or	2 my.jl	R	Barbary	182U	C p.l Desf. at 2.193
17588 altissima Lam.	tallest	■■■ un	3 my.au	R	Italy	1596.	D co Jac. vin. 1.93
hirelna Jac.							
17589 antiquum L.	ancients*	■ A un	1 jn jl	P S	Europe	1790.	D co Lob. ic. 28
spinifera y glabra Dec.							
17590 procarricans Wallr.	procurent	■ A or	1 jnjl	P	Europe	1820.	D co
17591 rcpens L.	creeping	■ A un	1 jn.au	Pk	Europe	...	D co Di. el. 25. 28
arvensis Lam.							
17592 spinosa L.	spiny	■ ■ un	1 J n a u	F	Britain	...	D co Eng.bot682
arvensis E. B.							
2 alba	white-flowered	£ A un	1 jn.au	W	Britain	sea si.	D co
17593 mitissima L.	mildest	O un	I ■■■	R	Portugal	1732.	S co
mm diffusa Ten.	diffuse	O or	I ■■■	P	Italy	1820.	S co
17595 hirta Desf.	hairy *	■■■ or	1 jnjl	B	S Europe	1816.	S co
17596 alopecuroides L.	Foxteil-Uke	un	5 jlau	P	Portugal	1696.	S co
17597 monophylla Desf.	one-leaved	O un	i jnau	P	Barbary	1800.	S co
17598 oligoprlfla Desf.	few-leaved	O or	1^my.s	W	Naples	1823.	S p.l
iv. BUGRANO ^V DES.— <i>Leaves simple or trifoliate; flowers sessile or subsessile, yellow.</i>							
17599 arragonensis Asso	Arragonese	M or 11	my.jl	Y	Spain	1816. C r>1	Mag. mo. 21
17600 capitate Cav.	capitate	■■■ i	jn.au	Y	Spain	1820.	
17601 striata Gou.	striated	■■■ i	jn.jl	Y	Spain	1817.	
recline Lam.							
17602 Columna's W.	Columna's	■■■ un	1 jnjl	Y	S. Europe	1732.	
17603 minutissima L.	smallest	■■■ or	1 jnjl	Y	S. France	1818.	
barbate Cav.							
17604 variegata L.	variegated	O un	jl.au	Y	Spain	1784.	S co Desf. at £.185
II. LOTONOI ^V DEB.— <i>Stipules scarcely or not adhering to the petioles; leafy, as in Lbtusj but with the stamens monadelphous, as in the rest o/Onkgs.</i>							
17605 parviflora Berg.	small-flowered	Q or	1 jnjl	■ C. G. H.	1818.		
17606 villusa Tjuu.	twin-flowered	I Alun	I J j U	Pk	C. G. H.	1820.	
17607 geminata H. K.	umbellate	AJ ■	1 jnjl	C. G. H.	1787.		
17608 umbellata L.	smooth	■■■ or	J jn.jl	C. G. R.	1816.		
17609 glabra Thun.	Cape	■■■ i	jn.jl	C. G. H.	1824.		
17610 capitata L.	emarginated	■■■ or	i jn.jl	C. G. H.	1800.		
17611 emarginata Boj.	drooping-poddedn.	■■■ un	2 jl.s	Mauritius	1825.		
17612 cernua L.				T	C.G. H. 1774.		
					co	Com. h. 2.82	
1967. REQUIETNJI* Dec. RBQUIENIA. (Af. Requien, of Avignon, a Fr. bot.) Leg.Pap.Lbt.Genls. 1—SL							
17613 obcordata Dec.	obeordate	JL Q or	1 JI	Y	Senegal	1825.	C s.p Dec. leg. 4.27
Podaff rin obcordata Lam.							
* 1968. JKTHY'LLIS L. KIDNEY VETCH. (<i>Anthos</i> , flower, <i>iyoylos</i> , down") Leg. Pap. Et. Genis. 19.—25.							
I. DoRYCHNOIDES.— <i>Calyx scarcely inflated; legemens l-seeded; herbs peppnial; roots smffrutescenti head* of flowerspedunculated, leafless; leavesgrinnate.</i>							
17614 Gerards L.	Gerard's	^t A or	i ^g au	W	Provence	1806.	S co Ger. gal. 18
17615 onobrychitides Cav.	St. Foix-like	^ A or	J jnjl	Y	Spain	1817.	D s.1 Cav. ic. 2.150
II. A8PALATHO ^N DE9.— <i>Calyx scarcely inflated; legumen 1 or 2 seeded; flowers in interrupted spikes, or in the axils of the upper leaves; leaves trifoliolate or simple; humble shrubs.</i>							
17616 cytiMftdes L.	tususlike	« Li J or	g apjl	W	Spain	1731.	C p.1 Bar. ic. 1182
17617 Hermifinnw? L.	crman's	«. « J or	IJ apjl	Y	Levant	1739.	C p.1 Alp.ex.Si6

- 17618 Apfilathi Dec. Aspalathus-like. \square or 1 ap.jl Y C. G. H. 1824. E \square Bot. cab. 1169
 17619 tenuifolia L. fine-leaved SL \backslash J or 1* my.jl Y C. G. H. 1818. E \square
- III. ERINA\RA.—Calyx after flowering evidently inflated; legumen 1 or 2 seeded, compressed, lanceolate longer than calyx; shrubs much branched, spiny, nearly leafless.
- 17620 erinacea L. prickly \square or 1 ap.my P Spain 1759. C s.p Bot mag. 676
 17621 echinata Hort. hedgehog n. \square or 1 ap.jl P? S. Europe? ... C p.l
- IV. VuL\KA'aia.—Calyx after flowering evidently inflated; legumen 1 or 2 seeded, shorter than calyx or equals heads of flowers bracteate; shrubs or herbs with pinnate leaves.
- 17622 Barba J. D. IL. Jupiter's Beard * i. J or 3 mr.my Pa. Y S. Europe 1640. C p.l Bot. mag. 1927 *
- 17623 heterophylla L. various-leaved a. \square J or 1 jn H " Pk & Europe 1768. C p.l
 17624 montana L. mountain \wedge A or \backslash jn,jl P S. Europe 1759. S s.l Bot. cab. 578
 2 alba white \wedge ^ or \backslash jn,jl W S. Europe 1818. S s.l
 17625 Vulneraria L. common Woundwort \square A or i my.au Y Britain ch.pas. S s.l Eng.bot104
 Srdbra red-flowered \wedge A or i my.au R Britain ch.pas. S s.l
 3 albiflbra white-flowered 5 A or i my.au W Britain ch.pas. S s.l
 rtistica Mil.
- 17626 Dillenii Schultz. Dillenius's \wedge A or i my.au R S. Europe 1816. S s.l Di.eL320.413
 Vulneraria y rubriflora Dec.
- 17627 polypylia Kit. many-leaved \wedge A or 1 my.jn Y S. Europe 1816. S s.l
 Vulneraria 8 polyphyllea Dec.
- 17628 alptna G. Don hairy alpine 5t A or i my.au Y Britain sc.mo. S s.l
- V. COENICIVVA.—Calyx inflated; legumen smooth, two or many celled.
- 17629 tetraphylla L. four-leaved O or 1 jl.au W S. Europe 1630. S co Bot. mag. 108
 17630 hamsa Desf. hooked O or ijl.au Pa.Y Barbary 1821. S s.l
 cornicina Poir.
- 17631 cornicina L. crow O or 1 jl.au W Spain 1759. S p.l Car. i cl. 39.
 17632 /otrides L. Lotus-like O or i jn,jl Y Spain 1739. S co Cav. i d. 40
1969. CYAMO'PSIS Hec. CYAMOPSIS. (*Kyamos*, bean, *opsis*, resemblance.) Legum. Pap. Lbt. Clit. 1.
 17633 psoralea Mdes Dec. Psoralia-like O or 1 jl.au P Arabia 181& S s.l Cav. ic. 1.59
 X)6ichos psoraloides Lam. /abaeiSrmis Herit. Psoralea tetragonoblobus L. I.upnus trifoliatus Cav.
1970. PUERA^RIV* Dec. PUERARU. (M. M. N. Puerari, professor at Copenhagen.) Leg. Lbt. Clit. 2.
 17634 tuberota Dec. tuberose J. J or 3 ... Y E. Indies 1806. D s.l
 Jfedf larum tubersum Box.
- 17635 Wallichii Dec. Wallich's A. LJ or 3 ... Y Nepal 1826. C s.l Dec. leg. 43
- " 1971. DA?LEA L. DALEA (*Thomas Dale*, an English botanist.) Legum. Pap. Lbt. Clitbr. 15.—35.
 17636 laxiflbra Ph. loose-flowered \square A pr 6 jl.au W Louisiana 1811. D co
 Psoralea laxiflbra Poir.
- 17637 atrea Nut. golden \square A pr S jlau Y Louisiana 1811. D co
 17638 filocerooides Nut. Foxtail-like \square \square 1 jl.au Pa.B Mississippi 1812. S co Mic. am. 2. 38
 Psoralea Dalea L.
- 17639 CliffbrWna W. Cliffbr's 8 pr jlj.au B VeraCruz 1737. S co H.cL22
 17640 fagopus IV. Hare's-foot 4 o.n V Mexico 1780. S co Cav. ic. 1.86
 Psoralea leporina H. K.
- 17641 sericea La* silky \square BSD or jlj.au V Mexico 1824. C s.p Kth. mim. 48
 17642 tuberculata Lag. tuberculated \square D \square 2 jl.au P Mexico 1824. C s.p
 17643 citriodora W. Citron-scented O or 1 o.n Pk N. Spain 1780. S co Cav. ic. 3.271
 17H44 mutabilis W. changeable \square iAI or ljo.n P Mexico 1818. C co Bot mag. 2486
 17645 bicolor W. en. two-colored \square C3 or 2 o.n Y.B S. Amer. 1817. C co Hook. ex. fl. 43
 17646 nutans W. nodding \square E3 or 1 jl.au Vi Mexico 1824. C s.p Fl. mcx. ic. in.
 17647 reclinata W. • ccilinate C C3 or ij.l.au Vi Mexico 1820. C s.p
 17648 Thouin's Thouin's QJ or 1 jl.au B 1816. S s.p Sch. mo. 9
 17649 phymatodes W. impostume-like. I J or li jl.au W.v Caraccas 1819. C s.p Jac.ic.3.563
 17650 cmneaphylla TV. nine-leaved I Q p r 5 jl.au Pk W. Indies 1772. S co
- * 1972. GALE^GA Tou. GOAT'S RUE. (*Gala*, milk; increases in animals eating it.) Leg. Pap. Lbt. Clit. 5.
 17651 officinalis L. officinal \wedge A or 4 jns B Spain 1568. D co Mil. ic. 137
 2 alba white-flowered \square \square \square Spain D co
 17A12 persica Pers. Persian \square \square \square Persia 1826. D co Sw. fl. gar. 244
 17653 biloba Swt. two-lobed \square \square \square 1823. D s.p Sw. fl. gar. 216
 17K54 tricolor Hook. three-colored \square \square \square 1823. D \square Hook. ex. fl. ic.
 17655 orientalis Lam. oriental \square Levant 1801. D p.l Bot mag. 2192
1973. AMO'RPHA L. BASTARD INDIGO. (Apriv., \square , form; deformity of cor.) Leg. Pap. Lbt. Clit. 8.
 17656 fruticosa L. shrubby II or 6 jn,jl P Carolina 1724. C s.p Bot recg. 427
 2 emarginata \square or 6 jn,jl P Carolina 1724. C s.p
 17657 microphilla Ph. small-leaved * J fra 2 jl.au P Missouri 1811. C s.p
 17658 herbacea Walt. herbaceous \square Ai or 3 jn,jl B Carolina 1803. C s.p Bot cab. 689
 pubescens W.
 17659 canescens Nut. canescent SI \square or 3 jlau B Missouri 1812. C s.p
 17660 nana Nut. dwarf \square \square jlj.au B Missouri 1811. C s.p
 17661 cruceo-lanata Wat. yellow-woollled a. or 5 jl.au P N. Amer. 1820. C s.p Den. br. 139
 17662 Lewis* Lo. C. Lewis's \square \square 3 jlau P N. Amer. 1820. C s.p
 17663 gKibra Desf. smooth SI \square or 3 jlau P N. Amer. 1810. C s.p
 17664 fragrans Sat. fragrant Si \square or 3 jn,jl P N. Amer. 1800. C s.p Sw. fl. gar. 241
 nana B. M. not of others.
1974. MULLER^ L. MULLERA. (O.F.Mutter, a Dane, one of editors of Flora Danica.) Leg. Pap. Lbt. CM. 1.
 17665 momiformis L. bracelet-formed* or 20 ... Y Guiana 1792. C Lp Men sur. 35
- ~~—V. 21. 22. 23. 24. 25. 26. 27. 28. 29. 30. 31. 32. 33. 34. 35. 36. 37. 38. 39. 40. 41. 42. 43. 44. 45. 46. 47. 48. 49. 50. 51. 52. 53. 54. 55. 56. 57. 58. 59. 60. 61. 62. 63. 64. 65. 66. 67. 68. 69. 70. 71. 72. 73. 74. 75. 76. 77. 78. 79. 80. 81. 82. 83. 84. 85. 86. 87. 88. 89. 90. 91. 92. 93. 94. 95. 96. 97. 98. 99. 100. 101. 102. 103. 104. 105. 106. 107. 108. 109. 110. 111. 112. 113. 114. 115. 116. 117. 118. 119. 120. 121. 122. 123. 124. 125. 126. 127. 128. 129. 130. 131. 132. 133. 134. 135. 136. 137. 138. 139. 140. 141. 142. 143. 144. 145. 146. 147. 148. 149. 150. 151. 152. 153. 154. 155. 156. 157. 158. 159. 160. 161. 162. 163. 164. 165. 166. 167. 168. 169. 170. 171. 172. 173. 174. 175. 176. 177. 178. 179. 180. 181. 182. 183. 184. 185. 186. 187. 188. 189. 190. 191. 192. 193. 194. 195. 196. 197. 198. 199. 200. 201. 202. 203. 204. 205. 206. 207. 208. 209. 210. 211. 212. 213. 214. 215. 216. 217. 218. 219. 220. 221. 222. 223. 224. 225. 226. 227. 228. 229. 230. 231. 232. 233. 234. 235. 236. 237. 238. 239. 240. 241. 242. 243. 244. 245. 246. 247. 248. 249. 250. 251. 252. 253. 254. 255. 256. 257. 258. 259. 260. 261. 262. 263. 264. 265. 266. 267. 268. 269. 270. 271. 272. 273. 274. 275. 276. 277. 278. 279. 280. 281. 282. 283. 284. 285. 286. 287. 288. 289. 290. 291. 292. 293. 294. 295. 296. 297. 298. 299. 300. 301. 302. 303. 304. 305. 306. 307. 308. 309. 310. 311. 312. 313. 314. 315. 316. 317. 318. 319. 320. 321. 322. 323. 324. 325. 326. 327. 328. 329. 330. 331. 332. 333. 334. 335. 336. 337. 338. 339. 340. 341. 342. 343. 344. 345. 346. 347. 348. 349. 350. 351. 352. 353. 354. 355. 356. 357. 358. 359. 360. 361. 362. 363. 364. 365. 366. 367. 368. 369. 370. 371. 372. 373. 374. 375. 376. 377. 378. 379. 380. 381. 382. 383. 384. 385. 386. 387. 388. 389. 390. 391. 392. 393. 394. 395. 396. 397. 398. 399. 400. 401. 402. 403. 404. 405. 406. 407. 408. 409. 410. 411. 412. 413. 414. 415. 416. 417. 418. 419. 420. 421. 422. 423. 424. 425. 426. 427. 428. 429. 430. 431. 432. 433. 434. 435. 436. 437. 438. 439. 440. 441. 442. 443. 444. 445. 446. 447. 448. 449. 450. 451. 452. 453. 454. 455. 456. 457. 458. 459. 460. 461. 462. 463. 464. 465. 466. 467. 468. 469. 470. 471. 472. 473. 474. 475. 476. 477. 478. 479. 480. 481. 482. 483. 484. 485. 486. 487. 488. 489. 490. 491. 492. 493. 494. 495. 496. 497. 498. 499. 500. 501. 502. 503. 504. 505. 506. 507. 508. 509. 510. 511. 512. 513. 514. 515. 516. 517. 518. 519. 520. 521. 522. 523. 524. 525. 526. 527. 528. 529. 530. 531. 532. 533. 534. 535. 536. 537. 538. 539. 540. 541. 542. 543. 544. 545. 546. 547. 548. 549. 550. 551. 552. 553. 554. 555. 556. 557. 558. 559. 559. 560. 561. 562. 563. 564. 565. 566. 567. 568. 569. 570. 571. 572. 573. 574. 575. 576. 577. 578. 579. 580. 581. 582. 583. 584. 585. 586. 587. 588. 589. 589. 590. 591. 592. 593. 594. 595. 596. 597. 598. 599. 599. 600. 601. 602. 603. 604. 605. 606. 607. 608. 609. 609. 610. 611. 612. 613. 614. 615. 616. 617. 618. 619. 619. 620. 621. 622. 623. 624. 625. 626. 627. 628. 629. 629. 630. 631. 632. 633. 634. 635. 636. 637. 638. 639. 639. 640. 641. 642. 643. 644. 645. 646. 647. 648. 649. 649. 650. 651. 652. 653. 654. 655. 656. 657. 658. 659. 659. 660. 661. 662. 663. 664. 665. 666. 667. 668. 669. 669. 670. 671. 672. 673. 674. 675. 676. 677. 678. 679. 679. 680. 681. 682. 683. 684. 685. 686. 687. 688. 689. 689. 690. 691. 692. 693. 694. 695. 696. 697. 698. 699. 699. 700. 701. 702. 703. 704. 705. 706. 707. 708. 709. 709. 710. 711. 712. 713. 714. 715. 716. 717. 718. 719. 719. 720. 721. 722. 723. 724. 725. 726. 727. 728. 729. 729. 730. 731. 732. 733. 734. 735. 736. 737. 738. 739. 739. 740. 741. 742. 743. 744. 745. 746. 747. 748. 749. 749. 750. 751. 752. 753. 754. 755. 756. 757. 758. 759. 759. 760. 761. 762. 763. 764. 765. 766. 767. 768. 769. 769. 770. 771. 772. 773. 774. 775. 776. 777. 778. 779. 779. 780. 781. 782. 783. 784. 785. 786. 787. 788. 789. 789. 790. 791. 792. 793. 794. 795. 796. 797. 797. 798. 799. 799. 800. 801. 802. 803. 804. 805. 806. 807. 808. 809. 809. 810. 811. 812. 813. 814. 815. 816. 816. 817. 818. 819. 819. 820. 821. 822. 823. 824. 825. 826. 827. 828. 829. 829. 830. 831. 832. 833. 834. 835. 836. 837. 838. 839. 839. 840. 841. 842. 843. 844. 845. 846. 847. 848. 849. 849. 850. 851. 852. 853. 854. 855. 856. 857. 858. 859. 859. 860. 861. 862. 863. 864. 865. 866. 867. 868. 869. 869. 870. 871. 872. 873. 874. 875. 876. 877. 878. 879. 879. 880. 881. 882. 883. 884. 885. 886. 887. 888. 889. 889. 890. 891. 892. 893. 894. 895. 896. 897. 897. 898. 899. 899. 900. 901. 902. 903. 904. 905. 906. 907. 908. 909. 909. 910. 911. 912. 913. 914. 915. 916. 916. 917. 918. 919. 919. 920. 921. 922. 923. 924. 925. 926. 927. 927. 928. 929. 929. 930. 931. 932. 933. 934. 935. 936. 937. 938. 939. 939. 940. 941. 942. 943. 944. 945. 946. 947. 947. 948. 949. 949. 950. 951. 952. 953. 954. 955. 956. 957. 958. 958. 959. 960. 961. 962. 963. 964. 965. 966. 967. 967. 968. 969. 969. 970. 971. 972. 973. 974. 975. 976. 977. 978. 979. 979. 980. 981. 982. 983. 984. 985. 986. 987. 987. 988. 989. 989. 990. 991. 992. 993. 994. 995. 995. 996. 997. 997. 998. 999. 999. 1000. 1000. 1001. 1002. 1002. 1003. 1004. 1004. 1005. 1006. 1006. 1007. 1007. 1008. 1008. 1009. 1009. 1010. 1010. 1011. 1011. 1012. 1012. 1013. 1013. 1014. 1014. 1015. 1015. 1016. 1016. 1017. 1017. 1018. 1018. 1019. 1019. 1020. 1020. 1021. 1021. 1022. 1022. 1023. 1023. 1024. 1024. 1025. 1025. 1026. 1026. 1027. 1027. 1028. 1028. 1029. 1029. 1030. 1030. 1031. 1031. 1032. 1032. 1033. 1033. 1034. 1034. 1035. 1035. 1036. 1036. 1037. 1037. 1038. 1038. 1039. 1039. 1040. 1040. 1041. 1041. 1042. 1042. 1043. 1043. 1044. 1044. 1045. 1045. 1046. 1046. 1047. 1047. 1048. 1048. 1049. 1049. 1050. 1050. 1051. 1051. 1052. 1052. 1053. 1053. 1054. 1054. 1055. 1055. 1056. 1056. 1057. 1057. 1058. 1058. 1059. 1059. 1060. 1060. 1061. 1061. 1062. 1062. 1063. 1063. 1064. 1064. 1065. 1065. 1066. 1066. 1067. 1067. 1068. 1068. 1069. 1069. 1070. 1070. 1071. 1071. 1072. 1072. 1073. 1073. 1074. 1074. 1075. 1075. 1076. 1076. 1077. 1077. 1078. 1078. 1079. 1079. 1080. 1080. 1081. 1081. 1082. 1082. 1083. 1083. 1084. 1084. 1085. 1085. 1086. 1086. 1087. 1087. 1088. 1088. 1089. 1089. 1090. 1090. 1091. 1091. 1092. 1092. 1093. 1093. 1094. 1094. 1095. 1095. 1096. 1096. 1097. 1097. 1098. 1098. 1099. 1099. 1100. 1100. 1101. 1101. 1102. 1102. 1103. 1103. 1104. 1104. 1105. 1105. 1106. 1106. 1107. 1107. 1108. 1108. 1109. 1109. 1110. 1110. 1111. 1111. 1112. 1112. 1113. 1113. 1114. 1114. 1115. 1115. 1116. 1116. 1117. 1117. 1118. 1118. 1119. 1119. 1120. 1120. 1121. 1121. 1122. 1122. 1123. 1123. 1124. 1124. 1125. 1125. 1126. 1126. 1127. 1127. 1128. 1128. 1129. 1129. 1130. 1130. 1131. 1131. 1132. 1132. 1133. 1133. 1134. 1134. 1135. 1135. 1136. 1136. 1137. 1137. 1138. 1138. 1139. 1139. 1140. 1140. 1141. 1141. 1142. 1142. 1143. 1143. 1144. 1144. 1145. 1145. 1146. 1146. 1147. 1147. 1148. 1148. 1149. 1149. 1150. 1150. 1151. 1151. 1152. 1152. 1153. 1153. 1154. 1154. 1155. 1155. 1156. 1156. 1157. 1157. 1158. 1158. 1159. 1159. 1160. 1160. 1161. 1161. 1162. 1162. 1163. 1163. 1164. 1164. 1165. 1165. 1166. 1166. 1167. 1167. 1168. 1168. 1169. 1169. 1170. 1170. 1171. 1171. 1172. 1172. 1173. 1173. 1174. 1174. 1175. 1175. 1176. 1176. 1177. 1177. 1178. 1178. 1179. 1179. 1180. 1180. 1181. 1181. 1182. 1182. 1183. 1183. 1184. 1184. 1185. 1185. 1186. 1186. 1187. 1187. 1188. 1188. 1189. 1189. 1190. 1190. 1191. 1191. 1192. 1192. 1193. 1193. 1194. 1194. 1195. 1195. 1196. 1196. 1197. 1197. 1198. 1198. 1199. 1199. 1200. 1200. 1201. 1201. 1202. 1202. 1203. 1203. 1204. 1204. 1205. 1205. 1206. 1206. 1207. 1207. 1208. 1208. 1209. 1209. 1210. 1210. 1211. 1211. 1212. 1212. 1213. 1213. 1214. 1214. 1215. 1215. 1216. 1216. 1217. 1217. 1218. 1218. 1219. 1219. 1220. 1220. 1221. 1221. 1222. 1222. 1223. 1223. 1224. 1224. 1225. 1225. 1226. 1226. 1227. 1227. 1228. 1228. 1229. 1229. 1230. 1230. 1231. 1231. 1232. 1232. 1233. 1233. 1234. 1234. 1235. 1235. 1236. 1236. 1237. 1237. 1238. 1238. 1239. 1239. 1240. 1240. 1241. 1241. 1242. 1242. 1243. 1243. 1244. 1244. 1245. 1245. 1246. 1246. 1247. 1247. 1248. 1248. 1249. 1249. 1250. 1250. 1251. 1251. 1252. 1252. 1253. 1253. 1254. 1254. 1255. 1255. 1256. 1256. 1257. 1257. 1258. 1258. 1259. 1259. 1260. 1260. 1261. 1261. 1262. 1262.~~

- | | | | | | | | | | |
|---|---------------------|--|--|--------------|---------|---------|--|-------------------------|------------------------|
| 1978. ZOOINJ^Gm. ZORNIA. (J. Zorn, apoth., at Kempten, Bavaria, bot.auth.) <i>Leg.Pap. Bed. Euh.</i> 7.—1L | | | | | | | | | |
| I. DIPHT'LLA.—Leaves <i>bifoliate</i> . | | | | | | | | | |
| 17670 | glochidiata | Rchb. | spear-pointed | O « u | J jt.au | Y | Guiana | 182a | S p.l |
| 17671 | reticulata | Sm. | netted | O c u | ^jl.au | Y | W. Indies | 1800. | S p.l |
| 17672 | diphylla | /3 ciliata Pers. | Ceylon | O c u | Jjl.au | Y | Ceylon | 1825. | S 81 |
| 17673 | zeylonensis | Pers. | narrow-leaved | O un | 1 jl.au | Pu | India | 1733. | S p.l Bur. zey. 50.1 |
| 17674 | angustifolia | Mx. | dictyocarpa Dec. | net-capsuled | O c u | A ji.au | y | N. HoU. | 1820. S 81 |
| II. TETIUPIIT'LLJ!.—Leaves <i>i-foliated</i> . | | | | | | | | | |
| 17675 | tetraphylla Mx. | Anonymos bracteata Woll. | four-leaved | Ilo JCU | * jilau | w | Carolina | • 1824. | S 81 Mic. am. 2.41 |
| 17676 | captiosa Pers. | Ledjrsarum tetraphyllum Thun. | Cape | iAl cu | J jl.au | Y | C.G. H. | 1824. | S 81 Th.up.6.3 |
| 1979. STYLOSA'NTHES Swx. STYLOSANTHES. { <i>Stylos, style, anthos, fl.</i>) <i>Leg. Pap. Heds. Euh.</i> 6.—10. | | | | | | | | | |
| 17677 | procumbens | Swx. | procumbent | 1 | Lau | Y | W. Indies | 1821. | S co SeJjam. 11002 |
| S/edysarum hamatum L. | | | | | | | | | |
| 17078 | erecta Beauv. | erect | | | ll.au | Y | Guinea | 1825. | s.p Beau. ow. 2.77 |
| 17679 | viscosa Swx. | clammy | | | ll.au | Y | W. Indies | 1818. | s.p SL. Jam. 1.119.1 |
| 17680 | mucronata W. | mucronate | | | ll.au | Y | Ceylon | 1817. | s.p Bur. zey. 106.2 |
| 17681 | elatior Swx. | hispida Mx. | taller | | jl.au | Y | America | 1816. | f.l.p Pluk. at 447.7 |
| 17682 | guianensis Swx. | Guiana | | EQor | i jl.au | Y | Guiana | 1820. | S s.p Aub. guia. 2.309 |
| Trifolium guianense Aub. | | | | | | | | | |
| • 198a | ADE'SMI A Dee. | ADESMIA.- | {A, without, <i>desnōs</i> * a bond; stain, free.) <i>Leg. Pap. Heds. Euh.</i> 3.—9. | | | | | | |
| 17683 | muriata Dec. | muriatated | Jt iOllcu | 11 jjj.jll | | Y | Patagonia | 1793. | S 811 Jdcis. 668 |
| /fedysarum muricatum Jac. | | | | | | | | | |
| 17684 | pendula Dec. | pendulous | Jk QJ cu | 1 jn.jl | | Y | B. Ayres | 1825. | S s.l |
| 17685 | papposa Dec. | dovny-podded | -* iAl cu | 1 jn.jl | | Y | Chile | 1823. | S B.1 |
| jg./schynomene papposa Lag. | | | | | | | | | |
| 1981. £'BENUS L. | EBENUS. | (Abnous, ebony, Arabic.) | | | | | Legwmnhs* <i>Pap. Heds. Euhedysa</i> ce. 2.-3. | | |
| 17686 | cretica L. | Cretan | n-L-Jor | ijjjnjJl | | | Pk " Candia | 1737. | p.l Alp. ex. 278 |
| jfrathylli crética Lam. | | | | | | | | | |
| 17687 | pinnata Desf. | pinnate | | OJ or | jl au | Pk | Barbary | 1786. | S p.l Bot mag. 1092 |
| W. | | | | | | | | | |
| 1982. ROTH'il Pen. | ROTHIA. | {A. W. Roth, of Bremen, a German botanist) | | | | | Leg.Pap.Phas. 1. | | |
| 17688 | trifoliata Pers. | trifoliolate | -* O cu | 2 Jn Jl Sul | | | | S s.1 | |
| Gl'cine humifasa W. en. Cleome prostrata Hort. | | | | | | | | | |
| 1983. TERAMNUS^r. | | {Teramnos, soft; delicate pod.) | | | | | Legum. Pap. Phas. 2, | | |
| 17689 | vulgaris Swx. | twining | % | or 10 | ... | H | Jamaica | 1824. | F.m |
| 17690 | uncinatus Sun. | hooked | | 10 | ... | R | Jamaica | 1822. | s.l P.ed.B.221 |
| Z)lichos rotundifolius Vahl | | | | | | | | | |
| • 1984. CANAVALIA Dec. | CANAVALIA. | { <i>Canavali</i> , its Malabar name.) | | | | | Legum.. Hap. Phas. 4 ^.*. | | |
| 17691 | obtusifolia Dec. | obtuse-leaved | J_ • or | 6 Jl.au | | | E. Indies laSO. C s.l | R. maL 8.43 | |
| Z)lichos rotundifolius Vahl | | | | | | | | | |
| 17692 | 2 emarginata Dec. | emarginate | P or | 6 jl.au | | P | E. Indies 1800. | ■ Jac.'sc. 2.221 | |
| 17693 | gladifolia Dec. | *word-podded | P or | 6 Jn.jl | | W.B | E. Indies 1790. | ■ Jac. ic. 560 | |
| />ensiformis W. | | | | | | | | | |
| 17694 | ensiformis L. | sword-formed | Wjun | 3 Jl.au | | P | E. Indies 1778. | ■ Jac. ic 3.559 | |
| />/Mlichos roscus Swx. | | | | | | | | | |
| * 1985. ZUPPNUS Tou. | LUPINE. | { <i>Lupus</i> , a wolf; supposed to | | | | | destroy fertility of soil.) <i>Leg. Pap. Phas.</i> 23.—40. | | |
| 17695 | albus L. | white | | 3 | jl.au | W | Levant | 1596. | S co |
| 17696 | Thermis L. | Thermis | | 3 | jn.jl | W | Egypt | 1802. | S co |
| 17697 | varius L. | variable | | 3 | Jau | B.w | S. Europe | 1596. | S co |
| 17698 | hirsutus L. | hairy | | 2 | jl.au | B | S. Europe | 1629. | S co |
| 17699 | bracteolaris Dcsv. | bracted | | 1) | Jl.au | B | Monte Vi. | 1820. | s.co |
| 17700 | pilosus L. | pilose | | 3 | jl.au | F | S. Europe | 1710. | s.co |
| 17701 | angustifolius L. | narrow-leaved | | 2 | jl.au | B | Spain | 1686. | s.co |
| 17702 | /infolius Roth | Flax-leaved | | 2 | jl.au | B | . | 1799. | co Roth abh. 14.5 |
| 17703 | luteus L. | yellow | | 2 | jl.au | Y | Sicily | 1596. | co Bot mag. 140 |
| 17704 | microcarpus B. M. | small-podded | H | 2 | jl.au | B | Chile | 1821. | co Bot. mag. 2413 |
| 17705 | pusillus Ph. | little | Hap | 1 | jl.au | Pa.B | N. Amer? | 1817. | s.p |
| 17706 | bicolor B. R. | two-colored | H | 1 | jl.au | Pa.B | N. Amer. | 1826. | co Bot. rog. 457 |
| 17707 | mexicanus Cav. | Mexican | H | 2 | jl.au | B | Mexico | 1819. | co Bot. ref. 1124 |
| 17708 | europaeus llug Don. | white-leaved | S.o | 2 | jl.au | Pk | Colombia | 1826. | D.p |
| 17709 | perennans L. | smooth perenn. | | 2 | my.Jl | B | N. Amer. | 165a. | D.p. Bot. mag. 202 |
| 17710 | nootkatensis B. M. | Nootka Sound | | 2 | jnau | P | Nootka So. | 1794. | D.p. Bot. mag. 1311 |
| 17711 | polyphyllus B. R. | many-leaved. | | 2 | jn.jl | B | Colombia | 1826. | Lp Bot rog. 1096 |
| 17712 | villous W. | villous | | 1 | Jl.au | Pk | Carolina | 1787. | D.s.l |
| 17713 | sericeus Ph. | silky | | 1 | my.jn | P | N. Amer. | 1828. | D.s.p |
| 17714 | argenteus Ph. | silvery | | 1 | my.ju | W | N. Amer. | 1&6. | S 81 |
| 17715 | arboreus B. M. | tree | | 6 | jl.au | Y | . | 1793. | Bot mag. 682 |
| 17716 | mutabilis Swt. | changeable | | 5 | Jl.s | B.v | Bogota | 1819. | C s.l Sw. fl. gar. 130 |
| 17717 | multiflorus Desv. | many-flowered | | 4 | Jl.8 | B | Monte VL | 1810. | C s.l |
| 1986. PTEROCARPUS L. | PTEROCA'PUS | { <i>Pteron</i> , wing, <i>karpos</i> , fruit.) <i>Legum. Pap. Dalberg.</i> 8.—22. | | | | | | | |
| T. MOUTOU'CHIA.— <i>Stamens tnodadcl.; legmnen roundish, subgrous, l-seeded, upper suture straight, not winged.</i> | | | | | | | | | |
| 17718 | draco L. | dragon | f • m | 40 | ... | W | W. Indies | 1820. | C s.l Jac. am. 183.92 |
| II. AMPHYMENIUM.— <i>Stam. monadel.; Jr. suborbicular, 1 or 2 seeded, girdled spund with a incmbraneous wine.</i> | | | | | | | | | |
| 17719 | Rbhrii Vahl | Rohr's | f C Jor | 20 | ... | G u S | 1816. | C s.l Aub. guia. 147.56 | |
| 17720 | marsupium Rox. | pouch | \$ • or | 40 | ... | W | E. Indies | 1811. | C s.l Box. cor. 2.116. |
| 17721 | dalbergioides Roth | Dalbergia-like | ± CU or | 10 | ... | Y | E. Indies | 1817. | C s.l |
| +im-→ 111- SANTALA^IA.— <i>Stamens unequally diadelphus; fruit suborbiculate, 2 or 3 seeded.</i> | | | | | | | | | |
| 17722 | Indicus W: | Indian | t • or | 30 | ... | W | E. Indies | 1818. | C s.l Ru.am.2.70 |
| 7723 | santalinus L. | RedSaunders Wood | ec | 60 | ... | Y | E. Indies | 1800. | C 81 |
| 17724 | santalifides L. | Sandal Wood-lkf | O or | 50 | ... | Y | S. Leone | 1837. | C tJ |
| 17725 | scandens Poir. | climbing | f • or | J5 | ... | Y | Caracass | | |

- | | | | | | |
|--------------------------------|---|--|---|------------------------|--------------|
| 1987. <i>iiMERI'MNUM</i> | <i>Br.</i> | <i>AMERIMNUM.</i> | <i>(A. priv., menmna, care; supLined, qua</i> U | <i>Leg.Pap.Dalb.</i> | <i>2.</i> |
| 17726 Browne* Swt. | Browne's | • Q j o r l (Q) ... | W W. Indies | C r.m Jac. am. | 180.58 |
| 17727 <i>Btrigulsum H\$B.</i> | strigulose | Id or 2020 ... | W fruudad | 1817. C r.m | |
| 1988. <i>BRY^A Br.</i> | BRTA. { <i>Bryo</i> , to sprout} seeds germinate before falling.) | | | <i>Leg. Pap. Dalb.</i> | <i>2.—3.</i> |
| 17728 <i>Ebenus Dec.</i> | Jamaica Ebony* | a or 12jl.au | Y.G Jamaica | 1713. C r.m Br. jam. | 31.2 |
| ilmerimum E'benus Sun. | | | | | |
| 17729 <i>leoniensis Lo. C.</i> | Sierra Leone | *CDor 12 .. | Y.G S.Leone | 1821. C r.m | |
| 1989. <i>TAMARI'NDUS L.</i> | TAMARIND TREE. | { <i>Tamarhmdy</i> , Indian date, Arab.) | <i>Legum. Pap. Dalb.</i> | <i>2.</i> | |
| 17730 <i>indica L.</i> | Indian | ± *m GO jnjl | V India | 163a S r.m R. mal. | 1.23 |
| 17731 <i>occWentalisDw.</i> | western | S O m 4 0 I | W W. Indies | ... S r.m Jac. am. | 10 |
| 1990. <i>PHYLLO'LOBUM Fit.</i> | PIVVLLOLOBUM. | { <i>Phytom</i> % a leaf, <i>lobos</i> > a pod.) | <i>Legum. PhyBol.</i> | <i>1..—2</i> | |
| 17732 <i>sanzibarlnse Boj.</i> | Zanzibar | £ ES cu. | Zanzibar | 1826. C i.i.p | |

CLASS XVI. ORDER 7.

MONADE'LPHIA DODECA'NDRIA.

- | | | | | | | | |
|---|--|--|--|---|---------------|---------|----------------------------------|
| 1991. MONSO*N/if I* | MONSONIA. {Lady Anne Monson, eminent for bot. acquirements.) | <i>Geranticea?</i> 6.-9. | | | | | |
| 17730 speciosa L.
fipalidaSiv. | speciosus pale pilose | aparit R C. G. H. 1774. R rm Bot. mag. 73
Kauy Pa.R C. G. H. 1774. R rm Sw. ger. 177
Kau W C.G.H. 1774 It.r.m Sw. ger. 199 | | | | | |
| 17734 pilosa W.
speciosa Sw. ger. 77
2C61U* | Colla's Vctoed-leaved O)or | jl.au PR.W C. G. H. 1820. R r.m Col. h. rip. 2
ap.my P C. G. H. 1774. R rm Sw. ger. 273
au W C. G. H. 1774. R r.m Her. ger. 40 | | | | | |
| 1992. SARCOCAU'LON Dec. | SARCOCAULON. | {Sax, flesh, kaulos, stem.) | | | | | |
| 17737 L'Heritteri Dec. | L'Heritteri's | <i>Geranideaf.</i> 2.-3. | | | | | |
| Monson's spinosa Herit. | Z3 or | I my.jn P C. G. H. 1790. R r.m Her. ger. 42 | | | | | |
| 17738 Patersoni Dec. | Paterson'a | n. LJ or | | | | | |
| 17739 Burmanni Dec. | Burmann's | tL LJ or | | | | | |
| Geranium spindsum Brm. | | 2J my.jn ... C. G. H. 1827. R s.l Patters. 14
1 my.jn P C. G. H. 1800. S r.m Bur. af. 31 | | | | | |
| * 1993. HELICTERES L. SCREW TREE. | | {Helix, a screw; carpels twisted.) | | | | | |
| 17740 Isbra L. | Isore | i C 3 or 12 Jnjl Y E. Indies 1733. Bombhceee. 8.—18.
17741 baruensis L. | Baru | * Q o r JJ 80° W W. Indies 1739. C p.l Bot mag. 2061
17742 jwajBi>meblafl:4r-a | Guazuma-lvdif | O or 10 | Orinoco 1820. C p.l Jac. am. 149 |
| 17743 jamaicensis Jac. | Jamaica | * CZ or 8 in.jl p Jamaica 1739. C Lo Jac. vin. 2.143 | | | | | |
| 17744 verbascifolia Herit. | Mullein-leaved | • • or 8 Br Brazil 1820. C S Bot. reg. 903 | | | | | |
| 17745 ferruginata Lk. | rusty | • o r 8 Br Brazil 1823. C H | | | | | |
| 17746 lancolata Dec. | hmce-leaved | * : or 6 Y- E. Indies 1818. | | | | | |
| 17747 spicata Colb. | spicate | * : or 6 E. Indies 1823. C p.l | | | | | |
| 1994. DOMBEPYI* Cav. DOMBEVA. | {Joseph Dombey, a celebrated French bot.) | <i>Byttneridceee.</i> 5.—10. | | | | | |
| 17748 <linf6lia Cav. | Tilia-leaved | t Clor 15 ... W Bourbon 1820. "C 81 . ■ ■ ■ 3.39. 2 | | | | | |
| 17749 ferruRinea Cav. | ferruginous | 5 CD or 15 ... W Mauritius 1815. C 61 Cav. dis. 3.42.2 | | | | | |
| 17750 cordiflora Dec. | cordate-fcaitfa' | T : or 16 ... R E. Indies 1820. C 61 | | | | | |
| 17751 punctata Cav. | dotted | j D or 16 ... W Bourbon 1820. C 81 Cav. dis. 3.40.1 | | | | | |
| 17752 ovata Cav. | ovate | 9 : or 16 ... W Bourbon 1822. C 81 Cav. dis. 3.412 | | | | | |
| 1995. MONTEZITMit M. & S. MONTEZUMA. | {Monexuma, a Mexican sovereign.) | <i>BombUCete.</i> 1. | | | | | |
| 17753 spcciosissnhaAf.%S.showiest | I Q o r 30 | ... R Mexico 1827. C | | | | | |
| 1996. ASSCTNIA Cav. ASSOMA. | {Ignatius de Asso, a Spanish botanist.) | <i>Byttncrfoceee.</i> S. | | | | | |
| 17754 x>pCilnca Cav. | Poplar-feamf | f CD or 10 ... W Bourbon 1820. C s.l Cav. dis. 3.42.3 | | | | | |
| 17755 viburno'de8 Dec. | Viburnum-like | \$ CD or 11 ... W Bourbon 1822. C 81 | | | | | |
| 1997. PENTA'PETES L. PENTAPETES. | {Panic, five, pelalon, petal; flower.) | <i>Byttnerihcece.</i> 2. | | | | | |
| 17756 phoencea L. | scarlet-flowered | C3 or 2 jlau S India 1690. ■ s.p Mil. ic. 201 | | | | | |
| 17757 ovata Dec. | ovate-leaved | <. J or 2 jns. S N. Spain 1805. ■ s.p Cav. ic. 433
Brotero ovata Cav. | | | | | |
| 1998. A STRAP^A Lindl. ASTRAP&A. | {Astrape, lightning; splendid color of flower.) | <i>Byttnerihcece.</i> 2.-3. | | | | | |
| 17758 Wallachii LindL | Wallich's | • spl 20 jlau rf Madagasc. 1820. C sfp Bot.1mag. 2503
?SS'. i. asasw - clammy J or 30 ... Pk? Madagasc. 1823. C Bp
17760 tlbefBua Swt. | Lime-tree-lvd J : or 20. . . Bourbon 1824, C s.p | | | | |
| 1999. KY'DIA Rox. | KYDIA. | {Col. Robert Kyd, first director of Calcutta garden.) | <i>ByttneHheete.</i> 2. | | | | |
| 17761 calycina Box. | large-■■■■■ed | J D or SO ... W E. Indies 1818. C s.p Rox. cor. 3.215 | | | | | |
| 17762 fraterma ifox. | brotherly | J D or 40 ... W E. Indies 1823. C Bp Rox. cor. 5.216 | | | | | |
| 17763 arcfolium W. | Maple-leaved | or 10 jlS E. Indies 1790. C p.l Bot. mag. 620 | | | | | |
| 17764 /Jlatanifolium Lod. | Plane-leaved | or 15 ... W E. Indies 1820. C p.l | | | | | |
| 17765 canescens Box. | canescent | or 1b W E. Indies 1823. C P4 | | | | | |
| 17766 fuberifolium W. | Cork-tree-lvd | or 10 s.o W E. Indies 1783. C p.l | | | | | |
| 17767 lanceaefolium Box. | lance-leaved | or 13 ... W E. Indies 1820. C p.l Bot. mag. 1526 | | | | | |
| 17768 gemisagittatum Box. | balf-sagittate | J : or 10 ... 'W E. Indies 1820. C p.l | | | | | |
| 9001. KLEINHOTIo* L KLEINQOFIA. | (M. Kleinhoff, formerly director of bot. gard., Java.) | | | | | | |
| 17769 h6spita L. | stranger | f a or 20 ... P E. Indies 1800. C p.l | | | | | |
| «005. PLAGIA'NTHUS Forst. PLACIANTHUS. | | oblique, anthos, flower.) | | | | | |
| 17770 divaricatus Forst. | divaricated | * 8 Jnj. N. Zeal. 1822. t s.l | | | | | |

CLASS XVI. ORDER 8

MONADE'LPHIA POLYA'NDRIA.

200a <i>MA'LOPEL</i>	MALQPE.	(<i>Malos</i> , tender j soft leaves.)	<i>Malvdcea?</i> 2.-4.
17771 <i>malacoides L.</i>	Mallow-like	jg OJ or 1 jn.jl P	Barbary 1710. S s.l Cav. dis. 2.27.1
17775 <i>trifida Cav.</i>	trifld	O or 2 jl P	Barbary ■ S B.I Cav. dis. 2.27.2
*2004. <i>Af'A'LVAZ.</i>	MALLOW.	(<i>Malasso</i> , to soften; emollient qualities.)	<i>Malvaceae</i> . 70.—86.
		I. MALVA'STRUM.—Carpels 1-loclar, 1-seeded.	
i. CUUSA'NTER.	—Flowers yellow, in the axes of the upper leaves, sometimes in spikes from the uppermost leaves being wanting.		
17773 <i>tricuspidata H. K.</i>	tricuspidate	jg Ol pr 1 jl.au Y	W. Indies 172a S co Cav. dis. 2.22.2
17774 <i>cororaandelliana W.</i>	American	O P' 1 jn.jl Y	W. Indies 1756. ■ go
17775 <i>scabra Cav.</i>	rough-stemmedn.	• pr 4 jn.jl Y	Peru 1798. co Cav. dis. 5.138.1
17776 <i>jeoparia Jac.</i>	Broom	■ Dp' 6 a.u.s Y	Peru 1782. C co Jac. ic. 1.139
17777 <i>borbónica W. en.</i>	Bourbon	■ 4 jlau Y	Mauritius 1816. C co
17778 <i>polystachya Cav.</i>	many-spiked	<-i Jpr 6 jl.au Y	Peru 1798. C co Cav. dis. 5.1S8.3
17779 <i>spicata Cav.</i>	simple-spiked	n. Lpr 2 s.o O	Jamaica 1726. C co Cav. dis. 2.20. 4
17780 <i>ovata Cav.</i>	ovate	it-dpr 2 jl.au Y	Brazil 1820. C s.p Cav. dis. 20.2
	striata L.		
17781 <i>tomentosa L.</i>	woolly	■ Dp' 3 s.o Y	£. Indies 1820. C co Pluk. am. 356.1
17782 <i>waltheriegfd/k Lk.</i>	Waltherie-lvd	jg EJ pr ■ Y	Java 1824. D co
17783 <i>trachelifolia Lk.</i>	Trachelium-lvd	O un 1J jl.au Y	■ 1821. S co
17784 <i>gangética L.</i>	Ganges	O un 1 jl.au Y	E. Indies 1823. S co Pluk.al.74.6
17785 <i>domingensis Spr.</i>	Domingo	£ E2 pr 2 jl.au Y	St.Domin. 1824. S co
ii. CYMBALVIE.—Flowers purplish or white; pedicels axillary, l-flowered, solitary; outer calyx 3-leaved; leaves roundish s stems herbaceous.			
17786 <i>leprosa Or.</i>	leprosus	jC E2 un 2 my.jl P	Cuba 1815. C co
17787 <i>Sherardifow L.</i>	Sherard's	■ A un 2 myjl P	Bithynia 1818. D s.p Cav. dis. 2.2T>.4
17788 <i>cretica Cav.</i>	Cretan	O un 1 myjl P	Candia 1825. S co Cav. dis. 5.138.2
iii. BIBRACTBOLOTJE.—Flowers purplish or white; outer calyx 2-lvd; l-flowered; stems herbaceous.			
17789 <i>hispanica L.</i>	Spanish	O un 1 jl F	Spain 1710. S co Desf. at. 2.170
17790 <i>stipulacea Cav.</i>	/Hf-stipulcd	O un 1 jn.au P	Spain 1815. S go Cav. dis. 2:15.2
17791 <i>sgyptia L.</i>	Egyptian	O un 1 jn.jl ■	Egypt 1815. ■ co Cav. dis. 5.137.2
17792 <i>trinida Cav.</i>	trinid	O un 1 jn.jl ■	Spain 1815. ■ co
iv. BISMA'LYE.—Fls. purplish or white; pedicels axUL, solit.; invol. 3-lvd s Ivs multipartet; stems herbaceous.			
17793 <i>Tournefortina L.</i>	Tournefort's	O un 1 jlau B	Spain 1759. S co Cav. dis. 2.17.3
17794 <i>ilicæa L.</i>	Alcea	£ A or ■ jl.o P	Germany 1597. Deo Bot. mag. 2297
17795 <i>alccoldes Ten.</i>	Alcea-like	5 A or + jl.o K	Italy 1820. D co Bot. mag. 2793
Moren Pall. ^lceæ/S Moren Dec.			
17796 <i>moschata L.</i>	musk	It A P' ■ jl.au F	Britain bor.fi. D co Eng. hot.7M
2 undulata SOTJ	waved	5 A pr ■ jl.au W	— D co Bot. mag. 2298
17797 <i>althaeoides Cav.</i>	Althaea-like	O un ■ jl.au P	Spain 1822. S co Cav. dis. 2.15.3
17798 <i>fastigiata Cav.</i>	fastigiate	O un ■ jl.nu R	S. Europe 1818. S co Cav. dis. 2.23.2
v. FASCICULA^TJE.—Flowers purple or whites leaves angular cordate, 5-nerved pedicels axillary, many-flowered; outer calyx S-leaved; stems herbaceous.			
17799 <i>mauritiana L.</i>	Mauritanian	6 jn.jl Pk	S. Europe 1768. S co Sw. fl. gar. 81
17800 <i>glabra Desv.</i>	smooth China	Vk	China 1818. S co Cav. dis. 2.25.4
17801 <i>glomerata W. Sf K.</i>	glomerated	■	Hungary 1817. S co
17802 <i>sylvestris L.</i>	wood	■	Britain was.pl. D co Eng.bot.671
2 albiflra Dec.	white-flowered	■	Europe ... D co
17803 <i>Henningif Goldb.</i>	Henning's	■	Russia 1820. S co
17804 <i>rotundifoliaZ.</i>	round-leaved	■	Britain ro.sid. S co Eng. bot. 1092
2 pusilla Sm.	small	■	Britain ro.sid. S co Eng.bot.L'41
17805 <i>brasiliensis Desr.</i>	Brazilian	■	R.Janeiro 1824. S co
17806 <i>nicaeensis All.</i>	Nice	■	Nice 1810. S co Cav. dis. 2.25.1
17807 <i>microcarpa Desf.</i>	small-fruited	■	Egypt 1823. D co
17808 <i>parviflra L.</i>	small-flowered	■	Barbary 1779. C co Cav. dis. 2.26.1
17809 <i>verticillata L.</i>	vhox\flowered	■	China 1683. ■ co Cav. dis. 2.25.3
glomerata Hart.			
17810 <i>mareticæa Del.</i>	Marcotic	■	Egypt 1822. ■ co
17811 <i>crispa L.</i>	curled	■	Syria 1573. ■ co
17812 <i>bonariensis Cav.</i>	Buenos Ayres	■	B. Ay res 1820. ■ ro
17813 <i>scaberrima Lag.</i>	roughest	■	Spain 1827. ■ co
17814 <i>flexuosa Horn.</i>	flexuous	■	Europe 1820. ■ co
vi. CAPB'NSES.—Flowers purple or white; leaves angulate-lobed; stem shrubby; outer calyx 34eaved; pedicels			
solitary, l-flowered, rarely in twos or threes			
17815 <i>amœna Sims</i>	pleasing	■ Jor ap.my P	C. G. H. 179G. C co Bot. mag. 1998
17816 <i>virgata Cav.</i>	twiggy	■ Jor myjl P	C. G. H. 1727. C co Cav. dis. 2.18.2
17817 <i>canænsis Cav.</i>	Cape	■ Jor : year R	C. G. H. 1713. C p.l Bot. reg. 295
17818 <i>balsamica Jac.</i>	balsamic	■ Jor my.s P	C. G. H. 1800. C co Jac. ic. 1.140
17819 <i>tridactylites Cav.</i>	three-fingered	■ 3 jn.au Pk	C. G. H. 1791. C p.l Bot. rep. 135
reflexa And. Rep.			
17820 <i>divaricata Andr.</i>	straddling	■ I-4 3 jn.s W.vy C.G. H. 1792. C co Bot. reg. 182	
17821 <i>retbsa Cav.</i>	blunt-leaved	* I or 4 mr.myPk C.G. H. 1803. C p.l Cav. dis. 2.21.1	
17822 <i>calycina Cav.</i>	tor'i-calyxed	1 J or 4 my.au R C. G. H. 1812. S co Bot. reg. 297	
17823 <i>trigrans Jac.</i>	fragrant	A ■ or 3 myjl S C. G. H. 1759. C co Bot. reg. 396	
17824 <i>stricta Jac.</i>	strict	• J or 3 my.au W.vy C.G. H. 1805. C co Jac.sc. 3.294	
17825 <i>bryoniæa L.</i>	Bryony-leaved	11 or 4 jl.au P C. G. H. 1731. C co W.h.b.1.4	
17826 <i>grossulariæa Cav.</i>	Gooseberry-lvd	■ 3 my.s Pk C. G. H. 1732. C p.l Bot. reg. 561	
17827 <i>yanacanthodes Hor.</i>	Oxyacantha-likeft	■ 4 my.au W C. G. H. 1818. C si	
17828 <i>asplirrima Jac.</i>	roughest	si I or 3 jn.8 R C. G. H. 1796. C co Jac.sc.2.39	
vii. MULTIFLOILE.—Flowers purplish or whites peduncles axillary, many-flowered.			
K. particled milk-wMte	LJ or 4 ja.f W Mexico 1780. C co Cav. ic. 1.20		
Cav. red i j o r 4 myjl Ve S. Amer. 1798. C p.l Cav. ic.3.273			

<i>I</i> 7831 Monroænii Dou.	Monro's	& A	or	2	jLau	S	Columbia	1828.	■	⊗	Bot reg. 1306
17832 pcruviana Z,	Peruvian	O	un	2	jn.au	P	Peru	1759.	■	⊗	Jac vin. 2,156
17833 fim&isüs J,	Lima	O	or	4	J	■	Peru	1768.	■	⊗	Jac. vin. 2,141
17834 capitata Cav.	capitate	• i_J	or	2	n.d	R	Peru	1798.	■	⊗	I av. dis. 5.137.1

II. SPHALEROMA. — Carpels unilocular, 2 or many-seeded, joined into a globose fruit; peduncles axillary, often many-flowered; stems shrubby.

17835 <i>umbellata</i> Cav.	umbelled	*	or	4' ja.inr	€	S. Amer.	1814.	C co	Bot. cab.	222
17836 <i>abutiloides</i> * L.	Abutilon-like	■	or	4 ins	R	Bahama	1.1725.	C co	Bot. mag.	2541
17837 <i>elegans</i> Cav.	elegant	■	or	3 my.au		C. G. H.	1791.	C co	Jac. c. 4.6.1	
17838 <i>angustifolia</i> Cav.	narrow-leaved	■	or	3 au	Pk	Mexico	1780.	C p.l	Cav. ic.	1.68

III. MODI^{*}OLA. — Carpels bivalved, 2-seeded; valves 2-awned, inflexed inwardly, and hence the carpels are half-bilocular; pedicels axillary, flowered; stems herbaceous, prostrate, or diffuse.

17839 caroliniana L.	Carolina	«* O or 1 jn.jl	R	Carolina	1723. S co	Cav. dis. 2.	15.1
17840 prostrata Cav.	trailing £L AJor 1	jnau Pk	Brazil	1806. S co	Bot mag.	2J	15
17841 decumbens IV. en.	decumbent fc,	LAJ or 1 ins.	Pk	S Amer.	1815. S co		
17842 obtusiloba Hook,	obtuse-lobed	£ ^ J or 4 s	R	Chile	1827. C p.l	Bot. mag.	2787

2005. NUTTA'LLW Dick. NUTTALLIA. (*Thomas Nuttall, prof. min., Cambridge, ttew England.*) *Malv.* 2.
17813 *digitata* Dick. finger-leaved t A or 3 au P. N. Amer. 1824. R p 1 Hook. ex fl. 171
17844 ~~pedata~~ Dick. pedata ^{sp} A or 3 au P. N. Amer. 1824. R p 1 Hook. ex fl. 171

2006. KITAIBETLI W. KITAIBELIA. (*Paul Kitaibel*, M.D., prof. bot. at Pest, Hungary.) *Malvace**. 1.
17845 vitifolia W. Vine-leaved. ^ A or 5 ilis. W. Hungary. 1801. D. pl. Bot. mag. 821.

*2007 ifL THIEA, L. MARSH MALLOW (*Althea*, to cure i. medical qualities). *Malydcete*, 16—22.

I.	ALTHYSTRUM	— Carpels emarginate, destitute of membranaceous margins; involucres often 8 or 9 cleft.					
17846 officinMis L.	officinal	i A ™	4	jls	F	Britain	saltm.
17847 taurinensis Dec.	Turin				R	Turin	1817.
17848 narbonensis Pou.	Narbonne				Pk	S. Europe	1780.
17849 cannibina L.	Hemp-leaved				P	S Europe	1597.
17850 hirsuta L.	hairy			jn.jl	W	Britain	chl. pi.
17851 L. 1. f. 1. I.	Ludwig's				Pk	Sicily	1791.

17851 Ludwig L. Ludwig's 81-342 jnjl 1K Stein 1951 S co. Cav. d.

II. ALCR ^a . — Carpels surrounded by sulcate tнемbranaceous margins; involucels 6 or 7 cleft.							
	stemless	leafy	jl.	P.	Aleppo	...	Cav. dis. 2. 27.3
17852 acanthifl. Cav.							
17853 carib. ^a B.M.	Caribbean	if fl. or	mr.ap	Pk	W. Indies	1816.	Bot. mag. 1916
17854 striata Dec.	striated	%	jl.au	V			
17855 pillida IV.8f.K.	pa-c-flowered	?	jl.au	P. a	Hungary	1805.	S
17856 rbsae Cav.	rosy Hollyhock		jl.s	R	China	1573.	3
2 biloba Dec.	two-lobed		jl.au	R	S. Europe	...	Cav. dis. 2. 28.1
grandiflora Hori,							
17857 sinensis Cav.	Chlna		jl	R	China	1818.	S co
17858 flexubus A.M.	flexuous		jn.au	Pk	E. Indies	1803.	D co
17859/ificiblia Cav.	Fig-lyd Antwerp Holl.		jn.a	O	levant	1597.	S co
17860 Froloviana Fis.	Frolov's		jl.au	W	Siberia	IA27.	S co
17861 leucontha EIU	white-flowered		jl.au	W	Siberia	1827.	co

² 2008 LAVA'TERI L. LAVATERA (The two *Lavaters*, friends of Tournefort's). *Malváceae*. 24-27.

I STKIIA — *Receptacle or axis of fruit at top expanded into a disk concealing the ovaries.*

L. STKJIA. — Receptacle, or axis of fruit, at top expanded into a disk concealing the ovaries.
 17862 *tristriatum* L. three-monthly O or 2 jls F S. Europe 1633. S co Bot. mag. 109
 17863 *Pseudo-Olbia Pair.* Bastard Olbia H (J or 5 jn.jl R S. Europe 1817. C 81

17864 phœnicioa Ven. II. O'LRIA. — *Receptacle of fruit central, conical, exserted.*
fine-red ■ i. 1 or 5 in il Pk Cameris 1816 C 61 Ven mal 120

17804 pincneea <i>Ven.</i>	<i>Junc-reu</i>	[?]	of	3 jnjl	Pk	Canares	1810.	C s.l	Ven. mal. 120
17855 hispidin <i>Desf.</i>	<i>hispid</i>	J or		6 jnjl	Pk	Algiers	1804.	C co	Bot. mag. 2541
17866 fccrifolia <i>Cav.</i>	Maple-leaved	[?]		5 jnjl	Pk	Tenerifie	1820.	C s.l	Her. am. 322
17867 africtaria <i>Cav.</i>	African	[?]		[?]	Pk	Spain	1820.	C s.l	Cav. dis. 5.139.1
17868 Olibia <i>L.</i>	Ollria	[?]	lor	3 in.o	R.p	France	1570.	C s.l	Cav. dis. 2.32.2
17869 unguiculifera <i>Desf.</i>	clawed			6 jls	Li		1807.	C co	
17870 micans <i>L.</i>	glittering	H L J or		3 jnjl	P	Spain	1796.		M.h. 1.17.9
17871 lusit.&nlca / ,	Portuguese	* i or		3 a.us	P	Portugal	1731.	C s.l	
17872 flava <i>Desf.</i>	yellow	<i>Si</i> A or		4 jnjl	Y	Sicily	1818.	D s.l	Desf. at. 2.172
Emptocoris Raue.									
17873 plebeia <i>Sims</i>	vulgar	[?]	[?]	[?]	[?]	N. Holl.	1820.	D co	Bot. mag. 2269
17874 thuringiaca <i>L.</i>	Thuringian	[?]	[?]	[?]	[?]	Germany	1731.	D co	Bot. mag. 517
17875 biennius <i>Rieb.</i>	biennial	[?]	[?]	[?]	[?]	Caucasus	1819.	S s.l	
17876 punctata <i>AU.</i>		[?]	[?]	[?]	[?]	Italy	1800.	S co	

III. AX6LOPIA. — Receptacle at top truncate, expanding in a crest.
 17877 *maritima Gou.* seaside m—or 2 ap.jn W S. Europe 1597. S. sl Cav. dis. 2.32.3

17878 *triloba* L. thre-lobed a____or 3 jn.jl L.P Spain 1759. C s.l Bo

17879	<i>arborea</i>	<i>L.</i>	Tree Mallow	CD or	6	jl.o	P	Britain	sea cl.	S	CO	Eng. bot.	1841
17880	<i>neapolitana</i>	<i>Ten.</i>	Neapolitan	■	OR	6	jl.s	P	Italy	1818.	S	CO	
17881	<i>cretica</i>	<i>Z.</i>	Cretan	A	OR	4	jl.s	L.B	Candia	1723.	S	CO	Jac. vin. 1.41
17882	<i>sylvestris</i>	<i>Brot.</i>	wood	O	OR	2	jl.s	P	Spain	1817.	S	CO	
17882	<i>ambigua</i>	<i>Dav.</i>	ambiguous	O	OR	2	jl.s	P	Italy	1824	S	CO	

17883 <i>ambigua</i> Dec.	<i>ambiguoas</i>	O or 2	jls	P	Italy	1824	S co
17884 <i>lanceolate</i> TV. en.	V. DUVBLE.—Doubtful to which of the sections they belong. <i>lance-leaved</i>	O or 2	jls	P	Europe	1817.	S co

2009 IVAN LACHRA *I.* MALACHRA (*Malachra mollis*, *mollis*) Malachra 8-14

2009. J. VIALACHRA L.
17889. *rotundifolia* Schrank
MALACIKA. round-leaved
(Malache, mallows; resemblance.)
O un 1 il.au Y Brazil 1821. S s 1 Sch me 56
Malgache. 8.—14

• 2010. <i>UBJTNA</i> L.	URENA.	(<i>L^{Tren}</i> , its name in Milabar.)	<i>Malvhcea?</i> . 12.—22.
I. INDIVINJS. — <i>Z<10</i> «undivided or hardly 3-lobed.			
17894 <i>lobata</i> L.	<i>lobed-leaved</i>	F China 1731.	■ p.l Di. el. 319.412
17895 <i>scabriuscula</i> Dec.	<i>roughish</i>	2 jl.au R E. Indies 1817.	■ p.l Cav. dis. 185.1
<i>lobata</i> Cav.		O un	
17896 <i>repanda</i> Sm.	<i>repand</i>	Q un 2 jl.au R E. Indies 18C0.	■ p.l
17897 <i>multifida</i> Cav.	<i>many-cleft</i>	tLOun 2 ja.o P E. Indies 1817.	■ p. Cav. dis. 6.184.2
17K98 <i>trie uspis</i> Cav.	<i>three-pointed</i>	• Oun 2 jl.au K I. France 1820.	■ p. Cav. dis. d 183.1
17899 <i>reticulata</i> Cav.	<i>netted</i>	tt. Qun 2 jl.au R Guyana 1819.	■ p. Cav. dis. 6.183.2
17900 <i>subtriloba</i> Schrank	<i>Bub-trilobed</i>	O un 2 jl.au R Brazil 1823.	■ p. Sch. ma 79
IL LODA^TJS. — <i>Leaves deeply 3 or 5 lobed s lobes stnately lobed.</i>			
17901 <i>Swartn</i> * Dec.	SwarU's tt.	O un 2 ... W Surinam 1816.	C P.I SI. jam. 1.11.2
<i>americana</i> Sm.			
17902 <i>Sieberi</i> CoU.	Siebert	O un 2 jl.au R I. France 1824.	S p.l Col. h. rep. 39
17903 <i>sinuata</i> L.	<i>sinuated</i>	tt. O un 3 jl.au F E. Indies 1759.	C p. Cav. dis. & 185.2
17904 <i>muricata</i> Dec.	<i>mupicated</i>	O un 2 jl.au R Bengal 1816.	S ■
17905 <i>heterophyrla</i> Sm	<i>various-leaved</i>	■ 2 Jn.jl R Ceylon 1818.	S p.l Bur. zey. 69.2
SOU. PAVOMA Cav. PAVOMA. (<i>Don Josef Pavon</i> , of Madrid, a bot author.) <i>Malvdcea?</i> . 16.—24.			
I. Tvpia*LEA. — <i>Carpels echinate from rigid spines.</i>			
17906 <i>spinifex</i> W.	<i>prickly-seeded</i>	x g i u y ^r. indies 1778.	C s.1 Jac. vin. 2.103
JETBiscus <i>spinifex</i> L.			
17907 <i>Typhalea</i> Cav.	Typhalea	■ 1jj1au W.B Trinidad 1824.	C 8.1 Cav. dis. 6.187
<i>Urina Typhalea</i> L.			
17908 <i>typhahrtia</i> JLS/B.	Typhalca-like	tt. O or 1 ■ Pa.R Trinidad 1824.	C p.l
17909 <i>tirens</i> Cav.	<i>stinging</i>	* O un 2 - R Mauritius 1801.	C s.1 Jac. ic. 3.522
17910 <i>leptocirpa</i> Cav.	<i>slender-capsuled*</i>	O un 2 Jn.au W Surinam 1818.	C s.1
II. MA'LACITE. — <i>Carpels unarmed s involucels shorter than calyx, 5 or 1S leaved.</i>			
17911 <i>cocclnea</i> Cav.	<i>scarlet</i>	HI lor 2 S StDom. 1816.	C s.1 Cav. dis. 3.47.1
Columell* Cav.	<i>Columella's</i>	it O un 2 jl. W.P Bourbon 1807.	C s.1 Cav. dis. 3.48.3
17913 <i>parviflora</i> Desf.	<i>small-flowcrcd</i>	tL O un 2 jlau Pa.Y S. Amer. 1818.	C s.1
17914 <i>racemosa</i> Sum.	<i>racemose</i>	tt. O un If Jn.au Y Jamaica 1817.	C 8.1 Cav. dis. 3.46.1
<i>spicata</i> Cav.			
17915 <i>pramdrsa</i> W.	<i>bitten-leaved</i>	A i J un 10 Jn.au W C. 6. H. 1774.	C s.1 Bot mag. 436
III. CANCELЛАVUA. — <i>Carpels unarmed ; involucels 8 or 15 leaved, longer than calyx.</i>			
17916 <i>paniculata</i> Cav.	<i>paniced</i>	■ Oun 2 jlau Y Caracas 1820.	S s.1 Cav. dis. 3.46.2
17917 <i>odortta</i> W.	<i>fragrant</i>	i Q un 2 - R E. Indies 1807.	C s.1
17918 <i>corymbosa</i> W.	<i>corymbose</i>	tt. J un 2 jlau Y Jamaica 1818.	S a.1
17919 <i>ridoldes</i> Horn.	<i>Sida-like</i>	tt. TBlun 2 Jn.au F Bengal 1819.	S s.1
17920 <i>zeylanica</i> W.	<i>Ceylon</i>	■ 3 Jls. W E. Indies 1790.	S s.1 Cav. dis. 3.48.2
17921 <i>cancellata</i> Cav.	<i>latticed</i>	■ 1 Jl.au Y Trinidad 1820.	S s.1
<i>hibiscus cancellatus</i> L.			
2012. MALVAVI'SCUS D'd.	MALYAVJSCUS.	(<i>Malva</i> , mallow, <i>viscus</i> , glue.)	<i>Malvaceae</i> . 3.—15.
17922 <i>arboreus</i> Cav.	<i>tree</i>	H Q or 12 year S Jamaica 1714.	C p.l Bot mag. 2305
Achânia <i>Malvavscus</i> L.			
17923 <i>mtillis</i> Dec.	<i>soft</i>	H Q or 12 au.s S America 1780.	C p.l Bot. reg. II
Achânia <i>molus</i> H. K.			
17924 <i>pil&pus</i> W.	<i>pilose</i>	• Q or 12 o.n R Jamaica 1780.	C p.l Bot cab. 829
Achânia <i>pilosa</i> Stuz.			
2013. LEBRETO*N/i4 Schrank LEDRETONIA. (<i>Manuel le Breton</i> , a French botanist.) <i>Malvhce?</i> . 1. — 3.			
17925 <i>cocclnea</i> Schrank	<i>scarlet</i>	» D o r 10 Jn.jl S Brazil 1823.	C s.1 Sch.mo.90
* 2014. J/IBI'SCUS L. HIBISCUS. (<i>Ibis</i> , a stork; said to chew and inject as a clyster.) <i>Malvace?</i> . 79.—125.			
I. CREM6NTIA. — <i>Corollas convoluto-cylindrical; stigma* 5 ; cells of fruH many-seeded.</i>			
17926 /liifl&rus Cav.	<i>Lily-flowered</i>	■ i lor 6 Jn.jl S Bourbon 1822.	C s.1 Cav. dis. 3.57.1
17927 pdcumulatas Cav.	<i>/ofif-peduncled</i>	■ J or 2 my.d R C. G. H. 1812.	C s.1 Bot. reg. 231
17928 lampas Cav.	<i>lamp</i>	• O or 10 ... Pk E. Indies 1801	C p.l Cav. dis. 3.56.2
17P29 membranaceus Gzc	<i>membranaceous</i>	Dor 10 ... Pk 1816.	C s.1 Cav. dis. 3.57.2
17930 spiralis Cav.	<i>spiral</i>	Dor 6 Jn.jl Y.R Mexico 1823.	C s.p. Cav. dis. 2.16.2
17931 tubulbus Cav.	<i>tubular</i>	O or 2 Jl.o Y E. Indies 1796.	S s.1 Cav. dis. 3.68.2
17932 senegalensis Cav.	<i>Senegal</i>	■ Dor 4 Jn.jl Y Guinea 1824.	C s.p. Cav. dis. 3 68.1
17933 oxahdiffl&rus Boj. t	<i>Oxahs-flwd</i>	■ l Jor 3 ... Y Mauritius 1825.	C s.p.
II. PENTASPE'EMUM. — <i>Corollas expanded ; valves of capsule in the middle septiferous ; cells I-seeded.</i>			
17934 pentacfrpos L.	<i>five-fruited</i>	£i A Jor 3 Jls L.R Venice 1152.	S s.1 Jac. ic. 1.143
17935 virginicus L.	<i>Virginian</i>	k A r 2 Jls R Virginia 1798.	D s.1 Jac. ic. 1.142
III. MA'NIHOT. — <i>Loculaments of carpels, many-seeded; seeds smooth ; involucels 4 or 6 leaved s calyx</i>			
17936 <i>Mdnuitot</i> L.	Manihotaceou	S-toothed longitudinally rhizomed 1712.	C p.l Bot mag. 1702
17937 <i>palmatus</i> Cav.	<i>palmated</i>	■ S. Amer. 1800.	C s.p. Cav. dis. 3. fi.3.1
17938/cnlnnea L.	<i>Tg-leaved</i>	■ Ceylon 1732.	C p.l Cav. dis. 3.52.2
17939/culneoides B. R.	<i>Ficulneus-like</i>	■ 1823.	C s.p. Bot.rcg.938
17940 tetraphyllus Box.	<i>four-leaved</i>	Bengal 1818.	S sp
IV. KE'TM IA. — <i>Loculaments of carpels many-seeded; seeds smooth; corollas expanded ; involucels 5 or 7</i>			
17941 <i>quinquefolius</i> Hort.	<i>five-lobed</i>	leaved: calyx S.Jobed	
17942 <i>cerifolius</i> Dec.	<i>Maple-leaved</i>	D or 10 Jn.jl Y	C s.p.
17943 <i>Absa sinensis</i> L.	<i>Chinese Rose</i>	i or 5 mr.jn Pk China 1798.	C 8.1 Par. Ion. c. ic.
2 rtbro.pl tna	<i>double red</i>	■ dor 10 Jl.au D.R E. Indies 1731.	C p.l Bot. mag. 158
3 flavo-plena	<i>double yellow</i>	D or 10 Jl.au D.R E. Indies ...	C p.l Bot. cub. 995
4 c&neo-plena	<i>double flesh</i>	D or 10 Jl.au F E. Indies 1731.	C p.l
5 variegato-plena	<i>dble-variegated</i>	D or 10 Jl.au St E. Indies ...	C Pi Bot. cub. 963
6 l'tea	<i>double yellow</i>	I or 10 Jl.au Y E. Indies 1823.	C p.l But. cub. 932
17944 <i>sylflacus</i> L.	<i>Syrian</i>	or 8 au.s P Syria 1596.	C CO Bot mag. 83
1 purpdrcus	<i>purple-flowered</i>	A or 8 au.s P	— L CO
S riüber	<i>red-flowered</i>	tt or 8 au.s R	— L CO
Sallbus	<i>white-flowered</i>	tt or 8 au.s W	... L CO
4 variegatitius	<i>variegated-flwf</i>	St or 8 au.s St	... L CO
5 bo-plena	<i>double white</i>	tt or 8 au.s P	... L CO
8 Purp&P-lfenut	<i>double purple</i>	tt or 8 au.s W	... L CO

7 marginatus 17915 prunens Box.	margined-tal tickling	■ □ □ 8 nus 3 jl.8	P.w Y	Syria — L co R Indies 1804. S s.1	Bot rep. 498
V. JURCA^RIA. — Ch	WT^ many^eededj seeds smooth ; corollas expanded ; leaflets ofinvoluteel at ton bifure.				
179» suratt'i nsis L.	Surat O or 2 jl.s	Y	R Indies 1731. S s.1	BoL mdg	7356
17947 radiatus Cav.	rayed O or 2 jn.au	Y	R Indies 1790. S s.1	Bot mag. 1911	
17948 furcMus Box.	folked-calyxed Elfor 2 jl.s	Y	R Indies 1816. S s.1	Bot mag. 1911	
17949 scaber Ar.	scabrous jf IAI or 2 jLs	Y	Carolina 1810. ■ □		
acuteatus Wait.					
17950 bifurcatus Cav.	two-forked ■ □ □ 2 jn.jl	■	W. Indies 1825. C s.p	Cav. die. 3.51.1	

VI. ABELM6SCHUB.— Carpels many-seeded; seeds smooth or with a subvius fine on the back; corollas expanded ; involucels constantly with 6 or 1B entire leaflets.

» TUBERULATI—Stems prickly from tubercles.

17951 strigulus And. /iculneus Guv.	strigose different-leaved *i_J or 6	6 jn.jl Pk	Peru 1820. C s.1	Bot reg. 860	
17953 trilobus Cav.	thre-lobed < or 2	jl.au Y	W. Indies 1818. C s.p	Cav. dis. 3.53. 2	
17954 heterophyllus Fen. /grandiflorus Sal.	various-leaved * i_J or 6 aus	W.R N. S. W. 1803. C s.p	Bot reg. 29		
17955 cannabinus L.	Hemp-feaiwd £ O i or 5	jn.■ W.p	R Indies 1759. S 81	Rox. cor. 2.190	
17956 wtifolius Z.	Vine-leaved £ OS or 2 jl.■	Y	R Indies 1690. S p.l	B, maL 6.46	
17957 obtusifolius W.	obtuse-leaved * OJ or 2 jkau	Y	R Indies 1820. S s.p		
ii. INR'RME9. — Stems unarmed.					
17958 esculentus L.	eatable EH cu 4	jn.jl Y	W. Indies 1693. ■ r.m	Cav. dis. 3. 61. 2	
17959 longifolius W.	long-leaved O or 4	jl.o Pa.Y	R Indies 1817. ■ s.p		
17960 Bditionia Lk.	Bammia O or Ujn.jl	2 au.o ■	1818. S 81	Toz.mu.d.flr.2.7	
17961 Afoscheita L.	Mallow Rose 3 A or 3	jl.au ■	Amer. 1816. D p.l	Cav. dis. a 63.1	
17962 paliflora L.	marsh 5 A or 3	jl.s Pk	N. Amer. 1759. D bog	Cav. dis. 3.65.2	
17963 aquaticus Dec.	aquatic 31 A or 2	jn.au W	& Europe 1819. D bog		
paliflora Sav					
17964 incanus W.	hoary ■ JLor 3 s	Y	Carolina 1806. D bog	W.h.b.24	
17965 militaris Cav.	military ■ JLor 3 aus	P	Louisiana 1804. D bog	Bot. mag. 2385	
17966 specidis H. K.	superb ■ JLor 2 jn.au	S	Carolina 1778. D bog	Bot mag. 360	
17967 grandiflorus Mx.	large-flowered if JLor	jl.au F	Georgia 1816. D bog		
17968 ferrugineus Cav.	rusty T Qor 15	■ S	Madagas. 1824. C p.l	Cav. dis. 3.69.1	
17969 lunariifolius UR W.	Lunaria-leaved LZor 15	■	R Indies ... C 81		
17970 icthi^picus L.	JEthiopian » or 15	■ au P	C.G.H. 1774. C p.l	Cav. dis. 3.60 1	
17971 mutabilis L.	changeable ■ JLor 15 o.d	W	R Indies 1690. p.l	Bot. rep. 228	
2 flore pleno	double-flowered ■ JLor 15 o.d	W	R Indies		
17972 clypeMus L.	Bhieh Ucapsuled ■ or 6	jl.au Y	Jamaica 1759. p.l	Cav. dis. 3.58.1	
17973 unidens Lindl.	one-toothed ■ JLor 3	Y	Brazil 1822. co	Bot. reg. 878	
17974 Abelmidochus L.	Musk Okro ■ JLor 3	jl.s Y	India 1640. r.m	R. mal. 2.38	
17975 velutinus //, ^ A.	velvet ■ O or 6	jl.au W	Cuman. 1818. s.p		
17976 erlocarpus Dec.	woolly-fruited ■ Q or 4	jl.s Y	Bengal 1823. C s.p		

VII. BOMBICE'LLA.—Canx7s many-seeded; seeds woolly: corollas most often expanded; involucels 5 or 10 Ivd.					
17977 phoenix IV. vnl-flowered f or 8	jl.au P	R Indies 1796. C p.l	Bot reg. 230		
17978 W malabarica J&Jr	Malabar rose ft O or 2	aus. R »Indies 1818. C-p.l	Bot reg. 337		
17979 rhombifolius Cav.	rhomb-leaved ■	jl.au R Indies 1823. C ap	Cav. dis. 3.69. 3		
J7980 elatus Sun.	tall ■ or	■ R Indies 179X. C p.l			
17981 ^ossijinns Thun.	Cotton-like ft J or	jlau ■ C. G. H. 1818. C p.l			
17982 micraithus Cav.	small-flowered a-r-l or	■ R Indies 1794. C s.p	Cav. dis. 3.66.1		
17983 clandestine Cav.	clandestine <! or	jl.au Crea Guinea 1823. C s.p	Cav. ic. 1.2		

VIII. TRI6NUM—Carpels many-seeded; seeds smooth; corollas expanded; involve, many Ivd: calyx inflated.					
17984 Trioura L.	Bladder Ketmia O or 2	jns. Y.Br Italy 1596. S co	Bot mag. 209		
J7985 vesickeijs Cav.	bladdery O or 1jl.jau	Y.Br Africa 1713. S co	Cav. dis. 3.64.2		
17986 hispidus Mil.	hispida ■ II jn.s Y.Br C. G. H. .	S co	Bot.reg.80t)		

IX. SADDARIFFA.—Cells of capsule many seeded ; seeds smooth ; involucell Meaved, many-toothed.					
17987 Sabdariffa L.	Sabdariffa jgO Jor 2	jns. Y R Indies 1596. S p.l	Cav. dis. 6.198.1		
17988 digitatus Cav.	digitate O or 2	jl.s W.R Brazil 1816. S co	Cav. dis. 3.70.2		
2 Kertfn'l Dec.	Kcr's £ O Jor 2	jl.s W.R Rio Jan. 1816. S s.1	Bot reg. 608		
digitktus B. It.					

X. AZA'NZA.—Same as preceding, but with the stems shrubby or arborescent.					
17989 circinatus W.	rounded ■	■ Caracas 182a C p.1			
17990 tilaceus L.	Lime-tree./w/ ■	R Indies 1739. C p.l	Bot reg. 233		
17991 tricospis Cav.	three-pointed ■	Society Is. 1820. S s.p	Cav. dis. 3.55.2		
17992 guinlnsis 2)ec. Guinea Umbrella tr.	■	Guinea 1822. C p.1			
17993 abutiloides W. en.	Abutilon-like ■	Jamaica 1818. C p.1	SI. jam. 1.234 4		

XI. DU*BII.—Doubtful to which of the sections they belong.					
17994 mododus B. R.	one-toothed ■ O or 2	jl.jl ■ 1820. C s.1			
17995 racemiflora B. B.	racemose ■ O or 8	jlau ■ 1820. C p.1			
17996 Richardsia Lindl.	Richardson's Jor 3	jns ■ 1820. S co	Bot reg. 875		
17997 macrophyllus Box.	long-li-aved ■ or 12	jl.jl ■ 1810. C p.1			
17998 borbonicus Lk.	Bourbon ■ or 10	jlau ■ Bourton 1820. C p.1			
17999 gangticus Lk.	Ganges ■ 6	jl.jn.au ■ 1800. C p.1			
18000 parviflorus Wein.	small.flowred ■ it.au ■	■ 1823. C p.1			
18001 scndncs Box.	scandent ■ n.au ■	R Indies 1810. C p.1			
18002 cancellatus Box.	latticed ■ n.au ■	E Indies 1817. C p.1			
18003 setibus Rox.	setose ■ lau ■	R Indies 1800. C p.1			
18004 pulchellus Lo. C.	beautiful ■ Lau ■	R Indies 1890. C p.1			

XII. LAOUNA*RIA.—Involucels scarcely any, or l-leaved deciduous.					
18005 Patersomi H.K.	Paterson's £ i_Jor 15	jn.au P Norfolk 1.1792. C s.p	Bot rep. 285		
Lagun <i>c</i> ?i Patersonia B. M.					

18006 cuneiformis Dec.	wedge-shaped f u Jor 15	jn.au P N. HolL ... C s.1			
18007 popdinea Corr.	Yap\ta-leaved J D or 30	.. W R Indies 1770. C p.1	Hu.am!aW		
i/ibiscus populinus L.					
2 guadalensis Dec.	Guadaioupe f O or SO	.. Guadalon. ... C p.1			
18008 grandiflora Dec.	great-flowered f O or 30	.. ■ PortoRicol827. C p.1			

- | 2016. GOSSYTIUM L. COTTON. | | (Gothnemsegiar, its Egyptian name.) | | Malvaccat. 11.—16. | | | | | |
|--|--|--|----------|----------------------|---------------------------|--|--|--|--|
| 18009 herbaceum L. | common herbaceous | O ag | 3 jl | Y | s.l. Indies 1594. | | | | |
| 18010 mirrunthum Cav. | small-flowered | £ OJ or | 3 jl.au | Y | S.p. Cav. dis. a 1649. | | | | |
| 18011 Mndicum Lam. | Indian | OJ ag | 3 jl.au | Y | S.p. Cav. dis. 6.183 | | | | |
| 18012 arboreum L. | tree | 5 CD or | 12 jl.au | Y | S.p. Hu. am. 4.12 | | | | |
| 18013 wtifohum Lam. | Vine-leaved | O ag | 3 ... | Y | S.p. Cav. dis. 6.195 | | | | |
| 18014 hirsMum L. | hairy | if OI ag | 3 jl.au | Y | S.p. Cav. dis. 6.166 | | | | |
| 18015 religibsum L. | religious | Jg EJ ag | 3 jl | Y | S.p. Cav. dis. 6.167 | | | | |
| 18016 latifolium Mur. | broad-leaved | * Q ag | 5 jl.au | Y | S.p. Cav. dis. 6.164.1 | | | | |
| 18017 acuminatum Rox. | acuminate | • • ag | 5 jl.au | Y | Mur. 177a 1 | | | | |
| 18018 barbadense L. | Barbados | if OJ ag | 5 s | Y | Barbadoes 1759. | | | | |
| 18019 obtusifolium Rox. | obtuse-leaved | I Q a g | 5 jn.au | Y | S. fl.p. Bot reg. 84 | | | | |
| 18020 heterophylla yen. | various-leaved | O or | 3 jn | Y | S. co Ven. eels 11 | | | | |
| 2017. REDOUTS'A Ven. | | REDOUTEA. (P. J. Redout? [*] , celeb. French bot. draughtsman.) | | Malysear. 1.—2. | | | | | |
| 18020 heterophylla yen. | various-leaved | O or | 3 jn | Y | & Amer. 1822. S co | | | | |
| 2018. LOPI'MIA Mart. | | LOPIMIA. (Lophnos, easy of decortication.) | | Malyhceee. 1 | | | | | |
| 18021 malacophylla Mart. | soft-leaved | ■ • op _r | 5 jl.s | R | 1823. S s.p L.fco.ab.1.30 | | | | |
| 18021 malacophylla Lk. | | | | | | | | | |
| •2019. PALA^VIA Cay. | | PALAVIA. (A. Palau Verdera, second prof. bot., Madrid.) | | Malvdcear. 2. | | | | | |
| 18022 malvarf lia Cav. | Mallow-leaved | O un | lijn.au | P | 1794. S co | | | | |
| 18023 moschktia Cav. | musk | O un | 1 jl.au | P | Cav. din. 1. II. 4 | | | | |
| | prostrate Hort. | | | | Cav. dis. 1.11.5 | | | | |
| 2020. CRISTA^RIA Cav. | | CRISTARIA. (Crista% rrestj form of capsules.) | | Malvdcea?. 1.—4. | | | | | |
| 18024 coccinea Ph. | scarlet | ■ A pr | i jl.s | S | Missouri 1811. D p | | | | |
| | A/lva coccinea Nut. | | | | Bot mag. 1673 | | | | |
| 2021. A'NODA Cav. | | ANODA. (Anodos, impervious; capsule cells more united than in Ada.) | | Malvdc. 7. | | | | | |
| I. STELLOTJJ.—Cells of carpels ending in subspiouse muerones, stellately disposed. | | | | | | | | | |
| 18025 hasftata Cav. | h&lbett-leavcd | O Jun | ljjn.jl | B | Mexico 1795. S s.p | | | | |
| 18026 triloba Cav. | three-lobed | LQJ un | la jl.s | P | Mexico 1720. S s.p | | | | |
| Sida cristate W. | | | | | Cav. din. 1.10.3 | | | | |
| 18027 Dillcnidae Cav. | Dillcnius's | iQJun | ljjn.n | B | Mexico 1725. S co | | | | |
| A'da cristate B. M. | | | | | Bot.mag.S30 | | | | |
| 18028 triangularia Dec. | triangular | [Qjun | 1 jl.au | B | Mexico 1820. S s.p | | | | |
| Sida dcltfoldes Horn. | | | | | | | | | |
| 18029 incarnate H.% | B. flesh-colored | OI un | 2 jlau | F | Mexico 1824. S s.p | | | | |
| II. MLVICA.—Carpels avunless. | | | | | | | | | |
| 18030 acer'fbUa Dec. | Maple-leaved | (Q/un | 1 jl.au | B | Mexico 1809. S s.p | | | | |
| Sida hastate B. M. | not of others | | | | Bot mag. 1541 | | | | |
| 18031 parviflora Cav. | smaU.flowered | iQJun | ljjl.au | Pa.Y | W.Indies 1820. S s.l | | | | |
| 51da crenatiflora Pers. | | | | | Cav.ic.5.431 | | | | |
| 2022. PERI'PTERA Dec. | | PERIPTERA. (Peripptera, a shuttlecock; flowers resembling.) | | Malvace*. 1. | | | | | |
| 18032 punicea Dec. | crimson Shuttlecock* | • pr | 3 myjn | C | N.Spain 1814. C co | | | | |
| | Sida Periptera B. M. A'noda punicea Lag. | | | | Bot. mag. 1644 | | | | |
| •2023. STDAL. | | SIDA. (A name used by Theophrastus.) | | Malvhceee. 98. —198. | | | | | |
| I. MALVI'NDA.—Carpels 5 or 12, Lseeded, not bladdery. | | | | | | | | | |
| \ BRSNES.—Pedicels not exceeding the petioles ; leaves oblong or ovate. | | | | | | | | | |
| 18033 /iniftia Cav. | Flax.leaved | MI un | 3 jl | Pk | Guinea 1822. S co | | | | |
| 18034 angustifbia Lam. | narrow-leaved | £ ES un | lftil.s | Y | Brazil 1726. S co | | | | |
| 18035 viminea Lk. | twiggy | Jg El un | 2 jn.au | O | Brazil 1821. S co | | | | |
| 18036 spinbsa JL | spiny | j£ OJ un | 1U1 ■ | Y | E. Indies 1680. S co | | | | |
| 18037 alba /.. | white^-otw? red | O un | 2 jjjl | W | E. Indies 1732. S co | | | | |
| 18038 stipulatae Cav. | stipulate | « I I un | 2 jl.au | y | Mauritius 1819. S co | | | | |
| 18039 jamaicensis Cav. | Jamaica | tt-Dun | 2 jl.au | Y | Jamaica 1817. S co | | | | |
| 18040 bracteolate Dec. | bracteolate | • i_J un | 2 ■ | Y | Chile 1824. C co | | | | |
| 18041 carpimflta L. | Hornbeam-lvd sh | I un | 3 ■ | Y | Canaries 1774. C co | | | | |
| 18042 carpinooides Dec. | Hornbeam-like tl | O un | 2 jlau | Y | S. Amer. 1800. S co | | | | |
| 18043 tridentata Cav. | three-toothed | tt I un | 1J jl.au | Y | W. Indies 1824. S co | | | | |
| 18044 Schrankf Dec. | Schrank's | j£ ESI un | 1 jl.au | Y | Brazil 1820. S co | | | | |
| brasUsa Lk. | | | | | Cav.ic.4.S12 | | | | |
| 18045 eroaal Jr. | bitten | ES un | 2 jl.s | Y | 1824. S co | | | | |
| 18046 ciffans X. | ciliated | O Jun | ljjn.s | Y | Jamaica 1759. S co | | | | |
| 18047 alnifuUa L. | Alder-leaved | EQ un | 2 jl.s | Y | E. Indies 1732. S co | | | | |
| ii. ■—Pedicels elongated, distinctly jointed ; leaves oblong or ovate. | | | | | | | | | |
| 18048 frut^scens Cav. | frutescent | n. -Tun | 2 jl.au | Y | Mauritius 1810. S co | | | | |
| 18049 compressa Dec. | compressed | Q un | 2 jns | Y | Nepal 1823. S co | | | | |
| 18050 canariensis W. | Canary | O un | 2 jns | W | E. Indies 1820. S co | | | | |
| 18051 rhombifilia W. | rhomb-leaved | tkt ■ | 2 jn.au | Y | India 1732. S co | | | | |
| 18052 rhomboidea Box. | rhomboid | ■ ljjl.un | Y | E. Indies 1818. S co | | | | | |
| 18053 retCisa L. | retuse | nQun | T Y | Mauritius 1818. S co | | | | | |
| philippica Cav. | | | | | Cav. dis. 1.3.4 | | | | |
| 18054 redsa Schrank | cut | ■ ■ ■ | 2 jn.au | Y | Brazil 1823. S co | | | | |
| 18055 calyxhymena Gay | membrane-cal. | LJ un | 2 jnjl | Y | M. Holl. 1820. S co | | | | |
| 18056 irificans Cav. | glittering | £ tAlun | ljjn.au | Y | E. Indies 182a. S co | | | | |
| 18057 maculata Cav. | spotted | ■ ■ | ji.au | Y | StDomin. 1818. S co | | | | |
| suberbsa Herit. | | | | | Cav. dis. 1.1.1 | | | | |
| 18058 acuminate Dec. | acuminate | - - - | T | W. Indies 1800. S co | | | | | |
| Hi. CountbLME.—Pedicels eh leaves at base cordate, toothed, not lobed. | | | | | | | | | |
| 18050 pilosa Cav. | pilose | ■ ■ ■ | T | St Domin. 1793. S co | | | | | |
| 18060 hiimilis W. | humble | ■ ■ ■ | T | E. Indies 1800. S co | | | | | |
| 18061 sup^na Heri^. | supine | ■ ■ ■ | T | Jamaica 1821. S co | | | | | |
| 18062 rotundifolia Cav. | round-leaved | ■ ■ ■ | T | Bourbon 1819. S co | | | | | |
| 18063 multiflora Cbv. | many-flowered ML | ■ ■ ■ | T | Brazil 1818. S co | | | | | |
| 18064 atgUta Sun. | | | | | Cav. dis. 1. a 6 | | | | |
| 18065 cordifolia L. | heart-leaved | ■ ■ ■ | T | W. Indies 1732. C co | | | | | |
| 18066 altheifolia Am. | Althauuleaved | ■ ■ ■ | T | C. G. H. 1732. S co | | | | | |
| 18067 virens i. | stinging | ■ ■ ■ | T | Jamaica 1820. C co | | | | | |
| | | | | | SL jam. 136.8 | | | | |
| | | | | | Jamaica 1781. C co | | | | |
| | | | | | Cav. dis. 1.2.7 | | | | |

18068 <i>verticiflora Cav.</i>	whorled	«. Q un 2 JUu Y	Brazil 1823. S co	Cav.dls. 1.Llt
18069 <i>dumosa Swx.</i>	bishy	• u n 2 jl.i. Y	Jamaica 1818. C co	
18070 <i>paniculata L.</i>	panicled	£ O un 1J jl.s P	Jamaica 1795. S co	Cav. dis. 1.12.5
atrosanguinea Jac.				
18071 <i>triloba Cav.</i>	iv. PALMATILOBAVJE.—Leaves palmately deft into 3,5,7 or 9 lobes.			
18072 <i>jatrophoides Herit.</i>	threelobed	t j u n 3 jl.s W	C. G. H. 1794. C co	Jac. sc. 2.142
18073 <i>graveolens Rox.</i>	Physic-nuW/Aff	O un 4 au V	S. Amer. 1787. S co	Her. at. 1.56
18074 <i>nicindoides Herit.</i>	strong-scented	tf Ol un 1 jlau V	E. Indies 1820. S co	
18075 <i>Napsai Cav.</i>	Ricinus-like	O un 4 au W	Peru 1818. S co	Cav. dis. 1.3.3
<i>NapaOa IsVis L.</i>	Napea	£ A un 4 au.s W	Virginia 1748. D co	Bot. mag. 2193
18076 <i>dioica Cav.</i>	dioecious	jt A un 6 au.s W	Virginia 1759. D co	Cav. dis. 5.132.9
<i>Napaei dioica L.</i>				
18077 <i>wialvaeflbra Dec.</i>	Mallow-flwd	^ A or 2 jl.au P.B	Colombia 1826. D p.l	Bot. reg. 1036
18078 <i>occidentalis L.</i>	II. ABUTILOPPES — Carpels 15 or 40, l-seeded, bladdery.			
18079 <i>fce'tida CAV.</i>	occidental	O un li jl.au Y	America 1733. S CO	Di.el.7.6.6
visctaa Herit.	stinking	O un 1} jl.au Y	Peru 1795. S co	Her. st. 1.53
18080 <i>brripes Dec.</i>	short-stalked	O un 1 jl.au Y	St Martha 822. S co	
III. ABU^TILON.—Carpels five to thirty, many-seeded, often bladdery*				
	i. OLIGOCARPIE — Carpels five or eight.			
18081 <i>periplocifolia L.</i>	Periploca-lvd	£ O un 2 jl.au Y	India 1691. T co	Di. el. 4.3.2
2 zeylanica	Ceylon	£ OJ un 2 jl.au Y	Ceylon --	Pluk.al.74.7
3 caribe*a	Caribbee	tf Ol un 2 jl.au Y	W. Indies ...	SI.jam. 139.3
18082 <i>ferruginea Dec.</i>	rusty	n. I. un 2 jl.au Y	Peru 1822. T co	
18083 <i>Lechcnaultidna Dec.</i>	Lechenault's	n. • un 2 jl.au Y	E. Indies 1820. T co	
18084 <i>hermandioite* Heril.</i>	Hernandia-/wl	O un 6 -- Y	Hispanic 1798. T co	Her. st. J. 58
18085 <i>luciana Dec.</i>	St. Lucia	a. I. un 2 jnau W	St Lucia 1810. T co	
18086 <i>nudiflora Herit.</i>	naked-flowered	ill un 3 my.jn Y	Peru 1731. T co	Her. st 1.59
18087 <i>polyfinta Uc.</i>	many-flowered	1 • un 3 my.jn Y	... 1821. T co	
18088 <i>aurua Wai.</i>	large-eared	3 my.jn Y	Bengal 1823. T co	
18089 <i>triquetra L.</i>	triangular	■■■ 2 jl.au Y.P	W. Indies 1775. T co	Jac. vin. 2.118
18090 <i>incana Lk.</i>	hoary	■■■ 3 jl.au Y	Sandw. Is. 1818. T co	
18091 <i>pubella Bonp.</i>	neat	■■■ 2 jlau W	N. Holl. 1824. T co	Bot mag. 2753
18092 <i>umbellata L.</i>	umbellcd	■■■ liiU Y	Jamaica 1788. T co	Jac. vin. 1.16
18093 <i>gigantea Jac.</i>	giant	■■■ 5 jl.au Y	Trinidad 1820. T co	Jac.sc. 2.141
	ii. POLYCA'RPJE.—Carpels nine or more.			
18094 <i>reflexa L.</i>	reflex-flowered	• CD un 3 jl.au R	Peru 1799. T co	Her. st 1.64
18095 <i>crspa L.</i>	curled	O un 1 jl.au W	Carolina 1726. T co	Cav. dis. 5.135.2
18096 <i>virgata Cav.</i>	twiggy	tt. • un 11 jn.au Y	Peru 1818. T co	Cav. ic 1.73
18097 <i>arborea L.</i>	tree	* • un 6 jl.au Y	Peru 1772. T co	Her. st 1.63
18098 <i>mauritana Jac.</i>	Mauritius	E2 un 2 jl.s Y	Mauritius 1789. T co	Jac. ic 1.137
planiflbra Cav.				
18099 <i>riliifolia Fis.</i>	Lime-tree-leaved	iQJun 2 jl.au Y	China 1821. T co	
18100 <i>americana L.</i>	woolly	□ un 1) jl.au Y	Jamaica 1730. T co	
18101 <i>Abutilon L.</i>	Abutilon	O un 14 jn.au Y	India 1596. T co	H.sys.61
18102 <i>asiatica L.</i>	Asiatic	0 un U jlau Y	E. Indies 1768. T co	Cav. dis. 1.7.2
18103 <i>Sonnerattni W.</i>	Sonnerat's	■ 01 un 2 jnjl Y	C. G. H. 1806. T co	Cav. dis. 1. & 4
18104 <i>populiflria W.</i>	Poplar-leaved	im un 1 jl.au Y	R. Indies 1796. T co	Cav. dis. 1.7.9
18105 <i>hiita Lam.</i>	hairy	OJun 2 jl.au* Y	E. Indies 1820. T co	Ru. am. 4.10
pilbsa Herit.				
18106 <i>mlliissima Cav.</i>	softest	■ OJun 2 jn.jl Y	Peru 1789. C eo	Cav.dis.2.14.1
dstiflbra Herit.				
18107 <i>grandifolia W. en.</i>	large-leaved	i Q u a 8 n.d Y	Peru 1816. C co	Bot Teg.360
m6Us Or.				
18108 <i>perm6Us W.</i>	very soft	* O un 4 jlau Y	S. Amer. 1817. S co	
18109 <i>orbiculata Dec.</i>	orbicular	• i J un 3 jn.jl Y	China 1820. C co	
18110 <i>indica Z.</i>	Indian	M O un 1) jl au Y	India 1731. S co	Cav.dis.1.7.10
18111 <i>vesicKria Cav.</i>	bladdery	* i J un 3 jlau Y	Mexico 1822. C co	Cav. dis. 2.14.3
18112 <i>glatica Cav.</i>	glaucous	■■■ Dun 2 jl.au Y	Guinea 1824. S co	Cav. ic. 1.11
18113 <i>dibida W.</i>	whitish	* i Jun 3 jlau W.Y	Canaries 1822. T co	
18114 <i>mollicmna W.</i>	soft-haired	B-O un 3 jlau Y 1816. T co	
18115 <i>pdchra Cott.</i>	fair	tLCUor 4 jlau Y	W. Indies 1824. T co	CoL h. rip. 34
18116 <i>aceriflria Lag.</i>	Maple-leaved	tk i J un 3 jlau B	N. Spain 1822. T co	
IV. DITBUE.—Form of fruit unknown, and therefore doubtful to which of the sections they belongs leaves linear, oblong-ovate or lanceolate, from No. 18122. cordate.				
18117 <i>Milleri Dec.</i>	Miller's	0 vin	— 1749. S co	
18118 <i>semicrenata Lk.</i>	half-crenate	E J un	Manilla 1823. C co	
18119 <i>ActiMina Horn.</i>	Birch-like	F Alun.	S. Amer. 1820. S co	
18120 <i>Oacriththa ZJfc.</i>	pointed-flwd	Dun	Brazil 1820. T co	
18121 <i>Jpiraeifolia Lk.</i>	Spirea-leaved	■■■ un	— 1824. T co	
18122 <i>brasilicensis Cav.</i>	Brazilian	■■■	Brazil 1818. T co	
18123 <i>villtea Mil.</i>	villoua	■■■	S. Amer. 1739. T co	
18124 <i>verruculata Dee.</i>	small-warted	■■■	Brazil 1822. T co	
18125 <i>purpurascens Lk.</i>	purpurascens	■■■	Pk Brazil 1822. T co	
18126 <i>iatatns Andr.</i>	spreading	■■■	Y Abyssinia 1806. T co	
18127 <i>contracta Lk.</i>	contracted	■■■	Y Madagascar 1823. T co	
18128 <i>conferta Lk.</i>	clustered	■■■	Y Brazil 1822. T co	
18129 <i>lasioscga Lk.</i>	hairy-topped	■■■	Y Brazil 1824. T co	
18130 <i>pentacarpus Rox.</i>	five-capsuled	■■■	Y E Indies 1820. S co	
2024. LAGUNJTif Cav. LAOUNBA. {Andreas Laguna, a Spanish naturalist, d. 1543.) Malvaceat. 2.-4				
18131 <i>krtApa W.</i>	lbed-leaved	O un 3jl.a.u W	Bourbon 1787. S co	Cav.dis.5 1S&1
18132 <i>sinuata Horn.</i>	sinuated	Ol un 2 jlau Y	E. Indies 1817. S co	«*•*•» i
angulata Hort.				
2025. nULZIA Cav. RDIZIA. (D&n Hippolito Ruiz, one of the authors of Flora Peruvian^				
18133 <i>vanabilis Jflft</i>	Variable-Wtawl	• or 6 my W	Bourbon 1792. C pl. Ja. S	
18134 <i>lobata Qw.</i>	lobato	— or 6 ... W	Bourbon 1816. C pl. S	
i*.8R ERIOOUTNA Ike. EWOUMTA. Carton, wool, lama, cloak; calyx woolly.)				
18135 <i>Wallich't Dec</i>	■ U or « ... R	i Indies 1823. C		

- | | | | | |
|--|--|--|-------------------|--------|
| 2027. CAROL'NE/I L. | CAROMNEA. | {Princess Sophia Caroline, of Baden.) | Bombœcia. | 4.—5. |
| 18136 alba <i>Lod.</i> | white | CDspl 20 j <u>U</u> W Brazil 1817. C pi Bot.cab.752 | | |
| 18J77 princeps <i>L.</i> | princely | Jspl20 ... R.Y.G. W.Indies 1787. C p.l Aub.gui.291.2 | | |
| 18138 infnor <i>H. K.</i> | less | J spl 20 jl.au K.Y.C Guiana 1798. C p.l Bot mag. 1412 | | |
| 18139 insignia <i>Swx.</i> | showy | J spl 20 ... It W. Indies 1796. C p.l Bot. cab. 1004 | | |
| 2028. ADANSO*N/i4 L. | ADANSONIA. | {Michel Adanson, a famous French botanist.) | Bombūcete. | 1. |
| 18140 digitata <i>L.</i> | digitated | Sour Gourd 1 • ec SO ... W Guinea 1724. C p.l Bot. mag. 2791-2 | | |
| 2029. BOWS AX L. | SILK-COTTON | TREE. (Bombax, cotton: wool in pods.) | Bombœce*. | 4.—11. |
| 18141 malabarium <i>Dec.</i> | Malabar | ± • tm 60 ... s Malabar ... S p.l Hox. cor. 3.247 | | |
| heptaphyllum <i>Hox.</i> | | | | |
| 18142 globosum <i>Ahb.</i> | globose | f p.tm 60 = — Guiiana 1824. C p.l Aub. gui. 2.281 | | |
| 18143 Ceiba <i>L.</i> | Cciba | I O tm 100 = W S. Amcr. 1692. C p.l Jac. au. 176.1 | | |
| 18144 quinatum <i>Jac.</i> | | | | |
| 18144 septenatum <i>Jac.</i> | aeven-leaved | 1 • tm 50 = W Carthage. 1699. S p.l Pluk. al. 188.4 | | |
| 18144 heptaphyllum <i>L.</i> | | | | |
| 2030. ERIODE'NDTON Dec. | ERIODBNHDON | {Erion,vioo\dcndron, tree: wool in pods.) | Bombūccce. | 4.—5. |
| 18145 lcanthérum <i>Dec.</i> | smooth-flwd | JU O tm 70 ... S Brazil 1818. S p.l Cav. dis. 5.152.1 | | |
| Bombax crianthos <i>Cav.</i> | | | | |
| 18146 anfractuosum <i>Dec.</i> | winding | £ • tm 100 ... S £. Indies 1739. S p.l R. ma. 3.49 | | |
| Bombax pentandrum <i>L.</i> | | | | |
| 18147 guineense <i>G. Don</i> | Guinea | t • tm 130 ... S Guinea 1826. S p.l | | |
| 18148 caribajum <i>G. Don</i> | Caribbean | f • tm 70 ... S W. Indies 17J9. S p.l Jac.am.pic.l7&70 | | |
| 2031. MYRO*DIA Schreb. | MVHODIA. | {Myron, fragrant balsam, odtn, smell; odor of plant.) | Bombūcece. | 1.—3. |
| 18149 turbinata <i>Swx.</i> | turbinate | ft • or 6 ... W W. Indies 1793. C p.l | | |
| 2032. COCHLOSPE'RMUM Kth. | COCHLOSPERMUM. | (Kochlo, to twist, sperma, seed) | Ternstrcnnidcete. | 2. |
| 18150 Gobspium <i>Kth.</i> | cottony | J a or 60 — Y E. Indies 1824. C p.l Son. it. 2.153 | | |
| Bombax 6?osspium <i>L.</i> | grandiflūrum | Sonn | | |
| 18151 serratifolium <i>Kth.</i> | serrated | ± • or 60 ... Y Mexico 1820. C p.l Fl. mex. ic. in. | | |
| Bombax witifolium <i>W. en.</i> | | | | |
| 2033. GORDO*N/i4 EL | GORDONIA. | {James Gordon, a nurseryman at Mile-End.) | Ternstrccmiūccee. | 4.—6. |
| 18152 Laatanthus <i>L.</i> | Hairy-flower | fflc or 6 au.n Y N. Amer. 1739. L P.l Bot. mag. 668 | | |
| 18153 Hammatuxylon <i>Swx.</i> | Jam. Ked-wood | 9 • tm 40 ... W Jamaica 1820. S Lp | | |
| 18154 pubfscens <i>IV.</i> | pubescent | i& or 4 au.n W Carolina 1774. L s.p Ven. mal. 1 | | |
| Lacathla flurida <i>Sal.</i> | | | | |
| 18155 Franklin/ Herit. | Franklin's | ■ — 4 au.n W N. Amer. 1774. L s.p | | |
| 2034. STUA'RIVM Cav. | STUARTIA. | {John Stuart, Marquis of Bute, a distinguished bot.) | Ternstr. | 1. |
| 18156 virginica <i>Dec.</i> | common Virginian | 3f or 10 my.au W N. Amer. 1742. L Lp Bot. rep. 397 | | |
| Malachodendron <i>L.</i> | | | | |
| 2035. MALACHODE'NDRON Cav. | MALACHODENDRON. | (Malakos, aot,t,dendron, tree.) | Ternstroemihcea. | 1. |
| 18157 ovatura <i>Cav.</i> | ovate | \$ or 9 J U u W N* Amer. -1785. L Lp Bot. reg. 1104 | | |
| Stuartia pentagynia <i>Herit.</i> | | | | |
| 2036. POLY'SPORA Swt. | POLVSPORA. | {Polys, many, spora, seed; fruit.) | Ternstrccmiūcete. | 1. |
| 18158 axillaris <i>Swt.</i> | axillary | * • or 3 f.mr W E. Indies 1818. C r.m Bot. reg. 349 | | |
| Camcilia axillaris <i>B. B.</i> and <i>B. M.</i> | | | | |
| 8037. THE'A L. | TEA. | (Tc/ia, the Chinese name for tea.) | CameUtecE. | 2.—4. |
| 18159 Bohia <i>L.</i> | Bohea | H1_Jclt 4 aud W China 1768. C Lp Bot mag. 998 | | |
| chinensis/3 Bohea <i>Dec.</i> | | | | |
| 18160 viridis <i>L.</i> | green | ■L_ clt 4 f.n W China 1768. C Lp Bot. cab. 227 | | |
| chinensis * viridis <i>Dec.</i> | | | | |
| •£038. CAME'LLML. | CAMELLIA. | {George Joseph Kamel, or Camellus, a Jesuit.) | Camell&ec. | 6.—7. |
| 18161 curvooides <i>B. B.</i> | Eurya-like | * J or 44 myjn W China 1824. C p.l Bot reg. 983 | | |
| 18162 oleifera Abel | oleiferous | i J or 4 myj W China 1819. C p.l Bot. cab. 1065 | | |
| 18163 Kissi Wai. | Kissi | i J or 10 myjl W Nepal 1823. C p.l | | |
| 18164 SasfinquaThun. | Lady Banks's Sasan. | J Lpr 4 f.n W China 1811. C p.l Bot. reg. 12 | | |
| 3 plena rubra | double red | J Jpr 4 f.n Pk China 1818. C p.l Bot. reg. 5*7 | | |
| 3 plena Alba | double white | J Jpr 4 f.n W China 1824. C p.l Bot. reg. 1091 | | |
| 4 semiplena Alba | semidouble white | J Lspl 4 f.n W China 1811. C p.I Bot. reg. 12 | | |
| 18165 reticulata A, Br. | reticulated | J Lspl 6 apjn R China 1824. C p.l Bot. reg. 1078 | | |
| 18165? japonica <i>L.</i> | common Japan | J t 1 spl 10 my R China 1739. C p.l | | |
| Garden Varieties. | | | | |
| 1 r̄bra | single red | J spl 10 f.my R China 1739. C | Bnt mag. | 42 |
| 2 f̄ba | single white | J Lspl 10 f.my W China | p. Bot. cab. | 653 |
| 3 semidūplex | semidouble red | J spl 10 f.my R China | p. Bot. rep. | 559 |
| 4 rubro-plena | double red | J spl 10 f.my R China | p. Bot. rep. | 199 |
| 5 cárnea <i>Middlecmis</i> Vs | flesh co/or | J spl 10 f.my R China | p. Bot. reg. | 1670 |
| 6ftyrtiflMia | Myrtlc/rtrwrf | J spl 10 f.my R China | Bot mag. | 1670 |
| 7 atrorfibcns | Loddiges' dark red | J spl 10 f.my R China | | |
| 8 anemoniflra War. | Anemone-flwd | J spl 10 f.my R China | | |
| 9 versicolor | variegated Waratah * | J spl 10 f.my S China | | |
| 10 pezoniasfibra | I»ony.flou cred | J spl 10 f.my Bh China | | |
| 11 varigckta | double striped | J spl 10 f.my Bh China | | |
| 12 Pompnia | Keut blush Poinponcl | J spl 10 f.my Bh China | | |
| 13 flavescens | flavescens Hume's blush* | J spl 10 f.my W China | | |
| 14 alba pl&a | double-white M | J spl 10 f.my W China | | |
| 15 Welbankii | Welbank's * | J spl 10 f.my W China | | |
| 16 Kinvolulta | Lady Long's involute* | J spl 10 f.my R China | | |
| 17 variabilis | variable invtite Waratahkt y | J spl ... f.my W China | | |
| 18 hexangulkris | hexangular | J V spl W f.my R China | | |
| 19 dianthiflra | Carnat.-flwd Warat. | J LJ spl 10 f.my R China | | |
| JO Kiteo&lba | pale yellow M LJ spl 10 f.my Pa. Y China | | | |
| 21 fimbriata | fringed white * J spl 10 f.my W China | | | |

22 crassincrvls	thick-nerved	• i J ^s P ¹	1° f.my	■	China	■	■
S3 exp&nsls	expanded	il [] spl	10 f.my	■	China	■	p;
21 conchiflora	shell-flowered	» i J spl	10 f.my	■	China	■	p;
25 rubricaulis	L. Campbell's red-stmd* ^t	[] spl	10 f.my	■	China	■	p;
S16 longifolia	long-leaved	m [] spl	10 f.my	■	China	■	p;
27 aucubefolia	Aucuba-leaved	* i J spl	10 f.my	■	China	■	p;
28 coccinea	scarlet	il [] spl	10 f.my	■	China	■	p;
29 grandiflora	great-flowered	m [] spl	10 f.my	■	China	■	p;
SO nana	dwarf	m L] spl	10 f.my	■	China	■	Bot. cab. 140
31 incarnata	incarnate	* [] spl	10 f.my	■	China	■	p;
32 blanda	blush Waratah	il [] spl	10 f.my	Bh	China	■	p;
33 Rossii	Ross's	il [] spl	10 f.my	R	China	■	p;
34spatuulta	spatulate	* i J spl	10 f.my	R	China	■	p;
35 straminea	straw-colored	• [] spl	10 f.my	Str	China	■	p;
36 Aitton*	Alton's large single redHt	[] dpi	JO f.my	R	seedling	■	Chan. cam. 3
37 filthasiflora	Hollyhock-flwd	il [] spl	10 f.my	R	seedling	■	Chan. cam. 4
38 corallina	coral-flowered	il [] spl	10 f.my	S	seedling	■	Chan. cam. 5
39 insignia	splendid	* [] spl	10 f.my	S	seedling	■	Chan. cam. 6
40 fl6nda	flowery	il [] spl	10 f.my	R	seedling	■	Chan. cam. 7
41 anemoniflora alba	white Anem.-fl.	i [] spl	10 f.my	W	seedling	■	Chan. cam. 8
42 Chandler*	Chandler's str. Waratah*	i J spl	10 f.my	St	seedling	■	Chan. cam. 1.
43 punctata	Gray's spotted	il L] spl	10 f.my	W.pk	seedling	■	p;
44Ausamfindi	Rose of the World*	i [] spl	10 f.my	W.c	seedling	■	p;
45 PTCSM	Press's single-red*	[] spl	10 f.my	R	seedling	■	p;
46clipsis	Press's eclipse	il [] spl	10 f.my	W.pk	seedling	■	p;
47 rubro-punctata	single red-spotted*	i J spl	10 f.my	W.a	seedling	■	p;

2039. BARRINGTCTNJ/f Font. BARRINGTONIA. (*Hon. Daines Barrington, F.R.S.A.S.*) *Myrtaceae*. 1.
18167 specifcsa Forst. showy £ • spl 20 ... 8 E. Indies 1786 S r.m Ru. am. 3. Hi

2040. STRAVA^DIJUM J. STRAVADIUM. (*Tsjeria Samstravadi*, its Malabar name.) *Myrtaceae*. 8.
18168 acutangulum J. sharp-angled f O or 20 ... P E. Indies 1822. L sp R. raal. 4.7

Barringtonia acutangula Rox. *Eugenia acutangula* L.
18169 racemusum Swt. racemose i^m O or 20 ... W E. Indies 1822. L s.p R. raal. 4. 6
Barringtonia racemosa Rox. *Eugenia racemosa* L.

S041. GUST A?VIA L. GUSTAVIA. (*Gustavus III. king of Sweden, patron of Lin.*) *Myrtaceae*. 1. — 2.
18170 augusta L. august £ O sp 11 O ... W •Guiana 1791. C r.m Aub. gui.l.iyj

2042. CARITY/f Rox. CAREYA. (*William Carey, bot and nat, its discoverer.*) *Myrtaceae*. 3.
18171 herbacca Rox. herbaceous £ El spl J.l.au R E. Indies 1808. D 1. Rox. cor. 3.217
18172 sphcTrica Rox. round ■• spl 3^m ... R E. Indies 1823. C pS
18173 arLbrea Rox. tree ■• spl 10 ... RY E. Indies 1823. C p.i Rox. cor. 3.218

2043. itfNDROSATMUM All. TUTSAN. (*Aner, a man, haima, blood; color of Juice.*) *Hyperidnece*. 1.
18174 officinale All. officinal ■ A or 2 jLs Y Britain woods. Sk co Eng. bot 1225
Hypericum jfadrossVnum L.

2044. il'SCYRUM L. ASCVRUM. (*A, priv., skyros, roughness; plants smooth.*) *Hypericetace*. 5.-6.
18175 pumilura Mx. dwarf £ A) P* I in au Y Georgia 1806. C l.p
18176 crux A'ndre^ L. St. Andrew's cross* [] pr 2 Jl Y N. Amer. 1759. C l.p
18177 Hypericoides L. Hypericum-like* [] pr . Y N. Amer. 1759. C l.p
18178 stans Mx. standing * [] pr Jl Y M. Amer. 1806. C l.p Ven.mal.90
18179 amplexicaule Mx. stem-clasping il i [] pr Jl.s Y N. Amer. 1823. C co

2045. MORISO'NJif Plu. MORISONIA. (*R. Morison, prof. of bot. at Oxford, d. 1683.*) *Capparide**. t
18180 americana L. American £ O or W? W. Indies 1824. C p.i Jac. am. 97
Capparis Morisni Sun.

CLASS XVII. ORDER 1.

DIADE'LPHIA PENTA'NDRIA.

2046. MONNIE^R/il W. MONMBRIA. (*M. le Mounter, an able bot. phys. to Louis XV.*) *Rubiaceae*. 1.
18181 trifolia W. three-leaved G2jun lijl.au W Guiiana 1792. 8 s.l Aub.gui.2.293

CLASS XVII. OKUKR 2.

DIADE'LPHIA HEXA'NDRIA.

*2047. CORYD.VLIS Dec. CORYDALIS. (*Korydalis*, ancient Gr. name for fumitory.) *Fumariacet**. 22.—S3.

I. CAPNI^TES.—Roots tuberous s stems simple ; caulin leaves few, alternate.

18182 pauciflora Pers.	few-flowered	A A or L	f.ap	P	Siberia	1819.	D p.l	Deless.*. 2.9. A
iPumaria pauciflora Step.								
18183 MarshalliaM Pers.	Marshall's	A A or	f.ap	P	Tauria	1824.	D p.l	Deless.2 10
18184 tuberdsaZto.	tuberous hollow-rooted	& A or	f.ap	P	Europe	1596.	D co	Bot. mag. 232
2 albiflora	white-flowered	A A or	f.ap	W	Europe	1596.	D co	Bot mag. 234
J <i>Pumaria cava albiflora B. M.</i>								
18185/abScea Pers.	Bean-leaved	A A or	f.fap	P	Germany	1815.	D co	Fl. dan. 1394
18186 caucasica Dec.	Caucasian	A A or	f.fap	P	Caucasus	1823.	D p.l	
Fum'ria/abacea Bieb.								
18187 bulbosa Dec.	bulbous	A A or	I f.my	Pk	Britain	groves.	D co	Eng. bot. 1471
FU'maria sfilida Sm.								
18188 angustifolia Dec.	narrow-leaved	A A or		P	Iberia	1819.	D p.l	
18189 bracteata Pers.	large-bracted	A A or	I f.ap	Pa.Y	Siberia	1820.	D p.l	
18190 nobilis Pers.	noble-flowered	A A or	I my	L.Y	Siberia	1783.	D p.l	Bot mag. 1953

11. CAPNOI^DES.—Roots fibrous ; stems branched ; caulin leaves numerous, alternate.

18191 piEontuefbia Pers.	Pacony-leaved	A or	1 f.ap	P	Siberia	1820.	D p.l	Gm.si.4.34
18192 glafica PA.	glaucous	O or	2 jl.au	Y.P	N. Amer.	1683.	S co	Bot. mag. 179
i^umaria sempervirens L.								
18193 stricta Step.	strict		1 my.jl	Y	Siberia	1827.	S p.l	
18194 siberica Pers.	Siberian		iJl.au	Y	Siberia	1810.	S p.l	
18195 impatiens Fis.	impatient		1 my.ap	Y	Siberia	1823.	S p.l	
18196 aurea W. en.	golden		1 my.jl	Y	N. Amer.	1812.	S co	Bot reg. 66
18197 Kitea Pers.	yellow		1) ap.o	Y	England	oldw.	D co	Eng. bot. 588
18198 capnoldes Pers.	Capnus-like		2 my.o	W	S. Europe	1596.	S co	Pluk.al.90.2
18199 acaulis Pers.	stemless		4 my.o	Pa.Y	Hungary	1825.	S co	Jac.ic.3.554
18200 uralensis Fis.	Ural		1 au	Pa.Y	Altai	1824.	S co	
18201 breviflora Dec.	short-flowered		2 jn.jl	Pa.Y	Kamch.	1824.	S p.l	
18202 claviculata W.	tendrilled	J.	6 jn.jl	W.Y	Britain	thick.	S co	Eng. bot. 103

*8048. CYSTICATNOS Boer. CTBTICAPNOS. (*Kystis*, a bladder, *kapnos*, fumitory.) *Fmnariikce**. 2,

18803 africana Gae.	African	± O or	2 jn.jl	K,w	C. G. H.	1696.	S s.l	Boer. 1.300
Furaaria vesicaria L.								

18204 alexandrina G. Don	Alexandrian	_i O or	2 jn.jl	W.R	Alexandr.	1827.	S co	
--------------------------	-------------	---------	---------	-----	-----------	-------	------	--

*fio49. DIE'LYTRA Bore. DIELYTRA. two, elytron, a sheath ; base of flowers.) *FumariacetB.* 9.

18205 Cucullaria Dec.	Monk's hood	or jn.jl		W	N. Amer.	1731.	D s.p	Bot mag. 1127
J^umaria Cucullaria L.								

18206 bracteola Dec.	bracted	or	1 jn.jl	W	N. Amer.	1823.	D p.l	
18207 form&sa Dec.	handsome	or	1 jn.jl	F	N. Amer.	1796.	D p.l	Bot. mag. 1335
18208 exlmia Dec.	choice	or	1* jn.jl	F	N. Amer.	1812.	D p.i	Bot. reg. 51

18209 s>ectifibill Dec.	remarkable	or	1 jn.jl	P	Siberia	1810.	D p.l	Am. ac. 7.457.7
18210 specibsa Nois.	showy	or	1 jn.jl	F	.	.	1810.	D p.l
18211 tenuifolia Dec.	fine-leaved	or	f jn.jl	Pk	Kamch.	1820.	D p.l	Deless. 2.9.13
Corydalis tenuifolia Ph.								

18212 canadensis Dec.	Canadian	S	jn.jl	Pk	Kamch.	1820.	D p.l	
Ibyrd&lis canadensis Go/.								
18213 lachenali^y2oraDec.Lachcnalia.flwd^		1 jn.jl		W	N. Amer.	1819.	D co	Bot. mag. 3031.

•2050. ADLITML4 Ran ² . ADLUMIA.	(Adlum, an American writer on Vines.)			P	Siberia	1826.	D p.l	Ac. pet 1.19
18214 cirrhosa Rafi.	cirrose	I C J o r	15 jn.s	W.p	N. Amer.	1778.	S s.l	Sw. fl. gar. 189

Corydalis fungbsa Ven.								
------------------------	--	--	--	--	--	--	--	--

2051. SARCOCA'PNOS Dec. SARCOCAPNOS.	(Sax, flesh, <i>kapnos</i> , fumitory; leaves.)	<i>Fumaridcete.</i> 1.—2.						
18215 enneaphyila Dec.	nine-leaved	^ ^ or	1 my.jl	P.v	Spain	1714.	D co	Boc. mu. 2.73.1

-Fumkria enneaphylla L.								
-------------------------	--	--	--	--	--	--	--	--

8052. JTJMA'RIA Tou. FUMITORY. (Fumus, smoke; disagreeable smell of plant) *Fumarihcete.* 8.—10.

I. PLATYCA'PNOS.—Silicles compressed.

18216 spicata L.	spiked	Q or	8jl.au	F	S. Europe	1714.	S co	M.h. 3.12.11
------------------	--------	------	--------	---	-----------	-------	------	--------------

II. SPHAROCA'PNOS.—Silicles globose.

18217 capreoflata L.	tendrilled	1 O o r	my.s	Europe	...	S	■■	Dec. ic. 1.34
2 Burchellii	Burchell's	O cu	1 mr	C. G. H.	1816.	S	■■	

18218 m^dia LoL	intermediate	O or	jn.jl	Britain	corn fi.	S	■■	Eng. bot. 943
capreolata E. B. not L.								

18219 omcinalis L.	officinal	O w	2 my.au	Pk	Britain	eul.gr.	S co	Eng. bot 589
18220 parviflora Lam.	small-flowered	O w	2 au.s	Pk	England	corn fi.	S co	Eng. bot 590

18221 micrEntha Lag.	small-flowered	O or	1 my.jl	Pk	Spain	1823.	S co	
18222 Vaillantii LoL	Vaillant's	O or	1 my.jl	Pk	England	1800.	S co	Vail. pa. 10.6

18223 densiflora Dec.	dense-flowered	O or	1 my.jl	Pk	Montpel.	1824.	S co	
-----------------------	----------------	------	---------	----	----------	-------	------	--

CLASS XVII. ORDER 3.

DIADEXPHIA OCTA'NDRIA.

• 2053. POLY'GALA Tou. MILKWORT. (*Polys*, much, *gala*, milk; supp. to increase.) *Polyghle**. 48.— 168.

I. PBTCHA'NTHIW.	—Keel much crested; capsules smooth, marginate; braeteas three at base of pedicels, often persistent; leaves opposite, from No. 18229, alternate
18224 oppositifolia L.	opposite-leaved I H Jor 3 my.au R C. G. H. 1790. ■ i.p Bot. reg. 636
18225 cordifolia Thun.	heart-leaved I or 3 mr.au P C. G. H. 1791. ■ s.p Bot. mag. 2438
18226 tetragoia Burc.	four-angled I or 2 year P C. G. H. 18'JO. ■ s.p
18227 Nummularia-A	Nummularia-A I 3 year P C. G. H. 1812. ■ ff Bot reg. 645
18228 latifolia Ker	broad-leaved I or 1) my.jn P C. G. H. 1820. ■ ff Bot reg. 645
corditha B. M.	
18229 borbonw'lio Burc.	Borbonia-lvd I 3 year P C. G. H. 1790. C B.p Bot. mag. 498
o;>posititola B. At.	
18230 myrtifolia L.	Myrtle-leaved I 3 my.au ■ C. G. H. 1707. C p.l Bot. reg. 669
18231 grandiflora Lod.	large-flowered I 4 rayo ■ C. G. H. 1818. c s7p Bot. cab. 1227
18232 tenuiflora Lk.	fine-leaved I 3 apjn ■ C. G. H. 1823. C p.l
18233 iliiflora Ma. C.	Lily-leaved I 4 my.o ■ C. G. H. 1820. C s.p
18234 ligustris Ker	trap-leaved I 1) iny.au ■ C. G. H. 1820. C s.p Bot. reg. 637
18235 intermedia Dec.	intermediate I 3 my.o ■ C. G. H. 1816. C s.p
18236 bracteolata L.	bractcoldte I 6 my.o ■ C. G. H. 1713. C 8.p Bot. mag. 345
18237 Burmanni Dec.	Burmann's I 3 my.o ■ C. G. H. 1713. C ap Bur. af. 73.4
18238 umbellata Thun.	umbellate I 1 my.o ■ CO. H. 1713. S s.p Bur. af. 73.5
1H2,9 simplex Burc.	simple-stemmed I 4 my.o ■ C. G. H. 1816. C s.p
18240 speciosa B. M.	showy I 6 my.o ■ C. G. H. 1814. C B.p Bot. reg. 150
18241 teretifolia Thun.	cylindrical-lvd I 3 my.au P C. G. H. 1791. C P.p Hot rep. 370
18242 pinifolia Lam.	Pine-leaved I 3 my.o P C. G. H. 1823. C e.p Lam. il 598.8
18243 gem&toddes Pair.	Broom-like I 3 my.o P C. G. H. 1823. C s.p
18244 Garcinia L.	Garcin's I 3 my.o P C. G. H. 1820. C s.p Bur. af. 73.3
18245 attenuata Lod.	attenuated I 3 my.o P C. G. H. 1820. C s.p Bot. cab. 1000
18246 lanceolate Lo. C.	lance-leaved I 3 my.o P C. G. H. 18*. C s.p

II. POLYGALON. —Keel crested; capsules smooth; braeteas three at base of pedicels, soon falling off.

18247 major Jac.	larger <i>Austrian</i> I 1 jl.au II Austria 1739. D 6.1 Jac. au. 5.413
18248 vulgaris L.	common ft my.jn B Britain dr.pl. D BI Eng. bot. 76
18249 amara L.	bitci jn B Europe 1775. D I.p Bot. mag. 2437
18250 monspellaca L.	Montpellier i jn,jl B Mediterr. ... S 8.1 Dec. ic. 1.9

III. TIMU'TEA. —Keel crested (crest often very minute); capsules smooth, oval-oblong, scarcely emarginate; racemes sometimes elongated, sometimes crowded, spike-foamed; flowers small; leaves alternate or verUcillate.

18251 incarnata L.	flesh-colored O or 1 'jnjl Fk N. Amer. 1812. S co
18252 crucifera L.	cross-leaved O or i jn,jl G.p N. Amer. 1739. S co
18253 breviflora Nut.	short-leaved O pr i jn,jl ... N. Amer. ... S p.l
18254 fastigata Nut.	peaked O pr i jn,jl ... N. Amer. ... S p.l
18255 lutea L.	yellow O or 1 jn,jl Y N. Amer. 1739. S co Pluk. am. 438. 6
18256 nana Dec.	dwarf O or i jn,jl G.Y N. Amer. ... S co
vindescens Nut.	
18257 purpurea Nut.	purple O or 1 jl.s G.p N. Amer. 1739. S co Pluk. am. 438.5
sanguinea Mx.	
18258 sanguinea L.	blood-flitaf O or i jn,jl ■ N. Amer. 1824. ■ P.1
viridescens Poir.	
18259 ambigua Nut.	ambiguous O or ij Lau N. Amer. 1739. ■ co Pluk. am. 438.4
18260 verticillata L.	vrhor-leaved O pr ij Lau PaP S Amer. 1822. ■ co Bot. reg. 761
18261 paniculata L.	panicled O pr W Carolina 1824. ■ s.p
18262 graininifolia Poir.	Grass-leaved A or i jn,jl W Carolina 1824. ■ s.p

IV. SE'NEGA. —Keel beardless i the three outer sepals of calyx subequal; leaves alternate.

18263 Senequa L.	Rattlesnake root^t A or \$ jl R.w N. Amer. 1739. S co Bot mag. 1051
V. CIAMJRBIXUB. —Lower sepal of calyx concave-cuculate; inside at base embracing a gland.	
18264 paucifolia W.	few-leaved Jl A or \ my.au P N. Amer. 1812. D s.l
18265 Chamebitus L.	Bastard Box C or J my.jn Y Austria 1658. Sk al Bot mag. 316

2054. MURA'L TJJ Neck. MURALTIA. (*John von Muralt*, K Swiss botanist.) *Polyghlett*. 14.— 39.

18266 Hcisterfa Dec.	Heister's Polygala Heisteria L. ■ 6 ja P C 6. H 1787. C s.p Bbtmag. 340
18267 filojuco'ides Dec%	Foxtaulike ■ 3 my.au P C. G. H. 1800. C i.p Bot mag. 1006
i alopcuroides L.	
18268 squar!>sa Dec.	squarrose ■ ILLJor 3 my.jl P C. G. H. 1820. C s.p
18269 stipulke'ca Burc.	stipuled ■ 3 aps R C. O. H. 1801. C s.p Bot mag. 11%
18270 juniperif'Mia Dec.	Juniper-leaved ■ L or 3 aps P C. G. H. 1810. C s.p
18271 diffusa Burc.	diffuse il t L or 3 year P C. G. H. 1800. C s.p
18272 virguta Burc.	twiggy m L or 3 year P C. G. H. 1812. C s.p
18273 /inophila'la Burc.	lax-leaved *i or 3 my.jl P C. G. H. 1816. C s.p
18274 macrocarpas Burc.	long-horned ■ 3 year P C. G. H. 1820. C s.p
18275 mixta L.	mixed ■ 3 year P C. G. H. 1791. C s.p Bot mag. 1714
Polygala mixte Andr.	
18876 humili Swt.	humble ■ i or 1 my.jl P C. G. H. 1818. C s.p Bot cab. 420
Polygala humili Lod.	
JBVTI ciharis Burc.	ciliated ■ I or 3 my.jn P C. G. H. 1824. C i.p
18278 filiformis Dec.	filiform ■ L-Jor lft my.d P C. G. H. 1812. C ap
Polygala filiformis Thun.	
18279 micrantha W.	small-flowered M i_J or lft year P C. G. H. 1800. C s.p Bot rep. 384

* 3055. MU'NDI A Kih. MUNDIA. (<i>Mundus</i> , neat; appearance.)	P, ■■■■■, L
18280 spinosa Dec.	spiny ■ 11_I or 3 ja.my P C. G. H. 1780. ■■■■■
Polygala spinosa L.	
8 angustifolia Dec.	narrow-leaved ■ luJor 3 ja.my P C. O. H. 1800. C s.p
Polygala viminea Hout.	

2056. SECURIDA*CA L. SECVMOACA. (*Securis*, a hatchet; form of end of pod.) *Tolyehlece*. 2.—8.
 18281 volubilis L. twining JD f r a l f l ... W W. Indies 1739. C p.l Jac. am. 183.38
 18282 virgata Swt. twiggy tC J f i r a l O ... W Jamaica ... C p.l P.ed.B.248.1

CLASS XVII. ORDER 4.

DIADE'LPHIA DECA'NDRIA.

2057. PLAGICLOBUM Swt. PLAQIOLBUM. (*Plagios*, transverse, *lobos*, a pod.) *Leg. Pap. Lbt. Genii. I*
 18283 ilicifolium Swt. Holly-leaved H t J or 5 mr.jl B N. Holl. 1824 C s.p Sw. au. S
 H6vefliciR>liaCkw. #
 18284 chorizemifolium Sal. Chonzema-lvd il i_J or S mr.jl B N. Holl. 1826. C s.p
 Hbvcfli chorizemifolia Dec.

2058. ACHYRO^NIA JP. ACRYRONIA. (*Achyron*, chaff; branches and leaves chaffy.) *Leg. Pap. Lbt. Genis. 1.*
 18285 villusa W. oi. villoiu A ^jfor 3 jlau Y K. Holl. 1819. C pi Wen.her. J. 12

2059. LIPA^RIA L. LIPARIA. (*Liparos*, brilliant; surface of leaves.) *Legum. Pap. Lbt. Genis. 1.*
 18286 sphac'rica X. globe-flowered *LJ or 4 jl.au O C. G. H. 1794. C p.l Bot mag. 1241

2060. PRIESTLEY Dec. PRIESTLEYA. (3/ Priestley, physiological botan.) *Leg. Pap. Lbt. Genis. 14.—15.*
 I. EI9OTHIB^A.—*Calyx thrust in at base.*

- 18287 myrtifolia Dec. Myrtle-leaved *LJ or 3 ... Y C. G. H. 1893. C p.l Dec. leg. 29
Liparia myrtifolia Thun.

- 18288 hirsuta Dec. hairy-stemmed *L-J or 3 ap.d Y C. G. H. 1792. C p.l Bot. reg. 8
Liparia hirsuta Thun.

- 18289 tevigata Dec. smooth-leaved m v_J or 3jl.au Y C. G. H. 1820. C pi Dec. leg. 30
, Liparia Isvigata Thun. Borbnia Isvigata L.

II. ANBISOTHE\.—*Calyx ovate or obconically attenuated.*

- 18290 capitata Dec. capitate *LJ or 3 jl au Y C. G. H. 1812. C p.l
Liparia capitata Thun.

- 18291 «apinifolia Dec. Grass-leaved *uJor 3 ju.jl Y C.G. H. 1800. C co
Liparia graminifolia L.

- 18292 tress Dec. cylinder-leaved *LJ or 3 ... Y C. G. H. 1816. C p.l
Liparia tress Thun.

- 18293 ericifolia Dec. Heath-leaved *i_J or 3 jn.jl Y C. G. H. 1812. C p.l Dec. leg. 31
Borbntia iricefbria L.

- 18294 sericea Dec. silky-leaved «Lj or 3 jnjl Y C. G. H. 179* C p.l
LipHria sericea L.

- 181295 axillaris Dec. axillary JI LJ or 3 jn.jl Y C. G. H. J822L C p. I Dec. leg. 32
Borbntia axillaris Lam.

- 18296 elliptica Dec. elliptic #J or 3 ... Y C. G. H. 1825. C s.p Dec. leg. 33
 18297 villbsa Dec. woolly #uJ or 3 jn.jl Y C. G. H. 1774. C co H.n.h.5.29.1
Liparia villbsa L. Borbn<*i tomentbsd L.*

- 18298 vestita Dec. clothed «uJ or 3 myjn Y C. G. H. 1800. C p.l Bot. rep. 382
Liparia vestita Thun.

III. DirBiJE.—*Calyx not known; and hence doubtful to which of the sections they belong*

- 18299 tomentbsa Dec. downy *LJ or 3 jl.au Y C. G. H. 1812. C co
Liparia tomentbsa Thun.

- 18300 umbellifera Dec. umbelliferous I J or 3 ... Y C. G. H. 1626. C p.l
Liparia umbellifera Thun,

2061. MEDICA^GO L. MEDICK. (*Medike*, by Diosc. to a Median grass.) *Leg. Pap. Lbt. Trif. 40.—76.*

- I. "HJPVLijRih.—Lcgumcn renifom, falcate or subcochleate, smooth or pubescent; margins entire; leaves trf foL

- 18301 rupestris Bib. rock Jc H or 1 jnjl Y Tauria 1820. D co

- 18302 brachycarpa Fts. short-podded Jc O cu } jnjl Pa.Y Tifliz 1823. S co

- Jelilotus brachiflora Fts.

- 18303 lupulinai. Hop-like TrefoUjk O ag 1 my.au Y Britain ■ S co Eng. bot 971

- 18304 mniicarpa IVallr. Moss-fruited Jc O cu 4 jn.au Y Switzerl. 1816. S co

- luplna y unguiculata Dec.

- 18305 Willdenbw Hent. Willdenow's Jc O cu 1 jnau Y Europe 1816. S co

- 18306 cretacea Bib. chalky Jc ^ cu 4 jl Y Tauria 1805. D s.l

- 18307 falcta L. falc&te-pods & A ag 2 jl Y England bor. fl. D co Eng. bot 1016

- 18308 proctimpbons Bes. trailing Jc ^ or 1 jnjl Y S. Europe 1H18. D co

- 18309 cancellata Bleb. laticet Jc ^ or 1 jnjl Y Caucasus 1818. D co

- 18310 suffruticosa Ram. suffruticose & £ or 2 jnjl V.Y Pyrenees 1820. D co

- 18311 arborea L. tree Moon-trefoil & or 8 my.n Y IÜly 1596. C 11 LoUic. 2.46.2

- 18312 sibArica Dec. Siberian O or 1 jnjl Y Siberia 1817. S co

- Aclifitius sibtrica Pair.

- 18313 media Pers. intermediate ^ A or 2 jn.jl B Europe 1817. D co

- 18314 sativa L. cultivated Lucern£ A ag 2 jnjl V England me.pa. D r.m Eng. bot. 1749

- 2 verecioior various-colored A A or 2 jnjl Y.B Britain ganl. D co

- 18315 nostrata Jac. prostrate 3 A cu f jnjl Y Hungary 1793. D s.l Jac. vin. 89

- 18316 glomerata Balb. clustered ^ A cu 1 jnjl Y Italy ... D s.l

- 18317 glutinosa Bib. glutinous Jc A cu 1 jnjl Y Tauria ... D co

II. SnsocA'BPOS.—*Legumen cockleate-orbicular, pilose or smooth; margins entire, rough or spiny; Ivs tr\^f.*

1. Legumens with thin unarmed margins.

- 18318 obschra lietx. doubtful Jc O cu 1 jlau ■ 1734. S co Rets. 1.24.1

- 18319 IffiVis Desf. smooth -* O cu i jn.au ■ S. Europe 1816. S co

- Jielix W.

- 18390 orbicularis Alt. round-podded Jc O cu 1 jlau Y S. Europe 1688. S co M.h.^15.1

- 1|321 marginato W. en. margined Jc O cu 1 jlau Y S. Europe 1816. S co M.h.2.15.8

- 18322 aplanaflata W. flat Jc O cu ijn.au Y S. Europe 1810. S co

- 18323 scutellata AU. dish-like Jc O cu 1 jn.au Y S. Europe 1562. S co M.h.2.15.3

- *18324 rugosot. rugose Jc O cu 1 jlau Y Sicily 1680. S co M.h.2.15.4

- elcans W.

ti. Racemes sessile.							
18396 monspeliaca <i>L.</i>	Montpelier	■ or 1	jnjl	■	Montpel.	1710.	S f.l W.&K.8 142
18396 pinnatifida <i>Cav.</i>	pinnatifid	■ or i	jn.au	■	Spain	1801.	S 81 Cav. i.d. 38
media <i>Del.</i>							
18387 polycerata <i>L.</i>	many-horned	O or	1	jl.8	■	S. Europe	1640. S 81
18398 egyptica <i>Poir.</i>	Egyptian	Jt O or	1	jn.au	■	Egypt	1818. S co
18399 ornithobathra <i>Horn.</i>	Bird's bill	O or k	jn.au	■	Russia	1818. S co	
IV. FALCA'LULA. — flowers umbellate, pedunculate ; legumes compressed, falcate, reticulate, mucronate.							
18400 ruthenica <i>L.</i>	Russian	-4 A or	1	jnjl	Y	Siberia	1741. S p.l Gm.si.4.8
18401 mhdia <i>Lag.</i>	intermediate	-* A or	1	jn.au	Y	1818. S co
18402 platycarpus <i>L.</i>	broad-podded	O or	1	jns	W	Siberia	1741. S co Gin. si. 4.9
18403 trifida <i>Pou.</i>	hybrid	<* A cu	1	jns	W.v	France	1806. S 81 Dec. ic. 1.29
18404 corniculata <i>L.</i>	small-horned	^ O fra	J	jnjl	Y	& Europe	1597. S 81 M. b. 2.16.11
18405 ornithopodus <i>Urtdes</i>	Bird's-foot-lik.*	O or ft	jnjl	R	Britain	bar.he. S co	Eng. bot. 1047
7'rifium omithopodioides <i>L.</i>							
18406 esculentum <i>IV. en.</i>	esculent	£2 dt 1\$	jn.au	■	E. Indies	1815. S s.l	
2064. POCOCKIJA Ser. POCOCUA. (Richard Pococke, a traveller in the Levant.) Leg.Pap.Ut.Trif. 1.							
18407 cr&ica Ser.	Crete	O, or	4	jn.au	Y	Candia	1713. S co M. h.2.Ü.3
2065. MELILO'TUS Tou. MELILOT. (Me/i, honey, lotos, lotus j bees gather.) Leg.Pap.Lbt Trif. 28.-29.							
I. COSLOBUVIS.—Legumes lacuno&ely rugose.							
18408 Kochina W. en.	Koch's	■ ■	3	jns	■	Germany	1816. S co
	TYibium Kochianum Hay.						
18409 dentata IV. en	toothed	^ A un	3	jn.au	Y	Hungary	1802. D co W.&K.1.46*
1H410 lintersi Cav.	linear	O un	2	jlau	Y	Spain	181ft. S co
18411 ruthenica Bieb.	Russian	^ Q un	1	jlau	W	Russia	1830. ■ co
18412 melanocephala Bes.	black-seeded	5 Q) un	1	jlau	Y	Tauria	1819. S co
18413 officinalis W. en.	officinal	O cc	1	jlau	Y	Britain	bus. pi. S Eng. bot 1340
	TVifolium JV/ellitus officinalis L.						
18414 pahistum Sur.	marsh	un	3	ji.au	Y	Hungary	1820. ■ ■ W.&K.266
18415 arbbrea Castag.	tree	■ or	15	jn.au	W	Turkey	1826. ■ ■
18416 altissima rAif.	Ullest	■ & Q or	1	jlau	W	S. Europe	1816. ■ ■ [K.M.]
&da Lam.							
18417 leucantha tfocA vulg*ris IV. en.	white-flowered	^ Q) un	4	jl.s	W	Europe	— S co Stur. deu. 1.15.
	7%ifi.						
18418 macrorhiza P*vy*.	long-rooted	■ A un	3	jl.au	Y	Hungary	1801. ■ co W.&K.1.2
18419 indica Pers.	Indian	O un	1	jl.jn.au	Y	India	1680. ■ co Pluk.al.45.4
18420 parviflora Desf.	small-flowered	O un	2	jn.au	W	Barbary	1798. ■ co
rugulosa W.en.							
18421 Segetalis Ser.	Sword.graas	O cu	liJUu	Y	Spain	182a	■ co
18422 polonica Pers.	Polish	2	jn.au	L.Y	Poland	1778.	■ co
18423 taurica Bieb.	Taurian	■ Q un	3	jls	W	Tauria	1818. ■ co
18424 italica IMW	Italian	un	2	jl.au	Y	Italy	1596. ■ co
18425 rotundifolia Ten.	round-leaved	un	3	jl.au	Y	Italy	1823. ■ co
18426 gracilis Dec.	slender	8 un	2	jl.s	Y	France	1820. ■ co
18427 pflida Bes.	pale	cu	3	jl.s	Y	Volhinia	1816. ■ co
18428 suaveolens Led.	sweet-smelling	* A or	3	jl.au	■	Dahuria	1824. ■ co
II. PLAGIORUVIB.—Legumens transversely and subarcuate rugose.							
18429 arvensis Wattr.	corn	O un	2	jlau	Y	Germany	1800. ■ co
18430 Pettipierreana Hay.	Pettipierre's	& Q) un	2	jns	W	Germany	1816. ■ co
18431 Beneridiana Ser.	Besser'i	5 Q) un	4	jlau	Y	Tauria	1816. ■ co
plicata Stev.							
18432 messanensis Desf.	Messina	O un	3	jn.au	Y	Sicily	1680. ■ co
18433 sulcata Desf.	furrowed	O un	2	jn.jl	Y	Algiers	1798. ■ co
mauritanica JV. en.							
III. DU'BIE.—Doubtful to which of the section* they belong.							
18434 Baumetti Horn.	Baumetti's	O un	1*	jl.au	Y	1818. S co
18435 neapolitana Ten.	Neapolitan	Q un	2	jn.au	Y	Italy	1823. S co
•2066. TRIFOLIUM Tdū. TREFOIL. (Tres, three/OIKWI, leaf; 3 on stalk.) Leg.Pap.Lbt.Trif. 114.—150.							
1. LAGOPIS.—Flowers spiked; spikes oblong, at base bracteate; calyx viUous, after flowering not inflated.							
18436 angustifolium L.	narrow-leaved	O pr	1	jn.au	Pa.R	S. Europe	1640. S s.l Bar. ic.608
18437 intermedium Cus.	intermediate	O pr	1	jnjl	W	Italy	1820. ■ co
18438 purpureum Lot.	purple	O pr	1	jnjl	P	France	1816. ■ co Loi. gal. 14
18439 rubra L.	red	^A pr	1	jns	D.R	S. Europe	163? D co Jac.au. 4.385
18440 circrullatum Bieb.	cwrulescent	O pr	f	jl	BE	Siberia	1^7. S co
18441 incarnatum L.	flesh-colored	O pr	1	jl	F	Italy	1^6. S co Bot. mag. 323
18442 Molineri Balb.	Mohner's	O or	1	jl.au	W.a	S. Europe	1820. S co
18443 Lagopus Pou.	Hare's-foot	O pr	1	jl.au	...	Spain	1827. S co
18444 gracile ThuU.	slender	O pr	1	jnjl	...	France	1816. S co Bar. ic. 901
arvensis var. gracile Dec.							
18445 arvensis L. field	Hare's-foot	O w	1	jl.au	F	Britain	san. fi. S s.l Eng. bot. 944
18446 lasiocephalum Lk.	hairy-headed	O pr	\	jn.au	P	C. G. H.	1823. S co
13447 ligustrinum Balb.	Lovage-like	O cu	J	jnjl	W.B	Spain	1816. S co Brot. ph. 63.2
aristatum Horn.							
18448 divaricatum Horn.	divaricate	O cu	4	jnjl	W.a	S. Europe	1B16. S co
18449 gemmifera Pou.	twin-spiked	O cu	{}	jnjl	Y.o	Spain	1818. S co
18450 ^hleofdes Pou.	Cat's-tail-hke	O cu		my jl	W.Y	Spain	1818. S co
18451 /appacum L.	Burr-like	O pr	i	jn.au	W	Montpel.	1787. S.i Bar. ic. 871
isr>2ccinatum Bieb.	prickly	O cu	J	jajl	W	Caucasus	1821. S co
18452 eriaceum Bieb.	prickly	O cu		jnjl	Pa.R	Iberia	1818. S co
18454 malacanthum Lk.	soft-flowered	O pr	1	jl	P	1824. S 81
18455 sylvaticum Ger.	wood	O cu	ijnjl	P	France	...	S co
II. PHLEASTBUM.—Flowers capitate; heads ovate, conical; calyx after flowering not infated.							
ttfSiBocc&niSavi	Boccon's	O cu	f	jnjl	Pa. P	Spain	1820. S co grot ph. (a 1
18457 stridulum L.	striated	Ow	i	jn	P	Britain	bar.gr. S s.l Eng. bot. 1843
IS tS tenuia'rcua ZVn.	fine-flowered	O cu	J	jnjl	Pa.R	Italy	1823. S co
18459 scabrum L.	rough	Jc O pr	J	myjn	W	Britain	san. fl. S 81 Eng. bot. 903

HI. EirraiPHYLLUM. — Flower* capitate heads ovate, pedunculate or sessile, often bracteate i calys vttlous,									
18460 maritimum Hud.	sea	O pr	i [*] jnjl	not inflated.	Pa.P	Britain	saLm. S	8.1	Eng. bot 220
13461 irregulkre Pou.	irregular	O pr	ft jnj.l		Pa.P	Pisania	...	S co	Savi pis. 2.1.1
rigidum Savi									
18462 suptuum Savi	supine	Jt O cu	ft jnj.l		Pa.P	S. Europe	1816.	S co	Savi trif. 46.2
18463 constantinopolitanum Ser.	Constantin.	O cu	ft jnj.l		Y	Turkey	1820.	S co	
18464 cinctum Dec	girded	O cu	1 jnj.l		Pa.Y	1820.	S co	
18465 alexandr'mum L.	Alexandrian	O pr	1 jnj.l		Pa.Y	Egypt	1798.	S s.1	
18466 ochroleucum L.	yellowish-white	H	1 my.jl		Su	England dr pa.	D s.1	Eng. bot 1224	
18467 cangscens W.	canescent	H	1 my.jl		W.v	Cappadoc.	1803.	S co	Bot mag. 1168
18468 pannonicum L.	Pannonian	H	1 myjn		W.H	Hungary	1752.	D co	Jac. ob. g.43
18469 oljrmicum Hook.	Olympic	H	1 myjn		Pa.Y	S. Europe	182a	S co	Bot. mag. 2790
18470 trichoccephalum Bicb.	hairy-headed	H	1 jnj.l		Pa.Y	Caucasus	1827.	D co	
18471 armeniuu W.	Armenian	H	1 jnj.l		W	Armenia	182G.	D s.1	
18472 squarrosum L.	squarese	H	1 jnj.l		Pa.P	Spain	1640.	S s.1	M. h. 2.13.1
2 fl&vicaus Ser.	yellowish	H	1 jnj.l		Pa.Y	Puania	1817.	S co	Savi trif. 65
squarrsum Savi									
18473 albilium Retx.	whitish	H	W.Y	—	—	—	1796.	S s.1	
18474 Lagascnum Ser.	Lagasc'a	H	W.Y	—	—	—	1818.	S s.1	
cōnicum Lag.									
18475 cōnicum Pers.	conical	H	Pa.Y	Spain	—	—	1816.	S s.1	
18476 KitaibeltfnKffl Ser.	Kitaibela	H	Pa.Y	Hungary	—	—	1818.	S s.1	
c<>niciu Horn.									
18477 aln̄stre L.	alpine	H	D.P	Europe	1789.	S co			Bot mag. 2779
18478 mēdiun L.	mediate Cow-grass	H	R	England dr. pa.	D h.1	Eng. bot 190			
18479 bractektum Schous.	large-bracted	H	P	Morocco	1804.	S s.1			
18480 exp̄nsuin Kit.	expanded	H	P	—	1830.	D s.1			
18481 v.igin'itum Schl.	sheathed	H	Pa.Y	Switzerl.	1819.	D s.1			
pratense /3 flavicans Ser.									
18482 pr.itlnsu /* meadow common Clover		H	P	Britain me. pa.	D h.1	Eng. bot 1770			
18483 n̄fricum Wul.	Noric	S A	Pa.Y	Alp. Eur.	—	S s.1			
18484 pennsyWanicum W.	Pennsylvanian	A	Pa.Y	N. Amer.	1811.	S s.1			
18485 n̄allidum IV. tf K.	lae-flowered	H	Pa.Y	Hungary	1803.	S s.1	W.&K.1.36		
1K481 ditTusum Ehrh.	diftiae	H	Pa.Y	Hungary	1801.	S s.1	W. * K. 1.50		
18487 hlrtuin All.	hairy	H	Pa.Y	Barbary	1817.	S s.1	Desf. at 209.1		
hispidum Desf.									
18488 pictum Roth	painted	H	Montel.	1800.	S s.1	Bar. ic.859			
18489 Chrleri L.	Chrleri's	H	Montel.	1750.	S s.1				
1K490 sphicror^phalon Dec.	round-headed	H	N. Afrca	1820.	S s.1				
18491 glo1^um L.	globe	H	Levant	1713.	S s.1				
18492 clypeXtum L.	shield-like	H	Levant	1711.	S s.1	Alp. ex. 306			
18493 dtellatum L.	starry	H	England so. co.	S s.1	Eng. bot 1546				
18494 leucanthum Bieb.	white-flowered	H	Tauria	1820.	S s.1				
obscCirum Cus.									
18495 obsdruum Savi	obscure	H	Pa.R	Italy	1824.	S s.1	Savi trif. 31.1		
18496 fimbnatum B. R.	fringed	H	Pa.R	Colombia	1826.	S co	Bot reg. 1070		
18497 cyathiferum B. R.	cup-bearing	H	Pa.R	Colombia	1826.	S co			
18498 tridentatum B. R.	three-toothed	H	Pa.R	Colombia	1826.	S co			
18499 saxatile All.	rock	H	Pa.R	Switzerl.	1816.	S 6.1	Alped.59.3		

IV. TRIPOLI A/trrauM. — Flowers capitate; heads globose, sessile or pedunculate s flowers often defixed; calyxes not inflated.

18500 suffbeatum L.	suffocated	* O w	1 jnj.l	W	England sea sh.	S	8.1	Eng. bot 1040
18501 congestum Cus.	crowded	Jt O or		W.R	S. Europe	1820.	S s.p	
1W02 glomer'tuin L.	cluster-Aemferf	* O w		Pk	England gra.pa.	S	s.1	Eng. bot. 1063
18503 parviflum Eht.	small-flowered	O or		W	Hungary	1820.	S s.1	
strictum Sturm								
J204 strictum L.	strict	H		W	S. Europe	1805.	S s.1	W.&K. 1.37
1B-05 ri'liens L.	creeping white Clover	Jf.		W	Britain mca.	D co	Eng. bot. 1769	
2 pcntaphyllum	five-leaved	*		W	Britain fields.	D co		
18506 anomaluu Schrank	anomalous	H		W.	—	D co	Sen. mo. 47	
18507 cOrnum Brot.	drooping	H		Pa.P	Spain 18J0.	S s.p	Brot ph. 62	
1S008 pall'scens Schreb	pall'escent	H		Pa.Y	Carinthia	1804.	D s.1	
18509 caespit&sum Reyn.	turfy	H		P	Switzerl.	1815.	D s.1	Vil.del.3.41
18510 angulatum IV. \$ K.	angular	H		R	Hungary	1803.	S 8.1	Vf.&K. 1.27
18511 suav&olens IV.	sweet-scented	H		P	Italy	1820.	S co	W. h. b. 108
18512 elegans 5ati	elegant	H		Pa.R	Europe	182a	D co	Savi pis. 2. L 2
18513 hybridum Savi	hybrid	H		P	Europe	1777.	D s.1	Mic.gen.25.2.6
18514 Midiclytum Savi	Michel's	H		W.G	Italy	1817.	S s.1	Mic. gen. 25.2
18515 reflexum L.	reflexed	H		P	Virginia	1794.	D 6.1	
18516 monUnum L.	mountain	H		W	Europe	178ti.	D co	Fl. dan. 1172
18517 rup'Mre Ten.	rock	H		W	Naples	1820.	D i p	
18518 laAnum Seb.	Latin	H		R.w	Italy	18J5.	D s.p	Seba 1.1.2

V. VKSICA'BTRUM. — Flowers capitate, dense ; lower lip of calyx unchanged, upper one after flowering rising up inflated and arched, covering the legume.

18519 subterraneum L.	subterraneous	O pr	\$ my	W	England bar.he.	S	—	Eng. bot. 1048
18520 vesiculoum Savi	bladdered	O or	3" jnj.l	R	Hungary	1805.	S 8.1	W.&K.2.1G5
recurvum W.								
18521 spumMum L.	frothy	2k O un	* jnj.l	R	France	1771.	S s.1	Cup. pa. 61.1
18522 resupinatum L.	retiupinate	* O un	i jnj.l	P	Germany	1713.	S 8.1	Bar. ic. 872
18523 fragiferum L.	strawberry-bg	* A w	i jnj.l	F	England moi. p.	D s.1	Eng. bot. 1050	
18524 tomentosum L.	woolly	* O pr	I jnj.l	P	S. Europe	1640.	S s.1	Mag. mo. 264
18525 physodes Stev.	bladdery	** A or	ft jnj.l	F	Iberia	1818.	D s.p	
18526 aratum Biv.	winged	JL A or	ft jnj.l	F	S. Europe	1820.	D s.p	Cup. pa. 97.1
Cupani Tineo								

VI. LUPINA'STER. — Flowers targe petals coriaceous, persistent, red, white, or yellow segment* of calyx subulate or erect; leaflets coriaceous, three or seven, with numerous nerves.

18527 uniAcrum L.	one-flowered	* A pr	i jnj.l	B	Italy	1800.	u s.p	Bux.c.3.31.1
2 Stembergtamt Ser.	Sternberg's	Jt A pr	i jnj.l	W	S. Europe	1822.	—	
18528 exilmium Step.	choice	Jc A pr	ft jnj.l	P	Dahuria	1820.	D sp	Bux.c.3.31.2
2 albifl&rum Fis.	white-flowered	^ A pr	ft jnj.l	W	Altai	—	5 ip	
18529 alpinum L.	alpine	k A pr	ft jnj.au	P	Europe	1775.	5	
^KO involucratum 7K	involucred	O pr	1 jnj.l	Pa.P	Mexico	1802.	D s.p	
				p co	Stur. deu. 1. IS			
				S co				

18531 Lupini'ister <i>L.</i>	LujRnaster	■ A el	lijl.au	P	Siberia	1741.	D co	Bot.mag.879
LupinAter pentaphyllum <i>Moen.</i>								
18532 purpurascens <i>Fis.</i>	purpurascent	■ A P*	1 jl-au	P	Siberia	1816.	D co	
Lupinaster y oblongifolium <i>Ser.</i>								
18533 albens <i>Fis.</i>	whitish	■ A pr	1 jlau	W	Siberia	1818.	D co	
Lupinaster /2 albiflrum <i>Ser.</i>								
18534 megacephalum M/A great-headed		■ A or	1 jl.au	Pa.P	Missouri	—	D p.l Ph. am. 2. 23	
Lupinaster macrocephalum <i>Ph.</i>								
18535 Cussonii Tinea	Cusson's	3c A pr	1 jl.au	B.a	Sicily	1826.	D co	Bonam.245
VII. CHRONO9EMIUM.—Flowers disposed into an ovate pedunculated head; petals scarious, yellow, after								
flowering bay^colored, deflexed, persistent.								
18536 badium <i>Schreb.</i>	brown	■ A pr		Pyrenees		D s.1	Bar. ic. 10.24	
	field	o pr	£ jn.au			S s.1	Fl. dan. 558	
18537 agrinnum <i>L.</i>	scarlet	■ A pr	jn.jl	Europe	1815.	D s.1	Bot. mag. 557	
18538 spariceum <i>L.</i>	deceiving	Q pr	4 jl.au			1778.	D s.1	
18539 decipiens <i>Horn</i>	showy-flowered	Jc O el		Candia	1752.	S s.1	Lab. syr. 5.10	
18540 speciosum <i>W.</i>	procumbent	Hop Jc O ag		Britain	drpa.	S s.1	Bot mag. 945	
18541 procumbens <i>L.</i>	field	Jc (j or		Switzerl.	1819.	S co		
18542 campastré Schreb.	Parisian	Jc O or		France		S co		
18543 iariucpsc Dec.	Sebastian's	Jc O or		Pa.Y	1800.	S co	Scba2.4	
18544 Sebastian://; S ivi	small-flowered	Jc O or	i	Italy	1816.	S co	Viv. lib. 19.3	
18545 inicranthuin Viv.	lesser y, llow	Jc O or		fryene	1816.	S co		
18546 minus H. JC	thread-shaped	■ Q ff	my.jl	Y	Britain	gra.gr. S s.1	Eng. bot. 1256	
18547 filiforme <i>L.</i>						gra.pa. S co	Eng. bot. 1257	
VIII. DUNIA.—Species not sufficiently known.								
18548 tenuifolium <i>Ten.</i>	fine-leaved	Jc Q or	4jn.jl	Italy	1826.	S co		
18549 clandestinum <i>Lag.</i>	clandestine	O or	i jn.jl	Spain	1826.	S co		
18550 combsum <i>L.</i>	tufted	■ Apr	4jn.jl	America	1798.	D 8.1		
2067. DORYCNIUM Ton.	DORYCNIUM. (Dory, spear; ancient plant poisoned.)	Leg. Pap. Ldt. Trtf. 8.—12.						
18551 rectum <i>Ser.</i>	upright	tt. or	2 jn.au	R	S. Europe	1640.	S co	Bar. ic. 544
Lotus rectus <i>L.</i>								
18552 latifolium IV.	broad-leaved	^ A or	1 jn.au	W	Iberia	1818.	D co	
ibéricum IV. en.								
18553 hirsutum <i>Ser.</i>	hairy	A or	3 jn.au	R.w	S. Europe	1683.	S co	Bau.his.2.360.ic
.Lotus hirsutus <i>L.</i>								
18554 tomentosum G. Don	woolly	M or	3 jn.au	R.w	S. Europe	1817.	S co	
hirsutum 0 inchnum <i>Ser.</i>	Lotus tomentosus Jiohde	scriccus Dec.	affinis lies.					
18555 parviflorum <i>Ser.</i>	small-flowered	Jc O or	1 jl.au	Y.u	S. Europe	1810.	S co	Desf. at 2. 211
/.tus parviflorus Desf.								
18556 subbiflorum <i>Ser.</i>	subbiflowered	O or	1 jl.au	...	Spain	1820.	S co	
.Lotus subbillbrus <i>Lag.</i>								
18557 herbaceum V'd.	herbaceous	& A or	1 jn.s	W	S. Europe	1802.	D e o	YU. del. 3.41
18558 sutrificosum VU.	subshruhy	n. -	■ jl.8	W	S. Europe	1640.	S p.l	Lob. ic. 2.51.1,2
monspche'nse W.	Zütus Dorycniun <i>L.</i>							
• 2068. LOTUS <i>L.</i>	BIRD'S-FOOT TREFOIL. (Probably of Egyptian origin.)	Leg. Pap. Lit. Trtf. 35—56.						
I. KROKBRIA.—Legumes turgid, curved; flowers one or two.								
18559 editlis <i>L.</i>	esculent	-* O clt	i jl.au	Y	Italy	1710.	S s.1	Cav.ic.2.157
II. LOTEA.—Legumes compressed, long; flowers subumbellate.								
18560 oriothopodioides <i>L.</i>	Bird's-loot-like	-* O pr	1 jn.au	Y	Sicily	168a	S s.1	Cav.ic.2.163
1&061 >ercprnu3 <i>L.</i>	foreign	-* O pr	i jl.au	Y	S. Europe	1713.	S s.1	
18562 indicus <i>L.</i>	Indian	-* O pr	1 jn.au	Y	E. Indies	179a	S 8.1	Pluk.al.200.7
Trigonella indica W.								
18563 flexuus <i>Lam.</i>	flexuous	-* A or	i jl.au	Y		—	1816.	S 8.1
III. EILUTS.—Legumes cylindrical, long; flowers corymbose; styles one-toothed from No. 18567. toothless.								
18564 glaucus H. K.	glaucus	■ OJ pr	1 jn.au	Y	Madeira	1777.	S s.1	
sessile-leaved	XL I or	1 jl.au	Y	Tenenffe	1810.	C s.1		
18566 rntthyllodes Ven.	Anthyllis-like	tt. pr	1 jn.au	Y	C. G. H.	1812.	S s.1	Vcn.mal.92
18567 atropurpurus Dec.	dark-purple	«, or	1 ap.d	D.P	Tcnemile	1820.	C s.1	
ic-fir	St. James's Islandtt.	i or	2 year	D.Br	C.Verdi.1714.	C r.m	Bot. mag. 79	
2Kitcus								
18569 pusillus Viv.	ye Pow-Jiowcred L.	b. or	jl.au	Y	S. Europe	1816.	S 8.1	
18570 crticus <i>L.</i>	Cretan	fU I or	lijn.s	Y	Levant	1680.	C p.l	Cav. ic. 2.156
18571 cytisoldes <i>L.</i>	Cytisus-Uke	Jc O pr	1 jl.au	Y	S. Europe	1702.	S co	Al.ped. 1.20.1
18572 sericeus <i>Ph.</i>	silky	O or	4 jl.au	Y	Missouri	1824.	S s.1	
18573 Dioscoridus All.	Dioscorides's	O pr	1 jnjl	Y	Nice	1658.	S si	AI. ped. 1. 59.1
18574 Geblia Ven.	mountain	* Al esc	my.jn	Pk	Aleppo	1773.	S 8.1	Jac vin. 2.155
18575 arabicus <i>L.</i>	Arabian	* O pr		Pk	Arabia		D co	Ven. eels 57
18576 docAmbara Pair.	dewy	O pr						
18077 pilofissimus Pair.	decumbent	Jc O or	i jl.au	Y	Europe	1816.	S si	
18578 hispidus Desf.	hairiest	Jc O or	i jl.au	Y	France	1818.	S	
18579 suaveolens Pers.	hispid.	Jc O or	i jl.au	Y	Corsica	1817.	S s.1	Loi. gal. 16
18580 anguitissimus L.	sweet-scented	■ or	i jl.au	Y	S. France	1816.	■	
18581 diffusus Sol.	narrowest-ptxWerf	* O pr	1 my.au	Y	Britain sea bca.	S	Bau. his. 2.356	
ia082 ciliatus Ten.	diffuse	-J 9 i	1J my.jn	Y	England rocks.	S 8.1	Eng. bot. 925	
18583 gracilis IV.8/K.	ciliated		ijjl.au	Y	Italy	1820.	S 8.1	
18584 coimbrensis Brot.	slender	Jc O or	1 jl.au	Y	Hungary	1812.	S s.1	W. & K. S. 229
18085 gladiémus L.	Coimbra	O pr	J in.jl	W..ii	Portugal	1800.	S 8.1	Brot ph. 53
185K6 odorMus B. M.	very smooth	Jc O or	i jl.a u	W	S. Europe	1816.	S 8.1	
18587 corniculatus E. B.	swct-scentel	Jc A "S	1J in.au	Y	Barbary	1804.	D s.1	Bot. mag. 1233
2 alpinus Schl.	small-horned	Jc A or	1J in.au	Y	Britain	pas.	D co	Eng. bot. 2090
3 crassifolius Pers.	alpine	Jc A or	i jl.au	Y	Switzerl.	1819.	S s.1	
4 villus ThuiL	thick-leaved	Jc A or	i jl.au	Y	Europe	1817.	S 8.1	
18588 Försterf Swt.	villous	Jc A or	h jl.au	Y	Switzerl.	1819.	S 8.1	
deedmbens Forst.	Forster's	Jc A "r	4JLau	Y	Britain heaths.	S co		
18589 tenuis W.S/K.	slender	Jc A pr	1 jn.au	Y	Hungary	1816.	D p.l	
18590 major Sm.	greater	Jc A ag	1 jn.au	Y	Britain	w.sh.g.	D s.1	Eng. bot. 2091
18591 pedunculatus Cav.	/wj-pcdwicled	Jc A P*	1 jn.au	Y	Spain	1814.	D s.1	Cav. ic. 2.164
18592 palAstris W.	marsh	Jc A or	ft jn.au	Y	Crete	1821.	S s.1	
IV. DUNII.—Doubtful to which of the sections they belong; the last species is probably an Indigdscra.								
185S3 argenteus H. C.	silvery	« Al or	j».au	Y	1827.	co	
2 angustifolius	narrow-leaved	■	■	Y			co	
18594 microphylla Hook.	small-leaved	■	* jn.jl	Y			co	Bot. mag. 2808

2069. <i>TETR&GONODOROBUS</i> o. TETRAONOL. (7WIW,4, angle, lobos, pod.) Leg.Pap.Lbt.Trif. 5.
18595 <i>purpureus Mae</i> , <i>purplewi Pea</i> Jk O clt 1 jlau D.F Sicily 179d. S co Botmag. 151
<i>Lotos Tetragonotobus L.</i>
18596 <i>biflorus Ser.</i> two-flowered Jk O ox J jlau Y N. Africa 1818. S 81 Desf.at210
<i>Lotos conjugates Poir.</i> <i>biflbrus Desr.</i>
18597 <i>sihqubus Roth</i> siliquose * A or } jlau Y S. Europe 16Sa D co Jac. au. 4.361
<i>Lotos siliquibus L.</i>
18598 <i>maritimus Roth</i> sea t A or } my.o Y Europe 168a D co Fl. dan. 800
18599 <i>conjugatus Ser.</i> <i>tmn-podded</i> Jk O or jlau V MontpeL 1754. S s.l
<i>.Lotus conjugates L.</i>

2070. CARMICH AE'LLJ R. Br. CARMICUAELIA. (Cap*. Diun ^h ! Cartnichael, F.L.S.) Leg.Pap.Lbt.Trif. 1.
18600 <i>australis R. Br.</i> southern ft or 2 xyli B N. Holl. 1800. C s.p Bot mag. 1364
<i>.Lotus australis Andr.</i>

2071. PSORA^LEA L. PSORA LEA. (*Psoraleos*, scurfy; appearance of calyx.) Leg.Pap.Lbt.Clit. 34.-62.

I. AXILLARIFLORAS.—Flowers single in the axils of the leaves, pedicellate or sessile.

18601 <i>odoratissima Jac.</i> most fragrant * i_J or 6 my.il Pa.B C. G. H. 1795. C pl Jacsc. 2.229
18602 <i>arborja B. M.</i> tree or my Pa.P C. G. H. 1814. p Bot.mag. 2090
18603 <i>pinnata L.</i> <i>vrlng-leaved</i> or my.jl B C. G. H. 1690. p Bot. rep. 474
18604 <i>verrucosa W.</i> waited or my.au B C. G. H. 1774. p J.i.sc.2.2261.
18605 <i>intermedia Bcs.</i> intermediate or my.jl B C. G. H. 1820. pl Jacsc.2.ic.226.2
18606 <i>axilKiris /.</i> <i>hneans Thun.</i> axillary-tud 2 Jnjl B C. G. H. 1820. 8.p
18607 <i>teniuflua L.</i> fine-leaved J.Jor 3 mr.jl W.B C. G. If. 1793. Jacsc. 2.225
18608 <i>multicaulis Jac.</i> mnny-Btalked J.Jor 3 au.o W.B C. G. H. 1793. Jac. BC. U. 230
18609 <i>aphylla L.</i> leafless J.Jor 3 jn.jl B C. G. H. 1790. Bot. mag. 1727
18610 <i>/athyriofolia Balb.</i> <i>Lathryus-lvd</i> ii J.Jor 3 jn.au B 1816. Bal. st. ic.
18611 <i>deuinbents//. if</i> trailing JU or 1 a p. my W.B C. G.'H. 1774. Bot cab. 282
18612 <i>repens /.</i> creeping JJJUU" B C. G. H. 1774.
18613 <i>hirta L.</i> hairy J.Jor 3 my.au W.B C. G. H. 1713. Jac. us. 2.228
18614 <i>aculeata L.</i> prickly L-Jor 4 jn.jl B C. G. H. 1774. Bot. imag. 2158

II. CAPITAVA.—Flowers axillary, subsessile, and at the top of branches capitate or spicate.

18615 <i>bractelata L.</i> brauted or jnjl P C. G. H. 1731. Bot. mag. 446
18616 <i>involucrata Thun.</i> involucred or jn.jl B C. G. H. 1818.
18617 <i>spicata L.</i> long-spiked J.Jor 3 jl.au B C. G. H. 1774. Bot rep. 411
18618 <i>Stachys L.</i> <i>StLcbs-leaved</i> or ap.my Br C. G. H. 1793.
18619 <i>sthta Thun.</i> stnaled J.Jor 3 my.jn B C. G. H. 1816.
18620 <i>ar^ntea Thun.</i> silvery J.Jor 3 my.jl B C. G. H. 1816.
18621 <i>tomentosa Thun.</i> woolly J.Jor 3 jnjl B C. G. H. 1820.
18622 <i>capitata L.</i> headed L-Jor 2 jl.au P C. G. H. 1793.

III. SPICAVJE.—Peduncles axillary, at top spicately capitate.

18623 <i>corylifolia L.</i> Hazel-leaved £ QJ or 2 jn.jf V "India" 173>. Bot. mag. 665
18624 <i>arcuaria Nut.</i> sand & ^I or 1 jnau P Missouri 182a
18625 <i>palastria Gou.</i> Palestine £ IAI or 2 ap.s Pa.B S. Europe 1570.
18626 <i>bituminosa L.</i> bituminous m \J or 4 ap.s V C. G. H. 1815. Lam. il. 614.1
18627 <i>Bericca Poir.</i> silky * J.Jor 3 au.o Bot. reg. 223
<i>pedunculata Ker</i>

18628 <i>esculenta PAV.</i> <i>Poikade*NI</i> <i>Peduncles axillary, at top loosely capitate</i> Ph. am. 22
18629 <i>cuspidataPA.</i> <i>cuspidataPA.</i> <i>cuspidate large-tooted A A <</i> 1 jnjl P Louisiana 1811. R pl
18630 <i>incina Nut.</i> hoary ft .Aj or 2 jl.au B Missouri 1824. R BI
18631 <i>Lupinella Mr.</i> Small Lupine i A or 2 jnjl P Carolina 1812. R pl
IIfi2 <i>mehlotoides Ms.</i> Melilot-hke 5 A or 3 au V Carolina 1814. R pi Bot. mag. 2063
18633 <i>Onobrychis Nut.</i> Saintfoin-/w & A or 3 au P N. Amer. 1818. R p1 Bot. reg. 453
18634 <i>glandulosa L.</i> glandulous Mexic. tea mt or 4 my.au Pa.B Peru 1770. C pl Bot. mag. 990
18635 <i>pubescens Balb.</i> pubescent * O or 2 au Pa.B Lima 1825. C pl Bot. reg. 968
18636 <i>canescens Mx.</i> cane&uent A J or 2 jn.au B Carolina 1821. C 81
18637 <i>divaricata H. % B.</i> divaricate m. • or 3 au P S. Amer. 1820. C pl
18638 dent ita Dec. toothed * J.Jor \$ jl.au P Madeira 1640. C pl Jac. sc. 2.227
<i>amorpha L.</i>
<i>americanica L.</i>

•2072. INDIGCVFERA L. INDIUO. (*Indigo*, a blue dyestuff, ffero, to bear.) Leg.Pap.Lbt.CM.48.—126.

I. SIMPLICIFLMI.—Leaves simple, sessile, or on very short footstalks.

18639 <i>tnifflbia Retx.</i> Flax-leaved * O or 1 jl.au P E. Indies 1792. S 81 Rox.cor.2.195
18640 <i>fflfbia Thun.</i> thread-leaved ft LJ or 1 jl.o P C. G. H. 1812. C s.l
II. OLIGOPHYLLA.—Petals elongated; leaflets one or three, terminal, one petioled.
18641 <i>diphylla Ven.</i> two-leaved * IQJ pr ijJ.au P Africa 1816. S s.p Ven. ch. 30
18642 <i>Lechenaultia Dec.</i> Lechenault's O or 1 jl.au P E. Indies 1820. S s.p
18643 <i>tnfoliU L.</i> trifoliolate JF O or \jl.au P E. Indies 1816. S e.p
18644 <i>glandulosa IV.</i> glandulous O or 1 jl.au P E. Indies 1820. S 8.p
18645 <i>denudata Jac.</i> stripped n,L or 1 J my.jl P C. G. H. 1790. C s.p Bot. cab. 500
18646 <i>amorpha H. K.</i> pleasing pur/ile-JIwd. \ or 1 mr.ap P C. G. H. 1774. C s.p Bot. reg. 300
18647 <i>microphylla Lam.</i> small-luaved tt. \ or 1 year P C. G. H. 1812. C s.p
J!S rIK'1 W,O, ri*ld • CD or 2 jl.au 11 E. Indies 1816. C B.p
18649 <i>virgata Dec.</i> twiggy a -Q o r \jl.au P E. Indies 1820 C s.p

III. MULTIJUGS.—Leaves impari-pinnate, two or many paired.

i. FRDTICOSJS.—Petioles elongated s stems shrubby.
18650 <i>argentea L.</i> hilver-leaved * a cit 2 jl.au P E. Indies 1776. E s.p Her. st 79
18651 <i>tinctdria L.</i> East Indian dyer's t CD clt 3 jl.au Pk E. Indies 1731. E s.p R.mal. 1.54
18652 <i>A*ml L.</i> Anil West Indian • CD clt 3 jl.au P W Indies 1731. E s.p SL jam. 176.3
18653 <i>GuatinMa Lunan Guatinula</i> it • clt 3 jl.au P W. Indies 1800. E B.p
tinctbris 3 brachycarpa Dec.
18654 <i>leptostachya Dec.</i> slender-spiked J. or 3 jnjl P E. Indies 1818. E s.p
18655 <i>atropurpurea Ham.</i> dark-purple J. or 3 jl.au P Nepal 1816. E s.p
18656 <i>Juncce Heteram.</i> Rush-like J. or 3 jl.au ... C. G. H. 1825. E B.p Bot. reg. 104
filifiliK'ker, aphylla J.k. l'ckck/a contaminata H. K.
18657 <i>nuda G. Don</i> naked JL Q or 1 jn.jl P C. G. H. 1820. C s.p Bot. mag. 2214
Lebeckia nuda B. M.
18658 <i>angulata B. R.</i> angled «i_J or 2 jl.jl R N. Holl. 1825. C s.p Bot. reg. 991

S081. CHJETOTALYX Jtee. CIUTTOCALYJC (*Chaita, bristle, kalyx, cai.; teeth.*) Leg.Pap.LbtQU. 1.—2.
18720 vincentina Dec. St Vincent's fc. CD or 6 my.au Y St. Vine. 1823. C p.1 Bot.reg.799
Gly-cine vincentina B. R.

2082. GLCYRRHrZA Tou.	LIQUORICE.	(G/yAy*. sweet, rhixa, root)	Leg. Pap. Lbt. Gal. 8.-9.
18721 glabra L.	common smooth^	A clt 3 jn.a	Europe 1562. R r.m Lam. il.625.2
Liquirifica officinMis Moen.			
18722 echinata L.	w'cky-headed &	A or S jn.8	Fa Italy 1596. R s.p Bot mag. 2154
18723 glandulifera W. &f K.	glanduliferous	5 A or 3 jSiau	Pa Hungary 1805. R l.? W-& K? 21
18724 lepidota Ph.	scaled suky-lvd	5 A or 3 jIau	Pa Missouri 1811. R s.l Bot.mag.S150
18725 fortida Desf.	fetid	5 A or 3 jIau	Pa.Y N. Africa 1817. R s.p Deaf. at. 2,199
18726 aapcrimma L.	roughest	jfc & clt 2 juS	LV Siberia''' ITO R «p Pal. it. ap. M. 3
aspera Pall.			
18727 uralensis Fis.	Ural	£ A or 3 jIau	Pa.B Siberia 1818. R s.p
18728 hirshta H.	hairy	5 A clt 3 jIau	Pa Levant 1739. R s.p

*2083. TEPHRCT3IA Ptvs. TEPIROSIA. (*Tephros, ash-colored; foliage.*) Leg.Pap.Lbt.Gal. 28.-79.

I. Mu3lDUVBA. — Leaves pinnate, with an odd one; racemes axillary, erect, often panicled s calyxes truncate or with five short teeth; stamens monadelphous; styles smooth, No. 18731. bearded,			
18729 sube-fsia Dec.	Cork-barked	• or 4 jn.au	R E. Indies 1818. C p.1
Robinia suberbsa Rox.			
18730 fruticosa Dec.	shrubby	* • or 6 jl.au	R E. Indies 1816. C p.l
18731 sertcea Dec.	silky	«L i J or 1 jl.au	R C. G. H. 1800. C s.l
18732 Candida Dec.	vwhite Jeaved	* O or 4 ...	Pa.R E. Indies 1816. C p.1
Robinta Candida Rox.			

II. BRISUNIA. — Leaves pinnate, with an odd one; lobes of calyx acuminate, at base broadish; stamens monadelphous; styles laterally bearded; legumes more or less velvety or hispid; flowers subfasciculate, disposed in terminal panicled racemes; peduncles axillary.

18733 toxicana Pers.	JbA-poison	m Q clt 3 ...	P.R S. Amer. 1791. C p.l Pl. ic. 135
18734 virginiana Pers.	Virginian	^ A or 4 jn.au	Pk N. Amer. 1765. C s.p Pluk.al.23.2 Galda virginica L.

III. CRACCorDES. — Leaves pinnate, with an odd one; lobes of calyx acuminate, at base broadish; stamens diadelphous; styles bearded.

18735 ochroleuca Pers.	pale-yellow	& Q or 3 ...	Pa.Y W. Indies 1799. C l.p Jac. ic 1.150 Galega ochroleuca Jac.
------------------------	-------------	--------------	--

IV. REINE%RIA. — Leaves pinnate, with an odd ones teeth of calyx linear, subulate s stamens sometime* submoitdrlphous, sometimes diadelphous.

18736 caribae^a Dec.	West Indian	i Q o t 3 jn.jl	ILW W. Indies 1786. C l.p Jac.am. 125
Galfega caribs^a Jac.			
18737 grandiflra Pers.	great-flowered	* L j or 4 my.s	Pk C. O. H. 1774. C p.l Bot. reg. 769
Galga grandiflbra Vohl			
18738 villosa Pers.	villous	■ ■ ■ ■ ■ ■	W E. Indies 1779. ■ ■ ■ ■ ■ ■ Pluk. al. 59.6
18739 purpurea Pers.	purple	■ ■ ■ ■ ■ ■	P E. Indies 1768. ■ ■ ■ ■ ■ ■ Bur. zey. 32
18740 mucronata Dec.	mucronate	■ ■ ■ ■ ■ ■	Pa C. 6. H. 1823. ■ ■ ■ ■ ■ ■
Galega mucronata Th.			
18741 lanceifolia Lk.	lance-leaved	■ ■ ■ ■ ■ ■	Pa.Y — 1820. ■ co
18742 piscatibia Pers.	fisher's	■ ■ ■ ■ ■ ■	P India 1778. ■ 1P
Galega piscatibia H. K.			
18743 capensis Pers.	Cape	■ i j or 1 jl.au	P C. G. H. 1825. C s.p.1 Jac.ic.574
18744 hypargyreia Dee.	subsilver	■ ■ OR 2 jn.jl	P E. Indies 1816. C p.1
Galega heterophylla Fa.			
18745 littoralis Pers.	shore	■ ■ ■ ■ ■ ■	P W Indies 1824. ■ s.l Jac. am. 124
18746 stricta Pers.	strict-podded	■ ■ ■ ■ ■ ■	Pk C. G. R 1774. ■ pi Scop.d.1.3
18747 linearis Pers.	\near-leaved	■ ■ ■ ■ ■ ■	R Guinea 1823. ■ s.p.l
18748 pallens Pers.	\a'e-flowered	■ ■ ■ ■ ■ ■	Pk C. G. H. 1787. ■ p.1
Galega pallens H. K.			
18749 apollinea Dec.	Apollinis	■ ■ ■ ■ ■ ■	Egypt 1816. ■ ■ ■ ■ ■ ■ Del. eg. 53.5
18750 biflora Dec.	two-flowered	■ ■ ■ ■ ■ ■	P 1816. ■ ■ ■ ■ ■ ■
Galega biflora Poir.			

V. DU'BIJ. — Doubtful to which of the sections they belong. — leaves pinnate, with an odd one.

18751 capitulata Lk.	small-headed	■ EH or ■ Ujl.au	R Owhyhc 1823. C co Jac. ic 3.572
18752 longifolia Jac.	long-leaved	■ CD or 3 jn.jl	R S. Amer. 1820. C s.l
18753 colonia /.. Tr.	Colonial	■ L or 3 jn.jl	P E. Indies 1818. C s.l
18754 pentaphylla Rox.	five-leaved	■ Oor 3 jn.jl	P E. Indies 1818. C s.l
18755 Heyneana Rox.	Heyne's	■ Oor 3 jn.jl	P E. Indies 1822. ■ s.l
18756 chinensis Lindl.	Chinese	■ ■ OR 2 jLau	P China 1826. ■ s.l

2084. NISSLPL/jf Jac. NISSOLIA. (*William Nissolle, a French botanist.*) Leg.Pap.Lbt.Gal. 7.—17.

I. NIS9OLIA^RIA. — Teeth of calyx 5, acute; stamens monadelphous.			
18757 fruticosa Jac.	shrubby	LCD or 15 JLn	Y S. Amer. 1766. S p.l Jac. vin. 2.167
18758 racemiflra Dec.	racemose	L D ° r 15 jnau	W W. Indies 1800. C s.l
18759 glabrata Lk.	polished	II c5 or 12 ...	W 1823? C s.1-

III. MACH^HIUM. — Calyx at base Mbracteolate; lobes 5, roundish; stamens diadelphous.

18760 röptera Poir.	small-winged	* or 10 jn.au	W Teneriffe 1820. C s.1
18761 Tobiniaflbia Dec.	Robinia-leaved	■ ■ or 6 ...	St. Vine. 1824. C s.1

IV. DU^BIJ{. — Doubtful to which of the sections they belong.

18762 aculeata Lam.	prickly	i. O or 12 ...	Rio Jan. 18A C s.l Lam. il. 600.2
18763 retssa W. en.	retuse	L O or 6 ...	S. Amer. 1819. C s.l

2085. LONCHOCA'RPIUS H.*B. LONCHOCARPIJS. (*Logche, Umce, karpos, fr.*) Leg.Pap.Lbt.Gal. 9.—19.

18764 macrophyllus H.8fB.	large-leaved	f O or 20	— P & Amer. 1818. C s.l
18765 domingensis Dec.	St. Domingo	• o r SO	R St Dom. 1820. C s.1
Dalbergia domingensis Pert.			
18766 violaceus H. S.B.	violaceous	J Q o r SO	— V Carthage. 1759. C s.l Jac. am. 177.46
Robinta violacea Jac.			
18767 sericeum H.8fB.	silky	2 O or 12 ...	W. Indies 1826. C s.l
Robtno serif cea Poir.			
18768 pyxidiarius Dec.	box, Palo de Caja	f O or 20	P Cuba 1820. ■ 81
18769 latifolius // \$ B.	broad-leaved	\$ O or 20	P W. Indies 180& ■ 81
Jmerimum latifolium W.	Pterocarpus* latifolius Poir.		

18770	<i>roseus Dec.</i>	<i>roae-colored</i>	<i>f</i> • or 1G	...	R	S Amer	1700.	C s.l
	<i>Robln/a hispida /9</i>	<i>rosea Mil.</i>						
18771	<i>scipium Dec.</i>	<i>hedge</i>	2 O or 20	...	P	W. Indies	1821.	C s.l Jac. am. IC9.101
	<i>Robfnia s&pium Jac.</i>							
18772	<i>pubescens Dec.</i>	<i>pubescent</i>	<i>I LJ</i> or 16	...	P	S. Amer.	1824.	C s.i
	<i>ifmerimnum pubescens IV.</i>							
2086.	ROBI'N/i* L.	ROBINIA.	<i>(Jean Robin, a French botanist.) Legum. Pap. Lbt. Gal.</i>					
18773	<i>Pseudacacia L. com.</i>	<i>Bastard Acacia*</i>	<i>tm</i> 40	<i>my.jn</i>	W	N. Amer.	1640.	S s.l Schn. ar. 1. 25
2	<i>inermis W.</i>	<i>unarmed large Jvd*</i>	<i>tm</i> 40	<i>my.jn</i>	W	N. Amer.	-	■ 81
3	<i>crispa Dec.</i>	<i>curled</i>	5	<i>or 40</i>	<i>my.jl</i>	W	N. Amer. ?	■ ■ MI
4	<i>umbraculifera Ifcw.</i>	<i>umbrella-bg J</i>	<i>or 40</i>	<i>my.jn</i>	W	N. Amer. ?	-	■ SI
	<i>Robinia inermis Dum</i>							
5	<i>tortu?sa Ztec.</i>	<i>twisted</i>	\$	<i>or 40</i>	<i>my.jn</i>	W	N. Amer. ?	■ ■ 81
18774	<i>dubia Fauc.</i>	<i>doubtful</i>	\$	<i>or 30</i>	<i>my.jn</i>	W.R	N. Amer. ?	■ ■ 81
	<i>hfbrida >< ambigua Poir.</i>	<i>echinata Mil.</i>						
18775	<i>monstrosa Hort.</i>	<i>monstrous</i>	\$	<i>or 30</i>	<i>my.jn</i>	W.R	N. Amer. ?	... S co
18776	<i>pratina 7.0. C.</i>	<i>tall</i>	\$	<i>or 30</i>	<i>my.jn</i>	W.R	N. Amer. ?	... S co
18777	<i>sophorifolia Hort.</i>	<i>Sophora-leaved</i>	Y	<i>or 30</i>	<i>my.jn</i>	W.R	N. Amer. ?	... S co
18778	<i>strida Lk.</i>	<i>stridt</i>	*	<i>or 30</i>	<i>my.jn</i>	W.R	N. Amer. ?	... S co
18779	<i>amorpho-folia Lk.</i>	<i>Amorpho-lvd</i>	*	<i>or 30</i>	<i>my.jn</i>	W.R	N. Amer. ?	... S co
18780	<i>visebsa Fen.</i>	<i>clammy</i>	tm 30	<i>jn.au</i>	W.R	N. Amer. 1797.	G s.l	Bot. mag. 5fl
18781	<i>pendula Or.</i>	<i>pendulous</i>	or 30	<i>jn.au</i>	Pk	N. Amer.	G s.l	
18782	<i>hfspida A</i>	<i>hispid Rose-A</i>	or 10	<i>my.s</i>	Pk	Carolina 1743.	G s.l	Bot. mag. 31
	<i>2 rosca</i>	<i>upright rose</i>	or 10	<i>my.s</i>	It	N. Amer. ?	G 81	
18783	<i>purensa Lk.</i>	<i>purple</i>	i	<i>or 15</i>	<i>jl.au</i>	Peru 1810.	G co	
18784	<i>nucrophylla Schr.</i>	<i>long-leaved</i>	i	<i>or 10</i>	<i>my.jn</i>	N. Amer. ?	S 81	
18785	<i>hfspida y macrophylla Dec.</i>							
18786	<i>davurica I.A. C.</i>	<i>Dahurian</i>						
18787	<i>guineensis IV.cn.</i>	<i>Guinea</i>	i • or 6	■■	W	Davuria 1820.	S 81	
						S. Leone 1822.	C p.I	
2087.	SESBAFNIA Pers.	SESTUNIA.	<i>(Sesban, Arabic name.) Legum. Pap. Lbt. Gal.</i>					
18787	<i>fflRvptica Pers.</i>	<i>Egyptian</i>	■ Oor	4	<i>jlau</i>	Y	Egypt 1680.	S co Alp. eg. 81.82
	<i>^Eschynmene S&han L.</i>							
18788	<i>occidental?</i> Pers.	<i>western</i>	■ • or	4	<i>jlau</i>	Y	W. Indies 1816.	S l.p P.cd.B.1?Al
18789	<i>aculeata Pers.</i>	<i>prickly</i>	rm or	4	<i>jlau</i>	Y	E. Indies 1690.	S co Jac.ic.a5G4
18790	<i>cannAbina Pers.</i>	<i>Heinp</i>	Oor	4	<i>jlau</i>	Y	E. Indies 1800.	S co
18791	<i>afnnis Shr.</i>	<i>kindred</i>	[QJor	3	<i>jlau</i>	Y	E. Indies 1800.	S p.I
	<i>jEschynmencn cannabina Hort.</i>							
18792	<i>punctata Dec.</i>	<i>ra-spotted</i>	O or	3	<i>jlau</i>	Y	Guinea 1825.	S p.I
18793	<i>picta Pers.</i>	<i>painted</i>	O or	6	<i>jlau</i>	Y	W. Indies 1823.	S co Botreg.873
18794	<i>marmic&rpa MM.</i>	<i>long-podded</i>	rcTior	3	<i>jlau</i>	Y	Ixmisia 1820.	S p.I
18795	<i>palutissa Jac.</i>	<i>marsh</i>	Qor	3	<i>jlaa</i>	Y	E. Indies 1816.	S p.I
18796	<i>gracilis Schr.</i>	<i>slender</i>	Oor	3	<i>jlau</i>	Y	E. Indies 1820.	S p.I
18797	<i>sericea Dec.</i>	<i>silky</i>	roior	3	<i>jlau</i>	Y	E. Indies 1818.	S p.I
18798	<i>uliginosa Rox.</i>	<i>bog</i>	O or	2	<i>jlau</i>	Y	E. Indies 1818.	S p.I
208a	<i>AGA^TI Rh.</i>	AGATI.	<i>(Agaty, its name in the Sanscrit language.) Legum. Pap. Lbt. Gal.</i>					
18799	<i>cocinea Desv.</i>	<i>scarlet</i>	fa or	26	<i>Jlau</i>	R	E. Indies 176a	C p.I Ru. am. 1.77
	<i>Sesbania coccinea Poir.</i>							
18800	<i>grandiflora Desv.</i>	<i>great-flowered</i>	■ • or 20	<i>jlau</i>	R.W.Y	E. Indies 1820.	C p.I R. mal. 1.51	
	<i>Sesbinia grandiflora Poir.</i>							
2089.	GLOTTI'DIUM Desv.	GLOTTID.	<i>(Glottis, superior opening of larynx; legume.) Leg. Pap. Lbt. Gal.</i>					
18801	<i>floridanum Desv.</i>	<i>Florida</i>	O or	2	<i>jlau</i>	Y	Florida 1816.	S s.p Jac. ic. 148
	<i>Phaca floridna W.</i>	<i>Sesbania displrma Ph.</i>	<i>vesicaria Spr. platyc&rpa Pers. iSchynomcn platyc&rpa Mx.</i>					
2090.	DAUBENTO*N/i4 Dec.	DAUBENTONIA.	<i>(3./Daubenton, celebrated naturalist) Leg.Pap.Lbt Gal.</i>					
18802	<i>punicea Dec.</i>	<i>red</i>	A O or	3	<i>jlau</i>	Vc	NewSpain 182a	S s.p Cav.ic.4.316
	<i>^Eschynbmene minitria Or.</i>							
18803	<i>longifolia Dec.</i>	<i>long-leaved</i>	A O or	3	<i>jn.au</i>	Y	NewSpain 1820.	S s.p Cav.ic.4.315
	<i>^Eschynbmene longifolia Cav.</i>	<i>Piscldia longifolia W.</i>						
2091.	CARAGA'NA Lam.	SIBERIAN PEA TREE,	<i>(Carachana, name in Tartary.) Leg.Pap.Lbt.Gal.</i>					
18804	<i>arborescens iMtn.</i>	<i>arborescent</i>	£ or	15	<i>ap.my</i>	Y	Siberia 1752.	S co Schm. ar. 1.33
	<i>Robinia Caragana L.</i>							
2	<i>inermis</i>	<i>unarmed</i>	A or	10	<i>ap.my</i>	Y	Siberia 1820.	S 81
18805	<i>arenaria Dow.</i>	<i>sand</i>	A or	1	<i>Jnjl</i>	Y	Siberia 1802.	Sk s.l Bot. mag. 188G
18806	<i>Altaligina l'oir.</i>	<i>Alugana</i>	A or	3	<i>apjn</i>	Y	Siberia 1789.	G 81 Her. st. 76
	<i>Roblin/a Altaligina Pall.</i>							
18807	<i>microphylla Dec.</i>	<i>small-leaved</i>	A or	3	<i>apjn</i>	Y	Russia 1819.	S 81 Pal. ros. 42.1.2
	<i>Robinia Altugana var. Pall.</i>							
18808	<i>Chamfigu Lam.</i>	<i>Chamlagu</i>	A or	4	<i>myjn</i>	Y	China 177a	G co Her. st. 77
18809	<i>frutescens Dec.</i>	<i>frutescent</i>	A or	5	<i>apmy</i>	Y	Siberia 1752.	L co Sw. fl. gar. 227
	<i>Robfnia frutescens L.</i>							
18810	<i>Redowski Fis.</i>	<i>Redowski's</i>	A or	3	<i>myjl</i>	Y	Siberia 1827.	S co Dec. leg. 11.45
18811	<i>mdllis Bes.</i>	<i>soft</i>	A or	2	<i>ap.my</i>	Y	Tauria 1818.	S 81
	<i>Robinia mollis Bicb. tomentosa Fis.</i>							
18812	<i>grndi flora Dec.</i>	<i>large-flowered</i>	A or	* <i>jl</i>	Y	Iberia 182a	G 81	
18813	<i>pygmae Dec.</i>	<i>pygmy</i>	A or	1	<i>ap.my</i>	Y	Siberia 1751.	Sks.p Schm. ar. 1. 37
18814	<i>spinosa Dec.</i>	<i>thorny</i>	A or	6	<i>ap.my</i>	Y	Siberia 1775.	L s.p Bot. reg. 1021
18815	<i>tragacantho'idesPair.</i>	<i>Goafs thorn-lkA</i>	A or	4	<i>ap.my</i>	Y	Siberia 1816.	G s.l Pal. p. 10.7
18816	<i>juKita Poir.</i>	<i>maned</i>	A or	1*	...	Y	Siberia 1796.	G 81 Bot. cab. 522
18817	<i>macrocantha Lo. C.</i>	<i>large-thorned</i>	A or	2	<i>myjn</i>	Y	Siberia?	... L 81
18818	<i>frox Lo. C.</i>	<i>fierce</i>	A or	2	<i>myjn</i>	Y	Siberia?	... L 81
2092.	HALIMODE'NDRON Fis.	SALT THEE.	<i>{Halimos, maritime, dendron, tree.) Leg.Pap.IM.Gal.</i>					
18819	<i>argentum Fis.</i>	<i>silvery</i>	A or	6	<i>myjn</i>	P	Siberia 1779.	R »l Bot.mag. 1016
	<i>Robinia Halodndron L.</i>	<i>Caragana argentica Lam.</i>						
S	<i>subvir&ccns Dec.</i>	<i>subvirescent</i>	A or	6	<i>myjn</i>	P	Siberia	.. R 81
	<i>Robinia triflora Herit.</i>							
3	<i>brachyseia Dec.</i>	<i>short standard</i>	A or	6	<i>myjn</i>	P	Siberia	.. R'e1

8093. DIPHY^SA Jac. DIPHYBA. (*Dis*, 2, *physa*, bladder •, Inflated append, to pods.) *Leg.Pap. Lbt.GaU* 1.
18820 carthagensis Jac. Carthagena i Q o r 10 ... Y Carthag. 1827. C 8f Jac.am. 180.51

9091 CALOTHACA Fa. CALOPOACCA. (*Kalos*, beautiful, *phake*, lentil.) *Legum. Pap. Lbt. Gal.* 1.
18821 wolgaria Fis. Wolga A or 2 myjn Y Siberia 1786. S 81 Pal. ros. 1.47
Cytinus nigreans Pall., pinnatus Pall., wolgaricus L.

8095. COLITTEA L. BLADDER SENNA. (*Kolu*, to amputate: iup. to die after.) *Leg.Pap.Lbt.Gal* 5.-7.
18822 ~~com.~~ arborescent* or 10 jn.au Y France 1568 S co Bot. mag. 81
18823 media W. en. intermediate A or 10 in.au O France 1568 S co Bot. mag. 81
18824 nepalensis B. M. Nepal or 5 aui Y Nepal 1822. S co Bot. mag. 2f22
18825 cruenta H. K. bloody II or 4 Jn.il S Levant 1710. L e o Schm.ar.119
18826 halopepla Dec. Aleppo II or 6 my.o Y Levant 1752. S co Schm.ar.120
VooSSuH.I.L

S096 SPHJEROPHRSA Dee. SPH.ER. (*Sphaira*, sphere, *phusa*, bladder; pods.) *Leg.* 1.82
18827 caapica Dec. Caspian k A or IJjl.au R Siberia 1818. fil

2097. SWAINSONA Sal. SWAINSONIA. (//WC Swainson, bot gard. Twickenham.) *Leg.Pap.Lbt.Gal.* 3.
18838 gallegifolia R. Br. Galega-leaved II Ljor 2' jlau R N. S. W. 1800. S ap Bot mag. 792
2 albiliflora R. Br. white-flowered II Ljor 2' jlau W N. S. W. 1826. S ap Botreg. 994
18829 coronillifolia 5a/. Coronilla-lyd II Ljor 2' jlau P N. S. W. 1802. S ap Bot mag. 1725
18830 lesserti^b/M Dec. LeaaertiaJvd II il.J or 2' jlau P N. Holl. 1824. S ap
atragahifolia Hort.

8096. LESSE'RTJA Dec. LEBBBRTIA. ~~Men Delessert~~, a French bot patron.) *I^g.Pap.J.bt.Gal.* 8.—18.
18831 annua Dec. annual IJ or 1 jnjl C. O. H. 1731. a1 Hook. ex. fl. 84
18832 diffusa R. Br. diffuse * IJ or 1 jl.au C. O. H. 1792. a1 Jac. ic. 3.576
18833 perennans D^c. lasting jf IAI C' 1 au C. G. H. 175a 8.1 Jac. vin. 3.3
18834 pdlchra A AT. pretty <- \ pr C. G. H. 1817. co Bnt. mag. 2064
18835 falciformis Dec. sickleformed <- i J or C. G. H. 1822. B1 Dec. leg. 6.46'
18836 brachytachya Dec. ahort-apikid <- i J or JLau C. G. H. 1821. 8.1
18837 annularia Bure. annular n. I J or JLau C. G. H. 1816. 8.1
18838 fruticosa B. R. shrubby & L J or jlau C. G. H. 1826. 8.1 Bot reg. 970

2099 SUTHERL A'NDI* H. K. SOTIERLANDIA. (J. Sutherland, author of bot. cat.) *Leg.Pap.Lbt.Gal.* 1.
18839 frut&cena H. K. fruitecent * J or 3 jnjl S C. G. H. 168a S s1 Bot. mag. 181
Cibatea frutscena L.

* 2100. PHA'CA X. BASTARD VETCH. (*Phago*, to eat. Adopted from Dioac) *Leg. Pap. Lbt. Astrag.* 10.—17.
18840 bse'tica L. Betic ^ A pr 4 jl W Spain 1640. k B1 M.h.2.8.1
18841 frigida L. frigid " A pr 1 jl Pa.Y Austria 1795. R al Jac.au. 166
18842 alpina Jac. alpine 5 A P' 2 jl Pa.Y Austria 17'9. R al Jac ic. 1.151
18843 glabra Oar. smooth 3c A or 1jl W S. France 1818. R B1
18844 membranacea Fig. membranaceous^t A or 1l. Pa.Y Siberia 1820. R al
alpina & dahurica Pis.
18845 australis L. southern ^ A P* imy.in Pa.P S. Europe 1779. R 8.1 Bot. cab. 490
18846 astragalina Dec. Astragalua-lik ^ A P' 1 jnjl W.B N.Europe 1771. R al Bot. cab. 429
18847 lappimica Wahl. Lapland 5 A or ijjl P N.Europe 1816. R al Fl. dan. 1.51
18848 oroboldea Dec. Orobus-like ^ A or 1' jl P Norway 1820. R al Fl. dan. 8.1396
18849 arenaria Pall. sand ^ A P' JLau P Siberia 1796. R al Pal. it3.ee. 1,2

* 2101. OXYTROPIS Dec. OXYTROPIS. (Oxys, pointed, tropis, keel; flowers.) *Leg.Pap.Lbt.Astr.* 2a —52.
I- ACAU'LBS. — Stems almost none ; stipules adhering to the petioles ; leaflets conjugate, not verticilate ; flowers

purple or white, from No. 18862. cream-coloreL
18850 montana Dee. mountain ^ A P' jl.au P Austria 1581. S al Bot mag. 483
Astragalus montanus L. Phaca montana Crx.
18851 cyknea Bieb. azureCaucasian& A °r B Caucasus 1818. a1
18852 uratensis Dec. Uralian 5 A P' P Siberia 1800. a1 Pal. as. 42
2 sericea Dec. silky 5 A or B Scotland aeaba. S al Eng. bot. 466
Astragalus urallensis E. B. not L.
18853 sordida Pen. mean ^ A pr i my.au Y.p Scotland .. S al Eng. bot 468
Astragalus uralensis L.
18854 ambigua Dec. ambiguous B , 1 myji P Siberia 1817. S al Pal. as. 55
18855 Lambertii Ph. Lambert's 1 aji P Missouri 1811. Bnt. mag. 2147
18856 grandiflora Dec. large-mf-fl. wd 1 aji R Siberia 1820. Bnt. mag. 2147
2 leptoptera Dec. narrow-winged B Siberia 1H18. Pal. as. 46
18857 songarica Dec. Songarian V Altai 1824. Bnt. mag. 2147
18858 longirostra Dec. long-beaked B Siberia 1820. Bnt. mag. 2147
Gmelinii Fia
18859 brevirrostra Dec. ahort-beaked B Siberia 1802. S co Pal.oa.45
altica Pers.
18860 uncta Dec. honkod B i or i jl.au W Aleppo 176B. S co
18861 leptophylla Dec. fine-leaved B i or i jl.au R Siberia 1818. S al Pal. as. 66
18862 campestris Dec. field B pr i jnjl P.v Germany 1778. S al W.&K.2.130
Astragalus campestris E-B/nötl. A or i jl.au Pa.Y Scotland moun. S 8.1 Eng. bot 2522
18863 fo'tida Dec. fetid B A pr i jUu Pa.Y Switzerl. 1819, S al Vil.del.a43
Astragalus Halleri All.

II. VEETICILL\ BE3. — Leaflets all or for the most part subverticilate, or two or three leaflets rising from the same base.
18864 microphylla Dec. small-tea wrd B A or i jlj.au Pa.Y Siberia 1819. S al Pal.aa.76
18865 verticillata Dec. v/hoxX-leaved B A or f jlj.au B.w Siberia 1819. S al Am.rut.19.1
18866 prostrata Dec. prostrate B A or f jlj.au B.w Siberia 1880. S al Am.Pal.aa.72
Astragalus dahuricus Pall.

18867 myriophylla Dec. myriad-leaved B A or i jl.au P.R.W Siberia 181& B al Pal. as. 71
18868 oxyphylla Dec. sharp-leaved B A or i jl.au P Siberia 1816. B al Pal. as. 74

HI. CAULESCENTES.—Stems elongated; stipules cauUne, not adhering to the petioles; leaflets conm/mie mat
verticillate ; flowers cream-colored, from No. 18870. purple.

18869 pilosa Dec. pilose i A P' I jnAM Pa.Y Siberia 1732. S al Rot cab 144
18870 Pallaati Pera Pallas's ^ A or ijl.au Pa. Y Siberia 1818. S al Pal as >i

18871	dcalbMa	Dec.	whited	A	pr	I	jkau	P	Caucasus	180a	S	s.1	Pal. as. 23.2,3
18878	gläbria	Dec.	smooth	A	or	£	jlau	P	Dnhuria	1823.	S	s.1	Dec.as 8
18873	dcflxra	Dec.	deflexed	A	pr	i	jn.jl	P	Siberia	18X.	S	s.1	Jacic. 1.153
18874	Fischéri*	Dec.	Fischer's	A	or	ijl.	au	Il	Altai	1817.	S	s.1	
18875	dich6ptera	Dec.	doubly winged	A	pr	4.	jn.jl	P	Siberia	1815.	S	s.1	PaLas.39

- 2102. *iiSTRA'GALUS L. MILK VETCH.* {Aster, star, gala, milk.) *LegunoPap.Ut.Astrdg.* 103. — 248.
I. *HYPOGLOTTI'DEI.* — Flowers purple; stipules free from the petioles, but between themselves joined at base; *thence* *so-called*; *red*.

18883 *liayon&isis* *Lot.* Bayonne ^ A. or  France 1816. D 8.1
 18884 *tauricus* *Pall.* Taunan ^ A or Tauria 1826. D 8.1 Pal. as. 38
 II DISSIMILATOR — *Stinules neither adhering to the petioles, nor joined together; flowers purple distant.*

II. DISSITIPLORI.—Siliques neither awning to the pedicels, nor joined together; flowers purple, distant;								
			legumes straight.					
18855 austriacus <i>L.</i>	Austrian		Pa.B	Austria	1640.	D	81	Jac au. 2.195
1K886 otopterus <i>Dec.</i>	ear-winged		Pa.B	Altai	1817.	D	s.1	
18887 sulcatus <i>L.</i>	furrowed		L.II	Siberia	1785.	D	co	Jac. vin. 3.40
18888 gracilis <i>Nut.</i>	slender		P	N. Amer.	1821.	D	81	
18889) subulatus <i>Bieb.</i>	subulate		P	Siberia	1820.	D	s.1	Pal. as. 20
18890 virgatus <i>Paf.</i>	twiggy		3 my.au	V	Siberia	1806.	C	co Pal. as. 18

III. ONOBRYCHOI'DEI. — *Stipules distinct from petiole, and not joined together; flowers purple or white, densely spicate or capitate; vexillum linear, elongated; roots perennial; legumes falcate, frotn No. 1880)2, straight.*
1880 leptophylla Desf. fine leaved *W. Barbary* *1811 D. H. Desf. 2207*

18891	leptophylla	Desf.	fine-leaved		1 jnjl	W	Barbary	1811.	D	Dest. at 2.207
18892	dahtircus	Pall.	Dahurian		1 fijnjl	P	Dahuria	1822.	D	s.l. Pal. it. 3. app.
18893	Imearifblius	Pers.	linear-leaved		1 flau	P	Siberia	1780.	C	Sw. fl. gar. 73
	teniufl. L.									
18894	Onobrychia	L.	Saintfoin			P	Austria	1640.	D	81 Jac. au. 1.38
18895	aduncus	Bub.	hooked			P	Caucasus	1819.	D	s.1 Biab. cen. 2.80
18896	vimineus	Pall.	twiggy			P	Siberia	1816.	D	s.1 Pal. as. 21
18897	fruticosus	Pall.	shrubby			V	Siberia	1804.	D	s.1 Pal. as. 19
18898	adsfirgens	W.	adsurgent	A A or	4 [njl]	p	Siberia	1820.	D	81 Pal. as. 31
	2 prostratus	Fis.	prostrate	3 A or	4 jnjl	p	Siberia	1818.	D	s.1
18899	Laxm&nni	Jac.	Laxmann's			p	Siberia	1804.	D	81 Jac. vin. 3.37
18900	caryocarpus	B. R.	nut-podded	iAor	1 jn.ju	p	Siberia	1811.	D	Bot reg. 176
	camdus	Nut			1 jn.jl	p	N. Amer.		D	s.1

IV. SESA'AIEI.—*Stipules not differing from the petioles, not joined together; flowers purple, densely spicate or capitate; legumes straight; roots annual.* 1759. 1. 1. Griseb. 2188.

18901	pentaglottis	<i>L.</i>	five-tongued	i	O or	<i>i</i>	jin,jl	P	Spain	1759.	■	s.1	Cav. ic. 2.188
18902	Glaux	<i>L.</i>	Miltwort	Jc	O or	4	jnjl	P	Spain	1596.	■	s.1	CLli. 2.241
18X3	oxyglottis	<i>Stev.</i>	sharp-tongued	Jc	O or	4	TM.	Tauria	1817.	■	8.1	
18904	cruciatus	<i>Lk.</i>	cruciate	Jc	O or	4	jinjl	■	...	1820.	■	8.1	
18905	Stella	<i>Gou.</i>	star-podded	Jc	O or	4	jlau	E.	S. Europe	1658.	■	co	Pluk.al.79.4
18906	/ribuloides	<i>Del.</i>	Tribulus-like	Jc	O or	4	jnjl	E.	Egypt	1817.	■	8.1	
18907	/sesamoides	<i>L.</i>	Sesame-like	Jc	O or	1	jujl	Pa.B	S. Europe	1616.	■	S.1	Gar. aix 12

V. VESICA^ARRI.—*Stipules not adhering to the petioles nor joined together; fls cream-colored; calyx bladdery.*
18908 *vesicarius* L. bladdery calyx Jc A or | jl.au W.Y Europe 1737. S s1 W.&K.1.4U

VI. ■■■—*Stipules neither adhering to the petioles nor together; flowers purple or white; legumes arched in roots annual.*

VII. BUCE' RATES.—*Stipules neither adhering together nor to the petiole,' flowers cream-colored legumes*

18916	contortu <i>pi</i> cotus	<i>L.</i>	twisted-plaited	<i>Jc</i>	O	or	1	jl.au	aftrn st arched j. roots annual	Pa.Y	Siligeria	1764.	S	co	Pal. as 79
18917	trime'stris	<i>L.</i>	three-monthly	<i>Jc</i>	O	or	*	jn.jl	Pa.Y	Egypt	1739.	S	co	Jac. vin. 2.174	
18918	hamosus	<i>L.</i>	hook-podded	<i>Jc</i>	O	■	■	jnl	Pa.Y	Spain	Iff.J3.	S	co	Lam. 1L 622.4	
	2 macrocarpus		large-fruited	<i>Jc</i>	O	or	4	jn.jl	Pa.Y	S. Europe	1820.	S	81		
18919	ffigceras	<i>W. en.</i>	goat's horn	<i>Jc</i>	O	or	■	jl.au	Pa.Y	1818.	S	co		
18920	baceras	<i>W. en.</i>	ox-homed	<i>Jc</i>	O	or	■	jnl	Pa.Y	1816.	S	co		
18921	epiglottis	<i>L.</i>	tongue upon tongue	<i>Jc</i>	Q	or	■	jnl	Pa.Y	S. Europe	17J7.	S	sl	Her. lug. 77	
18922	triangularis	<i>Dec.</i>	triangular	C	or	1	jl.au	Pa.Y	1818.	S	81			

VIII. SYNOCHIREAVL.—*Stipules free from the petiole, but joined together: flowers cream-colored.*

VIII. SYNOCHIEBEATI.							
Stipules free from the petiole, but joined together; flowers cream-coloured.							
18924 capitatus L.	headed	I	A ^o r	1	J ^j u ^u	Pa.Y	Leydant
18925 emarginatus Dec.	emarginate	A	O	U		1859. SS	s.l
18 ^a -26 mirrophysius L.	small-leaved				Pa.Y	S. Europe	1825. D
18 ^a -27 semibilocularis Dec.	semibilocular	H			Y	Siberia	s.l. Rauw. hod. 29
18528 r ^r pans W.	creeping	I		14	jn.au	1773. D	p.l
18 ^a -29 odoratus I.am.	sweet-scented			i	jnjl	1804. D	s.l. Dec. as. 23
18130 stipitatus D. Don	tor ^o -stipulated	A	o	1	in.jl	W	Mexico
						1818. D	s.l. W. h. b. 2,88
						1820. D	81 Dec. as. U
						1822. D	81 Bot. mag. 2380
					Pa.Y	Nepal	

IX. CICERO'DEI. — Stipules neither adhering together nor to the petiole; spikes pedunculate; flowers cream-colored; roots perennial; legumes sessile. — *Pl. 100, Fig. 203.*

Jc A or	3	jnjl	Y o	Britain	ch. wo.	D	sl	Eng. bot	203
Jc A or	1	jnjl	Pa.Y	Iberia	1818.	D	61		
Jc A or	1	jnjl	Pa.Y	Siberia	1824.	D	sl		
J: A or	\ myjn	Pa.Y	Europe		1772.	D	8.1	Bot. cab.	680
Jc A or	\ my.au	W.Y			1776.	D	sl	Bot. cab.	III

18936 Cicer L.	Vetch	Jc A or	2 jnjl	Y	Europe	1570.	D s.1	Jac. au. 251'
18937 uliginous L.	marsh	^ A or	2 jn.au	Pa.Y	Siberia	175*	D p.1	PaL as. 26'
18938 micranthus Desu.	small-flowered	^ A or	1 jnjl	Pa.Y	...	1800.	D s.1	
18939 carolinianus L.	Carolina	^ A or	1 jl.au	G.v.	N. Amer.	1732.	D s.1	Di. el. 39,45
18940 canadensis L.	Canadian	^ A or	1 jl.au	Pa.Y	N. Amer.	1732.	D s.1	Dod. m. 64
18941 falcatus L.	sickle-potifcd	5 A or	3 \$ jl	G.T	Siberia	...	D p.1	Dec. as. 26
vir&cens H. K.								
18942 falciformis-Dec.	sickle-formed	■ A or	yjinjl	Pa.Y	Algiers	18ia	D s.1	Desf. at. 2,206
18943 asper Jac.	rough	■ A or	3 jl.au	Pa.Y	Astracan	1796.	D p.1	Jac. ic 152
18944 palaescens Bleb.	palescent	■ A or	1 jnjl	Pa.Y	Siberia	1818.	D 8.1	

X. GALEGIF6RMIE8. — *Stipules neither adhering together nor to the petioles; flowers cream-colored or white;*

18945 chinensis L.	Chinese	^ ^ or	1 jnjl	Pa.Y	China	1795.	S p.1	L.f.dec.3
18946 galegiformis L.	Galega-formed	^ A or	2 in.au	Y.o	Siberia	1729.	D si	PaL as. 29

XI. ALOPECUBOI'DEI.—*Stipules not adhering to the petiole; stems erect; flowers yellow, crowded into dense*

18947 alopecia B PaU.	spikes at the axils of the leaves, either sessile or on short peduncles.							
	sea-fox			Y	Siberia	1800.	S 1	Pal. as. 9
18948 maximus L.	greatest Foxtail	■ ■ ■	3 jl.au	Y	America	...	S 6.1	
18949 filop-jeuroides L.	Foxtail-like	■ ■ ■	3 jlau	L.Y	Spam	1737.	S 1	Pal. as. 8
18950 narbonensis Gou.	Narbonne	■ ■ ■	2 jnjl	Pa.Y	S. Europe	1789.	S 6.1	PaL as. 10
18951 panticus W.	Pontic	■ ■ ■	3 jnjl	Pa.Y	Tauria	1820.	S 1	Pal. as. 11
18952 vulpinus W.	fox	■ ■ ■	2 jlau	L.Y	Caucasus	18151	S 8.1	Pal. as. 7

XII. CHRISTIA^NI.—*Stipules neither adhering together nor to the petiole; stems erect; flowers cream-colored, in subsessile glomerate racemes at the axils of the leaves.*

18953 christianus L.	Christian	^ A or	3 jl	Pa.Y	Armenia	1737.	D s.1	To. it 2. 254
18954 tomentosus Zatn.	woolly-leaved	^ A Jor	3 jl	Pa. Y	Egypt	1800.	S p.1	Dec. as. 29

XIII. TRAQACA^NTIjs.—*Petioles persistent, tiprescent; stipules adhering to the petioles.*

18955 caucasicus Pall.	Caucasian	tt. or	i jn.jl	W	Caucasus	IH24	S 8.1	Den. br. 84
18956 Tragacantha L.	gear-Goat's-Thorn	L	or 1 my.jl	Pa.Y	S. Europe	1840.	S 8.1	Par. th.2
18957 Poterium sm.G.T.n.	Poterium	tt. or	i jn.jl	P	Pyrenees	1791.	S 8.1	Gar. aix 104
18958 anstatis Herit.	awned	tt. or	1 my.jl	Pa.Y	Egypt	1816.	S 1	Rus. al. 5
18959 t&midub W.	tumid	tt. or	§ jn.jl					

XIV. ANTHYLLOI'DEI.—*Stipules adhering to the petioles; petioles unarmed; calyx bladdery, including the legumes.*

18960 ammodytes Poll.	sand viper	^ L or	\ my.jl	W	Siberia	1820.	S s.1	Pal. as. 5
18961 /folicacibus Lam.	kettlecalyxed	^ A or	i jn	Pa.Y	Armenia	1806.	D s.1	Schreb. dec. 3

Phaca vesicaria Schreb.

XV. CAPRINI.—*Stipules at base adhering to the petioles; petioles unarmed; flowers yellow; calyx not bladdery.*

18962 caprinus L.	goa.Uscented	■ ■ ■	1' jnjl	Pa.Y	Barbary	1683.	D s.1	M.h.2.24.3
18963 longiflorus Pall.	long-flowered	■ ■ ■	1 jnjl	Y	Tartary	1806.	D s.1	Pal. as. 80
18964 uttinger Pall.	bladder-bearing	■ ■ ■	* jnjl		Russia	1818.	D 8.1	Pal. as. 61
18965 lanigerus Desf.	wool-bearing	■ ■ ■	I my. jl		Egypt	1791.	D 8.1	Desf. at. 202

XVI. INCA VII.—*Slip, adhering to unarmed petioles; calyxes and legumes not bladdery; leg. perfectly bilocular.*

18969 monspessulanus B L.	MontpeUer	JU A or	1 jl.au	P	France	1710.	D s.1	Botmag. 375
2 Sibts	white	t A or	1 jnjl	W	S. Europe	...	S 8.1	

18970 incknus L.

XVII. RkmcJbm.—*Stipules adhering to the unarmed petioles; calyx not bladdery; ruptured at top after*

XVIII. DU^BII.—*Doubtful to which of the sections they belong.*

18975 neglectus Fis.	neglected	Jt A or	1 jnjl	...	Siberia	1826.	D s.1	
18976 acutifolius Sehl.	acute-leaved	5 A or	i jnjl	...	Switzerl.	1826.	D s.1	Pal. as. 68
18977 mehlotoldes Pott.	Melilot-like	J A pr	3 jnjl	P	Siberia	1785.	D co	Pal. as. 41
18978 arenarius L.	sand	■ A or	1 jnjl	B	Germany	1798.	D s.1	Rets. a 3

2103. GULDENSTO'DTM Fis. GuLDENsr.JLDIA. (J.A.Guldenstaedt, nat.) Leg. Pap. Lbt. Astr. 1.—2.

18979 paucialba Fis.	few.flowered	^ A or	* jnjl	R	Siberia	1827.	S s.1	Pal. as. 66
flstr&galus pauciflorus Pall.								

2104. BISE'RRUALA L. HATCHET VETCH. {Bis, twice, serrula, little saw; pods.) Leg. Pap. Lbt. Astr. 1.

18980 Pelecinus L.	Corn-weed	O pr	1 jl.au	P	S. Europe	1640.	S s.p	Lam. ll. 622
18981 lffigigita Sm.	smooth-fvatt	-* O or		Y	Archipel.	1818.	S co	Fl.gr. 718

18982 muricata L.

XVIII. LU^TEJE.—*Flowers yellow.*

18983 sulcata L.	prickly	Jr o pr	2' jnjl	Y	S. Europe	1640.	S co	M.h.2.11.4
18984 subvillbs L.	*furrowed	* O pr	2 jnjl	Y	S. Europe	1596.	S 8.1	Desf. at. 2.4
18985 acutifilia Via.	subvillous	Jc o i>	2 jnjl	Y	S. Europe	1731.	S co	M.h.2.11.2
18986 vermiculata L.	acute-leaved	Jf O or	§ jnjl	Y	Corsica	1825.	S co	Viv.lib.19.4
	worm-podded	Jc Q pr	2 jnjl	Y	& Europe	1621.	S co	M. h.2.11.3

* 2106. CORONILLA L. CORONILLA. (Corona, a crown; disposition of fls.) Leg. Pap. Hed. Cor. 16. — 20.

I. LU^TEJE.—*Flowers yellow.*

18987 E'merus L.	Scorpion Senna	tt or	3 apjn	R.Y	France	1596.	L co	Bot mag. 445
18988 jdncea L.	Rush	■ or	3 jnjl	Y	France	1656.	C r.m	Bot. cab. 235
18989 valentia L.	Valentia	■ or	2 rar.n	Y	S. Europe	1596.	C r.m	Bot mag. 185

stipularis Lam.

18990 pentaphyllo Desf.

five-leaved

18991 argentea L.

silvery-leavrd

18992 glatica L.

glaucous seven-lvd

18993 viminalis Sal.

twiggy

18994 minima L.

least

18995 coronata L.

crown-headed

18996 montana Sco.

mountain

coronata Bieb, not U or W, or

X 2

II. TRIPOLIA^TJB.—Leaves trifoliate, terminal ones petioled.

- 19044 *fegopodtes* Dec. Hare's-foot-lik *tilpr* *Hjn.il* P China 1790. S 81 Bur. in. 53.2
tfedfsarum lagopodioides L.
- 19045 *lagocphala* Dec. hare-headed *£ iAI un* 2 *jl* Li Brazil 1824. S co
Hed^aarrum lagocephalum Lk.
- 19046 *lagbius* Dec. Hare's-foot *ftlor* 7 *jnjl* P Nepal 1824. S s.l
J/cdysarum arbbreum Ham.

2119. NICOLSO'NIJI Dec. NICOLSONIA. (Af. Nicolson, wrote on Nat ILSt.Dom.) *Leg.Pap.Hed.Cor.Euh.I.S.*
 19047 *barbate* Z*c. bearded *< . D a* 1 *jnjl* P Jamaica 1818. S 81
Ifcdysarum barb&tum L.

•2120. DESMOMMUM Dec. DEBMODIUM. (*Desmos*, band; stam. joined?) *Leg. Pap.Hed.Cor.Euh. 60.—140.*

- L EuDESMBDiM—*Joints of legume coriaceous, indehiscent, subettipic; and at each extremity truncates leaves*
 ««» ^ m* *Pinnately trifoliate; peduncles axillary; flowers corybosely umbellate.*
 19048 *umbellatum* Dec. *uinbel* flowered * • un 3 ... W E. Indies 1801. C l.p Jac. sc. 3.297
2/edysarum umbellatum L.

II. PLEUKA^OBBIUM.—*Joints of legume membranaceous, nearly quadrate, lower side a little convex, afterwards dehiscent; leaves unifoliate; petioles winged; wings at top, on each side, ending in a tooth :from 19053. leaves innately trifoliate; petiole's not winged?*

- 19049 *auriculatum* Dec. eared *ITBlor* 5 *jt.au* P Timor 1819. ■ s.1
 19050 *triquetrum* Dec. triangular-rffc *O or* 1 *Jlau* P E. Indies 1802. ■ Lp Bur. in. 52.2
Ifcdysarum triquetrum L.
- 19051 *psclido-triquetrum* Dec. bast.Triquet.tf OS or 4 *jl.au* ■ Nepal 1820. ■ s.1
 19052 *alatum* Dec. winged *£ DD or* 4 *jl.au* P E. Indies 1817. ■ s.1
 19053 *gyrans* Dec. whirling-plan. *FPlcu* 3 *Jlau* P E. Indies 1775. ■ p.1 Jac. ic 3.566
1/edysarum gyrans L.
- 19054 *gyroides* Dec. Gyrans-like *tf ITBlor* ■ . . . E. Indies 1817. S s.1

III. CHALA^RIUM.—*Joints of legume membranaceous, indehiscent, oval or orbicular, rarely oblong, with both sides convex, and at each extremity attenuated; racemes loose, elongated; pedicels three from each bract; leaves simple, from No. 19062. trifoliate with the terminal one a little larger.*

- 19055 *gangeticum* Dec. Gangetic *Ejum* *Wfijam* Pa.V E. Indies 1762. S r.m Bur. scy. 49.2
//edysarum gangeticum L.
- 19056 *maculatum* Dec. spotted *Oun* 1 *jl.au* ■ India 1732. S l.p Di. el. 141.168
Ifed^sarum maculatum L.
- 19057 *reniforme* Dec. kidney-shaped *■ or* 3 *jn.au* ■ E. Indies 1820. S 8.1 Bu. in. 52.1
 19058 *sagittatum* Dec. arrow-leaved *i ■ Jun* 3 ■ E. Indies 1807. S IP
//ed^sarum sagittatum Pair.
- 19059 *angustifbhum* Dec. narrow-leaved *■ ■* 2 *jniau* P Mexico 1824. ■ s.1
 19060 *latifbhum* Dec. broad-leaved *i ■ ■* 2 *au* P China 1818. ■ Lp Bdtreg.355
//edysarum latifolium Box
- 19061 *lasiocarpum* Dec. hairy-podded *i ■ Qor* 4 *jnau* R Guinea 1823. S s.1 Beau. ow. 1.18
 19062 *canadlinse* Dec. Canadian *■ A un* 6 *Jlau* P N. Amer. 1640. D s.1 Cor. ca. 45
/fed^sarum canadense L.
- 19063 *canescens* Dec. canescent *■ A un* 2 *jl.au* W.P N. Amer. 1733. D s.1 Pluk. al. 308.5
Hedysarum canescens L.
- 19064 *manlandicum* Dec. Maryland *■ A* ■ ■ T N. Amer. 1725. D s.1 DL el. 144.171
fledyiarum manlandicum L.
- 19065 *obtusum* Dec. obtuse *■ ■* 2 *jl.au* ■ N. Amer. 1805. D s.1
//ed^sarum obtusum MM.
- 19066 *ciiKfe* Dec. ciliated *■ A or* 2 *jl.au* V N. Amer. 1823. D 8.1
 19067 *viridiflrum* Dec. green-flowered *■ A un* 3 *Jl.8* P N. Amer. 1787. D 8.1 Pluk. al. 308.2
ifedysarum viridiflrum L.
- 19068 *bracteatum* Dec. large-bracted *■ A or* 8 P N. Amer. 1818. D s.1
 19069 *cuspitatum* Dec. sharp-pointed *■ A un* *lift* *Jlau* V N. Amer. 1806. D s.1
//ed^sarum cuapidatum W.
- 19070 *glabrum* Dec. smoothish *■ A or* P N. Amer. ... D s.1
 19071 *paniculatum* W. panicled *■ Al>r* P N. Amer. 1781. D s.1 Pluk. al. 432.fi
/iedysarum paniculatum L.
- 19072 *acuminatum* Dec. acuminate *■ A un* *lift* *jl.au* P N. Amer. 1805. D a.1
^ed^sarum glutin6sum Mx.
- 19073 *nudiflrum* Dec. naked-flowered T N. Amer. 1823. D s.1
- 19074 *pauciflrum* Dec. few-flowered T N. Amer. 1818. D s.1
- 19075 *rotundifolium* Dec. round-leaved N. Amer. 1733. D 8.1
canescens IV.
- 19076 *stipulaceum* Dec. stipulaceous T Mexico 1824. Fl. mex. ic. in*
- 19077 *y4partnes* Dec. Bedstraw Mexico 1823.
- //ed^aarum i4par'nes Lk.
- 19078 *cinerium* Dec. grey T Mexico 1820. C 8.1
- 19079 *dubium* B. B. doubtful P Mexico 1824. C a.1 Bot Teg. 967
- 19080 *cq/anifbium* Dec. Cajan-leaved B Nepal 1824. C a.1 H. & B.n. (J.598
- 19081 *incanum* Dec. hoary B Caracas 1820. C 8.1 P.ecLB. 149.1
- 19082 *uptatum* Dec. supine B Jamaica 1818. C s.1 SL.jam. 1.118.2
- 19083 *tortuosum* Dec. twisted-podded W. Indies 1816. ■ s.1 SL.jam. 1.116.2
- //edysarum tortuosum Sun.
- 19084 *ascendens* Dec. ascending Jamaica 1818. C a.1
- i/ed^sarum ascendens Sun.
- 2 caruleum Undl. blue W. Indies 1818. C a.1 Bob rag. 815
- 19085 *trigonum* Dec. three-angled Jamaica 1733. D 8.1
- //cd^saruin trigonum Sun.
- 19086 *spirale* Dec. spiral Jamaica 1700. C 8.1
- 19087 *tenclium* Dec. delicate Caraccas 1818. S s.1
- 19088 *aspercum* Desv. rough — 18ia C 8.1
- 19069 *diversifolium* Dec. divers-leaved E. Indies 1816. C 8.1
- 19090 *heterophyllum* Dec. various-leaved E. Indies 182a ■ si Bur. xey. 54.1
- 19091 *triflrum* Dec. three-flowered India 1796. ■ 8.1 Bur. in. 54.2
- Hedysarum triflorum L.
- 19092 *diffusum* Dec. diffuse * *Qun* 2 *jl.au* P E. Indies 1818. ■ 8.1
- 19093 *tomentosum* Dec. woolly £ *LAJun* *li.in.jl* P China 1782. ■
- Hedysarum tomentosum Thun.
- 19094 *multiflrum* Dec. many-flowered JIFHun *lijl.au* P Nepal 1823. C 61
- *Hedysarum multiflrum D. Don

II. HVMENOBBY^CHIS.—*Legumes falcate or suborbiculate, rugose or somewhat prickly in the disk, with the whole back expanded into a membranaceous crest; scarcely subdentate.*

- 19156 Pallasia^Bieb. Pallas's ^ A or 1 ju.au Pa.Y Iberia 1890. ■ 8.1 Bieb. cen. 1.3?
 19157 radiate *Bieb.* radiated ^ A or U jn.au Pa. Y Iberia 1818. ■ 8.1 An. mu 12.13
 19158 picta Desv. painted \$ A or 1 jn.au Pa.R Levant 1820. ■ 8.1 Bux. c. 2.42
Michatixii Dec.
 19159 ptolemaica Dec. Ptolemais ■ A or 1 jn.au Y Egypt 1816. S 8.1 Del. teg. 39.1

III. DBNDROBRYV:HIS.—*Legumes smooth, unarmed, crestless; stems pubescent.*
 19160 corntoa Desv. homed & Jor 1 jn.au R Caucasus 1816. S s.l To. it 2.249. ic.
 orientals HU.

2125. ELEICTTIS Dec. ELEIOTIS. (*Heltios, dormouse, ou, ear; leaves.*) Leg. Pap. Hed. Cor. Euh. I.
 19161 sorbria Dec. sister whste-rooted £0 un 1 ... R E. Indies 1817. S 8.1 Desv. 3. fa. 31
 JSTed^sorum sororium L. HilUa sorbria W. Onobrychis sorbria Desv.

2196. LESPEDE^Zil Mx. LESPEDEZA. (*Uspedez, gov. of Florida, bot pat.*) Leg. Pap. Hed. Cor. Euh. 15.
 19162 Juncea Pers. Rush-branched ■. Opr 8 iUu W India 1776? D Lp I* f. dec. 1.4
 /fedysarum j&ncema L.

- 19163 reticulata Pers. netted ■ □ □ ■ jn.jl P N. Amer. 1816. D Lp
 19161 sessiliflora Nut. sessile-flowered ■ □ □ ■ jl.au P N. Amer. ... D Lp
 19165 eriocarpa Dec. woolly-fruited ■ □ □ ■ jl.au V Nepal 1819. D Lp
Anthylhs cuneata Ham.
 19166 Sttvej Nut. Stuve's ■ □ □ ■ UJLau P N. Amer. 1821. D Lp
 19167 frutescens Pers. frutescent ■ □ □ 4 jl.au P Virginia 1739. C Lp Jac.vin.3.89
 19168 capitata Mx. headed ■ □ □ 2 jn.jl Str N. Amer. 1789. I Lp
 19169 angustifolia EU. narrow-leaved ■ □ □ 2 jn.jl W N. Amer. 1800. D Lp
 19170 polystachya Mx. many-spiked ■ □ □ 3 jn.au W N. Amer. 1783. D Lp Mic am. 2.40
 19171 hirta Horn. hairy ■ □ □ 2 jl.au W N. Amer. 1819. D Lp
villte Pers.
 19172 violacea Pers. violet-flowered ■ □ □ 2 jl.au V N. Amer. 1789. D s.l
 19173 divergens Ph. diverging ■ □ □ 2 jl.au V N. Amer. 1800. D Lp
 19174 procumbent Mx. procumbent ■ □ □ I jn.au P N. Amer. 1816. D Lp
 19175 prostrate Ph. prostrate ■ □ □ i jn.au P N. Amer. 1810. D Lp
 19176 glomerata Horn. glomerate ■ □ □ 3 jl.au P E. Indies 1819. D Lp

2127. FLEMING/jf Rox. FLEMINGU. (*John Fleming, F.R.S. F.L.S. &c.*) Leg. Pap. Hed. Cor. Euh. 9.-10.
 I. FLEMINOIA'STRUM.—*Flowers naked or stipulate, with small deciduous bracteas; leaves trifoliate; leaJU's sessile.*

- 19177 stricta Rox. strict £ EJ un 2 jl.s P India 1798. S s.p Rox. cor. 3.248
 19178 semialta Rox. winged tt-Qun 3 jLau P Nepal 1805. ■ Rox. cur. 3.249
 19179 congesta Rox. crowded-spiked&Dun 3 jLs P India 1802. ■
 19180 prostrata Rox. prostrate JU ES or 1 jl.au P E. Indies 1816.
 19181 procumbens Rox. procumbent JL EJ or 1 jl.au P E. Indies 181a
 19182 nana Rox. dwarf ft-Qun 11 au P India 1804. ■
 19183 lineal Rox. lined jt DD un 2 jl.au P India 1793. ■ Bur. in. 53.1

II. OffravbDITM.—*Flowers furnished with a large, concave, leafy, persistent bractea; leaves simple, penninerved.*
 19184 strobilifera H. K. strobile-bearing CD un 3 jLau P E. Indies 1787. S p.l BoLreg.617
 /fedysarum strobilifertim L.

- 19185 guineensis G. Don Guinea ■ Q or 10 ... P Guinea 1825. S Lp

2128. ALHA^GI Tou. MANNA. (*Arabic name of the plant*) Leg. Pap. Hed. Cor. Euh. 2.-3.

- 19186 maurinum Tou. Moors' flL Jm 2 jLau R Egypt 1714. S si Rauw.it94ic
 ied^sorum Alhhgi L. Manna hebraica D. Don

19187 camel&cam Fis. camels' * ^ | m 1J jl.au R Siberia 1816. S s.l

ifedyiarum Yaehdo-Alh&gi Bieb. Manna cispica D. Don

■ 3129. ALYSIC A'RPUS Neck. ALTBICARFUS. (*Atysis, chain, karpot, fruit.*) Leg. Pap. Hed. Cor. Euh. 5.-7.

19188 Apulcurifulus Neck. Hare's-ear-lvd tf E3 or 1 jn.au P EL Indies 1793. S s.l Bot. mag. 172?

Hedysarum ftpleurifoliura L. gravaineTO. Retz.

19189 nummularifbhus IV. Moneywort-lvdjf ZS un 1 jl.s P India 1777. S Lp Pet g. 26.3

JYedjgarum nummularifolium L.

19190 vaginiklis W. sheathed O un fjl.au R E. Indies 1790. S Lp Bur. ley. 49.1

Hedysarum vaginale L.

19191 monilifer Dec. bracelet-bearer £ ED or g ^ au p ^ i,jic8 1816. S s.l Bur. in. 52. 2

19192 Styracifilius Dec. Styrax-leafed fl-Qun 2 ... P E. Indies 1796. C Lp

A ? ■ ■ ■ *styracifilium L.*

2130. BREMONTIETRJ Dec. BRBMONTIERA. (if. *Bremontier*, an agriculturists Leg. Pap. Hed. Cor. Euh. L
 19193 Ammoxylon Dec. Sand-wood • or 4 ... P Mauritius 1826. C s.l Bux. zey. 82

2131. CPCER Tou. CHICK-PEA. {Kikys, force qualities.) Leg. Pap. lie. 1.
 19194 arietnum L. ram's-Atarf O dt 1 jLau P & Europe 1548. S co Bot.mag.2J74

2132. FA^BA Tou. BBAN. (Phago, to eat; esculent) Leg. Pap. Vic 1.
 19195 vulgaris Moen. common O cul 3 Jn.jl W Egypt ... S co Black. 19

Hcia Kiba L. horse O ag 3 Jn.jl P S co

2133. n XI A Tou. VBTCB. (Vtocio, to bind together; twining tendrils.) Leg. Pap. Fie 72. —108.

I. PBDUNCULA^f.—*Flowers pedunculated.*
 Pea-shaped - A or 2 jl.au Pa. Y Austria 1739. R co Jac. au. 4.364
 Pleasing red 4 A or 2 jn.jl Pi Siberia 1818. R co Gm.sL4.3
 Carolina ttf/terf 3 A or 2 jn.jl W Carolina 1820. R co

hedge toothed + A or 3 my.jn P France 1752. R co Spr. hal 7

toothed A or 4 jn.jl R Siberia 1819. S co

American iwood A or 6 jLau W. B. Britain moi. w. R co

common wood A or 6 jLau W. B. Britain moi. w. R co

variegated A or 3 jn.jl Pa.P Caucasus 1816. S co Eng.bot.79

Cassubian A or 3 jn.jl L.B. Germany 1711. R co An. mu. 12.12

short-peduncled A or 2 jn.jl Pa.B Caucasus 1818. S co Jac. au. 3.229

Crarra tufted A or 2 jn.au V Britain hed. R co

white-flowercd 1 ^ or 2 jn.jl W Bntain gard. s co Eng. bot. 1168

S fidrlbus rhbrls	red-flowered	ft	A or	I	R	Britain	gard.	S	co	
19207 Gerfrdi Jac	Gerard'B	T	or	I	V	S. Europe	1810.	S	co	
19208 Pseudo-Crmwi Bert.	BastardCraccaJ	I	I	I	P	S. Europe	1880.	S	co	
tenuifolia Ten.									Jac. au. 229	
19209 polyphylla Desf.	many-leaved	I	I	I	Pa.P	Algiers	1816.	S	co	
19210 argentea Lap.	silvery	SL	I	I	Pk	Pyrenees	1827.	S	co	
19211 tenuifolia Roth	slender-leaved	I	I	I	V	Germany	1799.	R	co	
19212 consentina Spr.	Consentian	I	I	I	B	Italy	1818.	S	co	
19213 capensis Pert.	Cape	I	I	I	P	C. G. H.	1802.	R	co	
19214 canescens £.o6.	canescent	I	I	I	B	Libanus	1800.	S	co	
19215 ochrolefcca Ten.	yellowish white.	I	I	I	Pa.Y	Italy	1825.	S	co	
19216 onobrychides L.	Saintfoin-like	I	I	I	P	S. Europe	1789.	S	co	
19217 atropurpurea Desf.	dark-purple	I	I	I	P	Algiers	1815.	R	co	
19218 benghalensis L.	Bengal	I	I	I	D.P	E. Indies	1792.	S	co	
19219 perennis Dec.	perennial	I	I	I	P	a Europe	1820.	S	co	
19220 tongifolia Poir.	long-leaved	I	I	I	Pa.Y	Syria	1818.	S	co	
19221 villbsa FT.	yillous	I	I	I	D.P	Germany	1815.	S	co	
19222 disperma Dec.	two-seeded	I	I	I	W	S. France	1820.	S	co	
parviflora Loi.										
19223 altissima Desf.	tallest	I	I	I	P.B	Barbary	1820.	S	co	
19224 biennis X.	biennial	I	I	I	D.P	Siberia	1753.	S	co	
19225 Nissoltna L.	Nissole's	I	I	I		Levant	177a	S	Gm. si. 4.2 Bot reg. 871	
atropurpurea Lindt.										
19226 pellucida Jac.	pellucid	I	I	I	1 jn.au	P	C. G. H.	1812.	R	co
19227 biflora Desf.	two-flowered	I	I	I	11 jn.au	B	Algiers	1801.	S	co
19228 calcarata Desf.	spurred	I	I	I	2 jn.au	KB	Barbary	1790.	S	co
monanthica Reit.										
19229 syrlaca Httn.	Syrian	J, O or	2 jn.jl	V	Syria		1816.	S	co	

II. SESSILIFLURA.—*Flowers subsessile.*

19230 safiva Z.	common cultivated	± O	I	3 my.jn	P	Britain	com fl.	S	■ Eng. bot 334
Ssegetalis	corn	J O	I	S my.in	P	Britain	corn fl.	S	h.l
dhemorialis	grove	I O	I	3 my.jn	P	Britain	corn fl.	S	hi
19231 angUDtifulia Sibth.	narrow-leaved	" J O	I	1 my.jn	It	Britain	corn fl.	S	co
luganensis Schl.									
19232 leucospermum Moen.	white-seeded	Jfc O	I	2 jn.au	P	Europe	1810.	S	co
dilba Moen.									
19233 glabra Schl.	smooth	I	I	2 Jn.au	P	Switzcr.	1819.	S	co
19234 intermedia Viv.	Intermediate	I	I	3 jn.jl	P	S. Europe	1818.	S	co
19235 inrisa Bleb.	cut	I	I	2 jn.au	P	Caucasus	1820.	S	co
19236 globosa W.	globose	I	I	1 jn.au	B	1804.	S	co
19237 cordata Wul.	cordate	I	I	2 jn.au	P	Germany	1816.	S	co
19238 pilosa Bleb.	pilose	I	I	2 jn.au	P	Tauria	1818.	S	co
19239 megalospermum Bib.	great-seeded	I	I	2 jl.au	P	Tauria	1798.	S	co
19240 peregrina L.	foreign	I	I	1} jl.au	P	France	1779.	S	co
19241 Michauxia Spr.	Michaux's	I	I	1} jl.jl	W	1803.	S	co	Pluk. al. 233.6
19242 Thou W. Mart.	Thouin's	I	I	2 jn.au	P	Europe	1800.	S	co
19243 ba/tica Fis.	Baetic	I	I	2 jn.au	P	Europe	1820.	S	co
19244 Musquincx Bosc	Musquincz	I	I	1 jn.au	P	Europe	1818.	S	co
19245 amphicarpos Dor/.	both sides podded	I	I	1 my.jn	P	France	1815.	S	co
19246 pyrenaica Pou.	Pyrenean	I	I	1 my	Y	Pyrenees	1818.	S	co
19247 strata Bieb.	striated	I*	I	1* jn.au	P	Tauria	1823.	S	co
19248 /athyn Ades L.	Lathyrus-like	O or	I	1* ap.jn	P	Britain	fhU.fl.	S	h.l
-Ernum soloniense L.									Eng. bot 30
19249 Uvigata Sm.	smooth-podded	I	I	1 jn.au	Pa.Y	England	sea. sh.	S	co
19250 dubia Schulz	doubtful	I	I	2 jn.au	—	—	1820.	S	co
19251 lutea £.	yellow	I	I	1) jn.au	Y	Britain	scash.	S	co
19252 hirta Balb.	hairy	I	I	2 jn.au	W	Europe	1816.	S	co
19253 hibrida L.	hybrid	I	I	1) jnau	Y	England	thick.	S	co
19254 hirsute Fis.	hairy	I	I	1) jn.au	Y	Siberia	1818.	S	co
19255 grandi flora Sco.	great-flowering	I	I	1 jn.au	Y	S. Europe	1818.	S	co
19256 sordida Bieb.	sordid	I	I	1 jn.jl	Y	Hungary	1802.	S	co
Biebersteinia Bcs.	Bieberstein's	I	I	1 jn.au	Y	Podolia	1820.	S	co
sordida Bieb.									
19258 tricolor Seb.	three-colored	I	I	If jnau	P v	Italy	1818.	S	co
19259 bicolor IV. en.	two-colored	I	I	2 jn.au	P.w	182a	S	ro
19260 trifida Ten.	three-flowered	I	I	2 jn.au	P	Italy	1820.	S	co
19261 pannitica Jac.	Pannonian	I	I	1 jn.au	W	Hungary	J658.	S	ro
19268 spium L.	hedge	I	I	2 my.jn	V	Britain	hed.	S	h.l
19269 truncifolia Fit.	small truncate	I	I	1 jn.au	T.a.Y	Siberia	1818.	S	co
19270 narbonensis L.	Narbonne	I	I	3 jn.jl	P	France	1596.	S	co
19271 serratifolia Jac.	saw-leaved	I	I	3 jn.jl	P	Hungary	1723.	S	h.l
19272 platycarpus Roth	broad-podded	I	I	Ujl.jl	P	Germany	1723.	S	co
monadelpha Roth	monadelphous	I	I	3 jn.au	P	N. Amer.	1820.	S	co

2134. ERVUM L. TARE. (*Erw, tilled land, Celt.; pest ID.*) Leg. Pap. Vic. 10.—18.L LBNS.—*Legumes subinflated, broad, oblong, two-seeded.*

19268 /ens L.	Lentil	O	clt	I	my	Pa.B	France	1548.	■ co	Riv.tet.35
19269 nigricans Bieb.	blackish	I	O un	I	jn.jl	P	Tauria	1817.	■ co	
fctiblides Ten.										
19270 Lenticula Schreb.	Small Lentil	X O	un	1	jn.jl	B	Carinthia	182a	S	co
sotomencs WIU.										
19271 trifolia Ten.	Clertrrvoldes Brig.	Cler	trrvoldes	I	jn.jl					
19272 hirsutum L.	hairy	I O	un	2	jn.jl	W	Britain	corn fi.	S	h.l
19273 dispermum Bat.	two-seeded	± Q	un	I	li jn.jl	P	E. Indies	1802.	S	co

II. ERV'LJL.—*Legumes subinflated, oblong* linear, four or six seeded.*

19273 J&ryflia L.	Lentil	I	O clt	1	jn.jl	■	S. Europe	1596.	S	Black. 208.3
J4cia ^ryflia W. ETV\\SL saflva IJk.										
19274 monanthos L.	one-flowered	ft	O	I	1 jn.jl	P	S. Europe	1798.	S	Stur.ger.1.32 ic
ficia articulata W.										
19275 tetraspernum L.	four-seeded	J O	un	I	li jn.jl	W.B	Britain	corn fi.	S	Eng. bot. 1223
19276 grjcilic Dec.	slender	ft	I O	un	I jn.jl	Pa.B	S. Europe	1822.	S	Brot. ph. 54
Picia gracilis Loi.										
19277 pubescens Dec.	pubescent	J, O	un	1	jn.jl	-	Italy	1820.	S	co

	PBA.	(P ^{ss} , pea, Celt)	Leg.	Pap.	Vie.	7.
19235. PPSUM Ton.	com. cultivated	J	O cu 3 jn.s	W	& Europe	... S co Lam.il.633
19278 sativum L.	sugared	O cul	jns	W	—>	... S co
2 saccharatum Ser.	long-podded	O cul	jns	W	—	... S co
3 macrocarpum Ser.	umbellate	O cul	jns	p	—	■ co Tab. ic 495
4 umbellatum L.	squared	O cul	jns	W	—	M S co
5 quadratum L.	humble	O cul	jns	W	—	... S co
6 hbmile Pair.	tall	O cul	jl.au	D,P	1820.	S co
19279 elatum 2toA.	Jomardi's	O cul	jl.au	W	Iberia	1820. S co
19280 Jomardi Schrank	Theban	C r	3 jl.au	...	Egypt	1825. S co
19281 arvense Z.	field	O ag	3 jns	R	S. Europe	... S co M.h.2.2.1.4
19282 maritimum L.		O or	li jl	P	England sea sh.	R 81 Eng. bot 1046
19284 americanum Afl.	American sea	A or	1 jlau	P	N. Amer.	1800. R sp.1

•2136. ZATHYRUS L. LATHYRUB. (*La, augment thouros, anything exciting; qual.*) Leg. Pap. Vic. 50.-6*

I. EULA'THYRUS.—*Vexillum at base toothless; leaflets opposite, or from abortion none; petioles narrow-winged.*

	Petioles peduncles many-flowered; leaves one-paired, from No. JL929t. many paired.						
19285 sylvestris L.	wood Everlasting	A or	2 jl.s	P	Britain ^m moi. w.	S co	Eng. DOC. oÜft
19286 interm ^d ius WaUr.	intermediate	± A or	4 jl.s	R	N. Europe	1820. S co	Muhl. dan. 785
19287 magellanicus Lam.	Magellan	T A or	6 jl.s	P.B	Cape Horn	744. S co	
19288 latifolius L.	broad-lvd Everlasting	T A or	6 jl.s	Pk	England woods.	S co	Eng. hot 1108
19289 rotundifolius W.	round-leaved	T A or	ijjl. au	P	Tauria	1822. S co	Bieb. cen. 1.22
19290 grandiflbus B. M.	large-flowered	T A or	4 jn.au	P	S. Europe	1814. S co	Bot. mag. 1938
19291 prate ⁿ nei L.	meadow	H A or	3 jn.au	Y	Britain me. pa.	S co	Eng. bot II70
19292 tuberosus L.	tuberous	A cul	2 jl.au	R	Holland	1596. S co	Bot mag. III
19293 rbeus Stev.	rosy	T A or	jl.au	R	Iberia	1822. S co	
19294 /risifirmis L.	Pea-formed	T A or	3 jl.s	W, B	Siberia	1759. S co	L.f. dec. 20
19295 mutabilis Swt.	changeable	T A or	4 jl.s	P.R	Siberia	1825. S co	Sw. ft*, gar. 192
19296 californicus B. R.	Californian	T A or	4 jl.s	P	California	1826. S co	Bot. reg. 11-14
19297 paltistrisi.	marsh	JL A or	4 jl.au	Pa.P	Britain moi. w.	S co	Eng. bot 169
19298 myrtifolius Mhl.	Myrtle-leaved	J A or	3 jl.s	P	Philadelphia	18*2. S co	
19299 polymorphic Nut.	multiform	ft	A or	3 jl.s	Pa.P	Missouri	1824. S co
19300 heterophylus L.	various-leaved	T A or	4 jl.s	F	Europe	1731. S co	Bau. his. 2.3(4.1)
19301 stipubecus Lee.	/ar/jr-stipuled	J A or	3 jl.au	P	New York	1816. S co	

ii. A'NNUI.—Annual; peduncles one or three-flowered; petioles leafless, from No. 19304; leaves one-paired, from No. 19320. two or three-paired.

19302 '^phaca L.	Yellow Vetchling	O or	1 Y	England	san. fl.	S co	Eng. bot. 1167
19303 Nissolia L.	Nissolia	O or	1 my	C	England bus.pL	S co	Eng. bot. U.S
19304 inconspicuous L.	inconspicuous	O cu	1 jl.au	P	Levant	1739. S co	Jac. vin. 1.86
19305 sphs'B'ricus Rctx	round-seeded	O or	1 C	C	S. Europe	1801. S co	Dec. ic. 1.32
19306 coccineus Pers.	scarlet	i O or	2 jn.j	S	Italy	1800. S co	
19307 micranthus Ger.	small-flowered	O or	1 jl.s	P	S. France	1816. S co	
19308 angulatus L.	angular-seeded	O or	1 jl.s	R	S. Europe	168a. S co	Bux. c. 3. 42.2
19309 longipedunculatus Led.	long-pedund.	O or	1 jl.s	P	1817. S co	
19310 leptophyllus Bieb.	fine-leaved	O or	1 jl.s	P	Caucasus	1818. S co	
19311 setifolius L.	bristle-leaved	O or	1 jl	R	S. Europe	1739. S co	Bau. his. 3.308
19312 amphicarpos L.	both sides fruited	O cu	1 jl	Pk	Levant	1680. S co	Sw. fl. gar. 236
19313 sativus L. cultivated	Chickling Vetch	± A or	3 jl.s	W	S. Europe	1640. S co	Bot. mag. 115
19314 Cicera L.	Cicraflat-podded	J O or	1	R	S. Europe	1633. S co	Gcr. cm. 3
19315 finnuus L.	annual	J O or	4 jn.au	Y	S. Europe	1621. S co	Bux* c. 3.42!
JJSJS "tus TV.	hairy-podded	J O or	4 jl	P	England bor. fl.	S co	Bot. mag. 1255
^ Mrtus Lam.	hairy	J Q cul	4 jl.au	P	Europe	1800. S co	
19318 odoratus L.	fragrant Sweet Pea	O fra	4 jl.s	Va	Sicily	1700. S r.m	Bot. mag. 60
Tangier	-	O or	4 jn.au	D,P	Barbary	1680. S co	Bot. mag. 100
19320 bith'nicus Lam.	Bithynian	- O or	1 jlau	P	England san.sh.	S co	Eng. bot 1842
Ptea bith'nica L.							
19321 sasiflorus Ser.	sessile-flowered	L O or	1 jl.au	P	Switierl.	1819. S co	Stur. ger. 1.32. ic
tomidus L.	tumid	J O or	1 jl.au	R	Piedmont	1820. S co	Al. ped. 1.26.2

II. CLY'MENUM.—*Vexillum at base marked on each side with a conical swelling; superior leaves two or si* foliate; leaflets often alternate; petioles often winged.*

19322 inc&rvus Roth	curved-nodded	✓	✓	Russia	" 1802. T co	Bux.c.4.46
19323 afatua Ten.	winged	✓	✓	Italy	1823. S co	
19524 Clhraenum L.	Clymenum	✓	✓	Levant	1713. S co	Pluk. al. 114. R
1>-J-J articulatus L.	joint-podded	✓	✓	& Europe	1640. S co	Bot. mag. 253
ID J-J auriculatus Bert.	cared	✓	✓	S. Europe	1800. S co	
19J27 sp&rius W. en.	spurious	✓	✓	1815. S co	
19528 purpureus Desf.	purple	✓	✓	Crete S co	Desf cor. 61
19329 tenuifolius Desf.	fine-leaved	✓	✓	N. Africa	1820. S co	
IS330 corniiUiB Horn.	horned	✓	✓	1818. S co	

III. DU^BII.—Doubtful to which of the sections they belong.

19331 helbdes Lk.	marsh	JL O or	4 jl.au	P	18V7. S co
19332 iUlicus Fit.	Italian	T O or	3 jn.au	...	Italy	1827. S co
19333 luatinicua Mart.	Lusitanian	i O or	3 jn.au	...	Spain	1827. S co

2137. CTCHRUS Pers. OcHaua (Ochros% pale; pale muddy-colored flowers.) Leg. Pap. Vic. 1.
19334 nkWid* Pers. pale i O or 2 jn.au Pa.Y S.Europe ... S co Tou.ins.219

•2138. O'ROBUS Tou. BITTER VETCH. (*Oro, to excite, bous, an ox; nourishing food.*) Leg. Pap. Vic. 34.-48.

I. UNIJUGL—Leaves one-paired; leaflets ovate or linear.						
19335 laxiflbrus Desf.	loose-flwd hairy	A O or	1 myjl	V	Candia	1820. R pi
19336 hirs&tus L.	hairy	5 A or	1 my,jn	R	Thrace	1822. R co
19337 formbusus Stev.	beautiful	5 A or	f my,jl	P	Caucasus	1818. R pl
19338 tethyrides L.	Lathyrus-like	^ A or	2 jn	B	Siberia	1758. R pl

II. MULTIJUGL—Leaves many-paired; leaflets broad; stipules much smaller than the leaflets; from No. 19359.

J9339 vénus L.	spring	A or	mr.ap	P	Europe	1629. R s.1	Bot. mag. 521
19340 pauciflous Hort.	flowered	A	1 myjf	P	1820. R pl	
19341 yenhus JV.	veiny	A or	1 my,jl	B	Siberia	1816. R pl	Gin si. 5.1
19342 naultiflorus Sie	rock	A or	2 jl	P	Hungary	1817. R pl	W. & K.2L126
19344 variegatus 7V».	many-flowered	A or	2 jn.au	Pa.R	Italy	1820. R pl	
	variegated	A or	1 jn.au	P	Italy	1121. R pl	Fl. nap. 68

19345 <i>vlcoldes Dec.</i> <i>Fic'ia oroboldes WuL</i>	Vicla-like	■ A or ■ ■ ■	■ Hungary	1819.	R p.l	W. & K. 3.842	
19346 <i>luteus L.</i>	yellow	■ ■ ■ or ■ ■ ■	L.Y	Siberia	1759.	R h.l	
19347 <i>Tourneforti Lap.</i>	Tournefort's	■ ■ ■ or ■ ■ ■	Y.p	Pyrenees	182L	R p.l	
19348 <i>laevigatus W/JK.</i>	smooth	■ ■ ■ or ■ ■ ■	Y	Hungary	1820.	R p.i	
19349 <i>aurantius Siev.</i>	orange	■ ■ ■ or ■ ■ ■	Ift my.jl	Y	1818.	R p.i	
19350 <i>sylvaticus L.</i>	wood	■ ■ ■ or ■ ■ ■	2 my.jl	CP	Britain m.wo.	R p.i	
19351 <i>ochroleucus W.%K.</i>	yellowish white	■ ■ ■ or ■ ■ ■	Pa.Y	Hungary	1816.	R p.i	
19352 <i>rfigerZ.</i>	black	■ ■ ■ or ■ ■ ■	P	Britain woods.	R p.i	Eng.bot518	
19353 <i>hiimilis Ser.</i> <i>/athyrus hiimilis Fit.</i>	humble	■ ■ ■ or ■ ■ ■	\JLau	P	Dahuria	1825.	R p.i
19354 <i>tuberbus L.</i>	tuberous	■ ■ ■ cul	1 my.jn	P	Britain heaths.	R p.i	Pluk.aLS10.2
19355 <i>pyrenaicus L.</i> <i>PluknēU' Lap.</i>	Fyrenean	■ ■ ■ or ■ ■ ■	2 my.jn	P	Pyrenees	1699.	R p.i
19356 <i>tenuifolius Roth</i>	fine-leaved	■ ■ ■ or ■ ■ ■	I my.jl	P	Germany	1810.	R p.i
19357 <i>divaricatus Lap.</i>	divaricate	■ ■ ■ or ■ ■ ■	f my.jl	P	Pyrenees	1816.	R p.i
19358 <i>erectus Pair.</i>	erect	■ ■ ■	1 jl.au	P	—	R p.i	Mém. m. 2.11
19359 <i>varius Sol.</i>	various	■ ■ ■	11 my.jn	Y.a	Italy	1759.	R p.i
19360 <i>can&cens L.</i>	caneeccnt	■ ■ ■	If my.jn	W.a	France	1816.	R co
19361 <i>pallidissimis Bieb.</i>	pallescent	■ ■ ■	1 my	W	Tauria	1823.	R co
19362 <i>albus L.</i>	white-flowered	■ ■ ■	1 my.jn	W	Austria	1794.	R s.J
19363 <i>lacteus Bieb.</i>	milk-white	■ ■ ■	1 my.jn	W	Caucasus	1820.	R co
19364 <i>angustifolius L.</i>	narrow-leaved	■ ■ ■	1 my.jn	W	Siberia	1766.	R s.l
19365 <i>saxatilis Ven.</i>	rock	■ ■ ■	1 JLau	P	S. France	1820.	S p.i

III. Difrii. — *Species not sufficiently known.*

19366 <i>coccineus Mil</i>	scarlet	^ A or	1 ap.my	S	N. Amer. ? ...	D co
2 <i>unijugus Dec.</i>	conjugate	& A or	1 ap.my	S	N. Amer. ? ...	Bot. cab. 883
19367 <i>venetus Mil.</i>	Venetian	5 A or	1 mr.ap	P	Germany ...	Mild.ed.2.193.2
19368 <i>Gmeini Fis.</i>	Graelin's	5 A or	1	Siberia	1827.	D co

2139. PLATYSTYLIS Swt. PLATYSTYLIS. (*Platys*, broad, *stylis*% style.) Leg. Pap. Vic. S.19369 *cyanea Swt.* azure i A or 1 ray P Caucasus 1823. R co Sw.fl.gar.239P'robos *cyaneus Stev.*19370 *sessiliflolia Swt.* sessile-leaved 3t A or 1 my P Tauria 1823. R co Bot mag. 2796Cfrobos *sessilifolius Swt.* digitatusA^ pyrenaeus PalL2140. A'RRUS L. WILD LIQUORICE. (*Abros*, soft; extreme tenderness of leaves.) Leg. Pap. Phas. 1.19371 *precatiorius L.* prayer ± • or 12 mr.my Pa.P W. Indies 1680. S s.p R. mal. 3.392141. SWEE*TM Dec. SWEETIA (*Robert Sweet*, F.L.S., author of several bot periodicals.) Leg. Pap. Phas. 3.19372 *longifolia Dec.* long-leaved S O or 4 jl.au P S. Amer. 1818. C fl.p Jac. ic. 572Galega *longifolia Jac.* Tcphrbis *longifolia Pert.*19373 *filiformis Dec.* thread-shaped ft. CD or 3 JLau P S. Amer. 1820. C s.p Jac. ic. 573Galega *filiformis Jac.*19374 *lignosa Dec.* woody i Q o r 5 ... P StDōm. 1824. C s.pGly*cine *lignosa Turp.*2142. MICRANTHUS Poir. MICRANTHUS. (*Mikros*, small, *anthos*, flower.) Leg. Pap. Plat. L.19375 *cochininchensis Lou.* Cochinchinese \$ ES clt 3jn.au W China ... S co2143. AMPHICA'RPA.EW. AMP. (*Amphi*, on both sides, *karpot*, fr.j above and undground.) Leg. Pap. Phat. %19376 *sermentosa EU.* twiggy \$ O or 2 a Ap N. Amer. 1820. S fl.p Ab. ins. 21Gl*cine *filosa Horn*, *sermentosa Roth*19377 *monoica EU.* monoecious j O ot • 1 Ap N. Amer. 1781. S s.p Ree. ax. L 8Gl*cine *monoica L.*19378 *comosa G. Don* tufted J O u n 1} jLs B N. Amer. 1812. S p.iGlycine *comosa L.** 2144. KENNEPD/a Ven. KENNEDIA. (*Mr. Kennedy*, nurseryman, Hammersmith.) Leg. Pap. Phas. 12.

I. LONGICARTAE. — Leaves trifoliate; keel straight, rather longer than the vexillum.

19379 *rubicunda Ven.* rubicund ft. uJ or 10 mr.au Br N. S. w. 1788. C s.p Bot mag. 26819380 *prostrata R. Br.* prostrate £ | J or 4 mr.jn S N. S.W. 1790. C s.p Bot mag. 27019381 *moph'ua Cun.* nerve-leaved ft. iJ or 4 ... S N. Holl. 1824. C fl.p19382 *dilatata Cun.* dilated ft. uJ or 4 ... S N. Holl. 1824. C s.p

II. BREYCARME. — Leaves trifoliate; keel shorter than the vexillum or wings.

19383 *coccinea Ven.* sc&rlc U'loweredy, 1 or 10 my.au S N. Holl. 1803. C s.p Sw. au. 2319384 *Comptoniana Lk.* Compton's ft. iJ or 12 mr.jn B N. Holl. 1803. C s.p Bot rep. 29819385 *parviflora Ma. C.* small-flowered ft. J or 4 ... N. Holl. 1824. C s.p19386 *heterophylla Ma. C.* variable-leaved % \ or 4 ... N. Holl. 1824. C s.p

III. SiMPLICifbL*. — Leaves simple s keel shorter than vexillum or wings.

19387 *monophylla Ven.* 8irole-leaved % iJ or 10 mr.jn P N. S. W. 1790. C s.p Bot mag. 26319388 *ovata B. M.* ovate-leaved £ iJ or 6 my.au P N. Holl. 1818. C fl.p Bot mag. 2169

latifolia 2!mft

IV. B\BV\BISL*. — Species not sufficiently known.

19389 *apetala Lo. C.* apetalous fc iJ or 4 ... Ap 1824. C s.p19390 *sericea Cun.* silky i.L| or 4 N. Holl. 1824. C s.p2145. RHYNCHCTSIA Lou. RHYWCHOM* (*Rhynchos*, beak; keeL) Leg. Pap. PhateU. 18.—51.

I. MONOPHYLLS. — Leaves all or far the most part simple, roundish, on longfootstalkt.

19391 *reniformis Dec.* kidney-formed-fb* £ j± pr 2 jl.au Y Carolina 1806. D p.i* Gfcine *reniformis Ph.*19392 *dimorpha Dec.* two-formed £ A un 1ij ■ Y N. Amer. 1732. D p.l Di.cL26.29Glycine *tomentosa Ph.* var. *voldbilis Mx.*

II. PHASEOLOI'DE*. — Leaves trifoliate, on lone footstalks.

19393 *erecta Dec.* erect ^ A pr SJjns Su N. Amer. 1820. D p.i

Gfcineerdcta

19394 *caribea Dec.* Caribsan J_E3pr S 6.0 Y W. Indies 1742. C p.i Bot.reg.275Glycine *caribea Jac.*

19395 <i>minima</i> Bee.	imallest	% 23 un	1} Jlau	Y	Jamaica	1776.	S s.1	Jacob.	1.22
ZMlichos <i>minimus</i> L.									
19396 <i>phaseoloides</i> Dec.	Kidneybean-lik	JL ES un	2 jn.jl	Br	Jamaica	1818.	C p.l	Bot. mag.	2284
Glycine <i>phaseoloides</i> Swx.	·<lit	_ Q u n	4 JLau	Y	Mexico	1825.	C pj		
19397 <i>precaturia</i> Dec.	prayer	·<lit							
Glycine <i>precatoria</i> W. en.									
19398 <i>reticulata</i> Dec.	neUeaved	i p l q r *	6 JLs	W	Jamaica	1779.	C p.l		
Glycine <i>reticulata</i> Vahl									
19399 <i>rhombifolia</i> Dec.	rhomb-leaved	JLESun	6 my.au	...	R Indies	1815.	C p.l		
19400 <i>suaveolens</i> Dec.	sweet-scented	I I Q o r	3 jl.s	Y.B	R Indies	1816.	C p.l		
Glycine <i>suavolens</i> L.									
19401 <i>viscina</i> Dec.	clammy	£ C 3 u n	3 jl.s	Y	R Indies	1820.	C p.l		
Glycine <i>viscosa</i> Both.	glutintaa Lech.								
19402 <i>viscidula</i> Dec.	viscid	i n u n	3 in.au	Y	R Indies	1823.	C p.l		
Frederic's	I Q u n	4 Jlau	Y		a Amer.	1820.	C p.l		
19404 <i>Bcarab\$Eodes</i> Dec.	scarabeus-like	* O u n	2 jjjl	P	R Indies	1773.	S s.1	Pluk.aL53.3	
Dolichos <i>scarabaeoides</i> L.									
19405 <i>biflbra</i> Dec.	two-flowered	J Q u n	5 JLau	Y	R Indies	1816.	C p.l		
JJolichos <i>scarabaeoides</i> Rot.									
19406 <i>angustifolia</i> Dec.	narrow-leaved	\$L_Jun	6 jn.au	Y	C. G.H.	1795.	C p.l	Jac.sc.2.231	
Glycine <i>angustifolia</i> Jac.									
19407 <i>mfilif</i> Dec.	soft	S Q u n	3 JLau	Y	Guinea	182a	C p.l		
Glycine m6lis W.									

III. EBIO9E^NMA. — Leaves trifoliate, all on short footstalks / racemes or fascicles of flowers axillary ; vex

Ulum

silky villous; stems not climbing.

19408 <i>violetacea</i> Dec.	Violet-like	« Q u a	3 jl.au	Y	Guiana	1820.	C p.l	Aub. gui.	2.306
Glycine plecta Vahl,	Cy'tiaus uiolaceus Aub.								

2146. FAG&LIA Neck.	FAGHLIA.	(Not explained.)			Leg. Pap. Phased.	I.	
19409 <i>bituminosa</i> Neck.	clammy	i. C 3 o r	4 ap.s	Y.r	G.G.H.	1774.	
Glycine bituminbsa L.					€	Lp	Botreg.261

2147. WISTA ^{^R} /il Nut	WISTARIA.	(Caspar Wistar, a professor in Pennsylvania.)	Leg. Pap. Phasebl.	3.
19410 Consequam <i>li</i> Loudon	Consequa's	J; or 15 ray.Jn B	China	1818.
sinensis Dec.	Glycine sinensis B. R.		L r.m	Sw. fl.gar.
19411 fruticulosa Nut.	fruticosus	J or 10 jn.s	P	N. Amer.
Glycine fruticulosa L.	A ^{^p} ioa frutescens PA.		L s.p	BOL mag.
19412 floribunda Dec.	bundlcflwd	* or 10 ...	P.w	China
Z)6chos iiolystachyos Thun.	Glycine floribunda W.		L r.m	H.pfl.8.64.2

2148. iTPIOS Boer.	APIO9.	(Apion, a pear; form of roots.)	Leg. Pap. Phasebl.	1.
19413 tuberosa Moen.	tuberous-roofcZA & or	6 au.s	.Br.pk.N. Amer.	1640T
Glycine ifpios L.			R s.p	Bot mag.
				1198

2149. PH ASETOLUS L. KIDNETBKAN. (*Phaselus*, little boat; sup. resemb. in pods.) Leg.Pap.Phas. 46.-63.

I. EupiiASBN)Lua. — Legumes compressed.

L CABACA'LLA.—Stems pubescents roots fasciculately tuberous; leaflets entire; vex <i>xiilum</i> twisted.				
19414 <i>Caracilla</i> L.	Caracalla	Jfc ESI cul	If au.s	Li India
				1690.
ii. PERE'NNIS.—Stems herbaceous roots perennial; racemes subpaniculate ; leaflets entire.				

19415 <i>petennis</i> Walt.	perennial	-\$ £ or	3 jl.au	D.P Carolina
66lchos polystachyos L.				1824 D s.1
iiL MACROPODII.—Annual; leaflets entire; peduncles longer than leaves.				
winged				
19417 specibus Hum.	showy	J O u n	3 jl	P Carolina
				1732. S s.1 DleL235.303
19418 sylvestris Kth.	wood	J O or	6 jl.au	S Orinoco
				1820. S s.1
19419 multiflora W.	many-flowered	-\$ O or	6 jl.au	S Mexico
2albus	white-flowering	\$ A cul	12 jl.s	S a Amer. ? 163a
				S co Schk.han.2.199.a

iv. BRACHYFODH.—Annual; leaflets entire; peduncles shorter than leaves.				
19100 <i>vulgaris</i> L.	common	Ji; O cul	1 jn.I	W India
2 fawiatu	banded-seeded	i O cul	... jl	W India? 1597. S co
3 variegatus	variegated-seeded	& O cul	... jl	W India? 1597. S co
19421 compressus Dec.	compressed	* O cul	2 jl.au	W — S co Savi ph. 3.10. SO
19422 oblonguB Savi	cttlong-seeded	— O cul 1}JLS		— S co Savi ph. 3.10.14
19423 saponaceus Savi	saponaceous	J O cul	1 JLS	■ a. v #*— S co Savi ph. a 10.15
19424 tumidus Saw	tumid	Nain Flageolet	O cul 2 jl.s	W — S co Savi ph. 3.10.16
19425 hsmatoc&pus Savi	blood-ikMlded	^ O cul	4 jl.au	Pa.V — S co Savi ph. 3.10.17
19426 spharicus&avi round	Haricot d'Orl.J.	O cul	4 jl.au	Pa.V S co Savi ph. a 10.18
19427 gonospermum Savi	angled-seeded	^ O cul	4 jl.au	Pa.Vw — S co Savi ph. 3.10.19
19428 derlfius Schrank	acer&ped-offbk.seed.	J O cul	3 jl.au	W.a Brazil 1819. S co Seh. mo. 1.89
19429 lunatus L.	hay-moon-podded	& O cul	In.jl	G R Indies 1779. S co H.n.h. 10.63.1
19430 inatpo>nb L.	unpleasant	J O u n	6 jl.au	G.yr Africa 1794. S co Jac. vin. 1.66
19431 torbus Rox.	torose	JO or	4 jl.au	V Nepal 1818. S co
19432 Xuarecsf Jac.	Xuare's	J O or	4 jl.au	R a Amer. 1818. S s.1
v. HETEBOPHYLLI.—Leaflets all or some of them lobed.				
19433 microspcrmus Or.	small-seeded	O or	1 jl.au	D.P Cuba 1825. S s.1
19434 heteroph?us W.	variable-leaved	i O or	4 JLau	R Mexico 182a S s.1
19435 ocomitibliss Jac.	Acombe-leaved	\$ O cu	9 jn.au	Pk R Indies 1731. S s.1
				Jac. ob. a 52

II. STROPBO'STYLKs.—Legumes terete.

i. LOBATIFOLII.—Leaflets all lobed.				
19436 diversifolius Pers.	various-leaved	Jc O un	1 Jj.jl	P N. Amer. 1806. S s.1
trilobus A/x.				
19437 trilobus Roth	three-lobed	O un	2 jl.au	G R Indies 1777. S s.1 Bur. in. 50.1
19438 subtrilobus Lk.	sub-three-lobed	C3 un	1 jl.au	Y Brazil 1824. S co
19439 angulata Or.	angular-feauoi	J O or	3 jl.au	P.a.lt. . . . 1820. S s.1
19440 farinibus L.	mealy	3 O un	3 jl.au	Pk R Indies 1759. S co N. a. p. 1730. 42
19441 stipularis Lam.	largy-stipuled	O un	2 jl.au	Y.Br Peru 1805. S s.1
19442 namicis W.	O un	1 jn.s	W India	... S s.1
ii IXTEGRIPOLII.—Leaflets entire.				
pale red	J O cul	3 jl.au	Pa.R Carolina	1732. S co Di.cL233.300 .
Mungo	O cul	1jj.jl	Y India	17y0. S s.1

19445 chrysanthos Savi	golden-flowered	Q or	S	JLau	Y		f.1	
19446 radiatus L.	rayed	O un	1	jn.jl	P	China	1732.	8.1 M. el. 235.305
19447 scaber Steu.	scabrous	O cul	3	JLau	Y.o	E. Indies		co M. h. 2.5.8
19448 lathyroides L.	Lathyrus-like	ED or	2	JLau	S	Jamaica	1786.	s.1 SI.jam. 1.116.1
19449 semierectus L.	half-erect	ED or	2	JLau	R	W. Indies	1732.	co Bot. reg. 743
19450 violaceus Steu.	violet	ED or	3	JLau	R	Africa	1800.	8.1
19451 Max L.	III. Dttaii. — Doubtful to which of the sections they belong.	Max	O cul	JLau	India	17...	S 8.1	Ru. am. 5.140
19452 hirtus Retx	hairy	i O or	3	JLau	—	C. G. H.	1818.	S co
19453 tifolichoides Box.	Dolichos-like	i O or	3	JLau	R	E. Indies	1824.	S co
19454 amoe'nus ForsL	pleasing red	J O or	3	JLau	R	Society ls.	1820.	S co
19455 mesoleucus Forst.	white-middled	J O	4	JLau	—	Brazil	1818.	S co

2150. A'MPHODUSIUM. AUPHODUK (*Ataphi*, on both sides, *odous*, a tooth.) Leg. Pap. PhasebL 1.
19156 ovktus Lndl. ovate 2. IS] cu 10 JLau P Trinidad 1820. C s.p. Bot reg. 1101

2151. 8ft JA Moen.	SOJA.	(<i>Sooja</i> , its name in Japan.)				Leg. Pan. Vic. 1.	
19457 hispida Moen.	hispid	J O cul 3 JLau	V			E. Indies 1790? S s.1	Jac.ic.145

2152. DCLICHOS L. DOLICHOS. (*Dolichos*, long; length of climbing stem.) Leg. Pap. PhasebL 24.—51.

I. EUDOLICHOS.— <i>Legumes compressed, apiculated with the style.</i>								
19458 lignatus L.	woody	J 1 J or	12	JLau	P	E. Indies	1776.	S p.! Bot mag. 380
19459 Jacquinii Dec.	Jacquin's	J 1 J or	8	JLau	W	S. Amer.	1800.	C s.1
19460 hirsutus Thun.	hirsute	S JL or	10	JN	P	China	1802.	C s.1
19461 pilosus Klein	pilose-podded	J ED un	3	au	PR	Indies	1790.	S s.1
19462 tetraspermus W.	four-seeded	JLCQun	3	my.au	Pa.Y	E. Indies	1816.	S s.1
19463 capensis L.	Cape	t IAI or	6	JLau	Y	C. G. H.	1821.	S s.1
19464 lieterophyllus Horn,	variable-lvd	J O cul	5	JLau	—	Canaries	1810.	S co
19465 likteus Sun.	yeUovr.jtowered	CD un	8	JLau	Y	Jamaica US	12.	S s.1
19465 Irribius L.	two-flowered	f, [S3 un	3	JLau	Pa.Y	E. Indies	1776.	S s.1
19467 Catjang LL	lobed	[S3 un	3	JLau	P	E. Indies	1793.	s.1 R.mal.8.41
19468 monachalis Brot.	monks*	O cul	2	JLau	Pa.R	Spain	1816.	co
19469 sinensis L.	Chinese	j O P'	6	frau	Pa.R	India	1776.	i.1
19470 Litbia Forsk.	Lubia	O cul	U	JLau	W.B	Egypt	1818.	co
19471 vexillata* Plastostolus vexilatus L.	vexillated	JO cul	3	JLau	Y	W. Indies	1732.	co Di. el. 2. 302
19472 lobatus W.	lobed	J O or	3	JLau	Y	C 6. H.	1800.	co
19473 angulus Dec.	angulose	JL O or	2	JLau	Y	N. Amer.	1820.	pi
III. UNGUICULABIA.— <i>Legumes subcylindrical, ending in beaks; leaves entire.</i>								
19474 unguiculatus Jac.	clawed	J O un	3	jn.jl	Y	AV. Indies	1780.	S s.1 Jac. vin. 1.23
19475 tranquebancus Jac.	Tranquebar	f O un	3	jn.jl	W.p	E. Indies	1801.	S s.1 Jac. vin. 3.70
19476 seqquipedalis L.	1 ft. long-podded	J D un	6	au	W	W. Indies	1781.	S s.1 Jac. vin. 1.67
19477 inelanupthalamus Dec.	black-eyed	JO cul	3	JLau	—	Italy	1800.	S co
19478 gangeticus Rox.	Gangetic	% CS cu	6	jn.au	—	E. Indies	...	S co
19479 sphacelospermum Dec.	round-seeded	O cul	4	JLau	—	Jamaica	1816.	S co SL jam. 117
IV. Du'an.— <i>Species not sufficiently known.</i>								
19480 reticulatus H. K.	net-leaved	f, Lj un	3	jn.jl	P	N. S. W.	1781.	8.1
19481 setifolius Boj.	bristle-leaved	S O un	4	—	—	Mauritius	1826.	s.1
19482 frutescens Ham.	Nepal frutescent	m i_J or	10	jn.jl	Pa.Y	Nepal	1820.	8.1

2153. VI'GNIL Savi VIGNA. (*Dominic Vigna*, commentator on Theophrastus.) Leg. Pap. PhasebL 1.—■
19483 giSibra Savi smooth £ O P* 4 JLau Y M. Amer. 1085. S s.1 Jac. vin. 1. SO
261ichos lutéolus Jac.

2154. LA'BLAB Adan.	LABLAB. [<i>Lablab</i> , the Arabic name of the convolvulus.]	Leg. Pap. Phased. 8.
19484 vulgaris Savi	common	—\$ O cul 8 JLau V E. Indies 1794. S s.1 Bot.mag.8U6
Dulichos Lablab L.		
19485 purpureus G. Don	purple	% CS or 12 au.i P E. Indies 1790. S s.1 Bot reg. 830
vulgaris & purpureus Dec.	JDAlichos purpurcus Jac.	
19486 bengalensis G. Don	Bengal	J O cul 8 JLau W Kindles 1800. S s.1 Jac. vin. 2.124
vulgaris & albiflorus Dec.	Z?6lichos bengalensis Jac.	
19487 nankinicus Savi	Nankin	—\$ O cul 8 JLau W China 1820. S s.1 Savid.22. 1&dg
19488 leucarpus Saw	white-podded	\$ O cul 8 JLau W E. Indies 1816. S s.1 Savi d. 22.9.a.d
19489 culturatus Dec.	cu'traUi-podded	J O cul 8 JLau W Japan 181a S s.1 Ksm ic. 25
>6lichos culturatus Thun.		
19490 microcarpus Dec.	small-podded	JE O cul 8 JLau P E. Indies 1818. S s.1 Ru. am. 5.141.1
19491 nerennans Dec.	lasting white	Obit% i cu 8 JLau W China 1820. S s.1 Ru. am. 5.137
D6lichos albiis Lou.		

2155. PACHYRHPZUS Rich. PACHYEHIZUS. (*Pachys*, thick, *rhtxa*, root; *tuberous*.) Leg. Pap. PhasebL 1.-3.
19492 angulatus Rich. angled | . D o r 5 Jl P E. Indies 1781. C s.1 Ru.am.5.132
D6lichos bulbosus L.

2156. PARCYCHETUS Ham. PAROCHETUB. (*Para*, nigh, *ochetus*, a brook.) Leg. Pap. Phasol. 1.—2.
19493 communis Ham. common HW-AJor 1 jl P Nepal 1820. D s.1

2157. DIOCLEJ Kth. DIOCLBA. (*Diodes Carystinus*, an ancient Greek botanist) Leg. Pap. PhasebL 1.
1S494 mtilis Dec. soft \$ E2un 3JLau Pa.Y 1824. D s.1 Jac.fr. 88
Dolichos m6llis Jac.

2158. PSOPHOCA'RPUIS Neck. PSOPHOCAHPI^ (*Psophos*, a sound, *karpos*, a fruit.) Leg. Pqp. PhasebL 1.
19495 tetragon6lobus Dec. square-podded \$ O < • <>n B Mauritius 1816. S s.1 Ku. am. 5.133
Dolichos tetragonolobus L.

*i%9. MUCU^NA Adan. StVec COW-ITCH. (*Muama-Guaca*, name in Brazil.) Leg. Pap. Phas. 12.—20.

I. ZOOPHTII A'LMUH.—*Legumes with lamellae transverse furrows.*

19g urens Dee. burning i Q c u 12 Jn.jl Y W. Indies 1691. C l.p. PI. am. 107
"/irhmifrrni L. Stisoltbiiim fin>n<r"

II. STIZOBLOM.—Legumes destitute of transverse lamella.									
19497 altissima Dec.	tallest	Q o r SO	...	P	Martinico	1779.	C	I.p	Jac. am. 182185
Stizolobium altissimum Pen.									
19198 prhriens Dec.	common stinging	\$ • cu 12	...	P	India	1680.	C	I.p	Jac.anf122
Btzolobium prbriens Per*.									
19499 gigantea Dec.	giant	fc_	• un 20	jl.lau	O	E. Indies	1815.	C	Lp R.mal.8.36
Carpopogon giganteus Box.									
19500 imbricata Dec.	imbricated	S • or 10	...	P	E. Indies	1815.	C	Lp	
Carpopogon imbricatus Box.									
19501 mltis Dec.	mild	* Qun 10	...	P	Peru	■■■	C	Lp'	
19502 macrocerafldes Dec.	long-horned-lk	St ■ tun 10	...	V	Brazil	1824.	C	Lp	Raddim.13.5
III. DU'BIB.—Doubtful to which of the sections they belong.									
19503 monosprlma Box.	one-seeded	f • un 10	E. Indies	1816.	C	Lp	
19504 atropurpurea Box.	dark-purple	• cu 10	...	P	E. Indies	1820.	C	p.l	
19105 nlvea Box.	mow.white	f • un 10	...	W	E. Indies	1816.	C	a.P	
19506 bracteata Box.	bracteate	[L] cu 10	E. Indies	1826.	C		
19507 anguina Rox.	snake	■■■ 10	E. Indies	1817.	C	p.l	
21G0. CAJA*NUS Dec.	PIGEON PEA.	(Catiang, its name in Malabar.)			Leg. Pon. Phosebl	2.			
19508 tricolor Dec.	two-colored	■ D cu 1 4 jLau Y			K Indies	1800.	S	s.l	Jac. vin. 2.119
Cytus Pschdo-G	G W Jac.								
19509 flavus Dec.	yellow	■■■ Q cu 1 4 jLau Y			E. Indies	1687.	3	§.1	Jacob. LI
Cytus Chjan L.									
2161. CYLI'STA H. K.	CVLISTA.	(Kylstos, twining; plants.)			Leg. Pap. Phasebl.	4.-5.			
19511 tomentosa Box.	woolly	ft. O or 4	...	V	E. Indies	1806.	C	p.l	Rox. cor. 1.92
19512 albiflora B. M.	white-flowered	F'Q-O-F 4	...	W	E. Indies	1816!	C	p.l	Rox.cor. 201-
19513 villosa H. K.	villose	* or 6 ap.my	W		Mauritius	...	C	p.l	Bot. mag. 1859
					C. G. H.	1776.	S	p.l	Bot rep. 446
* 2162. ERYTHRn A L. CORAL TREE.	{Erythros, red; flowers.)	Leg. Pap. Phasebl.	31—40.						
1. ACAU'LES.—Stemless; fructiferous racemes and leafy branches annually rising from subterraneous trunks.									
19514 herbacea L.	herbaceous	Jc uJ or 3 jn.s	S		Carolina	1724.	C	I.p	Bot mag. 87/
19515 resupinata Rox.	resupinate	£ • spl 1	...	Pa.S	E. Indies	182J.	C	r.m	Rox. cor. 1220
19516 hrrida Dec.	rough	* a sp! 5	...	S	Mexico	1824.	C	r.m	Fl. mex. ic. in.
II. CAULEBCE'NTES.—Shrubs or trees, stems bearing leaves and racemes of flowers.									
19517 carnea H. K.	flesh-cafor**	* or '12 my	Pk		Vera Cruz	1733.	S	r.m	Tr. ehr. 8
19518 Corallodendrum L.	Coral-trec	or 20 my	in	S	W. Indies	1690.	S	r.m	Com. h. 1.106
19519 enneandria Dec.	mine-anthered						S	r.m	Jac. sc. 4.46†
velutina Jac.									
19520 macronyilia Dec.	long-leaved	• spl 20	—	S	Teneriff	1822.	S	r.m	
19521 muis Jac.	...old	(~) spl 20	...	S	Caracas	179a	S	r.m	Jac.sc. 216
19522 piscidina Ades Hart.	Piscidia-hke	(~) spl 15	...	S	E. Indies	1800.	S	r.m	
19583 abyssinica Lam.	Abyssinian	(~) spl 10	...	S	Abyssinia	1820.	S	r.m	
19524 incina W. en.	hoary	(~) spl 10	...	S	E. Indies	182a	S	r.m	
19525 portoricensis Desf.	Porto Rico	1 or 10	...	S	Porto Rico	1800.	S	r.m	
19526 poi&nthes Brot.	Grass-flowered	• spl 20 ja.f	—	S	S. Amer.	1820.	S	r.m	Lin. tr. 14.11
19527 secundiflora Brot.	side-flowering	• spl 20	—	S	Brazil	1820.	S	r.m	Lin. tr. 14.12
19528 speciosa Andr.	showy	• or 10 au.o	—	S	W. Indies	1805.	S	r.m	Bot rep. 4«
19529 rubrinervia H. & B.	red-nerved	—	—	S	Bogota	1821.	S	r.m	
19530 ffligena Hart.	shining	—	—	S	E. Indies	1801.	S	r.m	
19531 aculeatissima Desf.	prickliest	—	—	S	—	1800.	S	r.m	
19532 umbrbsa H.8f.B.	shady	—	—	S	Caraccas	1817.	S	r.m	
arborea Hart.									
19534 arborensis JIor.	Caffrarian				C. G. H.	1816.	S	Lp	Bot.reg.736
19535 Indica Lam.	arborescent				E. Indies	1818.	S	r.m	Rox. cor. 3.219
19536 stricta Box.	Indian				E. Indies	1814.	S	r.m	R.mal.ti.7
19537 ovalifolia Box.	strict				E. Indies	1816.	S	r.m	
19538 ovalis Wat.	oval-leaved				E. Indies	1816.	S	r.m	
19539 peta L.	oval				Nepal	1820.	S	Lp	
19540 spathacea Dec.	painted				E. Indies	1696.	S	r.m	Ru. am. 2.77
19541 nberosa Rox.	spathaceous				W. Indies	1824.	S	r.m	
19542 velutina W.	Cork-barked				E. Indies	1816.	S	r.m	
19543 glifica W.	velvety				Caraccas	1810.	S	r.m	
19544 flosca Lou.	glaucous				Caraccas	1819.	S	r.m	
19545 Crista-gaUi L.	brown-flowered				E. Indies	1800.	C	Lp	Ru.am.2.78
19546 feurifolia Jac	Cock's-Comb				Brazil	1771.	S	r.m	Sw.fl.gar.214
Laurel-leaved					& Amer.	1800.	S	r.m	Sw. fl. gar. 142
2163. RUDO'LPHM W. RUDOLPHIA. (IV. J. H. Rudolph, a botanist of Jena.) Leg. Pap. Phasebl 2. - 4.									
19547 dflbiaHum.	doubtful	. D « 6	...	3	Havannahl	815.	C	s.l	
Glycine sagittata Hum.									
19548 rosea Tut.	roseate	L Q o r 6	...	R	W. Indies	1826	C	s.l	PI. am. 102.1 ?
2164. BITTED Rox.	BUTRA.	(John. Earl of Bute, a munificent patron of botany.) Leg. Pap. Vie. 3.							
19549 frondbsa Rox.	leafy	t DM 30	...	S	ETIndies	179a	C	r.l	Rox. cor. 1.21
19550 superba Box.	superb	{ 1 spl 30	—	S	EEIIndies	179a	C	r.l	Rox. cor. 1.22
19551 parviflora Box.	small-flowered	or 20	—	S	Coroman.	1818.	C	r.l	
2165. PONGSMIA Ven.	PONGAMIA.	(Pongam, its Indian name.) Leg. Pap. Dalb.	4. - &						
19552 glabra Ven.	smooth-irram^f	J Q o r 30	...	W	E. Indies	1G99.	C	s.l	Ven.maL2&
Dalbergia arbdrea W.	Robinia muis L.								
19553 uliginosa Dec.	marsh	± CD or 90	...	W	W. Indies	1824.	C	s.l	
Galedipa uliginosa W.	Robinia uliginosa W.								
19554 grandi flora Ven.	large-flowered	f • or SO	...	IV	E. Indies	181&	C	s.l	
Piscidia Box.	Piscidia^itc	\$ • or 20	...	W	E. Indies	1818,	C	s.l	
8166. DALBRRGGY Rox.	DALBRRGIA.	Sichou Dalberg, a Swedish botanist Leg. Pap. Dalb.	19.—go						
19556 ougetensis Box.	Ougein	• op. 41	—	W	E. Indies	1850.	C	s.l	
19557 latifolia Rox.	broad-leaved	• or 30	...	W	E. Indies	1811.	C	s.l	Rox. cor. 2.113
19558 Sissoo Rox.	Sissoo	• or 30	...	W	E. Indies	1820.	C	s.l	
19559 Gom	Crowe's	» or 20	...	W	E. Indies	1823.	C	s.l	
uhimgJ^ai	rusty	» or 10	...	W	E. Indies	1811.	C	s.l	
robfisi Rox.	robust	f • or 20	...	W	E. Indies	1816.	C	s.l	Rox. cor. 9.115

- 19562 alkta *Box.* winged W E. Indies 1823. C s.l
 19563 cmarginata *Box.* emarginate W E. Indies 1823. C s.l
 19564 scandens *Box.* climbing R E. Indies 1812. C s.l Rox. cor. 2.192
 19565 vol&obilis *Box.* twining W E. Indies 1818. C s.l Rox. cor. 2.191
 19566 marginata *Box.* marginated W E. Indies 1823. C s.l
 19567 rimtoa *Box.* chinky W E. Indies 1823. C s.l
 19568 Barcfoyi Hook. Barclay's B Mauritius 1823. C s.l Hook. ex. fl. 188
 19569 Telfairfi Hook. Telfair's W Mauritius 182a C s.l
 19570 frondosa *Box.* frondose 'W E. Indies 1818. C s.l
 19571 paniculata *Box.* panicled W E. Indies 1811. C s.l Roz. cor. 2.114
 19572 stipulacea *Box.* stipulaceous W E. Indies 1820. C s.l
 19573 timonensis *Dee.* Timor W E. Indies 1826. C s.l R.maL.6.22
 19574 tamarmidifbha *Box.* Tamarind-lvd W E. Indies 1820. C s.l
2167. DREPANOCA'RPUIS Mey. DREPANOCARPUS. (*Drepanon*, sickle, *karpas*, fruit) Leg. Pop. Dobl. 1.-6.
 19575 lunatus Mey. half-moon-capsuled* £3 or 12 ... W a Amer. 1792. C sj PL am. 201.2 Pterocarpus lunatus L.
2168. ECASTAPHYIJ-yMtfr, ECABTAPHTIXUM. (*Hekastos*, each./Ag/lon, leaf: simple.) Leg. Pap. Dobl. 3.-7.
 19576 Br6weui Pers. Browne's A • or 10 ... W W. Indies 1733. C r.m Br. jam. 32.1 Pterocarpus Ecastaphyllum L.
- 19577 Sidberi Rdth.- Sieber's « Q or 10 ... W. IR Guinea 1824. C s.l
 19578 PlumieriPen. Plumier's u m or io - W S. Amer. 1820. C s.l P.ed.B.246.2
2169. A'RACHISL. EARTH NUT. 'A, without, *rhalris*, branch.) Leg. toes. Geafft 1.—2.
 19579 hypogs^a L. underground O clt 2 my.jn Y S. Amer. 1712. S s.l Tr. pi. 3.3
- 2170. VOANDZPIA Thou. VOANDZBIA. (*Voandzou*, its name in Madagascar.) Leg. Ctes. Geoff: 1.
 19580 subterranea Thou. subterranea-mi %, O clt ± jn.au Y Africa 1823. S r.m Declegij.20.10? [and L.tdec ? Gl^cine subterranea X.
2171. ANDPBA Lam. ANDIRA. (Its name in Brazil.) Leg. Cats. Geoff: 2. — 6.
 19581 racemua Lam. racemose J • or 20 ... P Trinidad 1818. C Lp Pis. br. 2 Geotrccevi raccratasa-Poir.
 19582 inermis H. % 3. unarmed J Om 20 - W. Indies 1773. C Lp Ph.tr. 1777.10 Geoffhe^a iner'rmls Swz.
2172. GEOFFRCYJ W. BAST. CABBAGE TR. (E.F. Geffroy, F.R.S.prof.botParis.) Leg. Ca^s. Geaff. 2—6.
 19583 spinosa Jae. spiny J D o r 30 ... Y S. Amer. 1818. C l.p Jac. am. 180.62
 19584 violacea Pert. violet \ O or 20 ... V Guiana 1823. C l.p Aub. gui. 3.301
- 217a PABIVO^A Aub. PARIVOA. (Its name in Guiana.) Leg. Ctes. Catnea. 1.
 19585 grandiflora Aub. large-flowered 2 O or 30 ... P Guiana 1824. C r.m Aub. gui. 303 Dimorpha grandiflora W.
2174. DI'PTERIX Schreb. TONQUIN BEAN. (*Dis*, double, *pterix*, wing j app. of cal.) Leg. Ca^s. Geaff. 1—SL
 19586 odorata W. sweet-scented J D e c f l O ... P Guiana 1793. C Lp Aub. 8uL2.896 Bary6sma T6ngo Gae.

CLASS XVIII. ORDER 1.

POLYADE'LPHIA DECA'NDRIA.

2175. THEOBRO^MA L. CHOCOLATE NUT. (*Theos*, a god, *brome*, food; delicious.) Byttneribctce. 3.-5.
 19587 bicolor H. \$ B. two-colored f O clt 16 ... Br NewGran. 1820. C r.m H. & B. 1. 30
 19588 Cacao L. common Cacao J • clt 16 ... Br S. Amer. 1739. C r.m Bot. cab. 54/
 19589 guianensis W. Guiana J • or 16 ... Br Guiana 1803. C r.m Aub. gui. 2.275
2176. GUAZU^MA Tin. BASTARD CEDAH. (Its name in Mexico.) Byttnerihctee. a
 19590 flrimifolia Lam. Elm-leaved J O t m 4 0 a.u.s Y Jamaica 1739. C p.l Tr. chr. 76
 Bubrdma Guaxiana Vf. Theobrbma Guaziana f.
 19591 polyb'trya Cav. many-racemed ± CD or 20 - - Brazil 1816. C p.l Cav. ic. 3.299
 Bubr'tma polyb6tryum W. en.
 19592 tomentbsa Kth. wooy f CD or 20 - - Cumana 1820. C p.l
8177. DU'BIO I* DURIOS. (*Durya* a thorn, *Malay*; spinous fruit) Malvdeca?. 1.
 19593 zebethinus L. civet J D o r 60 ... W £ Indies 1825. C s.l Ru.am. L29
2178. ABRCMPA L. ABROMA. (^privative, *brome*, food; unfit for.) Byttneriiicear. 2. - 3.
 19594 augusta L. August smooth-italkedf O or 10 au P E. Indies 1770. C l.p Jac. vin. 3.1
 19595 fastubsa A Br. di8dainlul/>r.-<ttd j Q or 10 jn.o P N.S.W. 1800. C Lp Par.loal02

CLASS XVIII. ORDKB. 2.

POLYADE'LPHIA POLYA'NDRIA.

*2179. MELALEUCA L. MELALBUCA. (Afc&w, black, taifow, white; bk. wood, *, branches.) *Myrticear.* 48.

		I. ALTERNIFOLUE.—Leaves alternate		W.		E. Indies		1796.		C	
19596	Leucadendron <i>L.</i>	White Tree.	J or 15	J!	...	R Indies	1800.	C	s.l.p	Ru. am.	2.16
19597	<i>Cajuputi</i> Rox.	Cajuputi Tree	J or 15	J!	...	N. S. W.	1798.	C	s.L _p	Ru.am*	2.17.1
19598	<i>vindiflora</i> Sm.	green-flowered	L-J or 10	G	...	N. Holl.	1803.	C	s.l.p	Cav.ic.4.333	
19599	paludosa R. Br.	marsh	L-J or 6	JL	R	N. Holl.	1803.	C	s.l.p		
19600	<i>globiflora</i> R. Br.	globe-bearing	L-J or	J	...	N.Holl	1803.	C	s.l.p		
19601	<i>diosmochilia</i> Andy.	Diosma-leaved	L-J or	jn.jl	G	N. Holl	1794.	C	s.l.p	Bot rep. 476	
19602	<i>styphelioides</i> Sm.	Styphelia-lvd	LJOT	my.jn	...	N.S.W.	1793.	C	fcl.p		
TX*13	<i>gnatostylis</i> Sm.	Broom-leaved	t-J or	J	...	N.S.W.	1793.	C	s.l.p		
19604	Tanceolata Otto	lance-leaved	i*(J or	J	...	N Holl	1816.	C	s.l.p	H. ber. 36	
11605	<i>striata</i> Lab.	striated	ii(J or	J	...	N.Holl	1803.	C	s.l.p	LpLab.n.h.2.165	
19606	<i>hymoides</i> Lab.	Thyme-like-/imf	i J or	3	P	N.Holl	1803.	C	s.l.p	Lab. n. h. S. 167	
19607	imbricate Lk.	imbricated	iM J or	4	jn.au	P	N.Holl	1803.	C	Blp	
19608	<i>aquimca</i> La.*	BcaXy-branched	L-J or	4	jn.jl	...	N. Holl.	1800.	C	s.l.p	Botreg.477
19609	<i>taxifolia</i> Schl.	Yew-leaved	L-J or	J	...	V.D. L.	1805.	C	s.l.p		
19510	<i>pentagyna</i> La6.	five-angled	i T or	J	...	N. Holl	1820.	C	s.l.p		
19611	<i>nodosia</i> Sm.	knotty	L-J or	2	jn.jT	M. Holl.	1820.	L	s.l.p	pEx.bot.1.35	
19612	<i>ericifolia</i> 5m.	Heath-leaved	L-J or	3	jl.8	P	N. S. W.	1790.	L	s.l.p	
19613	<i>armillaris</i> Sm.	bracelet	L-J or	2	jn.jl	G	N. S. W.	1788.	L	s.l.p	Ex. bot 1.34
19614	<i>uncinata</i> R. Br.	hook-leaved	L-J or	3	jn.s	G	N. S. W.	1788.	L	s.l.p	Bot rep. 175
19615	<i>scabra</i> R.Br.	rough-leaved	L-J or	3	l.8	P	N. Holl.	1803.	C	8.l.p	
19616	<i>pulchella</i> ii. & r.	neat	L-J or	2	jn.s	P	K. Holl.	1803.	C	s.l.p	
					P	N. Holl.	1803.	C	s.l.p	Bot cab. 200	
								C	Blp	Bot reg. 410	
II. OppositiFbLiE.—Leave* opposite or verticillate.											
19617	incana it. Br.	hoary	L-J or	3	jn.au	Y N.	Holl	1817.	C	Blp	
	canfocena L.										
19618	/hyrritifolia Sm.	Thyme-leaved	L-J or	2	jn.8	J	N.S.W.	1792.	C	s.l.p	Bot mag. 1868
	coronata Aruri.										
19619	eriocéphala Sib.	woolly-headed	L-J or	4	...	J	N. Holl	1824.	C	s.l.p	
19620	decausata R. Br.	decussate	L-J or	4	jl.s	J	N. Holl	1803.	C	s.l.p	Bot mag. 2268
19621	rotundifolia Hort.	lound-leaved	L-J or	3	jn.au	J	N. Holl	1816.	C	s.l.p	
19622	<i>fulgens</i> R. Br.	splendid	L-J or	6	jls	J	N. Holl.	1803.	C	a.l.p	Botreg. 103
19623	discolor Rchb.	various-colored	L-J or	3	...	J	N. Holl.	...	C	s.l.p	
19624	<i>ltmaritbia</i> R. Br.	Toadflax-lvd	L-J or	3	jn.au	J	N.S.W.	1793.	C	s.l.p	Ex. bot L 56
Iff125	<i>cuticularis</i> Lab.	thin-skinned	L-J or	3	...	J	N.Hol	...	C	s.l.p	
19626	<i>gibbosa</i> Lab.	gibbous	L-J or	3	...	J	N.Hol	1820.	C	s.l.p	
h 19627	<i>fascicularis</i> Lab.	fascicled	L-J or	3	...	J	N. Holl	...	C	s.l.p	
19628	<i>pendulina</i> Lod.	pendulous	*L-J or	3	...	J	N.Hol	1820.	C	s.l.p	
19629	<i>trincrvia</i> Sm.	three-nerved	L-J or	3	...	J	N.Hol	181&	C	s.l.p	
PHRO	<i>fimbriata</i> Hort.	fringed	L-J or	3	...	J	N.Hol	1817.	C	s.l.p	
Iff131	<i>grandis</i> Fts.	grand	L-J or	4	...	J	N.Hol	...	C	s.l.p	
1WJ12	<i>dumosa</i> Lod.	bushy	L-J or	2	...	J	N.Hol	...	C	Blp	
I J131	<i>cpacidea</i> Hort.	Epacris-like	*J or	6	...	J	N.Hol	1820.	C	s.l.p	
1.6.14	euphorbioides Lod.	Euphorbia-like	HJ or	3	...	J	N.Hol	1824.	C	s.l.p	
Ijj&J131	<i>virgata</i> Hort.	twiggy	JL J or	5	...	J	N. Holl	181&	C	s.l.p	
**to13	<i>tenuifolia</i> Ma. C.	fine-leaved	<L-J or	2	jn.au	J	N. Holl	1824.	C	s.l.p	
1%37	<i>Apicerifolia</i> Sm.	Hypocricum-lvdM	L-J or	3	jn.au	J	N.S.W.	1792.	C	s.l.p	Bot. rep. 500
19UJ18	<i>squarbaa</i> Sm.	squareose	>L J or	2	J	S	N. S. W.	1794.	C	s.l.p	Bot. mag. 1935
I'VU9	<i>tetragona</i> Hort.	four-angled	*J or	4	...	W	N. S. W.	1820.	C	Blp	
19640	<i>calycina</i> R.Br.	permanent-calyxeA	*J or	3	jn.au	J	N. Holl.	1803.	C	Blp	
19>41	parviflora Lk.	small-flowered	HJ or	3	jn.au	J	N. Holl.	1816.	C	s.l.pH.bcr.S7	
19642	<i>densa</i> R. Br.	dense-leaved	*J or	2	...	J	N. Holl.	1803.	C	s.l.p	
19643	<i>tomentosa</i> Coil.	woolly	L-J or	3	jn.au	Crea	N. Holl.	1810.	C	s.l.p	Col.h. rip.37

* 2180. TRISTA^NI A R. Br.	TRISTANIA. (<i>Treis</i> , three, <i>stao</i> , to stand; dispos. of flws and lvs.)	<i>Myrrhccce</i> . 6L
I <illincre <i>ñ</i> lului R. Br.	Oleander-leaved	* _Jor 6 jns. Y N. S.W. 1804. C s.p Bot mag. 1058
i. i. i. >*ri <i>cif61ia</i> Cun.	Peach-leaved	* t_Jor 6 . . . N. Holl. 1824. C s.p
i.ulii.irholscsns Hort.	arboreocent	f__Jor 10. . . N.HoL 1820. C s.p
Jk\>7/ <i>sunna</i> R. Br.	Laurel-leaved	i j o r 6 . . . Y N. S. W. 1798. C s.p
WA8confstr ViR.Br.	crowded	* t_J or 6 jl.s Y N. S. W. 1805. C s.p
19649 depresso <i>Lod.</i>	depressed	H _Jor 6 . . . N. Holl, 1820. C s.p

2181. CALOTHATVINUS Za&. CALOTHAMNUB. (<i>Kalos</i> , beautiful, <i>thummos</i> , shrub.)	<i>Myrthcete</i> . 4.—
19650 quadrifida R. Br.	four-cleft • <u>I</u> or 3 <u>jl.a'</u> S N. Holl 1803. C s.p Bot mag. 1506
19651 villbsa R. Br.	villus ftilor 3 <u>jl.8</u> S N. H. ^a L 1803. C ap Bot reg. 1099 l<Hi>2 clavday Ma. C. clubJeaved • <u>L</u> or 2 <u>JLs</u> S N. Holl 1824. C s.p
19653 gracihs R. Br.	*)endcr-leaved * <u>L</u> or 3 <u>jl.s</u> S N.Holl 1803. C s.p

<i>BEAUFORTIA.</i>	(<i>Mary, Duchess qf Beaufort, hot pat</i>)	<i>Myrtaceous.</i>	&
<i>i. L-TU88ta R. Br.</i>	<i>decussated</i>	<i>i. J. spl</i>	<i>S. N. Holl. 1803. C s.p.</i>
<i>t.,,, ^pdfrsa R. Br.</i>	<i>scattered-leaved</i>	<i>U spl</i>	<i>... R. N. Holl. 1803. C s.p.</i>
<i>I>j6-keearinata Cun.</i>	<i>keel-leaved</i>	<i>»t. J. spl</i>	<i>3 ... S. N. Holl. 182a C s.p.</i>

2183. *m̄hL̄TTJ̄ Coll.* BILLOTIA. {*Mad. Teciphila Billoti*, of Turin.) Myrtoce*. 1
19657 *ncer̄tosa Coll.* Maple-tamoi ■ L Jor 5 JLs R N.HoLL 1816. C s.p CoLh.rip.fi3

2184. EUDE'SMIXR.Br. EUDESMEA. (*Eu*, well, *desme*, bundle; stamens.) *Myrtacete.* 1.
19658 *tetragona* ii. Br. four-angled **J**l_Jor 16 jl W N.Hol 1824. C s.p Sw.au.21

2185. CANDO'LLÉ! Lab. CANDOLLEA. (*Aug.Py.de Candolle*, F.R.S. F.M.L.S. Geneva.) *Dilleniacea*; 1

-2486 SYMPLLOCOSi. SYMPLOCOS. {*Symploke*, connection; dyer's *LaureUvd**. L or 3 ... Y Carolina 1780. L pl. Cat. car. 1, 154
timoraeaff. Chinese. * L or 2 ... Y China 1780. L pl. Cat. car. 710
varicosa Kar.

19662 coecfina Kth. scarlet ft — por — — Mexico 1825. H. & D. 1.52
 19663 cratsegi Adcs D. Don Crataegus-like ft uJ or — — Nepal 1824.

2187. CITRUS L.	ORANGE TREE.	(Derivation unknown.)	Aurantidcea?	14.—15.	
19664 A/edica Rizzo	Median Uman ft i I ft 8 myjl	W Asia	... B	r.m Fer. hes. 39	
19665 Limetta Rizzo	Lemon Bergamot ft i I ft 8 myjl	W Asia	1648. B	r.m Black. 362	
19666 Limonum Rizzo	Lime ft i I ft 15 myjl	W Asia	1648. B	r.m Gae.tr. 2.121.2	
19667 Aurantium Rizzo	sweet Orange.* ft i I ft 15 myjl	W Asia	1595. B	r.m Lam. il 639.2	
19668 vulgiins Rizzo	common Seville ft i I fr 15 myjl	W Asia	... B	r.m "to"	
2 myrtifolia Hort.	Myrtle-leaved a I—fr 3 myjl	W Asia	... B	r.m Bot reg. 346	
19669 decumana L.	huge Shaddock < I J fr 15 myjl	W India	1724. B	r.m Ru. am. 2 24.2	
19670 hirstrix Dec.	porcupine a uJ fr 15 myjl	W E. Indies	... B	r.m Ru.am.2.28?	
19671 spinosissima Afey.	splinters IMU	T LJfr 15 myjl	Cayenne	... B r.m	
19672 japatrica TAuff.	Japan small-fruited LJfr 6 myjl	W Japan	... B	r.m Thun. jap. 15	
19673 nobilis Low.	noble Afamfarotil I J fr 15 myjl	W China	1805. B	r.m Bot. rep 608	
2 minor	smaller ft! LJ fr 15 myjl	W China	1805. B	r.m Botreg. 2.11	
19674 margarita ZOK.	pearl sweet Lemon \$ LJfr 15 myjl	W China	... B	r.m	
19875 angulata W.	angular-/rwii*? Lj fr 15 myjl	W E. Indies	... B	r.m Ru. am. 2.32	
19676 madurensis IMU.	Madura \$ i J fr 10 myjl	W China	... B	r.m Itu. am. 2.31	
19677 Auxifolia Poir.	Box-leaved	Jt	3 myjl	W China	... B

2188. XANTHOCHTPMUS Box.	XANTHOCHYmus.	(Xanthos, yellow, chymos, juice; fruit.)	Guttifree.	8.
19678 pictibius Rox.	painter's f LJ fr 20	mmm y	E. Indies	1796. S r.m Box. cor. 2.19ti
19679 ovalifolius Rox.	oval-leaved + LJ fr 12	... T	E. Indies	IH24. S r.m
19680 I&teus IM.	yellow	... Y	E. Indies?	1824. S r.m
19681 macroph^llus Lo. C.	long-leaved (ZKr 20	... Y	E. Indies?	1824. S r.m
19682 purpureus Lo. C.	purp>le Qfr 20	... Y	E. Indies?	1824. S r.m
19683 guineensis G.Don	Guinea fr 20	... Y	Guinea	1824. S r.m
19684 finetdrius Rox.	dyer's E fr 20	... Y	E. Indies	1818. S r.m
19685 dulcis Rox.	sweet fr 20	... Y	E. Indies	1820. S r.m

8189. PENTADE'SMA R. Br. PENTAOESMA. (Pente, five, desme, bundle; stamens.) Guttifer*. 1.
 19686 butyracea R.Br. butter t/tattoo.tree± O ft 30 n ... a Leone 1822. S s.1

• 2190. HYPERICUM L.	ST. JOHN'S WORT.	(Origin unknown.)	Hyperichne.	77.—138.
L ABCYRB^IA.—Sepals at base joined and unequal; stamens numerous; leaves often large; flowers com*	monly terminal, large, and few; styles commonly three, from No. 19696. styles commonly five.			
19687 elatum H. K.	tall ft or 5jl.au	Y	N. Amer. 1762. L.1.1	Den.br. 85
19688 frondosum Mx.	leafy	Y	NAmer. 1806. C	
19689 grandifolium Choi.	great-leaved	Y	Teneriffe 1818. C	Chois. hyp. 3
19690 amcB^um Ph.	pleasing	Y	Carolina 1812. L s.1	l)i. cl. 151.182
19691 hircinum L.	goat-scented	Y	S. Europe 1640. S	8.1 Den. br. 86
2 minus	smaller	Y	S. Europe ...	L co Den. br. 87
19692 foliosum H. K.	sfuivngAeafy	Y	Azores 1778. C p.l	
19693 flonbundum H. K.	bundle-flwd	Y	Madeira 1779. C p.l	Com. h. 2.68
19694 ol^mpicum L.	Olympian	Y	Levant 1706. S s.1	Bot. mag. 1867
19695 cananense L.	Canary	Y	Canaries 1699. C p.l	Bot. cab. 953
19696 chinense L.	Chinese	Y	China 1753. C p.l	Bot. mag. 334
monogynum L.				
19697 cordifolium Choi.	heart-leaved	Y	Nepal 1825. C co	
19698 pyramidalum H. K.	pyramidal	Y	Canada 1759. D p.l	Vcn.mal.118
19699 ^scyon L.	St. Peter's wort	Y	Siberia 1774. Skco	Gm. 6i. L
19700 ascyron^des W.	Ascyron-hke	jnjl	N. Amer. 1812. Skco	
19701 patum Thun.	spreading	jnjl	Nepal 1823. C co	Thun. jap. 295
19702 uralum Ham.	Ural	jnjl	Nepal 1823. C s.1	Bot mag. 2375
19703 oblongifolium Choi.	oblong-leaved	Hi	N. Amer. 1759. C s.1	Chois. hyp. 4
19704 Kalmiunum Lam.	Kalm's	Y	Ireland Skco	Eng. bot. 2017
19705 calycinum I*	Jary-calyxed	jns	Majorca 1714. C r.m	Bot. mag. 137
19706 balearicum L.	Balearic	1) mr.s	China 1821. C co	
19707 cochinchinense Zofi.	Cochinchinese	3 jl.au		

II. ELODEA.—Calyx five-sepaled; sepals equal, entire; styles three; filaments numerous, nine, fifteen, or eighteen; herbs purplish; flowers axillary, or crowded terminal.

19708 pauidsum Choi.	marsh	4 A pr 2 jl.au	K	N. Amer. 1821. D co
19709 virginicum L.	Virginian	S A or Ijl.Ls	R	N. Amer. 1800. D p.l Bot rep. 552

Elodda campanulata Ph.

III. PERFOR^UIA.—Cat. 5-sepaled; scp. equal; stain, numerous*; styles commonly three; herbs or subshrubs; flowers axillary or pascled; leaves rarely linear; sepals entire, from No. 19740. toothed glandular.				
19710 angustatum Mx.	&ngu^BC-toot/icd-fb	A — jnjl	Y	N. Amer. 1812. D p.l Pluk.al.245.6
19711 punctatoid Lam.	dotted	ft	Y	N. Amer. 1823. 1) co
19712 dolabriflume Ven.	hatchet-formed	A or 2 jnjl	Y	N. Amer. 1821. D co
19713 procumbens Mr.	procumbent	A or 1 au8	Y	N. Amer. 1822. D co
19714 rosmariniflbum Lam.	Rosemary-lvd	J or 2 jn.au	Y	Carolina 1812. L s.1
19715 virgatum Lam.	twig^y k. A or Ujn.au	Y	N. Amer. 1820. D co	
19716 wiryfolium Lam.	Myrtle-leaved ^t A or 1 Jl.au	Y	N. Amer. 1818. D co	
19717 prolificum L.	prolific	or 4 jn.au	Y	N. Amer. 1758. S 8.1 Den. br. 88
19718 glaucum Mx.	glaucous	ft J or U jl.au	Y	N. Amer. 1812. C p.l
19719 lasivagatum H. K.	smooth	£ A or U jl.au	Y	N. Amer. 1772. D p.l
19720 nudiflorum Mx.	naked-flowered	ft J or Ili.O	Y	N. Amer. 1811. C p.l
19721 quadrangulum L.	square-stalked	^t A or U jl.au	Y	Britain m. me. C p.l Eng. bot. 370
19722 d&blum W.	doubtful	A or 3 jl.au	Y	Britain m. thi. C p.l Eng. bot 296
19723 maculatum All.	spotted	2 jl.au	Y	N. Amer. 1789. C p.l
19724 unduatum Schous.	wave-leaved	1 jl.au	Y	Barbary 1802. D p.l
19725 veronense Spr.	Veronese	1 jl.au	Y	Europe 1816. D co
19726 attenuatum Choi.	attenuated-^-	U jl.au	Y	Dahuria 1822. D p.l
19727 ja Mnicum Thun.	Japanese	T jl.au	Y	Nepal 1K23. D p.l
19728 simplex Mx.	simple	jl.au	Y	N. Amer. 1826. ■■■ Boc. mu. 12
19729 crfepum L.	curly-leaved	5t 1 jl.au	Y	Greece 1688. ■■■ Boc. mu. 12
19730 setibsum H. K.	bilby, burbanknickel	J or jl.au	Y	Carolina 17K. D p.l
19731 heterophyllum Ven.	various-leaved	J or jl.au	Y	Persia 1H2. D p.l Ven. cel<G8
19732 acgyptiacum L.	Egyptian	i or injl	Y	Egypt 1787. C p.l Bot reg. 19*
19733 humiflora L.	trailing	Jfc A or ijLau	Y	Britain pas. D co Eng. bot. 1226
19734 Liott&rd Vil.	Liottard's	Jt Q or Jn.au	Y	Switzerl. 1819. D co Vil. del. 44

19735 pusillum Choi.	small	4 jn.au	Y	N. S.W.	1818.	D	s.1	Lab.n.h.2.175
19736 uivolutum Choi.	involted	5 jn.au	Y	N. S.W.	1822.	D	BI	Lab. n. h. 8.174
19737 per foratum L.	bcrforated	6 li.lau	Y	Britain	bu.pL	D	p.l	Eng.bot.295
19738 quinqueñervium Wall.	five-nerved	7 1 in.au	Y	N. Amer.	1759.	D	p.)	
19739 canacWnal.	Canadian	8 1)U	Y	N. Amer.	1770.	D	Pi	
19740 elbdea L.	marsh	9 ij.lau	Y	Britain	sp. bo.	D	p.l	Eng.bot 109
19741 toinqlbussum L.	woolly	10 1	Y	& Europe	1648.	D	r.m	
19742 hirsutuin L.	hairy	11 mil	Y	Britain	ch.ba.	D	p.l	Eng.bot. 1156
19743 nummularium L.	MonyworWvd.	12	Y	a Europe	1823.	D	co	Lam. il. 643
19744 elcgans Step.	elegant	13	Y	Siberia	1822.	D	co	Spr. haL 9
KohHitom Spr.								
19745 glandulbum H. K.	glandulose	14	2 my.au	Y	Madeira	1777.	C	p.l
19746 reflekuin L.	xetVavcd	15	11 jns	Y	Tenerife	1778.	C	p.l
19747 ptlchruiin L.	fair	16	ijjl	Y	Britain	woods.	D	p.l
19748 nerviflum D. Don	nervous	17	11	Su	Nepal	1820.	D	Eng.bot.1227
elodeoides Choi.								
19749 barbtum L.	bearded	18	2 jn.o	T	Scotland	sc. thi.	D	co
2 calabricum Spr.	Calabrian	19	Ujn.o		Calabria	1816.	D	co
19750 dentatum Lot.	toothed	20	2 m.o"		Mcditerr.	1890.	D	Loi.gal.17
19751 montanum L.	mountain	21	ljil.lau		Britain	m.wo.	D	Eng. pol. 371
19752 fimbriatum Lam.	fringed	22	2 ilau		Pyrenees	1821.	D	Vil.del.44
19753 dlynum IV. &K.	alpine	23	2 jl.au		Hungary	WS&.	D	W.&K.265
19754 sthiopicum Thun.	Ethiopian	24	1 jn.au		C. G. H.	1817.	C	p.l
19755 pcrfotatum L.	Kirfoliate	25	1 jl.au		Italy	1785.	D	p.l
<erpylliflium Lam.	Thyme-leaved	26	* 11 au		Levant	1688.	C	M.h.2.5.6.2
19756 cihamom Lam.	tuige-flowered	27	2 11		Levant	1739.	D	l.m
19758 triplindry Yen.	three-nerved	28	A P'		N. Amer.	1821.	D	Boc. mu. 127
19759 Ayssopifolium Fil.	Hyssop-leaved	29	A P'		S. Europe	1823.	D	Ven. eels 58
19761 empctribUum W.	Emptetrum-lvd	30	1 pr		S. Europe	1820.	C	Vil.del.44
19762 Chris L.	Conn-leaved	31	1 pr		Levant	1640.	C	Den. or. 141
19763 encolade L.	Heath-like	32	1 jn.jl		Spain	1821.	C	Bot mag. 178
aspalathoides W.	fascicled	33	1 jn.au		Carolina	1811.	C	Cav. ic. 122

2191. HARO'NGA Thou. HAROVGA. (Its name in Madagascar.) 1—5.
19764. madagascariensis Choi. Madagascar i or 10 ... Y Madagascar C p.l
paniculata Lo. C.

*2192. VI'SMEil Van.	VISMEA.	(M. de Visme, a Lisbon merchant)		Hwericinac.	2.—17.
19765 guianensis Per*.	Guiana wax tree*	EU or 8 jl.o	Y	Guiana	1824. C 8.1 Aub.gui.2.311
hypencum guianense Aub.					
19766 brasiliensis CAo.	Brazilian wax tree*	£3 or 8 jl.o	Y	Brasil	1824. C 8.1 Choi*, hyp. 2
* 2193. LOJCSA Adan.	LOASA.	(Meaning unknown.)		Lodsea.	5.—10.
19767 Place* Lindl.	Place's	O or 4 jns	Y	Chile	1822. S co Botreg.785
acanthifolia R. B.					
19768 nltida Lam.	shining	O or 2 jns	Y	Chile	1822. S co Bot reg. 667
tricolor B. R.					
19769 volubilis J.	twining	iQJel	1 i mr.s	Chile	1824. S s.1 An. mu.5.2
19770 grandiflbra Lam.	large-flowered	iQor	2 1	Caraccas	1825. S co An. mu. 5.4
19771 patula Grah.	spreading	O or 1 jUu	Y	Chile?	1827. S s.1

2194. BLUMENBA^CHJ/f Schr. BLUMBNBACIIA. (J. F. Blumenbach, F.R.S. prof. med. Gottingen.) Loöse*. L
19772 insignis Schr. remarkable -* O or £jl.n W Monte VL1826. S r.m Sw.fl.gar.170
Lohat palmata Spr.

*2195. SCYPHA'NTHUS Swt. SCVPIANTHUS. (Skyphos, a cup, anthas, a flower.) Loitsea?. 1.
19773 grandiflbris SIM/, great-flowered £ O «f 2 aus Y Chile 1824. S B.1 Sw.fl.gar.238
L. vobtibB Hort.

CLASS XIX. OBDBR 1.

SYNGENE^VSIA *equivalis*

2196. GEROPO%GON L.	OLD MAN'S BEARD. (Germ, old man, pogon, beard; of seeds.) Comp. Cich. 3.-6.
19774 glaber.	smooth
19775 hirs&tus L.	hirsute
19776 calycifatus Jac.	calyxed

2197. rR AGOPO ^X GON L.	GOAT'S BEARD. (Tragos, goat, pogon, beard; of seeds.) Comp. Cich. IS.—17.
I. SUPERA'NTBB.—Florets exceeding the calyx.	
19777 canus W. **.	hoary
19778 angustiflbiit Bel.	narrow-leaved
19779 orientalis L.	oriental
19780 flocebus W. % K.	woolly
19781 roseus Treu.	roseate
19782 pustulus Bicb.	gmal

II. JEQUA'NTES. — <i>Floret* equalling the calyx.</i>									
19783 <i>pratensis</i> L.	meadow	it	C>	r?	my.jn	Y	Britain	past	■ r.m Eng.bot. 43*
19784 <i>dubius</i> Horn.	doubtful	■	Q>	or	3	ray.jn	Pa.Y	Podolia	1818. ■ co
livescens Bes.									
19785 <i>mutibilis</i> Jac.	changeable	■	CD or	3	myjn	Pa	Siberia	1816. ■ ■	Jac. ic. 1157
19786 <i>undulatus</i> Jac.	■	■	or	2	myjn	W.Y	Crimea	1790. ■ ■	Jac. ic. 1158
III. MINORES. — <i>M-florets smaller than the calyx.</i>									
19787 <i>porriflius</i> X.	Leek-lvd	Salsify	Q>	cul	4	my.jn	P	England	m.me. S f.m Eng. bot 638
19788 <i>crocifolius</i> L.	Crocus-lvd	(jor	1	jnjl	P%	Italy	1739. S &o Col. c.c. 1.230		
19789 <i>villbsus</i> L.	villous	Q	or	4	myjn	P.Y	Spain	1794. S &e	
19790 <i>campstris</i> Bes.	field	5	Q	Q>fr	3	my.jn	Y	Podolia	1819. S co
19791 <i>nujor</i> Jac.	grater							Austria	1788. S co Jac. au. 1.29
2198. TRO'XIMON Gae.	Troxhnos	(<i>Troxhnos</i> , eatable^						Comp. Cichor.	2.—
19795 <i>glaticum</i> Ph.-	da	or	1	myjn	Y	Missouri	1811. D e o	Bot. mag	1667
19793 <i>cuspidatum</i> Ph.	cuspidate	■	or	1	jl.au	Y	Louisiana	1824. D co	
margmatum Nut.									
2199. ARNOPO'GON TV.	SHBEP'9 BEAHD.	(<i>Arnos</i> , lamb, pogpn,						Comp. Ciehor.	4
19794 <i>Dalechamp</i> */IV.	Urosphrniu	^, A-pr	2	jn.o	L.Y	S. Europe	1739. D e o	Bot. mag.	1623
Urosphrniu Dalechamp;/ Desf.									
19795 <i>filicroides</i> W.	Pi	ens-like	O	pr	1	jl.au	Y	S. Europe	1683. S co
19796 <i>asper</i> IV.	rough	O	pr	1	Ujl.au	Y	Montpel.	1774. S co	Larn.il 64d 3
19797 <i>capdnis</i> TV.	Cape	^	(J p t	1	jn.jl	Y	C. G. H.	1818. S co	Jac. ic. 3.577
Urosphrnm caplnse Spr.									
•2200. PODOSPE'RMUM Dec. PODOSPERMUM. (<i>Pous</i> , foot, sperma, seedy foot-stalked.) Comp. Ciehor. 7-11.									
19798 calcitrapifohum Dec.	Caltrop-lvd	■	■	■	■	■	Levant	1820. ■ co	Bux. c. 2.23
19799 ladniatum Dec.	lag-leaved	■	■	■	■	■	S Europe	ICAQ. ■ s.l	Jac. au. 4.356
19600 octangulare Dec.	octangular	■	■	■	■	■	S Europe	1818. ■ co	
19801 rescidif?ium Dec.	Beseda-leaved	■	■	■	■	■	S Europe	1818. ■ co	Boa sic. 7. fa. c.
19802 /araxaci folium Dec.	Dandclion-lvd	^	■	■	■	■	Bohemia	1820. ■ co	Jac. ic. 1.106
Scorzonira /araxaciflta IV. *									
19803 coronopifblum Deaf.	Buckhom-lvd	■	H	■	■	■	N. Africa	1818. ■ oo	Desf. at S. 212
19804 pamilum Cav.	dwarf	O	or	■	■	■	1816. ■ co	Cav. ic. 2.121. S	
•2201. SCORZON&RA L. VIPER*S GRASS. (&nir>m viper, Catalan; sup. med. qual) CSmjo.Cichor. 27.-35.									
19805 <i>tuberbsa</i> L.	tuberous	A A pr	*J"		Y	Volga	1825. D e o	Pal. it. ap. Y. 3	
19806 homilis L.	'humble	j^ pr	1	au	Y	Europe	1597. D co	Jac. au. 1.36	
auslrtaca W.									
•19807 tomentosa L.	tomentose	■	■	■	■	■	Armenia	1780. D co	
19808 crispa Bieb.	curled	■	■	■	■	■	Tauna	1817. S co	
19809 hispánica L.	Spanish	■	■	3	jn.s	Y	Spain	1576. D co	Lam.il. 617.5
19810 macrorhiza Schl.	long-rooted	■	■	■	1*jnjl	Y	Switzerl.	1819. S co	
19811 glastifolia IV.	Woad-leaved	■	■	■	jns	Y	Germany	1816. D co	
graininifbia Hofm.									
19812 cancifbia Pall.	Carex-leaved	^t A Pr	1 jn.s	Y	Siberia	1805. D co	Pal.it3.ap.J.I.I		
19813 purpurea L.	purple-Jiotvercd	ifc A Pr	2 myjn	P	Austria	1759. D co	Jac. au. L 33		
19814 rbsca Kit.	roseate	3L A pr	Ifjl	PK	Hungary	1807. D co	W.&K.2.121		
19815 grammifolia L.	Grass-leaved	5 A Pf	2 jn.au	L.Y	Portugal	1759. D co	Jac. ob. 4 100		
19816 fistulosa Brot.	fistulose	• ^t fr un	2 jnjl	Y	Spain	1817. Ti co			
19817 pinifflia Gou.	Pine-leaved	5 A "	1* jl.au	Y	S. Europe	1816. D co	Bur. ic. 496		
subulata Lam.									
19818 trachysperma GUnt.	rough-seeded	^ A un	H Jl.au	Y	Calabria	1818. S co			
scrullata Viv.									
19819 angustiflula L.	narrow-leaved	■ A pr	ijn.au	Y	& Europe	1759. D co	W.&K.2.1		
19820 eriosperma Bieb.	woolly-seeded	■ A pr	1 jn.au	Y	Siberia	1805. D co			
19821 chondrillo'ides Pou.	Chondnlla-like'	■ A un	1 jnjl	Y	Spain	1818. S co			
19822 stricta Horn.	strict	■ A un	2 jnjl	Y	Tauria	1818. S co			
villosa Bieb.									
19823 mollis Bieb.	soft	^ A un	1* jnjl	Y	Caucasus	1818. S co			
undulata Vahl									
19824 rumiciflia Hort.	Rumex.leaved	^ A un	1 Jn.au	Y	1820. D co			
19625 tatarica Bieb.	Taurian	'ik A pr	1 jn.au	Y	Tauria	182a S co			
19826 parviflbra Jac.	small-flowered	5 A pr	2 jlau	Y	Austria	1819. D co	Jac.au. 305		
19H27 lanaU Bieb.	woolly	A A Pr	1 Jlau	Y	Iberia	1824 D co	M.h.7.6.17		
19828 radiata Fis.	rayed	£ A un	* jnjl	Y	Siberia	1820. S co			
19629 ensifolia Bieb.	sword-leaved	it A pr	1 myjn	Y	Caucasus	1835. D co			
19830 himata L.	hairy	5 Apr	J myjn	Y	S. Europe	1818. D co	AI.ped.3L 33		
19831 villbsa Dec.	villous	^ Q or	J* jnjl	Y	S. Europe	1818. S co	Scop. car. 46		
* 2202. PICRIDIUM Pers. PICUIDIUM. (Pikros, bitter; taste.) Comp. Cichor. 3.									
19832 vulgare Def.	common	O cul	1} jn.au	Y	France	1773. S co	AI. ped. 1.16.1		
Stuichus jtricroides IV.									
19833 tingitnum Pers.	Tangier	O or	1 jn.s	Y	Baruary	1713. ■ ■	Bot. mag. 142		
19834 álbidum Pers.	vrhish-Jlowered^	A or	1 jl.o	W.Y	France	1781. ■ ■	Jac. ic 1.164		
Crépis álbida W.									
•2203. SournisTLE. (Somphos, hollow; stems.) Comp. Cichor. 33.— 40.									
19635 fruticosus Jac.	shrubly	L uJor	3 apjl	Y	Madeira	1777. S p1	Jac. ic. 1.161		
.1 irnn'ini Ofv.									
II* II K.	ving-leaved	■	3	...	Y	Madeira	1777. C co		
198 n\ en.	smooth	■	3	...	Y	Madeira	1816. C co		
198 " en.	lyrc-leaved	■	3	...	Y	M *--	— C co		
19839 r'.uiicutu* II. K.	long-rooted	■	3 jl	Y	("	C l.p		
V&W AyoscTift>lius Horn.	Hyoscris-lvd	"	4 myjl	Y	—	C l.p		
19841 abbreviatus Lk.	shortened	■	2 myjn	Y	—	C l.p		
	II.	■ ■ ■	—	Leaves undivided.					
19842 marUimus L.	sea	■	2 jl.s	Y	S. Europe	1748. D co	Al.ped.1.16.2		
\ 19843 sibiricus L.	Siberian	■	2 jl.s	L.B	Siberia	1759. D co	Gin. si. 2.3		
19844 cykneus Z. Zkm	Nepal blue	■	2 ilau	B	Nepal	1880. D co			
19645 cacalisBfblius Bieb.	Cácalia-leaved	■	liJLau	B	Ibena	1826. D Lp			

19847 *tartarius* L.

held	2	il.au	Crea	W. Indies	1818.	S co
Tartarian	4	jnjl	B	Siberia	1784.	D co



111. RUNCINAN I.—Leaves runcinate or lyrate.

19848 <i>pallistris</i> L.	tall marsh	^ A pr	England	riv.ba.	D co	Eng. bot. 935
19819 <i>arvensis</i> L.	corn field	A un	Britain	corn fi.	D co	Eng. bot. 674
19850 <i>oleraceus</i> L.	potherb	A un	Britain	fields.	S co	Eng. bot. 843
19851 <i>fefer</i> Lk.	rough	A un	Britain	fields.	S co	
19852 <i>/araxacifolius</i> J.F.	Dandelion-Leaved	O un	Guinea	1824.	S co	
\$19853 <i>Plumifer</i> L.	Plumper's	A un	Pyrenees	1794.	D co	
19854 <i>alpinus</i> L.	alpine	A un	Scotland	al. pas.	D co	
V19855 <i>Iapponicus</i> Fro.	Lapland	A un	Lapland	1804.	D co	Sm. ic. in. 21
\19856 <i>florictus</i> L.	Florida	A un	N. Amer.	1713.	S co	
19857 <i>caucasicus</i> Fit.	Caucasian	A un	Caucasus	1818.	D co	
19858 <i>acuminatus</i> IF.	acuminate	A un	N. Amer.	1812.	S co	
19859 <i>pallidua</i> W.	pale	A un	Canada	1704.	D co	Rob. ic. 148.151
19860 <i>clivicarius</i> ffor.	divaricating	A un	A. Ib23.	D co		
19861 <i>uliginibus</i> Bleb.	swamp	A un	Caucasus	1621.	S co	Schk. han. 256.
19862 <i>lacerus</i> W.	torn	A un	A. 18901	S co		
19863 <i>macrophyllus</i> L.	long-leaved	A un	N. Amer.	1823.	D co	
19864 <i>leucophafus</i> IV.	dusky	A un	N. Amer.	1821.	S co	

19866 *djondnwlides Desf.*19867 *ruseu* Bes.*

IV. PINNATI'FIDL—Leaves pinna tifid.

very tender	O un	2 jlau	Y	& Europe	1691.	S co	Pluk.al.93.3
Chondrilla-like	CD or	1 jnjl	Y	Spain	1729.	S 81	Boc.sic.13.7.4
rosy	5 A un	1 jnjl	R	Podolia	1827.	S co	

2204. ZACTITCA L. LETTUCE. (*Lac*, milk; milky juice exuding when broke) Comp. Cichoriace. 28.-37.

I. RUNCINA^TA.—Leaves runcinate or lyrate.

19868 <i>saftva</i> L.	cultivated	A un	Y-W	".....	1562.	S co	
19869 <i>crispula</i> L.	curled	A un	Y	1570.	S co	
19970 <i>palmata</i> W.	palmate	A un	cul	1683.	S co	
19871 <i>hirsuta</i> L.	cul	3 jn.au	Y	S. Amer.	1781.	S co	Jac. ic. I. Ki2
19872 <i>tfucreina</i> /.	cul	3 myjl	Y	Sweden	1686.	S co	
19873 <i>canadensis</i> L.	Canadian	3 jn.au	Y	Canada	1800.	S co	
19874 <i>stneta</i> Kit.	strict	3 jn.au	Y	Hungary	1805.	S co	W. ft K. 1.48
19875 <i>Chaisii</i> Vil.	Chais's	1f jn.au	Y	Europe	1818.	S co	Vil.rtel.a32
19876 <i>segusiana</i> Balb.	Segusiau	pr } jlau	P	Piedmont	1822.	S co	
19877 <i>fonchibbia</i> IV.	Sowthistic-lvd	2 jlau	Pa.B	Candia	1822.	D co	
19878 <i>elongata</i> MM.	elongated	3 jnjl	Y	Pennsylv.	1805.	S co	
19879 <i>Scariola</i> L.	Scanolia	3 jlau	Y	England	rubble.	S co	Eng. bot 268
19880 <i>maculata</i> Horn.	spotted	6 jlau	Y	Europe	1817.	S co	
19881 <i>viminea</i> Lk.	rusAg-tmggeA	1 jlau	Y	Austria	1789.	S co	Jac.au. 1.9
19889 <i>cichonifolia</i> Dec.	Succory-leaved ^ Q)	3 jlau	Y	Pyrenees	1821.	S co	
joncholdes Lap.							

II. []—Leaves undivided.

19883 <i>virbsa</i> L.	poisonous	A un	Britain	ch. ba.	S co	Eng. bot 1957
19884 <i>augustana</i> All.	Augustan	A un	Italy	1791.	S co	AL pod. 1.52.1
19885 <i>sagittata</i> Kit.		A un	Hungary	1805.	S co	W.&K.1.1
19886 * <i>aligna</i> L.		A un	England	ch. ba.	S co	Eng. bot 707
19887 <i>framinea</i> A.R.	W\oxy-leaved	A un	tt Amer.	1816.	S co	
19888 <i>Integrifolia</i> A.R.	Grass-leaved	A un	Missouri	1817.	S co	
19889 <i>villisa</i> Jfc.	entire-leaved	A un	E. Indies	1784.	S co	
19890 <i>indica</i> L.	villous	A un	Caucasus	1823.	S co	
19891 <i>altissima</i> Bich.	Indian	A un	1810.	D co	Jac. vin. 1.57
19892 <i>tuberosa</i> Jac.	tallest	A un				
	tuberous	A un				

11L PIXNATI'FID*—Leaves pinnatifid,

19893 <i>cretica</i> Desf.	Cretan	^ Q u n	2 jlau	Y	Crete	1818.	S co
19894 <i>tenerrima</i> Pou.	very tender	5 A un	\$jlau	P	S. Europe	1815.	S co
19895 <i>perennis</i> L.	perennial	5 A un	2 jn.au	L.B	Germany	1596.	D co

Bot. mag. 2130

2205. CHONDRI'LLA L. GUM-SUCCORY. (*Chondros*, lump; lumps of gufomy matter on stems.) Cbm.GieA.aJL

19896 <i>Junccea</i> L.	common Rush	£ A un	U s.o	Y	France	1633.	D co	Jac. au. 5.427
19897 <i>graminea</i> Bieb.	Graas-ieaved	5 A un	H < o	Y	Volga-	1824.	D co	
19898 <i>latifolia</i> Bieb.	broad-leaved	5 CD un	1 a.u.s	Y	Caucasus	1820.	S co	

•2206. PRENA'NTHESES. PRENANTBES. (*Pretie\$, drooping, anthos*, flower.) Comp. Cichoīūc&e. la—19.

I. INDIVIVE.—Leaves undivided.

19899 <i>purpurea</i> L.	purple-flowered	4 jl.8	P	Germany	1653.	D co	Jac. au. 4.317
19900 <i>racemosa</i> Mx.	racemose	2 jn.au	Y	N. Amer	1812.	D pl	
19901 <i>chondrifoloides</i> Ard.	Chondrilla-like	1 jn.au	Y	S. Europe	1817.	D co	Ard. sp. 2.7
19902 <i>tenuifolia</i> L.	fine-leaved	1 jlau	Y	Switzerl.	1819.	D pl	At ped. 33.2
19903 <i>hispida</i> Pall.	hispida	3 jlau	Y	Siberia	1825.	D pl	
19904 <i>filiformis</i> 7*	white-flowered	2 jlau	W	N. Amer.	1762.	D pl	Bot mag. 1079
19905 <i>rubricanda</i> PA,	reddish	1 jlau	R.W	N. Amer.	1816.	D pl	
h 19906 <i>cordata</i> IV.	hevt-leaved	4 jlau	Pa.Y	N. Amer.	1816.	D co	

II. TRI'LOBJE.—Leaves trifid or trilobed.

& 19907 <i>altissima</i> L.	tallest	i A or	6 jl.au *	LY	N. Amer.	1696.	D pl	Pluk.al.317.2
\19908 <i>serpentaria</i> PA.	snake	5 A un	1 jn.au	Y	N. Amer.	1824.	D pl	Ph. am. 2.24

III. LYRA Vs.—Leaves lyrate, runcinate, or pinnatifid.

19909 <i>muralis</i> L.	wall	3t A w	2 jl	Y	Britain	woods.	D co	Eng. bot 457
^ 19910 <i>vimenea</i> L.	twiggly	Q> or	1\$ jnjl	Y	Austria	IS 10.	S co	Jac. au. 9
i 19911 <i>virgata</i> 3/x.	rod	5 A un	2 jl. au	Y	N. Amer.	1823.	D pl	

19912 <i>tuberba</i> Stev.	tuberous	A A un	1 jl	...	Iberia	1817.	D pi	
19913 <i>deltoidea</i> Bieb.	deltoid	^ A un	2 jlau	Y	Iberia	1825.	D pi	

IV. FBUTICOS-E.—Stems shrubby.

19914 <i>spinosa</i> Forsk.	prickly	tt i un	3 mr.my	Y	Barbary	1640.	S co	Par th
19915 <i>pinnata</i> L.	pinnate	«t i j un	3 jnjl	Y	Teneriffe	182a	S co	
19916 <i>arborescens</i> Brou.	tree	tL LJ un	3 jnjl	Y	Tereriffe	1824.	S co	

8207. LECNTODON L. DANDELION. (*Leon*, lion, *odon*, a, tooth j tooth-like divisions of lvs.) *Comp. Cich.* 15.-18.

I. KEFLE'XI.— <i>Involucr refleted.</i>							
19917 <i>Taraxacum L.</i> common	A w	ap.jl	Y	Britain	me. pa.	D co	Eng. bot 510
19918 <i>erythrospermus Ander.</i> red-seeded	A un	my.jn	Y	Siberia	1820.	D co	
19919 <i>cornicuUtuS Horn.</i> small-horned	■	my.jn	Y	Europe	1819.	I) co	
19920 <i>sertinus Kit.</i> late-flowering	A un	14	jls.	Hungary	1816.	D co	W. & K. 2.114
19921 <i>alpinus Hoppe</i> alpine	A un	■	Y	Alp. Eur.	1810.	D co	
II. AFPRE'SSI.— <i>Involucr erect or oppressed.</i>							
19922 <i>pa16stris E. B.</i> marsh	A w	1} in jl	Y	Britain	moi. p.	D co	Eng. bot 553
Ilividus Kit.							
19923 <i>lievigatua W.</i> smooth-leaved	A un	■	Y	S. Europe	1816.	D co	
19924 <i>Stevens Spr.</i> Steven's	■	■	Y	Caucasus	J819.	D co	
<i>alpinus Stev.</i>							
19925 <i>nigricans W. & K.</i> blackish	A un	■	Y	Hungary	1816.	D co	
<i>Bessarabian</i>	A un	■	Y	Bessarabia	b1tf21.	D co	
III. IPATE'NTEB.— <i>Involucres spreading,</i>							
19997 <i>obovatus W.</i> obovate-leaved	& A un	1	Jl	Spain	1805.	D co	
19928 <i>eridopodus D. Don</i> woolly-footed	£ A un	jjl.au	Y*	Nepal	1818.	D e o	
19929 <i>bulbibus L.</i> bulbous	^ A un	1	my.jl	S. France	182&* D co		
Prenanthes bulbosus	Dec.	//ieracium	bulbaum IV.				
19930 <i>glauci-scens Jücb.</i> glaucescent	^ A un	1	jl	Volga	1823.	D co	
IV. Du^Biua— <i>Doubtful to which section it belongs.</i>							
19931 <i>cichoraceus Schl.</i> Succory-like	^ A un	1	jl.au	Switzerl.	1819.	D co	
•2208.-4PA'RG IA Sco. APARGIA. (A Greek name of a plant now unknown.) <i>Comp. Cichordcear.</i> 19.—25.							
I. SCAP'TGER/K.— <i>Scape simple onc-Itoived.</i>							
19932 <i>aurantiaca Kit.</i> Orange-colored	& A P ^r	i	my.jn	O	Hungary	1816.	I) co
19933 <i>alpinaJ/cw*</i>							
19934 <i>coronopifolia W.</i> alpine	^ A un	1	my.jn	Y	Austria	1816.	D co
199.5 <i>fasciculata Biv.</i> Buckhorn-lvd	A un	1	ray.jn	Y	Numidia	1816.	D co
<i>cichoracea Ten.</i>							
19936 <i>hastalis Host</i> fascicled	spear	■	1	jl.au	& Europe	1796.	D co
19937 <i>dubia Hoppe</i>	doubtful	■	1	au	Germany	...	D co
19938 <i>incana Scö.</i>	hoary	■	1	my.jn	S. Europe	1784.	I) co
19939 <i>7'araxaci W.</i>	Dandclion-iM*	■	1	au	Britain	scalps.	D co
19940 <i>crfspa W.</i>	curled	■	1	jl.au	France	1803.	D co
19941 <i>hSpida WI</i>	hispid	■	kils	O	Britain	ch.nas.	D co
19942 <i>crteea W.</i>	saffron	■	1	my.jn	Hungary	182a	D co
19943 <i>caudacia Bleb.</i>	Caucasian	■	1	jl!	Caucasus	1820.	D co
19944 <i>strigosa flifet.</i>	strigose	■	1	in.jl	Caucasus	1818.	D co
19945 <i>macrofizza Gae.</i>	long-rooted	■	2	jl	Apulia	1827.	D co
19946 <i>Villarsi W.</i>	Villars's	A un	1	in jl	Dauphinyl	1821.	D co
19947 <i>Gouani Schl.</i>	Gpuan's	A un	1	my.jn	Switzerl.	1824.	D co
II. CAULESCENTES.— <i>Stem or scape divided.</i>							
19948 <i>autumnalis Hoppe</i>	autumnal	■c	A w	1	au	Britain	me pa.
19949 <i>aspera Kit.</i>	rough	5	A un	1	jn.jl	Hungary	1805.
19950 <i>Ayosco'ids Sieb.</i>	Hyoseris-like	■c	Q>	un	1	jn.jl	Crete
2209. THIU'NCIA Roth THRINCIA. (<i>Thrigkos</i> , battlement; seed-crown of marg. florets.) <i>Comp. Cichor.</i> 4.—6.							
19951 <i>hlrtia Both</i> rimp/c-haired	A/ A un	1	* Jl.au	Y	Britain	gra.pa.	D co
19952 <i>hlspida Both</i> hispid	■ O	un	1	in.jv	S. Europe	1815.	S co
19953 <i>tuberosa Dec.</i> tuberous-roo/a/A	A un	*	mv	Y	Europe	1683.	D co
jp&rgiatuber?>sa W. Leontodon tuberctus L.					Lob. ic.	232.1	
19954 <i>marocc'a na Pars.</i> Morocco	O un	1	jn.au	Y	Morocco	1799.	S co
Ify6seris hispida Schous.							
2210. PrCRIS L. * OX-TONGUE. (Pihos, bitter: taste)							
19955 <i>Aleracioides L.</i> Hawkweed-hke	O un	1	Jjfa	Y	England	bor.fl.	S co
19956 <i>strigosa Bieb.</i> strigose	^ A w	1	jl.au	Y	Tauria	1819.	S co
19957 <i>altissima Del.</i> tallest	O w	2	jn.au	Y	Egypt	1800.	S co
Crépis/appacea W.							
19958 <i>rlgida Led.</i> rigid	O un	2	jl.au	Y	S. Europe	1817.	S co
1959 <i>dahurica/fr.</i> Dahurian	O un	1	jl.au	Y	Dahuria	1818.	S co
■ Splicewort-like^f	O un	1	Jjfa	Y	Barbary	1805.	S co
19961 <i>lyfata Del</i> lyfiate	O un	1	* il.au	Y	Egypt	1817.	S co
19962 <i>hlspida //, K.</i> Upid	A un	1	jlau	Y	levant	1789.	D co
19963 <i>SpreiKvnVina Chaix</i> Sprenger's	O pr	J	in.jl	Y	Portugal	1783.	S co
19964 <i>pauciflora Gae.</i> few-flowered	O un	1	jl.au	Y	S. Europe	1816.	S co
19965 <i>llhagad'iolus Pert.</i> Hhagadiolus	O un	Uj	jl.au	Y	Spain	1816.	S co
19966 <i>kmatschatica/o.t.</i> Kamtschatka	O un	1	jl.au	Y	Kamtsch.	1820.	S co
•2211. HIERA'CJUM L. HAWKWEED. (H<rx, hawk; sup. to sharpen sight with juice.) <i>Comp. Cichor.</i> 123.—143.							
I. UNIFLORA.— <i>Scapes simple, one-flowered; stolons none.</i>							
19967 <i>rupdstre All.</i> rock	• ^ A pr	4J>J1	Y	Switzerl.	1820.	D co	Al. au. 1.1.1
19968 <i>alpinum L.</i> alpine	5 A P	h J'-au	Y	Britain	al. roc.	D co	Eng. bot 1110
19969 <i>pusillum Ph.</i> small	^ t A or	^ jla	Y	Labrador	1800.	D s.p	
19970 <i>ptimilum Hoppe</i> dwarf	^ t A w	i J'-au	Y	Switzerl.	1819.	D s.p	
2 tubul&tum Schl	5 A or	jl.au	Y	Switzerl.	1819.	D s.p	
19971 <i>glabr&tum Hoppe</i> smooth	^ A or	} jl.au	Y	Switzerl.	1819.	D co	
2 tubul&tum Schl	5 A or	Jjl.au	Y	Switzerl.	1819.	D co	
19972 <i>alpestre Jac.</i> alp	5 A pr	i J'-au	Y	Switzerl.	1822.	D co	Jac. au. 191
J19973 <i>aureum Sco.</i> golden	5 A or	my.jl	D.Y	Italy	1769.	D pl	Jac. au. 3.297
II. STÖLONIFERA.— <i>Stoloniferous scapes many-flowered, seldom leafy.</i>							
19974 Pilos&la L. Mouse-car	9U A P'	i my.jl	Y	Britain	dry pa.	1> co	Eng. bot 1093
2 Pelecriflma Stew.	Pelcter's smooth	fl A or	4 my.jl	Y	Switzerl.	1819.	D co
3 Pscido-Pilos^Ua	Bast. Mouse-car	J^ A or	1 my.jl	Y	Switzerl.	IK19.	D co
19975 stolon iferum Kit.	stoloniferous	L ^ or	1 my.jl	Y	Hungary	1820.	D co
19976 aurantiacum L. Orange	& A or	jn.jl	O	Scotland	sc.wo.	J> p-1	Eng. bot. 1469
2flavum yellow	'Z, A or	ijn.jl	Y	Switzerl.	181P.	D co	
J9977 piloseihforme Hoppe	Mouse-car like j^ A or	my.jl	Y	S. Europe	1819.	D co	

19978 <i>d&rium L.</i>	doubtful	L A pr	ijl.au	Y	Britain	hills.-	D co	Eng. bot 2332	
19979 <i>auricula L.</i>	umbell-ed-eax	%, A pr	U jl.au	Y	England	moun.	D co	Eng. bot. 2368	
19980 <i>repens IV.</i>	creeping	& A or	Ujl.au	Y	Switzerl.	1819.	D co		
19981 <i>Gochnati Spr.</i>	Gochnati's	JW A or	■ ■ ■ .	Y	Switzerl.	1819.	D co		
	coltnum <i>Goch.</i> , cymbsum <i>Oed.</i>								
19982 <i>BauMni Bes.</i>	Bauhin's	& A or	■ jnl.	Y	Germany	1818.	D co		
19983 <i>fallax W. en.</i>	fallacious	t A pr	■ jl.au	Y		1816.	D co	
19984 <i>florenfinum VU.</i>	Florentine	AT A pr	■ jl.au	Y	Germany	1796.	D co	Bau. pin. 67	
19985 <i>prsalatum VU.</i>	very high	AT A or	2 jn.au	Y	Switzerl.	1819.	D co		
19986 <i>cymbsum L.</i>	cymose	■ ■	1 my.jn	Y	Europe	1739.	II co	Col. ec. 1. S49	
19987 <i>piloselloides VU.</i>	Mouse-ear-like	^ A or	* jn.jl	Y	Switzerl.	1819.	D co	ViLdel.3.27	
19988 <i>flagellare W. en.</i>	twiggy	% A pr	1 my.jl	Y		1816.	D co	
	collnum <i>Bes.</i>								
19989 <i>BesserYmkwi Spr.</i>	Besser's	■ A or	1 jn.jl	Y	Germany	1816.	D co		
	auricula <i>Bes.</i> collnum <i>W.</i>								
19990 <i>collnum Bernh.</i>	hill	■ A or f	my.jl	Y	Switzerl.	1819.	D co		
		<i>Scapes many-flowered, naked, sometimes few-leaved; stolons none.</i>							
19991 <i>angustifoliumfop/tt</i>	narrow-leaved	^ A pr ±	-my.jn	Y	"Switzerl."	182*	D co		
19992? <i>bifircum Bieb.</i>	two-forked	» J pr	11 jn	Y	Tauria	1820.	I) co		
19993 <i>bifidum Kit.</i>	bifid	■ ■	14 jn	Y	Hungary	1820.	D co		
19994 <i>Gmcfini L.</i>	Gmelin's	■ ■	11 jnjl	Y	Siberia	1798.	D co	Gm. si. 2.8.2	
19995 <i>praemorsum L.</i>	bitten	■ ■	Pa.Y		Switzerl.	1818.	D co	Gm. si. 2.13.2	
19996 <i>crbicum Lam.</i>	saffron	■ ■	Cop		Siberia	1818.	U co	Gin. <. 2.8. 1	
19997 <i>incarnatum Jac.</i>	Rush-colored	■ ■	Pk		Carniola	1815.	D co	Jac. ic. 578	
	monCnum <i>Sco.</i>								
19998 <i>successifolium All.</i>	lopi>cd.lcaved	* □ □	■ ■	Y	Switzerl.	1819.	D co		
	integriglbum <i>Hoppe</i>								
19999 <i>Lawsoni Vil.</i>	Lawson's	■ □	■ ■	Y	Britain	n.ofe.	D e o	Eng. bot. 2083	
20000 <i>venosum W.</i>	veiny	■ □	■ ■	Y	N. Amer.	1790.	D co		
20001 <i>Gronbii W.</i>	Gronovius's	■ □	■ ■	Y	N. Amer.	1798.	D co		
20002 <i>Bubntdum Schl.</i>	nakedish	■ □	■ ■	Y	Switzerl.	1819.	D co		
		IV. FOLIOS*. — Stems leafy; leaves entire, runcinate or toothed.							
20003 <i>paniculatum IV.</i>	panicled	■ ■	■ ■	■	Canada	1800.	D co		
20004 <i>glaucum All.</i>	glaucous	■ ■	■ ■	■	SL Europe	1807.	D co		
20005 <i>saxatile Jac.</i>	rock	■ ■	■ ■	■	Austria	1801.	D co	Jac. ic. 1.163	
20006 <i>prenanthoides VU.</i>	Prernanthus-lik	■ ■	■ ■	■	Scott. riv. ba.	D co	Eng. Dot 2235		
20007 <i>chondrilloides Jac.</i>	Cliondlnlla-like	■ ■	■ ■	■	Austria	1640.	D co	Jac. au. 5.429	
20008 <i>jonchiftlium Bteb.</i>	Sowthistle-lvd	■ ■	■ ■	■	Caucasus	1821.	D co		
20009 <i>cydoniseflium VU.</i>	Quince-leaved	■ ■	■ ■	■	France	1816.	D co		
20010 <i>molle Jac.</i>	aofst-tcaved	■ ■	■ ■	■	Scotland sc. wo.	D co	Eng. bot. 2210		
20011 <i>ccrinthoules L.</i>	Honey wort-lik	■ ■	■ ■	■	Scotland sc. roc.	D co	Eng. bot. 2378		
20012 <i>amplexicaule L.</i>	stem-clasping	■ ■	■ ■	■	Pyrenees	1739.	I) co	Al. ped. 15.1	
	2 pulmonarioides&A.	Lungwort-lik	■ ■	■	Switzerl.	1819.	D co		
20013 <i>croaticum Kit.</i>	Croatian	k	■ ■	■	Hungary	1 ■ ■	D co		
20014 <i>intybacum Jac.</i>	Endw-leaved	■ ■	■ ■	■	Pa.Y	Europe	1794.	D co	Jac. au. 5. ap. 43
2(X)15 <i>crassifolium Kit.</i>	thick-leaved	■ ■	■ ■	■	Hungary	1820.	D co		
20016 <i>Halleri Vil.</i>	Haller's	■ ■	■ ■	■	Britain	sc.mo.	D co	Vil.deL.3.26	
90017 <i>Irbbridum Chaix</i>	hybrid	■ ■	■ ■	■	Switzerl.	1826.	D co		
20018 <i>lafatum Kit.</i>	woolly	■ ■	■ ■	■	Hungary	1820	D co	W.&K.2.127	
S0019 <i>Hoppdnum Funk</i>	Hoppe's	■ ■	■ ■	■	Switzerl.	1819.	D co		
90020 <i>canewens Schl.</i>	canescent	■ ■	■ ■	■	Swizerl.	1820.	D co		
90021 <i>ovatum Schl.</i>	ovate	■ ■	■ ■	■	Swizerl.	1819.	D co		
20022 <i>Sternbergii Lk.</i>	Sternberg's	■ ■	■ ■	■	Swizerl.	1819.	D co		
20023 <i>maculatum E. B.</i>	bpotted/t<zverf	■ ■	■ ■	■	Britain	al.roc.	D co	Eng. bot 2121	
90024 <i>pulmonarium E. B.</i>	Lungwort-like	■ ■	■ ■	■	Scotland	ic. roc.	D co	Eng. bot 2307	
90025 <i>pulmonario'ides Vil.</i>	Lungwort-like	■ ■	■ ■	■	France	1819.	D co		
20026 <i>pallescens Kit.</i>	palcl>cent	■ ■	■ ■	■	Hungary	1818.	D co		
90027 <i>pomfTihum //.</i>	Lcek-leaved	■ ■	■ ■	■	Austria	1640.	D co	Jac.au. 3.286	
90028 <i>montanum Jac.</i>	mountain	■ ■	■ ■	■	S. Europe	1775.	I) co	Jac. au. 2.190	
20029 <i><taticifolium All.</i>	Thrift-leaved	■ ■	■ ■	■	Europe	1804.	D co	Vil. del. 3.27	
20030 <i>anchusa?folium J5?r/.</i>	Bugloss-leaved	■ ■	■ ■	■	Italy	1816.	D co		
	verbaaciflum VU.								
90031-verbasciflumP?rs.	Mullein-leaved	^ A or	1 my.au	Y	S. Europe	1732.	D co	Mil. ic 1.146.1	
	<i>Andrpala lanata L.</i>								
20032 <i>erioiphllum Lk.</i>	woolly-leaved	■ ■	■ ■	■	Britain	D co		
20033 <i>sylvaticum L.</i>	wood	■ ■	■ ■	■	Scotland	rocks.	D co	Eng. bot. 2031	
20034 <i>vilKisum L.</i>	villus alpine	■ ■	■ ■	■	Switzerl.	1819.	D co	Eng. bot. 2379	
20035 <i>valdepiolosum VU.</i>	very pilose	■ ■	■ ■	■	Switzerl.	1819.	D co		
20036 <i>elongatium Lap.</i>	elongated	■ ■	■ ■	■	Switzerl.	1819.	D co		
20037 <i>SchrSderi Schl.</i>	Schrader's	■ ■	■ ■	■	Switzerl.	1819.	D co		
20038 <i>andryaMdm VU.</i>	Andryala-like	■ ■	■ ■	■	Switzerl.	1819.	D co	Vil. del. 3.29	
	2 Leotardi Vil.				Switzerl.	1819.	D co		
90039 <i>pilocephalum Lk.</i>	Leotard's	■ ■	■ ■	■	Europe	1820.	D co		
90040 <i>trichocéphalum?P.Kit.</i>	hairy-headed	■ ■	■ ■	■		1823!	D co		
20041 <i>flexuofolium Kit.</i>	bcndinz-stal/cett	■ ■	■ ■	■	Hungary	1804.	D co		
20042 <i>specibsum Horn.</i>	showy	■ ■	■ ■	■		1818.	D co		
20043 <i>prostrutin Dec.</i>	prostrated	■ ■	■ ■	■	S. Europe	1822.	D co		
20044 <i>inctaum Bub.</i>	hoary	■ ■	■ ■	■	Caucasus	1817.	D co		
20045 <i>Kalmi I*</i>	Kalmi*s	■ ■	■ ■	■	Pennsylv.	1794.	I) co		
20046 <i>canndensc Mx.</i>	Canadian	■ ■	■ ■	■	Canada	1800.	D co		
20047 <i>virgatum Ph.</i>	twiggy	■ ■	■ ■	■	N. Amer.	1816.	D co		
20048 <i>spociosissimum V.</i>	showiest	■ ■	■ ■	■	S. Europe	1821.	D co		
20049 <i>denticulatum E. B.</i>	small-toothed	■ ■	■ ■	■	Scotland	sc.wo.	D co	Eng. bot. 2122	
	prenanthoides Sm.								
20050 <i>parviflrum Schl.</i>	small-flowcrcd	■ ■	■ ■	■	Switzerl.	1819.	D co		
20051 <i>M Uteri Lk.</i>	Miller's	■ ■	■ ■	■		1820.	D co		
20052 <i>sudeticum Ster.</i>	Sudctic	■ ■	■ ■	■	Switzerl.	1819.	D co		
20053 <i>whioWes Lum.</i>	Viper's-Bugloss-like	■ ■	■ ■	■	Hungary	1802.	D co	W.&K.2.145	
20054 <i>picridiflum Schl.</i>	Picris-leaved	■ ■	■ ■	■	Switzerl.	1819.	D co		
20055 <i>verruculatum Lk.</i>	warled	■ ■	■ ■	■		1821.	D co		
90056 <i>undulatum H. K.</i>	vravc-leaved	■ ■	■ ■	■	Spain	1778.	D oo		
90057 <i>macropVum Ph.</i>	long-leaved	■ ■	■ ■	■	Canada	1825.	D co		
90058 <i>dwitaturn Hoppe</i>	toothed	■ ■	■ ■	■	Swizerl.	1819.	D co		
	*M59 latiflbum Frol.	broad-leaved	■ ■	■	Croatia	1820.	D co		

2212. HAPALOSTE*PHIUM D.Don HAPAL. (*Hapalos*, soft, *stephos*, crown; hairy receptacle.) *Com.Cich.* & £0089 *sibiricum* D. Don Siberian ^ A pt 3 jl.au Y Siberia 1755. D co Gm. si. 2.10

	/fieracium sibiricum W.	Crepis sibirica L.				
20090	/prenaicum D.	Don Pyrenean	^ A « li	JLau	Y	Pyrenees 1723. D co
	/fieracium pyrenicum All.	blattarioldef.iL.				
£0091	pilsum D.	Don pilose	ft A Er	1 jl.au	Y	Pyrenees 1723. D co Pier. par. 184
	/fieracium olattarioides L.	Lepicadne multicatulis Lap.				
20092	austrlacum D.	Don Austrian	^ A pr	1 jl.au	Y	Pyrenees 1723. D co Jac.aus. 5.441
		Crepis austriaca Jac.				
20093	grandiflbrum G.	Don great-flwd	3t A pr *	i Lau	Y	Switzerl. 1791. D co W. & K. 1.99
	/fieracium grandiflbrum L.	Lepicadne grandiflbra Lap.				
£0094	paluddsum D.	Don marsh	■ A pr	li Jl.au	Y	Britain moun. D co Eng.bot.1094
	/fieracium paludsum L.					
20095	lyratum G.	Don lyre-leaved	■ A P ^d	1 jl.au	Y	Siberia 1777. D co Gin. si. £.9
	/fieracium lyratum L.					
20096	spinulsum Lap.	spinulose	■ A or '1	jl.au	Y	Pyrenees 1820. D co
	Lepicadne spinulosa Lap.					

2213. LAGCySERISI. LAGOSEHIIS. (*Lagos, a hare, seris, succory.*) Camp. Cichor. 2.
 20097 tadrica *Bieb.* Taurian O un 2 jl.au P Caucasus 1817. S co
 £0098 nemausensis *Bieb.* Nismcs O un 2 jl.au Y Caucasus 1794. S co Al. ped. 1.75.1
 Crénis nemausensis *Gou.*

£214. BORKHAU'S/if B&hm. BORKHAUSIA. (*Moritz Borkhausen*, a German botanist) *CompXichor.* 21.

I. <i>Leaves undivided.</i>										
20099	alpta L.	alpine	■	■	Y	Italy	1739.	S	co	Gm.si.2.5
20100	ftclidiflba Dec.	Daisy-leaved	O	un	1	jl.au	Y	Corsica	1822.	■ co
20101	macrophylla Spr.	long-leaved	O	un	2	jl.au	Y	Numidia	1819.	■ co
20102	Gandolff's Spr.	Dccandoll's	O	un	1	jl.au	Y	1817.	■ co
20103	juncio'ides Spr.	Seuccio-like	O	un	1	jl.au	Y	Egypt	18120.	■ co
	Crepis senecioides Del.									
20104	apargioides Spr.	Apargia-like	■	Q	un	1	jl.au	Y	SwitzerL	181& S co
	/fieracium stipitatum Jac.									
20105	Suffreniima Dec.	Suffren's	O	un	■	jn.jl	Y	Corsica	1818. S	s.l
20106	versicolor Spr.	party-colored	■	A	un	■	jn.jl	Y	Dauria	182a S co

II. RUNCINAVA. — *Leaves runcinate, It/rate, or pitmatid.*

20107	rabra Lk. <i>Crepis rhabra L.</i>	red	O pr	1*	jn.jl	P	Italy	1632.	S co
£0108	nicaeensis Lk. <i>Crepis niceensis</i>	Nice	O un	li	jn.jl	Y	Nice	1823.	S co
20109	to'tida Lk. <i>Crepis foetida L.</i>	fetid	■ Q>	un	li	jn.jl	Y	S. Europe	1824. S co M.h.7.4.4
£0110	faraxibifolia Spr. <i>Caroliniana Nut.</i>	Dandelion-leaved	O un	li	jl.au	Y	Italy	1818.	S co Bal.mis.△
20111	bursifolia Spr. <i>Shepherd's purse-lvd^</i>	Carolina	O un	1	jl.au	Y	Carolina	1824.	S co
20112	bursifolia Spr. <i>Lag6scris bursifolia Lk.</i>	Crepis bursifolia L.	Q un	■	jl.au	Y	Sicily	1817.	S co Boc.mu.106.112
£0113	leontodontoides Spr. <i>Lag6seris leontodontoides Lk.</i>	Dandelion-like^	Q un	1	jl.au	Y	Italy	1804.	S co
£0114	raphanifolia Spr. <i>Lagoseris rapnanifolia Lk.</i>	Raphanus-lvd	^ A un	2	jl.au	Y		1816.	S co
20115	intybaccea Spr. <i>Lagoseris intybacea Lk.</i>	Endive-leaved	^ Q u	n	ljl.au	Y	Portugal	1816.	S co
20116	gravicolens Lk. <i>Asperula Jt.</i>	strong-smelling	O un	li	jn.jl	Y		1825.	S co
20117	aspera/Jt. <i>hispida Lk.</i>	rough	O un	fi	jl.au	Y		1797.	S co
S0118		hispida	O un	2	jl.au	Pa.Y	Sicily	1798.	S co W. & K. 1. 43

*2215. CRE^PIS IF. CREPIS. (Name by Pliny to a plant of which he gave no description.) *Comp. Cich.* 19.-47.

2221. HYO'SERIS L.	SWINE'S SUCCOBY.		Fayt, a hog, scris, succory.)	Comp. Cichor. 4
20170 radinta L.	rayed	^ A un	Jn.jl Y S. Europe 1640.	D co Pluk.al.37.2
20171 Ibcida L.	shining	5 A un	Ijn.a u Y Levant 1770.	D co Schm. ic. 39.41
20172 sc ^c bra L.	rough	O un	*ii! ^{uu} X S ^c ily Y3 §	^{co} Boc.mu.UelOG
20173 arenaria Schous.	sand	O un	ijl.au Y Morocco 1800.	S co
2222 HEDY'PNOISL.	HEDVPNOIB. (<i>Hedys</i> , sweet, <i>JHUO</i> , to breathe; scents breath.)		Comp Cichor. 7—	
20174 monsiMjlicnsis W.	Montpelier	O un	1 jn.jl Y S. Europe 1683.	S co
20175 rhagadiolus(Mdes L.	Rhagadiolus-like	O un	1 Jl.jl au Y S. Europe 1773.	S co Cav ic 1 43

20176 erotica IV. SOW coronopi folia Ten. 80178 tubariformis Ten. 80179 mauritanica W. 20180 pendula JF.	Cretan Buckhorn-lyd tube-formed Moorish pendulous	8 un 1 jn.jl Y O un 1 jn.jl Y O un 1 jn.jl Y O un 1 jn.jl Y O un 1 jn.jl Y	Candia 1731. S co Italy 182a S co Naples 1824. S co Barbary ... S co S co
2223. ROBE'RT/^ Dec. 80181 foraxacrides Dec. 80182 pinnatifida Spr. ifypocheVia pinnatifida Cyr.	ROBEOTTA. Dandelion-like * A ■■■■■ f jn.jl Y pinnatifld 5 CD ■■■■■ 1 jl.au Y	{ M. Aobert, a Conican botanist) Corsica 1834. S co Naples 1827. S co	Comp. Cichor. □
•2224. SERROLA L. 80183 setnlsis L. 80184 ferens L.	SERIOLA. -Etna stinging □	(Sens. succor); resemblance.) Italy 176 i. S co Jac. ob. 4.79 S. Europe 1773. S co Schra. ic. 32	Comp. Cichor. 2—riL
2225. RODI'G/il Spr. 80185 commutata Spr. Crepis rhoeadifrlia Bieb. 80186 levigata Spr. Seriola lse ligate L.	RODIGIA. changed smooth Garlic	(Rodig, a friend of Sprengel's.) Caucasus 1816. S co	Comp. Cichor. 3.
80187 alliata G. Don Seriola alhata Biv.		Candia 1772. S co Desf.at8.lii	
8226. SOLDEVI'LLJ Lag. 80188 set&sa Lag.	SOLDEYILLA. bristly & A cu f my.jn Y	(SoUUvila, a friend of Lagasca's.) Spain 1822. D co	Comp. Cichor. 1.
8227. JTYPPOCHAIIISX. &20189 helvetica L. (20190 maculate L. 80191 minima Detf. 80192 hispida W. en. 80193 glabra L. 20194 macrorhla Glint. 80195 demgrpha Brot. 80196 arachnites Biv. \$80197 radicate L. 80198 canescens Stev. 80199 Balbisti Noc	CAT'S EAR. Swiss epotted-leaved least hispida smooth long-rooted deformed cobwebbed long-Tooted oanescnt Balbis's	for, choiris, a pig; eaten by.) Switzerl. 1779. D s1 Jac. ic. 1.165 England ch.hil. D s1 Eng. bot 225 Barbary 1797. S co S Europe 1804. D co W. h.b.1.46 Britain sa.hea. S co Eng. bot. 575 Calabria 1820. D co Spain 1819. S co Sicily 1827. S co Britain mepa. D s1 Eng. bot 831 Caucasus 1820. D co Italy 1824. D co	Comp. Cichor. 11.—14.
2228. ZAPSA^NA L. 80200 foetida Sco. Hy6seris fo/tida L.	NIPPLBWOBT. fetid J Apn □ jlau Y	[Lapaxo, to purge; qualities.) Italy 1722. D co W. & K. 1.49	Comp. Cichor. 8. —10.
60201 pusilla Sco. Hyōseris minima L.	Arn6seris fae/tida Lk. small O ' i my.jn Y	Britain gra. fl. S co Eng. bot ifS	
80202 commibis L. 80203 crispa W. 80204 pubescens Zry 80205 intermédia Bieb. 80206 lyrrata TV. en. 80207 grandiflura tif&	Arn6seris minima Lk. common O w jn.jl □ curled O un jf jl.au □ pubescent O or 2 jn.au □ intermediate O un U my.jn □ lyrate O un 22 jl.au □ great-flowered O or 2i jn.au □	Britain d t gr. S □ Eng. bot. 844 — 1799. S □ Europe 1816. S □ Tauna 1823. S □ Caspi. Sea 181 a □ Caucasus 1816. S □	
2229. ZACI'NTHA Tou. 80808 rerruedsa Gae. * Xapsana Zaointha L.	ZACINTHA. (Growing in the Island of Zactnthus, now Zante.) warty O un f jn.jl Y.Br & Europe 1633. S co Gac. fr. 2. 157.7	Comp. Cichor. I.	
8230. RHAGAD1\)LUS Tou. 80209 stellaris W. 80210 edblis Gae. /apsanoides Dcsf. Lapsana Rhagadlulus L.	RHAAODIOLUB. (Rhagas, a slit; divisions of calyx.) starred O un 1 jn.jl Y S. Europe 1633. S co	Comp. Cichor. 3	
£0211 Ko3lpnw W. Keelpnto lineáris Pail.	Koelpin's O un f jl □	Levant 1633. S co Schk. ban. 3.825	
2231. MOSCAMIIARjBfr. MOSCARIA. 80212 pinnatifida Fl.per.	MOSCHOS. [Moschos, musk; odor.) pinnatifid 5 i jUau ... Chile 1823. S co	Comp. Cichor. 1.	
* 8232. CATANA'NCHE L. CATANANCRE. 80213 caralea L. 80214 &teaZ.	Katanagke, strong incentive; used in philtres.) azure i A or 3 ilo B S. Europe 1633. D co Bot mag. 293 yeUow O or f jn.jl Y Candia 1640. S co Alp. ex. 286	Comp. Cieh. 2.—a	
8233. TRIPPI'LION Fl. per. TRIPPIUON. 80215 cordifiliun Lag. 80216 spinosum Lag.	(Treis, 3.ptilon, soft feather; seed-down.) cordate-leaved O pr i my.au W Chile 1824. S co Bot. reg. 8J3 spiny O pr 1 my.au B Chile 1827. S co	Comp. Labiot. 2. —4.	
8234. CICHO^RIUM L. Srecoav. 80217 Tntybus L. 80218 pitmñum Jac	(An Egyptian name adopted by the Greeks.) Intybus or wi!df A ■■■■■ 2 jn.au B Britain gra. so. S co Eng. bot. 539 dwarf O un jtl.au B □ co Jac. ob. 4.80	Comp. Cichor. S —7.	
20219 Endivia L. 80220 divaricatum Schous.	Endive O cul 2 jl.au B E. Indies 1548. S r.m		
20221 spinosum L.	divaricate O un 2 jl.au B Barbary 1798. S co		
2235. BACA^ZIA FL per. 80222 spinbsa Ft. per.	spiny ITBlun 2 jl.au B Candia 1633. S co Bau. pr. 62		
8236. SCCLYMSZ. 80223 grandiflonu Detf.	GOLHEN THISTLE. (Skolos, a thom; plants spiny.) large-flowered ^t A « 3 my.jn Y Barbary 1820. S co Desf. at. 218	Comp. Cichor. 3. -4.	
80224 maculatus L. 80225 hispfoicus L.	spotted O or 3 jl.au Y S. Europe 1631. S co Lam. il. GSt Spanish ^ or 3 jLi Y & Europe 1658. D co		
2237. il'RCTIUM L. 20226 Lappa /. major Schk.	BURDOCK. (Arktot, a bear; rough fruit) Lappa or smoothJvd Q) w 3 jl.au P Britain wa.gr. S co Eng. bot 1228	Comp6*it<r Cardudcea. 3 —4.	
20227 BardanaJF. ^ ^ tomentoNfa Schk	Bardana woolly-headed^). (3) w 3 jlau P Britain wa.gr* S co Eng. bot 2478		
20228 minus Bieb.	smaller ^Qw 2 jlau P Europe ... S co Schk. han. 3.227		

^^2238. CESTRUS NUSCass. *CESTRIMIS*. (*Cestrinus*, the wn of Helepus and Andromache.) *Cotop. CartL* 1.
20229 *carthamoides* & *Cats.* *Carduus* L. A or 1 jlau F *Barbary* 1797. D co DesCat2.223
Serratula acatilia Dec. *Cynara acacuis* X

8239. SERRATULA L. SAW-WORT. (Dimin. of *serra*, a saw; leaves sawed.) *CompCard.S8.-40.*

I. UNIFLOR*. — *Stem one-flowered.*

20230	<i>cynaroides</i> Dec.	Cynara-like	3	jl.au	P	Pyrenees	164a	D co	M.h. 7.25.1
	<i>Cnclus centaurio^des</i> L.								
20231	<i>cyanoides</i> Gae.	Cyanus-like	&	A or	2	jl.au	R	Siberia	1778. D co Scop. car. 53
	<i>alata</i> Poir.	<i>Cirduis cyanoides</i> L. <i>multis</i> Poll.							
20232	<i>transylvanica</i> Spr.	Transylvanian	&c	Q or	2	jLau	P	Transylv.	1818. S co
	<i>simplex</i> Baug.								
20233	<i>simplex</i> Dec.	simple	^	A pr	1	jl au	P	Nepal	182L D co Bot mag. 2482
20234	Kitaibeli G. Don	Kitaibel's	5	A or	2	jl.au	P	Hungary	1816. D co W.&K.1.52
	<i>Carduus nitidus</i> Kit.								
20235	<i>nudicaulis</i> Dec.	naked-stemmed	^	A or	2	jl.au	P	& Europe	1739. D c6 Cav.ic.3.226
	<i>Carduus cerithifolius</i> J'il.	<i>cerithoides</i> JV.							
£0236	<i>pannonica</i> Spr.	Pannonian	i	A or	2	jn.au	P	Hungary	1810. D co Jac au. 3. 5
	<i>Carduus pannonicus</i> L. <i>serratuloides</i> Jac.	<i>Cnclus pannonicus</i>							
20237	<i>depressa</i> Stev.	depressed	■	or	1	jLau	P	Caucasus	1818. D co Lin.tr. 11.58

II. MULTIFLOUIE. — *Stem many-flowered.*

20238	<i>tinctria</i> L.	common dyeing	■	or	1	jl.au	P	Caucasus	1823. S co Bux.c.2.15.1
20239	2 <i>alba</i>	white-Jfotrocot	■	■	■	■	P	Caucasus	1819. S co Lin.tr. 11.37
20239	<i>roronata</i> L.	crowned	■	■	■	■	P	Siberia	1739. D co Hot mag. 2SKI
20240	<i>quin'ueflbia</i> JV.	five-leaved	■	■	■	■	P	Persia	1804. D p. I Gm. si. 2.91
20241	<i>Behen</i> Dec.	Behen	■	■	■	■	P	Levant	1797. S co Bot mag. 1871
M	<i>Centauria Behen</i> I*.								
20242	<i>lutea</i> Spr.	shining	■	■	■	■	P	Caucasus	1823. S co Bux.c.2.15.1
	<i>Centaurea rittens</i> Bieb.								
20243	<i>elegans</i> Step.	elegant	■	■	■	■	P	Caucasus	1820. S co
20244	<i>pulchella</i> B. M.	neat	■	■	■	■	P	Siberia	1804. D co
50245	<i>centaurioides</i> L.	CenUury-like	■	■	■	■	P	Siberia	1804. D co Gm. si. 4.38.17
50246	<i>arguta</i> F\$.	Bhzrp-toothed	■	■	■	■	P	Hungary	1814. D co
20247	<i>nitida</i> Fu.	shining	■	■	■	■	P	Siberia	1H27. S co
20248	<i>caspica</i> Bieb.	Caspian	■	■	■	■	P	Caspia	1*20. S co
20249	<i>radiata</i> Bub.	rayed	■	■	■	■	P	Hungary	1800. S co W.&K.1.11
	<i>centaurioides</i> Bieb.								
20250	<i>multiflora</i> L.	many-flowered	■	■	■	■	P	Siberia	1816. S co
	<i>tineariifolia</i> Dec.								
20251	<i>xeranthemoides</i> Bieb.	Xeranth.-like	■	■	■	■	P	Caucasus	1825. D co Gm. si. 47.1
20252	<i>polyclonus</i> Bieb.	many-sprigged	■	■	■	■	P	Caucasus	1820. S co
	<i>ambigua</i> Dec.	<i>Carduus polyclonus</i>							
20253	<i>heterophylla</i> Desf.	various-leaved	■	■	■	2	jl.au	P	Dauphiny 1824. D co
20254	<i>riflcachadiflia</i> Bieb.	Steechas-leaved	■	■	■	1	jl.au	P	Tauria 181U D co
20255	<i>Plcns</i> Bkb.	<i>Picris</i>	■	■	■	1	jlau	P	Caucasus 1822. D co
20256	<i>alata</i> JV.	winged	■	■	■	2	jlau	P	
90257	<i>aspera</i> JV.	rough	■	■	■	1	jlau.s	P	Nepal" 1821. S co

2240. SAUSSU'REif Dec. SAUSSUBEA. (*Horace Ben.de Satasure*, a Swiss philo. andbot) *Com.Cantl* 11.

20258	<i>runcinata</i> Dec.	runcinate	■	■	■	2	jl.au	P	Siberia 1819. S co
20259	<i>lyrata</i> Fis.	lyrate	■	■	■	2	jl.au	P	Siberia 1827. S co
20260	<i>alata</i> Dec.	winged	■	■	■	2	jl.au	P	Siberia 1818. S co
20261	<i>elongata</i> Dec.	elongated	■	■	■	2	jl.au	P	Caucasus 1820. D co
20262	<i>salsa</i> Spr.	salt	■	■	■	1	jl.au.s	P	Caucasus 1816. S co
	<i>Serratula salsa</i> Bieb.								
20263	<i>amara</i> Dec.	bitter	■	■	■	1	jl.au	P	Siberia 1820. S co
	<i>Serratula amara</i> L.								
20264	<i>angustifolia</i> Dec.	narrow-leaved	■	■	■	2	jl.au	P	Siberia 1816*. S co Gin. si. 2.33
	<i>Serratula angustifolia</i> JV.								
20265	<i>racemifolia</i> Dec.	Willow-leaved	■	■	■	2	jl.au	R	Siberia 1796. S co Gm. si. 2.37
20266	<i>alpina</i> Dec.	alpine	■	■	■	1	jlau	P	Britain al.roc D p.l Eng. bot. 599
	<i>Serratula alpina</i> L.								
20267	<i>discolor</i> Dec.	discolored	■	■	■	1	jlau	P	Switzerl. 1818. D co Hal. h. 6
	<i>Serratula discolor</i> JV.								
20268	2 / apat hi folia Dec.	Lapathum-lvd	■	■	■	1	jlau	P.B	Europe 1816. S co Austria 1816. D co Jac.au. 440
	<i>pygmaei</i> Spr.	pygmy	■	■	■	1	jlau	P	
	<i>Serratula pygmaei</i> Jac.								

2241. CA'RDUUS L. THISTLE. (*Ard*, a point, *Celt.*; plants mostly spiny.) *Comp. Card.* 40.—100.

I.	<i>SoLVtk'm.</i>	—Flowers subsolitary	leaves decurrent						
90269	<i>irrgente</i> Pou.	■	H-i-u	R					
20270	<i>dubius</i> Balb.	■	■	P	1816. D co			
20271	crassifolius Horn,	thick-leaved	■	■	■	1805. D co		
2W73	alatus Swt.	winged	■	■	■	1812. S co Sw.fl.gar.103		
20274	<i>defloratus</i> L.	half-naked	■	■	■	■	Caucasus 1819. S co		
2(27)	<i>alm*strw</i> Kit.	deflorated	■	■	■	■	Austria 1570. D co		
»	JV.	alpine	■	■	■	■	Croatia 1805. D co		
&>	<i>ilius</i> L.	Burdock-like	■	■	■	■	Carniola 1804. D ca		
2(3)	< us L.	white-painted	■	■	■	■	Italy 1752. S co		
20275	hoary	silvered	■	■	■	■	Egypt 1789. S co		
	<i>iptny-hooked</i>		■	■	■	■	Hungary 1805. S co		
80281	<i>uncinatus</i> Bu-	hooked	■	■	■	■	Hungary 1802. S co		
20282	<i>corymbbus</i> Ten.	corymbose	■	■	■	■	Tauria 1817. S co		
20283	<i>cinericus</i> Bieb.	grey	■	■	■	■	Naples 1824. S co		
20284	<i>coinnus</i> Kit.	hill	■	■	■	■	Caucasus 1818. S co		
20285	<i>nigriscens</i> Vil.	nigrescent	■	■	■	■	Hungary 1818. D co		
20286	<i>nutans</i> L.	nodding	■	■	■	■	S. France 1819. S co		
20287	<i>podacanthus</i> Dec.	foot-spined	■	■	■	■	Britain gra.fl. S co		
S0288	<i>macrocephalus</i> Desf.	long-headed	■	■	■	■	France 1H19. S co		
20289	<i>montibus</i> Pall.	mountain	■	■	■	■	Numidia 1827. S co		
90290	<i>ciTU/nafoliu</i> Lam.	Carline-leaved	■	■	■	■	S. Europe 1820. S co		
90291	médium Got*.	intermediate	■	■	■	■	Pyrenees 1804. S co		
20292	lanuginosus JV.	woolly	■	■	■	■	Piedmont 1819. D co		
90993	<i>ocanthoides</i> L.	cant li us-like	■	■	■	■	Armenia 1820. S co	Gou.il. 2*	
	A						Bntain wa.gr. S co	Eng. bot. 973	

II. AGGRECAVI — Flowers aggregate; leaves decurrent.								
foreign	O or	2	jn.au	P	m.m.	1816L S co		
curled	& CD or	2	jl.au	P	Europe	1804. S co		
clavulate	Q or	2	jn.au	P	Canaries	1827. S co		
Onopordum-lk	^ A or	H	J-"	P	Iberia	1818. D co		
dense-headed	^ A or	If	jl.s	P	S. Europe	1739. S co		
whitish	O or	2	jn.au	P	Tauria	1816. S co		
Carlinsklike	& CD or	1	jl.au	P>rceees	1784. S co	Gou. il.23		
CarDna pyrenaica L.								
20301 tenuiflbrus Hud.	slender-flowered	O w	2	jn.au	P	Britain banks. S co	Eng.bot.4L2	
20302 arcticus Jac.	Arabian	O or	ijl.au	P	Arabia	1789. S co	Jac. ic. 1.166	
20503 Persoonata Jac.	Burdock	CD or	4	jl.au	P	Austria	1776. S co	Jac. au. 4.348
SUD4 onentalis Adam	oriental	^ A or	2	jl	P	Iberia	1804. D co	
2010J parviflbrus L.	small-flowered	^ A or	2	jn.jl	P	S. Europe	1781. D co	
III. SEBSILIFOLII.—Leaves sessile.								
20306 volgensis Bieb.	Volga	O or	2	jn.au	P	Volga	1820. S co	
20307 leucanthus Cav.	white-flowered	O or	2	jn.au	W	Spain	1816. S co	
IV. PETIOLA>ru&—Leaves petioled.								
20308 ftriplicifolius Trev.	Atriplex-leaved	^ A or	10	au	P	Siberia	1784. D co	
Onopordrum dcltoideum H. K.								

2242. SI'LYBUM Gae. SILTBUM. Used by Greek writers for a plant not now known.) *Comp. Card. 2 T*.
 2010J Mariantm Gae. Virgin Mary's £ CD or 5 jl P Britain banks. S co Eng. bot. 976
 CArduis Marianu L.
 20310 cernuum Gae. drooping £ A or 4 jn.jl Y Siberia 1755. D co Gm. si. 2.19
 Cnicus cernuus L.

*9343. CI'RSIUM Fail. CIRSIUM. (Kirsos, a swelled vein; supposed to heal) *Comp. Carduaceee. 89*

I. SOLITA^RJA.—Leaves decurrent; flowers red, subsolUary.						
20311 lanceolatum Sco.	com. lanceolate	^ CD	■ 3	jl.S	I	Britain bank's. S co Eng. bot. 107
Cicucus lanceolatus TV.						
20312 veriitum Spr.	dart-armed	CD or,	4	jn.s	P	Nepal 1820. S co
Cnlcus veritus D. Don						
20313 strictum Spr.	strict	■ A or	2	Jls	P	Naples 1819. D co
Cichicus strictus Ten.						
20314 cichoraceum Spr.	succory-like	■ A or	3	jl.s	P	Naples 1816. D co
Carduus cichoraccus Cyr.						
20315 italicum Dec.	Italian	CD or	3	jl.au	P	Italy 1815. S co
Carduus italicus Savi						
20316 canum Bieb.	hoary	■ A or	4	jl.au	P	Austria 1633. D co
20117 neglectum Fis.	neglected	■ A or	2	jl.au	P	Siberia 1827. S co
20318 laniflrrum Bibb.	woolly-flowered	^ A or	2	jl.au	P	Tauria 1819. S co
II. POLYA'NTHEMA.—Leaves decurrent / flowers red, aggregate						
90319 pafastre Sco.	marsh	CD w	3	jl.au	P	Britain m.pas. S co Eng. bot. 071
CSrdius pallistris L.						
23320 pungens Spr.	pungent	CD or	3	jl.au	P	S. Europe 182a S co
Cnicus pungens Seb.						
20321 polyanthemum Spr.	many-flowered	CD or	■	■	Pk	Rome 1739. S co Tri. ob. 103
Carduus polyanthemus L.						
20322 paniculatum Spr.	panicled	■ A or	■	■	P	S. Europe 1781. D co
Carduus paniculatus L.						
20323 monspessulanum AIL	Montpelier	^t A or	2	jl.jl	P	Montpel. 1596. D co
20324 pyrenaeicum Dec.	Pyrenean	^ A or	^	Jl.au	P	Pyrenees 1816. S co
C&rdius pyrenäicus Gou.						Jac. ob. 4.93
20325 uliginosum Bieb.	swamp	^c A or	&	■	P	Caucasus 1820. D co
20326 desertrum Fis.	desert	^ A	^	■	P	Siberia 1821. D co
20327 Aclxma Dec.	Acarna	O or	2	■	P	Spain 1683. S co
Cnlcus Acarna L.	CarDna Acárna Bieb.					Cav. ic. 1.53
III. LEUCA'NTHUM.—Leaves decurrent / flowers whitish.						
20328 nxfescens Ram.	rufescent	k A or	3	jl.au	P	Pyrenees 1816. S co
IV. ERIOPHORA.—Leaves sessile / flowers red, subsolUary.						
20329 anglicum Lob. Dec.	English	■ ^ w	3	jn	P	Britain m.pas. D co Eng. bot. 177
Carduus pratinus IV.						
20330 tuberbum All.	tuberous	■ A or	3	au.o	P	England woods. D co
Cilium TV.	ciliated	■ A or	3	au	P	Mur. 6.5.
20331 ciliatum Bieb.	prickly	■ A or	1	jl.au	P	* Baibary 1817. D co
20332 echinatum Dec.	wool-bearing	CD or	4	jl.au	P	Britain ch.pa. S co
20333 eriphorum Sco.	Cynara-hke	A, or	2	jl.au	P	Crete 1&7. S co
Cnlcus cynaroides TV.						
20335 ingrme G. Don	unarmed	■ A or	3	jl.au	P 1824. D co
Cnlcus in^rmis TV.						
20336 heteromallum Spr.	one-side-woolly	CD or	3	jl.au	P	Nepal 1820. S co
Cnlcus heteromallus D. Don						
20337 pazcuariae Spr.	Pazcuara	■ ■	■	■	P	Mexico 1827. S co
Cnlcus pazcuarensis Kth.						
20338 cernuum Lag.	drooping	■ ■	■	■	P	Mexico 1827. S co
Cnlcus nivalis Kth.						
20339 flmbn.tum Bieb.	fringed	■ ■	■	■	P	Caucasus 1816. S co
Carduus flmbnatus Bieb.						
20340 arachnoideum Bieb.	cobweb-like	■ ■	2	jl.au	P	Tauria 1818. D co
20341 hirridum Bieb.	horrid	■ ■	■	liau.s	P	Iberia 1823. S co
20342 infidum All.	intermediate	■ ■	3	jl.au	P	Piedmont 1819. S co
Cnlcus mudiis W.						Al. ped. 4J
20343 orientale Spr.	oriental	■ ■	■	jl.au	P	Asia Min. 1827. S co
Cnlcus orientalis W.						
20344 discolor Spr.	discolored	■ ■	2	jl.au	P	N. Amer. 1803. S co
20345 altissimum Spr.	tallest	■ ■	6	au.s	P	N. Amer. 1726. D co
20346 heterophyllum All.	various-leaved	■ ■	2	jl.su	P	Britain m.al.p. D co
20347 helenioides Hud.	Elecampane-lk	■ ■	6	jl.au	P	Siberia 1804. D co
20348 serratuloides Sco.	Saw-worUike	■ ■	3	jn.o	P	Siberia 1752. D co
20349 el&tius G. Don	taller	A or	6	jn.o	P 1823 D co
Cnlcus clatiior Lk.						Jac.au. 2.127

V. ERIOPHORA.—Leaves sessile / flowers red, subsolUary.						
20329 anglicum Lob. Dec.	English	■ ^ w	3	jn	P	Britain m.pas. D co Eng. bot. 177
Carduus pratinus IV.						
20330 tuberbum All.	tuberous	■ A or	3	au.o	P	England woods. D co
Cilium TV.	ciliated	■ A or	3	au	P	Mur. 6.5.
20331 ciliatum Bieb.	prickly	■ A or	1	jl.au	P	* Baibary 1817. D co
20332 echinatum Dec.	wool-bearing	CD or	4	jl.au	P	Britain ch.pa. S co
20333 eriphorum Sco.	Cynara-hke	A, or	2	jl.au	P	Crete 1&7. S co
Cnlcus cynaroides TV.						
20335 ingrme G. Don	unarmed	■ A or	3	jl.au	P 1824. D co
Cnlcus in^rmis TV.						
20336 heteromallum Spr.	one-side-woolly	CD or	3	jl.au	P	Nepal 1820. S co
Cnlcus heteromallus D. Don						
20337 pazcuariae Spr.	Pazcuara	■ ■	■	■	P	Mexico 1827. S co
Cnlcus pazcuarensis Kth.						
20338 cernuum Lag.	drooping	■ ■	■	■	P	Mexico 1827. S co
Cnlcus nivalis Kth.						
20339 flmbn.tum Bieb.	fringed	■ ■	■	■	P	Caucasus 1816. S co
Carduus flmbnatus Bieb.						
20340 arachnoideum Bieb.	cobweb-like	■ ■	2	jl.au	P	Tauria 1818. D co
20341 hirridum Bieb.	horrid	■ ■	■	liau.s	P	Iberia 1823. S co
20342 infidum All.	intermediate	■ ■	3	jl.au	P	Piedmont 1819. S co
Cnlcus mudiis W.						Al. ped. 4J
20343 orientale Spr.	oriental	■ ■	■	jl.au	P	Asia Min. 1827. S co
Cnlcus orientalis W.						
20344 discolor Spr.	discolored	■ ■	2	jl.au	P	N. Amer. 1803. S co
20345 altissimum Spr.	tallest	■ ■	6	au.s	P	N. Amer. 1726. D co
20346 heterophyllum All.	various-leaved	■ ■	2	jl.su	P	Britain m.al.p. D co
20347 helenioides Hud.	Elecampane-lk	■ ■	6	jl.au	P	Siberia 1804. D co
20348 serratuloides Sco.	Saw-worUike	■ ■	3	jn.o	P	Siberia 1752. D co
20349 el&tius G. Don	taller	A or	6	jn.o	P 1823 D co
Cnlcus clatiior Lk.						Jac.au. 2.127

20350 Casabbn* Dec.	Casabona's	■	■	S jn.au	P	S. Europe 1714.	S p.l	Schm.ic. 51,52
20351 stellatum All.	starred	Q or	S in.jl	P	Italy 1715.	S co	Tri.ob.96	
20352 mūticum Afr.	awnleas	A or	2 jl.au	P	N. Amer. 1820.	D co		
Cneus gl&her var. Nut.								
20353 virginianum Mx.	Virginian	A or	3 Jlau	P	Virginia 1824.	D co		
Carduus virginianus L.								
20354 setbum Bieb.	bristly	Q> or	■	■	Silesia 1822.	S co		
20355 arvnse Lam.	corn or way	A w	■	■	Britain ro.sid.	D co	Eng.bot.975	
Serratula arvensis L.	Carduus arvensis Sm.							
20356 serrulatum Bieb.	serrulated	■	■	4 jl.au	P	Tauria 1819.	D co	
20357 dealb&tum Bieb.	whited	■	■	3< jl.au	P	Caucasus 1810.	S co	
20358 echinoc&phalum2teo.prickly.hcaded		■	■	2 jl.au	P	Caucasus 1826.	S co	
20359 strig&sum Bieb.	strigose	■	■	2 a.s	P	Caucasus 1825.	S co	
20360 /appaceum Bieb.	Burdock-like	■	■	4 jl.au	P	Caucasus 1821.	S co	
20361 munUum W. en.	armed	■	■	3 jl.au	P	Caucasus 1816.	D co	
20362 obvallatum Bieb.	fenced	■	■	3 jl.au	P	Caucasus 1816.	D co	
20363 pinnatifidum Spr.	pinnatifld	■	■	2 jl.au	P	Spam 1820.	S co	
Crius pinnatifidus W.								
20364 ignarium Spr.	fire	■	■	2 jl.au	P	Russia 1826.	S co	
Carduus ignarius Pall.								
20365 acabile All.	stemlesj	■	■	■ Jl->	P	Britain gra.pa.	D co	Eng. bot. 161
V. AGGBEoVtA.—heaves sessile ; flowers red, aggregated.								
20366 rivulare AU.	brook	A or	3 jl.au	■	Hungary 180*.	S co	Jac. au. 1.91	
20367 salisburgense G. Don	Salisbury	A or	3 jl.au	■	Europe 1816.	D co		
Cficus salisburgensis IV.								
20368 paludosum G. Don	marsh	A or	3 jl.au	P	Switzerl. 1819.	S co		
Chicus paludosus SchL								
■ giganteum Spr.	giant	■ Q) or	6 jl.au	P	Sicily 1820.	S co		
Carduus giganteus Desf.	CMcus firmus Pres.			■ jl.au	P	Hungary 1816.	S co	W.&K. 3.161
20370 pauciflbrum Spr.	few-flowered	■	■	■ jl.au	P			
Crius pauciflcrus Kit.								
20371 niite Fis.	meek	■	■	■ jl.au	P	Siberia 1820.	S co	
20372 org'ale Spr.	six feet	■	■	■ jl.au	P 1823.	D co	
20373 Hallerii Hort.	Haller's	■	■	■ jl.au	P	S. Europe 1816.	D co	
£0374 squarrsum Fis.	squarrose	■	■	■ jl.au	P	Siberia 1818.	D co	
20375 incanuin Bieb.	hoary	■	■	■ jl.au	P	Caucasus 1820.	S co	
20376 ambiguum All.	ambiguous	■	■	■ jl.au	P	Mt.Cenis 1820.	D co	
20377 afruraX>cc.	African	■	■	■ n.jl	P	Barbary 1800.	S co	Bot. mag. 22S7
20378 diaenanthum Dee.	two-spined	■	■	■ n.jl	P	Syria 1800.	S co	Lab. syr. 1.2
VII. CHRYSA'NTHA.—Leaves sessile ; flowers yellow or white.								
20379 olerticum All.	oleraceous	■	■	3 jl.au	Pa.Y	Europe 1370.	D co	Fl. dan. 860
20380 tart&licum AIL	Tartarian	■	■	■ Ujl.au	IV	Siberia 1775.	D co	Jac. au. 1.90
20381 ferox Dec.	fierce	■	■	■	W	S. Europe 1683.	S co	Al.ped. 1.50
20382 syracum Gae.	Syrian	■	■	■	W	Levant 1771.	S co	Cam. h. 10
20383 marcoticum Spr.	Mareotic	■	■	■	W	Egypt 1827.	S co	
Cdthamus marc&ticus Del.								
20384 Bertol&ntu Spr.	Bertoloni'i	■	■	■	■	Italy 1820.	S co	
Cficus hurfidus Bert.								
20385 bse'ticum Spr.	Bastic	■	■	■	■	Spain 1824.	S co	
h&rridus Lag.								
20386 rigens Spr.	stiff alpine	■	■	■	Y	Switzerl. 1775.	D co	Ac.hel.4.16
20387 leucocyphalum Spr.	white-headed	■	■	■	Y.w	Crete 1816.	D co	
Crikus leucoclyphalus IV.								
20388 hf bridum G. Don	hybrid	■	■	■	■	Pa.Y Switzerl. 1819.	D co	
Cnclus h'bridus SchL								
20389 ochroliticum AU.	pale-yellow	■	■	2 jl.	Pa.Y	Switzerl. 1801.	D co	
20390 Erisithales L.	Erisithales	■	■	3 jn.au	Pa.Y	France 1752.	D co	Jac. au. 4.310
2U391 glutinosum Dec.	glulinous	■	■	2 jl.au	Pa.Y	S. Europe 1816.	I) co	
20392 lacteum SchL.	m\k.colored	■	■	■ jl.au	Pa.Y	Switzerl. 1819.	D co	
20393 scmipectin'tum Led.	half-pectinated	■	■	3 jlau	Pa.Y	Siberia 1819.	D co	
20394 carmolicum Sco	Carmolan	■	■	2 jlau	Pa.Y	Carniola 1792.	I) co	Srop.car.100r.r2
20395 spinosissimum Sco.	spiniest	■	■	3 jn.au	Pa.Y	Switzerl. 1759.	D co	Bot. mag. 1366
20396 scleranthum Bieb.	hard-flowered	■	■	2 jlau	Pa.Y	Caucasus 1820.	S co	
CarllnaEchinus Bieb.								
VII. DUX'BIA.—Doubtful to which of the sections they belong.								
20397 angulatum SchL.	angled	^ A or	2 jl.au	P	Switzerl. 1819.	D co		
20398 nudiflbrum SchL.	naked-flowered	& A or	2 jl.au	P	Switzerl. 1817.	I) co		
20399 Cervini Thomas	Cervini's	5 A or	2 jlau	...	Switzerl. 1825.	D co		

2244. ERYTHROLA^NA Swt. ERYTH. (*Erythros*, red, *tena*, cloak; scales of calyx.) Comp.Card. 1.—C
20400 conspicua Swt. conspicuous ^ Q) or 8 s.o Pa.Y Mexico 1825. S r.m Sw. fl. gar. 134

2245. ONOPO'R'DUM L. COTTON THISTLE. (*Onos*, ass, *perdo*, pedo; effects.) Comp. Card. 13.—14.

I. CAULEBCE'NTIA. Having stems.								
50401 jfegnthium L.	Acanthium	Q) or	6 il.au	P	Britain gra.ba.	■ eo	Eng. bot. 977	
20402 tafricum W.	Taurian	Q) or	12 jl.au	P	Tauria 1800.	■ co		
20403 vlrns Dec.	green	Q) or	■ jlau	P	Montpel. 1818.	■ co		
20404 viscosum Schr.	clammy	Q> or	■ jlau	P	S. Europe 1818.	■ co		
20405 macract&thnum Sehous	long-spined	Q) or	■ jlau	P	Barbary 1798.	■ co	Schou. 5	
20406 illyricum L.	Illyrian	jt Q) or	■ jlau	P	S. Europe 1648.	■ co	Jac. vii. S. 148	
20407 elitum Sm.	tall	5 CD or	■ jlau	P	Greece 1816.	■ co		
20408 gra^cum Gou.	Grecian	O or	10 n.jl	P	Levant 1799.	■ co	Gou. il.25	
20409 cynaroldcs Stev.	Artichoke-liko	2t Q> or	10 ■	P	Caucasus 1823.	■ co		
20410 arJibicum L.	Arabian	Q) or	8 ■	P	S. Europe 68G.	■ co	Jac vin. 2.149	
II. ACAU'LIA. — Stemless.								
20411 acatilon L.	fitcmless	3t Q) cu	■ jlau	W	Pyrenees 1739.	■ co	Jac. ic. 1.167	
20412 uniflfrum Cav.	one-flowered	jt Q) cu	1jlau	W	Spain 1626.	■ co		
20413 pyrcn&cum Dec.	Pyrencean	^ Q> cuy i	jlau	W	Pyrenees 1820.	■ co		

2246. BERA'R'DJ^ Vil. BERAADIA. {M. Berard, a botanist of Grenoble.) Comp. Card. 1.
20414 subaca'lis Fil. sub-stemless ■ lljl.nu P Italy 1791. ■ co Vil. del. 3.22
jl'rectum lanuginbnum Dec. Onopordum rotundifolium All.

- 2247. CVNARA L. ARTICHOKE. (*Kyon, dog; spines of involucrum like dogs' teeth.*) *Camp. Card.* 8.
 20415 *fc61ymus* W. *Scolymus* cul 8 au.s P s. Europe 1545. D co Black. 548
 20416 *horrida* H. K. *horrid* 6 au.s P Madeira 1778. 1) co
 20417 *ftrox* *Ten.* *fierce* A or *jlau* B Italy 1840. D r.m
 20418 *spinosisima* *Presl* *spiniest* A or *jlau* B Sicily 182a D r.m
 20419 *Cardunculus* *L.* *Cardoon* A cul 5 au.s B Candia 1658. D co Tab. ic. 696
 20420 *integrifolia* *Vahl* *entire-leaved* A cu 4 *jlau* B Spain ... D r.m
 20421 *humilis* *L.* *humble* A <n *jlau* B Spain 1613. D co Pluk.al.81.2
 20422 *pygma*^a W. *pygmy* A or *jlau* P Spain 1820. D co
2248. CARLTJM L. CARLINB THISTLE. (*Charlemagne; preserved his army from plague.*) *Comp. Card.* 11.-18.
 20423 *acaquifolia* AU. *ACanthus**^{lvd} or ² *jl** W Carniola 1816. co Al.ped. 54
 20424 *acaulis* L. *stemlesa* or *fjnl* W Italy 1640. D co Kno.th.2.1
 2 caulicēns Schl. *caulescent* or 1 *jnjl* W Switzerl. 1819. D co
 20425 *simplex* Kit. *clustered* A °f *jnjl* W Hungary 1816. D co W.&K.2.152
 20426 *aggregata* W. *woolly* Q or 3 *jnjl* p S. Europe 1682. S co Gar. aix 21
 20427 *lanata* /. *corymbosa* A or 3 *jlau* Y S. Europe 1640. D co Col. ec. 1.27. 1
 20428 *corymbosa* L. *common* Q w *ljnl.8* PY Britain dry pa. S co Eng. bot. 1144
 20429 *vulgāris* L. *racemosa* O or 3 *jnjn* Y Spain 1638. S co
 20430 *racemosa* L. *racemose* 3 *jnjn* Y Spain 1638. S co
 20431 *Bleberstein*^a AnaBcjnh. Bleberstein's^t A or ? *Jls* — Caucasus 1816. D co
 20432 *sicula* *Ten.* Sicilian ^a CD or 1 *jlau* — Sicily 1327. S co
 20433 *brartoata* *Presl.* *lyrate* £ Ol or — C.G.H. 1816. S s.l
 2249. ATR A'CTYLIS L. ATRACTTLIS. (*Atraktos, spindle; light stems fit to make.*) *Camp. Card.* 1—2.
 2(434 *hūmilis*Z. *humble* ^a A un 1 *jnjl* W Spain 1759. D e o Cav. ic.1.54
2250. jfCA'RNA TV. ACARNA. (In Theophrastus for a plant resembling a thistle.) *Camp. Card.* 2.—&
 2C435 *gummifera* IV. *gum-bearing* ^a ^a un f *jnjn* P & Europe 1640. D co Cav. ic. 3.228
 i0436 *cancellata* W. *latticed* O un J *jnjl* B S. Europe 1640. S co Lam. il 662.1
ylractylis cancellata L.
2851. STOKE^SIil Herit. STOKESIA. (*Jonathan Stokes, M.D, coadjutor of Withering.*) *Camp. Card.* 1.
 20437 *cyanea* Herit. *azure-flowered* £ LAJ pr 2 au B Carolina 1766. D co Her. ser. 27.38
- 2252, STOBiEM Thun. STOB*:A. (*Dr. Stobceus, of Lund, one of Linneus's first patrons.*) *Comp.-Card.* 1.—11.
 20438 *pinnkta* Thun. *pinnate* tt. i_j or 2 year Y C. G. H. 1812. C co Bot mag. 1788
- 2S53. ONOBRCTMA Gae. OWOBROMA. (*Onos, ass, brotna, food; grateful to.*) *Comp. Card.* 5.
 20439 *ceruleum* Gae. *azure-flowered* A or 1 *jnjl* B Spain 1640. D co Bot. mag. 2293
Cdrtamus cacruleus L.
 20440 *arborescens* Spr. *arborescent* 1 lor 6 *jlnu* Y Spain 1731. C s.p
Cdrtamus arborēscens L.
 20441 *cynaroldes* Spr. *Cynara-like* A or 2 *jnjl* W Caucasus 1820. S co
Cdrtamus cynaroldes Bieb.
 20442 *leucocafilon* Spr. *white-stemmed* ^a A or 1 *jnjl* W Greece 1800. D co
Cdrtamus leucocaulis Sm.
 20443 *gladicum* Spr. *glaucus* O or *ljnl.au* P Tauria 1817. S s.p
Cdrtamus glaucus Bieb.
2254. CARLOWI'Z/l/* Moen. CARLOWIZIA. (*Carlowiz, some obscure botanist.*) *Comp. Card.* 1.
 20444 *jalicitlia* Moen. *Willow-leaved* m *J* or 3 au Y Madeira 1784. C 81
- S&55. CA'RTHAMUSh. CARTHAMUS. (*Oortom, to paint, its Arabic name; flowers.*) *Comp. Card.* 2.—20.
 20445 *tinctbrius* L. *dyer's* or 3 *jnjl* O Egypt 1551. S s.l Bot. ree. 170
 20446 *Oxyacantha* Bieb. *sharivspined* O or 3 *jl.au* Y Caucasus 1818. S co
2256. HERACA'NTHA Lk. HER. (*Heros, nohlc, akantha, thorn; beauty.*) *Comp. Card.* 4.—6.
 20447 *armenia* G. Don *Armenian* *Cdrtamus armtnius* W. O or S! *jnau* Y Armenia 1816. S co
 20448 *lanata* Lk. *woolly* O or 2 *jl.au* Y & Europe 1596. S co Bot. mag. 2142
Cdrtamus lanatus L. *Cirsium lanatum* Dec.
 20449 *cretica* Lk. *Cretan* O or 2 *jnjl* W Candia 1731. S co
Cdrtamus crticeus L.
 20450 *taGrica* Lk. *Taurian* O or 2 *j'.au* Y Caucasus 1818. S co
Onobroma denfatum Spr. *Cdrtamus tadricus* Bieb.
2257. CARDUNCE'LLUS Dec. CARDIINCELLUS. (A diminutive of *cardunculus*, cardoon.) *Comp. Card.* 2.
 20451 *mitissimus* Dec. *meekest* £k A or \$ *jnjl* B France 1776. D co
Cdrtamus mitissimus L.
 20452 *vulgāris* Dec. *common* ^a A or f *myjn* B France 1734. D co
Cdrtamus Carduncillus JV.
2258. CARDCPATUM J. CAROOPATIM. (*Carduus, thistle, pa tos, beaten path ; habitat.*) *Comp. Card.* 1.
 20453 *cymbosum* Pers. *cymbose* f A <n 3 *jl.au* B Levant 1821. D co M.h.7.3S.17
2259. STAHELPNif Lag. STADBLINA. (*Benedict Stathelin, a Swiss botanist*) *Comp. Card.* 3.—13.
 20454 *dxbia* L. *doubtful* a *fra* 3 *jnjl* P S. Europe 1640. C p.l I*m. il. offil 4
 20455 *arborescens* L. *arborescent* I pr 6 *jl.s* P Candia 1739. C p.l Schreb. dec. 1. 1
 20456 *Chanurpeace* L. *Ground-Pine-urf** *Jpr* 2 *jl.n* P *Candia* 1660. C II^a Pluk.al.94.3
Pteronia Chamepebe Spr.
2260. PALAFO'XJJ Lag. PALAFOXIA. (*Palafbx, a Spanish general.*) *Camp. Eupatbrear.* I.
 20457 *linearis* Ziff. *linear-^a^d* jf lAjpr 2 *jnjl* W Mexico 1821. 8 co Bot mag. 2132
Stéfve hnearis W.
- 2261. PTERO^NIA L. PTBRONIA. (*Pteron, a wing; feathery scales of receptacle.*) *Comp. Card.* 9.—33.
 20458 *camphorata* A. *camphor-uctd* * *i* J or 3 *jnjl* Y C. G. H. 1774. C p.l Pluk.m S45 2
 20459 *stricta* Thun. *strict* * *l* J or 3 *ap.jn* Y C. G. H. 1774. C p.l
 20460 *flexicaO* Us Thun. *bending stalked** LJ or 3 *jnjn* Y C. G. H. 1812. C co
 20461 *oppositifolia* /,. *opposite-leaved** LJ or 4 *jl* Y C. G. H. 1774. C p.l Brey. pr. 17. 3
 20462 *scaribsa* Thun. *scarious* a i_J or 2 *jnavi* Y C. G. H. 1815. C co

20463 echinata Thun.	echinato	■	2)n.au	Y	C. G. H.	1818.	C	p.l
20464 fasciculata Thun.	fascicled	*■Jor	2	jnjj	Y	C. G. H.	1816.	C	p.l
20465 glomerata Thun.	glomerate	*■Jor	2	jnj	Y	C. G. H.	1817.	C	p.l
20466 pallens Thun.	pale	**T_j or	■	jnj.l	Y	C. G. H.	181ft	C	p.l

* 2262. VERNONIA Schreb. VERNONIA. (*William Vernon*, bot. trav. in N. Araer.) *Comp. Card. 13—44.*

I. CORYMBUSJB.— <i>Flovers vaginatlate or corymbose; herbaceous.</i>									
20467 noveborac&ua W.	New York	T	▲ off	Tlrl	P	N. Amer.	1710.	D	co
20468 prsalta JV.	very tall	■ A or	8	an	p	N. Amer.	1732.	D	co
20469 alt&sima Nut.	tallest	■ A or	12	s.n.	P	Ohio	1820.	D	co
20470 angustifolia Mx.	narrow-leaved	■ A or	4	s.n.	P	N. Amer.	1817.	D	co
20471 glauca JV.	glaucous-leaved	■ A or	4	<n.	p	K Amer.	1710.	D	co
20472 scaberrima Nut.	most scabrous	■ A or	4	s.n.	P	N. Amer.	1824.	I	co
20473 pandurata Jac.	fiddle-leaved	■ A or	4	s.n.	P		1825.	D	co

II. SPICATIFORMES.—*Herbaceous, * flowers sessile, spicate.*

20474 linearis Spr.	linear	—Q or	1	il	IP	S. Amer.	1H25.	\$	s.l
20475 nericea Rich.	silky	tf	^	E3 or	s d	Pa.P	Brazil	1823.	S co
20476 flexuosa B. M.	flexuose	js	[f]	or	1J s	P	Brazil	1823.	S co
20477 arborescens Sim.	arborescent	• CB	or	5	n.d.	P	Jamaica	1733.	C co
20478 odoratissima Kth.	sweetest-scented	* O or	4	au.n	P	Caracas	1817.	C co	Pl. am. 10.130.2
20479 fruticosa SUM.	shrubby	• o r	4	au.n	P	W. Indies	1818.	C co	Pl. ic. 95.1

2263. ASCARICPDA Cass. ASCARICIDA. (*Ascaris*. an intestinal worm, *eedo*, to kill.) *Comp. Card. Fern. 2.*

20480 anthelmintica Swt.	anthelmintic	O or	1	aus.	P	K Indies	1770.	S	co
Vernonia anthelmintica L.									
20481 triplinervia G. Don	triple-nerved	O or	1	n'	P	Brasil	1825.	S	co

Vernbnja tripliervia Spr.

226*. AMMOBIUM B. Br. AMMOBIUM. (*Anwios*, sand, *bio*, to live; habitation.) *Cqptp. Card. Fern. 1.*

20482 alatum R. Br.	winged	£ AJ	pr	2	mr.s	W	N. Holl.	1822.	S	co
20483 plantagineum G. Don	Plantain-Zyd	£ AJ	or	1	aus	W	N. Holl.	1827.	C	p.l

2265. LIJITRIS Gae.

LIATRIS.

(Meaning unknown.)

Cdmp. Card. Fern. 17. — £5.

I. TuBERRHIE.—*Flowers spicate or racemose; roots somewhat tuberous.*

20484 squarrbsa W.	tough-cupped	■	■	■	B	N. Amer.	1732.	D	■
20485 scandens IV.	scious-cujMvi	■	■	■	P	N. Amer.	1739.	D	p.
20486 sphroidea Mx.	■	■	■	■	P	N. Amer.	1817.	D	co
20487 elegans H. W.	elegant	■	■	■	P	N. Amer.	1877.	D	p.l
20488 turbinata Swt.	turbinate	■	■	■	P	N. Amer.	1823.	I	co
20489 tenuifolia Nut.	fine-leaved	■	■	■	P	Carolina	1820.	D	p.l
20490 pilosa W.	hairy-leaved	■	■	■	P	N. Amer.	1783.	D	p.l
20491 gracilis Ph.	slender	■	■	■	P	Carolina	1818.	D	Bot reg. 595
20492 spicata W.	long-spiked	■	■	■	P	N. Amer.	1732.	I	p.l
20493 cylindracea Ms.	cylindr'ic-cupped	■	■	■	Pk	N. Amer.	1811.	I	co
20494 intermedia B. R.	intermediate	■	■	■	P	N. Amer.	1823.	D	p.l
20495 heterophyllum Ph.	various-leaved	■	■	■	P	N. Amer.	1790.	I	p.l
20496 pyrenostichya A/x.	dense-spiked	■	■	■	P	N. Amer.	1732.	D	co
20497 pimila Lod.	dwarf	■	■	■	P	N. Amer.	...	R	s.p

II. FIBRUSA.—*Flowers corymbose or paniced; roots fibrous.*

20498 odoratissima W.	sweetest-scented	£ AJ	el	3	au.o	P	Carolina	1786.	D	s.p
20499 paniculata W.	paniced	^ Al	or	3	au.o	P	Carolina	1826.	D	p.l
20500 corymbosa Nut.	corymbose	^ Al	or	3	au.o	P	Carolina	1825.	D	p.l

2266. MIKA'NJM W.

MIKANIA.

(Joseph Mikan, prof. bot.

Prague.) *Comp. Eupatbrete. 9.-35.*

20501 houstonia W.	Houston's	■ Dun	8	jl.au	Pk	Jamaica	1733.	C	co	
20502 haitata W.	■	■	■	■	■	Jamaica	.	C	co	
20503 ainra W.	bitter	■	■	■	■	Guiana	1816.	C	co	
20504 suaveolens Kth.	sweet-scented	■	■	■	■	S. Amer.	1823.	C	co	
20505 onnocensis Kth.	Orinoco	■	■	■	■	Orinoco	1826.	C	co	
20506 scandens JV.	climbing	■	■	■	■	Pa.B	N. Amer.	1714.	C	co
20507 Gudco Hum.	Guaco	■	■	■	■	W	S. Amer.	1823.	C	co
20508 chenopodi folia W.	Chenopod.-lyd	J	123	un	■	W	Guinea	1824.	C	co
20509 opifera Mart.	opium-bearing	%	■■■	6	au	W	Brazil	1823.	C	co

2267. SP ARG ANO'PHORUS Gae. SPARC. (*Sparganon*, fillet, *phoreo*, to bear; seed.) *Comp. Eupat. 3.-4.*

20510 Vailianto Gae.	Vailiant's	O un	■ J	au	Y	India	1823.	S	co
20511 verticillata Mx.	■	■	■	■	■	Carolina	.	C	co
20512 Struchium Swz.	Struchium	■	■	■	■	Jamaica	.	C	co

Br. jam. 34.2

* 2268. EUPATO'R/r/Af L. EUPAT. (*Muhridates Eupator* used it arcotinter-poison.) *Com Eupat. 55.—143.*

I. BACCUAROIDEA.—*Calyx oblong cylindrical, imbricated, with many series of scales; shrubby.*

20513 triflirum Aub.	threeflowered	%	• un	4	au	W	Guinea	1823.	C	Aub. gu. 2.314
20514 iaccharokles Kth.	Baccharis-hkc	■	O un	3	...	P.T	NewGran.	1820.	C	p.
20515 conhexum Fa/it	coriaceous	i	Q un	3	au	...	Hispaniola	...	C	p.
20516 macranthum Swz.	lofii, NEWGAE TA	■	■	■	■	AT	AT	AT	C	p.l
20517 repandum W.	repanel	■	L	11	3 au	W	W. Indies	...	C	p.l
20518 populifolium Kth.	Poplar-leaved	■	un	4	au	W	Mexico	1826.	C	pi
20519 pulchellum Kth.	neat	tf.	Q un	2	aus	Mexico	18-6.	C	p.	
20520 floribundum Kth.	bundle-flwd	n.	CD or	1	au	■	S. Amer.	1K23.	C	p.
20521 chama'dryfolium A/J	(ermander-lytt.	• or	1	au	»	S. Amer.	1822.	C	p.	
20522 veronicae>uaiKth S)	eedwclJvd	tt.Qor	IJau	B	Mexico	1895.	C	p.l	H.&B.4.341	
20523 Dalea L.	Dalea	«	• or	6	au	Pk	Jamaica	1773.	C	Jac sc. 2.146
20524 acuminatum D.Don	acuminate	*	i_J or	2	au	..	Nepal	1819.	C	p.l
20525 nepalense Spr.										
20525 odordium L.	sweet-scented	11	• fra	3	au.o	Pk	Jamaica	1752.	C	Pluk. aL 177.3

II. IRESI.VOI1>EA. — *Calyx oblong, cylindrical, imbricated, with many series of scales j herbaceous*i. PAUCIFLORA.—*Calyx few-flowered*

20526 iresinoldcs <i>Kth.</i>	Iresine-like	1. (23 or tallest	2 au. _{no} St A or	W Pk	N. Grenad. 1820. D co	H.&B.n.34O
20527 alfafasimum <i>L.</i>					N. Amer. 1699. D co	Jac. vin.2.164
20528 lanceofatum <i>W.</i>	lanceolate	5 A or	3 au.e	W	N. Amer. 1819. D co	
20529 album <i>L.</i>	white	i A "	2 au.s	W	N. Amer. 1820. D co	

ii. MULTIFLORA.—*Calyx many-flowered j flowers corymbose or paniced.*

20530 purpureum <i>Z.</i>	purple-stalked	A or	s.o	Pk	N. Amer. 1640. D co	Cor. ca. 72
20531 /amiflrium <i>Kth.</i>	Dead-Nettle-lvd	(29 or	au.s	Pk	... 1823. D co	
20532 macrophjrrillum <i>L.</i>	long-leaved	£ ES or	jl.au	W	Jamaica 1823. D co	PL am. 10.129
	molle Sun.	4geritum guianense Auh:				
20533 <rtica?folium <i>Mutls</i>	Nettle-leaved	AJ or	1J jLau	Pk	S. Amer. 1803. D co	Sm. ic. in. 63
20534 squarrbus <i>Cav.</i>	squarrose	iAI or	1J jLau	P	Mexico 1827. D co	
20535 nutans <i>Kth.</i>	nodding	LAJ or	2 -	Pk	Mexico 1827. D co	
20536 /wei.lium <i>L.</i>	Iva-leaved	CZ3 or	3 jn.jl	Pk	Jamaica 17>4. D co	
20537 <alviel5lium <i>B. M.</i>	Sage-leaved	A or	4 au.s	W	N. Amer. 1814. D co	Bot. mag. 2010

III. DAV.EA. — *Calyx subimbricated, with few series of scales, or simple ; shrubby*i. PAUCIFLORUM.—*Calyx few-flowered.*

20538 parviflorum <i>Swz.</i>	small-flowered	*L • un	2 jLau	...	Jamaica 1826. C co	
-------------------------------	----------------	---------	--------	-----	--------------------	--

Dalea Hort.

ii. MULTIFLORA.—*Calyx many-flowered.*

20539 monanthum <i>Swz.</i>	mountain	tt.	• or 2	jLau	...	Jamaica 1820. C co
20540 canescens <i>Vahl</i>	canescent	«L Q or	3 jn.jl	...	Jamaica 1821. C pl	
20541 villosum <i>Swx.</i>	villous	tt. • or	2	.	Jamaica ...	C pl
20542 xalapense <i>Kth.</i>	Xalapa	a. CZ1 or	3 jl.au	W	Mexico 1826. C pl	

IV. LUXURIA'NTIA. — *Calyx, subimbricated, with few series of scales, or simple j herbaceous.*i. PAUCIFLGRA.—*Calyx few-flowered.*

20543 coronopifolium <i>W.</i>	Coronopus-lvd	3	jl.s	W	Carolina 1824. D co		
20544/cniculaceum <i>W.</i>	Fennel-leaved	4	jn.s	Pa.Y	N. Amer. 1807. D co		
20545 Aytssopiflum? <i>?</i>	Hyssop-leaved	1	aus.s	W	N. Amer. 1699. 1) co	Di. el. 115.140	
20146 lineariflum <i>Walt.</i>	linear-leaved	1	aus.s	W	N. Amer. 1830. D co		
20547 Fra'ser; Poir.	Fraser's	lja	W	Carolina 1820. D co			
20148 sessilifolium <i>L.</i>	seabile-leaved	1 s.o	W	N. Amer. 1777. D co			
2(549 trucatum IV.	truncate	1*	W	N. Amer. 1800. D co			
20550 teucTifflum <i>W.</i>	Teucrium-lvd	W	W	N. Amer. 1816. D co	W.h.b.32		
20551 meliso'ides <i>W.</i>	Melissa-Jike	2 au.n	W	N. Amer. 1811. D co			
20552 cordatum <i>Walt.</i>	2 s	W	N. Amer. 1699. D co	Pluk.al.88.4			
20553 rotundiflum <i>L.</i>	round-leaved	jl.au	W	N. Amer. 1824. D co			
20553 ceanothifolium <i>W.</i>	Ceanothus-lvd	a.us	W	... 1824. D co			
20554 pubescens <i>W.</i>	pubescent	jl.au	W	N. Amer. 1819. D co			
20555 cannabinum <i>L.</i>	Hemp Agrimony	jlLo	Pk	Britain wnt.pl. D co	Eng. bot. 428		
20556 triohatum <i>L.</i>	three-leaved	au.o	p	N. Amer. 1768. D co			
20557 syracum <i>L.</i>	Syrian	jl.s	W	Syria 1807. D co	Jac. ic. 1.170		
		Calyx	P man^flowered kmerm ^	D co			
20558 verticillatum IV.	hpottcA-stalked	3 au.s	P	N. Amer. 1656. D co	Her. par. 158		
20559 macuftum <i>L.</i>	dotted	4 aiLS	P	N. Amer. 1815. D co			
20560 punctiitum VV.	per foli ate	2 nu.o	W	N. Amer. 1691. D co	Pluk.al.87.6		
20561 perfoliatum <i>L.</i>	aromatic	4 jl.au	W	N. Amer. 1739. D co	Pluk.al. 88.3		
20562 aromaticum <i>L.</i>	Ageratum-like	4 au.o	W	N. Amer. 1640. D pl	Cor. ca. 21		
20563 ugeratoldes <i>L.</i>	late-flowering	3 o	V	N. Amer. 1824. D co			
20564 seritimum <i>Mx.</i>	deltoid	3 jlau	P	S. Amer. 1822. D co			
20565 deltofclum <i>Jac.</i>	climbing	6 au.s	Y	S. Amer. 1821. D co			
20566 scandens <i>Lk.</i>	panicled	23 or	6 au.o	Pk	S. Amer. 1818. D co		
20567 paniculatum <i>Mx.</i>	paniculata Hort.						

*2269. KITHN//J L. KUHNIA. (*Adam Kuhn*, of Pennsylvania, pupil of Linnaeus.) *Comp.Eupatbre<e. 9..—tI*
20568 eupatoriolirrit *L.* Eupatorium-lk ^ A or 1J jLau W N. Amer. 1812. D p.l L. f. dec. 11

Critonta eupatorioides Gac.

20569 Critonta *W.* Critonia ^ A or 1} jn.au W N. Amer. 1816. D p.i Gac. fr. 2.174

Critonta Kühnii Gac.

2270. DUMERI'L'il Lag. DUMERILIA. (*M.A. M. Constant DumerU*, a French naturalist.) *Comp. Eup. t.*
20570 paniculata Lag. panicled ** „ „ D o r 3 au P Colombia 1825. C 'co An.rau. l'j.7

2271. AGE'RATUM <i>L.</i>	AGERATUM. I pi priv. ,geras, old age; because it never grows old.)	<i>Comp. Eup. 6.-10.</i>	
20571 conyzoldcs <i>L.</i>	Conyz.i-like O or 1 jl.au	L.B America 1714. S pl Sclik. han.3.238	
20572 werCleuni Pars.	azure iQJ or 1 jn.au	B W. Indies 1800. S co Hook. ex. fl. 15	
20573 latiflum <i>Cav.</i>	broad-leaved O or Ii jl.au	W Peru 1800. S co Cav. ic. 4.357	
20574 strfctum <i>B. M.</i>	strict O or 2 jn.jl	W Nepal 1821. S co Bot. mag. 2410	
20575 mexicAimra <i>B. M.</i>	Mexican O or Jjljl B Mexico 1822. S co Sw. fl. gar. 346		
20576 angustiflum <i>Spr.</i>	narrow-leaved iQJor 1 jn.au	Ma Video 1827. S co	

2272. CIELESTINA Cass. CJELESTINA. (*Calestis*, celestial; sky-blue flowers.) *Comp. Eup. 3.*

20577 rtgeratoldes Cass. Ageratum-lk jc IAJ or 1 jl.o B New Spain ... C co

20578 ca?ulea Cass. azure £ AJ or 1 jl.s B N. Amer. 1732. D pl Di. el. 114.139

Eupat6rium oer&lcum *L.*

20579 tncr/lnta Spr. small-flowered tt. _ or H A* B ^ Amer. 1800. C p I Bot. mag. 1730

Eupat6rium micrijnithum Lag. Ageratum caelestlnum *B. M.*

• 2273. ST&VIA Cav. STEVIA. (*Peter James Esteve*, M.D. prof, bot Valencia.) *Comp. Eup. 26.-38.*

1. FavncbsJR.—*Stems sub-shrubby.*

20580 ralicifrlia Cav. Willow-leaved £ Al pr llau s Pk Mexico 1803. D pl Cav. ic. 4.3S4

20581 subpubescens Lag. subpubescent jg iAI or 2 jl.s Pk N. Spain 1820. D pl

90562 lacida Lag. shining £ iAI or 2 jl.s Pk N. Spain 1824. D pl

20583 angustilbua Kth. narrow-leaved £ AJ or 1 jl.s Pk Mexico 1823. D pl

II. HERBA*CEA —*Stems herbaceous.*

20584 pubescens Lag. pubescent k -Al P' H Jl.s P Mexico 1823. D co

80585 trifida Lag. infld ^c ^ or 1 jl.s W N. Spain 1827. D pl

90586 incanescens Lag. incanescens 5t A) or 1 jl.B W ? N. Spain 1827. D pl

canescens Kth.

20587 tomentbsa Kth.	woolly	or	jljl.s	V	Mexico	1824.	D	p.l
Monarda-lvd	or	l/jl	V	Mexico		D	p.l	
leaked			W	N. Sjwin	1826.	D	p.i	
		2	a.u.s	W	Mexico	1816.	D	s.p
				W	N. Spain	1823.	D	
20592 S-a6 Cns ^	Al or	1j	a.u.s	W	N. Spain	1824.	D	p.l
20593 rhombifltha Kth.	K-Alor		jl.s	W	Mexico	1827.	D	p.
»W ? et A 5 Ua / / A	3. Alor		jl.s	W	Mexico	1824.	D	p.l
Sage-leaved	* Al or	%	jl.s	W	Mexico	1827.	D	p.l
three-leaved	S-Alor		jl.s	W	Mexico	1824.	D	p.l
Eupatoria	S-Alor		jl.s	Pk	Mexico	1798.	D	Bot. mag. 1819
Hyssop-leaved	Al or	1	a.u.s	Fk	Mexico	...	D	co Bot. mag. ISril
Iva-leaved	5 Al or	2	Jls	W	Mexico	1816.	D	s.L
lanceolate	2 Al or	1	Jls	p	Mexico	1822.	D	co!
saw-leaved	2 Al or	1	Jls	F	Mexico	1799.	D	s.ifc Jac. ic. 3.300
purple	3 Al or	1	Jls	P	Mexico	1812.	D	Co? Bot. reg. 93
pilose	K Al or	ls	a.u.s	Pk	Mexico	1820.	D	p.l
viscid	* Al or	jl.s	1'	Mexico	1821.	D	p.l	
callous	ek A or	jl.s	R	Mexico	1826.	D	H. & B. n.6. 351	

■■■■■ ALOTHORA Cav. CEPIHALOPHORV (Kephale, head, pAorco, to bear; flowers.) Comp. Anth. 1.
 KWglauca Cfl. glaucous O un 2j.l.au Y Chile 1798. D e o Cav. ic. 6.599
 *ymenopappu8 glauc Spr. Grafmia aromatica Hook.

2275. AMPHE'REPHIS Kth. AMPIKREPHIB. (Amp'tereophis, well covered; double invol.) Com Card. Vern. 3.
intermedia Lk. intermediate O pr If jl.au P Brazil 1821. S co Sw. fl. gar. 225
aristata Mu. awned O or i jl.au P Caracas 1824. I i H.&B.n.&3d4

on?*? HYMENOPAPPUS/fert. HYMEVOPAPPUS (Hymen, membrane, pappos, pappus) Comp. Hel. 3.-5*
 22J0 tenuiff,jiu8 PA. slender-leaved jf jQ un 2jn.a.u W Louisiana 1811. S co
 gij11 scabiosas^us Herit. Scabiosa-like O cu 1jl.s W Carolina ISM. S B p Jo. h. n. 1. ic
 *¹S peditus Lur. pedate Q cu 1- jl.s W S. Amer. 1803. S co Bot. mag. 2040
 Steviapedita Cnw.

■■■■■ POLYTTERIS Nut. POLYPTERIS. (Polys, many, pteron, a wing.) Comp. Sup. 1.
intcgrifolia Nut. entire-leaved O cu 1 jlau W Georgia 1823. S co
Hymenopappus integrifolius Spr.

■■■■■ 2278. ALC^MIA Kth. ALOMIA. (A, privative, loma, a fringe.) Comp. Eup. 1.
 agratolda Kth. Ageratura-like jf iAI or 1jjl.au W N. Spain 1824. C s.l H.&B.4. 354

■■■■■ 0*279. MELANANTHE^RA Mx. MELANANTHIKA. (Melas, black, anthera, anther.) Comp. Helidnfh. 2.—5.
 *81S hastata Mx. hasUte £. AI un 2 jn.jL W N. Amer. 1732. D co Di. el. 47.55

Bldeis nivea L.
 2n/i/-i Pandurata fiddle-leaved £ ^1 un 2 jnjL W N. Amer. 1732. D co Di. cl. 4! £4
 *UWIG doltoidea Mx. deltoid-leaved jf OfL un 3 jl.au Y S. Amer. 1799. S co Jac. ic. 3.583
 Calea Aspera Jac.

■■■■■ 1^SJO- MARSH A'LL/il Ph. MARSHALLIA. (Humphrey Marshall, a botanical author.) Owy. //<*. 3.—5.
 lanqeolita/A. uquear./fwrf ik ^J pr 1] jnjL P Carolina 1812. D e o
 latitudPA. broad-leaved ^ J pr lijn.jL Pa.P Carolina 1806. D e o Micam. 2.43
 angustifolia Ph. narrow-leaved £-A1pr 2 jl.s P Carolina 1800. D p.l

■■■■■ L* SPILA'NTHES L. SPILANTHIES, (Spilos, spot, anthos, flower; brown disk of fls.) Comp. Hel 8.—14.

I. ABADIA>TAS.—Rays of flowers none.
■■■■■ lo. Acmella L. Bastard Acmella O un 1 jl Y Ceylon 1768. S s.l PIuk.nl. 159.
Acmella lanccolktka Lk.
alba Herit. white-flowered O un li jn.jL W Peru 178&*S co Her. st 7.4
oleraea L. potherb O TM\ 1 jl.s Y E. Indies 1770? S co Jac. vin. S. 135
brasiliensis Spr. Brazilian O un jn.au Y Brazil 1827. S co
II. UADIAVE.—Flowers rayed.
■■■■■ 625 dc'bils Kth. weak O un \ jl.au Y E. Indies 1824 S co Ru. am. 6.63
O un ^ jl.au Y S. Amer. 1821. S co
III. MUVICa.—Seeds at top mutic. *
20626 exasperata Jac. rough ArOJun i jl.au Y Caracoas 1818. S co Jac. ic. 3.584
IV. DU^BIA.—Not sufficiently known.
20627 uliginbsa Swx. marsh O un 4 jl.au Y Jamaica 1819. S co

■■■■■ 2282. PETRO^BIUM R. Br. PETROBIUM. (Petros, a rock, bio, to live; habitation.) Comp. Hcl&ntheas. 1.
 628 arbTireum R. Br. tree " m a or 6 ... Pa.Y Sfc Helena 1816. C s.l
 Bldens arbbrea Rox. Spilanthes arbbrea Forst.

■■■■■ 20631 cernua L. SALMKA. (Prince Charles of Sahn Dyke, in Holland.) Cotnp. Helidntheaf. 2. - 4
 M32 chrysanthemoides Mr. Chrysanth.-lk climbing ft pr 6 jnjL Y VeraCruz 1820. C co Bot mag- &OS
 20R33 Aclinanthoides Kth. Htpirkirja scandentia Spr. hirsute « J_ O pr G au Y Jamaica 182a S co

■■■■■ 2284. ■■■■■ BIDENS. (Bis, twice,rfe/15, a tooth; seeds.) Comp. Helidntheaf. 30. — 43.

I. I. IDIIVv.—Leaves undivided.
20631 cernua L. nodding O un 2 jl.s Y Britain dit S co Eng. bot. 1114
M32 chrysanthemoides Mr. Chrysanth.-lk O un 1 jlau Y N. Amer. 1820. S co
20R33 Aclinanthoides Kth. Sunflower like O un, 2 jlau Y Mexico 1825. S co
20f34 luxurians W. luxuriant ^ AI un 3 jlau Y Mexico . D co
JG35 foliosa W. leafy O un 3 jnjL Y 1818. S co
20636 hirtella Kth. small hairy O un 1 jlau Y "Mexico 1836. S co
IPJ37 procumbens Kth. procumbent -* O^ un } jl.au X Mexico 1827. S co
50 ^b 38 nodiflora L. knot-flowered O un 1 jlau Y E. Indies 1732. S co Di. el. 44. Si
II. TRIPARTIVE.—Leaves triparted and ternate.
■■■■■ Hpartite L. triparted O un 2 jl.s Y Britain wat.pl. S co Enc. bot. 1113
rtpens D Don creeping J t A or 2 jn.au Y Nepal 1819. D co *

20341 <i>bullata L.</i>	<i>Uistered-leaved</i>	<i>O un</i>	9	<i>jl.au</i>	Y	N. Amer.	1759.	S	co	Ard.sp.2.18
20642 <i>connMa TV.</i>	<i>connate</i>	<i>O un</i>	9	<i>jn.jl</i>	Y	N. Amer.	1817.	S	co	
20643 <i>hetiTopjh Ha Or.</i>	<i>various-leaved</i>	<i>£ iAJ or</i>	9	<i>au.s</i>	Y	Mexico	1803.	S	s.l	Or. d. 8.13
SW544 <i>luminWra IV.</i>	<i>small-flowered</i>	<i>O un</i>	1	<i>jn.jl</i>	Y	Baical	1893.	S	co	
20645 <i>angustissima Kth.</i>	<i>narrowest</i>	<i>* iAJ un</i>	1	<i>jl</i>	Y	N. Spain	1820.	S	co	
20646 <i>btnta ScAotf</i>	<i>striated-flowered</i>	<i>O un</i>	1	<i>jl.au</i>	W	Mexico	1824.	S	co	Sw. fl. gar. 937
90617 <i>frondbsa i.</i>	III. PINNA^TJE. — <i>Leaves pinnate, bipinnatifid, or bipinnate.</i>									
90648 <i>leucantha Z.</i>	<i>frondose</i>	<i>O un</i>	1j	<i>jl.au</i>	Y	N. Amer.	1710.	S	co	M.h.6.5.91
20649 <i>chinensis W.</i>	<i>white-flowercd</i>	<i>O un</i>	1jj	<i>jl.au</i>	W	S. Amer.		S	co	Ru.am. 6.15.2
20650 <i>chrysanthra Or.</i>	<i>Chinese</i>	<i>O un</i>	1j	<i>jl.au</i>	Y	China	1801.	S	co	
20651 <i>pilosa L.</i>	<i>yellow-flowered</i>	<i>O un</i>	1	<i>jl</i>	Y	1824.	S	co	Di. el. 43.51
90652 <i>fainbuc</i>	<i>hairy</i>	<i>A or</i>	3	<i>jl.au</i>	S	S. Amer.	1801.	D	co	Cav. ic. 3.229
90653 <i>reflex.i</i>	<i>Elder-leaved</i>	<i>iAJ un</i>	2	<i>jn.jl</i>	Y	Mexico	1824.	D	co	
50654 <i>proct-ra</i>	<i>reflexed</i>	<i>2</i>	<i>6</i>	<i>jn</i>	Y	Mexico	1824.	S	co	Hot. reg. 684
30655 <i>crithmifbi</i>	<i>tall</i>	<i>3</i>	<i>3</i>	<i>jl.au</i>	Y	Porto Rico	...	S	co	
90656 <i>Berteridmi</i>	<i>Benthirc-lvd</i>									
Coreopsis curon^a	<i>pteroflowered</i>	<i>O or</i>	9	<i>jl.au</i>	Y	S. Amer.	1800.	S	co	
20657 <i>grandiflori Biff</i>	<i>sweet-scented</i>	<i>8</i>	<i>fra</i>	<i>jn.jl</i>	W	Mexico	1825.	S	co	Cav. ic. 1.13
ödörMa <i>Horf</i>	<i>bipinnate-/atvd</i>	<i>8</i>	<i>©</i>	<i>jl.au</i>	Y	N. Amer.	1687.	S	co	Her. par. 123
20658 <i>ortorfita Cav.</i> % 4	<i>Cynapium-leaved</i>	<i>iQ cu</i>		<i>jl.au</i>	Y	Cuba	1827.	S	co	
20659 <i>bipimi'itu L.</i>										
20660 <i>cynapifuhu Kih.</i>										


2283. PLATYPTJSRIS *Kth.* PLATyITEMS. (*Platys*, broad, *pteron*, wing; margin of seeds.) *Comp.Hel. t*
20661 *crocata Kth.* saffVon-co/orcd £ ESI or 2 ja.mr O & Amer. 1812. D co Hot mag. 1687
Spilanthus crocatui B. M.

2936. LAGA'SCI* Cav.	LAGASCA. (<i>Don Mariana Lagasca</i> , prof, bot., Madrid.) <i>Comp. Catd. Vern.</i> 1. 20662 m61lii Cav.	soft	£ O or 2	jns	W	S. Amer.	1815.	S	co	liot mag. 1804
2287. VO'CCA Cav.	NOCCA. (<i>Dominic Nocca</i> , a professor at Ticin.) " <i>Comp. Card. Vern.</i> 4. S0663 rubra Swt.	red	» D o r 2	jl	R	Mexico	1823.	C	co	H.&U.n.3H
<i>l'agasca rubra Kth.</i>										
20664 rigula Cav.	rigid	* • or 4	aus.s	Pk	Mexico	1825.	C	61		
<i>Lifigasca Aelianthubflia Kth.</i>										
2065J suav'colvs Sivt.	sweat-scented	M • or 4	aus.s	W	Mexico	1825.	C	s.l		
<i>LaK&Ma suavl'olcns Kth.</i>										
90666 latifolia Ccr.	broad-leaved	1 J o r 3	au	W	Mexico	1826.	C	fi.l	Sw.fl.gar.215	

8288. LAV&NIA L.	LAVBNIA. (Supposed to be of Cingalese origin.) <i>Comp+Helidnthe.</i> 2. 20667 erecta Sivz.	upright	ED un 2	jls	Y	E. Indies	1739.	S	co	Bur. zey.42
20668 decambena Swx.	decumbent	J c O u p	jlS	Y	Jamaica	1818.	S	co	SI.jam. 1.155. J	

9289. CACA^LIA L.	CACALIA. (<i>Kakos</i> , pernicious, <i>jian</i> , exceedingly; supp. to the soil.) <i>Com.Jac.</i> 33—72. I. VRUTicbuM.—Shrubby; leaves fleshy.									
90669 carntaa //, K.	<i>Qeshy.U-aved</i>	2		Y	C. G. a	1757.	S	s.p		
90670 papilKris L.	iy				C. G. H.	1727.	s.p		Di.cl. 55.63	
20671 cylindrica Lain.	cylindrical	CL	cu	1 jnjl	Y	C. G. H.	1H18.	e.p	Dec. pi. ic.	
20672/locoides L.	cl.	L	lj	6 jn.	Y	C. G. H.	1710.	s.p	Dec. pi. 90	
90673 Kleinta L.	Klein's	«. L	Jcu	3 s.o	Y	C. G. H.	1732.	•P	Dec. pi. 12	
20674 //cuphorbfum L.	AntcuphorbiuntL	I	cu	3 f.mr	Y	Canaries	1596.	s.p	Di. el. 55.9.3	
20675 repens IV.	creeping	tt.	I J cu	1 jn.o	Y	C. G. H.	1759.	s.p	Dec. pi. 42	
20676 Hawcirthii Swt.	Haworth's	«. I	eu	—	Y	C. G. H.	1795.	co		
<i>Kleinia tomentosa Haw.</i>										
80677 longiflMia Haw.	long-leaved	CL i J cu	1	—	—	—	1820.	C	s.l	
pugioniformis LA.	jointed	CLUJCU	l s.n	Y	C. G. H.	1775.	C	s.p	Dec. pi. 18	
80678 articulata Thun.	i. SUDCORIA^CEA.—Leaves subcoriaceous.									
90679 reticulata Lk.	netted	tt.	I J cu	2	Y	Bourbon	1823.	C	s.p	
90680 tomentisa Thun.	ii. MsMBBAifa^Cbie.—Leaves membranaceous.									
90681 appendiculata L.	tomentose	a . J cu	2	...	Y	C. G. H.	1795.	C	«p	
90688 radicans Thun.	appcnndgil	tt.	I J cu	2	... m Y	Teneriffe	1815.	C	co	
	H.^HERDA^CEA.—Herbaceous / leaves fleshy.									
	rooting	^ I J cu	jl	Y	C. G. H.	1893.	C	* s.p		
20683 bicolor Ros.	1. INDIVIDUAL.—Leaves membranaceous, undivided.									
90684 ovMis B. R.	two. colored	•LCD or	5	my.s	P	E. Indies	1804.	C	co	Bot reg. 110
20685 <i>conchifolia L.</i>	oval-leaved	£ O or	3	my.s	Y	£. Indies	...	C	s.p	Bot reg. 101
20686 coccpea //, K.	SouthUtle-lvd	O or	1*	Jl	P	E. Indies	1768.	S	co	R. mal. 10.68
90687 sagittata W.	BeatXct-flowercd	O or	O		O	1799.	S	co	Bot. mag. 564
90688 hastata L.	sagittate	nm	or	3 jl.au	O.P	Java	1823.	S	co	
90689 suaveolcns L.	halbcrt-tazivrf	k A	1	au.o	• W	Siberia	1780.	D	co	Gm.sL2.66
90690 atnplicifolia L.	sweet-scented	£ k A	6	au.o	W	N. Amer.	1752.	D	co	
20691 rcmförmis W.	Orache-leaved	£ A	4	au	L.P	N. Amer.	1669.	D	co	Pluk.aLI01.2
50692 alpma L.	kidney-formed	£ A	or	1jlau	W	N. Amer.	1801.	D	co	
20693 cordibflia Kth.	alpine	£ A	or	2 jl.au	P	Austria	1739.	D	co	Jac.au. 3.234
20694 tuberba Nut.	heart-leaved	J	Acu	1 au	W?	Mexico	1823.	R	co	
20695 dblifrona L.	tuberous	J	A	1	—	N. Amer.	1812.	R	p.l	
20696 rhoiobifoha IV.	white-leaved	^ A	or	2 jl.au	Y	Austria	1739.	D	co	Jac au. 3.235
macroph^lla Bieb.	rhomb-leaved	^ A	or	3 au.o	Y	Siberia	1816.	D	co	
20697 leucoph^lla W.	white-leaved	^ A	or	3 au.o	P.Y	S. Europe	1819.	D	co	
tomentfsa Vil.	climbing	fc, iAJ or	6	ap	Pk	C. G. H.	1774.	D	co	
20698 sc&ndcns Thun.	ii. DIVJB.—Leaves membranaceous, divided.									
20699 runcinkta Kth.	runcinate	£ ^ u n S	...	Y	Mexico	1825.	D	co		
20700 radulefuha Kth.	rasp-leaved	t AJ un	2	...	Mexico	1827.	D	to		
20701 peltafu Kth.	peltate	£ x u n	3	...	Mexico	1895.	D	co	H.&B.4. 361	

2290. CULCITIUM Bonn. CITLCITIUM. (*Culcita*, a stuffed bed; heads of palea-)
20702 talfcincum Spr. WMow-leaved s» L or 6 jn.jl Y N. Holl. 1820. C co Bot. rep. 923
CacMia ralicina Lab.

ttjl. KLEI'N/^ Jac.	KLRINIA.	(James Henry Klein, a German botanist.)	Comp. Jacob.	6.—13.
8 KP _{rud} eralis Jac.	rubble	O" un* 1 jl.au	W Jamaica	1823. S co Jac. am. 127
6 S _! to & Mf _{os} Kth.	Tagetes-like	O" un 1 jl.au	W Mexico	1823. S co
gW _{di} colorata Kth.	colored	O un 1 jl.au	W.R Mexico	1823. S co
§(706 Porophyllum L.	dotted-leaved	O un 14 Jn.o	W N. Amer.	1699. S co Cav. ic. 3.222
20/07 viruliflora Kth.	green-flowered	tt-Dcu i Jn.o	G Mexico	1823. S s.l
2U/08 sutfructicata W.	Bufruticosc	tL I' lun Sjn.o	P Brazil	1820. C co Cav. ic. a 257
	ETHELIA L.	ETHULIA.	(Meaning unknown.)	Comp. Eup. 4.—7.
	Conyzia-like	O un f jl.au Pa.P	India	1776. S co Bot. reg. 695
	divaricate	O un i jl.au P	India	1815. S co Lam.il. 699
	Brazil	EJ un 2 jl.au P	Brazil	1823. S co
	Tifilia D. Don	iQJ un 1 jl.au P	Nepal	1819. S co
	QUEMIUCav.	PIQUERIA.	(Andreas Piquerio, a Spanish botanist.)	Comp. Eup. 4.
mi <i>stoides</i> Kth.	Wormwood-like	LAI un 2 my.s	W Mexico	1827. C 81
	birec-ncrved	LAI I* 2 jl.au	W Mexico	1805. C 80 CaV. IC. 3.235
ovate <i>three-nctched</i>		ZJ my.s	W Mexico	1800. C 81
pilose	jf iAI un 2 my.s	W Mexico	1824. C 81	
	GOLDYLOCKS. (<i>Chrysos</i> , gold, home, hair; tufts of yel. fl.)	Comp. Card. Vern. 13.—32.		
?^a (ireal.	Golden-hair	M i_J or 6 jl.or	Y C. G. H. 1731. C p.l Bot. mag. 1972	
	spreading	M i_J or 3 jl.or	Y C. G. H. 1810. C p.l	
	drooping	M i_J or 4 jl.or	W C. G. H. 1712. C p.l Com. h. 2.45	
	rugged	M i_J or 4 au.s	W C. G. H. 1759. C p.l Com. h. 2.48	
	toothed-leaved *	M i_J or 4 au.s	W C. G. H. 1732. C p.l Di. el. 88. 103	
	Flax-leaved	i A r 2 8.0	Y Europe 1596. C co Jac. sc. 3.368	
	twiggy	5 A or 1 au.o	Y N. Amer. 1821. D p.l Eng. bot. 2505	
Mata	Lam. Taragon-like	3t A or 2 8.0	Y Siberia ... D co	
	naked	3t A or 118.0	Y Carolina 1818. D p.l	
	two-flowered	^ A < 3 au.s	B Siberia 1741. D co Gm. si. 2 82.1	
	villous-leaved	ft A or 1) au.s	Y Hungary 1799. D co W.&K.53	
22p5. OZOTHA'MNUS R. Br.	OZOTHAM.VUS.	(Oxo5, branch, <i>thamnos</i> , shrub.)	Comp. Card. Fern. 3.	
cinereus It. Br.	grey	s% or 1 ap.8	Y V.D. L. 1820. C p.l Lab. n. h. 2.182	
Chrysocoma cinerea Lab.				
rosmarinifilius/f.fr. Rosmary-lvd	*	L-J or 1 ap.8	Y V.D.L. 1822. C s.p Lab. n. h. 2.181	
Alipatritum rosmarinifiliun Lab.				
ferruginosus R. Br.	rusty	aH i_J or 1 ao.8	Y V.D.L. 1822. C p.l Lab. n. h. 2.180	
cupuritifm ferrugineum Lab.				
22296 EUTHA'MIA Nut. EUTHAMIA.	IE «, well, <i>thames</i> , crowded; flowers.)	Comp. Card. Vern. 2.		
w/r/raminifolia Nut.	Grass-leaved	^ A or 3 s.o	Y N. Amer. 1758. D co Bot. mag. 2546	
rosmarinifilius/f.fr. Rosmary-lvd	*	L-J or 1 ap.8	Y V.D.L. 1822. C s.p Lab. n. h. 2.181	
Alipatritum rosmarinifiliun Lab.				
ferruginosus R. Br.	rusty	aH i_J or 1 ao.8	Y V.D.L. 1822. C p.l Lab. n. h. 2.180	
cupuritifm ferrugineum Lab.				
22296 EUTHA'MIA Nut. EUTHAMIA.	IE «, well, <i>thames</i> , crowded; flowers.)	Comp. Card. Vern. 2.		
w/r/raminifolia Nut.	Grass-leaved	^ A or 3 s.o	Y N. Amer. 1758. D co Bot. mag. 2546	
rosmarinifilius/f.fr. Rosmary-lvd	*	L-J or 1 ap.8	Y V.D.L. 1822. C s.p Lab. n. h. 2.181	
Alipatritum rosmarinifiliun Lab.				
ferruginosus R. Br.	rusty	aH i_J or 1 ao.8	Y V.D.L. 1822. C p.l Lab. n. h. 2.180	
cupuritifm ferrugineum Lab.				
22296 EUTHA'MIA Nut. EUTHAMIA.	IE «, well, <i>thames</i> , crowded; flowers.)	Comp. Card. Vern. 2.		
w/r/raminifolia Nut.	Grass-leaved	^ A or 3 s.o	Y N. Amer. 1758. D co Bot. mag. 2546	
rosmarinifilius/f.fr. Rosmary-lvd	*	L-J or 1 ap.8	Y V.D.L. 1822. C s.p Lab. n. h. 2.181	
Alipatritum rosmarinifiliun Lab.				
ferruginosus R. Br.	rusty	aH i_J or 1 ao.8	Y V.D.L. 1822. C p.l Lab. n. h. 2.180	
cupuritifm ferrugineum Lab.				
22296 EUTHA'MIA Nut. EUTHAMIA.	IE «, well, <i>thames</i> , crowded; flowers.)	Comp. Card. Vern. 2.		
w/r/raminifolia Nut.	Grass-leaved	^ A or 3 s.o	Y N. Amer. 1758. D co Bot. mag. 2546	
rosmarinifilius/f.fr. Rosmary-lvd	*	L-J or 1 ap.8	Y V.D.L. 1822. C s.p Lab. n. h. 2.181	
Alipatritum rosmarinifiliun Lab.				
ferruginosus R. Br.	rusty	aH i_J or 1 ao.8	Y V.D.L. 1822. C p.l Lab. n. h. 2.180	
cupuritifm ferrugineum Lab.				
22296 EUTHA'MIA Nut. EUTHAMIA.	IE «, well, <i>thames</i> , crowded; flowers.)	Comp. Card. Vern. 2.		
w/r/raminifolia Nut.	Grass-leaved	^ A or 3 s.o	Y N. Amer. 1758. D co Bot. mag. 2546	
rosmarinifilius/f.fr. Rosmary-lvd	*	L-J or 1 ap.8	Y V.D.L. 1822. C s.p Lab. n. h. 2.181	
Alipatritum rosmarinifiliun Lab.				
ferruginosus R. Br.	rusty	aH i_J or 1 ao.8	Y V.D.L. 1822. C p.l Lab. n. h. 2.180	
cupuritifm ferrugineum Lab.				
22296 EUTHA'MIA Nut. EUTHAMIA.	IE «, well, <i>thames</i> , crowded; flowers.)	Comp. Card. Vern. 2.		
w/r/raminifolia Nut.	Grass-leaved	^ A or 3 s.o	Y N. Amer. 1758. D co Bot. mag. 2546	
rosmarinifilius/f.fr. Rosmary-lvd	*	L-J or 1 ap.8	Y V.D.L. 1822. C s.p Lab. n. h. 2.181	
Alipatritum rosmarinifiliun Lab.				
ferruginosus R. Br.	rusty	aH i_J or 1 ao.8	Y V.D.L. 1822. C p.l Lab. n. h. 2.180	
cupuritifm ferrugineum Lab.				
22296 EUTHA'MIA Nut. EUTHAMIA.	IE «, well, <i>thames</i> , crowded; flowers.)	Comp. Card. Vern. 2.		
w/r/raminifolia Nut.	Grass-leaved	^ A or 3 s.o	Y N. Amer. 1758. D co Bot. mag. 2546	
rosmarinifilius/f.fr. Rosmary-lvd	*	L-J or 1 ap.8	Y V.D.L. 1822. C s.p Lab. n. h. 2.181	
Alipatritum rosmarinifiliun Lab.				
ferruginosus R. Br.	rusty	aH i_J or 1 ao.8	Y V.D.L. 1822. C p.l Lab. n. h. 2.180	
cupuritifm ferrugineum Lab.				
22296 EUTHA'MIA Nut. EUTHAMIA.	IE «, well, <i>thames</i> , crowded; flowers.)	Comp. Card. Vern. 2.		
w/r/raminifolia Nut.	Grass-leaved	^ A or 3 s.o	Y N. Amer. 1758. D co Bot. mag. 2546	
rosmarinifilius/f.fr. Rosmary-lvd	*	L-J or 1 ap.8	Y V.D.L. 1822. C s.p Lab. n. h. 2.181	
Alipatritum rosmarinifiliun Lab.				
ferruginosus R. Br.	rusty	aH i_J or 1 ao.8	Y V.D.L. 1822. C p.l Lab. n. h. 2.180	
cupuritifm ferrugineum Lab.				
22296 EUTHA'MIA Nut. EUTHAMIA.	IE «, well, <i>thames</i> , crowded; flowers.)	Comp. Card. Vern. 2.		
w/r/raminifolia Nut.	Grass-leaved	^ A or 3 s.o	Y N. Amer. 1758. D co Bot. mag. 2546	
rosmarinifilius/f.fr. Rosmary-lvd	*	L-J or 1 ap.8	Y V.D.L. 1822. C s.p Lab. n. h. 2.181	
Alipatritum rosmarinifiliun Lab.				
ferruginosus R. Br.	rusty	aH i_J or 1 ao.8	Y V.D.L. 1822. C p.l Lab. n. h. 2.180	
cupuritifm ferrugineum Lab.				
22296 EUTHA'MIA Nut. EUTHAMIA.	IE «, well, <i>thames</i> , crowded; flowers.)	Comp. Card. Vern. 2.		
w/r/raminifolia Nut.	Grass-leaved	^ A or 3 s.o	Y N. Amer. 1758. D co Bot. mag. 2546	
rosmarinifilius/f.fr. Rosmary-lvd	*	L-J or 1 ap.8	Y V.D.L. 1822. C s.p Lab. n. h. 2.181	
Alipatritum rosmarinifiliun Lab.				
ferruginosus R. Br.	rusty	aH i_J or 1 ao.8	Y V.D.L. 1822. C p.l Lab. n. h. 2.180	
cupuritifm ferrugineum Lab.				
22296 EUTHA'MIA Nut. EUTHAMIA.	IE «, well, <i>thames</i> , crowded; flowers.)	Comp. Card. Vern. 2.		
w/r/raminifolia Nut.	Grass-leaved	^ A or 3 s.o	Y N. Amer. 1758. D co Bot. mag. 2546	
rosmarinifilius/f.fr. Rosmary-lvd	*	L-J or 1 ap.8	Y V.D.L. 1822. C s.p Lab. n. h. 2.181	
Alipatritum rosmarinifiliun Lab.				
ferruginosus R. Br.	rusty	aH i_J or 1 ao.8	Y V.D.L. 1822. C p.l Lab. n. h. 2.180	
cupuritifm ferrugineum Lab.				
22296 EUTHA'MIA Nut. EUTHAMIA.	IE «, well, <i>thames</i> , crowded; flowers.)	Comp. Card. Vern. 2.		
w/r/raminifolia Nut.	Grass-leaved	^ A or 3 s.o	Y N. Amer. 1758. D co Bot. mag. 2546	
rosmarinifilius/f.fr. Rosmary-lvd	*	L-J or 1 ap.8	Y V.D.L. 1822. C s.p Lab. n. h. 2.181	
Alipatritum rosmarinifiliun Lab.				
ferruginosus R. Br.	rusty	aH i_J or 1 ao.8	Y V.D.L. 1822. C p.l Lab. n. h. 2.180	
cupuritifm ferrugineum Lab.				
22296 EUTHA'MIA Nut. EUTHAMIA.	IE «, well, <i>thames</i> , crowded; flowers.)	Comp. Card. Vern. 2.		
w/r/raminifolia Nut.	Grass-leaved	^ A or 3 s.o	Y N. Amer. 1758. D co Bot. mag. 2546	
rosmarinifilius/f.fr. Rosmary-lvd	*	L-J or 1 ap.8	Y V.D.L. 1822. C s.p Lab. n. h. 2.181	
Alipatritum rosmarinifiliun Lab.				
ferruginosus R. Br.	rusty	aH i_J or 1 ao.8	Y V.D.L. 1822. C p.l Lab. n. h. 2.180	
cupuritifm ferrugineum Lab.				
22296 EUTHA'MIA Nut. EUTHAMIA.	IE «, well, <i>thames</i> , crowded; flowers.)	Comp. Card. Vern. 2.		
w/r/raminifolia Nut.	Grass-leaved	^ A or 3 s.o	Y N. Amer. 1758. D co Bot. mag. 2546	
rosmarinifilius/f.fr. Rosmary-lvd	*	L-J or 1 ap.8	Y V.D.L. 1822. C s.p Lab. n. h. 2.181	
Alipatritum rosmarinifiliun Lab.				
ferruginosus R. Br.	rusty	aH i_J or 1 ao.8	Y V.D.L. 1822. C p.l Lab. n. h. 2.180	
cupuritifm ferrugineum Lab.				
22296 EUTHA'MIA Nut. EUTHAMIA.	IE «, well, <i>thames</i> , crowded; flowers.)	Comp. Card. Vern. 2.		
w/r/raminifolia Nut.	Grass-leaved	^ A or 3 s.o	Y N. Amer. 1758. D co Bot. mag. 2546	
rosmarinifilius/f.fr. Rosmary-lvd	*	L-J or 1 ap.8	Y V.D.L. 1822. C s.p Lab. n. h. 2.181	
Alipatritum rosmarinifiliun Lab.				
ferruginosus R. Br.	rusty	aH i_J or 1 ao.8	Y V.D.L. 1822. C p.l Lab. n. h. 2.180	
cupuritifm ferrugineum Lab.				
22296 EUTHA'MIA Nut. EUTHAMIA.	IE «, well, <i>thames</i> , crowded; flowers.)	Comp. Card. Vern. 2.		
w/r/raminifolia Nut.	Grass-leaved	^ A or 3 s.o	Y N. Amer. 1758. D co Bot. mag. 2546	
rosmarinifilius/f.fr. Rosmary-lvd	*	L-J or 1 ap.8	Y V.D.L. 1822. C s.p Lab. n. h. 2.181	
Alipatritum rosmarinifiliun Lab.				
ferruginosus R. Br.	rusty	aH i_J or 1 ao.8	Y V.D.L. 1822. C p.l Lab. n. h. 2.180	
cupuritifm ferrugineum Lab.				
22296 EUTHA'MIA Nut. EUTHAMIA.	IE «, well, <i>thames</i> , crowded; flowers.)	Comp. Card. Vern. 2.		
w/r/raminifolia Nut.	Grass-leaved	^ A or 3 s.o	Y N. Amer. 1758. D co Bot. mag. 2546	
rosmarinifilius/f.fr. Rosmary-lvd	*	L-J or 1 ap.8	Y V.D.L. 1822. C s.p Lab. n. h. 2.181	
Alipatritum rosmarinifiliun Lab.				
ferruginosus R. Br.	rusty	aH i_J or 1 ao.8	Y V.D.L. 1822. C p.l Lab. n. h. 2.180	
cupuritifm ferrugineum Lab.				
22296 EUTHA'MIA Nut. EUTHAMIA.	IE «, well, <i>thames</i> , crowded; flowers.)	Comp. Card. Vern. 2.		
w/r/raminifolia Nut.	Grass-leaved	^ A or 3 s.o	Y N. Amer. 1758. D co Bot. mag. 2546	
rosmarinifilius/f.fr. Rosmary-lvd	*	L-J or 1 ap.8	Y V.D.L. 1822. C s.p Lab. n. h. 2.181	
Alipatritum rosmarinifiliun Lab.				
ferruginosus R. Br.	rusty	aH i_J or 1 ao.8	Y V.D.L. 1822. C p.l Lab. n. h. 2.180	
cupuritifm ferrugineum Lab.				
22296 EUTHA'MIA Nut. EUTHAMIA.	IE «, well, <i>thames</i> , crowded; flowers.)	Comp. Card. Vern. 2.		
w/r/raminifolia Nut.	Grass-leaved	^ A or 3 s.o	Y N. Amer. 1758. D co Bot. mag. 2546	
rosmarinifilius/f.fr. Rosmary-lvd	*	L-J or 1 ap.8	Y V.D.L. 1822. C s.p Lab. n. h. 2.181	
Alipatritum rosmarinifiliun Lab.				
ferruginosus R. Br.	rusty	aH i_J or 1 ao.8	Y V.D.L. 1822. C p.l Lab. n. h. 2.180	
cupuritifm ferrugineum Lab.				
22296 EUTHA'MIA Nut. EUTHAMIA.	IE «, well, <i>thames</i> , crowded; flowers.)	Comp. Card. Vern. 2.		
w/r/raminifolia Nut.	Grass-leaved	^ A or 3 s.o	Y N. Amer. 1758. D co Bot. mag. 2546	
rosmarinifilius/f.fr. Rosmary-lvd	*	L-J or 1 ap.8	Y V.D.L. 1822. C s.p Lab. n. h. 2.181	
Alipatritum rosmarinifiliun Lab.				
ferruginosus R. Br.	rusty	aH i_J or 1 ao.8	Y V.D.L. 1822. C p.l Lab. n. h. 2.180	
cupuritifm ferrugineum Lab.				
22296 EUTHA'MIA Nut. EUTHAMIA.	IE «, well, <i>thames</i> , crowded; flowers.)	Comp. Card. Vern. 2.		
w/r/raminifolia Nut.	Grass-leaved	^ A or 3 s.o	Y N. Amer. 1758. D co Bot. mag. 2546	
rosmarinifilius/f.fr. Rosmary-lvd	*	L-J or 1 ap.8	Y V.D.L. 1822. C s.p Lab. n. h. 2.181	
Alipatritum rosmarinifiliun Lab.				
ferruginosus R. Br.	rusty	aH i_J or 1 ao.8	Y V.D.L. 1822. C p.l Lab. n. h. 2.180	
cupuritifm ferrugineum Lab.				
22296 EUTHA'MIA Nut. EUTHAMIA.	IE «, well, <i>thames</i> , crowded; flowers.)	Comp. Card. Vern. 2.		
w/r/raminifolia Nut.	Grass-leaved	^ A or 3 s.o	Y N. Amer. 1758. D co Bot. mag. 2546	
rosmarinifilius/f.fr. Rosmary-lvd	*	L-J or 1 ap.8	Y V.D.L. 1822. C s.p Lab. n. h. 2.181	
Alipatritum rosmarinifiliun Lab.				
ferruginosus R. Br.	rusty	aH i_J or 1 ao.8	Y V.D.L. 1822. C p.l Lab. n. h. 2.180	
cupuritifm ferrugineum Lab.				
22296 EUTHA'MIA Nut. EUTHAMIA.	IE «, well, <i>thames</i> , crowded; flowers.)	Comp. Card. Vern. 2.		
w/r/raminifolia Nut.	Grass-leaved	^ A or 3 s.o	Y N. Amer. 1758. D co Bot. mag. 2546	
rosmarinifilius/f.fr. Rosmary-lvd	*	L-J or 1 ap.8	Y V.D.L. 1822. C s.p Lab. n. h. 2.181	
Alipatritum rosmarinifiliun Lab.				
ferruginosus R. Br.	rusty	aH i_J or 1 ao.8	Y V.D.L. 1822. C p.l Lab. n. h. 2.180	
cupuritifm ferrugineum Lab.				
22296 EUTHA'MIA Nut. EUTHAMIA.	IE «, well, <i>thames</i> , crowded; flowers.)	Comp. Card. Vern. 2.		
w/r/raminifolia Nut.	Grass-leaved	^ A or 3 s.o	Y N. Amer. 1758. D co Bot. mag. 2546	
rosmarinifilius/f.fr. Rosmary-lvd	*	L-J or 1 ap.8	Y V.D.L. 1822. C s.p Lab. n. h. 2.181	
Alipatritum rosmarinifiliun Lab.				
ferruginosus R. Br.	rusty	aH i_J or 1 ao.8	Y V.D.L. 1822. C p.l Lab. n. h. 2.180	
cupuritifm ferrugineum Lab.				
22296 EUTHA'MIA Nut. EUTHAMIA.	IE «, well, <i>thames</i> , crowded; flowers.)	Comp. Card. Vern. 2.		
w/r/raminifolia Nut.	Grass-leaved	^ A or 3 s.o	Y N. Amer. 1758. D co Bot. mag. 2546	
rosmarinifilius/f.fr. Rosmary-lvd	*	L-J or 1 ap.8	Y V.D.L. 1822. C s.p Lab. n. h. 2.181	
Alipatritum rosmarinifiliun Lab.				
ferruginosus R. Br.	rusty	aH i_J or 1 ao.8	Y V.D.L. 1822. C p.l Lab. n. h. 2.180	
cupuritifm ferrugineum Lab.				
22296 EUTHA'MIA Nut. EUTHAMIA.</				

2306. OTA'NTIIUS Lk. OTANTHI/8. (*OUS, ear, anthos*, flower; app. at base of florets.) *L\>?;/>, Anthem. 1.*

20756 maritimus Lk. sea f A P^t f J^t England aea ah. C Bl Eng tut. MI

Santolina maritima 5m. Dibtis candidinsuna Deaf.

2307. CALEA'CTE R.Br. CALBACTK. (*KaJos. beauty, nlcte, sea-shore; ornament there.*) *Comp. Helidnthe**. 1.

20757 urticifolia 7t. 22r. Nettle-leaved * • or 2 jl.au Y VeraCruz 1740. C co

Solidago urticifolia MU.

2308. JTHANA^SIA L. ATUANASIA. (//, priv., thanatos, death; lasting flowers) *Comp. Anthem. 17—28.*

I. iFPE — Leaves undivided.						
20758 capitata L.	capitate	■■■ lor	U ja mr	T	C. G. H. 1774.	C l.p M.h.6.3.48
20759 longifolia /am.	long-leaved	■■■ or	2 jl.au	T	C. G. H. 1800.	C l.p
20760 pubescens L.	pubescent	■■■ or	(i jn.au	F	C. G. H. 1768.	C co Com. h. 2.47
20761 punctata Berg.	dotted	■■■ or	3 myjl	F	G. G. H. 1822.	C l.p Pet.g.81,<*
20762 dentata /.	toothed	■■■	Ujl.au	F	G. G. H. 1759.	C l.p Com. r. 41
20763 tomentosa 7%IMI	vicioYLy-leaved	■■■	2 myjn	F	C. G. H. 1774.	C l.p
20764 canescens Thun.	cancscent	■■■	3 jn.jl	F	C. G. H. 1820.	C l.p Cav. ic. 1. 3
lanuginosa Cav.						
20765 crenata L.	crenate	■■■	J or	2 jn.jl	C. G. H. 1816.	C l.p
20766 cuncifolia Lam.	wedge-leaved	■■■	or	2 jl.au	C. G. H. 1816.	C l.p Lam. il. 670.1
20767 filiflraais Thun.	liliiform	■■■	pr	2 au	Y C. G. H. 1787.	C J.p
II. DIVTdjR. — Leaves divided.						
20768 virgata L.	twiggy	il ■■■ pr	1	leaves divided	C. G. H. 1815.	C co Jac. sc. 2. 148
20769 crithimflba /.	Sampshire-lvd	* L J I*	il	jl.au	C. G. H. 1723.	C l.p Coin. h. 2.50
20770 parviflrbm L.	small-flowered	il L J P ^r	2 ap	Y	C. G. H. 1731.	C l.p Jac. sc. 2.149
20771 trifurc&tia L.	triflA-leaved	tt L J P ^r	3 jl.au	Y	C. G. H. 1710.	C l.p Com. h. 2.49
20772 tricuspis Pert.	three-pointed	• i J or	3 my	Y	C. G. H. 1816.	C l.p
20773 pectinata Thun.	pectinated	il i J pr	U myjn	Y	C. G. H. 1774.	C co
20774 pinnata 7%tm.	pinnate	■■■	[t] myjl	Y	C. G. H. 1818.	C l.p Lam. il. 670.4

2309. LO^NAS Adan. LONAS. (A name of Adanson's, probably without meaning.) *Comp. Anthem. 1.*

20775 inodbn Qae. scentless O or 1 Jl.au Y Barbary 1G86. S co Bot. mag. 2276

ittthanasia &nnua L.

2310. 27ALSAMPTA Detf. COSTMARV. (<i>Balsatnon, balm ; strong balsamic smell.</i>) <i>Comp. Anthem. 5.-4!</i>
20776 virgMa Desf. twiggy ■■■ 3 jn.jl Y. o Italy 1791. D co Jacob. 4.81
20777 Agratifolia Def. Agratum-lvd ■■■ un 2 jn.o Y.a Candia 1605. C co Alp. ex. 326
20778 grandiflora Detf. large-flowering ■■■ or 2 jl.au Y Algiers 1821. S p.1 Dcsf. 1.1
20779 vulghrid W. common ■■■ A or* 3 aus.s Y.o Italy 1568. D co Schk. han.3.240
20780 annua Lk. annual O un 2 Jl au Y.a Spain 1629. S co Mil. ic. 2.227.1
Tanacetum annuum L.

2811. PF/NTZ//Thun. PENTZIA. (*Charles John Pentx, pupil of Thunberg.*) *Comp. Anthem. 1.*

20781 flabelliflrmis W. fan.form./c/w// il uJ or 3 my.au Y C. G. H. 1774. C p.l Bot. mag. 212

crenata Thun. Tanacetum flabelliflrrae Herit.

CLASS XIX. ORDER 2.

SYNGENE'SIA SUPERFLUA.

8312. TANACETUM L. TANSY.	from <i>athanatia.</i>	Cmnp. <i>Anthemidem</i> , 14,—21.
20782 boreal //, C.	northern ■■■ 4 my.U Y	N.Amcr.M818. D co
20783 sibericum L.	Siberian ■■■ 3 jn.jf Y	Siberia 1820. D co Gm. si. 2.63.2
20784 purpitreum D Don	\myrlc-flowered ■■■ 1 jn.jl Pa.R	Nepal 1HIB D co
20785 myriophyllum W.	myriad-leaved ■■■ 1 myjl Y	Levant 18K. D co
Achillea bipinnata L.		
20786 granditribum Thun.	/reat-flowercd ■■■ myjl Y	C. G. H. 1820. C p.l
20787 /infolium Thun.	Flax-leaved ■■■ Jl au Y	C. G. H. 1774. C p.l
20788 vestUum Thun.	clothed ■■■ 2 aus.s Y	C. G. H. 1816. C p.l
20789 Hiffuticrsum L.	soft/uticose ■■■ 2 my.s Y	C. G. H. 1751. C p.l Com. h. 2.100
20790 canarilusc Dec.	Canary ■■■ 2 myjl Y	Canary Is. 1816. C p.l
20791 argentum W.	silvery ■■■ 1 my.s Y	Levant 1812. D co
20792 orientate TV.	Oriental ■■■ H Jn jl Y	Armenia 1827. D co
20793 vulgarc L.	common ■■■ 2 jl.au Y	Britain ro. sid. D co Eng. bot 1229
20794 crfapum Hart.	curled ■■■ 2 jl.au Y D co
20795 angul&tum IV.	angled ■■■ 2 jl.au Y	Levant 1820. D co W. ach. 2.3
filipendiliina Lam.		

• 2313. jfRTEMI'S/ii L. Woaiwoood. (*Artemis, one of the names of Diana.*) *Comp. Anthem.*

I. FWJTICU8*—Shrubby ; leaves undivided.

20796 chinensis Lou.	ChineseMoxa ^ L J TM 4 jnjl Y.a	China 1818. C co
20797 cana Ph.	hoary • J u n I J au Y	N. Amer. 1800. C sl
20798 casrufcscns L.	csrulcscnt sH or I J au.o Y	England 6ea sh. C co Eng.bot.24S6
20799 montevideensis Spr.	Montevidean • L J un 2 ... M. Video 1827. C sl	
20800 ambigua L.	ambiguous i I u J un 2 ... Y.o C. G. H. 1800. C sl	
20801 inodbra W. en.	inodorous II 2 jl.au Y.o	Siberia 11548. D co Gm.si.2.^",r,01
Redgw6kfi Led.		
20802 Draetfincus L.	Tarragon ■■■ A cul 2 jl.au W.G & Europe 1548. D co	
20803 japonica Thuy.	Japanese ■■■ A un 3 o.n. IV Japan 1804 D co	
20804 rracunculoides Ph.	Dracunculus-lk^ A un 4 jl.au W.G Louisiana 1820. D co	
nbtanB> Fna. cernua Nut.	serrated ■■■ A un 6 jl.s W.o N. Amer. 1821 D co	Gm.si.2.5S.1.2
20806 integiflma /..	entire-leaved ■■■ A un 2 jl.au Y.a Siberia 1759. D co	
20807 longiflma Nut.	long.leaved ■■■ A u. 3 jl.au Y.a N. Amer. 1827. D co	
S0807<<naphaloldcs Wiif.	Gnaphallum-lk A un 3 jl.au Y.o N. Amer. 1824. D co	
S0B09 parviflra Ham.	smalUflowerd ■■■ A un 1 jn.au Y.a Nepal 1818. D sl	

III. PAJJUTIFOLIM.—Leaves fur the most pinnate.

i. St'rrrLierj.—Stems simple.

SOfill hvivetira SeiL	icy	^w A or	t jl.au	Y.o	Bwlterri	ISift	1) CO	Jac au. 5. np. 31
W815 Muhtina Vil.	Swis»		t jl.au	Y.Q			1)	
Jfl <i>loculii</i> (I'M/ rupi'rflris All.	Jfl re Ulna S A or	i jl.au			Alp. Kur.	1815.	I) m	V.I .M.a35
Still.	Wulfen'i	^ A o f	i fl.an					
	pedunculate	^ A pr	jn.au		Swuorl	1819.	1)	
					Caucatu	ISIS.	D RI	

K 80915 rurcrfta Jfl. eff.	rorkwl	\ A pr	i jn.uu	v i	Siberia	ifKO.	1> LI	
8061c alphna Bffr.	chiantg-ahlc	5 A or	jnjl	Y.a	Caucatu	1804.	n CO	
^ f'auc <u>uic.i</u> If*	iiiknl	\ A or	ln.il	Br	SwitMtl		D CO	Jac. au. 5. ap. Si
20618 BorvegUsa Friet	Norway	\ A pr	i in	Y.o	Norwar	1819.	1) 1,1	11. dan. Mil
rup&trii /'i/i/i. not								
90619 gTotnlindica II urj». tlrocnland	;^ A or	i jl.atl	Y.u	Omn).		1810.	D a	

20890 juiluica/;.	Juttean	« _r 11	BU	Y	Lnaat		C CO	Pluk. al
308SI lii/punica Lam,	Spanish	tt 1 un	in.y.s	Y.k	Spain		C	
20882 valiMitmii Lam,	Vnicntia	tt 1 or	1 Jura				C to	ILir.lc.48,J
20823 arrafjouOnsiu Lnm.	Arragmi	• (un 11		Y	Spain		C	AHD ar. K 1

iv. PunuTi'nu.—Italics pinnati/lil.

3«Q4 coarcldfta IK	comprewed	J. Si'MPictLH. — W<n^ simple.						
S0855 denrtontn 8to\	di'sert	i A UH 11 au.*	Y.u	Siberia	ISIH.	S CO		
Jeucan Lilt in itoli.i lhsf.	Si A un B jl.au	Y.a	Siberia	181C	S	CO		

v. EIHAM

heel.

8098B hlntita /In/A	hairy E.India Mugw.b -At UN	3	V.i	E. hi.ii...	IWH.	ii		
Wg7 mexciina W.	Mexican Mugwort & A u<	3	eut.t	Y.P			II	c tt
S088S vulgar.* L.	common Mugwort & A w		au*	P	Hrit:ll I		D	Eng. bot. 9'S
*0659 Indica ff.	Indian vul-icavcd	4 JOJ of	S MO	E.	tiiliies		D BO	[G. HMI.
90830 India W. en.	vul-icavcd	5 A utl	1 j. Jl.au	V.:			It	CO
JptSI procuntwn* Dw.	procumboat	3 A um	11 j.l.uu	V ii	Siberia	IMii.	II	
SOSaj canndeult MJ. Caun				Y.i	Caosila	181(1	T^	CO
SOBit cautlliLi b		A "»	3 avis	VM	Carolina	1H17.	D	CO

V. GLAUru'r.K. • Isao't	stüttntne. i*» with, Or SHHIQthhh.							
emab-Ueed	O frt	1 jnjl	Br	1	18K7.	s CO	Pnl.it. 3. Hh.2	
Broom	O or	WM	Hungary			W.&K		
S.i(ji(jiica	• in	I II.	Siberia			Gm.fi.		
• alj/in.i	^ A ^	1 au	Y.P	y(iitl):iitl	iian. II.	Eng. lwt. S3S		
• ilpinp	A "»	1 au	Y.P		181B.	D		
Arw-flownad	J A un	i au	Y			D		
nurirtb	O um	1.1.1.1.	Y			Biberli	8	Om. »l. B. 52.1,0
ncpU ¹	:i A un	2 Jl.nu	O.Y			SHHTiL	1> CO	tim.ii
Ms	^ A Ul)	1 iii.n	C.'trniitla				C CO	

c.iiim>li(jr;it(-il	• um	5 jl.au	W.o			c CO		
:	(tu i i	1 uu	S jl.au	Y.o		C		
Saiuhlii:;-/;^	< un	1 au.»	Br	Portugal	L7»	C		

VI. INCA\W*,--/IVJi'fj hoary or tomftUou, jimmate.

rock	i A of	3 Jnjl	Hungary			D CO		
	5 A "	1 jri.au	SBxena	ISOB.				
IKilmijlvd	£ Jor	1 JitJI	a.v			p CO		
Minwy	^ A or	2 j' il	Siltorin	1S1&		D CO		
iva	4 A or	2 nun	Br	Brünniu	i	Eng. bot. r		
Trunch	S A of	S .in.-	We	Hutignry	IKAJ.	ii		
• alt	3 A M ill	Ullu.0	Br	Aii'.ji.i	1,117.	i) CO		
Austrian	S A or	N jl.:ill	Armenia	IS 10.		Jac. nu, MOO		
irik'i	5 A or	1 uu	Y.o	NIHTIU	1K2H.	D CO		
ullyk-irairit	5 A or	H jn.jl	Slwrin			n CO		
creeppls	3 A or	Br	Siberia			C CO		
lir.iih'liy	• t_c u	2 ..	IIIifi.					

VII. BIPINNATI'FII)*,—Liane bipinnatifid itnd bipinnaU:

i. Sr'Mi'l.irps.—SUMs simple j leaves smoothth.								
rock	i A or	au	MT	Denmark	1748	I) CO	Fl. dan. 801	
Vallosi	5 A "r	1 jn.au	Y.	Siberia	1880,	p CO		
green-feared	Ii un	4 nu.o	Y.;		rope 18			
Smil liLruwnotl	it or	4 an ii	Y.	s. Miropo	JnX	C CO	BI*ck. fiSS	
ell'ink'r-leaved	at I_ wr	10 .(1	Y.d	China		C CO	DLeL 33.37	

ii. IUMII&H.—Slews broached ; letwet tmaoitieh.								
tirlinrcdit	i 1 B HI.	1 Jj .in	Y.u	Silieriu		B CO		
• iictliuij.lvd	31 !)1 III.	111.S	Y.o	atboria				
t munioUc-Ird	S ,A] nu	i! [Uu	B. Euiopa	ma.		(CO		
annual	O Ull	4 jl.au	W.G	Silierlo		8 CO	Am ru, I'M 23	
tiiciimul	I O) un	2 jl.au	Y.u	MHHMII'1	J74t.	8 CO		
	uu	8	Y.u			C CO		

S06lt9 odoratlniima/)^	ttwuftui.KCuntrdt	_ J un	ij jl.uu	Numldfa		C l.p		
				Cmt : items litnptr.				
S0S70 tiuollnsiiM / J. linn	;n;(;t	N EH, — /				8		
20871 tan.t	-tuaved	5 CD nn	1 jl.in	Ubetia	ITIW.	S	AL ped. 1.IU. 3	
foQa nr.	i n . j . i .	^ Q " "	1 jl.au	Y.a		a CO		
StJKS cuuescen* W.	I'otcentill-lvd	£ A cu	1 jl.nu		1818.	D CO	Rchb.bot.1.y	

gW6 jtrgtntea //, K.	nilv-ry	Mi l_ or	1 jnjl	Y.o		3777.	C CO	
2J77 kfrsi 7(n;	Afric.ii!	all l_ or	»! au.s	Y.o			C CO	
S878 Inefilta ZW.	iiiiflU-ated	A un	£ Jl..U	Egypt	T.sis.	C CO		
§ Itdnecnii* SocA-		cu	S Jl.au	Vy	Carpa ma			
UfaofaKn. 2,	otburcident	• or	Y.jn.au	Y.o	I.I V.HI	KM	C CO	Lob. 1c. 755

80881 <i>moxa</i> Hort.	<i>moxa</i>					China	C	co	
20882 <i>Absinthium L.</i>	<i>Absinthium, com.</i>					Britain	rubble,	D	Eng. bot. 1230
20883 <i>Sieversina Ehrh.</i>	<i>Sievers's</i>					Siberia	1800.	S	
20884 <i>laciniata W.</i>	<i>tmn-leaved</i>					Siberia	—	D	Gin. si. 2.57
20885 <i>pontica L.</i>	<i>Pontinc</i>					Austria	1570.	D	Jac. au. 1.99
20886 <i>£alsamita W.</i>	<i>Balsamita</i>						1816.	D	
20887 <i>taurica W.</i>	<i>Taurian</i>					Tauria	1818.	D	
20888 <i>vallesiacia Lam.</i>	<i>Vallais</i>					Italy	1739.	D	
20889 <i>£lbida W.</i>	<i>whitish</i>					—	—	D	
20890 <i>fragrans W.</i>	<i>fragrant</i>					Armenia	1739.	D	
20891 <i>fasciculata Bieb.</i>	<i>fascicled</i>					Iberia	1823.	D	
20892 <i>nil tans W.</i>	<i>nodding</i>					Tartary	—	D	
20893 <i>monogyna Kit.</i>	<i>one-styled</i>					Hungary	1816.	D	W. & K. 1.70
20894 <i>Marachiniana Spr.</i>	<i>Marschall's</i>					Caucasus	1816.	D	
<i>indotira Eichb.</i>									

• 2314. GNAPH A^LIUM L. EVERLASTI.VG. {Gnap/talon, soft down j woolly herbage.) Comp.Card. Vern. 24.

I. GNAPHVLIUM.—Leaves broad.

20895 <i>Ljiteo-fblum L.</i>	yellowish white	Ow	2jl.1.au	Y.w	England	san. pi.	S	8.1	Eng. bot 1002
20896 <i>albescens Sun.</i>	albescent	Jamaica tt. • or 2	... W.Y	Jamaica	1793.	C	co		
20897 <i>sanguineum L.</i>	bloody	^ A or H m.y Jl	C	Egypt	1768.	S	co	Rauw.it285.37	
20898 <i>undulatum L.</i>	waved	^ O cu 1	In.au	Y.w	Africa	1732.	S	co	Di. el. 108.130
20899 <i>obtusifolium L.</i>	obtuse-leaved	O pr 1	Jn.au	Y.w	N. Amer.	1699.	S	co	Di. el. 108.131
20900 <i>purpureum L.</i>	purple-^/ouvrif^	O l or 1) jns	P	N. Amer.	1732.	S	co	Di. cl. 109.133	
20901 <i>involutatum Forst.</i>	involute	£ 1AI cu 1	Jl.au	Br.Y	N. Zeal.	1823	8	co	Bot mag. 2582

II. FILASO. —Leaves narrow; flowers at base naked or involucrated.

20902 <i>americanum L.</i>	American	£ E2 un	1 jl.au	Y.Br	Jamaica	1815.	S	co	
20903 <i>pennsylvanicum W.</i>	Pennsylvanian	Q un	1 jl.au	Y.B	Pennt>yv.	1815.	S	co	
<i>spatulatum Lam.</i>									
20904 <i>monenum IV.</i>	mountain	O un	i jn.au	Br.Y	S. Europe	1820.	S	co	
<i>Filago minntana Pers.</i>									
20905 <i>sylvaticum L.</i>	wood	^ A un	1 au	Y.Br	Britain al. pas.	S s.l	Eng. bot. 913		
<i>Filago syl&tica W.</i>									
20906 <i>rectum Sm.</i>	upright wood	£ A un	1 au	Y.Br	Britain sa.pas.	D	co	Eng. bot 124	
<i>Filago recta W.</i>									
20907 <i>rectatum W.</i>	compressed	O un	1 au	Br	M. Video	1819.	S	co	
<i>Filum Vil.</i>									
20908 <i>alpinum Light.</i>	supine	£ A un	i jn.jl	Y.Br	Scotland sc.alp.	D	co	Eng. bot 1193	
20909 <i>pisillum Ilav.</i>	small	£ A un	lin jn.jl	Y.Br	Austria	1820.	D	co	Kr.sil.41
<i>hub</i>									
20910 <i>spiniferum Lk.</i>	spherical	£ A un	1* jn.jl	W	N. Holl.	1819.	D	co	
<i>Filago sphे'rica Lk.</i>									
20911 <i>cephaloideum IV.</i>	head-like	O un	jn.jl	Y.w	N. Holl. ?	1823.	S	co	
<i>Filago cephaloidca I.k.</i>									
20912 <i>uliginosum L.</i>	marsh	O un	1 au	Y.Br	Britain watpl.	S	co	Eng. bot 1194	
<i>multicaule W.</i>									
20913 <i>germinicum Sm.</i>	German	O un	f jn.au	Y.Br	Britain san. fi.	S	co	Eng. bot 948	
<i>Filago germanica L.</i>									
20914 <i>gfilicum Sm.</i>	French	O un	jn.au	Y.Br	Britain san. fi.	S	co	Eng. bot. 2369	
<i>Filago gallica L.</i>									
20915 <i>pyramidalatum IV.</i>	pyramidal	O un	i Jn.au	Br.Y	S. Europe	1779.	S	co	
<i>Filago pyramidata L.</i>									
20916 <i>minimum Sm.</i>	least	O un	jl.au	Y.Br	Britain sa.pas.	S	co	Eng. bot. 1157	
<i>Filago alpestris Presl</i>									
20917 <i>aryense Pers.</i>	corn-field	O un	jl.au	Y.Br	Europe	1804.	S	co	
<i>Filago arvensis L.</i>									
20918 <i>Lagopus W.</i>	Hare's-foot	O un	jl.au	Y.Br	Siberia	1820.	S	co	
<i>Filago Zagpus Pers.</i>									

2315. SPIRALEPIST D. DAN SPIRALEPIST. {Speira, a spire, lepis, a scale; twisted.) Comp. Card. Vern. 4.

20919 <i>squarrtsa D. Don</i>	squarrose	tLi_Jor	1 jl.o	P	C. G. H.	1816.	C	s.p.l	Jac.fr.3.4
<i>Cnaphalium squarrdsum L.</i>									
20920 <i>glomerata D. Don</i>	glomerated	^ iAI or ft	mr.s	P.a	C. G. H.	1774.	S	s.p.l	
<i>Gnaphalium glomeratum W.</i>									
20921 <i>dcclinkta D. Don</i>	declinate	£	jl.o	Br	C. G. H.	1787.	S	s.p.l	
<i>Gnaphalium declinatum L.</i>									
20922 <i>modesta G. Don</i>	modest	£ tAJor	1 JL0	R	C. G. H.	18S6.	S	s.p.l	Bot mag. 2710
<i>Gnaphalium modestum Hook.</i>									

2316. ZEONTOPO'DIUM R. Br. LION'S FOOT. {Leon, Won, pout, foot; heads of fl.) Comp. Card. Vern. 1.-9.

20923 <i>helveticum J. Don</i>	Swiss	^ A cu	{ jn.jl	Y	Austria	1776.	S	p.l	Bot. mag. 1958
--------------------------------	-------	--------	---------	---	---------	-------	---	-----	----------------

2317. FILA*GO L. COTTON Rose. (Filum, a thread: covered with.) Comp. Card. Vern. 1.

80924 <i>pygmaea L.</i>	pygmy	O un	i jl.au	Br	S. Europe	1629.	S	co	Cav. ic. 1.36
<i>£Jax pygmaea Gae.</i>									

• 2318. ANTENNAHIA R. Br. ANT. (Awns of pappus like antennae of insects.) Comp. Card. Vern. 7.—11.?

90925 <i>contorta D. Don</i>	twisted-leaved	£ ^ 1 pr	2 jl	W	Nepal	1821.	1)	co	Bot reg. 605
90M36 <i>triplofervis D. Don</i>	three-nerved	j£ AJ pr	au	W	Nepal	1823.	D	co	Bot. mag. 2468
20927 <i>dioica R. Br.</i>	dioicinus	^ A pr	\ my,jl	Pk	Britain	...	1)	p.l	Eng. bot 267
20928 <i>alpina R. Br.</i>	alpine	^ pr	\ jn.jl	Pk	Alp. Eur.	1775.	D	p.l	
80929 <i>carpathica R. Br.</i>	Carpathian	5 A P'	i jn.jl	Pk	Carpa.mo.1775.	D	p.l	Fl. dan. 332	

2319. METALA^SIA R. Br. METAL A si A. (Meta, change, lastos, hairy; lvstumcdO Comp. Card. Vern J5.

20932 <i><eriphiolecs R. Br.</i>	Seriphium-like	tt. J pr	3 ... Y	C. G. H.	1825.	C	p.l		
20933 <i>muricata R. Br.</i>	muricated	tt. J br	2 jn.au	W	C. G. H.	1816.	C	s.p	
20934 <i>divergens D. Don</i>	diverging	tL L_J or	2 jn.au		C. G. H.	1816.	C	s.p	Bur. af. 73.2
<i>Gnaphalium divergens Than.</i>									
20935 <i>festigia Xta R. Br.</i>	peaked	tt. J or	3 my.au	W	C. G. H.	1812.	C	s.p	Pet. g. 12.7.3
<i>Gnaphalium fastigiatum Thun.</i>									
80936 <i>mucronata R. Br.</i>	pointed	tt. _ or	2 my.au	W	C. G. H.	1824.	C	s.p	Bur. af. 66.3

2320. ASTE'LMA R. Br. ASTELMA. (*A*, privative, *stelma*, crown; construe, of *tt.*) *Comp. Card. Vern.* 10.—

I. CONNIVE'NTIA.—*Involucres connivent.*

- 20937 exfimium *B.Br.* fine a. i_J spl 3jl.au C C. G. H. 179a S up Bot.mag.S32
 20938 milleflirum *D.Don* 1000-flowered a-L-jel 1 jus Pa.P C.G.H. 1802. C s.p
Gnaphium milleflorum Thun.
 20939 sparkle *D.Don* *suira* Ueaved tt-uJor 2 jl.o W CG.H. 1801. C s.p Bot.rep. 262
2?hchyrum spirale W.
 20940 Stehelina *D.Don* Stahelina Jtte tt.i_jor 1^year W C.G.H. 1801. C s.p Bot.rep. 428
*.flichry*um SteheFma W.*

II. ■■■■■.—*Involucres radiate.* . „ „ „ „

- 20941 varieghtum *D.Don* variegated Jelichrysum variegatum W. tt.t_Jor 2 my.jn Br.WCG. H. 1801. C s.p
 20942 speciosissimum *D.Don* showiest JEUchr&sum speciosissimum W. ■■■■■ 8 jU W CG.H. 1691. C s.p Bot. rep. 51
 20943 imbnneatum *D.Don* imbricated Jelichrysum imbricatum W. ■■■■■ 2 jLo W CG.H. 1820. C s.p Pet.g.5.10
 20944 can&cens *D.Don* canescent flichry*um canescens W. <t L—Jor 1\$ap.au P C G. H. 1794. C s.p Bot. mag. 420
 20945 retCirtum *D.Don* twisted back JSlichry*um returatum W. n-i_lor 1 jl.au W CG.H. 1732. C s.p Di.eL322.415
 20946 frägrant) *Swt.* fragrant fUchry*um frägrans And. rep. tt-uJor Iftjl Pk CG.H. 1803. C s.p Bot.rep.561

2321. A T H R F X I A Avr ATHHIIA. (*Apnv.*, *thrix*, a hair; absence upon receipt.) *Comp. Helidnthea.* 1.
 20947 capensis Ker Cape tt-Ljor 3 ap R C.G.H. 1821. C p.l Bot.reg.681

2322. XERA'NTHEMUM L. XBBANTHEMUM. (*Xeros*, dry, *anthemon*, a flower.) *Comp. Card. Vern.* &
 20948 finnum L. annual O or a jl.au P S. Europe 1570. S a.f Jac. au. 4.388
 20949 inapertum !V. unopened O or 2jl.au P S. Europe 1680. S co M.h.6.12.9
 20950 orientle IV. oriental O or 2^jl.au W Levant 1713. S co

•2323 /FELICHRYWU Pen HEUCHIB. (*Helios*, sun, *chrysos*, gold; blossoms.) *Comp. Card Tern.* 43.—

20951 grndifloruml) *D.Don* f?reat.flowered «. i_J el 3 jn.au W C.G.H. 1731. C s.p Bur. ul 76.1

Criaplium grandiflurum L.
 20952 fnMicana *D.Don* shrubby *L j o r 3jn.au Y CG.H. 1779. C co Bot reg.780*~
Astelma fr&ticans B.R. Crnaphium fr&ticans X. grandifloum And.H. 1809. C s.p

20953 crispum *D.Don* curled £.IAlcl - 6 ... Pk CG.H. 1809. C s.p
Gnaphium crispum L.

20954 arboreum >. *D.Don* tree tt L_jel 6 f.au W C. G. H. 1770. C s.p

Cfnaphlium arboreum X.

20955 divaricktum *D.Don* spreading a.L-Jel 3 jn.au W C. G. H. 1820. C s.p Brey. pr 18.3

GnaphaliuM divaricatum L.

20956 congestum *D.Don* close-headed Gnaphilium congestum Lam.

20967 patulum *D.Don* spreading (rnaphSdium pfitulum L.

20958 discolorbum *D.Don* discolored CnaphaliuM discolorbum L.

~0959 cephal.Mes 2). *D.Don* large-headed tt-Ljor 4 Ja.n Pk CG.H. 1789. C co Pluk.ph.410.2

GnaphaliuM cephalotos W.

20960 diosmaftum £w*. Diosma-leaved tt. LJ or lftmr.au W CG.H. 1812. C co Ven.mal.74

GnaphaliuM dtosmaftum Ven.

20961 cricoedes *D.Don* Heath-like tt. LJ or lft mr.au Pk C. G. H. 1774. C s.p B*ot mag. 435

GnaphaliuM pricoedes h.

»0962 teretiflum *D.Don* round-loaved tt. LJ or 1 mr.au Br CG.H. 1812. C co Bur.af.77.3

GnaphiliuM teretiflum L.

~0963 tephrodes *Swt.* cinereous tt LJ el 3 jn.au Y.w C. G. H. 182a C s.p

GnaphiliuM tephrodes Lk.

~0964 acuminitum *Swt.* acuminate tt. I el 3 jn.au W CG.H. 182a C s.p

GnaphiliuM acuminatum Lk.

■■■■■ lasiocatlon *Swt.* woolly-staid tt. ■■■■■ 3 jn.au W C. G. H. 182a C s.p

GnaphiliuM lasiocatlon Alon Lk.

■■■■■ Stoc chds *D.Don* %Stcechas, comjhr. tt. or 2 jn.o Y Europe 1629. C co Bar.ic.410

GnaphiliuM Stoc chaa L.

■■■■■ ct'micum *Swt.* conic tt. or 2 jn.o Y S. Europe 1824 C co

GnaphiliuM cönicum Lk.

20968 angustiflum Swt. narrow.leaved tt. _J or 2 jl.s Y Naples ... C co Bar. ic. 1125

GnaphiliuM angustiflum Lam.

20969 ignisces D. *fxeTV-Jioxvercd* it. LJ or 2 jn.o R CG.H. 1731. C s.p

GnaphiliuM ignisccens L

-0970 microphf Hum *D.Don* small leaved tL^J el lft jn.o Crete 1823. C s.p

20971 dasyathium Suit, thick-flowered tt. ^J or 4 jn.au Y CG.H. 1812. C co

GnaphiliuM dasy&nthum TV. en.

WJJ2 crassifolium *D.Don* thick-leaved ■■■■■ 1 jl.s Y CG.H. 1774. C s.p

GnaphiliuM crassifolium L.

20973 maritimum *D.Don* sea tt.i_Jor 4 jnau W.T CG.H. 1772. C co Bur.af.77.2

GnaphiliuM maritimum L.

120974 apiculatum *D.Don* apiculate GnaphiliuM apiculatum Lab.

20975 tiacidum G. *Don* flaccid GnaphiliuM flaccidum Wan.

20976 foetidum *D.Don* fetid GnaphiliuM foetidum L.

20977 cylindricum *D.Don* cylindrical j£iAJpr 1 jn.jl Y CG.H. 1780. C s.p Pluk.ph.298.4

20978 orientMe *D.Don* oriental ■■■■■ or lft ap.au Y Africa 1629. C s.p Com. h. 2.55

GnaphiliuM orientMe L.

20979 arenarium *D.Don* sand ■■■■■ A or 1 Jl.s Y Europe 1739. D co Bot. mag. 2159

GnaphiliuM arei*rium L.

20980 graveolens *Swt.* heavy-smelling ^ A el 1 jn.jl Y Tauria 1819. D co

M GnaphiliuM graveolens Bkb.

20981 odoratissimum *Swt.* sweetest-acld £ ^ fra 2 ap.au Y CG.H. 1691. C s.p Mil.ic. 1.1S1.8

~0982 candidissimum *D.Don* whitest ■■■■■ A or 2 my.jl Pa.Y Caspian 182a D co

20983	rbitans <i>D. Don</i>	Bhinng-jtiKi (<i>Jnaphalium rbitans</i> L.)	tf A lor	1 jn.jl	R.v	C G. H. 1731.	C s.p	D!. el. 107.127
20984	cymosum <i>D. Don</i>	cynose <i>Gnaphalium cymosum</i> L.	jp A lor	1 ap.au	Y	Africa	1731.	C co Di. eL 107.128
£0985	hehanthemifblum <i>D. Don</i>	Heli.-lvd <i>Gnaphalium helianthemifolium</i> L.	«LL_»or	1 Jl.o	W	C. G. H. 1774.	C co Vol. nor. 194	
20986	dealbatum <i>Swt.</i>	whitened	tf i or	1} year	W	V. Di. Isl. 1812.	C co	Lab.n.h.2.190
20987	ffigidum <i>D. Don</i>	shining	* I or	2 f.o	Y	C. G. H. 1774.	S s.p	Bot. mag. 414
20988	argenteum <i>Swt.</i>	silvery	n. I br	2 apil	W	C. G. H. 1800.	C co	Bot.reg.552
20989	rlgidum <i>D. Don</i>	rigid-leaved	tt. t or	limyjn	W	C. G. H. 1801.	C co	Bot. rep. 387
20990	herbaceum <i>Swt.</i>	herbaceous	f A id	11 jl.s	Y	C. G. H. 1802.	S co	Bot. rep. 487
	spindens <i>B. M.</i>							
20991	paniculatum <i>Swt.</i>	panicleil	tt.i J or	2 jn.s	W	C. G. H. 1800.	S co	Bur. af. 67.1
20992	bracteatum <i>D. Don</i>	bracted	O or	4 Jl.o	Y	N. Holl. 1799.	S co	Bot rep. 375
20993	italicum <i>G. Doy</i>	Italian	tt. J or	2 jl.o	Y	Italy	1826.	C co
	<i>Gnaphalium italicum</i> Wnl.							

2324. LEUCOSTE'MMA *D. Don* LEUCOSTEUMA. (*Leukos*, white, *stemma*, a crown; *fit.*) *Comp. Card. Fern.* 1.
20994 vestUum *D. Don*, clothed **tl.Jor** «2 **jk** » W w Cr>Gn Hx 1774. S s.p W. at **bb.1**
Jlichrysum vesttum L.

2325. PHCENO'COMA *D. Don* PHECOMA. (*Phoinos*, bloody, *home*, hair; invol.) *Comp. Card. Vern.* L
20995 prolifera J. Dwi proliferous • uJ el 4 au n C C. G. H. 1789. C s.p Bot. reg. 21
Jlichrysum proliferum W. Xerantheinum proliferum L.

2326.	APHELEQCIS <i>D. Don</i>	APHELEXLJ. (<i>Apheles</i> , simple, <i>ens</i> , habit: plants.)	<i>Comp. Card. Vern.</i> 4.
20996	humilis <i>D. Don</i>	humble «i_J or 2 my.jn Pk	C. G. H. 1810. S
		flichr^sum spect&bile hod. pinifiliura Pers.	s.p Bot. cab. 59
20997	*esamoldes <i>D. Don</i>	Sesamum-like tl\ or S ap.jn P.w C. G. H. 1739. S	s.p Bot. mag. 425
		flichrysum xesamoides W. Xeranthemum ^esamoides L.	
£0998	fasciculata <i>D. Don</i>	fascicled a. L_J or 2 mr.s P.v C G. H. 1799. S	s.p Bot. rep. 242
		.Elichrysum fasciculatum W.	
2	filba	white-flowcrcd tt. uJ or 2 mr.s	W C. G. H. 1799. S
3	r&bra	red-flowered tt. LJ or 2 mr.s	R C. G. H. 1799. S
4	versicolor	party-colored a. i.J or 2 mr.s	Va C. G. H. 1799. S
20999	cricoldes <i>Swt.</i>	Heath-like tt. L_J or i ap.jn IV	C. G. H. 1796. C
		Elichrysum ericoides Pers.	s.p Brey. ic. 16.2
			s.p Lam.il. 693.2

2327. (7ARPE)5IUM L. CARPESIUM. (*Karpesion*, a bit of straw: resembl. in lvs of invol.) *Comp. Card. Vern.* 2.
21000 cernuum L. drooping ^ u n i jlau Y Austria 1739. D co Jac. au. 3.204
£1001 abrotanoides L. Southernw.-like^ A u n 2 jl.au Y China 1763. D co Osb. 10

2328. BRACHYL^NA R.Br. BOACHYUSNA. (*Brachys*, short, *Itena*, cloak; calyx.) *Comp. Card. Vern.* 1.
21002 nereifolia Swt. Oleander-leaved* i.J or 4 aitn W C. G. H. 1752. C s.p
^accharis nereifolia L. Tarchonanthus dentatus Thun.

2329. EA'CCHARIS L. PLOUGHMAN'S SPIKENARD. (*Bacchus*, wine; odour.) *Comp Card. Vern.* 11.—101.
I. INTEGRIFOMA.—Leaves entire, flat s flowers corymbose, or paniculate.

£1003	angustifolia Ph.	narrow-leaved * i pr 2 jl.s	W N. Amer. 1812. C co
II. SBRRATTFOLI*.—Leaves serrated or toothed; flowers corymbose			
21004	ivafolia L.	Iva-leaved * i _J pr 3 ji.au	W America 1696. C Lp Schk. han. 3.244
21005	Aalimifolia L.	Halimus-lvd St or 4 o.n	W N. Amer. 1683. C co Schm. ar. 82
21006	DioscoridM W.	Dioscorides's * J or 4 au.n	Levant ... C co Rauw.it 54
21007	Indica L.	Indian sH(H)or 3 —	E. Indies 1819. C sl Brey. c. 70
£1008	glutinosa Pers.	clammy ID or 3 —	Peru 1824. C 81
21009	parviflora Pers.	small-flowered flkCDor 3 —	Peru 1820. C s.l
	Molina parviflora Fl. per.		
21010	adnata Hum.	adnate H • un 6 au.n P & Amer. 1823. C co	
III. SESSIMFL6R. —Leaves serrated or toothed; flowers sessile, glomerate, or spicate.			
21011	conferta Kth.	crowded I Q or 3. au. Mexico 1826. C s.l	
£1012	glomculiflura Mx.	cluster-flwd A or 3 au.o W N. Amer. 1817. C co	
IV. SOLITARI/EFLOHA.—Leaves entire, flat; flowers solitary.			
£1013	jcoparia Swx.	Broom i D o r 3 jlau Crea Jamaica 1820. C JI Br. jam. 34.4	
	Calea scoparia L.		

*2330. COWZAL. FLEABANB. (*Konis*, dust; powder sprinkled to kill fleas.) *Comp. Card. Vern.* 42.—62.

I. Fnjribss.—Shrubby; leaves petioled or sessile, oblong-lanceolate.

		i. CORYMBOSE.—Flowers corymbose or paniculate.	
£1014	odorata L.	sweet-scented H O fra 2 jn.au P India 1759. C pi PI. ic.97	
£1015	balsamifera L.	balsam-bearing * U un 2 jl.s Br E. Indies 1822. D co Ru. am. 6.24.1	
£1016	carolinensis Jac.	* 1 un 5 jl.o P Carolina 1821. C co Jac. ic. 585	
£1017	inclusa H. K.	cut-leaved * L J un 3 jn.au P C. G. H. 1774. C p.l	
		ii. PEDUNCULAR.—Peduncles solitary.	
21018	Candida L.	white * Jun 1 jn.jl Y Candia 1714. C co Bar. ic. 217	
£1019	verbascifolia W.	Mullein-leaved * U un 1 ... Y Sicily 1808. C co Boc. sic.31.2	
21020	spatulata Lfc.	spatulate £ E3 un 3 jn.au B 1825. D pl	
£1021	/moniliflora Sm.	Limonium-lvd * \ un 1 jn.jl Pa.Y Greece 1740. C s.l	
	Candida W. not L.		
		iii. SUBSESBILIFLORA.—Flowers subsessile, glomerate^ or spicate.	
£1022	arbore'sceni W.	arborescent si un 6 n.d B Jamaica 173). C p.l	
21023	rigida Swx.	IQun 2 . . . Jamaica 1820. C p.l	
		iv. ANOUSTIF&LIJE.—Flowers subsessile, glomerate, or spiked; leaves narrow, subUnear.	
21024	rnuloides H. K.	Inula-like si LJ un 1 jl.au P Tcnerriffe 1780. C co Jac. ic. 1.171	
£1025	rupestris L.	rock si un 1 . . . Arabia 17K. C co Schm. ic. Sri	
£1026	Tentoi Lk.	Trone's si un 1 jl.au ... S. Europe 1824. C co	
21027	sax'tius L.	stone si un 1 jl.au Br S. Europe 1640. C co	
21028	geminiflora Ten.	twin-flowered si un 1 a.u.s Br 1823. C co	
21029	sordida L.	sordid si un 1 jl.s Br & Europe 1570. C co	
21030	sericea H. K.	silky si un 1 jl.s Y Canaries 1779. C p.l	
		v. DBCURRE'NTIA.—Leaves decurrent.	
£1031	rugosa Vahl	rugose • L_J «n 6 n Y Brazil 1772. C .	

II. HBRBA\:\:E2B. - *Herbaceous ; leaves petiolate or sessile, undivided, broad, ovate, oblong, or lanceolate.*i. CoRYMBBSJS. — *Flowers corymbose and panicle.*

21032 squarrbsa L.	squarrose	3t O> w	2 jl.au	Y	Britain	ch. pa.	S	co	Eng. bot. 1195
21033 paniculata TV.	panicled	£ [29 un]	1 jlau	P	E. Indies	1818?	S	co	
21034 axillaris Z. <w>	axillary	■■■ un	1} au.o	Y.		182a	S	co	
21035 amce'na Lk.	pleasing	£ [23 un]	3 jl.au	P	Congo	1824.	S	co	
21036 panitaf.K.	spreading	O un	■■■	Y.p	China	1758.	S	co	Mil. ic. 2.247
21037 blfrons L.	two-fronted	■■■	■■■ au.8	Y	N. Amer.	1739.	D	p.l	Pluk. al. 87.4
21038 camphorht Ph.	camphor-sctd	2k A un	■■■ au.o	P	N. Amer.	1704.	D	co	Di. el. 87.4
21039 maryl&ndica Mx.	Maryland	tf Al un	■■■ au.o	P	N. Amer.	...	S	co	Di. el. 89. J05
21040 purpurascens Swz.	purpurascent	0 un	■■■ jl.au	P	Jamaica	1800.	S	co	SL jam 1.159.1
21041 cinerea L.	Py	rm un	■■■ jl.au	P	E. Indies	1816.	S	co	Ru.am. 6.14 1
21042 prolifera Lam.	Srohferous	ED un	■■■ a.us	Y.p	China	1767.	S	co	Ru.am. 5.104.1
21043 hirsuta L.	Egyptian	£ [20 un]	■■■	Y	Egypt	1778.	S	co	Jac. vin. 3.19
21044 sgyptaca H. K.	Gouan's	£ OI un	■■■ jl.au	Y	Canaries	1772.	S	co	Jac. vin. 3.79

ii. PAUCIFLOR. — *peduncles one or two-flowered.*

21046 amplexicaulis Lam.	stem-clasping	Qu n	1 jl.au	...	E. Indies	1820.	S	co	
21047 fastigiata IV.	peaked	O un	2 Jn.jl	P	Senegal	1820.	S	co	

iii. SESSHIPLORA. — *Flowers sessile.*

Chinese	£ Ol un	2 jl.au	Y.p	China	1796.	S	co	Itu. am. 6. Ik 2
---------	---------	---------	-----	-------	-------	---	----	------------------

iv. LINEARIF^LIA. — *Leaves narrow, linear.*

Sicilian	O un	1 au.s	Y	Sicily	1779.	S	co	Boc. sic. 31.4
----------	------	--------	---	--------	-------	---	----	----------------

v. PINNATI'FiDf. — *Leaves at base, or all pinnatifld.*

eared?	O un	1 au.o	W	E. Indies	1818.	S	co	
fetid	j£ iAl un	2 a.us	Y	Africa	1724.	D	co	Mil. ic. 2.233

pinnatifid £ E3 un 2 jl.au ... E. Indies 1816. S co

vi. DECURENTES. — *Leaves decurrent.*

Thapsus-like ^ A un	2 jl.s	P	Casp. Sea	1806.	C	co	
twiggly j£ IAI un	2 a.us	?	America	1783.	D	co	SL jan.: 1.152.1

III. GLOMERA^TA. — *Flowers glomerate.*

glomerate	O un	1} jn.au	P		1825.	S	co	
-----------	------	----------	---	--	-------	---	----	--

*2331. MJTDIA Mol.

21056 visciosa Coy.

sativa Af.

21057 mellosa ■■■

*2332. ERI'GERON L. ERIGERON. {Er, spring geron, old man; hoary in spring.) Comp. Card. Vern. 35.—56

I. MultiFLRi. — *S/« many-flowered; flowers corymbose, paniculate.*i. LATIF&LII. — *Leaves broad, oblong, or lanceolate.*

various-leaved i Q pr	1 Jl	■■■	N. Amer.	1640.	S	co	Fl. dan. 486
-----------------------	------	-----	----------	-------	---	----	--------------

various-leaved i Q pr 1 Jl ■■■ N. Amer. 1640. S co Fl. dan. 486

Chinese O or 1 jl.s W China 1818. S co Jac. sc. 3.303

Chinese O or 1 — — S. Europe 1827. S co

southern O or 1 — — — 1823. D co

Aster-like & A Pr Ii jl.au W Huds. Bay 1776. D co

purple Jfc A pr 1 jl.au P N. Amer. 1816. S co

purple Jfc A pr 1 jl.au W N. Amer. 1816. S co

strigose i CD pr 1 jl.au B Britain gra.pa. S co Eng. bot H58

sharp ^ C pr lijl.au B Britain gra.pa. S co

rock 5 A pr f jl.au P Switzerl. 1819. S co

great-flowered ^ A pr f jl.au P Switzerl. 1819. S co

Chile Q cu 2Jl au.o Y Chile 1816. S co

Conyza chillensis Spr.

Attic ■ A or UJl.au P Attica 1816. S co

Podolian ■ Q or 1 jn.au P Podolia 1818. S co

Canadian O pr 1 au.s W England rubble. S co Eng. bot. 2019

little O pr iau.s W N. Amer. 1823. S co

strong-smelling O pr Ii jl.s P S. Amer. 181 a S co

strong-smelling O pr Ii jl.s Y S. Europe 1633. S co

21073 graveolens L. Buenos Ay res P S. Amer. 1732. S co Di. el. 257. 334

21074 bonariensis L. Buenos Ay res P S. Amer. 1732. S co Di. el. 257. 334

21075 philadelphicus IV. Philadelphian ^ A pr Ii jl.au P N. Amer. 1778. D co

ii. LiSEARiFOLii. — *Leaves narrow, linear.*

twisted £ Q or 2 jlau PPP 1826. ■ co

smooth-leaved S Q u n 1 jl.au W Cayenne 1822. ■ co

Flax-leaved O pr 1 jl.au P S. Amer. ■ ■

long-leaved ■ A pr ■ jl.au P N. Amer. 1820. D co

divaricate ■ CD pr ■ jl.au W Mississippi 1818. S co

II. SUBMI'LTIFLORI. — *Stem many-flowered i branches one-flowered.*

Villars's ^ A pr 1 Jlau P Piedmont 1804. S co Bot reg. 583

Carolina 5 A pr 1 jlau P N. Amer. 1727. D s.p. Di. el. 306.394

III. PAUCIFLORI. — *Stem few-flowered.*

Daisy-leaved ■ A pr "w Jlau" P* N. Amer. 179tt D co

Caucasian ■ ^ pr f jl.au P Caucasus 1821. D co

naked, stalked ■ A pr 1 jl W N. Amer. 1812. D co

smoothish ■ Aor 1 jlau B N. Amer. 1825. D s.p.

Jamaica O pr 1 jl.s P Jamaica 1818. S co SL jam. 152.3

21088 m6nte videensi* Spr. Monte Videan O un 1 jl.au W? Montevid. 1827. S co

IV. UNIFLURI. — *Stem generally one-flowered.*

alpine ■ pr 1 jl P Scotland al.rills. D s.l Eng. bot. 464

one-flowered ■ ■■ Jl.aus P Scotland highl. D co Eng. bot 2416

composite ■ ■■ jl.aus W.R N. Amer. 1811. Deo

Grass-like ■ ■■ jn.au P Siberia 1824. D co Gm. si. 2. 76.2

V. FRUTICOSUS. — *Stem shrubby.*

21093 glaucus B. R. glaucous il LJ pr 1 year P S. Amer. 1812. C co Bot. reg. 10
<4'ster bonariensis Sm.

•2333. rUSSILA'GO £. COLTWOOT. (<i>Tussis</i> , a cough; med.qual.)	I. UNIFLBBA — <i>Scape one-flowered.</i>	Camp. Jacobite. 13.—15.
21094 <i>alpina L.</i> alpine	£ mr.my L.P	Austria 1710. D co Bot. mag. 84
21095 <i>discolor Jac.</i> discolored	i ap.my L.P	Austria 1633. D co Jac. au. 3.247
21097 <i>sylvestris Sco.</i> wood	1 ap.my L.P	Austria 1816. D co Jac. au. 5 ap. 12
21097 <i>Färfera L.</i> <i>Parfara, common</i>	i mr.ap Y	Britain moi.pL D co Eng. hot 429
2 folia variegatis variegated-lvd	i mr.ap Y	Britain gard. D co
	II. THYBSFL6R«. — <i>Soppes ihysiferous.</i>	
21098 <i>frigida L.</i> frigid	* A or J my Pa	Lapland 1710. D co Fl. dan. 61
2)099 <i>sagittata Ph.</i> sagittate	I ap.my ...	N. Amer. ... D co
21100 <i>laevigata JV.</i> smooth	J my Y	Bohemia 1816. D co Gm.sL2.146.49
bohémica <i>Hoppe</i>		
21101 <i>fragrans Vil.</i> sweet-scented		Italy 1806. D co Bot. mag. 1388
21102 <i>alba L.</i> white <i>Butter Bur</i>		Europe 165J. D co Fl. dan. 524
21103 <i>nivea VU.</i> snowy-leaved		Switzerl. 1713. D co lletz. 2. 3
paradoxa <i>Retx.</i>		
21104 <i>Petasites L.</i> Petasites, com. But. B.		Britain m.me. D co Eng. but 431
2 h̄ibrida E. B.	hybrid	Britain m me D co Kng.bot.430
21105 <i>spirin Retx.</i> spurious		Germany 1790. D co Refs. 1. 2
tomentosa <i>Ehrh.</i>		
21106 <i>palmata H. K.</i> palmate-fanwd		Labrador 1778. D co H. kew. 3.11
21107 <i>rigida Kth.</i> rigid	£ IAJ pr *	Quito 1828. D p.e
Doronicum peruvi&num Lam.		

2334. WERNEPRM Kth. WBBNBBIA. (A. G. Werner, the celebrated mineralogist) Comp. Jacobhe. 1.
21107 *rigida Kth.* rigid £ IAJ pr * Quito 1828. D p.e

* 2335. SENETCIO L. GROUNDSSEL. (<i>Senex</i> , old man; naked receptacle like a bald head.) Comp. Jizc.104.—	I. FBUTICOSL — <i>Shrubby ; rays spreading £ leaves undivided j flowers corymbose.</i>		
	I. LATIFOLH. — <i>Leaves broad, ovate, oblong, or lanceolate.</i>		
21108 <i>Ilicifolius L.</i> Ilex-leaved	■ L or 3 Jn.j, Y	C. G. H. 1731. £ Com. r. 42	
21109 <i>Aalimifolius L.</i> Halimus-leaved	■ L or 3 Jn.j, Y	C. G. H. 1723. £ Duel. 104. ICt	
21110 <i>sohdagineus Berg.</i> Solidago-hke	■ L or 2 Jn.j, Y	C. G. H. 1824.	
21111 <i>Panceus//.JC</i> ayteur-leaved	■ L or 3 Jn.j, Y	C. G. H. 1774.	
21112 <i>rigidus L.</i> haxA-leaved	■ L or 3 Jn.s, Y	C. G. H. 1704. £ Com. h. 2.75	
21113 <i>IIIicifolius Lk.</i> lac-Jiowcred	■ L or 6 myjl Li	C. G. H. ? 182H.	
21114 <i>rigescens Jac.</i> rigescenUleaved	■ L or 3 Jl.au, Y	C. G. H. 1815. £ Jac. c. 5.6.1	
21115 <i>argutus ICtA.</i> d. ■ L or 3 Jl.au, Y		Mexico 1827. £	
	ii. LINBARIFOLII. — <i>Leaves narrow, sublinear.</i>		
21116 <i>rosmarinifolius L.</i> Rosemary-lvd	■ L jp 3 Jl.au, Y	C. G. H. 1827. C l.p Jac. ic 3.587	
21117 <i>Isper H. K.</i> rough	■ L Jpr 3 Jl.au, Y	C. G. H. 1774. C p.l	
21118 <i>longifolius L.</i> long-leaved	■ L Jp or 3 au.n, Y	C. G. H. 1775. C p.l Com. h. 2.71	
21119 <i>calcarius J&A.</i> chalky	■ L J un 3 Jl.au, Y	Mexico 1824. C l.p	
21120 <i>cineraroides Kth.</i> Cineraria-like	■ L <Jor 2 Jl.au, Y	Mexico 1826. C l.p	
	II. LiDivifei. — <i>Herbaceous ; leaves undivided ; flowers corymbose ; rays spreading.</i>		
	i. LATIFLUI. — <i>Leaves broad, ovate, oblong, or lanceolate.</i>		
21121 <i>nemore'nsis Z.</i> grove	3 Jl.au, Y	Austria 1785. D co Jac. au. 2.184	
21122 <i>ovatus JV.</i> ovate	3 Jl.o, Y	Germany 1823. D co	
21123 <i>midropophyllus Bleb.</i> small-leaved	4 Jl.au, Y	Caucasus 1818. L co	
21124 <i>croaticus Kit.</i> Croatian	4 Jl.au, Y	Hungary 1805. I) co W.&K.2.143	
21125 <i>sarracnicus L.</i> Sarracnic	6 Jl.o, Y	Britain mol.pl. 1) co Eng. bot. 2211	
21126 <i>Tournefortii Lap.</i> Tournefort**	3 Jl.au, Y	Pyrenees 1818. 1) co Bui. ph. 43.146,	
nemoreñtis Gou.	Tersicxftilius Ram.		
21127 <i>palutifl8ua L.</i> marsh		England fens. D c.p Eng. bot 650	
21128 <i>umbribus Kit.</i> shady		Hungary 1815. D £	
21129 <i>coriaceus // K*</i> leathery-leaved		Levant 172a D 'l.p Di. el. 105.125	
21130 <i>Duria L.</i> Doria		Austria 1570. D co Jac. au. 2.185	
21131 <i>cacaliaster Dec.</i> Cacalia-like		France 1772. D s.l	
Cacklia sarracenia L.			
21132 <i>sinuatus Kth.</i> sinuate		Mexico ■ D 81	
21133 <i>oprinus W.</i> autumnal		C. G. H. 1817. D 81 Jac.sc. 3.204	
lanceus Jac.			
21134 <i>vdrnus Bio.</i> spring	O un 1 Jn.au, Y	Italy 1816. S co	
leucanthemiflbus Sib.			
21135 <i>trilobus L.</i> three-lobed	O un 3 Jl.au, Y	Spain 1728. S co	
21136 <i>rutundiflbus Lap.</i> round-leaved	A un 3 Jl.au, Y	Pyrenees 1823. D co Boa mu. 61. ic.	
21137 <i>Barrelito Gou.</i> Barretter's	A un 3 Jl.au, Y	Pyrenees 1819. D co Bar. ic. 146	
21138 <i>baldensis Pair.</i> Baldo	A un 3 Jl.au, Y	Mt. Baldol 1820. D co CL h. 4.23. ic.	
	ii. ANGUSTIPCXIUS. — <i>Leaves narrow ; flowers corymbose.</i>		
21139 <i>inifolius L.</i> Flax-leaved	^ A un 2 Jl.au, Y	Spain 1820. D co Boc. mu. 49	
	iii. PAUCIFLORI. — <i>Stemfevo-flowered ; leaves broad, ovate, or lanceolate.</i>		
21140 <i>DorSnicum L.</i> Leopard's Bane^ A or I jis, Y		S. Europe 1705. D e o Jac. au. 2. ap. 45	
21141 <i>uniflbus All.</i> one-flowered	5 A or Jjl.au, Y	Alp. Eur. 1799. D i p Al.ped.17.8	
21142 <i>specibus W.</i> showy	■ IAJ or ^ Jl.au, S	China 1789. D s.l Bot reg. 41	
III. DiviBiFbLii. — <i>Herbaceous ; leaves divided ; flowers corymbose ; rays spreading.</i>			
21143 <i>alrtinus G.</i> alpine	■ n or 2 Jn.au, Y	S' Europe 1683. D co Jac. au. 2.17'	
Cherkria alp'na L.			
21144 <i>cordiflbus G. Don</i> heart-leaved	■ A or 2 Jn.au, Y	Austria 1749. D co Jac. au. 2.17-	
Cineraria cordiflbia L.			
21145 <i>lyratiflbus Ilchb.</i> lyre-leaved	■ A or 2 Jn.au, Y	Austria 1749. D co	
Cineraria alptnaj3 L.			
£1146 aqu&ticus Hud.	water	■ A un 1 my.jn, Y	Britain mar. D bog Eng. bot. 11 -
21147 err&ticus Bert.	strayed	■ A un 1 rayjn, Y	& Europe 1800. D m.s
aqu&ticus LoL			
21148 rupestris Kit.	rock	■ A un 1 Jl.au, Y	Hungary 1805. D s.l W.A.K.2.1.
21149 rontanüs JV.	mountain	O un 1 Jn.au, Y	Switzerl. 1819. S co
parad6xus Hoppe			
21150 <i>ycrnalis Xif.</i> spring	O un 1 ap.jn, Y	Hungary 1803. S co	
21151 <i>JfelsamUaslF.</i> Balsamita	■ A un 2 Jl.au, Y	Pensylva. 1820. D co	
21158 <i>obovatus W.</i> obovate	■ A un 2 Jl.au, Y	N. Amer. 1818. D co	

21153	a ^{re} us <i>L.</i>	Elden	j i	A un	2	jl.au	Y	N. Amer.	1758.	D	L	-	
21154	hastatus <i>L.</i>	Elbert-leaved	l Alun	I Alun	1	ljmy.au	Y	C. G. H.	1722.	D	L	15	
	glutinosis <i>Thun.</i>											Di. el. 152.131	
21155	lyratus <i>Thun.</i>	lyrate	j f	I Alun	2	myjl	Y	C. G. H.	1816.	D	s p		
21156	elegans <i>L.</i>	elegant			2	jn.au	P	C. G. H.	1700.	S	c o		
2	pl&no-r&ber	double red-flwd	n.	O pr	2	year	F	C. G. H.	1700.	C	s l	Bot mag. 233	
3	flöre-älbo	white-flowered	tt.	L j e l	1)	year	W	C. G. H.	1700.	C	81		
4	pleno-albus	double white	tt	\ Jpr	1*	year	W	C. G. H.	1700.	C	81		
21157	incdnus <i>L.</i>	hoary	\	Jpr	1	jlau	Y	Alp. Eur.	1759.	D	s.p	Pluk. al. 39.6	
21158	leucophyllus <i>Dec.</i>	white-leaved	^	A un	1	jlau	Y	S. Europe	1816.	D	81		
	incanüs <i>Gou.</i>		A or										
21159	parviflbris if?.	small-flowered	^		1	jLau	Y	Piedmont	1819.	D	co	ALped.38.3	
21160	carniöficus <i>W.</i>	Carniolan	&		1	jnau	Y	Carniola	1817.	D	co		
21161	Jacobs <i>v</i> il.	common Ragwort	^	A w	2	Jlau	Y	Britain	dry pa.	D	s.l	Eng. bot. 1130	
21162	tenuiflbris <i>Jac.</i>	slender-leaved	f L	A un	2	jlau	Y	Britain	woods.	D	l.p	Eng. bot 574	
UII/65	squillirius <i>L.</i>	squalid	O w		1	jijn.o	Y	England	walls.	S	c o	Eng. bot 600	
21164	dentatus <i>Jac.</i>	toothed	tf	LAI	1	jlau	Y	C. G. H.	1820.	D	co		
21165	chrysanthemiflbris	Poir. Chrys.-lvd	^	A	3	jlau	Y	Sicily	1800.	D	co	Boc. sic. 67	
21166	glaucscnt <i>Spr.</i>	glaucescnt	0		1	jn.au	Y		S	co		
21167	Jycopifbhus <i>Poir.</i>	Lycopus-lvd	\	A un	2	jlau	Y		1820.	D	co	
	canescens <i>Schr.</i>												
21168	arenarius <i>Bieb.</i>	sand	^	A un	1	jn.au	Y	Caucasus	1818.	D	co		
	cinerarioides <i>Viv.</i>												
21169	Othmnae <i>Bieb.</i>	Othona-/ife	f	A or	S	jl.au	Pk	Iberia	1816.	D	co		
	Calacala pinnata <i>W. en.</i>												
21170	delphiniflrlsuraM	Delphinium-lvd	^	A or	I	jn.au	Y	Barbary	1800.	D	co	Boc sic. 51	
21171	adonidifilius <i>Hort.</i>	Adonis-leaved	\	A or	1	my.au	Y	Europe	1800.	D	co		
21172	abrotaniflbus <i>L.</i>	Southernwood-lvd	f	A un	2	jl.o	Y	Alp. Eur.	1640.	D	co	Jac. au. 1.79	
U73	rteiniisis/p<V>>Pers	Wonnow-lvd	f	A	14	jinjl	Y	France	1816.	D	co		
21174	prsalitus <i>Bert.</i>	very tall	\	A un	4	Jlau	Y	Italy	182a	D	co		
	chrysanthemiflbus <i>Mor.</i>												
21175	canadensis <i>L.</i>	Canadian	3t	A or	H	jn.au	Y	N. Amer.	1820.	D	co		

V. REVOLU'TI. — *Rays of flowers revolute.*

- | | | | | | | | | | |
|---------------------|-------------|-------|---|---------|-----------|--------|--------|-------------------|----------------|
| i. | Fauricbus. | — | Leaves pinnatifid; flowers panicled; shrubby. | | | | | | |
| c. | cinerascent | ftUun | 6 myjl | Y | C. G. H. | 1774. | C. p.l | Jac. sc. 2.150 | |
| ii. | iNDivrsi. | — | Leaves undivided; herbaceous. | | | | | | |
| livid | O | un | 1 jl.au | Y | Spain | 1801. | S | co Schk.han.a245 | |
| Nehrod | O | un | 1 jin.au | Pa.P. | S. Europe | 1704. | S | co Bar. ic. 401 | |
| Telephium-lvd | O | cu | 1 jl.au | Y | C. G. H. | 1820. | S | co Jac. fr. 1.1. | |
| sea-green | O | un | lijn.au | Y | Egypt | 1739. | S | co | |
| iii. | DIVI*SI. | — | Leaves divided; herbaceous. | | | | | | |
| thick-leaved | O | pr | i JUU | P | S. Europe | 1815. | ■ | co Bar. ic. 261 | |
| three-flowered | O | un | lj.lila | Y | Egypt | 1776. | ■ | co | |
| ██████ ■ | O | un | n jl.au | Y | Egypt | 1771. | ■ | co | |
| Vervain-leaved | O | un | 1 █ | Y | Lgypt | 1803. | ■ | co Jac. vin. 1.3 | |
| eared | O | un | 1 jn.au | Y | Barbary | 1816. | ■ | co | |
| giant | 3(A | un | 3 jl.au | Y | Barbary | 1827. | D | co Desf. at 2.234 | |
| wood | O | w | 1 jl.au | Y | Britain | | | Eng. but 748 | |
| clammy | O | w | 1 jno | Y | Britain | | | Eng. but 32 | |
| Eruct-leaved | ff. | O | un | 2 jl.au | Y | Europe | 1816. | D | co Bar. ic 153 |
| loose-flowered | O | un | 1 jn au | Y | Numidia | 1827. | S | co | |
| Buckhorn-leaved | O | un | 1 jl.au | Y | Barbary | 1816. | S | co | |

VI. JACOBRÆ — *Rays of flower none.*

- | VI. JACOBIÆ — <i>Rays of flower none.</i> | | | | | | | | |
|---|--------------------------------|-------------------------------|-------------------|--------|-----------|----------|------|-----------------|
| i. FBITICOSI. — <i>Shrubby.</i> | | | | | | | | |
| 21194 <i>reclinatus</i> £.
graminifolius <i>Jac.</i> | reciinate | H [J or | 2 jn.au | P | C. G. H. | 1774. | S co | Jac. Ic. 1.174 |
| 21195 <i>persicifolius</i> L.
2l19G/euuccdanift>liu9/. | Peach-leaved
Pcucedanum-lvd | * i [J or
• [] or | 3 jnjl
3 my.jn | P | C. G. H. | 1820. | E | E |
| CacXha peuredanifolia <i>Jac.</i> | | | | P | C. G. H. | 1816. | E | E |
| ■*197 <i>squamibus</i> W.
21198 ha?matoph^llu8 | scaly
W. en. bloody-lvd | * i [J un
* [] or
2 ap | 3 myjl
Y | Y | C. G. H. | 1820. | C pi | |
| | | | | | — | 1789. | C | co |
| ii. INDIVISI. — <i>Herbaceous; leaves undivided.</i> | | | | | | | | |
| 21199 <i>glomeratua</i> Desf. | clustered | 0 un | ljau | Y | N. Holl. | 1816. | S | co |
| 21200 <i>scaber</i> W.
2ia<)l <i>divaricatus</i> L. | scabrous | J iAl or | 4 jn.apu | Y | C. G. H. | 1700. | C | s-1 |
| 21202 <i>Aieracifolius</i> X. | straddling | J 01 or | J *11 | P | China | 1801. | 1P | |
| 21203 c^rnmuus L. | Hawkweed-lvd | O or | J au | W | N. Amer. | IKK | co | Her. par. S26 |
| | drooping | O or | 1 jLau | V | E. Indies | 1780. | ss | |
| | | | | | | | co | Jac. \in. 3.U8 |
| iii. DIVIJI. — <i>Herbaceous; leaves divided.</i> | | | | | | | | |
| 21204 <i>cacalioides</i> Fis. | Cacaha-hike | Opr | 1 au | Y | Brazil | 182a | S | co |
| 21205 <i>purpircus</i> L. | purple | jg iA) | 2 jl.s | P | C. G. H. | 1774. | D | Jac ic. 3.580 |
| 2120D erub'scens H. K. | erubescence | tf OJ or | 2 jn.o | Pk | C. G. H. | 1774. | 1P | |
| 21207 <i>Psefido-China</i> L. | Bastard China | j‡ iAl or | ijn.au | Y | E. Indies | 1732. | s | Di. el. 258.335 |
| 21208 <i>jaix.lpiuca</i> Thun. | Japan | k A pr | 1 nu | Y | Japan | 1774. | C | |
| 21209 <i>vulgaris</i> L. | common | Groundsel | O w | 1 year | Britain | rubbish. | ss | Eng. bot. 747 |
| 21210 <i>lanuginibus</i> Spr. | woolly | G | O or | 5 o.d | | 1826. | co | |
| ^21211 <i>valerianaf</i> Hivi* Lk. | Valerian-leaved | O or | 4 my.au | Y | Europe | 1800. | co | |
| 21212 <i>arabicus</i> L. | Arabian | [] | 1 il.au | Y | Egypt | 1804. | ss | co |

- | | | | |
|---|-------------------|--|---|
| „2356. LE*Rli< Dec.
21213 nectans Dec. | LEHIA.
nodding | (<i>Leri</i> , a friend of De Candolle's?)
^A•Bu n 1 jnj P | Comp. <i>Labiati</i> Obr* S.
W. Indies 1793. S co PUU1. f7 |
| QH russilago nutans L.
S214 albicans Dec
^A215 puinila Dec. | whitish
dwarf | O un f jlau W Jamaica 1820. S co
if E0 un { inil P Jamaica 1816. S co | |

*2337. il'STER L. STABWORT. {Aster, a star; resemblance in flowers.) Comp. Card. Vern. 156.—									
I. CYMBALAVUI.—Shrubby; leaves broad, ovate, oblong, or lanceolate.									
21216 rcfldxus <i>L.</i>	reflexed-leaved	A or 3	—	C	—	1759.	C	p.l	Bot. mag. 884
21217 tomentosus Schr.	woolly	■■■ or 1*	myjl	Pk	N. S.W.	1793.	C	p.l	Bot rep. 61
21218 erubescens Sieb.	ferruginea Wnl.								
21219 stellulatus Lab.	erubescent	* [J or 3	my.jn	R	N. Holl.	—	C	s.p.1	
21220 myrsinoides Aft.	stellulate	* [J or 2	my.jn	V	V. D. L.	1823.	C	s.p.1	Lab. n. h. 196
21221 sericeus Ven.	Myrsine-like	* [J or 3	my.jn	Pa.P	N. Holl.	1825.	C	s.l.p	Lab.n. h.202
21222 Cymbalaria L.	silky-leaved	[J or 3	my.jn	B	Missouri	1803.	C	s.p	Ven. cel< 33
21223 hratus A. Af.	Ivy-leaved	• [J or 2	my.jn	W	C. G. H.	1786.	C	p.l	Ven. mal. 91
21224 argophyllum Lab.	ridged-stemmed	A [J or 1	my.jl	W	N. S. W.	1812.	C	I.p	Bot. mag. 1309
21225 angustifolius Jac.	white-leaved	M [J or 1	my.jl	W	V. D. L.	1804.	C	s.p	Bot mag. 1563
II. ANGUSTIFMMI.—Shrubby; leaves narrow or small.									
21226 villosus Tittm.	narrow-leaved	■■■ or ■■■	my.jl	Pa.B	C. G. H.	1804.	C	I.p	Jac. sc. 3.370
21227 obtusatus 7%w.	villous	■■■ [J or ■■■	my.jl	W	C. G. H.	1812.	C		
21228 pluriflorus G. Don	obtuse-leaved	■■■ [J or ■■■	my.jl	W	C. G. H.	1793.	C	I.p	Bot. mag. 2288
fruticulosus B. M.	many-flowered	■■■ [J or ■■■	my.jl	W	C. G. H.	1759.	C	p.l	
21229 fruticulosus L.	rather shrubby	◀-1 [J qr	mr.jl	B	C. G. H.	1759.	S	p.l	Bot. mag. 2718
21230 filifolius fins.	thread-leaved	◀-1 [J or 3	mr.jl	W	C. G. H.	1812.	C	I.p	Ven. mal. 82
21231 aculeatus Lab.	wimpy-leaved	◀-1 [J or 2	mr.jl	W	N. Holl.	1818.	C	I.p	Bot. cab. 830
21232 exasperatus ZA:	roughened	tt- [J or 3	mr.jl	W	C. G. H.	1823.	C	I.p	
21233 carolinianus W.	Carolina	tti [J lei	8 au.s	p	Carolina	—	C	co	
III. PAUCIFLURI.—Herbaceous; stem one or few-flowered.									
21234 alpinus L.	alpine	■■■ my au	P	Alp. Eur.	1858.	D	p.l	Bot. mag. 199	
2 fibre albo	white-flowered	£ iny.au	W	Europe	1827.	D	p.l		
21235 pulchellus IV.	pretty	f niy.au	P	Armenia	...	D	co		
21236 caucasicus FT.	Caucasian	1 jl.au	P	Caucasus	1804.	D	co		
21237 lusitanus 26*tf.	Spanish	1 my.jl	B.P	Spain	1826.	D	co		
21238 strictus Ph.	start dwarf	§§.n	V	N. Araer.	1806.	D	co		
21239 biflorus /&?*.	two-flowered	I jl.s	V	Caucasus	1820.	D	co		
21240 pergrinus Ph.	foreign	%jn.au	B	N. Amer.	...	D	co		
IV. LINEARIPULII.—Leaves entire, narrow, sublinear; stem many-flowered.									
21241 Ayssoniiflorus /.	Hyssop-leaved	■■■ ■■■	W	N. Amer.	1683.	D	co	Dod.m.60	
21242 montanus Nut.	mountain	■■■ ■■■		Carolina	—	D	co		
21243 solidaginoides Mx.	Solidago-like	■■■ ■■■		N. Amer.	1699.	D	co	Pluk. al. 79. 2	
21244 tardiflorus Mi.	late-flowering	■■■ ■■■	W	N. Amer.	1820.	D	co		
21245 nemoralis H. K.	grove	■■■ ■■■		N. Amer.	1778.	D	co		
fcdfiflorus Ph.	Li								
21246 rigidus L.	stiff-leaved	■■■ ■■■	P	N. Amer.	1759.	D	co	Pluk. al. 14.7	
21247 paupiflorus Nut.	few-flowered	■■■ ■■■	W	Missouri	...	D	co		
21248 lineariflorus L.	Toad-ilax-lvd	■■■ ■■■	Pa.B	N. Amer.	1699.	D	co		
21249 jraminiiflorus Ph.	Gra>s-leaved	■■■ ■■■	Pa.P	Huds.Jay	...	D	co		
21250 Timiflius TV.	Flax-leaved	■■■ ■■■	V	N. Amer.	1739.	D	co		
21251 pilibus W.	pilose	■■■ ■■■	Pa.B	N. Amer.	1812.	D	co		
21252 foliolous W.	leafy	■■■ ■■■	Pa.B	N. Amer.	1782.	D	co	Di. el. 35.59	
21253 aubulmus Mx.	subulate	■■■ ■■■	Pa.B	N. Amer.	...	I)	co		
21254 tenuiflorus L.	slender-leaved	■■■ ■■■	W	N. Amer.	1725.	D	co	Pluk. al. 78.5	
21255 foilioides H. K.	leafy	■■■ ■■■	W	N. Amer.	1799.	D	co		
21256 dumosus L.	bushy	■■■ ■■■	W	N. Amer.	1734.	D	co	Pluk. al. 78.6	
21277 tricoides L.	Heath-like	■■■ ■■■	W	N. Amer.	1758.	D	co		
21278 multiflorus H. K.	mane-flowered	■■■ ■■■	W	N. Amer.	1732.	D	co	Di. el. 36.40	
21259 ciliatus W.	ciliated	■■■ ■■■	W	N. Amer.	...	D	co		
21260 capitescens Ph.	cancscnt	■■■ ■■■	V	N. Amer.	1812.	D	co		
biennius Nut.	marshy	■■■ ■■■	B	N. Amer.	1784.	D	m.s		
21261 paludibus H. K.	scattered-flwd	■■■ ■■■	Ba.P	N. Amer.	1798.	D	co		
21262 sparsiflorus Mx.	Coris-leaved	■■■ ■■■	B	N. Amer.	...	D	co		
21263 coridiflorus Mx.	shoaly	■■■ ■■■	B	N. Amer.	...	D	co		
21264 surculibus Mx.									
V. LATIFOLH.—Leaves entire, broad, ovate, oblong, or lanceolate; stem many-flowered.									
21265 squarrosus Walt.	squarrose	^ A el		N. Amer.	1801.	D	co		
21266 oblongiflorus Nut.	oblong-leaved	3l A or		N. Amer.	1797.	D	co		
21267 argenteus Mx.	bilver-leaved	A el		N. Amer.	1801.	D	co		
21268 concolor L.	self-colored	■■■ ■■■		N. Amer.	1759.	D	co		
21269 reticulatus Ph.	retctd-leaved	■■■ ■■■		N. Amer.	1812.	D	co		
21270 trinervis Rox.	three-nerved	■■■ ■■■		Nepal	1818.	D	co		
21271 filibus W.	white	■■■ ■■■		N. Amer.	?	D	co		
21272 corniflorus W.	Cornus-leaved	■■■ ■■■		N. Amer.	1811.	D	co		
21273 hhmilis Ph.	humble	■■■ ■■■		N. Amer.	1699.	D	co	W.h.b.67	
21274 divaricatus L.	divaricate	■■■ ■■■		N. Amer.	1800.	D	co		
21275 amygdalinus Lam.	Wmowd-leaved	■■■ ■■■		N. Amer.	1759.	D	co		
umbellatus H. K.									
21276 jaliofibus H. K.	Willow-leaved	■■■ ■■■		N. Amer.	1760.	D	co	Rob. ic. 207	
virgatus Nees									
21277 praecultus Pair.	very tall	■■■ ■■■		N. Amer.	1800.	D	co		
21278 auffivus // K.	summer	■■■ ■■■		N. Amer.	1776.	D	co		
21279 tortiflorus Mx.	twisted-leaved	■■■ ■■■		N. Amer.	...	D	co		
21280 rambus G. Don	smal-bnnchy	■■■ ■■■		N. Amer.	?1816.	D	co		
alpnus 3 rambus Hort.									
21281 Novas A'ngliffi L.	New England	■■■ ■■■		N. Amer.	1710.	D	co	Bot. reg. 183	
2 riiber	red-flowered	■■■ ■■■		N. Amer.	1812.	D	co	Bot.reg.183. inf.	
21282 sessiliflorus Hort.	sessile-flowered	■■■ ■■■		N. Amer.	1700.	D	co		
21283 spurius W.	spurious	■■■ ■■■		N. Amer.	1789	D	co		
21284 cyancus Nut.	bright-blue	■■■ ■■■		N. Amer.	1789.	D	co		
21285 rubricaulis Lam.	red-stemmed	■■■ ■■■		N. Amer.	1815.	D	co	Hof. ph. 1. B. 1	
glaticus Nees, cyancus Hafin.									
21286 grandiflorus L.	great-flowered	A or 8 o.n	B	N. Amer.	1720.	D	co	Bot. reg. 273	
21287 pulcherrimus hod.	fairest	A el 1 s.o	B	N. Amer.	?1810.	D	co	Bot cab. 6	
81288 phlogiflorus TV.	Phlox-leaved	A or 1jl.o	V	N. Amer.	1797.	D	co		
21289 patens H. K.	spreading hairy	A or 1j.s.n	P	N. Amer.	1773.	D	co	Sw.fl. gar. 234	
21290 punctatus Kit.	dotted	A or 3 au.s	V	Hungary	1815.	D	co	W. & K. S. 109	
21291 acm L.	acrid	A or 2 au.s	U	S. Eun>ic	1731.	D	co	Pluk. al. 271.3	

2J292 <i>canus</i> J.Bf.	<i>hoary-leaved</i>				Hungary 1816.	D co	W. & K. I. 3D	
21293 <i>pannonicus</i> Jac.	Fannonian				Hungary 1815.	D co	Jac. vin. I. 8	
21294 <i>ilmellus</i> L.	<i>Amellus</i>				Italy 1890.	I co	Bot. reg. 340	
2 angustifolius	narrow-leaved				S. Europe 1596.	D co		
21295 <i>ibericus</i> Bieb.	Iberian				Iberia ...	D co		
21296 <i>sallgnus</i> JV.	<i>S&Vovr-leaved</i>				Germany 1815.	D co		
<i>hungaricus</i> Poir.								
21297 Schreberrh Nees	Schreber's				N. Amer. ? ...	D co		
21296 <i>longifolius</i> Lam.	long-leaved				N. Amer. 1798.	D co	M. h. 7.22.26	
VI. SEHRATIFOLH. — Leaves lanceolate and ovate, lower ones serrated.								
21299 <i>araplexicaulis</i> JV.	stem-clasping		A or	3 s.n	B	N. Amer. ...	D co	
pennsylvanicus Poir.								
21300 <i>præmannthoides</i> JV.	Prenanthes-like				N. Amer. 1821.	D co		
21301 <i>adulterinus</i> JV. en.	false				N. Amer. ...	D co		
21302 <i>lavigatus</i> JV.	smooth-rfmrf				N. Amer. 1794.	D co		
<i>glaficus</i> Nees								
21303 <i>versicolor</i> JV.	various-colored				N. Amer. 1790.	D co		
21304 <i>mutabilis</i> L.	changeable				N. Amer. 1710.	D co	Her. lug. 67	
keVis L.	smooth				N. Amer. 1758.	D co		
<i>luculifl H. K.</i>								
21306 <i>concinrus</i> JV.en.	neat				N. Amer. 1600.	D co		
21307 <i>abbreviates</i> Nees	shortened				N. Amer. ...	D co		
21308 <i>wyrtifolius</i> JV.	Myrtle-leaved				... 1812.	D co		
21309 <i>puniceus</i> L.	<i>red-stalkcd</i>				N. Amer. 1710.	D co	Her. lug. 651	
<i>amoënius</i> Lam.								
21310 <i>firmus</i> Nees	firm				N. Amer. 1816.	D co		
21311 <i>rivalaris</i> Cos.	river				N. Amer. 182a	D co		
21312 <i>hispidus</i> Lam.	<i>rough-stalked</i>				China 1804.	D co	Kaem. ic. 29	
21313 <i>floribindua</i> JV.	bundle-flwd				N. Amer. —	D co		
21314 <i>Nbvi Belgii</i> L.	New York				N. Amer. 1710.	D co		
21315 <i>Aellidiflbris JV.en.</i>	Daisy-flowered				N. Amer. —	D co		
21316 <i>spectabilis</i> H. K.	showy				N. Amer. 1777.	D co		
21317 <i>serotinus</i> W.	<i>axE-flowering</i>				N. Amer. —	D co		
21318 <i>tardiflbris L.</i>	late-flowered				N. Amer. 1775.	D co		
21319 <i>marginatus</i> Kth.	marginata				N. Gran. 1827.	S s.l.		
21320 <i>amellidids A*</i> .	Amellus-like				Podolia 1824.	D s.l		
21321 <i>pyrenense</i> Detf.	Pyrenean				Pyrenees —	D co		
lfc2 blandus PA.	charming				N. Amer. 1800.	D co		
*21323 <i>acuminatus</i> JW.	acuminate				N. Amer. 1806.	D co	Bot cab. 959	
S2324 <i>conyzomes</i> FT.	Conyza-like				N. Amer. 1778.	D co	Bot mag. 2707	
<i>marilandicus</i> Afx.								
21325 <i>inuloides</i> D. Don	Inula-like	^{at} A or	au	R	Nepal	—	D co	
21326 <i>nudiflbris</i> tfirf.	naked-flowered	^{at} A or	au	P	N. Amer. ...	D co		
21327 <i>tatricus</i> L.	Tartarian	^{at} A or	au	B	Tartary 1818.	D co		
21328 <i>plantaginiflbris</i> Nut.	Plantain-lvd	^{at} A or	aus.s	W	N. Amer. ...	D co		
21329 <i>ridula</i> H. K.	<i>ra&p-leaved</i>	^{at} A or	s.n	W	N. Amer. 1785.	D co		
21330 <i>Tradescanti</i> L.	Tradescant's	^{atc} A or	jls	W	N. Amer. 1633.	D co	M.h.7.21.42	
21331 <i>recurvatus</i> JV.	recurved	^{at} A or	3 aus.s	W	N. Amer. 1800.	D co		
21332 <i>eminens</i> JV.	eminent	^{at} A or	2 s.n	Pa.B	N. Amer. ...	D co		
pubescens Nees				L				
21333 <i>xxus</i> JV.			A or	2 s.n	W	N. Amer. ...	D co	
leucanthemus Detf.								
21334 <i>simplex</i> JV. en.	simple-stalked		A or	3 au.o	W.P	N. Amer. ...	D co	
<i>virginicus</i> Nees								
21335 <i>bicolor</i> Nees	two-colored		A or	3 aus.s	W.Y	N. Amer. 1759.	D co	Pluk.aLU4.8
<i>Solidugo bicolor</i> L.								
21336 <i>artemviflbris</i> Foir.	Wormwood-flwd	^{ai} A or	3 auo	W	N. Amer. ...	D co		
<fracuncoloides W. not Lam.								
21337 <i>polyphllus</i> JV. en.	many-leaved		A or	3 au.o	W	N. Amer. ...	D co	
<i>iniflbris</i> Nees								
21338 <i>laxiflbris</i> Lie.	loose-flowered	^{at} A or	4 s.o		N. Amer. ...	D co		
213% <i>junceus</i> H. K.	Rush	^{at} A or	4 s.o	F	N. Amer. 1758.	D co		
21340 <i>lanceolatus</i> Lam.	lanceolate	^{at} A or	4 au.n	W.p	N. Amer. 1811.	D co		
21341 <i>rfracuncoloides</i> Lam.	Tarragon-like	^{at} A or	3 s.n	W	Turria 1811.	D co		
21342 <i>fragilis</i> JV.	brittle	^{at} A or	2 s	F	N. Amer. 1800.	D co		
<i>cancus</i> Nees								
21343 <i>miser</i> L.	mesigre-flwd	ji	un	3 i.o	W	N. Amer. 1759.	D co	
parvifl&rhus Nees								
21344 <i>divergens</i> H. K.	spreading downy		A "	3 s.o	W	N. Amer. 1758.	D co	
21345 <i>diffusus</i> H.K.			A or	2 s.o	W	N. Amer. 1777.	D co	
<coparius Nees								
2134G <i>pendulus</i> H. K.	pendulous	^{3c} A or	2 s.o	W	N. Amer. 1758.	D co		
21347 <i>altalicus</i> JV. en.	Altaiian	^{atc} A o*	my.au	B	Siberia 1804.	D co		
<i>pbxnilus</i> Fis.								
21348 <i>surculbus</i> Mx.								
21349 <i>gracilis</i> Nut.	slender		1 aiLS	P	N. Amer. 1800.	D co		
21350 <i>tenellus</i> Thmt.	delicate		1 JLau	O	N. Amer. ? ...	D co		
21351 <i>7Yip6Liun</i> L.								
21353 <i>sibiricus</i> L.	Tripoly		S au.s	B	C. 6. H. 1769.	C p.l		
21353 <i>inclusus</i> Ft*.	Siberian		2 jl.o	B	Britain scash.	D co		
Grindelia siberica Spr.	cut-leaved		Ifjn.au	B	Siberia 1768.	D co		
					Siberia 1818.	D co		
81354 <i>desertrum</i> Fis.	desert			B	Siberia 1830.	D co		
81355 <i>Regans</i> JV.	elegant			B 1790.	D co		
81356 <i>pklens</i> JV. en.				V	N. Amer. ...	D co		
21357 <i>rigidulus</i> Nees	rigidish			B	N. Amer. 1816.	D co		
21358 <i>luxurians</i> Spr.	luxuriant			B	N. Amer. 1816.	D co		
<i>expansus</i> Nees								
21359 <i>yimneua</i> W.	twiggy			B	N. Amer. 1800.	D GO		
p&tulus Lam. fltrctus Poir.								
21360 <i>pneox</i> JV. en.	early-flowering			V	N. Amer. 1800.	D co		
VII. CORDIFOLII. — Leaves serrated or toothed, cordate.								
21361 <i>undulatus</i> L.	mi'e-leaved	^{at} A or	3 au.o	P.B	N. Amcr. 1699.	D co	Her. iiar. 9%	
paniculatus H. K.	panicked	⁵ A el	4 s.o	B	N. Amer. 1640.	D co	Cbr.ca.6j	
*anguineus JV.	bloody	^{ft} A or	3 s.o	B	N. Amer. ...	D co		

21361 cordiflous <i>L.</i>	heart-leaved	■	■	P.B	N. Amer. 1759.	D co
81365 aagittaeblus <i>IV.</i>	sagittate-leaved	■	■	B	N. Amer. 1700.	D co
21366 corymbbus <i>L.</i>	corymbed	■	■	W	N. Amer. 1765.	D co
21367 macrophyllus <i>L.</i>	long-leaved	■	■	W	N. Amer. 1739.	D co
21368 heterophyllus <i>P.en.</i>	various-leaved	■	■	Pa.r	N. Amer. 1811.	D co
21369 alwartensis <i>hod.</i>	Alwart	■	■	R	Caucasus 1807.	D co
21370 graveolens <i>Nut.</i>	heavy-smelling	k A or 2	.	Arkansa	1826	Bot mag. 8321

VIII. DUNJIUS.—Doubtful to which of the sections it belongs.

21370 graveolens *Nut.* heavy-smelling k A or 2 . Arkansa 1826 D co

2338. CALLISTETMA Cos. CHINA ASTER. (<i>Kallistos</i> , prettiest <i>sterne</i> , stamen : fl.)		<i>Goppi</i> , <i>Card. Yerpp.</i> SL	
21371 hortense <i>Cos.</i>	garden	O el	B China 1731. S co
<i>A'ster chinensis L.</i>			Di. el. 34.38
2 rubrum	red	O el	R China 1731. S co
3 album	white	O el	1 st jls. W China 1731. S co
4 variegatum	variegated	O el	Va China 1731. ■ co
5 multiplex	double	O el	14 jls. Va China 1731. S co
6 brachyptithum	short-flowered	O el	IJJLs Va China 1731. S co
21372 Sndicum <i>O. Dun</i>	Indian	O el	1 jls. B E. Indies 1820. S co
<i>A'ster indicus L.</i>			

2339. SOLIDANO L. GOLDEN ROD. (*Solido*, to unite; vulnerary quality of plant) *Comp. Card. Vern.* 68.—

I. UNILATERALYES.—Racemes onesided.

i. TRINE'RVBS.—Leaves three-nerved.		ii. VENbas—Leaves veiny,	
21373 sp&ria Worst.	spurious	■ or 6 n	Y? St.Helenal 772. C p.1
Conyza rugosa <i>H. K.</i>	Canadian	■	Y N. Amer. 1648. D co Sch. han. 3.24G
81374 canadensis <i>L.</i>	fragrant	■	Y N. Amer. ... D co
21375 procris <i>H. K.</i>	Call	■	Y N. Amer. 1758. D co
21377 serotina <i>//. K.</i>	<i>Latc. II/uering</i>	■	Y N. Amer. 1758. D co
21378 amsinca <i>Lo. C.</i>	gigantic	■	Y N. Amer. 1815. D co
21379 granted <i>H. K.</i>	ciliated	■	Y N. Amer. 1758. D co
21380 ciñaris <i>IV.</i>	reflex-l'taved	■	Y N. Amer. 1811. D co
21381 reñexa <i>H. K.</i>	lateral-flowered	£ ^ or 3 aus	Y N. Amer. 1758. D co
21382 lateriflora <i>L.</i>			N. Amer. 1755. D co
21383 aspera <i>H. K.</i>	tough-leaved	■	N. Amer. 1732. D co Di.eL305.392
21384 asperata <i>Ph.</i>	roughed	■	N. Amer. 1808. D co
21385 altissima <i>L.</i>	tallobt	■	N. Amer. 1686. D co Mart.cen. 14
21386 recurvata <i>Ph.</i>	recurvate	■	N. Amer. 1686. D co
3 virginixa PA.	Virginian	■	K. Amer. 1686. D co
21387 pyramidata <i>Ph.</i>	pyramidal	■	N. Amer. 1790. D co
21387 rugosa <i>W.</i>	wrinkle-leaved	■	N. Amer. 1732. D co Di. el. 308.396
21388 villosa <i>Ph.</i>	villous	■	N. Amer. 1732. D co
21389 scabra <i>W.</i>	scabrous	■	N. Amer. 1811. D co
21390 nemoralis <i>H. JL</i>	grove	■	N. Amer. 1769. D co
21391 patula <i>W.</i>	spreading	■	N. Amur. 1H05. D co
21392 ulmifolia <i>W.</i>	flm.leaved	■	N. Amer. 1805. D co
21393 arguta <i>H. K.</i>	sharp-notched	■	N. Amer. 1758. D co
21394 uncea <i>H. K.</i>	Rush-stalked	■	N. Amer. 1769. D co
21395 virga <i>Mx.</i>	twiggy	■	N. Amer. 1800. D GO
21396 pulverulenta <i>Nut.</i>	powdered	■	K. Amer. ... D co
arg^ntea <i>Lo. C.</i>	Folium-leaved	■	N. Amer. 1826. D co
21397 nothoflida <i>Lo. C.</i>	cUiytic leaved	■	N. Amer. 1759. D co
21398 elliptica <i>H. X.</i>	recurved	■	N. Amer. ... D co
21399 recurvata <i>W. en.</i>			
rctr6hra <i>Mx.</i>	evergreen	■	N. Amer. 1699. D co Cor. ca. 169
21400 sempervirens <i>L.</i>	sweet^melling	■	N. Amer. 1699. D co Fluk.aL116.6
21401 odbra <i>H. K.</i>			

II. ERE'CTJB.—Racemes erect.

i. SBRBA>Tf.—Lower leaves serrated or toothed.		II. ERE'CTJB.—Racemes erect.	
21402 petiolaris <i>H. K.</i>	petaoled	■	N. Amer. 1758. D co
21403 stricta <i>H. K.</i>	strict	■	N. Amer. 1758. D co
21404 virgata <i>Mx.</i>	twiggy	■	N. Amer. ... D co
21405 squarrdosa <i>Mhl.</i>	rough	■	N. Amer. ... D co
21406 cai^ia <i>H. K.</i>	U	■	N. Amer. 1732. T) co DLeLS07.395
21407 livida <i>IV. en.</i>	hairy	■	N. Amer. ... D co
21408 hispida <i>W.</i>	hairy	■	N. Amer. 1800. D co
21409 hirta <i>W. en.</i>	Gronwell-lvd	■	X. Amer. ... D co
21410 /ithospermiflba <i>TV.</i>	smooth-leaved	■	N. Amer. 1811. D co
21411 lavigata <i>//. K.</i>	Mexican	■	N. Amer. 1699. D co
21412 mexicana <i>L.</i>	twiggy	■	N. Amer. 1K83. I) co Dod.m.4.219
21413 viimlnca <i>//. K.</i>	showy	■	N. Amer. 1759. D co
21414 specibi <i>Nut.</i>	long-leaved	■	N. Amer. 1817. D co
21415 macrophylla <i>Ph.</i>	New York	■	N. Amer. ... D co
21416 noveboracensis <i>Ph.</i>	glomerated	■	N. Amer. 1820. D co
31417 glomerata <i>Mx.</i>	crook-stalked	■	N. Amer. 1725. D co Her. par. 244
21418 flexicaulis <i>L.</i>	tall hairy	■	N. Amer. 1811. D co Pluk.al.235.4
21419 elata <i>PA.</i>	broad-leaved	■	N. Amer. 1725. D co
21420 latifolia <i>L.</i>	ambiguous	■	... 1759. D co
21421 ambigua <i>H. K.</i>	rigid-leaved	■	N. Amer. 1710. D co Her. par. 243
21422 rigida <i>L.</i>	axillary	■	N. Amer. 1811. D co
21423 axillaris <i>Ph.</i>	Golden Rod, com.	■	Britain woods. I) co Eng.bot.301
21421 Virgafrura <i>L.</i>	American	■	N. Amer. 1800. D co
2 ameriehna	alpine	■	Switzcr. 1819. D co
3 alpina <i>Seal.</i>	sand	■	& Europe 1816. D co
21425 arenaria <i>Pair.</i>	rock	■	Hungary 1816. D co
montana <i>Pair.</i>	naked-flowered	■	S. Europe 1820. D co
21426 alpestris <i>Kit.</i>	puberulous	■	N. Amer. ... I> co
21427 nudiflora <i>Viv.</i>	Welsh	■	Wales ... 5 co
21428 puberula <i>Nut.</i>	many-rayed	■	Nepal D co
21429 cambrica <i>Hud.</i>			Labrador 1776. D co
21430 nepalensis <i>D. Don</i>			
21431 multiradiata <i>A. K.</i>			

Di.eL306.

ORDER II.

SYNGENESIA SUPER FLU A.

21432 simplex <i>Kth.</i>	alpe	1	-	Y	Mexico	1896.	D co	
21433 mirfutal.,	minute	ijl.au	Y	Pyrenees	1772.	D co	Bet cab.	189
21434 decurrent		jl.o	Y	ChiilM	1823.	D co		
£1435 littoralis <i>Savi</i>	shore	£ jl.au	Y	Etruria	1827.	D co		
21436 humilis <i>Ph.</i>	humble	1 jl.au	Y	N. Amer.	1811.	D co		
	ii.							—Leaves all entire.
21437 pauciflosculosa <i>Mx.</i>	few-floreted	2 au.o	Y	N. Amer.	1811.	D co		
21438 angustifolia <i>Nut.</i>	narrow-leaved	3 s.o	Y	N. Amer.	...	D co		
21439 ericta <i>PA.</i>	upright	3 au.o	Y	N. Amer.	...	D co		

*2340. CINERARIA L. CINERARIA. (*Oneres*, ashes; down covering surfaces of leaves.) *Comp. Jacobke.* 51.

I. GEIFOLIA.—Flowers radiate.

								Leaves undivided s peduncles many-flowered ; shrubby.
21440 lactea <i>W. en.</i>	milk-co/orw/	tt i Jor	W		1816.	D	Lp	
21441 Petasites <i>B. M.</i>	Butterbur- <i>flM</i> 2	*i Al or	Y	Mexico	1812.	D	co	Bot. mag. 1536
21442 #cifolia£.	Geum-leavcd	tt i Jor	ap.au	C. G. H.	1710.	D	p.l	Com. h. 2.73
21443 canescens IV. <i>en.</i>	canescent	<-i or	ap.au	C. G. H.	1790.	D	co	Bot. mag. 1990
21444 aurita <i>Herit.</i>	eared	tt 1 Jor.	I	P	Madeira	1777.	C	l.p
21445 cruenta <i>Herit.</i>	bloody-leaved	tt 1 or	2 f.my	Canaries	1777.	H	p.l	Bot. mag. 406
21446 uopulli'Mia Lam.	Poplar-leaved	tt 1 or	2 jn.s	Canaries	1780.	C	p.l	Yen. mal. 100
21447 W t a W.	lobed	tt 1 or	3 jn.au	C. G. H.	1774.	C	p.l	
21448 pulchella <i>Suit.</i>		tt \ Jor	Y	Canaries	1818.	C	Lp	
21449 hybrida <i>W.en.</i>	hybrid	tt 1 or	Y	Y	Mexico	1824.	C	p.l
21450 praeNwx <i>Cav.</i>	early	tt 1 pr	2 f.my	W?	c. & H.	1885.	C	
31451 Mba I* <i>C.</i>	white	a. 1 pr	IJ f.ap	Y	Grenada	18*5.	C	
21452 aincnacna <i>L.</i>	American	CD or	f.ap	Y	Jamaica	1823.	C	
21453 incana <i>Swx.</i>	hoary	tt. 1 pr	24inaii	Y	Jamaica	1804.	C	Lp
21454 discolor <i>Swx.</i>	two-colored-tori*	O or	4 jnau	W	Jamaica	1822.	C	
21455 glabrkU Sun.	smooth	tt. O or	1} jn.au	Y	W. Indies	...	C	Lp
21456 lhcida Swx	shining	tt. CD or	2 jn.au	Y	C. G. H.	1816.	C	
21457 cacalioides <i>Thun.</i>	Cacalia-like	n. 1 or	2 jn.au	Y	W. Indies	...	C	
21458 rodifolia <i>ATM.</i>	Willow-leaved	tt. 1 or	2 jn.au	Y	Mexico	1827.	C	p.l
21459 angustifolia <i>Kth.</i>	narrow-leaved	tt. y or	lj jn.au	Y	Mexico	1825.	C	

ii. Dms.EiA.LU3 —Leaves divided or pinnatifid ; peduncles many-flowered

21460 bicolor L.	two-colored	tt-uJor	2 jl.au	Y	Austria	...	C	co
ambigua <i>Biv.</i> nebrodensis <i>Cav.</i>								
21461 AitonMna Spr.	Alton's	tt	• or	1 jn.au	Y			C
21462 maritima L.	Ben Ragwort	«. Jor	2 jls	Y	S. Europe	1633.	C	Lp

21463 lanata <i>Herit.</i>	woolly	tt.	1 Jor	33 myss	IP	Canaries	1780.	C p.l
----------------------------	--------	-----	-------	---------	----	----------	-------	-------

III. TUSSILAGIN&K.—Flowers radiate ; peduncles many-flowered i stems herbaceous.

i. CORDIFOLI*—leaves broad cordate.

21464 malvifolia <i>Herit.</i>	Mallow-leaved	2 au	Y	Azores	1777.	R	p.l
21465 elatior <i>Bouchi</i>	taller	5 jl.au	W		D	l.p	
21466 gigant'a 5m.	gigantic	1jjl.au	W	Cape Horn	1861.	I	Lp
21467 sibirica <i>L.</i>	Siberian	4 jn.au	Y	Siberia	1784.	D	co
21468 speciosa ScAr.	showy	6 jl.au	Y	Siberia	1815.	I	co
21469 lavigata i/orf.	smooth-team*	1 jl.au	Y	Siberia	1819.	D	l.p

ii. INTBGRIFOLIX.—Leaves undivided, entire.

21470 suddtica Koch	Sudetic	1 jl.au		Switzerl.	1819.	D	rro	Bot mag. 2263
21471 rivulXris Kit.	rivulet	1 jl.su		Hungary	18 in.	D	p.l	
21472 crasmifolia Kit.	thick-leaved	1 jl.au		Carinthia	1827.	D	rm	
ovirénensis Koch								
21473 crispa L.	curled	2 jl.au		Switzerl.	1818.	D	r.m	Jac.au. 2.178
21474 alpestris Hoppe	alpine	† my.jn		Switzerl.	1819.	D	p.l	
21475 alpina Hoppe	alpine	† jn.au		Switzerl.	1819.	D	co	Jac. au. 2.177
21476 aurant'aca Hoppe	orange	† my.jl		Switzerl.	1819.	D		Sw. fl. gar. ic.
21477 spatulaefolia Gm.	spatulate-lvd	† my.jl		Germany	1820.	D		
integrifolia Roth, longifolia Sturm								
21478 longifolia Jac.	long-icved	2 jn.au	Y	Austria	1792.	D	co	Jac. au. 2.181
21479 papposa Rchb.	pappose	1jn.au	Y	Galicia	1821.	D	p.l	
21480 campiflora Jtex.	wild	£ my.jn	Y	Europe	...	D	co	Jac. au. 180
21481 integrifolia E. B.	entire-leaved	1 my.jn	Y	England ch.p.A.	D	co		Eng. bot. 152
21482 palustris L.	marsh	3 jn.jl	Y	England mar.	D	m.s		Eng. bot. 151
21483 aurea L.	golden	2 jn.jl	Y	Siliena	1827.	D	r.m	
21484 parviflora Bieb.	small-flowered	2 jl.au	Y	Caucasus	1820.	D	l.p	
21485 racemosa Bieb.	racemose	1 jl.all	Y	Caucasus	1820.	D	p.l	

iii. DIVI^A.—Leaves divided.

21486 canadensis L.	Canadian	^ A or	2 jn.au	Y	Canada	1739.	D	co
---------------------	----------	--------	---------	---	--------	-------	---	----

III. PAUCIFLORX.—Flowers rayed; peduncles one-flowered.

21487 caucasica Bieb.	i. INDIVISÉ—Leaves undivided.							
21488 huraifiti Herit.	Caucasian	£ A or	U jn.au	Y	Caucasus	181G.	D	l.p
	trailing	£ Al or	\ jl.au	Y	C. G. H.	1704.	R	p.l

21489 viscosa Jlerit.	ii. LYRATJEFOLIA.—Leaves lyrate-pinnatifid.							
	clammy	jg OJ or	2 jn.au	Y	C. G. H.	1774.	C	p.l

2341. AGATHA TA Cas.	AGATHAA. {Agathos, excellent; beautiful flowers.)							
21490 cœlestis Cas.	heavenly blue	tt. i J pr	U fs	B	C. G. H.	1759.	C	p.l
Cineraria amelloides L.								BoLmag. 249

21491 /inifolia G.Don	Flax-kavcd	& J pr	S f.i.	B	C. G. H.	1800.	C	p.l
Cineraria /inifolia L.								

2342. SENECPOLLIS Gae.	SENBICILLIS.	(Perhaps a diminutive of Senecio.)						
21492 glauca Gae.	glaucous	A or	6 jk.jau	Y	Siberia	1790.	D e o	Gm. sl2.74*
Cineraria glauca L.								

21493 purpurata Gae.	purple	2 my.jl	P	C. G. H.	1816.	D	p.l	
Cineraria purpurata L.								

qt?!!*^3- CALOTIS R. Br.	CALOTIS. {Kalos, beauty, ous, an ear; paloa; of pappus.)							
H94 cun^IMiaifclfr.	wedge-leaved	£ i Al p	1 my.jn	B	N. Holt	1819.	D e o	Botreg. 504

2344. KAULFU'SS/i* Nees. KAULFUSSIA. (G. Fred. Kaunas, M.D., prof. bot, Halle.) *Comp. Jacobhe.* 1.
81495 amcelloldes Nees Amellus-like O pr 1 i^U B C. G. H. 1819. S co Bot.reg.4H0

2345. PNULA £. INULA. (Corrupted from *Helenium*.) *Comp. Card. Fern.* 41.—

I. SUFFRUTICOSUB.—*Stems subshubby.*

21496 tuberbsa Lam.	tuberous-rooted & A or	lftjl.au	Y	S. Europe	1640.	D eo	M.L.7.19,20
frgeron tuberosus	Cnfhmum-like *	A or 4	... Y	Greece	1800.	C co	
21497 crithmoules L.	Crithmum-lvd £ A or	2 au.s	Y	England	sal.ni.	D co	Eng. bot 68
21498 crithmifolia A/tf.	• • or 1	... Y	VeraCruz	1733.	C l.p	R.Hou.8.19	

II. SUBUNIFLORS.—*Seems generally one-flowered; leaves broad, oblong, or lanceolate; herbaceous.*

21501 glamlulbsa W.	great-flowered glandulose	A or 2	jl.au	Y	Caucasus	1810.	D r.m
21502 hirta Z.	hairy	A or 1	jn.s	Y	Georgia	1804.	D co Bot mag. 1907
21503 montana L.	mountain	A or	ljjl.au	Y	Austria	1759.	D co Jac. au. 4.358
21504 calyclna 5pr.	montana Biv.	A or lft	jlau	Y	S. Europe	1759.	D co Gar. aix 10
21505 provinciMis X.	Provence	A or 1	jl.au	Y	Sicily	1827.	D r.m

III. SUBMULTIKLOILE.—*Stems generally many-flowered; branches and peduncle* one-flowered; leaves broad oblong, or lanceolate; herbaceous.*

21506 odhra L.	fragrant	oblong, or lanceolate; herbaceous	&	Europe	1821.	D co	M.h.&7.21:6
21507 mariiana L.	Maryland			N. Amer.	1742.	D co	Mil. Ic 1.57.1
21508 quadridentata Lag.	four-tooth-Jtfurf			Spain	1820.	D co	
21509 salicina /.	Willow-fc<ii>*<*			N. Europe	648.	D co	Fl. dan. 786
21510 Zcibonum Jac.	Bubonium			Austria	1801.	D co	Jac. au. 5. ap. 19
21511 Vaillntv Vil.	Vaillant'a			France	1739.	D co	
21512 Pulicaria L.	Flea-wort			England	moi. h.	S co	Eng. bot 1196
Pulicaria vulgaris Gae.							
21513 arábica Lk.	Arabian	O un	lftau.s	Y	Arabia	1823.	S co
Pulicaria arábica Lk.							Pluk. al. 149.4
21514 undulata L.	wave-leaved	O un	1 jl.o	Y	Egypt	1739.	■ co
21515 indica L.	Indian	un	U jl.o	Y	E. Indies	1739.	■ CO
21516 paludosa Lk.,	marsh	O	* jlau	Y	Spain	1816.	■ CO

IV. CORYMBOUB.—*Flowers corymbose s leaves broad, oblong, or lanceolate; herbaceous.*

21517 britannica L.	Britannic			Germany	1759.	D co	Fl. dan. 413
21518 dysenUrca L.	dysenCbr			England	wat.pl.	D co	Eng. bot. 1115
Pulicaria dysenterica Lk.							
21519 mCuIsATjiA.	soft					D co	
21520 conyzoides Desf.	Conyza-like			Crete	1810.	D co	
21521 suavolens Jac.	sweet-scented			S. Europe	1758.	D co	Jac. vin. 3. 51
21522 O'culus Christ* L.	Christ's eye			Austria	1759.	D co	Jac. au. a 1223
21523 campestris Bes.	field			Podolia	182J.	O co	
O'culus Christ* Bich. Bich. he'eno"dc Dec.							
21524 verbascifolia Bich.	Mullcm-leaved			Caucasus	?1819.	D co	
/hapsoldes Spr.							
21525 grossypina Mx.	cottony			N. Amer.	1823.	D p.1	
21526 villosa L.	vUlous			N. Amer.	1811.	S p.1	
Pulicaria villosa Lk.							
21527 squarrtsa L.	squarrose			Italy	1763.	D co	Pluk.ai.16.1
21528 viscosa H. K.	clammy			S. Europe	1596.	C p.1	Jac. vin. 2. 2165
f rgeron viscosus Jac.							
21529 germanica L.	German			Germany	1759.	D co	Jac. au. 2.134
21530 hybrida Baum.	hybrid			Podolia	1818.	D co	
21531 blfrons L.	two-fronted			S. Europe	1713.	D co	Her. par. 187

V.* ANGUSTIF5LI£.—Leaves narrow.

21532 ensifolia Z.	sword-leaved			Austria	1793.	D co	Jac. au. 2.162
21533 graminifolia Mx.	Grass-leaved			N. Amer.	...	D p.1	
21534 fcB'tida L.	fetid			Malta	1688.	S co	Boc. sic. 26.13
21535 saxatilis Lam.	rock			& Europe	1816.	D co	
frlgeron glutinosus L.							

2346. COKVISA'RTIA Merat CORVISARTIA. (*M. Corvisart*, a man's name.) *Comp. Card. Fern.* 2.

21536 HcMnium Merat	Elecampane	^ A or	4 jl.au	Y	Britain	m. me.	D co
.Tnula HeXhnium L.							Eng. bot 1546

21537 caucasica G. Don Caucasian i A or

m 4 jl.au Y Caucasus 1818. D co

JTnula caucasica Pers.

2347. GRINDE^NL/J1 W. en. GRINDELIA. (*Grindel*, a German botanist) *Comp. Card. Fern.* 8.—9.

21538 glutinosa Dunal	glutinous	*1 Jor	2 year	Y	Mexico	1803.	C Lp
Dön/a glutinosa R. Br.							Botreg. 187
21539 imuloides W. en.	Inula-like	il Ljor	1	Y	Mexico	1815.	C Lp
21540 spatulata Lk.	spatulate	il Ljor	lft	Y	Mexico	1819.	C Lp
21541 squarrtsa Dunal	squarrose	jf lAlor	2	Y	Missouri	1811.	C Lp
Dön/a squarrtsa Ph.							
21542 Laraberti G. Don	Lambert's		2 jl.s	Y	Mexico	1816.	C Lp
21543 angustifolia Kth.	narrow-leaved		1 jl.s	Y	Mexico	1822.	C Lp
21544 Duv.Mii Spr.	Duval's	L or	lftjl.s	Y	Mexico	1820.	C Lp
angustifolia Duval							Botreg. 781
21545 ciÜjita W. en.	ciliated	Q or	lft jl.s	Y	N. Amet.	1821.	S Lp
D6nia ciliata Nut.							Hook. ex. fl. 45

•2348. PODCyLEPIS Lab. PODOLEPIS. (*POUS*, foot, lepis, scale; stalks of fl. covered with.) *Comp. Card. Tern.* 2.

21546 rugata Lab. wrixiUc-scaled £ iAJ or

1 jl.au W N. Holl 1803. C s.p

Lab. n. h. 2.208

21547 acuminata H. K. ih&rp-scaled £ j\] or

| my.au W N. S. W. 1803. C s.p

Bot mag. 956

Sealia jaceofides B. M.

2349. CH^ATANTHBPR A Fl. per. CHATAMTBRA. (*Chaita*, bristle, anthera, anther.) *Comp. Labiat.* 2.-4,

21548 ciliata Fl.per. ciliat^AT C IAJor

2 jl.au ... Chile 1822. D co

21549 chitenai* f^A w. Chile tfSjor

ljl.a u ... Chile 1827. D p.1

Perdicium chil^Anse W.

«, fS° - A'RNIC A L.	ARNICA.	<i>(Arnáhīs, lamb's skin; leaves.)</i>	<i>pomp. Card. Vern.</i> 9. —SB.
21550 montana L.	mountain	1. Ao r 1. jlau Y	Europe 1731. D p.l Bot. mag. 1740
31551 helvetica G. Don	Swiss	1. A or 1. jnjl Y	Switzerl. 1819. D p.m
montana Schl.			
21552 scorioides L.	scorpion-like	■ A or 1. jlau Y	Austria 1710. D p.l Bot cab. 913
21552 concta WW.	cordate	■ A or 1. tjl.au Y	Switzerland 1819. D p.l
21554 lanjera Ten.	wool-bearing	■ A or 1. jn.au Y	Italy 1837. D p.l
21555 DoUpicum Jac	Leopard's bane	■ A or 1. jjlau Y	AustraL 1816. D hp Jac. au. 1.98
21556 Cl&ssii All.	Clusius's	■ A or 1. jlau Y	SwitzerL 1819. D p.l Alped. 17.51.2
21557 cfrsiga ioi.	Corsican	■ A or 1. jlau Y	Corsica 1844. D c.p
21558 glacialis Jac.	ley	■ A or 1. jn.au Y	SwitzerL 1823. D p.l
2351. BELLI DIA'STRUM Cos. BELLID.	daisy, astrum. an affix signif. like.)	<i>Comp. Card. Vern. h</i>	
21559 Micheli Cas.	Micheli's	£ A or 1. jnjl W	Austria 1570. D p.l Bot. mag. 1196
Arnica Bellidistrum Vil.			
2352. GERB&RIA Brm. GERBERIA.	(<i>Gerber</i> , German nat. and trav. in Russia.)	<i>Comp. Card. Vern. l. &</i>	
21560 cramata J.-hell.	crenated	£ O I pr 1. jt.au P	C # G. H. 1822. D p.l Botreg. 85fi
2353. DORO'NICUM L. LEOPARD'S BANE.	(<i>Doronigi</i> , Arabic name.)	<i>Comp. Card. Vern.</i> 7. -8.	
21561 Pardahanches L.	Panther-strangler	△ or 2 my Y	Britain m. pas. D co Eng. bot 630
21562 scorioides W.	scorpion-like	1 ap.jn Y	Germany ... D co
21563 Colfon* Ten.	Columna's	2 ap.JQ Y	Italy 1824. D co
21564 austriacum Jac.	Austrian	1 ap.jn Y	Austria 1816. D co Jac. au. 130
21565 alticum Pall.	Altaic	1 jn.au W	Siberia 1783. D 8.1 Ac.pet. 1779.16
21566 caucasicum Bieb.	Caucasian	1 jn.au Y	Caucasus 1815. D l.p
orientale Adams			
• 567 plantagneum L.	PlantauWtrf	■ A or 2 my Y	S. Europe 157a D co,
*2354. DIPLCCOMA D. Don DIPL.	(<i>Diploos</i> , double, <i>koma</i> , hair; pappus two forms.)	<i>Comp. Card. Vern.</i> 1.	
21568 villbsafl.Don	villous	^ A or 1 my.jl Y	Mexico 1826. D e o Sw.fl.gar. 246
Doronicum villsum Sessi			
„ 2355. PERDI'CIUM Dec.	PERDICUM. (<i>Perdix</i> , a partridge; fond of original plant)	<i>Comp. Lobiat.</i> I.—2.	
21569 Anandria Dec.	Anandria	£ A un f mr ...	Siberia 1759. D eo Gm. sL & 68.1
Tusibilago Anandria L.			
*2356. MUTI'S/i Cav. MUTISIA,	(<i>Celestine Mutis</i> , the discoverer, a S. Amer. tot)	<i>Comp. bOiat.</i> 1.—2.	
21570 arachnoidea Mart.	cobweb-like	L C D or 6 jlau R	Brazil 1823. C p.l Bot. mag. 270a
speciata B. M.			
2357. TETRAGONOTHEPCA Herit. TBTBAO.	(<i>Tetragonos</i> , flaudnmgular, <i>theta</i> , case.)	<i>Comp. BeH&n. 1.</i>	
21571 hCuantholde8i/enA Sunflower-Ufee	^ A or 4 au.o Y	Virginia 1726. D p.l Schk. han. 3.263	
2358. XIMEME*SI4Cav. XIMENESIA.	(<i>Joseph Ximenes</i> , a Spanish apothecary.)	<i>Comp. He lidn. 5.</i>	
21572 ence/ides Cav.	Encelia-like	1 lor 3 jlau Y	Mexico 1795. S l.p Cav. ic. 2.178
21573 cordata Kth.	cordate	1. jln 3 jl.n Y	Mexico 1826. ■ IP
21574 Cavamleui Spr.	Cavanilles's	2 JLi Y	Mexico 1820. ■ co Cav. ic. 3.268
Coreopsis heterophylla Cav.			
fetid	O or 2 i jl.s	■ Mexico 1824. S co Cav. ic. 1.77	
SlmsiafcifbWaPers.			
21576 heterophylla Kth.	variable-leaved	O or 2 jLau Y	Mexico 1827. S co
2359. HELPXIUM L. HELENIUM.	(Named after the celebrated Helen.)	<i>Comp. Hethanthee.</i> 7. -8.	
autumnale L.	autumnal	■ A or 3 au.o Y	N. Amer. 1729. D p.l Schk. han. 4250
pubescens H. K.	pubescent	3 au.s Y	N. Amer. 1776. B p.l
plumum IV.	dwarf	1 au.s Y	N. Araer. 21818. D p.l
canahculatum Lam.	channelled	3 au.s Y	N. Amer. 1800. D p.l
quadridentatum Lab.	four-toothed	2 my.o Y	Louisiana 1790. S Fp Bot reg. 508
quadrifarium Lie.	fsur-parted	3 my.o Y	182a S l.p
mexichnum Hum.	Mexican	3 au.s Y	Mexico 1825. D p.l
*2360. TRICHOPHYLLUM tfirf. TuchoraYLLUM.	(r/m>, a hair.pAy/Zem.aleafO	<i>Comp. Heliahn.</i> 1.	
oppositifolium Nut.	opposite-leaved	^ or. . . . Y	Missouri ... D co
2361. HE'LLIS L. DAISY.	(<i>Bellus</i> pretty; flower.)	<i>Comp. Card. Vern.</i> 4.—6.	
perennis L.	common perennial	£ mir.au W	Britain past D co Eng. bot. 424
hort^ndis	garden, large-double	A Pr imr.au R	... D co Bot. mag. 228
3 fistulosa	piped, double-quilled	& pr \ mr.au R	... D co
4 prolferous, Hen SJChicksi	£ pr	imr.au St	... D co
21596 hibrida Ten.	hybrid	H A or i. mr.jn W.R	Italy 1824. D co
21587 sylvestris Cyr.	wood	£ A pr fmy.U W	Portugal 1797. D co Bot. mag. 25U
21588 annuaL.	annual	O pr 1. mr.il W	S. Europe 1759. S co Bot. mag. 2174
23R2. BE/LLIUM L.	BELLIUM.	(<i>Bellis</i> , daisy; similarity.)	<i>Comp. Card. Vern.</i> 2.
21589 tellioloides L.	Daisy-like	O pr ^ jn.8 W	Italy 179& S s.p Lam.il 634
Xtto minutum L.	Minute	■ A cu linjn.o W	Levant 1772. D co Ac. up. 1.5.2
9*iSP	DAILI.	(<i>Undrew Daht</i> , a Swedish botanist, pupil of Linn.)	<i>Comp. Bdidn.</i> 2. -3.
		Double Garden Varieties.	
Agathe imperiale	Eclipse	Marquis	Sans rival
Agathe royale	Excellent	Morison	Sovereign
Atropurpurea	Fawn-colored	Mutabilis	Sprclsa
Beauty of England	Fimbriata	Orange flag	Spring grove lilac
Beauts suprême	Floribunda	Princess Alexandria	Straw-colored
Belvidere	Flonbunda nna	Victoria	Sulph. fcrea grandiflbra
Bright purplic	F6lgida -	Princess Elizabeth	Sulph. hbrea specibsa
CuneUrijbra	Grand Alexander	Ptilchra	Sunflower
Chancellor	Grin de Lin	Purp&rea ritgra	Superb crimson
Changeable purple	Henriette	Royal olive	Tendre Agathe
Comet	Insignia	Royal purple	Victory
Crimson (Wells's)	Komng Aza	Royal sovereign	Waratah
Crimson (Young's)	L'Honneur d'Anvers	Ruby	White Dutch
Dodoneus	Lilac-striped	Sablni	Yellow Clifton
Duchess of Gloucester	Luttalu	Sanguinca	Yellow dwarf
B'legans	Ma Favorite		

Agathe imperiale Eclipse Marquis Sans rival
 Agathe royale Excellent Morison Sovereign
 Atropurpurea Fawn-colored Mutabilis Sprclsa
 Beauty of England Fimbriata Orange flag Spring grove lilac
 Beauts suprême Floribunda Victoria Straw-colored
 Belvidere Flonbunda nna Princess Alexandria Sulph. fcrea grandiflbra
 Bright purplic F6lgida - Princess Elizabeth Sulph. hbrea specibsa
 CuneUrijbra Grand Alexander Ptilchra Sunflower
 Chancellor Grin de Lin Purp&rea ritgra Superb crimson
 Changeable purple Henriette Royal olive Tendre Agathe
 Comet Insignia Royal purple Victory
 Crimson (Wells's) Komng Aza Royal sovereign Waratah
 Crimson (Young's) L'Honneur d'Anvers Ruby White Dutch
 Dodoneus Lilac-striped Sablni Yellow Clifton
 Duchess of Gloucester Luttalu Sanguinca Yellow dwarf
 B'legans Ma Favorite

21592 frust'anea ft K.	<i>barren rayed</i>	A A or	6 s.n.	S	Mexico	1802.	R h.l	Cav.icl.SS6
\$ 2 folgens	<i>scarlet</i>	ji ^ or	6 sn	S	Mexico	1802.	R h.l	Bot. mag. 762
3 aurântia	<i>orange-colored</i>	A A or	6 s.n.	O	Mexico	1802.	R h.l	
4lutea	<i>yellow</i>	A A or	6 8.n.	Y	Mexico	1802.	R h.l	
5cr6cea	<i>saffron</i>	A A or	6 s.n.	Su	Mexico	1802.	R h.l	
2364. BCE'BERil W.	BCEDERA.	(<i>Barber, a learned Russian botanist</i>)				Comp.JacobEaf.	2.-7.	
81593 chrysanthemlddes TV.	Chrysanth.-lk	O pr l.jo	Y	Carolina	1821.	S l.p		
Dyssbia glutinbsa Cav.								
21594 porophylla Kth.	Porophyllum	O u n li	Y	Mexico	1817.	S co		
Dysbodia porophylla Cav.								
•2365. TAGE'STES L.	TAGETES.	(<i>Tages, a Tuscan divinity; beauty of flowers.</i>)				Cotnp. Jacobi*	14.	
		I. INTEGRIFOLIA.—Leaves entire.						
21595 lucida Cav.	ahlningJeaved	j& iAJ or	1 jl.n	Y	S. Amer.	1798.	D p.1	Bot. mag. 740
		II. PINNAVC.—Leaves pinnated or bipinnated.						
21596 pretula Z.	spreading	f%*fricat Marigold	O or	2 jl.s	Y.o	Mexico	1573.	co Bot mag. 150
21598 corymbbsa Swt.	corymbose	O or	1}aus	P	Mexico	1825.	co	lAm. il. 684.
2 latca	yellow	8 or	1}aus	Y	Mexico	1825.	co	Sw. flgar. 151
21599 subvillbsa LAg.	subvillous	O or	1}aus	Y	Mexico	1823.	co	
21600 minita L.	minute-flowered	O or	2 8	au.o	Pa.Y	Chile	17*8.	co
21601 caracasana Hum.	Caraccas	O or	3 au.s	Y	Caraccas	1819.	co	
21602 tenuifolia Cau.	fine-leaved	O or	3 JL0	Y	Peru	1797.	co	
21603 angustifolia Kth.	narrow-leaved	O or	3 au.s	Y	Mexico	1826.	co	
21604 clandestina Lag.	concealed	O or	3 JL0	Y	Mexico	1823.	co	
coronopifolia W. en.								
21605 filifilia Lag.	thread-leaved	O or	3 au.s	Y	Mexico	1826.	co	
21606 micranthi Cav.	small-flowered	O or	3 JL0	Y	Mexico	1822.	co	
21607 glandulosa Sehrank	glandular	O or	3 JL0	Y	& Amer.	1819.	co	
2WU8 glandulifora Sehrank	glanduliferous	O or	6 o.n	Y	Mexico	1826.	co	
2366L HETEROSPERMUM Cav.	HETEROS.	(<i>Heteros, variable, spertna, seed; shape.</i>)				Comp.Hclii'in.	1.-3.	
2KM9 pinnatum Cav.	wing-leaved	O or	2 au.s	Y	NewSpain	1799.	S co	Cav.icS.2tTl
2367. SCHKU*HR/i* Roth	SCHKUIRIA.	(<i>Christian Schkuhr, acute German botanist.</i>)				Comp.Heliidn.	f	
21610 abrotanoides Both	Wormwood-like	O un 2	jl.s	Y	Mexico	1798.	S co	Schkhan.3.250.b
2368. PE'CTIS L.	PECTIS.	(<i>Pecten, a comb j teeth of pappus.</i>)				Comp.Helianthea.	5.—10.	
21611 humifusa Sum	trailing	* iQJ un }	jl.au	Y	Jamaica	1824.	S co	Pl.ic.95.a
21612 prostrata Cav.	prostrate	Jc iQJ un	i jUau	Y	W.Indies	1818.	S co	Cav. ic.4.324
21613 ciliâris L.	ciliated	O un 1	jl	Y	Hispania	1793.	S co	PI. ic. 151. 2
21614 /im folia £.	Flax-leaved	O un 1	jl.au	Y	Jamaica	1732.	S co	SI. jam. 1.149.3
21615 canescens Kth.	cane>cent	-* O un	jl.au	Y	Mexico	1826.	S co	
2369. LONGCHA'MFSIA W. LONOC.	(<i>J.L.A Loiseleur Deslongchamps, M.D., French bot.</i>)	Comp.Heliidn.	1.					
21616 capillifoha W.	hair-leaved	O pr J jnjl	Y	Barbary	1822.	S co		
Pectis discoidea Horn.	Leyson discoidea Spr.	Gnaphalium leysero)des Dcs£						
2370. LEY'SEIM L.	TRYBERA.	(<i>Frederick William Leyser, a German botanist.</i>)				Comp.Eup.	4.—10.	
21617 pohtolia Thun.	Folium-leaved	a. i_J pr	'Hjl.8	Q	C. G. H.	1820..	P.I.	
21618 gnaphalides L.	woolly	«L LJ P*	2 jl.s	O	C. G. H.	1774.	P.I.	
21619 nquarrbsa W.	squarrose	«. i_I pr	2 jl.s	O	C. G. H.	1615.	P.I.	Pluk.aL302.3
Staehellna gnaphalbes L.								
21620 ciliata Thun.	ciliated	n. \J pr	J	Q	C. G. H.	1816.	S p.1	
237L SEI.LO^ Spr.	SBLLOA.	(<i>Fred. Sello, a German botanist in Brazil.</i>)				Comp.Eupatrese.	1.—2.	
21621 glutinosa Spr.	clammy	• un	3 f	Y	Brazil	1819.	C co	Bot. reg. 462
2372. RELHA Wil Herit.	RELHANIA.	(<i>Rev. Richard Relhan, author F.J. Cantab.</i>)				Comp.Helian5.	—	
21622 squarrbsa Herit.	squarrose	tL i_J pr		Q	C. G. H.	1774.	P.I.	
21623 pftngens Herit.	pungent	tt.	LJ P'		C. G. H.	1820.	P.I.	Bot reg. 587
21624 lateriflora Herit.	side-flowering	<. i_J pr		Q	C. G. H.	1823.	P.I.	
21625 /enHifflua Herit.	Genista-leaved	*L I_J pr	1 my.jn	Y	C. G. H.	1823.	P.I.	
Athanasia fenistifdia L.								
21626 paleXcea Herit.	chaffy	tt.	_I pr	1 my.jn	Y	C. G. H.	1818.	C p.1 Gac. fr. 173.9
Lejaera mto'ides Berg.								
2373. ZI'NNi/f L.	ZINNIA.	(<i>John Godfrey Zinn, prof. bot. Gottingen.</i>)				Comp.Heliahthee.	8.—	
		I. SESSIMFLORA.—Flowers sessile.						
21627 pauciflbra L.	few-flo'reved	O or	2 jl.au	Y	Peru	1753.	S rxn Mil.ic.	1. 64
		II. PEDUNCULIFLU&K —Flower* pedunculate.						
21628 multiflbra L.	many-flowered	O or	1 jn.o	R	N. Amer.	1770.	S r.m	Bot mag. 14)
21629 verticillkta Andr.	whorl-leaved	O or	1 jl.au	R	Mexico	1789.	S r.m	Bot. rep. 189
21630 ^legans Jac.	elegant	O or	1 ins	S	Mexico	1796.	S rra	Botmag.fi27
iriofaca H. K.								
21631 tenuiflbra Jac.	slender-flowered	O or	2 jl.au	S	Mexico	1799.	P.I.	Bot mag. 555
21632 revoluta Cav.	revolute	O or	2 jl.au	S	Mexico	1817.	P.I.	
21633) angustifolia HUTU.	narrow-leaved	O or	2 jl.au	R	Mexico	1824	P.I.	
21634 h^brida B. M.	hybrid	O or	2 jnjl	S	S. Amer.	1818.	P.I.	Bot mag. 2123
*2374. CHRYSANTHEMUM L.	CHRYSANTHEMUM.	(<i>Chrysos, gold, anthemon, fl.</i>)				Comp.Anthem.	30.—dS.	
		I. FRUTICOSA.—Shrubby i rays white.						
21635 pinnatifidum L.	plnnatifid	tk \J or	3 my.au	W	Madeira	1777.	C p.1	
21636 anomalous Lag.	anomalous	n. J or	1 jn.jl	W	Spain	1816.	D co	
21637 radicans Cav.	xooWng-brancied	ft i_J or	1 my.au	W	Spain	1818.	C s.1	
		II. ALBIFLORA.—Herbaceous : rays white.						
21638 paludsum Desf.	marsh	^ A	11 j1	W	Barbary	1810.	S co	Desf. at. 2.239
21639 rotundiflrium W.^r-K	round-leaved		11 j1	W	Hungary	1817.	D co	
21640 mexicanum Bonp.	Mexican		11 j1	W	Mexico	1825.	D co	
21641 atratum TV.	blackened-tai		11 j1	W	Austria	1731.	D co	
2 lobatum Schl.	lobed		11 j1	W	Switzerl.	1806.	D co	
21642 heterophyllum W.	various-leaved		11 j1	W	Switzerl.	1806.	D co	
21643 leucanthemum L.	white-flowered		11 j1	W	Britain	pas.	D co	Eng.bot.601

III. FLAVIFLORA. — *Herbaceous; rays yellow.*

21658 <i>segutumL.</i>	corn	O or	1] Jnau	Y	Britain	com fl. &	co	Eng. bot. 540
21859 <i>Mychufc v.</i>	Mycon'a	Q or	1] jal.	Y	Italy	1775.	co	Jacob. 4.9*
21660 <i>italicum L.</i>	Italian	* A or	2 injl	Pa.Y	Italy	1796.	co	n
21661 <i>coronarium L.</i>	garland	O or	4 jls	Y	Sicily	1629.	co	Lam. il 678.6
21662 <i>indicum L.</i>	Indian	£ A or	2 Bn	Y	China	-	r.m.	

IV. Valliflora. — *Herbaceous; flower variable*

21663 <i>tripartitum</i> <i>Swt.</i> <i>indicum</i> <i>Pen.</i>	triparted	£ A or 3	an	T	E. Indies	1800.	C r.m Sw. fl. gar.	193
21664 <i>sinense</i> <i>Sab.</i>	Chinese	£ A or 3	o.n	Va	China	1764.	C r.	

Garden Varieties

1 Purple <i>Sot. mag.</i> 327	16 large quilled Orange <i>Hort.</i> to.	28 Pale Buff
2 Changeable White <i>Sot. mag.</i> 2042	* 	29 Early Blush
3 Quilled White <i>Sot. reg.</i> 4	17 Expanded light Purple	30 Blush Ranunculus-flowered
4 Superb White <i>Sot. reg.</i> 455	18 Quilled light Purple	31 Changeable pale Buff
'5 Tabled White	19 Curled Lilac <i>Sw. f. gar.</i> 7	32 Two-colored Red
6 Quilled Yellow	20 Superb, clustered Yellow <i>Sw. fl.</i> gxr. 14	33 Starry Purple
7 Sulphur Yellow	21 Semidouble quilled Pink <i>Hort.</i> tr. "5.17"	34 Brown Purple
8 Golden Yellow <i>Sot. teg.</i> 4*	22 Semidouble quilled White	35 Late quilled Yellow
9 Large Lilac	23 Semidouble quilled Orange <i>Hort.</i> tr. "5.17" *	36 Double Indian Yellow
10 Rose or Pink	-	37 Parks' small Yellow
11 But! or Orange	24 Late pale Purple	38 Tasseled Yellow
12 Spanish Brown	25 Quilled Salmon-colored <i>Hort. tr.</i> 5.17*	39 Tasseled Lilac
13 Quilled flamed Yellow <i>Hort.</i> Ir. 4.14	26 Small Yellow <i>Hort. tr.</i> 5.17**	40 Semidouble quilled pale Orange
14 Quilled Pink <i>Sat. reg.</i> 616	27 Paper White	41 Golden Lotus-flowered
15 Early" Crimson <i>Hort. tr.</i> 5.3		42 Two-colored incurved
		43 Yellow Waratah
		44 Double Indian White

2375. PYRFTHRUM Sm. FEVERFEW. {*Pyr*, firej hottasted root.

Compds & Anthemicida. 49,

I. FninchSA.—	<i>Shrubby j rays white.</i>						
2W65 crithmiflbumJK.en. Samphire-lvd	■ L or	3 year	W	Teneriffe	1815.	C	co
2(666anethiflbumJK.<i> Dill-leaved	« i o r	8 year	W	Teneriffe	1815.	C	co
&6R7 /oDniculaceum W. en. FenneUtaiwed	* LJ or	3 year	W	Teneriffe	1815.	C	co
[1668 frut&cens L. ftutescent	« U o r	3 year	W	Canaries	1699.	C	p1
■1669 «(ronoi) ifflbumJK.en. Buckhorn-lvd u or		2 year	W	Canaries	...	C	L-p
BJ2? T ^{an} d fl'rumH'. «gffcat-flowered	• U o r	8 year	W	Canaries	1815.	C	Lp
8flf! Broussoneti Cham. Broussonet's	* i o r	2 jla	W	Canaries	1817.	C	81
«672 speciosum W. en. showy	si LJ or	3 year	W	Canaries	1815.	C	Lp

H. ALBIFLORA *Herbaceous plants, white*

II. ALBIFLORA. — <i>Herbaceous</i>		<i>Days white.</i>		W. Indies 1817. D co	
1673 <i>simpliciflum</i> W.	simple-leaved			Pyrenees 1820. D co	
"1674 <i>latifolium</i> W. en.	broad-leaved				
grandiflorum <i>Lap.</i>					
1675 <i>maximum Dec.</i>	laTgest			S. Europe 1818. D co	
"1676 <i>seritum WK.</i>	Ax-flowering			N. Amcr. 1731. D co	Jacob. 4.90
1677 <i>aluiticum Kit.</i>	marsh			Hungary 1816. D co	Bot mag. 2704
1678 <i>Ailsamita IK.</i>	Balsamita			Levant 1779. D co	Jacob. 4.89
1679 <i>Halleri W.</i>	Haller's			Switzerl. 1819. D co	Bar. ic. 458.2
1680 <i>alpinum IV.</i>	alpine			Switzerl. 1759. D co	
2 pubescens	pubescent			Switzerl. 1819. D co	
1681 <i>diversiflum Grah.</i>	variable-leaved			N. Holl. 1823. D co	
1682 <i>Barrelieri Lk.</i>	Barrelier's			S. Europe 1820. D co	
1683 <i>tomentosum Lot.</i>				Corsica 1818. D co	
1684 <i>pulverulentum W.</i>	powdery			Caucasus 1806. D co	
"1685 <i>ceratophyllum W.</i>	Hornwort-like	£ A		Piedmont 1819. D co	Al.ped. 1.37.1
"1686 <i>pal Astro W.</i>	marsh	£ A or		Armenia 1820. D co	
iY\$ <i>Pinnatifidum W.</i>	pinguidifid	£ A or	2 my.jn 1823. D co	
HS88 <i>macrophyllum W.</i>	long-leaved	£ A pr	3 jl.au	Hungary 1803. D co	W.&K. 1.94
ilchillea j <i>Alpinicifolia Def.</i>					
1689 <i>corymbsum IV.</i>	corymbose	£ A	1 ln.au	W Germany 1596. D eo	Jac.au.4.379
1690 <i>cinerariaeflum hun.</i>	7>cvc. <i>Cineraria-lvd</i>	£ A or	2 jl.au	Dalmatia 1820. I) s.l.	
x1691 <i>Parchemum Sm.</i>	common Eellitory	£ A w	2 ns	Britain rubb. D co*	Eng. bot. 1231
2flbrpl:ino	double-flowered	£ or	£ hs	W. C r.m	
1692 <i>Mandrina Hort.</i>	Mandiana	£ A or	lijnuu	France 1816. S co	
"1693 <i>partheniiflum W.</i>	Parthenii-lvd	£ A pr	2 jtjl	Caucasus 1804. D co	Ven.cel.«43
1694 <i>caucasicum W.</i>	Caucasian	£ A P'	ijjl.W	Caucasus 1804. D co	
2b-95 <i>sericeum Bieb.</i>	silky	£ A F	1 jn.au	W. Iberia W2;l D co	
%W% <i>tenuiflum Ten.</i>	slender-leaved	£ A pr	1 jl.au	Caucasus IBC. D co	
21697 > <i>arvili-crum W.</i>	small-flowered	O un	2ijn.W	IS.O. S co	
21698 <i>pne Vn Bieb.</i>	early	O or	1 jnjl	Caucasus 1818. S co	
slifly <i>pne Vn Bieb.</i>	Sncezowort-lvd	£ A pr	lijLau	Caucasus 1803. D co	
700 " <i>thiodictiflum W.</i>	scentsless	O pr	1 au.s	Britain dryfl. S co	Eng. bot. 676
2 fibrous plenis	large-double	Jk A «	i jn.au	W. 1825. C 81	
grandifl. <i>Cirum Hort.</i>					
g	Aegant	Jc O or	1 jl.au	W Mt.Baldo 1816. S co	
niaritimum Sp.	sea	£ Apr	1 jn.o	W Britain seash. D co	
	shaded	O or	ijjl.W	S. Europe 1821. S gl	
	fine-leaved	£ A or	1 au.3	Caucasus 1821. S s.l	

A a

III. RUDRIFLORA.—Herbaceous; rays red.									
21705 <i>carneum</i> Bleb.	flesh-colored	£ A cl	Iftau.s	Pk	Caucasus	1804.	D co	Bot. mag.	1080
Chrysanthemum <i>coccinum</i> R. M.									
IV. FLAVIFL5RA.—Herbaceous; rays yellow.									
21707 <i>Indicum</i> Rox.	Indian	O un	I	E Indies	1810.	S	p.l	Bot. mag.	1021
21708 <i>breviradiatum</i> Led.	short-rayed	O or	I au	-	1818.	S	co		
21709 <i>Bocconii</i> W.	Bocconi <i>ii</i>	A or	I au	Spain"	182a	S	co	Bar. ic.	1153
21710 <i>trifurcatum</i> W.	three-forked	O or	I au	Barbary	1820.	D	co	Desf. at	2,235.2
21711 <i>bipinnatum</i> W.	bipinnate-/w*	£ pr	I	Siberia	1796.	S	co	Gm. si.	2,85.1
21712 <i>millefoliatum</i> IV.	thousand-lvd	2 my.8	I	Siberia	1731.	D	co	Mil. ic	1,9
Milfoil-leaved	*	A pf	2 au.s	Caucasus	1823.	D	co	Gm. sl	86.2
2376. MATRICA'RIA L. MATRICARIA. (<i>Matrix*</i> the womb; sup. mcd. qual.) Comp. Anthem. 4.—5.									
21714 <i>suaveolens</i> L.	sweet-Binelhng	O un	I jjn.au	W Europe	1781.	8	co		
21715 <i>Chamomilla</i> L.	Wlf/tfChamomile	O w	I my.jl	Britain	ro. sid.	S	co	Eng. bot.	1232
21716 <i>capt'-nsia</i> L.	Ca'de	JU Ol un	I jl.s'	W	*C. G. H.	1699.	S	Scba	1,16.2
21717 <i>pusflu</i> W. en.	small	O un	I	W	S	co		
2377. BOLTCN/i* Herit. BOLTONIA. (J. B. Button, an English botanist) Comp. Card. Vem. 2.									
21718 <i>astroledes</i> Herit.	Starwort-like	^ A pr	3 au.o	F	N. Amer.	1758.	D s.l	Bot mag.	2554
21719 <i>plastifolia</i> Herit.	Woad-leaved	3 A pr	Ih	Pk	N. Amer.	1758.	D s.l	Bot mag.	2381
2378. LIDBE'CK/4 Berg. LIDOECKIA. (R. G. Lidbeck, a Swedish botanist.) Comp. Anthem. 2.—4.									
21720 <i>pectinata</i> Berg.	pectinated	III Jpr	2 my.jn	Y	C. G. H.	1774.	C Lp	Berg. c.	306.5. 9
21721 <i>lobata</i> Thun.	fobed	II I J pr	2 my.jn	Y	~ C. G. H.	1800.	C l.p	Lain. il.	701.3
Cotula quinque'loba L.									
2379. CE*NIA Com. CENIA. (Kenos, empty; inflated calyx.) Comp. Ssitcc Anthemidece. 1.									
21722 <i>turbinata</i> Pert.	turbinated	O un	I jl.au	W	C. G. H.	1713.	S co	Lam.il.	701.1
Lidb'ckia turbinata Thun.									
2380. COTCLA L. COTULA. (Dim. of <i>cola</i> , old name for a species of j4'nthemis.) Comp. Anth. 7.—29.									
21723 <i>anthemicAdes</i> L.	Anthemis-like	cm un	I jl.au	Y	St.Helena	1696.	S co		
21724 <i>aurea</i> L.	golden	O or	I jl.au	Y	Spain	1818.	S co		
21725 <i>coronopifolia</i> L.	Buckhorn-lvd	KB un	I jl.au	Y	C. G. H.	1683.	S co	Lam. il.	700.1
21726 <i>viscosa</i> L.	clammy	2 un	I i au	W	VeraCruz	1739.	D l.p		
21727 <i>tanacetifolia</i> W.	Tansy-leaved	O un	f jn.au	Y	C. G. H.	1783.	S co	Pluk.m.	430.7
2H& <i>sphrantza</i> Lk.	round-flowered	ES un	2 my.jn	Y	Congo	1821.	D co		
21729 <i>nudicaulis</i> Thun.	naked.stemmcdj	iAJ or	i jl.au	Y	C.G. H.	1816.	C l.p	H. sys.	69.4
2381. GRA'NGE, 4 J. GRANGEA. (Grange, probably a man's name.) Comp. dsitac Anthemide* 7.—									
21730 <i>chin^nsis</i> Lam.	Chinese	O un	} jCs	Y	China	181&	S, co	Lam.il.	699.2
Cotula cuneifolia W.									
21731 <i>minima</i> W.	least	O un	Y	China	1768.	S co	Bur. in.	58.3	
21732 <i>decimbens</i> Deaf.	decumbent	O or	4 Jl.au	Y	N. Holl.	1816.	S co		
21733 <i>maderaspatica</i> Lam.	Madras	O un	\$ jl.au	Y	E. Indies	1780.	S co	Lam.il.	699.3
21734 <i>cinerea</i> Del.	cinereous	O or	i jl.au	Y	1818.	S co			
21735 <i>bifcolor</i> W.	two-colored	un	1 Jl.au	W, Y	ETindies	1804!	S co	Lam. il.	1599.1
Cotula latifolia Desf.									
21736 <i>wnchifolia</i> Bieb.	Sonchus-leaved	O or	A jl.au	I	Caucasus	1821.	S co		
2382. ANA'CYCLUS L. ANACYCLUS. (An, priv., <i>aniphos</i> , flower, <i>kyklos</i> , circle; ovaries.) Comp. Anth. 8.—									
21737 <i>anthemoides</i> Ztg.	Anthemis-like	^ A wh	I jatl	Pa, Y	S. Europe	1727.	D co		
Santolina <i>anthemoides</i> L.									
21738 <i>creticus</i> L.	Cretan	O un	I jn.au	Y	Candia	1759.	S co	An. mu.	11.22
21739 <i>orientalis</i> L.	Oriental	O un	li jn.au	Y	Levant	1731.	S co	Boer.	1,110
21740 <i>aureus</i> L.	golden-flowered	O pr	I jn.au	Y	Levant	1570.	S co	Lam.il.	700. 2
21741 <i>valentinus</i> L.	Valentia	O un	I jn.jl	Y	Spain	1656.	S co	Schk.han.	3,254.t>
21742 <i>radiatus</i> Lk.	rayed	O pr	I jlj.lau	Y	S. Europe	1596.	S s.l	Brcy. c.	75
if'nthemis <i>valentina</i> Jr.									
21743 <i>davatus</i> Lk.	clubbed	O cu	I jl.au	W	Barbary	1801.	S co	Biv. cen.	1.7
21744 <i>divaricatus</i> Cav.	divaricate	O un	1 f jl.au	W	Spain	1818.	S co		
* 2383. CLADA'NTHUS Cos. CLADANTHUS. (<i>Kados</i> , branch, <i>anthos</i> , Bower; on branches.) Comp. Anth. 9.—									
21745 <i>arabicus</i> Cas.	Arabian	O pr	1 jl.au	V	Barbary	1759.	S co	Sin. spic.	9.10
il'nthemis <i>arabica</i> L.									
2384. if'NTHEMIS L. CHAMOMILE. ("(Anthemon, a flower; plants covered with.) Comp. Anth. 45.—49.									
I. FLAVIFL5RA.—Rays yellow; leaves bipinnatifid or bipinnate.									
21746 <i>tinctoria</i> L.	LyF*.	^ A pr	I jn.n	Y	Britain	stopl.	D co	Eng. bot.	1472
21747 <i>Rudolphina</i> Bieb.	Rudolph's	A or	1 jn.jl	Y	Caucasus	1824.	D		
Pyréthrum orientale W.									
21748 <i>Marshalliae</i> W.	Marshall's	or	14 jn.jl	Y	Caucasus	1816.	D		
21749 <i>discoidea</i> W.	discoid	pr	1 jl.au	Y	Italy	1800.	S		
II. BIPINNATIFIDIF.—Rays white; leaves bipinnatifid or bipinnate.									
21750 <i>fallax</i> TV.	uncertain	O pr	I jl.au	W	.	1825.	S co		
21751 <i>Pyréthrum</i> L.	Pellitory of Siain	A or	I jn.jl	W	S. Europe	1570.	D s.p	Jac. sc. L.	371
21752 <i>globosa</i> W.	Globose	LAJel	I au.s	W	S. Europe	1570.	D co	Bot. reg. Sia	
21753 <i>apiifolia</i> B. Br.	L'arsly-leaved	I Alpr	I au.s	W	China	1819.	D co	Eiig. bot.	980
21754 <i>nobilis</i> L.	common noble	A ni	I	W	Britain	gra.pa.	D co		
2 fibre pleno	double	% A	I	W	Britain	gard.	D co		
21755 <i>grandiflora</i> Dec.	great-flowering	A or	jnau	W	S. Europe	1835.	D co	Eng. bot	602
21756 <i>aryensis</i> L.	corn-field	£ Q	Q w	jn.au	Britain	dryfl.	S co	Eng. bot.	177<
21757 <i>Cotula</i> L.	Cotula	O w	I jn.s	W	Britain	corn fl.s	S co	W.h.b.	1,62
All.	rigescnt	£ A pr	I ls	W	Caucasus	1805.	D co	Jac. au.	5,44#
Nico	Austrian	O un	I myau	W	Austria	1759.	S co		
	Triumfetti's	j£ Q or	I jl.au	Pa, Y	Switzcrl.	1819.	S co		
parefima	Nico	O un	I jl.au	W	Italy	1819.	S co		

21762 chla L.	Chian	O H	lijno	W	Chio	1731.	S co
21763 altissima L.	tallest	O P	4 au	W	& Europe	1731.	S co
21764 Cite L.	Cota	O	jl.au	W	Italy	1714.	S co
21765 punctate Vahl	dotted	U 1	aus	W	Barbary	1818.	S co
21766 publscens TV.	pubescent	A pr	jl.au	W	S. Europe	1803.	D co
21767 Chamomilla TV. en.	Chamomile	A pr	jl.au	W	S. Europe	1817.	D co
21768 fuscate Brot.	brown-scaled	O pr	jlau	W	Portugal	1805.	S co
21769 ruthenica Bieb.	Russian	O pr	my.jn	W	Tauria	1823.	S co
21770 fruticulata Bieb.	shrubby	A pr	H	W	Caucasus	1820.	D co
21771 retissa Lk.	rectuse	O un	jl.au	W	1824.	S co
21772 secundiramea Biv.	side-branched	O un	1 jl.au	W	Sicily	1833.	S co
21773 ramosa Lk.	branchy	O un	1 jl.au	W	1824.	S co
21774 coronopifia Mia TV.	Buckhorn-lvd	A pr	1 jn.jl	W	Spain	1818.	D co
21775 saxatilis Kit.	rock	A pr	1 jl.au	W	Hungary	1807.	D co
21776 anglica Spr.	English	X O un	i jlau	W	England sea co.	9	co
maritima E. B. not L.							Eng. bot. 237a

21777 incrassata toj.	thick-pc<uncldcf	A or 1	jl.au	W	France	1818.	D co
21778 pedunculate Desf.	peduncled	O un	1 jl.jlau	W	Spain	1816.	SS Co
incrassata Lk.							
21779 tomentosa L.	downy	X	+	+	Levant	1795.	D co
21780 carpatica TV.	Carpathian	*	+	+	Carpathia	1820.	D co
21781 montana L.	mountain	*	+	+	Italy	1759.	D co
21782 peregrina Bieb*.	foreign	*	+	+	Caucasus	1823.	S co
21783 ibdrica Bieb.	Iberian	*	+	+	Iberia	1820.	D co
21784 petraea Ten.	rock	*	+	+	Italy	1825.	S co
21785 Barrelieri Ten.	Barrelier's	*	+	+	Italy	1825.	S co
21786 mixta L.	mixed	*	+	+	France	1731.	S co
21787 mclampodia Del.	black-footed	*	+	+	Egypt	1819.	D co
21788 maritima L.	sea	*	+	+	Mcditer.	1800.	S.s
21789 alptaa L.	alpine	*	+	+	Austria	1824.	D co
21790 KUaibelli Spr.	Kitebel's	*	+	+	Hungary	1823.	D co
sax&tilis Kt.							Jac. au. 30

21385. CENTROSPE'RMUMSpr. CBNTRBPEBMUM. (*Keniron*, spur, <p>wa, seed.) *Gmp.Anthem.* 1.
21791 chrysanthum Spr. yellow-flowered O pr I JUu Y Spain 182J. S co

2386. SANVITA^LW Cav. SANVITALIA. (Most probably a man's name.) *Ctmp. Helidnthe**. 1.-2.
21792 procumbens Cav. trailing -* O or 1 jl.au Y Mexico 1798. S co Bot. reg. 707

•2387. ilCHILLEM L. MILFOIL. (*Achilles*, pupil of Chiron, first used the plant in med.) *Com.Anthcm.lt.*

21793 lingulata JPff.	tongue-leaved	A or 1	jl.au	W	Hungary	1815.	D co	W.&K.1.2
21794 Herba-rote All.	Herba-rota	A or 1	jl.jl	W	France	1640.	D co	Al.ped. 1.9.3
21795 grandiflora Bieb.	great-flowered	A or 1	jl.jl	W	Caucasus	1815.	D co	
21796 Ptarmica L.	Sheezewort	A or 1	jl.n	W	Britain moi. pi.	D co	Eng. bot. 757	
21797 Ptarmica L.	double-flowered	A or 2	JLn	W	Britain fields.	D co	Eng. bot. 2531	
21798 decolorans Schr.	staining	A or 1	jn.au	W.Y	1798.	D co	
21799 mongolica Fis.	Mongolian	A or 1	jljl.au	W?	Siberia	1818.	D co	
SJ99 ⁰ speciosa Hqe.	showy	A or H	JU	W	1804.	D co	
21801 biserrata 2 nd .*	biserrate	A or 1*in.jl	W	Albania	1820.	D co		
21802 cuneata Arte.	Sweet Maudlin	A or 2	au.o	W	Italy	1734.	D pl	
21803 ^ g 1	II. PINNATIFIDJE.—Leaves phmatMd, o				S. Europe	1570.	D s.p	
21804 alptna Z.	alpine	A or Jjl.n	W	Siberia	173K	D 8.p	Boc.mu.144.101	
21805 serrata Retx.	wn4eaved	A or 2	aus	W	SwitzerL	1686.	D co	
21806 coronopifolia W.	Buckhorn-lrd	li	jl.au	Pa.Y	Levant	1823.	D co	W. ach. 1.2
21807 Clareñrup L.	Clavenna's			W	Austria	1636.	D pl	Bot. mag. 1287
21808 impatiens L.	impatient	2	jns	W	Siberia	1759.	D co	Gin. si. 2.8a 1
21809 Gérberi W.	Oerber's	li	jl.au	Pa.Y	Siberia	1821.	D co	Om.si.83.3
21810 recurvifolia Lap.	recurved-lvd	U	jl.jl	W	Pyrrenees	1820.	D co	
21811 pectinata W.	comb-leaved	li	aus	Pa.Y	Hungary	1801.	D co	W. & K. 1.34
ochroleuca Wald.								
21812 ehamaemclibha Potf. Chaniomile>lvds	A or	C	W	France	1825	D co		
21813 asplonifolia Ven.	Asplenium-lvd	A or	lijn.au	Pk N. Amer.	1803.	D s.p	Ven. eels 93	
21814 squarrbsa H. K.	ToughJicdedd	A or 1	jl.au	W	1775.	D pl	

III. PINNAVJE.—Leaves pinnate.

21815 falcata L.	sickle-fcavof	A or 1	jns	Pa.Y	Levant	1739.	D co	Lam.il. 683.3
21816 imbricate Ven.	imbricated	A or li	jn.au	Y	Persia	1818.	D s.l	Ven. eels 54
21817 tenuiflolia Lam.	slender-leaved	n. or 1	jn.au	Y	Levant	1733.	C co	
21818 Santolina L.	Santolina	A or 1	jn.au	Pa.Y	Levant	1759.	D pl	
21819 santolinoides Lag.	Santolina-like	ji	A or 1	jn.au	W	Spain	D co
21820 albida JV.en.	whitish	A or 1	jl	Pa.Y	1819.	D co	
	iiL HORIZONTALS.—leaflets horizontal.							
21821 onthemo'idca TV.	Anthemis-lk		1	jn.au	Pa.Y	D co	
21822 dextiibcns I<am.	decumbent	** A or i	jn.au	Y	Kamtsch.	1816.	D s.l	
21823 moschata Jac.	musk	A or 2	jl.J	W	Italy	1775.	D co	Jac. au. 5. ap. 33
21824 atfateL.	black-cupped	A or 2	jl.8	W	Austria	1596.	I) co	Jac. au. 1.77
21825 nana L.	dwarf	A or 2	jl.8	W	Italy	1759.	I) co	Al. pcd. 1.9.3
21826 erotica L.	Cretan	A or ij	jl.jl	W	Candia	1739.	D pl	Boc. mu. 34
21827 acyptUca L.	Egyptian	**-i-Jor	jl.s	Pn.Y	Levant	1610.	C pl	To.it. 1.87
21828 holosericea Sm.	velvety	5t A or	jl.s	W	Parnassus	1817.	D co	
21829 auriculata TV.	eared	k A or	n.au	Y	A. Minor	1827	D co	
21830 vallesaca Suter	Vallicsian		n.au	W	Switzcr.	1819.	D co	
helvetica Schl.								
21831 macrophilla L.	long-leaved	A or 3	jlau	W	Italy	1710.	D co	Tri.ob:23
	IV. BIPINNAVAS.—Leaves bipinnate and bipinnatifid.							
21832 compacta TV.	compact	A or 1	jl.au	P.i.Y.	1803.	D co	
21833 glomerita ^wft.	glomerateil	A or 1	jl.au	Y	Caucasus	1818.	D ro	
	A a 2							

21834 punctata Ten.	dotted	£ A or	1	Jl.au	Str	Naples	1820.	D co
21835 leptophylla Bieb.	slender-leaved	£ A or	1	jl.au	Pa.Y	Tauria	1816.	D co
81836 tafrica Bieb.	Taurian	5 A or	1	jl.au	Pa.Y	Tauria	1818.	D co
81837 pauciflora Lam.	few-flowered	£ A of	1	jl.au	Y	Spain	1820.	D co
S1838 EupathTu?n W. /ilicifolia Bieb.	Eupatorium	£ A	4	jl.au	Y	Casp. sho.	1803.	D co
81839 coarctata Poir.	compressed	£ A or	4	jls	Y	S. Europe	1816.	D co
81840 atrica Lam.	go/den-flowered	£ A or	1	jns	Y	Levant	1739.	D co
81841 pubtans L.	pubescent	£ A or	1	jns	L.Y	Levant	1739.	D p1
81842 crithmifolia Kit.	Saxnphire-lvd	£ A or	1	Jl.au	W	Hungary	1804.	D p1
81843 micrθ W.	small-flowered	£ A or	1	jno	Y	Levant	1805.	D p1
81844 tanacetiflora W.	Tansy-leaved	£ A	1	jl.au	Pk	Switzerl.	16j.	D co
81845 rbsca W.SFK.	roseate	£ A	1	jn.au	R	Hungary	1816.	D co
81846 distans Kit.	distant	£ A	3	Jl.au	W	Italy	1804	D co
ambigua Pott.								ALped.53.1
81847 Afillefolium L. 2 rubra	Milfoil	£	2	jn.o	W	Britain	past	D co
81848 dubia Us.	red-flowered	£	2	jn.o	W			Eng.bot.758
21849 montana & «.	doubtful	£	1	jn.au	W	Siberia	1820.	D co
21850 hetroptiflora Spr.	mountain	£	1	jn.au	W	Switzerl.	1819.	D co
81851 dentifera Dec.	variable-leaved	£	1	jn.au	W			
81842 intermedia Schi.	tooth-bearing	£	1	ln.au	W	S. France	1818.	D co
81853 polypylia Schl.	intermediate	£	1	ln.au	W	Switzerl.	1819.	D co
81854 strfcta 5cA.	many-leaved	£	1	ln.au	W	Switzerl.	1819.	D co
2ia05 bambatica W. % K.	strict	£	1	ln.au	W	Switzerl.	1820.	IX. co
81856 magna L.	Bannatic	£	1	ln.au	W	Hungary	1816.	O co
21857 lanata Sjw.	great	£	3	jnn	W	S. Europe	1683.	D co
21858 tomentosa L.	woolly	£	1	jl.au	W			
21859 microphylla W.	tomentose	£	2	my.o	Y	Britain	1804.	D co
21860 ochroleuca £A	small-leaved	£	1	Jls	W	Spain	1800.	D co
21861 odorata L.	yellowish-white	£	2	jl.s	Pa.Y			Bar. ic. 1114
21862 setacea Jo*.	sweet-scented	£	1	Jn.au	W	Spain	1804.	D co
21863 rrbilis L.	bristly	£	1	jn.au	W	Hungary	1805.	D p1
21864 ligustica All.	noble	£	2	jn.au	W	Germany	1640.	D co
21865 myrioptiflora W. en.	Lovage	£	4	jn.au	W	Italy	1791.	D co
21866 abrotenuifolia L.	myriad-leaved	£	1	W				Schk. han. 3.255
Balbia elongata W.	Southern w.-lvd	£	2	jn.au	Y	Levant	1739.	D p1

2388. TRPDAX L. TBIDAX. (*Tridaknos*, thrice-bitten; 3 deep divis. of rays of fl.) *Comp.Helian.* 1.—2.
81867 procumbens L. procumbent O un i jl.au Y Mexico 1804. S co

2389. ilME'LLUS L. AUELLUS. (Name of a plant mentioned by Virgil.) *Camp. Card. Vent.* 3.—4.
21868 ZychnUis L. Lychnitis JUi_Jpr. f jn.jl V. C. G. H. 1768. C p1 Jac. c. 10.1
21869 vill'sus Ph. villous £ A F 1 au.s Y Missouri 1811. D co
Sideranthus villosus Suit.

21870 Bpinthus Ph. spinulose £ A F 2 au.i Y Missouri 1811. D co
Sideranthus spinulbus Suit.

2390. STA'RKE^ W. STARKEA. (*Rev. M. Starke*, of Gros Tschirna, Silesia, cryptog.) *Comp.Card. Vern.* 1.
21871 umbellata IK. umbel-flowered £ ES un I,jn.jl Y Jamaica 1768. D l.p Lam. IL 682.2
Amillyu umbellatus L.

2391. ANDROMA'CH IA Kth. ANDROMACIIA. (*Andromachus*, phys. to Nero.) *Comp.Card.Vern.* 1.—9.
21872 ignaria Kth. touchwood £ ES un 1 jn.au Y S. Amer. 1824. C s.l H. & B. 2.112

2392. COLUMELLA'M Jac. COMIMBLIA. (*L. J. M. Columella*, flourished 42 yean A. C.) *Comp.Helian.* I.
21873 biennis Jac. biennial £ iQJ un 1 jn.jl Y C. G. H. 1821. S co Jac.sc. 3.301
Nestlera biennis Spr.

2393. ECLIPTA L. ECLIPTA. (*Ekleipo*. to be deficient; want of seed-crown or wing.) *Camp. Bel.* 6.—
21874. erecta L. upright O un 2 jl.s W America 1690. Di. el. 114.137
21875 prostrate L. prostrate & un 1 jl.s W E. Indies 1732. Di.el.113.138
21876 punctata L. dotted OI un * jl.au W W. Indies 1818. Jac am. pic. 197
21877 procumbens Mr. procumbent Je O un jl.au W Carolina 1818.
21878 latifolia L. broad-leaved OI un jl.au W E. Indies 182a
21879 undulata IV. waved Jk OI un , jl.au W E. Indies 1826. ■■■

2394. TETRA'NTHUS Swx. TETRANTHIS. (*Tetras, four, anthos, flower.*) *Camp. Eupatrea.* 1.—2.
21880 littoralis Sun. shore ArEScu ft jl.s W W. Indies 1820. D s.I

2395. MEY'EIM Schreb. MEYERA. (*Gottlieb An&revo Meyer*, a German botanist.) *Comp.Helian.* I.—B.
21881 sessilis Sun. sessile £ OS un 1 jl.au Y W, Indies ... D co

2396. CHRYSANTHE'LLUM Rchb. CHIRVSANTHELLUM. (Dim. of chrysanthemum^) *Comp. Helidn.* I.
21882 procumbens Rchb. procumblent O un 1 jn.jl Y W. Indies 1768. S co Sw, ob. 314.8.1
Vcrbstna mftica IV. Co'lte'a proc6mbens Spr.

2397. PTILOSTE^PHIUM Kth. PTILOSTEPHIUM. (*Ptilon, feather, stephos, crown; pappus.*) *Comp.Helian.* 2.
21883 coronaphititmn Kth. Buckhom-lvd O cu 1 jn.jl Y Mexico 1823. S co H.&B.4.387
21884 trifidum Kth. triad O cu 1 jn.jl Y Mexico 1823. S co H. & B. 4.388

2398. SIEGESBE'Ck/4 L. SIEOESBECKIA. (*J. George Siegesbeck*, M.D., a German bot) *Comp.Helian.* 6.
81885 orientalis L. Oriental 2 au.o Y India 1730. S co Schk. han. 3.256
21886 triangularis Cav. triangular 2 au.s Y Mexico 1825. S co Cav. ic. a 213
£1887 droseroides Swt. Sundew-like 2 au.s Y Mexico 1825. S co Bot. reg. 1061
jorulfensis^M.? heart-leaved 2 au.s Y Mexico 1826. S co
81888 cordifolia Kth. flosculose 2 jnjl Y Peru 1784. S co Her. st. 16
81889 flosculosa Ilerit. Iberian 1 au.s W Iberia 1818. S co Bux. c. 3.62

i 2399. VERBESNA L. VERBESIKA. (Altered from *Verbena*.) *Comp.Helidntheee.* 15.—

I. ALTEBNIFOLUE. — Leaves alternate.

i. HERBA^CE. — Herbaceous.

21891 alkta /,. wing-stalked £ IAI pr 2 my.o QW N. Amer. 1699. C Lp Bot. mag. 1716
21892 J2irgfnica£ Virginian ^ A pr 2 jl.s

CLASS XIX. ORDER 3.

SYNGENE'SIA FRUSTRA^XNEA.

			sun, anthos, fl:	resemblance.)	Comp. Heliun. 28.-35.
21933 annuus L.	annual	6 jn.o	Y	a Amer. 1516.	S co Ken. spec. 63
21934 Indicus L.	dwarf, Indian	3 jn.o	Pa.Y	Egypt 1785.	S co Tab. IC. 764
21935 tubastriformis Jac.	tube-formed	5 jlau	Y	Mexico 1799.	S co Jac. sc. 3.375
21936 multiflorus L.	many-flowered	6 au.o	Y	N. Amer. 1597.	D co Bot. mag. 227
2 plenus	double-flowered	6 au.o	Y	N. Amer. 1797.	D co
21937 tuberbus L.	tuberous	8 s.o	Y	Brazil 1617.	R co Jac. vin. 2.161
21938 angustifolius L.	narrow-leaved	3 8.0	Y	N. Amer. 1789.	D co Bot. mag. 2051
21939 macrophyllus W.	long-leaved	6 au.o	Y	N. Amer. 1800.	D co W. h. b. 70
21940 m6lbs JV.	soft	4 jl.o	Y	N. Amer. 1805.	D co
21941 decapetalus L.	ten-petaled	6 au.n	Y	N. Amer. 1759.	D p.l Rob. ic. 235
21942 prostratus W.	prostrated	2 il.s	Y	N. Amer. 1800.	D co
21943 strumbus k.	strumous	2	Y	N. Amer. 1710.	D p.l Boc. sic. 27.4
21944 altissimus L.	tallest	2	Y	M. Amer. 1731.	D co Jac. vin. 2.160
21945 giganteus L.	gigantic	2	Y	N. Amer. 1714.	D co M.h.6.7.66
21946 pauciflbrus Nut.	few-flowered	2	Y	Louisiana 1824.	D p.l
21947 fongitlius Ph.	long-leaved	2	Y	Georgia 1812.	D co
21948 diffusus A.M.	diffuse	2	Y	N. Amer. 1821.	D co Bot. mag. 2020
21949 linearis Cay.	linear	2	Y	Mexico 1823.	D co Bot. reg. 523
21950 trachelifdhus W.	Trachellium-lvd	2	Y	N. Amer. 1825.	D co
21951 cxc'e'Uus FT.	lofty	2	Y	Mexico 1820.	D co Cav. ic. 219
21952 missirkicus Lk.	Missouri	2	Y	Missouri 1821.	D co
21953 parviflbrus Kth.	small-flowered	2	Y	Mexico 1826.	D p.l H.&B.4.578
21954 coroiflctus it/A.	Cornus-leaved	2	Y	Mexico 1825.	D p.l
21955 lactiflbrus Pers.	lively-flowered	2	Y	N. Amer. 1810.	D p.l
21956 trilobatus f.*.	three-lobed	2	Y	Mexico 1824.	D co
21957 divaricatus Ph.	divaricate	2	Y	N. Amer. 1759.	D p.l M.h.6.7.66
21958 pubescens Vahl	pubescent	2	Y	N. Amer. 1795.	D co Bot. reg. 524
21959 Hooker [#] G. Don	Hooker's	2	Y	— — D co Bot. mag. 2778	
pubescens Hook.					
21960 villosus JV.	villous	2	Y	N. Amer. 1820.	D p.l
21961 atrorubens L.		2	Y	N. Amer. 1732.	D p.l Bot. reg. 508
2410. GYMINOLO ^X MIA Kth. GTMN. (Gymnos, naked, l&ma,				border; margin of grains.)	Comp. Hclidn. 3.-5.
21912 maculata B. R.	spotted	« pr	3 jn.il.	Y	W. Indies 1821.
21913 connate Spr.	connate	H Q o r	4 o.n.	Y	Hra/il 1827.
21961 triplinervia Kth.	triple-nerved	H Q o r	3 o.n.	Y	NewSpain 1325.
21965 pinnata Ven.	pinnated	3	aus.	Y	W. Indies 1821. D p.l Botreg.66J
21966 digitata H. K.	digitate,feaved	6	aus.	Y	M. h. 6. a 54
21967 laciniata L.	faged-leaved	6	JLs	Y	M. h. 6.6.53
21968 columnaris Ph.	columnnar	3	au.a	Y	Bot. mag. 1G01
21969 subtomentosa Ph.	rather downy	3	au.8	Y	1802. D co
21970 triloba L.	three-lobed	4	aus.	Y	1699. D co Bot reg. 525
21971 hlrta L.	great-hairy	2	jn.n	Y	1714. D p.l Sw. fl. ear. 82
21972 radula Ph.	rasp	2	jn.n	Y	1825. S p.l
% 21973 sertitipa Sw.	lateflowered	2	au	P	N. Amer. 1823. D co Sw. fl. gar. 4
21974 Newmant Hort.	Newman's	1\$	jl.au	Y	S. Amer. 1821. I) co
21975 napifolia Kth.	turnip-leaved	2	jl.au	Y	NewSpain 1824. D p.l
21976 fuigida H. K.	bright	3	jl.au	Y	1760. D p.l Bot mag. 1996
21977 lavigata Ph.	smooth	3	jl.au	Y	Carolina 1812. D co
21978 amp/Textifolia Jac.	stem-clasping	5	jl.au	Y	Louisiana 1793. S co Jac. ic.3.592
4 21979 purpurea L.	purple	5	jl.o	D.P	N. Amer. 1699. D p.l Hot mag. 2
* 2112.G AILLA'R'D/ii Lam.GAHLAHDIA(A. Gaillard de Marentonneau, an amateur bot)					Comp. Hclidn. 2.
21980 blcolor Lam.	two-colored	^ A or	2 jl.o	O	Carolina 1787. D co Bdt. mag. 1602
VirgiUa helbdes HerU.					
21981 aristata Ph.	awned	k A or	1 jl.o	Y	N. Amer. 1812. D co
2413. TITHO*O/N/J* Deaf. TITHIONIA. (Tithonus, favorite of Aurora; flower couleur d'aurore.) Conip.Hcl. 1.					
21982 tagetyibra Desf.	Marigold-flwd	* • pr	10 jl.o	O	VeraCrue 1818. C co Bot reg. 591
2414. CO'SMEA Cav.	COSMEA. (Kosmos, an ornament; flowers.)				Comp. Helidn. 7. —
21983 liitea B. M.	yc\oyr-Jiowercd	iQl pr	2 o.n	Y	Mexico 1811. ■ co Bot mag. 1689
21981 sulphurea W.	sulphur-wtf	O pr	2 jl.au	Y	Mexico 179y. S co Jac.ic.3.5U5
Core't>psis artemisia-fdHa Jac.					
21985 bipinnata W.	bipinnate-/IMI	iQJpr	2 jl.au	P	Mexico 1799. ■ Lp Bot mag. 1535
21986 parviflbra W.	small-flowered	O pr	2 jl.au	W	Mexico 1800. ■ co Jac. sc. 3.374
Coreopsis parviflbra Jac.					
21987 crithmifilia Kth.	Sampshire-leaved	iQlor	UJLn	Y	Mexico 1825. ■ co
21988 chrysanthemifilia J&A Chrysanth.-lvd	2 o.n.	Y	S. Amer. 1826. ■ co		
21989 tenuilla Kth.	delicate	iQJor	Y	Mexico 1824. ■ co	
Coreopsis odoratissima Cav.					
• 2415. COREOPSI'S Jac. COREOPSIS. {Koris, a bug, opsis, resemblance j seeds.) Comp. Uelidn. 20. —					
	I. COMPOSITE.—Leaves compound.				
21990/eruhcfblia Jac.	Ferula-leaved	Jt A Jor	3 o.n	Y	Mexico 1799. D l.p Bot. mag. 2059
21991 grandiflora Hogg	large-flowering	& A or	3 aus.	Y	N. Amer. 1826. S r.m Sw. fl. gar. 175
21992 verticillata L.	whorLfmsnf	5 A or	3 jl.o	Y	N. Amer. 1759. D p.l Bot. mag. 156
21933 tenuifolia Ehrh.	slender-leaved	£ A or	2 jl.au	Y	N. Amer. 1780. D co Pluk.m.544.4
21194 chrysanththaZ.,	golden-flwd	£ Ol or	2 jl.s	Y	W. Indies 1752. S co Pl. ic.Sai
21W5 tdpferis fc.	Mirec-winged	^ A or	6 au.o	Y	N. Amer. 1737. D P.I M.h.7.3.44
21i>96 Irivhosju-rooa Mi.	hairy-seeded	5 Q or	3 av s	V	NewJerscyl818. S r.m

21907 <i>cenifflia Mx.</i>	six-leaved	^ A or	4 au.o	Y	N. Amer. 1812.	D co	
21998 <i>filba L.</i>	white climbing	J E3 or	6 Jn.jl	W	Jamaica 1699.	D Lp	Her. par. 124
21999 <i>incisa AA</i>	<i>cuUeaved</i>	t CD or	6 s.d	Y	W. Indies ..	D co	Dot reg. 7
22000 <i>reptansi.</i>	creeping	T O or	6 jLs	Y	W. Indies 1792.	S co	Sin. spic. 22
					II. — Leaves divided.		
22001 <i>auriculata L.</i>	ear-leaved	3 or	6 au o	Y	N. Araer. 1699.	D p.l	Pluk.al.83.5
22002 <i>aurea/f.JL</i>	golden	3	3 Jujs	Y	N. Amer. 1785.	D p.l	
					III. — Leaves undivided.		
22003 <i>latifolia/..</i>	broad-leaved				N. Amer. 1786.	D co	
22004 <i>arctita Ph.</i>	<i>asharp-notched</i>				Carolina ..	D co	
22005 <i>lanceolate L.</i>	lanceolate				Carolina 1724.	S co	Bot cab. 821
22006 <i>crassiflba H. K.</i>	thick-leaved				Carolina 1786.	D p.l	
22007 <i>angustiflbia H. K.</i>	narrow-leaved				N. Araer. 1778.	D P1	
22008 <i>integriflbia Poir.</i>	entire-leaved				Carolina F.	D co	
22009 <i>dichotoma Mx.</i>	dichotomous				Caroline 1827.	D p.l	
24ia V1GUIEMM Kth.	VIGUIERA. (L. G. A. <i>Vtgukr</i> , of Montpelier, botanist)				Comp. <i>Helidn.</i> 2.—3.		
22010 <i>helianthoWes Kth.</i>	Sunflower-like	f ES or	3 jl.au	Y	Cuba 1825.	C p.l	H.&B.4.379
22011 <i>denteate Spr.</i>	toothed	f (£) or	3 jlau	Y	Mexico 1826.	C p.l	Cav.ic.3.820
	<i>Helianthus dentatus Cav.</i>						
„ 2417. ACTINO'MERIS. Nut.	ACTINOMBRIS. (Akitn, ray, meris, part.)				Comp. <i>Helidntheat.</i> 4		
22012 <i>squarrosa Nut.</i>	square rose	k A or	3 jl.au	Y	N. Amer. 164a	D p.l	Jac.vin. 2.110
	<i>Coreopsis alterniflbia Jac.</i> Verbesina <i>Core6psis Mx.</i>				"		
22013 <i>procumbens</i> Sf.		f A or	8 s.o	Y	N. Amer. 1766.	D p.l	
2^Whe7unthode8^ Sf.	Sunflower-like	* A or	3 jl.au	Y	N. Amer. 1825.	D p.l	
22015 <i>alata Suit.</i>	wing-stalked	5 A or	3 jl.au	Y	N. Amer. 1803.	D p.l	Cav. Ic. 3.260
	<i>Coreopsis alata Cav.</i>						
* 2418. CALLIOTSIS Rchb.	CALLIOPSIB. (<i>Kallistos</i> , most beautiful, <i>opsis</i> , eye; of fl.)				Comp. <i>Helidn.</i> 3.—4		
22016 <i>bicolor Rchb.</i>	two-colored	Od	S5imy.au	Y	Arkansa 1822.	S co	Bot. reg. 846
	<i>Coreopsis tinctibia Nut.</i>						
22017 <i>palmata Spr.</i>	palmate	f A or	3 my.au	Y	Louisiana 1523.	D co	
	<i>Coreopsis palmata Nut.</i>						
22018 <i>rbsea Spr.</i>	roseate	i A or	2 jl.au	R	N.Casanal778.	D co	
	<i>Core6psis rbsea Nut.</i>						
, 2419. SrMSJ/f Pers. SIM si A. (John Sims, M.D. F.R.S. F.L.S., editor of Bot Mag.)	Comp. <i>HeUdn.</i> 1.						
ft2£oi9amplexaclflis/Vrr». stem-clasping	£ A un	4 jl.au	Y180& D p.l
	<i>Core6psis amplexicaalis Cav.</i>						
2420 OSMITESZ. OSMITES. (<i>Osmē</i> , per ftimej strong smell of camphor.)	Comp. <i>Helidn.</i> a—5.						
22020 <i>camphorlnai.</i>	camphor-, c«ifcrf'UP'	1* apjl	W	C. G. if.	1794.	C fp	Sebal.90.2
22021 <i>dentata Thun.</i>	toothed	J J p.r	J4 apjl	W	C. G. H. 1820.	C l.p	
22022 <i>BeUidiastrum L.</i>	Bellidiastrum ft	il or	1 apjl	W	C. G. H. 1816.	C p.l	
„ 2421. ENCEUA IA Cav.	ENCELIA. (Egchelion, a little eel; seeds.)				Comp. <i>BeUanthe*</i> . 2.—S.		
22023 <i>canifscens Cav.</i>	canescent	■ L J p r	I jjl	O	Peru 1786.	C l.p	Bot. reg. 909
Pallasia Aalamifolia H. K.							
22024 <i>Aalimifolia Cav.</i>	Halimus-lvd	* L _J or	I jjl.au	Y	Mexico 1826.	C p.l	Cav. ic. 3.216
Pallasia A grandiflbra TV.							
2422. SCEROCA'R'PUS W. SCLKB. (Skleros, hard, <i>karpos</i> , fruit; covering of grains.)	Comp. <i>Helidn.</i> 1.						
22025 <i>africknus W.</i>	African	O un	2 jlau	Y	Guinea 1812.	S co	Jac. ic. 1176
2423. CULLITMJ/1 IUB. CULLUMIA. (Sir Thomas Cyllum, Bart F.R.& F.L.a)	Comp. <i>Hettdnthe*</i> . &						
22026 <i>ciliatis R. Br.</i>	ciliated	• L J or	2 my.jn	Y	C. G. H. 1774.	C p.l	Bur. af. 54.1
22027 <i>setbsa It. Br.</i>	bristled	* i J or	2 jn.au	Y	C. G. H. 178a	C Lp	
22028 <i>squarrosa R. Br.</i>	squarrose	• L <J or	2 jn.au	Y	C. G. H. 1786.	C Lp	Ac.haf.a5
2124. BERCKHE^Yi< Ehrh. III EtcKHEYA. (John Lefranc de Berekhey, Dutch bot)	Comp. <i>HeUdn.</i> 10.—XX						
22029 <i>cynaroides W.</i>	Artichoke-like	£ iAJ or	1 jjn	Y	C. G. H. 1789.	D Lp	
i:2(500bov^ta W.	smooth obovate	L J or	2 jn.au	Y	C. G. H. 1794.	C Lp	H.n.h.6.34.2
HixXil incana IV.	hoary	41 _J or	2 jl.au	Y	C. G. H. 1739.	C EpJac. ic. a 591	
22032 cuneata W.	vredG-leaved	* _J or	2 Jn.au	Y	C. G. H. 1812.	C l.p	Ac. haf. a 10
22033 palmata W.	palmated	ft	^J or	3 jn.au	C. G. H. 1800.	C Lp	Ac. haf.3.13
22034 pectinata IV.	pectinate	H _J or	2 jn.s	Y	C. G. H. 1818.	C p.l	
22035 grandiflbra W.	large-flowered	* J J or	2 jn.au	Y	C. G. H. 1812.	C p.l	Bot. mag. 1844
22036 uniflura W.	one-flowered	M J J or	3 jn.au	Y	C. G. H. 1815.	C co	Ac. haf. a 7
22037 cdrnua R. Br.	drooping.^twd	^ O l or	1 my.jl	Y	C. G. H. 1774.	S co	Meerb. ic. 1.40
22038 spinolssima W	spiniest	^ Iator	li jn.au	Y	C. G. H. 1821.	D p.l	Ac. haf. 12
10. DIDE'LTA Herit.	DIDELTA. (Dis, double, delta. Greek letter A: receptacle.)				Comp. <i>Helidn.</i> 2.		
22040 <i>spinoaum J. X.</i>	spinoaum	• U un	3 jn.jl	Y	C. G. H. 1774.	C l.p	Her. st. 28
		• U un	3 jn.jl	Y	C. G. H. 1774.	C Lp	Wen.ob.4.32
2426. GORTETR/J L.	GORTBRIA.	(David Gorter% Dutch prof, bot Harderwych.)			Comp. <i>HeUdn.</i> 1.—a		
22041 personata L.	mask	O or	ft jl.au	Y	C. G. H. 1774.	S co	Jac. c. 4.21.1
2427. GAZVNIA Gae.	GAZANIA.	(Gaza, riches; splendor of flowers.)			Comp. <i>Helidn.</i> 5.—		
22042 rtgens R. Br.	itifl*	#. L J or	4 my.s	Q. C. G. H. 1755.	*C p.l	• Bot. mag. 90	
GorterT* rlgens L.							
22043 heterophylrla W.	variable-leaved	iG iA or	0	Y	C. G. H. 1812.	W. h. b. 97	
22044 uniflora A 3/.	one-flowered	n. i J or		Y	C. G. H. 1816.	C p.	Bot. mag. 2870
22045 pavunia J. /M	peacock	jf iAl or		Y	C. G. H. 1804.	P.1	Bot. reg. 35
22046 subfutte R. Br.	awUeaved	£ iAI or		Y	C. G. H. 1792.	S co	
2428. CRYPTOSTE'MMA R.Br.	CRYPT.	(<i>Kryptos</i> hidden, <i>stemma</i> , crown; seeds.)			Ciwnn. H		
22047 calenthidicernia A^ Marigold Yurt	O or	1 jn.au	Y	C. G. H. 1752.	S		
22048 hy x chondrlacum A^ melanchofy	O or	ljl.au	Y	C. G. H. 1731.	S		
29049 mncinatum R.Br.	sxw-leaved	O or	1 jl.au	Y	C. G. H. 1794.	S co	

6429. ARCTOTHETC A <i>WtU.</i> ARCTOTUECA. (<i>Aktos, bear, theke, capsule; hairy seed.</i>) <i>Comp. Helidn.</i> 9.							
22050 <i>repens JvnL.</i> creeping	IAI or	1	Jl.au	Y	C. G. H.	1793.	D co Jac. sc. 3.306
Arctitis <i>scapigera Thun.</i>							
22051 <i>hirta Lk.</i> hairy	jfiA or	1	Jl.au	Y	C. G. H.	1820.	Dpi
2430. SPHENO'GYNE R. Br. SpnESOc. (<i>Sphen, wedge, gyne, female stigma.</i>) <i>Comp. Helian.</i> 9. —11.							
22052 <i>anthemoides R. Br.</i> Anthemis-like	O el	J	jl.s	■	C. G. W.	1774.	S co Bot mag 544
£2053/æniculacea Swt. Fennel-leaved	iQJor	1	jl.a	■	C. G. H.	1825.	S p1 Jac. BC. 2.156
Uremia/cernicul?fcca?r. Arctotis/cenicula Kicca Jac.					C. G. H.		
22054 crithmifolia R. Br. Samphire-lvd	n. i J cl	1	ap.au	y	C. G. H.	1768.	I p Bur. af. 65.1
22055 scaribsa R. Br. scarious	«L i J el	1	apau	Y	C. G. H.	1774.	Lp
22056 rtrotanifolia R. Br. Southern w-l vd tt.	[J cl	1	my.au	Y	C. G. H.	1789.	■ p
£2037 den tat a 11. Br. tooth-leaved	XL [J el	1	ap.jn	Y	C. G. H.	1787.	■ P Bur.af.64
22058 odorata R. Br. sweet-scented	«. i J el	1	ap.jn	Y	C. G. H.	1774.	I P
22059 pilifera Ker hairy	«. i J el	1	ap.jn	Y	"C. G. H.	1821.	P Bot. reg. 604
22060 /eucanthemcAdes RBr. Leucanth.-lk	O ■ i	jl.8	Y	CG* H.	1825.	■ P Jac. sc. 2.164	1
Arctotis /eucanthemcAdes Jac							
3431. ZCE^GEA L. Z<EOEA. (J.Zoega, M.D., published a Flora Islindica.) <i>Comp. Card.</i> 1.							
22061 Leptaura L. Leptaura	O un	\	jl.au	O	Levant	1779.	S co Jac. ic. 1.177
2432. LEITZEi* Dec. LEUZEA. (De Leuze, a friend of Decandolle'a.) <i>Comp. Carduiceaf.</i> 5. - 6.							
22062 conifera Dec. cone-bearing	^ A or	%	jns	F	S. Europe	1683.	D Lp An.mu. 16.14
Centaurea conifera L.							
22063 altaica Lk. Altaiyan	^ A or	SJns	P	Siberia	1822.	D co	
22064 salina Spr. * salt	k A or	1	jl.au	P?	Siberia	1817.	D co
Cirsium salinum Fis. Cnetaurea altaica Bieb.							
22065 australis Swt. southern	ik A or	1	jl.au	P	N. Holl	1821.	D e o Sw.fl.gar. ic
22066 carthamoides Dec. Carthamus-like^ A or	2	jl.au	P	Siberia	1616.	D co	
Cnicus carthamoide^ W.							
2433. RHAPCNTICA Dec. RHAPONTICA. (Bha, rhubarb, Ponticus, of Pontus) <i>Comp. Card.</i> 3.							
82067 scariosa Dec. scarious	st A or	2Jjl.au	P	Switzerl.	164a	D co Bot mag. 1752	
Centaurea Rhapuntica L.							
2 lyrata Dec. lyrate	■ A or	2	jl.au	P	Switzerl.	1819.	D co
22068 umflora Dec. one-flowered	■ A or	lijl.au	P	Siberia	1796.	D co Gm.sl.2.38	
Cnicus uniflora L.							
22069 Pallas/i. G. Don Pal I as'a Centaurea llhapuntica fl.	■ A or	2J jl.au	P	Switzerl.	1818.	D co	
+2434. CENTAURETA L. CENTA. (The Centaur Chiron, who first used medicinally.) <i>Comp. Card.</i> 158.—182.							
L CENTAURB^A.—Scales of calyx ciliated, sometimes at top spinosely mucronate.							
i. PECTINX\JE.—Scales of calyx pectinately ciliated, plumose.							
22070 phrgia L. dry	^ A or	H jn.o	P	Switzerl.	1633.	D co	Fl. dan. 520
2 ambigua Thomas ambiguous	■	■	■	■	■	■	
22071 salicifolia Bieb. Willow-leaved	■	■	■	■	■	■	
22072 pectinata L. pettinated	■	■	■	■	■	■	
22073 austriaca W. Austrian	■	■	■	■	■	■	
22074 uniflbra L. one-flowered	■	■	■	■	■	■	
22075 flosculosa Balb. flosculosus	■	■	■	■	■	■	
22076 nervosa IV. en. nerved	■	■	■	■	■	■	
22077 trichoccephala Bieb. hairy-headed	■	■	■	■	■	■	
22078 capillata L. hairy	■	■	■	■	■	■	
22079 rivularis Brot. rivulet	■	■	■	■	■	■	
22080 Avesopiflia Vahi Hyssop-leaved	■	■	■	■	■	■	
22081 flmfelia L. Flax-leaved	ilt A	1	jlau	P	Spain	1812.	C co Bar. ic. 80*
linariafolia Lam.							
23032 procumbens Balb. procumbent	* A or	\jn.au	P	S. Europe	1820.	D co	
22083 nigrata L. black Knapweed^i Aw	1	my.au	P	Britain pas.	D co	Eng.bot.278	
nlgicans Lk.							
ii. FIMBRIAIVJE.—Scales of calyx fringed, ciliated; flowers mostly red; disks for the most part green.							
22084 paniciflra Kita L. panied	■	■	■	■	■	■	
22085 Cineraria L. Cineraria	3	jl.au	P	Europe	1640.	S co	Jac. fl. 4.320
S2086 declinata Bieb. decimate	■	■	■	■	■	■	M.h.7.2b.20
22087 Icucophylla Bieb. white-leaved	2	jl.au	P	Caucasus	1821.	D co	
22088 cincrica Lam. grey	1 jnjl	P	Caucasus	1823.	D co		
22089 dealbata L. whitened	If	jl.au	P	Italy	1710.	D co	Jac. vin. 1.92
22090 Triumfetti All. Triumfetti's	1	jn.au	P	Caucasus	1804.	D co	
22091 nigrescens W. nigrescent	ljn.au	P	M. Ccnis	1820.	D co		
22092 transalp'ma Schl. transalpine	S jn.o	P	Hungary	1805.	D co		
S2093 vochinensis Bernh. Vochin	2	jn.au	P	Switzerl.	1619.	D co	
22094 bracteata Balb. bractate	2	jn.au	P	Austria	1817.	D co	
22095 pratensis T/uitf. meadow	2	jn.au	P	S. Europe	1817.	D co	
22096 astracanica Spi. Astracan	£	jn.au	P	France	1817.	D co	
22097 Marschalliana Spr. Marschall'i	If	jl.au	P	Astracan	1818.	D co	
sibirica Bieb. not L.							
22098 Scabiosa L. Greater Knapw.	t jn.au	P	Caucasus	1804.	D co	Eng. bot 56	
22099 coriacea Kit. H jn.jl	P	Hungary	1804.	D co	W.&K.2.195		
22100 trinervia W. three-nerved	2 jn.o	P	Podolia	1816.	D co		
22101 arenaria Bieb. sand	■	■	■	■	■	■	
22102 tnybyhcea Lam. Scoryz-Zoiwi	14	jl.s	P	S. Europe	1778.	D co	
22103 leucantha Pou. white-flowered	2	jl.s	W	S. France	1816.	D co	
22104 maculosa Lam. f'oticA-culyx	1	jlau	P	Siberia	1816.	D co	Gm. si. 2.44.1,2
22105 dissecta Ten. dissected	1	jlau	P	Naples	1823.	D co	
22106 Stachys L. Stcbs	1	jl.jl	R.Y	Austria?	1759.	D co	
22107 Hmbata Lk. fringed	2Jjn.au	P	Portugal	1818.	D co		
22108 sempervirens L. evergreen	li	jl.au	R.Y	Spain	1683.	C pl	Boc.si.39.3
iii. Scari&iUE.—Scales of calyx scarious, fringed.							
22109 macrocephala W. long-headed	■	■	■	■	■	■	
22110 pulcherrima^ very fair	■	■	■	■	■	■	
22111 concreta W. neat	■	■	■	■	■	■	
22112 atropurpurea Kit. dark-purple	■	■	■	■	■	■	
22113 calophylla IV. beautiful-lvd	■	■	■	■	■	■	
22114 neglecta A.* sulphurea L*ig.	neglected	■	■	■	■	■	

V. SERI'DIA. — *Scales of calyx at top palmately spinose.*

L RUBRIFLORJE.—Scales of calyx palmately spiny; flowers red or purple,							
22195 sonchiftlia L.	Sowthistle-lvd	1 au.o	Mediterr.	1780.	D co	Pluk. ph	39.1
22196 Seridis L.	"Endwlc-leaved	1 jn.o	Spain	1686.	D co	Pluk.aL38.1	
22197 cniænta W. en.	blood-leaved	1 jn.au		1816.	D co		
22198 romana L.	Roman	3 jl.s	'Some	1739.	S co	Bar. ic.	504
Zannbh Seb.							
22199 ferox De*f.	fierce	^t A.jor	2 Jls	Barbary	1790.	S p.l	Oesf. at. 2. 242
22200 sphaerocarpala L.	globe-headed	5 A or	2 jlau	& Europe	1683.	D co	Bot. mag. 2551
				Britain	Jersey.	D co	
22201 Isnardi ff.	IsnaniTs	31 A or	1 jlau	France	182U	S co	Eng. bot. 2256
22202 myacanthia Dec.	Mouse-thorn	£t A or	1 jn.o	S. France	1824.	D co	
22203 l'ouz'ini Dec.	Pouzin's	^ O or	2 jn.o	Candia	1691.	S co	Her. par. 189
22204 napifolia L.	Turnip-leaved	O or	3 Jls	S. Europe	1772.	S co	Boc. mu. 35.26
22205 aspera L.p	rough	O or	2 jn.o	Portugal	1804.	S co	
22206 polyacantha IV.	many-spined	O or	i jlau	Iberia	1818.	S co	
22207 iberica Trev.	Iberian	3t Q) or	2 jlau				

ii. *CiiRY9A'NTiiiE*.—*Scales of calyx palmate*, *s yellow*.

22203	solstitialis	W.	solstitial,	Barnaby's	T.	O or	2	jl.au		mid	fields.	S	co	Eng. bot. 243		
22209	melitensis	L.		Malta		O or	i	jl.au		Malta	1710.	S	co	Boc. sic. 35		
22210	sulphurea	W. en.		sulphur-colored		O or	I	jl.au		1815.	S	co			
22211	slcula	L.				gl _{cl} H _{an}	^	1*	jl.au	Sicily	1710.	S	co	Boc. sic. a 1		
22212	Stevenii	Bieb.		Steven's		CD	or	2	jl.au	Caucasus	1820.	S	co			
22213	xanthina	Spr.				yellow	3	A	or	2	jl.au		S	co		
22214	pallescens	Del.				pallcescent		O	or	2	jl.au	Egypt	1816.	S	co	
22215	Adam*	W.				Adam's		O	or	2	jl.au	Siberia	1804.	S	co	
22216	stramfnea	W.				straw-colored		O	or	i	jl.au	Egypt	1801.	S	co	
23217	coarctata	Viv.				compressed		O	or	i	jl.au	N. Africa	1827.	S	co	W.h.b.26

VI. CROCODY'LIUM. — Scales of calyx at top one-spined.

SALACTITES (Calc. wavy veins of brown white.) *Caves Caves 2*

Он - 1. Ден - В

SYNGENE-SIA NECfSSA^NRJA

*2436. <i>VJED&LZA</i> Jac.	WBDBBLIA. (<i>George Wolfgang Wedel</i> , a Ger. bot d. 1721.)	<i>Coinp.Helidn.</i> 6.-22.
22230 <i>hispida</i> Kth.	hispid	j ^f -Ajl or ijjn Y N. Spain 1819. D e o Bl! reg. 543
22231 <i>radicans</i> Ker	many-rayed	n. ^r or 3 ap.n Y Brazil 1820. C co Bot. reg. 610
23333 <i>acapulcnbis</i> Kth.	Acapulca	£ [SJ or 3 j ^f Y Acapulca 1826. C co
22233 <i>I helianthoides</i> Kth.	Sunflower-like	O "r 4 nd Y Guleto 1827. S co
22234 <i>bengalensis</i> lic/ib.	Bengal	4 O un ijnjl Y R Indies ... S co
22235 <i>carnosa</i> Rchb	shy-leaved	3 (E)cu. } inil Y W. Indies 1820. D e o

2437. *AhCVSA* Car. ALCINA. (*Fr. Ign. Alcina*, a Spanish Jesuit and naturalist.) *Cotnp.Helidn.* 1.
22236 *perfoliata* Cav. *perfoliate* iQJun 3 jl.au Y Mexico 1796. S co Cav. ic 1.15

2438. MILLE* ^R /f L.	MILLERIA. (<i>Philip Miller</i> , F.R.S., auth. <i>Gardener's Dictionary</i> .)	<i>Comp.Helidn. 9.</i>
22237 quinqueflora L.	five-flowered O un 2 jl.o Y VeraCruz1731. S co	<i>Cav. ic. 1.82</i>
22238 biflora L.	two-flowered O un 1 jl.o Y Campeachy1730. S co	<i>Mart. cen. 47.1</i>

2439. BALTIMORE' R J* L. BALTIMORA. (*Isord Baltimore*, proprietor of Maryland.) Comp. Helidn. 2.
 22239 recta L upright O un 2 jnjl Pa Y Vera Cruz 1699. S co Schk.han.3.26L.C
 22240 Alba Pers. q white O un 2 jnjl W S. Amer. 1800. S co

22440. <i>SILPHIUM L.</i>	<i>SILPHIUM</i>	[<i>Zalaph</i> , Arab., given to <i>Asafoetida</i> plant.]	<i>Comp. Heliidn.</i> 13.—15.
22241 <i>lachitum L.</i>	<i>fagged-leaved</i>	& A w 12 jl.s Y N. Amer. 1781. D & co. I, ffa. 1.8	
22242 <i>compositum Mx.</i>	<i>compound</i>	& A w 6 jl.s Y N. Amer. 1789. D co	
22243 <i>crebinthineum X.</i>	<i>Turpentine</i>	^ A w 6 au.s Y N. Amer. 17a5. D co Jac. vin. 1.43	
22244 <i>Rutaceum Wein.</i>	<i>glaucous</i>	£ A or 4 jl.o Y N. Amer. 1786. D co	
Coreopsis <i>latifolia</i> Mx.			
22245 <i>hiumilium Mx.</i>	<i>humble</i>	£ A or 2J jl.o Y N. Amer. 1786. D co	
<i>tomentosum Ph.</i>			
22246 <i>perfoliatum L.</i>	<i>perfoliate</i>	^ A w 7 ll.o Y N. Amer. 17G6. D co	

22247 conjunctum <i>W. en.</i>	conjoined	A w	7	jl.o	Y	N. Amer.	1765.	D	co	
*2248 connatum /.	connated	A w	6	jl.o	Y	N. Amer.	1765.	D	co	
*2249 Asteriscus <i>L.</i>	Asterieca	A w	5	jl.s	Y	N. Amer.	1732.	D	co	
*2250 trifolrttum <i>L.</i>	thrc-leaved	A w	6	jl.o	Y	N. Amer.	1755.	D	co	
*2251 ternatum <i>Retz.</i>	terwae-leaved	A w	4	jl.o	Y	N. Amer.	1806.	D	co	
*2252 atropurpureum Jfe/z.	dk-purp.-frArr	A w	4	jl.o	Y	N. Amer.	1812.	D	eo	
*2253 crythrocaelon <i>Bernh.</i>	red-stemmed	A w	4	jl.o	Y	N. Amer.	...	D	co	
2*41. GASTROCA'RPHA <i>D. Don</i> GASTBOC. (<i>Gaster, belly, karphe, chaff; on receptacle.</i>) Comp. T^b. 1.										
22254 runcinata <i>D. Don</i>	runcinate-wJ	f. Al or	3	aus	W	Chile	1827.	S	co	
8442. TTUXIS Br. TRIXIS. (<i>Trixos, triple; three-eved triangular capsule.</i>) Comp. MB. 2.-8.										
senecioloides Hook.	Groundsel-like	O P*	1	J au.s	W	Chile	1821.	S	co	
Leuceria senecioloides <i>D. Don</i>										
auriculkta Hook.	eared tt.	• pr	1	au.s	W	Brazil	1827.	C	l.p	
Perdicium brasiliense Dec.										
2243. POLY'MNif L.	POLYMNIA. (<i>Polymlnia, the name of one of the Muses.</i>) Comp. Widnthea?. 4.									
*2257 canadlnsis L.	Canadian	f A or	6	jl.au	LY	N. Amer.	1768.	D	eo	
*2258 Uvedalia L.	Uvedale's	^ A or	8	au.o	Y	N. Amer.	1699.	D	eo	
*2259 abyssinica L.	Abyssinian	f Ol or	4	ap.my	Y	Africa	1775.	S	co	
*2260 maculata Cau.	spotted	^ A or	6	aus	Y	Mexico	1824.	D	co	
2261 MONTANOM Lai.	MONTANOA." (<i>Montana, a Mexican patriot.</i>) Comp. Hclidnheat. 2. —									
*2261 grandiflbra Lai.	great-flowered	* i j un	4	n.d	Y	Mexico	1826.	C	co	
22262 tomentbsa Lai.	woolly	j ij n	4	n.d	Y	Mexico	1826.	C	co	
2245. CHRYSO'GONUM /., CHRVSOGONUM. (<i>Chrysos, gold, gnu, knee; fl. at joints.</i>) Comp. Helton. 1.										
22263 virginianum L.	Virginian	f, A pr	*	my.jn	Y	N. Amer.	...	D	p.l	
2246. A/ELAMPCTDIUM L. MELAMPODIUM. (<i>Melas, blaok,pous, foot; seed.</i>) Comp. HeMn. 5. — Ik										
22264 ameri ^c AnumL.	American	O un	ljau.o	W	VeraCruz	1733.	S	co	R.Hou.9.21	
ggfis hispidum Kth.	hispid	O un	ljl.a	W	N. Spain	1820.	S	co	tt&B. 4.399	
22266 longifolium Brou.	long-leaved	O un	1 jl.au	W	N. Spain	1820.	S	co		
22267 paludosum Kth.	marsh	O un	1 jl.au	W	S. Amer.	1817.	S	co		
ovatifl.lum Rchb.	Weddwi minor Horn.	Alchna ovalifolia Lag.	Dysodium divancatum Rich.							
humble	O un	lijn.o	W	Jamaica	1782.	S	co			
*2247. CHAPTA^LJ^ Ven.	CHAFTALIA. (<i>M. Chaptal, a famous French chemist.</i>) Comp. Labiatifi. 1.									
*JW9 tomentbsa Ven.	woolly	A pr	i	my.jn	W	N. Amer.	1806.	D	eo	
Tussil^go integrifolia Mx.										
*2248. CALE'NDULA L. MARIGOLD. (<i>Calcutta; first day of the month; flws monthly.</i>) Comp. Hel. 20.—34.										
22270 carnisols: L.	corn-field	o or	9	my.a	pl.Y	Euro ^a	1597.	s	ca	
*22271 sculca IV.en.	Sicilian	o or	1	my.s	D.Y	Sicily	1816.	f	co	
22272 stellKta Clev.	starred	O or	2	jns	O	Barbary	1795.	S	co	
22273 -fl	officinal common	O or	3	jns	O	S. Europe	1573.	S	co	
o ^ 2 plena	double-flciuered	O or	3	jns	S	co	
g274 fractai.	holy pale-flvod	O or	2	my.s	O	Levant	1731.	S	l.p	
gote	hoary	O or	lijn.au	Y	Barbary	1796.	S	lp	Desf.at.2.845	
	rainy, Small Cape	O or	1	jn.au	Y	C. G. H.	1699.	S	s.l	
	hybrid, Gr. Cape	O or	1	fl	w.P	C. G. H.	1752.	S	s.l	
*g78 mudicatilis L.	naked-stalked	O	1	jn.au	W	C. G. H.	1731.	S	s.l	
*g79? rainimifolia L.	Grass-leaved	tf	1	my.s	W	C. G. H.	1731.	C	p.l	
ZSK «) <i>Wa</i> L.	Goat-rush, bendin-stkd	tt	2	my.jn	W	C. G. H.	1774.	C	p.l	
2 flScida	flacid	tt	2	myjn	W	C. G. H.	1774.	C	lp	
viscfsa ft Jt.	viscous	tt	2	jns	O	C. G. H.	1790.	C	p.l	
**82 dentata B. R.	toothed	tt	1	cr	Umyjl	O	C. G. H.	1790.	C	Bot. rep. ic.
3 oppositifolia HI K.	opposite-leaved	tt	2	au	Y	C. G. H.	1774.	C	p.l	
fruticoso X.	shrubby	tt	2	fl	Y	C. G. H.	1752.	C	pi	
chrysanthemifolia Ven.	Chrysan.lyd	tt	2	fl	Y	C. G. H.	1790.	C	Bot. reg. 40	
arborcsens Jac.	arborescent	tt	1	uJ	Y	C. G. H.	1774.	C	Jac. ic.3.596	
subfruticoso Vahl	feufruticos	tt	1	mr.au	Y	C. G. H.	1823.	C	p.l	
denticulata Schous.	denticulated	n. i	2	d	Y	Barbary	1821.	C	p.l	
muricata Thun.	murirated	tt	2	d	Y	C. G. H.	...	C	p.l	
2*49. ARCTOrriS L. ABCTOTIS. (<i>Arctos, a bear, oüs, an ear; fruit shaggy.</i>) Comp. Helidn. 31. — 40.										
I. INDIVISIFOLJE. — Leaves undivided s shrubby.										
decurrent	tt	J or	1	jn.jl	W.R	C. G. H.	1794.	C	lp	
panicled	tt.i	Jor	1*	jn.jl	W.R	C. G. H.	1816.	C	lp	
roseate	tt	_ or	1	jl.s	Pk	C. G. H.	1793.	C	lp	
II. Fixxxn'FiWE. — Leaves lyrate or ptnatifid: shrubby.										
dark-circled	tt.	i J or	11	Jn.jl	W.P	C. G. H.	1812.	f	lp	
					P	C. G. H.	1739.			
narrow-leaved	tt.	J or	1	jl.s	O	C. G. H.	1795.			
ear-leaved	tt	J or	1	jn.au	Y	C. G. H.	1795.			
golden	tt	J or	1	jl.s	O	C. G. H.	1710.			
broad roigh-ZixtfO	tt	J or	3	jl.s	Y	C. G. H.	1710.			
arborescent	tt	J or	1*	jl.ju	W.pk	C. G. H.	1818.			
two-colored	tt	J or	1	jl.s	W.R	C. G. H.	1812.			
showy	tt	J or	1	jjn.au	Y	C. G. H.	1812.	P	Bot. mag. 2182	
taller	tt	J or	1	jn.au	Y.p	C. G. H.	1820.	P	Jac. sc. 2. 172	
spotted	tt	J or	1	jl.s	U my.au	W.o	C. G. H.	1812.	Bot. reg. 180	
copK:r-colouretln.	tt	J or	14	jn.au	Y.p	C. G. H.	1823.	P	Jac. sc. 2.176	
squarrosa Jac.	tt	J or	1*	jl.ju	O.p	C. G. H.	1825.	Lp	Jac. sc. 2.177	
smooth ish tt.	tt	J or	1f	jl.ju	Y.P	C. G. H.	1820.	Lp	Jacsc. 2.175	
III. INDIVAU. — Leaves undivided; stem caulescent; herbaceous.										
Bilvcr.leaved	f OJor	au	O	C. G. H.	1774.					
Aaggmg-stalked	iQJor	my.il	W.R	C. G. H.	1794.	I-P	Jac. sc. 2.163			
twiggy	O or	jl.au	Y	C. G. H.	1816.	co	Jac. sc. 3.307			
Tlaautain-lyd	f Alor	jnau	O	C. G. H.	1768.					
decumbent	f Alor	jl.s	Y	C. G. H.	1790.					
disdainful	yf OJspI	rnjwjl	Q.R	C. G. H.	1795.	I-P	Jac. sc. 3.381			
small thorn)-Ivd	iQJor	rnjwjl	O	C. G. H.	1795.	co	Jac. sc. 2.166			
						co	Jac. sc. 2.1(17)			

IV. LYRATIFOLIA. —	<i>pmnat'yid</i> ; <i>herbaceous</i> .						
82313 r&ptans <i>Jac.</i>	creeping	£ iAJ or	jI.i	W.O	C. G. H.	1795.	R P.I Jac. sc. 3.382
22314 grandiflbra <i>Jac.</i>	great-flowered	£ OJ or	mr.my	Pa.X	C. G. H.	1774.	S Lp
22315 glaucophylla <i>Jac.</i>	sca.green.lvd	£ iAl or	1 my.au	Y.p	C. G. H.	1794.	D Lp Jac. sc. 2.170
22316 revoluta Joe.	revolute	Jg IAJ or	1 my.au	Y	G. H.	1820.	C Lp Jac. sc. 2.173
82317 Cineraria <i>Jac.</i>	Cineraria	jE iAJ or	1) jn.an	Y.o	G. H.	1824.	E J Jac. BC. 2.174
	V ACAULIJ.E. — <i>Stemless</i> .						
22318 acafilis Z.	stemless	g iAlor	* ap.jl	Y.R	G. H.	1759.	R p.l Bot reg. 122
22319 tricolor L.	three-colored	g iA) or 1)	my jl W.a	G. H.	1794.	D Lp Bot reg. 131	
22320 undulata <i>Jac.</i>	v/SLVcleaved	£	1 ap.jn	O	G. H.	1795.	C p.l Jac.sc. 2.160
• 2450. OSTEOSPERMUM L. OSTEOSPERMUM. (<i>Osteon</i> , bone, <i>sperma</i> , seed; hardness of.) <i>Comp. Hel.</i> 13.—27.							
22321 corymbosum L.	corymbose	** iJ or	3 au	Y	G. H.	1822.	C Lp
22322 in cWm Th.	hoary	• I or	3 a	Y	C. H!	1815.	S p.l
22323 spinulosum L.	spiny	• I or	3 f.o	Y	C. G. H.	1700.	S l.p Com. h. 2.43
22324 spinescens L.	Spinescont	M \ or	3 mr.jn	Y	C. G. H.	1793.	C l.p Jac. sc. 3. 377
22325 pisiferum L.	pca-bearing"	M L or	4 mr.my	Y	C. G. H.	1757.	S l.p Bot cab. 470
22326 monilijsum L.	necklace-bearing	M i) or	3 jLau	Y	C. G. H.	1714.	S l.p Di. el. 68.79
J2J27 ihcifolium L.	Holly-leaved	ilu Jor	jl.au	Y	C. G. H.	1816.	I.p Bur. af. 172. 62
22358 calendulaceum L.	Marigold-like	• I.J or	2 jl.au	Y	C. G. H.	--	C p.l
Bldrs calcndulacea Thun.							
82329 perforatum L.	per foliate	—	3 jl.au	Y	C. G. H.	1820.	S l.p
22330 nivculin L.	snow-white	— or	3 jl.au	Y	C. G. H.	1816.	S l.p
22331 rfgidum H. K.	rigid	— or	3 ap.jl	Y	C. G. H.	1774.	C l.p
22332 caerileum H. K.	huc-flowered	—	3 jn.s	B	C. G. H.	1774.	C l.p Jac. Ic. 1.179
82333 polygaloidcs L.	Milkwort-like	—	3 jn.8	Y	C. G. H.	1759.	C Lp Fluk. m. 382
• 2451. 0THO'NN A L. RAGWORT. (<i>Qthone</i> , linen; downy clothing of original plant.) <i>Camp. Jacobic</i> 25.—39*							
I. DIVIDE. —Leaves divided; <i>shrubby</i> .							
22334 digitata Thun.	digitate	tt-i Jor	If jn.s	Y	C. G. H.	1824.	♦ I
22335 pectinata L.	pectinate-/?	i LJ or	3" ap.jn	Y	C. G. H.	1731.	C p.l Bot mag. 306
22336 jfthanasic Thun.	Athanasia-tfa?	■ L or	3 n.d	Y	C. G. H.	1795.	C p.l Jaesc. 27 242
22337 abrotanifltha L.	Southernw.-vrf	■ L or	3 ja.mr	Y	C. G. ft	1692.	C p.l Bot. reg. 108
II. —Leaves undivided; <i>shrubby</i> .							
J2338 retrofracta Jac.	retrofracted	■ L or	2 mr.au	—	C. G. T1	1812.	E L
22339 crassifolia L.	thick-leaved	■ J or	2 s.o	—	C. G. H.	1710.	C p.l Mil. ic. 2.245.2
22440 denticulate Thun.	tooth-leaved	■ J or	2 ap.jl	—	C. G. H.	1774.	C p.l Bot mag. 1979
22341 heterophylla Thun.	various-leaved	■ J or	2 ap.jl	—	C. G. H.	1812.	C tp
22342 arborscens L.	arborecent	• L-Jor	2 Jlau	—	C. G. H.	1723.	C P1 DLeL103.123
22343 frutescens L.	frutescent	• i Jor	2 jn.s	—	C. G. H.	1816.	C Lp
22344 cacalioides Thun.	Cacalia-like	AiA Jor	—	—	C. G. H.	1774.	D1..
22345 coronopifolia L.	Buckhorn-lvd	■ J or	—	—	C. G. H.	1731.	E Com. h. 2.70
22346 chciritoWz L.	Wallflower-leaved	Jk. J or	—	—	Barbary	1752.	C p.l Bot reg. 266
III. Drift&FbMA. —Leaves divided; <i>herbaceous</i> .							
22347 pinnata Thun.	pinnate	jc iAJ or	3 ap.jn	Y	C. G. H.	1759.	C Lp Bot. mag. 768
22348 pinnatifida Thun.	pinnatifid	£ IAJ or	1 jl.au	Y	C. G. H.	1823.	C Lp
22349 T&gktes L.	Marigold-feaimZ	O or	1 ap.ip	Y	C. G. H.	1823.	S co
22350 flabellifolia B. C.	fan-leaved	tt. uJ or	1 ap.jn	Y	C. G. H.	1821.	C co Bot cab. 728
IV. INDIVISJEFOLIJ.E. —Leaves undivided; <i>herbaceous</i> .							
22351 lfngua Thun.	tongue-leaved	tf iAJ or	8 my.s	Y	C. G. H.	1787.	D)p Jac. sc. 2.238
22352 fihecaulis Jac.	thread-stmd	AiA Jor	14 ap.my	Y	C. G. R.	1791.	D Lp Jac. sc. 2.241
22353 bulbosa L.	bulbous	A iAJ	2 my.jn	Y	C. G. H.	1774.	D l.p Brey. c. 66
22354 perforata Jac.	perforate	£ iAJ or	11 my.jl	Y	C. G. H.	1789.	D l.p Bot. mag. 1312
22355 parviflra L.	small-flowered	n. t-J or	2 jl.au	Y	C. G. H.	1704.	C p.l VoL nor. 226
V. A NcusTifulJiJ. —Leaves narrow.							
22356 nfcoldes L.	Heath-like	it. L-J or	2 jl.au	Y	C. G. H.	1815.	C l.p
22357 tenuissima L.	fine&i-leaved	tt. L-J or	1 ap.jl	Y	C. G. H.	1759.	C Lp Joc.sc 2.239
22358 /inifbla Thun.	Flax-leaved	£ iAI or	8 jn.s	Y	C. G. H.	1824.	C Lp
• 2452. HI'PPIA L. HIPPPXA. (<i>Hippox</i> , horse; horses fond of original plant, chickweed.) <i>Comp. 4nth.</i> 2.—5.							
22359 frutescens L.	frutescent	M J un	2 f.au	Y	C. G. H.	1710.	C p.l Bot mag. 1855
22360 integrifolia L.	entire-lvd	d un	J jl.au	Y	E. Indies	1777.	S Lp H. n.h.67.2
2453. SOLV^ Fl. per. SOLIVA. (<i>Salvator Soliva</i> , a Spanish phys. and bot.) <i>Camp. Anth.</i> 2.—9.							
22361 anthemifolia R. Br.	Chamomilelvd	O cu ft	jn.jl	Ap	N. Holl.	1818.	S co An.mu.61.1
Gymnostyles anthemiflia J.							
82362 stolonifera B. Br.	stoloniferous	O cu ft	jn.jl	Ap	Portugal	1816.	S co
Gymnostyles stolonifera Brot.							
2454. PSI A^DIA Jac. PBIADIA. (<i>Psias</i> , a dew-drop; glutinous exudation on leaves.) <i>Comp. Card. Fern.</i> 1.							
82363 glutinosa Jac.	glutinous	«L CD un	2 jn.au	Y	Mauritius	1796.	C Lp Jac. sc. 2.152
Conya glutinosa Lam.							
• 2455. ERIOC'PHALUS L. ERIOCBPHALVB. (<i>Erion</i> , wool, <i>kephale</i> , head; plant in seed.) <i>Comp. Anth.</i> ■							
82364 africanus L.	African	■■■ 6	j&mr	Y	C. G. H.	1732.	C p.l Bot mag. 833
82365 decussatus Burc.	crossed	■■■ Hor	4 rnrjh	Y	C. G. H.	1826.	C p.l
82366 purpuricus Bui c.	purple	■■■ Hor	4 mr.jn	P	C. G. H.	1816.	C p.l
22367 spinescens Burc.	spinescent	■■■ Hor	4 mr.au	Y	C. G. H.	1816.	C p.l
82368 racemosus L.	racemose	■■■ Hor	3 mr.ap	Y	CG. H.	1739.	C p.l
2456. MI'CROPUS L. MICROPUB. {Mikros, small, pou's, foot; like Leontopodium.) <i>Comp. Card. Fern.</i> 2.—4.							
22369 supinus L.	supine	O un	J jn.s	Y	S. Europe	1710.	S Lp Schk. han. 3.267
22370 erfctus L.	upright	O un	f jn.s	Y	& Europe	1683.	S Lp Loef. it. 1.3
2457. PARTHEWIUM Z. PARTHBNIUM. (<i>Parthenos</i> , virgin j sup. med. qual.) <i>Comp. Ambros.</i> 3.							
22371 Hysterophorus L.	Hysterophorus	O un	1 jl.o	W	Jamaica	1728.	S Lp Bot. mag. 2275
22372 incanum Kth.	hoary	O un	1 jl	W	Mexico	1824.	S co
22373 integriflbum L.	entire-leaved	^ A un	3 jn.o	W	Virginia	1661.	D p.l W.h.b.4
(Perhaps from <i>Yua</i> , a name used by the older botanist O							
82374 finnua L.							
22375 cUfaU W.	annual	O un	2 jl.au	W	S. Amer.	1768.	S Lp Schm. ic. 15
82376 frutescens L.	ciliated	O un	2 jl	W	N. Amer.	1820.	S p.l
	frutescent	A or	4 au	W	N. Amer.	1711.	C co Pink. al. 27.1

90SS⁹ ICA'RPH A J. ACICARPIA. (*Akis*, point, *karphe*, palea; appendage spiny.) *Catycirce*. 1.—3.
22377 spatulJita R. Br. spatulate £ QS cu ft . . . Brazil 1824. D p.l

CLASS XIX. ORDER 5.

SYNGENESIA SEGREGATA.

ELEPHANTOPUS L.	ELEPHANT'S FOOT. (<i>Elephas</i> , elephant, <i>pous</i> , foot.)	<i>Comp. Card. Ech Mn. 8.</i>
*379 caroliniknus W.	rough-leaved £ E un 1 jn.s W E. Indies 1695. S s.p R. mal. 1a 7	
*w80 nudicaulia Poir.	Carolina £ IAIUJI Iftjls R America 1732. S l.p Di. cl. 105.126	
*	naked-stemmed £ EJun 1 jls R Jamaica 1816. S co SL jam.3.4.1.150	
g382 m61lis Kth.	woolly £ E un 2 jls W W. Indies 1733. S l.p *	
** angustifl Mius Swx.	soft £ E un 2 jls W Caracas 1820. S co *	
384 nudiflbrus W.	narrow-leaved £ E un 2 jls W Jamaica 1822. S co SL Jam. 1.148.4	
385 spicatus J.	naked-flowered £ E un 2 jls W W. Indies 1818. S co	
	spicate £ HI Jnn 1 JLS W Jamaica 1816. S co SL jam.ii.16803.4	
2461. G^DERI L. CEDERA. (<i>George</i> (<i>Eder</i> , a Dane, prof. bot at Copenhagen.) <i>Comp. Card. Echinop.</i> 1.—3.		
>s86 prolifera Thun.	proliferous m J pr 3 my.jn Y C. G. H. 1789. C s.l Bot mag. 1637	

FLAVEHII A J. FLAVERIA. (<i>Flaws</i> , yellow: used in Chile for dyeing yellow.)	<i>Comp. Helidn. 1.</i>
7 Contrayerba Pers.	Contrayerba £ 0 1 TM 1 jl.« Y Peru 179*. S l.p Bot. mag. 2400
90 Milleria Contrayerba Cav.	
388 angustiflora Pers.	narrow-leaved O un 1 jls Y Mexico 1820. S co Cav. ic. 3.223
oon Milliera angustifolia Cav.	
*89 maritima Kth.	maritime B3 um 1 jl.fl Y Cuba 1823. S Lp
linearis Lag.	

STCFBE L. STOJBE. (<i>Stibas</i> , a bed of leaves; original plant BO used.)	<i>Comp. Card. Echinop. 4.—21</i>
390 ethMatica L.	Ethiopian tL LJ pr 2 au ... C. G. H. 1759. C p.l
391 erio.des L.	Hairy £ B — C. G. H. 1816. C p.l M.h. 7.18.12
392 cinerea Thun.	«.tipr ■ 3ls — C.G. H. 1784. C p.l Pluk. m. 297.1
393 reflexa Thun.	reflexed «.tjor ■ ju — C. G. H. 1816. C p.l
Serphium alopecuroides Lam.	

NAUENBU'RG/4 W. NAUBNBURGIA. (<i>John Samuel Nauenburg</i> , bot author.)	<i>Comp. Card. Echin. 1.</i>
239 trinervata W.	thrcr-nerved O un 3 jlau Y S. Amer. 1799. S l.p Sc. b. j. 1800.2.5
Brütera Contrayerva Spr.	

CASSINXA L.	CASSINXA. (<i>M. Henr Cassin</i> , a celebrated French bot.)	<i>Comp. Card. Fern. 5.—11.</i>
R. Br.	golden £ iAl or 1 jlau T N. Holl. 1803. D l.p Wen. c. 48	
Br.	long-leaved n. i.J or 2 apjl — N. Holl. 1822. C l.p	
bilis R. Br.	showy O or 6 jlau E N. Holl. 1818. S co Bot reg. 678	
affili « R Br.	allied «L v_J or 2 apjl .. N. Holl. 1820. C l.p	
*99 leptophylla i?. J3^.	slender-leaved tt-i_Jpr 2 jl.o N. Zeal. 1821. C co	

ANGIANTHUS WNL.	ANGIANTHUS. (<i>Aggot</i> , vessel, <i>anthos</i> , flower.)	<i>Comp. Card. Echin. I.</i>
atireus FBH.	golden £ iAJ or 1 jlau Y N. Holl. 1803. D l.p Bot. reg. 764	
Cassinia afirea H. K.		

SPHJERA'NTHUS Z.	SPHJEBANTHHS. (<i>Sphaira</i> , globe, <i>anthos</i> , flower.)	<i>Comp. Card. Echin. 3.—8.</i>
dicus L.	Indian £ El un 2 aud. B E. Indies 1691. C p.l Bur. zey. 94.3	
Africanus L.	African iQJun 1 jlau B C. G. H. 1759. S co Pluk. m. 108.7	
hirtus W.	hairy £ El un 2 Jlau B 1823. C co Lam. il. 718.1	

CHRONOPSJ.	GLOBB THISTLE. hedijehog.	aspect; heads.) <i>Comp. Card. Echin. 16.—</i>
Bphsrocculus L.	round-headed 5 Jlau + Austria 15-X. D CO Fl.gr. 923	
exultat ScAr.	exalted 6 Jlau + Austria 1817. S CO	
spinifl Z.	snxny-headed 4 Jlau Egypt 1597. D 1P M. h. 7.35.4	
yuebsu Bieb.	clammy 3 18 W Caucasus 1818. S CO	
Iffro L.	Ritro 3 Jlf. Europe 1570. O CO Bot mag. 932	
tenuiflbus P.*.	fine-leaved 2 Jls B Russia 1820. S CO	
JO ruthenicus Bieb.	Russian 3 JIB B Russia 1816. S CO	
humilis tif&	humble 1 Jl. Caucasus 1816. S CO	
virgatus iflw.	twiggy 2 Jl. S. Europe 1820. S CO	
polyraphalus Gou.	many-headed 2 Jl. S. Europe 1816. S CO	
fabricus W.	Taunan 4 18 B Tauria 1816. S CO	
meagre 1.	2 Jls W Spain 1729. S Bot. mag. 2109	
lanuginosus Lam.	woolly 2 Jl. Levant 1736. D l.p*	
pamculMus Jac.	panclcd 6 Jlau B Spain 1815. D l.p* Bot. reg. 356	
strictua B.	strict 3 <.au Pa.B Europe 1822. I. l.p Bot mag. 2457	
horrid Desf.	horrid 3 Jlau B Persia 1818. S CO	
persicus Fis.	« Jlau B Persia 1818. S	

ROLAN'DNRilRtb.	ROLAN. (<i>Daniel Rolander</i> , pupil of Lin., visited Surinam.)	<i>Comp. Catd. Ech. 1.—2.</i>
argentea Rib.	Bdvcr-leaved n - Lj or 1 Jl W W. Indies 1714. C l.p Sl.jam. 1.7.3	

L ⁸⁴⁷⁰ BRCTERI* W. BROTERA.	(<i>Felix Avelar Brotero</i> , prof. bot. Coimbra.)	<i>Comp. Card. Echinop. 1.</i>
28421 corymbosa »K.	corymbose A or 2 Jn.jl B & Europe 1640. D l.p M. h. 7.3117	
Onobrbma corymbosa Spr.		

-2471. GUNDE^L7<4 L.	GONDELIA. (<i>Andrew Gundelsheimer</i> , Ger. bot, its discoverer.)	<i>Comp. Card Frhi* i</i>
22422 TournefdrtilL.	Tournefors £ A cu Hjn.au L.G Levant 1739. D t-p MIL fear ..	

SKSS ² - EUXETNIA Cham.	EUXENIA. (Eus, beautiful, xerws, stranger.)	
^423 grata Cham.	pleasant «.LJpr "2 — Y Chile	R- ° ? * Echinop. 1. C p.l Hor. ph:b.6

CLASS XX. ORDER 1.

GYNA'NDRIA MONA'NDRIA.

	DISA.	(Meaning unknown.)	Orchideec.	12.—57.
	I. ELOVGAVE.— <i>Spur elongated.</i>			
22424 grandiflora L.	large-flowered A iAI spl 1 jl.au	S C. G. H. 1825. R B.p Bot.reg.926		
22423 comikta Swz.	horned ——————	Pa.B C. G. H. 1805. R s.p		
22426 draconis Swz.	dragon A iAJ cu	W.p C. G. H. 1823. R s.p		
22427 ferruginea Swz.	ferruginous A iAJ cu	Br C. G. H. 1820. R 8.p		
22428 prasinata Ji. R.	greenish-yellow A iAJ cu	G.R C. G. H. 1815. R s.p Bot. reg. 210		
22409 chrysostachya Swz.*	yellow-spiked A iAJ cu	Y C. G. H. ... R s.p		
22410 laceria Swz.	laceria A iAI or f i i	■ ■ G. H. 1826. R B.p		
22431 flexuosa Swz.	flexuous A IAJ or t jn-jl	■ ■ G. H. 1823. R s.p		
	II. BRBVES.— <i>Spur short.</i>			
22432 bracteata Swz.	bracted small-flwd & iAJ cu	jn.jl G C. G. H. 1818. R s.p Bot reg. 324		
22433 maculata L.	spotted A iAJ or 1 jn.jl	B C. G. H. 1816. R e.p		
22434 yraminifolia Banks	Grass-leaved AiAJspl	■ ■ B C. G. H. 1825. R s.p Jo. sc. 6.1.2		
22435 spatulata Sum.	spatula-shaped A iAlcu	jn.jl Fa.B C. G. H. 1805. R a.p Jo.sc.4.5.3		
2474. PTERYGODIUM Suit. PTERYGODIUM. (<i>Pterogodes</i> , wing-like: sepals.) <i>Orchidece.</i> 2.-6.				
22436 alatum Swz.	winged A LAJ or 1 jn.jl	■ C. G. H. 182L R Lp		
22437 volucrc Swz.	winged A iAl or 1 jn.jl	■ C. G. H. 1797. R Lp		
2475. DISPEMUS Swz.	DISPERIS. (<i>Dis</i> , two, <i>pera</i> , pouch: outer lateral segments of perianth.) <i>Orc/i.</i> 3.-5.			
22438 cucullata Kita Swz.	hooded AiAJor f jn.jl	P C. G. H. 1822. R Lp		
22439 capensis Swz.	Cape A iAJ or § my.au	S C. G. H. 1816. R Lp		
22440 secunda Sjuz.	side-flowering A iAJ or f jn.jl	P C. G. H. 1797. R Lp		
2476. CORY'CIUM Swz. Co VTCIUM. (<i>Kort/s</i> , a helmet; form of flower.) <i>OrcTildece.</i> 2.-4.				
22441 orobanchoides Swz.	Orobanche-like A iAJ or f jn.jl	Y C. G. H. 18*5. R Lp		
22442 criapum Sum.	curled A iAJ or } jn.au	Y C. G. H. 1825. 11 Lp Bux.c.3.11		
2477. SATY'R IUM L.	SATYRUM. (<i>Satyrus</i> , a satyr: from aphrodtsiacal properties.) <i>Orcldete.</i> 5.—19.			
22443 cucullatum Swz.	hooded A TAJ cu i jn.s	Pa. Y C. G. H. 1787. R s.p Bot reg. 416		
22444 carneum A. Jr.	feath-colored A iAJ el f jn.s	PK C. G. H. 1787. R B.p Bot mag. 1112		
22445 coriifolium Stuz.	leathery-leaved A LAJ or 1 p	Y.B C. G. H. 1820. R s.p Bot reg. 703		
22446 membranaceum Sua. membranaceous A iAJ or 1 jn.s	■ ■ C. G. H. 1826. R p.l			
22447 parviflorum W.	small-flowered A LAJ or jn.jl	■ ■ C. G. H. 1789. R p.l Jac. sc. 2.179		
2478. PLATANTHE^R A Rich. PLATANTHERA. (<i>Plotys</i> , broad, <i>anthera</i> , an anther.) <i>Orchide*</i> . 3.—11.				
22448 bifolia Rich.	two-lvd, <i>Butterfly</i> A A P' 1 m^y J^n w	Britain woods. R p.l Eng. bot. 22		
O'rechis bifolia L.				
22449 dilatata Hook.	dilated A A pr H au	W Canada 1823. R s.p Hook. ex. fl. 95		
22450 orbiculata Hook.	round-leaved A A cu 1 ap.my	G Canada 1823. R 8p Hook. cx.fl.lti		
2479. BONAM(E) W.	BONATEA. (<i>Bonato</i> , professor of botany at Padua.) <i>Orchideec.</i> 1.			
22451 apecibsa W.	showy £ G S e f 2 au	W C. G. H. 1820. D s.p Bot cab. 284		
2480. GYNADE*NIA Rich. GVMN. (<i>Gymnos</i> , naked, <i>aden</i> , gland: of stalk of pollen masses.) <i>Orch.</i> G.—7.				
22452 conopsea Rich.	gnat-like-/u/rf A A pr 1 jn.jl	P Britain me pas. R h.l Eng. bot 10		
O'rechis conopsea L.				
2 £ba	white-flowered A A pr	I jn.jl W Scotland moun. R p.l		
22453 odoratissima Rich.	sweet-scented A A pr	f jn.jl W Switzcr. 1824. R p.l Jac.au. 264		
O'rechis odoratissima L.	suavolens Vil.			
22454 angustifolia Spr.	narrow-leaved A A pr	f jn.jl W Caucasus 1819. R p.l		
O'rechis angustifolia Bieb.	iberica IV.			
22455 vfridis Rich.	green, Frog Orchis A A pr	I jn.jl G Britain me.pas. R Lp Eng. bot. 94		
fat^rium viride L.				
22456 albida Rich.	small whitish A A pr	4 jn.jl W Britain sun.hi. R Lp Eng. bot 505		
22457 cucullata Rich.	hooded A A pr	f jn.jl W Podolia ... R p.l		
*248L O'RCHIS L.	ORCMS. (<i>Orchis</i> , testis; frequent shape in roots.) <i>Orchidete.</i> SI.—54			
	I. TUBERbsiC — Roots tuberous, undivided.			
i. iNDIVTaE. — <i>Labellum undivided.</i>				
22458 papilionacea L.	papilionaceous A A or } jn.jl	Pa.P S. Europe 1788. R Lp		
22459 laxiflora Lam.	loose-flowered A A or % jn.jl	P France 1820. R p.l		
ensifolia Vil.				
	ii. Divps*. — <i>Labellum divided.</i>			
22460 mascula X.	male early A A or	■ ap.my P Britain woods. R i.p Eng. bot 631		
22461 simia Lam.	apic A A or	■ my.jl P England ch.pa. R p.l		
22462 Nicodemii Cyr.	Nicodemus's A A or	P Italy 1825. II p.l		
22463 parviflora Ten.	small-flowered A A or	■ P Italy 1825. R p.l		
22464 palfistris Jac.	marsh A A or	f ap.my P Switzcr. 1625. R p.l	Jac. ic. 181	
22465 mōrio L.	buffbon A A or	f my.jn P Britain mc.pas. R p.l	Eng. bot 2059	
22466 pauciflora Ten.	few-flowered A A or	f my.jl P Italy 1825. R p.l		
22467 longicornu Poir.	long-horned A A or	■ ap.my P Barbary 1815. R p.l	Bot reg. 202	
22468 pällcns L.	pale A A or	f my.jl Pa.Y Switzcr. 1825. R Lp	Jac. au. 45	
82469 provincialis Balb.	Provence A A or	■ ap.jn Pa.Y Switzcr. 1825. R p.l		
£2470 Cyrilii Ten.	Cyrillo's A A or	■ Pa.Y Naples 1825. R p.l	Cyr. ne. ic. in.	
22471 longibracteata Bio.	long-bracted A A or	■ P Sicily 1818. R p.l	Bot. reg. 357	
lloertiaha Loi.				
82472 variegata Jac.	variegated A Al or	ap.my Pa.P S. Europe! 1818. R L.p Bot. rog. 367		
22473 tnihtdris L.	military A A or	1 my.jn P Switzcr. 1825. K h.l Sw. fl. gar. 163		
S2474 Smithi Swt.	Smith's A A or	1 myjn P England ch.pa. R p.l Eng. bot 1673		
32475 RiVni Swt.	Rivinus*s A A or	} my.jn P Switzerl. 1S25. U p.l Sw. fl gir. 162		

- 2491. SERA^VFIAS L. SERAPIAS. (*Scrapis*, an Egyptian divinity.) *Orchideee*. 2.-4.
 82529 lingua L. *tonguc-lippcd* A iAI cu 1 my.jn Br S. Europe 1786. R Lp Bot. cab. 655
 92530 cordigera L. heart-bearing A iAI cu 1 jl.au Br S. Europe 1806. R Lp Bot rep. 475
2492. GOODYE^R R. Br. GOODYKRA. (*John Goodyer*, an obscure British botanist.) *Orchidear*. 5.-9.
 22531 r̄e^Δns R. Br. creeping k. A pr S jl.au W Scotland al.wo. D Lp Eng. bot 289
Neottia repens Swz.
- S2532 pubescens R. Br. pubescent tf EJpr f 2.d W N. Amer. 1802. D Lp Lind. col. 25
 22533 discolor B. R. two-colored £ E) p̄r 2 jnjl W S. Amer. J815. D Lp Bot. reg. 271
 22534 procera B. R. tall □ I jnjl W Nepal 1821. D Lp Hook. ex. ft. 39
 82535 tessellata Lod. tessellated □ I jnjl W N. Amer. 1821. D Lp Bot cab. 952
Neottia repens PA.
2493. DICTRIS Sm. DiURis. (*Dis*, double, *oura*, tail; lateral lobes of column.) *Orchideee*. 7.—
 I. FLAVIFLÖRJL — *Flowers yellow.*
 22536 afirea Sm. golden-flowered^Δ. iAJ el 1* ... Y N. S. W. 1810. R Lp Ex. bot 1.9
 22537 maculata R. Br. spotted A IAJ cu 1 mr.my Y N. Holl. 1823. R p.i Ex. bot. 1.30
 22538 sulphurica il.Zf/ A sulphur-ea/ora/A IA) cu 1 jn Pa.Y N.Holl. 1823. R p.i
 22539 pedunculata R. Br. pedunculate A iAI cu 1 f.au Y N. Holl 1826. R p.i
- II. PURFU^REAO-A'LHJS. — *Flowers purple or white.*
 22540 elongate Swt. elongated A IAJ cu 1 ap.my Pk N. Holl. 1822. R p.i
 22541 longifolia R. Br. long-leaved A LAJ cu 1 jl Pk N. Holl. 1824. R p.i
 aa5mkbAR.Br. white □ 1 jnjl W N.Holl. — R p.i
- 2*94. PONTHIETVitf R. Br. Powni. (*De Ponthieu*, Fr. W.I. merch. who sent plants to Sir J. Banks) *Orc. 2.*
 22543 glandulosa R. Br. glandulosus jf EJ cu 1 ja.mr G W. Indies 1800. D Lp Bot. mag. 842
 22544 petioUita Undl. stalked £ 23 cu 1 au Br St.Vincent 1822. D Lp Bot. reg. 760
2495. PELE'XIA Pott. PELEXIA. (Pelex, a helmet; large.) *OrcMdete*. 1.—3.
 23545 spiranthoides B. R. Spiranthes-like ^ □ J ap W W. Indies 1823. D Lp Bot. reg. 985
Neottia adinata Swz.
- **2496. NEO'TTIA Siwz. NEOTTIA. (*Ncottia*, a bird's nest \$ interwoven fibres of roots.) *Orcldex*. 14.
 I. NKO'TTIA. — *Spike straight.*
 22546 nidus avis Swz. bird* nest £ A cu 1 my Br Britain ch.wo. R Lp Eng. bot 48
 22547 picta Sims painted □ pr ap.jn W Trinidad 1805. D s.p Bot. mag. 1562
Spiranthes picta Lindl.
 22548 grandiflora Hook. great-flowered O or ap.jl G Rio Jan. 1823. f.s.p Bot mag. 2730
 22549 elata Swx. tall * GS pr ap.jn W W. Indies 1790. f.s.p Bot. mag. 2026
Spiranthes elata Undl.
 S25. Wplantaginac//00ft. Plantain/vtf □ iAJ pr jnjl R Nepal 1824. f.pi Hook.ex. fl. 226
 22551 apitfla Hook. leafless Ed pr jnjl Br.R.a Trinidad 1826. f.f.p Bot mag. 27*J7
- II. SPIRANTHIE. — *Spike spiral.*
 22552 pudica Suit. modest & E) p r i n.d Pk China 1819. D s.p Lind col. 30
Spiranthes pudica Lindl.
 £3553 australis Lindl. southern ff [S] pr R N. Holl. 182a
 22554 bicolor Ker two-colored ff E1 P' □ p.1
Spiranthes bicolor Lindl.
 22555 cernua IV. drooping/Jtwf □ 1 jl W N. Amer. 1796. D Lp Bot mag. 1568 .
Spiranthes cernua Rich.
 22556 tortilis Swx. twisted ED pr *Jn.jl W W. Indies 1822. D p.i
 22557 spirali H. K. spiral, *Ladies' traces* — Apf □ W Britain mc.pa.D Lp
Spiranthes oestivalis Rich. CIPhrys spirali E. B. p Eng. bot 541
 22558 autumnalis Swz. autumnal ^ Apr I W Europe 1800. D p.i
 O'phrys spiral is a L
 22559 estivialis Dec summer jfc A P* i W N. Amer. 1822. D p.i Bar. fl. 35.2
 dtritis Ph. OYⁿ¹tl⁸ spiralis y L. W
2497. STENORHY'NCHUS Rich. STENORIIVCVNUS. (*Stenos*, narrow, *rhynchos*, beak; pointedstig.) □ □ □
 22560 specibus Rich. showy £ El cl 1 ap.jn S W. Indies 1790. D s.p Bot. mag. 1374
Neottia specittia Jac.
- 22561 orchidEB Rich. Orchis-like jg El el li my F Jamaica 1806. D s.p Bot. mag 1036
2498. CRYPTO'STYLIS R. Br. CRYPTom LIS. (*Kryptos*, hidden, stu^Δ □ □ □) *Orchide**. 2.-3.
 22562 crēcta R. Br. erect £ El cu 1 jnjl D.Br N. Holl. 1824. D Lp
 22563 longifolia R. Br. long-leaved £ El cu 1 jnjl D.Br N. Holl. 1822. D Lp Lab. n. h. 2. 212
2499. ORTHO'CERAS R. Br. OBTHOCERAS. (*Orthos*, straight, *keras*, horn j outer sepals.) *Orchidea*. 1.
 22564 striatum JZ J'r. striated A IAJ cu ft . . . N.Holl. 1826. D Lp
- 2*00. THELY'MITRA Forst. THELYMITRA. (*Wielys*, woman, *miltra*, cap; hooded fructif.) *Orc. 8.-10.*
 22565 fxioldes Sun. Ixia.like A IAJ cu ap.jn B N. Holl. 1810. R Lp Sm. ex. bot 1.29
 2Jj^Δ media R. Br. mediate A LAJ cu ap.jn B N. Holl. 1820. R Lp
 &2j^Δ pauciflora R. Br. few-flowered A IA) cu ap.jn B N. Holl. 1820. R Lp
 22038 angustiflora R. Br. narrow-leaved A IAJ cu ap.jn B N. Holl. 1825. R Lp
 22569 cárnea R. Br. tlech-colored A IAJ cu ap.jn F N. Holl. 1820. R Lp
 22170 venosa R Br. veiny A IAJ cu ap.jn B N. Holl. 1826. R Lp
 22571 Forsterii Swz. Forster's " A IAJ cu ap.jn — N. Zeal. 1N24. R Lp
 22572 longifolia Forst. long-leaved A IAJ cu ap.jn — V. D. L. 1824. R Lp
- £501. LI'STERjf R.Br. TWAT BLADE. (*Martin Lister*, M D., a celeb. Enplish phys. and nat.) *Orcldex*. 9.
 22573 ováta li. Br. oxatc-leavcd ^ A cu 1 my.jn G Britain woods. R Lp Eng.Jtot. 154S
Neottia 1 atifolia Rich.
- 22574 cordata R. Br. heart-leaved ^ A cu I jn.jl Br Britain xnoi. h. R Lp Eng. bot 358
2502. CALOCHILUS R.5r. CALOCHILUS. (*Kalos*, beautiful, cheUos, lip.) *Orchidear*. 2.
 29575 campestris R. Br. field A tAl or f ... Br N. Holl. 1824. D Lp
 S2576 paludus Jl. Br. marsh A IAJ or \ ... Br N. Holl. 1823. D Lp
2503. ARETHU^U L. ARETHISA. (*Arethusa*, nymph of Diana's, changed into fountain; habitat.) *Orc. <>.**
 22577 bulbosa H. K. bulbous A -Al el i my.jn Pk.v N. Amer. ... R Lp Bot. mag. 2204
 22578 plicata Andr. plaited A El or 1 jl.au ... E. Indies 1805. R p.i Bot. rep. 321

2304. CALOPOXON R. Br. CALOPOGOV. (*M*, beautiful, pqgwi, beard; fringe of lip.) *Orchideee*. 1.
 22579 pulchellus H. K. pretty A-Al el 1J jl.au P N. Amer. 1771. R l.p Bot mag. XI6
Linodbrum tuberdsum B. M. pulchellum *Sal.*
2505. POGCTNIA R. Br. POONIA. (*Pogon*, bean!; fringed nectary lip.) *Orchidete. & — 5.*
 £2580 ophioglossoWesJRff. Ophiogl.-like AiAJel 1 jn.jl Pk N. Amer. 1816. R l.p Bot reg. 148
Arethftsa ophioglossoides L.
- 22381 divancita R. Br. divaricated **A A pr** f jn.jl Pk N. Amer. 1787. Dip Lam.il. 729.3
 22382 pendula Spr. pendulous **A A pr** I au Pk N. Amer. 1824. D l.p BoL reg. 908
Arethftsa pendula IV.
2506. LYPERA'NTHUS R. Br. LYPERANTIIITS. (*Lype*, sadness, *anthos*, flower; **hue**.) *Orchide**. 3.
 22583 suaveolens R. Br. sweet-scented A IAI cu ... jn.jl D.Br N. Holl. 1822. R p.1
 22584 ellipticus R. Br. elliptic A IAI cu ... D.Br N. Holl. 1824. R p.1
 22585 nigricans R. Br. blackish A IAI cu ... D.Br N. Holl. 1824. R p.1
- 2507. SPIPA'CTIS Rich. EPIPACTIS. (*Epipegnuo*, to ~~plate~~; effect on milk.) *Orchide**. & —
 22586 latifolia Swz. broad-leaved ^ A or 1* jl.au Britain m. wo. D l.p Lng. bot 269
Setifnas latifolia L.
- M 22587 palustris Swz. marsh + A or | jl.au P Britain mar. D l.p Eng.bot.270
2508. CEPHALANTHE'RA Rich. CEPHALANTHERA. (*Kephale*, a head, *anthera*, an anther.) *Orchideg*. 3.
 22588 pollens Rich. pale ^ AA cor 1 jn W Britain ... D l.p Eng.bot.S71
Epiptaxis pollens Smn. Sennkiptis grandiflora L.
- 22589 ensifolia Rich. sword-leaved ^ A or 1 jn W Britain m.wo. D l.p Eng. bot. 491
Epiptaxis ensifolia Swz.
- 22590 rufra Rchb. red k A or 1 Jn.jl P Britain m. wo. D l.p Eng. bot. 437
Swz. SerXpia<<Ilkbra Z
2309. CALEFY: R. Br. CALEVA. (*George Caley*, sometime superint of bot. gard. at St. Vincent.) *Orchidece*. 2.
 22591 major A 5r. larger AiAJpr 1 Jn.jl G.Br N. S.W. 1810. D l.p
 *%m minor R.Br. smaller AiAJpr § jn.jl G.Br N. Holl. 1822. R p.1
2510. PRASOPHYLLUM if. 2fc. P~~«~~ASOPHILLUM. (*Prason*, leek, *phyllon*, leaf; similarity.) *Orchidee*. 7.—12.
 ^ I. INTEGHA. — *Lateral segments of column entire: germen obtongylindrical from No. 22596. obovate.*
- 22598 elktumfl 2*, tail ^ A or 2 ... {j-go 1fe4. R l.p
 22599 flfrumR.Br. yellow S A or 1 Y. g. Hoi 1. 1825. R l.p
 22595 stratum R. Br. striated A IAI or 1 ... g. Holl. 1824. R l.p
 22598 fuscum R. Br. brown A IAI or 1 ... Br N. Holl. 1824. R l.p
 22597 pitens R. Br. spreading A-Al or 1 ... N. Holl. 182a R l.p
- II. BI'FIDA — *Lateral segments of column bifid.*
 22598 rMum R. Br. rufous AiAJor 1 ... Ru N. Holl. 1824. R l.p
 22599 fimbriatum R. Br. fringed A LAI or 1 ... N. Holl. 1824. R l.p
2511. MICROPTISIJZ. Br. MICROTIS. (*Mikros*, small, ous, ear; app. to anther.) *Orchidece*. 2. - 6.
 22600 parvtflbra R. Br. smaU-flowered A IAI cu 1 my.jn G.w N. Holl. 1824. R l.p
 22601 llba R. Br. white A IAI cu 1 my.Jn W N. Holl. 1826. R l.p
2512. ACIA'NTHUS R. Br. ACIANTHUS. (*Acis*, point, *anthos*, flower; bristly tins.) *Orchidece*. 3.
 23602 fomic&tus R. Br. arched A IAI or 1 my.jn Br N. Holl. 1822. R l.p
 22603 exscrtus R. Br. exserted A IAI or 1 my.jn Br N. Holl. 1822. R l.p
 22604 caudatus R. Br. tailed AiAlor 1 my.jn Br N. Holl. 1324. II l.p
2513. CYRTO'STYLIS R. Br. CYHTOSTYLIS. (*Kyrtos*, convex, *stylos*, style.) *Orchideee*. 1.—SL
 22605 reniformis R. Br. reniform Wwt A IAI cu \ my.jn ... N. Hull. 1823. D p.1
- 2514. CHILOGLCTTIS R. Br. CHILOGLOTTIS. (*Cheilos*, lip, *gloita*, tongue; app. to lip.) *OrcMde**. 1.
 «606diphylla i.^r. two-leaved. A IAI or i ... R N. Holl. ... R p.1 Bauer n. h.8
- oo??^5- ERIOCHRUS R. Br. ERIOCHILUS. (*Erion*, wool, *cheilos*, lip; disk of labellum pubescent.) *Orch*. 1.
 22607 autumnalis R. Br. autumnal AiAlor 1 n.d R N. Holl. 1823. R p.1 Lab.n. h.2.211.2
-Epipactis cucullata Lab.
2516. CALADE^NIA R. Br. CALADENIA. (*Kalos*, beautiful, *aden*, gland; disk of labellum.) *Orch. 5.-15.*
 22608 &ba R.Br. white A IAI or - jlau W N. Holl. 1810 R p.1
 22609 cSrnea R. Br. tosh-colored A IAI or - - F N. Holl. 1826. R p.1
 22610 caerulca R. Br. blue A IAI or B N. Holl. 1824. R p.1
 f26H alata R.Br. winged A IAI or ... jn.jl B N. Holl. 1823. R p.1
 22612te8taceaAfr. testaceous AiAlor ... jn.jl - N. Holl. 1824. R p.1
- O-2517. GLOSSO^DIA iZ. J7r. GLOSSODIA. (*Glossa*, tongue, *eidos*, like: append, within flower.) *Orch*. 2.
 *2613 major R. Br. larger A IAI pr ... jn.au B N. Holl. 1810. R p.1
 22614 nrinor R. Br. smaller A IAI pr ... jn.au B N. Holl. 1824. R p.1
2518. PTERO'S^YLYS R.Br. PTEEOTVLIB. (*Pteron*, wing, *stylas*, style; col. at top winged.) *Orch*. 11.—17.
 — *Stems leafy; appendix of labellum at top pencilled.*
- 22617 outtisa R. Br. large, powdered AiAlor . . . Pa.Y N. H6U. 1824. R p.1
 wsdexed A IAI or ... jl.au Pa.Y N. Holl. 1826. R p.1
 obtuse A IAI or ... jl.au Pa.Y N. Holl. 1810. R p.1
- oo?^10 imm. x1/10 TM TM* - *Scape naked.* AiAlor ... jl.au Pa.Y N. Holl. 1824. R p.1
 22618 cuculata A Br. hooded *radical leaves stellate; appendices of labellum at top pencilled.*
 &b19nhtans R.Br. nodding A IAI or ... jn.jl PaiYV.D. L. 182J. R p.1
- 22620 acuminita R. Br. acuminate A IAI or ... Pa.Y N. Holl. 1883. R p.1
 22621 cfrito R. Br. shori'lipped A IAI or ... my.jn Pa.Y N. Holl. 1826. R p.1
 22622 ophiogWSSa R. Br. Adder's tongue A IAI or my.jn Pa.Y N. Holl. 1822. R p.1
 22623 concinnat?fr. -neat AiAlor . . . Pa.Y N. Holl. 1826. R p.1
- III. SQITAMA'TO. — *Stems scaly; appendix of labellum at top naked.* Pa.Y N. Holl. 1824. R p.1
- 22624 longifblia R. Br. long-leaved A IAI or ... jUu . . . N. Holl. 1823. R p.1
 22625 gibbosa R. Br. gibbosus A IAI or . . . N. Holl. 1824. R p.1
2519. CORYSA'NTHES R. Br. CORYSANTHES. (*Korys*, helmet, *anthos*, fl.; helmet of fl. large) *Orrh* ^
 22626 bicalcarata R. Br. two-spurred A LAJ or ... jl.au D.Br N. Holl. 1823. R p.1
 22627 unguiculata R. Br. unguiculate A LAJ or ... jl.au D.Br N. Holl. 1822. R p.1
 ^QBB fimbriata H. Br. fringed AiAlor ... jl.au D.Br N. Holl. 1821. R p.1 Par. Ion. 83?

- *2520. CORALLORRHPZA R.Br. CORAIXOIIIIIZA. (*Korallion*, coral, *rhixa*, root; branched.) *Orch.* 3.-4.
22689 imiflta R.Br. inborn A A cu *JH^G Scotland sc.wo. D Lp Eng. bot. 1554
25260 multifl.^{ra} Nut. many-flowered * A <m 3 jnjl G N. Amer. 182*. D p.l Hook. ex. fl. 174
22631 odontorrhiza Spr. tooth-rooted A A TM 4 J^J N. Amer. 1824. D p.l
0^xphrys Corallorrhiza R.Br.

2521. RODRIGUETZ/il Fl.por. RODBIUEZIA. (*Emanuel Rodriguez*, a Spanish phys. and bot.) *Orch.* 1.—2.
SSG33 se^cunda Kth. Bide-flowermg ^ (ZSe^l | o R S. Amer. 1818. D p.r.w Hook. ex. fl. 129
Pleurothallis coccinea Hook.

2522. GOMVZA R.Br. GOMEZA. (Senor Gomez, a Spanish apothecary.) *Orchidear.* 1.
22633 rectirva R.Br. recurved Elel \$ my.jn Y Brazil 1824. D p.r.w Bot. mag. 1748

2523. CYMBIDIUM Swz. (CYMBIDIUM.) (Kymbe, a boat i form of labellum.) *Orchidece.* 8.-58.
22634 tripterum Swz. three-winged./Fwi/tfjpf El cu *jnjl W Jamaica 1790. D p.r.w Sm.ic. pic 14
22635 albidum Swz. Alo-leafed jg ES or 1 my.jn Rr E. Indies 1789. D l.p Bot. mag. SB7
22636 ensifolium Swz. sword-leaved El fra 2 zjn.o Br China 1780. D l.p Bot. mag. 1751
22637 sinense W. Chinese £ El fra + n. Hr China 1793. I) l.p Hot. mag. 888
22638 lancifolium Hook. lance-leaved £ Eld 2 my Y.R E. Indies 1822. D t.p Hook. ex. fl. 51
22639 fiphiolium Lindl. Xiphium-lvd l<Elpr | my.au G China 1814. D l.p Bot. reg. 529
22640 suave R.Br. sweet jg El or = ... N. Holl. 1824. D l.p
22641 reflexum It.Br. reflexed jg El or = ... N. Holl. 1824. D l.p

2524. CIRRH^W B.R. CIRRIUB*. (Kirrhos, yellow; flowers.) *Orchideee.* 1.
22642 depéndens B.R. depending f jL Y.G China 1822. D p.r.w Bot. cab. 936
Cymbidium depéndens B.C.

2525. SARCOCHVLUS KBr. SARCOCHILUS. (Sax, flesh, cheilos, a lip.) *Orchideee.* 1.
22643 falcatus R.Br. falcate £ iAI or = jn.au ... N. Holl. 1821. D p.r.w

2526. BRA'SS74 R.Br. BBASBIA. (*William Brass*, who collected plants on W. coast of Africa.) *Orch.* 2.
22644 maculata R.Br. Sjtottcd Jliud ^ LS]fl 1 jnjl Y.R Jamaica 180b*. J)p.r.w Bot. mag. 1691
22645 caudata Lindl. long-tailed ^ (22 el 1 jnjl G. v.a W. Indies 1823. D p.r.w Bot. reg. 832
Epidendrum caudatum L.

2527. LISSOCHRLUS it. Br. LiacosciLUS. (Lissos, smooth, chculos, lip; Up of flower.) *Orchidea.* 2.
22646 speciosus 21. Br. showy tf G3 spl 2 my.jn Y C. G. H. 1818. D l.p Lind. col. 31
22647 luteus Swt. yellow T El or 1 np.jn Y C. G. H. 1822. D l.p

* 2528. DIPO'DIUM R.Br. DIPODIUM. (Dis, two, potts, foot; threads of pollen masses.) *Orchideee.* 1.—2
22648 punctatum R.Br. spotted ± El or l jn.au R N. Holl. 1822. D p.l Sm. ex. bot. 1.12
Dendrobium punctatum Sm.

0539. GEODO*RU M Jac. GEODOHUM. (Ge, earth, dot on, gift; blossoms lying on earth.) *Orchidee.* 3.-4.
22649 purpureum R.Br. purple El el l jn.au P E. Indies 1800. D l.p Rox.c.1.40
Linodbrum nutans Rox.

22650 citrinum H.K. Citron-colored f Q3 d 1 ad Y E. Indies 1800. D l.p Bot. mag. 2193
22651 dilatatum R.Br. dilated-tt/poi ^ E Sd 1 my.au Pk E. Indies 1800. D l.p Bot. reg. 675

* 2530. CAT ASETUM Rich. CATASETITM. (Kata, downward, seta, bristle; two horns of col.) *Orchidea.* 6.—7.
25652 tridenfatum Hook. three-toothed jg El or 2 jlj.au Y.Hr Trinidad 1822. Dp r.w Hoi.k.ex.fl.90
22653 Claverinj Lindl. Cap. Claverinj's jf El or 2 jlj.au Y.Br Brazil 1822. Dpr.wliot.reg. 840
22654 floribundum Hook. bundle-flwd £ El or 2 n. Y.Hr Trinidad 1824. D pr.w Hook. ex. fl. 151
22655 Hofikeri Lindl. Hooker's js El or J J n Y.Br Brazil 1818. D p.r.w
22656 cristatum Lindl. crested £ El cu 2 on G Brazil 182J. D p.r.w Bot. reg. 966
22657 semiapertum Hook. half-open jg El or 1 n Y.o Brazil 1824. D p.r.w Hook. ex. fl. 213

2531. ANGULO*xil R. & P. ANGULOA. (Fran, de Angulo, a Spanish naturalist.) *Orchideee.* 1.—3.
22658 grandiflbra Kth. large-flowered £ EJor 1 jlau ... S. Amer. 1823. D p.r.w H. & B. 1.27

* 2532. ZYGOPE'TALUM Hook. ZYGOPETALUM. (Zygos, yoke, petalon, petal; joined at base.) *Orchideee.* 2.
22659 Mackht Hook. Mackay's £ El or 1 jnjl B.o.v Brazil 1825. D p.r.w Bot. mag. 27+8
22660 rostratum Hook. beaked £ E J or f o W.p.br Demerara 1827. D p.r.w Bot. mag. 2819

2533. SOPHRONI*TIS B.R. SOPHRONITIS. (Sophron, modest; appearance.) *Orchidece.* 1.
22661 cernua B.R. drooping i? El pr \ jn.jl R Rio Jan. 1827. D p.r.w Bot. reg. 1129

2534. HETEROTA'XIS B.R. HETEBOTAXIS. (Heteros, various, taxis, series.) *Orchideee.* 1.
22662 cras8ifoliaff.it. thick-leaved /2E J or | my.jn Y Jamaica 1825. D p.r.w Bot. reg. 1028

2535. TRIZEIAXIS Lindl. TRIBUXIS. (Treis, three,xeuxis, union; 3 segments joined.) *Orchidece.* 1.
22663 falcate Lindl. falcate f Sd 1 fmr G W. Indies 1820. Dp.rw Lind.col.2

2536. XYLO^BIUM Lindl. XVLOBIUM. (A>/on, wood, Wo, to live; growing upon.) *Orchidets.* 2.
22664 squalens Lindl. dirty flowered £ E3 cu \ my.jn Y.Br Brazil 1822. D p.r.w Bot. reg. 732
Dendrobium squalens B.R.
22665 longiflbum Lindl. long-leaved / [El cu 1 my.jl — N. Gran. 1822. D p.r.w
Dendrobium longiflbum Kth.

* 2537. MAXILLA^RI A Ft. per. MAXILLABIA. (Labellum resembles the maxilla* of some insects.) *Orch.* 7.—
22666 Barrigjtrini^ Lindl. Barrington'B £ El cu i jnau Y.G W. Indies 1790. D p.r.w Hook. ex. fl. 119
22667 Harrison&e Lindl. Mrs. Harrison's? El snl ljs Y.B S. Amer. ... D p.r.w Bot. reg. 897
22668 párvalia Hook. small "gfA1<- i mr.irgr Br.pk Rio Jan. 1824. D p.r.w Hook. ex. fl. 217
22669 pallidiiflbra Hook. pale-flowered T El cu ls Pa.Y St.VmcntB26. Dpr.wBot.mag. 2806
22670 aromaticia Grah. aromatic £ El cu 1 my Y.G Mexico 1824. D p.r.w Hook. ex. fl. 219
22671 Parkkérii Hook. Parker's ft 171 or \$s Y DcmcraralB26. D p.r.w Bot. mag. 2729
S2672 racemosa Hook. racemose > El o f my.jn Y S. Amer. 1826. D p.r.w Bot. mag. 2789

2538. NOTYLIA Lindl. NOTYLIA. (Notos, back, tplos, hump; singular callosity on stigma.) *Orchidece.* 1.—
SS673 punctata Lindl. dotted "H" * au.s G Trinidad 1822. D p.r.w Bot. reg. 759
Pleurothallis punctata B.R. Gomeza temuliflra B. &

- Ex39. PLEUROTHA'LLIS R. Br. PLEUROTHALLH.** (*Pleura*, side, *thalleo*, to flower.) *Orechidea?* 3.-5.
 22674 racemiflora Lindl. raceme-flwd E] cu 1 ag G W. Indies 1823. D p.r.w Hook.ex. fl. 123
 ***75 ruscifolia R. Br. Ruscus-leaved <E] cu fmy,in G.v W. Indies 1791. D p.r.w Hook.ex. fl. 197
 22676 foli&sa Hook. leafy j| El cu f f Y Brazil 1825. D p.r.w Bot. mag. 2746
- 2540. ONCIDIUM Swz. ONCIDIUM.** (*Ogkidion*. a tubercle; two prominences on lip.) *Orechidec.* 17.—25.
 *2677 altestimum Swz. tallest Elor 4 aus Y W. Indies 1793. D p.r.w Jac. am. 141
 22678 carthaginense Swz. Carthaginian Elor 4 my,jn Ol W. Indies 1791. I) p.r.w Bot. mag. 777
 **79 divaricatum B. R. divaricated Elor 11 jl.au GR Brazil 1825. D p.r.w Bot. reg. 1050
 22680 pulchellum Hook. neat Elor i jl.au W.Y Demerara 1826. D p.r.w Bot. mag. 2773
 >>wl pilos B. R. downy Elor 1 jl.au Y.R Rio Jan. 1825. I) p.r.w Bot. reg. 1007
 22682 bifbhum H. K. two-leaved Elor 1 jl.lau S Amer. 1811. D p.r.w Bot. mag. 1491
 *81&3 ornithorhynchon Kth. bird's-beak Elor 1 jl.lau Y Mexico 18 & D p.r.w
 22684 GeboHMi Swz. Cebollet's Elor 1 jl.lau Y W. Indies 1825. D p.r.w Jac.am. 131.2
 ***85 tctrapctulatum W. four-petaled Elor 1 jl.jl Y Jamaica 1824. D p.r.w Jac. am. 142
 *868.unciflbum Spr. Rush-leaved Elor 1 jl Y W. Indies 1833. D p.r.w
 Epidindruim unciflbum L.
 *2687 trtquetrum R. Br. triangular-frd Elor 1 jl Y Jamaica 1793. D p.r.w
 gffis l&ridum Lindl. lurid Elor f.mr Ol S. Amer. 1822. D p.r.w Bot. reg. 727
 *89 variegatum Swz. variegated Elor 1 jl.il Y W. Indies 1824. D p.r.w SI. jam. 1.148.2
 WO barbulum Lindl. bearded Elor 1 ap.my Y S. Amer. 1818. D p.r.w Lind. col. 27
 W1 flexubsum 22. JM flexuous Elor 1 jl.jl Y Brazil 1818. I) p.r.w Bot. mag. 2203
 2692 pcfmilm L/nrl. dwarf Elor 1 jl.jl Ol Brazil 1824. D p.r.w lint. reg. 920
 32693 papilio Zfatt. Butterfly Plants Elor 1 O.Y.H Trinidad 1823. D p.r.w Bot. reg. 910
- 2541. CYRTOPCTDIUM fl. Br. CYRTOPODIUM.** (*Kyrtos*, convex, portion, little font.) *Orechidec.* &
 22694 Andersf.nfi R. Br. Anderson's f El el 2 my.au Y W. Indies 1804. I) p.l Bot. mag. 1800
 Sta? Woodfordi B. M. Woodford't g El el So Pk S. Amer. 1814. D p.l Bot. mag. 1814
 **696 glutinbum Mey. clammy z El cl 2 ... S. Amer. 1825. D p.l
- 2542. CGELCGYNE Lindl. C<ELOGYNE.** f hollow, gyne, female; form of stigma.) *Orechidec.* 3.-7.
 22697 punctulata Lsirfl. dotted->Iow<< El el 4 ... V E. Indies 1822. D p.r.w
 2a&8 nltida LiW/. slinhhVclvcrf El el 1 ... Y E. Indies 1822. D p.r.w
 92699 firabrltda L/mtt fringed Elpr i jl-o Y.br China 1824. D p.r.w Bot. reg. 868
- 2543. MACRADETNA R.Br. MACRADENIA.** (*Makros*, long, *aden*, gland.) *Orechidea.* 1.-2.
 22700 lutescens R. Br. lutescent f SELcu i d Ol Trinidad 1821. D p.r.w Bot. reg. 612
- 2544. ANISOPE'TALUM Hook. ANHOPETALUV.** (*Anisos*, unequal, *petalon*, petal; sepals.) *Orechidec.* 1.
 22701 C&reyunum Hook. Dr. Carey's f | S] cu | o Br.P Nepal 1823. D p.r.w Hook. ex. fl. 149
- 2545. MEGACLI'NIUM B. R. MBOACLINIUM.** (*Megas*, large, *klinio*, to bend; spike.) *Orechidecar.* 1.
 22702 falcatum B. R. falcate f El or 1 jl.s Y.aR S.Leone 1824. D p.r.w Bot. reg. 989
- 2540 GONGOMI/J R& P. GONCORA.** (*Anton. Caballeroy Oongora*, a Spaniard, and friend of Mutis.) *OrcA.* 2.
 2277 r a a t w p u ^ a // S? dark-purple f El or 1 Jn.jl T.P. Trinidad 1824. D p.r.w Hook. ex. fl. 178
 2Q7JL-E^K? 15: * Jjowy j| S or li Jn.jl Y Brazil 1826. D p.r.w Bot. mag. 2755
- *2547. DENDRO^BIUM Swz. DBNDROBIUM.** (Dendron, tree, Wo,to Vive; growing upon.) *Orechidea.* 14.-24.
- I. TERMINAYIA.—JrtdWM terminal.
- 22705 speciosum Swx. showy EJ spl 1 jn.au P N. & W. 1801. D p.r.w Ex. bot 1.10
 227W Pierfirdi Rox. Pierard's El el 2 mr Pk E. Indies 1815. D p.l Hook.ex. fl. 9
 22707 Calceolaria Hook. slipper El or 2 year Pk E. Indies 1820. D p.r.w Hook.ex. fl. 184
 22708 fimbriatum Hook. fringed El or 2 ap Y E. Indies 1823. D p.l Hook. ex. fl. 71
 22709 crumcnatum Swx. pursed El or 2 ap.my W Sumatra 1823. D p.l Ru. am. 6.47.8
 22710 fe mulum R. rival El or 2 ap.my W N. S. W. 1824. D p.r.w
 **7H canaliculatum R Br. channelled El or 2 ... N. Holl. 1824. D p.r.w
- II. —Scapigerous.
- 22712 MyosHrus Swz. Mouse-tail f El or 1 jn.au ... Society Is. 1825. D p.r.w
- III. SOLITANIA.—Peduncles solitary.
- 22713 cucullatum R. Br. hooded f El el 2 mr Pk E. Indies 1815. D p.l Bot. mag. 2242
 2*714 moniliforme Swz. bracelet-formed^ El pr f ... P China 1824. D p.r.w Kern. 865 .
- IV. LATKRA^IA.—Racemes lateral.
- 22715 lingusfrme Swx. tongue-leaved j| El cu ... P N.S.W. 1810. D p.r.w Ex. bot. 1.11
 g716 undulkum R. Br. waved f El or f year ... N. Holl. 1823. D p.r.w
 *717 teretifolium R. Br. round-leaved El or 1 jn.au P N. Holl. 1823. D p.r.w
 22718 rJgidum R. Br. rigid El or ft. ... N. Holl. 1824. D p.r.w
- c.2548. CAMARPDIUM Lindl. CAMARIDIUM.** (*Kamara*, arched roof; tip of stigma vaulted.) *Orech.* 1.—2.
 <2719 ochroleucum Lindl. yellowish-white A El pr 1 jl W Trinidad 1823. C p.r.w Bot. reg. 844
 Dendrobium filbum Hook.
- *9549. ORNITHI'DIUM Sal. ORNIT. (*Ornis*, bird, *eidos*, like; upper lip of stigma beak-like.) *Orcf.h*
 82720 coccineum Sal. K*v&et-Jlowered&E\ov 2 year R W. Indies 1790. D p.r.w Bot. mag. 1437
- 2550. ISOCHPLUS R. Br. ISOCHILUS.** (I*OS, equal, cheUos, lip; sepals and lipO *Orech.* 3.-5. ?
 22721 hneariBA5r. linear * El pr f my,jl R W. Indies 1791. D p.r.w Bot. reg. 745
 22722 prdlifei R. Br. nroliferous "Sm pr * ... W W. Indies 1793. D p.r.w Bot. reg. 82a
 22723 SraminidiuV Kth. SrasstfSSed 1 ^ o r \ my,jn G.v Jamaica 1823. D p.r.w Hook. ex. fl. 196
 Cymbidium /fram'tnio/ids Swx,
- 2551. PHOLIDO^TA L. PIHOLIDOTA.** (*Pholis*,* scale; flowers covered with a scate-like bracteaO *Orechidec.* h
 22724 imbricate Lindl. imbricated \ El or 1* ... Br.w Nepal 1824. D p.r.w Hook. ex. fl. 138
- * 2552. BROUGHTOW/* R. Br. BROUGHTONIA. (*Arthur Broughton*, of Bristol, bot. author.) *Orechidec.* 1.
 22725 sanguines R. Br. blood-colored f USI spl li jn.jl S Jamaica 1793. D p.r.w Bot. mag. 793
- *2553. CATTLE ^ Lindl. CATTLE Y A. (W. *Cattley*, member of the London Hort. Soc.) *Orechidea.* 3.-4.
 22726 labiate Lindl. <ar*-lipped AE J spl 1 jlau V S. Amer. 1818. D p.r.w Lind. col. 33
 VHZ² Lodilig*g*s*i* Lindl. Lodigges's £ 2.3 el 1 jlau V S. Amer. 1815. D p.r.w Bot. cab. 357
 -3728 Forbesi Lindl. Forbes'a f El or f jl au Pk S. Amer. 182a D p.r.w

•2554. EPIDE'NDRUM L. EPIDENDRUM. (*Epi*, upon; *dendron*, a tree; growing upon.) *Orchidete*. 19. — 67.

I. BULBOSA. — *Bulbous s trapes few or manyflowered.*

82729	cochleatum	£.	spiral	EI	cu	1	*Ed	Br.p	W. Indies	178a	D s.p Bot. mag. 572
22730	fragrans	Swz.	sweet-scented	[S]	fra	f	o	Y.G	Jamaica	1778.	D s.p Bot. mag. 1669
22731	angustifolium	Swz.	narrow-leaved	[C]	or	u	au.o	Y	Jamaica	ISJO.	D p.r.w
22732	punctatum	L.	dotted	E	or	1	o	- -	W. Indies	Ifttf.	D p.r.w
22733	anceps,/oft		two-edged	EI	or	1	au.o	-	£ Indies	1820.	I) p.r.w Bot. cab. 887
22734	ellipticum	Hook.	elliptic	ES			fmr.au	Pk	Rio Jan.	1844.	Dpr.wHook. ex fl. 207
22735	ciliare	L.	fringed	EI	or	1	mr.au	W	W. Indies	1790.	Dixr.wBot.reg.784
22736	cuspidatum	Lod.	pointed	EI	or	1	jn	w.y	W. Indies	1808.	D p.r.w Bot. reg. 783
22737	monophyllum	tfcoAr.	one-leaved	EI	cu	£	d	G	Jamaica	1823.	I) p.r.w Hook. ex. fl.IIOO
22738	polybilobum	Sua.	many-bulbed	EI	cu		d	W	Jamaica	1822L	D p. r.w Hook. ex. fl. 112

II. ABULBOSA.—*Destitute of bulbs; flowers solitary, racemose, aggregate, or panicled.*

22739	<i>scleranthum</i>	L.	side-flowering	£ E l o r	2	jn.jl	R	W. Indies	1793.	C p.r.w Jac. am. 137
82740	<i>fuscum</i>	Sibt	brown	£ V C	—	£ jn.jl	Br	W.	1793.	D p.r.w Bot. reg. 67
	<i>anceps</i>	Jac.								
98740	<i>elongatum</i>	Jac.	<i>long-stalked</i>							
82742	<i>uinbelftum</i>	Sivz.	umbellled							
82743	<i>n titans</i>	Swz.	nodding							
82744	<i>conspicuum</i>	Bartr.	gnat							
82745	<i>diffusum</i>	Swz.	diffuse							
22746	<i>verrucosum</i>	Swz.	warty							
22747	<i>pedicellatum</i>	L.	night							

2555. POLYSTA[%]CHYA Hook. POLYSTACHYHA. f (*Polys.*, many, *stachys*, a spike.) *Orchidete.* 2. — 5.
82748 luteola Hook. yellowish ligula cu f *jl.au* Y.G W. Indies 1818. D pr.w Hook. ex. fl. 103
Dendrobium polystachyum Swz

2556. CRYPTARRHETNA R. Br. CRYPTARRHETNA. (*Kryptos*, hidden, *arrēn*, a male). Oichidece. 1.
2259. *Brachystele* P. Br. *Brachystele* ~~is~~ ^{was} ~~not~~ ^{not} limited, if ES does not mean W. Indies. 1815. Druce, 1884, p. 152.

2557. ORNITHOCEPHALUS Hook. Onv. (*Ornis*, bird, *kephale*, head; upper end of column.) *Orcb.* 1.
82751 *gladiatus* Hook. sword-leaved /OSCU i ... G. Trinidad 1823. D. p. r. w. Hook. ex. fl. #7

*2,58.	BLRTIA.	(<i>Louis Blet</i> , a Spanish apothecary and botanist.)	<i>Orchidace.</i>	9.—14.
82752 Tankerville & Ir. Br.	Tankerville's	A ED spl	2 mr.ap	W.Br China 1778. R p.l Bot. mag. 1124
82753 verecunda <i>R. Br.</i>	modest	A El el	3 jamy	P V. Indies 1733. R p.l Bot. mag. 930
Limodorum altum				
82754 flurida <i>It. Br.</i>	florid *	A El el	2 jl.au	P W. Indies 1786. R p.l Red. 111. 83
22755 speciosa <i>Kth.</i>	showy	A El or	2	Mexico 1827. R p.r.w
9SHS5 Guincensis Cr. <i>Don</i>	Guinea	A El or	2 f ****	P S. Leone 1822. R p.r.w
9SHS1 Ayacanthi/w R Br.	Hyacinth-like	& E3 el	1 rnr.jn	P China 1802. R p.l Bot. mag. 1492
82758 capitata <i>R. Br.</i>	headed	A El d	—	W. Indies 1795. R p.l
82759 pSlida <i>Lod.</i>	pallid		jn-jl	W. Indies 1820. R p.l Bot. cab. 629
88760 Woodfordii Hook.	Woodford's		f	Pk Trinidad 1823. R p.r.w Bot. mag. 2719

2559. ETRIA Lindl.	ERIA.	(<i>Eriogonum</i> , wool; ^{jn} ^{jl} ^{G.v} flower.)	Orchideet.	3.-4.
82761 stellata Lindl.	shirred	ACS el 2 f	Br.Y E. Indies?...	D.p.r.w Bot. reg. 90+
82762 publscens Lindl.	pubescent	£ ES el 1J mr	Y E. Indies	1820. D.p.r.w Hook. ex. fl. 124
Dendrobium publscens Hook.				
22763 rosea B. R.	Rose-like	■ T ■	Id R.w China	1824. D.p.r.w Bot. reg. 978

8560. OCTOMERIA R. Br. OCTOMERIA. (*Okto*, eight; *meris*, part: pollen in eight manses.) *Orchidete.* 2.
 82764 [*raminiflora*] B. R. Grass-leaved $\frac{1}{2}$ EScu $\frac{1}{2}$ jn.jl Pa.Y W. Indies 1793. Dpr.w Bot. mag. 2764
 82765 spicata D. Don spicate $\frac{1}{2}$ CSor — Nepal 1822. $\frac{1}{2}$ p.r.w
 82766 serratifolia Hook. serrated-leaved $\frac{1}{2}$ EI or $\frac{1}{2}$ Pa.Y Rio Jan. 1826. D.p.r.w Bot. mag. 2823

8561. TRIBRA^CHIA B. R. TRIBRACHIA. (*Treis, three, brachion, arm; spikes*) Orchidee. 1.
82767 p&idula B. R. pendulous .£ E J C U } jn.n G & Leone 1823. D p.r.w Bot reg. 963

•2562. BRASAVCTLI R Br. BRASA\OLV (*Ant. Musa Brasavoto*, an Italian botanist.) *Orehidea*. I.—2.
82768 cuculfata R. Br. hooded £ ES el i jn.s W W. Indies 1793. D p.r.w Bot. mag. 543

* 2563. SARCA'NTHUS	<i>Lindl.</i>	SARCANTHUS. (Sara, a Geith, anthos, flower; texture of sepals.) <i>Orchidee</i> . 5.		
27690 panicuftus	<i>Lindl.</i>	panicled £ El el 2) my.au Y China C p.r.w Bot. reg. 220		
22770 premorsus	<i>Lindl.</i>	bitten 2 23 el 1 G E. Indies 1824 DD prw		
82771 teretifolius	<i>Lindl.</i>	cylindric-lvd "£ El cu 1 n Y.P China 1819. C p.r.w Lind. col. 6		
82772 rostratus	<i>Lindl.</i>	rostrate £ El P' In YRG China 1819. C p.r.w Bot reg. 981		
<i>Vdnida recdrrva Hook.</i>				
99713 suclsus	<i>B. R.</i>	lopped off £ El pr In Y.br China 1824. C p.r.w Bot. reg. 1014		

8564. VA 'NDA R. Br.	VANDA.	(Hindoo name of original species.)	Orchidee.	5.-6.
22774 multi flora Lindl.	many-flowered	E[e]l. 2 jn	Y China	1800. C p.r.w Lind. col. 38
82775 Cycnium Kua Lodd.	tessellated	H[ui] 1 jn.lau	W. P China	1816. D p.r.w Rox. cor. 1.4

22776	Roxburghii' R. Br.	Roxburgh's hairy-rooted	E l el au	W.p P.G	China E Indies	1810. 1822.	C p.r.w C p.r.w	Bot. reg. 506 Hook. ex. fl. 72
22777	trichoriza Hook.	Trichorh'za teretifolia Lindl.	E lpr					
22778	crucifera Lindl.	bloody-flowered	E l or 2 au	R	China	1819.	D p.r.w	

2883. *AE* *KELLS* Sw. *AIR FLENT*. (Ac., the air, live for many months when suspended). *Ctenacut.* 2.—11.
 82779 *odkrotum* *Lou.* fragrant £ El fra 1| ... Pk China 1800. C p.r.w
 82780 *arachnites* *Swz.* spider-like £ El or 1| ... Br.p Japan 1793. C p.r.w Kcm. 869.1

8566. RENANTHE'RA Lou. RENANTHERA." {Ren, kidney, antkera, anther.) Orchidee. 1.
8278 cocinea Lou. scarlet £ ES spl 8 mr.myS China 1816. C p.rw Bot. reg. 1131

*T-68 EULOPHIA i? flr. EULOPHIA (*Eulophus*, well crested; surface of middle lobe of lip) *Orchidaceae*. 8-10.

82783 *gracilis* liiffl. slender A El pr 2 my.n G.Y S. Leone IKS. R p.l Bot. reg. 742
 92784 *guineensis* R. Br. Guinea A El d 1 my.n Pk S. Leone 1822. R p.l Bot. reg. 686

- 22785 *barbita* B. R. bearded AiAJor 1 jn.au C G. H. 1825. D p.l
 22786 *streptoneta* B. R. narrow-petaled A iAJ or 1 jl.n Y C.O.H. 1820. D pJ Bot. reg. 1002
 22787 *virens* R. R. green-flowered A LAI or A jn.au G.v C. G. H. 1825. D p.l Rox. cor. 1.38
 22788 *trilatis* B. R. sad A IAI or H myin Dk C. G. H. 1825.
 22789 *longimis* B. R. long-leaved * A I or H ph.au ... C.G.H. 1825. D p.l
 22790 ~~ofo~~ *oblonga* B. R. sword/w* S [] ap.au Pa.Y C. G. H. 1826. D p.l.s Bot reg. 1147

* 2569. ANGRJPCUM Thou. ANG&BCUM. (Latinised from the Malay appellation *aneree*.) *Orchidea*. a—
 22791 *maculatum* Lindl. spotted jgiZSpr I a n Plk Africa 1819. D p.r.w Lind. col 15
 22792 *falcatum* Lindl. falcate tfESpr * n.d. W China 18Jf Dp.r.w Bot mag 2097
 22793 *luridum* Lindl. lurid i&GSpr ljs.n Br S. Leone 1822. Dp.r.w

" 2570. A ERA'NTHES IAndl. AERANTHES. (Aer, air, anthos, flower.) *OrchMe**. 2.— 3.
 22794 *grandiflora* Lindl. large-flowered fU [] f jnjl G.Y Madagascar 1823. Dp.r.w BoBreg 1817
 22795 *sesquipedalis* Lindl. 18-inch-flaid [] 1 jnjl W Madagascar 1823. D p.r.w Thou. or. 66

2571. CALA'NTHE R. Br. CALANTHE. (*Kalos*, beautiful, *anthos*, flower) *Orchide**. 2.— 7.
 22796 *veratrifolia* R. Br. Veratrum-lvd £ El or 2 Jnjl W E. Indies 1819. D p.l Bot reg. 720
 22797 *sylvestris* Lindl. wood £ 123 or ljl.au W? Madagasc.1823. D p.l

* 2572. STETLIS Sun. STELI& (Greek name of some parasitical plant) *Orchidea*. 2.— 10.
 22798 *ophioglossoides* &itz. Ophiogloss.-lk * E Jc u } myjn G W. Indies 4791. D p.r.w Bot. cab. 442
 22799 *microstoma* Sun. snail-flowered Jg BScu f n.d. G Jamaica 1805. D p.r.w Hook. ex. fl. 158

2573. MALA'XIS Sun. MALAXIS. (*Malaxis*, softness; delicate texture of genuine species.) *Orchid*. 1.—13.

22800 *paludosa* Swz. marsh ^ A ^ e i jl Y.a England tur.bo. R pg Eng.bot.72

*2574. PRESCO*T/i< Lindl. PREBCOTIA. (John Prescot, an Englishman residing at Petersburgh.¹) *Orch*. 1—2.
 22801 *plantaginea* LintU. Plantain-feau<rte Q3 cu 1] ap.au G.w Brazil 1822. R p.l Hook. ex. fl. 115

*2575. MICROSTYLYS Nut. MICHOBTYUS. (Mikros, small, styllos, a column.) *Orchide* f.** f.±

22802 *ophioglossoides* Nut. Ophiogloss.-lk & A ^ e * j» Y.o N. Amer. 1824. R p.s Pluk. am. 434.4

*|57R LI'PARIS Rich. LIPARIS. (*Uparos*, unctuous; surface of leaves of original species.) *OrchUc** 6^13.
 22803 *lily* T Ha [] Lily-leaved & A pr i injl G.p N. Amer. 1758. R p.s Bot. mag. 2004
 22804 L' Rfch ^ S e I's 5 A or * injl Y. ■ England sa ma. R p.s Eng.bot.47
 22805 Corre'W Nut Correa's 5 A or * injl Y. ■ England sa ma. R p.s Eng.bot.47
 22806 *bituberculata* Lindl. two-tuberclcd £ E3cu f jl G N. Amer. 1823. R p.s
 22807 foehsa Lindl. leafy £ El cu ft au G Nepal 182^ Dp.r.w Hook. ex. fl. 116
 22808 reflexa Lindl. reflexed ^ E 1cu 1 au G I. France 1823. D p.r.w Bot. reg. 884
 N. Holland 1824. Dp.r.w

2577 FERNANDETS/jtf R. & P. FERXANDGSI. (George Garcias Fernandez, a Spanish bot) *Orchidea*. 1.
 22809 Elegans B. R. elegant ^ E Jc u ft jnjl Y Trinidad 1817. Djxr.w Bot mag. 2715.

Lockh&rtia Elegans Hook.

2578. CALYTSO Sal. CALYPSO. (Kalypto, to conceal; habitat.) *Orchidea*. 2.
 22810 borchMis Sal. northern A DT 4 myjn Yi N. Amer. 1805. R s.p Par.Ion.89
 22811 americana R. Br. American A*or * myjn Pk N. Amer. 1820. R s.p Hook. ex. fl. 12

2579. VANILLA Sw*. VANILLA. J - M ^ in W W. Indii 180ft C p. Bot.«b.7»
 aromatic L O o r 10

aromatica Sun.

CLASS XX. ORDER 2.

GYNÀ'NDRIA DIA'NDRIA.

- | •2580. CYPRIPEMTHUM L. | | LADIES' SLIPPER. | (Ajw. #*, Venus, podion, a slipper.) | Orchidca. | 10.—10. |
|--|-----------------------------|-----------------------|--------------------------------------|-------------|--|
| I. FOLIUSA — Stems leafy. | | | | | |
| 22814 | <i>Calceolus L.</i> | common. Slipper Swiss | A or A | my.jl my.jl | England woods. Switzerland. 1825. |
| 22815 | <i>2 helveticum</i> | | or | T T | s.p Eng. s.p |
| | <i>Calbelous Schl.</i> | | | | |
| 22816 | <i>parviflorum Sum.</i> | small-flowered | jfc | A or | N. Amer. 1780. |
| 22816 | <i>pubescens W.</i> | pubescent, yellow & | A | my.jn | R sp Bot.mag. 9H |
| 22817 | <i>Recifebile Sun.</i> | remarkable | ·^L | A or | s.p Bot. cab. 895 |
| 22818 | <i>album // K.</i> | white | ^A | 11 | R N. Amer. 1731. |
| 22818 | <i>candidum IF.</i> | white | 5 | A or | R 8.p Bot. mag. 216 |
| 22820 | <i>arcticum R. Br.</i> | white nm't. head | k | A or | W N. Amer. 1800. |
| | | | k | A or | W N. Amer. 1808. |
| | | | j | my | R s.p Bot mag. 15G9 |
| II. SCAPITHEA. — <i>Scaptgerous.</i> | | | | | |
| 22821 | <i>hhmille Sun.</i> | humble | i | A or | * myjn Hkw N. Amer. 786. |
| 22822 | <i>venustum Wai.</i> | handsome | tfGSel | ijlau | R s.p Hot reg. 188 |
| 22823 | <i>insigne Wai.</i> | admirable | IS | el i l j u | G.P. ** ^{s.pri} Jfli. R g.p Lind. col. 32 |
| | | | | u | G.P Napal 1819. |
| •2581. STYLI'DIUM Sun. STVLIDIUM. (<i>Stylos</i> , column; stamens andt sttlc jogged.). | | | | | Syllidea. H.—45. |
| 2M24 | «paminorium5ta». | GraS.leaved | * | IAJor | a.p Bot reg. II |
| 22825 | <i>fruticellatum R. Br.</i> | shrubby | ■ | 1 ap.au | N. & W. Iaoa. 1803. |
| | | | ■ | 11 my.o | Pk N. Holl. 1803. |
| | | | | Pk | p.s Par. Ion. 77 |
| glandulsa Sal. | | | | | S.s.p |
| ■^2(2) | <i>Bcandens R. Br.</i> | climbing | · | S jlau | N. Holl. 1803. |
| 22827 | <i>tenuiflum R. Br.</i> | fine-Icaved | £ | 1 jlau | S.s.p Bot. mag. 2249 |
| | <i>/ariciflum Rich.</i> | | | Pk | N. Holl. 1818. |
| 22828 | <i>adnatum R. Br.</i> | adnate | | jlj.lau | S.s.p Bot. reg 914 |
| 22829 | <i>lineare Swt.</i> | linear | | E jn.jl | D p.l Ex. bot. 2. f. |
| | Ventenatae minor Sm | | | | |

CLASS XX. ORDER S.

GYNA'NDRIA HEXA'NDRIA.

•2582. JRISTOLO'CHIA L. BIRTHWORT. (*Aristos*, best, *lochia*, parturition; sup. med. qual.) *Asarina.SUiO.*

I. TM'FIDJE	—Limb <tf perianth flat, trifidj shrubby.
22830 slpho <i>Hcrit.</i>	tube A or 30 jn.jl Y.br N. Amer. 1763. L Hp Bot. mag. 534
22831 tomentosa B. M.	tomentose J or 20 jn.jl P N. Amer. 1799. L s.p Bot. mag. 1369
22832 Arkinsa La. C.	Arkansa J or 20 jn.jl P Arkansa 1824. L 61

II. UNILABIA*™.—LimA qf perianth unUbiate.

i. OBLO'NGJS.	—Lip ovate, oblong, or lanceolate.
22833 boe'tica L.	Beetic ^ o r 6 myjn Spain 1596. R l.p M. h. 12.17.6
22834 glafica Desf.	glaucous-fcam/JL i_J or 6 jl Barbary 1785. C p.l Rot. mag. 1115
22B35 maxima/..	greatest LCD or 20 jl N. Spain 1759. C l.p Jac. am. 146
22836 hasftta Kth.	hastate & O or 16 jlau Cuba 1822. C p.l H.&B.2.116
22837 indica L.	Indian A. Q or 10 jn.jl P E. Indies 1780. C s.l R.mal8.25
22838 acuminata Lam.	/on&'ixrinted fr I lor 10 ... P Mauritius 1822. ■■■
22839 bilobdta L.	two-lobed J I lor 10 ... P 1824. G p.l PI. am. 106

ii. SPATULAV*.—Lip spatulate, obtuse.

22840 baTbhta Jac.	bearded fl ES or 10 ... P Caracas 1796. R 8.1 Jac. ic. 3.608
22841 foe'tida Kth.	fetid fi_CZlor 20 jlau ... Mexico 1822. C p.l H.&B.2.119

iii. CORDAV/E.—Lip cordate, unguiculate.

22842 odoratissima L.	sweetct-stcd i. D o r 10 jl P Jamaica 1737. C p.l SI.jam. 1.104.1
22813 surinamensis W.	Surinam fl_Qor 20 ... Y.br Surinam 1823. C p.l Jacob. 1.3

iv. ODCORDA'VB.—Lip at top cordate.

22844 trilobata Z.	three-lobeJ ft. O or 6 jn.jl P S. Amer. 1775. C p.l
22845 grandiflora Swx.	large-flowered fl * or 20 ... Jamaican 1824. C p.l

III. I ■■■.—Limb of perianth bilabiata.

22846 ringens Vahl	ringent ■■■ or 20 jn.jl P.u.v Brazil 1820. C p.l Vahl sy. 2.47
22847 brasiliensis Mart.	Brazilian ■■■ or 20 ... Y.br Brazil 1820. C p.l
ringens Lk.	
22848 panduriformis W.	fiddle-shaped D o r 10 ... Caracas 1823. ■■■
22849 labibsa A A.	or 20 ji.au P.G.Y Brazil 1821. ■■■ [p] Bot.xeg.689
22850 arborescens W.	arborescent l_Jor 20 jn.jl Y.p America 1737. ■■■

IV. HKROAVE v.—Herbaceous.

i. U\LAoiA'r.ii	—Limb of calyx unUbiate.
92851 sempervirens L.	evergreen ■■■ Candia 1727. D p.l Bot. mag. 1116
22852 longa L.	long-rooted ■■■ S. Europe 1548. R co Mil. ic. 51.2
52853 Scrpntaria L.	Snake-root ■■■ N. Amer. 1632. R s.p Jac. sc. 3.385
22854 bracteata Jtete.	bracted ■■■ E. Indies 1793. R si
22855 rot&nda L.	round-roofa? ■■■ S. Europe 1596. R co
22856 sagittata MM.	sagittate jnj N. Amer. 1819. R s.l Pluk. m 78. 1
22857 paTida JK	p&v-e-jioivred my.au W.p Italy 1640. R s.l M. h. 12.18. 2
22858 Clematis L.	Clematis my.au Y England woods R co Eng.bot.3U8
22859 hirta L.	hairy)& IAI or 2 myjn P Chio 1759. R s.l To. it. 1.147
22860 Pistolochia L.	Pistoluchia -* & or 2 jn.jl P S. Europe 1597. R ■■■

CLASS XXI. ORDER I.

MONCE'CIA MONA'NDRIA.

• 2583. ARTOCA'RPUIS L. BREAD FRUIT.	(<i>Artos</i> , bread, <i>karpos</i> , fruit.) <i>Urticea.</i> 2.—a
22861 incisa 7* true, cut-lvd f Q clt 50 ... W.u	S. Sea I. si. 1793. Sk r.m Ru. am. 1.33
2nuclfera nut-bearing J Q o r SO ...	W.u E. Indies 1793. Skr.m
22862 integrifolia L. entirc-lvd, <i>Jaca Trce</i> ± clt 60 jn ... W.o E. Indies 1778. C r.m R. mal3.26.28	
2 heterophylla Xaw. variable-leaved j * or 60 ... W.u E. Indies 1778. C r.m	

' 2584. FOILSKCPEAL.	FORSKOLRA. {Peter Forskol, a traveller in Egypt.) <i>Vrticeie.</i> 3.-5.
22863 tenacissima L.	clammiest O cu 1 jlau W.G Egypt 1767. S Itl Jac.vin.L48
22864 Candida L.	white T LA cu U jn.jl W.G C. G. H. 1774. C Itl
22865 angustifolia Retz.	narrow-leaved O cu 2 jlau W.u Teneriffc 1779. S Itl Mur.24.4

S585. CASUARINA L.	CASUARINA. (Branches sup. like feathers of Cassowary.) <i>Amentiae.</i> 9.—10.
--------------------	--

	■■■s. — Flowers dioecious.
22866 stricta H. K.	strict L J or 15 f.n Ap N. & W. 1775. S s.p Bot rep. 346
22867 distyla Fen.	two-styled i L J or 15 .. Ap N. Holl. 1812. S s.p Yen. eels 62
22868 torulosa H. K.	twisted i J or 15 .. Ap N. S. W. 1772. S s.p
22869 quadrivalvisLfl*.	four-valved J or 18 .. Ap N. S.W. 1812. S s.p
22870 tenudissima Slub.	slenderest I J or 10 .. Ap N. Holl. 1825. S Bp Lab II. h. 2.218
22871 muricata Jtox.	muricated U Q r 15 ... Ap E. Indies 1822. S s.p

II. MONCEYI-E.—*Flowers monoecious.*
 22872 nodiflora *Font.* knot-flowered t i-J or 15 ... Ap N. Caled. 1823. S ep
 22873 ttuiscifolia *Font.* Equisetum-lvd f u or 15 o.n. Ap S. Sea Isl. 177R S sp Bot. cab. 607
 »874 glauca *Sieb.* glaucous t i-J or 15 ... Ap N. Holl. 1824. S sp

2586. CERATOCA'RPUS L. CERATOCARPUS. (*Keras, horn, karpos, fruit; calyx 2-bomed.*) *Chenopbd&e.* 1.
 22875 arenarius L. sand O un i jn.jl G Tartary 1757. S a Ac. pet. 1.9

2587. ZANNICHE'LLM L. PONDWEED. (*John Jerome ZannichelH, a Venetian apothec.*) *Ftividks.* 1.-8.
 22876 palustrUL. manh * O w 2 jl Ap Britain dit S aq Eng. bot 1844

CLASS XXI. ORDER 2.

MONCECIA DIA'NDRIA.

. 2588. LE'MNA L. DUCKWEED. (*Lepis. a scale; form of plants.*) *Fluvihles.* 4.—11.
 «77 trisfilca L. 3-furrowed, Ivy-lvd | O w ... my.in Ap Britain sta.wa. S lp Eng. bot. 926
 2288 minor L. less ^ O w ... Jnjl Ap Britain sta.wa. S lp Eng. bot. 1095
 2289 gibba L. gibbous A O w ... Jnjl Ap Britain sta.wa. S lp Eng. bot 1233
 ^SWw polyrhiza L. greater^ many-rooted ^ O w ... my.s Ap Britain dit S lp Eng. bot. 2458
 2589. -NGITRIA Tou. ANGURIA. (One of the Greek names for the cucumber.) *CucurbitUce**. 4
 22881 trilobitaX. three-lobed « E or 20 Jnjl Pk Carthag. 1793. R lp Jac.am. 156
 22882 pedata Z. pedate t [23 or 20 Jnjl Y S. Amcr. 1820. H lp Jac.am. 155
 22883 trifolikta L. trifoliate JL E2 or 10 Jnjl Y St. Domin. ... 11 lp PL am. 99
 22884 umbrosatf.^J?. shady fl C5 for 10 Jnjl Y S. Ainer. 1827. R lp
 2590. CERA1TOLA L. CBRATIOLA. (*Iteration, a little horn; stigma.*) *Empetre**. 1.
 22885 cricoldes L. Heath-like *L_lpr lijn.jl Br N. Araer. 1829. C sp Bot mag. 2758
 2591. AfELOTHRIA L. MELOTHRIA. (*Melothron, supposed to be briony j similarity.*) *Cucurbit.* 2.—■
 22886 p&idula L. pendulous -* O un 4 jns Y America 1752. S co Pl. ic. 66.2
 22887 partida Drsr. fetid -* O un 4 jns Y Guiana 1820. S co
 Trichosanthes fectidissima L.

CLASS XXL ORDER 3.

MONCECIA TRIA'NDRIA.

2. COMPTO^NIA Banks COMPTONIA. (*Henry Compton, Bishop of London, cult, exot*) *Ament&cea.* L
 asplenifolia H. K. Fern-leaved ill or 4 mr.my Br N. Araer 1714. Sk sp Den. br. 166

2593. HERNA'ND/il L. JACK IN A Box. (*Francisco Hernandez, M.D., Spanish bot.*) *Laurinete.* a
 sonoraz., Bounding t O or 50 ... W.u E. Indies 1693. C pl Ru. am. 2.85
 ovigera L. egg-bearing ¥ • or 50 ... W.G E. Indies ... C pl Ru. am. 3.123
 Guiana I D o r SO ... W.a Guiana 1820. C rm Aub.gui. 2.309

f A'XYRIS L. AXYRIS. (Axyros, rough; leaves.) *Chenopbdete.* 3.—5.
 ffimaranthoides L. Amaranthus-lk O un li jn.jl G Siberia 1758. S co Gm.ui. 2.2
 probrata L. hybrid O un 1 Jnjl au G Siberia 1780. S co Gm.si. 4.1
 prostrate «* O un 4 Jlj.au G Siberia 1798. S co Gm.si.4.2

TRA^GIA L. TBAOIA. (*Jerome Bock, generally called Tragys, Ger. bot.*) *Euphorblucete.* 9.—19.
 22901 volubilisX. twining i. [Z] un 6 Jnjl G W. Indies 1739. C co Tr.pl. 2.15
 involucrata L. involucrld ^ £D un 3 Jnjl G E. Indies 1759. S co Jac. ic. 1.190
 dunculata Beauv. peduncled JL un 6 Jnjl G Guinea 1823. C co Beau. ow. 1.54

II. ERB'CTJL — *Stems erect.*

stinging O un 3 au G Virginia 1699. ■ Pluk. al. 107.5
 Nettle-leaved O un 2 Jlj.au G N. Amer. ... ■ R.mal.10.82
 Mercury-like CD un Jlj.au G E. Indies ... ■ ■
 small-homed in 1 Jlj.an G Caraccas ... ■ ■
 troublesome CD un 2 Jlj.au G Brazil — ■ ■
 Hemp-leaved il*E2 un 2 Jnjl G E. Indies 1699. C lp Bur. in. 63.4

aof^ up^NEMID/ S T Mart. CNBM. (*Knetnis, spoke of wheel, stachys, spike.*) *Euphorbi&ccaf.* 1.—19.
 22904 Chamelaji Spr. Charaelasa O un 2 Jnjl G E. Indies 1703. S sp R.mal.2.34

2597. rr PHA L. CAT'S-TAIL. (*Tiphos, a marsh; habitation.*) *Typhntce.* 4. ^7.
 j^J latifolia L. broad-leaved J A ec 6 Jlj. Br Britain dit R lp Eng. bot. 1455
 j^06 minor 5m. smaller Jd ^ ec 2 Jlj. Br England mar. * R lp Eng. bot. 1457
 j^07 angustifolia L. narrow-leaved J A ec 4 Jnjl Br Britain pools. R lp En* bot 1456
 82908 minima W. least Jb A ec 1 Jlj Br Switzerl. 1822. R co

2598. SPARGA^NIUMi. BUR REED. (*Sparganon, fillet \ riband-like 1-aves.*) *Tfoh* ■ J c 4. ■ K
 22(K)ramJsuin.yto. branched ^ A un 2 Jlj.au Ap Britain dit S lp | J c 4. ■ K
 22910 simplex Swx. simple * A un ljjl.au Ap Britain sta.wa. S E, & * £ * 745
 22911 nRfans.. floating * A un lit jl Ap England fens. S lp En* bot 278
 aSaZi ■ D. Don alpine * A un fit jl Ap Britain sc.mo. S lp

*2599. CA'REX L.	CAREX. (<i>Careo</i> , to want; upper spikes without seeds.) <i>Cyper&cea.</i> 121.—280.
22913 dioica L.	I. DKEVIJE.— <i>Dioecious, spikes solitary.</i>
22914 Davalliana Sm.	dkecious Jb A cu ft my.jn Ap Britain sp. bo. Sk bog Eng. bot 543 Davall's Jb A cu *my.jn Ap Britain mar. Sk bog Eng. bot 2123
22915 puiicaris L.	II. PI/LICA*HI£.— <i>Androgynous, upper part of spike male.</i>
82916 leucoglabrin L.f.U.	white-spiked M A un UJ ₁ jl Ap Britain mar. Sk bog Eng. bot 1051 Pyrenean M A un UJ ₁ jl Ap Pyrenees 1820. Sk bog Schk. ca. 5. D.15
22917 pyrenaica Wald.	rock * A " n } Jn.jl Ap Switzerland 1819. Sk bog Al. ped. 9%. 1
22918 rupestris All.	few-flowered M A cu 1 Jn.jl Ap Britain bgs. m. Sk bog Eng. bot. 2011
22919 pauciflora Light.	
22920 cyperoides L.	III. CYPERQ'DEIE.— <i>Androgynous, capitule stigmas two, #0.22921 stigmas three.</i>
22921 Juncifolia ij.J.	Rush-leaved M A un £ Jn.jl Ap Switzerland 1819. Sk bog Schk. ca. A.5
22922 stenophylla Ua Jf.JUL	narrow-leaved Jb A un 2 Jn.jl Ap Austria 1822. Sk bog Schk.ca. G.Ii.32
22923 chordata Aza L.	cord-rooted Jb A un 1 Jn.jl Ap Sweden 1823. Sk bog Schk. ca. G.I i.31
22924 incitra Zfc.A.	curved Jb A un ft jl.au Ap Scotland san.sh. Sk bog Eng. bot. 927
22925 fce'tida Jfaf.	fetid Jb A un f jl.au Ap Switzerland 1791. Sk bog Schk. ca. Hh. 96
22926 lobata Vil.	lobed Jb A un ft my.jl Ap Switzerland 1819. Sk bog Schk. ca. Hh. 96
22927 cfirvula il//.	small-curved Jb A un ft Jn.jl Ap Switzerland 1824. Sk bog Schk. ca. D. 17
22928 arcuaria L.	IV. BRIZO'I'DEA.— <i>Androgynous, spikelets spiked, many-flowered.</i>
22929 ripens Bel.	sand Jb A un Jn.jl Ap Britain salKsh. Sk bog Eng. bot 928
22930 intermedia Good.	creeping Jb A un Jn.jl Ap Switzerland 182a. Sk bog Schk. ca. J. 11
22931 schamodes Dec.	intermediate Jb A un I Jf J Ap Britain mar. Sk bog Eng. bot. 2042
22932 Schreberi W.	Rush-like Jb A un 1 my.jl Ap Germany 1823. Sk bog
22933 frizoides L.	Schreber's Jb A un i Jl Ap Germany 1800. Sk bog Host gr. 1. 46
22934 ovMis Good.	Briza-like Jb A un Y my.jl Ap Germany 1815. Sk bog Host. gr. 36.47
22935 leporina L.	ov&spiked M A un 2 Jn.jl Ap Britain mar. Sk bog Eng. bot. 306
28936 fagopodioides W.	Swiss, hare Jb A un J J Ap Switzerland 1819. Sk bog Schk.ca.FFT.129
22937 jecofria W.	Harefoot-like m A un J J Ap N. Amer. 1805. Sk bog Schk. ca. Yyy.177
22938 nemorbsa Lam.	Broom Jb A un J J Ap N. Amer. 1812. Sk bog Schk.ca.Xxx.175
22939 vulpina L.	grove Jb A un J J Ap Germany 1824. Sk bog
22940 stipita MM.	fox Jb A un J my.au Ap Britain mar. Sk bog Eng. bot. 307
22941 divisa Hud.	crowded Jb A un J my.au Ap N. Amer. 1825. Sk bog Schk.ca.Hhh.132
22942 muricata L.	divided Jb A un J my.au Ap Britain sal. m. Sk bog Eng. bot. 1096
22943 norvégica Wahl.	greater prickly Jb A un 2 my.jh Ap Britain moi. p. Sk bog Eng. bot. 1097
22944 diviflora Good.	Norwegian Jb A un lft my.jn Ap Norway 1822. Sk bog Schk. ca. 8. 66
22945 stellulata Good.	parted Jb A un 2 my Ap Britain m.s.pl. Sk bog Eng. bot 029
22946 rusea Schr.	stellulate Jb A un ft my.jn Ap Britain mar. Sk bog Eng. bot. 806
22947 axillikris Good.	Hose Jb A un 2 my.jn Ap N. Amer. 1812. Sk bog Schk.ca.Zzz. 179
22948 remonta L.	axillary Jb A un 2ft my.jn Ap England bogs. Sk bog Eng. bot. 993
22949 elongate L.	remote Jb A un 2 my.jn Ap Britain groves. Sk bog Eng. bot. 832
22950 cGrtia Good.	elongated Jb A un 1 my.jn Ap England mar. Sk bog Eng. bot. 1920
22951 faetic* W.	short Jb A un 1 Jn Ap Britain pools. Sk bog Eng. bot. 1386
22952 tenella Schk.	fodder Jb A un 2 Jn Ap N. Amer. 1318. Sk bog
straminea Donn	slender Jb A un J J Ap Scotland xnoun. Sk bog Schk.ca.23.6.104
22953 longipes D. Don	
22954 /ohacea L.	long-peduncled Jb A I Jn Jl Ap Nepal 1823. Sk bog
22955 straminea W.	Hay-grass-/i*e Jb A un 2 Jn Ap Sweden 1810. Sk bog Schk.ca.P.p. 104
	straw Jb A un Jn.jl Ap N. Amer. 1803. Sk bog Schk.ca.Xxx.174
22956 multiflora MM.	V. PANICULAVJK.— <i>Androgynous, spikelets panicle.</i>
22957 teretiflora Good.	many-flowered Jb A Jn 1A my.jn Ap N. Amer. 1812. Sk bog Schk.ca. L11.144
22958 parad6xa W.	rather rounder M A un 2 my.jn Ap Britain bogs. Sk bog Eng. bot 1065
8SB59 paniculata L.	paradoxical Jb A un lft my.jn Ap Austria 1823. Sk bog Host gr. 1.57
22960 appressa R. Br.	panicle Jb A un 3 Jn.jl Ap England bogs. Sk bog Eng. bot 1064
	close-spiked M A un 2 my.au Ap N. S. V. 1802. Sk bog
22961 bicolor AU.	VI. RACEM&9E.— <i>Androgynous, spikelets racemose.</i>
22962 atrata L.	two-colored M A un If my.jn Ap M. Cenis 1810. Sk bog Schk. ca. Aaaa. 181
	blackened M A un lft Jn.jl Ap Britain al. me. Sk bog Eng. bot. 2044
22963 thuringiaca W.	VII. THIRINGI"ACA.— <i>Terminal spikes male, the others androgynous.</i>
Thuringian Jb A un lft my.jn Ap Germany 1810. Sk bog Schkxa.P.pp.155	
22964 GLAREO'B£.— <i>Terminal spikes androgynous, the others female.</i>	
22965 Buxbaumii Wahl	Buxbaum's Jb A un 1 my.jn Ap Sweden l>2l. Sk bog Schk.ca.X.Gg.76
22966 glarctsia Wahl	gritty 4b A un 1 my.jn Ap Norway 1816. Sk bog
22966 microstachya Ehrh.	small-spiked Jb A un 1 my.jn Ap Switzerland 1822. Sk bog Schk. ca. 6.11
IX. CLANDESTINE.— <i>Spikes of distinct sexes; male ones solitary, female ones sessile or subsessile: scapes sheathed with membranous bracts.</i>	
22967 filbae,	white Jb A un 1 my.jn Ap Austria 1818. Sk bog Schk. ca. 0.55
22968 clandestina Good.	clandestine Jb A cu ft ap.my Ap England sun ro. Sk bog Eng. bot. 2124
22969 inversa R. Br.	inverted Jb A cu 1 Jn.jl Ap N. Foll. 1818. Sk bog
22970 digitata L.	fingered Jb A un 1 Jn.jl Ap England woods. Sk bog Eng. bot. 615
22971 wlanitincta Lam.	Plantain-like Jb A un ft my.jn Ap N. Amer. 18/5. Sk bog Schk. ca. U. 170
22972 Frascaniifl H. K.	Eraser's Jb A or ft apjn Ap N. Amer. 1809. Sk bog Bot. mag. 1391
X. PILI/LIFERJE.— <i>Spikes of distinct sexes; male ones solitary, female ones sessile or subsessile: culm leafy.</i>	
22973 pilumfera L.	piluliferou Jb A un 1 ap.jn Ap Britain hea. Sk bog Eng. bot. 8
22974 fimbriata Schr.	fringed Jb A un 1 apjl Ap Switzerland 1819. Sk bog
22975 mucronata All.	mucronate Jb A un 1 Jn.jl Ap Switzerland 1819. Sk bog
22976 lucofum W.cn,	grove Jb A un U.ap.jn Ap N. Amer. 1825. Sk bog
22977 coquina IV.	•4flM Jb A un 1 ap.jn Ap Germany 1824. Sk bog Schk. ca F. 29
22978 cilifera W.	ciliated Jb A un 1* ap.jn Ap Germany 1812. Sk bog Schk. ca. I. 42
22979 prae'cox Jac.	early Jb A un 1 ap Ap Britain dr.pa. Sk bog Eng. bot. 1099
22980 tomentosa L.	downy-fruited Jb A un 1 Jn Ap England mea. Sk bog Eng. b.2046
22981 extensa Good.	extended Jb A un 1 my.jn Ap Britain sea co. Sk bog Eng. bot. 833
22982 flava L.	yellow Jb A un 1 my.jn Ap Britain bogs. Sk bog Eng. bot. 1294
22983 CE'deri EhrlW	GSder's M A un J M Ap England m.me. Sk bog Eng. bot. 1773
22984 fulvzGood.-	tawny Jb A un J J Ap Britain mar. Sk bog Eng. bot 1295
XI. PANI'CEA.— <i>Spikes of distinct sexes; male ones solitary; upper female ones sessile or subsessile, lower ones stalked: culm leafy.</i>	
22985 distans L.	distant Jb A un IJ Jn Ap Britain mar. Sk bog Eng. bot 1234
22986 binervis Sm.	twi'-nerved Jb A un 2 Jn Ap Britain dr.he. Sk bog Eng. bot. 1205
22987 saxatilis X.	rff Jb A un 2 Jn Ap Greenland 1812. Sk bog Schk.ca. I&Tt.40
22988 pilla Good.	ttwet Jb A un 1 Jl Ap Scotland sc.mo. Sk bog Eng. bot. 2046

J29\$P ferruginea Schk	rusty	Jk A un	I jl	Ap	Austria 1822. Sk bog Schk.ca. M. 48
22090 Mielichhoferi U		Mielichhofer's	1 ^ u n	1 jl.au	Scotland al. roc. Sk bog Eng. boL 2293
scotica Spr.					Scotl. raar.ra, Sk bog Fl. dan. 1049
22991 speirostachya Swz.	<v,-, d,-stans Fl. dan.	banded-spiked	A A un	1 jl.au	Scotland roam. Sk bog
22092 phaeostachya Sm.	sallina G. Don	brown-spiked	Jk A un	4 jn	Austria 1810. Sk bog Schk ca.Uuu.165
22993 umbra Jforf	shady	■ A un	IJ my.jn	Ap	Europe 1820. Sk bog Schk. ca. M. 49
229W ptibsa Sco.	pilose	■ A un	1 myjn	Ap	N. Amer. 1807. Sk bog Schk.ca. Vvv.169
*2995 granularis Mhl.	grain-seeded	■ A un	Ujnjl	Ap	Britain inoi.p. Sk bog Eng. bot. 1505
% panfcfa L.	P&nic-leaved	■ A un		Ap	Europe 1800. Skbog
22997 Micheli Host	Micheli's	■ A un	1 my.jn	A1>	Hungary 1812. Skbog
2**J8 conglobata Kit.	conglobated	■ A un	1 my.jn	■	N. Amer. 1816. Sk bog Schk.ca.Hhh.lsl
22<99 rostrata AM.	beaked	■ A un	I myjl	Ap	Austria 1805. Skbog Hostgr. 1.71
2^KX nitida Jforf	glossy	■ A un	1 my.jn	Ap	N. Amer. 1805. Sk bog Schk.ca.FitT 198
SgOOL inceps Mhl.	two-edged	■ A un	1 jl.au	Ap	Europe 1804. Sk bog Schk. ca. G. 35
23002 alpcstris ^ <	alpine	■ A un	f myjn	Ap	Britain bogs, Skbog Eng.bot. 1507
23003 cspitosa L.	tufted, bog	■ A un	1 my.jn	Ap	Britain mar. Sk bog Eng. bot 914
?UXtt stricta Good.	strict-l't'aved	■ A un	1 ap.my	Ap	Britain wood*, Skbog Eng.bot2315
2-3005 pendula Goof.	great pendulous	Jk A un	4 myjn	Ap	Britain moun. Skbog Eng.bot2047
2300G rfgida Good.	rigid	M A un	* jn.jl	Ap	England woods. Skbog Eng.bot 944

XII. USTULAYK. — <i>Spikes of distinct sexes ; male ones solitary, female ones stalked ; culm leafy.</i>					
23007 capillaris L.	capillary	Jk A un)jl.au	Ap	Britain sc.mo. Sk bog Eng. bot. 2069
23008 pallwens L.	pallescent	Jk A un	1 apjn	Ap	Britain moi.p. Sk bog Eng. bot 2185
21009 ustulata IVahl.	scorched Alpine	M A un	J Jnjl	Ap	Scotland al.riv. Sk bog Eng. bot 2404
2J010 rariiflora E. S.	rare-flowered	Jk A un	{ }^"	Ap	Scotland seal. Skbog Eng.bot.2516
23011 Hmosal.	mud	Jk A un	Ij	Ap	Britain sp.bo. Sk bog Eng. bot. 2043
230K Pseudocorypha 8L.	BaardtCyperusii*	A un	3 ln^jl	Ap	Britain mar. Sk bog Eng. bot. 249
23013 flxuJisa MM	bending	Jk A un	2 jnjl	Ap	N. Amer. 1807. Sk bug Schk.ca.Ddd. 124
2KJH sylvatica Hud.	wood	Jk A un	3 myjn	Ap	Britain woods. Sk bog Eng. bot. 99J
23pi5uncea W.en.	Rush-like	Jk A un	2 my.jn	Ap	N. Amer. 18.U Sk bog
23016 strigosa Good	strigose	Jk A un	2 ap.my	Ap	England woods. Sk bog Eng. bot 944

XIII. PALUDOS* <i>Spikes of distinct sexes ; male ones not solUarp, two or more ; culm leafy.</i>					
23017 recurva Good.	recurved	ilk A un	1 my.jn	Ap	England hca. Sk bog Eng. bot. 1506
23018 nfitans Host	nodding	Jk A un	2 jnjl	Ap	Austria 1815. Sk bog Host gr. 1.83
23019 acuminata IV.	acuminated	Jk A un	2 jnjl	Ap	Mna. 1818. Sk bog Host gr. 1.97
23020 filiformis Good.	filiform	Jk A un	2 jnjl	Ap	Britain bogs. Skbog ling.bot.904
^3021 aquitilis WahL	water	Jk A un	yjnjl.	Ap	Lapland 1813. Skbog
23022 aedta L.	acute	Jk A un	2 my.jn	Ap	Britain wat.pl. Sk bog Eng. bot. 580
23023 paludissa Good.	marsh	Jk A un	2 myjn	Ap	Britain watpl Sk bog Eng. bot. 807
23024 riparia Good.	river.bank	Jk A un	2 apjn	Ap	Britain riv.ba. Sk bog Eng. bot. 579
%*J5 vsickrial.	bladdery	Jk A un	2 myjn	Ap	Britain mar. Sk bog Eng. bot 779
23025 ampullacea Good.	cruet-shaped	Jk A un	2 myjn	Ap	Britain bogs. Sk bog Eng. bot. 780
23027 <ecilina IK	Rye-like	Jk A un	2 myjn	Ap	Scotland bogs. Sk bog Schk.ca. 5. &>
21028 Aordefirms Host	Barley-formed	Jk A un	2 Jnjl	Ap	France 1805. Sk bog Schk.ca.Ddd.121
■3029 stictocarpa Sm.	dotted-fruited	^ A un	1 Jnjl	Ap	Scotland moun. Sk bog
23030 angustifolia Sm.	narrow-leaved	3t A un	1 jnjl.	Ap	Scotland mar. Sk bog
23031 hirta L.	hairy	Jk A un	2 my.jn	Ap	Britain wat.pl. Sk bog Eng. bot. 685
yle	smooth-stalked	Jk A un	2 myjn	Ap	Britain bogs. Sk bog Eng. bot 1387
*83 cmnua Lam.	hairy	Jk A un	3 myjn	Ap	N. Amer. 1807. Sk bog Schk.ca.Eee.125
23034 aallma JForf.	salt-marsA	Jk A un	1 jn	Ap	Norway ... Skbog
■3035 amplexocarpa W.	obtuse-fruited	Jk A un	1 jnjl.	Ap	Britain ... Sk bog Mic gen 62,32.12
23036 bullata W.	blistered	Jk A un	1) jn	Ap	N. Amer. 1811. Sk bog Schk.ca.Uuul66

XIV. DIBBLE. — <i>Douffut.</i>					
■3036 brachystachya Schk.	short-spiked	1 A un	I jnjl	Ap	SwitzerT. 1819. Skbog
sphrorostachya ScfU.	round-spiked	W A un	I jnjl	Ap	Switzerl. moun. Sk bog

■ KOBRE*S/il W. KOBREBIA. (<i>De Kobres, a German, and great promoter of botany.</i>) <i>Cyperdcete. 1.</i>					
erfoina W.	Sedge-like	M E2 un	} jl	Ap	Britain ... Sk bog Eng. bot 1410
ScheneNius monolca E. S.					

2601. TJNCI'NIA Pers.	UNCISIA.	(Offkinos, a hook; awn hooked.)	<i>Cyperhcetc. 2.-9.</i>
EUO/>hlco'ides/ra.	Cat's-tail-like	Jk El un	1 jl
***J5 jamaicensis Pers.	Jamaica	Jk ES un	1 jl
Caxex uncinata Swz.			Ap
		Jamaica	1816. D bog Schk.ca.l3.G.3O

INDIAN CORN.					
Mays	Jk O ag	2 jnjl	Ap	America 1552. S rm Lam. il. 749	
Curagita Mol.	Jh O >g	1 jnjl	Ap	Chile 1824. S r.m	
■x L Joi r8 FeARS.	(Applied by Theophrastus to a reed-leaved plant.)				
■x rhyra Z.	common, tear	Jb ES cu	2 jnjl	Ap	E. Indies 15P6. S l.p Rot. mag 2479
agentris W.		Jb cu		Ap	E. Indies 1812. S l.p Ru. am. 6. 9. 1
e x i mu		Jb cu		Ap	China 1816. S l.p Jac. ic. gram.
Koenig's		Jb cu		Ap	R Indies 1818. S l.p
iteed-like		JiikEScu	Iau	Ap	Mexico 1818. S l.p
inacea Lam.					

■ TRIPASACUM L.					
TRIPASACUM.	(Tyipsis, a grinding; flour made from grains.)				
Dactylus-like	Jb A un	4 au	Ap	Virginia 1640. D pi Lam.il.750	
single-spiked	Jk A un	2 au	Ap	N. Amer. 1825. D pl W. h. b. 1	

2605. HETEROPOV3ON Jichb.	HBTEROPOOON.	(Heleros, varioua, pogon, a beard.)	<i>Gramineee. 2.</i>
Allidnii Bcauv.	Allioni'a	2 au	Ap
graberto**			Switzerl. 1800. D co Al. ped. 91,4
23052 contortus Bcauv.	twisted	JU QS cu	2 jl.au
Andropgon contortus L.			Ap E. Indies 1817. D l.p Schk han 3+2.A-

2606. OLY'R'A W.	OLYBA.	(A sort of grain mentioned by Greek authors.)	<i>Gramiivw* 9 *</i>
23053 latifoliaL.	broad-leaved	m ES un	3 jl
paniculata Swx.			Ap W. Indies 1783. Sk s.p
■J5 pauciflbra Swx.	few-flowered	JUb 2Sun]j1
			Ap W. Indies 1817. Sk.s.p

CLASS XXI. ORDER 4.

MONGECIA TETRA'NDRIA.

2607. *jil'LNUS Tou.* ALDER. (*Al*, near, /an, edge of a river, *Celt.*; habitat.) *Atnentacea.* 15 — 18.

23055	<i>glutinosa L.</i>	clammy	ec 25	mr.ap	Ap	Britain	wat.pl.	L m.s	Eng. bot 1508
2	<i>laciiniata</i>	fringed-taiwrf	or 25	mr.ap	Ap	Britain	■	L m.s	W. ar. 44
3	<i>ffeliis variegatis</i>	variegatcd-lvd	or 25	mr	A p	Britain	...	L m.s	
4	<i>miercifolia</i>	Oak-leaved	or 25	mr	Ap	Britain	...	L m.f	
5	<i>indsa</i>	<i>cut-leaved</i>	or 25	mr	Ap	Britain	■	L m.s	
6	<i>emargininta</i>	emarginate	or 25	mr	Ap	Britain	■	L m.s	
23056	<i>oblongsta JV.</i>	<i>oh\ong,lcaved</i>	or 20	jl	Ap	S. Europe	1730.	L m.s	
2	<i>cllpatica</i>	elliptic-taivfti	or 20	jl	—	—	—	L m.s	
23057	<i>incima JV.</i>	<i>hoary-leaved</i>	or 20	jn	—	Europe	1780.	L l.p	
2	<i>angulata</i>	angular-fcat/t'd	or 20	jn	Ap	—	—	L l.p	
3	<i>pinnate</i>	pinnate	or 26	jn jl	—	Europe	—	L co	
23058	<i>glaifica Mr.</i>	glaucous	or 20	my jl	Ap	N. Amer.	1820.	L co	
23059	<i>undulata JV.</i>	<i>vne-leaved</i>	or 20	my,jn	Ap	N. Amer.	1782.	L l.p	
23060	<i>serrulata IV.</i>	<i>Baxv,lcaved</i>	or 20	mr	Ap	N. Amer.	1769.	L l.p	Ab. ins. 2,92
23061	<i>cordifolia Ten.</i>	<i>heart-leaved</i>	or 20	my,jn	Ap	Naples	1818.	L co	Bot. cab. 1231
23062	<i>macrocarpa Lo. C.</i>	long-fruited	or 20	my,jl	Ap	—	—	L co	
23063	<i>subrotunda Desf.</i>	roundish./eawdj	or 23	my,jl	A p	—	—	L co	
23064	<i>rubra Dett.</i>	red	or 20	my,jl	A p	—	—	L co	
23065	<i>macroph^lla 2ej'l</i>	long-leaved	or 20	my,jl	A p	Naples	—	L co	
23066	<i>canadensis Lo. C.</i>	Canadian	or 20	my,jl	Ap	Canada	—	L co	
23067	<i>oxyacanthif6liaXo.C.</i>	Oxycanth.-lvd	or 20	my,jl	Ap	—	—	L co	
23068	<i>pumila Nois.</i>	dwarf	or 10	my,jl	Ap	—	—	L co	

2608. 5 ETULA L.

BiRen.

(Betu, its Celtic name.)

Amentieece.

la -19.
Eng. bot. 2198

23069	<i>filba L.</i>	<i>common white</i>	tm 40	ap,jn	Ap	Britain	moi.w.	S co	
2	<i>verrucosa Ehrh.</i>	warty	* tm 40	ap,jh	Ap	Britain	...	S co	
3	<i>dalecarlica L.</i>	Dalecarlian	tm 40	ap,jn	Ap	Europe	...	S co	
4	<i>macrocarpa Lod.</i>	long,fruited	tm 40	ap,jn	Ap	Europe	...	S co	
23070	<i>jendula Roth</i>	pendulous	or 40	ap,jn	Ap	Britain	woods.	S co	
23071	<i>populifolia H. K.</i>	Poplar,leaved	or 30	jl	Ap	N. Amer.	1750.	L co	Mic. ar. 2,2
23072	<i>excelsa II. K.</i>	tall	tm 60	my	A p	N. Amer.	1767.	S co	Den. br. 95
23073	<i>daurica PaU.</i>	Daurian	tm 30	jl	A p	Siberia	1786.	L co	Pal. ros. 1,39
23074	<i>rtgra L.</i>	black	tm 60	jl,au	A p	N. Amer.	1736.	L co	Den. br. 153
23075	<i>Ianul6sa Mx.</i>	woolly	tm 70	jl,au	Ap	N. Amer.	1817.	L co	Mic. ar. 2,3
23076	<i>rftbra 3/J.</i>	red	tm 60	jl,au	Ap	Canada	—	L co	Bot. cab. 1248
23077	<i>papyracea H. K.</i>	paper	tm 50	jn	Ap	N. Amer.	1750.	L co	W. ar. 2,1
23078	<i>fcnta L.</i>	pliant	tm 50	jl	Ap	N. Amer.	1759.	L co	Den. br. 144
23079	<i>carpinifolia fArA.</i>	Hornbeam-lvd	tm 50	jl	Ap	Scotland	moi.h.	L co	Eng. bot. 2326
23080	<i>nana L.</i>	<i>smooth</i> dwarf	cu 4	my	A p	Switzerl.	1819.	L co	
2	<i>macroph^ua Schi,</i>	long,leaved	or 6	my	A p	N. Amer.	1762.	L s.p	Jac. vin. 2,128
23081	<i>pumila L.</i>	<i>hairy</i> dwarf	or 6	ap,my	Ap	Turkey	...	L s.p	
23082	<i>p6ntica //or^.</i>	Pontic	or 12	ap,my	Ap	Hungary	1820.	L co	Den. br. 94
23083	<i>ovata Schrank</i>	ovate	or 6	ap,my	Ap	L co	Den. br. 96
23084	<i>fruticosa i>n/.</i>	shrubby	or 6	ap,my	Ap	Siberia	1818.	L co	Den.br. 97
23085	<i>pubescens Ehrh.</i>	pubescent	or 30	ap,my	Ap	Germany	1812.	L co	
23086	<i>tristis Lk.</i>	sad	or 10	ap,my	Ap	Kamtsch.	...	L co	
23087	<i>latea Mx.</i>	yellow	or 20	my	Ap	N. Amer.	...	L co	

2609. BXV8 L.

Box TREE.

(Pyknos, dense; wood.)

Euphorbihccc.

4.-5.

23088	<i>baleitrica Zam.</i>	Barlearic	or 8	jl	Y.G	Minorca	1780.	C co	
23069	<i>semperf'ircns L.</i>	<i>com.</i> evergreen	or 8	ap	Y.o	England	ch.hil.	C co	Eng. bot. 1341
2	<i>angustifolia</i>	narrow-leaved	or 8	ap	Y.G	C co	
5	<i>suffruticosa</i>	subshrubby	or 1	—	Y.C	Skco	
23090	<i>chinensis Lk.</i>	Chinese	or 3	n	Y.G	China	1802.	C co	
23091	<i>australis Cun.</i>	southern	L-J or 6	—	...	N. Holl	1820.	C l.p	

2610. SARCOCOCCA B. R. SARCOCOCCA. (*Sarx*, flesh, *kokox*, a berry : substance of.) *Euphorbiacce.* 1
230D2 prmf6rmis B. R. Plum-formed »i_J or 4 jn,jl Pa.Y Nepal 1820. C p.l Bot.reg.f.

9611. CI'CCiL L.

Cicc/i.

{Peter Cicca, who wrote in 1553.)

Euphorbiiceat

1.—2.

23093	distichaL.	two-rowed-/w*	± n	fr	20	...	G	K Indies	1796. C p.l Jac.sc. 2,194
-------	------------	---------------	-----	----	----	-----	---	----------	---------------------------

•2612. A/O'RUSZ.

MULBERRY.

(Alax, black, *Celt.*; color of fruit.)

Urtkear.

12.—17.

23094	<i>alba /.</i>	white	rtt 30	jn	A p	China	1596.	L co	Schk. han. 3,290
2	<i>macrophylla</i>	long-leaved	clt SO	jn	A p	China	—	L r.m	
3	<i>nana</i>	dwarf	clt 10	jn	A p	China	...	L r.m	
4	<i>ovalifolia</i>	oval-leaved	clt SO	jn,jl	A p	China	...	L r.m	
5	<i>fosc</i>	Rose-like	clt 20	jn,jl	Ap	China	—	L r.m	
23095	constantinopolitina De', Constanti.		clt 15	jn	Ap	Turkey	1818.	L r.m	
(23096	<i>sinensis Hort.</i>	Chinese	clt 20	—	A p	China	1825.	L r.m	
23097	<i>scabra L.</i>	scabrous	or 20	jn	Ap	N. Amer.	1817.	L r.m	
23098	<i>pebnsylvanica Not.</i>	Pennsylvanian	or SO	jn,jl	Ap	Pennsylva.	1820.	Is r.m	
23099	<i>franca L.</i>	Italian	clt 20	jn,jl	Ap	Italy	1817.	L r.m	
23100	<i>butinna Not.</i>	dwarf	clt	—	Ap	1824.	L r.m	
23101	<i>tatarica Pall.</i>	Tartarian	or 20	jn	Ap	Tartary	1784.	L co	Pal. ros. 9,52
23102	<i>nigra Pall.</i>	common black	ft SO	jn	A p	Italy	1548.	L co	Dcn.br. 159
23103	<i>raZ.</i>	red	or 10	jn,jl	A p	N. Amer.	ML	L r.ra	
23104	<i>mauritiana Jac.</i>	Mauritian	Lor SO	—	Ap	Mauritius	1821.	C l.p	Jac.ic.&617
23105	<i>indica L.</i>	Indian	LJ or 20	—	Ap	E. Indies	IS24.	C l.p	Ru. am. 7.5

*2613. BCEHME'R/^ Jac. BOBUMEBIA. (George Rudolph Bahmer, a German botanist.¹) *VrHcea?* 14. — 111

I. OpposmphiLi^, — Leaves opposite.										
23106 cylindrica W.	cylindrical	A un	4	jn.au	G	Virginia	1759.	Sk s.p	SI. jam. 1.	82.2
rotundi folia D. Don	round-leaved	SuMun	3	jl.au	G	Nepal	1818.	D s.l		
elongata Fis.	elongated	O un	1	au.a	G	Brazil	1817.	S co		
broad-leaved	* CD un	2	...	G	Jamaica	1820.	C s.p			
swill macrophyllea D. Don	* U un	4	...	G	Nepal	1820.	C e.p			
long-leaved	I U un	6	...	G	Nepal	1819.	C fl.p			

II. AvtERjiiFbum. — Leaves alternate.										
23112 rutescem Jac.	rubescent	* i_J or	10	f.ray	G	Canaries	1779.	C s.p	Jac.fr. 5.1	
oof. i/rteca arborea L.										
23113 ramiflora Jac.	branch-flwd	i Q un	8	f.my	G	Jamaica	1823.	C co	Jac.am. 157	
frondosa I. Don	leafy	i un	4	..	G	Nepal	1819.	C s.p		
fruticosa 77 <i>bi</i> .	hairy	tL O un	2	...	G	Jamaica	1817.	C s.p		
fruticosa 77 <i>bi</i> .	fruticosa	* i un	3	...	G	Nepal	1819.	C s.p		
U7wlicifoliai. i/	Willow-leaved	^ A un	6	s.n	G	Nepal	1819.	D co		
J18 lateriflora A/A.	Bide-flowring	^ A un	1	f my	G	N. Amer.	1820.	Sk co		
interrupta W.	interrupted	£ 23 un	1	au.s	G	E. Indies	1816.	D co		

PILEA. (Pilos, a cap; nature of one of the divisions of perianth.) <i>Urticete.</i> 1.—3.										
musebsa Lindl.	mossy	fcx	23 pr	ap.my	G	W. Indies	1793.	S co	Lind. col. 4	

•2615. NETTLE. (J7ro, to burn; stinging qualities.) *Urticee.* 46.—114.

I. OPPOSITIFOLIE. — Leaves opposite.										
23121 pilulifera L.	Roman, pill-bearing	O w	lijn.au	G	England	rub.	S co	Eng. bot. 148		
23122 balearica J.	Balearic	O un	li_jnl	G	Balearic	1.1733.	S co	Black. 321.1		
g 23 convexa Horn.	convex	O un	U_jl	G	1824.	S co			
Brlpta/*wi.	black.hnd	cu	2' j.l	Y	Nepal	1818.	D co			
'epressa Swx.	depressed	cu	Jamaica	1820.	D co			
Dflirtii L.	Dodart's	m	fl	* 8^u	S. Europe	1613.	S co			
crophilia Stvz.	small-leaved	O un	jl	Jau	W. Indies	1793.	D co	SL. jam. 1.93.2		
pumil L.	dwarf	£ 23 cu	N. Amer.	...	co			
23129 involucrata A	involucred	O un	I jl	Jau	W. Indies	1821.	D co	Bot mag. 2481		
30grandifolia L.	great-leaved	ig 23 un	1	Jau	Jamaica	1793.	L co	SL. jam. 1.83.2		
S! 27 A	fierce	p. un	3	—	N. Zeal.	1823.	s.p.l			
digitii Cun.	ant	p. J cu	20	—	N. Holl.	1823.	—			
23133 moridiles Cun.	torus-like	f J cu	30	—	N. Holl.	1823.	P.1			
23134 reticulata Swx.	reddish	S J cu	10	G	Jamaica	1793.	E co	Bot. mag. 2567		
23135 rdfa Swx.	small stinging	Q W	1 jns	—	Britain	cltgr.	S co	Eng. bot. 1236		
23136 arena L.	arena	1 jns	—	—	Britain	wa.gr.	D co	Eng. bot. 1750		
23137 dioica L.	common, diocious	A	1 jns	—	Siberia	1817.	D co			
23138 angustifolia Fis.	narrow-leaved	—	2 jls	—	Spain	1820.	D co			
33139 membranacea Poir.	membranous	A un	21	qn	Jamaica	1793.	E co			
23140 diffusa Swx.	diffuse	i 23 un	1	JlJ.S	Surinam	1803.	S co	Jac. sc. 3.388		
23141 crassifolia W.	thick-leaved	j 23 un	1	jls	Canada	1656.	D co	Pluk.al.239.2		
23142 ardens Lk.	burning	O un	1 ll-s	—	Canada	1816.	D co			
23143 chamaeroides Ph.	Germander-like	C un	ijl.au	—	China	1735.	D p.l	Jac. vin. 2.166		
23144 nummularifolia Stvz.	Moneywort-lvd	O un	1 jn.jl	Ap	Jamaica	1820.	S co	SL. jam. 1.131.4		
155* cannabina L.	Hemp-like	A cu	3 jls	Ap	Siberia	1749.	D co	Am. rut 249.25		
23145 rugosa Swx.	wrinkled.5/M	DQun	2 my.jl	Ap	Jamaica	1793.	S co			
23146 nudicaulis 8Sp.	naked-stalked	n. I I un	■ my.jl	Ap	Jamaica	1793.	C co			
		A un	3 jn.au	Ap	Huds. a	1782.	D co			
		—	—	Ap	Jamaica	1793.	C co	SL. jam. 1.93.1		
		tL un	1 jls	Ap	Jamaica	1815.	C co			
Pellia L.*.	ciliated	fl un	1 jls	Ap	E. Indies	1820.	C co			
scabrella Rox.	pretty	a un	—	Ap	E. Indies	1815.	C co			

11. ALTERNIFULUE. — Leaves alternate.										
23153 iestuans L.	raging	O un	1	jnjl	Ap	Surinam	1803.	S co	Jac. sc. 3.388	
	Canada	^t A un	3	au.o	Y	Canada	1656.	D co	Pluk.al.239.2	
	divaricate	5 A cu	2	o.n	Y	Canada	1816.	D co		
	movr-white-lvd	5 A un	6	au.s	Ap	China	1735.	D p.l	Jac. vin. 2.166	
	berry-bearing	tuQun	4	jl.au	—	S. Amer.	—	Sk.s.p	Bot rep. 454	
	small-headed	H CD un	8	jl.au	—	Java	1810.	C co		
	caracasana Jac.	Caraccas	—	—	Ap	Caraccas	1824.	C co	Jac. sc. 386	
	caravelLina Schrank	Caravellan	O un	—	Ap	S. Amer.	1825.	S co		
	dongataza Z.	lengthened	O un	3	jl.au	Philipp. Is.	1823.	S co		
g 23 T M Mal.*.	various-leaved	—	3	aus	—	E. Indies	1823.	S co		
hylla Wai.	variable-leaved	tL cu	2	jl.au	—	E. Indies	1819.	C co	R.mal.2.41	
g 6* horrida Z.*.	horrid	O un	3	au.a	Ap	Nepal	1821.	S co		
"borescens Lt.	arborescent	* I un	8	aus	—	Manilla	1822.	C co		
II in Wndra Rox.	pentandrous	£ 23 un	2	aus.s	—	E. Indies	1800.	D co		

(The wife of Cephalus.) <i>Urticea?</i> 2. — 19.										
23167 punctita Ham.	1AI un	1	jl.au	G	Nepal	1820.	D co			
23171 ciftii	iAJ un	1	jl au	G	Nepal	1820.	D co			

SSifte PA C? YSA'NDRA iV*. PACHYSANOBA. {Pachys, thick, aner, man; stamens.) <i>Euphorbidcea.</i> 2.										
23172 Pr? finbch8iv*.	trailing	A A P'	i mr.ap	W	N. Araer.	1800.	D s.p	Bot. reg. 33		
23170 conicea 7/ooA.	coriaceous	iIZIpr	4	jn.jl	W	Nepal	1822.	C co	Hook. ex. fl. 148	

ASfIS. DTCVTIS L. Duma. (Dis, double, ous, ear; two appendages at b.isc of florets.) <i>Chenophdce.</i> 3.-4.										
ceratodes W.	hom-like	»	or	2	mr	Ap	Siberia	1780.	L ep	Jac. ic. 1. 189
atriplicia Spr.	Acriplex-like	O w	1	jl.jl	Ap	Britain	fields.	S co	Eng. bot. 232	

^triplex pedunculata L.	triplex, pedunculata	O un	1	jn.jl	Ap	—	—	1826.	S co	
-------------------------	----------------------	------	---	-------	----	---	---	-------	------	--

o.2619. EMPLEURUM H. K. EMPLEURUM. {En, in, Pleuron, membrane; seeds atUched to) <i>Didsmete</i> I.										
^^ serrulatum H. K.	serrulated	• i_J or	3	jn.jl	Pk	C. G. H.	1774.	C pl	Ex. bot2.' 63	

AUCVB L.	AUCUBA.	(Its name in Japan.)								
75 japonica L.	Japan< blotch-lvd >	or	6	my.jn	Ap	Japan	—	—	Rhamneat? 1.	C co Bot mag. 1107

2621. LITTOR'ELLA L. SHORE-WEED. (*Imitus*, the shore j place of growth.) *Plantaginete*. I.
 23176 lacūstris L. lake, *Plantain-lud* * A P' i J^{n*}au " Britain w.s.a.p. D p.l Eng. bot. 468
2622. SERPI'CULA L. SERPICULA. (Dim. from *serpo*, to creep; habit and size.) *Halarbge**. 1.—3.
 23177 repens L. creeping *r IAJ P* i jUu W C. G. H. 1789. D p.l Lam. il. 758
2623. MACLITRif Nut OSAGE ORANGE. (*William MacLure*, a North American geologist) *Urticee*. 3.
 23178 aurantia Nut. common Orange Y fr 20 ... Ap N. Amer. 1818. L p.l Lainb. pi. sup.
 23179 tinctoria D.Don dyes', *Fustick* f. or 20 ... Ap W. Indies 1739. C r.m
MOTUB tinct&ria L. *Broussonetia* tinctoria Art.
- 23180 Plumiera G. Don Plunder's f. or 20 ... Ap W. Indies 1824. C r.m PI. ic. 204
 A/brus Plumiera Brm. *Broussonetia* Plumiera Spr.

CLASS XXI. ORDER 5.

MONCFIA PENTA'NDRIA.

2634. EXOCA'RPOS Lab. EXOCARPOS. (*EXO*, outside, *karpos*, fruit j seed on fleshy recept) *Osyrideas*. 3—6.
 23181 cypressiformis IM. Cypress-shaped t i J tm 40 ... Ap V. D. L. 1824. C p.l Lab. vo. 14
 23182 humifusus R. Br. trailing &, i | or 2 ... Ap V. D. L. 1824. C p.l
 23183 strictus R. Br. strict * i J or 5 ... Ap N. Holl. 1822. C p.l

2625. SCHIZ A'NDR A Mx. SCIIIZANDRA. (*Schizo*, to cleave, *aner*, man; stamens split) *Menisp̄mece*. 1.
 23184 coccinea Mx. acnct-flowered JU i J or jn.jl S N. Amer. 1806. L s.p Bot mag. 1413

2626. FRANZEMLM Cav. FRANZERIA. (*Antony Franzer*, M.D., a botanist.) *Comp. Ambrosidcear*. 2. - 4.
 23185 artem'sio)des W. Artemisia-like * J un 6 jl.s G Peru 1759. C p.l W. h. b. 2
 2J186 ambrosicAdes Cav. Ambrosia-like • U u n 4 jl.s G Mexico 1796. C p.l Cav. ic. 2. 200

2627. XA'NTHIUM L. XANTHIUM. (*Xanthos*, yellow; used by Greeks to dye the hair.) *Comp Ambr.* 7.
 23187 Strumuriūm L. Small Burdock O un 3 jl.s G England dungh. S co Eng. bot. 2544
 23188 orientate L. Oriental O un 4 jl.s G China 1685. S co Schk.han.3.291
 23189 spinijsum L. spiny O un 3 jl.s G S. Europe 1713. S co Her. par. 246
 23190 cchinatum Mur. hedgehog O un 3 jl.s G S. Europe 1713. S co Mur. 1784. cic.
 23191 canadēse Mil. Canadian O cu 1 jl. G N. Amer. 1700. S co
 23192 macrocarpum Dec. long-fruited O cu 1 jl. G France 1817. S co
 23193 catharticum Kth. cathartic O cu 1 jl. G SxAmer. 1824. S co

S628. ifMARA'NTUS L. AMARANTH. (*A,priv.,maraino*, to wither; durable flowers.) *Amarant/iuce*. 60.

- I. TRIA'NDRIA. — *Flowers triandrous.*
- | | | | | | |
|----------------------------|-------------------|-----------------|-------|-------------------|-----|
| 23194 tonuifolius W. | fine-leaved | O un 1 jl.s | G | E. Indies 1801. S | co |
| 23195 angustifolius Bieb. | narrow-leaved | O un 1)jl.s | G | Levant 1723. S | co |
| 23196 albus L. | white | O un 1} jl.s | G | N. Amer. 1778. S | co |
| 23197 graedzans L. | Greek | O un lftJU | G | N. Amer. 1759. 8 | co |
| 23198 melanchōticus L. | melancholy | O or U jns. | P | E. Indies 1731. S | co |
| 23199 tricolor L. | three-colored | O or 2.jns. | R.T. | E. Indies 1548. S | r.m |
| 23200 bfcolor Noc. | two-colored | O or 2.jns. | R.Y. | E. Indies 1802. S | r.m |
| 23201 poljeamus L. | polygamous | O or lfjl.8 | G | E. Indies 1780. S | co |
| 23202 gangticus L. | Gangetic | O un 2 jl.s | G | E. Indies 1778. S | co |
| 23203 mangostan Mdnus L. | Mangostan | Q un 2 jl.s | G | E. Indies 1801. S | co |
| 23204 polyst^chys W. | many-spiked | Q un 2 jl.s | G | E. Indies 1816. S | co |
| 2*205 tristis /. | sad, round-headed | Q un 2 jl.s | P | China 1759. S | co |
| 23206 inamac'nus IV. | unpleasant | Q un 2 jl.s | Tjl.8 | Japan 1820. S | co |
| 23207 inc6mtus W. en. | shabby | Q un 2 jl.s | G | ... 1823. S | co |
| 23208 lividus L. | livid | O un 2 jl.s | R | N. Amer. 1759. S | co |
| 23209 oleraceus L. | potherb | O un 2 jl.s | Pa.R | E. Indies 1764. S | co |
| 23210 lancea'folius Rox. | lance-leaved | O un 5 jl.s | R | E. Indies 1816. S | co |
| 23211 atropurpuricus Rox. | dark-purple | O dt 6 jlau | P | E. Indies 1820. S | co |
| 23212 fasciatus Rox. | banded | O or 3 jl.au | P | E. Indies 1816. S | co |
| 23213 bulfotus Bes. | blistered | O or 3 au.n | G | E. Indies 1822. S | co |
| 23214 2jlum L. | wild Blite | O or 2 jl.au | G | England dungh. S | co |
| 23215 prostratus Balb. | trailing | Q un 4 jlau | G | France 1739. S | co |
| 23216 spicatus Pert. | spik'd | Jk Q un 3 jl.s | G | Europe ... S | co |
| 23217 vfrldis L. > | green | J Q un 3 jl.s | G | Brazil 1768. S | co |
| 23218 /xilygonoldes L. | Polygonum-lik | Jc Q un 11 jl.s | G | Jamaica 1778. S | co |
| 23219 scandens L. | climbing | I Q un 2 jl.s | G | America 1796. S | co |
| £3820 deflexus /. | bending | 3 Q un 1 jl.s | G | ... 1805. S | co |
| 23221 pallidus Bieb. | pale | O un 1 jl.s | Pa.R | Caucasus 1817. S | co |
| 23222 lineatus R. Br. | lined | O un 1 jl.s | G | N. Holl. 1823. S | co |
| 23223 persicarioides Poir. | Persicaria-like | O un 1 jl.s | G | Nepal 1820. S | co |
| 23224 camp^stris W. | field | O un 1 jl.s | G | E. Indies 1818. S | co |
| 2*225 interruptus R. Br. | interrupted | O un 2 jl.s | R | N. Holl. 1818. S | co |
| 23225 f'acilis Poir. | slender | O un 1 jl.s | R | Guinea 1806. S | co |
| Chenopbdium caudatum Jac. | | | | Jac. ic. 2. ZU | |
| £3227 Berchtoldii Jac. | Berchtold's | O un 2 jl.s | R | Europe 1825. S | co |
- II. PENTA'NDRIA. — *Flowers pentandrous.*
- | | | | | | |
|--------------------------|------------------|-------------|-----|-------------------|-----|
| 23228 caulifldrus Lk. | stem-flowering | Q un 4 jns. | G | Nepal 1821. S | co |
| 93229 h'bridus L. | hybrid | Q un 4 jns. | G | N. Amer. 1656. S | co |
| 23230 strictus IV. | strict | O un 2 jl.s | G | ... 1793. S | co |
| 23231 celostodes Kth. | Celosia-like | O or 2 jl.s | R | S. Amer. 1818. S | co |
| 23232 caracastaus Kth. | Caracas | O or 2jl.s | R | Caracas 1818. S | co |
| 23233 paniculatus L. | panicled | O or 6 jl.s | G | N. Amer. 1798. S | co |
| 23234 sanguineus L. | bloody | O or 3 jl.s | R | Bahamal. 1775. S | co |
| 23235 retroflexus L. | bent-back-Aatred | O un 2 jl.s | G | Pennsylv. 1759. S | co |
| 23236 lseHus W. | joyful | O un 2 jl.s | R | 1799. S | co |
| 23237 flavus L. | yellow | O or 2 jl.s | L.Y | India 1759. S | co |
| 23238 chlorostachys W. | green-spiked | O un 3 jl.s | G | ... 1796. S | co |
| 23239 hypochondriacus L. | melancholy | O or 5 jl.s | D.R | Virginia 1684. S | co |
| 2*210 cruentus L. | bloody | Q un 3 jl.s | D.R | China 1729. S | r.m |

2241 hecticus <i>TV.</i>	hectic	O or 3 au	Pk	1796. S co	Wang. 25.7.13
ig ² pūmilus <i>Nut.</i>	dwarf	O un J jl.au	R	N. Amer. 1821. S co	
M curvifolius Spr.	curve-leaved	O un 1jl.au	G 1821. S co	
* ¹⁴ parisienis Schk.	Parisian	O un 1 jl.au	G	France 1817. S co	
<<45 caudatus L.	tailed, <i>Love lies bleeding</i>	O or 4 au.s	R	E. Indies 1596. S co	
o.o., 2 maximus	tree	O or 6 au.s	R 1820. S co	
WT rumentaceus Box.	frumentaceous	O or 2S au.s	R	E. Indies 1822. S co	
spinibus L.	spiny	O un 2 ls	G	India 16S3. S co	
lbus B. M.	snowy	O or 6 jl.au	R	Nepal 1819. S co	Wang. 38.4.8
ntbuadie.	giant	O un 4 jl.au	R 1817. S co	Bot mag. 2227
I rigidus Schulz	rigid	O un 1 jl.au	G 1817. S co	
Zanensis Horn.	Zano	O un 2 jl.au	R 1822. S co	
sylvestris Desf.	wood	O un 1 jl.au	G	Pyrenees 1819. S co	
aa3 specitata Bkb.	showy, <i>Caucasian</i>	O un 1} jl.au	R	Caucasus 1818. S co	

ACROGLCPCHIN Schr. ACROGLOCHIN. (*Akros*, summit, *glochij*, spear point.) *Amaranth Acea*. 1.
cfacopodioides Schr. Chenopodium-lk O un 1} jl.au G Nepal 1821. S co
ifmaranthus Acroglobchin Spr.

2330. LUFFACvr.	LUFFA.	(<i>Lovff</i> , its Arabic name.)		<i>Cucurbitacea</i> . 3.—7.
>foetida Cav.	fetid	J*. QJor 12 jn.o	P India	1812. S co Bot. mag. 1638
acuUngula Ser.	acute-angled	Jk O cu 10 jn.o	P India	1692. S co Jacvin. 3.73.74
Cicumis acufangulus L.	Egyptian	Jc fft cu 10 jn.o	P E. Indies	1739. S co Ru. am. 5.147

2.8 integrifolia MM.	entire-leaved	O un 1 ls	G N. Amer. 1816. S co
triM-teaved	O un 1 ls	G N. Amer. 1699. S co	M. h. 6.1. 4
iior/ L.	taller	O un 1.lau	G N. Amer. 1696. S co
paniculate Mx.	Mugwort-leaved	O un 1.lau	G N. Amer. 1759. S co
>>61 cumandnsc Kih.	panicled	O un 1.s	G N. Amer. 1811. S co
23265 heterowhfla MM.	sea	O un 1.lau	G Italy 1570. S co
absinthiiflbia Dettf.	Cumana	O un 1.lau	G Cumana 1825. S co
variable-leaved	O un 1.lau	G N. Amer. 1817. S co	Schk. ban. 3.292

2632. SECURKNEGA J. OrAnEITE MYRTLE. (*Securis*, axe, *nego*, to refuse; extremely hard) *Eup.* 1.—2.
83266 nitida W. shining-Zeavd f • tm 40 jnfjl W Mauritius 1793. C co Lind.col.9

2633. NAVAS W. NAJAB. (*Nadas*, a water nymph ; habitat.) *Fluviales*. 1.
23257 nfajor All. greater O cu jl.au ... Europe 1816. S co Mic. gen. 11.8. S
monosperma IV.

CLASS XXI. ORDER 6.

MON(E)VIA HEXA'NDRIA.

ZIZ A*NI A L. ZIZIVANIA. A. (Greek name of darnel, applied to this plant.)		<i>Graminea</i> . 3.-5.
iuatica iMtn. water'. <i>Canada Rice</i> + O cc 6 jl.s	G N. Amer. 1790. S co Lin. tr. 7.13	
rijicca A/x. Mill	illet-like * O ec 4 jl.au	G Carolina 1816. S 1
itans Mx.	floating O ec 4 jlau	G N. Amer. 1824. S 1
9125. PHARUS.	broad-leaved M 23 or 6 JLau	G Jamaica 1793. S r.m Br. jam. 38.3
71 latifWius L.		
SRDA L. GUETTAITOAO. (Eugène Guettard, a French botanist)		<i>Rubiaceae</i> . 7.—17.
rsiu Pers.	hairy	Peru 1820. C p.l Fl. per. 2.145. a.
R. ^{tom} confusa Pers.		
74 lucida Ai	woolly	Jamaica 1820. C p.l
ug'Aldcida Sum.	shining	Jamaica 1818. C p.l Vahlsy.3.57
longa Pers.	sweet-scented f • or 20	Jamaica 1818. C p.l Jac.am.pic.259.16
Laugerta odorata Jac.		
scabria Ven.	scabrous J • or 20	W.Indies 1818. C p.l Ven.ch. 1.1
MatthiolascabraL.		
specifsa L.	showy-flowered f n SP! 30	E Indies 1771. S p.l R. mal. 4.47,48
rugbsa Swz.	wriuMyJeaeed J Q or 20	W. Indies 1793. S r.m
		<i>Palme</i> . 4.
		Rumphius's f • clt SO ... G
		E Indies 1800. Sk r.m Ru. am. 1.17,18
		Ruffia f• or 60 ... G
		Madagascar. 1830. S r.m
		pedunculate £ O or 50 ...
		Madagascar. 1820. S r.m
		A
		Imfera Per*. wine-bearing * Q o 50 ... G
		Raphia Plinus Goe. Raphia vinifera Bcauv.
		C&COSL COCOA NUT TREE. (<i>COCO</i> , Port.; end of nut like a monkey's head.) <i>Pdnue</i> . 3.—11.
		L common, nut-bearing! O clt 50 ... Pa.G E. Indies 1690. S r.m Rox. cor. 1.73
		exubsa Mart.
		flexuous i• or 50 ... Brazil 1825. S r.m Mart palm, ic.
		plumbosa Lo. C.
		plumose I D o r 50 ... Brazil 1825. S r.m Mart.palm.ic.
		combsa Mart,/
2639. ELATE L.	ELATE.	(A name adopted from the Greeks.) <i>Palmar</i> i
S6 sylvestris L.	wood	Id or 14 ap.jn G E. Indies 1763. S r.m R. maL 3.22/25
2640. BA'CTRIS Jac.	BACTRIB,	(<i>Baktron</i> , a cane j stems made into walking-sticks.) <i>Palm*</i> 7 10
? minor Jac.	less	Q or 12 ... G & Amer. 1691. S rm JaTahm 171
Jac.	greater	D or 25 ... G Carthag. 1800. S xm J w . 2 1712

- | | | | | | | | | | | | | |
|--------|----------------------------|---------------------|---|---|---------|--------|-----|-----------|--------|---|-----|-----------------|
| 83289 | pectinata | Mart. | pectinate | £ O or | 15 | - | - | Brazil | 1825. | S | r.m | Mart. palm. ic. |
| 23290 | cuspidata | Mart. | cuspidate | £ CD or | 20 | - | - | Brazil | 1826. | S | r.m | Mart. palm. ic. |
| 23291 | macraanthia | Mart. | long-spined | £ CD or | 20 | - | - | Brazil | 1823. | S | r.m | Mart. palm. ic. |
| 23292 | guianensis | Lo. C. | Guiana | £ O or | 16 | ... | - | Guiana | 1820. | S | r.m | |
| S3293 | caryouefolia | Mart. | Caryota-leaved | £ CD or | 10 | - | - | Brazil | 1825. | S | r.m | Mart. palm. ic. |
| 2641. | LEOPOLDINJI | Mart. | LEOPOLDINA. | (The late Empress of Brazil.) | | | | Pidme. | 1.—2. | | | |
| 23294 | pulchra | Mart. | beautiful | £ CD or | 60. | . | . | Brazil | 1825. | S | r.m | Mart. palm ic. |
| 2642. | CENOCA'RPUS | Mart. | CSNOCARPUS. | (OHIO*, wine, karpos, fruit.) | | | | P&Ivye. | 1.— 8 | | | |
| 23295 | Batata | Mart. | Bataua | £ O or | 40. | . | . | S. Amer. | 1820. | S | r.m | Mart. palm ic. |
| 2643. | EUTE/RPE Gae. | EUTERPE. | (Euterpes, pleasing j | habit of trees.) | | | | Palmce. | 4.— 4. | | | |
| 23296 | globosa | Gae. | raand-fruited | £ Cjj or | 35 | - | - | Mauritius | 1819. | S | r.m | Gae.fr. 9.3 |
| | Areca | speciflta Lam. | | | | | | | | | | |
| 23297 | pisiformis | Gae. | Pea-formed | £ a or | 30 | - | - | Madagasc. | 1819. | S | r.m | Gae.fr. 193. 4 |
| 23298 | oleracea | Mart. | potherb | £ CD or | 30 | - | - | Brazil | 1800. | S | r.m | |
| 23299 | canbaAi Spr. | Canbaan | Canbaan | £ a or | 30 | - | - | W. Indies | 1656. | S | r.m | Jac. am. 170 |
| | Ai ica | oleracea Jac. | | | | | | | | | | |
| *2644. | W A L L I ' C H / J J Rox. | WALLICHLIA. | (Nathaniel IVallich, M.D., superint of bot gard. Calcutta.) | Palmce. | 1. | | | | | | | |
| 23300 | caryotoldes | Rox. | Caryota-like | i CD or | 25. | . | . | E. Indies | 1825. | S | r.m | |
| 2645. | ACROCO'MIA | Mart. | ACROCOMIA. | (Akros, top, kome, tuft; leaves.) | | | | Palmce. | 8. | | | |
| 23301 | sclerocarpa | Mart. | hard-fruited | £ o or | 40. | . | . | W. Indies | 1731. | S | r.m | Mart. palm. ic. |
| 23302 | fusiformis | Suit. | spindle-shaped | £ CD or | 40. | . | . | Trinidad | 1751. | S | r.m | |
| | Cocas fusiformis | Svn | | | | | | | | | | |
| 23303 | minor | Lo. C. | smaller | £ C] or | 20. | . | . | Trinidad | 1820. | S | r.m | Gae. fr. 9.1 |
| | Bactris | globosa minor | Gae. | | | | | | | | | |
| 23304 | globosa | Lo. C. | globose | £ o or | 20 | - | - | St. Vine. | 1824. | S | r.m | |
| 23305 | aculeata | Lo. C. | prickly | £ o or | 40 | - | - | W. Indies | 1796. | S | r.m | Jac. am. 169 |
| | Cbcos | aculeata | L. | | | | | | | | | |
| 23306 | guianensis | Lo.C. | Guiana | £ o or | 30 | - | - | Demerara | 1824. | S | r.m | |
| 23307 | fiorrida | Lo. C. | horrid | £ o or | 30 | - | - | Trinidad | 1825. | S | r.m | |
| 23308 | tenuifolia | Lo. C. | fine-leaved | £ o or | 30 | - | - | Brazil | 1824. | S | r.m | |
| 2646. | DESMO'NCUS | Mart. | DESMONCUS. | (Desmos, band, ogkos, hook; tendrils at apex of Ivs.) | Palmue. | 4. | | | | | | |
| 23309 | polyacanthus | Mart. | many-spined | £ CD or | 6. | . | . | Brazil | 1822. | S | r.m | Mart. palm. ic. |
| 23310 | ortnacanthus | Mart. | straight-spined | £ Q or | 6. | . | . | Brazil | 1822. | S | r.m | Mart. palm. ic. |
| 23311 | africanus | Lo. C. | American | £ CD or | 6. | ... | ... | St. Vine. | 1824. | S | r.m | |
| 23312 | dubius | Lo. C. | doubtful | £ CD or | 6. | . | . | Trinidad | 1824. | S | r.m | |
| 2647. | ASSIROICA'RYUMM | Mey My ASTROCARYUM. | (Aslron, star, karyon, nut.) | | Palmue. | 5.—10. | | | | | | |
| 23313 | Bacilla | Mart. | steinless | £ D or | 10 | . | . | Brazil | 1820. | S | r.m | Mart. palm. ic. |
| 23314 | acuticatum | Mey. | prickly | £ o or | 40 | - | - | Guiana | 1824. | S | r.m | |
| 23315 | Murumuru | Mart. | Murumuru | £ CD or | 40 | - | - | Brazil | 1825. | S | r.m | Mart. palm. ic. |
| 23316 | vulgare | Mart. | common | £ D o r | 30 | - | - | Brazil | 1825. | S | r.m | Mtr. palm. ic. |
| 23317 | campestris | Mart. | field | £ CD or | 10 | - | - | Brazil | 1826. | S | r.m | Mart. palm. ic. |
| *2648. | SYAGRUS Mart | SYAGRUS. | (The first who wrote the Trojan war in verse?) | | Palmce. | 1. | | | | | | |
| 83318 | coccolles | Mart. | Cocoa-like | £ O or | 20. | . | . | Brasil | 1824. | S | r.m | Mart. palm. ic. |

CLASS XXI. ORDER 7.

MONCECIA OCTA'NDRIA

2649. DUVAUM Kth. DUVAUA. (*M. Buvau*, a French botanist.) *TereUnthcca.* JL
 23319 deplndens Dec. dependent % i_J or 20 ... W.a Chile 1790. C sp Cav.Ic.&239
 ifmyris polygamia *Cao.* *Schlinsus dependens Or.*
 83320 dentata Dec. tooltied J>_Jor 20 myjl W Owhyhee 1795. C 61 Bot. rep. 620
Schlinsus dcnt&ta Andr.

CLASS XXI. ORDER 8.

MONCEXIA ICOSA'NDRIA.

- S650** ATHEROSPERMA Lab. ATHEROSPERMA. (*Ather., awn, sr. erma, seed*) *Calycantheaf* 1.
£321 moschata Lab. musk f L Jor 23 myjn W N. Holl. 1824. C sp.l

CLASS XXI. ORDER 9.

MONGSCIA POLYA'NDRIA.

2653. SAGITTA* RIA L. AIROW-HKAD. (*Sagitta*, arrow; form of lvs of some species.) *Alistnace**. 15.-2L

I. SAGITTATIFOLIE.—Leaves sagittate or hastate.

SES «g»tifMia L.	common, arrow-lyd	A or	2	jn.au	W	England rivers. D Lp	Eng. bot 81
23327 latifolia W.	broad-leaved	A or		jl.au	W	N. Amer. 1819. D Lp	
23328 Doni»a Swt	Don's	A or		jl.au	W	Nepal 1820. D Lp	
■ hastata D. Don							
*3f(32) obtiiHa W	obtuse	A Al or		jl.au	W	N. Amer. 1830. D Lp	
23330 heterophylla Ph.	various-leaved	A Al or		jl.au	W	N. Amer. 1822. D Lp	
2J331 hastate Ph.	hastate	A Al or		jl.au	W	N. Amer. 1818. D Lp	
23332 obtusifolia L.	blunt-leaved	& El or	2	jl.au	W	China 1804. D Lp	R. mal. 11.45
gaiancifolia Z.							
natans Mx.							
Af.							
337 falcate Ph.							
23338 acutifolia L.	acute-leaved	A or	1	in jl	W	Carolina 1812. D Lp	
^w^ angustifolia B. R.	narrow-leaved	A or	1	in jl	W	Carolina 1812. D Lp	
					W	America 1816. D Lp	
					W	Essequibo 1827. D s.p	Bot. reg. IUI

II. ANGUSTIFOLLIE.—Leaves subcordate, lanceolate, or linear.

gaiancifolia Z.	lance-leaved	= iAI or		Ijfn.jl	W	W. Indies 1787. D Lp	Bot mag. 1792
	rigid-leaved	*= A T		U jn.il	W	N. Amer. 1806. D Lp	Hot mag. 1632
	floating	= A or		jl.au	W	Carolina 1812. D Lp	
	Grass-leaved	= A or		ljl.jl	W	Carolina 1812. D Lp	
	falcate	= A or		1 jl.au	W	Carolina 1812. D Lp	
	acute-leaved	A or		in jl	W	America 1816. D Lp	
	narrow-leaved	A or		in au	W	Essequibo 1827. D s.p	Bot. reg. IUI

III. TBI'FIDA.—Leaves trifid or triperted.

Chinese	=iA Jbr	2	s.n.	W	China	1812. D Lp	Bot mag. 1631
---------	---------	---	------	---	-------	------------	---------------

2654. BEGO^NX L. BEGONIA. (*Michael Begon*, a French promoter of botany.) *Begoniacea*. 27.-47.

I. INEQUALICORDATE.—Leaves unequally cordate.

23341 njtidatf.*:	shining-feafeef	n Q or	ljmyd	W	Jamaica 1777. C s.p	Par. Ion. 72
kwér ri chotoma Jac.	forked		2 jlau	W	Caracas 1800. C s.p	Jac. ic. 3.619
23343 discolor H.K.	discolored		3 my.s	W	China 1804. R s.p	Bot mag. 1473
KvansVina Swz.						
4 renfurmis Dry.	kidney-shaped				Brazil 1818. C p.l	Lin. tr. 1.14.1.2
	long-leaved				Jamaica 1793. C ap	PL ic. 34.45.1
	tuberous				Amboyna 1810. C Lp	Ru. am. 5.169.2
	shining				W. Indies 181A C pi	
	doubtful				Brazil 1818. C p.i	
	two-winged				C G. H. 1822. C p.i	
	simulatc				W. Indies 1819. C Lp	Bot cab. 107
	painted				Nepal 1818. C Lp	Bot. cab. 571
	few-flowered			 1816. C Lp	Bot reg. 471
	spreading				W. Indies 1818. C p.l	
	waved				Brazil 1825. C p.l	Bot. mag. 2723
	sweet-scented			 1824. C Lp	
	small hairy			 1824. C Lp	
	two-rowed			 1824. C Lp	

II. SEMICORDATE.—Leaves semicordate.

23358 acuminaté Dry.	ttQor 1 my.d	W	Jamaica	1790. C 8?	Bot reg. 364
*9 acutifolia Jac.	acute-leaved	<, r~lor	jl.au	W	W. Indies 1818. C ?
SSS? hlkmlly. 1>L	humble	jf	D3or	W	W. Indies 1788. S
pulchellá Maf.	neat	O or	4jl.au	W	Brazil 182a S p.l
	shaggy-leaved	f IIPIor	my.jn	W	W. Indies 1789. S l.p
	sweet-scented	n. D or	jl.au	W	W. Indies 1816. C p.l
	Elm-leaved	tt r~lor	my.jn	W	S. Amer. 1822. C Lp
	silver-spotted	tt.r~or	jl.o	W	Brazil 1819. C Lp

III. OvxTMFBUS.—Leaves roundish^ ovate, or palmate.

climbing	L CD or	2	jl.au	W	Guiana	1822. C p.l	Aub.gui.2.349
----------	---------	---	-------	---	--------	-------------	---------------

palmate

		1	jl.au	W	Nepal	1819. D p.l	

BI'RNET. (*Potion*, a cup; infused in cooling drinks.) Rosdcete. 6.—7.

rimonif&liumClv.	Agrimony-Ivd	£ A P	3 >	Ap	1822. S co
unguisorba L.	Sangmsorba, c&m, fi	A ag	2 jl	G	England ch.hil. S co
>>ygamum Kit.					Eng. bot 860
hybridum / K.	polygamous	£ A or	3 jl.au	Br	Hungary 1803. S co
ludatum / K.	sweet hybrid	£ A or	2 in.jl	G	France 1683. S co
insumZ.	talled shrubby	n. LJ or	3 ja.ap	G	Bar. ic 632
	prickly shrubby	tt LJ or	2 ap.au	G	Canaries 1779. S p.l
					Levant 1595. S p.l
					M. h. 8.18.5

AMIRO^LA Pers. AMIROLA. (Meaning unknown; perhaps of Peruvian origin.) *Sapindacca*. 1.

nitida Pers.	shining-Zrawrd	% or	24	.	Peru	1824. C p.l
Lagunca nitida R.Sfp.						

ACIDO^TON Swz. ACIDOTON. (*Akidotos*, pointed; stinging pointed hairs of leaves.) *Euphorbidce*? 1.

stinging	i Qu n	8	...	G	Jamaica	179a C Lp	Sl. jam. L 83.1
----------	--------	---	-----	---	---------	-----------	-----------------

TOELY'GONUM L. THELYGONUM. (*Thelys*, woman, gam, knee; joints supp. to resem.) *Urtic*. 1.

Cynorambe L.	Dog's-cabbage	Jc O un	f jl	G	S. Europe	1710. S co	Lam. il 777
--------------	---------------	---------	------	---	-----------	------------	-------------

CASTA^NEA TOU. CHESTNUT. (A native of the territory of *Costarica*, in Thessaly.) *AmentHcecc*. 4.

vésca Gae.		tm	50	my.jn	G	England woods. S s.l	Eng. bot 886
------------	--	----	----	-------	---	----------------------	--------------

intermediate

Asplenium-lyd	tm	50	my.jn	G	Europe	—	L co
golden-leaved	tm	50	my.jn	G	Europe	—	L co
American	or	50	my.jn	G	—	—	S co
China	tm	50	xnyjn	G	America	—	S co
dwarf	or	12	jl	G.T	N. Amer. 1699. S co	Micar. 2.7	

HOP-HORNBEAM. (*Ostryos*, a scale; scaly catkins.) *Amenthcect*. 2. — .

common	%	or 20	my	Ap	Italy	1724. L 81	Dcn.br. 143
--------	---	-------	----	----	-------	------------	-------------

Virginian

	or	20	my.jn	Ap	N. Amer. 1622. L s.l	Ab. ins. 2.75	

HORNBEAM. (*Car*, wood, pin, head, *Celt*; made into yokes.) *Amenthcect*. 8. - 5

Birch, common	tm	30	mr.my	Ap	Brtein woods. S co	Eng. bot 2032	
cut-leaved	or	15	mr.my	Ap	—	L co	
variegated	or	20	mr.my	Ap	Britain gard. L co		
Oak-leaved	a	or 30	my.jn	Ap	Europe	—	L co
American	or 20	---	Ap		N. Amer. 1812. S co	Den. br. 157	
	or 12	---	Ap		Levant 1739. L co	Den. br. 98	

*2662. FA ^X GUS L.	BEIZCII. <i>(Phago, to eat; nuts used as food in early agea.)</i>	<i>Amentdcat.</i> 2.
83386 sylvatica L.	common wood Y tm 70 ap.my Ap Britain woods. S co Eng. bot. 1846	
2 ástro-rubens Duroi	dark-red-Zw* Y or SO ap.my Ap L co	
3 incisa JK.	cut-torvol S or 10 ap.my Ap G s.l	
83387 ferruginea L.	American, rusty £ or 30 my.jn Ap N. Amer. 1766. L 81 Mic. ar.2.9	
*2663. CWRYLVS L.	NIJTTTHBB. <i>(Korys^a helmet; calyx enwrapping fruit.)</i>	<i>AmcntflC'Sie. fi.—7.</i>
23388 ilvella & L.	Filbert \blacksquare fr 10 fap Ap Britain woods. S co Eng. bot. 723	
2 alba	white <i>FUber</i> a^* fr 10 fap Ap L co	
3 rubra	red <i>Filbert</i> ft fr 10 (ap) Ap L co	
4 grndis	great Cob ft fr 8 fap Ap L co	
5 glomerata	clustered ft fr 8 fap Ap L co	
6 crispata	frizzled ft fr 8 fap Ap L co	
7 barcelonensis	Barcelona ft fr 8 fap Ap Spain ... L co	
8 variegata	variegated ft fr 8 fap Ap L co	
9 ovata	ovAtc-fruited ft fr 8 fap Ap L co	
10p&mila	dwarf ft fr 6 fap Ap L co	
23389 tubulosa TV.	Lambert's, tubulose ft fr 10 mr.ap Ap S. Europe 1759. L co Lam. il. 780	
23390 americana Mx.	true American ft fr 10 mr.ap Ap N. Amer. ... L co Walt. ca. 21. GJ	
23391 hbmilis W.	humble <i>Cuckold</i> ^ ft fr 6 mr.ap Ap N. Amer. 1798 L co	
23392 rostrata // K. com.	beaked <i>Cuckold</i> Sk fr 5 mr.ap Ap N. Amer. 1745. L co W.ar.1.2	
23393 Coltima L.	Hazel, <i>Constantinople</i> ft 10 mr.ap Ap Constant 1665. L co Den. br. 99	
8664. JXTGLANS L.	WALNUT. <i>(Joviselans, the nut of Jove.)</i>	<i>Jugldndeee. 5.-6.</i>
23394 régia L.	common, royal \blacksquare tm 50 ap.my Ap Persia 3562. S co Lam. il. 781	
22395 n'gm L.	black \blacksquare tm 30 ap.my Ap N. Amer. 1629. S co Den. br. 158	
23396 cinerea L.	gley, <i>ButterNut</i> * \blacksquare tm 30 ap.my Ap N. Amer. 1666*. S co Jac. ic. 1.192	
23397 faxinitöha Lam.	Ash-leaved a^{v} tm 40 ap.my Ap N. Amer. ... S co	
23396 pterocarpa Mx.	winged, fruited \$ tm 40 ap.my Ap N. Amer. ... S co	
2665. CA ^X RYA Nut.	CARYA. <i>(Karyon, a nut.)</i>	<i>Jugldndeece. 10.—14.</i>
23399 tomentosa A. ^{dc} .	woolly \$ tm 30 ap.my Ap N. Amer. ... S co Mic. ar. 1.6	
23100 microcarpa Nut.	small-fruited If tm 30 ap.my Ap N. Amer. ... S co	
23401 olivaeförmis Nut.	olive-shaped 3f tm 30 ap.my Ap N. Amer. ... S co Mic. ar. 1.3	
23402 Juglans angustifolia // AT.	Juglans angustifolia // AT. olivsfürmis Mx. tm 30 ap.my Ap X. Amer. 1804. S co Mic. ar. J. 8	
Juglans bulcata TV.	furrowed % tm 30 ap.my Ap X. Amer. 1804. S co Mic. ar. J. 8	
83403 filba Nut.	white shell-bark Hickory^ tm 30 ap.my Ap* N. Amer. 1629. S co Den. br. 148	
Juglans squambsa Poir.	alba Afr. tm 30 ap.my Ap N. Amer. 1730. S co Mic. ar. 1.7	
83404 compressa Cr.	Don compressed-//^ £ tm 30 ap.my Ap N. Amer. 1730. S co Mic. ar. 1.7	
Juglans compressa A.X.	Juglans compressa A.X. tm 30 ap.my Ap N. Amer. 1800. S co	
23405 amhra Nut.	bitter Ah a^* £ tm SO my Ap N. Amer. 1800. S co	
Juglans amara ifar.	Juglans amara ifar. tm 30 my Ap N. Amer. 1812. S co Mic. ar. 1.9.3,4	
23406 obcordata Swt.	obcordate \$ tm 30 my Ap N. Amer. 1799. S co Mic. ar. 1.9.1,*	
Juglans obcordata Poir.	glabra YT. tm 30 my Ap N. Amer. 1799. S co Mic. ar. 1.9.1,*	
23407 porchna Tu.	hog nu/ 2 tm 30 my Ap N. Amer. 1799. S co Mic. ar. 1.9.1,*	
Juglans porchna Mx.	Juglans porchna Mx. tm 30 ap.my Ap N. Amer. 1799. S co Mic. ar. 1.9.1,*	
23408 laciniata Mx.	jagged ¥ tm 30 ap.my Ap N. Amer. ... S co	
*2666. QUE'RCUS L.	OAK. <i>(Quer, fine, cuex, a tree, Celt.)</i>	<i>Amentdcete. 62. — 110.</i>
I. INIEGRIFBÜJG. —Leaves entire or a little toothed.		
83409 Phellos L.	Cork tree, <i>Willow</i> P tm 60 my Ap N. Amer. 1723. S 81 Mic. ar. 1.12	
2 h \blacksquare P	humble * tm 20 my Ap N. Amer. ... S 81	
3 latikilius	broad-leaved \blacksquare tm 60 my. h Ap N. Amer. ... S co	
23410 maritima W.	sea or 6 my. h Ap N. Amer. 1811. S co Mic. quer. 13.1	
23411 imbrickaria Mx.	tiled tm 40 my.jn Ap N. Amer. 1781. S co Mic. ar. 2.13	
23412 /aurifolia Mx.	Laurel-leaved tm 50 my Ap N. Amer. 1786. S co Mic. quer. 17	
2 obtusa Mx.	b\xtent-leaved tm 50 my Ap N. Amer. 1786. S co Mic. quer. b	
23413 Idaea Nees	yellow tm 20 my Ap Mexico 1825. S co	
2 mirrocarpa	small-fruited or 20 my Ap --- S sl	
3 macrocarpa	long-fruited or 20 my Ap --- S al	
23414 cincerea Mx.	ash-colored or 10 my.jn Ap N. Amer. 1789. S co Mir. nr. 2.14	
83415 ambigua Hum.	ambiguous T tm 60 ... Ap Mexico 1820. S l.p H.&B.2.93	
83416 mexicana Hum.	Mexican J tm 60 ... Ap Mexico 1823. S l.p H.&B.2.82	
23417 sericea W.	silky running or 2 my.jn Ap N. Amer. 1724. S co Mic. ar. 2.15	
23418 ylrens H. K.	green tm 40 my Ap N. Amer. 1739. S sl Mic. ar. 2.11	
2 latifolia	broad-leaved J tm 40 my Ap N. Amer. 1739. S co	
II. \blacksquare T —Leaves toothed, spiny.		
23419 lanuginosa D. Don	lanuginous tm 20 ... Ap Nepal 1824. S !*	
23420 Phullata Ham.	Phullata \blacksquare tm 20 ... Ap Nepal 1822. S !*	
23421 -Hex /.	Holly, evergreen tm 60 my.jn Ap S. France 1581. S !* Den. br. 90	
1 integriflolia	entire-leaved tm 60 my.jn Ap S. France 1:181. S !*	
2 serrata	saw-leaved or 60 my.jn Ap S. France 1581. G 81 Duh. ar. 1.123	
3 oblonga	oblong-leaved or 60 my.jn Ap S. France 1581. G 81 Duh.ai. 1.124	
4/agifolia	Beech-leaved or 60 my.jn Ap S. Europe 1581. G co	
5 scripifolia	cur\et-leaved or 60 my.jn Ap S. Europe 1581. G co	
23422 SCibr L.	Cork tree dt 20 jn Ap S. France 1699. S 81 Den. br. 89	
23423 Pseudo atilier Sant	Bast Cork tree tm 40 my Ap S. Europe 1824. G co San. vi. 15R. 4	
23424 coccifera /.	kermes-nearings dt 10 my Ap S. France lf<3. G 81 Den. br. 91	
23425 lusitanica TV.	Portugal tm 40 jn Ap Portugal 1824. G 81 Cav. ic. 2.129	
23426 2?allbla Desf.	Ballota, <i>Barbary</i> tm 60 my.jn Ap Barbary S 81	
23427 rotundifolia TV.	round-leaved tm 40 my.jn Ap Spain 1581. S 81	
23428 gramnitria L.	Grammont or 30 jn Ap France 1730. G 81	
23429 infectoria Oliv.	staining tm 40 my.jn Ap Levant 1822. G co	
canariensis JV.		
23430 Turners'W.	Turner's ¥ tm 40 my.jn Ap » G co	
39431 rinUB X.J.	tm 60 my.jn Ap N. Amer. 1730. S 81 Mic. ar. 2.7	
fi montcolailfx.	mountain-inhabiting? tm 60 my.jn Ap N. Amer. 1730. S 81	
23432 Michauxia Nut.	Michaux's * tm 60 my Ap N. Amer. ... S 81	
Prtnus / discolor Mx.		
83433 Cartnea TV.	Yellow Chestnut* tm 60 my Ap N. Amer. 1822. S a1	
89434 spicata Hum.	spiked J O tm 60 — Ap Mexico 1822. S Lp H.&B. 2.89	

23487 maculatum <i>Lod.</i>	spotted	<. • or 6	mr	G	& Amer.	1820.	C s.p	Bot. cab. 608
segulnum /3 maculatum <i>B. M.</i>								
23488 xanthorlitzum <i>W.</i>	yellow-rooted	■ ■	■	■		1822.	R ap	Jacsc. 2.188
<i>A'um xanthorizum Jac.</i>								
23489 grandifolium <i>IV.</i>	great-leaved	■ ■	■ ■	■ ■		Caracas 1803.	R s.p	Jac. sc. 2.189
23490 arboréscens <i>Ven.</i>	arborescent	■ ■	■ ■	■ ■		W. Indies 1759.	R s.p	Pluk. am. 44.60
20491 arbreum <i>Kth.</i>	tree	■ ■	■ ■	■ ■		Cumana 1820.	C s.p	
23492 lacerum <i>W.</i>	torn	■ ■	■ ■	■ ■		Caracas 1822.	R s.p	

II. *Ackv'hik.* — Stemless.

i. INDIV'ISA. — Leaves simple, undivided.

23493 esculentum <i>Ven.</i>	esculent	■ Elclt			America 1739.	R fl.p	SI jam. 1.106. 1
23494 pumilum <i>D. Don</i>	dwarf	£ IAlor		W	Nepal 1820.	Sk l.p	
23495 cddlc <i>Mey.</i>	eatable	£ Elclt		W	Guiana 1800.	Skip	
23496 sagittif&hum <i>Ven.</i>	arrow-leaved	£ El or		W	W. Indies 1710.	It fl.p	Jac. vin. 2.157
23497 virgñicium <i>Hook.</i>	Virginian	ill A or	jn.jl	W	Virginia 1759.	R co	Hook. ex. fl. 182
iTrum virginicum i.							

23498 nymphelium <i>Ven.</i>	Water-hly-lvd	£ El or	4	W	E. Indies 1800.	R s.p	R.mal. 11.22
23499 bicolor <i>Ven.</i>	two-colored	£ El or	1	W	Madeira 1773.	R s.p	Bot. mag. 820
23500 viviparum <i>Box.</i>	viviparous	£ (2X or	limy	W	K. Indies 1817.	II Ip	Bot. cab. 281
23501 ovatura <i>Ven.</i>	ovate	£ [23 or	4	W	E. Indies 1818.	Sks.p	

23502 pinnatidium <i>TV.</i>	pinnatifid	£ (ZQ or	2	W	Caracas 1817.	R Bji	Jac. sc. 2.187
33503 tripartitum <i>W.</i>	three-parted-torf	BL CD or	3	W	Caracas 1816.	R s.p	Jac. sc. 2.190

iii. COMPÓSITA. — Leaves compound.

23504 pedatum <i>Hook.</i>	pedate	£ El or	2	W	Brazil 1824.	Sks.p	Hook. ex. fl. 206
23505 auritum <i>Ven.</i>	ear-leaved	n. or	3	W	America 1739.	R s.p	Jac. sc. 2.191
23506 Acleboriffrilium KTi.	Helleborc-lvd	£ El or	2	W	Caracas 1796.	R s.p	Jac. ic 3.613

2673. ^RUM L. ARUM.

(Supposed to be an ancient Egyptian word.)

Arotdet. 37.—

I. CBMPOSITA. — leaves composite,

23507 crintum <i>H. K.</i>	hairy-sheathed	3 AJ cu	1	mr.ap	Br	Minorca 1777.	R 61	Bot. reg. 831
23508 Dracimelus <i>L.</i>	Common Dragon	5 A or	3	jnjl	Br	S. Europe 1548.	R s.l	M.h. 13.5.46
23509 palmatum <i>Lo. C.</i>	palmate	^ A or	1			1825.	R s.l	
23510 Dracontium <i>L.</i>	Green Dragon	3 A or	2	jn	G	N. Amcr. 1759.	R s.l	Bot. reg. fff8
23511 vendsum // K.	veiny purple Jlwd	£ El cu	11	mr	P	1774.	R &l	Bot. reg. 1017
23512 pentaphllum <i>L.</i>	five-leaved	£ El or	1			1818.	R 81	Ru. am. 23
23513 pedatum <i>Fts.</i>	pedrite	£ El or	1			1820.	R s.l	
23514 polyphyllum <i>Lk.</i>	many-leaved	£ El or	1			1818.	R s.l	
23515 campanulatum <i>Rox.</i>	campanulate	■ El or	2	my	P	EL Indies 1817.	R s.l	Bot. mag. 2812
Tricca phalifera <i>Hum.</i>								

23516 ringens <i>Thun.</i>	ringent	■ ■	■ ■	I niy.jn	Japan 1800.	R ■	
23517 flagelliforme <i>Lod.</i>	whip-formed	■ ■	■ ■	I my	Bengal 1819.	R LP	Bot. cab. 396
23518 triphllum <i>L.</i>	three-claved	■ ■	■ ■	f myjn	Br N. Amer. 1664.	R SW	Bot. cab. 820
2 zebribium	zebra	■ ■	■ ■	I myjn	Br N. Amer. 1664.	R co	Bot. mag. 950
23519 atrofibrum // K.	dark-purple-sAtf	■ ■	■ ■	I jnjl	Br N. Amer. 1758.	R s.l	Pluk. al. 148.6
23520 ternatum <i>Thun.</i>	ternate-leaved	■ ■	■ ■	myjl	Japan 1774.	R s.l	
23521 bulbiferum <i>Rox.</i>	bulb-bearing	■ ■	■ ■	my	Bengal 1813.	R lp	Bot. mag. 2072

II. INDIVINA. — Leaves simple, undivided.

23522 proboscidium <i>L.</i>	proboscis-like	ife	■ ■	G	Apennines 1818.	R s.l	Boc. mu. 2. 53
23523 Colochia <i>L.</i>	Egyptian	Colocasia	A	■ ■	G levant	1551.	R s.p
23524 syracum <i>Spr.</i>	Syrian				Egypt	1820.	R s.f

gramineum <i>Rus.</i>	Gram-leaved	■ ■	■ ■			182a	R 81	
23525 spirale <i>Retx.</i>	spiral	■ ■	■ ■		Br	China 1816.	R l.p	Bot. mag. 2220
23527 pictum <i>L.fil.</i>	painted	■ ■	■ ■			Corsica 1800.	R 81	
23528 marginatum <i>Lod.</i>	marginated	■ ■	■ ■			E. Indies 1K20.	R s.l	
23529 macrophyllum <i>L.</i>	long-rooted	■ ■	■ ■			K Indies 1803.	R s.p	Her. par. 73
23530 divaricatum <i>L.</i>	divaricated	■ ■	■ ■			E. Indies 1759.	R s.p	R.mal. 11.20
23531 maculatum <i>L.</i>	common spotted	■ A or	ap.jn			Britain sha pi.	R co	Eng. bot. 1298
2,152/I orientale <i>Bieb.</i>	oriental	^ A or	1	ap.jn		Tauria 1820.	R s.l	
23533 it&licum <i>Lam.</i>	Italian	^ A or	11	my.jn	UY	Italy 1K83.	R co	Bot. mag. 248-
23534 minium <i>W.</i>	small	A EJ cu	1	my.jl	P	E. Indies 1812.	R co	R.mal. 11.17
23535 tenuifolium <i>L.</i>	fine-leaved	A cu	1	ap.jn	W	S. Europe 1570.	R s.p	Bot. reg. 512
23536 indicum <i>Lou.</i>	Indian	t-O cu	5		Br	China 1824.	R s.p	
23537 obtusilobum <i>Lk.</i>	blunt-lobed	£ El or	2			1824.	R l.p	
23538 sagittifolium <i>Lk.</i>	arrow-leaved	£ El or	2			1824.	R l.p	
23539 infirifolium <i>Lk.</i>	entire-leaved	tt-Dor 3	my.jn	G		1825.	R l.p	
23540 rambsum <i>Lk.</i>	branchy	tt-Dor 3	my.jn			1810.	R l.p	
23541 Attheraceum <i>L.</i>	Ivy-leaved	jgElcu	6	my.jn	P	W. Indies 1793.	R Ip	Jac. am. 153
23542 lingulatum <i>L.</i>	tongue-leaved	2fZ3cu	6			W. Indies 179a	R Ip	PI. ic. 26.37

III. SAGITTAVA. — Leaves sagittate or hastate, three Zohed.

23543 trilobatum <i>L.</i>	thre-lobed	A El cu	1	my.jn	P	Ceylon 1714.	R s.p	Bot. mag. 339
2 auriculatum	eared	A El cu	1	my.jn	P	Ceylon 1714.	R s.p	Bot. mag. 2324
Orixiyan	A (29 or	1	au,o	P	E. Indies 1S02.	R P.p	Bot. reg. 450	

2674. i*RISACRUM <i>Tou.</i>	FRIAE'B COWL.	(Adopted from the Greeks.)					Aroldetc. 1.	
23545 vulgare <i>Kth.</i>	common	fc AJ cu	1	ap.jn	L.Y	S. Europe 1596.	R 81	Jacsc.2.192

2675. CARYOVTai.	CARYOTA. (The Greek name of the	cultivated	date.)				Pulma.S.
23546 itrens <i>L.</i>	stinging, torn4eaved± or	20	...	W	E. Indies 1788.	S r.m	R. mal. 1.11
23547 mUis <i>Lou.</i>	mild	I D o r.		W	China 1820.	S r.m	
23548 horrida <i>Jac.</i>	horrid	£ d o r.	20	...	& Amer. 1820.	S r.m	

2676. DIPLOTHE'MIUM <i>Mart.</i>	DIPLOTBEMIUM. (<i>Diploos</i> , double, <i>thema</i> , snathe)	Palm*	2. - 4.
23549 maritima <i>Mart.</i>	maritime	I Q o r. 10.	Brazil 1823.
23550 campestre <i>Mart.</i>	field	1 CD or 10.	Brazil 1823.

2677. GOMIPTUS Rum.	GoMurua.	(Its name in Malabar?)	Palnue. 1.
23551 weehäfer Spr.	sugar-bearing	£ O or 40	Moluccas 1820.

2678. MANICA*RIA <i>Gae.</i>	MANICAHIA. (Mantra, a glovej spathe.)	Pdnuc. 1
23552 sacctfera <i>Gae.</i>	sack-bg, Winc-palm± Q o r 30.	& Indies 1823. b r.m Gac. fir. C. 1

2079. ATTA'LIA Hum.	ATTALIA.	(Attains, magnificent; trees.)	Pdlmec. 7.-9.
23553 htimilis Mart.	humble	i or 10.	Brazil 1820. S r.m* Mart. palm. ic.
23554 c6mptha Mart.	decked	* po r 22	Brazil 1820. S r.m Mart. palm. ic.
23555 funifera Mart.	rope-bearing	JLC Dor 40	Brazil 1824. S r.m Mart. palm. ic.
23556 Rossi; Lo. C.	Ross's	■ ■ ■ ■ ■	Brazil 1825. S r.m
23557 speciosa Mart.	showy	■ ■ ■ ■ ■	Brazil 1826. S r.m Mart. palm. ic.
23558 spectabilis Lg. C.	remarkable	2. or 70	Brazil 1824. S r.m
23559 extaa Mart.	tall	1d or 70	Brazil 1826. S r.m Mart palm. ic.
2680. PARIcNA Aub.	PARIANA.	(Its name in Guiana.)	Gramine*. 1.
23560 campestris Aub.	field	n.fZ]eu 2 ... Ap	Cayenne 1803. D co Aub.gui.2.337

CLASS XXI. ORDER X.

MON(EV)CIA MONADE'LPHIA.

, S681. NPPA Rum.	NIPA.	(Its name in the Moluccas.)	Palnue. 1.
23561 Mticans Thun.	shrubby	i d o r 10 ... W E. Indies 1822. S	rm Ilu. am. 1.16
2682. A RFC A L,	CABBAGR TREE.	(Called areec, in Malabar, when an old tree.)	PdlnuB. 10—14.
23562 Citechu L.	medicinal, Catagchuf	Q cit 30	W E. Indies 1690. ■ r.m Rox. cor. 1.75
23563 hhmilis W.	humilis	£ cit 6	W E. Indies 1814. ■ r.m Ru. am. 1.7
23564 olerkcea W.	pooherb	£ dt 40	W W. Indies 1656. ■ r.m Jac. am. 170
23565 crimita Sory	hairy-coated	£ O or 20	W I. France 1824. ■ r.m
23566 lutescens Bary	lutescent	£ O or 20	W I. France 1834. ■ r.m
23567 itcha ■■■■■	red	HI or 30	Mauritius 1833. ■ r.m
23568 Mdnicot Lo.C	Manicot	■ or 30	S. Amer. 1822. ■ r.m
23569 exUis Lo. C.	slender	■ or 30	W. Indies 1823. ■ r.m
23570 montana Lo. E	mountain	■ or 30	S. Amer. 1820. ■ r.m
23571 triandra Rox.	triandrous	£ Cl or 20	E. Indies 1825. ■ r.m

2683. CUNNINGHA* MIA R. Br. CUNNIKGHAMIA. (J. and A. Cunningham, trav. in N. S. W.) Conifer*. I.
23572 lanceolata It. Br. lance-leaved > i.J or 20 ... Ap China 1804. C p.l Bot. mag. 2743
Plnus lanceolita Lamb. Bélis jaculifolia Sal.

2684. DA'MMARA Rum. DAMMAR PINE. (Its name in Amboyna.) Conifer*. 2.-3.
23573 orientalis Lamb. oriental \$ CD or 50 ... Ap Amboyna 1804. C p.l Lamb. pi. 38
Plnus Dthmnara Lamb. A'gathis loranthifolia Sal.

23574 australis Lawzo. southern, CowdiePine± d] t m 100 ... Ap N. Zeal. 182L C p.l Lamb.pi.2.14.62

2685. PrNUSX.	PINE.	(Pin, a mountain, Celt.; habitat.)	Conflera. 39.—
	I. GEMINAVA.—Leaves in pairs.		
23575 sylvestris L.	wood, Scotch	tm 80 my Ap	Scotland sc. alp. S s.1 Lamb. pi. 1.1.1
2 r&bra Mil.	Ted-barked	or 20 ap.my Ap	Pyrenees ... S s.1
23576 genevensis Lo. C.	Geneva	tm 80 my Ap	Geneva 1820. S 6.1
23577 umilio Hae.	dwarf	or 20 ap.my Ap	Carniola 1779. S 8.1 Lamb. pi. 5.2
23578 ■■■■■	Corsican, Larch	tm 80 ... Ap	Corsica 1814 S s.1 Lamb.pi.2. 28.9
23579 ■■■■■	pungent	tm 40 ... Ap	N. Amer. 1804. S 8.1 Micar. 1.61.5
23580 ■■■■■ Banksia Sm^P/w	Banksia Sm^P/w	or 12 myjn Ap	Huds. Bay 1785. S s.1 Lamb. pi. 7.3
21181 Miighus Jac.	Miugo	or 10 myjn Ap	Austria ... S co Jac. ic. 1.193
23582 PinTister // K.	Cluster, Pinaster	tm 60 ap.my Ap	S. Europe 1596. S S.1 Lamb. pi. 9.4,5
*>83 uncinita Dec.	hooked	tm 50 my Ap	Pyrenees 1820. S 8.1
QQ-O- 8an8"inea Lap.			
ooiSf pyrenaean Lap.	Pyrenean	tm 50 ... Ap	Pyrenees ... S.1
23583 Pallawawa Lamb.	Pallas's	tm 40 my Ap	Siberia 1820. S.1 Lamb.pi.2.1.1
*^ ^ MassonMna Lamb.	Masson's	tm 40 my Ap	China 1824. S.1
23587 Pineq L.	Stone, Pine	(tm 40 myjn	S. Europe 1548. S s.1 Lamb.pi.13.9,10
J ^ o maritima Lamb.	maritime	tm 40 my Ap	S. Europe 1159. S s.1 Lamb.pi.2.30.10
23589 halepensis H. K.	Aleppo	tm 50 my Ap	Levant 1683. S.1 Lamb. pi. 18.13
JPD ^ nojw // K.	Jersey, poor	tm 50 my Ap	N. Amer. 1739. S.1 Lamb. pi. 20.14
resinifesa H. K.	pitchy	tm 50 my Ap	N. Amer. 1756. S.1 Lamb. pi. 20.14
riibra Mx.			
23592 varia Wis	variable, 2 and 3-ld	tm 40 myjn Ap	N. Amer. 1739. S.1 Lamb. pi. 22.15
23593 TsMa L:	Frankincense	tm 30 myjn Ap	N. Amer. 1713. S.1 Lamb.pi.23.16,17
**f>eehlnaU1IK	chcinated	tm 50 myjn Ap	N. Amer. 1739. S.1 co
	late	or GO myjn Ap	N. Amer. 1713. S.1 Micar. 1.86.7
	rigid	or 80 myjn Ap	N. Amer. 1759. S.1 Lamb.pi.25.18,19
	swamp	tm 20 ... Ap	N. Amer. 1730. S.1 14mh. pi. 27.20
23598 ponderosa Dou.	weighty-wooded	tm 50 ... Ap	N. Amer. 1828. S.1 co L.1nb. pi. ic. in.
^ nari f^n8 Schmidt	Canary	or 40 ... Ap	Canaries 1815. S.1 Dec. gen c. ic.
iftea Walt	voinw	tm 40 my Ap	N. Amer. 1821. S.1
	Chinese	i J or 40 ... Ap 1820. S.1
23603 longifolia Lamb.	long-leaved	i J or 40 ... Ap	China 1825. S.1
		E. Indies 1801. S.1 p.1	Lamb. pi. 29. 21

	III.	—Leaves in fives.	
23604 Strbus L	Strobos, Weymauthf	tm 100 ap	N. Amer. 1705. S.1 Lamb. pi. 31.22
*23605 Latnbertina Dou.	Lambert's	I tm 150 ... Ap	N. Amer. 1S27. S.1 Lamb. pi. ic. in.
23606 pygmari Hort.	pygmy	or 40 myjn Ap	1827. S.1 co
23607 excelsa Lamb.	tall	tm 100 myjn Ap	Nepal 1823. S.1 Lamb. pi. 2.5.3
23618 occidentalis Sun.	western	tm 80 myjn Ap	W. Indies 1826. S.1 PI. am. 17.161
23609 Ctmhra L.	Cembra, Siberian	tm 25 my Ap	Siberia 1746. S.1 Iam.iu.SI.23,24
2 helvetica	Swiss	tm 40 my Ap	Switzerl. 1819. S.1
3 sibirica	Siberian	tm 40 my Ap	Siberia 1825. S.1
IV.	—Species not sufficiently known.		
23610 rigensis TJ. C.	Riga	tin 40 my Ap	S. Europe 1824. S.1
?"? Lo. C.	Taurian	or 40 my Ap	Tauria 1820. S.1
"? Lo. C.	blue	or 40 my Ap 1820. S.1
"? novaza?lendica Lo.C.	New Zealand	tm 50 my Ap	N.Zeal. 1825. S.1

2086. FIR. *[Apios, a pear tree; cones like its fruit.]*

Conifene. 18.—

I. SOLITARLEFO^LIJS. —Leaves solitary, flat, subdistich.							
E H Fraseri Ph. Fraser's, <i>Dōuble Balsam</i> t or SO my Ap l'ennsylv. 1811.		81					
23615 Picea W. <i>Silver, pitch</i> f tm 30 my Ap Germany 1003.			Lamb. pi. 46.30				
pectinate Lam.							
23616 Douglass Hort. Douglas's tm 50 my Ap N. Amer. 1827.			Lamb. pi. ic. in.				
23617 balsamifera Mx. bilm-bearing or 50 my Ap N. Amer. 1696.			Lamb. pi. 48.31				
23618 canadensis Z/A. Can. <i>hemlock-spruce</i> . or 30 my Ap N. Amer. 1736.			Lamb. pi. 50.32				
23619 dumosa Lamb. bushy or 20 my Ap Nepal 1820.			Lamb. pi. ic.				
23620 sibirica Dirl. Siberian tm 40 my Ap Siberia 1820.							
23621 /axifolia Lamft. Yew-leaved tm 90 my Ap N. Amer. 1823.			Lamb. pi. 51.33				
23622 spectabilis Zam& remarkable tm 90 my Ap Nepal 1823.			Lamb. pi. 2.3.2				
II. SFARBIF&LLJE —Leaves scattered, angulate or subulate.							
23623 orientalis W. <i>oriental</i> t or 30 my Ap Levant 1825.		S co	Lamb. pi. 45.29				
23624 Clanbrasili <i>n</i> H.K. <i>Ld. Clanbrasili</i> s or 8 my Ap — L co			Lamb. pi. 37.25				
23625 cxcOsa <i>Pair</i> , tall, Norway spruce ± Plinus A^biez. L. tm 100 ap Ap N.Europe 1518 S s.l			Lamb. pi. 37.25				
2 fblus variegatis variegatcd-/Mi or 41 ap Ap N. Europe ... S si							
23626 alba H. K. white spruce J tm 50 my.jn Ap N. Amer. 1700. S £1			Lamb. pi. 39.26				
23627 rubra Lamb. red spruce J tm 50 my Ap N. Amer. 1755 S £1			Lamb. pi. 43.28				
23628 irtgra H. K. black spruce J tm 50 my Ap N. Amer. 17c0. S 81			Lamb. pi. 41.27				
23629 carpatica Hort. Carpathian £ tm 50 my 'Ap Carp. mo. ... S s.l							
III. DIAHJS. —Not sufficiently known,							
23630 româna H. Cel. Roman t tm 40 my Ap S. Europe 1820. S s.l							
23631 Pichta Fis. Pitch \$ tm 50 uiy Ap Siberia 1820. S s.l							
2687. LA*RIX Sal. LARCH. (Lar, fat, Celt.; producing abundance of resin.)							
S3632 europa^a Dec. European, com. white*£ tm 50 mr.ap Ap Germany 1629. S s.l			Lamb. pL 53.35				
Finns Zárix L. 2 sibirica Siberian tm 40 mr Ap Siberia 1834. S s.l							
23633 daurica Fis. Daurian tm 50 mr.ap Ap Davuria 1827. S co							
23634 plndula Lamb. Black, pendulous tm 30 my Ap N. Amer. 173». S s.l			Lamb. pi. 5fi.5fi				
23635 microcarpa Lamb. lied, amall-ftd \$ tin 80 my Ap N. Amer. 1760. S s.l			Lamb. pi. 58.5f				
2688. CE^DRUS Barrl. CEDAR. (Cedron, a brook in Judea; plentiful on its banks.) Coniferte. 2.							
23636 Libâni Barrl. Cedar of Lebanon ± or 60 my Ap Levant 1683. S &1			Lamb. pi. 59.37				
Pinus Cedrus L. 23637 Deodara Rox. Dcodara \$ tm 90 my Ap Nepal 1823 S 81			Lamb. pi. 2.8.4				
2689. SCHUBE'R T/J* Mir. SCHUBERTIA. (M. Schubert, a Polish botanist.) Conifene. 2.							
23638 distichal/ir. 2-rovrcd, deciduottsCyp. ^ or SO my Ap N. Amer. 1640. S s.p			Mic. ar. 3.4.1				
Taxbdium distich um Rich. Cupressus dfsticha L. 2 pe*ndula H. K. drooping \$ or 20 my Ap S. Amer. 164a S s.p							
23639 capensis Spr. Cape *t_jor 6 ap.my Ap C. G. H. 1756. S p.l							
CupSisus juniperoldcs L. Thrixus elongatus Thun.							
23643 n CicferP</>, nut-bearing J LJ or 20 — Ap Japan 1822. S l.p			Ram. 815				
7Vixu n heifer Kami.							
23644 Yacca G. Don Yacca t O tm 50 — Ap W. Indies 1818. S r.m							
23645 ncriffbia Lam. Nerium-leaved f • tm 40 — Ap E. Indies 1820. S r.m			Ru. am. 3.20				
23646 nsplcnifflus Lab. Fern.-leaved J i_ or 20 — Ap N. Holl. 1825. S r.m			Lab. n-fa. 2,221				
ThaUimia rr^plnifflia Spr.							
23647 feixiblius Kth. Yew-leaved t • or 40 — Ap S. Amer. 1820. S r.m							
23648 chinensis Swt. Chinese f LJ or 30 — Ap China 1800. S r.m							
Taxus chinensis Rox.							
23649 spinulbus Spr. spinulose £ _J or 20 — Ap N. Holl. 1820. S r.m							
* 2690. PODOC A'RPUS Herit PODOCARPUS. (POUS, a foot, karpos, a fruit; stalked.) Conifer*. 9.—							
23640 raacophyllus Swt. long-leaved f or 10. jl.au Ap China 1804. S			UankaK.il.				
23641 vrticillatus Hort. vrhor-leaved t or 10 ■ Ap Japan —							
23642 elongatus Herit. elongated J I_ or 10 jl Ap C. G. H. 1774. S. Lp							
TVixus elongatus Thun.							
23643 n CicferP</>, nut-bearing J LJ or 20 — Ap Japan 1822. S l.p			Ram. 815				
7Vixu n heifer Kami.							
23644 Yacca G. Don Yacca t O tm 50 — Ap W. Indies 1818. S r.m							
23645 ncriffbia Lam. Nerium-leaved f • tm 40 — Ap E. Indies 1820. S r.m							
23646 nsplcnifflus Lab. Fern.-leaved J i_ or 20 — Ap N. Holl. 1825. S r.m							
ThaUimia rr^plnifflia Spr.							
23647 feixiblius Kth. Yew-leaved t • or 40 — Ap S. Amer. 1820. S r.m							
23648 chinensis Swt. Chinese f LJ or 30 — Ap China 1800. S r.m							
Taxus chinensis Rox.							
23649 spinulbus Spr. spinulose £ _J or 20 — Ap N. Holl. 1820. S r.m							
* 2691. DACRY'DIUM Sol. DACRYDIUM. (Dakry, a tear: gummy exudation.) Cmfcraj. 1.							
23650 cupr&Binum Sol. Cypress-like £. _J or 20. . . . N. Zeal. 1825. C 81			Lamb. pi. 1.41				
Thalamia cupressina Spr.							
2692. CUPRE'SSUS L. CYPRBS9. /Kuo, to produce, parisos, equals, regular branches.) Conifene. 8.—12.							
23651 scmpervirens L. com, evergreen f or 20 my Ap Candia 1548. S co			Den.br. 155				
1 stricta strict . or 20 my Ap Mediterr. ... S co							
2 horizontMis horizontal . or 20 my Ap Mediterr. ... S co							
23652 lusitanica Ton. Portug., CedarofGoa J _J or 12 ap.my Ap Goa 1683. C p.l			Lamb. pi. 95.42				
gladca Lam.							
23653 hyôules L. Thya-like, While Cedar ± or 20 ap.my Ap N. Amer. 1736. L co			Den. br. 156				
2'hfja sphsrrroidca Rich.							
23654 australis Pers. south, slender-bran. ± \J or 10 apmy Ap N. Holl. — L p.l							
23655 pendula Thun. pendulous J LJ — 20 my Ap Japan 1808. L p.l							
23656 bacciformis JV. berry-khaped . or 20 my Ap — 1818. L p.1							
23657 triquetra Lo. C. triangular J I_J or 12 my Ap C. G. H. 18M. L p.l							
23658 torulbsaZ. Don twisted J _J or 20 ... Ap Nepal 1824. L p.l							
* 2693. IHUVA L. ARBOR VITA (77)ion, sacrifice; used in for fumigation.) Conifer*. 10.—5.							
23659 occidentalism western, American J or 25 ray Ap N. Amer. 1596. L co			Mic. ar. 3.29.3				
23660 orijentalis L. or'enU\Chinese] J or 25 my Ap China 1752. S co			Den. br. 149				
23G61 taturica Lo. C. Tartarian J or 20 my Ap Tartary 1820. L co							
83662 articulata Vahl jointed J or 15 f.my Ap Barbary 1815. S co			Bot.cab.844				
23663 v\chta Lamb. plaited J or 20 my Ap Nootka S. 1796. L co							
23664 ptndula Lamb. drooping J or 20 my Ap Tartary 1828. L co							
93665 pyramidMis Ten. pyramidal J or 20 my Ap Italy 1824. L co							
23666 nepalensis Lod. Nepal J or 20 my Ap Nepal 1824. L co							
23667 australis Bosc. southern J or £0 my Ap & Europe 1820. L co							
23668 cupressoldcs L. Cypress-Ik, Aftic. 5 LJ or 10 ... Ap C. G. H. 1799. L p.l							
2694. TRICHOSA'NTHES L. SNAKE GOJBBD. (Thrix, hair, anthos, fl.; inner divis. ciliated.) Cucurb. ft-15.							
23669 angulna L. common, snake * Ol or 4 my in W China 1755. S co			Botmfig.722				
Cilcumis anguincus L.							

23670 <i>cucumbrina</i> L.	Cucumber-like <i>Jk</i> .QJ un 4 jn.jl W	E. Indies	1804. S	co	R. rail 8.15
23671 <i>colubrina</i> Jac.	Berwnt-like <i>-*</i> QJ un 4 jn.jl W	W.	1817. S	co	Jac. cc 123
23672 <i>lobata</i> Rox.	lobate <i>-*</i> QJ un 6 Jlau W	E. Indies	1812. S	co	
23673 <i>cordata</i> Rox.	cordate <i>-*</i> QJ un 4 jl.au W	E. Indies	1818. S	co	
2695. LAGEN A*RI A Ser. BOTTLE	GOURD. (<i>Lagena</i> , a bottle; form of fruit.)	<i>Cucurbitidea</i> . I. — 4.			
23>74 <i>vulgäris</i> Ser. common	<i>-*</i> Q clt 10 JLs Y	India	1597. S	co	Ru. am. 5.147
Cucurbita Lagendria L.					
2 <i>Gordia</i> Ser.	Gourd <i>Jfc</i> o clt 10 jls Y	India	1597. S	co	M.h. 1.5.1
3 <i>depressa</i> Ser.	depressed <i>-*</i> O clt 10 jls Y	India	1597. S	co	
4 <i>turbinata</i> Ser.	turbaned <i>-*</i> O clt 10 jLe Y	India	1597. S	co	M.h. 1.5.2
5 <i>clavata</i> Ser.	clubbed <i>J±</i> O clt 10 jls Y	India	1597. S	co	M. h. 1.5.3
6 <i>Courgo&rla</i> Ser.	Courgourde <i>-*</i> O clt 10 jls Y	India	1597. S	co	Ru. am. 5.144
269R CERATOSA'NTHES Brtn. CERATOBANTHES. (<i>Keras</i> , horn, <i>anthos</i> , fl. j inner segments.)	<i>Cucurb. I.</i>				
23675 <i>tuberosa</i> J.	tuberous S C3 un 6 jl.au W	W. Indie*	1810. R	co	Bot mag 2703
Trichosanthes tuberosa IV.	cornicuua Lam.				
2697. MOMCRDIC A Z. MOMORDICA. (<i>Mordco</i> , to bite; seeds seem bitten.)	<i>CucurbUace</i> *. 10.—19.				
23676 <i>ifalsanuna</i> L.	Balsam Apple <i>\$</i> QJ un 4 jn.jl Y	India	1568. S	co	
23677 <i>Chardnia</i> L.	hair, Charantia <i>^</i> QJ un 4 jn.jl Y	E. Indies	1710. S	co	Bot. mag. 2455
23678 <i>operculata</i> L.	lid-covered, rough-tid <i>-*</i> QJ un jn.s Y	W. Indies	1731. S	co	Coin. r. 32
83679 <i>diocia</i> Rox.	diocious <i>\$</i> IIHun Jlau Y	E. Indies	1818. S	co	R.mal.8.18
23680 <i>muricata</i> W.	muricated <i>-*</i> QJ un Jlau Y	E. Indies	1817. S	co	K.mal.8.10
23681 <i>tubiflora</i> Rox.	tube-flowered <i>Jf</i> A un Jlau Y	E. Indies	1825. S	co	
23>82 <i>umbellata</i> Rox.	umbelled <i>J>A</i> un Jlau Y	K Indies	1878. S	co	
23683 <i>mixta</i> Rox.	mixed <i>\$</i> El un jj.lau Y	E. Indies	1820. S	co	
23684 <i>senegalensis</i> Lam.	Senegal <i>A</i> QJ un Jlau Y	Senegal	1822. S	co	
^3G85 Elaterium Zr.	Squirting Cucumber <i>Jc</i> A un Jn.jl Y	S. Europe	1548. S	r.m	Bot. mag. 1914
2698. CUCURBITA L.	GOURD. (<i>Curvitas</i> , crookedness; shape.)	<i>CucurbWacc</i> ? 10.—17.			
23686 <i>ovifera</i> L.	egg-bearing <i>-*</i> O clt 3 jls	Astracan	...	S co	
1 <i>pyrifirra</i> Ser.	pear-form-/-/ Jk O clt 3 jlau	S co	
2 <i>Hubgobbsa</i> Ser.	subglobosc/d -i o clt 3 jlau	S co	
3 <i>grisea</i> Ser.	greyfruited <i>-*</i> O clt 3 jlau	S co	
23687 <i>siccaria</i> Mol.	<i>intoxicating</i> <i>Jk</i> O clt 4 jn.au	Chile	1824. S	co	
23688 <i>aurantia</i> W.	<i>Or*nt>e-fruited</i> <i>Jk</i> QJ clt 3 jn.au	...	1802. S	co	
1 <i>ordigno</i> Ser.	<i>false</i> Orange <i>Jk</i> QJ clt 3 jn.au	...	1802. S	co	
2 <i>coccyntoides</i> Ser.	<i>Coloquinelle</i> <i>-*</i> QJ clt 3 jn.au	...	1802. S	co	
23689 <i>maxima</i> Duch.	largest <i>Jfc</i> O clt 4 jn.au	S co	
máxima Pepo Dec.					
23690 Pepo L.	Pumpkin <i>Jk</i> O clt 16 jn.au Y	Levant	1570. S	co	
1 <i>subrotunda</i> W.	subrotund-//rf <i>Jk</i> O clt 6* jn.au Y	Levant	1570. S	co	
2 <i>obl6nga</i> IV.	oblong-//d <i>jk</i> o clt 6 jn.au	...	1570. S	co	
23691 Pcrfira Duch.	Potiron, large-std <i>Jk</i> O clt 10 jlau	S co	
I viridis Ser.	green Potiron <i>Jk</i> O clt 10 jlau	S co	
	Gourd-bearing <i>Jk</i> O clt 10 jlau	S co	
23692 verruebsa L.	warty <i>J*</i> O clt 12 jn.jl	...	1658. S	co	
23693 subverrucosa W.	subwarty <i>Jfc</i> Q clt 12 Jlau	S co	
23694 Jtfetpepo L.	Melon-pumpkin, squash <i>O</i> clt 3 my.s	...	1597. S	co	M.h.1.81
23695 moschata Duch.	musky <i>Jk</i> O clt 4 jn.au	S co	Dalch. 60(1:)
2699. CITCUMISZr.	CUCUMBER. (<i>Curyus</i> , crooked; shape)	<i>CucurbUhcae</i> . 22.-25.			
23696 Colocynthis L.	bitter, Colocynth <i>Jk</i> ^ in 6 my.au Y	C.G. H.	1551. S	r.m	
...	prophets, globe <i>Jk</i> m cu 2 jn.s Y	Levant	1777. S	co	Jac. vin. 1. &J
B B L	Anguria, round prickly <i>-*</i> fJ cu Jlau Y	Jamaica	1692. S	co	Mil. ic. 1. &J
J. African	African <i>-*</i> XJ cu 2 jlau Y	C. G. H.	...	S co	Bot. reg. 980
^3700 muricata ■.	muricate JCQJCU 4 jn.au Y	E. Indies	1817. S	co	
23701 A/elo L.	Melon <i>JcX</i> cul 4 my.s Y	...	1570. S	r.m	Sab. rom. 65
1 <i>reticulatus</i> Ser.	netted <i>Jk</i> QJ cul 4 my.s Y	...	1570. S	r.m	Black. 529
2 <i>Cantalupo</i> Ser.	Cantalupc <i>-*</i> QJ cul 4 my.s Y	...	1570. S	r.m	
3 <i>maltensis</i> Srr.	Maltese <i>-*</i> IQJ cul 4 my.s Y	...	1570. S	r.m	
23702 <i>delicibus</i> Roth	delicious <i>-*</i> S? cul 4 jn.au Y	E. Indies	1818. S	co	
23703 <i>utilissimus</i> for.	most useAll <i>-*</i> IQ cul 4 jn.au Y	E. Indies	1820. S	co	
23704 <i>Dud&im</i> L.	Dudaim, <i>apple-sliapcd</i> <i>JK</i> cu 6 jUu Y	Levant	1705. S	r.m	Bot. rep. 1*8
23705 Chkte L.	Chate, /ai/ <i>Jk XII cu</i> 6 Y	Levant	1759. S	co	Alp. aeg. 117
^706 pubescens ff.	pubescent <i>-*</i> XII cu Y	...	1815. S	co	
23707 saftvus i.	common, cultivated <i>-*</i> O cul 4 Y	E. Indies	1573. S	r.m	Sab. rum. G3
1 vlfiflis & r.	green <i>Jk</i> O cul 4 Y	E. Indies	1597. S	r.m	
2 tkviss Ser.	yellow <i>Jk</i> O cul 4 Y	S r.m	
3 albus Ser.	white <i>-*</i> O cul 4 Y	S r.m	
4 variegatus Ser.	variegated <i>-*</i> Q cul 4 Y	S r.m	
5 fastigiatus Ser.	peaked <i>Jk</i> Q cul 4 Y	S r.m	
23708 flexuosa L.	flexuous, snake <i>-*</i> £J cu 6 Y	E. Indies	1597. S	r.m	Gr. li. 7K. 3
23709 jamaicensis A;;7.	Jamaica <i>-*</i> QJ cul 4 Y	Jamaica	1824. S	co	
23710 hnc/tub	lined <i>Jk</i> QJ cu 4 Y	Guiana	1825. S	co	
23711 maculata ff.	Mulrat <i>Jk</i> QJ cu 3 Y	E. Indies	1805. S	co	Pluk. al. 170.2
2J712 maculata ff.	spotted <i>-*</i> QJ ru 4 Y	E. Indies	1820. S	co	Scr. dis. 3
H K ¹ ₂ B ¹ < T ¹ ₂ Xox.	dingy,dewell <i>Jk DN cu</i> 4 Y	E. Indies	1820. S	co	
27714 Hommrdira Rox.	Klitrium-//Ac <i>Jk n cu</i> 4 Y	E. Indies	1820. S	co	
23715 Cucurbita Ser.	Citrus 3 ft. dt Y	E. Indies	1820. S	co	
Cucurbita CtrOllas L.	...	S. Europe	1597. S	co	Ru. am. 5.146
1 Pff Ser.	J>a8tcque <i>Jk</i> jQJ clt 6 my.s Y	...	1597. S	co	Black. 157
2 Jdrc Ser.	Water Melon <i>Jk</i> QJCU 6 Y	...	1597. S	co	
25718 laciniatus L.	...	W. Indies	1824. S	co	Bur. am. <
23719 parviflorus IV.	Hoover <i>-*</i> O cul 4 jlis Y	Quito	1823. S	co	
23720 microphillus Kth.	small-leaved <i>-*</i> O cul 4 jls Y	...	1823. S	co	
2701. EL ATEMHIUM L.	ELATERIUM. (<i>Elater</i> , an impeller;				
21721 carthagin'ns L.	Carthngena <i>^</i> O cul 5 jn.jl Y				
^722 amioole5 W.	Tamniw-like <i>£</i> o cul 5 jn.jl Y	Mexico	...	S co	Jac. am. pic. ^ i

2702. SETCHIUM Br. CHOKO. *(Sckizo, to fatten; given to hogs.)* Cucurbitacee. 1.—&
23723 e_<lule Br. eatable Jc.QJcul 6 Jn.jl Y W. Indies 1816. S r.m.

Sicyos edulis Jac.

2703. NEUROSPERMA Baft. NEUHOSPERMA. (*Neuron, a nerve, sperma. seed.*) Cucurbitacee. 1.
23724 cuspidata Rifi. cuspidate * O clt 5 jn.s Y Kentucky 1827. S co

Z10kBRYO^yUAL. BRYONY. (*Bryo, to sprout up; rapid growth of annual stems.*) Cucurbitacee. 23.—61.

I. ANGULAVJE.—Leaves subangular.

23725 scabra L.	scabrous, globe-fruited^ LAJ un	6 s.o.	W.o	C. G. H. 1774.	S p.l
23726 verruebsa H. K.	warty ft.iAJun	4 ...	W.o	Canaries 1779.	S co
23727 confita Thun.	cordate ft.	LAI un	6 JLau	Y.G C. G. H. 1816.	S r.m
23728 grandis L.	greatflowered ft_	ES un	8 my.au	W.G E. Indies 1783.	S p.l Ru. am. 5. Ifin. 1
23729 epigas'a W.	ground, umbel-flwd% LjiuY*	Leaves lobate	E. Indies 1815.	S co	
23730 trilobata Thun.	three-lobed ft.	IAJ un	s.o.	W.o C. G. H. 1825.	S p.l
23731 scabrella L.	ruughiah JE vjl un	my.jl	W.o	E. Indies 1781.	8 co
2J732 latebrfcsa 1/. IT.	shady ft.	LAI un	Jn	W.o Britain hcd.	S co
23733 dioica Jac.	dioecious, red-brd^ A m	inv.s	W.o Europe 1807.	S co Eng. hot. 4iJ	
23734 alba L.	white, black-brd^ A I	m	W.o	Europe 1834.	S co Lam. ll. 796
23735 nitula Lk.	shining % iAJ un	jnl	W.o	Candia 1759.	S co
£3736 cretica i.	Cretan £ iAJ un	jnl	W.o	An. mu. 12 17	S co
23737 quinquidoba TAtm.	flye-lobed £ iAJ un	3 jh.o	Br C. G. H.	co Bet. rcg. 82	
23738 acifolia Lam.	Fig-leaved fc. iAJ un	d	W.o	Buen.Ay. 172 <i>q</i> .	S co Di. el. 50.58

III. PALMAYB.—Leaves palmate.

23739 palmata L.	palmate % ES un	4 jl.au	W.o	Ceylon 1778.	S co
23740 divisa W. en.	divided-feared fe_ iAJ un	6 ji ai	Y.G	... 1819.	S r.m
£3741 Garcin* W.	Garcin's ft.	QS un	6 jl.au	Y.G Ceylon 1812.	S r.in Bur. in. 57.3
23742 pinnatifida Burc.	pinnatifid (. LAI un	6 jl.au	Y.G C. G. H.	1815.	S co
23743 lacimosa L.	laciniated \$ (Z3 un	4 jl.au	W.o E. Indies 1710.	S co Her. lug. P7	
23744 africana Thun.	African J. IAI un	3 JLau	W.c C. G. H.	1759.	S co Her. i>ar. 70S
23745 dissecta L.	&is6ccctcd-leavedt iAI un	3 JLau	W.a C. G. H.	1710.	S p.l

IV. DU*BIA—Doubtful to which sections they belong.

23746 tenella Box.	slender ft.	(22 un 6 jl.au	Y.o	E. indies 1820.	S co
23747 filiformis Box.	filiform ft_	ES un 6 jlau	Y.o	E. Indies 1817.	S co

2705. /INDRA'CHNE L. BASTARD OUPINE. (The Greek name for purslain.) Euphorbiacea. 1.—2.
23748 te^{ephw}des L. Telephium-like Ow |jl.a.u W Italy 1732. S co Lam. il.797

2706. STILLVSGIA L. STILLINUIA. (Dr. Benjamin StUUngesteen, an English bot.) Euphorbidcea?. 4.—■.

23749 sylvatica L.	wood tf iAI un	2 jLau	Y	Carolina 1787.	C s.p
23750 ligustrina W.	TrixUcaved tk un	5 ...	Y	N. Amer. 1812.	C s.p
23751 sebifera W.	tallow-bearing • ec	10 8	Y	China 1703.	C s.p Pluk.al.SPO. 2
23752 Dopulnea VaU	Poplar-like { Q o r	14 . . .		Ceylon 182a	C si

2707. OMALA'NTHES Grab. OMALANTHES. (Homalos, smooth, anttos, flower.) Euphorb. 1.
23753 populifolia Grah. Poplar-leaved * • or 6 jl.s VV N. Holl. 1825. C p.l Botfmag.2780

2708. PHYLLA'NTHUS L. PHYLLANTHUS. (*Phyllon, leaf, anthos fl.; on the alae of the lvs.*) Euph. 28.—71

I. HERA^CBI.—Flowers solitary; leaves subpinnate; annual.

23754 obovatus W.	obcordate f ITEfmr		Jau	M. Amer. 1803.	i.p
23755 obcordatus Box.	obcordate f		Jau	K. Indies 1817.	s.p
23756 gracilis Rox.	slender	a	ejpr	Q.Y	K. Indies 1818.
2X157 Virftri L.	Niruri	a	ejpr	E. Indies 1692.	s.p a mat. 10.1/i
23758 cantomensis Horn.	Canton	EDpr	L au.s	Canton 1H20.	s.p
S3759 cuneMus W.	vrc&gc-leaved	B3pr	W	China 1818.	s.p
23760 Uriifana L.	urinary Opr	au.s	G.Y	E. Indies 1819.	s.p Ru. am. 6.17.2
23761 simplex Betz.	staple Ol>r	au.s	G.Y	E Indies 1817.	s.p

11. AGGRBGAVI.—Flowers aggregate; leaves subpinnate.

23762 scandens Box.	climbing f pr	10 jls	G.Y	E. Indies J822.	s.p
23763 grandifolius Lam.	great-leaved HQiin	5 -		America 1771.	s.p
23764 juglandifolius W.	Walnut-feawd * • or	2 jl s	G.Y	... 1818.	fi.p
23765 ificencs Poir.	shining * i_ pr	jts	G.Y	China 1820.	s.p Jacsc.2.1"i
23766 virifus Rox.	venomous A • pr	jls	G.Y	E. Indies 1802.	s.p
23767 lanceofatu? Poir.	lance-leaved * pr	au.s	R	I.Bourbon 1822.	s.p
23768 reticulatus Hort.	netted	■ pr	G	E. Indies ...	s.p Bot. cab. 116
23769/raxinifolius Hort.	Ash.leavcd	■ pr	G	E. Indies 1819.	s.p Bot. cab. 89
23770 Conumi Swz.	Conami	6 jl	G	W. Indies 1791.	1p Aub.gui.2.354
23771 obscurus Rox.	obscure	3 jl.s	G.Y	E. Indies 1824.	s.p
23772 ndtans Swz.	nodding	1 au.s	G.Y	Jamaica 1820.	B.p

III. SuBBOLFTA*Ri.—Flowers subsolitary; leaves subpinnate.

23773 maderaspafensis L.	Madras m. • un	3 jl	Ap	E. Indies 1783.	C s.p
23774 microphyllus Kth.	small-leaved m. • pr	1} au.s	G.Y	Caraccas 1817.	C s.p
23775 longifolius Lam.	long-leaved O pr	3 jls	G.Y	I.Bourbon 1822.	C s.p Lam. il. 156.1
23776 turbinatus B.M.IUTbinatc,shinin*T-lvd* • pr	2 jl	G	China	... C s.p Bot. mag. 186*	
23777 rhamnoides Betz.	Rhamnus-like * • pr	3 jl au	G.Y	E. Indies 1820.	C s.p
23778 strictus Box.	strict - • pr	3 jl.s	G.Y	E. Indies 1824.	C s.p

IV. Pi^An.—Leaves truly triniate, and bearing flowers between the pinnae.

23779 mimosoides Swz.	Mimosa-like m Q p r	10 au.s	G	Carribbees 1817.	C s.p Bot. cab. 721
23780 polyph'Uus JT.	many-leaved i Q p r	3 jls	G	E. Indies 1805.	C sp

2709. XYLOPHYLLA L. XYLOPIYLLA. (Xylon, wood, phylion, leaf; rigidity of foliage.) EuphorUdeie. H

I. PHYLLO'NTHUS.—Bearing flowers on the margins of the leaves.

23781 longifolia W.	long-leaved I Q o r	3 f.n	Y.R	£ Indies 1816.	C i.p Ru.am.7. W
OTIQO f*yx4nthus ceramicus Pers.	broad-leaved * * < »	3 au.g	Y.R	Jamaica 1783.	C s.p Bot.jnag.lQK ¹
23782 tifolius L.	showy t Q o i	3 au.o	Y.R	Jamaica 1818.	C s.p Jacic.3.610
23784 angustiflThiaV.	narrow-leaved * • or	3 jl.au	Y.R	Jamaica 1789.	C s.p Bot. cab. 577
•Pnyllanthus angustifbhus Swz.	cpii>hyUanthus L.				

ORDER X.

MONCECIA MONADELPHIA.

23785 <i>linearis</i> Pers.	linear	• • or 3	jls.	Y.B	Jamaica	1819.	C	s.p
Phyllanthus linearis Swx.								
23786 <i>elongata</i> Lod.	elongated-tui	• • or	J1.8	Y.B	Jamaica	1820.	E	s.p Bot. cab. 1091
23787 <i>media</i> J.O. C.	intermediate	• • or		Y.B	Jamaica	1825.	E	• p
23788 <i>montana</i> IV.	mountain	• • or	U	Y.B	Jamaica	1819.	E	s.p Bot. mag. 2652
23789 <i>falcata</i> H. K.	falcate-leaved	• • or		Y.B	Bahama	1.1699.	E	fi.p Bot. reg. 373
Phyllanthus <i>falcatus</i> Swx.			jls					

11. RAMIFlBFlRi. — Bearing flowers on the branches.

23790 <i>ramiflora</i> Raff. tf.	branch-flwd	• J or 3	U.au	Y.B	Siberia	1785.	E	s.p Pal. it 3. E. 2
23791 <i>obovatoid</i> W.	obovate-leaved	• J or 3	jl.au	Y.a				

2710. <i>EMBLICA</i> Gae.	EMBLICA.	—	%.	T&nSti~»	nSW&Arii.			
23792 <i>officinalis</i> Gae.	officinal							
Phyllanthus <i>Emblica</i> L.								
23793 <i>racemosa</i> Spr.	racemose	ft • or 2	jlau	G.v	£ Indies	1793.	C	s.p
Phyllanthus <i>racemosa</i> L.								

2711. ALEUR^TES For. ^f	ALEUBITEB.	Mto ^r ific# mealy; appearance.)						
23791 <i>triloba</i> Forst.	three-Tbed	I • fra 10	—	Ap	Society Is.	1793.	C	r.m

2712. OMPHA^LEA L.	OPHALEA.	(<i>Omphalos</i> , navel)	umbilicated anthers.)	Euphorbi&cce.	2—			
23795 <i>triandra</i> L.	triandrous, long-lud	Q or 15	jnj	G	Jamaica	176*	C	p.l Bot. cab. 519
23796 <i>diandra</i> W.	diandrous, <i>Mulberry-lvd</i>	H	12	...	Ap	W. Indies	1820.	H p.l Aub. gen. 2388
2 diversifolia	various-leaved	H	12	...	Ap	Cuba	1823.	E pl

2713. JIPPOMANE L.	MAVINEEL.	(MW, horse, mania, madness; effects of original						
23797 <i>Mancinella</i> L.	Machineel, common	Dp 80	—	G	W. Indies	1690.	L	r.m

« 714 SA^PIUML.	SAPHTIM.	(<i>Sap.</i> fat, <i>Celt.</i> ; unctuous exudation from wounded trunk.)	Euphorb.	4.—8.				
23798 <i>acuparum</i> Jinc.		P or 30	...	G	W. Indies	1692.	C	p.l Jac. am. 158
23799 <i>Indicum</i> W.	Indian	CO	—		E. Indies	181a.	C	p.l
23800 <i>tfippomane</i> Mey.	Manchineel	24	—		Guiana	1823.	C	p.l
*23801 <i>flcitibium</i> W.	Holly-leaved	20	—		a Amer.	1820.	C	p.l
Aipp6manc spinbsa L.								

*2715. CRO^TON-L.	CBOTON.	(<i>Kroton</i> , a tick j seeds resemble.)	Euphorb&cce.	33.—158.				
I. INTEGRIFOLI^. - <i>Uavcs entire or serrated, Uncar, ^ncolate, oblong, or ovate.</i>								
23802 <i>varicigata</i> JV.	variegated	10	—	W.G	E Indies	1804.	E	R.mal.6.61
2 <i>crispa</i> Rox.	curled	(J) or 6	jn.au	W.G	E. Indies	1804.	E	Ru. am. 4.26.2
3 <i>media</i> Rox.	intermediate	6	jn.au	W.G	E Indies	1804.	E	Ru. am. 4.25
23803 <i>linearis</i> Jac.	linear	Dor	—	W.o	W. Indies	1733.	E	Bot. cab. 481
23804 <i>Cascarilla</i> L.	Camarilla	* ip	4	W.G	S. Amer.	1778.	E	Cat car. 2.46
23805 <i>maritima</i> JV.	maritime	* ljun	4	W.o	Carolina	1786.	E	
23806 <i>palustris</i> L.	marsh	E2 jun	3	W.G	Vera Cruz	1731.	E	Mart. cen. 4. 38
23817 <i>Rialeua</i> L.	smoothish, <i>Laurel-lvd</i>	I or 6	—	W.G	Jamaica	1778.	E	Sl.jam. 2.174.2
23808 <i>tinctibia</i> W.	dyer's	Hu	3	W.G	S. Europe	1570.	E	Ac. pet 1712.17
23809 <i>flatum</i> U.	Tiglum, purging	ni 10	"aus	W.G	E. Indies	179&	E	R. mal. 2.33
23810 <i>Eleuteria</i> L.	Eleut., Sea-sute	BalsamM	m	W.G	Jamaica	1748.	E	Lp
23811 <i>castanæfolia</i> L.	Chestnut-leaved	US un	6	W.G	Trinidad	1826.	E	Bot mag. 2794
23812 <i>picta</i> Box.	painted	• or 4	jl.au	W.G	E. Indies	1810.	E	Bot cab. 870

II. COBDA^TJG. — <i>Leaves cordate, entire or serrated.</i>								
23813 <i>xalapensis</i> Kth.	Xalapa	Mlyn	2	jl.au	W.o	Mexico	1824.	E
23814 <i>argentea</i> L.	silver-leaved	O un	2	jl.au	W.G	S. Amer.	1733.	E
23815 <i>nitens</i> Swx.	shining	• un	3	jl.au	W.G	Jamaica	1818.	E
23816 <i>licra Desf.</i>	jagged	• un	2	jl.au	W.G	1822.	E	
23817 <i>m'cans</i> Swx.	glittering	1-lun	3	—	W.G	Jamaica	1815.	E
23818 <i>macrophylla</i> Sum.	long-leaved	I-lun	3	—	W.G	Jamaica	1823.	E
23819 <i>gf9, g8, g11</i> Hic.	pungent	• un	4	—	W.G	Caracas	1791.	E
23820 <i>pencillata</i> Ven.	pencilled	El un	4	jl.au	W.G	Cuba	1799.	E
23821 <i>glandulosa</i> L.	glandulous	E un	2	au.s	W.G	W. Indies	1817.	E
23822 <i>flvens</i> L.	yellowish	H1 un	3	jl.au	W.o	Jamaica	1818.	E
23823 <i>aromatica</i> L.	aromatic	I un	6	—	W.G	Ceylon	1793.	E
Jghi	humble	I un	2	—	W.o	Jamaica	1799.	E
23824 <i>molucca</i> W.	Moluca	• un	10	—	W.o	Ceylon	1803.	E
AlcurJes Ambinax Pers.								
23826 <i>astrolotes</i> H. K.	star-like, woolly*	un	6	jl.au	W.o	W. Indies	1782.	C
23827 <i>mvea</i> Jac.	snow-white	• un	3	jl.au	W.G	Jamaica	1823.	C
23828 <i>lucida</i> Swx.	shining	• un	3	—	W.o	Jamaica	1820.	C
23829 <i>tomentosa</i> Lk.	tomentose	• un	2	iny.jn	W.o	1824.	E

III. LOBATIPOLIJE. — <i>Leaves lobate.</i>								
23830 <i>ijossypifolia</i> Vahl	Cotton-leaved	* O un	3	jl.au	W.G	Trinidad	1818.	C
23831 <i>Tobeta</i> L.		O un	2	jl.au	W.o	Vera Cruz	1730.	S co Mart. cen. 5.46
23832 <i>Aibiscifolia</i> Kth.		Hibiscifolia	• un	4	jl.au	W.o	Mexico	1810.
digitata Hart.								
23833 <i>digitata</i> Fu.	digitate	H un	3	jl.au	W.G	N. Holl.	1810.	C p.l
23834 <i>flabellifolia</i> L.	fan-leaved	H un	3	—	W.o	—	1825.	C co
8716. BRADLEV,* Gae.	BBADLEJA.	(Rich. Bradley, first prof. bot. Cambridge ?)	Euphorbuiiccto.	2.—7.				
23835 <i>sinica</i> Gae.	Chinese	• c u	...	jl.au	...	China	1816.	C s.l. Gae. fr. 2.109.1
23836 <i>nitida</i> Rox.	shining	• CZcu	—	—	E. Indies	1820.	C s.l.	

2717. JA'TROPHIA L.	PHYSIC NUT.	<i>iatros</i> , physician, <i>trophe</i> , food ; qualities.)	Euphorb.	13.—21.				
I. LoBATIPOLIE. — <i>Leaves lobed or multifid.</i>								
23837 napicifolia Desv.	Napsoa-	un	3	jn.au	G	Antilles	1825.	C p.l
25838 multifd Ada L.	multifid	or	3	jn.au	S. Amer.	1696.	C r.m	Par. Ion. 91
23839 frtgrans Kth.	sweet-scented	D or	10	—	Cuba	1822.	C p.l	
24140 /rossyptifolia L.	Cotton-leaved	• or	3	my.au	W. Indies	1690.	C l.p	Bot. cab. 117
54841 glandulosa Vahl	glandulous	• or	3	—	Arabia	1854.	C p.l	
23K42 <i>Circas</i> L.	Curcas, angular-lvd	• or	4	—	S. Amer.	1731.	C r.m	Jac.vin.a63
2M441 <i>peltata</i> Kth.	peltate	ftO	4	—	S. Amer.	1825.	C p.l	
2S844 <i>irens</i> L.	stinging	• or	4	my.jl	Brazil	1690.	C p.l	
S1845 <i>herbeea</i> I*	herbaceous	3	jl.au	G	Vera Cruz	1759.	S r.m	Bot cab. 478
S3846 <i>austrolis</i> Lo. C.	southern	HQun	1	—	N. HoU.	...	S r.m	R.hou.6.15
		C c 4			C p.l			

II. ANGITSTIFOLIA.—Leaves narrow.

23847 integerrima Jffse & most cut respicth-lvd D orr	3	xny.au	S	Cuba	1809. S r.m Bot. mag. 1464
23848 coccinea Lk.	scarlet	H J or	4	. Cuba	1824. C p.l
23849 pandurafolia And.	fiddle-leaved	H • or	4	my.au '9	Cuba 1800. S r.m Bot.mag.G04

* 2718. JANPPHA Kth. JAMPIU. (*Janipaba*, its name in Brazil.) *Euphorbulcete*. 4.

23850 Loefflingii Kth.	Lading's	■ CJ clt	3	jl.au	Br Carthag. 1820. C r.m Jac. am. pic. 244
Jatropha Jahipha L.	Cassava	• - clt	3	jl.au	Br S. Amer. 1739. C r.m SL jam. 1.85

83851 A/r/^{Ad}/Kth. ^A Jatropha Manihot L.

23852 osculifolia JT/A. Horsechestnut-lvd

23853 foetida Kth. fetid * D o r 3 ■ Br Mexico 1824. C r.m

2719. iZI'CINUS L FALMA CnRisi. (*Ricinus*, a tick j seeds resemble.) *Euphorbiicete*. 8.—10.

23854 commixnis L com., *Castor-oil Plant* D Jm 6 jl.au G E. Indies 1548. S co Hot. mag. 2219

23855 vridij W. green QJor 6 au G £. Indies 1802. S s.p W.h.b.⁴J

23856 africanus TV. African *i Jor 15 jl.au G Africa ... S s.p

23857 lividus Jac. Vivid-leaved H J or 8 jl P C. G. H. 1795. S e.p Jac. ic. 1.1%

23858 inrmis Jac unarmedc<HMtt/grf Dior 6 JLau P India 1758. S s.p Jac. ic. 1.195

23859 armatus B. B. armed-capsuled £21 or 6 JLs G Malta 1807. S s.p Bot.rcp.4J0

23860 tanarius W. Tanarian, scallop-lvd QJor 4 JLs G E. Indies 1810. S s.p Ru. am. 3 1121

23861 globosus FT. globose IQot 4 — — Jamaica 1826. S p.l

2720. SIPHCTNIA Rich. SIPHIIONIA. (*Siphon*, a pipe; use of exudation, Indian rubber.) *Euphorb&ccce*. 1.

93862 Calluchu Rich. Cahuchu • or 6 . Guiana 1823. C s.l Lam. il. 790

clastic Pers. Jatropha clastica L Hevia guianensis Aub.

2721. AGYNEM A L. AGYNEJA. {A priv. sync, womanj neither style nor stigma.) *Euphorbtice*e. 2.—1.

23863 impbbes L smooth O un 1 jlau G China 1820. S s.l

23864 pbber L. hairy H Q u n 3 ... G China 1823. C s.l

2722. KIRGANE^LIA J. KIBGANEIIA. (*KirganeU*, its Malabar name.) *Euphorbulcete*. 1.—5.

23865 elegans J. elegant HO or 6 jl.au ... Mauritius 1820. C 81

>hyllanthus Kirganilla W.

2723. HIPRA L. SANDBOX TREE. (Its South American name.) *Euphorbiicce*. 2.—3.

23866 strepens W.en. soundngvnequ-toothf • or 12 ... W.v S. Amer. ... C l.p

23867 crépitans L rattling, cqual-toothcd± O or 12 ... W.Y S. Amer. 1733. S p.l Lam. il. 793

*2724. STEItCITLIAZ., STBRCULIA. (*Stercus*, ordure; some species fetid.) *Bytnerihcece*. 17. —2*J.*

I. BALA'NGIAS.—Leaves ovate or oblong, entire, rarely three-lobcd.

23868 Baldnghas L. Balanghas | Q o r 20 jna G.v E. Indies 1787. C R. mal. 1.49

23869 pubescens G. Don pubescent H J or 20 — W Guinea 1793. ■ p.

23870 grandiflora Fen. great-flowered H dor 6 — E. Indies 1820. ■ p.

23871 coccinea Rox. 6carltc-/rufai t a or 20 — E. Indies 1817. ■ p.

23872 ndbilis Sm. noble J • or 20 — E. Indies 1787. ■ p. Bot. reg. 185

Balandgas H. K. Southwells n6bilis Sal.

23873 angustifolia Rox. narrow-leaved f J or 20 — Nepal 182a ■ P!

23874 guttata Rox. spotted f O or 20 — E. Indies 1810. ■ R. mal. 4.61

23875 acuminate Beauv. Cola, pointed f J or SO — W Guinea 1825. ■ Ueau.ow. 1. 24

23876 Iv'a Swz. Ivira | I or 20 — W. Indies 1793. ■ pi Aub. gui. 2.279

crinata Cav.

23877 diversifolia G. Don various-leaved f LJ or 14 — N.Holl. 1824. C p.l

heterophylla Cun.

23878 iirens Rox. stinging f CD or 10 — E. Indies 1793. ■ p.l Rox. cor. 1.24

t 6879 vildsa Rox. villous 5 • or 15 — E. Indies 180J. ■ p.l

2J880 Helictrcres Pers. Helictrcres-jfc aQor 8 — Ap Carthag. 1820. ■ p.l Jac. am. Ml. 97

Helicteres an^tala Jac.

23881 platanifolia L. Plane-tree-lvd t i J or 30 jl — China 1757. ■ Cav.dis.5.145

23882 acrifolia Cun. Maple-leaved f i J or 20 — N.Holl. 1824. ■

23883 colorata Rox. colored \$ or 30 — E. Indies 1818. ■ p.l Rox. cor. 1.25

III. PALMA^TA.—Leaves palmate.

23884 foytida L. fetid * Q or 8 ... E. Indies 1690. ■ p.l R. mal. 4.36

2725. HERITIE'Rif H. K. LOOKING-GLASS PLANT. (f. L. VHerUicr de Brut die. Fr. bot) *Byltneriiie*. 2.—S

23885 littoralis H. K. shore t • or 20 . . . E. Indies 1780. C p.l R. mal. 6.21

23886 minor Lam. smaller I Q or 12 . . . E. Indies 1824. C p.l Gae.fr.2.98.2

2726. ^CA'LYPHA L. ACALVPHA. {A, priv., *kalos*, pleasant, *aphe*, touch; in handling.) *Euph*. S0.-60.

I. ANDR6GTNA.—*Monococious*; spikes androgynous.

23887 virgfnica L. Virginian Q un ■ jl.au G N. Amer. 1759. ■ Sch.han. 3.311

23888 carolinifana Walt. Carolina Q un ■ jl.au G N. Amer. 1811. ■ I^in.il. 780.2

03889 ciliata Forsk. ciliated rm un ■ jl.au G E. Indies 1799. ■ Vahl sy. 1.20

23890 pauciflora W. en. few.flowercd O un ■ jl.au G China 1816.

23891 brachystachya W.c. short-spiked O un ■ jl.au G China 1816.

23892 prunifolia JBA. Plum-leafcd im un ■ jl.au G Mexico 1825.

23893 indica L. Indian QJ un ■ ft 1 G E. Indies 1759. ■ R. mal. 10.81

23894 r^ptans Sum. creeping Lf un ■ ft G Jamaica 1820. ■ Sl. jam. L82. 3

23895 rubra W. red JE O un 3 jl.au G ... 1820. ■

23896 hispida TV. hispid JE O un 3 jl.au G E. Indies ... ■

23897 cuspidata Jac. cuspidate Q. un 4 jdjl G Caracas 1819. ■

II. ALOPECUROIDEJL. Spikes of distinct sexes, female ones terminal.

23898 alopecuroidca Jac. Foxtail-like O un 2 jl.s G Vnezuelal804. S co Jac. ic 3.620

23899 polystachya Jac. many-spiked E) un 1 jl.au G W. Indies 1817. S co Jac. sc. 2.246

23900 mollis Kth. soft ED un 1 jl.au G Mexico 1825. S co

23901 Isvigata Swz. smooth-leaved I Q un 3 jn.jl G Jamaica 1819. C co

III. DKEVLJE.— Dioecious.

23904 integrifolia W. entire-leaved <1 • un 5 jn.s G Mauritius 1823. C co

23905 scabrosa Sum. rough il Q un 6 jn.jl G Jamaica 1820. C co

Jac.sc. 2.245

	V. AXILLANIIS.— <i>Spikes of both sexes axillary.</i>	
23906 vhrgJtal.	twiggy i i Q u n 6 jnjl G	Jamaica 1823. C co Dr.Jam.36.2
2727. DALECHA'MP/i! L. DALECHAMPIA. (<i>James Dalechamp</i> , French bot. d.1588.) <i>Euphorbidcap.</i> 3.—18.		
23907 scandens L.	climbing ai um 1p2 Jnjl G.v	W. Indies 1739. C l.p Jac. am. 252.160
23908 brasiliensis Lam.	Brazilian 6*-cu 6 jnjl *	G.v Brazil 1824. C l.p
23909/icifolia Lam.	Fig-leaved I L O c u 6 jnjl G.v	Brazil 1830. C l.p
o.2728. PLUKE NETJJ L. PLUKE NETIA. (<i>Leonard Pluknet</i> , an English botanist) & q*m*Acew. J. 5.5.		
23910 rol&bilis L.	twinning t O u n 6 jl.au G	W. Indies 1739. C p PL ic. 220.2*6
23911 verruebsa Sm,	warty * - d u n 6 J^1" G	Surinam 1820. C p.l
2729. GNETUM L. G.VETUM. (Altered from <i>gnemon</i> , its name in the island of Ternate.) <i>Urtfcar.</i> I.—		
23912 Gnmon L.	Gnemon tQor E. Indies 1815. C p.l R.am. L.71	

CLASS XXII. ORDER 1.

DHFCIA MONA'NDRIA.

2730. PANDJCNUUS L. fl. SCREW PINE	Panda Mg. to behold, Malay; conspicuous.	Pamliuric. 20. — L's.
23916 htimilis Hum.	dwarf lor 20	W E. Indies 1771. S r.m Rox. cor. 1.94fi
23917 amarylli/d/w Rox.	Amaryllis-lvd lor 20	W Bourbon *** S r.m Jac. fr. 13,11.1
23918 candelabrum Beauv.	candlestick lor 60	W N. S.W. 1805. S r.m
*23919 fascicularis Lam.	fascicled lor 20	W Mauritius *** S r.m
23920 intrmis Rox.	unarmed lor ...	W E. Indies 1820. S r.in
23921 integrifolius Lou.	entire-leaved lor ...	W Klndies IS18. S r.m
23922 reflexus Hort.	reflexed lor ...	W E. Indies 1818. S r.m
23923 alhus Hart.	white lor ...	W E. Indies 1820. S r.m
**W* 86881118 JSoj.	sessile lor ...	W E. Indies 1822. S r.m
23925 latifolius Hort.	broad-leaved lor ...	W E. Indies 1818. S r.m
23926 marginatus Rox.	margined lor ...	W E. Indies 1822. S r.m
23927 longifolius IA> C.	long-leaved lor ...	W E. Indies 1823. S r.m
23928 turbinatus Lo. C.	lor ...	W E. Indies 1820. S r.m
23929 furcatus Rox.	forked lor ...	W E. Indies 1823. S r.m
23930 pedunculatus R. Br.	pedunculate lor ...	W E. Indies 1824. S r.m R.mal.£. 8
23931 foetidus Rox.	fetid lor ...	W N. Holl. 1825. S r.m
23932 edulis Thou.	eatable lor ...	W E. Indies 1818. S r.m
		W Madagasc.182A. S r.m
S731. TRICHOCCLADUS P<r>. TRICHOCLADV9. (Trichion, hairy, klados, branch.) I.		
2%33crinUuaPer«.	haired ft _Jor	. C G. H. 1823. C p.18

CLASS XXI. ORDER 2.

DIOTCIA DIA'NDRIA.

* 2732. SA'LIX L.	WILLOW. (Sal, near, Its, water, Celt, j place of growth.)	Amcnatactu. u1.
I. SERRAS'JL — Leaves smooth, serrated.		
^2334 triandra L:	triandrous tm 30 my.au Ap	Britain riv.ba. C m.s Eng.liot.1435
—> Hoffmannia Sm. Hoffmann's	tm 30 my Ap	England riv.ba. C m.s Cur. Ion. 72
triandra Hfm.		
—> lanceolata Sm.	tm CO ap.my Ap	England mca. C m.s Eng. bot. UX>
Hoppeana W.	tm 30 ap.my Ap	Austria 1820. C m.8
—> wavy-leaved	tm 30 ap.my Ap	Gennany ... C m.s
—> amygdalina W.	or 6 ap.my Ap	S. France 1818. C m.s
2335 decipiens E. B.	deceptive or 6 ap.my Ap	Britain mar. C m.s Eng. bot. VXA
2336 Runyliana W.	or 8 my Ap	England woods. C m.s Eng. bxt. VW
2337 Kumboldt/VJwrl. W.	tuuhsrs tm 40 ap.my Ap	England mar. C m.s Eng. bot. 1S<
2338 tetraspLrma lioth	four-acceded fnor 20	Peru 1823. C m.s
23945 nigra Marsfr**	black or 20 my Ap	E. Indies 17Ux C m.s Rox. cor. I.*7
23916 pentandra L.	pentndrous or 15 mr.jn Ap	N. Amer. 1811. C m.s An. hot 2. J. *
&2347 nericans Sm.	blackish broadUvdm or 10 ap Ap	Britain riv.ba. C m.s Eng. bot. IW5
23948 viula'cea Ander.	violaceous A or 10 Ap	England os.hol. C m.s Eng.bot.12U
	Weigel's fil or io ; Ap	Silesia* !!! C m.s
23950 Phylicifolia L.	Phylica-leaved fil or Umy Ap	Scotland sc. alp. C m.s Eng. bot. 195S
23951 Borer terna Sm.	Borer's fil or 8 my Ap	Scotland moun. C m.s Fi. dan. 1052?
23952 nUens Ander.	shining fil or 10 ap Ap	Scotland moun. C m.s
23953 Davallimi Sm.	Davall's A or 6 my Ap	Scotland moun, C m.s
—> ylicifolia W.	Wulfen's or 6 my Ap	Carinthia 1818. C m.s
23955 silesaca fk	Silesian or 6 my Ap	Silesia 1816. C m.a
23956 Pontederana W.	Pontedcra* s or 3 my Ap	Switzerl. 1821. C m.a
23957 aurina Sm.	Laurel-like A or 8 apmy Ap	England sto.hi. C m.8
23958 teuuifulia IV.	tiuin-leavcii or 2 myjn Ap	Britain sto.hi. C m.8 Eng. bot. 18*ii
		m.s Eng. UtfilW

23959 Ammannntta W.	Ammann's	or	■ myjn	Ap	Austria 1821.	C	m.s Hof sal. 17, 18, 11)
23960 hastata L.	haXherUeaved	or	■ my	Ap	Lapland 1780.	C	m.s Fl. lap. 8. f. g.
23961 serrulata W.	serrulate	or	■ my	Ap	Lapland 1810.	C	m.s Fl. dan. 1&2
23962 /rino'ides Ph.	Prinos-like	or	■ mr.ap	■	N. Amer. 1811.	C	m.s
23963 discolor Mhl.	discolored	or	■ ap	■	N. Amer. 1811.	C	m.s An.bot. 2.5.1
23964 angustita Ph.	narrow-leaved	or	■ mr.ap	■	Pennsylv. 1811.	C	m.s
23965 petiolaris Sm.	dark, pctioled	or	10 ap	Ap	England mar.	C	m.s Eng.bot. 1147
23966 myncoledes Mhl.	Gale-like	or	8 ap	Ap	N. Amer. 1811.	C	m.s An. hot 2.5.2
23967 cordata Mhl.	heart-leaved	or	6 ap.my	Ap	N. Amer. 1811.	C	m.s An. bot. 2. n. 8
23968 rlgida Mhl.	rigid	or	15 ap.my	Ap	N. Amer. 1811.	C	m.s An.bot. 2.5.4
23969 lucida Mhl.	shining	or	8 my	Ap	N. Amer. 1811.	C	m.s An. bot. 2.5.7
23970 ambigua Ph.	ambiguous	or	20 ap	Ap	N. Amer. 1821.	C	m.s
23971 acutifolia JV.	sharp-leaved	or	8 ap	Ap	Casp. Sea 182J.	C	m.s
23972 vitellina L.	ye\ovr.branched	clt	15 mr.my	Ap	England mar.	C	m.s Eng. bot. 1389
23973 fragilis L.	brttue	or	15 ap.my	Ap	Britain mar.	C	m.s Eng. bot. 1807
23974 prsecox Hoppe	rfaphnoldes Vil.	■ ■	or 20 ap.my	Ap	Austria 1820.	C	m.s
23975 Meyeriufl W.	Meyer's	or	20 ap.my	Ap	Germany 1822.	C	m.s
23976 babylonica L.	Babylonian, weeping	el	30 my	Ap	Levant 1692.	C	m.s Rauw. it 183.25
23977 annularis Hort.	nng-icav'd	or	20 -	Ap	... 1823.	C	co
23978 Napoleona Hort	Naj>looon's, weep.	u.J or	30 -	Ap	St. Helena 1823.	C	m.s
23979 purpurea W.	bitter purple	or	8 -	Ap	England mar.	C	m.s Eng. bot. 1S88
23980 poinerina W. en.	Pomeranian	or	10 my	Ap	Pomeran. 1822.	C	m.s
23981 //Mix L.	Helix	or	10 mr.ap	Ap	Britain mar.	C	m.s Eng. bot. 1343
23982 Lambertiana Sm.	Lambert's	SI	clt 10	mr.ap	England riv. ba.	C	m.s Eng. bot. 1SSU
23983 tetrapla Lk.	four-fold	A	or 4 mr.ap	Ap	... 1825.	C	m.s
23984 rubra Hud.	red Osier	A	8 ap.my	Ap	England os.hol.	C	m.s Eng. bot. 1113
23985 Forbyana Sm.	Forby's	A	clt 8 ap	Ap	England os.hol.	C	m.s Eng. bot. 13 i4
23986 Crownna Sm.	Crowe's	A	clt 8 ap.my	Ap	England mar.	C	m.s Eng. bot. 114i
23987 malifolia W.	Apple-leaved	A	or 3 ap	Ap	England moun.	C	m.s Eng. bot. 1(3)7
23988 Houstonidna Ph.	Houston's	A	or 4 ap.iny	Ap	Virginia	C	m.s
2J989 falcta Ph.	sickle-leaved	A	or 4 ap.my	Ap	N. Amer. 1811.	C	m.s
23990 Starkeuna W.	Starke's	A	or 4 ap.my	Ap	Silesia 1820.	C	m.s
23991 prumfulia Sm.	Plum-leaved	A	or 3 ap.my	Ap	Scotland scalp.	C	m.s Eng. bot. 1361
23992 A/yrsinUesL.	Myrsinc-like	A	or 3 ap.jn	Ap	Scotland scalp.	C	m.s Eng. bot. 1360
23993 Waldsteinftta W.	Waldstein's	A	4 ap.jn	Ap	Croatia 1822.	C	m.s
23994 venuldsia E. B.	veiny-leaved	A	or 2 apmy	Ap	Scotland sc. alp.	C	m.s Eng. bot. 1302
23995 planifolia Ph.	flat-leaved	A	or 2 ...	Ap	Labrador 1811.	C	m.s
23996 fuscata Ph.	brown-s/rmmrrfa	or	2 ap	Ap	N. Amer. 1811.	C	m.s
23997 wicciinifolia Sm.	Bilberry-leaved	A	or 2 ap.my	Ap	Scotland s.of sc.	C	m.s Eng. bot. 2341
23998 carinata Sm.	keol-icavcd	A	or 3 ap.my	Ap	Scotland sc. alp.	C	m.s Eng. bot. 1363
23999 corfiscans W.	glittering	A	or 3 ap.my	Ap	Germany 1818.	C	m.s Jac.au. 408
24000 criantha Schl.	woolly-flowered	A	or 2 ap	Ap	Switzerl. 1823.	C	m.s
24001 arbfscula Sm.	little tree	A	or 2 ap	Ap	Scotland sc. alp.	C	m.s Eng. bot. 1366
24002 livida Wahl.	livid	A	or 1 my	Ap	England hea.	C	m.s Wa. lap. 16.6
24003 polaris Wahl.	polar	•	or 1 -	Ap	Spitzberg. 1820.	C	m.s
24004 humilis I>fc.	humble	A	or 1 ap	Ap	... 1820.	C	m.s
24005 herbacea JL.	herbaceous	J*	or 1, J"	Ap	Britain sc. alp.	C	m.s Eng. bot. 1907
24006 ulmif&lia Thuil.	Elm-leaved	J*	or 1 apmy	Ap	Switzerl. 1821.	C	m.s
24007 arbutifolia W.	Arbutus-leaved	jK	or 1 apmy	Ap	Switzerl. 1818.	C	m.s

II. INTEGRJS.—Leaves smooth, entire.

Kitaibel's	J1	oi	1 ap.my	Ap	Carpathi. 2823.	C	m.s
blunt-leaved	■	or imy	Ap	Italy 1763.	C	m.s	
Thyme-leaved	■	or 1 my	Ap	Switzerl. 1818.	C	m.s	
nefted	■	or Jjnjl	Ap	Britain sc. alp.	C	co Eng. bot. 1903	
Myrtle-like	■	or 2 my	Ap	Sweden 1772.	C	rns.Vil.dcl3.50.il	
recurred./ftu/	■	or 3 ap	Ap	N. Amer. 1811.	C	m.s	
Bearberry like	■	or } ap.my	Ap	Labrador ...	C	m.s	
cordate-leaved	■	or 3 ...	Ap	N. Amer. 1811.	C	m.s	
pedicellated	■	or 3 ap	Ap	N. Amer. 1811.	C	m.s	

III. ViLLoSf.—Leaves vtilous.

24017 glaAca L.	glaucous	■	or 2 my	Ap	Scotland sc. alp.	C	m.s Eng. bot. 1810
24018 bricea Vil.	silky	or	2 my	Ap	S. Europe 1820.	C	m.s VilTdL3.51.27
24019 Janata L.	woolly	or	2 my	Ap	Scotland moun.	C	m.s
24020 Lappfonum L.	Laplanders*	ft	or 2 my.	Ap	Lapland 1812.	C	m.s Fl.lap.8.T
24021 arenaria L.	sand	Sk	or 3 my.jn	Ap	Scotland sc. a.p.	C	rnsEng.bot.lgg
24022 cinerea L.	grey	or	15 my	Ap	Britain woods.	C	m.s Eng. bot. 1397
L24023 obtusa Lk.	blunt	or	4 my	Ap	Switzerl. 1820.	C	m.s Eng. bot. 1397
24024 bicolor Ehrh.	two-colored	■	or 5 my	Ap	Hercynia 1820.	C	m.s
24025 Muhlenbergirina W.	Muhlenborg'B	or	3 ap	Ap	N. Amer. 1811.	C	m.s
24026 Jacquinina W.	Jacquin's	A	or 2 ap	Ap	Austria 1818.	C	m.s
24037 trsfis Mhl.	dark/cavcZ	A	or 4 ap	Ap	N. Amer. 1765.	C	m.s An. bot. 2.-^Q
24028 foliosa W.	leafy	A	or 6 ap	Ap	I*pland 1818.	C	m.s Jac. au. 1.4tt*
24029 longifolia Mhl.	long-leaved	A	or 10 -	Ap	Pennsylv. ...	C	m.s
24030 argentea Sm.	silvery, sand	■	or 3 my"	Ap	Britain san.sh.	C	m.s Eng. bot. 13**
24031 Iucophylla W.	white-leaved	A	or 4 my	Ap	Europe 1824.	C	m.s
24032 tlxagnoidcs Schl.	Elaeagnus-Zw/	■	or 4 my	Ap	Europe 1824.	C	m.s
24033 ripens W.	creeping	■	or 2 my	Ap	Britain sa.hca.G	C	m.s Eng. bot. 185
24034 fusca L.	brown	A	or 2 my	Ap	Britain m.hea.	C	m.s Eng. bot. Wxj
24035 cassia Vil.	grey	A	or 3 my	Ap	S. Europe 1824.	C	m.s Vd.deL3..*5'
24036 prostrata Sm.	prostrate	■ ■	L my.jn	Ap	Britain m.alp.	C	m.s Eng. bot. l>5'
24037 Schradeiana W.	Schrader's	■	or 2 my	Ap	Germany 182a	C	m.s
24038 nyrcnica Gou.	Pyrenean	A	or 1 my	Ap	Pyrenees 1823.	C	m.s
24039 hirta Sm.	hairy-branched	Y	or 15 ap.my	Ap	England woods.	C	m.s Eng. bot. Ij
24040 Dicksonwhl Sm.	Dickson's	A	or 1 ap	Ap	Scotland sc. alp.	C	m.s Eng. hot W
244&1 parviflha Sm.	small-leaved	A	or 2 ap.my	Ap	England moi.h.	C	m.s Eng. bot. iyg
24042 adscendens Sm.	ascending	J*	or J ap.my	Ap	England moi.h.	C	m.s Eng. bot. 19^
24043 incubacea W.	trailing	A	or 3 my	Ap	Europe 1775.	C	m.s
24044 Dontfana Sm.	Don's	A	or 6 my	Ap	Scotland moun.	C	m.s
24045 roBinarinifolia TV.	llosmary.lvd	A	or 2 ap.my	Ap	Britain san.pl.	C	m.s
24046 riparia W.	bank	A	or 6 ap.my	Ap	Germany 1821.	C	m.s
24047 angustiflba W.	narrow.icavcd	A	or 3 ap.my	Ap	Caspian 1825.	C	in.s
24048 grisea W.	grizzly	A	or 6 ap.my	Ap	Pennsylv. 1820.	C	m.s

24049 spatulata W.	spatulate	A	or	5	ap.ray	Ap	Germany	1818.	C	m.s	
24050 aurita L.	cared	• A	dt	2	ap.my	Ap	Europe	1830.	C	m.s Hof. sal. 1.22 1	
24051 uliginosa W. en.	marsh	A	or	2	apjn	Ap	Britain	woods. C	m.s	Eng. bot 1487	
aurita £. £.											
T*052 aquatica Sm.	water	^	or	10	ap	Ap	Britain	w. thi. C	m.s	Eng. bot. 1437	
24033 oleifolia TV.	Olive-leaved	S*	or	4	mr	Ap	Britain	thick. C	m.s	Eng. bot. 1402	
*054 cotinifolia Sm.	Cotinus-leaved	A	or	2	ap	Ap	Britain	woods. C	m.s	Eng. bot. 1403	
*4055 sphacelata TV.	withered-point.	A	or	2	apmy	Ap	Scotland	sc. alp. C	m.s	Eng. bot. 2333	
24056 caprea L.	goat, great round-lod.^	A	or	30	ap.my	Ap	Britain	dr. wo. C	m.s	Eng. bot. 1488	
24057 Stuarttina E. B.	Stuart's <	A	or	4	jl.au	Ap	Scotland	sc. alp. C	m.s	Eng. bot. 2586	
*J058 acuminate //q/m.	acute	J	or	15	ap	Ap	Britain	moi.w. C	m.s	Eng. bot. 1434	
J059 conifera FFan.	cone-bearing	J	or	10	my	Ap	N. Amer.	1820. C	m.s	Wang. 31.72	
*4060 viminalis L.	twiggy, common <i>Ösier</i>		clt	12	ap.my	Ap	Germany	1800. C	m.s		
24061 mollissima EArA.	softest	Y	or	20	ap.my	Ap	Germany	...	C		
2*052 Smithia a W.	Smith's	\$	or	20	...	Ap	England	os. gr. C	m.s	Eng. bot. 1509	
ORINN	mollissima £. B.										
24063 stipitKrb ff.	stipular	■	clt	d	mr.ap	Ap	England	os. gr. C	m.s	Eng. bot. 1214	
24CM Candia IK	white	■	or	10	ap.my	Ap	N. Amer.	1811. C	m.s		
24065 FluggeAna W.	Flugge's	■	or	10	ap.my	Ap	S. France	1820. C	m.s	Vil. del. 3.51. 28	
ORAN	patulae & r.										
?JSS < «Mei JE. A	common, white	Y	clt	40	ap.my	Ap	Britain	woods. C	m.s	Eng. bot. 2430	
*168 rup&tris Sm.	blue	J	or	40	apmy	Ap	England	m. me. C	m.s	Eng. bot. 2431	
*169 Addorsandtff E. B.	silky, rock	A	or	3	ap	Ap	Scotland	sc. alp. C	m.s	Eng. bot. 2342	
Forstnera E. B.	A. Anderson's	A	or	3	ap.my	Ap	Scotland	sc. mo. C	m.s	Eng. bot. 2343	
Forster's	Forster's	Y	or	10	ap.my	Ap	Scotland	sc. wo. C	m.s	Eng. bot. 2344	
*J071 heterophylla Deb.	various-leaved	A	or	...	ap.my	Ap	Europe	1823. C	m.s		
*J072 finmarctica TV.	Finmark		or	10	ap.my	Ap	Sweden	1825. C	m.s		
2J73 holaoicea TV.	velvety		or	8	ap.my	Ap	Germany	1822. C	m.s		
24074 obtusifolia TV.	blunt-leaved		or	...	ap.my	Ap	Lapland	1818. C	m.s		
chrysanthos Oed.	golden-flwd		or	...	ap.my	Ap	Norway	...	m.s	FI. dan. 1007	
	IV. HELVEV.						Swiss species not yet described.				
J4076 atrofrens Schl.	dark-green	or	...	ap.my	Ap	SwitzerL	1824	C	m.s		
24077 australis Schl.	southern	or	-	ap.ray	Ap	SwitzerL	1824	C	m.s		
24078 carpiniMia Schl.	Carpinus-lvd	or	-	mr.ap	Ap	SwitzerL	Ib24.	C	m.a		
24079 cydonifMia Schl.	Quince-leaved	or	-	mr.ap	Ap	SwitzerL	1824.	C	m.s		
24080 subalpina Schl.	subalpine	or	-	ap.my	Ap	SwitzerL	1820.	C	m.s		
24081 QTolearfbia Schl.	Protea-lcaved	A	or	-	ap.my	Ap	SwitzerL	1820.	C	m.a	
24082 strpida Schl.	creaking	A	or	-	ap.my	Ap	SwitzerL	1820.	C	m.s	
24083 glabriata Schl.	smooth	A	or	-	ap.my	Ap	SwitzerL	1820.	C	m.s	
24084 decumbens ScA.	decumbent	A	or	-	ap.my	Ap	SwitzerL	182a	C	m.8	
24085 Jurtina Schl.	wall	A	or	-	mr.ap	Ap	SwitzerL	1824.	C	m.8	
24086 nervos.i ScA.	nervose	at	or	-	mr.ap	Ap	SwitzerL	1824.	C	m.s	
24087 pannosa Schl.	cloth-leaved	A	or	-	ap.my	Ap	SwitzerL	1824.	C	m.8	
24088 pallidescens Schl.	pallidescens	A	or	-	ap.my	Ap	SwitzerL	1823.	C	m.s	
24089 pallida ScA.	pale	A	or	-	ap.my	Ap	SwitzerL	1823.	C	m.s	
24090 Kilaternoldcs Schl.	Alatcmus-like	A	or	-	ap.my	Ap	SwitzerL	1824.	C	m.s	
24091 albdscens ScA.	albescent	tt	or	-	ap.my	Ap	SwitzerL	1824.	C	m.s	
24092 cerasifera ScA.	Cherry-leaved	A	or	-	ap.my	Ap	SwitzerL	1824.	C	m.s	
*J093 cinnambmia Schl.	Cinnamon-SC/f	A	or	-	ap.my	Ap	SwitzerL	1824.	C	m.s	
gWW clethraffMia ScA.	Clethra-lcaved	A	or	-	ap.my	Ap	SwitzerL	1824.	C	m.s	
24095 griseophylla ScA.	crcty-leaved	A	or	-	ap.my	Ap	SwitzerL	1824.	C	m.8	
24096 lacriflora ScA.	lake	A	or	-	ap.my	Ap	SwitzerL	1824.	C	m.8	
24097 mespilifolia Schl.	Mespilus-lvd		or	-	ap.my	Ap	SwitzerL	1824.	C	m.a	
24098 obtuso-serratd Schl.	blunt-serrated		or	-	ap.my	Ap	SwitzerL	1824.	C	m.s	
Si?? OTUWIA Schl.	Pyrus-leaved		or	-	ap.my	Ap	SwitzerL	1824.	C	m.s	
24100 Schleicherid/M Schl.	Schleicher's		or	-	ap.my	Ap	SwitzerL	1824.	C	m.8	
2733 CEC(TPIA	L. SNAKE WOOD.		(Cccrops, king of Athens, his legs fabled to be snakes.)				Urticae. 3.				
24101 peltKta L.	peltate		to 9r	30	...	Ap	Jamaica	1778.	C	pi Lam. 11.800	
24102 palmXta W.	palmate		•	20	...	Ap	Brazil	1820.	C	p:	
24103 concolor TV.	tcfl-colored		•	20	...	Ap	Brazil	1822.	C	p.	
2734. BCTRYi* ■	BORYA.	(A. Bory de St. Vincent, a French botanist.)					Euphorbiacea?. 6.-7.				
24104 porulosa TV.	small porous			-	-	G	Florida	1806.	C	m.s	
24105 ligustrina TV.	Privet-leaved			-	-	G	N. Amer.	1812.	C	m.s	
24106 acuminate TV.	pointed			-	-	G	N. Amer.	1812.	C	m.s Mic.am.2.28	
24107 prinoides TV.	Prinos-like			-	-	G	N. Amer.	1824.	C	m.8	
241f.8 nitida TV.	shining			-	-	G	N. Amer.	1824.	C	m.s	
24109 retissa TV.	retuse			-	-	G	N. Amer.	1824.	C	m.s	
VALLISNE^Ril L.	VALLISNERIA.	{Antonio VaUlsneri, an Italian bot)					Hydrochartdetc. 1.-4.				
H. spiralis L.	spiral	^ ^ Je u...	jl	Br	S. Europe	1818.	S	aq	Mic.gen.	12.10.1	

CLASS XXII. ORDERS.


DICE'CIA TBLA'NDftIA:

2739. HYPOTJEW A *It. Br.* HYPOLJENA. (*Ifypo*, under, *chaina*, cloak; base of fruit.) *Resttcea?*. 2.
 24115 fastigiata/f.fc. peaked JUILAICU 2 Jn.jl Ap N. Holl. 1820. R ap
 24116 exsu lea/*.f?/. furrowed M iAI cu 2 jn.jl Ap N. Holl. 1821. It s.p
3740. LEPYRO^XDIA *It. Br.* LEP. {*Lepyrodea*, scaly; bractes within scales of spike.) *Restiace?*. 1.—4
 24117 gracilis *R. Br.* slender Jn iAI cu 2 my.jl Ap N. Holl. 1824. It s.p
2741. THAMNOCHO'RTUS *Berg.* THAMNOC. (*Thamnos*, shrub, *chortos*, grass; habit) *Restiuccce*. 1.—3.
 24118 dichotomus A. i?r. two-forked M i_J CU 2 jn.jl Ap N. Holl. 1817. A s.p Rtb.gr. 4. 2. 1
Restio dichotomus Thun.
2742. RE'STIO L. ROPE GRASS. (*Restis*, cord; use at Cape of Good Hope.) *Restiicea*. 9.—52.
- I. SI'MPLICES. — *Culm simple*.
- 24119 tectorura 7%tf». thatch Jb iAJ un 3 my.jn Ap C. G. H. 1793. R s.p Rtb.gr. 10.3.2
 24120 virgitiu JJtf. i twiggly Jit TAJ un 3 my.jn Ap C. G. H. 1824. R s.l Rtb.gr. 5.1. 2
 24121 vaginthus *Thun.* sheathed Jb LAJ un 3 my.jn Ap C. G. H. 1820. R s.p
 24122 australis /«. 5r. southern Jb iAJ cu 3 my.jn Ap N. Holl. 1824. R s.p
 24123 gracilis *R. Br.* slender Jk TAJ cu 3 my.jn Ap N. Holl. 1824. R 8.1
- II. IU.Mosi. — *Culm branched*.
- 24124 tetraphf ilus *R. Br.* four, leaved Jib TAJ cu 3 my.jn Ap V. Di. Is. 1825. R ■ Lab. n. h. 2.226
 24125 paniculatus *Rb.* 'paniced Jb iAJ un 2 my.jn Ap C. G. H. 1824. R s.p Rtb.gr. 4. 2. 3
 24126 latoriflbrus *It. Br.* side-flowering Jt iAIcu 3 my.jn Ap N. Holl. 1824. R 8J Lab.n.h.2.228
Calopropus elongatus Lab.
 24127 festigatus ft *Br.* peaked JllriAIcu 3 my.jn Ap N. Holl. 1824. R 6.1
2743. LEPTOCA'RPUSS *R. Br.* LEPTOCARPUS. (*Leptos*, slender, *karoos*, fruit; *r*oointed.) *Restidcear*. 1.—7.
 24128 tēnax *R. Br.* tough M iel un 2 ... Ap N. Holl. 1813. D co Lab.n.h.229
Schoenbium tenax Lab.
2744. J2LEGPA *Thun.* ERLGIA. (*Elegos*, lamentation: sad color of plants.) *Restihcexe*. 2. - 3.
 24129/Giicea *Thun.* Rush-like JtiAlun 1 jl.au Ap C. G. H. 1789. D l.p Rtb.gr. 8.3.4
 24130 racemosa *Lam.* clustered JtfciAlun 1 my.jn Ap C. G. H. 1804. D i p Lam. il. 884.4
2745. PHOENIX L. DATE PALM. (The Greek name of the date.) *Palm**. 8.
- 24131 dactylifera L. com., date-bearing i_J fr 40 - WW Levant 1857. ■ r.m Kami. 686.1.2
 24132 reclinMa *Jac.* reclined £ i_I or 10 - W.o C. G. H. 1792. ■ l.p Jac.fr. 27.24
 24133 farinifera *Box.* small, meal-bg £ CH or 28 - W.o E. Indies 1800. ■ r.m Rox. cor. 1.74
 24134 aca filis *Box.* stemless IQor 6 - W.G E. Indies 1816. ■ rm
 24135 paludosa *Rax.* marsh £ O or 20 - E. Indies 1820. ■ r.m
 24135 pygmaAi *Lo. C.* pygmy £ • or 6 - Mauritius 1823. ■ r.m
 24137 sylvestris *Rox.* wood £ • or 20 - E. Indies 1823. ■ r.m
 24138 leonensis *Lo. C.* Sierra Leone * I or 30 - S. Leone 18:23. ■ r.m
2746. STILAN30 L. SriLAno. (*Stylos*, a style; probably so called from its length.) *AntuUsmcc*. 2.
 24139 JKiniua *L.* Bunius, Laurel-ldvf CD or 20 au Ap E. Indies 1757. C p.l R. inal.4. 56
 24140 diundra *Rox.* diaudrous f O or 20 ... Ap E. Indies 1800. C p.l Rox. cor. 2.166
2747. OSVMSLam. POF.TU CASSIA. (*Ozoz*, a branch: numerous supple branches.) *Osyrlde*. 1—2.
 24141 alba *Lam.* white ■ J o r 3 ... W S. Europe 1739. C l.p Lam. il 802

CLASS XXII. ORDER 4.

DI(E%)CIA TETRA'NDRIA.

- 2748. AU'LAX *Berg.* AULAX. *lAulax*, a furrow; furrowed leaves of first species) *Proleucete* '2.
 24142 pinifl'ia *Berg.* Pine-leaved ft L-I or 2 jl.s Y C. G. H. 1780. S l.p Hot. rep. 76
 24143 umbellati it. *Br.* umbelled « i_J or 2 jn.au Y C. G. H. 1774. S Lp Hot. rep. 248
2749. LEUCADE'NDRON L. LEUCADENDRON. (*Leukos*, white, *dendron*, tree; appearance.) *Prot. S9.—4*·
- I. A nc vHODE'Nnos. — *Xut ventricose s style and calyx persistent; leaves mostly silvery.*
 24144 argenteum *R. Br.* silver free il Jor 15 au Y C. G. H. 1790. ■ Lp Hot. reg. 979
 24145 plumosum *R. Br.* fcRther-Jiivd Box-loaved il I or 4 - Y C. G. H. 1774. ■ Lp
 24146 sericeum *R. Br.* silky ■ 3 apmy Y C. G. H. 1817. ■ Lp
 24147 spatulatum II. *Br.* spatnlate ■ i_J or 3 apmy Y C. G. H. 1818. ■ Lp
 24148 rectium *R. Br.* rectusc ■ L-J or 3 apmy Y C. G. H. 1810. ■ Lp
- II. LEVISA^XNUR. — *Ktt ventricose or lenticular, wingless, on both sides or margins pilose s style totally deciduous or wth the base alone refrainine; calyx a long time persistent, ^-parted.*
 24149 imbricatum *R. Br.* imbricated ** i_J or 4 - Y C. G. H. 1790. ■ Lp
 24150 AuxiflMiu *Ji. Br.* Box-loaved il I or 4 - Y C. G. H. 1812. ■ Lp
 24151 Levhdnus *R. Br.* Lewis's ft ■ or 4 aprjn Y C. G. H. 1774. ■ Lp Bur. af. 100.2
 24152 angustatum *R. Br.* narrowed il I or 3 JnJl Y C. G. H. 1K20. ■ Lp
 24153 /imflbrum *R. Br.* Flax-leaved il I or 4 ap.jn Y C. G. H. - ■ Lp Jac. sc. 1.26
 24154 fusciflorum *R. Br.* dusky-flowered il LJ or 4 my.jn Y C. G. H. - ■ Lp Bot. mag. 881
Prnte stellhris B. M.
 24155 tortum *R. Rr.* thtcd-leaved** i_J or 3 mr.my Y C. G. H. 1790. ■ Lp Bot.rog.82G
 24156 cinfrcum // *Br.* grey tt i_J or 3 Jl.au Y C. G. H. 1774. ■ Lp
 24157 corybosum *R. Br.* corymbed A i_J or 3 ap.jl Y C. G. H. 1790. ■ Lp
- III. SAMARIFERA. — *Samara smooth, winged or wingless s style and calyx deciduous ; scales of strobile distil'*
 24158 decbrum *R. Br.* decorous il LJ or 3 ... Y C. G. H. 1790. C l.p
 24159 squarrbsum ?*Br.* squarrose il I or 3 my.jl ■ C. G. H. 1824. ■ Lp
 24160 concolor 71. *Br.* self-colored il t- or 3 mr.jn ■ C. G. H. 1774. C Lp Bot. rep. 307
 2-1161 grandiflbrum *R. Br.* great-flowered it i- or 3 ap.jn ■ C. G. H. 1789. C Lp Par. Ion. 105
 24162 decurrents *R. Br.* decurrent il I or 3 ... C. G. H. 1812. ■ Lp
 24163 ovalc *R. Br.* oval il i_J or 3 ap.jl ■ C. G. H. 1818. ■ Lp
 24164 ygnsbun *R. Br.* veiny * i_J or 3 ap.jn ■ C. G. H. 1816. ■ Lp
 24165 K^{IND}ru « A *Br.* smooth il L-I or 3 ap.jn ■ C. G. H. 1810. ■ Lp
 24166 Ktrictum *R. Br.* strict il i_J or 3 ap.jn ■ C. G. H. 1795. ■ Lp Par. Ion. ■
 L'il67 virgatum if. *Br.* twiggy My \ or 3 ap.jn ■ C. G. H. ■ Lp

- 21168 adscendens R. Br. ascending <Ljor C.G.U. 1774. C Lp Pluk. m. 229. G
 24169 coecinnum R. Br. neat m LJ or C. G. H. 18X. C l.p
¹⁷⁰ faltum R. Br. Willow-treefir • iJ or C. G. H. 1774. C l.p Boer. 2.204
 swamp « uJ or C. G. H. 1795. C Lp Brey. c. 21.9
 M. A. - florid #LJ or C. G. H. 1795. C Lp Bot rep. 572
 smooth.
 IV. PLATYSPE/RMA. — Scales of strobile connate; samara Lp
 flat-scaded *i_ | or 3 apj Y C. (x. H. 18 C l.p
 SL te* 2 Hoi ijls⁸ Y C.G.H. 178J. C Lp Bot rep. 429
 rough » LJ or 3 ... Y C. G. H. 1812. C Lp Bot rep. 461
 C Lp
 V. DU'HA Doubtful to which of the sections they belong. C Lp Lam. il. 1.53
 Globularia « U o r 3 apjn Y C. G. H. 1810. C Lp
 inflexed » uJ 6 r 3 apjn Y C. G. H. 1800. C Lp
 tailed « L J or 3 apjn Y C. G. H. 1800. C Lp
 marginate » L J or 3 apjn Y C. G. H. 1800. C Lp
 pubescent *L J or 3 ap. h Y C. G. H. 1819. C Lp
 MISTLETOE. (*Viscus*, clammy j nature of berries.) *Lordnthece*. 1. — 25.
 comtnon, white J5 cu 2 my G England trees. S ma Eng. bot 1470
 C Lp
 CANDLEBERRY MYBTLB. (*Myro*, to flow; found on banks of rivers.) *AmeHdce**. 13-SL
 Sweet Gale » or 4 my Ap Britain sp. bo. L s.p Eng. bot. 562
¹⁴ 85 cerifera L. com. wax-bg N. Amer. If99. S. s.p Cat. car. UK
 QH* carolinensis wan. Carolina Azores 1777. L s.l Duh.no. 2.55
¹⁴ 86 mucalam. sylvanL. tiopian 1705. L s.l Pluk. al. 48.8
 saw-leaved ft C. G. H. 1793. L s.l Pluk. am. 424.3
 jagged Oah-lyd C. G. H. 1752. L s.l Jac. fr. 2. L 4
 Affl' Oak-lyd C. G. H. 1752. L r.m
 heart-leaved C. G. H. 1759. L p.l Pluk. aL 319.7
 Mexican Mexico 1823. L p.l
 separated S. Amer. 1824. L p.l Jac.ic.G25
 esculent Nepal 1817. L s.l
 2752. NAG&IA Gac. NACEIA. (NagiAts Japanese name.) *Amaphthca*: 1.—a
 24197 Putranflua Rox. Putranjiva « D « n IS » Ap E. Indies 1892. C r.m
 2753. DIY'PETES Vahl DUYPETBB. (to lacerate Rh&mnee. 1.-4.
^{2U98cr^ceaPo,7.} copper-cofom* *C D or 1820. C Lp
 Schafft'n a laterrilla Stuz.
 2754. SHEPHE'UD/J Nut SnEPHEttDIA. (ffk. Shepherd, curator of bot gard. IIverpool) *Elaagnex*. 2.
^{*liWcanadensis N<.} Canadian £ or 10 ap.my Ap N. Amer. 1759. L co
 T d * t J or 10 ap.my Ap Missouri 1818. L p.l
 argenteaPA.
 2755. HIPPO'PHAE L. SEA BucRthonN. (Hippos, horse, pAoo, to destroy.) *Elaagine**. 2.
 S Si Hiamnddes X. «>< Khamnus-lik^a or 12 ap.my Ap England sea co. Sk co tng.bot 425
 2*202 *alicif6lia D. Don Willow-leaved » or 8 ... Ap Nepal 1822. L Lp
 2756. BROUSSONETT/ll Ven. BUOUBSOXETIA. (P. N. V. Broussonet, a French naturalist) *Urtiae*. 2.
 Papryrlfera Ven. paper-bearing ^ »^r₁₂ fa ^ P^{III}P^{an} WSl. Skco Ktem.472
 spatulate/w* ^ or 12 fs Ap 1824. C co.
 2757. SCHJEFFE'R7^ Jac. SCHJSFFBJIIA. (James Christian Schaffer, * German naturalist) *JUtlmnca*. 1-SL
 aOS complete Swz. complete, white-fitw O or (i au G W. Indies 1793. C p.l Lam.il. 809
 BBUCBA. LJ»<< ^ aeg, celeb, traveller in Abyssinia.) rrwAfe/Aa<r«, a
 SIS⁵ ferruginea Herit. rusty, Ash-lyd mCD or 6 apmy C Abyssina 1775. C p.l Bot cab. 129^a
 4207 sumatranra Box Sumatra » or 10 ap.my G K Indies 1820. C p.l
 21208 gracilisLref Ssnder » or 6 ... Y.G E. Indies 1820. C p.l
 2759. TETRADODUMMIAL TETRADIUM. (Tetradion, a quaternion; ffuctification) *Terebinthdcete*. 1
 14U trichotomia I^. thre-forked ± [±] J or 20 China 1820. C p.l
 Bracea trichotoma Spr.
 *AV^{C10}, ANTHOSPE'RMUM L. AMBERTBEE. (*Anthos*, flower, sperma, seed.) *Rubiaceee*. 1.—G.
 24210 ffithiopicumX,. Ethiopian *uJ or 2 jn.jl G.w C. G. H. 1G92. C p.l Pluk. al. 183.1
 S76I Rwoov TREE. (Trophe, fodder; cattle oat in scarcity.) *Thi'melca*. 2. — fil
 24211 americana L. American 4 a or 20 ap.my G W. Indies 1789. C l.p Br.jam.37.1
 24212 asperaArfe. rough-leaved I D or 25 ... G E. Indies 1802. C l.p
 87G2. MONTI'N/^ L. MoMwiA. (Laurence Montin, a Swedish botanist) *lrite*. 1
 24213 cnyophylftcca /. PIBTACHIA TREE. (Fbustaq, its Arabic name.) *Terebinthheeee*. 5.-8.
 24214 officinarum H. K. L Te.nnm/TITHva. —Leaves impari-pinnat; deciduous.
 24215 reticulata W. narbonefs'w L wleaved j S 15' S?"y IS K2K BB¹ S¹ C Lp Rauvit.729
^{2*216} cerebinthus L. tree* 
¹² atlantica Dcsf. 

CLASS XXII. ORDER 5.

DIOECIA PENTA'NDRIA.

2763. PISTATIA /. PIBTACHIA TREE. (Fbustaq, its Arabic name.) *Terebinthheeee*. 5.-8.
 24214 officinarum H. K. L Te.nnm/TITHva. —Leaves impari-pinnat; deciduous.
 24215 reticulata W. narbonefs'w L wleaved j S 15' S?"y IS K2K BB¹ S¹ C Lp Rauvit.729
^{2*216} cerebinthus L. tree* 
¹² atlantica Dcsf. 

- II. LENTI'SCL'S.—leaves abruptly pinnate; remaining.
- 24818 7-entiscus L. Mastich tree • uJ cc 15 my Ap S. Europe 1664. L r.m Bot. mag. 1967
2 massiliensis Mil. Marseilles \$ uJ < 15 my Ap. S. Europe 1661. L r.m
2764. Z ANTHO'XYLUM L. TOOTHACHE TREE_{to} (*Xanthos*, yellow, *xylon*, wood.) *Rutdcea*. 14.—43.
S419 emarginatum Sun. notch-leaved • O or — G.W. Jamaica 1739. C l.p SI. jam. 2.168.4
24220 clava Herculw L. Hercules's club « uJ or 10 ap.my G.W. W. Indies 1739. C l.p Cat. car. 1.26
24221/raxineum Wf com, Ash-like ■ or 6 mr.ap G.W. N. Amer. 1759. L s.l Duh. ar. 1.97
24222 tricarpum Mx. three-capsuled ft or 6 ... G.W. N. Amer. 1806. L 18
24223 nitidum Dec. shining M Q or 6 ap.my G.W. China 1823. L r.m Bot mag. 2558
24224 Avicinna's " or 10 — China 1823. C l.p Lob. ic. 2.133.2
24225 acuminkatum Swz. acuminate or 10 — Jamaica 1818. L p
24226 nrte W. mild life or 10 — N. Amer. 1818. L p
24227 aromaticum W. aromatic i f l or 10 — W. Indies 1824. L p
24228 armatum Rox. armed if l or 20 — E. Indies 1816. L p Jac. ec. 1. 70
24229 sapindoides Dec. Sapindus-hke f l or 20 — Jamaica 1823. L p Br. jam. 20.3
24230 australie G. Don Figara australis Cun.
F d. Budrunga Rox. silvery • Dor 10 — 1825. C l.p
24231 Budrunga Dec. Budrunga !Dor 20 — E. Indies 1825. C l.p
- \$765. PICRA'MNI A Swz. PICRAMNIA. (*Pikros*, bitter, *thamnos*, shrub; whole plant.) *Cassuviv*. 2.
24233 Antidesma Sun. Antidisma il • un 4 ... G. Jamaica 1793. C p.l SI. jam. 2.208. S
24234 pentandra Suns. pentandrous H Q o r 5 ... G. W. Indies 1822. C p.l
2766. ANTIDE'SMA L. ANTIDESMA. (*Anti*, like, *desmos*, bond; bark used for ropes.) *Antidesmets*. 5.—11.
24235 alexiteria L. Alexiteria i Q u n 10 my.jn Ap E. Indies 1793. C p.l K.mal.5.11
24236 paniculata Box. panicled ■ Q ■ 10 — Ap E. Indies 1800. L p
24237 zeylanica Lam. Ceylon ■ CD or 6 — Ap Ceylon 1821. p. Bur. zey. 10
24238 pubescens Rox. pubescent ■ 6 — Ap E. Indies 1818. p. Rox. cor. 2.167
24239 guineensis G. Don Guinea ■ ■ 6 — Ap Guinea 1825. L p
2767. IRESRNE L. IRESINE. (*Ebros*. wool; woolly appearance of branches.) *Amaranthi*c*g*. 5.-8.
24240 celosio'ides L. Celosia-like tf Al pr 1jl.au W S. Amer. 1733. S l.p Lam. il. 813
24241 elongata Hum. <mg-leaved ■ Q ■ 2 jl.au W S. Amer. 1822. L p Pluk.al.26L1
24242 diffusa W. straggling ■ J pr n jl.au W S. Amer. 1818. L p
24243 elatior Rich. taller iQJor 3 jl.au W Antilles 1820. L p SLjam. L90
24244 flavescens W. flavescent ■ 1 jlau W S. Amer. 1824. L p
2768. SPINA" CIA"Z. SPINACH/ (*Spina*, a prickle; prickly integument of fruit) *Chenopdeia*. 1. — S.
24245 oleracea L. common, potherb O cul li mr.o G. 1568. S co Schk. han. 3.32*
1 spinosa prickly O cul H mr.o G. S co
2 glabra Mil. smooth O cul H mr.o G. S co
2769. FLU'GGGEI< W. FLITGGEA. (*John Flugge*, a German cryptogamic botanist.) *Euphorbiucca*. 1.
24246 leucop^rus W. white wheat H un 6 ... Ap "E. Indies 1825. C r.m
2770. ACNPDA L. VIRGINIAN HEMP. (A, priv., knide, nettle; nettle-like, without stings.) *Chnop*. 1.-&
24247 cannabinia L. common, Hemp-lik. O un 2 jn.jl G.v N. Amer 164a S co
2771. CA'NNABIS /,. HEMP. (*Can*, reed, *ab*, small, *Celt*; or *qanb*, its Arabic name.) *Urticeee*. 2...
24248 sativa L. common, cultivated O ag fi jn.jl G.Y India ... S h.l Schk. han. 3.32B
24249 indica L. Indian Tobacco O clt U au.s G E. Indies 1800. S co
- 2772 HITMULUS L. HOP. (*Humus*, the ground; if not supported creeps along it) *Urtlccar*. 1-
24250 Lbpulus L. common, Mop £ A *8 15 J n a u Y Britain hed. D r.m Eng.bot.427
2773. MODECCA Jac. MODECCA. (Its Indian name.) *Pasiifibrea*. 4.
24251 lobata Jac.)obeA-kayed B CD or 12 au G S. Leone 1812. C r.m Bot reg. 433
24252 tuberosa Rox. tuberous ■ " or 10 jls — E. Indies 1822. C 81 Ru.am.8.20
24253 dubia Rox. doubtful LD or 10 E. Indies 1826. C l.p
24254 trilobata Rox. three-lobed ■ ■ 10 jls . . E. Indies 1818. C Lp
2774. NEGU'NDO Moe. Box ELDER. (Meaning unknown.) *Acerinee*. 1.— 3-
24255 Jraxinfblium Nut. Ash-leaved ¥ or 35 ap G N. Amer. 1688. L s.l Schm. ar. 1.1*
Afcrc Negundo L. Negundium americanum Dec.
2 crispum curled *£ or 30 ap G N. Amer. 1688. L s.l
2775. NEMOPA'NTHES Raft. NEMOPANTHEB. (*Nemos*, grove, *ops*, eye, *anthe*, flower.) *Celastrine**.
24256 canadensis Dec. Canadian * or 6 N. Amer. 1812. L p.l Dec. gen. *
fascicularis Raft. /*lex canadensis Mx.

CLASS XXII. ORDER 6.

DIQECIA HEXA'NDRIA.

2776. XE'ROTES R. Br. XEROTES. (*Xerotes*, dryness; aridity of herbage.) 12--^{2*1}
- I. CAPITA'TJE. — Female flowers capitate i leaves at top entire.
- 24257 flexiblia R. Br. curled-leaved £ iAl or 1 ... W.G. N. Holl. 1824. D r.m , _ 17 \$
24258 mucronata R. Br. mucronate £ IAJ or 2 ... W.o N. Holl. 1824. D r.m Thun.4**'
- <rtc[^] " " RACE MOSE Flowers racemose, or female ones spcate, male ones PTM*cjrd.
24259 guesseffii Br. Bifoliate tf iAl or 1 ... WVGG N. Holl. 1824. Dr.m Thun.iu t ^
24260 gradiiflora R. Br. SchAn tf tftSoS Ur 22 ... WVGG N. Holl. 182a D r.m
deJticate | S or 1 ... WVGG NN.Holl. 1826. D r.m
loose £ ubJ or 1 .. W. N. Holl. 1823. D r.m

HI. DENTICULAR*. - Flowers spiked or panicle; branches opposite or whorled j male flowers sessile imbricated, with bracts; capsules smooth s leaves at 400 goutiulates stemless.

Montana R. Br.	mountain		N : H :	1824.	D r.m.
long-leaved	£ iAI or 3	■	G.w.	N. Holl.	179b. D r.m.
Istrix il.*t.	porcupine	£ iAI or Q	■	W.G.	N. Holl. 1824. D r.m.
S^67 arenina A ?;	sand	£ iAI or 2	■	W.G.	N. Holl. 1820. D r.m.
**268 ecbinata Ma. C.	hedgehog	£ iAI or 2	■	W.G.	N. Holl. 1824. D r.m.
ELJEMS Jac.	OILY PALM of Elaia, olive; natives of Guinea express an oil from the fruit) Pdlmte. 4.				
melanoœcea Chr.	black-seeded	£ Q or 36	... G.W.	fenad.	1821. b co
occidentalis W.	western	£ CD or 30	... G.W.	Jamaica	1820. r.m
Quilk Guineensis Jac.	Guinea	£ • or 30	■ G.W.	Guinea	1730. r.m. Jac. am. 172
**72 pernambucana Lo.	C. Pernambuca	£ Q or	■ G.W.	Brazil	1825. r.m
*. CHAMIEDE*REA W.	CHIAMASDOREA. (Chamai, ground, dorea, gift; flowers low.) TM "JJ" S^io				
grAcilis IV.	slender	£ a or r 10	... W.o.	Caraccas	1803. Sk r.m. Jac. sc.2, & 7, 5*
fragrans Mart.	sweet-scented	£ • or 8	... W.	Trinidad	1820. Sk r.m. Mart. palm. ic.
Nundzia frkgrans W.					
SdSp' HYPJH^NE Gae.	HYPILJENE. (Hypaino, to entwine; fibres of fruit.) Pdlmte. 1.				
***75 coriacea Gae.	le&thet-leaved	■ or 20	Egypt	1824. S r.m.	Gae.fr. 84.
CucSferathebaicaDei.					
A. 2 ⁸ 0- BORA'SSUS Z.	BORASBVS. (One of the names applied to the spatha of the date.) Palme. 1.—&				
2276 flabelliformis L.	fan-leaved	£ a or 30	... W.G.	E. Indies	1771. S r.m. Rox. cor. 1.71, 72
2781. MAUR'I'THil L. fil. MAUBITIA. (Prince Maurice, of Nassau, the patron of Pisso.) Pdlmte. 3. - 4.					
*277 flexuosa L.M.	flexuous-*m*xi	£ CD or 40	... W.o.	Surinam	1818. S F.M. Mart. palm. ic.
916 278 *Uwi Men.	armed	£ • or	■ Brazil	1824. S r.m. Mart. palm. ic.	
2429 Vintera Mart.	wine-bearing	£ St. or 0	■ Maranh.	1824. S r.m. Mart. palm. ic.	
, 2782. GE(yNOMA W.	GEONOMA. (Geonomos, skilled in agriculture; propagation.) Pdlmte. 6. — 13				
24280 Spixiella Mart	Spix's	1 • or 15	■ Pk	Brazil	1824. S r.m. Mart. palm. ic.
21281 Schotttim Lo. C.	Sl-hots	1 • or.	■ Ap	Brazil	1820. S r.m.
■ pmtifrons W.	pinnate-leaved	± • or	■ Ap	Caracas	1821. S r.m.
■ simplicifrons W.	simple-leaved	£ O or	■ Ap	Trinidad	1818. S r.m.
24284 acadis Mart.	stemless	IQ or	■ Ap	Brazil	182a S r.m. Mart. palm. ic.
2*285 macrdstachys Afari.	long-spiked	£ • or	■ Ap	Brazil	1823. S r.m. Mart. palm. ic.
278a SMrLAX L.	SMILAZ. {Smile, a scraper j stems rough from prickles.) SmilcLcete. 40.—70.				
I. ACULEA*T.E. — Stem prickly.					
i. ANGULA'YE. — Stem angular.'					
24286 Jispera L.	rough, Bindweed^	un 8 s	W.G.	S. Europe	1648. Sk.s.p Schk.han.a328
2auricuftta	each-leaved	un 8 s	W.G.	S. Europe	1648. Sk.s.p Pluk.al. 110.3
24237 mgra. W.	b*ck.berried	un 8 s	Y.G.	Spain	1817. Sk.s.p
***288 catalonica Pair.	Catalonian	un 8 s	Y.G.	Catalonia	1817. Sk.s.p
**289 mauritanica Pair.	Mauritanian	un 8 au.o	Y.G.	Mauritania	1820. Sk.s.p
21290 cxcSUai.	tall	IL-Jun 12 au.s	W.G.	Syria	1739. Sk.s.p Bux. c. 1.27
*4291 horrida Desf.	horrid	un 8 au.s	Y.G.	N. Amer.	1820. Sk.s.p
24292 zeylanica L.	Ceylon	un 10	W.G.	E. Indies	1778. Sk pl Ru. am. 5.161
qudrangulkriskrisMM.	squaw-stalked	± un 6 jn.jl	W.G.	N. Amer.	1812. Sk.s.p Den.br. 109
ingifolia Rie.	long-leaved	un 10 myjl	G.w.	Cayenne	1820. C pl
Sarsaparilla U	Sarsaparilla	m 4 jl.au	W.G.	N. Amer.	1664. Sk.s.p Den. br. III
Wat*Sni Swt	Watson's	m 4 jl.au	W.G.	N. Amer.	1811. Sk.s.l Den.br. 110
[ongifolia Wat.					
b 2616 IsS>r.	Brazil	un 6 jls	Y.G.	Brazil	1820. Sk.s.p
S hastata JF.	hastate	un ... jl.au	Y.o.	Carolina	1820. Sk.s.p
«W9 lanceolata Walt.	s*each-leaved	un 5 my.jn	G.w.	N. Amer.	1785. Sk.s.p Pluk. al. 111. 8
					Cat. car. 2.84
ii. TERETICAUL'LES. — Stems terete.					
24500 CMna L.	China	• m 6 m	W.G.	China	1759. Sk.s.p Ksm.782
SI? I n^n W.	acuminate	& • un 10 jn.au	Y.G.	W. Indies	1824. Sk.s.p Pl. ic 83
Pa6ca	glaucous-leaved	un 2 my.jl	G.w.	N. Amer.	1811. Skp.1 Bot mag. 1846
wW.Jr.	southern	un 6	W.G.	N. S. W.	1815. Sk.s.p
Sundibbia L.	round-leaved	un 6 jl.au	W.G.	N. Amer.	1760. Sk.s.p
SurimMa	Laurel-leaved	un 5 jl	W.G.	N. Amer.	1739. Sk.s.p Cat. car. 1.1
4306 siphilitica Ar.	siphilitic	in 10	Y.G.	Trinidad	1818. Sk.s.p
24307 tamnooides L.	Tamnus-like	un 6 jn.jl	W.G.	N. Amer.	1739. Sk.s.p Cat.car. 1.52
24308 cad&caL.	deciduous	un 6 jn.jl	W.G.	N. Amer.	1759. Sk.s.p
24309 havan^nsis J.c.	Havannah	10 jlau	Y.G.	Havannah	1823. Sk.s.p Jac.am. 179.102
24310 ovalifolia Rox.	oval-leaved	10 ...	Y.G.	E. Indies	1823. Sk.s.p
24311 maculate itox.	spotted	6 ...	Y.G.	Nepal	1823. Sk.s.p
II. TNERMES. — Stems unarmed.					
i. TNERMES. — Stems angular.					
24312 bona n6x L.	good night	JL un 6 jn.jl	W.o.	N. Amer.	1739. Sk.s.p Pluk. al. 111. 1
24313 prollfera Rox.	proliferous	JL un 10 !!!	E. Indias	1819. Sk.s.p	
21314 glabra-Aw.	smooth	t un 10 !!!	Y.G.	E. Indies	1820. Sk.s.p
24315 Alpini W.	Alpini's	A un 6 jn.au	Y.o.	Greece	1820. Sk.s.p
24316 latifolia R. Br.	broad-leaved	JL I un 8	G.w.	N. Holl.	179J. Sk.s.p
24317 cumanensis W.	Cumana	LC Dun 10	Y.G.	Cumana	1818. Sk.s.p
2*318 herbacea L.	herbaceous	A un 4 jl	G.	N. Amer.	1699. Sk.s.p Bot.mag. 1920
ii. TEKISTICAU'LES. — Stems terete.					
24319 peduncularis Mhl.					
24320 Psebdo-cMna L.	Bastard Chinese!	un 6 myjl	G.w.	N. Amer.	1812. Skp.1
24321 canariensis W.		un 6 my.jn	G.w.	S. Amer.	1739. Sjcp.1 SL jam. 1.143. 1
24322 glycyrhilla&wz. sweet-lvd. Bot.B.TT.	i Jun 5 jl.au	Y.a	Tcnrifle	1816. Sk.s.p	
24323 puber Mx.	downy	B un	G.w.	N. S.W.	1815. Sk.s.p
24324 rtibens Wat.	red	j un	G.w.	N. Amer.	1806. Sk.s.p
2*25 virginica Mil.	Virginian	± un	G.w.	N. Amer.	1812. Sk.s.p
					kp.1 Den.br. 108
2784. TA'MUS L.	BLACK				oKS.p
2*26 commfinis L.	common				
-i327créticaL.	Cretan	A un 5 JLJu G	ffla	739. R	Tamea. 2.-3.

- 2785 & TESTUDINA* RI A *Burc.* ELEPHANT'S FOOT. (*Testudo*, tortoise; outside of roots resembling.) *Diosc.* 2.
 24328 elephantiWM *Burc.* cam., cleph. foot_ iAI cu 8 jl.au Y C. G. H. 1774. R p.1 Bot. mag. 1347
 Tūmūs elephuntipes *Hcrit.*
- 24329 montaña *Burc.* mountain 1 tAJ cu 8 jl.au ... C. G. H. 1816. R p.1
2786. RAJSNIA L. RAJANIA. (*John Bay*, a distinguished English naturalist) *Dioscbreas.* 3.—14.
 24330 cordiata L. cordate Jeaved & El un 6 jl. G W. Indies 1783. R p.1 Pl. ic. 84.98
 24331 hastata L. hastate A El un 20 jl.au Y.o W. Indies 1822. R p.1 Pl. ic. 155.1
 24332 quinquefolia L. five-leaved JL El un 20 jl.au Y.o W. Indies 1818. K p.1 Pl. ic. 155.2
2787. DIOSCOMU^ L. YAM. (*Pedacius Dioscorides*, a Greek physician.) *Dioscbrace.* 32.-53.
- I. DIVISIFOLIA.—leaves divided.
- 24333 pentaphyla L. five-leaved A (23 cul 10 ... G £. Indies 1768. R r.ra R. mal. 7.35
 24334 triphylla L. three-leaved A (23 un 8- ... G Malabar 1820. R r.m Ru. am. 5.128
 24335 trifoliate Kth. trifolate AE Jun 4. ... Trinidad 1819. R r.m
- II. LOHATJISFOLUE.—Leaves lobed.
- 24336 quirqueloba Thun. five-lobed A IAJ un 4 ... G.w Jaijan 1822. R r.m Ksm. ic. 15
 24337 brasiliensis W. Brazilian A G2 esc 8 ... G Brazil 1823t R r.m
 24338 trifida L. trifid AE Jun 4. ... Trinidad 1818. R r.m
 24339 heteroplitilla Box. variable-leaved A LSI un 4 ... G.w E. Indies 1818. R r.m
- III. Connybump.—Leaves cordate.
- 84340 aculata L. prickly-stmd A (23 cul 10 ... G K Indies 1803. R r.m R. mal. 7.37
 24341 alata L. wing-stalked A (23 cul 15 ... G India 1739. R r.m R. mal. 7.38
 24342 bulbifera L. bulb-bearing A (23 esc 12 jl.au G E. Indies 1692. R r.m Par. Ion. 17
 24343 leonensis G. Don Sierra Leone A El un 10 jl. W S. Leone 1822. R r.m
 24344 Nummularia Lam. Moneywort A (23 un 10 — E Indies 1820. R r.m Ru. am. 5 168
 84345 crispata Box. curled A El un 5 ... W.o E. Indies 1818. R r.m
 24346 angulna Box. snake AE Jun 10 ... L. £. Indies 1803. R r.m
 24347 pulchilla Box. neat A 73. un 10 ... E Indies 1818. R r.m
 24348 atropurpurea Box. dark-purple A E Jun 10 ... D.P E. Indies 1820. R r.m
 24349 globusa Box. globe A (23 un 10 ... G.w E. Indies 1818. R r.m
 24350 fasciculata Box. fascicled A El un 10 ... G.w E. Indies 1817. R r.m
 24351 glabra Box. smooth A El un 10 ... G.w E. Indies 1803. R r.m
 24352 saffra L. com., cultivated A (23 ctt 20 au G W. Indies 1733. R r.m R. mal. 8.51
 24353 rubella Box. reddish A El un 6 ... R E. Indies 1803. R r.m
 24354 purpurca Box. purple A El un 8 ... P E. Indies 1803. R r.m
 84355 nepalensis Swt. Nepal A (23 un 6 ... G.w Nepal 1816. R r.m
 24356 pipcriftilla Hum. Pepper-leaved A El un 6 ... G.w S. Amer. 1817. R r.m
 24357 coriacea Hum. leathery A El un 8 ... G S. Amer. 1818. R r.m
 24358 schbra Hum. scabrous A (23 un 10 ... G.w Orinoco 1817. R r.m
 24359 altissima Lam. tallest A H un SO ... G Martiniq. 1821. R r.m Pl. ic. 117.1
 34360 angustifolia Lam. narrow-leaved A EJ un 10 ... G Peru 1821. R r.m
 24361 villbsa L. villous A A un 3 au R N. Amer. 1752. R s.p Jac ic. 3.626
 24362 quaternata Walt. four-leaved A A un 3 au R N. Amer. 1811. R s.p
- IV. OnLovfiiFOLIA.—Leaves oblong.
- 24363 oppositifolia L. opposite-leaved A (23 un 6 ... G E. Indies 1803. R s.p Pet. g. 31.6
 24364 cinnamomifoliaifolifft. Cinnamon-lvd* *Aior* 6 n ... G.v Rio Jan. 1827. R s.p Bot mag. 9X59
2788. CCCCULUS Bank. COCCULUS. (*COCCUS*, syst. name of cochineal; scarlet berries.) *Menisp.* 12.—50.
- I. OBLONGIKOLLI.—Leaves oblong, ovate, or oval.
- 24365 /aurifblus 2tar. Laurel-leaved fl. • or 10 ... W.o E. Indies 1816. R l.p Deless. 1.97
 24366 vUlosus Dec. villous a • or 6 ... G.Y E. Indies 1800. R l.p Plu. am. 384 ♀
 . 2 hirstitus Dec. hairy l. O or 6 ... G.Y E. Indies 1800. R Lp Plu. am. 384.7
- II. CorDiPbilii.—Leaves at base cordate.
- S4367 orbiculatus Dec. round-leaved ■ 6 ... G.Y E. Indies 1790. R Lp Pluk. al. 384.6
 24368 palmatus Dec. palmate ILCDr 10 ... W.o E. Indies 1800. R l.p As. res. 10 5.3\$
 S4369 crispus Dec. curled LD 20 ... W.o E. Indies 1822 R Lp Ru. am. 5.44.1
 94370 tomentiflous Col. woolly LD or 10 ... W.o E. Indies 1819. R Lp
 24371 incanus Col. hoary fl. O or 10 ... W.o E. Indies 1820. R Lp Lin.tr. 13.6.1
 84372 Plukenetii-Dec. Plukenet's, qfficinalt. ■ 10 ... G.Y E. Indies 1790. R Lp Plk.m.345.2
 Menispernum Coccus W.
 24373 suberbus Dec. Covk.barked JL HI or 20 ... W.o E. Indies 1800. R l.p Gae.fr. 1.70.1
 24374 cordiflbus Dec. cordate-leaved fl. ■ or 20 ... W.a E. Indies 18*0. R Lp R.maL7.21
 84375 rotundifolius Dec. round-leaved L ■ or 20 ... W.c 1820. R Lp
2789. COSCFNIUM Col. COSCINIUM. (*Koskinion*, a little sieve: cotyledons perforated.) *Menhpirmacea.* J
- 84376 fenestratum Col. windowed LCD or 10 ... YG Ceylon 1800. R Lp Gae. fr. 1.4& *
 Menispernum fenestratum Gae.
2790. TILIACO'BA Col. TILIACORA. (*Tiliakora*, its name in Bengal.) *Menispe'macea.* L
- 24377 racemosa Col. racemose ■ • or 20 ... Y E. Indies 1800. R r.m
 Menispernum polycirpon Box.
2791. MA^BA Forst MABA. (Its name in Tonga-Tabu.) *Ebenace**. 2.-9.
- 24378 Auxifbia Pers. Box-leaved * O p r U *** E. Indivc 1810. C s.p Rox. cor. 1 L
- Ferréola 6uxi flbia Box.
- 84379 /alurina B. Br. laureUccved il LJ or 3 jl.au ... N. Holl. 1824. C s.p *
2792. CHAMIELI'RIMUM W. CHAMJELIMI/M. (*Chamai*, ground, *leirion*, HyO Melanthhcc*. I
- 84380 caroliniānum W. Carolina tf A or 1 jl.au Y N. Amer. 1759. D p.l BoLmag. J
- FeratrumlētcumL. Helbnias ldeea B.M. dioica Ph. Afelanthium densum Lam.

CLASS XXIJ. ORDER 7.

DICEdA OCTA'NDRIA.

						24.—
I alba L.	white, Abele Tree	tm 40	mr.ap Ap	moi.w.		Eng. bot 1618
% TM hybridal.	SJSSi*1*	or 40	mr.ap Ap	S. Europe 1800.	C co	
SSI can<>cens An.	caneicent	or 40	mr.ap Ap	Caucasus 1816.	C co	Eng. bot. 1G19
3* «5 tremida Bl.	trembling, Americ.	tm 40	mr.ap Ap	England watpl.Sk co	N. Amer. 18 W. C co	Mic. ar. 3. 8.1
3* «5 tremida Bl.	trembling, Aspen	tm 30	... Ap	N. Amer. 18 W. C co		
	smooth	tm 50	mr.ap Ap	Britain moi.w.Skco	Eng. bot. 1909	
		tm 80	mr.ap Ap	N. Amer. 1769. G co	Mic. ar. 3.11	
	twiggy	or 40	mr.ap Ap	Canada 1820. C co		
	Maryland	or 70	mr.ap Ap	Maryland 1824. C co	Dull. ar. 184. Si	
	Greeks JF A'MU	tm 40	mr.ap Ap	Archipel 1779. C co	Eng bot 1910	
g^nlgrAL.	black	tm SO	mr.ap Ap	Britain wot.pl. C co	N. Amer. ... C co	Mic. ar. 3.10.1
g^nlgrAL.	92 AetuHfolia PA. Birch-lvd, Bl Amer.	tm 40	mr.ap Ap			
	dilated, Lombardy	tm 70	mr.ap Ap	Italy 175a C co		
	necklace-bearing	tm 70	my Ap	Canada 1772. C co	Dim. br. 102	
	grandidentata Mx.	large-toothed	tm 70	mr.ap Ap	N. Amer. 177& C co	
2 pendula Mx.	pendulous	or 40	mr.ap Ap	N. Amer. 182a C co		
Z*W angulata //, ISL	Lindl. twigfesc 1* /te<<n	tm 70	my Ap	N. Amer. ... C co		
Z*W angulata //, ISL	angular, Carolina	tm 80	mr Ap	Carolina 1738. C co	Mic.ar.3.302.12	
8 baUamifera Z.	bolsam-bearing	tm 70	ap Ap	N. Amer. 1692. C co	Mic. a.. 3.13.1	
g.^ rocawaAflea Mil.						
T^vS suavolens Pi*.	sweet-scented Y	or 70	mr.ap Ap	Russia 1825. C c		
24400 longifolia Jfc.	long-leaved	or 40	mr.ap Ap	Siberia 1826. C c		
*24401 macrophilla ZKH.	long-lvd.Off/lfrioY	tm 70	* Ap	N. Amer. 1820. C c		
24402 candidans/JJC	whitish, heart-lud \$	tm 50	mr Ap	N. Amer. 1772. C co	Cat car. 1.34	
canad^nsis Afocn.						
24403 heterop?Ua L.	various-leaved £	tm 70	ap.my Ap	N. Amer. 1765. C co	Mic.ar.3.9	
cordifolia Burg.						
2M04suplnhafo.(7.)	supine	\$	or ... mr.ap Ap	N. Amer. 1824. C co		

CLASS XXII. ORDER 8.

DIGETCIA ENNEA'NDRIA.

ItfERCUMjfxIS L. MERCURY. (<i>Mercury discovered the virtues of this plant</i>) <i>Euphorbitcece</i> . 5.-7.						
perennial k Aw I ap.my G	Britain woods. D	sl	Eng. bot 1872			
doubtful O un 1 jl.au G	Spam	1806. S	co L. f. dec. 1.8			
annual O w 1 jl.s G	Britain rub. S	co	Eng. bot. 558			
c\W\A\c-leaved tt_jun 1 my.jl G	Portugal 1802. C	co	Ven.cels12			
woolly tt_Jun 1 jU G	Spain 1640. C	co				
HYDRO'CHARIS L. FROG-BIT. (<i>Hydor, water, charts, grace j pretty.</i>) <i>Hydrocharidea</i> . I.						
frsus rime L. corn., frog's bite £ A cu i jn.jl W	Britain dit. Skco	Eng. bot 808				
TRI'PLARIS L. TRIPLARIS. (<i>Triplex, triple; parts of fructification In threes.</i>) <i>Polygbne</i> e. 1. — 2.						
American J El tm 40 ... PaY S. Amer. 1824. C r.m Aub. gui.37						

CLASS XXII. ORDER 9.

DIOTCIA DECANDRIA.

2797. CORIARIA.	CORIARIA. (<i>Corium, a hide; used in tanning.</i>) <i>Coriart</i> *. 2. — 7.					
	Myrtle-leaved *	or 6 my.au G	S Europe 1629> L co	Den.br. 103		
	twiggy ft LJ cu 3 roy.au G		N. Zeal 1823. C co	Bot mag. 2470		
24415 integifolia Jflc.	L. KIOOELARIA. (<i>Francis Kiggelar, a Dutch bot author.</i>) 2.					
	African t J or 10 my.jn W.a C. G. H. 1683. C s.l Lam.il.8SI					
	entire-leaved I LJ or 10 my.jn W.a C. G. H. 1683. C s.l Jacic.3.628					
•2799.SCHrNUSZi.	SCHINUS	(The Greek name for Pistacia ienttsca.) <i>TereMnthdcear</i> . 2. - 6.				
24416 mile U	Perwrt <i>d</i> /ijUor 12 il.au G	Peru W L r.m Mil. ic. 2.246				
24417 virgata Sui.	twiggy *i_jor 8 my.jl G	Lima 1822. C p.l				
2800. GYMNO'CLADUS Lam. GYMINOCLADUS. ((
24418 canadensis Lam.	Canadian *	or				
2801. CA'RICA L. PAPAW TREE.						
2*419 PflpiyA.	com:iOjiPapay.					
3)*20 citnformisJta.	Orange-formed					
3**21 pyriformis W.	Pear-formed	or 20 ...	Guinea	IS2J. s r.m		
		D d				

- | | | | | | | | | | | |
|--|-----------------------------|---------|----|----|------|----------|--------|---|-----|-------------------|
| 94422 cauiiflora <i>Jac.</i> . , | stem-flowering | f • or | 20 | -- | O | Caraccas | 1806. | S | r.m | Jac. sc. 3.311 |
| 24423 spinosa W.
digitata Aub. | prickly | \$ • or | 20 | -- | W.a | Guiana | 1821. | S | r.m | Aub.gui.34ti |
| 24424 microcarpa <i>Jac.</i>
2 monoca itef. | small-fruited
monoecious | ■ no r | 20 | -- | W.G | Caraccas | 1806. | S | r.m | Jac. be. a 309,10 |
| | | CD- | 20 | -- | W.a. | . . . | .1818. | S | r.m | |

CLASS XXII. ORDKR 10.

DICE^NCIA DODECA'NDRIA.

- | | | | |
|------------------------|----------------------|---|--------------------------------|
| 5802. STRATIOTES L. | WATER SOLDIER. | (<i>Strata</i> , army; sword-like leaves.) | <i>Hydrochardeae</i> . 1.—2. |
| 24^5 Saloidea i. | Aloclike | * A el 2 jn.jl W England dit. | Skip Eng.bot.379 |
| Qfai HYJENA'NCHE U.K. | HYJENA POISON. | (<i>Hyana</i> , hyaena, af>cho, to strangle; poisons.) | <i>Euphorbiacea</i> . 1. |
| 24426 Blobosa H. K. | globose | * 1_ or 8 ap.s W.u C. G. H. 1783. C l.p Lamb.ci.52.10 | |
| Toxicodendron capense | T/iun. | | |
| 2804. EUCLEA% A L. | EUCLEA. | (<i>Eukleia</i> , glory; neat evergreen foliage.) | <i>Euphorbulcear</i> . 2. |
| 24427 racembsa L. | racemose, round-lvd* | J o r 5 n.d W C. G. H. 1772. C pl Jac. fr. S. 1.5 | |
| 34428 undullita Thun. | viwe-leaved | * LJ or 5 ... W C. G. H. 1794. C pl | |
| 2805. DATI'SCA W. | DATISCA. | ' (Meaning unknown.) | <i>Batiscea</i> . 2. - 3. |
| 24429 cannabina X. | Hemp-like | ■ A or 4 jl.s Y Candia 1640. D co Alp. ex. 298 | |
| 24430 hirta L. | hairy | ■ A or 4 jn.jl G.v Pemulv. 1826. D co | |
| •2806. MENISPOEIMUM L. | MOON SEED. | (<i>Mene</i> , moon, <i>sperma</i> , seed; crescent-shaped.) | <i>Menisper</i> . 5.-7.- |
| 24431 canadense L. | Canadian | ■ or 10 ■ G.v N. Amer. 1691 11 s.p Hot. nag. JP10 | |
| 24432 dafuricum Dec. | Daurian | ■ or 10 jn.jl W.Y Dahuria 1810. K s.p Bdcicsp. Lit(x) | |
| 24433 virginicum W. | Virginian | ■ or 20 jn.jl G.Y N. Amer. 1732. R s.p Di. el. 178.219 | |
| 24434 «milacium Dec. | Smilax-like | ■ or 10 ... G.v Carolina 1776. 11 l.p Jac. ic. 3.629 | |
| 94435 Cissampelos Ph. | railacia W. | | |
| 94435 Lyra Ph. | Lyon's | JL or 10 jn.jl W.a N. Amer. 1823. R s.p | |
| 2807. ABUTTA Aub. | ABUTA. | (<i>Abutua</i> , its name in Guiana) | <i>Menispermhecc</i> . 1. — 2. |
| 24436 rufescens Aub. | rufescant | fl. O or 10 ... G.Y Guiana 182a RLp Aub. gui. 2.250 | |

CLASS XXII. ORDER 11. =

DKKCIA ICOSA'NDRIA.

- | | | | | |
|--------------------------------|-----------------------------|--|----------------------|----------------|
| 2808. <i>PETTMUS</i> Pers. | <i>PEUMUS</i> . | (<i>Peumo</i> , its Chilean name.) | <i>Urticat.</i> | 1. |
| 2M37 <i>fragrans</i> Per*. | fragrant | ■ O&a3° . . . Chile | 1824. C p.l | Feu.ch.3.6 |
| 2809. <i>GELONIUM</i> Rox. | <i>GELONIUM</i> . | (Meaning unknown^) | <i>Euphorbiacat.</i> | ■ |
| 24438 <i>bifarium</i> Rox. | S-ranked^ aiwadsHQun | 6 jn.au Ap E. Indies | 1793. C p.l | |
| 24431 <i>fasciculatum</i> Rox. | fascicled aQu n 6 | ... Ap £. Indies | 1818. C p.l | |
| 24440 <i>lanceolatum</i> Rox. | lance-leaved > Du n 6 jn.au | Ap £. Indies | 1818. C p.l | |
| 2810. <i>ROTTLETU</i> Rox. | <i>ROTTLEHA.</i> | (Rev. Dr. Rottler, a Danish missionary.) | <i>Euphorbiace</i> . | 3.—10. |
| 24441 <i>tinctoria</i> Rox. | dyer's | H un 15 ... Ap E. Indies | 1810. C p.l | Rox. cor. 2.16 |
| 24443 <i>brasiliensis</i> Spr. | Brazilian | I Qu n 5 ... Ap Brazil | 1827. C p.l | |
| 24443 <i>paniculata</i> Jac. | pamceled | I Qu n 4 ... Ap Ceylon | 1823. C p.l | |
| <i>Cruton paniculata</i> Lam. | | | | |

CLASS XXII. ORDER 12.

DIGETCIA POLYA'NDRIA.

- 2812. CLIFFO'RT*i** L. CLIFFORTIA.** (*G. Cliffbit*, of Holland, first patron of Linnaeus.) *Rosace**, I. MULTINE'BVLE.—*Leaflets solitary, at base, many nerved; stipules simple, but rather with a leaflet at bath sides, stipulate, Joined into one.*

24452 *ilicifolia* L. Ilex-leaved * *L*. *I* or ³ *my.s* G C. G. H. 1714. C p.l Di.cL31.35
24453 *cordifolia* Lam. heart-leaved * *Il*. *3* *jn.jl* G w C. G. H. 1820. C p.l
24454 *tridentata* W. three-toothed * *LJ* or ³ *my.s* G.w C. G. H. ... C p.l
24465 *ruscifolia* L. Ruscus-leaved • *UJ* or ³ *jn.jl* G.w C. G. H. 1752. C p.l H. cL.31
H. D. —Leaves at first sight solitary, one nerved *stipules bifid*, or rather leaflets *bisiplyate trifoliate*; lateral leaflets small, *dentiform*.

24456 *cuneata* H.K. wedge-foawl ft *LJ* or ³ *ap* G.w C. G. H. 1787. C p.l

8823. JUNI'PERUS L. JUNIPER. (*Jurwprus*, rough or rude, *Celt.*; stiff shrubs) *Conifer**. 24.-26.
- 94511 thurifera L. frankincense-bearing fill or 10 my.jn Ap S. Europe 1752. L s.l Her. lug. 347
- 24512 bermudiana L. Bermudas. *Cedar* tm 20 my.jn Ap Bermudas 1683. S
- 24513 chinensis L. Chinese a. *Jor* 10 my.jn Ap China 1804. L
- 24514 excelsa Bieb. tall t tm SO ... Ap Siberia 1806. L
- 9*515 Sab'ina, L. common Savin * or 1 my.jn Ap S. Europe 1548. L
- 2 variegata variegated ■ or 1 my.jn Ap Europe — L
- 34516 temanscifolia Hort. Tamansk-lvd ■ or 1 my.jn Ap S. Europe 1562. L
- 24517 cractvia Lo. C. Cracow ■ or 1 my.jn Ap 1820. L
- 24518 squamiflora D. Don scaled ■ or 1 my.jn Ap Nepal 1824. L
- 24519 prostrate Mx. prostrate ■ or 1 my.jn Ap N. Amer. — S
- rhpens Nut.* .
- 24520 daurica Pall Daurian ■ or 8 jn.au Ap Dauria 1791. L s.l Bot rep. 534
- 24521 virginiana L. Virginian, tied Cedar ± caroliniana Duroi ■ or 12 my.jn Ap N. Amer. 1800. L s.l
- 2 humiliis humble ■ or 80 my.jn Ap Canada 1820. L s.l
- S4522 canadensis Fis. Canadian ■ tm 15 my.jn Ap Britain heaths. S ,s.l Eng. bot. 1110
- 24523 communis L. common ■ or 15 my.jn Ap N. Europe ...
- 24524 suica Mil. Swedish ■ or 2 my.jn Ap Siberia S l.p Pal.ros.2.54.A.B
- 24525 nana W. dwarf ■ or 3 my.jn Ap Siberia 1800. L s.i
- 24526 bibirica Burg. Siberian ■ or 4 my.jn Ap Nepal 1817. L s.i
- 24527 recurva Ham. recurved ■ or 4 my.jn Ap Syria L 81
- 24528 drupacea Lab. drupaceous ■ or 1 my.jn Ap Britain scmo. L 81
- 24501 alpina Hort. alpine ■ or 4 my.jn Ap China 1816. h s.l
- 94530 glauca W. glaucous ■ or 15 my.jn Ap Spain 1739. C 81 Duh.ar.2.128
- 9&II Oxf cedar L. Sharp Cedar ■ or 15 my.jn Ap S. Europe 1683. C s.l Pal. ros. 2.57
- 24532 phoenicia L. Phoenician ■ or 10 my.jn Ap S. Europe 1693. L s.l Pal. ros. 2.56
- 94533 tycia L. Lycian ■ or 20 ... Ap Florida 1811. L Pluk. al. 197.4.
- 24534 barbadensis L. Barbadoes Cedar £
2898. TA'XUS L. YEW TREE. XToxon, a bow j much used in making.) *Confrere.* 3.
- 24535 baccata L. common, berried f or 20 f.ap Ap Britain m.wo. S co Eng. bot 746
- 2 humilis Irish \$ or 12 ... Ap Ireland ... S p.l
- 2 hibridica Hook. procumlynt fill or 8 f.ap Ap Europe ... S co
- 3 procumbens variegated t or 20 f.ap Ap Europe ... S co
- 4 variegata
- 94536 canadensis W. Canadian \$ or 20 f.ap Ap Canada 1800. S co
- 94537 nucifera Ktmt. nut-bearing £ LJ or 20 ... Ap China 182a S co
2824. f' PHE'DRA L. EFHEDRA. (Greek name of the herb horsetail; resembling.) *Confene.* 3. —5.
- 24538 distachya £. two-spiked fill cu 2 jn.jl Ap France 1570. L co Schk. han.3.339
- 24539 monostachya L. one-spiked fill cu 2 s.n Ap Siberia 1772. L co Den. br. 142
- 24540 altissima Desf. loftiest ft cu 24 ... Ap Barbary 1825. L co Dcsf.aU.253
2825. CISSA'MPELOS/. PABEIRA BHAVA ROOT. (*Kissos*, ivy, *ampelos*, vine.) *Menispermacea?* 6.—28.
- 24541 Pareira Lam. Pareira ■ 6 jl.au G S/Amer. 1733. C s.p Lam. il. 800
- 94542 mauritanica Thou. Mauritian * CZlor 6 — Mauritius 1824. C s.p Thorej.2,3,4
- 94543 mieroe. irpa Dec. small-fruited 6 ... Y.o W. Indies 1823. C s.p
- 24544 hirsute Ham. hairy ■ : Y.o NeNil 1819. C s.pj
- 24545 Caapeba L. Caapeba ■ jl.au G S. Amer. 173a C s.p PI. ic. C7. 2
- 94546 capensis Thun. Cape ■ v G C. G. R 1775. C p.l
2826. EXCIEC AERIAL. EXCACARIA. (*Exaeco*, to blind; juice causing loss of sight.) *Euphoibi&clear.* 3.-10.
- 94547 serrate //, K. suvr-teaved fill Qor 6 f.n W Chile 1796. C p.l
- 24548 agallocha L. Ceylon fill • or 5 f.n W E. Indies 18JO. C p.l Ru. am. 2.79.80
- 94549 glandulosa Swx. glandulous fill D or 5 f.n W Jamaica Ib21. C p.l bl jam. 2.1b.8. S
2827. ADBTLIA L. A DELIA /A, prir., delos, visible; part of fructification minute.) *EupJtorbidbear.* 3.-6.
- 94550 Bernardia K. Bernard de Jus'sicu's fill • or 6 jl.au G Jamaica 1768. C p.l
- 24551 Ricinella L. Iticinus-like fill • or 6 jn.au G.w Jamaica 1768. C p.l
- 94552 Acidoton L. minted, spiny fill CD or 3 jn.jl G.w Jamaica 1768. C p.l
- 282a LOUREPRI* Cav. LOUREIRA. (*Father John de Loureiro*, a Portug. bot. auth.) *Euphorbihcete.* 2.
- 24553 glandulosa Co* glandulous fill O or 6 Mexico 1759. C p.i Cav.ic. 5.430
- 24554 cuneifolia Cav. wedge-leaved fill • or 6 JLau W Mexico 1824. C p.l Cav.ic.5.4i9
2829. MYRISTICA L. NIITMEO. (*Muristikos*, sweet smelling; odor of fruit) *Myristice*c. 3. —16.
- 24555 moschata Th. tiucmutv. i • clt 30 ... Pa.Y K Indies 1795. C p.l Bot. mag. 2756-7
- officinalis L. aromatica Lam. tasteless J • or 30 — G.w Surinam 1812. L Pluk. al. 250.6
- 94556 fatua Swz. wax-bearing <Dor 10 Y.o Guiana — Aub. gui. 2.345
- 24557 schifra Swz. Virbla sebf fera Aub.
- i 9830. NEPE'NTHES L. PITCHER PLANT. (AV, *jtriy*, *penthos*, grief; qual. of original plant) *Cytthacee.* 2.-3.
- 94558 distillatrica L. distilling, Chinese fl. • cu 6 ap.my G.Y China 1789. C p.l Bot. mag. 2798
- 94559 Phyllophthora W. Pitcher-leaf % Q cu 6 jn.au Y.a China 1820. C p.l Bot. cab. 1017
9831. CLVY'TIA U CLIYTIA. (*Outgets Cluy*, a Dutchman, prof. bot. Leyden.) *Etiphorbitcet.* 12.—fil
- 94560 alaternoldcs L. Alaturnus-like fill |pr 2 mr.d W C. G. H. 1G92. C p.l Bot. mag. 1&21
- 24561 polygonoldcs L. Polygonum-like fill |pr 2 mr.d W C. G. H. 1790. C p.l W. h. b. 51
- 24562 raphnoldcs W. Daphne-like fill LJ pr 3 my.jn W C. G. H. 1731. C p.l W.li.b.52,
- 24563 tricodca Th. Heath-hke fill LJ pr 2 ap.jn W C. G. H. 1790. C p.l
- 24564 tenuifolia W. sclndcr-lcavcd fill i | or 3 my.jl W C. G. H. 1617. C p.l
- 24565 soliflria Jac. Poly-leaved fill LJ pr 2 ap.jn W C. G. H. 179a C p.l Jac sc 2.50
- 24566 tomentosa L. tomentose fill LJ pr 3 ap.jn W C. G. H. 1812. C p.l
- 24567 imbeacons Th. pubescent fill i | or 3 ap.jn W C. G. H. 1800. C p.l
- 245W heteroqit'lla Th. variable-leaved fill t | or 3 ap.jn W C. G. H. 1818. C p.l
- 2459 pulchella L. neat * i Jpr 2 ja.jn W C. G. II. 1739. L p.l Bot. mag. 1945
- 24570 coonna Box. hill fficOpr 3 ... E! fIndies 1807. C p.l ubx.cor.2J
- 24571 pitiful Box. spreading fill • or 3 ... W E. Indies 1812. C p.l
2832. JOLLIFY BoJ. JOLLIFU. (*Jolty*, a friend of Bojer's.) *Cucurbit&clear.* 1.
- 24572 africana Del. African * E! cu 20 jl P Zanzibar 1825. C p.l Bot. mag. 2/181
- Tellaria pedata Hook. FeuiUea pedata 8m.

CLASS XXIII. ORDER 1.

POLYGA^MIA MONG^CIA.

²⁸³³⁻ *rNGA* Plu. *INGA*. (A South American name adopted by Marcgraff.) *Leg.Mimbsea*. 25.—113.

I. VE^R& — Leaves simply pinnate; leaflets large, 2 to 9 pairs, upper ones larger; unarmed.

i. PTEROPODJS.	Common petioles evidently winged.									
1*/>IZ quassuffoKa W.	Quassia-leaved	f Hor 20	...	Pk	Brazil	1820.	■	p.i		
**^4 sitifera Dec.	bristle-bearing	f Hor 20	...	Pk	Guiana	1824.	■	p.i		
&I& *»» W.	white	I Dor 20	...	W	Cayenne	1804.	C	p.i		
Feuillee's	proud	IDor 8	...	W	Lima	1824.	C	p.i	Feu. ob. 3.19	
velutina IV.	velvety	IDor 16	...	R	Caracas	1820.	C	p.i	Jac.fr. 10	
kindred	IDor 20	...	Pk	Para	1820.	■	p.i			
spurious	IDor 30	...	Pk	Brazil	1800.	■	p.i			
common, true	I Dor SO jl.au	W	W	Cumana	1820.	C	p.i	Kth. mim. 12		
Pomegranate-lvsdl	rior 12	...	Pk	W. Indies	1730.	C	s.p	SL jam. 2.183.1		
				Brazil	1815.	C	s.p			
ii. APRDPGDB.	Common petioles scarcely winged or perfectly naked.									
Eh. margined	t D fr 20	...	Pk	S Amer.	1830.	C	s.p			
torgfmj Dec.	Bourgon's	\$ O or 20	—	Pk	Guiana	1752.	C	p.i	Aub.gui.2.358	
marginato W. Mimbsa/kgifi.lia 7-										
W. LzureLleaved	JO or 20	...	Pk	S. Amer.	1818.	C	p.i	Jac.am. 164		
r,ar Awif										
TM punctata W.	spotted	IDor 20	...	Pk	Caracas	1818.	C	p.i		
TM punctata arr.	Knotty	T,D or 20	...	Pk	Ceylon	1690.	C	s.p	Pluk.al.211.5	
sHLL roph^lla W.	long-leaved	\$ D or 20	...	Pk	Cumana	1815.	C	s.p		
?tox IP^ndens W.	splendid	f D or 20	...	Pk	Para	1824.	C	p.i		
?tox hymcnoides Desv.	Hyenia-like	\$ D or 20	...	Pk	Cayenne	1823.	C	p.i		
AlzWoldes IM. C.?										
HYMEN JSODEJL	— Leaves conjugately pinnate, scarcely bipinnate; pinna one-paired, bearing two, three, or many leaflets; spiny or unarmed.									
i. BIGEMINA. — Leaves bigeminate; leaflets often one pair in each pinna.										
microphylla W.	small-leaved	f D or 16	...	Pk	Cumana	1817.	C	p.i		
clAlcis W.	sweet	^ D or 20	...	Pk	E. Indies	1800.	C	n.i	Rox. cor. 1.99	
clAlcis W.	cat's claw	T,Q or 20	...	Pk	W. Indies	1810.	C	p.i	Jac. sc. 3.3y2	
IV.	honey-bearing	T,Q or 20	...	Pk	Arabia	1822.	C	p.i		
	fetid	J D or 20	...	Pk	AV. Indies	1816.	C	p.i	Jacsc. 3.319	
— Leaflets in three or five pairs, merely six or ten in each pinna, with the interior deficient of the inferior pair.										
IBemina W.	three-paired	I D or 20	...	Pk	W. Indies	1820.	C	p.i	Jas.cam. 17781	
pmarginata H. &B.	emarginate	I D or 20	...	P	Mexico	1825.	C	p.i	Kth. mim. 17	
broad-leaved	or 10 mrap	P	—	P	W. Indies	1768.	C	p.i	Pl.lic.9	
— Leaves conjugately pinnate; pinnae bearing two or many pairs of leaflets, allopposite.										
240PS circ>nali< W.	round-podded	ff or 10	...	P	W. Indies	1726.	C	p.i	Pl.lic.5	
purple, Soldier Wood*	purple	orr ffi mr.ap	I*	W. Indies	17A3.	C	p.i	Bot. reg. 129		
wood-fruited	wood-fruited	f Q or 60	—	Pk	£. Indies	1816.	C	p.i	Rox. cor. 1.100	
Rox.	two-paired	Q or 20	—	Pk	Malabar	1820.	C	p.i	R.mal.6.12	

III. SAMVNUJE. — Leaves duplikately pinnate.

— Pinna from two to nine; leaflets of the pinna from two to twenty; spines none.

Zygia FT. tufted J D or 20 ... Pk Jamaica 1818. C p.i Br. jam. 22.5

fairest f D or 20 ... Mexico 1822. S pi

Houston's Dec. Houston's or 10 my.au P Mexico 1729. C p.i Bot.reg.98

Houstbni Herit. Acclv Housthnl W. anomalous t D or 10 my.au R Mexico 1729. C p.i Col. h. rip. 9

Acacia grandiflora W.

*2834. V+MIMOSA. (Mimos, a mimic; mimics animal sensibility.) *Leg.Mmosea*. 21.— 7S.

U MIMOSA8. — Legumes compressed, moniliform, with the ribs at the articulations contracted; flowers red.

— Leaves conjugately pinnated; leaflets two pairs, interior pair smaller than the inferior; stamens four, rarely five.

Sensitive Plant & O.Jcu Haps Pk Brazil 1648. S l.p Bot. reg. 25

blunt-leaved B-Del 3 ... P Brazil 1816. S l.p

strigosa M-Dpr 1 jnjl Pk Trinidad 1818. S p.i

bundle-flwd n, D pr 1 Jnj Pk Cumana 1824. S p.i

uni'ci, —. Leaves conjugately or subdigitately bipinnated; partial petioles few-paired, approximate at top of common petiole; leaflets subequal j stamens four, rarely five.

viira r lively ft? GHor Ifjl.s P Jamaica 1818. S p.i SI jam. 2.182.7

chaste. Humble Plant n. D cl 1 aps. W Brazil 1638. S r.m Bot. rep. 54*

hispidish <, D pr 1 J-Jl P.I.R Trinidad 1820. S p.i

blush tt CD pr 2 .. Pa.R Balua 1H18. S p.i

many-fingered tt, D cl 1 jnjl P Brazil 1822. S r.m Kth. mim. 5

viscid a^Dpr 2. Brazil 1825. S p.i

microciphala //.* B. small-headed <- D pr 1 jnjl Pk Orinoco 1820. S p.i Kth. mim. 8

HABKA*SIA. — Legumes compressed, hispid; margins straight, not contracted at the articulations.

fil9 branched IV. branched and petioles prickly; leaves bipinnated flowers white.

H. & B. sleeping jnjl W S. Amer. 1818.

AuW-spined jnjl W VeraCruz 1713. S p.i

avperata W. roughened jnjl W W. Indies 182J. S p.i

Hrey. c. 20 Dec. leg. 63

HI. BATAUCAULON. — Legumes compressed, flat, smooth, or scarcely pubescent; ribs parallel, not contracted at the articulations, unarmed or prickly; leaves biinnate; flowers whitish or pale yellow.
 24623 c&sta L. chaste ** pr 2 jl Pa.Y f. Indies 1741. S p.l Com. h. 1.28
 24624 rubicaulis W. bramble-stalked* cl 3 jn.jl Pa. Y E. Indies 1799. S Fp Kox. cor. 2.200

IV. DIBBLE.—

S4625 abstergens Rox.	cleansing	« : pr	—	E. Indies 1826. S p.l
24626 lorruginea Rotl.	rusty	* : P*	—	E. Indies 1818. S p.l
24627 Barclayfrw Host	Barclay's	« t D pr	— 1824. S s.p
24628 latispinosa Lam.	broad-spined	ft • el	—	Madagascar 1823. S s.p

2835. SCHRA'NK/i W. SCHRANKIA. (*Francis von Paulo Schrank, a German bot*) Legum. Mim. 2. —5.
 24629 aculeata W. prickly ACScu 2 jlau Pk Vera Cruz 1733. S p.l Mil. ic. 2.182.1
 24630 uncinata W. hooked AiAlcu 2 jlau Pk N. Amer. 1789. S p.l Ven.cu. 28

2836. DESMA'NTHUS W. DBBMANTHIIUS. (*Desme, bundle, anthos, fl.; fascicles of fls.*) Leg.Mhn. 12. —19.

I. NEPTU^NIA. — Legumes oblong, 4 to 6 seeded; sterile filaments always petaloid; herbs aquatic, prostrate, often rooting; leaves sensible to the touch; pinnae two or three pairs; leaflets 10 to 12 pairs; peduncles axillary, solitary.

24631 nitens W.	floating	£ [D] un 2 jl.s	W	c. j. n. a. ^ Bot. rep. 629
24632 lacutris W.	lake	^ El pr 1jl	W	S. Amer. 1818. D s.p H. & B. 1.16
24639 plenus W.	double yellow	^ BH un 2 jl.s	Y	Vera Cruz 1733. C p.l Mil. ic 2.182.2
24634 triquetrus W.	triangular	* [Alpr 1 jl	W	E. Indies 1820. D p.l

II. DESMA'NTHEA. — Legumes linear 10-15 seeded; sterile stamens subfiliform; neutral flowers often apetalous; shrubs or subshrubs; leaves not sensible to the touch; pinnae 2-5 pairs; leaflets 12-15 pairs.

24535 diffusus W.	diffuse	t D un 3 jlau	W	W. Indies 1731. C p.l Pluk.al.307.3
24636 virgatus W.	Zong-twiggled	ft • un 3 jl.au	Y	W. Indies 1774. S p.l Bot. mag 2454
24637 stnetus Bert.	strict	* O pr 3 jl.au	W	W. Indies 1800. S p.l
24638 punctatus W.	spotted-stalked	ft • un 3 jl.au	W	Jamaica 1636. C p.l Com. h. 1.31

III. DICHRO'ST&CHYS. — Legumes linear, twisted or subfalcate; sterile filaments linear, elongated, those with the fertile anthers at top bearing a pedicellate gland; petals five, distinct or joined; shrubs with branches, often spinescent; leaves bipinnate, pubescent; pinna 5 to 10; leaflets many-pairs, linear, with glands between the pinnae, especially the lower ones; spikes of flowers oblong-cylindrical, two-colored; fertile stamens yellow; sterile white, or variously colored.

24639 cinereus W.	grey	ft • un	3 jn jl	W E. Indies 1739. C p.l Rox. cor. 2.174
24640 divergens W. en.	diverging	ft CD	ti jn jl	W Abyssinia 1816. C p.l Bruce tr. 6
24641 callitachys Dec.	beautiful-spiked	ft • pr	3 jl	R.Y Tenerifl 1824. S p.l
24642 leptostachys Dec.	slender-spiked	ft CDpr	2 jl	W Guinea 1825. S p.l

*2837. jfCA*CIA Neck. ACACIA. (*Akaxo, to sharpen; many species thorny.*) Leg. Mim. 245. —C98.

I. PHVLLOD!NEiB.—Leaves of twofoms; leaflets, especially in the adult plants, abortive; petioles dilated or filiform, changed into phyllodia; flowers yellow. Species all from New Holland.

i. CAPITAVJS. — Flowers collected into globose heads, with a solitary head on each peduncle.				
24643 afata R. Br.	vring-stalked	« i J or	6 apjl	N. Holl. 1803. C H.l.p Bot. reg. 396
24644 dolabriiformis Wnl.	hatchet-slipd-lvd*	i Jor	6 apjl	N. Holl. 1818. C S.l.D
24645 decipiens R. Br.	deceiving	■ Jor	3 mr.jn	N. Holl. 1803. C s.l.p Bot mag. 1745
24646 trapezidies Dec.	trapezium-like	* i Jor	4 mrjn	N. Holl. 1810. C s.Lp
24647 biflora R. Br.	two-flowered	H i J or	3 mr.jn	N. Holl. 1803. C s.Lp
24648 hastulata Sm.	hastulatc	* I Jor	4 apjn	N. Holl. 1824. C s.I.p
24649 nervosa Dec.	nerved	Mi Jor	4 apjn	N. Holl. 1824. C s.I.p
24050 ornithophora Swt.	bird-bearing	• I (or	6 apjn	N. Holl. 1824. C s.Lp Sw.au. 24
24651 arnata R. Br.	ittrple-lvd-axmed	ft i J or	10 apjn	N. Holl. 1803. C s.l.p Bot. mag. 1653
24152 htifrida Swt.	hybrid	My or	5 apjn	hybrid 1823. C s.l.p
24653 pendula Cun.	pendulous	ft i or	3 apjn	N. Holl. 1824. C s.l.p
24654 /jenistifolia Lk.	Genista-lvd	ft J or	3 mr.au	N. a w. 1825. C a.l.p
24655 unipcrina TV.	Jumper-leaved	ft J or	6 mrjn	N.S.W. 1790. C s.l.p Bot cab. 308
ulicifolia Wnl.				
24656 osparagoldes Cun.	Asparagus-like	ft i J or	5 mrjn	Y N. Holl. 1818. C s.l.p
24657 Brdwnti Steu.	Brown's	ft uJ or	6 mr.au	Y N. S. W. 1796. C s.l.p
acicularis R. Br.				
24658 echinata Dec.	prickly	■ i Jor	mrjl	Y N. HoH. 1824. C s.Lp
24659 pugioniformis Wnl	dai-T-shaped	ft t J or	mrjl	Y N. HoH. 1818. C s.l.p
24660 diffusa B. R.	dai-e	ftuJor	myjn	Y N.S.W. 1818. C s.Lp Bot. reg. 634
prostrata B. C.				
24661 sulcata R. Br.	furrowed-yd	■ i J or	2 my.au	Y N. Holl. 1803. C s.l.p
24662 *aligna Wnl.	Willow-leaved	■ i J or	10 mrjn	Y N. Holl. 1818. C s.l.p
24663 emarginata Wnl.	cmarginatc	■ i J or	8 mr.jn	Y N. Holl. 1824. C s.l.p
24664 mucronulata Ma.C.	mucronulate	■ i J or	4 apjn	Y N. Holl. 1824. C s.l.p
24665 strcta W.	strict	double-headed	■ i J or	2 f.my N. S.W. 1790. C s.Lp Bot rep. 53
24666 leprosa Sieb.	lepros	■ i J or	6 mr.jn	Y N. Holl. 1817. S s.l.p
24667 doAnorifolia W.	Dodouia-lvd	ft i J or	5 mr.jn	Y N. Holl. 1817. 8 s.l.p
vibcbsa Wnl.				
53468 in ul fiter via Dec	many-nerved	• L Jor	5 mrjn	Y N. Holl. 1824. C 81Q
24669 lanigra Cun.	woolly	ft i Jor	6 mr.my	Y N. Holl. 1824. C s.l.p
24670 eglandulosa Dee.	glandless	ft L Jor	6 mrjn	Y N. Holl. 1824. C s.l.p
24671 rigens Cun.	stiff	• i Jor	5 apjn	Y N. Holl. 1824. C s.l.p
24672 hispidula W.	rather hairy	ft L Jor	2 ap.my	Y N. S. W. 1794. C s.l.p Bot.cab.823
24673 lincata Cun.	lined	• i Jor	6 mr.my	Y N. Holl. 1824. C s.l.p
24674 cochlearw Wnl.	spoon-leaved	• i Jor	4 myjn	Y N. Holl. 1818. C s.l.p Lab. n.h. 2.2H
24675 coriacea Dec.	coriaceous	• i Jor	5 myjn	Y N. Holl. 1824. C s.l.p
24676 anccps Dec.	two-edged	ft L Jor	4 myjn	Y N. Holl. 182a C s.l.p
24677 Mangium W.	Mangium	• OR	—	E Indies 1820. C s.l.p Ru. am. 3.81
24678 aurifolia W.	kaurel-leaved	> dor	4 myjn	Tanna 1775. C s.l.p
24679 elongata Sicb.	elongated	> t Jor	6 apjn	Y N. Holl. 1824. C s.l.p
24680 trinervata Sicb.	three-nerved	ft L Jor	6 apjn	Y N. Holl. 1820. C s.l.p
24681 heben-phald Cun.	heavy-headed	• L Jor	6 mr.my	Y N. Holl. 1817. C alp
24682 verniciflua Cun.	varnish-flowing	LJ or	d mr.my	Y N. Holl. 1818. C s.l.p
24683 daviesia-lvda Cun.	Daviesia-lvd	ft LJ or	6 myjl	Y N. Holl. 1817. C s.l.p
24684 calamiflvia Swt.	Rcd-leavrd	• i J or	3 myjn	Y N. Holl. 1823. C s.l.p
24685 quadrilateralis Dec.	four-sided	■ L J	4 ir.yjn	Y N. Holl. 1820. € s.l.p
24686 ranaliculata Swt.	channelled	■ i Jor	4 ap.jn	Y N. Holl. 1824. C s.l.p
S4687 uncinata Lod.	hooked-leaved	■ J	4 apjn	Y N. S. W. 1819. C s.l.p Bot. reg. 859

JJ588 *texifolia Cun.* Yew-lea vell ftLjor 4 apjn Y N. Holl. 1823. C 81D Bot cab. 122J
 ■■■■■ *indulaefolia Cun.* wave-leaved ft i_J or 4 anjn Y N. Holl. 1824. S sli^s
 ■■■■■ *onfrcra Cun.* crowded ft i_J or 4 fjh Y N. Holl. 1824. C sAp

ii. HETEROUENEO-PHYLLODINEJE — *Kot known to which subdivision they belong.*

fjjgj Cyclopia Ma. C. Cyclopis-sce ft _J or 4 apjn Y N. Holl. 1824. C sl.p
 U^2 Aspera Ma. C. rough ft tj or 4 apjn Y N. Holl. 1824. C s.l.p
 !jj@ grunf.ni Ma. C. Brown's ft LJ or 4 apjn Y N. Holl. 1824. C s.Lp
 *W4 Esterhazyza Ma. C. *Prin. Esterhazyza* ftJod 4 apjn Y N. Holl. 1824. C 81.p
 3g5platyphillaS< broad-leaved ft Dor 10 myjl Y N. Holl. 1820. C e.l.p
 TM Kichardson/ Swt Richardson's «Q or 10 myjl Y N. Holl. 1822. C sAp
 *wyn-atiftifa M. Dillwynia-lvd ft i_J or 3 apjn Y N. Holl. 1824. C Bl.p
 cucophylla Sutf. white-leaved »i_J or 6 apjn Y N. Holl. 1822. C 81.p
 a ri ISM kindred «L_J or 5 apjn Y N. Holl. 1822. C Bl.p
 ■■■■■ Cun. heavy-smelling fti_J or 15 apjn Y N. Holl. 1820. S sAp

J.-WITAVO-RACEMOS* — *Flowers collected into globose heads: heads racemose along the axillary peduncles.*

■■■■■ tngonocarpa Ciin. triangular-poddedft LJ or 3 apjn Y N. Holl. 1824. C 81 p
 ■■■■■ Cup. Olf-e-leaved ft i_J or 6 apjn Y N. Holl. 1824. C sAp
 ■■■■■ Cun. Podalyria-lvd ft i_J or 6 mr.my Y N. Holl. 1824. G 81.p

■■■■■ Polygalacoccinea Cun. Polygala-lvd ft LJ or 6 mr.my Y N. Holl. 1824. G 81.p
 g705 felcata W. sickle-Zoiwri ft i_J or 6 myjn Y N. S. W. 1790. C bl.p

Sjjig falciforaiis Dec. sickle-shaped ft LJ or 6 apjn Y N. Holl. 1818. Bl.p
 ***W penninervis Sieb. pinnate-nerved ft J or 6 apjn Y N. Holl. 1824. C fl.p Bot mag. 2754

■■■■■ Presa Cun. inelanoxylon R. Br. black-wood ft i_J or 8 apjn Y V. D. I. 1808. C s.l.p Bot. mag. 1659
 heterophylla W. variable-leaved ft i_J or 5 apjn Y N. Holl. 1824. C s.l.p

wiyriffhlia W. klyrtle-leaved ft i_J or 5 apjn Y N. Holl. 1820. C s.l.p

■■■■■ puiformis Cun. clothed ft i_J or 3 f.my 1789. C sl.pSw.au. 49

■■■■■ davaricata Cun. marginatc-wt ft i_J or 6 apjl Y N. Holl. 1820. S eAp Bot. reg. 698

■■■■■ umbrida Cun. blister-shaped ft i_J or 4 apjn Y N. S. W. 1803. C 81.p

o5 k PjMtohL Dec. divaricated ft i_J or 6 mr.my Y N. Holl. 1824. C s.l.p

U^8 Wnervata D. shiny ft i_J or 6 mr.my Y N. Holl. 1824. C s.l.p

24720 lunjita Sieb. two-nerved ft i_J or 8 apjn Y N. Holl. 1824. C s.l.p

S421 brevifolia Lod. two-veined ft i_J or 10 apjn Y N. Holl. 1824. C s.l.p

«422 obtusifolia Sieb. lunate ft i_J or 2 apmy Y V. D. I. 1810. C sAp Bot. cab. 384

24723 crassiuscula Wnl. short-leaved ft i_J or 3 apjn Y N. Holl. 182a C s.l.p Bot. cab. 1235

24724 suavtolens W. blunted ft i_J or 10 apjn Y N. Holl. 1824. C 61.p

■■■■■ ambigua Sal. sweet-scented ft i_J or 10 apjn Y N. Holl. 1824. C s.l.p

■■■■■ priminae Cun. Box-leaved ft i_J or 4 fjin Y N. HoU. 1824. C 81.p

■■■■■ prminima Cun. prominent ft i_J or 1 fjin Y M. Holl. 182*. C s.l.p

21727 angustifolia Wnl. narrow-leaved ft i_J or 1 ap.my Y N. S. W. 1816. C s.l.p Bot calx 763

24248 /infolia W. Flax-leaved ft i_J or 1 my.jn Y N. S. W. 1790. C s.l.p Bot. mag. 3168

[Jig <bietina FT. Fir-like ft LJ or 1 apjn Y N. Holl. 1823. C s.l.p

1^30 rfcida Cun. reddish ft LJ or 1 mrjn Y N. Holl. 1823. C s.l.p

sublate ft LJ or 1 apjn Y N. Holl. 1824. C si.p

24732 virg. twiggy ft LJ or 1 apjn Y N. Holl. 1824. C s.l.p

IV. SPICVHE. — *Flowers disposed into cylindrical spikes.*

spur-fruited ft LJ or 6 mr.my Y N. Holl. 1824. C s.l.p

whitened ft LJ or 10 mrjn Y N. Holl. 1823. C s.l.p

thick-fruited ft i_J or 6 mr.my Y N. Holl. 1824. C s.l.p

vrhor/Leaved ft LJ or 10 mr.my Y V. D. I. T80 C sl.p

narrow ft LJ or 10 inr.iny Y N. Holl. 1780! C 81.p Bot. mag. 110

Herit. broad-leaved ft LJ or 10 nr.my Y N. Holl. 1780. C s.l.p Wen. c. 1.30

Yew-leaved ft i_J or 7 myjn Y N. Holl. 1824. C s.l.p

Sharp Cedar ft i_J or 7 apjn Y N. Holl. 1824. C sl.pSw.au.6

Ruscus-Icaved ft i_J or 10 mr.my Y N. Holl. 1824. C sl.p

linear ft i_J or 3 myjn Y N. S. W. 1820. C Bl.p Bot mag. 25R

mucromate ft i_J or 4 myjn Y N. Holl. 1818. C s.l.p Bot. mag. 2747

longest-tourerf ft O or 4 myjn Y N. S. W. 1819. C 81.p Bot reg. G80

bundle-flwd ft i_J or 6 myjn Y N. S. W. 1798. C s.l.pVen.ch.13

equal-woolley ft i_J or 6 apjn Y N. Holl. 1822. C s.l.p

l-side-woolly-urf ft • or 10 myjl Y N. Holl. 1818. C s.l.p

long-leaved ft i_J or 10 mr.my Y N. S. W. 1792. C s.l.p Bot. mag. 2166

spear-wood ft J or 10 mr.jn Y N. Holl. 1823. C s.l.p

interwoven ft i_J or 6 apjn Y N. Holl. 1824. C s.l.p

blunt-leaved ft i_J or 6 apjn Y N. Holl. 1823. C s.l.p

cinerascent ftJor 10 apjn Y N. Holl. 1824. C s.l.p

glaucescent «l_J or 3 fjin Y N. S. W. 1790. C s.l.p

Sophora/HM&critft i_J or 10 apjn Y V.D.I. 1805. C s.l.p Lab. n. h. 2.237

H. — *Leaves conjugately pinnate, leaflets bipinnate. An artificial series of heterogeneous species.*

Q&K r^6h. camelopard's i • or 40 — — C. G. H. 1816. C pl

g £* strombulifera W. Miiral-podded ft LJ or 8 — — Peru 1825. C s.l.p

S 2 ^ TxeRilla Lk. Rue-leaved ft Dor 4 ... Y 1810. C s.l.p

5 S &^ gophilla Hofm. few-leaved ft CD or 4 ... Y 1817. C s.l.p

tirr Summifera W. gum-bearing ft i_J or 30 — ... Guinea 1823. C s.l.p

^758 coronillifflia Desf. Coronilla-lvd ft • or 10 N. Africa 1817. C s.l.p

MimHa girdifc Brou. pilose f O or 30 — W Jamaica 1800. C s.l.p

4759 pilosa Bert. netted ft uJ or 10 — W 1816. C

24760 reticulata W. en. pretty ft>Jor 4 apjl Y 1803. C

24761 milchilla H. K. hairiest ftJor 3 ... WV 1800. C

24762 hispidissima Dec. detaining fti_jor 3 apjn Y 1824. C

24783 dentincis Burc. green-branched ft i_J or 4 ... y 1824. C

*J764 viridirhmis Burc. Cassia-like J • or 20 ... W C G. H. 1816. C

4765 cassioloides W. en. 1820. C

POLYGAMIA MONIECIA.

CLASS XXIII.

24766 dptera W.	two-winged	f • or 20	W	S. Amor.	1818.	C	s.l.p
24767 robusta Bure.	robust	ft uJ or 6	...	C. G. H.	1816.	C	s.l.p
24168 litakunensis Bure.	Litakun	ft i_J or 6	...	Litakun	1816.	C	s.l.p

III. SPICIFLURÆ.—Leaves bipinnate, many-paired flowers spiked.

i. INE'RME'S.—Unarmed.

24769 lophanthia W.	crest-flowered	ft L-J or 6	my.jl Y	N. Holl.	1803.	C	s.l.p
24770 arenosa W.	sand	*Q ⁰ * I	...	Caracas	1818.	C	s.l.p
24771 lophanthoides Zucc.	Lophantha-like	ft CD or 20	...	Jamaica	1820.	C	s.l.p
24772 guianensis TV	Guiana	D or «	W	Cayenne	1803.	C	s.l.p
24773 llicida Box.	shining	i U or D	...	E. Indies	1820.	C	s.l.p
24774 Sertsa Rox.	Shireesh	J or 20	...	E. Indies	1822.	C	s.l.p
24775 alatafistulosus Korth.	Axelpapakiped	L or 45	W	Mauritius	1824.	C	s.l.p
24777 elephantortifiza Bure.	elephant root	Y uj or 30	...	Acapulca	1825	C	s.l.p
24778 Rohriæ Vahl	Bohr ³	1 CD or 30	W	C. G. H.	1818.	C	s.l.p
24779 scleróxylus Tus.	hard-wood	■ ■ ■	...	W. Indies	1822.	C	s.l.p
24780 pulcherrima W.	fairest	H~lor	...	Brazil	1823.	C	s.l.p
24781 Wallichrffia Dec.	Wallich's	Q or	...	E. Indies	1820.	C	s.l.p

ii. ACULEAVA.—Prickly or spiny.

24782 S&ndra Dec.	Sundra	- t • or 15	— Y	E. Indies	1789.	C	s.l.p	
24783 ferruginea Dec.	rusty	ft Q or	...	E. Indies	1818.	C	s.l.p	
24784 Câtechu W.	Catechu	t • m 40	—	Pa.Y	E. Indies	1790.	C	p.l Rox. cor. 2.175
24785 viridiflora Kth.	green-flowered	J • or 20	—	W.o	S. Amer.	1823.	C	s.l.p Kth. mim. 25
24786 ckfra W.	Caffraria	Q L J or 12	—	Y.w	C. G. H.	1800.	C	s.l.p
24787 cornigera W.	hom-bearing	• or 15	—	Pa.Y	S. Amer.	1692.	C	fl.p.l Pluk.aL 3.122.1
24788 moncantha W.	one-spined	• or 20	—	W	Brazil	1818.	C	s.l.p
24789 Spin* Balb.	Spin's	J • or 15	—	R.Y	E. Indies	1824.	C	s.l.p Col. h. rip. ap. 5
24790 latrñnum W.	plunderers ¹	t CD or 20	—	W	Africa	1823.	C	s.l.p
24791 Senegal W.	Senegal	J • or 30	—	W	Africa	1823.	C	s.l.p Alp. eg. 15

IV. GLOBIFLORA.—Leaves bipinnate; flowers collected into globose heads.

i. AcIRLEisn'puuE.—Prickles all stipular, straight; legumes unarmed; stamens twenty or more.

24792 ebirnea W.	ivory-thorned	ft D or	—	E. Indies	1792.	C	s.p.l Rox. cor. 2.199	
24793 tortuosa W.	twisted	ft L or	—	Jamaica	1824.	C	s.l.p	
24794 Burmanimí Dec.	Burman's	*O or	Y	Ceylon	1818.	C	s.l.p	
Mimosa tortuosa Brm.								
24795 h6rrida W.	horrid	ft D or	6	Africa	1823.	C	s.pl Pluk.al. 121.4	
24796 vera W.	tmc Egyptian Thorn	t • or 12	jl	Egypt	1596.	C	s.p.l Pluk. al. 123.1	
24797 arabica W.	Gum Aralic tree	J • ec 20	—	E. Indies	1820.	C	s.p.l Pluk. al. 251.1	
24798 hebæclata Dec.	young-branched	J J or 40	—	C. G. H.	1816.	C	s.l.p	
stolonf cfrca Bure.								
24799 Icucophls'a W.	white	f • or 12	—	Pa.Y	E. Indies	1812.	C	s.p.l Rox. cor. 2.15
24800 mauroceana Dec.	Moorish	ft i_J or 6	—	...	Morocco	182-	C	s.l.p
24801 farnesina W.	Farnesian	t • fra 15	jn.au	Y	St.Domin.	1656.	C	s.p.l
94802 haematôxylo W.	bloody-wooded	J Q or CO	...	C. G. H.	181a	C	s.p.l	
24803 tomentosa TV.	woolly	ft Q or 20	—	E. Indies	1816.	C	s.l.p	
34804 india Desv.	Indian	ft CD or 10	—	E. Indies	1800.	C	s.l.p	
24805 macracanthides Bert.	long-spined	f • or 20	—	Jamaica	1820.	C	s.l.p	
24806 flexuosa H. #5.	flexuose	ftQ or 6	...	Cumana	1824.	C	p.l	

ii. AcANTiocoRPji.—Prickles stipular in pairs, also often petiolar, and along the ribs of the legumes ; stamens ten.

24807 acanthocarpa W.	spiny-nodded	ft • or 10	...	Pa.R	N. Spain	1822.	C	s.l.p
24808 acantholoba Hum.	spiny-lobed	ft L	—	W	S. Amer.	1823.	C	s.l.p

ill. AcANHOCLA'DIE.—Prickles on branches scattered ; stems not climbing.

24809 tamarindifb W.	Tamarind-lvd	ft I or	4	W	W. Indies	1774.	C	s.p.Uac. sc. 3.396
24810 conclina Dec.	neat	f • or 20	—	W	E. Indies	1823.	C	s.l.p
24811 Irtsia W.	Ir.tsia	f I or SO	—	Y.w	E. Indies	1778.	C	s.p.l R. mai. 6.4
24812 pennata W.	fc & thc T-leaved	J • or 20	—	E. Indies	1773.	C	s.p.l Bur. zey. 2.1	
24813 CJEP'sa IV.	grey	f • or £0	—	Y	E. Indies	1773.	C	s.p.l Pluk. 330.1
24814 centrophyila Dec.	spur-leaved	J • or SO	—	W	Jamaica	1818.	C	s.l.p
24815 Courantina Dec.	Courant's	f J or SO	—	Canaries	1818.	C	s.l.p Pluk. ic. 330.3	
24816 arrophul D. Don	Arrophul	ft L or 20	—	Nepal	1818.	C	s.l.p	

iv. SCANDENTES.—Prickles scattered ; stems climbing.

24817 scindens W. en.	climbing	fl • or 10	...	P	India	1780.	C	s.l.p R. mal 8.52.34
24818 Guilandina Dea	Guilandina-Mr	f • or 10	...	Y	Cayenne	1820.	C	s.l.p
24819 sarmcntbsa Desv.	twiggy	fl • or 10.	...	—	—	—	C	s.l.p

v. TRICHO'DEA.—Unarmed anthers hairy ; stigma penicilliform.

24820 trichodes IK	hairy	ftQ or	10	...	Pa.Y	Peru	1818.	C	s.l.p Jac. sc. 394
--------------------	-------	--------	----	-----	------	------	-------	---	--------------------

vi. JULIBH'SSHTC.—Unarmed ; anthers smooth ; stigma simple.

24821 nligricans R. Br.	blackish	ft L or	8	my.jl	Y	N. Holl.	1803.	C	s.l.p Bot mag. 2188
24822 formosa Kth.	beautiful	• D or	10	...	W	Mexico	1825.	C	s.l.p Kth. mim. 32
24823 ciliflta R. Br.	ciliata	ft L J or 8	mr.jn	Y	N. Holl.	1803.	C	s.l.p	
strigosa									
24824 Mbbeck W.	Loebach	t • or 20	mr.jn	Pk	Egypt	1893.	C	s.l.p Pluk. m. 331.1	
24825 v^ga IV.	common	• or 40	...	W	Brazil	1818.	C	s.l.p Pis. br. 80.2	
24826 odoratissima TV.	most fragrant	f • fia 40	—	E. Indies	1790.	C	s.p.l Rox. cor. 2. 120		
24827 procOra IV.	tall	f • or 60	—	Pa.Y	E. Indies	1816.	C	s.l.p Rox. cor. 2.21	
24828 speciosa W.	showy	» • or 10	aus.s	P	E. Indies	1743.	C	s.l.p Jac. ic. 1.198	
24829 glauca JV.	glaucous	i J or 5	jn.au	W	America	1690.	C	s.l.p Cat car. 2.4S	
24830 leucoccephala Pers.	white-headed	ft • or 5	jn.au	W	S. Amer.	1823.	C	s.l.p	
24831 lomatocarpa Dec.	fringed-podded	ft • or 30	...	Pa.Y	E. Indies	1824.	C	s.l.p R. mal. G 5	
Mimb ^a margin'ta Lam.									
24832 latisiliqua W.	broad-podded	ft I or 10	mr.jn	Pk	W. Indies	1777.	C	i.l.p PLic.S.6	
24833 portoricensis W.	Porto Rico	• D or 12	jn.au	W	S.Amer.	1824.	C	s.l.p Jac. ic. 633	
24J34 caracaskna W.	Caraccas	•• or 12	...	Caraccas	1817.	C	s.l.p Jac. ic. 632		
24835 quadranguUris Lk.	quadrangular	•• or 4	jls"	W	...	1825.	C	s.l.p	
124836 Lambertina D.Don	Lambert's	• ■ ■ 6	my.jn	P	Mexico	1818.	C	s.l.p Bot. reg. 721	

9*£I tottagôna W. ^38 discofor JT.	four-angled	H	A or	26	jn.jl	W	Caraccas	1820.	C alp	
\$43 angulcea Desv.	two-colored	w	flor	10	mr.jn	Y	N. S.W.	1788.	C s.l.p Bot. rep. 2:6	
SSJ? P ^{uf} escens R. Br.	angular	w	flor	26	jn.jl	Y	N. Holl.	1820.	C s.l.p	
SSJ! Ilk** JV.	dovny-stemmu	w	flor	10	mr.jn	Y	N. S. W.	1790.	C s.l.p Bot map. 1213	
SSJ! Ilk** JV.	Fern-leaved	1	flor	20	...	—	Mexico	1825.	C B.l.p Cav. ic. 1.78	
SSJ! Ilk** JV.	villous	w	flor	6	...	W*	Jamaica	1800.	C s.l.p	
SSJ! Ilk** JV.	leafy	w	flor	30	...	W	E. Indies	1816.	C s.l.p	
g.Jt stipulata MacDec.	/irge-stipuled	w	flor	20	...	W	Bengal	1800.	C s.l.p	
»43 divaricata W.	divaricate	w	flor	20	...	W	E. Indies	1820.	C s.pl Jac. sc. 395	
g.44 carbreae Baff.	tree	w	flor	40	...	Pk	Jamaica	1768.	C s.pl Pluk al. 6. 2512	
g.45 amara W.	bitter	w	flor	30	...	W	E. Indies	1816.	C s.l.p Rox. cor. 2.122	
g.46 polyphylla JD.	Silk tree	w	flor	20	au	W	Levant	1745.	C s.pl Scop. d. 1.8	
5*?? decurrens W.	many-leaved	m	•	20	...	W	W. Indies	1824.	C s.l.p	
5*80 mollis B. R.	decurrent ,	m	i	lor	6	my.jl	Y	N. S. W.	1790.	C s.l.p Bot. reg. 371
oiom »»Sima IT.	soft	m	•	6	jl.au	Y	N. Holl.	1810.	C s.l.p Sw.au. 12	
4452 dealbata L*	whitened	m	flor	4	—	Y	1824.	C s.l.p	
S*J pmsmatica /I^HJ.	prismatic	w	flor	6	—	Y	1818.	C s.l.p	
6\$1 intorta Dec.	twisted	w	flor	20	...	W	Brazil	18*5.	C s.l.p	
**has esculenta Drc.	eatable	w	flor	20	...	W	N. Spain	1825.	C s.l.p Fl. mex. ic. in.	

~~Species not sufficiently known~~ *sleaves bipinnated, prickly, or unarmed.*

<i>Jacaranda</i> Lo. C.	tall dusky	t • or • or	60 15	E. Indies	1820. C	s.l.p
<i>ra Rox.</i>	Kalkora	f • or	45	S. Amer.?	1825	C s.l.p
<i>ea Rox.</i>	shrubby	4 • or	4	E. Indies	1818.	C s.l.p
<i>idna Rox.</i>	Smith's	f • or	45	E. Indies	1820.	C s.l.p
<i>femcordata Rox.</i>	half-cordate	f • or	40	E. Indies	1818.	C s.l.p
	Ross's	f • or	40	E. Indies	1820.	C s.l.p
	<i>gum copal</i>	J • or	20		1822.	C 8.l. p
<i>ijit? Peregrina W.</i>	foreign, <i>whitc-fiwd*</i>	• or	8	W	S. Amer.	1825. C sl.p
<i>WW? an Sustifolia Wnl.</i>	narrow-leaved	f • or	20	W	W. Indies	1820. C s.l.p
<i>W.</i>	small-leaved	■	10		Caracas	1826. C s.l.p
	grateful	■	10		Brazil	1820. C s.l.p
	charming	■	6	Pk	S. Amer.	1816. C s.l.p
	eatable	■	20		S. Amer.	1820. C s.l.p
24-25 <i>bira chyacantha W. en.</i>	short-spined	■	4		S. Amer.	1824. C p.l
♦?^? <i>Peruviana W. en.</i>	Peruvian	■	6		Peru	1820. C p.l
♦S-1 ³ <i>'ei>toph'lla Dec.</i>	fine-leaved	■	6		S. Amer.	1824. C s.l.p
0T244 <i>vir^scens Dec.</i>	virescent	■	6		S. Amer.	1820. C s.l.p
^ < 5 /entisciflia Desf.	Lcntiscus-lvd	■	6		Mexico	1824. C s.l.p
^ 6 <i>patula Hum.</i>	spreading	■	6		S. Amer.	1818. C s.l.p
2u ⁸ ₂ <i>guaiquilensis Desf.</i>	Guaiquil	■	6		Guiaquil	1818. C alp
0u ⁷ ₂ <i>urachyacantha Hum.</i>	short-spined	■	6		S. Amer.	1820. C s.l.p
Sisam J ¹⁰ <i>dacantha Delt.</i>	rcd-spined	■	6		C.G. <	1816. C s.l.p
2 ¹¹ <i>petracanthia Burc.</i>	variable-spined	■	6		Jamaica	1820. C s.l.p
> <i>ang⁹ <ona Bert.</i>	Bancroft's	■	6		S. Amer.	1822. C s.p.l
	ciliated	■	6		E. Indies	1818. C s.p.l
	bushy	■	6		E. Indies	1818. C s.p.l
<i>Lpncordina Rox.</i>	Concord's	■	6		S. Amer.	1800. C s.p.l PI. ic.8
<i>stratina W.</i>	Ceratonia	■	6		E. Indies	1822. C s.p.l
<i>dunculata Rox.</i>	pedunculate	■	6		Brazil	1825. C B.n.l
<i>aliensis Spr.</i>	Brazilian	■	6			

DA. (Its name in Malabar.)

			single-spiked	L eo r 20	... W		
21SRn						Malabar	1800. C Lp
	vf	ftha	Dec 7	Pursatha A.	• or 20	-- W	E. Indies 1780. C Lp
24890	??&**	seandens	Rox.	"Schanthfera Dec.	Adenanthera-/*A.	• or 20	— W
24891				chanthfera seandens Forst.	Gigalobium	L O o r 20	S. Sea Isl. 1817. C Lp
	if Ra	QWm	Dec.	Sii^ascandens Swz.		— W	W. Indies 1819* C Lp
				"ystachya Dec.	many-spiked	a o r 26	W. Indies 1816. C Lp
				Mimdsapoly.achvaZ.			

(*Vere*, truly, *ater*, black; root)

		white	Vet., Nutt., etc.	black,	root,	Indicates, &c.
24895	<i>iæ H. K.</i> <i>Wⁿ Rn in.</i>	Am 5 jn.au	L.Y.	Europe	1548.	Spk.l Fl.dan. 1120
24896	<i>J. L.</i>	5 jlau	G	N. Amer.	1742.	Sk.p.l 'Böt.' mag. 1098
24897	<i>Jarvill. Mx.</i>	3 jn.jl	D.P.	Siberia	1596.	D.p.l Bot. mag. 963
"	<i>sngustifolium Ph.</i>	2 jn.jl	G	Carolina	1809.	D.p.l
	<i>narrow-leaved</i>	2 jn.jl	G	N. Amer.	1823.	D.p.l

Whi, meadow, *an/hos*, flower.) *Melanthoncete*.

Br N. Amer. 1768. D Lp

Hebniæ virginalia B. M.
N Amer. 1817 D. Jn

N. Amer. 1817. D. Lp.

inj. acembsum Mr. N. Amer. 1822. D Lp

pogon, beard: tufts of hairs on flower

Jl.au Ap N. Ainer. IS₁₈₁₀ 18 D co

A P Egypt F^{1810.}
A p Egypt F^{1824.} D co &c

— Ap Japan 1816. D co

1 au Ap S. Europe 1768 D. co

S. Europe 1768. D co
S. Europe 1818. D co

jau Ap S. Europe 1802. D co
il au Ap N. Spain 1820 B

Jl.au Ap N. Spain 1820. D co
Jl.au A|z Jamaica 1816. D co

2 au Ap Jamaica 1810. D co
Mexico 1825 D co

2842. RIPPIDIUM *Trin.* RIPIDIUM. {*Rhipidion*, little fan; application not evident) Graminea. 2.—
- 24911 Ravenna? *Trin.* Ravenna Jl A or 2 jLau Ap & Europe 1810. D co Host gr. 3.1
Andropogon Ravenna? *Hort. Saccharum* Ravenna W. Eriantlius Ravenna? Beauv.
- 24912 strictum *Trin.* strict Jl A or 2 jLau Ap Hungary 1802. D co Host gr.2.2
Andropogon strlctus *Host*, *Saccharum* adpressum *Kit.*
2843. POLLINIif Spr. FOLLINIA. {Cyrus Polling professor at Verona.) Graminea. 3.—16.
- 24913 distachya Spr. two-rowed Jl A un jlJ.lau Ap & Europe 1805. D co flgr.69
Andropogon distachyos W.
- 24914 striata Spr. striated * E J un 2 au Ap E. Indies 1793. D co
Andropogon stri&tus W.
- 24915 undata Spr. waved JllbCSlun 2 jLau* Ap Mauritius 1823. D e o Jac.ic.3.361
Andropogon undatus *Jac.*
2844. CYMBOPO'GON Spr. CYMBOPCKJOV. (*Kymhe, ho&t, pogon*, beard: valves of calyx.) Graminea. 1-6.
- 24916 Schoen&thus Spr. Lemon-grass jjii(ZQfra 2 ... Ap E. Indies 1786. D i p Ru. am. 5.72.2
Andropogon Schflenanthus L.
2845. HETEROPCTGON Beauv. HETEROPOOON. (*Heteros*, variable, *pogon*, beard; awns.) Graminea. 2.
- 24917 Allibnii Beauv. Allioni's * A un 2 jLau Ap S. Europe 1816. D e o ALped.91.4
Andropogon Allibnii W.
- 24918 contortus Beauv. twisted M El un 2 jla Ap E. Indies 1779. D co Schk.han.3.342.II
Andropopon contortus W.
2846. APLITDA L. APLUDA. (Apluda, chaff; involucres.) Graminea. L—5.
- 24919 aristata L. awned Jk El un 1 jlau Ap E. Indies 1820. D p.l Schrb. gr. 2.42
Andropogon involucratus *Kion*
2847. AN ATHETRUM Beauv. ANATH. (A, prlv, other, awn; valves of calyx awnless.) Gram. 3. —10.
- 24920 rouricatum Beauv. muricate M E) un 2 ... Ap E. Indies 1816. D co
Andropogon muricatus Retz. nquarensis W.
- 24921 virginicum R. & S. Virginian M A un 1 jlau Ap Virginia 1819. D e o SI. jam. 1.68.2
24922 miticum R. % S. awnless Jl iAJ un } jl.s Ap C. G. H. 1794. D co
Andropogon miticum L.
2848. ANTHISTFRIA TV. ANTHISTIRIA. (*Anthistemi*, to resist; harshness of stubble.) Graminea. 2.—13.
- 24923 ciliata Retz. ciliated M ED un 1 au.s Ap Jamaica 1826. S co Cav. ic. 5.45!)
- Andropogon quadrivalvis L.
- 24924 australis li. Br. southern M AJ un 1 au.s Ap N. Holl. 1818. D co
2849. CHLCTRIS Swx. CHLORIS. (Chloros, green; color of herbage.) Graminece. 15.—27.
- 24925 ciliata Swx. ciliated JtfTHpr | jl.s Ap Jamaica 1779. D co
- 24926 pallida W. pale - - Mi O pr * au s Ap S. France 1816. S co
- 24927 polydactyla Swx. many-fingered m O pr Jl.lau Ap Jamaica 1810. S co Jac. gram. 1.9
- 24928 elegans Hum. elegant • Jl rm R 1 jlau Ap S. Araer. 1818. S co H.&B. n. 1.49
- 24929 compressa Dec. compressed Mi O pr 1 jlau Ap & France 1820. S co
- 24930 barbata Swx. bearded PR 1 jn.jl Ap E. Indies 1777. S co Jac. gram. 1.8
- 24931 Roxburghjtina R. & S. Roxburgh's PR i jlau Ap E. Indies 1820. S co
- 24932 inflata lk. inflated i Apt 1 jlau Ap California 1826. S co
- 24933 fasciculata R.S/S. fascicled M O-pr | jl.au Ap Brazil 1827. S co
- 24934 radiata Sum. radiated PR } au.s Ap W.Indies 1739. S co M.h.8.3.15
- 24935 ventribsa R. Br. ventricose PR } jl.au Ap N. Holl. 1820. S co
- 24936 truncata R. Br. truncate IAI PR J jlau Ap N. Holl. 1818. S co
- 24937 dolichostachya Lag. long-spiked M O pr fjl.au Ap PhiIsl. 1822. S co
- 24938 retusa Lag. retuse m O P* % jlau Ap B. Ayres 1824. S co
- 24939 grettae Durand slender Mi O P' fjl.au Ap 1824. S co
- Durandsna R. & S.
2850. ATHEROPO'GON Mhl. ATHEROPOGON. (Ather, awn, *pogon*, beard; awns bearded.) Gram. 1.—17.
- 24940 apludoloides W. • Apluda-like it ^ pr f au Ap & Europe 1768. D e o Schrb. gr. 2. S3
Cnloris curtipendula Mx. *Dintbra* curtipendula. ■■■
2851. EU^STACHYS Desp. EUSTACHTS. (EU, well, *stachys*, a spike; large.) GranulhecB. 1.
- 24&41 petraea Desv. rock jUkQSpr jt* jl.au Ap Jamaica 1779. D co VahUy.2.27
Chloris petraea Thun. ilgrostis compianate //, A.
2852. S&RGHUM W. en. SORGHUM. {*Sorgho*, its Indian name, according to Bauhin.) Graminea. 10—12.
- 24942 bicolor W.en. two-colored ■ Θ clt 3 jl Ap Persia 1731. S co Ac.hel.8.4.4
- 24943 cernuum W. drooping Mi O clt 3 jl Ap 1810. S co Host gr. 4.3
- 24944 aven&ceum Beauv. Avenaa-Uke O Clt 3 jl.au Ap C. G. H. 1816. S co /fulcus avenaceus Thun.
- 24945 vulgare Pert, common IndianMilUm O clt 4 jl Ap India 1596. S co M. ache. 8.4.3
- 24946 n^grum R. & S. black Mi O clt 4 jl Ap India? 1816. S co Aid. sag. 1.5.1
- 24947 rrbens W. en. red-seeded Mi O clt 3 jl Ap Africa 1817. S co
- 24948 saccharatum Pen. sugared Mi O clt 6 jl.au Ap India 1759. S co Host gr. 4.4
- 24949 Califbrum R. S. S. Califves* Jir O clt 3 jl.au Ap C. G. H. 1816. S co Ard. sag. 1.1.1
- Ardu'int Jac.
- 24950 halepense Pert. Aleppo M A un ■ jl.au Ap Syria 1691. D co FLgr.1.68
- 24951 elongatum R. S. S. elongated Jb AJ un ■ jl.au Ap N. Holl. 1823. D co
2853. CHRYSOPCTGON Trin. CHRYBOPOGON. (Chrysos, gold, *pogon*, a beard; yellow.) Graminea. 1*
- 24952 grfllus Trin. cricket Mi A un jn.jl Ap S. Europe 1791. D e o Fl.gr. 1.67
Andropogon grf llus L. Holcus gry'llus R. Br. Pollinia gryllus Spr.
2854. HCKLCUS L. SOFT GRASS. (Helko, to extract: original plant supposed to extract thorns.) Gram. 2.—*
- 24953 m6llis L. soft, creeping JH A ag 2jl.au Ap Britain cornfi. D h.l Eng.botHTO
- 24954 lanatus L. woolly, meadowM A ag 3 jn.jl Ap Britain me.pa. D h.l Eng.bot.ll6y
2855. ARRHENATHETRUM Beauv. ARR. (Arrhen, male, other, point; male spikes awned.) Gram. 2.
- 24955 avehaeum Beauv. A vena-like M & ag 5 jn.jl Ap Britain me.pa. D co Eng. bot. 813
•Holcus avenaceus Sco. Avtna, clatior L.
- 2m&ticum awnless M A ^8 * jn.jl Ap Scotland hills. D co
- 8*96 tmjbtbaum Swt. bulbous Mi A ag 3 jn.jl Ap Germany ... D co
//ulcusbulbosa* Schr. ilvna bulbosa W.

2856. JSCHA?MUM L.	IACHISMUM	(<i>Tscho</i> , to stop, <i>haima</i> , blood; sup. ined. qual.)	Gram. 3. — 21.
24957 aristatum Sal.	awned	J C23 un 2 jn.jl Ap E. Indies 1803. 1) co	
24958 rugosum L.	wrinkly	J un O un 2 jl.au Ap K. Indies 1791. 1 co SdL st. 1.1	
24959 austral R. Br.	southern	J iAI un 2 jlau Ap N. Holl. 1822. Deo	
„ 2857. ^GILOPS L.	HARD GRASS.	(<i>Aix</i> , goat, <i>ops</i> , eye; sup. med. qual.)	Graminea. 7.
24960 ovata L.	ovate-apitaf	M O un f jnj.l Ap S. Europe 1683. S co Fl.gr. 1.93	
24961 triangularis L.	3-inchJipiked	M O un jjlau Ap S. Europe 1739. S co Schreb.gr. 1.10.1	
24962 triariifolia W.	threec-awned	M Q un 1 jLau Ap S. Europe 17 J9. S co Host gr. 2. 6	
24963 cyHindca Host	cylindrical	M O un 1 jn.jl Ap Hungary 1805. S co Host RT. 8. 7	
24964 squarrosa L.	tough-spiked	M A un 1J.jn.jl Ap Levant 1794. D co Schrb.gr. 2.27.3	
24965 hystrix Nut.	porcupine	Jlt A un 1 jn.jl Ap Missouri 1819. D co	
24966 caudata W.	tailed, Cretan	M Q un 1 jn.jl Ap Candia 1739. S co Fl.gr. 1.95	
2858. POTAMOTHILA A-Br.	POTAMOPHILA.	(<i>Potamos</i> , river, <i>phUeo</i> , to love.)	Gramfaea?. 1.
24967 parviflora B. Br.	smallflowered	M iAI un ... jl.au Ap N. Holl. 1820. D aq	
„ 2859. VALA'NXM L.	VALANTIA.	(<i>Sebastian Faillant</i> , a French botanist)	Bvbidcea*. 11.—12.
24968 muralis L.	wall	O un 4my.jl G S. Europe 1739. S co Col.ec. 297	
24969 hispida L.	bristly	O un 1 my.il G.T S. Europe 1768. S CO	
24970 filiformis H. K.	thread-shaped	O un Ijl.s G.v Canaries 1780. S co	
24971 Crucifolia L.	Crosswort	Jk A or I} my.jn Y Britain ... O co Eng.bot.143	
	Sm.		
24972 chersonensis W.	Cherson	-* A or 1 jn.jl Y Chenon 1817. D co	
24973 hispida W.	rough	Jk A un fjin.il G.Y Siberia 1804. D co	
24974 verna G. Don	spring	Jc A ^r 1 my.jn Y SwitzerL 1819. D co	
	glabra VU.	Galium vernum Hal.	
24975 pedemontana Bel.	Piedmont	O or ■ jl G.Y Hungary 1799. S co W. & K. 1. S3	
	Galium pedemontanum All.		
24976 articulata L.	jointed	O un 1 jl.au G.Y Egypt 1752. S co	
	Galium articulkum R. A. S.		
24977 humifosa IV.	trailing	■ A or i jn.jl Y Asia Min. 1816. D co	
24978 glabra L.	smooth	■ ^ u n 1 jlau G Germany 1371. D co W. & K. 1.32	
	Galium BauMni R. & S.		
„ 2860. BENINXA^&4 Savi BENINCASA.	(Count Benineasa, an	■ ■ ■ ■ ■	CunrbUhce*. 1.
24979 cerifera Savi	wax-bearing	-* iQ un 6 my.jl Y L. Indies ■ S co R.mal.8.3	
	Cucurbita cerifera Fis.		
2861. PARIETARIA L. PEBBLIORITY P <i>(Pebble) awl; commonly found on oMwalls.)</i> Urtke*. 10.—19.			
24980 indica X.	Indian	■ & (S) un 1jjapmy G Es Indies 1790! Deo	
24981 officinalis L.	official	■ A w 1 jns ■ Britain walls. D co Eng. bot 879	
24982 judaica L.	Jewish, Basil-lvd	■ A un 1 jns ■ Germany 1728. D co Schk.han.3.346	
24983 pennsylvanica MM.	Pennsylvanian	O un i jl ■ Pennsylv. 1821. S co	
24984 floridina Nut.	Florida	O un *jlau ■ Florida 1827. S co	
24985 urticifolia L.	Nettle-leaved	[HI un 1 jns ■ Bourbon 1700. S co	
	Bochineria tirticeflbia Spr.		
24986 lusitanica* Portug.	CMckweed-lvd	Jt O un i jlau ■ Spain 1710. S co Boc.sic.24. B	
24987 cretica L.	Cretan	O un i jlau ■ Crete 1818. S co	
24988 prostrate Lk.	prostrate	Jt A un ij.lau ■ 1823. D co	
24989 polygonoides W.	Polygonum-like	O un } jlau ■ Armenia 1728. S co	
2862. ifTRIPLEX L.	ORACHB.	(Ater, black; dark color.)	Chenopdea?. 35.—51.
	I. FRUTICBSIE.—Shrubby.		
24990 triflorum L.	Halimus	■ or 5 jl.au G Spain 1640. ■ ■ Par. th. 724.2	
24991 Anifilia Kth.	Flax-leaved	ti L-Jun + jlau ■ S. Amer. 1819. ■ ■	
^992 canescens Nut.	canescens	<L Jun 2 jlau G Missouri 1818. ■ ■	
	Calligonum canescens Ph.		
24993 portuloides L.	Portulaca-like	tt. or ■ Lau G Britain mud. s. C co Eng. bot. 261	
5 ^ 4 glauca L.	glaucous	tL i Jun ■ Lau G S. Europe 1732. C RJ Di. el. 40.46	
	white	■ Jun ■ jn.jl G C. G. H. 1774. C si	
	kidney-shaped	■ ■ ■ Lau G N. Holl. 1824. C sl	
	II. HERB^CRE.—Herbaceous.		
**&? rosea L.	Rose-like	O un 1* jn.jl G S. Europe 1739. S co Schk. han. a 350	
	white	O un lijlau G Italy 1820. S co	
	Siberian	O un 2 jlau G Siberia 1783. S co Schk. han. S ■	
	Tartarian	O un 2 jlau G Tartary 1778. S co Schk.han.3.539.	
	small-leaved	O un 1 jlau G Mexico 1827. S co [349]	
	Monte Video	O un 1 jlau G MonteVi. 1827. S co	
	diffuse	O un 2 jlau G Naples 1825. S co	
	garden	O cul ti jlau G Tartary 1548. S co	
	red, garden	O cul 6 jlau G Tartary 154a. S co	
	acuminate	O un 3 jlau G Hungary 1818. S co W. & K. 103	
25006 virgata Roth	twiggy	O un fj.lau G R. Indies 1810. S co	
25007 oblongifolia Kit.	oblong-leaved	O un 1 jlau G Hungary 1810. S co W. & K. 221	
25008 prostrata Bouch.	prostrate	i O un ij.lau G N. France 1820. S co	
25009 decumbens R. & 5.	decumbent	■ ■ ■ i jlau G N. Holl. 1820. S co	
	fringed	O w 1J jlau G Britain san.sh. S co Eng. bot 165	
	spreading	O w S jns G Britain dungb. S co Eng. bot 936	
	Venetian	O un ij.lau G Venice 1816. S co	
	hoary	O un 1 jlau G Spain 1818. S co	
	hastate	O un fj.lau G N. EuropelOO. S co Fl. dan. 1286	
	obtuse	O un 1 jlau G Siberia 1825. S co	
	narrow-leaved	O w } jn.ju G Britain rub. S co Eng. bot 1774	
25017 campestris Koch	field	O un 1 jlau G Germany 1816. S co	
	tatirica Schk.		

POLYGAM1A MOXCF.C1A.

(:LAHXXIII.

8864, TEUMIN.VUA7. TMvi». (*Tnmkm*, end) *Ivi In bunchC** At *vnUof brunch0* *OombrrAct**. !<—JI.

narrttw-lsavait	IQor	mflr/M,"W.ii"i;" ih.i'u-	naUfKuaira.	Jac.vin.3.100
S0030 molin-i) ni /"-".	I E3 ^{o'} 4 1 ! i	K .M W..	UM, ..	
45033 & ((' << ' I' SS03J inaitriti. ua / JIM , Mam	tftj ^{r"} tftj ^{...'}	SO ... BO w%		.1 Jack. 1.197 [·]
tft.uvaU or .	tbtcmprrtu-ii,	dry or battate; putamn round	anoint fly ittlratt.	
	Hor w	Y.i. E iii-lin 181	p.I Itos. cor. 3. SM	
	P o	M I Y.o KatafM tSSB, t	PJ	
	itr2+ti	Pair.		
It.JX	l)i> Mrrnbilanui	C i.r		
	alia	H rr		p-1
	IT,	SO	C 111	Rux. cot. 2.1ST
	Ir,MIK('tir	< 0 ...	C p.l	
		M ..	(` p.1	
SSatI ttitirtin />, C	twrbmwni	BO y o	C	
SJMi r.iiiuli(h.i / JJ. C.	i.....T.lavnl	32 -*ii	i: I.. r	
SSOH arnitucuS San.	dirub	Q or L1	i i c pi	
	ina	K	S. Ai. tan < p.l	

' MM. i r L COLFOOM. Ifultit, <||||||| - rr..., /r anil Icart like.) Sanali/tf'e. J.—5
S5WS cmmpr/uui /.. CainpfCTwrta/fid ft [l un]> U.w I.I. «ur/f.C.J8.1.1

SOU. B It Alii-". /, ArRiiiH ALKnNO, /lratieim, a > ; NCtmn or flower.) «w. 1.
2HHd ilrril.it uiu 7VJUH. ilirail 1 | Jur II lump W (1.P Brey. a i 1

" 2807. JPCER I.

faper* /ji/, C. »"« 1.vviyil.t II SSOTJ crunfjilia fitnt. •507B i- 8S07	or 20 ur 90 py V SO ajuny J thick. tared \ or SO ipi! dwarf	N. Anwr. IGSd. N. Ainrr. ... L Co N. Amor. ... L CO N. AIIHT, IH'T'i i N. Amur. IHJS. I 188). C pi II 188). C pi II 188). C pi II
Bo	Gil** M1. Hl I nj(n (...mri'.i. i mam	Guan, c.r.f m.t . MontpaBar.1 Hammem. tt — or 10 ... 180a < I'i 10 m W Indlc I , link. ,tl. SOI i OT III q Hi...lii IS/I < > 10 O.I V I,- i: [il Oac.fr, 3. Iftl 1 10 li. T II. V p.l • or 1U mjf.Jl O id H>K. cor.
SETS. GYROCA'RPU S Jne. CIY*oc*mni<- (* to luru round, koTpM, rrull t In sir) IMUT pffinf 4-t. O s af F Wit, koi. AIK'IMMN 1 Our . . . Y W7h, am 178	Attallo f CD or	C Lp Hut. tor 1.1
: boa Htam*. halj	unknown, i li i & II BOS ii Lp Bur.«C71.1	rtK ff, —a
- UBSgtannti TYTMN.	t 1Arlt I ted it iAJ«* .. JL Ifffl u 11. Bui LI	*. II. I7'H. D i p A
.uii Biiorui w. BuDBLU. jPrgftmorBn	MinhJ	C Lp
aoBeSSBd.n : uraiKlcit	10 SO)U Ap I mn. C li	
(50P7 mooUna if. W. iiiouitain	<>O ... Ap E. Itidle* C Lp	
• JVA Tiom. 1 lirs«; «npllr4t(in n<(r	J (r li ?d W C 1883. C p.1 fal	Temitrcnrl&oeit i— 4.
^BM. .(DPIC AtLitrrtM. [Atiania, tree of h w m ; lodif growth.) uitbu Of/ RlaiHloului	Terrblnihitcr,r, V. — +, or i» an I K I.p I' n bi —I or S*» ... 0 K. Imile* JMK, C <i> li>x. cur	
„ „877. CLU'SIA L. B*u*t i T ««. (Cartr// de rZcittx (Cfw/w. &5101 r.'4M L. SS09 ih* /.,	f O or tit ... W	
*SiW venvta L, uj.Ua*ed I Q n SO ... W & Ann I,, i I	S	
,, W7l OPHIOTCYLON /. OMBOXTUW. fo//Aii, serpent, fjrrow, wowl j lortn..		U- &
SiOS «iimittntmi /. icrpvlidHO ML O "" ^ Piyju W B. IsdtM MM I' n> Hot-1—*'''		
^m. MANOI'tEUAX, MuraoTwu. iVOHV, II. III. £IOS Ittdca /.. Indi.in fulid \ Lam.	W II IIH S > i»	
I VIM /;/. K. I ltii*«. flt, tphh, « «(11c i arutP awin if n.rc: A' I U I J IAJ ur li * ti	P, Itma>. 9. I t pj	
. SEAFO'RTM R. Br. van* B. Br. cltg	A. In i to ii lea I patron.) /Mfat*. 1. S Hull r.m	
MIUAN^Mart.	i MuteOKm PrinM Wdd-Kpiiwlnt) f t) U1 Mitt. B I'I <	I. — f.

CLASS XXIII. OJIDKK 9.

POLYGA'MIA DKFCIA.

low, Km i'altn t I J or '10 f,tm	the n run mi, r* ttk > in	r-Hm* 7
f<>, y* neilm trail	• ... <> vv '	
ijmcii un Uulan.i ? i or » or lor in ... 0	Nmwt Q w tiuuii* Ihvi I f "I	
OLKDITSCH/* L Mr Mil it	\ unuiikt, I*I[NJ].J s A.u. i. IVNt B <1 Dvn. In	Lrsim. f<>fj,r J1, -
lli(lino)	or 30 ijaji S a.) iXOO
		N. AN S *I
		N. Aiatr, I7fl. a > Cat car

POLYGAMIA DKECIA.

CLASS XXIII.

25128 h6rrida W. 2 purp6rca	horrid purple	or 10 or 20	jn.jl jn.jl	G G	China China	1774. 1774.	L L	p.i co	Dcn.br. 75
85123 ludica Pers.	Indian	or 20	jn.jl	G	Bengal	1812.	L	co	
25124 latiflqua Lo. C.	broad-podded	or 20	jn.jl	G			L	co	
25125 Hinensis Lam.	Chinese	Jor 20	... jn.jl	G	China	1812.	L	p.l	Dec. leg. 1.1
25126 ftrox Desf. orientalis Bosc	fierce	or 20	jn.jl	G			L	co	
25127 macracanthna Desf.	long-spined	or 20	in.jl	G			L	co	
25128 micracanthos Lo. C.	small-spined	or 10 or 20	jn.jl jn.jl	G G			L	co	
25129 c&tptica Desf.	Caspian			G	Caspia	1822.	L	co	
2885. CERATCTNIA L.	CAROB TBEE.	(Kras, a hom; horn-like pods.)					Legum. Cassiæ.	1.	
25130 siliqua L.	pod	1 Lj	cc 15	s.o.	R.Y.	Levant	1570.	S dJ Bot. rep.	567
2886. DALRYMPLA ROX.	DALRYMPIKA. (Alex. Dalrymple, auth. Oriental Repertory.)	Celastrinae.	1.22						
25131 nomifera Box.	apple-bearing	Si uJ or 10	ap.my	...	Nepal	1820.	C s.l	Rox. cor.	3.279
Turpinto pomifera Dec.									
2887. FRA'XINUS.	ASH TREE.	(Phraxis, a separation; used in formini.)					O&inte.	41.	
25132 americana L.	American, white%	tm 20	my	G	N. Amer.	1723.	G	co	
25133 acuminatula Lam.	pointed, green	tm 40	my	G	N. Amer.	1723.	(i)	CO	
25135 caroliniflta Lam.	Walnut-Tvd	or 40	my,jn	G	N. Amer.	17&.	G	co	Duroi 1.
95136 pubescens Walt.	Carolina	or 30	my,jn	G	N. Amer.	1783.	G	co	Cat car. 80
rfiiga Duroi	pubescent	or 20	my	G	N. Amer.	1811.	G	co	
25137 pannosa Fen.	cloth-taiwwf	V	tm 30	my	G	Carolina	1820.	G	co
25138 cplptra IV.	twing upon wing	x	tm 30	my	G	N. Amer.	1825.	G	co
discolor Mtc.									
25139 lancea Boce	lance-Avitt'rf		or 30	ap.my	G	N. Amer.	1820.	G	co
25140 quadrangulata Mx.	quadrang., blue		or 30	my	G	N. Amer.	1822.	G	co
25141 platycarpa Mx.	broad-fruited		or 30	my	G	N. Amer.	1820.	G	co
25142 expansa W.	expanded		or 30	my	G	N. Amer.	1824.	G	co
25143 mixta Bosc	mixed		or 30	my	G	N. Amer.	1824.	G	co
25144 pulvcrullnta Bosc	powdery		or 30	my	G	N. Amer.	1824.	G	co
25145 rubicincta Bosc	ruddy-veined		or 30	my	G	N. Amer.	1824.	G	co
25146 longifolia Bosc	long-leaved		or 30	my	G	N. Amer.	1824.	G	co
25147 vfridis Bosc	green		or 30	my	G	N. Amer.	1824.	G	co
25148 cinerea Bosc	grey		or 30	my	G	N. Amer.	1824.	G	co
25149 Jilba Bosc	white		or 30	ap.my	G	N. Amer.	1823.	G	co
25150 Rich&rdi Bosc	Richard's		or 30	ap.my	G	N. Amer.	...	G	co
25151 ovita Bosc	ovate		or 30	ap.my	G	N. Amer.	...	G	co
25152 cllpptica Bosc	elliptical		or 30	ap.my	G	N. Amer.	1825.	G	co
25153 irigra Bosc	black-branched		or 30	ap.my	G	N. Amer.	1825.	G	co
25154 fusca Bosc	fuscous		or 30	ap.my	G	N. Amer.	1823.	G	co
25155 rufa Bosc	rufous		or 30	ap.my	G	N. Ami-r.	1822.	G	co
25156 pâllda Bosc	pale		or 30	ap.my	G	N. Amur.	...	G	co
25157 exc6lsior W.	taller		tm 80	ap.my	G	Britain woods.	S s.l	Eng.bot.	4
crispia Bosc									
1 argentea	Wvcr.edged		or 20	ap.my	G	Britain	gard.	G	co
2 aurea	goldcn'aged		or 20	ap.ray	G	Britain	gard.	G	co
3 fungbsa	fungous		or 20	ap.my	G	Britain	gard.	G	co
4 hñontklis	horizontal		or 20	ap.my	G	Britain	gard.	G	co
5 nsua	dwarf		or 10	ap.my	G	Britain	gard.	G	co
6 verticillâris	whorled		or SO	ap.my	G	Britain	gard.	G	co
7 crtoa	gnawed		or 20	ap.my	G	Britain	gard.	G	co
8 strata	striated		or 20	up.my	G	Britain	gard.	G	co
9 pcndula Hort.	pendulous		or 20	ap.my	G	Britain	gard.	G	co
10 jaspidea Wro.	Jasper-lik.yel.-barked		or 30	ap.my	G	Britain	gard.	G	co
95158 atrovirens Pers.	dark-green		or 4	ap.my	G	Britain	gard.	G	co
25159 vrruuta LA.	warty	i	or 60	ap.my	G	England	Norf.	G	co
25160 lieteroph^lla Vahl	vario'us-leaved	i	tm 30	ip.my	G	England	woods.	G s.l	Eiig.bot.247G
simplicibflia W.									
25161 macroph^lla 7% ouii long.lcaved			or 40	ap.my	G		1823.	G	co
25162 parvifolia W.	small-leaved		or HO	apmy	G	Levant	1822.	G	co
25163 centiscifolia IV,	Lentiscus.lvd		or 6	my,jn	G	Aleppo	1710.	G	co
25164 argentea Lot.	silvery		or 15	my,jn	G	Corsica	1825.	G	co
25165 rambuciflta W.	Sambucus-lvd		or 30	my,jn	G	N. Amer.	1800.	G s.l	
25166 nanu Bosc	dwarf		or 6	my,jn	G			G	co
25167 oxyçarpa W.	sharp-fruited		or 20	—	G	Caucasus	1815.	G	co
2 oxyphylia Bleb.	sharp-leaved		or 20	...	G	& Europe	1821.	G	co
25168 lcidia Lo. C.	shining		or 20	ap.my	G			G	co
25169 amarissima Lo. C.	bitterest		or 20	ap.my	G			G	co
25170 appendica Lo. C.	appendaged		or 20	ap.my	G			G	eo
25171 appendiculata Lo. C.	appendiculate	i	or 20	ap.my	G			G	co
25172 ylrens Lo. C.	green		or 20	ap.my	G			G	co
2 variegata	variegated		or 20	ap	G			G	co
2888. BRO'SIMUM Sun.	BREAD NUT.	(Brositnos, good to eat; fruit eatable.)					Vrticea:	2.-4.	
25173 ijlicastruin Swx.	Alicastrum	M • or 6	...	Ap	Jamaica	1776.	C r.m	Swz. fl.	1.1.1
25174 sphrium Swx.	spurious, AJukwood	Q or 6	...	Ap	Jamaica	1789.	C r.m		
2889. DIOSPVBOS L.	DATE PLUM.	(Dios, divine, pyros, (err.) pear; resemblance.)					Ebenuctte.	22. — S	
1. GL&BRIfbuji.	— Leaves on both surfaces smooth.								
25175 virginikna L.	Virginian, American**	dt 20	jn.jl	Y.u	N. Amer.	1629.	S s.l	Dcn.br.	146
25176 llcida Lo. C.	shining	\$ or 15	jn.jl	Y.u	N. Amer.	1820.	L	B.I	
9BITJ Mabbia Rox.	Mabola	D f r	8	—	Y.a	Philip. Is.	1828.	C s.l	Bot reg. II f
25178 sylviflta Box.	wood	£ or £0	—	—	W	E. Indies	1801.	C s.l	
25179 /cyclo'idies Desf.	Lycium-hke	G o r	10	—	W	E. Indies	17U2.	C s.l	Ac.haf.2L5
25180 A'bicum Retx.	Ebony	f	tm 30	—	W	Bengal	1792.	C Bl	Ru.am. 1.6
251H Ebcnaster lictx.	Ebenaster	T	or 20	—	W'	E. Indies	Ihl7.	C s.l	
	wooded	O	or 20	—	W'	E. Indies	IS 18.	L rin	Cot. reg.41'9
	Einbryoptcris	I J or 25	jl	—	W.c	Mauritius	1824.	C s.l	
	netted	L J or 20	—	—	E	Indira	IS 24.	C <l	
25185 edulis Lo. c.	eatable	I clt fcQ	—	—	E	China	1822.	L s.l	
25186 lobitta Lou.		I J clt IS	—	—					

- 25187 Kiccinioldes *Litutl.* Vaccinium-likcii il pr 2 ap.my W China 1823. C r.m Hook. ex. fl. 139
 25188 discolor W. discolored f 7- fr 20. . . . Philippin. 1823. C r.m
 25189 montana Rox. mountain * i To r 6 ... W.a E. Indies 1822. C r.m
- II. PUBEBCE'NTBS. — *Leaves underneath pubescent.*
- 25190 L&tua L. Lotus, European* dt 20 jn.jl Y.a Italy 1596. L s.l Mil. ic. 116
 25191 pubescens Ph. pubescent \$ clt 20 ap Yc N. Amer. 1812. C s.l
 25192 Kuki L. Kaki, Japan i J fr IS - W.a Japan 1789. L r.m Kami. 806
 25193 hirsuta L. hairy - or 20 - - Ceylon 1820. L r.m Kami. 806
 25194 chlorAxyton Jto*. pale-wooded - or 20 - - E. Indies 1822. C s.l
 20195 cord) folia Rox. heart-leaved - or 15 - - W.a E. Indies 1794. C s.l
 25196 rugulosa A. Br. small-wrinkly - or 20 - - N. Holl. 1823. C R. Rox. cor. 150
 25197 obovata J.k. obovate - or 15 - - W.a W. Indies 1796. G co Jac. sc. 3.318
2890. CARGIXL/il JR Br. CAROILLIA. (*James CargUL M.D., of Aberdeen.*) *Ebenace**. 2.
 251CB australis R. Br. southern t J or 15 ... N. Holl. 1816. C 1.p
 25199 laxa R. Br. loose f I J or 15. . . . N. Holl. 1827. C 1.p
2891. AFYRSINE L. (A Greek name of the myrrh.) *Myrsine**. 19. — 26.
 25200 africana L. L.Jor 4 mr.myBr C. G. H. 1691. C pl Coin.h.l.M
 2J201 retusaL. L.Jor 2 jn W.a Azores 1778. C pl Ven. ecls 80
 2Vj02 scmiincrata JZox. L.Jor 40 Pk Nepal 1822. C p.l
 2sauJ capitulata Rox. L.Jor 30 G Nepal 1822. C p.l
 2W04 biliaria Rox. L.Jor 20 V./pk Nepal 1822. C p.l
 20205 subspinosa D. Don Kubspinoze L.Jor 20 - - Nepal 182a C p.l
 25206 coriacea A. Br. leathery L.Jor 8 - - W.a Jamaica 1823. C p.l
 Samara coriacea Swx.
- 25207 S&mara R. Br. Samara - 3 f.n W.a C. G. H. 177a C 1.p
 Samara pentandra H. K.
- 25208 melandrophicos B. Br. black paper Siderfxylon melanophleum L. - - W.a C. G. H. 1783. C 1.p Jac. vin. 1.71
- 25209 vari&bihs R. Br. variable L.Jor 3 Jnjl - - N. Holl. 1824. C 21
 25210 ilicifolia Lo. C. & Holly-leaved L.Jor 3 ... 18&L C p.l
 25211 venuibsa Spr. small-veiny L.Jor SO jn.au - - C. G. H. 181d C p.l
 Siderfxylon venulsum IV.
2892. NY'SSIL L. TUPELO. (The name of a water nymph (?); habitat) *Santalatca**. 7. — *.
 25212 villbsa Mx. hairy, Sour-gum - or 10 my G N. Amer. 1806. L s.l Mic.ar.21
 25213 bifbra Mx. two-flwd, mountain^ or 10 ap.my G N. Amer. 1739. C s.l Mic.ar.22
 aquatica L. caroliniana L.
 25214 capitata Walt. round-headed *f or 10 ... G N. Amer. 1806. F II
 25215 toment&s Mx. downy \$ or 50 ap.my G N. Amer. 1818. L s.l
 grandident^ta Mx.
 25216 candicansA/r. whitish, Oacchce tinted or 20 - G N. Amer. 1812. C s.l Mic. ar. 20
 25217 denticulata H. K. sraah-toothed \$ or 30 - G N. Amer. 1735. L s.l Cat car. L GO
 angunzans Mx. angulfts Pair.
 *25218 sylvatica Lo. C. wood ? or 15 ap.my G N. Amer. 1824. L s.l
2893. HAMILTO*N/if Mhl. OIL NUT. (*Mr. Hamilton, an American botanist*) *Santalcea?*. 1.
 25219 olefrera Mhl. olive-bearing il Jor 6 ... G.Y N. Amer. 1800. C s.l
- Pyrularia piibera Mx.
2894. LAUROPHYLUS Thun. LAUROPHYLMS. (*Lanrus, laurel, phylon, leaf: resemblance.*) - - - - -
 25220 captasis Thun. Cape i j o r 6 ... G C. G. H. 1801. L pl
2895. BU'RSERif Jac. BURBBRA. (*Joachim Burser, a disciple of Caspar Rauhin.*) *Terebinthccee*. 2. - 4.
 22222 gummifera Joe, Jamaica, gum-bgt • or 20 ... W.a W. Indies 1690. S pl Jac. am. G3
 22222 serrata Rox. serrated J or 30. . . . E. Indies 1818. C p.l
- ors5*- ANACA'RDIUM Rox. AVACARonpr. (*Ana, without, kardia, heart: seed external.*) *Terebinthuvar*. 1.
 ^223 occidentMc L. western f Q f r 20 ... G.B W. Indies Ni99. C r.m Jac. am. 1.18135
 2 indicum Dec. • Indian \$ • fr 20 ... G.B E. Indies 1699. C r.m Hu. am. 1. GO
- *297. SEMEC A'RPUS L. SEMRCARPUS. (*Semeio, to mark, karpos, fruit; use of juice.*) *Terebinthdcar*. 1.
 *** Anacardium L. Anacardium ± fr J0 ... G.Y E. Indies 1820. C r.m Itu. am. 1.70
 Anacardium longifolium F*ast. Cassitivum Spr.
 2 cuneifolium Dec. wedge-leaved f • fr 20 ... G.Y E. Indies 1824. C r.m
2898. ARCTO^PUS L. ARCTOPUB. (Arcto, a bear, pouss, a foot) *UmbeUiferar*. 1.
 2998 cchinatus L. heilgehog - 1 my.jn G C. G. H. 1774. D p.l Bot reg.70*
- PANAX. (*Pan, all, akos, remedy; medical qualities.*) *Aralidce**. 9. - 11.
 Ginseng, 5-lvd k A pr 1|jn LY N. Aincr. 1740. 1) s.p Bot. mag. 133.1
 prickly pr m 1^n G N. Amer. 1759. D s.p Bot mag I, II
 shrubby pr 6 au.s G China 1773. C sp Jac. ic. 3.634
 attenuated • or 20 - - W. Indies 1823. C p.l
 tree J L.J or 20 - - N. Zeal. 1820. C p.l
 golden-leaved J • or 20 - - Guiana 1822. C p.l Aub. gui. £. 3(10
 fragrant J Dor 10 - - E. Indies J815. C p.l
 shell-leaved f • or 20 - - E. Indies 1820. C p.l Ru.am.4.31
- *2900. FrCUS L. Fro TREE. (Derivation unknown.) *Urticca?* 74 - 1+
- I. CORUFOLIA. — *Leaves cordate, entire.*
- i. GLABRIKOLM!. — *Leaves smooth.*
- §£. £ "lacroph"lla Desf. long-leaved ABRI or 14 - Ap N. Holl. ... C p.l
 Vrf.2 wymphojRaia L. Water-lily-lvd or 10 - Ap E. Indies 1159. C p.l
 'C'gosa L. religious, Po,Jar-lyd 95 - Ap E. Indies 17-31. C p.l R. mal. 1 27
 22w. Myrtlc-leaved O o r 6 - Ap S. Amer. 1775. j. Pluk. al. r/i. |
 »»yrtiflbia Lk. thick-nerved \$ • or 10 - Ap S. Amor. 1823. p.l Thun. fir 8. c.
 Desf. rib-leavrtt or 10 - Ap E. Indies 17/23. p.l
 XVIR^U Vahl poUsed E 6 - Ap W. Indies 1823. p.l
 corUcca H. K. leathery-/i/wy/ or 10 - Ap E. Indies 1772. p.l Hook. ex. fl. 423
 'cida H. K. shining^/awrf oblong-leavcd J Q o r - An C. G. H. 1825. p.l

252*7 lontiginbu *Tahl*
 «, «« venfea //, K.
 *52W Infct&ria JK
 £5250 Iucatoma *Polr.*
 ventwa W. not //, if.
 SSS1 rUdfbtle 1* a
 SMW itipulita ?W

S5254 lomentbia A u,

2555 tinctoria *Font.*
 2S256 Be7fam)im L,
 95937 tx-tijhalciui* £•
 2SS55 pDmila L.
 SSSB popflnea ff^
 £SS1 cajculit* JW-

2SSGS butoprtfl* U.
 £5483 raso M 0. AM

«fla» long! folia *Lo.* C.
 £5265 allptira *Kth.*
 \$5306 dumiihi Lo. C.
 S527 return L.
 £52>i di.tica Ro*.
 S528) Af*IM Ci Don
 SM70«UbrfcU AM.
 SSB71 ntidu 7.Aun.
 £5279 Hookcrir riwt.
 III III 11 4 nOHi

2577 gloriorita *Rax.*
 74 minima /to*
 brtu »f.

, L*.
 10101 Li.
 SSS79 aiiirri.i.tii.i *Aubl.*
 SSSAt niiR.K.-,I>:1 /Vra/
 €SVH 111. HIIHIMHL AM.
 25283 obluiifbla JCU

£5284 auitrklU IT.

S598S Logfcmfi G. Don.
 SSB8B mart in iL-iwii IK
 i UPBRJ //, A.
 fiffIBV Initn-i Fit*
 •invife«« Hurt, indlca
 25285) rtrnaJa Rtut.
 2SSU0 Inicwillit,jj /Itur.
 S52U1 an.r w fulfil 7.0. C.
 £Wf p<iiidula Lk.
 9SS3 brulll&uil Li.

•, -i i-i)CQiii 21 w
 m^qj^ Tjchtecijjtcinij TrV

89sB0 fcncnt if*

SM6« BriM« ft. Itr.
 3SS99 oxiportU r<A/

25300 onpolitipjla /or.
 icftbra jnC
 85301 coreahha vrt

£5304 mtatairrti lir.

S0305 ipora *Font.*

KBOI ...fn-, (1 .
 ; Us i.

ipw.Wrf J a or 14 Ap W. Indies 1830. C >
 ntWT tifiwri J CII or in ... Ap E. Indie* C p B. mal, 3.64
 •ulnha f CD or ... Ap All; hilkH C p
 whiite-i IL(1 • LJ « BO ... Ap K. Indies 176U C £ W. h. 1x1.36
 CIJCIMKCU Spr. 10 ... Ap isai. C P.
 tainniy-leavtil ... Ap 1810. C P.
 •tpnteM i' ... Ap C P.

It. Susnras Liatrt utuh-rneath pttbetcnrt. C
 woolly t dor ... Ap K huliM 1816. r 1-

II. OVUTIVULIJC — I MM (vaU, entire.

1. L*tiuN r.*. —Leovet smooth.

•UitiUiff J LFI Of 15 myjn Ap Swiotti 1. C P
 itrnjatMB tnn f CD ts 10 ... Ap // 1 mnifft C P
 IkiitKtl i i or « Ap K. li. C P
 dwarf £. L J ... Ap (Mai C Pi
 tupomitiTiiu* *, 2 12 Ap S. Amer. 181V. C P
 c«Jyculato nt ... Ap N. Spain ICUO. C

ii. LASIOUHIVI 4MIVJ puberirt.

wrinkly**1 B S u » Ap S. Leone ten C P
 III. OB i oblong, entire.

1. 11'si S — I riivrt smooth C P
 long. li-av til * f) n pa... Ap 1825. C P
 (.liptic Ci or £0 MM Ap S. Ainer. 1884. C P
 tmny ' i-if 6 AP
 tiiiiu-/('i(^ 3 at (t Ap h 1 Nil 111 17WF C P
 elaitic K>m Ati'luit'i 1 I. II. Ap & .. C P
 MI II ii't d I] or in Ap Cam ecu C P
 uV*?y-tfaw.d i 3« 6 mr!jn A|i IndiLf C P
 Houker'i O or 6 ... Ap R moi. a r,r, C P
 i-iliTHT,lt- d i • or 10 Ap ' 1: JB18. C P cor. I.123
 UIAc'd T 1 J ur IP Ap ' 1: Irulii» in IN. C P Jtox. cot it \Sb
 pil'Ulli T t ' i Id .. Ap M.iiriilut C P Jtury 17
 piTOOIUMMnf « ; of 8 Ap H. Atuer. C P
 Artiutu*-leavedA Q or 3 Ap C P
 clh * : _ or 4 Ap C P
 A iii'ricaii : _ or 8 Ap IIU WII). C P
 eluv'. imjill.tniitul 1 D or 4 Ap 1 IIIHIM C P
 Cut Ullit 1 fit Vlf 90 K> Oula IBIJ. C P
 blunt-LMved 10 Ap Mi-k 1 Ap U. uial. 1. Ji
 90 Ap Ap I p I

ii. 1 it, —Leave* pvtKrt.

*outittn mr.jn Ap 1789. C P.I "Veil, msi.114

IV. IIAXCirbum. —Letters ianculaie. C P
 1>ij{iiii'i t CD or B .. Ap p 1
 Martinltue I CJ M B .. Ap W, Inditi IIM. C P. 1 SI. Jain.2.ia3
 Indian J CD 'T ID .. Ap R liuli« 1750. C P. R. mal. 3 03
 Wlluw-leaved r L i ^ 10 .. Ap Arutii.i C P
 •infe*ce* Hurt, indlca J CJ "T SO .. Ap R Indii-i C P
 25285) rtrnaJa Rtut. >r:rije: like T (*-) or ID .. Ap 1 IIIliM C P
 2SSU0 Inicwillit,jj /Itur. IM'DIIIIdIII f CD ° IS AII C P
 S52U1 an.r w fulfil 7.0. C. liraillian *D»r 4 Ap BcuU 1823. C P

V. Sean*ripbu,v.--Lⁿw'i serrated ur teoUucL

L Cpt • i | «t 4 Ap C G. H. 181U. C P.I
 Llchtietyl* B^ or AD C G. If. 1821. C P
 if. Si HMIII*LIJt. C P

rriirijii if:-»)!/* n of i Ap i: iMtiili'i ISQS. C P
 Mm Ieavod •1 CD or 4 Ap PrdBppin. C P
 lir.-imi'i • j T Ti 1! .. A. B LCOM 1 P
 nuHpmtdad Ap Guinea 1S00. t: C P

t* n 7/ur Up>HM)C.liaVcf • • or 4 Ap B. ItidlM 1802. C P. ROK. COT. 2, IS4

crowned • dor 6 myjn Ap IS00. C P. ((l. h.r)p.B

illuill • r*-l or 4 Ap C P

If until A IJ or 4 * Ap N. Boa C P
 K d. HIT VCl IDor 10 .. Ap Brutl C BO

II BMHI J ill. olnVj r^ VH'i Sit C P

rough -/^i* i-| ir 3 Ap ' N. Holl. 1807. C P

IV. J O or I.IT Ap ii i. 1750. C P. LI 1
 aquatic varutbie-iuTd III Ap labaT 1548. C P. n.nii!. 3. *32
 common, fig 15 Ju.Jl Ap 'Tr. clir. T-i- 4

CLASS XXIV. ORDER I.

CRYPTOGAMIA FILICES

TRIBE I. POLYP(S)DIA⁵ CEA

*2901 POLYBO'TRY A H.% B. POLYBOTRYA. (*Poii*s, many, *botrys*, raceme: fertile div. of frond.") 2.-6.
§509 cervina AT/IM/. hart's/oMgu* jg El or 4 ap.my Br Jamaica 1823. D l.p Petfl. 8.3
**310 vivipara Hook. viviparous f El or 1 my.au Br W. Indies 1883. D l.p Hook. etVfl. 107

*2X2 ACRCSTICHUM L. AcROSTICMNR. (*Akros*, highest, *stichos*, order; lines on reverse of lvi.) 9.-65

I. INPIVIN. — *Fronds undivided.*
J511 simplex Swz. simple jg El or 1 ... Br Jamaica 1793. D l.p Bot.cab. 709
J12 longifidium Jac. long-leaved ff El or 1 my.s Br Jamaica 1817. D l.p
Jw13 ermitum Swx. hairy ff El or J jl.au Br W. Indies 1793. I) fp Tl. fil. 12ii
**314 glandulosum Hook. glandulous ff El or 1 ... Br Jamaica 1825. I) l.p H.-&G.3.

II. DiVIMA. — *Fronds simple, divided.*
alcicurne Swx. elk's horn jfi iAJ CU f au.o Br N. & W. 1803. D s.p Bot. rec. 862-3
U Stcmaria Beauo. Stcmaria f El or 1 my.au Br Guinea 1823. D l.p Beau. ow. 1.2
§317 appendiculatum 7/oo Jlr.appendiculate jf Tl. vior lfjtn.au Br W. Indies 1824. D i.p H. & G. 100
§318 fimbriatum 11. L. fringed < S or 1 ... Br Brazil 1824. I) l.»
*319 aurcum L. golden jf El or 4 au Br W. Indies 1815. D l.p Pl. fil. 101

2903. J/EMIONITISL. HRMIONITIB. (*Hemionos*, a mule; supposed to be sterile.) 1.
§520 palmata A. palmate f (29 el } jn.au Br W. Indies 1793. D l.p Hook. ex. II. J3

* 2904. OYMNOGRA'MM A Dcs. GYMNOOR. (*Gymnos*, naked, *gramma*, writing; naked sort.) 10.—31.

I. HIRSUVA: — *Fronds underneath hairy or scaly.*
§21 pedata Kaulf. pedate f El br ft jn.jl Br K. Spain 1822. D l.p Swz. fil. 1.3
//cmiorittia pedata Swx.
§i myriophylla Swz. myriad-leaved f El or 1 ... Br Brasil 1824. D i.p H.&G.4.¹
§vfa Desv. red-haired jf El pr ft jn.au Br Jamaica 1793. D l.p Schk.fl. 17.21
7/emionitis rufa Swx. Acrostichum rufum L.

III. FAROIJJK. — *Fronds underneath mealy.*
xi<>24 trifoliata Desv. three-leaved jg El or 2 jl.au Br Jamaica 1810. D l.p Pl. fl. 144
Acrostichum trifolium L.
§jj*5 chrytophylla Kauff. golden-leaved f El or 1 jl.au * Br W. Indies 1824. D i.p Pl. fl. 44
Acrostichum chrysophyllum Swz.
rgj Bulphorea Desv. sulphur sulphur f El or 1 jn.jl Br Jamaica 1R0R. D l.p Schk. cryp. 4
§gj tartarea Desv. infinal f El or 1 au Br W. Indies 1817. D l.p
//emionitis dealbata W.
JDcruvka Desv. Peruvian jp El or 1 jn.nu Br Peru 1822. I) l.p
calomelanos Kauff. beautiful black f El el 2jl.au Br W. Indies 1790. D i.p W. h.b.41
Acr6stichum calomelanos L.

III. Nu^oA. — *Fronds without any hairiness or mealiness.*
§ctophylla Desv. slender-leaved ^ A or 1 jh au Br S. Europe 1819. 1) l.p H. A ii. '23
^oly)buim lupophyllum L. Acrostichum leptophyllum Dec.

■ MENI'SCIUM Sehreb. MENISCIUM. (*Meniskos*, a crescent; shape of fructification.) 3.-7.
proliferum Swx. proliferous f Qs or 1 ap.jr Br E. Indies 1820. D co
§Wp "W. Wionis prolifera Ilefx.
§Wp *orbifluit IV. Sorbus-leaved jf El or 1 ... Br Brazil 182 J. D co
4spli"niuim *orbifluit* Joe.
cticulatum Swx. netted f El el f np.iny Br Martinico 1793. D l.p Pl. fl. 110
•folypbdium reticulatum L.

■ XIPHO'PTERIS Kauft. SWORD FERN. (*Xiphos*, sword, *pfcris*, fern; form of fronds.) 2. — 8.
"errulata Kauft. serrulate f El pr * Jn.jl Br W. Indies 1823. D l.p Schk. cryp. 7
Stair JrammMis serrulata W.
" hctvrnphylla S/tr. variable-leaved ^ iAJ or \ jn.jl Dr N. 11 oil. 1824. D co
urammitis heterophylla Lab. Polypodium grammoides R. Br.

■ CF/TERACH W. CETERACIL. (Called *chetherak* by the Persian physicians.) 1.—4.
mijArum W. officinal <: A m f my.o Bj Britain cal. ro. D l.p Eng. hot 1244
^oplenium Cterach L. GrammUis Citerach Swz.

■ J*OLYPO*DIUM L. POLYPODY. (*Polys*, many, pow, foot; multitude of roots like.) 52-1'JO

I. iNmm*HA. — *Fronds simple, undivided.*
§337 DISCREPANTIA. — *Bearing fertile and sterile fronds.*
Mousc-car-hkeJuElor i AU Br W. Indies 1793. D l.p Pl. fl. 118
variable-leaved HW El or i jn.au Br W. Indies 1820. D l.p Pl. fl. 120
creeping JWEElpr ftmyjn Br W. Indies 1816. D l.p Pl. fl. 121

II. FKRTILIA. — *Fronds all fertile.*
■ i R. Br. attenuated * mrjl nr N. Holl. 1823. D l.p
itbliinn Swx. narrow-leaved ftinrau Irl W. Indies 1H90. 1) l.p
un Hum. banded 2 inyjn Br S. Amor. IH15. D l.p
creeping inyjn ff W. Indies 1810. I) l.p Pl. fl. 134
Club-moss ff W. Indies 1792. I) l.p Schk. fl. 8
lanceolate ff El or i Br W. Indies 1812. I) l.p IM III. 137
2-11, 11' l'y'ifulis/.. Hart'8-tongue ff El or 1 an Br W. Indies 1791. Sksp Pl. III. ISO
8-11' siKMiOMiin Swx. thick-leaved ? El or jn.s Br E. Indies 18s.B. ?? l.p
4-11' itol>um L. Plantain-like f El or 3 aus Br W. Indies 1823. D l.p p|. fl. i.22
Vahl Plantain-like f El or 1 jn.au Ur W. Indies 1817. D l.p Jaç c. lit

CRYPTOGAMIA FILICES.

CLASS XXIV.

11. PI.VNATI'FIDA.—*Fronds pinna/ifid,*
- I. IMTEGERRIMA.—*Leaves very entire, smooth, from No. 25 fil. pubescent underneath.*
- | | | | | | |
|---|--|---|--|--------------------------|--------------------------|
| 25350 phymatidps <i>L.</i> | tubercl. £ Q5 or | Jr | I* Indies 1823. | D l.p | Pluk. ph. 404.5 |
| 25151 pustulatum For*. /. | puited | £ [] or | mr.au Br | N. Zeal. IH20. | ■ |
| 25352 Billarditri R. Dr. | scändens Lab. | La Billardicre's JW | mr.au Br | V. U. I. 1823. | D fp Lab.n.h.2.240 |
| 8535.1 aurcum <i>L.</i> | golden | f El or | 3 mr.ap Br | W. Indies 1742. | SkB.p PI. fil. 76 |
| 25,r4 7ueriffT>hum fc. | Oak-leaved | £ El or | Br | E. Indies 1824. | I) l.p Ru. am. 6.36 |
| 25,lw arcolltinum //MOT. | areolate | £ 123 or | — Br | Brazil 1824. | I) l.p |
| 25,15i Catharine Fin. | St. Catharine's | £ El or | — Br | Brazil 1824. | I) l.p |
| 25,07 curvatum <i>Swz.</i> | curved | £ G5 or | — Br | Jamaica 1823. | I) l.p |
| 25,15b pectinatum <i>L.</i> | comb. | £ ED or | — Br | W. Indies 1713. | Sk Bp Bot.cab.748 |
| 25359 plbmula //<>. | feathered | E El or | 1 jn.nu Br | S. Amer. 1824. | D l.p |
| 23,M0 icoloKMHnemlesJ. | Scopelundrum-lkft | pg or | 1 jl.au Br | W. Indies 1820. | D l.p PI. fil. 91 |
| 253,il /irheimanoldes.Swz. | Trichumanes-lkf | LS or | 1 my.jl Br | W. Indies 1822. | I) l.p Schk. rryp. 10 |
| £ [] - p - [] | Spulenwort-lvd | [A] or | — Br | Martinico 175K. | Sks.p PI. fil. 102 A |
| 2536J incanum Sun. | hoary | £ (23 or | — Br | S. Amer. 1811. | D l.p Schk. m. 11. u. |
| £5,364 decumanum W. | ii. SBRRAVA.— <i>Leaves dentate or serrated.</i> | | | | |
| 9,105 incium Sws. | tall | £ El el | 5 au Br | Brazil 1818. | D l.p |
| 2d,96 trifurcatum L. | cut | £ El or | 1 au Br | W. Indies 1810. | D l.p |
| 25367 vulgoe W. | three-forked | £ 22 or | jau Br | W. Indies 1820. | I) l.p PI. fil. 138 |
| 2 cambricum | common | S A or | 1 my.o Br | Britain sha.ua. | D l.p Eng.bot. 114! |
| 25368 virginianum L. | Welsh | — | 1 my.o Br | Britain | D l.p Bolt.fil. 2.5. a |
| | Virginian | £ Z or | 1 jl Br | N. Amerc. | ... D l.p PL fil. 77 |
| | III. PINNAVA.— <i>Fronds pinnate.</i> | | | | |
| | L INTEQERRIMA.— <i>Leaves very entire.</i> | | | | |
| 25369 hastatum Swx. | hastate | />E J or | 2 my.au ... | Jamaica 1820. | D l.p |
| 25370 longifblum Presl | long-leaved | /f El or | 3 my.au ... | Brazil 1819. | D Lp |
| 25J71 aunculatum Presl | eared | fi | — El or 2 my.au ... | Brazil 1824. | D l.p |
| 25372 diastmleL. | dissimilar | £ El or | 2 my.au ... | Brazil 1820. | D l.p Pluk. al. 228.1 |
| | ii. SERBAVA.— <i>Leaves repand, crenate, serrate.</i> | | | | |
| 25,173/raxinifblum Jac, | A heleaved | ffff El el | 2 au Br | Caracas 1817. | D l.p Jac. ic. 639 |
| 25374 rc>aiidlii Sun. | repand | £ El or | 1* au Br | Jamaica 1820. | D Lp |
| 25,175 tenullum Furst. | slender | &riA or | 11 au Br | N. Holl. 1823. | D Lp |
| 25,J76 crenatin Swx. | crenate | 1} au Br | Jamaica 1823. | D Lp | |
| 25,J77 Juglandifblum //. | 4 B. Juglans-lvd | £ El or | 1 my.au Br | & Amer. 1822. | D Lp |
| | IV. BIPINNATIPIDA.— <i>Fronds bipinnatifid, pubescent.</i> | | | | |
| 25378 Pheopteris L. | Sun-fern | k A or | f jnj Br | Britain moun. | D l.p Eng.bot.2234 |
| 25,I79 hexajonopterum Mx | ct>anglw-wgd A | A or | 1 Jj | 5 N. Amer. 1811. | D l.p Pluk. al. 284.2 |
| Q'IKO CUMI>C ^{MT} | connected | 5 A or | 1 jn.il Br | Canada 18'23. | I) l.p |
| &WV0 coniectile Mi. | Jamaica | £ E J r 1 | IJL.au Br | Jamaica 1820. | D l.p |
| 25381 jamaicense Dcs. | V. BIPINNA*TUM.— <i>Fronds bipinnate.</i> | | | | |
| 25382 sanctum Swx. | holy | £ E) or | 1jjl.au Br | W. Indies 1820. | D Lp |
| | VI. TRIPINNAVA.— <i>Fronds Mpinneate or supra-decompound.</i> | | | | |
| 25383 DryAptcris L. | Dryopteris | ^ ^ or | 1 jn.s Br | Britain moi.pl. | D l.p Eng. bot Old |
| 25384 calcareum Sm. | snur.hranched | 5 A or | Jjl Br | Britain cal.ro. | D l.p Eng. bot 1525 |
| 25385 effusum Swx. | spreading | — El or | 3 n Br | Jamaica 1761. | SL Jam. 1.57.3 |
| 25386 pruinatum Swx. | froaUcavd | £ El or | 2 s Br | Jamaica 1793. | D Lp |
| | VII. DU'BIA.— <i>Doubtful to which of the sections they belong.</i> | | | | |
| 25387 polyfiathum H. L. | manly-flowered | £ [] & | : : Br | Brazil 1824. | D Lp |
| 25388 f & L | for>igd | £ [] & | : : Br | Brazil 1824. | D Lp |
| | 290!. T ^ NITIS Swx. | — | (Tainia, a fillet; long narrow frond.) | a—9. | |
| 25389 lancolato Kaulf. | lancolato | Ar El pr | 1 au Br | W. Indies 1818. | D Lp PI. fil. 132 |
| l'tens lanceolata L. | | | | | |
| 25,390 ^rammifolia Hook. | Grass-leaved | £ I or | ijn.au Br | Trinidad 1820. | D Lp |
| 25391 angustiflia Spr. | narrow.leaved | £ I or | i jn.au Br | Jamaica 1816. | D Lp Hook. ex. fl. 77 |
| Pteris angustifolia Swx. | | | | | |
| | *2910. NOTHOCHLfl^NA R.Br. NOTHOCYIL. (<i>Nothos, spurious, chlaina, cloak: seeming invol.</i>) 4.—13. | | | | |
| 25392 pilselfoldca Kautf. | Pilosella-like | £ El or | ijn.au Br | E. Indies. 1822. | D Lp H.pfl. 96.2 |
| /*teris piloselloidcs L. | adnascent | % El cu | \ mr.au Br | E. Indies | 1824. D Lp R. mal. 29 |
| 25393 Marant> II. Br. | Maranta's | £ IAJor | Jjn.au Br | N. Holl. 1820. | D Lp Spr. cryp. 2.18 |
| AcroHtchum MaranUr L. | distant | £ iAJ or | f jn.au Br | N. Holl. 1823. | D Lp |
| 25394 distans III. Br. | wodly | tf IAJ or | \ aus Br | Madeira 1778. | R i p Desf. at 2.256 |
| 25395 lanRintaa Desf. | | | | | |
| Acrostichum villicum //. | | | | | |
| Acrostichum villicum //. | Acrostichum villicum //. | Acrostichum villicum //. | Acrostichum villicum //. | Acrostichum villicum //. | Acrostichum villicum //. |
| | Acrostichum villicum //. | Acrostichum villicum //. | Acrostichum villicum //. | Acrostichum villicum //. | Acrostichum villicum //. |
| | | | | | |
| 291L NIPHCyBOLUS Kaulf. NIP. (<i>Nip/iobolos, covered with snow; indusia like flakes of snow.</i>) 6.—17. | | | | | |
| | 1. DiHCUPA'NTBfl. — Bearing fertile and sterile fronds. | | | | |
| 25396 adnascens Kaulf. | adnascents | % El cu | \ mr.au Br | E. Indies 1824. | D Lp R. mal. 29 |
| Polyp>klum adnascens Swx. | | | | | |
| 25397 pertukih Spr. | Iwred | H^Elcu | mr.au Br | China 1821. | D p.l Hook. ex. fl. 162 |
| Polypodium pertiflum Hook. | | | | | |
| 25398 rup'stris Spr. | rock | ^ El cu | mr.au Br | N. Holl. 1824. | D p.l |
| Polyptodium rupestre Jl. Br. | | | | | |
| | 11. CONF6RMES.— <i>Fronds all fertile.</i> | | | | |
| 25399 confluent Spr. | confluent | IWEJcu | f mr.au Br | N. Holl 1820. | D p.l |
| Poly>odium c6nfluens R. Br. | | | | | |
| 25400 lingua Spr. | tongue | HW El cu | } mujn Br | Japan 1817. | D p.l Thun. Jap. 38 |
| Acrostichum lingua Thun. | Acrostichum lingua Thun. | Polyptodium lingua Swx. | | | |
| 25401 linearc O. Don | linear | HWEJcu | mr.au Br | Japan 1822 | D p.l Thun. Jap. 1° |
| Polyptodium line^rc L. | | | | | |
| | 2912. ONOCLESVA L. | ONOCLBA. (Onos, a vessel, kleio, to enclose; apparent capsules.) 2. | | | |
| 25402 sensibili^i. | sensitive | ^ A or 1) au | Br | Virginia 1799. | I) Lp Schk. fil. 102 |
| Germfimica IV. | obtuse-lobed | S A or 1 jl | Br | N. Amer. 1812. | D l.p Schk. «L IO* |
| n n, G'ndnd <i>StruthWpteris L.</i> | | | | | |
| 25405 pennaylinica IV | Pennsylvanian | £ A or 2 an | Br | N. Amer. 1812. | D i p Schk. fil. III |

2914. ALLOSO^XRUS *Bernh.* ALLOSORUS. (*Alias, various, torus, heap; sort changing at different period**.) 1.-4.
2540S crisp^uB *Bernh.* curled cu i jLau lit Britain sto.hi. 1) l.p Eng. bot. 1160 /teris crfspa Sivz. OsmGnda crlspa L.

2915. ELLOBOCA'RPUS *Kaulf.* POD FEE V. (En. in, lobos, apod, karpos, fruit; div. of frond.) 1.—2.
25407 olerucus *Kaulf.* potherb £ fzs or 1J au Br Tranqueb. 1818. D l.p Pluk. al. 215. S Ceropt6pterw /halictroides Broeg.

2916. PAHKEMI/J Hook. PARKERIA. (C. S. Parker, who first found the plant in Esicquibo.) j.

25408 ptcTo\dcB Hook. Pteris-like ^ E] or J au Br Essequibo 1825. D l.p Hook.ex.fl.H7.4H

⁴ 2917. LOM A'RI A JV. LOMARIA (*Loma, an edge; marginal position of indusia.*) 5.-43.

25409 ^pfalcata Sp^r *falcata* R. Br. falcate' £ IAJOT 2 my.au ^"vISi. 1823. D pl Stegania falcata R. Br. Making. £ iAI or 2 my.au Br V.D.I. 1822. D pl Lab. n. h. 2.218 Stegania nuda R. Br. ftnoclea niida 7,a6. 25411 nAcant Desv. spike £ ^ or 1 jl Br Britain heaths. D pl Eng. but. 1159 Jlechnum bore'ile Sm. Suu. 25412 procnd Spr. tall U.VwsTJL-Sterile fronds pinnate; leaflets serrulated. 25413 Stegania procera R. Br. Osmfmda procera Forst. long-leaved JKEjor 2 jn.jl Br W. Indies 1810. D l.p Pl.fil.117.rfc.rf/. Acristicum torbifolium Vahl

• 2918. BLE'CHNUM L. BLBCITVUU. (*Bleehnon, one of the Greek names for a fern.*) 10 — ;i.

I. INTEGE'REIMA. — Fronds pinnated; leaves entire.
25414 austral^e L. southern £ IAJpr 5mr.s Br C. G. H. KJ01. R ip Schk.fll. 110. b.
25415 occident^e L. western jj CS3>r 1 mr.s Br S. Amer. 1777. 11 i.p Jac. ic. 3. 641
85416 glandulbhium Lk. glandilious i? C3 or } mr.my Br Brazil 182 i. II np
25417 pectinatum Presl >cctinated £ 123 or 1 ... Br S. Amer. 1827. R sp
25418 longitatum Hum. long-leaved £ El or 1 mr.s Br Caracas 18'0. R sp Bot mug. 28IH

II. DBNTICULA'TA. — Fronds pinnate; leaflets serrulated, crenulated, or denticulated.

25419 hevigatum Cav. smooth jg 1Alor 1 mr.s Br N. Holl. BLM. R s.p
25420 serrulatum Mx. serrulated £V A or f mr.s Br Florida 1818. R np
25421 stratum /(. Br. striated fl. iAi or { mr.s Br N. Holl. 1824. R sp stramneum Lab.
25422 brasiliote Desv. Brazilian jg El or mr.s Br Brazil 1820. R s.p
25423 cartilagineum Swx. cartilaginous tf 1Alor 1 jn.au Br N. Holl. 1820. R s.p

2919. WOODWA'R'D/f Sm. WOODWARDIA. (*Thomas Jenkinson Woodward, an English botanist.*) 1.

2.H24angustifolia Sm. narrow-leaved * A or 1 au Br N. Amer. 1812. I) p.l Schk.fll. III onoctorides IV. Osmfimla caroliniXna Walt. rootinp/cnwr. #, IAJ or 1 s llr Madeira 1779. R sp Schk.fll. 112 25424'i radicans Sivz. Virginian £ A or 1 au.s Br N. Amer. 1774. D l.p Pluk. al. 179 2 ifl'chnum virginicum L.

2920. DOO D7if R. Br. DOODIA. (*Samuel Doodt*, a London apothecary, and first Brit cryptogamist.) 3.
25427 aspera R. Rr. miffr-stalkcd f uM br } mr.s Br N. S. W. 1S08. R «p Bot.cab.3!)
£428 media R. Br intermediate £ uj or \$ mr.s Br N. Holl. 182J. 1) l.p
25429 caudata R. Br. tailed ^ iAl or 1 mr.s Br N. Holl. 1820. D l.p Hook. ex. fl. 25 Woodfirdia caudata Cav.

*2D21. ifSPLETNIUM L. SPLEENWORT. (/, priv., splen, the spleen; mod. qual.) 47. — 154.

I. INDIVIA. — Fronds undivided.
25430 Nidus L. JWrf* Nest K Indies 1820. l.p Brey. c. 99
*^>1 brasiliense Sun. Brazilian Brazil 18^>. l.p
*^>1 rhizophyllum L. taxi-leaved W. Indies 18'S. l.p Snhk. III t'A
attenuatum rontiiK-leaved N. Amer. 18^>. l.p l'luk. al. 105.3
attenuatcll N. Holl. 18'.k l.p

II. LODATIFR6NDKS. — Fronds angular, lobate, divided.
25431 p^amatum Lam. palmate jf iAi or { aus. Br S. Europe IHlf. I) l.p Bnt. cab. 8f8
^>*³⁶ septentrionUc Sun. northern 3t A or > jn.o Br Britain l.p Eng. bot 1017

III. PINNATIFROVDES. — Fronds pinnaW.
i. SHIRINTERPOLIUM. — Leaves subentire.

25432 angustifblum Mx. narrow-leaved ^ A ^ 1 jn.jl Br N. Amer. 1812. D l.p Schk.fll.C7. fi9
*4jf HhcphfaUJ apr. Knif-leaved Br W. Indies l.p PI. III. & 1
^>40 obtiiHritum Forst. Shepherd's obtuse Br Jamaica 1800. l.p
ob n i Lab. marnum L. one-floweret £ A or iAl cu Br Britain rocks. D s.p Eng. bot. W
mopanum Sm. falcatum Lam. falcate If 1Alor Br N. Holl. 18'S. l.p Sm. ic. in. 73
^> njeifMium IV. Zamia-leaved f EJcu 1 Jnjl Br Caracas HLO 1) Bot. cab. &4
M<>ctfum IV. biparted i A or 14 jug Br JnnailM 1820. l.p Schk.fll.73
S >cpnum II. K. ebony-stalked ic rXior 14 jn.jl Br N. Amer. 177>. 1) W. Indies 18'2.
formifnum IK. beautiful ^ A Pr 1 Jns Br Irritaiii sh.roc. l.p Eng. bot 57fi
TV afi L. Maiden hair h A br l. Br N. Amer. 1812. D
black-stalked £ A Pr f jn. Br Britain al. roc. D
Petrarch* Dec. Petrarch's l. Br France 1819. D Eng. bot. 2207

VilliB-elafisa; Req. toothed or 4 jn.jl Br W. Indies 1820. l.p
toothed or i jn.jl Br N. Holl. IRL0 l.p Hook.ex. fl S08

flabelliMium r*si*. fiin-leaved \$ EJ pr 1 an Jfr Jamaica 1793. l.p SL jam. 29.30.1
H rhiziphorum //. root-bearing g A cu > jn.o Br Scotland ... l.p Eng. bot. 2258
H alternumSm. germanicum It'eis

IV. BIPINNATI'FIDA.—*Fronds bipinnatifida.*

S5456	pbumilum <i>Sivz.</i>	dwarf	£ El pr	* J ⁿ J ^l Br	W. Indies 1823.	D l.p	PL fil. 66. A.
25457	diffirme <i>R. Br.</i>	two-formed	£ LAI or	J jn jl ^ A el	Br N. Holl. 1823.	D p.i	
25458	fontanuin <i>R. Br.</i>	fountain	£ El or	f jn.au	Br England w. &c r. D l.p	Eng. bot. 2024	
	Aspidiuu futitatum	IV. /olypodium fontanum L.					
25459	birtctuin <i>Swz.</i>	binected	£ El or	1) au	Br Jamaica 1821.	D l.p	
25460	praunonum <i>Sun.</i>	bittcnUeaved	£ E) e1	} au	Br Jamaica 1793.	K i p	Pluk. al. 73.5
25461	itnatum <i>L.</i>	striated	£ El pr	1 jn.au	Br W. Indies 1793.	It s.p	PL fil. 18.1!
25462	flicypteroides <i>Mx.</i>	Thelypteris-like	S A or	1 Jl	Br N. Amer. 1823.	D p.i	Schk.fil.7a6
25463	acrosticho'ides <i>Sun.</i>	rooting	£ El or	I jn.au	Br W. Indies 182a	D p.i	Schk.cryp.76
	rhizophoruin <i>Schk.</i>						

V. BIPIVNAVA.—*Fronds biminate.*

251C4H4llerri.R.Br.	HallerVi	* A or	1 jn.au	Br	Switzerl. 1819.	D p.i	Schk. cryp. 53
	Aapidium lontanum <i>Sun.</i>	Halleri W.	Athyrium Halleri	Roth			
25465	F\lx.fa/mina. Hernh	Female Fern	^ A or	2 jns	Br	Britain w.sh pi. D l.p	Eng. bot. 1453
	Polypodium niiix-fin'mina L.						
25466	Athy'riu Spr.	Athynum i A or	2 Jns	Br	N. Amer. 1823.	D p.i	Schk. cryp. 78
	Nephrediu flipnclioidi* Mx.	Aspfrium nupncloides W.					
25467	Michadixii Spr.	Michaux'u & A or	If jns	Br	N. Amer. 1823.	D p.i	
	Nephrediu /-iix-flu inina Mx.	Aapidium angibuum IV.					
25468	lanceoUtum Hud.	lanceolate	£ A pr	i J ⁿ J ^l	Br	England rocks. D l.p	Eng. bot. 240
25469	Hhia muraria L.	Wall Hue 5 A cu	4 Jn.o	Br	Britain sh.roc. D l.p	Eng. bot. 150	
25470	bulhferuin Ford.	bulb-bearing	£ tAl or	1 Jns	Br	N. Zeal. 1820.	D p.i
25471	viviparum // K.	viviparous	£ El or	1 Jns	Br	Mauritius IM20.	D p.i
25472	fragrant! Sun.	fragrant	£ El cl	f Jl	Br	Jamaica 1793.	D l.p
25473	/di&ntum ii^rum L.	Bk Adiantum^ A P'	I ^P-0	Br	Britain sha.pl. D l.p	Eng. bot 1950	
25474	mont'inum W.	mountain ik A pr	i jl	Br	N. Amer. 1812.	D l.p	
25475	cicutanum Sivz.	Cicuta-like	£Elor	1 Jns	Br	W. Indies 1820.	D p.i
						PI. fil. 48.a.	

V. TRIPLOCA^TO-PINNAVUM.—*Leaves triplicately pinnate.*

25476	acutum Bory	acute	£ iAJ or	2 ap.my	Br	Tcenenftb 1818.	D l.p
-------	-------------	-------	----------	---------	----	-----------------	-------

2922. CJENO'PTERIS Berg. CAWOPTERIS. (*Kain'os*, new, pteris, a fern.) 2.—13.

25477	Odontitis Thun.	Odonfites	£ tAl or	f jn.au	Br	N. Holl. 1822.	D p.i
	ifspfemum Odontitw H. Br.	Darl-a Cjdonfes W.					
25478	appendiculatum Lab.	appendaged £ LAJ or	# Jl.au	Br	N. Holl. 1822.	D p.i	Lab. n. h. 2.243

ifspfemum l&xum R. Br. Darca appcndiculata W.

2923.	ALLANTO%DIA B. Br.	ALLANTODIA.	{Allantos, a sausage: resemblance in industia.)	4.-5.			
25479	tñera R. Br.	tender	£ IAJ or	ll jns	Br	N. Holl. 1820.	D p.i
S5410	auNtralis R Br.	southern	£ IAI	2 Jns	Br	V. D. L. 1820.	D p.i
25481	axilTirif Kaulf.	axillary	£ IAJor	2 jns	Br	Madeira 1779.	D l.p
	PolypTidium axillire // K.						
23482	umbrosa R. Br.	Hhtdy	£	4 Jns	Br	Madeira 1779.	D l.p
	umbrfusin II. K.					Schk. fil. til	

2P24. SCOLEPO'KDUIUM Sm. HARTH TONOU. (*Scolopendra*, a centipede; underside of fronds.) 2—4.

25483	offlcinrum Swz.	officinal	■ A cu	liJl.au	Br	Britain m.B.pl. D l.p	Eng. but 1150
	ilapW-muin 5coloi>diidriuin L.						
2	criHiuum	cuTe&leaved	■ A cu	Jlau	Br	Britain ms.pl. D l.p	
3	undulatum	wave leaved	■ A cu	M Jlau	Br	Britain ms.pl. 1) l.p	Pluk. ph. 248.2
4	multifidum	many-clrt	■ A cu	H Jlau	Br	Britain ms.pl. 1) l.p	
5	ramoflum	brunchy	■ A cu	ll Jl.au	Br	Britain ms.pl. 1) l.p	Pluk. ph. 248.1
25484	Hemionitis Cav.	llemionitis	■ Jl or	1 aus.s	Br	Spain 1779.	1) p.i
	//einonlti* L. palmatum BTC.					Schk. cryp. 66	

2995. DIPLA^ZIUM Swz. DIFLAZIUM. (*Diplaxo*, to be double; indium double.) 9.—2aI. SIMPJCPR<NDfEa.—*Fronds simple.*

£5485	priantaglneum&t0z. Vlmta\n4eaved£ El or	2 au	Br	W. Indies 1819.	D i p
	ifspfemum plantaglneum L.				

25486	castanca?>lium Swz.	ChestnuUvd	£ El or	1 Jl.au	Br	Guinea 1824.	D l.p
-------	---------------------	------------	---------	---------	----	--------------	-------

11. PINNATIFR6NDES.—*Fronds pinnate.*

25487	scramporense Spr.	Scramporc	£ EJ or	3 ail	Br	Scramporcl820.	D l.p
25488	grandif Mium Sun.	large-leaved	£ El or	4 au	Br	Jamaica 1793.	D l.p
	//mon'itiM grandifflia Sun.						
25490	uglandiflium Swz.	Walnut-leaved	£ El or	3 au	Br	Jamaica 1820.	D l.p

25490 aurirul/atum Kautf. auricled ± CJ or 10 au Br Caraccas 1820. D l.p Schk. cryp. 85

111. BIPINNAVA.—*Fronds btpiwiate.*

25491	malabfcum>Vor.	Malabar	f • or	8 ...	Br	E. Indies 1818.	S l.p
	jfpfenl ambtgum Sun.						
25442	arbo'recens Viz.	arborescent	£	or 12	■	Mauritius 1826.	S l.p

25493 cfculentum Sivz. eatable

//cmionitis esculenta Retx.

• 2926. PTITRIS L. BRAKE. {Pteryx, a wing; appearance..) 15.—119.

25494	sagittafblia Rod.	sagittate-leaved	£ El or	I Jn.au	Br	Brazil 1825.	D p.i
-------	-------------------	------------------	---------	---------	----	--------------	-------

II. DiviNuB.—*Fronds divided.*

25495	peiUUL.	pedate	k A or	f Jl au	Br	Virginia 1820.	D l.p
25496	palmata W.	palmate	£ El or	f jn.au	Br	Caraccas 1821.	D l.p

III. TERNA\A.—*Fronds ternate.*

25497	arg^nte Gm. pedata var. L.	• livery	k A or	f jn.au	Br	Siberia 1816.	D p.i
-------	----------------------------	----------	--------	---------	----	---------------	-------

N. c. p. IS. I

IV. SIMPLICIPINNA TJE.—*Fronds simply pinnate.*

fSJOJrotundlfMlaFor*/.	round-leaved	£ El or	1) Jlau	Br	N. Zeal. 1824.	D p.i	
	long-leaved	£ El or	2 jl s	Br	W. Indies 1770.	Sk &p Jac. sc.3.399,400	
	falcate	£ IAI or	1 jn.jl	Br	N. Holl. 1820.	D p.i	
25501	grandifbia L.	large-leaved	£ El o r	2 au	Br	W. Indies 1793.	Sk p Schk.fil.89

- V. CoMP^{*}SaiTopiNVA^NrJi. — *Froonds pinnate; lower leaves compound.*
- 25502 denticulate *Hook.* toothed ^ [Q or S] ll Br Brazil 1824. Sk s.p
 25503 crdtica *L.* Cretan £ tA cu 1 jl.au Br Candia IH20. I) l.p Schk. cryp. 90
 88504 serrulata *L.* serrulated £ EJ pr 1 au.s Br India 1770. Sk s.p Schk. fil 91
 25505 umbrtaa *R. Br.* shady £ L&[or 3 jn.au Br N. Amer. 1870. D p.p Schk. fl. 101
 25506 atropurpiarea *L.* purple. 5 A pr i au.s Br N. Amer. 1870. D p.p Schk. fl. 101
- VI. BIPIVNATI'FIDA. — *Froonds bimarginatid; lower leaves 1/2.* D l.p
- 88507 Plumierfi *W.* Plumier's £ (23 or 2 jl Br S. Amer. 1818. D l.p
 88508 ncmorinlia *W.* grove £ El or 1 my.jl Br Bourbon 1823. D l.p Pet. g. 80.3
 25309 arguta /*^{*}/ tharp-notched £ iAi el 1 au.a Br Madeira 1778. D l.p Huk. «d. 2^oJQ. %
- VII. BIPINNAVA. — *Froonds bipinnate.*
- 25510 hastate *Sun.* hastate £ IAJor 2 Jl Br C.G. H. 1883. D l.p Pluk. ph. 403. 5
 15511 cordata *Cav.* heart-shaped £ (23 or 3 Jn.jl Br Mexico 1880. D p.l
 88512 sagittata *Cav.* arrow-shaped £ EJor 3 Jl Br S. Amer. 1826. D p.l
 88513 f legana *Pair.* elegant £ El or 3 au.s Br E. Indies 1824. D l.p
 25514 discolor *MIX?^{*} elegans Swz. discolored - 3 au.s Br Brazil 1825. D l.p
- 88510 heterophylla *Suiz.* variable-leaved £ [P] or 4 Jlau Br Jamaica 1820. D l.p Pl. fl. 37
 88516 aculeate *W.* prickly-rfmd £ [P] or 10 au.fl Br W. Indies 1793. D l.p Pl. tiL5. etll
- VIII. TRIMCA^{*}ro.nsitA^{*}TM. — *Froonds triplicately pinnated.*
- 25517 vespertilibnis *Lab.* bat-iiMit/pif tf tAJ or 3 my.au Br N. Holt. 1823. D l.p Lab.n.h.2.845
 25518 tremula *R. Br.* trembling 2 i£ or 3 my.au Br N. Holl. 1820. I) l.p
 25519 esculenta *Font.* eatable £ L&J or 3 au.s Br N. S. W. 1815. D l.p Lab.n.h.2.844
- IX. DECOMPOSITAC. — *Froonds ilecompound.*
- 25520 caudata *L.* tailed k A Br 2 s.d Br N. Amer. 1777. D l.p Jac. lc. 3.615
 88521 aquilina /.. aquiline ^ A or 3 Jl.au Br Britain hea.w. D l.p Eng. but 1679
 88522 leptophylla *Sua.* slender-leaved f [S or 2 Jl.au Br Brazil 1824. I) 1 p
 25523 Unuguia *Itory* woolly £ E3 or 3 Jl.au Br Bourbon 1819. D l.p
- X. DU^NBIJS. — *Doubtful to which section they belong.*
- 355>4 chintasia *Lo. C.* Chinese £ IAJ or 2 Jl.au Br China 1824. D p.l
 88525 crenulata *Lo. C.* circnulated f iAi or 2 Jl Br 1827. I) p.l
 25526 Cervantes^{*} H. L. Cervantes^{*} £ tA] or 1 jl Br Mexico 1824. D p.l
8927. VITTAMUA Sm. VITTARIA. (Vitta. a riband: shape of narrow fronds.) 2.—10.
- 2^o527 lincata *W.* YuicvrUeaved £ El cu 2 au Br America 179L D Lp Schk. fil 101. b
 88528 #ramiuifolia *Kaulf.* Grats-leavcd £ El or 1 jl.au Br Brazil 1820. D Lp
2028. LONCHIPTIS L. LONCHITIS. (Louche, a lance; form of fronds.) 8. — 5.
- 2VV29 himita *L.* hairy £ El pr 1 jn.s Br W. Indies 1793. O l.p Schk. fl. 86
 88530 pedluta *L.* pedate £ EJ or 1 jn.jl Br Jamaica 1793. D l.p Br. jam. 89.1
 Plbus podophyilla Swz.
8929. ANTROPHYUM. Ktiuf. ANTROPHYUM. (Antron, a cavern, phyto^o to grow; habitation.) I. — 8.
 88531 lancollitui Kaulf. spear-tawed £ [■] f jls Br W Indies 1793. D Lp Schk. fl. 6 //emionitis lanceofate *L.*
- * 8930. ADI A'NTUM L. MAIDENUAIB. (Adiantos, dry; plunged in water, yet remaining dry.) 20. — 63.
- I. UKNIPORMK. — *Leaves simple, rrntfonn.*
- 88532 reniforme *L.* ktdney-leaved £ iAi I* \$ jn.s Br Madeira 1699. II s.p Bot. cab. 841
- II. SIMPLICIPINNA'TA. — *Fionds simply pinnate.*
- 25533 macropli^Illum Swz. long-leaved 1 Jlau Br Jamaica 1793. I) l.p Br.jam.38.1
 25534 lunktum *Cav.* lunated 1 Jl Br Mexico 1823. D l.p
- III. COMPOSITO-PINNA^NTA. — *Froonds at base compound or bipinna/e, above simply pinnate.*
- * »55 deitoidcum Swz. deltoid £ El or 1 jl au Br Jamaica 1824. I) l.p
 f5. h.i(i Varium *Hum.* various £ El or 1 jl.au Br S. Amer. IK20. I) l.p
 * 5537 scrriatum L. serrulate £ El or 1 au Br Jamaica 1822. D l.p Pluk. aL 185.2
- IV. DIGITAVO-PIVNAVA — *Froonds triparted, digitate* or pedate; branches pinnate.*
- ^»38 ternatum 7/iim. ternate £ El or 1 jn.s Br S. Amor. IH19. I) l.p
 ^2. radiatum L. radiated £ iAI pr J ap.au Dr W. 1 mil^o 1776. D Lp Pl. filIOU
 * »41 pulverulentum L. pedate £ r. 1 au.s Br N. Amur. IK40 H s.p Schk. III. 115
 ^H patens F. spreading £ El or 1 au.s Br Brazil 1821. I) s.p
 ■ iiispidulum It Br. rather hairy £ -IAJor 1 iny.au Br N. Holl. 1822. I) l.p
- V BiPistia'r. — *Fionds all bipinnate.*
- ?M3 villsum /.. haWy-stalked £ El or 1 jns Br Jamaica 1775. D t.p Schk. fl. 120
 jij41 pulverulentum L. dusty £ E iJ 1 UJHB Br W. Indies 1793. D s.p Schk.fl. 119
 rhombochium Kth. rhomboid f El or 1 J" Br S. Amer. 1820. D l.p
 16 CapitulK Vcneri* L. Venus's hair £ iAJ cl \$ my.a Br Britain rocks. R s.p Eng. bot 1564
- VI. SUPRADKOMP6SITA. — *Froond triplicately pinnate or supradecotnpound.*
- cuncMum Hook. wedKCsha)ed £ iAJ or 1 au Br Brazil 1820. D p.l H. & G. 30
 Thomb-Jeaved £ tZ3el 1 jii,jl Br W. Indies 1793. It s.p Schk. fl. 112
 itsalmile Swx. tender £ EieI 1 Jl Br Jamaica 1793. D l.p Phlk. al. 354.1
 trigbnium Lab. assimilated £ iAI or 1 my.au Br N. Holl. 1823. D Lp Lab. n. h2jMft
 formbsum R. Br. handsome £ IAJ or 1 my.au Br N. Holl. 182a D Lp
- 893L LI NDS JEM Dry. LINDHAA. (Lindsay, an Englishman, who wrote on germination of mosses.) 5.—89.
- I. PINNA^Nr. — *Froonds pinnate.*
- linear tf iAI or \$ ap.jn Br N. Holl. 1820. D p.l
 tickleshaned £ El or } apjn Br Trinidad 1819. D p.l Lm.tr. a 7.8
- II. DECOMPOSITE. — *Froonds decompond.*
- ~*54 microphylla Stvz. small,lcaved £ AJ or { apjn Br N. Holl. 1820. D p.i Lin. tr. 3.9
 5 S 5 ^oHzaBf'.rmis Dry. Trapiuum-findtf El or 1 apjn Br S. Amer. 1819. D p.i Lin. tr. 3.9
 midia R. Br. Intermediate £ uQJ or 1 apjn Br N. Holl. 1823. D p.i
- * 2932. CHEILA'NTHES Sun. CHIBILANTHES. (Cheilot, lip, anthos, flower; form of induium.) 10.-38.
- I. BIPIVNATI'pinA. — *Froonds bipinnatifld.*
- «^o57 Kr.'icilis Kautf. slemler k & or # jl Br N. Amer. 1823. D p.l
 Pferis grucilis MJC.

9V ? r?	atomed Rhartian royal bulbiforous	A A or 5 A el 5 A el S A or	Br Br Br Br	N. Amer. 1820. D Britain rocks. D Britain al.roc. D N. Amer. 1638. D	I.p I.p I.p I.p
SSL	iferum Swx.	<i>tin</i> 1 jl.au			Eng.botISS
V.	SUPRADUCOMPOSITA.				
	anularc <i>Kit.</i>				
" "	angular				
" "	pubescent				
JW in Si w.	villous				
ae'inulmn <i>Sun.</i>	rival	1 Alor			
° J uecompöhithum Sjw.	decomposed	1 AJor			
	^ciirriidium decompositum 7L				
SAWS	alpmc	jnjl	Br	S. Europe 1825. D	I.p Jac. ic. 3.612
** <` montanum Suo.	mountain	my.au	Br	SwiUerl. 1819. D	I.p Vil.deL3.53
PLEOPE'LTIIS Ftfm. PLEOPBLTIS. (P/eoi, full./Wte, shield; form of indusium.) S.—8.					
SH12 f" '£: ^{ha} / ^{0/0*}	word,leaved	* EJor	ft my.jn	Tir	S. Amer. 1821. D I.p Hook. ex.fl. 62
2f ^{2*} latifoha /A £	broad-leaved	tf El or	J my.jii	Br	S. Amer. 1823. D I.p
W nhda D. <i>Don</i>	naked	j & EJor	my.jn	Br	Nepal Hoot ex.fl. 63
*JJ40. WOO'DS/^R.Br. WoonsiA.					
5S ¹ P ¹	pubescent	...	Br	Brazil 1820. 1)	I.p
hyperbrca H. Br.	northern	4)1.1	Br	Scotland al. roc. D	I.p Eng.bot.2023
Ilva		i Jnjl	Br	Britain 1812. D I.p	Eng. bot & 2b ² li
CYATHEVA Sm.	CYATHIBA.				
arbbra Sm.	tree				
exculta Swx.	tall				
arbbra Bory					
Woods, a British botanist) 3.—4.					
TRICHCM ANES L. TRICHOMAWM. (<i>Thrit</i> , a hair, <i>manos</i> , soft; stems like fine hnlr.) 8. — 46.					
revisione H. K. short-bmtcd	£ A el	i my.jn	Br	Britain moi.ro. D	I.p Eng. bot. 1417
Hymenoph Hym allMum £ A f					
mcmbran&ceum /., membranous	fc E3 or	ft my.jn	Br	W. Indies 1820. D	I.p Hook. ex.fl. 76
HYMENOPHY'LLUM Sm. FILMY LEAP. (<i>Hymen</i> , membrane, <i>phyUn</i> % leaf: thin leaves.) 1.—4(1					
tunbrugl^nse 5un. Tunbridge	£ A cl	4 my.jn	Jlr	Britain moi.ro. D	I.p Eng. bot. 162

TBIBEJL: QSMUNDRATE-JE-

<i>T. TCPDEA</i> W. TODEA. (<i>Henry Julius Tode</i> , of Mcklenberg, an experienced mycologist) 1. africtaa W. African £ 1AI or 2 my.au Br C. G. II. 1805. D Lp Sent fl. 147 Oimümla barbara Thun.
L. OHMUNDA. (<i>Osmunder</i> , one of the names of Thor, a Celtic divinity.) &—8.
I. CINNAMOMIA.— <i>Bearing both fertile and sterile fronds.</i>
<i>cinnamumca</i> L. Cinnamon ^ A or 2 jn Br N. Amer. 1772. D Lp Schk. fl. 146
II. REGALES.— <i>Fronds all fertile.</i>
& reRMisL royal 2 jl.au Br Britain sha.bo. D Lp Eng. bot. 20!)* *Jl*8 ClaytonMM I* S i "Aer6ruu J f. spect&bili W. 85645 gracii slender ^ A or 1 Jn.jl Br 1827. D p.l
Clayton's 2 au Br N. Amer. 1772. D Lp interrupted 2 jn.jl Br N. Amer. ... 1) Lp Schk. (11.144 showy • 2 jl Br N. Amer. 1811. D Lp Pluk.al.184.4

TRIBE III. GLEICHENIEM.

				(<i>Lygodium</i>)* flexible; twining plants.)	6.-23
oAndens Swx. Drit. & l. in Sum.	climbing circinate	I. E j el i E.jel	6 mys fi au	Br E. Indies 17M 1) l.p	Bot cab. 742
PalmMum Swx.	palmate		6 ou.s	Br N. Amer. ... D.lp	Ru am. 633
polymfirphum <i>Kth.</i>	multiform		6 au	Br S. Amcr. 1820. D Lp	Schk. fl. 140
Hynrogh'tMum hirsutum W.					Cav. 1c. 0.595.1
vulchiae Stiff.	twining				
P^am^nda sc&ndens Aub.					
hastitum ft.	hastate	JL El or	6 au	Br Maranh. 1820. D Lp	
	ANEMIA,			(<i>Anemone</i> , naked j naked spikes of inflorescence.)	5.— 27.
	hairy	£ E! or	3 ...	Br Jamaica 1794	D Lp
Giantifolia Sun.	Maidenhair.lvdv.	El or	■ aus.	Br W. Indies 1793. D p.l	PI. fl. 1<2
	Phillitis-like	£ El or	■ JLo	Trinidad 1820. D Lp	PI. fl. 158
	L. la Phillitidis				PI. fl. 156
	Hook.	lanceolate	£ E l o r	Jl.o	Br W. Indies 1820. 1) p.l
	dwarf	5 A or		Br M. Amer. 1823. D p.l	Hook. ex. fl. 28
	va humilis Cav.				
infiblia Hook.	Ash.lcaved	£ El or	■	Br Brazil 1825. D p.l	
				(<i>Schizophragma</i> , to cleave, appearance of aggregate fan-like spikes.) 4—12.	
	elegant	£ El or	■ S. smit	Hr Trinidair. 1H19. 1) p.l	Vahl sy. 2.50
Hum.	pencilled	£ E)	or f	Br S. Amcr. 1816. 1) p.l	
	rock	£ IAJ or f	■ 31	Br N. Holl. 1822. 1) p.l	
>ua swx.	bifid	£ LA) or	Br N. Holl. 1822. 1) pi	W.m.far. 3.2	
>u Hla Ph.	small	k ^ J or	Br N. Amcr. ... D p.l	H.&G.47	
OLEICHETNW Sm.	GLEICHENIA.			(<i>Baron P. F. von Gleichen</i> , a German botanist.)	5—28.
ub6bcnsX/A.	pubescent		3 Jn.au	Br S. Amer. 1822. 1) p.l	
pelfincip It. Itr.	cavern	i Alor	jn.au	Br N. Holl. 1KVA D p.l	
inichernhylo JL Br.	small-leaved		jn.au	Br N. Holl. 1824. 1) p.l	
flabellata R. Br.	fan-.caved		jn au	Br N. Holl. 1823. 1) n.l	
	pectinated	bm or	3 au	Br Trinidad 1S24. ■ Ac. hoi. 1804.4	

TRIBE IV. OFHIOGLCSSEJE.

- | | | | |
|-------------------------------|-------------------------------|---|--------|
| 2950. BOTRY'CHIUM | Sua. MOONWORT. | (<i>Botrys</i> , a bunch of grapes; form of fructification.) | 6.—10. |
| 25668 <i>Lunaria Stux.</i> | common, Moonwort [^] | A cu ft my.jn Br Britain hil.pa. D p.l Eng. bot 318 | |
| Osmundrt Lunaria L. | | | |
| 25669 <i>aUitrMe R Br.</i> | southern | fk 1AJ ^{or} i my.jl Br N. Holl. 1823. D p.l | |
| 25n70/ <i>umarioides</i> IV. | Fumitory-like | * A cu ftjl.ju Br Carolina 1806. D p.l Schk. fil. 157 | |
| 25071 <i>dissectum</i> 5 | cut-leaved | 5 A cu * ji Br N. Amer. 1806. D p.l Schk. fil. 158 | |
| gr&cle Ph. | | | |
| 25672 <i>virKincuin</i> Swx. | Virginian | ‡ A cu 1 au Br N. Amer. 1790. D p.l Schk. fil. 156 | |
| 25613 <i>obliquum</i> Mhl. | oblique | ‡ A cu ft au Br N. Amer. 1821. D p.l | |
| 2951. OPHIOGLO'SSUM L. | ADDER'S TONGUE | (<i>Ophis</i> , a serpent, <i>tfoassa</i> , a tongue; leaves.) | 7.—12. |
| 2ID74 <i>vulgatum</i> L. | common | & A cu my.Jn Br Britain' m. me. I) p.l Eng. bot. 108 | |
| SIJSU <i>ptciolatum</i> Hook. | Mlti:latcd | SE)cu Jja.mr Br Jamaica 1820. D p.l Hook. ex. fl.56 | |
| 25n75 <i>reticulatum</i> L. | netted | ‡ [SI pr i my.jn Ur W. Indies 1793. D p.l H.&G.26 | |
| VTriffi <i>bultmsum</i> Mr. | bulbous | A Al pr \jlau Br N. Amer. ... D l.p | |
| &nH7 luHifuiicium L. | FortKil | ‡ A cu , my.jn Br Portugal 1816. D p.l | |
| QTAYW "ramineum" IV. | GrmtKzAtr | fct LAJ cu f my.jn Br N Holl. 1820. D p.l | |
| 2S8U8 costikuum It ttr | ribbed | 3t IAJ cu f my.jn Isr N. Holl. 1820. D p.l W. erf. 1802.1.1 | |

TRIBE V. POItOPTE'HIDES.

2952. MARA'TTM Sm. MARATTIA. (*J. F. Marati*, of Vallombrosa, in Tuscany, who wrote on ferns.) 1-5.
25681 alaU Sm. • winged £ (Z3 or lft au Br Jamaica 1/93. D l.p. Sm.ic. in. 46

CLASS XXIV. ORDER 2.

CRYPTOGAMIA EQUISSETATE-ffi.

- | | | | | |
|----------------------|----|--------------------|--|----------------|
| •2954. A'QUISETTUM | L. | HOHSETAIL. | <i>(Equus, a horse, adeta, hair; fine branches.)</i> | 10.—18. |
| 85683 prathsef ArA. | | I. POLYSTAYHVA.— | <i>Fructiferous fronds many-spiked.</i> | |
| | | meadow | ^t A cu 1 jn.jl Br (icnn.my 1816. D pi | |
| | | marsh | ifc A w l Jn.jl Br Britain bog.pi. D p.l | Eng. bot. 2021 |
| 25645 umbrblflum W. | | II. MOVOSTA-HIVA.— | <i>Fructiferous fronds onspiked-</i> | |
| SHWG sylvanticum L. | | sliady | cu 1 Jujl Br Germany 1816. D p.l | |
| 256K7 hymale l. | | wooti | w 1 ap.my Br Britain m.s.pi. 1) p.l | Eng. bot. 1874 |
| Wm Intontum L. | | winter | pc l jl.au Br Britain moi.w. D p.l | Eng. bot. 915 |
| urmii fluviatilic L. | | nvirsh | w 2 Ju jl Br Britain wat.pl. I) p.l | Eng. bot. 929 |
| 25UM arvensis L. | | great-river | w d ap.my Br Britain wat.pl. I) p.l | Eng. bot. 2022 |
| 2DAH variegatum W. | | corn | w i mr.ap Br Britain moi.f. 1) p.l | Eng. bot. 2020 |
| 2GU2 acirpoide* Mx. | | variegated | or f jn jl Br Scotland sc.sh. I) p.l | Eng. bot. 1987 |
| | | Scirpus-like | cu ft jn jl Br N. Ajaer. 1820. D p.l | |

CLASS XXIV. ORDER 3

CRYPTOGAMIA LYCOPORI'NEJE-

- | | |
|---|----------|
| •2955. LYCOPODIUM L. CLUB Moss. (<i>Lykos, wolf, pouss, ftiot; resemblance in roots.</i>) | 18.—140. |
| 1. PEDUNCULAR.— <i>Spiked, pedunculate.</i> | |
| 2Mm caro?ini?num L Carolina t a cu { au Br Carolina 1812. D pi Di.muse.62. | |
| 25694 clavatum L common-club % A cu il:tu Br Britain hca. D pl Eng. bot. 224 | |

- | | | | | | | |
|---|---------------|--------------------------------|---------|-----------------------|-------|--------------------|
| 25&J5 complanatum L. | smooth(J/M/ | £ A cu \ jl.au | Br | N. Amer. 1770. | D pl | Fl.dan.78 |
| II. SBHSTLIA.— <i>Spiwls sessile.</i> | | | | | | |
| i. VA ^N HIA —Leaves in various directions. | | | | | | |
| 95R9R alpinum L. | alpine | tU. A cu i n u | Br | Britain albngs. | D pl | Eng. bot. 234 |
| 250J7 dendrofdeum Mx. | tree-like | £ A cu | ■ | Br N. Amer. 1770. | I) pl | Hook.ex.fi. 7 |
| SdJ98 dñpñnum Lab. | dense | £. A <u>J</u> Cu | £ au | Br N. Holl. 18'Jl. | I) pl | I-tb. n. h.2.251.1 |
| 85(FX) annutinum L. | year-old | JU <u>A</u> cu | I jn.au | Br Brit.'in nl.hca. 1 | I) pi | Kng. bot. 1727 |
| 257(X) inundhtum L. | overflowed | JU A cu | * jnjl | Br Britain turbo. I) | I) pl | Eng. bot &> |
| 25701 Alopecurfl's L. | Foxtail-hkc | JU. A cu | Jau | Br N. Amer. 18'Jl. I) | I) pi | l)i. muse. 6&6 |
| 25702 xelagino'ds L. | Sclago-hkc | Is A cu | 1 au | Br Britain w.al.hc. 1 | I) pl | Eng. bot. 1148 |
| 25703 rupéfite L. | rock | JL A cu | ■ | Br N. Amer. ... | D pl | Schk. fil. 165 |
| ii. Di'tn KJIA.—Leaves distich, stipulate | | | | | | |
| 2.170* ornithopodioides L. | Ornitliop-hkc | 1 st . A cu) jn.au | Br | 1812. | D pl | DLmusc.661.B- |
| 25701 denrésum Swx. | depressed | ■ iA cu 4 au | Br | C. Cl. H. 1816. | I) pl | |
| 2J7M helveticum L. | Swiss | >: A cu Tm .. | Br | Switzcrl. 1779. | D pl | Di. muse. 64.8 |
| 25707 denticlitum L. | toothed | JL. A <u>s</u> 1 in JL | Br | Swizcrl. 1779. | D pl | Dimusc.66.1-A* |
| S5708 spudum L. | stemless | III A cu i au | Br | N. Amer. 1819. | D pl | Di. muse. 64.3 |

- Selaginella** *JL. A cu | an Br Britain w.al.h. D p i Eng. bot 233
rather glittering JL. A cu | an Br N Amer. 1829. D p i Schk fil LEM*

(*Psilos*, naked; *destitute of leaves*.) D 1.—3.
■ ZQu 9 jiaii Br W. Indies 1793. D p.1 Schk. fil. 

CLASS XXIV. ORDER 4.

CRYPTOGAMIA MARSILEATEJE.

2837. ISOETES L. QUILLWORT. (*Iosos*, equal, *etos*, the year; remaining the same throughout.) 1-4
 aciistris W. marsh & A cu 1 my.o Br Britain al.lak. D p.l Lng. bot. 1084
- P1LUL A'RI A L. PILLWORT. (*PUula*, a pill: little heads containing the reproductive organs.) 1.
 lobulifera L. globule-bearing t A cu ± Jns Br Britain moi. h. D p.l Eng. Dot 521
- SJW. SALVI'NM Mich. SALVINIA. (*Antonio Maria Salvini*, Greek prof, at Florence.) 1-4.
 nkuaw All. floating ^ lit ^ my.au ... Italy 1818. D p.l Schk. cryp. 173
- M(x) MARSVLEA L. MARBILBA. (*Count L. F. Marsigli*, founder of Acad. of Sciences, Bologna.) 1.-8.
 quadnfolw L. four-leaved ^ cu 4 jn.au ... Europe 1820. D p.l Juss. sci. 1740

CLASS XXIV. ORDER 5

CRYPTOGAMIA MU'SCI.

TRIDBI. A'STOMI.

	English NUM.	Popular Character.			Locality.	Refrain to Figures.
PH A'SCUM L. PHASCUM. (<i>Phaskon</i> , an ancient Greek name for a moss.) 1-39..						
I. teOLOSi'nwM.-Shoots creeping, leafless, articulated, branched.						
serratum Schreb. serrated solitary & spring Pa. G shady banks Muse. brit. 5						
stolon ifcrum E. B. 2006						
II. ERE'CTA.—Creeping shoots none.						
alterniflum Dick. alternate-lvd solitary & spring Pa.G moitit banks Muse, brit 5						
crispin Hulw. erip solitary £ spring Pa.G banks and II. Muse, brit.5						
sulmaltuin L. 18 subulate small patches i spring L.G dry banks Muse, brit R						
axillare Dick. axillary lax. sol. sp.iuinmer Y.G moibt banks Muse, brit 5						
strictum E. B. 2093						
pateni Hedw. spreading solitary fa sp.summer Pa.G clay fields Muse.brit.5						
niuticum Schreb. pointless solitary * sp.summer Bt.G moist banks Muse. brit. 5						
1 inkjus Hook. larger solitary ft Hpsummer JUG moist banks Eng. bot. 2027						
2 minus Hook. smaller solitary JJ np.summer Bt.G sea coast Muse. brit. 5						
jiukitum Schreb. cuspidate solitary & sp.summer G hcd. moi. ba. Muse. brit. 5						
apiculatum Hook. small-tufted solitary sp.summer Gsh hcd. moi. ba. Eng. bot. 2025						
hrcberftititi li B. 2026. um E. B. 2959						
85.52 iiii %*. pilifcrd» patches sp.summer Hoa sandy downs Eng. hot. 1888						
ST.2-1 ^ynWes 7/c*. Bryum-like solitary } sp.sumtnr G banks und fi. Muse, brit 5						
jx^2 rertum ***. upright solitary ft sp summer L.G moist banks Muse, brit.5						
'-< curvicollum Hedw. bcnMiccked			-fa sp.summer	L.G	moist banks	Muse, brit 5

TRIBE II. GYMNO'STOMI.

SPH A'GNU L. Boo MOBS. (A name employed by Pliny to designate some kind of moss.) 4-11.						
blunteaved aquatic 7 year Y.G bogs Muse, brit 4						
common aquatic 7 year Y.G bogs Eng. bot. 1405						
Hook. viridiflum E. B. smaller aquatic 3 year Y.G bogs Schwfig. sup. 3						
lniflum E. B. floating aquatic 24 year Y.O bogs						
* minus Hook. Web. squarrose aquatic 7 year Y.G bogs Muse. brit. 4						
Turn. Ehr. sharp-leaved aquatic 6 year Y.G bogs Muse, brit 4						
S857a>epli<f;iliumi:A cuspidate aquatic 6 year Y.G bogs Muse. brit. 4						
<* cupidatum Ehrh. feathered solitary * spring L.G banks, Dev. Muse. brit. 8						
^ymn6stomum penfatum E.B.9319						

GYMNCYSTOMUM Hedw. GVMNOBTOMUM. (*Gymnos*, naked, *stoma*, mouth j orifice of theca.) 15.-47.

I. RAMOSA.—Slcfis long. In anched.						
Jfffw. Lapland dense tufts 1 spring I) G alpine rocks Muse, brit. 8						
Jffw. "in £.B.2201 ummr thick tufts \ spring BtG wet rocks Muse. brit. G						
viridflum E. B. greenest tufts f summer BtG trees & rocks Muse, brit 6						
S57J71 fimmOB Forster/ E. B. 2225 benUbcakcU tuffs 1} spring Pa.G moist rocks Muse. brit. 6						
2*^irg^tn'n Hedw. 857M. ill>min K.B. 2202 rock large tufts If sp.summer D.G moi.tnou.ro. Schweg. sup. 11						
schw. ttr <i>l</i> Schwtc.						
II. SIMPM'CIA.—Stems short, simple.						
Griflhssiriftum E. B. Griffiths'* little sjmts J summer Pa.G mountains Muse, brit. 7						
vatum Hedw. ovate broad jwttches \ year G ba. & wa.tops Muse.brit.7						
1 vulgdx Hook. common broad patches i year *G ba. & wa.tops Eng. bot. 1889						
3574.1 Kt&cile Hook. slender broad patches year G ba. & wa.tops						
"i trunc^tulum Hofm. small truncated patches iycar Bt.G fields *j ban. Muse. brit. 7						
>termfedium E. B. l!76						

25742*HcfmJJ Hedw.	Heim's	small patches	1 summer	Rsh	marit. banks	Muse. brit. 7
olijtusum E. B. 1407						
25743 rōmēcum Schuee.	conical	little spots	ft summer	Pa.G	fields, S. Ire.	Muse. brit. 7
2aJ744 lasncl.'rc Hedw.	bundled	patches	ft hummer	Y.G	clayey banks	Muse. brit. 7
2A745 pynlifinc Hedw.	pear-shaped	dense patches	J summer	Bt.G	moist places	Muse. brit. 7
'25146 tlnuc Hedw.	slender	little patches	lft spring	BtG	sand't. rocks	MubC. brit. 7
paucifolium £. A 2506						
25747 micrōstomum Hedw.	small-mouthed	little patches	ft spring	Pa.G	banks	Muse. brit. 7
25748 Donniunum Sm.	Donn's	thin patches	ft spring	Pa.G	Scotch rocks	Muse. brit. ^
2965. ANICTANGPUM Hedw.	AWICTAWHUM.	(Anoiktoi, open, aggeton, Taie: open theca.)	2.			
25741) ciliitum //c ^t u.	ciliated	depress, tufts	1 summer	Hoa	rocks	Muse. brit. 6*
Gymnobtoimim ciliatum £. A1179						
25750 hnbcrbc 7ooJt.	bear ^A 11? M> _m	d e P r e M . tuft*	1 summer	Pa.G	Irish moun.	Muse. brit 6
Gymnustum imbfrbc E. B. 2237						

TRIBE III. APLOPERI'STOMI.

9966. DIPHY'SCIUM Mohr	DIPHYSCIUM.	(Dfr, twice, phykton, a vesicle: shell of theca double.)	1.
S5751 loh>suin Mohr	leafy	matt patches	ft spring
I'suxbaAmif f'oliusa^.		D.G woods	Muse. brit 8
B. 329			

8967. TE'TRAPHIS Hedw.	TETHAPIIB.	(Tefra, four; theca four-toothed.)	2.-5.
25752 pellucidum Hedw.	pellucid	wide tufts	1 year
Vsfis. spl.u'hnoles Hook.		Pa.G dry banks	Muse. brit 8
25753 mowniūin Grey.	Brown's	solitary	ft year
ovāta Hook.	GrSmM/tf Brounidn/i E. B. 1422	OLG roofs of caves	Grev. cryp. 169

2968. DI'SSODON G. \$ A.	DISHOHOV.	(DM, twice, odous, a tooth.)	2.—4.
25754 fplachuoU's (I. \$ A.	Splichiuin-lk	wide tufts	2 summer
Vsfis. spl.u'hnoles Hook.		D.G alpine bogs	Grev. cryp. 145
25755 Ir'hu I><IM/M G. & A.	I'r'u'lich's	tufts	1 summer
TurneriuNum K. B. 1116	Aiphadlimmi i'ra'hchihittmj Hook.	reticulatum E. B. 2507	Pa.G mountains Muse. brit 9

25756 spha'ricum/	spherical	solitary	2 summer
		Pa.G dung of ani.	Muse. brit 9
25757 tōnuc Dick.	slender	subsolitary	lft summer
Grhnto clphaehnolden Fl. br.		Pa.G Scot. moun.	Muse. brit 9
25758 mnioKies L.	Mmmm-like	tufts	2 year
1 minus Hook.	smaller	tufts	1J year
2 miuH Hook.	larger	tufts	2 year
fa*tiji. ¹ tuin£.a78G		Pa.G mountains	Muse. brit. 9
25759 angulMtituin L.	rarrowed	tufts	ft spring
25760 anipultecuni L.	hotticheaded	solitary	3 summer, aut
TurneriuNum K. B. 1116	P&G bogs		Muse. brit. <J
25761 vasculusum Hedw.	vascular	tufts	3 spring
2 rugsum E. B.	wrinkled	tufts	2 spring
		Pa.G mountains	Grev. cryp. 17H
		Lu.G mountains	Grev. cryp. 311

2970. CONO'STOMUM Sivx.	CONOSTOMUM.	(Konos, cone, stoma, mouth; teeth of theca united.)	1.—4.
257(2) Iwrealc Sum.	northern	small tuft*	1 summer
Grlmrarw conf6stoma	E. B. 1135	Bt.G moun., Scot	Muse. brit 10

2971. POLY^TRICHUM L. POLYTEICHUM. (Pots, many, thrix, hair; calyptra hairy.) 10. —22.

	I. mu'DA.	— Calfflitrunaked.	gs^
525/3 undulatum Hedw.	vrnvc-lcavd	solitary	4 autumn
25764 herc^nicum Hedw.	Hercynian	solitary	3 autumn
		OLG mountains	Muse. brit 10
	II. HIIHHUHA.	— Calyptra hairy,	
25765 pillferum Schreb.	hair.bearng	solitary	# autumn
25766 junijērinuni W.	Juniper-like	solitary	# autumn
strctum E. B. 2435		01. G heaths	Muse. brit. 13
S3767 septentrionale Swx.	northern	solitary	OLG heaths
scxangulārc E. B. 1906		Muse. brit W	
25768 commine L.	rommon	broad masses	# year
1 yuccafitWum Ehrh.	Yucca-leaved	broad masses	# year
2 attenuat. Mrnx.	narrowed	broad masses	# year
gracile E. B. WL		OLG heaths	Muse. brit 10
257G9 alpinum L.	all>inc	patches	4 summer
iLVHO urnigeruin Mcnz.	urn-bearing	scattered	4 summer
*Mtni flcmli'N 7/fv. W.	Aloe-like	scattered	lft autumn
1 inajiiH Hook.	larger	scattered	lft autumn
rufelium E. B. 1939		Br.G heaths	Eng. bot. 1198
2 Dicksbn^ Turn.	Dickson's	scattered	1 autumn
85773 nhnuin Hedw.	dwarf	scattered	1 autumn
subrotiindum £. J. 1624		Br.G heaths	Eng. bot. 1609
		lir.G moist banks	MUM:, brit. 11

25773 fontinaloides Beauv.	Floating	5 summer	D.O1 in streams
Fontinalis nitnor E. B.			JEng.bot557

25774 enervis H. *G.	nerveless	subsolitary	ft year
rigida E. B. 180		D.G clay banks	Brew. jour. 1
25775 brevir6sts H.\$G.	short-beaked	subsolitary	ft year
25776 rtgida Turn.	rigid	small patches	# year
SI777 nuirilis lldcw.	wall	tufts	1 year
SB778 ruralis Elnh.	country	deep patches	2 year
2SVJ) subtila Hedw.	subulate	thirk tufts	lft year
SSIH0 eupifblia Roth	wciglcavated	solitary	ft spring
U57H1 stellata Sm.	stellate	Military	ft spring
>i782 trntubKa Hedw.	tortuous	broad masses	lft sp.summer
25783 fallax	deceitful	tufts	lft year
unsuiculata E. B. 2316,	imbeibis E. B. 2J29	L.G limebt. rocks	Muse. brit. 12
2V784 gracilis II. \$ G.	slender	lax tufts	1 summer
		R.Br clay banks	Muse. brit. bi'l

² < 85 revoluta? rftf. 27/11/1 unguiculata E. B. 2383 27/11/1 convoluta Swz.	revolute tufts	■ spring	L.G banks	Muse. brit. 13
27/11/1 unguiculata E. B. 1291, anstatae A. J. 392, barbata E. A. 2J91, humilis E. B. 16J1, apiculata B. B. 2494	unguiculate tufts	4 spring	Dp.G ban. & hedg. Muse. brit. 12	
27/11/1 convolute loose patches	wide patches	ft year	Y.G moist banks Muse. brit. 12	

27/11/1 ENCALYPTA Hedw. EVCALYPTA. (En. within, kalypter, covering; large calyptra.) 4.—7. 27/11/1 > treptocirpa Hedw. twisted-fruited tufts 1) year				
vulgaris Hedw.	common	ft year	Bt G moist rocks Muse. brit. 13	
7frum extinctorium E. B. 558	ciliated	tufts	Pa.G mountains Muse. brit. 13	
rihMa Hedw.	self-colored	tufts	Pa.G mountains Eng. bot. 1418	
1 concolor Hook.	alpine	tufts	Pa G Scotch alps Eng. bot. 1419	
2 alpina Hook.	bundle-fruited	tufted	D.G Scot, mount. Grev. cryp. 163	
1 raptocarpa Schwet.				

27/11/1 Glrl'MM/J HCJW. GBIMMIA. (J. F. C. Grimm, a German botanist.) 11.—29.

I. IMME'K&K. — FruUstalks scarcely any.				
spocirpa Hedw.	distant-fruited	dense tufts	D.OI rocks ft trees Muse. brit. 13	
1 nlKro.vlndis Hook.	dark-green	tuto	D.01 rocks tk trees Kng. bot. 1134	
2 stricta rar/t.	strict	loose tufts	Ilu mountains Muse. hib. 2.1	
maritima Turn.	sea-coast	tufted	Br?G marine rocks Muse. brit. 13	
£> S ^{8nx1} co'a Hook.	rock-dwelling	subsolitary	HtG rocks Muse. brit. 13	
£& P ⁸ Pulvinaria E. B.	cushioned	round tufts	Br.G house tops Musr. brit. 13	
W. v. in Dicranum ovule E. B. 2165	russet	broad tufts	D.OI subalp. rocks Wern. tr. 4.6	
Ha Grev.	ovate	tufts	D.G alpine rocks Muuc. brit. 13	
Worm*.	hair-leaved	tufts	Hoa stone w., Sc. Grev. cryp. 100	
oniuna Sin.	spiral	tufts	Hoa alpine rocks Grev. cryp. 203	
	twisted	tufts	Br. alpine rocks (irev. cryp. 199)	
	one-colored	tufts	B.G alpine rocks Grev. cryp. 1%	
	Don's	little tufts	D.G loose stones Muse. brit. 13	

27/11/1 VfEI'SSIA Hedw. WfMSIA. (F. W. Weis, a German cryptogamic botanist.) 19.—54.

I. FUNA^RIA. — Thcca wUh an apophysis.				
JeupletdnJ Hook.	Templton's	little patches	ft spring	L.G banks Muse, brit 14

unaria Tcmplctn E. B. 2524

II. GRI'MMIA. — Thcca without an apophysis.

I. LANCEOLAVJK. — Leaves ovate or lanceolate.				
mndaT/ooAr.	naked	little patches	ft summer	L.G clayey soil Muse. brit. 14
Grimm « « "uda E. B. 1421	dark-colored	tufts	f summer	Br.G mount bnn. Muse, brit 14
Shui & Ritta 77/11/1.	broad-leaved	tufted	i autumn	Pa.G Scot, mount. Grev. cryp. 149
Urimyw. nigrta E. B. 1825	Starke's	little patches	ft sprng	D.G banks and A. Muse. brit. 14
StarkebM Hedw.	K. B. 1490	kindred	subsolitary	Pa.G fields Muse, brit 14
(irfmif Starkeuna tnt'lris Hook.	lanceolate	lanceolate	subsolitary	L.G moist banks Muse, brit 14
Grimmia lancolata E. B. 1408				

III. SUBULAVA. — Isavcs linear or subulate.				
striata Hook.	striated	round tuft	■ spring	BtG alpine banks Muse. brit. 15
? minor Hook.	smaller	round tufts	■ spring	BtG alpine banks Hed. muse. 13
gv. nakjor Hook.	larger	round tufts	j spring	Bt.G alpine banks Schwuf. sup. 19
gv. trit'holes Hook.	hairy	inutinpatch.	ft spring	Bt G granite roc. Muse, brit. 15
2510 Grimm n ¹ trichodes E. B. 8163	tendrillrd	tufts	ft summer	L.G decay, wood Muse, brit. 15
v. rlmma/cirrata A. A. 2J5(2) Dicksbni E. B. U20	disputed	dense patches	ft year	BtG banks Muse. brit. 15
v. tenuirobtini 77. Xf T.	Blender-beaked lax tufts	1	summer	BtG shady rocks Muse, britsup.3
* curvirostra Hook.	bent-beaked	tufts	1 year	11.G roc. and ban. Muse, brit 14
* recurvirrostra E. A. 14J8	rrispish	dense tufts	ft summer, aut	D.G rocks Muse. brit. 15
Hedw.	crispula E. B. 2203			
controvrsia Hedw.	disputed	dense patches	ft year	BtG banks Muse. brit. 15
^rnm/i controvrsia E. A. 1J67	chalk	subsolitary	ft spring	01.G chalk cliffs Muse, brit 15
calciroa Hedw.		solitary	A spring	L.G rocks Muse, brit 15
'frum calcareum E. B. 191				
Juvirvata Hook.	recurred	dense patches	r spring	BtG calcar. rocks Muse. brit. 15
Stain nm/ recurvata E. R. 1489	dwarf	dense patches	ft summer	BtG moist rocks Muse. brit. 15
* pusilla Hedw.	whorled	tufts	ft summer	Bt.G moist rocks Muse. brit. lfi
rimm/n pusilla E. B. 2551				
verticillata Schwa.	verticillata	tufts	1 summer, aut	O.L.G moist rocks Muse, brit 15
rimm/Vi verticillata E. B. 1258	acute	tufts		
acute Hedw.				
acute acuta £. A. 1(44)				

IV. PTKUGO^NIUM Sin. PTBROUONIUM. {P/cron, a wing.gonos, a shoot; pinnated Nti'inO :} —7.				
Sinlh/JSwz.	Smith's	creeping	3 year	Bt ¹ trees, S. En* Mimir. brit 14
irile Sin.	slender	creeping	lf year	Bt G subalp. rocks Muse. brit. 14
>&tre Hedw.	filiform	creeping	lf year	Bt.G inoiiiitains Muse. brit. 14

F.A2526

1>FCR ANUM Hedw. DICRANUM. (Dikranos. two-headed; dlv. of teeth of capsulo fringe.) 20. — 47.

I. Di'a-nniA. — Leaves distich.

nroldesSwz. Bryum-like patcheM ft spring Pa. G nroist banks Muse, bnt 16

Pfcranum virfdulum E. B. lkW, mmundiohies II. I>i2

diantoldes Swz. Maidcnfair-lvd patches 2 summer L.G wet pastur. Muse. brit. 16

Jpmiin/r/diagtoidei E. A. 4J04 » * , . . . "

xif.huin Swz. Vew-leaved tufts \$ summer L.G moist banks Muse, brit 16

yppnum /uxifulium E. B. 416 11. AHTRITMAN-KA. — Thcca without a strum.

Blaticum Hedw. glaucous broad tufts 4 autumn W.G moors Muse*brit 16

lutifoiium Hedw. broad-leaved subsolitary \$ spring Bt.G mouu. banks Muse brit" 16

I richustoinum piliferum E A 53b

23R30	<i>longiflbum</i>	<i>Hedw.</i>	long-leaved	dense tufts	3	winter, sp	BtG	wet rock	Muse, brlt	16
258J1	<i>fluxiisum</i>	<i>Hedw.</i>	flexuewe	loose tufts	3	winter, sp	D.G	peat bogs	Mime. M	16*
85832	<i>fld've seiM</i>	<i>Sm</i>	flavescnt	tufts	3	winter, sp	Y.G	river banks	Muse. lint.	17
208-l	<i>btrj.irrb.jiim</i>	<i>Schwa.</i>	squarroc	large wastes	3	summer	Y	wet san. pi.	Muse. bnt.	17
258J4	<i>pellucidinum</i>	<i>Swx.</i>	pellucid	tu(U)	1J	sp summer	D.G	wctsan.pl.	Muse, brit	17
25835	<i>spunnum</i>	<i>Hedw.</i>	spurious	dense masses	4	summer	BtG	bogs	Mu*c.	17
25H4M	<i>crispinus</i>	<i>Hedw.</i>	crisp	loose patches	J	year	BtG	moist banks	Muse. briL	17
25837	<i>Scollinium</i>	<i>Turn.</i>	Scott'	large masses	H	summer, aut	Bt.G	moun. rocks	Muse. briL	18
	<i>flagi-lilArc E. B. 1977</i>									
SJ&B	<i>polycarpo</i>	<i>>on Ehrh.</i>	many.fruited	round tufts	t	year	BtG	rocks	Muse. brit.	18
25839	■	<i>/fort.</i>	w&ve Jeaved	tufts	2J	summer	BtG	woods & roc.	Muse. brit.	18
	■		rock	patches	3	winter, sp	Dp.G	woods & ban.	Muse. brit	18
	■		larger	patches	3	winter, sp	Dp.G	woods &r ban.	Eng. but	354
	■		fuscescent	tufts	2	spring	Brsh	heathy plac.	Kng. bot.	1597
25841	<i>A fuscscens</i>	<i>Turn.</i>	varium	loose patches	i	spring	II. G	moist banks	Muse. briL	17
	■		green	loose patches	i	spring	D.G	moist banks	Eng. bot.	1215
	■		rufescent	loose patches	i	spring	Rsh	moist banks	Eng. bot	1216
	■		lurid	loose patches	i	spring	Lu	moist banks		
2582 ful	<i>fulvum</i>	<i>Swi.</i>	rather tawny	dense tufts		sp.summer	BtG	crcv. of roc.	Grv. cryp.	188
25841	<i>heteromallum</i>	<i>Hedw.</i>	various-woolcl	large patches		spring	BtG	moist banks	Muse. bnt.	18
25844	<i>ibulbundum</i>	<i>Hedw.</i>	subulate	loose patches		spring	BtG	moist banks	Muse. brit.	18

III. STRUMA YEA.—*Theca with a struma.*

2581! cervicuw tum <i>Iedw.</i>	small-necked	small spots	ft	spring	Str	bogs	Muse. brit. 16
pusillus. E. II 24 ^{JK} , uncinatura E. B 22til							
25846 virem <i>HcUw</i>	green	tuft	1} year	BtG	mount, mar.	Muse. brit. 17	
20847 strumiffl <i>Tuin Sm.</i>	wen-bearing	tuft	1 year	BtG	mount, mar.	Muse. brit. 17	
25818 falciulum <i>Iedw.</i>	falcate	large patches	2 spring, aut	Bt G	alpine rocks	Muse. brit. 17	
9584! SelirellH. r/nsttw <i>Hedw.</i>	Schreber's	tufted	\ spring	BtG	moi, pl., Scot	Grev. cryp. llo	
25050 <i>Slrkrik</i> VV.&M.	SURke's	tul	1 spring	BtG	alpine rocks	Muse. brit. 17	

8071). TRICHO'STOMUM Hedw. TBICHIOM-MH M (*T/irir*, a hair, *storna*, the mouth.) 9. —18.
 95851 Nitens Schwa. spreading deep utches G year Hoa mountains Muse, brit 18.

Dicranum pAtens E. B. TWO, obtusum 'n'. in:
 2M852 iHuanglin Hium Hcdw. woolly deip't tufts 4 year Hoa stony moun. Muse, brit 18
 258.) can' Hecu M Ht'dw canoscent tufted creep. If year Y.G heaths Muse, bnt. 18

283. <i>Juncus acutus</i> L.	canescens	tufted crep.	1 year	AG	heads	Muse. brit. 18
rru-(Mitis E. II. Wft						

*li. W5/1 microArMju II dw. sinall-fruited deep patches 2 year 01 rocks Muse bnt 18
2585*l*culturale* IJruv. nctnile-potted loose tufts 1) summer OLG wet rocks Muse, bnt 1)**

STATI rascisticure *Sc/tr* bundled broad patches 2 year
 25858 *Kilyphyllum Scifwic*, many-leaved round tufts f summer
Dasyphyllum *Ms-brachytrum* *E.P.* 1217 circatum *VL* br.

2555) Hm>rhHmi/rw.jk. *Dicranum ellipticum* E. B. 1901 little tufts f sp. Butnraer BtG moun. rocks Muse. brit 19

2580. GLYPIIOMITRION *Briil.* GLYPMOMITHIOV. (*Glypho*, to emboss, *mitrk*, a little diadem.) 1
2580. Глипомитрион *Бриль.* Глыпомитхiov. (Глыфо, to emboss, митркъ, a little diadem.) 1

25860 Davichi Vi. Ijnd. Davies's tufts spring BtG marl rocks MTSC BfT 13
Grinnell Vi. Daviess Turn. Encalpita Daviesi E. B. 1281

28801 Schenk's Schizura. *Tumifascia* creeping 3 summer B.G. trunk of trees Muse Brit 90
Dicranum Beiulordes E. B. IIKB

25862 *purpi'rium* *Hedw.* purple large patches ft year Ksh moist rocks Muse bnt. 90
JTrfum bipartitum If. B., 2357, Dicranum strlctum £.£.2294, Dicranum Cflsii E. B. 2414, T~~90~~

S5863 *inrlnatnum Swz.* inclining spots f spring L.G moist heaths Muse brit. 00
(*irrimmIA inrlntha E. B.* 1824)

25S64	<i>ncrvum Hook.</i> (irrim/a atrocivres E. B. 2015)	nerved	loose patches	ft	spring	Dp.G	dry banks	Muse. brit. 90
25S65	<i>flaviflum Hook.</i>	bent-leaved	loose tufts	+	sp summer	BtC	banks	Muse. brit. 90

25865	<i>Trichostomum flexibilium</i> E. B. 2490	bent-leaved loose tufts		sp.summer	BtG	banks	Muse. brit. 90
258nn	<i>ripidulum Hedw.</i>	ngidiwh tufts	f	sp.summer	Br	walls & roc.	Muse. brit. 80

25867 Trifoliodiniu rigidiluui !. B. 2178
 tnt.irriui Sixv. three-rowed tufted I sp.summer BtG moist banks Muse, brit SO
 TurhoMeijinu trifarinis E.B. 1707 [no]ides E.B. 2215

Thrinaxodon triaurum E. B. 1707, includes E. B. 2215
25868 cupid'it'um Schr. hairy dense tufts 4 year
Trich'hoinum capill'cum E. B. 1152

25809 heterocrinum & lllum *Hook* virious-woollcd patches A spring Y.G mountains Muse, brit 80
tirilmna hetercromalla E. B. 189U

25870 *Brunitinia* *lutea* broad tars 1 spsummer E.G. alpine rocks Grev. sp. 1
Brunitinia Muse. brit. sup. 4
 25871 *glaucoccaens* *Grev.* gUuccsent closely tufted 1 summer, w Gl Scot moun.' Grev. cryp. i

TBIDE IV. DIPLOPERI'STOMI.

WAS. FUNAHUA <i>Iledw.</i>	FUN&RM.	(<i>Unit, a rope; twisted structure of fruitstalks.</i>)	3.
525872 hyfvrmltrica <i>Hrdw.</i>	hyi?rometrical tufts	winter Pa.G everywhere	Muse. brit. W.
25873 Mühlh>n>frKf Turn.	Muhlenberg's tufts	spring Pa.G rocks	Muse. brit. W.
25874 hibfrnra Hook	Irish tufts	f spring Pa.G cottage roofs	Muse. brit. W.

2984. **OKHTOTRICHUM** Hedw. **ORTHOOTRICUM** (*Orthos.* straight, *th** hair; i teeth of peristome.) 1 & ⁴⁹.

2984. ORTHOTRICHUM Hedw. CYTHOTRICUM. (Strios, straight, *infr.* hair & teeth of peristome.) I-
1. *Kv'a*. — Peristome without ciliary processes. 11

25R75	Drummondia H. & G.	Drummond's creeping anomalous small-capped	li summer broad tufts } tufted } 1 year	DG trunk. of trees DQ1 rocks & walls DQ1 wo. & stones	Grev. cryp- Muse. brit. Muse. brit.
2.1870	anoianolius Hedw.				
25277	cyanophyllum Huds.				

25877 cup-shaped *Hofm.* small-cupped tufted 1 year D.01 wo. & stones Muse, bnr
nhdum E. B. 1325, animal urn E. B. 1423

SJG Ospeciosum Mv*	showy	tufts	1	summer	L.G. trunk. of trees Grev. cryp. 137
g<<IrupSnoia Funk	rock-dwelling	branched, lax	1	myjl	Br. rocks & walls Grev. cryp. 105
2 maj	Miss Hutchins's	tufts	1	spring	Br.G. rocks Muse. brit. 21
3 pumilum E. B.	kindred	tufts	If	spring	Pn.G. trunk. of trees Muse. brit. 21
	larger	tufts	1}	spring	Pa.G. trunk. of trees Eng. bot 1323
	dwarf	tufts		spring	Pa.G. trunk. of trees Eng. bot 2KJ8
III. PLIRICLIA. — Peristome with sixteen ciliary processes.					
ss8 Wdiaphanum ScAr.	transparent	tufts	1	gn.summer	Iloa trees ft walls Muse. brit 21
. arlb'tum Muse.					Muse. brit. 21
> pulchillium Sm. hib. 9.2	pretty	tufts	1	year	L.G. trunk. of trees Muse. brit. 21
nvulhre Turn.	rivulet	floating	2	year	G. roc. in strea. Muse. brit. 21
"triMum Hedw.	striated	tufts	2	year	BtG. trees Muse. brit. 22
Lyellw Hook.	Lyell's	branched	3	year	Y.G. trees
ZY'GODON Hook.	ZVOODO.V.	(Zygott, a yoke, odous, a tooth; teeth yoking together by pain.)	1.	small tufts	Pa.G. trunk. of trees Muse. brit. 21
conoideum Hook.	conoid		f	spring	
MA um conoideum E. B. 1239					
ANO'MODON Hook. ANOMODOV.	(Anomos, irregular, odous, a tooth; peristome.)	2. — 8.			
cimip'e'dulum/lcMJr.	short pendulous pinnate		8	summer	D.OI. roc. & trees Muse. brit. 22
NecKer. curtipinula /, B. 1444					
vitellKrum Hook.	tendrilled	creeping	6	spring	Y.G. trees & roc. Muse. brit 22
tvynura vitulbum E. B. 265	feathered	flat-branched	3	sp.summer	BtG. trunk. of trees Grev. cryp. 109
JACKEKA Hedw.	NBCKERA.	(N. J. Necker, a German botanist.)	3.-24.		
P <i>n</i> sila Hedw.	dwarf	creeping	2	spring	Pa.G. woods Eng. not. 1443
criupa Hedw.	crisp	creeping	6	summer	BtG. tr. & roc. Eng. bot 61(G.H.^pnL
Pennata Hedw.	feathered	flat-branched	3	sp.summer	BtG. trunk. of trees Grev. cryp. 109
I?? ⁸ - DALTO^N/^ Hook. DALTOVIA.	(Rev. James Dalton, an English muscologist.)	2. - 5.			
plachiiodes Hook.	Splichnum-lk tufts	\ summer	JUG	Irish moun.	Muse. brit 29
Neckem jplathno'ides E. B. 2564					
hetromalla Hook.	various-wooled tufts	\ summer	L.G.	trunk. " of trees	Muse. brit 22
Neckera heteromalla E. B. 1180					
9. FONTINA^LISL. FONTINALW.	(Fans, fountain; place of growth.)	3.-9.			
antipyrltica L.	heat-proof	floating	12	summer	Dl.d rivers Muse. brit. 22
s'lauinbsa L.	scaly	floating	6	summer	OLG. rivulets MUST. brit. 22
capillkca Dick.	capillary	floating	6	summer	Br.G. alp. rivulets Muse. brit. 22
fc BUXB AU'M I A L. BUZBAUMIA.	(John Christian Buxbaum, a German botanist.)	1.			
aphylla L.	leafless	solitary	1	summer	Br. Fir woods Muse. brit. 22
BARTRA*M/itf Hedw.	BARTRAMIA.	(John Bartram, an American botanist.)	«. — 1.		
I. Lov <i><<PEOiiicuLA'rA</i> .	— Fruitstalks very long, not curved.				
8fi9* pomiformii Hedw.	apple-shaped	tufts	ii	minimcr	Ht.G. heaths Muse. brit. 2J
2 nrtnor Hook.	smaller	tufts	1	i	Ht.G. heaths Eng. bot. 998
? major Hook.	larger	tufts	2	summer	Ht.G. alp. hea. Eng. bot. WiG.B.crU.
< Utliy skifij. Brid.	straight-lvd	tufts	1	summer	BtG. dry banks Muse. brit. 23
KricinH/6.	slender	deep patches	3	summer	Ip.G. alpine rocks Muse. brit. 23
* J>>fana Stz.	fountain	thin tufts	3	summer	Bt.G. wet places MUST. brit. 23
* major Hook.	larger	thin tufts	6	summer	BtG. wet places Di. muse. 44.2
* ffrum fontanum E. B. JK					
z märchica Stox.	La Marche	tufts	1	summer	Bt.G. wet places Eng. bot. 207*
II. BRUVIPEDUNCILAVJR — Fruitstalks very short, curved.					
MrMna Hedw.	Haller's	deep patches	6	summer, aut	BtG. moun. rocks Muse. brit. 93
arcuate Brid.	arched	loose tufts	4	summer, aut	BtG. mountains Muse. brit 23
HOOKE*RLf Sm.	HOOKERIA.	(William Jackson Hooker, LL.D. rec. prof. bot. Glasgow.)	2. - 27.		
InCTw Sm.	shining	procumbent	3	summer	Pa.G. moist banks Muse. brit. 27
late-vlrens Hook.	bright-green	procumbent	3	summer	BtG. Irish bog Muse. brit 27
LE'SKEif Ehrh.	LRSKRA.	(N. G. Leske, prof. nat. hist. Marburg.)	10. — 43.		
cichomano'ideHMv/io.	richdoman-lk	entangled	2	spring	Y.G. trunk. of trees Eng. bot 1493
Utlw.	flattimifd	entangled	4	spring	Y.G. trunk. of trees Eng. bot 1492
pnlychrMi Ehrh.	many-fruited	ctntun[skud	3	spring	Lu. trim. of trees Eng. bot 1274
25b,io Vpnum medium E. B. iniuIUum E. B. 1922	downy	prostrate	3	spring	Y.G. ground Eng. bot 2525
ykka Mohr					
25tn K<<MiuiN? rotundiflum E. R	pretty	dense tufts	1	gp.summer	BtG. moist hanks Eng. bot. 2006
asSi puFc H<<H Hedw.	rufescent	creeping	4	sp.Bummcr	Rsh. moun. rocks Eng. bot. 2296'
***) rureK cu.is Schwa.	silky	entangled	3	sp.sumrocr	Y.G. roc. & trees Kng. bot. 1445
? " <<<>>a/ferfio.	tree-like	erect	3	sp.summer	Y. G wo. and bogs Eng. bot. 15b5
Jendrrfdes/erfw.	incurved	procumbent	3	sp.summer	D.G. trees & rocks Eng. bot 2422
w irurvata Hedw.					
" Pnumatotrichos E. B. attenuatum E. B. 2120	many-flowered creep,	tangled	3	summer	Y.G. trunk. of trees Grev. cryp. 151
Yyantha Hedw.					
S99* HY'PNUM L.	HVPNIW.	(Hypnos was a name among the Grooks for a moss.)	07. — 119.		
I. PLANICAO'LIA. — Stems plane; leaves distichous.					
river bank	loose patches	4	summer, aut	BtG. ban. of ditc.	Eng. hot. 200
wavy	lax masses	fi	summer, aut	W.(r) heathy pine.	Kng. bot. 1181
toothletcd	prostrate	1	summer, aut	Ht.G. roots of true	Kng. hat. I.'(0
narrow-leaved	prostrate	II	summer, aut	Ht.G. roots of trees	Hod sti. cr 4.31
btusifbhmu Hook.	blunMcaved	prostrate	1	summer, aut	Ht.G. mountains Eng. bot 144U
2496					
11. PATE'NTI A. — Leaves spreading on all sides of the stems.					
delicate	denwe patches	1	spring	up u roc.	Eng. bot JTA
creeping	Datche's	1	spring	Bt.G. roots of trees	Eng. bot 1037
Poplar	entang.patch.	2	spring	D.G. trees ft ston.	Muse. hib. 16
reflexed	loose masses	2	spring	D.G. mountains	
alpine	entangled	3	summer	R.G. alp. rivulets	Grev. cryp. 282
soft	much tufted	3	summer, aut	Lu. alp.nvulce	Grev. cryp. 283

2^092 turbinatum SVJX. nfricans E. B. 1528	top-shaped	thin tufts	2 rammer	Pa.G	wctaan.pl.	Musc.brit.S9
» nitans Schreb. compactum E. B. 1527?	nodding	little patches	3 summer	Bt.G	walls &hca,	Muse. brit. 29
alpinum £.	alpine	tufts	■ summer	P	subalp/rock*	Muse. brit. 28
Jugatum Die*. ventnebum Dick.	ii. CRABSIU'BCULA. — Leaves with a thickened margin.					
1518, cubitale £. 27.2554	long	aubsolitary	1* summer	Bt.G	mountains	Musc. one. ju
Ieri Grev.	ventricose	deep-tufts	4 summer	Br	mar. ground	Muse. brit. 30
feArto. rosti atum &*. SJJI! "" Kinatum Die*. htonum Schreb. cuspidatum ScAreA.	Tozcr's	solitary	summer	Rsh	clay banks	Grev. cry p. 285
	dotted	solitary	3 summer, aut	L.G	mar. places	MUHO. brit. 30
	ligulate	solitary	4 summer, aut	L.G	moist banks	Muac. brit 30
	rostrate	solitary	S summer	L.G	subalp.coun.	Muse. brit. SO
	edged	tufts	2 summer	Y.G	shady banks	Muse. brit. 31
	this year's	deep tufts	5 summer	Y.G	mar. places	MUAC. brit 31
	cuspidate	subsolitary	2 BU miner	L.G	wo. At walls	Muse. brit. 31

TRIBE V. SCHISTOCA'RPI.

ANDREJEM Hedw.	ANDREAA. (J. G. II. Andrea, a German botanist.)	4.
alpina Hedw.	alpine loose tufts	1) summer D.Br rocks Muse. brit 8
riesttris Hedw.	rock dense tufts	* hummer D.Br rocks & ston. Muse. brit. 8
HdM Mohr	Roth's dense tufts	{ summer D.Br rocks & ston. Muse. brit. 8
nivalis Hook.	snow deep patches	3' summer D.Br mountains Muse. brit H

CLASS XXIV. ORDXR 6.

CRYPTOGAMIA HEPATIC&

JUNGERMA'NN/itf L. JUNGEHMANNIA. (louis Jungermann, a German bot. died 1G53) 81. — I. ABTIPULAVTJL — Stipules none ; stems leafy.

Leaves inserted many ways.						
hair-leaved	loose tufts	1) summer	D	turfy heaths	Hook. jung. 7	
setaceous	dense tufts	2 spring	Pu.G	bogs	Hook. jung. 8 HI	
downy	dense patches	summer	Pa.OI	mountains	Hook. Jung. 2	
loose-leaved	cush.-hkpat	J sp.summer	Pa. G	mountains	Hook. jung. 59	
Juniper-like	crowded tufts	3 summer	11. Br	mountains	Hook. jung. 4	
Hooker's	small patches	g winter, sp	G	ditches	Hook. Jung. 54	
II. BIFA'RIA: — Leaves bifarious.						
i. INDIVIVI*.	— Leaves undivided.					
Asplenium-lk	loose patches	3 year	OLG	moist woods	Hook. jung. 13	
spinulosc	crowded tufts	3 year	Y.G	mountains	Hook. jung. 14	
deceitful	dense tufts	1 autumn	OLG	Irish heaths	Hook. jung. 50	
Don's	cntangl. tufts	2 September	P.Br	Scot mount.	Hook. jung. 39	
dwarf	small patches	winter, sp	01	rocks	Hook. jung. 17	
lanceolate	dense clusters	£ autumn	Pa. G	damp woods	Hook. jung. 18	
heart-leaved	dcilic tufts	2 august	DQI	mountains	I look. jung. 32	
Sphagnum	cnlang. pate,	3 autumn	Y.G	maTahypl.	Hook. jung. a.s.2	
crenulate	matted patch.	f o.november	K.G	tigts	Hook. jung. 37	
round-fruited	dense tufts	\ early spring	Pa.G	Irish bogs	Hook. jung. 74	
transparent	broad tufts	1 early spring	D.G	bogs		
compressed	dense tufts	4 June	rivulets,IrcL		Hook. jung. CS	
ii. EMARQINAVJK. — Leaves emarginated or h/fid ; segments equal.						
cmarginata Ehrh.	large atres	\ may, junc	Br	wet pi. on m.	Hook. lung. 27	
concinnata Hook.	thick tufts	may, June	Sil	wet pi. on m.	Hook. jung. 3	
arcadinaig Hook.	loose patches	1 may, junc	Bt.G	mountains	Hook. jung. 71	
Dick.	dense patches	\ july	OLG	boggy places	Hook. jung. 38	
Hook.	cut out	scatter, patch.	D.G	shady woods	Hook. jung. 9	
ata Hook.	dcntc patches	\ au.n	Pa.G	woods	Hook. jung. 28	
two-pointed	Turner's	small patches	Pa.G	Irish rivul.	Hook. jung. 29	
Bryssus-like	larRC tufts	\ march, april	Pa.G	damp bunks	Hook. jung. 11	
connivicit	dense tufts	\ march, april	D.OI	heaths	Hook. jung. 1	
curve-leaved	loose patches	1 april, may	Y.G	wet places	Hook. jung. 15	
	small patches	I april, may	Dp.P	mountains	Hook. jung. Ifi	
Hi. QIADRI'PIDJE. — Leaves three or four cleft; segments equal.						
capitate	very sml. pat.	4 s.Ja	Pa.	bogs	Hook. jung. 80	
cut	sml.dncspat	\ July	Pa.G	heaths	Hook. jung. 10	
dwarf	sol.orthk pat.	\ October, may	Bt.G	moist banks	Hook. jung. <9	
bristle-shaped	dense tufts	2 spring	G.Br	mountains	Hook. jung. HO	
io. — Leaves bifid ; segments unequal, folded together.						
grove	matted tufts	2) July, October	P	woods	Hook. jung. tl	
flat-leaved	crowded pate.	2	Din.Br	mountains	Hook. jung. 67	
shady	dense tufts	\ march, april	G.Br	shady places	Hook. Jung. 24	
wavy	large tufts	3 may, June	Bt.G	wet places	Hook. jung. 22	
recaupinate	very unl.tufts	J may, June	Ilr.G	heaths	Hook. jung. 23	
whitish	broad tufts	If april, July	Pa. G	hedge banks	Hook. Jung. 25	
blunt-leaved	little tufts	\ march, april	Pa. G	damp places	Hook. jung. 26	
Dickson's	dens.tut.fuf.	\ august	1. Br	mountains	Hook. jung. 48	
minute	loose patches	f sp.summer	Ol.Br	mountains	Hook. jung. 44	
cut out	small(Nitches	i	Pa.G	heaths	Hook. jung. 19	
L.	spoon-shaped	large nches	4 summer	R.Br	mount. bogs	Hook. jung. 68
Hook.	flattened	1) summer	Pa.G	trun. of trees	Hook. jung. 81	

11. STIPULAV*.—*Furnished with sttputet.*

L IVTEOIIHKOU*.	— L* nw ^a i'n/jrc or rarely emitt					
anomalous	IHKHC patthe ^c	I	o.ruivcmuc		jiiia.	JJ
Taylor*	largo patchet	3	tumuicr	Pk	IQUUltaill	Hcink, jung.
ladder	brood pjtchi	j	lummox	Pa.<J	loamy toil	Hook, jt..
many-flowered	loo*c pat-he*	!	april, may	Pa.O	wet placet	HDOIL jin
weilge-leaved	paraiitiral	£	tuminer	Br	oiJiinfi'r.nn	Iliuk, jung.
twiggy	loote patches	jl	anting	Y. (lr	par. dump pi,	Hwik, Jung.
TricKomape*	large patches	13	MUM*	V.Xt	nolit place*	Hook, jutig.

Trickmane	large patches	1.	MUM	
it.	Bid'tIM.	—	Leave* two or three cleft; mgment equal. crowded pat.	
		<1	ML.all tulf d cuih.-Ukc tuf.	li o.nnvcmbor tnpn.l.n
Ji	bearded		crowded pat. crowdedpntt'. It	J summer iaprll.july
rti'(ill){			loose [withes	M'niik
hriH.-lotml			J turn me r	P B .0
			1 Jimmii r	WIHHI\ IIM. limit. jlllh. 70
			3 pat int	r., i; ikn fm.. 0 wood*
				Ol.O rock* Hook. jung. 76

iii. HiVin<. — *fsnecii liyid*; *tfgmtnlt unequal, fittiied together.*
 P l i N ' C* i nli, or tmntlfr onft, fitti.
 lirn.nl-llMVIKJ wide pntcht'i 1 march, au ilr.li uld wa Hnok.jung.40.*.5
 liKne tufti 9) iiimiticr I : !) woinii H E K I K , J U N G .
 d>iiM.-patrbm 2" *>.jumner tLBr rocks A hea. Hook jui
 \Vi)ixl'ii cmwileil tutli 3 ip.juumner ILDr lltboun. Ilmik. Jimg. fO
 •mall ilowny broad patchet 3 march, oct. Pg.N maitt placet Hook jung. 36

<i>svoixm'nott.</i>	broad patches	3	march, etc.	Pn.G	mantl placet	Rook, jung. 53
<i>Latuer segments, or tmaller onet, invnltr.</i>						
<i>HaclLty'4</i>	dents patchei	1	tnovember	Blt.U	tree* & roc,	<i>Unok.</i> jung. 53
<i>Tliyine.leived</i>	ImbHctnuKt	^	april, junu	Pa.	6 trim, of trees	Rook. jung. 42
<i>hook-leaved</i>	very tmll.pat.	1	iprlng	(r	rocks	Hook. JIIIIK,51
<i>minuteit</i>	little patche*		april, may	V.C)	hrnk of trcM Hcmk. jung. SB	
<i>calyptra-lvd</i>	little lufU	t	^minmer	Pn.G	on f*lc* naa.Hook, jung, (*)	

iii. S**CCATl'nOJi	— /fiiu.T KifmenU, or smaller onet, saccate.
Jlifl Hutehlmii (OOM patchM	1 summer
MilIti'l)tr.P Inn:
TamarUk	large pfltrdei 3 april t
	sp. in 1 st birth lank Jung 1

HutcliliMbr Hook .il.itiilu /.. TmmariKl Umtk.

SS08I Vitiid. 1 L

48083 U
S *Schrank.*
S60S7 t,yi!l»i IIRH.k.
S6D68 htWrniw Hook.

M.A.i:< HATSTA
iwilyiiiiirih. /,
S(»9») hemitlik rica I-
9009Z aaid'iijrynn A'.

3000. I.U'CCI^ E.B.
idea ft //
* n Hum £, /(.
fltiltnn. K A

3001. ANTHOCER
SfIO37tnuUittdin *Dick.*
SfOtW |uiiri.iitis /.
9009D tnijor /; K.

anfia. TAIIGIO^NI/* KB. TAWHOKI*. (*John Anthony Targiimi, a Florentine ixil-uit.*) 1. — 3.
SSIOO hypophjlla /, *Unf* under I Laf broad patch la \ winter, »p U.U wtl plucw Eng. bot S87

30Ca. H X n Q A ; B. SniJiKoCirTft. , a globe, *karpoi*, fruit; jccd.vouel.j J. — *.
36101 terrJtrU g. B, curt h ipot* winter Bt.O (Uljni) Ucei Kng. bol, (NW

CtAfisXXIV.

CRYPTOGAMIA ALGAE

Tame t.

ACHNA'NTHES Ag. *Aenurnu H.* *i.Achnr*, the froth of the orean, *anthns*, a flry
i Ag. *lotiK-^t* *ttiked* fine down *Jy* *jl* (>*h* *diLica* coal Eng.bot.V4w, &
IftJ *lirevipei Ag.* **hort-italked* fine down *IJ* *KU* *Osh* orean *Cirpv,trz,*
EGIOi unipuncUlo Oirm. **inglec-«potl-^t* *Hmc* down *li* *iu* *Cih* ocean *Orer*, i

aotfi.	IXIII.A'RIA	Grew.	EXIUIHI*.	{ <i>Xkilh</i> , tlwider; <i>jiluitli</i> .)
	"")'.	circular	line man	Ar *-ri'tB BtO watery pi.
i	iilti: i fftcti	bundled	Ittik- tuft*	*f tit U.Y MM
	tluirring	littio	tiifti	-^ tu O lea
	hd	t<	t upjning	li te*

m.vroMA^r.	/nja <i>i,imt</i> M>paralion; ftuurnenU divided into juinls.)	7.-1(1
Ag. nje rioastcd	Hm & MI Y llr dltche* KtiK li.. r, l\l CunfC.	
MI] in 4J. in. tlnrt down 1j pting illih prrnN Ajr. >viui. hot. 491	riMhtdmr M *ting ill water t n i hyil lit	
>'ni"ui[h*jiiij Ag. &tin Btildutpli kloet down A n.d O Kil	little tnfu * f y.O >rtwn	K.
iff. oblique mlnuu bnuie.i « IX Br ocean Eng. ba ^ r i w	mlnuu bnuie.i « IX Br ocean Eng. ba ^ r i w	
L.T..VIUA Ag. iis, brittle; nature] 8L	.it; i..ml like limit tiiit- j mi- V o on w.lt plfJIL Kljii lut.t.till.	
9*nkBi Ag. winter dense n. tufu 3 ap O. Br rivulet* Lyng. liyd. (U	dense n. tufu 3 ap O. Br rivulet* Lyng. liyd. (U	
MKE.OSEI'RA As. MKUKHIU. i U./(in,au ant!* i, to divide, n-w.M. n fltament.) 2.—9.	vn-Uko v t^n* nu Brh smicoait Ejig,bot,£101 Curtjh-'t.	3.-5.
m	llu-i lown JV mr V.ii rivilli'tn DIW.ron, i short duivn , tit Drill ives of aquat. I>tw. COD. '~->. B t.'nuiii.	
•"Cyllilirn'j ri.iiliMilleUIjE, ii. •	(DnnmuM,a boml; parti palMlini whan in a itntc of dltoilution.) %	
nniupips tj MI i. niiw.it.Ti Gnnr.ervp.eaS	ttHWO IIIMIMI 1 ipring Y.U tlthct Gnr.ervp	
lax i- li <u>u</u> V u IH 4, A.cryp.30S	tUpp. nu Brh smicoait Ejig,bot,£101 Curtjh-'t.	
ttmall-tiincit iTirl tuft* \ »)iri.iH V.II MM I.r.,	filuiji I >i ltr>]) lakes	
ft. TV. 'J'llw. liihi-riiV	- MI OLCS MM	
QU Lilipunct mini Grev, four-dotted Lix tuttit 1! <u OL< tea	1! <u OL< tea	
GOMPHONFTMA 4f. ("s.....MtM> ;Ajj, i dab, rirma.n tlnment.) 1	Psiul6xitn Jf, pnradnxicil due down 1 HI HI	
KrmiuLlum Ag, twin round HUM f i)riiit U*h.Y nip. rivulet* lir.**. i <>riig Uih.V IHM:J . cryp. !M	toinuiisiijnmn Grev, miitulctt	
Trmin- II.		
S- CHLOROCO'CCUM A ** Cn- year O ttnn. afroet Orer. cryp, 208	*iljürr (. cum ii ii .it thin magi	
« J, I*prirla butryotdd £. H. ai*» yoar O Crev. cryp.325	it ill Diiti h	
A- linn liyi-r (Proio*. * berry.) J. rocks Grev,crvp,£31		
PAI-ME'LLA Ag, /Paltru, Tibration j Jelly-like nature.) 7.—12.	f t ' l n i h t " ill* 1 >prlng <+ utrenmii Orcr.cryi'	
JointLil thin Him ... jn.jl 0 luko» Eiig.lwl. LJ7S.t,rt:Wjiui	innta ep.auL G rookl King. bot. B 8a U'ftm	
Ag. ALPvostiniM. (ffnflywiftasthtfcwnoftb>wmihiltot) .T.—7,	lik thill <kiii ... nut 0 damp place* Orev. cryti	
trAm Kirvit flehy maw (i M> V txran Knj, bot. U(U. tl'in	ntintc gregariinii } aut Y.Hr mount.rock* Lynf. byd.tl!)	
rlavacetit flyhy mam 3 IU \ oana PI djin.ttia.UVM	ri.si Brcenridui ...MI Pk on litln	
broken vermicular 4 »u Y.Pk ocean Eng. bot. ICS& Viva	mountain iciiWiko H MI HCi alpine rock* Knj, IM< BW5. U'i-n	
wiul ^b IV. g. !. IIIM-II.I I/EchimiM,a. hedffehog; brlmly appcarttneo.) 1.—	thlncreuit ... year Jt.Br damp walU Grev,cryp.UU5	
•rtlcuLlu Ag. JointLil thin Him ... jn.jl 0 luko» Eiig.lwl. LJ7S.t,rt:Wjiui		
Ag. ALPvostiniM. (ffnflywiftasthtfcwnoftb>wmihiltot) .T.—7,		
trAm Kirvit flehy maw (i M> V txran Knj, bot. U(U. tl'in		
rlavacetit flyhy mam 3 IU \ oana PI djin.ttia.UVM		
broken vermicular 4 »u Y.Pk ocean Eng. bot. ICS& Viva		
common lobnl m.i» i m. OLO dampplacw Enu. tot.4fil. Trcm&a	(Without explanation.) 5.-32.	
ptum-ibaped little bmfi ra OLG laken hyrtfl		
HUtericnl littl< iu (11.0 Kill wateri III mu<c. II). 17		
w;irty (trcpirinuii] ap o n ; rwk* Di. muse. II. K.		
blue minute balls .., in t; (Srcv. eryj		
0 rov. (IL-v. Mr. BiTkrfy, an K.igllih crypt, bot.) I.		
•Upp. I'll i itirtnjj lit tea Orar.erjp		
ite Grrv. Mvill.lMiH*. (Myritu, injrl.nl, nrmt, filament.) v cryp. 300		
IS ^b RYNE'1 ¹ MORA At. ConVNKPURM». (Koryn?, n club,pAi^mi, tol>e>r j club.ihajNHI tlnmwuI> J, mmit ID i< 1i Htl iore Oref. cryp, 53		
lubwiWrmi 5. S. 1056		
unkted£>		
TOT'HORA j*ff. CHJrrronnoM*. tC***** brlrl^jpfoneo, to btari tlamcnli.) 4 —12	\ Utii/ux, n rh	A. —11.
fur-ditfl tiin liyt-r ^j aut MM ptanoi Bttg. bot IT98	•Mte J aut >on	
warly balli i mil HI Q ii.ii'hftH Kng. lw.t <W	Kiihill DUM. 1J	
Ktiave-loavjil liiniwhitl G till IV	b h d f tu	
*«*m		

3024. SCYTHYMETNIA *Am.* SCVTIIYMBWIA. (*Skyo* leather, *h*~~u~~*m*, membrane; substance.) 1.
86102 rupestris *Ag.* rock broad mass **24** su *B*~~r~~ rocks Eng. bot 2194

Tamii III. CONFERVOI'DE/E.

3025. BYSSOCLADIUM Ag. BYSSOCLADIUM.
86163 fcncstrlc Ag. window fine tuft
(*Byssot*, fine *fax,klados*, branch.) 1.—
ft year G on windows Dlw. con. 94

- | | | | |
|------------------------|-------------|---|---------------------------------|
| 3026. MYCINEWf A Ag. | MVCINEMA. | (<i>Myket</i> , a kind of minute fungus. <i>ne?na</i> , a thread.) | 5.-6. |
| 26164 arachnoitcum Ag. | cobweb-like | patch ft year p.,Q dead trees | Ulw.coti. C. <i>Conferva</i> |
| 26165 fiftvum Ag. | tawny | down-like ft aut Br rotten wood | |
| 26166 rubigibsuin Ag. | rusty | patch ft year Rust rotten wood | Dlw.con.G8.C0//VVM |
| 26167 nhosphircum Ag. | phosphoric | »iches ft year V rotten wood | Dlw. con. HI <i>VoijMm</i> |
| 26168 pulvireum Ag. | powdery | thin crust ft year G rotten wood | Dlw. con. 78. D. <i>Cotrif.</i> |

- | |
|---|
| 3027. CHITOO'LEFUS Ag. CHIIOOL. (<i>Chroos</i> , skin, <i>lepo</i> , to decorticate; inner memo, changing to powder.) 5-6. |
| 2616! Jithloches Ag. Hock Violet patches ft ft yg ^{an} P socks Fl. d ¹⁶⁰⁰ 899. 1 |
| 26170 odoratus Ag. sweet-scented patches ft ft winter Bbr trees Lyngbya hyd ¹⁷ 7 |
| 2 <i>if171</i> /ichenicola Ag. Lichen-dwelling down ft >u II.O on lichens Eng. lot. 1609 |
| 2 <i>if172</i> rubricidiis Ag. ruddy patches ft su It.Br barkofap.tr. |
| 20173 rrbencus Ag. Ebony patches ft year Ilk rocks Eng bot.702. <i>Bpssus nig-</i> |

- | | | | |
|--------------------------|----------------|--|---------------|
| 3028. TKENTEPO'HLJJF Ag. | TRENTEPOILIA. | (Trentepohl, an obscure German botanist) | 3.-4. |
| 26174 purpurca Ag. | purple | , patches | tW year |
| 86175 afirea Ag. | golden | patches | ft year |
| 2 ilicicola Ag. | Holly-dwelling | branch, pate, | ft spring |
| 86176 pulchella Ag. | pretty | downy tufts | ft spring |
| 2 chatfbecke Ag. | iron | tufts | s |
| | | D.01 | fresh water |
| | | | Eng. bot 1996 |

- | | | | | | | |
|-----------------------|--|-------------------------------------|-----------------|--|--|--|
| 3029. SCYTONtTM A Ag. | SCYTONEEMA. (<i>Skytos</i> , leather, <i>nema</i> , filament: nature of filamentous fronds.) 6.—17. | | | | | |
| 26177 compactum Ag. | compact tufts 1 su | Ilksh mountain* | Lyng. hyd. 28 | | | |
| 26178 byssiticum Ag. | Byssus-like tüfts * su | II rocks | Dl. muse. 1. 18 | | | |
| 26179 myochrouM Ag. | mouse-colored sjimy coat 1 su | D.Br rocks | | | | |
| 2 ocellatum Ag. | eyed silmy coat 1 su | D.Br aquat plants Eng. bot 2530 | | | | |
| A imundum Ag. | inundated fclimy coat 1 su | D.Br inund.places Eng. bot. 1555 | | | | |
| 26180 BällKf KypK- | Hang's compact tufts lft su | Ttrlug subalp.banks Lyng. hyd. 28 | | | | |
| 261K1 Sowrybu,u/n Ag. | Sowerby's short down ft su | Ol.Br ocean Eng. bot W19. C. mir&O- | | | | |
| 36182 comoides Ag. | hair-like broad patches fto | RBr ocean Eng. bot. 11 | | | | |

- SOSO.STIGONE-~~■■■~~. STIGONEMA. (*Stigon, dotted, nemb, filament; regular annular dots of filaments.*) I--³,
86183 atrovirens Ag. dark^recn bushy tufts ft su Bk.G rocks Dlw. con. 25. *Vonfirv**

- | | | |
|-----------------------|--|---|
| 3031. PROTONE'MA Ag. | PHOTONEMA. (<i>Protot.</i> , first, <i>nema</i> , | a filament; simplicity of structure.) 7.—10- |
| 26184 rr'jens Ag. | creeping | j>ches ft su G pots in hoth. |
| 261KJ umbrusum Ag. | shady | patches ft ft su G on the earth <i>Dlw. con. 61. Confirva</i> |
| StilHo yulutum Ag. | velvety | patches ft n G on the earth <i>Dlw. con. 77. Cotif<r**</i> |
| 2iiH7 fragrans Ag. | fragrant | patches ft n G on the earth Eng. bot. 1556 <i>Gryve</i> |
| SdKH crypt+ruin Ag. | vault | patches ft su (r caverns Kng.bot. 2 ¹⁸⁸) |
| 2iH8 Ord'K'ilnchi Ag. | <i>Orthotrichum</i> | dishM tufts ft su |
| 26190 7iuscicola Ag. | Moss-dwelling | minute down ^ ap Br on Orthotr. <i>KgthWiS</i> |

- | | | | |
|----------------------------|-----------------|---|---|
| 3032. HYGRO'CROCIS Ag. | HYUROCROCIS. | (<i>Hygros.</i> moist, <i>krokit</i> , a little tuft.) | Br. on mosses Eng. bot. 1701.
7.-9. |
| 27111 <i>bajit'tca</i> Ag. | Barytes | fructiflts | year Tran solofmbr.b. |
| 2til'2 atram. nti Ag. | ink | fine tufts | A year Whl surf. of ink Lyng hyd. 57 |
| 2til'w typhlod'rma Ag. | obscure-skinncd | tine tufts | ft year OI in sol. g.arab. Dlw. con. 83.C |
| 26194 pallida Ag. | pallid | fine tufts | ft year Y sol. of ochre Dlw. con. ■ |
| 86195 <i>Rhsai</i> Ag. | lime-water | fine tufts | <fa year Tran rose water |
| 26196 sanguinaca4g. | blooilcolored | fine tufts | ft year C ibinglass size |
| 86197 vini Ag. | wine | fine tufts | ft year V. in Med. wine |

- | | | | |
|-------------------------|-------------|--|-----------------------------|
| 3033. LEPTO'MITUS Ag. | LEPTOMITUS. | (<i>Leptos</i> , slender, <i>mitos</i> , a thread.) | 4.—15. |
| 86198 minutissimus Ag. | minutest | little tufts | • & year Tran on mar. alga; |
| 26153 (Lacteus Ag. | milky | patches | ft winter Tran pools |
| SiYKH <i>i</i> n'NM Ag. | dwarf | like down | ft aut G rotten algae |
| SOaoi clavatus Ag. | clavate | minute | ft aut Tran dead fishes |
| | | | Dlw. con. 79. Cm |
| | | | Dlw. con. 30. Coijij |
| | | | Lyng. hyd. 22 |

- | | | | | |
|--------------|--------------------|---------------------|---|---------------------|
| 3034. | MESOGDC! | A Ag. | MESO (<i>Mesot</i>), middle, <i>globos</i> , <i>viscid</i> : spines globose, brachaea solid mass.) | 5 — 8. |
| 86202 | <i>multifolia</i> | Ag. | many-cleft tufts | 3 aut |
| 86213 | <i>Hudsonia</i> | Ag. | Hudson's branched | 6 aut |
| 86204 | <i>coccinea</i> | Ag. | scarlet bushy | 4 su R |
| | <i>Uvularia</i> | <i>verticillata</i> | <i>E. li.</i> | ocean Eng. bot 2466 |
| 86205 | <i>capillaris</i> | Ag. | capillary tufts | 3 su It ocean |
| 86206 | <i>vermiculans</i> | Ag. | vermicular bushy | 5 au Brsh ocean |
| | <i>2 coriacea</i> | Ag. | leathery bushy | R au Brah ocean |
| | <i>Rivularia</i> | <i>vermiculata</i> | <i>E. B.</i> | Eng. bot 1819 |

- | | | | |
|--------------------------|----------------------------|---|-----------------------------------|
| 3036. BATRACHOS BEPRUNUM | <i>g.g. BATRACHACH</i> | (<i>Batrachos</i> , frgg, <i>tperma</i> , seed; grows in marshes.) | 2.— |
| 862G/ vagum Ag. | rambling fine tufts | 3 my | bsh ditches Lyng. Ind. 64 |
| 2 tenuifissimum Ag. | slenderest fine tufts | lft my | D.OI ditches Eng.bot.690. Coty*Ag |
| 26208 moniliformic Ag. | necklace-shaped fine tufts | 2 su | G fresh waters |
| 1 stagnic Ag. | pool fine tufts | S IU | G pools Di. muse 7.44 |
| 2 simplicius Ag. | simpler fine tufts | 2 su | B pools' Di. muse. 7.45 |
| 3 purpurascens Ag. | purpurascens fine tufts | 2 su | Pk sea shore Di. muse. 7.40 |
| 4 dactylosum^ | cleared fine tufts | 8 su | D.OI pools Dlw. con. 11 t. |

3036. DRAPARN ASDIA Ag. DRAPARNALDIA. *{Janet Philip Ralf. Draparnaud, a French bot. 3, J. S.!!^111''4 BT.* slender fine tufts * year DLG pools Dlw.con (T ^ P S
& 4 Kp Pm 4 *glomerata* *leafy* broad tufts 6 su Bt.G rivulets Engm. 107. J.
heal.M.m *gelatin* tuft* 4 su-su BLO pools

				(Plants having an <i>oscillatory motion.</i>)	13.—47.	
1. OSCILLATO^RIA Ag., OSCILUTORU.	winged bordered	patches thin layer	4 spring } spring	Hr wet rocks V.C wet rocks	Grev.cryp.3tt Grev. cryp. 246	
alata Grm.						
limbata Grcv.						
rupestris Grcv. cryp.						
14 tenuissima Ag.	slenderest	patches	su Pa.	G warm springs	Eng. bot 2584. Conf.	
autumnalis Ag.	autumnal	If su	01. G	on the earth		
2 vaginata Ag.	sheathed	11 su	O1.G	in the earth	Dlw. con. 99	
	black	4 su	D.G	still waters	Dlw.con.64.0/o/j/m,Vj,	
	lcather-iffe	broad layer	1 spring	Ysh rocks in wat		
	brownish	tufts	1 year	Br.V stones in riv.		
	Stendid	thin masses	1 year	Fa B wat in both.		
	Sender	slippery layer	* spring	Pa. G still waters	Dlw. con.'Ju.t///>*`	
	mud	G year	Erug inud. bot. po. Fl. dan. I.W. M			
	blue	floating mass.	... year	B church walls	Eng.uot 2578. <i>Conferv</i>	
	barking	thin film	... year	B.G damp wood	Dlw. COIL 25	
	ochre-colored	thin flakes	} mr	Och pools	Dlw. con. 62. <i>Conferv</i>	
		gelatmasscs	i year			
2. CA'LOTHRIX ^.	CALOTHIX.					
SVS. ^W	yellowish					
nWea 4K.	snowy					
infervicola ^.	Conferva-dwelling					
•'bulirum 4f.	rocks					
imr. ^{culata} 4g.	fascicled					
bilis Ag.	wonderful					
?frta if*,	distorted					
lanata ^.	woolly					
* fuscescens Ag.	fuscescent					
1 Y'NGBY ^ Ag.	LYNGBYA.					
mur. >> Ag.	patch					
crispa^ff.	wall					
prolifica Greu.	crisped					
BA'NO/if Ag.	BANOIA.					
Laminalic 4	(Christian Frederick Bang, a botanical author.)					
atropurpifca .	Laminaria					
2 ftscu-ii.	dark-purple					
la Carm.	brown-purple					
	beautiful					
3041. ZYGNE^MA ^.	ZYGNBMA. (Zygos, yoke, nema, filament; flam, jointed together in pain.)					
cruciatum Ag.	crossed	entangl.mass.	\ ap	Y.G ditches	Eng.bot S4G3.C.bipuncf.	
2 longi-articulatum	lg. long-jointed	entangl.mass.	\ ap	Y.G. ditches	Dlw. con. 2. A.	
3 brevi-articulatum	lg. short-joint.	entangl.mass.	\ ap	Ybh rivulets	Dlw. con. 2. B.	
leucossum Ag.	decussate	floating tufts	H su	G ditches	Dlw. con. 39	
color Ag.	two-colored	floating tufts	3 su	G sto. in rivul.	Eng.bot.2288. <i>Confdrva</i>	
	pectinate	patches	11mr	G rivulets	Eng.bot.24(3.B.(o; [^] Jr.	
	curved	patches] mr	G rivulets	Eng.bot.24ti3.A.C.<cl.	
(juinliuin Ag.	quinate	large masses	} sp.su	1L G still waters	Van. con. 5.1	
	decimate	large masses	1 su	BtG still waters	Dlw.con.4.A.B.Cnf/j<fti	
	shining	float.patches	2 su	BtG ditches	Eng. bot. 1656. <i>C.spiridlis</i>	
punctatui' Ag.	dotted	floating cloud	■ ■	BtG pools	Dlw.con.51. <i>Confithra</i>	
2. MOUGEOT^T/4 Ag.	MOUGEOTIA.	(J. B. Mouceot, a cryptogamic botanist)				
genuflexa Ag.	knee-bent	ctngLmaes.	A ap	Y.G ditches	Dlw. con. 6. <i>Conferv</i>	
tarulescens Ag.	ceruloscant	pale patches	1jj1	P.B ditches	Eng.bot.2457. Coij4rua	
3. HYDRODICTYON Ag.	HYDRODICTYON. (<i>Hydor</i> , water, <i>dictyon</i> , net; reticulated structure.)					
utriculatum Ag.	purse-like	floating web	6 jn.s	G riv. & lakes	Eng.bot.16B7. C.reticul.	
* CONFE'RVA Ag.	CONFERVA.	(Conferrumino, to consolidate; close coherence.)			52.—13a	
		1. SIMPLICES.—Filaments simple.				
		i. COLORANT.—Filating, cobwebbed, colored.				
Sug Jpcetorum illtfA	heatly	blue web	sp.su	Br.P dry bogs	Grev. cryp. 261	
	lpinc	fine web	1 sp.su	Br muntains	Grev. cryp. 261	
	banded	fine web	\ spring	P.Br pit. on rarr.	Dlw. con. B. <i>Conferv</i>	
	silky		\$ su	G pools & dit	Dlw. con. GO.C.s&rdida	
		ii. VI'RIDES — Floating, cobwebbed, green.				
	napped	float, masses	1\$ spring	G ditches	Eng.bot.2303. C sardida	
	mucous	float, masses	1} spring	G bogs	Dlw. con. B. <i>Conferv</i>	
	zoned	long tuft	3 year	G sto. in rivul.	Dlw. con. 47. C. licens	
	bursting	floating tufts	3 su	G ditches	Eng. bot. 2461	
	Dil.	entangled	broad mat	3 su	G sea-shore	Eng.bot.2WJ.C.wipl<fxa
		Hi. — Capillary or setaceous ; articulations filled with globose granules, when dry alternately				
	tumidish	fine film compressed.	1 mr	G pools	Eng. bot. 1670. <i>Cinflata</i>	
	blistered	float, masses	6 mr	G ditches	KIK bot.ZMH.C.a/fcn).	
	fuscescent	float, masses	6 mr	G ditches	Dlw. con. B.	
	rivulet	long tufts	24sp su	G rivers	Eng. bot 1654.	
	English	long tufts	24 HJSU	G ditches	Dlw. con. 79	
	capillary	long tufts	3 sp.su	G ditches	Di. muse. 5.35. B	
	Flax	long tufts	3 sp.su	G ocean	Lyng. hyd. 50	
	matted	small tufts	spring	G sea shore		
	tortuous	crisp masses	■ ■	G sea shore	Eng. bot. 2220	
	thick	crisp masses	ap	G salt marshes	Dlw. coa 9	
	Melagonium	tuft	■ su	G ocean	Dlw. con. II	
	brazen	long tufts	6 year	G ocean	Dlw. con. 80	
	Young's	minute tufts	4 ■	G sea shore	Dlw. con. 102	
	nefklae-Uke	minute tufts	1 su	G sea shore	Lyng. hyil. 49	
	slippery	floating tufts	■ su	G Germ, ocean	Eng. bot. l'fia C. *ea	
	flaccid	tufts	■ su	G Hutchi	Dlw. con. 49	
	equal-jointed	float patches	1 spring	G Twc on r	g Twc on r C.	
		tufts	isu		Eng. bot 2310	
		tufts	my			

96977 ferruginea Both	iv. ADNAVJK. — <i>Adnate, pencilled, fastigiate, colored.</i>	Dlw.con.66. <i>C. fucicola</i>
86378 cflra Dil.	rusty tufts 1 year Rus. on Fuel	Dlw. con. 76
26279 cornea Dtf.	short minute tufts ft HU Ollir on Fbel	Dlw. con. 84
	flesh tuft* lft aut Pk on ■■■■■	
26280 a>ruginea Hud.	11. RAMUUK. — <i>Filaments branched.</i>	
26281 Hriwnii Dil	verdigrise tufts lftsu BtG seashore	Dlw. con. E.
W2H2zii\SiTx*Fl.dan.	Brown's patches ■ spring G Irishcaverns	Dlw. con. D.
2H23 rip&ria ZW.	starry floating tufts 2 su G ins.ofwa ves. Fl. dan. 600. 1.	
&2H4 gloinocrata L.	bank floating tufts 3 su G salt ditches	Eng. bot. 2100
2f2H-JenspiU /fe/A	clustered bushy tufts 1 su BtG riv.on stones	Eng.bot.IH54.C.te/tft>r
SISG llfacta I//.	curled p itches 2 sp.su G lakes	Eng. bot. 2350
25287 patens Ag.	broken large tufts 6 sp.su ■ pools	Eng. bot. 2338
2 prolificra Ag.	spreading large tufts ft su ■ ditches	Eng. bot. 2338
	proliferous large tufta ft su ■ ditches	Dlw. con.I.O.C.Jlt'xuitka
26288 congregate i*g.	111. CUMULAVM. — <i>Filaments branched and heaped.</i>	
26289 lanbua y%.	heaped tufts 81 G roc. sea coo. Lyng hyd 5G.C.uncidli»	
26290 flavescent Atf.	woolly tufts ■ SU G on /J'igic	Dlw. con. E.
&J91 sertcea Hud.	flavescent tufts ■ SU Y.G salt ditches	Eng. bot 2088
9648 rfrActa /to/A	silken shining tufts ■ SU Y.G ea shore	
SlitkJ HHpcra 4?	broken crispent.tufts ■ jnjl Pa.G ocean	Eng. bot 2327. <i>C. albida</i>
26294 heterichloa ^.	rough tufts ■ year G ocean	Dlw.con.EX. nfgncani
SO21S rupi strih A.	various grass tufts i year D.G ocean	
2P < ■■■■■ L	rock dense tufts 4 BU Gl ocean	Eng. bot. 1699
	wild goat balls foott ball 3 su G lakes	Eng. bot. 1377
26297 arcta K. B.	close floating tuft 3 su Tran ocean	King. bot. 2oJS8
86298 vuueberiaformis Ag.	Vaucheria-ik branched 1ft su G ocean	Dlw. con. E. C. brcta
86299 catenae L.	chain-like patches 1 su lir ocean	Di. muse. 5.27
aiJO) Hutchins'e Dil.	aiJO) Hutchins's dense tufts 4 su Gl ocean	Dlw. con. K/y
26301 pellucida Hud.	pellucid finely branch. 5 au Pa.G ocean	Eng. bot. 1716
86302 di*tans Ag.	distant loose bundles 6 aut Pa.G ocean	Dlw. con. 21. ■■■■■
3045. BULBOCHJETTE Ag. BULBOCIJETE. (<i>Bolbos, bulb, chaite, bristle; primary filaments.</i>) 1.		
86303 setigera Ag.	setigerous delicate tufts 6 aut D1.G lakes & riv. Dlw.con.59.C.vlvlpava	
3046. NITE'LLA Ag.	NITBLLA. (Nito, to shine; shining plants.) 5.—11.	
26304 traingtlicens Ag.	transparent branched ■ su Y.OI pools	Eng. bot 1855. <i>ChUra</i>
86305 filixlis Ag.	flexible branched ■ su Y.OI pon. & rivul.	
Chra flexilis L.		
86306 opica Ag.	opaque branched 12 Jl.au Y.OI pools	Eng. bot. 1070. OMrfje*
26307 mlqlfca Ag.	nest-making branched 12 jl.au Y.OI pools	Eng. bot. 1703. <i>ChAra</i>
26308 gracilis Ag.	•lender muchbranch. 6 • Y.OI pools	Eng. bot. 8140. C ■■■■■
3047. CHA'RAL.	CHIARA. (Chairo, to delight; delights in water.) 3. —16	
86309 hispida L.	hispida branched 9 Jl.au Y.G ponds	Eng. bot 465
26310 vulgaris L.	common branched 12 Jl Y.G ponds	Eng. bot 336
86311 aspen W.	rough branched 5 su Y.G lakes	Grev. cryp. 339
. 3048. CERA'MIUM Ag. CRAMIUM. (<i>Keramion, a pitcher; appearance of capsules.</i>) 23.-38.		
	I. FABTIGIA^A. — <i>Filaments short, fastigiate.</i>	
26312 Ianugin.sum Ag.	woolly fine down &e y"ar Br on i"lg» Dlw. con. 45. <i>Confir*</i>	
26311 floridulum Ag.	flowering little tufts ft year Pa.G roc. sea shor. Dlw. con. F	
26314 mlqMs AR.	creeping short down &e Jl Pk on large^"lg. Eng. bot 1608. <i>Confer*</i>	
86315 pdcma Ag.	feather fine tufts ft su R on largeyjlg. Dlw. con. F.	
26316 DavitsM Ag.	Daviv's small tufts ft jl It sea shore Eng. bot. 2329	
86317 Rothi Ag.	Roth's broad tufts ft su V sea shor. roc. Eng. bot 1702	
	II. DICHTOMA. — <i>Filaments dichotomous ; branchlets forked ; Joints obscure j thec hvolured.</i>	
86318 dlaphanum Ag.	diaphanous diffuse 5 su Va ocean Eng. bot. 1742	
8 pilosum Ag.	pilose diffUse 5 su Va ocean King. bot. 2428 ■■■■■	
86319 rtbrum Ag.	red solitary weak 10 su P ocean Eng. bot.llffifi. ■■■■■	
	III. RAMbsA. — <i>Branches furnished with branchlets, which are more or less dense and shortened.</i>	
26320 tctragonum Ag.	square tufts 3 su R ocean Eng. bot 1690	
26321 pedccilmum Ag.	stalked dense tufts 4 su O sea shore Dlw. con. 108	
8 W2 Houkeri Ag.	Hooker's fine tufts 1ft su Pa.br sea shore Dlw. con. 106	
26323 ltens Grev.	spreading solitary? 3 spring Fi.R sea Grev. cryp. 318	
26324 urb(j)cula Ag.	little tree bushy tufts 3 year D.R sea shore Eng. bot. 1916	
	IV. ALTERNAVA. — <i>Branches pinnulate, pinnule alternate.</i>	
86325 corymbsum Ag.	corymbose little tufts 1ft jl R seashore Eng. bot 2352	
86326 pinnulatum Ag.	pinnulate tufts 2 R sea	
2(27 roscum Ag.	roiy finely branch, 1ft su R ocean Dlw. con. 17	
86328 /hbjUleVte.	Arbor-Vitoe-ik finely branch. 6 jl R ocean Eng. bot.2465.(■■■■■)	
ZIKJ-28 veraicolor Ag.	vannus.colorcd fine tufts 3 RU P.R. on Ffci Eng. bot.966. <i>C. rbsea</i>	
86330 Bomri Ag.	Borrer's little patches 1ft o O.R. ocean Eng. bot. 1741	
26331 dticum Ag.	rough tufts 6 spring Di.P sea shore Eng. bot. KMF	
86332 intrruptum Ag.	interrupted little tufts ft jl Di.P sea shore Eng. bot. 1838.	
	V. OPP6BITA. — <i>Filaments piniate s pinnae) opposite.</i>	
26333 Ttirneri Ag.	Turner's delicate bran. 2 sp.su Pk seashore Eng. bot. 2339	
26334 plCimula Ag.	featherJife delicate bran. 2 sp.su Pk seashore Eng. bot.ll37.C.2Vtf**	
3049. GRIFFITH'S/i Ag. GRIFFITHS™. (<i>Mrs. Griffiths, of Devonshire.</i>) 5-7.		
26335 multifida Ag.	many-cleft lf fine tufts 3 Jl R sea shore Eng. bot 1816. ■■■■■	
26336 eqiihCitolia Ag.	Equlscutm-lvd sponge-lk-tu. 6 su R sea shore Eng. bot. 1479	
26337 setdeca Ag.	bristly lax tufts 4 year R sea shore Eng. bot. 1<<9	
2/338 barbate Ag.	bearded flocculent 1ftjl C sea shore Eng. bot. 1814	
2(UJ9 corallina Ag.	coralline branch, tufts 3 Jl O.R sea shore Eng. bot. 1815	
3050. CHJETO'SPORA Ag. CIJETO'SPORA. (<i>ChaUe, bristle, spora, seed; on fine divisions.</i>)		
26340 Wfgfi Ag.	Wigg's finely branch. 5 su RBr seashore Eng. bot 1165.*■■■■■	
	VI. POLYSIPHO^NIA Grev. POLYBIPHONIA. (<i>Polys, many, siphon, tube; tubes for color.</i>)	
	I. SIJBPINNAVS. — <i>Purple or scarlet, flat, somewhat pinnated.</i>	
	parasitical small patches 1ft su R.Br on Fleet Eng. bot.142. ■■■■■	
	Bony-stnd small patches 1ft R.Br Ma shore Eng. bot	
2(K343 cocclinea Grev.	nearict bushy tufts 1 year S ocean	

OMAA.M . * ~ 111 SE	CUNDIRA % MBA. — Creeping j branches divaricating, often one-sided.			
» M divaricate Grrir.	divaricating tufts	3 su 11 ocean	Lyn. hyd. 34	
wu*5 gracihs Grev.	III. PURPU^REA. — Purple hole collated adhering to paper.			
	slender long tufts 4 year P ocean	"	Dlw. con. 40. C. stricla	
	I, Mack, generally rose-colored above, adhering to paper.			
	IV. p* -			
vfolkca Grev.	violet little bushes 9 str	Vi ocean	Lyn. hyd. 31	
26347	larger bushy tufts 6 su	D.P sea shore	Eng. bot. 2340. C. ngra	
	nigrescent fine tufts 6 su	D.P ocean	Eng. lwt. 1717. Conferva	
	pectinate little tufts 1 su	D.P ocean	Eng. bot. 123) C. Jibrida	
untill il Li Givo.	urceolate long branches 8 year	li.Br ocean	Dlw. con. G. Conferva	
JJx*»i>thiiii,iU	elongated shrubby 8 year	H.Br ocean	Dlw. con. 33. Conferva	
s>>< allochroa	different-cld small tufts 1) year	V ocean	Dlw. con. G. C. Jibrida	
	Brodie's large tufts 15 year	M.R. ocean*..	Dlw. con. 107. Conferva	
utrorubscens Grev.	dark-red long tuft* 6 su	B.R marine roc.	Dlw. con. 70. Conferva	
« NiJ RJT — Black or blackish-brown when dry, rigid, scarcely adhering to paper.				
* astigata Grev.	peaked little bushes 2 su	D.Jjr ocean	Kng. lwt. 176*. C. polym.	
* Agardhina Grev.	Agardh's bushy tufts 8 su	11 sea	Grev. cryp. SIO	
* hild Grev.	brown fine tufts 3 su	D.Br ocean	Dlw. con. G. Conferva	
va Grev.	recurved fine tufts 3 su	D.Br seashore	Dlw. con. G. C. pdtens	
W7bygoWesGrw.	VI. LATERAYES. — Branchlets lateral, short, fascicled.			
Jmticulbia Grev.	Uysalis-like slender tufts 6 spring R sea shore		Eng. bot. 597. Conferva	
nlamenfosa Grev.	small-shrubby finely branch. 4 su	Br ocean	Eng. bot. 1686. Fucus	
	filamentous bnnich. tufts 4 mr	II ocean	Eng. bot. 2312. C. GrifWA	
LJE Ag. RVTIPILKA. (Rytis, a wrinkle, phleo, to abound In.) 2.-3.				
Gwit	dyer's masses 6 year OLG ocean		Turn, fuci 224. Fucus	
	plane tufts 5 su R sea		Turn, fuci 23.h. Fucus.	
LECTOCA'RPUS Ag. EcorApapua. [Ektos, outside, karpos, fruit; theca not enclosed.] 4. - 8.				
Ritkousik Ag.	smalUiKhdcd bushy (spring D.G ocean		Dlw. con. E. Cotif-rva	
I atrovirens Ae.	dark-green bushy 6 spring Rus. ocean		Eng. hot. 21!<. C. si/cul.	
	ferruginous bushy 6 spring Rus. ocean		Eng. bot. VMIX. Mtothr.	
RanulususK. brachiitic floating tufts 3 ap	Pa.br ocean		EIK. hot.&V. I. Voujyra	
tomentibus Ag.	granular flocculent 3 jl OLG on Fbci		King. bot.&l>I. Conferva	
	downy fine down 2 jl Br ocean		Grev. cryp. 3IG	
Jffl- kPHACELLA % RIA Ag. SPH. (Sphakelos, gangrene; appearance of truncate extrem. of frond.) 6. - 14.				
ra gci m WG w.	racemose tufts 1 su OLG sea shore		Grev. cryp. 96	
W67 SHITWAG.	cirrhoose dense tufts 1 su OLG sea shore		Dlw. con. 86. C. confirm	
RTTQ 2 Intertissima Grev.	most spreading little tufts 1 su 01. ngr on Ffici		Grev. cryp. 317	
radicans Ae.	rooting fine tufts f su Br.Ol marine roc.		Dlw. con. C. Conferva	
	plumose flarcid 3 spring G.Rr ocean		Eng. bot. 2; > H. V. viinnta	
W* Wnii Ag.	Alcrtens's elegant tufts 4 su OLUR Bacchoro		Eng. bot. IM) Vonifrya	
J71 fcoparU Ag.	Broom dense tufts 3 su Br ocean		Eng. bot. 1552. Conferva	
CLADOSTEPHUS Ag. CLAD. (Klados, a branch, stephos, a crown; whorled branches.) 2. - 7.				
spongiosus Ag.	spongy rigid tuft 3 su G ocean		Eng. bot. 2427.1. Conf.	
myriophyUum Ag.	myriophyLeaved rigid tuft 5 su G ocean		Eng. bot. 1718. Cvertic.	

TRIBE IV. ULVA^CE-E.

VAUCHE'R/^ Ag.	VAUCHIERIA. (Rev. M. Voucher, of Geneva, a bot author.) 7. - 29.			
chitoma Ag.	two-forked large tufts ■ su DG ditches		Eng. bot 932. Conferva	
? "uhmarlna /f	submarine large tufts ■ su D.G submar.ditc.		Lyn. hyd. 20	
? p <i>rri</i> *M Ag.	Dillwyn's thin mat ■ sp.su D.G on the earth		Dlw. con. 16. C. frieida	
J*li*at1 4,7	ground creeping sp.aut G moist grou.		Vbu. con. 3. 3	
	rooting patches % D.G dry ditches		Eng. bot. 324. Trem.gr.	
	double large masses 6 su D.G still waters		Eng. bot. 1766	
	racemose large masses 4 su.aut D.G ditches		Lyn. hyd. 23	
	many-fruited patches & su.aut D.G dry banks		Dlw. con. 71. Confifrya	
CODIUM.	(Kodion, a skin; appearance of second s/cleclii) £. - 7.			
downy coralloid 6 Jn G ocean			Kng. bot 718. F. Ueues	
purse spong. mass 3 year G ocean			Eng. bot 2183. Fucus	
BRYO'PSIS Ag.	BBVOPBIB. (Bryon, a moss, oopsis, resemblance.) 1. - 7.			
uindsa^.	feathery fine tufts 2 i Dp.G ocean		Eng. bot. 2375. U7I./>/<0i.	
sp.	compound fine tufts 3 su D.G marine roc.			
Ag.	SOLENIA. (Solen, a tube; tubular nature of fronds.) 4. - 18,			
	intubinal simple' 24 su G ditches		Di. must 9.7	
	largest simple 24 su G ditches		Di. muse. 9. d	
	Linza simple 18 su G ocean		Di. muse. 9. 5	
	lanceolate simple 18 su G ocean		Eng. bot 1739. Viva	
	compressed nearly simple 12 su G ocean		Di. muse. 2. 7	
	haired compound 12 su G ocean		Dlw. con. F. Paraddxa	
	latticed branched 3 su Vsh fritch water		Eng. bot. 2137. U7if.nim.	
2 uncliiQta Ag.	hooked irreg. bran. 3 su Ysh ocean			
	ULVA.	(Ul, water, Celt.; place of growth.) 5. - 12.		
	broadest soft leaves 12 su.aut G ocean		Eng. bot 1551	
	Lettuce soft leaves 5 spring G marine roc.		Grev. cryp. 313	
	blistered soft leaves 6 su.aut (j ditches		Eng. bot 2320	
	crisp broad mass 1 iu (j on the earth		Di. muse. 10. 12	
	furturaceous minute leaves & aut G walls		Grev. cryp. 265	
	POBPIIYRA. (Porphyra, purple; plants.) 9. - 3.			
	torn soft leaves 4 IU P sea shore		Fl. scot 33	
	purple soft leaves 8 su P sea shorn			

TRIBE V. FLORI'DEJE.

PuLYinBa	{Polyides, multifarious; diversity of appearance.) 1.		
worm-like	coralloid 6 n Br ocean	Eng. bot. 1738. Fhc. rot.	
	V f 3		

306.1 PTIIJOTA Ag.	PIILOTA.	(<i>Ptilolos</i> , pinnated; form of frond.)	1. — 4.
2(J97 plumosa Ag.	feathery	fine tufts	4 hii.aut It ocean
2 tenuissima Ag.	most delicate	fine tufts	4 su.aut 11 seashore
Eng. bot 1308. <i>Fucus</i>			
9064. RHODCMELA Ag.	RIODOMELA.	(<i>Rhodon</i> , a rose, <i>melos</i> , a limb; color of fronds.)	5. — 21.
2ffW8 dentata Ag.	toothed	flat branched	4 su Ol.Br sea shore
S6&Pl lycopodioidci Ag.	Lycoxxdium-lk coraloid	6 su Ol.Br ocean	Eng. bot. 124L <i>Fucus</i>
LJKO subtuse Ag.	brownish	finely branch.	6 su Ol.Br ocean
26401 bcorpidides Ag.	scorpion-like	feathery	4 su R.P ocean
26402 /iuastro/ides Ag.	Pinaster.lkic	acicular	3 su Br ocean
			Eng. bot 1042, <i>Fucus</i>
3CKS. CHO'NDRIA Ag.	CIIONDRIA.	(<i>Chondros</i> , cartilage; substance of fronds.)	9. — 38.
26403 pinnatifida Ag.	pinnatifid	bushy	6 su Pan ocean
96404 obtiua Ag.	blunt	bushy	4 su Y.Pk ocean
96401 ovalis Ag.	oval	rigid branch.	4 su Br ocean
26406 thasyphylla Ag.	thick-leaved	4fclum.lkic	4 year P ocean
96407 tenuissima Ag.	slenderest	aspargnoid	6 year Pa.Ol ocean
96408 davallbsa Ag.	small-studded	gelatinous	!) jl.au Pa pk ocean
26409 kalifbrmis Ag.	kali form	coralloid	5 jn D.LP ocean
96410 articulata Ag.	jointed	much branch.	6 su H.Pk ocean
			Eng. bot 1574. <i>Fucus</i>
9066. SPHA:ROCO/CCUS Ag.	SPIUBROCOCCUS.	(<i>Sphaira</i> , orb, <i>kokkos</i> , fruit; theca round.)	23. — 128.
Mil renifl»mis Grev.	remform	broad leaves	4 aut Fi.R marine roc.
L412 rtibena Ag.	red	tufts	4 winter Psh ocean
J41J HrodiaO* Ag.	Brodic's	proliferous	4 su.aut Psh ocean
It 14 mcinbranifolius>4L	mcmbranous-lvd	branched	6 @ja R.P ocean
26445 Palmitta Ag.	Palmetto	flat branched	3 su Dl.P seashore
9A416 norv-gicus Ag.	Norwegian	little tufts	lftaut D.P marine roc.
96417 crlnpus Ag.	crisp	branched	4 s.my R.Br ocean
26418 mammillous Ae.	teatcd	branched	4 year R.Br ocean
26419 ciliatus Ag.	ciliated	flatlobed	G winter R.Br ocean
2 palmatus Ag.	palmated	flatlobed	6 winter R.Br sea shore
3fubutus Ag.	mand	finely cut	6 winter R.Br sea shore
4 angostus Ag.	narrow	finely cut	R winter II. Br sea shore
5 spinibus Ag.	spiny	finely cut	6 winter R.Br sea shore
96420 gigartinus /U.	grape-stone	coralloid	3 year R.Br seashore
&6W1 Tecdii ife.	Teed's	flat branched	4 Dp. 11 ocean
96422 crneus Ag.	horny	finely pinnat.	3 ■ dp.pk ocean
2 pinnatus Ag.	pinnated	finely pinnat.	3 ■ dp.pk sea shore
3 pulchillus Ag.	pretty	finely pinnat.	3 ■ dp.pk sea shore
4 Nrcfdeus Ag.	Nereid	finely pinnat.	3 ■ dp.pk sea shore
5 clavifer If Ag.	club-bearing	finely pinnat.	3 ■ dp.pk sea shore
6 clavatum J	clavate	entangled	I ■ K marine roc.
Ch6ndria pusilla Hook.			Lin. tr 3.17. 2. <i>Fucus</i>
96423 cartilagineous Ag.	cartilagineous	finely pinnat	8 year dl.br ocean
96424 coronopifulus Ag.	Buckhorn-lvd	rigid bushy	6 su Dp.R ocean
96425 laciniatus ite.	tagged	lobed	3 f.my Pk ocean
96426 bifidus Ag.	lid	blushy lobed	2 f.my p.nk sea shore
96427 cristatus Ag.	crested	small tuft	1 su Pk ocean
96428 confervoides Ag.	Conferva-like	much branch.	6 aut.w OLG ocean
CV29 plicatus Ag.	plaited	coarse bush	3 year Ol.Br ocean
96430 Griffiths** Ag.	A/rf.Griffiths's	coarse tuft	1* aut Br.R marine roc.
26431 acicularis Ag.	pointed	branched	3 aut Dl.R ocean
26432 purpur&scens ^g.	purpurascens	thinly branc.	6 year Pa.Y ocean
96433 Acheno'idcs Ag.	Lichen-lkic	thinly branc.	9 s II ocean
			Grev. cryp. 341
3067. HALYMETNIA Ag.	DULSE.	(///*, the sca, <i>hyemen</i> , a membrane j fronds.)	6. — 21.
26434 cdMis Ag.	eatable	broad leaves	8 ait Dl.lt ocean
2 media Ag.	intermediate	broad leaves	8 aut R sea shore
26435 palmata Ag.	palmate	broad leaves	8 o.ap psh.R ocean
2 marginifera Ag.	margin-bg	broad leaves	■ winter P ocean
3 sarniens Ag.	Guernsey	broad leaves	■ winter P sea shore
26436 ligulMa Ag.	tmm-shaped	lobed fronds	■ winter P ocean
SGt.ll furcelPita Ag.	small-forked	much lobed	■ winter P ocean
25438 Optunia Ag.	Indian fig	matted	■ sp.aut Pap ocean
26439 purpurascens Ag.	purpurascens	much branch.	6 su Pbh ocean
1 crispita Grev.	crisped	distorted	9 su dk.p sea shore
			Grev. cryp. 240
3068. BONNEMAISO^N// Ag.	BONNEMAISONIA.	(<i>M. Bonnmaison</i> , a French cryptogamic botanist) I-	
26440 asparagoWcs Ag.	Asparag-like	finely branc.	4 jLn Dl.pu sea shore
			Eng. bot 571. <i>Fucus</i>
3069. DELESSETR.M Ag.	DRLKTOERIA.	(<i>M. Benjamin Delessert</i> , a French bot. matron.)	11. — 24.
26441 sanguinea Ag.	YooA-colored	blushy	6 ja.my Ppk ocean
2ff422 sinuosa Ag.	indented	leafy tufts	6 ja.my Ppk ocean
WAW riwWwii Ag.	litHCiiflrmvrr	lint IOIMNI	7 su Fi.R ocean
'2M>V liy[M>gl <i>ii</i> Uiii Ag.	ton/fi/tf-uidcr-tonguuijlls		
26445 afata Ag.	winged	finely branc.	4 Ja.f Fi.p ocean
2 dilatata Ag.	dilated	finely branc.	3 Jns Fl.pk ocean
3 angUatiHsima Ag.	narrowest	finely branc.	6 ja.aut dp.pk ocean
26446 Plocamium Ag.	Plocamium	finely branc.	6 ja.aut dp.pk sea shore
26447 glandulbsa Ag.	glandular	finely branc.	4 su.aut dp.pk ocean
2f4H lacrata Ag.	torn	nearly simple	q alft Pa. II ocean
2H4J punetUto Ag.	dotted	very tender	4 su Bt.pk sea shore
26150 ocellata Ag.	spotted	very tender	5 su Fi.pk ocean
26451 Bonnmaisoni Ag.	Bonnmaison's	very tender	3 su I'i.pk ocean
			Grev. cryp. 322

TRIBE VI. FUCADEIE.

3070. LEMA^Nfi4 Ag.	LBMAHBA.	(<i>M. Leman</i> , a French botanist)	2. — 5.
26452 fluviatilis Ag.	river	lax tufts	6 su DLG stones in riv.
MM. « media Ag.	intermediate	lax tufts	6 su DIG rivers
2G453 torulbsa Ag.	torulosc	tufts	4 aut DLG rivers
			Eng. bot 1763. <i>Fucus</i>
			Ac. hoi. 1814.2. 1
-?fl CHORD A%R1 A Ag.	CIOBDARIA.	(<i>Chorde</i> , a cord j appearance of plants.)	1.-55
86454 flagelh6rmis Ag.	whip-shaped	long masses	24 su 01. G ocean
			Eng. bot 1222. <i>Fucus</i>

2072 SCYTO'SIPHON Ag.	SCYTOSIPIION. (<i>Skytos, leather, tiphon, tube; fronds coriaceous and tubular.</i>) 8.			
9 JUS 4f Jmrix Ag.	thread simple 240 su Br.Ol ocean Turn, fuci 86. <i>Fucus</i>			
» » » -i-iiuaum i*, /ccniculacus Ag.	hair simple 24 su Br.Ol ocean Stack, fuci 12. <i>Fucus</i>			
	downy simple 60 su Br.Ol sea shore Lyng, hyd. 19			
	<i>Fennel-leaved</i> simple 30 su Br.Ol ocean Turn.fuci XH.F.s&btiHs			
A- SPOROCHEMUS. (<i>Sporos, seed, chnos, wool; reproductive organs.</i>) 6. — 14.				
stalked downy 6 su L.G. seashore Eng. bot 545. <i>Fucus</i>				
acuteal much branc. 24 sp.su OLG ocean Turn, fuci 187. <i>Fucus</i>				
viridis 4* green very finely br. 18 su OLG ocean Eng. bot. 1659. <i>Fucus</i>				
vHK,BUg,r. rhi*,des Ag.	villous downy 6 su Pa.Y sea shore Eng. bot. 046 <i>Conferva</i>			
	rooty smth branch, 2 su Y.Br ocean Lyng, hyd. 13			
	larger smth.branch. 3 su Y.Br ocean Eng. bot.(J88 <i>C.vcrue.</i>			
	ligulate much branc. 48 su 01. G ocean Eng. bot 16J& <i>Fucus</i>			
HALISERIS. (<i>Hals, the sea, seris, lettuce; membranous fronds.</i>) 1. — 5.				
Polypodium-lk flat branch. 6 year OLG ocean Eng. bot 1758. F. <i>memo</i> .				
ENCO3LIUM. (<i>En, within, koilos, hollow; fronds tubular and bladdery.</i>) 2. — 4.				
blistered simple tubul. 6 su OLG sea coast Eng. bot 2570. U.Turn.				
Lmuan Grev. Lyngbyc's simple 12 aut OLG seashore Grev.cryp.290				
Iva Pstulbsa Eng. bot. 6*2.				
ZONARIA. (<i>Zone, a girdle j</i>				
peacock flatlobed 1 su Br.G ocean 4.-34.				
two-forked branched 4 su OLG ocean Eng. bot 127& U'va				
entangled branched 6 su OLG ocean Eng. bot 774. <i>U'lua</i>				
many-cleft flat cut 3 au Pu.Ol ocean Eng. bot 1913. <i>Viva</i>				
atom fan-leaved 5 su OLG seashore Eng. bot. 419. <i>Viva</i>				
Grev. STRIARIA. (<i>Stria, a groove; plants marked with.</i>) 1.				
iuata Grev. tapering branched 9 su OLG ocean Grev. cryp. 288. <i>Carm.</i>				
LAMINARIA. (<i>Sori upon the lamina of fronds.</i>) 8.—25.				
Agarum large masses 60 su Br ocean Turn, fuci 75. <i>Ficta</i>				
esculent large masses 60 su Br ocean Kng. bot 1759. <i>Ficut</i>				
digitate large masses 60 year OLG ocean Eng. bot 2274. <i>Fictus</i>				
bulbous large masses 60 year OLB ocean Eng. bot 170. <i>Fucus</i>				
saccharine large masses 48 year OLG ocean Turn, fuci 153. <i>Yicus</i>				
blistered large muses 48 year OLG ocean Eng. bot. 137ff. <i>F. sacch.</i>				
weak simple 8 su OLG ocean (Jrev. cryp. 277				
broad simple 16 su OLG ocean Fl. norv. 7.5				
Phyllitis simple 12 su BtG ocean Eng. bot. 1331. <i>Ffoitf</i>				
L1CHPNA LICHIVA.	(<i>Lichen, a lichen; resemblance.</i>) 2.			
Pygm rfa^. rgmy small patches fsu G.B. roc.onse.co. Grev.cryp.219				
confinis Ag. fhd small patches f G.B. roc.onse.ca Grev.cryp.221				
[0. FURCELLARIA Ag. FURCELLARIA. (<i>Furcella, a little fork; arrangement of fronds.</i>) L				
fastigiata Ag. peaked much branch. 9 aut. sp. II.Ol ocean Eng.bot824.F./>>iWd				
FUCUB. (<i>Phykos, a Greek name for sea-weed.</i>) 8—18.				
podusus L. knotty leatherybran. 36 d OLG ocean Eng. bot 570				
MacK'i Turn. Mackay's bladdercrd 24 d D.Br ocean Eng. bot 1927				
riffructus Ag. long-fruited loose masses 24 spsu OLG ocean Eng. bot 1056				
neAris Ag. linear loose masses 24 spnu OLG ocean				
J BubocostMus Ag. rather riblss loose masses 24 sp.su OLG ocean Esper fuel 146				
® "ateriffructus Grev. lateral-fruited short masses 2 su OLG salt marsh Grev. cryp. 181				
ceranpMes L. thunderbolt-like loose masses 30 su OLG ocean .Grev. cryp. 319				
	bushy 12 sp.su OLG ocean Eng. bot 2115			
	bushy 12 sp.su OLG ocean Turn, fuel 4			
	serrated masses 36 sp.aut U.OI ocean Eng. bot. 1921			
	channeled small masses 6 sp.aut Y.OI ocean Eng. bot. 823			
	warted branch, wart 6 jn Y.OI ocean Eng. bot 736			
	strap-Mr* masses 36 su D.OI ocean Eng. bot 569			
Ag. (Kystis, bladder, aelra, chain ; upper part of fronds.) 7.—45.				
Heath-like coralloid 6 su.aut 01.Ur ocean Eng. bot 1968. <i>Ficut</i>				
granulated bushy 12 spring OLG ocean Eng. bot 2169. <i>Fucus</i>				
bearded much branch. 6 su.aut Ol.Br ocean Eng. bot 2179. <i>Ficug</i>				
variable bushy 6 su.w OLG ocean Eng. bot 9151. <i>Fucus</i>				
southernwood bushy 14 spring OLG ocean Eng. bot 2130. <i>Fucus</i>				
fibrous bush, deform. 6 su.w Ol.Br ocean Eng. bot. 1969. <i>Fucus</i>				
podded loose masses 24 autsp D.OI ocean Turn, fuci 15». <i>Fucus</i>				
Himalier lnflMc mnwn 8 outup D.OI oconn Stack. Aid 11. ■				
naked loose masses 24 autsp D.OI ocean				

CLASS XXIV. ORDER 8.

CRYPTOGAMIA LICHENES.

THIEUI. IDIOTHA'LAMI.

(Spiloma, a spot; appearance of fructification.) 12.-S0.

tundish thin Am J year O bark of trees Kng. tot. 2151		
chaiigaeable-cld spotted crust 3 year Gy bark of trees Eng. bot. 2070		
small-branch, cloudy J year Wsh aged oaks Eng. bot. 2150		
black-faced eooty spots 2 year B apple trees Eng. tot. 2358		
obi. patches If year SeaG old boards Eng. bot. 239 <i>i</i>		
crust 3 year YBII old mortar Kng. bot. 2397		

26503 dUpIrsrum <i>E. B.</i>	scattered	even crust	1 year	Gy	old rails	Eng. bot. 2598
26504 decolorans <i>E. B.</i>	staining	lobed iwtcba	1 year	Gy	old wood	Eng. bot. 838i*
20505 punctatum <i>E. B.</i>	dotted	crust	2 year	Gy	old oaks	Eng. bot. £472
26506 varioloNUin <i>E. B.</i>	speckled	cracked crust	2 year	Wsh	old trees	Eng. bot. '2077
26307 auratum <i>E. B.</i>	gulden	tumid crowd.	If year	Wsh	old walls	Eng. bot. 2018
26508 tuberculsum <i>E. B.</i>	warted	even patch	3 year	Caes	sandst. rocks	Eng. bot 2556

3084. *SOLORINA Ach.* *SOLORINA.* {Solos, solid mass, rhinos, skin; texture of fronds.) 2.—10.
 26509 *croccea Ach.* yellow leafy frond If sp.su Ol.G tops of mou. Eng. bot. 498
 96510 *saccata Ach.* hollowed leafy frond 2 su Gыш on the earth Eng. bol. 288

3085.	LECI'DEA Ach.	LBCIDBA.	(/dfcia, saucer, riilos, like; shields.)	66. —183.
I. RBNIFÓRMES.— <i>Thallus crustaceous, reniform.</i>				
26511	atrocinnerea E. B.	I NIJBJE.—Apothecia constantly black, naked, not pruinose.		
26512	coracina Ach.	dark-grey close patches 1 year ISL rocks Eng. bot. 2096		
26513	atro-alba Ach.	raven tessellated 2 year Gy.B granite rocks Eng. bot. 2J35. L. corad.		
26514	flaco-Mra Ach.	black & white cracked crust 3 year Bl rocks Eng. bot. 180		
26515	fumata Ach. /lchm athrocarpus E. B. 1821)	brown-thin crust 2 year Bl rocks Eng. bot. 1734. Yudendril		
26516	lupicida Ach.	wnoky tessellated 3 su Br.Gy alpine rocks Eng. bot. 1830. Lar. Al./*		
26517	pifri'A Ach.	stone-splitting rock broad patches 3 year G brick walls Eng. bot. 821. L.a.J'fe*		
26518	cviil'Hucus Ach.	thin crust 1J year W roc. & stones Eng. bot. 245		
26519	parahema Ach.	confluent tartaricus 2 aut Gy.Br rocks Eng. bot. 194		
2(20)	sanguinaria Ach.	well-marked membranous 3 aut W hh bark of trees King. bot. 1450		
26521	obulvifruuin Ach.	hood-fruued gravel-pit rugose crust 2 year Whh rocks Eng. bot. 155		
26522	2 glichroa Ach.	carth-colored thin cuticle U year Wsh bark of trees		
26523	miscella dck.	mixed lobed crust 2 year Gy bark of trees Eng. bot. 1450. L.pat an-		
2(21)	clcharcAdcs E. B.	scar-like granul. crust 1/2 year Pa.Ol whinstocks Eng. bot. 1831		
2(22)	2 JfC aromatica jfc.*	aromatic lobed crust 1 year D.lir earth flocks King. bot. 1247		
2(23)	dolbba A.*:	deceiving broad cuticle 1 year Ol old walls King. bot. 1777		
26524	atro-virens Ach.	dark-green thin coat 1 year Ca's rocks Eng. bot. 2581		
2(24)	2 geographica jit/V	geographical figured crust 1 year B rocks		
26525	siltcea Ach.	yellow-dolored tessellated 1 year Y.Ol rocks King. hot 245		
26526	(E'dcri Ach.	USdr's membranous 2 year Y.R rocks Eng. bot. 1118		
MiWi	Alba Ach.	white tessellat.powd. 3 aut Wlsh rocks Eng. bot. 1117		
26528	citrinella jfcA.	cracked coat 3 spring Y.G sand, ground Eng. bot. 1877		
26529	uliginibia Ach.	marsh whole colored 3 spring B sandy heaths King. bot. 14G6		
26530	Jscabrbsa Ach.	Tuggyd-shield. lobed patches 2 year Pa.G tiled'roofs King. bot. 1878		
ii. MUTADILES.—Apothecia black, naked, when moistened becoming red or brown.				
26533	immersa Ach.	immersed even crust 4 yea! Pa.Y calcar. rocks King. bot. 193		
26534	rivulosa Ach.	rivulet broad incurbt 6 year Br.Ol rocks Eng. bot. 1737		
26335	allwvBcrufescens Ach.	iii. CINB REJf.—Apothecia black, with agicy bloom.		
2	tfrigida Ach.	A bluish white tartarc. crust 3 su Wsh Scotch alps		
26536	abitima Ach.	turgid simuated crust 3 su W bone walls King. bot. 820. L.tuit		
26537	Stcrica Ach.	Pine tree pruinoso 2 year GL trunks, 1/4" hics		
26538	vipolia Ach.	spiral simuated crust 4 spring W.Y. flint, pbebles King. bot. 1fffm		
26539	corticola Ach.	superficial tartarc. crust 2 su W Scotch ah,* Eng. bot. 117		
26540	couspuncia E. B.	bark-inhabiting small verruc. 1 aut COBS old trees Eng. bot. lb8<		
26340	defiled	defiled rimobc crust If Cas old walls King. bot. 964		
iv. RNV.—Apothecia blackishJirown, brownish, or deadened by some other color.				
26541	Lightfoot'Ach.	Lightfoot's oak sinuatu, patch, thin crust 3 year Pa.G smooth bark Eng. bot. 1451		
26542	outrencia Ach.	thin crust 1 year Y.G clefts of bark Eng. bot. 485		
26543	virid&ccns Ach.	thin crust 2 aut Pa.G dead trees		
26544	incana Ach.	hoary leprous 2 aut Gl trun. of trees Eng. bot. 1683		
26545	Hulnhiirca Ach.	sulphur cracked crust 2 aut Su rocks King. bot. UR6		
26546	orosthca Ach.	perennial toment crust 3 year LtG trees & pales Eng. bot. 1549		
26547	dcolobrins Ach.	discoloring granular 2 su Gysb on earth		
Z	2 granulosia Ach.	granular granular 2 su Gysb on earth Eng. bot. 1185. l^q		
26548	anomala Ach.	anomalous spotted patch. 3 aut Pa.Ol on earth Eng. bot. 2155.		
26549	ruprestrib Ach.	rock tartaricus 2 su Gysb rocks Eng. bot. 2243		
Lichen calvus E. B. 948				
26550	luteofola Ach.	yellowish thin crust 3 year Wsh bark of trees Eng. bot. 845. L.		
26551	carneola Ach.	horny-cupped papillobc crust 3 year Wsh on oaks King. bot. 95. I. rorn<%		
2	archentina Ach.	Jumper tree smooth coat 2 year W.Hrbark of trees King. bot. 1735.h.GrV		
26552	ftico-futca Ach.	yellow-brown thin crust 3 su Gysb mountains Eng. bot. 1007		
26553	cinerco-fasca Ach.	cinereous-brown cracked crust 3 year Gysb trun. of trees		
26554	anthraeoma Ach.	coal-like scaly crust 1 su D.Br rocks & trees Eng. bot. 432. L.ty*		
v. JfNRE.—Apothecia dark-red, or whitish flesh-color.				
26555	CliYio-Rufa Ach.	reddish grey tessellat.crust 3 su JXGY rocks & trees Eng. bot. 1650. L.		
26556	ichmadophila Ach.	moisture-loving leprous crust 2 year G.w on ear. in he. Eng. bot. 37^, L.		
26557	marmorea Ach.	marbled thincrust 3 year Gy.Whbark of trees Eng. bot. 739		
26558	alabastina Ach.	Alabaster thin crust 2 su Gy.W Scotland Eng. bot. J651.L.ro*/*M		
vi. hvVJL.—Apothecia pale yellowish, waxen, or orange-colored.				
26559	mcilzica Ach.	yc Wovi-shield. crackud crust // spring Y.Ol moss, truitks Eng. bot. 1263. L. /		
26560	EhrhartiINA Ach.	Khrh.ir'ta cartilng. crust 2 year Gsh rocks Eng. bot. 1136		
26561	pol'tropca Ach.	variable tessellated 2 year Pale rocks Eng. bot. 1134		
2a	2 fucida Ach.	shining soft crust 2 su G.Y rocks Eng. bot. 1550		
26563	atro-flava Ach.	dark-yellow ragged crust 2 year Ur expos, flints King. bot. 249		
26564	luteo-dlba Ach.	yellow-white smooth crust J * year W rocks Eng. bot. 1426		
II. LEPIUUMMA.— <i>Thallus crustaceous, of a regular flpiirc or leaf, not reniform.</i>				
26565	Candida Ach.	hpary sinuous 3 year Wsh old walls King. bot. 1138		
26566	vesicularis Ach.	blistered imbricated BU Br B 3 su High.rocks King. bot. 1139.L.c*-> n*jk		
26567	Ifrida Ach.	tfrid imbricated 3 su G.Ur Scotch alps Eng. bot. 13^9		
S6568	atro-rtifa Ach.	dark red imbricated 4 su Br red san. gro. Eng. bot. 1102		
26569	scakris Ach.	ladder imbricated 3 aut,w 1 aut,w 11 hardrockf King. bot. 2317		
26570	verruculbsa E. B.	small-warted irregular.patch. patches 2 spring Pa.G turfy earth King. bot. 2112		
26571	rubifermis Ach.	deceitful imbricated 2 spring F earth King. bot. 870		
26572	decimis Ach.	scaly leafy crust 4 spring COM quartz rocks King. bot. 1955JLg		
26573	phoudibta Ach.	smalUcaved broken patch. 2 spring Gy.G trees King. bot. 2128		

KHf canycens Ach.	canscent	round, patch,	If spring	W	bark of trees	Eng. bot. 582
*C76 dodalea E. B.	intricate	leafy lobed	spring	Y.G	rocks	Eng. bot. 2129
1. *? A LI, IUM Ach.	CALCIUM.	(<i>Kalykion</i> , a little cup; appearance of reproductive organs.)	17.—29.			
wn tigillare Ach.	rail	soft crust	2 aut	Gl	trees	Eng. bot. 1530
2f? SUKOMMum Ach.	small-dotted	pimpled	3 aut	Pa.Ol	on Lichens	King's bot. 2500.
2f? clavicular Ach.	small-headed	cloudystreaks	4 d	OLG	oak rails	Eng. bot. 1863
SKHO u>halumilicA.	club-headed	granul. crust	f aut	Gyh	naked wood	
2 ^ Q nypcrdlum Ach.	sphere-headed	thin crust	spring	Gyh	old pales	Eng. bot. 414
"ISRI b? cP. AlumicA.	yellow-headed	irregular coat	2 spring	BT.G	old oak	Eng. bot. 1832
2f? L A ab nll^m **.	small beam	atches	aut	I.cm	trun. of trees	Eng. bot. 2501
	small rafter	dense granul.	aut	Br.Ol	boards	Eng. bot. 1540
	small-headed	obscure crust	aut	Wsh	decay, wood	Eng. bot. 2537
	acicular	irregul. patch,		G.Y	sandy soil	Eng. bot. 1539
	rusty	irrcg. incrust	3 su	Ol	Scotch lira	Eng. bot. 2485
	fouCnlobbed	lobed crust	aut	Pa.Ol	pales	Eng. bot. 2473
	dewy	tesnelleratrust	aut	W.Br	dead wood	Eng. bot. 810
	weak	mealy coat	year	G	old boards	Eng. bot. 1464
	verdigrise	cloe-set pate.	aut	Br	old timber	Eng. bot. 2412
ruumoHUm E. B.	ahort-stalked	granular	aut	D1 G	old boards	Eng. bot. 2502
curium E. B.	rrowd. patch.	winter DIG	decay, wood	Eng. bot. 2503		
k 1 L 115 > PHORA Ach. G IIIWIOWIK A						
	smooth	leafy thallus	2 su	i phorcip	to bear; in disk of shields.)	8.—19.
	many-leaved	leafy thallus	2 su	D.OI	rocks	
	proboscis-like	netted frond	3 spring	Smo	rocks	Eng. bot. 1283
	arctic	smooth lobed	14 winter	Br	rocks	Eng. bot. 24H3
	cylindrical	folded frond	2 spring	Cir.Ol	rocks	Eng. bot. 522
	gnawed	ragged	3 year	OLBrrouks		Eng. bot. 2066
	scorched	rough leafy	3 year	Ol. Mr	rocks	Eng. bot. 2483
	pimpled	blistered frond	2 spring	Cm (J	rocks	Eng. bot. 1283
	furred	shunted	2 year	Gy.Brtnek*		Eng. bot. 5131
Aeh.	irregular lob.	irregular lob.	1 year	Br	rocks	Eng. bot. 2483
24. 8. ENDOCA'RPON Ach. END.						
>> nopicum Ach.	red	{ <i>Endon</i> , within, <i>karpos</i> , fruit; recept. embedded in leaf.)	10.—22.			
amaragdulum Ach.	emerald	tessellatmass.	1 su	O	schist	Eng. bot. 17h*
tcplofdci Ach.		little patches	f su	Y.G	rocks	Eng. bot. 1512
'lchen fuscillu		little patches	1 su	Gl	earth	Eng. bot. 2013
Hedwigif Ach.		crowd, patch,	J su	Ol	on the earth	Eng. bot. 51> htrapexif.
palldiu A-A		crowd, patch,	D G	on the earth		Eng. bot. 1698
		finely lobed	f year	Pa.Ol	rocks	Eng. bot. 2541
		round, patch,	18 U	Cop	on Lichens	Eng. bot. 1866
		thick crust	1f sprig	Gyh	nicks	Eng. bot. &3
		round patches	Br	rocks		Eng. bot. 2012
		coriaceous	f year	Gyh	rocks	Eng. bot. 593A L.atnpM.
		cartilaginous	f w.sp	G.Br	wet rocks	Eng. bot. 594. L.aqidic.

TUIDE II. CCENOThA'LAMI.

R < Y R ? , NU LA ilc A. PVRBNLLA		(Dimin. <i>otpyrcn</i> , a kernel; receptacle in thallium.)	2 4—34.
	nigrescent	tartarebus	If year
	tessellated	circular dots	2 year
	uoibonAta Ach.	even coat	1 & year
2@ Vn Y ^ R ^ LA ARIA Ach. VARIOLARIA.	veiled	sinuous surf.	1 aut.
	much-dotted	granular	2 winter
	globuliferous	uneven crust	\ year
	common	radiated	II year
	sprinkled	radiated	1} year
	bitter	cracked crust	2 year
	discoid	pulverulent	2
	milky	tartar, crack,	W winter
	greysih green	tubercular	14 aut
	whitened	cracked crust	3 year
	cinerous	tubercular	2 year
Acharius's		cracked crust	2 year
convex-shielded		tessell. crust	2 year
		fringedpatch	3 year
		concent. zones	3 year
		solid crust	1 year
		obscure crust	1J year
		crowd, warts	U year
		sinuous pate.	W year
			Cats roc. & stones
			Eng. bot. 1940
Ach. LE CANORA. (Lekanc, a basin; form of shields.)			46.—79.
I. RINODI'NJS.— <i>Thallus adnate, uniform,</i>			
i. KISRA.— <i>Disk of apothecia constantly dark or black.</i>			
dark	granulated	\ year	Gyh bark of trees
white-scaled	warted crust	2 su	Pa rocks
eyed	oinooth. crust	2 bpring	W roc. & earth
compressed	brood patches	4 year	Br brick walls
			Eng. bot. 949
			Eng. bot. 1833
			Eng. bot. 1034

718 diatrypa	bbddety	mulrtC	«mth.	2	Will	rock	Bag. bot	ISS
	pierced through	muki I	MM!L.	2	Gy.	wet rock*		
BO'RRERJ	B»BEKA.	{William Borrr. Y. L. S.,)	7. —23.
	•lender	branch, tcfn	1} year		hrjti. oftrrv*	EtiK-	boL	1351
	black and white	den*e tufts			i-jilin cirtli	l-jig.	IHIL.	SS18
,,,ji ialiiM	JcA. golden-eyed	buihy		li year	i-H> trun.	uf trees	Eiig.	boL 964
u	y-tlowUh	branched	I	yew	O	apple Irev*	Eng.	bot 1088
* atlantica vteft.	Atlantic	b)	taffii	II year	Y	trun. ortrces	Ens	hot 2113
		Ij	ap	Gl	trun. of trees	Eng.	Ivet	
				0.01	dull	Eug.	bat.	1715

(Cefrxt, a buckler; rece)

&.—14.

Junijior	bujby							
	tw*hy		11	year				
hedge-mhablting	btuhjr	waved			Ol.Br	st»n.		
guucoui		• shining		year	Gl	mn Che groo,	Eng.	bot. 1606
CHUJUH:				year	W	on the grou.	Eog.	bot £373
• UW		hjr tuft*		yemr	Su	rocks	Eng.	bot US*
Iceland MiHt	bu>h,		yc*r		OUBr	rocky place*	Eng.	bot 1330
SrtCT*.	ISstiMlo*	dotted	: under	jurfaw of fronds.)			7. —18.	
	ydlow	waiti	3	year	GI.Brrrock*	Eng. not		
gililod			6	year	Br	trun. oftree*	bng.	bot 9SSB
lung		reticulated		year	Oliv.i	truu.oftn		
tin—furrowed	roun Hah pat		ye>r		tiy*h	trun. of tree*	Eng.	bit'.
bordered	nipoCfl bfi MKI		year		GtBrrckf	E>g		
•tnutty	round patch,		year		X<ugy	moitt riM'ks	Eng.	bot
wood	pitted i		Vi>ir			hady wood*	Eng.	bot. SttA

uiKi.	(Pdte, • targt, f*rflo,	Twcooblwire; form	of thirfdi]	p. — .L
vctny	2 (U	Qth onthi'tMtlh	Fjit-	
ted	11 year	Cin bark of In-		
horiuantal	2 year	Br.G thadyrock-		
mrtjimm ilorv	2 aut	G ainong nn u	Eng. Ixm	
rufnevnt	iijcurred	ILBr on tin-cnri ti	Hug. bot. 2300	
membnuiotu	bmad4obed	2 year	Gy>h on tb>	.,. bot. S299
*purtou>	lobedfroad	1(jl	Ol.Br thatch	Eng. bot 1512
many. Angered	siuixh-hood	IJ jl	Gl on the earth	Jac c. :

6 polyductyla' Ach.	cA. Ksnikoiu.	(AVpArtu, a kidney; fonn of apotheci.)	£. —B.
NEPHHO\MA	renipinate	thorUtobcd	Gy. Bramong mou
	equal	foUaccom	Fs'l g. bot. 105

*8 nirllii Ach.			Eng. boL ftjfio
RfiCCETLLA Ach.			

>linct.ria Ach.	Iruc dyer'i	{Rocka, a rock, Port. ; place of growth.)	
Ick.	Futut'-funned	bu*hy tul»	1} year

EIEIMI.	t, well branched ; buihy plasta.)	1.-6.
Prun4*tri Ach.	PruoMter	inuiiii ¹ . tegm. 2 year G.W tree*
I «Jrtocro* Eng. bot. 13S3.		

Ach CKMOMVCE. (A>»t, empty, wpto*, fungui; hollow receptacle.) SO.— 43,

Pi i sv,i m;Yi i.— ThatJut mlcnutaccouM, uniform; vodetja hollow.

Ach, pimpled graDuL emit } winter Gjih uamp earth Eng. bot 907

SevfBo'ryob* — Thnillit fitildctmu ; podetia fiutular, dilattd upwards and Jertii; ur turtle and tabulate i apotkeda cbmd aM a membrane.

tCM. — Apatkcia Aunui or pallid.

til born	luiliifid tuft»		dry placet	Enjj bntKAit
StagVh»,rn	mult.ftl tufti	1J	rocky beatbi	Eng. be
lift*	tuiU		bank*	Eng. bot 1399
fringed	corallloid tuft.	1	noon ft hea.	Eng. bat M38
rw	corallotil tuft*	S	on the trou.	Eng. bol. 1835
eushil-n-sprouting	coraloid tu(U		moon ft IIM.	Eng. boL IMS
iii.in:-laf	brittle tuft*	1	Lin raounalot	
imitative	fine tuftj		hiUi Eat. bot 1807	
iltt	line tuftJ	<	Gy hn.fr raoun.	
			hea.& mouu. Eng toot. ISSt	
VOCCTXEIL — Ap	tcarttt or dari red.			
rod-ttk	brmc	3 y»« Wth woodi	t	
finaered	pow.lery	1(year Y.G wood*	Eng. boi. ?*»	
!(icd	branch, tuft*	3 ye-	rouuoirtt«* Eng. bt-	
berry-bearing	lmp.	3 winter Gy G moor*ft h*a. fcng. bot SW1		
Rolde*, cA cornuojHj-lk	thort tuft*	1 winter Or.G moon 4 hea.		
i Ach.	Hif.	Jc Mly nrnin. Ear bot. 181*		
nt — TkallmifOiaettM*; pylHajit tujgr, dilated upward*, and ftttUi; apotketa prrtiouu				
ML ttrrn	blanch, tuft, it	year Gl to wooi	Eng!buL 2TT	

drtk-ate	diptott i		r G	ratten ra i It Eng. bot SOSS
	mealy patch		eartilofimtiu,	rigid,JUlar, aU tapriuu* tuluteU
		If bordthrough	G*h	h<ath*
				Di. mtMC. l'i, £5
•UbulfC	hlybrS.	1	Ld.br	
Inch	r>id	winter Pa.G	moor*	16. Q
		S year Hoa	wood*	Eng. bot 174
	i-d tu/l	S ysar Gy	commc	Eng. bot S77
		tild, tubulate, tomeiahjit	branched	E>ff. bot 9<M

3103. BJEO'MYCES Ach. BJSOMYCES. (*Baios*, small, *mykes*, fungus; appearance of fructification.) 4.—10.
 26772 rōseus Ach. rosy granulated \ su Gsh heaths Eng. bot 374
 26773 ruful Ach. rufbus powdery \ su Gsh sandstone Eng. bot 373.6.ywol<fc*
 26774 microphyllus f. 2?. small-leaved imbri. patch. 3 winter D.G wet heaths Eug. bot. 1782
 26775 ca»pitifl. A. turfy leafy tuft 3 aut Pa. G oaks Eng. bot 1796
3104. ISI'DIUM Ach. ISIDIUM. (*ISOS*, equal; small differences which exist between the podetia.) 5.—11.
 26776 microsticticumflo*. small,dotted tartareous £ aut Brsh rocks Eng. bot 2243
 26777 corallinum Ach. coralline crowded pate. £ aut Gysk rocks Eng. bot 1541
 26778 Westringi Ach. Westring'a cracked crust \ aut Gysk rocks Eng. bot 2204
 26779 ihmatus Ach. bladdery powdery crust 3 winter Pa.Su stems,old tr.
 2 phragmoseum Ach. bushy powdery crust 3 winter Y.OI stems, old tr. Eng. bot. 1529. L. lutis*
 26780 coebes^cA. berried powdery crust 2 aut Pa.OI park pales Eng. bot 1511
3105. STEREOCAUXON Ach. STEREOCAULON. (*Stereos*, rigid, *kaulon*, a stem.) 1.—6.
 26781 paschale Ach. Easier branch, tufts 2 year Gysk mountains Eng. bot 282
3106. SPHiERO'PHORON Ach. SPILE a. (*Sphaira*, globe, *phoro*, to bear; form of fructification.) 3.—14.
 26782 coraloides Ach. coraUlike bushy 1* year Pa. Br rocks Eng. bot. 115
 26783 fragil Ach. brittle bushy 1 year Gysk rocks Eng. bot 2474
 26784 compressum Ach. compressed bushy 1 year Wsh rocks Eng. bot. 114. L fr&gw*

TRIBE III. HOMOTHA'LAMI.

3107. ALECTCTRIA Ach. ALISCTORIA. (*Alektor*, unmarried; uncertainty respecting male flowers.) 2—7«
 26785 jubata Ach. maned long tufts 3 winter Br on fir trees Eng. bot 1880
 2 chalybiformis Ach. steel-formed long tufts 3 winter Gy.B on fir trees
 26786 sarmentosa Ach. twiggy much branch. 2J winter Pa. Y mountains Eng. bot 2040

3108. RAMALPNA Ach. RAMALINA. (*Ramale*, a withered branch; habitat.¹) 5.—19.
 26787 /raxinea Ach. Ash-like loose tufts 2 year Gysk bran, of trees Eng. bot 1781
 26788 fastigiatä Ach. peaked loose tufts 2 year Gl rocks & trees Eng. bot. 890
 2 calfurta Ach. calyx-like loose tufts 1J year Gl rocks & trees
 26789 scopulifrum Ach. rocks loose tufts 11 year Y.G marine roc. Eng. bot. 688
 26790 farinacea Ach. • mealy bushy tufts 2 year Gysh trun. of trees Eng. bot. 889
 26791 pollinaria Ach. powdery bushy patch 3 year Bt.G old oaks Eng. bot. 1607

3109. CORNICULA'RIA Ach. CORNICULARIA. (*Corniculus*, a little horn; divisions of thallus.) 7.—I⁶*
 26793 tristis Ach. dingy shrubby 1| year Dp.Br alpine rocks Eng. bot. 720
 26793 aculeata Ach. prickly shrubby 1 year Ch Highl. mou.
 2 spadicea Ach. scarlet shrubby 1 year Ch Highl. mou. Eng. bot 452. "L.hisp W*
 26794 bicolor Ach. two-colored shrubby 1 year B Highl. mou. Eng. bot. 1853
 26795 orhroleuca Ach. pale-yellow shrubby 1} year Pa.Y Highl. mou. Eng. bot 2374
 26797 pubescens Ach. woolly shrubby f year Gy.B rocks Eng. bot. 846
 26797 pubescens Ach. pubescent entangl. tufts 3 aut B rocks Eng. bot. 2318
 26798 heteromalla E. B. wrious-woolloed rough patch 3 aut B bark of trees Eng. bot 2246

3110. U'SNEA Ach. USNEA. (*Achne*, lichens, Arab.) 3.—10.
 26799 fMrida Ach. flowering erect 2) winter Gsh old trees Eng. bot. 872
 26800 phæta Ach. plaited pendulous 4 winter Gsh old trees Eng. bot. 257
 2 hirta Ach. hairy nearly erect 2 winter Gsh old trees Eng. bot. 1354
 26801 barbata Ach. bearded pendulous 4 winter Gsh old trees Eng. bot. 258.2
 2 articulata Ach. Jointed pendulous 4 winter Gsh old trees Eng. bot. 258.1

Sill. COLLEGIA Ach. COLLEMA. (*KoUema*, glue; fronds gelatinous.) 27.—41.

- I. PLACYNTIUM. — *Thallus* crust-like, irregular or uniform.
 26802 nigrum Ach. black regular patch 3 wet w B.G calcar. rocks Eng. bot 1161
 II. ENCHYLIUM. — *Thallus* imbricated, plaited, roundish, composed of minute lobes, becoming very turgid when wet.

- 26803 rhelleum Ach. lipped round, patch 1\$ wet w B.G roots of trees
 26804 fragrans Ach. fragrant small patches i wet w D.OI trun. of elms Eng. bot 1912
 26805 cricpum Ach. crisp round patch, i wet w Gl on the grou. Eng. bot 834
 26306 tenax Ach. tough lobed tuft 1 wet w G moist places Eng. bot 2349
 26807 plictile Ach. plaited lobed tuft 1 wet w 01. G wet rocks Eng. bot. 2348
 S6808 fluvile Ach. river many-parted * wet w Br calcar. rocks Eng. bot. 2039
 26809 mcl u:ium Ach. blackish starry * wet w Br rocks
 2 margin-lie Ach. marginal imbrife. lobes 1 su Ol Highlands Eng. bot 1924
 2GS10 fascicular Ach. fascicled roundish 2 aut.w Br trun. of trees Eng. bot. 1162
 26811 creticeum Ach. cretaceous minute dots * & winter Br chalk stones Eng. bot 738
 26812 corrugatum Ach. wrinkled small patches \$ winter D.G rocks, sea co. Di. muse. 19.19

III. SCYTI'NIUM. — *Thallus* somewhat foliaceous, irregular, formed of naked, expanded, thick, turgid* naked lobes.

- 26813 palmatum Ach. palinated lobed patch 1 sp. su Br sand ground Eng. bot. 1635
 2(814 granulatum E. B. granular imbrrie. patch 1} wet w Br gravel walks Eng. bot. 1757
 26815 multipartum E.B. many-parted lobed patch 3 su OLG rocks & walls Eng. bot. 2582

IV. MALIA.THIM. — *Thallus* foliaceous lobes rounded, downy, or fibrous beneath.

- 26816 saturnnum Ach. dingy leafy 2 year B.G trun. of trees Eng. bot. 1930
 26817 Burgess's Ach. Burgess's leafy 2 year Gl trun. of trees Eng. bot. 300

V. LATTHIA\JRIUM. — *Thallus* foliaceous ; lobes somewhat membranous, lax, naked, dark green.

- 26818 nigro'cens Ach. nigrescent leafy 2 year D.G trun. of trees Eng. bot 345
 26819 flccidum Ach. flaccid leafy smooth 2 year D.G Scotland Eng. bot 1J>g
 26830 flirvum Ach. dusky rugose memb. 2 year D.G trun. of trees Eng. bot. 1/37
 26821 scotnum Ach. dark flat patches 1* su Ol old walls
 2 sinuatum Ach. sinuous flat patches 1J su Ol old walls Eng. bot 77*

VI. LEPTOUIUM.— *Thallus* foliaceous ; lobes rounded, membranous, thin, naked, cinereous, glaucoevs, somewhat transparent; apothecia slightly stalked.

- 26822 tremelldes Ach. Tremella-like half transpar. 1 spring Lead rocks Eng. bot. IS^
 26823 lacerum Ach. lacerated half transpar. 1} spring Gl earth Eng. bot. 1»»
 OCQO* , >. VII. Mra'coLUM. — *Thallus* very finely ladiated and branchlctted. ^.
 26824 subfile Ach. btle starry 14.su D.G earth Eng. bot. 1008
 ESS ft- Jmml11 Ach. finest flat patch 2. jl. au D.OI dry banks Eng. bot. 1077
 2f826 behradcri Ach. Schrader's small tufts * jn Y.G old walls Eng. bot.

S§. »*U8clcola Ach.* Moss-dwelling cushion-like § spring Br among moss Eng. bot. 22ii4
Pongium Ach. spongy large fruit 3 year 01. Br rocks Eng. bot. 1374

TRIBE IV. ATHALAMI.

	LEPRARIA.	(<i>Lepra</i> , leprosy; scurfy crust.)	4.—13
85	pale	cushion-like 2 winter Su rocks	Eng. bot. 8&J
K Ach.	ochracea E. B.	thin coat 2 winter Bt.Y old pales	Kng. bot. 1350
virescens B. it	virescent	scatter, warts \ winter G.Y old trees	Eng. bot. 2408
		granular winter Y.G elm trees	Eng. bot. 2149

TRIBE V. PSEITDO-LICHETNES.

3113. OPB/GRAPHIA Ach. OPEG. (*Ope*, a chink, *grapho*, to write; shields of apoth. crack-like.) 10 —35.

• HVSTERIVIA. —Disk of apotheca very narrow, crack-like, somewhat covered in by the conniving

	tumid margins.			
E.B.	cloudy variegated	1 year	Pa.Y old trees	Eng. bot. 2346
Ach.	veiny flat patch	U year	na.och beeches	Eng. bot 2454
	Persoon's tartaceous	\ year	Wsh stones	
	poreless leprous	2 year	Wsh slate & stones	
	calcareous angular dots	& year	B mort, old w. Eng. bot 1790	
Jjerpetica Ach.	spotted	largish spots	Brsh birk of trees Eng. bot. 22K2, <i>O. epiphe</i> .	
epipasta Ach.	eruptive	dotted crust	I'a.Ol bark of trees Enj. bot. 1789	
	different	mealy crust	Pa.Ol bark of Irccs King. hot. 2347, <i>O. rubella</i>	
	common	scaly	G.W bark of trees King. bot. 1811	
epipasta Ach.	sprinkled	smooth skin	Gy smooth bark King. bot. 1828	
2 microscopica Ach.	microscopic	smooth skin	Ol smooth bark King. bot. 1911	
Stomigrata Ach.	narrow-fruited	smooth patch.	Ol smooth bark Eng. bot. 1753	
	blackened	smooth patch.	Pa.G smooth bark Eng. bot. 1753	
• ALYXURIA. — Disk of apotheca concave channelled, or flat appearing between the separated margins.				
nótha Ach.	spurious	dotted crust	Wsh old trees Eng. bot. 1896	
2 diaphora Ach.	various-fruited	3 year	Gy trun. of trees Eng. bot. 2280	

3114. VERRUC A'RI A Ach. VERRUCARIA. (*Verruca*, a wart j warty nature of shields) 11.—56.

I. LICOPHLEA.—*Thallus cartilaginous, membranous, contiguous, polished.*

mafira Ach.	Moorish	cracked crust	2 year	JJ rocks	Eng. bot 2156
44 punctiformis Ach.	dot-shaped	thin coat	2 year	Br sm. ash bark Eng. bot 2412	
45 analcpta Ach.	not oily	thin coat	S year	Br sm. oak bark Eng. bot. 1848	
46 epidérmidis Ach.	cuticle	thin coat	1 year	W birch bark	
847 stigmatella Ach.	small-dotted	thin coat	3 year	Pa.Br smooth bark Eng. bot. 1891	

II. BLENNORIVIA—*Thallus nearly solid, somewhat gelatinous.*

268 48 ceuthocarpa Ach.	hidden-fruited	tessellated	4 year	Pa.Ol slate rocks	Eng. bot 2372
-------------------------	----------------	-------------	--------	-------------------	---------------

• LITHOCIA. — <i>Thallus subtarlaceous, crustaceous, contiguous, cracked into areoUe, or powdery.</i>					
Jir&deri Ach.	Schrader's	dotted crust	4 year	Wsh calca. stones	Eng. bot. 1711
Harmannii Ach.	Harmann's	small patches	1 year	Br.Ol hard rocks	Eng. bot. £539
pltimbea Ach.	lead-colored	lobcel patches	1\$ year	Ol limest rocks	Eng. bot 2540
striatula Ach.	striated	cloudy spots	\ year	Pa.G flints	
2 acrotella Ach.	small-peaked	cloudy spots	f year	Pa.G flints	Eng. bot. 1712

IV. INODE'RMA. — *Thallus soft, cottony, somewhat spongy, or thin and arachnoid.*

ground	mealy tessell.	1 year	G dry banks	Eng. bot 681.L.ferr&J,
--------	----------------	---------	-------------	------------------------

• PORINA Ach.	PORINA. (<i>Porinos</i> , any thing that crumbles away; nature.) 1. —19.			
*»» Pertbsa Ach.	perforated crust	i aut	Ciu bark of trees	Eng. bot 677

	ARTHOXIAX.	(Unexplained)	4. - r 14.
IB.	unpolished	spotted patch	1 year
Ach.	Swartz's	cracked crust	1} year
	star-like	membranous	\ year
	obscure	warty	3 year
	lynx-spotted	broad masses	3 year

	GRAPHIS.	(<i>Grapho</i> , to write ; apothecia like writing.)	5. — Id.
Jcr.Jpta Ach.	written	shining crust	1J year
* Pulverulenta Ach.	powdery	thin crust	Pa.Y trees
2 Cerasi Ach.	Cherry tree	thin crust	Y old chcr. tre. Eng. bot 2301
is Ach.	tree-like	smooth patch.	Y smooth bark Eng. bot. 1756
	serpentine	even crust	Pa.01 smooth bark Eng. bot. 1754
	LylePs	cracked crust	Pa.Ol rugged bark Eng. bot. INTI
	elegant	uneven crust	Pa.Y smooth bark Eng. bot. 1812

CLASS XXIV. ORDER 9.

CRYPTOGAMIA FU'NGI.

TRIBE I. HYMENOMYCFTES.

Class I. HYMENIVI. — Division I. PUedti.

• ^GA'RICUS L. AGARIC. (*Agaria*, a region of Sarmatia.) 314. — 715.

I. VOLVAYEI.—*Stem central, with a veil; gills unchangeable ; sporidia white,*

h 1. AMANI'TA Pers. — *Veil double, universal, separate, partial, annular, somewhat persistent*

i. — *Volva loose j edge of cap smooth ; unwholesome.*

vernal	fetid	«-6 sp su	W woody dam.pl. Bui. her 108
phallus-like	scentless	4 jl.o	W woody places Bui.her.2.577 bidbhsus
- verruebus Fl. Ion. warty	scentless	4 jl.o	Y woody places Fl. Ion 3» d 
-(Sri? ? Virg8Ce)n8 W. dan. virescent	scentless	4 jl.o	Pa.G woody places Fl. daai 1246
«<7 porph^rius Fries puq;le	scentless	4 jl.o	Ld among moss Mic"gen"76?3

	ii. — <i>Volva loose; edge of cap striated; eatable.</i>
26868 <i>vaginatus Bui.</i>	sheathed eatable 6 aut W waste places Bui. her. 512
1 <i>vimbicus Schaff.</i>	<i>cad-colored</i> eatable 6 aut Lead waste places Schaff. 85. 86
2 <i>hyahnus Schaff.</i>	transparent eatable 6 su Css waste places Schaff. 244
3 <i>pulvinatus Bol.</i>	cushioned eatable 6 su Br waste places Bolt. pit. 49
4 <i>fulvus Schaff.</i>	tawny eatable 6 su Taw waste places Bolt. pil. 38.2. <i>trUobatu*</i>
26869 <i>nivalis Grev.</i>	snow delicate 5 au W Scotch mou. Grev. cryp. 1.18

	iii. — <i>Volva obliterated; edge of cap striated; poisonous.</i>
28870 <i>muscarius L.</i>	fly-Mown poisonous 8 au.o O.R woods Grey. cryp. 1.54
26871 <i>pantherlini Dec.</i>	panther-like warded 3 au.o 01 moun.woods Schaff. 90. <i>maculatus</i>
26872 <i>rubescens Pers.</i>	rubescent nauseous 3 jl.s F. Col heaths Schaff. 91. <i>pustulatus</i>
26873' <i>toper A. #S.</i>	rough fetid 3 jl.o Rah open woods Bui. her. 31G. <i>verrucosus</i> U\$

12. LEPI&TA Pers. — Veil simple, universal, concrete, annular, somewhat persistent.

	i. — <i>Veil finally separated; gills distinct; eatable.</i>
86874 <i>procérus Sco.</i>	tall esculent 10 au.n W.Br woods Sow. 190
26875 <i>excoriatus Schaff.</i>	flayed esculent 7 jl.au Wsh fields Scheff. 18.19
	ii. — <i>Veil fixed; skin of the cap peeling off; gills separate.</i>
26876 <i>clypeolarius Bui.</i>	buckler insipid 2 au.o Wsh beech woods Sow. 14.
2 <i>fellinus Pers.</i>	cat insipid 2 au.o Wsh hot-houses
3 <i>Meliagris Sow.</i>	Guinea-hen insipid 2 au.o Wsh pine woods Sow. 171
26877 <i>cristatus Bol.</i>	crested fetid 1 au.n Wsh grassy places Grev. cryp. 3. 176
	iii. — <i>Veil fixed; gills separate; skin of the cap adhering.</i>
26878 <i>illinUus Fries</i>	besmeared mucilaginous 3 jl.o Wbh meadows Fl. dan. 600
	iv. — <i>Veil fixed; cap covered; gills somewhat united.</i>
26879 <i>granulbus Pers.</i>	granular muricated 2 jl.d Y heaths Grev. cryp. 2.104

13. ARMILLA'RIA Fries. — Veil simple, partial, separate, annular, somewhat persistent.

	i. — <i>(UspUose; cap smooth.</i>
26880 <i>racidus Schr.</i>	musty glutinous 2 jl.d W old trees Fl. dan. 773. <i>niidus</i>
	ii. — <i>Caspitose; cap not smooth.</i>
26881 <i>milieus Bol.</i>	<i>honey-colored</i> esculent 4 au.o DI Y trun. of trees Grev. cryp. 332
2 <i>Zaricus Bol.</i>	Larch-like esculent 4 au.o DI.Y trun. of trees Bolt. pil. 19
3 <i>elisticus Bol.</i>	elastic esculent 4 au.o DI.Y trun. of trees Bolt. pil. 15

14. LIMA*CIUM Fries. — Veil very fugacious, viscid; lamellae adnate, decurrent.

	L — <i>Cap smooth, floccose at edge; unwholesome.</i>
26882 <i>chrysodon Batsch</i>	golden.toothed noxious fs. o Fa.Y beech woods Batsch fun. 2.212
26883 <i>carneus Sow.</i>	fleshy noxious 3 s.o Pa.pk among grass Sow. 246
	ii — <i>Cap smooth; stalk scaly j eatable.</i>
26884 <i>ebfimeus Bui.</i>	ivory shining 4 au.n T woods Sow. 71. <i>nitens</i>
2 <i>nitens With.</i>	shining shining 4 au.n T woods Sow. 121. <i>cussus</i>
	iii. — <i>Cap finally depressed; stalk spotted.</i>
26885 <i>olivaceo-a'llbus Fries</i>	olive-white viscid 4 jl.o Ol pine woods Schaeff. 312. <i>limaculus</i>
26886 <i>hypothecus Fries</i>	sub-sulphur-cr/ clustered 4 o.d Ysh heaths Sow. 8. <i>limachius</i>
26887 <i>aromaticus Sow.</i>	aromatic glutinous 3 o.d Pa. Br woods Sow. 144

% 5. TaincoLBMA Fries. — Veil very fugacious, fiocculose, marginal; lamella emarginate or rounded.

i. — Cap truly fleshy, somewhat blunt, humid, viscid, with an involute downy edge; gills white or yellow, emarginate; stalk clothed, separate from the cap.

26888 <i>elbo-brtinneus Pers.</i>	whitish-brown glutinous 3 au. o Br fir leaves Schaeff. 58. <i>striatus</i>
26889 <i>fulvus Dec.</i>	tawny smells of flour 4 au.s Taw thickets Schaeff. 63. <i>incirtus</i>
26890 <i>usUlis Fries</i>	scorched scentless 3 au.o R.Br beechwo.&c.
26891 <i>RAssula Schaff.</i>	Russula delicious 2 aut Pk woods Schaeff. 58
26892 <i>aur&ntius Schaff.</i>	orange bitter 3 aut O pine woods Schaeff. 37
26893 <i>prásinus Schaff.</i>	leek-green tuberous 3 aut Y.G mossy places Schaeff. 218
26894 <i>fucatus Fries</i>	painted mild 2 s.o Lu way sides
26895 <i>luridus Schaff.</i>	lurid gregarious 2 s.o Di.Ii pine woods Schaeff. 69

ii. — Cap always dry, scaly, with the young edge involute, downy or villous; gills separate or emarginate i stalk scaly, separate from cap.

26896 <i>equestrus L.</i>	horse mild 2 s.d Y.Br way sides Schaeff. 41. <i>aureus</i>
26897 <i>rutilans Schaff.</i>	glittering splendid 3 au.o Y roots of trees Sow. 31. <i>xerampetinw</i>
26898 <i>vacculus Schaff.</i>	cow scaly 3 o.d Ru damp places Schaeff. 25
26899 <i>Myomycetes Pers.</i>	Myomycos smells of mice 3 o.n Ld plantations Sow. 76. <i>tirreus</i>
26900 <i>columbeta Bauh.</i>	small pigeon eatable 1} au.o W sandy places Bui. her. 428.1

iii. — Cap always dry, smooth, but often fibrillose, with a naked edge; gills separate or emarginate; solid, smooth, striated, separate from the cap.

26901 <i>a'stuans Fries</i>	burning very acrid 3 au.s Y among moss
26902 <i>scjdncutus Sow.</i>	disjoined bitter 3 s.o Pa.Y dry pine wo. Sow. 126
26903 <i>virgatus Fries</i>	streaked bitter 3 s Gy plantations
	iv. — Cap always dry, smooth, with a thin floccose, parted, involute edge; flesh soft j gills rounded, clustered, obliterated in front; stalk united with cap.
26904 <i>decastes Fries</i>	tenth stipes naked 3 au.s Cin beech woods
26905 <i>gambosus Clus.</i>	ox-hoof smells of flour 2\$ my.jn Pale chalk, mead,
26906 <i>personatus Fries</i>	mask variable 3 s.n P woods Sow. 209. i
26907 <i>nftidus Bui.</i>	naked wavy 2 s.n V gardens Bui. her. 439
26908 <i>álbus Schaff.</i>	white not spotted 3 s.n W woods Schaeff. 256
26909 <i>spéndens With.</i>	shining juicy 3 jl Cop alder stump

II. NUDICAU'LBS. — Stem central, naked; gills unchangeable; sporidia white.

16. RU'SSULA Pers. — Pueusfleshy, becoming depressed; lamella equal, juiceless.

	i. — <i>Gills all equal; sporidia yellow.</i>
26910 <i>alutaceus Pers.</i>	tanned leather eatable 2 au.s R shady woods Bui. her. 509
2 xanthopus Fries	yellow-stalked eatable 2 au.s R shady woods
auratus With.	
26911 <i>Ifateus Hud.</i>	yellow brittle 1 au.B Y beech woods Schaeff. 2D4
26912 <i>nitidus Pers.</i>	shining nauseous 1 au.o Rsh woods Sow. 201. xx**
26913 <i>emelicus Schaff.</i>	emetic acrid 3 su Rsh woods Sow. 201. xx**
2 Gebrig. L.	St. George's acrid S su Y woods liul. her. 50%

26914 depfUena Pers.

pallid ii. — Gills nearly equal; sporidia white.
nauseous I*jl.s K.Br heaths

	red	iii. — Gills forked, and many of them halved.	
	fetid	very bitter	2 } ls R woods Bul.her. 42. <i>sanguineus</i>
	forked	rigid	2 au.s Y woods Bui. her. 292. <i>iperatus</i>
26918 adfistus J.v.r., » elephanfinus Bol.	scorched	bitterish	2 au.s G woods Bui. her. 26. <i>bifidus</i>
	elephantine	very compact	2 jl.o Ol woods Bui. her. 212. <i>nigricans</i>

7. GAXORHJSHJS Fries. —Pileus becoming depressed; lamella unequal, milky.

2691Q rnn». ~ I	Edge of cap rotted, inwards downy; hot, poisonous.	„ „ „ ^{mM}
g g w controversus Per*, culiculus Sco.	controverted meteoric	2 s.o Va beech woods Bui. her. 538. <i>dcris</i>
	furrowed gigantic	4 au.o Y damp woods Schasff 227
	gripping dangerous	2 jn.o Pk way sides Sow. 103
«3 ciUcioides Frfe	destructive poisonous	2 au.o 01. Br woods Bui. her. 14

downy very downy 3 s Di.Pk pine woods
ii. — Cap smooth, viscid, with a naked edge; hot, poisonous.

	lurid	flattened	1 f s o Lu Heaths Sow 203. <i>zondrius</i>
	acid	very acrid	2 au.n Cin groves Bolt pil. 60
	moist	brittle	1} au.o Li.Pk damp groves
	withered	very acrid	2 au.n Ld woods
	Hysginum	variable	2 au.o Pk grassy places
	dull	very acrid	If jl.o Gsh beech woods
	pallid	gregarious	J au.o Pa.Y beech woods
	delicious	eatable	Ii jl.n O pine woods Sow. 202
	orange	acrid	3 au.o O woods Batarra, 16. A.

Omp ^{rubr.} Cap dry, naked at edge; gills close, when young white, afterwards yellow; eatable, but acrid.

£5°' mitissimus Fries	mildest	sweet	3 au.n O woods
W Fries	still	sweet	3 aut Pk oak woods Fl. dan. 1069. <i>Ribiscens</i>
	sweetish	nauseous	3 GU Brsh woods Sow. 204. <i>lacifflavus</i>
	sulphur-juiced	poisonous	2i s.o Ful shady woods Bui. her. 567. A.
Tithymallus-lk	very milky	3 s.o Pa. Y shady woods Batsch fun. 60. <i>ichorht.</i>	
rufous	scentless	2 jl.o Br pine woods	

- rub^scens With.

ggfc hiflvs Fries

SgW glycyosmus Fries

W plumbbeus Bui.

- lead-colored

With. Bui.

flexibus Pers.

piperatus Sco.

ellerius Fries

U6dulcis/W.

depressus With.

pale red acrid 2| jl.n R.Och damp places

sweet-smelling esculent 3 jl.o Lu thickets

insipid 4 au.s Lead damp places Sow. 245. *Listen*

very acrid If au.b 1. . groves Bui. her. 529.1

flexuous compact 1 jl.o Br grassy places Bui. her. 559.1. A. *axon.*

peppery eatable 2 s.o W woods Bolt. pil. 21

fleecy gregarious 2 au.n W thickets Sow. 104. *Listen*

depressed variable 2\$ au.o Pk grassy places

CLIT6CTBE Fries. —Pileus fleshy, when young convex; lamella unequal, juiceless.

A. AcuVi. — Gills equally narrowed backward, acute.

i. — Cap dry, smooth; gills close, decurrent, or acutely adnate.

Kiganieus Leys. gigantc very broad 6 s Wsh thickets Sow. 244

Kiganieus Pers. anti-colored gigantic 3 au.n Di.Y among moss Grev. cryp. 1.41

	flaccid	pretty	3 s.o W woods Bolt. pil. 185
	gibbous	fragrant	2 o Br plains Bui. her. 573. 1. 2
	turfy	scentless	1 n Br turf Sow. 210
	perfected	tough	2 s.n Pk woods

K'IT' JP rto*dy fleshy, convex, opening out flat; gills truly decurrent: stalk strong; eatable.

Iff.*** Batsch cloured gregarious 3 s.o Cin heaths Boit. pil. 40.-md/to

26955 U8 With.

**** ergidus Grev.

& mm Cap truly but not firmly fleshy, flattish or slightly depressed; gills adnate, not properly

decurrent; stalk slender.

2695 f1 c4b Wuh	green	slender	2 au.n Cin woods Bolt. pil. 12. <i>sterneleus</i>
	scented	seabore	3 au.n W dead leaves Bolt. pil. 17
	hoary	shaggy	3 au.n Wsh meadows Sow. 123

	lincar.stalked	shaggy	3 o W grassy grov. Sow. 281. <i>graveolens</i>
	dog's collar	depressed	3 ja Brsh woods Sow. 184
	unadorned	pretty	2 aut Ld.G upon earth Sow. 342
	fringed	gregarious	1 au.s Wsh rotten wood Bolt. pil. 61

lu. — Tufted, variable, some growing on wood, some on earth.

wood irregular 2 au.d Wsh rotten wood

adhesive irregular 3 s W.Br plantations

5cA «if:sweHed-sUlked tufted 2 sp.aut Ru woods SchaffT. 259.

ralloid <<8 Dick coral-like tufted 2 spaut Ru hollow trees Battarra, 9. F.

18 with squat tufted 2 ap.s W.Br pastures

p^aJue crackin S au W among grass Sow. 142

ii. — Cap thin, somewhat compressed dry; gills very distant, arcuated, decurrent.

us Fries arched-leaved brittle 4 au.o Sooty damp woods Sow. 172. *elexus*

ers. meadow eatable 1| au.n Ysh waysides Grev. cryp. 2.91

formis With club-shaped eatable 11 au.n W way sides Schaff 307

With. heath eatable 1| s.n W heaths Bui. her. 467. *erecturus*

Wul. virgin-tuAfc. eatable 2 s.n W heaths Grev. cryp. S. 166

M »» Cap thin, viscid, wet; gills variable; stalk hollow, terrestrial.

2ji wittactaus&faed? parrot-colored pretty 2 an Gsh.Y meadows Grev. cryp. 2 74

8675 us5ow. wax-colored gregarious 2 au.n Y pastures Sow. 20

a conicub Schasff: conical watery 4 my.o Ysh meadows Sow.S81. *aurantiacus*

86976 punfucus Fries	crimson	beautiful	3 au.o	O.R	among grass Bui. hcr. 202. <i>coccineus</i>
26977 coccineus Pers.	scarlet	beautiful	2 au.o	S	meadows Sow. 381
B. INEQUAV.ES.— <i>GUIS unequal at back, that is, toothed; or arcuated, decurrent, sinuate, emarginate.</i>					
iv.— <i>Cap dry, minutely scaly; gills generally arcuated, decurrent, rarely adnate; firm.</i>					
26978 laccatus Sco.	lake-colored	Jiandsome	2 jn.n	Ro.R	on earth Grev. cryp. 249
2 amethystinusftai.	amethystine	handsome	2 jnn	V	shady places Sow. 187
26979 ovinus Bui.	sheep	mild	2 au.o	Brsh	meadows Bui. her. 580
26980 sulphurous BuL	sulphureous	fetid	4 s.o	Test	trees&woods Sow. 44
SKW81 turtulis Bol.	twisted	distorted	\ aut	Br	gard. mould Bolt pil. 41
26982 ovalis With.	oval	satiny	2\$ s.o	It.	Br fir woods
v.— <i>Cap smooth, somewhat humid; gills arcuate at their connection with the edge, reticulated at their union with each other, with appendages at edge.</i>					
26983 pelianthinus Fries	dark violet	beautiful	3 aus	P	roots of trees Bolt pil. 4.1. dentieul.
vi.— <i>Cap thin, dry; gills emarginate, brittle,</i>					
i.— <i>Cap fleshy, smooth, and stem regular, terrestrial.</i>					
26984 macleaticus Pers.	black and white	elegant	S au.n	Sooty	damp places
it.— <i>Cap somewhat fleshy; stem rootless, irregular, terrestrial.</i>					
26985 compr̄ssus With.	compressed	pellucid	3 jn	Br	among grass Sow. 66
26986 murinaceum Bui.	sweet-scented	fragile	2 au.o	Cin	pastures Sow. 106
wi.— <i>Cap somewhat fleshy; stem rootless, regular, growing on wood.</i>					
26987 platiphyllus Pers.	broad-leaved	large	4 jl.o	Wsh	trun. of trees BuL her. 594. <i>gramma</i> .
G OBTU ^{VS} l.— <i>Gills equal, behind blunt.</i>					
vii.— <i>Cap fleshy, glutinous; gills somewhat united; stem rooted.</i>					
26988 radicMus Rel.	rooted	gigantic	£ jns	W	trun. of trees Grev. crypt. 4. 217
26989 gr̄acilis With.	slender	more slender	li jn.s	W	trun. of trees
26990 vcKitipes Fl. Ion.	velvet-stalked	caespitose	2 o.mr	Ful	trees Sow. 384. 3
viii.— <i>Cap tough, dry; gills separate, close, white.</i>					
26991 fuscipes Bui.	spindle-stalked	edible	2 jl.n	W	woods Sow. 129. <i>crdssipes</i>
26992 cunfluens Pers.	confluent	caespitose	4 au.o	Wsh	shady woods
26993 collinus Sco	hill	caespitose	4 s.o	Pallid	among grass Bui. her. 403. <i>arundi</i> *.
26994 ttryophilus Sow.	Oak-loving	oespítose	2 o.n	Wsh	heaps of Ivs Sow. 127
ix.— <i>Cap somewhat leathery, dry; gills separate, distant, pallid.</i>					
j.— <i>Cap fleshy; stem solid.</i>					
26995 peponatus Bol.	clasped	changeable	2£ jl.n	Test	dead leaves Sow. 37
26996 oreades Bol.	Oreades	eatable	3 my.n	Pa.Ru	grassy places Grev. cryp 323
it.— <i>Cap fleshy; stem fistulose.</i>					
26997 porreus Fries	Leek-scented	fetid	3 o.n	W	plantations Sow. 81. <i>alliacetus</i>
26998 fusco-purpiircusPers.	dark-purple	caespitose	2} jl.s	D.P	beech leaves Pers. ic. pict 4.1
\$ 9. COLLV'BIA Fries.— <i>PUeus fleshy, membranous, flatfish, small, dry.</i>					
i.— <i>Cap slightly fleshy, smooth, scarcely umbilicate; gills true; stem hollow, or somewhat fistulose.</i>					
26999 icorodonioides Fines	Garlic-scented	strong smell	1J au.o	Wsh	heaths Schaeff 99. <i>ajUitius</i>
27000 carnēus Bui.	flesh-colored	dwarf	1 aus	R	grassy places Bui. her. 533.1
punicus With.					
27001 cscullintus Wul.	eatable	esculent	1J ap.my	Clay	way sides Scharff. 59. <i>clodus</i>
27002 conigenus Pers.	cone-sprung	filiform	2 aut	Gy	on fir cones and dead leaves
47003 tenacillus Pers.	small tough	long	4 aut	Pa.br	fir woods Pers. ic. pict 1.2
27004 tuberibus Bui.	tuberous	gregarious	ft au n	W	on fungi Grev. cryp. 1.23
alfinnus Bol.					
27005 racemosuB Sow.	racemose	-	compound	J aut	Gy on fungi Sow. 287
27006 cftvus Bui.	club'	gregarious	1 au.o	O.R	dead branch. Bolt. pil. 39. B.
27007 rameal is Bui.	branch	Jiving	£ year	Wsh	dry branches Bolt pil. 39. D. <i>cdndidut</i>
27008 parasiticus Bui.	parasitical	meteoric	£ au.o	Gy	on fungi Sow. 343
ii.— <i>Cap thht, membranous, flat, becoming depressed, plaited, rugose; gills veiny, of the same substance as the cap; stem horny, black.</i>					
27009 Vaillantii Fries	Vaillant'a	tough	1 a-<	W	dead branch. Vail. pa. 11. SI-S4
Mcriiliu<imdroBveus With.					
27010 r̄tula Sco.	little wheel	gregarious	1 year	W	woods Sow. 95
27011 androstaeus L.	Androsace-lk	tough	1jyear	W.Br	dead leaves Bolt pil. 32
27012 fce'tidus Fries	fetid	gregarious	1 au.s	Br	fallen bran. Sow. 21. <i>Merhlius/aft-</i>
27013 perforans Fries	boring	fetid	1 year	Pallid	dead fir Ivs Sow. 94. <i>androsaceus</i>
27014 epiph^llus Pers.	leaf-borne	grrgarious	f.s.d	W	dead leaves Sow. S3. <i>squidmula</i>
27015 Hudsoni Pen.	Hudson's	hairy	f aut	Wsh	holly loaves Sow. 164. <i>pillbus</i>
\ 10. MYCR ^X NA Pers.— <i>Pileus membranous, campanulate, slender; stipes hollow.</i>					
L— <i>Dry cap, generally umbonate, not depressed; gills separate or adnate, not decurrent.</i>					
i.— <i>Stem rooting, smooth, Juicelrss; gills separate, whole-colored.</i>					
2701f alliaceus Jac.	GmWc^cented	fetid	6 jl.n	W.Br	dead leaves Jac.au. 82
27017 atro-albus Bol.	black and white	scentless	3 au.n	Bsh	damp places Bolt. pil. 137
27018 f Hopes Bui.	thread-stemmed	campanulate	3 aut		moist woods Bui. her. 320
ii.— <i>Stem smooth, juiceless, somewhat rooting; gills adnate, whole-colored.</i>					
27019 alcannus Fries	alkaline	fetid	2 my.o	Cin	on earth Vail. pa. 12.1,2
27020 galericulatus Sco.	small-capped	scentless	3 my.o	Brsh	trun. of trees Sow. 165
vitrius With.	proffcrus Sow.				
Hi.— <i>Stem juiceless, striated; gills whole-colored.</i>					
27021 polygrammus Dec.	much marked	cespítose	6 s.d	Cin	dead leaves BuLher.518.J7/!*A ⁵¹¹
iv.— <i>Stem smooth, milky, somewhat rooted; gills attenuated, united at edge.</i>					
27022 galopus Pers.	milky-stemmed	scentless	4 au.n	D.G1	woods Fl. dan 1550. 2
27023 heerhātopus Pers.	bloody-stemmed	caespitose	2 au.o	Pk	beech trunks
87024 cruentus Fries	bloody	solitary	3 au.n	R	fir leaves
v.— <i>Stem smooth, juiceless, somewhat rooting; gills adnate, discolored at edge.</i>					
27025 illegans Pers.	elegant	fennel-scented	2 au.n	Ld.Y	pine woods
27026 strobillinus Pers.	strobose-like	gregarious	2 au.n	C	pine woods Sow. 197. <i>coccineus</i>
27027 roseub Pers.	rosy	gregarious	2 au.n	Ro	pine woods Pers. syn. 3.5
vi.— <i>Stem smooth, juiceless, scarcely rooting; gills affixed, whole-colored; color pure.</i>					
27028 piius Pers.	pure	gregarious	3 jn.n	Ro	woods Sow. 72. <i>roseus</i>

S 5 S Adunli Bui.	Adonis various-dd	SJs.n	Rosy woods	Bui. her. 560. 2
S ? S J ^{te} bus Bol.	yellowish, white pretty	2 au.s	Y among moss	Bolt pil. 38. 1
Sf S lac te ^{us} Per.	milky gregarious	14 JIn	W.Y heaths	Sow. 385.5. Nuis
<7032 stylobates Fries	pedestal solitary	Ijsu	Pa dead lva& bran.	Neespilzel 89
9<*> * vi. — 5te/n juiceless, rootless, but swollen at the base into a globe ; cap blunt.				
ZU35 pihpes Sow.	hairy-footed caespitose	2 aut	Pa.Br dead agarics	Sow. 24
	ii. — Cap or stem viscid ; gills adnate or decurrent.			
Fern . borne variable		1\$ aiui	Cin among moss	Sow. 92. nhtans
common gregarious	1J au.n	Cin fir leaves	FL dan. 1678 2	
	iii. — Dry cap, finally depressed; gills decurrent.			
	i. — Firm, persistent, with a firm stem.			
27036 pellucidus Bui.	transparent thick	1) aut	Ru the ground	Bui. her. abO. 2
	ii. — Delicate, withering, with a capillary stem.			
70JB COTTERMIS Bul.	bark delicate	} o.f	Ru bark of trees	Sow. 243
k rosellus With.	Fern-sprung beautiful	} 1) au.o	Rosy among moss	
S ^o 39 "Pnipes &nu.	spiny-footed gregarious	■ au.o	Br pine cones	Sow. SOS
*7040/uncolca Fries	Rush-dwelling gregarious	■ o	Rsh rotten Jfnci	Mic.gen.80.9

L I OMPIA^XLIA Pers. — Pileus membranous or fleshy-membranous, when young umbilicated.

i. — Cap somewhat membranous; gills decurrent.

	i. — Small cap membranous.			
starred	gregarious	1 jl.au	W hollow trees	Sow. 107. buecmalis
button	slender	1) my.o	O.Y among moss	Sow. 45
box-like	variable	2 my.n	Test on earth	Bui. her. 568.2
wall	suhgregar.	f aut	Br among grass	Sow. 322
heaths	variable	1 my.n	W damp heaths	Bui.her. 2
thick-stalked	solitary	8 jl.o	Fer pine woods	Sow. 163
	ii. — Large cap, somewhat membranous.			
tunnel	tender	1 jlo	Cin will, trunks	Pen. ic. pict. 13. 1
oblique	solitary	1 aut	Pa.Ci on earth	Pen. ic. met. 13.3
	ii. — Cap fleshy, fleshy-membranous, gills admite.			
fragrant	assesscented	1)aud	Ld damm grass	Sow." 10
csspitose	.pellucid	1 au d	Y peat	Bolt. pil. 41. C
cup-shaped	club-shaped	3 o.n	D.Br earth	Sow. 363. sordidus
	iii. — Cap fleshy, coriaceous, somewhat corky, soft; gills decurrent.			
70a0 caapitteus Bol.	gregarious	2 s	G earth	Sow. 162
70S1 cyathifArmis Bui	gregarious	..	W will, trunks	
clavatus With.				
SP52 murlinus Sow.	sweet-scented	2		
27053 DunWt Dec.	Dunal's	gregarious	spring	W will, trunks
7054 Ignus Bul.	gregarious	1) my.jn	Wsh trun. of trees	Sow. 64
7055 lepidus Fries	scaly	variable	1 my.jn pa.och pine trunks	SchsfT. 2a squambsu\$
S monstrus Fries	monstrous	deformed	3 my.jn pa och pine trunks	Sow. 3S2. tu&tfrmis
*Q56 cocheatus Pers.	spiral	csspitose	3 sp.aut Wbh old trunks	Sow. 1u8. confluens

III. ACAU'LES. — Stem out of the centre none; gills unchangeable; sporidia white.

4 12. PLEUEOTU9 Fries. — Pileus out of the centre or lateral.

	i. — Veil universal; cap compact, horizontal.			
*fo57 rfrlynus Pers.	Oak	solitary	1 au.n	Wsh oak trees Scheff. 233. dimididtus
	ii. — Veil none; cap fleshy; gills decurrent.			
	i. — Cap always entire.			
twisted	very tough	1) jl.o	Rsh birch trees	Mees pilze 176
	ii. — Cap entire or halved.			
shell-shaped	gregarious	1 jLs	Cin birch trees	Bui. her. 298
oyster-shaped	eatable	sp.aut	Cin trun. of trees	Sow. 241
	iii. — Cap always halved, somewhat ascending.			
petal-like	gregarious	i <.o	Brah beechn trees	Bul. her. 226.557.2
stretching	imbricated	1) jl.n	W pine tree*	
fan-shaped	thin	7 Jl.n	I'o. Dr side of tree* Bolt pil. 157	
	eapflethy, when young horizontal; gills terminating in a determinate manner.			
Km	capitofie	3 ox o.d alc	trun. of trees	Sow. 67
palmate	caespitose	2 o.d	Br oak trees	Sow. 62
late	dwarf	1 o.ja	Ol trun. of trees Bux. c. 5.2.2	
Hiptic	gregarious	% o.ap	Tan	
	IV. — Cap fleshy, when young resupinate; gills running together in a point out of the centre.			
nidulans Pers.	nestling	imbricated	1 o.d	Ysh fallen trees
	furred	imbricated	1 s.n	Gy beech trunks Sow. 99. echindtus
	soft	gregarious	£ s.n.	Y.Br beech trunks Sow. 98
	variable	gregarious	s.n	W beech trunks Grev. cryp. 2S5
	lateral	imbricated	1) su	Umb birch trunks Fl. dan. 1556.2
	V. — Cap membranous; gills adnate or running together in one point.			
tremulous	almost sessile	\ aud	Gy earth	Sow. 242
putrefactive	delicate	\ au.o	W decay, trun.	Sow. 821. pubescent
applied	delicate	\ autsp	Cin decuy. trun.	Sow. 301

IV. ROSEI. — Stem central; veil none; gills changing color; sporidia rose-colored.

13. MOUCERO.V Bauh. — Pileus fleshy, becomes depressed; lamella long, decurrent; odor of new flour, prunulus Cats. little prune esculent

	lj.jn.o	W woods	Sovr. Ii3. pdllidus
ft 14. CuTOprLUS Fries. — Pileus fleshy convex.			
i. — Gills affixed, terrestrial.			
garden	clastic	2 ^h aut	Sooty gard. on ear.
rosy white	beautiful	3 jl.n	Ld damp places
proflic	gregarious	3 aut	P.Ld hedge rows
sinuated	fragrant	5 o	W.Y damp woods
seashore	small	1 o	W damp woods
ii. — Gills altogether distinct, upon wood.			
Won-colored	fragile	3 au.o	Y beech woods Schaeff 43
penthouse	variable	3 my.n	Sooty trim. of trees Sow. 106

27084 *phlebóphorus* *Bit.* vein-bearing gregarious 4 jl.o 01 decay, wood Grev. cryp. 3.173
reticulatus *With.*

§ 15. *LEPTONIA* Fries.—*Pileus fleshy, membranous, from convex becoming plane, small.*

27085 *griseocyancus* Fries blue-grey solitary 1| au.s Li grassy hills Bolt pil. 41. *purpureus*
 27086 *chalybeus* *Pen.* steel pretty 2 jLi B among grass Sow. 161. *columbarmus*

J 16. *NOLAVEA* Fries.—*Pileus membranous, campanulate; slender stipes hollow.*

27067 *majális* Fries hog csBspitose 4 spring Cin fir woods Sow. 174. *mollifaculus*
 27088 *páscuus* Pers. pasture variable 3 year Sooty everywhere Bolt. pil. 35. *Jissus*

% 17. *ECCI'LIA* Fries.—*Pileus umbilicate; lamella? adnate.*

27089 *asprillus* Fries roughish gregarious 1/ su Gy grassy places
 27090 *aquila* Fries swarthy subsoitary 1 au.s Umb riversides
 27091 *politus* Fries polished gregarious 3 au.o Ld among grass
 97092 *carneo-albus* *With.*, light-flesh gregarious 1 au.o W among grass

V. OCUR/SCR. —*Stem central; veil like cobweb; gills changing color, becoming dry; sporidia ochre-colored.*

& 18. *TBLAM&NIA* Fries.—*Veil annular, woven, somewhat persistent; lamella; distant.*

27093 *tórvus* Fries grim strong-scld 4 jl.o Br damp woods Bui. her. 6Q0. araneb \$U3
 27094 *brúnneus* Pers. *wown weak-scented 4 jLn Pa.Umb pine woods
spongibus *With.*

27095 *ev*nius* Fries well-branched solitary 5 jl.n P. Br pine woods Sow. 125. *impilber*
 27096 *sublanHus* Sow. half-woolly radish-scented 3 au.o Ol Br woods Sow. 224
 27097 *bulbosus* Sow. bulbous radish-scented 4 au.o Br among grass Sow. 230

§ 19. *I tiOLuM A* Fries.—*Veil fugacious; lamella emarginate; stipes bulbous; color something of violet.*

i. —*Cap always dry, scaly, or fibrous, obtuse, or mlnbonate, never depressed.*

27098 *violaceus* /,, violet showy 4 au.o V groves Bolt. pil. 52 *[cephalu]**
 27099 *pholidius* F//a scaly solitary 4 au.o Cin woods B\heT.586.l.psamtno-
 27100 *spilomodus* Fries spotted solitary 3 au.s Pa. Br woods Sow. 384.1. *aranedsus*

ii. —*Cap smooth, humid, viscid, always obtuse, finally depressed; stem blue, becoming white.*

27101 *scafirus* Fries club-footed soft 3 ja.o Ol woods Batsch fun. 2. 184
 27102 *cálluchrous* Per*, fine-skinned insipid 4 au.o Psh woods Batsch fun.J.14.subpur.
 27103 *glaécupus* Schteff. glaucous-footed gregarious 3 au.o Ol woods & hea. Sow. 223
 27104 *varius* Schaff. variable variable 4 au.o Y everywhere Sow. 102. *turbindus*
 27105 *turbinátus* Bui. turbinate soft 4 s.n Y damp woods Bui. her. 110

\$ 2a *DERMO'CYBE* Fries.—*Veil fugacious; lamella closely packed; stipes equal.*

i. —*Cap scaly or fibrous; stem same color as cap or paler; growing on earth**

i. —*Cap fleshy, at first convex.*

27106 *sanguineus* WuL bloody handsome \jn.n C woods Sow. 43
 27107 *cinnambmeus* L. Cinnamon variable 3 jn.d Cin everywhere Sow. 205
 27108 *helvolus* Pers. pale red dirty 2 jn.d Cin woods Sow. 173. *hinnheus*

U. —*Cap somewhat fleshy, at first campanulate.*

27109 *Cbumis* Pers. Cucumber-sclf strong smell. 3 au.oc. P.Br woods Sow. 344. *tugripes*

ii. —*Cap smooth, but with a few surface fibres; stem white; growing on earth.*

27110 *armeniacus* Schceff. Apricot-cW softish 3 jl.n Pa. Br woods Scheff. 81

27111 *castáneus* Bui. ChestnuU&J gregarious 2 jl.n Ch woods Bui. her. 268

iii. —*Cap smooth, dry; gills affixed.*

27112 *hytidoides* Sow. hybrid variable 2 my.n O fir leaves Sow. 221

27113 *testoëus* *With.* testaceous crooked 4 s Y plantations

27114 *flavidus* Sow. yellowish soft 2 s Ysh among grass Sow. 366

VI. DIOTI'NCTI. —*Veil distinct, not like a cobweb; gUs discolored, somewhat persistent; sporidia ferruginous.*

*> 21. *PHOLIOTA* Fries.—*Veil dry, annular.*

27115 *afreus* Sow. golden subcspitose 6 s.o Ful damp earth Sow. 77

27116 *caperatus* Pers. fendrilled solitary 5 jl.o Lem mount wo. Fl. dan. 1675

27117 *aunvillus* Batsch golden-fleeced solitary 3 o.n Y old trees SchefT. 209. *filament'*.

27118 *squarrosus* Pers. squarrose cspitose 2 aud. Fcr roots of trees Grev. cryp. a. *Jioccbsu**

27119 *flammans* Batsch tiamcolored elegant 3 jl.o Y pine trees Batsch fun. SO

rheo)des *With.*

27120 *muricatus* Fries muricated variable 2 jLo DLY commons

scaribus *With.*

2 inaequalis Batt. unequal variable 2 jl.o Di.Y commons Bolt pil. 50. *Itheus*

27121 *mutabilis* Scheffii changeable eatable 3 my.n Pa.Ci on trees Schsff. 9

27122 *constrictus* *With.* constrained watery f s Y.Br rotten wood

\ 22. *MYXAVIUM* Fries.—*Veil viscid, fugacious; lamella affixed.*

27123 *collinUus* Sow. besmeared solitary 5 jl.n O woods Sow. 9

27124 *longicaulis* Fries long-stemmed membranous 4 o Tan pine woods

flexuosus *With.*

I 2a *HEBEIAMA* Fries.—*Veil marginal, fugacious; lamella emarginate.*

27125 *fastibilis* Pers. haughty fetid \ jl.n Wsh everywhere SchaffT. 221. *gUvus*

VII. FUOA^CBS. —*Veil very fugacious or spurious, not like a cobweb; gills discolored, somewhat persistent' > sporidia ferruginous.*

\ 94. *FLA'MMULA* Fries.—*Pileus fleshy, convex, smooth, somewhat viscid; lamella not emarginate.*

i.—*Cap dry; gills adnate, tufted; growing on wood.*

27126 *flavidus* Schaffi yellowish ca»pitose 2 au.n Ysh trun. of trees SchaffT.35

27127 *Inopus* Fries. destitute subesspitose 2 s.o Ysh trun. of trees Bolt, pil.148. rad. connatus *With.*

ii. —*Cap viscid; gills adnate, not tufted; growing upon both wood and earth.*

27128 *spumbus* Batt. frothy gregarious 3 au.n Ysh on earth, &c. Battarra, 22. C

% 25. *INO'GYBE* Fries.—*Veil formed of the longitudinal fibres of the fleshy convex pileus; lamella whiti**

i. —*Stem fibrous or scaly with fibres.*

27129 *scaber* Sow. rough solitary 1j au Sooty pine woods Sow. 207

87130 *plumosus* Sol. feathery solitary 4 aut Gy woods Bolt pil. SS

87131 *lanuginosus* Bui. woolly solitary 3 jls Brsh waysides Bui. her. 370

ii. —*Stem near the top covered with white scales.*

27132 *nmbusa* Bui. crarked variable 2 jns Y.Br woods Grev. cryp. 3. I**

S 1 S p°l^h f»» Sow. earth-leaf variable 2 jLo Wsh woods Sow. 124

27134 *furfuribus* *With.* surfy watery 1 jn Y.Br hedges

27198 lottitus Fries	lobed	tough ft	spring Brsh	humid plac.	Bolt. pil.	<i>ITI.membranH.</i>
27199 lutescens Fries	lutescent	spirit-scented	2	jl.n. Y* ^h	humid plac.	Sow. 47.A. <i>cantharelo</i>).
3122. MERITLIUS Hal.	DRY-ROT.	{ <i>Meta</i> , pillar; resemblance in original fungus <i>metulius</i> .)	1.—10.			
27200 lachrymans Schum.	weeping	parasite	4 year	Y.Br	decay, wood	Sow. 113
2 obliquus Bol.	oblique	parasite	4 year	Y.Br	decay, wood	Bolt. pil. 74

3123. SCHIZOPHYLLUM Fries.	SCHIZOPHYLLUM.	(<i>Schizo</i> , to cut; <i>phyllon</i> , a leaf; appearance.)	1.
27201 commune Fries	common	gregarious	2 wet w Gyh trun. of trees

3124. DiEDA^LEA Pers.	D^DALEA.	(<i>Dcedaleus</i> , artificial; appearance of sinuosities.)	7.—30
27202 quercina Pers.	Oak	variable	0 year Pa. Y oak trees
27203 biennis Fries	biennial	thrcinch.br.	1 year Fer rotten wood
27204 tetulina Pers.	Birch	smaller	0 year Pallid birch trees
27205 confragans Pers.	rough	woody	0 year Brsh service trees
27206 unicolor Fries	one-colored	imbricated	0 aut Sooty trun. of trees
27207 gibbosa Pers.	gibbous	six inches br.	0 aut Wsh trun. of trees
27208 angustala Fries	narrowed	two inches br.	0 aut Cin poplar trees

3125. POLYPORUS Mich.	POLYPOMJS.	(<i>Polys</i> , many, <i>poros</i> , a pore; under surface.)	36.—143.
t 1. FAVOLUS Beauv.	—Pores ample, with four or six angles resembling a honeycomb.		
27209 squamulosus Fries	scaly	3-18 inc. wide	2 jn.n Och trun oft roes

27210 heteroclitus Fries	variously bent	2J inch, wide	0 aut O on earth
--------------------------	----------------	---------------	------------------

2. MICROPORUS Beauv. —Pores minute, roundish.						
27211 leptoccephalus Jac.	small-headed	one inch wide	1 aut Gy	on wood	Jac. m. 1.12	
27212 brunneus Pers.	winter	1-4 inch wide	2 year	Sooty trun. of trees	Sdaff. fe81. <i>pileus</i>	
27213 pcrmnis Fries	perennial	thin	3 aut	Cin. trun. of trees	Sow. 192	
27214 strophiliformis Dick	cone-like	lumpish	2 aut	Br trun. of trees	Dick. cr. 3.2	
27215 pellucidus With.	pellucid	two inch. br.	f au	Br old wood		
27216 varius Fries	variable	patches	2 year	Br trun. of trees	Grev. cryp. 202	
2 lateral. Bol.	lateral	patches	2 year	Gyh trun. of trees	Bolt. pil. 83	
3 nummularius Bui.	Moneywort	patches	2 year	Wsh trun. of trees	Sow. 368. fig. min.	
27217 lhcidus Fries	shining	variable	1 to 6 su	Ysh trun. of trees	Grev. cryp. S45	
27218 frondifus Fries	leafy	broad patches	2 s.o	Sooty roots of oaks	ScheffE 127	
27219 velutinus Fries	velvety	thin	2 sp.aut	Wsh trun. of trees		
27220 gigant&us Fries	gigantic	tufts	24 su	Pa.Br beech trees	Sow. 86. <i>imbricatus</i>	
27221 sulphureus Fries	sulphur-c/d	tufts	24 su	Rsh.Y oak trees	Grev. cryp. 113	
27222 teflinus Fries	Birch	acid	2 su	Brsh birch trees	Grev. cryp. 229	
27223 sphmeus Fries	frothy	thick	3 aut	Wsh trun. of trees	Sow. 211. <i>stipitatus</i>	
27224 cascius Fries	grey	very thin	2 aut	Bsh trun. of pines	Sow. 226. <i>t'Ulnodus</i>	
27225 hispidus Fries	hispid	spongy	6 su	Fer oak trees	Grev. cryp. 14	
Ifaletus velutinus Sow. 345						
27226 cuticularis Fries	cuticular	imbricated	3 aut	Fer trun. of trees	Sow. 195. <i>impuber</i>	
27227 adficius Fries	scorched	imbricated	2 aut	Pa.Br trun. of trees	Sow. 231. <i>Qurpinus</i>	
27228 tilmarius Fries	Elm	3-4 inch, wide	0 aut	Pallid elm trees	Sow. 88	
27229 suaveolens Fries	sweet-smelling	fragrant	3 aut	W willow trun.	Sow. 228	
2 <calicinus Fries	Willow	fragrant	3 aut	V willow trun.	Sow. 227	
27230 versicolor Fries	changeable	tufted	1 su.aut	Bsh trun. of trees	Sow. 22!	
27231 radi&tus Fries	radiated	imbricated	1 aut	Y.Br trun. of trees	Sow. 190	
27232 pallescens Fries	pallcescent	imbricated	2 aut	pa.orh trun. of trees	Sow. 250. <i>pcleporus</i>	
27233 abilitinus Fries	Pine tree	imbricated	1) aut	Wsh dead pines	Grev. cryp. 22i	
27234 fomentarius Fries	soft	spongy	6 year	Sooty beech trees	Sow. 133	
27235 igniarius Fries	fiery	hard	6 year	Fer trun. of trees	Sow. 132	
27236 Kibis Fries	currant	appressed	ft aut	Fer red curr. ro.		
27237 spongeosus Fries	spongy	tufts	2 aut	Fer trun. of trees	Boltpill.6i>.r4Y!fbf/id/tf<	
27238 medinae p&n Fries	bread-crumb	thick	4 aut	W fallen timber	Bolt. pil. 156. 1	
27239 vulgaris Fries	common	fragile	12 year	W fallen timber	Bolt. pil. 163. <i>Prteus</i>	
27240 ferruginosus Fries	rusty	unequal	1 6U	Fer fallen bran.	Grev. cryp. 155	
27241 muUicus Fries	molluscous	variable	3 year	W dead trees	Sow. 326. <i>tnedHl. pants</i>	
27242 incamatus Fries	flesh-colored	firm	3 su	Pk pine woods		

3. POLYSTICTA. —Dots superficial only.						
27243 reticulatus Nees	netted	su	W	pine wood	Nees pilze 225	
27244 CarmichslidW Grev	Carmichael's crust-like	very delicate	aut	W	decay, trun.	Grev. cryp. 224

3126. BOLETUS Dil.						
27245 luteus L.	BOLETUS.	(<i>Bolos</i> , a mass; globular form.)			9.—20.	
21246 lactifluus With.	yellow	3 inch, broad	2 aut	Y woods	Grev. crypt. 183	
27247 piperatus Bui.	milk-flowing	2-4 inches br.	2J aut	Buff' pastures		
27248 vancagatus Sum.	peppery	1/4 inch, broad	U su.aut	Ysh woods	Sow. 34	
27249 subtonmentosus L.	variegated	4 inches br.	3 aut	D1.Y fir woods		
2 & sanguineus B. With.	rather downy	cracked	2 jn.o	Ol woods	Bui. her. 393	
27250 Inridus SchaffT.	bloody	cracked	2 jno	C woods	Sow. 225. <i>commutatus</i>	
rubeolus Sow.	lurid	6 inch, broad	2 su.aut	OL.G groves	Grev. cryp. 121	
27251 edblis Bui	esculent	cracked	4 su.nut	Sooty woods	Sow. 111	
27252 scaber Fries	rough	3 inch, broad	4 su.aut	W woods	Bolt. pil. 86. <i>prdcervs</i>	
2 aurantiacus Sow.	Orange-cW	3 inch, broad	4 su.aut	Ru woods	Sow. 110	
3 bovinus Schaff.	ox-like	3 inch, broad	4 su.aut	Sooty woods	Sow. 175. <i>sciiber</i>	
27253 cyanescens Fries	bluish	frosted	3 su.aut	Str woods	Bui. her. 369	

3127. FISTUL^NA Bui.	FISTUUNA.	{ <i>Fistula</i> , a pipe; tubular.)	1.
27254 hepatica Bui.	liver	patches	6 aut C oak trees

S128. JIY'DNUM L. HYDNUM. (*Hydnon*, truffle. A Greek name adopted by Linneus.) 15.—87.

I. PERPENDICULARIA.—Stem perpendicular; cap distinct, round, nearly entire; growing on the ground.

27255 imbricatum L.	imbricated	esculent	1 s.o	Umb pine woods	Grev. cryp. 71
27256 repandum L.	repand	esculent	1J su.aut	Ysh woods	Grev. cryp. 4*
2 squamosum Fries	scaly	esculent	1} su.aut	Y ^h woods	Bolt. Pil. ^ ^ T^L
27257 rufataceum Pers.	rufescant	eatable	3 su.aut	Pksb beech woods	Bolt. pil. 68. <i>repanaiu</i>
27258 auriscalpium L.	ear-pick	curious	2} year	B pine cones	Grev. cryp. 190

II. HORIZONTALIA.—Stem simple, somewhat horizontal; cap halved or out of the centre, fleshy; growing on wood.

27259 gelatinosum Sco.	gelatinous	curious	i aut	Fus pine woods	Jar. au. 239
£7860 erinaceum Bui.	hedgehog	variable	3 o	W beech trees	Bui. her. 34

III. OBLITBRA'TUM. — Cap confounded with the stem, obliterated, fleshy, growing on wood.

27261 coraloides Seo.	coral-like tufts	1 aut	W trun. of trees	Sow. 252
IV. LATERA\IA. — Cap sessile, lateral, growing on wood.				
27262 erf spurn ScAatf:	crisp	4 inches wide	0 o	Brsh dead wood Schsffi 147. 1
✓2863 ochraceum Pers.	ochre-colored	variable	2 year	Ysh pine wood Sow. 15. Daviesil

x/264 minimum Bol.	least	curious	ft aut	Taw rotten oak Bolt. pil. 171
--------------------	-------	---------	--------	-------------------------------

V. RESUPINAVA. Cap resupinate, effuse; growing on wood.

✓27265 membranaceum Bui.	membranous	effused	0 su	Fer pine wood Sow. 327
27266 b&rba Jbv Bui.	Jove's beard	12 inch, wide	0 su	Wsh hollow trees Sow. 328
✓2767 spatulatum Fries	spatulate	very delicate	0 year	W pine trees Nccs pilze 231
✓228 r&dula Eries	rasp	irregular	ft aut	Ysh. W dead bir. trun. Grev. rryp. 278
✓27269 ferruginosum Pers.	rusty	bryssoid	ft aut	R>Br rotten trun. Nees pilze £48
3129. SISTOTREMMA Fries	SISTOTREMA. (Sisto, to place, Lot, trema, orifice; rows of pores.)	-		
*27270 c&nfluens Pers.	confluent gregarious	1 au.n	Wsh waysides	Grev. cryp. 248

PHLE^BIA Fries	PIILEBIA.	(Phleps, a vein; veiny branches.)	2.-4.
dro^v*fer Fries	wandering membranous	2 s.n	Su alder trees
*772 merismoldes Fries	Merisma-like confluent	1 aut	O mossy wood Grev. cryp. 280

3131. THELE'PHORA Ehrh. THELEPH. (Thele, nipple, phoreo, to bear; papillose surface of pileus.) 30.—75.

I. I'NTEGIE — Cap entire, with a stem, terrestrial.

✓TKJ, P^noBa Fries	ragged	gregarious	2 aut	Pallid on ground Sow. 155
*774 caryophylle'a Fries	clove	tough	lf aut	Psh among grass Schscheff. 325
*775 ftegans Mey.	elegant	erect	2" aut	P.Br on ground

II. LATERAYIJE. — Cap lateral, somewhat stipulate, terrestrial.

27276 terrestris Fries	terrestrial	gregarious	lft aut	Brsh on ground Bui. her. 268. caryophyl.
27277 laciniata Fries	ragged	gregarious	lft aut	Fer on ground Grev. cryp. 264

III. SE'SSILES. — Cap sessile, lateral; growing on wood.

27278 Padi Pers.	Bird Cherry	broad	4 aut	" P.Br" trun. of trees Grev. cryp. 234
27279 rubiginosa Schr.	rusty	woody	2 year	Bt.Br old oaks Sow. 26. Au.ferrugin.
27280 tahacina Fries	Tobacco	membranous	6 suaut	Fer bran. of trees Sow. 25
Auriculkria Nicotid/ia Bui. 174	elegant	thin	6 suaut	Fer bran. of trees Sow. 25
✓281 hepatica Fries	liver	imbricated	4 aut	Di.Br trun. of trees Sow. 38S. 2. Aur. lafuu
27282 hirsute W.	hairy	firm	3 year	Ysh trun. of trees Grev. cryp. 256
✓283 ochroleuca Fries	pale-yellow	membranous	3 aut	Wsh trun. of trees Sow. 349. An.yapyrinus
✓284 purpurea Schum.	purple	wavy	3 aut	P trim. of trees Sow. 383.1. A.persists
27285 Intybacea Pers.	Endive-like	very irregular	6 aut	Rsh stumps of tr
✓286 sruuans Pers.	sinuous	gregarious	1 au.w	Y.Br oak branch.
27287 coi-rllea Pers.	Hazel tree	imbricated	6 year	Och decay, hazel
27288 cbrium Pers.	leathery	thin	6 aut	Buff dead trunks Grev. cryp. 147
27289 ochracea Fries	yellow-ochre	very broad	12 aut	Och rotten trun.
27290 radiato-rimba Grec	ray-cracked	confluent	4 aut	R.Br damp fir tim.
27291 sanguinolenta Fries	bloody	very gregar.	2 n	Psh dead trees Grev. cryp. 225

IV. OBLITERAVA. — Cap obliterated, resupinate; growing on wood.

27292 fuercina Pers.	Oak tree	brittle	6 sp.aut	Bsh fallen oaks Grev. cryp. 142
✓293/raxinea Pers.	Ash tree	thin	1 aut	Gysh dead ashes
27294 niias Pers.	Lime tree	irregular	2 aut	Cin woods
✓295 epidemea Pers.	cuticular	irregular	3 n	Pa.BufTdead trunks
✓296 incrustans Pers.	incrusting	spreading	3 aut	Ysh earth,tr.,&c.
✓297 calcea Pers.	chalky	cracked	6 aut	W decay, wood
✓298 Samb&ci Pers.	Elder tree	membranous	4 aut	W decay, elders Grev. cryp. 242

V. PHYLACTE^RIA Pers.	Sporidia four in a row, resupinate; growing on the earth
mis Fries	biennial plaited

VI. HIMA'NTIA Pers.	Household smooth	6 wet w Br	da.pla. in ho.
domestica Pers.	brown rugose	2 aut	V.Br trun. of trees

i*a lactea Fries	milk-white very thin	9 aut	W trun. of trees Grev. cryp. 828
Himantia Candida Pers.			

LEIOSTROMA Fries.	Resupinate, somewhat contiguous, smooth, or with spurious papUUT; ascii none.
cmerea Fries	cinereous uneven

MERISMA Ptfr*. MERIBMA.	(Merismos, division: branched habit.)
ice'tida Pers.	fetid tufted

2 anthocephala Fries.	flower-headed stinking	2 aut	PBr woods Grev. cryp. 46
rhist*ta Greo.	crested crust-like	2 aut	Pallid damp places Sow. 151.

:uberbsa Greo.	tuberous solitary	1 aut	Pa damp groun. Grev. cryp. 178.
----------------	-------------------	-------	---------------------------------

Division II. Clavati.

313- CLAVA^RIA VaU. CLAVARIA. (Clava, a club; form.) 24.-66.

I. CRAS8ICAU'LE9. — Much branched; stems thick.

✓293/cinea Pers.	bunched much branch.	3 su	Wsh beech woods Schaeff 76. acroporph.
✓293/yunnana Pers.	yellow delicious	3 au	Ysh woods Schaeff 175
✓293/laevigata Pers.	cora like	3 aut	W on ground Sow. 278. fig. sup.
✓293/epicrateria Pers.	Tine-wood	3 aut	Och fir woods Grev. cryp. 117
✓293/epicrateria Pers.	crested polymorph.	2J aut	W woods Grev. cryp. 190
✓293/epicrateria Pers.	cinereous tufted	3 su	Cin damp places Grev. cryp. 64
✓293/epicrateria Pers.	corneous gregarious	ft aut	Y dead fir trees Sow. 40

II. TENUICAU'LES. — Branched; stems thin.

✓293/epicrateria Pers.	strict thin	1 au.o	Brsh trunks Schaeff 286. pallida
✓293/epicrateria Pers.	meadow viscid	1 aut	Y meadows Bolt pil. 114. muscoides

✓293/epicrateria Pers.	small horned solitary	3 aut	Y meadows Sow. 157. muscoides
✓293/epicrateria Pers.			

III. CLAVA^TA — Simple, clavate.

✓293/epicrateria Pers.	pestle-like largest	12 au.n	Ysh beech woods Sow. 277
✓293/epicrateria Pers.	Lady Arden's opaque	6 s	Brsh bran. of lime Sow. 215
✓293/epicrateria Pers.	fusiform regular tufts	3 s	Y among grass Sow. 204
✓293/epicrateria Pers.	thunderbolt-like much tufted	3 aut	Ysh upon trees Sow. 235. ruri*.

G ff 3 " rUg***

27322 <i>insequalis</i> Fries ■■■ <i>fragilis</i> Pers. <i>gracilis</i> Sow. 232	unequal brittle	gregarious 2 aut gregarious 1 aut	Y Ysh	meadows damp places	<i>Sov./253.yermicul&ris</i> Grev. cryp. 37
27324 <i>acuta</i> Sow.	acute	gregarious 2 aut	W	upon trees	Sow. 333
27325 <i>fimbriata</i> With.	fringed	polymor. 2 aut	W	upon trees	
27326 <i>hilvola</i> Pers.	pale red	flexuose 1J aut	Y	meadows	
27327 <i>vermicularis</i> Fries	worm-like	crowd, tufts 3 aut	W	meadows & pasture	
27328 <i>uncialis</i> Grev.	inch	very greg. i aut	W	rotten twigs	Grev. cryp. 98
27329 <i>setipes</i> Grev.	bristle-footed	gregarious taut	W	dead leaves	Grev. cryp. 49
3134. CALO'CERA Fries CALOCEBA. (<i>Ka/os</i> , beautiful, <i>keras</i> , a horn; divisions of plants.) 2. — 6.					
27330 <i>tuberosa</i> Fries	tuberous	root round. 2 aut	Y	under bark of trees	Sow. 199
27331 <i>cetrnea</i> Fries	homely	tufted ■ jd.d	■	various trees	Sow. 40
3135. GEOGLCSSUM Pers. EARTH-TONGUE. (<i>Ge, earth, glpssa</i> , tongue; form.) Clayhria Sow. 4.-9.					
27332 <i>hirsutum</i> Pers.	hairy	solitary 2 aut	B	bogs & meadows	Grev. cryp. 185
27333 <i>glabrum</i> Pers.	smooth	gregarious 1 su	Bsh	among grass	Bolt. pil. 111. 2
27334 <i>visctsum</i> Pers.	viscid	cylindric. If aut	B	moist meadows	Grev. cryp. 65
27335 <i>viride</i> Pers.	green	gregarious 2 aut	G	moist meadows	Grev. cryp. 211
3136. SPATULA*RIA Pers. SPATULARIA. (<i>Spathula</i> , a spreading knife; form of head.) 1. Clayhria Sow.					
27336 <i>flavida</i> Pers.	yellowish	variable 1J aut	Ysh	dead leaves	Grev. cryp. 165
3137. MI'TRULA Fries MITRULA. (<i>Mitra</i> , mitre; form) 1.—5. Clavhria Sow. Lebtia Yen.					
27337 <i>uliginosa</i> Pers.	marshy	hollow 1 my.au	Y	wet ditches	Grev. cryp. 312
27338 <i>minuta</i> Fries	minute	gregarious } su	Y	brae, of Dips, pilos. Sow. 391	
27339 <i>jibietis</i> Fries	Pine	dry i aut	Cin	fir woods	Sow. li.ferrugineum
<i>Lebtia Mitrula</i> Grev. crypt-%1					
3138. TY'PHULA Fries TYPHULA. (Dimin. of <i>pphaj</i> heads resembling in miniature.) 4.-11. Clavhria Sow.					
27340 <i>Phacorrhiza</i> Fries	Lentil-root	flexuose 2 aut	W	woods	Sow. 253
27341 <i>cyathiformis</i> Fries	rcd.foot	gregarious aut	W	sticks and leaves	Grev. cryp. 43.PAcor<
27342 <i>tenuis</i> Fries	slender	gregarious \ su	Bsh	on wood	Sow. 386.5
27343 <i>filiformis</i> Fries	filiform	creeping i aut	Cin	dead leaves	Grev. cryp. 93.P/wicor
3139. PISTILLA*RIA Fries PISTILLARIA. (<i>Pistillum</i> , a pestle; form.) 1.—7. Clavhria Sow.					
27344 <i>quisquiliaris</i> Fries	dead leaves	gregario. i aut	W	dead fern leaves	Sow. 3311. obtusa

Class II. UTERIVIV. ELVELLA^XCEAE. — Division I. *Mitrati*.

3140. MORCHELLA Dil. MORE** (Latinised from <i>morchel</i> , the German name of the plant) 3. — 14.					
27345 <i>esculenta</i> Pers.	esculent	'eatable 3 spring	Wsh	on the earth	Grev. cryp. 68
1 <i>rotunda</i> Pers.	round	eatable 3 spring	Wsh	on the earth	Sow. 51. fig. sinistr.
2 <i>vulgaris</i> Pers.	common	eatable 3 spring	Wsh	on the earth	Sow. 51. fig. dextr.
27346 <i>pattula</i> Pers.	spreading	eatable 3 spring	Ysh	on the earth	Sow. 51. fig. med.
27347 <i>hfbrida</i> Pers.	hybrid	cap brown 4 spring	Wsh	woods	Grev. cryp. 89
3141. //ELVE'LLA L. HELVELLA. (A name employed by Cicero, as the name of a fungus.) 5.—15.					
I. PLICA ^{ME} . — Cap waxy, membranous, at first united, afterwards wavy in plaits.					
27348 <i>leucophala</i> Pers.	grey	solitary 4 aut	Ysh	borders of fields	Grev. cryp. 143
27349 <i>mitra</i> L.	mitre***	solitary 4 aut	Ld	hedge banks	Grev. cryp. 36
27350 <i>esculenta</i> Pers.	esculent	eatable 3 mitra	Brsh	pine woods	Schaff. 160
27351 <i>infusaria</i> Scheeff.	raigent./f/c	eatable 4 aut	Cin	damp scorch.places Fl. dan. 835	
II. SEPARAVA. — Cap somewhat membranous, smooth* always separate.					
27352 <i>elastica</i> Fries	elastic	slender 4 su.aut	Bsh	damp places	Sow. 154. <i>J.uliginous</i>
3142. VE'RFA Sun.	VERPA.	(A Latin word synonymous with phallus.) 1. — 6.			
27353 <i>ctaica</i> Swx.	conical	fistular 3 aut	Br	on ground	Sow. 11. <i>JtelhAni</i>
3143. LE&TIA Hill LEOTIA.					
27354 <i>infundibuliformis</i> J.Wiffunnel-fmd	dwarf	solitary 3 o	Cin	on ground	Sow. 153
27355 <i>nana</i> With.	slippery	subsessile j aut	W	on trees	
27356 <i>lubrica</i> Pers.		gregarious 2 aut	Ol	moist woods	Grev. cryp. 5b.

Division II. *Cupuldti*.

3144. PEZPZA Dil. PEZIZA. (Pexica?, a tribe of fungi in Pliny.) 49.—300.					
I. ALEU ^{RI} RIA Fries. — Fleshy, or fleshy membranous, pruinose or surfy, with flocculent matter ; usually on earth.					

i. HELVELLOI'DEC —. Cupule always open, or when young conniving; veil superficial; sporidia with two smaller sporidia.

27357 <i>acetabulum</i> L.	saucer	clubcrd spring	Sooty damp woods	Sow. 59
27358 <i>badia</i> Pers.	brown	tufted 1 su.aut	Br	grassy places
27359 <i>leporina</i> Batsch	hare War	gregarious 1 au.o	Brsh	on ground
27360 <i>onotica</i> Pers.	asinine	gregarious 1& au.o	Brsh	dead leaves
27361 <i>aurantia</i> FL dan.	Orange	beautiful f aut	O	sandy places
27362 <i>concinna</i> Pers.	neat	very broad 1 su	Lem	dead leaves
27363 <i>cochlearia</i> Hud.	cochlearate	brittle 2\$ su.aut	Y.Br	fields
27364 <i>crebra</i> Sow.	waxen	gregarious} su	Sooty dunghills	Sow. 3
27365 <i>vesiculosus</i> Bui.	bladdery	crowded 2} aut	Wsh	dunghills
27366 <i>repanda</i> Fries	repand	fleshy 1 aut	Wsh	on ground
ii. GEOPY'XIB. — Cupule at first closed; veil innate ; sporidia simple.				
27367 <i>m&ropus</i> Pers.	long-footed	" solitary 2 su.aut	Cin	shady woods
27368 <i>tuberba</i> Bui.	tuberous	slender 2 mr.ap	Br	shady woods
27369 <i>cupulifera</i> L.	cupped	fringed i aut	Pa.Br	scorched earth
27370 <i>argillacea</i> Sow.	argillaceous	scattered \ aut	Ysh	clay
27^1 <i>granulata</i> Bui.	granular	gregarious A su	O.R	cow dung
21372 <i>reticulata</i> Grev.	netted	g f 4 spring Br	on ground	Grev. cryp. 107
iii. HUMA ^{RIA} . — Cupule a UtUe fleshy, small; veUfiocose only at the edge, or fugacious i sporidua u>*				
27373 <i>crecta</i> Soto.	erect	patches \ aut	shady woods	Sow. M9.10
2131* <i>humosa</i> Fries	earth	scattered Jt aut	C	damp earth

I. DACTYLISCHIUM , — <i>Membranaceous, bursting forth with a separating veil; sportidia simple.</i>	
27375 <i>than Fries</i> split solitary ± winter Br hazel bark	
II. LA'CUNEA Fries. — <i>Waxy, hairy or villous externally; usually on wood,</i>	
i. SABCOBCVPIE .— <i>Cupule fleshy or fleshy-membranous; crust none.</i>	
^376 <i>coccinea Sco.</i> scarlet subgrega. ft spring W dead branches Grev. cryp. 171	
a*cm P'dgndra Sow. black-mouthed solitary J f.mr Pa.Br heaths Sow. 149	
g377 <i>melastoma Sow.</i> black and red solitary faut Bk.R rotten sticks Grev. cryp. 315	
g378 <i>tro-riifa Grev.</i> rooting clustered \ su.aut Su earth in gardens Sow. 114	
radiculata Sow. hemispherical scattered 0 jn.d Brsh earth in woods Sow. 147. <i>hispidia</i>	
hemisph'rica Wig. hairy scattered & au.o Brsh earth Sow. 369.1. <i>hfbrida</i>	
hirta Schum. waxen much crowd. 0 sp.aut Ysh decayed dry wood Sow. 24	
deana Pers. shielded beautiful *sp.aut O old cow dung Grev. cryp. 83	
scutellata L. bird's-nest punctiform 0 aut Br decayed stems Bolt pil. 108.2	
27401 <i>*hirtula* Pers.</i> blue punctiform ft aut Bsh pine trees Sow. 352. <i>equina</i>	
27402 <i>plano-umbril>cata Grev.</i> plano.umbilicat hairy 0 su.aut W decayed nettles Sow. 352. <i>equina</i>	
plano. umbril>cata Grev. plano.umbilicat hairy 0 su.aut Taw cow dung Sow. 352. <i>equina</i>	
dung gregarious TO sp.su R.Br bare earth	
light red handsome TO aut Y decay.herbac stems Grev. cryp. 83	
sulphurea Per*. sulphur pretty 0 aut	
DASY'SCYPBIB. — <i>Cupule waxy, dry, villous; crust none; epiphytes.</i>	
virgin white solitary * aut W rotten Blcks Sow. 65. <i>nivea</i>	
two-colored beautiful * aut W larch twigs Sow. 17	
various-colored gregario. 0 year Ysh rotten wood Sow. 178. <i>hydnoidea</i>	
shapeless depressed 0 aut Och dead branches Sow. 177	
pimpled gregarious 0 year W upon wood Sow. 177	
villous crowded 0 aut W dead herbac. stems Sow. 389. 1. <i>sessilis</i>	
leaden crowded 0 aut Ful.ol rotten wood Grev. cryp. 11	
iii. TAPE^IA. — <i>Cupule waxy or coriaceous, seated on a downy crust.</i>	
anomalous crowded < year Di.Y fallen branches Sow. 309.3. <i>rugbsa</i>	
domestic minute 0 year Ru damp walls Sow. 351	
Wauch's beautiful TO aut Pa.Br dead wood Grev. cryp. 139	
brown spots 0 ap.my Gy dead branches Grev. cryp. 192	
III. PHIAYEAE Pers. — <i>Waxy or membranous, rarely gelatinous, smooth, naked; on wood.</i>	
I HYMENOSCYPIE .— <i>Cupule somewhat membranous, distinctly stalked; hymenium distinct.</i>	
27403 <i>finna Pers.</i> firm gregarious 1 aut ~ Och. Br rotten sticks Sow. 115. <i>ochroleuca</i>	
27402 <i>vinosa Alb.</i> » wine-like minute 0 aut R dead branches	
27403 <i>Persoonii Moug.</i> Persoon's aggreg. 1 aut R bogs on JEquisetum Grev. cryp. 162	
27404 <i>fructigena Bui.</i> fruit-borne clustered } aut Ysh nuts, &c. in woods Sow. 117	
27405 <i>serfina Pers.</i> late clustered 4 spring Y damp shady places Holt. pil. 98	
27406 <i>inflexa Bol.</i> in flexed gregarious } aut Wsh rotten sticks Sow. 306	
27407 <i>campanula Nees</i> bcW-shaped gregarious } aut W rotten sticks Nees pilze 295	
2/408 <i>pedicellata Sow.</i> stalked solitary J aut Wsh rotten sticks Sow. 369.4	
ii. CALYCPINE .— <i>Cupule fleshy, waxy, firm, obconical, somewhat stalked; hymenium distinct.</i>	
27409 <i>tiiba Bol.</i> tubular gregarious ft aut Y fallen branches Bolt pil. 106.1	
27410 <i>calculus Sow.</i> small-cupped gregarious ft aut O.Br rotten wood Sour. 116	
aruginosa Pers. verdigrise destruct 0 su.aut Bt.G damp wood Grev. cryp. 241	
A & R ^{Swed.} Aspergillus Aspergén's gregarious i aut Ysh damp wood Sow. 309.7. <i>blcolor</i>	
cyathoides With. Leinon-coior ^ crowded 4 a Y fallen branches Sow. 150. <i>aUrea</i>	
gl1* Pallescens Pers. pallescent numerous aut Pa.Y old trees Sow. 151. <i>cUrina</i>	
tricolor Sow. three-colored scattered 0 aut Sooty trunks of trees Sow. 369.6.	
> campanula Nees bell delicate & aut W dead twigs Nees pilze, 38.295	
Grev. sieve-like curious 1 aut Bl sandy places	
nettita Grf. bright-yellow punctiform 0 aut Y decayed wood	
herbarum Pers. dotted punctiform 0 aut Y dead leaves Grev. cryp. 63	
herbs crowded TO aut W dead herbac. stems	
iii. MOLLISIA. — <i>Cupule waxy, soft, watery, sessile, or obconical; hymenium confluent.</i>	
cone-borne gregarious 0" aut sp W pine cones Sow. 152	
yellow-haired crowded 0 aut Ful.O posts & rails Sow. 64	
cinereous crowded A aut Ash >osts & rails	
common patches 0 winter Wsh dry bark	
^424 <i>With.</i> transparent scattered 0 winter Tran rotten wood Sow. 389.7	
27491; * ^a 1 ^b <i>pnana Sow.</i> breaking forth scattered 0 aut Css stalks of sycamore Grev. cryp. 99	
iv. PATELLA — <i>Cupule waxy, dry, sessile, flat at base, or innately edged.</i>	
ochreous puckered 0 aut Och Br fallen trunks Grev. cryp. 5	
dark green crowded 0 aut G rotten wood Sow. 389.8	
Abbott's scattered 0 aut Sea G wood	
IV* HELOTIUM Pers. — <i>Plane convex; growing on wood.</i>	
button-shaped gregarious J aut Y trunks of elms Bolt pil. 176	
needle-like gregarious f aud. W hollow oaks Sow. 57 <i>agariciform.</i>	
ASCOBOLUS. (<i>Askos</i> , a bladder, <i>baUo%</i> to eject; seeds ejected from cases.) 1—11.	
scurfy gregarious 0 year Brsh old cow dung Grev. cryp. 307	
27430 <i>BU LGANRIA Fries</i> BULOVIA. (<i>Bulga</i> , a leather bag; its saccated form.) 2. — &	
27430 <i>inuinans Fries</i> staining gelatinous 0 aut.w Umb dead oaks Sow. 428	
** Barcoidea Fries flesh-like polymor. \ aut Pa.R decaying trees Bolt. pil. 101. 2.	
PLA Fries DITIOLA. (<i>Dittos</i> , double, <i>hulos</i> , down; nature of pubescence of velum.) 1. — 5.	
radicata Fries rooting gregarious i ap.jn Go barked pines Fl. dan. 1378.2	
STICTIS. (<i>Stiktos</i> , dotted; dot-like appearance of many species.) 3. — cu	
radiated spots 0 aut W seems & branches Grev. cryp. 227	
changeable oblong 0 year Y.W.Br dead wood Alb. & Schw. 1	
Fir parallel 0 year Bk dead fir wood Hyst i	
CBYPTOMYCES. (<i>Kryptos</i> , hidden, <i>mykes</i> , a minute fungus) 1	
Wauch's firm 0 su.aut Bf. willow branches Grev. cryp. 206	

Class III. TREMELLZ MI.

515a TREM E'LL A L. TREMELLA. (*Tremo*, to tremble; gelatinous, tender, tremulous substance.) 6.—13.I. CEREBRI*N^f Fries. — *Gelatinous; surface at length powdery; form various.*

27439	Mesente <i>Yica</i> Ret*.	Mesent.-like	subsoht	2 aut.sp	Y	fallen branches	Eng. bot 709
27440	albida Hud.	whitish	clustered	1 aut	Wsh	fallen branches	Eng. bot. 2117
27441	intum&ccens E. B.	tumid	twisted lobes	2 wetw	Br	trunks of trees	Eng. bot 170

II. CARYNE Nees. — *Fleshy, gelatinous, somewhat clavate.*

27442	sarco'ides Frfe*	flesh-like	clustered	£ aut	P	-rotten wood	Eng. bot. 2450
27443	clavarisfirms Pers.	Clavar.-lk	gregarious	1 su.aut	DLO	juniper stems	Jac. ic. 648.

III. PHYLLOPTA Fries. — *Somewhat cartilaginous, expanded, leafy.*

27444	biparasitica Fries	parasitical	deformed	i s	Bl	dead agaric	
-------	--------------------	-------------	----------	-----	----	-------------	--

3151. EXI'DI A Fries EXIDIA. (*Exidio*, to exude; sporidia from receptacle.) 4—14

27413	auricula Jbdie Fries	Jew's-ear	tufted	3 aut.w	Blsh	elder trunks	Bolt pil. 107
2	rubescens-fu8ca.fr/Vjf	redL-brown	tufted	3 autw	Rsh	elder trunks	Eng. bot 2447
27446	recls <i>Fries</i>	cutoff	gregarious	f	winter Brsh	dead willows	Eng.bot.1819.bo/rf08rfitis
27447	flaccid* E. B.	flaccid	thin	\ winter	Dark oak bark		Eng. bot 2452
27448	glandulosa Fries	glandular	verygelat	2 aut	Br	dead trees	Eng.bot 2448. <i>Tarbrea</i>

3152. DACRY'MYCES Nees DACRYMYCES. (*Dakry*, a tear, *myke*, a fungus; deliquescent nature.) 2.—7.

27449	monilifrmis Fries	mulberry-lk	sessile	0 aut	Bl	dead wood	Eng. bot 2446
27450	stillatus Nees	trickling	very soft	ft	year O.Y	rotten wood	Grev. cryp. 159

3153. PYRE'NIUMFris PYKENIUM. (*Pyren*, a kernel; appearance of plants.) 1.—2.

27451	lignatile Fries	wood	firm	ft	aut	Fk. W	on dead wood Grev. cryp. 276
-------	-----------------	------	------	----	-----	-------	------------------------------

3154. AGY'R IUM Fries AGYRIUM. (*Agyris*, a crowd; its clustered disposition.) 2.—6.

27452	ceYmm Fries	cessious	punctiform	0 year	Cæs	dead pine wood	
27453	rufum Fries	red	punctiform	0 year	R	dead wood	Grev. cryp. 232

3155. HYMENE'LLA Fries HYMEVEF.LV. (*Hymen*, a membrane; plants) 1.—4.

27454	vulgaris Fries	common	tumid	£ aut	Blsh	nettle steins	
-------	----------------	--------	-------	-------	------	---------------	--

3156. NJEMATE'LLA Fries NJEMATELLA. (*Nuima*, gelatine, *ello*, to contain; nucleus in recept.) 1.—3.

27455	enclphala. Fries	brain	deformed	aut	Flesh	dead pine wood	
-------	------------------	-------	----------	-----	-------	----------------	--

TRIBE II. PYRENOMYCETTES.

Class I. SPHJERIA'CEI.

3157. SPHA?RIA Hal. SFIIJERIA. (*Sphaira*, a globe; figure of plants.) 133.—350.I. CORDICEP9 Fries. — *Compound, club-shaped, simple or branched, stipitate.*

27456	militaris Ehrh.	military	simple	2 aut	R	woods	Sow. 60
27457	entomorizaDrAr.	insect-rooted	stalked	2 aut	Fus	dead larva? of insects	Dick. cr. 1.5.3
27458	gracilis Grev.	slender	simple	3 aut	Br	moist places	Grev. cryp. 86
27459	capitata Holmsk.	capitate	tufted	3 aut	Br	on Scler. cerftaum	Sow. 354
27460	ophioglossoides Ehrh.	Ophiogl.-lk	stalked	2 aut	Bk.Ol	woods	Bui. her. 440.2
27461	alutacea Pers.	tan-like	brittle	2 aut	Tan	dead pine leaves	Sow. 159
27462	digitata Pers.	fingered	tufted	2	year	Bk	stumps of trees Bui. her. 220
27463	polymorpha Pers.	multiform	variable	2 aut	Bk	stumps of trees	Grev. cryp. 237
27464	hypoxylon Ehrh.	on wood	gregarious	2 aut	Bk	old trunks	Sow. 55
27463	carpöpnla Pers.	fruit-loving	long-stmd	3 aut	Bk	on beech mast	Pers. obs. 1.3

II. PORUNIA Pers. — *Compound, cupulate, stalked or sessile, dotted.*

27466	punc&ta Sow.	dotted	gregarious	year	Bk.W	animal dung	Grev. cryp. 327
III.	Pui.viNA'ra Fries	Compound,	subhemispherical, sessile.				
27467	concentrica Bol.	concentric	subcrose	1 year	Bk	ash trunks	Grev. cryp. 324
27468	fragimors Pers.	strawberry-lk	clustered	0 year	K	dead branches	Grev. cryp. 136
27469	fiasca Pers.	brown	subcrose	0 year	Br	dead hazel	Nees pilze 310
27470	multifurmi* Fries	many-shaped	rough	0 year	Bk	dead branches	Sow. 355

IV. CONNIE'E Fries. — *Compound, effused, undefined, plane.*

87471	clrina Pers.	lemon-cld	downy	0 aut	Y	ground, trunks, &c.	Grev. cryp. 215
27472	rubiginosa Pers.	rusty	efflued	0 year	Br.R	dead trunks	Grev. cryp. 110
27473	alpreus Pers.	creeping	effused	0 year	Bk	dead branches	Sow. 372.11
27474	rambsa Sow.	branching	plane	0 aut	Bk	dead wood	Sow. 395.1

V. GLEBOSJE Schmidt. — *Compound, determinate, separating from the subjacent body; perithecia large,*

27475	deusta Hofm.	scorched	unequal	{ aut	Brsh	rotten stumps	Grev. cryp. 324
27476	succentuMa Tode	substituted	oblong	0 year	Bk	dead branches	Sow. 3H. 8
27477	nummulXria Dec.	money	effused	0 year	Bk	dead wood	Sow. 373

VI. LicNosJi Schmidt. — *Compound, determinate, adhering to the subjacent body; bursting forth.*

27478	bullMa Ehrh.	studded	gregarious	0 spring	Bk	dead branches	Sow. 216
27479	undulata Pers.	wavy	broad	0 year	Bk	dead branches	Grev. cryp. 223
27480	stigma Hofm.	dot	eftused	0 year	Bk	dead branches	Grev. cryp. 223
27481	disiformis Hofm.	disk-shaped	gregarious	0 year	Br.Bk	dead branches	Grev. cryp. 314
27482	vurrueafommis/fitrA.	wart-shpd	gregarious	0 aut	Bk	dead branches	

27483 flavovirens Hofm. yellow green gregarious 0 year Bk dead branches Grev. cryp. 320

27484	immersa Sow.	immersed	effused	0 year	Bk	dead hazel	Sow. 374.1
-------	--------------	----------	---------	--------	----	------------	------------

27485 uda Pers. moist sub-immers. 0 year Bk dead wood Sow. 374.4

VII. VERSA'TILKS Fries. — *Compound, determinate, adhering, at length bursting forth; perithecia scattered,*

27486	irregularis Sow.	irregular	prominent	0 aut	Brsh	dead wood	Sow. 374.9
-------	------------------	-----------	-----------	-------	------	-----------	------------

27487 quercina Pers. Oak gregarious 0 aut Bk dead oak branches Nees pilze 321

27488	ferruginea Pers.	rusty	gregarious	0 year	Bk	decayed hazel	• Pers. obs. 1.5.1
-------	------------------	-------	------------	--------	----	---------------	--------------------

27489 elliptica Grev. elliptical gregarious & year R.Br dead beech Grev. cryp. 114

VIII. CONCRESCB'NTES Fries. — *Compound, effused, indeterminate, adhering; the perithecia only emergi*R'*

27490	ftU Pers.	broad	effused	0 year	Bk	dead wood	Sow. 373.9
-------	-----------	-------	---------	--------	----	-----------	------------

27491 million a Fries thousand effused 0 year Gy dead wood Sow. 373.10

27492	velata Pers.	veiled	broad	0 aut	Bk	dead branches	Sow. 373.10
-------	--------------	--------	-------	-------	----	---------------	-------------

IX. CiReuiscBi'PTiE Fries. — *Compound; perithecia converging in an entire receptacle, distinct from epidcrvus.*

27493	smithsa Pers.	prickly	effused	0 year	Bk	rotten wood	Pen. syn. 2.9
-------	---------------	---------	---------	--------	----	-------------	---------------

27494 Prunastri Pers. Plum gregarious 0 year Bk dead sloe branches

27495	sr.rbi Schmidt	Mountain Ash	roundish	0 year	Bk	dead mountain ash	
-------	----------------	--------------	----------	--------	----	-------------------	--

27496	uornicul&ta Ehrh.	horned	sub-immers.	0 >nar	Bk	dead branches	
-------	-------------------	--------	-------------	--------	----	---------------	--

97 <i>in?</i> ^{INCUS FTICS}	<i>Compound</i> ; peritheda converging; stroma dimidiate, adnate with the epidermis.
VIZZ? ^{?ivea HoA*}	snowy pulverulent 0 year W dead branches Sow. 219
eucostoma P<<	pale-mouthed imbedded 0 year W dead branches Sow. 372.7
XI. OBVALLK X Fries.	<i>Compound</i> ; stroma spurious; orifices collected into a disk.
eiphania Fries.	pale cortical 0 spring Pa dead oak branches Sow. 218?
alicina Pers.	Willow circular 0 year Wsh dead willow Tode fun. 107
XII. CIRCINAVJE Pers.	Nearly simple; peritheda in a drce beneath the epidermis.
illa Per*, pretty depressed 0 year Bk dead beeches Grev. cryp. 67	
quaternata Pers.	quaternate circinate 0 year Bk dead branches Sow. 374.6
Igu* convergens Tode	convergent subinxmers. 0 year Bk dead branches Sow. 374.6
ow, . , XIII. GESPLITPOSJB Pers.	Peritheda tufted on a subjacent stroma.
cinnabrina Tode	vermilion prominent 0 year 11 dead branches Grev. cryp. 135
e	scarlet tufted 0 year R dead trunks Sow. 255
Tode	currant tufted 0 year Br.R dead currant Tode fun. 103
Per.	Laburnum tufted 0 year Bk dead laburnum Nees pilze 325
nii Sow.	Berberry tufted 0 year Bk dead berry Grev. cryp. 84
Pinaster	cup-shaped tufted 0 winter Bk dead branches Pers. syn. 1.9
Pinaster	nedt-like tufted 0 winter Bk dead whortleberry Sow. 373.1
Pi	tufted 0 year Bk bean roots Sow. 394.2
27511 ^{XIV} CONFLUE'NTES Fries.	Peritheda emerging from the stroma, at length confluent.
27512 ^{et on 8 Hta} Fries	long black cracks 0 year Bk furze branches Grev. cryp. 193
'n melogr&mma Pers.	black-lined lines 0 spring Bk dead beeches Bui. her. 492.
27iii: *-ir . XV. SERIA'NIVE Fries.	Petithecia immersed in the stroma, in parallel lines.
Sn%J <i>J</i> l <i>cxcl*</i> Fries	Fern lines 0 spring Bk dead brake stems Sow. 394.10
27513 ^{Hki} Fries	Rush interrupt lines 0 year Bk dead rushes
27*10 ^{strifc*} orpini Fries	channel-fmd crowd, lines 0 aut Bk dead herb stems
S% <i>a</i> arundinae Sow.	Reed conn, lines 0 spring Bk dead reed stems Sow. 336
nebulosa Pers.	cloudy dim spots 0 spring Cy dead herb steins Nees pilze 341
XVI. CONFERTC Fries.	Stroma little or none; peritheda innate in the substance of leaves, crowded.
Taminis Fries	Grass l'onfl.patch. 0 aut Bk dead grass leaves Nees pilze 314
fimbriata Pers.	fringed little mass 0 aut Bk hornbeam leaves
Coryli Batsch	Hazel circles 0 su live hazel leaves Grev. cryp. 330
ons Schmidt	two-fronted scattered 0 year Bk dead oak leaves Sow. 373.4
XVII. BXBBISE^DC Fries.	Simple; peritheda glabrous, on a subjacent tomentose stratum.
27514 ^{aureo-rr} Tosevita A.S.	Orange-dd byssoid 0 aut O decaying fungi, &c. Grev. cryp. 78
27515 ^{bysaidea} Tode	pink byssoid pat. 0 aut Pk moist wood, &c. Grev. cryp. 138
27527 tristis Tode	flax-like patches 0 spring Gy.Br dead wood Tode fun. 69
2528 lignaria Grev.	melancholy byss. spots 0 spring Bk dead trunks Pers. ic. pict 12. 5
*1529 ovina Pers.	wood emerging Simple peritheda hairy downy. dead wood rotten wood Grev. cryp. 82
*7030 canescens Pers.	fleecy moist 0 year Wsh rotten wood Sow. 373.12
Racodium Pers.	canescent hairy 0 aut Pa rotten wood
g133 hirsuta Fries	two-formed hairy 0 spring Bk ground, and rotten wood Sow. 37a 7
cruita Pers.	Racodium broad 0 aut Bk dead wood
Pilosa Pers.	hirsute aggregated 0 aut Bk dead wood Sow. 386.3 ?
nispida Tode	hairy imbedded. 0 year Bk soft decayed wood
Ilva Tode	pilosus short-hair. 0 aut Bk dead wood Pers. ic. pict 10.9
8240 XIX. DENUDA^TJE Fries.	seed-like crowded 0 aut Bk dead branches' Tode fun. 84
87*! f Grev,	bald shining 0 aut Bk dead branches Tode fun. 10.83
S8 S Pertla Pers.	golden crowded 0 aut O decayed large funei Grev. cryp. 47
27541 ^{ai} finea Sib.	blood-red minute 0 spring R dead wood * Grev. cryp. 175
27542 ^{all in} w.Grey.	allied minute 0 aut R B&ngia atrov'rens Grev. cryp. 186
pomiformis Pers.	appleleshued minute 0 year Bk dead wood Sow. 354
27543 ^{all in} ombidra Batsch	omnidra miniform 0 year Bk rotten wood Sow. 354
spermoldch Hofm.	musket long 4 year Bk rotten wood Noes pilze 357
moriformis Tode	seed-like crowded 0 aut Bk dead branches Sow. 3J. J7. 2
97K* v'ruchsa Grev.	mulberry-shpd wrinkled 0 year Bk dead Polyp, abietina Grev. cryp. 39
H-H9 puvis, pyrius Pers.	warty scattered 0 aut Bk dead wood Grev. cryp. 152
pulveracea Elrh.	chimney-dust crowded 0 year Bk dead wood
myriocarpia Fries	dusty crowded 0 year Bk dead wood Grev. cryp. 152
XX. PERTU*SA Fries.	Simple; peritheda glabrous, flattened at the base.
nucula Fries	small nut minute 0 year Bk oak trunks
27544 ^p XXI. CERATOSTOMA Fries.	double volva minute diokO spring Bk dead alder branches
27554 ^p ^{info} fa Fries	hair-bearing naked 0 year Bk dead pine wood Nees pilze 354
*** cirrus Pers.	tendrilled emerging 0 year Bk soft dead wood Pers. ic. pict. 24.3
27555 ² XII. OBTE'CTA Fries.	Simple, quite immersed, protruding a neck which contains the orifice.
27556 ^u ^{to} Pries	well marked minute spotu 0 year Bk dead branches Sow. S17
27557 ^a ^{to} lba Fries	double volva minute diokO spring Bk dead alder branches
27558 ^a ^{to} lba Fries	quivering concealed 0 aut Bk dead plum branches
S7 Ca ² ^{brut} >8 Flies	staining black spot 0 aut Bk dead branches Sow. 372.3
oo Inquinans Tode	inhabiting minute 0 year Bk dead branches
27559 ^{terci?} Sow.	Arbutus dwell, confluent 0 year Bk dead ^{rb.} UV TJrsi Sow. 370.6
strobillina Holm	Honey-suckle cracks 0 year Bk dead honeysuckle branches Sow. 393. b
XXIII. QBTURAV* Fries.	Simple, innate, at length bursting from the epidermis.
cone irregular 0 year Bk dead fir cones	
XXIV. SUBTE'CTJJ Fries.	Simple, innate, the upper part at length exposed.
57E* ¹ mariscinus Grev.	Tanisk immersed 0 year Bk dead ^{ingerm.} Grev. cryp. 45
**TM semiuimmersa Pers.	half-immers. shining 0 year Bk dead honeysuckle
1. Klinc-headed grcga.	0 year Ilk dead branches Sow. 375.2 ?
2. Sow.	inhabiting minute 0 year Bk dead branches Sow. 370.5
3. Sow.	Arbutus dwell, confluent 0 year Bk dead ^{rb.} UV TJrsi Sow. 370.6
yrene Tode (0)	Yew distinct 0 year Bk dead yew leaves Grev. cryp. 13
XXV. CAU'COLIB Fries.	Simple, concealed, at length exposed by the deciduous epidermis.
acuminata Sow.	pointed gregarious 0 year Bk dead thistle stems Sow. 3M 3
curvirostra &ni	crooked-beaked gregarious 0 year Bk dead Umbelliferae Sow. 373.5
acista Hofm.	acute dot-like 0 year Bk dead Uinbellifcrs Grev. cryp. 239

27571 millepunctata	Gren.	IOOO-dotted	minute	0	year	Bk	dead ash branches	Grev. cryp. 201
27572 complanata	Tode	flat	punctiform	O	year	Bk	dead herb stems	Tode fun. 88
27573 doliolum	Pers.	little-cask	conical	0	year	Bk	dead herb stems	Grev. cryp. 239
27574 Piai	Sow.	Pea	scattered	0	winter	Bk	dead peas haulm	Sow. 393.8
27575 herbirum	Pers.	herbs	punctiform	O	year	Bk	dead herb stems	
27576 nigrella	Fries	blackish	long spots	0	spring	Bk	dead hemp nettle	Fries obs. 4.8
XXVI. FOLI'COLJE Fries. — <i>Simple, adnate, concealed, not on a circumscribed spot; epiphyllous.</i>								
ZISTI tubsformis	Tode	tube-shaped	scattered	0	winter	Br	dead tree leaves	Grev. cryp. 335
27578 gnbmon	Tode	gnomon	punctiform	0	winter	Bk	dead hazel leaves	Grev. cryp. 3J5
27579 scftcea	Pers.	bristly	minute	0	winter	Bk	dead leaves	
27580 So\w\ Pers.	Potato	punctiform	O	aut		Bk	rotten potatoes	
27581 petioVorum	Schutein.	petioles	rigid	0	aut	Bk	dead leaf-stalks	
27582 duplex	Sow.	double	immersed	0	aut	Bk	dead reed-mace, &c.	Sow. 375.4. & 394.7
27583 /ISJera Sow.	Ivy	scattered	0	year	Bk	dead ivy leaves	Sow. 371.5	
27584 £'mpetri	Fries	Crake-berry	punctiform	0	year	Bk	dead crake-berry leaves	
27585 microscopica	Grev.	microscopic	dot-like sp.	0	year	Bk	dead Port laurel leaves	
27586 macuteformis/Vr*		spot-shaped	punctiform	0	year	Bk	dead leaves	Sow. 370.7
27587 punctifrmis	Pers.	dot-shaped	shining	0	year	Bk	dead oak leaves, &c.	
27588 iEgop&ddii Pers.	Gout-weed	spots	0	aut	Bk	dying gout-weed leaves		

3158. VERMICULARIA Tode VEEMICULARIA. (*Vermiculus*, little worm; arrangement of seeds) 1. — ▲

27589 trichella Fries small hairy minute 0 aut Bk dead ivy leaves — Grev. cryp. 345

3159. DOTH'DEA Fries	DOTH IDE A.	(<i>Dothien</i> , a tubercle, <i>eidos</i> , similar.)	8.—54.
27590 ribesia Fries	Currant	elliptical	0 year Bk
27591 fletiMina Fries	Birch	shining	0 su Bk
27592 t'l'mi Fries	Elm	convex	0 aut IronGy
27593 Ptc-ridis Fries	Brake	irregular	0 aut Bk
27594 Gertnii Fries	Geranium	spots	0 su Bk
27595 Himantia Fries	Himantia	broad	0 aut Bk
27596 dlnea Fries	Alder	numerous	0 aut Bk
27597 Roberta Fries	Herb Robert	Bhining	0 su Bk

3160. POLYSTI'GMA Pers. POLYSTIGMA. (*Polys*, many, *stigma*, a mark.) 2.

27598 rubrum Pers. red innate 0 aut R live sloe leaves Grev. cryp. 180

87599 fulvum Pers. fulvous innate 0 aut Br.R live bird-cherry leaves

Class II. PHACXDIVCEI.

3161. TY'MPANIS Tode TYMPANIS. (*Tympanon*, a drum; resemblance of the young velum.) 2.-8.

27600 a'lnca Fries Alder minute 0 spring Bk dead alder branches

27601 conspersa Fries sprinkled crowded 0 year Gy dead branches Grev. cryp. 3.38

3162. CENANGI'UM Fries CENANGIUM. (*Kenos*, empty, *aggeion*, vessel.) 5.-~30'

27602 Prunastri Fries Plum tufted 0 aut Bk dead plum branches

27603 Cerasi Fries Cherry variable 0 year Bk dead cherry branc. Pers. ic. pict 10. 1

27604 Aucu|aris Fries MountainAsh tufted 0 aut Gy dead mountain ash

27605 ferrugintosum Fries rusty tufted 0 aut RBr. on dead firs Grev. cryp. 197

87606 guercinum Fries Oak waved 0 year Bk.Gy dead oak branches Sow. 373. 3.

3163. HETROSPHf RIA Grev. HETROSPHARIA. (*Heterozoyarinableplana*, *sphaera*) 1.-T8.

87607 paella Grev. dish shining 0 year Bk dead herb stems Grev. cryp. 103

3164. LCTPHIUM Fries LOPHIUM. (*Lophion*, a hillock; completely evolved.) 2.—3.

27608 elatum Grev. tall scattered & year Bk pine bark Grev. cryp. 177

27609 mytilnum Fries limpeUshaped crustaceous O year Bk pine bark Grev. cryp. 177

3165. RHYTI'SMA Fries RHYTISMA. (*Rhytis*, a wrinkle; plants.) Xybmact authors, &—

27610 corrugatum Fries wrinkled gregarious 0 year Bk crust of lichens Eng.bot.464. L. gruith

27611 Andromedae Fries Andromeda shining 0 aut Bk live wild rosemary

27612 jal'cinum Fries Willow solid spots 0 aut Bk liv.SM capne^alvs Grev. cryp. 118

27613 acerinum Fries Maple broad spots 0 su Bk liv. sycamore lvs Grev. cryp. 118

27614 punctatum Fries dotted gregarious 0 su Bk liv. sycamore lvs

27615 trtla Fries Nettle broad spots 0 aut Bk dead nettle stems

3166. PHACI'DIUM Fries PHACIDIUM. (*Phakos*, a lentil, *eidos*, similar.) 4.—SO.

27616 coronatum Grev. crowned black spot 0 year Bk dead oak leaves Grev. cryp. 58

27617 denfatum Schmidt toothed white spot 0 year Bk oak leaves

27618 Pni Fries Fir disk-like 0 year Bk Scotch fir trunks Nees pilze 399

27619 Afibi Fries Bramble confluent 0 spring Bk raspberry leaves

3167. HYSTE*RIUM Tode HYSTEBIUM. (*Hysteresis*, penury; appearance of infested trees.) 16. - 5 *

27620 lineare Fries linear lines 0 year * Bk dead wood Grev. cryp. 167

27621 maculare Fries pale spotted blotches 0 aut Bk dead leaves * Grev. cryp. 129

27622 JZhbi Pers. Bnmh'e-stnd lines 0 aut Bk bramble branches Grev. cryp. 84

27623 folicolum Fries leaf-inhabiting dot-like 0 spring Bk common ivy leaf Grev. cryp. 189

27624 melale&cum Fries blk & white dots 0 au Bk Fac. m's dafli Grev. cryp. 88

27625 pulicare Pers. flea-like verygrega. 0 year Bk rugged oak bark Grev. cryp. 167

27626 FMxini Pers. Ash corneous 0 year Bk dead ash branches Grev. cryp. 78

27627 guercinum Pers. Oak gregarious 0 year Gy. Br dead oak branches

27628 angustatum Pers. narrowed minute 0 year DL.Bk dead wood & stumps

27629 Pinistri Fries Pinaster scattered 0 year Bk dead Scotch fir lvs Grev. cryp. 60

27630 Juniperi Grev. Juniper spots 0 year Bk dead juniper leaves Grev. cryp. 80

27631 gramneum Pers. Grass gregarious 0 year Bk dead grass leaves Grev. cryp. 87

27632 vrium Fries variable unequal 0 year Bk oak bark Grev. cryp. 233

27633 decgnerans Fries inconstant emerging 0 year Bk dead bog whortle-berry branches

27634 arundinkceum Schr. Reed innate 0 spring Bk dead reed culms Sc. sam. 2. 3.3

27635 conigenum Moug. cone-sprung punctiform O year Bk dead fir cones

3168. EXCI'PULA Fries EXCIPULA. (*Excipio*, to catch j roughness to the touch.) 1;-

S7636 A&bi Fries Bramble punctiform 0 aut Bk dead raspberry stem Grev. cryp. <

Class III. CTTISPO'REI.

3169. SPHERONJETMA Fries. SPBGERONJEMA. (*Sphaira*, sphere, *naima*, gelatiae j receptacle) 1.-1*

87637 subulatum Fries awl-shaped spiculiform & aut Cin on agarics Grev. cryp. "

- | | | | | |
|-----------------|------------------------------|------------------------------------|--|--------------------------------|
| 37IX | SEVT&RIAFHes | SEPTORIA. | /Septum, a division; sporidia.) | L *** *, |
| 27638 | tfimi JS^M | Elm-fetf stains | 0 aut Br elm leaves | Grev. cryp. 112 |
| 3171. | CYTISPORA | Ehren. CYTIBPORA. | (Kystis, a bladder, spora, a sporule.) | 4. --18- W ^a iaSow. |
| 27639 | chrysosperma | *W ^a yellow-seed, spots | 0 year Bksh poplar bark | Sow. 13 arrh Uta |
| 276W | W a r u s G T ^m . | É t o f c pustular | 0 Jut Pallid dead rose branches | Grev. cryp. SO |
| gejUeucosperma^ | pale-sVed& | scattered | 0 year W dead branches | Hof. cryp. I. |

3172 CEUTH(ys)PORA *Fries*. CEUTHOSPORA. (*Keutho,to* hide, *spora*, « JP^oTM^{le}; ») * - ?
 SS²e¹idioWesG*<1.*; Phacidium-lk innate 0 year Bk dead holly leaves Grev. cryp. 253
 2 S¹ kitiri *Grew.* Laurel-pustulate 0 year Bk dead laurel leaves Grev. cryp. 254
 27645 pha2cones *W* Laurel-haired minute 0 aut Bk dead H61cus mttlis Grev. cryp. 69

3TO. PHOBIA Fries
g.6.16 «aligna Fries
\$ Populi Fries
^wB pustula Fries

PHOMA. (PAcw, a pustule; appearance of plants.) *iz* L. *Lycoperdon*
Willow-felt pimpled 0 w.sp Brsh dead willow leaves Sow.
Poplar-^ pimpled 0 w.sp Test dead poplar leaves Sow. 374. 2
pustule-W^ prominent 0 ant Br dead oak leaves

Class IV XTLQOMA^CEX

- BJJE⁴**; ACTINOTHY'RUM *Kitnxē* ACTINOTHYRIUM. (*Aktin*, a ray, *tkyron*, a little door.) 1.
²w9 Graminis *Kunze* Grass gregarious 0 spring Bk culms of grasses Grey, cryp. 818

.Si' »™ 0 P Bk <JeJl»e.of.ricc. Frie.oto.a7-4

	SB	Kurd	wjtUn	0	aut	Bk	dead	laurel	leave*
SI7a	ASTEBO^MABw	ACTEROXI-		(i*r,»iUi.)	,				*• — ■
?	i n n	G H m	Pde.pot.	0 year	Bk	Mving elm leave.			
	Grev.	Lad?	Ubnble pale port	«y««	SJ	& ^ e i 3 £ " f t			

TRIBE III. GASTEROMYCFIATES.

Class I. ANGIOGA'STRES—Division I. *PhaUotdete.*

- | | | | |
|---|--|---|------------------------|
| 3179. PHA'TT nan/* | PHALLUS. | (<i>Phallos</i> . Gr.; resemblance.) | 2.-9. |
| ^I ¹ [^] foytidus PRS | ^I ¹ [^] stolid Mor% | fetid 8 su' Wsln woods fit thadges | Grev. corp. 213, & 214 |
| [^] SESSET | dog 4 smaller | aut aut Plpk rotten hazel trunks | Sew. 330 |
| 3180. BATT'A'RREA Pers. | BATTARREA. | { <i>A. J. Battarra</i> , a cryptogamic boUnist.) | 'I. |
| [^] 665 phallodes Pers. | Phallus-like long | 7 aut Br banks | San. spic. 1.12. |

Division II. *Tuberaceae*

3181. *TITBER* Mr. *TRUFFI**. (An ancient Roman name for a fungus.) S.—5.
666 *cibarium* *Sibth.* food esculent 1st winter Bui. her. 356
~~biduum~~ *Cas.* whitish less fragrant & su. aut. Wsh underground

RHIZOPCPGON *Fries* RHIZOPOCON. (*Rhhn.*, root, *pajon*, beard.) 1.—4.
 *albu*"*iffii* white flocculent \ aut W.R waysides Bui. her. 404
*Lyco*perdon *gibb*sum *Dick.*

Division III. *Nidulariaicei*.

- NIDULA**[^]**RRIA** *Bid. NIDULARIA.* (X/rfw.nest; planU consist of cups containing egg-like seeds.) S.-12.
 rtriaai?!!.. striated gregarious J aut Br rotten leaves Sow. 29. *htshtta*
Panulata *Sib.* bell-shaped flocculent t' aut Gjr rotten wood Saw. 28
Crucibulum *Pers.* crucible coriaceous j aut Ysh dead wood Grev. cryp-»
 ■ 3J84. **MYRCIOCCCCUM** *JWf** **MYRCOCOCCUM.** . (Myrio*, myriad, *o**o», berry.) 1.
 pr^coxj^i^riy confluent i spring W dead leaves
 ■ 3185..**POLYANGIWI** *Lk* **POLYANCIUM.** i^j^i^i^ ^ ! T! S r t t S! damp trunks
 eVStitellinL* yolk-<forth! gregarious 0 aut Y

Division IV. *Carp&boH.*

- 87^86. ATRACTO'BOLUS *Jbde* ATRACTOBOLL.B. C^Ajf*^aaplncJjJW*toc^ect) 1
 *7674 ubiquitaTius *Tode* universal powdery 0 year W wood, stonen.^{cc.} lode run, 9

3187. THELE'BOLUS 7bA; THIBLEDOLUS. (*Thele*, a nipple, *Wfljj* to eject.) 1 — 3.
fcte sturcSreuTpS2T dimg gregarious A ausp Ysh cow dung Neepsilpx363

3188. SPHJERO'BOLUS Tode SPIAROBOLUS. (*Sphaira*, a sphere, *boleo*, to eject.) 1. — 2-
27G76 stellktus Tode starry emerging ^ su.aut Pa. Y dead wood, &c. Grev. cryp. 158

3189. PILO'BOLUS Tode PILOBOLUS. (*Pilos*, a cap, *boleo*, to eject.) 1.
27677 crystallinus Tode pellucid fugacious \ year Bk horse dung Bolt pil. 133. 1
2 "xondus Pers. dewy fugacious } year PI horse dung Bolt. pil. 133.4.

Class II. TRICHOSPE'RMLI. — Division I. *Sclerodermat*

3190. SCLEItODETIMA Pers. SCLERODERMA. (*Skleros*, hard, *derma*, a skin; nature of coat) 4.—U-
27678 verrucosum Pers. warty handsome 5 aut Y.Br plantations Grev. cryp. 48
27679 Cipa Pers. Onion Burf.variab.2 aut Y.Br plantations Grev. cryp. 66
Tuber s6lidum With.
27680 dtrinum Pen. lemon-cld tessellated 2 aut Pa.Y about oak roots Bolt. pil. 116
27681 spadiceum Pers. light red tessellated 1 su Pa.Br beech trunks Schaeft: 188

Division II. *Lycoperdinei.*

3191. LYCOPE'RDON Mx. PUA-BALL. (*Lykos*, a wolf, *perdo*, crepto; application not known.) 5. — 11-
27682 Bovista Pers. Bovista turbinate 3 aut Wsh pastures Sow. 332. *Proteus*
27683 pratlnse Pers. meadow \ subterra. 2 su.aut W pastures Bui. her. 435.2
276K4 excipuliflme Pers. Excipua-lk chan. to br. 2 aut W pastures Bui. her. 430 2
276H5 pyriforme Pers. pear-shaped tufted 1} su.aut Pa.Br about tree stumps Grev. cryp. 304
27686 perlatum Pers. pearlized tufted 3 aut W woods Fl. dan. 1120

3192. BOV'ISTA Pers. BOVISTA. (Latinised from *bafist*, its German name.) 2. - 4.
27687 nigr&cens Pers. nigrescent becom. blk. 2 su.aut W pastures Sow. 331
27688 gigantea Nees gigantic cracking 12 su.aut Y.W pastures Grev. cryp. 336

3193. GEA'STRUM-Mx. GEASTRUM. (*Ge*, the earth, *aster*, a star; stellate appearance.) 4. - 5. -
27689 coliforme Pers. distaff-shaped subsolitary 2 aut Brsh pastures Dick cr. S. 4. *Lycoper**-
27690 Woodwards Pers. Woodward's subsolitary 1 aut D.Br dry banks *Bry.h\stA9.Lycoperdon*
27691 quadrifidum Pers. quadrifid subsolitary 2 aut Wsh pine woods Schaeff. 183. *L.sformicatutn*
27692 multifidum Pers. many-cleft subsolitary sp.aut Br moors Grev. cryp. 305
Lycoperdon recoilgens Woodw.

3194. TULO'STOMA Pers. TULOSTOMA. (*Tulos* a wart, *stoma*, the mouth; nature of orifice.) 1. — 3-
27693 bruraklc Pers. winter subsolitary 1 au.o W.Br pastures Grev. cryp. 340

Division III. *Physarei*

3195. LYCCGALA Mx. LYCOGALA. (*Lykos*, a wolf, *gala*, milk j early state like a mass of thick cream.) 1. — ?
27694 miniata Pers. vermillion granular 0 sp.aut R rotten wood Grev. cryp. 38

3196. DIDY'MIUM Schr. DIDYMIUM. (*Didymos*, double; inner and outer peridium.) 1. — ?
27695 globosum Lk. globose sessile ft aut Gy dead leaves Grev.cryp.122. *Dia"\,77;*

3197. PHY'S ARUM Pers. PRYSARUM. (*Physa*, a bladder; appearance of peridium.) 6. — ?
27696 sulcatum Lk. furrowed weak ± sp.aut Gy rotten wood
27697 nititans Pers. nodding weak i aut Gy rotten wood Bui. her. 407.3
27698 nigripes Lk. black-stnd firm ± aut D.Gy rotten wood Stur. dcu. 42
27699 ylide Pers. green rather weak } aut Y.G rotten wood Bui. her. 481. L
27700 lecopus Lk. white-stnd very stiff ft aut Cl dead beech wood
27701 adruum Pert. golden gregarious sp.aut Y decuyinff trunks Grev. cryp. 185

3108. LEQCA'lIPUS Lk. LKOCAKPUB. (*Leios*, smooth, *karpoa*, fruit j peridium.) 1. — ?
27702 vernicous Lk. varnished encrusted ft aut R.Br stems of grasses Grev. cryp. III
Lycoperdon fragile Sow.

3199. LEANGTUM Lk. LEANGIUM. (*Leios*, smooth, *aggeion*, a vessel; peridium.) 2. — ?
27703 florifbrme Lk. flower-like scattered & aut Y decaying trunks BuL her. 371
27704 Trevelyan's scattered A aut Pa.Br leaves of mosses Grev. cryp. 132

3200. PERICHJFNA Fries PBRICHJENA. (*Peri*, around, *ischan*, to compress.) 2.-4.
27705 strobillina Fries Fir-cone crowded & year Br cones of *P.A'bies* Grev. cryp. 275
27WJ p\puluia Fries Poplar gregarious A aut Br. Y dead trunks Grev. cryp. 252

3901. CRATETRIUM Trent CRATERIUM. (*Krater*, a cup; form of peridium.) 2. — 6.
27707 leucocephalum Trent, white-head pretty | aut R.Br mosses, &c. Grev. cryp. 65
27708 vulgare Dit. common pretty | aut R.Br mosses, &c. Sow. 239
Cyatthus minutus Sow.

3202. LI'CEA Schr. LICEA. (Meaning not explained.) 1. — ?
27709 fragiformis Nees strawberry-lk pulpy 0 aut DLR rotten wood Grev. cryp. 308

Division IV. *Angioridii.*

3203. ANGIORI'DIUM Grev. ANGIORIDIUM. (*Aggeion*, a vessel, e*Aos% like.1 1.
27710 sinubsum Grev. waved gregarious t aut W dead sticks, &c. Grev. cryp. 310
Reticularia sinuosa Sow.

Division V. *TrichidceL*

3204. TRI'CHI A Pers. Tn. (*Thrix*, hair; internal mass of elastic fibres expanding after head bursts.; 4.-
27711 reticulata Pers. netted pulpy 0 aut Ysh rotten wood Grev. cryp. —
27712 ovata Pers. ovate crowded 0 aut Y rotten wood Sow. 85. *turomuw*
27713 fallax Pers. deceitful variable 0 aut Rsh rotten wood Sow. 279
Spheroscirpus frigilis Sow. , OQI in i.art.
27714 nUens Pers. shining gregarious A aut Rsh.Y rotten wood Grev.cryp. 281. m I

- | | | | |
|---------------------------------------|-----------------------------|---|---------------------------------|
| 3205. ARC Y'RI A Pen. | <i>ARCYRIA</i> | (<i>Arkys</i> , a net; sporules fastened together by a network of fibres.) | 4-? |
| "lopuniceaPm. | crimson | gregarious ft su.aut C | rotten wood Grev. cryp. 130 |
| 277.fl T'chriadenu?itaScnw. 29 | | | |
| "#nava Pers. | yellow | weak ft su.aut Pa.Y | rotten wood Grev. cryp. 309 |
| *#47 mcarntka Pers. | fash-colored | conical ft aut Pk | rotten wood |
| * " « leucocephala a>/in. white-head, | gregarious ft aut | W | rotten wood Hofgr. 2.6.1 |
| ██████████ STEMON nms Pers. | STEMOXma. | (<i>Stemon</i> , a stamen; form of some species.) | 2.—' |
| p'culata Pers. | fascicled | dense f su.aut Bk Br | rotten wood Grev. cryp. 170 |
| Lncfianftda&n.u. | | | |
| ██████████ pimpled | scattered ft aut | D.Br | rotten wood Nees pilze 10.118 |
| ██████████ pRIBR A%RI A Schr. | <i>CRIRRARIA</i> . | (<i>Cribrum</i> , a sieve; upper part of peridium.) | 1-? |
| micropus Schr. | small-stmd pin-headed & aut | Br | pine trunks Sc. gen. nov. 2.1-2 |
| 3208. piCTY'DIUM | <i>Schr. DICTVDIUM</i> | (<i>Dikyon</i> , a net, <i>eidos</i> , similar; peridium.) | 1-? |
| "<ecernuumMv« | drooping pin-headed & year | Bk | rotten wood Grev. cryp. 153 |

Class III. TRICHODERMA'CEI. — Division I. (*Ethalinu*)

- | | | |
|---|--|-----------------|
| A Per*. SPUMARIA.
go Pert. Mucilage frothy | (<i>Spuma</i> , froth; appearance.) | 1.—? |
| Lk. JETHKUV.
yellow flocculent | (<i>AUhales</i> , splendid ; appearance.) | 1.—2. |
| 27701: KETICULALIA Bui. RBTICULARIA.
2770A ff Kcntea Bui silvery fragile | 1 aut Y.Br on bark in stoves | Grev. cryp. 272 |
| W ^s minuta Grew. | (<i>Reticulum</i> , a net; appearance.) | 2—? |
| | \ aut W rotten wood, &c. | Grev. cryn. 106 |
| | 0 aut W dead leave and sticks | Grev. cryp. 40 |

Division II. *Trichodermei.*

- | | |
|---|----|
| AMPHISPO^oRIUM Lk. AMPHISPORIUM. (<i>Amphi</i> , on either side, <i>spora</i> , a sporule.) | 1. |
| <i>versicolor</i> L*. changeable spots 0 winter Y hyacinths in glasses Nees pilze 100 | |
| LE Fries INOTITALE. (<i>TnttUa</i> , a fringe + j appearance of plants.) | 2. |
| <i>radiating</i> woolly ft aut Br on plastered walls Sow. 145 | |
| <i>fgariciftrml&Frto</i> Mushr.-like downy & «* B* rotten wood Sow. 146 | |
| <i>Mcooperdon radfttum</i> and agariciforme Sow. | |

Division III. PUicrinu

- ASTERO'PHORA** *Dit.* ASTEROPHORA. (*Aster, star, phoreo, to bear.*) 1.
ycopcrdoldes *Dit.* Puff-ball-like woolly 1 aut Rsh rotten mushrooms Bui. her. 519.1
3215. ON Y'GEN A Pers ONYGBNA. (*Onyx, hoof, genos, offspring; on hoofs and horns of animals.*) 1. — S.
~~equlni~~ *Pers.* horse-hoof minute ft aut Wsh decaying hoofs and Grev. cryp. 343
similar substances

Division IV. *Afyrothecin.*

27. ^{DI}*CHOSPO^YRIUM Nees DICHOSPORIUM*. (*Dicha*, double, *spora*, sporule; double coat of peridium.) 1.
" " ^{aggred}^{tum} *Nees* clustered *spoU* 0 aut Bk bark of trees *Neespilse* 99
Spumkria physartidea Pers.

[■] MYN OT III - rci UM 7W- MI miTiiKt HIM. (*Muru*, to tlatl, /AcJrf, tlioulim.) 1.-&
michaeliaUrcv. Carmelchad'N greffarlow* O out W rotten bog matter Grev. cryp. 140

[■] TRICHODE'RMA Pers. TRICHIODRRMA. (*Thrix*, hair, *derma*, skin; downy covering.) 1.-?
hide Pers. green pulvinate ft aut W.Gr dead branches Grev. cryp. 271

Class IV. SCLEROTIA'CEJE. — Division I. Rhizobnei.

9. RHIZOCTONIA Dec. RHIZOCTONIA. (*Rhixa*, root, *kto* to destroy; *root* it grows on.) 1—3.
Crocorum Itec. Crocus-blight clustered 0 year Rsh saffron roots Neespieli SJ

Division II. *Apiospbrei.*

- CH^TO'MIUM** *Kunxe* CHOTOMIUM. (*Chaita*, a bristle; hairy appearance of plants.) 1.—S,
 &tumjfimtf tall bristly &aut Bk damp straw, &c. Grev. cryp. 230

2 ^ 1. ILLOSPO^XRIUM *Mart.* ILLOSPIRIUM. (/to, to envelope, *sporos*, a sporule.) 1.—?
 **i* *roseum* *Fries* rosy minute 0 aut Pk on *Borrera cilians*, &c. Grev. cryp. 51

Division III. *SderbtU.*

- | | | | | |
|---|--------------|----------------------------------|--|-----------------|
|  PERKFLA | <i>Fries</i> | PERIOLA. | (<i>Peri</i> , about, <i>ioulos</i> , down; appearance.) | 1. |
| tomentbsa | <i>Fries</i> | downy scattered | ft spring W potato roots | |
|  SPERMCE^DI | A | <i>Fries</i> ERGOT. | (<i>Sperma</i> , <i>aed,oidro</i> , to swell i diseased seeds.) | 1. - 2. |
| clivus | <i>Fries</i> | spike, <i>horn-shpd</i> nauseous | 0 su Bksh glumes of grasses | |
|  ACROSP'RMUM | Tode | ACROSPBRMUM. | (<i>Akros</i> , the summit, <i>sperma</i> , seed; situation.) | 2. — 5. |
| ornbtum | <i>Fries</i> | eonnSn gregar | JL aut Rsh putrid agarics | Bui. her. 256 |
| compressfimSwc. | SSfpPrSised | Icattered | «* aut Bksh dead herb stalks | Grev. cryp. 182 |
| 27?S ⁵ | SCLERO'TIUM | Tode | SCLEROTIUM. (<i>Sklerotes</i> , hardness; texture of plants.) | 11.—54 |
| 27?S "ajellatum Alb. | shield-like | regular | 0 spring Br dead leaves | Grev. cryp. 144 |
| "«Suercinum Pers. | Oak | scattered | 0 aut Bk dead oak leaves | Grev. cryp. 77 |

27744 stamen <i>Tode</i>	hard	JL	aut	Br	dead leaves, stalks, &c. Grev. cryp. 144
2 Arassica <i>Boi.</i>	liard	°	0 winter	Br	dead cabbage leaves Sow. 393.
27745 <i>A/uscurum Pers.</i>	irregular	4	spring	O.Y	stems of mosses Grev. cryp. 101
27746 <i>Fungbrum Pers.</i>	roundish	0	aut	Br	gills of dead agarics Sow. 287
27747 <i>frictuum Grev.</i>	confluent	A.	aut	Bk	rotten fruit
27748 <i>bullatum Dec.</i>	blistered	confluent	0	aut	rotten gourds
27749 <i>difurum Pers.</i>	hard	oblong	0	winter	dead herb stems Grev. cryp. 1
27750 <i>paphlinum Pers.</i>	Poplar	patches	0	autw.	poplar leaves
27751 <i><alicinum Dec.</i>	Willow	patches	0	autw.	willow leaves
27752 <i>herbarum Fries</i>	herbs	cunate	0	aut	herb leaves

Division IV. *PerisjMu*

3226. ERY'SIPHE Dee.	MILDEW.	(Greek name of mildew.)	9.-24.
27753 <i>fuliginea Lk.</i>	dark	patches	0 aut D
27754 <i>communia Lk.</i>	common	powdery	0 aut Wsh
1 <i>liCuminosXrtn J/c.</i>	Legum.	powdery	0 aut Wsh
2 <i>Umbelliferarum Lk.</i>	Umbellif.	powdery	0 aut Wsh
3 <i>fichoraccarum Lk.</i>	Cichoric.	powdery	0 aut Wsh
4 <i>/fanunculaceJrum Lk.</i>	Ranunc.	powdery	0 aut Wsh
5 <i>Polygeanearum Lk.</i>	Polygfnics	powdery	0 aut Wsh
27755 <i>lamprocarpa Lk.</i>	shining-fid	patches	0 aut Blsh
27756 <i>depressa Lk.</i>	depressed	wide-spreading	0 su.aut Wsh
1 <i>Bardanas Lk.</i>	Jardana	wide-spreading	0 su.aut Wsh
2 <i>Artemauc Lk.</i>	Wormwood	wide-spreading	0 su.aut Wsh
adonca Lk.	hooked	patches	0 aut W
1 P6puli Lk.	Poplar	patches	0 aut W
2 Prunastri Lk.	Prunaster	patches	0 aut W
27758 <i>bicornis Lk.</i>	forked	powdery	0 aut Wsh
27759 <i>divaricata Lk.</i>	divaricated	powdery	0 aut Wsh
27760 <i>penicillata Lk.</i>	penicillate	powdery	0 su.aut Wsh
1 <i>Grossufrice Lk.</i>	Gooseberry	powdery	0 su.aut Wsh
2 <i>BerUridis Lk.</i>	Barberry	powdery	0 su.aut Wsh
27761 <i>guttata Lk.</i>	dropped	distinct	0 aut Rsh
1 <i>Coryli Lk.</i>	Hazel	distinct	0 aut Rsh
2 <i>Ulxbxum Lk.</i>	Elm	distinct	0 aut Rsh

S227. LASIO'BOTRYS Kunze LASIOPBOTBY& (*Lottos*, woolly, *botrys*, a bunch of grapes.) 1.
87762 Lonicene Kunze Woodbine spots 0 flu Bk live honeysuckle lvs Grev. cryp. 191

TRIBE IV. CONIOMYCE^TES.

Class I. TUBERCULA'RINI. — Division I. *Cephalohichei.*

3228. ISA^RI A Pers.	ISAHIA.	(/SOT, equal; flamenU.)	1.—?
27763 <i>microscopica Grev.</i>	microscopic	vermin.	0 spring W Trichia clavata Grev. cryp. 3
3229. A'NTHIN A Fries ANTHINA.	(<i>Anthinus</i> , florid; color.)	I.—?	
27764 <i>miniata Grev.</i>	scarlet	gregarious	1 aut O.R on dead leaves Purt. fl.3.18. <i>Clavelina</i>
3230. CEPH ALOTRICHUM Lk.	CEPHALOTRICHUM.	(<i>Kephale</i> , the head, <i>thrix</i> , hair.)	1.—?
27765 <i>nānum Lk.</i>	dwarf	gregarious	0 aut Bk dead branches, &c.
3231. AMPHICHO'RDA Fries AMPHICHORDA.	(<i>Amphi</i> , on either side, <i>chorda</i> , a cord.)	1.—?	
27766 <i>farindcea Grev.</i>	powdery	long	0 aut W rotten trunks under ground

Division II. *Scoriadei.*

3232. CER ATIUM Alb.	CERATIUM.	(Iteration, a little horn; form of plants.)	1.—?
27767 <i>Aydholdes Alb.</i>	Hydnnum-lk	fugacious	& aut W dead wood Grev. cryp. 169

Division III. *DermosporeL*

5223. JEGERITA Pers.	JEaEurrA.	(<i>Aigeiros</i> , a poplar; first discovered on it.)	2.-8.
27768 <i>cindida Pers.</i>	white	punctiform	0 aut W dead wood Grev. cryp. 268
27769 <i>set?JaCarm.</i>	bristly	minute	0 aut W dead wood Grev. cryp. 2»

Division IV. *Tuberculated*

3234. TUBERCULANRIA Tode TUBERCULARIA.	(<i>Tuberculum</i> , a pimple; appearance of irritants.)	4.—13.
27770 <i>vulgāris Pers.</i>	common	gregarious ^year Dp.R decayed sticks Sow. &H
Clavaria coccinea Sow.		
27771 <i>cōfluens Pers.</i>	confluent	gregarious 0 autsp F.R dead sycamore branches
27772 <i>granulata Pers.</i>	granulated	scattered ^aut Br dead branches Grev. cryp. 187
27773 <i>discoldea Pers.</i>	disk-like	crowded aut R sycamore branches

3235. FUSA'RIMUM Lk.	FUSARHJM.	(<i>FUSUS</i> , a spindle, shape of sporules.)	I.—?
27774 <i>tremolloides Grtfw.</i>	TremeUa-lk	very smau	0 spring Pksh dead nettle steins Grev. cryp. 10

Class II. MUCOBIVNL. — Division I. *Stilbinū*

3236. STILBUM Tode	STILBUM.	(<i>SHlbos</i> , shining; plants at first watery and gelatinous)	3.-?
Z75 Tode	erymjn.	0 aut Wsh decayed stems	Tode fun. 2.16
*76 cftrinum Pers.	lemon-cld	very min.	0 aut Y decayed stems
*77 tomentbum Schr.	tomentose	verymin.	0 aut W Trfchianlents Grev. cryp. 281

 *Tode* ASCOPHORA crowded 0 year Gl putref. paste, &c. Grev. cryp. 269

Division II. *Mucbrei.*

oij ³⁸ . AftTCOR Pen.	Mucon.	(<i>Muceo</i> , to be musty t. on musty bread, vegetables, Sec.)	2.—34.
*7179 stercdreas Grev.	dung	watery 2 winter W dung	
*o^A^ HydrAphora stercbrea Tode			

*780 canlinus Pers. dog minute } year Y dog's dung Grev. cryp. 305

- 3239. THAMNI'DIUM Lk. THAMNIDIUM. (7% fl>m<, atwig: appearance of plant under microscope.) 1.
*781 elegans Lk. elegant whorled & aut Pale putrid substances Ncespilz 75

- 3240. EUROPIUM Lft. EUROTIUM. (J?I>W was the Greek name of a sort of mouldiness.) 2.—
SZ2* herbari&rhum L*. herbarium punctiform O year Y dried plants Grev. cryp. 164
^83 ifasárum Grev. Roses patches 0 su W rosebushes Grev. cryp. 164

Division III. *Acrembnii.*

erSi¹ STiCHYLI'DIUM Lk. STAOIYLIDUM. (*Stachys*, spike, *eidos*, similar; disposition of sporules.) 1.—9.
*/84 terrestre Lk. ground spreading & aut W dead wood Grev. cryp. 257

ACREMCTNIUM Lk. ACRBMONIUM. (*Akrcmon*, a branch; thece clustered about the plants.) 2.
ecum Schmidt brown patches 0 aut OLG dead sticks Grev. cryp. 124
verticillatum Lk. whorled creeping 0 aut W dead trunks Grev. cryp. 124

Class III. MUCE'DINES.—Division I. *Trichoth&cii.*

tJS!& CHLORFDIUM Lk. CILOBIDIUM. (*Chloros*, pallid, *eidos*, appearance.) 1.—2.
*7 griseum Ehren. grey very min. 0 year Gy rotten wood Grev. cryp. 236. *Chetop.*

Division II. *Botrytidei.*

*- CORFMIUM Lk. COREMIUM. (*Korema*, filth; found there.) 1.—2.
glaucum Lk. glaucous powdery ft year Gl old paste, &c. Grev. cryp. 301. *Floccdr.*

PENICI'LLIUM Lk. PENICILLIUM. (*Peniculum*, a painter's pencil; form of filaments.) a — 4.
«Pfirsum Grev. scattered broad lines & aut W rotten herbac. stems Grev. cryp. 58.2
glaucum Lk. glaucous tufts A year Gl rotten substances Grev. cryp. 58.1

ASPERGI'Llus Mick. MOULDINESS. (*Aspergillum*, a brush for lustration j resemblance.) 4.—9.

Kladcus Lk. blue patches 0 year B rotten substances Mag. ber. 3.1.23
mausillk. woolly patches tW aut Wsh putrid fungi

vircns Lk. green broad spots% aut Gsh putrid fungi [barium

penicillatus Grev. pencilled spots 0 year D.Gy damp specimens in Her. Grev. cryp. 22

- 3247. BCTRYS TIS Mich. BOTRYTIS. (*Botrys*, a bunch of grapes; clusters of seeds.) la — 25.

U75 diffTisa Alb diffuse broad tufts £ aut W rotten herbac. stems Grev. cryp. 126

U'S ogaricina Lk. Mushroom wool coat 0 aut W decaying fungi Grev. cryp. KG

effusa Grev. ertUsed spots 0 aut P.Gy underside of live leaves

Par'a*itica Pers. parasitic lax 0 sp.su W on shepherd's purse Sow. 359

m graZ*. black very powdery J aut Bk rotten trunks Grev. cryp. 274

ensa Dil. dense roundish 0 aut W on mosses Stur. deu. 52

Pol^spora LA. many-seeded broad 0 aut W dead branches Stur. deu. 35

aJES ma « "spora Lk. long-seeded spreading 0 aut Ro.R dead leaves Stur. deu. 50

278S murtl>a -W*. mouse-crf tufts 0 aut Gy rotten branches Stur. deu. 36

•u* Ieucospora Lk. pale.seeded woolly tufts 0 year Gy rotten pears

ACROSPORIUM Nees ACROS. (*Akros*, summit, *spora*, sporule; upon summit of filaments) 2.—9

^mildoles Nees Monilia-like spots 0 sp. spoutVW Idæss off grasses Grev. cryp. 75*

> fasciculatum Grev. fascicled patches 0 spring Gl rotten oranges

Division III. *TrichSmyci.*

(Spora, a sporule, *thrix*, hair.) 9. —49.

8761 f*. brown spreading 0 aut Fer rotten trunks

278n TS * 0 ^ n 1 **. dun K tufts o y ear o damp cellars

878U S- toB m Lk. cavern broad \ year W rocks

golden spreading 0 year O.Y rotten trunks Bui. her. 504. 5

Window spots 0 year W damp glass Necspilce47

hita. U m Lk. f*. TRICHOTHECHIUM. (*Thrix*, hair, *theké*, *thecium*.) 2L —?

27817 ? m Lk. rot J tuftB o aut w w rotten wood Grev. cryp. 172

"eterosporum 5pvariouB-seeded spots 0 aut Br dead caps. Gentidna campestris

S7818 L?PE D. O'NIUM Lk. SEPEDONIUM. (*Sepedon*, putrescence; growing on putrid substances.) 1.
TOycophilum Lk. Fungus-loving cloudy 0 aut O dying fungi Grev. cryp. 198

Sfctal LT^D IUM ^k. FUSIDIUM. (*Fusus*, a spindle; fusiform shape of sporidia.) 2.—7.

27820 V^ren Grev. grey effused spo. 0 aut Wsh dead beech leaves Grev. cryp. 102

27821 vi DX. yellow-green effused spo. 0 aut Ysh dead beech leaves Grev. cryp. 102

*' candidum Lk. white effused spa 0 aut W dead oak leaves

Division IV. *SporSmyci.*

YTHRIINCTUM Kunxe POLVTHRINCUM. (*Polya*, many, *thrigkos*, a little division) 1.
Trefoil punctifonn 0 aut Bk clover leaves Grev. cryp. 216

3254. DEM A^vriUM Pen. DEMATIUM. (Diminutive of *Aetna*, a bundle; mode of growth.) 1.—
87823 articulatum Pert. articulated minute ^{n. * u.} Bks^h stems nf hnrha lwa *^{* k*}. ■

Class IV. BVSSA'CEJB. — Division I. Rhizoinorphace.

3255. RHIZOMO'RPHA Jto/A RHIZOMORPHA. (*Rhixa*, a root, *morphe*, form; resemblance.) 4. — ?
87824 subcorticalis Per*, subcortical net-like 72 year Br beneath bark Sow. 392. 1 & 2. pit/ens
87825 divulgans Grev. diverging creeping 24 aut Rah beneath bark Grev. cryp. 154
27826 subterranea Pen. subterranc. filament. 24 year Bk mines
27827 medullaris Sm. medullary much bran. 144 year W cellars Lin. tr. 12. 80

3356. OZO^NIUM Lk. OZONIUM. (OXOS, a branch; filaments branched.) 1 — ?
S7828 auricomum Lk. yellow-haired byssoid 3 aut O rotting wood Grev. cryp. 260

Division II. Racodiucece.

3257. R ACO*DIUM Pert. RACODIUM. (Rakos, a torn garment; appearance of plants.) 1. — ?
87829 cellare Pen. iw/u?-cellar shaggy 3 year Sooty cellars Grev. cryp. 259
Fibularia vinaria Sow.

3258. HELMISPO^XRIUM Lk. HELMISPORIUM. (Helmins, a worm, *spora*, sporule.) 2. — 8.
27830 macrocifpon Grev. long-fruited broad 0 year Bk dead branches Grev. cryp. 148
27831 velutinum Lk. velvet spreading 0 year Bk dead wood Grev. cryp. 148

3259. CLADOSPCPRIUM Lk. CLADOSPORIUM. (Klados, a branch, *spora*, a sporule; attached to.) 2.—&
87832 herbárum Lk. herbs very min. 0 su aut Ol G. dead stems Nees pilze 5. 64
87833 velutinum Grev. velvety patches 0 spring G.Bk rotten wood

3260. HELICOSPCPRIUM Nees. HELICOSPORIUM. (Helikos, twisted, *spora*, sporule; sporules curved.) 1.
27834 vegetum Nees vigorous cloud-like 0 o Gy foot of trees Nees pilze 66

Division III. Byssece.

3261. MONI'LIA Pm. MONILIA. (Mordle, a necklace; filaments articulated.) 1. — 12.
87835 antennata Pert. antenæ-lik tufts 0 aut Bk dead wood Grev. cryp. 855

3262. TCRUL A Lk. TORULA. (Diminutive of *torus*, a twisted cord; appearance of filaments.) 1. —
87836 herbárum Lk. herbs fragile 0 aut Bk dead stems

Class V. PHYLLERIA % C.E.S.

3263. ERI'NEUM Pers. ERINEUM. (Erinos, a hedgehog, erinaceous appearance.) 19.—40.
27837 aureum Pers. golden velvety spots 0 su Bt.Y Ivs P6pulua nlgra Grev. cryp. 33
87838 griseum Pers. ■ velvety spots 0 sp.su DI.P under oak leaves Grev. cryp. 141
87839 ocerinum Pers. Maple depres. tufts 0 sp.aut R.Br und. sycamore Ivs Ed. ph. jour. 6. 2.1 & ■
87840 joyrinum Pen. Pear tree depres. tufts 0 aut R.Br on crab tree lvs Grev. cryp. 22
87841 tortubsum Grev. tortuous depres. tufts 0 sp.su Wsh on birch leaves Grev. cryp. 94
27842 Jugländia Dec. Walnut depres. tufts 0 su Pa under walnut Ivs Grev. cryp. 263
87843 clandestinum Grev. concealed depres. tufts 0 su W.Pk und hawthorn Ivs Grev. cryp. 141
87844 rbeum Schultx rosy depres. tufts 0 su C on birch leaves Grev. cryp. 21
27845 Aetulinum Reb. Birch depres. tufts 0 sp.su Wsh on birch leaves Ed. ph. jour. 6.3.16
87846 purpur&scens Gae. purpurascen. depres. tufts 0 su P.Br maple leaves
27847 Stirbi Funk Mountain Ash depres. tufts 0 su Rsh.B mountain ash Grev. cryp. 283
87848 Pkdi Duvai Bird-Cherry depres. tufts 0 su D.Br on Primum Padus Grev. cryp. 157
87849 ilneum Pers. Alder depres. tufts 0 su O.Br alder leaves Grev. cryp. 157
27850/agtneum Pers. Beech depres. tufts 0 su Pk.Br beech leaves Grev. cryp. 250
27851 /mutulinum Pers. Pclar depres. tufts 0 su D.Br. on trembling poplar Grev. cryp. 350
87852 /flaceum Pers. Linden depres. tufts 0 su Pa lime tree leaves
87853 lanbum Grev. woolly hidden tufts 0 su Pa margin of pear lvs
87854 FUis Dec. Vine woolly tufts 0 su Rsh.Br vine leaves
87855 Pseido-platani Pers. Sycamore depres. tufts 0 su Rsh.B sycamore leaves

Class VI. EUTO'FBYTI. — Division I. Sporodtsmii.

3264. CORYNETJM Nees COBYNEUM. (Koryne, a club; form.) 1.—7.
87856 pulvinatum Nees cushioned minute 0 aut Bk dead branches Nees pilze 31

3265. SPORIDE'RMIULJt. SPORIDERMIUM. (Spora, a sporule, *derma*, a skin; situation.) !• rj? 87857 atrum Lk. dark parasitic 0 aut Bk on species of Thclephora Grev. cryp. »*.

3266. EXOSPO'RIUM Lk. EXOSPORIUM. (Ego, outside, *spora*, sporule.) 1. — ■
27858 niße Lk. Linden punctiform 0 year Bk linden branches Grev. cryp. SOS

Division II. Stilbosbrei.

3267. ASTROSPO^RIUM Kunxe ASTROSPORIUM. (Aster, a star, *spora*, sporule; form.) *.
87859 Hofmanni Kunze Hoffmann's minute 0 year Bk dead beech branches

3268. STILBO'SPORA Hofin. STILBOSPORA. (Stilbo, to shine, *spora*, a sporule.) 3. — ■
87860 teflinia Pers. Birch raised 0 year Bk dead birch trunks Grev.cryp.273.-D<W⁷¹,
87861 bilobulata Grev. two-celled emerging 0 year B dead furze bran,
87862 profusa Grev. profuse spots 0 year Bk dead sycamore bra. Grev. cryp. 912 ■

3269. MELANCO^NIUM Lk. MELANCONIUM. (Melas, black, konis, dust; appearance.) a o
87863 ovatum Lk. ovate minute spots 0 year Bk dead branches Grev. cryp. ■■■
87864 conglomeratum Lk. conglomerate spots 0 year Bk dead branches Lk. obs. 1.1.7

Division III. Namasp&reL

3270. NJEMA'SPORA Pers. NJSMASPORA. (Kema, a thread, *spora*, a sporule; resemblance.) & " ?
■>Ufirpi Sow. Hornbeam irregular 0 year Bk dead hornbeam branches Sow. >"
IKRHI ■■■. filamentous tortuous 0 year O dead branches
■■■. uttivon gregarious 0 year O dead beech branches

Division IV. Hypod&rmn.

JEL ¹	PUCCPNM Mich.	PUCCIHIA.	(T. <i>Puccinius</i> , a professor of anatomy in Florence.)	'21. — 50.
7868 ^sparagi Dec.	Asparagus	two-celled	0 aut	Bk dead asparagus
SSSJ Circæv Pers.	Circæa	two-celled	0 aut	Pk.Br under urc&i lvs
S.S.? Chr y<oiplenii Grev.	Chrysosple.	two-celled	0 my	Pa.Br under Chrys. opp. lvs
A <i>icularia</i> Dec.	Knot-grass	two-celled	0 aut	Bk under Polygonum lvs
Straus ^gopodium Grev.	tumid	two-celled	0 aut	B.Gr on /Egopodium lvs
— Pers.	Mint	two-celled	0 spring	Br.Gr on Zfuniq Bulbocastanum
— Pers.	Polygonum	two-celled	0 aut	Bk under 3/cntha lvs
Centaureæ Dec.	Centaury	two-celled	0 aut	IIBr under Po)fg. amphfb. lvs
Umbelliferarum Dec. Umbellif.		two-celled	0 aut	Bk on Centaurtangra
aniculaj Grev.	Sanicle	two-celled	0 aut	Ve.D on Umbellifera; lvs
S2Z? variabilis Grev.	variable	two-celled	0 aut	Bk Br under Sanicula lvs
i/eraclei Grev.	CowParsnep	two-celled	0 su	Bksh on Leont. Taraxacum Grev. cryp. 75
Epilobii Dec.	Epilobium	two-celled	0 jn	Bksh under Spindlylum lvs Grev. cryp. 42
— cae Dec.	Betony	two-celled	0 aut	Br under Epilobium palustre lvs
pulverulenta Grev.	powdery	two-celled	0 su	Fer under ifcton. offic. lvs
* Adoxas Dec.	Moschatel	two-celled	0 su	D.Br under Epilobium lvs
— G	Primrose	two-celled	0 su	D.Br on Adoxa Moschata Uina
/oob Smis Dec.	Violet	two-celled	0 su	D.Br under primrose lvs
887 Valanti* Pers.	Cross, wort	two-celled	0 su	D.Br under /tola canina lvs
	heaped	two-celled	0 sp.su	Dp.Br on Galium verum
	Ulmaria	two-celled	0 aut	Dp.Br under Senecio JacobaAi lvs
	Grass	two-celled	0 aut	P.Bk under Ulmaria lvs
	globose	one-celled	0 aut	en tifex leaves
	Box	two-celled	0 su	Bk on corn & grasses Sow. 140. U./<—
	Anemone	two-celled	0 su	Bk on bean leaves Grev. cryp. 29
				leaves of box Grev. cryp. 17
				on Jnemonc leaves Sow. 43. —

3272. PHR AGMT'DIUM Lk. PIIRAGMIDIT'M. (Phragma, a division or partition, e'tdos, similar.) 4.-5.				
895 Aosae Grev.	Rose	many-cell.	0 aut	Bk under rose leaves Grev. cryp. 15
896 flubi Dec.'	Bramble	many-cell.	0 aut	Bk under bramble lvs
fracile Grev.	slender	many-cell.	0 aut	Bk under raspberry lvs
Potentilla Pers.	Potentilla	many-cell.	0 aut	Bk under Potentilla lvs Grev. cryp. 57

			Uikion, a wheal, cidos, like; pustules.)	26. — G*.
	Pine	scattered	4 su	Pa.0 on Ptaussylvestris Grev. cryp. 7
	Epilobium	beautiful	0 su	W on Epilobium montanum lvs
	Violet	crowded	0 su	Wsh under flota cartna lvs
	albescens	beautiful	0 ap	W Ad6xa Moschatellina
	Dandelion	spreading	0 su	under Leontodon Taraxacum lvs
	Woodbine	large spot	0 su	under woodbine lvs
	Ground Nut	deformed	0 spring	Ysh on 2?unum leaves
	Ragwort	agglomerat.O	0 su	Ysh under Sen. Jacoba?z lvs
	Prenanthes	spots	0 su	W under, //icrac. sylvat. lvs
	Nettle	clusters	0 su	Ysh on ITrfica dioica
	dense	clusters	0 su	Wsh on Ficaria
	Gooseberry	bright sp.	0 su	R under gooseberry lvs Grev. cryp. 62
	Crowfoot	clusters	0 bu	Wsh under 7/lmine, lingua lvs
	Caltha	bright	0 spring	O under Caltha palustris lvs
	Berberry	spreading	0 su	Q on Bctb. vulgaris Grev. cryp. 97
	ragged	swollen	0 su	Br on hawthorn Grev. cryp. 209
	cornut	spots	1/6u	Y.Br on mountain ash Grev. cryp. 180
	Coltsfoot	concentrical	0 su	Psh under Farfara lvs Sow. 397. 1
	reddish	concentrical	0 su	C under Atimex lvs Sow. 405 Rhinitis
	Garlic	concentrical	0 su	Y under jfl. urs'inum lvs
	Thalictrum	clustered	0 su	O Thalictrum alpinum Grev. cryp. 4
	Buck-thorn	clustered	0 su	Hah. Y buckthorn leaves Pers. oba. 1.2.4
	Colocynthrum Dec.	pale-seeded	0 scattered	W anemone leaves
	PrSmuls Dec.	Primrose	0 scattered	Pa.Y primrose leaves
	Mint Yue Dec.	Mint	0 deformed	O. Y mint stalks & leaves
	liminosarum Grev.	Lcguin.	0 scattered	Wsh leaves of Leguminosse

3274. URVSDOPers.				40. — 105.
7925 Geranii Dec.	Geranium	scattered	0 su	D.Br on Geranium lvs Grev. cryp. 8
5J26 Ficaria! Alb.	Pile-wort	pulverul.	0 su.aut	D Br under Ficaria lvs
VSM suav^olens Pers.	odoriferous	fragrant	0 sp.su	P.Br Cricus arvensis lvs
ow oligognathum Dec.	Polygonum	spreading	0 su aut	Pa.Br under 7'olygonum lvs Grev. cryp. 80
9 Pmliu Dec.	Primrose	scattered	0 su	Pa Br under primrose lvs
	Cichoraceæ	spots	0 su	D.Br on Composite leaves
	CowParsnep	patches	0 su	Pa.Br under Sphondylium lvs
	two-fronted	spots	0 su	aut Pa.Br both sides of sorrel lvs
	Dock	small	0 su	Br on Abmx leaves
	Bean	patches	0 su	Pa.Br on bean leaves Grev. cryp. 95
	Labiate	pustular	0 su	Y.Br on mint leaves
	intruded	scattered	0 su	aut R.Br on Liizula leaves Grev. cryp. 12
	oblong	pustular	0 su	under <i>Alchnilla</i> leaves
	Willow	mottled	0 su	under Liizula leaves Grev. cryp. 12
	GoldenOsier	pimpled	0 su	under <i>Salix vitellina</i> lvs
	mealy	mealy	0 su	under Salix caprea lvs
	Colt's-foot	gyrose	0 su	under colufoot lvs
	Ragwort	blotches	0 su	under Senecio lvs
	confluent	gyrose	0 su	Y on il/ercuriati* lvs
	Cinquefoil	powdery	0 su	Go.Y on Fragaria sterilis lvs
	Rose	mottled	0 su	under rose leaves
	Bramble	very powd.	0 su	under bramble lvs
	effused	spreading	0 su	under itosaceac lvs Grev. cryp. 19
	circled	gyrose	0 su	sp.PU Y on raspberry leaves
7Q2! St Aewif//* Pers.	Lady's Mantle	spreading	0 su.aut	under Alchemilla lvs
27QVT- ² hilanthrum Dec.	Rhinanth.	spots	0 su	Scrophularines
279M c ⁱⁿ ^ c -	Flax	shining	0 su	Lnum catharticum Grev. crvn 31
27QM 5 ^{ax} fragarum Dec.	Saxifrage	brilliant	0 su	saxifrage
3 Campnulosa Pers.	Campanula	bright	0 su	under Campanula lvs

£7954 Pfroia Grev.	Winter Green minute	0	su	Go	under Pyroia lvs
27955 //eliosuipis Dec.	Euphorbia round	0	aut	Go	under Euphorbia lvs
87956 linearis Pers.	linear, very com.	0	sp.aut	Y	on grass leaves Sow. 140. frumnti
S7957 JBCidiiformls Grev.	Efcidium-lk	pustular	0	spring	Y on Sphond'lium lvs
27958 Cerastii Grev.	Cerastium punctiform	0	su	Go	on Ccrast. visebsum lvs
£7939 pustulata Pers.	pimpled	punctiform	0	spring	Y on Epilobiun palustrc lvs
£7960 Sbuchi Pers.	Sow Thistle	spreading	0	BU	R.O under Sunch. olerac. lvs
S7961 PetasUes Dec.	Petasites gyroae	0	aut	O	under Petasites lvs
£7962 populin Pers.	Poplar	beautiful	0	aut	Go under Populus nigra lvs. An. wett 2.11.5
£7963 ovata Straus	ovate	spots	0	aut	Taw on Popus tr^mula lvs. An. wett 2.11.6
£7964 Candida Pers.	white	spreading	0	aut	Cruciteras . Grev. cryp. £51

3275. I7STIL ANGOLA. UBTLAGO.	(Ustus, scorched; appearance.)	5 . -■
£7965 Sdgetum Pers.	corn	spreading 0 su Bk within grains of corn
£7966 tfrccolarum Dec.	Pellitory	spreading 0 su Bk on fructif.of Crrex
£7967 caries Dec.	canker	destroying 0 aut Bk.Br within grains of wheat Stur. deu. 34
£7968 antherarum Dec.	flowers	spreading ✓ su P on Caryophyllea ²
£7969 flosculbsa Dec.	floret	spreading ✓ su P.Br on Scabiosa arvensis Sow. 396.2. Scabibta:

3276. CYLINDROSPOTUUM CYLINDROSPOBIBL (Kylindros, cylinder, spora, sporule; form.) 1.
 £7970 concentricum Grev. concentric speck.lk.hea. 0 my.jn w cabbage leaves Grev. cryp. 20

SUPPLEMENT.

To save unnecessary reference from the body of the work to this Supplement, and at the same time to indicate when a reference to it is necessary, an asterisk (*) is placed in the body of the work before such of the genera as are divided or repeated here, and also in those places where new genera should have come in. Those genera or species in this Supplement, which are merely alterations of the names, or distribution in the Natural System, of genera or species in the body of the work, are distinguished by a section (\$). The same mark, in the body of the work, indicates that the name, or its place in the Natural System, is altered in the Supplement

CLASS I. ORDER I.

MONANDRIA MONOGY'NIA.

S | £ 1 * 1 .
I I I U H

 S7 ^{37a}	^L CA'NNA. 5a discolor Lindl.	various-clad.-/rfjg ED or 10 n S Trinidad 1827. R Fam Dot. reg. 121a
 51a	grandifolia Lindl macricta Lindl	great-leaved £ ES or 2 year Y Rio Jan. 1826. D 81 Bot ree 1210 meagre £ BS or 2 year W li St Rio Jan. 1826. D r.l Bot cab. 1781.
 .5.	PHRY'NIUM.	
 59a	setosum Rose.	bristly £ E2 or 2 my.jl P Rio Jan. 1824. R s.1 Rose, scit icon.
 9	HELLENM. i _{a2fl} abn ^r raisZihf _{f1} :	irregular jg E3 or 10 s W China 1824. D s.1
 H7a	Roscoea ⁿ allMe.	Roscoe's £ E] or * o W E. Indies 1827. R s.1 Bot reg. 1212

CLASS I. ORDER 2.

MONA'NDRIA DIGY'NIA.

 27.	CORISPE'RMUM. 239a sabulosum Led.	sandy O cu 1 jl.au Ap Siberia 182L S co
---	--------------------------------------	---

CLASS II. ORDER 1.

DIA'NDRIA MONOGY'NIA.

 27 ^{37a}	MILLINOTO^N/il Rox. MILLINGTONIA. (Sir T. MMingtm, Szy. Prof. Oxford.) Oltinat. 1.—5. - simplicifolia Rox. simple-leaved f CD or 20 ... Y E. Indies 1828 C p.1
 . ^IGUSTRUM.	spic&fum D. Don spike-flowered A or 8 jn.jl W Nepal 182a C co Bot mag. 21)21
 J** ³	nepalnse Wai
 304a	«ASMrNUM. lanceolarwf J. Q or 10 ... W Silhet 182& C s.p
 87982	S 2 ^o ca, wtachya Led. prettiest-spiked^ A or 2 ln.il B Podolia 1829. D co
 JJ ¹¹	stolonifera Ish. * A or 1 Jn.jl B « D co
 87983	dianthifolia Cun. Pink-leaved £ IAJ or my.jn B N. Holl 182S. D p.l
87984	«4a?oiif r?rX ⁴ * ² poUshed i O w I mrjl B Britain dtgar. S co Eng.bot783
87984	ii ⁵⁸ - JUST1'C7.4. ventricose * Q or S Jajl W.a China 182& C p.l Bot mag. 2766
 S?"	ve«ric6sa Wai oblongata L*.
	H h 2

59. DICLI'PTERA.
 27987 558a spintoa *Lod.* spiny 2 apjl Y Mauritius 182*. C p.l Bot.cab.1244
- *65. CALCEOLARIA
 27988 577 integrif&lia.
 2 angustifolia *Lindl.* narrow-lvd n. or 2 au.s Y Chile 1822. C Lp Bot. reg. 1083
 27989 577a asclndens B. R. ascending or 1 jl.s Y Cordiller. 1826. C Lp Bot. reg. M*
 27990 - thysifl&ra *Grah.* thyrse-flowered a. or 1 jijas Y Chile 1827. C p.l Botmag.29#
 27991 - polif 61ia *Hook.* Poly-leaved f iAI cu 1 jl. Y Chile 1826. D Lp Bot.mag.28g
 27992 - connate Hook. connate-leaved f OJor 3 jl.au Y Chile 1824. D l.p Bot. mag. 2887
 floribunda *Lindl.* Bot reg. 1214. paniculate Herb.
 27993 578a purpura *Grah.* purple/*floM/ma*"f LAJ or 1 jl.s P Chile 1827. D Lp Bot mag. 2775
 27994 - arachnoidea *Grah.* cobweb-like f iAI or 1 jn.s P Chile 1827. D Lp Botmag.287⁴
 tinctebria *Gill.*
 27995 579a plantaginea *Sm.* Plantain-fcawdjc iAJ or 1 au Y Chile 1827. D Lp Bot mag.
74. MONA'RD^.
 27996 621a mcnthatfblia *Grah.* Mint-leaved ^ 3 jn.s Li N. Amer. 1827. D co Bot mag. i.
3278. 74a. BLEPHI'LIA *Raf.* BLEPHILIA. (*Blep harts*, eye-lash j sepals fringed?) *Labiates.* 2.—3-
 L - " ciliata Aq*. ciliated 3t A < 3 jni R N. Amer T ... D co
 Mondrda ciliate X. Na 630. Mirsute No. 632
- *76. SA'LVIA.
 27997 686a involucrata *Cav.* involucred 14 jls R Mexico 1824. C p.l Bot. mag. 287\$
 27998 717a desertbrum *Sthan.* desert A or 2*o B Siberia 1829. S co
 27999 721a Simstfna B. R. Sims's A or 3 jn.jl Pa,B Russia 1820. D p.l Bot reg. 1003
 bracteate *Bot. Mag.* 2320.
3279. 78a. STREPTOCA'RPUS *Lindl.* STREPTOCARPUS. (*Streptos*, twisted, *kagoo*s, fruit) *Bignoni&cea?*. \) . . *Röü Lindl.* Rex's f E 2 p r i year B C. 6. H. ... S p.l Bot.reg.1173
Didymocarpus Rtex Bow. Na 15434.
82. AC^NA.
 £8000 775a myriophilla *Lindl.* many-leaved A cu 1 myjn G Mendoza 1828. D co
3280. 82a. MARGYRICA'RPUS R.\$P. MARGYRI. (*Margaron*, pearl, *Jcarpos*, *frmt*; resemblance.) *Ro3(ict*-)*
 28001 - *BetuuBRSP.* bristly I Q c u 2 ... G Peru 1829. C s.p FLper. 1.8.*>
- *87. PIMELETA.
 £8002 805a davate *Lab.* clubbed 3 ap.my W N. Holl 1824. C sp
 28003 796a humilis *R. Br.* humble Jor 1 my.jn W N. Holl 1824. C s.p Bot reg. 1

CLASS III. ORDER 1.

TRIA'NDRIA MONOGYNIA.

96. VALERIAN&LLA.
 \$8004 922a cong&ta *Lindl.* crowded-Jhwered O or 1 Jl R Columbia 1826. S co Bot reg. 109*
- *116. CRO^CUS.
 £8005 1014a argenteus *Sab.* silvery tf A or *f.mr W.Br -- -- O co
 28006 1019a lacteus *Sab.* milky A or i f.mr Pa,Y -- -- O co
- *3281. 117a. STREPTANTHE'RA *Suit.* STRBPTANTHERA. (*Streptos*, twisted, *anthera*, anther.) *Iridceo.* 1-^
 28007 - Regans Suit. elegant tf^Jpr I my.jn W.B.Y C. G. H. 1827. O s.p.lSw.fl.gar.8*
127. MELASPHJE'RULA.
 28008 1155a parviflora *Lod.* small-flowered tf iAI or ft ap.au Y C. G. H. ... O s.p.lBotcab.443
3282. 141a. LEUCOCO'RYNE *Lindl.* LEUCO. (*Leukas*, white, *koryne*, a club; sterile anthers.)
 28009 - odorata *Lindl.* sweet-scented X Ai or 1 au W Valparaiso 1826. O pl Bot reg.
 Na 1228. and 1230. of Brodtuez belong to this genus.
3283. 1416. TRITELEPA *Lindl.* TRITELEIA. (*Treis*, three: stigmas.) Asphodelaea?. 1.—3.
 28010 - grandiflora *Lindl.* great-flowered tf or 2 jl.au W? N. Amer. 1826. O pl
- *148. TRIS.
 28011 1303a ténax *Don.* tough f A or 1 ap.my P California 1826. D p.l Botreg.1^{10J}
 28012 1313f dandestina *Le* clandestine f ftClor Brazil 182P. O s.l.p
 28013 1255a tafrica *Lod.* Taurian ^ A or f ap.my Y Tauria 1827. O co Bot cab. I^{eng}
 I^{**1}
143. MORTEM. •
 28014 1327a catenulata *Lindl.* ch&mjotted 1 my W.B Mauritius 1826. D s.p Bot reg. 107*
- *144. MA'RICA.
 28015 1339a celestis *Leh.* sky-6/<<< 1 ... B Brazil 1829. O p.l
- *145. SISYRI'NCHIUM.
 28016 1343a chilense *Hook.* Chile 1 jn.au B Chile 1826. D p.l
 28017 1352a raminif blium *Lindl.* Grass-leaved h ap.my Y Chile 1825. D fl.p
 28018 pedunculatum *Gill.* st.ked->fur 1 s.o Y Chile 1827. D s.p
 28019 1353a odoratissimum *Lindl.* sweetest-sctd 1 jn W S. Amer. 1828. D s.p Bot reg.
162. TRICHO'PHORUM.
 ■ 1433a montevideInseG.Don MontevideanJBli ES un 4 my.n Ap SL Amer. 1827. D bog
 ■ . . Eriophorum montevideense Lk, Lockharti G. Don Lockharti's JBkEJun 4 my.n Ap Trinidad 1827. D bog;

CLASS III. ORDER 2.

TRIA'NDRIA DIGY'NIA.

- ^{88^2} 1576a' etiog&nia Schr. woolly-angled M O un 1 jn.Jl Ap Germany 1827. S co
221. DESCHA'ML'SZ'F
^{88023 1} 850< paludbsa Marsh, marsh Jk A un 3 Jajl Ap Germany 1829. S co
 •dirapaludbsaiZoM
 oiS^{8"} MOL'I'NL
 •128 caeritea. .
 2 depauperata LmdL starved I A un 1 au Ap Scotland Clamo D co

CLASS III. ORDER S.

TRIA'NDRIA TRIGY'NIA.

- c** c ®°- HOLCSTEUM.
 8487a sperguloides Leh. Spurrey-like O un | jl W Europe 1829. S co
293. ERIOCAU'LON.
 2415a decangulare Mx. ten-angled * A cu 3 au W N. Jersey 1825. D bog Bot cab. 1310

CLASS IV. ORDER 1.

TETRA'NDRIA MONOGY'NIA.

- 311a. SI'MS74 R. Br. SIMSIA. (70in AAM, M.D., an English liotanist) Protehce*. L
 - anethifolia B. Br. Fennel-leaved «L J or 1} ... Y? N. Holl 1825. C p.1
-
-
- i ?^{yJl} O S i s l i t s si
- 314<. ANADETNIA/?». ANADBNIA. Cf priv., a fe[n], aglandi nectariferous wanting.) Prote&cca. I-
 - pulchella B. Br. neat * i_J or 2 ... Y N. Holl 1824. C p.1
- ^a. QUA'DR/^ R. & P. QUADRIA. (Antoinc de la Quadra, a Spanish cultivator.) Proteedcee. I
 - heterophylla A^P. variable-leaved j L j f r 40. . . . Chile 1826 C p.1 FL per. 1.99.6
 G MoL
- * 324a. EMBOTHR1UM Fora. EMBOTHR1UM [En, in, bothrion, a little pit; anthers.) Proteedcee. 1.— 2.
 - strobilnum Loft, strobile-like s< LJ or 3 ap G.Y N. Holl 1824. C s.p Lab. n. h. 265
-
 8803* 2714a' /etifolia 6<
 Cumming&mu Sieb. Ledum-leaved * LJ or 6 year Pa N. Hofc 1824. C s.p
- *331. PO'THOS.
 2781a longifotta/.*/O. long-leaved jgCScu 1 ... Ap Mexico 1829. SkLp
 - fflauca Lk. 8c. 0. glaucous ± USI cu 1 ... Ap Mexico 1829. Sk Lp
 - microphylla Hook. smalMeaved & E] cu ^s ... Ap Brazil 1829. Sk Lp Bot. mag. 2953
- 339 DORSTE^NTii
 2836a' tubicina B. % P. trumpet->ltwf ^ C S cu \ au G .Trinidad 1817. D p.1 Bot mag. 2804
- * 3*3. HOUSTO^N/iL.
 *857a serpyllifolia Mx. WildThyme-/>rff A pr ijn.a u W N. Amer. 1826. D i p Bot mag. 2828
- 346a. SUCCI'SA Vail. DEVIL'S BIT. (Succisus, lopped; appearance of roots.) Dfspdceie. 18.—24.
 - pratinensis Moen. meadow & A c* * <" v Britain pas. D w En& bot 878
 Scabibsa Succls L. No. 2887 and 2888; also those species from the beginning to Na S881 of Ccpnalana
- *?*• 3466. ASTEROCE'PHALUS Vail. ASTEROCEFHALUS. (ifefer, star, *<>*<>, head: seedfc) A ^ - ^ r^{8 0} -
 * - canteens Spr. canescens ^ A or 1 jl.au U Hungary 180i D co W.&K.53
 Scabibsa canescens Kit. Na 2906.; and from this to the end, Na 2923. excepted.
***** 346c\?TEROCE'PHALUS r<>7. PTEROCBPHALUS. (Pteron, wing, ArepAafe, head; seeds.)
 - Vaillantit Lag. Vaillant's O or 1 jl Aleppo 1822. S co
 - Nai905L and 2923.
-
 Kentucky 3t A cu J jnjl W N. Amer. 1829. S co
\$012 S S :? S £ ^ f P. connate-feawrf^nor 5 my O Peru 1826 C Lp Bot mag. 2853
⁸⁸⁰¹³ * 3226a heterophylla B.B. variable-leaved • or 10 my Y & Amer. ... S Lp Bot reg. 1259
- 3240< ffi^Gr**. imbricate-/!*! «l_Jpr 2 jajl Pk C. G. H. 1824. C p.1 Bot mag. 2809
 H h 3

28044 3278a *atalis Wai.* *teasile-flowered* • or 4 ... W E. Indies 1828. C p.1

28045 3895a" *caffra A.w.* Caffrarian « i J o r • ... W C. G. H. 1823. C p.1

3291. 388a. LIPO'STOMA *D. Don* LIPOS. (*Leipo*, to fall from, *stoma*, mouth; lid from capsule) *Rubia*¹
28046 3308a *campanulifl6raD.Do»bell-floweredl.IZ2pr_{AN}** jn.au B BrazU 1825. C Lp Bot mag.
Mghxhtia capitata Grah. *Hedyutis campanuliflyra Hook.*

397. AMMA'NN/i<
28047 3327a *Wormskioldii H.ber.* WormskioWs O u n i jl R Brazil 1827. S co

405. ALLKFN.M.
28048 3402a *ovata Ph.* ov&teJeaved O or lft jl.au F N. Amer. 1827. S 00

CLASS IV. ORDER 4.

TETRA'NDRIA TETRAGY'NIA.

•414. POTAMOGERRON.

28049 3444a *acutifblium Lk.* acute-leaved A A cu fit jn.jl GY Britain dit D co Eng. botsu.x

CLASS V. ORDER 1.

PENTA'NDRIA MONOGY'NIA.

422. ABRO'NIA.

28050 3467a *mellifera Dou.* honey-bearing-* A pr ijn.au W California 1826. D s.p. ■■■■■

423. PLUMBAGO.

28051 3469a *micrantha Led.* smalUflowcred ^ A or 3 ■ W? Siberia 1829. D co
28052 3473a *mexicana Lk. & O.* Mexican • • or 3 jl W Mexico 1829. C p.l
28053 3475a *rhombif olia Hook.* rhomb-leaved DD or ft's B S. Amer. 1826. S p.l Bot mag. 8917

426. ECHINOSPE'RMUM.

S519 *pátlum Leh.* is squarrbsum, No. 3519.

448. NOLA*NA.

28054 3748a *taitalia Lindl.* slender Jt Q or ft JL B Chile 1824. S p.1

28055 . . *grandiflbra Leh* greaUflowered Jk Q or ft jl B Chile 1829. S p.1

449. ARETI.

28056 3750a *publscens Dec.* pubescent ■ ■ ft my.jn W SwitzerL 1824. D 8p Bot cab. I&⁵

*450. ANDRO'SACR
SSUS8 3763a *carinatu Torrey* keeled ft ap.Y N. Amer. 1826. D pf
linv&ris Grah. linear-leaved ■ A ■ ft ap.my W N. Amer. 182a D l.p

•451. PRI'MULA.

28059 577a *verticillVt&Forak.y>hoT\ed.JlwltS(lydf iAl P' ft mr Y* Egypt 1826. D s.p. Botmag. 28g
\$ 3798a *glaucBcns Morel,* glaucesccut £ A or ft JIJI SwitzerL 1826. D pf Sw. ft gar. 25*

calycina Gau. Na3798.

•454. DODECATHEON.

28060 3816a *intgrifblium Mx.* entire-leaved ■ ■ or ft ap.jn LP N. Amer. 1829. D p.1 ■■■■■

*455. (TY'CLAMEN.

3820 *plrsicum*
4 *laciniatum* ■■■■■ AiAJpr \ apjn R.W S p.1 Bot reg. 1095

4flk VILLA'RSM

28061 3867a *geminata R. Br.* tmn-pcduncld &\Jfox fit jn.jl ■ N. Holl 1828. O p.l

465 CHIRC*N/i4.

28062 3872 *terpyllifbia Leh.* Wild lliyme-lvdtt. u.j or ■ ■ . CG. H. 1829. C Bp

•472. PPILOX.

28063 3905a *Sickmanni Leh.* Sickmann's ^ A or 2 au W N. Amer. 1826. D p.1 Sw.fl.gar.n.«

280H4 3907a *cordata Ell.* coxAaic-leavcd % A or 3 ... Pa.P Carolina 1827. D p.1 ■■■■■

£8065 3908a *reflexa Swt.* reflexed 5 A or 3 jls It hybrid. - I) p.l Sw!n gar.g\$

28066 . . *sclbra Su/.* scabrous ^ ^ or 3 au Li N. Amer. ... I) p.l ■■■■■

28067 . . *odorata Swt.* sweet-scented ^ t A or 3 au Li N. Amer. ... D p.1 ■■■■■

28068 3920a *procumbens Leh.* procumbent £ A or ft my F N. Amer. 1827. D p.1 ■■■■■

473. COLLO'MIA.

28069 3924a *heterophylla Hook.* various-leaved O cu ft jn.au Pk Columbia 1826. S co Bot mag SJ⁵

28070 . . *grandiflora Dou.* greaUflowered O or 2 jn.au Pk Columbia 1826. S co Bot. reg

*474. POLEMCIUM.

28071 3926a *lacteum Leh.* mWk-cooored ■ A or lftmy.jn W 1829. D co

28072 3927a *moschatum Worm.* musky ■ A or lft my.jl Bk N. Amer. 1827. D co

|j-73 . . *Richardsoni Grah.* Richardson's ■ A or ft jl.o Pa.B N. Amer. 1826. S co

w74 . . *villKisum Rudol.* villcus ■ A or ft jno Pa.B Siberia 1826. S co Swr.

28075 . . *hymile R. f. S.* humble ■ A or ft jno B N. Amer. 1826. S co

<<^ *78. NEMO'PHILA.

28076 3938a *parviflora Don.?* small.flowered ^ A or 1 ■ ! X. Amer. 1836. S p.1

Phacdha parviflura Ph. Eutoca patviflfra It Br

- [^]J292. 478a. EUVTOC A R. Br. EUTOCA. (*Eutokos*, fruitful; number of seeds.) *Hydrophyllaceae*. 2.
 ZZL' - multiflora Dow. many-flowered 3 el 1 myjl Pk N. Amer. 1826 S co Bot reg. 1180
 ZW8 - Franklin R. Br. Franklin's Q el 1 apjl Pk N. Amer. 1827. S pl
- ⁴⁸⁷ ffYOSCVAMUS.
 D₂ dura for 8k nucrithus Led. small-flowered Datunu JMte tt Jor l myjl Y Egypt m 1829. C pl
 40Gla glatica Grah. glaucous O or SO au.o Y.o Bu. Ayr. 1887. S co Bot mag. 2837
 - noctiflora Hook. night-flowering O or 2 au W.P Chile 1826. S co Bot mag. 2785
 - alata L*. winged O or 2 au.o W? & Amer. 182a S co
- ^{PEIPNIA.}
 acuminata Grah. acuminate-/* £ .A] or 2 jns W Chile ^{IMW B} 1827. S co
- ² I S ^ 6 Pa.Pk Algoa Bay 1823. C pl Bot reg. 1116
- ^{SIRR AJ^9, GL'IA.}
 ^ 426a gracil Dou. slender O or } jnjl Pk N. Amer. 1826. S co Bot mag. 9924
- ⁴⁹² IPOMO PTS Mx. IPOMOPSIS. (*Ipo*, to strike forcibly, *optis*, sight; dawling flowers.) *PoUmonidee*. L—.
 >^a Regans Mx. elegant £ Q spl 2 jl S N. Amer. 1826. S co Bot. reg. 1281
 Guia pulchilla Dou. Gilia coronopifolia No. 4265.
- ^Q S12B. SPHENOTOMA Swt. SPHENOTOMA. (*Spheno*, to bind, *tome*, a slice or section.) *Epacridra*. L—2.
 " * gracilis 5w/. slender * i_J or 2 ap.my W N. Holl. 1823. C sp Sw. au. 44
 Dracophyllum gracie R. Br. No. 4314.
- ^{25S} 4330a iw Vt SSiv? J? H? many smked «i_J or Z my.au W N. Holl. 182a C Lp Bot cab. 143&
 88089 4341a Jctifl Ua ^ a*. Ledum-leaved »U or 8 mr.my W China 1819. C pl Bot. reg. 811
 4347a 1 f a ^ / % N a 4 : 1 | W . • or 4 my.jn F hybrid ... L sp Sw.fl.gar.n.s.lO
 2 prae'stans Swt. excelling • or 4 my.jn Cop hybrid ... L up
- ² I or 20 ... W Caracca. 182a C r.m
- ² cymose t^ orlo. R Indies 182a C pl
- ^{4j} 4. j £_g Ta ^ S_g my.s W Kindle. 18% C ? ! jj ^{jj}
 - densiflbra Wai dense-flowered • or 4 jn W K Indies 1824. C pl Bot reg.
- ^{550a.} ANTHOCLEPSTA Ajz. ANTHOCLEISTA. (*Anthot*, a flower, *kleistos*, shut up.) *Apocynace*. L
 - macrophylla G. Don Jong-leaved 1 • or 20 * W S. Leone 1820. C pl
- ^{457a} hymenandra Wai. membrane-ant*i*_J or 8 myjl Pk Silhet 1883. C pl
- ^{459a} obovata Wai obovate-taww* 1 • or 20 ... W Silhet 1816. C «pl
- ^{581a.} ULLOM Pers. ULLOA. (*Antonio Uuaa*, a Spanish naturalist.) Ataaew. L
 - pararitica Pers. parasitical £ • or. R Peru 1824. C hr m FL per. S. 185
 Juanulloa paruitica R.\$P.
- ² Alaternus-like i 1 ^ o r 6 f.ap Pa.Y Trinidad 1824. C pl Bot mag. 2929
- ^{V&S&SSJM} whitened J J or 3 Jn ... Chile 1825. C U
 - wiyrif ilium Lad. Myrtle-leaved * LJ or 3 ... B C 81 Bot cab. 1431
- ⁴⁸⁸⁵ «a Sm L? Jrf/C az ^ • o r 4 J a Pa.B Nepal UBS? C Lp Bot reg. 1235
- ^{^n⁶⁰⁵}- ADENO'PHORA.
 4016a reticulata Fis. netted k A or 2 ... Pa.B Siberia 1829. D co
 4965, liliiflbia w. Lily-leaved ^ A or 3 my.s B Siberia 1784. D pl
 Adenuphora ril'liflbia Na 4917.
- ^{sis ?@} K₁ K₂ K₃ K₄ K₅ K₆ sharp, notched ffeIor 2 s. Y Chile 1824 D i.p Bot reg. 973
 SSSSaK₁K₂K₃K₄K₅K₆ fcspr Sjljl Pa.B Nepal 1825 D co
 ^ o r S jl.o B N. Amer. 1825. D sp
 Baxteri, No. 5182. -uJpr 1 ap.jl O N. Holl. 1824. C pl Sw.au. 46
- ² «Si ^ u S T S 1 d E ^ Douirlas's A or 20 Jls O N. Amer. 1824. C co
 S219[«] SSSSi¹ Sal toSSsowered] or 20 jU Y.w China J826. C co Bot reg. 1238
- ² ^ ta Trt J S tr. involucrated & cu 3 my Y Huds.Jay 1824. C pl Bot reg. 1179
 campanidbra bell-flowered ft or 2 my Y N. Amer. ... C co Bot cab. 1361
 XylOsticum campaniflorum Ifsd.

646. WE'BER/f.
SS110 5336a coriacea *Wai.* coriaceous-tod sh • or 6 ... W E. Indies 1828. C p.l

32g7. 649a. UNCA*RIA *Gae.* UNCABIA. (*Uncys*, a hook; old petioles.) *Rubihcea.* 2.—10.
28111 . sessilifrctus *Rox.* sessile-fruited I O^or 10 . . . E. Indies 1839. C p.l
& . *GdmnbierWz.* Gambier L D^or 10 . . . Pa.R E, Indies 1825. C p.l Lin.tr.9.82
Nauclea *Gdmnbir.* No. 5347. *

660. ZI'ZYPHUS. ^
28112 5136a melastomffidesGrm. Melastoma-like*i_J or 6 . . . N. Holl 1824. C p.1

3298. 667a. RETANI'LLA Brong. RETANILLA. (Its name in Peru.) *Rhdmne**. 2.
& . obcordata Brong. obcordate/wf * p^or 2 . . . Y p_eru 1822. C Lp Ven.cels92
CollUVz obcordCta Koi. No. 5488, also 5489.

•3299. 6G7& TREVOM Cav. TRRVO*. (r#wo, the name of some botanist) *Rhdmnete.* 1—2.
28113 - triplintfrvis Gif. triple-nerved *|_j or 4 . . . G.Y Chile 1828. C p.l Bot mis. 55

3300. 668a. COLUBRrNA Brong. COLUBBINA. (*Coluber*, a snake; twisted stems.) *Rhdmnear.* *—■

SJ01. 6636. WILLEME*T/f Brong. WILLEMETIA. (*P. A.tff/feJ?<?, auth. of Herbarium Mauritianum.*) *Rhanmeee.*
4 - - africana Brong. African • L_ or 6 mr.ap B_a C. G. H. 1712. C p.l
Ceanthus africanus, No. 5510.

673. DIO'SMA.
28114 5562a sphserocphala lad. round-headed *|_J or mr.my W? C. G. H. . . C p.l Bot cab. 1490

3302. 673a. ACMADETNIA B. \$ W. ACMADENIA. (*Ahne*, a point, *aden*, a gland; anthers.) *Ruthcece.* 1: ^
\ . . tetrogoia Hil. four-angled «i_| or 2 apjl W C. G. H. 1798. C p.l Thun. v.4.5
Adenindra ttrabgna Swt. No. 5573.

t 675. ADENA'NDRA.
linearis Hil. YmezrJeav'ed n. i_J or 1 mr.jl W C. G. H. 1800. C p.l
Diósma linearis Thun. No. 5541.

3303. 675a. COLEONEWF A B. & TV. COLBONEMA. (*Koleos*, a sheath, *nema*, a filament) *Rutulece.* L—&
& . . "alba 5. # W. white flLi_Jor 2 mr.jl W C. G. tt 1800. C p.l
Ditoma alba Thun. No. 5544.

676. BARYO'SMA.
\\ 5579a pulchilla Hil. neat * L_J or 2 f.s P C. G. H. 1787. C p.l Bot mag. *I* ■
Agathosma pulchella W. No. 5581, also 5582.

3304. 676a. MACRCSTYLIS B. % W. MACBOSTYLIS. (*Makros*, long, *stylis*, a style.) *RiittVea.* %—*•
\ . . barthta//%/. bearded *i_J or 2 ap.jn W C. G. H. 1810. C p.l
Agathosma barb^ta Spr. No. 5589, and 5583, belong to this genus.

681. THOUA^SIA.
28115 5622a dumosa *Cun.* bushy * i_j or 2} my.jn W N. Holl. 182a C p.1

687. ESCALLO^N7il.
28116 5639a glandulbsa 5m. glandular < J or 3 s R Chile 1827. C l.p Bot cab. *T* ■
28117 - - *rubraPers.* rcd-floated i j or 3 I R Chile 1827. C Lp *B* m a g
28118 - - bitidaZ.ft.4r.O. bifid ft LJ or 3 . . . S. Amer. 1827. C Lp

3305. 692a. BERZETLI^ Brong. BERZEMA. (*Berzelius*, a celebrated chemist) *Brutudcae.* 9-T'
{ . . abrotanoides Brong. Southernw.-lk*|_J or 1* my.jl W C. G. H. 1787. C p.l Bot cab. 55?
Brinia abrotanoides, No. 5658., and also 5651. belong to this genus.

3306. 6925. RASPA?U4 Brong. RASPALIA. (*Raspail*, a French botanist) *Bruniacta?* 1.
\$ - - microphylla Brong. small-leaved tl|_J or 1 jn.au W C. G. H. 1804. C 81
Brinia microph^lla, No. 5660.

3307. 692c. BERA'DIA Brong. BEHABDIA. (*Berard*, professor of chemistry at Montpellier.) *Bruniacee.* *•
\ . . palecea Brong. chaffy i J or 2 ji..au W C. G. H. 1791. C p.l
Brinia paleoca, No. 5667., and also 5661. belong to this genus.

3308. 6924. AUDOUI'N/J Brong. AUDOUINIA. {*Audouin*, an entomol. and friend of Brongniart) *Brvn** /
% . . capitata Brong. headed «. LJ or U my.il P C. G. H. 1790. C p.l
Diosma capitite L. No. 5552.

3S09. 698. STAA^V/il Thun. STAAVIA. (*Martin Statf*, a correspondent of Linneus's.) *Brunttce**. *T''
28119 - - radiata Thun. radiated a-Lj or 1 myjn W C. G. H. 1787. C p.l Brey.c»
Phljica radiata L. Brinia radiata L.

£8120- - glutinbsa Thun. clammy tt.i_Jor Ijap.my W CO. H. 179J. C p.l BoL ■
BrumVi glutinbsa L.
& . . ciliata Brow.; ciliated-feawd tt. i_J or 1| my.jl W C. G. H. 1812. C p.l
Brinia ciliata, No. 5667.

*695. CLAYTON/iL.
£8121 5681a cuWnsis H. \$B. Cuba O c u } my.jl W Cuba 1812. S co

*701. JTOLA.
28122 5728a prsm6rsaa Dou. bxten.rooted ^ A or ± my.jl R Columbia 1826. D co ^{Af} refi.12^{5*}
28123 5752a flavicornis Sm. yellow.horned £ A or \ my.jl Y.B Britain ch. pi. D co
Bot res
28124 5783a epipsHa Led. naked above £ A or \ my.jl Y Livonia 1826. D co

3310. 709a. HYMENANTHE'RA R.Br. HYMENANTHERA. (*Hymen*, membrane, *anthera*, an anther.)
28125 - - tenuita R.Bt. toothed-fcawrrf * i_J or 6 ap.my Y N. Holl 1820. C p.1

&J11. 710& SOULANGif Brong. SOI^ANCIA. (*Soulange Rodin*, a nurseryman near Paris.) Jtrf***
{ . . axillem Brong. axiflowered « Lj or 2 myjn W C. G. H. 1812. C p.l
Ph^hca axillaris, No. 5860., also 5856, 5854, 5854, and 5855, belong to this genus.

CLASS V ORDBH 2.

PENTA'NDRIA DIGY'NIA.

RPBES.

	alplnum												
1819	2 pumilum <i>Lindl.</i>	dwarf	A	cu	2	ap.my	G	N.	—	—	C co		
28127	toniKjrum <i>Lindl.</i>	slender-flwd	A	cu	6	ap.my	Y	N. Amer.	...	C co	Bot reg.	1274	
	flavum <i>Coll.</i>	yellow	**	cu	6	ap.my	Y	N. Amer.	...	C co	Col. h. rip.	J. B.	
	aareum sanguineum <i>Lindl.</i>												
	atropurpureum Lefr.	dark purple	sit	or	4	ap.my	G	Siberia	1826.	C co			
181S	cereum Dow.	wax	Ss	or	2	ap	W	N. Amer.	1827.	C co	Bot reg.	1263	
28131	punctatum A * P.	dotted	A	-J cu	3	ap.my	G.v	Chile	1826.	C co	Bot reg.	1278	
28132	decumbens Led.	decumbent	flk	fr	1	ap.my	G	Siberia	1829.	C co			
	setosum <i>Lindl.</i>	bristly	A	or	4	ap.my	W	N. Amer.	1810F	C co	Bot reg.	1237	

88133  GONO'LOBUS.
viridiflurus Mty. green-flowered ^Qcu 10 s G S. Amer. 1826. C p.l Bot reg. 1126

776. ~~Lindl.~~ Pallida Lindl. pale JL • or 6 jl.au W E. Indies 1815. C r.m Bot cab. 1309

28134 lanceolate *Haw.* № 6204. accta *Haw.*
BRACHYSTE'LMA.
lanceolata *Inf.* A. A. Linn. 1. in ill. C. C. H. 1826. C. 1. R. et reg. 1113

-**PALLIUM**-

white-flowered k A or 1 au.s W N. Amer. 1826. D p1

28137  Richardson's £ A cu 1 ... G N. Amer. 1827. D co

carpinifolia Lindl. Hornbeam-lvd tm 40 ap.my Br Britain hed. L co.
stricta LindL strict tin 60 ap.my Br Britain hed. L co.

Na 6582, and 6577, of *Cherophyllum* belong to this genus.

2814A* 821a. CONOPCTDIUM Koch CONOPODIUM. (*Konos*, a cone; *podion*, a little foot; disk.) *Umbelliferae*. 3.
 - denudatum Koch denuded & A un 2 jnjl W S. Europe ... D co
 - •*Canadense* Koch Canadian ^ A un If jl.au W N. Amer. 1699. De o
 - Chaerophyllum canadense Pars. Na 6599, and also 6600, belong to this genus.

* 826a. ORLA^Y Hofm. OHLAYA. (*Orlay*, probably the name of some botanist) *Umbelliferae*, a
platyc&rpos; *Koch* broad-fruited O un 1 jn.jl W S. Europe 1800. S co M. h.9.14.2
Cadicalis platyc&rpos;, Na 6659., also 6663. and 6656. belong to this genus.

8266. TURGE^NIA Hofm. TuuaENia. {*Turgeo*, to swell; seeds.) UmbeUlerts. 1.
 latifolia Hofm. broad-leaved O un 3 jLau R England ch.fi. S co Ene_k bot 198
 Caticalis latifl.Ua, Na 6657.

835. ^THAMA'NTA.
Ua macedonica 5/r. Macedonian £ Q cu 2 jl.au Pa,Y Greece 1596. S co Black. 382
macedonicum L. Na 6745.

8H. SE'SELI.

No. 6746, 6747, 6748, 6749, 675a of iftbon, and 6703. of jfthamanta, belong to this genus.

Atfseli, belong to this genus.

podóhium *Hori.* Podolian
viréscens *Spr.* greenish + un 1 jl W Podolia 1827. D co
- 8a- HELOSCIADIUM Koch HELOSCIADIUM, (*Helos*, marsh, *skiadion*, umbeL) UmbeWfene. 5.-7.
- nodiflbrum Koch knot-flowered *W A w f jl.au W Britain rivul D aq Eng. bot 639

51um nodiflorum L. No. 6796., also 6797.6804.; and Na 6809. and 681 a of Sison, belong to this genus.

**&c- DREPANOPHYLLUM Hofm. DHEPANOPH. (*Drepanon, scythe, phyllon, leaf; form.*) UmbeL J,
agreste Hofm. field ^ A un S jn W Europe 1726. D m.s Jac. au. & 257

^{33 19} ;85*». ZI'ZMKoch Zizu. (J. B. Zizii, a German botanist) UmbeMlferte L
 golden i A or 2 ray.jn Y N. Amer. 16S9. D'ram

1

860. GNI DIUM.

{ 6851a *a/vio/des Spr.* Apium-like f A un 2 jnjl P.Y ConstantL 1816. D e o Ven.cels⁹⁷
Zaserpitium rilaftlum Jac. No. 701&, auro 6766. under G. venusum.

*8G7. TRACHYME'NE.

28144 6922a *csriilea Grah.* sky-blue O pr 1J jl.s B N. Holl 1827. S s.p Bot recg. 12»
Didfscus csrbleus Dec. Bot Mag. 2075

333a 873a. MELANOSELPNUM *tifin.* MELANOSEL. (*Melas*, black, *tfi'lio/i*, parsley; appearance.) 
 & . . *decipiens Hiffm.* deceiving j i A u n 2 jnjl W Madeira 1785. I co Set han.
Sellnum decipiens mrf. Na 6941.

3321 874A. ARCHANGE'LICA *Hofm.* ARCH ANGELIC A. (Supposed archangelic virtues.) *UmbeUfertr^J*
 & - officinalis Hq>>. officinal k CDcul 4 Jnau G England watpL S ms FI. dan. 206
Angelica Archangelica L. No. 6957., and also 6968. belong to this genus.

3322. 8746. OSTFRICUM *Hqftn.* OBTERICUM. (*Osteon*, a bone; seeds ?) *Umbetyferee.* I-
 h . - pratense *Hofm.* meadow f A un 3 jn.au F Galicia 1818. D co
Angelica pratensis Spr. No. 6964.

875. LIGU'STICUM.
 & 6971a */erufaceum All.* Ferula-like 9t Q un 1 jl Pa.P Switzerl. 1775. S co
/ascrptium lucidum W. No. 7018. and No. 6938. of Sellnum.

3323. 875a. MOLOPOSPE RMUM Koch MOLOPOSPERMUM. (*Mobops*, a wheal, *sperma*, a seed.) *UtnbeUfere*⁻¹
 \ - - pelopponesiacum Koch Peloponneses. ^ A or 4 my.jl Pa.Y Switzerl. 1598. D co Jac au.5.1⁹
lifisticum pelopponesiacum L. Na & X&.

3324. 8756. LEVISTICUM Koch LEVISTICUM. (*Levo*, to assuage: relieves flatulency.) *UmbeWfera.* 1.
 t . . officinale Koch officinal 4 A cul 6 jn.jl Y Italy 1596. D co Black. 275
Ligusticum Levisticum L. No. 6966T

3325. 875c. TROCHISA'NTHES Koch TBOCHISANTHES. (*Trochiskos*, a small wheel, *anthos*, a flower.) *Undgl-*⁻¹
 § - - nodiflbra Koch knot-flowered ^ A un 3 jlau G Alp. Eur. 1810. D p.1 Alped.72
Ligusticum nodiflbum VU. Na 69667 belongs to this genus.

876. PHYSOSPE'RMUM.

No. 68291 of *Smyrnium* belongs to this genus.

88a FE'RULA.

28145 6987a *cannestris Bes.* field jl A or 3 jn.jl Y Tauria 1829. S co
 28146 - *sylvatica Bes.* wood k A or 3 jn.jl Y Podolia 1829. S co

3326. 880a. FERULA\SO Koch FERULAGO. (*Ferio*, to strike: used as rods.) *Umbellifer*^{*}. 2.
 ^ - - *nodiflbnKoch* knot-flowered 9t A un 3 jn.jl Y S. Europe 1596. D 81 Boc.mu.70
Ferula nodiflora L. Na 6990., also 6988.

3327. 881a. SVLER Sco. SILER. (*Salio*, to shoot up; quick growth.) *UmbeWferte.* 3.
 § - - trilobum Sco. three-lobed/pd ^ A un 4 my.jl W Austria 1796. D e o Jac. au. 
Laserpitium trilobum L. aquilciforum 5>r. Na 7006.

882. PEUCEDANUM.

No. 6934,69J5,6937,694a 6914,6945,6947. of ftñnam, Na 6932. of *Thyssellnum*, and Na  belong to this genus.

3328. 882a. SILA'US Bes. SILAUS. (A name used by Pliny.) *UmboUferte.* 3.
 § - - prat'nsis Bes. meadow ^ A un 2 jn.au Y England mea. D co Eng. bot 21⁴⁴
JPeucedanum fflalus L. Gnidiun Silaiis Spr. No. 6852.; also Na 7052. and 7036. of PeucSdanum.

3329. 883*. OPO'PANAX Koch OPOPANAX. (*Opos*, juice, *panax*, the plant y'elding it) *Utnbellifer*^{*}. 
 | - - Chirum'cutn Koch Chiron's f A m 6 jn.jl Y S. Europe 164a D co Gou. il 19.
Pastinaca Opupanax L. No. 7049.

3330. 885a. PETROSELNUM Hofm. PARSLEY. (*Petros*, a rock, *selinon*, parsley: habitat) 
 § . . satvum flo/wi. cultivated f O cul 2 jn.jl L.Y Sardinia 1*48. S co
^Tpium Petrosellnum L. Na 6833.; also Na 6807. of Siñon, and Na 6975. of Ligulsticum.

3331. 8856. CONDYLOCA'RPUS Hofm. CONDYLOCARPUS. (*Kondylc*, a knob, *karpos*, fruit; margin.) *U*
 \ - - officinalis Koch officinal Ow 2<jl F England cor. fi. S co Eng bot.
rordylum officinale L. No. 7082., also Nos. 7084 and 7087. belong to this genus.

S332. 885a PLATYSPE'RMUM Hofm. PLATYBPERMUM. (*Platys*, broad, *spcrma*, seed.) *UmbelW*
 4 - - muricatum Hofm. xtxcMysseeded O un 2 jn.jl Pk Barbary 1683. S co M n. 9.
 i - - pulchfrrium Koch fairest O un 2 jn.jl W Levant 1699. S co
Caucalis orientalis But. Na 6664. and also Na 6667. belong to this genus.

CLASS V. ORDER 3.

PENTA'NDRIA TRIGY'NIA.

903. DRYMA'RIA.

28147 7254a *gracilis Li:* slender ^ (23 un } jn.s W Brazil 1829. S co

. . . 906. PHARNATEffAf.

S8148 7259a Ho«VnanseggiatiimR.&a Hoffman.^ iQJ cu ijL.au W Brazil 1829 D e o


CLASS V. ORDER 5.

PENTA'NDRIA PENTAGY'NIA.


mall-pierced fctAlor 1 an W CG.R 1824 C .1
 wavy.,*, -tU or | au W C ft H. IK C P»


CLAW VI. Oftsn 1.

HEXA'NDRIA MONOGY'NIA.

iU« W M«, W. O « 
 Pancratium incarnatum iUA.

bloody f El or 2 aus B.w Rio Jan. 1834. Skr.m


 TLLA'NDSJM. , , " , , w

 WLLBERGJil. , , , m i w , , n. Rio Jan. 182& Sk r.m Bot.reg. 1130

 ruseus Swfc A m  i A P *
 ruseo-ecologica

 Imatoph^llum Aitbni Hook. Bot Mag. 2836.

*967. BRUNSVI'GM
 S " " El cu 1 " Pa C. G. R 1826? 0 r.m Bot reg. 12W
 Amar^UU coranica var. pallida Lindl.

 ffig^uncinata^tong-pedtf tAJor 2 mr.ap 0 Mexico 1826. 0 r.m Bot reg. 1188
 acuminate var. longipedunculata Bat. reg. 534

 Sr* A Jt2 <^ S ?^ u a ro b u <t * j U o r U n j l R Ba Ayres 1887. 0 r.n Sw.fl.gar.n.il*

*979. ALSTRffME^RLI.
 Hookerff SwL » pulchfla Sims Bot cab. 1272, Na 80W.
 Sims/Swt. is rbsea Hook. Sw.fl. gar. 207 No. 8W4.
 nallida Grah pale AiAJel 2 <o PkR 'Chile?' 1827. 0 Lp
 ^ S ^ S n i f A . parrot *! Se 1 6 <o Cr Mexico 1829. 0 Lp Sw.fl.gar.n. 15
 ^ 980o. CDMMI'NGWD. Don. CUMMINOU Ady Gordon Cumming, of Altyre, nearForres, NB.) *Apto.* 1.-2.
 " campanulataD/jo) cainpanulate tf JEl \ jln B Chile 182a 0 Lp Sw.fl.gar. 257
 Conanthra campanulas Hook. Na 805a

88164 8100a n ^ S ^ 'SS ^ noddingifwd f E 3 c u 1 n Ld E Indie* 1822. R Lp Bot reg. 1223

 Cape myUun 1 myjn Ap C.G.H. 1829. D mi

 lEIzaor lmy.au W Brazil 182, D co B *~»»

 ny.uGT C.G.H. 1824 m*

 or 4 MP W China ... O co

 SteJML Pe<.«Vy, «l 1 TM SBk Penia » ° » Batit«.ll«

28@' 108o. CYCLOBCTHRA SwL (A'^fb, acircle,>ottr@s,apit; each sepal) Tuiiphce*. 2.

E^reaSuJ pu^le S i5Jor 3 aus P.G Mexico 1827. Op.1 Sw.fl.gar.n. 2U
 88170 - tetf^TC^Xl «^lor 3 a,, Y Mexico 1837. O pi Sw. fl. gar. 273
 Pitthna barbata Kth.

- | | | | | | | | | | | | |
|---|--------------|----------|------|-------|-------|-----------|-----------|-------------|-----------|------------|------------------|
| 1020. DRAC£?NA. | | | | | | | | | | | |
| 28171 8461a brasiliensis <i>Parm.</i> | Brazilian | * s* or | 6 | Jl | ... | W | BrazU | 1825. S | p.1 | | |
| 28172 8471a surculosa <i>Lindl.</i> | surculose | s* pr | 4 | | | W | S. Leone | 1821. Skp.1 | Bot reg. | 1169 | ,tea |
| •1027. YU'CCA. | | | | | | | | | | | |
| 28173 8518a puWrula <i>Haw.</i> | small downy | £ A or | 7 | aus.s | | W | | .. | Skri | Sw.fl.gar. | 25 ¹ |
| 28174 8525a graciilis <i>Lk.</i> | slender | £ ES or | . | . | . | W | Mexico | 1829. | Skr.l | | |
| •1046. A'UAVM. | | | | | | | | | | | |
| 28175 8708a nudicaule <i>Leh.</i> | naked-stmd | tf | A or | 1 | Jn,jl | — | | 182a | O co | | |
| 28176 - eu6smoN Lk. \$f. 0. | welUscented | V | A or | 1 | jn,jl | W | | 1829. | O co | | |
| 28177 - glandulidsum Lk. \$ O. glandular | | tf | A or | 1 | jn,jl | W? | | 1829. | O co | | |
| 3338. 1050a. HESPEROSCO'RDUM <i>Lindl.</i> HESPERO (Hesperos, evening, rtordott, garlic; smell P) <i>Asphodel</i> **- | | | | | | | | | | | |
| 28178-A y Ayatihimw Lindl. Hiatihimlike \$ A Jl or 11 JLau BP | | | | | | | | 1826 | O p.1 | | |
| Brodug> grandiflura Nutt. | | | | | | | | | | | |
| •1054. SCI'LLA. | | | | | | | | | | | |
| 28179 8821a obtusifolia <i>Poir.</i> | blunt-leaved | tf | A P» | .4 | Cap | R | S. Europe | 1829. | O c | | |
| 28180 8822a rosea Uh. | rose-colored | tf | A Pr | ■ | — | R | Nuroidia | 1827. | O c | | |
| 1085. CYANE'LLA. | | | | | | | | | | | |
| 28181 9035a odoratissima <i>LindL</i> most fragrant | ft | tAl or | 1 | jlau | L | C. G. H. | 1826. | O s.p | Bot. reg. | UU | |
| •1091. MAHO% N£4. | | | | | | | | | | | |
| 28182 9069a repens G. Don | creeping | ft | or | 2 | ap | T | N. Amer. | 1824. | R r.m | Bot reg. | H7 ^{fi} |
| Berberis repens LindL | | | | | | | | | | | |
| 1099. FRANKFNM. | | | | | | | | | | | |
| 28183 9089a pauciflora <i>Dec.</i> | few-flowered | tt.i_Jpr | 1 | jaau | Pk | N. Holl. | 1824. | C s.p | Botmag. | 289& | |
| 1100 .PE^PLIS. | | | | | | | | | | | |
| 28J84 9091a occidentalis <i>Spr.</i> | western | ■ | — | \ jl | O | Guadalou. | 1829. | S | bog | | |

CLASS VI. ORDER 3.

HEXA'NDRIA TRIGY'NIA.

- | | | | |
|--------------------------|-------------------------|---|---|
| 3339. 111a. CALOCHCKRTUS | <i>Ph. CALOCHORTUS.</i> | (<i>Kalos</i> , handsome, <i>chortos</i> , grass.) | <i>Lilihccce. 3.</i> |
| 28185 - - nitidus | <i>Dou.</i> | shiping tf [A] pr | 1 au P Columbia 1826. O s.p. Hort tr. ^l ₉ |
| 28186 - - macrocarpus | <i>Dou.</i> | long-fruited 9 ^J spl | 2 au P Columbia 1826. O s.p. Bot reg. 11 ^{**} ₉ |
| 28187 - - degansPA. | | elegant tf ^J pr | .in.jl W Columbia 1826. O s.p. Horttr.7- |

CLASS VIII. ORDER 1.

OCTA'NDRIA MONOGY'NIA.

- | | | | | | | | | | | | |
|--|--------------------|------|---------|--------|------------|---------|----------|---|-------|---------------------------|------|
| • 114a TROP-ffIOLUM. | | | | | | | | | | | |
| 28188 9317a tricolbrum Swt. | three-colored | I | A el | 12 | jas | O.p | Valpara. | 182a | C p.1 | Sw.fl.gar.tf ⁰ | |
| 1172. BJE'CKIA. | | | | | | | | | | | |
| 28189- 9397a sax i cola Cun. | rock-inhabiting | II | i_J or | 2 | jLau | W | N. HolL | 1824. | C p.1 | | |
| • 1173. ERPCA. | | | | | | | | | | | |
| 28190 942°1 rigida Lod. | stiff | * | L j o r | 2 | jl.au | Pk | C. G. H. | 1820. | C s.p | Bot cab. | 12° |
| 28191 - - rubida Lod. | ted | | | 2 | myjn | W.R | C. G. H. | 1826. | C s.p | Bot cab. | 1\$° |
| 28192 - - crinlo Lod. | hairy | | | 2 | ... | R | C.G. H. | 1826. | C p.1 | Bot cab. | 1** |
| 1177. MICHAU'X/J. | | | | | | | | | | | |
| 28193 9951a decandra Fit. | decandrous | ^ | Qb or | 3 | jlau | LB? | Persia | 1829. | S s.l | | |
| 1179.  | | | | | | | | | | | |
| 23194 9969a  Cua Paullinia-lvd | il t_J un | 4 | jn.jl | G | | N. HolL | 1826. | C p.1 | | | |
| • 1181. OSBE'CKIA. | | | | | | | | | | | |
| 28195 9978a glomerata Dec. | g'amerate. JlwD J* | O pr | lfjLau | Pk | Trimidad | 1818. | S p.1 | Bot mag. | 5 | | |
| Xhtxifl glomerata | Bot. cab. | 3M. | | | | | | | | | |
| • 1184 ffI NOTHETRA. | | | | | | | | | | | |
| 28196 10019a dec6mljen8 Dou. | decumbent | O pr | 1 jan | P | California | 1827. | S co | Bot. mag. | | | |
| 28197 - - vineinea Dou. | twiggy | O pr | 3 jas | P | California | 1826. | S co | Bot mag. | | | |
| 88198 - - tenuifolia 11. # P. | fine-Leaved | O b | I jns | P | Chile | 1828. | S co |  | | | |
| 28199 - - miadriyflchraDott. | 4.*potcd.pctalcld | O or | II 8 | Pk | N. Amer. | 182a | S co | Bot. reg. I | | | |
| 28200 10020a Lindleyii Dou. | Lindley's | O or | U_jn.jn | P | N. Amer. | 1826 | S co | Bot mag. | | | |
| 28201 10020a pallida LindL. | ra,jlowercrd | ^ ^ | I*jas | W.R | America | 1826. | D p.1 | Bot reg. I | | | |
| 28202 10037a  araxacifolia Horst. | Dandolion-lvd | A or | W | Persia | 1825. | | |  | | | |

- | | | | | | | | | | |
|--------|------------------|-------------------|---------------|------------|-----------------------------|-------|------|---------|-----------------|
| | grandiflora A^f. | | | | | | | | |
| •1186. | EPILO^BIUM. | | | | | | | | |
| 28203 | 10053a | crassifilum Leh. | thick-leaved | ■ A or 1 | jn.j1 | R | - - | 1829. | S co |
| *1188. | FU'CHSW. | | | | | | | | |
| 28204 | 10077a | microphylla Kth. | small-leaved | * LJ pr | 6 | jn.i | S.P. | Mexico | 189R C |
| 28205 | 10079a | thymifolia X/A. | Thyinlcavcled | ft LJ or | 6 | my.o | R | Mexico | 1827. C p.1 |
| 3340. | 1192a. | PIERA'RD/JJ | Rox. | PIERARDIA. | (Mr. Pierard of Kew Green.) | | | | Sapindheeatt 1. |
| | | sávida Rox. | well-tasted | f • fr | 20 | ... | Y | Sumatra | 1820. C p.1 |
| 3311. | 1195a. | MA' RLE A | Rox. | MARLUA. | (Marliya, its Bengal name) | | | | |
| 28907 | - - | begon/d'o/ra Rox. | Begonia. | leaved | III | ior 4 | ... | Y China | - C p.1 |

ogo^ 1199a. GNI'DIA.

^*w 10174a ochroleuca Laf. yellowish white* i_J or 1J ap.my Pa.Y C. O. H. 1820. C p.l Bot. cab. 1184

* } * 1204a. APLOPHY'LLUM J. APLOPHIYLLUM. (*Haploos*, simple, *phyllon*, lcaf j leaves simple.) *IlutHcett.6*—a
» - - patavnum.J. Paduan * A un l^jns G.v Italy 1819. C co Mic.gen. 19
fluta patavna L. Na 10920, and aboTfoi 10921,10922,10923,10924,10925. belong to this genus.


CLASS VIII. ORDER 3.

OCTA'NDRIA TRIGYNIA.

 i!Si^{7a} "ijuctndum B.H. disagreeable <. _] un 2 my G.w Valparaisol 825. C co Bot reg. 1250
*^{1U} 10279a acutatum Leh. sharpened O or 2 jLau Pk 1838. S co


CLASS X. ORDER 1.

DECA'NDRIA MONOGYNIA.

- *1237. SOPHCWA.
104^{Jo} velufina Lindl. velvety *i_|or 6 jn Pk Nepal ... C p.l Bot. reg. 1185
- 1239a. PIPTA'NTHUS Surf. PIPTANroua (*Pipto*, to fall, *anthos*,* flower; soon falling off.) Leg. Pap Soph. 1.
* - nepalensis Swt. Nepal a or 10 my.jn Y Nepal 1821. Lr.mSw.fl.gar.SW
ilnagyris fndica Wai. So. 10162.
-  fine-leaved «L J or 4jl.au Y C G. H. 1829. C p.l
-  smaUer k A or lijnjl B N. Amer. 1829. D co
-  i lor 1 my Y N. Holl 1820. C s.l.pBot.raag.28G0
tjor 2 apjl Y N. Holl 1824. C l.p
- & S i ^ o p h y S ^ . slender-leaved ; L J or 2 apjn Y N. Holl. 182J C s.Lp
phys&des Cun. bladdery • i_J or 2 apjn Y N. Holl. 1824. C s.Lp
-  * . . « 20 jl W Cumana 1826. C p.l Bot reg. 1133
- 1080a¹ &TM² ROYNA, royal I [I]sp140 ... C liadagasc.1828. C r.m Bot mag. 2881
- 1308a WALLSITRA Rox. WALLSUIUL³ (^a/te&i,itsTeUnganame) ¥ d t e . 1.-2.
robusta Ao*. robust t O or 20 E. Indies 1827. C p.l
-  1100a prostrata Swt. prostrate a. or J myjn W N. Amer. ... L p.s Sw.fl.gar. ic in,
- 2829 ni?? *HODEDE'NDRON.
^ U015a Gowenianum Swt Gowen's « or 4 myjn R hybrid ... L p.l Sw.fl.gar.263
- ^* ^I43a Banksii Cun. Banks's II LJ or 1j. . . . N. Holl. 1824. C p.l
- Stel^{5g} 62a- CHJETOGA'STRA Dec. CHATOGAOTRA. (*Chaite*, hair, /o<^r, belly; ovary.) *Melastbmear*. 1-28.
* lanceolita Dec. lanceolate/pft O P' 1 j* W Trinidad 1820. S p.l Bot. mag. 2835
JZh&xia lanceolita Bon. Rhex. 21
- 1364a. MOTJRTRIA J. MOURIRIA. (*Mouriri*, its name in Guiana.) Memecpleet. 1—5.
guianensis^<ft. Guiana * f a o r 6 jls Y Guiana 1817. C p.l Aub. guL 1.180
Petalbma Mourtri Swz. Na 11189, and also 11188. belong to this genus.
- *Y*⁵ 13646. POIV'RE^ Com. POIVREA. (*N. Poivrc*, intendant of the Mauritiis in 176&) Combretace. 6.-14.
coccinea Dec scarlet tf * or 20 jn.d S Madagas. 1818. C r.m Bot. mag. 21C2
Combretum purpureum Vahl, Na T0197, also 10198,10199,10201,10207, and 10208. belong to this genus.
- 34S⁶ 372a. DE/CODON Gm. DECODON. (*Dekas*, ten, *odous*, a tooth; calyx.) Salicariite. 1.
" " - verticillatus Ell. ytYtorX-flowered ^ A " 2 jLi P N. Amer. 1759. D p.l
Nesa;vi verticiUata Kth. No. 12199.
- 88\$J⁷- 13726. HORKETLJM Cham. HOBKBLIA. (*John Horkel*, profcMor of Pfywologyat BerUn.) 
congeita Duu. crowded-ytfur⁸ i A cu 8 au W N. Amer. IH26 D co

CLASS X. ORDER 2.

DECA'NDRIA DIGYNIA.

-  119S^a/AXI,FRAOA . A. A or * B W N. Amer. 1827. D p.l
-  BffiSaMFSS- i i S **myjn_w Europe 1826. D co
-  coloring £ A un * jnjil W N. Amer. 1827. D co
-  &JS5SE;LL pentandrous k A < * > Y g. Amer⁸ 1827. D co Botmg.2933
trifida Grah. trifld-teawed 5t A cu * my w K. Amer. IBZ. V co

- | | | | | | | |
|--|-------|------------------------------|-----------------------------------|----------------|-------------|--------------|
| • 3350. 1383A. TE'LLIM A it. Br. TELLIMA. (Anagram of Mitella; separated from it) Sarifrbgete. I—* | 28231 | grandiflbra Dou. | great-flowered f ^ e u 1 ap.my Pk | N. Amer | 1826. D s.p | Hot reg. U7» |
| 1384. GYPSO'PHILA. | 28232 | 11353a pubdscens <i>Led.</i> | pubescent | 3t A or 1 jnjl | W | Siberia |
| *186. DIA'NTHUS. | 28233 | 11393a giganteus <i>Urv.</i> | gigantic | £ A or 3 jlau | F | Greece |
| | 28234 | 11449a siculus <i>Pret.</i> | Sicilian | £ A or {jn.au | R | Sicily |
| | | | | | 1824. S co | Sw.fl.gar. |
| | | | | | 1829. C &l | |

CLASS X. OJLDER 3.

DECA'NDRIA TRIGY'NJA

- *1388. SILE'NE.
 28235 11489a Stevent'Spr. Steven's ^ A un I jnjl W Caucasus 1827. S co
 84236 11538a cernua Thun. drooping £ OJ un 1 ... W C. G. H. 1829. S co
 3351. 1389. LARBREYⁱ Hit. LARBRBA. (*Abbe Larbry*, a botanical author.) Paronychiles 1-
 28237 - - aquatica Hit. water O un | jail W Britain rivul S bogs Eng. bot 10^b
 Stellaria uliginbsa Cur. aquatica Poll. *JLarb&a aquatica*, No. 1417., is a true Cerastium.

1390. ARENA*RIA.
 £8238 11684a rubella Wahl. redduh_r; flwd. lg A.P'. k ny.&u^R ScotLBenlawers. D s.p Wa.lap.6
 cherlerrfilia G.Don (discovered in Scotland 1790), hirta Fl.dan. 1646

3352. 1398a. THRY ALLIS L. THRYALLIS. (Greek name for a plant of the mullein kind.) *Malpighidice**.
 28239 - . brachjftetachysLind. short-spikedfl. O el 10 s.o Y Rio Jan. 1823. C p.s Bot reg.

*3353. 1398b. GALPHI'MLI Cav. GALPHIMIA. (An anagram of *Malptgkia*.) *Malpighidice**. L-
 28240 - - glifica Cm. glaucous J. D " 8 — Y Mexico 1829. C l.p Cav. i 5-

CLASS X. ORDER 4.

DECA'NDRIA PENTAGY'NIA.

CLASS XI. ORDER 1.

DODECA'NDRIA MONOGY'NIA.

1434. EL^OCA'RPUS.
 2L253 12117a *grandiflorus* *Wai.* large-flowered 1 • or 20 W E. Indies 1829. C p.l
 • 1447. PORTULA^NCA.
 28254 12158a *grandiflora* *Hook.* great-flowered A lAl or ft jnj1 Y.P Chile 1827, S »1 Botnia»!
 3357. 1448a. CALANDRVNL* H. & B. CALANDBINIA. (*J. L. Calandrmi*, a Genevese botanist)
 28255 . . *grandiflra Lindl.* great-flowered
 \ - - *ascididens Hort.* []
 \ - - *paniculata Dec.* p. []
 Tannum paniculatum
 1449. ifNACA'MPSEROS.
 28257 12183a *intermedia* *Haw.* intermediate «. LJ CU * jLau Pk C. G. R 1824. C s.l

J < a HEIMM.

*2200n tnyrtiblia flbr/. Myrtle-leaved M J ■ ■ ■ Y S. Amer. 1826 C Lp
 - • huarisfolia ffort. Linaria-leaved * J ■ ■ ■ Y S. Amer. 1829 C Lp

CLASS XI. ORDER S.

DODECA'NDRIA TRIGY'NIA.

vina	Pentag&na a w.	five-angled	« ZD gr	3	-	Y	C O. H. 1824. C pi
i6^i ^a	"P^deni By	shining	ft	4	Jn.s	S	1. France 1886. C p.l
Jr,Tm ^d	angustifolia Lock	narrow-leaved	• CD cu	4	-	Y	Trinidad 1827. C p.l
2422a	KönzeLch.	Kunze'a	O un	1	■ ■ ■	Ap	Europe 1829. S co

CLASS XII. ORDER 1.

ICOSA'NDIUA MONOGY'NIA.

1470a.	MELOCACTUS C. Bauh.	MELONTHISTLE. {Melotya melon, cactus; shape.)	Cârte*	6.—10.
-	commimis IJc. \$0.	common tt.	ZD gr	jl.au ft W. Indies 1688. O s.p Dec. pi. 112
-	Cactus Melocactus L. No. 12511., and also 12512.	belong to this genus.		
-	pyramidalis Lk. \$ O. pyramidal	■ ■ ■ R	Caracas	— O sp L.&O.dis.25
-	placentif6nnis Dec. placiita-shpd n.	■ ■ ■ R	Brazil	— O sp L & O. dis. 21
-	Büser/ Lk. & O.			
-	Sellwt ^o Dec.	Sellow's tt.	ZD gr	— O sp L.&O. dis.22
-	Echinocactus SellwtY Lk. & O.			
-	polyacanthus Dec. many-spined tt-ZDgr	■ ■ ■ Brazil		— O sp L.&O.dis.1&l
-	Echinocactus polyacanthus Lk. \$ 0.			
1470ft.	ECHINOCACTUS Lk.&0.	ECUINOCAC. (Echinus, a hedgehog, cactus; spin..)	CActee. 22.—40 ?	
-	gibb ^a sa Dec	gibbous tt.	ZD gr	\ jLau W. Jamaica 1808. O s.p But reg. 1S7
-	Cereus gibbibus Swt. No. 12545.	■ ■ ■ and ViStil. of Cereus belong to this genus.		
-	ovaliflata Dec.	fenced round tt.	ID gr	— P Mexico — O sp Dec cac. 9
-	melocactiformis Dec.	McIon-th.-lktt.	ZD gr	— W Mexico — O s.p Dec. cac. 10
-	cornigera Dec.	hum-bearing tt	ZD gr	— P Mexico 1823. O i.p L. & O.dis. 14
-	platacantha Lk. \$ 0.	Cactus latispina Haw. No 12517.		
-	crisp ^a ta Dec.	curled it	ZD gr	— P Mexico — O s.p Dec. cac. 8
-	tenuispina Lk. % 0.	fine-spined tt.	ZD gr	— P Mexico — O s.p L.&O.di. 19.1
*	recurva Dec.	recurxeiSpindca. ZD gr	1	— Y Brazil — O s.p L.&O.di. 19.1
-	Cactus recurva Mil. nobilis L. No. 12519.			
-	inforta Dec.	twi&tcft-spined tt.	IZ3 gr	1 Jn.jl P Antigua 1768. O &p PL am, & 8
-	Cactus intorta Haw. No. 12520.			
-	SahmTwa Lk. & O. Salm's	tt.	— gr	— Curacao — O a.p L.ftO.dis.13
-	tubercuftta Lk. \$ O. tubercled	tt-ZUGr		— Mexico — O s.p KftO.dis.2fi
*	gladifata Lk. \$ 0.	gladiated tt.	— gr	— Mexico — O &p L. & O. dis. 17
*	subuliflra Lk. \$ 0.	awl-bearing a.	ZLJgr	— Mexico — O s.p L.&O.dis.27
*	depr ^a ssa Lk. A. 0.	depressed tt.	— gr	— S. Amer. 1798. O <>p
Cactus depr ^a ssa Haw. No. 12513.				
"	orthacinthal./f. ^a 0.	erect-spined tt.	ID gr	— M. Video — O s.p I., & O. dis. 18
*	arcuata Lk. & 0.	arched tt.	ZD gr	— M. Video — O s.p L.&O.dis.23
*	parvispina Lk.&O.	small-spined tt.	ZZ1 gr	— S. Amer. — O s.p
Cactus parvispina Haw. Na 12514., and 12515.				
-	O. intricate			— M. Video — O <>p L.ftO.dis.84
*	meonacantha Zir. lchs-spined			— Jamaica — O sp L.fcO.dii.15
"	OtfonuLk.	Otto'g		— Mexico 1829 O s.p
*	dC-n&aLk.&O.	dense		— Mexico 1829. O s.p
*	Déppnei Lk.& O.	Deppc's		— Mexico 1829. O s.p
*	sulcata Lk. Sf O.	furrowed		— Mexico 1829. O s.p
*J471.	MAMMILLA RIA.			
1^57a	fulvispina Haw.	brown-spined	■ ■ ■	— Brazil 1829. O &p
1^57a	CE'REUa			
1^57a	Haw. signus Haw.	large tt.	3 gr.	
1^57a	seniis Sain	old tt-ZJgr	i	SLDomin. 1829. C s.l
1^57a	«rox Haw.	ferocious «.ZDgr.		Brazil 1827. C s.l
Jog**	undulatus Haw.	wavy ttZDgr.		China 1629. C 81
Jog**	three-winged Haw.	three-winged tt.	ZD gr.	
Jog**	quamultaus&r/m small scaly		tt.ZDgr.	Brazil ... O gp
Jog**	equaceus Salm	bristly tt.	ZD gr	Brazil ... O s.p
Jog**	pnsmaticus Salm	prismatic tt.	Z3 gr.	Brazil ... O s.p
Jog**	coccineus Salm	scarlet tt.	ZD gr.	S Brazil ... O s.p
Jog**	extended tt	ZD gr.		R Brazil ... O s.p
Jog**	slender tt	ZD gr.		R Brazil ... O s.p
Jog**	bristle-bearing « ID gr.			Brazil 182a C s.l
Jog**	Ethiopian	tt-ZDgr.		Brazil 1829. C s.l
S9P	gW.^HVLLUM.			
oyi	Hooperi Haw.	Hooper's tt.	ZD or	2 jn W S. Amer. — C s.p Bot.mag.SG9S
oyi	Wctus Phyllanthus Hook.	but not of Haw.	2 jn	S Mexico 1829. C s.p
oyi	Ackermannii Haw.	Ackermann's tt.	ZD or	Brazil 1829. C 81
oyi	curled tt.	m gr		
oyi	ruhescent	«.rDgr	1	Brazil O 81
oyi	long-spined tt.	ZD gr.		Brazil 1829. C 81
oyi	glomerated	tt-ZDgr.		Brazil 1829. C 81
oyi	cylindrica Dec.	cylindrical tt-ZDgr.	3	Peru 1799. C 81
oyi	lericus cylindricus Haw.	No. 12578, and also 12576 belong to this genus.		
oyi	he body of this catalogue, several Chilean species of C&cetes have been referred to eenom frnm			
oyi	the lung state, without any knowledge of their fructification or habit when full growa			

M475. RHIPsalis.

28304 12630a cerefuscula Haw. small Cerew-ftfcftt. ZD cu - - - - Brazil 1829. C 81

1479. PHILADE'LPHUS.

28305 12640a Zeyheri Schr. Zeyher's ■ or 4 jn W N. Araer. - L co Sc. phiUe-

28306 . . floribiflndus Schr. bundle-flwd ■ or 6 jn W N. Amer. - L co Sc. phil. ic

28307 . . verruchsus Schr. warty ■ or 6 jn W N. Amer. - L co Bot. reg. 5

28308 . . grandiflbus B. R. broad-leaved ■ or 6 jn W N. Amer. ... L co Her. am. 28

28309 12645a specibus Schr. showy ■ or 10 jn W N. Amer. ... L co ScphiLic

28310 . . grandiflcrus Hort. and laxus Lewis's ■ or 6 jn W N. Amer. - L co

336a 1482a. ANGCPHORA Cav. ANGOPIORA. (Aggos, a vessel, photo, to bear; form of fruit) Myrt&cea. 2- ■

& - cordifolia Cav. heart-leaved ft I_J or 6 my.au Y N. Holl 1789. C s.l Bot. inag. ■

Metrosideros hispidus Sm. No. 12G79, hirsutus Andr. ■

& - lanceolata Cav. lanceolate-tod *I_J or 6 my.au Y N. Holl 181& C s.l Gae. fr. 1. ■

Metrosideros costatus Pen. No. 1268J. ■

S361. 1482ft. ACME^Nil Dec. ACMENA. (Acmena, a nymph of Venue's.) Myrthcece. 1"-2L ■

& - flonbiinda Dec. bundle-flwd *I_J or 6 jlau W N. Holl 1788. C s.l V ■

Metrosideros floribundus Sm. No. 12680. ■

1483. CALLISTETMON.

28311 12693a saltgnus Dec. Wllow4eaved * LJ or 6 my.jl Y N. Holl 1788. C p.l Bot mag. 18*

Metrosideros jaigngus Sm. ■

28312 12701a scaber Lod. scabrouB ■ kLjor 4 jl R N. HoU. 1820. C p.l Bot cab. ■

3362. 1484a. JOSSFNIA Com. JOSSIMA. (Not known.) Myrthcea. 1.-8.

28313 . . orbicular&tua Dec. orbicular-fri ■ - 6 ... W Bourbon 1823. C p.l

3363. 1487a. JAMBPSA Rum. JAMBOSA. (Jambos, aboriginal name.) Myrtacea. 6.-1* ■

t - vulgkris Dec. common il D fir 25 fil W E. Indies 178a C s.p Bot. mag. ■

-Eugenia Jambos L. No. 12732.; also 12745.12721.12729.12738. and 12725. belong to this genus. ■

3364. 1489a. MY'RCM Dec. MYECIA. (A surname of Venus.) Myrtdecar. 2.- ■

t - heria Dec. sharp II CD or 10 my.jl W W. Indies 1759. C s.p PluL ■

Myrtus aterris No. 12766; also 12768, and perhaps several of the South American Eugenias. ■

33G5. 14BQa. SYZYGIUM Gae. SVZYGIUM. (Xuxygiunja aboriginal name.) Myrticeof. 3.-24.- ■

k - caryophylliflbum Ztec. Clove-lvd 2 O or 20 ... W E. Indies 1822. C s.p Ru.am.** ■

Calypt&nthes caryophylliflbum Na 12773. j also 12771. and Eugenia 12741. belong to this genus. ■

1502. CERASUS.

28314 12863a serrulata serrulate-Zvo' i j o r 4 apmy W China 1822. O co

Prinus serrulata Lindl. ■

28315 . . salicina Willow-leaved* ■ - 4 apmy W China 1822. O co

Prinus salicina Lindl. ■

3306. 1503a. VU'RSHIA Dec. PURSHIA. (F. Pursh, author of FL Amer. Sept.) ■

28316 . . tridentata Dec. 3-toothed-leavedA or 2 ... Y N. Amer. 1826. C ■

Tigheea tridentata Ph. ■

CLASS XII. ORDER 2.

ICOSANDRIA DI-PENTAGY'NIA.

1506. HtATJE'GUa

12919 Oxyacanthia

7 punicea scarlet ■ or 15 my.jn S - - - B co Bot cab. 1 ■

• 1507. PY'RUS.

28317 12957a sinensis B. R. Chinese A or 4 my.jn W China - L co Botreg. ■

12965 jfria ■

3 undulata vravy-leaved \$ or 30 my W S. Europe - G co

4 rugbsa wrinkled/w* \$ or 30 my W S. Europe - G co

5 cretica Cretan \$ or 30 my W Crete - G co

6 bullata blistered-leavcd* or 30 my W S. Europe - G co

12979 arbutiflbia 2 intermedia Lindl. intermediate A or 4 my.jn W - - G co

3 serbtina Lindl. late ift or 4 Jap* W - - G co

28318 12979a grandiflbia Lindl. great leaved A or 5 my W N. Amer. - gS Botreg. 11**

28319 - piibeps Lindl. downy HI or 5 my W - - B co

28320 12980a depresso Lindl. depressed tt or U my.jn W - - L co

1513. COTONEA'STER.

28321 13006a laxiflora Jac. lax-flowered A or 4 ap R.w - - L co Bot fij. ■

28322 frifida Lindl. cold A or 10 ap.my W Nqial 1820? L.co Botreg. ■

28323 13007a UVa-ursi Hort. Bear.berry.lvd A or 4 ap.my W Nepal 1825. L.co Botreg. ■

microphyllea var. UVa-ursi Lindl. ■

* 1515. SPIRAEA.

28324 13013a vacciniflbia D.Don Strawberry-lvd* or 2 jlau W Nepal 1820. C p.l Bot cab. ■

* 1520. MESEMBRYA'NTHEMUM.

13177a lacerum Salm jagged JL LJ or | jn.au Pk C. G. H. 1811. C s.l Bot ■

dimidiatum Haw. No. 13177. ■

13298a alsinifolium Hort. Chickweed-lvd iQJ or J jl ... N. Holl? 1827. S s.l ■

CLASS XII. ORDER 3.

ICOSA'NDRIA POLYGY'NIA.

CLASS XIII. ORDER 1.

POLYA'NDRIA MONOGY'NIA.

peltata loc.t. peltate.fcwai Ai_ior 6 Trinidad 1827. C pl
 SANGUINA^RIA.
 grandiflora Rose, great-flowered A A or £ apjn W M. Amer. ... R pl
 GLAU'CIUM.
 tricolor Bernh. three-colored O*n 1 jn.s R Persia 1828. S co
 Persicum Dec. Persian O or 1 jl.a R Volhynia 1829. S co
 & ISTS! - ARGEMO^NE.
 " » ochroleuca Swt. yellowish white O or 2 jLau Su Mexico 1827. S co Sw fl car 245
 Milpurea, No. 1379&
 1555a - J.J.W.P.S / A Phu LEW? SI* - (MIP< M. Lewis, who accompanied Clarke to Rocky Mountains.) Crasstd. I.
 redivivaPA. revived % A or * ... W N. Amer. 182& D s.p
 Hunneman, a zealous botanist) Ppnverueca*. J.
 »* Mexico 1827. S r.m Sw fl gan 276

CLASS XIII. ORDER 2.

POLYA'NDRIA DIGY'NIA.

CLASS XIII. ORDER 3.

POLYA'NDRIA TRIGY'NIA.

		1599.	i>ELPHFNIUM.								
£8365	14133a	Menziesif	Dec.	Menzies's	A	A	spl	2	jn.jl	B	N. Amer.
£8366	14135a	altaicum	Fis.	Altaian	£	A	or	4	jn.jl	B	Altaia
										D	1826.
										p.co	Bot reg. U#
		* 1600.	JCONI^TUM.								
28367	14188a	Lamfrckii	Rchb.	Lamarck's	■	A	or	2	jn.au	Pa.Y	Pyrenees
		pyreniacum	Lam.	not L-						1817.	D co
		pallirinn	Nutt.	pale	■	A	or	3	jn.au	Pa	N. Amer.
										182a	D co

CLASS XIII. ORDER 4.

POLYA'NDRIA TETRAGY'NIA.

337a 1602a. ESCHSCHO'L TZ/i* Cham. ESCHSCHOLTZIA. (*Dr. Eschscholtz*, a botanist) *Papaverhcea.*
californica Cefwra. Californian A A or 1 jlo. Y California 1826. S co S

CLASS XIII. ORDER 5.

POLYA'NDRIA PENTAGY'NIA.

1607. REAUMTTRJJ. 28370 14311a *vermiculata* L., vermiculated tt- J pr 1 jLo Pk Sicily 182& C i.p.

CLASS XIII. ORDER 6.

POLYA'NDRIA POLYGY'NIA.

*161& MAGNCLTLA.							
S8371	14348a odoratissima	Aeinw.	sweetest scd ±	O or 10Jl.au	Y.w	Java	1829. 6 p.l
* 1629. i*NEM(FNR)							
88372	14430a Hudsonina	Rich.	Hudson's	■ A or \jn.jl	W	N. Amer.	1827. D co
S8373	- - Nuttallana	Dec.	Kuttall's	■ A or i jn.jl	W	N. Amer.	1827. D co
*1631. CLFMATIS.							
14460	grandiflora	Dec.	is Chlorantha Lindl.	Bot reg. 1234.			
£8374	14468a railacifelia	Wai	Smilax.leaved ft.	O un 20 ...	■	E. Indies	1824. C p.
• 1637. FLANUNCULUA							
28375	14629a affinis R. Br.	allied		^ A un my.jl	■	N. Amer.	1827. D co
28376	- - ovalis Aq!.	oval		k A un % my."	■	N. Amer.	1827. D co
£8377	- - Schlechtendollia	Hook.	Schlechtend.5	A un % my.jl	■	N. Amer.	1827. D co
£8378	14640a laciniatus	Bang,	fringed	O un 1 jn.au	■	Transylv.	1827. S co
1638. TRVLLIUS.							
£8379	146500 oconitifolius	Lk.	Aconitunulvd	■ A or 2 my	Y	1829. D p.l

CLASS XIV. ORDER 1

DIDYNAMIA GYMNOSPERMIA.

1648. *TEITCB.JUM.*
 28380 14698a *orchideum Lindl.* Oreohii-like «*•_or 1 jLau Pk Chile 1826. S co Bot. reg.
 heterophylous Cav. 125^a

3571.1653a. LOPH A'NTHUS Benth. LOPBAN. (*Lophas*, crest, *anthos*, flower; appearance of flower.) *Labut**
 2 S i " - wticiflvia Dm. Nettle-leaved ^ A or 2 jl.au B K Amer. 182«. S co ^,
 S8382 - - anatis *Benth.* Anise-scented ^ ^ or 3 jl.s B N. Amer. 1825. S co Bo f^#n
 Hyss5pu8 anistikus *Nut.* *discolor J5esf.* as well as No. 14772. 14773, and 14774. of *flyssopu8>*
 14812. of NÆpu8 belong to this genus.

1659. ME'NTHi*.
 1467a brevispicata Leh. short-spiked ^ u n I ... P Siberia 1829. D co
 *3372. 1659c APHANOCHRLUS Benth. APHANO. (*ApAanos*, obscure, *cheilos*, & lip lip of flower.) LfWoto. 1.-&
 *** - inclusus Benth. cut i ^ o r ! « W Nepal 1824. D co
 Jlfenthä blanda Lindl. but not Wat.
- 1659c. DYSOPHYLLA Blume. DYSOPHYLLA. (*Dysodes*, fetid, j>tetto» leaf i smell ?) Labiate. 2.—12.
 quadrifolia 2fc/A. four-leaved «, LJ or 2 jlau P Nepal 1820. C s.l
 - M'nha quadrifolia Box. No. 1*859.
 pfamila Benth. dwarf i A cu | jl8 " " . io- ~ ^ A
 frenthapnum GraA. verticillata Don
 Ifi63tf. -fORMPNUM Pers. HORMINUM. (Jfemao, to incite; qualities.) LaTM***: Jj
 - pyrenaicum Pers. Pyrenean £ A or 1 jn.jl B Pyrenees 1820. Deo bw. fl. gar. 252
- OD 1669 STATHY! 7
 } Fhirauta JftA. hairy £ ^ | or S ... R Moxico 1329. C pl
 germinica a pubescens ZindY. purple »? A *pl 2 au.o P Germany 1825. D p.l Bot reg. 1289
1669. SPHA'CELE &»A. SPHACELB. (*Sphacck*, Greek name for Sage.) LaMbia. L— 1
 " - Lindleyif Benth. Lindley's *Li_Jun 2 au Pjt Chile 1825. S co Bot.reg. 122b
 ftachys Salvis Lindl.
- 17& 1671a. ROY'LEil WaL ROYLGA. (*Dr.Boule*, superintend, of the botaa gard. at Saliarumpoor.) Labitx.L—3.
 * - - elegans Wai. elegant fe A or 2 jlau P Nepal 1824. C s.l
 •Ballta cinerea DXy» No. 15028.
- ^ If72a. BERINGE^R/^ Neck. BERINGERIA. (*Beringer*, probably a man's name.) Labitoe. a—9.
 - cincerea Benth. cinereous ^ A o* 1 Jn.jl Pa.P Spain 1823. D co
 Afariibium cinereumS<r. Na 150+*. 15048,15043,1504a 15045,15043,150-16,15047. belong to this genus.
- U ? flocky * _ or 2 jl.o Y Egypt 1828. D s.Lp Bot rep. 1300 ,
- * 1677a. CHASMO^NIA Lindl. CHASMONIA. (*Chasmao*, to gape wide; calyaO Labiate. \
 - inclusa Lindl. cuUeaved O or 2 jlau Pk Levant U93, S co Bot reg. 1244
 Moluccella spinosa, Na 15101.
1680. ORIGANUM.
 15H6 stoloniferum Bes. stoloniferous ^ A or 1 jnjl " " tan^o ^
 1681a. MICROME^RIA Benth. MICROMERIA. (*Mikros*, small, *meris*, a part) Labitce. 2.—?
 - Teneriffae Benth. Teneriffie <- J 1 jnjl P Tenenffo 1829. C co
 TTymusTeneriffie P^r*. as well as Na lol&. and several others.
- 33J-1685a. PHYSOSTEXGIA Jso/A. PHVSOSTECIA. (*Physa*, & bladder, *stege*, a covering; calyx.) Labiate. 5.
 > ' - virginiana Benth. Virginian ^ A or 2 jls R N. Arcer. 1683. D Lp Bot mag. 467
 Dracocephalum virgininum L. No. 15192,15193,15194,15195,15196, belong to this genus.
- ^1* 1688a. MOSCHO'SM A Rchb. MOSCHOSMA. (*Moschos*, must, *osme*, smell: odor.) Labiate. 2.
 ocytoides Benth. Ocytum-like ED or 1 jls W 1823. S co
 Lummitzera ocytoides Jac. Na 15237. and 15238.
- ^^•16886. ACROCEPHALUS27R/A. ACRO. (*Akron*, summit, *kephale*, head; flowers on branch tops.) Za&l.—3.
 ^ caputatus Benth. headed O ec lfjLau W China 1806. S t.\
 Lummitzera capitata W. No. 15240.
- ^J^ 1689a. CO^LEUS Lou. COLBUS. (*Koleos*, a sheath; united stamens.) Labiate. 2.—11.
 barbatus Benth. bearded tt-Oor 3 o.n B Abyssinia 1806. C Lp Bot mag. 203
 Plectranthus barbatus Bot. Rep. 6. Forskbhlfi B. M. No. 15251. and 15242. belong to this genus.
- *?*• 1689c. ANISOCHPLUS Wai. ANISOCHILUS. (*Jnisos*, unequal, *cheilos*, a lip.) iMbiUe. 1.—4.
 carnosia 5w/A. fleshy jg O or lfjn.8 Li K Indies 1788. S co Am. ac. 10.3
 Lavandula caribea W. Na 14828.

CLJLSS XIV. ORDER 2.

DIDYNA'MIA ANGIOSPE'RMIA.

8335. ISSS" raacrwrachya' Lindl. long-spiked A E3 spl 2 year S Rio Jan. 1825 C p.l BO. reg. 180J
 ^" x-329a rufila Lmdl. brilliant ? S & 2 au.s SY BrazU 1825. C p.l Bot reg. 1158
- BIGNOWi* amenaJFa. pleasing f O or 20. K Indies 182a C pl
- IjS!212^{1M81a}p aSBied O<pl 2 myjn Va Chile 1820. 8 Jo Sw.fl.gar.258
 - atropurpurea GraA. dark-purple O or 1*• P CoidiUer. 1826. S co Sw. fl. gar. 271
- 1715«CALA'MPELIS D .Don. CALASvtls. (*Kofi**, pretty, *ampelis*, a vine.) Jg>onrtctf». L
 schbra I>. Zlow scatfrous «L JV pr 6 jlau O Chile 1824. C Ls Sw. fl. gar.n.s.50
 &ccremocjrpus scaber Jl * J*. Na 1&488
- PENTSTE>MON. PulchraiusXim//. pretty / Al pr l.jn.au Li Mexico 1827. D p.l Bot. reg. IMI
 Chelbne «legans Kih. Pcntotemon elegans No. 15444.
 I i 2

28399	.	Scotileri Dou.	Scoulcer's	£ A or	3 my.jn	P.B	N. Amer.	1827.	D co	
28400	15447a	diffusa Dou.	diffuse	£ □ or	1) jn.au	P	N. Amer.	1826.	D p.l	Hot reg. 1245
88401	-	triphylla Dou.	three-leaved	£ A or	U jl.lau	Pk	California	1827.	D p.l	Bot reg. 1262
28402	15457a	glandulosa Dou.	glandular	■ A or	2 jnjl	Pa.P	N. Amer.	1827.	D co	Bot mag. 206*
28403	-	ovata Doti.	ovate/-eoi'cd	£ □ sp	4 jn.au	B	N. Amer.	1826.	D p.l	Bot reg. 127C
28404	-	speciosa Dou.	showy	■ A or	3 jls	11	N. Amer.	1827.	I) co	
28405	-	acuminata Dou.	acuminate	if A or	1 jn.au	P	N. Amer.	1827.	D co	
28406	-	venibliti Dou.	graceful	■ A or	2 jls	P	N. Amer.	1827.	C co	Botreg.I3U*
28407	-	pruinosa Dou.	frosted	■ A or	1 jnjl	B	N. Amer.	1827.	I) co	Bot reg. 12W
28*08	-	defista Dou.	blasted	£ A or	J	P	N. Amer.	1827.	D co	
28409	-	attenuata Dou.	tapering	£ A or	1 ijl.au	Pa.Y	N. Amer.	1827.	1) co	Bot. reg. 1295
S8410	-	glauca Grah.	glaucous	£ A or	1 jnau	Pa. Lin	N. Amer.	1827.	I) p.l	Bot. reg. 1286
S8411	-	procifra Dou.	tall	■ A or	1 jn.au	P	N. Amer.	1827.	D p.l	Kot mag. g*
28412	-	confirta Dou.	crowded-Jlw	£ A or	2 jl.au	Pa.Y	N. Amer.	1827.	D p.l	Bot reg. 12*
1718. CHELCTNE.										
28413	15460a	nemorosa Dou.	grove	& A or	1 jls	P	N. Amer.	1827.	S r.m	Bot reg. 12H
•1727. RUE'LLI/f.										
28414	15507a	Sabinidna Wai	Sabine's	ttflor	£ year	B	Silhet	1826.	C p.l	Bot reg. 1238
*1738. LANTA^NA.										
28415	15584a	m6Us Grah.	60ft	• *or	4 mr.au	R.w	Mexico.	1828.	C co	
1740. LITPIif.										
28416	1588a	dulcis Hurt.	sweet	£123 pr	1 jn.s	W	Trinidad	?1827.	D p.l	
1744. CHLOA^NTHESL										
28417	15622a	rosmarinifoliaCfrn.	Rosemary-lvd	■ □ □	2 jaaau	G.Y	N. HolL	1823.	C p.l	
•1749. fERBE^NA.										
28418	15638a	barbata Grah.	bearded	£ A m	1 jn.s	Pk	Mexico	1826.	S co	
28419	-	alata Lk. t	wing-stemmed	£ A or	1 jls	F	S. Amer.	1829.	D co	
28420	15641a	pulchella Swt.	neat	£ Ul el	1 jn.s	P	B. Ayres	1827.	S co	
S8421	15642a	chamaedrifolia&n. Germander-lvd	£ LAJ el	1 jn.s	S	B. Ayres	1827.	C p.l		
Mellndres Gill.										
1752. CLERODE^HDRON.										
28422	15691a	emirntase Bo).	Emire	■ O or	3 f.mr	W	Madagascar	1822.	C p.l	Bot. mag. J
1762. STENOCHrLUS.										
28423	15737	viscosus Grah.	clammy	■ J or	2 o	Y	N. Holl.	1824.	C ep	Bot mag. S
•1774 ANTIRRHINUM.										
28424	15779a	monteviddenseL*.	Monte Vidcan	O or	1 jl	R	M. Video	1829.	S co	
1775. LINA^RIA.										
S8425	15786a	asquitriloba Spr.	equal 3-lobed	t A or	1 jnjl	P	Sardinia	1829.	S co	Bot. mag.
S8426	15818a	purpur&cynafernA.	purpurascen	^ Q or	1) jnjl	P	S. Europe	1829.	S co	
1789. DIGITA^LISi										
£8427	15922a	laciniata Lindl.	fringed-leaved	^ A □	■ jnjl	Y	Spain	1827.	D r.m	Bot reg. 12 ⁰¹
3386. ? 1789a. ADA^M7J Wai. ADAMIA.										
88428	-	cycfnea Wai.	blue	■ CD or.	E. Indies	1829,	C p.l		
3387. 1811a. FRAISCI'SCEA Pohl. FRANCISCEA.										
28429	-	unifloraPoA/	one-flowered	A + or	3 jn.au	W.p	Brazil	1826.	C l.p	Soldnccp. 1.—1—
Hopdena Hook. Bot mag. 2828.										
1813. ALONSONM.										
28430	16071a	intermedia Lod.	intermediate	tt. I J or	2 year	S	hybrid?	... C s.p	Bot.cab.I	*56

CLASS XV.

TETRADYNA^{MIA}.

TETRAG. TRA. NAR.									
28431	16132a	pAtula <i>Grah.</i>	spreading	O un		my.jn	W	N. Amer.	1827. S co
28432	- -	<i>strictaGrah.</i>	strict	O un		my.jn	W	N. Amer.	1827. S co
1838. VESICA^RIA.									
28433	16241	ärctica <i>Rich.</i>	arctic	± Jf	pr		au.s	Y	N. Amer. 1828. S sp FL dan. ¹⁵²⁰
		A\$sum ärcticum	Fl. dan.	1520.					
28434	• -	arenbsa <i>Rich.</i>	sandy	£ Al	pr	£	au.s	Y	N. Amer. 1836. S sp Bot mag-
		j41y*sum Srcticum	<i>Hook.</i>						
184a DRA'B'A.									
28435	16292a	crassifJilia <i>Grah.</i>	thick-leaved	£ A or		myjl	£	N. Amer.	182a S 8.1
28436	16315a	gracilis <i>Grah.</i>	slender	O cu		jnjl	£	N. Amer.	1827. S co
		lutea ,3 16ngipes <i>Rich.</i>							
28437	16317a	nummulaxia <i>Lk.</i>	Moneywort-lvd	O un	\	myjn	W 1828. S co
•1863. JFIE/SPERI&									
28438	16429a	exclsa <i>Hort.</i>	lofty	■	A or	3	my	W	.
								 1828. S co
1865. SISY'MBRIUM.									
W439	1646 ⁸⁰	brachycarpumfccA.	short-fruited	O un	1	ijnjl	Y	N. Amer.	1829. S co
28440	- -	canescens <i>Rich.</i>	canescent	O un	1	jnjl	Y	N. Amer.	1827. S co
1867. URY'SIMUM.									
28441	16487a	exaltatum <i>Amir.</i>	exalted	^ CD	un	1	myjl	Y	Podolia 1829. S ro
28442	16493a	Marschall'hånumAndr.	Marschall'sji Q)	un	1	myjl	Y	Podolia	1829. S co

3877.	<i>oblongata</i>	<i>Dec.</i>	oblong	jg	Q	or	2	my	Y	Baical	182a	S	co	
2841:	-	<i>variegata</i>	<i>Led.</i>	variegated	f	Q	or	2	my	Y	Siberia	1828.	8	co
SeT ^A	-	<i>megacarpa</i>	<i>Fis.</i>	large-fruited	^	CD	or	2	my	Y	Persia	1829.	S	co
	-	<i>latisiliqua</i>	<i>Fis.</i>	thick-fruited	^	CD	or	2	my	Y'	Siberia	1828.	S	co
	-			broad-podded	^	CD	or	2	my	Y	Persia	1829.	S	co

CLASS XVI. ORDER 1.

MONADE'LPHIA TRIA'NDRIA.

atrata *Lod.* darkened 1 cu i my.jl D.P C.G.R . 0 e.p Bot cab. 1356
I97fo. SEATALA'NTHUS Swt. SPATALANTHUS. (*Spatalos*, delicate, *anthos*, flower.) *Iridace*. 1.
speciosa Swt. showy ff ^1 spl i jaau R C.G.H. 1825. O 81 Sw. fl. gar. 300
Trichon&ma monadelphum Swt. No. 1064.

CLASS XVI. ORDER 2.

MONADE'LPHIA PENTA'NDRIA.

28*30 R1EDLEV.M.
 borb6nica Dec. Bourbon tt.Oun 2 Bourbon 1829. C p.1 Cav. dis. 174.1
 28451 i!*⁹²³ PASSIFLO^RA.
 16898a Raddtfnna Dec. Raddi's j L O o r 10 jLs W Brazil 1825. C p.1
 338^b 1923a. DISE'MMA Lab. DISEMMA. (*Dis*, double, *stemma*, a crown: crown double.) *Passifibrae*. 2.—4.
 | - • Herberti^brt Dec. Herbert's £ d o r 30 jLau G.w N. Holl 1821. C Lp Botreg.737
 Muruciüa Herberti^bna Swt. No. 16925, and also 16926. belong to this genus.
 J225. TACSCFNIA.
 pinnatistfpula J. pinnate-stipuled j£ • or I r - - Chile 1828. C p.1 Cav. ic 5.428
 1927a. CLINTO^N/J Dou. CLINTONIA. (*De Witt Clinton*, late governor of the State of N. York.) *Lobelid.* L
 elegans Dou. elegant O pr * jLau B Colombia 1827. S co Bot reg. 1241

CLASS XVI. ORDER 4.

MONADE'LPHIA HEPTA'NDRIA.

M929. PELARGONIUM.

abutilSides Sw. ger. n. s. 2.
 acetabulsumSw. ger. n. s. 40
 acudientatumSw. ger. n.s. 36.
 alatumSw. ger. n. s. 25.
 anacamptonSw. ger. n. s. 61.
 amsoduntonSw. ger. 398.
 bipinnatifidumSw. ger. n. s. 62.
 calamiflratumSw. ger. n. s. 32.
 callistonSw. ger. n. s. 8
 carbassimumSw. ger. n. s. St.
 colophijllumSw. ger. n. s. 3.
 compactumSw. ger. n. s. 70
 commixtumSw. ger. n. s. 71.
 concretumSw. ger. n. s. 23.
 cordiflarmeSw. ger. n. s. 45.
 cratsgifulumSw. ger. n. s. 18.
 Cumming?VrSw. ger. n. s. 35.
 DarnlyiinuumSw. ger. a.s. 63.
 DebigrnheSw. ger. n. s. 20.
 decbrumSw. ger. n. s. 14.
 divergensSw. ger. n. s. 35.
 dissimileSw. ger. n. s. 60.
 eratulumSw. ger. n. s. 28.
 flabelliflriumSw. ger. n. s. 44.

Garden Varieties.

flagrans Sw. ger. n. s. 15
 glabrum^scens Sw. ger. n. s. 41.
 graphicum Sw. ger. n. s. 11.
 hannastictum Sw. ger. n. s. 17.
 heteromallum Sw. ger. n. s. 30.
 (hymnus?)? Sw. ger. n. a. 47.
 insciliptum Sw. ger. n. s. 65.
 Kenrichie Sw. ger. n. s. 58.
 laciniatum (ml.) Sw. ger. n. s. 2.
 latidentatum Sw. ger. n. s. 27.
 Lawrenceum num. Sw. ger. n. s. 22.
 Littlednium Sw. ger. n. s. 72.
 Loudonidnum Sw. ger. n. s. 16.
 magnifolium Sw. ger. n. s. 9.
 malacophyllum Sw. ger. 397.
 «talachra?» felium Sw. ger. n. s. 1.
 megalanthum Sw. ger. n. s. 3.
 mpgalclon Stir. ger. n. s. 4.
 megalostictum Sw. ger. n. s. 7.
 melanostictum Sw. ger. n. s. 5.
 mirabile Sw. ger. n. s. 57.
 nodutum Sw. ger. n. s. 68.
 nphilum Sw. ger. n. s. 19.

ntans Sw. ger. n. s. 66.
 Gbienidnum Sw. ger. n. & 10.
 ananthiftlium Sw. ger. n. s. 12.
 papyrdeum Sw. ger. n. s. 21. -
 prEclarum Sw. ger. n. s. 67.
 penasticum SIP. ger. n. s. 24.
 poramplum Sw. ger. n. s. 61.
 rctybula: Sw. ger. n. s. 42.
 picturatum Sw. ger. n. s. 31.
 plectophyllum Sw. ger. n. s. 37.
 poculifolium Sw. ger. n. s. 31.
 rhodopetalon Sw. ger. n. s. 13.
 saturatum Sw. ger. n. s. 33.
 succuktum Sw. ger. n. s. 69.
 suffusum Sw. ger. n. s. 43.
 tinctum Sw. ger. n. s. 29.
 tortuosum Sw. ger. n. s. 39.
 translucens Sw. ger. 395.
 urbanum Sw. ger. n. s. 46.
 vestiflum Sw. ger. n. s. 26.
 Wictorianum Sw. ger. n. s. 6.
 Yeatmann^num Sw. ger. n. s. 59.
 xingibdrinum Sw. ger. n. s. 38.

CLASS XVI. ORDER 6.

MONADE'LPHIA DECA'NDRIA.

17274a	<i>grandioips</i>	<i>Jac.</i>	large-headed	« C pl «» •»	R	Caracas	1829.	C	81	Jac. coi & 22. a.
28455	^JpW.	HO^VEil.								
17282	«	<i>apiculata</i>	<i>Cun.</i>	small-pointed	• LJ or	3	mrjl	P	N. Holl.	1824. C s.p
28456	T	T	« ^s COTT/i<							
			«ngustifoliaZiVutt	narrow-leaved	* LJ or	6	i***	Pk TN.	N. Holl.	1826. C p.1 Bot reg. 1268
28157	M	^*!* CROTALARIA.								
*387c	dichotoma	<i>Grah.</i>	dichotomous	tL uJ or	T 5	1	jl	Y	Mexico	1824. »"

8655. POTETRIUM.
£8550 23369a *errucosum Lk. warty & A un 2 Jl O . . . 1828. S co
2672. CALYDIUM.
28551 - - zamisfluum hod. Zamia-leaved £ E l c u 2 ... Y Brasfl ... R i p Bot cab. 1408

CLASS XXI. OBDER 10.

MON(E*CIA MONADE'LPHIA.

3416. 2693a. CA'LLITRIS Yen. CALLTPRIB. 'Kallos, beauty; appearance.) Conifer*. 1—2.
285/2 - - cupreB8if6rmiBr». CypresB-flnd. £ LJor 20 ... N. HolL 1826. S s.l
*27W. CROUTON.
88553 23804a rosmarinifoliaCtfft. Itosmary.JTd m or 5 jn.jl ... N. HolL 1824. C co
3417. 2715a. CROZOTHORA Neck. CaozopnoRA. (Not explained.) Euphorbiaceat. 1—1&
\$. - tinctoriaJ. " Jg «C-Qno O dy 3 jl W.o S. Europe 1570. S p.! Ac. pet 1712. " "
Croton tinctdn JP. No. 23806. * " " "
*2724. STEHICITLIA.
28554 23873a lanceolata Cav. lanceolate-tod 2 LJ un 20 my.jn Br.R China 1822. C p.l Bot reg. 1256

CLASS XXII. ORDER 7.

DICBXJIA OCTA'NDRIA.

- 3H8. 2793ir. RHODIH)LA L. Ron ROOT. (Rhodon, a rose; roots smelling like roses.) Crassutocae. 8,
C8555 - - rbseaL. Rose-smelling A or 1 rnyjl Y Britain moun. D co Eng.bot508
28536 - - sibcneAJforf. Siberian i* A * imyjl Y Siberia 1827. D co

CLASS XXII. ORDER 9.

DIGTCIA DECA'NDRIA.

2799. SCHrNUS.
£8557 24416 ferebinthiR)liaiJfl<f.Tercbinthu5JrdJ uJ or 20 . . . Brazil 1829. C s.l

CLASS XXII. ORDER 10.

DICFCIA DOODECA'NDRIA.

2806. MENISPE'RMUM.
28558 24435a planifoUum WaL flat-leared fl_Oun 20 ... Y E. Indies 1828. C p.l

CLASS XXIII. ORDER 1.

POL7GATMIA MONO/CIA.

- *2837. >fCA^CIA
98559 24740a pendula Cuif' pendulous * LJ or 8 ap.my Y N. HolL 182a C p.l
3419. 2873a. FENTACRY*PTA Leh. FENTAcYPTA. (Pente, five, krypte, a vault) UmbeUtfer*. t
28560 . . atropurpurea Leh. dark-purple £ E 3 un 2 ... F Mexico 1829. S co

CLASS XXIII. ORDER 2.

POLYGAMMIA DICECIA.

3420. 2688a. GALACTODE'NDRCN Hum. Cow THIEE. (Gala, milk, dendron, a tree.) Urtlcx. 1*
28561 - ' ttitle/fawi. useful 3 50 . . . Caracas 1829. S lp
3421. 2890a. SCRINDETA Thou. SORINDEIA. (Unknowna) TerebinthUcecc. I
■ ■ ■ madagascarie'nisis Thou. Madagasc.ft • o r 8 ... P Madagasc. . . C p.l
« 3422. W,ola. MEL ANORRKCE'A Wal. BLACK VARNISH TREE. (Melas, black, rheo, to flow; juice.) Tercbi*-
28563 - - usitJita Wal. common £ • us 50 ... R E. Indies 1829. C lp WaLpLas.ra.IV
• 2900. FIXVS.
28564 25271a luUscns Nois. Hitescent ft • un 3 my.jn Ap . . . 1824. C p.l

CLASS XXIV. ORDER 1.

CRYPTOGAMIA FI'LICEfc

2940. WOO'DSJA
38565 25633a PerrinViRA Torrcy Perrin's £ A el | jn.jl Br N. Amer. . . 1) I p



CLASS XVIII. ORDER 2.

POLYADE'LPHIA POLYA'NDRIA.

- 88186 *toS&i thea*6rn*»Cu**. 'Tea-shaped « U o r S ... » N.HoLl 1884. C s.lp
 2179a. ASTARTEM Dec. ABTARTRA. - - {Astarte, the Syrian Venus.} *Myrlhce**, t
fascicularis Dec. *fascicled* * LJ or 8. . . . N. Holl ... C & LpLab.ah.2.170
Melaleuca fascicularis Lab. No. 19627.

CLASS XIX. ORDER 1.

SYNGENE'SIA 

- , 2300. PODOSPE'RMUM.
 19803a *angustifolium* Led. narrow-leaved ^ Q> un 1 jn.jl Y & Europe 1828. S Co
 , 2201. SCORZONVRA.
 19630a *julia* Vest July & A un 1 my.jn Y Austria 182a S co

 belong to this genus.
 2203ft. AT AL A'NTHUS D. Don ATALANTHUB. (*Atalos*, soft, *anthos*, .ajower.) Comp. Cieh. 3.
 - *pinnatus* D. Don pinnate ftjvn 3 jnjl Y Tenenff 1820. S co
 Pren&nthes pinnkta L. No. 19915., and also Nos. 19914. and 19916. belong to this genus.

 2208a. OPORI'NEA D. Don OPORINEA. (*Oporinos*, autumnal; fæm of flowering) Comp. Cich. 1.
 - *autumnalis* D. Don autumnal £ A w 1 au Y Britain mea.pas. D co Eng. hot 830
Apirgia autumnalis IV. No. 1994a
 2211a. CALLIOPEM D. Don CAIXIOPEA. (*Calliope*, one of the Muses.) Comp. Cich. 1.
 - *aftrea* D. Don golden £ A or ft my.jl D.Y Italy 1789. D e o Jac.au. 3.297
Leptodon afireum L. -*Hieracium aureum* Seo. No. 19973.
 3399. 22116. JETHCTNW D. Don ^JTHONIA. (*JEthon*, one of Phcebus's horses.) Comp. Cich. 2.—a
 h frutiedsa D. Don shrubby «-i Jor 2 jn.jl Y Mndeira 1785. C co
JTieracium fruticosum W. 20072., and also No. 20144. of Crfpis belong to this genus.
 * 2215. CRE*PIS.
 20142a *croatica* W^K Croatian O un 1 jn.au Y Hungary 1829. S co
 SERPOLA.
 opargiBlides *Tineo* Apargia-like £ A or 1 ap.my Y Sicily 1829. S co
 gladica *Tineo* Glaucous £ A or 1 ap.my Y Sicily 182a S co
 libicans *Tineo* whitening £ A or 1 ap.my Y Sicily 1828. S co
 * 2227a. ACHYROTHORUS D. Don ACHYROPHORIJS. (*Achyron*, ch&ff, *phoreo*, to bear; receptacle.) Com.Cfeh. 8.
 I maculalus D. Don spotted, leaved & A un 1 jn.jl Y England ch.hi. D s.l Emr bot 225
 fTypochaeVis maculkta L. No. 20190., and also Nos. 20189. and 20197. belong to this genus.
 2247. CY'NARA.
 20422a *glomerata* Thun. clustered £ ^J or. i jLs P C. G. H. 1824. D co
 Stobea' *glomerata* Spr.
 2261a. CRASPEM3IA Font. CR*8PEDIA. (*Craspedon*, a fringes flowers.) Comp. Card. 1.—2.
 plebeia Cun. plebeian * L_I cu. . . . N. Holl 182a C s.1
 2269. KIPHNI/l
 20569a *rosmarinifl61ia&ft*. Rosemary-lvd£ iAi or 1 jLau W Cuba 182a D p.1
 linearifolia Spr. linear-leaved IAJor 1 . . . Brazil 1829. C p.1 An. mu. 12.61.1
Klcinia hnearifilia J. Jadmea linearis Pert.
 BPDENs
nacrosperma Fit. long.aeeded O or 1 jnjl Y Siberia 182a S co

CLASS XIX. ORDER 2.

SYNGENE'SIA SUPE'RFLUA.

- 
- ^ A un | Jaau Y.Br New York 182a S co
-
- 
- £iAlpr 2 my P.T V. D.L. 182a S sp Botma^SSSI
-
- 88800 20131a* SSSSS&tOsm. Genista-like • • » 4 ... Y & Amen 1824. C p1
-
- I i 4

- 2332. JERI'GERON. *SB501 21090a humille Grah.* humble ■ A Pr I au>* F Amer. 1828. D co
2333. TTJSSILA^GO. *28502 21106a purpurea H. K.* purple ■ J or 1 jn.au P C. G. H. 1824. D co
3402. 2335a. ADENOTRI'CHIA *LindL* ADENOTRICMA. ■ ■ ■ thrix, hair.) *Comp. Jacobhe.* 1.
28503 • . amplexicaulis L'tnd/. stem-clasping £ LAI pr 2 my Chile 1826. S co Bot.reg.ll!*>
- 2337. iTSTER. 21234a alpinus 3 ramboſus branched ■ A or } my.au B Europe .. D co Bot cab. 1829
- 234a CINERARIA. 28504 21457a vestita *Haw.* clothed ■ ■ ■ Y C. G. H. 1824. C pl
28505 21488a scapiflora *Herit* escape-flwd ■ ■ ■ Y C. G. H. 1829. C Ip
2347. GRINDE*LM. 28506 21543a coronopifolia *Leh.* Coronopus-lvd £ iA) or ij ji-s Y Mexico 1826. C l.p
3403. 2348a. STYLCLEPIS *Leh.* STYLOLEPIS. {Stylos, a style, lepis, a scale; form.) *Comp. Card. Vern.**
28507 21547a ■ ■ ■ *Leh. slender & A pr 3 jLs Pk N. & W. 1826. S co Sw. fl. gar. 285
lepis gr&cilis Grah.
3404. 2354a. N&JA D. Don NEJA. (Without meaning.) ■ ■ ■ Aster. 1.
28508 . . gracilis D. Don slender jg. A) un 1 au Y Mexico 1828. S co
3405. 2360a. ERIOPHYLLUM *Lag.* ERIOPHYLLUM. (*Erion, wool, phyllon*, a leaf; woolly foliage.) *Comp. Helta.* 1.
28509 . . caespitiflum *Lag.* turfy t A or 1 my.jn Y N. Amer. 1826. D eo Bot reg. 1167
Trichophyllum lanatum Nut. Actinia lanata Ph.
2363. DA'H利亚 change to GEORGVNA.
- * 3406. 2363a. GEORG!il W. GEORGINA. (J. G. Georgi, a Russian botanist.) *Comp. Helianth.* 4.
28510 - - crocata *Lag.* saffron-colored A or 6 jLn S Mexico 1802. R pl Sw.fl.gar. 282
Dishlia frustacea 2 i x a g e i, No. 21592.
- 28511 . . CervantéBii Lag. Cervantes's ^ A or 7 n S Mexico ... 11 pl Sw.fl.gar.n.s.4
2365. r A G E r ^ ■ 28512 21595a florida Swt. florid JI.AJor 1 jl ■ Y Mexico 1827. D co Sw. fl. gar.n.*3*
2374. CHRYSANTHEMUM. 28513 21649a rutheniacum *Fis.* Russian k A or } jn.au Pk Russia 1827. D co
- 2409. HELIA'NTHUS. 28514 29134a ovktus *Leh.* ovate-leaved O or 4 jn.o Y Mexico 1829. S co
& 515 21935a lenticul&ris *Dou.* lenticular O or 6 au Y N. Amer. 1827. S co Bot reg. 285
28516 21961a patens *Leh.* spreading k A or 3 jl.o Y N.Amer.? 1829. D co

CLASS XIX. ORDER S.

SYNGENESIA FRUSTRANEA.

- 3407. 2411a. ERICOMA *Kth.* EHIOCOMA. (Eriom, wool, tome, hair; p ■ ■ Comp. Hel. 1.
28517 . . fragrans D. Don fragrant i j o r 3 i W Mexico 828. C co
- * 3408/ 24116. ECHINAX3EA Moen. ECHINACEA. (Echinax, a hedgehog; receptacle.) *Comp. Hel.* 3.
^ . . heterophylla D. Don variable-?d ^ A or 1 J s o P Mexico 1829. D eo Sw. fl. gar. ■
Coreopsis heterophylla Cav. and No. 21973. and 21979. of Rudbeckia belong to this genus.
2415. COREO'PSIS. h 21997a amplexicaulis Cav. stem-clasping ■ A un 4 jl.au Y S. Amer. 1800. D pl
Simsia amplexicaulis Pers. No. 22019.

CLASS XIX. ORDER 4.

SYNGENESIA NECESSA*RIA.

2436. WEDETLIJ. 28518 22235a afirea D. Don golden A-AJor 2 s T Mexico 1829. S co
3409. 2436a. ZEXMETNU Lai ZBXMENIA. (An anagram of Ximenesia.) *Comp. Hel.* 1.
28519 . - tagetijibra D. Don Tagetes-flwd £ j \ | o r 2 s Y Mexico 1829. S co
2448. CALE'NDULA. 28520 22272a mexicana *LJc.* Mexican O or I jas ■ Mexico 1829. S co

CLASS XIX. ORDER 5

SYNGENESIA .SEGREGA'TA.

5410. S*7S<.PODA'NTHU8log. PODANTHW. (POW, a foot, anthos, a flower; stalked.) *Comp. Card. Ech.**
■ 521 - - Mitt)qttiUnd. Mitiqui ft or 3 au.n Y Chile 1824. C pi

CLASS XX. ORDER 1.

GYNA'NDRIA MONA'NDRIA.

- 8488 (TPHRYNS.
 22520 aranferra
 2 limbata Lindk. limbate * . Alcu 1 ap.my Br Rome 1826. R s1 Bot reg. 1197
 2°lon^ealaLiiMif. long-petaled A IAI cu 1 ap.my Br Rome 1826. R Lp Bot reg. 1189
 ^ S ^ i n S K S 1 W great-flowered ^ El or 1 jl.s W StVincent 1829. D pl Bot. mag. 2956
 2528a. GASTRO'DIA B. Br. GASTRODIA. (*Gaster*, a belly, *odous*, a tooth: top of column.) *Orchidca*. I.
 - - fesamoldes R. Br. Sesamum-Uke £ E] cu \ ap.my W N. Holl 1826. D pr.w
 12537. MAXILLA'XRIA.
 Oa ciliata Lindl. fringed £ El cu i my G S. Amer. 182R D pr.w Bot reg. 1198
 squalens Hook. Bot. mag. 2955. is Xylbium squalens Bot. reg. 1897. No. 22664.
 12539. PLEUROTHA'LLIS.
 Xa prolixa Herb. proliferous £ El i jl R & Amer. 1827 P D pr.w Bot. reg. 1298
 82714a" anceps Lindl. two-edged-friffig E) cu * year Y.o Bengal 1826? D pr.w Bot. reg. 1239
 am. " pygnse um Cuiu pygmy £ El cu \ N. Holl. 1825. D pr.w
 Ssyx^L" Beccrndm Zhnfc secund 2 El or *jnjl Pk Sumatra 1828. D pr.w B-t reg. 1291
 7^A^S a chrysanthum Wai. golden-flwd J H or If Y Nepal 1828. D pr.w Bot. reg. 1299
 2547a. CERATOCHILUS Lod. CERATOCHILUS. /K"TM>*£om*£oil%>?X\|L^A^
 - - grandiflorus Lod. great-flowered £ El cu 1 JLo P Tn^n!dad *% ^ K'I Snt m< S<
 - - insignis LifioV.; noble ^ IS or 1 jl.o P Tnmdad 1826. D pr.w Bot. mag. 2948
 Stanhopea insignis Hook.
 2553 CATTTT^
 »728a" intermedia Gr'ah. intermediate «E I or 1 ap* Pk Rio Jan. 1824. D ^ 5 ^ - 2 8 5 1
 - - crispa Lindl. curled-lu>d £/*& 1 au W.Pk Rio Jan. 1826 D p.r.w Bot. reg. 1172
 2553a. EKYCLIA Hook. ENCYCLIA. (Ekykleo, to wrap round column byaip.) OrcMdet. 1
 - - green-flowered £ El cu 1 f G Rio Jan. 1827. D pr.w Bot. mag. 2830
 22768a tuberculata Hook. tubercled £ E3 P* * JI W. Y Bontaf. B. 1827. D pr.w Bot. mag. 2878
 2555. MICRO'STYLIS.
 ?ieS?i?«iM. Mexican *. Alcu ftjl. G Mexico 1829. R pl Bot. reg. 1290
 eAtt& Lindl. taU ^ El cu 2 jlau Br Rio Jan. 1826. D p8 Bot. reg. 1175

CLASS XX. ORDER 2.

GYNA'NDRIA DIA'NDRIA.

- CYPRIOPDIUM:
 S^gg ventricosum Swz. bellied tf A or f ... P Siberia 182a R ip Sw.fl.garns.1
 macr&nthum Sun. long-flowered tf A or f ... P Siberia 1828. R s.p Bot mae. 2938
 guttatum Swz. spotted tf A or f ... Spot Siberia 182& R s.p Am. rut'22

CLASS XXI. ORDER 4.

MONCE^CIA TETRA'NDRIA.

- S^612a. CONOCEPHALUS Btume CONOCEPHALUB. (*Konos*, a cone, *kephale*, a head; form.) Urticeq 1-12.
 * naucleiflbrus Lin**/. Naudea-flwd » Q c u 10 year Y Chittagon. 1820. C pl Bot. reg. 1203
 l/rthca naucleiflbra Rox.

CLASS XXI. ORDER 6.

MONCRCIA HEXA'NDRIA.

- KU'NTH/^ Hum. KUNTHIA. (C.S. Ktn*h, an acute French botanist) Pahna. 1.
 - moutana Hum. mountain i CD or 10 . . . N.Grenad. 1829. S r.m

CLASS XXI. ORDER 9.

MONCECIA POLYA'NDRIA.

- bloody tt.Oor 1 . . . Brazil 1829. C l.p
 ever-flowered n. • or 2 ap.s Pk Brazil 1829. C Lp
 vanous-leaved £ El or 1 o Pk Mexico 1829. C Lp
 papillose B. D < w 3 ap.s Pk Brazil 1826. C l.p
 remarkable < • or 2 d Pk S. Amer. 1826. C l.p
 villous EQ w i Jl W. & Amer. s Lp
 bulb-bearing £ i_Jor 1 jlau W.Pk Peru 1827. C U
 rah. two.petalfd < • or 3 aps Pk Bombay 1827. C Lp Bot. mag. 2849

17MI eurojV*
| rt.r. J.I*;'''

i.r. 0 •Pjr, V JlriUln 8 oo

HW. CAXAVfLtA.
» * » HOMa botwrUMB U*4t, rttttM* Ar m o l (it 10 jn.Mi P B. AT ret ... C Lp Botntl

f.t'Pl'SU*
177WU mirrinlliu. Oo
nm&lui />DH.
int.itiii. /;*.
H4ffl I plut^«i« Aw.
flWH I was**** **
WMU 17717- rhanrwllwl

dMMe.llovmd 1 O ... 1 mijjl P.I s AMMt.un s co Bot.rt!'
1 A i.r. 1 u X l't i ;,..n.li.» lh.ti co I
1 MIL* (öttiintim UM en
1 n 01 1 In.a it n s co lk*, ft*
1 A M i CaUJtanUlKVti ti Co
ur J n.jl n.ia ... a co
or S nr l'k.ii s co
J <ir S Vil Pk.w .. co
J w 4 it n Aym UM b». B. Ju

>u XVI. Om>8,

I.1.IJ1IA POLYA-NDRIA.

* « 01 JTA'I.VA.

rrnu HUM am, dLTHJCA.

:a n&M SM*T» H iMp ii i o or 5 JLU P MA 8 <

17WU I hftc

ITB Mi rnoui

hybrid

k A or 4 H

Tk

M.iuritimi!i<<.

{tot. m

Sw. n

7i liOOtn glotur
i*» MwUIS&ra Hook

nf 3 n

W

7. C 1.p Rot
C to Hot.

3391. HKBVfc-SM Lin.ll JtcE<r.it (Join ller*ti <

IWL C p1

CAMVJXM

JipAnca

CLAM XVII. Ou u3.

TA'

larjtf

3 my.Bii I' C.O.H

CI.*MXVII. O<>r.ii A.

DIADE'Ll'IN I Ht I A.

• I J- A <" (ft* V

a at i ln.au Y N, Amcpr. 1W4. 8

a i e t; \ i i

li

TM

29IX DESMCTOITM.

• •or jLul P K IndiM 18M C t.p Bnt-rw*¹

• 81.HI LATHYRUH.

1WA ffii&.ui J/«

1 & or 4 injl W.« IVntwylv, ... i> i i

UBtti

(UpuUcaotu

J- A «

I

P

SlbMtff'

...

R rn

N.Ai7iw.r...

Deo

. CUM. No. IU387. «nd HBW. rtould t» pUeed »n « fubtefiwU, Na IMTtt

In tw> than a tenth orthewe spaOM. In Awonkwwhich will ihf.rty appear *, will be found • *•**
of plan* for h̄erbwoutjroumU and arbor' neox|KMcd hi tin- nutirw defcrilxil, a»| Tur |>lantinK nil i—
initadl ..! | M idMM fti i urdiik 1' the N itural Kjrt* .
n wa ammen U o n i thoipeCjt < insin ib iker?ffii mM.T. >:::LOOW • — my.n i i w
U» bOtkliC lltiUloCUO. ii , th . . M . ' " in. Tl.int ami varied 11. , rr.i,
I'uHure, regftabtp gaosraph"
we preturao to be to oUiou* t> every gardtnrr arid botanut at not to require explanation

INTRODUCTION TO THE JUSSIEUEAN, OH NATUKAI-, SYSTB

MOO, VAS(

(t with npiral'
flJ ip.

DICOTLUKf XU: WM, tWO,

SuMiriiion f. Wchlamyds.r A*, two,
calyx ami

claw I. Ttialjmtabrw *fi/wliimui*, a bed,
men*underpiihllura).—1. [tain-
dMt, 7. I'oduphyllir™, 8. t!ydn>(*ltnlnr.
SuMII 9. Calrriflto* frity* and jlm ; •*-
itCJT, 01 r, r w, SS.
uiithacoL¹, 87. PCKun

flan 3, Cornilm'.n- 'evrotla Kivijtotj •!<
— 117 Kj iDutfticv, ll'i.
•In'iip, 191. >tii rax-iiftcca ,
US..I „cal, ii-,to IAI

Subdlrlriioti II •ifrUi*
I Dorolla

IVJ. Aina.
•i m .
JWionri Ctua,
ITU. IttiTuulijirtili'ir, IKi Ali>
Juic>(tin.-ti-. EM OrciSdme, he..

IQnd Diviitm, < i MM
tiMUC Olli)

1. FlUet*. tt, KqulMMeam, 3. Lycopodlnc*, Ac

CUM, APIU'LUI /a, priv.,
7. A'Ktt, R, Lichenc*

and chlaatfii. a coat w Dovarlur j <* .w^S

yicv, a flower; nit.
t iillenlftcM!, 1

t m E] J

tmtni on fln<

ni4.i-i,

1

5Et

17DJ

DP. r/i/umi/tt a coat.j

t f Art

- *{*. @a

(tnuidj, one, anil koyhiliin

I. BuliV „ T/

•i. a illtr I

;/i<<tr^ui, iraiy; habi¹.

j

Ithyltvn, a leaf) IMSMI).

1 111: li.iif, l'„< cMitUK-tnl mi'.

At the

MMnL... dkM can In bnw
H probably upawnri, or of it


• illiistr.iifiMi i.i l,iitiiniM|
■ Eurui*, and in Nr.rtJi An

rden Archill

In Imperial iiiid[u.(i qiuuru-rl}, SI <


" / i hi h; * ilfscovend ih*it hLi tilanl tiMunrt to RuUicOa

their general stnulu <> pmuJ ifivUioiM c*ll
 .>.n riiri.>ti.l. ~1 >..L. ILrl Aa*hdhM: tin. VUCU
 it. Celluli



{that it, flowers funuit)et] either with ttntnlhu, or pMtlUum, or both), which will slway* prevent their I with the >ijfh<< ti ilif> of Ct'lilujre*.

Slonucotylfdimius arc Che first remove frum CuilUres, and hold nil iricrmittliate I boiwfim them ami EI^III »*** or >>>itt/ed(tmuu plants, in whiel vegetation acquire* lu highrat Conn oft' lopemuii, Tlmy were formerly eha raft Priced by" having it jingle cotyledon, hut this circumstance It Of* not abnuutu, but difficult >r ricti'rroination, accept after minute aiply»U, The real iifflrlinj't in the them and Dicotyledon! U thi, that in Monocotyledon! there it only nuc cotyledon {Jig. 3. t)j or, if 1




iuu oT • mcmocotjli item

. Section of (tnnnluuind tnibrjni of * m i . Is ihow t

f, HKton of ditto, to 4how tww tatstmiaa immatctng In lh* nprwnt thMt, tww middle lob* utteiiluinula.

they are tiltmtato with i-ni-h other (), while in Dicotyledonl they are nlwayi opposite, and more than <"jj MiltM iteveral, nt in /' Unix (ff. 4. y). The pli of tin- tun chbac* it, however, it'! Lj whii li they are (tmili their lir.t irijlii'ipk'. thr toed from which they origuittv. In K ifonoculj/i-ilmti, than > no dWinettm balHIWJ trOod .tul 1Mrk y/ '}./*; in / wood mill bark are dUtingt] cftutntd 'Jig.i >. In HonocotThuoof the wood nud CLJul.it thucare t d l • without any dritinet winuhi lnyi-n off the furmr HHHK in Dicotyladonl tht wood cellular ti>tuc h n ato their >irticulie limiu uMigne-d them, a distinct layer of the former betas dffpottdad. In Monocotyladotu than ate no radiation* oedula lu tin- 1Mrk ; in Dicotyledon i,hinEuiuN an ih^inicy nitirli'd. In Molineotyedcni! therti '^ ^i^i<-rilly ni> artiruLiti'li U-iut'^ii il .1 u I 0)g stem, winc* in Dicotyledon! Uli lening n tear behind. In Mono tlieyetm i I >aH in parallel QHM from tin- a>ex, in Uicollcuone they ilwi'KL- ii'tmi the inilril tuwurds tlir margin ut vrious .siujh'; in the fuiiH'r iro unbrsmeheo, the principal veins 1' sted bj m-ary itniple secondary velni; in tin- Inner t(** much britiuttc, ramifying in many directions, and ^iviu^ the sumTM of the lca! a rjetlfd a>iiranace.



B, T n i m i H lection o* • dkotjUdeoou tUm
it, An unbkt with two rotjlwinu.
a, An mibrjo wlUi tbor cotjlsfaM*.

a, An *mbrro with mm; cotylidou.
x. Stem and Ua*n < a dloMhdamtu illani.

Such ore. lh> very obvloui dtitnctionj of the two great cbufet of phanogamjgs, or law to far U tt frdm && w bring any neccawly for dUctedLnf! a >seed In order tu atevrtin lt.

i one it the IIUKI md about which there cannot U in niic HK ".
tymnae & M. Coobtobtordifficility. It hi almc* Impoutble tc il mriMcl of a plant without U
iti ptk'M>n.

HEM • i ir lieave we :n
reeof fenowledgooraptdlwtion li regidatte than what ou^ht to lk< pen re>j a
y p ...l upon the inn

y the utriuin mul (HWihiu u(he *•
tttOtgh SKIH ;escher- i Omit: iiii the nature of the ti*Jiji;< and Itta' TH* '•

he chataeten < in thi plaf, neither li It uoMMurf to advance farther in nn esp»TM
Qio niEiiiu-i of di ilu'in; upon Unit point each order >roala, roqulw i particular note.
-ity lh'licvefl, th'hoH! that tum b an already retrnovud ; mid thuit olltonijh llllnaT »ic **

"the list of urnen under each order. In the fullwinK arrangement, the column or figure* shows the number of hiwdj li Mtdj hertMueoiM ipLties and varletie*, beli-
»at've Ijritih ipeciet, a lUr (*) is prefixed to II- ro low. Where a genui includ..

FutST GRAND DIVISION, VASCU.IA IM-S.

i 3040, *Spueit* S77S"; IM.hmat .*
Ihirilij Iwract'ouii Sj .

I. DICOTTLEDCTNBJB, OR EXO'GENJE fo», ouisi.lf, gtfnumoj, to grow).

*Qotra £+07, Specfet 5.10CT; HoLhout,- Spt ek*4S ii, //nufy/ Uaneou**
^MOfttMR] ILinly ILibticrotts Specie >®n. f 4HJ feet; £ 214 |.*

SUBDIVISION L DrCHILAMY'DI

livliion comprehenilti all the Dicotyledonous planttj that have both a cilix and urtilla, by which ulshed tripui & deir, in which thr> calyx only exiits. It i. In eon of this

LASS I. Ti ALAMI \ l. < yRJE. Petals and Stamens mierud in the Btxptaett.
« 517 f 74 UU 1st In W. 3

insertion of the petal* and stamens into the receptacle is the RIMT character of this cubrains, which, g. «brcont» in all the polymidrous plant* of Linnseu*, a* the Calyciflurc contain the ioeandrous gen

SXCTION I. *CarpBa* numerous, or Stattmis oppotite the Petals[^]

1. OBBBRI. KANUNCULA*CEJE.

i 29, Specif* *ffll; Iht.house Speciet ft; ffntm tour 8p <i^ 10] Itarify l.fsrneota SpecitiSi
I fartjy Herbaceous Species 561. f LSfeet) f*

uf ihp R^eres^r P⁰¹^A of the plant of this order U I'nterwl wish gardenen, contjiunnjr. * ii <•
nost^r eleg Bnl or thaw; *! the trilir^r U^r limdry plutiti. It in here fdil the (ir.n. ; flic luvly
mbliiMby ti'd charactei of tin i mi. tin re T, that thi
jy > nnn, .. planb i neuv*i* |^{1***1} i* in il;jn •' that
J-| mi' ; tint ir> in honey,
in n
in liu Mitt 'H.*

7h1lr, filch grt^c
Ir i<tic ; i
pep^fH^gt [in] s'1d III If...]
gratt number have t.

Europe, but they are so abundant in all parts of the world that an order can scarcely be found more universally and equally dispersed. It is singular, that, with the exception of the climbing species of *Clematis* and of *Xanthorrhiza*, scarcely an instance occurs in Ranunculaceae of a shrubby stem. All the perennial species may be propagated by division, and those which are woody by that mode or by layers or cuttings: the annuals and most of the perennials bear seeds freely.

2. Tribe 1. CLEMATIDEJE. lig. 31. herb. 7. £9 ft.; £9 ft.	4. Tribe 3. RANUNCULEA. lig. O. herb. 131. £ 18 ft.; $\frac{1}{2}$ 3 ft.	1639 <i>sopfum L.</i> 0 2 1392 <i>Garidilla TOIL</i> 0 1 1606 <i>Nigella Tou.</i> 0 14 1601 <i>Aquiltgia /.</i> 0 21 1599 <i>Delphinium Tou.</i> 0 *61 1637 <i>/ianftnculus Bauh.</i> 0 *126 1636 <i>Ficaria Dit.</i> 0 *2 1630 <i>Ceratocephalus Moen.</i> 0 2 1635 <i>/ianftnculus Bauh.</i> 0 *126 1636 <i>Ficaria Dit.</i> 0 *2
3. Tribe 2. ANEMONES lig. O. herb. 12*. £22\$ feet	5. Tribe 4. HELLEBG*REJS. Ug. 59	6. Tribe 5. P*O.VIAYEM. lig. 12. herb. 4& f10\$ ft.; £ 20ft.
1633 <i>rhalictrum £.</i> $\frac{1}{2}$ 59	1643 <i>Caltha /.</i> 0 *8 1638 <i>Trollius L.</i> 0 *10 1640 <i>Eranthis Sal.</i> 0 1641 <i>/elMborus L</i> 0 $\frac{1}{2}$	1604 <i>Cimicifuga £.</i> 0 4 1546 <i>Macrbtys Ilafl.</i> 0 1 1545 <i>Act's'a L.</i> 0 *3 932 <i>Xanthorrhiza Ilceril.</i> 1 0 1596 <i>Pasbnia L.</i> *J1
1628 <i>Hepatica Dil.</i> $\frac{1}{2}$ 45	1642 <i>Coptis Sal.</i> 0 $\frac{1}{2}$	
1645 <i>Hydrastis L.</i> 10		
1635 <i>Knowltoni Vi Sal.</i>		
1634 <i>Adonis L.</i> 0 *10		

7. ORDER II. DILLENIAXE-K.

Genera 12, Species 29; Hot-house Species 11; Green-house Species 18; Hardy Ligneous Species 0;

Hardy Herbaceous Species 0. f 0 feet; £ 0 feet j $\frac{1}{2}$ 0 feet.

Fine plants, almost exclusively confined to tropical countries. *Dillenia speciosa*, a native of India, is a most noble tree with large yellow flowers, rivalling those of a Magnolia. *Hibbertia volbilis* is a green-hued plant well known for the beauty of its blossoms, and their powerfully fetid smell. The medical properties of this order are scarcely known; a decoction of the leaves or bark is astringent, and used for gargles; and the acid juice of the fruit of some of the species of *Dillenia* is used in India, mixed with water, as a pleasant beverage in fevers. The foliage of many of the species is extremely scabrous, whence the dried leaves are used for the same purposes as fish-skin and sand-paper in Europe; those of *Trachytella aspera* are even employed in China for polishing works of metal. Most of the species strike readily by cuttings of the ripened wood in sand under a bell-glass.

8. Tribe 1. ■■■■■	9. Tribe 2. ■■■■■	1610 <i>Würmw Rtb.</i>
1609 <i>Tetracra L.</i>	1376 <i>Pachynema R. Br.</i>	1608 <i>Colbérta Sal.</i>
1584 <i>Doliocarpus Sol.</i>	1595 <i>Pleurändra Lab.</i>	1614 <i>Dillenia L.</i>
1583 <i>Delima L.</i>	2185 <i>Candölca Lab.</i>	
1594 <i>Curatilla L.</i>	1597 <i>Hibbertia Andr.</i>	
1603 <i>Trachytella Dec.</i>		

10. ORDER III. MAGNOLIA*CE£

Genera 5, Species 37; Hot-house Species 3; Green-house Species 11; Hardy Ligneous Species 23;

Hardy Herbaceous Species 0. 1 62 feet; £ 0 feet; $\frac{1}{2}$ 0 feet

them is said to have a bitter flavor without any astringency, and combined with a hot aromatic prime. In the United States, the bark of *Magnolia glauca* and *Liriodendron tulipifera* is employed for the same purpose as Jesuit's bark, and from the fruit of *Magnolia acuminata* a tincture is prepared which has some reputation for removing attacks of rheumatism. The fruit of *Ilicium anisatum* is the material which flavours the liqueur called Amisette de Bourdeaux. The Magnolias are exclusively inhabitants of Asia and America, no species having hitherto been found either in Europe or in Africa. The shoots of most of the species do not root freely, and therefore they are generally propagated by layers, grafting, inarching, and sometimes seeds; two or three of the species by cuttings.

11. Tribe 1. ILLICIE.	12. Tribe 2. MAGNOLIE.	1618 <i>Magnbifl L.</i> 21
1615 <i>Ilicium L.</i>	lig. 23. herb. O. \pm 62 feet	1617 <i>Liriodendron</i> 2
1602 <i>Drimys Forst.</i>	1619 <i>Micheliifl L.</i>	

la ORDER IV. ANONA*CE£

Genera 9, Species 45; Hot-house Species 40; Green-house Species 1; Hardy Ligneous Species 4;
Hardy Herbaceous Species 0. f 11J ft.; £ 0 feet; $\frac{1}{2}$ 0 feet.

The plants of this order are closely allied to Magnolaceæ, from which they are principally distinguished by the absence of stipule, and by the structure of their anthers and seeds. The latter consist of a hard maw albumen, ruminated, as the botanists call it, that is to say, perforated by the substance of the seed-coat, every direction. They are all trees or shrubs, and chiefly inhabitants of the hottest parts of the tropics. A few have been discovered straggling into the temperature zones of America. The fruit of the *Aniba* is many species highly esteemed as an article for the dessert, especially that of the Cherimoyer, which has a reputation of being the finest fruit in the world, next to the mangosteen. The hard fruits of the species of *Uvaria* are highly aromatic; those of one of them furnish the ipecacuanha of the shops. The *Asimina* is the only one which contains any hardy species, and these are so delicate as to be seen only in this country. In Brazil, the bark of *Xylopia sericea* is used for cordage; for which it is admirably adapted. Cuttings or layers.

1622 <i>Anona Adan.</i>	1626 <i>Asimina Adan.</i>	1624 <i>Artdbotrys R. Br.</i>
1623 <i>Monodora Dunal.</i>	ISA) <i>Uvaria L.</i>	1627 <i>Xylbia L.</i>
1538 <i>Eupomatia R. Br.</i>	1621 <i>UnonaZ.,</i>	1625 <i>Guattifia R. & P.</i>

14. ORDER V. MENISPERMA*CEJE

Genera 8, Species 29; Hot-house Species 20; Green-house Species 5; Hardy Ligneous Species 4;
Hardy Herbaceous Species 0. £ 4 feet; £ 0 feet; $\frac{1}{2}$ 0 feet

The order of Menispermaceæ consists entirely of twining shrubs with minute flowers. They are dissimilar in habit from the orders which are placed near them, and occupy their present station on account of certain minute but important characters in their fructification. With the exception of *J. J. la arccocci nea* none of them are worth cultivating as plants of ornament. The berries of *Lardizabal* are sold in the markets of Chile, under the name of *Aguilboqui*, *Guibogui*, or *Cogull-Vochi*, according to the travellers. The bitter, diuretic, and aperient roots of *Pareira brava* are the produce of the *Cissampet*.

NATURAL ARRANGEMENT.

4D?

aiuf i * and the famous: Col iiribn root, to much ertcemod for Its Intense bitternes-t. and for its use in diarrhoea
 illini y. the produce of the Cocettan palms. The poeouuk ttruq called *Cocouus indicus* in the
 the aed off oie* nf yu-ri , probably tuberous, Several Brazilian spesies of Coecalmnd
 we ure "A to PO'e" powerful (ebirimrn) proifit(?)s. So S)cic* of Muctperniaceai 1* found in
 they we chiefly native* of tropical America and Asia. Cutting*.
 Tribe 1.
 BICO *Tiliacura Col*
 1129 *Wendlhndia W.*

17. OanEitVI. BERBER I'D E-fc.

, Spies 33; Hot.home Specie 10 •, GrcnJumse Specirs 10; Kany Li&wu* Specie* 20

/Arrrfly llahaceuns Sjxclct 3,f 35) feet; £ £ feet; * 0 reet

NuL •19 108(1) 1088 **m I.**
Then. 1 1057 Mr. Mr.

la. Ontif.ii vii.
Gerura 2, Species 2, Uot-houtr Species 0 j. G) <m-A<iiw Specirj 0; IIlirty IJgneotu Speci'tQ;
Uot-houtr Species 0 j. G) <m-A<iiw Specirj 0; IIlirty IJgneotu Speci'tQ;

^{Dn} *Uic* other to the *hwbaecon* genera of *Btberheit*. Their juice is held to be purgative. *Pl nson*

1548 | 1178 Je8erAn*g*ji Bart.

19. OatEn VIII. HYDROPELTIDEA

G«it» 1 Speiei I ; HaUmts Specifi 0; QwWt house Bpedtt 1 ; llarffi / Lirneut
trrii/ O M M N M Bptcsfa a J 0 feet | f U feet; * U feell

The order is formed by Nymphipice* chiefly in having a definite number of seeds, it consists* of only two families containing a single species. Both are little floating plants of tropical and northern America.

Hydroptilii L.

£0. OBDER JX.

*Genera 4, SpreiW Si; IJtff.hmse Speiet 19; Green-house Species i; Wmfff Ligneut Sprits 0;
JAifry Herbaceous Speciei 18. }0 feet; 10 feet; £ 3 J ft*

ilce the last, these are all flo.tinft plnnts, and to gardener, possessed of great interest, on account of the
***, form mid various hue* of their flower*. Three species are known as the lilies of our own uretanu and
"s am) the remainder occupy similar e tat ions in otier countries. Some of the Indian j s of
Jc de Umuuy frarfrat. The Jly Cy-nmu*, or Phihaporeainbean of antiquity, is the produce of the Ni-lum-
u " " " > stately aquatic, which buiinds in all the hotthr couiiLries of the Vast, where iU roots are frequently
It* l'sg an Jm* of tow, ditchis aiuxi Pokin and other Chinese cities, are literally choaked up with
J' w, un, ailee. Tin-petfoatyta or teant are dhling, hard, nnith t>dLi>, and jotmetw thi; power of veseting
J' w having been triili (or wun thirty yt-jirt. The flowers and rootiit of the common white Ayuia? have
long[cerebral tebl] ir sedative and antiaphritisical similitie, white are however, now conridertd
uttl in K. of tcorcity, the roots of A'liphor Mittea are pounded into cake* alone with the
park in IntM aylditrii.

In order to settle the question of the true nature of the seed of *Xelumkium*, it is necessary to consider the structure of the embryo. The embryo consists of two thick cotyledons, which are enclosed in a thick integument. The seed is described as having a thick outer layer, which is removed by the action of water. The seed is germinated in water, and the resulting plant is found to be a monocotyledonous plant, similar to *Imyedalini*. The author concludes that the seed of *Xelumkium* is a monocotyledonous seed, and that the plant is a member of the family *Zingiberaceae*.

but observer compared the seed of *Kuanum* to the seed in Jinygudam, and also t
aUo demonstrnted that the structure of the storn waa anainKOu¹ to that of
H* (limits. A very illirri-ni i : tly afterward² held by M I
I order, "do admitted iuti
il iffan taken hy previous obeer
were M entlreli
to the ulii upinkm, that Nyit.
ill i I laid, a nans for >
1 material
litnn, arnl

“‘*«nflirt of opinion**, I have detrnuii
I^t, because the structure of their steift w that
it- luMiv,” “in’lm, ‘il within the little h
---ay entyledoni. which i. BonflnBwi by *ou* |
“Ucame of the itrudui flower, whith has a *ji*
IV W; 4thly “on Aci*iu*Vnt’ot” the “fimiliirity *Ictweeti* their fruit mid etiginn and *that of* *lapAvi*”r; Sthly
“Ute oftheir” milkv juice *ntui* convolute leave*, two Kharartars which are not known to t-si-t among *lind*,*J*
ThoeewhonrelInterLitted in pursuing this curicm* itiseuwion any farUicT,wLll find mjinjy remark); anc
in the English edition of the *Anaivtc tiu Frail*, published by Mr. Xliidjiev in “*

NATURAL AURANGEME

Table 1. NHLU»(BO\SI
lt!13 Neltmbtum 3

2. Tribe 2. Nr^pn^BA-

tig. 0. kerb. IS. * Si feet.

ISM Barfwii Sal

MM Ayinphac'a Mae*. 0 *T

1557 Afiphar A'm, 0 *5

S3. ORDER X.

Gentu 1 i Species 4; Hot-house Specie* 0; Green-koute Species 4; Hardy Ligneous Species 0;
Hardy Herbaceous Specie* 0. • 0 feet; £ 0 feet; ^ 0 feet

Plant* remarkable for the singular form of their leaves, which are tubular and bold water, and some species have lids or covers, which, it is alleged, shrink and close over the mouth of the tube in dry weather, so as to prevent the exhalation of the water. The rarer it chiefly distinguished from Pajmvlracct and Nyropeapa** having a broad peltate leafy stigma. It consists only of one genus, containing 4 sin-cits, all inhabiting warm parts of North America. Division.

1555 Sarracen/o L.

SECTION II. Carpetta solitary or connate; Placenta parietal.

24. ORDER XI. FAPAVERA»CK»L

Genera 12. Specie* (& ; Hot-house Species; Green-koute Species 1; Hardu Ligneous Specie* d; Hardy Herbaceous Species (& 10 feet; £ IS ft; ■ Fret

These plants are better known for their medicinal properties than for their beauty. Some of them are the common pests of corn fields, and with grain have been disseminated over all the world. Sangulnkrila is a neat little American plant well known for its eritisHi juice, and the emetic purgative powers of its roots. The peculiar power of the poppy is, as a well known narcotic, a property which pervades all the order, although in a less intense degree in all than in the final *P.* (omniferous, from whence exclusively the drug opium is obtained. The Mexicans use the expressed oil of the seeds of Argimbiie inuxicar for polishing furniture-Division, seeds, or cuttings.

1552 Papaver	»38	I-JU" Sanguin'iria Z.	0	2	.1770 KschschVitzm Cham, 0
1554 Argentine Ton.	4	<B Boocinii i.,			n.tt. o
1553 Mecon6j < Vie.	*1	1423 Macleftyu R. Hr.	0	1	1516 nu-liiunnini /,,,/, /
3368 HuBBcmSu'A Sw t.		155 Uumeri'u Med.	0		MM iyilomM /.. o

K OHDSR XII. FUMAIKA'CILE.

Species U; Hat.house Species 0; Green-house ftwefetO] Hnrtly Ligneous Species 0;
Hardy Herbaceous Species U. 1 t) feet s £ 13 ft.; * U feet

Tender herbs, with finely cut leaves and annual stems, abounding in watery juice; -without any ant* of milkiness. They are reckoned slightly diaphoretic and aperient, but their medicinal properties trifling. Formerly they were combined with 1'aimveracefl, from which they are now unifinal di>ti>guish. The greater part of them are natives of hedges or thickets in the cooler parts of the northern hemisphere two are natives of the Cajie of Good Hope. Many of the species are beautiful ornaments of the flower-garden Division, seeds.

SOW Dicylrya Sore	0	9 2048 Cystic&pnos Boer.	0	2 1	2051 SarcocSpos Dec. 0
8050 Adlumia Jfiafi.	0	1 SH7 torydiilis Dee.	0	*23 £052 fumurik Tuu.	0

28. ORDER XIII. CRUCIFER.E {erut, a ctoa>fen}, to bear; form of corolla}.

Genera 88, Species 7G1; Hot-house Species 0; Green-house Species 77; Hardy Lignuous Species IS;
Hardy fferftsjiru Species GO5. J (J ft; £ 50 feet; £ 3 fat,

The importance of this order to mankind, and the singular nature of its botanical characters, render expedient to speak very fully upon it: in which time (remark) at the initial M Di-camhiHi, who has "Cruciferie particular attention, will be chiefly followed. This older omnibus wholly of annual or a binomial, herbs, occasionally assuming a subvirescent habit; then, however, never exceeding the height of three feet. The roots are either thick and perennial, or annual or biennial and slender, undivided and perpendicular and undivided. The young roots are t'pea with a little tilciijih, called the i-oleorliia, produced by the extended ruptured coat of the epidermis when the rootlet first appears. This is a character, and deserves attention. The stems are round or somewhat angular, branched, and often, two-valved the annual species, indurated at the base. The branches proceed from the axillae of the leaves*, but the most ones are abortive in most cases*. The racemes are always opposite to the leaves; sometimes* the terminal branch is abortive, when the raceme appears to be terminal; but this is merely owing to that circumstance. The leaves are simple, generally radical or alternate, i.e. opposite. The flowers are either white, yellow, or purple, or in a few rare species bright blue. The fruit is called either a silique or silicula, the former being a linear pod containing many seeds, the latter a roundish pod containing one or very few seeds, whence this



Older, which is the same as the Linnean class Tetradyrhmia. Is divided by TJnsns into two parts, L-alitil Siliquatus, and Siliruthbaw. In the Mod, the raiu'al ami cityliitnm arc sjnliet to emcb obar in dittront wt>, from which the subord LTD or M. IJoonJulU itt-rive their characters. When the edge of the cotyledons is pressed against the radiculus, so that a cross section would be an at fig. 5. a, that cotyledons are said to be decumbent, as in all I'ciirnrhlxea'; where the side of the stem; riifly:luij> i-jrrtwel to the radicula as at b, the R are called incurvate, as in Nutrliziee. If the cotyledons are inculcate, and at the same time half folded together or LDiuiplicate, as at c, the suborder Orthoijur<e is formed; when the cotyledons are iu-untbriit and spirally psw that a section would be as at d, they constitute the suborder BpionUBesi; and finally, when the cot cot> are incumntiit, itd doubled twice as their length, as at e, we have DilecolMttB

The whole order is preeminently European; 166 species are found in the north and middle of Europe, and 150 on the seashores of the Mediterranean; 45 are found between Mogadore and Alexandria; 18^g in the British Isles; 11 in the Canaries; 2 in Saint Helena; 2 in the West Indies; 11 in America; 48 in North America; 5 in Kamtchatka and the bordering islands; and finally, 33 are common to several parts of the world. From this it appears that there are about 100 species in the southern hemisphere, and about 800 in the tropics. 111; or, if they are considered with reference to the zones of temperature, 255 are natives of the frigid zone, the northern hemisphere; 30 of the whole of the tropics; 548 of the temperate zone of the northern hemisphere; many in open sandy places, some form the vegetation about the limits of the perpetual snows of lofty mountains, and many follow the footsteps of man through all parts of the world.

The useful qualities of the turnip, the radish, the rape, and the cabbage and its multiform varieties, are all known. The greater part of the order consists of plants possessing high antiscorbutic powers. These appear to depend upon a certain acrid, volatile, oily principle, the chemical nature of which is imperfectly known.

Herbage of the scurvy-grass, *Succowia*, is **mildly** when taken inwardly; thus, when any cruciferous plants are found to be eatable, either from exposure to other circumstances, it is to be understood to depend upon a reduction of this acrid principle. The Usual Powers of the last are what render the horseradish, the scurvy-grass, and others, so remarkably viui ag antiscorbutics; they are also believed to possess diuretic and diaphoretic properties. It is to be radix, &c., *terra*, &c., always eatable when the textures succulent and watery, as in the roots of the *Potato*, and the *Turnip*, and the leaves of the cabbage tribe. A further diminution of the acrid principle is caused by blanching. Cruciferae are said to possess a greater share of azote than any other tribe of plants; this is apparent in their fetid smell when fermented. The embryo of all the order abounds in oil, whence many species are employed with much advantage for expressing, either for eating or for feeding lamps. Some *Cruciferae* are extremely beautiful and fragrant, as the Stocks, the Gillyflowers, the Hespenses, the Candytufts, and many others. The Hutchinsias, Drabas, Cardamines, &c., are among the most interesting of *Wile* plants. Almost all the species are propagated by seeds, but some of them grow by cuttings, and the Perennials by division.

¹² f. ^{B<mi>} PLEURORHIZA	31. Tribe 5. ANASTATICEJJ., or PLEURORHIZEJS SEPTULAT ^B (dim. of <i>septum</i> , a partition).	1884 SiliKipis Ton. 0 *32
f. ^{& septum, a side or flat aa} root; radicle at the side of the cotyledons).	1885 MoncindVa Dec. 0 1	1886 Diplotaxis Dec. 0 *11
*. Tribe 1. ARABI'DEJE, or PLEURORHIZA	1887 frCica 7b. 0 10	
KORHIZEJS SILIQUOVK [siliquott, a long Pod].		
>> herb. 149. £19 ^s ft;	1888 Anasta L. 0	42. Tribe 1& VE'LLEJC, or .OR. THIOPLO'CEJE LATISEPTJB. lig. O. herb. 2. glfcet.
Matritoli R. Br. 0	1889 B6lcum Dcsv	1890 Carrichti-ra Adan. 0 *1
ChirAnthus L.	1891 Succuw'a Med. 0 1	1892 Psychline Desf. 0 1
Nasturtium R. Br.		43. Tribe 14. PSYCHI'NUC, or OUTUOPL'CEIE ANGUBTISE'FI'. lig. 0. herb. 1. ±2 feet.
Leptocarpax* a Dec.		1893 Z7/a Fojsk.
Notoceras R. Br.		4894 Muricaria Desv. 0 1
Or B.		1895 Calpacia Adan. 0 1
Ste. shirt Adan.		44. Tribe 15. Z'LLEK, or ORTHO. THOM'ET. lig. 0. her. b. 33. £ Hi feet
Ster.		1896 Trambe Zbit. 0 9
L.		1860 /Taptrum Bauh. 0 3
cropodium R. Br.		1898 Pidčinisi i».
S. dae L.		1809 Enarthrocarpus Lab. 0 2
		1897 Aphanus L. 0 17
ALYSSI'NEA, or LATISE PT. fl-		46. SUBORDER IV. SPJROLO\ BEM (speira, a spire, lobos, a pod; cotyledons).
	36. Tribe 7. SISYMBREJB, or NOTORII'ZEJB SILIQUOVA'C.	1872 Scnchiéra Poir. 0
lig. 1. herb. 123. f 3 ft.; £24) ft.	1873 Zcpodium L. 0	47. Tribe 17. DUMA'DEJG, or SPI-ROLO^BEJS Nt'CAMENTA'CEJI.
1862 Malcolm/a R. Br. 0	1874 Bivon'a Dec. 0	lig. 0. herb. 3. £3ifeet.
1863 /Kspelis L. 0	1875 Eunbmia 0	1900 Žunias Z. 0 3
1864 Andreuskia Dec. 0	1876 JEthoncma R. Br. 0	48. Tribe 18. ROLO'DEJI. lig. 0. herb. 22. £ Hi ft.
1865 Siftymbrium L. 0	1877 Mtis Bauh. 0 *18	1901 Eruckria Gae. 0
1866 Al/tfmAdan. 0	1878 Tauschérija Fis. 0	49. SUBORDER V. DIPLECO-LO^BEJE (diplex, double, lobos, apod; cotyledons).
1867 Efsimum L. 0	1879 3/^aprum L. 0	50. Tribe 19. HKIlopin'i.FjK, or P. lig. O. herb. 11. jf<jiueut.
1868 LeptMcum Dec. 0	1880 Sobolewbia Bich. 0	1902 HcUophilax. 0 17
1869 Stanlyca Nut		51. Tribe 20. SUBULARIK^, DIPLECOLO'BEJE. lig. 0. herb. 1. 1 foot.
		1903 Subularia /> 0 4
37. Tribe 9. LEPIDI'NEJE, or NOTORII'ZEJE ANGSTISE'FI' JS.		52. SUBORDER VI. SCHIZOPE-TJCLEJE.
lig. 2. /MT^36. t3ft. f II\$ft		1816 Schizopetalon Sims.
1872 Scnchiéra Poir. 0		
1873 Zcpodium L. 0		
1874 Bivon'a Dec. 0		
1875 Eunbmia 0		
1876 JEthoncma R. Br. 0		
38. Tribe 10. ISATIDEJI, or NOTORII'ZEJE Nt'CAMENTA'CEJK.		
lig. 0. herb. 22. £ Hi ft.		
J877 Mtis Bauh. 0 *18		
1878 Tauschérija Fis. 0		
1879 3/^aprum L. 0		
1880 Sobolewbia Bich. 0		
39. Tribe 11. ANCHONHAE, NoTomii'/K.: LOMENTA'CEJE. tie. 0. herb. S. £ 4 feet.		
1881 OoldbachwVi I>cc. 0		
1882 Sterigma Dec. 0		
4a SUBORDER IIT. ORTHO-PLÖ^CEJE (orthos, upright, ploke; cotyledons).		
41. Tribe 12. BRASSICE.E, or R. IUVLOVLO'PE. SILIQUOVK lig. 0. herb. 94. £23* feet.		
1883 A:ssica I.. 0 *40		

53. ORDER XIV. RESEDA^CEJE.

Genus 1, Species 23; Hothouse Species 0; Green-house Species 6; Hardy Ligneous Species 0;
Hardy Herbaceous Species 17. ± 0 feet j jf 10 ft.; £ 0 feet

This order differs from Cruciferae in the capsules being unilocular and the stamens indefinite, as well *Rt* in the seeds being destitute of an umbilical cord. Weeds of no interest, except the *Reseda odorata* for its delicious fragrance. *R. Luteola*, a common annual in waste places, yields a yellow color fit for dyeing. A" natives of Europe and North Africa. Seeds.

1459 *iZeseda L.**Q *YI*

54. ORDER XV. DATISCEAE.

Genus 1, Species 2; Hot-house Species 0; Green-house Species 0; Hardy Ligneous Species 0;
Hardy Herbaceous Species 2. £ 0 feet; £4 feet; ^ 0 feet

Differs from the preceding order in having an inferior ovary, and in the seeds being furnished with albumen. This order consists only of one genus, containing three species, natives of the south of Europe* Nepal, and North America. They are strong coarse hardy perennials, with the appearance of hemp.

DIVISION,

2805 *Datisca W.* 0 2

55. ORDER XVI. CAPPARDEJE.

Genera 11, Species 72; Hot-house Species 51; Green-house Species 5; Hardy Ligneous Species 0;
*Hardy Herbaceous Species 16. £ 0 feet; £ 8 ft feet; * 0 feet*

These are nearly related to Crucifers, of the properties of which they partake. Many are very pretty plant*, especially Cleome rufsea, and the various species or CrataeVa. The common caper is an elegant bush, remarkable for its large white flowers and long purple stamens. The species are found occasionally in various parts of the world. The different kinds of Capparis are reputed to be stimulating, antiscorbutic, and aperient. The bark of the root of the common caper passes for a diuretic medicine. Several species of Cleome have an acrid taste, which has been compared by travellers to that of mustard. The root of Cleome dodecadandra is employed as a vermifuge in the United States; and the leaves produce an inflammation of the skin, whence they are used in Cochin-China as a sinapism. Dec. Cuttings or seeds.

56. Tribe 1. CLEO^MEA. <i>lig.O herb. 16.</i> £ 8 ft feet	1455 <i>Polanisia Rqfi.</i> 0	1540 <i>Niebtihra Dec.</i> 1442 <i>Boscirt Lam.</i>
1905 <i>Peritoma Dec.</i> 0	57. Tribe 2. CAPPA^REJB.	1539 <i>Ctpparis JL</i>
1906 <i>Gynandropsis Dec.</i> 0	1440 <i>CrateV* L.</i>	1101 <i>Stephanha Dec.</i> 2045 <i>Morisonia Flu.</i>
1904 <i>Cleome W~</i> 0	1441 <i>Richie* R. Br.</i>	

5a ORDER XVII. FLACOURTIA'NEAI

Genera 3, Species 11; Hot-house Species 8; Green-house Species 3; Hardy Ligneous Species 0;
Hardy Herbaceous Species 0. ± 0 feet j jf 0 feet; ^ 0 feet

A very small order, formerly comprised in Tiliacee. It is remarkable on account of the structure of fruit, to the inner lining of which the seeds are attached upon a branched placenta. Nothing is known of the properties of the Flacourtiacae. The berries of Flacourtia Ramontchi are eaten in Madagascar. The order consists entirely of small tropical trees or bushes. Cuttings.

59. Tribe 1. PATRISIEAE. I 60. Tribe 2. FLACOURTIE^E. 61. Tribe & KIGGELARIE'A
1541 <i>Ryania Vahl</i> 2811 <i>FlacoUrta Herit</i> 1 2798 <i>Kiggeldria L.</i>

62. ORDER XVIII. BIXEJE.

Genera 4, Species 9; Hot-house Species 9; Green-house Species 0; Hardy Ligneous Species 0;
*Hardy Herbaceous Species 0. 10 feet; £ 0 feet j * 0 feet*

The plants of this order are few in number, and not remarkable either for beauty or use. The *Bixa Orellana* is chiefly known for producing the seed catted in the shops Arnotta (JZoon, FT.) and used for coloring cheese; the properties of the Arnotta are slightly purgative and stomachic. They are all bushes or 10*4 trees, and mostly tropical Azaras, Chilian shrubs with fragrant flowers, are not yet known in the garden* Europe. Cuttings.

1565 <i>Bixa L.</i>	I 1567 <i>Pr6ckia L.</i>	I 1568 <i>Ludia Lam.</i>
1569 <i>Lartia L.</i>		

63. ORDER XIX. CISTI'NEAL

Genera 4, Species 187; Hot-house Species 0; Green-house Species 55; Hardy Ligneous Species 111;
*Hardy Herbaceous Species 21. ± 22 feet; < 6ft feet; * 0 feet*

The common rock roses of our gardens give an accurate idea of this order, which contains little else. They are all very ornamental, and particularly well calculated for covering rockwork. They are natives of most parts of the world in dry elevated places. The gum called Ladanum is the produce of some kinds of Cistus. It exhales a fragrant perfume when burnt, and possesses slightly tonic and stomachic properties. Cutting* Europe.

1589 <i>Cistus Ton.</i> 40 0 I 1445 <i>Hudsonia L.</i>	I 292 <i>Lechea L.</i> 0 4
1590 <i>Helianthemum Tou.</i> *71 17	

64. ORDER XX. VIOLARIEAK.

Genera 12, Species 135; Hot-house Species 13; Green-house Species 13; Hardy Ligneous Species 0;
Hardy Herbaceous Species 109. ± 0 feet; £ 11 feet; £ 0 feet

This is one of the most favorite orders with gardeners; consisting, as it chiefly does, of the Violet £*nu? from which most of the others are recent dismemberments. The greater part are hardy herbaceous plants, some of which are remarkable for their perfume, others for their brilliant colours, and all for their fragrance. They are natives of the temperate or cold zones of both hemispheres, often growing at great elevations above the sea. Among them is a tribe called Alsodines, consisting of aromatic tropical plants; but none of them have been introduced into the gardens of this country. The attention of collectors should be directed to the shrubby Violaceae of Brazil, some of which possess great interest. The medical properties of the order are found principally in their roots, which appear to possess, in all cases, emetic properties, in a F^a or less degree. One of the Ipecacuanhas is the root of a Brazilian violet. M. Decandolle has the observations upon the affinities of the Violaceae; — They are very nearly akin, he observes, to the Rosaceae, and especially to the Passiflora?. From the first they are distinguished by their "Vibrantia", fruit, leaves furnished with stipules, and two-celled anthers j from Drosaceae by their solitary style, lengi

*%? embryo, and stipulate leaves, the vernation of which is involutive, not circinate. From Passiflora the Jęzler in their fruit being capsular, not berried; in their albumen being compact and shining, not pitted; in their stamens being hypogynous, not perigynous; in their anthers being attached along their whole length, not fixed by their middle; finally, in their stigmas being one and not three. The genus Calyptrion approaches Passiflora in its twining stem, and Hymenanthera borders upon Polygala on account of its monospermous Pericarpium with solitary pendulous seeds. Divisions, seeds, and cuttings.

65. Tribe 1. VIOYEA. -0. herb. 109. £ 11 feet	703 SMea Ging. 704 Pombalfa van. 705 Ionidium Yen.	0	709 Ceranthera Beauv. 700 Lavrtdia Vill. 3310 Hymenanthesra R. Br.
?1 folia Tou. << Erp&tion Swt.	66. TribeS. ALSODI'NEA 708 Alsodela Thou.	67. TribeS. SAUVA'SE & 689 Sauvag&sia Jac.	

68. ORDER XXI. DROSERACEAL

Genera 5, 5jKde<16; Hot-house Species 0; Green-house Species 5; Hardy Ligneous Species 0

Hardy Herbaceous Species 11. f 0 feet; £ 2 feet; £2}feet

L **P** **order of sun-dews** is a small group of plants, natives of marshes or inundated grounds in all the temperate parts of the world.* The species are very remarkable for the abundance of glandular hairs with which all parts of the forage are covered. Only two species are in any degree frutescent. The young leaves are always rolled up in the circinate manner, so remarkable in ferns. Their medicinal properties appear to be "liling; the leaves have the power of curdling milk. Divisions or seeds, some by leaves.

§22 DnSsera L. 0 *6 I 923 B?bli« Sal. I 910 Pamaßsia L. 0 «1
925 Aldrovanda L. 0 1 | 1331 DionaeVi L. I

69. ORDER XXIT. POLYGA[^]LEJE.

Genera 6, Species 64; Hot-house Species 5; Green-house Species 41; Hardy Ligneous Sproies 1;

Hardy Herbaceous Species 17. 2 1 foot; £ 5 feet; £ 0 feet

Most of the plants of this order are interesting, and deserving the attention of the gardener, some for their neatness, some for their beauty, and some for their use in medicine. They are natives of most countries, and are either low herbaceous plants, occasionally less than an inch in height (small specimens of *Polygala purpurea*), or shrubs varying from a dwarf, rigid, spiny habit, to a tall, graceful, drooping appearance. *Polygalaceae* are remarkable for the union of their stamens into a single body, their one-celled anthers opening with a pore, just their irregular flowers, one of which is often keel-shaped, and beautifully crested or bearded. The leaves have generally a bitter astringent taste, which is much more abundant in the roots, combined with an acrid and somewhat resinous flavor: these properties are particularly sensible in *P. Sénega*, which is reputed a

cularly used as a remedy for dysentery. The well known Ratanha root, of Chile, is the Produce of a plant of this order, and possesses powerful tonic and astringent qualities. According to the analysis of a trench chemist, it contains gallic acid, but neither tannin nor resin. Cuttings or seeds,

£053 i>Okgalia Tou. *1 17 I 2055 MAndia Kth.
205* Murflit/a Neck. ' I 3056 Securidaca L. I 1703 Kramen'a I*

70 ORDER XXI1L TREtfA'NDBEiE

Genus 1. Species 6 j Hot-house Species 0 ; Green-house Species 6; Hardy Ligneous Species 0 ; Hardy Herbaceous Species 0. ± 0 feet; £ 0 feet; ^ 0 feet;

▪ A very small order, containing only seven species, all small bushes, natives of New Holland, and remarkable J* the peculiar neatness of their appearance. In habit, they may be compared to heaths, with which thoſ agree in the anthers bursting by a pore at the end. Nothing is known of their properties. Cuttings or seeds! ▪

51 ORDER XXIV PITTOSEO%P&S

Genus 4. Species 82: Hot-house Species 2; Green-house Species 20; Hardy Lignaceous Species.

Hardy, *Hardyaceous Species* 9, + 9 feet; if 9 feet, Δ 9 feet $\Delta\Delta$ *

Hardy Herbaceous Species. ± 0 feet; jf 0 feet; ^ 0 feet ***
Beautiful shrubs and small trees, with simple alternate leaves destitute of stipulas and
flowers, chiefly natives of New Holland and the African islands. The order is distinguishable
by estimation of the sepals and petals which in ^{the} t, as well as the stamens, are & w S \$
innumerate embryo. Cuttings and seeds.

671 Billardiæ Sm. I **689 Bursaria Cav.**
679 Pittosporum Ban. | **665 Seneca Com.**

72 ORDER XXXV FRANKENIA*CEF

Genus 1, Species 10; Hot-house Species 0; Green-house Species 4; Hardy Ligneous 
Hardy Herbaceous Species 6 * 0 feet fV2*feet; *0 feet

Hardy Herbaceous Species 6-10 feet; 1/2-2 feet; 1-6 feet
distinguished from *Caryopteris* by the fruit not having a central separate placenta, but bearing the seeds on a single central column. They have no medicinal properties. Besides the genus here recorded, there are two others mentioned by W. Decandolle. Cuttings, division, or seeds.

1999 FrankemVz L. 8^6

SECTION III. *Ovarium solitary; Placenta central.*

73. JP RDBR XXVI. CARYOPHY'LLEÆ

Genera 26, Species 531: Hot-house Species 11; Green-house Species 27; Hardy Ligneous Species 1; Hardy Herbaceous Species 492. f 1 foot; £ 38* feet; JklJ feet

In these consist of herbs or low undershrubs, inhabiting the mountains and pastures of all parts of the world, and Siberia they are particularly abundant, and least so in Africa and South America. Many are annual weeds, as most of the Cerastiums, Spigularias, and others. Several of the Silenes are very ornamental plants, the *Arenaria* are to be found some dwarf species of considerable elegance. But it is in *Dianthus* that the order consists: this genus is almost unequalled for the brilliancy of its colours, the neatness of its flowers. From the finest of its species the title of the order has.

K k S

74. Tribe 1. SiLEMkifi. tig. 1. herb. 315. f 1 ft; £34* ft.	75. Tribe 8. ALHI'NE.E. lig. 0. kerb. 177. £ 16 ft	290 tftotslcum /. 1418 Splrgula /. 1417 1 ^{er} bre/i Hil.	0 .2 0 .9 0 .1
1384 GysAphila L.	28	114 Ortegfe L.	OS
1386 Dianthus L.	*93	406 Buffbnia IV.	0 *S
1385 Saponaria L.	■	416 SagtaazR.	0 *6
1387 CucubalusL.	0 *1	1208 MoehringiaL.	0 2
1388 Silenc L.	0 *153	1223 flatinC L.	0 * 2
1415 Zycbui L.	0 *28	1419 Borgia L.	0 1
802 Velezitz W.	0 1	295 Jollbgo L.	0 1
901 JDrypis L.	1 0	906 7'harnaceum W.	0 1

76. ORDER XXVII. LINNEAE.

Genera 2, Species 43; Hot-house Species 0; Green-house Species 6; Hardy Ligneous Species 2;
jftmfy Herbaceous Species 35. £ 3 feet; £ 12fcet; A 0 feet

Separated by If. Decidolic from Caryophylleae, from which it is well distinguished by its fruit having several cells, or in the language of the botanist just named, being formed by the cohesion of several carpella. Most of the species are pretty plants, bearing yellow, blue, or white flowers. They are of immense importance in the world, on account of the tenacity of their fibres when made into flax. The seeds of common flax are between mucilaginous and oily; the leaves of Linum catharticum and L. selaginoides, the latter a native of Peru, are purgative. Cuttings, division, of seeds.

* 921 Zlnura Bauh. *2 34 419 RadAola Dil.

■ ■

77. ORDER XXVIII. MALVA*CJF&.

Genera 22, Species S91; Hot-house Species 203; Green-house Species 97; Hardy Ligneous Species 8;
Hardy Herbaceous Species 83. f 28J feet; £ 25 feet j £ 0 feet.

Before this order was dismembered of Bombacce and Byttneriace, it contained most of the grandest flower* in nature. Even now, the splendor of the various species of J/Alva, yithseVi, to which the hollyhock belong'f and //ibiscus, renders it one of the most remarkable groups of plants. With the exception of the numerous genus Sida, nearly all Malvaceae are objects worthy of the gardener's care, particularly those which are hardy. In stoves or green-houses, the softness of their branches and leaves renders them peculiarly liable to the attacks of the red spider, mealy bug, and scale, from which few collections are free; a circumstance which makes them less generally esteemed than the surpassing beauty of many of them merits. The greater part of the order is clothed with stellate pubescence, and a reniform one-celled anther is a character common to the whole. These two peculiarities, together with the alternate stipulate leaves, distinguish Malvaceae from all the rest of Dichlamy'dcr. All the species abound in a nutritive mucilage; a quality which renders the young heads of the Ochro, or Hibiscus cscullant, an object of great value within the tropics, as an ingredient in soups. In Brazil, the Abutilon, esculentum serves the same purposes. The emollient properties of //Ithaca officin&li'f or Guimauve of the French, are well known to physicians, as a remedy for catarrhs and pulmonary complaints. A decoction of the leaves of Sphaerxcea cisplathna is used for similar objects in Brazil. A species of Laybmw* is employed in the same country as a diuretic in the form of a decoction. The straight shoots of SW* micrantha are employed as rocket sticks at Ilio Janeiro. The chewed leaves of Sida carpinifolia allay the inflammation occasioned by the stings of wasps. The tough fibres of many Malvaceae are manufactured into cordage. Their petals are astringent; whence those of J/ibiscus rbsa sinensis are used in China to blacken the eyelashes and the leather of snoe&. The fibrous threads in which the seeds of Gossypium are enveloped furnish the valuable cotton, an article of immense importance to the world; these threads, when examined by the microscope, will be seen to be finely toothed, which explains the cause of their adhering together with greater facility than those of Bi'mba and several Apocynaceae, which are destitute of teeth, and which cannot be spun into thread without jgan admixture of cotton. Division, cuttings, or seeds.

78. Division 1. Calyx double.	2010 Uriia L.	2018 Lopifmia Mart.
2003 Afalope L.	0 1	2011 PavbmViCav.
2004 Mfilva L.	0 *38	2012 Malvaviscus Dil.
2005 Nuttillia Dick.	- 0 2	2013 Lebrctbnfo Schrank
2006 KitaibMTTW.	0 1	2014 7'bfocus L.
L007 ifltha^a L.	0 *16	2015 Thesptzia Corr.
2008 Lavitem L.	0 *J2	2016 Soti Gossypium L.
2009 Jfafachra L.		2017 Redoutta Ven.
79. Division 2. Caly* simple'		
2019 PalKyta Cav.	0	
2020 Cristiria Cav.	0	■
2021 A'noda Cav.		
2022 T'erpptcra Dec.		
2023 Sida L.	0	■
2024 Lagun^a Cav.		

80. ORDER XXIX. BOMBA'CRX.

Genera 14*Species 28; Hot-house Species 27; Green-house Species 1; Hardy Ligneous Species 0 :
Hardy Herbaceous Species 0. }0 feet; £ 0 feet; *0 feet

Distinguished from the last by the imbricate aestivation of the calyx, and the arrangement of the stamens in five sets, or, in Linnean language, brotherhoods. The species are mostly fine trees with large showy fl*we^A and natives of the tropics. Some of them are among the largest trees in the world; Adanso'mr, the Baobab of Senegal, has been seen with a diameter of twenty-five feet, and specimens of Btmbax Cciba and Eriodcn^dron anfractibus are not uncommon an hundred feet in height. The wood of all the species is light an< soft, as in Malvaceae, from which this order probably does not differ in its medical properties. Cuttings or seeds.

Vm Hciicteres L.	1995 Montexdima M. & S.	1921 OchrarbmaSw*.
£031 Myrbdia Schreb.	2027 Caroline** L.	2177 Diirio L.
2002 Plagi&nthus Font	2029 Bomrax L.	1H34 Cheiroitemon H.%B.
2028 Adanso'ma L.	9050 Eriodendron Dec.	

81. ORDER XXX. BYTTNERIAXEJE.

Genera 28, Species 151; Hot-house Species 78; Green-house SpecimIS; Hardy Ligneous Species 0;
Hardy Herbaceous Species 0. £ 0 feet; j 0 feet; h. 0 feet

Much the same kind of plants as those of the two last orders, from which they were not formerly distinguished; and from which they scarcely differ, except in their bilocular anthers. Many of the Sterculia are fine umbrageous trees, the seeds of which are large and eatable; especially those of the famous ^ojia* which possess the property, being chewed, of rendering bad water pleasant to the palate. The seeds of tat Chicha, another and very noble species of the genus, are highly esteemed in Brazil for the dessert. Astrap<> in gruirnaca. uutztina timiona has us fruii muca wun a picasani mucuagr, which is swcv*^{**} agreeable; an extract of the bark of the same plant is used in Martinique to clarify sugar; its old bar* employed, in the form of a strong decoction, as a sudorific. WaRlu;r< Douradinha contains a great w* of mucilage, and is employed by the Brazilians as an antisyphilitic. Cuttings or freeds.

82. Tribe 1. STKRCULIKK.
 52t StercMia L.
 ☐ J Recvēsfe Lindl.
 *725 Heritiira H. K.
 f1^83n Tribe 2. BYTTNERIIJE.

5178 Abrbm JL
 Guaxhma Plu.
 ☐ Coinmcrabfa W.
 Rulmgif R. Br
 Byttmfrw Loe
 AtewWIL.
 Kleinbdifl L.

84. Tribe 3. LABIOPBTA*LBI£.
 682 Ser'ng/a Gay
 680 Lasioi'taluin 5m.
 681 Thomasi'a Gay
 8J. Tribe 4. HERMANNIE£.
 1918 Melbchia L.
 1919 Riedleta Ven.
 1915 Waltherra L.
 1916 Hermann/Vi L
 1917 Mahernia L.

86. Tribe 5. DOMBBTAVBA
 2025 Ru{»aCav.
 1997 J'entipetes L.
 1996 Assbt'a Cav
 1994 Dombeva Cav.
 1920 Melhania Fortk.
 2000 Pterospermum Schrb
 1998 Astraps'a Lindk
 1999 K^dra Rox.

87. Tribe 6. ☐ ☐ ☐ ☐

88. ORDER XXXL TILIA^CEJE.

Genera 12, S/wfes 77; Hot-house Species 61; Green-house Species 5; ifanty Ligneous Species U;
 Ain/y Herbaceous Species 0. 114 feet; £ 62 ft.; £ 0 feet.

These shrubs, or herbs, in geneial not remarkable for their beauty, the greater part of the last being the win[er]est weeds of the tropics. The Lime, from which the order derives its name, is a genus of fine trees Dime F'ab'ant. flowers, and Sparmannia and Entelea are handsome broad-leaved green-house arborescent trees. The inner bark of Tilia is tough and separable, and supplies the material whence the Russia mats vUrLv. gardeners and others are prepared. Cúrchorps olitribus is cultivated in Egypt as a kitchen-garden of the bark of Cúrchorua capsularis are twisted into fishing lines; and the roasted nuts of the Lime tree arc reported to bear some resemblance to chocolate. Cuttings or seeds.

JSH E"telea R. Br.
 KI? Sparmannia L.
 E'E Heliocarpus L.
 «77 Corchorus Z..

1443 Triumftta L.	1574 MuntingrVi L.
1575 Grtwm J.	1571 Apelba Marg.
1568 Brownb/fl Rox.	1570 Sloknea L.
1576 Tilia L.	+11 0 1566 Berrya L.

89. ORDER XXXII. ELJEOCALPE^L

Genera 4, Specks 6; Hot-house Species 3; Green-house Species 3; Hardy Ligneous Specks 0;
 farfly Herbaceous Species 0. ± 0 feet; j 0 feet; ^ 0 feet

These differ from Tiliaceae in nothing except their lobed petals and anthers opening by two pores at the Apex. The flowers of some of the species of Ekeocarpus are fragrant, the fruit eatable, and the hard rugose tones manufactured into necklaces. Cuttings.

1434 Elaoocarpus L.	I 1436 Dicra Font. A
1435 Aceratium Dec	I 1437 Friesia Dec A

9a ORDER XXXIII. CHLENATEJE. -

Genus 1, Species 25 Hot-house Species 2; Green-house Species 0; Hardy Ligneous Specie* 0»
 Hardy Herbaceous Species 0. ± 0 feet 5 £ 0 feet j ^ 0 feet

This order, according to Petit Thouars, is allied to Malvaceae, in having an involucrum to the flower, and the stamens being monadelphous; but, according to Jussieu, it is allied to Ebenaceae, from the petals being connected at the base, and in the seeds being furnished with albumen. Shrubs and trees, with simple alternate stipulaceous leaves, chiefly natives of Madagascar. The genus Hugelia does not agree with the rest of the genera of this order, in being destitute of the involucrum to the flower, therefore it is placed in Chlenaceae affines by Decandolle. Cuttings.

* 1935 HugtwViL.

*

91. ORDER XXXIV. TERNSTRGZMIA'CEiE.

Genera 11, Specks 19; Hot-house Species 10; Green-house Specks 41 Hardy Ligneous Specks 5;
 Hardy Herbaceous Specks 0. f 18 ft 3 £ 0 feet; * 0 feet

A very small order, consisting wholly of trees or shrubs, bearing handsome white or yellowish flowers. They are nearly related to Camelliae, from which they do not differ at all in habit. Nothing is known of their properties. Noronha states that a species of Sauralja found in Java has a subacid fruit, in flavor resembling the Tomato, and that is eaten by the Javanese under the name of Koleho. Inarching, cuttings, or layers.

SL Tribe 1. TRRNSTR(EMIE£.
 159 TertMtraVnia L.

1562 Letts5m(aR.ftP.

Ug.5. herb.0. ± 18 feet

93. Tribe 2. FBEZIB^BE&
 ☐ Freritria Sw.
 2875 Eurya Thun.

94. Tribe 3. SAUBAU'JEJS.

2035 Malachodlron Cav. 1	0
2034 Stuartha Cav. 1	0
2033 Gordbnia EL 3	0
2036 Pol'spora SwL	

95. Tribe 4. LAPLA^CE.«

2032 Cochlospermum Kth.

97. ORDER XXXV. ☐ ☐ ☐ ☐

Genera 2, Species 59; Hot-house Species 0; Green-house Specks 59; Hardy Ligneous Specks 0;

Hardy Herbaceous Specks 0. £ 0 feet; £ 0 feet; £ 0 feet

The claaaarc too weU known in our gardens to render it necessary to say much upon their peculiarities. V-t-amii'lia is one of the most beautiful, and the tea one of the most useful, plants in the world. Both are in[*]es of China, Japan, or Nepal. The tea is well known for the stimulating influence of its decoction of the nerves, which is attributed by Cullen to the presence of a narcotic principle. The seeds of Camellia circa yield a fine oil. None of the species bear fragrant flowers. Their nearest affinity is with Ternstroem*. uce<£, from which they probably ought not to be separated. Cutting, grafting, or inarching.

2037 Thea L. I 2038 Camellia L.

98. ORDER XXXVI. OLACI'NE£

Genera 4, Specks 6; Hot-house Specks 5; Green-house Specks 1; Hardy Ligneous Specks 0;

Hardy Herbaceous Specks 0. ± 0 feet j ± 0 feet; ^ 0 feet,

trees or shrubs, with simple stalked exstipulate alternate entire leaves, and little axillary flowers, s doubt whether what is called a calyx is not rather an involucrum, in which case the corolla would a calyx, and the station of the order among Monochlantdee, rather than in this place. Cuttings.

104 OMax L.

I 1319 Heistcria L.

105 Spcmixyron Lab.

I 1171 Ximenia L.

Kk 4

NATURAL ARRANGEMENT.

99. ORDER XXXVII. AURANTIA*CEA

Genera 11, Species 32; Hot-house Species IS; Greenhouse Species 20; Hardy Ligneous Species 0; Hardy Herbaceous Species 0. 1 0 feet; £ 0 feet i & 0 feet*

These are also known under the name of *Hesperidete*. They consist of trees or shrubs of the greatest beauty and utility. The well-known orange and lemon are the representatives of the order, the characters of which are so well defined that there is no material deviation from the type afforded by those species. The thick leaves, articulated with the petiole, and abounding in transparent reservoirs of odiferous oil, are the most obvious peculiarities. The flower* are fragrant, and the fruit in all cases fleshy, and generally eatable. The wood is particularly close-grained. The volatile oil contained in the reservoirs of the leaves and fruit possesses powerful tonic and stimulating properties. M. Decandolle thus explains the singular structure of the fruit of the orange. In the opinion of this learned botanist it consists, first, of a thick, valvate, indusiate indusium or coat, which is most likely to be considered a continuous torus. Secondly, of several carpella, verticillate around an imaginary axis, often separable without laceration; membranous, and either containing seeds only, or filled with pulp, lying in innumerable little bags proceeding from the inner coats of the cells. Cuttings, and sometimes by layers, budding, grafting, and inarching.

1322 Atalant'a Corr.
654 Triphasia Lou.
1320 Limonia L.
1327 Coukia Sonn.

1326 Murraya Ron.
1321 Bergia Kon.
1324 Clausina Brm.
1323 Glyc6smis Corr.

1325 Ferbnia Corr.
1585 JE^gle Corr.
2187 Citrus L.

100. ORDER XXXVIII. HYPERICI'NE.

Genera 7, Species 90: Hot-house Species 4; Green-house Species 34; Hardy Ligneous Species 14; Hardy Herbaceous Species 38. 1 9} ft. j 14 ft. j fe 0 feet*

The whole of these abound in a resinous juice, and are in most cases glandular in some degree. Their leaves are all dotted, and, which is very remarkable, the dots are often black, even upon the yellow petals. These latter have a singular obliquity, which is not indicated by their outline, but by the arrangement of their veins. The juice just noticed as abundant in this order is yellow, viscid, rather bitter, often purgative or anthelmintic, and so very analogous to Gamboge, that the juice of *Hyppocrycum baccatum*, and some other Guiana species, has received the name of American Gamboge. Most Hypericines are bitter, and slightly astringent, whence they have been used as febrifuges. A small part of the order is tropical; but in its most genuine form it consists of herbaceous or undershrubby plants, delighting in the shade of groves and thickets in the cooler parts of Europe and Asia. Nearly all the flowers are yellow; those of *H. cochininchinense* are dull red. Seeds, division, or cuttings.

101. SUBORDER I. HYPERICF-
KEJE V&RJE.
108. Tribe L ■■■■■
2191 Har&nga Thou.
2192 Vfsmea Van.

103. TribeS. HYPERI'CES.
iff. 14. herb. 37. £9* ft. ;£ 14) ft.
2043 j*ndrose*mum AU, 0 »1
2190 flypericum L. *14 37
2044 /Tscyrum L.
300 V Elodea MM.

104. SUBORDER II. HYPERICI''
NEJE ANO*MALJE.
1611 Carpod6ntos Lab.

105. ORDEB XXXIX. GUTTI'FERJL

Genera 11, Species 28 j Hot-house Species 26 j Green-house Species 2; Hardy Ligneous Species 0; Hardy Herbaceous Species 0. ± 0 feet; £ 0 feet; £ 0 feet

Trees or shrubs found in the hottest parts of the world, and well known by their thick entire opposite leaves and resinous juice. In the countries where they grow they are of great importance. One, the Garcinia *Mangosthina*, bears a fruit, the equal of which is supposed not to exist. The well known Gamboge is the impregnated juice of *Garcinia Gambelia*, and, perhaps, other species; the juice of others is found an efficacious vermifuge, and also a remedy for the chiggers, one of the worst pests of equinoctial America. The bark and fruit of many Garciniæ are astringent. The unripe fruits of *Grias cauliflora* are pickled. The dowers of all the order being showy, the foliage good, and the properties interesting, every species deserves cultivation*. Cuttings or layers.

106. Tribe 1. CLUSIB'A.
2877 Clisia L.
3369 Godbya R. & P.
1581 Bonnetta Schreb.
107. Tribe 2. GARCINIB'B.
1429 Garcinia L.

108. TribeS. CALOPHYLLEE.
1580 Mammea L.
2188 Xanthochymus Box.
1579 Calophyllum L.
2189 Pcntadclena R. Br.

109. Tribe 4. SYMPHONIES.
1439 CanillaP. fr.
110. DOUBTFUL.
1578 GrlasZ..
1533 ? Rheædta U

111. ORDER XL. MARCGRAAVIA"CEEL

Genera 4, Species 6; Hot-house Species 6; Green-house Species 0; Hardy Ligneous Species 0; Hardy Herbaceous Species 0. f 0 feet j £ 0 feet; £ 0 feet

Very curious half-climbing shrubs, all natives of hot countries. Some of them bear among the flower*, which are large and showy, singular hollow bodies, like the pitchers of *Sarraceno*. The order has been well illustrated by Professor Hooker, in the 160th article of his *Exotic Flora*. Cuttings.

112. SUBORDER! MARCGRAVI&JE.
1543 AnthophmaZo&
1542 Marogravia Plu.
i»

113. SUBORDER II. NORA'NTEM.
1544 Koranæa Aub.
688 Räysch/a Jac.

114. ORDER XLI. HIPPOCRATEA^CEiE.

Genera 5, Species 13; Hot-house Species 13; Greenhouse Species 0; Hardy Ligneous Species Oj; Hardy Herbaceous Species 0. ± 0 feet; £ 0 feet; £ 0 feet

Little is known of this order. The species are tropical arborescent or climbing shrubs, with opposite leaves, and small inconspicuous flowers. The genus *Tonsdilla* contains some species known in Sierra Leon*.. bearing poisonous fruit. Cuttings.

100 Hippocratta L.
102 A'nthodon B. \$ P.

I 101 Tonsitta Vahl
\\ 10S Johnfa Rox.

I HIPPOCRATEA:E< sr</Ri*
| 19S6 Trigbnia AubU

115. ORDEBXLIT. ERYTHROXY'LEJR

Genera 8, Specks 4j Hot-house Species, Green-house Species 0; Hardy Ligneous Specks 0; Hardy Herbaceous Species 0. ± 0 feet; £ 0 feet; * 0 feet*



iJS*?. order is distinguished from Malpighiaceae in the presence of albumen, and in the petals being jyrnwhed at the base with a scale; the habit of the plants is also different Shrubs and trees, with simple Alternate leaves furnished with axillary stipulas. Native of South America, Mauritius, and Madagascar, and one from India. Cuttings.

1400 *Erythroxylon L.*| 1401 *Sethwi Kth.*

116. ORDER XLIII. MALPIGHIA*CEAL

*Genera 13, Specks 78; Hot-house Species 72; Green-house Species 0 j Hardy Ligneous Specks 0; Hardy Herbaceous Specks 0. ± 0 feet; £ 0 feet; * 0 feet*

*i?n?ulated unguiculate spreading petals form one of the most obvious characters of this order, the species of ft' i?l' o? a'e ill tropical, and are either trees or shrubs, often climbers. Many of the Malpighias are well known JIII the prurient hairs produced on the surface of their leaves; their fruit is eatable, their timber of a deep red r?o?i and their bark a febrifuge. Their showy pink or yellow flowers, and firm neat foliage, render all this r?" worthy of cultivation, except Aspicarpa, which is a weed. Cuttings.

"7. Tribe 1. MALPIQHIB**.	118. Tribe 8. HIFIVQEJL	1396 <i>Tripteris L.</i>
	1388 <i>Gertnera Rox.</i>	1401 <i>Tetrapteris Cav.</i>
	3352 <i>Thryallis L.</i>	1395 <i>Banister/a L.</i>
	31 <i>Aspicarpa Rich.</i>	1398 <i>Heteropteris H. * B.</i>
	119. Tribe 3. BANIBTERIE\B.	1165 <i>pVittmannia Vahl</i>
	1399 <i>HiraVi Jac.</i>	

ISO. ORDER XLIV. ACERI'NEJEL

Genera 2, Species 88; Hot-house Species 0; Green-house Species 1; Hardy Ligneous Species 87; Hardy Herbaceous Specks 0. f 73 ft. j ^ 0 feet; £ 0 feet*

oble trees, natives of the woods of Europe, Siberia, and North America. Their flowers are in all cases am? Bi?col M i the breadth and rich color of their leaves constituting their beauty. All the larger species found in very saccharine sap, from which sugar is prepared in North America: it is chiefly made from but? saccharin? um an? *Hegiindium*, but may be obtained from many others. Commonly by layers or seeds, au the species will grow by cuttings in sand.

8867 *iTcer L.*<85 0|8774 *Negundo Mce*

181. ORDER XLV. HIPFOCASTA^NEJE.

*Genera 8, Species 14; Hot-house Species 0; Green-house Species 0; Hardy Ligneous Specks 14; Hardy Herbaceous Specks 0. 141*ft; £ 0 feet; * 0 feet*

all the order » much valued for the grandeur of the foliage and flowers of most of the species, which are Hardy trees. Their bitter fruit has sometimes been used as a sternutatory; it contains a large quantity of starch, and an abundance of starch. The bark is astringent, bitter, and febrifugal, and has been recommended - a substitute for Cinchona. Seeds or layers, and by grafting.

1137 ^'scului L. 6 01 JIS8 *Pavia Boer.* 8 0

188. ORDER XLVI. RHIZOBO^LEJE.

Genus 1, Species 3; Hot-house Species 3; Green-house Specks 0; Hardy Ligneous Specks 0; Hardy Herbaceous Specks 0. J 0 feet; ^ 0 feet; £ 0 feet

and a ver. di ti nct order closely allied to the Terebinthaceae, but, from the hypogynous insertion of the Details in th? mma? and the structure of its fruit, differs essentially from that order. It agrees with Hippocastanace a laUL in * Aon & L Petals and stamens, and in bavin* opposite palmate leaves, but differs from it in having con* rad? and small cotyledons, not with a small radicle and large cotyledons as in that order. The order for £! ns on vone Kenusg the Carycar, the Rhizobolus of Gsertner and Ptkea of Aublet: a genus remarkable for Bw? ringe Butter and Saouari, or Suwarow, nut. The last is not unfrequent in our fruiterers' shops. It tree, and palatable, containing a rich oil, and the pulp is dry, not buttery as in the Butter.nut. Large ' natives of Guiana and Essequebo; propagated in our stoves by cuttings, in sand under a bell-glass

1601 *CarycarZ.*

183. ORDER XLVII. SAPINDA^CEJE.

*Genera 19, Species 88; Hot-house Species 78; Green-house Species 15; Hardy Ligneous Species 1; Hardy Herbaceous Specks 0. ± 10 ft.; * 0 feet; £ 0 feet*

t ^*ne of * inside. nr. distinctive peculiarities of this order consists in the petals having an additional lobe in the few of f if m hairs instead. Nearly all the plants have compound leaves, and bunches of white flowers; we wi-i em are twining herbs, but the greater part are trees or shrubs, all natives of the warmer parts of Afric? and in a great proportion, of the East. The only genus which will bear the climate of England is ignor? a fine shrub or small tree, with panicles of white or pale yellow flowers. Nephelium and am? ar? are both genera bearing excellent fruit. The rind of the berry of Sapindus Saponaria is of a ex2Si tility as the name of the plant indicates. The pulp of Melicocca and the arillus of Bughia sapida tient eating. Cuttings, layers, or seeds.

	1163 <i>Talisia Aub.</i>	1160 <i>Melicocca L.</i>
	162 <i>Maturba Aub.</i>	1108 <i>Stadmanni Vi Lam.</i>
	1W7 <i>Schnidell L.</i>	
	2159 <i>Euphoria Com.</i>	
	5340 <i>Pierardta Rox.</i>	
	1164 <i>Thouinia Poit.</i>	126. TribeS. DODON^ACE*. lig.l. herb. ti. £ 10 feet.
	1158 <i>Cupinia L.</i>	1166 <i>Kolreuteria Lax.</i>
	1093 <i>Cossignia J.</i>	1179 <i>Dodoive Vi L.</i>
		8656 <i>Atnirbla Pen.</i>

NATURAL ARRANGEMENT.

127. OADBBXLVIII. MELIA'CEJE.

Genera 14, Species 85; Hot-house Species 29; Green-house Species 6; Hardy Ligneous Species 0; Hardy Herbaceous Species 0. f 0 feet; j 0 feet; ^ 0 feet

The nearest affinity of this order is probably with Sapindaceæ. It is particularly distinguished by the tamens being united into a tube bearing the anthers. The leaves are usually pinnated, and most often species, which are all either trees or shrubs, are natives of tropical forests. *Jatilia* bears bunches of fine lilac-colored flowers, but few of the genera are interesting on account of their inflorescence. The qualities of which different species are little known. The bark of *Gaura trichilioides* is said by Aublet to be purgative** emetic. The pulpy fruit of *Melia Axedardch* is said to be poisonous; both this part and the inner bark have been used as anthelmintics either in substance or in decoction. It is asserted by Michaux, that the pulp surrounds the kernel is considered in Pekin a specific in scrophulous cases. The oil expressed from the seeds of the same plant is said to have strong antispasmodic powers. Cuttings or layers.

128. Tribe 1. MBLIB\B.

- 1295 Turned L.
1296 Quivisia Cav.
1294 Sandoricum Cav.
1293 A/elia L.

12a Tribe 2. TRJCIULIB\&

- 1292 Trichilia L.
1301 Ekeblrgia Spar.
1167 Gudrea L.
1302 Heynea Rox.

ISO. Tribe a CRDBBYS*

- 690 Cedrela X.
1297 Swietfenni Til*
1298 Chloroxylon Bee.
1299 Flindérsia R.Br.
1300 Carapa Aub.
3344 IValisura Rox.

131. ORDER XLIX. AMPELI'DEJE.

Genera 4, Species 53; Hot-house Species 31; Green-house Species 6; Hardy Ligneous Species 16; Hardy Herbaceous Species 0. ± 13 ft j 0 feet; ^ 0 feet

The vine is the type and representative of this order. *Cissus* and *Arapelopsis* differ little from it in botanical characters, and not at all in habit. The common grape is the only species that bears really good fruit; tv* American kinds, with large fleshy berries, being spoiled by a disagreeable foxy flavor, which is not found to be removed by cultivation. Cuttings.

132. Tribe 1. VINI'FERA, or SAR-MENTA'CEA.

- % 16. herb.0. ±13 feet.
400 Cissus L.

656 Ampelopsis Mx.
655 n'tis Z.

4
12

133. Tribe 2. LEEA'CEJB.
594 Leea L.

134. ORDER L. GERANIA*CE.SL

*Genera 5, Species 366; Hot-house Species S; Green-house Species 276; Hardy Ligneous Species 0; Hardy Herbaceous Species 87. 2 0 feet; £ 15 feet; * 0 feet*

The Geraniaceæ are well known to all gardeners for their beauty, and the facility with which hybrid varieties are produced among them. *Geranium* and *Eridium* are chiefly natives of the northern hemisphere and *Pelargonium* of the southern. Different as they appear from *Vinifera* in most respects, there are so many points in which a curious resemblance may be found between the two orders. The young stems of both are articulated and separable at the articulations; and the lower leaves are opposite, while the upper alternate. In Geraniaceæ no tendrils are produced, but the peduncles are opposite to the leaves, as in *Malvaceæ* and occupy the place of tendrils. M. Dcandolle observes, that of the true Geraniaceæ, some are acid, especially those of which the leaves and bark are succulent; several exhale a resinous smell which is sometimes agreeable, but occasionally so powerful as to be unpleasant. The resinous principle is so abundant

which are DUVI dcunneiu vulnerary, ana in £. moscnaiuin, tr. praisnc, anu others, in wchicn it IS Unneu slight aromatic principle, whence they have been recommended for various purposes, and among others i removing calculous disorders. The astringent property of the geranums is also present in *G. maculatum*, which grows in much abundance about Philadelphia; the root of this plant, boiled in milk, is used for cholera in children. Barton is of opinion, that it would be a good substitute for gum kino in nephritis ft cholesterina. Cuttings, division, or seeds; or sometimes by cuttings of the roots.

135. Tribe 1. GERANIE.JE.
lig. Q. herb. 87. £15 feet
1992 Sarcocallion Dec.

- 1991 Monsbn/j I*
1932 Geranium Herit.
1927 Ertidium Jfer*.

136. Tribe S.
1929 Pelargonium Herit.

137. ORDER LL THOPAO'LEJE.

*Genus 1, Species 11; Hot-house Species 1; Green-house Species 7; Hardy Ligneous Species 0; Hardy Herbaceous Species 3. ± 0 feet; £ 3 feet; * 0 feet*

In which the peculiar acrid flavour of Cruciferae is found to exist Tropeolum pentaphl. Hum, with P other species, is a powerful antiscorbutic. All are natives of shady places in various parts of South Africa. The roots of some are fleshy and eatable. Seeds or cuttings.

1148 Tropafalum L. 0 3

138. ORDER LIL BALSAMI'NFJE.

*Genera 2, Species 19; Hot-house Species 2; Green-house Species 9; Hardy Ligneous Species 0; Hardy Herbaceous Species 8. ± 0 feet; £ 7 feet; * 0 feet*

The flower of this order has been remarked by a learned botanist to be that of Fumariacca?, the Cxalis, the embryo of Z. Inum, and the habit peculiar. The well-known elastic spring with which are ejected constitutes a principal character of the order. All the species are annuals, with the exception of *Impatiens fruticosa*; they delight in moist hot situations, generally within the tropics, and are remarkable for the singularity and varied colors of their flowers. Seeds, and sometimes by cuttings.

697 Jfolsamlna Use.

I 696 Impatiens Riv.

0 *8

139. ORDER LII. OXALIDE^A.*Genera 3, Species 112; Hot-house Species 8; Green-house Species 93; Hardy Ligneous Species 0**Hardy Herbaceous Species 11. 10 feet; £ 6 feet; * 0 feet*

*m*l¹⁰?TM confounded with the last order. It is the opinion of modern botanists, that the species are more nearly allied to Rutaceae or Zygophyllaceae, and that their character and peculiar habit are quite sufficient to distinguish them. The beauty of the genus *Oxalis* is very great, and the readiness with which the species are cultivated and caused to flower, would have been expected to make them universal favorites; they are however, much less popular than many others. Their properties are well known; all of them have a slightly acid taste, and depends on that exists in great abundance. Several species are employed in Brazil as medicine. Most commonly by division, offsets, or seeds, but also by cuttings.

0 *10

140. ORDER LIV. ZYGOPHY'LLEJE.

*Genera 7, Species 32; Hot-house Species 8; Green-house Species 22; Hardy Ligneous Species 0;**Hardy Herbaceous Species 2. ± 0 feet; £ 3) feet; ^ 0 feet*

The hardness of the wood of the shrubby species of this order is most remarkable, if the softness of the beech and herbaceous ones is remembered. To this the extreme difficulty of propagating *Gualacum* is to be attributed. *Zygophyllum Falago* is employed as an anthelmintic, but it is in the *Guaiacum* that the great medicinal virtues of the order are found: all the genus is extremely exciting; the wood and bark of *Guaiacum officinale* and *sancutum* have a rather bitter acrid flavor, and are principally used as sudorifics, *guaiacum* or *aloes* alternatives; they have been found to contain a particular substance differing both from gum *guaiacum* which has been called guayacine. Many of the species bear beautiful flowers, especially the *Nobilissimum*, which, with their brilliant yellow Cistus-like blossoms, enliven many an arid waste in the tropics, "are found in the colder latitudes of the world. Cuttings, division, or seeds.

J ^{ulus} TOU. agouti Tou. Urrea Cav.	1304 Zygophyllum L.	1155 Porlier/o R. & P.
	1303 Guaiacum L.	1704 Meli&ithus L.

141. ORDER LV. RUTACE-E.

*Genera 37, Species 162; Hot-house species 18; Green-house Species 137; Hardy Ligneous Species I;**Hardy Herbaceous Species 4. ± 22 feet; £ 5 feet; * 0 feet.*

Extensive, but rather heterogeneous, group of plants, natives of all countries and all situations. The species are either fetid northern herbaceous plants, as the garden rue, or neat heath-like southern shrubs, with aromatic odor, as the Cape Diosmas; broad or long-leaved Australian shrubs, with a stellate inflorescence, as *Phedium*, or tropical trees with panicles of pallid minute flowers, as the *Cuspanas* and *Zanthoxylum*. Medical properties of many genera are considerable. Ifuta and *Vaganum* are emmenagogue, stimulant, and sudorific. Cape Diosmas abound in a volatile oil of an agreeable smell, but acrid flavour; other species are reputed antispasmodics. The *Zanthoxylums* are said to possess acrid, stimulating, *W* qualities; *Clava Herculis* and *Rraxineum* are said, in America, to be powerful sudorifics and diabetics. According to Barton, they possess a remarkable power of exciting copious salivation, not only *H* *PPK* to the mouth, but even when taken internally; they have both been found powerful remedies *VQ* *ara*ysis of the muscles of the mouth. *Zanthoxylum caribaeum* is regarded in Guiana as a detergent *bals* and febrifuge. *Bruxa* is used as an astringent in dysentery. The famous febrifugal *Angostura* is the produce of *Cusparia febrifuga*. Cuttings, division, or by inarching.

Tribe 1. 3ft; <i>g</i>	3303 Colconema B. Sf W.	144. Tribe 3. CUSPARIEAE.
	3304 Macrostylis B. % W.	678 Spirantha St. Hel.
	2619 Empleurum H. K.	48 Galipea Aub.
3. DISSME AUSTRALISICA.	1134 Corr^a Sm.	2046 Monnieria W.
132 Phellium Ven.	1933 Philotheca Rud.	145. Tribe 4. ZANTHOXY'LEJK.
1310 Crowes Sm.	1310 Eriostemon Sm.	lig. 5. herb. 0.1 & 1 feet
1152 Borbini Sm.	399 Zieria Sm.	2758 Bruxa Herit
909? Poranthera Rud.	1191 Melicope Forst.	2764 Zanthoxylum L 3
§ 4. DISSME AMERICANA.	396 Evodia Forst.	398 Faglura L.
1190 Elkphrium Jac.	1190 Elkphrium Jac.	686 Toddahaj.
1313 Choisyia H. & B.	1317 Simaruba Aub.	391 Ptelea L. 1
		2876 Aildntus Dcsf. 1
		2759 Tetradium Tou.

147. ORDER LVI. SIMARUBACEAL

*Genera 3, Species 5; Hot-house Species 5; Green-house Species 0; Hardy Ligneous Species 0;**Hardy Herbaceous Species 0. f 0 feet; £ 0 feet; ^ 0 feet*

Plants, found in equinoctial America, constitute this order. They are trees or shrubs with an aromatic and bitter taste, a milky juice, and pinnated leaves. *The Simaruba officinalis* is well known as the most *gr* *oil* *U881a* hitherto discovered; the same property exists, in a milder degree, in the rest of the 1316 r. *amara* is a very ornamental plant, but rare, at present, in collections. Cuttings.

1317 Simaruba Aub. | 1318 Simaruba Hil.

IV. Fruit (gynobasic) inserted into a fleshy Receptacle, with which the Style is continuous,

148. ORDER LVI. OCHNATEJE.

*Genera 3, Species 15; Hot-house Species 14; Green-house Species 1; Hardy Ligneous Species 0;**Hardy Herbaceous Species 0. ± 0 feet; £ 0 feet; * 0 feet*

Kalmia yellow-flowered tropical shrubs with lucid leaves. The roots and leaves of *Walkeria serrata*, a *antifl* *an* *ter*; a decoction of them, either in water or milk, is used in Malabar as tonic, stomachic *h* *h* *2* *u* *c*. The bark of *Gymnia hexasperma* is found useful in healing sores produced in cattle *inf* *nesting* of insects. Cuttings.

1582 <i>Lycn</i> * Schreb.	1315 <i>Gomphia</i> Schreb.
744 Walkeria Schreb.	1151 Castek Turp.

149 ORDER LVII. CORIARIE\&

**Genus 1, Species 2; Hot-house Species 0; Green-house Species 1; Hardy Ligneous Species 1:
Hardy Herbaceous Species u, fl foot; £10 feet; £ 0 feet**

Five species constitute the whole of this order, distributed in South Europe, New Zealand, Peru, and Mexico. They possess no beauty, and are only interesting on account of their problematical station in a botanical arrangement. The leaves of *C. wyrtofilla* are astringent, and are employed in dyeing black. The berries are very poisonous. On one occasion, during the Spanish war, fifteen French soldiers were taken up after eating them, and three died from their powerful narcotic effects. Cuttings and suckers.

2797 *Coriaria* L. 1 0

SUBCLASS II. CALYCIFLO'IE. Petals separate, inserted in the calyx.

*Genera 1149, Soecies 11435; Hot-house Species 2089; Green-house Species 3486; Hardy Ligneous Species 1&c;
Hardy Herbaceous Species 4552. ± 283 feet j f 15dfee^ * 7 feet*

150. ORDER LIX. CELASTRI'NE-E.

Genera 18, Species 114; Hot-house Species 19; Green-house Species 53; Hardy Ligneous Species 42; Hardy Herbaceous Species 0. f 46} feet; £ 0 feet; ^ 0 feet

This order differs from the succeeding, in having the stamens alternate with the petals; the sepals imbricated in aestivation; and the ovarium wholly superior. It consists entirely of shrubs or small trees, with simple, rarely compound, alternate or opposite leaves, and inconspicuous flowers of a greenish or white colour. Several are favourite ornaments of our shrubberies, as the *Staphylea*, the *Celastrus*, and the *Euonymus*; the latter of which is valued on account of its beautiful-colored fruit. The fruit of *Euonymus europaeus* is a purgative, as is also the inner bark, and in strong doses powerfully emetic. The famous Paraguay tea *Urtica* foliage of a species of *Urtica*. The bark of *Prinos verticillatus* possesses such active, astringent, bitter, tonic and febrifugal qualities, that it is used in North America, with success, as a substitute for *Cinchona*. A decoction of the twigs of *Maytenus boaria* is used to bathe the swellings produced by the poisonous shade & the tree *Litsea*. Cuttings, suckers, layers, and seeds; sometimes by grafts.

151. Tribe 1. STAPHYLEAEJS. <i>lie.</i> 2. <i>herb.</i> 0. 114feet	[^] j <i>Maytenis</i> MoL & Feu.	664 <i>Hartftgfe</i> Thun.
897 <i>Staphylifa</i> L. *2	663 <i>Alzatta</i> R. & P.	393 <i>CurtWa</i> H. K.
2886 <i>Dalfr</i> mplea Kox.	672 <i>Eckoldndrum</i> Jac	410 <i>MygindJaC</i> .
152. Tribe 2. EUONY'MEJL <i>tig.</i> 15. <i>herb.</i> 0. ± 8 feet	394 <i>Ptelidium</i> Thou.	?804 <i>Bumalda</i> Thun.
666 <i>Euonyminus</i> Tou. •IS	9 1496 <i>FitBtidia</i> Com.	411 "/lex <i>L</i> •13
«62 <i>Celastrus</i> L. 2	153. Tribe 3. AQI;IFOLIA*CE£. <i>lig.</i> 95. <i>herb.</i> Q 143Jfeet	1089 <i>Vinos</i> L. 10
712 <i>Plectrdnia</i> L.	895 <i>Cassine</i> L.	2775 <i>Nemop&nthes</i> Rafi. 1

154. ORDER LX. RHA'MNEAL

*Genera 18. Species 171; Hot-house Species 32; Green-house Species 87; Hardy Ligneous Specks 511
Hardy Herbaceous Species 1. f 79*feet; £ 3 feet; £ 0 feet*

In habit, this altogether agrees with the last, from which the medical properties of the species are not ~~J2~~ different. Throughout the order, as far as it has been examined, there is a remarkable agreement betw^{een} the fruit and the inner bark, especially in *Ichnamnus catharticus*, *Frdngula*, and others, in which they both are purgative and emetic. Some, as the Jujuba and the African Lote, nevertheless, yield a wholesome and "Sree" fruit; and the berries of the greater number yield^{*} under the chemist's hands, green or yellow dyes of m^{any} importance in manufactures. The leaves of *Ahamnus Theeians* are substituted for tea by the poorer, uⁿ among the Chinese. The bark of *Ceanththus cstrtus* is esteemed in Mexico as a good febrifuge. Cutting*

zyphe	4	0	3298	<i>Retanula Brong.</i>			3311	<i>Soul&ngia Brong.</i>
659 <i>Paphirus Tou.</i>	2	0	668	<i>Ceanothus L.</i>	7	L	711	<i>Cryptandra Sm.</i>
658 <i>Betchem/a Neck.</i>	1	0	3300	<i>Colubrina Brong.</i>			3299	<i>Trebla Cav.</i>
657 <i>Ahamnus L.</i>	•37	0	3301	<i>WillemettVi Brong.</i>			2871	<i>Gouan/a L.</i>
661 <i>Conch'a Cav.</i>			670	<i>Pomadlris Lab.</i>			691	<i>Hovtnia Thun.</i>
667 <i>Colletia Com.</i>			710	<i>Phyicula L.</i>			2757	<i>Schceffelia Jac.</i>

155. ORDER LXI B R U N I A ^ E J E .

Genera 7, Species 30: Hot-house Species 0; Green-house Species 30; Hardy Ligneous Species 0; Hardy Herbaceous Species 0. £ 0 feet; £ 0 feet; £ 0 feet

Small heath-like shrubs, all natives of the Cape of Good Hope, and extremely ornamental, both in and foliage. Their properties are unknown. Cuttings.

6f2 Brtmm L. &00 Berzelia Brong. 3306 Raspalfa Brong.	I 3309 Staavia Thun. I 3307 Berford/a Brong. I	I 803 Lincknia I* I 3308 Audouinia Brong.
---	--	--

156. ORDER LXII. SAMY'DEJE.

Genera 2, Species 14; Hot-house Species 14; Green-house Species 0; Hardy Ligneous Species 0;
Hardy Herbaceous Species 0. ± 0 feet; £ 0 feet; £ 0 feet

Haray Herbaeae Species 6. ± 0 feet; ± 0 feet; ± 0 feet
 Tropical shrubs or small trees, with entire, stipulate, alternate leaves, covered with pellucid dots, *
 axillary flowers of little show. Some of the species of Samyda are pretty, but very rare. Their P^dP^jJ^l on
 are unknown. M. Decandolle remarks, that in their fruit they approach Bixineae and Flacourtiaceae i z^gt.
 account of the position of their stamens must be arranged in the vicinity of Rhamneae and Rosaceae. Cum **

1368 Samvda L. 11369 Casearia Jac.

157. ORDER LXII I. HOMALI'NEA.

Genera 4, Species 8; Hot-house Species 5; Green-house Species 2; Hardy Ligneous Species 1; Hardy Herbaceous Species 0. J 3 feet; £ 0 feet; ^ 0 feet

Evergreen handsome shrubs, with alternate leaves and deciduous stipule; they are readily known by ^{the} *Parietal* placenta?, an unusual character among the orders that surround them. Blackw^{ell}i/71/agif ^{U.S.} bunches of starry white fragrant flowers. *Anstotelia* is an evergreen half-hardy shrub, with eatable Little is known of their medicinal properties; the root of *Homalium Racauca* is used in Guiana as a c fonorrhœa. Cuttings or layers.

1595 Homálium Jac.	11146 Astranths Lou.
1465 Blackw<lli7z Com-	11*38 AristoteJw L. 1 0

158. ORDER LXIV. CHAILLETIA^CEJE.

Genus 1, Species 1; Hot-house Species 0; Green-house Species 1; Hardy Ligneous Species 0;
Hardy Herbaceous Species 0. ± 0 feet; £ 0 feet; & 0 feet

This order contains only three genera, Chailletia, Leucosia, and Tapiura. It agrees with Teiebinthacea in having 8 furnished with a corolla and calyx. Fruit furnished with a dry covering, including a two or three celled

Cuttings.

815 Chailletia Dec.

159. ORDER LXV. AQUILARI'NE.

Genus 1, Species 1; Hot-house Species 1; Green-house Species 0; Hardy Ligneous Species 0;
Hardy Herbaceous Species 0. f 0 feet; £ 0 feet; £ 0 feet

the plant* contained in this order are very imperfectly known. It differs from the preceding in the seeds being erect, not inverted, as well as in the capsules being 2-valved. Trees with alternate entire leaves, natives of Asia. Cuttings.

1370 Aquilaria Z.

16a ORDER LXVI. TEREBINTHA^CEJEL

Genera 27, Species 142; Hot-house Species 58; Green-house Species 62; Hardy Ligneous Species 22;
Hardy Herbaceous Species 0. 140 ft.; £ 0 feet; £ 0 feet

in the order notwithstanding the labours of several botanists, in a very confused state; few of them of knowledge of many of the genera, which have been hitherto imperfectly described, it is difficult to determine the value of the characters assigned to the tribes, or the dignity of the tribes themselves. In this order species are shrubs or trees, with alternate exstipulate leaves, and inconspicuous flowers. They oalsamiferous resin, which is chiefly present in the leaves and bark, and from which J. M. JJ. wder has been derived. Notwithstanding the minuteness of their flowers, many of the species are valuable as ornamental plants, on account of the beauty of their foliage, others for the sake of their utility in arts or JSJd'tme, and others for their fruit. The Casewh and the Pistachio are valuable for their nuts, which are well known articles in the markets of Europe. The Spondias and Mango are equally famous in the tropics, iff. well-known balsam of Mecca is the produce of the Balsamodindron gileadensis and balm of Acouchi, ati. heterophylla; gum comes from Amyris elemifera and Icica leptophylla; mastich from Pistacia lentiscus and *L. lentiscus*; resin from Euphorbia cantharoides; Echinus Molle produces a resin, which the order chiefly consists, he concealed others in which acrid and poisonous qualities abound. Such are several species of herbs, the juice of which produces blisters upon the skin. To conclude this long list of the uses and dangers of Terebinthaceæ, the bark of *Chios galba* is used as a rubefacient and as a mordant for red dyes, and that of Ahus Cibaria as a powerful means of tanning the skins of animals. Cuttings, layers, and seeds.

161. Tribe 1. ANACARDIJE or CASSUVIEE. W. a herb. 0. 117* ft.	2649 <i>Duvatia</i> Kth 2799 <i>Echinus</i> L. ? 1408 <i>RobergM</i> Schub.	3421 <i>Sorindeia</i> Thou. 1332 <i>Gariga</i> Rox.
Anacardium Box.	163. Tribe 3. SPONDIA^CEA	165. Tribe 5. AMYRIDEJB.
Anafera L.	1405 <i>Spindias</i> L. 1407 <i>Poup&ru'a</i> Com.	1168
Buchanania Roz.	164. Tribe 4. BURSBVUVBA.	166. Tribe 6.
Pistacia L.	1333 <i>Boswellia</i> Rox. 1169 <i>Balsamodindron</i> Kth. 1334 <i>Icica</i> Aub. 2895 <i>Birserm</i> Jac. 1170 <i>Hedwigia</i> Sws.	106 <i>Chehrum</i> L. 896 <i>Spathelia</i> L.
Comocladia L.		167. Tribe 7. CONNARAYEC
Picritmia Sum.		1937 <i>Connarus</i> L. 1936 <i>Omphalibium</i> Gae. 1403 <i>Crestis</i> J.
Tribe 2. SUMACH(NE& g. 19. herb. 0. ± 37 feet.	19	

168. ORDER LXVII. LEGUMINO'SAL

Genera 244, Species 2626; Hothouse Species 886; Green-house Species 675; Hardy Ligneous Species 177;
Hardy Herbaceous Species 888. f 117 ft.; £ 58) feet; i 0 feet*

the various kinds of pulse belong to one of the most familiar to the world, and at the their full one of the most useful to mankind. Their papilionaceous flowers characterise a large number, and As oblong and pinnate leaves the remainder, with a few exceptions which it is not necessary to particularise, the RAVT of ornament, many are possessed of unrivaled beauty, for example, among hardy flowering trees, tribe of the Xabdrnum; among shrubs, for decorating the borders of the flower-garden, the various Chilii, *Caragana*, *Cbitica*, *Amorpha*, and others; among hardy climbers, the far-famed *Wistaria* of last! * and its sister of North America, with the species of the herbaceous genera *Picia* and *Xathyrus*; and, among hardy herbaceous plants, the numerous species of *Lupinus* and *Astragalus*. Great, however, a beauty of the Leguminosa which can brave the inclemencies of the seasons of Northern Europe, it must way before the splendor and elegance of their brethren of the tropics. The flowers of the *Erythrina*, or *Mimosa*, with their twin leaves, hang in festoons of flowers from branch to branch, and are only less vigorous and elegant, but more richly colored blossoms of the *Mucunas*. But all these,

While the forests of not countries are in a state of wildness, the meadows and pastures of the same latitudes are enamelled with the flowers of myriads of *Hedysarums*, *Eschscholtzias*, and animated by the wonderful motion of sensitive plants. As in our own country, the Ravest *Hu* energy is in many places indebted to the yellow flowers of our furze and broom, to in other nes the same effect is produced by other genera of Leguminose; by *Lipana*, *Borbonia*, and *ipsa* Uthut

at the Cape of Good Hope, and by the Pultenaas, Daviosias, Aotuses, and multitudes of similar genera to New Holland. The wood of the order is very hard and durable, with a yellow tinge, sometimes changing into green, as in the Laburnum of Europe, and in the better known Brazil wood of commerce, produced by Caesalpinia. The following useful remarks upon the properties of the order are made by M. Decandolle: —

" The family of leguminous plants, though established upon characters of primary importance, offers, nevertheless, so large a number of species and such singular botanical anomalies, that it is easy to foresee that their properties will exhibit little uniformity. Still more exceptions may be anticipated, if one reflects that the chemical principle which is found most abundantly in every part of leguminous plants, and to which we must attribute their principal properties, is the extractive. It is probable that this principle, either from its own nature, or from its peculiar power of uniting with different matters, or perhaps instead of being a simple principle, it is rather a compound of different matters; it is probable, I say, that the extractive principle exhibits much less uniformity in its results than any other. It is, without doubt, to the presence of the extractive principle, in considerable quantities, that many leguminous plants owe their purgative properties, which are common to several extracts, and which many chemists attribute to the acetate of potash, which they are almost universally found to contain. Thus the leaves and foliaceous pods of *Cassia Senna* Lin., of *Cassia lanceolata* Forth., of *Cassia emarginata* of the Antilles, of the *Cassia marylandica* employed in the United States, of *Colutea arborescens*, of *Genista phragmites*, and perhaps also of *Coronilla Eremus*. act as brisk purgatives, and often cause wind and pain in the bowels. The juice of *Coronilla varia* excites vomiting, and may even become

Juicy pulp that exists in the *Cassia fistula* Lin., in the *Tamarindus indica* Lin., in the *Ceratonia siliqua* Lin., and probably in the *Inga vera* and *I. punctata*, which are eaten in small quantities in the Antilles, but which, taken more copiously, would have the same effect as our Carobs. There are some fruits of Leguminosae, for example, the *Sophora* and the *Gleditschia*, with tumid pods, in which is found a juice which surrounds, it is true, the seeds, as in the plants just mentioned, but which differs from them altogether in a very astringent and nauseous flavour; the nature and properties of this juice deserve to be examined by chemists, and would undoubtedly throw some light upon the nature of Leguminosae. I am induced to think that the astringent Juice of the *Sophora* is a secretion of the pericarp, whilst the sweet and purgative juice of the *Cassia* would be a secretion of the external part of the seed; but this hypothesis requires to be verified, what leads me to this opinion, is the flavour commonly found in husks in the Carobs, for example, the hull is astringent, and the pulp sweet and laxative. But let us return to the properties which may be attributed to the extractive principle. It is undoubtedly from some one of these modifications that the singular property of the Piscidia and many Galegas is derived, which are employed in America to stupefy fish, which are taken when this means as readily as Nux vomica. The decoction of the root of *Galega virginiana* is considered in America as a powerful vermicifuge. It is, perhaps, indeed, to the very same cause that the rubefacient power of the fresh leaves of some Leguminosae are to be ascribed, which act readily upon the skin if applied as plasters I

found in small quantities, the seed may serve as food for man and animals, as we see in French beans, pea-lentils, chick peas, beans, and many others, &c. If found in a more considerable quantity, it will render them purgative or emetic, as in the *Cytisus Laburnum*, the "nagris fœtida", and even in most Coronillas. It is remarkable that the botanical characters of Leguminosae should so strictly agree with the properties of their seeds: the latter may be divided into two sections; namely, first, those of which the cotyledons are thick and filled with fascula, and destitute of cortical pores, and which, moreover, in germination do not undergo any change, but nourish the young plant by means of that supply of food which they already contain; secondly, those of which the cotyledons are thin, with very little fascula, and furnished with cortical pores, and which change at once into leaves at the time of germination, for the purpose of elaborating food for the young plant. All the seeds of the first section are employed as food in different countries; none of those of the second section are ever so employed: the Cajan, which has long been classed among the Cytisi, was apparently an exception to this general rule; but observation has proved the contrary. Bearing in mind its known properties, I formerly paid particular attention to its structure, and I have shown in a note which accompanies my catalogue of the Montpellier garden, that the Cajan forms a particular genus much more nearly allied to the French bean than to the Cytisi, and that it, in fact, belongs to the first of the sections which I have just described. The seeds of Leguminosae present also many other anomalies more difficult to reduce to any

or *Coumarina odura* of Aublet, which, under the name of Tonquin bean, is used for perfuming snuff: there are others which, like the chick pea, have rather a bitter taste and exciting properties, and are so bitter as to be used in Java and Brazil as tonic, alexiteric, and vermicifuge. In a word, are not the aperient and diuretic properties which are observable in the herbage and the roots of many leguminous plants, such as broom, beans, *Ononis*, *Moringa pterygosperma*, *Ifathyllis cretica*, &c., to be attributed to a modification of the extractive principle? There are, in another view, roots which are furnished with tubercles, that is to say, with reservoirs of faecula, which furnish mankind with wholesome food, as we see in the *Lathyrus tuberosus*, which is eaten in Holland, the *Dolichos tuberosus*, and the *D. bulbosus*, which the Indians use as food. The roots of the liquorice have a sweet and mucilaginous taste, which is well known by every body, and which, according to Vauquelin, an acrid and rather exciting principle, causes it to be employed as a pectoral; the analysis of this root published by M. Robiquet, proves that, independently of its woody skeleton, the same kind of amyloseous fascula is found as in the tuberous roots of which we have just been speaking; it is thence seen that the acrid flavour of decoctions of liquorice depends on the small quantity of resinous oil which it contains, and that

for their bitterness, and are used as febrifuges; the different kinds of *Geotryba* and *Andira* possess this bitter and febrifugal quality in a remarkable degree; in India, the bark of the *Agati grandiflora* and of *Guilandina Bondiella* are employed for the same purpose. The barks of many leguminous plants are also remarkable for their astringent qualities, caused by the quantity of tannin which they are found to contain; this is observable in the *Acacia Catechu*, and in the *Acacia arabica*, which is used for tanning leather and elsewhere. It is well known that almost all colouring matter proceeds from the extractive principle; as it appears that this principle abounds in Leguminosae, we ought to find in them a considerable number of the colors which are used by dyers: to this family, in fact, belong the principal blue colors, such as the name of indigo, extracted from every kind of *Indigofera*, and from some Galegas; and the red color which are yielded by all the species of *Caesalpinia* and of *Haematoxylon*. We may add the red juice, drawn from the *Pterocarpus* and *santalinus*, under the name of sandal and of dragon's blood; *Biklea frondosa*, under the name of gum lac; and also from *Dalbergia Monetaria*. These juices differ in many particulars, but their history and analysis are at present so far from being known, that it is impossible to form a true estimate of the nature of their differences. But anomalies of this nature are far from being confined to the plants just mentioned. Among the exotic drugs employed in the art) they are common: such, for example, are the balsam of Capivi, produced by the *ropoifera*; the balsam of Peru,

Jfuttg says, If obtained from Myrosp^rrum perufferum, and the balsam of Tolu from M. tolulenims th* Cachou, which has been found to be almost pure tannin; and which is supposed to be produced by Jctfcia SSF*, of the "ame character is that remarkable resin that is yielded by Hymen* *aCoHrbaraj* gum Arabic, Produced by the bark and roots of *Acacia vera*, Senegal, árabica, and others; gum tragacanth obtained from ~~and~~ *gaulis criticus* » ?emmer, and vdrus; and finally, manna, secreted by *Utaga Maurtrum*.¹ Cuttings

The arrangement of Yhifo tribe of plants has been found to be attended with much difficulty. By Linnaeus, and the writers who succeeded him the number of genera was much smaller than those of the present age; many additions have been made in consequence of the discovery of New Holland, and a large number of subdivisions in old genera have been from time to time introduced. To combine these scattered improvements under one uniform system has lately been attempted by the learned Sⁿist, from whom the foregoing extract has been taken. M. Decandolle's system is to be useful for explaining the principles upon which it is founded. He divides Leguminosae into two families, Jivwions, the first of which consists of plants the radicle of whose seed is curved, and Jevyedons, and the second of those whose radicle and cotyledons are straight. The latter RECTIMBRIE. In the Curvibrute, certain diversities in the structure of the calyx and corolla again divide into two principal forms: one of which, comprehending all the genera with papilionaceous flowers, is Papilionaceae, and the other, consisting of a very small number of species, with one or two petata OT, and an obscurely lobed calyx, is called Swartzieae. The last is not subdivided, but the Papilionaceae themselves into the two great tribes pointed out by M. Decandolle, namely, those with fleshy cotyledons and eatable pulse, Sarcolobat; and those with foliacous cotyledons and seeds which are not eatable, globae. Each of these is divisible by three, upon slight differences in the fructification. In Recumbritae suborders, Mimicæ and Cesalpiniæ, are founded upon variations in the estivation of the calyx and corolla. In the former, it is valvate, in the latter, imbricated; the first constitute a single tribe, the Utitae, distinguished by less momentous peculiarities of structure. Having premised thus much, the following tabular explanation will be intelligible: —

i59. Div. I. CURVE'MBRI ^{ffi.}	1962	5partium L.	2	0	2101	Oxytropis D. c.	*0	29
j70. SUBORDER I. PAPILIO-	1963	Genista L.	38	0	2102	Astragalus /.	*\$	98
Subdiv. i. Phylobas.: Tribe]. SOPHO'BEJR.	1964	Cytisus L.	33	0	2103	Uvidenbta ^{dtni} FIJI	0	1
Jac.	1965	Adenoctrpus Dec	4	0	2104	Biserrula L.	0	1
gg JUwardsi Vi Sal.	1966	Ononis L.	5	29	17ft	Tribes. ■■■■■		
gg Orpin&cia Jack.	1967	Requienia Dec.			180. Subtribe 1. Coronillex.			
log Y?**" Lam.	1968	Jahillis L.	0	14	lig. St. herb. £9. % 4 ft. j £ 9 ft			
Iffca *faer <itropis Dec.	175. Subtribe 2. ■■■■■				2105	Scorpiurus L.	0	6
SJ? U'Panthus Swt.	176. Subtribe a Clibria.				2106	Coronilla L.	2	6
Fenn&psis R. Br.	lig. O. herb. 1. 10 ft; £ &				2107	Astrolomia Dcvs.	0	4
SMsia Yen.	2061 Medicago /..	•1	75		2108	Ornithopus /.	»0	4
Ven.	2062 Hymenorraps Sam	0	3		2109	Hippocratea /.	*0	8
J** Br.	2063 Trigon'illa /.	0	27		2110	Secuniera Dec.	0	1
Ven.	2064 Pocockia Ser.	0	1	181. Subtribe 2. Euhedyturea.				
R. Br.	2065 Jilelliptus Tou.	•1	27	lig. 2. herb. 75. m ft; £ 19 ft.				
^ompholobium Sm.	2066 7Yilghun Tou.	•0	119	2111 Diphaca JLou.				
?urtbn ^a R. Br.	2067 Jorficum Tou.	3	4	2112 Pictetii Dec.				
Jackson/o R. Br.	2068 Lbus L.	0	27	2113 Ormocarpum Beauv.				
24R	2069 Tetragonalobus Sco.	0	5	1976 Amichi H. ft B.				
1257	3595 Hoskcka Dou.	0	2	1977 Poiretii Ven.				
Ma R. Br.	2070 Carmichaelia R. Br.			2114 Myriadenum Desv.				
clerothamnus R. Br.	1969 Cyam&psis Dec.	u	x	1978 Zom/o Gm.				
Gastrok.ium R. Br.	176. Subtribe a Clibria.			1979 Stylosanthes Szrz.				
Euchilus R. Br.	2071 Psoralea L.	T)	5	1980 Adesmia Dec.				
Pulten ^a Sm.	2072 Indigofera X.			2115 JEschyn6mene Z.	0	1		
pavieata Sm.	2073 Clitoria L.			2116 Smiths H. K.				
Mirbelia Sm.	2074 Nucrocalpum Desv.			2117 Uurea Neck.				
J73. Tribe 2. ■■■■■	2075 Colgania Kth.			2118 Uraria Desv.				
Subtribcl. Genistcar.	2076 Gal.ktu Br.	0	2	2119 Nicolsbim Dec.				
«. f 49 ft j 1ft10ift	2077 Vilinorin/fl Dec			2120 Desm ^c diump Dec.	0	1	L	
R. Br.	2078 Barbierw Dec			2121 Dic ^c raia Dec.				
1972	1970 Pueraria Dec			2122 Taverneira Dec				
1257	2079 Dumasia Dec			2123 //cdjsarum L.	1	28		
24R	2080 Gifcinc L.			2124 Onobrychis Tou.			SO	
1257	2081 Chictocalyx Dec.			2125 Kleiberts Dec.				
1257	177. Subtribe 4. Ga&gcte.			2126 Lepidopoda Mx.	1	11		
1257	lie. 45. herb. 24. f 841 a ;£13 Jft			1981 f'bénus L.				
1257	1922 Petalostemnum Mx.	0	4	2127 Flemingia Rox.				
1257	1971 Dalea L.	0	4	182. Subtribe a Alhdgea.				
1257	2082 Glycyrhiza Tou.	0	8	2128 Alhagi Tou.				
1257	1972 Gafrag Tou.	0	6	2129 Alysicarpus Neck.				
1257	2083 Tephrosia Pert.	0	1	2130 Bremontiera Dec				
1257	1973 Amorphal.	3	0	183. Subdiv. it Sarcobolte.				
1257	2084 Nisibliri Jac			184. Tribe 4. VICIE ^a .				
1257	1974 Mtllera L.			lig. a herb. 190. ■■■■■ ft. j £ 28 ft.				
1257	2085 Lonchocarpus H. & B.			2131 Cleer Tou.	0	1		
1257	2086 Robinia L.	17	0	2132 /'aba Tou.	0	2		
1257	2087 Sesbania Pera.			2133 Ticia Tou.	»0	74		
1257	2088 Agtii Rh.			2134 ^rvun /.	»0	10		
1257	2089 Glottidium Desv.			2135 Fisum Tou.	»0	12		
1257	1975 Piscidia /.			2136 7-athyru L.	»0	51		
1257	2090 Daubent5nia Dec			2137 OVhrus Pert.	0	1		
1257	2091 Caragana Lam.	16		2138 C'rohus Tou.	»0	97		
1257	2092 Halimodindron Fit.	3		2139 Plat ^c btylis Swt.	0	9		
1257	2093 Diphsysa Jac.			185. IYibefi. PHASEOF ^a .				
1257	2094 Cabphaca Fit.	1		lig. 3. A ^c ft. 71. 1SJ ft.; £ 19 ft				
1257	2095 Colutea L.	5		2140 AM ^c X				
1257	2096 Spherosphysa Dec.	0		2141 Sweel/Vi Dec.				
1257	2097 Swainson/o Sal							
1257	2098 Lesserti/i Dec.							
1257	2099 Sutherland ^a H. K.							
1257	178. Subtribe 5. Astraghe [*] .							
1257	lig. 8. herb. 139. t 5 ft.; £ 20* ft							
1257	2110 Phaca /..	0	10					

NATURAL ARRANGEMENT.

2142 <i>Micranthus Poir.</i>	187. SUBORDER II. <i>SWARTZI&A3., or Tribe 7.</i>	2884 <i>Gleditschia L.</i>
1982 <i>Rothia Pers.</i>	1591 <i>Swartzia W.</i>	2800 <i>Gymnocalamus Lam.</i>
1983 <i>Terfimus Br.</i>	1291 <i>Baphia A/.</i>	1278 <i>Gullandina J.</i>
2143 <i>Amphicarpa &/&</i>		1277 <i>Coulteria Kth.</i>
2144 <i>Kenned/a Ten.</i>		1275 <i>Cssalpinii Plu.</i>
2145 <i>Rhynchosia Lou.</i>	188. Div. II. <i>RECTE'MBRI.E.</i>	1285 <i>Erythrophleum A/.</i>
2146 <i>Fagilia Neck.</i>	189. SUBORDER III. <i>MIMO*-SEM, or Tribe 8</i>	1274 <i>Poinciana L.</i>
2147 <i>Wistaria Nut</i>	<i>lig.i. herb.O. 110 feet</i>	1276 <i>Reichardia Roth</i>
2148 <i>^ i o s Boer.</i>	2838 <i>Entiida Adan.</i>	1280 <i>Hoffmannseggii Cav.</i>
2141 <i>Phaseolus L.</i>	2834 <i>Mimosa L.</i>	1287 <i>Haematoxylon L.</i>
2150 <i>A'mphodus Lindl.</i>	1286 <i>Gagnebinia Neck.</i>	1273 <i>Parkinsonia L.</i>
2151 <i>Sbla Moen.</i>	2833 <i>I'nga Plu.</i>	1282 <i>Cddia Forsk.</i>
2152 <i>Dolichos L.</i>	1939 <i>Parke R. Br.</i>	2885 <i>Ceraonia L.</i>
2153 <i>Vigna Savi.</i>	2835 <i>Schrankskfo W.</i>	1290 <i>Hardwickia Rox.</i>
2154 <i>Lablab Adan.</i>	743 <i>Darlingtonia Dec</i>	1139 <i>Jonesta Rox.</i>
2155 <i>Pachyrlizus Rich.</i>	2836 <i>Desmanthus W.</i>	1289 <i>Tachyleglin Aub.</i>
2156 <i>Parochetus Ham.</i>	1281 <i>Adenanthera L.</i>	1989 <i>Tamarindus L.</i>
2157 <i>Dibclea Kth.</i>	1283 <i>Prosopis L.</i>	1471 <i>Cassia L.</i>
2158 <i>Posphearpus Neck.</i>	1284 <i>Lagonychium Bieb.</i>	1267 <i>Omphalibium Jac.</i>
1984 <i>CaTwalia Dec</i>	2837 <i>Acacia Neck.</i>	1266 <i>Schiitia Jac.</i>
2159 <i>MutAna Adan. ft Dec.</i>		1288 <i>Copaifera L.</i>
2160 <i>Cajinus Dec.</i>		1270 <i>Cynomitra L.</i>
1985 <i>7uplin Tou.</i>	190. SUBORDER IV. <i>CJESAL-</i>	2173 <i>Pariba Aub.</i>
2161 <i>Cylsta H. K.</i>	<i>Tribe 9. GE6FFREJ&</i>	109 <i>OUtea Aub.</i>
2162 <i>Erythrina L.</i>	2169 <i>i'rachis L.</i>	108 <i>Vouipa Aub.</i>
2163 <i>Rudolphia W.</i>	2170 <i>Foandxeia Thou,</i>	1269 <i>Hymeievj L.</i>
2164 <i>Bdte Rox.</i>	2171 <i>ifad/ra Lam.</i>	1268 <i>Bauhinia Plu.</i>
186. Tribe 6. DALBSRGIB'A.	2172 <i>Geoffftya W.</i>	1265 <i>«rcjs /..</i>
2165 <i>Pongamia Ven.</i>	1938 <i>Br6wnca Jac</i>	34 <i>Codarium TaM</i>
2166 <i>DalbVrgia Rox.</i>	2174 <i>Dipterix Schrb.</i>	
1986 <i>Pterocarpus L.</i>		192. DOUBTFUL.
2167 <i>Drepanocarpus Mey.</i>		1990 <i>Phyllolbbium Fis.</i>
2168 <i>Ecastaphy' Hum Br</i>		
1987 <i>jfmerimnum.Br.</i>		
1988-Brya-Br.		
	191. Tribe 10. CASSIB'A.	
	<i>lig. 17. herb. 3. £ 51 ft; £ 5 ft</i>	
	1279 <i>Moringa Dec.</i>	

19a ORDER LXVIII. ROSA^CEJE.

Genera 46, Species 783; Hot-house Species 12; Green-house Species 73; Uarddr Ligneous Species 478;
/forffy Herbaceous Species 219.1199Jfeet; £28* feet ^ 0 feet.

With the exception of Chrysobalanaceae and Sanguisorbaceae, this order is so uniform in its appearance, tb>*, JZosa, the type from which all the other genera are to be considered variations, when justly understood, *>* be found to contain every form of structure which is essential to the order. Having stated this, it will be > once obvious, that if the other genera have such close affinity to /dsas, they must also bear a great analogy f^ beauty. And this is, indeed, the fact. *rfmvgdalu* and *Prnhus* among trees, and *Potentilla*, *f'fem*, *ft"* others, among herbaceous genera, rival the rose in their blossoms, and, in many particulars, surpass that 0^*2 lovely of all flowers in foliage and general appearance. But it is not for charms alone of smell, or blossom, ^ foliage, that this order has fixed itself so high in the estimation of mankind. It has also the rare merit, comprehending all the most important of the fruits of the temperate regions of the world. Thus the apple and the pear belong to *Pyrus*, the apricot to *Armenaca*, the plum to *Prunus*, the peach and the nectarine to ?***?1 *Eriobotrya* produces the loquat, *Mispilus* the medlar, and, finally, the quince is borne by the *Cydonia*. TP* medical powers of many plants of this order are not less active than their fruit is excellent. The principal** these is the well-known Prussic acid, which exists in abundance in the leaves and kernels of many K***?(especially of *Prunus* and *yfm?gdalus*: it is the basis of Laurel water, which, when taken in small doses, ac* either as a violent purgative or as an emetic; and, in stronger doses, is said to destroy irritability without exciting inflammation; these properties, however, although thus dangerous in the distilled water of the l*** and other similar plants, can scarcely be said to exist in any important quantity in the plants in a state J* nature. The kernel of the bitter almond, for example, in which the Prussic acid is more abundant than usu<>, is used for many culinary and other purposes without any bad effect. There are, however, cases in which >> if said to be dangerous to eat the fresh leaves or kernel; as in the *Clarus virginiæ*, the leaves and fruit o\ which are reputed in the United States to be poisonous to certain animals. Besides the Prussic acid, there aij several other principles which abound in the order. All *Drupinæ* yield a gum which is nearly allied to f*** Arabic, and which affords a strong evidence of the affinity that exists between Rosaceæ and Leguminosæ. A great deal of astringency is found in many species, whence different parts have been occasionally employ^ as febrifuges, and as remedies for hemorrhage, diarrhoea, and dysentery. The root of the *Tomentilla* is y* Z fox tanning in the Ferro Isles; and that of the *Capollum* cherry in Mexico. The bark of *Clarus virginia* la mati na a tñrific HTP. in the United States. *Potentilla* fntan» has ho»» grñimi fur thump properties A of tanmn; it has been used DQUL in America ana Europe as a sunstitute tor Jesuit's baric, Inc Dras octopetalia in the north of Europe, of *Jtobus ircticus* in Norway, of *Prunus spinbsa* and *Crasus avia* and of *Aosa pubigibnsa*, have been manufactured into a sort of subsitute for tea. The bark of the *oot

It must not be omitted, that the order Rosaceæ nearly answers to the eosandria of Linneus.

194. Tribe 1. CHRYBOBALAVEJE.	1531 <i>Kerrifl Dec.</i>	10	1526 <i>Fragaria Tou.</i>	0
1503 <i>Chrysobalanus L.</i>	1515 <i>Spiraea L.</i>	*35 14	1529 <i>Tomentilla L.</i>	0
1143 <i>Parin&rium J.</i>	1516 <i>Gilltnia Moen.</i>	0 2	1528 <i>Potentilla L.</i>	4
1431 <i>Grangeria Com.</i>			1527 <i>Cómarum L.</i>	0
653 <i>Hirtella W.</i>			924 <i>Sibbalda L.</i>	6
195. Tribe 2. AMYGDALA*LRE. hg.96. herb. 0. 1	197. Tribe 4. NBURA'DEA.		1458 <i>Agrimonia Tou.</i>	0
1498 <i>i<m^gdalus Tote</i>	1412 <i>Grifelum L.</i>		1209 <i>Aremonia Dec.</i>	0
1499 <i>P^rsica Ton.</i>	198. Tribe 5. DRYA'DBA.		3349 <i>Horkelta Cham.</i>	0
1500 <i>j4rmenaca Ton.</i>	199. Tribe 6.			
1501 <i>Prunus Ton.</i>	<i>Hg.64 herb. 171.</i>		<i>herb. 32.</i>	
1502 <i>Ceraaus J.</i>	1535 <i>Dryas Z.</i>		<i>AlchemUlaj.</i>	
	1530 <i>Geum L.</i>	*	1467 <i>Cephalbus Lab.</i>	
	1537 <i>Siev^rsia W.</i>	0	82 <i>Acs^naL.</i>	
	1536 <i>Coltria R. Br.</i>	0	3280 <i>Margyrocarpus R.</i> 9^	
	1514 <i>Waldsteinia W.</i>	0	338 <i>Sanguisgrba L.</i>	
	1525 <i>Coraaropsis Rich.</i>	0	2655 <i>Potiriura L.</i> J	
	1523 <i>Jttbus L.</i>	*58	2812 <i>Cliftortia L.</i>	
196. Tribe 3. SPIRJEANTELE. ^ .37. herb. 16. £18ft.; £1uft	1524 <i>Dalibarda L.</i>	0		
3366 <i>Porshia Dec</i>				

800. Tribe 7. ROSE*	901. Tribe 8. POMADE*.	1513 Cotone&ster Med.
** 145. herb. 0. f 81 ft.	iig. 137. * ** 0. jt 1». It	1513 AmelAnchier Med.
Aosa	1506 ftategus L. *J8	1505 JWspilus L.
	1509 Photinia Lindl.	1507 ixynis L.
	1510 Eriobotrya Litatt.	1508 Cydmania Tbw.

202. ORDER LXIX. CALYCA NTHEJE.

Genera 2, Species 8; Rot-house Species 0; Green-house Species 0; Hardy Ligneous Species 8; Hardy Herbaceous Species 0. £21 feet; £ 0feet; 0feet.

* XMC wood, leaves, ana nower* or uaiycanthus noricus are sweet-scenica, nence it is twicu AUIWIUSH
The order differs from Kosacs in the form of the embryo, and from Gran&tee in the imbricate
Cutti calyx, from both in the absence of Peta H and in the numerous diviuons of the calyx.

1532 Calycanthiu L. 5 0] 1534 Chimonanthus Lindl. 3 0

203. ORDER LXX. GRANA^TEJE.

Genus 1, Species 5; Hot-house Species 0; Green-house Species I; Hardy Ligneous Species 4;

Jonier consists of only one genus, the well-known pomegranate, which is well worth a place in every
Pell. account of its beautiful scarlet flowers. It differs from Myrtaceas in the leaves being destitute of
m »? as well as in the seeds being enveloped in pulp from Calycantheae in the calyx being w Ivate
estivation; and from Memecylee in the stamens being indefinite. Cuttings, layers, or seeds.

1197 Punica W. • 4 0

004. ORDER LXXI. MEMECYLEJEL

Genera 3, Species 4; Hot-house Species 4; Green-house Species 0; Hardy Ligneous Species 0;
Hardy Herbaceous Species 0. 1 foot j £ 0 feet; ^ 0 feet

? pical shrubs or trees, with opposite entire smooth exstipulate leaves destitute of dots, axillary or terminal
violet or white flowers, and edible fruit. It is a very doubtful order. Cuttings.

195 Afemecylon L. | 3346 Uourlia J. | 1364 Petal&ma Swx.

205. ORDER LXXXII. COMBRETA^CEJE.

General, Species 39: Hot-house Species 37*, Green-house Species 2; Hardy Ligneous Species 0;
Hardy Herbaceous Species 0. f 0 feet; £ 0 feet; ^ 0 feet

Brefum and Quisquallis are among the most splendid of the climbing plants of the tropics, adorning the
is n ch they depend with garlands of white and crimson, and yellow. The bark of Buclda Bbercas
used with success in Guiana for tanning leather. The juice of Terminalia lirnix is employed by the Chinese
A varnish: it is, however, caustic, and its exhalation dangerous; benzoin is the produce of Termin&lin
excdn. The kernel of several species is eaten as a nut, and the expressed oil has the remarkable quality of
becoming rancid. Cuttings.

06. Tribe 1. TEHMINALIEB. 713 Conocarpus Joe.

1367 Buclda L. | 207. Tribe 2. COMBHE^TEJE.
2864 Terminalia L. | 1203 Combretum L.
1354 Getnia Rox. | 1355 Quisquallis L.

208. ORDER LXXXIII. VOCHYSIE.

Genus 1, Species 1; Hot-house Species 1; Green-house Species 0; Hardy Ligneous Species 0;
Hardy Herbaceous Species 0. f 0 feet; £ 0 feet; ^ 0 feet

tne trees and shrubs, natives of tropical America, with terminal panicled or thyrsoid racemes of yellow
and opposite or whorled entire feather-nerved leaves, each furnished with two stipulas at the base
Agree with Combretocete in having convolute cotyledons and inverted seeds, and with Onagria*
early with the genus Loptzw, in the flowers being monandrous, or, if triandrous, two of the anthers
ways sterile. Cuttings.

21 Vochysia Poir.

209. ORDER LXXXIV. RHIZOPHO'REJE.

Genera 2, Species 2; Hot-house Species 2; Green-house Species 0; Hardy Ligneous Species 0;

Hardy Herbaceous Species 0. f 0 feet; £ 0 feet; ^ 0 feet

nn-oves are plants of arborescent stature, which are remarkable, in tropical countries, for growing
snores of the sea and rivers, even as far as low water. The seeds have the singular property of
ng, while enclosed within the capsule and adhering to their parent, and pushing forth a long fusiform
zim lengthens till it reaches the soil, where it takes root, and forms a new individual. The bark of
pnora gymnorhiza, which is very astringent, is used in India for dyeing black. Cuttings.

1488 Rhizophaora L. | 1430 Carallia Rox.

210. ORDER LXXXV. LOPHI'REJE.

Genus 1, Species 1; Hot-house Species 1; Green-house Species 0; Hardy Ligneous Species 0;

Hardy Herbaceous Species 0. I 0 feet j £ 0 feet; ^ 0 feet

Cor ymh7 *oubtful order, containing only one genus. A beautiful shrub, native of Sierra Leone, with terminal
or white flowers, and a one-leaved permanent calyx. Leaves long, entire, and coriaceous. Seeds.

1504 Lophra Banks

Gene

a single style. From this form there are some anomalous variations, such as *Circsea* and *Lopezia*, which are, however, easily reconciled to the usual structure of the order. Most of the genera are preeminently beautiful; as *Epilbium*, *Himothera*, and *Fuchsia*, which are old favorites among gardeners. The properties of *Oidgrkrice* are little known, and probably very weak. The leaves of *Jussiaeia peruviana* are used as an emollient poultice, the seed of *Trapa nitens* as an eatable nut, and the root of (*Enothra biehnis* as a sort of salad. Cuttings and seeds.

212. Tribe 1. MONTINIEAE.	1184 <i>Gaura L.</i>	0	4	21& Tribe 5. CIRCSIA
2762 <i>Montinia L.</i>	1183 <i>Himothera L.</i>	0	58	<i>lig.O. herb.T</i> £4 Jft
? 1462 <i>Visnea L.</i>	1185 <i>Clarkta Ph.</i>	0	1	18 <i>Lopezia Cav.</i> 0
? 1187 <i>Gfela Lour.</i>				85
213. Tribe 2. FUCHSIEAE.	215. Tribe 4. JUSBIEAE.			
1188 <i>Fuchsia L.</i>	<i>lig.Q. herb. 5.</i> £3 Jft.			
2H. Tribe a ONAVJB	1353 <i>Jussiaea L.</i>			217. Tribe 6. HYDROCAVITBS.
<i>lig. 0. herb. 86. £ 22 ft.; £ 1</i>	404 <i>Ludwigia L.</i>	0		<i>lig.O. herb. I.</i> ^ Jft.
1186 <i>Epilbium L.</i>	340 <i>Isonarda L.</i>	«Q		403 <i>Trapa L.</i> 0 1

218. ORDER LXXVII. HALORA'GEAE.

Genera 7, Species 12; Hot-house Species 0; Green-house Species 3; Hardy Ligneous Species 0; Hardy Herbaceous Species 9. \$ Ofect; £ 3 feet; *4*feet

Obscure weeds, chiefly distinguished from *Onagraceae* by their naked and solitary ovula. They are natives of moist places or ponds, in various parts of Europe and North America. Some of the species of *Halorhizis* are tropical. They are not known to possess any medicinal properties. Division and setos, sometimes by cutting*

219. Tribe 1. CERCODIA'EJS.	299 <i>Proserpinaca L.</i>	0	2	221. Tribe 3. HIPFURI'DK*
<i>lig. 0. herb. 5. £ 3 ft. j £ 3 ft</i>	2652 <i>Athyriophyllum</i>	«0	2	<i>lig. 0. herb. 1.</i> *3 ft.
2622 <i>Serpula L.</i>	220. Tribe 2. CALLITRICH'NEJE.			
420 <i>Gonocarpus Thun.</i> 0 1	<i>lig.O. herb. 3.</i> J	1	24 <i>Jippuris L.</i> *0	
1224 <i>Halorhizis Forst.</i>	28 <i>Callitricha W.</i>	«0		

222. ORDER LXXVII I. CERATOPHY'LLEJE.

Genus 1, Species 2; Hot-house Species 0; Green-house Species 0; Hardy Ligneous Species 0; Hardy Herbaceous Species 2. £ 0 feet; £ 0 feet; *3 feet.

Aquatic plants, with the habit of *Athyriophyllum*. The order consists only of *Ceratophyllum*, which contain* two species, both natives of lakes and ditches in Britain. Seeds.

2651 *Ceratophyllum X.* *0 2

223. ORDER LXXIX. LYTHRARIEAE or SALICARIEAE.

Genera 11, Species 61; Hot-house Species 23; Green-house Species 13; Hardy Ligneous Species 0; Hardy Herbaceous Species 25. ± 0 feet; £ 10 J ft.; ^ 1 foot

Most of these are very showy plants, in particular the genera *Lythrum* and *Lagerstroemia*, which are *Jf representatives of the order. They are chiefly natives of temperate climates, on mountains and among bushes. *Pelargonium* is a common bog plant in England. *Heimia* is remarkable for its yellow flowers. Little is known » the properties of *Salicariae*; they are mostly astringent; the common *Salicornia* is used in inveterate diarrhoea**; a species of *Lythrum* is used in Mexico as a vulnerary and astringent; and *Lawsonia*, which is used by the Turkish women to stain their nails, is also supposed to possess similar properties. There is a plant of this order, called *Hanchinol* in Mexico, which is said to possess much more remarkable powers than any of the preceding; its expressed juice, taken in doses of four ounces, excites violent perspiration and secretion of urine, and » said to cure venereal disorders in an incredibly short space of time. Cutting, division, and seeds.

224. Tribe 1. SALICARIEAE.	1454 <i>Cuphea P. Br.</i>			1180 <i>Lawsonea I.</i> *
<i>lig. 0. herb. 25. £ 10* ft. ± 1 ft</i>	1365 <i>Acisanthera J.</i>			1150 <i>Grislea L.</i>
1100 <i>Pdplis L.</i> *0 2	1453 <i>Heimin L.</i>			
397 <i>Ammannia L.</i> 0 12	1452 <i>Nesaea Com.</i>			225. Tribe 2. J
1451 <i>lythrum X.</i> «0 9	3348 <i>Dlcadon Gm</i>			1587 <i>Lagerstroemia I.</i> *

226. ORDER LXXX. TAMARISCPNEAE.

Genus 1, Species 6; Hot-house Species 3; Green-house Species 0; Hardy Ligneous Species 3; Hardy Herbaceous Species 0. ± 17 J ft.; £ 0 feet; *0 feet

Shrubs, rarely herbs, with twiggish branches, small entire scale-like leaves which are usually g'a and small white or rose-coloured flowers which are disposed in spikes or racemose spikes. This order differs from *Onagraceae* in the ovary being free, and in the imbricate estivation of the calyx; and from *Lythrarieas* » of petals being inserted at the bottom of the calyx, as well as in its aestivation. The species are > at tlv ^ 0 Europe, Asia, and Africa. Some of the kinds of *Tamarix* have an astringent tonic bark, and yield, - burnt, a large proportion of sulphate of soda. Cuttings.

899 *T&maiixL.* *3

227. ORDER LXXXI. MELASTOMA*CE.

Genera 12, Species 76; Hot-house Species 69; Green-house Species 3; Hardy Ligneous Species 0; Hardy Herbaceous Species 4. JO feet; £ 2 feet; *0 feet

All these are remarkable as handsome tropical shrubs or trees, with large purple or white ^{owen} fur* flowers with several cost*, or nerves as they are incorrectly termed. The genera admitted in the body of ¹¹J pty are those received by the greater part of previous writers; they have been much increased, and ¹¹aply not with great propriety, by Mr. D. Don. The species are generally ill treated in collections, where ¹¹fj ¹¹gadin unfrequently to be found under the form of sickly stunted plants, instead of noble broad-leaved ¹¹shrub¹¹s, with masses of brilliant flowers. To be grown well they require much heat, much moisture ¹¹pe TV summer, and much pit-room and head-room. The fruit of true *Melastomas* is a fleshy insipid ¹¹J ¹¹e who which is for the most part eatable, and is often so deep a black as to dye the teeth and mouth pr ¹¹the eat it. They are nearly related to *Myrtaceae*, from which they differ in the want of essential oil £ urea dot-like reservoirs of the leaves which contain it. The juice of the leaves of *Micromia succosa* ana ¹¹wrea as a lotion for recent wound* by the inhabitants of Guiana. Cuttings and seeds.

<p>*•• Tribeg. RHEXIB\</p>	<p>SSO. Tribe 3. OSBE'CKJA.</p> <p>1181 <i>Olfbleckw L.</i> 1356 <i>Melastoma X.</i> 1360 <i>Pler&ma D. Don</i> 1362 <i>Acibitis D. Don</i></p>	<p>J?'.. Tribe*. MICOMB\E.</p> <p>1359 <i>Clidfemia D. Don</i> 1361 <i>Tocda Aub.</i> 1362 <i>Acibitis h. D. Don</i> 3345 <i>Chastogastradeft,</i> 1357 <i>Miconia R. & P.</i> 1425 <i>Blakea h.</i></p>
----------------------------	--	--

232. ORDER LXXXII. ALANGIETJE.

Genus l. Species 9. i Hot-house Species 2; Green-house Species 0; Hardy Ligneous Species 0:
~~&& H. W.~~ —, £ CIM 0# £ 32* ft£ £ 0 fcetj £ 0 feet

ng more numerous and in the anthers being adnate, as well as in the fruit being one-celled. Cuttings.

23a ORDER LXXXIII. PHILADE'LPHE^.

*Genera 2, Species 18; Hot-house Species 0; Green-house Species 0; Hardy IJgneout Species 18;
Herbaceous Species 0.* \pm 7 feet j \leq 0 feet; \geq 0 feet

K. t. f. 6118 which was formerly referred to as *Myrtilloides*, but which has lately been shown to be a hardy ornamental shrub, native of North America.

1479 *Philadelphia L.* 15 0 J1433 *Decumaria L.* 3 0

234. ORDER LXXXIV. MYRTACEAE.

G*ü*ra 37, Sjucia 293; Hot-house Species 85; Green-house Species 208; Thor& Ligneous Species 0;
2far^{ty} Herbaceous Species 0. i 0 feet j £ 0 feet k * 0 feet

Arytus & *SL* le *SX* es with marginal ribs, and an inferior ovary and single style, are the great features of
fructifying ~~plants~~ ^{are} evergreen shrubs or trees, generally bearing white flowers, and in the first section
of the Jy ^l y de the common myrtle. The section with capsular fruit comprehends, with the exception
of *Eucalyptus*, almost wholly handsome hard-wooded New Holland or South Sea shrubs, with
yellow flowers and stamens: yellow flowers are very uncommon. The volatile oil contained in the
leaves of the bark ^{the} Jeaves, and the floral envelopes, gives these plants the fragrance which has
been celebrated by poets of all ages. It is very aromatic, a little acrid, and slightly tonic and
astringent. In the clove this oil is so abundant as to constitute nearly a fifth of the whole weight of
the plant. Produce it. There is also a considerable proportion of astringent principle in these plants - in
Arytus Ugni and *multiflora* of Chile, and *Jambosa* malaccensis, it is so abundant as to render a
large proportion of those plants of great use in cases of dysentery. *Eucalyptus resinifera* produces an astringent
substance resembling gum kina. The leaves of the Chilean myrtles, *Leptospermum scoparium*,
some other species, have been used as substitutes for tea. Cuttings and seeds.

Tribes. LEPTOSPB'RMES.
 Subtribe I. *Melaleuceae*.
 2188 *Syzygium* R. Br.
 2187 *Jambus* L.
 2181 *Vitex amnus* L.
 Melaleaca L.
 Subtribe Euleptosperme*.
 Eucalyptus Herit.
 Hud.
 1483 Callistemon B. Br.
 1482 Metrosidtros Gae.
 3360 Angophora Cav.
 3361 Acmena Dec.
 1480 Leptospermum Forst.
 1481 Fabricia Gae.
 1172 Biflckm L.
 239. Tribe & [REDACTED]
 1485 Neiitis Gae.
 1484 Psidium L.
 1489 A/yrta L.
 3364 Myrica Dec
 3365 Syzygium Gae.
 1492 Olinchia Lindl.
 1491 Pimenta Lindl.
 1490 Calyptranthes Sun.
 1488 Caryophyllus Z.
 1487 Eugenia L.
 3363 Jambbsa Rum.
 3362 Josslnia Com.
 240. Tribe 4. [REDACTED]
 2039 Barringtonia Forst
 2040 Stravadium J.
 2041 Gustavia L.
 41. Tribe 5. LECYIUI'DE
 iS £6cytnis L.
 J586 Bertholetia H. &
 2042 Careya Rox.
 242. DOUBTFUL.
 1427 AgathophyUum^.

243. ORDER LXXXV. CUCURBITA[^]CEJE.

*G*nera* 18, *Species* 123; *Hot-house Species* 27; *Green-house Species* 54; *Hardy ligneous Specks* 0
Hardy Herbaceous Species 42. £ 0 feet! * 41* ft.; £ 0 feet

station of the gourd, the melon, and the cucumber, succulent climbing vegetables, the fruit of many of our comforts and necessities. The importance of the gourd in hot countries, where, from the nature of the climate, few of those culinary vegetables that are so

^{atabo} Pur⁸⁴t Jye qualities, which are, indeed, to be found, in a slight degree, even in the mildest of
^{ed by} ^{gourd}. Decandolle observes, that as the violent action of the Colocynth resin is much
the ^W mixture with it of gum, it is probable that the difference in the fruits of the order depends
^{Blo} ^{Fent} Proportions between these two substances. The seeds of the gourd, like those of the
^{n.fw} ^{Wer} POSSIBLY Af the ^{proportioning after} they may ^{and} ^{and} ^{really} find

2701	flatatrium L.	2696	Ceratosanthes Brnu
2697	Momordica&t;	2832	Johllsa BoJ.
2703	Neurospirma Rafi.	2698	Cuc&rbto L
	Vim Sechium Br.	2589	jfngoxia Ton,
2591	JV/el&htria L.	720	Gronfvia L.
2694	Trichosanthes L.	??2801	CarienL.

NATURAL ARRANGEMENT-

SH. OapRS LXXXV

<, Sptcie M; nothmtr SpeeinlS; Grrē^haute Sprirt 5 ; Unrdt Ltgnxoiu Spftif.

The beauty Of PMofloru it well known: they *re remarkable for the tiffluUr uranmnem of the »unt*jj and uUiilliitn, upon a column mrrounded by *e*eral lino uf circuntirallatian, funned by ai nuns ro*i* » barren thrtuililc colored Manteo*, which are popularly called the ray*. The fruit of aevnl • « • uufion-flomr U ailed with pta**ant acidulated pulp, on which account they are eatea a* daun i not known that they poMM* any medical properbtt. The tUUon of the order it not *etU*d; I vexy new CucurbiicQ*². Cutting* >>d i

9(1 Tribe L PAKorsIK'm

Tribe!. Fitunbui

198*

1925 Taabua J.

192J

>>7. OROBS LXXXVII. LOA-SEE.

\$ Ltgtwmt

Hard* RethateQUL Spetic* 5. X 0 feet P2 5| & 0 feet

NatMni U known of the quAhii* of this order. It oncf*ta of .mxuknl cuUmmid pUnu, tCMndly <**|jj with uperliw or rigid Hinging hain, atid yellow or white Haven. They are all native* of Arif hanlrfiff* *'''***. A rjer few of them are climben. Lulling and *e*dt, UT7 Hutbnt* Sim* 1 2193 toijj AtUn. 1) 3 I 1*78 llentittia L. Slft Blumenbkctua Selir. 0 I I 2195 &7pU°tnui Swt. 0 1 |

SU OtDuLXXVIII TUBSEB1A1CE.E

Spetic* IS; mt-kinur Sprcirt 10; Ow>Jk(m)f5^riTMIui Hardy Lfpteoit Specie*0-, llsry Utbrteeemt Syrcift 2. JO feet; £ 3 feet; f> d fort.

This order couliU on); of two genera, Turticmr and Ptri*mHa; thej arc imall tuffrutkoae or I planu, chiefly iatiivw of trofjica Annrici, with »]tmie te euUulate Iravei, and isUlary ydhnr Utij thbi* of fH*ththbum. Thla OPJCT U* kbledd dir tMudr>r>m Lo4*Mt in the I U M equal in number with the petal*, and iiu**tod at the bottoo) of the Calyx, not in the ttroat Of the I that order. Cutting* or toed*.

900 Turnena I* 6 2

249. OatHtxLXXXIX. POHTULA"CE.E.

Otmra 10, Specie* 35; JtotJumte Sptiet Et; OrcctJuiute Specie* I*; liarty Ltg* tlanfy Htriaeewu Specie* Si. f o feet; £ 5 feet; fe H feet,

In the exception of Tallnum and Calandrinu, and a few tpeccinof CUyt; run, the wl coniat* of itiugnffluot weedy plant*, of no beauty, and little u». Oaytnfr perfotliu and cumma which arc ocratioria)ly u*ed as labwu, being the only (peciet of » utenil kind. They are chiefly planu, frequeeting dry bunt lituatiofu, or the *h*ire of all part* of the world; all are modorou*, and destitute, ** far a* 1* known, ofsuiedlcina propriea. Cutting* *nd MMB

137* Triinthenu L I 3357 Calandrinu H. & a I W Cloytu*ci W.
IW7 AvtuDca 2.. 0 * 1 1*50 Phacotjirroa How. Midilia L.
1<H ,nacimp>eToaf*r*. I 90} portuafecria Jac, I "Jit* /. uieuui/*
IMS Tallnum J&m.

SHI OKDKKXC. PARONYC'HIK'E

Gncral6,Sf*tic*5H; Ittl-ktmuc Sptru* 5 ; GtYoi4o*u- SprtM 15; Wodr IJgmeotu S/xt.
HJ Hrbaceous Specie*557. II 1 foot[(fM t; ft 11fo6l,

Thit order eomktt chirBj of null InsignAcant wwdy, hnt>n>au, or ui pbnU, with exception off Crttoota, TWalalU/Ufobrum, Mid a teve perkifTf Hrk. life flo< whktr or gnenub-whtte, KMvnmoc* axillary and totmetnMa diajdaul ta tennitul c-ytrm, the l and entire. The order diffen Iron PortuUce* in the rtMacw being wpporite the lobe* of alternate with them. Cutting*, teeda.

; Tribe 1. Taxeraii*x
iig.Q. lurb.1 fillSO* TUephrtM L u
80S Corriglob L. .

MI Tribei IU.ICB<<ML f S i t l

813 Hemiuira L
730 Anfchu W*, 0
796 7IW<brura J.
7W /arottfehia Dw.
UrfaroaHiL

253. Tribe a

T. 6. £3|tt
1U Ortegid L O
29t Polycirpn L. O
»9 Polich*i* W,
30 Hnrknin ATVE,
7W 0 2

25*. Tribe*.

1375

SSS. TriboS. QIMJ*

% 0. Afrfi. 1. ft 1 ft.
S97 CuirtfiLot

237. O B M » XCL

into or nn*n thrubiIOi ttai rum, and bMutirui rad, orwjfp, yellow, or white Bowen. put* of the world. The (rder i* chrtSf dutinfublMid stow Fkoide* ID the embryo being *tr><M> eurified. Cutting, <Md, (tfnetlnw* by dhrttbo.

Ssa Tribe 1. CKutu*!**; or
I UtOfitMX./jr. a
417 TUWt L. 0
418 Bull lird< Dec <
11<; Septa* i.SU Crt*.
tIS TiM>6Ma Hair.

019 Vauittthe. Umm.

3SS PU<nni<flnr.
19U X I d

U<> BryophV Uum & >L

MIS BboAou'L.
I<B Settpetmum L.
I4tB Mon>U>c* tf"

AJtMMiU.

HU

90a OuwmXCIL FICOI'DEX.

n. lirirtr ItT. HetJmmmt Sprit* Si O n M J t w ^ d M t s i

Tbeat >rt an plant* with a creeter or Urndhaw* of Mttine, Oof it art n B t o m <f> t—n >U&U, rf whiteb u*toutJbj In the

le Bower* arc of all colon, many of tb< mot vivid hues, and rrururkaUlt- for exuandlnft
r" *!!!!*'. T¹.itiiniMiiBnii. liitwil IT_____*."'-rir mdn Tetratⁿnu cxDitua.
PortiaCirtnia, and U^S'Sy^Smn^S^m ae^tm tib^M^ tM ^mm^tm^A
y? w aaline matter 1* contained in all of them ; in tuaumnrw vermlcuUta a iub>.br/>**«becn (bund by chemical aoalriu to cmuiat of muriate of torfa and nitrate of mta*h. Xfce whole**
TM¹⁰» tverj dry or *alifie place*, in the temperate region* of Uif^w u Fiu rillfi, »i the whole are
***ther^{pe}of & wodHope! TT, el<>e.oft Kdrrcr<il*peri<of Me*aabrraath<rnnum oflbr the T**
**** utanucan of figure known to the vegetable world. Cutting*, oblett, division, or aeed***

L. 0 3	1519 <i>Tmaii'.nla L.</i> 1517 <i>Setirmm L.</i> 1518 <i>A.tb, m L.</i> <i>Mmuu/</i>	“& Snrtux. Xfc. frcro A J3Jn. I SIT Rf Muradrw L 1+46 Niliriru /.	3
--------	---	--	---

961 OuwtXCM! rACTEXor OFLTNTIA'CELE.

I It:i, PerfekM Haw.

i IpiuhVliuD //•*•
1471 Opuitua U w . fliUi Tritefl.
HIS

ML OKDCB Starr, OB.OSSCLAUIE-IL
W; Htrf /hiwif Sptrif^n; Grrfw ^ 1H *piv I ; Hardy Ligacout Specie* 18;
Hflrfy f/rrtDmni '^A-CIVJ O. j 3» feet; t 0 feet j ^ 0 feet.

from the lart by the definite number of their tUmeu* and woody tnfly item*. The utility and
of the gooMbomr and currant are known to every one. None of the other specif* equal ihwe.
If xrenj po*a^{ce}aea conridwablif eKcritena. The berry of too*t of these U «w»i, water},
that of AkAet nlgrum, and a few more, I* tonic and tbnnUat, which appear* to have NOW
the presence of gland* upon the leave* of tho*e specie*. Cutting* or H-eds.

Rita La *j8 0

S67. Own XCV. fiBCALLO'ytAl
tirtr*Jhna* Specie i; Hard; Lijpwvw* Speci 0;
• £ 0 fret; ^ 0 fecL

^{^t_{tb}} ^{HB} ^{ertTyreen} ^{hrtibt}, natives of Chile, with the h*H< of Andrvmoda, and which probably should be ^h with terminal raceme* of mull white or rait dower*. Nothing *a* known of their proprur*.
" & and *aed.

687 E^{cc}cal!..nijj Kut

T_{cra}

Siftienifti Iliit

t> cocutut? the glorj and deBgbt of the culm.-
— aaM perpetual verdure of thjoT lasTn, and the nqutaite tAmiffi* * and *!...*•
rather than to any ttr&tng attractow, of which they are <x L-
whtte oc pale pink,
iptonta, ahfcrt i ng n^o
upon dry walk. AD am watt*** of edU regtau, or of the smt taaMWaV?
a if them a* 1

* *b a weal in cancerous den
**c Arubbys

•0 Mitella L.
3m Tellina A Br.
1379 Cfarytotplenum L. 0 0 A .0 1
0 1 6 0 1

Quin

• : «t; A O Alt

which Mr. Brawn fint djatnmkhed thTM 1-faey are
and white or rnl fl. >crana and
bark of a tpnun of WViinUttu* is
be «l>o owed far adwHerting tb^ontqnn*. Notting t*

f • " - S ^ ISSYZSAI: **t** . Ifgl IU,,rj H K
«a Otnu XCV11L r*BE1.MTER.K

' ffluBpi of plant*, ant at the » T
a* teal than they an dain
venja, hot under the brtw! r Tentm, jnkd
food for »'lll TbffY an geocraUy rccogBbed by their hollow stem* and rot leave*,
that f* to nav. with a petiole, the buo of whtrh w

NATURAL ARRANGEMENT.

natives of damp ditches or way-sides, in cool parts of the world; In the tropics they are extremely rare. The simplicity of their structure, and uniformity of their appearance, have rendered their classification a matter of very great difficulty. It has been attempted in modern days by Sprengel, Lagasca, Koch, and Decandolle, all of whom have added much to our knowledge. The arrangement of Koch and IK² candolle is here adopted as the most perfect, upon the whole, of any yet published. The culinary and agricultural importance of many species is well known; the parsnip and carrot form a large part of the staple winter store of the inhabitants of Europe, as the Arracachas do of those of South America; and the Prangos of Thibet is supposed to be the most important and productive of any in the whole world, as a forage plant. The medicinal properties of Umbelliferae are not more powerful than they are at variance with each other. While the seeds of some are aromatic, and stimulating in the highest degree, the fresh roots and leaves of others are not less narcotic. This has been supposed to arise from the difference in the state of the sap in different parts of the plant; and it has been thought that the narcotic principle is only to be found in the ascending sap, while the aromatic stimulant properties are found in the juices, which are fully elaborated and matured. It has been already observed, that their dangerous properties are often removed by cultivation; the common celery is a familiar instance of this; but the most remarkable, that of Cenanthe pimpinelloides, a most dangerous species when wild* which is cultivated about Angers for the sake of its roots, which are there called Jouanettes, and about Saumur, where they are known by the name of Muchons. The roots of some Umbelliferae contain a large proportion of sugar; those of the carrot, when dried, more than an eighth; those of the parsnip just an eighth; and those of the chervil about eight parts in 100. Galbanum, Opopanax, and Assafretida, are all the produce of different species of Umbelliferae. Division or seeds, sometimes by cutting**

871 <i>SIBORDERI. ORTHOSPHER-</i> <i>MM</i> [prthos, straight, spcma, seed; form].	838 <i>Phellandrium L.</i> «0 1	283. Tribe 10. <i>THAPSIEV</i> Koch. <i>lig.O. herb.</i> 28. £16ft.
869 <i>iEthfisal.</i> *0 4	844 <i>Seseli L.</i> *0 27	284. <i>Thipsia L.</i> 0
844 <i>Seseli L.</i> *0 27	860 <i>Chlidium Cus.</i> 0 7	881 <i>/aserpitium L.</i> 0 *
860 <i>Chlidium Cus.</i> 0 7	3325 <i>Trochianthes Koch</i> 0 1	3320 <i>MelanoseUnumHqfw.</i> 0
833 <i>^thamanta L.</i> 0 18	875 <i>i,jig'isticum L.</i> *0 13	879 <i>Artedia L.</i> 0
875 <i>i,jig'isticum L.</i> *0 13	3317 <i>Brignolia licht</i> 0 1	284. Tribe 11. <i>DAUCINEJR</i> Koch-
863 <i>Hydrochyle L.</i> *0 4	3323 <i>MolopspL-rmum Koch</i> 0 1	<i>lig.O. herb.</i> 23. £12} ft "
867 <i>Trachymene Rud.</i>	3324 <i>Lcvisticum Koch</i> 0 1	3313 <i>Orfaya Hofm.</i> 0 "
864 <i>Bowlcs/a R. & &</i>	3328 <i>Silius Bes.</i> *0 3	825 <i>Dabucus L.</i> *0 1j
874 Subtribe 2. <i>Mutims.</i>	832 <i>Wallruthia Spr.</i> 0 8	3332 <i>Platyspermum Hqfm.</i> 0
866 <i>Drtst Deo,</i>	857 <i>3/i.um Tou.</i> «0 4	285. SUBORDER II. <i>CAMPYLO-</i>
865 <i>Spananthe Joe.</i>	834 <i>CrithmumL.</i> 0 1	<i>SPE/RMM</i> [kampylos, curved spcma, seed; form].
875. Tribe 2. <i>SANICITLE*</i> Koch. <i>lig.O. herb.</i> 35. £14} ft.	878. Tribe 5. <i>ANGEL</i> (CEF Koch.)	88a Tribe 12. <i>CAUCALINEJE</i> Jff**- <i>lig.O. herb.</i> 16. £9} ft.
846 <i>Actin.Mus Lab.</i>	<i>lig.O. herb.</i> 25. ± 13} ft.	826 <i>Caucalis /.</i> «0 f
823 <i>Sanlcula L.</i> *0 3	874 <i>Angelica L.</i> *0 7	3314 <i>Turgenia Hofm.</i> «0 J
886 <i>Astrantia f.</i> 0 6	3321 <i>Archangelica Hofm.</i> *0 2	827 <i>Torilis Gae.</i> «0 f
839 <i>DondiaSpr.</i> 0 1	3324 <i>Ostericum Hqfm.</i> 0 1	7840 <i>Trachyspermum Lk.</i> 0 *
822 <i>JFrngium L</i> *0 25	873 <i>Scllnum L.</i> 0 14	887. Tribe 13. <i>SCANDICINE*</i> Koch. <i>lig.O. herb.</i> 31. £15 ft.
ST.6. Tribes. <i>AMMI'NKS</i> Koch, dist.	279. Tribe 6. <i>PEU</i> [] <i>LIC</i> Koch, dist.	819 <i>Scndix L.</i> *0 ^
tig. 1. <i>herb.</i> 113. f 3ft; £ 24 ft.	<i>lig.O. herb.</i> 93. [] <i>LIC</i>	820 <i>ifnhrfscus Pcrs.</i> «0 ^
888 <i>ilamia Hqfm.</i> 0	3329 <i>Op6anax Koch</i> 0 1	821 <i>Ctetrophllum L.</i> «0 ^
851 <i>Cicuta L.</i> *0 []	880 <i>Firula L.</i> 0 18	830 <i>Ar^rbhU Mor.</i> *0
3319 <i>Ztz'u Koch</i> 0	3326 <i>Ferulago Koch</i> 0 2	888. Tribe 14. <i>SMY'RNEJE ATHI*</i> <i>lig.O. herb.</i> 89. £13ft
855 <i>iTpium L.</i> «0	882 <i>jPeucfalum L.</i> *0 24	717 <i>LagoeVia L.</i> 0
3330 <i>/ctrosclnum Hqfm.</i> *0	870 <i>Imperatoria L.</i> *0 3	828 <i>Oliverra Ven.</i> 0
847 <i>Trtma Hofm.</i> *0	871 <i>Callisace Fin.</i> 0 1	824 <i>Echindphora L.</i> *0
3316 <i>Helosciadium Koch</i> *0	842 <i>Jftibon L.</i> 0 6	2898 <i>Arctpus L.</i>
3318 <i>Drepanoprklura Hqfm.</i> []	858 <i>jfn-thum L.</i> 0 6	889 <i>Cachrys L.</i> 0
3315 <i>Ptychbitis Koch</i> 0	853 <i>Capnoprtf Hum Gae.</i> 0 1	891 <i>iippomarrathrum Lk.</i> 0
849 <i>Saqn L.</i> *0 5	883 <i>Pastinaca L.</i> «0 8	890 <i>Agas'Uis Spr.</i> 0
850 <i>Schultzfa Spr.</i> 0 1	884 <i>//cracleum L.</i> *0 88	287 <i>Hernias Thun.</i>
841 <i>A'mmi L.</i> 0 8	887 <i>Zosimja Hofm.</i> 0 1	852 <i>ConiumL.</i> *0 *
856 <i>iEgopbdium L.</i> *0 1	880. Tribe 7. <i>TOHOYLINE*</i> Koch. <i>lig.O. herb.U.</i> ^7*ft	3419 <i>Pentacrypta £<A.</i>
859 <i>Carum L.</i> *0 3	868 <i>Ulosp'nnum Lk.</i> 0 1	877 <i>Pleurosperrum Hqfm.</i> 0
831 <i>Aanium /.</i> «0 4	878 <i>Hasselquist'a L.</i> 0 8	854 <i>An^rnium L.</i> *0
836 <i>PimpinellaZ.</i> *0 11	885 <i>7*ord?lum L.</i> «0 5	876 <i>Physosperraum Cus.</i> *0
837 <i>Tragium Spr.</i> 0 7	3331 <i>Condylcarpus Hqfm.</i> *0 3	289. SUBORDER III. <i>COILOSP&%</i> <i>MAC</i> [koilos, channelled, sperm"] seed; form).
829 <i>Ledeburn Lk.</i> 0 1	881. Tribe 8. <i>SILERINEJE</i> .	290. Tribe 15. [] <i>NP</i> []
828 <i>Oliveria Ven.</i> 0 1	<i>S'g.O. herb. I.</i> £ 3 ft.	<i>lig.O. herb.</i> 1. £ S It
848 <i>Slum L.</i> *0 9	<i>S'fler Sco.</i> 0 1	818 <i>Bfforis Spr.</i> 0
3312 <i>Conopodium Koch</i> 0 3	282. Tribe 9. <i>CUMINEJE</i> Koch, diss.	817 <i>Cortandrum L.</i> 0
861 <i>iTupieurum L.</i> *0 33	<i>lig.O. herb.</i> 1. £ 1 ft.	
862 <i>Tenona Spr.</i> 1 0	843 <i>CumlnumL.</i> 0 1	
277. Tribe 4. <i>SESELI'NEC</i> Koch. <i>lig.O. herb.</i> 93. £24\$ ft.		
833 <i>ffihanthie L.</i> *0 15		

291. ORDER XCIX. ARAlia*CEF

Genera 7, Species 43; Hot-house Species 26; Green-house Species 5; Hardy Ligneous Species 7 i
Hardy Herbaceous Species 5. ± 15 feet; £ 6 feet; * 0 feet

Arallacea are a slight divergence from the well-known Umbellifer, with which they nearly agree *** except in being usually frutescent, and from which they are obviously distinguished by their 5-celled ir>. Their flowers have no beauty, but the foliage of many is extremely fine, especially of the species of & i^aj^bf lum; that of our common ivy must not be omitted. Their medicinal properties are much the same as Umbelliferae, except the fruit, which differs in virtues as it does in botanical structure. Their bark e^cudl^d aromatic gum resin, as in Aralia umbellifera. Their roots are tonic, with, in some cases, the flavor of paTM, aphrodisiac qualities; but it is probable that these have been greatly exaggerated. Cuttings, layers, awne by division.

£ fanax L.

8J6 CussiniB L.
912 Sciodaphliura B,

911 Aralia L.

1466 GastonW J.

3386 Adama WaL

718 iederia L.

NATURAL ARRANGEMENT.

CAPKIFOLIA 'CEJE

^o*t*tra* 11, *Spectkt* 118; *Uot.hoiuc Speck** 3 j *Grecm-honte Sjpopfm* 10 ; *Hardy Ligntmu Specia* 96;
*Hardy Htrtxcemu Specie** 9, \$ 7V feet; £ 7 WC ; * 0 M

- 15 S I (S» nii-rvilln Tou.
- 13 3 I tKl (.'-ipriltiliuin H AS •16
- 48 0 r>3! l, onl(. -era |;
- 3 0 I 623 Tridactum L.

6S6 y
100G Schrader Vaill
170* Unive'd tiro.

233. OMOBR CL UJBA'NTHE*

Grwra 2, S/trc^n If; Hot-hotuc Specie* Q; Grte*~hott»e Specia fi j B m b l f

th^r⁰¹¹⁶ of th^{TME} > re ***> rt*B; they are all /tenuine itirjULUM, looting beneath the bark of the tree
h...; Kfuu, and devching from their jijee, the whole of their palmit-ii. like Win
TAWIX, but the ijarAKtin are nKau th- most lovely. Of plant*, hunting in rufoton,
>c branches of trees in the tropic* uhirh thi lit-willi a IJKMIV mil
Oruid* ig - n», the common 'In-um never
ic oak. f/h
" t[. kin though now extint T thei irr. Li t
• itt ilu It ngocitoq, that ill veatigei nr tl
iy held he Kiranthus* having diaii[K, tritl

2750

S9i

Specirt 3; Hoi-housr Sj ; *llafdg Ligntout Specia Oi*
Hantp Herbaceous Specie 0. ± 0 feet. (.. t J » 0 ftc-t.*

Cul?^1 'ncoetwjtluou "hrut> with terminal epikc* oT sfreoi flower*, wholly destitute of ij>terett for garden*. ^uuuia^{gf jncfcgi}*

26 Chloriillhu-

SS6L OIMICIII. KUBIA'CEJEL

Centra 73, Specie 3x71 Haxjuxue Species MI; GrenJtoute Sxeiei 3t; Hardy Ugneotu Sptciet:
^ Manly Herbaaxnu Sptciet W. il(4ft.; £19 t HA

^fPposite eatin Un a with interi-pning itipulac, a monopetalou* upcri(ir corolla, with a drtinit,
and a bUocular ovaiitun, -in- 'in- peat rlmnicteristics of Ruhilc^c; nn urili- or pitirh exl
a very large proportion of the whole 04 -im imjrainoiH plants, iin-Kidinjr within
and lofty tree*, plant* with important noted drMt 'ili
which Die (irder baa bMB dlvda are nil m of Sjermart G4lc
it ho*, which are the representatives t if in i irtbern regioni. Amot^ tteae lh> Au
it Lhe mart ImporUnt on account of it* dye: GfcUtna ab^a potMMH Mar^c (uaMicsif JJUK,
the planti. . iiiiuinerah(c: of the rommer data
Kcnra ixor: iutur'nda V*xniUiU CetiiU
y'v¹ u² u³ u⁴ u⁵ u⁶ u⁷ u⁸ " " " tu* ha* a contribution tit ◇
" " " I or amitier. llw n>[>i Hnlyhii, urn!
La ill the Moturm, ami ut Moi lia in Indi;
t T¹ fi-brifudl proptsties of th< I :<tit on ; it u ISM gmenll} Known tltut tl
" " of Fiiicknt'Ya [IUU-II-, Mjcoinecneum ruryrbmtNm, WTUO iccinea, aiisi i nrwuiHora,
it weaker, [ifjvrr*. The tjark and root* of AMirhbx are ued, in the Ide of Bourbo,
and that of Morfnd - ued fir inV. Antrigent irrn(MTii)B of a v^r^
" -I-T arc found iii tin nil i of S; " " ' of Hunter, the Unchria tidmbr of tt<txliu(h, whit
confiuided with Gum kino, th nt plant
idi'd under the p i ec separatt'd by IhltiUMoI 1.Mwtt-mm.i, po*teu
I/Int, and
utv_(s) cu¹ Uuu w² w³ w⁴ w⁵ w⁶ w⁷ w⁸ ul the Citfia iirnishi-j the valuable
i> *u much esteemed in Europe u : UUT the DBN of coltec. Utvliion, >eed^s, w

M Schreib.
tin It. lit.
&6 Coccor^pvlunt A. Br.
L.

* Baa

" Amb

T7 Adlpp S₂f.

equidium *CuI*

301. Section >•
: *s HctlyMii IF,

301. Section >.

:*s HctlyMii IF,

inn *Ii.*

10 of 10

1996-1997 Annual Report

10 of 10

—
—
—

Digitized by srujanika@gmail.com

NATUKAL AUKANOEMI.'

I ,JJ.
- 'o j i M U a Sir:

J 1 ft.

ii riutit tiof-

6JL j
Sit Ntma Wi*
8S) PitK-ftrta A
05] CrAphila />
BS Chtodcc W.
Ml Pttfemu'/. A'

30* fertlonR
tig. t. Ar-ft S. J 3 ft
riif liii I^
•v.iiiiKth,
nibdu;..

355 *Bigt\&ia* j
358 *Bpernwdkv* /..

30t	Section n,	
//*	ft herb.	ISA

	AntbcMpinnnn i,	
	.Mill* /*	
y>/	Mus>r.ir-tt< U	"0
MB	BttMa -	4
	• j L.	M
118	OUtAOD f.	II
		*0

306. OIMICIV. AIUT.E,

, Marl* related to BubthenD. Their pro^erti
tlnl their illiiifi to Valortaneat la tummrntl by thr oil
of Uic Oi*stUo»UiU ii do time of tin; c um-ulalit. BM*

S29 P_{<r>}

WL QITYCV. VALEEIA^XKEJ£

**5, Spain ft) t Uot'lmiut Sperk-i (I; Orcn-Ai-tut- .Vjnjvv I; Hardy Lintout Spctki Q i
Hardy Herbueta >• ? » } (t 1 ft. 0 fte*.**

U il be riKni
DM Vtterriani, ami all thr I
• II und M I
of tuUrri 8cin41U, /*<!, Md uthrrt,
NFP OOMRHHI V u*ed m* riijfUJM. The odor of Valcrtan b not arnmltr u miUi, but v
n*ixrl*, the roouor *i* .. with wh.
1 thu IUy, wnjiliij tit- *I'nti-ri&na jatam&iut*, the iiiiikmard • . .*

3f) r^itrlinlliiw ZJrr.
!»5 fal

330 Patrlm.i J

80 Valtrian Wa Uufr. «0 ..

507. OBOKVI. DIPSA'CEJEL

**Oftiira 8, Sprclfj KM I Hot-kmitr Spedrt 0; OrrmJumu S/x-nVi !>, Airrfy Ligwotu 8ptci,
m<<1>>irrtVI!>, 1 V M | (£25|r<<>>: * tfaC

**Very nearly akin to Compfälur, of wbleh UMH ba.
flowen rrowlnB In hrada. So..**

HI KOrtlM L

I S*H *»,, M rVitt

*0

SciblbM IW/-

*Q 17 L

..L

308. OIDMCVII.

J 1 i UoUiauuif Specie\$ t; | <4rrsn . /tout? S)X-cu-i 0; /Mnfy /«^M
/;,,.// Uertocum Spdttt u. JU fwi. t

(MMFFirr WFittj weedy South American plnt, diffrcing from CinnplAttEt chiefly in the pxellion of U

DBOBECVUI **COMPtySITiE.**

• 009; *Hardy Igitti-ou* H*)

NATURAL AUK AN »• I.Mi

flf'l'T rooU of HeMnthetai tuberbfui. * vtwImM* food far narcotic, and hi* been «v*n nnployni « *—*n to mention the utility of i>n. It " JW Svrthmh and SalaaO*. u culiuu * IwanUr to .111 MII reader*, >i Bl t. t.^TMhI **P<*cl* of Ilir [']imfr «"«o of tinder; ihax- of thr arti lofCompftt*. ^{i riuiu,}
«n toor.hr In. *-tint NM MM M I hv i

FATU&AL ARftANGEMENT.

3405	Iainium L.	0	3407	Ertooom* Kih.	0	MM	.InilirMia I»
	Kiif,j.livltim I At.	0	H14	< Vanrai tin*	0		AVnithium A.
	Trir/iophflum Nut.	0		Dahlia Cav.	0		Framer/o Cav,
ma	Zinnia L.	0	:WM	Geora-Uo W.	8		
	Helhinm HeliL			Corepaia Jnr.		in	
	Aihrixla Kcr				0	3	
BMB	ijOiiKChlnpsiu W.	0			»0	89	
							SfSORMtR IX.
	Trldax /.	0					M'DF.JP.
1302	CutumIlia Jac.						{971*}
	Knllnta /.						
S3»i	Motfern Sehwb.						
8396	y						
SW-1	SraeaMckta I.						
83S	Vcrbcina /..						
S47S	Kuxenia Chant,						
	"(xUrthtu / {f- → lir.tr.-hi War.						
	g.v. IL ft P.	0					
SHU	Zjhi'iliixi Tcri.	0					
MM	'i.vilm Of.	0					
	,KU /•**.	0					
OiomWirt CM.							
MN	to IUura.	0					
	Jfpulifltnuii £.						
S4.W	W I t l Jot	0					
	i M.						
	Sut.						
Sil!)	s *						
	IMitnilhui L.						
24I	VlguMra Kih.						
9412	U.il(in!«t Ijim.						
I'U I	Idol:***-Via I.						
M00	Bohlnto*. HNM.						

ail, OBPEB CIX. 1.OBKLI,V-
6VIKT « 6,flriiM 90; Tlot-hnut Shrift H ; (irtrn-hmtt., , Jl*i*,l, l.^aeaut SpedM 0 j
Jurdg llerbiifnni 9ptci: U f 0 fi-vl; j£ il fet-t; i K ft-eL

ornamental herbaceous plant, with blue, white, or K>rM flowers. The anthers are of the
strurli're »• UIOM nf Cum(i6tita. The jute-t of tome ul'th obelied il highly c.n.
taken niirrn.illy, jirmluiiu yumKtDK, nii'l ivm death: IK-M-IIIK II '•, tin- rout of Ixibelm aiphltitia, nt no**"
dote*, acta ai a ttmi. Rreatoi '(ianlly, aa dlurlli' or purgative; and, if taken m
quantitttt, a a MII •nit-lic. An infiuiioi (if LobfcUa inflita La uied in North America a* a remedy fur li-unirrh**.
and the root of liitbt'd c^rdlnaJti U employed In the tame country M a vermifuge. Stedi, division.

8» LobjUo L. •0 17 1 3390 ClintoDM Don. 0 1 I 714
610 Monfippti Sal. 0 S I 614 LwhchidHto H. Br.

331. Oaina CX. STYTA'UEJL.

1, Specie* fi; Hoi.hotttr Hprrit (>; O1W JMM Sircin (; Hardy Jgneaunt Spfcifi 0;
lurrfy Hfrtaenw. S>xi.'j II. fo tat) j£ t feet; A, 0 f<t

Like Hii; l.T.t, the propertys of tiiti very triall bul cnrlnui onler «rc, it any, u d. All are inh».
JOT* b^f Sew 1 liiltaiui, ami nlhiT Ju'fir* ai : tuua ahru)*. Tl link Aower#, i I ^ h
tUterin* itlandi: their atameoa are united into n oohtmn. irhcti ii timmiatnl by a teulc (ti^ina, anil
fa imUblc in ao hlf b * degree, that, if touched with a pin, It luautitly tURU from lu place with ttre.it clu

2581 Styldiutn Swz.

33\$. OHnRK CXI. 0OODENO'VIJE.
i, Spdn 21; Hot.hotue Spedet 1; Grtn.kouar Sjretes IS ; Hnrdt Lt&tfoiu SpecifO;
Hardy Htrbactotu Specw 0. I 0 feet; f <> i-

Now Holland and South Sea herbt or underatirabu, vory np'erly akin to the latt, rrow which th.y '•'JJ
more in artificial charrrera than ID habit. All of them ant pretty, and dmrvvtng culture. Nothing u F™
of their propertied ('utting, ilu HnHI, or aeda.

CIS GoaStnia R. Br.
Bid KUthakt H. Br.

fil'i\yUHn Km.
680 Scar'vola It. Br.

G17 Damjiiera IL Br.

833. O»nR* CX11. CAMPANI I. \< KJE.

Genera II, Iftttom Op&tm 1; Grtn-hmt Species33 j //drffy Liewuv Specif
hrbaceoui Species 188. 1 () h^ . ,

Thoc differ Del lining Uu lioyrt-w lii lieadii, in tbtti u.u.illy diattaal a) I
poly>iHTiinjis / nut, .in<l.il"> tn exuding i milky htk*. All the fti-'icra arc- prott
mental. They are inewtly herlwccou*, and by far the gtentr numU'r <tre extra-trnjral, abouodin
tn the wDdk and cupptco (if the north. The rood of tjmpiuula Aapdiuuhiin an uaed aa a > .
the name of lUinpnu Citttinga, divUon, and H-div,

J09B Canarlnn ... VRIVI Sehr. 0 3 (iM /Mhyoitnta //.
1177 MyhiiDtM Merit. •0) lhi'titnii /..
715 Lightfo Ua ll.-nt fil« Prlamatnrtiua HerU I Krc /..
003 Aituni'ili-Ti ffa 0 16 i;f Kurlik L, .0

33t. O w n CXIII. OESNERIE¹ JE.

Genera S, Specie 38; Hot-Aoute Specie* 38 j Green-home Specie* 0 j 7<m;/ Upowl Spfdt* ° i
llnrtiy Herbaeau* Specie* ft 1 0 lhet; tf 0 feet j A <) ntt

Fine tni>iil ln>riM, with brond, fl*i**, tlowny Iraven, and juri*jili* or iorlet ttuwprt, Tlmy »H fCI
heat, and decayed vegetable aoll; In thir Dative i-imity, wliih >> cliufly rqluncttal Anin
foiuid (trwinjt In the womla, where the earth ta little- more than a bed of rotten leaves and bark. CW ««P

346. ORDER CXIX. STYRACI'NEJE.

Genera 2, Species 7; Hot-house Species 0; Green-house Species 0; Hardy Ligneous Species 7;
*Hardy Herbaceous Species 0. f 23 feet; jg 0 feet; * 0 feet*

Elegant trees with white flowers, chiefly natives of North America. The *Styrax officinalis* afford the officinal storax of Asiatic Turkey, which issues from incisions made in the bark. The *Halesia tereptens* » the well-known snowdrop tree, with its pendulous white flowers. Cuttings, layers, or seeds.

1353 <i>Styrax L.</i>	4 01 1432 <i>Halesia L.</i>	8 0
-----------------------	-----------------------------	-----

347. ORDER CXX. MYRSI'NEJE.

Genera 8, Species 52; Hot-house Species 31; Green-house Species SI; Hardy Ligneous Species 0;
Hardy Herbaceous Species 0. ± 0 feet; £ 0 feet; ^ 0 feet

Showy shrubs, with evergreen undivided leaves, and cymes of white or red flowers. The Ardisias are common in collections. None are natives of Europe, but are found in the hot parts of Asia, Africa, and America. Nothing is known of their properties. Cuttings and seeds.

581 <i>Ma^sa J.</i>	I 669 <i>Corynocarpus Forst.</i>	I 570 <i>MangIWa J.</i>
560 <i>Jucuinfo I*</i>	I 572 <i>Emblema Brm.</i>	I 527 <i>ClavTja Fl Per.</i>
571 <i>Ardisia Swz.</i>	I 2891 <i>A/yrssinc L.</i>	\

348. ORDER CXXI. SAPO^XTEJE.

Genera 11, Species 44; Hot-house Species 31; Green-house Species 6; Hardy Ligneous Species 7;
*Hardy Herbaceous Species 0. f 25 feet; £ 0 feet; * 0 feet*

These are also shrubs and trees, which are mostly evergreen, and natives of the warmer regions of the world. Some of the Burnt-las are found in the southern states of N. America, but none of the order exists in Europe. They are chiefly valuable for their fruit, which, in many cases, contributes richly to the dessert. *Mimusops Eungi*, *Imbricaria malabirica*, and *Argania Sideroxylon* are all of this description; the star apples of the W» Indies, the produce of several species of *Chrysophyllum*, and particularly of *C. Chumia*, are esteemed delicious; and the Medlars, Lucumas, and Sapotillas of equinoctial America, all the fruit of different kinds of yChras, are among the most valuable productions of the western world. The seeds of all the order are oily; those of yChras *Saputa* are accounted diuretic and aperient. Their oil is not fluid, but so concrete as to have the appearance and consistence of butter, whence the name of butter-tree has been applied to different species both in Africa and India. The most famous of this description is the Indian maya, mahva, or madhuca, the *B. issia butyracea* of botanists; the seeds of which are so oleaginous, that a single tree has been known to produce three quintals of oil; the dried flowers of the same tree are mixed by some Indians with their food and a kind of spirit distilled from them by others. The juice of all the sapotas is milky, but not acrid ana poisonous like that of most other lactescent orders, but, on the contrary, yielding a wholesome beverage or food. According to Brown, the bark of some of the Achras and Lucumas is so astringent and febrifugal » to be substituted for quinquina. Cuttings and seeds.

1351 <i>Inocertrpus Forst.</i>	■ ■ 559 <i>Arguinia R. & S.</i>	562 <i>Luchma J.</i>
555 <i>Ifcimëha Stoic</i>	■ ■ 556 <i>Chrysophyllum L.</i>	1156 <i>Mimusops L.</i>
569 <i>Sersalisia It. Br.</i>	■ ■ 557 <i>Nycterisition R. & P.</i>	1424 <i>Bassfo L.</i>
558 <i>Sideroxylon L.</i>	■ ■ 561 <i>-yChrasZ.</i>	

349. ORDER CXXII. EBENA*CEAL

Genera 4, Species 38; Hot-house Species 12; Green-house Species 22; Hardy Ligneous Species I;
*Hardy Herbaceous Species 0. ± 6 feet; £ 20 feet; * 0 feet*

Some of these are hardy trees or shrubs, with deciduous leaves and white flowers, natives of woods, mountains, and banks of streams in North America, Europe, and New Holland; others are tropical evergreens. Of the latter, many of the Diospyrus produce eatable fruit; as, for example, the Mabolo of the Phitppin^{inc} Islands, which is as big as a peach, and the Kaki of Japan, which resembles an apricot. All these fruits are remarkable for their extreme austerity before maturity, and the necessity of letting them decay, like our medlars, before they are fit for table. These are also distinguished for the excessive hardness of their wood, and for the black color it sometimes acquires when old, as the Ebony. The bark of *Diospyros virginiana* » used in North America in intermittent fevers. Cuttings and layers.

S791 <i>Maba Forst</i>	12889 <i>7>iospfros L.</i>	4 0
2890 <i>Cargillfe U. Br.</i>	11373 <i>Royena L.</i>	

350. ORDER CXXIII. BREXIE&Eacute;.

Genus 1, Species 3; Hot-house Species 3j Green-house Species 0; Hardy Ligneous Species 0,
*Hardy Herbaceous Species 0. ± 0 feet; £ 0 feet; * 0 feet*

A very obscure order, its place in the Natural System is extremely doubtful. Fine trees with the habit of Thcoplir&Bta, and with axillary bunches of white flowers. Cuttings.

523 *Bræxia Nor.*

351. ORDER CXXIV. OLEMN&

Genera 11, Species 124; Hot-house Species 11; Green-house Species 23; Hardy Ligneous Species 90;
*Hardy Herbaceous Species 0. f 1374 ft. £ 0 feet; * 0 feet*

The olives are known by their monopetalous corolla, with a valvular aestivation, two stamens alternate with the segments, a bilocular ovary with no discus at the base, and pendulous collateral ovula. They were formerly known with the jasmines. They have all similar omnissito lavrav. their flnwpra either *

to the order. The seed of the olive contains so large a proportion of fixed oil, that it has long been one of the most important objects of cultivation in the south of Europe. The bark and leaves of many olives are bitter and astringent; these properties are particularly apparent in the ash, which has often been successfully used as a febrifuge. From the exudation of many species of that genus, the mild purgative called *lilac* is formed; it is most commonly found upon the *Olea europaea*. M. Decandolle remarks, that in pi oof natural affinity of the plants here combined, and of the propensity of separating the jasmines from the olives. It has been found that all the olives, as now restricted, will bud or graft upon one another, but not on the *lilac*. Thus the lilac will graft on the ash, the *Chionanthus*, and the *Fontanesia* and even upon *flnilyrea* la and the olive will take upon the *lilac*, and even on the ash Cuttings, layers, and by grafts or buds.

•R&xinua L.	*S3	o	81 Linocifera Swz.	37 Phillfrcia L.	13
Ormu8 Pers.	5	0	80 FontanesTi Lab.	40 ZigQstrum L.	7
Chionanthus L.		20	39 Notelac'a Yen.	41 Syringa L.	10
Millingttonia Rox.			36 (Pica L.		

352. ORDER CXXV. JASMPNEAL

**Genera 2, Species 40: Hot-house Species 22; Green-house Species 12; Hardy Ligneous Species 6;
Hardy Herbaceous Species 0. ± 6 feet j * 0 feet; * 0 feet**

of *Vk* ^{abundance} to the predominant property of the *jasmine*, and has made it for ages the favourite of poets and fine people; this arises from the presence of an oil which can be extracted so as to retain its perfume. In *bölf*-^{qualities}, the *jasmines* do not differ materially from the last; they are neatly distinguished by *wmists* by the direction of their *ovula*, which are erect in *Jasmine**, and pendulous in *Olcinc* Cuttings.

42 Nyctanthes L. | 43 Jaunlnum L. 6 7

353. ORDER CXXVI. STRY'CHNEJE.

Genera 3, Species 10; Hot-house Species 9; Green-house Species 1; Hardy Ligneous Species 0; Hardy Herbaceous Species 0.* f 0 feet j £ 0 feet; ≡ 0 feet

is il2P^{ic} al trees » nearly allied to Apocyn'e, and like them truly poisonous. The Stry*chnos ndx v6mpica the ^{mark}~~blc~~ for its bitterness and acrid deleterious effects, which are indicated not only when introduced into tovo? ¹ ma~~■■■■■~~ b ut still more violently when introduced into the system by inoculation. Theophrastas are fine whnri-^E with simple stems, and undulated spiny-toothed leaves, which are disposed in something like wⁿ orls. Cuttings and seeds.

574 Strychnos L. | **575 Fagraea Thun.**

354. ORDER CXXVII. APOCY'NEA

Genera 36, Species 145; Hot-house Species 115; Green-house Species 14; Hardy Ligneous Species 9;
Hardy Herbaceous Species 7. f 7 feet; ff 8 feet; fff 10 feet.

Haray Herbarious Species. J. 7 feet; jg. 8 feet; 200 feet.
Now turn from the contemplation of plants endowed with mild and agreeable properties, and fragrant y...
I and often hearing food for man, to others which are among the most dangerous and fatal poisons;
whose juices though milky are not a wholesome and delicious beverage like those of Sapteal, but, on the contrary,
are caustic, or bitter. They are readily known by the twisted direction of the segments of the corolla, which
is compared to the rays of a Catherine's wheel, whence they were called by Linnaeus, *Contort**. By
Vin... greatest part, of the order consists of tropical trees and shrubs; a few Apocynums, Amsonias, and
are natives of the colder zones of the earth. Many are elegant climbers, as the different species of
and Melodinus. The splendid Oleander belongs to JVtrium; the different species of Plumitria, Cam-
Strophanthus, and Arduino are stove plants of the greatest beauty. The medicinal action of these
is highly powerful. In general, the Apocynes are acrid, stimulating, and astringent; these principles,
act so powerfully on the nerves as to produce stupefaction. The root of Ophiophyton is very
Th... and purgative; under the name of snake-root it is used in India as an antidote to the bites of serpents,
the bark of Cerbera Mdnchias is purgative; of ifchltcs antidysentrica, and the Wrightia of the same name,
the Pkt-ut and febrifugae. The leaves of the Finca are so astringent, that they have been used successfully in
fining; those of A'trium Oleander are said to abound in free gallic acid. The inspissated juice of a species
of w... known in Mexico under the name of Ycotli, is a fatal poison. Cuttings or divisions.

Alyxia R. Br.	535 Wrightia R. Br. 546 Strophanthus Dec. 529 Cérium L. 548 Taberçrmontana L. 547 Cameraria L. 549 Amsbini Ili Walt 544 Plumiera L. 534 Presidnta R. Br. 530 Alstünia R. Br. 531 Cryptolepis R. Br. 533 Thenardia Kth. 537 fchtes L.	536 Beaumontia Wai. 541 Hsmadietyon Lindl. 538 Valfaris R. Br. 540 Ichnoökarpus R. Br. 539 Pardhsiví R. Br. 553 Willughbëtæ Sea 545 Lyonsm. R. Br. 748 Melodinus Forst. 543 Carpodnus R. Br. 542 Wpucynum L. 3295 Anthoclcsta Afx. 392 MonetiA L.
JS? Cebera L.		
£1 Ochromsiaj;		
Moñena Lou.		
Allamanda L.		
F	•8	

355. ORDER CXXVIII. ASCLEPIA^{DE}iE.

Genera 44, Species 264; Hot-house Species 187; Green-house Species 32; Hardy Ligneous Species 2;
*Hardy Herbaceous Species 43. ± 3 feet; £ 18feet; * 0 feet*

789	Piar&nthus <i>R. Br.</i>		772	Stylandra <i>Nut.</i>
790	Hulrnia <i>R. Br.</i>		770	Gomphocarpus <i>R. Br.</i>
791	Brachyst'lma <i>R. Br.</i>		754	Harrnbtin Hook.
792	Caralltiva <i>R. Br.</i>		763	Oxystclma <i>R. Br.</i>
776	Huya <i>R. Br.</i>		762	Metastelma <i>R. Br.</i>
777	Tylophora <i>R. Br.</i>		768	Xysmalubium it. <i>Br.</i>
778	Ceropogia <i>L.</i>		766	Calotropis A A-
774	Pergularia <i>L.</i>		761	Cyrancnum <i>L.</i>
775	Marsiltenh <i>R. Br.</i>		759	Da'mta <i>R. Br.</i>
767	Dischidilia <i>R. Br.</i>		760	Uiplolepis <i>R. Br.</i>
764	Gymnfema <i>R. Br.</i>		757	Sarcostcmma /{. <i>Br.</i>
765	Sarcobolus <i>R. Br.</i>		758	Eustegia A. £.
773	Gonlobus <i>Mx.</i>	6	753	Microtoma <i>R. Br.</i>
Til	jisclepija <i>I.</i>	23	756	Astéphanus <i>R. Br.</i>
769	Anamherix <i>Nut.</i>	1		

NATURAL AHUANGKMK!

386.

OEKTIA'NEJT

O M M B, Sprin 1H; HoUuut Spreitt 9, j Grfi-n-fumt Sprift IS; Hardy Ugnetu Sf
Hrdy Hhmanu SpeSte* V&. t " `` . f M frf ; i * `` *.

An orditr In tome defiwe tntrnncd Mir M M M PoUmonice* and ferophiiMrlaw, from both whkl>
I boUi brhabhandfnt Ki ton** of h* `` `` * bordt dotarr upon ApocfiK*. Thai
mountaloM* nnoM or pooli In all wit* of th, wottd. ^Tb* Crntwn, arc m,
, tr.thdMfiblurlowm; Tho latter eefar, and dUfcfint ihaitoa of oranar, bring Uw

Itnpal..

* root of 0WjtUM
«td Mil

• 1,41c fogil/I III IIIia!
IM(*IM `` J `` a v<tnfuit.
r a< anthflmtntle, and /WU/M amara,>
AichM ttawiaMI ftwaUw» utu[^]ainndi (bmiui, ^

1.
7
AdatL
W

•0
1
•0
Autt,

371 Mi
543

t

D

« t

W

.J tvtrrum t, S*	14	0	9ML (tention 1 (>s'«n* • .aria 1HII ltruwnU.	CrMC^i.tui t- raft>'cium Airi i id
---------------------	----	---	---	--

» O H M ' V I . I SCROPHVLA'BINJE.

Uts.

1 (,

A fraal put of lirmanw'i DtdjaUsla Artgi«p/pnnia U found her*,
bail* among tfw nott obrKua cfcfMtarMm of the ontn. Th« tn
IMUM yii II MM| * *#utu - u ^ j^rikdA ar HdoaAiftl ...
world. Th* f>r*wai*(n»«U»»IH

Mtttari

RfK-uUrt*,

mia, ad trtd STUMPIBIMH rgggb*a* i t*t t*U* colour b MN
tb» t*fU f-rvrwhata: t» Wlaauha, lOto**.
bacon* BBollhtnf to mm AMlffchMMN. Tb* RbMaMbanjA

SB utar fruit and didpunMM* ilaw
•r« gMMral
- T p T «aj 'y u
.u «FmjBpn^» in KM |

fwjMth Tti* li»»* thw plant, roduNd to fomtm, «B*tt i umlli
Mffloof Uw poaa: In M m M M W tb n cam* danUi

taacud» iwiMhof taw*.
I, and U* orOi riary arrld
w4» (br thmr aatnofrni
Mb aad *MW*k-M. and C
taaktfhakv

notaat* ihaw lbvjan titsful in ala,

371. SartJon 1. 8rO«xa (4) AX.
nustnota.

M. II. .1.(1'i J.
JAt Seopi r 1 a./.

IRKt V;, r/in Itiitnl
17IV M.UK-pi.) 1.
11 SlivnUla f.

7W1 Twirarw W. tn.
i 1.
i .niljfl Jar.

INI - Cjrmrl*riji /
17Ti it IIfft.

t il. *0 18
1 Aria Tun.
17SO 8rrig>iliultrk* L.
17KI IMgtlali* /,
- leowterwo If.

1718 Chriont L

1HII Hmrftlll H. Br.
Maim /«.

>l<>nirmi.itu* W. ML

17K1 •nttio»ini.

lftt) l.unl. rrnrj L,

<<.. Kill

V P.

> / >

fi 1

.0 1

o

/ 1

.4

urtala L.

.0

17« Euebrbma Nit.

0 1

litr, t iutil*|i HutU.

0 8

1771 >>>iawu)uuui /., Z.
1741 Z. .0 5
->l.1rtii.i 1, 0
l'h, 0
lot 40 Nut 11

37* SortioAjl Sr*
THKalraai.

A/r6 ISJ • Id- tf Vjt • A

Cilnolkru
nthw Ft. per. 0
SellW(i)l'ktLJa,

M 0 *

M 0 0
44 LapUmri 0
i. IVTOHI 0
4(1 Ulplm>hJHum /<* 0 1

llemf mvrii i.

379. OwutCXLIL LAI

Mw«V Htfbastou Spetitt 52*. f VJJ ftr*, f { frtr i .

1 lortium of DiamIrU MooogfnIA, AFJII thf whulr nt [iir]J.
LAttte, which

-tonfmnj. Hi the IMTH, lumr limr
TIIIM are << rrmatkaair fat ltrir Ionic, wt
•ruuvMtc priotfile, ln ddfWrni p<

tlawM MrltafM: SoMmldia
tdniulativt lVov tha dlfevmt 4M

11 a* ttiotyshynm, |

1 ijMTiiiw, tliat i' Mm, and woda.

574 Tribt L HUTROfOML

it L. *0 fi

WI l<>riilim Mi, « |

.0 SI

IAD /VrU/a W. 0 1

JUS KUM.'Itlid W. 0 1

3J7* Aphanotnllu* Jbad. 0 1

337J Dlauih^lla Burnt 0]

ITH t owbndkM fM

973. Trio* it b m u l « u .
tig. 34. *rt-A.S9. f »i R ;)C J3IL

mm lf* 0 7

i la Bfntk. J ii

...it/, *SS 1

3TM LojpWmhui ft-iKA. 0 ii

IjM flWatwii /. 3 0

low Wvttiinfifit SnL

37 Tribe .1. AJV^IOR*.

lfr.7. *crA.41. Jafl; tf.Jft., AlSft.

um 7>ti .7

.0 1

.7 I'hyn., 1 8

ISM A hii .0 1

1(17 Aiilv.rofle* A Br.

77 Coliin*|nAi L. 0 "

977. Tribt 4 MnN*ilPia.

lig.S, Jkrbb.W llftft

n.fj. 0 W

. f< 9

n

378. Tribe fi. V i n vnut

1^. 30. hrrb, a4>, f 11 it j £44 ft.

l(F7fi Loonitit /.

1671 .

3377 Oriii||*rtn Kwk. 0 "

tint, balm* Atkj ri r11.IJn -V

• vim In iucfa rjn.r

U in lflit IK cuuMT an object of -MM

378. Tribe fi. V i n vnut

1^. 30. hrrb, a4>, f 11 it j £44 ft.

l(F7fi Loonitit /.

1671 .

3377 Oriii||*rtn Kwk. 0 "

MS]

ion Uttdula £

i |>ialiiii

U 1 0

a \V

n /Yri 0

MM 7'ljjrmrl* /.

PA.

MI

•u

in

NATURAL ARRANGEMENT.

an obtain mluu wted, and the Utter luu itu-	the BoriiL Tte ptominrnt botanical charaewr a	o— otled fruit, with a central ptacrnU, ami ur little coruiqueoCT; Ihry an	roriovu to thtt botanltt but ur
.he	t aM faabl* ami ur little coruiqueoCT;	Ihry an	
til* primriMo am	i (raffrant, and mikljljr udortrk and topoflc. CortOaa Malhlola Itu ttccn lucd I		
nervoui dlaontor	Dñ [ejcj], oSEata, and aeod*.		
455 Cyclamen /,,	•0	4M nt-tifflili /_	0 4 458 jfnay&illi /.
151 i>.i...iiu..n /	0	it/.	47' Din M.
0		IHUMIM I.	Sit i
4 H * OfttM I.	0 1	457 i.jwmichu f.	0 1 Janthu* Rot 1
401 Primula It.	•0 00	W3 Lutiriu* Com.	
450 Andt6ucm I.	0 IU		

317. OawaCXLIX. OLOBULVRI.T.

M, Ih^uifSptinn; Green-kotut Spretri; Ilmrty Jignfomt Sper*-4Q;
Hardy //roocMM 8p- I Obat] »<rf! i * « * «
rrrtt alpln* plant* with bluff (wnw, Ttia la a m of tibdiuVirta Atjuum are very bllrr and now
purple, gilly at the vanc time a more to the blanch and tele. Mrlkou and nting.

342 OlobullrU I n o

r
•It.,

am OtDutCL. i|i.i;MiiA(.rs'K£
OfMrTd 3, Sptieit ftj ; itv-ktmu Spetic-i I; (MC-nAtmc S'^rin 77 ; //nri/y lif*f,m, SaedrtOi
Iforrry tUrUceout Spetie, la f u : ,
•It.,

tin. It* m* at SUUc* /4mbnium l*«*«nnge*.
•nW ud MMK aod <a^4oywl a> vaacatWiam.
OSB mttof £. «0 W t W8 Anntru E. •0 18 | 423 Fhimbafu L.

SIKI.AU IV. M' I LA M Y'DEvK.

; Gresn-houte Spptiet T7 ; ttnto ligntxm* SpraftWi

The »b_cnce of corolla chanctrty* IhU tuhJivitum of dloMjiladomout vegatation j but, u the term
I* luMect to ftwnumt mitunaarftaodmt, II ihuuU !> burnr that whrnrvn thm it onlj
•tiTclnpr that rnenvelo li to bo ronahteted ulfx, »hrth^x gnti^u In mott u m , or colored, ai In the H*
of Peru.

SSit OaowCU PLANTAI
Ot**TO J, *-/ Holhotu, Specif 0 (firm km., 3/r, I! Mb Ufnrrw Spfde* 9j

k tneonapktUM* herb, found In waata placM all orrr tir *;nid
II the pÜbeMror* la Jotnlnt; the flu wen, «*!
M l n and umrnrii, their aeedt mwIugii
IN 14i||r iUanlti<+ (ruin tin; tuuth "I li.)in>, lm [hi- [iul]Mjt* ut formiUK an IIIILIMIPU IN which n
waihed. I*, medka 1* MMIM timrt rulUvatnl by Ijnnrr* under the namr Of nifstirM. MivuLom aud <-rit*.

Mi Omnt: KYCTAOI'

1., Hardy Lignmui %<

With ih» exctvitmt of MlriMIlt, In which the cwtond eatn bat a thaw* evAvt, all DM ord<
tf »*.->, ROlimi/ aAta iherlnewuiH acoa<to/tha>troptc*aj>d<
ffutnl in Kurijpr and nit. ii lr>|traict. lIn- ri>t &(Miibiln,
firtin 1D i-miincrH the JaUp, »I «*air error t It la how<yn iurjatir>r, aithiin,. I
ilegrcr, locrcia4vrn lulirrbu 1* alto a rrputad |orgaUve. Cutcimn», itlvitloni, axl Hntlt.

1. xJb* i.h.u.A*i'. 0 S / a1. 0 3 i II3S Piif*M	09 Boldoa
431 Hlrmhlli. A. 0 1 U BowlUaMtt U	
U 8 I	

301. O am a CMU. AMAflAVI Ifa

Ilit till/ Iterbaeram Spet-lei iA. Jo A*t; ,
IhU ontrt br vnn M'rthut h»» ilit- fntliwirific remark*;— U » n , nrrel „,r i la*
tn'turv, itmundiiK n tnd thereftm At EonL H".
are hrlnaeMW, eoaoiating chW< it 'ttrlr virtu** an> .
lh» root of Oomphr< na offlcnlilu l₂ Utuu- and dimuUnL Thr t|twii< are .
aaaUTdimwcand vtlMu, and <aUtln| in dry Monj <ipoaæ VAttr* or err. t and miiruitg on
: uni thi*. 1. ..rrati; a few are found tn ut
!in»llv, •

>>i.l. Aiii.iNK an

*« 01 7S5 (Usddataeha fx Am, WI A<hyrcnthea L 0	TS3 l'tth;x.-nti fi Br.
7.« Oattita /.. IH Deamocha>ta Ht< 0	757 (rtinf>hrkna /..
!«* 11 Br. Aitri.iTitjc (I. Br. rtik.	v.- Opiota* i.VK/

US CtiatnMbn II * IV

8

ATURAL ARRANGEMENT

til. OKDIB CLXIX. ANTIDE'SMKT.

*Genera J, Spffiei' : llal-ttmie Sarrlrl: GfWii-*o>u* \$p*tit* 0: Ifardg I.igiewu SprciciO;
lUrdy lhrbac- . . . i - fee*; A U frH*

Tr**«, native* of the Ea>I Indie*, with •mfile leave* and liwwitnlruout diixriou* (lower*).
an acid eatable fruit, a> well at the Pi*frlt of Atiliiti'mi, the*hang In Hmlcr like currant*
ili-cutt mi! of[heleave* i> rri'miM] to be an autdote igahtit the bite of*CTpen(t. liming*,
nod AiitukMi L (S7< SWago /.

CLXX. UHT'CJCt.

*if Hft<K*na Specie* 38. f.14*

Few art Ilm riHtyU in i"> 'Tlrr doemng the rare of tbr cultivalx; il II* nather nlraardiury, ha
that thoM- fctwt abundantlr •& A twio» worthIM w^d* and ahsby half.h nfaoou* *hrub*.
which aev ooewa** WKB jt . . . > JHM*HWO I P^g aMwaj w# And the rig, the i
iicip, ami tlMbrad.fr i

iBtMdlti

<i>*jH**aj</i> IftMi	<i>JUH^IIIJI, >HTJ.II, tillrtjtti</i> AHKASHIEM* . TV not oTU itet d	<i>IIIIMI, brytihally mi</i> own intniutnit TurkiUi drug called Bang or Haktakeli Th , tetii alag i..it f In OOMTft* ' r . . . rir; . . . ; in, . . . ill. * rniarin*. *tiimiiing-netili*, anJ other*. Cutting*, !*«*, dirblan, and Mod*.
--------------------------------	--	--

<i>iUaVta, i D</i> Hint niumf CrtVca /.	<i>•rlraria /-. I /.. I</i>	M (iniitr.i K ,* I. <i>... Mi rrirlo.l Kill.</i> <i>r> H rm» Ijii</i>
---	-----------------------------	--

D H I K CLXXI. I. I.MA (iii:

llrmfa -I. Spedr M ; llo-kauf Specif 5; Orffn.ioMtf Upton I I liarry /ipCMM ^wfVn »j
//u ry Jrrbaetm Hpetim a J Mj hat; C 0 feet j A 0 led.*

Many of llir oha<>atknu upoa lh# but order at* aim •pnble to thli, whirh iiiffrii iah(T
UflinlOtl rlurael*!*, than In any arrangement of nature. Jtie l'lui if il* reprimMaUtrt, ffin
< thm only ilifflly difflbr. B—Ui, layer*, or by graft*,
413 Plinara MI. 9 0 | nil L' imu* t. *)9 0) 8870 tVltU i.

ni oaririt n.xxii, riri.u \ i KJC

, Jtorrw-J 71 i ll.x.hmiir ft <rlf>ltoue Specif* 0; //<i^fv tjiuimi Spfciet Oj
;/i> i^v llrbartmu Specif X J , . , >. 4 feet.

|ir[i]r-ii m far tnnre valuable tn cotnawtct Lhan inlemtng in rult.t.v
i.fin HI a
ftt^r^r^i>jj*# a* ihry are aoajNrtlatti

the (inter i
-t u- c- ||||>K i, dtv MI ", ml JIMtI
Saulitmt ; n M /'i|ar /.

OanKftCUCXIII H BU

Orneru S, £ "i/- Hprtn 01 drtm-hovtr Sprfrt 0 ; Hardy I ignfuii Kp<*

H,i,f, Hrbat tarn Sprefi (L. f 77 frt; ^ 0 frt; ». D i*

U t n tre**, rhienjr n idm nf S'.mh Ami-ric*, with pinnat IsmrM and nnall liulmlficant
kamel of III* nut* of all are ralatlr. J<i>dal<> regia I* I ho mm man walnuL Seed*.

«IM Jiltlui* I. S 0 | MBS fferja S'ut to o

lift Oa>R* CLXXIV AMENTA'CK.lt

Genera I ii, t.kmtt Spn-ift f; (irf).hatut Rptrt i*

✓ HrrbtretiH Specie* 0. J UM flMt; f 'i ·*

1	Mg c n.....	0	M L bi.o.iir. IV	PLATS.
	MI OfeyfiiJ	1	U	
	SkluinR* 111.		MJ	« 0
	Ug.m. trfcIX 1171 a		t'1	8 0
	•it t.	0		1 fi
		7	H	•4 u

MS. ORDKVCLXXV.

UMJum* Sutrtft <) (!refn-k>u^se <>pc. > JJg»MU i
iSpri'it.v I I > >ifwt; iOfWrL

iw OHillrrad tjuite ort'i&l. NuUtuig ii

S 0 | JJSJ l«dlir(tUL(I, 4 0

4S& O a o u < i >-B.

7, tyfrtrJ 1 <(AIUOMC Sjxtitt **B**, OrtemJkimtr l^fit* <>. «<><> J>>,
itordy //rMa... f WBib* (< 0 (r.-t, A " (m.

III 1-1
mil k<
i> now t m t M l the

of r< •

in

"(Jr. T\ e

A few <rr Uta w.utlirn

Ar/i. a 1 JSft.

Bfflu

, o
,,nil Ct'PRKS-

W

i ann /<nr.7.

'>W. Krrb. a f 6.1 ft.
i3 O KM t,g ex her, a 1 m n.

It It

4ST CLXXVII.

J/orrtj/ //, i/^ ? i •

• V" *! rhUh~~h~~* with otMcur* flotrrD and twrriw, tMtivo of l-Uicq* and Nortfi Amwka.

JT1 •1 0 | S737 J. /) <>

It MONOCOTYLEDO'NEJE, o* BNDO'OSHJB (m-i.l.,
to grow).

4TOB i //ui-Autuf Rpeci- i fn.AiJim' Specie* Hardy TJgmnm
I It-I Uii ft Mi Sprit?* SfWL J l'ij i

pcl li
it '1 <
in it i- lii
I.tittt, and •

utuilr*,

cillu-i

Li mill A»Jn*i**li

tss, O IDH CLXXM11 *,

v ruriri* i

ijwrit tint

MJ nr<

« Um Uui.k. »1

B[ij>,n..im iJ]

ilm

M • jr<<

«'1, Wihili II

•*»ntaitinl, **tujb**«

MATERIALS & APPENDICES

49b) Outi Cl-XXIX. HYUBOCHARI'DKJt

(!(*•« 4, Spetict 5[// i. "•" " * > •; & 3| FWT
 Floating whlrl. fowmd plant., of whin I roajtUc They frntn.ti no known prop"***
 ml)M<>I > ' if II lar chat.. * tetnf In HEMOMI lartcKVit. Ttrr ♫*Cr> .
 of vartout |«t(> uf ilia world OBtrU and d>
 7735 v. | Damaahnium ScJtrkr.
 iT*. Hffeedla <O III «S SUatUaa JL *0 1

43a Oun CLUX. ALISHA-CJ

3. Sptnst Wt lir*^t Ugmtimtaprcifti
 lUD *vfir. I I J I nūt i ATlceL
 flaittbnm* water plant*, with whlrl Bowrrt, am.l nunt Maria. Urnir are rtmnnon In oür KIIK1»II till
 other an """.I in limutl iltuatum* lit the tñpli-* ItnVu, itMdilMt, ami amta.

9VJ tmthI

III O.n

Ofmrt ft. J M n 91 fkttnmmt %f*nn I, t. *<•'> *nūr .4* ;r-*0;
 // N ' .:mi Sprirri 11 • ,

M* I'LIJIU, OfwkM i>li(i)>iut, ljr pw<r4ls)Mmt Itir mnct brsuliru' J*nU,)IM
 flutrot; aiul Liuuui'luru, a n.iit r of the nurvoei of Acutl, luu Ttfccnr OHM 1 >D i.icn ami WCUL
 12» Atgmu /.. 8

r

«mcrn 2,

.niAcant bof |OarΦ, wMh p w f UBTM, and crnirtl «plk» or nu*tnm of ((Twtilth

-O- I ! , -O- II .It/. .:|

UUCXIII OttLIU'p

Otter* 111, ij.Kcmi* Mmttift irll; OrrtmJmut Spet iei 111 i Hardy LtgmtMu SmrcnQi

1 itiix-i or ptani
 fitmrtrarp t
 , ami lnr lhr .ti.nigr 1
 n anaaf (ruac«, and aooMtinn

•iiii-t pmnun . ! v.
R MH, on th* rtmtrarj, A

iVh Ma. i
 that th> pollr

A 1 4 i t: * i. i, i, ; *i, i. ati M I H M I *** \jtiiii.i'i MM A M I I tin* I
 VantiU. rrM M I h. IMUHI alea* of WOW •/ Uir llr&illun rpai-lr* a kna<<u. yluc ».
 Al pun-wca. UtoUkoo and MI

K5

THbe L N(t.-i 1

lf.Q. hrrb. in. tf H

n 11 Or. <0 3

mltsAraL

/fr.

< vi|a W. Br.

illii.vn It llr

um /' <r.

•0 B

Wit. lrl.'»U /'

ho* JSCL

431 Tribe t Aaicrnit'aa.r EAMB
 W. 0. hrrb. n *

SS13 Arrlidii l.

tSM , '(. flr.

CBIS Pof<nU W. tr. U

' .11 Mi-

logAith. A. Or.

MM U. Br

u.ilia / kV

, <r.

•0 j

lialailhrra /VA

R flr. »0 a

Sf. CM yit It. irr.

iraanthn R. Br.

•ibeo a am en ri

b

tf.7* Prr

437. Tribe 4. OFIIT'IMSJ: Aiwr/
 r. a tar* Sl ft,

him S, :
 " II

Mi

««M Anarainjitt AkA.

flWB V)thtflL* #fr*.

WK7 A*t*ra* H. hr.

Y4vt b* phry# i.

•Hi <•".!. Mini 7./C

»r

rtulib4lanl It fir.

t .• i. 1. . II . (,

fl. beilferiu :

fl., ft * i

•II 3

11 Mr <0 1

; l,

HIIM //

/

»r

rtulib4lanl It fir.

t .• i. 1. . II . (,

fl. beilferiu :

fl., ft * i

•II 3

11 Mr <0 1

; l,

HIIM //

/

FholblbL. I.

M r * 1 p...

Off..15 . .

SSM Cmx

»r

rtulib4lanl It fir.

t .• i. 1. . II . (,

Tnbt 0.

»r

rtulib4lanl It fir.

t .• i. 1. . II . (,

Tnbt 0.

»r

rtulib4lanl It fir.

t .• i. 1. . II . (,

Tnbt 0.

»r

rtulib4lanl It fir.

t .• i. 1. . II . (,

Tnbt 0.

»r

rtulib4lanl It fir.

t .• i. 1. . II . (,

Tnbt 0.

»r

rtulib4lanl It fir.

t .• i. 1. . II . (,

Tnbt 0.

»r

rtulib4lanl It fir.

t .• i. 1. . II . (,

Tnbt 0.

»r

rtulib4lanl It fir.

t .• i. 1. . II . (,

Tnbt 0.

»r

rtulib4lanl It fir.

t .• i. 1. . II . (,

Tnbt 0.

»r

rtulib4lanl It fir.

t .• i. 1. . II . (,

Tnbt 0.

»r

rtulib4lanl It fir.

t .• i. 1. . II . (,

Tnbt 0.

»r

rtulib4lanl It fir.

t .• i. 1. . II . (,

Tnbt 0.

»r

rtulib4lanl It fir.

t .• i. 1. . II . (,

Tnbt 0.

»r

rtulib4lanl It fir.

t .• i. 1. . II . (,

Tnbt 0.

»r

rtulib4lanl It fir.

t .• i. 1. . II . (,

Tnbt 0.

»r

rtulib4lanl It fir.

t .• i. 1. . II . (,

Tnbt 0.

»r

rtulib4lanl It fir.

t .• i. 1. . II . (,

Tnbt 0.

»r

rtulib4lanl It fir.

t .• i. 1. . II . (,

Tnbt 0.

»r

rtulib4lanl It fir.

t .• i. 1. . II . (,

Tnbt 0.

»r

rtulib4lanl It fir.

t .• i. 1. . II . (,

Tnbt 0.

»r

rtulib4lanl It fir.

t .• i. 1. . II . (,

Tnbt 0.

»r

rtulib4lanl It fir.

t .• i. 1. . II . (,

Tnbt 0.

»r

rtulib4lanl It fir.

t .• i. 1. . II . (,

Tnbt 0.

»r

rtulib4lanl It fir.

t .• i. 1. . II . (,

Tnbt 0.

»r

rtulib4lanl It fir.

t .• i. 1. . II . (,

Tnbt 0.

»r

rtulib4lanl It fir.

t .• i. 1. . II . (,

Tnbt 0.

»r

rtulib4lanl It fir.

t .• i. 1. . II . (,

Tnbt 0.

»r

rtulib4lanl It fir.

t .• i. 1. . II . (,

Tnbt 0.

»r

rtulib4lanl It fir.

t .• i. 1. . II . (,

Tnbt 0.

»r

rtulib4lanl It fir.

t .• i. 1. . II . (,

Tnbt 0.

»r

rtulib4lanl It fir.

t .• i. 1. . II . (,

Tnbt 0.

»r

rtulib4lanl It fir.

t .• i. 1. . II . (,

Tnbt 0.

»r

rtulib4lanl It fir.

t .• i. 1. . II . (,

Tnbt 0.

»r

rtulib4lanl It fir.

t .• i. 1. . II . (,

Tnbt 0.

»r

rtulib4lanl It fir.

t .• i. 1. . II . (,

Tnbt 0.

»r

rtulib4lanl It fir.

t .• i. 1. . II . (,

Tnbt 0.

»r

rtulib4lanl It fir.

t .• i. 1. . II . (,

Tnbt 0.

»r

rtulib4lanl It fir.

t .• i. 1. . II . (,

Tnbt 0.

»r

rtulib4lanl It fir.

t .• i. 1. . II . (,

Tnbt 0.

»r

rtulib4lanl It fir.

t .• i. 1. . II . (,

Tnbt 0.

»r

rtulib4lanl It fir.

t .• i. 1. . II . (,

Tnbt 0.

»r

rtulib4lanl It fir.

t .• i. 1. . II . (,

Tnbt 0.

»r

rtulib4lanl It fir.

t .• i. 1. . II . (,

Tnbt 0.

»r

rtulib4lanl It fir.

t .• i. 1. . II . (,

IL AKITAM.I MI \ I

Trite7. Utiiiiii /..*«	AV In .Vin
r,7. *3ft	I lift.

IK; *ti*
Hmrdp He,
w' 'L » .t.,.,.,.,.,.,l ^ .mtl.r
unluUr ntarium Ti

LiMiyutn Kim

f 13 OttMtCl XXXV

Hengt Ilrhacr. U f "f" I
»mt ?**** from the privedirvc, in tin- aliamrr & si i (it
•tre crt rftbrir .nil ..n.l.l.,
"I nutt h I M M ...:, MM MM OF

IKI. III Ji.]<ni,iii. r, 1 **talurr, iii!**

445 Own ri-xxwii. pit hi

«4d QHDCHCJ XXXVII) BJEHODOBAT1

21; Ifitt.hiniif ifftyta 11 QrttnAtmt* Sp Hardy J.iftiQiu
) tartyt Uerbmcctmi Specif* 1. JO fin't, » % 0 IM.
, .^ m * * ^ J * * <w. dwWwl from Jtsnfin, Wood. Indii tl -(riinn;t (MHuliarity j U
" " ilx .. whir

NATURAL AKIANU

«S7 OMDEK CXCIX. PONTEI>K'HE&

fr. Unrty tllrbace < nr**HW Spftun 1; Hardytiq
 it aquatic illant", usually with Cotl. d In w, and central *plko" or raceme* at Wu#
 Nativn of North and South "f'l
 it | US IMmnthi-ra A***. 0 1| 1» Uptaothui Us.

m

omtzui

r Ugntota Spftitt 0;

* IT lufe u

M<tlr Inhabitant* of manor mound ta
 AnHrrki U thru ,<>! .l>U,<t K>J••• i, but mil known In Europe rccrfit in cultivation)
 H eat MpUMa,v*waf!> with blue, rvddlah, or whu* Aftwflt Dkhart-
 MM or the order, with large Ihjrac* of Nu* Dovtn. tht* color la tha prera

JQDI Ufahariiindf* AK*

0 10 4rtw>fcn*I
 ill CBMJE

0 1 118 Andltm* It Ih.
 IIS CummHiiii I.

*JStt Oanu CCI. FA'LVJE.

Ofmrr* <, *># -ijtuw Sprftf 119; tirr**if AwrVi II, //^rfy Ugnttm Spteki 0;
 //•itjfTtHM ,1pTlft O. f Ofret; £

The*c wef* well n pt ur iln- r*, far *urn>ti all .
 iii HH p a M i a M M M M [nf n m farr. rbindrlifly •Grit, m'nwrM iy t H M M i d f 4 H M M r.<.^.
 .. horn lUelf at the nrrumfWrence , tt it alwaT uitltranevod. bearing a (uit of intn at Ih* I
 awar* found, and It I m . m l " hy a *ityt* bud , br Ih* fall of the pr>»U of the MMMW, whMt *J
 In • irratr or foai o*gra" at Ih* late, tt la rovernl with largv *an 'Itir k m , HrhiluiBj sattod
 I pinnal* or tatliefonn, wUom *Uapl*[and, In a youna »*te, tifare UMIJ «S)

MI *9*a* «a>it, of a pallid color, and coitainod in a large bag r*llt-

H H N W c*iwd a "f*!*) bunbi m M M M y Uuwufb, tp*, wUhw *u*
 r Mi (rafrant odon pbea rapidrl, ihroua-li I u*

upon th* turfare of th* tUgwa*. which gap* open to r*
 Kind riryin* R •t* monUtt a jrvr; when rtur » is a drupe or bt-nrj. with elibar a t
 MM of IU kvrnd nt Mr alhutnm, whkb, in theriaeof the roroa nut, H <n
 I M h t n i f ", the cvlrbrated Ira..
 ruMitml p all itiAl ii ki>own ui tliMv trljut^, cobrudaw h

b# fuir.o, Mith HJLI Jitmtjii/H, ofalHud*, nor HiUMnnrthrT.

C M M to put that wherever a dMfk

CHN nu- Ut*. ArtiK-fni* MkNttrM
 nMlnaonUie DM* of in* *arth.

MM of them Unr the mar

I f*t a *mltrr num

•rallred ..

I K- MM " I .."

^x elf •

that I

il amount i> at many at ll

I U ll..n,M I.I., Mi, II., ...!

MMtn . MM rolle. I In .f.

9UI L

Hum, I fll

VII 1

III., li I I

I * tit-Miiuin fimi.

'alia fimm.

vm Gomblut hum.

i iMitm CM *

Hum,

Itort.

«!U Oii.. < CII

Genera t, tpttitt 9); tit k-we Speeft* 11l| (ir*t*k<.*" Hardy JJgiumu Sptiirt n,
 Hardy lirrimrfout f » frei i A « <*<

AII *rv UfijileM].

mo

I ii

| SSW Phytitoutw. /T. t/P

OaantCCItl rv-i

^& fr<v threat**! Spnrt 8. j^O fert j £ 0 ftvt; ft Tftet

RaedJlk* pUnu, common rv*ry where the »it Id In l•b•• and iii. ln-i, u>u.i!!> »ill>
 nx>t>, and kn< *oft, or tuund iitauriw-niri. All iialim fr Lntaln.

NATURAL AR1

47t. U*DM CCX. CnAMI'NK.C

Otmtrm ISt, Speting 10711 Jfrtr-iMIK 4rVt*» 119 i f7rrt*.iM«K- & uvi/j W; /Ajfrfy f^IM M.Wei
Herbmnm* Sftttt* 8ftt J 0 fm, , ^

Thr order of j r u m it brond ttoubt Un m ri natural of all thai UM inavtiui* of -ftnnatic bolanli

ami iir tural tmrlpm, which an bradaai la a pragaaWa atata to tlm In ..(»m ait1 i-rul.. offw
ticikinf character* by whi.h the t|t^cmcan bacbntiUrkmL Mmcr aU that the A i l t f l W u Uicon.
ami IU divUlw Into «mrr». ha* not tmrl b**M fcund MiUvMrr difilcuH, but ha* g'-n'-i— mnrh I
^opinion aiftai-bi4'nli«i MM of »lwm, »fh(Tin_N to the •muWtcal arr*n-t*arat of Uunm u., .
• tault number of n M n, vhl* oftern, admittiu rbr inalTUral prtoripto of Modem artMna,
Into a rart nuntacr. Th» nMai> •ow aairtk H caAM, U mtabii UM-IUll
•ruionof ih-<-ur-Ur Int. tnb» h«l|m uirt>, *r mountiM. KaaMif. X

be paMcD over)>tUn« i > ><..1 twa ra'r, prop*!-!* Utat ir< acti
aoaout, poMMtagant, -litWr t|U..

^k^f1J1-13B1Mij In hit'r M-
i-t....» |ta.....*...*...*
ry#, barWy.
Ffuijtkni(ihji if Oka-ttiaRtr I
-ml MI ml. t.... . . .

<HI* trhhttaIM»
») inij-.H HM i|o> . ! thr *.-<..
^frt kin-ct cil lf rmh It niutt ho
t .. imrllynoacc.i

N M

'htruta »»»

II » si* -
«f, < r nut |- I nur M
i flum bar «fl

i4rfanmt fliliilill the utbm, W

*hr<t, (.

th.tt.lofi'

thr trrii inf *

o lwi umiti.

that thru • p**tly **fTT***

nunui

tt Hul erim niypata» ull INJt

Mil- l.,(.-r-t, fr ... »l IT. ..

IH M V M b M I' & a M d I B j (.

i, tl., MB) 14

• «a M «Li li E abjS mat • ' <.

....la(. .1., U i,-stl'i thr

tU terminates-

471 Btettan 1. Omrirain*, iAr- 0 * T * , ft < a ft.

• 9

III! l.

litui llraut. o 1

471 Jtectwf I lauttaM
Ig 0, hrrb. % < 10 ft.

*71 .tilimn /.. *0

474. faction & NA'RIMKM.

W* ft herb. I. C 1 fl-

178 Mkrd» f« *0

475. Stction 4. C;«'frti»t«JiL
(<. 0. herk. 4. * 4 ft.

170 Oiwhrut L, n 3

MM I nttnthfc* Dftv. o 1

MQ UM-littr. il 1.

SB1 A111 li ' jilura SeMrtb,

474 Section 8. I.*rr*f<(MUL

/i<. U Acrtf J. f_ *ft.

I< f,a]i|4go *A 0 1

477. Berloti tt, AxitLdn»»..
ii». (i. Arft 7. £ 0 ft

W7 X tftttKn J 0 7

47K. Hoc ii tm 7, 1 lu»nr *(>..

% A Ai-ri (. ,

Hi f/inl<-tim /

470. Sadlcm H. Tmhityi.

tig. 0. kerb. Ul tf £31 ft.

B73 .far tlv / . A fl

V70 A llitur. m M

, JUT. tt

SOt Wnjtrnli.jrow Trln. 0

Silicitr tubtern itiolet.

1/C Strlinn t. Si*»' a.

tig. II hsrh, 7. %

174 ttpartlna Schreb. »0 7

MI. Bction 9. P u n t a i

%0 Acr6. I> i

178 Mitaluni 1 0 10

1H Crritte l'm

4K. Sactioii ;t, CrwnDO.vnu1

tig. 0, *«r*. (i f 4) ft.

«0 Cfndon ftoSi 0

481 Stetton4, Cnioinii

//jr, 0. *»». B5. # l'l ft.

hlna lh-ttu*. 0 3

. II'. 4

fj 1

n 7

'f. O 3

0 1

1 - lu. 'killfliuum lmt 0

t

>u-ttlat ftt(»y?(W.

4H1 ntIMim J t'haUirkIMt

1 't' H ft. A

*0

-tl

tl

4M IllvULotI £ Agroi

tig. 0 Vrfr to ij ft.

ill] lilypbftiin •0

1

•0

33 Clini /

0 i

1

•0

0

0

4ML nwttttm 3. NiiUmr.

ft Arrft. ft f 71 ft

[H] .l/lliiin /.. •0

1'J1 1

1< %

IT; i

497. Uirilun 4

Kg. a hrrb. vi \$ <|

4ftL DtrtWon S. vrnrwrfi

Ar. a Av. t. 16. f K> tv

Kit rfclatriagnW. •U

Up BchwkwrM fikwa
ull trtrut.

MU.
» 0

SH

fchmk
tU CWabrtM fatt

tawnttmt

HI

lie t. 1

«7 Ketüncrta /

fil

£ S .-

179 Aoiphi|>-'(»i> it- Mr.

m ft,

S87)iti|<rt<l < 0 I

IHt AndriH*." *
tt.u.WWx 0 t

B 1

0

0 1

nm Tilm. 0 S

m OS

9Kif. I A.

II. FESTVC4*. CEJK.

Hi. o. Arrft 7». * WJ ft
^iftm /< lit

lit. a *& h. 1,

ritox.

Kg. a herb. 7

ik'Mi> n i iiun.i Ji

ii H

SM ClirhIrt 11

Microl<"ti> IWDK

** L..
! tmli li. 8r

J ASI

Hg.n

). LffMM C

Divtsiov, CI'I-I-I'I.A RES.

57(1, Sptciti MB| | BfcfcMi v>TM.»» l/gtum Spittit*Ot

Hardy thrb*

I fiT

fh>r>rtniilirt tf thit JIV.HLN i union tht
rrtiurki>t-i ti <m> i>li>ly ltrvn
tin- L.HIJ of lh< wirv
m M II> hM mill rdrrm. » a Mirl f<MI

FO1.I

fln-< AMF*- Am to Mt H Ugnrtmt SprtUt 0[

i w n i iri.n

*fti Mouie SaretfiBQ; ttndu JJjpuvmt I

jrtti I

tSfrnml IT

M I L I M >rr(ltl.!!<1 hy

"ong rarer .. *o
-in.illy iiKli trppi I),,,
H<<< 1 "m< hnvp a.
roan

BAI. AtlRANCi

WOTtilOI* an article of faod | ben la obtained AnttIIM reoUrf oUiencj ftwl, finally.
luw M M d u H Diruiwi and Mcda.

SOU TrUw I. J'iLvi-niti t V «*

S90I Pdjrbfc™. S 4 A
ichum L,

oprTMtrium 3< «O
'umAa* •0

CtfU 41 antum L.

WIS 0 Mdt» L
hrlt
AfWm J

0
*U

um A •!•//.
• M. ,. ,. L. « //..
llr.
.•(• /n M M M <
m.r>,plVltumS<

Tritwi Ormout6»UL

V>7. TtitwS.
Mariltid Dm.

UM. OUMUIL KOT ISfcTAXEJE.

i 101 Mfla— • ifTfft 10 tODI .

Md H* nmuiuibto for a hjrironutria*! mayMMtit
j'urninljr (lj) wric **UMI** In nwddinr M
>'(itmpkijred for tea, and a* a |w*Uhlfli matalial tot

BM

Hi LYCOFODI'inUE.

(hry hJv* lh Mod pf fr
ptimn ranbndl Vtfc
ui b* rmplitrnl
aaunabtr nature. 'Itm

n and SetkfQ U

•0 In

£10. O w n IV. H.A

Urmrrm 4, h d n 4; Hot-hitvtr Smrtr* 01 Ortm-*ui*f Sprcin U; Hmria Ugntmu Spnift 01

Iini or mrt (IwpIrr.Wf^t |.Unli. now The Mir.il. «, which «r» lo «inc C>M.
I tl U>lhil, UtfrOtkBDWII In nillinh...), 1<

S057 IJOHM JL • f i l l saw fulgtiiii V 0 1
ilidtrU/* *U 1 I «(» Mar.il 0 1

CtkuEi V. MI

ftwrwa :17, Bprtit* 3X1,

WtnrVT piano, nvlrtai in humid air, »i.umunt Jbou ihr |«.ln, tM m Hn- rquaMr. Thry rovtlhr I
talna of lhc r{id ai hiyh .. l'M- tir.....

*pnta mill
and flh up bufi *xl nwaaMt with vtgttabU a< • DM remany of nalun
lubitTTlrril than to th* |Arua*Mai nun. Mnlniuit ailiniKcnl |ir^x'ittin wert lirtmerif «*CIII
but ttirjr iff now ovclactad or AiryDttai.

611 Oasu Vt HKPATltJS.

tiilg troall itlanu, wild then Uavm arraiitrd in an ttirkatrd manner. They differ ftum Lulu'''*
luit. color, and fruit; fr<n M<>. In the Milnmntt) at Iiulr cjuile Thnr <jtiall(ic> 4tr nil,], n »>J'
•ratal aMflawnat

It APHTXLJK.

Oi-aurn IT7I, 4MWta * SS.

under (hr foltnuring nnlrrt b HKK<arnmnd undaf the tame ordri In the Linnr-iii
r natural attlnltic*, nr »<t lire rrjM M

513. OaiKK VII, A XOJE.

UtHcru 7°, Sprciet 4it

rianta Meandlnf from the nnrjlnt ftwm known in *«*Matinn to a very rninpnnrt *t>te. 'Hie
lllform, Inlw, with lhrl frw tncalleli Imaieranl, On, hlfhMt U.
•n Indebiamnt wwt.lik< prn.*jt, nun. Bowa rnpuiit> lib siititui
the lowest arm* frtrn, la the hlghest IM or ptt

i *Tinii4l at ixrotiQifitL U)4 Ottim f ftilnd to lb* length o/ mini fathom* T ^ T jrow at th^{**}
 ? h with iiicf jnm <• *• •*•?••> <* ~ <* ntf><-
 L o**"1 <* Keotad Ilka vloclu. Tbmr butr I

61*. O m i VIM.

wAtl in ib» Moacnr of Mtui*. M P»«V<²A II»- "•••* urlh
 but they «IT, uori

4 I t

MM

SIS, Om u !

iirmra 159, Uperit* 1157.

now reached the Jovm ttaUM of *t>> PH in »rrivin(»), vhlrh

4 Bi ii» of » - jfy ctmiuwvi. M'^>

SE

• • * • «PP»ch Uw njuatur thrji Men'

Uir inf^t prfcct cIMuflrlion which tiir tirrtrni IUI C r^f

kMntMf

•twlfhrd to itit

K yrt bwn dm.

873) TfirlrnVliuluj Pen. | « H Utophfluj 7»ML

GENERAL INSTRUCTIONS

COMrmftttENCtKG

THE SYSTEMATIC AND ENGLISH GENERIC NAMES, AND THE ENGLISH AND SCIENTIFIC SYNONYMS IN COMMON USE.

Or. « tjf. *tjnonftae*, 9. *vanetf.*

C KNUUAL INDE

S<.	BJ	la*	Cm	Lin.	KM
* to B Ajtmeja	m al£4	a Attfrau	isn	if1S 511	A. tee Ma. ;
a ^Crfa.-	4159 550	« Al^pom BM	in	in	A ice Piptunifiiu &+!:
XT 5iS J(r»	am £57	Al%vifin>ti<	sien	itH 539	Aamtmtun
BJ jf ice Airdptu	am	v. «> AutiriWd	Sil	Anittiihri.i	215611 B30
B A. ace /r<na	SSU	A. »* K<>*ti/	(*)	AnAntherix	
v*. K« CaUbri*!	Si8	f. MW Kont(ta	£50	SM Jnaarhiiuum	
A. tee Corynephorus*	257	A. «ee	BM 4<W	Auaitaik-a	MM
A tec DncliamptMi	•~	«f tec Schivfrtckifl	410	Annthferum	
3 A WKhrMrw	Hi]	A. wee Wviciria	21B	Aiirhovypear	un
A lee Eria-hun	237	.fflwm 1508C	tsn	527 iHMM	IM
A we Kwlcrti	B7 U	Aljxis	53 A. MC Cvno^uttum	I*	
^ .«> PcnbilU	S3	AI wt;:a	53 A. tee l,harptronu ii	MB	
543 Airtjui.	BBM	Amada ginger IK	55 A. »ee :culT	444	
vir plant	13	Amnliigu HE	55 A. tee ttnoa	•43	
	[930]	S8	»» A. tut OmplialWes	437	
i «7 AiOca	117	JauA»Mn 7thi	BO	AlicitTocarpui	
Ajtak	14124	Amaranth	6123 « .801		
A > # M S BM	1599	.fwIRa-vTHiVLt, Or. 151	523 AnchtnbnNi	510	
** .VJUI-.	MM	linartiitiu	512 Andra		
A. »ee Ank*^mOk*	1161	». we Acn^thrifiin	3P0 SJ3 .ImIhjhne		
Akc*tree	121	AaH0*ni BSM 7'i	AndntVa	2W7	
f 15 Atmgia^ Or. 81	133	AmarfuU	an «B		
flatfmu	14M	A. »cc J-llium	156 S39 Androctmbium	UM	
SI Altta	tH7	104S	5^1 Androm*rhia	1117	
jftite	177% 1	A. «ee Knin*Tigia	173 na An.lrnixsla	BM	
512 fffitmn	337	121	178 A. ne LyatU*	1341	
LtelM	S437	A. ValRtta	543 &Bdnpk(AD	BSU	
la Alrvaiidium	133	^r:n-	410 A-M? All; •l)<Tnm	SJM7	
JS AUlfia	3017	thr*	410 A. H» Cymbopfgon	SM8	
Alder	t 477	AininJnM	410 A. mSMMiAcDa	s&ts	
** 501 Alrirnvithtt	397	Amber rree	410 A. tee InilKrw	SS43	
	3107	5760	410 A. «ee Illipidium	MH	
SS ^ isrec	356	SOT	50 Sxt	^, ice Aretut	
i «8 Alfeirl*	AM	A. tec	MB	A. ice ZlrUtt	
^33 Al-uritic*		..< 7163	504 Jndrow^mum	isn;	
A. keeCrMcM		N. rriiMn rtm^hp	27 jji Atutryaia	231JI	
Alnani!		57 CM pnmboRi	3BS	., inn	
Alexandrian Imi-		American ::	15 540	ani 133	
rel		• M 1 * B	27 49D		
541 A'VOM Or. 7.		MB US7	827 Anrbmbnc		
311 51) Jtjifgi	10811	HM	A. IK Hepfitij	UM	
• .	1383	..< 7163	A. tee rhalutruio	.	
JIS SSI JisiMJ	25Y7Z	57	105 RIS Atexhum		
US ' >re Ai-tinorarpoi	m l	504 MN CM pnmboRi	618 At.Rtl.Oi	5	
Tl FT A OtekHfi		512 A. tee Loncbocax-	A in ArchangiJka	3881	
SSS Attaminda	BM	JUI	47+ 105 A. tee fnidium	Mt	
544 Allantbdia	MI	cr<	474 51 A. ee OfWricum		
.	BBB3	6SS Amethfttm	107 474 A. we Stalin urn		
8 530 Allifator pew	10\$	5)1 AmfrUi	115 115 Angelicn.tne	7884 !U1	
1 w Orfbaphuj	iIS	5<5 Ban	BS 450 Angelicn.tne		
53P jfiltiim	103	A. tee Traehyiper-	3& 3& Angtantliui	ttalfla	
544 ALLwru		tn um	450 MI Angiorfdiuiu	IBM	
Allotfa	333	521 AlldtjtM>idTII	373 M tngUttotm	Mil	
AlUecd	1SI	AmaCenarii	ASSVM		
Atlipicc		171 T877 4 7S79	7* ^inrwtM-	4Sli	
X Al'ipkeiree		M Amm<r."inc	BM AnfnKtt	fi&	
ABughM of Cey-	133	19193	515 fnrirria	SS	
Icn 120	S3	171 M Amm<r."inc	sn AnicUnxluBi		
.. Almond	506 Ampd	133]	Ani(roz4ntJ)<£		
ft) Y* J'hmi	52 3	521 Amphi^ri'phU	A/v Aniraal art	ma	
HI -Moc	3	K75	42 Avkm**tm	a<	
«>ee A plrrm	W	S143	87 AttUintfaut	ua	
A. tec HawtrthM	*G3	Atnphchfrda	103 Aniw AniMMdtm	an	
* »ee PithitJ*ii.	9	AtnphiAbium	MB BB AniKdiUui		
(triii)	143	Aroptiuptmm	61 M		
* »e Ph vIK.ma	m	Ainphi<^rium	5J: A^SPHalW-	1<7	
	1010	3S12	5J: fnltum	S!H4	
	1009	SUO	Annrl,I*j	BIT	
*	eps	549	BM MB A'noda	MCI	
X laj ** A Wiaa	S19	140	537 A.^u,>U<>n	M	
Alflne BUtort	aS	13 BM	466 .In.na	MSB	
9 ST, I Itel!?	1MB	A. we BattuaodID-	2S7 *, *ct Atimifta	MM	
III 110	B	drao	2S4 A. *M Mooodftra	IS	
80		M	2S4 AnAninifM 17675 it	isn	
Hi Si! ^ ^ * >	an	Aatbi	2S7 Aaotti Amrrlna	19BSS	
	530	51< .DKtni a<TH	1410 4rti	S318	
	BBM	m 118*3	9S7 Anafimpit		
	717	#15 ffp	am mn	61 OnffiMwli	MM
	AM	A. «ee Swnecifpii	354 BB <.ae< Aft*r>	4M	
		Anicrlut			
		AnadenU			
		HI			

Lin.	Nat.	ftp.	Gen.	Lin.	Nat.	\$p.	Ocn.	Lin.	Nat.	ftp.	Ocn.
55'	A. see Clad&nthua	2383	328	521	A rectum	8237	1S2	Askalon	8619	1046	
M2	Anthlphora	281	331	A. sec BerardiVi	2246	143	Asbca	-	9&90	1139	
539	Anthericum	1064	415	518	Arcti.pys	2898	279	511	/fsp&lathus	1989	
133	A. see Albica	1047	173	523	Arctostaphylos	1346	152	A. sec Cycfbia	1243		
136	A. see Arthropb-	dium	1065	522	Arctopheca	2429	401	Asiwn	24386	2793	
135	A. sec Eulitiue	1060	663	522	Arctutis	2449	35	Asjerella	8351	27?	
136	A. see ChloropJiy-	tum	1067	260	A. sec Arctotheca	2429	463	Aspcrgillus	3246		
133	A. sec Czackia	1062	253	SHO	A. sec Sphnogyne	£430	55	itfpferugo	442		
131	A. sec Gagca	1052	47il	Arcturua	16062	18W	45	Asperula	350		
1.%	A. sec Phalangium	1063	70	Arcy'Yia	3205	49	A.seeErbdea	383			
462	A'nthina	3229	387	524	Ardisia	571	13*	Asphodel	1061		
410	543 Anthistiria	2848	382	525	Ardulna	573	5C.9	^SU10DE^LEt Or.	19?		
553	527 Anthoeicia	1814	30U	512	Aremunia	1209	4	505 Aspicarpa	31		
432	Anthocroa	30U	158	502	Arenaria	1390	157	411 Aspidistra	1204		
471	525 Anthoelclsta	3295	368	536	Arethusa	SEA>3	422	544 Aspidium	£938		
14	504 A'nthodon	102	367	A. sec Barthollna	<2484	420	A.hceyfpleniun	2921			
515	504 AnthoK.ma	1543	SG9	A. sec Pognia	S505	419	544 itfpleniun	££gj			
18	537 Antholyza	131	56	530 Areha	449	420	A. sec Csn6pteris	£922			
S97	520 Antiospermum	2760	69	524 Argania	BB9	417	A. sec Citerach	S9W			
12	543 Anthoxanthum	90	216	498 Argemde	1559	420	A. sec Diplazium	2925			
101	518 yfnthnskus	820	118	538 Argolasia	948	417	A. sec Mcmsciun	2J05			
102	^uthriscus	6668	64	527 Argyrela	49 i	4LX	A. sec Sculop^n.				
282	511 Jnfhitis	1968	£01	AYik.	18965	1507	driun	£924			
284	A. sec £1cnus	1981	386	541 Amarum	2674	107	Assaf'a'tida	C997			
311	A. see Lcspedeza	2126	18	537 Arista	138	285	503	1996			
534	ANTIDE'SMEJS, Or.	169.	85	542 Aristida	195	487	515	3593			
99*	534 Antidesma	2766	85	A. see Chactaria	19ri	341	521 Astelma	8320			
180	Antirrhlna	11558	374	533 aristolchbia	8582	341	A. see Heichrysum	23g			
549	528 Antirrhinum	1774	188	508 Arutotl'm	1428	93	525 Astéphanus	75g			
849	A. sec Linaria	1775	374	Arkansa	88832	8582	521 A'Hér	8337			
250	A. see A'temesia	1777	199	512 jfrmenaca	1500	348	A. see Callistema	S3SJ			
421	544 Antrbphyum	8929	114	530 Ann'eria	928	343	A. see £ngcron	2S&			
567	Antwerp Holly-	258	140	Armoracca	16323	1848	"6	Asterias	6230		
91	516 Anfchia	17859	2007	581 A'rnic	2350	363	Astensrus	82849			
516	%"		351	A. sec Bellidiastrum	2;15j	469	Asterocéphalus	S&9			
1G1	511 Abutus	1256	322	521 Arnopbgon	2199	44	A. sec Scabiøsa	340			
32*	521 Apargia	2208	328	Arnoscria	£0800	2228	Asterolinon	3841 sy.			
487	A. sec Oporfnea	3597	500	Arnotta	12 <i>425</i>	1506	459	Astcibma	31/J		
324	A. see Thrinacia	2209	541	ARoDEJEy Or. 204.	Arbnia	12 <i>425</i>	461	Astcr6phora	3V1*		
309	Aparinea	19077	2120	Arraccha	6821	852	517 Atob	IS8J			
218	503 Apiciba	1571	410	543 Arrhenathferum	2855	306	511 Astragalus	£1U*			
158	550 Aphananthe	1206	140	Arrow-grass	1107	307	A. see Gylden-	..			
421	Aphancs	{gg^ w3	383	Arrow-head	£653	SOS	stiE'dtm	2109			
152	j		1	Arrow-root	2	305	A. see Ox^tropia	2101			
483	528 Aphanochlus	1659	£27	496 Artabotrya	1684	305	A. see Phaca	2100			
545	529 Aphelandra	1731	107	518 Artedia	879	162	A. see Sophbra	&#P			
342	521 Aphetfxis	2826	338	522 Artemhia	8313	144	*08 Astranthus	I'II			
544	APIY'LLIE, Cl. 2.		445	Arthf.nia	3116	109	518 Astrantia	g^			
132	539 Aphylanthca	1044	136	539 Arthropidium	1065	103	A. see Di'mdia	3%			
128	5.8 A'picra	1015	136	A. see Stypandra	1066	285	Astraps'a	K'S			
315	512 A'vios	2148	332	Artichoke	8847	382	Astroaryum	**			
315	A. see Wistaria	2147	374	534 Artocarpua	2583	308	A strol biun	**Vg			
105	518 A'pium	855	386	541 A'^rum	2673	65	523 Astrolobma	..			
474	A. see PetroseTinum	3330	386	A. see Arisarum	8674	464	Astrosr* rium	<xt			
477	507 Alopophyllum	3342	585	A. sec Colidium	8672	487	521 Atalaanthus	3^			
410	543 Apluda	2846	203	Arufcua	13038	170	504 Atalant/a	1^g			
525	IPOCY'NEJ?, Or.	127.	36	543 Arundinaria	£89	123	Atamasco Lily	8019			
68	525 jtpdcyntim	542	89	543 yfrtido	830	103	518 Athamanta	g^			
93	A. see Eustegia	758	89	A. see Calamagrftia	231	103	A. see J'ubon	AS			
68	A. see Ichnocarpua	540	88	A. see Danthbnta	SS/	105	A. see Cnidium	its			
141	541 Aponogeton	1123	87	A. see Degrphnia	817	107	A. see Zig6sticum	J74			
166	Apoucouita	10718	89	A. see Phragmletea	229	107	A. see Sellnum	SIA			
84	Apple-berry	671	32	A. see PU	854	SS8	582 A. see fescli	XI			
fi.01	Apple tree	12955	27	A. see Psamma	210	3C8	Athanasia	g^			
199	Apricot	VAX)	187	Asarabacca	1421	358	A. see lhnat	o<n			
46	527 AquariVi	358	249	Asarlna	15776	1774	410	543 A. see Rejhama	ZzL		
5(9	AQLARI'NEJI, Or.	64.	533 j4s'aaiNjt, Or.	165.	382	534 Athcrobgon	iskn				
175	509 Aquilaria	1370	187	533 f'sarum	1421	341	522 Athcrosperma	*g			
825	4DT Aulegia	1605	333	521 Ascariclda	2263	420	Athrxiia	^ iSi			
2.05	429 Alabis	1827	525 ifscLEPiAVnr, Or.	128.	180	A thorium	85466				
857	A. see Aubrietra	IH37	525 ifsclepins	771	459	379	A. see Dibertia	*#.			
856	A. see Neurolbma	18S2	94	A. see Anäntherix	769	71	A'tropa	Rg5			
855	A. see Turrilts	1826	94	A. sec Cal6tropia	766	61	A. see J./andragor	Jg			
918	512 A rachii	2169	94	A. sec Cynancnum	761	228	Atraginc	ig^			
83	Arachne	1569 \$y.	94	A. see Dafmia	759	£88	A. see Karaviha	g/#			
112	518 Aralia	911	94	A. sec Gomphocárpus	770	159	531 Atrapaxia	£&			
112	A. see Sciodaphjl.	lum	94	A. ace Gymni-ma	764	411	531 il'triplex	fig			
		912	93	A. ace Sarcostemra	757	379	A. see Dibertia	*#.			
			95	A. ace Stylandra	772	71	527 A'tropa	Rg5			
			94	A. see Xysmalbbium	768	71	A. see J./andragor	Jg			
			104	Asclepium	6786	61	A. ace Xicandra	Jgg			
			455	Asc6bolus	3145	71	A. see Ph^salis	g?			
			462	Asc6phora	3237	72	587 540 A. see Sarah*	aff9			
			380	A'scyron	19699	519	A. see Atrapaxia	ipS7			
			293	504 J'scyum	£191	379	A. see Atrapaxia	50A			
			215	A. see Kordntca	1544	379	A. see Sarah*	aff9			
			414	Ash trc	£8b7	472 S<8 Audouim'a	574S				
			190	AsiaUc Poison-		472 S<8 Audouim'a	574S				
				bulb	7833	504 A 6 lux	Si				
					964	504 Aurstntmra					
					VolSQ						
						320					

tin.	Nat.	fip.	Gen.	Lin.	Nat.	■ Gen.	Lin.	Nat.	gp.	Gen.	
57	Auricula	3779	451	429	Bartramia	29'H	160	Bct'gigba	10376	1225	
456	AuriculaJbds	27445	3151	249	BartsiVi	1772	188	Bejaria	1426		
235	Auricularia	14874	1659	248	B. see Castilleja	1767	387	Bélis	23572 sy.	268-3	
453	Auriculhria	27280 sy.	3131	248	B. see Euchroina	1768	71	Brittadonna	4640	584	
381	Jvellna	23388	2663	249	B. see Rhinanthus	1771	121	Belladonna	Lily		
28	Ābna	823	&?	507	Barydsma	676			7897	969	
410	A. see Arrhena.			318	B. sec Dipteric	2174	835	Belveisle	Cress	16123	
	thérum	2855		111	Bassilla	908	412	Belleric		1822	
28	A. sec Danthbnif	226		240	Basil	1687	412	BeMrica		2861	
28	A. see Gaudinta	227		187	Bassio	1424	135	BellcvalfTi		2864	
28	A. see Tricha^ta	225		20*		75		Bell-flower		1055	
n ²⁸	A. see Trisetum	224				351	521	Bellidiastrum		607	
214	Avens	1530		335		3-9		Bellidiastrum		2351	
183	Āverrhixi	1404				351	521	Bellidiastrum	22022	2420	
2+7	Āvicēnma	1751	,80					Jellis		2420	
266	Awl wort	1903				219		Ielium		2361	
23	Ax6nopus	180		240	Bastard Balm	189*	411	Bengal Quince		2362	
375	A'xyris	2594		295	Battard Box	18263	2053	Benincasa		1585	
86	Ayhnia	684		318	Bastard Cabbage	3172	416	Benjamma		2860	
163	Azadirdchta	10865			tree		161	Benjamin	C2525 <i>i</i>	2900	
523	Azalea	521				j		tree	110402	1226	
	A. see Chamelédon	522					26	Bent-grass		203	
1	500 Azaras						161	Benzoin		1226	
201	Azarole	12924	1506					BciañUa		2246	
168	Azarbvs	12924	1506	M»			331	sn			
49	Axerdach	10658	1293				497	Berberidea, Or. 6.			
A	Azhma	3313 sy.	392	384			497	BJberu		1090	
527	Babuna	125					138	B. see Mahbnfa		1091	
521	Bacvzia	2235					476	Jerberry		1090	
342	521 JMcharis	2329		n			138	Brcchemfc.		678	
342	B. see Brachyle'ha	2328		,77			83	BerkhCyrt		2424	
381	Bâctris	2640					259	licgamot	14889	659	
224	Badger's-bane	14245	1600				170	Bergera		1321	
145	515 Ba'ckia	1172					187	Bdrgia		1419	
411	Ba6myces	31(13					483	Bermgena		3377	
392	Baldnghas	23868	2724				433	Berkeleya		3019	
422	Balantium	2935					401	Bermudas Cedar			
356	Balbisia	21867 sy.	2388				24512		2822		
528	A/7/fil	6050	745	100	Bastard Ground	14700	1648	Bcrnhärdia	25711 sy.	2956	
PX	ffalbla	1671		233	Pine		424	Birrya		1566	
237	B. see Leonurus	1673			Bastard Hare's ear	816	217	Birrya			
237	B. see Afarrdbium	1672			Ba-to rdHy<>y wi8ifiM		257	Btererb/i		1836	
483	B. sec Roy'ea	3376		283	Bastard Tndigo	1973	219	Berthol fetch		1586	
384	JBallta	83426	2666	56	Bastard Jasmine	450	472	BcrzeliA		3305	
239	Balm	1684		300	Bastard Lupine		253	liesluria		1809	
	BalmofGilead						100	5S1 Bbia		811	
		15201	1685	69	Bastard Manchineel	£47	13	Bctle-nut		826	
	Balsam	697		324	Bastard Manchineel		236	2tet6nica		93	
	Balsam apple	23676	2697	287	Bastard Manchineel		236	Betony		1668	
	Ifalsamna	23676	2697	390	Bastard Olbia	2008	233	Betony		1668	
	506 fALSAM(N&B, Or. 52,			47	Bastard Orpine	2705	378	/Jetula		14733	
338	522 ifelsairita	2310		202	Bastard Pimpernel	363	335	Bdens		1648	
340	522 Iflsamita	20886	2313	264	Bastard Quince	12981	1507	B sec Melanthëra		2284	
145	Balsamodendron	1169			Bastard Rocket		335	B. gce Osteospti-		2279	
168	Balsam of Capevi	1288			16614	1883	526	mum		S450	
413	Balsam tree	S877			Bastard Toad-flax	740	B. see Petrbrium				
362	SS2 Baltimbra	2439	305		Bastard Vervain	68	335	B. see Salmca			
124	Bamboo cane	984			Bastard Vetch	2100	101	518 Biforis			
124	543 Bambus	984		98	Bastard Wind.		46	520 Bigelv7.			
124	B. sec Mclocanna	985		233	flower	6355	242	Bignbnia			
124	B. see Mtstus	986			sage	794	242	B. see Ampin ir biuin			
	Bdmmea	17960	2014	62	Batas	147S1	243	B. see CheilApsis			
	Banana tree	6059	746	221	Batas	4126	71	B. see Gelsitiun			
48	Bandhooka	3272	379	53	Batata	491	243	B. see Jacardnda			
48	Bandhde'a	3272	379	527	Batata	23295	242	B. see Spathbdca			
216	Bane-berry	13757	1545	53	Batrachosplrmum	30'5	243	B. see Te'coma			
435	Bangui	3040		527	Batschia	430	526	BiGNONticor, Or. 130.			
182	505 Banisteri	1395		221	Battarrai	3180	93	Biiai	6148	745	
182	B. see Heterdpterif	1398		517	Batierra	1592	156	Bilberry	10118	1194	
41	532 Bñksm	325		165	Bauhinfo	1268	183	Bilimbi tree	11783	1404	
168	512 Bøbab	311		105	Bawd-money	857	84	SOL Billanifera		671	
163	512 Bøphia	1291		161	Bay tree	10387	119	Billberg/a		957	
162	B. sec Baptisia	1244		168	Bead tree	1293	319	Bilkitia		2183	
118	to* 5- ^{sec} fnagyris	1239		162	Bean	2132	63	Bindweed		*92	
404	Barbacfema	54		64	Bean Caper	1304	399	Bindweed		24986	
	Barbadoes Cedar			244	Bean Trefoil	1239	507	Brphytum		24986	
182		24534	2822	57	Boarbind	494	378	Birch		24986	
106	Barbadoes Cherry	1393		231	Itear's-breech	1722	383	Birch		1413	
	Barbadoes Goose.			156	Brar'g-eat Saniclc	452	200	Bird .cherry		2608	
122	terry	1476		173j	Bear's-foot	14665	74	Bird-pepper		1708	
283	Barbadoes Lily	7986	969	223	Bear's crane f	1641	298	Bird's-bile		1808	
255	S'rdM JJ	176J2	168	319	Beast's-bane	14186	57	Bird's-eye		2063	
302				515	Beauftrtz	1600	308	Bird's-foot			
328	Barbitrfo	2078		67	Beauftrtz	2182	191	Bird's-foot			
244		2237		525	Beaum6nt'a	536	300	TrefolT2			
35	Barleria	1725		525	Beccabniga	397	419	Bird's-nest			
155	5ade:y	875		7		45	25430	2121			
S ³	Barn&rdfo	MIM		32	Beckmannia	250	140	Bird's-tongue			
293	15 Isarringtonfo	20^9		100		348	374	Birthwort			
301	^ B. see Stravadium	5040		383	Bed-straw	19077	2120	259	Biscutella		
	55 Barthollna	2484				2120	326	R. see Megacarpic^a			
	16 Unrtbm.i	1177				2662	380	ST>5			
						811	383	Ilia^rrula	"io"		
								JJWiojTi-wewl	OMD		
								•m	MM		
								2^6 rta	10286	1210	
								412	Bitcoria	25040	
								385	Bitter Oak	23458	
									Sigif		

GENERAL INDEX.

.WI

GENERAL INDEX.

Lin.	Nat.	Bp.	Gen.	Lin.	Nat.	Bp.	Gen.	Lin.	Nat.	Bp.	Gen.
415	<i>Cassitrium</i> 25221 sy.	2897		335	<i>Cephalophora</i>	2274		412	<i>Chifbula</i>	85037	2864
161	532 <i>Cassitha</i>	1230		462	<i>Cephalotrichum</i>	3230		44	<i>Cheese Rennet</i>	2985	348
383	535 <i>Castanea</i>	2659		193	512 <i>Cephaelitus</i>	1467		421	544 <i>Chcilnthes</i>		29.S
144	507 <i>CasteUf</i>	111		436	<i>Cerramium</i>	3048		254	499 <i>Cheiranthus</i>		1818
248	528 <i>Castilleja</i>	1767		89	501 <i>Ceranthera</i>	709		260	C. see <i>Andre6skia</i>	IHGJ	
392	<i>C. sec Euchrbma</i>	1768		186	502 <i>Cerastium</i>	1416		255	C. see <i>Barbara</i>		
	<i>Castor-oil plant</i>		23954 2719	187	<i>C. sec Larbrea</i>	f1417		2	C. see <i>Cmcllna</i>		1870
374	535 <i>Casuarlna</i>	2585		478 J				2	C. sec <i>r'simum</i>		1867
3	<i>Casumhnar</i>	127	10	1P0	512 <i>Cerasus</i>	13351		266	C. see <i>Heli.phila</i>		1902
2	<i>Casupo</i>	G7	5	375	535 <i>Ceratola</i>	1502		260	C. sec <i>Malcbmia</i>		186*
88	543 <i>Catabrftsa</i>	220		462	<i>Cerdtium</i>	2590		263	C. see <i>Sterfgma</i>		1886
12	526 <i>Catalpa</i>	78		375	531 <i>Cerato6rpus</i>	3232		254	<i>CheWi</i>	16095	1818
328	521 6'atananche	2232		115	4%' <i>Ceratocephalua</i>	2496		277	502 <i>Cheirostemon</i>		19
412	<i>Catappa</i>	25032 2H64		489 5:/T>	<i>Ceratochillus</i>	931		216	<i>Chlidbnium</i>		15*9
412	<i>Catappan</i>	25032 2864		32	543 <i>Ceratochloa</i>	3412		216	C. sec <i>Kom&riYs</i>		1550
234	<i>Cataria</i>	14791 NvH		414	512 <i>Ceratonia</i>	247		244	528 <i>Cht'Kine</i>		1718
370	536 <i>Catasetum</i>	2iW0		175	517 <i>Ceratoetalum</i>	2885		243	C. sec <i>Pcnts'emon</i>		1717
179	<i>Catchfly</i>	1388		514 <i>CKRATOPHY'LLEJB</i> OF	78.			91	531 <i>Chenr.lea</i>		729
387	<i>Ciuechu</i>	22562 2682		382	514 <i>Ceratophyllum</i>	2651		279	<i>Chotdpoca</i>	17435	■59
307	<i>Caterpillar</i>	2105		389	515 <i>Ceratonnthes</i>	2096		531	<i>CHENOPD»Ejt.</i> Or. 155.		
49	519 <i>CatesturVi</i>	381		68	525 <i>Cerbra</i>	550		99	531 <i>Chenopodium</i>		810
167	<i>CathartOcarpus</i>	1272		68	C. sec <i>Dissolcna</i>	552		99	C. see <i>Kilchin</i>		809
316	<i>Catiang</i>	19497 2152		165	512 <i>Cercis</i>	1265		412	C. see <i>Rhagbdia</i>		2861
233	<i>Catmint</i>	1654		23	542 <i>Cercsfo</i>	182		227	496 <i>Chermoyer</i>		
328	<i>Cat's Ear</i>	2227		195	517 <i>Cereus</i>	1472		182	502 <i>Cherleria</i>		
375	<i>Cat's Tail</i>	2597		479	C. see <i>Echinocactus</i>	3359		199	<i>Cherry</i>		
27	<i>Cam-tail Grass</i>	213		479	C. see <i>Opuntia</i>	1474		101	<i>Chervil</i>		
2:33	<i>Cat Thyme</i>	14737 1648		55	527 <i>Cerfnthc</i>	439		102	<i>Chervil</i>	6646	825
571	536 <i>Catlycyt</i>	2553		95	525 <i>Cerptogia</i>	778		383	<i>Chestnut</i>		\$659
102	518 <i>caucalis</i>	826		93	C. sec <i>Mirrolbma</i>	753		502	<i>Chlca</i>		
473	C. see <i>Orlaya</i>	3313		419	<i>Crocpt4pteris</i>	25407		200	<i>Chickasaw Plum</i>		12875 1508
474	C. see <i>Platyspermum</i>	3332				sy. 2915		313	<i>Chickling Vetch</i>		19313 2136
102	C. see <i>Tonis</i>	827		3R5	<i>Orris</i>	23458 2666		311	<i>Chick Pea</i>		2131
473	C. see <i>Turgenia</i>	3314		103	<i>Crvaria</i>	6704 835		377	<i>Chickweed</i>		[902]
264	<i>Cauliflower v.</i>	1883		329	521 <i>Cestrlnus</i>	2238		1115	<i>Childchloa</i>		2
138	497 <i>Calophyllum</i>	1087		71	528 <i>Centrum</i>	583		27	542 <i>Chingl6ttU</i>		SS1*
84	508 <i>Ceanbthus</i>	668		417	541 <i>Ceterah</i>	2907		369	536 <i>Chi6psis</i>		17U
472	C. see <i>Colubr'.na</i>	3300			542 <i>Cetraria</i>	3096		243	526 <i>Chimphila</i>		1350
84	C. see <i>Pomaderris</i>	670			518 <i>Cethchosnora</i>	3172		173	523 <i>Chimonanthus</i>		1584
472	C. see <i>Willcmetia</i>	3301		459	<i>Cethrophllum</i>	821		214	513 <i>China Aster</i>	F&	
395	534 <i>CectripVi</i>	27.13		101	C. see <i>Wnthriscus</i>	820		288	519 <i>Chinese Rose</i>	17943 2014	
388	<i>Cedar of Goa</i> 23652 2692			473	C. see <i>Coimpbdium</i>	3312		221	519 <i>Chinese tree</i>	14094 1596	
388	<i>Cedar of Leba-</i> non	23636 2688		350	521 <i>Chaetanthera</i>	2349		80	520 <i>Chioc6coa</i>		628
86	506 <i>Ccdrela</i>	690		25	542 <i>Chietaria</i>	196		5	525 <i>Chionanthus</i>		38
388	535 <i>fdrus</i>	26B3		91	542 <i>Chiftcalyx</i>	2081		58	526 <i>Chirbn/a</i>		4615
292	502 <i>Ceba</i>	18143 2029		477	515 <i>Chaetogastra</i>	3345		58	C. see <i>Erythrae*a</i>		467
216	<i>Celandine</i>	1549		461	541 <i>Chet&ium</i>	3220		59	C. see <i>Sabbaiia</i>		468
387	<i>Celandine</i>	12092 1422		433	<i>Chaedphora</i>	8023		174	515 <i>Chibubia</i>		1358
412	SOS <i>CEASTInEJR</i> , Or. 59.			25	543 <i>Chxtbrus</i>	197		174	<i>Chives</i>	8620	1046
	<i>Celastris</i> 25069 & 70		sy. 2868	91	<i>Chaf-flower</i>	5967	727	132	153 <i>CHLENA^CEA</i>	Or. 33.	
83	508 <i>Cel&strus</i>	662		100	509 <i>Chailletia</i>	815		117	538 <i>Chhdanthus</i>	r	939
M	C. sec <i>Ceanbthus</i>	668		509	<i>ChallEILLETtcor</i> , Or. 64.			246	529 <i>Chlbra</i>		U44
84	C. see <i>Senacfo</i>	665		295	<i>Chambwuxs</i>	18265 2053		153	526 <i>CHLORA'NTHEJE</i> , Or. 102		U76
:i05	<i>Celeri</i>	6834	855	200	<i>ChamsccrasuBl28W</i>	1502		519	519 <i>ChlorAnthus</i>		S0
	543 <i>CELLULA'BES</i>			171	<i>ChamiEcfts</i>	11028 1339		463	519 <i>Chlorfdium</i>		3243
92	530 <i>Celtoia</i>	735		167	<i>Chamscrfsta</i>	10778 1271		410	542 C. sec <i>Athernpbgon</i>		2849
91	C. sec <i>M'rua</i>	731		337	<i>Chanuccyparisus</i>	20745 2m		410	C. see <i>Etutachys</i>		
92	C. see <i>DeeringtVi</i>	733			133	<i>Chamaembly</i>	8(9) 1046	410	C. sec <i>Rhabdchloa</i>		6U
92	C. sec <i>Lestlbud&fo</i>	732			212	<i>Chamiembrs</i>	13558 1523	410	<i>Chloroedcum</i>		7W
853	527 <i>Celsia</i>	1812			332	<i>Chamaepetucc</i>	20456 22.9	200	539 <i>Chlor/phtym</i>		1057
253	C. sec <i>Alonsba</i>	1813		157 j	<i>Jasme</i>	110177 1200		136	539 <i>Chlortxylon</i>		1P*
412	534 <i>Celtis</i>	2870		66	523 <i>ChamslMon</i>	522		169	506 <i>Chlorbxyon</i>	10384 Wf	
387	<i>Umbra?</i>	23609 2685		400	539 <i>Changuilium</i>	2792		318	506 <i>Chocolatcnut</i>		12
458	<i>Cenangulum</i>	3162		202	<i>Chamnmespilus</i>	12P81 1507		318	507 <i>Choisryt</i>		13W
22	542 <i>Cenchrus</i>	170		413	540 <i>Chumaj'rol's</i>	^883		200	507 <i>Choke</i>	12871 & 1287	18%
56	c. see <i>Anthlphora</i>	281		191	<i>Chamseyce</i>	17347 14C0		390	507 <i>Choko</i>		27W
29	C. sec <i>Cntotheca</i>	228		375	541 <i>Chamels'a</i>	22904 2596		48	520 <i>Chomelia</i>		*J
34	C. see <i>Echinaria</i>	267		91	530 <i>Chamisbtff</i>	722		438	520 <i>Chondria</i>		SOg
SD4	522 <i>Cinia</i>	2379		304	C. sec <i>Cladostachys</i>	725		163	521 <i>Chordaria</i>		308
443	<i>Cnomyco</i>	3102		354	<i>Chamomite</i>	2381		438	499 <i>Chnfrspora</i>		Wf
360	521 <i>Centaurea</i>	2434		355 j	<i>Chamom- la</i>	21767 2384		163	511 <i>Chorizema</i>		1412
362	C. sec <i>Gaiactites</i>	2415		367	536 <i>Chamfirchis</i>	2489		163	C. sec <i>Podolubium</i>		W7
360	C. sec <i>Leuzea</i>	2432		363	521 <i>Chaptkl/a</i>	2H7		231	<i>Christmas Rose</i>		
360	C. sec <i>Rhapontica</i>	2433		363	<i>Châra</i>	3047		350	<i>Christ's Eye</i>	21522	659
321)	C. sec <i>Serratula</i>	223		436	C. see <i>Nitella</i>	3046		83	<i>Christ'sThorn</i>		
587				436	<i>CseeNitella</i>	3046		434	<i>ChrO/olepus</i>		2396
SGI j				355 j				356	522 <i>Chrysanthellum</i>		
£60	<i>Centaury</i>	24j4		355 j				356	522 <i>Chrysanthemum</i>		237*
58	<i>Centauria</i>	3878 407		367	<i>Champaca</i>	14349 1619		354	C. see <i>Pyrdthrum</i>		940
29	542 <i>Cntotheca</i>	228		226	<i>Champaca</i>	14349 1619		117	538 <i>Chrysiphala</i>		
4	520 <i>Cntr&nthui</i>	20		363	<i>Chaptkl/a</i>	2H7		200	512 <i>Chrysotnlanus</i>		
355	522 <i>Centrosptrum</i>	2385		436	<i>Châra</i>	3047		357	512 <i>Chrysotnlanus</i>		
47	550 <i>Centunculus</i>	363		363	<i>Charles's Sceptre</i>	15882 1781		337	551 <i>Chrysotnlanus</i>		
112	<i>Cepa</i>	8601 1046		192	<i>Characias</i>	12454 1460		337	537 <i>Chrysotnlanus</i>		
184	Ceps'a	11854 1410		389	<i>Chardntia</i>	23677 2697		337	537 <i>Chrysotnlanus</i>		
	520 <i>Cephelia</i>	65!		850	<i>Charles's Sceptre</i>			337	<i>C tee Euthamia</i>		
566	<i>Cephalanthem</i>	2508									
47	619 <i>Cephalanth</i>	361		264	<i>Charlock</i>	16630 1884					
43	£20 <i>Cephaftria</i>	345		130	5S9 <i>Chailwo6di7»</i>	1019					
469	C. see <i>Pterocphalus</i>	3290	48-3	528	<i>Chasmbnia</i>	3378					
4(0)	C. see <i>Succisa</i>	3288	389	Châte		23704 2699					

GENERAL INDEX.

5£

Un.	Nat.	6p.	Gen.	Lin.	Nat.	Sp.	Gen.	Lin.	Nat.	Bp-	Cm*	
337	C. sec Ozothamnus	2295		Clown-				356	Columbia	2398		
C63	Chrys6gonum	2445		Club Grass	218			522	ColumnM	1797		
138	ro. <\$'y'B>num	9036	1086	Club Moss	2965			528	C. nee Stemodia	1795		
69	52* Chrysophyllum	556	424	Club Moas	25344	2908		512	Coluria	15-36		
r/j	C. <ee iftimlia	555	417	Club Rush	158			Colierna	83393	8663		
(>	..< C. << Nycterisition	557	21	Club	2877			511	Colutea	2095		
410	2*2 Chrysopbgon	2853	413	Clasia	2877			C. see Sphserosphysa	2096			
175	f. 17. Chrysoplfeium	1379	413	C. sec Bridle/a	2877			C. sec Sutherlandia	20W			
	Chrysurus	232	404	Cat'stia	2877			Colymbia	24507^A.	2820		
42)		12772	1491	Cla'menum	19524	2136		532	Comandra	741		
378		29. J6	21	Clypeola	1843			512	Comar.-p*ia	1525		
311	icer		375	Cnemidofatachys	2596	213		512	Comarum	1527		
32			15	Cnorum	106	337		Comafirca	80717	229*		
301		2131	63?	Cne.%rum	4229	4f12		513	6'OMURFIA*CEJE, Or. 72.			
318	C see £'rvum	2134	157 f	Cne.%rum	4109	1197		513	Combretum	1203		
1\$		2102	2113	C14091	1595	477		C. see Poivrea	3.147			
107	Cicuta	851	183	Cnëstis	140.3	276		515	Comespnna	1931		
S2	£-rula	880	330	Cricus see drsium	2242	54		Comley	43S			
	Cinna	600	360	C. see Leizea	24.32	145		Comifa	9351	1157		
81	C. sec Hymenodfctyon	599	360	C. see Rhapontica	2433	15		540	Commellmi	118		
426	C. eee Luculia	643	330	C. sec Scrr-tula	2239			540	CoMELINAfI Or. 200.			
3.9	Cindidotus	2972	105	C. sec fflybum	2242	114		503	CommerutvVi	926		
340	Cineraria	2340	106	C. sec /aserpttum	86(1	86		C. MCC Hyttneña	683			
3H	C. see Agath's'a	2341	474	C. sec Peuc^danum	881	15		509	Comorlacia	107		
	MTICQJU	77	474	C. sec Silaus	3328			520	CoMPAHIG, Or. 108.			
3]	C. sec Senecio	8335	64	Cotxri	390			535	Cointpbnia	2592		
	Cineraria	22085	475	CoBaffea, Or. 131.	498			Com, mi	23770	2708		
	Cinna	'''	49	Cobitphria	123			539	Conanthén*.	H80		
	L. sec Trichochloa	204	49	Coccorpactum	475			C. sec Cuimngta	&136			
	Cinnambm	1225	49	C. sec Fernclw	83			508	CondU/Yi	6GI		
213	Cinnamon	1225	159	Cgecoloba	474			518	Cond>lofarpus	33.31		
		1528	400	Cocculus	1212			Conf'ra	SHU			
		85	497	C. see WendlandVi	433			C. sec Desmi<lium	3009			
		2524	142	C. see Cochinigal Fig	1129			535	CoNIPBnJi, Or. 176.			
		2243	195	Cochingal Fig	12608	105		518	C. nium	a^2		
Jf10	C see Herac&tha	2256	258	Coghlaria	1474	105		C. Capnoph^lliim	K53			
404	c. eee Lci>ea	24.32	262	Coco-nut tree	1848	105		C. sec ffinanthe	8J3			
	CtA	2825	262	C. sec /epodium	1873	103		C. see Trachyspc.				
9	t. see xMenispérnum	2806	292	Coohospérnum	2032			mum	840			
	Cissus	400	317 j	Cock's-comb	135	106		C. sec Ulosprfmum	8(9)			
59	C see Ampelopsis	656	25	Cock's-foot	16C7	188		509	(bn)árus	1937		
82	CisTINE^i, Or. 19.		29	Cock's-foot Grass	236	90		C. BCeOmphalbbium	IJU6			
248	Cftstus	1589	200	Cockspur	12897	1506		513	ConocarpuN	71J		
3K9	Cltharexylum	1758	381	Cocoa-nut tree	26.38	473		534	Couocephalus	3414		
9^o n	19664	8187	200	Cocoa Plum	1503	474		518	Conopottiinn	3318		
320	Lfr&flus	83715	199	Cocomtilo	12855	118		532	Conosu^rinum	.312		
35.	£U'''	2187	381	Cbcos	12855	118		538	Conostylis	8970		
641	Cladfnthus	2383	382	C. sec Acrocdmia	26.38	117		C. see Lophlola	941			
	Cl-idiun	88	403	C. sec Lodoicea	2815	150		Constantia	9729	1173		
1^530	Cladoslibrium	3259	5	Codárium	34	48		Contrajerva root				
155	Cladi'istachys	725	512	Codium	3057			28.37	339			
	Cladóstephus	3035	155	Codlins and Cream	10054	1186		Contraye'rba	28J87	2468		
		U85				131		Convallaria	1038			
J7	Clary	721	170	Codon	1314			C. see Ophiopogon	1041			
	Clavseña	721	241	Codontiphora	16P9			C. sec Polygonatum	1040			
	Clkva HerculU	1324	186	Cce'li 7'osa	12009	1415		526	CoNvOLMILANeEJi, Or. 136			
453		24220	2764	371	537 Ceclbgync	2542	63		527	Convdlvulus	412	
	Clavarla	2133	240	CoMsya	15262	64		C. sec Calysfegia	494			
524	C. see Tubercularia	3234	80	Caffia	1689	62		C. sec Ipomoe'a	491			
516	Clavljrt	527	49	C. lCC Tetramerium	3H0	159		Convalvulus	10276	1210		
	ClaylñrVr	696	80	Coffee tree	627	342		521	Conyzá	2331)		
	Clearing.nut	4587	496	Cogweed tree	10384	343		C. see £r(geron	23S2			
	Cleavers	3030	160	Cogweld	1226	364		C. sea Psiidia	2454			
		348	377	Clix	2603	170		C. see Solid.fo	23S9			
254	C. see Jfnefone	1631	392	Cola	S3875	2724		504	Cookia	I>27		
374	C. sec Jfffragenc	1630	225	Colbe*rtfo	1608	168		518	Copai/cra	1288		
217		22858	141	Colchicum	1119	451		Coprinus	3119			
		131	141	C. sec nulbocodium	1031	238		Cóptis	1642			
			51	Csec Mercnbta	1120	317		Coralloidendrum				
			13	Cold^nh	413	370		5C6	Cornillorrliza	8520		
			287	CoUanthus	817w.	317		Corai tree	19518	2162		
			472	Colchro6kca	1750	121		Coranic Poison-				
			483	Colconéma	3'03	Coron		bulb	7879	97		
			529	d leus	3383	503						
			1695	Ca/za? 21882 m.	<23>16	214		Ctrchonts	1577			
			356	Calléma	2.353	527		C. see Kerr/Vi	1531			
			1752	444	2673	69		Crt!Vr	563			
			1347	508	23699	67		C. sec Patacónula	SG*			
			141	ColletiVi	667	70		C. see Varron/a	564			
			472	ColifnsiVi	1804	527		Connulcrte, Or. 138.				
			11	Collins'mia	77	539		539 Connlylnc	1078			
			59	ColK.mia	47.3	3V5		535 Corbina	2737			
			386	Colocasia	2.353	463		Coremum	3244			
			389	Coloc'nths	23696	358		522 Coreopsis	2415			
			302	511 Colognfh	2075	359		C. see Actinõraeris	2417			
			S89	Colquhuitelle	23688 v.	366		C. sec Bldens	2284			
			412	Eblion	2698	359		C. sec Calli6psis	£'418			
			111	Colpoon tree	7221	£58		C. sec Cosmea	2414			
			344	Col'ts-foot	2333	488		C. sec Kchinacea	3408			
			478	508 Culibrina	3300	362		C. sec Alphium	2440			
			285	Colubrlnus	5496	359		C. see Simsia	2419			
			1488	Columbine	1605	351		C. sec Ximenes/fl	2.3.<			
				497 Couiniboo-root		101		Coriander	817			

Lin.	Nat*	Sgt.	Hen.	Liu.	Nat.	Sp.	<<n.	Lin.	Nil.
101	518 Corlandrum	817		113	C. ice Curtdyne	918	399	Cucffera	24275 sy. 2779
101	C. see ISifurls	818		113	C. ice Globula	917	1862	Cuckoo	f 14018 1415
401	508 Cortina	2797		160	C. ice Kalanchuc	121t)	256 J	F l o W e r {	gg
	508 CORIAHIEB, Or. 58.			112	C. ice Kalosunthes	914	179	502 Cucobalui	
159	Conndum	10340	1217	112	C. ice J.arbchc/i	915	179	C. we Silene	
58	530 Corp	4fil		113	C. bw TurjehHia	916*	294	Cucullaria	18205
4	531 Compcrmura	27		113	C. we VauanthM	919	398	Cucumber	jS
384	Cork tree	8J422 206<1		■■■ CSUMULAY-EJi, Or. ■■■			389	515 Cucumii	>Si
826	Corkwood	143(Ki 1622		200	513 Crata^gui	1506	381	C. We Liffii	JSS
JHj	(orn.cockle	12019 141t		202	C. see I^ottnia	1509	388	CwqTnchosantheffJg	
50	Cornelian Cherry			201	C. ice /*yru	1507	389	£15 Cucurbita	9S
		3380	401	188	500 Cratas'va	1440	411	C. see Benincisa	gg
17	Corn Flag	1*8		815	C. see Niebfhrfa	1540	389	C. we Cucupii	*Sl
444	Cornlrfaria	3109		188	C. ice Hichiea	1441	389	C. we l^igciaria	JJ»
852	Cornish Moneywort			460	Craferium	3201	515 CUCURBITA'CEA, Or. v^		
		15982	1792	26	Creeping Fiorin	1712	442	Cu^ear il^BSR #E	
				193	CREPITIANA	12465	248	Cvjite	15745
22	548 Cornucopia	16)					336	521 Cilrftiuin	
12	Cornucopia*	793 8i						Cillaban	10379
50	512 CV.riis	401		327	521 Crepis	2215	359	522 Culliimto	
8	589 Corndlfia	52		327	C. we Antlyala	2819	104J	Cumin	
8	C. we H6sU	52		QUA	C. we lorkfiaOH/a	2214	104	104 518 Cumlnum	
307	Cornweed	18980 2104		QxO	C. ice Ilapalobtc.		475	539 Cummfnfia	
	523 COROLLIFLOR* SHDC. 3.			326	phium	8812	10	528 Cunlla	
307	511 Coronilld	210.1		386	C. ice Laguierii	8213	387	535 Cunningham/a	
£08	C. sec Securfgera	2110		322	C. we Picrdium	8802	175	517 Cunhflvi	
				324	Cieei*Icrii	8810	517 CvNotuuccar, Or. 97.		
262	j Coronoput	J 1 6 W 2 1 872		328	C. we Kodlgo	8225	145	505 Cupania	
144	507 CorreVi	1154		327	C. ice TolpiM	2218	189	514 Cupha	
111	516 Corrigiola	905		528	Creventia	1765	388	535 Cupreuu	
57	530 CorfiIM	452		248	C. we TansVium	176C	388	C. sec Schubert/fl	
350	521 Corvihirtfa	23UI		862	Cress	16539 1873	377	Curagua	
12	543 Coryfirpus	91		98	527 Crebia	800	821	496 Curatella	
366	536 Corycium	8476			Cress Rocket	1888	391	Circat	83848
89*	498 Corydalii	2047		265	Cretan Carrot	6711	98	527 Ctscuta	
894	C. ice Adlumla	2050		103	Cribaria	3207	99	518	
894	C. sec Dlctytra	2049		120	538 Crlnum	964	886	Custard Apple	
384	535 Crifyhui	Mil		117	C. we Eurycole*	937	883	511 Cyam6psii	
433	Coryncphora	3021		167	Crilita	10814 1275	49	508 Curtfsia	
87	543 Corynphorui	218					113	516 Curtdyne	
464	Coryneum	Siii'A		248	502 Criitarla	2020	159	Curura	10312:
84	524 Corynoc&rus	609		290	CYithmum	834	98		
185	540 Curypha	996		103	Critmlra		99	518	
413	C. we ChanWVop*	2883		334			886	Cyathaea	
141	C. tec Siabal	1124					422	C. sec Aspfdlum	
185	C. lee Taiirira	997		362	Crocotyl-ium	82225 2434	65	523 Cyathbdci	
369	536 Corysanthe*	2519		15	537 rrbeus	116	423	Cjathui	27708 sv.
400	497 Cosc(nlum	2789		302	C. ice Neurocarpum	WT15	535		
358	522 Cdsmea	2414		278	C. ice lkfnia	1947	403	535 cycai	^25
65	523 Cosmelia	517		278	C. ice Xmuixi	1948	535	Cyclamen	J#
159	505 Colignia	1093		46	Crosswort	356	475	539 Cyclanthui	S«
338	CIMtnary	2J10		411	Croisiwort	24971 2859	162	511 Cyclobdthra	39g
3	537 CFwtui	11		278	511 Crotalaria	19A2	202	513 Cyf.pia	J*5
355	Cota	21761 2584		279	C. ice Hypocaly*ptUB	1953	466	538 Cydnbia	JgJ
110	C6tinui	7145 894		302	C. ice Neurocarpum	WT15	317	512 Cylindrosporium	**L
802	513 Cotoneaster	1513		278	C. ice lkfnia	1947	140	Cylitta	JJS
890	Cotton	2016		278	C. ice R6tlera	2810	849	Cymation	V JIg
21	Cotton Gran	161		402	C. we Crozophora	3417	853	Cymbria	15783
340	Cotton Roue	J&I7		490	C. we Crozophora	3417	370	Cymbbaria	
331	Cotton Thistle	2245		395	Crowberry	27XJ	370	Cymbia	
354	588 C6tula	8380		169	507 Crdwa	1310	371	C. ice	
354	C. we Grange*	8381		230	495 Crowfoot	1637	372		
354	C. ice Lidbeckfl	8378		495			410	543 C. ice Vdnda	Jfa
354	Citula	21757 8384		498	CRI: CIFER®, Or. 13.		155	5W Cymboubgon	Jjj
183	516 Cotyledon	1409		361	Cryptoa	22167 2434	104	525 Cyninsuma	Jaji
160	C. sec Bryophyllum	1220		25	Crow's.fnot	1666 188	94	525 Cymlnum	6754 gi
478	C see Eclicvria	3356		490	533 Craxphora	3417	95	Cynanrhnum	JfA
478	C. ice PistorInla	3355		46	520 Crucinella	356	94	C. jtfclopia*	k#
478	C. ice Umbillfcui	3354		411	Cruciata	84971 2859	95	C. we Dafnria	ifis
34	Couch Grass	2299	870	498	CRI: CIFER®, Or. 13.		93	C. we Gonol	
167	512 Coulterfo	1277		361	Cryptoa	22167 2434	332	521 Cynapium	
165	Courbard	10683 1269		200	Crus.gilli	12897 1506	383	Cynara	-->
389	CourtgotrdazMTHv.	2695		893	Crux A'droe	18176 2044	34	542 Cynoc&mbc	23376 ^5
75	519 Cautaria	603					54	527 Cynodon	455
48	526 Coutoibta	367		27	542 CryptiN	211	54	6'ynogldssum	tfl
188	Cdwa	12107 1429		90	508 Cryptandra	711	54	C. sec Omphaludci	^
105	Cowbane	851		372	5.'A) Cryptarrhena	2556	166	512 C. we Rlndem	Wfi
156	Cowberry	10123 1194		161	532 Cryptocarya	1229	90	Cynometra	719
387	CowdlePine	23574 2684		417	545 C. FiucEH, Or. 1.		89	543 Cyndbatl	B9& &
899	Cow Grass	18478 2066		417	543 C. FiucEH, Or. 1.		32	Cynoidrui	ofO
177	Cowherb	11368 1355		424	544 C. IQJIIHETAVEJE,		29	C. see Beckm&mfif	Z&
316	Cow Itch	21.09		439	541 C. LvcoopoofNEJJ, Or. 3.		33	C. we Chryiurui	*
108	Cow Parsley	7059 884		425	544 C. MARHiLrdcc*, Or. 4.		33	C. ice Dactyloct-	ggi
108	Cow Panncp	884		425	544 C. My HCl, Or. 5.		33	mium	Qb9
57	Cowslip	3773 451		431	544 C. HKPA'TKJE, Or. &		21)	C. ice RhåMochloa	gj
460	Cow tree	.3420		432	544 C. ALGK, Or. 7.			C. ice Sclicer/n	
£46	Cow Wheat	1741		439	545 C. AICIBVES, Or. 8.				
911	Crdcca	19206 8133		445	545 C. FV'NOI, Or. 9.				
865	499 Crfcnbc	1896		445	545 Crypt8lcpii	531			
865	C see SobolewakW	1880		67	Crypt8lomycci	3149			
155	Cranberry	1193		455	Crypt8lomycci	329			
876	Cranc'i Bill	1932		41	520 Cryptosplrmum	750			
244	686 Craniolaria	1721		93	&25 Cryptobtesla	2428			
*?7	621 Craptldia	.1401		359	Cryptost<!mnia	2498			
118	516 Crtwula	§15		338	53d Crypt6stylis	2498			

GENERAL INDEX.

555

Lin.	N.M.	8p.	Gen.	Lin.	Nat.	*P.	«•	Lin.	Nat.	fiy	U<<
192	Cyparissiai	12142	14M	162	D. MONCKJY'NIA, Or. L			145	505 Dimocarpui	9358	
268	537 Cypella	1912		175	D. DIUV'NIA, Or. 2.				U< M & J ty.	1159	
	CH CYP Mi\K,M, Or. 209.			179	I). TRIOV'NIA, Or. A			318	Dlm6rphtt	10585 ty.	2173
22	541 Cyperuf	16)		183	D. PBNTAOY'NIA, Or. 4.			34	542 Dtnbra		265
22	C. sec Papyrui	164		187	D. DKCAHY'NIA, Or. 5.			410	D. sec Atheroiibgon	2850	
ft)	522 Cephia	714		388	Deciduous C y , ^ ^			44	D. sec Wangeu.		
3)8	Cypress	2692							hemm ,		266
192	Cyriess	12442	14H	188	515 Decumaria	113)		64	527 DJLMUI		4&7
373	537 Cypripedium	2580		92	530 Dceriwgia	7W		316	512 Dirclea		2157
o7	523 Cyrtillii	695		6ti	Dégraphis	217		46	51D Dibilia		353
852	C sec Trevirana	1796		27	D'tleuerMi	306)			DIKYIA, (1.22.		
121	538 CyrUnthus	965		438	496 Della	1583		393	1). MO.NA'NIRIA, Or. 1.		
121	C. sec Gastronema	966		222	496 yjclphnium	LOW		393	I) DIA'NDUIA, Or. 2.		
121	536 Cyrtobodium	2541		464	Dembitium	3^54		395	IX TRIA'NUUIA, Or. 3.		
&H	&H Cyrtotylis	2513		537	Dendrbium	9547		396	D 'IVIRA'NIIIIA, Or. 4.		
%	CyrticApnos	204H		371	1). see Cainarfodium	254K		397	D. PKNI-A'NIRIA, Or. 5.		
	3082			371	D. sec Diphdium	W2H		398	D. HKXA NIRIA, Or. 6.		
	Cystmelra			370	D. sec KVia	i5.09		401	D. Oi IA'NDIIA, Or. 7.		
	533 CIT(NJK), Or. 166.			378	D. sec Polystachya	257.0		401	D. KNNI'A'NDUIA, Or. 8.		
itjl	Cytfspora	3171		372	D. sec Xylbium	i5.56		401	D. DKCA NIRIA, Or. 9.		
281	511 C'tisus	1914		370	Di's Chnis	85J6		402	1). DoDucVNURIA.Or.IO.		
C	C. sue Adenocarpus	1965		131	491) Dent'ria	JK Jl		402	D. lrtMA'MiRA, Or. 11.		
2P	C. see Cluanus	2160		£56	519 Dentiflta	H^i		402	1) POM A'NDIA, Or. 12.		
2P	G see Ca)6phaca	2094		74	Deodina	2XKJ7		403	IX Mu.NAIK'LPIIA,		
J77	C. sec Goodia	1944		388	Dechamia	2-21			Or. 13.		
315	C. sec Illychnbhsla	2145		28	DechamiaM/a	2-21		357	522 DiomMia	94J7	
16J	C. see Therm6p*is	1241		406	Dckinlhthus	28:>i		170	501 Dioiu>%A	I.»1	
S<	533) Czack/a	1062		433	Demiidium	MW		400	5.K DiiNnWa	2787	
3HH	535 Darr'dium	2691		91	1) csmorhala	724		538 DmiHoRi^, Or. 192.			
JSJ	Dacrymyccs	3152		91	I). see Cladustachys	725					
1H	Dactylanthes 12285			309	All Dminudium	2120		85	507 Di6>ina	673	
	& 12267 sy.	1460		382	Dsmeönicus	2fUi		472 j	D. nee Adcmundra	675	
FJ	543 Aactylis	236		469	Devil's Bit	3288		85	D. sec Agath6ima	677	
	D. see Rostraria	242		191	Devil's Milk	12158		472	D. sec Audoufnirt	3308	
52	542 Ixictyloctenium	261		212	Dewberry	1350H	1523	85	D. see Harrysina	676	
94	523 Daphalea	3124		293	DIADK'LPHIA, Cl. 17.			472	I). see Culfonicina	3303	
117	523 D'mia	759		293	D. PUNTA'NDRIA, Or. 1.			85	D. Me Dichoama	674	
117	Daffodil	7599 IM3		294	D. HXXA'NDHIA, Or. 2.			99	D. Ki'o Liticunia	803	
351	522 Dahlia	2363		295	D. OCTA'NDHIA, Or. 3.			414	524 /Jiospyro*	28h9	
488	D. ice Georgtaa	3406		296	D. DKCA'NDMIA, Or. 4.			379	531 Di'tis	2t18	
51	DahoOn	3425	411	5	DIA'NDRIA, Cl. 2.			338	D. sec Otinthus	2KNi	
531	532 Dals	13f6		5	D. MONOU'NIA, Or. 1.			308	511 Diphara	2111	
317	Daisy	Weil	12	5	D. DIUY'NIA, Or. 2.			138	497 Diplillycia	1JB8	
317	512 Dalbergia	2Ui6		13	I). TRIUY'NIA, Or. 3.			305	511 Diphysa	2093	
	UbccDiphaca	2111		137	539 Dia<>7a	1071		426	DipMscium	2966	
	D. see Lonchocar-			177	502 DiRiUhns	1381		31	543 Diplachnc	246	
)>us	2085		58	y^apciiaia	45!		23	541 Diplasla	175	
	511 IKlea	1971		252	Diftsia	1802		420	544 DipKixum	2925	
	D. see Petaloste.			433	DIAToma	SIM16		351	521 Diplocoma	2354	
333	mum	1922		188	503 Dfrera	1436		379	531 Di'tis	2t18	
JOJ	Dalca	20523	22Pt	310	511 Dirrrma	2121		338	D. sec Otinthus	2KNi	
212	512 Dalech/impia	2727		279	511 Dichilus	1957		308	511 Diphara	2111	
	Dahb&rda	1524		4L0	4L0 DICMILAMY'I)KilS,Subd.			138	497 Diplillycia	1JB8	
	D. see Comarunsls	1525		407	Dirhuidra	799		305	511 Diphysa	2093	
*14	508 DalmCmpica	LHWi		404	Dichorizifindra	1004		426	DipMscium	2966	
	DdlUnifVi	2)H8		507	DirlriHma	674		31	543 Diplachnc	246	
H	536 Damas.nium	1130		461	Dichnsp6rium	3216		23	541 Diplasla	175	
	D. sec Actiioearims	11>1		544	Dickbbiwi			420	544 DipKixum	2925	
	535 Minmra	26H4		422	DickbbiwiVi			351	521 Diplocoma	2354	
	Dammar Pine	i(iK4		422	1). HOC HalAntium	9&f1		379	531 Diptcrlx	8174	
.022	Dampitira	617		422	I). sit' Clbbtum	2936		41	520 Jfpsacus	344	
5-14	Danu,a%	2953		521	Dicliptera	59		157	520 Jfptcrlx	8174	
	Daitling Girls	194	16	294	Diclytra	2049		157	&2 Dfrcra	1198	
	Dandelion	2207		495	DICOTYL'DONEB, Cl. 1.			0J6	D)s	2473	
	Danthbnia	226		427	Dferanum	2978		494	51(i DiM'pmma	33H9	
	532	1197		428	D. see Didy*modon	2)K2		360	53> DiHRTs	2475	
	532	1197		428	D. see Grimmia	2)175		131	539 DIHporum	103J	
	532	1197		428	D. sec Ilicidodon	2981		426	5f8 Dodfrtfo	17)0	
	Darfa 2527 & 8^.	2922		428	D. sec Trich6sto.			69	525 Dissolirna	552	
	512 Darlingtiiia	743			mum	2979		455	Ditiola	3147	
	Darnel	2328	271		507 7)ictfmnu	1:08		238	Dittany of Crete		
	515 Darwfn/a	1372			JJictfmnu	15132			15132 1H80		
	Date Palm	2745		238	JJictfmnu	1681		368	536 Dior is	2493	
	Date Plum	2881)		4(i	Dict^ndium	3208		141	Dock	1126	
	500 Ihitiixa	2805		522 Didi'lta	2425		252	58 Dodfrtfo			
	Or. 15.			499 DicdHmis	IHW				500 Dodder	798	
	627	481		474 Didsctis	28144jry.	W7		187	Doii'M'A'Nmia, CL 11.		
	542 DriiKm&nfa	482		4W	D^mum	3VX>		187	D. MONIKY'NIA, Or. 1.		
	518	2090		468	Didymocfirpus	3279		190	D. DKJY'NIA, Or. 2.		
		825		428	Didymixlon	2982		190	D. T'nna, NIA, Or. 3.		
	X). sec Platyspr-			232	D>YNA*MIA, Cl. 14.			193	D. Ti TRAtv'NIA, Or. 4.		
103	mmum	S332		232	D.GYMNOSPK'IMIA, Or. 1			133	D. PI.NTALY NIA, Or. 5.		
	D. see Trachyspr-			241	D. ANGIOSPB'UMU, Or. 2.			14K	D. IIIAAUY NIA, Or. 6.		
				80	519 Dicryfla	624		19,J	1). DoDi-i-AiiY'NIA, Or. 7.		
	5*4 DavMlia	2933		251	528 Digitalis	17N		57	530 iJnilCMIcon	451	
	r.n JJ^w^&v	10505 sy.	1247	251	I), sec iHop^xii	178H		104	505 DodoiM-Vi	1179	
	°H pavitbia	1263		24	543 Dlgit^rla	IK		106	Dogliramble	5K9 719	
	p. sec. Jacksbsnf0	1253		18	538 Dilatris	139		68	Dog Poisen	ti9'J4 861	
	pay Lily	1007		105	Dill^nwi	858		68	DoK's.banc	542	
71	Deadly Carrot	845		226	D. sec ColWrtja	1614		383	Dog'u-hune	14179 II>00	
	Deadly Nightshade	225		225	D. sec Tetr&cra	Km		29	Dog's cabbage	25376 26.08	
	4640	584		225	D. sec W6rmia	1610		131	Dog's-tail Grass	234	
	jjed Nettle	1667		164	511 Dillwfinfo	1257		50	Dog's-tooth Violet	1028	
	r, w {^"tflla	2523^.	S06	164	D. sec Eutaxia	1258		512 Dogwood		401	
	ii,j I)*^10*10,,	3348						512 iJ61choa		21&	
	in2 DECA'Nnau, CL 10.							512 D. see Canavulia		I'J64	

GENERAL INDEX.

tin.	Nat.	Sp.	Gen.	Lin.	Nat.	6p.	Gen.	Lin.	Nat.	Sp.	&".
283	D. see <i>Cyamopsis</i>	1969		281	Dyer's green weed	70			524 <i>Emhllia</i>	378	
316	D. sec <i>Diuelco</i>	2157			^	17503	1963	391	<i>E mbhea</i>	8710	
316	J. see <i>Ldlab</i>	2154		190	Dyer's little yellow	469		532 <i>Embdthrium</i>	3287		
316	D. see <i>Muchna</i>	2155		352	weed.	414		Embrydpterus	25183 28J9		
316	X). see <i>Pachyrhlzus</i>	2155			<i>Dyssbiua</i>	307		E'merous	lb>87 21\$		
315	D. see <i>Phaskolu</i> *	2149			XII><J3& 21594 sy.	142		EHX	1127		
316	D. see <i>Psophocarpus</i>	2158		363	Dyssbdium	531		EMPB'THEML	Or. 177. m9R		
315	D. see <i>Ilhynchbsia</i>	2145		528	Dyssophytu	395		JS,mpetruni	2737		
316	D. sec <i>Sija</i>	2151		31HJ	Earth Nut	395		E. see <i>Corema</i>	219		
316	D.socVigna	2153		313	Earth Pea	379		507 <i>Empleiirum</i>	1999		
315	J) sec <i>Wistarta</i>	2147		454	Earth Tongue	26		Enarthrocirpus	1999		
219	496 <i>Doliocarpus</i>	1584		339	East India Mug.	427		Kncaly'pta	2974		
285	503 <i>Dombcy/a</i>	1994			wort	428		E. sec <i>Glyphoml-</i>			
844	D.sectourrett'a	1719			20836	SP60		trion			
29	Dbnax	1909	2:10		2313			522 <i>Encelia</i>			
103	518 D6i.dia	839		414	£BE\A:E£, Or.	122		Enchanter's Night-			
350	Dbnia 21538 21541 &			414	Ebcmaster	12		shade			
	21545 sy. 3347			414	£'benum	439		Encu%lium	5075		
419	544 Doddfo	2920		284	£'uenus	349		536 <i>Encyclia</i>	3413		
344	Dòria	21150 2335		285	A'btifus	489		Endive			
351	521 <i>Durdnicum</i>	2353		318	Ecasaphyllum	328		Endivia	20219 «g		
351	D. sec <i>Diplocoma</i>	2334		8	Ecasfium	318		Endocarpon			
344	J). see <i>Wcrier/a</i>	2334		243	Eccremocarpus	441		EnglibhSlercury			
351	i			483	E. sec Calampelis	(468)					
42	534 Dorstenra			478	Echcvèria	33H5					
123	538 Doryanthes	976		468	Echinacca			533 <i>Enkianthus</i>	1***		
300	511 <i>DorScnium</i>	2067		34	Echinaria			ENM:A'URIA, Cl. 9.			
63	/>orjcnium	4233	4:2	433	Echinilla			160	E. MONOGVNIA, Or. 1*		
458	Dothidea	3159		479	Echinocactus			161	E. TRIGY'NIA, Or. £		
	502 <i>Douradin/ia</i>			479	E. sec Melocactus			162	E. HEXAOYMA, Or. 3...		
876	Dove's-foot	17249	1932	543	Echindchloa			409	512 <i>Entiida</i>	*&	
858	499 <i>Draba</i>	1846		468	Echinophora			818	503 Entelea	1^	
262	Jrâba	16534	1873	25	Echinopbgon			523 EpAcuDEC, Or.	117. ^		
£57	D. sec <i>Aubrittia</i>	1837		,165	fch'.nojjo			64	E'pacris	SI	
858	J). see <i>Erdplilia</i>	1847		5	Echmos^rmum			65	E. see Lysinema	j>\	
858	D. sec <i>Petrocallis</i>	1445		67	AuMtes			404	535 £'phedra	S?2	
1:0	539 Dracaena	1020		68	E. sec Haemadictyon			58	phemerum	3828 fl	
130	D. sec <i>Charwlwd/a</i>	1019			E. sec Parsons/a			145	Ephiehs	9367^. J'S	
137	I). secDianrf/a	1071			E. see Stroph&nthus			372	536 Epilendrum	#£	
130	I). sec <i>Phylluina</i>	1021		527	Echium			370	E. sec Braswa	S	
130	Drâco	8459	1020	522	Eclipta			371	E. see Oncidium	*31	
839	528 Dratocephialum	1685		511	Ectostriuma			369	E. see Cephalan-		
483	D. sec <i>Physostegia</i>	33S)		511	Edwardsia			thera	2508		
143	541 JDracutium	1140			Egg-bearer			369	E. see Eriochillus		
143	J). sec Calla	1141			4808	591		103	spipactis		
48	I). see <i>Symplocarpus</i>	332						195	517 Epiphyllum		
386	7Jardnium	83510 8673			13415	15*2		98	Apithymum	6409 7»	
65	523 Dracophyllum	512			Egyptian				£44 /UJIKKTA;KJ		
471	D. see Sphem.toma	3294			Lotus			484	544 fquisetum		
386	Draefinculus	23508 2673			13809	1556		33	543 Kragrdstis	256	
143	Dragon	1140			Egyptian Thorn			33	Eragrbtis		
386	Dragon	23508	2673		84796	8837		9	529 Eranthemum		
239	Dragon's Head	166>						244	E. see Uarsterfe	lrfig	
130	Dragon tree	8459	1010	527	Ehret/rt			231	496 Era nth is	ios	
42	DrakVna	2838	339	527	Pi we Bourrera			135	539 Eremurus	#g	
434	Draparnald/a	80W		527	Ehrhart*			461	Ergot		
318	512 Drcpanorâniis	2167		528	E. sec Microlae^na			372	537 E'ria	2559	
473	518 Drcpanophyllum	3318		506	Eckberg/Vz			28	543 EriAchne	92*	
137	539 Drifnia	1077		558	ELAA'UNKM, Or.			36	543 EriAnthus	285	
824	496 Drimys	1602		164.	AteAgnus			410	E. sec Kipfdium		
70	Drinker's Nut	4587	574		538	Elw'is		146	583 2'rca		
114	501 Drôsera	922			540	E. sec Fries/a		146	E. sec Menziesra	111*	
35	501 DKOSERA%CE£, Or.	21.			540	Eleagnus		583	ERICEJE, Or.	115.	
	501 Drunken Daniel				541	Elecampane		593	E. see 7'uua		
					541	Elat. tis		581	Engerion		
					541	Elett-haris		343	E. see Enncum		
					541	Elephantopus		350	E. see Enncum		
					542	Elephant's-foot		202	513 Eriobdrya		
					542	Eleusine		104	Eriocalia		
					542	Eleuteria			541 Erioradlon		
					83810	2715		36	542 Erioc'phalus		
					543	Elliottm		364	Eriochillus		
					543	Ellisia		536	Eridroma		
					544	Elini		595	Endodendron		
					544	Eliogryphus		161	531 Krigdonum		
					545	Elytrria		291	503 Eriof'na		
					545	Elytrtria		334	Eriopippus		
					546	Elephant's-foot		21	541 Eridphorum	M 105	
					546	Elephant's-foot			R see Trichdphorum	ig	
					547	Elephant's-foot		468	522 Eriophy-luin		
					547	Elephant's-foot		539 Eriopbpermum	jo'ij		
					548	Elshdtrza		169	507 Eriostemon	7>\$	
					548	Elymus		188	E. sec Dfcrra	1/1\$	
					549	Elyna		277	E. sec l'hiotheca	£g	
					549	Elytrria		237	E. sec Sthch}•	\$0	
					549	Elytrtria		1655	l'risJthalcs	20390 Sg	
					549	Elytrtria		331	519 frfthahs	qSt	
					549	Elytrtria		160	E-rium	7818 V-	
					549	Elytrtria		66	Ernodea	£3	
					49	520				^	

GENERAL INDEX.

557

** Nat	6p. Gen.	Lin.	Nat.	8p ^{ss}	Gen.	Lin.	Nat.	Sp. Gen.
\$0 506 Erf-dium	1927	48	<i>E. sec Microcale</i>	368		21	541 Fimbrifstylls	155
*g. 439 E.6phila	1847	48	<i>E. see Sebe'a</i>	70	24		Finger Grasi	185
501 Eruption	702	404	533 Exoccaria	28-6	26		Fiorin	1712 203
g£ 499 £racq	1887	458	Excfpula	3168	388		Fir	2686
2te A^ Erucago	1609° 19(K)	456	Exfdia	3151	263		Fish Poison	16553 1873
+g 499 Erucana	1901	432	Exiliria	3005	167		Fistula	JOT99 1272
312 JSI, £ villa	19273	380	532 Exocfpros	452			Fistulina	3127
		495	ExoduENJi, Cl. 1.	402	500		FlacourtfVi	2811
2g 511 f'rvum	2134	464	Exosbrium	500			FLAcojIRTitinor, Or.	17.
lf* E. see Galactia	2076	75	519 Exo-stmrna	541			Flagclflrla	1105
A see A cia	2133	249	Eye-bright	277			Flat Pea	Iphi
518 tfringium	822	311	511 fviba	2132	365		Flavéria	2162
SRI Eryngio	822	169	Fabtgo	10886	113		Flax	921
499 £rynum	1867	196	F. see Leptospér.	342			Flax Lily	1084
R see Alliaria	186ri	50	507 Fagara	398			Flax Star	3841 457
502 Ab. see Barbarea	1822	155	F. sec Ephyrum	311			Fcabane	2380
27. sec Nof'iccras	1821	49	F. see Evbdia	502			Fleawort	[21512 2345]
37 fS E. sec Symbrium	1H&	49	F. see Xanthoxy-	311			Flemngta	2127
512 Erythrina	3226	398	lum	350			Flindlrwf	1299
		467	276<	261			Flix.weed	164G3 1865
55? f. 2162	f12432 1460	315	512 Fagilla	2146	186		Flos Cuculi	12018 1415
-1 ro, « i 176 W 1975	159	169	507 Fagbnfo	1305	186		Flos Jbvi	12007 1415
31 g,j Erythrote^na	2244	70	515 Fagopyrum	10280	123		Flos MRthni	8042 979
Erythronium	1028	384	520 Fagus	1481	167		Flower Fence	1274
S? Erythrophilum	12K5	38*	F. see Castanca	2659	12		Flowering Ash	83
100 30° ERYTHROXY'LLE, Or. 42.	104	104	FalcXria	6798	162		Flowering Rush	12J4
505 Erythr'xylon	1400	98	527 F.ilkfh	848	186		Flower of Jove	
E. see Sethia	1401	177	False Armeria	1383	12007		1415	
511 1- Erythroxylon	16S48	125	Fan Palm	996	2769		FLUVUEB, Or.	905.
86 S 7 Escallbnii	687	413	Fan Palm	25112	541		Fly	5225 622
4R9 A/J EacALLiiNi< r. Or. 95.	257	344	4ittrfra	2883	105		/Vnfeulum	6817 858
<* 498 Eschsch61ZM	3370	257	499 Fanetia	1835	297		fw num grc % cum	18387 20(13
uS J^sculus	23456 2666	257	F. sec AubrietVi	1837			Footdia	1496
3g E'sula	12430 1460	412	F. sec Bcrterba	1836	199		FOLIA^CE£ ^	
1& ?? * * *	2292	397	Fatrafra	9M& 2864	543		Fontanfaia	80
IS 5J Eucalyptus	1493	25	Feather Grass	24188 2751	525		Fontinalis	2989
24ft r1 Euchillus	1261	12	Fidia	86	426		F. see Cnu-lidötus	S972
JJS l2 Euchrbma	1768	97	Felwort	793	106		Fool's Parsley	869
533 Euclea	2804	420	Female Fern	2>465 2921	374		Fonkblivi	2584
499 EucKdiura	1856	105	Fennel	6847 858	221		Fothergillrt	1593
117 IS^ Eucomis	1012	225	Fennel Flower	1606	174		F. sec Chitonia	1358
31J £38 Eurbisia	936	297	Fenugreek	18387 206J	223		Foxbane	14178 1G00
197 ? ! Eudesmia	2184	373	Fernandesia	2577	551		Foxglove	17H9
j^ 515 EugenTi	1487	49	Fernel/a	2>65	27		Fox-tail Grass	213
E. see Calypttran-		170	Fernbia	1325	212		Fragaria	1526
athes		267	Ferrari*	1909	512		Fragillaria	3107
E. see Jambbsa	1490	267	F. sec Tigridia	1910	484		Franciscca	3387
4H E. see Stravodium	2040	20	F. sec ViesscfxiVi	147	528		Frangula	5409 657
372 *i ^{ss} £^*ety*& '»»»	3365	400	Ferrarlo/a	16756^ 1909	139		Frankenin	1009
536 Euphelia	2568	107	Ferrtola	24J78< y. 2791	501		YRANHEmdcte, Or.	25.
4J Eunbmia	1875	104	Ferula	880	387		Frankincense	23593 2685
1SJ x ^{ss} £u6nymus	666	474	F sec Aubon	842	380		Franzéria	2624
33S? Eim*ftr.V. f 12221	14C8	108	F. sec Ferulago	3326	48		Fräsem	372
a1 ^ t ^ 120597	2273	107	F. sec Pastinaca	8H3	169		Fraxint'lla	10908 1308
je? 521 £upati>rtim	2268	474	F. sec Sellnuia	873	414		rtaxinus	2887
«S £. see CscsIlna	2272	107	Foralua	3326	W		F. see O'rヌus	83
3?1 E. see Ozothamnus	2^95	30	F. sec Stellnbi	6991	&£		French Marigold	21596 2365
i^ Eupatbr/ttm	21838	30	Fescue Grass	880	387		Frexitm	1561
533 /?uph-/rbm	1460	543	Fathca	238	386		Friar's Cowl	2674
3^ 3. gce Pedilanthus	1461	31	F. sec Drachypodium	243	503		Fries/a	1437
533 A'UPiiORBMco, Or.	167.	31	F. sec frbmus	241	503		Fringe tree	38
50 > Euphoria	1159	31	F. sec Ceratichloa	247	5		Fritillfria	1018
628 Euphrasia	1773	31	F. sec Corycarpus	91	129		F. see Cyclorfthra	3337
49 G Eummattia	1538	29	F. sec Diplachne	246	129		Fntillary	1018
JS Eurbtium	3240	SO	F. sec Myjalurus	240	129		Frog-bit	2795
53 Eurya	2875	30	F. sec Stihedonbrus	239	366		Frog Orchis	£2455 2480
132 Euryx	1564	S2	F. see Schismus	248	187		Frutescent Celan-	
5/0	1933	33	F. see Scler6chloa	257	dine		dine	12092 1432
39 Ei E. xrys	3259	404	FcuUea	24572 sy. 2832	514		Fuchsia	1188
	2643	353	Feverfew	2375	139		F. see Schradfcra	1096
522	616	80	Feverwort	(25	439		JParus	3C81
	2296	461	Fibrillaria	27828 sy. 3257	22		Fuirena	171
	32P2	2J0	Ficaria	1636	498		J'umaria	2052
	248	415	FicoJOEJE, Or.	516 294	294		F. sec CorydMis	2047
	243	534	Fleus	£900	294		F. see Cysticapnos	2048
	45	Fieldia	1709	294	294		F. sec Dfelytra	2049
	203	Fig Marigold	351	294	294		F. sec Sarcocapnos	2051
	205	Fig tree	1520	294	498		FVMATH^CZM, Or.	12.
	251	Figwort	2900	428			Fumitory	2052
	340	521 Filago	1790	427			Funaria	2183
	340	F. see Gnaphalium	2317	545			Furze	
	340	F. see Gnaphalium	2314	545			Funkra	1008
	3b4	Filbert	23388 2663	539			Furcellaria	S
	543	FILICES, Or. 1.	2943	545			Furce'a	
				545			Furze	
				545			Fusiumu	
				545			4x52 Fusarium	3035
				463			Fuidium	3252
				470			Fustick-wood	23179

m

GENERAL INDE5

Un.	Nx.	ts>	«».	Uo.	It*	*>	On.	Un.	KM.		
170	505	<i>CAttntta</i>		ISSJ	BS	Gennamler	1048	BH	Goat's Beard	OB	
134	MS			10M	BB	tuTii.Tti Mndwort	443	908	Goil's Beard	13038 I	
MB		<i>Grtgwitjw</i>		S81	521	GvropA'oti		BB	Goat's Foot	11UG4 1414	
302	52A	<i>GaTaclia</i>		SLJTR	533	Gestvertu		S39	Goat's Origanum		
302	G. MO	Baxhierwi	9078		441	G. MX Codondhnhra	10BB			15161 If*	
BBS	531	<i>OakwUUi</i>		MSB	941	G. see Pentar&phia	(700)	BBS	Gout'* limp'		
490	BM	<i>Galactodfodron</i>	MDO	SES	<i>OexKBUiac</i> , Or, 113.			307	Goat's Thom	189S6 ^	
S	1		C 98	1S3	Clch Jills	977			Goat's Wheat	I*, U	
3		Galangale	IS	214	<i>Ulmn</i>			thl	Goaweed	1O)12 U	
	538	<i>Galantliut</i>		215	(i. tec Coihri*	153ij		504	<i>Ciotbya</i>	S»	
jm	sat	<i>OattnWa</i>	Uifi	215	O. see Sevfetio	1537		4KJ	(ioldlt&chfn		
58	517	<i>GUas</i>		yy,	(j. him	11244	20717	897	O o U Hair	SB}	
		via	r.vi	is	<i>BS</i>	1980			<;.iiiiin Bod) MJB	
	an	G. tee Ijpcjprofiscn		848	G. see <i>Tmaunia</i>	1757		SMi			
	JM	(jallianum	6J51	1C	Giant 1 cniu-1	880		138	Golden Rod	OMC W£	
	397	<i>Otto</i>	24184	40U	iinyuihiuIT	94801		175	<i>Gulden tiaKifragu</i>		
	317	>a	1USS3.V.	iii	<i>BS i ilui</i>	MS			<i>Gulden Tliiitlc</i>	Bfl	
•si	511	i. rite >J	1 Vrs		ft. tee Ipnotnoptil	SS83		aa	Golden Tbirtte	£479 3	
		O. «* Swett/o	9141	512	ft. tee Ipnotnoptil	1.>Iri		337	tfojj cf Pjtuni-rc	j*ES	
	SOB	G. lecTephrrtU	B083	571	.MI	1988		370	GoUyloctu		
	BB	531 flalcn/i	1905	Mi	1.I.1 USIrI, Or. flt*7			536	Gototia		
	•98	ir.JteiJudOIOll		BM	i:n	(uNyHriwr 1GO78	J817	.71	G. «; Notfli*	M	
	toT	m		a	Niu_k,r	10		507	1f(>ii).u.i		
3-J7	522	G. see Verbetliu	EtO!								
8	BD	<i>Ualipia</i>	SJU9	143	" * " "	9800	11+3	rq	<i>CiomploK.illum</i>	1.,",	
44	BBC	Otitun	4x	IBS	Cttijffilium	6646	BS		G. see liurlbn/ft		
45		Anpt'rula	348		Ginkgo	£3476 (ty.	BSSB	977	G. set' Hatycbllum	1!	
45		(/. tecSherardi'a	350		Gln teng	BBBW	BB	IBS	GOULJIII dlt'-N.I	3()	
411		G. tee Val&nUa	8G1	ii ;	531	Gistik'd	£90		G. tee Oplotheca	J3n	
478	SOS	GalpMmla	MSa	188	H	Glatitilim	198		4.1		
<::		(itlinhxr	3359	537	Oitht<o	19019		3SS	540 tutmuttis		
as	ftK	[Gamboge { , g g	3997	18	G. see Anlt4nthus	125		371	fim^hrtt	BBJ	
IKS	5D4	<i>Gumbhpa</i>	12105	17	G. see Babima			52	srput		
188	504	<i>Gironji</i>	USfi	18	(il.nlwyn	1371		B5			
9				19	Glattwort	S3		79	Gonotemon		
80	519	G. ir.i.ii fiaUam	5S1	£01	Illiustiitmrj Thnrn			79	^f volft		
81		OK			ijjij li Ø,	LWf		BB	GOO*K>0>1.1. or. III	in	
81		G. see (itnipa		SIR	493	Glasticium		99	Good Hew*	6168 g	
81		G. MO Kindia		92	BM	OUUi	18902	511	QooxMi		
138		(iurilt-n Rocambole		41J	BS	Glcch.'.mn		877	Good night	4245 *5	
US		OaviMffil	SM	423	SIS	Glciclienjn		368	&&G G nod y era H		
1	Garland	\$ inifn	j iKifli	193	SM	QUUi		SO	Oowwunj SBW< '	'	
BSI	flower			517		Globe Amaranth		EO	Goo*e Corn Bl	E	
1ii	Garlic					(;ii.lw Flower		279	Goosfuot	1743S <S	
188		Garlic P»r				Globe ThiKle		214	Linsky	13658 K	
171	SOB	Gm-iiga						es	Gordt>n/«		
	538	Gtxt&fl.i						W	Gorttrw		
	542	Gutrlidum						m	G. MW GM4JII>	V	
	MB	t'.T-uk'&riiha						MB	DOMfpion		
	tM	"IIA						So	OoarfHuiu	b>150 ggj	
1.1									See Goujinia		
98	\$48	nio								cd	
173	BIS	Gaultheris									
156		G. fe Oxyccoui									
155	514	1. iii.r.i									
	BOB	(Mikola									
	aaa	Gc Ait rum									
MS	£39	Gcitcomi-ria	Ea								
18	537	GeUtอร'tia	•00								
	514	OUa	SU								
MS	581		893								
71	wir, <i>Gfhimium</i>		RM								
8			303	811	Glytyrrhlia						
81	510	0Mh>i	BS		Glypnomlriot						
8L		1.11.1. tree	ISO		lijn'l'iifl						
9ft>	511	Gcmfits	IM		m an	Gnaphiliuum					
		Cr. sco Adencoc&r-put	SI0			O. tee Antennaria					
	Ifi2	G. BPC Cyclbia	ISM			G. <AtMnn					
97			ISO								
97	596	GontiiHd	IM								
		CrBTNTidn^, Or. 129.	303	811	Glytyrrhlia						
	058	(i<nt>rum	BOB		Glypnomlriot						
	51S	<co(DH)yrt	H4		lijn'l'iifl						
		fl. t.M Anifira	318		m an	Gnaphiliuum					
	454	1.ri. /1.i.sum	316			O. tee Antennaria					
						G. <AtMnn					
	NO	'-ma									
	SSO	(t.t.'iihil	em								
	488	(ieorglina	9406								
	SAS	fi'Ehxii'cO^ Or. 50.									
	276	Geranium	E8B								
		O. sec KrOfiium	SSI								
	jgl	G' tec IVlargftpium	53*								
		G. *w SarcocablOR									
	598	Ottfltlfc	1								
		G. nee Seymirij	1780								
	lit	GcrA^hb nth us 4517	S23								
		SSI GwMr*.	BJH								

GENERAL INDEX.

Kat.	Sp.	Gen.	Lin.	Nat.	Sp.	Gen.	Lin.	Nat.	Sp.	Gen.	Hen.
1	Indian Shot		414	Jamaica Milkwood			175	Knawel			1375
	Indian Tobacco	24249	277		25173 & 4	2888	40	Knight/a			324
	Indigo	107	292	Jamaica Redwood			91	Knot Grass			725
	Indigofera	207	174	Jamaica Rose			229	Knowltonra			1635
	511 Indigofera	207	198	Jambolana tree			46	Knōxia			354
	512 Inga	28ft	155	Jambolifolia			S77	KobresVi			2600
	524 Inocarpus	135		10084 8c 5 sy.	1189		99	Kbch/a			809
	Institale	3215		Jahbot	12732	1487	29	Kceleria			235
	Intybus	2234	197	Jambosa			531 Kcemgia	Keelpin/a	20211 jy.		2230
	350 521 'nula	2345	480	Jambosilla			531 Kola				298
	/. see Corvisartf a	2346	197	Jan'pha			503 Koleho				
	601 Ionfdium	705	533 Jan'pha			505 Kdfreuttr/a					
	I. see Cal'ptrion	707	372	Jantha	22782 jy.	2567	499 Konigfl				
	LseePombaUa	705	90	Jasbiné			501 KramenVi				
	536 Ionopsis	2567	5	Jasmine			327 521 Krigia				
			525 JASMINEJE, Or. 125.			334 521 Kthnia					
			391	Jatropha			489 540 Kunthia				
	61 527 Ipome'a	49	392	J. see Jan'pha			285 503 Kfdia				
	I. sec Argyrela	493	392	J. see Siphonia			541 KyUnga				
	471 526 Ipomopsis	3293	487	Jafiraca	28497 sy.	2269	528 LhBikM, Or. 142.				
	/. see Gflia	499	153	Jeffersón/a			512 hdblak				
	530 Ircisine	2767	497	Jersey Thistle	22201	2434	Labrador Tea				
	537 /IUDJEJS, Or. 187.		358	Jerusalem Artichoke			Zabrusca				
	'rio	16452	237	Jerusalem Sage			1336	/. abarnum			
	19 537 /Vis	142					17519	Zacath'a			
	/. see Vieussexfxa	147					19M	18154 w			
	153 Irish Heath	9944	456	Jew's Ear			2035	Lacebark			
	49 Iron Tree	384	377	Job's Tears			10136	1196			
	69 Iron.wood	558	14	504 Jhñnia							
	69} Iron-wood	C 4496	434	J6lithus	26169	3027	539 LachenaliVi				
	i12678	1482	404	515 Jollif/a			1083	L. sec Drl.jia			
	83* 175 Ironwort	1857	143	512 Jonésto			1077	L. sec Zucc6gnia			
	Iroucdna	11201 sy.	116	Jonquil	7561	933	1081	532 Lachnanthes			
	25 543 Isâchna	192	116	Jonquilla	7561	933	1202	532 Lachnanthes			
	833 528 Isânthus	1651	480	515 Jossbia			140	500 Zadanum	14953	IGU6	
	Isâria	3298	453	Jove's Beard	27266	3128	Ladies' Bedstraw				
	499 /sâtis	1877	161	Jove's Fruit	10400	1226	Ladies' Mantle				
	409 543 /scha*mum	2856	15	Joutay			337 Ladies' Slipper				
	/scha^mum	24905	2841	Jtianulloa	*28097	sy.	2580 Ladies' Smock				
	189 519 Istf70	1097	165	Judas tree			1838 Ladies' Traces	22.157			
	Istfium	3104	277	Juga	17276	1939	2496 Lady Fern	25601	2D38		
	.48 538 Isfmene	935	534 JUGLA'NDEA, Or. 173.				217 500 Lst/a				
	514 Isnarda	340	384	Juglans			15f9	336 521 Lagasca			
	521 IsoArphha	2299	384	J. see Cary'a			<2186	L. see N6cca			
	536 Isochilliu	2550	83	Jijuba	5433	660	2287	L. sec N6cca			
	536 Isoletes	2957	83	Jujube	5433	660	2695 Lagcnaria				
	541 Isolepis	"	157	Julibrissin	24848	2837	1589 LagerstræWa				
	558 Isoplexis	1788	409	July-flower	10840	1283	1582 Lagitita				
	558 Isopbgon	303	168	K. see Trichonema			165H LagacVia				
	496 /sopyrum	1639	536 JuNCAGNEIG, Or. 182.				717 Lagonychium				
	Ishra	17740	1993	541 JUGLA'NDEA, Or. 206.			1284 Lagopus				
	78 Isdtoma	5141 fy.	124	541 Juncus			16352 1921	Lagopus			
	47 Ispaghilla	3215	125	J. see Bclhzula			2213 Laggscris				
	Ispagool	3215	364	J. see Jungermannia			2214 L. sec Borkliausio				
	Italian Beech		404	Juniper			2215 Crepis				
		23456 2666	404	535 Juniperus			2215 L. sec Lagitita				
P	523/'tea	694	283	Jupiter's Beard			2024 L. sec Juniperus				
	//VAa	5814	173	17622	1968		2014 L. sec Juniperus				
	522 Tva	704	514 Jussietm				2214 L. sec Juniperus				
		2458	8	1353			2214 Lambert				
	14677 1646		529 Justi'cto				319 Lambi's lettuce				
	/trira	23876 2724	245	58			9f1 Laminariu				
	Ivy	718	244	J. sec Aphelandra			5078 Laminaria				
	537 /'xia	118	8	1731			26286 3040	Laminaria			
	/. see Bdbihna	135	96	J. sec Barler/a				26286 3040	Laminaria		
	J. see Hesper&ntha	121	264	1725							
	16 /. see Sparaxis	122	8	J. sec Elytraria			528 -Lámium				
	18 /. see Synnþta	129	96	6228	779	237	1665 Lampwick				
	536 ^ I. see Trichonema	119	264	Kdber	16634	1884	15072 1674	Landria	7656 s/.		
	520 Ixbro	379	415	Kaki	25192	2889	942 Lancashire Asphodel				
	522 Chomelia		171	523 Kalmia			8923 1070	Lancewood	14383 1625		
	522 Chomelia		512 Kalosanthes				Landra	16684 1897			
	522 Tcfamerium	382	160	KanguruVine	3360	400	227 Lantana				
	522 Tcfamerium	380	50	Kar&tas	7688	951	1738 Lantana				
	522 *lx>tapita	10947	118	KarAtto	7671	948	7129 Lantana				
	Jacaranda	1712	145	Kauki	9347	1156	126 Lapeyrotista				
	J. sec Bignbnfo	1706	350	Kaulfust'a			126 Zappa				
	J. sec Bignbnfo	22862	314	2344			20226 2237 Zappago				
	522 *lx>tapita	237150	214	512 Kehnedta			193 " Ppula	(12130	1443		
	J. sec Cea	22151	315	2144			521 Zapsana				
	J. sec Cea	2434	282	512 Kerria			2228 Z. see Rhagadolus				
	522 *lx>tapita	237150	214	1531			2230 Z. see Zacitha				
	J. sec Cea	2434	315	2149			2229 Z. see Zacitha				
	J. sec Cea	2434	282	478			516 Larbreia				
	J. sec Cea	2434	401	Kiggelarw			3351 Larch				
	J. sec Cea	2434	500	2798			2687 Lardizabala				
	J. sec Cea	2434	533	Kirganiila			23578 Lario				
	J. sec Cea	2434	292	Kissi			2685 Zârix				
	J. sec Cea	2434	287	18163	2038		2687 Larkspur				
	J. sec Cea	2434	502	Kitaibeto			1599 Lascbea				
	J. sec Cea	2434	285	2006			913 Larrca				
	J. sec Cea	2434	503	Kleinfl&fla			1397 Lârrca				
	J. sec Cea	2434	337	2001			518 Fascrptium				
	J. sec Cea	2434	521	Klelnia			881 Z. see Cai-hrys				
	J. sec Cea	2434	336	2291			889 Z. sec 6'ndliim				
	J. sec Cea	2434	521	K. see Cacalia			860 Z. sec Zigisticum				
	J. sec Cea	2434	487	2289			875 Z. sec Slier				
	J. sec Cea	2434	24	542 Knapp/a			3387 Z. see Tüftpsia				
	Jamaica Dogwood		17666	1975	r22083			81j			
	Jamaica Etony		360	Knapweed	22098	2434					
					€.22151						
			44	520 Knautia							
			347								

GENERAL INDEX.

Lin.	Nat.	Bp.	Gen.	Lin.	Nat.	8p.	Gen.	Lin.	Nat.	Sp.	Gen.
108	Laserwort	881		445	Lepraria	3113		49	Lily Thorn		
104	Laserwort	6785	845	433	L. see Chlorococcum	3013		320	Lime	19666	2187
462	Lasiobotry8	3227		■ -499	Leptalium	1868		218	503 Lime tree	9665	§
337	522 Lasiostpermum	2305		6	528 Leptandra	44		320	Limetta		
86	503 Lasiopetalum	680		18	540 Leptanthus	134		143	516 Ximeum		
86	L. see SeringtVv	682		18	L. see Heteranthera	135		217	536 Limnocharis		1558
86	L. see Thomasia	681		360	Leptaurea	22061	2431	170	504 Limonia		13#
403	540 Latania	2818		255	Leptocarpae*a	1820		170	X. see Clausina		13**
248	529 Lathrs'a	1770		396	Leptocarpus	2743		170	X. see Glycosmis		13*
313	511 Xathyrus	2136		33	Leptochloa	262		82	X. see Triphasia		
313	X. see'rynum	2134		92	Leptomeria	742		115	Ximonium	7480	918
314	X. see O'robus	2138		434	Leptifimus	3033		170?	Ximō. f 10963 sy.	1S&	
192	Xathyrus	12369	1460	196	Leptospermum	1480		320 J	nium I	19666	2187
381	Laugera/a	2636		459	Leptostruma	3176		252	528 Limosella		1559
160	Laurel	1226		36	Lepturus	23804Y	278	249	528 Linaria		1775
200	Laurel Cherry	12887	1502	396	Lepturus	2740		99	508 Linchnia		809
156	Laureola	101% 1197		541	Lepyrbdia			252	528 Lindernwi		1800
109	Laurestinc	7109	892	521	Lēru*	£336		251	L. see Mfeus		17\$
	531 ZAuaNEiE, Or.	158.		429	Leakea	2993		431	644 Lindsay		2931
200	Laurocrasus	12887	1502	311	Lapsidea	2126		502	LiseJE, Or.	27.	
415	545 Laurophylus	2894		305	Lassettifl	2098		71	Ungun	4598	577
160	532 Xaurus	1226		92	Lestibudta'a	732		433	Lfkia	26157 sy.	30\$
160	X. see Cinnambum	1225		217	Letsbmta	1562		242	519 Linn*i		1703
161	X. see Persea	1228		64	L. see Argyrela	493		147	Linnadna	9477 sy.	U7^
234	528 Lavändula	1656		323	Lettuce	2204		12	525 Linociera		
483	L. see Anisochillus	3384		396	Leucadendron	3749		3697			
287	502 Lavátero	2008		319	Leucadendron			370 y	Fym	122649	25g
234	Lavender	1656						3723		622753	25JJ
337	Lavender Cotton	2:504		238	Lehcias	1675					
336	521 Lavinia	2288		363	Leucetria	12255	2442				
87	501 Lavradia	700		468	Leucocoryne	3281					
154	514 Lawsonia	1180		38	Leucodendron						
134	539 Laxmannfe	1049				2509 sy.	305				
52	Leadwort	423		428	Leiicodon	2981					
460	Leanglum	3199		119	Zeucoujum	962					
157	Leather wood	1196		475	L. see A:i:i*	3334					
279	511 LebeckiYi	1956		65	Leucopbgon	518					
.101	L. see Indig6ferra	2072		247	Leucoscpétrum						
288	502 Lebrethiūfa	2013				15695	1752				
441	Lecanbra	3093		38	Leucospermum	305					
36	500 Lecher	292		342	Leucostēmma	2324					
79	522 Lechenia61tfo	614		360	Leacea	2432					
440	Leefdea	3085		474	Levfsticum	3324					
219	515 Lécythis	1588		481	Lewftist'a	3367					
102	518 Ledebfuria	829		352	Lefsera	2370					
219	Ledon Gum	13950	1589	352	L. see Longchämpsta						
171	523 Xēdum	1336				2369					
171	X. see Ammylnle	1337		353	L. see Relhan'a	2372					
74	594 Leeā	594		80	Leycestena	626					
145	Lee Chee	9358	1159	333	521 Lihtris	2265					
132	Leek	8602	1046	103		16703	835				
36	543 Leersia	288		108	Libanfetis	\ 7007	881				
	509 LEGUMINOSJE, Or.	67.		109	C7097	889					
409	539 Leimanthium	2840		460	Licea	3202					
171	Leiophy'Um		11004 sy.	441	Lichen	5^,^3088	C3055				
	"Lemknea	3070		443		C3101					
375	541 Xēmna	2588			545 XCDE^NEB, Or.	8.					
320	504 Lemon	19665	2187	439	Lichlna	3079					
410	Lemon Grass	24916	2844	140	539 Lichenstefnja	1108					
312	Zens	19268	2134	125	540 Licdua	998					
109	Lctn&go	7119	892	354	522 LidbeckiVi	3378					
	529 LENTIBULA^RUE, Or.	147.		354	I^* see Cenia	S379					
313	Xenfcula	19270	2134	90	522 Lightfootfe	715					
312	Lentil	19268	2134	169	Lignum-vite tree	1303					
398	Xentfcus	24218	2763	123	Lijgtu	8041	979				
460	Leocarpus	3198		107	518 /ig&sticum	875					
238	528 LeonbtU	1676		103	L. see jthamfinta	835					
138	497 Xē6ntice	1086		108	Ir- see Zaserptfium	881					
	Z. see Caulophyllum	1087		474	/.. sec Lcvisticum	3324					
324	521 Ledodont	2207		105	L. see A/eum	857					
487	L. see Callopœa	3398		474	L. see Molopspér.						
324	L.sceThrfncia	2209			mum	3323					
138	Xeontoptalmon	9038	1086	474	L. see Petrosellnum	3330					
340	521 Leontopodium	2316		107	L. see Physosper-	876					
237	528 Leonurus	1673			mum	876					
	Leopard's-bane	2353		107	L. see Pleurosper-	877					
344)	Leopard's, f	21140	2335		mum	877					
3515	bane	121555	2350	474	L. see Trichosa'nthes						
382	540 Leopoldliw	2641		108	L. see Wallröth/a	832					
454	Lcbta	3143		5	525 Zigusttum	40					
	L. see Mfrtula	3137		5	Lilac	41					
236	528 Lepchlnia	1663		136	Liliago	8881	1063				
326	Lepicanne		20077, &c.jy.	136	Liliarstrum	8879	1062				
			2212	129	Liliaster	8879	1062				
43	Lepic^phalus	2879 & 2880 ay.	345	135	Liliura	8879	1016				
	X. see Euñmia	1875		129	L. see Fritillaria	1018					
244	529 Lepidagathis	1724		135	Lilio-Hyachthuf						
262	499 Xepidium	1873				8814	1054				
	Z. see Hutchinsia	1851		129	Lily	1016					
262	X. see Senebiera	1872		135	Lily Hyacinth	8814	1054				
259	L. bee Teesdalia	1852		131	Lily of the Valley	1038					
31	541 Lepidosperma	152		132	Lily Pink	1044					

GENERAL INDEX.

(553)

F	N.t.	Sp. (Jen.)	Un.	Nat.	Sp. *	* ..	Lin.	Nat.	ftp. Gen.
50	L. «ee Symphbrla	623	65	523 Lyslnema	511		188	7 Mangostan	12104 142.9
52	Looking-glass Plant	2725		514 LYTHRARI BX	Or. 79.		188	504 Mangosthna	12104 142i)
58	Loose-strife	457	189	514 Lythrum	1451		188	Mangosteen	1429
3	514 Loptafa	18	189	L. see Nesa>fl	J4J2		413	Mango tree	2879
528	Lophanthus	3871	400	524 Miba	2791		188	Mangrove	1428
538	Lophiola	911	414	Mabbla	25177 2889		386	Manicaria	2678
513	Lophlra	1504	382	Macaw tree	23302 2b45		387	Mdnicot	23568 2682
513	LOPIIREJL Or. 75.		502	Machucunha			392	Mânihot	2718
3	Lepimla	2018	187	Macleayo	1J23		288	Mdnihot	17936 2014
3	Loquat	J511	380	Madura	2623		96	£42 Mansurus	282
au	LORA'NTHEJE, Or. 101.		188	Mdgui	12121 U\$		311	Manna	2128
Lord Anson's Pea			371	Macradtnia	2543		12	Manna Ash	786 83
83	Lotetree	19287 2136	74	Macrocnemum	597		3	537 Mantisia	16
83	Lbtua	5421 66)	15	M. see Musae'nda	612		253	528 Manilea	1806
300	L. see Carmichaell	2070	256	Macropbdium	1828		412	Maple	2867
300	L. see Dorjrcnium	2067	472	Macristylis	3304		1	537 Marantar	2
301	L. see Hosackia	3392	162	Macrotropis	1240		424	544 Maratia	2952
301	L. see Tetragnod-		216	Macrbtys	1546		215	504 Marcgravta	1542
lobus	lobus	2069	188	Madagascar Nutmeg			504		
217	Lbtus	J 5421 (66)	74	Madagascar Potato			432	Marchantia "	2999
217	Lbtus	J 13809 1556			4815 591		4	Mare's Tail	24
2	Loirea	25190 2889	73	Mad Apple	4809		468	512 Margyricarpus	3280
S3	Loureira	21J7	45	Madder	349		20	537 Mârica	144
107	Lousewort	17H1	243	Madia	2331		20	M. see Bob&rtia	146
I<ovage			257	Madwort	1840		363	M. sec Sisyrinc'iux	145
72	Lovage	S1864 2387	71	Ma'sa	581		22	541 A/arfcus	166
33	Ix've Apple	4703 590	326	Magnolia	1618		238	Marjoram	1680
Love Grass	Z55	200	496	MAGNOL&car, Or. 3.			476	545 Mdrlea	1341
Love Grass	2222 256	268	503	MaMUB	12879 1502		219	Marmifos	13916 1585
Love-lies-bleeding		168	503	Mahernia	1917		2372	Marrubias-	C15052 1673
		168	503	Mahdgoni	10870 1297		238	trum	115103 1677
316	Liibia	23245 2628	168	Mahogany tree	1297		237	528 A/arrubium	1672
530	Lubmla	19470 2152	139	Mahbnta	1091		483	M. see Beringeria	3377
S	Lucern	503	421	Maiden Hair	2930		95	525 Marsden/Vi	775
519	Luculia	18314 2061	270)	Maiden Hair	C16912 1923		335	521 Marbhall/a	2280
524	Luchma	643	419J	Hair	(25448 2921		213	Marsh Cinquefoil	
500	Liidia	562	385	Maiden Hair tree	vane	1527			
55?	I* see Prdccta	1568							
	Lud61tya	2406 sy.	53	Maiden Lip	23476 2669		287	Marsh Mallow	£007
36	LudJ.via	23481 sy.	15	Maiden Plum	3518 426		232	Marsh Mangold	1643
	Lmfwfgia	2670	238	Majordnia	107		106	Marsh Pennywort	
514	Lmfwfgia	404	108	Malabafia			485	544 Manilas	6900 863
515	Lftfh	2630		7046 ft 7047 sy.	883		544	MAKSILIA CAP	2960
529	Lumnltzera	1688	160	Malabar Leaf	10372 1225		235	Marsypianthus	
L. see Acrocephalus			111	Malabar Nightshade	908			14020 sy.	1661
L. see Mosch6sma	3382	9	289	Malabar Nut	559 58		229	Mdrtagon	8408 1016
Luna	7553 933	160	160	Malabar Rose	17978 2014		244	526 Martnia	1720
499	Lunaria	18&J	292	A/alabatrum	10372 i2&		244	M. see Craniofaria	1721
257	L. see Farsetia	1835	292	MalachodOndron	2035		238	Mdru	15128 1C180
142	L. see Ricbtia	IS34	287	M. see Stuartia	2034		233	ATarum	14737 1648
55	Lunaria	9222 n a g	184	Malachra	2009		52	Marvel of Peru	421
325	Lungwort	438		Malacophy'llum			135	539 Massbnfh	1058
300	Lungwort	20024 2211	373	Malaxis	11833 1409		106)	Masterwort	fJJJ!
284	Lupinaster	18531 2066	373	M. see Lfparis	2573		109	j	
Lupine	19H5	197	373	Malay Apple tree	2576		238	Mastic	15128 1681
Lupinella	18631 2071		12721	1487			239	MasUch	15157 1681
24	7,uplnuB	1985	260	Malctanfo	1862		239	Mastich'ma	15157 1681
8?	Lysam6ps16250	1992	422	Male Fern	25609 2918		398	Mastich tree	24218 276J
398	Lysam6ps16250		218	Mallofcica	13857 1575		145	505 Matafha	1162
190	Luteola	12235 1459	286	Mallow	2004		210	Mat Grass	0 76
125	Luzula	995	288	Mallow Rose	2014		239	Mathloia	1817
137	Luzurfaga	1076	286	502 Afalope	2003		381	M. see Guett&rdra	2626
250	Lychnfdea	15891 1782	182	505 Malpighia	1393		354	522 Matricaria	2376
186	Lychnis	1415	170	M. see Bunchbsia	1329		53	527 Matt/Vi	427
	r 3 % 9	480	182	M.sceByrs6nimia	1394		250	528 Maurandy	1778
	lychnltis	415072 1674	201	505 MALPICRACOE, Or. 43.			399	540 Mauritifl	2781
	C21868 2389		286	A/alus	12955 1507		111	Maurocenia	7219 895
	Lycoserfssa	589	502	A/âlva	2004		316	Max	19451 2149
	J-y'icum	588	290	M. see Cristaria	2020		370	536 Maxillaria	2537
J60	Lycrtgala	3195	502	JIAALVA'CEA, Or. 28.			413	540 Maximilians	13762 1548
460	Lycoperdon	3191	502	Malvaviscus	2012		216	May Apple	23042 2602
461	L. see Geastrum	3193	218	504 Mammea	1580		377	Mays	
460	L. see SQC Institale	3213	218	Mamme tree	1580				
	* see Leucarpus	3198	194	517 Mammillaria	1471		412	J508 Mdytenus	35
B27	* see Rhi*opbgon	3182	391	Manchinel	2713		251	528 Mazus	12868
g44	Lycopersicum	590	391	Manchinel	23800 2714		32	Meadow Grass	1786
424	Or. a		391	Mancinilla	2J797 2713		229	Meadow Rue	16J3
SS		2955	320	Mandarin	19573 2187		141	Meadow SafiVon	1119
55	J	444	353	Mandiana	21692 2375		104	Meadow Saxifrage	84)
122	see Nf,neu	443	71	527 Mandrfgora	585		203	Meadow Sweet	
	Lyc'pus	69	71	Mandrake	585				
	Lycbrus7889&90fy.	968	49	519 Manlttia	389		216	498 Mecon6ps:s	1553
	L'geum	id8	68	Manghas	4463 & 4 sy. 550		140	538 Medeofo	1114
	Lycfdium	2946	413	509 Mangifera	2879		140	M. see Myrsiphy'tiuin	
	Lysbdil	577	183	M. see Buchanftnw	1406				
	Grass	272	85	M. see Etaedolndrum	672		296	511 .. .	1109
	S	183	85	M. see Spndrias	1405		297	Medicago	2061
		1342	406	Mdngium	24677 2837		296	M. see Hymenocar-	
		545	188	Mdngle	12103 1428		JP*?U	2062	
		530	2506	524 Mangilla	570		Medick	2Q5I	
		70	3	Mango Ginger	182 14		Medium	5022 607	
				£00			Medlar	1505	

GENERAL INDEX.

J.Jn.	Nat.	Sp.	Gen.	Lin.	N ^t at.	Sp.	Gen.	Lin.	Nat.	Sp.	Gen.
190	Medusa	12281	ty.	1460	401	Mercury	2794	405	M. see <i>Tnga</i>	£834	
190	Medusa's Head			141	539 <i>Merendaria</i>	1120	168	M. see <i>Lagoit-</i>			
		12281	1460	141	<i>M.</i> see <i>Cölchicum</i>	1119	K8	<i>chium</i>	1284		
510	490 Megacarpa^a	1855		174	515 <i>Meriana</i>	1363	250	M. see <i>Prōsopig</i>	I**		
371	536 MegacUnum	2345		80	<i>Meriana</i>	5280	528	A/imulus	1783		
.175	Megasea	11239*y.	1380	413	<i>Merisma</i>	2132	145	<i>Mimusops</i>	I ⁵⁶		
33	543 McGastachya	2:16		412	534 <i>Mertensf*</i>		197	<i>Mini</i>	12736 1487		
33	M. see Scleruchloa	257		452	<i>Merilius</i>	3122	235	Mint	J659		
435	MeUgbium	26268	3044	448	<i>M.</i> sec <i>yfgaricus</i>	3118	37	<i>Minuartia</i>	296		
319	515 McLaleúca	2179		451	<i>M.</i> sec <i>Cantharillui</i>	3121	2	<i>Mibga</i>	124 10		
487	M. sec Astarte <i>i</i>	3393		203	517 <i>Mesembryanthem-</i>		52	<i>Mirabilis</i>	421		
246	528 A/elampyrum	1741		208	<i>M.</i> see <i>Hymendgyne</i>		165	<i>Mirbelfo</i>	1264		
363	522 A/elampbdura	2446		513	<i>Mcsoglia</i>	3034	397	Mistletoe	2750		
335	£21 McLanthera	2279		434	<i>Aespilus</i>	1505	49	<i>Mitchell*</i>	387		
464	McLancbnium	3269		200	<i>M.</i> see <i>Amelanchier</i>	1512	520	<i>Mitella</i>	1380		
1197	519 Melanopsfrium	1456		202	<i>M.</i> sec <i>Cotoncaster</i>	1513	48	<i>Mitrasa'cmc</i>	371		
491	509 Melanorrht'ca	S422		200	<i>M.</i> sec <i>Crats'gus</i>	1506	454	<i>Mitpila</i>	3137		
474	518 McLanosellnum	3;520		202	<i>A/espriobutrya</i>	1511	4	<i>Mniarum</i>	30		
269	Melan6xylon	16849	1920	202	<i>M.</i> see <i>Photinia</i>	1509	429?				
	539 A/ELANTHA'CKJ ^a , Or. 198.			56	527 <i>Messerschmid/d</i>	447	193	<i>Mocanh-a</i>			
140	A/eUnthium	1111		56	<i>M.</i> see <i>Tourncfurba</i>	446	483	<i>Mosch6sma</i>	12471 3,381		
140	M. sec Androc^mbium	III		340	521 <i>Mctaftsia</i>	2319	398	<i>Modicca</i>	277		
400	M. see Chanoilirium	2792		94	525 <i>Metastelma</i>	762	158	502 <i>Mehrtingh</i>	1S08		
409	M. see Lchnaithium		2840	60	<i>Mitel</i>	4012	52	<i>Moenchia</i>	3452 & Gsy. #16		
				110	<i>A/etbpium</i>	7146	239	Moldavian Balm			
140	M. sec Ornithogl6s-sum			1113	515 <i>Mctrosideros</i>	1482	342	Mor.n/i	15206 1685		
140	A/.seeWürmbea	1116		1116	<i>M.</i> sec <i>Acmena</i>	3361	124	538 <i>Molineria</i>			
17	537 Melasphas'rula	127		480	<i>M.</i> sec <i>Angphora</i>	3360	32	543 <i>Molinia</i>			
173	515 Meldstoma	1356		197 \	<i>M.</i> sec <i>Calhste.</i>		145	<i>M. sej Cui^nia</i>			
174	M. sec Acibitis	1362		4H0J	<i>mon</i>		401	<i>M6l/s</i>	24416		
174	M. see Chitbnia			196	<i>M.</i> see <i>Leptospér-</i>		92	<i>Mullin</i>	6024 & 5 sy.		
174	M. see Cliddinia	1359		381	<i>Mctr6xylon</i>	23279	37	502 3/oilugo			
174	M. see MichnrVi	1357		105	<i>A/feum</i>	857	111	<i>M.</i> see <i>Pharnaceuwi</i>	906		
154	M. see Osbuckia	1181		105	<i>M.</i> see <i>ifnethum</i>	858	441		34f		
174	M. see Plerbma	1360		473	<i>M.</i> see <i>Ptychbtis</i>	3315	474	518 <i>Molopspérimum</i>			
	M. see <i>Tochba</i>	1361		104	<i>M. sectfson</i>	849	238	<i>Molucca Balm</i>			
514	MELASTOMA'CEA, Or. 81.			122	<i>Mexican Lily</i>	7937	238	528 <i>Moluccilla</i>	1677		
269	503 Melhania	1920		339	<i>Mexican Mugwort</i>		483	M. see <i>Chasmbnia</i>	5'JJ		
168	506 A/elia	1293		301	<i>Mexican Tea</i>	18634	133	JJtby	8700 1046		
242	507 Melianthus	1704		267	<i>Mexican Tiger-flower</i>	16760	183	<i>Mthubin</i>	m*5sy. ^L		
32	543 Melica	251		356	522 <i>Meyera</i>	2395	389	515 <i>Mom6rdica</i>	& M%		
32	M. see MolntVi	252		337	<i>M.</i> see <i>Castilia</i>	2302	881	M. see <i>Ltfa</i>	\$&		
32	Melic Grass	251		156	<i>Mczereon</i>	10137	25	542 <i>Monachne</i>	W		
65	523 Melichrus	506		156	<i>Mixeum</i>	10137	267	MONADE'LPHIA, Cl. 16			
145	505 Melic6cca	1160		156	<i>M.</i> see <i>Calhste.</i>		267	M. TRIA'NDRIA, Or. 1.			
155	507 McUope	1191		347	<i>Michaelmas Daisy</i>		268	M. PENTA'NDRIA, Or. *			
298	Mcilot	2065		21330	<i>Miranthemum</i>	47	271	M. HEXA'NDRIA, Or. 3.			
298	511 A/ellibutus	2063		314	511 <i>Micranthus</i>	2142	285	M. HEPTA'NDRIA, Or. 7.			
296	M. see Medicago	2061		48	522 <i>Michaùxia</i>	1177	286	M. POLYA'NDRIA, Or. 9.			
298	M. see Pocdkui	2064		226	<i>Microcâle</i>	368	468	MONAD'KDRIA, CL 1.			
297	M. see Trigonilla	2063	8	36	524 <i>Microchloa</i>	276	1	M. MONOGY'NIA, Or. 1.			
239	528 Melfssa	1684		36	525 <i>Microbta</i>	1357	1	M. DECA'MIRIA, Or. 6.			
239	M. see Calamfntha	1684		51	543 <i>Microfflnm</i>	409	285	M. DODECA'NDRIA, Or. 7.			
233	M. see Mpeta	1654		4	<i>Microlbma</i>	753	194	516 <i>Monanthes</i>	14g		
240	528 Melittis	168f		93	525 <i>Micromeria</i>	3379	10	528 <i>Monarda</i>	7J		
389	A/lo	23701	2699	93	525 <i>Micromeria</i>	3379	433	M. sec <i>Blephflia</i>	S7»		
194	Mvlocáctus	12511	1470	483	528 <i>Micromeria</i>	3379	4	542 <i>Monerma</i>	" J£		
479	517 Melocactus	3358	183	1779	<i>Micropetalon</i>		36	M. see <i>Ophiurus</i>	27g		
124	543 Melocanna	985		1779	<i>Jy.</i> 1402		49	525 <i>Monetta</i>	%		
269	503 Melochia	1918		364	521 <i>Micropus</i>	247J6	58		3844 - g[
269	M. see Riedl&a	1919		373	537 <i>Micr6stylis</i>	2575	238	Moneywort	A(g)J2 31g		
93	525 Melodinns	748		98	531 <i>Microtea</i>	801	30U				
398	Melon	23701	2699	369	536 <i>Micritis</i>	2511	4	Monilia			
73	Melongnha	4808 sy.	591	190	536 <i>Mignonette</i>	12241	464	Monkey Flower	*		
398	Melon Pumpkin			333	521 <i>MikanVr</i>	2266	227	Monilia			
		23694	2698	462	<i>Mildew</i>	3226	374	Monk's-hood	f14S,3,1		
479	Melon Thistle	3358		355	<i>Milfoil</i>	2387	49	2247	118203 2		
194	Melon Turk's-cap	1471		356	<i>Milfoil</i>	21847	294J	293	507 Monnrla		
389	JV/leopeo	23694	2698	23	542 <i>A/lillum</i>	2387	501	501 Monnlna			
433	Meloselra	3008		24	<i>M.</i> see <i>Gastridium</i>	202	530	530 MoNOCHLAMY'DEF*			
375	515 JI/eldthria	2591	25	25	<i>M.</i> sec <i>Monachne</i>	190	Subc. 2.	Subc. 2.			
189	Molvilla	12213ty.	1454	25	<i>Milk Parsley</i>	873	535 MoNOCOTYLE'DONKSf				
156	513 A/emeclyon	1195	US	256	<i>Milk Vetch</i>	2102	CI 2	CI 2			
513	JI/EMECY'LEJK, Or. 61.	106		256	<i>Milkwood</i>	25174	227	496 Monod'bra	J		
257	499 Menibcus	1841		257	<i>Millkwort</i>	2053	374	MONCEYIA, Cl. 21.			
417	544 Menfrium	2905	414	259	<i>Millra</i>	1078	374	M. MONA'NDRIA, O			
496	MEMSPI:RMA\EE, Or. 5.	295		359	<i>A/ilcf blium</i>	21847	375	M. MONA'NDRIA, O			
402	497 Menisiërmum	2806		356	<i>A/ilcf blium</i>	2387	375	M. TRIA'NDRIA, Or. »			
400	M. see Cbcculus	2788		362	528 <i>Milleria</i>	2438	375	M. TETRA'NUKIA, Or. *			
400	M. see Coscinium	2789		365	<i>M.</i> see <i>Flavèria</i>	2462	375	M. PENTA'NDRU, Or.^			
400	M. see Tiliacbra	2790		365	<i>Millet</i>	1590	375	M. HEXA'NDRIA, Or. *			
235	528 A/entho	1659		24	<i>Millet Grass</i>	181	375	M. TETRA'NUKIA, Or. *			
483	M. see Aphanochllus			33	<i>Millingtania</i>	3277	378	M. PENTA'NDRIA, Or.^			
483	M. sec Dyssophph lla	3373		372	525 <i>Mimetes</i>	306	380	M. HEXA'NDRIA, Or. *			
196	51C Mcntzelin	1478		38	532 <i>Mimeta</i>	2834	381	M. OCTA'NUBIA, Or. *			
58	521 A/enyanthes	463		405	512 <i>Mimbsa</i>	2834	382	M. ICOSA'NDRIA, Or. *			
58	Af. see Villarsia	464		407	<i>M.</i> see <i>Acacia</i>	2837	382	M. POLYANTRIA, OJ »			
153	521 MenztalTv	1174		92	<i>M.</i> see <i>Darlingtania</i>	743	382	M. POLYANTRIA, OJ »			
401	53* MercuTidUs	2794		409	<i>M.</i> see <i>Enthda</i>	2838	387	M. Gagnebna	1286		
				168	<i>M.</i> see <i>Gagnebna</i>	1286		Or. 10.			

**• Nat.	Sp. Gen.	Lin. Nat.	6p. Gen.	Lin. Nat.	Sp. Gen.
2 522 MonApsii	610	263 499 3/yagrum	1879	398 505 NegHndium	84855 ty. Sm*
65 523 Mon6toca	515	2&? M. see Calep)na	1895	398 505 Negiindo	8774
JJO 523 Mon6tropa	1330	262 3/. see Camellna	1K70	246 Negundo	15616 sy. 1743
gf 506 Monsonta	1991	258 M. see Cochlic'ria	1848	488 521 Npia	3404
** M. see Sarcocaulon	1992	266 M. see Didēmis	1898	197 515 Nelitria	1485
g* 522 Montanba	2444	262 M. see Ne'sm	1871	9 529 Nelsonta	60
g* 502 Montezfma	1995	260 M. see Aapfstrum	1860	226 498 Nelumbium	1613
J2 516 MAntia	094	434 Mycinēma	3026	250 528 Afcmfsja	1777
22 514 Montimā	2762	30 543 MygaKtrus	240	Nemolapathum 9184	1126
2* Moon-seed	2806	51 508 Myginda	410	398 508 NemopAnthes	2775
2J Moon Trefoil	18311	173 523 Mylocaryum	1348	60 527 Nem6phila	478
** Moonwort	2950	446 My6myces	26899 3118	368 536 Nc6ttia	8496
j3* Moonwort	9222 1126	529 MVOPIRINA, Or. 144.		368 N. see Goodyera	2492
2J 537 MoraeV.	143	848 529 My6porum	1761	368 N. see seLfstera	2501
3* M. see Bob&rta	146	327 521 Myosoris	2217	3 18 N. see Pelexia	8495
£* M. see Cypella	1912	53 527 A/yosotis	425	368 N. see Scenorhy'nchus	
SI M. see Homērta	1914	54 M. see ^nchd̄sa	432		
4? f M. see Vieusselxtia	147	S3 M. see Echinospes-			
IE Mochilla	3140	mum	426	404 533 Nepenthes	8497
*£ Morel		58 M. see Aeliotrd-		233 588 JVepcta	2830
lfts 527 Morenba	495	pium	424	232 N. see Anis6meles	1654
«? Mordgdnna	10891 1304	115 496 Myoēirus	930	239 N. sec Calamfntha	1684
^ 499 Moric&ndia	1885	371 Myosbrus	82713	235 N. sec Hyptis	1661
J* 520 Morina	;	480 515 Myrcia	3364	482 N. see Lophanthus	3371
1Rf f19 Morinda	648	308 511 Mynahēnus	2114	145 505 A^ephelium	9360 1159
JJJ 512 ikfor/iigi	1279	397 535 Myr.m	2751	4201 C2921	
\$1 500 Morisfnta	8045	459 Myriococcum	3184	422J Nephrbdium	^.-C2932
2* 534 Affars	2612	433 Myriōnēma	3020	4223 C2938	
4? 3/. see Hroussonftia	2756	382 514 A/yrioph'allum	2652	413 Nephrbma	3099
3% r* J/ - see Madura	2623	404 532 Myrfstica	2829	121 538 Nerline	968
IS 21 Mosckria	22.11	827 532 Myrftica	14380 1623	121 N. see Brunsvigla	967
lft Moschatel	1222	199 MyRisfrEA, Or. 159.		67 525 JVtrium	529
^ Moscatellina	10364 1222			67 N. sec Alstbnia	530
7g A/osecheutus	17961 2014			68 N. see Tabernsmon-	
528 Moschfsmas	3381	12858v. 1501		tkna	548
5\$ Moth	3992 480	1831 Myroba- lanus V.J 25035	1405	N. see Wrlghtfe	535
7? Motherwort	1673	413 I 2864		189 514 NcseVi	1453
7^ Mougebtia	S048	892 502 Myrtidia	2031	N. sec Dēcodon	3348
££ Mould	87778 3237	8 Myrdsma	64 sy. 5	N. sec Helm̄ka	1453
oV Mouldiness	3246	162 511 Myrospermum	1236	262 499 Ni'slia	1871
** Mountain Ash		461 Myrothdium	3217	Nestlcrta 21873^.	2392
IGS ,	12973 1507	102 Myrrh	830	356 Nettle	'2615
102 Mountain Ebony	1268	102 518 Mfnhis	830	412 Nettle tree	2770
Mountain Parsley		101 3/. see ChacropyI- lura	821	302 511 Neurocnrpum	707+
Itt G940 873		415 524 JVfrsine	2891	337 521 Neurolse'iia	2300
iX Mountain Sorrel	1104	70 3/. see Ardfsia	571	256 499 Neurolbma	1832
121 MoureUa	11738 1394	70 M. see Manglula	570	390 515 Neurosprma	8703
477 e.« ^flwW U189 1364		524 JVFRSINEA, Or. 120*		New Jersey Tea 5502 668	
JJ 513 Mouraria	3346	192 jl/yrhntles	12444 1460	New Zealand Spinach	13036 1519
!<* Mouse-ear	19974 8811	140 539 Myrsiphylum	1109	196 New Zealand Tea	
*		515 MYHTA'SCEM, Or. 81.			
^ ^ Mouse-ear Chick. weed	1416	156 Myrtillus	10118 1194		
113J Mouse-tail	f ^ J	197 Myrtle	1489	61 Nicandra	485
3?i Mouse-tail	£930	156 Myrtle Bilberry	10118 1194	N. sec Anisodus	486
52 Mouse-thorn	22712 2547	197 515 Mfrtm	1489	Nicker tree	1278
«?r Mouse-thorn	22202 2414	197 M. see Eugēnia	1-187	309 511 Nicolsonia	2119
Si {foi,te}	14094 1596	480 M. see Mf rcia	3364	527 Nicot;Vwi	488
2g Moving Plant	19053 2120	198 M. see OlVntha	1495	61 N. see Icthnia	489
£? 3/cfm 2OK81 2311	198 M. see Pimenta	1491	«?9 Nidularia	SI 83	
Jg Jlfucedo	27778 3237	4510 563 Mfxa.		iJ? *TM £ idU8 y*® ^Mi	
53 MixcoT	3238	464 Nsmaspora	3270	S15 500 Nicththra	1^>40
5if 3/-^na	2159	456 Nematella	3156	496 Nigella	i (nr
2g Mudwort	1793	397 535 Nageitt	2752	182 Nigellastrum	11720 1392
^7 Mugham	83581 8685	535 Nain Flageolet		72 Nightshade	501
5^ Mugwort	80828 3131	19424 2149		367 536 Nigrictlla	2482
£ Mulberry	2618	381 541 Najas	2633	387 540 Mpa	2681
2Ha Mulin	480	139 497 Nandlna	1092	418 544 Niphdbolus	2911
SI S! Mundora	1974	291 NaiME'a	18075 & 6 2023	328 Nipplewort	8888
3ls *1 Mundia	2055	224 495 Napēllus	14235 1600	590 Kiriit i	83757 2708
G6 - J 7 ^ ^ 1 M 44 8149		393 Napoleon's Weeping Willow	23978 2732	303 511 NissMia	8084
4lf ^ « ^ 0 5	4358 524	264 A?ipus	16593 1883	313 Nissnra	19303 8136
218 ftm, ?[i] «« 85303 2900		228 4% Naravelia	1632	317 Nitaria	1446
106 5[2] Mjngingia	1894	116 538 A'arcfssus	933	328 Nitella	1939
		23 542 A'ardus	176	39 532 Nivenia	308
		136 541 Narthēcium	1070	100 MMA	6559 816
485 16 X»« C«FVI	1924	226 IStrum	14356 1621	3.36 521 S6cca	2287
388 VV Becc Disēmma	3389	254 499 Nasberry tree	4504 561	89 501 Noik'tt'a	706
93 507 Mjil /rttlw&ra	2331S 267	255 Nasturtium	1819	56 527 NolJlna	448
138 5[2] Mjil	1082	11. Sec Notdceras	1821	141 539 Nollui	Heg
		124 543 Mstus	986	87 Nolitangere	5698 &/8
		882 Matrix	17560 1966	80 520 Nonatelia	632
451 Muscpula	10986 1331	519 Nauclea	649 55	587 Nbnea	443
289 Jfuahroom	27153 3118	48 N. see Adlna	377 55	N. see Zyc6psis	444
i? M Usk Okro	17S74 2014	478 N. see Uncaria	3297 296	Nonesuch	18303 2061
519 5J Usk'ul ne*	19244 2133	365 522 Nauenburgio	2464 215	504 Nordantea	1544
61	638	ja3 Navelwort	1409 403	Norfolk Island Pine	
		429 Neckera	2987 24509 2881		
		K see Anomodon	2986 388	Norway Spruce	
		X. see DaltuntVi	2988		
		199 Nectarlna	12847 sy. 1499 5 525 Notelei		
		1P9 Nectarine	12847 1491 544 Nothochla;*na		
		47 Nrfmrt	3230 365 418	N. sec Cheilanthea	
		47	422		

GENERAL INDEX.

Lin.	Mat.	Sp.	Gen.	Lin.	Nat.	Sp.	Gtn.	Ijn.	Nat.	8 <i>j.</i>	Oci-
139	<i>Nithria</i>	9082	1099	311	<i>O. see Elclbtis</i>	2125		419	<i>O. see Allosbrus</i>	S9I*	
255	499 Notoceraa		1881	306	<i>Onobrychis</i>	18894	2102	423	<i>O. see Anelmia</i>	2947	
370	536 Notylia		2538	418	544 Onoclea		2912	424	<i>O. see Botrychium</i>	2^50	
167	Nyga	10806	1275	419	<i>O. see Lomkria</i>		2917	419	<i>O. see Lomaria</i>	2917	
400	Nummularia	24341	2787	282	511 Onbnis		1966	423	<i>O. see Lygbdium</i>	2946	
999	Nunfelia	24274	qy. 2778	331	521 Onopodium		2245	418	<i>O. see Struthiop-</i>		
817	498 A'phar		1557	331	<i>O. sec Brardia</i>		2246	tcris		S9I3	
404	Nutmeg	2829		330	<i>O. see Carduus</i>		2241	423	<i>O. sec Tbdea</i>	2944	
227	Nutmeg	14380	1623	53	527 Onosmodium		431	419	<i>O. see Woodwrdia</i>	9919	
287	502 Nutallwi		2005	54	527 Onysmodium		434	S64	522 Osteospermum	2450	
	Nut tree		2663	461	Onygcnia		3215	474	518 Ostericum	2SS*	
70	Nux-v6mica	4585	574	412	O'palus	25058	2867	106	Ostruthium	6928	870
530	NYCTACINEA, Or. 152.			445	Opegrapha		3113	383	535 Ciftrya	26<>	
5	525 Nyct&thes			3	Ojera Girls			10	Oswego Tea	628	74
70	Nyctelea	4556	568	41	520 Opercularia		328	532 OSYRnEAS, Or. KGL			
69	524 Nyctcrisftion		557	41	<i>O. see Cryptospér.</i>			396	532 O'tyris	2747	
74	527 NyctErium	tm		mum			329	183	Otaheite Apple		
217	498 NymphseVi			520	OPERCULARIEB, Or. 104.					11787	1405
217	N. see Nfrphar	1556		524	OPhioglossum		2951	173	Otaheite Chestnut	1351	
	497 29yunuuSxM, No. 9557			131	538 Ophiogbgon		1041	381	Otaheite Myrtle	2£#	
415	538 Nyssii	2892		66	519 Ophiophila		524	338	522 Otanthus	2306	
384	Oak	2666		132	Ophiocrodon	8592	1046	364	521 titohna	2451	
28	Oak Grass	223		413	525 Ophioxylon		2878	179	OtUes	11493	1368
97	525 Obesia	785		36	542 Ophiarus		278	15	512 Outea	109	
411	Obibne	24999 sy.	2862	36	<i>O. sec Pholidrus</i>		280	507 OxALIDZM, Or. 53.			
219	507 Cchna	1582		367	536 CTphrys		2488	185	507 CKxalis	141^	
170	O. see Glmpbia	1315		357	<i>O. see Chamrelis</i>		2489	184	<i>O. sec Biophytum</i>	141	
507	OCINA'CKJK, Or. 57.			370	<i>O. sec Corallorrhiza</i>		29W	357	Oxeyc	24CP	
269	502 Ochrbma	1921		368	<i>O. see Herminium</i>		2490	352	Oxeje Daisy	21643	2374
68	525 Ochrtsia	551		25	<i>O. see Ne6ttia</i>		2496	57	Oxlip	3775	45)
313	511 OYhrus	2137		92	530 Oplotliéca		736	175	Ox's Horn	11192	1567
260	499 Ochthbdium	1857		474	518 Op6panax		3329	324	Oxtongue	S210	
161	Octea	10393 sy.	1226	487	521 OIN>rnea		3397	110>			
144	OCTA'NDHIA, Cl. K.			1C9	O'puslus		7133	2015			
144	O. MONOU'MA, Or. 1.			195	517 Gpntia		892	81	519 Oxyiinthus	6*2	
158	O. DIGY'NIA, Or. 2.			438	Op&ntia	26438	1474	14	530 Oxybaphus	97	
108	O. TRIGY'NIA, Or. 3.			411	Orache		2862	404)	C24531	28*	
160	O. TETRAGY'NIA, Or. 4.			3H0	504 Orange tree		2187	4073	OxJcedruB	^24038	2837
372	536 Octomeria	2560		96	5& O'reba		784	155	523 Oxydceus	1193	
350	Cculus Christi			443	Orchall		3100	163	511 Oxylbium	12*2	
		21522	2345		536 ORCHIDEJE, Or. 188.			93	525 Oxyptalam	75?	
£40	529 O'cymum	1687		566	536 O'rehis		2481	140	531 Oxyria	110*	
240	O. sec Lumnitzera	1688		367	<i>O. see Anacamptis</i>		2486	94	525 Oxyst'lma	763	
240	O. sec Plectranthus	1689		366	<i>O. see Gymnadenia</i>		2480	305	511 Oxftripib	210†	
C8	Odllam	4463	550	367	<i>O. see Habenaria</i>		2483	464	Ozonium	325J	
		6885	8t1l	367	<i>O. sec Platanthera</i>		2478	337	521 Ozothamnus	2293	
249	Odontites	1576.1	1772	448	Oreads	20&6	3118	126	538 Pachidendron	10J	
430	Odontites	25477	2922	217	SCO Orelluna	15835	1565	175	496 Pachyneina	13?o	
h65	521 Cedew	2461		422	Orcopteris	25594	2938	316	512 Pachyrhlzus	Slig	
102	518 tffnlithe	833		107	Oreosclnum	6J40	873	379	533 PachysAndra	26W	
382	£40 (Enocarpus	2642		2J8	528 Origanum		1680	200	P.dus	12880	1#g
83	Cenöpha	5430	660	473	.018 Orlay/i		3313	71	520 Pa'dferia	577	
83	CE sec Berchemia	658		308	511 Ornioe&pxn		2113	9	528 Palavta	2019	
154	514 finothera	1183		162	.011 OrnUia		1238	83	508 Palicar/a	659	
415	Ogechee Lime			371	586 Ornithidium		1052	359	Palifurus	689	
		25216	2892	372	536 Ornithocéphalus		2549	392	Pallasio 22023 Scisy.	24-i	
415	Oil Nut	2893		134	539 Grnithogalum		1113	540 Valma Christi		2719	
357	Oil Seed	21899	2329	133	<i>O. see AstrolMrium</i>		2108	381	PALMC, Or. 201.		
243	Oily Grain	1716		134	<i>O. sec Eriospermum</i>		2107	433	Palma-Pmus 23282 i		
S99	Oily Palm	2777		134	<i>O. see Gheea</i>		2114	433	Palmella	S0V	
		503 OlkcisEM, Or. 26.		140	5C9 Ornithoglossum		1052	414)	Palmetto	J4IS }g	
14	503 CPLax	104		3C8	511 Ornithopus		2108	108	Pinacei	7059 #0	
287	O'lbia	17868	2008	308	<i>O. see Myriadenus</i>		2107	112	Pannx	2w»	
49?	Olden- .C5302&G	388		308	<i>O. see Myriadenus</i>		2114	117	P. sec Aralia	£JJ	
74J	1 a n d w ^ 1	4686	596	191	Ornithopus	12296	1460	117	Pancratium	Sn	
321	Old Man's Beard	2196		145	Ornf trophc	9349	&	475	P. sec Chliuithus	»§	
5	525 Clea	36			51 sy.	1157		117	P. sec Cobburghui	**2l	
67	Oleander	4367	529	12	525 O'rurus		83	117	P. sec Eikrycle*	^	
42	Oleaster	341		248	5S9 Orobanché		83	540 P. sec Ismtenc	**		
		524 OM^NA, Or. 124.		529	5ODANcnE, Or. 146.			S93	549 Pandanete, Or. 202.	^	
171	Olibanum tree	1333		313	511 O'ruber			24	Pandunus	*!%*	
5	Olive	36		314	<i>O. see Platystylis</i>				Panic Grass	£5j	
175	Olive Bark tree	1367		124	541 O'rentium			24	543 Panic Grass	6097 JIg	
102	518 Olivéria	828		124	<i>O. see Rhbdea</i>			36	Pinicium	TS	
85	Olive-wood	672		249	6rntium	15778	1774	34	P. sec Arundiniana	»g	
105	Glustarum	6826	854	184	Orristachys	11833 sy.	1409	25	P. sec C'ndon	29J	
198	515 OWnthia	1492		111	Orpine		904	25	P. sec Echin.chloa	1J	
377	543 Olyra	2606		184	Orpine	11841	1410	83	P. HCO Pnnlbttum	****	
390	533 Omalanthes	2707		15	502 Oritfia		114	80	P.BCTncholie'nii	,gS	
240	Online Plant	15263	1689	368	636 Orthoceras		2499	837	PanzenVi	15063 sy.	1^
391	533 Ompheala	2712		25	543 Orthopögon		189	81G	Papaver	JJS	
165	512 Omphalibium	1267		428	OrtMtrichum		2984	498	P. Bee Meon^pua	14'	
277	509 OmphaK.bium	1936		267	537 OrthroHftnthaloi		1IM8	401	ohOl		
M b'el	Oui>liu>bdtci	487		59	Orvala	14937	1665	397	*****		
		fil3 ONAUIIA'IILB, Or. 76.		139	543 Oryza		1102				
971	536 Oncidium	2540		23	5*2 Oryzopsis		177				
132	Onion	8601	1046	380	Osase Orange		2623				
838	Onites	15125	1680	154	515 Osb&ldVi		1181				
332	521 Onobrbma	2253		173	<i>O. see MeUistoma</i>		1356				
365	O. see Brftera	2470		435	OscilUtrbia		3037				
332	O. see CariowifziTs	2254		394	Oiser	24060	2732				
W2	O. see Heracantha	£256		559	5S2 Oraifste*		2420				
310	511 Onobrychis	2124		42J	^4* Osmunda		2945				

GENERAL INDEX.

567

Lin.	Nat.	gp. tien.	Lin.	Nat.	Sp. U.«.	Lin.	Nat.	Sp. Gen*
22	541 Papyrus	164	25	543 Penicillaria	191	27	542 Phalaris	216
51	Paraguay Tea	SMS 411	461	Penicillium	9245	27	P. sec Achnoddnton	214
192	Paraffias	12,193 1460	23	543 Pennisetum	173	27	P. see Aim	219
*»1	Pardalianches	215612353	259	PennyCrew	16336 1849	27	P. see Chilochloa	215
20	537 Pardanthus	149	235	Pennyroyal	14901 1659	27	P. see Ctfpsis	211
404	496 Pareira Brava Root	2825	233	P.....10yal	"74751 1650	27	P. see o/raphis	217
337	543 Portina	2680	IS	SSSJE	i m i 5	27	Phallus	5179
U1	534 Parietaria	2861	499	518 Pentacipta	3419	111	502 Pharnacetim	906
JS	512 Parinarium	1143	320	504 Pentadsma	2189	381	543 Pharus	2635
J60	figa pj^jg	1221	52	PENTA'DRIA, CL 5.		425	Phascum	2961
3 8	512 Partvda	2173	52	P. MONOGYNIA, Or. 1.		315	512 Phnseolus	2149
Jg	54* Parkrta	2916	93	P. DIGVYNA, Or. 2.		316	P. see Dulichoa	2152
?<2	512 Parkia	1939	109	P. THIGY'NM, Or. 3.		244	529 Phayl6psis	1726
lp	512 Parkinstialw	1273	112	P. TETRAOT'NIA, Or. 4.		229	Pheasant's Eye	
442	Parmelia	3094	112	P. PENTACY NIA, Or. 5.		14537	1634	
J2	501 Parnassia	910	115	P. POLYUV'NIA, Or. 6.		507 PhebMium	1312	
16	512 Parchetus	2156	285	Pcnt&petes	1997	418	Phegopteris	25378 2IK8
?1	516 Paroifrhia	728	241	Pentapharia	1700	518 Phelandrium	&8	
91	P.sec/Ulcerebrum	726	184	Penthorum	1411	103 P. BCC 3/^um	857	
?»1	516 PAKONYCIUB ^X J _E Or. 9883	4621	243	Pcntstemon	1717	23409	2666	
Jij8	499 Parrya	1825	338	Péntza	2311	515 PHILAOE'LPIEJS, Or. 83.		
47+]	Parsley	[ggó	13	534 Peperomia	94	515 Philadelphia	1479	
421	Parsley.	C2824 337	139	514 PdpUs	1100	525 Phill^rea	37	
Jg J	pier	19822 1173	191	P&plus	19349 1460	507 Philotheca	1933	
?<2	Parsnep	883	191	Pdplus	12358 1460	530 Philoxdrus	723	
67	525 Parsons/A	539	389	Pépo	23690 2698	541 Phlydrum	17	
261	5>2 Parthenium	2457	13	Pepper	93	Phlebia	3130	
*>2	Parthénium	21691 2375	438	Pepper Dulse	26403 3065	542 Phleum	213	
^7	522 PñcMia	2404	425	Pepper Grass	25713 2958	P. see Chilochloa	215	
23	542 Pásplum	178	235	Peppermint	14863 1659	528 Phl6mis	K>74	
?»	p. gee Crctsia	18*	198	Peppermint tree	238	P. sec Leon6tis	1676	
	Pasque Flower		82	Pepper Vine	5376 656	P. see Zcōcas	1675	
		14402 1629	262	Pepperwort	1873	526 Phl6x	472	
J^	532 Passerlna	1201	51	Perldo	3419 411	540 Phoenix	2745	
J2	p. see Daphne	1197	S51	Perdfcium	2355 138	521 Ph»16coma	2325	
J2	P. see Lachn^a	1202	521	P. sec Chectanthra	2349 202	536 Phohddta	2551	
TM7	p. see Stellera	i20f	350	P. sec Trixn	2442 465	542 Fholitirus	280	
I>	Passerlna	9765	363	Pereskia	1476 29	Phhma	3173	
S19	516 Passiflbra	1923	95	Pergularia*	774 241	539 Ph6rmium	1084	
*20	P. see Muruekia	1924	525	P. sec Vallaris	538 2	513 Photinia	liO!)	
270	P. see TVradilla	1925	67	Peribalia	209	327 Phragmidium'	3272	
oo»	PASSIFIAUEA, Or. 86.		26	Periccia	3200 123	543 Phragmtes	229	
?29	Passion Flower	1923	460	PericheVia	3202	528 Ph'yma	1697	
?J4	Ptfrc<<	23715v. 2699	79	Pericl^menum	5216 621	537 Ph^niun	5	
gg	Pasteque	23715*. 2699	235	Perilha	1660 472	Phullida	23420 26.16	
108	518 Pastinaca	883	461	Periola	3222 472	538 Phycella	J>70	
?Z*	P. see Op6panax	3329	93	P. sec Microloma	753 390	508 Ph^lica	710	
70	527 Patag. nult	565	290	Pcriptera	2022 391	P. see Soulangto	3	
267	537 Patersfrnla	1907	267	Peritoma	1905 89	P. see Staav/o		
I*	Patience	9188 1126	67	Periwinkle	528 390	Phyllanthophora	24559	
MI	Patientia	9188 1126	36	Pfrotis	283 100	533 Phyllanthus	2768	
41	520 Patrinte	330	161	Persea	1228 417	P. see E'mblica	2710	
215	P. see Ryaiva	1541	475	Persian Sun'6-eye v.	1017 439	P. see Khgnllia	1770	
165	Pauleitia	10G18 i'63	199	Plrsica	1499 390	P. sec Xylophylla	W)	
159	505 Paullima	1214	158	Persicaria	10232 1210	520 Phyllis	816	
49	520 Pavitta	382	150	Persdulta	9720 1173	Phyllitidis	25S46 2i68	
49	^m Pav4tta	3282	39	Petodma	13848 1571	Phyllitidis	26478 3078	
143	505 Pkvfo	1138	174	Perso6no	314 116	512 PhylloKibium	1!!90	
288	502 Pav&nto	2011	335	Petriblum	2282 391	539 Phyllobium	1021	
313	Pea	21&5	258	PetrocalHs	1845 390	Phyl6gyne	7570*^ 933	
199	Peach	1499	143	Fretocarya	9301 sy. 1143	527 Phfsalis	586	
201	Pear	12944 1507	37	PetrAphila	302 71	P. sec Nicändra	485	
52	Pearlwort	416	513	Petrosclnum	3130 78	Ph'sarum	3197	
101	Pecten	6563 819	247	Pdtrea	1755 417	Physospermum	876	
97	525 Pectinaria	788	529	Pfetribium	2282 391	518 Physostégia	S380	
352	521 Pectis	£868	335	Petcates	21104 2333	403 Phytelphas	2816	
?5*	P. aee Longchampsia	23G9	499	Petcates	21104 2333	522 Phycima	fill	
O40	526 PEOALINEJE, Or. 132.		511	Petalostcmum	1922 391	1901 'hyteuma	12242 1459	
**8	526 Pedalium	1760	512	Petasites	21104 2333	187 531 Phytolæcca	1420	
1S	f? ^edicularig	1781	513	Petiv&rtta	1136 390	531 PIYTOLAVBJE, Or. VA.		
?53	533 Pedianthus	1491	514	Petodma	13848 1571	97 525 Piaranthus	789	
188	PeijAa	2778 331	474	Pfetribium	2282 391	PicBua	36201	
238	u w E 2 * 16 0	^TS 331	474	Petcates	21104 2333	&Ssu	437	
22	524 P. wpm	1441	402	Petcates	21104 2333	Pichta	23631 2686	
J30	Pcfntinus	12890 12934	454	Petcates	21104 2333	509 Picrdmia	2765	
	Pcfntinus	8040 979	150	Pezza	9714 1173	521 Pierdiuum	2202	
132	S		305	Plrta	2100 322	521 Plcris	2210	
%4	Pcfhtory	10*1	511	P. sec Astragalus	2100 324	P.sceHelmfnthia	2216	
^	Pellitory	21691 2975	306	P. sec Cluttfidium	2089 327	Plcris	20255 S239	
***	Pellitory of Spain		304	P. sec Oxytropin	2101 329	511 Picti-tVi	2112	
240	«	«761 284	a25	P. sec Sphsrosph^a	2101 329	505 Pierardia	3S40	
oS	Pelfria	15845 v. 1775	509	PhacUia	477 308	Pigeon Pea	Clio i'R.J+	
g	499 Pelt&rtta	1844	470	P. sec Nemrphila	478 476	53* Pfca		
4 <i>i</i>	F- ^{8c} Heliophila	1901	458	Phaodium	3166 3166	Pfiewort	na\	
48	500	116. 3098	454	Phacorhlsa	27340 3138	Pillwort	2P58	
			180	Phacosplma	1450 3138	400 Pilobolus	SI 89	
			141	P. sceNollna	1122 3098	384 PloB^lla	19974 ii'21†	
						544PiJuaria	i<n	
						532 PitneR-a	«	
						515 Phnhta	145ij	
						103 518 Pimpinilla	?S	

GENERAL INDEX.

Lin. Nat.	Sp.	G	Lin. Nat.	Gen.	Lin. Nat.	Ep. Gen.
104 P. see <i>Solson</i>	84		163 511 <i>Podalyria</i>	124	317 512 <i>Pongamia</i>	2165
103 P. see <i>Tragium</i>	83		163 P. see <i>Baptisia</i>	1244	540 P. see Or. 199.	959
104 P. see <i>Trt nia</i>	84		282 P. see <i>Requiento</i>	1967	540 Pontederta	2494
387 Pinaster	23582	2685	162 P. see <i>Thermopsis</i>	124	368 536 <i>Ponthifcva</i>	2793
81 519 Pinckneya	644		96 525 <i>Podanthcs</i>	789	Poplar	
387 Pine	2685		488 522 <i>Podanthus</i>	341	216 498 Poppy	1552
118 Pine-apple	952		419 Pod-fern	291	401 535 <i>Pupulus</i>	2793
9 529 Pingufcula	66		388 535 <i>Podocarpus</i>	64	527 Porana	496
118 <i>Pinguin</i>	7*5 H5		350 521 <i>Podolepi8</i>	2348	1*. sec <i>Dinetus</i>	497
177 Pink	138d		488 P. Bee <i>Styl61epis</i>	3403	112 <i>Poranthra</i>	909
195 Pin Pillow	12617	1471	163 511 <i>Podolbium</i>	1247	216 497 <i>Porcupine</i>	C1696 186
387 535 <i>Pinus</i>	2685		497 PODOPHYLLA^CE*. Or. 7	251	1245 507 <i>Porlira</i>	12373 275
388 P. sec ifbl	2681		216 497 <i>Podophyllum</i>	1548	445 <i>Porlina</i>	3115
388 P. see <i>Cedrus</i>	2688		153 P. sec <i>Jeffersbnha</i>	1178	85 <i>Porophyra</i>	3061
387 P. see <i>Cunninghamia</i>	2688		142 531 <i>PtkMpterus</i>	1128	437 <i>P6rrum</i>	8602 1046
					5535ff. Portenschlagia	
387 P. see <i>Dhmmara</i>	2684		322 521 <i>Podospermum</i>	2200	519 Portland/a	672
388 P. see <i>Lkrix</i>	2687		95 Podostigma6178w.	772	75 P. see <i>Coutarell</i>	603
13 534 Piper	93		396 Poet's Cassia	2747	200 Portugal Laurel	
13 P. see <i>Peperomia</i>	9*		369 536 <i>Pogonia</i>	2505	5535ff. 12886 1502	
S39 534 PIPERAYEJJ. Or. 172	15164	1681	248 P. sec <i>Myfforum</i>	1761	Portland/a	
36 Pipewort	293		240 528 <i>Pogostemon</i>	1690	75 P. see <i>Coutarell</i>	603
477 511 Piptanthus	3343		167 512 <i>Poincidna</i>	1274	200 Portugal Laurel	
23 542 Piptatherum	183		167 P. see <i>Coultaria</i>	1277	189 516 <i>Portutica</i>	1447
337 521 Piquerw	2293		283 511 <i>Poiretto</i>	1977	189 P. see <i>Tallnum</i>	1448
111 <i>Piriqueta</i>	7247sg.	900	277 P. see <i>Hvca</i>	1940	111 516 <i>Portulacaria</i>	SU7
283 511 Piscidia	1975		121) Poison f 7878* 9	967	516 PORTULA^CEJE. Or. 89.	
1 304 P. sec DaubcnthmVi	2090		120j <i>Budb</i> I 7833	064	203 Portulacstrum	
, 263 Piscidium	16553	1873	70 Poison Nut	1585	13047 5y. 1517	
143 530 Pisbw/a	1135		110 Poison Oak	7170	80 519 <i>Posoquib-i'a</i>	635
397 509 Pistacia tree	2763		477 513 Poirea	3347	80 P. see <i>Gardenia</i>	688
374 Pistofchia	22860	2582	189 500 Polanisia	4485	52 541 <i>Potamogeton</i>	414
478 516 Pistorinia	3355		526 POLEMONIA^CEJE, Or. 134.	411 543 <i>Potamciphila</i>	2858	
313 511 Plum	2135		233 Polembnium	474	Potato	4706 591
313 P. see Othrus	2137		123 538 Polianthes	978	512 Potentilla	1528
118 539 Pitcafrne	954		233 Pblium	14717 1648	213 P. sec C'omarum	1527
404 Pitcher Leaf	24559	2830	4 516 Pollichia	22	383 512 <i>Pottrum</i>	2655
404 Pitcher Plant	2830		410 543 Pollinia	2843	41 541 <i>Pothos</i>	331
501 PittosrbKEf, Or. 24			319 POLVADE'LPHIA, Cl. 18.		42 P. see Symploc^rpus S&	
86 501 Pittsdporura	679		318 P. DECA'NDRIA, Or. 1.		Pofra 23691 26&	
' 48 526 Pladera	369		319 P. POLYA'NDRIA, Or. 2.		S89 Potiron 23691 2698	
285 502 Plagianthus	£002		215 POLYA'NDRIA, Cl. 13.		207 Pot Marigold	1580
296 Pjagiflobura	2057		215 P. MONOGY'NIA, Or. 1.		183 509 Poupart/a	1407
51 534 Planera	412		221 P. DIG^MA, Or. 2.		119 559 <i>Ponrrhita</i>	9&
385 Plane tree	2668		222 P. TRICY'NIA, Or. 3.		241 529 <i>Prasium</i>	1696
277 Plank Plant	17294	1943	224 P. TETRAGY'NIA, Or. 4.		369 536 <i>Prasoplflum</i>	2510
47 530 PLANTAUINEJJ. Or. 151.			225 P. PENTAGY'NIA, Or. 5.		245 529 <i>Prēmina</i>	1787
			226 P. POLYGY'NIA, Or. 6.		323 521 <i>Prcnathes</i>	2206
			326 POLYGLUM	3185	487 P. see Atalanthua	388
			327 P. see Crepis		357 P. see Lefatodon	2215
			328 P. see Harpalycce		324 P. see Prescttia	2574
			329 P. see Hrpalycce		373 536 Prescttia	
			330 P. see Prestonia		67 525 Prestonia	534
			331 Prickly Cedar		Prickly Cedar	519
			332 Prickly Grass		25 Prickly Grass	188
			333 Prichlyta		296 511 Prichtya	2C60
			334 Primrose		57 Primula	451
			335 PRIMLAVEJE, Or. 148.		57 530 <i>Primula</i>	451
			380 Prince's Feather		529 PRIMLAVEJE, Or. 148.	
					23239 2628	
					138 508 <i>Prlnos</i>	1089
					384 <i>Prlnus</i>	23431 2666
					244 <i>Prionitis</i>	15484
					77 522 <i>Prismatocarpua</i>	60S
					246 529 <i>Prwa</i>	1746
					246 P. see Strlptium	1747
					217 500 <i>Prickia</i>	1567
					379 534 <i>Prhri*</i>	2616
					217 Prophys 7644 sy.	
					37 514 Procrpinaca	
					168 512 Prosppis	
					240 529 Proftanth&ra	
					459 Prosthēmum	3175
					37 532 <i>Vxltca</i>	S0*
					P. see Isopbgon	
					P. 6ce Leucaden-dron	2749
					396 P. sec Leucoper-mum	
					38 P. sec Mimetes	3C5
					P. sec Nivtnwi	COS
					P. sec Petr6phila	S0*
					P. see Scrfria	307
					532 PhorKæte, Or. 160.	
					433 Protococcus	
					1'rotomina	
					Prunaster 26751	
					529 Pi unella	1694
					512 Prunus	
					P. see //mgdalui	
					P. see ilrménla	

Lin.	Nat.	gp.	G.n.	Lin.	Nat.	Sp.	Gta.	Lin.	Nat.	Sp.	GIB.
2007	P. see Cerasus	1502		480	512	Purshia	*Jg	147	Red Nightshade	9510	1173
480J	Ps&mma	210		54	P. see Onosmbdium	Jg	S94	Red Osier	2398*	2732	
27	Psammatis	11580	1388	189	Purslane	1447	890	Redoutca	2017		
180	Pseiid-ifacia	18773	2086		Purslane tree	967	284	Red Saunderg Wood	17723	1986	
304	Pseiul-Armēria	111			Puschkinia	Mg	168	Red Water tree	1285		
177	Pseiul-Armēria	11383	1386	397	Putrat'iva	24197	84	Red Wood	668		
335	PseiLdo-Acmēlla	20620	2281	238	Pycnanthemum	1679	285j	Ked Wood	J _{16 M 8 19 2 0}		
180	Psetudo-^tucion	11625	1383	236	Pycnatachys	166^	29	Reed	229		
COO	Psetudo-Census	12663	1502	200	Pyracantha	12895		Rccd-upon.Rccd			
		14700	1648	201	Pyrastrer	129^* v. laffl	1912		1912	231	
232	Psetudo-ChamaVpitys	19208	2133	451	Pyrénium	31^	486	Rcevesfo	3391		
312	Psetudo-Crēcca	19208	2133	441	Pyrnula	3(>0	503	Reichardfo	1276		
377	Psetudo-Cypfenw	2J012	2599	353	Pyrfethrum	2J75	S55	Rclham'a	2372		
265	Pseudo-Cytisus	16662	1888	354	P. see il'nthemia		22	Reniūria	167		
237	Pseudo-Dictamnus	15094	1672	523	P^rola	1349	372	Rcanthéra	2566		
264	Pseudo-ErucSstruin	16614	1883	415	P. see Chimaphila	1350	20	Ilenealmirt	1J8		
233	Pseudo>Ifysspus	14718	1648	201	Pyrularia	25219 sy.	282	Rcqquenfa	1967		
117	Pseido- C	7599	933	513	P^rus	1507	190	500Acseda	^ , 1459		
287	Psefdco-O'lbia	17863	2008	201	p'ee Amcttinchic	1502	500	7xSEI)AVE*, Or. H.			
324	Pseudo-Pilosella	19974v	2211	497	Pythagorean Bean		282	Rcstharrow	196		
412	Pseudo-Platanus	25061	2867	526	Pyxidanthera	460	541	REMTIAVEA, Or. 206.			
98	Pseudo-Pneumonanthe	6155	79*	458	Pyxidaria	16002	5*1	Rēstio	2742		
233	Pseudo-Scorodnia	14732	168*	532	Pyridifl	3286	396	R. sec Tliamnoch6rtus			
384	Pscido-Süber	23*23	2666	135	Quaking Grass	2a3	472	Rctantilla	32>8		
364	Psidādia	2454		170	Quash	8832	461	Reticulāria	3211		
197	Psidium	1484		507	(.) see Sitnartha	1317	460	R. see Aitfiorfdium	3203		
515	P. see Nēlitris	1485		116	Quillia	7590 sy.	64	Ketzia	502		
4	Psillurus	249 jy.	32	385	Querciton	23*^4	33	Rhabd6chloa	260		
424	Psilbium	2956		384	Cjuercus	2666	51	7fhacbm	S413	410	
316	Psophocarpus	2158		37	Queria	297	328	Hhagadolus	223f		
301	Psoralea	2071		91	Y. see Anfchia	2939	324	Rhagadolus	19965	2210	
283	P. see Cyamōpsis	1969		425	Quince	1508	412	Rhagbdia	2863		
283	P. see Dalea	1971		202	Quiniba	6477	508	AIIA'MNEJE, Or. GQ			
265	Psych'ine	1892		173	Quisqualis	1355	508	/ihamus	657		
80	Psychotria	631		518	Quisqualis	129ti	83	R. see Berchēmia	658		
82	P. see Geophila	651		506	Quisvia	3257	84	R. see feanuthus	(i)8		
80	P. sec Nonatilia	632		46*	Racodium	27532 3157	83	R. see Zizyphus	660		
80	P. sec Paicurfo	633		52	Radolla	419	413	Rhispis	2850		
511	Palium	3209	364	266	Radish	1897	360	Rhaptinctica	2*33		
355	Ptarmica	21796	2387	499	Raftia	1947	521	Rhapticum	10422	12S2	
49	Ptfel	391		511	R. see Vascha		214	Rhcēd/a	1533		
49	P. sec Ptelidium	39*		278	Ragged Robin	12018	161	531 /iheum	1232		
49	Ptelidium	394		278	Ragwort	1415	515 Tjhexia	1182			
420	Ptfcris	2926		512	Ragwrt	21161	174	R. see Acibitis	1362		
419	P. see Allosbrus			400	Rajflto	2786	174	R. see Acisanthfcra	1365		
421	P. see Cheilanthes			518	Rambutan	9360 sy.	477	R. see Chutogastrā	3345		
421	P. sec Lonicltis	J928		527	Ramuductai	24444	476	R. see Osbi'ckia	1181		
421	P. sec Lonchltis			397	Ramoont tree	2761	174	R. see Plrbm	1360		
418	P. see Nothochloe'na	2910		78	Itarapion	611	126	Rhinflnthus	1771		
418	P. see Tienitis	2909		76	" .	C4963	196	538 Rhipidodendron	1009		
284	Pterocarpus	1980		90	" .	607	517 Ripsahs	1475			
318	P. see Drepanocarpus			519	Rampion		505 RmzoBbLEX, Or. 46.				
318	P. see Ecastaph?llum	2IG8		519	Rampion	^ ss	505 Rhizobolus				
303	P. see Lonchocarpus	2085		519	Randi'a	6	461 Rhizoctbnia	3219			
469	Pterocephalum	3290		230	R. sec Ficaria		464 Rhizomfrpha	3255			
43	P. sec Cephalaria	345		264	R. sec Chompora		188 Rhizophrora	142U			
44	P. see Scabiosa	346		266	II. sec Enarthrocárpus		513 RmzoPiureik, Or. 74.				
•27	Pterogonium	2977		266	R. sec Goldbhchio		459 Rhizopbgon	3182			
429	Pterognbium	25912	299J	381	Kaphia	123Wsy.	490 516 RhckJola	3418			
ass	Pteronearon	1830		260	Rai-hi61cpis	1510	523 y^hodod^ndron	1339			
521	Pteronia	2261		260	Rapitrum	1860	438 Rhodbmela	3064			
332	P. scCStaehellina	2259		260	R. sec Ochthfriium	1857	171 523 7thod?ra	1338			
285	Pterosplrmum	2000		76	Apticulus	4963 607	40 532 Rhopilla	323			
356	Ptcrl6tylii	£518		472	Raspal'a	330H	161 Rhubarb	1232			
366	Ptcrygbdium	2474		212	Raspberry	13505 1523	110 509 fihls	894			
356	Ptilostephium	2397		100	Ratanhia Root		277 R. see Omphalibium	1936			
438	Ptiba	3063		424	Rat Poison	6558 815	314 512 Rhynchbsia	2145			
47J	Ptychbtis	3315		295	Rattany		21 541 Rhynchospora	154			
445	Puccinii	3271		295	Rattlesnake Fern		458 Rhtfsma	3165			
SIR	Puccoon	1547		295	R. sec Brassica		90 517 Kibes	719			
483	Pueraria	1970		295	R. sec Chompora		47 Rib Grass	3160			
460	Puff Ball	S191		295	R. sec Enarthrocárpus		432 Riccia	3000			
235	PuU	14901	1659	71	R. sec Goldbhchio		139 Rice.	1103			
350	Pulicaria	21512	23*5	118	Rauwolffo	18263 2053	143 541 Kichfrdh	1142			
55	Pulmonaria	438		225	RavenMa	579	541 R. scCRichardsonw	1095			
227	PulsatUla	14-102		161	7663 sy.	433	520 Richardson'a	1095			
164	Pulten <i>vi</i>			404	Reaumima	1607	188 500 Richiea	1441			
J25	Pūmos	8172	U96	110	Rattlesnake Root		533 ficiinus	£719			
398	Pumpkin	2J(i90 2698		110	Red Bay	10396 1226	257 499 Ricott'a	1834			
					Red Cedar	24521 2822	269 50r Riedle/a	1919			
					Red Gum tree	12787 1493	54 527 Rindcra	436			
					Red Lac	7164 89*	410 53* Ripedim	28*2			
							131 538 Ripogoumum	1037			
							365 Tfiro	22408 2468			
							42 5S1 Rvtaa	334			
							533 Rivularia	3022			
							433 R. see Corynlpbora	3021			
							434 R. sec Mcsoglbia	3034			
							183 509 Iloblrgia	1408			
							328 521 Roblrrta	5223			

Un.	Nat.	Sp.	Gen.	Lin.	Nat.	Sp.	Qen.	Lin.	Nat.	Sp.	Gen.
304	511 Robin/a	Sp.	2086	852	528 Russelia	Sp.	1798	159	505 Saponaria	10342	1218
304	R. sec <i>Caraghna</i>		8091	169	507 J?uta		1309	69	Sapta	4503	#1
304	R. sec Halimoden-		8098	477	R. see <i>Aplophyllum</i>			524	Sapteas, Or. 121.		
303	R. see <i>Lonchocar.</i>			507	RVT^CBX, Or. 55.			167	Sappan	10810	1875
	pus		2085	420	iZuta muraria			72	Saracha		
162	R. see <i>Ormosia</i>		1238					372	Sarcinthus		
SOB	R. see <i>Pictetia</i>		8112	86	504 Royschia		688	394	Sarcopanos		
317	R. see <i>Pongdmia</i>		2165	815	500 Ryanea		1541	285	Sarcocatilon		
303	R. see <i>Tephrosia</i>		2083	35	Rye		873	82	Sarcocephalus		
162	R. see <i>Virgflia</i>		1242	35	Rye Grass		871	370	Sarcophilus		
385	Rbbur	83464	2666	437	Rytiphalafr		3052	378	Sarcococca		
132	Rocambole	8595	1046	141	540 Subal		lli^4	94	Sarcobatus		
443	Roccua	3100	59	526 Sabatia		468	279	511 Sarcophyllum			
SfaO	Rocket	1863	889	527 Sabdariffa	17987	2014		91	Sarcostemma		
865,L	Rocket		1887	36	543 Saccharum		884	216	528 Sarraadiua		
819	Rock Rose	1589	36	S. sec Erianthus		885	498 SARHACEMW, Or.	498	SARHACEMW, Or.	10	
63	Rock Rose	4233	492	36	S. sec Imperita		287	399	Sarsaparilla	24295	#7&3
434	Rock Violet	26169	3027	410	S. sec Rhipidium		2842	293	Saidqua	18164	2"*
				36	S. sec Tricholatia		886	161	Sassafras tree	10403	1\$
■ 521	Rodigio	2225	36	S. sec				2.3	528 Satur&t;		
370	Rodriguezes	2521	826	Sacred Bean		1613	366	536 Satfrium			
79	Roella	613	16	Saffron	1024	116	366	S. sec Gymnndenia			
77	R. sec Campanula	607	10	Sage		76	367	S. sec Nigellella			
124	541 Rohdea	991	52	502 Sagina		416	367	S. sec O rchis			
365	Koliidra	2469	383	536 Sagittaria		2653	225	503 Saurauja			
816	498 Homer/a	1550	381	Sago Palm		2637	144	534 Saururus			
75	519 RondeKtfVs	602	381	540 Sagus		2637	329	521 Sauvarea			
396	Rope Grass	8742	893	St Andrew's Cross			87	501 Sauvagdsta			
806	513 Ausa	1522		310	Saintfoin	2124	404	404	Savin	84515	88A
	iiosa:A:KJE, Or. 68.			306	Saintfoin	18894	803	803	Savory		log
889	Rbsa. malabarica	17978	2014	414	St John's Bread	2102	264	264	Savory Cabbage v. 3.	188?	
				320	St. John's Wort	25130	2885	136	Savory Spiderwort		
888	JZha sinensis	17943	8014	320	St. John's Wort	2190		329	Saw-wort		
8	537 Roscea	7	320	123	St Martin's Flower			175	517 Saxifraga		
808	Rose	1522		80	St. Peter's Wort	O23		175	Saxifrage		
304	Rose Acacia	18782	2086	80	St. Peter's Wort			103	Saxifrage	6716	**>
155	Rose Bay	10048	1186	80	8042	979		517 SAXIPRA^GE, Or. 96.			
186	Rose Campion	1415		80	8042	979		43	Scabibsa		
10	103) Rosemary	75		80	8042	979		43	S. sec Cephalaria		
109j	Rosemary	f ^ 835		80	8042	979		469	S. sec Suctisa		
186	Rose of Heaven	12009	1415	80	8042	979		43	Scabious		
860	Rose of Jericho	IB58		80	8042	979		79	522 Sca/vola		
893	Rose of the World	v.	44 20-8	80	8042	979		350	Scalia	21547 sy.	9Z»
				80	8042	979		329	Scallion	8619 v.	1^
490	Rose Root	3418		80	8042	979		63	Scammbnia	4184	^
109	Rose Snowball tree	7133v.	892	80	8042	979		63	Scammony	4184	490
10	528 Jtosmarinus	75		80	8042	979		101	ftandix		
31	543 Rostraria	842		80	8042	979		101	S. sec ifothriscus		
884	512 libithia	19S2		80	8042	979		101	S. sec Cha?ophyllum		
36	542 ltottboell/rt	279		80	8042	979		102	S. sec TorlUs		
4	R. sec Monerma	32		80	8042	979		323	Scarcola	19879	22" I
36	R. sec Ophiurua	278		80	8042	979		129	Scarlet Pomponc		
402	533 Rottlera	2810		80	8042	979		850	Sc^ptrum Carol.	8413 10i"	
101	Rough Chervil	820		80	8042	979		380	Sc^ptrum Carol.	1582 1781	
40	Ruujo/a	2697 *y	323	80	8042	979		382	S. sec DuvaGfl		
144	538 Roxburghita	1149		80	8042	979		32	543 Schismus		
161	Royal Bay	10388	1226	80	8042	979		425	Schist6stega		
175	524 Royena	1373		80	8042	979		857	Schivereckia		
483	528 Roylea	3376	6	80	8042	979		423	544		
81	Royoc	5342	648	80	8042	979		380	497 Schizandra		
85	Rubentasy.	672		80	8042	979		8	528 Schizanthus		
86j	Rubentasy.	jKg 686		80	8042	979		433	Schizon&ma		
45	519 Jlubia	349		80	8042	979		254	499 Sibopralton		
519	JUBIA*CE», Or. 103.			80	8042	979		452	Sibopralton		
811	512 JZbus	1523		80	8042	979		452	Schizophrillum		
358	582 Jlubc?cks	8411		80	8042	979		352	Schkbhria		
488	R. sec Echinacea	3408		80	8042	979		13	542 Schmideba		
317	512 Rudolphia	2163		80	8042	979		450	Schm(dtm		
169	529 Rue	1309		80	8042	979		596	SchcpAnthus	24916	254*
244	529 Ruellia	1727		80	8042	979		132	Schoenhdum		
845	R. sec Aphelandra	1731		80	8042	979		21	541 Schu'nus		
845	R. sec Crossandra	1733		80	8042	979		81	& see Cjaet6spora		
845	R. sec Hygrphila	1728		80	8042	979		12	S. sec Cladium		
381	RCifU	23280	267	80	8042	979		377	5. see Kobrestra		
891	503 Ruzia	2025		80	8042	979		SI	S. see Scfrpus		
114	503 RulfngTi	927		80	8042	979		145	505 SchbfVi		
189	R. sec [^] nacampseros			80	8042	979		165	S. see Omphal6bium2		
				80	8042	979		165	S. see SchbfVi		
				80	8042	979		139	519 Schrädera		
141	531 itdmex	1126		80	8042	979		406	512 Schrfinkia		
142	R. sec JTmor	1127		80	8042	979		406	512 Schrebcr		
140	R. sec Oxyria	1104		80	8042	979		388	535 SchubfTfn		
109	518 Jtumia	888		80	8042	979		104	518 Schultzia		
105	R. sec Capnophyllum			80	8042	979		118	Schwagrichema		
52	541 Ruppia	853		80	8042	979		8	588 Schwtnckra		
100	Rupturewort	813		80	8042	979		177	8042	7657 sy.	9g
403	538 /tuscus	819		80	8042	979		502	Saponaria		
124	Rush	994		80	8042	979		502	Saponaria		
323	Rush	19896	8205	80	8042	979		502	Saponaria		
164	Rush Broom	1254		80	8042	979		502	Saponaria		
22	Ruth Nut	1460	163	80	8042	979		502	Saponaria		

GENERAL INDEX.

571

Nat.	Sp.	Gen.	Lin.	Nat.	Sp.	Gen.	Nat.	Sp.	Gen.
135	539 Scilla	1054	246	529 Selago	1742	21K	502 Silda	2023	
J36	& see ^a 'HIum	8677	1046	155	Selas	10070 iy.	230	S. see A'node	2021
	s. see Anthericum	1061	241	Selfheal	161H	5	5 see Lopimia	2018	
	S. see Bellevallo	1055	106	518 Sellnum	873	290	S. sec Pemptera	2022	
155	s. see Ormthgalum	1053	104	S. see ^a fbon	842	336	Sideranthus		
	S. see Uropetalon	1079	105	S. see Cnulum	860		21869 ft 70 ty. S'39		
518	Sciadophyllum	913	474	S. see Ligisticum	875	234	528 Sideritis	1657	
541	Scirpus	158	474	S. see Melanosellnum		236	S. see Stichyg	1669	
81	S. sec Eleocharis	159	474	S. see Puccedanum		49	520 Siderodendrum	384	
	S. see Isdlepis	157	474	S. see & seli	882	69	524 Sideroxylon	558	
J1	537 SCITAMINEA, Or. 184.	7*1	76	104	S. see Thyssclinum	872	S. see ^a rcula	559	
175	Scleranthus	1375	106	S. see Selloa	2371	70	S. see ^a umella	555	
359	522 Clerocarpium	2422	382	509 Scraecarpus	2897	415	S. sec Mangliula	570	
&	543 Sclerethloa	257	415	516 Scmpervvum	1468	70	S. sec M\$nvic	2891	
f60	Scleroderma	3190	193	S. see Monanthes	1469	216	S. see Seralisia	569	
164	511 Sclerothtanus	1259	194	S. sec Sedum	1410	356	Sidesaddle Flower	1555	
461	Sclerbium	3225	184	84	501 Senacia	665	522 Siegesbeckta	2398	
£20	544 Scolopendrium	2924	85	S. see Elseode'nruin	672	S. see Verbesina			
<77	Scolopendrium		262	521 Senebiera	1872	215	Sieviersia	1537	
		17294	249	349	Sene <i>c</i> Uis	2342	518 Silixus	3328	
328	521 Scolymus	2236	344	521 Scnecio	23J5	105	Siliatis	6852	
388	Scolymua f (20415	2479	2247	295	Stnega	18263	179	Silene	1388
47	528 Scoparia	362	166	Senna	10702	474	Slier	3327	
61	527 Scopolia	490	418	Sensitive Fern	25402	518	SiliKus		
86	S. see Toddhlia	686	405	Sensitive Plant		1041	1041		
232	Serdium	14706	1648	24608	2834	108	Slier	16812	
234	Scordbtis	14811	1654	Scpdnum	3251	109 J	C7107	881	
933	Scorobnia	14731	1648	516 Septas	1147	165	Siliquistrum	10G31	
132	Scorodprasum	8595	1046	214	Septfoil	1329	Silk Cotton tree	12H	
280	Scorpion	17485	1963	459	Soptoria	3170	409	Silk tree	24818
53	Scorpion Grass	425	368	536 Sapri/M	2491	364	Silver-weed	28J7	
307	Scorpion Senna			S. see Cephalanthera		330	Slybum	2242	
		18987	2106			170	Simuba	1318	
						507	Simarba	1317	
						507	Simarub&cea, Or. 56.		
511	Scorpiurus	2105	369	S. see fpipactis	2508	359	Slmefia	2419	
	Scrprius	17485	1963	159	505 Seriana	1215	488	S. see Coreopsis	2415
322	Scorzonh-a	2201	86	Seringt'a	682	351	S. see Ximcnnesia	2358	
322	5. see Podospermum		49	S. sec Ptelidium	394	264	599 Sinapis	1884	
140	Scotch Asphodel	2200	328	521 Scrlola	2224	266	S. sec Didesmis	1898	
281	Scotch Laburnum	17520	328	S. see RidiRia	2225	266	S. see Diplotifxis	IKH	
			365	SeriSubi	223	389	S. see Eructria	1901	
			80	520 Scrissa	623	Single-seeded Cu-			
			574	Scrpentia	S2853	cumber	2700		
278	511 Scettflo	1945	350	Scplcula	2622	523 SimnRivz	1701		
393	Screw Pine	27*:	239	514 Scrlcula	2622	247	Siphonanhlsus lf^73	1752	
285	Screw tree	1993	47	511 Serraria	15147	392	Siphonia	2720	
S51	528 Scrophularia	1790	329	Seringt'a	1681	403	Sribba	833	
	528 SCHOPHULARIN.B., Or. 141.		329	S. sec ferratula	3197	Sir Joseph Banks's	93		
900	Scrubby Oak	12892	329	S. sec Ccstrlnuf	2209	Pine	24507		
42	Scunkweed	2803	38	S. sec Crtsium	2238	2820	Sfsarum	6799	
958	499 Seurvy, Grass	1848	5.32	S. sec SausBurca	2240	848	Sison	849	
240	529 Scutellaria	1693	70	Scrfuria	307	5. see C&rum	859		
233	S. seq TctLcrhtm	1648	524 Sersalis'a	569	5. see Cherophallum	821			
321	516 Scyanthus	2195	201	Service	12975	S. see Helosciadium	3316		
			526	Or. 133.	1507	101	S. see Ligusticum	875	
			243 B26			473	5. see Pctrosellum	3.30	
			304	511 Scsfrnia	2087	107	S. see Schfritz/a	850	
			304	S. see Aguti		474	5. see Smf rnum	854	
			301	S. see Glottfdiura	2189	105	S. see Jragium	837	
			104	518 Selseli	844	317	Sissoso	19558	
			103	S. see JTubon	842	453	Sistotrfema	S12)	
			473	S. see Ptychjtitis	3315	261	5. see SifSfmbrum	1865	
			107	S. see Sellnum	873	260	S. see Andreoskia	1864	
			29	543 Scslria	233	255	S. see Barbar^a	1822	
			34	S. see Echinaria	267	104	S. see Z:ras*ica	1883	
413	540 Sea6rtia	2881	203	Seshyfum	1517	264	S. see DiplotAlvi3	1886	
139	Sea Heath	1099	24	543 Setaria	187	265	5. see Leptahum	1868	
101	Sea Holly	6612	183	Sethia	1401	262	S. see Leptocarpac'a	1820	
			850	Seymeria	1780	254	5. see Nasturtium	1819	
			528	Shaddock	19669	255	S. see Andreoskia	1864	
			320	Shuttleton	2187	104	S. see Z:ras*ica	1883	
			172	Shallot	8619	104	S. see DiplotAlvi3	1886	
			132	C 24331	1046	104	5. see Smf rnum	854	
			404	124638	2837	104	S. see Brignblia	3J17	
			407 j			473	S. see Drepanophyl-lum	a318	
			171	Sheep Laurel	10995	473	S. see Hclosciadium	3310	
			90	Sheep's Scabious	1335	473	S. sec Pastinaca	883	
			142	Sheen's Sorrel	1126	108	S. sec Sison	849	
			397	Shepherd/a	2754	104	Skinnera	10081	
			322	Shepherd's Beard	2199	155	1188	ty.	
			60	Shepherd's Club	3940	104	Skirret	6799	
					480	193	Slipule	848	
			2581	Shepherd's Purse f }	1^	193	Slipw>rwort	65	
			2593	Sheridfti	351	503 Sloanea	1570		
			45	Shield Fern	2938	199	Sloe tree	12852	
			422	Shore-weed	2621	188	Small Bur	1501	
			380	Shrubby Trefoil	391	380	Small Burdock	1213	
			49	Shuttlecock	18032	238		1443	
			114	SibbeldVi	924	25187	2627		
			201	Siberian Crab	12960	3	Small Cardamom	58	
			304	Siberian Pea tree	2091	22!	S ^{H!!} ! t. P ^{ine} 18631	13	
			252	Sibthorpja	1792	238	Small Monarda		
			389	Sfcyos	2700				
				S. sec Sschium	2702				
			902						

Lin.	Nat.	Sp.	Gen.	Lin.	Nat.	6p.	Gen.	Lin.	Nat.	Rp.	(ieii.	
239	Small Peppermint			SO	Spanish Nut	1327	143	472	508	StadyM	3.W	
	15164	1681		16	537	Sparaxis	122	463		Stachylidium	3241	
	516	Smeathmähn/Vi	1616	375	540	Spargnium	2598	236	528	Stachys	1669	
	538	SMILA^CBJE, Or. 194.		333	522	Sparmannophorus	2267	237	9	S. sec Sphäcele	3375	
131	538	Smilacina	1039	218	503	Sparmanni <i>Ti</i>	1572		S. see Zielenfia	1670		
599	518	Smilax	27a)	157		Sparrow-wort	1201		S. see Zietenfia	68		
308	511	Smithia	2116	1511		Sparrow.	T 9765	111	533	StackhoiWa	898	
105	518	£my>nium	854	157 J		wort	110176	1200	533	STACKHOUSIVP, Or.	168.	
105		S. see Thildium	860	23	542	Spartina	174	155	505	Stadmännm	1192	
107		S. see /igubiticum	875	280	511	Spartium	1962	332	521	Stshellna	2259	
108		S. see Pastinaca	883	281		S. sec Adenocarpus	1965	352		S. see Lef sera	2370	
474		S. see Physosperaum	876	281		S. see Ciftis	1964	83		Staff tree	662	
473		S. see Zizia	3319	280		S. see G'cnfsta	1961	443		Stag's-horn	26755	
315		Snail Flower	19414	2149	22	Spa rt urn	1503	168	489	Stanhbpc	28532 sy.	
388		Snake Gourd	2694	485	537	Spatalanthus	3388	262	499	Stanl'ya	3412	
66		Snake-root	524	39	532	Spatalla	310	95	525	Stapèlia	779	
374		Snake-root	22853	2582	111	Spathelia	896	97		S. see CaraUhma	792	
131		Snake's Beard	1041	242	526	Spathodea	1708	97		S. see Caruncularia	787	
423		Snake's Tongue	2946	454		Spatularia	3136	97		S. sec Duval'i'a	786	
159		Snakeweed	10286	1210	230	Spearwort	14596	1637	96		S. sec Gonostemon	781
395		Snake wood	2733	77		Speculum	£060	608	97		S. sec Hulrnia	790
84		Snake wood	5496	668	6	Speedwell	45	97		S. sec Obesia	785	
549		Snapdragon	1774	34		Spelt	2285	269	96		S. see O'rbea	784
180		Snapdragon	11558	1388	187	Spērgula	1418	97		& see Pectinaria	788	
9		Snap tree	544	58	183	Spērgulastrum	1402	97		S. see Piaranthus	789	
3.05		Sneczewort	21796	2387	46	Spennai-bee	302	96		S. sec Podanthes	782	
109		Snowball tree			46	S. sec BigelvzVz	355	96		S. sec Trid^nteal	780	
			7133 v.	892	49	S. sec Hedjbtis	388	96		S. sec Trom6tnche	7H3	
80		Snowberry	628	46	S. sec Knöxta	354	223		Staphisigna	14107 l>99		
119		Snowdrop	961	74	519	Spermadictyon	595	111	508	StaphyRa	897	
£.8		Snowdrop	14427	1629	15	Spermamaxyon	105	99		& sec Bumaldfl	804	
188		Snowdrop tree	1432	■		SpermceMia	3223	69		Star Apple	&->6	
119		Snowflake	962	483	528	Sphäcele	3375	96		Starfish	6230 779	
150		Soap-berry	1218	437		Sphacellnria	3054	356	521	Stärkea	2390	
177		Soapwort	1385	365	521	Sphaeranthus	2467	134		Star of Bethlehem	1053	
203	499	Sobolewskia	1880	456		-Sphagna	3157	47		Star of the Earth		
99		Söda	6458	808	460	Sphaerobolus	3188				3200 364	
410		Soft Grass	2854	432		Sphaerochrypus	3003	■		Star Thistle	22189 2434	
316	512	Sbjä	2151	460		S. see Trichia	3204	346		Starwort	2337	
71	528	Soländra	582	438		Spha?rococcus	3066	115	530	A'fatice	92!	
	527	SohA'NK/e, Or. 140.		164	511	Sphaerolbium	1255	114		S. see Armēria	928	
72	527	Solanum	591	458		Spharonēma	3169	280	511	Staurocanthus	1961	
72		5. see Lycopersicum	590	444		Spharophoron	3106	223		Stavesacre	14167 1599	
74		S. see Nyctériura	592	305	511	Sphrosphysa	2096	419		Step'nia	2917	
57	530	Soldanelia	453	425		Sphagnum	2962	373	537	Stella	2572	
«4		Soldandlia	4255	494	360	Sphenbgync	2430	180	502	Stellaria	1389	
328	521	SolddevilUs	2226	471	523	Sphen6toma	3294	1872			1417	
4(15		Soldier Wood	24600	2833	108	Sphondflium	7053	478 J		S. see Larbrea		
89	501	Sflea	703	103	367	Spider Oprhys	22520	157	532	Stellem	1200	
		S. see Ionfdium	705	125	367	Spiderwort	1000	417		Stcmaria	25316 2902	
80		Solena	5H78&9sy.	635	246	Spielmänne	1748	252	528	Stemodia	J79J	
437		Solēma	3059	61	526	Spigelflirt	484	461		Stcmonitis	W	
348	521	Sohuago	2339	103	526	Spicncll, see Spick.	835	65	523	Stcnanthera	2io	
347		S. see ^ster	2387	103	527	Spiñacia	2768	248	529	Stencarpus	248	
338		S. see Caleacte	2307	nell	527	Spinach	2768	117		Stenomlson	7651 sy. 9W	
337		S. see Euthāmia	2296	SI	527	Spike Rush	159	368	536	Stenorhifinchus	2497	
364	522	SopIva	2453	335	527	Spilanthes	2281	139	501	Strphnra	1101	
131		Solomon's Seal	1040	357		S. see Acmllla	2402	S92	503	Sterculia	2724	
440		Solorina	3084	335		S. see Petrōbium	2282	444		Stercoa6lon	3105	
322	521	Sönchus	2203	336		S. sec Platyptrcis	22K5	263	499	Sterfgma	1882	
487		S. sec Agathjrsus	3394	439		Spiloma	2768	123	538	Stcrbergia	973	
322	5. see Picerdium	2202	398		527	Spinach	2768	255	499	Stevénia	1823	
261		Sophia	16463	83	527	Spinach Christi	5424	334	521	Stevill	2273	
162	511	Sophbra	1237	398	527	Spinacia	2768	335		S. see Hymenopippus		
162	5	see Cyclbia	1243	84		Spindle tree	666				2276	
162	S. see Edwards/a	1235	202	527		S' sec Gilttnia	1516	443		S. see Palafoxia	22IO	
162	6	see Thermopsis	1241	203		Spiralepis	2315	455		SMcta	3097	
162	S. see Virglia	1242	340	527		Spiranthera	678	434		Stictis.	3148	
370	536	Sophronitis	2553	86	507	Spirhntics	sy. 2496	396	534	Stigonema	3030	
201	536	S6rbu8	12975	1507	368	Spirchnum	969	316		Stilago	2746	
410	543	Sorghum	2852	426		Splachnum	2968	464		Stibuspora	326B	
490	509	Sonnea	3421	426		S. sec Dfssodon	2968	iGi		Stilbum	3236	
39	532	Sorocphalus	309	419		Spleenwort	2921	390	533	Stillingia	2706	
		Sorrel	1126	183	509	Spöndias	1405	237		Stinking Horchound	-1671	
			1414	183	5. sec Buchanan/a	140G	25	542		Stipa	194	
172	508	Somed tree	11055	1341	408	Si>ong tree	24801	47		Stishwort	3195 36*	
472		Soulang	3311	464		Sporidermium	3265					
292		Sour Gourd	18140	2028	26	Sporobolus	3265	180				
415		Sour Gum	25212	2892	439	Sporochnus	5073	316				
226		Soursop	14363	1622	463	Sporotrichum	3249	332	521			
339		Southernwood	20861	2313	12	Sprengelfa	509	487				
51		South Sea Tea	3429	411	461	Spring Grass	90	254	499			
392		Southwells			461	Spumaria	S209	365	521			
			23872 w	2724	190	Spurge	1450	341				
105		Sbwa	6846	858	156	Spurge Laurel	10138	332	521			
132	539	Sowerb <i>ai</i>	1045	89		Spurless Violet	2197					
322		Sowthistle	2203	187		Spurrey	1702	184				
316		Soy	2151	105		Square Parsley	6840	173				
106	518	Span6nthe	865	398		Squash	23694	271				
280		Spanish Broom			135	Squill	60	111				
			17468	1962	135	Squilla	8810	402	556			
263		Spanish Cress	16549	1873	389	Squirting Cucumber	1053	556	5triatutes			
69		Spanish Elm	4517	563		Squirt	23685	293	515	Stravadiūm	204C	
						Strawberry	2697	212				

'in. Nat.	Sp.	Gen.	Lin. Nat.	Sp.	Gen.	Lin. Nat.	Sp.	Gen.
Strawberry Blite	29		42 541	Symplocarpus	332	106 518	Tenor/a	862
172 . Strawberry tree	1:345		523	SYMPLOCINEA, Or.	118.	103	T. see Crthmmum	834
-Si 537	Strelitzia	747	319 52;	Symplocos	2186	303 511	Tcpbrsia	2083
Streptanthra	3281		357 522	Synedrella	2400	314	T. see Sweetta	2141
Streptium	1747		321	SYNGENE'SIA, Cl. 19.	.	284 512	Teramus	1983
468 537 Streptocarpus	8279		321	5 JEQUAYIS, Or. 1.	.	509	7EREUNTHACEJE,	
246 529 Streptopus	10:34		338	S. SuPE'BFLirA, Or. 2.	.	Or. 66.		
J31 538 Striaria	5077		358	6 FRUSTRA^NKA, Or. 3.	.	397	Terebinthug	24216 2763
439 Strbus	23604 2685		362	& NECESSARIA, Or. 4.	.	412 513	Terninlia	2864
387 Strophanthus	546		365	S. SEGREGATA, Or. 5.	.	302	Ternatea	18688 2073
68 525 Struchium	20512 2267		18 537	Synnbfto	129	217 503	TernstrceWrt	1559
333 Strumpuria	96:3		188	Syrian Rue	12147 1444	503	TERNSTRIKMldctP.	
120 538 Strumarium	23187 2627		I 525	Syringa	[U*J	Or. 34.		
380 Struthiola	327		480 515	Syzygium	3365	400 538	Tcrpanthus5602^y.	678
Struthiopteris	2913		61	Tabdeum	4050	378	Testudinana	2785
Struthium	11341 1384		68 525	Tabernsmontana	548	32	Tita	8588 1043
STRYCHI.VEJE, Or. 125.			68	T. see AmsumVi	541	225 496	Tetracra	1609
Strfchnos	574		171	Tnrmnn	fl 01**3 MM	219	T. see Dellma	1583
Stuart/Vi	2034		218	hacd	C243i)8*y.2793	219	T. see Doliocarpus	1584
S. see Malachoden.			401	Tdcca	993	597 507	Tetradium	279
dron	2035		386	FacetedTrum	5201 2628	254	TETRADYNA'MIA, Cl. 15.	
95 525 Stylandra	772		71	Tachylgia	1289	203 517	Tetragonia	1519
STYLIDEIE, Or. 100.			168	T. see Calandrfmifl	3357	301 511	Tetragon6lobus	206!
373 522 Stylium	2581		270	Tacsbia	1925	351 521	Tetragonotheca	2^57
488 521 Stylilepis	3403		517	fte'da	23593 2685	236	Tetralit	149." 9 166n
Stylosanthes	1979		418	Tsntis	2909	151	Tetralix	9782 1173
136 539 Styphandra	106*		387	Tdgera	10713 1271	49 520	Tetramerium	380
fit 523 Styphelia	505		352	Tg*ghc*	2365	2	TETRA'NORIA, Cl. 4.	
S. see Cyathodes	519		521	Taliera	997	37	T. MONO'Y'NIA, Or. I.	
65 S. see Leucopbgon	518		540	Taliera Palm	997	51	T. DIGY'NIA, Or. 2.	
65 S. see Monstoca	515		270	Taligulia	15529 sy. 1730	51	T. TKTRAGY'NIA, Or. 3.	
65 S. see Trochocarpa	513		516	Tar.num	1448	161 532	Tetranthera	1227
173 SJ STYRACNEE, Or. 119.			189	Tarnum	10713 1271	532	Tetraithus	2394
Styrax	1352		189	T. see lnacampseros	1449	426	Tetraphis	2967
Saber	23422 2666		478	T. see Calandrfmifl	3357	35	Tetrapbgon	274
499 Subularia	1903		145	Talisia	1163	505	Tetn'pterus	1397
520 Succisa	3288		505	Tallow tree	23751 2706	232	Tetraitheca	1153
S. see Cephalaria	345		510	Tamaquan'a	235	233	Teicriutu	1G18
43 S. see Scabiosa	346		68	4455*%, 548		T. see H^ptis	1661	
Succory	2334		111	Tamhra	14323 1613	388	T. sec A^petea	
265 499 Succbiwa	1891		226	Tamarind tree	1989	123650 2691	146 2690	
Sugar Cane	284		285	Tamhndus	1989	495	TIALAMIFL6RIB _r SubC. 1.	
108 Sulphurwort	882		512	Tamarisk	899	229 496	ThMia	4
110 Sumach	894		514	TA^MEJE, Or. 193.	899	483	Thalictrum	1633
Summer Cypress	6466 809		515	Tamnia	1757	396 541	Thamnfdium	3239
US TM Sundew	922		529	Tamndia	11163 1358	104 518	Thamnoch6rtU3	2741
358 Sun Fern	25378 2908		538	Tiimus	2784	292 503	Thapsia	845
104 Sunflower	2409		538	Tanacetum	2312 459	290	Thapsus	3940 480
S19 Sun Rose	1&90		538	Tanacmlta	2310	2	Tua	2037
159 Supple Jack	10322 1214		338	T. see Pehtzaa	2311	iheeanTea	5411 67	
305 511 Sutherland^ Nut	209!)		338	Tanace'cum	1766	318	Thcleblus	31W
505 Swarrowsnt			248	Tanace'cum	1766	318	ThU'phora	3131
305 511 Swainsbnta	2097		527	Tanier Pea	19319 2136	453	Thclorema	3089
94 Swallow-wort	771		191	Tanna	12334 1460	422	Thelypteris	25601 2938
104 Swallow-wort	6786 845		68	Tdnquin	4462 550	67 525	Thcnfrdm	533
413 Swamp Locust tree	25121 2884		338	Tansy	2312 459	318 503	Theobrftima	2176
385 Swamp-post	23461 2666		82	Tapachmia	5349 & 53 650	318	T. see Guaxitma	2176
221 512 Swartzm	1591		188	Tupia	12125 1440	67 55-	Theophrasta	526
901 Swedish Bean tree	12966 1507		342	T. see BrachyleVia	2328	162 511	Themn'ipsis	1241
264 Sweet Turnip	v. 1883		342	Tare	2134	532	T. see jfnagyris	1259
1 Sweet Bay	10387 1226		432	Tarantan	3002 K71	92	Thesium	740
210 Sweet Briar	13448 1522		432	Tarfxacum	19917 2B07	92	T. see Com&ndra	741
Sweet Calabash	16858 1923		157	Tarfxacum	2297	92	T. see Leptomeria	742
124 Sweet Flag	8101 989		60	Tartan	2297	289 502	The&p^sia	2015
Sweet Gale	24184 2751		499	Tauchera	1878	329	Thistle	2241
385 Sweet Gum	23470 2667		263	Tauzain	23469 2666	278 499	Thlaspi	1849
511 Sweeti/i	2141		385	Taverniera	2122	263	T. see JEthionl'ma	1876
Sweet Marjoram	15127 1680		310	Taxanthema	7521 sy. 929	263	T. see Bivon^a	1874
555 Sweet Maudlin	21803 2387		388	Taxbdium	23638 sy. 2689	263	T. see Capsella	1850
313 Sweet Pea	19318 2136		404	Taxus	2823	262	T. see /epfdium	1873
Sweet Potato	4126 491		404	T. see Podocarpus	2690	265	T. see Psychlnc	1892
361 SwcUop	14370 1622		69	Taxitta	7546 933	86 503	Thom^sia	681
Sweet Sultan	22163 2434		43	Teak Wood	554	230	Th&ra	14574 16J7
Sweet Wil.	01385 1386		292	Teasel	344	60	Thorn-Apple	481
Ham	111626 1388		503	Tea tree	2037	105	Thorough Wax	
Swfrtia	793		526	Ticomia	1710	145 505	ThoumfiVi	1164
Swiefema	1297		529	Tfcomaca'	9385 116S	114	Thrift	928
S. see Chloroxylon	1297		253	Tictona	554	125 540	ThrInax	999
327 Swine's Succory	2221		259	Teidia	1810 76	324 521	Thrtncia	2209
117 Sword Fern	2906		499	Teesdalflo	1852 76	79	Throatwort	612
Syagrj*	2648		111	Teghre	4720 591	76	Throatwort	4981 607
5 f 19 Sycamore	25061 2867		516	Tefephthum	904	478 505	Tiiryallic	152
Symphtaia	623		404	Tefevium	11841 1410	388 535	Thdja	2693
*8 523 gymphytum	433		478	Telfairia	24572^y. 2832	388 535	T. see Cupressus	2194
			511	Telpea	321	245 529	Thunblrgsa	
			278	TemplettiniVi	1946	157	T. see Thymelaa	10155 117
			125	Tenageia	6145 994	532	THYMELJ^JJ, Or. 161.	
			375					

Lin.	Nat.	Sp.	Gen.	Lia	Nat.	Sp.	Gen.	Un.	Nat.	Sp.	Oen.		
238	528	Myinuia	1681	5	Tree of Sadness	293	42	S3	T. sec	Set	273		
239		T. see ^' cynos	1682	132	Tree Union	8600	1046	130	558	Tritoma	1023		
239		T. see t'alamftha	1683	142	Tree Sorrel	9222	1126	17	537	Tritbnia	123		
483	71	see Micromeria	3379	298	Trefoil	2066		188	503	Triumfetta	1443		
233		T. see JVepeta	1654	191	Trehsia	f ¹²²⁷³	1469	249	TYxago		15761 1771		
238		T. see Pycnанthe-	mum	i	TREMA'NDREAS, Or.	23.		363	521	Trixia	2442		
			1679	501	Trembling American			370	536	Trizeuxis	2535		
134	539	Thysanftus	1050	401	tree	24385	2793	518	Trochi&anthes		3325		
106	518	Thyssennum	872	456	Trcmella		3150	523	Trochocarpa		513		
474		T. see />eucfdanum	882	444	Trcntepuhlia		3028	231	49i	TrdUius	1638		
177	517	Tiarélla	1381	252	Trcviraua		1796	96	525	Trom6triche	783		
177		T. see Astflbe	1382	472	Trevda		3299	74		Trongum	4820 591		
53	527	Tiarfdium	428	403	Trewta		2817	506	TaoPIubLEf, Or.	51.			
218		Tibourbou	13817	1571	533			144	502	TronkLolum	1148		
4		Ticksseed	27	14	TKIA'NDRIA, a.	3.		397	532	Trpbphis	2761		
480		TigHrea	28317 sy.	3366	14	T. MONOGY'NIA, Or.	1.	322	521	Tr6ximon	2198		
267		Tiger Flower	1910	23	T. DIUY'MA, Or.	2.		327	T. sec Krfgfo		2220		
391		Tiglum	23809	2715	36	T. TRIGY NIA, Or.	3.	98	True Dodder	6409	798		
267	537	Tigrfdia	1<10		I!?	516	Trianthema	{ ^	105	True Parsley	6835	855	
268		T. see Cypella	1912		372	536	Tribrachia	2561	201	True Service	12975	1507	
16		Tile Root	120	169	Tribulus		1306	459	Truffle		3181		
218	503	Til'm	1576	28	TYichsta		225	242	Trumpet Flower		1706		
503		TILIAYEJE, Or.	31.	460	Trichia		247	Tube Flower	15673	1752			
400	497	TiUachra	2790	461	T. see Arcy-ria			459	7'uber				
52	516	TilbeVi	417	461	T. see StemonUis			460	T. see Sclerodrlma		3190		
119	5J9	Tillandsia	956	461	Trichilia			3026	Trubcrclaria		SL'34		
119		T. sec BiÜbergVi	937	168	Trichofcladus			462	Tuberose		91&		
119		T. sec Caraguata	958	393	Trichofcladus			130	538	Tulbagh/n	1026		
430		Tfmmia	2995	461	Trichod^sma			129	Tulip		1017		
109		Tinus	7109	55	Trichod^sma			129	539	Tulipa		1017	
191		TirucdUi	12299	1460	26	Trichidium		140	T. see Jtfelanthium		1111		
358	522	Titlbnia	2413	36	Tricholae^na			539	TulipLcea, Or.	197.			
191				26	Trichohcloa			296	Tulp tree		1617		
				423	Trichotnanes			460	Tulostoma		3194		
				544	Trichotnanes			195	Tuna		12595 1474		
271	528	Tittmannia	1785	422	T. see DavalhTi								
249		Toadflax	1775	4197	7Yich6. C	25448	2921	771					
61		Tobacco	488	432J	manes	126089	2998	78J					
86		Tobtra	5605	16	Trichonéma			415	Tupelo		2892		
174	515	Tuch'a	18P1	485	T. sec Spatalanthus			124	Tupfstra		993		
80	519	Tocoyina	636	21	Trichdphorun			63	Turbeth		4102 491		
86	507	Todd&lia	(86)	351	Trichophfilum			473	518	Turgfenia		3314	
423	544	Tödea	2944	488	T. see Eriophyllum			113	516	Turgosia		916	
140	539	Tofieldk	1110	372	Trichorhlza	22777 sy.		3	Turmeric		1		
327	521	Tdpis	2218					256	H1	Turner*		900	
1		Tonchdt	42		388	515	Trichosanthes		516	TuRNERIucar, Or.	88.		
318		Tdng'o	19586 sy.	2174	474			264	499	lumip	16593	1883	
318		Tonquin Bean	2174					52	Turnsole		424		
14	504	TonsiUa	101	375	T. see Ceratosanthes			362	Turnpentine		22243 2440		
86		Todna	5643	240	528	Trichostema		397	Turpentine tree				
398		Toothache tree	2764	233	T. see Isanthus			24216	27g3				
248		Toothwort	1770	428	Trichostomum			414	Turpithum		4102 J9J		
116		Tbra	10710	427	T. sec Dicranum			283	Turpinto	25131 sy.	28b^nmsy.m		
109	518	Tbntylum	885	428	T. sec Dldymodon			168	506	TurreVi	1^?		
474		T. see Condylco&rpus		463	Trichothcium			256	Turrila		16167 18g		
251	528	Torenfo	1787	136	Tricdryne			499	Turritis		1^26		
102	518	Torllis	827	52	Tricratus	3467 W.		255	T. see Arabia		1^b17		
214	512	TormentlBa	1529	31	TricutpU			255	T. sec Chaptalita		2493		
426		Törtula	2973	556	Trldax	"		344	591	T. sec IptocBipas^a	1820		
464		Torula		96	Tridctea			363	T. sec Chaptalita		2336		
87		Touch me not	698	240	Tridctea			345	r.sccLcria		2336		
56	527	Tournefortia	446	248	Trichostemum			351	T. sec Perdfcium		2355		
53		T. seefeltiotorphium	424	284	T. sec Stylosanthes			293	Tutsan		2043		
56		T. see Messercmni diet	446	298	T. Bee Trigonlla			368	Twayblade		250J		
244	526	TourrrtUa	1719	140	536	Trigluchin		458	Tympans		3101		
255		Tower Mustard	1826	297	511	Trigonella		375	540	2Jpn		2; ^	
256		Tower Mu&tard		300	T. see Lbtus			288	Typhalea		17907 2011		
				270	504	Trigonia		540	TYPHINJS	Or.	203. ^		
1107		Toxico	f 7170	141	538	Trillium		454	T^phula		3138		
4025		ndñron.	f 24426 sy. 2803	104	518	Trinia		280	511	£/1ex		!<>	
79	522	Trachelium	61S	32	543	Tnodia		280	U. see Staurocanthus				
106	518	Trachymlc	867	289	TYibnum	17984	2014	471	527	Ullbl			
103	518	Trachyspcrmum	840	182	505	Triorteris		534	524	ULMARA	Or.	171.	
225	496	Trachytilla	1603	182	T. see Tetrapteris			203	534	Ulmana	1^040	wfc	
125	540	Tradescantla	1000	80	519	Triosteum		100	534	Ulmus	81*		
125		T. see Campelia	1001	82	504	Triphasia		106	518	U. see Plinero	J/j		
125		T. see Cyambris	1002	248	Triplnnaria	15748 sy.		437		Ulospelimum	J/J		
307		7Yagacfintha	18956	2102				439	517	U. see EncoB^lum	3075		
375	533	Tragia	2595	401	531	Trfplaris		478	516	UMBELLIFEIUB,Or.	98-		
103	518	TYagium	837	347	TYipblum	21351	2337	289	511	Umbircus	\$**		
102		T. see Ledebur/a	829	347	Tropoly	21351	2337	14		Umbrella tree	*-*		
239		7Yagor/ganumal	1681	377	543	Tripsacum		472	519		17U92 20J*		
321	521	Tragopogon	2197	36	T. see Anthephora			377	541	Umbrella-wort	A^1		
159	531	Tragopyrum	1211	328	521	Triptilion		377		Uncaria	f>1		
WS7				28	5^3	Triscutrum		472		Uncinia	2gJ		
363	j	^<Xo>	f 6443 606	28	T. see TricbaHa			377	541	Umbrella-wort	2gJ		
51	514	Trapa	403	319	515	Tristania		377		Ctkiedo	H071 J3g		
00gf		Traveller's f	14448	26	543	TrietegIB		377		Uncinia			
"^"		Joy	114466	1631	468	539	Triletefa		377		Urnula		
257		Treacle Mustard	1843	34	542	TYfticum		31	441	Urania			
187		Tree Celandine		34	T. see Agropyrum			226	496	Urcaria			
				31	T. see Brachy	podium		118	537	Urcoccarlia			
887	"	Tree Mallow	12092	1422	33	r. seeSclerochloa		308	511	Urbnba			
				33				441		Urania			
										Uraria			
										Urcoccarlia			

GENERAL INDEX.

575

Lin.	Nat.	Sp.	Gen.	Un.	Nat.	Sp.	Gen.	Lin.	Nat.	Bp.	Gen.		
465	f/redo	3274		257	499	Vpslcaria	1838	254	Water Cress	16106	1819		
502	Urina	2010		59	527	Vestia	475	141	Water Dock	9179	UitS		
	U. see Pavbnia	2011		311	Vetch	2133	102		Water Dropwort	833			
137	Urinaria	23760	2708	307	Vetch	18936	2102	232	Water Germanander	14706	16<		
	Uropetalon	1079		313	Vetchling	19302	213d						
	Urospérnum			279	511	Viborgux	1954	103	Water Hemlock	838			
	19794 & Isy.	2199		109	519	Hburnum	892	9	Water Horehound	69			
360	Ursfnia	220535V	2430	311	511	Ffcia	2133	59	Water Leaf	476			
379	friftica	2615		312	V. see £'irum	2134	269		Water Lemon	16863	1923		
379	U. see BoehraenTi	2613		311	V. see Aba	2132	217		Water Lily	1556			
489	U. see Conociphalus			313	V. see Lathyrus	2136			Water Melon				
		3414		133	Victorialis	8703	1046			23715 v.	2699		
217	534 ZtortcBJC, Or. 17a			20	537	Vicussexif'a	147		Water Milfoil	mm			
159	Urucur&na	13837	1565	19	V. see/Vis	142	104		Water Parsnep	848			
444	UrviUVa	1216		316	512	Vigna	2153	1587	Water Pep.	C10237	1210		
466	Vsnea	3110		359	522	Viguiera	2416	1603	per	I10365	1223		
125	ftstiligo	3275		58	526	VillarsiVi	464	142	Water Plantain	1132			
9	UHan	8170	996	302	511	Vilmorfnf'a	2077	139	Water Purslane	1100			
90	Utnculária	67		164	511	Viminária	1254	29	Water Reed	230			
226	UVa-crispa	5925	719	67	525	rinc'a	528	402	Water Soldier	2802			
227	Uvaria	1620		67	Vincentifxicum	6112	761	4	Water Starwort	28			
130	U. see Guatteria	1635		82	Vine	655	225		Water Vine	14314	16W9		
173	Uvaria	8484	1023	228	Vine Bower	14473	1631	58	Water Violet	456			
131	ITva-drsi	11081	1346	132	Vine Leek	8603	1046	160	Watcrwnrt	1283			
131	539 Uvularia	1032		87	501	nola	701	17	Watson/a	124			
131	U. sceDisporum	1033		89	V. see Calfrtrion	707	109		Wayfaring tree	7129	892		
177	Vaccaria	11365	1385	89	V. see Erpetion	702	81	519	Webenz	646			
1.%	523 VACCINIE**, Or. 114.			89	r.scelonfdium	705	362	522	Wudela	2436			
155	racfnium	1194		89	V. see NoisCttia	706	3G3		W. sec fl/elampb-				
1.%	523			89	V. see Sblea	703		dium	2446				
411	520 Valantia	2859		89	500	riOLARIEX Or. 20.		158	Weiimfnnfa	1217			
45	V. see GMium	348		89	Violet	701	427		Wcluaia	2976			
79	Valerandi"	5193	618	257	Violet Thlaspi				W. sec Dfssodon	2968			
U	Valerian	95				16277	1843	132	Welsh Onion	8624	1046		
U	520 Valeridna	95	228		Viorna	14466	1631	142	WendlAndta	1129			
U	V. see Patrinia	330	55		Viper's Bugloss	445	344	521	Wernēn'a	2334			
41	520 Valerihnete, Or 105.				Viper's Grass	2201	233	528	Westrlngia	1649			
14	580 ValerianUla	96			Virgaurea	21424	2339	34	Wheat	269			
71	525 Vallaris	538			511	Virgilia	1242	309	Whirling Plant				
395	525 Vallesia	580	358		V. see Gal&rtda	2412		19053	2180				
123	536 Vallisneria	2735			Virginian Creeper		201		White Bean tree				
372	538 Vallotta	971				5377	656		White Cedar	12965	1507		
372	Vdnida	2564			Virginian Hemp	2770	388		White Clover	23653	2692		
372	/. see Sarcanthus	2563	187		Virginian Poke		299		White Fiorin	18505	2Utti		
232	528 Vandellia	1791			12083	1420	26		White Spruce	23626	2tK6		
80	519 Vanguutra	637			Virgin's Bower	1631	388		White Tree	2179			
373	536 Vanilla	2597			Virōla	24557 sy.	2829	319	White Vine	14448	1631		
441	VariolAria	3091			Hscum	2750	228		White Wood	2282			
70	527 Varr&fill	564			504	Vfsmea	2192	335	Whitleya	4032 sy.	486		
278	511 Vascfca	1948			Visn'ga	6737	841	61	Whitlow Grass				
495	VASCULAVIES, DIV. I.				514	Visnea	1462		16531	1873			
115	516 Vauanthes	919			Vitalba	14448	1631		Whole Schoenus	1396	157		
437	Vaucheria	3056			Fltex	1743	21		Whortle Berry	1194			
99	502 Velezfa	802			Viticella	14473	1631		Widow-wail	106			
265	499 Fella	1888			Fitis	655	155		Wild Basil	1678			
265	V. see Bbleum	1889			ntis Idia:a	10123	1194		Wild Beet	7480	929		
265	V. sec Carrkhtira	1890			Vitm&nnta	1165	238		Wild Blite	23214	2628		
260	V. see Eub'dium	1856			Vittkria	2927	115		Wild Bugloss	444			
o ⁷⁹	522 Velléia	619			Voandzia	2170	380		Wild Chamomile				
385	Velonia	23443	2666		152	Vochfsia	21	55	21715	2376			
130	538 Velthelmi'a	1024			526	Vohiria	469	354	Wild Clove tree	12766	1489		
373	Vcntcnatai		22829 sy.		247	Volkamefa	1753			Wild Cumin	717		
101	Venus's Comb	6563	819		247	V. see Cleroddn- drum	1752	198		Wild Liquorice	2140		
170	Venus's Flytrap	1331			4585	574	90		Wild Oat	1864	223		
421	Venus's Hair	25546	2930		512	Vouupa	108	314		Wild Olivie	J 7145 89* J 10155 1197		
77	Venus's Looking-glass				283	Vulncaria	17625	1968		WildPurslane	2, U9	1460	
		5060	608		283	Vulparia	14178	1600		Wild-Radihh	1G683	1897	
4	Venus's Navelwort	437			223	Wachendorffia	136	157 J		Wild Rosemary			
539	Keratrum	2839			75	Wahlcmblrgia	606	191		11036 V.	1341		
400	r.see Chamslfrium	2792			202	Waldstclnia	1514	266		Wild Service	12967	1507	
409	V. see Leimanthium	2840			212	W. see Comaropsis	1526	172		Wild Tamarind	34		
60	527 J'erbascum	480			507	Walkena	744			Wild Tansy	13658	1528	
60	y. see Ramonda	479			424	Walking Fern	25701	2955		Wild Thyme	15147	1681	
246	y. see Terbtma	1749			425	Wall Cress	1827	5		Wild Vine	5369	655	
	V. see Aloysia	1739			254	Wallflower	1818	214		Willdenbyui	2738		
	y. see Priva	1746			2537	Wallflower	1806	239		Willemētfa	3301		
	y. sec Zapaljt'a	1745			264	j. Wallflower	1883	82		Willow	2732		
	FERBENAYEJI, Or. 143.				382	Wallichm	2644	395		Willow Herb	12187	1451	
359	Verbesina	2399			540	Wallrbthto	832	472		Willugheta	553		
356	V. see Actindmeris	2417			518	W. see nt ex	1743	393		W. see Fagre'a	575		
	V. see Chrysanthem-				420	Wallsbra	3344	155		Wind Flower	6354	794	
357	um	2396			477	Walnut	2664	69		Wine Palm	23552	2678	
160	V. see Synedrella	2400			384	Walnut	1915	70		Winged Pea	18595	2068	
458	Vferen	10354 & 5 sy.	1219		268	503	Walthferfo	1872	231		Wlntera	14280 sy.	1602
333	Vermicularia	3158			170	Wampee tree	1527	98		Winter Aconite	1640		
	1 Vernbn/a	2262			34	542	Wangenhelmofo	266	386		Winter Berry	1089	
	2 yerdnie	2263			40	Warratap	321	301		Winter Cherry	586		
	£ sec Diplophyllum	46			293	Warrat	v. 8	2038					
	P. see Leptandra	44			262	Wart Cress	1872	224					
	Y^rpa	3142			182	Wartwort	12409	1460	1*8				
SS	Jerrucaria	3114			37	Water Caltrops	403	71					
	vervain	1749				Water Chickweed	159			10336	1217		
						2418	294						

GENERAL INDEX.

Lin.	Nat.	Sp.	Gen.	Lin.	Nat.	Rp.	Gen.	Lin.	Nat.	Rp.	Gen.
255	Winter Cress	1822		341	521 Xeranthemum	2322		375	541 Zannichellia	2587	
82	Winter Grape	5372	655	342	X. sec Aphelexis	2326		398	507 Zauthoxylum	2764	
143	Winter Green	J	1349	342	X. sec Phoendcoma	2325		529	Zapanz'a	1745	
173				140	539 Xerophyllum	1115		69	Zapot'dla	4504	561
238	Winter-sweet	1680		398	541 Xerutes	2776		377	543 Zei		
93	Wise Men's Banana tree	6059	746	145	503 Ximenest'a	2358		34	Zea	2285 sy.	269
315	512 WistanVx	2147		18	538 Xiphidium	133		1	Zebra Plant	49	3
51	Witch Hazel	407		417	544 Xiphopteris	2906		3	Zedouria	166	14
46	527 Witherfngia	359		370	536 Xylbium	2536		3	Zedoary	166	974
108	Withy	7010	881	40	532 Xylolemum	320		123	538 Zephyrinthes	C126	10
16	537 Witsenfo	117		390	533 Xylophylla	2709		«	Zerumbet	£169	14
263	Woad	1877		227	496 Xylophia	1687		a			
223	WolPs-bane	14184	1600	71	Xylopicron	4596	576	488	522 Zembnia		
79!	Woodbine	JTM	621	83	Xylopyrus	5431	660	507	ZieriVi		
1595			1210	79	Xylosteum	{**\$}		237	528 Zietenia	leTO	
313	Wood Everlasting			471	Xylosteum	{**\$}		131	539 Zigad'nub	* T030	
	Pea	19285	2136	15	541 Xyris	110		265	499 ZUla	189J	
45	Woodroof	350		94	525 Xysmalobium	768		2	537 Zingiber	JJ	
233	Wood Sage	14731	1648	400	Yam	2787		352	522 Zinnia	2373	
423	544 WoddsiVi	2940		536	Yarrow	21847	2387	381	543 Zizania	2634	
185	Wood Sqrrel	1414		501	Yelhoi	10		543	Zizir	3319	
419	544 WoodwardtVi	2919		170	Yellow Bird's-nest	1330		528	Ziziphora	71	
419	W. see Dobdf'a	2920		384	Yellow Chestnut	83		508	Zizyphus	6C0	
61	Worm Grass	484				83		Z. see Condal/Vi	661		
225	496 Wörmii	1610		236	Yellow Dead Nettle	23433	2666	83	Z. see Palitirus	659	
338	Wormwood	2313				360		360	521 ZceY'a	2+31	
283	Woundwort	17625	1968	248	Yellow Rattle	14961	1667	439	Zonaria	3l)H*	
4	Wrack Grass	25		115	Yellow Root	1771		284	511 Zornia	1^	
67	525 Wrightfo	535		360	Yellow Sultan	932		310	Z. see Dicerma	2121	
68	W. see Cameraria	547				109		109	518 Zostmia	887	
9	528 WulfeniVi	63		313	Yellow Vetchling	22166	2434	4	541 Zoster'a	&	
140	539 Wtinnbtfa	1116				137		539 Zucragm'a	1081		
100	Wych	6544	814	153	Yellow-wort	19302	2136	198	Zuzygium	12770	149J
101	WyTia			100	Yervamora	6530	812	170	Zwfngera	10955 sy.	1318
				404	Yew Tree	2823		405	Zygia	24603	2833
						239		405	Zygis	15143	16S1
380	622 Xaiithium'	2627	130	589 Yucca		1027		435	Zygnema	S041	
320	504 Xanthochymus	2188		328	521 Zaciitha	2229		429	Zygodqn	29>>	
115	496 Xanthorrhiza	932	124		Zalaccq	8094	987	370	536 Zygoietatum	25ft	
134	539 Xanthorrhæ'a	1048		357	522 Zaluzania	2403		507	ZYUOPHY'LLEA, Or.	54	
398	507 Xanthoxylum	2764	5		Zambac	294	43	169	507 Zygophylluin	1S9J	
50	X. see Fædra	398		403	Zamia	2814		169	Z. sec Guaiacum	I***	

THE END.

**FIRST
ADDITIONAL SUPPLEMENT.**

ADDITIONAL NAMES OF BOOKS REFERRED TO.

Chab.sc	Chabrey (Dominicus). <i>Omnium Stirpium Sciagraphia</i> . Fol. 1677.	Hern. mex.	Hernandap (Franciscus). <i>Nova Plantarum, Animalium, et Mineralium Mexicanorum Historia</i> . 1 vol. folio, with wood cuts. Rome, 1651.
Chan. ill.	Illustrations, by Alfred Chandler, and Descriptions, by William Beattie Booth, of the Camelhee. 1 vol. imperial 4to. London, 1831.	Hook. am.	Hooker (William Jackson). <i>Flora Boreali-Americana</i> . Published in 4to numbers, 4 of which have appeared.
FLbr.	Pohl (Jo. Imman.). <i>Icones Plantarum Brasiliensium riariorum</i> . Fol. 2 vols. Vienne, 1828—1831, continued.	Led. alt.	Lebedour (Carolus Fredericus). <i>Icones Plantarum Altaicarum</i> . Fol. Berol. hne, 1830.
FIJav.	Blume (Carolus Ludovicus), M. D. <i>Adjutore Joanne Baptista Fischer, M.D. Flora Javas necnon Insularum adjacentium, &c.</i> Folio. Brussels, 1828, continued.	Rchb. rar.	Reichenbach (H. G. L.). <i>Icones et Descriptions Plantarum</i> . 4to. Lipsiae, 1822, et seq.
FI. lap. ed. 2.	Linnaus (Carolus). <i>Flora Lapponica</i> . Edito 2., curante J. E. Smith. Londini, 1792.	Rich. con.	Richard (Louis Claude). <i>Me'moires sur les Conifères</i> . 4to, 1 vol. Stuttgart.
FL med.	Tenore (Michel). <i>Saggio sulle Qualità medicinali delle piante Neapolitane</i> . 8vo. Napoli, 1808.	Ster. den.	Sternberg in <i>Denkschriften der Königlich-Baierischen Botanschen Gesellschaft</i> in Regensburg, 4to. Regensburg, 1818, continued.
Gus. pi.	Gussone (Joannes), M. D. <i>Piante rariores, quas in Itinere per Oras Ionii ac Adwatici Maris et per Regiones Samnii ac Aprutii collegit Joannes Ousaone</i> , M. D. 4to. Napoli, 1826.	Sw. flgar. 2.8.	Sweet (Robert). <i>The British Flower Garden</i> . Second Series. 1830—1832, continued.
H.ftH.	Hoppe et Hornschuch. <i>Reise nach d. Adriat. Meere, c iconae</i> .	Sw. guide.	Sweet (Robert). <i>The Florist's Guide and Cultivator's Directory</i> . 8vo, with plates. London, 1831.

ADDITIONAL AUTHORITIES FOR' GENERIC AND SPECIFIC NAMES.

A, Dee,	<i>Alphonse Decandolte</i> . Author of <i>Monographie des Campanulas</i> , and son of the distinguished Decandolle.	Guss.	<i>Joannes Gussone</i> , M. D. Director of the royal botanic garden at Palermo, and a botanical author.
Bane, John Bancroft,	M.D., of Jamaica.	Her. B.	<i>Sir Joseph Banks's Herbarium</i> , preserved in the British Museum.
BaxtMSS.	<i>Manuscripts of Mr. William Baxter</i> , a collector of plants in New Holland in 1827—1829.	Her. Doug.	<i>Douglas's Herbarium</i> , in the keeping of the Horticultural Society.
Bertero.	<i>Carolus Josephus Bertero</i> . Author of <i>Specimen medicum nonnullas indigenas Stirpes continens exoticis succedaneas</i> . 4to. Taurini, 1811.	Jack.	<i>William Jack</i> . A writer in the Botanical Miscellany on the plants of the Malay Islands.
Bigel Jacob	<i>Bigelow</i> , M.D. Professor of botany at Boston, U. S., and author of <i>American Medical Botany</i> , and of <i>Florula Bostoniensis</i> .	Kautfuss.	<i>G. Fred. Kaulfu&s;</i> , M. D. Author of » work on Ferns, and demonstrator of botany at Halle. He died 1831.
Bishop.	<i>David Bishop</i> . Curator of the botanic garden, Belfast, and author of <i>Causal Botany</i> .	Lessing.	<i>C. F. Lessing</i> . A writer on Composite and resident at Berlin.
Booth.	<i>William Beattie Booth</i> . Describer of the <i>Camellias</i> figured in Chandler's Illustrations of the <i>Camellias</i> .	Nectoux.	<i>H. Nectoux</i> . A French botanist and traveller.
Borrer.	<i>William Borrer</i> . A writer on British plants, and one of the authors of <i>Lichenographia Britannica</i> .	P.B. Webb.	<i>Philip Barker Webb</i> . A traveller in the Canaries, who sent thence species of plants not previously in Britis ⁷ gardens.
Bree.	<i>W. T. Bree</i> , A. M. A discriminative amateur botanist	Penny.	<i>George Penny</i> , A. L.S. Botanical cultivator in the Epsom Nursery.
Caley.	<i>George Caley</i> . For ten years a botanical collector in New South Wales, and afterwards curator of the botanic garden at St. Vincent.	Richards.	<i>John Richardson</i> , M. D. A traveller in the northern parts of British America, and author of the Appendix on Natural History to Franklin's Travels.
Den*.	<i>John Demon</i> , A.L.S. Curator of the botanic garden, Bury St. Edmunds, from 1821 to 1829.	Riedel.	<i>L. Riedel</i> . A botanical collector attached to the botanic garden at St. Petersburg.
O.Anders.	<i>George Anderson</i> . Author of a monograph on Peonies, in vol. xii. of the <i>Linnsan Transactions</i> , and a collector of willows.	Sermge.	<i>N. C. Seringe</i> . A Swiss writer on systematic botany in Decandolle's <i>Prodromus</i> , &c.
G.E. Sm.	<i>Gerard Edwards Smith</i> . Author of a <i>Flora of South Kent</i>	Vahl.	<i>Martin Vahl</i> . Professor of botany ac Copenhagen, and a botanical author.
Gow.	J. R. Gowen. Originater of hybrid Aialeas and Rhododendrons at Earl Caernarvon's, at Highclere.	fV.QN.	<i>A. Weihe and Ch. G. Nees von Esenbc*</i> . Authors of <i>Rubi Germanici</i>
		Wray.	<i>Dr. Wray</i> . Resident at Augusta, in Georgia.
		Zuccarini J.	<i>G. Zuccarini</i> . Author of a monograph on the genus <i>Oxalis</i> , and of other papers.

ADDITIONAL ABBREVIATIONS.

Hybrid. Known or presumed to be hybrid, but where raised not precisely known.
 English hybrid, Scotch hybrid, &c. Hybrid, with the country in which it has been raised stated.
 Gardens. Varieties which have appeared and are preserved in gardens.

FIRST ADDITIONAL SUPPLEMENT.

The alphabetic mode of arrangement has been adopted to avoid the necessity of a second Index. The number prefixed to each genus and species declares the relative place for each in the Hirtus; and such genera as have not been inserted there have here a star (*) prefixed to them, their etymology given, and their Linnean class and order, and Natural order, stated. In the copies of the Hirtus Britanicus which may hereafter be printed, stars will be inserted into the plates wherever the additions made in the Supplement come in, in order that the reader may be apprised of their existence, and lose as little time as possible in turning to them.

The species to which this character (\$) is prefixed are those which have been raised in the Exotic Nursery of Mr. Joseph Knight, King's Road, Chelsea, from seeds collected and brought home by Mr. Baxter. The names of these species have been verified and authenticated by Mr. Knight's very able and intelligent foreman, Mr. Alexander Scott, who, while he trusts that each may be found accurately denominated, nevertheless reminds us that Proteaceous plants are subject to great variation in foliage, in their progress upwards from a seedling state. Besides the above, Mr. Baxter's &c &c have produced various other species of Banksia and Dryandra, and species of Petropilum, Oxylobium, Mirbelia, Gomphiolema, Hovea, Burtoa, Pultenaea, Brachysema, Kennedyi, Daviscia, &c.: many of which, when farther advanced in growth and age, may prove to be new species.

Whoever wishes to ascertain the additions and improvements made subsequently to the present Additional Supplement may consult the *Gardener's Magazine!* in which, under the article, which will henceforth be contained in every number, entitled, "Floricultural and Botanical Notices of New Plants, and of Old Plants of Interest, supplementary to the latest editions of the Encyclopaedia of Plants, and of the Hirtus Britannicus," will be found the name of every plant newly introduced or originated, and every recent improvement in botanical nomenclature. At the close of every year, or of every two or three years, according to the accumulation of matter, these names and improvements will be collected and published in additional Supplements similar to the present one.

Systematic Name and Authority.	English Name.							Reference to Figures.
2837. jfCAXJIA. (Sect. II. Conjugate)								
24763a sapindifides Cun.	Sapindus-like							MoretonB.1830. S p1
2867. ^{2867.}	hybrid							
25055a hirbridum Bosc	Austrian	or 20 ap.my	G	Amcr.hyb.1790. L co	Bot cab.			
%68 25058a aquatracum Lk.	Georgian	or 40? my.jin	G	Austria	L co	Trat.ar.1		
*89 25052a ibridicum Bieb.	xowad-leaved	or 40? ...	G	Asiatic 0.18ii.6.	L co			
^570 95060a circinatum Ph.		or 40 ap.my	-	Columbia 1827. L co	Hook. am. 39			
2387. ACHILLVA.								
**>? 2180 acuminate Led. ?	acuminated	■ A or 3 aii.8	W	1830. D co			
*154*a. A^HLY5 DM. ACHILLYB. (^CWM, the goddess of obscurity. genus obscure.) I3. I. Podoptoltoeg.								
tripbillia Dec.	A A or ... apjn	W	NT Amer. 1827. D i.l.p	Hook. am. S				
1600. ACOXVTVM.								
NuttalliiSwt.	Nuttall's	£ A or 5 jn.o	Pa.B	N. Amer. 1829. D co				
pallidum Nut., and Loud. hort. brit.		No. 28368.	Pah	Moldavia 1830. D co				
moldavicum Bes.	Moldavian	^ A or 5 jn.o						
2857* ^{Q e 2 0 i} ACRO'STICHUM.								
8857? ^{Q 2 1 5 a} fffande Cun.	magnificent	■ IAIor 23 or	=	Br	MoretonB.1828. D Lp			
^ 25317a flagelliflrum Wat.	rod-bearing			E. Indies	1828. D I.p	H. & 0.23		
1378a ADA'M/i4 Wai. (J-Adam, M. D., of Calcutta.)		10. 5.						
28428 cySuctw. hue.berricd		<1 J or 4 ...	Pk	Caprifoliictej Hj/drangcdcte Lindl.				
1880. ADE/SMIA.				rfepal 16". C Lp	Bot mag. 3046			
microphl lla Hook.	small-leaved	rt. J or 2 va.sea.	Y	TiParia. TPA				
clammy.-wfxi i-jor		2 ..	Y	Chile 1831. S s.l				
a. ²⁹³⁰ ^DIA'NTUM.	(Section IV. DigitMo-pinnata;)							
25540a pubtoens ScA*.	downy	xiAIor 1 ap.my	Br	N. Holl. 1830. D pi				
1634. ^{1634.}								
- sibirica Pat.	Siberian	■ A or 2 ap.my	Y	Siberia 1827. D co				
- davurica Rchb.	Dahurian	■ A or 1 ap.my	Y	Siberia 1827. D co	Rchb. ic. S21			
- distorta Ten.	distorted	■ A or 1 ap.my	Y	Naples 1827. D co	FL med. 448.1			
88589 ^{SS65*} AE'RIDES.	hornJipped	■ ■ 1 jl.au	F	India 1820. D p.r.w	Bot. reg. 1485			
" cornutum Ztz.								
^EGOPO^DIUM	6839 Podagrillria	A or 1 my.jl	W	England ih. pi	Sk co			
2 variegata Dens,	d							
Swt.	small-	rk pr	J jn.au	Li	Persia	ISS IS		
		rk pr	J jn.au	Li	Persia			
		Pp	2					

1774. iiNTIRRHR NUM 15770 majus " 5 variegatumita* variegated-/*£ A «* liJn.au R Engl. gardens. C ru
3372. APHANOCHPLUS. - blandus Benth. mild £g QS] fra 2| su " G Nepal 1827. S co Bot mag. 3091
- *2520a. APLETTRUM Suit. APLECTRUM. {A, without, plektron, a spur; flower.) 20.1. Orchidect. « - hiemale&v. wintry * A cu 1 •• Br N. Amer. 1827. O s.p Corallorrhiza hiemalis Nut,
3342. APLOPHYLLUM. Buxbaumti J. Buxbaum's £ A un 11 Jn.au Y Levant 1828. ^{Orfa} C co 10921a tuberculatum J. warded & A un If jn.au Y Egypt 1824. C co
- *2547a A'PORUM Blume (Apox, a running shoot; appearance.) 20.1. Orchidear. anceps Lindl. two-edged, rfmfd E cu 1 year Yu Bengal 1826. ? D p.r.wBot. reg. 1239 Dendrbium anceps Bot. reg. 1239. Hort. brit. Mo. 28526. p. 489.
1605. AQUILETGIA. 14288a Garnier[†]aa Swt. Miss Garnier's £ A or l*my.jl P.Str Eng.hyb. 1829. D e o Sw.fl.gar.2.1.103
1827. i4'RABIA retrofracta Grdh. bent back jf A *** apjl Bh N. Amer. 1827. S ru Hook am. Turrifis retrofracta Hook. am. 17. f A.
- 16156 prs^cox W.fK. [?procirrens IV.SfK.] 2 variegata Dens, variegated-/* A or | jn.jl W gardens ... D Lp
- 16178 llicida 2 variegata Dens, variegated-/** £ A or | jn.jl W gardens ... D Lru
- 7289a *ambuciflia Cto. Elder-leaved i j o r 5 au G N. Holl. 1823. C si 7299a macrophlla Cun. large-leaved • LJ or 6 Norfolk 1.1831. C s1
1345. -4'RIBUTUS. 11079a mucronata L.JU. mucronate-/*d U LJ CU 1 my W Magellan 18% L Lp Bot. mag. 3093 , MilleriSwt Miller's * or hybrid 1825. L s1
- - proceraitoff. tall * or W N. Amer. 1827. L s1
- - Menzefisii Ph. Menzies's m or W N. Amer. 1827. L s1
1554. ARGEMO^NE. - Barclaidia Penny Mr. Barclay's £ Jsl or 5 Jn.n Crea Mexico ^{IRBM} 1827. S co intermedia Swt. hort. brit. ed. 2. p. 585. j but subsequently to the publication of Mr. Penny's name in Gard. mag. vi. 115.
493. ARGYREPA. 4244a pomacea&uf. amle-fruited * CDor. Pk E. Indies 1828. C r.m Gae. fr. 2.178 Lettsomia pomacea Box.
2582. iRISTOLO\:\H. 22844a caudata Unit. UaledJipped ^J (23 cu 5 jn Ld Brazil 1828. Sk It.l.r Bot reg. 1453
- . I43B. ARISTOTE'L'il 12121 MAcqui Eloocarpe* D. Don, HomaUne* Dec. 2 variegata Dens. variegated-/*i _J o r 4 ap.my W.a gardens ... C 1
- *%52a. AREACA'CIA Bancroft. (Euphonised from Arracacha, the name of the plant in Santa Fe) Umbel. esculent Dec. esculent & E esc 3 jn.jl lirsh.R Santa Fe 1823. D r.m Bot. man [^] 3092 Conium Arrachcha Hooker, and Loud. hort. brit. No. 6821.
2313. jiRTEMISIA. 82 lactiflora Wai. pale-flowered tt-L-Jeu 2 n P&W Nepal 182& C s1
- &&*1182a. ARTHROSTE'MMA PaiKm.(ifr/AronJoint,«fmma, crown; anthers jointed upon filam.) 8.1. Melost.23» mitidum (?)A. glossyJcaved n. LJ or 2 jl Pa,Li B. Ayres 1830. S p.l Bot. mag. 3418
- T. - - Green's « n p r 2 all sea W. 1828. C p. 1
1074. ^SPA'RAGUS. Smithicintu P. B. Webb, Smith's LJCU Teneriffe 1829. ...pi " plocamOldesP. A HW,Plocama.lk LJCU Teneriffle 1829. ... pl
- 88610 350: *SE£?A' glowy ?. A A or * jU Pk Greece 1829. D co Fl.gr. M4
- P. - - proliferou Ocu J jn.o Wsh Armenia 1824. S co Bux.c2.36.J
- thicket jfeAor f jl Br Britain sh.ro. D p
- S. - - various-leaved « 1Alor 2 f Br Norfolk 11831. D Lp madertnsePimiy Madeira *iAlcu 1 jLd Br Madeira 1828. D p1 Related to pram6rsum Swx. ?

- "3. BCEHME'Riu.
 23117a castaneefbla Cun. Chestnut-leavedi LJ un 15 n Gsh Norfolk 1.1831. C s.l
- *430, BOLIVA'R'il Schlec. BOLIVABIA. (In ~~l~~ ~~l~~ ~~l~~ 2.1. *Jasmine**, i
 trifida Schlec. trifld «L J or 2 jl 182a C Lp
- liSaaetiamiiThun.'* Heath-leaved *i_J or 2 ja.f Pk C.G.H. 1821. S p.l
791. BRACHYSTI/LMA.
 28134a crispum Grah. curled-forf A 2] el \$ au.o Br.Y C. G. H. 1829. C s.l Bot mag. 3016
- 2562; BRASAVO^XL,^{..} nodbsa Lindl. knotty £ESfra 1 o Ysh.G Mexico 1828. D p.r.w Bot reg. 1465
- 16130a pili5sa Hook. hairy * A cu \ ap.jn P N. Amer. 1827. S p.l Hook. am. 17. A
- *2170a. BRONGNIA'RTM H. B. & Kth. (*M. Adolphe Brongniart*, a Fr. bot) 17.4. *Legum. Getifrea*. 1. podalyrioYv Kth. Podalyria-like *L _J or 1 s.n F NewSpain1827. C Lp H.&B.n. 6.588
2552. BROUGHTON'^.
 . nvid& Herb, glossy £ ES or li jn,jl R £. Indies 1824 D p.r.w
- *1794. BROWA'LLW.
 15986a grandiflora Grah. large-flowered iDJor £ jn,d LLY Peru 1829. S p.l Bot mag 30(9)
- 193a BRO'WNEA
 17275a racemosa Jac. cluster ■■■■■ 6 ... Ro Caracas 1826. C s.l Jac ft. 25.16
967. BRUNSVI'G/A
 - grandiflora Lindl. large-flowered tf iAI or 1 au.fl Pk C. G. H. 1827. O r.m Bot reg. 1335
- *1135a. BUGINVILLE^il Commerson. (*Bougainville*, a French navigator.) 8.1. *Nyctaginece*. spectabilis J. showy ■■■■■ 10 - Pk S. Amer. 1829. C 1 -Lam.il. 294
1060. AULBrNE.
 - glauci?. Sf/S. glaucous tf U or 1} jn W Chile 182a Dip Bot cab. 1580
947. BUONAPA'RTJ?il.
 . gracilisSuf. slender £ O or 2 - ... Mexico 182a C s.l
1275. CASALPI'NM.
 . Gillidsii Wai. Gillies's SluJor ... - ... Mendoza 1829. C Lp
2672. CALA^DIUM.
 . lividumAC. livid ■■■■■ 1 s Din. W.Indies 182a D Lp Bot.cab. 1590
1683. CALAMI'NTHA 15179 grandiflora
 2 variegata f^n y variegated-/vtf ^ A or 1 jn.s R gardens ... D co
65. CALCEOARIA.
 5786 Wheeler Swt Wheeler's £ J^ot 1 my.o P Eng. hyb. 1831. D p.l Sw.fl.gar.2. SI3C
 279886 bfcolor Grah. tw<Mcoured n-LJor ft aus. Y.Wsh Peru 1829. C It Bot. Reg. 1374
- diffrcs Lindl. in Bot. reg.
- 27990 angustiflora g. & P. narrow-flwd tL LJ or 1 jn T Peru 1830.? C p Bot mag.
- 695 27905a HerbertinaLindl. Herbert's ■■■■■ Jor 2 my.o Y Chile 1828. S p ■■■■■
- GeiantfiRd Penny Gellan's £ LJ spl 4 all sea Och.&D Eng.hyb.1830. D r.m
 Intermediate between corymbbsa and purpurea.
- 8897 - h l a n s / W. gpJnS, £ LJ or 1 my.o Y Eng. hyb. 1830. D r.m
 - Hopeana Morrison's MSS. Hope's jg LJ or 1 jnmy.o Y Eng. hyb. 1830. D r.m
 Intermediate between corymbbsa and jalantaginea.
- Morrison* Penny Morrison's »i_i or 2 my.o Och Eng. hyb. 1830. C ID
- 28700 - polyantha Penny many-flowered £ | or 3 my.o Y Eng. hyb. 1830. D r.m
 Intermediate between corymbbsa and petiolafirs.
- *0701 578a Youigf Penny Young's £ LJ spl 2 my.o Och.&D Eng.hyb. 1830. D r.m Bot res 144S
 2 pallidior Penny paler £ LJ or 2 my.o Pa Eng. hyb. 1830. D r.m
 3 difcta Penny beloved £ LJ or 2 my.o Bt.Br Eng. hyb. 1830. D r.m
 4 atra Penny dark £ i_J spl 2 my.o Dk Eng. hyb. 1830. D r.m
 Intermediate between corymbbsa and petiolafirs.
- &>,- - chiloensis Lindl. Chiloe n. _J or 2 su Y Chiloe 1830. C r.m Bot. reg. 1476
2418. CALLIOTYSIS. "
 22016 AtkinsonaTW Doug. Atkinson's O or 2 su Y.Br Columbia 1826. L co Bot reg. 1376
- 28704 98. CALU'TRICHE.
 241a pedunculate Dec. peduncled ■■■■■ I ap.o W Scotland dit. S aq Eng.bot.2C06
1643. CA'LTHA.
 - biflora Dec. two-flowered ■■■■■ or * my.jl W N. Amer. 1827. D aq
 - leptose'pala Dee. small-sepaed ■■■■■ or 1 my.jl W N. Amer. 1827. D aq Hook. am. 10,
- 203a CAME'LL/d 18166 japdnica
 var.gpeci5saHor./fr.ilaw;«* showy * uJsp1 10 mr Dp.B Chfn? 1824. ■ Lp Chan.iLL32
 flagens Booth elegant «UsplOja Ro seedling ... Lp Chan. ill. 26
 imbricated Booth imbricated HJsp1 10 mr CB Chlna 1824. Lp Bot reg. 1398
 concinna Booth neat LJsp1 10 mr.ap W Chma 1822. ■ Lp Chan. ill 33
 Colvillf Swt Colvill's * U or 10 ap Bh.Spot Eng. hyb. 1824. ■ Lp Sw.fl.gar.2.i.2
 Sweet's Sweet's « U or 10 f Ro.Va Eng. hyb. 1824 P ■ Lp Sw.fl.gar.2.«.133
- JJS- CAMPANULA.
 49S7u Nuttallti Penny Nuttall's ■■■■■ or 1 jl.o B N. Amer. 18S9. D co
 Pp 4

- | | | | | | | | |
|--------|--|---|-------------------------------|----------------------------|---|----------------------------|---------------------------|
| 2<709 | .. Achraq Gill | Achiraa | £ LJ or | 4J au | D.R Mendoza | 1839. | D r.m Bot reg. 1SSS |
| 28710 | - lagunensis Lindl. | Laguna | £ or | 5 8 | Y.Spot Laguna | 1828. | D r.m Bot reg. 1311. |
| | 1539. CA'PPARIS. | | | | | | |
| 88711 | 13721a chinensis Q.Don | Chinese | * Q o r | 4 jn.au | W | a Amer. | 18S7. C s.1 Bot reg. 1390 |
| 28712 | - acutifolia Swt. | acuminate | Lindl. | in Bot reg. not of others. | | | |
| | (521. CAPRIFO^LIUM, | | | | | | |
| 88713 | 281(XSa occidentalis Lindl. | western | J or | 20 jn.au | O Ft | Vancouver!824. | C co Bot. reg. 1458 |
| | hirsutum Dens. | hdvycrafted | * | or 20 my.jn | Y | Canada 1822. | C co Bot. mag. 3103 |
| | JLoneco hirs&t;a Eaton | in his Manual of Botany. | Hooker in Curt bot mag. 3103. | Caprifoliu | pub&tions or Loudon's hort. brit. No. 5213. | | |
| | 1217. CARDIOSPE'RUMUM. | | | | | | |
| 88714 | 103360 canesuens Wai. | hoary | 5 liQj or | SO all seaGsh. | WE. | Indies 1828. | S l.p Wal.pl.as.ra.1* |
| | 2599. CA'REX. | | | | | | |
| 28715 | - podocarpa R.Sr. | fruit-stalked | £ A cu | Jn.jl | Br | N. Amer. 1827. | Dp |
| 28716 | - VahliaSchir. | Vahl's | £ A cu | f au | Br | Scotland moun. | D p' Eng. bot. 2666 |
| 28717 | 23014a depauperata Good, | few-flowered | Jt A cu | 1 my.jn | Ap | Eng. dr. wo. | D &1 Eng. bot 1096 |
| | 1271. CA'SSIA. | | | | | | |
| S8718 | 10690a HerbertaLindl. | Herbert's | ■■■ or | 9 n | Y | Barbadoes!828? S | ■■■ 10d |
| 23719 | - brevifolia Lam. | short-leaved | ■■■ or ... jn jl | Y | | Madagaa. 1824. S 8.1 | |
| | 2659. CASTA^NEA. | | | | | | |
| 88720 | . In&ica Jlox. | Indian | 1O o r | 40 | -- | E Indies | 1887. |
| | 3437.*1269a. C ASTANOSPE'RUMUM Cun. | MORETON BAY CHESTNUT. (Castanea, chestnut, sperma, seed.) £<• | i L J f r | 40 | ... Saf | N. Holl. 1828. | LI Bot mis. 51,» |
| 88721 | - austr&le Cun. | southern | | | | | |
| | 8232. CATANA'NCHE 20213 csrblea | | | | | | |
| | 2 bf color Penny two-coloured | | ■ A or | 3 jl.o | W.B | gardens | 1827. D co |
| | 2553. C ATTLE^. | | | | | | |
| 88722 | - gut&ta LindL | epatted-ftwd | ■ ED or | 1 ... G.Bd.w.p | Brazil | 1827. Dp.r.w Bot reg. 1406 | |
| 88723 | - elatior Lindl. | taller | ■ CS3 or | 1 ... G.Bd.w.p | Brazil | 1827. Dpjr.w | |
| | 1812. CE'LSU. | | | | | | |
| 28724 | . heterophylla Pert. | various-leaved | ■ or | ... jl.s | Ysh | — | 1829. S co |
| | 2434. CENTAUREV*. | | | | | | |
| 88725 | . argutaAfees | ahiTp-notched | ■ — | ... jU | Str | Canaries 1829. | S s.1 |
| | 3438*. 2411a. CENTROCA'RPHA D.Don. (Kenton, sharp point, karphe, chaff; paicee bristly.) 19.3. Camp. | | | | | | |
| | aubtomenosfaD. Don rather downyfr A or 3 aus Y N. Amer. 1802. D co | | | | | | |
| | triloba D. Don three-lobed | ED or '4 aus | Y | N. Amer. 1699. S co | | | |
| | Hudb't'-ckia triloba L. | | | | | | |
| | hirta D. Don hairy | ■ A or | 8 jn.n | Y | N. Amer. 1714. D co | | |
| | Rudbtfeksa hirta Sw.Jl.gar. | ■ il.au | Y | N. Amer. 176a D co | | | |
| | fulgida D. Don fulgid | £ A or | ■ il.au | Y | Bot mag. 1821. D co | | |
| | chrysomela D. Don yell.&dark.^tvdf | A or | ■ ■ | Y | | | |
| | acutifolia Swt's hort. brit. ed. 2 p. 310. | Riudbe'ekia Newm&hi Hort. | Loudon's hort brit p. 358. | | | | |
| 88726 | 2 chrys&ntha Dens, yellow-flowered | ■ A " S jl.s | Y | gardens | ... D co | | |
| | spathufta I. Don <pathulate->rf | Alor | S au.n | Y | N. Amer. 182% B p.1 | | |
| | gracilis D. Don slender | Jt A or | 1 au.n | Y | N. Amer. 1825. D ■ | | |
| | radula D. Don rasp | Q) or | | | | | |
| 88728 | . grandiflbra Swt, | great.flowered | £ A " | ■ ■ | Y | N. Amer. 1830. D e o | Sw.fl.gar.2.«87 |
| | Rudbeckfa nudicaulis Nut. | not of others. | | | | | |
| | 3439. •2147a. CENTROCLI'NIUM D.Don. (Kenton, sharp point, kline, bed.) 19. 2. Comptditaj Labiatiflor. | | | | | | |
| 88729 | - reflidrum Hook. | TcAxedscaled | (O or 2 au | Ro | Peru | 1830. S It | Bot mag. 311* |
| | Onfseris salicifolia of Hum. \$ Kth. | is thought to be near akin to this. | | | | | |
| 88730 | - apprcssum Hook. | appressetUclf/crf* | Q o r | 2 jn | Ro | Peru | 1830. S It |
| | " Differs from C. albicans D. Don, in its entire leaves; and from Onoserisangustifolia Hum. \$ A** | | | | | | |
| | in its larger and broader foliage." | | | | | | |
| | 3412. CERATOCHrLUS. | | | | | | |
| 88731 | 28531a oculatus B. C. | eyed | ■■■■■ | 1 jn | Y.Spot Xaiapa | 1829. | D p.r.w Bot cab. 176* |
| | 1472. CE'REU& | | | | | | |
| 88732 | 12565a Colvillt Swt | Coivtl's | ■■■■■ | jn.o | Ro | Eng. hybrid | ■ S 1 |
| 88733 | I'257a polygnous Isam. | many.angled | ■■■■■ | 10 | W | StDomin.1826. | |
| 28733* | 1257^ rokitceus Lk. S/0. | rosy | ■■■■■ | | Ro | ... 1826. | |
| 28735 | VKlic Depjet Lk. & O. | Deppe's | ■■■■■ | | Mexico | 1826. | P.J |
| 28736 | 12572f incrustatus Lk. | \$0. incrusted | ■■■■■ | | | 1826. | P.1 |
| 88737 | 12372U valdivia Haw. | stott | ■■■■■ | 23 gr 1* | S, Amer. | 1826. C s.p | |
| | Plants of three species of this genus were sent, in 1830, by M. Decandolle, to the collection <» of the Horticultural Society; their names are cincrascens, leptacanthus, and leptophis. See under Aiaw millaria, below. | | | | | | |
| | 778. CEROPETGIA. | | | | | | |
| 88738 | • illegans Wai. | elegant | J-Et QF 20 | au.o | P | v Et Indies, H28 U Dn 1 ih | Bot mag. »W' |
| | 583. CEySTRUM. | | | | | | |
| 88739 | . acuminatum Swt. | acuminated | *1 Jpr | 6 s.o | Gsh.Y Mexico | 1824. C co | |
| | 3440. •23560.CH.ET A CfLt L^N A D.Don. (ChaueM'H,cMaina, coverng; points of invol. scales.) 19.2. Com, | | | | | | |
| 28740 | - odorata D. Don | sweet-scented | £ i. or | 1 aus | R | Chile 1830. S s.1 | |

- | | | | | | | | | |
|--|-----------------|--|-------------|-----------------|----------------|--------------|--------------|---------------------|
| CHEILA'NTHES. | | | | | | | | |
| - crenulata Lk. Sf 0.? crenulated | £ (29 or | L | .. | Br. | .. | .1831. | D | p.i |
| - cuneata Jt * 0. ? wedge-leaved | £ [A]or | .. | .. | Br. | .. | .1831. | D | p.l |
| - repens Kaulf. creeping | £ 2 or | .. | .. | Br. | W. Indies | 1824. | D | p.l |
| - spectabilis Kaulf. showy | £ | In.o | .. | Br | Brazil | 1829. | D | p.l |
| *1691a. CHILOCTDIA R. Br. CHILODIA. (<i>ChcUos</i> , a lip, <i>odous</i> , a tooth; lip toothed.) | 14.1. Labiates. | | | | | | | |
| - scutellaria-pulidae &?r. Scutellaria-lik | J or 2) o.n | V | N. Holl. | 1829. | C | p.1 | | |
| - southern | *i or 3 | jn.au | N. Holl. | - | C | Ep | | |
| Forster's | A w | 31jl.au | C | Eng. | bogs,wds. | D co | | |
| CITRIO'BATUS CWK. ORANGE THORN. (<i>Cureum</i> , a citron, <i>batis</i> , a thorn; fruit and habit.) Pittos.2. | | | | | | | | |
| 287g | multiflora CUR. | many-flowered | £ L J or 3 | n.d. | .. | N. S. W. | 1818. | |
| pauciflora Can. | few-flowered | a J | .. | d | .. | N. Holl.tro. | 1822. | |
| Spiny shrubs, of which the first here enumerated is an inhabitant of shaded subhuinid thickets near Port Jackson, and of dark woods on the coast of New South Wales, where, in consequence of its bearing an abundance of small orange-coloured fruit, it is called by the colonists Orange Thorn. | | | | | | | | |
| 287*1 CLADA'NTHU& | | | | | | | | |
| - canescens Swt. | whitish | <t_J or 1 | mr.au | Y | Canaries | 1829. | S s.l | Sw.fl.gar.2.s.ic.in |
| CLADIA'NTHUS | | | | | | | | |
| acutiflora Swt. | acute-petaled | A A or \ | mr.au | W | N. Amer. | ... | O s.l | Pluk. aL 102.3 |
| virginica Bot. mag. | 941. | Hart. brit. No. 5675, not of <i>Lin.</i> — C. virginiana acutiflora Dec.prod. 3. 361 | | | | | Bot. mag. | 941 |
| longifolia Penny | long-leaved | A A or i | jajn | W | N. Amer. | 1827. | S p | |
| polyphylla Penny | many-leaved | A A or \ f | jn Pa.Pk | N. Amer. | 1827. | S p | | |
| Two species hitherto confused with <i>virginica</i> and <i>grandiflora</i> . | | | | | | | | |
| 28757 i.L? U CLE'MATIA | | | | | | | | |
| 28750 ^{i. L? U} <i vionioldes Schr. | Viorna-like | I | or 8 | jn.s | Li | N. Amer. | 1828. | S co |
| - odorata Wai. | fragrant | Wwf | Luor | - - - | - | E. Indies | 1831. | S 1 |
| Oat... IQHJ. ^/T V/^HJW | | | | | | | | |
| MJ/SQ ** " " VzJLJELJL IUX* | | | | | | | | |
| - speciorfssimaDqapi? most showy | O or ... | jn.s | P | Mexico | 1827. | S co | Bot reg. | 1312 |
| SB*- U19. COLCHICUM. | | | | | | | | |
| * ^o iw 9160a chionense Haw. MSS. | Chio | tf | A or \ o | P | Chio | .. | O co | |
| - 2X15a. COOLIGIGIA Mnl. | COOLIDCUJA. | | | | (Native name.) | | Euphorbitce. | |
| - odorifera Mol. | odoriferous | j | £ L J or 6? | .. | ;Gsh.W Chile | 1831. | Bot mis. | 40 |
| *1101rt. COLOPHO»NIA Kth. | COLOPHONIA. | (Yielding a resin like <i>Colophon</i> .) | 6.1. | Tercbinthce* 1. | | | | |
| mauritiafna Kth. | Mauritian | t C D e c. | P | Mauritius | 1826. | S 8.1 | | |
| 112. COMMEL'Nif. | | | | | | | | |
| 978a gracilis R.\$P. | slender | j | £ iAl or 1 | jl | B | Lima | J1 | Bot. mag. 30*7 |
| cucullata L. | hooded | iQJ or .., | jn.s | B | Brazil | Lp | | |
| 312. CONOSPE'RUM. | | | | | | | | |
| - t cajruleum R. Br. | blue | m v J or. | .. | .. | N. Holl. | 1830. | S Lp | |
| - \$ triplinervum/i. 2fr. three-nerved M i_j or. | .. | .. | .. | .. | N. Holl. | 1830. | S Lp | |
| 2106. CORONI'LLA 18992 glafica L. | | | | | | | | |
| 2 variegata Dot*. variegated-/i * LJ or 4 | my.s | Y | gardens | .. | C r.m | | | |
| *2330a. CORYA'NTHES Hook. HELMET-FLOWER. (<i>Korys</i> , helmet, <i>anthos</i> , flower; shape of lip.) Orchide* | | | | | | | | |
| * maculata Hook. spotted-lipped | J <E]spl | 1 | jn | Y.p | Demcraral | 829. | D p.r.w | Botmag. 3102* |
| J)r. Hooker refers to this genus also the Gongbra speci5sa Hook. Bot mag. 2755., and Ciongbra macrantha Hook. Bot. misc. 80.; but, as to the name of this genus, <i>Coryanthes</i> , seems to have overlooked its inadmissible nearness to <i>Corysanthes</i> of Brown. | | | | | | | | |
| 2047. CORYDA'LIS. | | | | | | | | |
| GeblerLcd. Gebler's | A? pr | .. | my.y | Pa.Y | AlUi | 1827. | S co | |
| 187a bicalcarate Haw. MSS. | 2-spurred | A A or i | f.y | Pk | - - | - | O co | |
| CORYDALIS Fis. | various-leaved | ■ | fir | sp | Y.R | Danube | 1829. | L co |
| II CO'STUS | | | | | | | | |
| *J9a discolor Rose. | 2-co\ouTei-lyd | £ ESI or 4 | my.s | W | Maranh. | 1823. | D s.1. | Rose, scit 81 |
| 28772 151a COTONEA'STER Wai. | | | | | | I | | |
| g773 nielanocarpa Fis. | dark-fruited | St | or 8 | | | | | |
| " nifflifera Jaz. | loose-flowered | St | or | | | | | |
| " frigida Wai. | frigid | St | .. | | | | | |
| " Nfosis Ham. in Prod. fl. nep. | round-leaved | * | or 3 | ap.my | W | Nepal | 182a G 1 | Bot reg. 1187 |
| " rotundiflta Wai. | round-leaved | * | or | ap.my | W | Nepal | 182a G 1 | Bot reg. 1187 |
| " taicrrphfla 2 U a 6rsi Lindl. in Bot reg. | taicrrphfla | 2 U a 6rsi Lindl. in Bot reg. | 187. | | | | | |
| " vulgaris Lindl. | common | a | or 4 | my.jl | Pk | Wales | ch.cl S 1 | |
| " Nummularia Lindl | NummuL-fvtf | * | or 10 | ap.my | W.G | Nepal | 1824. G 1 1 | |
| 88777 TPIS. | | | | | | | | |
| macrorhiza Her. B. | long-rooted | Jls | Y | Madeira | 1829. | S s.1. | Bot mag. | 1968 |

964. CRPNUM.										
28778	7807a nūrpurasclerfc	purplish	EJ or	... my.au	F	Fernan.	Pol82&	O	r.m	
S8779	7821a Lancei Herb.	Lance's	EJ or	... jn.o	R.w	Surinam	1825.	O	r.m	
28780	7850a algoense Herb.	Algoa Bay	Ig or	... au.o	R.w	C. G. H.	1826.	O	r.m	
28781	78506 califrum Herb.	Caffe	BS or	... au.o	R.w	C. G. H.	1825.	O	r.m	
28782	- BacbniHerb.	Bacon's	E1 or	... jn.8	R.w	Eng. hyb.	1826.	O	r.m	
116. CIUTCUS 1010 v^rnus										
	5 leucorhyncus Sab.	white-beaked	A or	J mr.ap	W.B	seedling	...	O co	Bot reg.	1416
	5 pictus Sab.	painted	A or	§ mr	P.w.			O co	Bot reg.	1440
28783	- . Thomaaii Ten.	Thomas's	A or	J sp	B	Naples	1830.	O co		
28784	- . odbrus Ten.	odorous	A or	} sp		Naples	1830.	O co		
28785	- . suaveolens Ten.	sweet-scented	A or	§ sp		Naples	1830.	O co		
28786	- . Imperati Ten.	Imperati's	A or	Jf.mr	Li	Naples	1830.	O 81	Sw. fl. gar.	
	" recurvus James Donn,	in his Hort cant,	seven years before							Mr. Haworth.
28787	1015a varieg'itus Hoppe	variegated	A or	*f.ap	Y.p	Levant	1829.	O co	H.&H.	
28788	- . aureus&w.	golden	A or	*f.ap	Go	Greece	-	O co		
3336. CUMMI'NG/i*.										
28789	- . trimaculata Dbn	three-spotted	If	.A el	f d	B	Chile	1829.	O p.I	Sw.fl.gar.2.* ⁸⁸
28790	- . tenella D. Don	delicate	tf	_Al or	\$ n.d	B	Chile	1829.	O p.l	
115a CUPA ^X NM.										
S8791	9352a setifera Cun.	bristle-bearing	I	i_J or	20 nd					MoretonB.1830. C p.l
1454. CITFHEA.										
28792	12213a Llavea Dec.	LUve's	i	■ _J cu	1} jn	C	Mexico	1830.	S Lp	Bot reg. 138b'
14, CURCITMA.										
28793	- . parviflora Wai.	small-flowered	■	■ ■	f of	W.v	Prome	1828.	D s.l	Wal.pLas.ra.5
455. CY'CLAMEN.										
38794	- . ibericum Goldie	Georgian	A A or	I	-					Asiatic G. 1831. S Lp
761. CYNA'NCHUM.										
28795	- . A'rgchl Del	Arghel	i	ll_	cu	Gsh.W	Egypt	1831.	C l.p	Del. eg. 20.8
28796	- . oleiflum Nectoux									
163. CYPE ^X RUS.										
28797	- . kyllingoidej Vahl	Kyllinga-like	M A cu	} jl.au	I		N. Amer.	1828.	D p.l	Rtb. gr. 4.5
965. CYRTA'NTHUS.										
28798	786Sa carneus Lindl.	flesh-coloured	tf	tAJ or	1 au	I	C. G. H.	...	O r.m	Bot reg. I* ⁶⁸
2541. CYRTOP(TDIUM.										
28799	- . flavum Lk. £ 0. f	yellow	■	■ ..	I ..	T				1831. D p.l
1964. CYTISUa										
28800	17538a elegans Ait.	elegant	■	LJ or	3 flu	I	C. G.H.	1821.	S s.l	
2691. DACRY'DIUM.										
28801	23650a dātum Wai.	lofty	■	LJ or	20 ..	-	Pulo Pen.	1830.	C l.p	
346. 2048a. DACTYLCA'PNOS WaL {Duktylos, finger, kapnot, fumitory; berries finger-shaped.) <i>Futnafi</i> ^{'''}										
	- . halicrifolia Wai.	£. _A or	3 au.o	Y.Br	Nepal	1831.	S It			Sw.fl.gr. ^{ar4}
										Di'lytra scandens Don prod. fl. nep.
236. DA'CTYLIS 1958 glomcrāta										
	2 variegāta Dent,	variegated-tod"	Jk A or	I jn.jl	Ap	gardens	...	D al		
1263. DAVIE!*S/il.										
28803	. . lineāris B. C	linear-leaved	«L LJ or	1 m y	Y N.	S.W.	1827.	C 81	Bot cab.	161 ^s
1583. DELrMA.										
28804	. . nftida Vahl	ihinlŋ4eaved	£. CD or	I -	I		Trinidad	1830.	C l.p	.
9511. DENDRCTBIUM.										
28805	. . longicōrnu Lindl.	long-spurred	^ (23 or	1 my	W	Nepal	1828.	Dp.r.w	Botreg.lfl ⁵	
28806	- . macrostachyum Lindl.	long-spiked	* E3 or	-	G	Ceylon	1829.	Dpr.w		
28807	. . moschMum Wai.	musk-scented	£ E\ or	-	Pa.Y	Pegu	1828.	Dp.r.w		
1386. DIA'NTHUS.										
11385	barbatus									
	2latiflbus Ser.	broad-leaved	£ A >pl	I J ^{'''}	s					
	pumilus Hort.									
28808	1139fa nknu Suit.	dwarf	£ A or	gjl.s	C	Switzerl.	1820.	S BI		
28809	1144/1 ciliatus Guu.	ciliated	£ A or	1 jn.au	Pk	Naples	1829.	S co	Gus.	
2S810	1147 ilabuchormeflsir.<?d.	Buchormian	£ A or	...jl.au	R	Russia	1826.	S s.l		
28811	11467< erubescens (?)	blushing	£ A or	...jl.au	Bh	Pyrenees	1825.	S s.l		
	11469 Fischer* Spr.									
	2 albus Penny	white	£ A or	I jn.o	W	gardens	1830.	C co		
1004. DICHORIZA'NDRA.										
S8812	. . pfcta B. C.		£ O or	I so	I	S. Amer.	1830.	D Itl	Bot cab. I ⁶⁶⁷	
2934. DICKSOPSIA.										
£8813	. . fldUntoldes Hum.	Adiantum-like	* El or	2 au.d	Br	W.Indies	1828.	D p.l		
1308. DICTA'MNUS.										
S6814	10909a angustifillus Swt.	narrow.leaved	■ A or	,2 my.jl	Li.	Altai	1821.	S S.1.S		

FIRST ADDITIONAL SUPPLEMENT.

537

- | | | | |
|--|---|---|--------------------|
| 2447. | *!48a. DI ^V ETES Sat. | (Dis, twice, etes, an associate; related to /Via and Mor*^r.) | J. Mo* |
| * ^{OB15} | - bicolorSttf.drt* A/S. 55. two-coloured. Allor 2 jlo Y.D. | | > Lp Bot reg. 1404 |
| | /Vis bicolor Lindl. in Hot reg. | | |
| | fridoldes Swt. Iris-like £ AJ or ap.my W.B. C. G. H. 1758. D i n Bot mag. 693 | | |
| 88817 | Pris m o r a l e s Ker& Hort brit p. 19. No. 1267.; Mor<x iridioides 2?ol. mag. Mora? a odorata Hook. | | |
| | catenulata Swt. chAn-dotted £ AJ or If ap.au W.B. Madagas. 182f. D Lp Bot. reg. 1074 | | |
| | More i catenulata Bot. reg. | | |
| 454. DODECA'THEON. | | | |
| 3816 | McadVi | | |
| 1 | IU'dnum Swt. | \A*c-flowered A or 1 ap.jn Li Virginia 1744. D 1 Bot mag. 12 | |
| 2 | albiflorum Swt. | white-flowered A or 1 ap.jn W gardens 1824. D 1 Bot cab. 1489 | |
| 3 | elegans Swt. | elegant A or 1 ap.jl Ro gardens 1827. D 1 Sw.fl.gar.2.s.60 | |
| 4 | giganteum Swt. | gigantic A or 2 ap.jl Li gardens 1819. D 1 | |
| 28060 | integrifolium Mx. | entire-leaved A or 2 ap.jn L.P. N. Amer. 1829. D Lp Pluk.aL76.6 | |
| 1685. DRACOCETHALUM. | | | |
| 15204 | altailnse Lax. | Altaic L A or ^ jl B Altai 1759. D r.m Sw.fl.gar.2.e.57 | |
| | grandiflurum Hort. brii. No. 15204. | | |
| 512. DRACOPHY'LLUM. | | | |
| | t capifatum R. Br. headed | H-Jor. N. Holl 1830. C Lp | |
| 1077. DRI'MIA. | | | |
| | villosa Lindl. villous tf A or my.jn G.St C. G. H. 1826. O s.p Bot reg. 1346 | | |
| | lanceolata (R. S.) spear-leaved tf L A or au.o Y.o C. G. H. 1774. O s.p And. rep. 2 ^o o | | |
| | l'chenal/Vi reflexa Andr. bot. rep. 299. | | |
| | longipedunculata long peduncled tf A or s.o G.p C. G. H. 1800. O r p Sch. ma 10? | | |
| 1602. DRPMYS. | | | |
| | chile'nsis Mien | Chilian ± Ljfral2 ... W Chile 1829. C l.p | |
| 8054. DRUMMCND/J Dec. DRUMMONDIA. (Mr. Drummond, botanical Collector.) 5.2. Sax/fragrabe<r. 1. | | | |
| | mitelloides Dee. | Mitella-like £ A cu i jn Ysh Rocky Mo. ... D p Bot. mag. 1829 | |
| | Mitella pentandra Hook. Hort. brit. No. 28229 | | |
| 326. DRYA'NDR.*. | | | |
| | tfoliolata? R.Br. leafleted *i_J or - - - N. Holl. 18%. C l.p | | |
| | t arctotoides R. Br. Arctotis-like il or - - - N. Holl. 1830. C l.p | | |
| | t calophylla R. Br. beautiful-lvd • i_J or - - - N. Holl. 1830. C l.p | | |
| 335a ECHEVETOM. | | | |
| | larida/iaw. | dingy-leaved ■ 23 or 1 jl S Mexico 1830. C s.l | |
| 34G8. ECHINA'CEA Moen. | | | |
| | 4)uprirea D. Don purple-Jftwl itudbeckw purpurea L. | ■ ■ ■ 4 jl.o R N. Amer. 1G99. D co Bot mag.S | |
| | serotina D. Don late-flowering illudWckiri serotina Sw.fl.gar. 4. | ■ ■ ■ 3 au.o R N. Amer. 1816. D e o Sw.fl.gar4 | |
| | fiapifolia Swt. Rape-leaved Rudbt/Gkin napifolia Kth. | ■ -AJor 2 jl.au Y N. Spain 1824. D p.l | |
| | heterophy"lla D.Z.0/i various-lvd intermedia Penny intermediate | ■ ■ or ljs.n R Mexico 1828. D e o Sw.fl.gar.2.8.32 | |
| | Intermediate between purpurea and serotina, and excels both in beauty.—Penny. | A or 4 au.o R Eng. hyb. 1826. D co | |
| 3359. ECHINOCA'CTUS. | | | |
| | i tephraanthos Lk. ashy-spined | ■ ■ ■ - - - Brazil 1825. O s.p | |
| | i inflate Gill. | ■ ■ ■ - - - Chile 1828. O s.p | |
| | subgibbsa Haw. subgibbous ■ 23 gr - - - Valpar. 1830. O s.p | | |
| 300. ELODE ^V A. | | | |
| | pulch&lajf.C. pretty | ■ - - - - E. Indies 1831. | |
| 254. EPIDE'NDRUM. | | | |
| | 30odoratis8imumiumfl.6weete9t-gctd^ El^ra 1 su G,Y RIO Jan. ^{mettem} 1827. D p.r.w Bot. reg.1415 | | |
| | Encyclia ptkeni Hook. in Bot mag. 3013., Macradenia lutescens Bot. cab. 1556., but not of Bot. reg. | | |
| | rlgidum Jac. rigid-petaled £/S/c\x 1 o Yen.G Dominical 826. D p.r.w Bot. cab. 1600 | | |
| 1 EPILO ^V BIUM 10054 hirsMum | | | |
| | 2 variegatum Den*, variegated-/^ ^ A or 4jl.au Ro England wat pi. Sk co | | |
| S3to4 390. J?PIME*DIUM. | | | |
| 3311a hexandrum Hook. | six-stamened | ^ A cu I ^p*m y " N. Amer. 1827. D s.l Hook. am. U | |
| | + | | |
| | purple | k A or I i" " " En <- *htL Wot D l | |
| " ^ t S S S B H ^ | Drummond'sar | A cu "1 ap Br Scotland aL bog. D p.1 | |
| 10b5ndJac. | charming | « U o r 2*myjn Pk C. G. H. 1827. C <p Bot.caUWW | |

9814v&gans


- 5450*. 1224a. FRANCOM Cav. (F. Franco, of Valencia, a promoter of botany in the 16th cent) & 4. *Galadnece*.
 - appendiculata Cav. appended £ A :: 2 jLau P Chile 1830. D p.1
 - sonchifolia W. Sowthistle-lvd £ A :: 2 jLau P Chile 1830. D p.1
 - rambsa D.Don branched £ A :: 2 jLau W Chile 1831. D p.1
1018. FRITILLA'RIA 8447 meleagris
 2 multiplex Penny full-flowered tf A or 1 mr.my P gardens ... O co
1188. FU'CHS/*.
 28204a bacillaris Lindl. vr&nd-branched ft LJ or 5 su Ro Mexico 1829. C p.1 Bot reg. 1480
1008. FU'NK/i*.
 - lanceefolia Dens, lance-leaved i A or 2 jl.au Li Japan 182 . D 8.1 Bot cab. 1658
 /femeroocallis lanceae&Jia Bot. cab. Iba8.
1052. GA'GE/f
 " podolica R. \$ S. Podolian tf A or :: my P Podolia 1827. O co
 Liotardi R. fr S. Liotard'u tf A or :: my S. Europe 1825. O co Ster. den. 2.3
 Ornithogalum fistulbrum fled. /ft. 221. bulbifera it. \$S. bulb-bearing tf ^ or :: my Taurica 1829. O co Rchb.ic.117
- 2897 2 4 1 1 G A L A 'R D / J * Richardson's Penny Richardson's 1J my.o O N. Amer. 1829. D p
1972. GALFGA 17652 pénica
 2 Uldcina Penny Lilac 4 or 3 jns. Li Persia 1830. D co
- 48 GALIPETA.
 462a odoratissima Lindl. most fragrant :: f r a 2 my W Rio Jan. ... C l.p Bot. reg. 1420
- 88879 248 GALLIUM hairy £ _J cu 1 au.o Gsh.W Teneriffe 1830. D co
- 225 GALPHI'MU. hairy t Q o r ... mr.jn Y Mexico 1824. C l.p
 hishta Cav.
- 1683a. GARDOQUIM R. & P. GARDOQUM. (D. Diego Gatdoau, a noble Spaniard) Labiate.
 - Gilliesti Grab. Gillies's «i_J... 2 ... Li Chile 1828. S ...
 - discolor Kth. two-coloured, apji P Caraccas 1827. S ...
 G. origanoides of Reichenbach is a species of Lantana (*Brahamini* in Bot. reg. 1300.)
- 8a. GASTROCHPLUS Wai. GASTROCHFLUS. {Gaster, belly, f^i/o*, lip; bellied-lip.) 1.2. ScUaminecc.)
 - - G. prettissimus Wai. pretties ^ El or 2 au Y.Pk Rangoon 1828. D s.1 Wal.p.Las.rar.i
- OJJS*. 747a. GAUDICHAU'D/J H. B. & Kth. (Charles Gaudichaud, accomp. Freycinet) 5.1. Mahtigitice'tB. 4.
 *"** - - cymachodes/LA*JL7A. Dogsbanc-likj. a or 10 ... Y Mexico 1824. C r.inH.&B. n.s. 445
- OOQD, I¹⁸*. GAU'R.A.
- 4*8510042a angustifolia Mx. narrow-leaved £ A P' 2 jLs Pk - - - S s.l
1. 1073a. GEITONOPLE'SIUM Cun. (Geiton, neighbour, plesion, near; in affinity and habitat)
 8966 cymtoum R. Br. cymflowered \$ i_J cu 1 ... G N. S. W. 1825. C p.1 Bot. xmag. 3131
 Luzunagi cymbosa Brown in Prod. 1782. Hort brit No. 8967.
- 8967 montanum R. Br. mountain £ iAI or 1 ... G N. Holl 1820. C s.p
 Luzurilaga montana Brown in Prod. Mont. brit No. 8967.
- iscrum Cun. rough-angled ^ iAI cu N. Holl. 1831
 The Peruvian genus Luzurilaga, No. 1U7&, should now be removed from Hort brit, no species of it having ever existed in Britain.
1187. GETLA. Nat Ord. Olacine*, not Onagritria as in Hort brit. p. 155.
 10070a oblongifolia Cun. oblong-leaved f LJ or 6 jn Wsh.G N. Holl 1823. C s.p
3406. GEORG P^ 21591 sup^rflua
 Blood red anemone-flowered variety, Maund's Botanic Garden, 297.
 Painted lady anemone-flowered variety, Maund's Botanic Garden, 329.
1932. GERANIUM.
 1/23ta albilobatum Hook. white-flowered ^ A or 1] su Wsh N. Amer. 1827. D co Bot. mag. 3J24
 Has been called *G. maculatum*, and a variety of *G. angulatum*: was brought home at the return of Franklin's second expedition.
- apicata L. \ 0. spiked ACS or 1831. O p.1
1530. GENUM chilense Balbis
 2 grandiflorum Lindl. large-flwd £ A 8Pl 11 jn.o S S s.l Bot reg. 1348 *
499. GI'L.U.
 " pdngens Don. prickly O cu 11 jn.au B N. Amer. 1827. S s.l Bot. mag. 2l⁸⁷⁷
 cuLeaved O .. i jl l*k Chile 1831. S pi
- iiJ? GLADPOLUS.
 1w3a cochleatus Shv. spoon-lipped tf uMor 1} mr W.R C.G.H. 1829. O s.p.l Sw.fl.gar.Cs 14A
 870? natalensis Rinnwardi Natal tf iAJ H* 1} au S.v Natal Riv.1830. O p.l Dot. mag. K V2. A
 psittaculus Lindl. in Bot reg. 1442. [in note 3084
- GLECHOMA 14935 Aederacea
 2 rbsea Dent. rosy-/itec %. A or i mr.my RaLi Eng Ips D co
 3 variegata D*«i»*. variegated Witf JL, A or \ mr.my B England
- S080. GLY'CINE
 " biloba Lindl. two-lobed L J or 20 n V Mexico 1827. S l.p Bot. reg 1418
- 1251 GOMPHOLO^BIUM.
 * - X KnightidRum LindL Knight's tt i_J el au. Ro N. Holl 1830. S p.1 Bot reg. 1168

8455. ^2558 GOVE^xNJi* Lindl. GOVENIA. (*J. R. Gowen*, an originator of splendid hybrids in Ericce.) *Orchidef-*
 28896 - - suplrba Lindl. superb A El or 1 mr Y Xalapa 1828. D >lt Bot ~~ab.~~ 17W
 Probably the *Maxillaria superba* oTLaLlave.
53. GRATPOLA.
 28897 - - tetragona Hook. squarestmd ■ □ □ 1 au B B. Ayres 1830. D p.l Bot.mag. ^{313*}
316. GREVI'LLEi*.
 28898 2617a planifblia R. Br. MSS. flat-leaved* J or 2 ap.my Ro N. Holl. 1823. C p.l Bot reg. ¹³⁸³
 concinna Lindl. not of others, *Seymoixrie* Swt in Gard. mag. vii. p. 506.
- 28899 6635a Calfci R. Br. Caley's * i_J or 70 ... O MoretonB.1830. C Lp
Alechnidlia Cun. MSS. Blcchnum-lvd
- 28900 2636a robusta Cun. robust, or *Stlk oakf.* i_J or 5 jn.s R Port.Jack. 1829. C p.l Bot. mag. <>>"
2904. GYMNOGRA'MMA.
 28901 - - tomentbsa Desv. tomentose ■ EJor - - - Brazil 1831. D p.l
2964. GYMNO'STOMUM.
 28902 - - Wilstlm'Hook. Wilson's small patches □ ja Y.c Britain fields - - -
3456. >1645fl. GYROSTETMON Desf. (*Gyros*, circle, *stemon*, stamen j concentric arrangement.) 22. 12. *TUideat*
 28903 - - attenuatu Hook. attenuated f LJ C U 25. MoretonB 1830. C Lp Bot. mis. 53
Codonocarpus australis Cyp. The male flowers of this plant have never been seen, and when they should prove unlike those of *Gyrostemon*, the plant will be the type of the genus *Codonocarpus* ^
2483. HABENA^NRIA.
 28904 22057a cordata Hook. cordate ACS or - - G Madeira 1830. O p.l
3457. *729a. HABLITZL* Bieb. (*M. C. Von Hablitzx*, author of travels in the Crimea.) 5. 1. *Chenopdex.*
 28905 - - famnoldes Bieb. Tamnus-like □ A - - jLo W Caucasus 1828. S
975. HABRA'NTHUS.
 28906 - - BagnAldi Herb. Bagnold's tf ^ jor | n Y Chile 1829. O s.l Bot reg. Hg
 28907 - - Andersön* Herb. Anderson's tf | Alor ap.my Y.R M. Video 1829. O s.l Bot. reg. ^{1383*}
 28908 - - r&seus Swt. rosy f Al or | jn Ro Chiloe 1828. O r.l Sw.fl.gar.2.*
 28909 - - pall id us B. C. pale-flowered tf | Alor | jn Pk Valparai. 1830. O r.m Bot cab. ^{115*}
 28910 - - phycellttides Herb. Phycella-like tf | Alor | o S Chile 1805. O r.m Bot. reg. If.
 28911 - - pdmilus B. C. dwarf fl | Alor i | au R Chile 1631. O s.l Bot. cab. ^{177*}
17. H ■ TFL.
 28912 - - X/aurina R. Br. LaureUfeaveo* II i | or - - - N. Hoi. 1830. C Lp
 28913 - - X Bixterf R. Br. Baxter's * i | or - - - N. Hoi. 1830. C Lp
 28914 - - X tuberculata R.Br. tubercled il i | lor - - - N. Hoi. 1830. C Lp
3458. *2337a. H AXTOW// Caley. (*John Haxton*, gardener attached to Macartney's expedition to China.)
 argophylla Caley. A ster argophy illus Lab. Mr. David Don describes and defines this genus i! V^
 Edinburgh New Philosophical Journal, Oct. 1831, p. 272., and refers to it the ^ster argopW
Lab., *viscbsus* *Lab.*, *phlogopappus* *Lab.*, *stcllulatus* *Lab.*, and *tomeutulus* *Willd.* and *Hart*, kew-
3459. <974a. HAYLO'CKM Herb. HAYLOCKIA. (*Matthew Haylock*, gard. to Mr. Herbert) 6.1. *Amaryllidef.**
 28915 - - puflillaHerb. dwarf tf^Jor | 6 Str B. Ayres 1829. O s.l Bot.reg."
6. HEDYCHIUM.
 28916 - - urophyllum Wai. ? tailed-lvd ■ ES fra 5 s Y India 1828. D r.l Bot cafctf^{8*}
2409. HELIA'NTHUS.
 28917 - - pcfolfrtx Nut. long-petioled O or 3 au.n Y Arkansa 1826. S co
1993. HELI'CTERES.
 28918 17745a virgata Wai. twiggy Dor | - L E. Indies 1831. S Lp
 28919 - - elongata H. C. lengthened <<Dor | - L E. Indies 1831. S Lp
3460. *32&r. HEMICLI'DI A R. Br. (Probably from *heminu*, half, *kleio*, to shut up.) 4. 1. *ProteHcea*,
 Baxteri R Br. Baxter's << Lj or 3 jn Y Lucky Bay ... C p.l Botreg. ¹¹⁵
Dryandra falcata R. Br. Hort brit No. 2736.
1863. J/ESPEKIS.
 28920 1644a specibsa Swt. showy jg A or i ^P R o P Siberia 1829. S Iru
805. HEITCHERIi.
 289S1 6439a micrantha Dou. small-flowered £ A cu 2 jn.jl Ysh.G Columbia 1827. D e o Bot reg.
1597. HIBBE'R'T/J.
 28922 14117a flexuosa Cun. zigzag il LJ or 2 my.au Y N. Holl. 1823. C s.p
2014. HIBISCUS.
 28923 17926a splendens Fra. splendid #a pi | □ Ro N. Holl. 1828. C r.m Bot mag-
 28924* 1794/1 pentaspalmrus/fcrtero five-seeded ill - or 3 jLo Y Jamaica 1825. S r.m
 28925 nmSa Lindlef Wai. Lindley's ill - or 3 d P India 1828. C Lp
 28926 1794ito crinitus Wai. long-haired ■ Expi 3 s o YR Prome 1828. S p.1
 28927 179&UI africinus Roth. African C or 2 jno W.p Africal 1826. S co
 28928 17998a Macleay Ante Bane. MacLeay's ■ ... au.s Y W.Indies 1827. C Lp
3461. *2897A. HOLIGA'RNA Rox. HOLIGARXA. (Name in Karnata.) 23.1. - - -
- 28929 - - longifolia Rox. long-leaved J • ec 60 ... W E. Indies 1828. C r.m Rox.
3392. HOSA'CKZ/I.
 28930 - - decambens Benth. decumbent)£ A cu | i jl.s Y N. Amer. 1827. ■ s.l
 28931 - - parviflura Benth. small-flowered £ A cu | i jl.s Y N. Amer. 1827. ■ s.l
343. HOUSTON/A
 29932 2858a longifolia Gae. long-leaved]£ A cu ■ - my.au W N. Amer. ... S p " ■ mag
 28933 2858A albi flora Graft. ? white-flowered j£ A P' . my.au W N. Amer. 1828. D s.p

1340. *HOVEA* 17281 lanceolate.
 2 *linehris* *Lindl.* *lineatitwoö* *i j o r* *3* *mr* *P* *N. Holl.* *-* *S p.l* *Cot. reg. 1427*
 Is this the *H. linehris* of *Browns the Forresta linearis* of *Smith*?
- 108^o *flyACI'NTHUA*
 -- *bruma* *is ifau*. *MSS.* winter *tf A or f f.mr* *Various* *--* *O co*
863. *HYDROCOTYL*
 6907a *prolifera* *Lk. \$ b.* *proliferous* *%, 23 un ft apjn G Brazil 1824. D p*
- 934^a, *HYMENOCALLIS*,
 - - *Staples** *Herb.* *Staples** *tf ^ or ... jnjl W Mexico 1826. O r.m*
2943. *HYMENOPHYLLUM*,
 - - *Wilsöni Hook.* *Wilson's* *f A el \ my.jn Br Brit moi. ro. D l.p*
2190. *ffTYPE'RICUM*,
 - - *Gebleri Led.* *Gebler's* *HI or ... jn.au Y Altai 1829. Sk s.l*
- ijij³ *16te. *HYPOE'LYTRUM* *Rich.* (*Elytron*, covering, *hypo*, under; bracteas under glumes.) 3.1. *Cyperieeat.* 1.
 B . . *argenteum* *Vahl* silvery *M GS cu ... jt.au W.P E. Indies 18ii4. D p.l*
- G98. *IMPATIENT*
 5697 *paMida Nut.* *p&le-Jlowercd* *O or 2ft jn.o Pa.Y N. Amer. 1812. S p Eng. bot. 937*
Noli me tangere Eng. bot. not of others.
- 5697o *borealis* *Surf.* *northern* *O or 3 Jn.o Y Denmark ... S p Fl. dan. 582*
 5698 *Noli me tangere L.* *Touch-me-not* *O or 3 jn.o Y.B EngL Essex... S p*
2072. *INDIGOTERA*,
sylvatica Sieb. *wood* ** |J or 8 Jn.jl Ro.Li N. Holl. 1825. C s.l Bot mag. 3000*
 Identical with No. 18658, but distinct from No. 18659, and is *I. spectosa Hortui*.
491. *IPOM(E'A.*
 - - *Sellbwñ Penny* *Scllow's* *| - - 10 - - - - 18J1. C pi*
142. *Hooker Penny* *Hooker's* *f A or 1| my.jn P N. Amer. 1826. D co*
tripetala B. M. 2886. not of Walter.
- 38913 2718. *JANVPHA*.
 23853a *angustifdiaPoAi* narrow-leaved *H • or 3 ... Br Brazil 1829. C r.m Fl. br.*
- *885fl *JABORO SA J. JABOROSA.* (*Jaborose*, Arabic name for the mandrake; afflBnity.) 5. 1. *Solu*
 - - *runcinata Lam.* *runcinate* *f LJ P' f plata 1831- D PJ*
- 99915 5889a *hdmilis A. Dec.* dwarf *f A or \ jn.s B S. France 1824, D p.l*
- ^ *JASMrNUM.*
 323a *pubgerum* *2. Dan* down-bearing *f. or 10 mr.o Y Nepal 1827. C 1 Bot ret. 14fig*
Wallachianam UndL
 Its old leaves are glabrous, but those recently expanded are pubescent, as ia, according to the *Bot. reg.*, its calyx as well. The general smoothness which, in the open air of Britain, the *wlan* assumes, led Professor Lindley to distinguish it from Mr. D. Don's *publgerum*.
994. *JUNCUS*.
 28947 8126 *capitatus Weigel* headed *- O?cu \ my.jl Ap Jersey san.gr. S s*
58. *JUSTI'C/y.*
aspéruula Wai. *roiiighish* *i D o r 2 su Pk India 1829. C Itl Bot. eah. 1681*
*quadrangularisflw**. *Bot mag. 2845., Hortbrit No. 27985.*
- 2 - *cirnea LindL* flesh-coloured ** O upl 4 aus F Rio Jan. 1827. C l.p Bot re* 1397*
 - - *guttata Wat* spotted *J E o r H ap.au Y.Spot E. Indie, 1828. D Lp Bolre* 133**
 - - *venüslaH'a/.* beautiful *< Q o r 5 s P Bengal ... C Lp Bot. reL 1380*
 - - *amabilis Hook.* lovely ** or No. 546. in Hort brit is but a synonyme of Na 547.*
- Dis^{5*} *2810a. *KAGENE'CK/i< R. & P. KAORNECKIA.* (*Conn/ Kageneck, a patron of botany.*) 22. 11. *2810a*
 - - *crat<goWesD.DofiHawthotn.hkel J or 60 ... W Chile 1831. LI*
- 2808, 12. *élégansWa.* elegant *2810a 1 ... P Pegu 1828. D s.l Wal.pl. as. ra.27*
- KENNE'DYif 19387 monophylla
 2 *longeracembsa Lindv. Ig-racemed J* *2810a 3 mr.au Pk N.S.W. 1828. S l.p Bot. reg. 1336*
- ^ISfl *plnlda*TW pale.flmoered* *tf ^ o r ft sp Pa.B C.G.H. 1782. O p.l Bot. reg. 1350*
 2 *minor Ltndl.* *smaller V 1Alor i sp Pa.B C.G.H. J782P O p.l Bot. reg. 314*
 Professor Lindley has remarked, *Bot. reg. 1350.*, that he has proof that 11350. represents the *I* pall-Hda* of *Hortus Kewensis*, which he identifies also with that of *Thunberg*; and he considers t. 314. of *Bot. reg.* to represent but a smaller variety of the same species; but remarks, "the supposed variety figured at t 287. of this work is altogether another species." Agreeably with this view, a new specific epithet, with the name of its author, are wanting for No. 9015. of *Hortus Britannicus*.
1887. *LAGERSTR<E*M7i< 13918 indica*
 2 *rosea B. C.* *rosy ft 2810a 12 aus Ro Chiha 1825. C r.l Bot. cab. 1765*
- 18955 319^c *1 h l u c u l*
 **** *a E. Br. related* *»i_J or - - - - N. Holl. 1830. C Lp*
R. Br. Heath-leaved *• LJ or - - - - N. Holl. 1830. C Lp*

FIRST ADDITIONAL SUPPLEMENT.

593

- | | | | | | | | | | |
|---|--|--|---------------|-----------|----------------|--------------|------------------|----------------|-----------------------------|
| S575. | MICRO'STYLIS. | | | | | | | | |
| . | versicolor Lindl. | changeable | jfc | EJ cu | 1 | Jn.o | O | China | 1830. D p.r.w Bot cab. 1751 |
| Lfpars priochillus B. C. | | | | | | | | | |
| 2834. MIMCTSA. | | | | | | | | | |
| £9083 . | . stipulata H. C. | stipuled | | - | - | - | - | E. Indies | 1831. S p.1 |
| 1783. MI'MULUS. | | | | | | | | | |
| 99024 . | . propinquus Lindl. | related | ■ | A or | ap.o | Y | | N. Amer. | 18S7. D p.l Bot. reg. ■■■ |
| 1156. MIMU^SOPS. | | | | | | | | | |
| 29025 9345a cyanocarpa Cun. | blue-fruited | i 1 Q or 6 s | | Gsh | N.co.N.H.1821. | C | p.l | | |
| 347a «2055a. MONNPNP^ R. & P. | MONNINA. | {Monnino, Count de Flora Blanca.) | 17.3. | PolyeM* | & J | | | | |
| 89026 . | - obtusifolia HgftA. | obtuse-leaved | * u J or 2 jn | Rsh.PPeru | 183a | S | p.1 | Bot. mag. 31** | |
| 2612. MTRUS. | | | | | | | | | |
| 29027 23105a calcar-galli Gun. | cock-spur | * i J ec | | | | | | N.S.W. | 1850. L r.m |
| ** Yellow wood vine " of the colonists. | It is a shrub which extends itself to a great length, and ma | | | | | | | | |
| eventually be found to belong to the genus Madhra (T6xylon Raf.), rather than to Mbrus. | | | | | | | | | |
| 746. HL | | | | | | | | | |
| 29028 . chinensis Swt. | dwarf | Chinese f • or 4 | | ■ | | China | 1829. | Sk r.l | |
| 1082. MUSCANRI. | | | | | | | | | |
| 29029 8993a pedunculare Haw. | peduncled | tf A or i | mr.ap | B | | | | O co | |
| 29030 89936 parviflorum Detf. | small-flowered | tf A or ... | ap.my | B | Sicily | 1827. | O co | | |
| 425. AfYOSO^TIS. | | | | | | | | | |
| 29031 3506a collina Hgfm. | hill | O pr 1 ap.my BY | Britain | dstpl | S co | Eng.bot | 2538 | | |
| From No. 3506. remove the figure cited, and substitute Eng. bot. 2629. | | | | | | | | | |
| 1489. JFY^RTUS. | | | | | | | | | |
| 29032 12761a melastomoides dm. | Melastonuulk | ± LJ or 15 | - | - | - | Moreton B... | C r.l | | |
| 933. JEVARCISSUS. | | | | | | | | | |
| 29033 7592a cernuus Haw. | drooping | If A or 1 mr.ap | Crea. | W Spain ? | - | O co | Sw.fl.gar.2.s.l^ | | |
| 2 corona plena Haw. | full-crowned | tf A or 1 mr.ap | Crea. | W Spain? | - | O co | Sw.fl.gar.2.8.1^ | | |
| II. A^jax Haw. | | | | | | | | | |
| 29034 7546a ddbius Pen. | doubtful | tf A or ... | ap | W | S. France | - | O co | Red. liL 420 | |
| 29035 75466 Cypri Haw. | Cyprian | tf A or 1 | mr.ap | W.Y | Cyprus | - | O co | Sw.fl.gar.2.« | |
| 2 corona plena Haw. | full-crowned | tf A or 1 | mr.ap | W.Y | gardens | - | O co | | |
| 890% 7558a intermedium Red. | intermediate | tf A or 1 | ap.my | Pav | Pyrenees | - | O co | Red. liL 427 | |
| 29037 - neglectus Ten. | neglected | tf A or ... | ap my | W.Y | Naples | 1830. | O co | | |
| XIV. Hermione Sal. & Haw. | | | | | | | | | |
| 29038 7537a stellkris Haw. | staxy-sepaled | tf A or 1 | my | We | - | - | - | 1629. | O co Sw.fl.gar.2.*^ |
| We place these species under Narcissus to i | | | | | | | | | |
| but Mr. Haworth, who has studied the species | | | | | | | | | |
| entitled Narcissinearum Monographia, a second | | | | | | | | | |
| into sixteen genera, three of which we here exhibit sectionally. | | | | | | | | | |
| 1819. tfASTURTIUM. | | | | | | | | | |
| £9039 16114a natans Dec. | floating | A A cu | 4 | my.s | Y | Siberia | 1827. | D co | Deles*. & 1* |
| 2496. NECTTIA. | | | | | | | | | |
| 29040 22557a gemmifera Sm. | butl-bearing | A A or | ■ | ■ | W | Ireland | mar. | O bo | |
| 488. NICOTJ/TJM. | | | | | | | | | |
| 29041 4010a nepal^n sis LA. ^-0. | Nepal | O or | ■ | ■ | Ro | Nepal | 1829. | S r.m | |
| 29042 40506 brasiliensis Lk.8/0. | Brazilian | O or | ■ | ■ | Ro | Brazil | 1825. | S r.m | |
| 2P(43) 4050c petiolata As. | fooUstalked | O or | ■ | jls | Ro | S. Amer. | 1829. | S r.m | |
| 29044 4050a sanguf nea Lk. 8f 0. | & car'et-flowered | O or | 4 | jls | S | S. Brazil | 1829. | S r.m | |
| 28081 glafica Grab. | glaucous | il J or | 20 | au.o | Y.G | RAyres | 1827. | C l.p | Bot mag. 2837 |
| 3474. *490a.? NIEREMBE RG./4 Kth. | NIEREMB. | {John Eusebius Nuremberg, a Spanish Jesuit.) | 5.1. | Soldnf | | | | | |
| 29045 . - Unari83fbliaGraA.Toadflax.lvd | O L_Jel | } | jl | | Wsh | Uruguay | 1831. | S p? | Bot. mag. 31^ |
| 2910. NOTHOCHLJFNA. | | | | | | | | | |
| 29046 . tenera Gill. | tenier-textured | OS or | } | - | Br | Mendoza | ... | D s.p | Bot. mag. 3055 |
| 29047 . sinuata Kautf. | sinuated | jf (23 or | ... | - | ... | Peru | 1831. | D p. | I |
| 1687. O'CYMUM. | | | | | | | | | |
| 29048 . montanum Hook. | mountain | O c u | | my.au | W | W.Indies | 1825. | S p.1 | Bot.mag. ■■■ |
| 1183.  | | | | | | | | | |
| 89049 10015a Wirona Lindl. | two-faced | O or | ■ | ■ | Pc | hybrid | 1830. | S co | Bot reg. 14^ |
| 29050 10039a Nuttallii Swt. | Nuttall's | k A or | ■ | my.au | W | N. Amer. | 1811. | D co | |
| 29051 10038a anis61oba Swt. | unequal-lobed | tf A or | ■ | W | | Chiloe | 1828. | S co | Sw.fl.gar.2.*^ |
| 29052 . - Sellowia Lk. & O. ? | Sellow's. | ... | ... | ... | M. Video | 1831. | ... co | | |
| 2540. ONCI'DIUM. | | | | | | | | | |
| 29053 . biconratum Hook. | two-horned | ■■■■■ | 1 | jn | Y.Br | Brazil | 1830. | D p.r.w | Bot. mag. Sti* |
| 1966. ONO'NIS section KATRI'DIUM. | | | | | | | | | |
| 29054 . . pedunculkrisLmrfl. | peduncled | tt-Jfr | 1 | ap | W.RoTeneriffel | 829. | S s.1 | Bot. reg. 1446 | |
| 431. ONO'SMA. | | | | | | | | | |
| 29055 . Gmelini Led. | Gmelin's | jg A or | 1 | ap.au | * Str | Altai | 182a | C s.1 | Led. alt.^2% |
| 29056 . polypliflum Led. | many-leaved | £ A or | 1 | j1.a | u Str | Tauri | 1829. | C s.1 | Led. alt. S» |
| 29057 . rfgidium Led | rigid | £ A or | 1 | il au | J Pa.Y | Tauri | 1826. | C s.1 | Led. alt. S» |

««*.	OPU'NTIA.						
*9058	12603a leucantha Lk. * 0. white-flowered tt. 23 gr	-	... W	S. Amer.	1824.	C 8.1	
2909	13603b Hernapdezi* Dec. Hernandez's tt. 23 gr	-	... W	Mexico	1827.	G 8.1	Hern.mex.78.ic
-	- myriacanthal.&.^O. myriad-spinedtt. Z3 gr	-	... Y?	Mexico	1830.	C 8.1	
-	- rbsa Dec. to y-towered tt. Z3 gr	-	... Ro	Mexico	1830.	C 8.1	Dec. cac. 15
-	- imbricata Dec. imbricated tt. Z3 gr	-	-	-	1820.	C -P	

Cereus imbricatus Haw.

Plants of four species of this genus, under the names decpiens, Stapelfe, exuvia, and foliata, were sent by M. Decandolle, in 1830, to the collection of the Horticultural Society. See under *Mammillaria*, above.

2481.	O'RCHIS.						
22471a	quadripunctata TVii. 4-spotted	A A or I	myjn P	Italy	1828.	O l.p	FLnap. 89
22471b	eaccata Ten. saccate	A A or f	myjn P	Sicily	1828.	O l.p	
22471c	pseudosambCicina TVn. Elder-stnelli&	A or f	...	P Italy	1888.	O l.p	FL nap. 86a
-	2 lutescens Ten. pale-yellow	A A or	...	Italy	1828.	O Lp F)	nap. 866
-	- fuscescens B. C. drying-brown	A Al cu	jn	Ysh Pensylv.	1831.	O p.lt	Bot. cab. 1748

1053.	ORNITHOGALUM.						
29067	- notatum R. S. brown-marked	tf Alor	jn.au	C. G. H.	1825	O r.m	Jac. c. 2.1& 13
-	- maculatum Jac. not of <i>Thun.</i>	-	-	-	-	-	-
-	- triglypum Red.	three-styled tf	A] or ... jn.jl	W.a.	1825.	O r.m	Red. lil. 417
-	- miniatum Jac.	red-stained tf	A] or ... jn.jl	Y	C.G. H.	1790.	O r.m Jac. ic. 2.438
-	- brachystachys F&.	short-spiked tf	A r	mr.ap	W Dauria	1821.	O r.m
8W78	- Brgfr Schle.	Bergius'fl	AJ or	mr.ap	W.G.	1816.	O r.m
-	- concinnum Sal.	neat tf	A] or f	ap.my	W Portugal	1797.	O r.m Bot. mag. 953

1764.	OROBA'NCHE.						
29073	- caryophyllacea Sm. dove-scented	/5 A cu	-	Din.P England	-	S 8.1	Eng.bot.263i

1181.	OSBE'CK/ii						
28195	glomerata Ztec.	-	-	-	-	-	-
-	2 albiflora Dec.	white-flowered M • or U dap	W	Braail	1821.	S p.l	Botcab.3&t
9978	nepalensis Hook.	-	-	-	-	-	-
-	2 albiflora Lindl.	white-flowered J* • or U au	W	Nepal	1829.	C p.l	Bot reg. 1475

29U74	2945. OSMUNDS.						
-	- palustris Lk. \$ 0. ? marsh	-	2 	-	1831.	D p.l	

2451.	OTHCNNA.						
22350	virginea L. virgin-tite flabelliflta 2tof.caA. 728.	81 L J or	3 Bn	Y	Africa	1821.	C Lp Bot cab. 728

11902a	crenata Jac.	notched/)*/**	AiA Jesc	3 s	Y	Lima	1829.
-	- Arracacha G. Don, Syst bot and gard. i. 756.	-	-	-	-	-	Sw.fl.gar.2..l&
976	11906a chinensis Haw. MSS. Chinese	SUiA Jpr	1 my.o	Y	China	-	D co
-	11909 Sims/iSwt Sims's	£ IAJor ft	ap.jn	C	Chile	1822.	O s.p Bot mag. 2415
-	- rosea Sims, Bot mag., not of others.	-	-	-	-	-	-
11910	rbsa Jac.	rosy floribanda Bot. reg. 1123.; not of Lvhmann or of Sweet.	JCiAlpr	i mr.jn	Ro	Chile	1826.
4077	11926a cyprea B. C.	copper.cld	LAj pr	ap.jn	Cop	CG.H.	1822.
S778	11932a mauritiana B. C.	Mauritian	tf	IAJor *	Pa.Ro	f. France	1810.
29079	11933a bipunctata Grah.	2-spotted	tf	Alor	5	Li Brazil	1825.
-	11933c Dépper B. C.	Deppe's	tf	Alel	mr.n	Cop.R Mexico	1827.
-	11933c floribunda Leh.	many.flowered	tf	JM or	ap.8	Ro Brazil	1829.
-	11933d palustris Hit.	marsh papilionacea Zuccarini	£	SI pr	... my.d	Li Brazil	1828.
-	11959a glandulosa Jac.	glandulous	tf	Alpr	h o.n	W C O. H.	1822.
-	11961a breviflropa Jac.	short-scaped	tf	LAJ pr	U f	W C G. H.	1823.
-	11962a rigidula Jac.	stiffish	tf	Al pr	1 Bn	W C G. H.	1822.
-	11963a laxula Thun.	loose	tf	Al pr	o.d	W C G. H.	1820.
-	H96tia strumbosa Jac.	swollen-styled	tf	IAJ pr	o.f	W C G. H.	1821.
-	11971a fallax Jac.	deceptive	tf	Alpr	-	Y C G. H.	1825.
-	11975a cruentata Jac.	bloodied	tf	Alpr	sd	P C G. H.	1826.
-	tortubosa Lindl.	twisted	tL	^Jor	1 jn.au	Y Chile	1825.
-	tiriba Hil.	city	£	Al or	... my.a	Li Brazil	1828.
-	Martiana ZuccariQi.	-	-	-	-	-	D s.l Bot. reg. 1249

1299a.	OXLE^Yil Cum. OXLEYA. {John Oxley, late Surveyor.general of N.&WJ	10.1.	McIttcca. 1.
-	- xanthdryxylon Cum. yellow-wooded £ LJ tin 100.	N. Holl. 1828. C It Bot. mis. 54

1193.	OXYCOTCUS 10090 macrocarpus						
2 variegatus Dens.	variegated/w/ tt fc.fr f my.jn Pk gardens	-	L bog	-	-	-	-

H7s-	»537a. PACHYPONDIUM Lindl. (Pachys, thick, podion, peduncle: thick footstalks.) & 1-4 w f t t f e £ tuberbum Lindl. tuberous-rarfd J> LJ or 1 su W.R C. G.H. 1813. C s.p Bot cab. 1676 Whites tuberbsa Haw. Hort brit No. 4392.						
-	«culentum Li/17. succulent j*i Jor 1 apjn W.R C. G. H. 1820. C s.p Bot.reg.151S	-	-	-	-	-	-
-	•Echites succulent Thun. Hort. brit No. 4391.	-	-	-	-	-	-

2 fl&repisno Goldie	dWir-flowered X A or	I my.in R	RussU	1831.	D co	-	-
o albibiflra Pall.	10 Pdtstf Sab.	Potts's	A A V ¹ 3 J»	c	china	1822>	D Bl Bot reA 143b
PALAVV/^.	rhombifolia Grah.	rhomb-leaved	-.Q]or	1 jn.s	Ro	Lima	1829 S co Bot mag. 3100
Safton-fW	i A or	liny.	Saf	Altai	1829,	S s.l	tot. mag. 3035

728. PARONY'CHIA.

29091	- canarie ⁿ sia J.	Canaries	■ tAI cu 1 au.o W	Teneriffe 1829. C co
	1923. PASSIFLO ^V RA.			
29092	16868a Cavanillfcstf Dec.	Cavanilles's	■ ■ ... jl.o	Cop Caribees 1822. C l.p Cav.dis.273
£9093	16874a AndersohnY Dec.	Anderson's	■ ■ ... jLo	Str St Lucia 1823. C Lp
29094	- cephaleima Bory	head-wreath	■ ■ ...	Brazil 1826. C Lp Boryann.22.*
29095	- albicans ft C.	whitish	2. CD or 20 jn.o W	Mexico 182a C r.m

1929. PELARGONIUM.

	staphisagroides Swt.	Stavesacre-lktt. LJ or	ljmy.o P	C. 6. H. 1827. C r.ra Sw. ger. 2. s. ■
Garden Varieties.				
	la Alexandertdnum Penny	122a Hislopifnum Penny		206a Richmondfaum Penny
	42a Camerbnianum Penny	154a melanichlcura Penny		260a Webbsantroi Penny
	46a Carelcssidnum Penny	183a Pennyiinum Young MSS.		

1476. PERE'SK/i*.

29097	12631a BlloH.8cK.	Bleo	* 13 fir 5 o.ja	Ro Mexico 1827. C p.l Hot. reg. 1473
-------	-------------------	------	-----------------	--------------------------------------

3477. *1670a. PERILO^NMIA H. \$ K. (*Peri*, around, *loma*, margin; fruits with a membranous border.) ■■■■■
- - ocmiflides Kth. Basil-like 41—or 3 aus.s P Peru 1829. C s.l Bot reg. 1594

3478. *2530a. PERISTETRIA Hook. DOVE FLOWER. (*Peristera*, a dove; its column resembles.) 20.1. *Orc/iide<e.J-*
29099 - - d'a taffoo*. lofty jgEJ or 4 su Ysh Panama 1826. Dp.r.w Bot. mag. S1H|

302. PETROTHILA.

29100	2438a came'scens Cun	canescent	Si i_J or 5	— N. Holl 1830. S &p
-------	----------------------	-----------	-------------	----------------------

126. PEYROU'S/il: substitute this word for Lapeyrou's/7/.

2100. PHA^CA.

29101	18844a lutea Led.	yellow	^ A pr 1 JLau	Y SiberIA 1827. R co
29102	188446 densiflilia Sm.	crowded-leaved	£ A Pr 1 JLau	R California 1822. R co
	18846a astragalina Dec.	Astragalus-like	& pr 1 jn.jl	W.B Scott. Clova mo. S 81 Bot cab. 489 ^M
29103	18848a trianguläriss Fis.	triangular	^t A Pr 1 jl>au	B Siberia 1824. R co
29104	188486 exaltata Led.	tall	^ A pr 1 Altai 1828. R co

1063. PHALA'NGIUM.

29105	8882a longifblium Grah.	long-leaved	■■■■■	Lima 1829. D p.l
1674. PHLOWIIS.				

29106	15073a agraria Led.	field	■■■■■	Siberia 1827. D co
-------	---------------------	-------	-------	--------------------

472. PHLCX.

29107	3906a acutifolia Swt.	acute-leaved	■■■■■	N. Amer. 1825. D Lp Swtfl.garic ¹¹¹
29108	3907c excelsa Penny	tall	■■■■■	N. Amer. 1824. D Lp
29109	3907c Wheelerftna Swt	Wheeler's	■■■■■	hybrid. 1824. D Lp
29110	3907a* Shepherdrt Penny	Shepherd's	■■■■■	hybrid. 1824. D l.p
29111	3918a crassifolia B. C.	thick-leaved	■■■■■	N. Amer. 1825. C co
29112	3920a speciosa Don.	showy	■■■■■	Columbia 1826. C 61 Bot. reg. 1135
29113	3922a aristata B. C.	awned	■■■■■	Carolina 1828. C p.l Bot. cab. 157
29114	- - penduliflora Swt.	pendulous.flwd	■■■■■	N. Amer. 1824. D l.fl
29115	- - disticha Sab.	two-rowed	■■■■■	N. Amer. 1826. D co
29116	- - involucrata Nut.	involucrated	■■■■■	N. Amer. 183a D co
29117	- - &a. Penny	charming	■■■■■	N. Amer. ... D co
29118	- elata Penny	tall	■■■■■	N. Amer. 1828. D co
	S8063 tardiflbra Penny	late-flowered	■■■■■	" "
	longiflbra Swt. B.	KG. 2. 8.31.		Sw.fl.gar.2.s.3r

5. PHRY'NIUM.

29119	64a set&sum Rose.	bristly	■ E3 2 fmr	. Brazil 182J. b r.ni Rosc.scit4l
Maranta secunda Grah.				

29120	. . coloratum Hook.	coloured	■ El or 2 ap.my O	Brazil 1828. D 81 Bot. mag. 3010
-------	---------------------	----------	-------------------	----------------------------------

970. PHYCE'LLA.

29121	8006fgnea			
	2 pulchra D. Don	pretty	tf iAI or 11 o	R Valparai. ... O r.m Sw.fl.gar.2.8- ¹
	glauca B. C.	glaucous-vrf.	tf IAI or 1 jn	R Valparai. 1824. O l.p Bot cab. 174&

29122 - - Herbertftna Lind.J. Herbert's tf TAI or 1 my.jl R.Y Andes 1825. O r.m Bot. reg. 1341

3479. *2690a.PHYLLOCLADUS Rich. PIYLLOCLADUS. (*Phyllon*, a leaf, *klados*, a branch.) 21.10. *Conifer**^h
rhomboidalis Rich. rhomboidal f i_J or 20 ... An V.Di L. 1825. S r.m Rich. con. 3
Podocarpus Heri U asplenifolius Lab. n. h. 2. 221. No. 23646.

2202. PICRI'DIUM.

29123	- ihaft>iamP.B. Webb	Holly-leaved [^] AJor	1 jl.o	Y Teneriffb 1829. C co
-------	----------------------	--------------------------------	--------	------------------------

87. PIMELifA.

29124	7966 intermedia Lindl.	intermediate	n. i_ or 2 mr	Wsh Kg.6.'sSdl825. C sp Bot...
29125	799a diosmifolia B. C.	Diosma-leaved	n. i_ or 1 su	Ro N. Holl. 182a C sp Bot c
29126	. . sylvstris R. Br.	wood N. Holl. 1830. C Lp

679. PITTO'SPORUM.

29127	5606 undulatum	variegatumDen<	variegated-urf	Ht. ■■■■■ 5 f.jn W.v gardens ... C p1
29128	5615a /gustrifolium Cun.	Privet-leaved	* ■■■■■ 6 s	... W.co.N.H.182a C Lp
29128	5615b cornifolium Cun.	Coriaceous-leaved	■■■■■ LJ or 3 my	Br N. Zeal. 1827. C p

3480. *1708a. PLATYCARPUMKth. PLATVCARPM (Platys, broad, kurpos, seed-pod.) 14.2.2% ■■■■■

29189 - - orinocense Kth. Orinoco 2 O or 20 ... Pa.Ro Orinoco 1813. C p.l ■■■■■

3481. *2434a. PLECTOCEPHALUSD Don (Plctos, plaited, kephale, head; scales of involucre.) 19.4. ■■■■■

(americknus D. Don American O *v 5 aus.s Li Arkansa 1824. S s.l Sw.fl.g*r*—

No. 25180. of Hort brit. p. S61.

- 2539. PLEUROTHA'LLIS.**
 - - saurocéphalus AC. lizard-head-/Zuro'/f ESJcu § my G Rio Jan. 1829. D p.r.w Bot. mag. 3030
 - - Lancedna B. C. Lance's _£E)cu j au Y.u Surinam 1831. D p.r.w Bot. cab. 1767 J
- 544. VLVMIE^yRIA.**
 - - LamberttfnalindL Lambert's •• o r 8 jl.au W Mexico 1819. C r.m Bot reg. 1378
 44360 parvifoliaifaw. small-leaved *CJor ... jlau W W. Indies 1813. C r.m
- So, ^ 2690. PODOCA'RPU'S.** - - latifolius WaL broad-leaved ± t_J or 20 m Ap Pundua ^M 1828. S p.1 Wal.pLas.ra.30
- a,,, 474. POLEMO^NIUM.**
 *"S . . pulcherrimum Hook. prettiest £ A or f jlau B ^{vt} N. Amer. 1827. S co Bot. mag. 2ii?9
- 2q_{loe} 2901. POLYBO'TRYA.**
 'k_{loe} . . acuminate Lk.\$0. acuminated £ E] or 1831. D co
- T 2053. POLY'GALA.**
 - - rubella PA. reddish £ A or £ jn.jl Pa.R N.Amer. 1828. S si
- 2908. POLYPCTDIUM.**
 253580 Schukhri Raddi Schkuhr's £ i A | or 1 ... Br Brazil 1824. D p.1 Haddi in. 27. 2
 No. 25388. tetragonum Sum. belongs to Division IV.
- 670. POMADE'RRIS.**
 5514a fietulina Cun. Birch-leaved m LJ or 5 my.jl Y N. S. W. 182a C p./
 w!* 5517a andromeda^{ta} Cun. Androm.-lvd*< i*J or 5 my.jl Y N.S.W. 1824. C p.1
- On,, 1447. PORTULAXJA.**
 ***! . . Gilliesn Hook. Gillies's a. | AJ or } jn.jl R.p Mendoza 1827. S si Bot. mag SQtii
- W 1528. POTENTILLA.**
 13578a dahurica Nest. Dahurian or 2 jl.8 Y Dauria 1824. L co
 13637a Hopwood MNaSwt Hopwood's or 1 jn.jl B.Ro Eng. hyb. 1829. D co Sw. fl. gar. 2.8.61
 13646a missQrica Horn. Missouri cu 1 jn.au Y N. Amer. 1827. D co Bot reg. 1412
 13646g argute Ph. sharp-toothed 2 jl Ysh.W N. Amer. 1818 D co Bot zeg. 1379
 13579 nivea JL cu f jn.au Y N. Amer. 1827. S co Bot mag. 2982
 2g_{u..} 2 macrop'hlla Ser. long-leaved cu 1 jn.au Y N. Amer. 1827. S co Bot mag. 2984
 13637a MackayAna Swt Mackay cu 1* su.aut Y.Pk Eng.hyb.fotm.sp D 1 Sw. fl. gar. 2.8.43
- 331. POTTHOS.**
 - - refl£xa/J4 f reflexed £ (23 cu - s.0 Br Brazil 1828. O p.1
 - - reptansI-*O. creeping LCDcu - 8.0 Br Brazil 182& C p.1
 - - rubaceans/JK. *O. reddish jg O c u - s.0 Br Brazil 1828. D p.1
- 955. POURRETW.**
 p\Q2 . - cerClea Miers blue ig O or ap.au B Chile 1827. Skp.l
 ^153 . . rubricates Miers red^temmed £ (23 or ap.au B.R Chile 1827. Skp.l
- 1737. PRE/MNA.**
 - - escuUnte Rox. ? esculent - or 25 - W E. indies 1824 C s 1
- 2574. PRESCOTTJA**
 3)155 - - lancesfblia Lk.SfO. spear-leaved ■ ES or — 1331. D p 1
- 451. PRI'MULA.**
 3785 farinosa L. mealy-Uaved 1 small red-flowered A or 5 i. 5^{nr*} Br bog. D p.1 Sw.fl.gar.2.s.65a
 2 large pale-flowered A or Pa.R England bog. D p.1 Sw.fl.gar.2.s.65b
 3 white-flowered A or W England bog. D p.1 Sw.fl.gar.2.s.65c
- 1694. PXUNE'LLA.**
 15306a Brownianna Penny Brown's aust&lis Swt. hort. brit. ed. 2. p. 410. O or f Jlo B N.S.W. 1826. S co
- 2J157 ■ 1. PSORA^xLEA.**
 I86?8a brachi'te Dou. brachiate A A esc 1| jn.jl W.B Pl.RkyM.1828. D p.1 Hook. am. 53
 " Allied to P. esculenta, and like that species used as food by the natives of the Plains of the Rocky Mountains."—Drum mond.
- 631. PSYCHOTRIA.**
 5263a rfaphnoldes Cun. Daphne-like i j o r 3 jn W MoretonB.1830. C l.p
- MiSa otlz^z -PTETRIS.**
 ^155 f 125 a calomelanos Sws. neat-dark if Elcu f aut Br C. G. H. 1830. D p.1 Bot cab. 1689
 291RI f 125a peruviana Lk. % o Peruvian £ CSor 1 o Br Peru 1830. D sp Bot cab. 1665
 Sq g 25500a latiflina Cun. "broad-zoned" £ LAI or 1 j jn Br MoretonB.1831. D p.1
 ^125 flexubsa Lk. Sc 0. flexuose (SI or . . 1831. D 60
 291(2 " - intramarginalis MM. intra-margin. J^mJ or 1 jl.d Br Mexico 1828. *D co
 * - 8subverticillata Gau. n. n. n. n. n. n. Al or 1 . . Mexico 1831. D co
- 2916S 1262. KJLTENJTil**
 - - mucronate B. C. mucronate Ivd tL i_J or 2 ap Y N.Holl. 1826. S p.1 Bot. cab. 1711
 Qq 3

LOUDON'S HO'RTUS BIUTA'NNICUS.

- | | | | | | | | | | | |
|----------------|--|--|---|-----------|---------------|-------------|---------------|-----------|--------------|------------------|
| 2C1G | • - ovatuin Sett. | ovate | A | cu | 3 | Jl.o | W | N. Amer. | 1829. | D co |
| | 1507. PVRUS. | | | | | | | | | |
| 29167 | 18953*; Pdtka Ham. | V&tXn^rough^fruitni f | cu | W | my.jn | W | Nepal | | | |
| S01C8 | 28355a pneix)* Stet | early, Faradite J | ft | 10 ap | Bli | Ruwia | | | | |
| 3*82. | *21a. QUA^Lfisi Aub. | QtULR*. (Name given to the tree by the Indians of Guiana.) | 1. | L V | | | | | | |
| 291(K) | - - utol&cca ,V/or. | VioleUcolourcdf • Ira V | | | | | ISrazi] | ISSfcS | | |
| | 66G. QUE'RCUS. | | | | | | | | | |
| SU7CI | >>UMJ dealbuta Li->d7. | whitened | f | i_J or | 20 | | | Nepal | 1838. | |
| | 1687. RASWaciJI.VS | lt< im'tinis L. | | | | | | | | |
| | S macrophylliular/ | I iirgt--leaved | * | A | i | my | Y | Teneriffe | 1658. | |
| 29171 | 1+5H7a P&Llidufc £>cc. | ixdv.flowered | i | A | ji | Pa.Y | | | | |
| 2917a | MRIHa iu^jeurfilde* Brat | Hare's ear-Jri*. i | 1 | ati.jn | Y | Portugal | 1896. | | | |
| 2917.1 | 1+5H9N l i ' JF | manh | **8 | i jn.it | Y | NViH-rifle | 1826. | | | |
| 89174 | 14f99f polypMIUu W. | minny-leaved | | | liiy.jii | Hungary | IS!!. | | | |
| 29175 | l+fitte Unptekim *. | tort | t, A | | jiP LIU | Iipland | 1857. | | | |
| 29176 | 146080 SiliJti It. Dr. | pedatitid | 3tA | | 1 my.jti | K. Amer. | 187? | | | |
| 29177 | 146086 p<intiHiun 8m. | northern | A A | | 5tA ... jn.au | Siberia | 1857. | | | |
| S9J78 | t-tjOKc ucUcui llktmrds | | ^ A | | S jn.au | K. Amer. | 1827. | | | |
| 7 | | | | | V | N. Araer. | 1827. | | | |
| S9JS0 | IKIIsa n'lfulus | | | | | | | | | |
| 29181 | l-tolita IS'tyti.j->f. Cr. | reddish-Anim/ | 4 | A | ... my jl | Switseirla. | 161& | | | |
| | afireus .VW. villosus Anton's | Brynniu's H:7 | ^ A | | y | | | | | |
| S9Sttl | HGSHa reeurnAtii Pair. | rw.utxetLsreded i | | | | N. Amer. | 1827. | | | |
| 29183 | - rhomboiit'iis/ZiW. | rhomboioid | | | | N. Amer. | 1835. | | | EA jour.d I |
| £9184 | - giaberrimuH /uoA- | smoothest | | | | N. Amer. | 1827. | | | .im. 5. |
| WI85 | - brevica'iilu 7/twi. | short-stalked | | | | N. Atner | | | | IliKik.:LIU. 7. |
| 99188 | IKSSa cantopWBm Boo*. | heirLtd | | | | Canada | 1889. | | | Dot. |
| <>H7 | - EfceliictiottKit Honk. | Bccbichofts'i ^ | | | | N. Amer. | 18S7. | | | |
| 14S36 | 14S36) null;L-ILH | £ whrolei'it'iis/fijAoyi jiale yellow | j | A or | f au | Pa.V | Englaad | mcs. | O co | |
| | 1S3S. JIHE^UM. | | | | | | | | | |
| 29188 | - leucorhlzltm Fit, | white-rooted | £ | A | mv,jD | St | Siberia | 1827. | D co | |
| | 1475. KH1TKAI.IS. | | | | | | | | | |
| S9189 | 13625a Hookrtina Dec. | Hooker'i | <> cu | 1 | jlo | W | W. Indies | | | |
| | IS39. fiHOODEDENUHON. | | | | | | | | | |
| 29190 | 1101.id FarrTir Swt. | Mrs. Farrer'g | ts | or | :3 f.mr | Li | China | 1829. | M: | *r.nr.? |
| S91S1 | 1103a itsut's] 1MH/(HI Swt. | BnnWi | < | or | 4 mr | Ko.H | hybrid | IH27. | | |
| 29193 | HOSSo alt;7/onSc Lindl | Hlghden | 1i | <pl | 15 ap.my | V, | Kng. hyb. | IH27. | p.] | Ritr. rej.; HJ-i |
| S91S8 | III*lic (GmteO hybrid) | Carton's Lindl. | 1t | or | 3 jn | LI | Y.ng. hyb. 1» | | | |
| S9194 | - in'ilchrmn Swt. | Smith's beautifiH | 1J or | 3 ap,Jl | llo | Eitg. hyb. | IH27. | l.p | | |
| | Smithri Swt. | Smith's | 4 | spl | 8 <p | C | ling. hyb | I8S6. | LP | |
| | R. Indiana j Smthtf Sweet'i hort. | brit ei 2. p. 3*i, and' Anklea Indie | | | | | | | | |
| | is from seeds or A. fodifolia Hbot | i d | | | | | | | | |
| | Indicum, No. 43+1. as Axileu Indies. | | | | | | | | | |
| | var, igattcem Suit, | fiery | <> spl | 2 | mt.my BtC | | | t: p.) | Sw.ti.gar.J | |
| | KM. ABU'S, | | | | | | | | | |
| 39195 | 7S12O paniculO ff</. catiational Book, | jranicisl sl caustic | • i | i or | 3]n | G | Kepal | 1823. | C Lp | |
| 29196 | - La&nu c <i>!T</i> ii:i Molina | | J or | 40 | ... G*h.W | Chile | 1S2H. C 1 | | | |
| | 719. nrsEs. | | | | | | | | | |
| | div:rio; jtun Hint. | str.ijt^liliK | tt | or | 7 ap | W.R | N. Airier. | 18S7. | C CO | Bat. rcjr. 13S |
| | infебriHDS Limit. | intoxicating | ft | cu | : | GshW | N. Amer. | 1827. | C co | But rug. 147 |
| | 33*. | | | | | | | | | |
| 29199 | £3a'<i tincthri: lamiUvn | MSS. dyeing Dr. Hamilton'* " Rouge Plant" | • • Ay | 4 | my | W | Caraccal | 18,m | S p.1 | |
| | | | May Ue ilUttntc from all fpcie* in Britain, but new | | | | | | | |
| | 15S& SO^SA 13437 vllK.ssa | | | | | | | | | |
| | 2 Le&lna Swt. | Lee's glo!« hip* | or | C | | | | | | Id^ w. |
| I3470P | fOardeo variety) | Clare's "wall or | 6 | all >a | Dp.C | Italy | " | C co | »* > « * { * | |
| | (.ii.r-h-11 hybrid) | Ruga Limit. | ± ... or | SO jl | Bfi | HaLhybrid | 18S7. | C co | Dot. reg. 1* | |
| MttSnuitlfiteB | py | 7 | | | | | | | | |
| | Annl<teP.B.WeM> | Arrahla's | | | | | | | | |
| | 15S3. | | | | | | | | | |
| | fltvil.1 // | | cu | | | W | | 18E7. | I. co | |
| | | | cn | 2 | my.jl | W | oar. | 1887. | I | |
| I3ILVMI | Dntkftnni afee | Nootka Sound | or | 4 | roy.o | \W | N. Amt-r. | 183a | Hit CO | |
| | | | or | 4 | ai),my | Dp,R | C'olinnl | | | |
| 29207 | E I luriibeam-lvd t> | | cu | 1U | JLau | W | hw1. | I | | |
| | 2411. RUDBE'CK/J. | | | | | | | | | |
| | numAta Fe<. | | | | | | | | | |
| | diflitM^ B. K. | | | | | | | | | |
| | Isrini&tn | | | | | | | | | |
| | i-'>linin:iri-i PA | cdttmnat | | | | | | | | |
| | | leaved | | | | | | | | |
| | | | 3 | * | Y | | | | | |

FIRST ADDITIONAL SUPPLEMENT.

SAGTNA 3451 procAmlxms
2 pluripi'Ula Pent, rony pot. or <&>ff, cu i my.* Gih.W Engtani) gard. D t.i

tenuior *Jhrer* slenderer ii tti 15 my Ap Scotland riv. ba. C m.s Eng. bot. 2650
in Wooigaritunti Bower Wonlgar's cu IS ap Ap England riv. La. C in.a Eng. bot.
femigtnca <3. Wnf. ruvit/tMtf'd cu 12 ap Ap Britain Jiw. 0 00 Eng. bot SS85
//hillyreiBfiliajitoj-wr Ph " cu 5 ap Ap Scotland high.v. C m.i Eng. bot 26dO
» 240H8a petra.-a G. Anders, rock cu 7 ap.my Ap Britain < BO

171*. SALPIGLO'KSIS.

SBSCId Bar. i Ol or 3 Jat Br.Y 3 It Sw.fl.g4r.ai.ns
- m l J *
28397* im v. entire-leaved BaaurHHwl 8 i- or ||| Ro Uruguay J83L 8 Bot. mag. 3113
7 liniurU Book 1 or 1|| Urugmty 1831. 8

SV I.VIA.

Hira erect-flowered i D o r 3 d Br.R Peru 1831. C It
cardinlits H. li. Sf Kth. aptt&de& rat now finrrf, k<v. vol. v. i>. fi77. tig. llili.; an aXCeUen .iml
original figure, mptXW&l 'lie lirst published in Hritain.
Grahmli Benth. Graham's 1 or 4 JLlo Mi'MCu 1899. C 61 Hot. refl. J
foliuu Benth, leafy H all tea. B Mexico J8J7. S co Hot. reg. » «)

338. SANGUISO'KBA.

S8270 praj'tox Bt-s. eaxlj-fcnvering 3 my.jl Pk Siberia n co
2fcf74 angu>ti(Tilia J7*. narrow.lt-itvtl 3 jn.au Pk Siberia iaaa 1) ru

1385. SAPONA

USiln cai&bric* Gutt. Calubrian O or i au.a * R Calabria 1830. S co Sw. (I. gar.2. »7

f 2561 SARCA'STHUI

guttatuii Lindl. spottm! 1 ap W.V.Ro Dacca 1818. D p.r.w Bot reg. 1443
Aerldes guttatum liox. MSS.

1555.

minor Mtf. sm.iltiT i ror.my P.O Carolina 1829. D bag Sw.fl.gar.2.i.13

1650. S A Tl¹ It I. J.A.

mutlii P. B. h'ebb tort 1 | jlo W IVifriUt 18J9. C co

138a AXI-FIIAGA.

spiked i A i mv.in Spot. S. Amer. 1R27. • D pl
U253rt argfta /J. fijw ilia., * A i my.jn Spot N. Amer. 1BS7. D pl
lia>anuilica(klie),J)» nal * A fmyjii W K. Amer. 1887. D pl
112586 pyrotefblia D. Don PyroLa.leaved £ A | my.jn W N. Amer. 1827. D pl
titlil. Hun: A run- mblinf, in habit. but fin* rwts mil graiulnti!
A*w)iire(Hu not ft coonlltaif! of S. otibMea flm., an wa doubtingly suppowd by Mr. Honorth
when be tVumed the genus Leptiucn irmn them.

S5 SCHIZ.VNTHIUS.

iT7a HoAkeri 0XI. Bfookefl O or 2 te.O Ro.Li Chile 18S8. S LI Bod (Mf, 8070
mi Hnfc. Orraham'n O or if jti.o Va ChUt ... S It BOL mag. 30*4
47HA ic-UlMii Hixik. leXu&v-petuttd Q or 2 jn.o Va Chile ... S It Mut, nug 8M{

, •3350a. SCHIZOME^flia D.Don. SCMI/OVRRIA. (frA^.tocut, mtrft, port: cut petah.1 in 9,
ovataD. Don oval-fcoerf it t_J or 8 ... W N. Holl. wK C tl " i

SCI'LLA.

• ffwef-centc<1 U A or ... uiy.in B Portuital 1818. O co
prntf*E*i) W, *K. meniow my.jn B -7. O co u K. 18)
plfiraboa >n<tf. Iend-co/oumf 9 tAJ gr Lc
uiimil.1 /Juf. dwarf S A Pfl 1819. O t.l

rose-coloured ff A or i ap.my Ro Kumillin ... O co Brot uh 4A
. mag. 17*6. left hand figure. Caruea Suit, in his Hort. brit ed. S. p. 59S, "

1790, SCHOPHULA'RIA.

»>73a laitata Fh. woolly 1 i a r S ji.rt IMi Siberia 1JK8. S co
elotigHts P. R Wthl) elongated .__! or 6 jLo INii TciierM)! 1829. S co

: LA'RIA 15390 alpina L.

2 ViirtegbtA Hart. lailajltin,IM if A or J gu P.v Switier). D <J Bot reg. 14C0

H10, SPDUM.

S72,, subclavatum Haw. lubclnvdtc £ A N. Amer. IKS. C ru
•itSI All- A W Engl, GBoue. C ru

S. tnit'riuitlitmi Han. in Phil map. folium /,m. Lippr.Ir.ived A. A P^r W Eng. Heref. C rm
lure /,.. S diiniintum Haw. 1CM * A Y hea. D ru
T cloogatum Evt*rvLPH. Evers'a t A y England Suf. D ru
Siberia I) co

liELA'GO.

GilW Hook. a. | or 1 tut Pn.no S. Africa 18Ea C t.l Hot.

¹⁴ra. sEMPERrvvuM.

JitiL-riffe C i.l

it. ^ Q

CD at 5 a So Braill 18SP, C p1 Hot mag.

1537. SIEVE'RSJM.
 29250 . . *rbseaGrah.* rosy my Rocky mo. 1827. D p.l
1388. SILETNE 11480 *maritima*
 2 fibre pléno *Penny* double-flwd my England sea sh. S ru
 29251 11620a *laciniata* *Cav.* cut-petaled Aj or I} 4Jl.o W Mexico 1823. S s.l Bot. reg. li
1884. SINA'PIS.
 29252 16619a *amplexicaulis*; Dcc. stem-clasped O cu 2 jns Y Algiers 1829. S co Desf. at If
145. SISYRI'NCHIUM.
 29253 1344a *glaucophyllum* *Penny* glaucous-lvd A or I} jn.o W N. Amer. 1830. D co
 29254 - - *grandiflorum* *Dou.* large-flowered A or I} ray.jn D.P N. Amer. 1826. D p Bot. reg. 13

2783. SMPLAX.
 29255 § i. ^ ii. *sagittifolia* *B.C.* arrow-leaved A |_J or If aut ^,W China 1820? D Lp Bot. cab. ll

453. SOLDANE'LLA.
 29256 3812a *creniata*. *Penny* crenated f A or } ap - - D p.l

3484. *671a. SCKLLYJ Lindl. SOLLYA. (*Richard Horsman Solly*, versed in veg. phys. and anat.) 5.1. *Pittoi*
 29257 5532a *heterophylla* *Lwf.* various-leaved \$, L.J or 5 jl B N.Holl. 183a C p.l Bot. reg. If
 55326 *angustifolia* *Lindl.* narrow-leaved J, i.J or 8 jn.au B V. Di. L. 1823. C p.l
 Billardiera Sin. *fusiformis* *Lab.* Hort brit No. 5530. p. 84.

1237. SOPHCTRA.
 29258 10451a *macrocarpa* *Sm.* large-fruited * - | o r 6 ap.my Y Chile 1822. L 1 Bot. cab. lj

122. SPARA'XIS.
 29259 . . *pendula* *Ker* pendulous tf AJ or 1 jn D.P. C.6.H. 1825. O si Bot. reg. IJ
 29260 . . *lineata* *Swt.* red-lined if iA) or j sp W.PkC. G. H. ... O sp.1 Sw.fl.gar.8J

3485. *1761a. SPARTOTHA'MNUS *Cun.* (*Sparfos*, broom, *thamnos*, a shrub; habit.) Myoporineee.
 29261 . . *jfinceus* *Cun.* rushy iTi_jor 3 au.e W N.Holl. 1819. C From the interior of the colony.

3375. SPHA'CELE.
 29262 - - *campanulas* *Benth.* bell-flowered tt. i_J pr 2 jLau Pa.B Chile 1800. C s.l Bot. reg. li

3486. *554a. SPHENODE'SME Jack. (*Sphenoo*, to wedge up, *desme*, fascicle; flowers by bracteae.) 5. 1. *Verl*
 29263 - - *pentandra* *Jack.* five-stamened A_CZ or E. Indies 1823. C s.l

1515.
 29264 13036a *arifolia* *Sm.* Whitebeam-lvdft or 8 jnjl Ysh.W N. Amer. 1827. C co Eotreg.U
 29265 13009a *capitata* *Ph.* headed A or 3 jnjl W Columbia 1827. Sk co
 29266 13027a *cratygifolia* *Lk.* Hawthorn-lvd A or I} W 1824. L co
 29267 13037a *grandiflora* *Swt.* large-flowered A or 4 jLau W Kamtsch. 1817. Skco Pal. ros. 1.
 29268 13031a *sororia* *Penny* sister A or 2 JLlo W N. Amer. 1829. L co

1669. STA^CHYS.
 29269 - - *sericea* *Swt.* silky ^ A cu 1 jls Li Nepal 183a D co

2572. STE^L1S.
 29270 - - *tubata**. C tube-flowered /f El cu | mr.my O N.Spain 1828. Op.rwBot.cab. 1J

1795. STEMO^DIA.
 29271 - - *chilensis* *Benth.* Chilean ^ J5J or 1| au.o B Chile 1829. D l.p Bot. reg. 1

2724. STERCITLIA.
 23869 TYagacantha *Lindl.* Tragacanth J Q m 40 my RBr S. Leone 1793. C l.p Bot. reg. i:
 pub^scens *G. Don* in Syst bot and gard. i. 515.

2973. STETVW.
 29272 205R9a *adenophora* *Lag.* gland-bearing £ AJ or 2 aus. W Chile 1822. D co
 29273 20695a *micropitilla* *Kth.* email-leaved ^ iAl or 2 au.o Bh Mexico 1828. D p.l

3281. STREPTANTHFRA
 29274 28007 *cyprea* *Swt.* copper-coloured tf IAJor f jnjl Cop C. G. H. 1825. O p.l Sw.fl.gar.i

54ii. STROPH'A'NTHUS.
 4438 *dichotomus* *Dec.* forked A • or 3 fd Ro E. Indies 1816. C r.ra Bur. in. (i)
 89275 4438 *adivergens* *Grah.* spreading i D o r 3 fd Y China 1816. C r.m Bot. reg.
dichotomus *Bot. reg.* not *Dec.*

2581. STYLI'DIUM.
 29276 X 22328a *fasciculatum* *R.Br.* bundled tt i_J pr i au Pk N.Holl. 1830. S s.p Bot. reg.J
 29277 - - *tjuncum* *R. Br.* rushy £ LJ or \ N. Holl. 1830. O l.p

3487. *607<. SYMPHI'A'NDRA Alph. Dec. (*Sympyo*, to grow together, *aner*, anther; anth. connate.) 5. l.Ca*
 pendula Alph. Dec. Sw. fl. g. 2. s. 66.; the *Campanula pendula* *Hort. brit.* 5505. p. 4.

623. SYMPHO'RIA.
 29278 * . *montana* *Kth.* mountain ■ ■ 4 au.o Pk Mexico 1829. C co U.&B.9

5483. *1618a. TALAVMA3. TALAUMA. (Native name of a kindred plant)"13. 6. A 1.
 29279 - - *Plumieri* Dec. Plumier's A Q f r a 60 .. W Antilles 1822. L p.
 - - *Candollii* Blume Decandolle's A O fra ... ap.jn St Java 1827. L p.
Magnolia *odoratissima* *Hort.*

929. STA'TICE.
 29281 7480a *binervosa* G.E.Sm. 2-nerved j£ A or 1 jns B England ch.cl. S Lp Eng. bot.
 898Q2 7506a *publula* *Webb* Aowny-leaved £ LJ CU { m y V.w I.Graciosal 830. S co |

" Apparently near *S. furfuracea* of *La Casca*."