

Indian Botanic Garden Library
BOTANICAL SURVEY OF INDIA


CLASS NO.....

BOOK No. *STI-C*.....

ACC. NO... *B-10,127*.....

*in Histoire Naturelle
de la part*

*de H. Bouquet & N. de
Melle*



MORT.BQT.REC;
S|CAICVTr. ^{Ma} 103

STIRPES CRYPTOGRAMÆ

VOGESO-RHENANÆ

QUAS

IN RHENI SUPERIORIS INFERIORISQUE, NEC NON
VOGESORUM PRÆFECTURIS

COLLEGERUNT

J. B. MOUGEOT Bruyericnsis Med. Doct. et C. NESTLER Argentincaais
Facult. Med. Argent. Botan. Prof.



—DEUS O. M. *et* natura nihil frustra creaverit.
Postea: "tanni Int JTJt 'iitiius i'urv uttlilates ex musd»
augurortiror quot ex reliquis vegetabilius.—

Linn. *Cui bono Amæn. Acad.* torn. III, p. 241.

FASCICULUS IX.



BRUYERII Vogesorum, 1826.



[Faint, illegible text, likely bleed-through from the reverse side of the page]

INDEX SISTERS SPECIERUM NOMINA FASCICULI LX.

Artmoneta Ci-aarpi Pers.		OrihoOiricum rivulare Turn.	824.
Anahama nil inLrauna Jlorv;	806.	OrliioiLricuni lupincola Funck.	825.
vvkuU pi aril is Scfintrgr.	819.	Oscillaria urbica fiorf*	897.
Borbuh Hnriis. Imrluina Schiiltz.	820.	Oscillaria viiidis Boij.	898.
haum Prevostii Fries.	848.	Peliidca spuiia Achar.	837.
W*?noria Bnxi Fries.	882.	Pertiisaiia Icioplaca Schtvr/	847.
Ijnmii enihrorarpon Brid.	832.	Peziza a rife Pers,	888.
lijuin Ludwiggii Schkwgr,	831.	Pcziza Frangidse Pers.	887.
BB i, m rosetim Schreb.	833.	Pliacorliiti^ttICTotioidcii Pers.	885.
(-iiltims \ioljicea Jlorv.	8(,5.	Phascnm ifctkum Schreb.	802.
Calirinum 5*atfQnclliuu Achar.	85(1;	Pliascura rectum TVith,	804.
Caliciurt' ivniji;icUiim Achar.	859.	Pliasrum scrratum Schreb.	803.
Onur^hnu dSbrme MrUs	889.	Fohlia cutullata Bruch.	830.
Lenanglum nepandnm Fries.	890.	Pohlia indinata Swarti.	829.
Cenomyce canosa Achar.	8JO!	Pierogcrium gracile Swartz.	817.
Cenomyce futrata Achar.	852.	Pirroyonium repem Hedw.	816.
tenomj cc grucilis et v. In bridla Flærke.		Ptylioslomuni compacium Ilornscli*	828.
Cenomyce raeranosa ^ c W.		Jihyisina tiiucae Fries,	865.
Cornicularia ocbroleaca Achar.	853.	Riccia natans Linn.	836.
Cytispora chrysosperma Fries.	881.	Salvinin natans Haffhl.	801.
Dicranum anibiguttii Heihv.	822.	Srlerotium areolaium FrieSi	883.
Din' ami m rongtstun B'i.iii.	821.	Sclootium roseimi Kneiff'.	884.
Difrairum iian-vum JV. et Molir,	823.	Smtaria I Inii Fries.	893.
Biclyni(«!on luittitis jjornseh,	818.	SpWria ambiens Pers.	872.
Uotljidci Heder^ Fries.	861.	SphcEria AfjuHulii Fries.	879.
Dothidea Utiberiani Fricst	86a.	Sphasria Berbericls. Persi	874.
Dothidea gtdlary F«'«.	863.	S}»li,Tiifi hullata Pers.	866.
Excipula aspt-ra Fries'.	88G.	Sphseria Corjli Bdtscfu	877.
Fusidium p-LSpuni Dittm;		S_ h_ i_ elougatii Fries.	875.
Grimmia leucophara Cr«'r.	81 :t.	Sphjeria praminis Pcr\$:	876.
Grimmia tridboph^Ua Grec.	814.	Spboerifl Laburni Pers.	873.
Gymnostimnw mtermednnn JTurn.	809.	Sphaeria oedema Fries.	880.
G^ ninostomuni tonue Schrad.	810.	Spba?ria palhiii Fries.	878.
H^ p^ m salrbiosum 7/o^»/.	83.	SpiicCtia profusa Fries.	871.
Hypmim veluinum Linn.	83:5.	S}lia?ria tjnerdna Pers.	868.
i^isiobotj^s Lonirers KUIZP.	860.	SjiliBeria Sorbi Fries.	869.
Lecanora sophodes /?. piTina Achar.	839.	Spbafria syugeiicsia FHft	870.
Lecanora yaria ^cAan	89.	Spliicria ^nencsfbrniis Pers.	867.
Lecidea qiboEa c. tr;ib. et fur, Flverhe.	8-J4.	Sphngnum compactum Z). C.	805.
Lecidea	843.	Sphagnum contortum Schmltzi	807.
Lecidea pimctata Flarkt.		Sphagnum molliscum Bruch.	808.
Lecidea saDgiinaria ^;Aflr.	842.	Sphagnum suhsecundum Rees.	806.
Lecidea vemalis ^cAar.	845k	Tetraphi* Browniana Gre&&	811.
Leptomitus inyestiens Botf.	699.	Tbelatiirema chiism Scha-rer.	846.
Mycinema Pteridis jQhrdh.	900.	Trirhostomum Fimale Schwægr.	815.
Nephroma p;iriis ^ a r .	838.	Triphragmiuin Isopyri Nob.	892.
Opegrapha Uilyi-ga Jchar.	856.	TriphiHgnium Ilmaria? Link.	891.
Optraplia nodia ^c/urr,	857.	Vernicaria byssacea Achar.	854.
wthothritum Brucliii Uornsch.	826.	"Verrucalia rann&i Achar.	855.
*ibLricum crispUitun Brush,	827.	Weissia crispula Hedw.	812.





80i- *Salvinia natara* *Jfoffht. Ft. germ. 2. p. 1.*

Mursil^a nataiii *Linn.*

In aquis stagnantibus et fossis ad ripas Rheni, æstate.

802. *Phascum muticum* Schreb. — *Bryol. germ.* p. 46. tab. 5. f. 6.
In campis sylvisque ad terram nudam argillosam vel arenosam. Februario, Martio.


803, *Phascum sarattim Schrrb.* — *firrol. perm. p. 35. tah, ,\f. r.*
In pralb et campb ad terrirn SSKWUS- An turn MO ac Uyemc.

804. Phascum rectum *Kith. ex Hook. and Tayl.* *Musc. hrlf* 9. tab. 5.
In arvis ad terram lutosam raro apud *ms.*, .1 fr^quens cum W ^ j *
Starkeana circa Andegavum *u, Jc* «b anico G«pfa .ecqgW VCre.

805. Sphagnum compacting De CailL Fl.fr. n^o, f. g. — Hedw. sp. Musc. suppl. 3.
Sphagnum condensatum Schleich.
↳ Iolictis Vogciorum, ^tate.



flor. *Sphagnum subsecundum* N. ab *Esenb. ex B.* [^] germ. p. 17. tab. 3. f. 7.
In paludibus **UU&ttebfIU** Vogesorum, astate.





807. Sphagnum contortum Schults ex Bryo., germ. p. 15. tab. 2. f. 6.
I nterfacis et paludosis Vogesorum cum precedenti.
Forma major Sphagni subsecundi, coloribus foliorum variis at non specificè
distincta, ex Bruch in Litt.

80S. *Sphagnum molluscum* Thuch in Litt.

In Yagesonim torfecen, festaLe fructiforme.

ff>Jij canlium longe ovalis, cellutii msioribus, foliis perichseif niinus lalis acnloribus,
maximo subsecundis, slaini hpmiliori i Sp>i. Sphscante divert. TetUate eel.

Persoon in lilt, species a"5uo Sph. tnelb yti ae ne viz divctsa.



809. *C^nmosiomum inicniediom Ttrn. Musc. hibtrn.* p. 7. tab. 1. a-c.
In agris, Bum, hyeme.

810. Gymnostoma tejuR SchrttiL in Heihv* Sp. Hfitsc* p. 37. lab. \$J", i—].
In cailibus iiJictii Bruscar vaJJb jirftpc Mulzi(> Y*-'re ac istate.



Si i- Tetfaphis Browniana Grev. scott. crypj. Fh. tah. 10Q.
Bryum firoii nianum Dicks. Teiraphii ovat* HooV, el Toy!- Tetrodoplium Brovminpuni S^MveegTt
Ad cAma iiiictas circa Bi'poutefn observavit el benigiie coratminicavil amicijj Urucli. Vere.

8ia. Welssia crispla *Hethv. Sp. Muse. p. 68. tab. 13. f. i—6.*
Ad rtips momis Hohneck Vogearum. Æstate.



813. Grlminiji <•iirnphaca Grev, in JFermS Trans. 4. y.,8r. tab. 6,
Dicranum piliferum Scfdeick* ?•* Mmott Disp. mrth. tins esp. de mousses > It.
In camihui liliccis reiro Mutzrg *1 ptojic Custl ad rupts LaJaUcos. Vere.

#

••4- Grimmi, tricaophylloides Gw. Fl. edin. p. 35
C«»pilpa. affinis Hornst. DHj>iodor» Schului Brut T" crypt. Fl. 100,
In camibui jJiceis* moutii Scm«l»e7 » • "«5ltmum decipiens Schultz.
Bipontem. Vere.



815. *Trichostomum Funale* Schwægr. in *Hedw. Sp. Musc. Suppl.* 1. p. 150. tab. 37.
Dryptodon Funalis Brid. *Bryol. univ.* *Campylopus funalis* Brid. *Meth.*
Ad Vogesorum cautes siliccas granitosasque. Maio et Junio.

816. *Pterogonium reptans* Hedw. *Sp. Mitsc. Suppl. i. p. 100. tab. 27.*

Pterogonium Britl.

Ad truncos Pini a Jaleslrii tica Argonitorauini. Hyemc ac pdniD vcre.





817. *Pterogonium gracile* Swartz *Musc. Succ.* p. 26.
Hypnum gracile Linn. *Pterigynandrum gracile* Hedw. *Stir. crypt.* 4. tab. 6.
Ad rupes Vogesorum raro fructiferum et in Gallia occidentali ubi sat frequens thecas profert. Æstate.



818. *Didymodon luridus* Hornsch. in Litt.
Ad moenia Argentorati, et in caulis siliceis ad ripam dextram Mosellae infra Spinalium. Vere.

Didymodon tritricum Hook.



819. *Barbula grab* Schwægr. in *Hedw. Sp. Musc. Suppl.* t. p. 1a J. (34.
Tortula gracilis Hook. et Grev.
Ad saxa « ad terram orenosam. Vere.

Barbula gracilis Hook. et Grev.

1. M. Barbula aomadfticbs.TM Scullz &* > dt Barb, in Jet. n. cur. u. p. 217. f. 25.
hl «••«• id tciram tigiilonm et «hulwi,rii. Tew.
Hacc ut et Barbula obtusifolia Schwagr. ad Tort. revolutam Hook. retulit cel. Arnott.

821. Dicnimmi CODgestum Brid.—Sclav, in Hedw. Sp. M. S. r. p* \GP>. (, j.
Dicraill lu&ccsciiiii Turit varielas usqicda in BiLautiia jir,rfrv'tsa ex OitviNe JQ Lin.
At! Ifna putiida in monU Tluhticck torfjeis. jEslalc.



nr 821. Forstan mera varietas
Dicranum longicaule.



Saa. DicramuTi amlagum Hedw. Stirp. Crypt. 3. tab. 36.
 Trematodon ~~iguum~~ ^{iguum} Brid. Meth. Musc. p. 52.
 Ad terram madidam circa Homburg in Vogesis inferioribus legit amicus Bruch. Æstate.

823. *Dicranum incurvum* *Web. et Mohr Crypt. Germ. p. 162.*
Fissidens incurvus *Schwægr. in Hedw. Sp. Musc. Suppl. 1. tab. 49.*
Varietas *Dicrani viriduli Swartz.* In argillosis vere.

#24. Ojiluiik-btini rivuiare *Turn. Muse hiberrt.* />. QG. tab. A.
In unit *d ripas rivuloium ia Vogett's infetioribas tt in (iilKa occitlenl»li, XkUi«.





825. *Orthotrichum* ^{n1J)(J1co)a} *Flmck* in *Grev. scott. crypt. Fl. tab. 105. ex Grev. in Litt.*
Orthotrichum Schlmeyeri Hornsch. in Litt.
Ad rupes circa Bruyeriam et in Vogesis inferioribus unde misit amicus Bruch. Vere.



826. *Orthotrichum Bruchii* Hornsch. in Litt.
Orthotrichum coarctatum Pal. Beauv. — Schwagr. in Hedw. sp. musc. ex Bruch in Litt.
Habitat promiscue cum *Orth. crispum* ad arborum truncos. A vere ad autumnum.





837. *Orthotrichum* *crispulum* *Bruch in Litt.*

1 arborum truncos cum *Orthotr. crispum* et *Bruchii*. M. » junior.

Speciem hanc *Bruchianam* tanquam *Orth. crispum* varietatem estimavit cel. Greville, sed ut specificè distinctum *Orthotrichum Bruchii Hornsch.* salutavit amicus noster scotinus musci, *Orth. Bruchii* nempe et *crispulum*, in aliquot exemplaribus collectionis nostræ n.º 30 cum *Orth. crispum* inseruimus.



828. *Ptychostomum compactum* Hørsch.—*Hedw. Sp. M. S. 2. p. 56. tab. 115.*
Nos in muris vetustis circa Argentoratum, amicus Bruch in collibus et locis humidis
sabulosis prope Bipontem promiscue cum Bryo coespitio, pseudotriquetro,
Pohlia inclinata, æstate fructiferum, observavimus.

829. *Poldia inclinata* Swartz in Hedw. Sp. Sfusa. StppL t. sect. a. p. *}\$, tab. 63.
In truitLs Jnitndii iaiLsqut iiaJrilis Vogcsorum. Estate.



830. *VoU* *h* *ocuf* *lata* *Btuch* *in* *Liu*,
la prafrupiU moulis UuLueci tmu *Brya* UUw^i. *Estate.*

Bryum *sp.* *gizcum* *Duby* *GA.* *gall* *2.p.* *552.*

S31. Bryum Ludwigii Schwmgr. in Hethv. S{>. muse, supph r. sect, -J, p. 95. r. 03-

Urjfuin gj-acit Schlcidt- Bryutn Beimaai PrM. in I-itt, ft OryiL n«tV, p, Ji,

Webera Ludwigii wtr. h. angoftirolla llruch in List.

- A<1 terrai in pra/mptis moijUi HOJJULCI cum lliiKliii> TOpAWL /Estate.



Bryum Ludwigii Schwmgr.





83a. Dryum erjOiroearpon *Brid, Brjol. unit', p. 6*^{J^}- (*e& %rtd. in Liti.*)
Br^um atropurpurn *ffib. At filolw.*
Jo igi* argiiJosii propc Argetjioraium et in ijlvis ciica Bipontem, Maio.

833. *Biyum posewm Schrcb. SpiciL B.*\.
Muiim serpillifnliuni g. *Linn.* Miiuni msentn *Ueifr.*
It* »jlvaruiu 100E Immiiij faro cum \heei quu> fert autumio.





8
 i- »M»"»n_iSH[L.bmsHm Hoffuu-Gvev. Tcoil, crypt. Fl tab. 184.
 Hjprtum plumosum lh-dw. St.rp- crypt. ^ 10^ 15i
 orum terramque in sylvis humidis, vere.
 Hypnum salebrosum collectionis nostræ fee. V.,r. 428 est Hypn. pseudoplumosum Brid.



835, *Hypnum velulum* Linn.
Ad icram *el* ad radices arborum in *jlv_{is}, Ve_{re}.



836.. Aiccli naUuis *Linn.*
In fa-sjs ei arjiris quictU, æstate.



837. *Peltidea spuvia* Achar. *Meih, Licit, p. a83.*

Pckigerii ^purJa *D. C. Fl. fr< ex Dttfiur in Liii.*

ID sylvis propc Argcn^{*TM¹} cum Pelt, canina a qua consumer dffert stature
liujuilioi-i, laciDiis Thalli **oauubm** pelligtrls.



838. *Nephroma pumilum* Ach. *Sjn. Zich. „ 242.*
Lichen parilis Achur.

Ad nigrum, *supra maTM, arborum corticem.
Ano luera varians *Wepbronae* resupinatae?

839. Lecanora so Rhodes v. h. p̄jriia Acltar. Syn, Lick- p. 153.
Lie he n pyrius 'Atihnr.
Ad trncos Fagi , praacrlim iu (iallia occidentali.

840. ! « « « varia Achar. Syn. Lich. p. 161.
Lichen varia D. C. Fl. fr.
, Ad tsseres ct KgHnenu Jignejl.

ftfr. Lecidea punctata *ill.*
Lecidea parajema b. punctata *Ackar. Sm.* (•) TM Lichen punctatus *Scop.*
In corticibus l. *is.*

84³« *Lecidea sangomaria* Achar. *Sjn. Lich* p. IQ.*
Lichen 5iiriguiiioL-hiS *Linn.*
Ad arboruiu trmicos ct in **rupibus.**

843. *Letidea gelatinosa* Fl&rke—*Jchar. Syn. Lich. p. 26.*
Supra tjueram *pm\>c* *Bruyerium.*



b



b

844. *Lecidea epipolia* v. *b. trabinella* Flotow. et v. *g. farinosa* Floerke.
Var. *b.* habitat ad sepimenta lignea; var *g.* *Lecidea corticola* *b.*
farinosa Achar. *Syn. Lich.* p. 32. in truncis Quercuum.



b



b

84^r. *Lecidea vema* Ks Achar. *Sin. Lkh. p. 30.*
Lichtn vema Linn.
Supra muscos q_u < in crahat' in satij d ad pedts arborum.



846. *Thelothrema clausum* Schærer *Spicil. p. 68 et Lich. exsicc. n^o. 122.*
Lichen clausus Hoffm. Lichen volvatus Vill. Thelothrema exantematicum Achar.
In rupibus calcareis Birsæ vallis Jurassæ

847. *Pertusaria* *Wbptaca* *Schaerer Spicil. p. 66 et Lich. exsicc. n^o. 119.*
Porina leioplaca *Achar.*

In conice l'yl arborum prTitrim F_{ar}F. P_{ar}t_{ar}isariae communis status abortivus
ex AT₉w Eatw_{ktl}. *Metamorph. etc. der Flechten p. 199.*

ly. An Indium. *Pertusaria*
*«w_{ar}isariae?



848. *Biatora Prevostii* Fries in Litt.
Ad rupes calcarias circa Rothomagum primus observavit amicus Le Prevost, postea
amicus Prost prope Mimatum, nos in Jurasso. Affinis *Gyalecta epulotica* Achar.

Biatora Prevostii Duby. *Gyalecta Prevostii* Fries in Litt. p. 197



*% - *Cenomyce gracilis* et var. *tybrida* *Flaerhe in Liu.*
a- *Lichen graciis Linn.* *Ceohijec femorjTin v, gracilis Ackar.*
b. *Cenom. ecmocjia var. tyjjida Ackar.*
Ad tmitu saxaquc in Jocis mouiosb et umbrosis.



850.

Cenomyce canosa Achar. *Sjn. Liclu p. 273 ex Flærk_e in Zit.*
Cettom. Vaillantii modificatio CJ; Dittbur in Liu,
Li gylvü ad terrdvi secus vias.





851. *Cenomyce racemosa* Jchar. Syn. Lirh v 5-8 ex Flerke in Litt.
Clado P. 39
ad lenam, s»« in locis sterilibus et ljlvlis.



852. *Cromyria furcata* Jchar. *Syn. Lich. p. 276.*
Lichen furcatus Huds. **U M r t a u, ^ . c i**, fruticosa var. *furcata* Schärer *Spicil. p. 41.*
in ericeti, ad terram.



853, Cerniculariit nrhroleuea *Jchar*, SVn r J, 301.
.....eucus Ehrh.
In pascuij summii cum *G'lraria islandica* et *cucullata* raro in Vogesis at
frequens In Delobiuau *uade* ab amko de Miribel accepimus.

854. *Verrucaria byssacea* Achar. *Syn. Lich.* p. 97.
▲d corticem Quercus vetustam in sylvis circa Argentoratum.



85- Vermifaria ca^{linea} Achar. Syn. Lich. p. 88.
Ad corticem Carpii.-

8*6. **Opegrapha** liifyrgh *Jchar. Sjn. Lkh. p. 7a.*
Opegrapha tmaiifis *U. C. Fl. fi: O. saicoU Ach. ex Dufow in Lkt.*
Ad mania argi.-mhiciisia **latcribni** esslructa. , copiose-

Sj;. Opegrapha nnJi:i *Jtchar*, Sin* *Lick*: />. •;<'>. ex *Fhrrkc* in *lalL*
Opegrapha lichenoidp* *Pers.* *Q'g.* *populimk* *Pert.* *J? i *htuillVun'*
Ad *LjiiicoS* *Populi* nigr.
Opegrapha uoilia collccuioiiii qclrx- *Jiwd* VI. n.* 55G *est* O. noiJu var. *sigitala* *Aclair*, I. C

858. *Calicium stigonellum* Adutr. Syti. Hcl_h p. tfc^j. p_res Ltch. cxsic.

Calicium so^sjlc Pen. in Lit I. *Calicium scutellaria* Desfour in Lit.
Ad corticem arborum parce apud ues, abundanter circa Rothomagunrubi observavit amicus Le Prevost.



Calicium tympanellum -7.W. Syn. Lich. „„>“. ex Fries Lich. exsic.
Succ. n.º 18..

Calicium leucomelas Pers. i Ja. d A char. \$%-». ! lich. p. 333.
In cortice et ligno Abioli nc» 1J- circa RcMororBr r Vegetarum.

A? hanc speciem (cuiusmc reductionem Cal. leucomelas Pers. Achar. affirmat exl. Fries l. c., alij
forsan aliter iudicent) sed si istud Cal. leucomelas Pers. spec. tympanello Achar. distinctum
esset, hic addere deberemus, errore typographico in Soc. Wett. r:L up. leucomelas denominatum
fuisse, nam scripserat ccl. Persoon Calicium leucomela ob m... theciorum albicantem.

660. *Lasiobolus* **Idnieene** Kunze—Griseb. *Fl. tab. 102.*
Xyloma Loakam Schrick. *Xyloia* Xyloia Clutii. in D. C. Ft. Jr.
In f. l. LomceTH Xyluiei, astate.





861. *Dothidea Hederae* Fries *Syst. mycol.* 2. p. 564.
Xyloma Hederae Pers. in Litt.
Ad folia viva Hederae in sylvis, vere.



862. Doihid» Robertiani Fries — CIVI? • scoll • rry)(• Fl. tah • 146.

Ad fol' *Doihid» Robertiani* in *Flora* *ver.*

8C3. Dmkklen sieliayis *Fries*'*Syst. mycal.* ♀♂ p. SGo.

Xyloina stelliit *Pers.* *Asteroma* ('lit U-UMJUI D. C
la fbtis vivii j'!)j'teuuiaruiu el drnpaniikt-uju , astate.





864. *Actinonema Cratægi* Pers. *Mycol. eur.* 1. p. 52.
Ad folia adhuc viridia Cratægi torminalis in Jurasso legit et communicavit amicus Chaillet.
Dothideæ incunabulam ex Fries in *Lit.* ad Chaillet.

865. *Rhytisma Urtice* Fries *Syst. mycol.* 2. p. 570.
Ad caules aridas Urticarum, hyeme ac vere.




865. Sphaeria bullata Pers. Icon. pict. tab. 3. f. 6.

AJ ratnos iaXlclsOr

867. *Sphaeria verruciformis* Pers. *Syn. Fung.* p. 26. *tab.* 1. *f.* 5—6.
Ad ramos Coryli, Quercus, Salicum, etc.





80S. *Sphaeria quercina* Pers. Sjn. Fimg.jZ ^ (ah. i.J. 7. b.
In ramii quercis.

869. *Sphaeria Sorbi* Fries *Syst. mycol.* 2. p. 380.
Sphaeria acutangula Pers. in *Litt.*
Ad ramos sorbi Aucupariae.

8jo. Spharria sptgenesu *Frits S/st. mycoU* a. p. '382

Spharria Frangulae Pen. in Lin.

AjJ RiBiH Hlumni PranguLr.



87)- Spbairia profiisa Fries Sfst. inrcol. p. 392.

^s ¹ ^{MI} ^{Ad} ^{J3} circumsnpta Schn. i Sou. Sj.li. torqiau Pets, tn IM. Sph. irreg'ilaris Fl. fr. ex Chaillet.
moi ttobiiiiia; PiCLiJ^cacilC , atJLC vaiiaihS quamobrtin Jiversij iaduUur nOfimLuus.

8; 2. S hpima amfeens p m % ^ Fimgt ^ /f 4.
Ad now, F, gi, Crala-gi, PrUqj sp i QWSj PjTJ ac Mali.

s-i. Sphaeria Laburni *PerSr* Syn. *Fung.* p. 50.
Ad ramos Cytisi Laburni apud nos in hortis et septis.


874. Splwria Berbprtdis Pers, Sjn. -Fung. p. 52.
Cucurp i i Berim-ilk Grev. scott. cry/.i. Fl, tab. B4.
Ad no w Berberidis vulgarb.



Fig. 5. Sli.riia elon^itg *Fries Sjst. mjcoh 2. p. fan**

Corenbiuha don^ii *Grev. scott. ajpt- H- *&. '9^**

In r>tnk RoLinux PifWincii, CdnmilJ* Emwi, Cjtisi sesiiffcJ", UJkis europai.



876. *Spharia graminis* Pers. *Syn. Fung.* p. 30.
In foliis graminum languescentibus, autumno.


877- Spheria COJJI Balsck Elench. Fung. tab. 231.

Sj)ha'ia finilriata var. A- Coryli Ptrs. Sjrn. h'tutg, p* 36.

Ad foJia adhuc viridia Cetyli, aestate.



878. *Sparia patna* Fries Syst. mycol. 2. p. 404* ex Fries in Lin.
Ad ranot Pad! emortuos , vtre.




579- SpWia Aqitfofil *Fries in Lilt,*
Ad ramos Uch A^nifulii, vere.




S60, Sji) i. n. i. l. (i-) (nla Fries in Lt'l-
In Pjiii Ulmi dL-tiJui., , Uremc ac vere.

no. 880. Varietas sphaerica
m. 2. culiformis Pers.

88i. *Cytispora chryosperma* Fries' *Sysf. mrenh* 2. p. 3 fr.
Nemaspora chryosperma Pers. *Hypoxylon cirraium* Ball.
la *Lattice irukcarum* Populi.



882. Blennosporium Buii Fries Srsi- orb. pag. p, 3GS.
Sclerotium Buii Ptris. in Lilt.
in foliis e«iccatis Buii tictetil jiucis Cur-pin.



883. *Xyloma areolatum* Fries Syst. mycol. a. p. 263.
Xyloma areolatum Iries Obs. mycol. a. p. 358. t. 5. Sclerotium pdivup /W* m Lit.
In fulij subeintruis decluisque Padi, atilumoo.



ROYAL BOTANIC
* LIBRARY *
GARDEN CALCUTTA.

884 C. ~~longissima~~ ~~maxima~~ *Walt.* --Pers. — ^{ies, in Litt.} *rr*
In culmis marcescentibus
iuenit oculuu. Kutill i autumn.



885. *Phacorrhiza sclerotioides* Pers. *Mycol. eur.* p. 193. t. 11. f. 1—2.
Ad caules exsiccatos *Sonchi alpini*, *Cacaliæ albifrontis*, etc. in *Vogesis summis*, vere ac æstate.

886. *Eicipuia aspera* Fries *Sysl. mycol.* a. p- '597.

Peiiza asfu-ra &t&*

AJ Ha, in tinliuni iiridoium O^muudx r^galis , veic.

SS;. PMBM Frangute *Pers. Jffycol. eur. p. 3a4.*
Tynpanis Fiaiigu!* *hriei Sysi- mj-cot. a. p. 174.*
Ad mmos dXiccatos Rlurani l'Vangut* , *vere.*

88!). PwJza Aria; *Pers. Mjcol cur. p. 325.*
Tympanis *Aria Fries* *st. mycol. 2. p. 175.*
In rarai, Cr*t*gi Arw et Sorbi Aucuparue, vert.

889. *Cenangium difforme* Fries in Litt.
Triblidium Jil'orne P*rs. in Litt,
Ad ramos Salk's CJprt*, verc ac astate.

cena 5) fuliginosa von Fries Abhandl. 2. p. 23.

Faint, illegible handwriting in red ink, possibly a library or archival stamp.

'8f)0. Cen.'tn^iLim rrjuinlmn P'rii'S in lAtt,
Ad ttiatw RiLis pcirai in pi-a;[iipii& m<vatU Hotmeck, v«re.



891- Tripiiragmiim Minan'FB Lnk in Linn. Sp. pi. §. sect. i. p. 8f.

Pocčhiä Ullian' D. CC. FF. fr. 6. n. 56.

In foiti Uimarm, autumnno.

Sr>3. -mphragmium Isopyri Nob.—Fries &st. orb. veg. p. ?JU.

• Puccinia Isopyri Gaertn in Litt.

In fullis Isopyri tfulikt. primus obittvavU area Andegavum am. Gaepin. Vere.





m



Sc>3. *Sentaria Ulmi* Fries—Crtv. Scott, cnpt. Fh tab. n5.

Fusidium ~~...~~

In illi> UJmi., auumno.



&>f. Fuskliim grwii.n i Dittm. — Gnx: scatt. crypt. Fl tab. 102.
lo fblis Quercus aridis, aulumuo.

895. Cadmus vio
Confer C, 3.
Vefe ad rolas tataracla-iqud muktKlima*
pi freqwns iu GaiUa mi ioua i uulle au

Specim. missis.
Agardh Syst. Alg. p. 76.
ad nos



8n& Anabaina membranina Bory (æ .V> l"mcattj)iet. clqss. / 7-
Kostoc Ftos atjtit Lyi>^>. Hyilrcph. ilm. tab. 68.
In aijnii sug¹¹ntibus, Maio.

i

8



897. Os(ill;iii;i whiai fiotr (I? S.^L ftncerit in Lift.
Ad terrain, pavjjuctiia, ijism murprum, toto armo lemjesuce huinitia.



898. *Oscillatoria vridis* Eory de S.¹ Vincent in Litt.
Oscillatoria vridis Vauch. *Oscillatoria leimii* Agardh.
Ad terram fitosam et in aquis quietis »c Ilueitti'Lus, verc.



890. *Leptomitus* *iiMt-5tiens* *Bory de S.¹* *t^ulucent in Lilt.*
Oscillatoru iavtaUma jtgardh in Liu. Calothrti tincioria A%arJh Srst. Mg. p. 72.
Θafcrnt HlptfTik.ffa'W'''. ABIDO Lejit. Liherii Agtirilk.*
In rivulis circa Brujerium lierjjs inveslicns, var A oJ fulia granium t
nar b in loljii RuuLiijc<jjuu'titJi, eit. hjeme ac vere-

f)oo. Myrinema Pccridis Jgardh fyst. Jfg. p. 33.

Conferva Pteridk Agard—Lyngh, ifyHrophjri. dan. tab. 48.

Ad rhizoinam Ptfriidii aquilinar , tt>lo auno.

Pars orgmiica filicis test* Bory de S.' ViñcLj-jt in Lftt. ei a Ctl. Agardh e(Lyngbye
J- c. ad nlas idatuni : adiiioe sub iudice lis eit.

INDIAN BOTANIC GARDEN.
LIBRARY.

PROCESSED

✓

ACCESSION

NUMERATION

CLASSIFICATION

DATE

BOOK CARD
Indian Botanic Garden Library
BOTANICAL SURVEY OF INDIA

Call No. gJ.f-g. _____ Ace. NJ&rLP.t&l.....

Author..... *tim\$dt>falLk&tfi'*

Vogelso-144-

Name of tiottaiPeWCade So,	• Date of Issue	Date of Return

A

#

