

Indian Botanic Garden Library
BOTANICAL SURVEY OF INDIA

CLASS NO..... 581.9(595)

BOOK NO....RHE-nV.7.....
ACC.NO..... B2401



Geburts

HORTI MALABARICI PARS SEPTIMA, DE VARII GENERIS FRUTICIBUS SGANDENTIBUS;

*Latinis, Malabaricis, Arabic is, Belgicis, Eramannum charafferibus
nominibusqne exj>rejjis,*

*Adje&a Florum, Fru&uum, Seminumque native magiikudinis vera
delineatione; colorum viriumcjuue accuraca defcripcionc,*

A T > O R N A T A

Per Nobilissimum ac GeneroJiJJimum C. D. 2).

HENRICUM VAN RHEDE TOT DRAEKESTEIN^s


Toparcham in Mydrccht, quondam Malabarici Regni Gubernatorcm, fupremi Confeffus apud
Indos Belgas Senatorem Extraordinarium, nunc vero Equeftris Ordinis nomine Ulufribus
ac Praepotentibus Provincial Ultrajeftin[^]: Proceribus adfcriptum,

Not is adauxiti & Comment ariis illujlav it

J O A N N E S C O M M E L I N O S

In ordinem redigit, & latinitate donavit

A B R A H A M U S à P O O f M. D.



A H S T E L A E D A M L

Sumptib^u* Vidua JOANNIS Van SOMEIHEN, Harcdum JOANNIS van
DYCK, HENRICI Sc Vidux THEOD ORI BOOM,
Anno clo Ioc LXXXVIII.

G P A R S S E P T I M A .

H O R T I
f M A L A B A R I G I ,

D E

V A R I I G E N E R I S


F R U T I C I B U S S C A N D E N T I B U S .

N A t S J A T A M ;

X A T S f A CM-C % V 4;

A U T

5 A T T A * V A L L lf * 8240 /



T A B U L A I

NauJAtam; apud Bramannes ^arundo - *Pala*; Lufitanos, *Fruit A ^Matta Teire*; Belgis, IVaterquaad'i frutexeft* cuius *folia*, in brevibus ac viridibus furculis₅ plura proveniunt, formâ cordis, anterius cupidata₃ veletiairi obtufiora, superficie plana₃ lcnia» texturâ duriori, in parte anteriori ex fulcô viridia, adversâ, diluta & furda> faporis subamari. *Floret* in racemis oblongis viridiорibus hinc inde ex craffioribus erumpunt ramis, pendunculis brevioribus albicantibus mlidentes: monopetalii, quinque partiti, candidi, exteriora verlus reflexi₅ apicibus flavis alperfi, odoris fambucini. (*emma* rotundas, virides. *FruBus* conferti, petiolis craffis, ex racemis oblongis cubitalibus pendent₅ ex albo virides ? uvas rotundas formam referunt; primò albi₅ dein rubri, ad maturitatem provefHj nigri; nafcitur in arenofis circa *tfarapoly*, *Ange-*

A

caimali

H O ,R T U S

caimah & aliis locis , femper virens. Integra Planta cum gingibere trita pannoque oblita, pedum clavos toUit. Radix cum bubuli fascibus & arundine Indicaconcremata, per modum fumigii, idem prافتat. Fru&us capiendis pifci- bus & amovendis pediculis inferviunt.

T A B . I.

Nobilillimus Dominus a *Rhecde* primus eft, ; bus non male convenit. Secundum notas gene* qui perfe~~tam~~ nobis hujus planta: dedit descri- jricas doftiffimi *Raji*, *Hift.* ?Unt. Tom. L lib. 1\$. ptionem & delineationem , quotquot enim antea deilladixerunt, formam ignoravere, utvi- part i. cap. n. Ipécies *Solani*; *Flores Monofctal't*, qum- dcrecftapud *Mattbiol.lib. 3. Bptji.pag.156.* kalis que parrui dut fteUati (unt, apicibus in umbiutcQ oblon- vocatur *Cuculo Indiano* & *Galuzza di Levante*; gis, &c. cum quibus figura florum juftè convenit j Pharmacopasis , *Coculus Indi. Caff. Baubhius' in Pinac** nomen dedit *CoccuUoficmanw*, *Cordus in ibis Commentariisin Diofcoridem*, *Solani furiofi Fruflus*, nccabfonum, cum boUni enigmenci- part i. cap. n. Ipécies *Solani*; *Flores Monofctal't*, qum- que parrui dut fteUati (unt, apicibus in umbiutcQ oblon- gis, &c. cum quibus figura florum juftè convenit j commodè etiam, noftro judicio, inter illas re- ferri poteft. De ufu & opinione variorum Au- thorum , vide *Job. Baubin. Hjil. Plant, lib. 3. Cap. 155.*



TSJE.

GENEROSISSIMO 6^s NOBILISSIMO

V I R O D I D E R I C O

BARONI DEBAER, ADSCRIPTO NOMINE DUCATUS
GELDRIy£ CONSESSLJI POTENTISSIMORUM ORDI-
NUM GENERALrUM FOEDERAT[^] BELGIC£, ADSES-
SORI JUDICII, CUI IN VELAVIENSI AGRO CURA
SALTUUM ET REI VENATION EST COMMISSA.



*itio Tibi Septimum Malabarici horti Tomum, quern , turn ablega;
tus caufja public* rei apud extremos hominum Indos ago, meo
juſſu Jmilerodami ediderunt, qui & priores tomos publicarunt
tBibliopoU. Nemini mortalium hoc muniu accepting quam Tibi
accidere pojfe , mihi eſt exploratſlimum. Nol>i Te euros
& loUicitudines, qu<vs maxima* fuſtines in rerum publkarum
procuration ^ detergere quaſi jucundiftimo ſpecclaculo florum G?
fruſluum, quos fert hortus iUe THUS Venuftus & per quam delicatus. Qui ergo non
poſſunt non grati ejfe tarn eleganti rerum pukhrarum ſpeſtatori, quas TeUus omnium
procreatrix exſocundoſuofinu ejfundit benigniflime, tot mir* & Europ<eis adhuc inYifm
arbores & fruBus , qui in ultimis Indorum terris proVenint, quiqefidem faciunt, Vix
aliagentem effe > cī largius exquifitiūma munera indulferit ISlatura, quam Mala-
bares. Ne non dicam quam omnis, tarn feVerioris, quam amoenioris \$> politioris do-
ſlrin<e fis cultor > & quanto cum dileſlu genus omne librorum , qui noniratis Mufis &
Gratiis nati funt 5 pares, eVofoas , tuamque copioſifūmam inftruas bibliothecam. Hortus
ijfe Malabancus non dubitat quin (§ locum fit inenturus in tarn illuftri Mufieo. !Prv?
terea neceſſitudinis & affinitatis jura > qua femper coluimus religiofe, quxque nos dudum
junxerunt arSliflime , nonpaſſa funt hoc arnitici* monument um > ad omnem poſleritatis
memoriam duraturum, Tibi diutius debere. (Poterit, credo, hoc tibi perfuadere nihil^
quamyis longi/Jime jam fim à Vobis nmotus, memoria Tui mihi carius & jucundius ejfe.
Quodut hie profiteor in oculis orbis ten'arum, fie femper , quandocunque dabitnr ft-
cultas, teſtatumfaciam , ejfe me*

Gencrofiffimo Nomini Tuo

** CALCUTTA*
addidtiflimum & obſervantiilmT&m-

HENRICUM VAN RHEDE,
TOT DRAEKESTEIN.

AD


ABRAHAMUS *k* POOT M.D.

L. B. S. R



Andem, poft alios, HORTI MALABARICI cura in me devoluta eft: fuatis verborum lenociniis animos captare, multiimjaſtare > montes polliceri, arenam dare, meum non eft ; *Septimam* tibi tradimus *Tartem*, *OBavam* daturi, modovitam protrahat Deus. Tu interim, C. L., fruere noſtro labore ac vale.





Murex - Kelingū - Līg

مُورِّكْ كَلِنْجُو لِيْج

مُورِّكْ
كَلِنْجُو

KATU-NUREN-KELENGU.

TAB. XXXIV.



Atu-Nuren-Kelengu apud Bramannes *Tiava-Karandi*; Lufitanos (*ftabofa rodonda*; Belgas *Keylen met ronde watten*; species *Kjntu*, ceu *Z(uren Kfkngu)* nafcitur in petrofis terra *Candanate & Cotate*; *%adix magna & crafla* inftar Bryonias; *Caules* spinofi. *Folia* iena? quina, quaterna veltrina in uno pctiolo. Praeter verrucas bulbofas, intus carne denfa, aquea? vifcidarepletas? qu^u caulibus circa exortum foliorum infident > nullos habet frudtus, femper novas propagines antequam terram attingant, ejicientes. Ignotius.

R

N



Kia-Pura-Kelingin Linn.
Ph. 84004 D. 9.63. M.L.

قَامُورْ بِرْجِمْ خَمْرْ وَنْجَا



T J A G E R I-N U R E N.

T A B . X X X I I I .



fagen-Nunn Malabarenles: *Carandi* Bra-
mannes; (*/raboja* Luiitani; *Key ten Be]g*«: dicunt, frutexeft, cujus *radix* ut rapha* niHifpanici; *Catties* tenues₃ rotundi, glabri, **fpinofij perinde ut 3**(uren-*letengu*. *Folia* in petiolis oblongis -, terna proveniunt? interiori parte viridi fufca&furda, aduersa^t glabra ac nintentia -> coftulis albicantibus, quae in adverfa parte eminent? obfita. *Floret* fert nullos. *FruBus* plures congregati fpectantur in petiolis oblongis ?, ad exortum foliorum erumpentes? adlongitudinem pollicis, triangulares, in vertice cufpidem ferentes 3 virides & cinerea Ianugine amidtij adpetiolumconverfi; finguli anguliduosveficulas feminalescontinents cartilaginofb fepimentofejundli, in quibus unum velduo reperiuntur lèmina oblongo rotunda₅ plana? fubrubra, lèpimento accumbentia, tenui membrana infstructa. Prater fructus, verrucas ceu iþurios habet fruþbus, oblongos, fcabros ad pedem foliorum quafi oppofitos > è quibus perinde ac è radice & femine nova planta exfurgit. Natalefolumeit *Cotate* & alialoca, iemper virens & frugens. *Folia* millcpedurti morfui me* dentur. %adix cocta pro balneo ufurpata velpotu afliimta hsemorrohides fanat.

Tyrian Pepper
Indigofera.

Indigofera.

Cane.

جافر بوزة

بوزة



W A T T O U - V A L L L

T A B . X X X I I .

N Alabarenfium *fVauou-Falli-*, a Bramannibus *Maroi*; à Lufitanis *Torque*; à Belgis *Temhout* vocatur. *Caules* rotundi iunt, virides ac glabri. *Folia* in petiolis oblongis, craflioribus, interius fulcatis proveniunt, mollia, lenia, viroris pleni > faporis subamari, odoris gravioris. *Flores* fafciali erumpunt in communi petiolo -> pentapetali, viroris ex flavo diluti > interius pun

- &
- ul

 nigris anticibus adiperfi; in meditullio globulus conlpicitur, capitulo viridi inftru⁶lus, quinque angularis, viridi vena ftriatus, quinque foliolis parvis > rigidis, viridi dilutis ftipatus, odoris subgravis; nafcitur circa urbem *Cochin*, floret tempore pluviali; femper virens. *Fru* Sur raro inveniuntur. %adix funiculofa eft longe lateque fe dit pergens, faporis amari. *Folia* cum cortice *Lanja* contrita ulcera inveterata exficcant: Cortex cum Sandalo & Muliebri lafte in formam noduli adhibitus* prseftantiflum Caufonis remedium eft.

Pithecellobium

Le

جبل طارق
المرسلي



dicisbulbus cumCurcuma tritus &in oleo fergalini coitus? catharrorum fluxum impedit; cum gingibere vel pipcre, oleo foliorum de Emferno, vel melle intindlus eundurn habet effedtum. Loco vero Salfaparillo Peruviano à Lufitanis ufurpatur.

TAB. XXXI.

Smilax hsec afpera noflri Au&oris* pro fpe-
cic Zalfaparillohabetur, acejus loco apudMa-
iabarcnes optimo cum {icelii in ufu eff. sicut
& omnia genera *Smilacis Afpert*, in illo tertio
conveniunt, Secundum relationem plerorumque
Botanicorum: hujus plantx divers'e inveniuntur
ipecies-, vide *Kechium Hift. Plant. Mexic. lib. 8. Cap.
41. ftfi infua JtijlSr'mNatuu Braf. 6^4, tap. 6\$.*

duas species enumerat fub titulo *Ivapetanga*^A quse
non male cum *Macapatli-Paralta Recpii* convenit
& huic *Carl Vilandi* quam maximè affimilatur,
exceptòquod capreolis caret fecundum notam
chara^lerifericam doftijfimi *Raiji* in *Hif. Plant.*
lib. q. p. i.cap. 5, quorfum Leftorem ivirum
removemus.



WAT-

C A R I-V I L L A N T > L

T A B . X X X I



Ari-Villandi; Bramannes *CaraP^u* dicunt; Lufitani %qizJa^cBoba\ Belgae!Po^«^; nafcitur in planis & arenofis locis inter *Calicolumgen* & *Qranganoor*, & per totam ^idabariam^ floret tempore pluviali, frugens menfc Novembri & Decembri , femper virens.

Radix craffa , lignofà , dura, hinc inde fibras tenues lignofas acrigidasfpargit, coloristerreofufci. %amitennes, nodofi, duris, acutis, viridibus, ac inferiora verfus incurvatis fpinis inftrudi, hinc inde capreolos ejiciunt. *Folia* in furculis ac pedunculis brevioribus, craffis, viridibus, interius fulcatis proveniunt , oblongo rotunda > anterius brevi cuspide, rigida, texturâ fragili, viroris in reda parte fufci & nitentis, in adverfa dilutionis, odoris nullius, faporis fubastringentis. *Cola* in adverfa parte eminent, fubtilionbus ac transverfis venuis intertext#. *Flores* hinc inde in furculis foliofis, fafciatim confpiciuntur, nodulum ceu capitulum petioli in vertice pedunculis brevioribus 8c incurvis circumfidentes , hedla petali, quorum trcs interiores valde angulti & ftridi, exteriores non item, fed inmediovenula ftriati, decuflatim interpofiti; in medio *flamiM* eminent viridia , fex albis ac inflexis apicibus inftruda j nullum odorem habent. Poft flores fequuntur *fruSus* parvi, circa capitulum petioli, fafciatim difpofiti, fubrotundi & nonnihil oblongi, vertice parum ftrido, primò virides, dein fubnigri, intra corticem unicum tantum femen continentes, faporis expers. Radix cephalica cumGingibere&Curcuma, pannooleo Sergalini tindo infperfà, per modum fufHmigii capitis dolores mitigat; cum Glyrhirrhifa, cortice Caniram & fiocco Tamarindorum , loco balnei adhibita , idem prasftat, ut&in oleo cocta & capiti inunda vel auribus immiflâ. Concretus radicis



نیپارجی
نیپارجی
نیپارجی



T I R LITT LCA A' N I.

TAB.

XXX,



Iri-Itti-Canni Malabarenfium; Bramannum *Satayo*; *Lufitanorum Ftt/a da Njüira*; Belgarum *Smiltbout*; planta eft in corticibus craffiorum & vetuftiorum arborura initium fumens? ficut *IttiCanni* & omnes {pecks tSXdaravara. %ami cortice cinereo ac craflis nodulis hinc inde extuberant₃ circa nodos fpinis geminacis? acutis & exadveriō oppofitismuniti. *Folia* inde dunculibrevioribus, craffis emergentia, oblonga funt, texturacrafliori? fpiffa ac ibiida. *Flores* in communibus petiolis, oblongis folitarii₅ bini, ternive ex nodulis ramorumfpicatim erumpunt, ieriatim difpofiti, pentape-tali? coloris aurancei. *Stamina* in illis conipicua quinque breviflma, apicibus albicantibusdotata. (femm*rotunda mndem ut flores colorem referunt. *FruBus* nulli, nee vires.

P

CARL.



لہے کریں

Triplinervia

ITTI-GANNI.

T A B . X X I X .



Rutex *Itti-Canni*; apud Bramannes *Medefanni*; Lufitanos *Fula* *5\4orifca; Belgzs *Trifyout*; ramos habet quadrangulares, geniculis hinc inde interlindtosj coloris cinerei. *Folia* ex geniculis ramorum, petiolis longis, craf- fioribus, interius planis provenientia? oblongo rotunda funt? textura craffiori coriacea₅ non-nihil rigida, fragilia₅ fra6ta gratum odorem ipirant, laporis fubamari. *Flores* ad exortum foliorum plures congregati in uno petiolo vifuntur, flavi, pentapetali? ob' longo & flavo collo, intus fubtiliflima flava lanugine; excujusorificiopluraftaminulaemicant, flava, tenuia, fine apicibus, quorum aliqua in vertice bifida. (*femma* florum oblongse, exunaparte nonnihil inflex[^]e, vertice obtufiori, coloris flavi. Foliis florum marcefcentibus ex pfa florum bafi *frullm* nafcuntur, rotundi₅ globofi, in vertice in angustum fe ftringentes > coloris ex viridi f lavi; in medio cavitas eft? nucleum rotundum, album acparvum? fporis amari in fe continens. Nafcitur fupra arbores ^eque ac *5\daravaras, continuo radices cmittens in corticibus arborum, nunquam in terra; femper virens, & floresvelfrudlushabens; natale folum circa *Locbien* 8cin multisaliiseftlocis, indenfis nemoribus <k proceris arbo- ribus. Radicis cortex dentes figit.

T A B . X X I X

Species *CaprifofHi* ycl *Tericljmemi* hie frutcx eft.

جعفری
جعفری
جعفری

نامه

V S

U N J A L A.

T A B . X X V I I I .

Alabarenfium *VnjakveWnbala*; àBraman-
Mnibus *Zendarli*; àLufitamis *FruiaTanaoi* a
 Belgis *mmKruid* vocatur; frutex eft naf-
 cens in terris *Zettniene*, *Calgolano* & alibi,
 iemper virens , florens ternpore pluvmli;

Pr̄viridi flavefcētibus, tñna, quaterna,
 cuſ-
 n race-

matim ſpectantur, quini, ſeni & plures, brevibus inſi-
 dentes petiolis; gemmæ rotundæ ſunt & virides, plani,
 capitulo albicante inſtruēti, pentapetalæ, cuspidatæ, in-
 terius carne viridi, transversim diſiectæ, ſtellam referunt,
 fanoris ſubacris & aromatici. *Fructus* floribus ſuccedentes,
 ex rotf^S ^{1 lu" T} flavefcēte unicum emul- f»m_{gn},
 albicans, continent, in cuius
 nda, tenuis, albicans & trans-
 rens Tfrporis ſubamari. Vires adhucdum incognitæ.

O

ITTI-



Qaro intusdenfa, flava ac humidior, faporis dulcis , lin-guam supra folia, landnans, numerolā oflicula ex flavo albicantia, plana oblonga & angufta in fē habet₅ putami-na duriori, nucleo albicante. Succus expreflus & hau-ftus inteitinorum tormina tollit. Nafcitur circa *Tarue* &c in aliis locis; fructus menlē Januario maturi.

T A B . X X V I I

Error eft quod planta *hxc* species *Cada-?ilava* J latius viderc eft in annotationibus ibi adje-fit₉ quoniam nulla inter illas convientia₅ prout j &is.




بـ
بـ

بـ
بـ

بـ
بـ

بـ
بـ




M O L A G O-C O T > I.

T A B . X I I .



O/agd quatuor funt Jpecies: *HMoUgd - Codl j CattH-£Molag6| Cdtpi-Tirpati; Capb-| Molago.* Bramannes *CAiir̄t* vocant; Lufitani *Piamenta;* Belgico Idiomate **Peper;* Piper eft ro.tufl² dum, %adice fibrofa₅nigricante; *Caulæ.* rotundi funt, virides, lignofi, duri, nodofi, ex nodulis novos hinc inde caulinosejientes. *Folia* quas brevibus* craffis, viridibus & interius ftriatis petiolis ex nodulis erumpunt₅ oblongo rotunda, anteriuscufpidata₅ in medio latifima? texturâcraifiori&folidf? superficie planar viroris anteriori parte fufci & nitentis, exteriori clari. *Floret* fpicatiin petiolis viridi dilutè confpiciuntur, monopctalij circa extrema tripartite; (*emma* viridi dilute & flavefcentes₃ intus acinum parvum, viridem, ac in vertice nigricantem, rudimentum frudtus -> habent. *FruBut* 5 acini rotundi, racematim in petiolo viridi₃ viginti imo triginta simul proveniunt, primum virides & £quali superficie₅ dein rubri; ad maturitatem perdufti decerpuntur, folique per septem odiove dies exponuntur, à quo tempore cortex nigrefcit. Semel quotannis floret hxc planta, veletiambis, si vigeſtor eft, menic Julio; quanto menfe poftflores, colliguntur frudtus. Nafcitur per totam Malabarium. Cortice nigro ablato fit piper album t Clolitur per fragmenta ramorum terra intrufa. Folia & fructus faporis funt acrioris, vegeti₅ iicci. Tbeta life plantain pulverem redacta ac cum butyro cocta, puſtulis aurium confert. Radix trita & in aqua communi decocta, pro lavamento oris adhibetur, catharros enirrl & gingivse affea



Catharanthus
ثعلب العصافير
R. Frazer Miller 1855.

C A T T U - d M O ' L A G O .

T A B . X I I I .



Alabarenfium *Cáttu-CMo'lago*[^] Bramannum *TlavO'CAdiri*; Lufitanorum *Tiementa-Qomatd* eft; Belgse dicerent *Wine Tsper*; planta plane fy 1 veftris, nec unquam apud Indigenas pro pi* pere venit> infipida nempe eft? quamvis quo- ad formam cum vero pipere maximè conve- niat. *Caules* sunt virides 5 ftriati 5 minus lignofi. *Folia* exgeniculiscauliumoriuntur, ad petiolum lūlcata, ob- longoangufta? parum rotundiora, lata? textura craffa? dura; virōrisinteriusfufci? exterius clari. *Flores* iidem? ut & fru&us, fed majores > femel quotannis florens. Suc- cusut& *Cattt'Tirpali* fimul exprellus cum bu tyro fumptus tuffi convenit.

T A B . X I I I .

Figura *Cáttu-Mo'lago*, imperfecia eft, floresl *Piperefdtnineo Pifonis* convenire videtir; cnim & fmftus defunt[^] non male autem cum I



No. 7 552




Catt.-nervosa
Catt.-nervosa
جعفر طهري
جعفر طهري

C A T T U-T I R T zA L I

T A B . X I V .

 *Attu-Tirptdt*, tertia species eft MoJago, Bra-
mannibus "Bipali; Lufitanis *Timenta Longa*
dicitur; Belgice *Lange <Peper.* %adix fibroia,
nigricans. *Cades* rotundi, virides, nodulati?
minus lignofi, novos hinc inde ejiciunt
furculos clavicuktos. *Folia* circa nodos caulum , lon-
giffimis interne ftriatis petiolis erumpunt, forma ut Cattu-
Malogo, aft longiora₃ minus cupidata ? fuicioris viroris
quam Molago-Codi •> circa bafm incifa , textura valde
tenui & molli₅ fuperficie lenia₅ viroris in re6ta parte
fufci, in finiftra clari; differunt aCattu-Molago? quod
cum cofta media duas vel tres alias tenuiores excurrant,
quarum inferior arcuato du6tu nervulos transverfos ver-
fus marginem emittit, utrinque eminentes. *Flares* mo-
nopetali , quinque vel fex partiti₅ arctiffime *fmttibus*
nfidentes, qui oblongo rotundi funt, fpiris obliquis pa-
allelis in fuperficie ex diverfis partibus ftriati, in fpirarum
nterfeclionibus velutiparvis rotundis foliolisfcutiformi-
-r obdu6ti; inter fpiras gemmae apparent quibus flores
federunt , eminentes & nigricante pundfco notate,
rimo ex viridi fkvte, carneintusexflavoalbicante; dein
ridifufcae, carne viridi fufca ; maturas & ficca[^] exci-
reo nigricantes. In fructibus transverfim diflectis in-
ftitia videntur, in quibus femina oblonga & nigrican-
continentur. Colitur multis in locis , femel quot-
iis₅ tempore nimirum pluviofo , floret. Piper tri-

turn



卷之三

B E ÉT L A ^C O T > I.

T A B. X V.

M Alabarenfium *Bēetla-Coih* Bramannum 8t
 Lufitanorum *Bekla-*, notissima ilia apud
 omnes exteros herba "Bethel eft* quas per
 totam Indiam supra Oryzam 8c frumen-
 turn comeditur? ac maxima cumcuraexc
 colitur. %adicem habet fibrofām; *laules*
 virides* rotundi 5 glabri, nodulati ? supra nodulos ex*
 tuberantes. *Folia* petiois longiufculis 3 craffis*? interius
 ftriatis? ex nodis caulum proveniunt? oblongo-rotunda?
 cufidata, texturācraflora, mollia, glabra 3 virorisfufci
 8c nitentis; faporis aftringentis; majora nullius fuit usus;
 minoraquaeintenris transverfis furculis nafcuntur mol-
 liafint, & cum *Jreequa* addito momento calcis abindi-
 genis quotidie manducantur; faporis fubacris 8c grati, labia
 Scfalivamcolore rubro tingentia; fi minus calcis addatur
 falivaex flavofufcaeft, fapor aftringens & nonnihil adu-
 ren; fin plus minus rubra5ardens. Genus eft /[^]e^«*, Malab-
 barenfibus *Scborikftna-Adeca*, Bramannibus [^]reequa-Bebeda
 hoc eft? inebrians di6his; medulla interior rubefcens
 cum *Bethel* manfa cerebrum offendit; cum radice vel
 fructu *BeUaca*, vel cortice *Elenji* & *OiU-Tulli* & exigua
 ^uantitate calcis falivam tindlurā rubrā implet. Eft &
 ecularisXpecies *Beētla Codi*, fed rarior > cuius folia fapo-
 rem garyophillorum aemulantur* *Floret* monopetalii,
 quinque₃ fex, septemve partiti 5 in brevibus petiois ex
 nodis erumpentes proveniunt₅ albi, at cum exficiati fuit
 nigri; capitula odons funtamoeni> cum tempore magis
 magifque viridefentia.

TAB.


XV,

Edparm Btubinus fruticem hunc Tembtl, A vice-
 ma Taming Garziasab Horto Eetre, Clufius Betle no-
 minavit. Confspecies eft Piperislongi. Differntiam
 iater folia riperis & Bethel doctiffimus Fabius de-
 •nontravit ut & Ant. Realms in fuis Annota-

tionibus in *Nardum*, ^37.pag 876. ubi prolix^
 de eo traftat; netnoautem meliorcm unquam
 delincationcm exhibuit, prater nobihffimum
 Do. vanRheede.

H

A.MO.



Digitized by
Digitized by
Digitized by

Digitized by
Digitized by
Digitized by

M A L A B A R I C U S .

31

A M O L A G O.

TAB. XVI.



U*i* frutex Malabarenibus *Amolago*, Bra-
mannibus *JumJJo*, Lufitanis *Pimento Ma-*
cho, Belgis *Peper het mannek*, n VOCatur^s ecce
est Malógo; *Caulibus* rotundis, geniculat^t Wn
exortu internodiorum erura^s Pentibus. *Foliis*
ut Mòlago, sed spissiorib^s us, virorefaturis,
odore Piperis, sed minus vegeto. Loco florum virides
habet ligulas, ad longitudinem extensæ manus, bullulis
cuspidatis obsitas; duobus albicantibus apicibus, qui postea
nigrescunt, instructus, sanoris acris & aromatis^s asci-
ofo.
Iyl
us.

TAB. XVI.

Amolago. species sylvestris est *Piperis longi*, & se- | les species porro recensentur. Vide *Casp. Bauh.*
cundum nostrum Authorem extra usum. Simi- | *Pin. lib. XI. sect. 3. pag. 412.*

Kun.-Kun. M. L.
212. 66
J. H. M.
Mr. فیصل



K ATU*R A'R A-WA L L I.

T A B . X V I I.



Atu-Kara-Walli Malabarenfes; *Fasil & Sato-*
Val-G'alt Bramannes; *Fruita Tatao* Lufitani;
Salk-Doorn Belgse hunc fruticem nominant;
%amot habet rotundos, geniculatos, geminatis
 supra geniculos oblongis, cupidatis
 lignofis spinis initructos; spinse teneriores in
 tenerioribus ramis, superius furgunt; in denfioribus re-
 flexas font. *Folia* ijeminate transverfim in geniculos infra
 spinas proveniunt? oblonga, in cufpidem ftrid:a, tenuia?
 interiori parte lenia, viroris nitentis > exteriori in qua ex
 cofta media, laterales obliquodud:u egrediuntur ac extu-
 berant, fcabra, coloris diluti & furdi. *Flom* supra ex-
 ortum foliorum, proxime ad spinas in petiolo brevi &
 nonnihil lanuginofo pendent, monopetali, quinque par*
 titij cufpidati, coloris viridis ac flavefcenis; ex foliorum
 interftitiis quinque apices fufci emicant; in medio ftylus
 eft > ad pedem lanagine alba obdu<Stus, cum capitulo fla-
 viuculo in vertice; nullius odoris. Gemma; virides.
FruBus* iidem cum *IWalli^ara*. Nafcitur cirrafyein-tSXJan
 gatti & aliis locis fublimioribus : femper vel flores vel
 frudtus habet & extra ufum eft.

T A B , X V I I

Frutcx hie commode inter species *Rhtmni* pojuitur.

I

V d L.



Digitized by
Digitized by

Digitized by
Digitized by

Digitized by
Digitized by

P[?] Ul L L LK c7' R A.

T A B . X V I I I .



Alli-Kara Malabarenfes > *Fruitado Cao*, Lufitani, *HondsbeJJen* Belgse dicunt; frutexeft finuofis ramis fe arboribus aiSgens ac circumvolvens. %adix cortice rubro lāguid neo & cinereà cuticula cruflatus , lignum habet album 5 interius ex casruleo rubefcens. Stipes è radice exiens craflus , lignofus 3 cortice cinereo, intus medullâ albicante fungosâ gaudens; furculi vetuftiores etiam cinerei > cortice viridi 3 teneri fpadiceo fufci 3 villofâ & mucilaginofâ lanugine induti. *Folia* ex petiolis longifculis tenuibus 5 viridi dilutis acmufcofis pendent, adversâ parte lenia&villosâ > interioribreviffimispilisinftrucla; odoris? cumteruntur acutis&noningrati. Cum cofta media utrinquelateralis prodit; exterioriparte emt nentes. *Flores* supra originem foliorum fpiratim in brevibus petiolis cernuntur, monopetali, quinque partiti, exteriora verflis reflexi 5 funtque gemmula: rotund[^] mufcofe) ex viridi flavefcetes; circa difcum flavefcen* tern quatuor fele breviffimaerigunt ftamina, singulafuis nodulis flavis dotata₅ ac farinaceo rore afperiâ ; odoris grave olentis. Poft deciduos flores fequuntur *FruBar*, globulivrides&mufcofij cortice, cum vetuftioresfunt, ob punctula eminentia alpero > intus viridi, bini mutuo uniti, ac in vertice transverlo communi ex riiffo fufco umbilico confuti & ftriati; intus unicum rotundum & totamcapiactatemreplensj femen habent; quod cortice fie-

ficcato ac fiflb lponte prodit Natale folum eft prope *Cochin* circa *Taloerti*, & in aliis locis, femper florens vel frugens. Semen cum oleo & croco coctus, rabidi canis morfum frufra facit, modo in tempore adhibeatur.

T A B . X V I I I .

<i>Valli-Kara</i> pertinet inter <i>Frutic. bacc'tf. scanden-</i> <i>tts</i> , a nemine quad feciam descriptus, inter <i>Solani</i> species rcccaferi posse, ut <i>Dulc-Atnara</i> . Vide	Dodifim. <i>Eajum HiJlor.fUm. Tom. U B. i?</i> , pan. 2. Cap. it. do <i>SU</i> no.
---	---





Bidens - L.
Bidens
Mimosa trifolia
পুরাতন খেজা
পুরাতন খেজা

পুরাতন খেজা


P E E-A' M E RT> U.

T A B . X I X . X X .



*Ei-Amerdu apud Malabarenfes, quern Bra**
*mannes Fallu-Amerdu-Falli) & (Enam no**
minant; Lufitani Trapadeir de Ceo; Beigse*
fPindtoortel, species irftnerdu eft. Ttydix fi-
brofa, capillata corticerufo-fufco₅ intus*
lignofo, asque intraac extra terram? imo
funiculo fulpenfa, &aëri expofita, nontantum crcfdt
fed & floret & fruget quod etiam nudi faciunt rami.*
taules acfurculi rotundi iiint₉, tortuofi, lenti, flexiles,
teneri virides ? albicantibus & veluti lanuginofis pilisob-
fiti; vetuftiores cortice exteriori cinericeo, interiori vi-
ridi > filamentis intus lignofis albicantibus feu flaviufculis,
laxis, carne fufcofa ac mucofa interpolati₅ pertexti,
tamefuperioriquaminferiori parte radicis propagantur,
primò ut tenuiflima filamenta, molles ? flexiles & cinc-
reovirides; dehincad pollicemcraffi. Folia hincindead
cauliumfeuramorum flexuras, petiolisoblongis, rotun-
dioribusj pilofis > ad exortum parum extuberantibus₅ pro-
veniunt, formalata? anteriuscuſpide angufto&oblongo
eminentia > ad petiolum finuolà? &pilis₃ maximèinad-
verfa parte veftita > lanuginolà ? in anteriori pili afperiores
funt j textura laxa & fragili. Coitas tarn media quam
laterales adfunt₅ poſteriori parte eminentes. Floret in
petiolis viridi dilutis₃ fupra ex origine loliorum prodeunt >
numeroplurcsj tenuinfmis ac viridi dilutis & albicanti-
*bis pasdunculis; monopetali? tripartiti₅ parvi, nitentes**
in mediotenui venulâ ftriati; inmedioquafi rofa fpedta-
tur > in qua fex ftamina albicantia emicant > nodulis flavis
& rufits, quas in gemmis albefcunt. Gemmse rotundio-
res ? fuperne non nihil planar₅ ad petiolum ftridiores,
virides; nuUius odoris. FruBus racematum lpedfantur ac
terni pedunculis brevioribus, transverfis₅ viri dilutis,
flavefcenibus infident&ex fngulis floribus oriuntur,

oblongo rotundi, primùm virides, post flavefcentes, dein corallini ruboris & nitentes, intus vifido, mofcofo hu^more repleti, non femper seque magni, fed differentes : in medio officulum habent grande, oblongum , parum complanatum, fcabrum & exafperatum , primum albicans, dein nigrelcens : Nucleus albus eft. Nafcitur in fFarapoli & fMouta> temper florens & frugens. Balneum ex hac planta & NjUSPanna-Scelenzu prasparatum abfceflus inveteratos curat: *Folia cum Carique de 'Bengala & Carari* mixta, vel cum cortice tAk 8c *Sergelim* in lade vaccifio co&a , exanthemata refolvunt. Pulvis ut & nuclei *Cteanga-Salt** fiffuras pedum confolidat; coda cum aqua *Ory%oe* pedum ulcera mundat.





جعفرية
جعفرية
جعفرية

جعفرية
جعفرية
جعفرية

C I T - A M E R T > U.

T A B . X X L



It-Amerdu, fpccies Pèe-Amerdu, nihil diffe
rens, mfi quod flores& frcdtus fint mino-
rū. L Decottum cum fummo frudu in fe-
bnbus ardētibus, arthridite, & ift_{ero}
adhibetur, vires reficit. Exiuocco & Co/utpa
ac *Tirra-tali* cum ladle balneum antispafmā-
dicum pp'patatur. *Teneritina folia* cum *BamJcSmt:*
hiie tnta, hnicntum dant in *Phlegmone* & *Eryfipe* late
Succus, ut & *Mulcenti*, *TsjiuJPulla* & *Vtmja* expreflus',
punsgenerationem pr[^]pedit & ulcera fanat. Succuscau-
liū ablato cortice, cum lafte & aqua decoctus, & ad
ficcitatemevaporatus, ac oleofoliorum de *Bnfermo* mixtūs
hiimenturn Antiarthriticumeft: cum facchāro potus cāl
chexiam tollit; cum Pipere longo pituite intemperiem-
cum faccharo & piperelongobihs dyfcrafiam & morbum
articularem corngit.*



U**L A.

TAB. X X I L

Verba, Bramannes *OulaJJo* vel *tZMacuda Jan-*
goli; Lufitani *FruitaT emacboj* Belgas *Harlaar*
 dicunt; *%qamos* habet lignolbs 5 albicantes,
 durosgeniculatos, nodoibs> extuberantes;
 cortex exterior niger eft & cinericeus ma-
 culis hinc inde notatus > interior viridis >
 tener. *Folia* exfurculis rotundis, viridi fufcis, genicu-
 latis gemina₅ decuffato ordine, oblongo rotunda, textu-
 rà spiffa &c fblida₅ superficie plana & glabra₅ antérori partē
 fufco viridia* nitentia? exteriori virore latura, parvis &c
 crafloribus pedunculis, interiori parte planis proveniunt;
Florum loco funt ligulae oblongas, rotundas₅ piper oblon-
 gum referentes₅ annulis nigris feriatiam interftinbas, in
 interftitiis viridi dilutae; annuli tenuiflmia, breviffima₅
 nigra habent ftaminula > apicibus minutiflimis &c viridi
 dilutis dotata; exgeniculis ramorum vel folkarias vel tres
 simul erumpunt; floret menfe Januario; llaminulis deci*
 fis parvae virides gemmse, circa annulos in orbem prodeunt
 ac in *FmBuj* excrefcunt₅ forma^ exafte oblongo rotundi,
 in vertice minutiflimum cupidem habentes, calice viridi
 fufco & craflb: caro interior ex viridi albicans, nucem
 oblongo rotundam in fe continet •> duro putamine₃ pilo-
 fam ac hirfutam; nucleus ex albo ilavefcit, quern mem-
 rana albicans interius ambit, faporem refert caftaneas ac
 nter edulia recenfetur. Nafcitur circa Calicut & in aliis
 ocis.



253 m.
T.M.




T E E»U L A.

T A B . X X I I I .

Pee-Vla , Bramannes (*fāngoli* 6c *Oulaffjani*) Lufitatini *Tenacbo Macho* dicunt; nofrates *Harlaar iSManne^en*. **F**rutex eft radice lignosa longe prorepente > ligno duro ex albo flavefcnte₅ medulla ruffa cum tenui corticula teneri&fquammosa. %ami nodofi, cortex fpadiceus -, vetuftior ex cinereo fufcus j circa lignum ex viridi flaveicens , ipfum lignum albicans. *Folia* paffim geminataac etiam quaterna₅ innodis ramorum brevibus petiolis, interiuslatofulcoltriatis, proveniunt; oblongo rotunda funt \$ craffa fpifia & folida ?, fuperficie plana 6c glabra₅ viroris fufci 8c nitentis₅ in parte adverfa dilutions. (*Jemmte* rotundas funt, fuperne in cuſpidem ftriſta3, ex viridi fufese. *Flores* \$ fupra originem foliorum in ramis oblongis racematis proveniunt, 6c pedunculis parvis viridibus appendent j ex viridi diluti, oblongi, paffimodlo petalis cuſpidatis conftantes > exterius quatuor oblongoribus 6c viridi dilutis ftelliformiter expands, interius fimi-liter quatuor alblicantibus 6c obtule cuſpidatis₅ qux fingula inter duos cuſpides externos interjacent, intus flavo circulo fignati. *Stamina* lēx vel oſto babent pro numero petalorum , alblicantia₃ croceis apicibus inftrudla : ftilus oblongior* craflor ac viridi dilutus; marcefcentibus floribus ftamina rubefcunt. *FruBus* nulli: ufuſ hue uſque incognitus. Naſcitur circa 'Bellange 6c in aliislocis, florens tern pore pluviali.

T A B . XXII. 6c XXIII.

t^iaravis ha: *dust* plantx idem cognomen ha- flicx diffcrant ; fed in defcriptione Authoris
beant, tamen nuliaincer illas convenientia, ni- | noſtri acquiescentcs^ de illarum **ulteriori** inda-
xiuum enim quantum nota: illarum charaſteri- | gine ſuperfedebimus.




Bufellia ex
Digit.
पातो - गुण
केला - दी

Digit. Linn.

B A S E L L A.

T A B . X X I V .



Rutex qui Malabarenibus 'Bafella; Bra*
mannibus Wali\ Lufitanis 2?ra/0-7V/; Belgis
<Beet{lim dicitur. Cau/es aquei funt, ex una
parte viridcs, alterā verfms folum ex pur-
pureo rubelcentes. *Folia* in pedunculis
craffis, brevioribus ? interius ftriatis & viridi
dilutis proveniunt, ampla , cuipide obtuso. *Cofta* in
adverfa partc craffie ? viridi dilutee & albicantes. *Flores*
cyathiformes, tetra vel pentapetali, informs ex viridi al-
befcentcs -, fuperius ex purpureo rubefcentes. *Stamina*
quatuor vel quinque ; quorum apices in medio ruffi.
FruBts caulinis infident , supra cxortum foliorum,
numero fex₅ feptem vel odto₃ iuntque bacc^e piano ro-
tundse, quatuor quinqueve rotundioribus interfittiis in-
flrudix, primo virides , maturas ex rubro nigrantes
& nitentes₅ faporis aquei fubdulcis ; intus carne fuc-
culenta? cuius fuecus expreffus? tin&uram ex purpureo
ubram praebet. Officulum in medio albicans, in quo
i ucleus albus. Lignum cum foliis tritum carbunculos
x>erit: inter edulia habetur: Colitur in hortis.

T A B . X X I V .

Bafella hxcaZanonio *Bafila* dicitur. Semen j cus ex fruSHhus expreffus commodiffime pro
A. L. 1685. ~ex „Ceylon transmiflum in horto miniatura utitur. Planta hicq; longieva eft, ^{fed}
Medico Amftelodamenfi , plantam produxit; quam cautiffine pro frigore confevanda. Flo-
non Tub BetV fed Bryonia recenferi poteft. Sue- rum figura in dclineatione plus debitò major.

M

C V-



CURINIL vel CURIGINIL.

TAB.

XXV.



Vrinil vel *Curigil* Malabarenfes; *Thali* & *Kenny** Bramannes; *Crica da Velba Tequen** Lufitani; *KleinHondepint Belgxdicaat*. Species eft *Terim-Curigil*; vide defcriptionem •*Tab. 24. Tart: 6. Stir ados* habet rotundos, ocniculatos, in geniculis nonnihil extuberrantes, intusiignofos, cum corculo viridi. *Folia* ingeniculis geminata, decuffato ordine videntur in petiolis oblongis ac flaccidis, oblonga, cupidata, texturâ molha, flacciða, superficie plana & glabrâ, viroñsm redaparte faturi, in adverfa dilutioris odore & fapore deftituta. *Cotta* media ex viridi albefcens & in adversa parte eminens, plures laterale semittit. *Born* fafciatim proveniunt pen-tapetalii ex albo flavefcentes, extensus vindes, mtenus cāndido, lanuginoib uncinulo vcftiti; in med.o globulus eft rotundus, quinque ftaminibus parvis fuccmclus. IS ulliusfunt odoris. *Florum (femm*) oblongo* rotundas, m vertice albicantes. *BuBm* oblongi iunct, cortice vmdi claro, came albicarite fubhumida, lapons lubaman; officulum in fe continent durum & albicans, quod nucleum album includit, cujus fapor fubamarus & nonnihil adftringens. Natale folum eft infula. *Tlayptm*, femper frugens. Foliorum fucus loco balnei adhibetur, adverfustumorcuquem Malabarenfes *Tsjddcmir*, nuncupant; radix in aqua oryzse cofba & pro potu ufurpata, idem prceftat.



C U R I G L T A L I.

T A B . XXVI.



Vrigi-Tali; apud Bramannes Tali dau K,tinapa;
Lufitanos (/teā davelha Trapadeira ; Belgas
(froot ^limmend Hondepint. Taly species eft,
nafcens in arenofis prasfērtim in Jngiecaimal.
%adix fibrofa •> prof unde fe demittens ? cor-
tex nigricans & maxime rubicundus. C^{a^m}
liculi lignofi cortice fufco ex radice prodeuntes hinc inde
in nodulos protuberant. Folia ex furculis lateralibus bina
proveniunt, tertiumque in vertice emittunt, oblongo
rotunda₅ cuipidata ? inperficie glabra₅ textura⁵ deniori,*
ex viridi fufca. Cofta media in utraque parte eminens,
nervos aliquot annulari du6tu producit Flores in petiolis
tenuibus plures congregatim convenient? pentavelhex-
tapetali , oblongo rotundi₅ cupidati, candidi₅ odoris
*expertes. Stamina in medio plura₅ tenuia> parva₅ albi**
cantia₅ apices crafiores ac ex flavo albicantes habent.
FmBm oblongo rotundi, nonnihil incurvi -> &c in parte
convexa futura ftriati, primum virides₃ ex in rubefcentes;
unicum femen in flavori membrana continent₅ oblon-
gum , planum, convexum. ex cinereo fufco rubefcens.
*Succus foliorum cum laſte ebutyrato > & chalybeato, hse**
morroidibus medetur : Fruſtus in febribus malignis₅ ad
fedandos ardentiflimos calores adhibentur.

T A B . XXV. & XXVI.

Fru&uum fortnam fi fpeftes pro Ipecie *Per'tm-* | turn differant. Semina autem *quxc in vcficuii^*
Curigil part. V I. fag. 45. Tab. 24. deſcriptum | funt magna ex parte conveniunt, dadylum refe-
haberi poſlunt ^ quamvis in aliis partibus mul- rentia quamvis non ita dura.



دہلی
دہلی
دہلی

۷



T S J O R I - F A L L I

T A B , IX.



Sjóri-Falli; lingua Braminiim Cerfori+Vdiiij Lufitanorum Vvas D'atjofare -, Belgarum Tkt Timper/in^%. adix carnofa₃ albicans? humida? vifcofa, transverfim incifa? ftriata, fibrofa. Caulir viridis, vifcofus? intus carnofus? clavicularius. Folia fcrrata₅ craffiora? mollia? glabra, lenia> ftriata, in brevibus fulcatis con- Ipiciuotur petiolis. Floret racemati -> monopctali? quadripartiti, ex rubro-fufci. Stamina quatuor₅ subflava, apicibus dotata, ftylum medium ar<5te amplefuentia. Fru&us uvas piano rotundas, eadem forma ut^e Bekta Tsjo'ri-Fat/i> primo virides, tum albicantes & nitentes, dein rubefcunt & casrulefcunt, ultimò ex fufco in totum nigrefcunt. Caro vifcidaelt & mucilaginofa₅ fucco rubro casruleo repleta; Officula triangularia funt, quorum duo latera plana tertium rotundum, rubefcentia. Locus nativitatis eft nJroer & CAdoutan. Ex foliis cum medulla 2s(ucif Indie* contritis, fit unguentum contra Colicam. T\$adix in aqua Oryfe co6ta& expreffa, ulcerafanat.*

Betula sp. - will
o. m. g. h. n. t. v. m. a.
Kashmir - will
نیالہ جاہیں
بَلْوَانَ

بَلْوَانَ

BELÚTTA-TSJÓRLVÁLLL

TAB. X



*Elutta-Tsjori-Fdlli; Bramannes vocant CerberU
Valli; Lufitani Vvas cfJljafare CMacbo; Belgas
Timperling z5Manneken, species Tsjori-Vdlli|
eadem radice gaudens. Cau/icuH rotundi, pi-
lofi 5 intus lignofi, fragiles, claviculari.*

*Folia ex adverio clavicularum prodeuiit, in
medio unum 5 ad latero duo, tria, quatuor, imo quinque,
oblongorotunda? cufpidata, ferrata, pilola? afpera, an-
teriori parte ex viridi fufca, faporis acrioris & ardentis.
Flores monope'cali 5 quinque partiti 5 racematim ex parva
& viridi fufca gemma oriuntur, odoris expertes. *Stamina*
quatuor parva albicantia, juxta orbitas ambitum erumpunt?
flaviulculis apicibus inftrudta; *Stylus* brevior è medio
centri emicat. *FruBm* racematim pendent -> uvarumin-
ftar, forma^rotundo plani, fpiculo muniti, in principio
virides, postea albicantes, eburnei, splendentes, mol-
lem5 untuo famacpelludicam carnem continent? fapore
admodum acri; officula unum, duo, quatuorve, una
parte rotunda, altera plana; nucleus ex eseruleo albicans.
%epoli & aliis in locis invenitur. *Folia* matura cum rafura
U^ucisIndiae icabiem tollunt; cumoleomixta&concodla
vulnera fanant; fuccus calci admixtus, populas curat.*

SCHV-



دہلی جنگل
کے ٹوپی

SCHUNAMBU.FALL

T A B . X L



Cbundmbu'VdlU •> Malabarice ; Bramannibus
Cada-Pdlli; *Lufitanis Vvas iEmfermot;* **Belgis**
Snoepdruiven. *'Radix* admodum fibrofa* filamen-
tis contexta & albicans? plures emittit
caules viridesj hinc inde velut cake feu
farinâ albaâ afperfos •> nodulofos₃ claviculatos , molles ,
albiffimo fucco inftrudtos. *Folia* , longis ac tortuofis
petiolis è nodulis prodeunt > vel caulinis transver-
fis infident, grandia? oblongo - angusto - culpide mucro<
nato₅ tenuia₅ fragilia₃ anteriori parte fufca₅ adverfa ni-
tente, male orentia. *Floret* racematin in caulinorum
transverlbrum fuperiori parte vifuntur, monopetali, qua-
dripartiti, albicantes; gemmas parviūcute , ftylum in
medio habent, ftaminanulla. *FruBut*, primiim oblongo
rotundi, virides; poft^ecrotundi-, valde nigri? nitentes,
molles 5 iucculenti, undtuofa carne repleti; faporis inten-
fiflime fervidi. *Ofikulum* ex cinereo > nigricans eft 8c
rotundum; nucleus ex albo casruleus. U bique locorum
invenitur modo denfa adfint nemora. Expalmitibusejus
corbes conficiuntur in quibus edulia refervantur > unde
nornen *Cada-Va\li* > nzm *Cada* Bramannibus corbemnotat.
Infigne medicamentum hie frutex eft; fuccus expreflus
:um oleo codtus₅ ac formâ emplaftri adhibitus craflbs
ligerit humores. Decodlum cum faccharo, in febribus
rdentibus & pleuritide fummo cum frudlu exhibetur:
Vqua extrunco ffillatitia cum faccharo idem praeftat? tut
im lenit₃ fanguinem purificat₅ vomicam pulmonum
F curat.

22

H O R T U S

curat. Radix trita & aquâco&a odontalgiam amobet *I*
fcetorem gingivarum: cortex tritus incarnationem ulcc
rum promovet.

T A B . VIII. IX. X. XI.

Omnis hx species, ad genus hederxt per-1 enim, *Tm tices bacciferi scandentes, non secundum*
tinent, & inter illud recenteri debent; funt |



CMU

M A L A B A R I C U S, IJ

V A L L I A-T Sf '0%I-V A L L I.

T A B . V I I I .

W *Attid*Tsj6ri4"dlU apud Malabarenfes > frutex eft Bramannibus Sjora/Io; Lufitanis Vvatde Tove, Nofratibus Timper/ing di&us. %adix lenta eft & admodum fibrofa₅ cortice nigricante₅ intus ex ruffo albicante : Cauliculi rotundi₅ teneriores₅ virides, geniculati, afperi, claviculati. Folia in furculis afpcris ex geniculis radicum oriuntur₅ terna > quaterna -> ac etiam folitaria, oblonga₅ angulis acutis eminentia; texturâcrafla, denfa, >lana, glabra, anteriori partefufco vindia, adversâ dilu-a₅ furda; fapor lancinans & adurens. Flores in furculis bliaceis racematis videntur > parvi, monopetali₅ quadri-Dartiti? cuspidati₅ ex viridi albicantes. Stamina quatuor laviufcula, nodolofa. (jèmm* albicantes, planas. FruUut iv^e rotuudas₅ grandiores, racemati₅ primo virides₅ ex in Ibicantes₅ eburneas. Caro mucilaginofa₅ transparens. Acini oblongicupidati, bini ternive indiftin&is cellulis èparati, pelliculâ albicante ardte obtefti, unâparteconvexi, altera^plani, duriores > ex ruffo rubefcentes; faporis in totum lancinantis. Nafcitur in Berfycour , & fylvis denfis, femper frondens. Floresmenfe Junio & Julio; Frudtus Odlobri & Novembri leguntur; afthmati convenient.*

TSfO-



VAL LIA-TIR A-PITICÆ.

T A B . VII.




Allia-Tira-Thica Malabarenfes; Catulama Bra-
 mannes ; *Vvas iElipbanti Lulitani; fruti-*
cem hunc nominant; Belgae sStearnsdruivem
Cau/is craflus , nonnihil viicofus, adftrin-
gens, exviridifufcus. Folia impedunculis
oblongis? craflis> ex viridi fufcis; tortuo-
fisacrigidis? intus fulco profundo flriatis > amplafunt&
triculpidata ? ad pedunculum femi rotunda ? profunde
*excifa? rigida, alpera ? fraſtu facilia ? anteriori parte fulco**
viridia? poſteriori magis diluta, ubi & cofbe ramofe plus
fblitò eminent? iaporisfubamari? aftringentis. Flores ra-
cematim provenientes ? monopetali fuit ? qiiinque partiti?
cufpidati? ex viridi albicantes. Stamina quinque tenuia;
apicibus albicantibus dotata: Stylus cum capitulo albicans.
FruHus uvae praegrandes ? rotundas? virides; intus pulpa
fucculenta, adurens ? duobus vel tribus acinis inftruſta.
*Pro loco fibi eligit denia nemora in Tar be *ZMangatti &*
aliislocis, plerumque flores & frudus habet.

T A B . V I I .

Duae hac species fuit ex gcnere uvarum fyl- ncm infeit > in India & prxfertim Batavia^s
 vcftnum & inter vites recenferi debent, nam fe- fecundum cnarrationcm *Xieihovri* in suo *Itine-*
 cundum earum notas genencas fuit *Trutic. [can- ratio , pag. 2x7. vites fcmpre virent, & diverfi-*
baccifer, foliis deciduis non fp'mofis, caprealis feu temporibus maturos proferunf tru&us, ut fub
daviculisinfrutti, fruttus racematingerentes-y quam- illo calidiori ca:lo pluribus plantis accidit j nos
 vis noſter Author non dicat quod folia annuatim ergo pro vite fylveſtri habemus. Vide *Rajnm*
 decendant. Differentia enim Climatis mutatio- *Method. Flantar. Fol. 46,*

Sialia pumila Linn.
Gmelin
M. A. C. D. 1758
نیلی سیالیا



Ifterum ocu) mim toUit. Radix trita, cum oleo & ladte
Qoam-Indid-> [Liftulas , carbunculos & apoitemata fanat:
Succus radicis cum faccharo , catharticum eft ₅ humores
pituitofos evacuans.



SGHÈMBRA-VALLÌ


T A B . VI:




Cbembra-Pdlli; linguâ Braminum %am-Diacq;
Lufitanorum Vvas dos "Bugios, Belgarum >
sJpe Druiven. %adix nodofa quattior quin-
*que ve craflas diffundit prdpagiries * molles j*
aquofas; exterius nigricantes, intus albefeen-
tes, poftea magis magisque rubefcentes?
*faporis'aquei aft parum acriorisi *Caules* tenues, lignofij*
*nodofi, lanuginofi, fub Ianugine viridcs. *Folia* in nodulis*
longiufculis, rubefcentibus -> craffiolis petiolis prodeunt,
anterius cupidata •> in margine denticulis incifa, anteriori
parte viridi fufca 5 nitentia, Candida Ianugine obfefla; ex-
*teriori viridia, splendida. *Flores* racematum in petiolis*
craffis, rigidis rubris, lanuginofis, è nodulis foliorum*
oriuntur ac claviculos emittunt. (/emma* primo virides,*
*dein ex viridi diluto albefcentes & flavelcentes; *elt* autem*
flosmonopetalus, quinque partitus •> cupidatus, rubeus,
exteriora verfus reflexus; gemma media Tubrubra, apice
rubro in vertice notata: ftaminula quinque fubrubra? f lavis
apicibus dotata, irttra folia & medianam gemmaffi orientes?
*è medio foliorum eminent. *FruBus* uvse planè rotunda,*
initio virides, poft h^c nigrse, craffiori tunica, medulla
fucculentâ, acidâ, intus quatuor trilaterata habentes gra-
nula> circa tunicamrotundiora, de castero plana, colon's
dum immatura funt, fubtlavi & nitentis, nucleo albo
*inftrudta -> faporis fubamari 6c aftringentis. Circa *Oedeam-**
*pere, *Candanata*, & aliis in locis invenitur, Continuo iloret*
& fruset > femper virens. Succus cum oleo mixtus

i z H O R T U S

Idterumocolorumtollit. Radix trita •> cum oleo& ladle
Coqut-Indici, puflulas , carbunculos & apoltemata fanat;
Succusradicis cum faccharo , catharticum eft» humores
pituitofos evacuans.



PJL-



جعفری
کاظمی
علی
علی
علی

SCHERU-KATU.FALLCANIR^M,

V E L

T S F E R O - K A T U - V A L L I - K A N S F E R A M.

T A B. V.



Cheru-K[^]atu-Falli-Cdniram, vel *Tsjt'ru Katu-Válli-Cdniram*, Malabarenfes; Bramannes *T>avo Cdro* dicunt; Belgse *Wild' KjimmendKraanoo?*, i species eft *Cdniram*. %ami cinericei flint. *Folia* obfcuro viridia; *flores* monopetali •> quinque partiti; *Fmttm* curcis pedunculis ramis infident₅ globoli, deformes -, minores quam *C[^]niram*, cortice glabro, maturi primiim flavefcunt ? exinderubet cunct, Arantiorum inftar, came intus mucilaginofa₅ oleofa ac vifcida, ex rufo viridi flavefcens, tribus quatuorve gaudens cellulis, in quibus includuntur *Semina*<, forma peneut *Q[^]ram* > parum oblonga₅ plana₃ albicantia, cufpidata -> quorum pulpa durior, nervo(a₅ transparens. *FruBus* amari, *femina* minime. Nafcitur in infula *Ttaypyn* & intra *Qochien* 6c *Toaha* femper virens. Floret men fe A uguftto; Fru&usmenfe Januario leguntur; contufi & capite impofiti₃ maniacis profunt. Radix trita cum pipere > diarhasam fiftit & ventriscruciatus; cumoleo co6ta& illita artuum dolores lenic

T A B . II. III. IV. -V.

Omnes has species nofter Author fub cognoscione *Cávram* defcripfit, cuius etiam mentio in animadverfionem in partem fequaentem difcrcit, cujus etiam mentio in mus, ubide vero *dimiram* Janus diffcrendi non prima pane hujus Horci fada. Noftram vero deerit occafio.

بِسْمِ اللَّهِ الرَّحْمَنِ الرَّحِيمِ

اللَّهُمَّ

كَرِيمُ

رَحْمَةَ رَبِّ الْعَالَمِينَ

أَنْتَ أَنْتَ الْمُكَفِّفُ

وَالْمُنْجِي

وَالْمُنْهَى

وَالْمُنْجِدُ


وَالْمُنْهَى

ر

/



/



nucis guflum refert. Nafcitur circa *Tanani* -> *Catour* & alialoca; semperviret, florens tempore pluviali. Succus cumoleodecodtus & capiti impofitus humores pituitofos elicit.



SGHER TJ-VAL L LCANIR AM,

V E L

S FÉROU-VALLI-KAN S FERA M.

TAB. IV.



Cbérū-FaUi-Cdmram > Bramartnibus Ttav Roma-
Cárō; Lufitanis Fruita Tardao Trapadeira;
Belgicē KJein KJirnmend Kraanoo[^], fpecies eft
Cdniram. Ttydix longa > crafla, fibrofa; cortex
cinereus, fubdulcis ; lignum fubamarum.
(julls craflus, tortuofus ad altitudinem duodecim pedum
aflurgit, ramos quatuor cubito longiores transverfim ejiciens,
cortice viridi, cinerea & fcabra crufa obdu&o. 2^w/,
duri, lignofi 5 cortice ex viridi fufco, nitente, longioribus,
viridibus, ipinofis eminentiis minutis > faporis amari. Fo/ia
furculisramorum, ac étiā spinolis eminentiis? petiolis
brevibus & viridibus infident, oblongo rotunda, anterius
contracT:a&in cufpidem definentia, crifpa, texturâ
craffiori, denfa₅ superficie, plana, anteriori parte fufco
viridia? [nitentia amara. Flores monopetali 5 fupraori-*
ginem foliorum 5 in brevibus viridibus petiolis fpicatim
proveniunt, exteriora verfus quadripartiti, reflexi. Sty-
lus viridi dilutus cum capitulo viridi fufciori in medio
eminet; (lamina quatuor parva ? tenuiffima albicantia >
flaviufculisapicibus dotata. FruUm, qui bini ternive
petiolis parvis infident, baccas funt oblongo rotundas? fpi-
culâ parvâ in verticeeminentes 5 primòvirides& nitentes,
tumflavi) faporis fubamari: Nucleus in officulo albus,
nucis



طَبَرِيَّةٌ حَلَقَةٌ

بِرْهَنُ الدِّينِ بِرْهَنُ الدِّينِ
أَبْدُ اللَّهِ بْنُ مُحَمَّدٍ

أَبْدُ اللَّهِ بْنُ مُحَمَّدٍ

كِتَابُ الْمُجَاهِدِ


VZALLI-CANIRAM.


T A B . III .



Ui Frutex Malabarenibus *Pdlli-Cdniram*, Bramannibus *Roma-Pdlli-Caró*, Lufitanis *Fruita Tardao da Sena*, Belgis *t(lein Kjaan oog* dicitur; specieseft *Cdniram* feu *U\jicis vomic**, nafcens in declivi montium ; circa *Taracavo* & alia loca. *%adix* fibrofa eft , faporis amari. *Cauhculi* tenues, rorundi; virides : *cortex amarus*. *Folia* > hinc indeex tenuibus petiolis viridi dilutis provenientia, oblonga funt adpediculum rotunda, anteriora verfusin cupidem contradta , superficie plana ac glabra , parte fuperiori nitentia, ex fufco viridia, advcrsà communis viroris, faporis amari & fylvefris. *Flores* fafciatim in longiufculis ac viridi-dilutispetiolis, ex baft foliorum nafcuntur, monopetali parvi₅ flavi, trifoliati, rotundâ orâ in cupidem confadli. *Stamina* in medio fex₅ parva; ex flavo albicantia, nigro apice interius notata, inferius foliolis valde parvis in orbem cingba. (*jemma* rotundas ex viridi flavefcunt; apertce, oblongaefunt & flaviores. *FruBm* oblongo rotundi, parum plani, inferiori parte conftri&i & angufti, primum virides* deinflavi, minus ftricti, crafiores & formâ magis oblongo rotundi; decidente flore racemiatim proveniunt, feptem vel oclo ; ofliculum in fe continent oblongo rotundum, ftri&um , superficie afperum , colonis primùm candidi, exficcatum ex rufofufci, faporis in totum amari. Floret mtnfe Majo > Junio &c Julio, femperfondens. Morbum curat facrum. Decodtum foliorum , puftulas & ulcera tollit, externè nempe applicatum. Folia trita, idem prseftant. Succuscumoleo *Coqui India ana* paptes sequales con-

co&us, capitis & dentium dolores mitigatj fpontaneam corporis lafEtudinem toUit; cum cera ad unguenti confl-
fftentiam reda&us, ulceragravia&inveteratafanat. Radix
trita & aqua *Oryt** permixta -> humores pituitofos ex-
tenuat






جعفری
جعفری
جعفری
جعفری

TSJEREXJ-CANIRAM

V E L

TSJEROU-CzANSJERzAM.

T A B . II.



Sjereu-Caniram Malabarenfibus, Bramannibus
Tilo-Caro, Lufitanis Fruita Tardopequeno, Bel-
gis KJeinKjraanoog) species eft *Caniram* > *ramit*
 rotundis > lignolis; *cortice ex viridi fusco %*
ligno albicante, medulla^ viridi. Folia 9 ob-
Iongo angulta in cuspidem acutum deiQnen-
tia 5 in furculis propriis? ex brevibusproveniunt pedun-
culis, craffa, Iblida, superficie glabra, faporis amari &
ardentis. Flores monopetali quadripartiti > in iurculis fo-
lioifis extuberantes, petiolis viridibus, bini ac terni trans-
verfimerumpuntj & racematim excrecentes, numerum
plerumque duodenatum implent , gemmae speciem re-
ferentes> ex viridi albicantesj anterius ftri<5ti 5 cuspidati
stelliformes, exteriora verfus reflexi. Stylus viridiufculus^
cum sua gemmula viridi, in medio eminet, ftamina nul-
la5 neccalix, odoris expertes. Fru&us baccas rotundas ex
viridi fuscae, cuspidem breviorem in vertice grentes, in-
tus nucleus album habent, faporis subamari. Nafcitur
*circa Toiga^ Ca/ello & alia loca > femper virens > floret tern**
*pore pluviali. Succus ut & *Tsjetti Mulla*, ulcerofam fca-*
biem & artuum dolores tollit

VALLh



جعفری
کوهستانی

دسته گل

گل

۱۲۷



Pl. 42

D. - Will. Linn.
Wallacea
سرمه و زالی


TSJANGELAM-PARENDA.

TAB. XL I.

»  *SjangeUm-f'Parenda')* a Bramannibus *5M.am\ II a Lufitanis Bahofa Qmnado; a BeJgis Quyl/tfim ! nuncupatur. *Caulet* quadrangulati, angulfs \$ acutis 5 ac lateribus planis •> quorum duo I adverfa latiora funt, geniculati > cortice vi-
 ridi fufco & glabro, intus carnoii, carnē
 humida? denfa& vifcoia, minimē lignofi, exgeniculis
 clavicularis emittentes. *Folia*, quae in geniculis caulum
 proveniunt, lata, antenus ftri^tiora & obtusa ora 3 acutis
 fubtilibusque cuipidibus eminentia? texturacrafla, ipifla,
 moJlia3 fupcrficie Ieni& giabra3 virorisin utraque panc
 iaturij aft in adverfa fufcioris. *Cofla* nncdia duos et nitit
 lateralesj inre^h & adveria parte eminentes? & plures
 minores difTudentes. *fyimi* exgeniculis dependent? in
 furculos plures ramos fe divaricantes ; ex viridi fufcf,
 rotundi, intus aquei & virides; iiccati materiam nigerri-
 mam atramenti & carbonum inftar in ie continent. *Flora*
 parviunt, ut & *femina*, Nafcitur circa *Coykn* & *Tu'evan-*
toor, fruget tempore pluviali. *Folia* contrita aurium fu-
 ligines muncfant; fuccus cum Cardamomo, Coriandro
 ic Gingibere fumtus, in teftinorum flatus difcutit.

TAB. XL I.


Planta hæc inter fpt:acsrepentes s<& numa-andae fl.



• *Phaseolus Parvulus* L.

A N A-T A' R V A.

TAB. XL



N-a-Párvia; apud Bramannes 'Bender It; Lufitanos To/Ai <*/^wr*; Belgas *Trangwortel-*) profecunda specie *Tinda-parva* habetur, cuius defcriptio *Tomoprimo* videri potest. ^&e/ fibro-/x ex caulis inferioribus/uper terram repentibus hinc inde proveniunt. *Catties* rotundi virides, claviculas, quibus fe arboribus affigunt₃ emittentes. *Folia* quce caulis & furculis infident, velut ex duobus foliis diiFormibus, conformata funt; quorum inferiora loco petioli; fuperiora oblongo rotunda₅ in cufpidem contradta , venulis obliquo & arcuato du6bu verfus cufpidem ftriata; inferiora oblongo-angufta, anterius ubi foliis fuperioribus annexa ? nonnihil dilatata, ac in Iongum venis ftriata -> textuña fblida₅ fupeřkie plana, glabra₅ viroris communis ? laporis fibamari & fubaitnn-gcntis. *Floret* nulli. *Fru&us* hinc inde in caulis & iurculis ex origine foliorum , petiolis incurvis -> brevibus^ craffiufculis ex rubicundo fufcis proveniunt; capitula oblongo rotunda₅ paralellis lineis le mutuo intercxantia? ftriata, pluribus parvis gemmulis extuberantibus? è quibus minuti culpides emicant? interfincia; colore primura albicante ? dein viridefcente; ultimò rubrococallino ac parum nitente: infrafructus in petiolo unicum & quidem minutum confpicitur folium, rotundum ex rubicundo-obfcurum, verfus exteriora reflexum, interiora contradtum, quo frudtus tencr & albicans, ab una parte obtegitur. *Semina* in frudlibus numerofa₃ velut femina pifcium continentur, colons fubflavi, viridi dfluti.

^r

diluti. Locus nativitatis eft *Chanottl*, Tar OH & alia loca; femel quotannis fruget tempore pluviofo. Vires esedem cum *Tinda-Tarva*. In febribus ardentibus & ma iignis adhibetur.

T A B . X L.

In prima parte hujus horti f*g. 88, *Ana* I totocado aſc inviccmdiſſerunt, quamvis nomi-
Tarva ut ſpecies *Tinda Ptrva*, mentio faſta ne poenē convenient, id quod pluribus in locis
fuit 3 quae hie pleniffimc defcribitur. Vcruci, | in hoc opere videtur, 6cannoſQtumcft.



TSFAN.



Chlorophytum Topur
तूपरी च्लोरोफिटम्
M
०८२४०१. श्र.
उत्तर प्रदेश
वन विभाग
काशी वन्यजीव संग्रहालय

É R i M A · T Á L I.

T A B . XXXIX .



Alabarenium *Erima-Tait*; Bramannum
T>ou Tarvati; Lufitanorum ^aix da foheira
eft; Belgas vocant *Kjeeftbout*. fydix lon-
ge lateque fe circa terras superficiem diiper-
git. *Caules* rotundi funt ? tortuofi? cinerei,
mufcofi? Ipadicei. *Folia* pedunculis bre-
vioribus ? intenus ftriatis 5 ex viridi flavefcentibus &
mufcofis erumpunt, oblongo rotunda 5 texturâ craffa,
lpiila 9 Iblida. *Co/la* media in adverfa parte eminens plures
futiliores venas ? in adverfâ parte extuberantes producit;
nee odorem, nee faporem habent. *Fkm* in oblongis pe-
t; »lis 5 plures congregatim & racematim confpiciuntur
d plices, pentapetali> interiores primò candidi & niten-
t .5 dein cum nitore flavefcentes. Exteriores ex ruffo
befcentes, incifi* in medio plica^ftriati; in orificio api-
c J candidi, lati, foliofi 5 cufpidati emicant 5 in meditullio
globulus ftellatus > quinque cuſpidibus inter apices inter-
pofiti5 videtur; qui primordium frudhiseft; odorisgra-
tiffimi. *Florum* (*femnx* rotundas , fpadiceac , villofe,
infra villofam tunicam virides; ex quibus hiantibus 5 capi-
tulum fpadiceum fe exferit, inftar ferici fulgcns; *Frifdus*
mufcofi 5 oblongo rotundi, in quorum medio globulus
oblongus? ex viridialbicans, continetur. Nafcitur circa
(*ranganoor*, *<fMangatu* 8c aliis in locis, florens tempore plu-
vizll Succus expreffus cum oleo Nucis Indicas & pulvere
Cumini coitus, vilcera *finis* mitigat & fanat.

M



Entom - 1st. L.
sp. sp. sp. sp. sp.

جبل طه - 1st.

1st. Entom - 1st.

جبل طه - 1st.

KATSJI KELENGU.

TAB. XXXVIII



dtsji I|élengu. Caules habet quadratos 5 vi-
 ride_s amgulatos? alatos? criipatos, acin
 alarum ora nonnihil rubefcentes_s geniculis
 interftru&os, intus lignofbs. *Folia* in geni-
 culis geminata & ordine decuflato in pedun-
 culis oblongis quadrilateratis proveniunto
 oblonga, lato finu excifa, molliaj lenia> viroris in re&a
 partefaturi. *Co/la* media utrinquetres laterales emittit
 alatas ? candicantes Sc nitentes , venuis transverfis fub-
 rubefcentibus intertextas. *Floret* nulli. *FruBus* cortice
 fpadiceojinteriusfanguineorubri? carnedenfa₃ albicante,
 vifcida? caulibus ad exortum foliorum infident? funtque
 velut radices_s quas folia ut & novellos foliorum caules
 emittunt, & radiculas ceu fibras albicantes > ubi terras in-
 cumbun"yfundentes. Foliorum fuccus fcorpcionum morfui
 medeiur. Radicis pulvis cum *Catu-Tanna-Kjelengu*, ulce-
 ribus malignis fummo cum fucceffu infpergitur ; fotus
 loco adhibiti idem prasftant.

TA B. XXXIII. XXXIV. XXXV. XXXVI. XXXVII. XXXVIII.

Frutices hi fpecies funt *Batatti fylveftris* àne- | A⁰. 1686 exCeylone **transmifli**. Radices ibi-
 mine unquam , quod fciam, defcripti 5 omnes | dem à porcis & aliis **animalibus comeduntur**.
 inHorto Medico Amftelodamenfi inveniuntur,



K Á TTU-K E L A' N GU.

T A B . X X X V I I .




Alabarenium *fytu-l{elangu*; Bramannum *Carando*; Lulitanorum *Frutta burruga penacbojo*; Belgarum *Tluymkgylen* > species est *J^uren Kflengu* vel Kalangu. %*i'aix* major & craf fioreltexterius exipadiceonigricans* hinc indetenues emittens fibras, intus carne alicante > fubflava? denlā₅ humida₅ nonnihil undtuofe c laxis filamentis pertexta₅ faporis aquei, odoris nullius, ang[^]vaacfemperderivo furculos emittens. *C^{al^es te}" mes*, rotundi > duris acutisque fpinis hinc inde obfiti? eniculati> inferius nigricantes. *Folk* inftar cordis hu^{*} nani> ad petiolum profunde incifa, planalunt, glabra, enia, mollia. Cum *cofta* media utrinque tres transveralesj duftu valde obliquo, excurrunt, venistransverfis, n adverfa parte nonnihil extuberantes concomitata[^]. *Ftorer* Jures in petiolis brevioribus fupra exortum folioRim fpiati oriuntur, funtque gemmulas parv[^]e, nullum fru<3:um antes. Cum hiiče fimul proveniunt *Fru&ur* rotundiuf jli, irregulares, triplo vel (juadruplo grandiores quam *uren f^elengu*, cortice tenui 5c fubamaro, intus carne lenfa₃ folida, humida, unfstuofa₅ fubflava> faporisaquei> lllumfemen habent; femper viret, florens menfcJulio k fequentibus. Nafcitur circa *Cocbien* & in multis aliis Deis. Fruclus à plebe comeduntur. Radix cum (*joco* mixta&appofitacrurum tumores diffipat; fuccus foliorum cum urina vaccas dolores aurium fiftit.

Kella - Achaea - L.

بَرْبَادُوس . الـ

قَطْلَعَةِ الْأَنْجُو

Gossypium
कान्दो - *Bom*




K AT U-K AT S J I L.

TAB. XXXVL



*Am E^ujil; Bramannibus Karendi; Lufitanis
Grabofa & Fmita'Barruga; Belgis tfylwrni/y
Key ten; Ipecieseft TsjagariJ\uren i reipedtu
radicis, frudtuum, & verrucarum ? qux
& totam producunt plantam. Caulet te-
nues > nonnihil lignofi, fe invicem finuofe
impleflientes. Folia petiolis longis, angulatis & interius
blcatis, ad exortum nonnihil extuberantibus, infident?
blonga, mollia, lenia? tenuia? ex uno latere longius
xporrefba > lato finu ad petiolum profunde incifa. (o/la
media, quatuor alias annulari duftu emittit₅ in adverfa
^arteeminentes. Floret nulli. FruBus plurescongrega*
<im proveniunt ? triangulares , angulis late extenfis ac
lalde extenuatis₃ in unoquoque duo reperiuntur fernina
h fe invicem decumbentia, fine ullo interlepto, rotunda,
?na,lataac tenui rotunda pellicilla circumcindta, pri-
'm viridia deinde ad fpadiceum vergentia. Prater hoice
?tiam fruftus habet fpurios , caulinis fupra exortum
foliorum infidentes, utin -KJtu Iselengu, fpadiceos? cortice
fcabro, carnedenfa? circa corticem viridi, intus albican-
:e: humor nonnihil vifcidus. Nafcitur in denfis ncmo-
ribus circa Candanata, &c Oedampere & alibi* Fru&usmen-
tC Septembri maturefcunt, femper virens. Radicis de-
:o<5lum cum pulvere radicum China ulcera mundificat
3c fanat.*



काली
खातो खात

• *Arcocci*

• *Riccia*

Riccia
Ricciaceae

Riccia
Ricciaceae

T A B . X L I I.



Rutex *Tu'Falli* > à. Bramantiibus ?'allay; & Lufitanis *Salique da Serre*; à Belgis *Slagblom* vocatur* *I^ami* corticecinereo, lignoalbi* cante. *Folia* in petiolis brevioribus folitaria proveniunt, oblongo rotunda₅ anterius cu-
spidata, texturâcraflâ ac folida? superficie plana & glabra, viroris in re&a parte viridis & nitentis, in adverâ lûrdi; odoris nullius > faporis valde acidi. *Co/la* media, inadverfaparte extuberans, obfcuratradtuvanas emittit? folium intus fubeentes. *Floret* in communibus ac brevioribus petiolis plures hincinde congregati videntur₅ iimtque gemmae parvae, quibus apertis > tetrapetali funtj obtufeculpidati> nullius odoris. *Stamina* iniisqua-
tuor cum fiis apicibus* *FruUus* baccp parvae rotundas, coloris fubruiH, fpiculotenui & nigricantein vertice in-
ftruflxc, carne intus fucculentaac dulci; inmedio *ofiicu** km eft yalde durum & albicansj inillounum? duo&(à>
pius tria femina inveniuntur₅ dura, glabra* coloris ex ruffo-fufci acnitentis. Natalefolumeft *Catour*, *Tetrawide* Scaliasloca; floret menfe Januario. Nullius in medicina; ufu.

T U - ? A L - V A L L I


T A B . X L I I I .



Upal-Vel: tamannice *T' allay*; Lufitani-
ce *Fula daz5|4ef*; Belgicè *Kjoeuy-nortel %a**
mi ipadiceo mufcofi, ligno duro, cortice
ladeo. *Folia* inpedunculisbrevioribus, fpa-
^iccis, gemina, decuffato ordine prove-
niunt > oblongo rotunda₅ anterius cupidata,
viroris in recla parte faturi, adversà clari. *F/ores* racemati,
in petiolis exfpadiceorubefcentibus, tenuibus, rotundis,
qui supra exortum foliorum emicant, vifuntur, penta-
petali? parvi, candidi, interius lanuginofis pilis obfiti,
in medio expundtati • > circa orificium iibvirefcentes. *Calix*
arcT:us₅ viridis, circa bafim pmordii — fructus conti-
nens. *FruBut* globuli lanuginofi, cinti[^]i, exterius fpi-
culis xutis, quibus vftibus adherent, obfiti; (*Lappas*
noftratumeflēdiceres) *femen* in fe continent unum, figu-
rāovalij glabrum, durum, nigrum, admodum nitens.
Globuli aperti odoris funt fuavis & mellei. Nafcitur poft
Calao/an & aliis in locis, pra'iertim arenofis & petrofis.
Floret temporepluvii, femper virens. Vires adhucdum
incognitas.

T A B . X L I I I

Plants: hujus fem.cn magnam habet flmilitudinem cum *Akxa Carp'tm Folia, femiuerojirato.*



A C A T S J A > F A L L I

œu

M O D I L A T A L I.

T A B . X L I V ,



Qatsja-Falli; Bramannes dicunt *Medicata Talii* Lufitani *%amos dajevi;* Nofrates *^AieervkchmorteL* (*auliculi rotundi?* tortuofi, valde tenues? lend? fuper plantas lato tradtu excurrentes, ramofque quos arripiunt ita adherentcs & obfidcetes ut vifum ac acceflum impedian. *Folia* in tenerioribus viridibus caulinis rara ac fparfa, valde parva 5 ex viridi flavefcentia, plura fimul jun<5ta initident. *Flores* in viridibus 5 rotundis 5 nonnihil tortuofis ac ftriatis petiolis 5 congregati feu fpicati videntur? ad exortum brevibus foliolis fuccincl:i; hextapetali funt ?, albantes, quorum tres exteriores lati & cufpidati, interiores fuperne bicufpidati, odoris nuUius. In medio *flylur* craffior, oblongus, cufpidatus; (*lamina* fex quse flylum interiore minlecompletduntur. *Fruttm* baccas rotundas 5 primum virides 5 exin albantes 5 umbilico latiori, fex foliolis coronatis primò viridibus, poftnigricantibus, fub tunica crafla exteriore globulum continentes ; officulum in fē conti* nent rotundum, cortice tenuiffimo 5 viridi -> polt nigricante ?, intus nucleo albo repletum : Natale folum eft *Cochien* &c multa alia loca ; denfa amat nemora. Integra planta cum gingibere exficcato, & *^Qrvala-Tulk* tofta? pulverifata& butyromixta? inveterata ulcera munificat. Cum oleo *Sergalin* capiti illita capillorum radices firmat; cum cortice *Arekgra* & *Tsjangelam-Tarendl* tofta, pulveriata & *Opio* mixta capitis ulceribus medetur; cum Cardamomo₃ lacT:e&oleo *Sergelim* ad liquidam confiften-


Y

tiam

tiatn redacta, capitis asftum tollit. Succus cum faccharo oculorum ardorem & fuffFufionem amovet; per fe loco balnei adhibitus hemicraniam mitigat.

T A B . X L I V.

Species haec eft Cufcutas, crefcens fupra arbores 6c plantas.



Kerria-Japonica L.

Bryonia n.


Zizania Ziz.

مُحْرِيَّةٌ جَمِيعَ الْأَلْفَيْنِ.



K<*iARETA-TSJORhV[']ALL*

T A B . X L V .



*dreta-Tsjori-Fdlti; lingua Bramannum 2)o<-terberi-Valli ; Lufitanorum tW & 2?o? ; Belgarum WildTimperlingen. Cauiculi exunia parte ex fufco rubefcunt. Folia numero tres, in petiolis ex adverfo clavicularum iitis proveniunt, oblongo rotunda? cufpi-data, medium majus eft, lateralia minora, texturâ craflâ, denfa, viroris in redta parte fufci & niteptis. Tsjori-Valli folia effe diceres, maximè enim cum illis conveniunt? fed oblongiora funt, necferrata, aft in oris apicibus, quse coilularum transverfarum funt producliones, eminentia Floret racemati in pedunculis yiridibus & pa-rum nitentibus proveniunt; tetrapetali? interius cochleato-cavi , ex viridi albefcentes, exterius rubefcentes. Stamina habent quatuor, flavis apicibus nodulata: inme-dio jlyhs viridis, craffior acdepreffusfefe erigit. FruBm uvse rotunda, in medio vertex eft, qui maturis frudtibus decidit; in principio virides, poft rubefcentes, maturi nicmcantes, carne intus viridi fucculenta, lancinante, qux oisculum grandius rotundum ex rubro nigricans comple-clitur, cujus nucleus candidus. Sylvas Trigolin, Catati & alia loca pro natali folo habentur, femper frugens vel flo*rens. Folia (iccata, pulverifata, zingibere ficco, vetufto pipere & faccharo commixta afthmaticis profunt.*

T A B . X L V .

*Summojureplantahsjc HedeuTtifoliaBaccifera I omnes notx charaderifticac convenientia.
MaUbarntfs dici p o f c mirum enira quantum!*

سُلَيْمَان
الْجَنِيدِي
جَنِيد

M 0 7) IR A-V A L L L

TAB. X L V I.



Odira-Falli; apud Bramannes Caro-Crodi; Lu-
fitanos ^tBarba-Castilhano; Belgas fyievel{fim.
[^]*ami-, teneriores virides> cortice ipadiceo*
fufcefcente, ligno albicante. Folia in fur-
culis transverfis? fpinis lignofis muniti? plura
in arctis intervallis congregata proveniunt,
longa> cufpideanguftoeminentia> textura crafla ₅ Ipilla,
ac folida > fuperficie plana & glabra ₃ viroris in refta partc
nitentis, in adverfa dilutions; Co/la media in adverfa par-
te eminet₅& plures tenuiores transverfim emittit; tota hxc
plantaadfumma magnitudinem excrefcit,. arboribus e
*affigens atque illas circumvolvens. Nafcitur circa Antigen**
*fr*a, Recare & alia loca > arenofa amans; iemper viret, fed*
•/lores nee frvSus habet, ut& nullum in medicina aufum.

Z

VALL



VALLI-dMO'DAGCAM.

T A B . XLVII.



5. **utex**^U J^Malab^aenf. bus *Vdlli-Modágam-*
finmanDibus Carotba-Tali; Lufitanis Folbl-
daTaoi Belgis Qorblad dicitur. %ami viri-
des, ff» g^ei", cu^J at¹, lignoalbicante, medulla¹
rulta. /b/M mgemculisgeminatainpedun.
culis craffis fed brevioribus vifuntur; ampla,
oblonea, texturf admodum craffa, carnofa, coriacea,
fuperhce plana, glabra; viroris in redla parte nitentis^{^i}
*adverfa furdi. *Cofla* media in adverfa parte alte extubmr,*
mdorfum acutumdefinens, tenuiter etiamin re&i emi
nens, exquanerviannulares, obfcurotra<3u excurrunt'
*Fkrer ac fmBm habet nulos. Nafcitur circa <*ParaUo**
^Mangam, &inaluslocis, femper virens. Cortex radi
cis cum cortice limonum in pulverem redaclus, ulceri"
bus, qu^MalabanbusTsjeu-Barapenvocantur, cum/uc-
ceffu perlinitur.

TA B . XLVI. & XLVII.

Folia harum hcederarum, optimè cum ilia Bela-Moda
gam conveniunt.



النهرة المبردة

النهرة المبردة

النهرة المبردة

النهرة المبردة

CM E R I 'AM-? U L L I.

T A B . XLVIII



Eriam-Tulli; a Bramannibus \$tafil; a Lufitanis Folbal-kredilha; à Belgis Smindelblad vocatur; frutex eft repens fupra terram. *Caukt* habet virides, glabros geniculatos intus aqueos & fragiles. *Folia* distincStis intervallis, in petiolis oblongis proveniunt, claviculis hinc inde comitata, oblonga, cufpidata, texturâ fpifla, superficie glabra & lenia, viroris vividi & nitentis, circa ora rubefcentia & apicibus cufpidatismunita; faporacidus eft & linguam vellicans, odor fubgravis. *Cofiam* medium albican tem in adveiià parte extuberantem, du^ lateraJes concomitantur, multis raris venulis interfperfe. *Floret* umbellatiin petiolis, quiad exoitum foliorum è geniculis erumpunt, vifuntur, tetra velpentapetali, ex viridi albicantes. In medio *capitulum* habent planum & rotundum cum suo ftylo rofaceo. *Stamina* quatuor funt vel quinque, prouero petalorum, albicantia, flavisapicibus inftrucSta. *Florum* (/èmmte oblongas, ex rubro-purpure^e & viridefcentes. *FruUus* baccas rotundas faciatim pendentes, glabras? ex viridi rubras; caro mollis, valde fucculenta, ex viridi fulca, faporis fubacidi, odoris gravis, ut folia t5\andaru. OjfEculum quod continet, valde durum eft, nonnihil rotundum, ex viridi-fufcum; nucleus albus. Nafcitur circa *Warapoli Sc* in aliis locis > florens & frugens ad finem anni. Ex foliis balneum antiparalicum conficitur. Eorundem & J^el'tca fuccus ophtalmiam tollit*

Rubus *leucodermis* Schlecht.
Morinda-purpurea Schlecht.
Rubus *leucodermis* Schlecht.

• Part 24

T A T> tA-F A L L I;

ceu

TAT) A-K, E L E ^ £ V.

T A B . X L I X.



Rutex *Tdda-Fdlli*; aut *Tdda-Kelengu*; qui lingua Bramannum *Tdda-VdUi^ & t^Muts-jegaTada*; Lufitanorum *%oaijLe(^amara*. dicitur; à Belgis *Stoetoortelvocatur*. *7(adix longa*, craffa₅ fibris terrce affixa; *cortex* fucus eft &amarus. *fyules* tenues₅ lignofi, virides? pilofi. *Folia* in pilofis & ad originem extuberantibus petiolis proveniunt, oblonga, triangularia, ad exortum latiora, in cufpidem unius duorumve pollicum extenfa, texturâ den fa & craffiufcula > fuperficie interiori fcabra & fuhaftpera₅ viroris communis₅ exteriori leni, admodum viiloia, viroris vividi. *Co/la* media plures alias emittit, in adverfa parte eminentes. *Floret* admodum minuti, albicans? racematis fupra ex origine foliorum producuntur. *Fruttus* in racemis, acini funt minores₅ primò oblongo-rotundi & parum compreffi₅ dein rotundiores ; *colonis* in principio viridi-diluti & minutis pilis obdudti, poftea candidi &c nitentis₅ faporis fubamari & parum pungentis; finguli acini officulum habent rotundum, planum₅ bilaterale, in unoquoque latere tribus ordinibus denticulorum minutorum obfitum; *ofiictda* primumflava, poftea fufca fine ullo nitorc *Pa/oerti*, *Cocbien* 6c alia loca, pro natali folo agnoscit, floret menfe Augufto, Septembri & Octobri; frudlus circa finem anni ad maturitatem perducuntur? femperviret. Radix cum la&e ebutyrate&femine Cumini fumta difenterias & hsemorroidibus infervit; cum aqua calida fatus difcutit. Succus foliorum expreffis rubedinem oculorum folvit₅ & capitidis dolores mitigat.

T A B . X L I X.

Species eft Bryonise Icvis Malabarienfis,

KAP.

N U R E M - K E I G . U .

T A B . X X X V ,

Rutex, qui Malabarenibus *Z\|ttrem-If{e/e?igu;*
Bramanriibus *Conmtti* & *Tito Carandi*; Lu-
fitanis (*frabofa Ovada* | *Belgis Ey- Key /en* dici-
tur. ^ 4 ^ ut *K^tH'^Qtren-K^elenguy* led major.
rW<r spinofi ac tenues. *Folia* ex petiolis₅
intus ftriatis quina proveniunt₃ minora
quam prascedentia? texturatenui, plana? glabra. *Co/la*
media plures tranverfales emittit₃ nervulisinftru&ai, ad-
verfam foliorum partem cancellatim pertexentes. *Flora*
ut *E^atm-^ekngu*, in vertice frudfcuum pentapetali j colons
^avis dein nigricantes? qui umbilicum frudtus confi-
:uunt; & *FruBus*, fed minores; praster illos & alii vifun-
:ur, plures fimul provenientes 5 virides, oblongi, trian-
gulati₃ in unoquoque angulo faba valde plana & com-
>reTa Eft & alia species *J\£uren Kfkngu* cum priori
ronveniens, praeter quod folia tantum trina prodeant,
& fruclius oblongo anguftiores fint. Nafcitur in terris
Jngecaimaei & ubique locorum, femper viret, floret₃ fru-
get & verrucas habet, inter edulia recenfetur; in medicina
nullius uetus.



Herbarium - Collingwood
No. 30983. 1883.
Canna
S. W. T.

قناة عجمان



KATTA-KELENGU'

œu

"BATATAS.

T A B . L.




Jppa-Kele'ngu ; Bramannicè Cananga ; LufitonichHatatas; Belgicè Tattater, Itydixmul-
turn fibrofa , magna , craflà , carne intus
carnofa, aquea, cortice fanguineo - rubro.
Cautes^ rotundi, virides,pilofi,afperi, intus
aquei, cum filamento lignofo albicante. Folia
in petiolis oblongis, rotundis, viñdibus, fulcatis, pilofis,
fubafpensvidentur, radiculis ad ortum ftipata, cufpidata,
circa bafim angulatim excifa, tenuia, mollia, lenia, elabra,
coloreviridifaturo in refta, in adverfa parte clarof
pore aqueo fubdulci. Cofla media duas laterales oblique
fpargit, e quibus alias exeunt, ad oras ardluatim inflex^,
in adverfa parte eminentes. Excolitur hsec planta in hor-
tis& agns, radix enim culinaria eftut Pafiinacas nofraturn,
grataacdulcis; inmedicinanullius uetus.


T A B . L.

Planta kceft Emotes Hifpanorum Clufii, It,ca dcferibicur.
 Bauh.rn. ccaPifone sub nomine Brafiliano Ge- |

Bb

TO-





July 1852
R. H. Dyer
कोलेज द्वारा दिया गया

PÓDAVAKELENGU.

T A B . L I . L 11



ódava *fyg* fecundum Bramannes %aro.
 di; Lufitanos *1{ai^da* <*Bot/a*; Belgice "Bcurs-
 wtrtel. <*Radix* ex pluribus craflis, ruffis, ob-
 longis, fpadiceo dilutis bulbis conftat vari-
 cofis, quibus minores adjuncli iunt, carne
 intus albicante dcnfa > Knguam acriter vel-
 licante. *Cauki* rotundi, tortuofi, nodulati, fpinis acu-
 tis, ngidfs, cufpidatis, rubefcentibus, transverfisobfiti,
 verfus interiora nonnihil reflexi, ex parte virides, ex parte
 rubefcentcs, caeruleo rore tin6ti. *Folia* in petiolis craffis,
 fpinofis, viri dilutis, ftriatis, circanodulos caulumex'
 tuberant, term, oblonga, anterius cuipide longiori emi-
 nentia, viridi fatura, pilofa lanugine veftita, ad tadtum
 parura hirfuta. *Floret* inpetiolis oblongis, craffioribus,
 lpinofis, viridi fufcis, fupra exortum foliorum erum-
 punt₅ heptavel ofto petali, ftellatim cufpidati, viridi
 diluti, flocculofa lanugine obfiti, fuperne oblongis & te-
 nibusfilamentis cufpidati, ac oblongis pilis barbati, o-
 doris expertes, ut & fru&uum. *Candanata*, *Warapoh* &
 alia loca pro natalifoloagnolcit, florens tempore pluvio-
 fo. Folia cum ^or/Wro in aqua decocta pr^fentaneum
 proalgida febrifuntremedium/

TAB. L I . til.


Pódaya-Kelen ell *Kafium* Brafiicaimm feu A- I dicic Pifo *lib.* 4. 1 ... 64 i pu(j Ljfi tanos sub
 mricianutnaltcrtn E.r.th. Fin. lib. j./iff. i. Cajif! nomine Jft *sama* venit.

% Sf

Ranunculus

odoratus

Pl. 22



PANtA MBTJ-VALL

TAB. LIU.

P | Andmbu-Valli\ aBramannibus Silona; aLufitanis %gtao Faljo; vocatur; a Belgis Wilde %otting. %adix nodoia, albicans, tenui corticula ruffa obdudh, qua? fibras longas tegit &craffaS) exalbofub flavas; OcuJiadlate-
 ra novos edunt caules? faporis Tub aquci, Ipfi
Qmh rotundi, glabricxviridifufci, parumiigno/i. *Folia* arundinacea? oblongo augufta, in cuipidicm obJongum valde crifpoiuin & intortum flrid:a ? venis in longum ftri-
 ata₅ viror fufcus eft & nitens, fulcior cum vetuftiora fiint; *CMMkni* foliis prorfus {imiJia? fcdhascparumfuf^ ciora. *Fiores* infatigiiscauliculorumracmatimvifuntur? primo albicantes₃ exin virefcentes₅ petali rotundiores funt, parum cupidati, circa lemutuoconvoluti: *Stylur* craffior, oblongus_s viridis> qui germcn frudtus eft, ex vertice tria oblonga alba folioJa ejiciens ; *Stamina* fa tc-
 nuiilima, Candida, oblongis, craffis, lubflavis apicibus dotata; odor gratus. *VruHut* in cacumine caulum race-
 matim pendent* funque bacc^ rotundx, Ipiculum & cum illo folia /ex in vertice gcrentcs, primumviridesj poſtrubicundi; carointus vix ulliusmomcnti, nonnihil iucculentaj faporisubfalfi; officulum unicum infecon-
 tinens₅grandius,rotundum>nucleoalbicanteinftruclum.
 Naſciturcirca *Qranganoor^Tarou* dcalialoca, plerumque vel florens vel frugens. Vires apud Medicos ignotie.

T A B . L 111.

Multe (pecks Kot*gg in India inveniuntur, Ipoteſl; Bn»b.V'm. afcendeps eiiim arundoefit.
 pro genere Aiundinis indict Voluhulis haberi I

Cc

PL



Diplo
Diplo
Diplo
Diplo
Diplo

P i R IT a

T A B . L I V.



Ruṭex *Thrift*; apud Bramannes *Ogoranim*;

Luiitanos ḥrva "Bendta; Relans " "

rotundi,

li.sveft.ti. *Folia* ex geniculis caulfum pro
veniunt, m pedunculis ab interiori _{PS}
plains, ph'ago rotunda funt ^{cusp.^{^L}} *A* ^{crenata} lenia, vi-
circa oram * [?] *lenia* roris in recta parte _{laturi}; odoris nullius, *A* lenia, vi-
gentis.

videntur

ad exortum foliorum

tapetali,

albicantibus instructi, quæ cæruleis apicibus donata. Odor

nuuus. florum \$?m»* oblong*, anterius ftrial ^{ores},

cand,d[^], nrtentes. *Fn&u* gemmutefuntfem ^S es, co-

nic[^], valdedur[^]e, coloriscinereofufci; du o ST illis semi-

na, parva, roṭunda, dura, nigra. Colitur circa *P* _a.

Jmacon Scmzlmloas; floret menfe Decembri, m u S

xftitmtus. Succuspro balneopr[^]paratur contr Therr T

t em.& .mpetigmem. Radix alexip Wmaca e i t, pr Sr'

timinaqua *Ory[^]* decocta,trita,&umbilicoilira.

TSfE-

ft



Tyria-papuanae
نیپالیہ کے
گلے اور پھولے
کے
نیپالیہ کے
گلے اور پھولے
کے

TSJERIA-PU-PAL-VALLI.

T A B . L V .



Sjeria-fu-fal-Vatti; Bramannes dicunt *Fanualla-VaUi*; Lufitani *'Fu/a Leiticai* nofrates *Spar-ling*. *Cau/er* corticem habenc teneriorem; fpadiceo-dilutum, mufcofiimac villofum. *Folia* binain nodulis ramorum erumpunt* oblongo rotunda, anterius cufpidata? tex-turâcradâ? fpifla, lolida, superficie lenia & glabra; viro-ris in refla parte nitentis, in adveria levioris > pilis fparfim obfita. *Floret* racemati in fiimmitate furculorum fpe6lan-tur> pentapetali? oblongi, cufpidati? ftelliformiter ex-panfi₅ exteriore parte candidi₅ interiori circa unam oram albicantes > circa aliam ex albo dilute rubefcentes , ladlei, odoris grati. *Stylus* in medio conicus₅ cufpidatus₅ quin-que lignofis oblongis apicibus confans. *FmBus* fertnul-los. Natalefolum *IVarapoll* eft₅ arenofaamansacfemper florens. *Folia* cum fandalo citrino & cumino in aqua ma-cerata&melli permixta, febresexarticulare morbo ortas? tollit.

D d

TA-

Dr. C. A. M. L.
B. ۱۸۷۴
Dendrocnide
تُنَرْ-جَنِيد
تُنَرْ-لَهْلَهْ



T A L U · D A M A.

T A B . L V I .



dlu-'Dama; Bramannicè Tunarnava; Lufita[^]
nice Folbas do Titao; Belgice Trat Wbrtel. Cauks
rotundi, rubefcentes, geniculati, nodofi.
Folia rotundiora, craffa, abunaparte viri-
dię, ab alia viridi clara; circa coftas plicis
«ata, circa oram rubefcentia. Flolculiūupra
exortum foJiorum exftant, cyathiformes, perfico rubri,
in oris leviter crenati, in orificio interior!, purpurei stilus
rubro purpureus, rubra gemtnula dotatus. Stāmina kidem
purpureo rubra flavis, gemmis nodulata. gemma feminales,
oblongorotunda, coftulis quinqueeminences ac pilofe-
fcmen in illis unicum, oblongum. &4mm nativitatis
locus eft, floret tempore pluviofo, arenofa amans & pe-
trofa. Decoclum hujus plants, febres ex endemio
morbortos, quemmidigena; Titdo vocant, tollit: %adkis
cortex cum cumino & fero laflis pro potu adhibitas al-
vum laxat.

MEN



لیلیوم
Lilium
لیلیوم
Lilium

<J|4 E N T> O N I

T A_#B. L V 11.

iJBndom; a Bramannibus *zJrty*; a Lufitams
tyK d'empoje; a Belgis *Goes/ken* vocatur.
%adix in duo finditur brachia, longitudinem
 spithamasadasquantia, nonnihil plana, unum
 tamenmajus altero, cuticulatenuiruffa ve-
 ftita; carodenfa, albicans, subflavaeſt. *Cauli*
 eradice exfurgit unus, hinc in detres quatuorve caulin-
 los similem mittens, virides, tenues, ipferigidus, minus
 durus, vindis, medulla aquea, filamentis lignofis pef-
 texta. *Folia* ad cymas caulinorum prodeunt, oblongo
 angulta, incufpidem acutissimum definentia, venis den-
 fis inftrucla; inſtar foliorum arundinis, ftriata, coſtula
 inadverfaparte extuberante inftrucla, viroris communis
 ac vividi, faporis adstringentis. *Floret* in petiolis propriis
 proveniunt, rotundis, longis, furreftis, viridibus, hebla-
 petah, oblongo angulti, cufpidati, vermiculatim in oris
 cnfpaci, inadverfampartem inflexi, textura denfi, ex-
 tcrius coſtula eminente infrudti; colon's funt ſpeciofi*
 primo in totum ferme dilute viridelcentes, fblo culpe
 rubore aliquo perfuib, dein viroris inferius dilutions,
 fu-
 penus obfcure rubefcentes, poft inferius flavefcntes, fu-
 penus colore miniato venuſto tindi, quo demum etiati
 in fenori partefuffunduntur. *Stylus* in medio viridi-fufcus^
 oblongus, craftus, furrebus, trilateralis; in cōrtice *fta-*
men eft unicūm oblongum, trilaterale, in finguſis lateri-
 bus fulco ftriatum, fuperne in tres cufpides divifum, circa
 bafin lex ftaminibus longioribus comitatūm, quas primum
 viridi diluta, ex in flavefcntia, poll rubefcentia, perinde
 ac infloribus vidimus, in cufpide lingulas habent trans-
 verfas, nonnihil nutantes, latas, planas, virides, duo-
 bus limbulis flavis circa oras obduda, flavo farinaceo rore
 confperli. *Florum gemma* conicea^ [ceu pyramidales; ob

— e

quin-

quinq[ue]coftulas eminences, quinq[ue] angulares effe dices, viroris fulci. Floribus deciduis remanetfylus, oblongas viridi fufcas conilituens capfulas ieminalcs, qua: ad digit[i] longitudinem cxcrefcunt, trilaterales₅ laxo cortice in tres oblongos loculos divifas. Seminibus maturis iē relaxant, interius futura ftriati? quse tandem etiam *pvx* fccitate folvitur. *Semiwa* dupl[ic]ilerie in hifce loculisiveniuntur₅ rotunda, primò albicantia, dein rubra&cnitentia. Nafcitur circa *Cochien* & in aliis locis; (ed propter floris venuftatem etiam in hortis excolitur₅ plerumque pene florens. Indigence ifti radicis parti? quae ad Boream vergit? magnam vim adfcribuntin fympathia, magia & aliis Mctaphyfics operationibus. Fruclus alioquin radici adnatus luxuriantem ulcerum carnem impedit; cum oleo *Sergelim* & hauftus & pudendis applicatus hujuspulvis, partum facit facilem.

TAB. L V 11.

Planta hacc ex Ceylon transmifla eft fub nomi- floruit. Doftiflimus D°: *Taulus Hermnns*, in eccl* ne *NiugbdUj* & anno prccidenti 1686, in horto berrimma Lcydenflum Acadmcnfia Botanices Nobilliflimi ac Clariflimi Domin Gafparis Fagel, Profcflbr, nominat *Liltuinfupnbm Ccylanicum*.

VEET.



Vitex - castus L.
विटेक्स कस्टस . लै.
Vitex agnus-castus
विटेक्स कस्टस
Linn.

F E ET L A-C A ITU.

T A B. LVIII.



Eetla-Caitu; apud Bramannes arlē fondo; Lufitanos <5\>darchato\ Belgas *Himpbla* dix fibrofa₅ tenuis > albicans. *Cauies* in nodulosfeugeniculosdivifi, rotundi, virides? rubelcentes? pilofi, venofi, ènoduliscau* liculos ejicientes. *Folia* quse cauies in nodis inveftiunt? oblongo rotunda > inferius lata, anterius in cuipidem definentia 5 viroris in re<5ta parte fufci, in adverla non item, in ea tamen cofta eminet -> nullius faporis. *Floret* fimul in cacumine caulium proveniunt, parvi₅ ex* rulei, hexta petali₅ quorum tres oblongo rotundi > interius concavi₅ ex petiolo transverfim exeuntes. *Stamina* habent fex? flava, quorum duo apicibus casruleis, reliqui fubflavis dotati iiint; ftylus in illorum medio emicat, virididilutus? globulo inftrucl:us. Poft flores (*/emma feminates* oblongas vifuntur , trilaterales & rotundas; in uno quoque latere in longum ftriatas, primò albicantes & nitentes , exin flavefcentes. *Semina* continent fex ? duo intra fingula? in longum jacentia, parva₃ oblonga? in principio viridia? turn Candida, tandem ex rubro fufca; faporis nullius. Natalefblumeft *Cochien* & alialoca, fem-pervirens&perterrā rependo cum flore& frud:ibus^ ulterius fe extendens. Medici ufum ignorant.

PON.



Datura L.
Datura L.
D. m. of S. I. B. M. H.
B. M. H.

T O N G O L A M

T A B . L I X.

Dongolam*) Bramannes dicunt *Sammana-*, Lu-
fitani *itgi^da t5A4oftarda*; Belgas [*ferswortel*.
Rami cinerei. *Folia* viridiafiint* oblongo
angulia, cufpidata? in oris acutis denticulis
crenata₅ rigida > plana, glabra, viroris fufci;
praster coftam medium nervulos habent
transverfoS) annulatim infeinvicem incurrentes; faporis
ejufdemut *Thlmpi* alterum, ceu *Kaphanut* major; belgse
dicunt *sUMyredil^* Floret nee fruget. Circa *CMangatti*
invenitur? & in paucis aliis locis, arenofa amans. Ufus
ujus plantae in medicina magnus eft j catefacit enim, ex-
ccat, difcutit omnia vitia ex frigore orta₅ ut & humores
uituitofosacfebres.



F f

IN-

I N D E X.

Prior numerus Tabulam, posterior paginam indicat.

A.

Cats jaValli, seu Mudi	laTaIi.	44-83
j -Amdlago.		16-31
I Ana-Parva.	<i>Pothos</i>	44-83

B.

R-Afelh.	<i>Loranthaceae</i>	
•"Bana-Valli, vel NatsjamCrua.		24-45
Beetla-Codi. /h^- ><,- -		1-1
Belutra-Tsiori-Valli	<i>Oii*/'</i>	if-2 9 10-19

C.

O Ari-Villandi.	<i>Smilax</i>	31-59
I Jr Cattu-i-Molago.		13-25
Cattu-Tirpal. <i>Piper longum</i>		14-27
Cit-Amerdu. <i>Menispermum</i>		21-29
Curio-Talv <i>Lauraceae</i>		26-49
Urnil, vel Curignil.		25-47

I.

SiRima-Taly.	<i>f'wai</i>	39-73
--------------	--------------	-------

1.

Tci-Canni.	<i>Loranthaceae</i>

K.

^"Appa-Kclengu, vel Batatas.		50-95
^Kareta-Tsiori-Valfi.		45-85
itsji-Kelengu	<i>Stenocarpia</i>	38-71
attu-Katsjil.	<i>/ Stenocarpia</i>	36-69
utu-Kelanrtti.	<i>Dioscorea</i>	37-71
tu-Kara-Walli.		17-33
tu-Nuren-Kelengu.	<i>Dioscorea</i>	34-63

M.

M Endoni.	<i>Globoseae</i>	57-107
Meriam-Pulli.		48^1

Mo lira-Valli.		46-87
Molago-Codi. <i>ftc/u* *%tis?~i+**i.</i>		12-2;

N.

TAtsjam-Crua, vel Batta-Valli, j	-1	
Nurem-Kelengu.	2^-^-^<-/t	35-67

P.

PAda-Valli vel Pada Kelengu.		49-93
Pada-Vara		27-51
Panambu-Valli.		53-99
Pee-Amerdu.	<i>Menispernum</i>	19. 20-37
Pee-Ula.		23-43
Piripu		54-101
Podava-Kelengu.		55-97
PongoJam.		50-105
Pu-Pal-Valli.		43-81
Pu-Valli.		42-79

S.

CCembra-Valli.		6-II
k^Sciieru-Vajji-Caniram, reJ		vitis in dir.
Sjerou-Vaili-Kansjeram.		4-7
Schem-Karu-Valli, Caniram, vel,		<i>Cassiera</i>
Tsjeni-Karu-Valli-Kansjeram.		S'9
Scimjambu-Valli.		11-21

T.

X'Alu-Dama.		56-105
*• Tjageri-Nurem.		33-63
Tjori-Valli.		9-17
Tiri-Itti-Canni.		30-57
Tsjanglam-Paren da.		4i-77
Tsjera-Pu-Pal-Valli.		55-103
Tsjereu-Caniram, vel		
Tsjerou-Cansjeram.		

U

^ Unjala.

22-41	<i>Goeatum scandens</i>
18-53	

V.

VAlli-Caniram.		3-5
Valli-Kara.		18-35
Valli-Modagam.		47-89
Vallia-Pira-Pitica.		7-1
Vallia-Tsiori-Valli.		8-15
Veetla-Caitu.		58-109

W.

Attou-Valli.

32-61

INDIAN BOTANIC GARDEN LIBRARY
BOTANICAL SURVEY OF INDIA