

Indian Botanic Garden Library
BOTANICAL SURVEY OF INDIA

CLASS No...£. 51.9/—**!
BOOK No....RHE.....
ACC. NO..... 2.5403



Hausk Empoyy Boluccii racem
H O R 1* | ^k^£-
M A L A B A R I C I
P A R S N O N A,

D E

H E R B I S
ET DIVERSIS ILLARUM
S P E C I E B U . S

Latinis, Malabarkisy Arabicis, Brachmannum char after thus nominibusque exprejjs,

Adjedta Florum, Fruftuum, Seminumque native magnitudinis vera delineatione,
colorum viriumque accurata defcripcione,

A D O R N A T A

Ter Nobilissimum ac Generosissimum T>. 1).

HENRICUM VAN RHEDETOT DRAKESTEIN,

Toparcham in *Mydrecht*, quondam Malabarici Regni Gubernatorem supremi Confeflus apud
Indos Belgas Senatorem Extraordinarium, nunc vero Equeftris Ordinis nomine Illuftribus
ac Prasotentibus Provincie Ultrajedinse Proccribus adfcriptum,

In ordinem redegit 6? latimtate donavlt

A B R A H A M U S A P O O T M . D -


Nott's adauxit) & Commentariis illuftravit

J O A N N E S C O M M E L I N U S .



A M S T E L J E D A M I .

Sumptibus Viduce JOANNIS van SOMEREN, Hxredum JQ ANN IS vad
DYCK, HENRICI & Vidux THEODORI BOOM,
Anno CID IOC'LXXXIX.



Gehlerofffimo Viro

D. GODARDO WILHELMO
V A N T U Y L L ,
VAN SEROSKERCKEN,

**Domino yan Wei I ant, Solchgrckgn & *c. Olim Ordinibus primi loci dioscefeos TrajeBina adfcrypto, Caponico Feteris <^Klona* flerii > TrafeBo aggeris ad Leccamjuperiorif.*



N Tibi munufculum, VirGene-rofiflime, quod abfens Tibimitto prfefentififfimum mei in Te ftudii & amorispignus. Namet filongif-fime disjun&us a Te in extremis Indorum ter-
ris, & alio, quod ajunt, fole calentibus agam
hoc tempore, femper tamen mihi obverfatur
jucundiffima Tui imago. Ea tam a!te infixa
eft animi mei medullis, ut nulla temporise ulla
locorum longinquitas earn poffit obfcurare.
Temihi conjunxit non tantum fanguinisne-
cefliudo, fed & mirifica morum fuavitas, fin-
gula quadam gravitate temperata; nee non
omnium egregiarum artium, viro Nobili di-
gnarum, fcicntiapneclar. Horas enim Tu
vacuas non rebus inanibus, fed rerum divina-
rurn & humanarum perveftjgationi impendis:
de quibus apud amicos differis fubtilius, quam
ii, qui omnem setatem in illis uniee confum-

2

fe-



ferunt. Cum igitur tanto pulcherrimarum artium studio tencaris, nullus dubito quin iftud Horti Malabarici in quo colendo tantam olim operam pofui, volumen, quod tot ignotas orbi noftro arbores, frutices, herbas detegit, Tibi non ingratum fit futurum, quique id eo iis excepturus animo, quo Tibi à tan tis terrarum intervallis remoto cognato, fed qui ubique te femp̄er in oculis geftat, offertur, hoc eft amoris & benevolentias pleno, Velim enim tibi perfuadeas

ihCobilidimi TtiiNominis

Studio fiflimum cflē

H. VAN RHEDE TOT DRAKESTEIN.



Geestes
1655. F. M. V.
H. A. RHEEDE
HENRICUS ADRIANUS VAN RHEEDE

Tot Denck van den Nederlantse
Landschappen en Gezichten. Enige
Belangen der Landen. Dierende
naren, en Cadeaux van den Oosten. Historie
D'Uytvalck van de vryheidt des Tempore
Leyden en Leiden. Beladen met vaste Landen.



Celebemmo > Slngukri, &- undequadque Eruditifimo

V I %, 0,

Don> ABRAHAMO VAN FOOT

Med. To6i. Eximio,

HORTUM MALABARICUM

Bono Tublico edenti, Tomino & Evergette suo
aternum Fenerando.



NGENIUM MENTISque JUBAR penetrantius omni
FULGUME, cuidatafunt, ARDUA ferrecupit:
Divina hinc POOTf Sapientia novit acuto
Vifu, quâ frendens valle moretur A P E Rj
Ingenio cuius MEDICUS non sufficit ORBIS 5
Sicut ALEXANDRO non fatis ORBIS erat:
Difflando Invidias fumum sublimior exftat
INGENII & CALAMI dexteritate SUI.

Mira videre mihi videor quotiens TUA VIRTUS:
CURA, LABOR, MERITUM, DEXTERiTAS que monenr.
Arvatenes, VIR MAGNE OEOT-facrata- AOTIAS :
Nee fugit amplexus HISTORIA Alma TUOS:
Nuper enim EXITII HUNGARICI quos vidimus Annos
Infaustos, fauftx dispofuere Manus.
Teftis fum majorum O P E R U M: Victoria parta eft.
Police de multis, quas TUA Theca tenet.
Ardua PARNASSI juga tendis adire bicornis :
Quid vetat eximum munus amoris ? Iter ?
Credo, quod Herculeus LABOR, & rnoeftia Vitae
Non levis exiftat, durâque CURA TIBI:
Scd perfa ! Quoniam LABOR improbus omnia vincit /
Per MARE, per TERRAS, omnibus OMNIS, adeft t
Ac tantos noſtris propter conatibus addit
Succeflus, quantos ſcribere penna nequit j
Euge ! LABORANTfUMquiaNOX, FENIT: Haec TUA SPARTA eft,
Pafce Gregem CHRISTI qualibet ARTE potes.
Nos etiam STUDIO non femper fungimur uno j
Officia ^tati funt data cuique fua.
Difcolor eft MUNDI facies, diverfaque RERUM
NATURA : Ipfe vices quattuor ANNUS habet.
BRUMAM VER fequitur, VER excipit .ESTUS & JESTUM
AUTUMNUS, quern mox frigida claudit HIEMS.
Hunc fervare modum novit NATURA : Quid obftat
Affiduè partes fie variare T U A S ?
Hinc TIBI LAUS & HONOS nuljo delebilis Mvo,
Inque CHORO, inque FORO furget adufque POLUM.
Non puto majus ONU S 3 fed nee puto majus HONORIS
PONDUS in hac vita poſle manere VIRUM,
Publica quâ VIRTUS fi quern manifeſtat , ut ARTE

Sana

Sanā , aliis grātā, pandere vela queat.
 Nee dubitare juvat: SACRORUM CURA ferax eft :
 Per FRETA, per FLO RES & THYMA fumma volat
 Inftar APIS, redolētque THYMO fragrantia mella,
 HOC E ST , Virtutum Proemia corde fovet:
 I R R A D i AT T E R R A S 5 neque enim latitare valens eft j
 Qac MARE, jungit OPES, DETERIO RA fugit:
 Nifibus occukis & furgit ut Arbor in auras :
 'Sic neque SACRA filenc 5 nee latitare finunr,
 Non nōrunt AQJJIL^E imbelles generare COLUMBAS*
 SACRA nee immundas progenerare MANUS:
 Hinc PQOTI CURA & MERITUM, LABOR improbaVirtus*
 Permittos FLÓRES audet adire foràs.
 Die ubi praccipuè ? MALABARI fuftinet HORTUS
 NiTnc, nimium faciies POOTII utrafque MANUS.
 Pcrvigiles jbi quid per'agunt ? peteretur ab HERO
 Praecellens venia ? Nobilis HORTUS enim eft?
 Sifte gradum ! Quoniam cum quo rem gerit adftat
 Ipfe DEUS, largas qui tulit Almus opes :
 Hinc eft, quòd glomerare Rofas, Lapathum atque Melanthium:
 S E MIN A mox fecum fingula ferre, queat.
 Adrachnem, Daicum, Amnum, Apiafrum, Urticam, Alicafrum,
 Diverfi generis S E MI N A fie liabeat :
 S E MIN A qu# mixtim confufa videbat acervo
 Seligat, & Nomen cuique fuum tribuat.
 Difficilis LABOR eft fateor cui fubjicit UNUS
 Ipfe CAPUT j quoniam Pondus Onúsque tenet:
 Quo fuper innumerī MEDICI fudando periti
 Multa dabant fcriptis , SOLUS adimplet. Atat!
 DignumLaude VIRUM PIETAS, SOLERTIA, CVKJE
 Ingeminant: Laudes hoc OPUS ARTE dabit.
 Nolo OPUS hoc multis laudari: Non OPERANTEM:
 Aft OPUS infpicito. Vis quod ? Utrimque dabit.
 Perge bonis Avibus, DOCTOR CELEBERRIME POOTI
 Sic TIBI CURA SOLI, fit TIBI CURA POLL
 Interea vigeas PIETATE, recepta LABORUM
 BLANDITIES, DOCTOR Purus & Harmonicus.
 Sedquianon FELIX CONATUS ET UTILIS UNQUAM-
 CONSILIU M SI NON DETQUE JUVETQJUE DEUS*:
 Tu DEUS, Omnipotens GENITOR, Mow & T^ unà,
 ARBITER Sterne, fis fine fine prope .
 Profper adefto Tuas LAUDES, Tua DONA canenti,
 Extollàsque inter nubila denfa CA'PUT!
 VivnK longxvos DOCTOREM Neftoris Annos
 Fac Supreme PATER 1 Nam Tua Téque cupit .
]>UC ME, NEC SINE ME PER ME DEUS OPTIME DUCI
 Ingemiiiat: Quoniam SE DUCE pulvis erit.
 O Dcus exaudito VIRUM! Servato LABORES
 Ingenuo amplexu, quos meruère dies !
 O Deus hunc tueare VIRUM! Tua voce canentem
 Sufcipias , foveas nettare AMORIS! AMEN!

*L. M. fU Levidenfem hunc Carminum Appamum
 AMSTELODAMI fudit*

NICOLAUS APATI P. D.

E N C O <£|4 I A S T I C O N.

Tro Nobili iflhac circa

HORTUM MALABARICUM

R U S T I C A T I O N E ,

Viri Eruditiffirni juxta ac Laborioffifimi

DN. ABRAHAMI VAN POOT

Med. Dodloris Celeberrimi, celebranda.



*U quantum mortalepoteft comprehendere peēl
Ingenii quantum latebra ! quam maxima prwjas,
Plurimdq[ue], in v[er]a rias divifa induftria cur as !
Qu<e non incept ant> incept dque fine coronant*
Queis largos infudit opes natura , quibusq[ue]
Dulcia virtutum stimulos dant prcemia ; nulla
*Po(l cineres avo moritura gloria famx ?
Vnusadtnifrarerit SCRIPT OR Celeberrimfts, HORTVM
Qtti tantd aMALABARICVM alitque cohtque labored
Vt toti ille fuas mundo communicet Her has.*

*Jam multos laffavit onus, cui fiibjkit unus
Ipfe humeros collumque novijfimus : unde, refumpto
*Progreffu latatur opus, gaudētque Tatrono
Foelici, cujus manibus labor ardet in ipfis ;
Strufitura tarn magnified , cum tempore , Utd
Idem ornaturus fublime coronide culmen*
Tefuper.atqueTuvs, VIR MAGNE, laboribus, eft quod
t^Attonitus flupeat) qnifquis videt: HERCVLIS ifte
Eft ^Animus , mult is natique, labor ibus artus i
Mir a cui in partes Induftria (tifficit omnes.
Undique (iipatus cur is, & rebus agendts,
t^Accerfts Tu Jponte novai: ut pondere^Uta
*Palma viret, vivis cumulando labor? labor em.
Tu quutefolus agis junffim, vix fingula plures
ComrHode agant: quant ddigna hac fblertia laude?
^Arva colis Mtdica fane fee licit er Art is \$
Sedne fbla Tuo fanatis ere Teat ab agris
Gloria corporibus decori, non ultima circa
Infirmos Tibi cur a Animos ; ruditate refettd
Vi/ofies ratio, & mens fana in cor pore fano,
Et vegetans habit etprgclard fcientia vitam.
Hinc eft, quod Libris, noBefque diefque labor ans,
Scribendis aut edendvs , incumbere gaudes.
Inter quos primam prabets tibi fcriptio palrnam
Hiftoria : rebus quā non dimtaxat inheres,
(Vt multi faciunt:) prafentibus atq[ue] propinquis ;
Ad longinqua oculos minis /oca : 'PANNONIS' eras
H*ec inter luftras, illuftratrtus easdem**

% *

Tra-

*Trater eos \ quos lj>fe Libros componis & edis \
Mult a Tibi lucent debent aliena, vetuftas
Qua fortaffe opera , aut alii de Bibliothecis
Abftulerant cafus s Tua nutricatio rurfs
Curaque refituit: vel tranjlatoria mundo
*Publicat ilia mantis, diffundens latiñs uſutn.
għidrefertpluraaddere? ES IN^CDEFESSVS AGENT>0:
*tanca quies oculis, Animo vix ulla ; retraffas
*Pcrvigites oculos, pfrque ipfa infbmnia cur as,
^ueis quoquāque modd Respublica facra juvari
*Fofii, ^ his meritis Tibi debit a gloria quart.
Euge ! lucrum capt as vel tniquo judice pulcrum :
Cut confer re nefas argenti aurique faburras >
^Decant at a Tagi, multorum, munera , scriptis ;
Nee, quas dives opum mittitprocul India, gemmas.
Incept urn virtutis iter , quod ducit ad aftra >
"Profequere: Invidia nilguttura rauca morare ;
Virtuti comes Invidia eft: Trafentiasfama
*Demit j at ab/entes ultrb comitatur honores.
Jam mult is toti scriptis fat cognitus orbi,
fsfois ttbique : magis porrb, pojiquām hora quiet em
Clef* Tibifera fōnet!) fejjis dabil ultima membris,
Vithirus 9 pretiōque operarum perfrūtiturus.
Hungara (fl quid id e(l:) gratanter Natiojēmper
Agnoſcet, quicquid fūmpitura eft muneris à Te,
Nee pietatis erit tanta immemor : Acdpe fignay
Qua Tibi vota die at , fmcerō fota favore,
Ejjfujks tenui Vefla inpraconia Mufd,*

NICOLAUS KIS M. TOTFALUSI Hungary

Artium litcur'wum Mtgħifler



G. C. H.
Botanist
Drawing
1908

HORTI
MALABARICI
PARS NONA
DE
HERBIS
ET
DIVERSIS ILLARUM SPECIEBUS.

TSJOFANNA-zALERI

VEL

FVLJ & IESTICA INCARNATA.

T A B U L A I,



Relij Braman̄nes dicunt *Caneru*, duas fūnt species; *Sjovdnna-Areli*, & *TSelutta*,* *aAreli*. Priorplanta humilis eft, *Jtipite* medioredto affurgente, quiramos iirfum verfus obliquè emittit? craffties ferme brachium ad^quat, colorealbo, corculo item albicante , dein cinereo; *Rami* fuperioresgeniculatifunt? cortice viridi, primiim triangulati, poftea magis magisque ad rotunditatem vergentesj angulisevanefcentibus. *Folia* in geniculis terna furredta prodeunt 5 fingula ex angulis lateralibus , pendunculis parvis> craffiolis* viridi albicantibus 5 interius planis, ac femutuo eo ordine decuflantia, ut in eadēni ierie ceu linea fibi mutuo rēpondeant; oblongo angufta, cuſpidata₅ in inferiori parte ftričl:iora, longitudine unius circiter fpitham^) pollicis Jatitudine, texturā craflā^

A

fpiffa

S H O R T U S

fpiſla > folida \ flexilia 3 superficie plana? glabra, in reſta parte viridi fufca, nitentia 3 in adverfa clara. *Cofta* media rotunda eft, crdla ac Candida? in adverfa parte eminens. Foliorum fapor amarus, nullius odoris. *Flores* in fummitate furculorum in petiolis, qui terni fimul ex angulis lateralibus jerumpunt, ac ad exortum parvo cuſpidato viridi foliolo, quod dein decidit, fuſcunguntur, plures congregati, feptem, o&o videntur; funtque rofas plenas, foliolis rotundiolis₅ anteriusleviter cuſpidatis» incarnato rofaceis > tribus ordinibus deſtinftis₅ quorum finguli quinque conſtant pqtalis; exterioris ordinis intenfiusrubefcunt; medii latiora funt ac rofaceo dilutiora > exterius magis albicantia & nitentia; intimivalde crifpata? inme dio interius tenui zonula albicante obdudta₅ ex qua floſculos ejiciunt candidos₅ collum in collo foliorum medii ordinis includitur. *Stamina* in medio numerantur decem flocculofa? inferius plana? glabra-> Candida? fu perne pilis barbata & tortuosa. *Calix* ex quinque anguftis> viridibus? cuſpidatis foliolis conficitur. Odor fragrans ac am^eniffimus. (*Yemrnx* oblongas funt₃ foliis fupra fe invicem convolutis, ac nibro fufco nitentibus. In floribus prasterea animadvertisendum, quod primò magis rubefcant; aedeinde magis ad incarnatum vergant; item quod rarius dent *fruBum*, ac non nifi in vetuftioribus plantis; cuius *germen* turn in &ore, utfy/ur candidus, oblongus? lanuginofus appetet; funtque oblongiac angufti, fefquidigitales, lateribus duobus complanatis, virides? ex petiolo verticem verfuſ in cortice lyris denfis ftriati, quarum duas in medio laterum profundiores > ubi in duas fe relaxant capfas oblongas? quarum fingulae unica lata obducuntur futura, in qua fe p̄ae ficcitate aperiunt; cortex interior lanuginofus eft rigidus & albicans, verfuſ interiora fe infledens, ac ibi duos veluti loculos feminales efficiens. Floret hsec plantatoto anno; nullius in medicina uſus eft.

T A B. I.

Tsiovanna-Areli eft *Xerium Indicum latifolium*, flo- pertinet *Apocyni*, in omni enim parte notas il- vhus flems odoratis. *Hermannii Cat. Hort. Lugd. Batav.* lius Chara&erifticas oitendit.
Primum ab Auftore hue delata eft; ad species

Bellissima
نیکیتی

Cana - canna
کانہ - کانہ

لارج - لارج
لارج - لارج



B E L U TTA-zA R ELI
 # A U T
 F V L A-*cME STIC A* sAL<BA.

T A B U L A II.



priori planta hasc fblum differt, quod flore fit simplici pentapetalo, ac colore candido, dein ad incarna^im dilutissime vergente: interior florum tunica flaviucula, venis in longum radiata ex ora fuperne flocculos plures oblongos albantes ejiciens. *Stamnula* flocculofa funt oblonga, capillata, crifpatim convoluta; *fyium* in medio habent candidum, capitulo dotatum; gratiimogratissimodorisflores funt; fed nullum in medicina habent usum.

T A B. I I.

Bclutta-Areli confpecies est prascedentis, acad idem pertinet genus, secundum notas illarum Charaderifticas > & est *Nerium Indicum angustifolium*, floribus odoratis simplicibus. Hermann: Catal. Hort. lugd. Batav.) haec & omnes *Apcyni* species pro noxiis habentur j & quamvis arbustus has ligorem non tollent ladeum, ut reliqu *Afocyni* species, aft clarum & limpidum, nihilominus tamen prsecipua indicia cum illis *Apcyni* conuenientiis nempe pentapetalus filique, & femina, qua: langue luntobduita 3 & ulterius differuntur quia arbustulæ evadunt, cum *Apocynum* tantum fit herba, quæ inter *Nerium* five *Rhododendron* numerari debet.

B

NE-

B E L U T T A ~ z A R E L I

A U T

F V L A - e M E S T I C A * A L < B A .

T A B U L A II.



priori planta hasc fblum differ*; quod flore
it fimplicipentapetaloj ac colore candido,
dein ad incarna%im dilutiiEme vergente:
interior florum tunica flaviuicula , venis
in longum radiata > ex ora fuperne flofculos
plures oblongos albicantes ejiciens. *Stami-*
nula flocculofa funt? oblonga, capillata, crifpatim con-
voluta; *ftylum* in medio habent candidum? capitulo do-
tatum; grati imo gratiffimi odoris ilores funt; fed nullum
in medicina habent ufum.

T A B. I I.

Bcltitta-Areli confspecies eft pr^cedentis, acad
idem pertinet genus, fecundum notas illarum
Charaderifticas; & eft *Nerium Indicum angulii fo-*
lium, ftoribus odoratisfmplicibus. Hermann: Catal. Hort. *Lugd. Batav.* } haeut & omnes *Apocyni* species pro-
noxiis habentur j & quamvis arbufculae has li-
quorem non ftillent lacleum, ut reliqux *peponi*

species , aft clarum 8c limpидum , nihilominus
tamen priEcipua indicia cum illis *Apcym* conve-
niuntjfloſ nempc pentapetalus,filiuſe, & femina,
nuntur quia arbufculae evadunt, cum *Apocynum*
tantuni fit herba, quas inter *Nerium* five *Rhodo-*
dendron nuxnerari deber.

B

N E -



1850



Leptophyllum
Gardner

1860 - 1861

Botanical
Gardens

N E L E M < P A L A .

T A B U L A I I I I V ,



Elm Tala;^ Bramannice *Vtago Culo*, plants eft nafcens in arenofis, ad altitudinem trium ferme hominum ; *caudex* unius hominis crafftiem adasquat. *Flores* petiolis craffis, viridibus, pilofis, ad exortum foJiolis exceptis, in vertice furculorum, plures congregati proveniunt, pentapetali, oblongi, craffioli, cupidibus verfus unam partem inflexi > exteriora verfus crifpati, intusflavo rubefcentes, exterius viridefcentes ; *umbilicus* in medio eminet cavus, rotundus, rubefcens^ in ipfa ora foliofa lacinia annulari rubra, qu# ad interftitia foliorum cufpides furre&os protendit, velut collari furre&o circumdatus, quod fuperne fuscefcit. Ex hujus meditullio *Jylus* fefe erigitoblongus, conicus, lignofus, albicans, parum pilofus, theciformis, conftans ex quinque ftamunculiscolumnaribus unitis? intus globulum recondens viridem, rudimentum frudlus. *Calix* viridis ex guinquerotundiolisconftatfoliolis. *gemma* conioe funt, inquibusflorum petala fpiratim ad verticem convolvuntur, exterius viridi fufca fubcasrulefcientia ceu hyali coloris, interius viridi dilution, & Ia6te turgentia. Balneo ex foliis& cortice arthritic^e curantur febres; algidse vero piluh's ex fru&ubus formatis,* limimento ex radice & virente *Curcuma* in *Oriye* lotura & *fero laBis* contrita, calida depellitur arthritis ; ex *cortice* radicis levigato fit globulus ad magnitudinem Arek^, qui devoratus pr^efen'taneum in moriu ferpentis remedium eft.

T A B . 3 . 4 .

Quamvis Nclem Pala non fit herba,fed in arbo- rem evadat,tamen ē generc Apocyni excludi non poteft, omnia enim indicia id pro vero affirmanr,

atq[ue] idco ipfi nomen dedi <i>Apcynium Malebatkum</i> <i>Arborefientis</i> , <i>Limonii folio,flore exflayg ruManu</i> , fitf. <i>quis maximif.</i>

دیگر از این طرف
و چشمکه ای خواهد بود

نام: زلیخا

B. camelliaefl.



Blaue - L - ke L.



BELUTTA-KAKA-KODI.

T A B U L A V . VI.



Elutta-Ka^a pKiddia, n t a repens eft ; rami craffioli, rotundi, geniculati, ingeniculis nodis extuberantes , lignoalbo, cujus medulla crafla , viridufcula ; cortex viridis eft &flavefcens, valde lignofus , uterquevaldc Jaćteus. *Folia* geminata&ordinedecuffato videntur ingeniculfs ramorum, quorum petioli oblongi, rotundi, craffi? viridiflavefcentes&lanuginofi; funtque ampla, parum oblonga?anteriuscuipideanguftoeminentfa) texturā craffa, in adveria parte valde lanuginoia, leni/Gma) viridialbicantia? in re< Sapilis minutisiparfim obfita, ac ob conum hir/utum femi afpera -> viroris fufci ac nitentis. *Co/la-* media plures laterales emittit, qu^ omnes in adveria parte eminent. *Flores* iimul plures feexhibit in iürcu-Jorum cacumine, quorum pediculi? craflioli, rotundi, virides ac rubefcentes, ad exortum parvis culpatis rubef centibus foliolis ibccinfli; pentapetali lunt & candidissimi, petala ipfa oblonga₅ unguiculis ftrida, anterius latiora, ac rotunda ora, ad unam partem incurvata ? ubi dilutē ex croceoflavefcunt; illorum *collum* oblong₆*rotundum eft ac lacleturgens, ad calicem ftri&ius ac velut quinqueco-ftulis extuberans , exterius nitens ac ex collo flavefcens > interius flavalanugrueobfitum. *Calix* collum arde excipitj colon's fanguineo rubri > confitans ex quinque rotundē cupidatis foliis. Odor fragrantiflimus eft, ip/à Caryophilla, & cum flavefcunt flores, Melones maturos ipirans. *Stylus* illorum conicus & flavefcens, ex quinque rigidius culis, lignofls, cupidatis, albo flaviufculis componitur itaminuijs, interius bifurcatim divifis, includitque capitulum craffiuſculum, viride, ex meditulliocalycis emergens. (*femma* florū conyae, foliis Ipiratim convolutis confitantes , primò rubro fufco ^nguinei colon's; dein cum nitore albefcentes, & ante quam fcapient flavefcen-

tcs; quorum *calyx* rubro-fanguinco-fuicus. *FruBus* gcominati prodcunt» longi, rotundi, anguiti, in cuiuspidem definentes, ex una partc viridesjcxaltcraquafoliexpofiti rubefcentcs, fulco ftriati i cortice valSe laiifceo, conger iem numeroibrum in ie continentes *Jemhmm*, quae plana funt ruffa , ac ie mutuo fcuteLLatim obtcgentia, penicillo argenteorum inftrucia i tarn mum} quse /e tandem in or- bem difiundunt. Vires ignorantur.

T A B . 5. 6.


tc littii-Kikjt-Kedi inter confspecics 01D11 mm maxtm»e(t, atqueidcoamc vocatur *Apyimm S'.di-*
din *mjaxittium*, *Jalit tiush* , *ijtsiüido*, *fierietwtiic*, *fitiquii Uxpt*,



Calliolaria L.

வாங்கிரி மலை

காஷ்ட் புதையெடு



of 2) A-K O V I E N.

T A B U L A VII.



7)a-Kjodien planta eft ramis fc vinciens finuofis. %adix albicans fibrolā; intus n̄ lamento lignofb > nullius faporis. *Caules* rotundi virides, nodulati, intus parum lignofiaclaftei. *Folia*, bina&bina, ordine fe mutuo decuflantia? in rotundis₅ longis, tortuoſfis petiolis proveniunt, oblongo rotunda, inferius latiora> anterius rotundis oris in cupidem contraſta? molliaj la&ea₅ lenia,inadverfaparteplurimiscoftis, quas refl:amftriant> cancellatimobdufta, viroris clari, inre<5la partefufci, faporis lylyveſtris. *Flores*, qui nullum fpirant odorem, plures congregati ex origine foliorum in nodulis proveniuntj cyathiformes? pentapetali₅ cupidati, rigidi> furrefſti ? interius nonnihil cochleato cavi > viridi flaveicentes & albefcentes ? oblonga, purpureo rubra ftria in medio verlūsculpidem notati, exterius magis albicantes) Scab utraque parte velut rubro limbulo circumdati, exquorum meditullio *flylus* exfurgitcrafliſj viridi flavefcens albicans, qui ad verticem in quinque acutos ftringitur angulos₅ quinquecupidibusnigrisin vertice inftru6tus; *Calyx* planus, quinque foliis flores arcle fuccingit, in fundo rubris arcibusfriati. (*jemma* conioe funt; *FruBus-* oblongi, anterius cupidatij tres eminentes habent coſtulas > acinfecongiem oblongo rotundorum ac planorum continent femi- num₅ qux mutuo fe invicem obtegunt, & anterius ex ftriſto orificio *penicdlum* oblongorum argenteorum filamentorum emittunt, funque primò viridi flava," dein rubefcentia, in totum hi frudtus fimiles font frudtibus *Ericu*. %adix pulverifata & oculis infperfa nubeculasau- fertj vel cùm butyro coda & epota. *Xulbm* cum radice *Talma* agreftis& *Scelengu* tritus & cum pulvere *JdaKjdien* in panno confutuſjadjedomodicdji'W^/i, feribili nigras gc recenti butyro lento igne ad ceras confiftentiam reda- dtus,

io

H O R T U S

Etus 5 pro quovis oculorum vitio mane & vesperi fumitur: Idem cytrino *Sandalo* ac *faccharo* mixtus₃ in pulverem redadlusj & in pillulas formatus oculorum pathemata ex bile ianat: vel denique si cum foliis *Ficuum Inferit* > radice *Talu-Dama* primum in aqua, dein in vaccino *UUe* decoquatur & *faccharo* mifceatur₅ quovis oculorum morbo opitulatur: vires enim hujus plantas plane ophtalmicae iunt.

T A B . 7.

Ddincatio Scdefcriptio hujus phntir diffcrre rounct eft: unde tti;im liquetbanc plsntam illij Tidentur retpe&u foliorum, quit Viina Cvbimfe I adnunoeiaridebcrc; ego illam vocai «n *Apocynum* invjecmnon ticcuflant; ut omnibus *Apatjai* i'pc- I fitn&tiii, fictc vasirgMe, [Uiquii Lricu sim libus. cicbus, quod deferiptio per benecxpri miii com- 1




K A.

دیل-لری لی
جعفریه
کلینیک
دکتر



KARA K O 2) /.

T A B U L A V I I I



Aka-Kodi; vel fecundum Bramannes *Kiti*, planta eft naſcens in arenofis, fcandens & finuofisramisarboribus fc circumvinciens. "Radix fibrofa; y %/ pedenTcraffus , cum veſuſtionbus famis, corticem habet cinereum, qui in filamento deducitur, i⁷ ^ infurculis pnmō vñdibus., qui poſtca nigrefcunt, ficcxramorum nodis onuntur, ad exortum extuberant & geniculati funt, gemma femutuodecuffant, oblongo anguſta, adpediculum rotuñdiola , cum cuſpide brevi in vertice, t̄cxtura craſla, fpiffa, denfa, glabra, viroris in reda parte vividi, fuſaonsac valdcmtentis, faporis fubaſtringentis fubacidi & vinoli. *Cofta* media cum lateralibus utrinque eminent *Floret* in umbe Jlisproveniunt, vel ex cacumine furculo^ rum foliaceorum, & turn funt longiores, vel ex illortim latere, & turn breviores funt, funtque parva flavefcentes gemmuk, conftantes quinquefoliis oblongis an \triangle yufis ac flavefcen̄ibus interius pilis candidis Januginoſis* veſtitse, maxime in orificio ſuperioris colli, tenuius ac rarius in folns; illorum collum flavum eft & ſuperiori parte frictius; *calyx* ex quo emerget flavefcit. *Fructus* ex uno flore bini eduntur, oblongo rotundi, anterius friftiores , in primaparte ar&euniti & mutuo obverfi , cortice viridi, glabro & craflb, qui cum ficcus eft lignofus redditur, exterius nigricans, & rugis in longitudine frtiatus; *femina* verfuſ partem inferiorem vel craffiorem continent pi urima 5 quas in frith'ori parte vel cuſpide *penicillurn* emitunt oblongorum argenteorum filamentorum; funtque plana ? valde comprefJa & tenuia -> oblongo rotunda, īcutellatim dirpofita, primum viridia, dein albefcentia, tandem rubrofufca, oculo albicante inftruſta; flores illos Warn* *Caca-Codi* magnitudine ſuperant, ut & *penicillus*. *Folia* in fuppreffis lochiiſ uſui veniunt; eadem cum lacte *Nucis* In-

ia

H O R T U S

Indies contrita & pedlori illita, dolorofam cordis palpitationem j atquehinc promanantem prsefbeatonem fpafm unique corrigunt. "Radix cum teneris Jretjueir[^]Ta/mte filve frisac Toadeli radicibus in ebutyrato UBe ac maccte loth decofta> affedtasque parti illita > renum doloribus ac Ifcaticse; vel etiam cum Toncfo, Viittja aliisque radicibus in UBis firo, & oleo foliorm de Inferno codta gonagras, **vel** denique cum radice Talu-Dama, primo in aqua, dein in oleis Sergelim, 'Bepa-[^]deis-Elui ac poftea in ebutyrato L8e> Tamarindortim &- Muringu iiccis ad ceras confiftentiam cotta, in univerfum arthritidi medentur, quarum o6toginta lpecies incolas enumerant; quemcunque enim dolorem flatus corpori affiant) arthritidis tanquam morbiprincipisj nomine falutant.

TAB. 8.

Eft Apocynum Indicum floribus parvis, flavescentibus

in dispositis, filiquat nigricantibus, oblongè striatis,



KV-



K U T> I C I K O T> I.

T A B U L A IX.



Vdici-Kjodi, alio nomine *T)uda-Fal/i* planta repens eft nafcens in petrofis.£'auks rotundi, tortuofi, geniculati, fuper fe in vicem convoluti, parum lignofi & virides, quorum *medulla* viridiaquea. *Folia* pediculis craffis, viridibus, interius planis geminata prove- niunt? oblonga, adpediculum, rotundoarnbitu, incuf- pidem anterius ftri&a » texturâ fpiffa, crafla, superficie plana & glabra , viroris in redla parte iaturi, in adverfa dilutij iaporis filveftris. *Cofta* media in adverfa parte emi- net, ac non nihil in recl:a. *Flores* in faciis umbellatis? oblongo rotundo petiolo, in geniculisad exortum foliorum proveniunt, ftelliformes , pentapetali, oblongi, cuipidati? viridi-dilutiexalboflavefcentes, inmediopeta- lorum vena viridiufcula ftriati, collo craffo, orificio ro- tundo ac profundo₅ exterius in collo venis rubefcentibus inftrudti? odoris expertes. Ex illorum meditullio *fiyulus* emicat craffus ? albicans₅ theciformis₅ ex quinque *flaminulis* candidis fpiratim convolutis compofitus, quas fingula apice oblongo, cuipidato funt dotata? & fimuljunbla capitulum flyli eiformant₅ in quo *filamentum* reconditur viride > capitulo oblongo, viridi prasditum, capitulum fuum cum ftaminibus tanquam propriam theculam emittens. *Sedes* cui ftylus infidet ftelWa eft, quinque radiorum > qui collo ad interftitia petalorum funt affuti. *Calyx* florum collum arde excipiens 5 quinque habet cuipides, in cuipidum oris rubefcens. (*Yemm** oblonge funt, primum virides, dein viridi flavefcentes & albefcentes. Linimentum ex hac planta cum vaccino *laSe* , oleo *Sergelim* paratum, temporibufque applicatum, narium fiftit h^morrhagiam.

T A B . 9.

Defci ipto hxc non eft perfe&a, deficit enim] fit *Apocjni*, cui hoc nomen dedi > *Apocynum ludium*, forma fucci, fructuum vel filiquarum, ex deli- *folia* oblongo , gUBio , flore ex albofiaeftente. neatione tamen facile dciumi potcfi: quod species |

Riccia palustris L.
ریچیا پالاستر لایکن
and كاربون يعلان ماليزي



WALLIA-PAL-VALLI.

T A B U L A X



Allia-Pal-Valli, eadem eft planta ac praxe-densj in eo tantum differens ? quod *fores* ftamunculiscareant. *FruBus* oblongi₅cuf pidati, feminibus repleti, quas anterius filamentis argenteis quafi barbata funt. *Folia* recenti *butyro* commixta pro linimento ad verfus exiles adhibentur carbunculas.

T A B . 9.


Etiam hac descriptio imperfe&a eft, itautdeadivinari liccat, nihilominus tamen inter *Petipa-*
(Us vel species *Apgejni*, ob faciem fuam externam, recenferi debet.

H

KA-

کار پالی اگر
بیان کنید که
این میوه

Karpalill
Karpalill
Karpalill



KATU-PAL-VALLI.

T A B U L A X I



Atu-Pal-Valli, planta eft ramis fe vinciens fine ullis claviculis; *furculi* rotundi, glabro viridi *cortice*, geniculis extuberantibus interftindli, medulla la&ea eft & vifcofa > ficut & cortex. *Folia* petiolis flaccidis? craflis, interius fulco ftriatis ac viridi dilutis? gemina fe invicem decuflant, oblongo rotunda, in vertice cuſpidata, oris verfus exteriora crifpatiſ, texturā craffiuſcula? fpiffa? folida ac maximē mollia, fuperficie plana&glabra? viroris in rcſla parte umbroii & nitentis? in adveria furdi & viridi albicantis? odoris agreſtis. *Coſla* media in adverfa parte altiirn eminet. *Flores* plures congregati quafi faſcia videntur, hinc inde brevibus fuccinti foliolis cuſpidatis, pentapetali? oblongi, angufti, cuſpidati> viridi diluti, parum verfus exteriora crifpati; umbilicus *ftylum* emittit thcciformem > ex quinque furreclis quafi ftilulis conftantem, interftitiis petalorum obverfis 5 capitulo albicantc dotaris; ac ipfe viridem recondit gemmulam, ad latera parum compreflam 5 in medio per verticem futura viridi-fufca ftriatam 3 ubi fe folvit, & germen fruſtuum eft. *Calyx* quinque foliatus florem arſte excipit. (*feſmm*<e oblongas in conum ftrid:a3, foliis fpiratim convolutis. *FruBm* ex uno flore in commune calice proveniunt quinque, oblongi, reſli:i> in cuſpidem definentes, ad baiin jundli, quorum medium longo ftriatur fufco ? cortice exteriori valde laſteo& viridi, interioricuticula viridi flaveſcente, cartilaginofa, carne viridi corticis exterioris obftipata. In hifce continentur *femina* oblongo rotunda, plana, ruffa, fcuteſſatim difpoſita; in fuperiori parte penicillo argenteo tenuiſſimorum filamentorum, inſtar fericeſ lenium 5 inftru6h, in medio interioris fuperficiei futuram habent albicantem & flaveſcentem; intra corticem *nucleus* abſconditur can-

didus, admodum tenuis; funtque femina hacc primum viridia, dein rubro ruffeiccntia. Planta haec *oieo* inco&a oculorum vitia emendat; pedum tumores? quos ab incantamentioccultisviribus> derivatpopuli hujus iliperftitio, cvanefcere facit; ubi trita bujus plant* folia affeclis parti- bus applicant ur.

T A B . II.

PI i nta hate i me v ocat u r *Ajt<) uum Indiatm f antm, Urife Hum, Jlu it ri; idi d ilstt, fiUqmi Mtugii, tKf spit Utit.*





جعفرية
جعفرية

جعفرية

Page 221

T A L - F A L L L

T A B U L A X I I.



AUVdli planta fcandens eft , cujus *cau/es* tenues, rotundi, lignofi, geniculati, cortfce ipadiceo. *Folia* in geniculis geminata proveniunt , oblongo rotunda₅ anterius obtufo cuiipse₃ craila , viroris ex una parte umbrofi & nitentis₃ exakeracJari. *Cofta* media unà cum lateralibus in adverla parte eminet. *Flores* ad exortum foliorum racemati ? non nihil villofi videntur, pentapetali, angufti, interiusvirididiluti, exterius rubefcentes, colluloalbicante, in cujus inferiori orificio *globulm* appetet? in vertice *fyllo* inftruūtus parvo. *FruBus* ex fingulis floribus gemini eduntur, rigidi, oblongi, angufti, parum complanati, Ipadiceo rubefcentes, exunapartenonnihil virides; intra corticem tenuem & cartilagineum, *femina* includuntur oblonga₅ angulta, albicantia, fulco ftriata, ac ibicarneviridirepleta, anterius penicillo argenteorum fta-minum dotata. *Folia* cum *Ori^e* lotura trita earbunculos exficcantexiles: *radix* antefebriliseft, fumtacum *Corian-dro* & *Zingiberē*.

T A B . 12.

Mihi ell, *Apacjnum Malabicumfctndens, Utifolium^flire rubfcnte,filiquis, longifimls & dnguflifimis.*

K

NANS-



Smith

NANSJERA-TolTSja.

T A B U L A X I I I .



Insjera-Tatsjajiramannes dicunt *I)adu-Falli*, naſcitur in arenofis. [^]adix albicans, fibrofa> intus filamentolignofoj odoris debilis, ingrati. *Caules* rotundi > viridi diluti, glabri, geniculati, teneriores fpadiceò diluti & villofi. *Folia* in geniculis, quorum pediculi craffi, geminata erumpunt, oblonga, in cufpidem ftricta₅ crāmola, fpiflā* mollia, glabra* viroris diluti & nitentis in recta, in adverfa parte clari) fiocco limpido repleta. *Cofia* in utraque partenonnibil eminent *Flores* fafciatim geniculis caulum adhasrent, pentapetalis oblongi, cufpidati > viridi diluti, qui ad unguiculos rofaceo rubefcunt; in medio *capitulum* appet rotundum₉ feneſtratum, ex *fyllo* conflatum₉ quern quinque cufpidata > nervoia ? crafla, rofaceo rubefcentia > verfuſ ftyli verticem rigide inflexa foliola y cufpidibus arripiunt ([^]emma piano rotundas, parum cufpiaatas & rubefcentes. *Calyx* quinque conftat foliis. *FruBus* ex fingulo flore prodeunt gemini j oblongi, angufti, in cufpidem defincntes, virides > ad pedem transverfimfiti; *Semina* in le continent numerofa* plana₅ cutellatim difpofita ₅ *fenicillum* argenteum anterius includentia. *Integra* hsec planta cum aqua *Lanja* venenum pervomitum expellit; idem prſeftatfī cum *Sandalo* vero, ovo ac muliebri *laBe*, temporibus atque pe<3:ori inungatur. **Trita cum radice** *Car-E/u-regon*, atque *Orifcincodfr radix*, ac lxfxpartiillita, rabidifelismorfum fruftrafacit. **Succus cum Codageny t!Manga-!**[^]rii *Carinta-Tali ScSandah* niixtus ac temporibus illitus, toxicī vires infringit. Cum floribus ^cPalm<eagreilis, *Colupp**, &: *Sam/iru-vari* cum muliebri latte fumtus, plane medicamentum eft Alexiphanniacum; **cum radice** *Jngolam*, *Tiru-uli* &cdmel-^c*Podi* vacino *laBi* fi incoquatur, capitique ac peitori illinatur, omne

virus idternum» & inde ortam fyncopen ac prsecordiory m
aeftum toUit; ita ut hujus plantae ergeia in totum ftc
alexicacsu

T A B . 13.

Nanjeta-Patsja, planta scandens est *Apocyni* CniiJis, succo limpido turgens:





جعفری
جعفری
جعفری
جعفری

M A L A B A R t C U S , ¶

K A M El · T L V A L L L

T A B U L A X I V .




nAmetti-Vatti; lingua Branlanum Vlu-Valli
planta eft nafcens in arenofis ac finuofisramis fe arboribus circumvolvens₅ cuius altera
*species *Tsjera (ametti-Fallii %adix brevis,**
*craffiufcula₅ intus lignofa₅ fibris veftita**
*quam plurimas fpargens laterales. *Stipites**
ad radicem craffi, cortice cinereo, ratni tenues, genieulati^
*cortice fpadiceo fusco rubente? funtque fubladtei. *Folia**
cum fuis pedunculis bina, trinave in geniculis ratnorum
proveniunt, oblongorotunda? cufpideoblongoangufto,
exteriora verfus refiexo, denfa funt, craffiufcula, mollia^
*laftea, plana > glabra> viroris in reffo partenitentis. *fofta**
media in adve daparte eminet) fature rubefcens; in rcfra
*tenuiter exftat viridi albicans. *Flores* umbellatim in apice*
furculorum, cum fuis pediculis rubrisvidentur, pentape-
tali, oblongo angufti, cufpidati > coloris interius candidi >
exterius rubefcentes^ collulo oblongo, rubro circa me-
*dium turgefcenti; *Stamina* in illo continentur breviffima,*
quinque flavis> oblongis? ftriatis ac cufpidatis₅ {lirreclis
*apicibus dotata; *Stylus tenuk*, viridi albicans: f>//*: parvus*
viridiufculus₅ quinque cufpidibus collum florum ardle ar-
ripiens. ^emm^e oblongo rotundas, anterius rionnihil ftri-
*fiores₅ primo virides, dein rubefcentes. *Siliqux* ex fin-*
gulis flonbus prodeunt binse •> pofteriori parte invicem an-
nexx, oblongo angufte, anterius ftridliores, venis fecun-
dum longitudinem ftriatas; primum rotundiote ac fur^
reftse, dein ad latera compreffie cortice exteriori fusce*
*rubefcentes, intus cortice albo fubrufo. *Filamentum* in*
hifce continetur craffum > cortice cartilagineo, corculo
viridi, humido > fuscefcente; exterius tenui, fusce rubef-
*centi membranula> veftitum. *Semina* in illo reperiuntur*
valde plana> tenuia₃ oblongo rotunda, tenui > membri anofa

ac transparente lacinia fuccin&a, in unaguaque filiqua numeroflex, primum viridia > laciniis albicantibus, deinde cum laciniis rufFefcentia. *Succus* cum fucco *limonis* narinbusinje&us fpafmo& epilepticis medetur. *TUnta* cum fru&ubus *Sabac* contrita ac vivario injedum pifcibus anodynum eft toxicum. Cum feminibus *Avanacu*, *zAfJa F<z-tida^ & (jngibere in oleo Sergelim* ceu *T^t Elu*, & *dvanacn* ac *Ori%t lotura* > vel etiam cum *M'trabolanis* & *Jero* lafbis, in unguenti formam redadka, dolores arthriticos lacinantes obtundit, pforam, fcabiemque tollit? per peptiduum ufurpata. Eadem cum *floribus*, (*inamomo* & cortice *Cyno» gloffa* pulverifata jacaqua *Orifk U¿ique* ebutyrato? *melli-que* incopta, pro potu adhibetur in cachexia> lichene atque lepra.

T A B . 14.

Hujus plantai exteriores *G Cpe&cs* notas, etiam *bifeente*, *interim candido^ filiquis tillongit, angustis*,
inter *Apocyna numerali* debet, & quidetrub *vtnis in htifitudinifliatis.*
hocnomHic. *Afocpum Indicum Utiflum, fltrtru-*



Mitchella repens
Linn.
विष्वासी गुलाबी फूल

WATTA-KzAKA-COTtL

T A B U L A X V .



Aita-Kakacodi planta eft arboribus lē vinciens ac arenofisgaudensj *radicem* habet cujus cortex interior albicat, exterior flave(cit₅) faporis f>eramari. (*aules* nodulis extuberantes primo virides funt, ac deinde cinerei.

Folia bina ac bina ex nodulis erumpunt* quorum petioli mediocres, rotundi, viridi diluti, interius ftriati, oblongo rotunda funt, ad petiolum quam maximē curvata, texturāmollia, superficie plana, glabra > lenia, viridi fufca, faporis fubamari. *Qofla* media plures laterales albicantes emittit. *Flores* umbellatim in petiolo commu* ni viridi dilutopronviunt, ex caulum nodis supra originem foliorum erumpente, pedunculi quibus appendent, tenues funt₃ viridi diluti > ipfi flores pentapetalii, viridi-diluti? anteriusnonnihilcontraferti, nūUius odoris: *StelU* in medio videtur quinque angularis? viridi diluta, plana, ex quinque ftylis columnse mediae circumfositconfor-mata. (*jlix* quinque parvis, viridibus, cuipidatis conflat foliolis. (*jemmx* planse, viridifulc^, foliis iuperiori parte fuperfe convolutis & circa medium contraitis. *Frucfus* furculis feu petiolis appendent la6teis , funtque oblongo rotundi 5 verfus fummitatern contrafli, bini in inferiori parte fibi invicem jundti, fecundum longitudinem profun-dis lyris ftriati, viridi flavefcentes, in lyris flavo rore afperfi, cortex craSiufculus eft & lafteus; caro parum fungofa, albicans; interior membranacartilaginofa> albicans, fub* flava. *Semina* in illis continentur valde plana, oblongo rotunda? ex viridi-flavodiluta, limbo flavo rufefcente circumdata; fcutiformiter ar6te fibi mutuo junfta, in anteriori parte oculum habent, qui *penicil/um* emittit cujus filamenta argentea funt. *Ifydx* cum aqua *Orifye* parti appofita i6tum colubri cobra cabella fruſtra facit: *Qortex*

radicis tritus ac cum aqua fumtus doloricapi*Us* infervit, quo
mulieres poft partum iunt obnoxiae.


T A B. I5.

CoJoreminternihu*us* plants; fucri, quamvis
Authoroofter non exprimere, extra dubium ta-
men Aftjtium eft, ficquidem *Apocynum Malabar-*
icum, folio cardis humani formia, flore viridi dilato,
umbellatim disposito, filiisque oblongis, latis, obtusis. Quod
ſi totus hic Hortus Malabaricus in totum à principio
mili fuisset exhibitus; omnes *Apocyni* species

ordine disposuiffem, cum iam per omnes ferme
partes fint dispersae. Verbi gratia *Pala*, *Curatu-*
Pala & *Codago Palo*; quarum descriptio Lib. I.
pag. 81. 83. & 85, videri potest, quae omnes
species sunt *Apocyni Arborescenti*; ut & *Erica* &
Bell Erica, quae sunt Beidel Offar Alpini & Veslin-
gii in obſervat. &c.



*Un-purified
Bitter melon
Seed*



NJOTA.NJOVEM.FALL

T A B U L A X V I



fota-R(JQden-Vdlli; Bramannke ~~Ziru~~ Puti-
Falli, ipecies eft Inota «InoSen, planta finuo-
fis caulis arboribus fc vinciens; funtautern
canliculi illi tenues, rotundi, virides, ex
parte rubefcentes > geniculati. *Folia* in ge-
niculis gemina proveniunt, oblongo anguita,
ad petiolum rotunda, anterius in cufpidem contrafta?
texturadenfa, fblida, mollia, fuperficie utraque plana₅
glabra. *Co/la* media in adverfa parte eminet. *Floret* plures
congregativifuntur incommuni viridi petiolo, inftar co-
rona? coUo oblongo rubro purpureo, fufco & nitente,
inferius ad calicem ventre inBato? veficulato, albicante,
venis rubro purpureo fufds in ventre & coUo, fecundum
longitudinem ftriati ?, pentapetali, ex flavo viridi diuti,
cumoris exterioraverfuscrifjpati; cufpidati; quicufpides
oblongi? rubro-fufco-purpurei ac pilofi, fibi mutuo in
vertice junguntur, inter folia eorumque cufpides fene-
frati; interius pilis oblongis raris parfimobfiti) inodorcs.
Capitulum in medio ventre habent flavum -> quinque angu-
latum 5 ftellas inftar expanfum, ex cujus vertice *Stylus* {efe
erigit oblongus, flavus? theciformis? qui in quinque *Jla-*
mina refolvitur. *Calix* quinque cufpidata habet folia; ve-
ficulato florum ventre ex compreffione rupto, aër cum
fbnitueliditur. *FruBus* oblongo angufta^ filiquas; *femhu*
in iis continentur oblonga? in una ferie difpofita, in verti-
ce filamentis candidis argenteis penicillata. Hujus & *JārU*
%epu, Ela-Cai!i, 73ala, & ^Mirabolani folia trita innodulo
colligantur, qui in aqua *Oritf* caleiäclus & affectoloco
applicatus, ventris ex flatibus tumorem minuit, yel fi
cumfoliorum *Tamarindorum* fibris & *ZMuringu cortice in

murta excoquantur₃ Iinimentum contra articularem *ex-*
hibit morbum.

TAB. I 6.

Pcrcgrina hate pJanta corifpecics eft *Ajatjm* ic-
peniis, quoad formsin flomm mirabilis & fine
fariibencamen pcrfpca, OCIDM *Aptjm* nois
Charaitcritica:, in ca reperiuntur: folia cnim
lina&binafe mucotjecutrant; fios fupcrne in
^uinque Hruia dividitur foliola; turn fcquuntur

filique Jnnigerx, tjuibtu femna includur.^{tur;}
& quamvis entiles fuccum fuadanE limpiUum, non
laftcmn y 8c hoc quaint: in quibusdani *Afecyst*
fpcicibus reperiatur, ^hnu hxc ncutiquim ab
ilJislcparsi poterit.



Paracram Lec.
البَرْكَةُ مَل.

بَارِيَا هَجَانَمْ
arab.



T A R T A R zA M.


T A B U L A X T I L



Lantula eft nafcens in locis humidis & petrofis. *Caules* graciles > rotundi, geniculati, virides. *Folia* in geniculis , pedunculis brevibus, albicantibus > interius parum pilofis proveniunt, parva, oblongo-angufta, cut pidata, adpetiolum latiora, rotundo ambitu, texturâ craffiulcula, mollia, fuperficie lenia, viroris in re&a parte faturi; *Co/la* media albioat, in adverfa parte confpicua > re&am ftriangs. *Flores* plures congregati prodeunt in petiolis , qui ad exortum foliorum emergunt> fuitgue parvi, tetrapetali, cuſpidati, coloris purpureo rubric qui *calke* quatuor minutorum excipiuntur foliorum. *CapfuU* feminales filiquofe ? oblongo-rotundse, anterius in cuſpidem ftrid^e, duo latera convexa funt, quorum unum planius, cum cuſpide verfus latus convexius inflexas. *Semina* quas in iis continentur oblonga funt, tenuia, complanata, unolaterecavo, itadifpolita, ut inteiriora , inferiora fcutellatim obtegant, in ora inferiore cuſpidibus crenata, anterius ftriſtiora, ac in ilia parte, filis penicillatis, candidis, filamento medio communi, craffiufculo, carnofo, albicanti annexa, primum albicantia, exinrufelcentia. *Foliorumfuccus* cumfucco *Limonis*^ infantibus ventre dolentibus , aliisque fpasmodica febre segrotantibus exhibetun *Oleo* mixtus atque ettrinfecè adhibitus, ocolorum vitia corrigit, efficacius autem, fi quando decoolum, ex eodem fucco prseparatum, affumatur.

T A B U L A 17.

Si *Parpaum* hxc plenarie eft defcnpta'; vix I album velviridem funderet fuccum, inter *Apocyn.t* inter *Apocyna* recenferi poteft. Flos enim tetra- numcrari deberet. Alt ex defcriptione hocnon petalus eft, filique vero cum illis *Apocpi* conve- liquet. niunt 5 fi flos eflēt pentapetalus > Sc caules vel



N E L I-T A L L

T A B U L A XVIII.



Eli-Tali; lingua Bramannum 3\(*alaM*, naf-
citur in udis & paludofis, & paflim inter
Orizam, unde ei nomen. %*adix* fibroia,
densè capillata, carne fungofa, denfa, molli,
levi, Candida, corculo pennas haud diflimilis,
ac in teneris fibris corticula flava & pilis fla-
vefcenibus obducia. *Caules* rotundi, glabri, virides, in-
tus carne in totum fungofa albicante; teneriores ex una
parte rubefcunt. *Folia* petiolio medio rotundo viridi fuf-
co, qui ad exortum viridi diluto ventre extuberat, ac unico
novello furculo paflim comitatur, denfo ordine in duabus
adverfis feriebus infident, funtque valde parva, oblongo
rotunda, exterius in medio vena ftriata, viroris fufci, hyalo
ad unam one partem cum tenera fnt rubefcentia. Admi-
rationedignum est, quod fi folia hasc paulo preflius ftrin-
<rantur, ftatim fc fcutiformiter ad fe invicem in duabus
adverfis feriebus claudant, ut & in furculis cum radice ex-
tral disceuavulfis. *Flores* ad exortum petiolorum folio-
rum in pedunculis videntur tenuibus, papilionacei, parvi,
flaviufculi, tetrapetali; exterius foliolum maximè interius
rubefcit, circa oras flavefcens; duo interiora flaviufcula
funt, quartum viridi albicans, in roftro quo cum verfus
folium latiuſculum inflexum eft, rubefcens, onficio fu-
perne aperto, includens decem. viridi albicantia *pmmula*,
flavis nodulis data; nullius odoris. *Calix* quatuor fo hola
habet viridia. *Siliqu** oblong* fnt, plan*, PTMTM "TM
inikxx, cortice primiim viridi. poftruto fufco & fcabro.
Semina aux per corticem extuberant, parv^ fnt fabap,
oblong*, faporis fabacei, in longitudine jiliquarum in
propriis loculis difpo fits, futuris inter medns, in corticis
extuberantibus diftinfti; color fpadiceo fufcus eft ac nitens.


Q,

La-


Latvamentum ex hac planta conficitur antivulnernum.

T A B . 18.

Stliuti plants eft in totum convenicns cum 1 no/*, *filiqws ttlicuUtiş-* f_{cj}j ex omni pane major
inn *Bqmnt Gttmtnity*, *filiqun infummiute*, *B. Pin.* eft , atque ideo vocatur *To rum Unmutn m'ajin*
Jrpertinct inter flrrfcu /lowfafitimttf ftu Ltgumi- \ AuUhrimft, filiijuis in (ummiuit.



IT, ltd - <(!) lat
ବୁଦ୍ଧିମୁଖ ପ୍ରକାଶନ
T1Z& tab.
ସ୍କରାଟ୍ର ମୀଟିଂ



T O T>T>A«VA T>T>I.

T A B U L A X I X.



Odda>Valli duas sunt species, una ejus nominis, altera [^]*iti-Todda Valli*; prior à Bramannibus *Lacberi* vbcatur; plantula est humilis udo & umbrofo gaudens folofemper florens. *l^adix* albicans in annulares nodos ad unam partem contracta, de csetero fibrofa. *Caulis* inferiori parte supra terram repit, fibras hinc inde emittens; superiori parte furredus, rotundus, viridi-clarus, una parte in annulares ftrias eorrugatus; intusaquofus & viridis, in nodulos extuberans. *Folid* brevifimis pedunculis bina & bina ordinatim proveniunt, in caulinis tenuibus viridibus, vel etiam in totum rubefcentibus? ac tenuiter pilofis, qui novem, decem, plures paucioresve ex singulis caulinis nodulis in orbem exēunt > verbus folium hinc inde oblique fe demittentes, ad exortum craffo tubere pullulantes; suntque folia hasc parva, oblongo-angufta, uniformisfermelatitudinis, anterius rotundiola, circa medium petiolum minora, ut & in fummitate, inutraqe parte viridi-furda. In medio *venuld* est apice eminens, quas folium in duo insequalia distinguit latera; ex attractu clauduntur. *Flores* in propriis furreclis petiolis foliaceis, oclo, decem, plus, minus ex eodem caulin libdulo oriuntur» atque etiam inferius ad originem extuberant, pluresque congregati? ac in orbem pediculis albicantibusceuvirididilutis in orbem appendentes? confipiuntur, suntque campanulas pentapetal[^], vel etiam tetrapetate; ipsa petala flava sunt, in medio radio rubro in longumftria; in collo nonnihil albicantia ac venulis rubris in longumftria, ad fundum viridi-diluti sunt coloris, nulliusodoris; circa vefperam & tota nocte ard:è clauduntur. *Stamina* habent decem plus minus brevia, tenuia, albicantia, quorum apices etiam albicant. *Calix*

R

par-



NITI-TO T> T> A-VAT> 7) I.

T A B U L A XX.



Ltera *{pecks linguâ Bramannum vocatur Lauri}*; nafcitur in aquis dulcibus, cum caulis aquas innatans, femper florens, maxime tempore pluviofa. "Radices excaulis Jordinatim hinc inde prodeunt, maxima S parte prope inflexuras funtque condensae ac breves facie ex tenuibus fibris condensatis ac fibi mutuo implexis conitantes, coloris perfici. *Caules* aquis innatantes rotundi, duriusculi, viridi-clari, aet Soli expofiti purjuirafcentes, fungoſintus & albantes, fiftulofi, ad corticem lignofiores, angulisin contrarias partes alterna-tim inflexi, proxime circa inflexuras fungoſa, fpongiofa? humida > albante tunica circumveftiti. *Folia* bina ac bina in petiolis mediis ac ad exortum extuberantibus & rubefcentibus proveniunt; forma angusto oblonga, non tam lata quam illa *Todda-Faddi*, in fummitate rotundiola, latitudine uniformia? mollia? tenuia, plana & valde lenia, coloris hyali, in adverfa parte *venulam* habent fubtile mad fui exortum non nihil eminentem > folium in duo in asquili dividens latera; folia haſc non in eodem piano, utilia primas lspeciei, fed diversi planis fibi mutuo paralellis, petiolum medium oblique lecantibus? & verfus fummitatem non nihil inflexis jacent, tali modo fita ut aperta cum exteriori fuperficie aquas obverfa fint; ex attadtu clauduntur? ut & vespere ac de node, *Flores* iopicatim ac condensè in orbem proveniunt in fummitate priorum petiolorum j qui rigidi, rotundi, non nihil inflexi, coloris perfici diluti, & ex inflexuris caulis vel foliarii, vel cum caulis, prodeunt; funtque inferiores plurium foliorum valde oblongo-anguiforum & crilpatorum, coloris in totum flavi, in *calic%nulh fiamina* vel filiquofum germen eden-

S tes;


tes; fuperiorcs pen tape tali, virides, cufpidati, hidecem
habent *lamina* locgiufcula > albicantia, cum fuis flavis,
Itriatis & oblique furreCtis *apidbus*[^] inter ilia unicUmlon-
geeminet *Momentum*, ex filiquofo viridi germinc inum-
bilico florisoriens. *Siliqute* parvas> plan[^], cotnprefJk&
nonnihil falcate mfummitate furculorum, fex₃ feptem,
plus minus confpiciuntur, ad caulicum transverfim *dtx*,
primum virides₅ dein coloris hepatici rubri, tandem
fufcefcentcs. *Semina* quasin iiso£to_a novem velpauciora
pro magnitudine filiquas continentur, oblongo-rotunda,
plana₃ contra&a, oblique in latitudine filiquarum proxi-
mc adfe mutuo nullis interfepimentis fejundh, coloris
fpadicei-fufci-nitentis, ad oras fufciora & magis nitcntia.
In dcco£to data flomachum corroborat, in feribus uri-
nam provocat & arthriticum fedat dolorem; trita cum
bntpo apoilmata digerit.

T A B . 20.

PerfpcSis omnibus bujus flanta; paitibusj | *tigttit Utu, tamptffu,m'mmhus*, Breyni *Ctnwi* U
prajlbrtim quoad fcmina&filiquas, nulfibi con- foL 47, videctiam Rayum U^J. *Flint.lib.*, 18/V&
venientius potui refcre quam ad *JEffbmentiutt*, I 1, (gf. j.
tuif.stamik vei poims *iltri/jm wffjm ZrrUnhitm*, \$• I



6 Malam-Toda-yadoi L.



>JMALAM-TOT>^rDA-FAT>T)L

T A B U L A X X I



Lantula eft nafcens in petrofis , caulinis
I uper terram expanfis. *%qadix* ruffa eft, fi-
brofa 5 oblonga, refte fe in terram demittens.
Caules tenuesjrotundi, lignofi, fuperno latere
ubi terras expofiti, rubefcentes, inferiori viri-
des. *Folia* in petiolis communibus, brevibus,
ad exortum nonnihil extuberantibus > ac parvo angufto,
viridi\ cuſpidato foliolo fuccinfis > induabusferiebus, ia
fingulis ofto, novem? plus minus ac geminata proveniunt*
parva ^ oblongo angufta, tenuiflima, viroris ex una parte
vividi, alterafurdi; *Cojlulam* habent anterius eminentem,
obliquo traſtu folium dividentem; taſtaiè claudunt? aft
nontam fubitoquam ilia *Tadda-Faddi*; atque turn eo ja-
cent ordine, ut fibi mutuo cum interioribus accumbant
lateribus. Gentis fuperftitio plants huic occultas tribuit
vires, nempe, (fi credere fas fit) ad amiflā virginitatem
reftituendam, & mulierum amorem conciliandam, fi
fanguifuga intents virginis pedi finiftroad faturitatem ap-
plicata > fruſtulatimque difcifla & cum planta hac trita
exhibeatur, vel etiam pette foeminas eadem mifcella con-
fricetur manu. Virenti *Curcuma*] *Camphoraque* commixta
impetigini medetur.

TAB. 21.

Cum Author noſter huius *Malam-Toddi-raddi* neceflores addiderit, nequēliqua\$, ncqucfemi-
na, nihil certi de iſla diči potuit. Aft quoniam fcbjmemene ntty, & meo judido, non male
coaveni Tet cum Mimofadijamau*. Zapopi, nempe
cum prima ipfius Figura. Dę ea vxde IUIUS Hifto-
mimme fpinofa fit, ncceflario adnumeratur ^£- riam Cap. 77.


T

MAN:

P. t'. ro

O M P I F S . O. }
a a - daa- raoor lat

Done?
Lund, Aug
1862. 2. 1



M A N - T O D D A - V A D D I .


T A B U L A X X I I .



Afcitur inarenofis. %^{dix} fibrofa, albicans. (*jules* parum lignofi, rotundi, virides, in furculos divaricati. *Folia* in duabus feriebus geminata petiolis infident, obliquè verfus anteriora converfa, oblongo-rotunda) anguftaj ad exortum paulo ftri&iota, tenuia, lenia, in oris breviflimis ac tenuibus pilis obfita, *cojlula* in adverfa parte, quas brevi cufpide anterius eminet, pilofaftriata> viroris utrimque diluti; ex attadtu confertim fe claudunt & quidem furfum verfus, aliter ac ilia *Todda-Faddi*, cuius latera interiora, deorfum verfus fe claudunt; claufafcutellatim fibi mutuo accumbunt. *Floret* in propriis furculis proveniunt, quorumpediculibrees, tripetali funt, coloris pallido flavi, fubtiliflimis venulis flriati. *Stylus* in medio ruber. *Calix* cyathiformis ac viridifufcus. *Siliqute* oblongo anguftasfunt ac compreffae, in cufpidem definentes, maturas coloris rubro fufci, parvis? rotundis gram's ceu *feminibus* confertse, quse primum viridi rubelcunt, poftea alblicant. *Balneum* ex hac planta pr[^]paratum, lepras ac pruriginofo acrique ardori medetur; cum oleo *T[^]it-Elu* ceu *Sergelim* ex eadem linimentum conficitur antipafmodicum. *Succus* cum *Sandalo* tritus frontale eftadverfus casphalalgiam calidam.

T A B . 22.

HJEC H<tawt*fpecies, anemine, quodfciam, ac filiquas articulatsvs, ideo optimojure vocar unquam defcripta eft. Parva quidem confpecies poteft; *Mitnofa Malabarica*, foliispilofts, mm spinofa, videtur *Mimoфа tionspinof* majoris Zeylanica* Breynii> minor, flort papilionaco lutto, fili<gt;as angufUs, articu Sc cum nullas habeat fpinas Scilores papilionaceos latis.



A R I A - F E E L A.

T A B U L A X X I I I .



'Rta-*Feela*; velfecundum Bramannes T/Aw/, planta eft altitudine quatuorpedum , nat cens in arenofis. *Ifyrix albicans*, fibrofa 5ccapillata, *Cantes eorumque firculi* virideshincinde parum rubefcentes, ftriati6c capillati. *Folia* inpetiolis terna5quaterna, quina congregatim proveniunt, parva, oblongo rotunda* anteriuscupidata, tenuia, mollia> fuperficie plana > brevibuspilisobfita, faporis fubacris, odoris fortis & gravis. *Cojlam* habent in medio plures laterales emittens. *Flores* qui ex origine foliorum, in viridibus? tenuibus pilofisque petiolis, & ex fummitate furculorum plures proveniunt, parvi funt, tetrapetali ? rotundi, parum oblongi, tenues, flavi leu lutei 5 fingulivenulaftrati. *Stamina* in illis videnturplura? tenuia, viridi flava, furrefta; *apices* in vertice habent nigros, *Stylum* craffiufculum. *Siliqute* viridi fufcas funt 5 quatuorpollices, & quod fuper eft longas> recT:a3, oblongo anguft^, parum comprefe, in lateribus ftriatae. *Semina* qu^denlaferieinfiliquisinveniuntur, futuris adherent? funtque rotundo plana, primum viridia, dein flavefcntia ac rufefcentia ? tandem nigricantia, *oculum* habent in medio, ex quo tanquam centro, in reclum ftriantur; probe ex ficcata lufca funt, ftriis turn minus confpi- cuiis. *Succus* naribus affumitur ne venenum in corpore latens caput inficiat Ad idem conduceit tota *planta* aqua; inco&a & epota; *folia* appetitum promovent, expeclo- rantia funt & confortantia; phlegmata & flatus diffipant; trita cum radice & in globulos formata, fub axillis in prin- cipio paroxysmi febrilis applicantur? & turn frigus & mi- nuunt & tollunt. Foliorum *fuccus* cum oleo frixus & auribusimmiflūs furditatiopitulatur; eadem folia trita & capiti appofita, dolorem ex frigore ortum fedant.



Cissus rotundifolia
Linn.
Cissus rotundifolia
Linn.

CAR A-P E E L A.

T A B U L A XXIV.



Jara-Feela; vel uti Bramannesdicunt *FeeU* & ut *Tiloni*, planta eft fortis & gravis admodum odoris ⁵ nafcens in arenofis. %adix ut ilia *aJria-Feeli*. *Caules* furculique viridi clari* dein ad unam partem rubefcentes ? rotundi, non ftriati, magisque capillati. *Folia* quina simulproveniunt in petiolis viridi-albicantibus, capillatis, quorum pediculi breves, ftriati, casteris quidem majora, aſt in reliquo quam maxime convenientia. *Flores* in fuperiori parte furcotorum proveniunt, paulo majores, can* didi, tetrapetali, oblongi, corrugati, nullis venuſis ftriati. *Calix* quatuor foliorum, ex cuius petioli nodulo *ftylus* fefe erigit nitens, fuperiori parte nodofus, ubiquinque eminent *flamina* longiufcula, rubefcentia? *apicibus* flavis ⁵ furreflis, ftriatisdotata. *Siliqua* longiores & viridi-clariores, non ftriatse, fed afperse & fcabrse, pilisque minutis obfitse. *Semina* plane eadem, fed majora. *Integra* hsec *planta*- contritacum fuccode *Tfyca-Talam*, fadlapernunctione puftulas amovet; in *oleo* decora, cutaneis medetur morbis, pra?-lertim lepr^e ; Vide quas de viribus *Jria-Feela* diximus? cum qua omni ex parte convenientiunt.

T A B . 24.

Plantahsec, meo judicionulli coramodiusad jungipoteft, nifi *Pentapbyllofiliqiffo*, &: interconfpccics *Tamiaja Brtflienfium*. Margg, atquc ideo reae vœacur *PemafbijllumfiliqHofam MjUturiei,fe*.



تاجیکستان
گلستان
میوه
بادام
جعفری

T A N T) A L E - C O T T I

T A B U L A X X V .



Uinque bujus plants funtipecies, quatuor ejufdem nominis ; quinta .*Tee-Tandalecotti*. Prasfens hasc à Bramannibus vocatur *Sckama-Tufpi*, nafcens in arenofis & udis. ^c*B^dix* fibrolâ₅ albicans? intuslignofa. *Stipites* qui plures exradice exeunt, ut& rami virides, teneriores, angulati. *Folia* in ramis oriuntur, oblongo-rotunda, linguiformia> tenuia₅ texture fiibtilis, lenia, colonis ex flavo viridi diluti. *Co/la* media in adverfa parte cminet. *Flores* fpicatim videntur, aureo flavi, papilionacei, tetrapetalii, odoris nullius. *Stylus* viridi dilutus, craTus, in decem viridi diluta dividitur *jlamina*lula, quorum quinque, qui supra cetera *emicmt, apices* gerunt flavos feu croceos; reliquorum apices flavi sunt *dc* quafl noduli. *Calix* quatuor acuminata habet folia, (*Etnma* florum virides , oblongo-rotund[^]e, unguli-formes. *Siliqua* parvaefuntj recT:[^], oblongo rotundas j glabra₂ circa ventrem futura verfus inferiora contradlsB) *calice* excipiuntur cavo₃ in quinque acumina foliola divifo. *Fab a* parvas magnam filiquarum partem vacuam relinquunt 5 umbilicis binas & binx future ventris adhasrentes, suntque planse, reniformes, primum viridesj deinfubrufo ecu ex flavo fiibfufcse, &inflar *Veluti* nitentes. Ex decofto lavamentum pra3paratur, pro febribusj tertianis praecipue chronicis. [%]*adix* in decodlo data pituitamevocat₅ flatus difcutit. *Tritacum yino pr& ftantiffimum* eft anticolonum. *Foliorum* fuccus epotus febres symptomaticas ex colico dolore aufert, per vomitum nempe; hepar & fanguinem purificat.

T A B . 2 5-

Omnrs hxquinque fucccdentespiantx pertin-
ent inter *HrtLifftwp.ipitH/.t.ttetftHqttifM.* Do&ti-
iimus D°. Hermans VOCOE ilia* in i'uo *Cjtdogv*
<*TetoUrU*) quia In Jorum infantbus in ludo pro
crotakinierviunr; Cingalenks dicunt *Rili'ilippe*
id eft, Crolum j Vide piiudidiirn *C*tM. pig.*
196. fed ut debicus inter phmas, il'.is aalignetur
locus, inter *Icgum'tm noa {cjudentia» hliqms bul-
fozis , fminc rcnifwmi colloca-i debeni, V'cra*

quia apudnullos Auctorej hucufquc occurrcre,
ut novafspecies inferend^c funt. Hxc igitur no-
ftia *TAnitlc cwi*, nan male convenirc videcur,
vcl forceeadem eft.quam D.D. Hermans in *CMJI.*
Hart. Acid L<>!•. sm*, nominat, *Cwt.ddiu. ifuiuj*,
folio *jinguUTtoi at far mi, fluribus iaicit.* MM iciitcoiii
eft, hanc & rcliquas pro *Genifl** generibus ha-
bendasede, atqucadco imevocatur *GmijtstdA**
Ulutict, foliofugnlAii tfisribtu liatisfitifmti/mtsU.





Botanical
Illustration
of
Acanthus
sp.

Botanical
Illustration
of
Acanthus
sp.

T A ^ © A L E-G O T T U N o. 2.

T A B U L A X X V I ,



Ecunda species a Bramannibus *Daco/i/ageri*
In nominatur. Cum priori in omnibus pene
convenit) præterquam quod *JblU* longiora
fint, in redta parte colons viridi fulcioris,
in adverb hyali. *C°ft^* niedia akius emi-
neat. *Flores, filiqua , faba , vires > plane*
idem & eaedem.

T A B . 26.

Lmut-TaniUle-Cotii, vt1 i"iJMrfjifCotirifyLi'eflris,
apud ncmincm mcntio fafta, folia cuani habet
ftngula , **qux** , "fi **CrfWiiri*** nomen placerct ,
vocari poffct, *Creulau M4*b*ri(* [j]vefirit, foliit* |
fingUribati mtjtriitus, totfcwj iuit'u. Mcliusauwm
Gmifl M,d*b4ti(4, folio ftn£tdini tbtwgs, fart fl<t¥t-*
dilutkTt, [tStquis bulUltt.



Aa

TAN-



نیوجنگلند
جگہ پریمیون
کوئن سٹی، نیو جرسی
جے. ایل. مارک

NELL/A<7ANDAL&COTri

T A B U L A X X V I I .



Ertia Ipecies; lingua Bramannum *Gageri* di<Sa; altiusaflurgit, adalitudinemf'ermc hominis; nafcens in arenofis & udis. 'Radix ut prsecedentes. *Stipites* ac *rami* viridēs? angulati,ligniofiorescasteris. *Folia* numero terna in uno petiolo , rotundiola funt> reliquis ipeciebus breviora , colons in ref̄ta parte viridi-fubflavefcents ; advcrfa hyali. *Flores* iidem , nifi quod illorum ungula paulo major fit. *Siliqua* cum illis primas Ipecieiconveniunt, ut& *Semina* & vires.

TA B. 27.

ConfpecieshxvKtetur *CrttoUriaAftaticafrutef-* | illamultolatiora &rotundioradelineavit, Sores eentis trifolioft, fiaribus luteis Amplis: Hermann: Catal; | alioquin8e filiqua: non multum difFerunt j ego difFerentia tamen in foliis eft, quae cum illis j illam dicerem *Genifta trifolU MaUbaric**, fionbus Hermanni non conveniunt, Author enimnotfer I luteis amplis 3filiquisbullatis.

Bb

WEL-



W. H. Bishop, Jr.
G. W. Brinkley
J. C. Ellsworth
J. C. Ellsworth
J. C. Ellsworth

WE L LIA-TA^T) ALE-COTTI.

T A B U L A X X V I I I .



Uarta species à Bramannibus vocatur *garo*, altitudinefermetriumpedum. %*adix* & *flifites* convenient. *Folia*, numero quina, oblongo-anguftiora funt, anterius rotunda & ad petiolum contrada, colon's in refta parteviridis, in adverlā hyali: *Flores* iidem; filique longiores, brevi pedunculo calici infidentes, latiores., ventre extuberantes, quo velut anguftas funt veficas, multoaëre replete. *Faba** vires utfupra.

T A B . X_8.

Nemo hujus speciei quinque foliorum hue ufqiic mentionem fecit \$ rocari, idcirco poflet *Genijl** *Malabar tea pentaphytidis, flo rt amflo tHteo flavefcente, filquis buUatis.*

Cc *TEE-*



ପ୍ରକଳ୍ପିତ
ଫୁଲ
ମାତ୍ର
D. 125

<P BE-TJ^T) AL E-COTTi.

T A B U L A XXIX.



Uinta species apud Bramannes "Butta-*Cja-*
gcri audit ; *radix*, *Jlip̄s* à prioribus non
differunt. *Folia* laciora funt, oblongo-ro-
tunda₃ maxima latitudine in medio, inoris
crifpa, texturâ crafla, tenuiffime pilofâ 6c
fubinde lanugine albicante obdu&a. *Flores*
purpureo-casruleo-diluti & albicantes, vetuftiores magis
colorati; latiffimum folium exc^cruleo maxime albeïcit?
& fubinde in dorlb nonnihil pilofum eft: duo furre6ta co-
Ioris funt purpureo>c«ruleo faturi; ungula viridi albicans?
ac in cufpide cum roftro nonnihil c^aerulefcens; apices *fylfi*
oblongi) flavi, crocei. *Calix* exterius nonnibil pilofus.
Siliqua ut *illx* primse Ipeciei, fed pilis decumbentibus te-
nuiterobdu(3:2B; *Semirne* e^adem; ut & *yires*, prsenter quod
hujus *radix* trita & oculis appofita, memoriam reducat
& corrigat: fuccus foliorum Mercuriifluorem diminuat:

T A B. 29.

Crotalaria hacc fpecies eft *Afuttica*, *folio fingulari verrtuofo*, *floribus camleis*; *Herm. CattL Hort. Acad.*
Lugd-BtVayx fecHndummc, Gmiflalnditta) *Alnifolio*, *floribus caruleis*, fuitquis bttllatis.

Dd


KAT.

Klein-Boeket

1891

عَالْمُونَ لِلْمَعْتَدِلِ

Doubl. Teil 10.



K A T T U - L A G E R A .

T A B U L A X X X .



Lanta eftaltitudineduumtriumve pedum* nafcens in arenofis. *ifyJix albicans*, fibrofa. *Surculi lignofi*, rubefcentes -> pilis longiuC culis rubefcentibus obdufti, teneriores ex una parte virides, ex alterafanguineo-rubefcentes, vetuftiores cortice ruffo-fubrubefcente; corculum interius album eft. *Folia* in communi brevi pilolb petiolo trina & quina proveniunt, oblongo rotunda, utrinque pilis albicantibus denfe veftita, tenuia, mollia? fuperficielenia? forma ut *Tagera*, fed pauloma-jora₅ laporis adftringentis & fubacris, cum *cojla* in adverfaparte eminente, & reftam ftriante. *Florès* plures congregati, ordinata ierie in cacumine furculorum > & hinc inde alternatim vifuntur, parvi, papilionacei, coloris rolaceo-laturrimi. *Stylus* qui folio ungulato accumbit, theciformis, viridi dilutus, fupernein brevia filamenta, apicibus flaviufculis nodulata ₃ fiilus. (*jtlx* folia habet viridia anguftiffima, fex numero, pilis veftita. *Siliqux* breves funt, brevibus pedunculis feriatim appendentes, rotundiolse, valdepilofe₃ ad appofitas partes duabus futuris viridi-fufcisinlonguminterftin61:fe, inter futuras pa-rum extuberantes? illarum cortex primoviridis, dein ex ruffo-fubrubefcens. *Semina* quseiniis continentur parvse, ad maturitatem provedlas₅ longas, in longitudine filiquarum fitse» interfepimentis a fe mutuo interfintinft^, albicanter & fubflavefcentes. Ex hac planta decoctum conficitur antiphrenitidum.

T A B . 30.

Specierum *TageM^* mentio fada eft *Tarufe-iund. bujus operis, pag. 102.* ubi etiam noſtra *hxcrnemoratu*_i Quamvis autern omnes plants fine filiquofo, cumſiac tamen nulla liuerceditcoiv venientia, conſpecies quippe *Aſtralagi* eft 3 quid jamnomen *Tggra* propri[^] ſignificet, nonnovi,

hocquidem, quod *Katu-Tagera* tantudem dicat, quam *Tagra filviftris*. Plantam hanc in Herbario meo vivoCeylanico poflea inveni[^] à D.D, Hermans vocata, *Aſiralagus Indicus* (*ptcatus*, filiquis copioſide orfum[pedantibus, nwfabatis; ſeu *Polylobos*, à Zingalenfibus *PiUgbas*.


Ee

KON-

Kondam pulli *Vaccaria*

தெங்கு மலை.

جُنْطَلْ قُبَّاعْ a. 10.



K O NT> A M-T A L L U.

T A B U L A X X X I .



Lanta humilis; arenofaamans; *tydix albicansj fibrofa*. *Caules quadrilateri, viridi-clarij aquei> geniculati; filamentum medium carnofum eft*, viride ac in tubulos diftin[<]5tum. *Folia geniculis geminata?* & ordinedecuffatoinfident₅ oblongo-angufta, angulis breviflimis raris iparfim crenata₅ rigida, craffaj glabra, viridi-clara₅ adveria parte in medio altè extuberantia. *Flores ad exortum foliorum in petiolis prpveniunt brevibus, c^ruleo purpurei, bipetali, uno ad ventreni colli, in tria foliola rotunda infedii*, ad medium ufque ftriisalbicantibusnotato, altejrofurre&o, cuipidato, verfus primum reflexo. In medio unicum habent *flamen* candidum, fupernerubro-purpureum. *Calix foliahabet <juinque viridia , culpidata*. *CapJuU feminales oblongo-rotund^*, in quibus lemna arenacea₃ ut iflapilcium₅denlbj confertoquenumeroj confident.

Niv-a-rii Linn.

QlaQ mJ.

Saniti

سَلْكَلَةَ بَرَم.

نَبِرْ مَقْصِبْ نَبِرْ

Arab.



N I R-CM U R R t

T A B U L A XXXII.



Etricis & aquofis gaudet₅ altitudineduum pedum & famplius. [^]*ami* parum lignofi, virides. *Folia* quse pedunculis brevibL7s proveniuntj oblonga? tenuia, Viridi-dara. *Flores* papilionacei, rubefcentes₅ in niediq ex rubra ad caeruleum vergentes? odoris nullius. *Siliquae* parvas, oblongas? re<3:a3, anguftse, compreto, ficcse rubefcentes; *Semina* longse? quadratasalbantes. *ThnU* hase excepta radice 5 in olla bene obturata aquas incoquitiir? cuius vapor tumorem ilium cacochyto'cum, quem endemius ille morbusj *Pitao*₃ diffusus inducitj diffipat.

T A B . 32.

Nir-Afini, feccndum Bramanncs j*i*//₅ perrinctfriier *Utr'Wsflare papilionacea* rea. *terri* fpecies eft:

Gg

CUPA.




Fig. 1
C. - 200
Galapagos
Isles
1889

CUT A-V E L \a.

T A B U L A XXXIII.




XJx planta Malabaribus *Cupa-Fela*; J3ramannibus *Itynuquela* dicitur, altitudine eft duorum triumve pedum, nafcens in arenofis. *tydix* albicans? fibrofa. *%irni* quadrangulati, viridi-diluti₅ geniculati. *Folia* in geniculis geminata & ordine decuflato, pedunculis proveniunt brevibus, oblongo-angufta, in cuipidem ftridta? viroris communis. *Co/la* media cum lateralibus in adverfa parte eminent₅ reftam ftriant. *Flores* in furculis ad exortum foliorum > duo tresve congregati ipe&antur, parvi> pentapetali₅ albi, rotundi, ftelliformiterexpanfi, cuipidibus minutiffimis eminentes; collo oblongo₃ craffiole? viridi albicante, ex *calice* arflo, viridi-diluto, quinque foliis cupidatis exftrufo emergentes. Illisdeciduis *filiuae* fuccedunt otriongx, tenues, rotunda, incufpidemftrid^e, leviterintort^₅ primo virides₅ dein purpureo-fufce-rubr^j tandem cinerefcentes. *Semina* in iicontinenturoblonga, fulculisftriata, lacunam habent unicam oblongam₅ profundam> atram. *Folia* indigene adhibent in concoftione *Caril*

Tsjeru-véla le.

ବାଲ୍ମୀକି ମାଳ

جَبْرُونْ جَلْمَلْ

aub



T S J E R U-F E L A.

T A B U L A X X X I V .



Reno/à amat. *fydix* albicans > fibrofa * filamentum in medio habens albicans & lignofum. *Caules* virides pilofi, intus parum lignofisi. *Folia* in petiolis pilosis 5 viridibus provenientia, oblongo-angufta 5 in cufpi* deminteriusftrid:a,pilofa. *Co/la* media cum reliquis in adverfa parte eminet, redlam ftriantes. *Flores* supraexortum foliorumvidentur, tetrapetali> oblongo-rotundi? angufti, rubro-diluti, fiib purpurei- *Stamina* iniisquinqueoblonga, albantia, quorum apices oblongij viridi-oerulei; cum illis fefe prodit filiquofum germen* pilofum, viride 5 cuius cufpis rubefcens verfus eandem partem cum ftaminibus incurvatur. *Calyx* ex quatuor conftat oblongo-anguftis, cuipidatis 5 rubefcentibus foliis. *Siliqute* oblongo anguftas ? rotunda, in cufpidem defihentes, pilofe, lacunulis ftriatas, virides ? ut illas *Cupa-Fela** Nervorum vulnera ac pun&uras fanat hxc planta₃excepm radice, cum aqua *Ori^e* trita.

T A B . 33. 34.

Utraquchaecplanta, quamvis quoad formam J tapetalus fir? Sc ex defcriptione non fatis contgeti foliorumdifFerant, hoc tamen commune habent, I utrum in totum decidac nee ne i planta luce- quod flores & filiquas ex ortu foliorum prove J anomala eft. niant, & cum flos ieeundum delineacionempen-1



ପୁରୁଷ କାଳି
ପୁରୁଷ କାଳି
ପୁରୁଷ କାଳି
ପୁରୁଷ କାଳି

V A L L I A ~ C A < P O - M O L A q 0.

T A B U L A X X X V .



Ecunda haec species cum priori, in fecunda
partedeicripiaj convenit; eft Sctertui ipe-
Icies, in eo tantum diiFcrcns 5 quod fru6hi.s
digitum ferme oblongf, ac multo crafEorcs
evadant. Vide ejus defcriptionem *part. z.*
TaL 56. p&g 109.* Furibundi cam's virus
enervaturj ubi l^iaparsj trito hujus plan tie *fru\$u_y* vel
ctiam *udice ijlinatur.*

T A B . 35.

Inter fpcciej *Ctffe virix* fm£htum formar | Hernandum *nijf*. *Plant. Mexic.* lib. 5. cap. 3. at-
fuat>utividcrec[lapudCluCum *Bifl. ri**>*t.txeti-* que inici ilias hsc ctiasn *Valid-Caps-Molayo*
**r, *ftl*, j^p. & j_n car. pejto, f/. j i > ut & apuJ J comprchenduur.



Kk

NIR.

Perseus

Nir-Gulári L.

Malabar

نیر گلاری



مار

ل

N I R-? U L L A R I

TABULA XXXVI.



Lantula eftarenofisgaudens. *Caules* tenues* virides, capillati > fuperne rubefcentes. *Foliorum* petioli rubefcunt) ad exortum duo* bus minutis, cupidatis foliolis succindij folia ipia oblongo-rotunda > tenuia, Jenia, viroris in refta parte viridis, adverfa hyaii > ubi *corta* media eminet. *Floret* plures congregate lupra exortum foliorum vifuntur in tenui Emis pedubculis, papilionacei, tetrapetali, coloris rofaceo-faturi-purpurei. *Stylus* ungu latus, oblongiusculus, craffiolus > candid us^ fuperne in tenua filamenta fiffus, quas apicibus flavis nodulata funt, odoris nullius. Integra hasc *planta* cum *ferola*(3;is, vel quoque per fe in balneum prasparatur, antido* tale contra morfum ferpentis Po lega : trita & cum *oleo* 3\(*ucis Indiea* fortiter frixa, in colica ac dyfuria cum fuc* ceflii afluxit; mingendi difficultatem tollunt *folk* ex *Aqua Lanj** cum *Calamo Iromatico* exhibita. *Succus* cum *jure* (*jdlimceo* epotus lithonthripticus eft.

T A B . 36.

Quxhicdefcibicur *Kir*rullari*, pertinet inter legumina trifoliataj aft turn de vera filiqmm forma nulla fiat mentio, vix certi quid de iliajudicari potefh

Li

MAN.



I



1
Mallotus

M A N N E L I


T A B U L A XXXV II.

NlAfcitur in arenofis. *fymi* dtiri •>, viridcs* ex parte rubefcentes. *Folia* plura congregata proveniunt ? quaterna > iena ? plus minus, parva, oblongoangufta, tenuia, vi-[ridi-fuica. i^orw papilionacci , pulchre rubefcentesjodorisnullius. *Siiiqme* valde parvx, oblonga-anguftse, *fibulas* in fe continent virides, quie poftea rubefcunt.- *Folia* ab indigenis adhibentur in concoftione *CariL* Exintegra hac *planta cum UBeebw tyrato* lavamentum conficitur, femorum refolvens tumores ? phlegmonodes, CEdematodes. In *orifye* lotura apocicma pr^paratur ventris inflationem auferens altumque emolliens. Ex *cineribus* Jixivium elicitor, quo cum fordes capitis expurgantur. *Folia* trita abfccfTus refblvunt vel aperiunt. Ex *radice* oleum cxtrahitur > quo caput in eryfipelate perungitur.



M m

•TSfO.



جیوکل پاکستانی

جیوکل
جیوکل پاکستانی
جیوکل پاکستانی

جیوکل پاکستانی

T S f O V A ^ Q J N ^ I - < £ M A H ^ C E L L .

T A B U L A XXXVIII.



Lanta humilis eft? furculis numerous, ift latum diffufa ? aquofa amans. "Radix albicans j fibrofa. 'Surculi tenues , ex omni parte rubro-perfici 5 rigidiufculi. *Folia* in singulis petiolis geminata proveniunt> duobus tribusve paribus; parva, & ubi veterafcit planta? in minorem formam contracta 5 oblongo-rotunda, colon's furdi 5 ex attaftu clauduntur 5 ut & de nocfe. *Cofta* in adverfa partc eminet 5 ubi folia, maxime in cofta & oris, pilis albicantibus & nitentibus obfita funt. *Flores* qui tres quatuorve iSmul ad exortum petiolorum folia-[<]eorum prodeunt > papilionacei funt 5 flavi, inodores> tetrapetali. *Stylus* , candicans , theci-formis > fuperne in filamenta, croceis apicibus dotata ? fiflus, quorum una fupra apices emicat. *Calyx* quatuor conftat foliis. *Capful & feminales* oblongas, rotundiolas, breviufcul^e, tria, quatuorve incidentes *femina*, rotundiola, nonnihil plana> primum viridia? dein fuscentia 5 poftnigra acnitentia; feriatim filamentoviridi, quodinflore emicat, acfeextra capfulam crilpatim exferit 5 intra capfulam annexa & ad^ hasrentia. *Unguentum* ex tritis hujus *foliis* & *recenti bmyro*, fecundum artem pr^eparatum ablceflius tollit phlegmonodes , quos fi concomitentur febres 5 decorum pr^fcribitur? ex hujus *radice* (fingibere dc "Pipere.

Tjérw-mannet *lax.*

جَرْجَرْ جَرْجَرْ *Mali*

jata

جَرْجَرْ جَرْجَرْ *Arab.*



TA 30



T S J E R U - M A N N E L I

T A B U L A X X X I X .



Afcitur hsec plan tula in aquofis, caulinis ramofis fiiper terram decumbentibus repens, ac hinc inde radiculis fe affigens. *'Radix albicat, aquea, fibrofa, capillata eft.* *Caules virides, aquei, geniculis interftindti, ingeniculis foliis paribus, quas paffim novellis caulinorum partibus ftipantur, veftiti. Folia valdeparva, cuipidata? craffa? rigida? lpiculis candidis nitentibus, longiufulis) obquse&afperaflint, munita. Floret in canibus fupra exortum foliorum₅ sex calicibus oblongis? ardtisj collulo candido cmicant, pentapetali, cufpidati₅ ftelli-formes, in totum candidifimi. Stamina quinque parva, nodulisflaviufculisdotata, in medio flylus ieieerigit. Calix quinque oblongorum eft ctifpidum, fpinis *candidis*, fpi-culatis, afperis munitus. (*flobuli* feminales oblongi, virides₅ parvisarenaceis/^///ferepIeti; prirnoviridibusjdein flavefcantibus, tandem nigricantibus. Integra planta exceptaradice? pulverifata&in *ferolaSis* cum *arugine* con-trita? pedum clavos cradicat: idem pulvis cum modico *'Piperis* & (*jlycirrbife* *Jucco* ex aqua calida propinatur₅ vel cumrecentibutyroexhibiturtuffi (Luiitanis Cogreira de Garganta) Jaborantibus.*

T A B. 39.

Tresquashic, fub cognomine *Manneli*, def- | *GenifleUtin8ori&. Tsjovanna-Manneli, MirnoJA gt^us*
cribuntur plantar> à fc invicem differunt : *hUn-* | *eft 5 ScTsjetu-Manneli* planta eft aquatica, omnes
ncll cnirn eft inter Papiliomteasfiliquots, species | ergo ad finguarem pertinent *fpecieſ*.



بِسْمِ اللّٰهِ الرَّحْمٰنِ الرَّحِيْمِ

سُلْطَانُ الْعَالَمِينَ

سُلْطَانُ الْعَالَمِينَ

S V E 3^T) J T> I-<P V L L V.

T A B U L A XL.



Rcnofo h̄iEcplantagaudetfolo, alritudine d̄uum triūmve pedum. *adlx* albicans, nbrola. (*j, uks rqtundi*, virides, parum lignofi. *Folia in petiolis tcnibus paffim tcrrna onun̄tur*, oblongo-rotunda, mollia, Jenia, virons in re \angle a parte fufci & furdi. *Co/la* mediainadverfaparteeminet, redlam ftrians. *Flores* in propriis petiohs viridibus, fupra exortum pediculorum fōhaceorum, plures congregati vidcntur, inferiora verfus inrlexi, parvi, papilionacei, tetrapetalii. *Stylus in (lamina*
S i T M W ne Y „ „ *ffillamæmtum*
 oblonp, m̄gricantes, fingute unicum /&, & contiocntesoblongo-rotundum, flavum, fubrufum. *Yi_{Kas} hxc* plantanullius eft odoris, ficca grati. Per defillationem aquae hctur fragrantifliina. *FruBus* in pulverem redadi, artuum Jaffitudmem adjuvant.

T A B . 40.

Jem! ^{fej}! ^{te!} ^{liura<ll^A*} ramnofrocommoni m.gnam habaa convenicKian, finon

P P CO-



Chamaesyce

Burm. f.

CO L E TT A. V E E T L A.

T A B U L A X L I .



Inguâ Bramannum £W^, planta eft trium, quatuorve pedum , arenofa & uda amans loca. "Radix numerofis lignofis fibriſ unde- quaque circumdata & nigricans. *Stipites* ut &: *rami* rotundi, viridi nitente, glabro *cortice veftiti, geniculati, in geniculis fpinis transverfis viridibus, tenuibus, acutis, longiufculismu- niti. *Folia* in illis bina & bina proveniunt , fingula fpinis utrinque quaternis comitata, ordine fe mutuo decuffantia, oblongo-rotunda, nonnihil cuipidata ? iſpiffa, lenia, gla- bra, colon's iiifci &c nitentis> faporis amari. *Cojla* in ad*; verlaparte eminent; ut-& *Flores* ternijquaternii maxime ad fummitatem furculorum ? *collo* flavo longiufculo ex apertis calicibus, qui plurium funt foliorum prodeuntes, flavo-rubefcentes > pentapetali, odoris expertes. *Stylus* in medio tenuis 5 & albicans 5 ac cum eo *lamina* quatuor flaviufcula 5 ac colli inferiori parti inferta, duo furrediora & craffiola 3 flavis , furreclis > craffis *apicibus* nodulata> duo breviufcula ac demiflā , in collo inferius recondita. *Floribus* deciduis fuccedunt *Capful<efeminales* oblonga^com- preffo, lateribusparum convexisj & incuipidemoblon- gumftrid:^., rigidas, lignese? cortice intus albicante, qui viridi inveftitur corticula; in medio laterum comprefTo- rum& inoris, futuriftriat2B. In fingulis capfulis, duo reperiuntur *Jemina*, in propriis loculamentis? interfepi- mentolignofo, craflb, à fe mutuo fejundia, rotunda, parum oblonga,unaparte qua interfepimento accumbunt, plana, altera nonnihil convexa, colons ex albicante iiib- rufi, nitentis, faporis amari.- *Folia* indigenis vicem fo- liorum '*Bethel* in comeftione, *Jrequa* prſeftant. *Succus* aphtis conducit? in teftinorumque ex flatibusintumefcentk.

T A B. 41.

PURNA hacc nū!nm habet conrpccicm; atio-
nwla ergo crit, ul^ucdum major ad nos illius
pervencrit notitia- Anno 1688. illam in Barto
Medico -Amftdodamenfi *ex icmine produxi, nec*
Cpcrehradere potui, illam anwtwe fuiffc dzcri-

piatn; polka autem in Herbnrio vivit *Ceylanico*
obviam mihi ia&:i eft, ubi a DD. Hermans vo-
tan sevijas *toribus lateis*
Cingolcnirs dicuri *Kalnikarabini.*



L

A

1795
1795

1795
1795

1795
1795

V A T> A-K O T> I

TABULA XLII.



*Jda-Kpdi; Bramannice Kjdb-Adulujjb, plan**
taeftaltitudine trium quatuorve pedum,*
arenofum eligens & terreum folum'. T^aqadix
fibrofa, albicans. Stimes ut & rami rotundi,
atro ceu etiam Ipadiceo nigro nitente cortice,₉
geniculati, in articulorum initii extube-
rantes; corculum interius fungofum albicans. Folia in
geniculis bina & bina videnturin pediculis brevibus? in-
terius planis, fpadiceo nigris & rubefcentibus, ordine fe-
mutuo oblique decuflantia₅ oblongo-angufta₅ in culpidem
anSfcumoblongumanguftata? textura^a folida, mollia> fu-
perficieglabra, viroris diluti. Cofia media₃ i_padiceo nigra,
rubefcnes&nitens utrinque eminet? maxime in adveriā
parte. Flores in fummitate furculorum plures congregati
in geniculis proveniunt, in unoquoque tres quatuorve,
collum illorum album eft & bifolium , quorum unum
furre&umanguftius&cufpidatum, exterius coftula emi-
nente extuberans₅ albicans₅ venis rubro-purpureis interius
flriatum; alterum latius, exteriora verfus reflexum , an-
terioris in tres lingulas diffe&um •> exterius ex colli ventre
tribus coftis inftrudli, quarum media craflior & alteemini-
ens, & verfus lingulam folii Jatioris medium excurrit;
duas alias verfus lingulas laterales producuntur; interius
plicis ftriati, venis ex cofta media reticulatim pertexti, in-
terius lineis rubro purpureis anteriora verfus notati. In
medioduo ftamina habent alba, folio furrefto accumberitia,
apicibus flavodilutisnodulata. Stylus ex purpureo albcat,
ex oblonga gemmula , colon's atro-purpurei ac nitentis
emicans. Calyx quinque cufpidata habet foliola. In uno-
quoque calice femen deprehenditur unicum, oblongum,
nonnihil planum, in medio velut nervulo extuberante
obfitum, fufce tandem rubefcens. Foliorum fuccus ih

so

H O R T U S

mini *finapi* admixtuse metica virtute asthmati adverfatur.
Ex iisdem balneum conficitur, vel fi facculus ex illis con-
iiciatur, & admifto falenigro incalefcat, egregium in arti-
culari morbo remedium fuppeditat.

TAB. 42.

Folia coquentiam aliquam habent cum lutum Herb* -, flos cftgajcatus & labiatus) quisque mu-
cum fecundoprefbt; imenomina uft: *Persicaria Malabarica*, flore galeato & labiato monospermon.





Stellaria

stellata

var. glauca

Stellaria

stellata

var. glauca

AT) E L-O T) A G A M

T A E U L A X L I I I




Ingua Bramannum *Adulajfo*, planta eft ex genere virgultorum, nafcens in arenofis ad altitudinem ftatura humane. *Stipites* ex *radice* plures exfurgunt, qui, *ut tkrami* vc-tuffiores) cinerei , teneriores , vjrides 6c. geniculati, lignum albcat. *Folia* in ramis ad geniculos bina & bina exoriuntur , pediculis parvis, interius fulcatis, oblongo-angufta, in oris, raris, tenuifimiscrenulis fiflajin cufpidem oblongum coardlata, mollia>lenia, plana₅ virorisfufci, faporis amari. *Co/la* media in adverfaparteeminet *Flores* in petiolis viridibus, interius ftriatisj qui ex geniculis fupra originem foliorum emergunt ? plures congregati proveniunt, folii_s brevibus, cufpidatis fuccin<5ti; in totum candidi funt, conflantes *eollulo* brevi, lato6cnnonnihilpiano, acduobuspetalisoblongis, furre6lis ac mutuo obverfis, unum illorum cuipidatur&inmedioexcavatur, exterius *cojla* Candida,quas* inculpidemincurrir> obfitum? alterum planumcft, fu perne in tres lacinias feu *lingulas* profunderiffedlumj in medio *coftula*, quas verfus mediam lingulam excurrit, prx-ditumj transverfis venis reticulatim llriatum, criipatum & tranparens. *Stamina* in illis duo Candida? craffiola, *apicibus* magnis > viridibusdotata₅ folio cavo accumbentia; *Stylus* cum illis craffiolus? candid us ex *globulo* oblongo, viridiemergit,in *calice* fitus,quiviridi-dilutusj quinque oblongo, cufpidata habens folia. *Foliorum* engrgeia in *aftbmate* lunari, tuffi, fanguinis fputo atque atrophiae lu-cefcit^fi cum aqua calida trita propinentur; fi vero in manipulumcolligataaffeo lococalide` applicentur, velpro fuffimigio adhibeantur, in arthridite. Ex *filiis* fimul6c *radice* parum toftisfuccusextrahitur antiafthmaticus.

T A B . 43.

Defcriptio hujus plantx imperfefta eft, nam nee capfularum femimlium me femipura ulla fit msntio. Caulesquadratilunt, flosmonopeulosdiffimis.

S s

I'd'



مَعْلُوْفٌ فِي الْأَصْنَعِي

2 100 mm

Al-Mutanabbi
Mutanabbi

الْأَصْنَعِي

< A T v -K ^ % ^I r i-

T A B U L A X L I V .



Lanta hsec à Bramannibus vocatur *Rana*.
T^ddumari^ nāicens in arenofis. *Cantes* quadrati, geniculati , in exortu articulorum turgefcentes, pilis hirfutis, afperis obfiti.
Folia ad geniculos geminata oriuntur, iē mutuu decuffantia ₅ caules alatim fuccin* gentia, oblongo-angufta? cufpidata, tenuia? mollia₅ leniffimainftarhololérici, in oris pilis minutiflimis obfita. *Cofia* in adverfa parte eminens, rubelcit. *Flores* fummitibuscaulium cum fuis pediculis angulatis ₅ pilofisinfident? monopetali? diiFormes, in tres lingulas infecT:i, ac inillisex *rubronigricantes*, in collo in terius albicantes, ut & exterius, fed ibidem venis rubris ftriati, in dorfo colli exterius rubefcentes. *Stylus* furredlus ₅ planus ₅ quo folium anguftius refpicit albicans₅ in terius venis rubris ftriatus₅ *Capitukm* ipfiusnigrum eft&adlateracomplanatum₅ ad verticerncriftulaflavaobfitum₅ inferne pilis albicantibus barbatum. (*/eṁma* florū oblongae ₅ pilofas.

T A B . 44.

XmperfedehecccutprKcdensdefcripta , frudusenim & femina defunr.

T t



ପୁରୁଷ ପାତା

ମହାକାଳ

ପାତା

P A L L L U T U-7) A L L

T A B U L A X L V.



Ltcra ipecieseft, tortuofis &procumben* tibus furculis repens, arenofo gaudens folo. *tydtx albicans*, fibrolà, duriola 5 lignofa. *Caules quadrangulati*, geniculati, cymis inflexi, fuprageniculos extuberantes, viridi fufciores, glabri> pilis rarioibus fparfim obfiti. *Folia* bina & bina ex geniculis erumpunt, inpetiolis longiufculis ? interiusplanis, majora utilia prim^ ipeciei, pilis rarioibus confperfa, viridi fufciora. *Flores* una velut ferie 5 in fummitate furculorum appendent, anterioraverfusconverfi, ad pedunculorum exortum in nodulis ? folioliis breviffimis₃ cuipidatis comitati, campani formes, pedunculo breviffimo purpurafcente ex calice exfilientes, quafi pentapetali, duobus ad dorfum collipetalisbrevisbus, qu^eexterioraverfusrefledlunt, iè aperientes ,* ad ventrem tribus, quorum medium latius, ac velut lingula emicans, pr^ casteris altè purpurafcens; nervuloin medio obdupta, ad lingulam ftriataj in dorfo colli purpureo-rubro-fatura₅ in ventre dilutiora & albeP centia₅ pundluliique purpureo-rubris interius adiperfa. *Stamina* habent quatuor 5 albicantia 5 quorum apices bifidi, albefcentes,adfemutuoinflexa. *Stylus* albicans, tenuis, fuperne inflexus. *Calyx* quinque fólius eft. *CapfuU* fe minalesutilise *Vpu-Ttali*, fed majores, pilofas, exterius ob *femina* extuberantia lacunofe j quasfermeeadem, fed majora. Adhibetur hascplanta in concofione *Sails S^itrL*

T A B. 45.

Pertinethacplanta inter berbas flore monopetab, difformi-, fares (dice birAlfi, congenus *Digitalis* vi- d^eur & vocariöffet, *Digitalis ^msMMhricaJoi...r...l...*)

V v

TEE.

I

I

P f E - r u M B A.

T A B U L A X L V I .



Vtmio planta hsc à Bramannibus vocatur
nascens in loas humidis ad akitu Sm
trium crater pedum. *%adi_x* *Mk*ff*
b_{ropia} *TM* quadrati, geniculati, virides,
pilis c *lon* *Siu^c ^ obCm*, parum
lignosi. " " *8₈* *8₈* *obCm*, parum
geminata & ordine decussat
rigidiuscula, pilosa & aspera, versus interiora non nihil
clausa. *Costa* media crassa, viridi albicans, cum laterali-
bus utrinque eminet, maxime in *ad* *ad* *ad*
Flores in
geniculis
Capulæ
complanatae, lignosæ, virides, futura
lacunosa striatae, quinque constant foliolis oblongis, ap-
guftjffimis, lurre_a, sviridibus ac rubebentibus, candidis
mten^bbus, longifculis pilis obfita,. J J ? i n S S !
confident loculis, oblongo-rotunda, plana. In *bu*
ergo planta he pdp& eft, odoris gratihortenfk totum
Folia
c *s rabidi canis toxifi-*
c *caerulea. Succus specificum*
quid est in algidis febribus.

T A B . 46.

Proximo hcc planta accedit ad Lyflmachiam Virginianam,



Fig. 22

Stems
leaves

Flowers

Pods

Seed

O N A T U.

TABULA XLVII.



N'apu; Bramannicè *Tambido* tres daritiir species; unae jusnominis, altera *Vdli-Onapu* tertia *Tsjeria-Ompu*; planta eft hafcens in udis ad altitudinem duorum pedum. %adix cauli haud ab fimilis, aquea, in geniculis fibris albis veftita. *Cauliculi* rotundi ac geniculati, primum viridi-clari, dein rubicundopulchri, transparentibus venu lis albicantibus interftin&i, intus fungofi. *Folia* in geniculis bina &: bina, fe mutuo decuflant, oblongo-angufta, in margine denticulis acutis, rubefcentibus inciā, denfa, rigida, lenia, exterius viridi-diluta; r^/^inre & a parte, ubifufca funt, non nihil eminet. *Floret* qui duo vel tres ac etiam plures, petiolis rubicundis, languidis oriuntur, galeati, rofaceo-faturi, tetrapetali, illorum unum furrendum eft à poftica parte capituli petioli emicans, parvum & rotundiolum, in parte exteriori coloris rubri & nitentis, ac cofta viridi in medio obdudtum: duo oblongo-rotunda & uniformia, galeam integentia, qua cum capitulo petioli ex transverfo anguftis unguiculis adftri&a fiint, fingula unguiculam emittentia; quartum conftituit galea, eftque ilia oblonga, rofrata, membranacea, transparens, venulis intensè rubris, exterius fuocolore transparentibus, transverfim ftriata, à parte poftica unguiculam emittens. In capitulo petioli *fructus* refidet culloobtedus, fuperne rubens, in medio albicante farinaceo rore coniperfus, quinque fubtilibus rubris membranulis petiolo adftri&fus; funtque hi fru&us, qui floribus deciduis etiam cucullo nudantur, oblongo-rotundi, angufti, primū viridi obfcuri, dein viridi clari, non nihil transparentes, quinque ftriis, in quinque oblongos cancelllos diftin<ai> ex genere fenfilum, cancellis namque iuis ex Yy levi

leviattaſtuceu comprehione relaxatis, impetu in quinque fefeintorquentannulos, acfemina fuaexecutiunt? expli- cantqueie cuipides fuperiores cancellorum in latum? ad inferioresillorumpartesj quaepeſiolo annexſe funt; qui annuli ita rigefcunt,' uteorum cuſpidesnon nififrangendo a partibus inferioribus divelli poffint. In fruttibus prasterea *Stylus* reſtus *Me* erigit & albicans, cui femina adherent; queſeparvaſrotundiola₃ primo albicantia; poft nigerrima & nitentia.

TAB. 47-

Balsamine feminine genus eſt.





St. J. L.

24

25

26

27

V A L L I - O N A T U.

T A B U L A X L V I I I



Ltera ippecies *THo* à Bramannibus dicitur. *Ifyclix* eadem. *Caides* qui in ventofiores nodosextubrant viridiclarū funt, non nitentes, & in geniculis obfcurè rubefcentes, minus rotundi, in partibus fuperioribus quadfangulares, intus filamentis magis lignofis pertextij atque adeo duriores. *Folia* quse in nodis pctiolis interius planis bina, & paſſim novellis caſiculis comitata, proveniunt, oblongo-rotunda > majora, oris anterius in cuipidem anguſtumcontraſtis, inmargine multum incifa, in redta fblum parte pilis raris & furreddis oblita, minus folida. *Cofta* media in adverfa parte maximè, in recla alliquantiuum eminet. *Flores* iupra ex origine foiiorum oriuntur, duo, pluresve, veletiam lbitarii, infummitate caſium, etiam galeati, colons non ita faturi pro tempore magis minusve albicantes, maximè exteriori parte, heptapetalii, unum è poftica parte capituli fitum, latius eft & interius cochleato cavum, exterius vena viridi obduſtum, verfus alia quatuor inflexum, qu^ rotunda, paruna oblonga, galeam obtegentia, ac cum ilia transverbi capitulopetioliadfrida; fextum galeam conſtituit, quas albicans^ eft & minus oblongum > cum fuo calcari albicante^ verfus roſtrum crifpatum inflexo. (*/emm** rotundas & nonnihil cuſpidate. *FruBus* eodem modo cucullo obteguntur, fed illerubefcit, & fuperius slavefcit, dec^tero fimiles ac etiam fenfiles, fed illiprioris fpecie iangufhores. *Semina* rotunda funt, formâ nonnihil plſtna, primiūm Candida, pilis veftita.

T A B . 48.

Secundahxc *On^u* ippecies eft, fed ex orani parte major. Inter fpccies *Uſfamin** *Umh** proximi-


TSJE.

دہلی - بھارت

تھ

مکان

بہار، بنگال وہ
وہ جھری



T S F E R I A - O N A P T U.

T A B U L A X L I X .



Ertia species à Bramannibus vocatur *Tilo*.
 ^adix eadem- *Caules* magis rubefcunt. *Folia*
 interiores species media₅ pilis minutiffi-
 mis, maxime in venularum traſtibus obfitā,
 paffim duabus gemmulis viridi dilutis ob-
 dudta, fufciora. *F/ores* ad illos *Onapti* ac-
 edunt 5 ied pallidiores, tetrapetali; *galea* albicans eft,
 ^car viride; *capitukm* petioli fuperne flavum, *FruBtts-*
etiam feniiles» obJongiores & anguftiores, e quibus *fimina*
 permembranam interiore hinc inde valde protuberant.

T A B . 49.


Tertist hujus fpccici folia latiora, flores minores.



A aa

MAN-

Man-enaria L.



C M 4 N - O N A T U.

T A B U L A L.



St & *Onapu* minor, in omnibus conueniens, prater quod *caules* magis fint rufefcentes, *filia* majora 6c viridi-fulciora; *fiorcs* induobus foliis purpurdb ceu rubro-J casrulcfcentes; *fruUus* virides & venis viridi. fufcioribus ftriati₅ majores & oblongiores.

T A B . 50.

Plantahtecquamvi's non plenarie defcripra, coofccies eft prscedentium, exteris tantum in omnibus partibus minor.



B b b

BEL-

B E L L U T T A N A P U.

T A B U L A L I.

Peci	s	<i>Onopru</i> mi	g	1	f	3	T
sis.	-	ex albi-	-	s	-	-	-
stat.	is.	<i>Caules</i>	O	3	r?	g	X
t	1	albicat.	s	nit	<	g	Q
ex <		iculis novos			i	ri	it
<i>Fo</i>		caulicum			m	m	tent
		geniculicis caulum bin			a	ur	
		longo-angusta, in	3		d	a	


idem desinentia, ad pec **ti** *Tum strida,*
in oris minutis, venulis incisa, pilis n **x** *item in vena-*
rum trachu ob **st** *ra* *stra* *viridi-fusca est ac nitens, etiam*
in recta parte no **a** *51s* *albantibus, incurvis congregatis*
terni vel etiam q **51s** *petiolis oblon-*
giisculis, tenuib *ni, supra exortum foliorum erum-*
punt, candidi, unormes, tripeta; *duo, verbis que ca-*
pitulum petioli in flexum, latiora sunt, unguiculis trietis,
ac ibi punctulo rubro notata; *tertium interius cochleato*
cavum, exterius pilosum ac subviride **cens**; *ad ejus exor-*
tum sacculus retiformis subviridis, calcar viridi-albicans,
in curvum dimitens, a petiolo versus duo folia transversa
incurvatum, anterior parte folia ad mutum contractum
luffulciens. *Gemmæ virides*, *in cuspide strigae, ad la-*
tera complanata; petioli capitulum viride, umbilico flavo
in vertice dotatum. *Fraga* *qui petioli viridi albicanti-*
bus, demissis appendent **0** *pitula lunt oblonga, viridia,*
latu, ut illa prioris speciei.

T A * »I.

Hec minor species *Onopru*, cum illa **1** coarent.

O n

T I L O.





ج. س. م.

1800

ج. س. م.

1800

ج. س. م.

TIL O-O N A P O : cœu NJ) TE N G A.

T A B U L A L I I .



Lanta hsec duum triuñive pedum nafcitur in arenofis: *Cauñes* viridkiiluti, aquei. *Folia* oblongo-angufta, cuipidata?in orisar£tècre. nata₅tenuia₃ fufca₅in reila parte pilis obfefla, artiara. *Flores* plures simul fupra exortum foliorum confpiciuntur tetrapetali; uno furrecto, obtuiè cuspidato, interiora verfus nonnihil claufo, ex albofubrubefcente; duobus rotundiolis?unguiculis ftridtis, rofacco faturis, inferius rubefcentibus, transverfim fupra quartum expands, quod galeatum, ex adverfo petali furrefli fitutrij albicansac dilutiffimiruboris, calcaremittit viride, a'petiolo crilpatum. *Capitulum* in medio emicat rotceo apice obte&um, germen *fructus*, ex genere fcnfilium, qui ex attadtu fecum impetu laxant & femina excutiunt, ut illi *Tiloy* à quibus tamen differt, quod denfis.fint albicantibusque pilis pbfiti.

T A B . 52.

Ad priores hxc planta pertinet *Onapu* fpecies, f Balfamina: feminiE haberij hanc vcro ego atqueillarum confspecies eft, paucotamen cum j repuco pro *Perficaria filiquo*f* *Swtca*, flare tofacto n*dj* *gr* *mine* *ficaplularum* lemnalium, port at- I *lore*, *fructupilo*fa. *ta* *ctum*, *diffra* *ctio* *ff* *et* *jomnes* pro eonfpiciebus I

Ddd



Lacapu
Coch-
on

دہلی
کھنڈی

K A K A-? U.

T A B U L A L.I.I.I.



Ingua Bramannum *Caela-*, plants humilis .. repens , arenofa 6c uda amans Joca. *%adix* fibrofa, albicans. *Caihculi* geniculati hinc inde ex geniculis radīcu* las fpargunt tenues, albicantes> teneri funt, quadrangulati , pilofi, in angulis pilis niagisconipicuisobfiti, intushumidi. *Folia* ordinenon-nihil decuffato bina & bina* ? ex geniculis, petiolis brevibus, teneris, pilofis 5 interius planis, fulcatis, glabris, nitentibusproveniunt* parvula> cufpidata, inorisdenticulis majufculis ex sequo ferratim inciia 5 texturā tenuia inollia, superfielenia?. utraque parte pilofa; *cojlulis* in adverfa parte ftriata. *Flores* in geniculis caulum erumpunt, monopetali, campaniformes, quadripartiti, coloris ex parte venufti? partim purpureo-rubro-fufci, vel albo rubri dilute purpurafcentis. *Stamina* in illis quatuor purpureo-rubro-fufca, bina ac bina arcuatim ad iē invicem inflexa, fingula apicibus, qua colli dorfum repiciunt candidis & nitentibus, quā ventrem c^rulseo-dilutis inftru6ta; riorum duo fupereminent 5 colloventris inferta, ubi fingula unguiculo tenui 5 purpureo-rubro-fufco concomitantur; duo depreiliora funt, dorfo campan^ inierta. *Stylus* candidus eft ac nitens > dorfo proximè accumbens, fupern& quafi in furculum fiiTus. *Calyx* oblongus 5 viridfs quinque angulis eminet 5 fupernequinquecupidesfiflus) adlatera profiil iius hians. Planta cum *Sandalo*, *Cariopbillis*, *U\jice* *%ihsc a, & aquarofarum* contrita? linimentum prasbet *exanther*. iiticum* *Foliorum fuccus* cum *Saccharo* gonorrhæam fiftit.

T A B . 53.

Flos eft <i>MonQptalusydifformis</i> ; fcmina & capfula;	<i>Heder&temfliis</i> speciesrecenferi, & pro confpc ^z
içminalcs non funt defcripta, atq[ue] idco reftium	cic <i>Afarina</i> five <i>Hederua faxtilis Lobeli</i> , prout cx
«ellis judicium vix ferri poteil, alioquin inter	dclincationcdcfumitur, haberि poſler.



جعفر

O. J. D.

S C H I T-E L U.

T A B U L A L I V .



Lu dusefuntfspecies, *Schit.Elu* & *Car.Elu*, prior Bramannibus *T>avo-Tilo*'e dicitur, planta eft altitudine quatuor vel quinque pedtim, arenofum amans folum. %*tdix* parva, fibris capillaceis veftita. *Caulis* ex radice exfurgit unicusj in raroſ caulincoſ divifus, quivel bini vel folitarii è nodulis prodeunt, quadrangulati, in unoquoquelaterefulcoſtriatiſ pilofi, virididiluti. *Folia* queſe petiolismediocriterlongis, viridi claris, pilofis; ftriatis> hincindecaulibusinfident, oblongo-rotundafunt, cufpi-data, pilola, nervis in adveriā parte eminentibus ftriata, in orismagisminusvècrenata; in inferiori parte caulum folia quasdam majora? iimplici numero, confpiciuntur. *Flores* ex origine foliorum oriuntur, cucullati, oblongij monopetali, fuperius quinque partiti, tenuiter incifi, la-Duginofi, prsecipue in exteriori fuperficie? candidi, in odores. *Stamina* habent quatuor, tenuia, rotunda •> Can-dida; flavisacfurreclisapicibusdotata; *fiylus* tenuifeſt& bifurcatus. *Calyx* parvus, furreiſlus, in quinque anguſta & cufpidata foliola incifus. (*jemce* oblongo-rotundx, ad primiim ex albo ſubflav^e, poſtea magis albefcentes •> quinque angulis velut *Helix* formam repraſtentantes. *Siliqu* ^ ieminibus plense, ubi viſferme, ubi folia funt? petiolis brevibus caulincoſinfident, reſte, oblongs, quadrilate^ rat<> duobus lateribus adverſis planiores •> omnes vero fulcoſtriatiſ, virides, pilofe, in quatuor capfulas feminales, interfeſimentisliſnofisdiſtindte. *Semina* in fingulis capfulis feriatim ac fimplici ordine confident, umbilicis columnute medkaffixa, plana, rotundiola, ad umbilicum cufpidata, lateribus fuis planis in longitudine capfularum di/pofita, coloris, cum ficca funt, ex rubro fpadiceofufci, intus albicantis, faporisfubamari. *Oleum* ex fcminibus extra-

Fff

ctum

£lum₅ quodetiam oleum *Sirgdim* vocantj pro quib[if.
cunque pene exhibetur doloribus₅ eoque in lotione r[if.
quentiffime utuntur Indigense ? ad perungendum corpus*
& quotiefcunque dolore aliquo ineffontur) additis in de
cocto nonnullis fimplicibus: cum z*Jmphione* furntut
mortem inducit, ejus antidotum eft fuccus radicis *Tinted*
Lufitanice *C^{ove}A* vel fuccus ex fru&u *TSarttdem* diclo
Lufitanice *Uringella*.

T A B , 54.

Planta hxc *Sefdimm* rectrum eft, *S. Pin.* Al- bum, *femenenimj* *UilludSefamn* *vulgaris, album*
pious *d:fUmti.Zitfi.* vocatillam *\$emffim|* Ma- eft.
UccenTcs Bid-^jjMttietth, quod valet. *Btk.J*mal-*





C A R - E L U.

T A B U L A L V.



Ecunda species *Car-Elu* > dicitur; lingua Bramannum, *Caro-Tilu*, cum prima maxime con venit, nifi quod paulo akius exfurgat. *Caules* etiam nonnihil crafliores funt; *folia*, *viores*, *femina*, majora, color magis fuičusj iapor fubamarus. *Flores* comefti oculorum vitiis medentur; cum *fruttibus* triti & *butyro* in formam emplaftri redadli, apoftemata ad maturitatem perducunt. *Oleum* ex *feminibus* extra&um .(*Oleum Sirgelim* nomine Venit) humoribus phlegmaticis ex flatibus conductis fi nempe corpus cum illo perungatur; & fi Malabarienfib⁹ habenda fides, pinguiores extenuat ac iiiigcentos obas- fos reddit; oculorum caliginem pellit, vifum debiliorem confortat > capiti nempe illito; ex eodem etiam unguenta conficiuntur pro vulneribus & ulceribus: cum *Amphwne* fumtum mortem infert: *Semen* contritum & cum fucco (*jjennhm* exhibitum > vertiginem tollit: eadem bene lota & excorticata variis modi's condiuntur.

T A B. 55.


CarElu secunda species eft *Sefami*; à nemine, *ferrato, fiore ffiajcre, fem'me nigricatite* vocari potest; quod fciam, ante hac memoratutn 5 ex omni partemajus, quare *sefemum Indicum, folio amplio,* *Malaccenfes dicant Bidzjate Bitém, idque lore feminis.*

Carex Ciliolata L.

காரே கிலோலி

காரே கிலோலி

தாவி 200



C tA R A-C A fi IN AM.

T A B U L A L V I .



Ecunda species eft *Qaniram*; Bramannicè *IBoin-Caro* vocatur, planta elt akitudine duum pedum? arenofa amans, florens temporc pluviofo. *tydix* plurimis conftat fibris lignofis) capillatis, nigricantibus, cujus cortex maarus. *Caules* quivel fimplices, velduo tresve ex radice exfurgunt > virides, quadrangulati, in nodosdiftinfti, cauliculos paffim binos, qui le mutuodecujffant, einittenteS. *Folia* è nodis caulum petiolis brevibus bina&binaproveniunt, oblongo-angufta, inferius contraftiora, tenuia, mollia, fiipdrficie plana, coloris viridi-fufci,faporisamariflimi; nervulus in utraque parte eminetj magis autem in adverla. *Flores* in petiolistenuibus, viridibus, quadrangulatis, interdum etiam nonnihil ramofis videntur, qui è caulum nodis, fupra originem foliorum, paffim bini&bini ex adverfo oriuntur, n̄fque pedunculis brevibus, tenuiffimis, hinc inde, pr^fertim in fuperioribus parvibus adherent: bipetali funt, u-Ho angufto & reflexo, in totiim albo; altero latiori, furreflo, nonnihil reflexo, ac in tres cufpides ^quèaltè emicantesdivifo, quodetiarrl album, & circa oras duobus, rubro obfcuris radiis, qui ad duos extremos cufpides cxcurrunt, variegatum eft, & in medio macula rubra notatum. *Calyx* laxus eft, in quinque angufta & pilofa foha divifus, cui tenuiori pede infident. *Stamen* habent longiufulcum, albicans& pilofum. *Style* tenuifeft& ahquantulum rubefcens, ex germine capfularum feminalium Sh *Capful*feminale*; angufta, quadrangula: tra ens tæ, compressæ, phdS&duriol^, in longum fulcoftnate, intersepimento afulcis excurrerit, induasquafi came rulas feminales oblongo anguftas diftintf*. *Semina* qua, in singulis cameruB s tria repeñuntur quatuorve, parum ob-

H h h

bloangi, prtriM > vir idig, dcin al bican t fa, ficca ru ffa ceuf:
 lavo rufadreflm* fublpra- TotaplantacuniLnfusion'
 * roaccrit* moriu i Vrulento ierpcntLs, Co pra Capcl la >
 dict- njedttur,cptanempe, quod & per unia prafat.

T A B , 56,

Cannas, I (quamvis hoc ex descriptione non patet) radicis
 plantas. Cannas dicta, prout
 I*MB «) te er, jimiailJ in gnwrr KrptriiHrj, KBtt «» Cannas omnes plantae panifices sunt
 cum eam filique, se uijamie. nil tS-
 lineationem vericalis videtur



• Typha - pugn

• Typha - pugn

شایعه نویسنده

M A L A B A R I C U S, i n

TSJANG A-TUSToIM.

T A B U L A LVII.



Lantula eft naſcens in arenolis. *%adix* viridi-albicans? acalbisveftita fibris. *Caules* craſ fioli 5 quadrangulares₃ fuperne ftridliores* geniculati₃ ex geniculis novos caulinios emittentes, ac in illis foliis geminatis₅ it mutuo decuflantes veftiti; in totum aquofi. *Folia* parvulafunt? oblongo rotunda, tenuia; *co/iula* media cum duabus latreralibus in adverfa parte ftriata. *Flores* qui in petiolis tenuibus₅ longioribus fupra exortum foliorum ègeniculisiuntur, caerulei fint? papilionacei, bipartiti, inferius duabus casruleo*purpleo-fufcis maculis notati; *Jlamina* in illis duo albicantia, ad fe mutuò inflexa* apicibus albicantibus inftrudla; *C^ab^x* quadrifolius eft. Tota planta *butyro* incočla gonorrhasse adverfatur, vel etiam cum *Calamo aromatico*, *Pipere longo* infantibus propinatur adverfus Epilepfiam, viridi diarrhasa comitata: *radix*, cum fucco ex foliis expreſſo? & *Sandalorubro*, trita₅oculorumvitioj Malabaribus *EdJaTadelle* diſlomedetur.

T A B . 57.

Species *GentianilU hxc* plantula videtur.

I i i

K&-

Katú-peé-tojánga-puspám lat.

کاتو پي تو جانگا پوسپام

کاتو پي تو جانگا پوسپام

كَاتُوْ جِنَانْجَةَ يُونْتِيَّاْمْ Arab.



K A T U - P E E - T S J A N G A - P U S P A M.

T A B U L A L V I I t



Lantula eft arenofo gaudeñs folo. *S_{adix}* albicans > fibroia. *Caules* virides, quadran-
gulati, geniculati, fulco in oppofitis lateri-
bus, afternato in internodiis ordine, ftriati.
Folia ingeniculis ad latus planum inferioris
internodii geminata videntur, oblongo-an-
gufta, tenuia, leriia, glabra, in oris acutis ornatacufpi-
dibus. *t°ft?* media in adverfaparteeminet. *F lores* plures
congregati in vertice caulum proveniunt * papilionacei,
albo purpurei, inferius rubro-purpurei. *Stamina* in iis
duo furredla, albicantia, ad fe mutuo inflexa, albicanti-
bus apicibus inftru6la; *flylus* albicans > fuperiori parte com-
planatus, ex orificio calycis exfurgit, qui oblongus eft Sc
quinque cufpidatus*. *CapJuUfeminales* oblongas, *virides* i
fubruffeſcentes, in illis numerofa, minuta ? arenace^
fernina, primiim viridi albicantia, communi viridi ftyluld
adh^rentia.

T A B . 58.

Folia & flores hujus herbulas, illis *Tcucrīi* penç affimilantiir.

K k k

TEE.

C/b.

tffit/l/ <* - purp a m > at.

வாய்வு மல்

காப்பும் ஓம்.

R

a ij>- 9
(s?) جانغبو

cP E E-T f A 5^_g A.T V L < P A M.

T A B U L A L I X .



Repola amat & uda, cum priori in omnibus fere convenit, pñeter quod *john* minus longaj ktiora fint 6c rotunda; *fiores* ifdem led infra flavefcentes; *calyx* quadri-partitus, *lamina* in illis quatuorcum apicibus ad fe invicem intorta. (*apftdee feminaks* oblongo rotunda, numerofa includentes *femim*.

T A B . 59,

Plintutnm bane P<-TiJ4n>J- P~~g~~^{on} cindem *fitKiiufiitii f*eiiniunitii**, vocatm, nullnsdubiin, clif(c*ciimillil,cjUi:nb* Vitama in Lit. *Ssu.Mti*. excepcoquod iarts aliquomodo tolrc *mitin:u::: Ut₂ E,u. Tiuiimni JsmijU/mum ffinAim, Veniuu*

LIJ

N E



NE L AM-TzAR END cA.

T A B U L A L X



Lanta humiliseft; furculisdcumbentibus
fupra terram expansis. %adix fibrofa₅ al-
bicartSj intus Jignofa, odoris grati ac acn's.
^ww/tenues, Jignofi} viridi corticc vctiti.
Folia furculis infidentj oblongo angulta,
cufpidataj inoris tcnuiflimecrcnata, mol-
liaj Ienia> *fuita* furda. *Co/la* media utrimque eminet*
laterales in adverJā tantiūm parte. *Fl̄kes* pediculfs te<
nuibusj viridibus & verfus inferiora infiexisj iupracxortum
foliorym videntu'r -, monopctali, rofaceo-faturi > *coftam*
in adverfa partf habent medium > in rc6ta tres rubicundio-
res venulas, necnon duas foliofasac falcatas uncinulas ē
calycc emergentesjquiE Candidas ac venisin longum-ftriatar,
cuipiderofaceoiaturo. *Calyx* quinquefolius, culpadiis,
in cuius raedio, *lamina* conlpiciuntur albicantia? lurrepta,
uncinulis obtcēta, quasin ruffos, croceos ac pianos defi-
nunt apices, in quorum medioviridis *gkbulfm flyiulum* al-
bicantem ē verticeemittit. (*fl̄obilfemmales*, cum culpide
albicante in vertice, forma nonnihil trilateralis funt, cortex
cartilaginofus, ftriatus, prasfertimin rnedio, **ubiglobuJl**
fe laxant, qui primum **viridi-diluti**» dcin ruffeicents.
Semina in fe continent odo> oblonga rotunda, aqueocan-
dida> nitentia.

T A B . 60.

Cum *Hillihiat* phnnihharcquandam Jlmilitudinerahaberei

Katu-vistna-clandi L.

குஞ்சிலூஷ் mal.

خَاطُوْ بِشْنُوْ فَحَلْ رَضَيْ arab.



Bridges

K JTV'F 1ST U^A-C L A ^ Z) /.

T A B U L A L X L



Lanta eftnafcensinarenofis. *fydix* ruffa* fibrofa & lignofa. *fymi* tenues, rotundi, Jignofi, virides, pilis obfiti. *Folia* pedunculisparvis, viridi-fufcis, crafSolis₃ ramis infident, oblongo rotunda ? pilis tenuiter obfita₃ rnollia ? flaccida > lenia, viroriſ laturi. *Gfa* media in adverfa parte eminetjillorum fapor fubacris, odor nullus. *Flores* furculis transverfis infident papilionacei tripetali, quorum duo rofaceo purpurei 5 tertium folium ungulatum eft& cseruleum, in dorfo ftaminibus purpureo c^aruleis bifurcatis ceu trifurcatis criftatim dota- turn. *Stilus* theciformis eft, inferne purpureo ruber* fuperneintenuiffima, albicantia, longiuſcula filamenta, queſe fingula gemmulis albis nodulata funt ? fifTus; filamen- tumtransmittensalbicans, incurvum, capituloj longo, oeruleo inftrucSum. *C^ab^x* quadrifolius eft: quorum duo majora angulata? duo minora, majoribus dccuflatim interpofita.

T A B . 61.

Imperfetta defcriptio eft ₅ quarc de bac planta nen ex a:qao dijudicari potefit • figura cnim frudus non memoratnr.

Nna

MAN.



سید جعفر

الطباطبائی

مکتبہ طباطبائی

جی ۴۵

C MANJA-KURINL

T A B U L A LXII.

BRamannicè *nAbuli*, nafcitur *in* arenofis. %*ami* rotundi, geniculati, cortice viridi fufco, glabro; ligno albicante •> in cujus medio medulla. *Folk* geniculis > petiolis oblongis 5 viridi dilutis, interius planis ac nonnihil alatis? paffim quaterna? furredta infident^ oblongo rotunda₅ tenuia, mollia, inadverfaparte, plurimis coftulis annularibus 5 quas ex *cojla* media exeunt acvalde eminentinfrud:a; inredlaparte? velutrugis, in traſl:u coftarum, ftriata; faporisfubacris&fubdulcis?illi Raphanorum nonnihil referenda. *Flores* in faftigis furculorum, fpicatim videntur, in petiolis qui recla ex eo* rum vertice prodeunt, ceu qui transverfi ac laterales, fiipra exortum foliorum ex geniculis erumpunt, monopetali* intreslaciniasfupernediftindi, colon's in totum aurancei floridi; collo oblongo flavefcente ex *calyce* emicantes; *fiaminuk* habentquatuor₅ flavaoblonga. *CapfuUjeminales* oblongse₅ culpidata^j lignofe, parumcomplanatfe,infe^ rius ftridliores ? primò virides 5 dein albicantes, foliolis parvis obte<3:2B, quse primum in calyce oriuntur > dum ad* huc ex fuperiori parte flores ex gemmis erumpunk

T A B . 62.

ManJ+Kmih plants ett anoraala feu fui generis; cum ullis cnim nullam habeot convehkmkrirt

Ooo

T A .

Tali-pallū L.

ତାଳି ପାଲୁ ମାଲ.

طَلَّالِيْ بُلَلِيْ

Arab.



T A L I - ? U L L U .

T A B U L A · L X I I I



Renofo gaudetfolo. *'Radix albicans, fibrofa.*
'Caules virides, rotundi? geniculati, aquei,
'foliis grami formibus 5 quas ex geniculis
 I prodeunt, veftiti *Flores* tripetali lun₁ 5 ro-
 tundioli 5 cupidati, cseruleo-purpurei, fcin-
 tillantes. *Stamina* in illis quinque 9 exalbo
 c^rulefcentia 5 oblongis casruleis pilis barbata; tria tenuiora
 flavis apicibus nodulata 5 reliqua duo craffiora > croceis.
Calyx trifolius eft. (*femmafemales*, viridi dilutee 3 nitentes > in vertice cufpis eminet nigricans > de castero convexi,
 trilaterati, in laterum medio leviter extuberantes > atque
 ibi fe aperientes ac relaxantes. *Semina* in illis continentur
 lèx 5 interfepimentis deftin&a, trilaterales 9 uno convexo,
 duobusplanis > fuperficie cinerea ac fcabra. Planta cum
 lafleo *Zh(ucif Indica* fucco capiti fuperfufa capillos pro-
 movet «\ . *Succus* cum oleo *Tzjt-Elu* deco&um *lepr**, Ma-
 labaribus BemCuttam₃ Carin-Cuttamj adverfatur.

T A B . 63.

pcrtinct inter confpecies <i>Tbalangii vrlgmiani</i> . A». 1688. plantamhancexfemine, abAuthore noftro fub nomine <i>Rak*lk*z.«J BtngaUnfis</i> mihi	transmiflb, produftam vidi, atque eodem anno in horto noftro Medico floruir, menfe Augufto.
--	--



Digitized by
Digitized by
Digitized by

U T U-T> A L I

T A B U L A LXIV.



Ux sunt species una ejusdem nominis, altera *FaMa-Vpn-Dali*. Prior plantula humilis est, furculis tortuofis procumbens, ac con volvuli infarrepens. *Ifyrix lignosa?* fibrofa, albicans, *fcaules* tenues? viridi fufci, quadragulati, durioli, geniculis interftindti, supra geniculos in singulorum internodiorum principio, ventriculo turgentesj cymisinflexi? capillis albicantibus tenuiter obfiti. *Folia*, guas in geniculis in longis, viridi dilutisj capillatis, interiusplanispetiolis> bina&binaproveniunt, oblongo rotunda sunt & cuspidata, capillis albicantibusobfita, tenuia, mollia₅ fubaipera, viridi fufca. *Co/la* media cum nervulis annularibus in adveria parte eminet. *Flores* in geniculis ad exortum foliorqm proveniunt, catnpani formes, monopetali, quinque partiti, nervis percollum exterius ftriati ₃ ex albo casrulei dilutiiSmi; (*lamina* habent quatuor albicantia, quorum duo altius emicant 5 furrettis ac albicantibus apicibus dotata. *Stylus* albicat, invertice intortus. *Calyx* quinque anguftiflimisj pilofis) oblongis foliis₅ florum collum excipit, qui ad calycis exortum 5 duobusfoliisminoribuscomitantur, fine ulloodore. (*apfuU feminales* oblongas, futurisftriat^jin orisnempe, induosloculosdiftincT:^, in medio laterum ubife relaxant₅ tenuiori futura exaratse; primiim virides &capillafce; deinlignose, albefcentes. *Semina* valdecomprella, tenuia, orbiculata,cinerea,inoris]imbuloparum extuberante & fubrufo circumdata₅ in singulis capsulis paflim quatuor fcutellatim confita, interfepimento medio accumbentia. Foliorum[^] *fuccus* (ut&folia) inco&ione falis adhibetur. *Folia cum file* coclra&in fuccum compreffa? humorum cachexiam corrigunt.

T A B . 64.

Quaravisdux hoc loco species *Upu-Da:t* me- fe habent convenientiam, nee quoad formam, nee morentur, quarum hxc prior 5 altera vero tabula quoad genus satque ideo hte fui generis eft.
45. hujus partis fitdefcripta, nullatn tamen inter

Qq q

SO-

SONBR./ILc^.

T A B U L A I X V .



Afcitur infrenofis.' UUUT ajbicatms) ro*
funds geniculati, pilis oblongiiTiculis cbfiti>
intusaqud. Fd/* ineenictifos petiole ob*
longjia, lurreibs, albanticibus.intcriusJlil-
cati•i, gniimta & ordjnc dteuffato video tur>
dilangorotundii, pilis oblongiSt nonnifnl
hirfutiaobtai ox a>jli media crafRi, toftulis craflis cmi>
nentibus? tJLixoblk|uo duitu anterior3 vt-rius excurrunti
itriataj in totuni vanegatt s cstcrius ad "mditim ufquoma-
ximj parte rubro-parpurai uti & in-i, c coftis; interius
ex adveria viridi ^uTea, ad Jpadjeeuin very ;centia; dc extero
virons vividf, punttuJis rorund^i ii libis , numcrofw, ex
ijuurum iingulii iDteriu.9i unicLLJcmitatpiJus^inter/pcrfa.
fhrt 1 in Jummftate cauliim in pcdunculis n 1 bo-pur puraf
ccntibm * pil ofis, pi urcs coigre^iti, ad icx, Jcptern, pJ us
niinm proveniunt₅ triptaii, oblongf j ciipidutt >incerfua
roiacco ktur'i * o.yteriufi roficeo dili it lores 1 pilis riris obfiri.
Stimtiu in ill is tri:i 1 roiicca 1 npidbus Jurrcclis» culpidatisi
flavis , dutJtJ. Siifitr oblon ^{gus, re} ^J-TIR in media ex
calyce LKluryit t t|dorifico ^{cit protus}"JLii jc [hum bre*
vium tuipiJiitorum foJiorui.

T A B . 65.

Planta huc proxime accedit ad Pichonaria monilifera, & non aliter dista pertinet, perfring
visari Pichonaria fico monile & Drosa fico, fico, certain, refecta fico.

R r r

Cela

Kalu-polapen Linn.
കാലുപലപൻ mal.
Corosjerá
സ്രവനം Ban
قَالُوْتَلَكَ عَرَبَ



KALU-TO LATEK

T A B U L A L X V I .



Lantula humiliseft/pitharnam vix excedens* arenofa a mans & petrofa. %adix nigricans* (lignofa, rigida, fibris veftta. Surculi craijufcnli) virides, quadrati, in fingulo latere profunde ftriati; brevibus albicantibus pilis obfiti. FoliA in furculis abique pcdiculis gcominata prodeunt > le mutuo decuffantia j parva, oblongo-angufta , craffiufcula, rigida > pilis veftta , in advcrfa parte unica coftukj fine ullis nervulis lateralibus , obdutta ; ubi detrahuntur , fimul pars corticis , inftar petioli folio ad hffirentis,deglobit. Flores ubique ferme Jupra fingulorum foliorum exortum furrefJ:im prodeunt, ipfis furculis accumbentes, conftantque collo albicante iibviridi & piloib, quo ex calyce emicant, ac quatuor candidifliim foliis difformibus, in transverfum ad collum. iitis; unum illorum breviffimum aft latiuiulum , ora rotundiufcula₅ & inmediobrevi cufpideemineiis, exte-riora verfus crifpatum, furculo > cui flores adhasrent, ob-verfum; tria oblongiora, & circa adverfara partem folii brevioris extenfa , oblongo rotunda, (jdyx oblongus₅ viridis > ad orificium quinque fuſcorum cuſpidum; in quo prodeunt Capfuſfeminatei oblongae, intramembranoum corticem Jiamim includentes numcrofe, minuta, arenu-laruminttar, ftyluloviridiintermedio circumpoſta, pri-inum viridiaj dein casruleo-nigricantia , & ac perficum vergentia. Datur&species CaluTolafi, ineotantuma priori differens» quod flores ex flavo virides fint.

T A B . 66.

Siii gcnuseft, &iioroin>ripoffct lUntxUfdimawi: 4^{mm} fd'apl^{ofo}, fiore monopetalo difformi Malakaiu.

Sss

KO*



پٹی جاہنی

کھنڈلہ

G. M. Holmes

دہلی

G. M. Holmes

K OT>ATS J E R L

T A B U L A LXVII.



lEtrofaamat. *Cuhs* tenues, rotundi, viridesj lignofi. *Folia* caulibus hinc inde infident, pblongo rotunda > craggiola, superficie plana 6t glabra, in adverfaparte *cojula* viridi obdufta, in cujus tra&u interius ftriatafunt, odoris cum teruntur fuavis. *Floret* fupra exortumfoliorumbini, terniifimul, in uno parvo comniuni petiolo proveniunt, papileonacei, candidi ? difformes ; unum illorum ungulatum eft? ad cufpidem dilute casrulefcens, ac infra verticem in dorfo crifta ceu plumula flaminea fubcasrulefcente donatum. *Stilus* candidus eft > fuperne in filamenta tenuia, apicibus parvis nodulata, fiffus, ac in medio filamentum candidum oblongius, quod annulatim crilpatur > recondens. *Calyx* quinque folius eft & viridis₅ quorum duo oblongiora₃ tria minora? unum inter duo folia oblongiora, intermedio folio ungulatoflorisobverfum eft; *Semina* oblongo-rotunda, plana, fcabra, ruffo-fufca. *Folia* cum *Ory^e* lotura lasvigata, vel etiam *cléo* incodta&exhibitaabdominis cruciatus mitigate, menfes promovent.

T A B . 67.

Confpcicmnonhabcthseeplantula, vocaripotcf: *rm*kc4 filiitfimlisflanu, jlore «|hog*U«to & Ubiato, femine ohlwga, rot undo, rujfufco color t.*

T 11

CO:

Corosinam L.

கொரோசினம் Mal.

Corosinam

كُوروسِينَامَيْهَمَّ لَهَامَ

كُوراجِنِيَعَهُ عَذَابَ



C O R O S I N c* M.

T A B U L A L X V I I L



Lantula bumilis -> Ipithamse longitudincm
ivixexcedens. 'Radix fibrofa? colons crocei
faturrimi, manum ex atta&u rubore tingens,
intus ligneo filamento. (*jiuliculi* virides,
rigidiufculi > pilis longiufculis denfe obfiti.
Folia quas caulinulos denfa ferieinveftiuntj
geminata proveniunt & fe mutuo decuflantia₅ oblongo an-
gufta 5 valde craffa ac pilofa, viroris in refta parte nonnihil
nitentis, inadverfafurdi, *cqfta*, re<3:am ftrians, obdu&a.
Flores monopetali funt, campani formes > candidi > quin-
quepartiti? intuspilofi, in coUo abutraque parterubro
perfici coloris. *Stamina* in illis quatuor, craffis albicanti-
bus apicibus nodulatajpilis candidis barbatim veftita: *Stilus*
viridis eft & imberbis; odor nullus. *CaffuU Jeminales* in
vaginulis florum pilofis fuccedunt floribus deciduis; ob-
longx rotundas funt > nonnihil cuſpidatas₃ lyrisinlongum
ftriat^e ? in lyrarum träftibus interius futura eminente in-
frudte> ubificcas funt cortex cartilaginofus eft j albicans,
fcaber. *Semina* in illis continentur denfa ac minutiiima,
oblonga, una parte acuminata₅ & cumcuſpideftilulome-
dio piano ac membranaceo affixti, primiim viridia , dein
flavefcientia, tandem furufa. Refpirandi difficultas tol-
litur, ubi plants *hujis-Succus* cum *Curcuma* prascordiis il-
linitur.

T A B . 68.

Planta lisec anoialaeft & fui generis; fineullo> quod fciam, congenere.

207

نیزه

نیزه

نیزه



T U l-C O L L f.

T A B U L A L X I I



Ux hujus plants funt ipēcies, *Tuhcolli* & *Cdtu-Tul-Qilli*. Prior lingua Bramannum *Dadumāri* vocatur, fuffrutex eft > ex genere virgultorum, altitudine trium quatuorve pedum, nafcensinarenofis. ^{mc}*S^dix* lignofa & fibrofa > cortice iiibflavo. *Stiptes* ut 6c *rami* geniculati? teneriores virides , vetuftiores cinerei* corculo intus albicante. *Folia* brevibus, viridibus > & interius planis proveniunt petiolis, oblonga? culpidata^ tenuia j in adverfa parte maxime circa coftulas villofis pilis tenuiter obfita. *Co/la* media cum lateralibus utrimque eminetj maxime autem in adverfa parte. *Flores* in petiolis ramofis qui in vertice emicant, vel exgeniculis ad latus fli- pitum erumpunt, plures congregati vifuntur > longiufcul^ virides, ftriati? pilofi, angufto collo, oblongo-anguftij candidi, tetrapetali; horum triamagis uniformiafunctj acad collum è transverfo alatim protenfà; medium paulò latius eft > ac in una parte ad collum rubris pundiulis confperfum; quartumexadverfoiftiusfitum^ anguftiflSmumeft? inftar plumule criftatim ere<3:um, in apice bicufpidatum. *Stamina* habent duo albicantia, orificio fuperiori colli inferta, rubro- fufcis apicibus dotata; cumiis *ftylus* albicans fuperne bi- cufpidatusex *calyjis* orificio emergit; quiparvus> viridi- fufcus, fublaxus, quinque folius; in quo clauio unum reperitur *femen* oblongum -> rotundum, ex cuſpide filamentum ejtciens albicans ac ruffefcens. *Folia* lasvigata impetiginem? pforam fcabiemque tollunt.



N E L I T U.

T A B U L A . L X X .

Udo & paludofo gaudet folo , prasfertin ubi *Oryfy* nafcitur, quamconvolvoliinftar,finuofo cauleampleflit. *%adix* tenuis, albicans₅ ex brevibus capillatis confitans fibris; unicumtan* turn cmittens caulem, in nulos alios fe dividentem, qui rotundus ? tenuis, viridis, nullis foliis veftitus. *Floret* in caule hinc inde folitarii j petiolis planis > uncialibus ? viridi dilutis, quia dexortumparvocufpidato fuccinguntur folio? proveniunt, casrulei, bipetali, difFormes; illorum unum retiformiter cavum, cum orificio inferiora refpi- ciens ; alterum rotundum, furre&um, folio retiformi obverfum; quodretiforme cseruleumeft, in fundoalbi- cans, caeruleis purpurafcentibus venis in longum ftriatum; alterum in totum cseruleum purpurafcens ; cornu illius pendulum albicat&nitet, incufpide autem ex viridi cas- rulefcit *Stamina* in iis duo parva, albiantia > fuperne c^erulea •> cornu inferta, & in illius orificio foliis obte&a; cum illisieprodit *fylus* latus₅craffus, viridiufculuS) qui fuperne nigricat ac nitet: flores prasterea circa cornu & fo- lium rotundum > duobus fuifulciuntur foliis cupidatis, viridibus, c^rulelcentibus & purpurafcentibus. Ex *planu* hacadditis confimilis energies nonnullis aliis potioprapa- ratur, abdominis tumorem ex flatibus diminuens; in eun- dem finem ex ilia fola balneum conficitur capitii inun- guendum.

T A B . 70.

PhniahaeccUaquacka, lnefoliiss vocari poftct fknt a aquatic* ifhjlbt rtſins, flore caeruleo diformi.

Y y y

/OT,y:


سیفیں / سیفیں

کوئی دیگر نہیں

Ridder

نام پالی

بُنْجَانِ بُنْجَان



T A B U L A L X X I.




Afcitur in aquofis. [^]*qdix* fibrofa, albicans*
Tauks ex radice oriuntur, angulati, viridif-
 fufci, infra adradicem albicantes & transpa-
 rentes > angulis interdum spiratim contortis,
 tenues funt, rigidiusculi, re&i, intusa quei.

Cum Mis fk folia proveniunt? oblonga, cau-
 les primiim ampledentia > lateribus claula, latitudine pol-
 licem transverfum vix fuperantia, craffiola? viridia? in-
 terius nitentia, inferius aqueo albicantia ₅ venis in longum
 flriata, apertura[^] laterum fuperne evanefcente , faporis
 aquei. *Flores* m capitulis videntur oblongo-rotundis>rubro-
 tpadiceis ac nitentibus, foliis imbricatis, &fcuteJJatis, qu[^]
 iliquo modo rotunda & cochleato cava, obfiti," ex fingulis
 Horum orificiis, brevi * piano, Jatiufculo •> rotundiolo
 pctiolo erumpentes, primò ex inferioribus, dein ex fupe-
 rioribus; funtque tripetali, unguiculis ftridbioribus, finulis
 ceu crenulis nonnihil crifpati, colon's flavi? acinfarferici
 ^i tent is. *Stamina* in illis iex₅ plana ? tria latiora > apicibus
 oblongis, furrediis inftruda₅ unguiculis foliorum inferta,
 ftaminulis denfecapillatis flipata: triatenuiora₅ ex medio
 florum prodeuntia .. tribus latioribus interpofita , planis
 apicibus nodulata. (*femma* ferninalis ₅ inter fingula capi-
 tulicochleatafolia₅ oblongavidetur, qu[^]interius planaj
 exterius[^] convexa ₅ futuris viridibus > uno in medioplani,
 duobus in convexiori latere , ftriata , continetque in fe-
 numerofa *femina*, per membranaceum corticem transpa-
 r6ntia, parva, nonnihil oblonga, flava. Foliorum fuccuf
 cum *aceto* mixtus impetigini refiftit. *Folia* cum *radke*
oleo incoda? contra lepram fummutur; cum *ZMungo*
 decoda & epota fomnum conciliant.

T A B. 71.

Videtur hxc planta aliquam cum *Gladio lo pa-If ucai* vocatur, cum ex fguotamendifcrimjne[^]aft
 '4ri habere communionemj flos enim xque ac [quoniam ipfius defcnpno, ut & noftri authoris,
 illius tripetalus eft. *Pifo lib.6. ctp.56.* fimilis plan-] nullo modo fit perfetta, non ulterius niflex dc-
 ^dat dehneationem, qaseap Hd Brafilianos Ju- 1 lincatione judicure potin.



•JM IN-A N G AN L

T A B U L A LXXII



lingua Bramannum nZAdatsjadada vocatufj plantula eft lēfquicubitalis, arenoib gaudens folo- %tdix fibrofa > capillata > Jignofa, ni- ricans. Cauks quadrangulatij geniculatij Jignofi, longiufculis albanticibus pilfs obflti, in geniculispaffim caulinis lateralibus binst & ordine fē decuffantibus coronatim accindti. Folia in geniculiscaulibusinfidentbina, parva, oblongo-angufta, anonstenuitercrenata₃ tenuia₃ mollia₅ pilisrarisveftita. Cola media cum lateralibus in adverfa parte eminefc Flores numerofi & congregati in capitulis oblongo rotundis> quas caulinisinfident₃ acfoliis fuccinftafunt, proveniunt, ex caulinis foliofis ac pilofls erumpentesj iuntque minus piloii, monopetalii > quinque cupidati > candidi , dein ca;rulefcentes. Stilus in medioadeft in quo quinque fia~miniiU confpiciuntur mīnūtissimaj coilulo interiori florū inferta, apicibus, primum albanticibus₃ flaviufculis, ex in nigricantibus dotata. In ipfis gem mis flores claufi funt, cupidati •> viridiufculi; illorum odogravis ac male olens. In ftngulis tria quatuorve continentur Jemina parva, ex albo nigricantia. In butyro cumaliis antinephriticiscon^ coquitur, & mente alienatis pro potu exhibetur.

T A B . 72.

Plants IIIC anomai* eft fine confpecie, flos 1 depiaxit ; racari potefit sUmafik. (Mtfifc^ē₃
lurncn bene oVerraviEj quadripartitutn enim | dido.

Aa a a

TSfB.

Tsjeu-nien lae.

وَرْقَةَ الْمَلِكَةِ مَل.

جُهْرُفُ وَرَقَّةِ أَصَابُ.



Malva sylvestris (L.)

T S J E R U-U R E..N.

T A B U L A L X X I I I ..



Lant_#a eftaltitudinetriumpectum, nafcers
in arenofis. *R^dix-* fibrofa , albicans. *^ami*
ex parte virides -> ex parte rubefcentes. *Folia*
inramishincideproveniuntj oblonga, *in*
oris acutè crenata. *Cofla* media cum latera-
libus in adverfa parte eminent, redtam ftrians.

Ftores plures congregati cacumini ramorum in fidentjcyathiformes, monopetali > quinque partiti, venis in longum ftriati, in quorum tra&u rofacei funt coloris₉, inferius ad unguiculos flavi. *Stamina* jn illis novem furrect:a> flavis apicibus dotata cum breviori *flylo*. (*jlobuli* feminales virides ,* pilofi> dein ruffo-rubefcentes ? in vertice parum *cufpidati*? quinquelacunofis futurism in quibus fē relaxant? ftriati. *Sem'ma* inbilcecontinentur quingue, ad fingula latera inter futuras unum > interfepimentis fejunc:a> trilateralia funt₅ uno exteriori rotundo, duobua planis. Integra planta₅ excepta radice, cum fimilibus ejufdem energise in *oleo* decoquitur , & efHcax , pro morfu ferpentis aquatici, remedium prasftat:

T A B . 73.

Diverfe ejufdem coffnominis *Uren* ', in hoc I aliquomodo affimilantur ; vocari potefl *flams* bperc inveniuntur plant_s;, qu« nullam tamen | *sideritis folio infipido, flare pentdpaoit», rofacet «/;,,, inter fe habent convenientiam, & omnes plants | *futiu rotunds in quinque parm dmsafemne tmngulat**. funt anomaly. FoJia *Tsjem-Uren* illis *Sideritis* I*

B b b b

P E.



جعفر
دیگر
بـ

I R I B E L I

T A B U L A LXXIV.



Ux .hujus plant[®] funt species; unaejusnd-
 minis di^{cta} *Iribeti nigra*, altera *Vgmacciam*,
 ditta • *Irtbelt alba*. Prior à Bramannibus *Cd.*
rabalo voca tur, & eft plan tula duum trium vc
 pedum₅nafcensinarenofis. %*dix* admodurii
 fcbrofa&capillata, tenue intus & Jignoñurii
 continens filamentum, cortice amaro, exficcata colon's eft
 cinereo-obfcun, a quo ad differentiam %*tmacciam* ceu *Iribeli*
albx, nigra nominatur; odors vegeti eft & grati, *Iribelique aU*
*b**, quae minus odonfera eft, prxfbrtur; ilia vero qu[^] in re-
 gno Rapolin invenitur, odoris pr[^] *cxtcris* eft veg[^]tioris, &
 maximi facit; aft qu[^] in promontorio Comorim non
 tanti. *Cadex* cum cauhculis *virides* funt ac rubefcentes, intus
 valde aquei & virides, raris pilis obfiti. *Folia* in petiolis lon-
 gis, viridibus, languidis proveniunt, rotundipla ac parum
 oblonga? crappa, aquea, mollia , lenia, rarioribus pilis
 obiita, viridi obfcura. *C°P* media cum lateralibus in ad-
 verfaparte eminet, redam ftriants. Sapor aqueus eft &;
 amarus. Planta proved:ior fa&a caules emittit in quorum
 fummitate *flores* Ipicatim proveniunt. Propagatur ex in-
 fitione caulum₅ /eriturqueJ rrjenie Majo, imo & totomno
 inarenis albicantibus diflblutis, quo radices, ob odorc^m
 in mercaturis expetit[^], liberius fc diffundant, quin ut
 odore vegetiori & abundantiores proveniant radices, irfdi-
 gen[^] amputatos caules vafe teftaceo ? plurimis *in locis per-*
*forate** & arenulis humidis repleto, quo radicum demiffioni
 aditusperviusfit, imponunt, atqueadhunc modum tria
 <juatuorve vafacumfuis infitis caulibusfibi mutuofuper
 imponunt? quantum, arenulis incumulum congeftisex
 omni parte circumvallant. Septimo à fatione menfe radices
 eximuntur. %*dix* in decodisadhibetur anticacheticis, prfC-
 fertimprofefrecalida&ardorejecorisj ut 8c manuumac
 pedum tumoribus? etiam pro flatibus; dolores Capitis lèdat,

fitritacumlaae muliebri mifceatur & capiti inungatur-
ctc«m cum aqua calida epotum febri calid* medet. MS *
 don! jecoris'plurimisque malis; acnon difcnwrficpurown
 u materia p^rmixt^ habitum infuper coxript gravem
 & fetidum Vprodcfque in oculorum dolonbus. 0/^» «
 radicc per deftiUationem extraclum, pnxnum turbidum
 cftacnigricans, dein, foce alb.cante ceu c.nerestadtun
 dumdemifla, clarum & transpareos, pulchrc rub^cn
 & rutilans, odoris pergrati, fapoirts : *fubacri* s, i
 tione fimul cum aqua aicendens eique innatans.


T A B . 74.

Multa, in descriptione & teUnettfonchuj. plow, *defunt, atque ideo de illa certi quid vix*
dki pot; ft.



Dau-nay L
Dau-nay L
Dau-nay L
Dau-nay L

દાનાય



T E R I M-M U N J A

T A B U L A L X X V /



Renoifā amat. 7{*ami* rotundi, virides₅ pa-
rum lignofi & glabri. *Folia* in petioJis ob-
longisy rotundisj tcnuibusproveniunt, ad
exortum duobus parvis anguftiffimis foliolis
comitata, funtque oblonga? adpetiolumlato
ac rotundo ambitu₅ anterius anguftiora₃ in
culpidem ftrida, in orisferrata, tenuia fcabra₅ odoris
grati. *Cofta* media cum lateralibus in adverfa parte extu-
berant> etiam in refta₅ fed non tam alte. *Flores* nulli
funt. *Folia* cum *Calamo aromatico* & *Allio* in aqua *oryfye*
tritalinimentumpreftant antiipafmodicum, quod tanto
efficaciusjfi *radicis* pulverifatus *cortex* cum *orizje* lotura
hutyroque indies afolumatur.

T A B. 75.

Mores quoniam hie defintut&fruaus, incertum eft ad quod genus planta haecreferridebeat.
Folia affimilanturillis *Lamii*.

Dddd

TAR-

Turkestan

Salvia

جەزىئەر

تىرىپ



TAR7)AFEL.

T A B U L A LXXVI



Lanta humilis eft, arenofo gaudens fold.
 %adix ruffa? fibrofa? intusligno/a. *Caules*
 quadrangulati, nodofi>pilofi, virides, dein
 rubefcentes > adulti rufe/centes, exficcati ci*
 nerei. *Folia* bina & bina iē mutuo decu/Tant,
 caulesin nodis inveftiunt) parva funtj ob*
 longa? craifiola, ridigiufcula, criipa, piloia? fuba/^era*
 maxime in margine jin adverfa parte *cofta* & *co/iulis* ftriata*
 fufca. *Flores* in furculis fupra ex origine foliorum qua-
 tuor, fex, etiam bini & terni proveniuuit, purpureo-
 pallidi^ tetrapetali, cuipidati, non nihil pilofi> venuis
 in longum ftriati > odoris expertes. *Stamina* in illis quatuor
 albiantia, apicibus furre<3:is, atque ex viridi c^rulefcen-
 tibus ceupurpurafcentibus; *flylus* longiufculuseft, albi-
 cans, in vertice bifurcatus. *Calyx* quadrufolius eft, ex-
 terius pilofus. *CapfuU* feminales oblongas, rotunda, crat-
 fiote ac pilofej colon's perfici, funtque primi imipfiffimi
 florum petioli; in fngulis femina continentur duo oblon-
 ga> tenui membranaceo interfepimento fejuncSa_x exuna
 parte convexa* alterapiana? fulcata, colons ruffo fufci.
Succus exprelTus cum *Hutyro* decodtus lienterite prodeft.
 %otdix antidotumeft, fitritape<3:us cum illa JInatur; qui
 humor in pe< Sore remanens nulla veneni eft fuipicio₅ fin *
 certiffimum eft fignum venenum fubefle.

T A B, 7d

Delineator caules non bencffigiavit, quadrangulati enim funt₅ ille veto rotundo depinxit₀

E e ee

EN.

E N T A 2) A.

T A B U L A L X X V I J .



Uffrutex eft, nafcens in arenofis* %ami virides, rotundi, lignofi> rigidi, ftriati Vcūm furculis quos diffundunt, undiquaque foliolis parvis, viridibus, anguftiflimfs, cuipidatis > rigidis veftiti, quas fe in ramis majoribus verfus inferiora refle&unt: rami cum foliis viorem longe fervant. *Flores* capitula funt oblonga ex foliorum imitatione penniformia? viridi albicantia? furculorum cacumini infidentia.

T A B . 77.

Imperfecta hujus planrx defcriptio in caufa eft quare non re&c poffit agnofci. Floculi qui in (icas proveniunc , fecundum delineauonem,	pcntapetali funt, ftaminibus in medio dotati; fed qua: forma fie feminnm vel capfularum femi* naliu, dc illis nc \$v> quidem*
---	---

P O L A T S J i R A.

T A B U L A LXXXVII1.



Sjira species est & planta aqua

Iplurimis & quidrm nfbis *:ojiirt:int tibris, pe.
uics caulfum, ((ufous super terram repit in
Igtrnicijjs circurrwdbcntcs. Cvltt totundi,
IcriiJsjij v*crist:u(pid)cm (enlImitrichtitjrcs_T
viriJi-cJarJiniJ.Cntcs,aqtfc, (unbolt, |-cnk
transparenlibus & parafeUj's itrtatt^ gcaicuJjatii mttjrmodiis
sensim icycriis apjeeni minucntfSus, ex pedibui novoa
cau Jcs ejicicntesiutus iater(tip!n^:atis> qua: f^ % iu mcJin^
tenui » pfaoo , rufib-rubro cveunt & t-orti^i c-stcriorflc
affiguntf J(l ctulliceCK tubulosdeftinGt. FitU CiUlcalbr
tim in tjenicuJii gentinata circuinvc/IJUot * oblongo-mi*
gp{(a, cuspidiirkacuthdiinticuLisocdLnate fctcta, icnta^
glabra, Fu-idinicentia. t o^ media ip atlvcrfa. parec cmi-
a et. Ffortt pi u res con ^regiti tn {jen icutis tup ra txortunt
folii>Fumjprodunt,CJuJibufjrtritcniidentcs, parvi, penra-
pectali , albo dt^uci i Jit:Jjiformes, (^uirt^ut fibvividitoiw
Fohis cuJpidattsfucpnndti, *Stamina* in iUis III-CCIII albican-
tia^ gtmmulrs flaviufciilis rwlukca* ghtiuluai iiicdiuin
virVJeoi ampj^entii ijinbtdi Jcnunncls iubrubri , tjuin-
fjuc JUurus ftcuti* ifttcrius tjuin^uc vifiditifSjCuIptiktia
foliis cstepti *Stums-* infUfinumerohn par^j) primutti
albicantii Li dctQcxftlibLcanterubcfccntta.

T A B . ?S-

Planta hrc est aquatica, vocari p-
nida, juncos ministrano, ex alio rufi, cente,

G g g

P E E.



Pee-tardáve Linn.

وَلِلْمُرْسَلِينَ مَل.

جَنَاحَةُ طَارِخَانَوْهَلَهُ

Arab.



? E E.TA R T> A r E L.

T A B U L A LXXIX.



Lanta hasc arenofa amat. *'Radix albicans*⁵ fibrofa, fubflavefcens, intus lignofa. *Stipites* nonnihil tortuofi accymis inflexi; *vividi* & hincindefubrubefcente corticc, fuperiores rami virides, fpiculis brevibus, afperis, fparfim obfiti, ad cymas turgefccntes, venifque *ihkti* *Folia*, cumfuisbrevisbus₅interiusplanis,petiolis₅inramis proveniunt, rotundiola, parum oblonga, tenuia, mollia, lenia. *C^oftula* media etiam invertice eminet. *Flores* in capitulis globofis fpicatis₃ fafligiis ramorum infident, ecu etiam ad latera hinc inde adherent, funtque obJong^, cut pidatas fpic^i 5 pentapetali, candicantes, tran/parentes, femivirides, in medio cuipidum viridi-fufca vena ftriad *Stamina* in illis quinque, oblonga, albicantia, *furre&z*, purpureo-rubrisgemmulisnodulata; *Stylus* fuperneppureo-ruber, ex gemmula viridi albicante emergehs. *Flores* prasterea ceu fpic₂ tribus cuipidatis viridiufculis foliolis, qu^ein filamenta oblonga & crifpata definunt, fuccinguntur, odoris expertes; fiorum feminibus excuffis, jpcicas apert^feclaudunt, gemmulamceu capfulam oblongam feminalem intra fe recondentes, quas in vertice filamento oblongoeftdotata; in fingulis fpicis? intra propriam capfulam membranaceam *femen* unicum includitur parvum, lateribus utrinque convexis, qu^e in oram angulatam coëunt, colonis atri & nitentis. Incantatis opitulatur h^cplanta.

T A B. 79.

Tce-Tardavel nullam habet convenientiam cum *TArdavel* jamanteadefcriptai Scplantafui gen-
ris eft, fine confpeeie.

Hbhh

KA.



Kara-Pito l.

Mal.

سْ قَحْلُو طَفَاعِي

K c * L V-T A L L

T A B U L A L X X 3 C



Afcjtur inarenofk %idix alb/cans, fiiro fa. C-mlis aijuci, rotundi, rubcicujr^{cs}, pUis h iriu ti's */pcri. fotu o Wo oga, an tcri us culpidaM 9 "no Jatcrc mfcritus htius ex por^r^ito, .it- rotimdiorjs ambitus 3 piJis Jurfutis, qui afpcrttstem inducuntj obfitQ, viro-

pctjolis iiipraexortum foliorum prodcunc plufcs conErc^ati, pcduncuJis brcviffimirs **infidencesj** i'untque in gem*tnisvirides, cupidari &: Bngulati> cv una parrc inferius in a [am flncti; jpcrti bipenfj ijnt, in :tcj uaJcs > rubro purpurei, gcmmuia cat qua falij prodcunt,i:oii/hintcs. tflAv cui jpfmmula iniidct quingucFoItaeft, cu/pidatusj **vin-dis**. *Fruktus* noo tcr. Qtdictum ccphatfttgiamopomtt forus ex hoc frutice præparatus.

T A E. 30.

Kara-Pito sive defens. inv off I*** Miftti* in pilosa, ad quod delinctor non bene expedit.


lxxi

EAE.

Ené-páel lat.

போக்குவரதை Mad.

البيهقيّة عرب.



Botanische Zeitschrift

E N E-T A E L.

T A B U L A L X X X I



LTmiliſ eft planta arenoſo humidoq;^
Igaudensiblo. *Caules* tenues? rotundi>
Igeniculati, virides₅ in geniculis oſto
ſfilis oblongo-anguftis₃ cuſpidatis₅ ri-
* gidiuſculis > ftellifortniterexpanfis, ac
cum cuſpidibus verius pedem caulfum
infledlis yerticillatim cin^fci. Cauſes
foljoi ex geniculis geminati prodeunt; adfmgulos geo-
culosuper folia quatuor *gemma* rotundiolae? viridi-albi-
cantes caulibus adbſerent. %utyro recenti commixta ab-
fceffus haec planta ad maturitatem perducit.

T A B . 81.

*Imperfecta hujasplant*defcripvo ineaufa* eft, quare nihil ccertidcſpfapoflitaiKrmarij quugd
externam phafin iſe c i ſte* refcrcm.



K k k

NE.

Nélam-múri lat.

وَسْوَسَةً مَال.

سِنْجَارِي

سِنْجَارِي بَنَم.

سِنْجَارِي تِلِّي عَرَب.



Sesbania aculeata

N E L ^A M-MtA R I

T A B U L A L X X X 11.



Lanta humiliseft, furculis in latum diffusis* arenofo gaudens fundo. %adix albicans* fibrofa & Tignofa. *Stipites* in plures fparguntur furculos virides & dein rubeicetites. *Folia* in furculis hinc inde in brevibus₅ tenuibus, viridibus petiolis_? bina & binaproveniunt₉ anterius con verfa, oblongo rotundo-angufta? parum cuipi-data₅ craffiuſcula, denia? plana, lenia, viroris in re&a pane fufci > in ad verfa clari, pundtulis viridi-fufcioribus interlperfi ;. *Co/la* media in adverfa parte altiim eminet; redlam ftriāns. *Flores* in propriis videntur petiolis, qui paffim ad exortum petiolorum foliaceorum erumpunt, funtque in illis feriatim confiti, ex calicibus furre6lis emicantes -> bipetali_? fibi mutuo accumbentes. *Globuli* ex finguliscalicibus proveniunt plani₅ viridiuſculi, parum oWongi, tres quatuorve in una ferie, viridi fufco filamento finibriasannexi, pilislongiuſculisobſfiti, cortice rigiduP culo₅ unicum finguli continentes *femen*, rotundum, non-nihil oblongum •> primūviride&nitens) dein nigricans₀ Apozema ex hacplanta pro algidis prasparatur febribus.

T A B . 82.

In fepius jam memorato HerbariovivoCey- j viUofu* art'uulatis; fecundum defcriptionem talanico , flmilis farinas plantam invenio fub no- men, aliquod videtur incerecdrc dicerimen. mine l>L:nu bifolU bumirepa, Arifolio, floſculisaurás, |

Mallam-tsjulli Lat.



dMALLAM-TSJULLI

T A B U L A LXXXIIT.



Lanta hurailfs eft, arenofa amans. *fydix*
albicans •> fibrola. *%mi* Jignofi j tenues,
virides. *Folia in petiolis viridibus* •> ac fer-
ine pojlicera transverfum loogis > qui ad
exortuna duobus valde brevibus oblongo-
anguftis fuccinguotur foliis, geminata pro-
venfunts oblongo-rotunda, angufta, tenuia, nervis per-
duS a virorisiaturi&obfcurL De viribus nihil conflat.

T A B . 83.

Campraccdentipiantulahate *bUUtm-TsjuUi* videtur aliquid modo convenire, fed imperfe&a *de*
feiptio facie j ut aon niC conjefturando de ilia judicari poflic.



M m m m

BE.

B E L I ' T S J I R

T A B U L A LXXXJV.



Atcitur in arenofe & umbrofj>. Radix fi-
brofa > albicaos, Qtodei, qtii ex radice **piures**
provenfunt, kte juper tcri jm xpantluntufj
craflilDt8c gcnicuan* cxgensculisnovos
caules ejictepesi«queij vindkfifuti, nitcntcs,
& tx parte rubefcentcs. *foit* gcniculis
cdulium gcrninaci infiJcnt, oblongo-rotunJaj »ngulU<
in vcriic obrule cupidata , cenuia s Jcnia» niollia, vfro-
ria in *Tcia*. partc fijfciorisj in advcrla cjiEutiofs> ubi wjU
modincmialt. *FhretQonl* gregati Jcptem* otbjDovnn*
lupra cxoriginc tbliorum provciiunt i virides* cyathu
formes» inorificio quadranguJatij tctrapctdijinfijeuJis
ora: angu 1 is v i ridi cupidc cm me n tcs. *Fntdmf* crt n ullc*.?
B.ilneum ex hac phnra confidtur dexkacum ndvcrfus
omnigenorumfcrpen turn virus, ft/w cunifuccocortias
L,iH/f iivigata illi't^quc» car brici s , [>hlcgtnique
medentur.

T A B . 84.

ffidtcspia.iiiJrrr;illIJLLUlt' fcuiuliliu hujl phante ceri quid coelitae, de sic ex argo quid
jafivUij^jVrLj jij;:i...luUfi 111* iflus; , ptemoribus habens relinguo.

N n n 11

TS/^-

TSJERIA-MANGA-NARI.

T A B U L A LXXXV.



RenoJa a mat. fydix albincs , iibrofia
Tfji/fj trianguJatb viridcs, intusaqud- /olia,
t[us caulibus indent, oblonga, a ilpidj
in onsan^ulisajcutis (iirrata-t teauti, krxin,
in advtrJa parte coflia cm incur if) us obiitaj
n u JL us faporiti. Fhrcts, q u i;in pctiol is ob-
longis iupra cKortum i"oliorum» in cacumiac caulium
crumpunt) can-ulti fufit, monopctai» tripartite StyInt
ickksm casrokus eft capituo rigidiufculo, Ceijx bxus,
cjuiiH]uc folius. Linimomum ex hac pjjmta sjtque Olea
Nucis Indie;u t-onfectum, elephantiacis opituhtur, Suaus
cum BinXfbrc & Cumiw exhibit us, vcltiumcum Of to
Scrgtlirit & Cf&amt ArwutitQ corporf iJItus, lummopcre
in pdtilcentiali tonducit icfar.

T A B . 85.

10 caripelle herc plantas. : "UJ MUHJ. 200 Veronic India-s, flore carmine, Cnn^Jrf i;r, riji 1 omnes
in UJ gbb ouue noue cb milcfilbcM, <Ju't x delineatione tu; jiuav x tci scriptione apparent.



Oaoo

TSJE.

1 t



جعفری - بیانیه

۳۱۴

ساقی

جعفری - بیانیه

TSf E%I A-H^A%lH%A<£MT^eO Li.

T A B U L A LXXXVI.



Afcitur iiiarenofis&petrofis. %adix aquea* carnofa, albo-ruffefcens, fibris denfis veftita, faporis aquei & fubamari. *Canles* rotundi, Ianguidi, carnofi, aquei, fufcè rubefcentes, nodulis turgefcentes, vetuftiores fpadicei, pilis longiufulis, raris fparfim obduftij fubaſpcri? faporis acidi. *Folia* pedunculis craffiolis, ru^ befcentibus, qui parvis, cuſpidatis? albicantibus? membranaceisfoliolisad exortum fuccinguntur, prōveniunt> oblonga, in cuſpidem definentia, uno latere latiori, ac ad petiolum verfus inferiora longius exporredip, quo ex petioſo annularidu<5luexcurrunt> funtque in oris crenafa, mollia, inrefta&adverfaparte) in qua Qftd* eminent, quſeetiam redlam ftriant, pilis hirfutis, longiufulis albi° cantibus obſita, nonnihil afpera, viroris floridi & nitentis* *Flores* in petioliſ ſufce rubefcentibus qui fuſra exortum foliorumerumpunt* binivel terniin unoquoque videntur petiolo, tripetali > candidi»fcintillantes, pilis longiuſculis fparfim obſiti > ex arflo *calycis* orificio erumpentes* quioblongus eft&triangulatus) angulis alatim expanfis, inalaruminterftiis, venarubrafriatus, ut & in alarum Jateribus. *Tenicillus* ex floris orificio emicat trifidus, cuius fingula portio iterum bipartita, ac globulo viridi inftruſta. *Gemmce* florum rotundiolas, parum oblonge, complanatse, exterius piloſo. Prater hofce etiam flores habet tetrapetalos? fine calyce in quorum medio *ftylus* ftamunculis, qu<< flavis funt inftru&a nodulis, penicillatim circurredatus. *Baccte* in fingulis alarum calycis lateribus, circa medios illorum angulos) ubi alae albefcunt, inveniuntur copioſifſim^, membrana calycis obteſtas> & in illis copioſiflima arenacea, primum albicantia, dein rubefcentia *Semina*. Indigene hac planta ? excepfia radice, utuntur In concoftione *Card,*

P P P P

Fo|

*Folia oleo inco**b**> liniment urn fuppeditant vulnerarium:
ex iifdem calefadtis > addito *Sale* fit nodulus > pro putridis
purulentifque inferviens dentibus.*

T A B . 85.

Pertinet hxc pluu inter herb** btcciferu; Acmeojudiciocomflari poteft* JtU«*fl^{1M*}
tic*, flvt&ttciii i>iifutmiiui.



بَعْدَ بَعْدَ

بَعْدَ بَعْدَ

بَعْدَ بَعْدَ

B A H E L - T S J U L L I.

T A B U L A LXXXVIL



Lantula eft nafcens in fulcis agrorum, duutn pedum plus minus altitudine. *%adix* oblonga j albicans ₅ fibris veftita. *Caules* digiti craffitie; *Surculi* quadrangulati & geniculati> in internodiis alternato ordineplani, fulco ftriati, *cortice* glabro & viridi. *Folia* inge- inculis ? iupra plana internodii latera > geminata prodeunfr ordinefemutuodecuffantia? furculisinfidentia, oblongo- angulta, in oris cupidibus crenata, fpifla & mollia. (*ofta* media in adverfaparte altum eminet? laterales nonnihil, qu2B ad oras delatse evanefcunt. *Flores* plures congregati in cacumine furculorum proveniunt? in fingulisgeniculis gemini > fupra ex lateribus planis ₅ ordine decuflato, brevi, craflb, furre6to₅ viridi-diluto, appendentes petiolo *t* qui parvo, cupidato > viridi fuccingitur folio; tetrapetali funt> rotundioli, exterioraverfusinflexi, cum collo brevibus piliscapillatijceulanuginofi; colons perfico rubri; petalorumunum oblongiuseft, & interius perfici dilutions > exterius cum collo albefcens; alterum huic oppofitum brevius &c latius> ejufdem coloris, fed magis faturi, exterius perfico fanguinei ₅ ut & in collo ₅ quod ad partem folii oblongioris, intusvenis ftriatur purpureo-rubris. In collo quatuor conlpiciuntur (*lamina* Candida; illorum duo arcuatafunct&velutrofac^ruleaj *in* medio arcta &ftrida, quas folio latiori accumbit; duobus veluti brachiolis incurvis, quorum extrema in latum ac plane purpureiim defi- nunt nodum > fe mutuo obverfa, paulo infra brachii nodos collo affuta; reliqua duo ex medio collo à latere folii latioris₅ ac paulò inferius emicant, ad folium latius accum* bentia > furredla, arcuata, verfus fe invicem inflexa > ac rolà casrulea adftridla; per quorum infertionem ad folii latioris interfittia ? albicans in collo tra<Stus fe fignat. *Stylus* ex umbilico colli florum exfurgit candidus , in vertice

Q q q q

bi-

r HORTUS MALABARICUS.

bi/idus, folio htiori ar<5tc accumbens , cum ftaminibus arcuatisroiamamplectens. *Calyx* quinque folius eft, cu-
pidatus oblongus, cra/lloJus. In (/cmmis iupra folium oblongum, folia *quuxci* ad latera confidents convoJuta jaccnt, *ik* fupra ea folium latiusj funtqucquoadmaxi-
mamillorum partcm, purpureo fufcas, adfblii oblongi
partcm albicantes. (*jemmx* fcminales grandiufulcas₃ ro-
tunda;, viridcs, calycilateccxpJanatoardlc infidcntcs, in
verticc tilamento rulFo dotatas, vel illodcciduo> cuipidc
parvo, durioio, acuto, perverticcmiubtili fulcofafutu-
raftn'ati, qua reJaxata induas ic apcnunt partcs: cortex
rigidiuiculus& cartilaginous cvadit: *Semhia* in illisconfi-
dent numerofa> minutiillmaj nonnihiloblonga, globu-
li medii rotundi, qui viridis ac intus luimiduscft, fuperfi-
ciem obfidcntia > primum candidiffima imo fcintillantia,
dcinrufFa; globulus iJle mcdius intcrfepimcnto cingitur
annulari, viridi-aJbicantc ac tranfparente , ex % Io pro-
dicns, & futuras mediae gemmx ic annclens ; annulus
proximo ad vcrticem globuJi liliatu intcrmedio fempcr
apertus eft. Notandum prxterca quod gemmae hae iemi*
nalcs, flores? quiiniurculisinferioresiuntfubfequantur*
dum adhuc ex iurculorum vertice flores noveJji produ-
cuntur. *Fo/u* tn'ta 6c in formam cataplalmatis uiurpata
ab(ceflusmatnrant. Ex *radice* abluendis capitis ibrdibusj
ccphaliciim prasparatur lixivium.

T A B. 87.

Pbnum hint ego nomicuicm *Digitalis agni India, Eleatris folia, flere rabiunde.*



INDEX

'Prior numerum paginam, posterior tabulam dejignai.

A.		
	Da-Kodien.	9-7
	Adel-Odagam.	81-43
	Aria-Veda.	41-23
B,		
^Bahel-Tsjulli.	171-89	
Beli-Tsjera.	165-84	
Belutta-Areli.	3-*	
Belutta-Kaka-Kodi.	maanapay: 7-f. 6	
Belutta-Onapu.	99-51	
C.		
r Car-Elu.	J07-5S	
Cara-Carinam.	109-56	
Cara-Veela.	43-24	
«Coletta-Veetla.	77-41	
Corofinam.	133-68	
Cupa-Vela.	61-33	
E.		
Ene-PdeL	159-81	
Entada.	scandens	IJi-77
F.		
Futa-Meftica-Incarntata.	1-1	
Fula-Meftica-Alba.	3-2	
I,		
Iribeli.	145-74.	
K.		
Kaka-Kodi.	n-8	
Kaka-Pu.	103.53	
Kalu Tali.	157-80	
Kametti-Valli.	z ₃ -i ₄	
J [^] alu-Polapen.	IZ9-66	
Katu-Karavi.	83.44	
Katu-Pal-Valli.	17-11	
Katu-Pee-Tsjanga-Pulpam.	113.85	
Katu-Viftna-Klandi.	119-61	
Kattu-Tagera:	55-30	
Koda-Tsjeri.	131-67	
Kondam-Palu.	57-31	
Kotsjiletc-Pullu.	139-71	
M.		
^lan-Onapu.	95-50	
Man-Todda-Vaddi.	39-iz	
Manja-Kurini.	1x1-62	
Malatn-Todda-Vaddi.	37-ii	
Mallam-Tsjulli.	I63-83	
-Manneli.	69-38	
Miri-Angani.	I4.I-72	
N.		
Nansjera-Patsja.	21-13	
rNelam-Muri.	J6I-82	
Nelem-Pala.	• 5-H	
Nelem-Parenda,	117.60	
O.		
Neli-Pu.	137-70 [^]	
Neli-Tali.	31-13/	
Nellia-Tandali-Cotti.	49-2.7 [^]	
Njbta-Njodem-Valli.	1.7-16	
Nir-Murri.	59-3*	
Nir-Pullari.	67-36	
Niti-Todda-Vaddi.	rectores 35-ao	
Nottenga.	101-52	
P.		
Pal-Valli.	19-12	
Parparam.	29-17*	
Pee-Tandale-Cotti.	53-*9.	
Pee-Tardavel.	155-79-	
Pee-Tsjanga-Pulpam.	"\$-59}	
Pee-Tumba.	.87-46,	
Perim-Munja.	'47-7*	
Pola-Tsjsira.	153-78/	
Pul-Colli.	135-69,	
S.		
Schit-Elu.	105-54	
Soneri-lла.	127-65	
Suendudi-Pullu.	75-40	
T.		
Tali-Pullu.	123-63 ⁶	
Tandale-Cotti.	45-25 ⁴	
Tandale-Cotti.	47-	
Tardavel.	149.76 [^]	
Tilo-Onapu.	101-52	
Todda-Vaddi.		
Tsjanga-Pulparti.	111-58 j	
Tsjera-Ufen.	143-73 •	
Tsjeria-Manga-Nari.	167-85 i	
Tsjeria-Narinam-Pulli.	169-86 ^	
Tsjeria-Onapu.	93-49 ^	
Tsjeru-Manneli.	.73-39	
Tsjeru-Vela.	63-34 <	
Tsjeru.Urem.		
Tsiovanna-Aleri.	M3-73	
Tsiovanni-Manneli.	71-38	
V.		
Vada-Kodi.	79-4* [^]	
Valli-Onapu.	91-48 £	
Valli-Upu-Dali,	85-45/	
Vallia-Capo-Molago.	65-35 c	
Upu-Dali.	125-64	
W.		
Wallia-Pal-Valii.	15-10	
Watta-Kaka-Codi.	*5->5 [^]	
Wellia-Tandali-Kotti,	51-zC	

fa HORTUS MALAEARICUS.

bifidus, folio Jatiori ar&c accumbens , cum ftaminibus arcuatisro&mampectens. *Calyx* quinquefoliuseft, cu-
pidatus oblongusj cra/Eolus. In (*jrcmmis* fupra folium oblongum, fo/jia qux ci ad htera confident, convolura
jaccntj *ik* fupra ea folium Jatusj funtquc quo ad m^{*}
jnamiUorum partcm, *purpureo fufcse^ adfolti oblongt*
partcm albicantes. (*fe»m** fcminalcs grandiufcula:, ro-
tundx, viridcs, calycilateccxplanatoardlc infidentes, in
vcrtice filamcntoruffo dotatae) vel illodcciduo> cuipidc
parvo, duriolo, acuto? per verticcm fubtili fulcofa futu-
raflriati, qua rclaxata induas fe apcriunt partcs: cortex
rigiditiiculus&cartilagineus evadit: *Stmin** in **illisconfi-**
dent numerofa» minutifllma? nonnihiloblonga, globu-
Ji medii rotundi, qui viridis ac intus humidus cit, fuperfi-
ciem obfidcntia , primum candidiflma imo fcintillantiaj
dcinruffa; globulus iilc mcdius intcrfepimcnto cingitur
annulari, viridi-albicantc ac tranfparente , ex ftylo pro-
liens, & futuras media; gemmx ie anncdlens ; annulus
proxil I vcrticem globuli lliatu intermedio fern per
apertus eft. Notandum prasterca quod gemma: has /emi-
nalcs, flores, qui in Iurculis inferiores func fubfcquantur»
dum adhuc ex lurculorum vertice iiorcs novelli produ-
cuntur. *fo/ia* trita&informam cataplafmatis uiurpata
abfceilus maturant. Ex *rttiict* abluendis capitis/brdibusj
ccphalictim prasparatur lixivium.

T A B. %J.

Plenum hant *ego iominarem Digitalis agna Indica, Blattaria folia, flote radicula,*



INDEX

Trior numerus paginam, posterior tabulam dejignal:

 A. Da-Kodien. 9-7 Adel-Odagam. Si-43 Aria-Veela. 41-23	B, <-*Babel-Tsjulli. 171-39 Beli-Tsjera. 165-84 Belutta-Areli. 3- Belutta-Kaka-Kodi. 7-f. 6 Belutta-Onapu. 99-51	C. f Car-Elu. J07-55 Cara-Carinam. 109-56 Cara-Veela. 43-24 «Coletta-Veeda. 77-4 ¹ Corofinam. 133-68 Cupa-Vela. 61-33	E. Ene-Pdel. 159-81 Entada. I51-77	t. Fuh-Meftica-Incarnata. I-1 Fula-Meftica-Alba. 3-2	F* Iribeli. * 45-74	K. Kaka-Kodi. 11-8 Kaka-Pu. 103-5* Kalu Tali. 157-80 Kametti-Valli. X3-I4 Kalu-Polapen. IZ9-66 Katu-Karavi. 83.44 Katu-Pal-Valli. 17-11 Katu-Pee-Tsjariga-Pulpam. 113.85 Katu-Viftna-Klandi. 119-61 Kactu-Tagera. 55-30 Koda-Tsjeri. iji.67 Kondam-Palu. 57-31 Kotsjiletti-Pullu. 139-71	M. Man-Onapu. 5,5-50 Man-Todda-Vaddi. 39-iz	N. Manja-Kurini. 1x1-62 Malam-Todda-Vaddi. 37-21 Mallam-Tsjulli. 163-83 •Manneli. 69-38 Miri-Angani. 141-72	21-13 361-82 5-H II7-60	Neli-Pii. 137-70 Neli-Tali. 31-18, Nellia-Tandali-Cotti. 49-27 Njota-Njodem-Valli. 27-KS Nir-Murri. 59-32- Nir-PullarL 67-36 Niti-Todda-Vaddi. recte 3 5-io Nottenga. 101-52	O. Onapu. 89-47	P, Pal-Valli. 19-12, Parparam. 29-17*	3. Pee-Tandale-Cotti. 53-29 Pee^Tardavel. 155-79 Pee-Tsanga-Pulpam. 115-59 Pee-Tumba. .87-46' Perim-Munja. M7-75 Pola-Tsijira. 153-78 Pul-Colli. 135-69	T, Schit-Elu. 105-54 Soneri-Ua. 1x7-65 Suendudi-Pullu. 75-4°	1x3-634 45-*5 47- 149.76 Tilo-Onapu. 101-5X Todda-Vaddi. 33-19 Tsjanga-Pulpadi. 111-53 Tsjera-Ufen. 143-73 Tsjeria-Manga-Nari. 167-85 Tsjeria-Narinam-Pulli. 169-86 Tsjeria-Onapu. 93-49. Tsjeru-Manneli. .73-39 Tsjeru-Vela. 63-34 Tsjeru-Urem. M3-73 Tsiovanna-Aleri. 1-1 Tsiovanni-Manneli. 71-38	V. Vada-Kodi. 79.4* Valli-Onapu. 91-4S Valli-Upu-Dali. 85-45 Vallia- Capo-Mola go. 65-35 Upu-Dali. 125-64	W. Wallia-Pal-Valii. 15-10 Watta-Kaka-Codi. X5-15 Wellia-Tandali-Kotti. 51-28
--	--	---	---	---	-------------------------------	--	--	---	--	--	---------------------------	--	---	--	---	---	---

1 INDIAN 60		TANIC GfKOEN I
LIBRARY		
PROPS 60	✓	
ACCESS TO 4	✓	
CATALOGUE 60	✓	
CLASSIFICATION	✓	
DATE	15.3.64	2006

BOOK CARD

Indian Botanic Garden Library
BOTANICAL SURVEY OF INDIA

581.9(595)

Call No..... *RHE-1* Acc. No. *B2403*

Author *Rhede, H. van*

Title. *Hortus Indicus Malabaricus*
161.9.

Name of Borrower/Code No.	Date of Issue	Date of Return
.....
.....
.....
.....
.....
.....
.....

INDIAN BOTANIC GARDEN LIBRARY
BOTANICAL SURVEY OF INDIA